



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/861, C07K 14/075, C12N 7/00	A1	(11) International Publication Number: WO 00/15823 (43) International Publication Date: 23 March 2000 (23.03.00)
--	----	--

(21) International Application Number: PCT/US99/20728

(22) International Filing Date: 10 September 1999 (10.09.99)

(30) Priority Data:
60/099,851 11 September 1998 (11.09.98) US
60/136,529 28 May 1999 (28.05.99) US

(71) Applicant: GENVEC, INC. [US/US]; 65 West Watkins Mill Road, Gaithersburg, MD 20878 (US).

(72) Inventors: WICKHAM, Thomas, J.; 1203 Arista Manor Way, Germantown, MD 20876 (US). KOVESDI, Imre; 7713 Warbler Lane, Rockville, MD 20855 (US). ROELVINK, Petrus, W.; 17502 Gallagher Way, Olney, MD 20832 (US). BRUDER, Joseph, T.; 10243 Royal St. Andrews Place, Ijamsville, MD 21754 (US).

(74) Agents: KILYK, John, Jr. et al.; Leydig, Voit & Mayer, Ltd., Suite 4900, Two Prudential Plaza, 180 North Stetson, Chicago, IL 60601-6780 (US).


(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: ALTERNATIVELY TARGETED ADENOVIRUS

(57) Abstract

The present invention provides a recombinant protein having an amino terminus of an adenoviral fiber protein and having a trimerization domain. A fiber incorporating such a protein exhibits reduced affinity for a native substrate than does a wild-type adenoviral fiber trimer. The present invention further provides an adenovirus incorporating the recombinant protein of the present invention.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

ALTERNATIVELY TARGETED ADENOVIRUS

TECHNICAL FIELD OF THE INVENTION

The present invention relates to an alternately targeted adenovirus and
5 includes methods for producing and purifying such viruses as well as protein
modifications mediating alternate targeting.

BACKGROUND OF THE INVENTION

The various physiological responses of a host animal to the presence of a
10 virus depend on the different ways such viruses interact with the host animal, each
of which is first mediated by the surface of the virus ("the virion"). The
adenoviral virion is a non-enveloped icosahedron about 65-80 nm in diameter
(Horne *et al.*, *J. Mol. Biol.*, 1, 84-86 (1959)). It comprises 252 capsomeres -- 240
hexons and 12 pentons (Ginsberg *et al.*, *Virology*, 28, 782-83 (1966)) -- derived
15 from three viral proteins (proteins II, III, and IV) (Maizel *et al.*, *Virology*, 36, 115-
25 (1968); Weber *et al.*, *Virology*, 76, 709-24 (1977)). Proteins IX, VI, and IIIa,
also present, stabilize the virion (Stewart *et al.*, *Cell*, 67, 145-54 (1991); Stewart *et al.*,
EMBO J., 12(7), 2589-99 (1993)).

The hexon provides structure and form to the capsid (Pettersson, in *The*
20 *Adenoviruses*, pp. 205-270, Ginsberg, ed., (Plenum Press, New York, NY, 1984)),
and is a homotrimer of the protein II (Roberts *et al.*, *Science*, 232, 1148-1151
(1986)). The hexon provides the main antigenic determinants of the virus, and it
also contributes to the serotype specificity of the virus (Watson *et al.*, *J. Gen.*
Virol., 69, 525-35 (1988); Wolfort *et al.*, *J. Virol.*, 62, 2321-28 (1988); Wolfort *et al.*,
25 *J. Virol.*, 56, 896-903 (1985); Crawford-Miksza *et al.*, *J. Virol.*, 70, 1836-44
(1996)).

The hexon trimer is comprised of a pseudohexagonal base and a triangular
top formed of three towers (Roberts *et al.*, *supra*; Athappilly *et al.*, *J. Mol. Biol.*,
242, 430-455 (1994)). The base pedestal consists of two tightly packed eight-
30 stranded antiparallel beta barrels stabilized by an internal loop. The predominant
antigenic and serotype-specific regions of the hexon appear to be in loops 1 and 2
(i.e., LI or *l1*, and LII or *l2*, respectively), within which are seven discrete
hypervariable regions (HVR1 to HVR7) varying in length and sequence between
adenoviral serotypes (Crawford-Miksza *et al.*, *supra*).

35 The penton contains a base, which is bound to the capsid, and a fiber,
which is non-covalently bound to and projects from the penton base. The penton
base, consisting of protein III, is highly conserved among serotypes of adenovirus.

and (except for the enteric adenovirus Ad40 and Ad41) it has five RGD tripeptide motifs (Neumann *et al.*, *Gene*, 69, 153-57 (1988)). These RGD tripeptides apparently mediate adenoviral binding to α_v integrins, a family of a heterodimeric cell-surface receptors that also interact with the extracellular matrix and play important roles in cell signaling (Hynes, *Cell*, 69, 11-25 (1992)). These RGD tripeptides also play a role in endocytosis of the virion (Wickham *et al.* (1993), *supra*; Bai *et al.*, *J. Virol.*, 67, 5198-3205 (1993)).

The adenoviral fiber is a homotrimer of the adenoviral polypeptide IV (Devaux *et al.*, *J. Molec. Biol.*, 215, 567-88 (1990)), which has three discrete domains. The amino-terminal "tail" domain attaches non-covalently to the penton base. A relatively long "shaft" domain, comprising a variable number of repeating 15 residue β -sheets motifs, extends outwardly from the vertices of the viral particle (Yeh *et al.*, *Virus Res.*, 33, 179-98 (1991)). Lastly, about 200 residues at the carboxy-terminus form the "knob" domain. Functionally, the knob mediates both primary viral binding to cellular proteins and fiber trimerization (Henry *et al.*, *J. Virol.*, 68(8), 5239-46 (1994)). Trimerization also appears necessary for the amino terminus of the fiber to properly associate with the penton base (Novelli *et al.*, *Virology*, 185, 365-76 (1991)). In addition to recognizing cell receptors and binding the penton base, the fiber contributes to serotype integrity and mediates nuclear localization. Moreover, adenoviral fibers from several serotypes are glycosylated (see, e.g., Mullis *et al.*, *J. Virol.*, 64(11), 5317-23 (1990); Hong *et al.*, *J. Virol.*, 70(10), 7071-78 (1996); Chroboczek *et al.*, *Adenovirus Fiber*, p. 163-200 in "The Molecular Repertoire of Adenoviruses I. Virion Structure and Function." W. Doerfler and P. Böhm, eds. (Springer, NY 1995)).

Fiber proteins from different adenoviral serotypes differ considerably. For example, the number of shaft repeats differs between adenoviral serotypes (Green *et al.*, *EMBO J.*, 2, 1357-65 (1983)). Moreover, the knob regions from the closely related Ad2 and Ad5 serotypes are only 63% similar at the amino acid level (Chroboczek *et al.*, *Virology*, 186, 280-85 (1992)), and Ad2 and Ad3 fiber knobs are only 20% identical (Signas *et al.*, *J. Virol.*, 53, 672-78 (1985)). In contrast, the penton base sequences of Ad5 and Ad2 are 99% identical. Despite these differences in the knob region, a sequence comparison of even the Ad2 and Ad3 fiber genes demonstrates distinct regions of conservation, most of which are also conserved among the other human adenoviral fibers (see, e.g., Figures 1A-2B)

One interaction between the adenoviral virion and the host animal is the process of cellular infection, during which the wild-type virion first binds the cell surface by means of a cellular adenoviral receptor (AR) (e.g., the coxsackievirus

and adenovirus receptor (CAR), the MHC class I receptor, etc. (Bergelson *et al.*, *Science*, 275, 1320-23 (1997); Tanako *et al.*, *Proc. Nat. Acad. Sci. (USA)*, 94, 3352-56 (1997)). Hong *et al.*, *EMBO J.*, 16(9), 2294-06 (1997)). After attachment to an AR, the virus binds α_v integrins. Following attachment to these cell surface proteins, infection proceeds by receptor-mediated internalization of the virus into endocytotic vesicles (Svensson *et al.*, *J. Virol.*, 51, 687-94 (1984); Chardonnet *et al.*, *Virology*, 40, 462-77 (1970)). Within the cell, virions are disassembled (Greber *et al.*, *Cell*, 75, 477-86 (1993)), the endosome disrupted (Fitzgerald *et al.*, *Cell*, 32, 607-17 (1983)), and the viral particles transported to the nucleus via the nuclear pore complex (Dales *et al.*, *Virology*, 56, 465-83 (1973)). As most adenoviral serotypes interact with cells through broadly disseminated cell surface proteins, the natural range of host cells infected by adenovirus is broad.

In addition to cellular infection, host animals react defensively to the presence of adenoviral virions through mechanisms that reduce the effective free titer of the virus. For example, host immune systems, upon exposure to a given adenoviral serotype, can efficiently develop neutralizing antibodies, greatly reducing the effective free titer of the virus upon repeat administration (see, e.g., Setoguchi *et al.*, *Am. J. Respir. Cell. Mol. Biol.*, 10, 369-77 (1994); Kass-Eisler *et al.*, *Gene Ther.*, 1, 395-402 (1994); Kass-Eisler *et al.*, *Gene Ther.*, 3, 154-62 (1996)). Interestingly, such antibodies typically are directed against the same determinants of adenoviral serotype specificity, and are primarily directed to the hypervariable hexon regions and, to some extent, fiber and penton base domains (Watson *et al.*, *supra*; Wolfort *et al.* (1988), *supra*; Wolfort *et al.* (1985), *supra*; Crawford-Miksza *et al.*, *supra*). Of course, the presence of adenoviruses agglutinates red blood cells in humans in a serotype-dependent manner (Hierholzer, *J. Infect. Diseases*, 123(4), 541-50 (1973)). Additionally, adenoviral virions are actively scavenged from the circulation by cells of the reticuloendothelial system (RES) (see, e.g., Worgall *et al.*, *Hum Gene Ther.*, 8, 1675-84 (1997); Wolff *et al.*, *J. Virol.*, 71(1), 624-29 (1997)). In such a response, Kupffer cells, endothelial liver cells, or other RES cells scavenge the virus from the circulation (see generally, Moghini *et al.*, *Crit. Rev. Ther. Drug Carrier Sys.*, 11(1), 31-59 (1994); Van Rooijen *et al.*, *J. Leuk. Biol.*, 62, 702-09 (1997)). For example, virions can become opsonized, possibly through interaction between collectins and glycosylated viral proteins, triggering recognition by such RES cells; alternatively, such cells may recognize charged amino acid residues on the virion surface (see Hansen *et al.*, *Immunobiol.*, 199(2), 165-89 (1998); Jahrling *et al.*, *J. Med. Virol.*, 12(1), 1-16 (1983)).

Based on the popularity of adenoviruses as gene transfer vectors, efforts have been made to increase the ability of adenovirus to enter certain cells, e.g., those few cells it does not infect, an approach referred to as "targeting" (see, e.g., International Patent Application WO 95/26412 (Curiel *et al.*), International Patent Application WO 94/10323 (Spooner *et al.*), U.S. Patent 5,543,328 (McClelland *et al.*), International Patent Application WO 94/24299 (Cotten *et al.*)). Of course, while the ability to target adenoviruses to certain cell types is an important goal, far more desirable is an adenovirus which infects only a desired cell type, an approach referred to as "alternative targeting." However, to exclusively target a virus, its native affinity for host cell ARs must first be abrogated, producing a recombinant adenovirus incapable of productively infecting the full set of natural adenoviral target cells. Efforts aimed at abrogating native adenoviral cell affinity have focused logically on changing the fiber knob. These efforts have proven disappointing, largely because they fail to preserve the important fiber protein functions of stable trimerization and penton base binding (Spooner *et al.*, *supra*). Moreover, replacement of the fiber knob with a cell-surface ligand (McClelland *et al.*, *supra*) produces a virus only suitable for infecting a cell type having that ligand. Such a strategy produces a virus having many of the same targeting problems associated with wild-type adenoviruses (in which fiber trimerization and cellular tropism are mediated by the same protein domain), thus decreasing the flexibility of the vector. Moreover, due to the necessity of having a propagating cell line, and the integral connection between the fiber trimerization and targeting functions, obtaining a mutant virus with substituted targeting is difficult. For example, removing the fiber knob and replacing it with a non-trimerizing ligand (e.g., Spooner *et al.*, McClelland *et al.*, *supra*) results in a virus lacking appreciable fiber protein.

Aside from the broad natural tropism of the virus noted above, the non-infectious interactions between adenovirus and the host also pose problems for using adenovirus as gene transfer vectors. Such interactions effectively reduce the free titer of a given dose of adenovirus beneath that which is clinically effective. As such, there is currently a need for an adenovirus exhibiting reduced affinity for such natural interactions with a host animal (e.g., target cell affinity, innate or acquired immune surveillance, etc). Moreover, there is a need for such a virus which is able to deliver and express a desired transgene within a predefined tissue – an alternatively targeted virus.

BRIEF SUMMARY OF THE INVENTION

The present invention provides a recombinant protein having an amino terminus of an adenoviral fiber protein and having a trimerization domain. A fiber incorporating such a protein exhibits reduced affinity for a native substrate than does a wild-type adenoviral fiber trimer. The present invention further provides an adenovirus incorporating the recombinant protein of the present invention.

The present invention is useful in a variety of gene-transfer applications, *in vitro* and *in vivo*, as a vector for delivering a desired gene to a cell with minimal ectopic infection. Specifically, the present invention permits more efficient production and construction of safer vectors for gene transfer applications. The present invention is also useful as a research tool by providing methods and reagents for the study of adenoviral attachment and infection of cells and in a method of assaying receptor-ligand interaction. Similarly, the recombinant fiber protein can be used in receptor-ligand assays and as adhesion proteins *in vitro* or *in vivo*. Additionally, the present invention provides reagents and methods permitting biologists to investigate the cell biology of viral growth and infection. Thus, the vectors of the present invention are highly useful in biological research.

DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B sets forth a comparison of the amino acid sequences of the non-group B serotype fiber knobs (SEQ ID NOs: 5-18) using the Clustal method with PAM100 residue weight table. The height of the bars at the top of each row of sequence comparison correlates to the degree of homology. Consensus and majority sequences are indicated as SEQ ID NOs: 29 and 30, respectively.

Figures 2A and 2B sets forth a comparison of the amino acid sequences of the group B serotype fiber knobs (SEQ ID NOs: 19-25) using the J. Hein method with PAM250 residue weight. The height of the bars at the top of each row of sequence comparison correlates to the degree of homology. Consensus and majority sequences are indicated as SEQ ID NOs: 31 and 32, respectively.

DETAILED DESCRIPTION OF THE INVENTION

Recombinant Protein

The present invention provides a recombinant adenoviral fiber protein having an amino terminus derived from an adenoviral fiber protein and having a trimerization domain. A trimer including such a recombinant protein exhibits reduced affinity for a native substrate, such as an antibody, collectins, opsins, a cellular binding site, etc. (i.e., native to the serotype from which the shaft, and

particularly the amino-terminus, is drawn) as compared to a native adenoviral fiber trimer. The trimer can be a homotrimer or a heterotrimer of different fiber monomers. Any modification of the monomeric units reducing the affinity of the resulting trimer for its native cell surface binding site (i.e., a native AR) is within the scope of the invention. Preferably, the reduction in affinity is a substantial reduction in affinity (such as at least an order of magnitude, and preferably more) relative to the unmodified corresponding fiber.

As mentioned, where a trimerization domain is itself a ligand for a native cell surface binding site, fiber proteins possessing such trimerization domains present some of the same problems for targeting as native adenoviral fiber trimerization domains. Therefore, the trimerization domain of the inventive protein invention preferably is not a ligand for the CAR or MHC-1 cell surface proteins. Most preferably, the non-native trimerization domain is not a ligand for any native adenoviral cell-surface binding site, whether the site is an AR or other cell surface binding site. As is discussed herein, adenoviruses incorporating such proteins exhibit reduced ability to appreciably infect cells via native AR proteins, and can serve as efficient source vectors for engineering alternatively targeted vectors. Therefore, while the trimerization domain preferably is not a ligand for a cell surface binding site, the entire trimer can be such a ligand (e.g., by virtue of a non-native ligand as discussed herein). Moreover, the trimerization domain can be a ligand for a substrate other than a native cell surface binding site, as such trimerization-ligands do not present the same concern for cell targeting as do trimerization domains which are ligands for cell surface binding sites. Thus, for example, the non-native trimerization domain can be a ligand for a substrate on an affinity column, on a blood-borne molecule, or even on a cell surface when it is not a native cell-surface binding site (e.g., on a cell engineered to express a substrate cell surface protein not native to the unmodified cell type).

The recombinant fiber protein can lack a sizable number of residues, or even identifiable domains, as herein described. For example, the protein can lack the native knob domain; it can lack one or more native shaft β -sheet repeats, or it can be otherwise truncated. Thus, a recombinant fiber protein can have any desired modification so long as it trimerizes when produced by a eukaryotic cell. Furthermore, a recombinant fiber protein preferably is not modified appreciably at the amino terminus (e.g., the amino-terminus of a monomer preferably consists essentially of the native fiber amino-terminus) to ensure that a fiber incorporating the recombinant fiber protein interacts properly with the penton base. Hence, the present invention also provides a composition of matter comprising a recombinant

fiber protein of the present invention and an adenoviral penton base. Preferably, the recombinant fiber protein and the penton base associate much in the same manner as wild-type fibers and penton bases. Of course, the penton base can also be modified, for example, to include a non-native ligand, for example as is
5 described in U.S. Patent 5,559,099 (Wickham *et al.*).

In one embodiment, the fiber is modified to render it less able to interact with the innate or acquired host immune system. For example, one or more amino acids of the native fiber protein can be mutated to render the recombinant fiber protein less able to be recognized by neutralizing antibodies than a wild-type fiber
10 (see, e.g., International Patent Application WO 98/40509 (Crystal *et al.*). The fiber also can be modified to lack one or more amino acids mediating interaction with the RES. For example, the fiber can be mutated to lack one or more glycosylation or phosphorylation sites, or the fiber (or virus containing the fiber) can be produced in the presence of inhibitors of glycosylation or phosphorylation.
15 Similarly, the fiber (or other protein within the virus) can be conjugated to a lipid derivative of polyethylene glycol (PEG) comprising a primary amine group, an epoxy group, or a diacylglycerol group (see, e.g., Kilbanov *et al.*, *FEBS Lett.*, 268, 235 (1990); Senior *et al.*, *Biochem. Biophys. Acta.*, 1062, 11 (1991); Allen *et al.*, *Biochem. Biophys. Acta.*, 1066, 29 (1991); Mori *et al.*, *FEBS Lett.*, 284, 263
20 (1991)) to avoid collectin and/or opsonin binding or scavenging by Kupffer (or other RES) cells.

A recombinant fiber protein lacking one or more amino acids, as herein described, can optionally comprise a non-native residue (e.g., several non-native amino acids) in addition to (i.e., insertions) or in place of (i.e., substitutions) the
25 missing native amino acid(s); of course, alternatively, the native amino acid(s) can be deleted from the knob. Preferably, the amino-acid is substituted with another non-native amino acid to preserve topology and, especially, trimerization. Moreover, if substituted, the replacement amino acid preferably confers novel qualities to the recombinant fiber protein. For example, to maximally ablate
30 binding to the native substrate, a native amino acid can be substituted with a residue (or a plurality of residues) having a different charge. Such a substitution maximally interferes with the electrostatic interaction between native adenoviral knob domains and cellular ARs or interferes with a conformational change required to efficiently bind an AR or elements of the RES. Similarly, a native
35 amino acid can be substituted with a residue (or a plurality of residues) of differing weight, where possible. For example, substitution with a heavier residue

maximally interferes with the steric interaction between adenoviral domains and native substrates, by virtue of the longer side-chains on such heavier residues.

Any native amino-acid residue mediating or assisting in the interaction between the knob and a native cellular AR is a suitable amino acid for mutation or deletion from the recombinant fiber protein. Such amino acid need not itself be the site of contact between the fiber and the receptor. For example, the native amino acid might be involved in a conformational change associated with receptor binding. The inventive fiber protein can lack any number of such native amino acids, so long as, in the aggregate, the recombinant fiber protein can associate to form a trimer. The amino acid can be within a β -sheet of the knob or within a loop connecting two β -sheets (such as, for example, the AB, BC, CD, DE, EF, FG, GH, HI, or IJ loops). Indeed, the amino acid can be within 10 (e.g., within 5) residues of a β sheet or a loop. In the mature, folded trimer of the present invention, the amino acid can be within about 10 nm (e.g., within about 5 nm or even within about 2 nm) of a β sheet or a loop.

Native amino acid residues for modification or deletion can be selected by any method. For example, the sequences from different adenoviral serotypes (which are known in the art) can be compared to deduce conserved residues likely to mediate AR-binding. Alternatively or in combination, the sequence can be mapped onto a three dimensional representation of the protein (such as the crystal structure) to deduce those residues most likely responsible for AR binding. These analyses can be aided by resorting to any common algorithm or program for deducing protein structural functional interaction. Alternatively, random mutations can be introduced into a cloned adenoviral fiber expression cassette. One method of introducing random mutations into a protein is via the *Taq* polymerase. For example, a clone encoding the fiber knob (see, e.g., Roelvink *et al.*, *J. Virol.*, 70, 7614-21 (1996)) can serve as a template for PCR amplification of the adenoviral fiber knob, or a portion thereof. By varying the concentration of divalent cations in the PCR reaction, the error rate of the transcripts can be largely predetermined (see, e.g., Weiss *et al.*, *J. Virol.*, 71, 4385-94 (1997); Zhou *et al.*, *Nucl. Acid. Res.*, 19, 6052 (1991)). The PCR products then can be subcloned back into the template vector to replace the sequence within the fiber coding sequence employed as a source for the PCR reaction, thus generating a library of fibers, some of which will harbor mutations which diminish native AR binding while retaining the ability to trimerize.

The amino acids of knobs from strains other than Ad5 that correspond to these listed residues are apparent upon a comparison between the sequences of the

fibers of different adenoviral strains, and any suitable method of determining such correspondence can be employed (e.g., Clusal method with PAM100 residue weight table, J. Hain method with PAM 250 residue weight table, etc.). Examples of such sequence comparison of the knobs of Ad fiber proteins (SEQ ID NOs:5-5
25) are set forth in Figures 1A-2B. By such comparison, residues (e.g., conserved) from other serotypes which, mutated as described, result in fiber trimers with reduced AR binding can be identified (see, e.g., SEQ ID NOs: 29-32). Thus, for example, for CAR-binding fibers, preferably, the amino acid(s) to be mutated is within 10 (e.g., within about 5) amino acids or within about 10 nm
10 (e.g., within about 5 nm) of an amino acid corresponding to residues 404-406, 408, 409, 412-417, 420, 439, 441, 442, 449-454, 456, 458, 460, 462, 466, 467, 469-472, 474-477, 482, 485, 487-492, 505-512, 515, 517, 519, 521-528, 533, 535, 537-549, 551, 553, 555, 559-568, 580, or 581 of the native Ad5 fiber protein (SEQ ID NO:1). More preferably, the amino acid(s) to be mutated correspond to at least
15 one of these residues, such as amino acid 189, 190, 198, 201, or 262 of the native Ad9 fiber protein (SEQ ID NO:3) or amino acid 395, 396, 404, 407, or 470 of the native Ad41 long fiber protein (SEQ ID NO:2). Even more preferably, the mutant fiber protein comprises at least one replacement mutation of a residue corresponding to residues 408, 409, 412-417, 420, 477, or 487-491 of the native
20 Ad5 fiber protein or at least one deletion mutation of a residue corresponding to residues 474-477 or 489-492 of the native Ad5 fiber protein. Similarly, for group B fibers, the amino acid(s) to be mutated is within 10 (e.g., within about 5) amino acids or within about 10 nm (e.g., within about 5 nm) of an amino acid corresponding to residues 136, 155, 177, 181, 198, 210, 211, 215, 233, 234, 236,
25 238, 248, 257, 260, 261, 276, 284, 302, 303, 317, or 318 of the native Ad3 fiber protein (SEQ ID NO:4).

The recombinant fiber protein of the present invention can be produced by any suitable method. For example, the mutant fiber protein can be synthesized using standard direct peptide synthesizing techniques (e.g., as summarized in
30 Bodanszky, *Principles of Peptide Synthesis* (Springer-Verlag, Heidelberg: 1984)), such as via solid-phase synthesis (see, e.g., Merrifield, *J. Am. Chem. Soc.*, 85, 2149-54 (1963); and Barany *et al.*, *Int. J. Peptide Protein Res.*, 30, 705-739 (1987)). Alternatively, site-specific mutations can be introduced into the recombinant fiber protein by ligating into an expression vector a synthesized
35 oligonucleotide comprising the modified site. Alternatively, a plasmid, oligonucleotide, or other vector encoding the desired mutation can be recombined with the adenoviral genome or with an expression vector encoding the

recombinant fiber protein to introduce the desired mutation. Oligonucleotide-directed site-specific mutagenesis procedures also are appropriate (e.g., Walder *et al.*, *Gene*, 42, 133 (1986); Bauer *et al.*, *Gene*, 37, 73 (1985); Craik, *Biotechniques*, 12-19 (1995); U.S. Patents 4,518,584 (Mark *et al.*) and 4,737,462 (Mark *et al.*)).

5 However engineered, the DNA fragment encoding the recombinant fiber protein can be subcloned into an appropriate vector using well known molecular genetic techniques. The fragment is then transcribed and the peptide subsequently translated *in vitro* within a host cell. Any appropriate expression vector (e.g., Pouwels *et al.*, *Cloning Vectors: A Laboratory Manual* (Elsevier, NY: 1985)) and

10 corresponding suitable host cells can be employed for production of recombinant peptides. Expression hosts include, but are not limited to, bacterial species, yeast, mammalian or insect host cell systems including baculovirus systems (e.g., Luckow *et al.*, *Bio/Technology*, 6, 47 (1988)), and established cell lines such

15 HEK-293, COS-7, C127, 3T3, CHO, HeLa, BHK, etc. An especially preferred expression system for preparing modified fibers of the invention is a baculovirus expression system (Wickham *et al.*, *J. Virol.*, 70, 6831-38 (1995)) as it allows the production of high levels of recombinant proteins. Of course, the choice of expression host has ramifications for the type of peptide produced, primarily due to post-translational modification.

20 Once produced, the recombinant fiber proteins are assayed for fiber protein activity. Specifically, the ability of recombinant fiber protein to form trimers, interact with the penton base, and interact with native substrate's (e.g., antibodies, ARs, opsonins, collectins, RES cells, etc.) is assayed. Any suitable assay can be employed to measure these parameters. For example, as improperly folded

25 monomers are generally insoluble (Scopes, "Protein Purification" (3d Ed., 1994), Chapter 9, p. 270-82 (Springer-Verlag, New York)), one assay for trimerization is whether the recombinant fiber is soluble. Determining solubility of the fiber is aided if an amount of radioactive amino-acid is incorporated into the protein during synthesis. Lysate from the host cell expressing the recombinant fiber

30 protein can be centrifuged, and the supernatant and pellet can be assayed via a scintillation counter or by Western analysis. Subsequently, the proteins within the pellet and the supernatant are separated (e.g., on an SDS-PAGE gel) to isolate the fiber protein for further assay. Comparison of the amount of fiber protein isolated from the pellet vis-à-vis the fiber protein isolated from the supernatant indicates

35 whether the mutant protein is soluble. Alternatively, trimerization can be assayed by using a monoclonal antibody recognizing only the amino portion of the trimeric form of the fiber (e.g., via immunoprecipitation, Western blotting, etc.). Another

measure of trimerization is the ability of the recombinant fiber to form a complex with the penton base (Novelli and Boulanger, *Virology*, 185, 1189 (1995)), as only fiber trimers can so interact. This propensity can be assayed by co-immunoprecipitation, gel mobility-shift assays, SDS-PAGE (boiled samples migrate as monomers, otherwise, they migrate as larger proteins), etc. Yet another measure of trimerization is to detect the difference in molecular weight of a trimer as opposed to a monomer. For example, a boiled and denatured trimer will run as a lower molecular weight than a non-denatured stable trimer (Hong and Angler, *J. Virol.*, 70, 7071-78 (1996)). A trimeric recombinant fiber also can be assayed for its ability to bind native substrates. For example, modification of fiber to interfere with its interaction with the host innate or acquired immune system can be accomplished by measuring the free titer of the virus over time. This can be assessed by measuring serum half life, tropism to organs associated with the RES (e.g., liver in mice and humans, lung in pigs, etc.), by agglutination of red blood cells, or by detection of adenoviral genetic material in cell samples.

A trimeric recombinant fiber also can be assayed for its ability to bind native ARs. Any suitable assay that can detect this characteristic is sufficient for use in the present invention. A preferred assay involves exposing cells expressing a native AR (e.g., HEK-293 cells) to the recombinant fiber trimers under standard conditions of infection. Subsequently, the cells are exposed to native adenoviruses, and the ability of the viruses to bind the cells is monitored. Monitoring can be by autoradiography (e.g., employing radioactive viruses), immunocytochemistry, or by measuring the level of infection or gene delivery (e.g., using a reporter gene). In contrast with native trimers which reduce or substantially eliminate subsequent viral binding to the HEK-293 cells, those trimers not substantially reducing the ability of native adenoviruses to subsequently bind the cells are trimers of the present invention. The reduction of interference with subsequent viral binding indicates that the trimer is itself not a ligand for its native mammalian AR, or at least binds with reduced affinity.

Alternatively, a vector including a sequence encoding a mutated fiber (or a library of putative mutated fibers, such as described herein) can be introduced into a suitable host cell strain to express the fiber protein, and mutants can be identified by assaying the inability to bind the soluble CAR protein (e.g., by probing a replica lift with radiolabeled CAR or by other suitable method). Because a reduction in CAR-binding could be due to either selective ablation of the ligand or structural modification affecting trimerization, mutant fibers

identified as non-CAR binding by such a library screen must be assayed for the ability to trimerize, as described above.

Virion and Virus

5 The present invention provides an adenoviral virion incorporating the recombinant fiber protein of the present invention. The virion does not interact with native substrates (e.g., innate and acquired immune systems, cell-surface proteins, etc.) as readily as the wild-type serotype, due to the above-mentioned reduction in affinity of the fibers present in the virion. Moreover, the virion can be
10 further modified to reduce interaction with native substrates through the inclusion of other recombinant proteins. Thus, for example, the virion can include one or more recombinant penton base proteins lacking a native RGD sequence to reduce cell binding via α_v integrins (see, e.g., U.S. patents 5,559,099 (Wickham *et al.*) and 5,731,190 (Wickham *et al.*)). Similarly, the virion can include one or more
15 recombinant hexons lacking a native sequence (e.g., HVR) to reduce its ability to be recognized by a neutralizing antibody (see, e.g., International Patent Application WO 98/40509 (Crystal *et al.*)). Also, the virion can be modified to reduce its ability to interact with the RES. For example, the virion proteins can be mutated to lack one or more glycosylation or phosphorylation sites, or it can be
20 produced in the presence of inhibitors of glycosylation or phosphorylation. Similarly, the virion proteins can be conjugated to a lipid derivative of PEG comprising a primary amine group, an epoxy group, or a diacylglycerol group, as discussed above, to reduce collectin and/or opsonin affinity or scavenging by Kupffer cells or other cells of the RES. Such modifications reduce the ability of
25 host animals to develop neutralizing antibodies to the virions, thereby permitting repeat administration of the virions.

 While the virion exhibits reduced affinity for natural adenoviral substrates, it can include one or more non-adenoviral ligands, for example, to effect targeted infection of a population of cells other than that for which adenoviruses are
30 naturally tropic. Additionally, the non-native ligand can be used to purify the virus, to inactivate the virus (e.g., by adsorbing it to a substrate for the ligand), or to grow the virus on cell lines having receptors recognizing the non-native ligand, for example, as described in International Patent Application WO 98/54346 (Wickham *et al.*).

35 The virus can include any suitable ligand (e.g., a peptide specifically binding to a substrate). For example, for targeting the adenovirus to a cell type other than that naturally infected (or a group of cell types other than the natural

range or set of host cells), the ligand can bind a cell surface binding site (e.g., any site present on the surface of a cell with which the adenovirus can interact to bind the cell and thereby promote cell entry). A cell surface binding site can be any suitable type of molecule, but typically is a protein (including a modified protein such as a glycoprotein, a mucoprotein, etc.), a carbohydrate, a proteoglycan, a lipid, a mucin molecule, or other similar molecule. Examples of potential cell surface binding sites include, but are not limited to, heparin and chondroitin sulfate moieties found on glycosaminoglycans; sialic acid moieties found on mucins, glycoproteins, and gangliosides; common carbohydrate molecules found in membrane glycoproteins, including mannose, N-acetyl-galactosamine, N-acetyl-glucosamine, fucose, and galactose; glycoproteins such cell adhesion molecules (CAMs) (e.g., ICAM-1, ICAM-2, ICAM-3, VCAM-1, NCAM), selectins (e.g., E-selectin, P-selectin, L-selectin, etc.), CD, cadherins, TNF family receptors, GPI-linked receptors, receptors that are efficiently internalized (e.g., CD44, CD31 on endothelial cells, CD34 on high endo-venules), endoglin, growth factor receptors, PSA, androgen receptors, glucocorticoid receptors, prostate-specific membrane antigen (PSMA), MUC1, MUC234, MUC5AC, MUC5B, MUC7, KSA carcino-embryonic antigen (CEA), HER2/NEU (erbB2), folate receptor, corionic gonadotropin- β , (Zhang *et al.*, *Clin. Cancer Res.*, 4, 2669-76 (1998); *Cancer Res.*, 58, 4055 (1998)), and others are known in the art.

A particular cell surface binding site can be present on a narrow class of cell types (e.g., cardiac muscle, skeletal muscle, smooth muscle, etc.) or a broader group encompassing several cell types. Through integration of an appropriate cell-specific ligand, the virion can be employed to target any desired cell type, such as, for example, neuronal, glial, endothelial (e.g., via tissue factor receptor, FLT-1, CD31; CD36; CD34, CD105, CD13, ICAM-1 (McCormick *et al.*, *J. Biol. Chem.*, 273, 26323-29 (1998)); thrombomodulin receptor (Lupus *et al.*, *Suppl.*, 2, S120 (1998)); VEGFR-3 (Lymboussaki *et al.*, *Am. J. Pathol.*, 153(2), 395-403 (1998); mannose receptor; VCAM-1 (Schwarzacher *et al.*, *Atherosclerosis*, 122, 59-67 (1996)), or other receptors); blood clots (e.g., through fibrinogen or aIIbb3 peptide), epithelial (e.g., inflamed tissue through selectins, VCAM-1, ICAM-1, etc.), keratinocytes, follicular cells, adipocytes, fibroblasts, hematopoietic or other stem cells, myoblasts, myofibers, cardiomyocytes, smooth muscle, somatic, osteoclasts, osteoblasts, tooth blasts, chondrocytes, melanocytes, hematopoietic cells, etc., as well as cancer cells derived from any of the above cell types (e.g., prostate (such as via PSMA receptor (see, e.g., Schuur *et al.*, *J. Biol. Chem.*, 271, 7043 (1998); *Cancer Res.*, 58, 4055 (1998))), breast, lung, brain (e.g.,

glioblastoma), leukemia/lymphoma, liver, sarcoma, bone, colon, testicular, ovarian, bladder, throat, stomach, pancreas, rectum, skin (e.g., melanoma), kidney, etc.). Thus, the inventive virions can be targeted to cells within any organ or system, including, for example, respiratory system (e.g., trachea, upper airways, lower airways, alveoli), nervous system and sensory organs (e.g., skin, ear, nasal, tongue, eye), digestive system (e.g., oral epithelium and sensory organs, salivary glands, stomach, small intestines/duodenum, colon, gall bladder, pancreas, rectum), muscular system (e.g., skeletal muscle, connective tissue, tendons), skeletal system (e.g., joints (synovial cells), osteoclasts, osteoblasts, etc.), immune system (e.g., bone marrow, stem cells, spleen, thymus, lymphatic system, etc.), circulatory system (e.g., muscles connective tissue, and/or endothelia of the arteries, veins, capillaries, etc.), reproductive system (e.g., testis, prostate, uterus, ovaries), urinary system (e.g., bladder, kidney, urethra), endocrine or exocrine glands (e.g., breasts, adrenal glands, pituitary glands), etc.

In other embodiments (e.g., to facilitate purification or propagation within a specific engineered cell type), the non-native ligand can bind a compound other than a cell-surface protein. Thus, the ligand can bind blood- and/or lymph-borne proteins (e.g., albumin), synthetic peptide sequences such as polyamino acids (e.g., polylysine, polyhistidine, etc.), artificial peptide sequences (e.g., FLAG), and RGD peptide fragments (Pasqualini *et al.*, *J. Cell. Biol.*, 130, 1189 (1995)). The ligand can even bind non-peptide substrates, such as plastic (e.g., Adey *et al.*, *Gene*, 156, 27 (1995)), biotin (Saggio *et al.*, *Biochem. J.*, 293, 613 (1993)), a DNA sequence (Cheng *et al.*, *Gene*, 171, 1 (1996); Krook *et al.*, *Biochem. Biophys. Res. Commun.*, 204, 849 (1994)), streptavidin (Geibel *et al.*, *Biochemistry*, 34, 15430 (1995); Katz, *Biochemistry*, 34, 15421 (1995)), nitrostreptavidin (Balass *et al.*, *Anal. Biochem.*, 243, 264 (1996)), heparin (Wickham *et al.*, *Nature Biotechnol.*, 14, 1570-73 (1996)), cationic supports, metals such as nickel and zinc (e.g., Rebar *et al.*, *Science*, 263, 671 (1994); Qui *et al.*, *Biochemistry*, 33, 8319 (1994)), or other potential substrates.

Examples of suitable ligands and their substrates for use in the method of the invention include, but are not limited to, CR2 receptor binding the amino acid residue attachment sequences, CD4 receptor recognizing the V3 loop of HIV gp120, transferrin receptor and its ligand (transferrin), low density lipoprotein receptor and its ligand, the ICAM-1 receptor on epithelial and endothelial cells in lung and its ligand, linear or cyclic peptide ligands for streptavidin or nitrostreptavidin (Katz, *Biochemistry*, 34, 15421 (1995)), galactin sequences that bind lactose, galactose and other galactose-containing compounds, and

asialoglycoproteins that recognize deglycosylated protein ligands. Moreover, additional ligands and their binding sites preferably include (but are not limited to) short (e.g., 6 amino acids or less) linear stretches of amino acids recognized by integrins, as well as polyamino acid sequences such as polylysine, polyarginine, etc. Inserting multiple lysines and/or arginines provides for recognition of heparin and DNA. Also, a ligand can comprise a commonly employed peptide tag (e.g., short amino acid sequences known to be recognized by available antisera) such as sequences from glutathione-S-transferase (GST) from *Shistosoma manosi*, thioredoxin β -galactosidase, or maltose binding protein (MPB) from *E. coli*, human alkaline phosphatase, the FLAG octapeptide, hemagglutinin (HA) (Wickham *et al.* (1996), *supra*), polyoma virus peptides, the SV40 large T antigen peptide, BPV peptides, the hepatitis C virus core and envelope E2 peptides and single chain antibodies recognizing them (Chan, *J. Gen. Virol.*, 77, 2531 (1996)), the c-myc peptide, adenoviral penton base epitopes (Stuart *et al.*, *EMBO J.*, 16, 1189-98 (1997)), epitopes present in the E2 envelope of the hepatitis C virus (see, e.g., Chan *et al.* (1996), *supra*), and other commonly employed tags. A preferred substrate for a tag ligand is an antibody directed against it or a derivative of such an antibody (e.g., a FAB fragment, single chain antibody (ScAb)).

As mentioned, a suitable ligand can be specific for any desired substrate, such as those recited herein or otherwise known in the art. However, adenoviral vectors also can be engineered to include novel ligands (e.g., in protein II, III, IIIa, IV, IV, VI, and/or IX) by first assaying for the ability of a peptide to interact with a given substrate. Generally, a random or semirandom peptide library containing potential ligands can be produced, which is essentially a library within an expression vector system. Such a library can be screened by exposing the expressed proteins (i.e., the putative ligands) to a desired substrate. Positive selective binding of a species within the library to the substrate indicates a ligand for that substrate, at least under the conditions of the assay. For screening such a peptide library, any assay able to detect interactions between proteins and substrates is appropriate, and many are known in the art. However, one preferred assay for screening a protein library is a display system (e.g., using an adenovirus or a bacteriophage), which employs a virus expressing the library (e.g., Koivunen *et al.*, *Bio/Technology*, 13, 265-70 (1995); Yanofsky *et al.*, *Proc. Nat. Acad. Sci. U.S.A.*, 93, 7381-86 (1996); Barry *et al.*, *Nature Med.*, 2(3), 299-305 (1996)). Binding of the virus to the substrate is assayed by exposing the virus to the substrate, rinsing the substrate, and selecting for virus remaining bound to the substrate. Subsequently, limiting dilution can identify individual clones

expressing the putative ligand. Thereafter, the insert present in such clones can be sequenced to determine the identity of the ligand.

Once a given ligand is identified, it can be incorporated into any location of the virus capable of interacting with a substrate (i.e., the viral surface). For example, the ligand can be incorporated into the fiber, the penton base, the hexon, protein IX, VI, or IIIa, or other suitable location. Where the ligand is attached to the fiber protein, preferably it does not disturb the interaction between viral proteins or monomers. Thus, the ligand preferably is not itself an oligomerization domain, as such can adversely interact with the trimerization domain as discussed above. Preferably the ligand is added to the virion protein, and is incorporated in such a manner as to be readily exposed to the substrate (e.g., at the terminus of the protein, attached to a residue facing the substrate, positioned on a peptide spacer to contact the substrate, etc.) to maximally present the ligand to the substrate. Where the ligand is attached to or replaces a portion of the penton base, preferably it is within the hypervariable regions to ensure that it contacts the substrate. Furthermore, where the ligand is attached to the penton base, preferably, the recombinant fiber is truncated or short (e.g., from 0 to about 10 shaft repeats) to maximally present the ligand to the substrate (see, e.g., U.S. Patent 5,559,099 (Wickham *et al.*)). Where the ligand is attached to the hexon, preferably it is within a hypervariable region (Mikszta *et al.*, *J. Virol.*, 70(3), 1836-44 (1996)).

When engineered into an adenoviral protein, the ligand can comprise a portion of the native sequence in part and a portion of the non-native sequence in part. Similarly, the sequences (either native and/or nonnative) that comprise the ligand in the protein need not necessarily be contiguous in the chain of amino acids that comprise the protein. In other words, the ligand can be generated by the particular conformation of the protein, e.g., through folding of the protein in such a way as to bring contiguous and/or noncontiguous sequences into mutual proximity. Of course an adenovirus of the present invention (or a blocking protein) can comprise multiple ligands, each binding to a different substrate. For example, a virus can comprise a first ligand permitting affinity purification as described herein, a second ligand that selectively binds a cell-surface site as described herein, and/or a third ligand for inactivating the virus, also as described herein.

The protein including the ligand can include other non-native elements as well. For example, a non-native, unique protease site also can be inserted into the amino acid sequence. The protease site preferably does not affect fiber trimerization or substrate specificity of the fiber ligand. Many such protease sites are known in the art. For example, thrombin recognizes and cleaves at a known

amino acid sequence (Stenflo *et al.*, *J. Biol. Chem.*, 257, 12280-90 (1982)). The presence of such a protease recognition sequence facilitates purification of the virus in some protocols. The protein can be engineered to include the ligand by any suitable method, such as those methods described above for introducing
5 mutations into proteins.

The virion can be used by itself, for example in studies of viral tropism or binding kinetics. In other embodiments, the virion can be used as a genetic vector. For example, the virion can be used in conjunction with lipids and/or liposomes to deliver exogenous genetic material to target cells, in accordance with well-
10 documented methods. In other embodiments, the virion contains a genome derived from an adenovirus; thus, the invention provides an adenoviral vector including the inventive virion and an adenoviral genome.

The adenoviral vector of the present invention can include one or more non-native amino acid sequences for expression (e.g., "expression cassettes" or
15 "genes") as well. Preferably, the non-native amino acid is capable of being transcribed in a cell into which the vector has been internalized. The non-native amino acid can encode a product that effects a biological (e.g., therapeutic) response either at the cellular level or systemically; alternatively, the non-native nucleic acid sequence can encode a product that, in some fashion, can be detected
20 in a cell (e.g., a "reporter gene"). The non-native amino acid can exert its effect at the level of RNA or protein. For instance, a protein encoded by the non-native amino acid can be employed in the treatment of an inherited disease, such as, e.g., the cystic fibrosis transmembrane conductance regulator cDNA for the treatment of cystic fibrosis. Alternatively, the protein encoded by the non-native amino acid
25 can exert its therapeutic effect by effecting cell death. For instance, expression of the non-native amino acid in itself can lead to cell killing, as with expression of the diphtheria toxin. Alternatively, the expression of the non-native amino acid, can render cells selectively sensitive to the action of certain drugs, e.g., expression of the HSV thymidine kinase gene renders cells sensitive to antiviral compounds
30 including acyclovir, gancyclovir, and FIAU (1-(2-deoxy-2-fluoro- β -D-arabinofuranosil)-5-iodouracil). Moreover, the non-native amino acid can exert its effect at the level of RNA, for instance, by encoding an antisense message or ribozyme, a protein which affects splicing or 3' processing (e.g., polyadenylation), or a protein affecting the level of expression of another gene within the cell (i.e.,
35 where gene expression is broadly considered to include all steps from initiation of transcription through production of a processed protein), perhaps, among other things, by mediating an altered rate of mRNA accumulation, an alteration of

mRNA transport, and/or a change in post-transcriptional regulation. Of course, where it is desired to employ gene transfer technology to deliver a given non-native amino acid, its sequence will be known in the art.

Where the inventive adenoviral vector includes a non-native amino acid
5 and a non-adenoviral ligand in its virion, the non-native amino acid can be operably linked to any suitable promoter, such as a promoter native to the adenoviral genome or a non-adenoviral promoter. Where the ligand is employed to deliver the vector to a desired cell type, preferably the non-adenoviral promoter is active within the cell type, and more preferably, the non-adenoviral promoter is
10 a tissue-specific promoter (e.g., specific for the cell type to which the ligand binds), such as those cell types discussed above. For example, expression in targeted endothelial cells can be mediated using the E-selectin promoter (see, e.g., Whelan *et al.*, *TIBS*, 21, 65-69 (1996)); passenger gene expression in targeted prostate cancer cells can be mediated using the PSA promoter (see, e.g., Schuur *et al.*, *J. Cell Biol.*, 271, 7043 (1996). Pang *et al.*, *Cancer Res.*, 57, 495 (1997)) or the
15 E2F promoter. Furthermore, the promoter can be that for a tissue-specific receptor, such as those receptors mentioned herein, still other tissue specific promoter systems are known in the art. Alternatively, the non-native amino acid can be placed under control of a regulable promoter (e.g., metallothionein
20 promoter, tetracycline-responsive promoter, RU486-responsive promoter, etc.).

The altered protein (e.g., the recombinant fiber protein or the coat protein having the ligand) and the non-native amino acid where present) can be incorporated into the adenovirus by any suitable method, many of which are known in the art. As mentioned herein, the protein is preferably identified by
25 assaying products produced in high volume from genes within expression vectors (e.g., baculovirus vectors). The genes from the vectors harboring the desired mutation can be readily subcloned into plasmids, which are then transfected into suitable packaging cells (e.g., HEK-293 cells). Transfected cells are then incubated with adenoviruses under conditions suitable for infection. At some
30 frequency within the cells, homologous recombination between the vector and the virus will produce an adenoviral genome harboring the desired mutation.

Adenoviruses of the present invention can be either replication competent or replication deficient. Preferably, the adenoviral vector comprises a genome with at least one modification therein, rendering the virus replication deficient
35 (see, e.g., International Patent Application WO 95/34671 (Kovesdi *et al.*)). The modification to the adenoviral genome includes, but is not limited to, addition of a DNA segment, rearrangement of a DNA segment, deletion of a DNA segment.

replacement of a DNA segment, or introduction of a DNA lesion. A DNA segment can be as small as one nucleotide and as large as the adenoviral genome (e.g., about 36 kb) or, alternately, can equal the maximum amount which can be packaged into an adenoviral virion (i.e., about 38 kb). Preferred modifications to the adenoviral genome include modifications in the E1, E2, E3, and/or E4 regions. An adenovirus also preferably can be a cointegrate, i.e., a ligation of adenoviral genomic sequences with other sequences, such as other virus, phage, or plasmid sequences.

The virion and adenoviral vector of the present invention have many qualities which render them attractive choices for use in gene transfer, as well as other applications. For example, in many embodiments, the adenovirus does not infect its native host cells as readily as does wild-type adenovirus, due to the recombinant fiber protein. Moreover, by virtue of additional modifications, such virions and vectors are less readily cleared from the host by the innate or acquired immune responses, thus boosting effective free titer and lengthening serum half-life. Furthermore, the virions and vectors have at least one non-native ligand specific for a substrate which facilitates viral propagation, targeting, purification, and/or inactivation as discussed herein. The presence of such ligands can effectively confine expression of non-native amino acids within a predefined cell type or tissue. Linking the non-native amino acid to a tissue-specific or regulable promoter further minimizes expression of the non-native amino acid outside of the targeted tissue. The ligands and the trimerization domains can be separate domains, thus permitting the virus to be easily be reengineered to incorporate different ligands without perturbing fiber trimerization.

Of course, for delivery into a host (such as an animal), a virus of the present invention can be incorporated into a suitable carrier. As such, the present invention provides a composition comprising an adenovirus of the present invention and a pharmacologically acceptable carrier (e.g., a pharmaceutically-acceptable carrier). Any suitable preparation is within the scope of the invention. The exact formulation, of course, depends on the nature of the desired application (e.g., cell type, mode of administration, etc.), and many suitable preparations are set forth in U.S. Patent 5,559,099 (Wickham *et al.*).

Cell Line

As mentioned herein, an adenovirus of the present invention does not readily infect its native host cell via the native AR because its ability to bind ARs is significantly attenuated (due to the incorporation of the recombinant fiber

protein of the present invention). Therefore, the invention provides a cell line able to propagate the inventive adenovirus. Preferably, the cell line can support viral growth for at least about 10 passages (e.g., about 15 passages), and more preferably for at least about 20 passages (e.g., about 25 passages), or even 30 or
5 more passages.

For example, the adenoviruses can be first grown in a packaging cell line which expresses a native fiber protein gene. The resultant viral particles are therefore likely to contain both native fibers encoded by the complementing cell line and non-native fibers encoded by the adenoviral genome (such as those fibers
10 described herein): hence a population of such resultant viruses will contain both fiber types. Such particles will be able to bind and enter packaging cell lines via the native fiber more efficiently than particles which lack native fiber molecules. Thus, the employment of such a fiber-encoding cell line permits adenovirus genomes encoding chimeric, targeted adenovirus fibers to be grown and amplified
15 to suitably high titers. The resultant "mixed" stocks of adenovirus produced from the cell lines encoding the native fiber molecule will contain both native and chimeric adenovirus fiber molecules; however, the particles contain genomes encoding only the chimeric adenovirus fiber. Thus, to produce a pure stock of adenoviruses having only the chimeric adenovirus fiber molecules, the "mixed"
20 stock is used to infect a packaging cell line which does not produce native fiber (such as HEK-293 for E1-deleted non-group B viruses). The resultant adenoviruses contain only the fiber molecules encoded by the genomes (i.e., the chimeric fiber molecules).

Similar fiber-complementing cell lines have been produced and used to
25 grow mutant adenovirus lacking the fiber gene). However, the production rates of these cell lines have generally not been great enough to produce adenovirus titers of the fiber-deleted adenovirus comparable to those of fiber-expressing adenovirus particles. The lower titers produced by such mutants can be improved by temporally regulating the expression of the native fiber to more fully complement
30 the mutant adenovirus genome. One strategy to produce such an improved cell line is to use of a regulable promoter to permit fiber production to be controlled and activated once the cells are infected with adenovirus. Alternatively, an efficient mRNA splice site introduced into the fiber gene in the complementing cell line improves the level of fiber protein production in the cell line.

35 When the adenovirus is engineered to contain a ligand specific for a given cell surface binding site, any cell line expressing that receptor and capable of supporting adenoviral growth is a suitable host cell line. However, because many

ligands do not bind cell surface binding sites (especially some of the novel ligands discussed herein), a cell line can be engineered to express the substrate for the ligand.

5 The present invention provides a cell line expressing a non-native cell-surface receptor (a pseudo-receptor) to which a virus having a ligand for said receptor binds. Any cell line capable of supporting viral growth is a suitable cell line for use in the present invention. If the virus lacks genes essential for viral replication, preferably the cell line expresses complementing levels of such gene products (see, e.g., International Patent Application WO 95/34671 (Kovesdi et al.),
10 U.S. Patents 5,658,724 (DeLuca) and 5,804,413 (DeLuca)). When the virus is an adenovirus, preferably the cell line of the present invention is derived from HEK-293 cells. When the virus is a herpesvirus, preferably the cell line of the present invention is derived from VERO cells.

15 The non-native cell surface binding site is a substrate molecule, such as described herein, to which an adenovirus having a ligand selectively binding that substrate can bind the cell and thereby promote cell entry. The binding site can recognize a non-native ligand incorporated into the adenoviral coat or a ligand native to a virus. For example, when the non-native viral ligand is a tag peptide, the binding site can be a single chain antibody (ScAb) receptor recognizing the
20 tag. Alternatively, the ScAb can recognize an epitope present in a region of a mutated fiber knob (if present), or even an epitope present on a native adenoviral coat protein, (e.g., on the fiber, penton, hexon, etc.). Alternatively, if the non-native ligand recognizes a cell-surface substrate (e.g., membrane-bound protein), the binding site can comprise that substrate. If the substrate binding site is native
25 to a cell-surface receptor, the cell line can express a mutant receptor with decreased ability to interact with the cellular signal transduction pathway (e.g., a truncated receptor, such as NMDA (Li *et al.*, *Nat. Biotech.*, 14, 989 (1996))), attenuated ability to act as an ion channel, or other modification. Infection via such modified proteins minimizes the secondary effects of viral infection on host-
30 cell metabolism by reducing the activation of intracellular messaging pathways and their various response elements. The choice of binding site depends to a large extent on the nature of the adenovirus. However, to promote specificity of the virus for a particular cell type, the binding site preferably is not a native mammalian AR. Moreover, the binding site must be expressed on the surface of
35 the cell to be accessible to the virus. Hence, where the binding site is a protein, it preferably has a leader sequence and a membrane tethering sequence to promote

proper integration into the membrane (see, e.g., Davitz *et al.*, *J. Exp. Med.* 163, 1150 (1986)).

The cell line can be produced by any suitable method. For example, a vector (e.g., an oligonucleotide, plasmid, viral, or other vector) containing a nucleic acid encoding the non-native receptor can be introduced into source cell line by conventional means. Preferably, the vector also encodes an agent permitting the cells harboring it to be selected (e.g., the vector can encode resistance to antibiotics which kill cells not harboring the plasmid). At some frequency, the vector will recombine with the cell genome to produce a transformed cell line expressing the non-native receptor.

EXAMPLES

While it is believed that one of skill in the art is fully able to practice the invention after reading the foregoing description, the following examples further illustrate some of its features. In particular, the examples demonstrate the construction of several recombinant fiber proteins, each exhibiting reduced affinity for native adenoviral substrates. The examples further demonstrate the inclusion of such recombinant fiber proteins into adenoviral vectors, and the retargeting of such vectors using non-native ligands. The examples also demonstrate the successful construction of a pseudoreceptor cell line able to propagate the alternatively targeted viruses. As these examples are included for purely illustrative purposes, they should not be construed to limit the scope of the invention in any respect.

The procedures employed in these examples, such as affinity chromatography, Southern blots, PCR, DNA sequencing, vector construction (including DNA extraction, isolation, restriction digestion, ligation, etc.), cell culture (including antibiotic selection), transfection of cells, protein assays (Western blotting, immunoprecipitation, immunofluorescence), etc., are techniques routinely performed by those of skill in the art (see generally Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989)). Accordingly, in the interest of brevity, experimental protocols are not discussed in detail.

EXAMPLE 1

This example describes mutant fiber trimers exhibiting reduced affinity for the CAR protein.

Using standard site-directed mutagenesis, mutations were introduced into
5 nearly every major sheet and loop in the native Ad5 fiber knob sequence (SEQ ID NO:1). In a first series of mutagenesis, replacement mutations were designed in which between 3 and 6 contiguous amino acids within a loop were replaced by the same number of glycine residues. In a second series of mutagenesis, mutations
10 were designed in which between 1 and 4 amino acids were deleted from the native sequence. Extensive point mutations also were conducted. One additional mutant was designed in which 12 amino acids were deleted and replaced with a tetrapeptide sequence.

Respective baculovirus clones, each containing one of the recombinant mutant protein genes, were created and used to produce recombinant mutant knob
15 proteins in insect cells. The baculovirus-infected insect cells were freeze-thawed at 3 days post-infection to release any soluble recombinant mutant protein (approximately 10^7 cells per ml of PBS). The freeze-thawed lysate was centrifuged and the soluble fraction and the insoluble pellet were collected. Western analysis of the soluble and insoluble fractions revealed that similar levels
20 of the mutant and native fiber knobs were present in the soluble fraction. Mutants which retained solubility represent proteins which folded properly and trimerized, and these are set forth in Table 1. In the table, mutations are indicated by noting the location of the mutated residue or residues of the Ad5 fiber within parentheses. The identity of the native residue or residues is set forth to the left, and the identity of any substituting residue or residues is to the right of the parentheses. Deletions
25 are further delineated using the "Δ" symbol.

Table 1

Mutation Location	Mutations	
AB Loop (403-418)	T(404)G P(405)G A(406)K S(408)E S(408)G P(409)A R(412)G	RLN(412-414)GGG N(414)G A(415)G RLNAEK(415-417)SLNNGG E(416)G K(417)G K(417)L
B Sheet (419-428)	K(420)A	
C Sheet (431-440)	L(439)S	
CD Loop (441-453)	V(441)S Δ SG(449-450)	SGTVQ(449-453)GSGSG
D Sheet (454-461)	S(454)N + R(460)Q H(456)E + R(460)E	I(458)E + R(460)E
DE Loop (462-478)	D(462)A V(466)S L(467)S NNS(469-471)GGG Δ F(472)	DPE(474-476)GGG DPEY(474-477)GGGG Y(477)A Y(477)T
E Sheet (479-482)	N(482)A	
F Sheet (485-486)	L(485)G	
FG Loop (487-514)	E(487)G T(489)G A(490)G EGTAY(487-491)GGGGG Y(491)A Δ TAYT(489-492)	P(505)G Δ K(506) H(506)A Δ KT(510-511) SHGKTA(507-512)GSGSGS
G Sheet (515-521)	N(515)S + V(517)S V(517)S + Q(519)S	Y(521)H
GH Loop (522-528)	NGDKT(523-527)GSGSG D(525)K KTK(526-528)RSR	K(526)E K(528)S
H Sheet (528-536)	T(535)E	T(533)S + T(535)S
HI Loop (537-549)	N(537)E	GTQETGDTTPSA(538-549)GSGG
I Sheet (550-557)	S(551)N + S(555)N S(551)E	S(553)E
IJ Loop (558-572)	SGHN(559-562)GSGS Δ HN(561-562) Y(563)H	INEI(564-567)GSGS E(566)K F(568)H
C-Terminus (573-578)	Q(580)G	E(581)G

To determine whether a given mutant fiber had reduced affinity for CAR, competition experiments were performed by preincubating A549 cells with either the trimeric mutants or native fiber knobs followed by incubation with radiolabeled Ad5 virus. Either 1 or 10 μ l volumes of the native knob preincubated with A549 cells blocked 90% or more of the labeled Ad5 binding to cells measured in the absence of a competitor. In this assay, any soluble, trimeric mutant less efficient in blocking fiber-mediated Ad5 cell binding or gene transduction than the native knob was considered to have reduced affinity for CAR. Those trimeric mutant fibers exhibiting reduced affinity for CAR in this assay are indicated in Table 2.

The trimeric mutant fiber proteins were mass produced by infecting roughly 15 million insect cells each with the baculoviral vectors (MOI = 10) and culturing them for 3 days. The cells were harvested and freeze-thawed, and the cell debris was removed via centrifugation. NaCl was added to the supernatant to a final concentration of 750 mM, and then the supernatant was added to 500 μ l TALONTM resin. After one hour at 25 °C, the resin was centrifuged at 2,500 for two minutes. The supernatant was removed, and the resin resuspended in 10 ml 750 mM NaCl. After 30 minutes incubation, the resin suspension was run through a column. The mutant protein was eluted using 2 ml of elution fluid (20 mM TRIS, pH 8.0, 100 mM NaCl, 150 mM imidazole). The eluate was dialized once against PBS with 750 mM NaCl, once against PBS with 500 mM NaCl, and once against PBS with 250 mM NaCl. Protein concentration was determined by standard methods and protein integrity verified by Western analysis.

The purified proteins were subjected to a competition assay with Ad5 capsids to assess the degree to which each mutation decreased interaction with CAR. Serial dilutions of each mutant protein, as well as wild-type Ad5 fiber, were added to A549 cells (10^5 cells/well) in 24-well plates. Following this preincubation, an Ad5 vector containing the lacZ gene were added to each well (MOI = 10). After a one hour incubation at 37 °C, the inoculum was removed, the cells were washed with culture medium, and then a culture medium (DMEM with 5 % FCS) added. The cells were incubated overnight, lysed 18 hours post infection, and assayed for β -galactosidase activity by standard methods. Plotting the degree of β -galactosidase activity against concentration of preincubation protein permitted assessment of each protein's IC_{50} value (the concentration of the competing protein at the 50% level). The degree to which each mutation reduced CAR-binding as calculated by this method is set forth in Table 2.

Table 2

Mutation Number	Mutation Location	Mutation Sequence	Competition
F5K	-	Native	100%
F3K	-	Native	< 0.1%
Ad5-1	AB Loop	S408E	< 0.1%
Ad5-2	AB Loop	P409A	< 1%
Ad5-3	AB Loop	RLNAEK(412-417)SLNGGG	< 0.1%
Ad5-4	AB Loop	K(417)G	< 0.1%
Ad5-5	B Sheet	K(420)A	< 0.1%
Ad5-6	DE Loop	ΔDPE(474-476)	< 20%
Ad5-7	DE Loop	ΔDPEY(474-477)	< 0.1%
Ad5-8	DE Loop	Y(477)A	< 0.1%
Ad5-9	FG Loop	EGTAY(487-491)GGGGG	< 0.1%
Ad5-10	FG Loop	ΔTAYT(489-492)	< 0.1%

EXAMPLE 2

5 This example describes recombinant fiber proteins exhibiting reduced affinity for the CAR protein.

10 The Ad9 and long Ad41 fiber proteins corresponding to mutations Ad5-1, Ad5-2, Ad5-4, Ad5-5, and Ad5-9 (see Figures 1A and 1B) were generated. The resultant mutant proteins were soluble, and each was used in competition assays against wild type Ad5, as described in Example 1, to assess whether the mutations affected CAR binding. The results of these experiments (presented in Table 3) reveal that residues important for CAR binding are conserved among adenoviral serotypes.

Table 3

Mutation	Mutation Sequence	Corresponding Ad5 Mutation	Competition
Ad9-1	S(189)E	Ad5-1	No
Ad9-2	P(190)A	Ad5-2	No
Ad9-3	K(198)G	Ad5-4	No
Ad9-4	K(201)A	Ad5-5	No
Ad9-5	Y(262)A	Ad5-8	No
Ad41-1	S(395)E	Ad5-1	No
Ad41-2	P(369)A	Ad5-2	No
Ad41-3	L(404)G	Ad5-4	No
Ad41-3	T(470)A	Ad5-8	No

EXAMPLE 3

This example describes the production of a pseudo-receptor for
 5 constructing a cell line able to replicate adenoviruses lacking native cell-binding
 function (but targeted for the pseudo-receptor). Specifically, the exemplary
 pseudo-receptor includes a binding domain from a single-chain antibody
 recognizing HA.

Anti-HA ScFv was constructed as an N-Term-VL-VH fusion protein. RT-
 10 PCR was performed on RNA obtained from hybridomas producing HA antibodies
 using primers specific for κ - or $\gamma 2\beta$ - and C-terminus of the VL and VH genes (see
 Gilliland *et al.*, *Tissue Antigens*, 47, 1-20 (1996)). After sequencing the resulting
 PCR products, specific oligonucleotides were designed to amplify the VL-VH
 fusion in a second round of PCR. The final PCR product was cloned to create a
 15 plasmid for production of anti-HA ScFv in *E. coli*. The expressed protein has a C-
 terminal E peptide for detection of binding to HA-tagged penton base via Western
 analysis of ELISA assay. Upon transformation of bacterial cells with the plasmid,
 Western analysis using an antibody recognizing the E peptide revealed a protein of
 the expected size.

20 To determine whether the anti-HA ScFv was functional, it was used in
 protein A immunoprecipitation assays using adenoviral coat proteins (recombinant
 penton base) containing the HA epitope. The anti-HA ScFv was able to precipitate

HA-containing penton base proteins. These results indicate the successful construction of the extracellular portion of a pseudo-receptor for binding an adenovirus having a non-native ligand (i.e., HA).

To create an entire anti-HA pseudo-receptor, the anti-HA ScFv was cloned
5 in frame with sequences encoding a C-terminal pair of myc epitopes followed by the PDGF receptor transmembrane anchor. The entire sequence of this pseudo-receptor is indicated at SEQ ID NO:28. A eukaryotic expression plasmid containing this sequence, pSc(HA), was transfected into HEK-293 cells. The following day the pSc(HA)-transfected cells or cells transfected with a control
10 ScFv construct were incubated for 30 min on ice with a fluorescein-tagged HA peptide (HA*) or with a fluorescein-tagged scrambled HA peptide (scrHA*). Following the incubation of HA* with the pSc(HA)-transfected cells, a discrete population of cells was found to brightly fluoresce specifically around the cell membrane. The pSc(HA)-transfected cells incubated with the scrHA* peptide did
15 not display this fluorescent pattern, nor did the cells transfected with the control plasmid and then incubated with HA*. Enhanced fluorescence of the pSc(HA)-transfected cells incubated with HA* was also demonstrated by FACS analysis. Moreover, preincubation of the anti-HA pseudo-receptor cells with excess unlabelled HA peptide, but not unlabelled FLAG peptide, blocked the fluorescent
20 pattern observed on cells incubated with HA* alone.

These results demonstrate the successful construction and expression of a cell line consisting essentially of cells expressing a functional pseudo-receptor.

EXAMPLE 4

25 This example describes an alternatively targeted adenovirus having recombinant fiber proteins exhibiting reduced affinity for the CAR protein and having a non-native ligand.

The Ad5-10 mutant described in Example 1 was subjected to further site directed mutagenesis to introduce a polypeptide including the HA epitope into the
30 HI loop of the fiber knob (between amino acids 543 and 544 of SEQ ID NO:1). The resultant fiber has the TAYT deletion in the FG loop and an HA epitope sequence inserted into the HI loop.

The gene encoding this mutant fiber was combined into a plasmid that contains a full length, E1- and E3-deleted adenovirus genome carrying the above
35 fiber mutation plus a CMV-driven *LacZ* reporter gene in the E1 region. This plasmid was then linearized and transfected into HEK-293 cells expressing the anti-HA pseudo-receptor described in Example 3. After 5 days the cells were

freeze-thawed three times, and the virus-containing lysate was passaged onto fresh anti-HA-293 cells.

The resultant adenoviruses were further amplified in the anti-HA-293 cells and then purified using standard methods. The vector (AdZ.F*fg(HA)hi) exhibits
5 reduced binding capacity to CAR on standard HEK-293 cells due to the TAYT deletion; however, it binds with high affinity via its HA epitope to the anti-HA pseudoreceptor present on the anti-HA-293 cell line.

EXAMPLE 5

10 This example describes an alternatively targeted adenovirus having recombinant fiber proteins exhibiting reduced affinity for the CAR protein and having more than one non-native ligand.

The Ad5-10 mutant described in Example 1 was subjected to further site directed mutagenesis to introduce a polypeptide including the HA epitope and a
15 high affinity RGD ligand into the HI loop of the fiber knob (between amino acids 543 and 544 of SEQ ID NO:1) . The resultant plasmid encodes a fiber with the TAYT deletion in the FG loop and an RGD sequence inserted into the HI loop.

The gene encoding this mutant fiber gene was then combined into a plasmid that contains a full length, E1 and E3-deleted adenovirus genome carrying
20 the above fiber mutation plus a CMV-driven *LacZ* reporter gene in the E1 region. This plasmid was then linearized and transfected into HEK-293 cells expressing the anti-HA pseudo-receptor described in Example 2. After 5 days the cells are freeze-thawed three times and the virus-containing lysate is passaged onto fresh HEK-293 cells.

25 The resultant adenoviruses were further amplified in the anti-HA-293 cells and then purified using standard methods. The vector exhibits reduced binding capacity to CAR on standard HEK-293 cells due to the TAYT deletion; however, it efficiently infects cells expressing α_v integrins (such as tumor cells) via the RGD ligand present in the HI loop.

30

EXAMPLE 6

This example describes an alternatively targeted adenovirus having recombinant fiber proteins exhibiting reduced affinity for the CAR protein and having a non-native ligand.

35 The Ad5-3 mutant described in Example 1 was subjected to further site directed mutagenesis to introduce an 18 amino acid polypeptide including the HA epitope into the HI loop of the fiber knob (between amino acids 543 and 544 of

SEQ ID NO:1) . The resultant fiber has the RLNAEK mutation of the AB loop and an HA epitope sequence inserted into the HI loop.

5 The gene encoding this mutant fiber was combined into a plasmid that contains a full length, E1- and E3-deleted adenovirus genome carrying the above fiber mutation plus a CMV-driven *LacZ* reporter gene in the E1 region. This plasmid was then linearized and transfected into HEK-293 cells expressing the anti-HA pseudo-receptor described in Example 3. After 5 days the cells were freeze-thawed three times, and the virus-containing lysate was passaged onto fresh anti-HA 293 cells.

10 The resultant adenoviruses were further amplified in the anti-HA 293 cells and then purified using standard methods. The vector (AdZ.F*ab(HA)hi) exhibits reduced binding capacity to CAR on standard HEK-293 cells due to the RLNAEK mutation; however, it binds with high affinity via its HA epitope to the anti-HA pseudoreceptor present on the anti-HA 293 cell line.

15

EXAMPLE 7

This example describes an alternatively targeted adenovirus having recombinant fiber proteins exhibiting reduced affinity for the CAR protein and having more than one non-native ligand.

20 The Ad5-3 mutant described in Example 1 was subjected to further site directed mutagenesis to introduce a polypeptide including the HA epitope and a high affinity RGD ligand into the HI loop of the fiber knob (between amino acids 543 and 544 of SEQ ID NO:1) . The resultant plasmid encodes a fiber with the RLNAEK mutation of the AB loop and an HA epitope and RGD sequence inserted into the HI loop.

25 The gene encoding this mutant fiber gene was then combined into a plasmid that contains a full length, E1- and E3-deleted adenovirus genome carrying the above fiber mutation plus a CMV-driven *LacZ* reporter gene in the E1 region. This plasmid was then linearized and transfected into HEK-293 cells expressing the anti-HA pseudo-receptor described in Example 3. After 5 days the cells are freeze-thawed three times, and the virus-containing lysate was passaged onto fresh anti-HA 293 cells.

30 The resultant adenoviruses were further amplified in the anti-HA 293 cells and then purified using standard methods. The vector exhibits reduced binding capacity to CAR on standard HEK-293 cells due to the RLNAEK mutation; however, it binds with high affinity via its HA epitope to the anti-HA pseudoreceptor present on the anti-HA 293 cell line. Moreover, the virus also

efficiently infects cells expressing α_v integrins (such as tumor cells) via the RGD ligand present in the HI loop.

EXAMPLE 8

5 This example describes an alternatively targeted adenovirus having recombinant fiber proteins exhibiting reduced affinity for the CAR protein and having a non-native ligand.

A mutation was introduced into the Ad2 fiber knob, deleting the Asn-Pro residues in the FG loop (residues 90 and 91 of SEQ ID NO:7). Additionally, the high-affinity RGD motif was introduced into the HI loop of this protein. The sequences encoding the knob domain were fused to sequences encoding the Ad5 shaft, resulting in a nucleic acid encoding a chimeric Ad5-Ad2 fiber. This construct was cloned into an Ad5 genome also containing the lacZ gene (the AdZ virus), replacing the native fiber sequence. The resultant viruses are termed AdZ.F*(RGD).

10 Increasing particle doses of either AdZ or AdZ.F*(RGD) were incubated with either SKOV-3 cells (which express both CAR and α_v integrins) or Ramos cells (which express CAR but not α_v integrins) in suspension (10^6 cells/300 μ l medium) for one hour at 36 °C, following which the cells were washed and incubated overnight. Following the incubation, the cells were assayed for lacZ activity using conventional methods.

20 The SKOV-3 cells were transduced by both viruses, while the Ramos cells were transduced by AdZ, but only poorly transduced by AdZ.F*(RGD). These results demonstrate that the native CAR-binding ability of the vector can be blocked by mutating selective residues of the fiber knob and the virus retarded by the addition of a non-native ligand to the viral coat protein.

EXAMPLE 9

This example demonstrated the reduced affinity for the CAR protein of recombinant fiber proteins.

30 Various cell types (A172, HuVEC, HCAEC, A549, HeLa, HEK-293, and HS68) (10^6 cells/300 μ l medium) were preincubated for 30 minutes at 37 °C with either soluble Ad5 fiber protein (3 μ g/ml) or penton base protein (100 μ g/ml). Following this incubation, either AdZ, AdZ.F*ab(HA)hi or AdZ.F*fg(HA)hi (100 viral particles/cell) were added to the cells. After a one hour incubation at 37 °C, the cells were twice washed and incubated overnight, again at 37 °C. Following the incubation, the cells were assayed for lacZ activity using conventional

methods. Except for the HS68 fibroblast cell line, the results indicate that preincubation with Ad5 fiber blocked AdZ transduction, but preincubation with penton base did not. In contrast, the viruses containing the mutant fibers were not blocked by preincubation with fiber, but were blocked by preincubation with
5 penton base. These data are consistent with the ablation of native fiber-based infection through mutating the fibers as indicated.

EXAMPLE 10

This example demonstrated the alteration of viral targeting *in vivo*, using an
10 alternatively targeted adenovirus.

The jugular veins of Balb/C mice were injected with either AdZ, AdZ.F*ab(HA)hi or AdZ.F*fg(HA)hi (10^{10} particles/animal in 100 ml, eight animals each). The experiments were run in duplicate, and two animals served as a control (100 ml saline). At one day post inoculation, the animals were sacrificed
15 and the liver of each was snap-frozen in liquid nitrogen. The livers were then pulverized, and lacZ activity was assayed by conventional methods to determine enzymatic activity/mass of tissue.

The livers from the AdZ.F*ab(HA)hi- or AdZ.F*fg(HA)hi-inoculated animals exhibited about 10% of the lacZ activity as those inoculated with AdZ,
20 while control animals exhibited background levels of activity. These results indicate that fiber mutations ablating native cell-receptor binding are effective in greatly reducing native tropism *in vivo*.

All references cited herein are hereby incorporated by reference to the same
25 extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

While this invention has been described with an emphasis on preferred embodiments, it will be obvious to those of ordinary skill in the art that variations of the preferred embodiments can be used and that it is intended that the invention
30 can be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the following claims.

WHAT IS CLAIMED IS:

1. A recombinant fiber protein comprising an amino terminus of an adenoviral fiber protein and a trimerization domain, wherein said trimerization domain comprises an adenoviral fiber knob domain having a mutation affecting at least one amino acid residue within the region corresponding to the AB loop, B sheet, DE loop, or FG loop of the wild-type Ad5 fiber protein, and wherein said recombinant fiber protein trimerizes when produced in a eukaryotic cell.
5
2. The recombinant fiber protein of claim 1, wherein said region is the AB loop.
- 10 3. The recombinant fiber protein of claim 1, wherein said region is the B sheet.
4. The recombinant fiber protein of claim 1, wherein said amino acid residue corresponds to a residue selected from the group of residues consisting of 408, 409, 412-417, 420, 474-477, and 487-492 of the wild-type Ad5 fiber protein.
- 15 5. A recombinant fiber protein comprising an amino terminus of an adenoviral fiber protein and a trimerization domain, wherein said trimerization domain comprises an adenoviral fiber knob having a mutation affecting at least one amino acid corresponding to residue 404-406, 408, 409, 412-417, 420, 439, 441, 442, 449-454, 456, 458, 460, 462, 466, 467, 469-472, 474-477, 482, 485,
20 487-492, 505-512, 515, 517, 519, 521-528, 533, 535, 537-549, 551, 553, 555, 559-568, 580, or 581 of the wild-type Ad5 fiber protein, and wherein said recombinant fiber protein trimerizes when produced in a eukaryotic cell.
- 25 6. The recombinant fiber protein of claim 5, wherein said amino acid residue corresponds to residue 189, 190, 198, 201, or 262 of the native Ad9 fiber protein.
7. The recombinant fiber protein of claim 5, wherein said amino acid residue corresponds to residue 395, 396, 404, 407, or 470 of the native Ad41 long fiber protein.
- 30 8. The recombinant fiber protein of claim 5, wherein said amino acid residue corresponds to residue 136, 155, 177, 181, 198, 210, 211, 215, 233, 234, 236, 238, 248, 257, 260, 261, 276, 284, 302, 303, 317, or 318 of the native Ad3 fiber protein.
9. The recombinant fiber protein of any of claims 1-8, wherein said mutation alters the charge of said residue.
- 35 10. A trimer comprising the recombinant fiber protein of any of claims 1-8, wherein said trimer has an affinity for a native adenoviral cellular receptor of at least about an order of magnitude less than a wild-type adenoviral fiber trimer.

11. An adenoviral virion comprising the trimer of claim 10.
12. The adenoviral virion of claim 11, comprising a penton base having a mutation affecting at least one native RGD sequence.
13. The adenoviral virion of claim 11, comprising a hexon having a mutation affecting at least one native HVR sequence.
14. The adenoviral virion of claim 11, lacking a native glycosylation or phosphorylation site.
15. The adenoviral virion of claim 11, which is conjugated to a lipid derivative of polyethylene glycol comprising a primary amine group, an epoxy group, or a diacylglycerol group.
16. The adenoviral virion of claim 11, which elicits less immunogenicity in a host animal than does a wild-type adenovirus.
17. The adenoviral virion of claim 11, comprising a non-adenoviral ligand.
18. The adenoviral virion of claim 17, wherein said non-adenoviral ligand is conjugated to a fiber.
19. The adenoviral virion of claim 17, wherein said non-adenoviral ligand is conjugated to a penton.
20. The adenoviral virion of claim 17, wherein said non-adenoviral ligand is conjugated to a hexon.
21. The adenoviral virion of claim 17, wherein said non-adenoviral ligand is conjugated to protein IX, VI, or IIIa.
22. The adenoviral virion of claim 17, wherein said non-adenoviral ligand binds a substrate other than a native mammalian adenoviral receptor.
23. The adenoviral virion of any of claim 17, wherein said non-adenoviral ligand binds a substrate other than a native cell-surface protein.
24. The adenoviral virion of claim 17, wherein said substrate is present on the surface of a cell.
25. An adenoviral vector comprising the adenoviral virion of claim 11 and an adenoviral genome.
26. The adenoviral vector of claim 25, which is replication incompetent.
27. The adenoviral vector of claim 25, which does not productively infect HEK-293 cells.
28. The adenoviral vector of claim 25, wherein said virion comprises a non-adenoviral ligand, and said adenoviral genome comprises a non-native nucleic acid for transcription.
29. The adenoviral vector of claim 25, wherein said non-native nucleic acid for transcription is operably linked to a non-adenoviral promoter.



30. The adenoviral vector of claim 25, wherein said ligand binds to a substrate present on the surface of a cell and wherein said non-adenoviral promoter is active within said cell.

5 31. The adenoviral vector of claim 29, wherein said non-adenoviral promoter is a tissue-specific promoter.

32. The adenoviral vector of claim 29, wherein said non-adenoviral promoter is a regulable promoter.


33. A method of infecting a cell, comprising contacting a cell with an adenoviral vector of claim 25.

10 34. The method of claim 33, wherein said adenoviral genome comprises a non-native nucleic acid encoding a protein, and wherein said nucleic acid is expressed within said cell to produce said protein.



Consensus #1: When 57% (8) match the residue of F5K, otherwise show ' '.

Figure 1B



Consensus 'SEQ ID NO: 31': When 57% (4) match the residue of the Consensus show the residue of the Consensus, otherwise show . . .

Figure 2B

SEQUENCE LISTING

<110> GenVec, Inc.
 <120> Alternatively Targeted Adenovirus
 <130> 202033
 <140> WO 99US20728
 <141> 199-09-10
 <150> US 60/099,851
 <151> 1998-09-11
 <150> US 60/136,529
 <151> 1999-05-28
 <160> 32
 <170> PatentIn Ver. 2.2
 <210> 1
 <211> 581
 <212> PRT
 <213> Human adenovirus serotype 5
 <400> 1

Met	Lys	Arg	Ala	Arg	Pro	Ser	Glu	Asp	Thr	Phe	Asn	Pro	Val	Tyr	Pro
1				5					10					15	
Tyr	Asp	Thr	Glu	Thr	Gly	Pro	Pro	Thr	Val	Pro	Phe	Leu	Thr	Pro	Pro
			20					25					30		
Phe	Val	Ser	Pro	Asn	Gly	Phe	Gln	Glu	Ser	Pro	Pro	Gly	Val	Leu	Ser
		35					40					45			
Leu	Arg	Leu	Ser	Glu	Pro	Leu	Val	Thr	Ser	Asn	Gly	Met	Leu	Ala	Leu
	50					55					60				
Lys	Met	Gly	Asn	Gly	Leu	Ser	Leu	Asp	Glu	Ala	Gly	Asn	Leu	Thr	Ser
65				70						75				80	
Gln	Asn	Val	Thr	Thr	Val	Ser	Pro	Pro	Leu	Lys	Lys	Thr	Lys	Ser	Asn
				85					90					95	
Ile	Asn	Leu	Glu	Ile	Ser	Ala	Pro	Leu	Thr	Val	Thr	Ser	Glu	Ala	Leu
			100					105					110		
Thr	Val	Ala	Ala	Ala	Ala	Pro	Leu	Met	Val	Ala	Gly	Asn	Thr	Leu	Thr
		115					120					125			
Met	Gln	Ser	Gln	Ala	Pro	Leu	Thr	Val	His	Asp	Ser	Lys	Leu	Ser	Ile
	130					135					140				
Ala	Thr	Gln	Gly	Pro	Leu	Thr	Val	Ser	Glu	Gly	Lys	Leu	Ala	Leu	Gln
145					150					155					160
Thr	Ser	Gly	Pro	Leu	Thr	Thr	Thr	Asp	Ser	Ser	Thr	Leu	Thr	Ile	Thr
				165					170					175	
Ala	Ser	Pro	Pro	Leu	Thr	Thr	Ala	Thr	Gly	Ser	Leu	Gly	Ile	Asp	Leu
			180					185					190		
Lys	Glu	Pro	Ile	Tyr	Thr	Gln	Asn	Gly	Lys	Leu	Gly	Leu	Lys	Tyr	Gly
		195					200					205			
Ala	Pro	Leu	His	Val	Thr	Asp	Asp	Leu	Asn	Thr	Leu	Thr	Val	Ala	Thr
	210					215					220				
Gly	Pro	Gly	Val	Thr	Ile	Asn	Asn	Thr	Ser	Leu	Gln	Thr	Lys	Val	Thr

225		230		235		240									
Gly	Ala	Leu	Gly	Phe	Asp	Ser	Gln	Gly	Asn	Met	Gln	Leu	Asn	Val	Ala
				245					250					255	
Gly	Gly	Leu	Arg	Ile	Asp	Ser	Gln	Asn	Arg	Arg	Leu	Ile	Leu	Asp	Val
			260					265					270		
Ser	Tyr	Pro	Phe	Asp	Ala	Gln	Asn	Gln	Leu	Asn	Leu	Arg	Leu	Gly	Gln
		275					280					285			
Gly	Pro	Leu	Phe	Ile	Asn	Ser	Ala	His	Asn	Leu	Asp	Ile	Asn	Tyr	Asn
	290					295					300				
Lys	Gly	Leu	Tyr	Leu	Phe	Thr	Ala	Ser	Asn	Asn	Ser	Lys	Lys	Leu	Glu
305					310					315					320
Val	Asn	Leu	Ser	Thr	Ala	Lys	Gly	Leu	Met	Phe	Asp	Ala	Thr	Ala	Ile
				325					330					335	
Ala	Ile	Asn	Ala	Gly	Asp	Gly	Leu	Glu	Phe	Gly	Ser	Pro	Asn	Ala	Pro
			340					345					350		
Asn	Thr	Asn	Pro	Leu	Lys	Thr	Lys	Ile	Gly	His	Gly	Leu	Glu	Phe	Asp
		355					360					365			
Ser	Asn	Lys	Ala	Met	Val	Pro	Lys	Leu	Gly	Thr	Gly	Leu	Ser	Phe	Asp
	370					375					380				
Ser	Thr	Gly	Ala	Ile	Thr	Val	Gly	Asn	Lys	Asn	Asn	Asp	Lys	Leu	Thr
385					390					395					400
Leu	Trp	Thr	Thr	Pro	Ala	Pro	Ser	Pro	Asn	Cys	Arg	Leu	Asn	Ala	Glu
				405					410					415	
Lys	Asp	Ala	Lys	Leu	Thr	Leu	Val	Leu	Thr	Lys	Cys	Gly	Ser	Gln	Ile
			420					425					430		
Leu	Ala	Thr	Val	Ser	Val	Leu	Ala	Val	Lys	Gly	Ser	Leu	Ala	Pro	Ile
		435					440					445			
Ser	Gly	Thr	Val	Gln	Ser	Ala	His	Leu	Ile	Ile	Arg	Phe	Asp	Glu	Asn
	450					455					460				
Gly	Val	Leu	Leu	Asn	Asn	Ser	Phe	Leu	Asp	Pro	Glu	Tyr	Trp	Asn	Phe
465				470						475					480
Arg	Asn	Gly	Asp	Leu	Thr	Glu	Gly	Thr	Ala	Tyr	Thr	Asn	Ala	Val	Gly
				485					490					495	
Phe	Met	Pro	Asn	Leu	Ser	Ala	Tyr	Pro	Lys	Ser	His	Gly	Lys	Thr	Ala
			500					505					510		
Lys	Ser	Asn	Ile	Val	Ser	Gln	Val	Tyr	Leu	Asn	Gly	Asp	Lys	Thr	Lys
		515					520					525			
Pro	Val	Thr	Leu	Thr	Ile	Thr	Leu	Asn	Gly	Thr	Gln	Glu	Thr	Gly	Asp
	530					535					540				
Thr	Thr	Pro	Ser	Ala	Tyr	Ser	Met	Ser	Phe	Ser	Trp	Asp	Trp	Ser	Gly
545					550					555					560
His	Asn	Tyr	Ile	Asn	Glu	Ile	Phe	Ala	Thr	Ser	Ser	Tyr	Thr	Phe	Ser
				565					570					575	
Tyr	Ile	Ala	Gln	Glu											
			580												

<210> 2
 <211> 562
 <212> PRT

<213> Human adenovirus serotype 41.LONG

<400> 2

Met Lys Arg Ala Arg Leu Glu Asp Asp Phe Asn Pro Val Tyr Pro Tyr
 1 5 10 15
 Glu His Tyr Asn Pro Leu Asp Ile Pro Phe Ile Thr Pro Pro Phe Ala
 20 25 30
 Ser Ser Asn Gly Leu Gln Glu Lys Pro Pro Gly Val Leu Ser Leu Lys
 35 40 45
 Tyr Thr Asp Pro Leu Thr Thr Lys Asn Gly Ala Leu Thr Leu Lys Leu
 50 55 60
 Gly Thr Gly Leu Asn Ile Asp Glu Asn Gly Asp Leu Ser Ser Asp Ala
 65 70 75 80
 Ser Val Glu Val Ser Ala Pro Ile Thr Lys Thr Asn Lys Ile Val Gly
 85 90 95
 Leu Asn Tyr Thr Lys Pro Leu Ala Leu Arg Ser Asn Ala Leu Thr Leu
 100 105
 Ser Tyr Asn Ala Pro Leu Asn Val Val Asn Asn Asn Leu Ala Leu Asn
 115 120 125
 Ile Ser Gln Pro Val Thr Val Asn Ala Asn Asn Glu Leu Ser Leu Leu
 130 135 140
 Ile Asp Ala Pro Leu Asn Ala Asp Thr Gly Thr Leu Arg Leu Gln Ser
 145 150 155 160
 Ala Ala Pro Leu Gly Leu Val Asp Lys Thr Leu Lys Val Leu Phe Ser
 165 170 175
 Ser Pro Leu Tyr Leu Asp Asn Asn Phe Leu Thr Leu Ala Ile Glu Arg
 180 185 190
 Pro Leu Ala Leu Ser Ser Ser Arg Ala Val Thr Leu Lys Tyr Ser Pro
 195 200 205
 Pro Leu Lys Ile Glu Asn Glu Asn Leu Thr Leu Ser Thr Gly Gly Pro
 210 215 220
 Phe Thr Val Ser Gly Gly Asn Leu Asn Leu Thr Thr Ser Ala Pro Leu
 225 230 235 240
 Ser Val Gln Asn Asn Ser Leu Ser Leu Val Ile Thr Ser Pro Leu Lys
 245 250 255
 Val Ile Asn Ser Met Leu Ala Val Gly Val Asn Pro Pro Phe Thr Ile
 260 265 270
 Thr Asp Ser Gly Leu Ala Met Asp Leu Gly Asp Gly Leu Ala Leu Gly
 275 280 285
 Gly Ser Lys Leu Ile Ile Asn Leu Gly Pro Gly Leu Gln Met Ser Asn
 290 295 300
 Gly Ala Ile Thr Leu Ala Leu Asp Ala Ala Leu Pro Leu Gln Tyr Arg
 305 310 315 320
 Asp Asn Gln Leu Gln Leu Arg Ile Gly Ser Thr Ser Gly Leu Ile Met
 325 330 335
 Ser Gly Val Thr Gln Thr Leu Asn Val Asn Ala Asn Thr Gly Lys Gly
 340 345 350
 Leu Ala Val Glu Asn Asn Ser Leu Val Val Lys Leu Gly Asn Gly Leu

355 360 365

Arg Phe Asp Ser Trp Gly Ser Ile Thr Val Ser Pro Thr Thr Thr Thr
 370 375 380

Pro Thr Thr Leu Trp Thr Thr Ala Asp Pro Ser Pro Asn Ala Thr Phe
 385 390 395 400

Tyr Glu Ser Leu Asp Ala Lys Val Trp Leu Val Leu Val Lys Cys Asn
 405 410 415

Gly Met Val Asn Gly Thr Ile Ser Ile Lys Ala Gln Lys Gly Ile Leu
 420 425 430

Leu Arg Pro Thr Ala Ser Phe Ile Ser Phe Val Met Tyr Phe Tyr Ser
 435 440 445

Asp Gly Thr Trp Arg Lys Asn Tyr Pro Val Phe Asp Asn Glu Gly Ile
 450 455 460

Leu Ala Asn Ser Ala Thr Trp Gly Tyr Arg Gln Gly Gln Ser Ala Asn
 465 470 475 480

Thr Asn Val Ser Asn Ala Val Glu Phe Met Pro Ser Ser Lys Arg Tyr
 485 490 495

Pro Asn Gln Lys Gly Ser Glu Val Gln Asn Met Ala Leu Thr Tyr Thr
 500 505 510

Phe Leu Gln Gly Asp Pro Asn Met Ala Ile Ser Phe Gln Ser Ile Tyr
 515 520 525

Asn His Ala Leu Glu Gly Tyr Ser Leu Lys Phe Thr Trp Arg Val Arg
 530 535 540

Asn Asn Glu Arg Phe Asp Ile Pro Cys Cys Ser Phe Ser Tyr Val Thr
 545 550 555 560

Glu Gln

<210> 3
 <211> 362
 <212> PRT
 <213> Human adenovirus serotype 9

<400> 3

Met Ser Lys Arg Leu Arg Val Glu Asp Asp Phe Asn Pro Val Tyr Pro
 1 5 10 15

Tyr Gly Tyr Ala Arg Asn Gln Asn Ile Pro Phe Leu Thr Pro Pro Phe
 20 25 30

Val Ser Ser Asp Gly Phe Gln Asn Phe Pro Pro Gly Val Leu Ser Leu
 35 40 45

Lys Leu Ala Asp Pro Ile Ala Ile Val Asn Gly Asn Val Ser Leu Lys
 50 55 60

Val Gly Gly Gly Leu Thr Leu Gln Asp Gly Thr Gly Lys Leu Thr Val
 65 70 75 80

Asn Ala Asp Pro Pro Leu Gln Leu Thr Asn Asn Lys Leu Gly Ile Ala
 85 90 95

Leu Asp Ala Pro Phe Asp Val Ile Asp Asn Lys Leu Thr Leu Leu Ala
 100 105 110

Gly His Gly Leu Ser Ile Ile Thr Lys Glu Thr Ser Thr Leu Pro Gly
 115 120 125

Leu Arg Asn Thr Leu Val Val Leu Thr Gly Lys Gly Ile Gly Thr Glu
 130 135 140
 Ser Thr Asp Asn Gly Gly Thr Val Cys Val Arg Val Gly Glu Gly Gly
 145 150 155 160
 Gly Leu Ser Phe Asn Asn Asp Gly Asp Leu Val Ala Phe Asn Lys Lys
 165 170 175
 Glu Asp Lys Arg Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys
 180 185 190
 Lys Ile Asp Gln Asp Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys
 195 200 205
 Cys Gly Ser Gln Ile Leu Ala Asn Val Ser Leu Ile Val Val Asp Gly
 210 215 220
 Lys Tyr Lys Ile Ile Asn Asn Asn Thr Gln Pro Ala Leu Lys Gly Phe
 225 230 235 240
 Thr Ile Lys Leu Leu Phe Asp Glu Asn Gly Val Leu Met Glu Ser Ser
 245 250 255
 Asn Leu Gly Lys Ser Tyr Trp Asn Phe Arg Asn Glu Asn Ser Ile Met
 260 265 270
 Ser Thr Ala Tyr Glu Lys Ala Ile Gly Phe Met Pro Asn Leu Val Ala
 275 280 285
 Tyr Pro Lys Pro Thr Ala Gly Ser Lys Lys Tyr Ala Arg Asp Ile Val
 290 295 300
 Tyr Gly Asn Ile Tyr Leu Gly Gly Lys Pro Asp Gln Pro Val Thr Ile
 305 310 315 320
 Lys Thr Thr Phe Asn Gln Glu Thr Gly Cys Glu Tyr Ser Ile Thr Phe
 325 330 335
 Asp Phe Ser Trp Ala Lys Thr Tyr Val Asn Val Glu Phe Glu Thr Thr
 340 345 350
 Ser Phe Thr Phe Ser Tyr Ile Ala Gln Glu
 355 360

<210> 4
 <211> 319
 <212> PRT
 <213> Human adenovirus serotype 3

<400> 4

Met Ala Lys Arg Ala Arg Leu Ser Thr Ser Phe Asn Pro Val Tyr Pro
 1 5 10 15
 Tyr Glu Asp Glu Ser Ser Ser Gln His Pro Phe Ile Asn Pro Gly Phe
 20 25 30
 Ile Ser Pro Asp Gly Phe Thr Gln Ser Pro Asn Gly Val Leu Ser Leu
 35 40 45
 Lys Cys Val Asn Pro Leu Thr Thr Ala Ser Gly Ser Leu Gln Leu Lys
 50 55 60
 Val Gly Ser Gly Leu Thr Val Asp Thr Thr Asp Gly Ser Leu Glu Glu
 65 70 75 80
 Asn Ile Lys Val Asn Thr Pro Leu Thr Lys Ser Asn His Ser Ile Asn
 85 90 95
 Leu Pro Ile Gly Asn Gly Leu Gln Ile Glu Gln Asn Lys Leu Cys Ser

				100						105					110			
Lys	Leu	Gly	Asn	Gly	Leu	Thr	Phe	Asp	Ser	Ser	Asn	Ser	Ile	Ala	Leu			
		115					120					125						
Lys	Asn	Asn	Thr	Leu	Trp	Thr	Gly	Pro	Lys	Pro	Glu	Ala	Asn	Cys	Ile			
	130					135					140							
Ile	Glu	Tyr	Gly	Lys	Gln	Asn	Pro	Asp	Ser	Lys	Leu	Thr	Leu	Ile	Leu			
145					150					155					160			
Val	Lys	Asn	Gly	Gly	Ile	Val	Asn	Gly	Tyr	Val	Thr	Leu	Met	Gly	Ala			
				165					170					175				
Ser	Asp	Tyr	Val	Asn	Thr	Leu	Phe	Lys	Asn	Lys	Asn	Val	Ser	Ile	Asn			
			180					185					190					
Val	Glu	Leu	Tyr	Phe	Asp	Ala	Thr	Gly	His	Ile	Leu	Pro	Asp	Ser	Ser			
		195					200					205						
Ser	Leu	Lys	Thr	Asp	Leu	Glu	Leu	Lys	Tyr	Lys	Gln	Thr	Ala	Asp	Phe			
	210					215					220							
Ser	Ala	Arg	Gly	Phe	Met	Pro	Ser	Thr	Thr	Ala	Tyr	Pro	Phe	Val	Leu			
225					230					235					240			
Pro	Asn	Ala	Gly	Thr	His	Asn	Glu	Asn	Tyr	Ile	Phe	Gly	Gln	Cys	Tyr			
				245					250					255				
Tyr	Lys	Ala	Ser	Asp	Gly	Ala	Leu	Phe	Pro	Leu	Glu	Val	Thr	Val	Met			
			260					265					270					
Leu	Asn	Lys	Arg	Leu	Pro	Asp	Ser	Arg	Thr	Ser	Tyr	Val	Met	Thr	Phe			
		275					280					285						
Leu	Trp	Ser	Leu	Asn	Ala	Gly	Leu	Ala	Pro	Glu	Thr	Thr	Gln	Ala	Thr			
	290					295					300							
Leu	Ile	Thr	Ser	Pro	Phe	Thr	Phe	Ser	Tyr	Ile	Arg	Glu	Asp	Asp				
305					310					315								

<210> 5
 <211> 179
 <212> PRT
 <213> Human adenovirus serotype 12

<400> 5

Thr	Leu	Trp	Thr	Thr	Pro	Asp	Pro	Pro	Pro	Asn	Cys	Ser	Leu	Ile	Gln
1				5					10					15	
Glu	Leu	Asp	Ala	Lys	Leu	Thr	Leu	Cys	Leu	Thr	Lys	Asn	Gly	Ser	Ile
			20					25					30		
Val	Asn	Gly	Ile	Val	Ser	Leu	Val	Gly	Val	Lys	Gly	Asn	Leu	Leu	Asn
		35					40					45			
Ile	Gln	Ser	Thr	Thr	Thr	Thr	Thr	Val	Gly	Val	His	Leu	Val	Phe	Asp
	50					55						60			65
Gln	Gly	Arg	Leu	Ile	Thr	Ser	Thr	Pro	Thr	Ala	Leu	Val	Pro	Gln	Ala
					70					75					80
Ser	Trp	Gly	Tyr	Arg	Gln	Gly	Gln	Ser	Val	Ser	Thr	Asn	Thr	Val	Thr
				85					90					95	
Asn	Gly	Leu	Gly	Phe	Met	Pro	Asn	Val	Ser	Ala	Tyr	Pro	Arg	Pro	Asn
			100					105					110		
Ala	Ser	Glu	Ala	Lys	Ser	Gln	Met	Val	Ser	Leu	Thr	Tyr	Leu	Gln	Gly
		115					120					125			

Asp Thr Ser Lys Pro Ile Thr Met Lys Val Ala Phe Asn Gly Ile Thr
 130 135 140
 Ser Leu Asn Gly Tyr Ser Leu Thr Phe Met Trp Ser Gly Leu Ser Asn
 145 150 155 160
 Tyr Ile Asn Gln Pro Phe Ser Thr Pro Ser Cys Ser Phe Ser Tyr Ile
 165 170 175

Thr Gln Glu

<210> 6
 <211> 179
 <212> PRT
 <213> Human adenovirus serotype 31

<400> 6

Thr Leu Trp Thr Thr Pro Asp Pro Pro Pro Asn Cys Thr Leu Arg Gln
 1 5 10 15
 Glu Leu Asp Ala Lys Leu Thr Leu Cys Leu Thr Lys Asn Glu Ser Ile
 20 25 30
 Val Asn Gly Ile Val Ser Leu Ile Gly Val Lys Gly Asp Leu Leu His
 35 40 45
 Ile Gln Pro Thr Thr Thr Thr Val Gly Leu His Leu Val Phe Asp Arg
 50 55 60
 Gln Gly Arg Leu Val Thr Thr Thr Pro Thr Ala Leu Val Pro Gln Ala
 65 70 75 80
 Ser Trp Gly Tyr Lys Gln Gly Gln Ser Val Ser Ser Ser Ala Val Ala
 85 90 95
 Asn Ala Leu Gly Phe Met Pro Asn Val Ser Ala Tyr Pro Arg Pro Asn
 100 105 110
 Ala Gly Glu Ala Lys Ser Gln Met Leu Ser Gln Thr Tyr Leu Gln Gly
 115 120 125
 Asp Thr Thr Lys Pro Ile Thr Met Lys Val Val Phe Asn Gly Asn Ala
 130 135 140
 Thr Val Asp Gly Tyr Ser Leu Thr Phe Met Trp Thr Gly Val Ser Asn
 145 150 155 160
 Tyr Leu Asn Gln Gln Phe Ser Thr Pro Ser Cys Ser Phe Ser Tyr Ile
 165 170 175

Ala Gln Glu

<210> 7
 <211> 183
 <212> PRT
 <213> Human adenovirus serotype 2

<400> 7

Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro Asn Cys Arg Ile His Ser
 1 5 10 15
 Asp Asn Asp Cys Lys Phe Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Val Leu Ala Thr Val Ala Ala Leu Ala Val Ser Gly Asp Leu Ser Ser
 35 40 45
 Met Thr Gly Thr Val Ala Ser Val Ser Ile Phe Leu Arg Phe Asp Gln

<212> PRT
 <213> Human adenovirus serotype 8

<400> 9

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Arg Ile Asp Gln
 1 5 10 15
 Asp Lys Asp Ser Lys Leu Ser Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Asn Val Ser Leu Ile Val Val Ala Gly Arg Tyr Lys Ile
 35 40 45
 Ile Asn Asn Asn Thr Asn Pro Ala Leu Lys Gly Phe Thr Ile Lys Leu
 50 55 60
 Leu Phe Asp Lys Asn Gly Val Leu Met Glu Ser Ser Asn Leu Gly Lys
 65 70 75 80
 Ser Tyr Trp Asn Phe Arg Asn Gln Asn Ser Ile Met Ser Thr Ala Tyr
 85 90 95
 Glu Lys Ala Ile Gly Phe Met Pro Asn Leu Val Ala Tyr Pro Lys Pro
 100 105 110
 Thr Thr Gly Ser Lys Lys Tyr Ala Arg Asp Ile Val Tyr Gly Asn Ile
 115 120 125
 Tyr Leu Gly Gly Lys Pro His Gln Pro Val Thr Ile Lys Thr Thr Phe
 130 135 140
 Asn Gln Glu Thr Gly Cys Glu Tyr Ser Ile Thr Phe Asp Phe Ser Trp
 145 150 155 160
 Ala Lys Thr Tyr Val Asn Val Glu Phe Glu Thr Thr Ser Phe Thr Phe
 165 170 175
 Ser Tyr Ile Ala Gln Glu
 180

<210> 10
 <211> 182
 <212> PRT
 <213> Human adenovirus serotype 9

<400> 10

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Lys Ile Asp Gln
 1 5 10 15
 Asp Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Asn Val Ser Leu Ile Val Val Asp Gly Lys Tyr Lys Ile
 35 40 45
 Ile Asn Asn Asn Thr Gln Pro Ala Leu Lys Gly Phe Thr Ile Lys Leu
 50 55 60
 Leu Phe Asp Glu Asn Gly Val Leu Met Glu Ser Ser Asn Leu Gly Lys
 65 70 75 80
 Ser Tyr Trp Asn Phe Arg Asn Glu Asn Ser Ile Met Ser Thr Ala Tyr
 85 90 95
 Glu Lys Ala Ile Gly Phe Met Pro Asn Leu Val Ala Tyr Pro Lys Pro
 100 105 110
 Thr Ala Gly Ser Lys Lys Tyr Ala Arg Asp Ile Val Tyr Gly Asn Ile
 115 120 125

Tyr Leu Gly Gly Lys Pro Asp Gln Pro Val Thr Ile Lys Thr Thr Phe
 130 135 140
 Asn Gln Glu Thr Gly Cys Glu Tyr Ser Ile Thr Phe Asp Phe Ser Trp
 145 150 155 160
 Ala Lys Thr Tyr Val Asn Val Glu Phe Glu Thr Thr Ser Phe Thr Phe
 165 170 175
 Ser Tyr Ile Ala Gln Glu
 180

<210> 11
 <211> 187
 <212> PRT
 <213> Human adenovirus serotype 15

<400> 11

Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro Asn Cys Lys Ile Ile Glu
 1 5 10
 Asp Lys Asp Ser Lys Leu Thr Leu Ile Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Gly Ser Val Ser Leu Leu Val Val Lys Gly Lys Phe Ser Asn
 35 40 45
 Ile Asn Asn Thr Thr Asn Pro Asn Glu Ala Asp Lys Gln Ile Thr Val
 50 55 60
 Lys Leu Leu Phe Asp Ala Asn Gly Val Leu Lys Gln Gly Ser Thr Met
 65 70 75 80
 Asp Ser Ser Tyr Trp Asn Tyr Arg Ser Asp Asn Ser Asn Leu Ser Gln
 85 90 95
 Pro Tyr Lys Lys Ala Val Gly Phe Met Pro Ser Lys Thr Ala Tyr Pro
 100 105 110
 Lys Gln Thr Lys Pro Thr Asn Lys Glu Ile Ser Gln Ala Lys Asn Lys
 115 120 125
 Ile Val Ser Asn Val Tyr Leu Gly Gly Lys Ile Asp Gln Pro Cys Val
 130 135 140
 Ile Ile Ile Ser Phe Asn Glu Glu Ala Asp Ser Asp Tyr Ser Ile Val
 145 150 155 160
 Phe Tyr Phe Lys Trp Tyr Lys Thr Tyr Glu Asn Val Gln Phe Asp Ser
 165 170 175
 Ser Ser Phe Asn Phe Ser Tyr Ile Ala Gln Glu
 180 185

<210> 12
 <211> 182
 <212> PRT
 <213> Human adenovirus serotype 17

<400> 12

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Arg Ile Asp Lys
 1 5 10
 Glu Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Asn Val Ser Leu Ile Val Val Ser Gly Lys Tyr Gln Tyr
 35 40 45

Ile Asp His Ala Thr Asn Pro Thr Leu Lys Ser Phe Lys Ile Lys Leu
 50 55 60
 Leu Phe Asp Asn Lys Gly Val Leu Leu Pro Ser Ser Asn Leu Asp Ser
 65 70 75 80
 Thr Tyr Trp Asn Phe Arg Ser Asp Asn Leu Thr Val Ser Glu Ala Tyr
 85 90 95
 Lys Asn Ala Val Glu Phe Met Pro Asn Leu Val Ala Tyr Pro Lys Pro
 100 105 110
 Thr Thr Gly Ser Lys Lys Tyr Ala Arg Asp Ile Val Tyr Gly Asn Ile
 115 120 125
 Tyr Leu Gly Gly Leu Ala Tyr Gln Pro Val Val Ile Lys Val Thr Phe
 130 135 140
 Asn Glu Glu Ala Asp Ser Ala Tyr Ser Ile Thr Phe Glu Phe Val Trp
 145 150 155 160
 Asn Lys Glu Tyr Ala Arg Val Glu Phe Glu Thr Thr Ser Phe Thr Phe
 165 170 175
 Ser Tyr Ile Ala Gln Gln
 180

<210> 13
 <211> 181
 <212> PRT
 <213> Human adenovirus serotype 19

<400> 13

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Thr Ile Ala Gln
 1 5 10 15
 Asp Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Asn Val Ser Leu Ile Val Val Ala Glu Arg Tyr His Ile
 35 40 45
 Ile Asn Asn Lys Thr Asn Pro Glu Ile Lys Ser Phe Thr Ile Lys Leu
 50 55 60
 Leu Phe Asn Lys Asn Gly Val Leu Leu Asp Asn Ser Asn Leu Gly Lys
 65 70 75 80
 Ala Tyr Trp Asn Phe Arg Ser Gly Asn Ser Asn Val Ser Thr Ala Tyr
 85 90 95
 Glu Lys Ala Ile Gly Phe Met Pro Asn Leu Val Ala Tyr Pro Lys Pro
 100 105 110
 Ser Asn Ser Lys Lys Tyr Ala Arg Asp Ile Val Tyr Gly Thr Ile Tyr
 115 120 125
 Leu Gly Gly Lys Pro Asp Gln Pro Ala Val Ile Lys Thr Thr Phe Asn
 130 135 140
 Gln Glu Thr Gly Cys Glu Tyr Ser Ile Thr Phe Asp Phe Ser Trp Ser
 145 150 155 160
 Lys Thr Tyr Glu Asn Val Glu Phe Glu Thr Thr Ser Phe Thr Phe Ser
 165 170 175
 Tyr Ile Ala Gln Glu
 180

<210> 14
 <211> 179
 <212> PRT
 <213> Human adenovirus serotype 28

<400> 14

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Lys Met Ser Glu
 1 5 10 15
 Val Lys Asp Ser Lys Leu Thr Leu Ile Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Gly Ser Val Ser Leu Leu Ala Val Lys Gly Glu Tyr Gln Asn
 35 40 45
 Met Thr Ala Ser Thr Asn Lys Asn Val Lys Ile Thr Leu Leu Phe Asp
 50 55 60
 Ala Asn Gly Val Leu Leu Glu Gly Ser Ser Leu Asp Lys Glu Tyr Trp
 65 70 75 80
 Asn Phe Arg Asn Asn Asp Ser Thr Val Ser Gly Lys Tyr Glu Asn Ala
 85 90 95
 Val Pro Phe Met Pro Asn Ile Thr Ala Tyr Lys Pro Val Asn Ser Lys
 100 105 110
 Ser Tyr Ala Arg Ser His Ile Phe Gly Asn Val Tyr Ile Asp Ala Lys
 115 120 125
 Pro Tyr Asn Pro Val Val Ile Lys Ile Ser Phe Asn Gln Glu Thr Gln
 130 135 140
 Asn Asn Cys Val Tyr Ser Ile Ser Phe Asp Tyr Thr Cys Ser Lys Glu
 145 150 155 160
 Tyr Thr Gly Met Gln Phe Asp Val Thr Ser Phe Thr Phe Ser Tyr Ile
 165 170 175

Ala Gln Glu

<210> 15
 <211> 181
 <212> PRT
 <213> Human adenovirus serotype 37

<400> 15

Thr Leu Trp Thr Thr Pro Asp Thr Ser Pro Asn Cys Thr Ile Ala Gln
 1 5 10 15
 Asp Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Asn Val Ser Leu Ile Val Val Ala Gly Lys Tyr His Ile
 35 40 45
 Ile Asn Asn Lys Thr Asn Pro Lys Ile Lys Ser Phe Thr Ile Lys Leu
 50 55 60
 Leu Phe Asn Lys Asn Gly Val Leu Leu Asp Asn Ser Asn Leu Gly Lys
 65 70 75 80
 Ala Tyr Trp Asn Phe Arg Ser Gly Asn Ser Asn Val Ser Thr Ala Tyr
 85 90 95
 Glu Lys Ala Ile Gly Phe Met Pro Asn Leu Val Ala Val Ser Lys Pro
 100 105 110
 Ser Asn Ser Lys Lys Tyr Ala Arg Asp Ile Val Tyr Gly Asn Ile Tyr

115 120 125
 Leu Gly Gly Lys Pro Asp Gln Pro Gly Val Ile Lys Thr Thr Phe Asn
 130 135 140
 Gln Glu Thr Gly Cys Glu Tyr Ser Ile Thr Phe Asn Phe Ser Trp Ser
 145 150 155 160
 Lys Thr Tyr Glu Asn Val Glu Phe Glu Thr Thr Ser Phe Thr Phe Ser
 165 170 175
 Tyr Ile Ala Gln Glu
 180

<210> 16
 <211> 179
 <212> PRT
 <213> Human adenovirus serotype 4
 <400> 16

Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro Asn Cys Gln Ile Leu Ala
 1 5 10 15
 Glu Asn Asp Ala Lys Leu Thr Leu Cys Leu Thr Met Cys Asp Ser Gln
 20 25 30
 Ile Leu Ala Thr Val Ser Val Leu Val Val Arg Ser Gly Asn Leu Asn
 35 40 45
 Pro Ile Thr Gly Thr Val Ser Ser Ala Gln Val Phe Leu Arg Phe Asp
 50 55 60
 Ala Asn Gly Val Leu Leu Thr Glu His Ser Thr Ser Lys Lys Tyr Trp
 65 70 75 80
 Gly Tyr Lys Gln Gly Asp Ser Ile Asp Gly Thr Pro Tyr Thr Asn Ala
 85 90 95
 Val Gly Phe Met Pro Asn Ser Thr Ala Tyr Pro Lys Thr Gln Ser Ser
 100 105 110
 Thr Thr Lys Asn Asn Ile Val Gly Gln Val Tyr Met Asn Gly Asp Val
 115 120 125
 Ser Lys Pro Met Leu Leu Thr Ile Thr Leu Asn Gly Thr Asp Asp Thr
 130 135 140
 Thr Ser Ala Tyr Ser Met Ser Phe Ser Tyr Thr Trp Thr Asn Gly Ser
 145 150 155 160
 Tyr Ile Gly Ala Thr Phe Gly Ala Asn Ser Tyr Thr Phe Ser Tyr Ile
 165 170 175

Ala Gln Gln
 <210> 17
 <211> 176
 <212> PRT
 <213> Human adenovirus serotype 40LONG
 <400> 17

Thr Leu Trp Thr Thr Ala Asp Pro Ser Pro Asn Ala Thr Phe Tyr Glu
 1 5 10 15
 Ser Leu Asp Ala Lys Val Trp Leu Val Leu Val Lys Cys Asn Gly Met
 20 25 30
 Val Asn Gly Thr Ile Ser Ile Lys Ala Gln Lys Gly Thr Leu Leu Lys
 35 40 45

14

Pro Thr Ala Ser Phe Ile Ser Phe Val Met Tyr Phe Tyr Ser Asp Gly
 50 55 60
 Thr Trp Arg Lys Asn Tyr Pro Val Phe Asp Asn Glu Gly Ile Leu Ala
 65 70 75 80
 Asn Ser Ala Thr Trp Gly Tyr Arg Gln Gly Gln Ser Ala Asn Thr Asn
 85 90 95
 Val Ser Asn Ala Val Glu Phe Met Pro Ser Ser Lys Arg Tyr Pro Asn
 100 105 110
 Glu Lys Gly Ser Glu Val Gln Asn Met Ala Leu Thr Tyr Thr Phe Leu
 115 120 125
 Gln Gly Asp Pro Asn Met Ala Ile Ser Phe Gln Ser Ile Tyr Asn His
 130 135 140
 Ala Ile Glu Gly Tyr Ser Leu Lys Phe Thr Trp Arg Val Arg Asn Asn
 145 150 155 160
 Glu Arg Phe Asp Ile Pro Cys Cys Ser Phe Ser Tyr Val Thr Glu Gln
 165 170 175

<210> 18
 <211> 176
 <212> PRT
 <213> Human adenovirus serotype 41LONG

<400> 18

Thr Leu Trp Thr Thr Ala Asp Pro Ser Pro Asn Ala Thr Phe Tyr Glu
 1 5 10
 Ser Leu Asp Ala Lys Val Trp Leu Val Leu Val Lys Cys Asn Gly Met
 20 25 30
 Val Asn Gly Thr Ile Ser Ile Lys Ala Gln Lys Gly Ile Leu Leu Arg
 35 40 45
 Pro Thr Ala Ser Phe Ile Ser Phe Val Met Tyr Phe Tyr Ser Asp Gly
 50 55 60
 Thr Trp Arg Lys Asn Tyr Pro Val Phe Asp Asn Glu Gly Ile Leu Ala
 65 70 75 80
 Asn Ser Ala Thr Trp Gly Tyr Arg Gln Gly Gln Ser Ala Asn Thr Asn
 85 90 95
 Val Ser Asn Ala Val Glu Phe Met Pro Ser Ser Lys Arg Tyr Pro Asn
 100 105 110
 Gln Lys Gly Ser Glu Val Gln Asn Met Ala Leu Thr Tyr Thr Phe Leu
 115 120 125
 Gln Gly Asp Pro Asn Met Ala Ile Ser Phe Gln Ser Ile Tyr Asn His
 130 135 140
 Ala Leu Glu Gly Tyr Ser Leu Lys Phe Thr Trp Arg Val Arg Asn Asn
 145 150 155 160
 Glu Arg Phe Asp Ile Pro Cys Cys Ser Phe Ser Tyr Val Thr Glu Gln
 165 170 175

<210> 19
 <211> 188
 <212> PRT
 <213> Human adenovirus serotype 3

<400> 19

Thr Leu Trp Thr Gly Pro Lys Pro Glu Ala Asn Cys Ile Ile Glu Tyr
 1 5 10 15
 Gly Lys Gln Asn Pro Asp Ser Lys Leu Thr Leu Ile Leu Val Lys Asn
 20 25 30
 Gly Gly Ile Val Asn Gly Tyr Val Thr Leu Met Gly Ala Ser Asp Tyr
 35 40 45
 Val Asn Thr Leu Phe Lys Asn Lys Asn Val Ser Ile Asn Val Glu Leu
 50 55 60
 Tyr Phe Asp Ala Thr Gly His Ile Leu Pro Asp Ser Ser Ser Leu Lys
 65 70 75 80
 Thr Asp Leu Glu Leu Lys Tyr Lys Gln Thr Ala Asp Phe Ser Ala Arg
 85 90 95
 Gly Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe Val Leu Pro Asn Ala
 100 105 110
 Gly Thr His Asn Glu Asn Tyr Ile Phe Gly Gln Cys Tyr Tyr Lys Ala
 115 120 125
 Ser Asp Gly Ala Leu Phe Pro Leu Glu Val Thr Val Met Leu Asn Lys
 130 135 140
 Arg Leu Pro Asp Ser Arg Thr Ser Tyr Val Met Thr Phe Leu Trp Ser
 145 150 155 160
 Leu Asn Ala Gly Leu Ala Pro Glu Thr Thr Gln Ala Thr Leu Ile Thr
 165 170 175
 Ser Pro Phe Thr Phe Ser Tyr Ile Arg Glu Asp Asp
 180 185

<210> 20
 <211> 193
 <212> PRT
 <213> Human adenovirus serotype 7

<400> 20

Thr Leu Trp Thr Gly Val Asn Pro Thr Thr Ala Asn Cys Gln Ile Met
 1 5 10 15
 Ala Ser Ser Glu Ser Asn Asp Cys Lys Leu Ile Leu Thr Leu Val Lys
 20 25 30
 Thr Gly Gly Leu Val Thr Ala Phe Val Tyr Val Ile Gly Val Ser Asn
 35 40 45
 Asp Phe Asn Met Leu Thr Thr His Lys Asn Ile Asn Phe Thr Ala Glu
 50 55 60
 Leu Phe Phe Asp Ser Thr Gly Asn Leu Leu Thr Ser Leu Ser Ser Leu
 65 70 75 80
 Lys Thr Pro Leu Asn His Lys Ser Gly Gln Asn Met Ala Thr Gly Ala
 85 90 95
 Leu Thr Asn Ala Lys Gly Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe
 100 105 110
 Asn Val Asn Ser Arg Glu Lys Glu Asn Tyr Ile Tyr Gly Thr Cys Tyr
 115 120 125
 Tyr Thr Ala Ser Asp His Thr Ala Phe Pro Ile Asp Ile Ser Val Met
 130 135 140

Leu Asn Gln Arg Ala Leu Asn Asn Glu Thr Ser Tyr Cys Ile Arg Val
 145 150 155 160
 Thr Trp Ser Trp Asn Thr Gly Val Ala Pro Glu Val Gln Thr Ser Ala
 165 170 175
 Thr Thr Leu Val Thr Ser Pro Phe Thr Phe Tyr Tyr Ile Arg Glu Asp
 180 185 190

Asp

<210> 21
 <211> 193
 <212> PRT
 <213> Human adenovirus serotype 11A

<400> 21

Thr Leu Trp Thr Gly Ile Asn Pro Thr Glu Ala Asn Cys Gln Met Met
 1 5 10 15
 Asp Ser Ser Glu Ser Asn Asp Cys Lys Leu Ile Leu Thr Leu Val Lys
 20 25 30
 Thr Gly Ala Leu Val Thr Ala Phe Val Tyr Val Ile Gly Val Ser Asn
 35 40 45
 Asn Phe Asn Met Leu Thr Thr Tyr Arg Asn Ile Asn Phe Thr Ala Glu
 50 55 60
 Leu Phe Phe Asp Ser Ala Gly Asn Leu Leu Thr Ser Leu Ser Ser Leu
 65 70 75 80
 Lys Thr Pro Leu Asn His Lys Ser Gly Gln Asn Met Ala Thr Gly Ala
 85 90 95
 Ile Thr Asn Ala Lys Ser Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe
 100 105 110
 Asn Asn Asn Ser Arg Glu Lys Glu Asn Tyr Ile Tyr Gly Thr Cys His
 115 120 125
 Tyr Thr Ala Ser Asp His Thr Ala Phe Pro Ile Asp Ile Ser Val Met
 130 135 140
 Leu Asn Gln Arg Ala Ile Arg Ala Asp Thr Ser Tyr Cys Ile Arg Ile
 145 150 155 160
 Thr Trp Ser Trp Asn Thr Gly Asp Ala Pro Glu Gly Gln Thr Ser Ala
 165 170 175
 Thr Thr Leu Val Thr Ser Pro Phe Thr Phe Tyr Tyr Ile Arg Glu Asp
 180 185 190

Asp

<210> 22
 <211> 192
 <212> PRT
 <213> Human adenovirus serotype 16

<400> 22

Thr Leu Trp Thr Gly Ala Lys Pro Ser Ala Asn Cys Val Ile Lys Glu
 1 5 10 15
 Gly Glu Asp Ser Pro Asp Cys Lys Leu Thr Leu Val Leu Val Lys Asn
 20 25 30
 Gly Gly Leu Ile Asn Gly Tyr Ile Thr Leu Met Gly Ala Ser Glu Tyr
 35 40 45

Thr Asn Thr Leu Phe Lys Asn Asn Gln Val Thr Ile Asp Val Asn Leu
 50 55 60
 Ala Phe Asp Asn Thr Gly Gln Ile Ile Thr Tyr Leu Ser Ser Leu Lys
 65 70 75 80
 Ser Asn Leu Asn Phe Lys Asp Asn Gln Asn Met Ala Thr Gly Thr Ile
 85 90 95
 Thr Ser Ala Lys Gly Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe Ile
 100 105 110
 Thr Tyr Ala Thr Glu Thr Leu Asn Glu Asp Tyr Ile Tyr Gly Glu Cys
 115 120 125
 Tyr Tyr Lys Ser Thr Asn Gly Thr Leu Phe Pro Leu Lys Val Thr Val
 130 135 140
 Thr Leu Asn Arg Arg Met Leu Ala Ser Gly Met Ala Tyr Ala Met Asn
 145 150 155 160
 Phe Ser Trp Ser Leu Asn Ala Glu Glu Ala Pro Glu Thr Thr Glu Val
 165 170 175
 Thr Leu Ile Thr Ser Pro Phe Phe Phe Ser Tyr Ile Arg Glu Asp Asp
 180 185 190

<210> 23
 <211> 191
 <212> PRT
 <213> Human adenovirus serotype 21

<400> 23

Thr Leu Trp Thr Gly Ile Lys Pro Pro Pro Asn Cys Gln Ile Val Glu
 1 5 10 15
 Asn Thr Asp Thr Asn Asp Gly Lys Leu Thr Leu Val Leu Val Lys Asn
 20 25 30
 Gly Gly Leu Val Asn Gly Tyr Val Ser Leu Val Gly Val Ser Asp Thr
 35 40 45
 Val Asn Gln Met Phe Thr Gln Lys Ser Ala Thr Ile Gln Leu Arg Leu
 50 55 60
 Tyr Phe Asp Ser Ser Gly Asn Leu Leu Thr Asp Glu Ser Asn Leu Lys
 65 70 75 80
 Ile Pro Leu Lys Asn Lys Ser Ser Thr Ala Thr Ser Glu Val Leu Gln
 85 90 95
 Pro Ala Glu Ala Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe Asn Thr
 100 105 110
 Thr Thr Arg Asp Ser Glu Asn Tyr Ile His Gly Ile Cys Tyr Tyr Met
 115 120 125
 Thr Ser Tyr Asp Arg Ser Leu Val Pro Leu Asn Ile Ser Ile Met Leu
 130 135 140
 Asn Ser Arg Thr Ile Ser Ser Asn Val Ala Tyr Ala Ile Gln Phe Glu
 145 150 155 160
 Trp Asn Leu Asn Ala Lys Glu Ser Pro Glu Ser Asn Ile Ala Thr Leu
 165 170 175
 Thr Thr Ser Pro Phe Phe Phe Ser Tyr Ile Arg Glu Asp Asp Asn
 180 185 190

<210> 24
 <211> 193
 <212> PRT
 <213> Human adenovirus serotype 34A

<400> 24

Thr Leu Trp Thr Gly Val Asn Pro Thr Glu Ala Asn Cys Gln Ile Met
 1 5 10 15
 Asn Ser Ser Glu Ser Asn Asp Cys Lys Leu Ile Leu Thr Leu Val Lys
 20 25 30
 Thr Gly Ala Leu Val Thr Ala Phe Val Tyr Val Ile Gly Val Ser Asn
 35 40 45
 Asn Phe Asn Met Leu Thr Thr His Arg Asn Ile Asn Phe Thr Ala Glu
 50 55 60
 Leu Phe Phe Asp Ser Thr Gly Asn Leu Leu Thr Arg Leu Ser Ser Leu
 65 70 75 80
 Lys Thr Pro Leu Asn His Lys Ser Gly Gln Asn Met Ala Thr Gly Ala
 85 90 95
 Ile Thr Asn Ala Lys Gly Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe
 100 105
 Asn Asp Asn Ser Arg Glu Lys Glu Asn Tyr Ile Tyr Gly Thr Cys Tyr
 115 120 125
 Tyr Thr Ala Ser Asp His Thr Ala Phe Pro Ile Asp Ile Ser Val Met
 130 135 140
 Leu Asn Arg Arg Ala Ile Asn Asp Glu Thr Ser Tyr Cys Ile Arg Ile
 145 150 155 160
 Thr Trp Ser Trp Asn Thr Gly Asp Ala Pro Glu Val Gln Thr Ser Ala
 165 170 175
 Thr Thr Leu Val Thr Ser Pro Phe Thr Phe Tyr Tyr Ile Arg Glu Asp
 180 185 190

Asp

<210> 25
 <211> 191
 <212> PRT
 <213> Human adenovirus serotype 35

<400> 25

Thr Leu Trp Thr Gly Ile Asn Pro Pro Pro Asn Cys Gln Ile Val Glu
 1 5 10 15
 Asn Thr Asn Thr Asn Asp Gly Lys Leu Thr Leu Val Leu Val Lys Asn
 20 25 30
 Gly Gly Leu Val Asn Gly Tyr Val Ser Leu Val Gly Val Ser Asp Thr
 35 40 45
 Val Asn Gln Met Phe Thr Gln Lys Thr Ala Asn Ile Gln Leu Arg Leu
 50 55 60
 Tyr Phe Asp Ser Ser Gly Asn Leu Leu Thr Glu Glu Ser Asp Leu Lys
 65 70 75 80
 Ile Pro Leu Lys Asn Lys Ser Ser Thr Ala Thr Ser Glu Thr Val Ala
 85 90 95
 Ser Ser Lys Ala Phe Met Pro Ser Thr Thr Ala Tyr Pro Phe Asn Thr

			100						105						110			
Thr	Thr	Arg	Asp	Ser	Glu	Asn	Tyr	Ile	His	Gly	Ile	Cys	Tyr	Tyr	Met			
		115					120					125						
Thr	Ser	Tyr	Asp	Arg	Ser	Leu	Phe	Pro	Leu	Asn	Ile	Ser	Ile	Met	Leu			
	130					135					140							
Asn	Ser	Arg	Met	Ile	Ser	Ser	Asn	Val	Ala	Tyr	Ala	Ile	Gln	Phe	Glu			
145					150			145		155					160			
Trp	Asn	Leu	Asn	Ala	Ser	Glu	Ser	Pro	Glu	Ser	Asn	Ile	Ala	Thr	Leu			
				165					170					175				
Thr	Thr	Ser	Pro	Phe	Phe	Phe	Ser	Tyr	Ile	Thr	Glu	Asp	Asp	Asn				
			180					185					190					

<210> 26
 <211> 156
 <212> PRT
 <213> Human adenovirus serotype 40SHORT

<400> 26

Thr	Ile	Trp	Ser	Ile	Ser	Pro	Thr	Pro	Asn	Cys	Ser	Ile	Tyr	Glu	Thr			
1				5					10					15				
Gln	Asp	Ala	Asn	Leu	Phe	Leu	Cys	Leu	Thr	Lys	Asn	Gly	Ala	His	Val			
			20					25					30					
Leu	Gly	Thr	Ile	Thr	Ile	Lys	Gly	Leu	Lys	Gly	Ala	Leu	Arg	Glu	Met			
		35					40					45						
Asn	Asp	Asn	Ala	Leu	Ser	Val	Lys	Leu	Pro	Phe	Asp	Asn	Gln	Gly	Asn			
		50				55					60							
Leu	Leu	Asn	Cys	Ala	Leu	Glu	Ser	Ser	Thr	Trp	Arg	Tyr	Gln	Glu	Thr			
65					70					75				80				
Asn	Ala	Val	Ala	Ser	Asn	Ala	Leu	Thr	Phe	Met	Pro	Asn	Ser	Thr	Val			
				85					90					95				
Tyr	Pro	Arg	Asn	Lys	Thr	Ala	Asp	Pro	Gly	Asn	Met	Leu	Ile	Gln	Ile			
			100					105					110					
Ser	Pro	Asn	Ile	Thr	Phe	Ser	Val	Val	Tyr	Asn	Glu	Ile	Asn	Ser	Gly			
		115					120					125						
Tyr	Ala	Phe	Thr	Phe	Lys	Trp	Ser	Ala	Glu	Pro	Gly	Lys	Pro	Phe	His			
	130					135					140							
Pro	Pro	Thr	Ala	Val	Phe	Cys	Tyr	Ile	Thr	Glu	Gln							
145					150					155								

<210> 27
 <211> 156
 <212> PRT
 <213> Human adenovirus serotype 41SHORT

<400> 27

Thr	Ile	Trp	Ser	Ile	Ser	Pro	Thr	Pro	Asn	Cys	Ser	Ile	Tyr	Glu	Thr			
1				5					10					15				
Gln	Asp	Ala	Asn	Leu	Phe	Leu	Cys	Leu	Thr	Lys	Asn	Gly	Ala	His	Val			
			20					25					30					
Leu	Gly	Thr	Ile	Thr	Ile	Lys	Gly	Leu	Lys	Gly	Ala	Leu	Arg	Glu	Met			
		35					40					45						
His	Asp	Asn	Ala	Leu	Ser	Leu	Lys	Leu	Pro	Phe	Asp	Asn	Gln	Gly	Asn			

```

                    50                    55                    60
Leu Leu Asn Cys Ala Leu Glu Ser Ser Thr Trp Arg Tyr Gln Glu Thr
65                    70                    75                    80
Asn Ala Val Ala Ser Asn Ala Leu Thr Phe Met Pro Asn Ser Thr Val
                    85                    90                    95
Tyr Pro Arg Asn Lys Thr Ala His Pro Gly Asn Met Leu Ile Gln Ile
                    100                    105                    110
Ser Pro Asn Ile Thr Phe Ser Val Val Tyr Asn Glu Ile Asn Ser Gly
                    115                    120                    125
Tyr Ala Phe Thr Phe Lys Trp Ser Ala Glu Pro Gly Lys Pro Phe His
                    130                    135                    140
Pro Pro Thr Ala Val Phe Cys Tyr Ile Thr Glu Gln
145                    150                    155

```

<210> 28
 <211> 354
 <212> PRT
 <213> Anti-HA ScFv fused in frame with 2 C-terminal myc epitopes and PDGF receptor transmembrane anchor (Anti-HA pseudo-receptor)

```

<400> 28
Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1                    5                    10                    15
Gly Ser Thr Gly Asp Gly Ala Gln Pro Ala Asp Ile Val Met Thr Gln
                    20                    25                    30
Ser Pro Ser Ser Leu Thr Val Thr Ala Gly Glu Lys Val Thr Met Ser
                    35                    40                    45
Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser Gly Asn Gln Lys Asn Tyr
                    50                    55                    60
Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile
65                    70                    75                    80
Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val Pro Asp Arg Phe Thr Gly
                    85                    90                    95
Ser Gly Ser Gly Arg Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
                    100                    105                    110
Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn Asp Asn Ser His Pro Leu
                    115                    120                    125
Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg Ala Gly Gly Gly
130                    135                    140
Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Gln Leu
145                    150                    155
Val Glu Ser Gly Gly Asn Leu Val Asn Pro Gly Gly Ser Leu Lys Leu
                    165                    170                    175
Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr Gly Met Ser Trp
                    180                    185                    190
Val Arg Gln Thr Pro Asn Lys Arg Leu Glu Trp Val Pro Thr Ile Ile
                    195                    200                    205
Arg Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Ser Val Lys Gly Arg Phe
210                    215                    220
Thr Ile Ser Lys Asn Asn Ala Lys Asn Thr Leu Tyr Leu Gln Met Ser
225                    230                    235                    240

```

Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys Ala Lys Arg Glu
 245 250 255
 Thr Phe Asp Glu Lys Gly Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val
 260 265 270
 Thr Val Ser Ala Ala Ala Ala Glu Gln Lys Leu Ile Ser Glu Glu Asp
 275 280 285
 Leu Asn Gly Ala Val Asp Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu
 290 295 300
 Asn Ala Val Gly Gln Asp Thr Gln Glu Val Ile Val Val Pro His Ser
 305 310 315 320
 Leu Pro Phe Lys Val Val Val Ile Ser Ala Ile Leu Ala Leu Val Val
 325 330 335
 Leu Thr Ile Ile Ser Leu Ile Ile Leu Ile Met Leu Trp Gln Lys Lys
 340 345 350

Pro Val

<210> 29
 <211> 218
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Consensus
 sequence from the comparison between non-group B
 adenoviral knobs as indicated in Figures 1A and
 1B. Xaa is any amino acid or no amino acid as
 indicated in Figures 1A and 1B.

<400> 29
 Thr Leu Trp Thr Thr Pro Xaa Pro Ser Pro Asn Cys Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Lys Asp Xaa Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Xaa Val Ser Xaa Xaa Xaa Val Xaa Xaa Xaa Xaa Gly Xaa
 35 40 45
 Xaa Xaa Xaa Ile Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80
 Xaa Xaa Xaa Xaa Phe Asp Xaa Asn Gly Val Leu Xaa Xaa Xaa Ser Xaa
 85 90 95
 Xaa Xaa Xaa Leu Xaa Xaa Xaa Tyr Trp Asn Phe Arg Xaa Gly Xaa Xaa
 100 105 110
 Xaa Xaa Xaa Xaa Xaa Tyr Xaa Asn Ala Val Gly Phe Met Pro Asn Xaa
 115 120 125
 Xaa Ala Tyr Pro Lys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 130 135 140
 Ala Xaa Xaa Xaa Xaa Ile Val Xaa Xaa Xaa Xaa Tyr Leu Xaa Gly Xaa
 145 150 155 160
 Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Thr Xaa Asn Xaa Xaa Xaa Glu
 165 170 175

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Tyr Ser Xaa Xaa Phe Xaa Xaa
 180 185 190
 Xaa Trp Xaa Xaa Xaa Xaa Xaa Tyr Xaa Asn Xaa Xaa Phe Xaa Thr Xaa
 195 200 205
 Ser Xaa Thr Phe Ser Tyr Ile Ala Gln Glu
 210 215

<210> 30
 <211> 215
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Majority sequence from the comparison between non-group B adenoviral knobs as indicated in Figures 1A and 1B. Xaa is any amino acid or no amino acid as indicated in Figures 1A and 1B.

<400> 30
 Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro Asn Cys Thr Ile Asp Gln
 1 5 10 15
 Asp Lys Asp Ser Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Ile Leu Ala Thr Val Ser Leu Ile Val Val Xaa Ala Xaa Xaa Gly Lys
 35 40 45
 Leu Leu Ile Ile Asn Asn Thr Thr Asn Pro Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Lys Xaa Phe Thr
 65 70 75 80
 Ile Lys Leu Leu Phe Asp Ala Asn Gly Val Leu Leu Glu Asn Ser Asn
 85 90 95
 Xaa Xaa Xaa Leu Gly Lys Ala Tyr Trp Asn Phe Arg Asn Gly Asn Ser
 100 105 110
 Thr Val Ser Thr Ala Tyr Glu Asn Ala Val Gly Phe Met Pro Asn Leu
 115 120 125
 Val Ala Tyr Pro Lys Pro Thr Gly Xaa Ser Xaa Xaa Xaa Xaa Xaa Xaa
 130 135 140
 Ala Lys Asp Xaa Xaa Ile Val Tyr Gly Asn Val Tyr Leu Gly Gly Asp
 145 150 155 160
 Pro Asp Gln Pro Val Val Ile Lys Ile Thr Phe Asn Xaa Xaa Gln Glu
 165 170 175
 Thr Xaa Xaa Gly Ser Gly Tyr Ser Ile Thr Phe Asp Phe Ser Trp Ser
 180 185 190
 Lys Xaa Xaa Thr Tyr Ile Asn Val Glu Phe Glu Thr Thr Ser Phe Thr
 195 200 205
 Phe Ser Tyr Ile Ala Gln Glu
 210 215

<210> 31
 <211> 248
 <212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Consensus sequence from the comparison between non-group B adenoviral knobs as indicated in Figures 2A and 2B. Xaa is any amino acid or no amino acid as indicated in Figures 2A and 2B.

<400> 31

```

Thr Leu Trp Thr Gly Xaa Asn Pro Xaa Xaa Ala Asn Cys Gln Ile Xaa
 1           5           10
Xaa Xaa Xaa Xaa Ser Asn Asp Cys Lys Leu Thr Leu Xaa Leu Val Lys
      20           25           30
Asn Gly Gly Leu Val Asn Gly Tyr Val Xaa Leu Xaa Gly Val Xaa Ser
      35           40           45
Xaa Xaa Xaa Asn Xaa Leu Xaa Xaa Phe Thr Xaa Lys Asn Xaa Xaa Xaa
      50           55           60
Xaa Xaa Xaa Asn Ile Xaa Xaa Glu Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      65           70           75
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Phe Asp Ser Thr Gly Asn
      85           90           95
Leu Leu Thr Xaa Xaa Xaa Xaa Xaa Leu Ser Ser Leu Xaa Xaa Xaa Lys
      100          105          110
Thr Pro Leu Asn Xaa Lys Ser Xaa Gln Asn Met Ala Thr Gly Ala Xaa
      115          120          125
Thr Xaa Ala Lys Gly Phe Met Pro Ser Thr Thr Ala Tyr Xaa Xaa Pro
      130          135          140
Phe Asn Xaa Xaa Xaa Arg Glu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Glu Asn
      145          150          155
Tyr Ile Tyr Gly Xaa Cys Tyr Tyr Xaa Ala Ser Xaa Asp Xaa Thr Leu
      165          170          175
Phe Pro Leu Xaa Ile Ser Val Met Leu Asn Xaa Xaa Xaa Arg Xaa Ile
      180          185          190
Xaa Ser Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Thr Ser Xaa Xaa Tyr Xaa Ile
      195          200          205
Xaa Phe Xaa Trp Ser Leu Asn Ala Xaa Gly Xaa Ala Pro Xaa Xaa Glu
      210          215          220
Thr Xaa Xaa Xaa Xaa Xaa Xaa Thr Leu Xaa Thr Ser Pro Phe Thr Phe
      225          230          235          240
Ser Tyr Ile Arg Glu Asp Xaa Asp
      245

```

<210> 32

<211> 248

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Majority sequence from the comparison between non-group B adenoviral knobs as indicated in Figures 2A and 2B. Xaa is any amino acid or no amino acid as

indicated in Figures 2A and 2B.

<400> 32
 Thr Leu Trp Thr Gly Ile Asn Pro Xaa Glu Ala Asn Cys Gln Ile Met
 1 5 10 15
 Glu Ser Ser Glu Ser Asn Asp Cys Lys Leu Thr Leu Val Leu Val Lys
 20 25 30
 Asn Gly Gly Leu Val Asn Gly Tyr Val Tyr Leu Ile Gly Val Xaa Ser
 35 40 45
 Asp Thr Val Asn Met Leu Xaa Xaa Phe Thr Asn Lys Asn Xaa Xaa Xaa
 50 55 60
 Xaa Xaa Ile Asn Ile Thr Ala Glu Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 65 70 75 80
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Phe Phe Asp Ser Thr Gly Asn
 85 90 95
 Leu Leu Thr Xaa Xaa Xaa Xaa Ser Leu Ser Ser Leu Xaa Xaa Xaa Lys
 100 105 110
 Thr Pro Leu Asn His Lys Ser Gly Gln Asn Met Ala Thr Gly Ala Ile
 115 120 125
 Thr Ser Ala Lys Gly Phe Met Pro Ser Thr Thr Ala Tyr Xaa Xaa Pro
 130 135 140
 Phe Asn Thr Asn Ser Arg Glu Xaa Xaa Xaa Xaa Xaa Xaa Lys Glu Asn
 145 150 155 160
 Tyr Ile Tyr Gly Thr Cys Tyr Tyr Thr Ala Ser Xaa Asp His Thr Leu
 165 170 175
 Phe Pro Leu Asp Ile Ser Val Met Leu Asn Xaa Xaa Ser Arg Ala Ile
 180 185 190
 Ser Ser Glu Xaa Xaa Xaa Xaa Xaa Xaa Thr Ser Xaa Xaa Tyr Ala Ile
 195 200 205
 Arg Phe Thr Trp Ser Leu Asn Ala Xaa Gly Glu Ala Pro Xaa Xaa Glu
 210 215 220
 Thr Ser Xaa Xaa Xaa Ala Ala Thr Leu Val Thr Ser Pro Phe Thr Phe
 225 230 235 240
 Ser Tyr Ile Arg Glu Asp Xaa Asp
 245

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/US 99/20728

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/861 C07K14/075 C12N7/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 26281 A (WICKHAM, TJ; GENVEC INC; CORNELL RES FOUNDATION INC (US)) 29 August 1996 (1996-08-29) example 2 ---	1,4,5,8, 9
P,X	WO 98 44121 A (TRANSGENE SA; CNRS) 8 October 1998 (1998-10-08) page 8, line 4 - line 9 ---	1
P,X	WO 98 54346 A (GENVEC, INC.) 3 December 1998 (1998-12-03) page 3, line 3 - line 13; examples 1-4 ---	1
A	WO 98 13499 A (CIBA GEIGY AG ; SCRIPPS RESEARCH INST (US); MEMEROW GR (US); VO) 2 April 1998 (1998-04-02) page 59, paragraph 3 ---	1,9-11
	-/--	

Further documents are listed in the continuation of box C. Patent family members are listed in annex.

° Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>
--	--

Date of the actual completion of the international search 10 January 2000	Date of mailing of the international search report 17/01/2000
---	---

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Cupido, M
--	--

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/20728

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 97 06826 A (BOTH GW; COMMW SCIENT IND RES ORG (AU)) 27 February 1997 (1997-02-27)</p> <p>page 5, line 4; figures 3,4</p>	<p>1, 4, 5, 9-11, 22-27, 33, 34</p>
A	<p>WICKHAM T J ET AL: "INCREASED IN VITRO AND IN VIVO GENE TRANSFER BY ADENOVIRUS VECTORS CONTAINING CHIMERIC FIBER PROTEINS" JOURNAL OF VIROLOGY, US, THE AMERICAN SOCIETY FOR MICROBIOLOGY, vol. 71, no. 11, page 8221-8229 XP002911344 ISSN: 0022-538X page 8223, right-hand column -page 8224, left-hand column</p>	<p>11-33</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 20728

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 33 and 34 insofar they concern an in vivo method are directed to a method of treatment of the human or animal body, the search has been carried out and based on the alleged effects of the adenoviral vector.
2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter: International Application No
PCT/US 99/20728

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9626281	A	29-08-1996	US 5770442	23-06-1998
			A 698254	29-10-1998
			AU 4980496	11-09-1996
			CA 2213343	29-08-1996
			EP 0811069	10-12-1997
			JP 11500315	12-01-1999
WO 9844121	A	08-10-1998	FR 2761688	09-10-1998
			FR 2761689	09-10-1998
			AU 7054798	22-10-1998
WO 9854346	A	03-12-1998	AU 7604998	30-12-1998
WO 9813499	A	02-04-1998	AU 4624197	17-04-1998
			EP 0937150	25-08-1999
WO 9706826	A	27-02-1997	AU 708870	12-08-1999
			AU 6696696	12-03-1997
			CA 2229631	27-02-1997
			EP 0851769	08-07-1998
			JP 11511139	28-09-1999
			NZ 315295	29-09-1999