

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Crystalline form of fatty acid amine hydrolase (FAAH)

(51)⁶ International Patent Classification(s)
C12N 9/78 (2006.01) 20060101AFI2005100
C12N 9/78 8BMEP
PCT/US2003/036125

(21) Application No: 2003290780 (22) Application Date: 2003 .11.14

(87) WIPO No: W004/044169

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/426,788	2002 .11.14	US

(43) Publication Date : 2004 .06.03
(43) Publication Journal Date : 2004 .07.08

(71) Applicant(s)
The Scripps Research Institute

(72) Inventor(s)
Stevens, Raymond C., Bracey, Michael H., Hanson, Michael A., Cravatt, Benjamin F.

(74) Agent/Attorney
Pizzeys, Level 2, Woden Plaza Offices, Woden, ACT, 2606

CORRECTED VERSION

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 May 2004 (27.05.2004)

PCT

(10) International Publication Number
WO 2004/044169 A2

- (51) International Patent Classification⁷: C12N (74) Agent: HALE, Lisa, A.: Gray Cary Ware & Freidenrich
L.L.P., Suite 1100, 4365 Executive Drive, San Diego, CA
92121-2133 (US).
- (21) International Application Number: PCT/US2003/036125 (81) Designated States (national): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, GN, GU, HT, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MY, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 14 November 2003 (14.11.2003) (84) Designated States (regional): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/426,788 14 November 2002 (14.11.2002) US (15) Information about Correction:
see PCT Gazette No. 27/2004 of 1 July 2004, Section II
- (63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/426,788 (CIP)
Filed on 14 November 2002 (14.11.2002)
- (71) Applicant (for all designated States except US): THE
SCRIPPS RESEARCH INSTITUTE [US/US]; 10550
North Torrey Pines Road, La Jolla, CA 92037 (US).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): BRACEY, Michael,
H. [US/US]; 4946 Niagara Avenue, San Diego, CA 92107
(US). HANSON, Michael, A. [US/US]; 1703 Coast Blvd.,
Del Mar, CA 92014 (US). STEVENS, Raymond, C.
[US/US]; 7096 Caminito Valverde, La Jolla, CA 92037
(US). CRAVATT, Benjamin, E. [US/US]; 5555 La Jolla
Hermosa Avenue, La Jolla, CA 92037 (US).
- Published:
— without international search report and to be republished
upon receipt of that report
- (48) Date of publication of this corrected version:
1 July 2004
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2004/044169 A2

(54) Title: CRYSTALLINE FORM OF FATTY ACID AMINE HYDROLASE (FAAH)

(57) Abstract: The present invention is directed to FAAH crystals in complex with the inhibitor methoxyarachidonyl fluorophosphate (MAFP) and to the use of these crystals to determine the three-dimensional structure of FAAH. This invention is further directed to the use of this structure for the modeling or determination of the structures of related proteins. This invention is further directed to the use of this structure in the pursuit of drug design to identify, characterize, or optimize agents which bind to the active site, substrate channels, product channels, or regulatory sites of FAAH, and to the evaluation of these agents to identify agents which may stimulate, inhibit, relocalize, stabilize, or destabilize FAAH and/or its activity. This invention is further directed to the use of this structure in the development of engineered FAAH variants which display altered solubility, catalytic profiles, or substrate specificity. This invention is further directed to the use of this structure in the development of engineered heterologous proteins with altered membrane tropism.

CRYSTALLINE FORM OF FATTY ACID AMIDE HYDROLASE (FAAH)**GOVERNMENT SUPPORT**

[0001] This application was partially supported by National Institutes of Health Grant Nos. DA13173 and F32 MH12414-03. Accordingly, the U.S. Government may have an interest in this application.

FIELD OF THE INVENTION

[0002] The present invention relates generally to structural biology and medicine, especially in the interception of endocannabinoid influence and allied physiological processes. More specifically, the present invention relates to the crystalline form of fatty acid amide hydrolase (FAAH) and the use of these crystals to determine the three-dimensional structure of this protein.

BACKGROUND OF THE INVENTION

[0003] Fatty acid amide hydrolase (FAAH) is the only characterized mammalian member of the amidase signature (AS) family of serine hydrolases. This family is an ancient and ubiquitous group of enzymes that share an amino acid motif known as the amidase signature. The representatives of this family have a highly diverse array of substrates though they may share a common reaction mechanism. Most AS enzymes described to date hydrolyze amides of small metabolic intermediates, such as acetamide, opiates, propionamide, and malonamide. Those AS enzymes described to date also appear to be exclusively soluble proteins.

[0004] Though a member of the AS family, FAAH possesses unique features which distinguish it from lower homologues. FAAH behaves as an integral membrane protein when isolated from native sources or when expressed recombinantly. The enzyme cannot be separated from membrane fractions with the use of high salt concentrations or alkaline sodium carbonate. It can only be separated from membranes

with the use of detergents. FAAH protein extracted with the aid of non-denaturing detergents retains catalytic activity and hence presumably its native structure.

[0005] FAAH is present in mammals in various tissues throughout the body, including the brain, liver, duodenum, kidney, and testis. The enzyme is noticeably absent from the heart. In these tissues, the enzyme appears to reside on extensive intracellular membrane systems, likely the smooth endoplasmic reticulum (SER). This conclusion is supported by confocal immunofluorescence data from each of these tissues as well as immunogold electron microscopy of the rat liver. The mechanism by which FAAH is inserted into these membranes and how it is anchored there is currently a subject of active investigation.

[0006] Unlike its metabolic counterparts in lower organisms, FAAH retains an important role in nervous system function. The substrates of FAAH, the fatty acid amides, have been demonstrated to exert powerful neuromodulatory effects in test animals relating to the physiologies of pain, locomotion, memory, cognition, pyresis, and sleep. The mechanism by which these compounds exert their influence is not yet fully characterized, though a subset of these effects can be abrogated *in vivo* by antagonists of the trimetric G-protein coupled receptor CB1. The fatty acid amides are hydrolyzed by FAAH in an expeditious manner to their pharmacologically inactive acids. Therefore, FAAH acts to terminate the signaling of these molecules and to establish their baseline levels in the cell. As a result, FAAH intersects the physiologies associated with its substrates.

[0007] *In vivo* demonstrations of the critical role that FAAH plays in mammals has recently been documented with the aid of a genetically engineered mouse model that lacks a functional FAAH gene. These mice possess greatly enhanced sensitivities to the neurological effects of the fatty acid amides when administered by intraperitoneal injection. Further, these mice exhibit an increased tolerance for pain even in the naïve state. Concomitantly, brains from these mice exhibit highly elevated levels of fatty acid amides, suggesting that an absence of FAAH leads to disturbed endogenous fatty acid amide function and results in altered baseline physiology. These observations suggest

that a pharmacological intervention of the catalytic activity of FAAH may result in similar effects as the present genetic ablation.

[0008] The catalytic mechanism by which FAAH hydrolyzes its substrates has been investigated and several important findings have resulted. The amino acid residues that are responsible for the active site chemistry have been identified, as has at least one amino acid residue which resides in the substrate binding pocket. One particular finding of interest is the observation that FAAH is able to catalyze the hydrolysis of amides in the presence of equivalent amounts of comparable esters. This feature is contrary to solution chemistry and the action of the unrelated serine hydrolases of the catalytic triad family. How FAAH maintains this preference is currently unknown, though at least one amino acid residue, lysine 142, has been identified which contributes to this effect.

SUMMARY OF THE INVENTION

[0009] The present invention provides a crystallized mammalian fatty acid amide hydrolase (FAAH) in complex with the active site-directed inhibitor methoxyarachidonyl fluorophosphonate (MAFP). The present invention further provides derivatives of invention FAAH crystals including various heavy metals and methods for the collection of the X-ray diffraction patterns of both native and derivative crystals. The invention also provides methods for analyzing diffraction patterns produced by invention native and derivative crystals by multiple isomorphous replacement (MIR) and single- and multiwavelength anomalous diffraction (SAD/MAD).

[0010] In one embodiment, there is provided a three-dimensional model for the protein structure of FAAH at the secondary, tertiary, and quaternary levels. Identification of this structural model allows the analysis of FAAH's physicochemical properties and its application to understanding the enzyme's function and mechanism *in vivo*.

[0011] In another embodiment, there are provided methods for the identification of the active site of FAAH as well as its substrate binding residues. The invention provides the unambiguous localization of the MAFP molecule within the FAAH protein structure, and therefore allows the assignment of amino acid residues that interact with the enzyme's natural substrates as mimicked by the arachidonyl chain of the inhibitor.

[0012] In a further embodiment, there are provided methods for characterizing the physical aspects of the enzyme which allow heterologous agents to enter, interact with, or otherwise perturb the native structure of the enzyme. The invention describes the unpredicted presence of a second, alternate route of entry to the active site distinct from the cellular membrane. Further, the invention provides for the identification of a *src* homology (SH3) binding domain on an aqueous surface of the enzyme at a quaternary interface.

[0013] In a still further embodiment, there are provided methods for identifying, characterizing, and optimizing agents that interact with the internal channels of FAAH and therefore stimulate, inhibit, relocalize, stabilize, or destabilize FAAH and/or its activity. Such an agent might interact with FAAH, for example, at the active site, the substrate binding pocket, the membrane port, the cytosolic port, the dimerization tunnel, the membrane-binding domain, the alkyl tunnel, the head group tunnel, and the like. In another embodiment, there are provided methods for identifying agents that interact with the SH3-binding domain and the surface helix-loop-helix. Such an agent may be further identified, characterized, or optimized by comparison with the MAFP molecule. An interaction between FAAH and such an agent may be investigated with the aid of manual or computerized simulation of the interaction between the agent and FAAH. Such a simulation may then be optimized based on the structure of FAAH and observed complementarities and incompatibilities between FAAH and the agent under consideration. The effects of such an agent may then be determined by obtaining the agent and bringing it in contact with FAAH to measure its effects.

[0014] In yet another embodiment, there are provided methods for the elucidation of the three-dimensional structure of a protein or protein complex which is structurally related to FAAH. In this embodiment, there is provided a method by which a molecule or complex may be crystallized and X-ray diffraction data obtained from the crystal. The resulting diffraction data may then be compared with the present known three-dimensional structure of FAAH, and the structure of the molecule may be determined by the method of molecular replacement.

[0015] In a further embodiment, there are provided methods for identifying the domain of FAAH responsible for the enzyme's association with membranes (membrane-binding domain, MBD). In this embodiment, there are provided methods wherein the MBD may be combined with other molecules or molecular complexes to cause a novel membrane association in the assembly. Additionally, this method provides a means for removing or mutating the MBD in order to produce FAAH variants that have altered or abolished membrane-binding characteristics.

[0016] In yet another embodiment, there are provided methods for altering the active site of FAAH and its substrate recognition mechanism so as to affect a change in the enzyme's substrate selectivity. The method provides for the development of FAAH variants that degrade heterologous compounds. Such FAAH variants may prove useful as novel chemical catalysts for enantioselective amide hydrolysis.

[0017] In another embodiment, there are provided methods for screening an agent for the ability to modulate the activity of FAAH, comprising contacting FAAH with the agent to form a FAAH-agent complex and measuring the activity level of the FAAH-agent complex relative to un-complexed FAAH, thereby screening the agent for the ability to modulate the activity of FAAH.

BRIEF DESCRIPTION OF THE FIGURES

[0018] Figure 1 illustrates an overview of the three-dimensional structure of fatty acid amide hydrolase represented in ribbon diagram format. A number of relevant structural features are highlighted. Parts A and B are related by a 90 degree rotation about the vertical axis.

[0019] Figure 2 depicts the active site of FAAH in complex with the arachidonyl inhibitor MAP.

[0020] Figure 3 depicts the predicted membrane-binding cap of FAAH.

[0021] Figure 4 sets forth proposed modular adaptations that convert a soluble enzyme to an integral membrane enzyme based on differences in the structures of FAAH and MAE2.

[0022] Figure 5 depicts exemplary internal channels of FAAH which may be targeted by a subset of agents which interact with the enzyme.

[0023] The amino acid sequence of FAAH (SEQ ID NO: 1) is included herein.

DETAILED DESCRIPTION OF THE INVENTION

[0024] The present invention provides crystalline mammalian fatty acid amide hydrolase (FAAH) of sufficient quality to produce interpretable X-ray diffraction data. As used herein, the phrase "fatty acid amide hydrolase (FAAH)" includes any protein from any mammalian source that displays a catalytic activity to hydrolyze the amide bond of fatty acid amides and which further contains the amidase signature (AS) sequence. Such a protein may be derived from primary, recombinant, or synthetic sources. As used herein, the term "protein" refers to a protein, polypeptide or peptide.

[0025] FAAH has an amino acid sequence as set forth in SEQ ID NO: 1, including conservative variations thereof. As used herein, the phrase "conservative variations" refers to a replacement of an amino acid residue by another, biologically similar amino acid residue. Examples of conservative variations include the substitution of a hydrophobic residue such as isoleucine, valine, leucine or methionine for another, or the substitution of a polar residue for another, such as the substitution of arginine for lysine, glutamic for aspartic acid, or glutamine for asparagine, and the like.

[0026] Crystalline FAAH may be produced by combining FAAH protein with a mother liquor (for example, 100 mM sodium citrate pH 5.0, 100 mM lithium sulfate, 8% polyethylene glycol (average molecular weight 6000)) and allowing the mixture to vapor equilibrate with a reservoir of mother liquor for a sufficient time to afford crystals of FAAH. Suitable crystals are introduced into a collimated X-ray source such that the

crystal may be rotated and a complete diffraction pattern may be recorded. Such a diffraction pattern may then be analyzed and compared to independent patterns generated from isomorphous crystals derivatized with heavy metals. Heavy metals contemplated for use in the practice of the present invention include, osmium, platinum, and the like.

[0027] Alternately, such derivatized crystals may be exposed to X-rays of suitable wavelength(s) such that anomalous signals may be obtained for comparison. Such comparisons may then yield a molecular model of the three-dimensional structure of the FAAH protein. Such a model may be represented by several means, including computer records such as protein data bank (pdb) files (see, e.g., Table II). The invention described herein includes any representation, including but not limited to, binary records, text records, graphic representations, and virtual representations, each either physical or electronic. The present invention also provides for the introduction of an inhibitory substrate analogue to the protein so that its interactions with the protein may be determined.

[0028] FAAH crystals may be expected to take a wide variety of forms, all of which are included in the present invention. In one embodiment of the FAAH crystal, the crystal displays primitive monoclinic symmetry P21 and contains sixteen FAAH molecules per asymmetric unit. The monoclinic embodiment further exhibits pseudomerohedral twinning as the crystallographic unit cell axes *a* and *c* may be interchanged. In an alternate embodiment, the crystal displays C centered orthorhombic symmetry C2221 and contains eight molecules per asymmetric unit. In a third embodiment, the crystal displays primitive hexagonal symmetry P6322 and contains two molecules per asymmetric unit.

[0029] The structure further reveals that the FAAH protein does not exist in the crystal as a monomeric, independent unit, but instead it forms a dimer with simple two-fold symmetry about an axis parallel to the long axis of the monomer. This arrangement results in the concerted presentation of helices 15 through 19 from each monomer on the same face of the assembled dimer. Furthermore, these helices form a protruding, highly hydrophobic plateau on this face of the dimer. As such, this structural motif provides a

means for interacting with cellular membranes in a specific and stable manner. This face of the enzyme also displays a route of entry to the active site through which substrates or other agents might pass from the membrane. As used herein, "active site" refers to a region of a FAAH that, as a result of its shape and charge potential, interacts with an agent (including, without limitation, a protein, polypeptide, peptide, nucleic acid, including DNA or RNA, molecule, compound, antibiotic or drug).

[0030] A second route of access to the active site is found on a lateral face of the enzyme near the dimer interface. This second port is marked by histidine 449 and possibly tryptophan 445, each from the polypeptide chain of the dimer mate. A possible mechanism exists by which the enzyme may undergo structural rearrangements so that these ports may be opened and closed in a regulated manner in concert with the progress of the enzymatic reaction or as a means for controlling the enzyme's activity. Such a means of control may result from any number of mechanisms, including, but not limited to, the binding of protein partners or ligands. By means of example but not as a complete explanation, such an effect might be triggered by a representative of the SH3-domain class of proteins. The present structure demonstrates that FAAH displays at its apical surface a consensus polyproline sequence for binding SH3-domain proteins, as well as those of the Homer family. This motif resides on the loop between helices eleven and twelve and includes the amino acid sequence proline-proline-leucine-proline. A potential mechanism exists whereby a regulatory protein such as Homer or an exogenously supplied agent may bind the polyproline sequence on FAAH to introduce a conformational change which might alter the cytoplasmic ports of the enzyme or other structural features to change active site geometry or substrate/product transit.

[0031] In another embodiment, there are provided methods for identifying agents which interact with the internal channels of FAAH, based on the knowledge of the three-dimensional structure of FAAH, and thereby stimulate, inhibit, relocalize, stabilize, or destabilize FAAH and/or its activity. As used herein, the term "internal channels" refers to several sites within the protein, such as, the active site, the substrate binding pocket, the membrane port, the cytosolic port, the dimerization tunnel, the membrane-binding domain, the alkyl tunnel, the head group tunnel, and the like. The amino acids which

comprise the internal channels of FAAH are set forth in Table 5. In still another embodiment, there are provided methods for identifying agents which interact with the SH3-binding domain and the surface helix-loop-helix of FAAH.

[0032] Such agents may be identified, developed, and characterized based on an investigation of possible interactions between the agent and the FAAH molecule. This process might occur by visual inspection of the FAAH structure or, more quantitatively, with the aid of various computer algorithms designed for such purposes, such as AUTODOCK, InsightII, and QUANTA. The present invention allows for the virtual binding of candidate agents *in silico* so that they may be characterized as possible leads for further study. The effects of such agents may then be tested for activity against FAAH by bringing the agent into contact with FAAH and determining any effect on the enzyme's normal function. Such agents identified as candidates by invention methods will aid in identifying lead compounds for further chemical optimization and/or for evaluation *in vivo* as potential therapeutics. The invention provides that such optimized agents may be used in a pharmacological preparation designed to exert physiological effects when administered to living organisms. Such medically relevant uses for agents identified or optimized with the use of the present invention include, for example, the treatment of pain, sleep, addiction, fertility, anxiety, anorexia, fever, cognitive processes, and the like.

[0033] A candidate agent can be any type of molecule, including, for example, a peptide, a peptidomimetic, a polynucleotide, or a small organic molecule, that one wishes to examine for the ability to act as a therapeutic agent, which is an agent that provides a therapeutic advantage to a subject receiving it. It will be recognized that a method of the invention is readily adaptable to a high throughput format and, therefore, the method is convenient for screening a plurality of test agents either serially or in parallel. The plurality of test agents can be, for example, a library of test agents produced by a combinatorial method library of test agents. Methods for preparing a combinatorial library of molecules that can be tested for therapeutic activity are well known in the art and include, for example, methods of making a phage display library of peptides, which can be constrained peptides (see, for example, U.S. Patent No. 5,622,699; U.S. Patent

No. 5,206,347; Scott and Smith, Science 249:386-390, 1992; Markland et al., Gene 109:1319, 1991; each of which is incorporated herein by reference); a peptide library (U.S. Patent No. 5,264,563, which is incorporated herein by reference); a peptidomimetic library (Blondelle et al., Trends Anal. Chem. 14:8392, 1995; a nucleic acid library (O'Connell et al., *supra*, 1996; Tuerk and Gold, *supra*, 1990; Gold et al., *supra*, 1995; each of which is incorporated herein by reference); an oligosaccharide library (York et al., Carb. Res., 285:99128, 1996; Liang et al., Science, 274:1520-1522, 1996; Ding et al., Adv. Expt. Med. Biol., 376:261-269, 1995; each of which is incorporated herein by reference); a lipoprotein library (de Kruif et al., FEBS Lett., 399:232-236, 1996, which is incorporated herein by reference); a glycoprotein or glycolipid library (Karaoglu et al., J. Cell Biol., 130:567-577, 1995, which is incorporated herein by reference); or a chemical library containing, for example, drugs or other pharmaceutical agents (Gordon et al., J. Med. Chem., 37:1385-1401, 1994; Ecker and Crooke, Bio/Technology, 13:351-360, 1995; each of which is incorporated herein by reference). Accordingly, the present invention also provides a therapeutic agent identified by such a method.

[0034] The route of administration of a candidate agent will depend, in part, on the chemical structure of the candidate agent. Peptides and polynucleotides, for example, are not particularly useful when administered orally because they can be degraded in the digestive tract. However, methods for chemically modifying peptides, for example, to render them less susceptible to degradation by endogenous proteases or more absorbable through the alimentary tract are well known (see, for example, Blondelle et al., Trends Anal. Chem. 14:83-92, 1995; Ecker and Crooke, Bio/Technology, 13:351-360, 1995; each of which is incorporated herein by reference). In addition, a peptide agent can be prepared using D-amino acids, or can contain one or more domains based on peptidomimetics, which are organic molecules that mimic the structure of peptide domain; or based on a peptoid such as a vinylogous peptoid.

[0035] In another embodiment, there are provided pharmaceutical compositions including agents identified by invention methods. Invention pharmaceutical compositions can administered in a variety of ways including, for example, orally or parenterally, such as intravenously, intramuscularly, subcutaneously, intraorbitally, intracapsularly,

intraperitoneally, intrarectally, intracisternally or by passive or facilitated absorption through the skin using, for example, a skin patch or transdermal iontophoresis, respectively. Furthermore, the invention pharmaceutical compositions can be administered by injection, incubation, orally or topically, the latter of which can be passive, for example, by direct application of an ointment, or active, for example, using a nasal spray or inhalant, in which case one component of the composition is an appropriate propellant.

[0036] The total amount of pharmaceutical composition to be administered in practicing a method of the invention can be administered to a subject as a single dose, either as a bolus or by infusion over a relatively short period of time, or can be administered using a fractionated treatment protocol, in which multiple doses are administered over a prolonged period of time. The pharmaceutical composition can be formulated for oral formulation, such as a tablet, or a solution or suspension form; or can comprise an admixture with an organic or inorganic carrier or excipient suitable for enteral or parenteral applications, and can be compounded, for example, with the usual non-toxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, suppositories, solutions, emulsions, suspensions, or other form suitable for use. The carriers, in addition to those disclosed above, can include glucose, lactose, mannose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition auxiliary, stabilizing, thickening or coloring agents and perfumes can be used, for example a stabilizing dry agent such as triulose (see, for example, U.S. Pat. No. 5,314,695).

[0037] In another embodiment, there are provided methods for treating a pathological condition, including administering to a subject in need thereof an invention pharmaceutical composition. In one embodiment, the subject is a mammal. In another embodiment, the subject is human. Pathological conditions that can be effectively treated by invention methods include, for example, anxiety, pain, hunger, sleep, fertility,

cognition, immunological disorders, fever, tremor, glaucoma, intestinal disorders, and the like.

[0038] In a still further embodiment, there are provided methods for determining the molecular structure of a molecule or molecular complex whose structure is unknown, comprising:

(a) obtaining crystals of the molecule or molecular complex whose structure is unknown;

(b) generating X-ray diffraction data from the crystallized molecule or molecular complex;

(c) comparing the X-ray diffraction data from the molecule or molecular complex with the three dimensional structure determined from the crystalline form of FAAH; and

(d) using molecular replacement analysis to conform the three dimensional structure determined from the crystalline form of FAAH to the X-ray diffraction data from the crystallized molecule or molecular complex. This method allows for the determination, for example, of a protein which is structurally related to FAAH. As used herein, "structurally related" refers to a protein whose structure can be determined by comparing its X-ray diffraction data to the X-ray diffraction data of FAAH. Such a use of the present invention would include the crystallization of the target protein or protein complex and collection of its X-ray diffraction data. The target data may then be interpreted by means of molecular replacement using the FAAH structure as a template. Such methods obviate the need to collect data from derivatized crystals and greatly simplify the process of solving the structure of the target protein.

[0039] In a still further embodiment, there are provided methods for engineering proteins with altered physicochemical properties using the structural information obtained from the FAAH crystal. The invention allows for the identification of the membrane-binding domain of the FAAH protein. As such, this domain may be mutated or deleted, in whole or in part, to result in a protein with altered or absent membrane tropism. Further, the identified membrane-binding domain may, in full or in part, be transferred to a heterologous protein so as to affect its membrane tropism. Moreover, the identification

of the active site and substrate binding pocket of FAAH allows for the targeted generation of variants with altered substrate specificity and/or altered enzymological properties. Such variants may prove useful as catalysts of amide hydrolysis unrelated to the fatty acid amides, and as such may have chemical or industrial applications.

[0040] The invention will now be described in greater detail by reference to the following non-limiting example.

EXAMPLE

Crystallization

[0041] Protein crystals were obtained by recombinantly expressing rat FAAH with 29 amino acids at the amino-terminus deleted. Though the deleted region is predicted to participate in membrane binding (β), the truncated FAAH variant (residues 30 - 579) retains the wild-type protein's association with membranes, detergent requirement for solubilization, and ability to degrade fatty acid amides in mammalian cells (15). Crystals of FAAH complexed with methoxy arachidonyl fluorophosphonate, an active site-directed irreversible inhibitor, were obtained in three space groups, P2₁ with sixteen molecules in the asymmetric unit, C222₁ with eight molecules in the asymmetric unit, and P6₃22 with two molecules in the asymmetric unit (Materials and Methods (16)). The structure of the FAAH-inhibitor complex was first determined to 3.3 Å using a combination MIRAS / MAD phasing strategy followed by phase extension with multi-crystal and NCS averaging (Table 1). The resulting electron density maps displayed density for over 90% of the side chains with no main chain breaks, yielding an exceptional search model for molecular replacement phasing of twinned monoclinic data. The final model, built in P2₁ at 2.8 Å resolution, was of excellent quality.

Data Collection

[0042] Table 1 includes a summary of data collection and refinement statistics. Initial phases were calculated with DmMulti (24), incorporating all MAD and MIRAS

contributions. Ramachandran distributions of amino acids in the final model were calculated with PROCHECK (24) from a single protein chain after phase refinement under tight NCS restraints. Side chains for which no density was observed were set to zero occupancy during refinement, and no sigma-based rejection of reflections was employed. Numbers in parentheses refer to data in the highest resolution shell. Two residues were in disallowed regions: glutamate 122, which is in the $i+1$ position of a gamma turn, and glutamine 189, which is also in a tight turn immediately followed by a second turn. Table 2 includes the complete data collection and refinement statistics.

Table 1

Data set	TWIN		MAD		
	T1	Ta peak	Ta inflection	Os peak	Remote
Beam line	ALS 5.0.2	SSRL 9-2	SSRL 9-2	SSRL 9-2	SSRL 9-2
Wavelength (Å)	0.97910	1.255	1.2584	1.14025	0.925
Space group	P2 ₁	C222 ₁	C222 ₁	C222 ₁	C222 ₁
Resolution (Å)	50-2.8	65-3.3	65-3.3	60-3.3	48-3.3
Unique reflections	173615	81567	81574	81609	81511
Redundancy	2.1	14.5	14.5	14.5	14.5
Completeness*	76 (76)	99.7 (99.1)	99.7 (99.2)	99.8 (99.5)	99.7 (98.7)
$I(\sigma)$ *	4.4 (2.7)	8.2 (2.3)	6.9 (1.2)	7.6 (1.7)	8.5 (2.2)
R _{sym} (%)*	10.4 (26.4)	7.9 (33.3)	9.7 (63.2)	8.5 (45.5)	7.4 (33.4)
Phasing from Ta₃Cl₁₂ + OsCl₂ MAD data			Refinement against detwinned data set T1		
Resolution (Å)		96-3.3	Resolution (Å)	50-2.8	
Number of Os sites		8	Reflections used	173615	
Number of Ta sites		0	Test reflections	9297	
Os peak isomorphous phasing power (centric/acentric)		0.948/1.43	Number of nonhydrogen atoms	65744	
Os peak anomalous phasing power (acentric)		2.18	R _{int} (%)	19.7	
Remote isomorphous phasing power (centric/acentric)		0.288/0.471	R _{free} (%)	24.9	
Remote anomalous phasing power (acentric)		1.88	Average B value (Å ²)	44.829	
Ta inflection anomalous phasing power (acentric)		0.871	MAFP average B value (Å ²)	64.26	
Ta peak (reference) anomalous phasing power (acentric)		1.02	rmsd bond length (Å)	0.079	
			rmsd bond angle (°)	2.500	
Combined figure of merit before NCS and multi-crystal averaging (3.3 Å) (centric/acentric)		0.47/0.60	Ramachandran distributions (%)		
			Most favored	79.9	
			Additionally allowed	17.7	
Twinning operator		(l, -h, k)	Generously allowed	2.0	
Twinning factor (α)		0.28	Disallowed [†]	0.4	

Table 2

Data set	NATIVES				SAD			
	N1	N2	N3	T1	(NH ₄) ₂ PtCl ₂	OsCh	Ta peak	
Beam line	SSRL 9-2	SSRL 11-1	SSRL 11-1	SSRL 9-2	SSRL 11-1	SSRL 9-2	SSRL 9-2	
Wavelength (Å)	1.03317	1.07185	1.07185	0.87946	1.07185	1.13973	1.235	
Space group	C222 ₁	C222 ₁	C222 ₁	P6 ₃ 22	C222 ₁	C222 ₁	C222 ₁	
Resolution (Å)	60-5.0	70-4.0	64.5-3.8	50-3.5	70-4.0	63-4.0	65-3.3	
Unique reflections	19722	46254	53415	18446	45171	46181	81567	
Redundancy	4.1	4.3	4.7	20	2.1	9.5	14.5	
Completeness*	83 (83)	98.7 (100)	99.7 (99.7)	99.6 (99.9)	76 (76)	99.3 (99.0)	99.7 (99.1)	
I/(σI) [†]	5.0(2.8)	5.9 (2.6)	9.1 (4.0)	4.6 (1.9)	4.4 (2.7)	7.0 (3.2)	8.2 (2.3)	
Rsym (%) [‡]	10.5 (27.2)	9.7 (26.5)	6.7 (18.6)	12.6 (48.8)	10.4 (26.4)	9.4 (22.4)	7.9 (33.3)	
Phasing from MIRAS data								
Resolution (Å)			N1	N2	N3			
Number of Pt sites			48-5.0	46-4.0	48-4.0			
Number of Os sites			13	13	13			
Pt anomalous phasing power (acentric)			0.062/0.173	0.107/0.178	0.123/0.181			
Pt isomorphous phasing power (centric/acentric)			0.583	0.987	0.548			
Pt anomalous phasing power (acentric)			0.555/0.632	0.436/0.638	0.408/0.568			
Os isomorphous phasing power (centric/acentric)			0.767	0.736	0.739			
Os anomalous phasing power (acentric)								
Phasing from Ta ₂ Cl ₁₀ • OsCl ₄ MAD data								
Resolution (Å)				96-3.3				
Number of Os sites				8				
Number of Ta sites				0				
Os peak isomorphous phasing power (centric/acentric)				0.948/1.43				
Os peak anomalous phasing power (acentric)				2.18				
Remote isomorphous phasing power (centric/acentric)				0.268/0.471				
Remote anomalous phasing power (acentric)				1.88				
Ta inflection anomalous phasing power (acentric)				0.871				
Ta peak (reference) anomalous phasing power (acentric)				1.02				
Combined figure of merit before NCS and multi-crystal averaging (3.3 Å)								
				0.470/0.60				
Ramachandran distributions (%)								
				Most favored				
				0.948/1.43				
				2.18				
				0.268/0.471				
				1.88				
				0.871				
				1.02				
Twinning operator								
				0.470/0.60				
Twinning factor (σ)								
				0.470/0.60				

*Numbers in parentheses refer to data in the highest resolution shell.

†Two residues were in disallowed regions: glutamate 122, which is in the H+1 position of a gamma turn, and glutamine 169, which is also in a tight turn immediately followed by a second turn.

[0043] The crystal structure of FAAH reveals a dimeric enzyme (Fig. 1), consistent with chemical crosslinking studies indicating that the enzyme is a dimer in solution (15). The dimer interface buries approximately 1560 Å² of molecular surface area per monomer (17). The protein core is characterized by a twisted beta sheet consisting of eleven mixed strands which is in turn surrounded by twenty-four alpha helices of varying lengths. The overall fold of the monomer closely resembles that of malonamidase (MAE2) from the nitrogen fixing bacterium *Bradyrhizobium japonicum* (12) and peptide amidase (PAM) from *Stenotrophomonas maltophilia* (REF: Labahn J, Neumann S, Buldt G, Kula M, Granzin J. J Mol Biol 2002 Oct 4;322(5):1053).

[0044] However, key structural features make FAAH unique, and these will be discussed below. Further, these variations can be quantified by measuring the root mean squared deviation (rmsd) between FAAH and its bacterial homologues (Tables 3 and 4). If a comparison is drawn between domains of the proteins which obey amino acid sequence alignment, the rmsd difference increases as more atoms are included in the calculation. This is true both in the case of restricting the calculation to an expanding sphere centered on the active site and in the case of expanding the calculation from C α s to all atoms. In any case, it is also of note that only a fraction of FAAH can be included in the calculation due to sequence considerations, 42% in the case of PAM and 36% in the case of MAE2 (Table 3). Further, if protein fold topology is taken into consideration, the comparable fractions of FAAH which can be included fall even further, to 22% for PAM and 34% for MAE2 (Table 4).

Table 3. Average rmsd_{xyz} of sequence-based equivalent residue ranges after superposition of catalytic triad C α s. The number of amino acid residues included in the calculation is indicated.

	5Å	10Å	20Å	100Å
PAM-CHY*				
C α (222)	1.217	4.443	6.442	8.525
BACKBONE	2.361	4.044	6.452	8.494
ALL ATOMS	2.195	4.486	6.752	8.737
MAE2*				
C α (191)	0.433	2.191	5.487	9.092
BACKBONE	1.272	2.088	4.634	8.702
ALL ATOMS	1.253	2.363	4.827	8.783

Table 4. Average rmsd_{xyz} of topology-based equivalent residue ranges. The number of amino acid residues included in the calculation is indicated.

	5Å	10Å	20Å	100Å
PAM-CHY*				
C α (116)	0.415	1.136	5.473	5.449
BACKBONE	0.553	1.126	5.350	5.488
ALL ATOMS	0.504	1.126	5.147	5.290
MAE2*				
C α (178)	0.561	1.044	1.146	1.384
BACKBONE	0.661	1.028	1.221	1.429
ALL ATOMS	0.606	1.134	1.341	1.664

*Insertions/deletions and non-identical amino acid side chains are not included in the calculation. PAM-CHY is the complex of peptide amidase and the inhibitor chymostatin.

[0045] The active site of FAAH was identified based on the location of core catalytic residues defined previously by mutagenesis (10, 18) and from the density of the inhibitor adduct methoxy arachidonyl phosphonate (MAP) (Fig. 2). The catalytic nucleophile, serine 241, is covalently bonded to the phosphorous of the MAP molecule,

and the neighboring density can be modeled to accommodate the arachidonyl chain in an energetically favorable conformation (Fig. 2A). As observed for MAE2, the serine nucleophile in FAAH forms part of an unusual serine-serine-lysine catalytic triad with serine 217 and lysine 142 (Fig. 2B), and the relative orientation of the catalytic residues is nearly superimposable between the structures of FAAH and MAE2 (rmsd=0.27 Å). In contrast, the substrate-binding pockets of FAAH and MAE2 differ significantly. The tunnel leading from the surface of FAAH and containing the buried arachidonyl chain is lined with a preponderance of aromatic and aliphatic amino acids (Fig. 2B), many of which arise from sequence insertions in FAAH relative to MAE2. These residues include isoleucine 491, identified by UV cross-linking and mutagenesis studies to participate in substrate recognition (14). In contrast, the active site of MAE2 is lined with mostly hydrophilic amino acids that accommodate its soluble substrate, malonamide (12).

[0046] Amino acids 410 to 438, another sequence insertion in FAAH relative to MAE2, form a helix-turn-helix motif that interrupts the AS fold. The two helices, $\alpha 18$ and $\alpha 19$, cap the active site and present several hydrophobic residues that likely compose FAAH's membrane binding face (Figs. 1, 3A). The *N*-terminus of the intact enzyme, predicted by sequence analysis to form an additional membrane-binding helix (amino acids 7-29), would be properly positioned to reinforce the membrane interactions of the $\alpha 18$ and $\alpha 19$ membrane cap (Fig. 1B).

[0047] A potential substrate entryway is adjacent to $\alpha 18$ and $\alpha 19$, and the arachidonyl chain of MAP contacts phenylalanine 432 of $\alpha 18$, which may indicate direct access between the FAAH active site and the hydrophobic membrane bilayer. The putative substrate entry is amphipathic in nature with hydrophobic residues covering three sides of the rim and charged residues arginine 486 and aspartate 403 completing the remaining side (Fig. 2A, 3B). This arrangement of residues may accommodate the admission and movement of polar fatty acid amide head groups towards the active site. Overall, the intimate relationship between the membrane binding surface and active site of FAAH is similar to that of squalene cyclase (19) and prostaglandin H₂ synthase (20) which also act upon lipid soluble substrates and have hydrophobic caps surrounding their respective active site entrances. In all three enzymes, the hydrophobic cap is surrounded by basic amino acids (Fig. 3B) that may interact with negatively charged phospholipids.

[0048] A second major tunnel emerges from the active site at an approximately 80 degree angle from the arachidonyl-filled cavity. This tunnel bifurcates to create both a solvent-exposed cytosolic port and a route blocked by tryptophan 445 (Fig. 2A), a residue that forms a lock-and-key intersubunit contact. Thus, the FAAH active site appears to simultaneously access both the aqueous environment of the cytoplasm and the lipid milieu of the bilayer. This architecture may provide an exit route to the cytosol for the polar amine substituents liberated from the fatty acid amide substrates and could also provide entry for a water molecule required for deacylation of the FAAH-fatty acyl intermediate (18).

[0049] Overall, a comparison of FAAH's structure to that of the soluble AS enzyme MAE2 reveals how a single member of a large clade of proteins has adapted to perform a specialized cellular function through the addition of discrete folding modules and without gross changes in catalytic mechanism or fold architecture (Fig. 4). For example, although FAAH and MAE2 dimerize about roughly the same face, they have different relative monomer orientations. These distinct quaternary orientations produce an antiparallel and parallel alignment of the MAE2 and FAAH monomer active site entrances, respectively. The parallel orientation of FAAH monomers has important biological implications, as it should permit both subunits to function concurrently by recruiting substrates from the same membrane. Furthermore, this parallel alignment places another key structural module, the α 18- α 19 hydrophobic cap, on the same face of the dimer, thereby enhancing membrane binding (Fig. 4). Additional sequence insertions in the FAAH protein account for other specialized features of the enzyme's structure, including its cytoplasmic channel and apolar substrate binding pocket (Fig. 4).

[0050] The intimate relationship between the active site of FAAH and cell membranes revealed by the enzyme's structure raises the possibility that fatty acid amides need not be transported through aqueous cellular compartments in order to proceed from site of action to site of degradation. As a consequence, proper endocannabinoid tone may rely on both the expression levels of FAAH and its localization relative to CB receptor systems *in vivo* (21). Finally, the structure of FAAH may facilitate the design of inhibitors that could have therapeutic benefit for a variety of nervous system disorders (22), including not only agents that bind to the enzyme acyl chain pocket, but also

compounds that target the active site via the cytoplasmic channel. This latter class of inhibitors might display exceptional selectivity for FAAH relative to the more than one hundred lipid hydrolases present in the human proteome.

[0051] Figure 1 depicts the structure of the integral membrane protein fatty acid amide hydrolase (FAAH). The enzyme is a homodimer assembled from 63 kDa subunits. The inhibitor adduct methoxy arachidonyl phosphonate (MAP) is depicted in the active site with van der Waal surfaces rendered in yellow (carbon), gray (phosphorous), and red (oxygen). (A) Front view of the enzyme, highlighting the central twisted beta sheet that forms the core of the structure. (B) Side view produced by a 90° rotation from view A. The amino (N) and carboxyl (C) termini of the green subunit are labeled. Proline 129 is indicated. A genetic polymorphism conferring substitution of threonine at this position has been implicated in drug abuse and renders the enzyme more vulnerable to proteolytic attack (25). All figures were produced with Molscript (26), GRASP (23), and Raster3D (27). The membrane model was generated by Molecular Dynamics simulation of a palmitoyloleoylphosphatidylethanolamine (POPE) bilayer (28).

[0052] Figure 2 depicts the active site of FAAH in complex with the arachidonyl inhibitor MAP. (A) The FAAH dimer is shown with the protein surface rendered in gray (hydrophilic) and green (hydrophobic) for one subunit and pale yellow for its mate. A portion of the protein surface has been removed to highlight the continuous internal channel leading from the membrane binding face at arginine 486 (blue) and aspartate 403 (red) to the active site and on to the cytosolic port. Electron density corresponding to the arachidonyl inhibitor is shown in violet and lies in the hydrophobic (green) substrate-binding pocket. Tryptophan 445 from the dimer mate is rendered in van der Waals surface and colored cyan to demonstrate the effective plugging of this potential port. (B) Aromatic and aliphatic residues in the substrate binding pocket surrounding the arachidonyl chain and their interactions are indicated. Also shown is the unusual serine-serine-lysine catalytic triad of FAAH (ser241-ser217-ser241), with the serine 241 nucleophile covalently bonded to the phosphorus atom of the MAP inhibitor.

[0053] Figure 3 depicts the predicted membrane-binding cap of FAAH. (A) The hydrophobic helices $\alpha 18$ and $\alpha 19$ that comprise the putative membrane-binding cap are

shown in green. The primary sequence of this domain (404-433) is indicated using amino acid one letter code (29) except for hydrophilic amino acids, which are indicated by *x*. Five of these seven hydrophilic residues are arginine or lysine; the remaining two are serines. (B) The molecular surface of FAAH viewed from the membrane face; the observed structure demonstrates the presence of a hydrophobic cap (*top, green*), surrounded primarily by positive electrostatic potential (*bottom; blue, basic; red, acidic*). The entrance to the active site from the membrane face is indicated, as are arginine 486 and aspartate 403, which form one side of this access port.

[0054] Figure 4 sets forth proposed modular adaptations that convert a soluble enzyme to an integral membrane enzyme based on differences in the structures of FAAH and MAE2. The soluble enzyme, if oligomeric, reorganizes so that all active sites can concurrently access the bilayer for maximal efficiency. The addition of discrete structural elements, represented here in red, confer the new oligomerization domain (tryptophan 445 and residues 299-314; *lower left panel*) as well as a membrane binding face (*lower middle left panel*) and cytoplasmic access channel (*lower middle right panel*). The final monotopic integral membrane enzyme must also undergo mutation of key substrate binding residues in the active site to effectively recruit its hydrophobic targets from the lipid bilayer (*lower right panel*).

Table 5. Residues expected to line the interior channels of FAAH. The coordinates of the C α s in the present structure are provided as a guide only.

ATOM	0	CA	LYS	A	142	12.558	-31.943	29.679	0.00	0.00
A	C									
ATOM	0	CA	SER	A	190	17.871	-20.245	32.364	0.00	0.00
A	C									
ATOM	0	CA	MET	A	191	17.760	-22.751	29.499	0.00	0.00
A	C									
ATOM	0	CA	LEU	A	192	19.766	-20.186	27.640	0.00	0.00
A	C									
ATOM	0	CA	SER	A	193	22.793	-22.387	27.483	0.00	0.00
A	C									
ATOM	0	CA	PHE	A	194	24.110	-25.405	25.603	0.00	0.00
A	C									
ATOM	0	CA	SER	A	217	19.643	-27.223	27.046	0.00	0.00
A	C									
ATOM	0	CA	THR	A	236	12.999	-27.716	22.998	0.00	0.00
A	C									
ATOM	0	CA	ILE	A	238	15.579	-22.197	19.986	0.00	0.00
A	C									

ATOM	0	CA	GLY A 239	19.319	-22.114	19.282	0.00	0.00
A	C							
ATOM	0	CA	GLY A 240	20.185	-25.826	19.177	0.00	0.00
A	C							
ATOM	0	CA	SER A 241	19.006	-27.246	22.499	0.00	0.00
A	C							
ATOM	0	CA	PHE A 244	23.393	-30.169	20.863	0.00	0.00
A	C							
ATOM	0	CA	THR A 257	9.128	-26.186	12.131	0.00	0.00
A	C							
ATOM	0	CA	ASN A 259	4.442	-24.354	14.717	0.00	0.00
A	C							
ATOM	0	CA	LEU A 261	3.547	-28.979	17.435	0.00	0.00
A	C							
ATOM	0	CA	SER A 262	1.827	-27.120	20.226	0.00	0.00
A	C							
ATOM	0	CA	LYS A 263	3.541	-25.099	22.924	0.00	0.00
A	C							
ATOM	0	CA	SER A 264	0.514	-24.586	24.939	0.00	0.00
A	C							
ATOM	0	CA	LEU A 266	5.685	-24.877	27.877	0.00	0.00
A	C							
ATOM	0	CA	GLY A 268	9.936	-20.496	29.403	0.00	0.00
A	C							
ATOM	0	CA	CYS A 269	11.633	-17.639	31.102	0.00	0.00
A	C							
ATOM	0	CA	VAL A 270	10.747	-14.992	28.633	0.00	0.00
A	C							
ATOM	0	CA	TYR A 271	7.552	-14.969	26.504	0.00	0.00
A	C							
ATOM	0	CA	GLY A 272	6.363	-13.140	23.340	0.00	0.00
A	C							
ATOM	0	CA	GLN A 273	9.614	-13.020	21.432	0.00	0.00
A	C							
ATOM	0	CA	THR A 274	8.660	-14.154	17.873	0.00	0.00
A	C							
ATOM	0	CA	ALA A 275	11.611	-13.454	15.736	0.00	0.00
A	C							
ATOM	0	CA	VAL A 276	13.533	-16.679	15.922	0.00	0.00
A	C							
ATOM	0	CA	GLN A 277	11.336	-19.629	16.496	0.00	0.00
A	C							
ATOM	0	CA	LEU A 278	10.970	-21.950	19.360	0.00	0.00
A	C							
ATOM	0	CA	ASP A 306	-4.793	-26.348	20.414	0.00	0.00
A	C							
ATOM	0	CA	THR A 308	-3.495	-22.832	16.561	0.00	0.00
A	C							
ATOM	0	CA	VAL A 309	-1.332	-25.679	15.400	0.00	0.00
A	C							
ATOM	0	CA	TYR A 335	32.027	-18.555	10.993	0.00	0.00
A	C							
ATOM	0	CA	LEU A 372	23.776	-14.742	12.937	0.00	0.00
A	C							
ATOM	0	CA	GLU A 373	24.892	-11.299	14.242	0.00	0.00
A	C							
ATOM	0	CA	SER A 376	19.587	-14.121	15.269	0.00	0.00
A	C							

ATOM	0	CA	ALA A 377	21.364	-12.481	18.109	0.00	0.00
A	C							
ATOM	0	CA	LEU A 380	17.049	-12.894	20.827	0.00	0.00
A	C							
ATOM	0	CA	PHE A 381	18.126	-9.780	22.708	0.00	0.00
A	C							
ATOM	0	CA	PHE A 388	15.472	-12.482	30.704	0.00	0.00
A	C							
ATOM	0	CA	LEU A 401	26.491	-19.448	32.271	0.00	0.00
A	C							
ATOM	0	CA	GLY A 402	29.874	-17.666	32.101	0.00	0.00
A	C							
ATOM	0	CA	ASP A 403	29.566	-13.889	31.936	0.00	0.00
A	C							
ATOM	0	CA	LEU A 404	26.384	-13.628	29.944	0.00	0.00
A	C							
ATOM	0	CA	ILE A 407	27.407	-8.419	30.439	0.00	0.00
A	C							
ATOM	0	CA	ARG A 423	32.279	-10.875	16.349	0.00	0.00
A	C							
ATOM	0	CA	LEU A 429	32.511	-9.785	19.879	0.00	0.00
A	C							
ATOM	0	CA	PHE A 432	27.625	-9.601	20.031	0.00	0.00
A	C							
ATOM	0	CA	TRP A 445	9.219	-4.401	13.406	0.00	0.00
A	C							
ATOM	0	CA	GLN A 448	10.948	-9.045	11.948	0.00	0.00
A	C							
ATOM	0	CA	HIS A 449	10.701	-7.903	8.430	0.00	0.00
A	C							
ATOM	0	CA	ILE A 451	13.771	-11.849	8.960	0.00	0.00
A	C							
ATOM	0	CA	GLU A 452	10.849	-12.835	6.872	0.00	0.00
A	C							
ATOM	0	CA	MET A 453	12.858	-11.529	3.862	0.00	0.00
A	C							
ATOM	0	CA	ARG A 455	13.826	-16.561	4.859	0.00	0.00
A	C							
ATOM	0	CA	GLN A 456	12.307	-15.855	1.397	0.00	0.00
A	C							
ATOM	0	CA	PRO A 484	29.797	-23.817	26.313	0.00	0.00
A	C							
ATOM	0	CA	GLY A 485	29.335	-20.718	28.271	0.00	0.00
A	C							
ATOM	0	CA	ARG A 486	32.477	-19.218	26.838	0.00	0.00
A	C							
ATOM	0	CA	THR A 488	28.751	-18.591	22.179	0.00	0.00
A	C							
ATOM	0	CA	GLY A 489	29.978	-18.362	18.590	0.00	0.00
A	C							
ATOM	0	CA	ILE A 491	25.976	-22.214	18.037	0.00	0.00
A	C							
ATOM	0	CA	SER A 492	26.470	-20.646	14.610	0.00	0.00
A	C							
ATOM	0	CA	VAL A 495	20.892	-21.229	15.354	0.00	0.00
A	C							
ATOM	0	CA	CYS A 499	16.329	-18.125	12.495	0.00	0.00
A	C							

ATOM	0	CA	LEU	A	500	15.055	-19.509	9.162	0.00	0.00
A	C									
ATOM	0	CA	ASP	A	501	12.509	-21.783	10.900	0.00	0.00
A	C									
ATOM	0	CA	ILE	A	530	38.543	-15.553	22.466	0.00	0.00
A	C									
ATOM	0	CA	TRP	A	531	35.687	-17.843	21.343	0.00	0.00
A	C									
ATOM	0	CA	ILE	A	534	38.136	-21.645	24.267	0.00	0.00
A	C									
ATOM	0	CA	TRP	A	556	6.679	-28.127	7.026	0.00	0.00
A	C									

SEQ.ID. NO 1:

```
1 mvlsevwtll sgvsqvcilac silsaavvlr wtgrqkarga atrarqkqra sletmdkavq
61 rfrlqnpdld sealltlp1l qlvqklqsge lspeavffty lgkaweavng tncvtsyltd
121 cetqlsqapr qglllygvpvs lkecfisykgh dstlglslne gmpsesdcvv vqvlklggav
181 pfvhtnvpqs mlsfdcsnpl fgqtmnpwks skapggsggg egaligsqgs plgltdigg
241 sirfpaafcg icglkptgnr lsksglkqcv yqqtavqlsl gpmardvesl alcliallce
301 hlftldptvp plpfreevyr ssrplrvgyy etdnytmpsp amrralietk qrleaaghtl
361 ipflpnnipy alevlsaggl fsdggrsflq nfkqdfvdpq lgdlililrl pswfkrllsl
421 llkplfpzrla aflnsmrpre aeklwklqhe iemyrqsvia qwkamldvl ltpnlgpald
481 lntpgzatga isytlvyncl dfpagvvpvt tvtaeddaqm elykgyfgdi wdiilkamk
541 nsvglpvavq cvalpwqeel clrfmreveq lntpqkqps
```

References

1. W. A. Devane *et al.*, *Science* **258**, 1946 (1992).
2. B. F. Cravatt *et al.*, *Science* **268**, 1506 (1995).
3. A. Calignano, G. La Rana, A. Guiffrida, D. Piomelli, *Nature* **394**, 277 (1998).
4. F. Rodriguez de Fonseca *et al.*, *Nature* **414**, 209 (2001).
5. A. H. Lichtman, E. G. Hawkins, G. Griffin, B. F. Cravatt, *J. Pharmacol. Exp. Ther.* **302**, 73 (2002).
6. R. I. Wilson, R. A. Nicoll, *Science* **296**, 678 (2002).
7. N. Fukushima, I. Ishii, J. J. Contos, J. A. Weiner, J. Chun, *Annu. Rev. Pharmacol. Toxicol.* **41**, 507 (2001).
8. B. F. Cravatt *et al.*, *Nature* **384**, 83 (1996).
9. B. F. Cravatt *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **98**, 9371 (2001).
10. M. P. Patricelli, M. A. Lovato, B. F. Cravatt, *Biochemistry* **38**, 9804 (1999).
11. K. Gomi, K. Kitamoto, C. Kumagai, *Gene* **108**, 91 (1991).
12. S. Shin *et al.*, *EMBO J.* **21**, 2509 (2002).
13. A. W. Curnow *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 11819 (1997).
14. M. P. Patricelli, B. F. Cravatt, *Biochemistry* **40**, 6107 (2001).
15. M. P. Patricelli, H. A. Lashuel, D. K. Giang, J. W. Kelly, B. F. Cravatt, *Biochemistry* **37**, 15177 (1998).
16. Materials and methods are available as supporting online material on *Science Online*.
17. Buried van der Waals molecular surface area was calculated in GRASP (23).
18. M. P. Patricelli, B. F. Cravatt, *Biochemistry* **38**, 14125 (1999).
19. K. U. Wendt, K. Poralla, G. E. Schultz, *Science* **277**, 1811 (1997).
20. D. Picot, P. J. Loll, R. M. Garavito, *Nature* **367**, 243 (1994).

21. M. Egertova, D. K. Giang, B. F. Cravatt, M. R. Elphick, *Proc. R. Soc. Lond. B. Biol. Sci.* **265**, 2081 (1998).
22. D. L. Boger et al., *Proc. Natl. Acad. Sci. U.S.A.* **97**, 5044 (2000).
23. A. Nicholls, K. Sharp, B. Honig, *Proteins Struct. Funct. Genet.* **11**, 282 (1991).
24. Collaborative Computational Project, Number 4, *Acta Crystallogr.* **D50**, 760 (1994).
25. J. C. Sipe, K. Chiang, A. L. Gerber, E. Beutler, B. F. Cravatt, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 8394 (2002).
26. P. J. Kraulis, *J. Appl. Crystallogr.* **24**, 946 (1991).
27. E. A. Merritt, D. J. Bacon, *Methods Enzymol.* **277**, 505 (1997).
28. D. P. Tieleman, H. J. C. Berendsen, *Biophys. J.* **74**, 2786 (1998),
<http://indigo1.biop.ox.ac.uk/tieleman/downloads.html>, pope.pdb.
29. A, alanine; F, phenylalanine; L, leucine; I, isoleucine; P, proline; W, tryptophan.

While the invention has been described in detail with reference to certain preferred embodiments thereof, it will be understood that modifications and variations are within the spirit and scope of that which is described and claimed.

WHAT IS CLAIMED IS:

1. A crystallized mammalian fatty acid amide hydrolase (FAAH).
2. The crystallized FAAH of claim 1, wherein said enzyme has an amino acid sequence of SEQ. ID. NO.1, or conservative substitutions thereof.
3. A three-dimensional model for the structure of FAAH at the secondary, tertiary, and quaternary levels.
4. A method for determining the three-dimensional structure of FAAH from the crystals of claim 1, comprising collecting X-ray diffraction data from said crystal, and analyzing said data by multiple isomorphous replacement (MIR) and single-and multiwavelength anomalous diffraction (SAD/MAD), thereby determining the three-dimensional structure of FAAH.
5. A method for determining the molecular structure of a molecule or molecular complex whose structure is unknown, comprising
 - (a) obtaining crystals of the molecule or molecular complex whose structure is unknown;
 - (b) generating X-ray diffraction data from the crystallized molecule or molecular complex;
 - (c) comparing the X-ray diffraction data from the molecule or molecular complex with the three dimensional structure determined from the crystal of claim 1; and
 - (d) using molecular replacement analysis to conform the three dimensional structure determined from the crystal of claim 1 to the X-ray diffraction data from the crystallized molecule or molecular complex.
6. A method for identifying an agent that interacts with an internal channel of FAAH, comprising performing manual or computer assisted fitting analysis of the agent with the three-dimensional model of claim 3, thereby identifying an agent that interacts with an internal channel of FAAH.
7. The method of claim 6, wherein the internal channel is the active site.

8. The method of claim 7, wherein the active site comprises amino acid residues 142, 217, 218, 236, 238, 239, 240 and 241.
9. The method of claim 6, wherein the internal channel is the substrate binding pocket.
10. The method of claim 9, wherein the substrate binding pocket comprises amino acid residues 491, 495, 335, 372, 377, 194, and 244.
11. The method of claim 9, wherein the interaction of the agent with the substrate binding pocket is identified, optimized, or compared to the position of arachidonyl inhibitor MAFP.
12. The method of claim 6, wherein the internal channel is the membrane port of FAAH.
13. The method of claim 12, wherein the membrane port comprises amino acid residues 403, 407, 428, 429, 486, 530, and 534.
14. The method of claim 6, wherein the internal channel is the cytosolic port of FAAH.
15. The method of claim 14, wherein the cytosolic port comprises amino acid residues 266, 271, 272, 445, 449, and 453.
16. The method of claim 6, wherein the internal channel is the dimerization tunnel of FAAH.
17. The method of claim 16, wherein the dimerization tunnel comprises amino acid residues 257, 259, 261, 262, 263, 264, 306, 308, 309, 448, 451, 452, 455, 456, 499, 500, 501, and 556.

18. The method of claim 6, wherein the internal channel is the membrane-binding domain of FAAH.
19. The method of claim 18, wherein the membrane-binding domain comprises amino acid residues 383-440.
20. The method of claim 6, wherein the internal channel is the head group tunnel of FAAH.
21. The method of claim 20, wherein the head group tunnel comprises amino acid residues 268, 269, 270, 273, 274, 275, 276, 277, and 278.
22. The method of claim 6, wherein the internal channel is the alkyl tunnel of FAAH.
23. The method of claim 22, wherein the alkyl tunnel comprises amino acid residues 190-194, 244, 335, 372, 373, 376, 377, 380, 381, 388, 401, 402, 404, 432, 484, 485, 488, 489, 491, 492, 495, and 531.
24. A method for identifying an agent that interacts with the SH3-binding domain of FAAH, comprising performing manual or computer assisted fitting analysis of the agent with the three-dimensional model of claim 3, thereby identifying an agent that interacts with the SH3-binding domain of FAAH.
25. The method of claim 24, wherein the SH3-binding domain comprises amino acid residues 297-315.
26. A method for identifying an agent that interacts with the surface helix-loop-helix of FAAH, comprising performing manual or computer assisted fitting analysis of the agent with the three-dimensional model of claim 3, thereby identifying an agent that interacts with the surface helix-loop-helix of FAAH.

2003290780 14 Apr 2009

WO 2004/044169

PCT/US2003/036125

31

27. The method of claim 26, wherein the surface helix-loop-helix comprises amino acid residues 511-546.




FIG. 1B

FIG. 1A

SUBSTITUTE SHEET (RULE 26)




FIG. 2B




FIG. 2A

SUBSTITUTE SHEET (RULE 26)





FIG. 3

SUBSTITUTE SHEET (RULE 26)

4 / 5



404-LILILxLPxWFxxLLxLLxPLFPxLAAFL-433

FIG. 4A




FIG. 4B

SUBSTITUTE SHEET (RULE 26)




FIG. 5

SUBSTITUTE SHEET (RULE 26)

SEQUENCE LISTING

<110> THE SCRIPPS RESEARCH INSTITUTE
 BRACEY, Michael H.
 HANSON, Michael A.
 STEVENS, Raymond C.
 CRAVATT, Benjamin F.

<120> CRYSTALLINE FORM OF FATTY ACID AMIDE HYDROLASE (FAAH)

<130> SCRIP1590WO

<150> US 60/426,788
 <151> 2002-11-14

<160> 1

<170> PatentIn version 3.1

<210> 1
 <211> 579
 <212> PRT
 <213> Rat

<400> 1

Met Val Leu Ser Glu Val Trp Thr Thr Leu Ser Gly Val Ser Gly Val
 1 5 10 15

Cys Leu Ala Cys Ser Leu Leu Ser Ala Ala Val Val Leu Arg Trp Thr
 20 25 30

Gly Arg Gln Lys Ala Arg Gly Ala Ala Thr Arg Ala Arg Gln Lys Gln
 35 40 45

Arg Ala Ser Leu Glu Thr Met Asp Lys Ala Val Gln Arg Phe Arg Leu
 50 55 60

Gln Asn Pro Asp Leu Asp Ser Glu Ala Leu Leu Thr Leu Pro Leu Leu
 65 70 75 80

Gln Leu Val Gln Lys Leu Gln Ser Gly Glu Leu Ser Pro Glu Ala Val
 85 90 95

Phe Phe Thr Tyr Leu Gly Lys Ala Trp Glu Val Asn Lys Gly Thr Asn
 100 105 110

Cys Val Thr Ser Tyr Leu Thr Asp Cys Glu Thr Gln Leu Ser Gln Ala
 115 120 125

Pro Arg Gln Gly Leu Leu Tyr Gly Val Pro Val Ser Leu Lys Glu Cys
 130 135 140

I
 145 ly His Asp Ser Thr Leu Gly Leu Ser Leu Asn Glu
 150 155 160

Gly Met Pro Ser Glu Ser Asp Cys Val Val Val Gln Val Leu Lys Leu
 165 170 175

Gln Gly Ala Val Pro Phe Val His Thr Asn Val Pro Gln Ser Met Leu
 180 185 190

Ser Phe Asp Cys Ser Asn Pro Leu Phe Gly Gln Thr Met Asn Pro Trp
 195 200 205

Lys Ser Ser Lys Ser Pro Gly Gly Ser Ser Gly Gly Glu Gly Ala Leu
 210 215 220

Ile Gly Ser Gly Gly Ser Pro Leu Gly Leu Gly Thr Asp Ile Gly Gly
 225 230 235 240

Ser Ile Arg Phe Pro Ser Ala Phe Cys Gly Ile Cys Gly Leu Lys Pro
 245 250 255

Thr Gly Asn Arg Leu Ser Lys Ser Gly Leu Lys Gly Cys Val Tyr Gly
 260 265 270

Gln Thr Ala Val Gln Leu Ser Leu Gly Pro Met Ala Arg Asp Val Glu
 275 280 285

Ser Leu Ala Leu Cys Leu Lys Ala Leu Leu Cys Glu His Leu Phe Thr
 290 295 300

Leu Asp Pro Thr Val Pro Pro Leu Pro Phe Arg Glu Glu Val Tyr Arg
 305 310 315 320

Ser Ser Arg Pro Leu Arg Val Gly Tyr Tyr Glu Thr Asp Asn Tyr Thr
 325 330 335

Met Pro Ser Pro Ala Met Arg Arg Ala Leu Ile Glu Thr Lys Gln Arg
 340 345 350

Leu Glu Ala Ala Gly His Thr Leu Ile Pro Phe Leu Pro Asn Asn Ile
 355 360 365

Pro Tyr Ala Leu Glu Val Leu Ser Ala Gly Gly Leu Phe Ser Asp Gly
 370 375 380

Gly Arg Ser Phe Leu Gln Asn Phe Lys Gly Asp Phe Val Asp Pro Cys

3 390 395 400

Leu Gly Asp Leu Ile Leu Ile Leu Arg Leu Pro Ser Trp Phe Lys Arg
 405 410 415

Leu Leu Ser Leu Leu Leu Lys Pro Leu Phe Pro Arg Leu Ala Ala Phe
 420 425 430

Leu Asn Ser Met Arg Pro Arg Ser Ala Glu Lys Leu Trp Lys Leu Gln
 435 440 445

His Glu Ile Glu Met Tyr Arg Gln Ser Val Ile Ala Gln Trp Lys Ala
 450 455 460

Met Asn Leu Asp Val Leu Leu Thr Pro Met Leu Gly Pro Ala Leu Asp
 465 470 475 480

Leu Asn Thr Pro Gly Arg Ala Thr Gly Ala Ile Ser Tyr Thr Val Leu
 485 490 495

Tyr Asn Cys Leu Asp Phe Pro Ala Gly Val Val Pro Val Thr Thr Val
 500 505 510

Thr Ala Glu Asp Asp Ala Gln Met Glu Leu Tyr Lys Gly Tyr Phe Gly
 515 520 525

Asp Ile Trp Asp Ile Ile Leu Lys Lys Ala Met Lys Asn Ser Val Gly
 530 535 540

Leu Pro Val Ala Val Gln Cys Val Ala Leu Pro Trp Gln Glu Glu Leu
 545 550 555 560

Cys Leu Arg Phe Met Arg Glu Val Glu Gln Leu Met Thr Pro Gln Lys
 565 570 575

Gln Pro Ser