

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 January 2004 (22.01.2004)

PCT

(10) International Publication Number
WO 2004/008099 A2


- (51) International Patent Classification⁷: G01N
- (21) International Application Number: PCT/US2003/021590
- (22) International Filing Date: 11 July 2003 (11.07.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:

60/396,290	15 July 2002 (15.07.2002)	US
60/480,043	20 June 2003 (20.06.2003)	US
- (71) Applicant (for all designated States except US): GENETECH, INC. [US/US]; 1 DNA Way, South San Francisco, CA 94080 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): KOLL, Hans [DE/DE]; Blütenanger 45, 85247 Oberroth (DE). BOSENMAIER, Birgit [DE/DE]; Horst-Wolfram-Geissler-Weg 10, 82229 Seefeld (DE). MÜLLER, Hans-Joachim [DE/DE]; Ruhe am Bach 5 b, 82377 Penzberg (DE). SLIWKOWSKI, Mark, X. [US/US]; 42 Oak Creek Lane, San Carlos, CA 94070 (US). KELSEY, Stephen, Michael [GB/US]; 360 14th Street, Montara, CA 94037 (US).
- (74) Agent: DREGER, Ginger, R.; Heller, Ehrman, White & McAuliffe LLP, 275 Middlefield Road, Menlo Park, CA 94025 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ (utility model), DE (utility model), DK (utility model), DK, DM, DZ, EC, EE (utility model), EE, ES, FI (utility model), FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK (utility model), SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
 — without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR IDENTIFYING TUMORS THAT ARE RESPONSIVE TO TREATMENT WITH ANTI-erbB2 ANTIBODIES



(57) Abstract: Tumors are identified as responsive to treatment with anti-HER2 antibodies by detecting the presence of a HER2/HER3 and/or HER2/HER1 protein complex or for HER2 phosphorylation in a sample of tumor cells. Patients suffering from a tumor comprising HER2/HER1 and/or HER2/HER3 heterodimers and/or HER2 phosphorylation are treated with anti-HER2 antibodies, such as rhuMAb 2C4.

WO 2004/008099 A2

METHODS FOR IDENTIFYING TUMORS THAT ARE RESPONSIVE TO TREATMENT WITH ANTI-ErbB2 ANTIBODIES

Field of the Invention

The present invention relates to methods of identifying a tumor as responsive to
5 treatment with anti-HER2 antibodies, as well as methods of treating patients suffering
from such tumors.

Background of the Invention

The ErbB family of receptor tyrosine kinases are important mediators of cell
growth, differentiation and survival. The receptor family includes four distinct members
10 including epidermal growth factor receptor (EGFR or ErbB1), HER2 (ErbB2 or p185^{neu}),
HER3 (ErbB3) and HER4 (ErbB4 or tyro2).

EGFR, encoded by the erbB1 gene, has been causally implicated in human
malignancy. In particular, increased expression of EGFR has been observed in breast,
bladder, lung, head, neck and stomach cancer as well as glioblastomas. Increased EGFR
15 receptor expression is often associated with increased production of the EGFR ligand,
transforming growth factor alpha (TGF- α), by the same tumor cells resulting in receptor
activation by an autocrine stimulatory pathway. Baselga and Mendelsohn *Pharmac. Ther.*,
64:127-154 (1994). In addition, an epidermal growth factor receptor related protein
(ERRP) wherein a cDNA fragment clone of 1583 base pairs with 90-95% sequence
20 homology to mouse EGFR and a truncated rat EGFR has been described (US Patent No.
6,399,743; and US Publication No. 2003/0096373). Monoclonal antibodies directed
against the EGFR or its ligands, TGF- α and EGF, have been evaluated as therapeutic
agents in the treatment of such malignancies. See, e.g., Baselga and Mendelsohn., *supra*;
Masui et al., *Cancer Research*, 44:1002-1007 (1984); and Wu et al., *J. Clin. Invest.*,
25 95:1897-1905 (1995).

The second member of the ErbB family, p185neu, was originally identified as the
product of the transforming gene from neuroblastomas of chemically treated rats. The
activated form of the neu proto-oncogene results from a point mutation (valine to glutamic
acid) in the transmembrane region of the encoded protein. Amplification of the human
30 homolog of neu (HER2) is observed in breast and ovarian cancers and correlates with a
poor prognosis (Slamon et al., *Science*, 235:177-182 (1987); Slamon et al., *Science*,
244:707-712 (1989); and US Patent No. 4,968,603). To date, no point mutation analogous
to that in the neu proto-oncogene has been reported for human tumors. Overexpression of
ErbB2 (frequently but not uniformly due to gene amplification) has also been observed in

other carcinomas including carcinomas of the stomach, endometrium, salivary gland, lung, kidney, colon, thyroid, pancreas and bladder. See, among others, King et al., *Science*, 229:974 (1985); Yokota et al., *Lancet*, 1:765-767 (1986); Fukushigi et al., *Mol Cell Biol.*, 6:955-958 (1986); Geurin et al., *Oncogene Res.*, 3:21-31 (1988); Cohen et al., *Oncogene*, 4:81-88 (1989); Yonemura et al., *Cancer Res.*, 51:1034 (1991); Borst et al., *Gynecol. Oncol.*, 38:364 (1990); Weiner et al., *Cancer Res.*, 50:421-425 (1990); Kern et al., *Cancer Res.*, 50:5184 (1990); Park et al., *Cancer Res.*, 49:6605 (1989); Zhau et al., *Mol. Carcinog.*, 3:354-357 (1990); Aasland et al., *Br. J. Cancer*, 57:358-363 (1988); Williams et al., *Pathobiology*, 59:46-52 (1991); and McCann et al., *Cancer*, 65:88-92 (1990).
10 ErbB2 may be overexpressed in prostate cancer (Gu et al., *Cancer Lett.*, 99:185-9 (1996); Ross et al., *Hum. Pathol.*, 28:827-33 (1997); Ross et al., *Cancer*, 79:2162-70 (1997); and Sadasivan et al., *J. Urol.*, 150:126-31 (1993)). Overexpression of ErbB2 may lead to tumor growth via ligand-independent activation of ErbB2 or ErbB2 homodimers.

Antibodies directed against the rat p185neu and human ErbB2 protein products
15 have been described. Drebin and colleagues have raised antibodies against the rat neu gene product, p185neu. See, for example, Drebin et al., *Cell*, 41:695-706 (1985); Myers et al., *Meth. Enzym.*, 198:277-290 (1991); and WO94/22478. Drebin et al., *Oncogene*, 2:273-277 (1988) report that mixtures of antibodies reactive with two distinct regions of p185neu result in synergistic anti-tumor effects on neu-transformed NIH-3T3 cells implanted into
20 nude mice. See also U.S. Patent 5,824,311 issued October 20, 1998.

Hudziak et al., *Mol. Cell. Biol.*, 9(3):1165-1172 (1989) describe the generation of a panel of anti-ErbB2 antibodies which were characterized using the human breast tumor cell line SK-BR-3. Relative cell proliferation of the SK-BR-3 cells following exposure to the antibodies was determined by crystal violet staining of the monolayers
25 after 72 hours. Using this assay, maximum inhibition was obtained with the antibody called 4D5 which inhibited cellular proliferation by 56%. Other antibodies in the panel reduced cellular proliferation to a lesser extent in this assay. The antibody 4D5 was further found to sensitize ErbB2-overexpressing breast tumor cell lines to the cytotoxic effects of TNF- α . See also U.S. Patent No. 5,677,171 issued October 14, 1997. The anti-
30 ErbB2 antibodies discussed in Hudziak et al. are further characterized in Fendly et al., *Cancer Research*, 50:1550-1558 (1990); Kotts et al., *In Vitro*, 26(3):59A (1990); Sarup et al., *Growth Regulation*, 1:72-82 (1991); Shepard et al., *J. Clin. Immunol.*, 11(3):117-127 (1991); Kumar et al., *Mol. Cell. Biol.*, 11(2):979-986 (1991); Lewis et al., *Cancer Immunol. Immunother.*, 37:255-263 (1993); Pietras et al., *Oncogene*, 9:1829-1838 (1994);
35 Vitetta et al., *Cancer Research*, 54:5301-5309 (1994); Sliwkowski et al., *J. Biol. Chem.*,

269(20):14661-14665 (1994); Scott et al., *J. Biol. Chem.*, 266:14300-5 (1991); D'souza et al., *Proc. Natl. Acad. Sci.*, 91:7202-7206 (1994); Lewis et al., *Cancer Research*, 56:1457-1465 (1996); and Schaefer et al., *Oncogene*, 15:1385-1394 (1997).

A recombinant humanized version of the murine anti-ErbB2 antibody 4D5
5 (huMAb4D5-8, rhuMAb HER2 or HERCEPTIN[®]; U.S. Patent No. 5,821,337) is clinically active in patients with ErbB2-overexpressing metastatic breast cancers that have received extensive prior anti-cancer therapy (Baselga et al., *J. Clin. Oncol.*, 14:737-744 (1996)). HERCEPTIN[®] received marketing approval from the Food and Drug Administration
September 25, 1998 for the treatment of patients with metastatic breast cancer whose
10 tumors overexpress the ErbB2 protein. However, not all ErbB2 overexpressing tumors respond to HERCEPTIN[®]. (Brockhoff et al., *Cytometry*, 44:338-48 (2001)). In addition, preclinical data suggest that HERCEPTIN[®] may be therapeutically effective in treating non-small cell lung cancer (NSCLC). HER2 protein is overexpressed in 20-66% of resected NSCLC tumors and has been shown to predict poor patient outcome in multiple
15 series (Azzoli, C.G. et al., *Semin. Oncol.*, 29(Suppl 4):59-65 (2002)).

Other anti-ErbB2 antibodies with various properties have been described in Tagliabue et al., *Int. J. Cancer*, 47:933-937 (1991); McKenzie et al., *Oncogene*, 4:543-548 (1989); Maier et al., *Cancer Res.*, 51:5361-5369 (1991); Bacus et al., *Molecular Carcinogenesis*, 3:350-362 (1990); Stancovski et al., *PNAS (USA)*, 88:8691-8695 (1991);
20 Bacus et al., *Cancer Research*, 52:2580-2589 (1992); Xu et al., *Int. J. Cancer*, 53:401-408 (1993); WO94/00136; Kasprzyk et al., *Cancer Research*, 52:2771-2776 (1992); Hancock et al., *Cancer Res.*, 51:4575-4580 (1991); Shawver et al., *Cancer Res.*, 54:1367-1373 (1994); Arteaga et al., *Cancer Res.*, 54:3758-3765 (1994); Harwerth et al., *J. Biol. Chem.*, 267:15160-15167 (1992); U.S. Patent No. 5,783,186; and Klapper et al., *Oncogene*,
25 14:2099-2109 (1997). Monoclonal antibody 2C4 is described in WO 01/00245, which is hereby incorporated by reference. 2C4 has been shown to disrupt dimerization of HER2 with other ErbB receptor family members (WO 01/00245).

Homology screening has resulted in the identification of two other ErbB receptor family members; ErbB3 (U.S. Patent Nos. 5,183,884 and 5,480,968 as well as Kraus et al.,
30 *PNAS (USA)*, 86:9193-9197 (1989)) and ErbB4 (EP Pat Appln No 599,274; Plowman et al., *Proc. Natl. Acad. Sci. USA*, 90:1746-1750 (1993); and Plowman et al., *Nature*, 366:473-475 (1993)). Both of these receptors display increased expression on at least some breast cancer cell lines.

The ErbB receptors are generally found in various combinations in cells and
5 heterodimerization is thought to increase the diversity of cellular responses to a variety of

ErbB ligands (Earp et al., *Breast Cancer Research and Treatment*, 35:115-132 (1995)).

However, the mechanism by which these receptors aggregate and how this contributes to signaling is not fully understood (Brennan, P.J. et al., *Oncogene*, 19:6093-6101 (2000)).

EGFR is bound by six different ligands; epidermal growth factor (EGF), transforming growth factor alpha (TGF- α), amphiregulin, heparin binding epidermal growth factor (HB-EGF), betacellulin and epiregulin (Groenen et al., *Growth Factors*, 11:235-257 (1994)). A

family of heregulin proteins resulting from alternative splicing of a single gene are ligands for ErbB3 and ErbB4. The heregulin family includes alpha, beta and gamma heregulins

(Holmes et al., *Science*, 256:1205-1210 (1992); U.S. Patent No. 5,641,869; and Schaefer

et al., *Oncogene*, 15:1385-1394 (1997)); neu differentiation factors (NDFs), glial growth

factors (GGFs); acetylcholine receptor inducing activity (ARIA); and sensory and motor

neuron derived factor (SMDF). For a review, see Groenen et al., *Growth Factors*, 11:235-

257 (1994); Lemke, G., *Molec. & Cell. Neurosci.*, 7:247-262 (1996) and Lee et al., *Pharm.*

Rev., 47:51-85 (1995). Recently three additional ErbB ligands were identified;

neuregulin-2 (NRG-2) which is reported to bind either ErbB3 or ErbB4 (Chang et al.,

Nature, 387:509-512 (1997); and Carraway et al., *Nature*, 387:512-516 (1997));

neuregulin-3 which binds ErbB4 (Zhang et al., *PNAS (USA)*, 94(18):9562-7 (1997)); and

neuregulin-4 which binds ErbB4 (Harari et al., *Oncogene*, 18:2681-89 (1999)) HB-EGF,

betacellulin and epiregulin also bind to ErbB4.

While EGF and TGF α do not bind ErbB2, EGF stimulates EGFR and ErbB2 to

form a heterodimer, which activates EGFR and results in transphosphorylation of ErbB2

in the heterodimer. Dimerization and/or transphosphorylation appears to activate the

ErbB2 tyrosine kinase. See Earp et al., supra. Likewise, heregulin does not bind ErbB2,

but when ErbB3 is co-expressed with ErbB2, an active signaling complex is

formed (Nagy et al., *Cytometry*, 32:120-31 (1998). Antibodies directed against ErbB2 are

capable of disrupting this complex (Sliwkowski et al., *J. Biol. Chem.*, 269(20):14661-

14665 (1994)). ErbB3 is tyrosine kinase defective and thus requires heterodimerization,

preferably with ErbB2, for signal transduction capabilities. (Graus-Porta et al., *EMBO J.*,

16:1647-55 (1995)). Additionally, the affinity of ErbB3 for heregulin (HRG) is increased

to a higher affinity state when co-expressed with ErbB2. See also, Levi et al., *Journal of*

Neuroscience, 15:1329-1340 (1995); Morrissey et al., *Proc. Natl. Acad. Sci. USA*,

92:1431-1435 (1995); and Lewis et al., *Cancer Res.*, 56:1457-1465 (1996) with respect to

the ErbB2-ErbB3 protein complex. Indeed, ErbB2 is the preferred heterodimerization

partner for both EGFR and ErbB3. (Graus-Porta et al., supra). ErbB4, like ErbB3, forms

an active signaling complex with ErbB2 (Carraway and Cantley, *Cell*, 78:5-8 (1994)).

Ligand-dependent heterodimerization of ErbB2 with EGFR or ErbB3 may promote the growth of tumors that express ErbB2.

The expression of the ErbB receptors and heregulin and the phosphorylation status of HER2 has been investigated in tumor specimens from primary breast cancer patients and in urinary bladder carcinoma (Esteva et al., *Pathol. Oncol. Res.*, 7:171-177 (2001); Chow et al., *Clin. Cancer Res.*, 7:1957-1962 (2001)). Correlation between active signaling through Her2/neu and clinicopathology and patient outcome in breast cancer has been reported by Thor et al., *J. Clin. Oncology*, 18:3230-3239 (2000), and DiGiovanna et al., *Cancer Res.*, 62:6667-6673 (2002).

Summary of the Invention

In one aspect, the present invention relates to a method of identifying a tumor that is responsive to treatment with an anti-HER2 antibody. Preferably the anti-HER2 antibody blocks ligand activation of an ErbB heterodimer comprising HER2. In one embodiment the antibody is monoclonal antibody 2C4, more preferably rhuMAb 2C4.

A sample of the tumor is obtained and the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein complex is detected in the sample. A tumor is identified as responsive to treatment with the anti-HER2 antibody when a complex is detected.

In one embodiment, the presence of a complex is detected by immunoprecipitating any protein complexes that comprise HER2 with an anti-HER2 antibody. The immunoprecipitated complexes are then contacted with an antibody selected from the group consisting of anti-HER3 antibodies, anti-HER1 antibodies, and anti-HER4 antibodies, and any binding is determined. A HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 complex is detected if it is determined that anti-HER3 and/or anti-HER1 and/or anti-HER4 antibodies bind to the immunoprecipitated complexes.

In another embodiment, the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein complex is detected by contacting the tumor sample with an anti-HER2 antibody that comprises a first fluorophore. The tumor sample is then contacted with an antibody selected from the group consisting of anti-HER3 and/or anti-HER1 and/or anti-HER4 antibodies, wherein said antibody comprises a second fluorophore. Measurements of fluorescence resonance energy transfer are then made to determine if the first fluorophore and the second fluorophore are in close proximity. The presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein complex is detected if the first and second fluorophore are determined to be in close proximity.

In yet another embodiment, the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 complex is detected by contacting the tumor sample with a first binding compound. The first binding compound comprises a first target binding moiety that specifically binds HER2. The first target binding moiety is preferably an anti-HER2 antibody or antibody fragment. The first binding compound further comprises a detectable moiety that is linked to the first binding domain by a cleavable linker.

The tumor sample is contacted with a second binding compound. The second binding compound preferably comprises a second target binding moiety that specifically binds HER3 or HER1 or HER4 and preferably does not bind HER2. In another embodiment, the second binding compound binds HER3 or HER1, and does not bind HER2 or HER4. In further embodiment, the second target binding moiety comprises an anti-HER3 or anti-HER1 or anti-HER4 antibody or antibody fragment. The second binding compound is capable of cleaving the cleavable linker in the first binding compound upon activation, thus producing free detectable moiety if the first binding compound and the second binding compound are in close proximity. The presence of a HER2/HER3 or HER2/HER1 or HER2/HER4 protein complex is detected when the presence of the free detectable moiety is identified. In one embodiment, the presence of the free detectable moiety is identified by capillary electrophoresis.

In another embodiment, the first binding compound comprises a first target binding domain that specifically binds HER1 or HER3 or HER4, and the second binding compound comprises a second target binding domain that specifically binds HER2.

In yet another embodiment, the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein complex, and resultant HER2 activation, is detected by assessing phosphorylation of ErbB receptor, for example by immunoprecipitation of the HER2 protein followed by Western blot immunodetection of phosphotyrosine.

The tumor sample that is analyzed for the presence of HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein complexes is preferably obtained from a patient that is suffering from the tumor. The sample can, for example, be obtained by biopsy. In another embodiment the sample is obtained by purifying circulating tumor cells from the patient. In yet another embodiment, the sample is obtained during surgery to remove the tumor from the patient.

In another embodiment, the sample of the tumor is obtained from a mammal other than the patient that originally developed the tumor. Preferably, the sample is obtained from a mouse, or another rodent. More preferably, the tumor is a xenografted tumor. The

xenografted tumor is preferably produced by transplanting a fragment of a human tumor into a mouse, or another rodent.

In one embodiment, the tumor is a lung tumor, more preferably a non-small cell lung cancer tumor. In another embodiment, the tumor is a mammary tumor.

5 In another aspect, the invention concerns a method for identifying tumor cells as responsive to treatment with an antibody inhibiting the association of HER2 with another member of the ErbB receptor family, comprising the steps of (a) providing a biological sample comprising HER2-positive tumor cells; and (b) detecting the phosphorylation of an ErbB receptor in the biological sample, wherein said phosphorylation indicates that the
0 tumor cells are responsive to treatment with the antibody. In one embodiment, the phosphorylation of an ErbB2 (HER2) receptor is detected.

Just as before, the other member associated with HER2 is HER3, HER1, and/or HER4, such as HER2 and/or HER1. The method can additionally comprise a step of detecting the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein
.5 complex, essentially as described above.

In another aspect, the invention further concerns a method for predicting the response of a subject diagnosed with a HER2-positive tumor to treatment with an antibody inhibiting the association of HER2 with another member of the ErbB receptor family, by detecting the formation of HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein
20 complexes and/or the phosphorylation of an ErbB receptor in a biological sample obtained from the subject, comprising HER2-positive tumor cells. The presence of such protein complexes and/or said phosphorylation indicates that the subject is likely to respond to treatment with the antibody. In one embodiment, the detection of the phosphorylation of ErbB2 (HER2) receptor indicates that the subject is likely to respond to treatment with the
25 antibody.

In yet another embodiment, the invention concerns a method for identifying a subject responsive to treatment with an anti-HER2 antibody, by detecting phosphorylation of an ErbB receptor in circulating tumor cells of the subject. The presence of such phosphorylation indicates that the subject is likely to respond to treatment with an anti-
30 HER2 antibody. In one embodiment, the ErbB2 (HER2) phosphorylation is detected. In another embodiment, the subject is a human. In yet another embodiment, the method further comprises treating the subject with an anti-HER2 antibody, preferably rhuMAb 2C4.

In another aspect, the invention provides an article of manufacture comprising a
35 container comprising an antibody which binds HER2 and instructions for administering

the antibody to a patient suffering from a tumor. Preferably, the tumor has been determined to comprise HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 heterodimers.

5 In one embodiment, the container comprises an antibody that blocks ligand activation of an ErbB heterodimer comprising HER2. In another embodiment, the container comprises monoclonal antibody 2C4, more preferably rhuMAb 2C4.

10 In a further aspect, the invention provides a method of treating a patient comprising administering to the patient a therapeutically effective amount of an antibody which binds HER2. Preferably, the patient is suffering from a tumor which has been determined to comprise HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 heterodimers.

In one embodiment, the antibody blocks the ligand activation of an ErbB heterodimer comprising HER2. In another embodiment, the antibody is monoclonal antibody 2C4, more preferably rhuMAb 2C4.

15 In another aspect, the invention provides a method of treating a patient comprising administering to the patient a therapeutically effective amount of an antibody which binds HER2. Preferably, the patient is suffering from a tumor which has been determined to have a phosphorylated ErbB receptor.

20 In one embodiment, the phosphorylated ErbB receptor is HER2. In another embodiment, the antibody blocks the ligand activation of an ErbB heterodimer comprising HER2. In yet another embodiment, the antibody is monoclonal antibody 2C3, more preferably rhuMAb 2C4.

Brief Description of the Drawings

25 Figures 1A and 1B depict alignments of the amino acid sequences of the variable light (VL) (Fig. 1A) and variable heavy (VH) (Fig. 1B) domains of murine monoclonal antibody 2C4 (SEQ ID Nos. 1 and 2, respectively); VL and VH domains of humanized 2C4 version 574 (SEQ ID Nos. 3 and 4, respectively), and human VL and VH consensus frameworks (hum κ 1, light kappa subgroup I; humIII, heavy subgroup III) (SEQ ID Nos. 5 and 6, respectively). Asterisks identify differences between humanized 2C4 version 574 and murine monoclonal antibody 2C4 or between humanized 2C4 version 574 and the human framework. Complementarity Determining Regions (CDRs) are in brackets.

30 Figures 2A and 2B show the effect of monoclonal antibody 2C4, HERCEPTIN[®] antibody or an anti-EGFR antibody on heregulin (HRG) dependent association of ErbB2

with ErbB3 in MCF7 cells expressing low/normal levels of ErbB2 (Fig. 2A) and SK-BR-3 cells expressing high levels of ErbB2 (Fig. 2B); see Example 2 below.

Figure 3 is an immunoblot showing the presence of HER1/HER2 and HER2/HER3 heterodimers in protein extracts from non-small cell lung cancer xenograft explants.

5 Figure 4 is an immunoblot showing the presence of HER2 phosphorylation in protein extracts from non-small cell lung carcinoma (NSCLC) xenograph explants.

Detailed Description of the Preferred Embodiment

The present invention is based, in part, on the experimental finding that responsiveness to the anti-HER2 antibody rhuMAb 2C4 correlates with the presence of
10 HER2/HER3 and/or HER2/HER1 and or HER2/HER4 heterodimers, and/or the phosphorylation of an ErbB receptor in tumor cells. Thus, a tumor may be identified as responsive to treatment with an anti-HER2 antibody, particularly an anti-HER2 antibody that has one or more of the biological activities of the anti-HER2 antibody 2C4, based on the presence of HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 heterodimers,
15 and/or the phosphorylation of an ErbB receptor. HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 heterodimers, and/or ErbB receptor phosphorylation may be identified by any method known in the art. By identifying specific tumors and tumor types that are responsive to treatment with anti-HER2 antibodies, patients can be identified who will likely benefit the most from such treatment. In addition, patients that would likely not
20 benefit from therapy with monoclonal antibody 2C4 can be identified.

Definitions

An "ErbB receptor" is a receptor protein tyrosine kinase which belongs to the ErbB receptor family and includes EGFR (ErbB1), ERBP, ErbB2, ErbB3 and ErbB4 receptors and other members of this family to be identified in the future. The ErbB receptor will
25 generally comprise an extracellular domain, which may bind an ErbB ligand; a lipophilic transmembrane domain; a conserved intracellular tyrosine kinase domain; and a carboxyl-terminal signaling domain harboring several tyrosine residues which can be phosphorylated. The ErbB receptor may be a "native sequence" ErbB receptor or an "amino acid sequence variant" thereof. Preferably, the ErbB receptor is native sequence
30 human ErbB receptor. Accordingly, a "member of the ErbB receptor family" is EGFR (ErbB1), ErbB2, ErbB3, ErbB4 or any other ErbB receptor currently known or to be identified in the future. Preferably, the member is EGFR (ErbB1), ErbB2, ErbB3, or ErbB4.

The terms "ErbB1", "epidermal growth factor receptor", "EGFR" and "HER1" are used interchangeably herein and refer to EGFR as disclosed, for example, in Carpenter et al., *Ann. Rev. Biochem.*, 56:881-914 (1987), including naturally occurring mutant forms thereof (e.g., a deletion mutant EGFR as in Humphrey et al., *PNAS (USA)*, 87:4207-4211 (1990)). erbB1 refers to the gene encoding the EGFR protein product. Antibodies against HER1 are described, for example, in Murthy et al., *Arch. Biochem. Biophys.*, 252:549-560 (1987) and in WO 95/25167.

The term "ERRP", "EGF-Receptor Related Protein", "EGFR Related Protein" and "epidermal growth factor receptor related protein" are used interchangeably herein and refer to ERRP as disclosed, for example in U.S. Patent No. 6,399,743 and U.S. Publication No. 2003/0096373.

The expressions "ErbB2" and "HER2" are used interchangeably herein and refer to human HER2 protein described, for example, in Semba et al., *PNAS (USA)*, 82:6497-6501 (1985) and Yamamoto et al., *Nature*, 319:230-234 (1986) (Genebank accession number X03363). The term "erbB2" refers to the gene encoding human ErbB2 and "neu" refers to the gene encoding rat p185neu. Preferred ErbB2 is native sequence human ErbB2.

"ErbB3" and "HER3" refer to the receptor polypeptide as disclosed, for example, in U.S. Patent Nos. 5,183,884 and 5,480,968 as well as Kraus et al., *PNAS (USA)*, 86:9193-9197 (1989). Antibodies against ErbB3 are known in the art and are described, for example, in U.S. Patent Nos. 5,183,884, 5,480,968 and in WO 97/35885.

The terms "ErbB4" and "HER4" herein refer to the receptor polypeptide as disclosed, for example, in EP Pat Appln No 599,274; Plowman et al., *Proc. Natl. Acad. Sci. USA*, 90:1746-1750 (1993); and Plowman et al., *Nature*, 366:473-475 (1993), including isoforms thereof, e.g., as disclosed in WO 99/19488, published April 22, 1999. Antibodies against HER4 are described, for example, in WO 02/18444.

Antibodies to ErbB receptors are available commercially from a number of sources, including, for example, Santa Cruz Biotechnology, Inc., California, USA.

By "ErbB ligand" is meant a polypeptide which binds to and/or activates an ErbB receptor. The ErbB ligand may be a native sequence human ErbB ligand such as epidermal growth factor (EGF) (Savage et al., *J. Biol. Chem.*, 247:7612-7621 (1972)); transforming growth factor alpha (TGF- α) (Marquardt et al., *Science*, 223:1079-1082 (1984)); amphiregulin also known as schwanoma or keratinocyte autocrine growth factor (Shoyab et al., *Science*, 243:1074-1076 (1989); Kimura et al., *Nature*, 348:257-260 (1990); and Cook et al., *Mol. Cell. Biol.*, 11:2547-2557 (1991)); betacellulin (Shing et al., *Science*, 259:1604-1607 (1993); and Sasada et al., *Biochem. Biophys. Res. Commun.*, 190:1173

(1993)); heparin-binding epidermal growth factor (HB-EGF) (Higashiyama et al., *Science*, 251:936-939 (1991)); epiregulin (Toyoda et al., *J. Biol. Chem.*, 270:7495-7500 (1995); and Komurasaki et al., *Oncogene*, 15:2841-2848 (1997)); a heregulin (see below); neuregulin-2 (NRG-2) (Carraway et al., *Nature*, 387:512-516 (1997)); neuregulin-3 (NRG-3) (Zhang et al., *Proc. Natl. Acad. Sci.*, 94:9562-9567 (1997)); neuregulin-4 (NRG-4) (Harari et al., *Oncogene*, 18:2681-89 (1999)) or cripto (CR-1) (Kannan et al., *J. Biol. Chem.*, 272(6):3330-3335 (1997)). ErbB ligands which bind EGFR include EGF, TGF- α , amphiregulin, betacellulin, HB-EGF and epiregulin. ErbB ligands which bind ErbB3 include heregulins. ErbB ligands capable of binding ErbB4 include betacellulin, epiregulin, HB-EGF, NRG-2, NRG-3, NRG-4 and heregulins. The ErbB ligand may also be a synthetic ErbB ligand. The synthetic ligand may be specific for a particular ErbB receptor, or may recognize particular ErbB receptor complexes. An example of a synthetic ligand is the synthetic heregulin/egf chimera biregulin (see, for example, Jones et al., *FEBS Letters*, 447:227-231 (1999), which is incorporated by reference).

“Heregulin” (HRG) when used herein refers to a polypeptide encoded by the heregulin gene product as disclosed in U.S. Patent No. 5,641,869 or Marchionni et al., *Nature*, 362:312-318 (1993). Examples of heregulins include heregulin- α , heregulin- β 1, heregulin- β 2 and heregulin- β 3 (Holmes et al., *Science*, 256:1205-1210 (1992); and U.S. Patent No. 5,641,869); neu differentiation factor (NDF) (Peles et al., *Cell*, 69: 205-216 (1992)); acetylcholine receptor-inducing activity (ARIA) (Falls et al., *Cell*, 72:801-815 (1993)); glial growth factors (GGFs) (Marchionni et al., *Nature*, 362:312-318 (1993)); sensory and motor neuron derived factor (SMDF) (Ho et al., *J. Biol. Chem.*, 270:14523-14532 (1995)); γ -heregulin (Schaefer et al., *Oncogene*, 15:1385-1394 (1997)). The term includes biologically active fragments and/or amino acid sequence variants of a native sequence HRG polypeptide, such as an EGF-like domain fragment thereof (e.g., HRG β 1177-244).

An “ErbB hetero-oligomer” herein is a noncovalently associated oligomer comprising at least two different ErbB receptors. An “ErbB dimer” is a noncovalently associated oligomer that comprises two different ErbB receptors. Such complexes may form when a cell expressing two or more ErbB receptors is exposed to an ErbB ligand. ErbB oligomers, such as ErbB dimers, can be isolated by immunoprecipitation and analyzed by SDS-PAGE as described in Sliwkowski et al., *J. Biol. Chem.*, 269(20):14661-14665 (1994), for example. Examples of such ErbB hetero-oligomers include EGFR-ErbB2 (also referred to as HER1/HER2), ErbB2-ErbB3 (HER2/HER3) and ErbB3-ErbB4 (HER3/HER4) complexes. Moreover, the ErbB hetero-oligomer may comprise two or

more ErbB2 receptors combined with a different ErbB receptor, such as ErbB3, ErbB4 or EGFR (ErbB1). Other proteins, such as a cytokine receptor subunit (*e.g.*, gp130) may be included in the hetero-oligomer.

By "ligand activation of an ErbB receptor" is meant signal transduction (*e.g.*, that
5 caused by an intracellular kinase domain of an ErbB receptor phosphorylating tyrosine residues in the ErbB receptor or a substrate polypeptide) mediated by ErbB ligand binding to a ErbB hetero-oligomer comprising the ErbB receptor of interest. Generally, this will involve binding of an ErbB ligand to an ErbB hetero-oligomer which activates a kinase domain of one or more of the ErbB receptors in the hetero-oligomer and thereby results in
10 phosphorylation of tyrosine residues in one or more of the ErbB receptors and/or phosphorylation of tyrosine residues in additional substrate polypeptides(s). ErbB receptor activation can be quantified using various tyrosine phosphorylation assays.

A "native sequence" polypeptide is one which has the same amino acid sequence as a polypeptide (*e.g.*, ErbB receptor or ErbB ligand) derived from nature. Such native
15 sequence polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. Thus, a native sequence polypeptide can have the amino acid sequence of naturally occurring human polypeptide, murine polypeptide, or polypeptide from any other mammalian species.

The term "amino acid sequence variant" refers to polypeptides having amino acid
20 sequences that differ to some extent from a native sequence polypeptide. Ordinarily, amino acid sequence variants will possess at least about 70% homology with at least one receptor binding domain of a native ErbB ligand or with at least one ligand binding domain of a native ErbB receptor, and preferably, they will be at least about 80%, more preferably, at least about 90% homologous with such receptor or ligand binding domains.
25 The amino acid sequence variants possess substitutions, deletions, and/or insertions at certain positions within the amino acid sequence of the native amino acid sequence.

"Homology" is defined as the percentage of residues in the amino acid sequence variant that are identical after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology. Methods and computer programs for the
30 alignment are well known in the art. One such computer program is "Align 2," authored by Genentech, Inc., which was filed with user documentation in the United States Copyright Office, Washington, DC 20559, on December 10, 1991.

The term "antibody" herein is used in the broadest sense and specifically covers intact monoclonal antibodies, polyclonal antibodies, multispecific antibodies (*e.g.*,

bispecific antibodies) formed from at least two intact antibodies, and antibody fragments, so long as they exhibit the desired biological activity.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations which include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al., *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, *e.g.*, U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson et al., *Nature*, 352:624-628 (1991) and Marks et al., *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include "chimeric" antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; and Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include "primatized" antibodies comprising variable domain antigen-binding sequences derived from a non-human primate (*e.g.*, Old World Monkey, Ape etc) and human constant region sequences.

"Antibody fragments" comprise a portion of an intact antibody, preferably comprising the antigen-binding or variable region thereof. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies;

single-chain antibody molecules; and multispecific antibodies formed from antibody fragment(s).

An "intact" antibody is one which comprises an antigen-binding variable region as well as a light chain constant domain (CL) and heavy chain constant domains, CH1, CH2
5 and CH3. The constant domains may be native sequence constant domains (e.g., human native sequence constant domains) or amino acid sequence variant thereof. Preferably, the intact antibody has one or more effector functions.

Antibody "effector functions" refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an
10 antibody. Examples of antibody effector functions include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g., B cell receptor; BCR), etc.

Depending on the amino acid sequence of the constant domain of their heavy
15 chains, intact antibodies can be assigned to different "classes." There are five major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into "subclasses" (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional
20 configurations of different classes of immunoglobulins are well known.

"Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g., Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for
25 mediating ADCC, NK cells, express Fc γ RIII only, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.*, 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an in vitro ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such
30 assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed in vivo, e.g., in a animal model such as that disclosed in Clynes et al., *PNAS (USA)*, 95:652-656 (1998).

"Human effector cells" are leukocytes which express one or more FcRs and
35 perform effector functions. Preferably, the cells express at least Fc γ RIII and perform

ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, *e.g.*, from blood or PBMCs as described herein.

The terms "Fc receptor" or "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an "activating receptor") and Fc γ RIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (See review M. in *Daëron, Annu. Rev. Immunol.*, 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.*, 9:457-92 (1991); Capel et al., *Immunomethods*, 4:25-34 (1994); and de Haas et al., *J. Lab. Clin. Med.*, 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.*, 117:587 (1976) and Kim et al., *J. Immunol.*, 24:249 (1994)).

"Complement dependent cytotoxicity" or "CDC" refers to the ability of a molecule to lyse a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (*e.g.*, an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, *e.g.*, as described in Gazzano-Santoro et al., *J. Immunol. Methods*, 202:163 (1996), may be performed.

"Native antibodies" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (VH) followed by a number of constant domains. Each light chain has a variable domain at one end (VL) and a

constant domain at its other end. The constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light-chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light chain and heavy chain variable domains.

5 The term “variable” refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions both in the light chain and the heavy chain
10 variable domains. The more highly conserved portions of variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β -sheet configuration, connected by three hypervariable regions, which form loops connecting, and in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close
15 proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of
20 the antibody in antibody dependent cellular cytotoxicity (ADCC).

 The term “hypervariable region” when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region generally comprises amino acid residues from a “complementarity determining region” or “CDR” (e.g., residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain
25 variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)) and/or those residues from a “hypervariable loop” (e.g., residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy
30 chain variable domain; Chothia and Lesk, *J. Mol. Biol.*, 196:901-917 (1987)). “Framework Region” or “FR” residues are those variable domain residues other than the hypervariable region residues as herein defined.

 Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, each with a single antigen-binding site, and a residual “Fc”
35 fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an

F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

“Fv” is the minimum antibody fragment which contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The “light chains” of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

“Single-chain Fv” or “scFv” antibody fragments comprise the VH and VL domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Plückthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994). Anti-ErbB2 antibody scFv fragments are described in WO 93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458.

The term “diabodies” refers to small antibody fragments with two antigen-binding sites, which fragments comprise a variable heavy domain (VH) connected to a variable light domain (VL) in the same polypeptide chain (VH - VL). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding

sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

“Humanized” forms of non-human (*e.g.*, rodent) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992).

Humanized anti-ErbB2 antibodies include huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and huMAb4D5-8 (HERCEPTIN®) as described in Table 3 of U.S. Patent No. 5,821,337 expressly incorporated herein by reference; humanized 520C9 (WO 93/21319) and humanized 2C4 antibodies as described herein below.

An “isolated” antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes

the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

5 An antibody "which binds" an antigen of interest, *e.g.*, ErbB2 antigen, is one capable of binding that antigen with sufficient affinity such that the antibody is useful in targeting a cell expressing the antigen. Where the antibody is one which binds ErbB2, it will usually preferentially bind ErbB2 as opposed to other ErbB receptors, and may be one which does not significantly cross-react with other proteins such as EGFR, ErbB3 or ErbB4. In such embodiments, the extent of binding of the antibody to these non-ErbB2
10 proteins (*e.g.*, cell surface binding to endogenous receptor) will be less than 10% as determined by fluorescence activated cell sorting (FACS) analysis or radioimmunoprecipitation (RIA). Sometimes, the anti-ErbB2 antibody will not significantly cross-react with the rat neu protein, *e.g.*, as described in Schecter et al., *Nature*, 312:513 (1984) and Drebin et al., *Nature*, 312:545-548 (1984).

15 An antibody which "blocks" ligand activation of an ErbB receptor is one which reduces or prevents such activation as hereinabove defined, wherein the antibody is able to block ligand activation of the ErbB receptor substantially more effectively than monoclonal antibody 4D5, *e.g.*, about as effectively as monoclonal antibodies 7F3 or 2C4 or Fab fragments thereof and preferably about as effectively as monoclonal antibody 2C4
20 or a Fab fragment thereof. For example, the antibody that blocks ligand activation of an ErbB receptor may be one which is about 50-100% more effective than 4D5 at blocking formation of an ErbB hetero-oligomer. Blocking of ligand activation of an ErbB receptor can occur by any means, *e.g.*, by interfering with: ligand binding to an ErbB receptor, ErbB complex formation, tyrosine kinase activity of an ErbB receptor in an ErbB
25 complex and/or phosphorylation of tyrosine kinase residue(s) in or by an ErbB receptor. Examples of antibodies which block ligand activation of an ErbB receptor include monoclonal antibodies 2C4 and 7F3 (which block HRG activation of ErbB2/ErbB3 and ErbB2/ErbB4 hetero-oligomers; and EGF, TGF- α , amphiregulin, HB-EGF and/or epiregulin activation of an EGFR/ErbB2 hetero-oligomer); and L26, L96 and L288
30 antibodies (Klapper et al., *Oncogene*, 14:2099-2109 (1997)), which block EGF and NDF binding to T47D cells which express EGFR, ErbB2, ErbB3 and ErbB4.

An antibody having a "biological characteristic" of a designated antibody, such as the monoclonal antibody designated 2C4, is one which possesses one or more of the biological characteristics of that antibody which distinguish it from other antibodies that
35 bind to the same antigen (*e.g.*, ErbB2). For example, an antibody with a biological

characteristic of 2C4 may block HRG activation of an ErbB hetero-oligomer comprising ErbB2 and ErbB3, ErbB1 or ErbB4; block EGF, TGF- α , HB-EGF, epiregulin and/or amphiregulin activation of an ErbB receptor comprising EGFR and ErbB2; block EGF, TGF- α and/or HRG mediated activation of MAPK; and/or bind the same epitope in the
5 extracellular domain of ErbB2 as that bound by 2C4 (*e.g.*, which blocks binding of monoclonal antibody 2C4 to ErbB2).

Unless indicated otherwise, the expression "monoclonal antibody 2C4" refers to an antibody that has antigen binding residues of, or derived from, the murine 2C4 antibody of the Examples below. For example, the monoclonal antibody 2C4 may be murine
0 monoclonal antibody 2C4 or a variant thereof, such as humanized antibody 2C4, possessing antigen binding amino acid residues of murine monoclonal antibody 2C4. Examples of humanized 2C4 antibodies are provided herein and in WO 01/00245, which is incorporated herein by reference in its entirety. Unless indicated otherwise, the
5 expression "rhuMAb 2C4" when used herein refers to an antibody comprising the variable light (VL) and variable heavy (VH) sequences of SEQ ID Nos. 3 and 4, respectively, fused to human light and heavy IgG1 (non-A allotype) constant region sequences optionally expressed by a Chinese Hamster Ovary (CHO) cell.

Unless indicated otherwise, the term "monoclonal antibody 4D5" refers to an antibody that has antigen binding residues of, or derived from, the murine 4D5 antibody
0 (ATCC CRL 10463). For example, the monoclonal antibody 4D5 may be murine monoclonal antibody 4D5 or a variant thereof, such as a humanized 4D5, possessing antigen binding residues of murine monoclonal antibody 4D5. Exemplary humanized 4D5 antibodies include huMAb4D5-1, huMAb4D5-2, huMAb4D5-3, huMAb4D5-4, huMAb4D5-5, huMAb4D5-6, huMAb4D5-7 and huMAb4D5-8 (HERCEPTIN®) as in
5 US Patent No. 5,821,337, with huMAb4D5-8 (HERCEPTIN®) being a preferred humanized 4D5 antibody.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially an ErbB expressing cancer cell either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one which
0 significantly reduces the percentage of ErbB expressing cells in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxanes, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents
5 that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such

as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami et al. (WB Saunders: Philadelphia, 5 1995), especially p. 13.

Examples of "growth inhibitory" antibodies are those which bind to ErbB2 and inhibit the growth of cancer cells overexpressing ErbB2. Preferred growth inhibitory anti-ErbB2 antibodies inhibit growth of SK-BR-3 breast tumor cells in cell culture by greater than 20%, and preferably greater than 50% (e.g., from about 50% to about 100%) at an antibody concentration of about 0.5 to 30 $\mu\text{g/ml}$, where the growth inhibition is 10 determined six days after exposure of the SK-BR-3 cells to the antibody (see U.S. Patent No. 5,677,171 issued October 14, 1997). The SK-BR-3 cell growth inhibition assay is described in more detail in that patent and herein below. The preferred growth inhibitory antibody is monoclonal antibody 4D5, e.g., humanized 4D5.

An antibody which "induces cell death" is one which causes a viable cell to become nonviable. The cell is generally one which expresses the ErbB2 receptor, especially where the cell overexpresses the ErbB2 receptor. Preferably, the cell is a cancer cell, e.g., a breast, ovarian, stomach, endometrial, salivary gland, lung, kidney, colon, thyroid, pancreatic or bladder cell. In vitro, the cell may be a SK-BR-3, BT474, Calu 3, MDA- 20 MB-453, MDA-MB-361 or SKOV3 cell. Cell death in vitro may be determined in the absence of complement and immune effector cells to distinguish cell death induced by antibody-dependent cell-mediated cytotoxicity (ADCC) or complement dependent cytotoxicity (CDC). Thus, the assay for cell death may be performed using heat inactivated serum (*i.e.*, in the absence of complement) and in the absence of immune 25 effector cells. To determine whether the antibody is able to induce cell death, loss of membrane integrity as evaluated by uptake of propidium iodide (PI), trypan blue (see Moore et al., *Cytotechnology*, 17:1-11 (1995)) or 7AAD can be assessed relative to untreated cells. Preferred cell death-inducing antibodies are those which induce PI uptake in the PI uptake assay in BT474 cells (see below).

An antibody which "induces apoptosis" is one which induces programmed cell death as determined by binding of annexin V, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies). The cell is usually one which overexpresses the ErbB2 receptor. Preferably, the cell is a tumor cell, e.g., a breast, ovarian, stomach, endometrial, 35 salivary gland, lung, kidney, colon, thyroid, pancreatic or bladder cell. In vitro, the cell

may be a SK-BR-3, BT474, Calu 3 cell, MDA-MB-453, MDA-MB-361 or SKOV3 cell.

Various methods are available for evaluating the cellular events associated with apoptosis.

For example, phosphatidyl serine (PS) translocation can be measured by annexin binding;

DNA fragmentation can be evaluated through DNA laddering; and nuclear/chromatin

5 condensation along with DNA fragmentation can be evaluated by any increase in

hypodiploid cells. Preferably, the antibody which induces apoptosis is one which results

in about 2 to 50 fold, preferably about 5 to 50 fold, and most preferably about 10 to 50

fold, induction of annexin binding relative to untreated cell in an annexin binding assay

using BT474 cells (see below). Sometimes the pro-apoptotic antibody will be one which

10 further blocks ErbB ligand activation of an ErbB receptor (*e.g.*, 7F3 antibody); *i.e.*, the

antibody shares a biological characteristic with monoclonal antibody 2C4. In other

situations, the antibody is one which does not significantly block ErbB ligand activation of

an ErbB receptor (*e.g.*, 7C2). Further, the antibody may be one like 7C2 which, while

inducing apoptosis, does not induce a large reduction in the percent of cells in S phase

15 (*e.g.*, one which only induces about 0-10% reduction in the percent of these cells relative to control).

The "epitope 2C4" is the region in the extracellular domain of ErbB2 to which the antibody 2C4 binds. In order to screen for antibodies which bind to the 2C4 epitope, a

routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*,

20 *Cold Spring Harbor Laboratory*, Ed Harlow and David Lane (1988), can be performed.

Alternatively, epitope mapping can be performed to assess whether the antibody binds to the 2C4 epitope of ErbB2 (*e.g.*, any one or more residues in the region from about residue 22 to about residue 584 of ErbB2, inclusive; see Figs. 1A-B).

The "epitope 4D5" is the region in the extracellular domain of ErbB2 to which the antibody 4D5 (ATCC CRL 10463) binds. This epitope is close to the transmembrane

domain of ErbB2. To screen for antibodies which bind to the 4D5 epitope, a routine

cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, *Cold Spring Harbor Laboratory*, Ed Harlow and David Lane (1988), can be performed.

30 Alternatively, epitope mapping can be performed to assess whether the antibody binds to the 4D5 epitope of ErbB2 (*e.g.*, any one or more residues in the region from about residue 529 to about residue 625, inclusive; see Figs. 1A-B).

The "epitope 3H4" is the region in the extracellular domain of ErbB2 to which the antibody 3H4 binds. This epitope includes residues from about 541 to about 599, inclusive, in the amino acid sequence of ErbB2 extracellular domain; see Figs. 1A-B.

The "epitope 7C2/7F3" is the region at the N terminus of the extracellular domain of ErbB2 to which the 7C2 and/or 7F3 antibodies (each deposited with the ATCC, see below) bind. To screen for antibodies which bind to the 7C2/7F3 epitope, a routine cross-blocking assay such as that described in *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, epitope mapping can be performed to establish whether the antibody binds to the 7C2/7F3 epitope on ErbB2 (*e.g.*, any one or more of residues in the region from about residue 22 to about residue 53 of ErbB2; see Figs. 1A-B).

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cows, etc. Preferably, the mammal is human.

A tumor which is "responsive to treatment" shows statistically significant improvement in response to anti-ErbB antibody treatment when compared to no treatment or treatment with placebo in a recognized animal model or a human clinical trial, or which responds to initial treatment with anti-ErbB antibodies but grows as treatment is continued.

The terms "treat" or "treatment" refer to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) an undesired physiological change or disorder, such as the development or spread of cancer. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (*i.e.*, not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already with the condition or disorder as well as those prone to have the condition or disorder or those in which the condition or disorder is to be prevented.

A "disorder" is any condition that would benefit from treatment of the present invention. This includes chronic and acute disorders or diseases including those pathological conditions which predispose the mammal to the disorder in question. Non-limiting examples of disorders to be treated herein include benign and malignant tumors; leukemias and lymphoid malignancies, in particular breast, ovarian, stomach, endometrial, salivary gland, lung, kidney, colon, thyroid, pancreatic, prostate or bladder cancer; neuronal, glial, astrocytal, hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, angiogenic and immunologic

disorders. A preferred disorder to be treated in accordance with the present invention is malignant tumor

The term "therapeutically effective amount" refers to an amount of a drug effective to treat a disease or disorder in a mammal. In the case of cancer, the therapeutically effective amount of the drug may reduce the number of cancer cells; reduce the tumor size; inhibit (*i.e.*, slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (*i.e.*, slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic. For cancer therapy, efficacy can, for example, be measured by assessing the time to disease progression (TTP) and/or determining the response rate (RR).

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. A "tumor" comprises one or more cancerous cells. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia or lymphoid malignancies. More particular examples of such cancers include squamous cell cancer (*e.g.*, epithelial squamous cell cancer), lung cancer including small-cell lung cancer, non-small cell lung cancer ("NSCLC"), adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, as well as head and neck cancer.

An "ErbB-expressing cancer" is one comprising cells which have ErbB protein present at their cell surface. An "ErbB2-expressing cancer" is one which produces sufficient levels of ErbB2 at the surface of cells thereof, such that an anti-ErbB2 antibody can bind thereto and have a therapeutic effect with respect to the cancer.

A cancer "characterized by excessive activation" of an ErbB receptor is one in which the extent of ErbB receptor activation in cancer cells significantly exceeds the level of activation of that receptor in non-cancerous cells of the same tissue type. Such excessive activation may result from overexpression of the ErbB receptor and/or greater than normal levels of an ErbB ligand available for activating the ErbB receptor in the cancer cells. Such excessive activation may cause and/or be caused by the malignant state

of a cancer cell. In some embodiments, the cancer will be subjected to a diagnostic or prognostic assay to determine whether amplification and/or overexpression of an ErbB receptor is occurring which results in such excessive activation of the ErbB receptor. Alternatively, or additionally, the cancer may be subjected to a diagnostic or prognostic
5 assay to determine whether amplification and/or overexpression an ErbB ligand is occurring in the cancer which attributes to excessive activation of the receptor. In a subset of such cancers, excessive activation of the receptor may result from an autocrine stimulatory pathway.

In an "autocrine" stimulatory pathway, self stimulation occurs by virtue of the
10 cancer cell producing both an ErbB ligand and its cognate ErbB receptor. For example, the cancer may express or overexpress EGFR and also express or overexpress an EGFR ligand (*e.g.*, EGF, TGF- α , or HB-EGF). In another embodiment, the cancer may express or overexpress ErbB2 and also express or overexpress a heregulin (*e.g.* γ -HRG).

A cancer which "overexpresses" an ErbB receptor is one which has significantly
15 higher levels of an ErbB receptor, such as ErbB2, at the cell surface thereof, compared to a noncancerous cell of the same tissue type. Such overexpression may be caused by gene amplification or by increased transcription or translation. ErbB receptor overexpression may be determined in a diagnostic or prognostic assay by evaluating increased levels of the ErbB protein present on the surface of a cell (*e.g.*, via an immunohistochemistry assay;
20 IHC). Alternatively, or additionally, one may measure levels of ErbB-encoding nucleic acid in the cell, *e.g.*, via fluorescent in situ hybridization (FISH; see WO 98/45479 published October, 1998), southern blotting, or polymerase chain reaction (PCR) techniques, such as real time quantitative PCR (RT-PCR). Overexpression of the ErbB ligand, may be determined diagnostically by evaluating levels of the ligand (or nucleic
25 acid encoding it) in the patient, *e.g.*, in a tumor biopsy or by various diagnostic assays such as the IHC, FISH, southern blotting, PCR or in vivo assays described above. One may also study ErbB receptor overexpression by measuring shed antigen (*e.g.*, ErbB extracellular domain) in a biological fluid such as serum (see, *e.g.*, U.S. Patent No. 4,933,294 issued June 12, 1990; WO 91/05264 published April 18, 1991; U.S. Patent No.
30 5,401,638 issued March 28, 1995; and Sias et al., *J. Immunol. Methods*, 132: 73-80 (1990)). Aside from the above assays, various other in vivo assays are available to the skilled practitioner. For example, one may expose cells within the body of the patient to an antibody which is optionally labeled with a detectable label, *e.g.*, a radioactive isotope, and binding of the antibody to cells in the patient can be evaluated, *e.g.*, by external

scanning for radioactivity or by analyzing a biopsy taken from a patient previously exposed to the antibody.

The tumors overexpressing HER2 are rated by immunohistochemical scores corresponding to the number of copies of HER2 molecules expressed per cell, and can

5 been determined biochemically: 0 = 0-10,000 copies/cell, 1+ = at least about 200,000 copies/cell, 2+ = at least about 500,000 copies/cell, 3+ = at least about 2,000,000 copies/cell. Overexpression of HER2 at the 3+ level, which leads to ligand-independent activation of the tyrosine kinase (Hudziak et al., *Proc. Natl. Acad. Sci. USA*, 84:7159-7163 [1987]), occurs in approximately 30% of breast cancers, and in these patients,
10 relapse-free survival and overall survival are diminished (Slamon et al., *Science*, 244:707-712 [1989]; Slamon et al., *Science*, 235:177-182 [1987]).

Conversely, a cancer which is “not characterized by overexpression of the ErbB2 receptor” is one which, in a diagnostic assay, does not express higher than normal levels of ErbB2 receptor compared to a noncancerous cell of the same tissue type.

15 A “hormone independent” cancer is one in which proliferation thereof is not dependent on the presence of a hormone which binds to a receptor expressed by cells in the cancer. Such cancers do not undergo clinical regression upon administration of pharmacological or surgical strategies that reduce the hormone concentration in or near the tumor. Examples of hormone independent cancers include androgen independent prostate cancer,
20 estrogen independent breast cancer, endometrial cancer and ovarian cancer. Such cancers may begin as hormone dependent tumors and progress from a hormone-sensitive stage to a hormone-refractory tumor following anti-hormonal therapy.

The term “cytotoxic agent” as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to
25 include radioactive isotopes (e.g., At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32 and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

A “chemotherapeutic agent” is a chemical compound useful in the treatment of
30 cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN™); alkyl sulfonates such as busulfan, improsulfan and pipsulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and
35 trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine,

cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, 5 bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, 10 zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone 15 propionate, epitiostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; 20 mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, *e.g.*, paclitaxel (TAXOL®, 25 Bristol-Myers Squibb Oncology, Princeton, NJ) and docetaxel (TAXOTERE®, Rhône-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; 30 xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition, are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)- 35 imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and

toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

As used herein, the term "EGFR-targeted drug" refers to a therapeutic agent that binds to EGFR and, optionally, inhibits EGFR activation. Examples of such agents include antibodies and small molecules that bind to EGFR. Examples of antibodies which bind to EGFR include MAb 579 (ATCC CRL HB 8506), MAb 455 (ATCC CRL HB8507), MAb 225 (ATCC CRL 8508), MAb 528 (ATCC CRL 8509) (see, U.S. Patent No. 4,943, 533, Mendelsohn *et al.*) and variants thereof, such as chimerized 225 (C225 or Cetuximab; ERBUTIX®) and reshaped human 225 (H225) (see, WO 96/40210, Imclone Systems Inc.); antibodies that bind type II mutant EGFR (U.S. Patent No. 5,212,290); humanized and chimeric antibodies that bind EGFR as described in U.S. Patent No. 5,891,996; and human antibodies that bind EGFR, such as ABX-EGF (see WO 98/50433, Abgenix). The anti-EGFR antibody may be conjugated with a cytotoxic agent, thus generating an immunoconjugate (see, *e.g.*, EP 659,439A2, Merck Patent GmbH). Examples of small molecules that bind to EGFR include ZD1839 or Gefitinib (IRESSA™; Astra Zeneca), CP-358774 (TARCEVA™; Genentech/OSI) and AG1478, AG1571 (SU 5271; Sugen). A "tyrosine kinase inhibitor" is a molecule which inhibits to some extent tyrosine kinase activity of a tyrosine kinase such as an ErbB receptor. Examples of such inhibitors include the EGFR-targeted drugs noted in the preceding paragraph as well as quinazolines such as PD 153035, 4-(3-chloroanilino) quinazoline, pyridopyrimidines, pyrimidopyrimidines, pyrrolopyrimidines, such as CGP 59326, CGP 60261 and CGP 62706, and pyrazolopyrimidines, 4-(phenylamino)-7H-pyrrolo[2,3-d] pyrimidines, curcumin (diferuloyl methane, 4,5-bis (4-fluoroanilino)phthalimide), tyrophostines containing nitrothiophene moieties; PD-0183805 (Warner-Lambert); antisense molecules (*e.g.*, those that bind to ErbB-encoding nucleic acid); quinoxalines (U.S. Patent No. 5,804,396); tyrophostins (U.S. Patent No. 5,804,396); ZD6474 (Astra Zeneca); PTK-787 (Novartis/Schering AG); pan-ErbB inhibitors such as CI-1033 (Pfizer); Affinitac (ISIS 3521; Isis/Lilly); Imatinib mesylate (Gleevac; Novartis); PKI 166 (Novartis); GW2016 (Glaxo SmithKline); CI-1033 (Pfizer); EKB-569 (Wyeth); Semaxanib (Sugen); ZD6474 (AstraZeneca); PTK-787 (Novartis/Schering AG); INC-1C11 (Imclone); or as described in any of the following patent publications: U.S. Patent No. 5,804,396; WO 99/09016 (American Cyanamid); WO 98/43960 (American Cyanamid); WO 97/38983 (Warner Lambert); WO 99/06378 (Warner Lambert); WO 99/06396 (Warner Lambert);

WO 96/30347 (Pfizer, Inc); WO 96/33978 (Zeneca); WO 96/3397 (Zeneca); and WO 96/33980 (Zeneca).

An "anti-angiogenic agent" refers to a compound which blocks, or interferes with to some degree, the development of blood vessels. The anti-angiogenic factor may, for instance, be a small molecule or antibody that binds to a growth factor or growth factor receptor involved in promoting angiogenesis. The preferred anti-angiogenic factor herein is an antibody that binds to Vascular Endothelial Growth Factor (VEGF).

The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor- α and - β ; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; a tumor necrosis factor such as TNF- α or TNF- β ; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to tumor cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active parent form. See, *e.g.*, Wilman, "Prodrugs in Cancer Chemotherapy" Biochemical Society Transactions, 14, pp. 375-382, 615th Meeting Belfast (1986) and Stella et al., "Prodrugs: A Chemical Approach to Targeted Drug Delivery," Directed Drug Delivery, Borchardt et al., (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention include, but are not limited to, phosphate-containing prodrugs, thiophosphate-

containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, β -lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described above.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as the anti-ErbB2 antibodies disclosed herein and, optionally, a chemotherapeutic agent) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products.

A "cardioprotectant" is a compound or composition which prevents or reduces myocardial dysfunction (*i.e.*, cardiomyopathy and/or congestive heart failure) associated with administration of a drug, such as an anthracycline antibiotic and/or an anti-ErbB2 antibody, to a patient. The cardioprotectant may, for example, block or reduce a free-radical-mediated cardiotoxic effect and/or prevent or reduce oxidative-stress injury. Examples of cardioprotectants encompassed by the present definition include the iron-chelating agent dexrazoxane (ICRF-187) (Seifert et al., *The Annals of Pharmacotherapy*, 28:1063-1072 (1994)); a lipid-lowering agent and/or anti-oxidant such as probucol (Singal et al., *J. Mol. Cell Cardiol.*, 27:1055-1063 (1995)); amifostine (aminothiols 2-[(3-aminopropyl)amino]ethanethiol-dihydrogen phosphate ester, also called WR-2721, and the dephosphorylated cellular uptake form thereof called WR-1065) and S-3-(3-methylaminopropylamino)propylphosphorothioic acid (WR-151327), see Green et al., *Cancer Research*, 54:738-741 (1994); digoxin (Bristow, M.R. In: Bristow MR, ed. *Drug-Induced Heart Disease*. New York: Elsevier 191-215 (1980)); beta-blockers such as metoprolol (Hjalmarson et al., *Drugs* 47:Suppl 4:31-9 (1994); and Shaddy et al., *Am. Heart J.*, 129:197-9 (1995)); vitamin E; ascorbic acid (vitamin C); free radical scavengers such as oleanolic acid, ursolic acid and N-acetylcysteine (NAC); spin trapping compounds such as alpha-phenyl-tert-butyl nitron (PBN); (Paracchini et al., *Anticancer Res.*, 13:1607-1612 (1993)); selenoorganic compounds such as P251 (Elbesen); and the like.

An "isolated" nucleic acid molecule is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the antibody nucleic acid. An isolated nucleic acid molecule is other than in the form or setting in which it is found in nature.

5 Isolated nucleic acid molecules therefore are distinguished from the nucleic acid molecule as it exists in natural cells. However, an isolated nucleic acid molecule includes a nucleic acid molecule contained in cells that ordinarily express the antibody where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

10 The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

15 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation.

20 Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

25 As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent

30 mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

Methods of Identifying Tumors that are Responsive to Treatment with Anti-HER2 Antibodies

Sources of Tumors and Tumor Cells

Tumors can be characterized as responsive to therapy with 2C4, or functionally
5 equivalent antibodies, that is, antibodies having one or more of the biological
characteristics of antibody 2C4, *e.g.*, based on the presence of EGFR-ErbB2 and/or
ErbB2-ErbB3 heterodimers on the cell surface, as a measure of HER2 activation. Tumor
samples may be assayed for the presence of heterodimers by any method known in the art.
Preferably, the presence of heterodimers is determined by one or more of the methods
10 described below.

Since HER2 activation is the result of receptor heterodimerization and
phosphorylation, a particularly important method for identifying tumors responsive to
therapy with 2C4, or functionally equivalent antibodies, is the detection of
phosphorylation of ErbB receptor, such as phosphorylation of ErbB2 (HER2) receptor, as
15 described below.

Sources of tumor cells that may be assayed include, but are not limited to, tumor
biopsies, circulating tumor cells, circulating plasma proteins, ascitic fluid,
xenotransplanted tumors and other tumor models, and primary cell cultures or cell lines
derived from tumors or exhibiting tumor-like properties, as well as preserved tumor
20 samples, such as formalin-fixed, paraffin-embedded tumor samples. The screening of
panels of various tumor cell types for EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers,
and/or phosphorylation of ErbB receptor is contemplated by the present invention. Tumor
cells of the same type as tumor cells that test positive for heterodimers, and/or
phosphorylation of ErbB receptor, such as ErbB2 (HER2) receptor, may be subjected to
25 therapy with 2C4. The tumor models described below are provided as examples and
should not be construed as limiting the invention.

In one embodiment, tumor cells that originate with a patient currently suffering
from a tumor are assayed for responsiveness to therapy with 2C4. If the cells are
determined to be responsive, based on the presence of HER2/HER3 and/or HER2/HER1
30 heterodimers or by demonstrating the phosphorylation of ErbB receptor, the patient may
subsequently be treated with an antibody with one or more of the biological characteristics
of 2C4. Preferably, the patient is treated with rhuMAb 2C4.

In another embodiment, tumor cells from particular type of tumor or cells that are
believed to have the characteristics of a particular type of tumor are assayed for the
35 presence of EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers, or for the phosphorylation

of ErbB receptor, preferable ErbB2 (HER2) receptor. If EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers and/or phosphorylation of ErbB receptor is detected, that type of tumor is considered to be a good candidate for treatment with an anti-ErbB2 antibody with one or more of the biological characteristics of 2C4. Patients suffering from that type of tumor may then be treated with such an antibody.

Cell Line Xenografts

In vitro propagated tumor cells, such as tumor cells grown in culture and tumor cell lines, may be xenografted into mice by implanting cells directly into a site of interest. Such methods are well known to one of skill in the art. The cells are assayed to identify the presence of EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers, or for the phosphorylation of ErbB receptor, such as the phosphorylation of ErbB2 (HER2) receptor.

In one embodiment, tumor cells are implanted subcutaneously into a mouse, preferably an athymic nude mouse. In another embodiment, tumor cells are implanted into a physiologically relevant location to create an appropriate in situ tumor model. For example, cells from a breast cancer cell line may be implanted at various concentrations into the mammary fat pad of athymic nude mice to more accurately model the biology of breast cancer. Tumor cells may be assayed for the presence of EGFR-ErbB2 or ErbB2-ErbB3 heterodimers, or for the phosphorylation of ErbB receptor either before or after implantation. Preferably, tumor cells are assayed after the implanted cells have developed into a tumor of a predetermined size. The mice may also be subjected to therapy with 2C4 or a functionally equivalent antibody, with untreated mice serving as a control.

Similar models may be established for any type of tumor from which cultured cells or cell lines have been derived. Tumor types include, but are not limited to, bladder, brain, breast, colon, esophagus, kidney, leukemia, liver, lung, melanoma, ovary, pancreas, prostate, sarcoma, stomach, testicle, and uterus. In one embodiment, tumor cells or cell lines that overexpress ErbB2 are used for implantation, while in another embodiment, tumor cells or cell lines that express normal or below normal amounts of ErbB2 are used for implantation. In yet another embodiment, tumor cells or cell lines non-responsive to HERCEPTIN® are used for implantation.

In a specific embodiment, approximately 20 million MDA-175 breast tumor cells are implanted into the mouse gonadal fat pad. Expression of HER2/HER1 and/or HER2/HER3 dimers on the surface of xenografted cells is determined, such as by one of the methods described below. Mice thus implanted may also be subjected to treatment

with 0, 3 mg/kg, 10 mg/kg, 30 mg/kg, or 100 mg/kg 2C4. Other dosage regimens would be within the determination of one of ordinary skill in the art.

While the present invention is suitable for the classification of any HER2 expressing tumor, solid tumors, like breast cancer, ovarian cancer, lung cancer, prostate cancer and colorectal cancer, are particularly suitable for analysis and treatment following the present invention.

Tumor Xenografts

Mammalian tumor specimens, preferably human tumor specimens, may be obtained and implanted into mice, preferably athymic nude mice. The tumor specimens may be obtained by any method known in the art. In one embodiment the tumor specimens are surgically resected, such as in a biopsy or in the process of surgery to remove the tumor from the mammal. In another embodiment the tumor specimen is obtained by purifying circulating tumor cells from the mammals blood.

In a specific embodiment, solid human tumor slices of approximately 5 x 5 x 0.5 to 1 mm in diameter are implanted into the flanks of athymic nude mice, generally four fragments per mouse. When the implanted tumors reach a median diameter of about 10-15 mm, they may be serially passaged, generally using smaller tumor fragments. Methods of generating and studying human tumor xenografts are described in the following references, herein incorporated in their entirety: Fiebig et al., "Human Tumor Xenografts: Predictivity, Characterization and Discovery of New Anticancer Agents," in Contributions to Oncology: Relevance of Tumor Models for Anticancer Drug Development, Fiebig & Burger, eds. (Basel, Karger 1999), vol. 54, pp. 29-50; Berger et al., "Establishment and Characterization of Human Tumor Xenografts in Thymus-Aplastic Nude Mice," in Immunodeficient Mice in Oncology, Fiebig & Berger, eds. (Basel, Karger 1992), pp. 23-46; Fiebig & Burger, "Human Tumor Xenografts and Explants," in Models in Cancer Research, Teicher, ed. (Humana Press 2002) pp. 113-137.

Human xenografts are considered highly predictive of tumor behavior within the donor patient, as the xenograft grows as a solid tumor, differentiates, and develops a stroma, vasculature, and a central necrosis. In most cases, xenografts retain most of the molecular, histological, and pathophysiological characteristics of the fresh patient-derived tumor. Tumor cells from mice containing first or serially passaged tumors may be analyzed for the presence of EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers, or for the phosphorylation of ErbB receptor. Mice may also be subjected to therapy with 2C4 or a functionally equivalent antibody.

In one embodiment, a newly created or established panel of human tumor xenografts is screened for the presence of EGFR-ErbB2 or ErbB2-ErbB3 heterodimers, or for phosphorylation of ErbB receptor. Fiebig & Burger, supra, describe a panel of over 300 human tumor xenografts established from a variety of common cancer types, such as bladder, brain, breast, colon, esophagus, kidney, leukemia, liver, lung, melanoma, ovary, pancreas, prostate, sarcoma, stomach, testicle, and uterus. In one embodiment, the entire panel is screened for heterodimers, or for phosphorylation of ErbB receptor, such as ErbB2 (HER2) receptor. Subsets of this panel may also be screened for heterodimers, or for phosphorylation of ErbB receptor, wherein subsets are based on, for example, tissue type, over-, under-, or normal expression of ErbB2, or failure to respond to HERCEPTIN®. In this manner, tumors may be categorized as candidates for therapy with 2C4 based on the presence of heterodimers, or by demonstrating the phosphorylation of ErbB receptor, such as ErbB2 (HER2) receptor. Likewise, patients possessing tumors thus categorized may be more rapidly deemed eligible for therapy with 2C4 or a functionally equivalent antibody.

Tumor specimens may be assayed for the presence of EGFR-ErbB2 or ErbB2-ErbB3 heterodimers, or for the phosphorylation of ErbB receptor either before or after implantation. In one embodiment, approximately one gram of tumor from a first and/or serially passaged xenograft is further characterized molecularly for heterodimers or snap frozen in liquid nitrogen and stored at -80°C for later characterization. Xenograft tumors may be further analyzed by a double layer soft-agar assay, also called a clonogenic assay, as described, for example, in Fiebig & Burger, supra. Solid human tumor xenografts are mechanically and proteolytically disaggregated into a single-cell suspension, which is plated into multiwell plates layered with soft agar as described. Tumor cell growth in vitro leads to the formation of colonies, which may be further analyzed for molecular characteristics, such as heterodimers, or for phosphorylation of ErbB receptor, or for other histochemical or morphological characteristics.

B. Detection of EGFR-ErbB2 and ErbB2-ErbB3 Heterodimers

Any method known in the art may be used to detect EGFR-ErbB2 or ErbB2-ErbB3 heterodimers in tumors. Several preferred methods are described below. These methods detect noncovalent protein-protein interactions or otherwise indicate proximity between proteins of interest. The methods described below are provided as examples and should not be construed as limiting the invention.

Co-Immunoprecipitation and Immunoblotting

Immunoaffinity-based methods, such as immunoprecipitation or ELISA, may be used to detect EGFR-ErbB2 or ErbB2-ErbB3 heterodimers. In one embodiment, anti-ErbB2 antibodies are used to immunoprecipitate complexes comprising ErbB2 from tumor cells, and the resulting immunoprecipitant is then probed for the presence of EGFR or ErbB3 by immunoblotting. In another embodiment, EGFR or ErbB3 antibodies may be used for the immunoprecipitation step and the immunoprecipitant then probed with ErbB2 antibodies. In a further embodiment, ErbB ligands specific to HER1, HER3, HER2/HER1 complexes or HER2/HER3 complexes may be used to precipitate complexes, which are then probed for the presence of HER2. For example, ligands may be conjugated to avidin and complexes purified on a biotin column.

In other embodiments, such as ELISA or antibody "sandwich"-type assays, antibodies to ErbB2 are immobilized on a solid support, contacted with tumor cells or tumor cell lysate, washed, and then exposed to antibody against EGFR or ErbB3. Binding of the latter antibody, which may be detected directly or by a secondary antibody conjugated to a detectable label, indicates the presence of heterodimers. In certain embodiments, EGFR or ErbB3 antibody is immobilized, and ErbB2 antibody is used for the detection step. In other embodiments ErbB receptor ligands may be used in place of, or in combination with anti-ErbB receptor antibodies.

Immunoprecipitation with EGFR, ErbB2, or ErbB3 antibody may be followed by a functional assay for heterodimers, as an alternative or supplement to immunoblotting. In one embodiment, immunoprecipitation with ErbB3 antibody is followed by an assay for receptor tyrosine kinase activity in the immunoprecipitant. Because ErbB3 does not have intrinsic tyrosine kinase activity, the presence of tyrosine kinase activity in the immunoprecipitant indicates that ErbB3 is most likely associated with ErbB2. Graus-Porta et al., *EMBO J.*, 16:1647-55 (1997); Klapper et al., *Proc. Natl. Acad. Sci. USA*, 96:4995-5000 (1999). This result may be confirmed by immunoblotting with ErbB2 antibodies. In another embodiment, immunoprecipitation with ErbB2 antibody is followed by an assay for EGFR receptor tyrosine kinase activity. In this assay, the immunoprecipitant is contacted with radioactive ATP and a peptide substrate that mimics the *in vivo* site of transphosphorylation of ErbB2 by EGFR. Phosphorylation of the peptide indicates co-immunoprecipitation and thus heterodimerization of EGFR with ErbB2. Receptor tyrosine kinase activity assays are well known in the art and include assays that detect phosphorylation of target substrates, for example, by phosphotyrosine

antibody, and activation of cognate signal transduction pathways, such as the MAPK pathway.

Variations on the above methods and assays would be readily apparent and routine to one of ordinary skill in the art.

5 Chemical or UV cross-linking may also be used to covalently join heterodimers on the surface of living cells. Hunter et al., *Biochem. J.*, 320:847-53. Examples of chemical cross-linkers include dithiobis(succinimidyl) propionate (DSP) and 3,3'-dithiobis(sulphosuccinimidyl) propionate (DTSSP). In one embodiment, cell extracts from chemically cross-linked tumor cells are analyzed by SDS-PAGE and immunoblotted
10 with antibodies to EGFR and/or ErbB3. A supershifted band of the appropriate molecular weight most likely represents EGFR-ErbB2 or ErbB2-ErbB3 heterodimers, as ErbB2 is the preferred heterodimerization partner for EGFR and ErbB3. This result may be confirmed by subsequent immunoblotting with ErbB2 antibodies.

FRET and Fluorescence-Based Methods

15 Fluorescence resonance energy transfer (FRET) may also be used to detect EGFR-ErbB2 or ErbB2-ErbB3 heterodimers. FRET detects protein conformational changes and protein-protein interactions in vivo and in vitro based on the transfer of energy from a donor fluorophore to an acceptor fluorophore. Selvin, *Nat. Struct. Biol.*, 7:730-34 (2000). Energy transfer takes place only if the donor fluorophore is in sufficient proximity to the
20 acceptor fluorophore. In a typical FRET experiment, two proteins or two sites on a single protein are labeled with different fluorescent probes. One of the probes, the donor probe, is excited to a higher energy state by incident light of a specified wavelength. The donor probe then transmits its energy to the second probe, the acceptor probe, resulting in a reduction in the donor's fluorescence intensity and an increase in the acceptor's
25 fluorescence emission. To measure the extent of energy transfer, the donor's intensity in a sample labeled with donor and acceptor probes is compared with its intensity in a sample labeled with donor probe only. Optionally, acceptor intensity is compared in donor/acceptor and acceptor only samples. Suitable probes are known in the art and include, for example, membrane permeant dyes, such as fluorescein and rhodamine,
30 organic dyes, such as the cyanine dyes, and lanthanide atoms. Selvin, supra. Methods and instrumentation for detecting and measuring energy transfer are also known in the art. Selvin, supra.

FRET-based techniques suitable for detecting and measuring protein-protein interactions in individual cells are also known in the art. For example, donor

photobleaching fluorescence resonance energy transfer (pbFRET) microscopy and fluorescence lifetime imaging microscopy (FLIM) may be used to detect the dimerization of cell surface receptors. Selvin, *supra*; Gadella & Jovin, *J. Cell Biol.*, 129:1543-58 (1995). In one embodiment, pbFRET is used on cells either "in suspension" or "in situ" to
5 detect and measure the formation of EGFR-ErbB2 or ErbB2-ErbB3 heterodimers, as described in Nagy et al., *Cytometry*, 32:120-131 (1998). These techniques measure the reduction in a donor's fluorescence lifetime due to energy transfer. In a particular embodiment, a flow cytometric Foerster-type FRET technique (FCET) may be used to investigate EGFR-ErbB2 and ErbB2-ErbB3 heterodimerization, as described in Nagy et
10 al., *supra*, and Brockhoff et al., *Cytometry*, 44:338-48 (2001).

FRET is preferably used in conjunction with standard immunohistochemical labeling techniques. Kenworthy, *Methods*, 24:289-96 (2001). For example, antibodies conjugated to suitable fluorescent dyes can be used as probes for labeling two different proteins. If the proteins are within proximity of one another, the fluorescent dyes act as
15 donors and acceptors for FRET. Energy transfer is detected by standard means. Energy transfer may be detected by flow cytometric means or by digital microscopy systems, such as confocal microscopy or wide-field fluorescence microscopy coupled to a charge-coupled device (CCD) camera.

In one embodiment of the present invention, ErbB2 antibodies and either EGFR or
20 ErbB3 antibodies are directly labeled with two different fluorophores, for example as described in Nagy et al., *supra*. Tumor cells or tumor cell lysates are contacted with the differentially labeled antibodies, which act as donors and acceptors for FRET in the presence of EGFR-ErbB2 or ErbB2-ErbB3 heterodimers. Alternatively, unlabeled antibodies against ErbB2 and either EGFR or ErbB3 are used along with differentially
25 labeled secondary antibodies that serve as donors and acceptors. See, for example, Brockhoff et al., *supra*. Energy transfer is detected and the presence of heterodimers is determined if the labels are found to be in close proximity.

In other embodiments ErbB receptor ligands that are specific for HER2 and either HER1 or HER3 are fluorescently labeled and used for FRET studies.

30 In still other embodiments of the present invention, the presence of heterodimers on the surface of tumor cells is demonstrated by co-localization of ErbB2 with either EGFR or ErbB3 using standard direct or indirect immunofluorescence techniques and confocal laser scanning microscopy. Alternatively, laser scanning imaging (LSI) is used to detect antibody binding and co-localization of ErbB2 with either EGFR or ErbB3 in a

high-throughput format, such as a microwell plate, as described in Zuck et al, *Proc. Natl. Acad. Sci. USA*, 96:11122-27 (1999).

In further embodiments, the presence of EGFR-ErbB2 and/or ErbB2-ErbB3 heterodimers is determined by identifying enzymatic activity that is dependent upon the proximity of the heterodimer components. A ErbB2 antibody is conjugated with one enzyme and an EGFR or ErbB3 antibody is conjugated with a second enzyme. A first substrate for the first enzyme is added and the reaction produces a second substrate for the second enzyme. This leads to a reaction with another molecule to produce a detectable compound, such as a dye. The presence of another chemical breaks down the second substrate, so that reaction with the second enzyme is prevented unless the first and second enzymes, and thus the two antibodies, are in close proximity. In a particular embodiment tumor cells or cell lysates are contacted with an ErbB2 antibody that is conjugated with glucose oxidase and an ErbB3 or ErbB1 antibody that is conjugated with horse radish peroxidase. Glucose is added to the reaction, along with a dye precursor, such as DAB, and catalase. The presence of heterodimers is determined by the development of color upon staining for DAB.

eTag™ Assay System

Heterodimers may be detected using methods based on the eTag™ assay system (Aclara Bio Sciences, Mountain View, CA), as described, for example, WO 83502 and in U.S. Patent Application 2001/0049105, published December 6, 2001, both of which are expressly incorporated by reference in their entirety. An eTag™, or “electrophoretic tag,” comprises a detectable reporter moiety, such as a fluorescent group. It may also comprise a “mobility modifier,” which consists essentially of a moiety having a unique electrophoretic mobility. These moieties allow for separation and detection of the eTag™ from a complex mixture under defined electrophoretic conditions, such as capillary electrophoresis (CE). The portion of the eTag™ containing the reporter moiety and, optionally, the mobility modifier is linked to a first target binding moiety by a cleavable linking group to produce a first binding compound. The first target binding moiety specifically recognizes a particular first target, such as a nucleic acid or protein. The first target binding moiety is not limited in any way, and may be for example, a polynucleotide or a polypeptide. Preferably, the first target binding moiety is an antibody or antibody fragment. Alternatively, the first target binding moiety may be an ErbB receptor ligand or binding-competent fragment thereof.

The linking group preferably comprises a cleavable moiety, such as an enzyme substrate, or any chemical bond that may be cleaved under defined conditions. When the first target binding moiety binds to its target, the cleaving agent is introduced and/or activated, and the linking group is cleaved, thus releasing the portion of the eTag™ containing the reporter moiety and mobility modifier. Thus, the presence of a “free” eTag™ indicates the binding of the target binding moiety to its target.

Preferably, a second binding compound comprises the cleaving agent and a second target binding moiety that specifically recognizes a second target. The second target binding moiety is also not limited in any way and may be, for example, an antibody or antibody fragment or an ErbB receptor ligand or binding competent ligand fragment. The cleaving agent is such that it will only cleave the linking group in the first binding compound if the first binding compound and the second binding compound are in close proximity.

In an embodiment of the present invention, a first binding compound comprises an eTag™ in which an antibody to ErbB2 serves as the first target binding moiety. A second binding compound comprises an antibody to EGFR or ErbB3 joined to a cleaving agent capable of cleaving the linking group of the eTag™. Preferably the cleaving agent must be activated in order to be able to cleave the linking group. Tumor cells or tumor cell lysates are contacted with the eTag™, which binds to ErbB2, and with the modified EGFR or ErbB3 antibody, which binds to EGFR or ErbB3 on the cell surface. Unbound binding compound is preferably removed, and the cleaving agent is activated, if necessary. If EGFR-ErbB2 or ErbB2-ErbB3 heterodimers are present, the cleaving agent will cleave the linking group and release the eTag™ due to the proximity of the cleaving agent to the linking group. Free eTag™ may then be detected by any method known in the art, such as capillary electrophoresis.

In one embodiment, the cleaving agent is an activatable chemical species that acts on the linking group. For example, the cleaving agent may be activated by exposing the sample to light.

In another embodiment, the eTag™ is constructed using an antibody to EGFR or ErbB3 as the first target binding moiety, and the second binding compound is constructed from an antibody to ErbB2.

Detection of Phosphorylation of ErbB Receptor

The presence of the phosphorylation of ErbB receptor may be used to demonstrate HER2 activation.

In one embodiment, phosphorylation of ErbB receptor is assessed by immunoprecipitation of one or more ErbB receptors, such as ErbB2 (HER2) receptor, and Western blot analysis. For example, positivity is determined by the presence of a phospho-HER2 band on the gel, using an anti-phosphotyrosine antibody to detect phosphorylated tyrosine residue(s) in the immunoprecipitated ErbB receptor(s). Anti-phosphotyrosine antibodies are commercially available from PanVera (Madison, WI), a subsidiary of Invitrogen, Chemicon International Inc. (Temecula, CA), or Upstate Biotechnology (Lake Placid, NY). Negativity is determined by the absence of the band.

In another embodiment, phosphorylation of ErbB2 (HER2) receptor is assessed by immunohistochemistry using a phospho-specific anti-HER2 antibody (clone PN2A; Thor et al., *J. Clin. Oncol.*, 18(18):3230-3239 (2000)).

Other methods for detecting phosphorylation of ErbB receptor(s) include, but are not limited to, KIRA ELISA (U.S. Patent Nos. 5,766,863; 5,891,650; 5,914,237; 6,025,145; and 6,287,784), mass spectrometry (comparing size of phosphorylated and non-phosphorylated HER2), and e-tag proximity assay with anti-HER2 antibody (e.g., using the eTag™ assay kit available from Aclara BioSciences (Mountain View, CA). Details of the eTag™ assay are described hereinabove.

Production of Antibodies

A description follows as to exemplary techniques for the production of the therapeutic and diagnostic antibodies used in accordance with the present invention.

While the description is generally directed to the production of anti-ErbB2 antibodies, one of skill in the art can readily adapt the disclosure to produce antibodies against any of the ErbB receptors.

The ErbB2 antigen to be used for production of antibodies may be, e.g., a soluble form of the extracellular domain of ErbB2 or a portion thereof, containing the desired epitope. Alternatively, cells expressing ErbB2 at their cell surface (e.g., NIH-3T3 cells transformed to overexpress ErbB2; or a carcinoma cell line such as SK-BR-3 cells, see Stancovski et al., *PNAS (USA)*, 88:8691-8695 (1991)) can be used to generate antibodies. Other forms of ErbB2 useful for generating antibodies will be apparent to those skilled in the art.

Polyclonal antibodies

Polyclonal antibodies are preferably raised in animals by multiple subcutaneous (sc) or intraperitoneal (ip) injections of the relevant antigen and an adjuvant. It may be useful to conjugate the relevant antigen to a protein that is immunogenic in the species to

be immunized, *e.g.*, keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues), N-hydroxysuccinimide (through lysine residues), glutaraldehyde, succinic anhydride, SOCl₂, or R₁N=C=NR, where R and R₁ are different alkyl groups.

Animals are immunized against the antigen, immunogenic conjugates, or derivatives by combining, *e.g.*, 100 μg or 5 μg of the protein or conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the animals are boosted with 1/5 to 1/10 the original amount of peptide or conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later the animals are bled and the serum is assayed for antibody titer. Animals are boosted until the titer plateaus. Preferably, the animal is boosted with the conjugate of the same antigen, but conjugated to a different protein and/or through a different cross-linking reagent. Conjugates also can be made in recombinant cell culture as protein fusions. Also, aggregating agents such as alum are suitably used to enhance the immune response.

Monoclonal antibodies

Monoclonal antibodies are obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Thus, the modifier "monoclonal" indicates the character of the antibody as not being a mixture of discrete antibodies.

For example, the monoclonal antibodies may be made using the hybridoma method first described by Kohler et al., *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567).

In the hybridoma method, a mouse or other appropriate host animal, such as a hamster, is immunized as hereinabove described to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the protein used for immunization. Alternatively, lymphocytes may be immunized *in vitro*. Lymphocytes then are fused with myeloma cells using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986)).

The hybridoma cells thus prepared are seeded and grown in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival

of the unfused, parental myeloma cells. For example, if the parental myeloma cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine (HAT medium), which substances prevent the growth of HGPRT-deficient
5 cells.

Preferred myeloma cells are those that fuse efficiently, support stable high-level production of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. Among these, preferred myeloma cell lines are murine myeloma lines, such as those derived from MOPC-21 and MPC-11 mouse tumors
10 available from the Salk Institute Cell Distribution Center, San Diego, California USA, and SP-2 or X63-Ag8-653 cells available from the American Type Culture Collection, Rockville, Maryland USA. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); and Brodeur et al., *Monoclonal Antibody Production*
15 *Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)).

Culture medium in which hybridoma cells are growing is assayed for production of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or
20 enzyme-linked immunoabsorbent assay (ELISA).

The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson et al., *Anal. Biochem.*, 107:220 (1980).

After hybridoma cells are identified that produce antibodies of the desired specificity, affinity, and/or activity, the clones may be subcloned by limiting dilution
25 procedures and grown by standard methods (Goding, *Monoclonal Antibodies: Principles and Practice*, pp. 59-103 (Academic Press, 1986)). Suitable culture media for this purpose include, for example, D-MEM or RPMI-1640 medium. In addition, the hybridoma cells may be grown in vivo as ascites tumors in an animal.

The monoclonal antibodies secreted by the subclones are suitably separated from
30 the culture medium, ascites fluid, or serum by conventional antibody purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding
35 specifically to genes encoding the heavy and light chains of murine antibodies). The

hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as E. coli cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce antibody protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra et al., *Curr. Opinion in Immunol.*, 5:256-262 (1993) and Plückthun, *Immunol. Revs.*, 130:151-188 (1992).

In a further embodiment, monoclonal antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in McCafferty et al., *Nature*, 348:552-554 (1990). Clackson et al., *Nature*, 352:624-628 (1991) and Marks et al., *J. Mol. Biol.*, 222:581-597 (1991) describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks et al., *Bio/Technology*, 10:779-783 (1992)), as well as combinatorial infection and in vivo recombination as a strategy for constructing very large phage libraries (Waterhouse et al., *Nuc. Acids. Res.*, 21:2265-2266 (1993)). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

The DNA also may be modified, for example, by substituting the coding sequence for human heavy chain and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; and Morrison, et al., *Proc. Natl Acad. Sci. USA*, 81:6851 (1984)), or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody, or they are substituted for the variable domains of one antigen-combining site of an antibody to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for an antigen and another antigen-combining site having specificity for a different antigen.

Humanized antibodies

Methods for humanizing non-human antibodies have been described in the art.

Preferably, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of

Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)), by substituting hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Patent No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some hypervariable region residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is very important to reduce antigenicity. According to the so-called "best-fit" method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework region (FR) for the humanized antibody (Sims et al., *J. Immunol.*, 151:2296 (1993); Chothia et al., *J. Mol. Biol.*, 196:901 (1987)). Another method uses a particular framework region derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (Carter et al., *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta et al., *J. Immunol.*, 151:2623 (1993)).

It is further important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *i.e.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the hypervariable region residues are directly and most substantially involved in influencing antigen binding.

Exemplary humanized anti-ErbB2 antibodies which bind ErbB2 and block ligand activation of an ErbB receptor are described in WO 01/0245, which is incorporated herein by reference. The humanized antibodies of particular interest herein block EGF, TGF- α and/or HRG mediated activation of MAPK essentially as effectively as murine monoclonal antibody 2C4 (or a Fab fragment thereof) and/or binds ErbB2 essentially as effectively as murine monoclonal antibody 2C4 (or a Fab fragment thereof). The humanized antibodies herein may, for example, comprise nonhuman hypervariable region residues incorporated into a human variable heavy domain and may further comprise a framework region (FR) substitution at a position selected from the group consisting of 69H, 71H and 73H utilizing the variable domain numbering system set forth in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991). In one embodiment, the humanized antibody comprises FR substitutions at two or all of positions 69H, 71H and 73H.

An exemplary humanized antibody of interest herein comprises variable heavy domain complementarity determining residues GFTFTDYTMX, where X is preferably D or S (SEQ ID NO:7); DVNPNSGGSIYNQRFKG (SEQ ID NO:8); and/or NLGPSFYFDY (SEQ ID NO:9), optionally comprising amino acid modifications of those CDR residues, *e.g.*, where the modifications essentially maintain or improve affinity of the antibody. For example, the antibody variant of interest may have from about one to about seven or about five amino acid substitutions in the above variable heavy CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below. The most preferred humanized antibody comprises the variable heavy domain amino acid sequence in SEQ ID NO:4 (Figure 1B).

The humanized antibody may comprise variable light domain complementarity determining residues KASQDVSIGVA (SEQ ID NO:10); SASYX1X2X3, where X1 is preferably R or L, X2 is preferably Y or E, and X3 is preferably T or S (SEQ ID NO:11); and/or QQYYIYPYT (SEQ ID NO:12), *e.g.*, in addition to those variable heavy domain CDR residues in the preceding paragraph. Such humanized antibodies optionally comprise amino acid modifications of the above CDR residues, *e.g.*, where the modifications essentially maintain or improve affinity of the antibody. For example, the antibody variant of interest may have from about one to about seven or about five amino acid substitutions in the above variable light CDR sequences. Such antibody variants may be prepared by affinity maturation, *e.g.*, as described below. The most preferred humanized antibody comprises the variable light domain amino acid sequence in SEQ ID NO:3 (Figure 1A).

The present application also contemplates affinity matured antibodies which bind ErbB2 and block ligand activation of an ErbB receptor. The parent antibody may be a human antibody or a humanized antibody, *e.g.*, one comprising the variable light and/or heavy sequences of SEQ ID Nos. 3 and 4, respectively (*i.e.*, variant 574; Figure 1A and B). The affinity matured antibody preferably binds to ErbB2 receptor with an affinity superior to that of murine 2C4 or variant 574 (*e.g.*, from about two or about four fold, to about 100 fold or about 1000 fold improved affinity, *e.g.*, as assessed using a ErbB2-extracellular domain (ECD) ELISA). Exemplary variable heavy CDR residues for substitution include H28, H30, H34, H35, H64, H96, H99, or combinations of two or more (*e.g.*, two, three, four, five, six, or seven of these residues). Examples of variable light CDR residues for alteration include L28, L50, L53, L56, L91, L92, L93, L94, L96, L97 or combinations of two or more (*e.g.*, two to three, four, five or up to about ten of these residues).

Various forms of the humanized antibody or affinity matured antibody are contemplated. For example, the humanized antibody or affinity matured antibody may be an antibody fragment, such as a Fab, which is optionally conjugated with one or more cytotoxic agent(s) in order to generate an immunoconjugate. Alternatively, the humanized antibody or affinity matured antibody may be an intact antibody, such as an intact IgG1 antibody.

20 *Human antibodies*

As an alternative to humanization, human antibodies can be generated. For example, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (JH) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, *e.g.*, Jakobovits et al., *Proc. Natl. Acad. Sci. USA*, 90:2551 (1993); Jakobovits et al., *Nature*, 362:255-258 (1993); Bruggermann et al., *Year in Immuno.*, 7:33 (1993); and U.S. Patent Nos. 5,591,669, 5,589,369 and 5,545,807.

Alternatively, phage display technology (McCafferty et al., *Nature*, 348:552-553 (1990)) can be used to produce human antibodies and antibody fragments in vitro, from immunoglobulin variable (V) domain gene repertoires from unimmunized donors.

According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimics some of the properties of the B-cell. Phage display can be performed in a variety of formats; for their review see, e.g., Johnson, Kevin S. and Chiswell, David J., *Current Opinion in Structural Biology*, 3:564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson et al., *Nature*, 352:624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., *J. Mol. Biol.*, 222:581-597 (1991), or Griffith et al., *EMBO J.*, 12:725-734 (1993). See, also, U.S. Patent Nos. 5,565,332 and 5,573,905.

As discussed above, human antibodies may also be generated by in vitro activated B cells (see U.S. Patents 5,567,610 and 5,229,275).

Human anti-ErbB2 antibodies are described in U.S. Patent No. 5,772,997 issued June 30, 1998 and WO 97/00271 published January 3, 1997.

Antibody fragments

Various techniques have been developed for the production of antibody fragments.

Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., *Journal of Biochemical and Biophysical Methods*, 24:107-117 (1992); and Brennan et al., *Science*, 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from E. coli and chemically coupled to form F(ab')₂ fragments (Carter et al., *Bio/Technology*, 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner. In other embodiments, the antibody of choice is a single chain Fv fragment (scFv). See WO 93/16185; U.S. Patent No. 5,571,894; and U.S. Patent No. 5,587,458. The antibody fragment may also be a □linear antibody□, e.g.,

as described in U.S. Patent 5,641,870 for example. Such linear antibody fragments may be monospecific or bispecific.

Bispecific antibodies

Bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of the ErbB2 protein. Other such antibodies may combine an ErbB2 binding site with binding site(s) for EGFR, ErbB3 and/or ErbB4. Alternatively, an anti-ErbB2 arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g., CD2 or CD3), or Fc receptors for IgG (Fc γ R), such as Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16) so as to focus cellular defense mechanisms to the ErbB2-expressing cell. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express ErbB2. These antibodies possess an ErbB2-binding arm and an arm which binds the cytotoxic agent (e.g., saporin, anti-interferon- α , vinca alkaloid, ricin A chain, methotrexate or radioactive isotope hapten). Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies).

WO 96/16673 describes a bispecific anti-ErbB2/anti-Fc γ RIII antibody and U.S. Patent No. 5,837,234 discloses a bispecific anti-ErbB2/anti-Fc γ RI antibody. A bispecific anti-ErbB2/Fc α antibody is shown in WO 98/02463. U.S. Patent No. 5,821,337 teaches a bispecific anti-ErbB2/anti-CD3 antibody.

Methods for making bispecific antibodies are known in the art. Traditional production of full length bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein et al., *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. Purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in WO 93/08829, and in Traunecker et al., *EMBO J.*, 10:3655-3659 (1991).

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH₂, and CH₃ regions. It is

preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations, as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in WO 94/04690. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

According to another approach described in U.S. Patent No. 5,731,168, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 domain of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g., tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g., alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known

in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

Techniques for generating bispecific antibodies from antibody fragments have also been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science*, 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Recent progress has facilitated the direct recovery of Fab'-SH fragments from *E. coli*, which can be chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.*, 175: 217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.*, 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (VH) connected to a light-chain variable domain (VL) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the VH and VL domains of one fragment are forced to pair with the complementary VL and VH domains of another fragment, thereby forming two antigen-binding sites. Another strategy

for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See Gruber et al., *J. Immunol.*, 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al. *J. Immunol.*, 147:60 (1991).

5 *Other amino acid sequence modifications*

Amino acid sequence modification(s) of the anti-ErbB2 antibodies are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibodies. Amino acid sequence variants of the anti-ErbB2 antibodies are prepared by introducing appropriate nucleotide changes into the
10 anti-ErbB2 antibody nucleic acid, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the anti-ErbB2 antibody. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes also may alter
15 post-translational processes of the anti-ErbB2 antibodies, such as changing the number or position of glycosylation sites.

A useful method for identification of certain residues or regions of the anti-ErbB2 antibodies that are preferred locations for mutagenesis is called "alanine scanning mutagenesis" as described by Cunningham and Wells, *Science*, 244:1081-1085 (1989).
20 Here, a residue or group of target residues are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with ErbB2 antigen. Those amino acid locations demonstrating functional sensitivity to the substitutions then are refined by introducing further or other variants at, or for, the sites of
25 substitution. Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation per se need not be predetermined. For example, to analyze the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed anti-ErbB2 antibody variants are screened for the desired activity.

30 Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an anti-ErbB2 antibody with an N-terminal methionyl residue or the antibody fused to a cytotoxic polypeptide. Other insertional variants of the anti-

ErbB2 antibody molecules include the fusion to the N- or C-terminus of the anti-ErbB2 antibodies to a reporter molecule, an enzyme (e.g. for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

Another type of variant is an amino acid substitution variant. These variants have at least one amino acid residue in the anti-ErbB2 antibody molecule replaced by a different residue. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. Conservative substitutions are shown in Table 1 under the heading of "preferred substitutions." If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Table 1

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; asp, lys; arg	gln
Asp (D)	glu; asn	glu
Cys (C)	ser; ala	ser
Gln (Q)	asn; glu	asn
Glu (E)	asp; gln	asp
Gly (G)	Ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	tyr
Pro (P)	Ala	ala
Ser (S)	Thr	thr
Thr (T)	Ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- 5 (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

10 Any cysteine residue not involved in maintaining the proper conformation of the anti-ErbB2 antibody also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the antibody to improve its stability (particularly where the antibody is an antibody fragment such as an Fv fragment).

15 A particularly preferred type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (*e.g.*, a humanized or human antibody). Generally, the resulting variant(s) selected for further development will have improved biological properties relative to the parent antibody from which they are generated. A convenient way for generating such substitutional variants involves affinity maturation using phage display. Briefly, several hypervariable region sites (*e.g.*, 6-7 sites)
20 are mutated to generate all possible amino substitutions at each site. The antibody variants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed variants are then screened for their biological activity (*e.g.*, binding affinity) as herein disclosed. In order to identify candidate hypervariable region sites for
25 modification, alanine scanning mutagenesis can be performed to identify hypervariable region residues contributing significantly to antigen binding. Alternatively, or additionally, it may be beneficial to analyze a crystal structure of the antigen-antibody complex to identify contact points between the antibody and human ErbB2. Such contact residues and neighboring residues are candidates for substitution according to the
30 techniques elaborated herein. Once such variants are generated, the panel of variants is subjected to screening as described herein and antibodies with superior properties in one or more relevant assays may be selected for further development.

Another type of amino acid variation alters the original glycosylation pattern of the antibody. By altering is meant deleting one or more carbohydrate moieties found in the

antibody, and/or adding one or more glycosylation sites that are not present in the antibody.

Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original antibody (for O-linked glycosylation sites).

Nucleic acid molecules encoding amino acid sequence variants of the anti-ErbB2 antibody are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the anti-ErbB2 antibody.

It may be desirable to modify the antibodies of the invention with respect to effector function, *e.g.*, so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp Med.*, 176:1191-1195 (1992) and Shopes, B., *J. Immunol.*, 148:2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al., *Cancer Research*, 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis

and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3:219-230 (1989).

To increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an antibody fragment) as described in U.S. Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (e.g., IgG1, IgG2, IgG3, or IgG4) that is responsible for increasing the in vivo serum half-life of the IgG molecule.

Screening for antibodies with the desired properties

Techniques for generating antibodies have been described above. One may further select antibodies with certain biological characteristics, as desired.

To identify an antibody which blocks ligand activation of an ErbB receptor, the ability of the antibody to block ErbB ligand binding to cells expressing the ErbB receptor (e.g., in conjugation with another ErbB receptor with which the ErbB receptor of interest forms an ErbB hetero-oligomer) may be determined. For example, cells naturally expressing, or transfected to express, ErbB receptors of the ErbB hetero-oligomer may be incubated with the antibody and then exposed to labeled ErbB ligand. The ability of the anti-ErbB2 antibody to block ligand binding to the ErbB receptor in the ErbB hetero-oligomer may then be evaluated.

For example, inhibition of HRG binding to MCF7 breast tumor cell lines by anti-ErbB2 antibodies may be performed using monolayer MCF7 cultures on ice in a 24-well-plate format essentially as described in WO 01/00245. Anti-ErbB2 monoclonal antibodies may be added to each well and incubated for 30 minutes. ¹²⁵I-labeled rHRG β 1177-224 (25 pm) may then be added, and the incubation may be continued for 4 to 16 hours. Dose response curves may be prepared and an IC₅₀ value may be calculated for the antibody of interest. In one embodiment, the antibody which blocks ligand activation of an ErbB receptor will have an IC₅₀ for inhibiting HRG binding to MCF7 cells in this assay of about 50nM or less, more preferably 10nM or less. Where the antibody is an antibody fragment such as a Fab fragment, the IC₅₀ for inhibiting HRG binding to MCF7 cells in this assay may, for example, be about 100nM or less, more preferably 50nM or less.

Alternatively, or additionally, the ability of the anti-ErbB2 antibody to block ErbB ligand-stimulated tyrosine phosphorylation of an ErbB receptor present in an ErbB hetero-oligomer may be assessed. For example, cells endogenously expressing the ErbB receptors or transfected to express them may be incubated with the antibody and then assayed for ErbB ligand-dependent tyrosine phosphorylation activity using an anti-

phosphotyrosine monoclonal (which is optionally conjugated with a detectable label). The kinase receptor activation assay described in U.S. Patent No. 5,766,863 is also available for determining ErbB receptor activation and blocking of that activity by an antibody.

In one embodiment, one may screen for an antibody which inhibits HRG stimulation of p180 tyrosine phosphorylation in MCF7 cells essentially as described in Example 1 below. For example, the MCF7 cells may be plated in 24-well plates and monoclonal antibodies to ErbB2 may be added to each well and incubated for 30 minutes at room temperature; then rHRG β 1177-244 may be added to each well to a final concentration of 0.2 nM, and the incubation may be continued for 8 minutes. Media may be aspirated from each well, and reactions may be stopped by the addition of 100 μ l of SDS sample buffer (5% SDS, 25 mM DTT, and 25 mM Tris-HCl, pH 6.8). Each sample (25 μ l) may be electrophoresed on a 4-12% gradient gel (Novex) and then electrophoretically transferred to polyvinylidene difluoride membrane. Antiphosphotyrosine (at 1 μ g/ml) immunoblots may be developed, and the intensity of the predominant reactive band at Mr ~180,000 may be quantified by reflectance densitometry. The antibody selected will preferably significantly inhibit HRG stimulation of p180 tyrosine phosphorylation to about 0-35% of control in this assay. A dose-response curve for inhibition of HRG stimulation of p180 tyrosine phosphorylation as determined by reflectance densitometry may be prepared and an IC50 for the antibody of interest may be calculated. In one embodiment, the antibody which blocks ligand activation of an ErbB receptor will have an IC50 for inhibiting HRG stimulation of p180 tyrosine phosphorylation in this assay of about 50nM or less, more preferably 10nM or less. Where the antibody is an antibody fragment such as a Fab fragment, the IC50 for inhibiting HRG stimulation of p180 tyrosine phosphorylation in this assay may, for example, be about 100nM or less, more preferably 50nM or less.

One may also assess the growth inhibitory effects of the antibody on MDA-MB-175 cells, *e.g.*, essentially as described in Schaefer et al. *Oncogene*, 15:1385-1394 (1997). According to this assay, MDA-MB-175 cells may be treated with an anti-ErbB2 monoclonal antibody (10 μ g/mL) for 4 days and stained with crystal violet. Incubation with an anti-ErbB2 antibody may show a growth inhibitory effect on this cell line similar to that displayed by monoclonal antibody 2C4. In a further embodiment, exogenous HRG will not significantly reverse this inhibition. Preferably, the antibody will be able to inhibit cell proliferation of MDA-MB-175 cells to a greater extent than monoclonal antibody 4D5 (and optionally to a greater extent than monoclonal antibody 7F3), both in the presence and absence of exogenous HRG.

In one embodiment, the anti-ErbB2 antibody of interest may block heregulin dependent association of ErbB2 with ErbB3 in both MCF7 and SK-BR-3 cells as determined in a co-immunoprecipitation experiment such as that described in Example 1 substantially more effectively than monoclonal antibody 4D5, and preferably substantially more effectively than monoclonal antibody 7F3.

To identify growth inhibitory anti-ErbB2 antibodies, one may screen for antibodies which inhibit the growth of cancer cells which overexpress ErbB2. In one embodiment, the growth inhibitory antibody of choice is able to inhibit growth of SK-BR-3 cells in cell culture by about 20-100% and preferably by about 50-100% at an antibody concentration of about 0.5 to 30 $\mu\text{g/ml}$. To identify such antibodies, the SK-BR-3 assay described in U.S. Patent No. 5,677,171 can be performed. According to this assay, SK-BR-3 cells are grown in a 1:1 mixture of F12 and DMEM medium supplemented with 10% fetal bovine serum, glutamine and penicillin streptomycin. The SK-BR-3 cells are plated at 20,000 cells in a 35mm cell culture dish (2mls/35mm dish). 0.5 to 30 $\mu\text{g/ml}$ of the anti-ErbB2 antibody is added per dish. After six days, the number of cells, compared to untreated cells are counted using an electronic COULTER™ cell counter. Those antibodies which inhibit growth of the SK-BR-3 cells by about 20-100% or about 50-100% may be selected as growth inhibitory antibodies.

To select for antibodies which induce cell death, loss of membrane integrity as indicated by, *e.g.*, PI, trypan blue or 7AAD uptake may be assessed relative to control. The preferred assay is the PI uptake assay using BT474 cells. According to this assay, BT474 cells (which can be obtained from the American Type Culture Collection (Rockville, MD)) are cultured in Dulbecco's Modified Eagle Medium (D-MEM):Ham's F-12 (50:50) supplemented with 10% heat-inactivated FBS (Hyclone) and 2 mM L-glutamine. (Thus, the assay is performed in the absence of complement and immune effector cells). The BT474 cells are seeded at a density of 3×10^6 per dish in 100 x 20 mm dishes and allowed to attach overnight. The medium is then removed and replaced with fresh medium alone or medium containing 10 $\mu\text{g/ml}$ of the appropriate monoclonal antibody. The cells are incubated for a 3 day time period. Following each treatment, monolayers are washed with PBS and detached by trypsinization. Cells are then centrifuged at 1200 rpm for 5 minutes at 40C, the pellet resuspended in 3 ml ice cold Ca²⁺ binding buffer (10 mM HEPES, pH 7.4, 140 mM NaCl, 2.5 mM CaCl₂) and aliquoted into 35 mm strainer-capped 12 x 75 tubes (1 ml per tube, 3 tubes per treatment group) for removal of cell clumps. Tubes then receive PI (10 $\mu\text{g/ml}$). Samples may be analyzed using a FACSCAN™ flow cytometer and FACSCONVERT™ CellQuest software

(Becton Dickinson). Those antibodies which induce statistically significant levels of cell death as determined by PI uptake may be selected as cell death-inducing antibodies.

In order to select for antibodies which induce apoptosis, an annexin binding assay using BT474 cells is available. The BT474 cells are cultured and seeded in dishes as discussed in the preceding paragraph. The medium is then removed and replaced with fresh medium alone or medium containing 10 $\mu\text{g/ml}$ of the monoclonal antibody. Following a three day incubation period, monolayers are washed with PBS and detached by trypsinization. Cells are then centrifuged, resuspended in Ca^{2+} binding buffer and aliquoted into tubes as discussed above for the cell death assay. Tubes then receive labeled annexin (*e.g.*, annexin V-FTIC) (1 $\mu\text{g/ml}$). Samples may be analyzed using a FACSCAN™ flow cytometer and FACSCONVERT™ CellQuest software (Becton Dickinson). Those antibodies which induce statistically significant levels of annexin binding relative to control are selected as apoptosis-inducing antibodies.

In addition to the annexin binding assay, a DNA staining assay using BT474 cells is available. In order to perform this assay, BT474 cells which have been treated with the antibody of interest as described in the preceding two paragraphs are incubated with 9 $\mu\text{g/ml}$ HOECHST 33342™ for 2 hr at 37°C, then analyzed on an EPICS ELITE™ flow cytometer (Coulter Corporation) using MODFIT LT™ software (Verity Software House). Antibodies which induce a change in the percentage of apoptotic cells which is 2 fold or greater (and preferably 3 fold or greater) than untreated cells (up to 100% apoptotic cells) may be selected as pro-apoptotic antibodies using this assay.

To screen for antibodies which bind to an epitope on ErbB2 bound by an antibody of interest, a routine cross-blocking assay such as that described in Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory, Ed Harlow and David Lane (1988), can be performed. Alternatively, or additionally, epitope mapping can be performed by methods known in the art (see, *e.g.*, Figs. 1A and 1B herein).

Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (*e.g.* a small molecule toxin or an enzymatically active toxin of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof), or a radioactive isotope (*i.e.*, a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Conjugates of an antibody and one or more small molecule toxins,

such as a calicheamicin, a maytansine (U.S. Patent No. 5,208,020), a trichothene, and CC1065 are also contemplated herein.

In one preferred embodiment of the invention, an antibody is conjugated to one or more maytansine molecules (*e.g.*, about 1 to about 10 maytansine molecules per antibody molecule). Maytansine may, for example, be converted to May-SS-Me which may be reduced to May-SH3 and reacted with modified antibody (Chari et al., *Cancer Research*, 52:127-131 (1992)) to generate a maytansinoid-antibody immunoconjugate.

Another immunoconjugate of interest comprises an anti-ErbB2 antibody conjugated to one or more calicheamicin molecules. The calicheamicin family of antibiotics are capable of producing double-stranded DNA breaks at sub-picomolar concentrations. Structural analogues of calicheamicin which may be used include, but are not limited to, γ 1I, α 2I, α 3I, N-acetyl- γ 1I, PSAG and θ 1I (Hinman et al., *Cancer Research*, 53:3336-3342 (1993) and Lode et al., *Cancer Research*, 58:2925-2928 (1998)). See, also, U.S. Patent Nos. 5,714,586; 5,712,374; 5,264,586; and 5,773,001 expressly incorporated herein by reference.

Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, *sapaonaria officinalis* inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. See, for example, WO 93/21232 published October 28, 1993.

The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (*e.g.*, a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

A variety of radioactive isotopes are available for the production of radioconjugated anti-ErbB2 antibodies. Examples include At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32 and radioactive isotopes of Lu.

Conjugates of an antibody and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as

toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science*, 238:1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO 94/11026. The linker may be a □cleavable linker□ facilitating release of the cytotoxic drug in the cell. For example, an acid-labile linker, peptidase-sensitive linker, dimethyl linker or disulfide-containing linker (Chari et al., *Cancer Research*, 52:127-131 (1992)) may be used.

Alternatively, a fusion protein comprising an anti-ErbB2 antibody and cytotoxic agent may be made, e.g., by recombinant techniques or peptide synthesis.

In yet another embodiment, an antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionucleotide).

Antibody Dependent Enzyme Mediated Prodrug Therapy (ADEPT)

The antibodies of the present invention may also be used in ADEPT by conjugating the antibody to a prodrug-activating enzyme which converts a prodrug (e.g., a peptidyl chemotherapeutic agent, see WO 81/01145) to an active anti-cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278.

The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form.

Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or

penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes," can be used to convert the prodrugs of the invention into free active drugs (see, *e.g.*, Massey, *Nature*, 5 328:457-458 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes of this invention can be covalently bound to the anti-ErbB2 antibodies by techniques well known in the art such as the use of the heterobifunctional crosslinking reagents discussed above. Alternatively, fusion proteins comprising at least 10 the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using recombinant DNA techniques well known in the art (see, *e.g.*, Neuberger et al., *Nature*, 312:604-608 (1984)).

Other antibody modifications

Other modifications of the antibodies are contemplated herein. For example, the 15 antibody may be linked to one of a variety of nonproteinaceous polymers, *e.g.*, polyethylene glycol, polypropylene glycol, polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol. The antibody may also or alternatively be linked to one or more of a variety of different moieties, such as a fluorescent label, a moiety with a known electrophoretic mobility, or a moiety that is able to cleave a specific 20 linker molecule.

The antibody also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, 25 albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, 16th edition, Oslo, A., Ed., (1980).

The anti-ErbB2 antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in 30 the art, such as described in Epstein et al., *Proc. Natl. Acad. Sci. USA*, 82:3688 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA*, 77:4030 (1980); U.S. Patent Nos. 4,485,045 and 4,544,545; and WO 97/38731 published October 23, 1997. Liposomes with enhanced circulation time are disclosed in U.S. Patent No. 5,013,556.

Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in
5 Martin et al., *J. Biol. Chem.*, 257:286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent is optionally contained within the liposome. See Gabizon et al., *J. National Cancer Inst.*, 81(19):1484 (1989).

Vectors, Host Cells and Recombinant Methods

10 The invention also provides isolated nucleic acid encoding the antibodies, including humanized anti-ErbB2 antibodies, vectors and host cells comprising the nucleic acid, and recombinant techniques for the production of the antibody. For recombinant production of an antibody, the nucleic acid encoding it is isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for
15 expression. DNA encoding the monoclonal antibody is readily isolated and sequenced using conventional procedures (*e.g.*, by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). Many vectors are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker
20 genes, an enhancer element, a promoter, and a transcription termination sequence.

Signal sequence component

The anti-ErbB2 antibodies may be produced recombinantly not only directly, but also as fusion polypeptides with a heterologous polypeptide, which is preferably a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the
25 mature protein or polypeptide. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells that do not recognize and process the native anti-ErbB2 antibody signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase,
30 lpp, or heat-stable enterotoxin II leaders. For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader, α factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders), or acid phosphatase leader, the *C. albicans* glucoamylase leader, or the signal described in WO 90/13646. In mammalian

cell expression, mammalian signal sequences as well as viral secretory leaders, for example, the herpes simplex gD signal, are available.

The DNA for such precursor region is ligated in reading frame to DNA encoding the anti-ErbB2 antibody.

5 *Origin of replication component*

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

Selection gene component

Expression and cloning vectors may contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for Bacilli.

One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin, mycophenolic acid and hygromycin.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the anti-ErbB2 antibody nucleic acid, such as DHFR, thymidine kinase, metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, etc.

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity.

Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding anti-ErbB2 antibody, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, *e.g.*, kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 (Stinchcomb et al., *Nature*, 282:39 (1979)). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1. Jones, *Genetics*, 85:12 (1977). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Alternatively, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis*. Van den Berg, *Bio/Technology*, 8:135 (1990). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed. Fleer et al., *Bio/Technology*, 9:968-975 (1991).

Promoter component

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the anti-ErbB2 antibody nucleic acid. Promoters suitable for use with prokaryotic hosts include the *phoA* promoter, β -lactamase and lactose promoter systems, alkaline phosphatase, a tryptophan (*trp*) promoter system, and hybrid promoters such as the *tac* promoter. However, other known bacterial promoters are suitable. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding the anti-ErbB2 antibody.

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CNCAAT region where N may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be the signal for

addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

5 Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

10 Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657. Yeast enhancers also are advantageously used with yeast promoters.

15 Anti-ErbB2 antibody transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the action promoter or an immunoglobulin promoter, from heat-shock promoters, provided such promoters are compatible with the host cell systems.

20 The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. The immediate early promoter of the human cytomegalovirus is conveniently obtained as a HindIII E restriction fragment. A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Reyes et al., *Nature*, 297:598-601 (1982) on expression of human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus.

30 Alternatively, the rous sarcoma virus long terminal repeat can be used as the promoter.

Enhancer element component

Transcription of a DNA encoding the anti-ErbB2 antibody of this invention by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Many enhancer sequences are now known from mammalian genes (globin, elastase,

albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

- 5 See also Yaniv, *Nature*, 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the anti-ErbB2 antibody-encoding sequence, but is preferably located at a site 5' from the promoter.

Transcription termination component

- 10 Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments
- 15 transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding anti-ErbB2 antibody. One useful transcription termination component is the bovine growth hormone polyadenylation region. See WO94/11026 and the expression vector disclosed therein.

Selection and transformation of host cells

- 20 Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as Escherichia, *e.g.*, E. coli, Enterobacter, Erwinia, Klebsiella, Proteus, Salmonella, *e.g.*, Salmonella typhimurium, Serratia, *e.g.*, Serratia
- 25 marcescans, and Shigella, as well as Bacilli such as B. subtilis and B. licheniformis (*e.g.*, B. licheniformis 41P disclosed in DD 266,710 published 12 April 1989), Pseudomonas such as P. aeruginosa, and Streptomyces. One preferred E. coli cloning host is E. coli 294 (ATCC 31,446), although other strains such as E. coli B, E. coli X1776 (ATCC 31,537), and E. coli W3110 (ATCC 27,325) are suitable. These examples are illustrative rather
- 30 than limiting.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for anti-ErbB2 antibody-encoding vectors. Saccharomyces cerevisiae, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and

strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*; Kluyveromyces hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070); *Candida*; *Trichoderma reesei* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolyposcladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

Suitable host cells for the expression of glycosylated anti-ErbB2 antibody are derived from multicellular organisms. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts.

However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., *Annals N.Y. Acad. Sci.*, 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transformed with the above-described expression or cloning vectors for anti-ErbB2 antibody production and cultured in conventional nutrient media modified

as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences.

Culturing the host cells

The host cells used to produce the anti-ErbB2 antibody of this invention may be
5 cultured in a variety of media. Commercially available media such as Ham's F10
(Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and
Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the
host cells. In addition, any of the media described in Ham et al., *Meth. Enz.*, 58:44 (1979),
Barnes et al., *Anal. Biochem.*, 102:255 (1980), U.S. Patent Nos. 4,767,704; 4,657,866;
10 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re.
30,985 may be used as culture media for the host cells. Any of these media may be
supplemented as necessary with hormones and/or other growth factors (such as insulin,
transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium,
magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and
15 thymidine), antibiotics (such as GENTAMYCIN® drug), trace elements (defined as
inorganic compounds usually present at final concentrations in the micromolar range), and
glucose or an equivalent energy source. Any other necessary supplements may also be
included at appropriate concentrations that would be known to those skilled in the art. The
culture conditions, such as temperature, pH, and the like, are those previously used with
20 the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

Purification of anti-ErbB2 antibody

When using recombinant techniques, antibodies can be produced intracellularly, in
the periplasmic space, or directly secreted into the medium. If the antibody is produced
intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, is
25 removed, for example, by centrifugation or ultrafiltration. Carter et al., *Bio/Technology*,
10:163-167 (1992) describe a procedure for isolating antibodies which are secreted to the
periplasmic space of *E. coli*. Briefly, cell paste is thawed in the presence of sodium
acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min.
Cell debris can be removed by centrifugation. Where the antibody is secreted into the
30 medium, supernatants from such expression systems are generally first concentrated using
a commercially available protein concentration filter, for example, an Amicon or Millipore
Pellicon ultrafiltration unit. A protease inhibitor such as PMSF may be included in any of
the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the
growth of adventitious contaminants.

The antibody composition prepared from the cells can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fc domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark et al., *J. Immunol. Meth.*, 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss et al., *EMBO J.*, 5:15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a CH3 domain, the Bakerbond ABXTMresin (J. T. Baker, Phillipsburg, NJ) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSETM chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered.

Following any preliminary purification step(s), the mixture comprising the antibody of interest and contaminants may be subjected to low pH hydrophobic interaction chromatography using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g., from about 0-0.25M salt).

Pharmaceutical Formulations

Therapeutic formulations of the antibodies used in accordance with the present invention are prepared for storage by mixing an antibody having the desired degree of purity with optional pharmaceutically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular

weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG). Preferred lyophilized anti-ErbB2 antibody formulations are described in WO 97/04801, expressly incorporated herein by reference.

The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide antibodies which bind to EGFR, ErbB2 (e.g., an antibody which binds a different epitope on ErbB2), ErbB3, ErbB4, or vascular endothelial factor (VEGF) in the one formulation. Alternatively, or additionally, the composition may further comprise a chemotherapeutic agent, cytotoxic agent, cytokine, growth inhibitory agent, anti-hormonal agent, EGFR-targeted drug, anti-angiogenic agent, and/or cardioprotectant. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic

acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid.

The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

5 **Treatment with Anti-ErbB2 Antibodies**

It is contemplated that, according to the present invention, anti-ErbB2 antibodies may be used to treat various diseases or disorders. Exemplary conditions or disorders include benign or malignant tumors; leukemias and lymphoid malignancies; other disorders such as neuronal, glial, astrocytal, hypothalamic, glandular, macrophagal, epithelial, stromal, blastocoelic, inflammatory, angiogenic and immunologic disorders. Preferably, anti-ErbB2 antibodies are used to treat tumors that are identified as responsive to treatment with such antibodies by the methods disclosed herein.

Generally, the disease or disorder to be treated is cancer. Examples of cancer to be treated herein include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia or lymphoid malignancies. More particular examples of such cancers include squamous cell cancer (*e.g.*, epithelial squamous cell cancer), lung cancer including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, as well as head and neck cancer. Preferably, the cancer to be treated is identified as responsive to treatment with anti-ErbB2 antibodies based on the identification of HER2/HER3 and/or HER2/HER1 heterodimers or the phosphorylation of ErbB receptor in a tumor sample. A particular group of cancers where HER2/HER3 and/or HER2/HER1 heterodimer formation and/or phosphorylation of ErbB receptor is expected to be detected includes, without limitation lung breast cancer, lung cancer, ovarian cancer, including advanced, refractory or recurrent ovarian cancer, prostate cancer, colorectal cancer, and pancreatic cancer.

The cancer will generally comprise ErbB2-expressing cells, such that the anti-ErbB2 antibody herein is able to bind to the cancer. While the cancer may be characterized by overexpression of the ErbB2 receptor, the present application further provides a method for treating cancer which is not considered to be an ErbB2-

overexpressing cancer. To determine ErbB2 expression in the cancer, various diagnostic/prognostic assays are available. In one embodiment, ErbB2 overexpression may be analyzed by IHC, e.g. using the HERCEPTEST[®] (Dako). Paraffin embedded tissue sections from a tumor biopsy may be subjected to the IHC assay and accorded a

5 ErbB2 protein staining intensity criteria as follows:

Score 0

no staining is observed or membrane staining is observed in less than 10% of tumor cells.

Score 1+

10 a faint/barely perceptible membrane staining is detected in more than 10% of the tumor cells. The cells are only stained in part of their membrane.

Score 2+

a weak to moderate complete membrane staining is observed in more than 10% of the tumor cells.

Score 3+

15 a moderate to strong complete membrane staining is observed in more than 10% of the tumor cells.

Those tumors with 0 or 1+ scores for ErbB2 overexpression assessment may be characterized as not overexpressing ErbB2, whereas those tumors with 2+ or 3+ scores may be characterized as overexpressing ErbB2.

20 Alternatively, or additionally, FISH assays such as the INFORM[™] (sold by Ventana, Arizona) or PATHVISION[™] (Vysis, Illinois) may be carried out on formalin-fixed, paraffin-embedded tumor tissue to determine the extent (if any) of ErbB2 overexpression in the tumor.

In one embodiment, the cancer will be one which expresses (and may, but does not

25 have to, overexpress) EGFR. Examples of cancers which may express/overexpress EGFR include squamous cell cancer (e.g., epithelial squamous cell cancer), lung cancer including small-cell lung cancer, non-small cell lung cancer (NSCLC), adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer,

30 glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, as well as head and neck cancer.

The present invention is specifically suitable for the identification of breast cancer, prostate cancer, such as Castration-Resistant Prostate Cancer (CRPC), and ovarian cancer patients that are likely to respond well to treatment with an anti-HER2 antibody that blocks ligands activation of an ErbB heterodimer comprising HER2, such as monoclonal antibody 2C4 or rhuMAb 2C4.

The cancer to be treated herein may be one characterized by excessive activation of an ErbB receptor, e.g. EGFR. Such excessive activation may be attributable to overexpression or increased production of the ErbB receptor or an ErbB ligand. In one embodiment of the invention, a diagnostic or prognostic assay will be performed to determine whether the patient's cancer is characterized by excessive activation of an ErbB receptor. For example, ErbB gene amplification and/or overexpression of an ErbB receptor in the cancer may be determined. Various assays for determining such amplification/overexpression are available in the art and include the IHC, FISH and shed antigen assays described above. Alternatively, or additionally, levels of an ErbB ligand, such as TGF- α , in or associated with the tumor may be determined according to known procedures. Such assays may detect protein and/or nucleic acid encoding it in the sample to be tested. In one embodiment, ErbB ligand levels in the tumor may be determined using immunohistochemistry (IHC); see, for example, Scher et al., *Clin. Cancer Research*, 1:545-550 (1995). Alternatively, or additionally, one may evaluate levels of ErbB ligand-encoding nucleic acid in the sample to be tested; e.g., via FISH, southern blotting, or PCR techniques.

Moreover, ErbB receptor or ErbB ligand overexpression or amplification may be evaluated using an in vivo diagnostic assay, e.g., by administering a molecule (such as an antibody) which binds the molecule to be detected and is tagged with a detectable label (e.g., a radioactive isotope) and externally scanning the patient for localization of the label.

Where the cancer to be treated is hormone independent cancer, expression of the hormone (e.g., androgen) and/or its cognate receptor in the tumor may be assessed using any of the various assays available, e.g., as described above. Alternatively, or additionally, the patient may be diagnosed as having hormone independent cancer in that they no longer respond to anti-androgen therapy.

In certain embodiments, an immunoconjugate comprising the anti-ErbB2 antibody conjugated with a cytotoxic agent is administered to the patient. Preferably, the immunoconjugate and/or ErbB2 protein to which it is bound is/are internalized by the cell, resulting in increased therapeutic efficacy of the immunoconjugate in killing the cancer cell to which it binds. In a preferred embodiment, the cytotoxic agent targets or interferes

with nucleic acid in the cancer cell. Examples of such cytotoxic agents include maytansinoids, calicheamicins, ribonucleases and DNA endonucleases.

In a particular embodiment, the antibody administered is rhuMAb 2C4, or a functional equivalent thereof. RhuMAb 2C4 is a humanized monoclonal antibody based on human IgG1 framework sequences and consisting of two heavy chains (449 residues) and two light chains (214 residues). RhuMAb 2C4 differs significantly from another anti-HER2 antibody HERCEPTIN® (Trastuzumab) in the epitope-binding regions of the light chain and heavy chain. As a result, rhuMAb 2C4 binds to a completely different epitope on HER2. The present invention provides sensitive methods for identifying cancers responsive to treatment with rhuMAb 2C4 or functional equivalents thereof. It is noted that such cancers responsive to rhuMAb 2C4 treatment are not required to overexpress HER2.

The anti-ErbB2 antibodies or immunoconjugates are administered to a human patient in accord with known methods, such as intravenous administration, *e.g.*, as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Intravenous or subcutaneous administration of the antibody is preferred.

Other therapeutic regimens may be combined with the administration of the anti-ErbB2 antibody. The combined administration includes coadministration, using separate formulations or a single pharmaceutical formulation, and consecutive administration in either order, wherein preferably there is a time period while both (or all) active agents simultaneously exert their biological activities.

In one particular embodiment, the patient is treated with two different anti-ErbB2 antibodies. For example, the patient may be treated with a first anti-ErbB2 antibody which blocks ligand activation of an ErbB receptor or an antibody having a biological characteristic of monoclonal antibody 2C4 as well as a second anti-ErbB2 antibody which is growth inhibitory (*e.g.* HERCEPTIN®) or an anti-ErbB2 antibody which induces apoptosis of an ErbB2-overexpressing cell (*e.g.*, 7C2, 7F3 or humanized variants thereof). Preferably such combined therapy results in a synergistic therapeutic effect. One may, for instance, treat the patient with HERCEPTIN® and thereafter treat with rhuMAb 2C4, *e.g.*, where the patient does not respond to HERCEPTIN® therapy. In another embodiment, the patient may first be treated with rhuMAb 2C4 and then receive HERCEPTIN® therapy. In yet a further embodiment, the patient may be treated with both rhuMAb 2C4 and HERCEPTIN® simultaneously.

It may also be desirable to combine administration of the anti-ErbB2 antibody or antibodies, with administration of an antibody directed against another tumor associated antigen. The other antibody in this case may, for example, bind to EGFR, ErbB3, ErbB4, or vascular endothelial growth factor (VEGF).

5 In one embodiment, the treatment of the present invention involves the combined administration of an anti-ErbB2 antibody (or antibodies) and one or more chemotherapeutic agents or growth inhibitory agents, including coadministration of cocktails of different chemotherapeutic agents. Preferred chemotherapeutic agents include taxanes (such as paclitaxel and docetaxel) and/or anthracycline antibiotics. Preparation and dosing schedules for such chemotherapeutic agents may be used according to
10 manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in Chemotherapy Service Ed., M.C. Perry, Williams & Wilkins, Baltimore, MD (1992).

The antibody may be combined with an anti-hormonal compound; *e.g.*, an anti-estrogen compound such as tamoxifen; an anti-progesterone such as onapristone (see, EP
15 616 812); or an anti-androgen such as flutamide, in dosages known for such molecules. Where the cancer to be treated is hormone independent cancer, the patient may previously have been subjected to anti-hormonal therapy and, after the cancer becomes hormone independent, the anti-ErbB2 antibody (and optionally other agents as described herein)
20 may be administered to the patient.

Sometimes, it may be beneficial to also coadminister a cardioprotectant (to prevent or reduce myocardial dysfunction associated with the therapy) or one or more cytokines to the patient. One may also coadminister an EGFR- targeted drug or an anti-angiogenic agent. In addition to the above therapeutic regimes, the patient may be subjected to
25 surgical removal of cancer cells and/or radiation therapy.

The anti-ErbB2 antibodies herein may also be combined with an EGFR-targeted drug such as those discussed above in the definitions section resulting in a complementary, and potentially synergistic, therapeutic effect.

Examples of additional drugs which can be combined with the antibody include
30 chemotherapeutic agents such as carboplatin, a taxane (*e.g.*, paclitaxel or docetaxel), gemcitabine, navelbine, cisplatin, oxaliplatin, or combinations of any of these such as carboplatin/docetaxel; another anti-HER2 antibody (*e.g.*, a growth inhibitory anti-HER2 antibody such as HERCEPTIN®, or an anti-HER2 antibody which induces apoptosis such as 7C2 or 7F3, including humanized or affinity matured variants thereof); a farnesyl
35 transferase inhibitor; an anti-angiogenic agent (*e.g.*, an anti-VEGF antibody); an EGFR-

targeted drug (e.g., C225 or ZD1839); a cytokine (e.g., IL-2, IL-12, G-CSF or GM-CSF); or combinations of the above.

Suitable dosages for any of the above coadministered agents are those presently used and may be lowered due to the combined action (synergy) of the agent and anti-
5 ErbB2 antibody.

For the prevention or treatment of disease, the appropriate dosage of antibody will depend on the type of disease to be treated, as defined above, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the
10 discretion of the attending physician. The antibody is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 $\mu\text{g}/\text{kg}$ to 15 mg/kg (e.g. 0.1-20mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. A typical daily dosage might range from about
15 1 $\mu\text{g}/\text{kg}$ to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. The preferred dosage of the antibody will be in the range from about 0.05mg/kg to about 10mg/kg. Thus, one or more doses of about 0.5mg/kg, 2.0mg/kg, 4.0mg/kg or 10mg/kg (or any combination
20 thereof) may be administered to the patient. Such doses may be administered intermittently, e.g., every week or every three weeks (e.g., such that the patient receives from about two to about twenty, e.g., about six doses of the anti-ErbB2 antibody). An initial higher loading dose, followed by one or more lower doses may be administered. An exemplary dosing regimen comprises administering an initial loading dose of about 4
25 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the anti-ErbB2 antibody. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

In a particular embodiment, rhuMAb 2C4 is administered in a fixed dose of 420 mg (equivalent to doses of 6 mg/kg for a 70-kg subject) every 3 weeks. Treatment may
30 start with a higher loading dose (e.g., 840 mg, equivalent to 12 mg/kg of body weight) in order to achieve steady state serum concentrations more rapidly. Specific dosing regimens are also provided in the Examples below.

Aside from administration of the antibody protein to the patient, the present application contemplates administration of the antibody by gene therapy. Such
35 administration of nucleic acid encoding the antibody is encompassed by the expression

“administering a therapeutically effective amount of an antibody”. See, for example, WO 96/07321 published March 14, 1996 concerning the use of gene therapy to generate intracellular antibodies.

5 There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient’s cells; *in vivo* and *ex vivo*. For *in vivo* delivery the nucleic acid is injected directly into the patient, usually at the site where the antibody is required. For *ex vivo* treatment, the patient’s cells are removed, the nucleic acid is introduced into these isolated cells and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes which are implanted into the patient
10 (see, *e.g.*, U.S. Patent Nos. 4,892,538 and 5,283,187). There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection,
15 cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and lipid-based systems (useful lipids for lipid-mediated transfer of the
20 gene are DOTMA, DOPE and DC-Chol, for example). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting
25 and/or to facilitate uptake, *e.g.*, capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.*, 262:4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA*, 87:3410-3414
30 (1990). For review of the currently known gene marking and gene therapy protocols see Anderson et al., *Science*, 256:808-813 (1992). See also WO 93/25673 and the references cited therein.

Articles of Manufacture

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided.

The article of manufacture comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an anti-ErbB2 antibody. The label or package insert indicates that the composition is used for treating the condition of choice, such as cancer. In one embodiment, the label or package insert indicates that the composition comprising the antibody which binds ErbB2 can be used to treat a patient suffering from a tumor in which the presence of HER2/HER1 and/or HER2/HER3 and/or HER2/HER4 complexes have been identified, and/or phosphorylation of ErbB receptor has been detected. Moreover, the article of manufacture may comprise (a) a first container with a composition contained therein, wherein the composition comprises a first antibody which binds ErbB2 and inhibits growth of cancer cells which overexpress ErbB2; and (b) a second container with a composition contained therein, wherein the composition comprises a second antibody which binds ErbB2 and blocks ligand activation of an ErbB receptor. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the first and second antibody compositions can be used to treat cancer characterized by the presence of HER2/HER1 and/or HER2/HER3 and/or HER2/HER4 heterodimers, and/or by the phosphorylation of ErbB receptor. Moreover, the package insert may instruct the user of the composition (comprising an antibody which binds ErbB2 and blocks ligand activation of an ErbB receptor) to combine therapy with the antibody and any of the adjunct therapies described in the preceding section (*e.g.* a chemotherapeutic agent, an EGFR-targeted drug, an anti-angiogenic agent, an anti-hormonal compound, a cardioprotectant and/or a cytokine). Alternatively, or additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

The antibodies may also be used in diagnostic assays. Generally, the antibodies are labeled as described in the methods above. As a matter of convenience, the antibodies of the present invention can be provided in a kit, *i.e.*, a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where
 5 the antibody is labeled with an enzyme, the kit will include substrates and cofactors required by the enzyme (*e.g.*, a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (*e.g.*, a block buffer or lysis buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of
 10 the reagents which substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients which on dissolution will provide a reagent solution having the appropriate concentration.

Deposit of Materials

The following hybridoma cell lines have been deposited with the American Type
 15 Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, USA (ATCC):

Antibody Designation	ATCC No.	Deposit Date
7C2	ATCC HB-12215	October 17, 1996
7F3	ATCC HB-12216	October 17, 1996
20 4D5	ATCC CRL 10463	May 24, 1990
2C4	ATCC HB-12697	April 8, 1999

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the constructs deposited, since the deposited embodiments are intended to
 25 illustrate only certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to
 enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims. Indeed, various modifications of the
 30 invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

It is understood that the application of the teachings of the present invention to a specific problem or situation will be within the capabilities of one having ordinary skill in the art in light of the teachings contained herein.

Further details of the invention are illustrated by the following non-limiting
5 Examples. The disclosures of all citations in the specification are expressly incorporated herein by reference.

Example 1

HRG Dependent Association of ErbB2 with ErbB3 is Blocked by Monoclonal Antibody 2C4

10 The murine monoclonal antibody 2C4, which specifically binds the extracellular domain of ErbB2 is described in WO 01/89566, the disclosure of which is hereby expressly incorporated by reference in its entirety.

The ability of ErbB3 to associate with ErbB2 was tested in a co-immunoprecipitation experiment. 1.0 x 10⁶ MCF7 or SK-BR-3 cells were seeded in six
15 well tissue culture plates in 50:50 DMEM/Ham's F12 medium containing 10% fetal bovine serum (FBS) and 10 mM HEPES, pH 7.2 (growth medium), and allowed to attach overnight. The cells were starved for two hours in growth medium without serum prior to beginning the experiment

The cells were washed briefly with phosphate buffered saline (PBS) and then incubated
20 with either 100 nM of the indicated antibody diluted in 0.2% w/v bovine serum albumin (BSA), RPMI medium, with 10 mM HEPES, pH 7.2 (binding buffer), or with binding buffer alone (control). After one hour at room temperature, HRG was added to a final concentration of 5 nM to half the wells (+). A similar volume of binding buffer was added to the other wells (-). The incubation was continued for approximately 10 minutes.

25 Supernatants were removed by aspiration and the cells were lysed in RPMI, 10 mM HEPES, pH 7.2, 1.0% v/v TRITON X-100™, 1.0% w/v CHAPS (lysis buffer), containing 0.2 mM PMSF, 10 µg/ml leupeptin, and 10 TU/ml aprotinin. The lysates were cleared of insoluble material by centrifugation.

ErbB2 was immunoprecipitated using a monoclonal antibody covalently coupled to
30 an affinity gel (Affi-Prep 10, Bio-Rad). This antibody (Ab-3, Oncogene Sciences, USA) recognizes a cytoplasmic domain epitope. Immunoprecipitation was performed by adding 10 µl of gel slurry containing approximately 8.5 µg of immobilized antibody to each lysate, and the samples were allowed to mix at room temperature for two hours. The gels were then collected by centrifugation. The gels were washed batchwise three times with

lysis buffer to remove unbound material. SDS sample buffer was then added and the samples were heated briefly in a boiling water bath.

Supernatants were run on 4-12% polyacrylamide gels and electroblotted onto nitrocellulose membranes. The presence of ErbB3 was assessed by probing the blots with a polyclonal antibody against a cytoplasmic domain epitope thereof (c-17, Santa Cruz Biotech). The blots were visualized using a chemiluminescent substrate (ECL, Amersham)

As shown in the control lanes of Figs. 2A and 2B, for MCF7 and SK-BR-3 cells, respectively, ErbB3 was present in an ErbB2 immunoprecipitate only when the cells were stimulated with HRG. If the cells were first incubated with monoclonal antibody 2C4, the ErbB3 signal was abolished in MCF7 cells (Fig. 5A, lane 2C4 +) or substantially reduced in SK-BR-3 cells (Fig. 5B, lane 2C4+). As shown in Figs 2A-B, monoclonal antibody 2C4 blocks heregulin dependent association of ErbB3 with ErbB2 in both MCF7 and SK-BR-3 cells substantially more effectively than HERCEPTIN[®]. Preincubation with HERCEPTIN[®] decreased the ErbB3 signal in MCF7 lysates but had little or no effect on the amount of ErbB3 co-precipitated from SK-BR-3 lysates. Preincubation with an antibody against the EGF receptor (Ab-1, Oncogene Sciences, USA) had no effect on the ability of ErbB3 to co-immunoprecipitate with ErbB2 in either cell line.

Example 2

Responsiveness of Cell Line and Human Tumor Xenograft Models to 2C4

Approximately 40 tumor models have been tested for responsiveness to 2C4. These models represent major cancers such as breast, lung, prostate and colon. 50-60% of the models responded to 2C4 treatment. Table 1 below lists selected tumor models tested for responsivity to 2C4. Briefly, human tumor xenograft fragments of about 3 mm size were transplanted underneath the skin of athymic nude mice. Alternatively, human tumor cells grown in vitro were detached from the culture dishes, resuspended in phosphate-buffered saline and subcutaneously injected into the flank of the immuno-compromised mice. The growth of tumors was monitored every 2 to 3 days using an electric caliper. When the tumors reached a size of about 30 to 100 mm, animals were randomized into different treatment and control groups. 2C4 was administered by intraperitoneal injection once every week. Control animals received equal volumes of vehicle solution containing no antibody on the same schedule as the treatment groups. The study was terminated after approximately 3 – 6 weeks, when the tumors of the control group reached a size of about

1000 – 1500 mm³. Responsiveness to treatment was defined as $\geq 50\%$ tumor volume reduction.

Table 1: Xenograft models

5

Model #	Tumor model	Responsiveness to 2C4	Reference
1	LXFA 289	No	(Fiebig et al., 1999)
2	LXFA 297	Yes	
3	LXFA 526	No	(Fiebig et al., 1999)
4	LXFA 629	Yes	(Fiebig and Burger, 2002)
5	LXFA 1041	No	
6	LXFE 211	No	(Fiebig et al., 1999)
7	LXFE 397	No	(Fiebig et al., 1999)
8	LXFL 529	No	(Burger et al., 2001)
9	LXFL 1072	Yes	(Fiebig et al., 1999)
10	Calu-3	Yes	(Stein et al., 2001)
11	NCI-H522	Yes	(Yamori et al., 1997)
12	NCI-H322	No	(Zou et al., 2001)
13	NCI-H441(KAM)	Yes	(Gridley et al., 1996)
14	MAXF MX1	No	
15	MAXF 401	No	(Fiebig et al., 1999)
16	MAXF 449	Yes	(Burger et al., 2001)
17	MAXF 713	No	(Berger et al., 1992)
18	MAXF 857	No	(Fiebig et al., 1999)

The models represent two major tumor types, namely non-small cell lung cancer (NSCLC; models #1-13) and mammary cancer (models #14-18). Nine of the NSCLC models (#1-9) and all breast cancer models were derived by serial in vivo passage of human tumor fragments in immunodeficient mice. The remaining NSCLC models (#10-13) are cell-based models in which in vivo tumor growth is induced by implantation of in vitro propagated cells into immunocompromised mice.

Berger, D. P., Winterhalter, B. R., and Fiebig, H. H. (1992). Establishment and Characterization of Human Tumor Xenografts in Thymus-Aplastic Nude Mice. In Immunodeficient Mice in Oncology, H. H. Fiebig and D. P. Berger, eds. (Basel: Karger), pp. 23-46.

Burger, A. M., Hartung, G., Stehle, G., Sinn, H., and Fiebig, H. H. (2001). Pre-clinical evaluation of a methotrexate-albumin conjugate (MTX-HSA) in human tumor xenografts in vivo. *Int. J. Cancer*, 92:718-24.

Fiebig, H. H., and Burger, A. M. (2002). Human Tumor Xenografts and Explants. In Tumor Models in Cancer Research, B. A. Teicher, ed. (Totowa, New Jersey: Humana Press), pp. 113-137.

20

- Fiebig, H. H., Dengler, W. A., and Roth, T. (1999). Human Tumor Xenografts: Predictivity, Characterization and Discovery of New Anticancer Agents. In Contributions to Oncology: Relevance of Tumor Models for Anticancer Drug Development, H. H. Fiebig and A. M. Burger, eds. (Basel: Karger), pp. 29-50.
- 5 Gridley, D. S., Andres, M. L., Garner, C., Mao, X. W., and Slater, J. M. (1996). Evaluation of TNF-alpha effects on radiation efficacy in a human lung adenocarcinoma model. *Oncol Res.*, 8:485-95.
- Stein, R., Govindan, S. V., Chen, S., Reed, L., Spiegelman, H., Griffiths, G. L., Hansen, H. J., and Goldenberg, D. M. (2001). Successful therapy of a human lung cancer
10 xenograft using MAb RS7 labeled with residualizing radioiodine. *Crit. Rev. Oncol. Hematol.*, 39:173-80.
- Yamori, T., Sato, S., Chikazawa, H., and Kadota, T. (1997). Anti-tumor efficacy of paclitaxel against human lung cancer xenografts. *Jpn. J. Cancer Res.*, 88:1205-10.
- Zou, Y., Wu, Q. P., Tansey, W., Chow, D., Hung, M. C., Charnsangavej, C., Wallace, S.,
15 and Li, C. (2001). Effectiveness of water soluble poly(L-glutamic acid)-camptothecin conjugate against resistant human lung cancer xenografted in nude mice. *Int. J. Oncol.*, 18:331-6.

Example 3

Detection of Heterodimers in 2C4 Responsive Tumors by Immunoprecipitation

20

2C4 responsive and non-responsive tumors were subjected to immunoprecipitation with anti-ErbB2 antibodies to assay for the presence of ErbB2-ErbB3 and EGFR-ErbB2 heterodimers. Unless otherwise stated the methods were performed according to Maniatis T. et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, New York, USA,
25 Cold Spring Harbor Laboratory Press, 1982.

Anti-HER2, anti-HER3 and anti-HER1 antibodies were selected that did not cross react. To determine if antibodies cross reacted, HER1, HER2, HER3 and HER4 receptors were expressed in human embryonic kidney (HEK) 293 cells. The cells were lysed using a Triton™ X100 (1% weight per volume) containing HEPES buffer (pH 7.5).

30

Approximately 20 µg of total cell protein from control cells and HER1, HER2, HER3 and HER4 expressing cells was separated on an SDS gel and transferred to a nitrocellulose membrane by a semi-dry blotting procedure. After blocking with gelatin, a variety of anti-HER1, anti-HER2 and anti-HER3 antibodies were tested for their cross reactivity against other ErbB receptors. Antibodies selected for use in the experiments described below did
35 not show any significant cross reactivity.

Fresh tumor samples were crushed mechanically on ice and lysed in a buffer containing 50 mM HEPES pH 7.5, 150 mM NaCl, 1.5 mM MgCl₂, 1 mM EDTA, 10 % (w/v) glycerol, 1 % (w/v) Triton™ X-100, 1 mM PMSF, 10 µg/ml aprotinin and 0.4 mM orthovanadate. Completely lysed tumors were centrifuged several times until the supernatants were completely clear. Immunoprecipitation proceeded by combining cleared tumor lysates (5-7 mg protein per lysate), 5 µg anti ErbB2 antibody (ab-3, mouse monoclonal; Cat.# OP15, Oncogene Inc., USA) and 50 µl protein G-coupled agarose in 1.5 ml Eppendorf reaction tubes. Upon addition of a one to twofold volume of 50 mM HEPES buffer, pH 7.5 containing 0.1 % (w/v) Triton™ X-100 the tubes were rotated for 3-4 hours at 4 °C followed by centrifugation. The pellets were washed two to three times with 500 µl of 50 mM HEPES buffer pH 7.5 containing 0.1 % (w/v) Triton™ X-100. An equal volume of 2x Lämmli sample buffer was added to the washed immunoprecipitate and the samples were heated for 5 min at 95°C. The samples were separated by SDS-PAGE and transferred to nitrocellulose. The presence of EGFR-ErbB2 and ErbB2-ErbB3 heterodimers was assessed by probing the blots with antibodies against EGFR (rabbit polyclonal antibody against EGFR, Upstate Inc., USA; Cat. # 06-847) and ErbB3 (polyclonal antibody against ErbB3; Santa Cruz Inc., USA; Cat. # SC-285). A peroxidase (POD) labeled anti-rabbit Fc antibody (BioRad Laboratories Inc. USA) was used as secondary antibody. The blots were visualized using a chemiluminescence substrate (ECL plus, Amersham).

Figure 3 shows the results of these experiments. The presence of ErbB2-ErbB3 and/or EGFR-ErbB2 heterodimers can be seen. A correlation was observed between 2C4 responsiveness, as shown in Table 1, and the presence of ErbB2-ErbB3 and/or EGFR-ErbB2 heterodimers.

Example 4

Correlation of Responsiveness to rhuMAb 2C4 with HER2 Phosphorylation

The effect of rhuMAb 2C4 on tumor growth has been studied in 14 human tumors explanted into mice (9 lung cancer and 5 breast cancer). The explantation of tumor and treatment were performed as described in Example 2. HER2 heterodimers were detected as described in Example 3.

HER2 phosphorylation was assessed by immunoprecipitation of HER2 and Western blot analysis. Positivity was determined by the presence of the phospho-HER2 band of the gel. Negativity was determined by the absence of the band. HER2

phosphorylation was confirmed by immunohistochemistry using a phospho-specific anti-HER2 antibody (clone PN2A, Thor et al., *J. Clin. Oncol.*, 18:3230-9 (2000).

In 5 of the tumors tested (3 lung and 2 breast), a significant inhibition of tumor growth was observed, which correlated with the presence of detectable heterodimers of HER2 with either HER1 or HER3, and with strong HER2 phosphorylation in all cases. In 9 tumors in which no significant response to rhuMAb 2C4 treatment was observed, heterodimers were not detected and HER2 phosphorylation was absent. The presence of HER2 heterodimerization or significant HER2 phosphorylation is a strong predictor of response to rhuMAb 2C4 treatment in nonclinical models. Similar observations have been made with xenografts generated from tumor cell lines.

Example 5

Detection of HER2 Phosphorylation in 2C4 Responsive Tumors

The efficacy of rhuMAb 2C4 was assessed in nine established non-small cell lung carcinoma (NSCLC) xenografted tumor models (LXFE 211, LXFA 289, LXFA 297, LXFE 397, LXFA 526, LXFL 529, LXFA 629, LXFA 1041, LXFL 1071, Oncotest GmbH, Freiburg, Germany). Human tumor xenografts are being considered as the most relevant models for anticancer drug development since the patient's tumors are growing as a solid tumor, develop a stroma, vasculature, a central necrosis and show some differentiation. In addition, the xenografted tumor models resemble very closely the original tumors in histology and chemosensitivity.

Growth inhibition was assessed as described in Example 2. Significant growth inhibitory activity was defined as >50% growth inhibition of the treatment group relative to the control group. In three of the NSCLC models (LXFA 297, LXFA 629, and LXFL 1072), a significant growth inhibitory response to rhuMAb 2C4 treatment was observed. Several hallmarks of ligand-activated HER2 have also been investigated at the protein level. As shown in Figure 4, HER2 could be immunoprecipitated from tumor extracts from eight of nine NSCLC tumors. To determine the activation status of HER2, these blots were then probed with anti-phosphotyrosine (anti-PY) antibody. As shown in the lower panel of Figure 4, the three responsive tumors (LXFA 297, LXFA 629 and 1072) displayed strong HER2 activation.

Example 6

Clinical Study to Identify Lung Cancer Patients for Treatment with rhuMAb 2C4 by Detecting HER2 Heterodimers

A patient is identified as having non-small cell lung cancer (NSCLC) that is responsive to treatment with rhuMAb 2C4 if a tumor from the patient is found to comprise

HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 complexes. A tumor sample is obtained by biopsy or during surgical resection of the tumor. The sample is then analyzed for the presence of HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 heterodimers.

Tumor cells or cell lysates are contacted with an eTagTM that specifically binds HER2. The eTagTM comprises a detectable moiety and a first binding moiety that is specific for HER2. The detectable moiety is linked to the first binding moiety with a cleavable linker. After allowing time for binding, excess eTagTM is removed.

The tumor cells or cell lysates are contacted with a second binding compound that specifically binds HER1 or HER3 or HER4. Unbound compound is removed by washing. The second binding compound is then activated. If the first binding compound and the second binding compound are in close proximity, the activated second binding compound cleaves the cleavable linker in the eTagTM to produce a free detectable moiety. The identification of free detectable moiety in the sample indicates that the tumor comprises HER2/HER1 or HER2/HER3 or HER2/HER4 heterodimers.

Upon determination that the patient is suffering from a tumor that comprises HER2/HER1 and/or HER2/HER3 and/or HER2/HER4 heterodimers, rhuMAb 2C4 is administered intravenously (IV) weekly or every three weeks at 2 or 4 mg/kg, respectively, until disease progression. The antibody is supplied as a multi-dose liquid formulation (20mL fill at a concentration of 20mg/mL or higher concentration). Primary endpoints for efficacy include response rate, and safety. Secondary efficacy endpoints include: overall survival, time to disease progression, quality of life, and/or duration of response.

Example 7

Clinical Study to Identify Cancer Patients for Treatment with rhuMAb 2C4

A biological sample comprising cancer cells is obtained from candidates for the treatment, *e.g.*, by biopsy of tumor tissue, aspiration of tumor cells from ascitic fluid, or any other method known in clinical practice. The biological sample is analyzed for HER2 phosphorylation, *e.g.*, by immunoprecipitation and Western blot analysis, and/or for the presence of HER2/HER3, HER2/HER1 and/or HER2/HER4 heterodimers by any of the

techniques described above. Subjects whose biological sample was positive for HER2 phosphorylation and/or the presence of HER2/HER3, HER2/HER1 and/or HER2/HER4 heterodimers, are likely to show a better response to treatment with rhuMAb 2C4 than patients whose tumor sample showed no HER2 phosphorylation or where no heterodimer was detected.

For example, subjects diagnosed with ovarian cancer will undergo a biopsy of tumor tissue or aspiration of tumor cells from ascites fluid. This tissue will be analyzed for HER2 phosphorylation by immunoprecipitation and Western blot analysis. This will require a minimum of about 250 mg tumor tissue.

Upon determination that the patient suffers from cancer (such as, Castration-Resistant Prostate Cancer - CRPC, or ovarian cancer) that is positive for HER2 phosphorylation, the patient will receive a loading dose of 840 mg of rhuMAb 2C4 on day 1 of cycle 1 (first 21-day treatment period), followed by 420 mg on day 1 of each subsequent 21-day cycle, as continuous intravenous infusion. Treatment continues by intravenous infusion every 3 weeks for up to one year (17 cycles), for patients who show no evidence of disease progression. Treatment can be discontinued any time earlier, for lack of response, adverse effects, or other reasons, at the discretion of the physician.

All references cited throughout the disclosure, and references cited therein, are hereby expressly incorporated by reference.

While the present invention is described with reference to certain embodiments, the invention is not so limited. One skilled in the art will appreciate that various modifications are possible without substantially altering the invention. All such modifications, which can be made without undue experimentation, are intended to be within the scope of the invention.

WHAT IS CLAIMED IS:

1 1. A method of identifying a tumor as responsive to treatment with an anti-HER2
2 antibody comprising:
3 a) detecting the presence of a HER2/HER3 and/or HER2/HER1 protein complex in
4 a sample of said tumor;
5 c) identifying a tumor as responsive to treatment with anti-HER2 antibody when a
6 complex is detected.

1 2. The method of claim 1 wherein the anti-HER2 antibody blocks ligand activation
2 of an ErbB heterodimer comprising HER2.

1 3. The method of claim 1 wherein the anti-HER2 antibody is monoclonal antibody
2 2C4.

1 4. The method of claim 1 wherein the anti-HER2 antibody is rhuMAb 2C4.

1 5. The method of claim 1 wherein the presence of a HER2/HER3 and/or
2 HER2/HER1 protein complex is detected by:

3 a) immunoprecipitating any protein complexes that comprise HER2 with an anti-
4 HER2 antibody;

5 b) contacting the immunoprecipitated complexes with an antibody selected from the
6 group consisting of anti-HER3 antibodies and anti-HER1 antibodies; and

7 c) determining if an anti-HER3 and/or anti-HER1 antibody binds to the
8 immunoprecipitated complexes,

9 wherein a HER2/HER3 and/or HER2/HER1 complex is detected if it is determined that
10 anti-HER3 and/or anti-HER1 antibodies bind to the immunoprecipitated complexes.

1 6. The method of claim 1 wherein the presence of a HER2/HER3 and/or
2 HER2/HER1 protein complex is detected by:

3 a) contacting the tumor sample with an anti-HER2 antibody that comprises a
4 fluorophore;

5 b) contacting the tumor sample with an antibody selected from the group consisting
6 of anti-HER3 and anti-HER1 antibodies, wherein said antibody comprises a second fluorophore;

7 c) determining if the first fluorophore and the second fluorophore are in close
8 proximity by measuring the fluorescence resonance energy transfer,

9 wherein the presence of a HER2/HER3 and/or HER2/HER1 protein complex is detected
10 if the first and second fluorophore are determined to be in close proximity.

- 1 7. The method of claim 1 wherein the presence of a HER2/HER3 and/or
2 HER2/HER1 protein complex is detected by:
- 3 a) contacting the tumor sample with a first binding compound, wherein said first
4 binding compound comprises a first target binding moiety that specifically binds HER2 and
5 further comprises a detectable moiety linked to the first target binding moiety by a cleavable
6 linker;
- 7 b) contacting the tumor sample with a second binding compound, wherein the
8 second binding compound comprises a second target binding moiety that specifically binds HER3
9 or HER1 and an activatable cleaving agent;
- 0 c) activating the cleaving agent such that if the first binding compound and the
1 second binding compound are in close proximity the second binding compound cleaves the
2 cleavable linker in the first binding compound to produce a free detectable moiety; and
- 3 d) identifying the presence of the free detectable moiety,
4 wherein the presence of a HER2/HER3 or HER2/HER1 protein complex is detected when
5 free detectable moiety is identified.

1 8. The method of claim 7 wherein the first target binding moiety comprises an anti-
2 HER2 antibody or antibody fragment.

1 9. The method of claim 7 wherein the first target binding moiety comprises a HER2
2 receptor ligand.

1 10. The method of claim 7 wherein the second target binding moiety comprises an
2 anti-HER3 antibody or antibody fragment.

1 11. The method of claim 7 wherein the second target binding moiety comprises a
2 HER3 receptor ligand.

1 12. The method of claim 7 wherein the second target binding moiety comprises an
2 anti-HER1 antibody or antibody fragment.

1 13. The method of claim 7 wherein the second target binding moiety comprises a
2 HER1 receptor ligand.

1 14. The method of claim 7 wherein the sample is obtained from a patient suffering
2 from the tumor.

1 15. The method of claim 14 wherein the sample is obtained by a biopsy of the tumor.

- 1 16. The method of claim 14 wherein the sample is obtained by purifying circulating
2 tumor cells from the patient's blood.
- 1 17. The method of claim 14 wherein the sample is obtained during surgery to remove
2 the tumor from the patient.
- 1 18. The method of claim 1 wherein the sample of the tumor is obtained from a mouse.
- 1 19. The method of claim 18 wherein the tumor is a xenografted tumor.
- 1 20. The method of claim 19 wherein the xenografted tumor is produced by
2 transplanting a fragment of a human tumor into a mouse.
- 1 21. The method of claim 1 wherein the tumor is a lung tumor.
- 1 22. The method of claim 1 wherein the tumor is a mammary tumor.
- 1 23. A method for identifying tumor cells as responsive to treatment with an antibody
2 inhibiting the association of HER2 with another member of the ErbB receptor family
3 comprising:
4 (a) providing a biological sample comprising HER2-positive tumor cells; and
5 (b) detecting the phosphorylation of an ErbB receptor in said biological sample,
6 wherein said phosphorylation indicates that said tumor cells are responsive to treatment with said
7 antibody.
- 1 24. The method of claim 23 wherein the phosphorylation of an ErbB2 (HER2)
2 receptor is detected.
- 1 25. The method of claim 23 wherein the other member is selected from the group
2 consisting of HER3, HER1 and HER4.
- 1 26. The method of claim 23 wherein the antibody binds HER2.
- 1 27. The method of claim 26 wherein the anti-HER2 antibody blocks ligand activation
2 of an ErbB heterodimer comprising HER2.
- 1 28. The method of claim 27 wherein the antibody is rhuMAb 2C4.
- 1 29. The method of claim 25 wherein the antibody binds HER3.

1 30. The method of claim 25 wherein the antibody binds HER1.

1 31. The method of claim 25 wherein the antibody binds HER4.

1 32. The method of claim 23 additionally comprising detecting the presence of at least
2 one protein complex selected from the group consisting of HER2/HER3, HER2/HER1, and
3 HER2/HER4 in the sample.

1 33. The method of claim 32 wherein the presence of said protein complex or
2 complexes is detected by:

3 a) immunoprecipitating any protein complex that comprises HER2 with an anti-
4 HER2 antibody;

5 b) contacting the immunoprecipitated complex with at least one antibody selected
6 from the group consisting of anti-HER3, anti-HER1, and anti-HER4 antibodies; and

7 c) determining if said anti-HER3 and/or anti-HER1 and/or anti-HER4 antibody
8 binds to the immunoprecipitated complex,

9 wherein a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 complex is detected if it
10 is determined that anti-HER3 and/or anti-HER1 and/or anti-HER4 antibodies bind to the
11 immunoprecipitated complex.

1 34. The method of claim 32 wherein the presence of said protein complex or
2 complexes is detected by:

3 a) contacting the tumor sample with an anti-HER2 antibody that comprises a
4 fluorophore;

5 b) contacting the tumor sample with an antibody selected from the group consisting
6 of anti-HER3, anti-HER1 and anti-HER4 antibodies, wherein said antibody comprises a second
7 fluorophore;

8 c) determining if the first fluorophore and the second fluorophore are in close
9 proximity by measuring the fluorescence resonance energy transfer,

10 wherein the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein
11 complex is detected if the first and second fluorophore are determined to be in close proximity.

1 35. The method of claim 32 wherein the presence of said protein complex or
2 complexes is detected by:

3 a) contacting the tumor sample with a first binding compound, wherein said first
4 binding compound comprises a first target binding moiety that specifically binds HER2 and

5 further comprises a detectable moiety linked to the first target binding moiety by a cleavable
6 linker;

7 b) contacting the tumor sample with a second binding compound, wherein the
8 second binding compound comprises a second target binding moiety that specifically binds HER3
9 or HER1 or HER4 and an activatable cleaving agent;

10 c) activating the cleaving agent such that if the first binding compound and the
11 second binding compound are in close proximity the second binding compound cleaves the
12 cleavable linker in the first binding compound to produce a free detectable moiety; and

13 d) identifying the presence of the free detectable moiety,
14 wherein the presence of a HER2/HER3 or HER2/HER1 or HER2/HER4 protein complex
15 is detected when free detectable moiety is identified.

1 36. The method of claim 35 wherein the first target binding moiety comprises an anti-
2 HER2 antibody or antibody fragment, or a HER2 receptor ligand.

1 37. The method of claim 35 wherein the second target binding moiety comprises an
2 anti-HER3 antibody or antibody fragment, or a HER3 receptor ligand.

1 38. The method of claim 35 wherein the second target binding moiety comprises an
2 anti-HER1 antibody or antibody fragment, or a HER1 receptor ligand.

1 39. The method of claim 35 wherein the second target binding moiety comprises an
2 anti-HER4 antibody or antibody fragment, or a HER4 receptor ligand.

1 40. The method of claim 23 wherein the biological sample is tissue obtained from a
2 tumor biopsy.

1 41. The method of claim 23 wherein the biological sample is a biological fluid
2 comprising circulating tumor cells and/or circulating plasma proteins.

1 42. The method of claim 23 wherein the tumor is selected from the group consisting
2 of breast cancer, prostate cancer, lung cancer, colorectal cancer and ovarian cancer.

1 43. The method of claim 23 wherein ErbB receptor phosphorylation is determined by
2 immunoprecipitation of the ErbB receptor and Western blot analysis.

1 44. The method of claim 43 wherein ErbB receptor phosphorylation is indicated by
2 the presence of a phospho-ErbB receptor band on the gel.

1 45. The method of claim 43 further comprising the step of confirming ErbB receptor
2 phosphorylation by immunohistochemistry using a phospho-specific anti-ErbB receptor
3 antibody.

1 46. The method of claim 23 wherein ErbB receptor phosphorylation is determined by
2 immunohistochemistry.

1 47. A method for predicting the response of a subject diagnosed with a HER2-
2 positive tumor to treatment with an antibody inhibiting the association of HER2 with another
3 member of the ErbB receptor family comprising:

4 (a) providing a biological sample obtained from said subject, comprising HER2-
5 positive tumor cells; and

6 (b) detecting phosphorylation of an ErbB receptor in said biological sample,
7 wherein said phosphorylation indicates that said patient is likely to respond to treatment
8 with said antibody.

1 48. The method of claim 47 wherein said ErbB receptor is ErbB2 (HER2).

1 49. The method of claim 47 wherein the other member is selected from the group
2 consisting of HER3, HER1 and HER4.

1 50. The method of claim 47 wherein the antibody binds HER2.

1 51. The method of claim 50 wherein the anti-HER2 antibody blocks ligand activation
2 of an ErbB heterodimer comprising HER2.

1 52. The method of claim 51 wherein the antibody is rhuMAb 2C4.

1 53. The method of claim 49 wherein the antibody binds HER3.

1 54. The method of claim 49 wherein the antibody binds HER1.

1 55. The method of claim 49 wherein the antibody binds HER4.

1 56. The method of claim 47 additionally comprising detecting the presence of at least
2 one protein complex selected from the group consisting of HER2/HER3, HER2/HER1, and
3 HER2/HER4 in the sample.

1 57. The method of claim 56 wherein the presence of said protein complex is detected
2 by:

3 a) immunoprecipitating any protein complexes that comprise HER2 with an anti-
4 HER2 antibody;

5 b) contacting the immunoprecipitated complexes with an antibody selected from the
6 group consisting of anti-HER3, anti-HER1, and anti-HER4 antibodies; and

7 c) determining if an anti-HER3 and/or anti-HER1 and/or anti-HER4 antibody binds
8 to the immunoprecipitated complexes,
9 wherein a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 complex is detected if it
10 is determined that anti-HER3 and/or anti-HER1 and/or anti-HER4 antibodies bind to the
11 immunoprecipitated complexes.

1 58. The method of claim 56 wherein the presence of HER2/HER3 and/or
2 HER2/HER1 and/or HER2/HER4 protein complex is detected by:

3 a) contacting the tumor sample with an anti-HER2 antibody that comprises a
4 fluorophore;

5 b) contacting the tumor sample with an antibody selected from the group consisting
6 of anti-HER3, anti-HER1 and anti-HER4 antibodies, wherein said antibody comprises a second
7 fluorophore;

8 c) determining if the first fluorophore and the second fluorophore are in close
9 proximity by measuring the fluorescence resonance energy transfer,

10 wherein the presence of a HER2/HER3 and/or HER2/HER1 and/or HER2/HER4 protein
11 complex is detected if the first and second fluorophore are determined to be in close proximity.

1 59. The method of claim 56 wherein the presence of HER2/HER3 and/or
2 HER2/HER1 and/or HER2/HER4 protein complex is detected by:

3 a) contacting the tumor sample with a first binding compound, wherein said first
4 binding compound comprises a first target binding moiety that specifically binds HER2 and
5 further comprises a detectable moiety linked to the first target binding moiety by a cleavable
6 linker;

7 b) contacting the tumor sample with a second binding compound, wherein the
8 second binding compound comprises a second target binding moiety that specifically binds
9 HER3, HER1, or HER4 and an activatable cleaving agent;

10 c) activating the cleaving agent such that if the first binding compound and the
11 second binding compound are in close proximity the second binding compound cleaves the
12 cleavable linker in the first binding compound to produce a free detectable moiety; and

13 d) identifying the presence of the free detectable moiety,

14 wherein the presence of a HER2/HER3 or HER2/HER1 or HER2/HER4 protein complex
15 is detected when free detectable moiety is identified.

1 60. The method of claim 59 wherein the first target binding moiety comprises an anti-
2 HER2 antibody or antibody fragment, or a HER2 receptor ligand.

1 61. The method of claim 59 wherein the second target binding moiety comprises an
2 anti-HER3 antibody or antibody fragment, or a HER3 receptor ligand.

1 62. The method of claim 59 wherein the second target binding moiety comprises an
2 anti-HER1 antibody or antibody fragment, or a HER1 receptor ligand.

1 63. The method of claim 59 wherein the second target binding moiety comprises an
2 anti-HER4 antibody or antibody fragment, or a HER4 receptor ligand.

1 64. The method of claim 47 wherein the biological sample is tissue obtained from a
2 tumor biopsy.

1 65. The method of claim 47 wherein the biological sample is a biological fluid
2 comprising circulating tumor cells and/or circulating plasma proteins.

1 66. The method of claim 47 wherein the tumor is selected from the group consisting
2 of breast cancer, prostate cancer, lung cancer, colorectal cancer and ovarian cancer.

1 67. The method of claim 47 wherein ErbB receptor phosphorylation is determined by
2 immunoprecipitation of the ErbB receptor and Western blot analysis.

1 68. The method of claim 67 wherein ErbB receptor phosphorylation is indicated by
2 the presence of a phospho-ErbB receptor band on the gel.

1 69. The method of claim 67 further comprising the step of confirming ErbB receptor
2 phosphorylation by immunohistochemistry using a phospho-specific anti-ErbB receptor
3 antibody.

1 70. The method of claim 47 wherein ErbB receptor phosphorylation is determined by
2 immunohistochemistry.

1 71. A method for identifying a subject responsive to treatment with an anti-HER2
2 antibody comprising

3 a) detecting phosphorylation of an ErbB receptor in circulating tumor cells of said
4 subject, and

5 b) determining that said subject is likely to respond to treatment with an anti-HER2
6 antibody if said phosphorylation is detected.

1 72. The method of claim 71 wherein ErbB2 (HER2) phosphorylation is detected.

1 73. The method of claim 72 wherein said subject is a human.

1 74. The method of claim 73 further comprising treating said subject with an anti-
2 HER2 antibody.

1 75. The method of claim 74 wherein said anti-HER2 antibody is rhuMAb 2C4

1 76. A method of treating a patient comprising administering to the patient a
2 therapeutically effective amount of an antibody which binds HER2, wherein the patient is

3 suffering from a tumor which has been determined to comprise HER2/HER3 and/or
4 HER2/HER1 and/or HER2/HER4 heterodimers.

1 77. The method of claim 76, wherein the antibody blocks ligand activation of an ErbB
2 heterodimer comprising HER2.

1 78. The method of claim 77 wherein the antibody is monoclonal antibody 2C4.

1 79. The method of claim 77 wherein the antibody is rhuMAb 2C4.

1 80. An article of manufacture comprising a container comprising an antibody which
2 binds HER2 and instructions for administering the antibody to a patient suffering from a tumor
3 wherein the tumor has been determined to comprise HER2/HER3 and/or HER2/HER1 and/or
4 HER2/HER4 heterodimers.

1 81. The article of manufacture of claim 80 wherein the antibody blocks ligand
2 activation of an ErbB heterodimer comprising HER2.

1 82. The article of manufacture of claim 81 wherein the container comprises
2 monoclonal antibody 2C4.

1 83. The article of manufacture of claim 81 wherein the container comprises rhuMAb
2 2C4.

1 84. A method of treating a patient comprising administering to the patient a
2 therapeutically effective amount of an antibody which binds HER2, wherein the patient is
3 suffering from a tumor which has been determined to have a phosphorylated ErbB receptor.

1 85. The method of claim 84 wherein the ErbB receptor is HER2.

1 86. The method of claim 84 wherein the antibody blocks ligand activation of an ErbB
2 heterodimer comprising HER2.

1 87. The method of claim 84 wherein the antibody is monoclonal antibody 2C4.

88. The method of claim 87 wherein the antibody is rhuMAb 2C4.

1/4

	10	20	30	40
2C4	DTVMTQSHKIMSTSVGDRVSITC	[KASQDV SIGVA]	WYQQRP	
	** * * * *	*	*	
574	DIQMTQSPSSLSASVGDRVTITC	[KASQDV SIGVA]	WYQQKP	
		* * * *		
hum xI	DIQMTQSPSSLSASVGDRVTITC	[RASQSI NYLA]	WYQQKP	
	50	60	70	80
2C4	GQSPKLLIY [SASYRYT]	GVPDRFTGSGSGTDFTFTISSVQA		
	** * * *	* *	* *	
574	GKAPKLLIY [SASYRYT]	GVPSRFSGSGSGTDFTLTISLQP		
	* * * * *			
hum xI	GKAPKLLIY [AASSLES]	GVPSRFSGSGSGTDFTLTISLQP		
	90	100		
2C4	EDLAVYYC [QYYIYPYT]	FGGGTKLEIK	(SEQ ID NO: 1)	
	* *	* *		
574	EDFATYYC [QYYIYPYT]	FGQGTKVEIK	(SEQ ID NO: 3)	
	* * * *			
hum xI	EDFATYYC [QYNSLPWT]	FGQGTKVEIK	(SEQ ID NO: 5)	

FIG. 1A: VARIABLE LIGHT

	10	20	30	40
2C4	EVQLQQSGPELVKPGTQSVKISCKAS	[GFTFTDY TMD]	WVKQS	
	* * * * *	*	* *	
574	EVQLVESGGGLVQPGGSLRLS CAAS	[GFTFTDY TMD]	WVRQA	
		* * * *		
hum III	EVQLVESGGGLVQPGGSLRLS CAAS	[GFTFSS YAMS]	WVRQA	
	50	60	70	80
2C4	HGKSLEWIG [DVNPNSCGSIYNQRFKG]	KASLTVD RSRIVYM		
	* * *	* * * *	* * * *	
574	PGKGLEWVA [DVNPNSCGSIYNQRFKG]	RFTLSVDRSKNTLYL		
	* * * * *	* * * *		
hum III	PGKGLEWVA [VISGDGGSTYYADSVKG]	RFTISRDN SKNTLYL		
	abc	90	100ab	110
2C4	ELRSLTFEDTAVYYCAR	[NLGPSFYFDY]	WGQGTTLTVSS	(SEQ ID NO: 2)
	* * *		* *	
574	QMNSLRAEDTAVYYCAR	[NLGPSFYFDY]	WGQGLTVTVSS	(SEQ ID NO: 4)
		* * * * *		
hum xI	QMNSLRAEDTAVYYCAR	[GRVGYSLYDY]	WGQGLTVTVSS	(SEQ ID NO: 6)

FIG. 1B: VARIABLE HEAVY

FIG. 2A

MCF7
Low/Normal HER2






FIG. 2B

SK-BR-3
High HER2



IP: αHER2
WB: αHER3

HER2 containing dimers in protein extracts from
Oncotest NSCLC xenograft explants



HER2/HER1 dimers and HER2/HER3 dimers are elevated in the rhuMAb 2C4 responsive versus rhuMAb 2C4 non-responsive NSCLC xenografts.

FIG. 3

4/4

HER2 expression/activation in protein extracts from
Oncotest NSCLC xenograft explants




FIG. 4

39766-0114PCseq.txt

SEQUENCE LISTING

<110> Genentech, Inc.
 Koll, Hans
 Bossenmaier, Birgit
 Muller, Hans-Joachim
 Sliwkowski, Mark
 Kelsey, Stephen

<120> Methods For Identifying Tumors That Are
 Responsive To Treatment With Anti-ErbB2 Antibodies

<130> 39766-0114PC

<150> US 60/396,290

<151> 2002-07-15

<150> US 60/480,043

<151> 2003-06-20

<160> 6

<170> FastSEQ for windows version 4.0

<210> 1

<211> 107

<212> PRT

<213> Homosapiens

<400> 1

```

Asp Thr Val Met Thr Gln Ser His Lys Ile Met Ser Thr Ser Val Gly
 1          5          10
Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Ile Gly
          20          25          30
Val Ala Trp Tyr Gln Gln Arg Pro Gly Gln Ser Pro Lys Leu Ile
          35          40          45
Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
          50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala
65          70          75          80
Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln Tyr Tyr Ile Tyr Pro Tyr
          85          90          95
Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
          100          105
    
```

<210> 2

<211> 119

<212> PRT

<213> Homosapiens

<400> 2

```

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Thr
 1          5          10
Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Phe Thr Phe Thr Asp Tyr
          20          25          30
Thr Met Asp Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
          35          40          45
Gly Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr Asn Gln Arg Phe
          50          55          60
Lys Gly Lys Ala Ser Leu Thr Val Asp Arg Ser Ser Arg Ile Val Tyr
    
```

39766-0114PCseq.txt

65 Met Glu Leu Arg Ser 70 Leu Thr Phe Glu Asp Thr Ala Val Tyr Tyr Cys 80
 85 Pro Ser Phe Tyr Phe Asp Tyr Trp Gly Gln Gly 95
 Ala Arg Asn Leu Gly 100 Thr Thr Leu Thr Val Ser Ser 110
 115

<210> 3
 <211> 107
 <212> PRT
 <213> Homosapiens

<400> 3
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Ser Ile Gly
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Tyr Ile Tyr Pro Tyr
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys 100 105

<210> 4
 <211> 119
 <212> PRT
 <213> Homosapiens

<400> 4
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr Asp Tyr
 20 25 30
 Thr Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Asp Val Asn Pro Asn Ser Gly Gly Ser Ile Tyr Asn Gln Arg Phe
 50 55 60
 Lys Gly Arg Phe Thr Leu Ser Val Asp Arg Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asn Leu Gly Pro Ser Phe Tyr Phe Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ser 115

<210> 5
 <211> 107
 <212> PRT
 <213> Homosapiens

<400> 5
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

39766-0114PCseq.txt

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Ser Asn Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Ala Ala Ser Ser Leu Glu Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Tyr Asn Ser Leu Pro Trp
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
 100 105

<210> 6
 <211> 119
 <212> PRT
 <213> Homosapiens

<400> 6
 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Val Ile Ser Gly Asp Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Gly Arg Val Gly Tyr Ser Leu Tyr Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ser
 115