



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C11D 3/386, C07K 5/08, 5/06</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/13460</p> <p>(43) International Publication Date: 2 April 1998 (02.04.98)</p>
<p>(21) International Application Number: PCT/US97/16623</p> <p>(22) International Filing Date: 19 September 1997 (19.09.97)</p> <p>(30) Priority Data: 60/026,632 24 September 1996 (24.09.96) US</p> <p>(71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): McIVER, John, McMillan [US/US]; 832 Cedarhill Drive, Cincinnati, OH 45240 (US). HUBER, Alan, Carl [US/US]; 4815 Trudy Lane, Hamilton, OH 45013 (US). McKILLOP, Kristen, Louise [US/US]; 1324 Duncan Avenue, Cincinnati, OH 45208 (US).</p> <p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).</p>		<p>(81) Designated States: BR, CA, CN, JP, MX, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: LIQUID DETERGENTS CONTAINING PROTEOLYTIC ENZYME AND PROTEASE INHIBITORS</p>		
<p>(57) Abstract</p> <p>Aqueous liquid detergent compositions are described which comprise a proteolytic enzyme wherein the proteolytic activity is reversibly inhibited by a peptide protease inhibitor selected from the group consisting of aldehydes and trifluoromethyl ketones.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

LIQUID DETERGENTS CONTAINING PROTEOLYTIC
ENZYME AND PROTEASE INHIBITORS

5

TECHNICAL FIELD

This invention relates to liquid detergent compositions containing enzymes. More specifically, this invention pertains to liquid detergent compositions containing a deterative surfactant, a proteolytic enzyme, and a peptide protease inhibitor selected from the group consisting of peptide aldehydes and peptide trifluoromethyl ketones.

BACKGROUND OF THE INVENTION

Protease-containing liquid aqueous detergents are well-known, especially in the context of laundry washing. A commonly encountered problem in such protease-containing liquid aqueous detergents is the degradation phenomenon by the proteolytic enzyme of second enzymes in the composition, such as amylase, lipase, and cellulase, or on the protease itself. As a result, the stability of the second enzyme or the protease itself in the detergent composition is affected and the detergent composition consequently performs less well.

In response to this problem, it has been proposed to use various protease inhibitors or stabilizers. For instance, various references have proposed the use of the following compounds to aid in the stabilization of enzymes: benzamidine hydrochloride, lower aliphatic alcohols or carboxylic acids, mixtures of a polyol and a boron compound, aromatic borate esters, and calcium, particularly calcium formate. Recently, it was discovered that certain peptide aldehydes and peptide trifluoromethyl ketones act to stabilize protease enzyme.

Although these compounds have been used to varying success in liquid detergents, they are not free of problems. For example certain peptide aldehydes can be rather expensive and create complexities for the formulators, especially for liquid detergents. Other inhibitors such as calcium and boric acids are less expensive but do not stabilize enzymes as well as peptide aldehydes. It is thus an object of the present invention to provide alternate peptide aldehydes and trifluoromethyl ketone protease inhibitors which are effective and suitable for use in a liquid detergent compositions.

BACKGROUND ART

It has been proposed to use various protease inhibitors or stabilizers. For instance, US 4,566,985 proposes to use benzamidine hydrochloride; EP 376 705 proposes to use lower aliphatic alcohols or carboxylic acids; EP 381 262 proposes to use a mixture of a polyol and a boron compound; and EP91870072.5 proposes to use aromatic borate esters. See also U.S. Pat. No. 5,030,378 issued July 9, 1991. Also see US4,261,868; US4,404,115; US4,318,818; and EP130,756.

The use of peptide derivatives for the inhibition of proteins appears to have been disclosed in therapeutic applications. EP 293 881 discloses the use of peptide boronic acids as inhibitors of trypsin-like serine proteases. EP 185 390 and US 4,399,065 disclose the use of certain peptide aldehydes derivatives for the inhibition of blood coagulation. J 90029670 discloses the use of optically active alpha amino aldehydes for the inhibition of enzymes in general. See also "Inhibition of Thrombin and Trypsin by Tripeptide Aldehydes", Int. J. Peptide Protein Res., Vol 12 (1978), pp. 217-221; Gaal, Bacsy & Rappay, and "Tripeptide Aldehyde Protease Inhibitors May Depress in Vitro Prolactin and Growth Hormone Release" Endocrinology, Vol. 116, No. 4 (1985), pp. 1426-1432; Rappay, Makara, Bajusz & Nagy. Certain peptide aldehydes have also been disclosed in EP-A-473 502 for inhibiting protease-mediated skin irritation.

In particular see EP185,390, WO94/04651, published 3 March 1994, WO94/04652, published 3 March 1994, EP 583,536, published February 23, 1994, EP 583,535, published February 3, 1994, EP 583,534, published February 23, 1994, WO 93/13125, published July 8, 1993, US4,529,525, US4,537,706, US4,537,707, and US5,527,487.

SUMMARY OF THE INVENTION

The invention herein is a liquid detergent composition comprising:

- a) an effective amount of a deterative surfactant;
- b) an active proteolytic enzyme; and
- c) a peptide protease inhibitor having the formula:



wherein B is a peptide chain comprising from 2 to 5 amino acid moieties; X is hydrogen or CF₃; Z is an N-capping moiety selected from the group consisting of sulfonamides, phosphoramidates, thioureas, sulfenamides, sulfonic acids, phosphinamides, thiocarbamates, amidophosphates, sulfamoyl derivatives, and phosphonamides; and R is selected from the group consisting of straight or branched

C₁ - C₆ unsubstituted alkyl, phenyl, and C₇ - C₉ alkylaryl moieties. Preferred compositions further comprise a source of calcium ion or boric acid.

Preferably, the liquid detergent compositions herein comprise, by weight of composition:

- 5 a) from about 1 to about 95%, preferably from about 8% to about 70%, of said deterative surfactant;
- b) from about 0.0001% to about 5%, preferably from about 0.0003% to about 0.1%, of an active proteolytic enzyme;
- c) from about 0.00001% to about 5%, preferably from about 0.0001% to about 10
10 1%, more preferably from about 0.0006% to about 0.5%, of the described peptide protease inhibitor;
- d) optionally, from about 0.01% to about 1%, preferably from about 0.05% to about 0.5%, of calcium ion; and
- e) optionally, from about 0.25% to about 10%, preferably from about 0.5% to
15 about 5%, of boric acid or a compound capable of forming boric acid, preferably with a diol.

The proteolytic enzyme useful herein is preferably a subtilisin-type protease and may be selected from the group consisting of Alcalase[®], Subtilisin BPN', Protease A, Protease B, and mixtures thereof.

- 20 The source of calcium ion for use herein is preferably selected from calcium formate, calcium xylene sulfonate, calcium chloride, calcium acetate, calcium sulfate, and mixtures thereof.

The dishcare compositions herein may contain further deterative adjuncts, including but not limited to, one or more of the following: suds boosters, chelants,
25 polyacrylate polymers, dispersing agents, dyes, perfumes, processing aids, and mixtures thereof. Moreover for dishcare compositions, the liquid detergent compositions may further comprise an effective amount of amylase enzyme. Additionally, the dishcare compositions may optionally comprise an effective amount of a source of boric acid and a diol. Typically dishcare compositions will
30 optionally, but preferably, comprise from about 0.25% to about 10%, preferably from about 0.5% to about 5%, more preferably from about 0.75% to about 3%, by weight of boric acid or a compound capable of forming boric acid and a diol, e.g. 1,2-propaneidiol.

In a preferred embodiment for heavy duty detergent compositions useful in
35 laundry care, the liquid detergent composition further comprises an effective amount one or more of the following enzymes: lipase, amylase, cellulase, and mixtures thereof. Preferably for laundry compositions, the second enzyme is lipase and is

obtained by cloning the gene from Humicola Lanuginosa and expressing the gene in Aspergillus Oryzae. Lipase is utilized in an amount of from about 10 to about 18000 lipase units per gram, preferably from about from about 60 to about 6000 units per gram.

5 In another preferred composition useful for laundry care, the second enzyme is a cellulase derived from Humicola Insolens and is utilized in an amount of from about 0.0001% to about 0.1% by weight of the total composition of said cellulase.

The compositions herein may contain further detergent adjuncts, including but not limited to, one or more of the following: suds boosters, builders, soil release
10 polymers, polyacrylate polymers, dispersing agents, dye transfer inhibitors, dyes, perfumes, processing aids, brighteners, and mixtures thereof. Additionally, for laundrycare compositions, the detergent surfactant is typically present in an amount of from about 10% to about 70%, by weight of total composition. Moreover, the laundry compositions may optionally comprise an effective amount of a source of
15 boric acid and a diol. Typically laundry compositions will optionally, but preferably, comprise from about 0.25% to about 10%, preferably from about 0.5% to about 5%, more preferably from about 0.75% to about 3%, by weight of boric acid or a compound capable of forming boric acid and a diol, e.g. 1,2-propanediol.

All percentages and proportions herein are by weight, and all references cited
20 are hereby incorporated by reference, unless otherwise specifically indicated.


DETAILED DESCRIPTION OF THE INVENTION

Definitions - The present detergent compositions comprise an "effective amount" or a "stain removal-improving amount" of individual components defined herein. An "effective amount" or "stain removal-improving amount" is any amount
25 capable of measurably improving soil cleaning or stain removal from a substrate, i.e., soiled fabric or soiled dishware, when it is washed by the consumer. In general, this amount may vary quite widely.

The liquid aqueous detergent compositions according to the present invention comprise three essential ingredients: (A) a peptide protease inhibitor selected from
30 the group consisting of aldehydes and trifluoromethyl ketones, or a mixture thereof, as described herein, (B) a proteolytic enzyme or a mixture thereof, and (C) a detergent surfactant. The compositions according to the present invention preferably further comprise (D) a source of calcium ions, (E) a detergent-compatible second enzyme or a mixture thereof, (F) boric acid and a diol, and may further comprise (G)
35 other optional ingredients.

Peptide protease inhibitors - The detergent compositions according to the present invention comprise, as a first essential ingredient, a peptide protease

inhibitor selected from the group consisting of aldehydes and trifluoromethyl ketones, or mixtures thereof, having the formula:



5

wherein B is a peptide chain comprising from 2 to 5 amino acid moieties; X is hydrogen or CF₃; Z is an N-capping moiety selected from the group consisting of phosphoramidate [(R''O)₂(O)P-], sulfenamide [(SR'')₂-], sulfonamide [(R''(O)₂S-], sulfonic acid [SO₃H], phosphinamide [(R'')₂(O)P-], sulfamoyl derivative [R''O(O)₂S-], thiourea [(R'')₂N(O)C-], thiocarbamate [R''O(S)C-], phosphonate [R''-P(O)OH], and amidophosphate [R''O(OH)(O)P-], wherein each R'' is independently selected from the group consisting of straight or branched C₁-C₆ unsubstituted alkyl, phenyl, C₇-C₉ alkylaryl, and cycloalkyl moieties, wherein the cycloalkyl ring may span C₄-C₈ and may contain one or more heteroatoms selected from the group consisting of O, N, and S; and R is selected from the group consisting of straight or branched C₁ - C₆ unsubstituted alkyl, phenyl, and C₇ - C₉ alkylaryl moieties.

Preferred R moieties are selected from the group consisting of methyl, isopropyl, sec-butyl, iso-butyl, -C₆H₅, -CH₂-C₆H₅, and -CH₂CH₂-C₆H₅, which respectively may be derived from the amino acids Ala, Val, Ile, Leu, P Gly (phenylglycine), Phe, and HPhe (homophenylalanine) by converting the carboxylic acid group to an aldehyde or trifluoromethyl ketone group. While such moieties are therefore not amino acids (and they may or may not have been synthesized from an amino acid precursor), for purposes of simplification of the exemplification of inhibitors useful here, the aldehyde portion of the inhibitors are indicated as derived from amino acids by the addition of "H" after the analogous amino acid [e.g., "-AlaH" represents the chemical moiety "-NHCH(CH₃)C(O)H"]. Trifluoromethyl ketones are similarly represented by the addition of "CF₃" after the analogous amino acid (e.g., "-AlaCF₃" represents the chemical moiety "-NHCH(CH₃)C(O)CF₃").

Preferred B peptide chains are selected from the group consisting of peptide chains having the amino acid sequences according to the general formula:



such that the following amino acids, when present, are :

- A¹ is selected from Ala, Gly;
- A² is selected from Val, Ala, Gly, Ile;
- A³, if present, is selected from Phe, Leu, Val, Ile;
- A⁴, if present, is any amino acid, but preferably is selected from Gly, Ala;
- A⁵, if present, is any amino acid, but preferably is Gly, Ala, Lys.

35

The present invention aldehydes may be prepared from the corresponding amino acid whereby the C-terminal end of said amino acid is converted from a carboxylic group to an aldehyde group. Such aldehydes may be prepared by known processes, for instance as described in US 5015627, EP 185 930, EP 583,534, and
5 DE 32 00 812.

The present invention trifluoromethyl ketones may be prepared from the corresponding amino acid whereby the C-terminal end of said amino acid is converted from a carboxylic group to the trifluoromethyl ketone group. Such trifluoromethyl ketones may be prepared by known processes, for instance as
10 described in EP 583,535.

While not wanting to be bound by theory it is believed that the peptide protease inhibitors according to the present invention bind to the proteolytic enzyme in the liquid detergent composition, thereby inhibiting said proteolytic enzyme. Upon dilution in water, the proteolytic activity is restored by dissociation of the
15 proteolytic enzyme/peptide protease inhibitor complex.

The N-terminal end of said protease inhibitors according to the present invention is protected by one of the N-capping moiety protecting groups selected from the group consisting of sulfonamides, phosphoramidates, thioureas, sulfenamides, sulfonic acids, phosphinamides, thiocarbamates, amidophosphates,
20 and phosphoramidates. However, in a highly preferred embodiment of the present invention, the N-terminal end of said protease inhibitor is protected by a methyl, ethyl or benzyl sulfonamide [CH_3SO_2^- ; $\text{CH}_3\text{CH}_2\text{SO}_2^-$; or $\text{C}_6\text{H}_5\text{CH}_2\text{SO}_2^-$], and methyl, ethyl or benzyl amidophosphate [$\text{CH}_3\text{O}(\text{OH})(\text{O})\text{P}^-$; $\text{CH}_3\text{CH}_2\text{O}(\text{OH})(\text{O})\text{P}^-$; or $\text{C}_6\text{H}_5\text{CH}_2\text{O}(\text{OH})(\text{O})\text{P}^-$] groups.

25 Synthesis of N-capping groups can be found in the following references: Protective Groups in Organic Chemistry, Greene, T., Wuts, P., John Wiley & Sons, New York, 1991, pp 309-405; March, J., Advanced Organic Chemistry, Wiley Interscience, 1985, pp. 445, 469, Carey, F. Sundberg, R., Advanced Organic Chemistry, Part B, Plenum Press, New York, 1990, pp. 686-89; Atherton, E.,
30 Sheppard, R., Solid Phase Peptide Synthesis, Pierce Chemical, 1989, pp. 3-4; Grant, G., Synthetic Peptides, W. H. Freeman & Co. 1992, pp. 77-103; Stewart, J., Young, J., Solid Phase Peptide Synthesis, 2nd Edition, IRL Press, 1984, pp. 3,5,11,14-18, 28-29. Bodansky, M., Principles of Peptide Synthesis, Springer-Verlag, 1988, pp. 62, 203, 59-69; Bodansky, M., Peptide Chemistry, Springer-Verlag, 1988, pp. 74-
35 81, Bodansky, M., Bodansky, A., The Practice of Peptide Synthesis, Springer-Verlag, 1984, pp. 9-32.

Examples of protease inhibitors for use herein are: CH₃SO₂Phe-Gly-Ala-Leu-H, CH₃SO₂Val-Ala-Leu-H, C₆H₅CH₂O(OH)(O)P-Val-Ala-Leu-H, C₆H₅CH₂O(OH)(O)P-Val-Ala-Leu-CF₃, CH₃CH₂SO₂-Phe-Gly-Ala-Leu-H, C₆H₅CH₂SO₂-Val-Ala-Leu-H, C₆H₅CH₂O(OH)(O)P-Leu-Ala-Leu-H, 5 C₆H₅CH₂O(OH)(O)P-Phe-Ala-Leu-H, CH₃O(OH)(O)P-Leu-Gly-Ala-Leu-H.

In the Synthesis Examples hereinafter methods are disclosed to synthesize certain of these peptide protease inhibitors.

Synthesis Example 1

Synthesis of the tripeptide trifluoromethylketone Moc-Phe-Gly-Ala-LeuCF₃

10

(a) N-trityl-leucine methyl ester: To a solution of 2.50 g (13.8 mmol) of Leu-OMe.HCl in 100 ml CH₂Cl₂ is added 3.86 ml TEA (27.5 mmol) dropwise. After the addition is complete 3.76 g (13.5 mmol) of triphenylmethyl chloride in 15 ml CH₂Cl₂ is added dropwise. The mixture is stirred for 4 H. The solution is diluted 15 with 5% EtOAc/petroleum ether and washed with water. The organic phase is dried (MgSO₄) filtered and the solvent removed. The residue is chromatographed on silica to give 4.8 g of pure product (90% yield).

(b) N-trityl-leucinal: To a cold (0°) solution of 4.70 g (12.2 mmol) of N-trityl-leucine methyl ester in 100ml THF is added 28.1 ml of a 1.5M solution of diisobutylaluminum hydride (42.2 mol) in THF dropwise. The solution is stirred for 6 h at this temperature and the reaction quenched with saturated Na-K tartrate, extracted with EtOAc, dried (MgSO₄), filtered and the solvent is removed. Recovered 4.13 g of the desired material that is used without purification. To a 25 solution fo 1.29 g (14.9 mmol) of oxalyl chloride in 20 ml CH₂Cl₂ at -78°C is added 2.26 ml DMSO (29.8 mmol) in 5 ml CH₂Cl₂ dropwise. After the addition is complete, 4.13 g (11.5 mmol) of crude N-trityl-leucinol in 10 ml CH₂Cl₂ is added. The solution is warmed to 0°C and poured into a mixture of water and ether. The phases are separated and the ether phase dried (MgSO₄) and evaporated to afford 30 1.37 g of the desired compound.

(c) 5-Methyl-3-tritylamino-1,1,1-trifluoro-2-hexanol: To a solution of 1.37 g (3.83 mmol) of N-trityl-leucinal and 0.653 ml (4.59 mmol) of CF₃TMS in THF is added 0.121 g (0.383 mmol) of tetrabutylammonium fluoride trihydrate in one portion. The 35 solution is stirred for 3 h at room temperature and the solvent removed. The residue is dissolved in EtOAc, washed with water, dried (MgSO₄), and the solvent removed

to afford 1.20 g of the product that is chromatographed on silica (0.760 g pure product).

5 (d) 3-(N-(Cbz-Gly-Ala))-5-methyl-1,1,1-trifluoro-2-hexanol: To a solution of 1.21 g (2.83 mmol) of 5-methyl-3-tritylamino-1,1,1-trifluoro-2-hexanol in 10 ml dioxane is added 5 ml of 4.0 M HCl in dioxane. The solution is stirred for 2 h at room temperature and the solvent removed. The residue is triturated with ether and the solid material filtered. The resulting HCl salt (0.627 g, 2.83 mmol) is suspended in 10 ml CH₂Cl₂ and Z-Gly-Ala-OH added (0.793 g, 2.83 mmol). To this mixture is added 0.870 ml (6.23 mmol) TEA followed immediately by the addition of 0.473 ml (3.12 mmol) of DEPC. The mixture is stirred overnight and the solvent removed. The residue is dissolved in EtOAc and washed with 1N HCl, saturated NaHCO₃, and brine. The solution of product is dried (MgSO₄), filtered and the solvent removed to give 1.06 g product.

15 (e) 3-(N-(Ms-Phe-Gly-Ala))-5-methyl-1,1,1-trifluoro-2-hexanol: To a solution of 1.06 g (2.37 mmol) of 3-(N-Cbz-Gly-Ala)-5-methyl-1,1,1-trifluoro-2-hexanol in 5 ml of MeOH is added 0.35 g Pd/C. The slurry is degassed and hydrogenated under a positive pressure of hydrogen overnight. The slurry is filtered through Celite and the solvent removed. The residue is dissolved in CH₂Cl₂ and 0.898 g (22.37 mmol) of Ms-Phe-OH added. To this mixture is added 0.732 ml (5.22 mmol) of TEA, followed by the addition of 0.395 ml (2.61 mmol) of DEPC. The solution is stirred overnight and the solvent removed. The residue is chromatographed on silica to afford 0.720 g pure product.

25 (f) Ms-Phe-Gly-Ala-LeuCF₃: To a slurry of 1.59 g (3.75 mmol) of Dess-Martin periodinane in 15 ml CH₂Cl₂ is added 0.650 g (1.25 mmol) of 3-(N-Ms-Phe-Gly-Ala)-5-methyl-1,1,1-trifluoro-2-hexanol in 5 ml CH₂Cl₂ and the slurry stirred for 3 h. To this mixture is added 6.51 g (25.2 mmol) of Na₂S₂O₃ in saturated NaHCO₃ and the resulting solution stirred for 10 min. The solution is extracted with EtOAc and the organic phase dried (MgSO₄), filtered and the solvent removed. The residue is chromatographed on silica to afford 0.445 g of pure product.

35 Synthesis Example 2:
Synthesis of Ms-Phe-Gly-Ala-LeuH

- (a) Ms-Phe-Gly-OH: To a solution of 2.0 g (9.0 mmol) Phe-Gly-OH, which is dissolved in 9 ml 1N NaOH and cooled to 0°C, is added simultaneously 0.766 ml (9.9 mmol) of methane sulfonyl chloride and 9 ml 1N NaOH, in separate addition funnels. After addition is complete the reaction is stirred 15 minutes at 0°C and 1 h at room temperature. At this point the solution is cooled to 0°C, the pH adjusted to 9.5 and is washed with EtOAc (1X, 50 ml). The aqueous phase (0°C) is then adjusted to pH = 2.5 (2N HCl) and extracted with EtOAc (3X, 50 ml), dried (MgSO₄), filtered, and the solvent removed to afford 2.0 g pure product.
- 10 (b) Ms-Phe-Gly-Ala-Leucinol: A solution of is prepared by dissolving 0.500 g (1.67 mmol) N-Ms-Phe-Gly-OH in 15 ml THF, cooling to -15°C, and adding 0.366 ml (3.33 mmol) NMM followed by 0.216 ml (1.67 mmol) isobutyl chloroformate. This solution is stirred 5 minutes and 0.374 g (1.67 mmol) Ala-Leucinol.HCl, in a mixture of 10 ml THF and minimal DMF, are added. Stirring is continued at 0°C for
- 15 15 minutes and 2 h at room temperature. The solution is quenched with 5 ml 1N HCl, extracted with EtOAc (3X, 50 ml), the combined extracts are washed with sat'd NaHCO₃ and sat'd NaCl. The resulting organic phase is then dried (MgSO₄), filtered, evaporated and chromatographed on silica to yield 0.260 g of the desired material.
- 20 (c) Ms-Phe-Gly-Ala-LeuH: A solution is prepared by adding 0.337 g (0.798 mmol) Dess-Martin periodinane to 5 ml CH₂Cl₂ and stirring for 10 minutes. To this solution 0.125 g (0.266 mmol) N-Ms-Phe-Gly-Ala-Leucinol is added in one portion. The reaction is continued until TLC showed complete conversion at which time the
- 25 solution is poured into 25 ml sat'd NaHCO₃ containing 1.8 g (5.586 mmol) Na₂S₂O₃. After stirring for 10 minutes the mixture is extracted with EtOAc (3X, 50 ml). The combined extracts are dried (MgSO₄), evaporated, and chromatographed on silica to afford 0.048 g of the product.
- 30 Gly = glycine
Ala = alanine
Leu = leucine
Phe = phenylalanine
OMe = methyl ester
- 35 TEA = triethylamine
DECP = diethylcyanophosphonate
TLC = thin layer chromatography

MeOH = methanol

Pd/C = palladium on activated carbon

EtOH = ethanol

THF = tetrahydrofuran

5 Ms = methanesulfonyl

Proteolytic Enzyme - Another essential ingredient in the present liquid detergent compositions is active proteolytic enzyme. Mixtures of proteolytic enzyme are also included. The proteolytic enzyme can be of animal, vegetable or
10 microorganism (preferred) origin. The proteases for use in the detergent compositions herein include (but are not limited to) trypsin, subtilisin, chymotrypsin and elastase-type proteases. Preferred for use herein are subtilisin-type proteolytic enzymes. Particularly preferred is bacterial serine proteolytic enzyme obtained from Bacillus subtilis and/or Bacillus licheniformis. Protease enzymes are usually present
15 in such liquid detergent compositions at levels sufficient to provide from 0.005 to 0.1 Anson units (AU) of activity per gram of composition.

Suitable proteolytic enzymes include Novo Industri A/S Alcalase[®] (preferred), Esperase[®], Savinase[®] (Copenhagen, Denmark), Gist-brocades' Maxatase[®], Maxacal[®] and Maxapem 15[®] (protein engineered Maxacal[®]) (Delft, Netherlands),
20 and subtilisin BPN and BPN' (preferred), which are commercially available. Preferred proteolytic enzymes are also modified bacterial serine proteases, such as those made by Genencor International, Inc. (San Francisco, California) which are described in European Patent 251,446, filed April 28, 1987 (particularly pages 17, 24 and 98), and which is called herein "Protease B", and U.S. Patent 5,030,378,
25 Venegas, issued July 9, 1991, which refers to a modified bacterial serine proteolytic enzyme (Genencor International) which is called "Protease A" herein (same as BPN'). In particular see columns 2 and 3 of U.S. Patent 5,030,378 for a complete description, including amino sequence, of Protease A and its variants. Preferred proteolytic enzymes, then, are selected from the group consisting of Alcalase[®]
30 (Novo Industri A/S), BPN', Protease A and Protease B (Genencor), and mixtures thereof. Protease B is most preferred.

Another preferred protease, referred to as "Protease D" is a carbonyl hydrolase variant having an amino acid sequence not found in nature, which is derived from a precursor carbonyl hydrolase by substituting a different amino acid
35 for a plurality of amino acid residues at a position in said carbonyl hydrolase equivalent to position +76, preferably also in combination with one or more amino acid residue positions equivalent to those selected from the group consisting of +99,

+101, +103, +104, +107, +123, +27, +105, +109, +126, +128, +135, +156, +166, +195, +197, +204, +206, +210, +216, +217, +218, +222, +260, +265, and/or +274 according to the numbering of *Bacillus amyloliquefaciens* subtilisin, as described in WO 95/10615 published April 20, 1995 by Genencor International.


5 Useful proteases are also described in PCT publications: WO 95/30010 published November 9, 1995 by The Procter & Gamble Company; WO 95/30011 published November 9, 1995 by The Procter & Gamble Company; WO 95/29979 published November 9, 1995 by The Procter & Gamble Company.

10 Deterative Surfactant - An effective amount, typically from about 1 to 95, preferably about 8 to 70, weight %, of deterative surfactant is yet another essential ingredient in the present invention. The deterative surfactant can be selected from the group consisting of anionics, nonionics, cationics, ampholytics, zwitterionics, and mixtures thereof. By selecting the type and amount of deterative surfactant, along with other adjunct ingredients disclosed herein, the present detergent
15 compositions can be formulated to be used in the context of laundry cleaning or in other different cleaning applications, particularly including dishwashing. The particular surfactants used can therefore vary widely depending upon the particular end-use envisioned.

The benefits of the present invention are especially pronounced in
20 compositions containing ingredients that are harsh to enzymes such as certain detergency builders and surfactants. These include (but are not limited to) anionic surfactants such as alkyl ether sulfate linear alkyl benzene sulfonate, alkyl sulfate, etc. Suitable surfactants are described below.

25 Anionic Surfactants - One type of anionic surfactant which can be utilized encompasses alkyl ester sulfonates. These are desirable because they can be made with renewable, non-petroleum resources. Preparation of the alkyl ester sulfonate surfactant component can be effected according to known methods disclosed in the technical literature. For instance, linear esters of C₈-C₂₀ carboxylic acids can be sulfonated with gaseous SO₃ according to "The Journal of the American Oil
30 Chemists Society," 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm, and coconut oils, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprises alkyl ester sulfonate surfactants of the structural formula:



35 wherein R³ is a C₈-C₂₀ hydrocarbyl, preferably an alkyl, or combination thereof, R⁴ is a C₁-C₆ hydrocarbyl, preferably an alkyl, or combination thereof, and M is a soluble salt-forming cation. Suitable salts include metal salts such as sodium,

potassium, and lithium salts, and substituted or unsubstituted ammonium salts, such as methyl-, dimethyl-, trimethyl-, and quaternary ammonium cations, e.g.

tetramethyl-ammonium and dimethyl piperdinium, and cations derived from alkanolamines, e.g. monoethanol-amine, diethanolamine, and triethanolamine.

- 5 Preferably, R^3 is C_{10} - C_{16} alkyl, and R^4 is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R^3 is C_{14} - C_{16} alkyl.

Alkyl sulfate surfactants are another type of anionic surfactant of importance for use herein. In addition to providing excellent overall cleaning ability when used in combination with polyhydroxy fatty acid amides (see below), including good
10 grease/oil cleaning over a wide range of temperatures, wash concentrations, and wash times, dissolution of alkyl sulfates can be obtained, as well as improved formulability in liquid detergent formulations are water soluble salts or acids of the formula $ROSO_3M$ wherein R preferably is a C_{10} - C_{24} hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C_{10} - C_{20} alkyl component, more preferably a C_{12} -
15 C_{18} alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g., sodium, potassium, lithium), substituted or unsubstituted ammonium cations such as methyl-, dimethyl-, and trimethyl ammonium and quaternary ammonium cations, e.g., tetramethyl-ammonium and dimethyl piperdinium, and cations derived from alkanolamines such as ethanolamine, diethanolamine, triethanolamine, and mixtures thereof, and the like. Typically, alkyl chains of C_{12} - C_{16} are preferred for lower wash
20 temperatures (e.g., below about 50°C) and C_{16} - C_{18} alkyl chains are preferred for higher wash temperatures (e.g., above about 50°C).

Alkyl alkoxyated sulfate surfactants are another category of useful anionic surfactant. These surfactants are water soluble salts or acids typically of the formula
25 $RO(A)_mSO_3M$ wherein R is an unsubstituted C_{10} - C_{24} alkyl or hydroxyalkyl group having a C_{10} - C_{24} alkyl component, preferably a C_{12} - C_{20} alkyl or hydroxyalkyl, more preferably C_{12} - C_{18} alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a
30 metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl-, trimethyl-ammonium and quaternary ammonium cations, such as tetramethyl-ammonium, dimethyl
35 piperidinium and cations derived from alkanolamines, e.g. monoethanolamine, diethanolamine, and triethanolamine, and mixtures thereof. Exemplary surfactants are C_{12} - C_{18} alkyl polyethoxylate (1.0) sulfate, C_{12} - C_{18} alkyl polyethoxylate (2.25)

sulfate, C₁₂-C₁₈ alkyl polyethoxylate (3.0) sulfate, and C₁₂-C₁₈ alkyl polyethoxylate (4.0) sulfate wherein M is conveniently selected from sodium and potassium.

Other Anionic Surfactants - Other anionic surfactants useful for deterative purposes can also be included in the compositions hereof. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono-, di- and triethanolamine salts) of soap, C₉-C₂₀ linear alkylbenzenesulphonates, C₈-C₂₂ primary or secondary alkanesulphonates, C₈-C₂₄ olefinsulphonates, sulphonated polycarboxylic acids prepared by sulphonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isothionates such as the acyl isothionates, N-acyl taurates, fatty acid amides of methyl tauride, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinate (especially saturated and unsaturated C₁₂-C₁₈ monoesters) diesters of sulfosuccinate (especially saturated and unsaturated C₆-C₁₄ diesters), N-acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, alkyl polyethoxy carboxylates such as those of the formula RO(CH₂CH₂O)_kCH₂COO-M⁺ wherein R is a C₈-C₂₂ alkyl, k is an integer from 0 to 10, and M is a soluble salt-forming cation, and fatty acids esterified with isethionic acid and neutralized with sodium hydroxide. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

Nonionic Detergent Surfactants - Suitable nonionic detergent surfactants are generally disclosed in U.S. Patent 3,929,678, Laughlin et al., issued December 30, 1975, at column 13, line 14 through column 16, line 6, incorporated herein by reference. Exemplary, non-limiting classes of useful nonionic surfactants are listed below.

The polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols. In general, the polyethylene oxide condensates are preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 12 carbon atoms in either a straight chain or

branched chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 5 to about 25 moles of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal[®] CO-630, marketed by the GAF Corporation; and Triton[®] X-45, X-114, X-100, and X-102, all marketed by the Rohm & Haas Company. These compounds are commonly referred to as alkyl phenol alkoxylates, (e.g., alkyl phenol ethoxylates).


The condensation products of aliphatic alcohols with from about 1 to about 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from about 10 to about 20 carbon atoms with from about 2 to about 18 moles of ethylene oxide per mole of alcohol. Examples of commercially available nonionic surfactants of this type include Tergitol[®] 15-S-9 (the condensation product of C₁₁-C₁₅ linear secondary alcohol with 9 moles ethylene oxide), Tergitol[®] 24-L-6 NMW (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; Neodol[®] 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), Neodol[®] 23-6.5 (the condensation product of C₁₂-C₁₃ linear alcohol with 6.5 moles of ethylene oxide), Neodol[®] 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), Neodol[®] 45-4 (the condensation product of C₁₄-C₁₅ linear alcohol with 4 moles of ethylene oxide), marketed by Shell Chemical Company, and Kyro[®] EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company. This category of nonionic surfactant is referred to generally as "alkyl ethoxylates."

The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol. The hydrophobic portion of these compounds preferably has a molecular weight of from about 1500 to about 1800 and exhibits water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially-available Pluronic[®] surfactants, marketed by BASF.

The condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetric® compounds, marketed by BASF.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; water-soluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.


These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides.

Alkylpolysaccharides disclosed in U.S. Patent 4,565,647, Llenado, issued January 21, 1986, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g., a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to

about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6- positions on the preceding saccharide units.


Optionally, and less desirably, there can be a polyalkylene-oxide chain joining the hydrophobic moiety and the polysaccharide moiety. The preferred alkyleneoxide is ethylene oxide. Typical hydrophobic groups include alkyl groups, either saturated or unsaturated, branched or unbranched containing from about 8 to about 18, preferably from about 10 to about 16, carbon atoms. Preferably, the alkyl group is a straight chain saturated alkyl group. The alkyl group can contain up to about 3 hydroxy groups and/or the polyalkyleneoxide chain can contain up to about 10, preferably less than 5, alkyleneoxide moieties. Suitable alkyl polysaccharides are octyl, nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl, and octadecyl, di-, tri-, tetra-, penta-, and hexaglycosides, galactosides, lactosides, glucoses, fructosides, fructoses and/or galactoses. Suitable mixtures include coconut alkyl, di-, tri-, tetra-, and pentaglycosides and tallow alkyl tetra-, penta-, and hexa-glycosides.

The preferred alkylpolyglycosides have the formula



wherein R^2 is selected from the group consisting of alkyl, alkyl-phenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4- and/or 6-position, preferably predominantly the 2-position.

Fatty acid amide surfactants having the formula:




wherein R^6 is an alkyl group containing from about 7 to about 21 (preferably from about 9 to about 17) carbon atoms and each R^7 is selected from the group consisting

of hydrogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, and -(C₂H₄O)_xH where x varies from about 1 to about 3.

Preferred amides are C₈-C₂₀ ammonia amides, monoethanolamides, diethanolamides, and isopropanolamides.

- 5 Cationic Surfactants - Cationic deterative surfactants can also be included in detergent compositions of the present invention. Cationic surfactants include the ammonium surfactants such as alkyldimethylammonium halogenides, and those surfactants having the formula:



- 10 wherein R² is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R³ is selected from the group consisting of -CH₂CH₂-, -CH₂CH(CH₃)-, -CH₂CH(CH₂OH)-, -CH₂CH₂CH₂-, and mixtures thereof; each R⁴ is selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, benzyl, ring structures formed by joining the two R⁴ groups, -
15 CH₂CHOHCHOHCOR⁶CHOH-CH₂OH wherein R⁶ is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not O; R⁵ is the same as R⁴ or is an alkyl chain wherein the total number of carbon atoms of R² plus R⁵ is not more than about 18; each y is from 0 to about 10 and the sum of the y values is from 0 to about 15; and X is any compatible anion.

- 20 Other cationic surfactants useful herein are also described in U.S. Patent 4,228,044, Cambre, issued October 14, 1980, incorporated herein by reference.

- Other Surfactants - Ampholytic surfactants can be incorporated into the detergent compositions hereof. These surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of
25 heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight chain or branched. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-solubilizing group, e.g., carboxy, sulfonate, sulfate. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column
30 19, lines 18-35 for examples of ampholytic surfactants.

- Zwitterionic surfactants can also be incorporated into the detergent compositions hereof. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary
35 sulfonium compounds. See U.S. Patent No. 3,929,678 to Laughlin et al., issued December 30, 1975 at column 19, line 38 through column 22, line 48 for examples

of zwitterionic surfactants. Ampholytic and zwitterionic surfactants are generally used in combination with one or more anionic and/or nonionic surfactants.

Polyhydroxy Fatty Acid Amide Surfactant - The liquid detergent compositions hereof may also contain an enzyme-enhancing amount of polyhydroxy fatty acid amide surfactant. By "enzyme-enhancing" is meant that the formulator of the composition can select an amount of polyhydroxy fatty acid amide to be incorporated into the compositions that will improve enzyme cleaning performance of the detergent composition. In general, for conventional levels of enzyme, the incorporation of about 1%, by weight, polyhydroxy fatty acid amide will enhance enzyme performance.

The detergent compositions herein will typically comprise about 1% weight basis, polyhydroxy fatty acid amide surfactant, preferably from about 3% to about 30%, of the polyhydroxy fatty acid amide. The polyhydroxy fatty acid amide surfactant component comprises compounds of the structural formula:



wherein: R^1 is H, C_1 - C_4 hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, or a mixture thereof, preferably C_1 - C_4 alkyl, more preferably C_1 or C_2 alkyl, most preferably C_1 alkyl (i.e., methyl); and R^2 is a C_5 - C_{31} hydrocarbyl, preferably straight chain C_7 - C_{19} alkyl or alkenyl, more preferably straight chain C_9 - C_{17} alkyl or alkenyl, most preferably straight chain C_{11} - C_{15} alkyl or alkenyl, or mixtures thereof; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxylated or propoxylated) thereof. Z preferably will be derived from a reducing sugar in a reductive amination reaction; more preferably Z will be a glycityl. Suitable reducing sugars include glucose, fructose, maltose, lactose, galactose, mannose, and xylose. As raw materials, high dextrose corn syrup, high fructose corn syrup, and high maltose corn syrup can be utilized as well as the individual sugars listed above. These corn syrups may yield a mix of sugar components for Z. It should be understood that it is by no means intended to exclude other suitable raw materials. Z preferably will be selected from the group consisting of $-CH_2-(CHOH)_n-CH_2OH$, $-CH(CH_2OH)-(CHOH)_{n-1}-CH_2OH$, $-CH_2-(CHOH)_2(CHOR')(CHOH)-CH_2OH$, and alkoxyated derivatives thereof, where n is an integer from 3 to 5, inclusive, and R' is H or a cyclic or aliphatic monosaccharide. Most preferred are glycityls wherein n is 4, particularly $-CH_2-(CHOH)_4-CH_2OH$.

R^1 can be, for example, N-methyl, N-ethyl, N-propyl, N-isopropyl, N-butyl, N-2-hydroxy ethyl, or N-2-hydroxy propyl.

$R^2-CO-N<$ can be, for example, cocamide, stearamide, oleamide, lauramide, myristamide, capricamide, palmitamide, tallowamide, etc.

Z can be 1-deoxyglucityl, 2-deoxyfructityl, 1-deoxymaltityl, 1-deoxylactityl, 1-deoxygalactityl, 1-deoxymannityl, 1-deoxymaltotriosityl, etc.

5 Methods for making polyhydroxy fatty acid amides are known in the art. In general, they can be made by reacting an alkyl amine with a reducing sugar in a reductive amination reaction to form a corresponding N-alkyl polyhydroxyamine, and then reacting the N-alkyl polyhydroxyamine with a fatty aliphatic ester or triglyceride in a condensation/amidation step to form the N-alkyl, N-polyhydroxy
10 fatty acid amide product. Processes for making compositions containing polyhydroxy fatty acid amides are disclosed, for example, in G.B. Patent Specification 809,060, published February 18, 1959, by Thomas Hedley & Co., Ltd., U.S. Patent 2,965,576, issued December 20, 1960 to E. R. Wilson, and U.S. Patent 2,703,798, Anthony M. Schwartz, issued March 8, 1955, and U.S. Patent 1,985,424,
15 issued December 25, 1934 to Piggott, each of which is incorporated herein by reference.

Second Enzyme - Preferred compositions herein further comprise a performance-enhancing amount of a detergent-compatible second enzyme. By "detergent-compatible" is meant compatibility with the other ingredients of a liquid
20 detergent composition, such as deterative surfactant and detergency builder. These second enzymes are preferably selected from the group consisting of lipase, amylase, cellulase, and mixtures thereof. The term "second enzyme" excludes the proteolytic enzymes discussed above, so each composition which has a second enzyme contains at least two kinds of enzyme, including at least one proteolytic
25 enzyme. The amount of second enzyme used in the composition varies according to the type of enzyme. In general, from about 0.0001 to 0.3, more preferably 0.001 to 0.1, weight % of these second enzymes are preferably used. Mixtures of the same class of enzymes (e.g. lipase) or two or more classes (e.g. cellulase and lipase) may be used. Purified or non-purified forms of the enzyme may be used.

30 Any lipolytic enzyme suitable for use in a liquid detergent composition can be used in these compositions. Suitable lipase enzymes for use herein include those of bacterial and fungal origin.

Suitable bacterial lipases include those produced by microorganisms of the Pseudomonas groups, such as Pseudomonas stutzeri ATCC 19.154, as disclosed in
35 British Patent 1,372,034, incorporated herein by reference. Suitable lipases include those which show a positive immunological cross-reaction with the antibody of the lipase produced by the microorganism Pseudomonas fluorescens IAM 1057. This

lipase and a method for its purification have been described in Japanese Patent Application 53-20487, laid open on February 24, 1978. This lipase is available from Amano Pharmaceutical Co. Ltd., Nagoya, Japan, under the trade name Lipase P "Amano," hereinafter referred to as "Amano-P." Such lipases should show a
5 positive immunological cross-reaction with the Amano-P antibody, using the standard and well-known immunodiffusion procedure according to Ouchterlony (Acta. Med. Scan., 133, pages 76-79 (1950)). These lipases, and a method for their immunological cross-reaction with Amano-P, are also described in U.S. Patent 4,707,291, Thom et al., issued November 17, 1987, incorporated herein by
10 reference. Typical examples thereof are the Amano-P lipase, the lipase ex Pseudomonas fragi FERM P 1339 (available under the trade name Amano-B), lipase ex Pseudomonas nitroreducens var. lipolyticum FERM P 1338 (available under the trade name Amano-CES), lipases ex Chromobacter viscosum, e.g. Chromobacter viscosum var. lipolyticum NRRLB 3673, commercially available from Toyo Jozo
15 Co., Tagata, Japan; and further Chromobacter viscosum lipases from U.S. Biochemical Corp., U.S.A. and Disoynt Co., The Netherlands, and lipases ex Pseudomonas gladioli.

Suitable fungal lipases include those producible by Humicola lanuginosa and Thermomyces lanuginosus. Most preferred is lipase obtained by cloning the gene
20 from Humicola lanuginosa and expressing the gene in Aspergillus oryzae as described in European Patent Application 0 258 068 (Novo Industri A/S), commercially available from Novo Nordisk A/S under the trade name Lipolase®.

From about 10 to 18,000, preferably about 60 to 6,000, lipase units per gram (LU/g) of lipase can be used in these compositions. A lipase unit is that amount of
25 lipase which produces 1 mmol of titratable fatty acid per minute in a pH stat, where pH is 9.0, temperature is 30°C, substrate is an emulsion of 3.3wt % of olive oil and 3.3% gum arabic, in the presence of 13 mmol/l Ca⁺⁺ and 20 mmol/l NaCl in 5 mmol/l Tris-buffer.

Any cellulase suitable for use in a liquid detergent composition can be used in
30 these compositions. Suitable cellulase enzymes for use herein include those from bacterial and fungal origins. Preferably, they will have a pH optimum of between 5 and 9.5. From about 0.0001 to 0.1 weight % cellulase can be used.

Suitable cellulases are disclosed in U.S. Patent 4,435,307, Barbesgaard et al., issued March 6, 1984, incorporated herein by reference, which discloses fungal
35 cellulase produced from Humicola insolens. Suitable cellulases are also disclosed in GB-A-2.075.028, GB-A-2.095.275 and DE-OS-2.247.832.

Examples of such cellulases are cellulases produced by a strain of Humicola insolens (Humicola grisea var. thermoidea), particularly the Humicola strain DSM 1800, and cellulases produced by a fungus of Bacillus N or a cellulase 212-producing fungus belonging to the genus Aeromonas, and cellulase extracted from
5 the hepatopancreas of a marine mollusc (Dolabella Auricula Solander).

Any amylase suitable for use in a liquid detergent composition can be used in these compositions. Amylases include, for example, amylases obtained from a special strain of B.licheniformis, described in more detail in British Patent Specification No. 1,296,839 (Novo). Amylolytic proteins include, for example,
10 Rapidase^R, International Bio-Synthetics, Inc. and Termamyl^R Novo Industries.

From about 0.0001% to 0.55, preferably 0.0005 to 0.1, wt. % amylase can be used.

Calcium - The compositions herein may optionally comprise a calcium ion source. Any water-soluble calcium salt can be used as a source of calcium ions,
15 including calcium acetate, calcium formate, calcium xylene sulfonate, and calcium propionate. Divalent ions, such as zinc and magnesium ions, can replace the calcium ion completely or in part. Thus in the liquid detergent compositions herein, the source of calcium ions can be partially substituted with a source of another divalent ion.

The calcium useful herein is enzyme-accessible. Therefore, the preferred
20 compositions are substantially free of sequestrants, for example, polyacids capable of forming calcium complexes which are soluble in the composition. However, minor amounts of sequestrants such as polyacids or mixtures of polyacids can be used. The enzyme-accessible calcium is defined as the amount of calcium-ions
25 effectively available to the enzyme component. From a practical standpoint the enzyme-accessible calcium is therefore the soluble calcium in the composition in the absence of any storage sequestrants, e.g., having an equilibrium constant of complexation with calcium equal to or greater than 1.5 at 20°C.

Boric Acid - The compositions herein also optionally contain from about
30 0.25% to about 10%, preferably from about 0.5% to about 5%, more preferably from about 0.75% to about 3%, by weight of boric acid or a compound capable of forming boric acid in the composition (calculated on the basis of the boric acid). Boric acid is preferred, although other compounds such as boric oxide, borax and other alkali metal borates (e.g., sodium ortho-, meta-, pyroborate, an sodium
35 pentaborate) are suitable. Substituted boric acids (e.g., phenylboronic acid, butane boronic acid, and p-bromo phenylboronic acid) can also be used in place of boric acid.

The compositions of the present invention can also contain polyols, especially diols, containing only carbon, hydrogen and oxygen atoms. They preferably contain from about 2 to about 6 hydroxy groups. Examples include propylene glycol (especially 1,2 propanediol, which is preferred), ethylene glycol, glycerol, sorbitol, mannitol, glucose, and mixtures thereof. The polyol generally represents from about 1% to about 15%, preferably from about 1.5% to about 10%, more preferably from about 2% to about 7%, by weight of the composition.

Optional Ingredients - Detergent builders can optionally be included in the compositions herein, especially for laundry compositions. Inorganic as well as organic builders can be used. When present, the compositions will typically comprise at least about 1% builder and can be either an inorganic or organic builder. Liquid laundry formulations preferably comprise from about 3% to 30%, more preferably about 5 to 20%, by weight, of detergent builder.


Inorganic detergent builders include, but are not limited to, the alkali metal, ammonium and alkanolammonium salts of polyphosphates (exemplified by the tripolyphosphates, pyrophosphates, and glassy polymeric meta-phosphates), phosphonates, phytic acid, silicates, carbonates (including bicarbonates and sesquicarbonates), sulphates, and aluminosilicates. Borate builders, as well as builders containing borate-forming materials that can produce borate under detergent storage or wash conditions (hereinafter, collectively "borate builders"), can also be used. Preferably, non-borate builders are used in the compositions of the invention intended for use at wash conditions less than about 50°C, especially less than about 40°C.

Examples of silicate builders are the alkali metal silicates, particularly those having a $\text{SiO}_2:\text{Na}_2\text{O}$ ratio in the range 1.6:1 to 3.2:1 and layered silicates, such as the layered sodium silicates described in U.S. Patent 4,664,839, issued May 12, 1987 to H. P. Rieck, incorporated herein by reference. However, other silicates may also be useful such as for example magnesium silicate, which can serve as a crispening agent in granular formulations, as a stabilizing agent for oxygen bleaches, and as a component of suds control systems.

Examples of carbonate builders are the alkaline earth and alkali metal carbonates, including sodium carbonate and sesquicarbonate and mixtures thereof with ultra-fine calcium carbonate as disclosed in German Patent Application No. 2,321,001 published on November 15, 1973, the disclosure of which is incorporated herein by reference.

Aluminosilicate builders are useful in the present invention. Aluminosilicate builders are of great importance in most currently marketed heavy duty granular

detergent compositions, and can also be a significant builder ingredient in liquid detergent formulations. Aluminosilicate builders include those having the empirical formula:




5 wherein M is sodium, potassium, ammonium or substituted ammonium, z is from about 0.5 to about 2; and y is 1; this material having a magnesium ion exchange capacity of at least about 50 milligram equivalents of CaCO₃ hardness per gram of anhydrous aluminosilicate. Preferred aluminosilicates are zeolite builders which have the formula:



wherein z and y are integers of at least 6, the molar ratio of z to y is in the range from 1.0 to about 0.5, and x is an integer from about 15 to about 264.

Useful aluminosilicate ion exchange materials are commercially available. These aluminosilicates can be crystalline or amorphous in structure and can be
 15 naturally-occurring aluminosilicates or synthetically derived. A method for producing aluminosilicate ion exchange materials is disclosed in U.S. Patent 3,985,669, Krummel, et al., issued October 12, 1976, incorporated herein by reference. Preferred synthetic crystalline aluminosilicate ion exchange materials useful herein are available under the designations Zeolite A, Zeolite P (B), and
 20 Zeolite X. In an especially preferred embodiment, the crystalline aluminosilicate ion exchange material has the formula:



wherein x is from about 20 to about 30, especially about 27. This material is known as Zeolite A. Preferably, the aluminosilicate has a particle size of about 0.1-10
 25 microns in diameter.

Specific examples of polyphosphates are the alkali metal triphosphates, sodium, potassium and ammonium pyrophosphate, sodium and potassium and ammonium pyrophosphate, sodium and potassium orthophosphate, sodium
 30 polymeta phosphate in which the degree of polymerization ranges from about 6 to about 21, and salts of phytic acid.

Examples of phosphonate builder salts are the water-soluble salts of ethane 1-hydroxy-1, 1-diphosphonate particularly the sodium and potassium salts, the water-soluble salts of methylene diphosphonic acid e.g. the trisodium and tripotassium salts and the water-soluble salts of substituted methylene diphosphonic acids, such
 35 as the trisodium and tripotassium ethylidene, isopropylidene benzylmethylidene and halo methylidene phosphonates. Phosphonate builder salts of the aforementioned types are disclosed in U.S. Patent Nos. 3,159,581 and 3,213,030


issued December 1, 1964 and October 19, 1965, to Diehl; U.S. Patent No. 3,422,021 issued January 14, 1969, to Roy; and U.S. Patent Nos. 3,400,148 and 3,422,137 issued September 3, 1968, and January 14, 1969 to Quimby, said disclosures being incorporated herein by reference.

5 Organic detergent builders preferred for the purposes of the present invention include a wide variety of polycarboxylate compounds. As used herein, "polycarboxylate" refers to compounds having a plurality of carboxylate groups, preferably at least 3 carboxylates.

 Polycarboxylate builder can generally be added to the composition in acid
10 form, but can also be added in the form of a neutralized salt. When utilized in salt form, alkali metals, such as sodium, potassium, and lithium, or alkanolammonium salts are preferred.

 Included among the polycarboxylate builders are a variety of categories of useful materials. One important category of polycarboxylate builders encompasses
15 the ether polycarboxylates. A number of ether polycarboxylates have been disclosed for use as detergent builders. Examples of useful ether polycarboxylates include oxydisuccinate, as disclosed in Berg, U.S. Patent 3,128,287, issued April 7, 1964, and Lamberti et al., U.S. Patent 3,635,830, issued January 18, 1972, both of which are incorporated herein by reference.


20 A specific type of ether polycarboxylates useful as builders in the present invention also include those having the general formula:



 wherein A is H or OH; B is H or -O-CH(COOX)-CH₂(COOX); and X is H or a salt-forming cation. For example, if in the above general formula A and B are both H,
25 then the compound is oxydisuccinic acid and its water-soluble salts. If A is OH and B is H, then the compound is tartrate monosuccinic acid (TMS) and its water-soluble salts. If A is H and B is -O-CH(COOX)-CH₂(COOX), then the compound is tartrate disuccinic acid (TDS) and its water-soluble salts. Mixtures of these builders are especially preferred for use herein. Particularly preferred are mixtures of TMS and
30 TDS in a weight ratio of TMS to TDS of from about 97:3 to about 20:80. These builders are disclosed in U.S. Patent 4,663,071, issued to Bush et al., on May 5, 1987.

 Suitable ether polycarboxylates also include cyclic compounds, particularly alicyclic compounds, such as those described in U.S. Patents 3,923,679; 3,835,163;
35 4,158,635; 4,120,874 and 4,102,903, all of which are incorporated herein by reference.

Other useful detergency builders include the ether hydroxypolycarboxylates represented by the structure:



wherein M is hydrogen or a cation wherein the resultant salt is water-soluble, preferably an alkali metal, ammonium or substituted ammonium cation, n is from about 2 to about 15 (preferably n is from about 2 to about 10, more preferably n averages from about 2 to about 4) and each R is the same or different and selected from hydrogen, C₁₋₄ alkyl or C₁₋₄ substituted alkyl (preferably R is hydrogen).

Still other ether polycarboxylates include copolymers of maleic anhydride with ethylene or vinyl methyl ether, 1, 3, 5-trihydroxy benzene-2, 4, 6-trisulphonic acid, and carboxymethyloxysuccinic acid.

Organic polycarboxylate builders also include the various alkali metal, ammonium and substituted ammonium salts of polyacetic acids. Examples include the sodium, potassium, lithium, ammonium and substituted ammonium salts of ethylenediamine tetraacetic acid, and nitrilotriacetic acid.

Also included are polycarboxylates such as mellitic acid, succinic acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, and carboxymethyloxysuccinic acid, and soluble salts thereof.

Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are polycarboxylate builders of particular importance for heavy duty liquid detergent formulations, but can also be used in granular compositions.

Other carboxylate builders include the carboxylated carbohydrates disclosed in U.S. Patent 3,723,322, Diehl, issued March 28, 1973, incorporated herein by reference.

Also suitable in the detergent compositions of the present invention are the 3,3-dicarboxy-4-oxa-1,6-hexanedioates and the related compounds disclosed in U.S. Patent 4,566,984, Bush, issued January 28, 1986, incorporated herein by reference.

Useful succinic acid builders include the C₅-C₂₀ alkyl succinic acids and salts thereof. A particularly preferred compound of this type is dodeceny succinic acid. Alkyl succinic acids typically are of the general formula R-CH(COOH)CH₂(COOH) i.e., derivatives of succinic acid, wherein R is hydrocarbon, e.g., C₁₀-C₂₀ alkyl or alkenyl, preferably C₁₂-C₁₆ or wherein R may be substituted with hydroxyl, sulfo, sulfoxy or sulfone substituents, all as described in the above-mentioned patents.

The succinate builders are preferably used in the form of their water-soluble salts, including the sodium, potassium, ammonium and alkanolammonium salts.

Specific examples of succinate builders include: laurylsuccinate, myristylsuccinate, palmitylsuccinate, 2-dodecenylsuccinate (preferred), 2-pentadecenylsuccinate, and the like. Laurylsuccinates are the preferred builders of this group, and are described in European Patent Application 86200690.5/0,200,263, published November 5, 1986.

Examples of useful builders also include sodium and potassium carboxymethyloxymalonate, carboxymethyloxysuccinate, cis-cyclo-hexane-hexacarboxylate, cis-cyclopentane-tetracarboxylate, water-soluble polyacrylates (these polyacrylates having molecular weights to above about 2,000 can also be effectively utilized as dispersants), and the copolymers of maleic anhydride with vinyl methyl ether or ethylene.

Other suitable polycarboxylates are the polyacetal carboxylates disclosed in U.S. Patent 4,144,226, Crutchfield et al., issued March 13, 1979, incorporated herein by reference. These polyacetal carboxylates can be prepared by bringing together, under polymerization conditions, an ester of glyoxylic acid and a polymerization initiator. The resulting polyacetal carboxylate ester is then attached to chemically stable end groups to stabilize the polyacetal carboxylate against rapid depolymerization in alkaline solution, converted to the corresponding salt, and added to a surfactant.

Polycarboxylate builders are also disclosed in U.S. Patent 3,308,067, Diehl, issued March 7, 1967, incorporated herein by reference. Such materials include the water-soluble salts of homo- and copolymers of aliphatic carboxylic acids such as maleic acid, itaconic acid and methylenemalonic acid.

Other organic builders known in the art can also be used. For example, monocarboxylic acids, and soluble salts thereof, having long chain hydrocarbyls can be utilized. These would include materials generally referred to as "soaps." Chain lengths of C₁₀-C₂₀ are typically utilized. The hydrocarbyls can be saturated or unsaturated.

Other optional ingredients include soil release agents, chelating agents, clay soil removal/anti redeposition agents, polymeric dispersing agents, bleaches, brighteners, suds suppressors, solvents and aesthetic agents.

The detergent composition herein can be formulated as a variety of compositions, for instance as laundry detergents as well as hard surface cleaners or dishwashing compositions.

The compositions according to the present invention are further illustrated by the following examples.

EXAMPLE I

The following compositions are made by combining the listed ingredients in the listed proportions.

<u>Compositions</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>
Linear alkyl benzene sulfonic acid	8.5	15	6.5	10	12.5	4
Sodium C ₁₂₋₁₅ alkyl sulfate	1	2	1	2	--	--
C ₁₄₋₁₅ alkyl 2.5 times ethoxylated sulfate	10	5	10.5	--	11	9
C ₁₂ glucose amide	--	--	9	--	--	5
C ₁₂₋₁₅ alcohol 7 times ethoxylated	3	10	4	7	2.5	--
Fatty acid	2	5	5	4	2	2
Citric acid	6	7	4	6	4	5
C ₁₂₋₁₄ alkenyl substituted, succinic acid	--	6	--	5	--	6
Sodium hydroxide	2	6	2	4	1	1.5
Ethanol	2	1.5	2	4	2	1.5
Monoethanolamine	6	5	4	--	--	--
1,2-Propanediol	12	10	5	5	4	6
Amylase (143 KNU/g)	--	--	0.1	--	--	0.2
Lipolase [®] (100KLU/g commercial solution)	0.5	0.2	0.5	0.5	0.4	--
Protease B (34 g/L commercial solution)	0.9	--	0.5	--	1.2	--
Savinase [®] (commercial solution)	--	0.3	--	0.4	0.2	0.3
Carezyme [®]	0.5	1	0.8	--	0.2	0.8
Peptide aldehyde ¹	0.009	0.005	0.001	0.0005	0.0003	0.01
Calcium Ions	0.01	0.5	0.1	0.05	0.9	0.25
Water and minors	Balance to 100%					

- 5 1) N-(methylsulfonyl)-Phe-Gly-Ala-LeuH which is prepared according to Synthesis Example 2.

EXAMPLE II

The following formula is prepared.

<u>Component</u>	<u>Wt (%)</u>
Alkyl, 1.4 ethoxylated, sulfate	30
Amine oxide	6
Polyhydroxy fatty acid amide	4
Nonionic surfactant (C11E9)	5
Mg ion from MgCl ₂	1
Ca ion from CaCl ₂	0.2
Peptide aldehyde*	0.0025
Sodium xylene sulfonate	4
Solvent	6
Water	to 100%
pH	to 8

*Peptide Aldehyde of Synthesis Example 2.

5

EXAMPLE III

The following compositions are made by combining the listed ingredients in the listed proportions.

<u>Ingredients</u>	<u>A (wt %)</u>	<u>B (wt %)</u>	<u>C (wt %)</u>	<u>D (wt %)</u>
LAS	0	0	0	12
AExS ¹	22.1	24.7	33.5	3
Polyhydroxy fatty acid amide	4.6	1.2	4.2	0
Amine Oxide	4.6	1.2	4.8	0
Betaine	0	1.2	0	0
Nonionic Surfactant	6.7	4.1	0	0
Mg(OH) ₂	0.5	0.5	0.7	0
Ca ion from CaCl ₂	0.1	0.3	0.4	0.1
Calcium xylene sulfonate	4.5	0	4	0
Polyethylene glycol	3	0	0	0
Polypropylene glycol 2000	1.5	0	0	0


Balance, water	to 100%	to 100%	to 100%	to 100%
Protease A or Protease B	0.001-0.01	0.001-0.01	0.005-0.01	0.0003-0.01
Peptide Aldehyde ²	0.00025- 0.0025	0.00025- 0.0025	0.00025- 0.0025	0.00125- 0.0025

¹ x= the degree of ethoxylation. The average degree of ethoxylation for the compositions are: A=2.2, B=0.6, C=1.4, D=2.2.

² The peptide aldehyde of Synthesis Example 2 is used herein.

WHAT IS CLAIMED IS:

1. A liquid detergent composition comprising:
 - a) from 1% to 95%, by weight of composition, of a deterative surfactant;
 - b) an active proteolytic enzyme; and
 - c) a peptide protease inhibitor having the formula:



wherein B is a peptide chain comprising from 2 to 5 amino acid moieties; X is hydrogen or CF₃; Z is an N-capping moiety selected from the group consisting of sulfonamides, phosphoramidates, thioureas, sulfenamides, sulfonic acids, phosphinamides, thiocarbamates, amidophosphates, sulfamoyl derivatives, and phosphoramidates; and R is selected from the group consisting of straight or branched C₁ - C₆ unsubstituted alkyl, phenyl, and C₇ - C₉ alkylaryl moieties.

2. A liquid detergent composition according to Claim 1 wherein the R is selected from the group consisting of methyl, iso-propyl, sec-butyl, iso-butyl, -C₆H₅, -CH₂-C₆H₅, and -CH₂CH₂-C₆H₅.
3. A liquid detergent composition according to either of Claims 1 or 2 comprising:
 - a) from 8 to 70% of said deterative surfactant;
 - b) from 0.0001% to 5% of an active proteolytic enzyme;
 - c) from 0.00001% to 5% of a peptide protease inhibitor.
4. A liquid detergent composition according to any of Claims 1-3 further comprising a source of calcium ions.
5. A liquid detergent composition according to any of Claims 1-4 wherein the N-capping Z moiety is selected from the group consisting of (R''O)₂(O)P-, (SR'')₂-, R''(O)₂S-, SO₃H, (R'')₂(O)P-, R''O(O)₂S-, (R'')₂N(O)C-, R''O(S)C-, R''-P(O)OH, and R''O(OH)(O)P-, wherein each R'' is independently selected from the group consisting of straight or branched C₁-C₆ unsubstituted alkyl, phenyl, C₇-C₉ alkylaryl, and cycloalkyl moieties, wherein the cycloalkyl ring may span C₄-C₈ and may contain one or more heteroatoms selected from the group consisting of O, N, and S.

6. liquid detergent composition according to any of Claims 1-5 wherein said proteolytic enzyme is a subtilisin-type protease.
7. A liquid detergent composition according to any of Claims 1-6 wherein said composition is a light duty detergent composition suitable for dishcare comprising amylase enzyme.
8. A liquid detergent composition according to any of Claims 1-6 wherein said composition is a heavy duty detergent composition suitable for laundrycare further comprising an effective amount one or more of the following second enzymes: lipase, amylase, cellulase, and mixtures thereof.
9. A liquid laundry detergent composition according to any of Claims 1-8 further comprising from from 0.25% to 10% by weight of boric acid or a compound capable of forming boric acid and a polyol.
10. The peptide compounds $\text{CH}_3\text{SO}_2\text{Phe-Gly-Ala-Leu-H}$, $\text{CH}_3\text{SO}_2\text{Val-Ala-Leu-H}$, $\text{C}_6\text{H}_5\text{CH}_2\text{O(OH)(O)P-Val-Ala-Leu-H}$, $\text{C}_6\text{H}_5\text{CH}_2\text{O(OH)(O)P-Val-Ala-Leu-CF}_3$, $\text{CH}_3\text{CH}_2\text{SO}_2\text{-Phe-Gly-Ala-Leu-H}$, $\text{C}_6\text{H}_5\text{CH}_2\text{SO}_2\text{-Val-Ala-Leu-H}$, $\text{C}_6\text{H}_5\text{CH}_2\text{O(OH)(O)P-Leu-Ala-Leu-H}$, $\text{C}_6\text{H}_5\text{CH}_2\text{O(OH)(O)P-Phe-Ala-Leu-H}$, and $\text{CH}_3\text{O(OH)(O)P-Leu-Gly-Ala-Leu-H}$.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/16623

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C11D3/386 C07K5/08 C07K5/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C11D C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 583 535 A (PROCTER & GAMBLE) 23 February 1994 cited in the application see the whole document ---	1-10
A	EP 0 583 534 A (PROCTER & GAMBLE) 23 February 1994 cited in the application see the whole document ---	1-10
A	WO 93 00418 A (GENENCOR INT) 7 January 1993 see claims ---	1-9
A	WO 95 24914 A (MYOGENICS INC) 21 September 1995 compound no 124, see page 47 ---	10
	-/--	

Further documents are listed in the continuation of box C. Patent family members are listed in annex.

° Special categories of cited documents :

<p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p>	<p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&* document member of the same patent family</p>
--	--

Date of the actual completion of the international search 21 January 1998	Date of mailing of the international search report 03.02.98
--	--

Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer Pelli Wablat, B
--	---

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/16623

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 03529 A (NOVONORDISK AS) 5 March 1992 see claims	1-9
A	--- US 4 537 706 A (SEVERSON JR ROLAND G) 27 August 1985 cited in the application see column 6, line 65 see column 7, line 60 - line 68 see column 8, paragraph 2 see claims -----	1,3,4, 7-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/16623

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0583535 A	23-02-94	AT 145666 T	15-12-96
		AU 4789593 A	15-03-94
		CN 1085941 A	27-04-94
		DE 69215514 D	09-01-97
		DE 69215514 T	12-06-97
		JP 8500143 T	09-01-96
		WO 9404652 A	03-03-94
		US 5582762 A	10-12-96

EP 0583534 A	23-02-94	AT 149562 T	15-03-97
		AU 4789493 A	15-03-94
		CA 2142297 A	03-03-94
		CN 1085943 A	27-04-94
		DE 69217934 D	10-04-97
		DE 69217934 T	04-09-97
		ES 2098483 T	01-05-97
		JP 9503793 T	15-04-97
		WO 9404651 A	03-03-94
		US 5576283 A	19-11-96

WO 9300418 A	07-01-93	US 5178789 A	12-01-93
		CA 2111970 A	07-01-93
		EP 0591445 A	13-04-94
		JP 7501350 T	09-02-95

WO 9524914 A	21-09-95	AU 682600 B	09-10-97
		AU 2228395 A	03-10-95
		CA 2185326 A	21-09-95
		EP 0804216 A	05-11-97
		FI 963602 A	04-11-96
		JP 9511501 T	18-11-97
		NO 963814 A	14-11-96
		US 5693617 A	02-12-97

WO 9203529 A	05-03-92	DE 69101557 D	05-05-94
		DE 69101557 T	14-07-94
		EP 0544777 A	09-06-93
		ES 2062812 T	16-12-94
		JP 6500142 T	06-01-94

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. onal Application No

PCT/US 97/16623

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4537706 A	27-08-85	CA 1244362 A	08-11-88
		EP 0162033 A	21-11-85
		IE 58048 B	16-06-93
		JP 2107071 C	06-11-96
		JP 7116471 B	13-12-95
		JP 61057697 A	24-03-86
