



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82	A2	(11) International Publication Number: WO 99/64614 (43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/US99/13559 (22) International Filing Date: 10 June 1999 (10.06.99) (30) Priority Data: 60/089,043 12 June 1998 (12.06.98) US (71) Applicant (for all designated States except US): CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): KNUTZON, Debbie [US/US]; 6110 Rockhurst Way, Granite Bay, CA 95746 (US). (74) Agents: SCHWEDLER, Carl, J. et al.; Patent Department Central, Monsanto/G.D. Searle, P.O. Box 5110, Chicago, IL 60680-5110 (US).		(81) Designated States: CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: POLYUNSATURATED FATTY ACIDS IN PLANTS (57) Abstract <p>The present invention relates to compositions and methods for preparing polyunsaturated long chain fatty acids in plants, plant parts and plant cells, such as leaves, roots, fruits and seeds. Nucleic acid sequences and constructs encoding fatty acid desaturases, including $\Delta 5$-desaturases, $\Delta 6$-desaturases and $\Delta 12$-desaturases, are used to generate transgenic plants, plant parts and cells which contain and express one or more transgenes encoding one or more desaturases. Expression of the desaturases with different substrate specificities in the plant system permit the large scale production of polyunsaturated long chain fatty acids such as docosahexaenoic acid, eicosapentaenoic acid, α-linolenic acid, gamma-linolenic acid, arachidonic acid and the like for modification of the fatty acid profile of plants, plant parts and tissues. Manipulation of the fatty acid profiles allows for the production of commercial quantities of novel plant oils and products.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

POLYUNSATURATED FATTY ACIDS IN PLANTS

This application claims the benefit of U.S. Provisional Application Number 60/089,043 filed June 12, 1998.

5 Technical Field

This invention relates to modulating levels of enzymes and/or enzyme components capable of altering the production of long chain polyunsaturated fatty acids (PUFAS) in a host plant. The invention is exemplified by the production of PUFAS in plants.

10

INTRODUCTION

Background

Three main families of polyunsaturated fatty acids (PUFAs) are the 3 fatty acids, exemplified by arachidonic acid, the ω 9 fatty acids exemplified by Mead acid, and the ω 3 fatty acids, exemplified by eicosapentaenoic acid.

15 PUFAs are important components of the plasma membrane of the cell, where they may be found in such forms as phospholipids. PUFAs also serve as precursors to other molecules of importance in human beings and animals, including the prostacyclins, leukotrienes and prostaglandins. PUFAs are necessary for proper development, particularly in the developing infant brain,
20 and for tissue formation and repair.

Four major long chain PUFAs of importance include docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA), which are primarily found in different types of fish oil, gamma-linolenic acid (GLA), which is found in the seeds of a number of plants, including evening primrose (*Oenothera biennis*),
25 borage (*Borago officinalis*) and black currants (*Ribes nigrum*), and stearidonic acid (SDA), which is found in marine oils and plant seeds. Both GLA and another important long chain PUFA, arachidonic acid (ARA), are found in filamentous fungi. ARA can be purified from animal tissues

including liver and adrenal gland. Mead acid accumulates in essential fatty acid deficient animals.

For DHA, a number of sources exist for commercial production including a variety of marine organisms, oils obtained from cold water marine fish, and egg yolk fractions. For ARA, microorganisms including the genera *Mortierella*, *Entomophthora*, *Phytium* and *Porphyridium* can be used for commercial production. Commercial sources of SDA include the genera *Trichodesma* and *Echium*. Commercial sources of GLA include evening primrose, black currants and borage. However, there are several disadvantages associated with commercial production of PUFAs from natural sources. Natural sources of PUFAs, such as animals and plants, tend to have highly heterogeneous oil compositions. The oils obtained from these sources therefore can require extensive purification to separate out one or more desired PUFAs or to produce an oil which is enriched in one or more PUFA. Natural sources also are subject to uncontrollable fluctuations in availability. Fish stocks may undergo natural variation or may be depleted by overfishing. Fish oils have unpleasant tastes and odors, which may be impossible to economically separate from the desired product, and can render such products unacceptable as food supplements. Animal oils, and particularly fish oils, can accumulate environmental pollutants. Weather and disease can cause fluctuation in yields from both fish and plant sources. Cropland available for production of alternate oil-producing crops is subject to competition from the steady expansion of human populations and the associated increased need for food production on the remaining arable land. Crops which do produce PUFAs, such as borage, have not been adapted to commercial growth and may not perform well in monoculture. Growth of such crops is thus not economically competitive where more profitable and better established crops can be grown. Large scale fermentation of organisms such as *Mortierella* is also expensive. Natural animal tissues contain low amounts of ARA and are difficult to process. Microorganisms such as *Porphyridium* and *Mortierella* are difficult to cultivate on a commercial scale.

Dietary supplements and pharmaceutical formulations containing PUFAs can retain the disadvantages of the PUFA source. Supplements such as fish oil capsules can contain low levels of the particular desired component and thus require large dosages. High dosages result in ingestion of high levels of undesired components, including contaminants. Care must be taken in providing fatty acid supplements, as overaddition may result in suppression of endogenous biosynthetic pathways and lead to competition with other necessary fatty acids in various lipid fractions *in vivo*, leading to undesirable results. For example, Eskimos having a diet high in ω 3 fatty acids have an increased tendency to bleed (U.S. Pat. No. 4,874,603). Unpleasant tastes and odors of the supplements can make such regimens undesirable, and may inhibit compliance by the patient.

A number of enzymes are involved in PUFA biosynthesis. Linoleic acid (LA, 18:2 Δ 9, 12) is produced from oleic acid (18:1 Δ 9) by a Δ 12-desaturase. GLA (18:3 Δ 6, 9, 12) is produced from linoleic acid (LA, 18:2 Δ 9, 12) by a Δ 6-desaturase. ARA (20:4 Δ 5, 8, 11, 14) production from DGLA (20:3 Δ 8, 11, 14) is catalyzed by a Δ 5-desaturase. However, animals cannot desaturate beyond the Δ 9 position and therefore cannot convert oleic acid (18:1 Δ 9) into linoleic acid (18:2 Δ 9, 12). Likewise, α -linolenic acid (ALA, 18:3 Δ 9, 12, 15) cannot be synthesized by mammals. Other eukaryotes, including fungi and plants, have enzymes which desaturate at positions Δ 12 and Δ 15. The major poly-unsaturated fatty acids of animals therefore are either derived from diet and/or from desaturation and elongation of linoleic acid (18:2 Δ 9, 12) or α -linolenic acid (18:3 Δ 9, 12, 15).

Poly-unsaturated fatty acids are considered to be useful for nutritional, pharmaceutical, industrial, and other purposes. An expansive supply of poly-unsaturated fatty acids from natural sources and from chemical synthesis are not sufficient for commercial needs. Therefore it is of interest to obtain genetic material involved in PUFA biosynthesis from species that naturally produce these fatty acids and to express the isolated material alone or in

combination in a heterologous system which can be manipulated to allow production of commercial quantities of PUFAS.

SUMMARY OF THE INVENTION

5 Novel compositions and methods are provided for preparation of poly-unsaturated long chain fatty acids and desaturases in plants and plant cells. The methods involve growing a host plant cell of interest transformed with an expression cassette functional in a host plant cell, the expression cassette comprising a transcriptional and translational initiation regulatory
10 region, joined in reading frame 5' to a DNA sequence encoding a desaturase polypeptide capable of modulating the production of PUFAs. Expression of the desaturase polypeptide provides for an alteration in the PUFA profile of host plant cells as a result of altered concentrations of enzymes involved in PUFA biosynthesis. Of particular interest is the selective control of PUFA
15 production in plant tissues and/or plant parts such as leaves, roots, fruits and seeds. The invention finds use for example in the large scale production of DHA, Mead Acid, EPA, ARA, DGLA, stearidonic acid GLA and other fatty acids and for modification of the fatty acid profile of edible plant tissues and/or plant parts.

20 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows possible pathways for the synthesis of Mead acid (20:3 Δ 5, 8, 11), arachidonic acid (20:4 Δ 5, 8, 11, 14) and stearidonic acid (18:4 Δ 6, 9, 12, 15) from palmitic acid (C_{16}) from a variety of organisms,
25 including algae, *Mortierella* and humans. These PUFAs can serve as precursors to other molecules important for humans and other animals, including prostacyclins, leukotrienes, and prostaglandins, some of which are shown.

Figure 2 shows possible pathways for production of PUFAs in addition to ARA, including taxoleic acid and pinolenic, again compiled from a variety of organisms.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

5

In order to ensure a complete understanding of the invention, the following definitions are provided:

Δ 5-Desaturase: Δ 5 desaturase is an enzyme which introduces a double bond between carbons 5 and 6 from the carboxyl end of a fatty acid molecule.

10

Δ 6-Desaturase: Δ 6-desaturase is an enzyme which introduces a double bond between carbons 6 and 7 from the carboxyl end of a fatty acid molecule.

15

Δ 9-Desaturase: Δ 9-desaturase is an enzyme which introduces a double bond between carbons 9 and 10 from the carboxyl end of a fatty acid molecule.

Δ 12-Desaturase: Δ 12-desaturase is an enzyme which introduces a double bond between carbons 12 and 13 from the carboxyl end of a fatty acid molecule.

20

Fatty acids: Fatty acids are a class of compounds containing a long-hydrocarbon chain and a terminal carboxylate group. Fatty acids include the following:

Fatty Acid		
12:0	lauric acid	
16:0	Palmitic acid	
16:1	Palmitoleic acid	
18:0	stearic acid	
18:1	oleic acid	Δ^9 -18:1
18:2 $\Delta^5,9$	Taxoleic acid	$\Delta^5,9$ -18:2
18:2 $\Delta^6,9$	6,9-octadecadienoic acid	$\Delta^6,9$ -18:2
18:2	Linoleic acid	$\Delta^9,12$ -18:2 (LA)
18:3 $\Delta^6,9,12$	Gamma-linolenic acid	$\Delta^6,9,12$ -18:3 (GLA)
18:3 $\Delta^5,9,12$	Pinolenic acid	$\Delta^5,9,12$ -18:3
18:3	alpha-linolenic acid	$\Delta^9,12,15$ -18:3 (ALA)
18:4	Stearidonic acid	$\Delta^6,9,12,15$ -18:4 (SDA)
20:0	Arachidic acid	
20:1	Eicosenic Acid	
20:2 $\Delta^8, 11$		$\Delta^8, 11$
20:3 $\Delta^5, 8, 11$	Mead Acid	$\Delta^5, 8, 11$
22:0	Behenoic acid	
22:1	erucic acid	
22:2	Docasadienoic acid	
20:4 γ^6	Arachidonic acid	$\Delta^5,8,11,14$ -20:4 (ARA)
20:3 γ^6	γ^6 -eicosatrienoic dihydro-gamma linolenic	$\Delta^8,11,14$ -20:3 (DGLA)
20:5 γ^3	Eicosapentaenoic (Timnodonic acid)	$\Delta^5,8,11,14,17$ -20:5 (EPA)
20:3 γ^3	γ^3 -eicosatrienoic	$\Delta^11,16,17$ -20:3
20:4 γ^3	γ^3 -eicosatetraenoic	$\Delta^8,11,14,17$ -20:4
22:5 γ^3	Docosapentaenoic	$\Delta^7,10,13,16,19$ -22:5 (γ^3 DPA)
22:6 γ^3	Docosahexaenoic (cervonic acid)	$\Delta^4,7,10,13,16,19$ -22:6 (DHA)
24:0	Lignoceric acid	

Taking into account these definitions, the present invention is directed to novel DNA sequences, and DNA constructs related to the production of fatty acids in plants. Methods and compositions are provided which permit modification of the poly-unsaturated long chain fatty acid content of plant cells. Plant cells are transformed with an expression cassette comprising a DNA encoding a polypeptide capable of increasing the amount of one or more PUFA in a plant cell. Desirably, integration constructs may be prepared which provide for integration of the expression cassette into the genome of a host cell. Host cells are manipulated to express a sense or antisense DNA encoding a polypeptide(s) that has desaturase activity. By "desaturase" is intended a polypeptide which can desaturate one or more fatty acids to produce a mono- or poly-unsaturated fatty acid or precursor thereof of interest. By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification, for example, glycosylation or phosphorylation. The substrate(s) for the expressed enzyme may be produced by the host cell or may be exogenously supplied.

To achieve expression in a host cell, the transformed DNA is operably associated with transcriptional and translational initiation and termination regulatory regions that are functional in the host cell. Constructs comprising the gene to be expressed can provide for integration into the genome of the host cell or can autonomously replicate in the host cell. For production of taxoleic acid, the expression cassettes generally used include a cassette which provides for $\Delta 5$ desaturase activity, particularly in a host cell which produces or can take up oleic acid. For production of $\Delta 6,9$ 18:2 or other $\omega 9$ unsaturated fatty acids, the expression cassettes generally used include a cassette which provides for $\Delta 6$ desaturase activity, particularly in a host cell which produces or can take up oleic acid. Production of oleic acid, taxoleic acid, or $\omega 9$ unsaturated fatty acids in a plant having $\Delta 12$ desaturase activity is favored by providing an expression cassette for an antisense $\Delta 12$ transcript, or by disrupting a $\Delta 12$ desaturase gene. For production of linoleic acid (LA), the expression cassettes generally used include a cassette which provides for $\Delta 12$

desaturase activity, particularly in a host cell which produces or can take up oleic acid. For production of ALA, the expression cassettes generally used include a cassette which provides for $\Delta 15$ or $\omega 3$ desaturase activity, particularly in a host cell which produces or can take up LA. For production of GLA or SDA, the expression cassettes generally used include a cassette which provides for $\Delta 6$ desaturase activity, particularly in a host cell which produces or can take up LA or ALA, respectively. Production of $\omega 6$ -type unsaturated fatty acids, such as LA or GLA, in a plant capable of producing ALA is favored by inhibiting the activity of $\Delta 15$ or $\omega 3$ type desaturase; this is accomplished by providing and expression cassette for an antisense $\Delta 15$ or $\omega 3$ transcript, or by disrupting a $\Delta 15$ or $\omega 3$ desaturase gene. Similarly, production of LA or ALA in a plant having $\Delta 6$ desaturase activity is favored by providing an expression cassette for an antisense $\Delta 6$ transcript, or by disrupting a $\Delta 6$ desaturase gene. For production of ARA in a host cell which produces or can take up DGLA, the expression cassette generally used provides for D5 desaturase activity. Production of $\omega 6$ -type unsaturated fatty acids, such as ARA, in a plant capable of producing ALA is favored by inhibiting the activity of a $\Delta 15$ or $\omega 3$ type desaturase; this is accomplished by providing an expression cassette for an antisense $\Delta 15$ or $\omega 3$ transcript, or by disrupting a $\Delta 15$ or $\omega 3$ desaturase gene.

TRANSGENIC PLANT PRODUCTION OF FATTY ACIDS

Transgenic plant production of PUFAs offers several advantages over purification from natural sources such as fish or plants. Production of fatty acids from recombinant plants provides the ability to alter the naturally occurring plant fatty acid profile by providing new synthetic pathways in the host or by suppressing undesired pathways, thereby increasing levels of desired PUFAs, or conjugated forms thereof, and decreasing levels of undesired PUFAs. Production of fatty acids in transgenic plants also offers the advantage that expression of desaturase genes in particular tissues and/or plant parts means that greatly increased levels of desired PUFAs in those

tissues and/or parts can be achieved, making recovery from those tissues more economical. For example, the desired PUFAs can be expressed in seed; methods of isolating seed oils are well established. In addition to providing a source for purification of desired PUFAs, seed oil components can be
5 manipulated through expression of desaturase genes, either alone or in combination with other genes such as elongases, to provide seed oils having a particular PUFA profile in concentrated form. The concentrated seed oils then can be added to animal milks and/or synthetic or semi-synthetic milks to serve as infant formulas where human nursing is impossible or undesired, or
10 in cases of malnourishment or disease in both adults and infants.

For production of PUFAs, depending upon the host cell, the availability of substrate, and the desired end product(s), several polypeptides, particularly desaturases, are of interest including those polypeptides which catalyze the conversion of stearic acid to oleic acid, LA to GLA, of ALA to
15 SDA, of oleic acid to LA, or of LA to ALA, oleic acid to taxolic acid, LA to pinolenic acid, oleic acid to 6,9-actadeca-dienoic acid which includes enzymes which desaturate at the $\Delta 6$, $\Delta 9$, $\Delta 5$, $\Delta 12$, $\Delta 15$, $\Delta 5$, or $\omega 3$ positions. Considerations for choosing a specific polypeptide having desaturase activity include the pH optimum of the polypeptide, whether the polypeptide is a rate
20 limiting enzyme or a component thereof, whether the desaturase used is essential for synthesis of a desired poly-unsaturated fatty acid, and/or co-factors required by the polypeptide. The expressed polypeptide preferably has parameters compatible with the biochemical environment of its location in the host cell. For example, the polypeptide may have to compete for
25 substrate with other enzymes in the host cell. Analyses of the K_m and specific activity of the polypeptide in question therefore are considered in determining the suitability of a given polypeptide for modifying PUFA production in a given host cell. The polypeptide used in a particular situation therefore is one which can function under the conditions present in the intended host cell but
30 otherwise can be any polypeptide having desaturase activity which has the desired characteristic of being capable of modifying the relative production of

a desired PUFA. A scheme for the synthesis of arachidonic acid (20:4 Δ 5, 8, 11, 14) from palmitic acid (C_{16}) is shown in Figure 1. A key enzyme in this pathway is a Δ 5-desaturase which converts DH- γ -linolenic acid (DGLA, eicosatrienoic acid) to ARA. Conversion of α -linolenic acid (ALA) to stearidonic acid by a Δ 6-desaturase is also shown. Production of PUFAs in addition to ARA, including EPA and DHA is shown in Figure 2. A key enzyme in the synthesis of arachidonic acid (20:4 Δ 5, 8, 11, 14) from stearic acid (C_{18}) is a Δ 6-desaturase which converts the linoleic acid into γ -linolenic acid. Conversion of α -linolenic acid (ALA) to stearidonic acid by a Δ 6-desaturase also is shown. For production of ARA, the DNA sequence used encodes a polypeptide having Δ 5 desaturase activity. In particular instances, this can be coupled with an expression cassette which provides for production of a polypeptide having Δ 6 desaturase activity and, optionally, a transcription cassette providing for production of antisense sequences to a Δ 15 transcription product. The choice of combination of cassettes used depends in part on the PUFA profile of the host cell. Where the host cell Δ 5-desaturase activity is limiting, overexpression of Δ 5 desaturase alone generally will be sufficient to provide for enhanced ARA production.

SOURCES OF POLYPEPTIDES HAVING DESATURASE ACTIVITY

As sources of polypeptides having desaturase activity and oligonucleotides encoding such polypeptides are organisms which produce a desired poly-unsaturated fatty acid. As an example, microorganisms having an ability to produce ARA can be used as a source of Δ 5-desaturase genes; microorganisms which GLA or SDA can be used as a source of Δ 6-desaturase and/or Δ 12-desaturase genes. Such microorganisms include, for example, those belonging to the genera *Mortierella*, *Conidiobolus*, *Pythium*, *Phytophthora*, *Penicillium*, *Porphyridium*, *Coidosporium*, *Mucor*, *Fusarium*, *Aspergillus*, *Rhodotorula*, and *Entomophthora*. Within the genus *Porphyridium*, of particular interest is *Porphyridium cruentum*. Within the

genus *Mortierella*, of particular interest are *Mortierella elongata*, *Mortierella exigua*, *Mortierella hygrophila*, *Mortierella ramanniana*. var. *angulispora*, and *Mortierella alpina*. Within the genus *Mucor*, of particular interest are *Mucor circinelloides* and *Mucor javanicus*.

5 DNAs encoding desired desaturases can be identified in a variety of ways. As an example, a source of the desired desaturase, for example genomic or cDNA libraries from *Mortierella*, is screened with detectable enzymatically- or chemically-synthesized probes, which can be made from DNA, RNA, or non-naturally occurring nucleotides, or mixtures thereof.

10 Probes may be enzymatically synthesized from DNAs of known desaturases for normal or reduced-stringency hybridization methods. Oligonucleotide probes also can be used to screen sources and can be based on sequences of known desaturases, including sequences conserved among known desaturases, or on peptide sequences obtained from the desired purified

15 protein. Oligonucleotide probes based on amino acid sequences can be degenerate to encompass the degeneracy of the genetic code, or can be biased in favor of the preferred codons of the source organism. Oligonucleotides also can be used as primers for PCR from reverse transcribed mRNA from a known or suspected source; the PCR product can be the full length cDNA or

20 can be used to generate a probe to obtain the desired full length cDNA. Alternatively, a desired protein can be entirely sequenced and total synthesis of a DNA encoding that polypeptide performed.

Once the desired genomic or cDNA has been isolated, it can be sequenced by known methods. It is recognized in the art that such methods

25 are subject to errors, such that multiple sequencing of the same region is routine and is still expected to lead to measurable rates of mistakes in the resulting deduced sequence, particularly in regions having repeated domains, extensive secondary structure, or unusual base compositions, such as regions with high GC base content. When discrepancies arise, resequencing can be

30 done and can employ special methods. Special methods can include altering sequencing conditions by using: different temperatures; different enzymes;

proteins which alter the ability of oligonucleotides to form higher order structures; altered nucleotides such as ITP or methylated dGTP; different gel compositions, for example adding formamide; different primers or primers located at different distances from the problem region; or different templates
5 such as single stranded DNAs. Sequencing of mRNA can also be employed.

For the most part, some or all of the coding sequence for the polypeptide having desaturase activity is from a natural source. In some situations, however, it is desirable to modify all or a portion of the codons, for example, to enhance expression, by employing host preferred codons.
10 Host preferred codons can be determined from the codons of highest frequency in the proteins expressed in the largest amount in a particular host species of interest. Thus, the coding sequence for a polypeptide having desaturase activity can be synthesized in whole or in part. All or portions of the DNA also can be synthesized to remove any destabilizing sequences or
15 regions of secondary structure which would be present in the transcribed mRNA. All or portions of the DNA also can be synthesized to alter the base composition to one more preferable in the desired host cell. Methods for synthesizing sequences and bringing sequences together are well established in the literature. *In vitro* mutagenesis and selection, site-directed
20 mutagenesis, or other means can be employed to obtain mutations of naturally occurring desaturase genes to produce a polypeptide having desaturase activity *in vivo* with more desirable physical and kinetic parameters for function in the host cell, such as a longer half-life or a higher rate of production of a desired polyunsaturated fatty acid.

25 Desirable cDNAs have less than 60% A+T composition, preferably less than 50% A+T composition. On a localized scale of a sliding window of 20 base pairs, it is preferable that there are no localized regions of the cDNA with greater than 75% A+T composition; with a window of 60 base pairs, it is preferable that there are no localized regions of the cDNA with greater than
30 60%, more preferably no localized regions with greater than 55% A+T composition.

Mortierella alpina Desaturases

Of particular interest are the *Mortierella alpina* $\Delta 5$ -desaturase, $\Delta 6$ -desaturase, $\Delta 12$ -desaturase and $\Delta 15$ desaturase. The gene encoding the *Mortierella alpina* $\Delta 5$ -desaturase can be expressed in transgenic plants to effect greater synthesis of ARA from DGLA, or pinolenic acid from LA, taxoleic acid from oleic acid or Mead and from $\Delta 8, 11-20:2$. Other DNAs which are substantially identical in sequence to the *Mortierella alpina* $\Delta 5$ -desaturase DNA, or which encode polypeptides which are substantially identical in sequence to the *Mortierella alpina* $\Delta 5$ -desaturase polypeptide, also can be used. The gene encoding the *Mortierella alpina* $\Delta 6$ -desaturase can be expressed in transgenic plants or animals to effect greater synthesis of GLA from linoleic acid or of stearidonic acid (SDA) from ALA or of 6,9-octadecadienoic acid from oleic acid. Other DNAs which are substantially identical in sequence to the *Mortierella alpina* $\Delta 6$ -desaturase DNA, or which encode polypeptides which are substantially identical in sequence to the *Mortierella alpina* $\Delta 6$ -desaturase polypeptide, also can be used.

The gene encoding the *Mortierella alpina* $\Delta 12$ -desaturase can be expressed in transgenic plants to effect greater synthesis of LA from oleic acid. Other DNAs which are substantially identical to the *Mortierella alpina* $\Delta 12$ -desaturase DNA, or which encode polypeptides which are substantially identical to the *Mortierella alpina* $\Delta 12$ -desaturase polypeptide, also can be used.

By substantially identical in sequence is intended an amino acid sequence or nucleic acid sequence exhibiting in order of increasing preference at least 60%, 80%, 90% or 95% homology to the *Mortierella alpina* $\Delta 5$ -desaturase amino acid sequence or nucleic acid sequence encoding the amino acid sequence. For polypeptides, the length of comparison sequences generally is at least 16 amino acids, preferably at least 20 amino acids, or most preferably 35 amino acids. For nucleic acids, the length of comparison sequences generally is at least 50 nucleotides, preferably at least

60 nucleotides, and more preferably at least 75 nucleotides, and most preferably, 110 nucleotides. Homology typically is measured using sequence analysis software, for example, the Sequence Analysis software package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wisconsin 53705, MEGAlign (DNASStar, Inc., 1228 S. Park St., Madison, Wisconsin 53715), and MacVector (Oxford Molecular Group, 2105 S. Bascom Avenue, Suite 200, Campbell, California 95008). Such software matches similar sequences by assigning degrees of homology to various substitutions, deletions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine and leucine; aspartic acid, glutamic acid, asparagine, and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine. Substitutions may also be made on the basis of conserved hydrophobicity or hydrophilicity (Kyte and Doolittle, *J. Mol. Biol.* 157: 105-132, 1982), or on the basis of the ability to assume similar polypeptide secondary structure (Chou and Fasman, *Adv. Enzymol.* 47: 45-148, 1978).

EXPRESSION OF DESATURASE GENES

Once the DNA encoding a desaturase polypeptide has been obtained, it is placed in a vector capable of replication in a host cell, or is propagated *in vitro* by means of techniques such as PCR or long PCR. Replicating vectors can include plasmids, phage, viruses, cosmids and the like. Desirable vectors include those useful for mutagenesis of the gene of interest or for expression of the gene of interest in host cells. The technique of long PCR has made *in vitro* propagation of large constructs possible, so that modifications to the gene of interest, such as mutagenesis or addition of expression signals, and propagation of the resulting constructs can occur entirely *in vitro* without the use of a replicating vector or a host cell.

For expression of a desaturase polypeptide, functional transcriptional and translational initiation and termination regions are operably linked to the DNA encoding the desaturase polypeptide. Transcriptional and translational initiation and termination regions are derived from a variety of nonexclusive sources, including the DNA to be expressed, genes known or suspected to be capable of expression in the desired system, expression vectors, chemical synthesis, or from an endogenous locus in a host cell. Expression in a plant tissue and/or plant part presents certain efficiencies, particularly where the tissue or part is one which is easily harvested, such as seed, leaves, fruits, flowers, roots, etc. Expression can be targeted to that location within the plant by using specific regulatory sequences, such as those of USPN 5,463,174, USPN 4,943,674, USPN 5,106,739, USPN 5,175,095, USPN 5,420,034, USPN 5,188,958, and USPN 5,589,379. Alternatively, the expressed protein can be an enzyme which produces a product which may be incorporated, either directly or upon further modifications, into a fluid fraction from the host plant. In the present case, expression of desaturase genes, or antisense desaturase transcripts, can alter the levels of specific PUFAs, or derivatives thereof, found in plant parts and/or plant tissues. The $\Delta 5$ -desaturase polypeptide coding region is expressed either by itself or with other genes, in order to produce tissues and/or plant parts containing higher proportions of desired PUFAs or in which the PUFA composition more closely resembles that of human breast milk (Prieto *et al.*, PCT publication WO 95/24494). The termination region can be derived from the 3' region of the gene from which the initiation region was obtained or from a different gene. A large number of termination regions are known to and have been found to be satisfactory in a variety of hosts from the same and different genera and species. The termination region usually is selected more as a matter of convenience rather than because of any particular property.

The choice of a host cell is influenced in part by the desired PUFA profile of the transgenic cell, and the native profile of the host cell. As an example, for production of linoleic acid from oleic acid, the DNA sequence

used encodes a polypeptide having $\Delta 12$ desaturase activity, and for production of GLA from linoleic acid, the DNA sequence used encodes a polypeptide having $\Delta 6$ desaturase activity. Use of a host cell which expresses $\Delta 12$ desaturase activity and lacks or is depleted in $\Delta 15$ desaturase activity, 5 can be used with an expression cassette which provides for overexpression of $\Delta 6$ desaturase alone generally is sufficient to provide for enhanced GLA production in the transgenic cell. Where the host cell expresses $\Delta 9$ desaturase activity, expression of both a $\Delta 12$ - and a $\Delta 6$ -desaturase can provide for enhanced GLA production. In particular instances where 10 expression of $\Delta 6$ desaturase activity is coupled with expression of $\Delta 12$ desaturase activity, it is desirable that the host cell naturally have, or be mutated to have, low $\Delta 15$ desaturase activity. Alternatively, a host cell for $\Delta 6$ desaturase expression may have, or be mutated to have, high $\Delta 12$ desaturase activity.

15 Expression in a host cell can be accomplished in a transient or stable fashion. Transient expression can occur from introduced constructs which contain expression signals functional in the host cell, but which constructs do not replicate and rarely integrate in the host cell, or where the host cell is not proliferating. Transient expression also can be accomplished by inducing the 20 activity of a regulatable promoter operably linked to the gene of interest, although such inducible systems frequently exhibit a low basal level of expression. Stable expression can be achieved by introduction of a construct that can integrate into the host genome or that autonomously replicates in the host cell. Stable expression of the gene of interest can be selected for through 25 the use of a selectable marker located on or transfected with the expression construct, followed by selection for cells expressing the marker. When stable expression results from integration, integration of constructs can occur randomly within the host genome or can be targeted through the use of constructs containing regions of homology with the host genome sufficient to 30 target recombination with the host locus. Where constructs are targeted to an

endogenous locus, all or some of the transcriptional and translational regulatory regions can be provided by the endogenous locus.

When increased expression of the desaturase polypeptide in the source plant is desired, several methods can be employed. Additional genes
5 encoding the desaturase polypeptide can be introduced into the host organism. Expression from the native desaturase locus also can be increased through homologous recombination, for example by inserting a stronger promoter into the host genome to cause increased expression, by removing
destabilizing sequences from either the mRNA or the encoded protein by
10 deleting that information from the host genome, or by adding stabilizing sequences to the mRNA (*see* USPN 4,910,141 and USPN 5,500,365.)

When it is desirable to express more than one different gene, appropriate regulatory regions and expression methods. introduced genes can be propagated in the host cell through use of replicating vectors or by
15 integration into the host genome. Where two or more genes are expressed from separate replicating vectors, it is desirable that each vector has a different means of replication. Each introduced construct, whether integrated or not, should have a different means of selection and should lack homology to the other constructs to maintain stable expression and prevent reassortment
20 of elements among constructs. Judicious choices of regulatory regions, selection means and method of propagation of the introduced construct can be experimentally determined so that all introduced genes are expressed at the necessary levels to provide for synthesis of the desired products.

Constructs comprising the gene of interest may be introduced into a
25 host cell by standard techniques. These techniques include transfection, infection, bolistic impact, electroporation, microinjection, scraping, or any other method which introduces the gene of interest into the host cell (*see* USPN 4,743,548, USPN 4,795,855, USPN 5,068,193, USPN 5,188,958, USPN 5,463,174, USPN 5,565,346 and USPN 5,565,347). For convenience,
30 a host cell which has been manipulated by any method to take up a DNA sequence or construct will be referred to as "transformed" or "recombinant"

herein. The subject host will have at least have one copy of the expression construct and may have two or more, depending upon whether the gene is integrated into the genome, amplified, or is present on an extrachromosomal element having multiple copy numbers.

5 The transformed host cell can be identified by selection for a marker contained on the introduced construct. Alternatively, a separate marker construct may be introduced with the desired construct. as many transformation techniques introduce many DNA molecules into host cells. Typically, transformed hosts are selected for their ability to grow on selective
10 media. Selective media may incorporate an antibiotic or lack a factor necessary for growth of the untransformed host, such as a nutrient or growth factor. An introduced marker gene therefor may confer antibiotic resistance, or encode an essential growth factor or enzyme, and permit growth on selective media when expressed in the transformed host cell. Desirably,
15 resistance to kanamycin and the amino glycoside G418 are of interest (see USPN 5,034,322). Selection of a transformed host can also occur when the expressed marker protein can be detected, either directly or indirectly. The marker protein may be expressed alone or as a fusion to another protein. The marker protein can be detected by its enzymatic activity: for example β
20 galactosidase can convert the substrate X-gal to a colored product, and luciferase can convert luciferin to a light-emitting product. The marker protein can be detected by its light-producing or modifying characteristics; for example, the green fluorescent protein of *Aequorea victoria* fluoresces when illuminated with blue light. Antibodies can be used to detect the
25 marker protein or a molecular tag on, for example, a protein of interest. Cells expressing the marker protein or tag can be selected, for example, visually, or by techniques such as FACS or panning using antibodies.

 The PUFAs produced using the subject methods and compositions may be found in the host plant tissue and/or plant part as free fatty acids or in
30 conjugated forms such as acylglycerols, phospholipids, sulfolipids or glycolipids, and may be extracted from the host cell through a variety of

means well-known in the art. Such means may include extraction with organic solvents, sonication, supercritical fluid extraction using for example carbon dioxide, and physical means such as presses, or combinations thereof. Of particular interest is extraction with hexane or methanol and chloroform. Where desirable, the aqueous layer can be acidified to protonate negatively charged moieties and thereby increase partitioning of desired products into the organic layer. After extraction, the organic solvents can be removed by evaporation under a stream of nitrogen. When isolated in conjugated forms, the products are enzymatically or chemically cleaved to release the free fatty acid or a less complex conjugate of interest, and are then subjected to further manipulations to produce a desired end product. Desirably, conjugated forms of fatty acids are cleaved with potassium hydroxide.

Surprisingly, as demonstrated more fully in the examples below, expression of the *Mortierella* $\Delta 6$ desaturase leads to the production of stearidonic acid in the oil extracted from seed tissue of host plant cells. Furthermore, expression of the $\Delta 6$ desaturase with additional desaturases provided for the enhanced production of SDA in the seed oil.

Thus, the present invention provides methods for the production of stearidonic acid (C18:4) in host plant cells. The methods allow for the production of SDA in host plant cells ranging from about 0.3 wt% to at least about 30 wt%, preferably, from about 5 wt % to at least about 25 wt %, more preferably from about 7 wt % to at least about 25 wt %. The SDA is preferably produced in the seed oil of host plants containing one or more expression constructs as described herein.

Furthermore, the present invention provides a novel source of plant oils containing stearidonic acid. The oils are preferably obtained from the plant seed tissue. The seed oils contain amounts of SDA ranging from about 0.3 wt% to at least about 30 wt%, preferably, from about 5 wt % to at least about 25 wt %, more preferably from about 7 wt % to at least about 25 wt %.

PURIFICATION OF FATTY ACIDS

If further purification is necessary, standard methods can be employed. Such methods include extraction, treatment with urea, fractional crystallization, HPLC, fractional distillation, silica gel chromatography, high speed centrifugation or distillation, or combinations of these techniques. Protection of reactive groups, such as the acid or alkenyl groups, may be done at any step through known techniques, for example alkylation or iodination. Methods used include methylation of the fatty acids to produce methyl esters. Similarly, protecting groups may be removed at any step. Desirably, purification of fractions containing ARA, DHA and EPA is accomplished by treatment with urea and/or fractional distillation.

USES OF FATTY ACIDS

The uses of the fatty acids of subject invention are several. Probes based on the DNAs of the present invention may find use in methods for isolating related molecules or in methods to detect organisms expressing desaturases. When used as probes, the DNAs or oligonucleotides need to be detectable. This is usually accomplished by attaching a label either at an internal site, for example via incorporation of a modified residue, or at the 5' or 3' terminus. Such labels can be directly detectable, can bind to a secondary molecule that is detectably labeled, or can bind to an unlabelled secondary molecule and a detectably labeled tertiary molecule; this process can be extended as long as is practical to achieve a satisfactorily detectable signal without unacceptable levels of background signal. Secondary, tertiary, or bridging systems can include use of antibodies directed against any other molecule, including labels or other antibodies, or can involve any molecules which bind to each other, for example a biotin-streptavidin/avidin system. Detectable labels typically include radioactive isotopes, molecules which chemically or enzymatically produce or alter light, enzymes which produce detectable reaction products, magnetic molecules, fluorescent molecules or molecules whose fluorescence or light-emitting characteristics change upon binding. Examples of labelling methods can be found in USPN 5,011,770.

Alternatively, the binding of target molecules can be directly detected by measuring the change in heat of solution on binding of probe to target via isothermal titration calorimetry, or by coating the probe or target on a surface and detecting the change in scattering of light from the surface produced by binding of target or probe, respectively, as may be done with the BIAcore system.

The invention will be better understood by reference to the following non-limiting examples.

10

Examples

Example 1

Expression of ω -3 desaturase from *C. elegans* in transgenic plants.

The $\Delta 15/\omega$ -3 activity of *Brassica napus* can be increased by the expression of an ω -3 desaturase from *C. elegans*. The fat-1 cDNA clone (Genbank accession L41807; Spychalla, J. P., Kinney, A. J., and Browse, J. 1997 P.N.A.S. 94, 1142-1147) was obtained from John Browse at Washington State University. The fat-1 cDNA was modified by PCR to introduce cloning sites using the following primers:

15

Fat-1forward:

5'-CUACUACUACUACTGCAGACAATGGTCGCTCATTCCCTCAGA-3'

Fat-1reverse:

5'- CAUCAUCAUCAUGCGGCCGCTTACTTGGCCTTTGCCTT - 3'

25

These primers allowed the amplification of the entire coding region and added PstI and NotI sites to the 5'- and 3'-ends, respectively. The PCR product was subcloned into pAMP1 (GIBCOBRL) using the CloneAmp system (GIBCOBRL) to create pCGN5562. The sequence was verified by sequencing of both strands to be sure no changes were introduced by PCR.

A once base pair difference was observed in the fat-1 coding region from pCGN5562 vs. the fat-1 Genbank sequence. The C at position 705 of the fat-1 sequence was changed to an A in pCGN5562. This creates a change of a GAC codon to GAA, changing the Asp residue at position 231 of fat-1 to a Glu residue. This identical change was observed in products of two independent PCR reactions using fat-1 template and most likely is not a result of PCR mis-incorporation of a nucleotide. For seed specific expression, the Fat-1 coding region was cut out of pCGN5562 as a PstI/NotI fragment and inserted between the PstI/NotI sites of the binary vector, pCGN8623, to create pCGN5563. PCGN5563 can be introduced into *Brassica napus* via *Agrobacterium*-mediated transformation.

Construction of pCGN8623

The polylinker region of the napin promoter cassette, pCGN7770, was replaced by ligating the following oligonucleotides:

5'- TCGACCTGCAGGAAGCTTGC GGCCGCGGATCC -3' and
5'- TCGAGGATCCGCGGCCGCAAGCTTCCTGCAGG-3'. These oligonucleotides were ligated into SalI/XhoI-digested pCGN7770 to produce pCGN8619. These oligos encode BamHI, NotI, HindIII, and PstI restriction sites. pCGN8619 contains the oligos oriented such that the PstI site is closest to the napin 5' regulatory region. A fragment containing the napin 5' regulatory region, polylinker, and napin 3' region was removed from pCGN8619 by digestion with Asp718I. The fragment was blunt-ended by filling in the 5' overhangs with Klenow fragment then ligated into pCGN5139 that had been digested with Asp718I and HindIII and blunt-ended by filling in the 5' overhangs with Klenow fragment. A plasmid containing the insert oriented so that the napin promoter was closest to the blunted Asp718I site of pCGN5139 and the napin 3' was closest to the blunted HindIII site was subjected to sequence analysis to confirm both the insert orientation and the integrity of cloning junctions. The resulting plasmid was designated pCGN8623.

To produce high levels of stearidonic acid in *Brassica*, the *C. elegans* ω -3 desaturase can be combined with Δ 6- and Δ 12-desaturases from *Mortierella alpina*. PCGN5563-transformed plants may be crossed with pCGN5544-transformed plants expressing the Δ 6- and Δ 12-desaturases, described below.

The resulting F1 seeds can be analyzed for stearidonic acid content and selected F1 plants can be used for self-pollination to produce F2 seed, or as donors for production of dihaploids, or additional crosses.

An alternative method to combine the fat-1 cDNA with *M. alpina* Δ 6 and Δ 12 desaturases is to combine them on one T-DNA for transformation. The fat-1 coding region from pCGN5562 can be cut out as a PstI/NotI fragment and inserted into PstI/NotI digested pCGN8619. The transcriptional unit consisting of the napin 5' regulatory region, the fat-1 coding region, and the napin 3'-regulatory region can be cut out as a Sse8387I fragment and inserted into pCGN5544 cut with Sse8387I. The resulting plasmid would contain three napin transcriptional units containing the *C. elegans* ω -3 desaturase, *M. alpina* Δ 6 desaturase, and *M. alpina* Δ 12 desaturase, all oriented in the same direction as the 35S/nptII/tml transcriptional unit used for selection of transformed tissue.

20

Example 2

Over-Expression of Δ 15-desaturase Activity in Transgenic Canola

The Δ 15-desaturase activity of *Brassica napus* can be increased by over-expression of the Δ 15-desaturase cDNA clone.

A *B. napus* Δ 15-desaturase cDNA clone was obtained by PCR amplification of first-strand cDNA derived from *B. napus* cv. 212/86. The primers were based on published sequence: Genbank # L01418 Arondel et al, 1992 Science 258:1353-1355.

25

The following primers were used:

Bnd15-FORWARD

5'-CUACUACUACUAGAGCTCAGCGATGGTTGTTGCTATGGAC-3'

5 **Bnd15-REVERSE**

5'-CAUCAUCAUGAATTCTTAATTGATTTTAGATTTG-3'

These primers allowed the amplification of the entire coding region and added SacI and EcoRI sites to the 5'- and 3'-ends, respectively

10 The PCR product was subcloned into pAMP1 (GIBCOBRL) using the CloneAmp system (GIBCOBRL) to create pCGN5520. The sequence was verified by sequencing of both strands to be sure that the open reading frame remained intact. For seed specific expression, the Δ 15-desaturase coding region was cut out of pCGN5520 as a BamHI/SalI fragment and inserted between the BglII and XhoI sites of the pCGN7770, to create pCGN5557.

15 The PstI fragment of pCGN5557 containing the napin 5'-regulatory region, *B. napus* Δ 15-desaturase, and napin 3'-regulatory region was inserted into the PstI site of the binary vector, pCGN5138 to produce pCGN5558. pCGN5558 was introduced into *Brassica napus* via *Agrobacterium*-mediated transformation.

20 To produce high levels of stearidonic acid in *Brassica*, the Δ 15-desaturase can be combined with Δ 6- and Δ 12-desaturases from *Mortierella alpina*. PCGN5558-transformed plants may be crossed with pCGN5544-transformed plants expressing the Δ 6 and Δ 12-desaturases. The resulting F1 seeds are analyzed for stearidonic acid content. GC-FAME analysis of F1

25 half-seeds revealed a significant accumulation of SDA in the seed oil of the *Brassica* lines. SDA levels (18:4) of greater than approximately 25% were obtained in hemizygous lines and are provided in Table 1. Selected F1 plants can be used for self-pollination to produce F2 seed, or as donors for production of dihaploids, or additional crosses.

Table 1

Strain ID	16:0	18:0	18:1	18:2	18:2	18:3	18:3	18:3	18:3	18:4	20:0	20:1	20:2	22:0	22:1	22:2
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	6	1.34	2.97	9.58	9.58	52.79	34.76	18.03	C6912	25.21	0.7	0.56	0.16	0.41	0.03	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.45	0.86	10.42	9.06	9.06	49.08	25.68	23.4		23.45	0.5	0.84	0.55	0.49	0	0.03
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	5.8	2.36	12.5	11.13	11.13	47.47	18.86	28.61		17.55	1.01	0.86	0.3	0.85	0	0.07
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	3.65	0.66	14.26	14.97	14.97	50.94	23.3	27.64		13.22	0.43	0.88	0.23	0.48	0.04	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.86	2.42	18.74	14.23	14.23	46.22	23	23.22		10.67	0.89	0.92	0.18	0.7	0.02	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	6.57	1.07	16.79	14	14	48.98	32.88	16.09		10.24	0.52	0.94	0.22	0.39	0.02	0.01
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	5.85	2.09	8.81	19.12	19.12	50.89	12.03	38.86		9.09	1.39	0.78	0.45	1.23	0	0.04
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.69	2.04	17.46	21.1	21.1	43.38	24.28	19.1		8.5	0.73	0.96	0.37	0.56	0	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	5.43	1.69	16.59	22.2	22.2	44.4	16.57	27.83		6.34	0.9	1.03	0.32	0.81	0.03	0.05
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.28	1.34	18.83	27.24	27.24	40.54	20.91	19.63		5.03	0.73	0.88	0.27	0.7	0	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.47	1.38	21.43	26.89	26.89	39.04	18.78	20.26		4.06	0.73	0.91	0.41	0.48	0	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.77	1.12	18.4	31.1	31.1	38.51	19.62	18.88		3.52	0.64	0.8	0.21	0.7	0	0

(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.34	1.86	24.73	35.49	35.49	28.79	10.79	18	1.91	0.67	1.07	0.45	0.48	0	0.02
Strain ID*	16:0	18:0	18:1	18:2	18:2	18:3	18:3	18:3 C91215 - ALA	18:4	20:0	20:1	20:2	22:0	22:1	22:2
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.71	1.75	20.72	34.68	34.68	34.01	4.65	29.36	1.62	0.71	0.89	0.1	0.63	0	0
(5558-SP30021-26 X 5544-LP30108-6-16-1-3)	4.3	0.8	40.79	14.34	14.34	37	36.88	0.12	0	0.43	1.47	0.29	0.29	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	6.61	1.5	22.09	11.23	11.23	39.9	25.84	14.06	16.25	0.75	0.75	0.27	0.57	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	4.64	1.89	22.73	15.12	15.12	44.48	32.21	12.27	8.78	0.73	0.89	0.23	0.43	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	5.51	1.45	24.82	17.79	17.79	41.84	27.46	14.38	6.45	0.59	0.82	0.23	0.39	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	4.06	1.67	26.39	16.93	16.93	42.64	32.65	9.99	6	0.64	0.96	0.24	0.41	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	5.24	1.44	22.2	20.02	20.02	42.76	28.69	14.07	5.98	0.67	0.79	0.26	0.45	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	5.34	2.2	22.68	18.6	18.6	43.14	31.45	11.69	5.5	0.82	0.87	0.25	0.53	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	3.98	2.9	25.23	21.21	21.21	38.78	24.6	14.18	4.98	1.02	1.04	0.24	0.57	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	3.94	1.77	28.92	20.89	20.89	37.02	21.71	15.32	4.96	0.64	1.09	0.3	0.43	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	5.12	1.24	27.7	19.02	19.02	40.2	31.05	9.16	4.76	0.48	0.77	0.23	0.35	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	4.16	1.52	28.59	21.99	21.99	36.85	23.33	13.53	4.55	0.6	0.98	0.27	0.41	0	0
(5558-SP30021-19 X 5544-LP30108-6-16-1-1)	4.91	1.32	30.46	18.01	18.01	38.59	30.23	8.36	4.34	0.58	0.93	0.25	0.4	0	0

Strain ID	16:0	18:0	18:1	18:2	18:2	18:3	18:3	18:3	18:3	18:4	20:0	20:1	20:2	22:0	22:1	22:2
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.66	1.52	29.52	20.52	20.52	36.61	20.09	16.52	5.63	0.67	1.12	0.14	0.52	0	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	5.09	1.81	25.81	21.54	21.54	38.2	22.52	15.68	4.92	0.75	0.96	0.12	0.57	0.02	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.77	1.5	29.79	22.36	22.36	35.46	14.84	20.62	4.39	0.74	1.17	0.18	0.59	0.02	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.71	1.45	32.18	23.86	23.86	32.32	17	15.32	3.92	0.63	1.12	0.15	0.5	0.02	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.55	1.56	33.27	25.21	25.21	30.69	16.63	14.06	3.08	0.68	1.2	0.16	0.54	0.03	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	4.04	1.52	33.63	24.47	24.47	30.72	18.19	12.53	3.07	0.63	1.17	0.14	0.46	0	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.67	1.58	31.98	26.13	26.13	30.89	15.92	14.97	3.05	0.69	1.21	0.16	0.51	0	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.58	1.8	30.2	27.22	27.22	31.42	15.48	15.94	2.85	0.79	1.21	0.17	0.61	0.02	0.01	0.01
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	4.68	1.41	28.32	28	28	32.22	14.92	17.3	2.74	0.65	1.1	0.18	0.53	0.01	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.5	1.46	34.13	25.92	25.92	29.7	16.77	12.93	2.65	0.67	1.26	0.15	0.51	0.01	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.9	1.68	33.44	26.18	26.18	29.43	16.11	13.31	2.6	0.72	1.23	0.18	0.5	0.02	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.82	1.71	31.84	27.78	27.78	29.49	15.28	14.2	2.59	0.73	1.19	0.16	0.55	0.02	0	0
(5558-SP30021-36 X 5544-LP30108-6-16-1-1)	3.6	1.78	29.45	28.14	28.14	31.64	12.83	18.81	2.57	0.76	1.21	0.17	0.58	0	0	0

An alternative method to combine the *B. napus* $\Delta 15$ -desaturase with *M. alpina* $\Delta 6$ and $\Delta 12$ desaturases is to combine them on one T-DNA for transformation. The transcription cassette consisting of the napin 5'-regulatory region, the $\Delta 15$ -desaturase coding region, and the napin 3'-regulatory region can be cut out of pCGN5557 as a *Swa*I fragment and inserted into *Swa*I-digested pCGN5544. The resulting plasmid, pCGN5561, contains three napin transcriptional units containing the *M. alpina* $\Delta 6$ desaturase, the *B. napus* $\Delta 15$ -desaturase, and the *M. alpina* $\Delta 12$ desaturase, all oriented in the same direction as the 35S/nptII/tml transcriptional unit used for selection of transformed tissue. In addition, the *C. elegans* ω -3 desaturase coding sequence was also cloned into pCGN5544 to create the construct pCGN5565.

Pooled T2 seeds of plants containing 5561 contain significant amounts of SDA (18:4), shown in Table 2. Levels of greater than about 7% SDA are obtained in pooled 5561 segregating seed. Furthermore, significant levels of SDA were obtained from seeds of 5565 *Brassica* lines, also shown in Table 2. As shown in Table 2, with constructs 5561 and 5565, levels of SDA ranging from about 0.8 wt% to greater than about 7 wt% can be obtained.

Table 2

STRAIN ID	16:0	16:1	18:0	18:1	18:2 C6,9	18:2 C9,12	18:3 C6,9,12	18:3 C9,12,15	18:4	20:0	20:1	20:2	22:0	22:1	22:2
5561-6	4.46	0.21	3.5	22.85	0	18.33	18.71	21.61	7.79	1.04	0.76	0.19	0.47	0	0
5561-4	4.14	0.15	2.62	33.07	0	21.07	17.61	14.56	4.39	0.87	0.92	0.14	0.39	0	0.02
5561-2	4.26	0.15	2.21	30.42	0	22.02	21.06	12.88	4.25	0.89	0.98	0.2	0.51	0	0.02
5561-8	4.29	0.18	2	33.05	0	22.44	16.23	15.3	3.95	0.84	0.96	0.19	0.43	0	0.04
5561-3	3.95	0.12	2.04	32.93	0	24.48	17.42	13.33	3.27	0.79	0.94	0.21	0.4	0.03	0.03
5561-7	4.26	0.17	2.02	38.4	0	23.3	13.35	13.3	2.73	0.75	1.06	0.16	0.41	0	0
5561-13	4.38	0.18	1.86	58.94	0.65	13.98	7.1	8.26	1.88	0.77	1.27	0.29	0.35	0.03	0
5561-15	4.29	0.15	2.3	40.96	0	26.63	8.58	12.98	1.51	0.83	1.07	0.19	0.45	0	0.02
5561-1	4.25	0.15	1.91	47.41	0	24.46	5.56	12.81	1	0.72	1.14	0.15	0.39	0	0
5561-5	4.07	0.16	1.96	52.29	0	20.88	5.02	12.17	0.97	0.72	1.16	0.21	0.28	0	0.06
CONTROL	3.89	0.21	1.65	58.48	0	22.44	0	11.03	0	0.6	1.15	0.16	0.27	0.01	0

Table 2 Continued

STRAIN ID	16:0	18:0	18:1	18:2_C69	18:2-LA	18:3-GLA	18:3-ALA	18:4	20:0
5565-SP30021-7	4.03	1.93	41.24	0.43	14.46	21.39	6.62	7.38	0.68
5565-SP30021-12	3.95	2.46	40.19	0	30.35	7.3	10.92	2.57	0.7
5565-SP30021-9	4.03	1.82	35.76	0	33.49	8.63	11.58	2.54	0.51
5565-SP30021-3	3.86	1.8	32.3	0	35.57	11.3	10.05	2.37	0.61
5565-SP30021-1	3.98	1.92	59.99	1.84	11.24	8.07	7.46	2.32	0.76
5565-SP30021-8	4.67	1.72	38.95	0	30.38	8.99	10.83	2.25	0.52
5565-SP30021-10	4.03	1.43	47.04	0	26.96	5.97	11.1	1.35	0.52
5565-SP30021-6	3.87	1.77	46.73	0	28.79	5.31	10.4	0.79	0.56

Control	3.89	1.65	58.48	0	22.44	0	11.03	0	0.6
---------	------	------	-------	---	-------	---	-------	---	-----

Example 3

Expression of $\Delta 5$ Desaturase in Plants Expression in Leaves

5 Ma29 is a putative *M. alpina* $\Delta 5$ desaturase as determined by sequence homology. This experiment was designed to determine whether leaves expressing Ma29 (as determined by Northern) were able to convert exogenously applied DGLA (20:3) to ARA (20:4).

10 The Ma29 desaturase cDNA was modified by PCR to introduce convenient restriction sites for cloning. The desaturase coding region has been inserted into a d35 cassette under the control of the double 35S promoter for expression in *Brassica* leaves (pCGN5525) following standard protocols (*see* USPN 5,424,200 and USPN 5,106,739). Transgenic *Brassica* plants containing pCGN5525 were generated following standard protocols (*see* USPN 5,188,958 and USPN
15 5,463,174).

 In the first experiment, three plants were used: a control, LPO04-1, and two transgenics, 5525-23 and 5525-29. LP004 is a low-linolenic *Brassica* variety. Leaves of each were selected for one of three treatments: water, GLA or DGLA. GLA and DGLA were purchased as sodium salts from NuChek Prep and dissolved
20 in water at 1 mg/ml. Aliquots were capped under N₂ and stored at -70 degrees C. Leaves were treated by applying a 50 μ l drop to the upper surface and gently spreading with a gloved finger to cover the entire surface. Applications were made approximately 30 minutes before the end of the light cycle to minimize any photo-oxidation of the applied fatty acids. After 6 days of treatment one leaf from each
25 treatment was harvested and cut in half through the mid rib. One half was washed with water to attempt to remove unincorporated fatty acid. Leaf samples were

lyophilized overnight, and fatty acid composition determined by gas chromatography (GC). The results are shown in Table 3.

Table 3
Fatty Acid Analysis of Leaves from Ma29 Transgenic Brassica Plants

Treatment	SPL #	16:00 %	16:01 %	18:00 %	18:01 %	18:02 %	18:3g %	18:03 %	18:04 %	20:00 %	20:01 %
Water	33	12.95	0.08	2.63	2.51	16.76	0	45.52	0	0.09	0
	34	13.00	0.09	2.67	2.56	16.86	0	44.59	0	0.15	0
	35	14.13	0.09	2.37	2.15	16.71	0	49.91	0	0.05	0.01
	36	13.92	0.08	2.32	2.07	16.16	0	50.25	0	0.05	0
	37	13.79	0.11	2.10	2.12	15.90	0.08	46.29	0	0.54	0.01
GLA	38	12.80	0.09	1.94	2.08	14.54	0.11	45.61	0	0.49	0.01
	39	12.10	0.09	2.37	2.10	14.85	1.63	43.66	0	0.53	0
	40	12.78	0.10	2.34	2.22	15.29	1.72	47.22	0	0.50	0.02
	41	13.71	0.07	2.68	2.16	15.92	2.12	46.55	0	0.09	0
	42	14.10	0.07	2.75	2.35	16.66	1.56	46.41	0	0.09	0.01
DGLA	43	13.62	0.09	2.22	1.94	14.68	2.42	46.69	0	0.51	0.01
	44	13.92	0.09	2.20	2.17	15.22	2.30	46.05	0	0.53	0.02
	45	12.45	0.14	2.30	2.28	15.65	0.07	44.62	0	0.12	0.01
	46	12.67	0.15	2.69	2.50	15.96	0.09	42.77	0	0.56	0.01
	47	12.56	0.23	3.40	1.98	13.57	0.03	45.52	0	0.51	0.01
	48	13.07	0.24	3.60	2.51	13.54	0.04	45.13	0	0.50	0.01
	49	13.26	0.07	2.81	2.34	16.04	0.04	43.89	0	0.59	0
	50	13.53	0.07	2.84	2.41	16.07	0.02	44.90	0	0.60	0.01

Table 3 - Continued
Fatty Acid Analysis of Leaves from Ma29 Transgenic Brassica Plants

Treatment	SPL #	20:02	20:03	20:04	20:05	22:00	22:01	22:02	22:03	22:06	24:0	24:1
		%	%	%	%	%	%	%	%	%	%	%
Water	33	0	0	0.29	0	0.01	0.09	16.26	0	0	0.38	0.18
	34	0.01	0	0.26	0	0.14	0.10	16.82	0.02	0.05	0.36	0.27
	35	0.01	0	0.25	0	0.12	0.06	11.29	0.04	0.05	0.29	0.25
	36	0	0.01	0.26	0	0.07	0.04	11.82	0.03	0.36	0.28	0.21
	37	0.02	0	0.21	0	0.18	0.08	15.87	0.06	0.20	0.30	0.17
GLA	38	0.01	0	0.24	0	0.15	0.07	13.64	0.09	0.08	5.89	0.23
	39	0.02	0.01	0.27	0	0.10	0.08	16.25	3.42	0.19	0.37	0.17
	40	0.01	0	0.27	0	0.10	0.10	14.74	0.05	0.10	0.36	0.14
	41	0	0	0.27	0	0.20	0.10	13.15	0.13	0.29	0.33	0.20
	42	0	0	0.28	0	0.11	0.11	12.60	0.02	0.24	0.38	0.13
DGLA	43	0.01	0	0.28	0	0.10	0.03	14.73	0.01	0.24	0.34	0.14
	44	0.02	0	0.26	0	0.13	0.07	14.43	0.05	0.16	0.33	0.17
	45	0.06	1.21	0.26	0	0.07	0.07	18.67	0.02	0.21	0.36	0.13
	46	0	1.94	0.27	0	0.11	0.09	17.97	0.09	0.39	0.41	0.11
	47	0.01	0.69	0.96	0	0.11	0.07	17.96	0	0.22	0.49	0.20
	48	0.01	0.70	0.74	0	0.14	0.09	17.14	0.05	0.32	0.52	0.10
	49	0	0.35	1.11	0	0.10	0.07	17.26	0.07	0.23	0.39	0.18
	50	0	0.20	0.87	0	0.21	0.07	15.73	0.04	0.15	0.37	0.18

Leaves treated with GLA contained from 1.56 to 2.4 wt% GLA. The fatty acid analysis showed that the lipid composition of control and transgenic leaves was essentially the same. Leaves of control plants treated with DGLA contained 1.2-1.9 w% DGLA and background amounts of ARA (.26-.27 wt%).

5 Transgenic leaves contained only .2-.7 wt% DGLA, but levels of ARA were increased (.74-1.1 wt%) indicating that the DGLA was converted to ARA in these leaves.

Expression in Seed

10 The purpose of this experiment was to determine whether a construct with the seed specific napin promoter would enable expression in seed.

The Ma29 cDNA was modified by PCR to introduce *XhoI* cloning sites upstream and downstream of the start and stop codons, respectively, using the following primers:

Madxho-forward:

15 5'-CUACUACUACUACTCGAGCAAGATGGGAACGGACCAAGG

Madxho-reverse:

5'-CAUCAUCAUCAUCTCGAGCTACTCTTCCTTGGGACGGAG

20 The PCR product was subcloned into pAMP1 (GIBCOBRL) using the CloneAmp system (GIBCOBRL) to create pCGN5522 and the $\Delta 5$ desaturase sequence was verified by sequencing of both strands.

25 For seed-specific expression, the Ma29 coding region was cut out of pCGN5522 as an *XhoI* fragment and inserted into the *SalI* site of the napin expression cassette, pCGN3223, to create pCGN5528. The *HindIII* fragment of pCGN5528 containing the napin 5' regulatory region, the Ma29 coding region, and the napin 3' regulatory region was inserted into the *HindIII* site of pCGN1557 to create pCGN5531. Two copies of the napin transcriptional unit were inserted in tandem. This tandem construct can permit higher expression of the desaturases per genetic loci. pCGN5531 was introduced into *Brassica napus* cv.LP004 via *Agrobacterium* mediated transformation.

The fatty acid composition of twenty-seed pools of mature T2 seeds was analyzed by GC. Table 2 shows the results obtained with independent transformed lines as compared to non-transformed LP004 seed. The transgenic seeds containing pCGN5531 contain two fatty acids that are not present in the control seeds, identified as taxoleic acid (5,9-18:2) and pinolenic acid (5,9,12-18:3), based on their elution relative to oleic and linoleic acid. These would be the expected products of $\Delta 5$ desaturation of oleic and linoleic acids. No other differences in fatty acid composition were observed in the transgenic seeds.

10

Example 4

Production of $\Delta 5$ -desaturated Fatty Acids in Transgenic Plants

The construction of pCGN5531 ($\Delta 5$ -desaturase) and fatty acid composition of T2 seed pools is described in Example 3. This example takes the seeds through one more generation and discusses ways to maximize the $\Delta 5$ -desaturated fatty acids.

Example 3 describes the fatty acid composition of T2 seed pools of pCGN5531-transformed *B. napus* cv. LP004 plants. To investigate the segregation of $\Delta 5$ -desaturated fatty acids in the T2 seeds and to identify individual plants to be taken on to subsequent generations, half-seed analysis was done. Seeds were germinated overnight in the dark at 30 degrees on water-soaked filter paper. The outer cotyledon was excised for GC analysis and the rest of the seedling was planted in soil. Results of some of these analyses are shown in the accompanying Table 4. $\Delta 5,9-18:2$ accumulated to as high as 12% of the total fatty acids and $\Delta 5,9,12-18:3$ accumulated to up to 0.77% of the fatty acids. These and other individually selected T2 plants were grown in the greenhouse to produce T3 seed.

25

Table 4
Composition of T2 Pooled Seed

	16:0	16:1	18:0	18:1	Δ5,9 18:2	18:2	Δ5,9,12,18:3	18:3	20:0	20:1	20:2	22:0	22:1	24:0
	%	%	%	%	%	%	%	%	%	%	%	%	%	%
LP004 control	3.86	0.15	3.05	69.1	0	18.51	0.01	1.65	1.09	1.40	0.03	0.63	0.05	0.42
5531-1	4.26	0.15	3.23	62.33	4.07	21.44	0.33	1.38	0.91	1.04	0.05	0.41	0.03	0.27
5531-2	3.78	0.14	3.37	66.18	4.57	17.31	0.27	1.30	1.03	1.18	0	0.47	0.01	0.30
5531-6	3.78	0.13	3.47	63.61	6.21	17.97	0.38	1.34	1.04	1.14	0.05	0.49	0.02	0.26
5531-10	3.96	0.17	3.28	63.82	5.41	18.58	0.32	1.43	0.98	1.11	0.02	0.50	0	0.31
5531-16	3.91	0.17	3.33	64.31	5.03	18.98	0.33	1.39	0.96	1.11	0	0.44	0	0
5531-28	3.81	0.13	2.58	62.64	5.36	20.95	0.45	1.39	0.83	1.15	0.01	0.36	0.05	0.21

Table 4
Fatty acid analysis of selected T2 half-seeds from pCGN5531-L-P004 events

CYCLE ID	SPL NO	STRAIN ID	12:0	14:0	16:0	16:1	18:0	18:1	18:2 Δ5,9	18:2 Δ9,12	18:3 Δ5,9,12	18:3 Δ9,12,15
97XX1539	93	5531-LP004-6	0.03	0.07	3.92	0.17	3.5	61.32	12.22	15.36	0.77	1.36
97XX1539	29	5531-LP004-6	0.01	0.04	3.6	0.09	3.23	63.77	10.63	14.47	0	1.22
97XX1539	38	5531-LP004-6	0.01	0.05	3.71	0.09	3.02	65.13	10.57	13.98	0	1.06
97XX1539	41	5531-LP004-6	0.01	0.05	3.64	0.07	3.22	62.51	9.7	16.63	0	1.28
97XX1539	18	5531-LP004-6	0.02	0.06	3.69	0.09	3.36	63.79	9.63	15.29	0.63	1.15
97XX1539	85	5531-LP004-6	0.01	0.06	3.6	0.09	3.54	64.81	9.54	13.69	0.6	1.26
98GC0023	98	5531-LP004-23	0.01	0.05	3.5	0.09	3.12	64.97	9.92	13.62	0.55	1.25
98GC0023	32	5531-LP004-23	0.01	0.05	3.43	0.08	2.62	65.21	9.83	14.28	0.59	1.15
98GC0023	78	5531-LP004-23	0.01	0.05	3.45	0.07	2.78	64.97	9.34	14.69	0.58	1.17
98GC0023	86	5531-LP004-23	0.01	0.05	3.32	0.08	2.7	64.18	9.08	15.99	0.68	1.18
98GC0023	67	5531-LP004-23	0.01	0.04	3.49	0.08	3.03	64.14	8.78	15.95	0.62	1.08
98GC0023	52	5531-LP004-23	0.01	0.03	3.38	0.07	2.56	67.44	8.65	13.55	0.5	1.02

To maximize the accumulation of $\Delta 5,9$ 18:2 in seed oil, the pCGN5531 construct could be introduced into a high oleic acid variety of canola. A high-oleic variety could be obtained by mutation, so-suppression, or antisense suppression of the $\Delta 12$ and $\Delta 15$ desaturases or other necessary co-factors.

5 To maximize accumulation of $\Delta 5,9,12$ 18:3 in canola, the pCGN5531 construct could be introduced into a high linoleic strain of canola. This could be achieved by crossing pCGN5531-transformed plants with pCGN5542-(*M. alpina* $\Delta 12$ -desaturase) transformed plants. Alternatively, the $\Delta 5$ and $\Delta 12$ desaturases could be combined on one T-DNA for transformation. The
10 transcriptional unit consisting of the napin 5'-regulatory region, the *M. alpina* $\Delta 12$ -desaturase coding region, and the napin 3'-regulatory region can be cut out of pCGN5541 as a NotI fragment. NotI/XbaI linkers could be ligated and the resulting fragment inserted into the XbaI site of pCGN5531. The resulting plasmid would contain three napin transcriptional units containing the *M. alpina*
15 $\Delta 12$ desaturase, and two copies of the napin/*M. alpina* $\Delta 5$ desaturase/napin unit, all oriented in the same direction as the 35S/nptII/tml transcriptional unit used for selection of transformed tissue.

Example 5

20 Expression of *M. alpina* $\Delta 6$ Desaturase in *Brassica napus*

A nucleic acid sequence from a partial cDNA clone, Ma524, encoding a $\Delta 6$ fatty acid desaturase from *Mortierella alpina* was obtained by random sequencing of clones from the *M. alpina* cDNA library. The Ma524 cDNA was modified by PCR to introduce cloning sites using the following primers:

25

Ma524PCR-1

5'-CUACUACUACUATCTAGACTCGAGACCATGGCTGCTGCT
CCAGTGTG

Ma524PCR-2

5'-CAUCAUCAUCAUAGGCCTCGAGTTACTGCGCCTTACCCAT

5 These primers allowed the amplification of the entire coding region and added *Xba*I and *Xho*I sites to the 5'-end and *Xho*I and *Stu*I sites to the 3' end. The PCR product was subcloned into pAMP1 (GIBCOBRL) using the CloneAmp system (GIBCOBRL) to create pCGN5535 and the $\Delta 6$ desaturase sequence was verified by sequencing of both strands.

Construction of pCGN5544

10 Plant expression constructs were prepared to express the *Mortierella alpina* $\Delta 6$ desaturase and the *Mortierella alpina* $\Delta 12$ desaturase in a plant host cell. The constructs prepared utilized transcriptional initiation regions derived from genes preferentially expressed in a plant seed. Isolation of the cDNA sequences encoding the *M. alpina* $\Delta 6$ desaturase and *M. alpina* $\Delta 12$ desaturase are described in PCT Publications WO 98/46763 and WO 98/46764, the
15 entireties of which are incorporated herein by reference.

 For seed-specific expression, the Ma524 coding region was cut out of pCGN5535 as an *Xho*I fragment and inserted into the *Sal*I site of the napin expression cassette, pCGN3223, to create pCGN5536. The *Not*I fragment of
20 pCGN5536 containing the napin 5' regulatory region, the Ma524 coding region, and the napin 3' regulatory region was inserted into the *Not*I site of pCGN1557 to create pCGN5538.

 The 5542 cDNA, encoding the *M. alpina* $\Delta 12$ desaturase, was modified by PCR to introduce cloning sites using the following primers:

25

Ma648PCR-for

5'-CUACUACUACUAGGATCCATGGCACCTCCCAACT

Ma648PCR-for

5'-

CAUCAUCAUGGTACCTCGAGTTACTTCTTGAAAAAGAC

These primers allowed the amplification of the entire coding region and added a BamHI site to the 5' end and KpnI and XhoI sites to the 3' end. The PCR product was subcloned into pAMP1 (Gibco-BRL, Gaithersburg, MD) using the CloneAmp system (Gibco-BRL) to create pCGN5540, and the $\Delta 12$ desaturase sequence was verified by sequencing of both strands.

A seed preferential expression construct was prepared for the $\Delta 12$ desaturase cDNA sequence. The Ma648 coding region was cut out of pCGN5540 as a BamHI/XhoI fragment and inserted between the BglII and XhoI sites of the napin expression cassette, pCGN3223 (described in USPN 5,639,790), to create pCGN5542.

In order to express the *M. alpina* $\Delta 6$ and $\Delta 12$ desaturase sequences from the same T-DNA, the following construct for seed-preferential expression was prepared.

The NotI fragment of pCGN5536 containing the napin 5' transcriptional initiation region, the Ma524 coding region, and the napin 3' transcriptional termination region was inserted into the NotI site of pCGN5542 to create pCGN5544. The expression cassettes were oriented in such a way that the direction of transcription from Ma524 and Ma648 and the nptII marker is the same.

For seed-specific expression, the Ma524 coding region was cut out of pCGN5535 as an XhoI fragment and inserted into the SalI site of the napin expression cassette, pCGN3223, to create pCGN5536. The NotI fragment of pCGN5536 containing the napin 5' regulatory region, the Ma524 coding region, and the napin 3' regulatory region was inserted into the NotI site of pCGN1557 to create pCGN5538. pCGN5538 was introduced into *Brassica napus* cv.LP004 via *Agrobacterium* mediated transformation.

Maturing T2 seeds were collected from 6 independent transformation events in the greenhouse. The fatty acid composition of single seeds was analyzed by GC. Table 5 shows the results of control LP004 seeds and six 5538

lines. All of the 5538 lines except #8 produced seeds containing GLA. Presence of GLA segregated in these seeds as is expected for the T2 selfed seed population. In addition to GLA, the *M. alpina* $\Delta 6$ desaturase is capable of producing 18:4 (stearidonic) and another fatty acid: $\Delta 6,9-18:2$.

5

Table 5

Fatty Acid Analysis of Seeds from Mas24 Transgenic *Brassica* Plants

SPL	#	16:0	16:1	18:0	18:1	6,9 18:2	18:2	18:3ga	18:3	18:4	20:1	22:0	22:1	24:0	24:1
		%	%	%		%	%	%	%	%	%	%	%	%	%
	LPO04-1	4.33	0.21	3.78	72.49	0	13.97	0	1.7	0	1.34	0.71	0.02	0.58	0.27
		4.01	0.16	3.09	73.59	0	14.36	0.01	1.4	0	1.43	0.66	0.02	0.5	0.2
		4.12	0.19	3.56	70.25	0	17.28	0	1.57	0	1.28	0.5	0.02	0.39	0.2
		4.22	0.2	2.7	70.25	0	17.86	0	1.61	0	1.31	0.53	0.02	0.4	0.24
		4.02	0.16	3.41	72.91	0	14.45	0.01	1.45	0	1.37	0.7	0.02	0.51	0.26
		4.22	0.18	3.23	71.47	0	15.92	0.01	1.52	0	1.32	0.69	0.02	0.51	0.27
		4.1	0.16	3.47	72.06	0	15.23	0	1.52	0	1.32	0.63	0.03	0.49	0.23
		4.01	0.17	3.71	72.98	0	13.97	0.01	1.41	0	1.45	0.74	0.03	0.58	0.23
		4.04	0.16	3.57	70.03	0	17.46	0	1.5	0	1.33	0.61	0.03	0.36	0.24
	5538-1-1	4.61	0.2	3.48	68.12	1.37	10.68	7.48	1.04	0.33	1.19	0.49	0.02	0.33	0.13
		4.61	0.22	3.46	68.84	1.36	10.28	7.04	1.01	0.31	1.15	0.48	0.02	0.39	0
		4.78	0.24	3.24	65.86	0	21.36	0	1.49	0	1.08	0.46	0.02	0.38	0.22
		4.84	0.3	3.89	67.64	1.67	9.9	6.97	1.02	0.36	1.14	0.53	0.02	0.5	0.18
		4.64	0.2	3.58	64.5	3.61	8.85	10.14	0.95	0.48	1.19	0.47	0.01	0.33	0.12
		4.91	0.27	3.44	66.51	1.48	11.14	7.74	1.15	0.33	1.08	0.49	0.02	0.34	0.13
		4.87	0.22	3.24	65.78	1.27	11.92	8.38	1.2	0	1.12	0.47	0.02	0.37	0.16
		4.59	0.22	3.4	70.77	0	16.71	0	1.35	0	1.14	0.48	0.02	0.39	0.15
		4.63	0.23	3.51	69.66	2.01	8.77	7.24	0.97	0	1.18	0.52	0.02	0.3	0.11
		4.56	0.19	3.55	70.68	0	16.89	0	1.37	0	1.22	0.54	0.02	0.22	0.03
	5538-3-1	4.74	0.21	3.43	67.52	1.29	10.91	7.77	1.03	0.28	1.11	0.5	0.02	0.35	0.14
		4.72	0.21	3.24	67.42	1.63	10.37	8.4	0.99	0	1.12	0.49	0.02	0.36	0.15

Table 5

Fatty Acid Analysis of Seeds from Ma524 Transgenic *Brassica* Plants

SPL #	16:0	16:1	18:0	18:1	6,9 18:2	18:2	18:3ga	18:3	18:4	20:1	22:0	22:1	24:0	24:1
	%	%	%		%	%	%	%	%	%	%	%	%	%
	-3	4.24	0.21	3.52	71.31	0	16.53	0	1.33	0	1.12	0.45	0.4	0.14
	-4	4.64	0.21	3.45	67.92	1.65	9.91	7.97	0.91	0.33	1.14	0.47	0.37	0.14
	-5	4.91	0.25	3.31	67.19	0	19.92	0.01	1.39	0	1.05	0.48	0.37	0.14
	-6	4.67	0.21	3.25	67.07	1.23	11.32	8.35	0.99	0	1.16	0.47	0.33	0.16
	-7	4.53	0.19	2.94	64.8	4.94	8.45	9.95	0.93	0.44	1.13	0.37	0.27	0.12
	-8	4.66	0.22	3.68	67.33	0.71	12	6.99	1.1	0.24	1.18	0.48	0.36	0.17
	-9	4.65	0.24	3.11	67.42	0.64	12.71	6.93	1.16	0.25	1.08	0.45	0.32	0.17
	-10	4.88	0.27	3.33	65.75	0.86	12.89	7.7	1.1	0.24	1.08	0.46	0.34	0.16
	5538-4-1	4.65	0.24	3.8	62.41	0	24.68	0	1.6	0.01	0.99	0.45	0.33	0.13
	-2	5.37	0.31	3	57.98	0.38	18.04	10.5	1.41	0	0.99	0.48	0.3	0.19
	-3	4.61	0.22	3.07	63.62	0.3	16.46	7.67	1.2	0	1.18	0.45	0.29	0.14
	-4	4.39	0.19	2.93	65.97	0	22.36	0	1.45	0	1.17	0.41	0.32	0.15
	-5	5.22	0.29	3.85	62.1	2.35	10.25	11.39	0.93	0.41	1.04	0.6	0.47	0.17
	-6	4.66	0.18	2.85	66.79	0.5	13.03	7.66	0.97	0.22	1.28	0.42	0.31	0.14
	-7	4.85	0.26	3.03	57.43	0.26	28.04	0.01	2.59	0.01	1.13	0.56	0.4	0.23
	-8	5.43	0.28	2.94	54.8	1.84	13.79	15.67	1.36	0.53	1.1	0.55	0.35	0.19
	-9	4.88	0.24	3.32	62.3	0.58	14.86	9.04	1.34	0.29	1.13	0.52	0.37	0.19
	-10	4.53	0.2	2.73	64.2	0.07	24.15	0	1.52	0	1.09	0.39	0.27	0.17
	5538-5-1	4.5	0.15	3.35	66.71	0.88	11.7	8.38	1.04	0.3	1.24	0.49	0.29	0.17
	-2	4.77	0.23	3.06	62.67	0.68	15.2	8.8	1.31	0.28	1.15	0.46	0.3	0.19
	-3	4.59	0.22	3.61	64.35	2.29	9.95	10.57	1.01	0.45	1.21	0.48	0.26	0.16

Table 5

Fatty Acid Analysis of Seeds from Mas24 Transgenic Brassica Plants

SPL #	16:0	16:1	18:0	18:1	6,9 18:2	18:2	18:3ga	18:3	18:4	20:1	22:0	22:1	24:0	24:1
	%	%	%	%	%	%	%	%	%	%	%	%	%	%
-4	4.86	0.26	3.4	67.69	0.65	12.24	6.61	1.09	0.23	1.07	0.45	0.02	0.32	0.14
-5	4.49	0.21	3.3	69.25	0.04	16.51	2.18	1.2	0	1.11	0.44	0.02	0.33	0.16
-6	4.5	0.21	3.47	70.48	0.08	14.9	2.19	1.22	0	1.13	0.49	0.02	0.33	0.16
-7	4.39	0.21	3.44	67.59	2.38	9.24	8.98	0.89	0	1.18	0.44	0.02	0.28	0.14
-8	4.52	0.22	3.17	68.33	0.01	18.91	0.73	1.32	0.01	1.08	0.45	0.02	0.29	0.17
-9	4.68	0.2	3.05	64.03	1.93	11.03	11.41	1.02	0.01	1.15	0.39	0.02	0.21	0.15
-10	4.57	0.2	3.1	67.21	0.61	12.62	7.68	1.07	0.25	1.14	0.43	0.02	0.25	0.15
5538-8-1	4.95	0.26	3.14	64.04	0	23.38	0	1.54	0	0.99	0.42	0.02	0.38	0.17
-2	4.91	0.26	3.71	62.33	0	23.97	0	1.77	0	0.95	0.53	0.02	0.42	0.19
-3	4.73	0.25	4.04	63.83	0	22.36	0.01	1.73	0	1.05	0.55	0.02	0.45	0.16
-4	5.1	0.35	3.8	60.45	0	24.45	0.01	2.13	0	1.07	0.65	0.03	0.53	0.24
-5	4.98	0.3	3.91	62.48	0	23.44	0	1.77	0	1.01	0.51	0.01	0.43	0.21
-6	4.62	0.21	3.99	66.14	0	20.38	0	1.48	0	1.15	0.53	0.02	0.48	0.19
-7	4.64	0.22	3.55	64.6	0	22.65	0	1.38	0	1.09	0.45	0.02	0.41	0.19
-8	5.65	0.38	3.18	56.6	0	30.83	0.02	0.02	0	0.98	0.55	0.03	0.39	0.26
-9	8.53	0.63	6.9	51.76	0	26.01	0	0.01	0	1.41	1.21	0.07	0.96	0.33
-10	5.52	0.4	3.97	57.92	0	28.95	0	0.02	0	0.95	0.52	0.02	0.41	0.16
5538-10-1	4.44	0.19	3.5	68.42	0	19.51	0	1.32	0	1.14	0.45	0.02	0.31	0.16
-2	4.57	0.21	3.07	66.08	0	21.99	0.01	1.36	0	1.12	0.41	0.02	0.31	0.16
-3	4.63	0.21	3.48	67.43	0	20.27	0.01	1.32	0	1.12	0.46	0.02	0.21	0.08
-4	4.69	0.19	3.22	64.62	0	23.16	0	1.35	0	1.08	0.46	0.02	0.33	0.2

Table 5

Fatty Acid Analysis of Seeds from Ma524 Transgenic *Brassica* Plants

SPL #	16:0 %	16:1 %	18:0 %	18:1	6,9 18:2 %	18:2 %	18:3ga %	18:3 %	18:4 %	20:1 %	22:0 %	22:1 %	24:0 %	24:1 %
-5	4.58	0.2	3.4	68.75	0	20.17	0.01	0.02	0	1.1	0.45	0.02	0.34	0.17
-8	4.55	0.21	0	73.55	0.05	14.91	2.76	1.21	0.07	1.24	0.51	0.02	0.19	0
-9	4.58	0.21	3.28	66.19	0	21.55	0	1.35	0	1.12	0.43	0.02	0.33	0.16
-10	4.52	0.2	3.4	68.37	0	19.33	0.01	1.3	0	1.13	0.46	0.02	0.35	0.18

Crosses were made between transgenic *Brassica* 5544 lines producing GLA and standard non-transformed canola varieties. Crosses between 5544 lines with Quantum, Eagle, and Ebony were conducted.

5 F1 half seeds were analyzed for SDA content and selected plants were grown and allowed to self-pollinate to produce F2 seeds. GC-FAME analysis of both single seed and half-seed samples from such crosses revealed accumulation of significant levels of SDA (Table 6). Half-seed analysis of 5544-LP108-6-16 with canola variety Eagle yielded a level of approximately 6.3% SDA. Analysis of F2 seed from a cross of 5544-LP108-12-1 with the canola variety Ebony
10 produced levels of SDA as high as about 7.4% SDA.

Table 6

STRAIN ID	16:0	16:1	18:0	18:1	18:2	18:3	18:3	18:3	18:4	20:0	20:1	20:2	22:0
(SP30035-46 x 5544-LP30108-6-16)	6.34	0.84	1.9	4.7	0	14.81	56.73	C6912	3.78	2.12	0.66	0.59	1.04
(SP30035-46 x 5544-LP30108-6-16)	10.18	1.43	4.23	4.34	0	15.96	48.78	C912	3.79	2.65	0.72	0.77	1.32
(SP30035-46 x 5544-LP30108-6-16)	4.81	0.45	2.53	12.2	0	21.61	46.74	C91215	4.83	0.98	0.79	0.4	0.43
(SP30035-46 x 5544-LP30108-6-16)	4.74	0.48	3.33	16.06	0	20.68	43.02	C912	4.82	1.25	0.7	0.33	0.75
(SP30035-46 x 5544-LP30108-6-16)	6.02	0.53	1.25	17.29	0	27.34	33.97	C6912	7.52	0.85	0.77	0.27	0.59
(SP30035-46 x 5544-LP30108-6-16)	3.68	0.13	1.99	19.75	0.09	22.75	39.98	C912	5.76	0.8	0.87	0.27	0.44
(SP30052-7 x 5544-LP30108-12-1)	8.92	0.96	1.64	14.61	0	18.69	36.98	C912	7.44	1.01	0.49	0.49	0.95
(SP30052-7 x 5544-LP30108-12-1)	9.02	0.89	1.88	10.69	0	16.73	43.39	C912	6.8	1.05	0.57	0.75	1.07
(SP30052-7 x 5544-LP30108-12-1)	7.76	0.59	1.86	8.15	0	16.04	52.24	C912	4.65	1.04	0.59	0.69	0.83
(SP30052-7 x 5544-LP30108-12-1)	9.21	0.87	2.23	17.44	0	18.77	36.87	C912	6.79	0.84	0.64	0.31	0.71
(SP30052-7 x 5544-LP30108-12-1)	5.76	0.31	1.6	20.38	0	24.36	29.94	C912	10.9	0.69	0.78	0.23	0.48
(SP30052-7 x 5544-LP30108-12-1)	4.03	0.22	1.3	16.87	0	19.3	46.67	C912	5.33	0.53	0.75	0.35	0.37
(SP30052-7 x 5544-LP30108-12-1)	4.66	0.29	4.47	18.09	0.05	19.07	41.92	C912	5.06	1.13	0.73	0.35	0.57
(SP30052-7 x 5544-LP30108-12-1)	4.91	0.26	3.13	18.16	0	18.53	43.99	C912	4.64	1.01	0.79	0.37	0.66

Example 6

Production of Δ 6,9 18:2 in Canola Oil

Example 5 described construction of pCGN5538 designed to express the
5 *M. alpina* Δ 6 desaturase in seeds of transgenic canola. Table 4 in that example
showed examples of single seed analyses from 6 independent transgenic events.
Significant amounts of GLA were produced, in addition to the Δ -6,9 18:2 fatty
acid.

A total of 29 independent pCGN5538-transformed transgenic plants of
10 the low-linolenic LP004 cultivar were regenerated and grown in the greenhouse.
Table 7 shows the fatty acid composition of 20-seed pools of T2 seed from each
event. Seven of the lines contained more than 2% of the Δ -6,9 18:2 in the seed
pools. To identify and select plants with high amounts of Δ -6,9 18:2 to be taken
on to subsequent generations, half-seed analysis was done. Seeds were
15 germinated overnight in the dark at 30 degrees on water-soaked filter paper.
The outer cotyledon was excised for GC analysis and the rest of the seedling
was planted in soil. Based on results of fatty acid analysis, selected T2 plants
were grown in the greenhouse to produce T3 seed. The selection cycle was
repeated; pools of T3 seed were analyzed for Δ -6,9 18:2. T3 half-seeds were
20 dissected and analyzed, and selected T3 plants were grown in the greenhouse to
produce T4 seed. Pools of T4 seed were analyzed for fatty acid composition.
Table 6 summarizes the results of this process for lines derived from one of the
original transgenic events, 5538-LP004-25. Levels of Δ -6,9 18:2 have thus been
maintained through 3 generations.

To maximize the amount of Δ -6,9 18:2 that could be produced, the pCGN5538 construct could be introduced into a high oleic acid variety of canola either by transformation or crossing. A high-oleic variety could be obtained by mutation, co-suppression, or antisense suppression of the Δ 12 and Δ 15 desaturases or other necessary co-factors.

5

Table 7
Fatty Acid Compositions of 20-seed Pools of pCGN5538 T2 Seeds

SPL #	event	12:0		14:0		16:0		16:1		18:0		18:1		Δ6,9		Δ6,9,12		Δ9,12,15		18:4		20:0		20:1		20:2		22:0		22:2	
		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	
31		0.02	0.06	4.07	0.07	0	59.4	5.4	10.07	15.93	1.2	0.6	0.98	1.16	0.0	0.44	0.03														
29		0.01	0.05	3.81	0.14	0	60.7	4.53	10.9	14.77	1.03	0.55	1.09	1.26	0.0	0.46	0.02														
19		0.02	0.07	4.27	0.13	0	62.9	4.17	10.03	13.14	1.02	0.59	1.18	1.25	0.0	0.53	0.02														
14		0	0	5.29	0.24	3.8	49.1	1.02	23.44	11.21	2.26	0.34	1.45	0.93	0.0	0.76	0														
22		0.02	0.05	3.87	0.09	0	64.1	2.59	12.57	11.18	1.27	0.6	1.18	1.08	0.1	0.56	0														
9		0.01	0.06	4.57	0.16	0	62.9	3.4	12.05	11.15	1.27	0.6	1.28	1.18	0.0	0.56	0.03														
25		0.01	0.06	4.17	0.14	0	62.4	2.49	14.42	11.03	1.2	0.46	1.18	1.15	0.0	0.53	0.01														
15		0.01	0.05	3.94	0.11	0	65.2	2.08	12.77	10.9	1.04	0.43	1.1	1.24	0.0	0.48	0.01														
18		0	0.06	5.34	0.29	0	58.4	1.42	18.19	10.53	1.8	0.49	1.2	1	0.0	0.58	0.02														
20		0.01	0.04	3.95	0.1	0	65.6	1.31	13.83	10.22	1.09	0.39	1.06	1.3	0.0	0.46	0.01														
7		0.02	0.07	4.04	0.11	0	62.1	0.92	18.12	8.72	1.77	0.35	1.26	1.19	0.0	0.58	0														
11		0.01	0.06	4.23	0.17	0	62.9	1.6	17.19	8.58	1.48	0.38	1.16	1.03	0.0	0.49	0.01														
27		0.02	0	3.99	0.14	0	65.3	0.64	17.85	7.89	1.36	0.31	1.08	1.21	0.0	0	0														
2		0.01	0.05	4.02	0.14	0	66.4	1.2	15.74	7.58	1.22	0.32	1.06	1.19	0.0	0.45	0														
28		0.01	0.04	3.77	0.11	0	67.5	0.79	15.56	7.58	1.12	0.28	0.97	1.23	0.0	0.44	0														
3		0.01	0.05	3.96	0.12	0	68.5	1.81	13.23	7.44	1.1	0.35	1.12	1.21	0.0	0.46	0.01														
21		0.01	0.05	3.74	0.1	0	66.9	1.16	15.9	6.97	1.35	0.28	1.15	1.27	0.0	0.52	0														
5		0.01	0.04	3.81	0.12	0	69.1	0.74	14.58	6.95	1.14	0.28	1.06	1.18	0.0	0.45	0														
6		0	0	2.84	0	3.06	62.5	1.55	18.44	6.94	1.21	0.39	1.04	1.33	0.1	0	0														
4		0.01	0.05	3.88	0.11	0	66.9	0.64	16.21	6.89	1.52	0.31	1.09	1.21	0.0	0.5	0														

SPL	5538-LP004	12:0		14:0		16:0		18:0		18:1		Δ6,9		Δ6,9,12		Δ9,12,15		20:0		20:1		20:2		22:0		22:2			
	30	0.01	0.04	3.89	0.12	0	68.6	0.72	15.58	6.47	1.17	0.23	1.03	1.07	0.0	0.46	0												
	16	0.02	0.05	3.75	0.13	0	70.4	0.91	13.56	6.39	1.13	0.28	1.04	1.2	0.0	0.44	0.01												
	26	0.01	0	3.77	0.12	0	67.6	0	21.08	3.61	1.37	0.13	0.96	1.16	0.0	0	0												
	23	0	0	4.92	0.22	0	65.2	0	22.23	3	1.79	0.11	1.28	1.11	0.0	0	0												
	24	0.01	0	3.84	0.13	0	68.4	0.36	21	2.09	1.53	0.08	1.06	1.27	0.0	0	0												
	10	0.01	0	3.74	0.11	0	70.4	0	20.82	0.65	1.3	0.03	1.05	1.18	0.0	0.46	0												
	12	0.01	0	3.83	0.12	0	69.9	0	21.61	0.34	1.34	0	1.06	1.12	0.0	0.47	0												
	17	0.01	0	0	0.13	0	72.6	0	23.03	0.24	1.51	0	1.18	1.21	0.0	0	0												
	8	0.01	0	4.54	0.2	0	64.9	0	25.65	0.22	1.94	0	1.38	1.01	0.0	0	0												
	13	0.01	0	3.99	0.16	0	65.8	0	25.9	0	1.58	0	1.17	1.16	0.0	0	0												
	LP004 control	0.01	0.04	3.46	0.09	0	69.9	0	21.95	0	1.37	0.01	0.9	1.25	0.0	0.42	0												

Table 7

STRAIN ID	T2 Pool		T3 Pool		T3 selection		T4 Pool	
	GLA	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2
5538-LP004-25	11.03	2.49						$\Delta 6,9,12$ 18:3
5538-LP004-25-3			9.1		11.92			
5538-LP004-25-3-31						13.61	7.82	11.02
5538-LP004-25-3-30						6.51	7.93	10.27
5538-LP004-25-3-29						13.35	11.23	9.42
5538-LP004-25-3-28						9.92	24.1	9.37
5538-LP004-25-3-25						5.3	30.34	7.95
5538-LP004-25-2			3.87		11.08			
5538-LP004-25-2-29						13.63	7.41	9.6
5538-LP004-25-2-27						5.02	22.04	6.95
5538-LP004-25-2-26						1.21	26.84	4.31
5538-LP004-25-2-25						5.83	34.16	8.77
5538-LP004-25-13			10.53		11.19			
5538-LP004-25-13-27						14.65	11.46	7.86
								10.49

STRAIN ID	T2 Pool		T3 Pool		T3 selection		T4 Pool	
	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2	GLA	$\Delta 6,9$ 18:2	$\Delta 6,9,12$ 18:3
5538-LP004-25-13-26					11.18	13.04	9.33	10.01
5538-LP004-25-13-25					4.18	36.78	7.2	12.22
5538-LP004-25-1			3.05	11.16				
5538-LP004-25-1-41					0	0	0.01	0.04
5538-LP004-25-1-28					3.43	19.98	4.63	6.53
5538-LP004-25-1-27					5.52	20.13	8.35	11.21
5538-LP004-25-1-26					0.1	25.16	5.52	8.59
5538-LP004-25-1-25					6.5	31.83	9.85	10.88

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or
5 patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the
10 scope of the appended claims.

Claims

What is claimed is:

5

1. A method for producing stearidonic acid in a plant seed, said method comprising:

growing a plant having integrated into its genome a first DNA construct comprising, in the 5' to 3' direction of transcription, a promoter functional in a
10 plant seed cell, a DNA sequence encoding a delta-six desaturase, and a transcription termination region functional in a plant cell, and

growing said plant under conditions whereby said delta-six desaturase is expressed.

2. The method according to claim 1 wherein said plant has a second
15 construct integrated into its genome, wherein said second construct has in the 5' to 3' direction of transcription, a promoter functional in a plant seed cell, and a DNA sequence encoding a delta 12 desaturase.

3. The method according to claim 1 wherein said plant has a second
20 construct integrated into its genome, wherein said second construct has in the 5' to 3' direction of transcription, a promoter functional in a plant seed cell, and a DNA sequence encoding a delta 15 desaturase.

4. The method of Claim 1 wherein said desaturase encoding sequence is from the genus *Mortierella*.

5. The method according to Claim 1, wherein said promoter is a napin promoter.

6. The method according to Claim 1, wherein said promoter is from the soybean β -conglycinin 7S subunit transcription initiation region.

5 7. The method according to Claim 1, wherein said method further comprises extracting oil from said plant seed.

8. The method of Claim 7, wherein said oil comprises about 5 weight percent or greater stearidonic acid.

9. The method of Claim 7, wherein said oil comprises about 10 weight
10 percent or greater stearidonic acid.

10. The method of Claim 7, wherein said oil comprises about 15 weight percent or greater stearidonic acid.

11. The method of Claim 7, wherein said oil comprises about 20 weight percent or greater stearidonic acid.

12. The method of Claim 7, wherein said oil comprises about 25
15 weight percent or greater stearidonic acid.

13. A seed comprising about 5 weight percent or greater stearidonic acid as a component of total fatty acids found in the seed oil.

14. The seed according to Claim 13 comprising about 10 weight percent or greater stearidonic acid as a component of said seed oil.

15. The seed according to Claim 13 comprising about 15 weight percent or greater stearidonic acid as a component of said seed oil.

5 16. The seed according to Claim 13 comprising about 20 weight percent or greater stearidonic acid as a component of said seed oil.

17. The seed according to Claim 13 comprising about 25 weight percent or greater stearidonic acid as a component of said seed oil.

18. Seed oil obtained from seed of Claim 13.

Pathways of PUFA Metabolism





FIGURE 1

Pathways of PUFA Metabolism

$\Delta 5$ Desaturation



2/2

FIGURE 2