



US 20080057067A1

(19) **United States**

(12) **Patent Application Publication**

**Ruelle**

(10) **Pub. No.: US 2008/0057067 A1**

(43) **Pub. Date: Mar. 6, 2008**

(54) **POLYPEPTIDES AND POLYNUCLEOTIDES  
"BASB040" FROM NEISSERIA  
MENINGITIDIS AND VACCINE  
COMPRISING SAID POLYPEPTIDES AND  
POLYNUCLEOTIDES**

(30) **Foreign Application Priority Data**

Dec. 7, 1998 (GB)..... 9826886.5

**Publication Classification**

(75) Inventor: **Jean-Louis Ruelle, Rixensart (BE)**

(51) **Int. Cl.**

Correspondence Address:  
**RATNER & PRESTIA- SB DIVISION  
ONE WESTLAKES  
SUITE 301  
BERWYN, PA 19482 (US)**

**A61K 39/395 (2006.01)  
A61K 39/095 (2006.01)  
A61P 31/04 (2006.01)  
C07H 21/00 (2006.01)  
C07K 14/22 (2006.01)  
C07K 16/12 (2006.01)  
C12N 1/00 (2006.01)  
C12N 15/63 (2006.01)  
C12P 19/34 (2006.01)  
C12Q 1/04 (2006.01)**

(73) Assignee: **GlaxoSmithKline Biologicals S.A., Rixensart (BE)**

(21) Appl. No.: **11/770,421**

(52) **U.S. Cl. .... 424/139.1; 424/250.1; 435/243;  
435/320.1; 435/34; 435/69.1;  
435/91.1; 530/350; 530/387.9;  
536/23.7**

(22) Filed: **Jun. 28, 2007**

**Related U.S. Application Data**

(63) Continuation of application No. 10/833,884, filed on Apr. 28, 2004, now abandoned, which is a continuation of application No. 09/857,669, filed on Sep. 21, 2001, now Pat. No. 6,770,284, filed as 371 of international application No. PCT/EP99/09560, filed on Dec. 2, 1999.

(57)

**ABSTRACT**

The invention provides BASB040 polypeptides and polynucleotides from *Neisseria meningitidis* encoding BASB040 polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are diagnostic, prophylactic and therapeutic uses.

Identity to SeqID No:1 is indicated by a dot.

```

                *           20           *
Seqid1 : ATGATGATCAAACCGACCGCCTGCTCCTG : 30
Seqid3 : ..... : 30
Seqid5 : ..... : -

                40           *           60
Seqid1 : CCGGCTTTATTTTCTTCCGCACGCATAC : 60
Seqid3 : ..... : 60
Seqid5 : ..... : -

                *           80           *
Seqid1 : GCGCCTGCCGCCGACCTTCCGAAAACAAG : 90
Seqid3 : ..... : 90
Seqid5 : ..... : 24

                100           *           120
Seqid1 : GCGGCGGGTTTCGCATTGTCAAAAACAAA : 120
Seqid3 : ..... : 120
Seqid5 : ..... : 54

                *           140           *
Seqid1 : AGCCCGACACCGAATCAGTTAAATTAAAA : 150
Seqid3 : ..... : 150
Seqid5 : .....C..... : 84

                160           *           180
Seqid1 : CCCAAATCCCCGTCCGCATCGACACGCAG : 180
Seqid3 : ..... : 180
Seqid5 : .....T..... : 114

```

Figure 1A: Identity to SeqID No:1 is indicated by a dot.

```

                *           20           *
Seqid1 : ATGATGATCAAACCGACCGCCCTGCTCCTG : 30
Seqid3 : ..... : 30
Seqid5 : : -

```

```

                40           *           60
Seqid1 : CCGGCTTTATTTTCTTTCCGCACGCATAC : 60
Seqid3 : ..... : 60
Seqid5 : : -

```

```

                *           80           *
Seqid1 : GCGCCTGCCGCCGACCTTCCGAAAACAAG : 90
Seqid3 : ..... : 90
Seqid5 : ..... : 24

```

```

                100           *           120
Seqid1 : GCGGCGGGTTTCGCATTGTTCAAAAACAAA : 120
Seqid3 : ..... : 120
Seqid5 : ..... : 54

```

```

                *           140           *
Seqid1 : AGCCCCGACACCGAATCAGTTAAATTAAAA : 150
Seqid3 : ..... : 150
Seqid5 : .....C..... : 84

```

```

                160           *           180
Seqid1 : CCCAAATTCCCCGTCCGCATCGACACGCAG : 180
Seqid3 : ..... : 180
Seqid5 : .....T..... : 114

```

Figure 1B: Identity to SeqID No:1 is indicated by a dot.

\* 200 \*

Seqid1 : GATAGTCAAATCAAAGATATGGTCGAAGAA : 210  
Seqid3 : ..... : 210  
Seqid5 : ..C..... : 144

220 \* 240

Seqid1 : CACCTGCCGCTCATCACGCAGCAGCAGGAA : 240  
Seqid3 : ..... : 240  
Seqid5 : ..... : 174

\* 260 \*

Seqid1 : GAAGTATTGGACAAGGAACAGACGGGCTTC : 270  
Seqid3 : ..... : 270  
Seqid5 : ..... : 204

280 \* 300

Seqid1 : CTCGCCGAAGAAGCACCGGACAACGTTAAA : 300  
Seqid3 : ..... : 300  
Seqid5 : .....G..... : 234

\* 320 \*

Seqid1 : ACCATGCTCCGCAGCAAAGGCTATTTTCAGC : 330  
Seqid3 : ..... : 330  
Seqid5 : ..G..... : 264

340 \* 360

Seqid1 : AGCAAAGTCAGCCTGACGGAAAAAGACGGA : 360  
Seqid3 : ..... : 360  
Seqid5 : ..... : 294

\* 380 \*

Figure 1C: Identity to SeqID No:1 is indicated by a dot.

Seqid1 : GCTTATACGGTACACATCACACCGGGCCCG : 390  
Seqid3 : ..... : 390  
Seqid5 : ..... : 324

400 \* 420  
Seqid1 : CGCACCAAATCGCCAACGTCGGTGTCGCC : 420  
Seqid3 : ..... : 420  
Seqid5 : .....C..... : 354

\* 440 \*  
Seqid1 : ATCCTCGGCGACATCCTTTCAGACGGCAAC : 450  
Seqid3 : ..... : 450  
Seqid5 : ..... : 384

460 \* 480  
Seqid1 : CTCGCCGAATACTACCGCAACGCGCTGGAA : 480  
Seqid3 : ..... : 480  
Seqid5 : ..... : 414

\* 500 \*  
Seqid1 : AACTGGCAGCAGCCGGTAGGCAGTGATTTC : 510  
Seqid3 : ..... : 510  
Seqid5 : .....C..... : 444

520 \* 540  
Seqid1 : GATCAGGACAGTTGGGAAAACAGCAAAC : 540  
Seqid3 : ..... : 540  
Seqid5 : ..... : 474

\* 560 \*  
Seqid1 : TCCGTCCTCGGCGCGGTAACGCGAAAAGGC : 570  
Seqid3 : .....C..... : 570

Figure 1D: Identity to SeqID No:1 is indicated by a dot.

Seqid5 : .....C....C. : 504

580 \* 600  
 Seqid1 : TACCCGCTTGCCAAGCTCGGCAACACCCGG : 600  
 Seqid3 : ..... : 600  
 Seqid5 : .....T..G.A. : 534

\* 620 \*  
 Seqid1 : GCGGCCGTCAACCCCGATACCGCCACCGTC : 630  
 Seqid3 : ..... : 630  
 Seqid5 : .....C. : 564

640 \* 660  
 Seqid1 : GATTTGAACGTTCGTGGACAGCGGCCGC : 660  
 Seqid3 : ..... : 660  
 Seqid5 : ..... : 594

\* 680 \*  
 Seqid1 : CCCATCGCCTTCGGCGACTTTGAAATCACC : 690  
 Seqid3 : ..... : 690  
 Seqid5 : ..... : 624

700 \* 720  
 Seqid1 : GGCACACAGCGTTACCCCGAACAAATCGTC : 720  
 Seqid3 : ..... : 720  
 Seqid5 : ..... : 654

\* 740 \*  
 Seqid1 : TCCGGCCTGGCGCGCTTCCAACCGGGCAGC : 750  
 Seqid3 : ..... : 750  
 Seqid5 : .....T.....T.....G..C..T.T. : 684

Figure 1E: Identity to SeqID No:1 is indicated by a dot.

```

              760                *                780
Seqid1 : CCCTACGACCTCGACCTGCTGCTCGACTTC : 780
Seqid3 : ..... : 780
Seqid5 : ..G..... : 714

```

```

              *                800                *
Seqid1 : CAACAGGCACTCGAACAAAACGGGCATTAT : 810
Seqid3 : ..... : 810
Seqid5 : .....G..... : 744

```

```

              820                *                840
Seqid1 : TCCGGCGCGTCCGTACAAGCCGACTTCGAC : 840
Seqid3 : ..... : 840
Seqid5 : ..... : 774

```

```

              *                860                *
Seqid1 : CGTCTCCAAGGCGACCGCGTCCCCGTCAA : 870
Seqid3 : ..... : 870
Seqid5 : ..C..... : 804

```

```

              880                *                900
Seqid1 : GTCAGCGTAACCGAGGTCAAACGCCACAAG : 900
Seqid3 : ..... : 900
Seqid5 : .....A : 834

```

```

              *                920                *
Seqid1 : CTCGAAACCGGCATCCGCCTCGATTCGGAA : 930
Seqid3 : ..... : 930
Seqid5 : ..... : 864

```

```

              940                *                960
Seqid1 : TACGGTTTGGGCGGCAAAATCGCCTACGAC : 960

```

Figure 1F: Identity to SeqID No:1 is indicated by a dot.

Seqid3 : ..... : 960  
Seqid5 : ..... : 894

\* 980 \*  
Seqid1 : TATTACAACCTCTTCAACAAAGGCTATATC : 990  
Seqid3 : ..... : 990  
Seqid5 : ..... : 924

1000 \* 1020  
Seqid1 : GGCTCGGTCGTCTGGGATATGGACAAATAC : 1020  
Seqid3 : ..... : 1020  
Seqid5 : ..T..... : 954

\* 1040 \*  
Seqid1 : GAAACCACGCTTGCCGCCGGCATCAGCCAG : 1050  
Seqid3 : ..... : 1050  
Seqid5 : ..... : 984

1060 \* 1080  
Seqid1 : CCGCGCAACTATCGGGGCAACTACTGGACA : 1080  
Seqid3 : ..... : 1080  
Seqid5 : ..... : 1014

\* 1100 \*  
Seqid1 : AGCAACGTTTCCTACAACCGTTCGACCACC : 1110  
Seqid3 : ..... : 1110  
Seqid5 : ..... : 1044

1120 \* 1140  
Seqid1 : CAAAACCTCGAAAAACGCGCCTTCTCCGGC : 1140  
Seqid3 : ..... : 1140  
Seqid5 : ..... : 1074

Figure 1G: Identity to SeqID No:1 is indicated by a dot.

\*                    1160                    \*

Seqid1 : GGCATCTGGTATGTGCGCGACCGCGCGGGC : 1170  
 Seqid3 : ..... : 1170  
 Seqid5 : ...G..... : 1104

1180                    \*                    1200

Seqid1 : ATCGATGCCAGGCTGGGGGCAGAGTTTCTC : 1200  
 Seqid3 : ..... : 1200  
 Seqid5 : .....G..A..... : 1134

\*                    1220                    \*

Seqid1 : GCAGAAGGCCGGAAAATCCCCGGCTCGGAT : 1230  
 Seqid3 : ..... : 1230  
 Seqid5 : .....C. : 1164

1240                    \*                    1260

Seqid1 : ATCGATTTGGGCAACAGCCACGCCACGATG : 1260  
 Seqid3 : ..... : 1260  
 Seqid5 : G..... : 1194

\*                    1280                    \*

Seqid1 : CTGACCGCCTCTTGGAACGCCAGCTGCTC : 1290  
 Seqid3 : ..... : 1290  
 Seqid5 : ..... : 1224

1300                    \*                    1320

Seqid1 : AACAACTGTGCTGCATCCCGAAAACGGCCAT : 1320  
 Seqid3 : ..... : 1320  
 Seqid5 : ..... : 1254

\*                    1340                    \*



Figure 1H: Identity to SeqID No:1 is indicated by a dot.

Seqid1 : TACCTCGACGGCAAAATCGGTACGACTTTG : 1350  
Seqid3 : ..... : 1350  
Seqid5 : ..... : 1284

1360 \* 1380  
Seqid1 : GGCACATTCCTGTCCTCCACCGCGCTGATC : 1380  
Seqid3 : ..... : 1380  
Seqid5 : ..... : 1314

\* 1400 \*  
Seqid1 : CGCACCTCTGCCCGTGCAGGTTATTTCTTC : 1410  
Seqid3 : ..... : 1410  
Seqid5 : ..... : 1344

1420 \* 1440  
Seqid1 : ACGCCCGAAAACAAAAACTCGGCACGTTC : 1440  
Seqid3 : ..... : 1440  
Seqid5 : ..... : 1374

\* 1460 \*  
Seqid1 : ATCATA CGCGGACAAGCGGGTTACACCGTT : 1470  
Seqid3 : ..... : 1470  
Seqid5 : ..... : 1404

1480 \* 1500  
Seqid1 : GCCCGCGACAATGCCGACGTTCTTCAGGG : 1500  
Seqid3 : ..... : 1500  
Seqid5 : ..... : 1434

\* 1520 \*  
Seqid1 : CTGATGTTCCGCAGCGGCGGCGGTCTTCC : 1530  
Seqid3 : ..... : 1530

Figure 11: Identity to SeqID No:1 is indicated by a dot.

```
Seqid5 : ..... : 1464

                1540      *      1560
Seqid1 : GTGCGCGGTTACGAACTCGACAGCATCGGA : 1560
Seqid3 : ..... : 1560
Seqid5 : ..... : 1494

                *      1580      *
Seqid1 : CTTGCCGGCCCGAACGGATCGGTCCTGCCC : 1590
Seqid3 : ..... : 1590
Seqid5 : ..... : 1524

                1600      *      1620
Seqid1 : GAACGCGCCCTCCTGGTGGGCAGCCTGGAA : 1620
Seqid3 : ..... : 1620
Seqid5 : ..... : 1554

                *      1640      *
Seqid1 : TACCAACTGCCGTTTACGCGCACCCCTTTCC : 1650
Seqid3 : ..... : 1650
Seqid5 : ..... : 1584

                1660      *      1680
Seqid1 : GGCGCGGTGTTCCACGATATGGGCGATGCC : 1680
Seqid3 : ..... : 1680
Seqid5 : ..... : 1614

                *      1700      *
Seqid1 : GCCGCCAATTTCAAACGTATGAAGCTGAAA : 1710
Seqid3 : ..... : 1710
Seqid5 : ..... : 1644
```

**Figure 1J: Identity to SeqID No:1 is indicated by a dot.**

```

                1720                *                1740
Seqid1 : CACGGTTCGGGACTGGGCGTGCGCTGGTTC : 1740
Seqid3 : ..... : 1740
Seqid5 : ..... : 1674
    
```

```

                *                1760                *
Seqid1 : AGCCCGCTTGCGCCGTTTTTCCTTCGACATC : 1770
Seqid3 : ..... : 1770
Seqid5 : ..... : 1704
    
```

```

                1780                *                1800
Seqid1 : GCCTACGGGCACAGCGATAAGAAAATCCGC : 1800
Seqid3 : ..... : 1800
Seqid5 : ..... : 1734
    
```

```

                *                1820                *
Seqid1 : TGGCACATCAGCTTGGGAACACGCTTCTAA : 1830
Seqid3 : ..... : 1830
Seqid5 : ..... : 1764
    
```

Figure 2A : Identity to SeqID No:2 is indicated by a dot.

	*	20	*		
Seqid2 :	MMIKPTALLLPALFFFP	HAYAPAADLSENK	:	30	
Seqid4 :	.....		:	30	
Seqid6 :		.....	:	8	
	40	*	60		
Seqid2 :	AAGFALFKNKSPDTE	SVKCLKPKFPVRIDTQ	:	60	
Seqid4 :	.....		:	60	
Seqid6 :	.....	L.....	:	38	
	*	80	*		
Seqid2 :	DSEIKDMVEEHLPLIT	QQQEEVLDKEQTGF	:	90	
Seqid4 :	.....		:	90	
Seqid6 :	.....		:	68	
	100	*	120		
Seqid2 :	LAEEAPDNVKTMLRS	KGYFSSKVS	LTEKDG	:	120
Seqid4 :	.....			:	120
Seqid6 :	.....			:	98

Figure 2B : Identity to SeqID No:2 is indicated by a dot.

\* 140 \*

Seqid2 : AYT VHITPGPRTKIANVGVAILGDILSDGN : 150  
Seqid4 : ..... : 150  
Seqid6 : ..... : 128

160 \* 180

Seqid2 : LAEYYRNALENWQQPVGSDFDQDSWENSKT : 180  
Seqid4 : ..... : 180  
Seqid6 : ..... : 158

\* 200 \*

Seqid2 : SVLGAVTRKGYPLAKLGNTRAAVNPDTATV : 210  
Seqid4 : ..... : 210  
Seqid6 : .....A.....Q.....A : 188

220 \* 240

Seqid2 : DLNVVVDSGRPIAFGDFEITGTQRYPEQIV : 240  
Seqid4 : ..... : 240  
Seqid6 : ..... : 218

\* 260 \*

Figure 2C : Identity to SeqID No:2 is indicated by a dot.

Seqid2 : SGLARFQPGTPYDLDLLLDFQQALEQNGHY : 270  
Seqid4 : ..... : 270  
Seqid6 : .....M..... : 248

280 \* 300

Seqid2 : SGASVQADFDR LQGDRVPVKVSVTEVKRHK : 300  
Seqid4 : ..... : 300  
Seqid6 : ..... : 278

\* 320 \*

Seqid2 : LETGIRLDSEYGLGGKIA YDYYNLFNKG YI : 330  
Seqid4 : ..... : 330  
Seqid6 : ..... : 308

340 \* 360

Seqid2 : GSVVWDMDKYETTLAAGISQPRNYRGNYWT : 360  
Seqid4 : ..... : 360  
Seqid6 : ..... : 338

\* 380 \*

Seqid2 : SNVSYNRSTTQNLEKRAFSGGIWYVRDRAG : 390  
Seqid4 : ..... : 390

Figure 2D : Identity to SeqID No:2 is indicated by a dot.

Seqid6 : ..... : 368

400 \* 420

Seqid2 : IDARLGAEFLAEGRKIPGSDIDLGN SHATM : 420

Seqid4 : ..... : 420

Seqid6 : .....A..... : 398

\* 440 \*

Seqid2 : LTASWKRQLLNNVLHPENGHYLDGKIGTTL : 450

Seqid4 : ..... : 450

Seqid6 : ..... : 428

460 \* 480

Seqid2 : GTFLSSTALIRTSARAGYFFTPENKKLGTF : 480

Seqid4 : ..... : 480

Seqid6 : ..... : 458

\* 500 \*

Seqid2 : IIRGQAGYTVARDNADVPSGLMFRSGGASS : 510

Seqid4 : ..... : 510

Seqid6 : ..... : 488

Figure 2E : Identity to SeqID No:2 is indicated by a dot.

520 \* 540  
Seqid2 : VRGYELDSIGLAGPNGSVLPERALLVGSLE : 540  
Seqid4 : ..... : 540  
Seqid6 : ..... : 518

\* 560 \*  
Seqid2 : YQLPFTRTLSGAVFHDMGDAAANFKRMKLLK : 570  
Seqid4 : ..... : 570  
Seqid6 : ..... : 548

580 \* 600  
Seqid2 : HGSGLGVRWFSPLAPFSFDIAYGHSDKKIR : 600  
Seqid4 : ..... : 600  
Seqid6 : ..... : 578

Seqid2 : WHISLGTRF : 609  
Seqid4 : ..... : 609  
Seqid6 : ..... : 587



**POLYPEPTIDES AND POLYNUCLEOTIDES  
"BASB040" FROM NEISSERIA MENINGITIDIS  
AND VACCINE COMPRISING SAID  
POLYPEPTIDES AND POLYNUCLEOTIDES**

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

[0001] This application is a continuation of U.S. patent application Ser. No. 10/833,884, filed Apr. 28, 2004, which is a continuation of U.S. patent application Ser. No. 09/857,669, filed Sep. 21, 2001, which is the National Stage Application of International Application No. PCT/EP99/09560, filed Dec. 2, 1999, which claims the benefit of Great Britain Patent Application No. 9826886.5 filed Dec. 7, 1998. The disclosure of these applications are incorporated herein by reference in their entirety.

**FIELD OF THE INVENTION**

[0002] This invention relates to polynucleotides, (herein referred to as "BASB040 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB040" or "BASB040 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

**BACKGROUND OF THE INVENTION**

[0003] *Neisseria meningitidis* (meningococcus) is a Gram-negative bacterium frequently isolated from the human upper respiratory tract. It occasionally causes invasive bacterial diseases such as bacteremia and meningitis. The incidence of meningococcal disease shows geographical seasonal and annual differences (Schwartz, B., Moore, P. S., Broome, C. V.; Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Most disease in temperate countries is due to strains of serogroup B and varies in incidence from 1-10/100,000/year total population sometimes reaching higher values (Kaczmarek, E. B. (1997), Commun. Dis. Rep. Rev. 7: R55-9, 1995; Scholten, R. J. P. M., Bijlmer, H. A., Poolman, J. T. et al. Clin. Infect. Dis. 16: 237-246, 1993; Cruz, C., Pavez, G., Aguilar, E., et al. Epidemiol. Infect. 105: 119-126, 1990).

[0004] Epidemics dominated by serogroup A meningococci, mostly in central Africa, are encountered, sometimes reaching levels up to 1000/100,000/year (Schwartz, B., Moore, P. S., Broome, C. V. Clin. Microbiol. Rev. 2 (Supplement), S18-S24, 1989). Nearly all cases as a whole of meningococcal disease are caused by serogroup A, B, C, W-135 and Y meningococci and a tetravalent A, C, W-135, Y polysaccharide vaccine is available (Armand, J., Arminjon, F., Mynard, M. C., Lafaix, C., J. Biol. Stand. 10: 335-339, 1982).

[0005] The polysaccharide vaccines are currently being improved by way of chemical conjugating them to carrier proteins (Lieberman, J. M., Chiu, S. S., Wong, V. K., et al. JAMA 275: 1499-1503, 1996).

[0006] A serogroup B vaccine is not available, since the B capsular polysaccharide was found to be nonimmunogenic, most likely because it shares structural similarity to host

components (Wyle, F. A., Artenstein, M. S., Brandt, M. L. et al. J. Infect. Dis. 126: 514-522, 1972; Finne, J. M., Leinonen, M., Makelai, P. M. Lancet ii.: 355-357, 1983).

[0007] For many years efforts have been initiated and carried out to develop meningococcal outer membrane based vaccines (de Moraes, J. C., Perkins, B., Camargo, M. C. et al. Lancet 340: 1074-1078, 1992; Bjune, G., Hoiby, E. A. Gronnesby, J. K. et al. 338: 1093-1096, 1991). Such vaccines have demonstrated efficacies from 57%-85% in older children (>4 years) and adolescents.

[0008] Many bacterial outer membrane components are present in these vaccines, such as PorA, PorB, Rmp, Opc, Opa, FrpB and the contribution of these components to the observed protection still needs further definition. Other bacterial outer membrane components have been defined by using animal or human antibodies to be potentially relevant to the induction of protective immunity, such as TbpB and NspA (Martin, D., Cadieux, N., Hamel, J., Brodeux, B. R., J. Exp. Med. 185: 1173-1183, 1997; Lissolo, L., Maitre-Wilmotte, C., Dumas, p. et al., Inf. Immun. 63: 884-890, 1995). The mechanisms of protective immunity will involve antibody mediated bactericidal activity and opsonophagocytosis.

[0009] A bacteremia animal model has been used to combine all antibody mediated mechanisms (Saukkonen, K., Leinonen, M., Abdillahi, H. Poolman, J. T. Vaccine 7: 325-328, 1989). It is generally accepted that the late complement component mediated bactericidal mechanism is crucial for immunity against meningococcal disease (Ross, S. C., Rosenthal P. J., Berberic, H. M., Densen, P. J. Infect. Dis. 155: 1266-1275, 1987).

[0010] The frequency of *Neisseria meningitidis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Neisseria meningitidis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

**SUMMARY OF THE INVENTION**

[0011] The present invention relates to BASB040, in particular BASB040 polypeptides and BASB040 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB040 polynucleotides or polypeptides.

[0012] Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

**BRIEF DESCRIPTION OF THE DRAWINGS**

[0013] FIG. 1 Alignment of SEQ ID NOs:1, 3, and 5.

[0014] FIG. 2 Alignment of SEQ ID NOs:2, 4, and 6.

## DESCRIPTION OF THE INVENTION

[0015] The invention relates to BASB040 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB040 of *Neisseria meningitidis*, which is related by amino acid sequence homology to *Neisseria meningitidis* D15 outer membrane protein. The invention relates especially to BASB040 having the nucleotide and amino acid sequences set out in SEQ ID NO:1, 3, 5 and SEQ ID NO:2, 4, 6 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

## Polypeptides

[0016] In one aspect of the invention there are provided polypeptides of *Neisseria meningitidis* referred to herein as "BASB040" and "BASB040 polypeptides" as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

[0017] The present invention further provides for:

[0018] (a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2, 4, 6;

[0019] (b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1, 3, 5 over the entire length of SEQ ID NO:1, 3, 5 respectively; or

[0020] (c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity, to the amino acid sequence of SEQ ID NO:2, 4, 6;

[0021] The BASB040 polypeptides provided in SEQ ID NO:2, 4, 6 are the BASB040 polypeptides from *Neisseria meningitidis* strains ATCC13090 and H44/76.

[0022] The invention also provides an immunogenic fragment of a BASB040 polypeptide, that is, a contiguous portion of the BASB040 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2, 4, 6. That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB040 polypeptide. Such an immunogenic fragment may include, for example, the BASB040 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB040 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most

preferably at least 97-99% identity, to that of SEQ ID NO:2, 4, 6 over the entire length of SEQ ID NO:2, 4, 6. A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB040 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

[0023] Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2, 4, 6 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

[0024] Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2, 4, 6, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2, 4, 6.

[0025] Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments may be employed as intermediates for producing the full-length polypeptides of the invention.

[0026] Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

[0027] The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

[0028] In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

[0029] Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0030] The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognized by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

[0031] Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesizes an N-acetyl-L-alanine amidase, amidase LytA, (coded by the *lytA* gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LytA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188-305.

[0032] The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

[0033] Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[0034] It is most preferred that a polypeptide of the invention is derived from *Neisseria meningitidis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

Polynucleotides

[0035] It is an object of the invention to provide polynucleotides that encode BASB040 polypeptides, particularly polynucleotides that encode the polypeptide herein designated BASB040.

[0036] In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB040 polypeptides comprising a sequence set out in SEQ ID NO:1, 3, 5 which includes a full length gene, or a variant thereof.

[0037] The BASB040 polynucleotides provided in SEQ ID NO:1, 3 and 5 are the BASB040 polynucleotides from *Neisseria meningitidis* strains ATCC13090 and H44/76. As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB040 polypeptides and polynucleotides, particularly *Neisseria meningitidis* BASB040 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

[0038] Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a BASB040 polypeptide having a deduced amino acid sequence of SEQ ID NO:2, 4, 6 and polynucleotides closely related thereto and variants thereof.

[0039] In another particularly preferred embodiment of the invention there is a BASB040 polypeptide from *Neisseria meningitidis* comprising or consisting of an amino acid sequence of SEQ ID NO:2, 4, 6 or a variant thereof.

[0040] Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1, 3, 5 a polynucleotide of the invention encoding BASB040 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Neisseria meningitidis* cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1, 3, 5, typically a library of clones of chromosomal DNA of *Neisseria meningitidis* in *E. coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E. F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1, 3, 5 was discovered in a DNA library derived from *Neisseria meningitidis*.

[0041] Moreover, each DNA sequence set out in SEQ ID NO:1, 3, 5 contains an open reading frame encoding a

protein having about the number of amino acid residues set forth in SEQ ID NO:2, 4, 6 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

[0042] The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1828 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

[0043] The polynucleotide of SEQ ID NO:3, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1828 of SEQ ID NO:3, encodes the polypeptide of SEQ ID NO:4.

[0044] The polynucleotide of SEQ ID NO:5, between the first codon at nucleotide number 1 and the stop codon which begins at nucleotide number 1762 of SEQ ID NO:5, encodes the polypeptide of SEQ ID NO:6.

[0045] In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

[0046] (a) a polynucleotide sequence which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1, 3, 5 over the entire length of SEQ ID NO:1, 3, 5 respectively; or

[0047] (b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2, 4, 6 over the entire length of SEQ ID NO:2, 4, 6 respectively.

[0048] A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Neisseria meningitidis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions (for example, using a temperature in the range of 45-65° C. and an SDS concentration from 0.1-1%) with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO: 1, 3, 5 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

[0049] The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in SEQ ID NO: 1, 3, 5. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or pre-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the

marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz et al., *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson et al., *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

[0050] The nucleotide sequence encoding BASB040 polypeptide of SEQ ID NO:2, 4, 6 may be identical to the polypeptide encoding sequence contained in nucleotides 1 to 1827 of SEQ ID NO:1, or the polypeptide encoding sequence contained in nucleotides 1 to 1827 of SEQ ID NO:3, or the polypeptide encoding sequence contained in nucleotides 1 to 1761 of SEQ ID NO:5, respectively. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2, 4, 6.

[0051] The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Neisseria meningitidis* BASB040 having an amino acid sequence set out in SEQ ID NO:2, 4, 6. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

[0052] The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2, 4, 6. Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

[0053] Further particularly preferred embodiments are polynucleotides encoding BASB040 variants, that have the amino acid sequence of BASB040 polypeptide of SEQ ID NO:2, 4, 6 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB040 polypeptide.

[0054] Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB040 polypeptide having an amino acid sequence set out in SEQ ID NO:2, 4, 6, and polynucleotides that are complementary to such polynucleotides. In this regard, polynucleotides at least 90% identical over their entire length to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

[0055] Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1, 3, 5.

[0056] In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB040 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1, 3, 5.

[0057] The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42° C. in a solution comprising: 50% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5×Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1×SSC at about 65° C. Hybridization and wash conditions are well known and exemplified in Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

[0058] The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1, 3, 5 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1, 3, 5 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

[0059] As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB040 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB040 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

[0060] A coding region of a BASB040 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1, 3, 5 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

[0061] There are several methods available and well known to those skilled in the art to obtain full-length DNAs,

or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, et al., *PNAS USA* 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

[0062] The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

[0063] The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NOS:1-6 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

[0064] The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case in vivo, the additional amino acids may be processed away from the mature protein by cellular enzymes.

[0065] For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

[0066] A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

[0067] In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in

describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

[0068] In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

[0069] In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

[0070] The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* (1992) 1: 363, Manthorpe et al., *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol. Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* (1989) 243: 375), particle bombardment (Tang et al., *Nature* (1992) 356: 152, Eisenbraun et al., *DNA Cell Biol* (1993) 12: 791) and in vivo infection using cloned retroviral vectors (Seeger et al., *PNAS USA* (1984) 81: 5849).

#### Vectors, Host Cells, Expression Systems

[0071] The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

[0072] Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

[0073] For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Labora-

tory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

[0074] Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Moraxella catarrhalis*, *Haemophilus influenzae* and *Neisseria meningitidis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

[0075] A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picomaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, (supra).

[0076] In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0077] Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

[0078] The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and in vivo infection with this live vector will lead to in vivo expression of the

antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), *Listeria*, *Salmonella*, *Shigella*, *Neisseria*, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

#### Diagnostic, Prognostic, Serotyping and Mutation Assays

[0079] This invention is also related to the use of BASB040 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB040 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB040 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

[0080] Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB040 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers et al., *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, Vi and S1 protection assay or a chemical cleavage method. See, for example, Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

[0081] In another embodiment, an array of oligonucleotides probes comprising BASB040 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype,

taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee et al., *Science*, 274: 610 (1996)).

[0082] Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1, 3, 5, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2, 4, 6 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2, 4, 6.

[0083] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, among others.

[0084] This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferable, SEQ ID NO:1, 3, 5, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

[0085] Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB040 polypeptide can be used to identify and analyze mutations.

[0086] The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying BASB040 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

[0087] The invention further provides a process for diagnosing disease, preferably bacterial infections, more prefer-

ably infections caused by *Neisseria meningitidis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1, 3, 5. Increased or decreased expression of a BASB040 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

[0088] In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB040 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB040 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

[0089] The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probe obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Neisseria meningitidis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1, 3, 5 are preferred. Also preferred is a grid comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2, 4, 6.

#### Antibodies

[0090] The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

[0091] In certain preferred embodiments of the invention there are provided antibodies against BASB040 polypeptides or polynucleotides.

[0092] Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor et al., *Immunology Today* 4: 72 (1983); Cole et al., pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

[0093] Techniques for the production of single chain antibodies (U.S. Pat. No. 4,946,778) can be adapted to produce

single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

[0094] Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB040 or from naive libraries (McCafferty, et al., (1990), *Nature* 348, 552-554; Marks, et al., (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson et al., (1991) *Nature* 352: 628).

[0095] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

[0096] Thus, among others, antibodies against BASB040-polypeptide or BASB040-polynucleotide may be employed to treat infections, particularly bacterial infections.

[0097] Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

[0098] Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

#### Antagonists and Agonists—Assays and Molecules

[0099] Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, e.g., Coligan et al., *Current Protocols in Immunology* 1 (2): Chapter 5 (1991).

[0100] The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in



screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB040 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB040 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB040 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and/or functionally related polypeptides (see D. Bennett et al., *J Mol Recognition*, 8: 52-58 (1995); and K. Johanson et al., *J Biol Chem*, 270(16):9459-9471 (1995)).

[0101] The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

[0102] The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB040 polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB040 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a BASB040 agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the BASB040 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, i.e., without inducing the effects of BASB040 polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in BASB040 polynucleotide or polypeptide activity, and binding assays known in the art.

[0103] Another example of an assay for BASB040 agonists is a competitive assay that combines BASB040 and a potential agonist with BASB040-binding molecules, recombinant BASB040 binding molecules, natural substrates or

ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. BASB040 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB040 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

[0104] Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB040-induced activities, thereby preventing the action or expression of BASB040 polypeptides and/or polynucleotides by excluding BASB040 polypeptides and/or polynucleotides from binding.

[0105] Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, Fla. (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB040.

[0106] In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0107] Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

[0108] The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to

interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB040 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

[0109] In accordance with yet another aspect of the invention, there are provided BASB040 agonists and antagonists, preferably bacteriostatic or bactericidal agonists and antagonists.

[0110] The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

[0111] In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognized by antibodies which recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

[0112] Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the purposes of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier protein. Thereby presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a beneficial derivative, for example to enhance stability of the peptide.

[0113] Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

#### Vaccines

[0114] Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB040 polynucleotide

and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Neisseria meningitidis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB040 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB040 polynucleotide and/or polypeptide, or a fragment or a variant thereof in vivo in order to induce an immunological response, such as, to produce antibody and/or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

[0115] A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB040 polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant BASB040 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB040 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

[0116] A BASB040 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-5-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

[0117] In a vaccine composition according to the invention, a BASB040 polypeptide and/or polynucleotide, or a fragment, or a mimotope, or a variant thereof may be present in a vector, such as the live recombinant vectors described above for example live bacterial vectors.

[0118] Also suitable are non-live vectors for the BASB040 polypeptide, for example bacterial outer-membrane vesicles

or "blebs". OM blebs are derived from the outer membrane of the two-layer membrane of Gram-negative bacteria and have been documented in many Gram-negative bacteria (Zhou, L. et al. 1998. FEMS Microbiol. Lett. 163: 223-228) including *C. trachomatis* and *C. psittaci*. A non-exhaustive list of bacterial pathogens reported to produce blebs also includes: *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella melitensis*, *Brucella ovis*, *Escherichia coli*, *Haemophilus influenza*, *Legionella pneumophila*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*.

[0119] Blebs have the advantage of providing outer-membrane proteins in their native conformation and are thus particularly useful for vaccines. Blebs can also be improved for vaccine use by engineering the bacterium so as to modify the expression of one or more molecules at the outer membrane. Thus for example the expression of a desired immunogenic protein at the outer membrane, such as the BASB040 polypeptide, can be introduced or upregulated (e.g. by altering the promoter). Instead or in addition, the expression of outer-membrane molecules which are either not relevant (e.g. unprotective antigens or immunodominant but variable proteins) or detrimental (e.g. toxic molecules such as LPS, or potential inducers of an autoimmune response) can be down-regulated. These approaches are discussed in more detail below.

[0120] The non-coding flanking regions of the BASB040 gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the shine-dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences. A region directly upstream of the BASB040 gene is given in Sequence ID No: 11 This sequence is a further aspect of the invention.

[0121] This sequence information allows the modulation of the natural expression of the BASB040 gene. The upregulation of the gene expression may be accomplished by altering the promoter, the shine-dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modification. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or it may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation include, but are not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

[0122] Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be carried out in vivo by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating

the region of interest and modifying it by random mutagenesis, or site-directed replacement, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome. A non-exhaustive list of preferred promoters that could be used for up-regulation of gene expression includes the promoters *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *1st*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*; *ompCD*, *copB*, *lbpB*, *ompE*, *UspA1*; *UspA2*; *TbpB* from *M. Catarrhalis*; *p1*, *p2*, *p4*, *p5*, *p6*, *lpD*, *tbpB*, *D15*, *Hia*, *Hmw1*, *Hmw2* from *H. influenzae*.

[0123] In one example, the expression of the gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream sequence of the gene, in vitro modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.

[0124] In other examples, the described approaches can be used to generate recombinant bacterial strains with improved characteristics for vaccine applications. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, strains with modulated expression of immunodominant proteins, strains with modulated shedding of outer-membrane vesicles.

[0125] Thus, also provided by the invention is a modified upstream region of the BASB040 gene, particularly the part of the upstream region identified in Sequence ID No: 11, which modified upstream region contains a heterologous regulatory element which alters the expression level of the BASB040 protein located at the outer membrane. The upstream region according to this aspect of the invention includes the sequence upstream of the BASB040 gene. The upstream region starts immediately upstream of the BASB040 gene and continues usually to a position no more than about 1000 bp upstream of the gene from the ATG start codon. In the case of a gene located in a polycistronic sequence (operon) the upstream region can start immediately preceding the gene of interest, or preceding the first gene in the operon. Preferably, a modified upstream region according to this aspect of the invention contains a heterologous promoter at a position between 500 and 700 bp upstream of the ATG.

[0126] Thus, the invention provides a BASB040 polypeptide, in a modified bacterial bleb. The invention further provides modified host cells capable of producing the non-live membrane-based bleb vectors. The invention further provides nucleic acid vectors comprising the BASB040 gene having a modified upstream region containing a heterologous regulatory element.

[0127] Further provided by the invention are processes to prepare the host cells and bacterial blebs according to the invention.

[0128] Also provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. et al. *Science* 273: 352 (1996).

[0129] Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Neisseria meningitidis*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Neisseria meningitidis* infection, in mammals, particularly humans.

[0130] The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

[0131] The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

[0132] An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

[0133] Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

[0134] It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

[0135] The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4+ ve T cell clones by Mosmann and Coffman (*Mosmann, T. R. and Coffman, R. L. (1989) TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173*). Traditionally, TH1-type responses are associated with the production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2-type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

[0136] It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2-type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes in vitro after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

[0137] Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen in vitro, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

[0138] Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

[0139] 3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Mont. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

[0140] Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22 micron membrane (European Patent number 0 689 454).

3D-MPL will be present in the range of 10  $\mu$ g-100  $\mu$ g preferably 25-50  $\mu$ g per dose wherein the antigen will typically be present in a range 2-50  $\mu$ g per dose.

[0141] Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of Quillaja *Saponaria* Molina. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with a carrier.

[0142] The method of production of QS21 is disclosed in U.S. Pat. No. 5,057,540.

[0143] Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

[0144] Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

[0145] Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21:3D-MPL will typically be in the order of 1:10 to 10:1; preferably 1:5 to 5:1 and often substantially 1:1. The preferred range for optimal synergy is 2.5:1 to 1:13D-MPL: QS21.

[0146] Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

[0147] A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

[0148] Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1 µg-200 µg, such as 10-100 µg, preferably 10 µg-50 µg per dose. Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene:alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the vaccines of the present invention will further contain a stabiliser.

[0149] Non-toxic oil in water emulsions preferably contain a non-toxic oil, e.g. squalane or squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may be, for example, phosphate buffered saline.

[0150] A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in WO 95/17210.

[0151] The present invention also provides a polyvalent vaccine composition comprising a vaccine formulation of the invention in combination with other antigens, in particular antigens useful for treating cancers, autoimmune diseases and related conditions. Such a polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore described.

[0152] While the invention has been described with reference to certain BASB040 polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions,

deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

[0153] The antigen can also be delivered in the form of whole bacteria (dead or alive) or as subcellular fractions, these possibilities do include *N. meningitidis* itself.

Compositions, Kits and Administration

[0154] In a further aspect of the invention there are provided compositions comprising a BASB040 polynucleotide and/or a BASB040 polypeptide for administration to a cell or to a multicellular organism.

[0155] The invention also relates to compositions comprising a polynucleotide and/or a polypeptide discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

[0156] Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0157] The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

[0158] In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic. In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0159] The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injec-

tion, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

[0160] For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

[0161] The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject.

[0162] A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

[0163] Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

[0164] Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

[0165] Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analy-

sis, open reading frame determination, nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

[0166] A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

[0167] A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

[0168] All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

#### DEFINITIONS

[0169] "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A. M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D. W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, N.J., 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN (Altschul, S. F. et al., *J. Mol. Biol.* 215: 403-410 (1990), and FASTA (Pearson and Lipman Proc. Natl. Acad. Sci. USA 85: 2444-2448 (1988)). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, Md. 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). The well known Smith Waterman algorithm may also be used to determine identity.

[0170] Parameters for polypeptide sequence comparison include the following:

[0171] Algorithm: Needleman and Wunsch, J. Mol. Biol. 48: 443-453 (1970)

[0172] Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,

[0173] Proc. Natl. Acad. Sci. USA. 89: 10915-10919 (1992)

[0174] Gap Penalty: 8

[0175] Gap Length Penalty: 2

[0176] A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison Wis. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

[0177] Parameters for polynucleotide comparison include the following:

[0178] Algorithm: Needleman and Wunsch, J. Mol. Biol. 48: 443-453 (1970)

[0179] Comparison matrix: matches=+10, mismatch=0

[0180] Gap Penalty: 50

[0181] Gap Length Penalty: 3

[0182] Available as: The "gap" program from Genetics Computer Group, Madison Wis. These are the default parameters for nucleic acid comparisons.

[0183] A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

[0184] (1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein  $n_n$  is the number of nucleotide alterations,  $x_n$  is the total number of nucleotides in SEQ ID NO:1,  $y$  is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and  $\cdot$  is the symbol for the multiplication operator, and wherein any non-integer product of  $x_n$  and  $y$  is rounded

down to the nearest integer prior to subtracting it from  $x_n$ . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

[0185] By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y)$$

wherein  $n_n$  is the number of nucleic acid alterations,  $x_n$  is the total number of nucleic acids in SEQ ID NO:1,  $y$  is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc.,  $\cdot$  is the symbol for the multiplication operator, and wherein any non-integer product of  $x_n$  and  $y$  is rounded down to the nearest integer prior to subtracting it from  $x_n$ .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y)$$

wherein  $n_a$  is the number of amino acid alterations,  $x_a$  is the total number of amino acids in SEQ ID NO:2,  $y$  is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and  $\cdot$  is the symbol for the multiplication operator, and wherein any non-integer product of  $x_a$  and  $y$  is rounded down to the nearest integer prior to subtracting it from  $x_a$ .

[0186] By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a * y)$$

wherein  $n_a$  is the number of amino acid alterations,  $x_a$  is the total number of amino acids in SEQ ID NO:2,  $y$  is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and  $*$  is the symbol for the multiplication operator, and wherein any non-integer product of  $x_a$  and  $y$  is rounded down to the nearest integer prior to subtracting it from  $x_a$ .

[0187] "Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

[0188] "Isolated" means altered "by the hand of man" from its natural state, i.e., if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

[0189] "Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

[0190] "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar

overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

[0191] "Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, upper respiratory tract infection, invasive bacterial diseases, such as bacteremia and meningitis.

#### EXAMPLES

[0192] The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

##### Example 1

#### Discovery and Confirmatory DNA Sequencing of the BASB040 Gene from two *N. meningitidis* Strains

A: BASB040 in *N. meningitidis* serogroup B strain ATCC13090.

[0193] The BASB040 gene of SEQ ID NO:1 was first discovered in the Incyte PathoSeq database containing unfinished genomic DNA sequences of the *N. meningitidis* strain ATCC13090. The translation of the BASB040 polynucleotide sequence, shown in SEQ ID NO:2, showed significant similarity (21% identity in a 623 amino acids overlap) to the *Neisseria meningitidis* D15 protective surface.

[0194] The sequence of the BASB040 gene was further confirmed experimentally. For this purpose, genomic DNA was extracted from  $10^{10}$  cells of the *N. meningitidis* cells (strain ATCC 13090) using the QIAGEN genomic DNA extraction kit (Qiagen GmbH), and 1  $\mu$ g of this material was submitted to Polymerase Chain Reaction DNA amplification using primers D15b-21-ctg (5'-GGG CCG CAA CCT CCG AAA TA-3') [SEQ ID NO:7] and D15b-22-ctg (5'-CGA GCC AGC CGA GGAAAC ATA C-3') [SEQ ID NO:8]. This PCR product was gel-purified and subjected to DNA sequencing using the Big Dye Cycle Sequencing kit (Perkin-Elmer) and an ABI 373A/PRISM DNA sequencer. DNA sequencing was performed on both strands with a redundancy of 2 and the full-length sequence was assembled using the SeqMan program from the DNASTAR Lasergene software package. The resulting DNA sequence and deduced polypeptide sequence are shown as SEQ ID NO:3 and SEQ ID NO:4 respectively.

B: BASB040 in *N. meningitidis* Serogroup B Strain H44/76.

[0195] The sequence of the BASB040 gene was also determined in another *N. meningitidis* serogroup B strain, the strain H44/76. For this purpose, plasmid DNA (see example 2A) containing the gene region encoding the mature BASB040 from *N. meningitidis* strain H44/76 was used as a PCR template. This material (1  $\mu$ g) was then



submitted to Polymerase Chain Reaction DNA amplification using primers D15b-01 (5'-CAT AGC ACC ATG GCC CC CGAC CTT TCC GA-3') [SEQ ID NO:9] and D15b-02 (5'-CTA GTC TAG ATT AGA AGC GTG TTC CCA AGC-3') [SEQ ID NO:10] specific for the BASB040 gene. The PCR amplicon was then submitted to DNA sequencing using the Big Dyes kit (Applied biosystems) and analyzed on a ABI 373/A DNA sequencer in the conditions described by the supplier. As a result, the polynucleotide and deduced polypeptide sequences, referred to as SEQ ID NO:5 and SEQ ID NO:6 respectively, were obtained.

[0196] Using the MegAlign program from the DNASTAR software package, an alignment of the polynucleotide sequences of SEQ ID NO:1, 3 and 5 was performed, and is displayed in FIG. 1; a pairwise comparison of identities is summarized in Table 1, showing that the three BASB040 polynucleotide gene sequences are all similar at identity level greater than 98.0%. Using the same MegAlign program, an alignment of the polypeptide sequences of SEQ ID NO:2, 4 and 6 was performed, and is displayed in FIG. 2; a pairwise comparison of identities is summarized in Table 2, showing that the three BASB040 protein sequences are all similar at an identity level greater than or equal to 99.0%.

[0197] Taken together, these data indicate strong sequence conservation of the BASB040 gene among the two *N. meningitidis* serogroup B strains.

TABLE 1

Pairwise identities of the BASB040 polynucleotide sequences (in %)		
	SeqID No: 3	SeqID No: 5
SeqID No: 1	99.9	98.3
SeqID No: 3		98.4

[0198]

TABLE 2

Pairwise identities of the BASB040 polypeptide sequences (in %)		
	SeqID No: 4	SeqID No: 6
SeqID No: 2	100.	99.0
SeqID No: 4		99.0

### Example 2

#### Construction of Plasmid to Express Recombinant BASB040

##### A: Cloning of BASB040.

[0199] The NcoI and XbaI restriction sites (underlined) engineered into the forward D15b-01 (5'-CAT AGC ACC ATG GCC GCC GAC CTT TCC GA-3') [SEQ ID NO:9] and reverse D15b-02 (5'-CTA GTC TAG ATT AGA AGC GTG TTC CCA AGC-3') [SEQ ID NO:10] amplification primers, respectively, permitted directional cloning of a BASB040 PCR product into the commercially available *E. coli* expression plasmid pBADgIII(A) (Invitrogen, USA, ampicillin resistant). This plasmid provides the signal peptide from the bacteriophage fd pII protein such that a mature

BASB040 protein could be targeted to the periplasm of *E. coli*. The BASB040 PCR product was purified from the amplification reaction using Wizard PCR Prep™ (Promega) according to the manufacturers instructions. To produce the required NcoI and XbaI termini necessary for cloning, purified PCR product was sequentially digested to completion with NcoI and XbaI restriction enzymes as recommended by the manufacturer (Boehringer Mannheim). Digested BASB040 PCR products and pBAD were gel-purified and ligated together using an approximately 5-fold molar excess of the digested fragment to the vector. A standard ~20 µl ligation reaction (~16° C., ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units/reaction, Boehringer Mannheim). An aliquot of the ligation was used to transform electro-competent *E. coli* Top10 cells according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37° C. in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates containing Ampicillin (50 µg/ml). Individual ampicillin-resistant colonies were selected and analyzed by whole cell-based PCR to verify that transformants contained the BASB040 DNA insert. Transformants that produced the expected PCR product were identified as strains containing a BASB040 expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB040.

##### B: Expression Analysis of PCR-Positive Transformants.

[0200] For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing ampicillin (50 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37° C. with shaking (~250 rpm). An aliquot of the overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 ml of LB Kn broth and grown at 37° C. with shaking (~250 rpm) until the culture turbidity reached O.D. 600 of ~0.5, i.e. mid-log phase (usually about 1.5-2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB040 protein induced by the addition of L-Arabinose to a final concentration of 0.2% (w/v). Incubation of both the arabinose-induced and non-induced cultures continued for an additional ~4 hours at 37° C. with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by centrifugation in a microcentrifuge at room temperature for 3 minutes. Individual cell pellets were suspended in ~50 µg of sterile water, then mixed with an equal volume of 2x Laemelli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for 3 min to denature protein. Equal volumes (~15 µl) of both the crude arabinose-induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels, Novex). The induced and non-induced lysate samples were electrophoresed together with prestained molecular weight markers under conventional conditions using a standard SDS/Tris/glycine running buffer. Following electrophoresis, one gel was stained with comassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB040 arabinose-inducible protein(s).

## Example 3

## Analysis of the Non-Coding Flanking Regions of the BASB040 Gene, and its Exploitation for Modulated BASB040 Gene Expression

[0201] The non-coding flanking regions of the BASB040 gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the shine-dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences.

[0202] This sequence information allows the modulation of the natural expression of gene BASB040. The upregulation of the gene expression may be accomplished by altering the promoter, the shine-dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modifications. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or may be uncoupled from this regulation. In another approach, the expression of the gene can be put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation include, but are not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

[0203] Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be done in vivo by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random mutagenesis, or site-directed mutagenesis, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome. A non-exhaustive list of preferred promoters that could be used for up-regulation of gene expression includes the promoter *porA*, *porB*, *lbpB*, *tbpB*, *p110*, *1st*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*.

[0204] In one example, the expression of the gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream sequence of the gene, in vitro modification of this sequence, and reintroduc-

tion into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium. In other examples, the described approaches can be used to generate recombinant bacterial strains with improved characteristics for vaccine applications. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, strains with modulated expression of immunodominant proteins, strains with modulated shedding of outer-membrane vesicles.

[0205] A region directly upstream of the BASB040 gene is given in the sequence of SEQ ID NO:11. This sequence is a further aspect of the invention.

## SEQUENCE INFORMATION

BASB040 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

*Neisseria meningitidis* BASB040 polynucleotide sequence from strain ATCC 13090

ATGATGATCAAACCGACCGCCCTGCTCCTGCCGGCTTTATTTTTCTTTCC

GCACGCATACGCGCTGCCGCCGACCTTTCCGAAAAAAGGCGGGGGTT

TGCGATTGTTCAAAAACAAAAGCCCCGACACCGAATCAGTTAAATTAATA

CCCAAATTCCTCCGTCGACATCGACAGCAGGATAGTGAAATCAAAGATAT

GGTCGAAGAACACCTGCCGCTCATCAGCAGCAGCAGGAAGAAGTATTGG

ACAAGGAACAGACGGGCTTCCTCGCGAAGAAGCACCGGACACGTTAA

ACCATGCTCCGAGCAAAGGCTATTTTCAGCAGCAAAGTCAGCCTGACGG

AAAAGACGGAGCTTATACGGTACACATCACACCGGGCCCGCACCAAAA

TGCGCAACGTCGGTGTGCGCATCCTCGGCGACATCCTTTTCAGACGGCAAC

CTCGCGAATACTACCGCAACGCGCTGGAAAACAGCAGCAGCCGGTAGG

CAGTGATTTTCGATCAGGACAGTTGGGAAAACAGCAAACCTCCGCTCTCG

GCGCGGTAACGCGAAAAGGCTACCCGCTTGCCAAGCTCGGCAACACCCGG

GCGGCGGTCAACCCGATACCGCCACCGTCGATTTGAACGTCGTCGTGG

CAGCGGCCGCCCATCGCCTTCGGCGACTTTGAAATCACCGGCACACAGC

GTTACCCCGAACAAATCGTCTCCGGCTGGCGCGCTTCCAACCGGGCACG

CCCTACGACCTCGACCTGCTGCTCGACTTCCAACAGGCACTCGAACAAAA

CGGGCATTATCCGGCGCTCCGTACAAGCCGACTTCGACCGTCTCCAAG

GCGACCGGTCCTCCGTCGAAAGTCAGCGTAACCGAGGTCAAACGCCACAAG

CTCGAAACCGGCATCCGCTCGATTGGAATACGGTTTGGGCGGCAAAAT

CGCCTACGACTATTACAACCTTTCAACAAGGCTATATCGGCTCGGTCCG

TCTGGGATATGGACAAATACGAAACCACGCTTGCCCGGGCATCAGCCAG

CCGCGCAACTATCGGGGCAACTACTGGACAAGCAACGTTTCTACAACCG

TTTCGACCCCAAAACCTCGAAAAACGCGCCTTCTCCGGCGGCATCTGGT

ATGTGCGCGACCGCGGGGCATCGATGCCAGGCTGGGGGAGAGTTTCTC

GCAGAAGGCGGAAAAATCCCGGCTCGGATATCGATTGGGCAACAGCCA

CGCCACGATGCTGACCGCTCTTGGAAACGCCAGCTGCTCAACAACGTGC

-continued

TGCATCCCGAAAACGGCCATTACCTCGACGGCAAATCGGTACGACTTTG  
 GGCACATTCTGTCTCCACCGCGCTGATCCGCACCTCTGCCCCGTGACGG  
 TTATTTCTTACGCCCCGAAAAAATAAAGCTCGGCACGTTATCATACGCG  
 GACAAGCGGGTTACACCGTGTCCCGCGACAATGCCGACGTTCCTTCAGGG  
 CTGATGTTCCGACGCGCGCGCGTCTTCCGTGCGCGGTTACGAACTCGA  
 CAGCATCGGACTTGCCGGCCCCGAACGGATCGGTCTGCCGGAACGCGCCC  
 TCCTGGTGGGCGAGCTGGAATACCACTGCCGTTTACGCGCACCTTTCC  
 GCGCGGTGTTCACGATATGGGCGATGCCGCCCAATTTCAAACGTAT  
 GAAGCTGAAACACGGTTCCGGACTGGGCGTGCCTGGTTACGCCCGCTTG  
 CGCCGTTTTCTTCGACATCGCCTACGGGCACAGCGATAAGAAAATCCGC  
 TGGCACATCAGCTTGGGAACACGCTTCTAA

SEQ ID NO: 2

*Neisseria meningitidis* BASB040 polypeptide  
 sequence deduced from the polynucleotide of  
 Seq ID NO: 1

MMIKPTALLLPALFFPHAYAPAADLSENKAAGFALFKNKSPDTESVKLLK  
 PKFPVRIIDTQDSEIKDMVEEHLPLITQQEVEVLDKEQTGFLAEEAPDNVK  
 TMLRSKGYFSSKVSLETEKDGAYTVHITPGPRTKIANVGVAILGDIILSDGN  
 LAEYRNALENWQPPVGSDFDQDQSWENSKT SVLGAVTRKGYPLAKLGNTR  
 AAVNPDATVLDLNVVDSGRPIAFGDFEITGTQRYPEQIVSGLARFQPGT  
 PYDLDLLDFQQALEQNGHYSVASVQADFDRLQGRVVPVKVSVTEVSRKH  
 LETGIRLDSEYGLGGKIAYDYNFLNKGYIGSVVWMDKYETTLAAGISQ  
 PRNYRGNWYTSNVSYNRSTQNLKRAFSGGIWYVRDRAGIDARLGAEFL  
 AEGRKIPGSDIDLGNSHATMLTASWKRQLLNNVLPENGHYLDGKIGITTL  
 GTFLSSTALIRTSARAGYFFTPENKLGTFIIRGQAGYTVARDNADVPSG  
 LMFRRSGASSVRGYELDSIGLAGPNGSVLPERALLVGSLEYQLPFRITLS  
 GAVFHDMDGAAANFKRMKLBKHSGLVGRWFSPLAPFSFDIAYGHSDDKIR  
 WHISLGRF

SEQ ID NO: 3

*Neisseria meningitidis* BASB040 polynucleotide  
 sequence from strain ATCC 13090

ATGATGATCAAACCGACCGCCCTGCTCCTGCCGCTTATTTTTCTTTCC  
 GCACGCATACGCGCTGCCGCGACCTTCCGAAAACAAGCGCGCGGTT  
 TCGCATGTTCAAAAAACAAGCCCGACACCGAATCAGTTAAATTAATA  
 CCCAAATCCCGCTCCGCATCGACACGCAGGATAGTGAAATCAAAGATAT  
 GGTGGAAGAACACCTGCCGCTCATCACGCAGCAGCAGGAAGAAGTATTGG  
 ACAAGGAACAGACGGGCTTCTCGCGAAGAAGACCGGACAACGTTAAA  
 ACCATGCTCCGACGAAAGGCTATTTACGACGAAAGTACGCTGACGGA  
 AAAAGACGGAGCTTATACGTTACACATCACACCGGGCCCGCGCACAAAA  
 TCGCCAACGTCGGTGTGCCATCTCGCGACATCCTTTCAGACGGCAAC  
 CTCGCCGAATACTACCGCAACGCGCTGGAAAATGGCAGCAGCCGGTAGG  
 CAGTGATTTTCGATCAGGACAGTTGGGAAAACAGCAAAACTTCCGCTCTCG

-continued

GCGCGGTAAACGCGCAAAGGCTACCCGCTTGCCAAGCTCGGCAACACCCGG  
 GCGGCGGTCAACCCGATACCGCCACCGTTCGATTGAACTCGTCTGGA  
 CAGCGCCGCCCCATCGCCTTCGGCGACTTTGAAATCACCGGCACACAGC  
 GTTACCCCGAACAAATCGTCTCCGGCTGGCGCGTTCACACCGGCGACG  
 CCTACGACCTCGACTGCTGCTCGACTTCCAACAGGCACTCGAACAAAA  
 CGGGCATTATTCGGCGCGTCCGTACAAGCGACTTCGACCGCTCCAAG  
 GCGACCGCGTCCCGCTCAAAGTACGCGTAAACGAGGTCAAACGCCACAAG  
 CTCGAAACCGGCATCCGCTCGATTTCGAAATACGTTTGGGCGGCAAAAT  
 CGCCTACGACTATTACAACCTCTTCAACAAAGGCTATATCGGCTCGGTGC  
 TCTGGGATATGGACAAATACGAAACACGCTTCCCGCGGCATCAGCCAG  
 CCGCGCAACTATCGGGCAACTACTGGACAAGCAACGTTTCTTACAACCG  
 TTCGACCCCAAAACCTCGAAAAACGCGCCTTCTCCGGCGGCATCTGGT  
 ATGTGCGGACCGCGCGGCATCGATGCCAGGCTGGGGCGAGGTTTCTC  
 GCAGAAGCGCGAAAATCCCGGCTCGGATATCGATTGGGCAACAGCCA  
 CGCCACGATGTCACCGCCTCTTGGAACCGCAGCTGCTCAACAACGTGC  
 TGCATCCCGAAAACGGCCATTACCTCGACGGCAAATCGGTACGACTTTG  
 GGCACATTCTGTCTCCACCGCGCTGATCCGCACCTCTGCCGTGACGG  
 TTATTTCTTACGCCCCGAAAAAATAAAGCTCGGCACGTTATCATACGCG  
 GACAAGCGGGTTACACCGTGTCCCGCGACAATGCCGACGTTCTTTCAGGG  
 CTGATGTTCCGACGCGCGCGCGTCTTCCGTGCGCGGTTACGAACTCGA  
 CAGCATCGGACTTGCCGGCCCCGAACGGATCGGTCTGCCCGAACGCGCCC  
 TCCTGGTGGGCGAGCTGGAATACCAACTGCCGTTTACGCGCACCTTTCC  
 GCGCGGTGTTCACGATATGGGCGATGCCGCCCAATTTCAAACGTAT  
 GAAGCTGAAACACGGTTCCGGACTGGGCGTGCCTGGTTTCAGCCGCTTG  
 CGCCGTTTTCTTCGACATCGCCTACGGGCACAGCGATAAGAAAATCCGC  
 TGGCACATCAGCTTGGGAACACGCTTCTAA

SEQ ID NO: 4

*Neisseria meningitidis* BASB040 polypeptide  
 sequence deduced from the polynucleotide of  
 Seq ID NO: 3

MMIKPTALLLPALFFPHAYAPAADLSENKAAGFALFKNKSPDTESVKLLK  
 PKFPVRIIDTQDSEIKDMVEEHLPLITQQEVEVLDKEQTGFLAEEAPDNVK  
 TMLRSKGYFSSKVSLETEKDGAYTVHITPGPRTKIANVGVAILGDIILSDGN  
 LAEYRNALENWQPPVGSDFDQDQSWENSKT SVLGAVTRKGYPLAKLGNTR  
 AAVNPDATVLDLNVVDSGRPIAFGDFEITGTQRYPEQIVSGLARFQPGT  
 PYDLDLLDFQQALEQNGHYSVASVQADFDRLQGRVVPVKVSVTEVSRKH  
 LETGIRLDSEYGLGGKIAYDYNFLNKGYIGSVVWMDKYETTLAAGISQ  
 PRNYRGNWYTSNVSYNRSTQNLKRAFSGGIWYVRDRAGIDARLGAEFL  
 AEGRKIPGSDIDLGNSHATMLTASWKRQLLNNVLPENGHYLDGKIGITTL  
 GTFLSSTALIRTSARAGYFFTPENKLGTFIIRGQAGYTVARDNADVPSG

-continued

LMFRSGGASSVRGYELDSIGLAGPNGSVLPERALLVGSLEYQLPFRTRLS  
 GAVFHDMDAAANFKRMKCLKHGSGLGVRWFSPLAPFSFDIAYGHSDKKIR  
 WHISLGTRF

SEQ ID NO:5

*Neisseria meningitidis* BASB040 polynucleotide  
 sequence from strain H44/76

GCCGCCGACCTTTCCGAAAACAAGCGCGGGTTTCGCATTGTTCAAAA  
 CAAAAGCCCCGACACCGAATCAGTCAAATTAACCCAAATTCCTCCGTC  
 TCATCGACACGCAGGACAGTAAATCAAAGATATGGTCGAAGAACACCTG  
 CCCTCATCACGCAGCAGCAGGAAGAAGTATTGGACAAGGAACAGACGGG  
 CTTCTCCGCGAAGAAGCGCCGCAACCGTTAAACAGTATGCTCCGACGA  
 AAGGCTATTTTCCAGCAGCAAAGTCCGCTGACGGAAGAACGAGCTTAT  
 ACGGTACACATCACACCGGGCCCGCACCAAAATCGCCAACGTCGCGCT  
 CGCCATCCTCGGCAGATCCTTTCCAGACGGCAACCTCGCCGAATACTACC  
 GCAACGCGCTGGAAGTGGCAGCAGCGTAGGCAGCGATTTCCGATCAG  
 GACAGTGGGAAAACAGCAAACTTCCGCTCCTCGGCGCGTAACGCGCAA  
 AGCCTACCCGCTTGCCAAGCTCGCAATACGCAGGCGCCGCTCAACCCCG  
 ATACCGCCACCGCGATTGAACTCGTCTGTCGTCGACAGCGCCGCCCATC  
 GCCTTCGGCGACTTTGAAATCACCGGCACACAGCGTTACCCCGAACAAAT  
 CGTCTCCGGCCTTGCGCGTTTCCAGCCCGGTATGCCGTACGACCTCGACC  
 TGCTGCTCGACTTCCAACAGGCGCTCGAACAAAACGGGCATTATTCCGGC  
 GCGTCCGTACAAGCGACTTTCGACCGCTTCAAGCGACCGCGTCCCGT  
 CAAAGTCAGCGTAACCGAGGTCAAAACGCCAACAACCTCGAAACCGCATCC  
 GCCTCGATTGCAATACGCTTTGGCGGCAAAATCGCTACGACTATTAC  
 AACCTCTTCAACAAGGCTATATCGTTCGGTCTGTCGGGATATGGACAA  
 ATACGAAACCACGCTTCCGCGCGCATCAGCCAGCCGCGCAACTATCGGG  
 GCAACTACTGGACAAGCAACGTTTCTTACAACCGTTTCGACACCCAAAAC  
 CTCGAAAACGCGCCTTCTCCGGCGCGTCTGGTATGTGCGGACCGCGC  
 GGGCATCGATGCCAGGCTGGGGCGGAATTTCTCGCAGAAGCCGGA  
 TCCCGGCTCGGCTGTGATTGGGCAACAGCCACGCCACGATGCTGACC  
 GCCTCTTGAAACGCCAGCTGCTCAACAACGTGCTGCATCCCGAAAACGG  
 CCATTACCTCGACGGCAAAATCGGTACGACTTTGGGCACATTCCTGTCT  
 CCACCGCGTGTATCCGCACCTCTGCCCGTGCAGGTTATTTCTTACGCCCC  
 GAAAACAAAAAATCGGCACGTTTATCATACACGCGGACAAGCGGTTACAC  
 CGTTGCCCGGCAATGCCGACGTTCTTCCAGGGCTGATGTTCCGACGCG  
 GCGGCGCTTCTCCGTGCGGGTTACGAACTCGACAGCATCGGACTTGGC  
 GGCCCGAACGGATCGGCTTCCGCGAACCGCCCTCTGTTGGGCGAGCTT  
 GGAATACCAACTGCCGTTTACGCGCACCTTTCCGGCGGGTGTCCACG  
 ATATGGGCGATGCCCGCCCAATTTCAAACGATGAAGCTGAAACACGGT  
 TCGGACTGGGCGTGCCTGTTTCCAGCCGCTTGCCTGTTTCTTCCGCA

-continued

CATCGCCTACGGGCACAGCGATAAGAAAATCCGCTGGCACATCAGCTTGG  
 GAACACGCTTCTAA

SEQ ID NO:6

*Neisseria meningitidis* BASB040 polypeptide  
 sequence deduced from the polynucleotide of  
 Seq ID NO:5

AADLSENKAAGFALFKNKSPDTEVSKLKPFPVLIDTQDSEIKDMVEEHL  
 PLITQQQEEVLDKEQTGLAEEAPDNVKTMLRSKGYFSSKVSLETKDGAY  
 TVHI TPGPRTKIANVGVAILGDI LSDGNLAEYRNALLENWQQPVGSDFDQ  
 DSWENSKTSVLGAVTRKAYPLAKLGNLQAAVNPDTATADLNWVVDVSGRPI  
 AFGDFEITGTQRYPEQIVSGLARFQPGMPYDLDDLLDFQQAELQNGHYSG  
 ASVQADFDRLQGDRVPVKVSVTEVVKRHKLETGIRLDSEYGLGGKIAYDYY  
 NLFNKGYIGSVVWMDKYEETTLAAGISQPRNRYGNWYTSNVSYNRSTTQN  
 LEKRAFSGGVWVVRDRAGIDARLGAFLAERKIPGSAVDLGNSHATMLT  
 ASWKRQLLNNVLPENGHYLDGKIGTTLGTFLLSSTALIRTSARAGYFFTP  
 ENKKGTFIIRQAGYTVARDNADVPGLMFRSGGASSVRGYELDSIGLA  
 GPNGSVLPERALLVGSLEYQLPFRTRLSGAVFHDMDAAANFKRMKCLKHGS  
 SGLGVRWFSPLAPFSFDIAYGHSDKKIRWHISLGTRF

SEQ ID NO:7

GGG CCG CAA CCT CCG AAA TA

SEQ ID NO:8

CGA GCC AGC CGA GGA AAC ATA

SEQ ID NO:9

CAT AGC ACC ATG GCC GCC GAC CTT TCC GA

SEQ ID NO:10

CTA GTC TAG ATT AGA AGC GTG TTC CCA AGC

SEQ ID NO:11

*Neisseria meningitidis* polynucleotide  
 sequence up-stream of the BASB040 gene sequence,  
 in strain ATCC 13090

AAACGGATAATCCATTTACGACGGTGGAAACCGCTTCCGCCAAATCGGC  
 AACGACCTGCCGCGTAACGTCGGAACCGTGATTCCGCAACCGCCGCGCCA  
 AAACCAAAGCCCAAGCAAATGCCGATATAGTTGGCATTGGCAATCGCG  
 TTAATCGGGTTGGCGACAGGTTTATCAGCAGCGATTTCACACTTCCAC  
 AATGCCGGAAGCGCGCGCGGACACATCGCCCGCCCGCCAAAACAA  
 TGTGCGTCCGGAAAACCATACCGCGATGACGGCGGTGAGGCTGCGGAA  
 AACGTACCAATGAGGTAAGGATGATAATCGGCTGATATGCGCCTTGT  
 GCCTTTTGGTGTGCGGATTTGGCCGCCACCAAAATAAATACCAAAA  
 CCGGCGGACCGCTTTGAGCGCGCCGACAAACAGGCTGCCGAACAAGCCT  
 GCCGCAAGCCAGTTGCGGGGAAACCGAACCGATTACGATGCCAACCGC  
 CAAACCGGCGCAATCTGCCTGACCAGGCTGACGCGGCCGATCGCATGAA  
 ATAAGGATTTGCCGAACGCATAATTTCTCTTATGTTGTGATATGTTAA  
 AAAATGTTGTATTTTAAAGAAAACCTATTCTCTGTGTTTTTTTTATTT  
 TCGGCTGTGTTTTAAGGTTGCGTTGATTTGCCCTATGACGTCGCCGACG  
 GCTTTGCTTTATCATTCGGCGCAACGGTTAATTTATTGAACGAAAATAA

- continued

ATTTATTTAATCCTGCCTATTTTCCGGCCTATTCCGAAACGCAGCCTGT  
 TTTCATATGCGGATTGGAACAAAATACCTTAAACAAGCAGATACATT  
 TCCGGCGGGCCGCAACCTCCGAAATACCGCGGCGAGTATGCCGTCTGAAG  
 TGTCGCCGCCCGTCCGAACAACACAAAACAGCCGTTTCGAAACCTGTCC  
 GAACAGTGTTAGAATCGAAATCTGCCACACCGATGCACGACACCCGTACC

#### Deposited Materials

[0206] A deposit containing a *Neisseria meningitidis* Sero-group B strain has been deposited with the American Type Culture Collection (herein "ATCC") on Jun. 22, 1997 and assigned deposit number 13090. The deposit was described as *Neisseria meningitidis* (Albrecht and Ghon) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *N. meningitidis* isolate. The deposit is described in Int. Bull. Bacteriol. Nomencl. Taxon. 8: 1-15 (1958).

[0207] The *Neisseria meningitidis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

[0208] The deposited strain contains the full length BASB040 gene. The sequence of the polynucleotides contained in the deposited strain, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

[0209] The deposit of the deposited strain has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strain is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 1830

<212> TYPE: DNA

<213> ORGANISM: *Neisseria meningitidis*

<400> SEQUENCE: 1

```

atgatgatca aaccgaccgc cctgctcctg cgggctttat ttttctttcc gcacgcatac      60
gcgctgcccg ccgaccttcc cgaaaacaag gcgggcgggt tcgcattggt caaaaaaaaa      120
agccccgaca ccgaatcagt taaattaaaa cccaaattcc ccgtccgcat cgacacgcag      180
gatagtghaa tcaaagatat ggtcgaagaa cacctgccgc tcatcacgca gcagcaggaa      240
gaagtattgg acaaggaaca gacgggcttc ctgcgccgag aagcaccgga caacgttaaa      300
accatgctcc gcagcaaaag ctatttcagc agcaaagtca gcctgacgga aaaagacgga      360
gcttatacgg tacacatcac accgggcccg cgcacaaaaa tcgccaacgt cgggtgtcgc      420
atcctcggcg acatccttcc agacggcaac ctgcgccgag actaccgcaa cgcgctggaa      480
aactggcagc agccggtagg cagtgatttc gatcaggaca gttgggaaaa cagcaaaaact      540
tccgtcctcg gcgcggtaac gcgaaaaggc taccgccttg ccaagctcgg caacacccgg      600
gcgcccgcca accccgatac cgcaaccgtc gatttgaacg tcgtcgtgga cagcggcccgc      660
cccatcgctc tcggcgactt tgaatcacc ggcacacagc gttaccccga acaaatcgtc      720
tccggcctgg cgcgcttcca accgggcaac ccctacgacc tcgacctgct gctcgacttc      780
caacaggcac tcgaacaaaa cgggcattat tccggcgcgt ccgtacaagc cgacttcgac      840
cgtctccaag gcgaccgcgt ccccgtaaaa gtcagcgtaa ccgaggtaa acgccacaag      900
ctcgaaccgg gcatccgcct cgattcggaa tacggtttgg gcggcaaaaat cgcctacgac      960
tattacaacc tcttaacaaa aggetatata ggetcggctg tetgggatat ggacaaatac     1020
gaaaccacgc ttgcgcggc catcagccag ccgcgcaact atcggggcaa ctactggaca     1080
agcaacgttt cctacaaccg ttcgaccacc caaaacctcg aaaaacgcgc cttctccggc     1140

```

-continued

---

```

ggcatctggt atgtgcgcgca ccgcgcgggc atcgatgcca ggctgggggc agagtttctc 1200
gcagaaggcc ggaaaatccc cggctcggat atcgatttgg gcaacagcca cgccacgatg 1260
ctgaccgcct ctggaaacg ccagctgctc aacaacgtgc tgcattcccga aaacggccat 1320
tacctcgacg gcaaaatcgg tacgactttg ggcacattcc tgtcctccac cgcgctgatc 1380
cgcacctctg cccgtgcagg ttatttcttc acgcccgaaa acaaaaaact cggcacgttc 1440
atcatacgcg gacaagcggg ttacaccggt gcccgcgaca atgccgacgt tccttcaggg 1500
ctgatgttcc gcagcggcgg cgcgcttctc gtgcgcggtt acgaactcga cagcatcggg 1560
cttgccggcc cgaacggatc ggtcctgccc gaacgcgccc tcctggtggg cagcctggaa 1620
taccactgc cgtttacgcg cacccttctc ggcgcggtgt tccacgatat gggcgatgcc 1680
ggcccaatt tcaaacgtat gaagtgaaa cacggttcgg gactgggctg gcgctggttc 1740
agcccgttg cgcgcttttc ctctgacatc gcctacgggc acagcgataa gaaaatccgc 1800
tggcacatca gcttgggaac acgcttctaa 1830

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 609

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Neisseria meningitidis

&lt;400&gt; SEQUENCE: 2

```

Met Met Ile Lys Pro Thr Ala Leu Leu Leu Pro Ala Leu Phe Phe Phe
  1          5          10          15
Pro His Ala Tyr Ala Pro Ala Ala Asp Leu Ser Glu Asn Lys Ala Ala
  20          25          30
Gly Phe Ala Leu Phe Lys Asn Lys Ser Pro Asp Thr Glu Ser Val Lys
  35          40          45
Leu Lys Pro Lys Phe Pro Val Arg Ile Asp Thr Gln Asp Ser Glu Ile
  50          55          60
Lys Asp Met Val Glu Glu His Leu Pro Leu Ile Thr Gln Gln Gln Glu
  65          70          75          80
Glu Val Leu Asp Lys Glu Gln Thr Gly Phe Leu Ala Glu Glu Ala Pro
  85          90          95
Asp Asn Val Lys Thr Met Leu Arg Ser Lys Gly Tyr Phe Ser Ser Lys
 100          105          110
Val Ser Leu Thr Glu Lys Asp Gly Ala Tyr Thr Val His Ile Thr Pro
 115          120          125
Gly Pro Arg Thr Lys Ile Ala Asn Val Gly Val Ala Ile Leu Gly Asp
 130          135          140
Ile Leu Ser Asp Gly Asn Leu Ala Glu Tyr Tyr Arg Asn Ala Leu Glu
 145          150          155          160
Asn Trp Gln Gln Pro Val Gly Ser Asp Phe Asp Gln Asp Ser Trp Glu
 165          170          175
Asn Ser Lys Thr Ser Val Leu Gly Ala Val Thr Arg Lys Gly Tyr Pro
 180          185          190
Leu Ala Lys Leu Gly Asn Thr Arg Ala Ala Val Asn Pro Asp Thr Ala
 195          200          205
Thr Val Asp Leu Asn Val Val Val Asp Ser Gly Arg Pro Ile Ala Phe
 210          215          220
Gly Asp Phe Glu Ile Thr Gly Thr Gln Arg Tyr Pro Glu Gln Ile Val
 225          230          235          240

```

-continued

---

Ser Gly Leu Ala Arg Phe Gln Pro Gly Thr Pro Tyr Asp Leu Asp Leu  
 245 250 255  
 Leu Leu Asp Phe Gln Gln Ala Leu Glu Gln Asn Gly His Tyr Ser Gly  
 260 265 270  
 Ala Ser Val Gln Ala Asp Phe Asp Arg Leu Gln Gly Asp Arg Val Pro  
 275 280 285  
 Val Lys Val Ser Val Thr Glu Val Lys Arg His Lys Leu Glu Thr Gly  
 290 295 300  
 Ile Arg Leu Asp Ser Glu Tyr Gly Leu Gly Gly Lys Ile Ala Tyr Asp  
 305 310 315 320  
 Tyr Tyr Asn Leu Phe Asn Lys Gly Tyr Ile Gly Ser Val Val Trp Asp  
 325 330 335  
 Met Asp Lys Tyr Glu Thr Thr Leu Ala Ala Gly Ile Ser Gln Pro Arg  
 340 345 350  
 Asn Tyr Arg Gly Asn Tyr Trp Thr Ser Asn Val Ser Tyr Asn Arg Ser  
 355 360 365  
 Thr Thr Gln Asn Leu Glu Lys Arg Ala Phe Ser Gly Gly Ile Trp Tyr  
 370 375 380  
 Val Arg Asp Arg Ala Gly Ile Asp Ala Arg Leu Gly Ala Glu Phe Leu  
 385 390 395 400  
 Ala Glu Gly Arg Lys Ile Pro Gly Ser Asp Ile Asp Leu Gly Asn Ser  
 405 410 415  
 His Ala Thr Met Leu Thr Ala Ser Trp Lys Arg Gln Leu Leu Asn Asn  
 420 425 430  
 Val Leu His Pro Glu Asn Gly His Tyr Leu Asp Gly Lys Ile Gly Thr  
 435 440 445  
 Thr Leu Gly Thr Phe Leu Ser Ser Thr Ala Leu Ile Arg Thr Ser Ala  
 450 455 460  
 Arg Ala Gly Tyr Phe Phe Thr Pro Glu Asn Lys Lys Leu Gly Thr Phe  
 465 470 475 480  
 Ile Ile Arg Gly Gln Ala Gly Tyr Thr Val Ala Arg Asp Asn Ala Asp  
 485 490 495  
 Val Pro Ser Gly Leu Met Phe Arg Ser Gly Gly Ala Ser Ser Val Arg  
 500 505 510  
 Gly Tyr Glu Leu Asp Ser Ile Gly Leu Ala Gly Pro Asn Gly Ser Val  
 515 520 525  
 Leu Pro Glu Arg Ala Leu Leu Val Gly Ser Leu Glu Tyr Gln Leu Pro  
 530 535 540  
 Phe Thr Arg Thr Leu Ser Gly Ala Val Phe His Asp Met Gly Asp Ala  
 545 550 555 560  
 Ala Ala Asn Phe Lys Arg Met Lys Leu Lys His Gly Ser Gly Leu Gly  
 565 570 575  
 Val Arg Trp Phe Ser Pro Leu Ala Pro Phe Ser Phe Asp Ile Ala Tyr  
 580 585 590  
 Gly His Ser Asp Lys Lys Ile Arg Trp His Ile Ser Leu Gly Thr Arg  
 595 600 605  
 Phe

<210> SEQ ID NO 3  
 <211> LENGTH: 1830  
 <212> TYPE: DNA

-continued

<213> ORGANISM: *Neisseria meningitidis*

&lt;400&gt; SEQUENCE: 3

```

atgatgatca aaccgaccgc cctgctcctg ccggtttat tttctttcc gcacgcatac    60
gcgctgccc ccgaccttc cgaacaacaag gcggcgggtt tcgcatgtt caaaaaacaaa   120
agccccgaca ccgaatcagt taaattaaaa cccaaattcc ccgtccgat cgacacgcag   180
gatagtgaat tcaaagatat ggtcgaagaa cacctgccgc tcatcacgca gcagcaggaa   240
gaagtattgg acaaggaaca gacgggcttc ctgcgccgaag aagcaccgga caacgttaaa   300
accatgctcc gcagcaaagg ctatttcagc agcaaagtca gcctgacgga aaaagacgga   360
gcttataccg tacacatcac accgggcccgc cgcacaaaaa tcgccaacgt cgggtgctgcc   420
atcctcggcg acatccttcc agacggcaac ctgcgccgaat actaccgcaa cgcgctggaa   480
aactggcagc agccggtagg cagtgatctc gatcaggaca gttgggaaaa cagcaaaact   540
tccgctctcg gcgcggaac gcgcaaaagg taccgcttg ccaagctcgg caacacccgg   600
gcggcctgca accccgatac cgcaccgctc gatttgaacg tcgtcgtgga cagcggcccgc   660
cccctcgcct tcggcgactt tgaaatcacc ggacacagc gttacccga acaaatcgtc   720
tccggctcgg cgcgcttcca accgggcacg ccctacgacc tcgacctgct gctcgaactc   780
caacaggcac tcgaacaaaa cgggcattat tccggcgcgt ccgtacaagc cgacttcgac   840
cgtctccaag gcgaccgctt ccccgtaaaa gtcagcgtaa ccgaggtcaa acgccacaag   900
ctcgaaaccc gcacccgctt cgattcggaa tacggtttgg gcggcaaaat cgcctacgac   960
tattacaacc tcttcaacaa aggctatata ggctcggctg tctgggatat ggacaaatac  1020
gaaaccacgc ttgccgcccg catcagccag ccgcgcaact atcggggcaa ctactggaca  1080
agcaacgttt cctacaaccg ttcgaccacc caaacctcg aaaaacgcgc cttctccggc  1140
ggcatctggt atgtgcgcga ccgcgccggc atcgatgcca ggctgggggc agagtctctc  1200
gcagaaggcc ggaaaatccc cggctcggat atcgatttgg gcaacagcca cgcaccgatg  1260
ctgaccgcct cttggaaacg ccagctgctc aacaacgtgc tgcatcccga aaacggccat  1320
tacctcgacg gaaaatcgg tacgaacttg ggcacattcc tgcctccac cgcgctgatc  1380
cgcacctctg cccgtgcagg ttatttcttc acgcccgaaa acaaaaaact cggcacgttc  1440
atcatacgcg gacaagcggg ttacaaccgt gcccgcgaca atgccgacgt tccttcaggg  1500
ctgatgttcc gcagcggcgg cgcgtcttcc gtgcgcggtt acgaactcga cagcatcgga  1560
cttgccggcc cgaacggatc ggtcctgccc gaacgcgccc tcctggtggg cagcctggaa  1620
taccaaactgc cgtttacgcg caccctttcc ggcgcggtgt tccacgatat gggcgatgcc  1680
gccgccaatt tcaaactgat gaagtgaata cacggttcgg gactgggctg gcgctggttc  1740
agcccgcttg cgccttttcc cttcgacatc gcctacgggc acagcgataa gaaaatccgc  1800
tggcacatca gcttggaac acgcttctaa                                1830

```

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 609

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Neisseria meningitidis*

&lt;400&gt; SEQUENCE: 4

```

Met Met Ile Lys Pro Thr Ala Leu Leu Leu Pro Ala Leu Phe Phe Phe
  1             5             10             15

```



-continued

---

Pro His Ala Tyr Ala Pro Ala Ala Asp Leu Ser Glu Asn Lys Ala Ala  
 20 25 30  
 Gly Phe Ala Leu Phe Lys Asn Lys Ser Pro Asp Thr Glu Ser Val Lys  
 35 40 45  
 Leu Lys Pro Lys Phe Pro Val Arg Ile Asp Thr Gln Asp Ser Glu Ile  
 50 55 60  
 Lys Asp Met Val Glu Glu His Leu Pro Leu Ile Thr Gln Gln Gln Glu  
 65 70 75 80  
 Glu Val Leu Asp Lys Glu Gln Thr Gly Phe Leu Ala Glu Glu Ala Pro  
 85 90 95  
 Asp Asn Val Lys Thr Met Leu Arg Ser Lys Gly Tyr Phe Ser Ser Lys  
 100 105 110  
 Val Ser Leu Thr Glu Lys Asp Gly Ala Tyr Thr Val His Ile Thr Pro  
 115 120 125  
 Gly Pro Arg Thr Lys Ile Ala Asn Val Gly Val Ala Ile Leu Gly Asp  
 130 135 140  
 Ile Leu Ser Asp Gly Asn Leu Ala Glu Tyr Tyr Arg Asn Ala Leu Glu  
 145 150 155 160  
 Asn Trp Gln Gln Pro Val Gly Ser Asp Phe Asp Gln Asp Ser Trp Glu  
 165 170 175  
 Asn Ser Lys Thr Ser Val Leu Gly Ala Val Thr Arg Lys Gly Tyr Pro  
 180 185 190  
 Leu Ala Lys Leu Gly Asn Thr Arg Ala Ala Val Asn Pro Asp Thr Ala  
 195 200 205  
 Thr Val Asp Leu Asn Val Val Val Asp Ser Gly Arg Pro Ile Ala Phe  
 210 215 220  
 Gly Asp Phe Glu Ile Thr Gly Thr Gln Arg Tyr Pro Glu Gln Ile Val  
 225 230 235 240  
 Ser Gly Leu Ala Arg Phe Gln Pro Gly Thr Pro Tyr Asp Leu Asp Leu  
 245 250 255  
 Leu Leu Asp Phe Gln Gln Ala Leu Glu Gln Asn Gly His Tyr Ser Gly  
 260 265 270  
 Ala Ser Val Gln Ala Asp Phe Asp Arg Leu Gln Gly Asp Arg Val Pro  
 275 280 285  
 Val Lys Val Ser Val Thr Glu Val Lys Arg His Lys Leu Glu Thr Gly  
 290 295 300  
 Ile Arg Leu Asp Ser Glu Tyr Gly Leu Gly Gly Lys Ile Ala Tyr Asp  
 305 310 315  
 Tyr Tyr Asn Leu Phe Asn Lys Gly Tyr Ile Gly Ser Val Val Trp Asp  
 325 330 335  
 Met Asp Lys Tyr Glu Thr Thr Leu Ala Ala Gly Ile Ser Gln Pro Arg  
 340 345 350  
 Asn Tyr Arg Gly Asn Tyr Trp Thr Ser Asn Val Ser Tyr Asn Arg Ser  
 355 360 365  
 Thr Thr Gln Asn Leu Glu Lys Arg Ala Phe Ser Gly Gly Ile Trp Tyr  
 370 375 380  
 Val Arg Asp Arg Ala Gly Ile Asp Ala Arg Leu Gly Ala Glu Phe Leu  
 385 390 395 400  
 Ala Glu Gly Arg Lys Ile Pro Gly Ser Asp Ile Asp Leu Gly Asn Ser  
 405 410 415

-continued

---

His Ala Thr Met Leu Thr Ala Ser Trp Lys Arg Gln Leu Leu Asn Asn  
                   420                                  425                                  430

Val Leu His Pro Glu Asn Gly His Tyr Leu Asp Gly Lys Ile Gly Thr  
                   435                                  440                                  445

Thr Leu Gly Thr Phe Leu Ser Ser Thr Ala Leu Ile Arg Thr Ser Ala  
                   450                                  455                                  460

Arg Ala Gly Tyr Phe Phe Thr Pro Glu Asn Lys Lys Leu Gly Thr Phe  
                   465                                  470                                  475                                  480

Ile Ile Arg Gly Gln Ala Gly Tyr Thr Val Ala Arg Asp Asn Ala Asp  
                                   485                                  490                                  495

Val Pro Ser Gly Leu Met Phe Arg Ser Gly Gly Ala Ser Ser Val Arg  
                                   500                                  505                                  510

Gly Tyr Glu Leu Asp Ser Ile Gly Leu Ala Gly Pro Asn Gly Ser Val  
                                   515                                  520                                  525

Leu Pro Glu Arg Ala Leu Leu Val Gly Ser Leu Glu Tyr Gln Leu Pro  
                                   530                                  535                                  540

Phe Thr Arg Thr Leu Ser Gly Ala Val Phe His Asp Met Gly Asp Ala  
                   545                                  550                                  555                                  560

Ala Ala Asn Phe Lys Arg Met Lys Leu Lys His Gly Ser Gly Leu Gly  
                                   565                                  570                                  575

Val Arg Trp Phe Ser Pro Leu Ala Pro Phe Ser Phe Asp Ile Ala Tyr  
                                   580                                  585                                  590

Gly His Ser Asp Lys Lys Ile Arg Trp His Ile Ser Leu Gly Thr Arg  
                   595                                  600                                  605

Phe

<210> SEQ ID NO 5  
 <211> LENGTH: 1764  
 <212> TYPE: DNA  
 <213> ORGANISM: Neisseria meningitidis

<400> SEQUENCE: 5

```

gcccgcgacc tttccgaaaa caaggcggcg ggtttcgcat tgttcaaaaa caaaagcccc   60
gacaccgaat cagtcaaatt aaaacccaaa ttccccgtcc tcatcgacac gcaggacagt   120
gaaatcaaag atatggtcga agaacacctg cgcctcatca cgcagcagca ggaagaagta   180
ttggacaagg aacagacggg cttcctcgcc gaagaagcgc cggacaacgt taaaacgatg   240
ctccgcagca aaggctatct cagcagcaaa gtcagcctga cggaaaaaga cggagcttat   300
acggtacaca tcacaccggg cccgcgcacc aaaatcgcca acgtcggcgt cgccatcctc   360
ggcgacatcc tttcagacgg caacctcgcc gaatactacc gcaacgcgct ggaaaaactg   420
cagcagccgg taggcagcga tttcgatcag gacagttggg aaaacagcaa aacttccgtc   480
ctcggcggcg taacgcgcaa agcctaccgg cttgccaagc tccgcaatac gcaggcggcc   540
gtcaaccccc ataccgccac cgccgatttg aacgtcgtcg tggacagcgg ccgccccatc   600
gccttcggcg actttgaaat caccggcaca cagcgttacc ccgaacaaat cgtctccggc   660
cttgcgcggt tcacgcccgg tatgcggtac gacctcgacc tgctgctcga cttccaacag   720
gcgctcgaac aaaacgggca ttattccggc gcgtccgtac aagccgactt cgaccgcctc   780
caaggcgacc gcgtccccgt caaagtcagc gtaaccgagg tcaaacgccca caaactcgaa   840
accggcatcc gcctcgatcc ggaatacggg ttgggcccga aaatcgccca cgactattac   900
    
```

-continued

---

```

aacctcttca acaaaggcta tatcggttcg gtcgtctggg atatggacaa atacgaaacc 960
acgcttgccg ccggcatcag ccagccgcgc aactatcggg gcaactactg gacaagcaac 1020
gtttcttaca accgttcgac caccctaaac ctcgaaaaac ggccttctc cggcggcgtc 1080
tggtatgtgc gcgaccgcgc gggcatcgat gccaggtcgg gggcggaatt tctcgcagaa 1140
ggccggaaaa tccccggctc ggctgtcgat ttgggcaaca gccacgccac gatgctgacc 1200
gcctcttga aacgccagct gctcaacaac gtgctgcatc ccgaaaacgg ccattacctc 1260
gacggcaaaa tcggtacgac ttggggcaca ttctgtctct ccaccgcgct gatccgcacc 1320
tctgcccgtg caggttattt cttcagccc gaaaacaaaa aactcggcac gttcatcata 1380
cgggacaag cgggttacac cgttccccgc gacaatgccg acgttccttc agggctgatg 1440
ttccgcagcg gcggcgcgct ttccgtgcgc ggttacgaac tcgacagcat cggacttgcc 1500
ggcccgaacg gatcggctct gcccgaaacg gccctcctgg tgggcagcct ggaataccaa 1560
ctgccgttta cgcgcaccct ttccggcgcg gtgttccacg atatgggga tgcgcgcgcc 1620
aatttcaaac gtatgaagct gaaacacggt tcgggactgg gcgtgcgctg gttcagccc 1680
cttgcgccgt tttccttcga catcgctac gggcacagcg ataagaaaat ccgctggcac 1740
atcagcttgg gaacacgctt ctaa 1764

```

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 587

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Neisseria meningitidis

&lt;400&gt; SEQUENCE: 6

```

Ala Ala Asp Leu Ser Glu Asn Lys Ala Ala Gly Phe Ala Leu Phe Lys
 1          5          10          15
Asn Lys Ser Pro Asp Thr Glu Ser Val Lys Leu Lys Pro Lys Phe Pro
 20          25          30
Val Leu Ile Asp Thr Gln Asp Ser Glu Ile Lys Asp Met Val Glu Glu
 35          40          45
His Leu Pro Leu Ile Thr Gln Gln Gln Glu Glu Val Leu Asp Lys Glu
 50          55          60
Gln Thr Gly Phe Leu Ala Glu Glu Ala Pro Asp Asn Val Lys Thr Met
 65          70          75          80
Leu Arg Ser Lys Gly Tyr Phe Ser Ser Lys Val Ser Leu Thr Glu Lys
 85          90          95
Asp Gly Ala Tyr Thr Val His Ile Thr Pro Gly Pro Arg Thr Lys Ile
100          105          110
Ala Asn Val Gly Val Ala Ile Leu Gly Asp Ile Leu Ser Asp Gly Asn
115          120          125
Leu Ala Glu Tyr Tyr Arg Asn Ala Leu Glu Asn Trp Gln Gln Pro Val
130          135          140
Gly Ser Asp Phe Asp Gln Asp Ser Trp Glu Asn Ser Lys Thr Ser Val
145          150          155          160
Leu Gly Ala Val Thr Arg Lys Ala Tyr Pro Leu Ala Lys Leu Gly Asn
165          170          175
Thr Gln Ala Ala Val Asn Pro Asp Thr Ala Thr Ala Asp Leu Asn Val
180          185          190
Val Val Asp Ser Gly Arg Pro Ile Ala Phe Gly Asp Phe Glu Ile Thr
195          200          205

```

-continued

---

Gly Thr Gln Arg Tyr Pro Glu Gln Ile Val Ser Gly Leu Ala Arg Phe  
 210 215 220  
 Gln Pro Gly Met Pro Tyr Asp Leu Asp Leu Leu Leu Asp Phe Gln Gln  
 225 230 235 240  
 Ala Leu Glu Gln Asn Gly His Tyr Ser Gly Ala Ser Val Gln Ala Asp  
 245 250 255  
 Phe Asp Arg Leu Gln Gly Asp Arg Val Pro Val Lys Val Ser Val Thr  
 260 265 270  
 Glu Val Lys Arg His Lys Leu Glu Thr Gly Ile Arg Leu Asp Ser Glu  
 275 280 285  
 Tyr Gly Leu Gly Gly Lys Ile Ala Tyr Asp Tyr Tyr Asn Leu Phe Asn  
 290 295 300  
 Lys Gly Tyr Ile Gly Ser Val Val Trp Asp Met Asp Lys Tyr Glu Thr  
 305 310 315 320  
 Thr Leu Ala Ala Gly Ile Ser Gln Pro Arg Asn Tyr Arg Gly Asn Tyr  
 325 330 335  
 Trp Thr Ser Asn Val Ser Tyr Asn Arg Ser Thr Thr Gln Asn Leu Glu  
 340 345 350  
 Lys Arg Ala Phe Ser Gly Gly Val Trp Tyr Val Arg Asp Arg Ala Gly  
 355 360 365  
 Ile Asp Ala Arg Leu Gly Ala Glu Phe Leu Ala Glu Gly Arg Lys Ile  
 370 375 380  
 Pro Gly Ser Ala Val Asp Leu Gly Asn Ser His Ala Thr Met Leu Thr  
 385 390 395 400  
 Ala Ser Trp Lys Arg Gln Leu Leu Asn Asn Val Leu His Pro Glu Asn  
 405 410 415  
 Gly His Tyr Leu Asp Gly Lys Ile Gly Thr Thr Leu Gly Thr Phe Leu  
 420 425 430  
 Ser Ser Thr Ala Leu Ile Arg Thr Ser Ala Arg Ala Gly Tyr Phe Phe  
 435 440 445  
 Thr Pro Glu Asn Lys Lys Leu Gly Thr Phe Ile Ile Arg Gly Gln Ala  
 450 455 460  
 Gly Tyr Thr Val Ala Arg Asp Asn Ala Asp Val Pro Ser Gly Leu Met  
 465 470 475 480  
 Phe Arg Ser Gly Gly Ala Ser Ser Val Arg Gly Tyr Glu Leu Asp Ser  
 485 490 495  
 Ile Gly Leu Ala Gly Pro Asn Gly Ser Val Leu Pro Glu Arg Ala Leu  
 500 505 510  
 Leu Val Gly Ser Leu Glu Tyr Gln Leu Pro Phe Thr Arg Thr Leu Ser  
 515 520 525  
 Gly Ala Val Phe His Asp Met Gly Asp Ala Ala Ala Asn Phe Lys Arg  
 530 535 540  
 Met Lys Leu Lys His Gly Ser Gly Leu Gly Val Arg Trp Phe Ser Pro  
 545 550 555 560  
 Leu Ala Pro Phe Ser Phe Asp Ile Ala Tyr Gly His Ser Asp Lys Lys  
 565 570 575  
 Ile Arg Trp His Ile Ser Leu Gly Thr Arg Phe  
 580 585

&lt;210&gt; SEQ ID NO 7

&lt;211&gt; LENGTH: 20

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 7

gggccgcaac ctccgaaata                20

<210> SEQ ID NO 8
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 8

cgagccagcc gaggaaacat a                21

<210> SEQ ID NO 9
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 9

catagcacca tggccgccga ctttccga        29

<210> SEQ ID NO 10
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 10

ctagtctaga ttagaagcgt gttcccaagc      30

<210> SEQ ID NO 11
<211> LENGTH: 1000
<212> TYPE: DNA
<213> ORGANISM: Neisseria meningitidis

<400> SEQUENCE: 11

aaacggataa tccatttcac gacggtggaa accgcttcg ccaaatcggc aacgacctgc    60
cgcgtaacgt ccgaaccgtg attccgcaac gccgcgccca aaaccaaaagc ccaagccaaa    120
atgccgatat agttggcatt ggcaatcgcg ttaatcgggt tggcgcaccag gttcatcagc    180
agcgatttca acacttccac aatgccggaa ggccggcggg cggacacatc gcccgcgccc    240
gccaaaaaaa tgtgcgctcg gaaaaccata ccggcgatga cggcggtcag ggetgcggaa    300
aacgtaccaa tgaggtaaag gatgataatc ggctgatat gcgccttgtt gcctttttgg    360
tgctgcgcga ttgtggccgc caccaaaaata aatacaaaa ccggcgcgac cgctttgagc    420
gcgcgcgaaa acaggtgtcc gaacaagcct gccgccaagc ccagttgcgg ggaaaccgaa    480
ccgattacga tgcccaacgc caaacggcg gcaatctgcc tgaccaggt gacgcggccg    540
atcgcatgaa ataaggattt gccgaacgcc ataattcttc cttatgttgt gatatgttaa    600
aaaaatgtgt attttaaaag aaaactcatt ctctgtgttt tttttatfff tcggctgtgt    660
ttaaaggttg cgttgatttg ccctatgcag tgccggacag gctttgcttt atcattcggc    720

```

-continued

gcaacgggtt aatttattga acgaaaataa atttatttaa tctgcctat tttccggcac	780
tattccgaaa cgcagcctgt tttccatag cggattggaa acaaaatacc ttaaaacaag	840
cagatacatt tccggcgggc cgcaacctcc gaaataccgg cggcagtatg cctctgaag	900
tgccccgcc cgtccgaaca acacaaaaac agccgttega aacctgtcc gaacagtgtt	960
agaatcgaaa tctgccacac cgatgcacga caccctgacc	1000

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

3. The polypeptide as claimed in claim 1 comprising the amino acid sequence selected from the group consisting of: SEQ ID NO: 2, SEQ ID NO:4 and SEQ ID NO:6.

4. An isolated polypeptide of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic activity of said immunogenic fragment is substantially the same as the polypeptide of SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6.

6. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 85% identity to the amino acid sequence of SEQ ID NO: 4 or 6 over the entire length of SEQ ID NO: 4 or 6 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

7. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO: 4 or 6 over the entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.

8. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO: 3 or 5 over the entire length of SEQ ID NO: 3 or 5 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

9. The isolated polynucleotide as claimed in any one of claims 6 to 8 in which the identity is at least 95% to SEQ ID NO: 3 or 5.

10. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:4 or SEQ ID NO:6.

11. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3 or SEQ ID NO:5.

12. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:4 or SEQ ID NO:6, obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:3 or SEQ ID NO:5 or a fragment thereof.

13. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2.

14. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1.

15. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.

16. An expression vector or a recombinant live microorganism comprising an isolated polynucleotide according to any one of claims 6-15.

17. A host cell comprising the expression vector of claim 16 or a subcellular fraction or a membrane of said host cell expressing an isolated polypeptide comprising an amino acid sequence that has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:4 or SEQ ID NO:6.

18. A process for producing a polypeptide comprising an amino acid sequence that has at least 85% identity to the amino acid sequence selected from the group consisting of SEQ ID NO:4 or SEQ ID NO:6 comprising culturing a host cell of claim 17 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.

19. A process for expressing a polynucleotide of any one of claims 6-15 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of any one of said polynucleotides.

20. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 5 and a pharmaceutically acceptable carrier.

21. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 6 to 15 and a pharmaceutically acceptable carrier.

22. The vaccine composition according to either one of claims 20 or 21 wherein said composition comprises at least one other *Neisseria meningitidis* antigen.

23. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 5.

24. A method of diagnosing a *Neisseria meningitidis* infection, comprising identifying a polypeptide as claimed in any one of claims 1-5, or an antibody that is immunospecific for said polypeptide, present within a biological sample from an animal suspected of having such an infection.

25. Use of a composition comprising an immunologically effective amount of a polypeptide as claimed in any one of claims 1-5 in the preparation of a medicament for use in generating an immune response in an animal.

26. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one

of claims 6-15 in the preparation of a medicament for use in generating an immune response in an animal.

27. A therapeutic composition useful in treating humans with *Neisseria meningitidis* disease comprising at least one antibody directed against the polypeptide of claims 1-5 and a suitable pharmaceutical carrier.

\* \* \* \* \*