



US 20060229273A1

(19) **United States**

(12) **Patent Application Publication**
Gold et al.

(10) **Pub. No.: US 2006/0229273 A1**

(43) **Pub. Date: Oct. 12, 2006**

(54) **HIGH AFFINITY OLIGONUCLEOTIDE
LIGANDS TO GROWTH FACTORS**

(75) Inventors: **Larry Gold**, Boulder, CO (US);
Nebojsa Janjic, Boulder, CO (US);
Nikos Pagratis, Boulder, CO (US)

Correspondence Address:
SWANSON & BRATSCHUN L.L.C.
1745 SHEA CENTER DRIVE
SUITE 330
HIGHLANDS RANCH, CO 80129 (US)

(73) Assignee: **Gilead Sciences, Inc.**, Foster City, CA
(US)

(21) Appl. No.: **11/444,181**

(22) Filed: **May 31, 2006**

Related U.S. Application Data

(60) Continuation of application No. 10/223,666, filed on Aug. 19, 2002, which is a continuation of application No. 09/818,237, filed on Mar. 27, 2001, now abandoned, which is a division of application No. 08/973,124, filed on May 11, 1998, now Pat. No. 6,207,816, filed as 371 of international application No. PCT/US96/08014, filed on May 30, 1996, which is a continuation-in-part of application No. 08/465,594, filed on Jun. 5, 1995, now Pat. No. 5,846,713, and which is a continuation-in-part of application No.

08/465,591, filed on Jun. 5, 1995, now Pat. No. 5,837,834, and which is a continuation-in-part of application No. 08/479,725, filed on Jun. 7, 1995, now Pat. No. 5,674,685, and which is a continuation-in-part of application No. 08/479,783, filed on Jun. 7, 1995, now Pat. No. 5,668,264, and which is a continuation-in-part of application No. 08/618,693, filed on Mar. 20, 1996, now Pat. No. 5,723,594, and which is a continuation-in-part of application No. 08/458,423, filed on Jun. 2, 1995, now Pat. No. 5,731,144, and which is a continuation-in-part of application No. 08/458,424, filed on Jun. 2, 1995, now Pat. No. 5,731,424.

Publication Classification

(51) **Int. Cl.**
A61K 48/00 (2006.01)
C12N 15/87 (2006.01)
(52) **U.S. Cl.** **514/44; 435/455**

(57) **ABSTRACT**

Methods are described for the identification and preparation of high-affinity nucleic acid ligands to TGFβ, PDGF and hKGF. Included in the invention are specific RNA and ssDNA ligands to TGFβ1 and PDGF identified by the SELEX method. Also included in the invention are specific RNA ligands to hKGF identified by the SELEX method. Further included are RNA ligands that inhibit the interaction of TGFβ1 and hKGF with their receptors and DNA ligands that inhibit the interaction of PDGF with its receptor.




FIGURE 1






FIGURE 2

Helix II




Helix III


Helix I

SEQ ID NO: 171


FIGURE 3



SEQ ID NO: 173



SEQ ID NO: 172



SEQ ID NO: 174

FIGURE 4




Figure 5A




Figure 5B




Figure 5C




FIGURE 6




FIGURE 7






FIGURE 8



$K_d=0.065$ nM

SEQ ID NO: 175



$K_d=0.097$ nM

SEQ ID NO: 176

FIGURE 9




FIGURE 10A




FIGURE 10B




FIGURE 11

Class 1 SEQ ID	Kd, nM	Ki, nM	Clone NO:	Seq	Seq	Seq	Seq	Seq	Seq	Seq	Seq	Seq	Seq	Seq	Seq
231	0.001	3.3	*14F	gggaga	acggaugcggUGG	UCUCCC	AAUUCUAAACUUUCU	CCAUCGUAUCUGGG							
223	0.05	1.3	*6F	gggagaga	cgaugcggG	AUCUUUU	GUGGGUCUUUGUAGCCCC	CUCGUUGUCCCCC							
232	0.07	6.7	*15F	gggagagacg	augcggUCAUGG	UGUCUUUCC	ACAGCUCUUC	CCAUGAUCGCCCCG							
259	0.07	0.3	*56F	gggagagacg	augcggGGACAAYAGCGGUGUCUUUUC	CCcagac	AUUUNKAUC	CUCCGACRUCCCag							
240	0.23			AACAGUCUG	GCAAUGGCCAUUAUGGCGG	VACUGCAUU	gacuc	gccccg							
246	0.46	6.7	37F	acgagucgg	GCCCGAUC	UACUGCAUU	ACCGAAAC	GAUUUCCCCACUGU							
233	0.83		*16F	gggag	gacgaugcggGAA	UUCCC	AGCGCUUGACUGAUAACAACN	UUCCCUGCCCCcag							
249	0.9		42F	gggagag	acgaugcggCGACACAUAAGCCUUU	UUCCC	GAUCCCCAUGAUGGCCUCG	CCGUGCCCcagacga							
244	1.0		31F	gggagagag	acggaugcggACGGAGAUG	UUCCC	CUCAAMAUUUAC	CGUCUCGUAUUGCG							
245	1.1		35F	gggagag	acgaugcggCGAAU	UAGCUUCUU	AUGACUCAC	GUUUCCUUUGCCGCGC							
257	1.23		54F	gggagag	acgaugcgg	AAGAUCUCCC	AACUGCUGGCUAAUAUUUCUCU	CCGCGUCCcagac							
255	1.27		51F	gggagag	acgaugcggCAG	UCUCCC	UAGUUUCAU	CUCUGCAGCAUUCC							
220	1.77		2F	gggagagacga	ugcggGCAUUCGUC	AAUACCUGUUUAUUCCUUUUUCUAGCGGCCca	UAGAUUACAGCACUU	gacgacucggccga							
239	2.02		24F	gggagag	acgaugcggUGG	UCUCCC	ACGGGCCCCGAUCCUGCCAG	CCAUCGCAUUGGGC							
262	2.1		60F	gggagag	acgaugcgg	UGACAUGGAAUUUUUCU	CUCUACUGUUA	CCGUGUGcagacga							
238	2.52		23F	gggagagag	cgaugcgg	CCCUACCACUUUUUCC	GAUAACCUUCUCUGUGCCcagacgcccga	UCCUGUC							
222	2.53		5F	gggagagacg	augcggCGUCCCGAGUC	ACGCUGUCUU	ACUAGCUA	UCCUA CCCUGC cag							
225	2.63		8F	gggagagagacg	acgaugcggUUUCGAAUAGGG	CCAUUUCUC	AACUAAGUCCCUCAUUUGCCUCGC	UCCUGUGCCcaga							
243	3.24		29F	gggagag	acgaugcggGGA	CUUC	GCCAGCUCUCCCAAGCGCGAC	CUGUGCCCcagacga							
224	3.69		7F	gggagag	acgaugcgg	CGGUACUCUUC		acgacucggcccga							
252	4.7		45F	gggagagacga	ugcggGUCCU	uYucYy	RRYuy	RRYyS							
268			Consensus	gGgagg	Sgsgg										

Summary Structure




FIGURE 12A

Class 2 SEQ ID	Kd, nM	Ki, nM	NO: Clone
230	*13F 0.03	10.0	
254	*50F 0.12		
247	38F 0.33	0.2	
227	10F 0.47	10.0	
226	9F 0.83		
256	53F 0.7		
253	49F 1.0		
250	43F 1.13	16.7	
260	57F 1.2		
248	41F 1.44		
241	27F 1.52		
235	20F 2.05		
258	55F 2.52		
261	58F 2.52		
221	3F 4.47		
236	21F ND		
242	28F ND		
269	Consensus		

ggacgaugcga	CACCUUAGACCUGUCCUCCNAGCCGUGAGUUG	CGGCGU	CUGUGCCcaga
cgaugccgUGA	UCAAUC	NCAGUGUA	UUACUCUUGCGCUCACC
ggaggaacgaug	cggNGACUGAUUUUCCUUG	UGGCCGA	AUUUCCUG
ggacgaugcga	acgaugcggGACCAGA	ACUAGC	UUUUUCAGCAAUCCUCCCGCUGCCcag
gaggaacgaugc	AUAAUGGCUAGA	AGCACUCA	UCGCAUC
ggacgaugcga	gacgaugcggAUCAAA	UUJACAG	UUGCC
ggaggaacgaugc	gaggacu	augcggUGUAG	GCAUUGAUU
ggacgaugcga	gacg	GGCUGGAU	UUUCC
ggaggaacgaug	UGACUAUCU	GACAGG	CCAAUCACC
ggaggaacgaug	cggGGACUUU	UGGAUCUC	CAUUGAUUUGA
ggaggaacgaug	cggCAGUC	AGUCAG	UAUUUGA
ggaggaacgaug	UUCCAGCCUGGAUUCAU	GGCAUAAACUGCGAUA	DGCUUCUGGCGCagacgacgccc
ggaggaacgaug	gacgaugcggUCCGUAUAAC	UGCCCGUGAUUAAACCAUGCAGGCUUCC	UGCCcagacg
ggaggaacgaug	acgaugcggGACUAA	GCCACUACUA	UGCGcagacg
ggaggaacgaug	ggaggaacgaugcggAUCGUAUC	GCAGGAUUAUCC	GUGCCcagacg
ggaggaacgaug	augcggUCCUA	UUCGGCAAGUGAAAAUA	CGUUGCCCCca
ggaggaacgaug	cgaugcggGAUUCU	rsBrs	GCUJCCCGAGC
rgGrrg	RaYY		YyyYyb




FIGURE 12B

Class 1

SEQ ID NO:	Clone	K _d , nM	K _i , nM	SEQ
195	14N	0.4	2.0	UCCAGCGGAU <u>UGAACUGUCGGGUAGGAACA</u>
199	25N	0.9		<u>augcggga</u> <u>CGGAUUG</u> AGGAA A
191	4N	0.7	23.3	<u>augcggguggug</u> <u>AAGAGG</u> UACCGGAA UUGC
189	1N	0.5	16.7	<u>gGAAGGG</u> <u>ACGAUAAAG</u> AGGAA UCGAA
205	36N	8.27		<u>gga</u> <u>ggacgaugcggg</u> <u>AAGGAACA</u> GGA
203	34N	0.8		<u>augcgggCGGG</u> <u>AAAGACU</u> AAUGGAA GUGGAU
207	42N	0.8	266.7	<u>augcgggGAG</u> <u>ACGU</u> <u>AGACGGGAACAUAGAACGAUCAAC</u>
196	16N	1.4		<u>cgaugcggAAAGUUCUAACAAGUUAGUGAA</u> GGUUC CA CU
270	Consensus			<u>GGG</u> <u>aa</u> Yaa RCRR

GCGGCca	<u>gacgac</u>	<u>ucgcccga</u>
GCAGAUACCGAGCUGGC	<u>cgagacg</u>	<u>cucgcccga</u>
UAAAGAUACCCAGG	<u>CCcagac</u>	<u>gacucgcccga</u>
GUGGCUGGCcaga	<u>cgac</u>	<u>ucgcccga</u>
ACAAGUCGAACAAAG	<u>CCGAGGUGcaga</u>	<u>cgacucgcccga</u>
ACAGUGC	<u>GGcagac</u>	<u>gacucgcccga</u>
GCGGCcag	<u>acgacuc</u>	<u>gcccga</u>
UGAA	<u>UGUAcaga</u>	<u>cgacucgcccga</u>
RCRR	<u>gA</u>	<u>CCC</u>

Summary Structure




FIGURE 12C

Class 2	SEQ ID NO: Clone	Kd, Ki, nM							
	209 47N	1.8		ACGAAGGUCAAGAGAAA	CAGC	CCUU	CGGG	GUGcagacga	<u>cucgcccga</u>
	198 24N	1.2		<u>gacgaugcgg</u>	GUGGAAGA	CCGGUCG	GCAU	AAUGUGAC	<u>ACUCCGGca</u>
	202 29N	0.43	13.3	AGAAAGAUCCA	GGAACAGCGAAA	UGCGU	GUGG	Ccagacgac	<u>ucgcccga</u>
	204 35N	2.3		<u>ugcggcuiag</u>	GGAAA	CUGAG	GUGG	Ucagacgac	<u>ucgcccga</u>
	192 6N	0.7	26.7	<u>accguggggc</u>	AGGGAGCAA	UGAACUCAAGUCAAGCCGGUGCACGU	GGGc	agacgacuc	<u>giccccga</u>
	190 2N	0.8	66.7			ggggaggacgaugcggCGGGAAAGGUC		CGAA	<u>GACC</u>
	211 54N	5.3		<u>GCCGA</u>	AAGGAA	CGAGA	UUGCCcagacgacucgcccga		<u>AGCCCCUUGGUGGU</u>
	271	Consensus							ygay

Summary Structure




FIGURE 12D





FIGURE 13

HIGH AFFINITY OLIGONUCLEOTIDE LIGANDS TO GROWTH FACTORS

RELATEDNESS OF THE APPLICATION

[0001] This application is a continuation of U.S. application Ser. No. 10/223,666, filed Aug. 19, 2002, which is a continuation of U.S. application Ser. No. 09/818,237, filed Mar. 27, 2001, now abandoned, which is a divisional of U.S. application Ser. No. 08/973,124, filed May 11, 1998, now U.S. Pat. No. 6,207,816, which is a 35 U.S.C. §§371 national phase application of International Application No. PCT/US96/08014, filed May 30, 1996. International Application No. PCT/US96/08014 is a continuation-in-part of U.S. application Ser. No. 08/465,594, filed Jun. 5, 1995, now U.S. Pat. No. 5,846,713, a continuation-in-part of Ser. No. 08/465,591, filed Jun. 5, 1995, now U.S. Pat. No. 5,837,834, a continuation-in-part of U.S. application Ser. No. 08/479,725, filed Jun. 7, 1995, now U.S. Pat. No. 5,674,685, a continuation-in-part of U.S. application Ser. No. 08/479,783, filed Jun. 7, 1995, now U.S. Pat. No. 5,668,264, and a continuation-in-part of U.S. application Ser. No. 08/618,693, filed Mar. 20, 1996, now U.S. Pat. No. 5,723,594. PCT/US96/08014 also claims priority from U.S. application Ser. No. 08/458,423, filed Jun. 2, 1995, now U.S. Pat. No. 5,731,144, and U.S. application Ser. No. 08/458,424, filed Jun. 2, 1995, now U.S. Pat. No. 5,731,424. Each of these references is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] Described herein are methods for identifying and preparing high-affinity nucleic acid ligands to TGF β , PDGF, and hKGF. The method utilized herein for identifying such nucleic acid ligands is called SELEX, an acronym for Systematic Evolution of Ligands by EXponential enrichment. This invention includes high affinity nucleic acid ligands of TGF β , PDGF and hKGF. Further disclosed are RNA and DNA ligands to TGF β 1 and PDGF and RNA ligands to hKGF. Also included are oligonucleotides containing nucleotide derivatives chemically modified at the 2'-positions of pyrimidines. Additionally disclosed are RNA ligands to TGF β 1 and hKGF containing 2'-NH₂-modifications or 2'-F modifications and RNA ligands to PDGF containing 2'-F modifications. This invention also includes high affinity nucleic acid inhibitors of TGF β 1, PDGF and hKGF. The oligonucleotides of the present invention are useful as pharmaceuticals or diagnostic agents.

BACKGROUND OF THE INVENTION

TGF β

[0003] The transforming growth factor- β (TGF β) polypeptides influence growth, differentiation, and gene expression in many cell types. The first polypeptide of this family that was characterized, TGF β 1 has two identical 112 amino acid subunits which are covalently linked. TGF β 1 is a highly conserved protein with only a single amino acid difference distinguishing human from mice forms. There are two other members of the TGF β gene family that are expressed in mammals. TGF β 2 is 71% homologous to TGF β 1 (de Martin et al., (1987) *EMBO J.* 6:3673-3677), whereas TGF β 3 is 80% homologous to TGF β 1 (Derynck et al., (1988) *EMBO J.* 7:3737-3743). The structural characteristics of TGF β 1 as determined by nuclear magnetic reso-

nance (Archer et al., (1993) *Biochemistry* 32:1164-1171) agree with the crystal structure of TGF β 2 (Daopin et al., (1992) *Science* 257:369-374; Schlunegger and Grutter (1992) *Nature* 358:430-434).

[0004] Even though the TGF β 's have similar three dimensional structures, they are by no means physiologically equivalent. There are at least three different extracellular receptors, type I, II and III, involved in transmembrane signaling of TGF β to cells carrying the receptors. For reviews, see Derynck (1994) *TIBS* 19:548-553 and Massague (1990) *Annu. Rev. Cell Biol.* 6:597-641. In order for TGF β 2 to effectively interact with the type II TGF β receptor, the type III receptor must also be present (Derynck (1994) *TIBS* 19:548-553). Vascular endothelial cells lack the type III receptor. Instead endothelial cells express a structurally related protein called endoglin (Cheifetz et al., (1992) *J. Biol. Chem.* 267:19027-19030), which only binds TGF β 1 and TGF β 3 with high affinity. Thus, the relative potency of the TGF β 's reflect the type of receptors expressed in a cell and organ system.

[0005] In addition to the regulation of the components in the multifactorial signaling pathway, the distribution of the synthesis of TGF β polypeptides also affects physiological function. The distribution of TGF β 2 and TGF β 3 is more limited (Derynck et al., (1988) *EMBO J.* 7:3737-3743) than TGF β 1, e.g., TGF β 3 is limited to tissues of mesenchymal origin, whereas TGF β 1 is present in both mesenchymal and epithelial cells.

[0006] TGF β 1 is a multifunctional cytokine critical for tissue repair. High concentrations of TGF β 1 are delivered to the site of injury by platelet granules (Assoian and Sporn, (1986) *J Cell Biol.* 102:1217-1223.). TGF β 1 initiates a series of events that promote healing including chemotaxis of cells such as leukocytes, monocytes and fibroblasts, and regulation of growth factors and cytokines involved in angiogenesis, cell division associated with tissue repair and inflammatory responses. TGF β 1 also stimulates the synthesis of extracellular matrix components (Roberts et al., (1986) *Proc. Natl. Acad. Sci USA* 83:4167-4171; Sporn et al., (1983) *Science* 219:1329-1330; Massague, (1987) *Cell* 49:437-438) and most importantly for understanding the pathophysiology of TGF β 1, TGF β 1 autoregulates its own synthesis (Kim et al., (1989) *J Biol Chem* 264:7041-7045).

[0007] A number of diseases have been associated with TGF β 1 overproduction. Fibrotic diseases associated with TGF β 1 overproduction can be divided into chronic conditions such as fibrosis of kidney, lung and liver and more acute conditions such as dermal scarring and restenosis. Synthesis and secretion of TGF β 1 by tumor cells can also lead to immune suppression such as seen in patients with aggressive brain or breast tumors (Arteaga et al., (1993) *J Clin Invest* 92: 2569-2576). The course of Leishmanial infection in mice is drastically altered by TGF β 1 (Barral-Netto et al., (1992) *Science* 257:545-547). TGF β 1 exacerbated the disease, whereas TGF β 1 antibodies halted the progression of the disease in genetically susceptible mice. Genetically resistant mice became susceptible to Leishmanial infection upon administration of TGF β 1.

[0008] The profound effects of TGF β 1 on extracellular matrix deposition have been reviewed (Rocco and Ziyadeh, (1991) in *Contemporary Issues in Nephrology* v23, Hormones, autocooids and the kidney. ed. Jay Stein, Churchill

Livingston, N.Y. pp 391-410; Roberts et al., (1988) *Rec. Prog. Hormone Res.* 44:157-197) and include the stimulation of the synthesis and the inhibition of degradation of extracellular matrix components. Since the structure and filtration properties of the glomerulus are largely determined by the extracellular matrix composition of the mesangium and glomerular membrane, it is not surprising that TGF β 1 has profound effects on the kidney. The accumulation of mesangial matrix in proliferative glomerulonephritis (Border et al., (1990) *Kidney Int.* 37:689-695) and diabetic nephropathy (Mauer et al., (1984) *J. Clin Invest.* 74:1143-1155) are clear and dominant pathological features of the diseases. TGF β 1 levels are elevated in human diabetic glomerulosclerosis (advanced neuropathy) (Yamamoto et al., (1993) *Proc. Natl. Acad. Sci.* 90:1814-1818). TGF β 1 is an important mediator in the genesis of renal fibrosis in a number of animal models (Phan et al., (1990) *Kidney Int.* 37:426; Okuda et al., (1990) *J. Clin Invest.* 86:453). Suppression of experimentally induced glomerulonephritis in rats has been demonstrated by antiserum against TGF β 1 (Border et al., (1990) *Nature* 346:371) and by an extracellular matrix protein, decorin, which can bind TGF β 1 (Border et al., (1992) *Nature* 360:361-363).

[0009] Too much TGF β 1 leads to dermal scar-tissue formation. Neutralizing TGF β 1 antibodies injected into the margins of healing wounds in rats have been shown to inhibit scarring without interfering with the rate of wound healing or the tensile strength of the wound (Shah et al., (1992) *Lancet* 339:213-214). At the same time there was reduced angiogenesis, reduced number of macrophages and monocytes in the wound, and a reduced amount of disorganized collagen fiber deposition in the scar tissue.

[0010] TGF β 1 may be a factor in the progressive thickening of the arterial wall which results from the proliferation of smooth muscle cells and deposition of extracellular matrix in the artery after balloon angioplasty. The diameter of the restenosed artery may be reduced 90% by this thickening, and since most of the reduction in diameter is due to extracellular matrix rather than smooth muscle cell bodies, it may be possible to open these vessels to 50% simply by reducing extensive extracellular matrix deposition. In uninjured pig arteries transfected in vivo with a TGF β 1 gene, TGF β 1 gene expression was associated with both extracellular matrix synthesis and hyperplasia (Nabel et al., (1993) *Proc. Natl. Acad. Sci USA* 90:10759-10763). The TGF β 1 induced hyperplasia was not as extensive as that induced with PDGF-BB, but the extracellular matrix was more extensive with TGF β 1 transfectants. No extracellular matrix deposition was associated with FGF-1 (a secreted form of FGF) induced hyperplasia in this gene transfer pig model (Nabel (1993) *Nature* 362:844-846).

[0011] There are several types of cancer where TGF β 1 produced by the tumor may be deleterious. MATLyLu rat cancer cells (Steiner and Barrack, (1992) *Mol. Endocrinol.* 6:15-25) and MCF-7 human breast cancer cells (Arteaga et al., (1993) *Cell Growth and Differ.* 4:193-201) became more tumorigenic and metastatic after transfection with a vector expressing the mouse TGF β 1. In breast cancer, poor prognosis is associated with elevated TGF β (Dickson et al., (1987) *Proc. Natl. Acad. Sci. USA* 84:837-841; Kasid et al., (1987) *Cancer Res.* 47:5733-5738; Daly et al., (1990) *J Cell Biochem* 43:199-211; Barrett-Lee et al., (1990) *Br. J Cancer* 61:612-617; King et al., (1989) *J Steroid Biochem* 34:133-

138; Welch et al., (1990) *Proc. Natl. Acad. Sci.* 87:7678-7682; Walker et al., (1992) *Eur J Cancer* 238: 641-644) and induction of TGF β 1 by tamoxifen treatment (Butta et al., (1992) *Cancer Res* 52:4261-4264) has been associated with failure of tamoxifen treatment for breast cancer (Thompson et al., (1991) *Br. J Cancer* 63:609-614). Anti TGF β 1 antibodies inhibit the growth of MDA-231 human breast cancer cells in athymic mice (Arteaga et al., (1993) *J Clin Invest* 92: 2569-2576), a treatment which is correlated with an increase in spleen natural killer cell activity. CHO cells transfected with latent TGF β 1 also showed decreased NK activity and increased tumor growth in nude mice (Wallick et al., (1990) *J Exp Med* 172:1777-1784). Thus, TGF β 1 secreted by breast tumors may cause an endocrine immune suppression.

[0012] High plasma concentrations of TGF β 1 have been shown to indicate poor prognosis for advanced breast cancer patients (Anscher et al. (1993) *N Engl J Med* 328:1592-8). Patients with high circulating TGF β before high dose chemotherapy and autologous bone marrow transplantation are at high risk for hepatic veno-occlusive disease (15-50% of all patients with a mortality rate up to 50%) and idiopathic interstitial pneumonitis (40-60% of all patients). The implication of these findings is 1) that elevated plasma levels of TGF β 1 can be used to identify at risk patients and 2) that reduction of TGF β 1 could decrease the morbidity and mortality of these common treatments for breast cancer patients.

PDGF

[0013] Platelet-derived growth factor (PDGF) was originally isolated from platelet lysates and identified as the major growth-promoting activity present in serum but not in plasma. Two homologous PDGF isoforms have been identified, PDGF A and B, which are encoded by separate genes (on chromosomes 7 and 22). The most abundant species from platelets is the AB heterodimer, although all three possible dimers (AA, AB and BB) occur naturally. Following translation, PDGF dimers are processed into \approx 30 kDa secreted proteins. Two cell surface proteins that bind PDGF with high affinity have been identified, α and β (Heldin et al., (1981) *Proc. Natl. Acad. Sci.* 78:3664; Williams et al., (1981) *Proc. Natl. Acad. Sci.* 79:5867). Both species contain five immunoglobulin-like extracellular domains, a single transmembrane domain and an intracellular tyrosine kinase domain separated by a kinase insert domain. The functional high affinity receptor is a dimer and engagement of the extracellular domain of the receptor by PDGF results in cross-phosphorylation (one receptor tyrosine kinase phosphorylates the other in the dimer) of several tyrosine residues. Receptor phosphorylation leads to a cascade of events that results in the transduction of the mitogenic or chemotactic signal to the nucleus. For example, in the intracellular domain of the PDGF β receptor, nine tyrosine residues have been identified that when phosphorylated interact with different src-homology 2 (SH2) domain-containing proteins including phospholipase C-g, phosphatidylinositol 3'-kinase, GTPase-activating protein and several adapter molecules like Shc, Grb2 and Nck (Heldin, (1995) *Cell* 80:213). In the last several years, the specificities of the three PDGF isoforms for the three receptor dimers ($\alpha\alpha$, $\alpha\beta$, and $\beta\beta$) has been elucidated. The α -receptor homodimer binds all three PDGF isoforms with high affinity, the β -receptor homodimer binds only PDGF BB with high affinity and PDGF AB with approximately 10-fold lower affinity, and the $\alpha\beta$ -receptor heterodimer binds PDGF BB and PDGF AB with high

affinity (Westermarck & Heldin, (1993) *Acta Oncologica* 32:101). The specificity pattern results from the ability of the A-chain to bind only to the α -receptor and of the B-chain to bind to both α and β -receptor subunits with high affinity.

[0014] The earliest indication that PDGF expression is linked to malignant transformation came with the finding that the amino acid sequence of the PDGF-B chain is virtually identical to that of p28^{sis}, the transforming protein of the simian sarcoma virus (SSV) (Waterfield et al., (1983) *Nature* 304:35; Johnsson et al., (1984) *EMBO J.* 3:921). The transforming potential of the PDGF-B chain gene and, to a lesser extent, the PDGF-A gene was demonstrated soon thereafter (Clarke et al., *Nature* 308:464 (1984); Gazit et al., *Cell* 39:89 (1984); Beckmann et al., *Science* 241:1346; Bywater et al., *Mol. Cell. Biol.* 8:2753 (1988)). Many tumor cell lines have since been shown to produce and secrete PDGF, some of which also express PDGF receptors (Raines et al., *Peptide Growth Factors and Their Receptors*, Springer-Verlag, Part I, p 173 (1990)). Paracrine and, in some cell lines, autocrine growth stimulation by PDGF is therefore possible. For example, analysis of biopsies from human gliomas has revealed the existence of two autocrine loops: PDGF-B/ β -receptor in tumor-associated endothelial cells and PDGF-A/ α -receptor in tumor cells (Hermansson et al., *Proc. Natl. Acad. Sci.* 85:7748 (1988); Hermansson et al., *Cancer Res.* 52:3213 (1992)). The progression to high grade glioma was accompanied by the increase in expression of PDGF-B and the β -receptor in tumor-associated endothelial cells and PDGF-A in glioma cells. Increased expression of PDGF and/or PDGF receptors has also been observed in other malignancies including fibrosarcoma (Smits et al., *Am. J. Pathol.* 140:639 (1992)) and thyroid carcinoma (Heldin et al., *Endocrinology* 129:2187 (1991)).

[0015] In view of its importance in proliferative disease states, antagonists of PDGF may find useful clinical applications. Currently, antibodies to PDGF (Johnsson et al., (1985) *Proc. Natl. Acad. Sci., U. S. A.* 82:1721-1725; Ferns et al., (1991) *Science* 253:1129-1132; Herrenet al. (1993) *Biochimica et Biophysica Acta* 1173:204-302) and the soluble PDGF receptors (Herrenet al., (1993) *Biochimica et Biophysica Acta* 1173:204-302; Duanet al., (1991) *J. Biol. Chem.* 266:413-418; Tiesman et al., (1993) *J. Biol. Chem.* 268:9621-9628) are the most potent and specific antagonists of PDGF. Neutralizing antibodies to PDGF have been shown to revert the SSV-transformed phenotype (Johnsson et al., (1985) *Proc. Natl. Acad. Sci., U. S. A.* 82:1721-1725) and to inhibit the development of neointimal lesions following arterial injury (Ferns et al., (1991) *Science* 253:1129-1132). Other inhibitors of PDGF such as suramin (Williams et al., (1984) *J. Biol. Chem.* 259:5287-5294; Betsholtz et al., (1984) *Cell* 39:447-457), neomycin (Vassbotn et al., (1992) *J. Biol. Chem.* 267:15635-15641) and peptides derived from the PDGF amino acid sequence (Engström et al., (1992) *J. Biol. Chem.* 267:16581-16587) have been reported, however, they are either too toxic or lack sufficient specificity or potency to be good drug candidates. Other types of antagonists of possible clinical utility are molecules that selectively inhibit the PDGF receptor tyrosine kinase (Buchdunger et al., (1995) *Proc. Natl. Acad. Sci., U. S. A.* 92:2558-2562; Kovalenko et al., (1994) *Cancer Res.* 54:6106-6114).

hKGF

a) Biochemical Properties of hKGF

[0016] Human Keratinocyte Growth Factor (hKGF) is a small (26-28KD) basic heparin-binding growth factor and a member of the FGF family. hKGF is a relatively newly identified molecule, which is also known as FGF-7 (Finch et al., (1989) *Science* 245:752-755). It is a growth factor specific for epithelial cells (Rubin et al., (1989) *Proc Natl Acad Sci USA* 86:802-806), and its main function is in development/morphogenesis (Werner et al., (1994) *Science* 266:819-822) and in wound healing (Werner et al., (1992) *Proc Natl Acad Sci USA* 89:6896-6900). The major in vivo source of hKGF is stromal fibroblasts (Finch et al., (1989) *Science* 245:752-755). Microvascular endothelial cells (Smola et al., (1993) *J Cell Biol* 122:417-429) and very recently, activated intraepithelial gd T cells (Boismenu et al., (1994) *Science* 266:1253-1255) have also been shown to synthesize hKGF. hKGF expression is stimulated in wounds (Werner et al., (1992) *Proc Natl Acad Sci USA* 89:6896-6900). Several cytokines are shown to be hKGF inducers (Brauchle et al., (1994) *Oncogene* 9:3199-3204), with IL-1 the most potent one (Brauchle et al., (1994) *Oncogene* 9:3199-3204; Chedid et al., (1994) *J Biol Chem* 269:10753-10757). Unlike bFGF, hKGF has a signal peptide and thus is secreted by producing cells (Finch et al., (1989) *Science* 245:752-755). hKGF can be overexpressed in *E. coli* and the recombinant protein (~19-21 KD) is biologically active (Ron et al., (1993) *J Biol Chem* 268:2984-2988). The *E. coli* derived recombinant protein is 10 times more mitogenic than the native protein (Ron et al., (1993) *J Biol Chem* 268:2984-2988). This difference may be due to glycosylation. The native protein has a potential Asn glycosylation site (Ron et al., (1993) *J Biol Chem* 268:2984-2988).

[0017] The hKGF bioactivity is mediated through a specific cell surface receptor (Miki et al., (1991) *Science* 251:72-75). The hKGF receptor is a modified FGF receptor resulting from alternative splicing of the C-terminal extracellular region of the FGF-R2 (Miki et al., (1992) *Proc Natl Acad Sci USA* 89:246-250). NIH/3T3 cells transfected with the hKGF receptor express high affinity (~200 pM) binding sites for hKGF (Miki et al., (1992) *Proc Natl Acad Sci USA* 89:246-250). The approximate number of specific binding sites per NIH/3T3 cell is about 500,000 (D. Bottaro and S. Aaronson, personal communication). The hKGF receptor binds hKGF and aFGF with similar affinities, and bFGF with about 20 fold less affinity (Miki et al., (1991) *Science* 251:72-75; Miki et al., (1992) *Proc Natl Acad Sci USA* 89:246-250). A variant of the hKGF receptor has been found to be an amplified gene (i.e., one gene, multiple copies), designated K-SAM, in a human stomach carcinoma cell line (Hattori et al., (1990) *Proc natl Acad Sci USA* 87:5983-5987).

[0018] Heparin has been reported to be an inhibitor of hKGF bioactivity (Ron et al., (1993) *J Biol Chem* 268:2984-2988). This is in contrast to the agonistic effect of heparin for aFGF (Spivak-Kroixman et al., (1994) *Cell* 79:1015-1024).

b) Role of hKGF in Human Disease

[0019] The recombinant hKGF molecule has been available only since 1993. Therefore, there is limited information on the role of hKGF in human disease. The published literature, however, contains evidence that strongly suggests

a role for hKGF in at least two human diseases, namely psoriasis and cancer. hKGF has also been implicated in inflammatory bowel disease (P. Finch, personal communication).

[0020] Psoriasis

[0021] Psoriasis is a skin disorder which can be debilitating (Greaves et al., (1995) *N Eng J Medicine* 332:581-588), characterized by hyperproliferation of the epidermis and incomplete differentiation of keratinocytes, together with dermal inflammation (Abel et al., (1994) *Scientific American Medicine* III-1 to III-18; Greaves et al., (1995) *N Eng J Medicine* 332:581-588). There is not yet an effective treatment for psoriasis (Anonymous, (1993) Drug & Market Development 4:89-101; Abel et al., (1994) *Scientific American Medicine* III-1 to III-18; Greaves et al., (1995) *N Eng J Medicine* 332:581-588). Psoriasis occurs in 0.5 to 2.8 percent of the population with the highest incidence in Scandinavia. In the US in 1992, it was estimated that 4-8 million people affected with psoriasis spent about \$600 million for various drugs and related therapies, none of which is very effective. Most of the expenditure was made by about 400,000 patients with severe psoriasis spending \$1,000-1,500 annually on treatment. There are about 200,000 new cases of psoriasis every year.

[0022] The basic cause of the disorder is not known, but it results from a primary or secondary defect in the mechanisms that regulate epidermal keratinocyte cell division (Abel et al., (1994) *Scientific American Medicine* III-1 to III-18). Psoriasis responds to steroids and cyclosporine and in that sense is characterized as an immune disease (Abel et al., (1994) *Scientific American Medicine* III-1 to III-18). Since hKGF is the primary specific growth factor for keratinocytes, its overexpression and deregulation are primary candidates as the cause of keratinocyte hyperproliferation in psoriasis. The demonstration that the immune system is a prime regulator of hKGF release (Boismenu et al., (1994) *Science* 266:1253-1255; Brauchle et al., (1994) *Oncogene* 9:3199-3204; Chedid et al., (1994) *J Biol Chem* 269:10753-10757) strengthens the notion that hKGF deregulation is the cause of psoriasis. Furthermore, application of hKGF in porcine wounds creates a histological appearance resembling psoriasis (Staiano-Coico et al., (1993) *J Ex Med* 178:865-878); keratinocyte derived hKGF in transgenic mice causes pathology reminiscent to psoriasis (Guo et al., (1993) *EMBO J* 12:973-986); in situ hybridization experiments demonstrated a moderate and a strong upregulation of hKGF and hKGF receptors respectively in psoriasis (P. Finch, personal communication). In situ hybridization experiments also demonstrated involvement of hKGF in another immune disease namely, inflammatory bowel disease (P. Finch, personal communication).

[0023] Cancer

[0024] It is well established in the literature that deregulation of the expression of growth factors and growth factor hKGF and/or its receptor is expected to be the transformation event in some human cancers. The transforming ability of the hKGF system has been demonstrated in vitro (Miki et al., (1991) *Science* 251:72-75). In another study, carcinoma cell-lines have been found to express the hKGF receptor and to respond to hKGF but not to aFGF, while sarcoma cell-lines do not express hKGF receptors and respond to aFGF but not to hKGF (Ishii et al., (1994) *Cancer Res* 54:518-522).

[0025] Gastrointestinal Cancer

[0026] Several poorly differentiated stomach cancers have an amplified gene, designated K-sam, which is an isoform of the hKGF-receptor (Kato et al., (1992) *Proc Natl Acad Sci USA* 89:2960-2964). In vivo administration of hKGF to rats causes proliferation of pancreatic ductal epithelial cell (Yi et al., (1994) *Am J Pathol* 145:80-85), hepatocytes, and epithelial cells throughout the gastrointestinal tract (Housley et al., (1994) *J Clin Invest* 94:1764-1777).

[0027] Lung Cancer

[0028] Administration of hKGF to rats causes type II pneumocyte hyperplasia similar to the bronchoalveolar cell variant of lung carcinoma (Ulich et al., (1994) *J Clin Invest* 93:1298-1306).

[0029] Breast Cancer

[0030] In vivo, hKGF causes mammary duct dilation and rampant epithelial hyperplasia, both of which are common features of breast cancers (Ulich et al., (1994) *Am J Pathol* 144:862-868; Yi et al., (1994) *Am J Pathol* 145:1015-1022). However, the ductal epithelium of breastfeeding rats is resistant to the growth promoting effects of hKGF and this is of interest in regard to epidemiological observations that pregnancy in women decreases susceptibility to breast cancer and that dairy cows almost never develop breast cancer (Kuzma, 1977, *Breast in Pathology*, Mosby Co.). There is additional supporting evidence implicating hKGF in breast cancer. hKGF mRNA has been detected recently in normal human breast tissue and in 12 of 15 breast tumor samples tested (Koos et al., (1993) *J Steroid Biochem Molec Biol* 45:217-225). The presence of hKGF mRNA in breast tumors considered in conjunction with the observation that hKGF is present in nonneoplastic mammary glands and that hKGF causes rampant proliferation of mammary epithelium suggests that hKGF may be an autocrine or paracrine growth factor important in the regulation of the growth of normal and neoplastic mammary epithelium (Ulich et al., (1994) *Am J Pathol* 144:862-868). Infiltrating ductal mammary adenocarcinoma is characteristically enveloped by a desmoplastic stroma that has been postulated to represent a defensive host response to the carcinoma (Ulich et al., (1994) *Am J Pathol* 144:862-868). Since hKGF is stroma derived it is possible that the desmoplastic stroma contributes rather than inhibits the growth of the tumor.

[0031] Prostate Cancer

[0032] The growth promoting effect of androgens on prostate tumors appears to be mediated through hKGF (Yan et al., (1992) *Mol Endo* 6:2123-2128), as androgens induce the expression of hKGF in prostate stroma cells. Prostate tumors that are androgen dependent in vivo, are androgen independent in vitro, but hKGF dependent (Yan et al., (1992) *Mol Endo* 6:2123-2128). In agreement with the role of hKGF as andromedin is the observation that hKGF functions in epithelial induction during seminal vesicle development, a process that is directed by androgen (Alarid et al., (1994) *Proc Natl Acad Sci USA* 91:1074-1078). Furthermore, hKGF causes aberrant activation of the androgen receptor, thus probably contributing to the failure of androgen ablation therapy in prostate cancer (Culig et al., (1994) *Cancer Res* 54:5474-5478). Based on this information, it is possible that genetic alterations cause hKGF to escape androgen regulation and thus convert the androgen dependent tumor

into an androgen independent, highly malignant tumor. Such tumors would still be able to express the androgen regulated marker PSA, as hKGF also causes the aberrant activation of the androgen receptor. It is also likely that hKGF might be responsible for Benign Prostate Hypertrophy (BPH), a common health problem in older men (D. Bottaro, personal communication).

d) hKGF Competitors

[0033] To date, a monoclonal antibody and a short hKGF-receptor derived peptide (25-mer) have been described as hKGF competitors (Bottaro et al., (1993) *J Biol Chem* 268:9180-9183). The monoclonal antibody, designated IG4, has a Kd of 200 pM for hKGF. The short peptide inhibits hKGF binding to the cell surface of NIH/3T3 cells expressing the human receptor with a Ki of about 1-5 μM. Bottaro et al. (WO 94/25057) provide hKGF-receptor peptides which inhibit binding between hKGF and its receptor. Also provided is a method of assaying test compounds for the ability to inhibit hKGF receptor-mediated cell proliferation.

e) Assaying for Receptor-Growth Factor Interaction

[0034] Blocking the interaction of growth factors and lymphokines with their cell surface receptor using antagonists has been an approach for disease treatment. The discovery of such antagonists requires the availability of biochemical assays for the receptor-growth factor or lymphokine interaction. A classic assay has been the competitive inhibition of radiolabeled growth factor or lymphokine (tracer) to its cell surface receptor. These types of assays utilize cell lines that express the relevant receptor on their surface and determine the amount of cell bound tracer in the presence of various concentrations of potential antagonists. Additionally, other assays utilize membrane extracts from cell lines that express the relevant receptor, and tracer binding is followed by filter binding (see Nenquest Drug Discovery System: Human Tumor Necrosis Factor-Alpha, NEN Research Products, E. I. DuPont de Nemours & Co. (Inc.), Boston, Mass.) or by immobilizing the membrane extracts onto solid supports (Urdal et al., (1988) *J Biol Chem* 263:2870-2877; Smith et al., (1991) *Bioch Bioph Res Comm* 176:335-342). Receptor induced electrophoretic mobility shift of tracer has been applied to identify the presence and size of cell surface receptors by crosslinking the receptor to the tracer and then analyzing on denaturing gels (for example see Kull et al., (1985) *Proc natl Acad Sci USA* 82:5756-5760; Hohmann et al., (1989) *J Biol Chem* 264:14927-14934; Stauber et al., (1989) *J Biol Chem* 264:3573-3576). The use of native gels and non-crosslinked complexes has not been described for growth factors or lymphokines and their receptors, but has been widely applied to study nucleic acid protein interactions (Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.).

[0035] Screening of various cancer cell lines for the presence of hKGF receptors by PCR, revealed that all carcinoma cell lines express hKGF receptor mRNA while sarcoma cell lines do not. The presence of mRNA does not necessarily mean that hKGF receptor will be present on the surface of these cells. For hKGF, only cell based assays have been described using Balb/MK keratinocytes (Weissman, (1983) *Cell* 32:599-606) or NIH/3T3 cells transfected with the hKGF receptor (Miki, (1992) *Proc. Natl. Acad. Sci. USA* 89:246-250).

SELEX

[0036] A method for the in vitro evolution of nucleic acid molecules with highly specific binding to target molecules has been developed. This method, Systematic Evolution of Ligands by EXponential enrichment, termed SELEX, is described in U.S. application Ser. No. 07/536,428, entitled "Systematic Evolution of Ligands by Exponential Enrichment," now abandoned, U.S. application Ser. No. 07/714,131, filed Jun. 10, 1991, entitled "Nucleic Acid Ligands," now U.S. Pat. No. 5,475,096, and U.S. application Ser. No. 07/931,473, filed Aug. 17, 1992, entitled "Nucleic Acid Ligands," now U.S. Pat. No. 5,270,163 (see also PCT/US91/04078), each of which is herein specifically incorporated by reference. Each of these applications, collectively referred to herein as the SELEX Patent Applications, describes a fundamentally novel method for making a nucleic acid ligand to any desired target molecule.

[0037] The SELEX method involves selection from a mixture of candidate oligonucleotides and step-wise iterations of binding, partitioning and amplification, using the same general selection scheme, to achieve virtually any desired criterion of binding affinity and selectivity. Starting from a mixture of nucleic acids, preferably comprising a segment of randomized sequence, the SELEX method includes steps of contacting the mixture with the target under conditions favorable for binding, partitioning unbound nucleic acids from those nucleic acids which have bound specifically to target molecules, dissociating the nucleic acid-target complexes, amplifying the nucleic acids dissociated from the nucleic acid-target complexes to yield a ligand-enriched mixture of nucleic acids, then reiterating the steps of binding, partitioning, dissociating and amplifying through as many cycles as desired to yield highly specific, high affinity nucleic acid ligands to the target molecule.

[0038] The basic SELEX method has been modified to achieve a number of specific objectives. For example, U.S. application Ser. No. 07/960,093, filed Oct. 14, 1992, entitled "Method for Selecting Nucleic Acids on the Basis of Structure," describes the use of SELEX in conjunction with gel electrophoresis to select nucleic acid molecules with specific structural characteristics, such as bent DNA. U.S. application Ser. No. 08/123,935, filed Sep. 17, 1993, entitled "Photoselection of Nucleic Acid Ligands," describes a SELEX based method for selecting nucleic acid ligands containing photoreactive groups capable of binding and/or photocrosslinking to and/or photoinactivating a target molecule. U.S. application Ser. No. 08/134,028, filed Oct. 7, 1993, entitled "High-Affinity Nucleic Acid Ligands That Discriminate Between Theophylline and Caffeine," describes a method for identifying highly specific nucleic acid ligands able to discriminate between closely related molecules, termed "Counter-SELEX." U.S. application Ser. No. 08/143,564, filed Oct. 25, 1993, entitled "Systematic Evolution of Ligands by EXponential Enrichment: Solution SELEX," now abandoned (see U.S. Pat. No. 5,567,588), describes a SELEX-based method which achieves highly efficient partitioning between oligonucleotides having high and low affinity for a target molecule. U.S. application Ser. No. 07/964,624, filed Oct. 21, 1992, entitled "Nucleic Acid Ligands to HIV-RT and HIV-1 Rev," now U.S. Pat. No. 5,496,938, describes methods for obtaining improved nucleic acid ligands after SELEX has been performed. U.S. application Ser. No. 08/400,440, filed Mar. 8, 1995, entitled

“Systematic Evolution of Ligands by EXponential Enrichment: Chيمي-SELEX,” now U.S. Pat. No. 5,705,337, describes methods for covalently linking a ligand to its target.

[0039] The SELEX method encompasses the identification of high-affinity nucleic acid ligands containing modified nucleotides conferring improved characteristics on the ligand, such as improved in vivo stability or improved delivery characteristics. Examples of such modifications include chemical substitutions at the ribose and/or phosphate and/or base positions. SELEX-identified nucleic acid ligands containing modified nucleotides are described in U.S. application Ser. No. 08/117,991, filed Sep. 8, 1993, entitled “High Affinity Nucleic Acid Ligands Containing Modified Nucleotides,” now abandoned (see U.S. Pat. No. 5,660,985), that describes oligonucleotides containing nucleotide derivatives chemically modified at the 5- and 2'-positions of pyrimidines. U.S. application Ser. No. 08/134,028, supra, describes highly specific nucleic acid ligands containing one or more nucleotides modified with 2'-amino (2'-NH₂), 2'-fluoro (2'-F), and/or 2'-O-methyl (2'-OMe). U.S. application Ser. No. 08/264,029, filed Jun. 22, 1994, entitled “Novel Method of Preparation of Known and Novel 2' Modified Nucleosides by Intramolecular Nucleophilic Displacement,” describes oligonucleotides containing various 2'-modified pyrimidines.

[0040] The SELEX method encompasses combining selected oligonucleotides with other selected oligonucleotides and non-oligonucleotide functional units as described in U.S. application Ser. No. 08/284,063, filed Aug. 2, 1994, entitled “Systematic Evolution of Ligands by Exponential Enrichment: Chimeric SELEX,” now U.S. Pat. No. 5,637,459, and U.S. application Ser. No. 08/234,997, filed Apr. 28, 1994, entitled “Systematic Evolution of Ligands by Exponential Enrichment: Blended SELEX,” now U.S. Pat. No. 5,683,867, respectively. These applications allow the combination of the broad array of shapes and other properties, and the efficient amplification and replication properties, of oligonucleotides with the desirable properties of other molecules. Each of the above described patent applications which describe modifications of the basic SELEX procedure are specifically incorporated by reference herein in their entirety.

BRIEF SUMMARY OF THE INVENTION

[0041] The present invention includes methods of identifying and producing nucleic acid ligands to transforming growth factor beta (TGFβ), platelet-derived growth factor (PDGF), and human keratinocyte growth factor (hKGF), and homologous proteins, and the nucleic acid ligands so identified and produced. For the purpose of this application, TGFβ includes human TGFβ1, TGFβ2, and TGFβ3 and TGFβ's that are substantially homologous thereto. By substantially homologous it is meant a degree of amino acid sequence identity of 70% or more. For the purposes of this application, PDGF refers to PDGF AA, AB, and BB isoforms and homologous proteins. Specifically included in the definition are human PDGF AA, AB and BB isoforms. In particular, RNA sequences are provided that are capable of binding specifically to TGFβ1, PDGF, and hKGF. Also provided are ssDNA sequences that are capable of binding specifically to TGFβ and PDGF. Specifically included in the invention are the RNA ligand sequences shown in Tables 3,

13, 16, and 23 (SEQ ID NOS:12-42, 128-170, 189-262, 272-304). The RNA ligand sequences of TGFβ shown in Table 3 include both pre and post SELEX modifications. Also included in the invention are ssDNA ligands of TGFβ and PDGF shown in Tables 6, 8, 9, and FIGS. 3, 4, and 9 (SEQ ID NOS:55-89, 93-124, 171-176). Also included in this invention are RNA ligands of TGFβ1 and hKGF that inhibit the function of TGFβ1 and hKGF, presumably by inhibition of the interaction of TGFβ and hKGF with their receptors. Also included in this invention are ssDNA ligands of PDGF that inhibit the function of PDGF, presumably by inhibition of the interaction of PDGF with its receptor.

[0042] Further included in this invention is a method of identifying nucleic acid ligands and nucleic acid ligand sequences to a target selected from the group consisting of TGFβ, PDGF, and hKGF comprising the steps of (a) contacting a candidate mixture of nucleic acids with the target (b) partitioning between members of said candidate mixture on the basis of affinity to the target and (c) amplifying the selected molecules to yield a mixture of nucleic acids enriched for nucleic acid sequences with a relatively higher affinity for binding to the target.

[0043] More specifically, the present invention includes the RNA and ssDNA ligands to TGFβ1 identified according to the above-described method, including those ligands shown in Tables 3 and 6 (SEQ ID NOS:12-42, 55-89). Also included are nucleic acid ligands to TGFβ that are substantially homologous to any of the given ligands and that have substantially the same ability to bind TGFβ and inhibit the function of TGFβ. Further included in this invention are nucleic acid ligands to TGFβ that have substantially the same structural form as the ligands presented herein and that have substantially the same ability to bind TGFβ and inhibit the function of TGFβ.

[0044] Additionally, the present invention includes the ssDNA and RNA ligands to PDGF identified according to the above-described method, including those ligands shown in Tables 8 and 13, and FIGS. 3, 4, and 9 (SEQ ID NOS:93-124, 128-176). Also included are DNA and RNA ligands to PDGF that are substantially homologous to any of the given ligands and that have substantially the same ability to bind PDGF. Further included in this invention are nucleic acid ligands to PDGF that have substantially the same structural form as the ligands presented herein and that have substantially the same ability to bind PDGF.

[0045] In addition, the present invention includes the RNA ligands to hKGF identified according to the above-described method, including those ligands shown in Tables 16 and 23 (SEQ ID NOS:189-264, 268-304). Also included are RNA ligands to hKGF that are substantially homologous to any of the given ligands and that have substantially the same ability to bind hKGF and inhibit the interaction of hKGF with its receptor. Further included in this invention are nucleic acid ligands to hKGF that have substantially the same structural form as the ligands presented herein and that have substantially the same ability to bind hKGF and inhibit the interaction of hKGF with its receptor.

[0046] The present invention also includes other modified nucleotide sequences based on the RNA ligands identified herein and mixtures of the same.

[0047] Further included in this invention is a method of assaying a test compound for the ability to inhibit hKGF

receptor-mediated cell proliferation comprising the steps of (a) contacting the test compound with a hKGF nucleic acid ligand and a keratinocyte growth factor; and (b) detecting the ability of the test compound to inhibit binding between the hKGF nucleic acid ligand and the keratinocyte growth factor.

[0048] Also included in this invention is a method of assaying a test compound for the ability to inhibit the interaction of a growth factor with its plasma membrane bound receptor comprising the steps of (a) solubilizing cells containing the plasma membrane bound receptor; (b) creating a plasma membrane extract of the cells; (c) reacting the extract with labeled growth factor alone and in the presence of the test compound thereby creating complexes; (d) analyzing the complexes by electrophoresis under native conditions; (e) visualizing the complexes by imaging; and (f) comparing the image of the extract with labeled growth factor alone to the image of the extract in the presence of the test compound to determine whether the test compound inhibited the interaction between the growth factor and its plasma membrane bound receptor.

[0049] Further included in this invention is a method for assaying cells to determine whether they express a growth factor plasma membrane bound receptor comprising the steps of (a) solubilizing the cells; (b) creating a plasma membrane extract of the cells; (c) reacting the plasma membrane extract with a labeled growth factor; (d) analyzing the reaction between the plasma membrane extract with the labeled growth factor by electrophoresis under native conditions; (e) comparing the electrophoresis of step (d) with electrophoresis of labeled growth factor; and (f) visualizing the results of the electrophoresis to determine whether a complex is formed with altered mobility relative to the mobility of a labeled growth factor alone.

BRIEF DESCRIPTION OF THE FIGURES

[0050] **FIG. 1** shows the binding analysis of the 40D7 DNA library for TGF β 1. Binding data obtained from Round 19 (triangles) and Round 0 (circles) are shown.

[0051] **FIG. 2** shows the results of the PAI-luciferase assay of TGF β 1 (10 pM) incubated with oligonucleotides (0.1 μ M) or anti-TGF13 (60 μ g/ml).

[0052] **FIG. 3** shows the consensus secondary structure for the sequence set shown in Table 9. R=A or G, Y=C or T, K=G or T, N and N' indicate any base pair.

[0053] **FIG. 4** shows the minimal ligands 20 t, 36 t and 41 t folded according to the consensus secondary structure motif. [3'T] represents a 3'-3' linked thymidine nucleotide added to reduce 3'-exonuclease degradation.

[0054] **FIGS. 5A, 5B and 5C** show the binding of minimal high affinity DNA ligands to PDGF. The fraction of 32 P 5' end-labeled DNA ligands bound to varying concentrations of PDGF was determined by the nitrocellulose filter binding method. Minimal ligands tested were 20 t (o), 36 t (Δ), and 41 t (\square). Oligonucleotide concentrations in these experiments were \approx 10 pM (PDGF-AB and PDGF-BB) and \approx 50 pM (PDGF-AA). Data points were fitted to eq. 1 (for binding of the DNA ligands to PDGF-AA) or to eq. 2 (for binding to PDGF AB and BB) using the non-linear least squares method. Binding reactions were done at 37 $^{\circ}$ C. in binding buffer (PBSM with 0.01% HSA).

[0055] **FIG. 6** shows the dissociation rate determination for the high affinity interaction between the minimal DNA ligands and PDGF AB. The fraction of 32 P end-labeled ligands 20 t (o), 36 t (Δ), and 41 t (\square), all at 0.17 nM, bound to PDGF AB (1 nM) was measured by nitrocellulose filter binding at the indicated time points following the addition of a 500-fold excess of the unlabeled competitor. The dissociation rate constant (k_{off}) values were determined by fitting the data points to eq. 3. The experiments were performed at 37 $^{\circ}$ C. in binding buffer.

[0056] **FIG. 7** shows the effect of DNA ligands on the binding of 125 I-PDGF-BB and 125 I-PDGF-AA to PDGF a-receptors expressed in PAE cells.

[0057] **FIG. 8** shows the effect of DNA ligands on the mitogenic effect of PDGF-BB on PAE cells expressing the PDGF β -receptors.

[0058] **FIG. 9** shows the 2'-O-methyl-2'-deoxy- and 2'-fluoro-2'-deoxyribonucleotide substitution pattern compatible with high affinity binding to PDGF-AB. Underlined symbols indicate 2'-O-methyl-2'-deoxynucleotides; italicized symbols indicate 2'-fluoro-2'-deoxynucleotides; normal font indicates 2'-deoxyribonucleotides; [3'T] indicates inverted orientation (3'3') thymidine nucleotide (Glen Research, Sterling, Va.); PEG in the loops of helices II and III indicates pentaethylene glycol spacer phosphoramidite (Glen Research, Sterling, Va.).

[0059] **FIG. 10A** shows the saturation binding of radiolabeled hKGF on the surface of the PC-3 cells. TB (total binding) is the binding observed in the absence of competing unlabeled hKGF, whereas NSB (nonspecific binding) is the binding observed in the presence of 100 fold molar excess of unlabeled hKGF. SB (specific binding) demonstrates the specific binding, and this curve was derived by subtracting the NSB curve from the TB curve. **FIG. 10B** is the Scatchard analysis of the data points shown in 10A for the SB curve.

[0060] **FIG. 11** shows the shift of the electrophoretic mobility due to plasma membrane extracts from PC-3 cells. In lanes 1-8, the membrane extracts were reacted with various concentrations of radiolabeled hKGF as shown under each lane. In addition to the radiolabeled hKGF (as shown under each lane) for lanes 9-12 a 100 fold molar excess was included of unlabeled hKGF. C1 and C2 represent two observed complexes due to the presence of hKGF binding moieties in the PC-3 plasma membrane extracts.

[0061] **FIGS. 12A-D** show the proposed alignment of 2'F and 2'NH $_2$ ligands. Lower case, italicized sequence residues indicate the constant region of the template. In the consensus sequences, capital and lower case letters are used for residues found in greater than or equal to 80% and 60% of the members of each family respectively. K_d and K_i values are also shown next to the designation of each ligand. The K_i values shown here were calculated using the formula $K_i = IC50 / (1 + (C/K_d))$, where IC50 is the measured half maximal inhibitory concentration of each ligand in the PC-3 cell assay as described in Example 16; C is the concentration of 125 I-KGF; and K_d is the equilibrium dissociation constant of KGF for its receptor, (about 150 pM). The ligands marked with stars show biphasic binding curves.

[0062] **FIGS. 12A and 12B** show the proposed alignment of 2'F ligands. The majority of 2'F ligands can be folded into

pseudoknot structures. Two classes are proposed as shown. The summary structure for each class is also shown. Bases participating in stem 1 (S1) are underlined with single lines while bases of stem 2 (S2) are underlined with double lines. Spaces were introduced for alignment of the various elements of the pseudoknots.

[0063] FIGS. 12C and 12D show the proposed folding of 2'NH₂ ligands. These ligands are assigned into two classes. As shown in the summary structures, class 1 and class 2 ligands can form a stem-loop and dumbbell structure, respectively. Spaces were introduced to allow sequence alignment. Residues participating in stems are underlined. In the summary structures, periods (.) indicate a variable number of residues. Ligands 2N and 54N are circular permutations of the same dumbbell structure. For alignment of the corresponding loops these ligands are wrapped around two lines.

[0064] FIG. 13 shows the minimal sequence requirement for binding of ligand 6F and 14F to hKGF. The predicted folding of each ligand is shown. Constant regions of the ligands are shown in lower case. Conserved sequences are underlined. Circles and triangles mark the 3' ends of active and inactive truncates respectively.

DETAILED DESCRIPTION OF THE INVENTION

[0065] This application describes high-affinity nucleic acid ligands to TGF β , PDGF, and hKGF identified through the method known as SELEX. SELEX is described in U.S. application Ser. No. 07/536,428, entitled "Systematic Evolution of Ligands by EXponential Enrichment," now abandoned, U.S. application Ser. No. 07/714,131, filed Jun. 10, 1991, entitled "Nucleic Acid Ligands," now U.S. Pat. No. 5,475,096, and U.S. application Ser. No. 07/931,473, filed Aug. 17, 1992, entitled "Methods for Identifying Nucleic Acid Ligands," now U.S. Pat. No. 5,270,163, (see also WO 91/19813). These applications, each specifically incorporated herein by reference, are collectively called the SELEX Patent Applications.

[0066] In its most basic form, the SELEX process may be defined by the following series of steps:

[0067] 1) A candidate mixture of nucleic acids of differing sequence is prepared. The candidate mixture generally includes regions of fixed sequences (i.e., each of the members of the candidate mixture contains the same sequences in the same location) and regions of randomized sequences. The fixed sequence regions are selected either: (a) to assist in the amplification steps described below, (b) to mimic a sequence known to bind to the target, or (c) to enhance the concentration of a given structural arrangement of the nucleic acids in the candidate mixture. The randomized sequences can be totally randomized (i.e., the probability of finding a base at any position being one in four) or only partially randomized (e.g., the probability of finding a base at any location can be selected at any level between 0 and 100 percent).

[0068] 2) The candidate mixture is contacted with the selected target under conditions favorable for binding between the target and members of the candidate mixture. Under these circumstances, the interaction between the target and the nucleic acids of the candidate mixture can be

considered as forming nucleic acid-target pairs between the target and those nucleic acids having the strongest affinity for the target.

[0069] 3) The nucleic acids with the highest affinity for the target are partitioned from those nucleic acids with lesser affinity to the target. Because only an extremely small number of sequences (and possibly only one molecule of nucleic acid) corresponding to the highest affinity nucleic acids exist in the candidate mixture, it is generally desirable to set the partitioning criteria so that a significant amount of the nucleic acids in the candidate mixture (approximately 5-50%) are retained during partitioning.

[0070] 4) Those nucleic acids selected during partitioning as having the relatively higher affinity to the target are then amplified to create a new candidate mixture that is enriched in nucleic acids having a relatively higher affinity for the target.

[0071] 5) By repeating the partitioning and amplifying steps above, the newly formed candidate mixture contains fewer and fewer weakly binding sequences, and the average degree of affinity of the nucleic acids to the target will generally increase. Taken to its extreme, the SELEX process will yield a candidate mixture containing one or a small number of unique nucleic acids representing those nucleic acids from the original candidate mixture having the highest affinity to the target molecule.

[0072] The SELEX Patent Applications describe and elaborate on this process in great detail. Included are targets that can be used in the process; methods for partitioning nucleic acids within a candidate mixture; and methods for amplifying partitioned nucleic acids to generate enriched candidate mixture. The SELEX Patent Applications also describe ligands obtained to a number of target species, including both protein targets where the protein is and is not a nucleic acid binding protein.

[0073] The nucleic acid ligands described herein can be complexed with a lipophilic compound (e.g., cholesterol) or attached to or encapsulated in a complex comprised of lipophilic components (e.g., a liposome). The complexed nucleic acid ligands can enhance the cellular uptake of the nucleic acid ligands by a cell for delivery of the nucleic acid ligands to an intracellular target. U.S. application Ser. No. 08/434,465, filed May 4, 1995, entitled "Nucleic Acid Ligand Complexes," now U.S. Pat. No. 6,011,020, which is incorporated in its entirety herein, describes a method for preparing a therapeutic or diagnostic complex comprised of a nucleic acid ligand and a lipophilic compound or a non-immunogenic, high molecular weight compound.

[0074] The methods described herein and the nucleic acid ligands identified by such methods are useful for both therapeutic and diagnostic purposes. Therapeutic uses include the treatment or prevention of diseases or medical conditions in human patients. Diagnostic utilization may include both in vivo or in vitro diagnostic applications. The SELEX method generally, and the specific adaptations of the SELEX method taught and claimed herein specifically, are particularly suited for diagnostic applications. SELEX identifies nucleic acid ligands that are able to bind targets with high affinity and with surprising specificity. These characteristics are, of course, the desired properties one skilled in the art would seek in a diagnostic ligand.

[0075] The nucleic acid ligands of the present invention may be routinely adapted for diagnostic purposes according to any number of techniques employed by those skilled in the art. Diagnostic agents need only be able to allow the user to identify the presence of a given target at a particular locale or concentration. Simply the ability to form binding pairs with the target may be sufficient to trigger a positive signal for diagnostic purposes. Those skilled in the art would also be able to adapt any nucleic acid ligand by procedures known in the art to incorporate a labeling tag in order to track the presence of such ligand. Such a tag could be used in a number of diagnostic procedures. The nucleic acid ligands described herein may specifically be used for identification of the TGF β , PDGF, and hKGF proteins.

[0076] SELEX provides high affinity ligands of a target molecule. This represents a singular achievement that is unprecedented in the field of nucleic acids research. The present invention applies the SELEX procedure to the specific targets of TGF β , PDGF, and hKGF. In the Example section below, the experimental parameters used to isolate and identify the nucleic acid ligands to TGF β , PDGF, and hKGF are described.

[0077] In order to produce nucleic acids desirable for use as a pharmaceutical, it is preferred that the nucleic acid ligand (1) binds to the target in a manner capable of achieving the desired effect on the target; (2) be as small as possible to obtain the desired effect; (3) be as stable as possible; and (4) be a specific ligand to the chosen target. In most situations, it is preferred that the nucleic acid ligand have the highest possible affinity to the target.

[0078] In co-pending and commonly assigned U.S. application Ser. No. 07/964,624, filed Oct. 21, 1992 ('624), now U.S. Pat. No. 5,496,938, methods are described for obtaining improved nucleic acid ligands after SELEX has been performed. The '624 application, entitled "Nucleic Acid Ligands to HIV-RT and HIV-1 Rev," is specifically incorporated herein by reference. Further included in this patent are methods for determining the three-dimensional structures of nucleic acid ligands. Such methods include mathematical modeling and structure modifications of the SELEX-derived ligands, such as chemical modification and nucleotide substitution.

[0079] In the present invention, SELEX experiments were performed in order to identify RNA and DNA ligands with specific high affinity for TGF β 1 from degenerate libraries containing 40 or 60 random positions (40N or 60N) (Tables 1 and 5). This invention includes the specific RNA ligands to TGF β 1 shown in Table 3 (SEQ ID NOS:12-42), identified by the methods described in Examples 1 and 2. This invention further includes RNA ligands to TGF β which inhibit TGF β 1 function, presumably by inhibiting the interaction of TGF β 1 with its receptor. This invention includes the specific ssDNA ligands to TGF β 1 shown in Table 6 (SEQ ID NOS:55-89) identified by the methods described in Examples 5 and 6.

[0080] In the present invention, two SELEX experiments were also performed in order to identify ssDNA and RNA with specific high affinity for PDGF from degenerate libraries containing 40 and 50 random positions (40N and 50N), respectively (Tables 7 and 12). This invention includes the specific ssDNA and RNA ligands to PDGF shown in Tables

8, 9 and 13 and **FIGS. 3, 4, and 9** (SEQ ID NOS:93-124, 128-176), identified by the methods described in Examples 7 and 15.

[0081] In the present invention, a SELEX experiment was also performed in search of RNA ligands with specific high affinity for hKGF from degenerate libraries containing 40 random positions (40N) (Table 14). This invention includes the specific RNA ligands to hKGF shown in Tables 16 and 23 and **FIG. 12** (SEQ ID NOS:189-262, 268-304), identified by the methods described in Examples 16 and 17. This invention further includes RNA ligands to hKGF which inhibit the interaction of hKGF with its receptor.

[0082] The scope of the ligands covered by this invention extends to all nucleic acid ligands of TGF β , PDGF, and hKGF, modified and unmodified, preferably those identified according to the SELEX procedure. More specifically, this invention includes nucleic acid sequences that are substantially homologous to the ligands shown in Tables 3, 6, 8, 9, 13, 16, and 23 and **FIGS. 3, 4, 9 and 12** (SEQ ID NOS:12-42, 55-89, 93-124, 128-176, 189-262, 268-304). By substantially homologous it is meant a degree of primary sequence homology in excess of 70%, most preferably in excess of 80%. A review of the sequence homologies of the nucleic acid ligands shown in Tables 3 and 6 (SEQ ID NOS:12-42, 55-89) for TGF β , Tables 8 and 13 (SEQ ID NOS:93-124, 128-170) for PDGF, and Tables 16 and 23 (SEQ ID NOS:189-262, 272-304) for hKGF shows that sequences with little or no primary homology may have substantially the same ability to bind a given target. For these reasons, this invention also includes nucleic acid ligands that have substantially the same structure and ability to bind TGF β , PDGF, and hKGF as the nucleic acid ligands shown in Tables 3, 6, 8, 9, 13, 16, and 23 and **FIGS. 3, 4, 9 and 12** (SEQ ID NOS:12-42, 55-89, 93-124, 128-176, 189-262, 268-304). Substantially the same structure for PDGF includes all nucleic acid ligands having the common structural elements shown in **FIG. 3** that lead to the affinity to PDGF. Substantially the same ability to bind TGF β , PDGF, or hKGF means that the affinity is within one or two orders of magnitude of the affinity of the ligands described herein. It is well within the skill of those of ordinary skill in the art to determine whether a given sequence—substantially homologous to those specifically described herein—has substantially the same ability to bind TGF β , PDGF, or hKGF.

[0083] This invention also includes the ligands as described above, wherein certain chemical modifications are made in order to increase the in vivo stability of the ligand or to enhance or mediate the delivery of the ligand. Examples of such modifications include chemical substitutions at the sugar and/or phosphate and/or base positions of a given nucleic acid sequence. See, e.g., U.S. application Ser. No. 08/117,991, filed Sep. 9, 1993, entitled "High Affinity Nucleic Acid Ligands Containing Modified Nucleotides," now abandoned (see U.S. Pat. No. 5,660,985), which is specifically incorporated herein by reference. Other modifications are known to one of ordinary skill in the art. Such modifications may be made post-SELEX (modification of previously identified modified or unmodified ligands) or by incorporation into the SELEX process.

[0084] Example 20 describes post-SELEX procedure modification of a nucleic acid ligand to basic fibroblast

growth factor (bFGF). The nucleic acid ligand was modified by the addition of phosphorothioate caps and substitution of several ribopurines with 2'-deoxy-2'-O-methylpurines.

[0085] As described above, because of their ability to selectively bind TGF β , PDGF, and hKGF, the nucleic acid ligands to TGF β , PDGF, and hKGF described herein are useful as pharmaceuticals. This invention, therefore, also includes a method for treating TGF- β -mediated pathological conditions by administration of a nucleic acid ligand capable of binding to TGF β , a method for treating PDGF-mediated pathological conditions by administration of a nucleic acid ligand capable of binding to PDGF, and a method for treating hKGF-mediated pathological conditions by administration of a nucleic acid ligand capable of binding to hKGF.

[0086] Therapeutic compositions of the nucleic acid ligands may be administered parenterally by injection, although other effective administration forms, such as intraarticular injection, inhalant mists, orally active formulations, transdermal iontophoresis or suppositories, are also envisioned. One preferred carrier is physiological saline solution, but it is contemplated that other pharmaceutically acceptable carriers may also be used. In one preferred embodiment, it is envisioned that the carrier and the ligand constitute a physiologically-compatible, slow release formulation. The primary solvent in such a carrier may be either aqueous or non-aqueous in nature. In addition, the carrier may contain other pharmacologically-acceptable excipients for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the carrier may contain still other pharmacologically-acceptable excipients for modifying or maintaining the stability, rate of dissolution, release, or absorption of the ligand. Such excipients are those substances usually and customarily employed to formulate dosages for parental administration in either unit dose or multi-dose form.

[0087] Once the therapeutic composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready to use form or requiring reconstitution immediately prior to administration. The manner of administering formulations containing nucleic acid ligands for systemic delivery may be via subcutaneous, intramuscular, intravenous, intranasal or vaginal or rectal suppository.

[0088] The following examples are provided to explain and illustrate the present invention and are not intended to be limiting of the invention. Examples 1-4 describe initial experiments to identify RNA with specific high affinity for TGF β 1. Example 1 describes the various materials and experimental procedures used in Examples 2-4. Example 2 describes a representative method for identifying RNA ligands by the SELEX method which bind TGF β 1. Example 3 describes the affinities the ligands have for TGF β 1 and demonstrates that the ligands are capable of inhibiting the function of TGF β 1, presumably by inhibiting the interaction of TGF β 1 with its receptor. Example 4 describes which regions of the ligands are believed to be necessary for TGF β 1 binding and inhibition of TGF β 1 receptor binding. Example 5 describes another representative method for identifying RNA and DNA ligands by the SELEX method which bind TGF β 1. Example 6 reports on the binding

analysis, bioassay, and sequences of a ssDNA SELEX library. Example 7 describes the various materials and experimental procedures used in evolving ssDNA ligands to PDGF described in Examples 8-13. Example 8 describes the ssDNA ligands to PDGF and the predicted secondary structure of selected nucleic acid ligands. Example 9 describes the minimal sequence necessary for high affinity binding. Example 10 describes the kinetic stability of PDGF-Nucleic Acid Ligand complexes. Example 11 describes the thermal melting properties for selected ligands. Example 12 describes photo-crosslinking of nucleic acid ligands and PDGF. Example 13 describes the inhibition by DNA ligands of PDGF isoforms on cultured cells and inhibition of the mitogenic effects of PDGF in cells by DNA ligands. Example 14 describes the modification of nucleic acid ligands to PDGF with modified nucleotides. Example 15 describes the experimental procedures used in evolving RNA ligands to PDGF and shows the ligand sequences. Example 16 describes the various materials and experimental procedures used in evolving nucleic acid ligands to hKGF described in Examples 17-19. Example 17 describes the RNA ligands to hKGF, the affinities the ligands have for hKGF, and the specificity of the RNA ligands to hKGF. Example 18 describes inhibition of hKGF binding to cell surface receptors. Example 19 reports on the inhibition of mitogenic activity of hKGF by a selected ligand. Example 20 describes the modification of nucleic acid ligands to bFGF with 2'-deoxy-2'-O-methylpurines.

EXAMPLES

Example 1

Experimental Procedures

[0089] This example provides the general procedures followed and incorporated in Examples 2-4.

[0090] A. Materials.

[0091] Human recombinant TGF β 1 used in this SELEX procedure was acquired from Genentech. Human recombinant TGF β 1 can also be purchased from R&D systems, Minneapolis, Minn., USA.

[0092] Biotinylated TGF β 1 was prepared by reacting TGF β 1 at 3.6 μ M with an 11 fold molar excess of sulfo-NHS-biotin (Pierce, Rockford, Ill., USA) in 50 mM NaHCO₃ for 3 hr in an ice bath. The reaction was acidified with 0.036 volumes of 10% acetic acid and applied to a 40 mg Vydac (The Separations Group, Hesperia, Calif., USA) reverse phase column made in a siliconized pipet tip to separate unreacted biotin from biotinylated TGF β 1. The column was prewashed with 200 μ l ethanol followed by 200 μ l 1% acetic acid, the biotinylation reaction was applied, free biotin was washed through with 200 μ l of 50 mM sodium acetate pH 5.5, followed by 200 μ l of 20% acetonitrile and finally eluted with 200 μ l of 60% acetonitrile. The sample was lyophilized and resuspended in 50 mM sodium acetate pH 5.0 at 40 μ M and stored at 4 $^{\circ}$ C. The TGF β 1 was spiked with 100,000 cpm iodinated TGF β 1 in order to follow recovery and to assess the success of the biotinylation reaction by measuring the fraction of the radioactivity that would bind to streptavidin coated agarose beads (Pierce) before and after biotinylation. An aliquot of the TGF β 1 before and after biotinylation was subjected to analytical

reverse phase chromatography. The biotinylated TGF β 1 substantially ran as a single peak which was retarded with respect to the unbiotinylated TGF β . A small amount (5%) of unreacted TGF β 1 could be detected. The efficiency of binding of the iodinated, biotinylated TGF β 1 to streptavidin (SA) agarose beads (30 μ l) was 30% under the binding conditions used for SELEX partitioning.

[0093] Iodinated TGF β 1 was prepared by the lactoperoxidase method (50 mM sodium phosphate pH 7.3, 0.16% glucose) with BioRad Enzymo beads (BioRad, Richmond, Calif., USA) and the bound iodine separated from the free iodine by gel filtration on G25 Sephadex in 50 mM sodium acetate 0.01% Tween.

[0094] The mink lung cell line expressing the luciferase reporter gene under the control of PAI 1 promoter (Abe et al. (1994) *Anal. Biochem.* 216:276-284) was a gift from Dr. Dan Rifkin (Department of Cell Biology, New York Medical Center, New York, N.Y. 10016). Luciferase was assayed by reagents purchased from Analytical Luminescence Laboratory, San Diego, Calif., USA.

[0095] 2'-NH $_2$ modified CTP and UTP were prepared according to the method of Pieken et al. (1991) *Science* 253:314-317. DNA oligonucleotides were synthesized using standard procedures either at NeXstar Pharmaceuticals, Inc. (Boulder, Colo., USA) or by Operon Technologies (Alameda, Calif., USA). All other reagents and chemicals were purchased from standard commercial sources and sources have been indicated.

[0096] B. SELEX Procedure

[0097] SELEX ligands that bind to TGF β 1 were derived essentially as described in U.S. Pat. No. 5,270,163 (see also, Tuerk and Gold (1990) *Science* 249:505-510). To generate the starting pool of PCR template, PCR product from twenty separate PCR reactions each containing 16.1 pmol of unpurified, single stranded DNA (at least a total of 2×10^{12} to 2×10^{10} different molecules) were pooled before the first transcription. PCR conditions were 50 mM KCl, 10 mM Tris-HCl, pH 9, 0.1% Triton-X100, 1.5 mM MgCl $_2$, 0.2 mM of each dATP, dCTP, dGTP, and dTTP, 2 μ M each primer and 0.075 units/ μ l of Taq DNA polymerase, 100 μ l per reaction in a siliconized microfuge tube. All PCR cycles took advantage of hot start using Ampliwax (Perk and Elmer, Norwalk, Conn., USA). Duration of the initial PCR was 10 cycles; a PCR cycle was 94 $^\circ$ C.-1', 52 $^\circ$ C.-1', 72 $^\circ$ C.-2'. An initial denaturation was 94 $^\circ$ C. for 4' and the final extension at 72 $^\circ$ C. for 5'. PCR reactions were combined, phenol/ chloroform extracted, and isopropanol precipitated (2.0 M ammonium acetate, 50% isopropanol) to remove primers.

[0098] Transcription reactions contained 200 nM DNA, 0.9 mM GTP, 0.9 mM 2'-NH $_2$ -UTP, 0.9 mM 2'-NH $_2$ -CTP, 0.5 mM ATP, 87 mM Tris-HCl pH 8.0, 17 mM MgCl $_2$, 4.4 mM spermidine, 22 mM DTT, 100 μ g/ml acetylated BSA (Promega, Madison, Wis., USA) and 4 units/ μ l T7 RNA polymerase. (2'-F-UTP and 2'-F-CTP (United States Biochemical, Cleveland, Ohio, USA) were used at 3.0 mM, whereas UTP and CTP were used at 0.9 mM each). Transcription reactions were incubated overnight at 28 $^\circ$ C. (at least 10 hours). After transcription the template was digested by addition of 2 μ l RQ1 Dnase (Promega) for 15' at 28 $^\circ$ C., and then extracted with phenol/CHCl $_3$, followed by three ethanol precipitations from ammonium acetate (3.9 M ammonium acetate, 72% ethanol).

[0099] The RNA molecules were incubated with TGF β 1 bound to SA agarose beads as described below in Krebs-Ringer solution (KR) (120 mM NaCl, 4.8 mM KCl, 10 mM Na phosphate buffer pH 7.4, 1.2 mM MgSO $_4$, 2.6 mM CaCl $_2$) modified to include 20 mM Na-Hepes pH 7.5 and 0.2% Triton X100 (Pierce). This buffer is referred to as KRHT.

[0100] TGF β 1-RNA complexes were separated from unbound RNA by washing the beads. Recovery of the selected 2'-NH $_2$ or F pyrimidine modified RNA from the agarose beads required guanidine thiocyanate extraction (5M GnSCN, 10 mM Tris-HCl, 0.1 mM EDTA, pH 7.0, 0.1 M beta mercaptoethanol) or from Seradyne SA coated beads by 2% SDS (0.1 M Tris-HCl pH 7.5, 50 mM NaCl, 1 mM Na $_2$ EDTA, 2% SDS, 1.5 mM DTT). Regular 2'-OH RNA was easily recovered under less harsh conditions with the same buffer used for the Seradyne beads containing only 0.2% SDS. After extraction and precipitation to purify and concentrate the RNA, the sample was reverse transcribed with a cloned MMLV RT with the RNase H sequence deleted. The reaction contained less than or equal to 16 nM RNA, 10 μ M 3' primer, 50 mM Tris-HCl pH 8.3, 75 mM KCl, 5 mM MgCl $_2$, 10 mM DTT, 0.5 mM dNTP's. Prior to addition of buffer the RNA and the primer were boiled together. After addition of buffer and salts the reaction was annealed for 10 min at 28 $^\circ$ C. before addition of 600 units of Superscript reverse transcriptase (Bethesda Research Labs, Gaithersburg, Md., USA) and synthesis at 50 $^\circ$ C. for 1 hour.

[0101] PCR amplification of this cDNA (<1 pmol) resulted in approximately 250 pmol double-stranded DNA, of this, 40 pmols was transcribed and used to initiate the next round of SELEX.

[0102] C. Partitioning Method for SELEX.

[0103] 2.5 pmols biotinylated TGF β 1 were bound to 30 μ l SA agarose beads (Pierce) in 200 μ l KRHT. The mixture was incubated on a rotator at 37 $^\circ$ C. for 15 to 30 minutes. The beads were washed three times by centrifugation and resuspension in 200 μ l cold KRHT to remove unbound TGF β 1, and resuspended in a final volume of 500 μ l KRHT. RNA containing 2'-NH $_2$ pyrimidines was heated at 70 $^\circ$ C. for three minutes (RNAs containing 2'-OH or 2'-F pyrimidines were heated at 95 $^\circ$ C.) and diluted into KRHT containing TGF β 1 bound to SA beads. The final concentration of RNA is 1 μ M and the TGF β 1 was 5 nM. Binding occurs with rotation at 37 $^\circ$ C. for 30 minutes. Beads were washed by centrifugation and resuspension three times with 200 μ l binding buffer to remove unbound RNA. RNA was eluted from the beads as described above.

[0104] D. Binding Assays.

[0105] Two binding assays for ligands to TGF β 1 gave equivalent results wherever tested. In the SA bead assay the biotinylated TGF β 1 was serially diluted in KRHT in polypropylene tubes (Linbro, ICN, Irvine, Calif., USA) and bound to the beads as described above. After unbound TGF β 1 was washed away, trace quantities of 32 P-labeled RNA (<0.1 nM) were added to each tube and vortexed to mix. After static incubation at 37 $^\circ$ C. for 30 minutes, the unbound RNA was removed by washing three times with 200 μ l of KRHT.

[0106] In the nitrocellulose filter binding assay, TGF β 1 was serially diluted in KRH containing 0.1% defatted BSA

(Fluka radioimmunoassay grade, Fluka, Hauppauge, N.Y., USA) as carrier instead of Triton X-100. Incubation with RNA tracer was as above. Samples were pipetted with a multiwell pipettor onto a multiwell manifold holding a sheet of wet BioRad 0.45 micron nitrocellulose, aspirated, and washed three times with 200 μ l KRH (containing no BSA). The filters were air dried and counted in a liquid scintillation counter (Beckmann Instruments, Palo Alto, Calif.)

[0107] The equation used to fit the binding of ligands to TGF β 1 describes the binding of a ligand to a receptor (in this case TGF β 1) that follows the laws of mass action and for which there is a single binding site: $Y=B_{max} * X / (Kd + X)$; where Y is the fraction of the ligand bound, B_{max} is the maximum fraction of the ligand bound, X is the concentration of TGF β 1 and Kd is the dissociation constant of TGF β 1 and the ligand. Data points were fit by nonlinear regression using the computer program Graphpad Prism (Graphpad Software, San Diego, Calif.). The algorithm minimized the sum of the squares of the actual distance of the points from the curve. Convergence was reached when two consecutive iterations changed the sum-of-squares by less than 0.01%.

[0108] E. Cloning and Sequencing.

[0109] SELEX experiments are described in Table 2. Primers for SELEX experiments 1 and 2 shown in Table 1 contain recognition sites for the restriction endonucleases SacI (5' primer T7SacBam; SEQ ID NO:7) and XbaI (3' primer 3 XH; SEQ ID NO:9). PCR products from SELEX experiments 1 and 2 were cloned directionally into SacI/XbaI digested pGem 9 zf (Promega). 5' primer T7SB2N (SEQ ID NO:8) and 3' primer 3 XH (SEQ ID NO:9) (Table 1) were used for SELEX experiments 3-9. PCR products from SELEX experiments 3-9 were cloned directionally into the BamHI/XbaI site of a modified pGem9zf :BamHI cloning vector. The BamHI site was engineered into pGem9zf in the following way. A clone isolated from library 2 (lib2-6-2) that did not bind to TGF β 1 (sequence not shown) was digested with BamHI and XbaI. The sequence flanking the cloning site of the modified pGem9zf vector is shown in Table 1 (SEQ ID NOS:10-11).

[0110] After digestion of the plasmid with restriction endonuclease and dephosphorylation with CIP (calf intestinal phosphatase), vectors were gel purified. Inserts were ligated and recombinant plasmids were transformed into *E. coli* strain DHIOB (Bethesda Research Labs). Plasmid DNA was prepared by alkaline lysis, mini prep procedure. Twenty-two clones representing 9 unique sequences were sequenced at random from libraries 1 and 2. 50 clones were sequenced from libraries 3-9 using a single dideoxy G reaction (called G track). The sequencing ladders were compared and organized for similarities. Selected clones from each family were chosen for complete sequence analysis. TGF β 1 binding assays were performed on transcripts representing different G sequences in each library. Out of a total of 140 binding assays, 27 ligands bound with a Kd less than 10 nM, and 21 of these were sequenced. Clones were sequenced with the Sequenase sequencing kit (United States Biochemical Corporation, Cleveland, Ohio).

[0111] F. Ligand Truncation.

[0112] Truncation experiments were carried out to determine the minimal sequence necessary for high affinity binding of the RNA ligands to TGF β 1. For 3' boundary

determination, RNA ligands were 5' end-labeled with γ - 32 P-ATP using T4 polynucleotide kinase. 5' boundaries were established with 3' end-labeled ligands using α - 32 P-pCp and T4 RNA ligase. After partial alkaline hydrolysis, radiolabeled RNA ligands were incubated with TGF β 1 at concentrations ranging from 1 nM to 50 nM and protein-bound RNA was separated by nitrocellulose partitioning. RNA truncates were analyzed on a high-resolution denaturing polyacrylamide gel. A ladder of radioactively labeled ligands terminating with G-residues was generated by partial RNase T1 digestion and was used as markers.

[0113] G. Inhibition of TGF β 1 Function.

[0114] TGF β 1 signal transduction begins with binding to a cell surface receptor and results in the induction of transcription of a variety of genes. One of these genes is Pai1. The TGF β 1 assay utilizes the mink lung epithelial cell (MLEC) carrying the luciferase reporter gene fused to the Pai 1 promoter. The MLEC has TGF β 1 receptors on its cell surface. Thus one can measure the response of the cells to TGF β 1 and the effective concentration of TGF β 1 in the culture media by measuring the luciferase enzyme activity after a period of induction.

[0115] Mink lung epithelial cells (MLEC) carrying the Pai1/luc construct were maintained in DME containing 10% fetal bovine serum and 400 μ g/ml G418. MLEC-Pai1/luc cells were plated at 3.2×10^4 cells per well in a 96 well Falcon plate, in 100 μ l of DME+10% fetal bovine serum overnight. Media was made from autoclaved water. The cells were washed three times (100 μ l) in serum free DME plus Solution A (1:1). Solution A is 30 mM Hepes pH 7.6, 10 mM glucose, 3.0 mM KCl, 131 mM NaCl, 1.0 mM disodium phosphate. Samples (100 μ l) were added in DME containing 20 mM Hepes pH 7.5, and 0.1% BSA (Fluka, radioimmunoassay grade). All samples were in triplicate. After six hours at 37 $^{\circ}$ C. in a 5% CO $_2$ incubator the media was removed and cells were washed three times (100 μ l each) in cold PBS. Lysis buffer (75 μ l) (Analytical Luminescence Laboratory) was added and the plates incubated on ice for 20 min. The plates were sealed and frozen at -80 $^{\circ}$ C. until assayed. Samples (25 μ l) were assayed for luciferase activity with the Enhanced Luciferase Assay Kit from Analytical Luminescence Laboratory (San Diego, Calif., USA) according to the manufacturer's instructions using the Berthold Microlumet LB96P luminometer. Luminescence is reproducibly a function of TGF β 1 concentration added to the media.

[0116] Ligands tested for inhibition of TGF β 1 activity were tested at a minimum of five concentrations. The ligands were serially diluted in DME, 20 mM Hepes pH 7.5, 0.1% Fluka BSA in polypropylene tubes and an equal volume of media containing 12 pM TGF β 1 was added to each tube, vortexed and transferred to the cells without further incubation. From the TGF β 1 standard curve included on every plate the effective concentration of TGF β 1 in the presence of the inhibitory ligands was determined by the reduction in luminescence measured. Some ligands were tested at both 3 pM and 6 pM TGF β 1 with the same results. To determine the IC $_{50}$ (the concentration of SELEX ligand necessary to reduce the TGF β 1 activity 50%), the five values obtained for each ligand were plotted and the value at 50% inhibition was determined graphically using Graphpad Prism assuming a hyperbolic fit of the data and using non-linear regression.

Example 2

RNA Ligands to TGF β 1

[0117] A. SELEX Experiments

[0118] In order to generate RNA ligands to TGF β 1, nine SELEX experiments, as summarized in Table 2, were performed using the methods described in Example 1. As shown in Table 1, the RNA pools differ in the number of random bases present in the central portion of the molecules: 40 nucleotides in the 40N6 (SEQ ID NO:2) SELEX and 64 nucleotides in the 64N6 and lib2-6-1RN6 (SEQ ID NOS:1, 3) SELEX experiment. Since the goal was to select RNA ligands that not only bound to TGF β 1 but also blocked receptor binding, the large random region (64N) was chosen. In two experiments, a shorter random region (40N) was also included. Ligands to TGF β 1 were very rare with 40N and were qualitatively the same as the 64N6 ligands selected.

[0119] The sequences of clones from the SELEX experiments are shown in Table 3 (SEQ ID NOS:12-42). The pyrimidines used in the various SELEX experiments differed at the 2' position of the sugar (Table 2). In the first two SELEX experiments, ligands were evolved as 2'-OH pyrimidines. Ligands were post-SELEX modified with 2'-NH₂ or 2'-F-substituted pyrimidines to see if they retained TGF β 1 binding. Since the 2' substitutions rendered the ligands resistant to RNase A they were also tested in the cell culture assay for inhibition of TGF β 1 activity. One such ligand lib2-6-1 (Group D, Table 3; SEQ ID NO:35) when substituted with 2'-NH₂-UTP and 2'-F-CTP was shown to inhibit TGF β 1 receptor mediated activity. To select more ligands, six more independent SELEX experiments (lib3-7 and lib9) were performed using the 2'-F and 2'-NH₂-substituted pyrimidines during the evolution process. In experiment lib8 the biologically active clone lib2-6-1 (SEQ ID NO:35) was randomized and subjected to re-selection to see if the binding and inhibition behavior of the clone could be improved. Lib8 was evolved as a 2'-OH pyrimidine RNA. In some cases, post-SELEX modification of TGF β 1 ligands derived from experiments 3-9 were performed, e.g., to determine if a ligand evolved with 2'-F pyrimidine substitutions would also bind with 2'-NH₂ substitutions.

[0120] Each starting pool for a SELEX experiment contained 3×10^{14} RNA molecules (500 pmol). The affinity of the starting RNA for TGF β 1 was estimated to be greater than 50 nM. After 4 rounds of SELEX, the affinities of the evolving pools had improved to approximately 10 nM and did not shift significantly in subsequent rounds. RNA was bulk sequenced and found to be non-random and cloned.

[0121] Lib1 took 20 rounds to evolve since optimum concentrations of TGF β 1 were not used until round 15 and libraries 5, 6 and 7 took longer to evolve because optimum conditions for recovery of bound ligands during the partitioning step in SELEX were not introduced until round 8. Optimum TGF β 1 concentrations and partitioning conditions are described in Example 1.

[0122] B. RNA Sequences

[0123] Many clones in a SELEX library are identical or similar in sequence. The libraries were screened by G track and only representatives of each G track type were tested in a binding assay. The binding assay was five points (16.5 nM, 5.5 nM, 1.8 nM, 0.6 nM, and 0.2 nM) and could detect only

those SELEX clones with a Kd less than or equal to 10 nM. RNA ligands that bound well (Kd < 10 nM) in the binding assay were sequenced. The sequences were inspected by eye and analyzed using computer programs which perform alignments and fold nucleic acid sequences to predict regions of secondary structure. Ligands were classified into five groups (A, B, C, D, and orphans) by sequence homology. Each group has characteristic allowable 2' substitutions.

[0124] 58 clones were identified by G track from 7 separate SELEX experiments to belong to group A ligands (Table 3; SEQ ID NOS:12-42). 15 clones were sequenced; 13 were similar but not identical, whereas 3 clones, lib3-13 (SEQ ID NO:12), lib5-6 and lib5-13, were identical. Group A ligands were recovered from seven of the eight SELEX libraries which included libraries evolved as 2'-NH₂, 2'-OH or 2'-F-substituted pyrimidines as well as a library evolved as 2'-F-UTP, 2'-NH₂-CTP. Post SELEX modification indicates that 2'-NH₂-UTP, 2'-F-CTP does not disrupt binding of lib8-9 to TGF β 1, thus the structure of Group A ligands appears to not require a specific 2' moiety on the pyrimidine sugar in order to maintain binding.

[0125] Group B ligands bind both as 2'-NH₂ and 2'-F pyrimidine substituted RNA. 28 Group B clones were detected by G track analysis from 3 libraries. Two of the libraries were evolved as 2'-NH₂ and one as 2'-F library. Four clones were sequenced, two were identical (lib5-47 and lib4-12; SEQ ID NO:28). An internal deletion can occur in group B, as in lib 3-44. The 40N orphan, lib3-42 was placed in Group B on the basis of secondary structure. The internal deletion in lib3-44, the binding affinity, the bioactivity and boundary experiments all support the placement of lib3-42 in this group.

[0126] Group C ligands bind as 2'-OH or 2'-F ligands as expected, since members of this group were evolved as 2'-OH ligands in lib 1 and as 2'-F pyrimidine substituted ligands in lib 6. Lib1-20-3 (SEQ ID NO:32) was post SELEX modified and as 2'-F derivative. Lib1-20-3 did not bind with 2'-NH₂ pyrimidines incorporated.

[0127] Group D ligand, lib2-6-1 (SEQ ID NO:35), was isolated after 2'-OH SELEX but was post SELEX modified and binds well as a 2'-NH₂-UTP and 2'-F-CTP pyrimidine derivative. Lib2-6-1 does not bind well to TGF β 1 with 2'-NH₂, 2'-F or 2'-F-UTP, 2'-NH₂-CTP-substituted pyrimidines. Variants of Group D were only reselected in two other SELEX experiments, lib8, a 2'-OH library, and lib 9, a 2'-NH₂-UTP, 2'-F CTP library, supporting the observation that there is specificity for the 2' pyrimidine position in this ligand.

[0128] The group labeled orphans are not homologous to each other and no variant sequences for these ligands have been determined. G track indicates that eight 40N clones similar to lib3-45 were isolated from two libraries. Two of the eight were sequenced and are identical. Lib3-45 (SEQ ID NO:39) binds whether it contains 2'-NH₂ or 2'-F substituted pyrimidines or the 2'-F-UTP, 2'-NH₂-CTP combination. Lib1-20-5 (SEQ ID NO:40) isolated as a 2'-OH ligand binds as a 2'-F, whereas lib1-20-12 (SEQ ID NO:41) and lib2-6-8 (SEQ ID NO:42) bind well only as 2'-OH pyrimidines and will not tolerate 2'-NH₂ or 2'-F post SELEX modifications.

[0129] As it was unusual that similar sequences were obtained from different SELEX experiments containing dif-

ferent modifications, another set of SELEX experiments was performed in search of RNA and ssDNA ligands to TGF β 1 as described in examples 5 and 6 infra.

Example 3

Inhibition of TGF β 1 Receptor Binding

[0130] The Kds and B_{max} values reported in Table 4 for Group A ligands are for the 2'-NH₂ substituted version of the ligand unless otherwise noted. B_{max} for the Group A ligands was 0.38±0.12 (n=14) which is in agreement with the measured retention of TGF β 1 on the nitrocellulose filters. The Kd's for Group A ligands were all similar, 2.2±1.1 nM (n=14). Where measured nitrocellulose and SA agarose bead binding assays gave equivalent results.

[0131] The IC₅₀'s in Table 4 for Group A ligands were all tested with the 2'-NH₂ pyrimidine substituted ligands except where indicated. 2'-NH₂ ligands were used in the tissue culture bio-assay since they exhibited the greatest stability under the conditions of the bio-assay. Five out of ten Group A ligands tested inhibited TGF β 1 receptor activity. IC₅₀ values for the inhibitors were typically 25 fold above the Kd for TGF β 1. The data are reproducible; the Kd for ligand lib3-13 was 0.83±0.11 nM (n=3) and the IC₅₀ for lib3-13 (SEQ ID NO:12) was 25±14 nM (n=4). RNA concentrations in the bioassays are all estimates based on an assumed extinction coefficient and 100% purity of the ligand. The RNA concentrations may, therefore, be overestimated during the bio-assay which in turn would overestimate the IC₅₀.

[0132] Another five Group A ligands did not inhibit TGF β receptor binding activity. One obvious difference between the non-bioactive ligands, lib2-6-4 (SEQ ID NO:20), lib5-48 (SEQ ID NO:19), and lib6-23 (SEQ ID NO:21), and the bioactive ligands is the substitution at nucleotide 72. Lib7-21 (SEQ ID NO:23) and lib7-43 (SEQ ID NO:24) were tested as 2'-F-UTP, 2'-NH₂-CTP ligands for bio-activity. These ligands were not bio-active despite their high affinity to TGF β . In conclusion, binding and bioactivity are separable functions of the TGF β Group A ligands.

[0133] Group B ligands have different binding properties than Group A ligands (Table 4). Both the Kd (0.63±0.5 nM, n=4) and B_{max} (0.14±0.04, n=4) are lower for Group B ligands. One Group B inhibitor, lib4-12 (SEQ ID NO:28), actually appears to stimulate TGF β 1 activity in the tissue culture bio-assay at low concentrations. The basis of this mixed agonist/antagonist behavior has not been determined. The best inhibitor in this group, lib3-42 (SEQ ID NO:30) has an IC₅₀ of 22 nM and had no agonist behavior over the concentration ranges tested.

[0134] Group C ligands were tested as 2'-F derivatives and were not bio-active. Neither was the 2'-F orphan lib1-20-5 (SEQ ID NO:40). The 2'-NH₂, 40N orphan, lib3-45 is an example of another ligand with high affinity for TGF β 1 and no ability to inhibit TGF β 1 receptor binding.

[0135] Group D ligands were tested in the bio-assay as 2'-NH₂-UTP, 2'-F-CTP derivatives. Both lib2-6-1 (SEQ ID NO:35) and the truncated version lib2-6-1-81 (SEQ ID NO:36) can inhibit TGF β 1 receptor binding; however, a single mutation from a C to a G at position 53 decreases bio-activity in clone lib8-23. Similarly a 2 base pair deletion in clone lib6-30 (SEQ ID NO:34) at positions corresponding

to nucleotides 67 and 68 in lib2-6-1 (SEQ ID NO:35) increases binding by 10 fold but eliminates bio-activity.

[0136] Lib2-6-1 (SEQ ID NO:35) was shown to be fully effective only against TGF β 1 and not TGF β 2 and TGF β 3. Lib2-6-1 (SEQ ID NO:35) was biologically active in the presence of 10% horse serum in the cell culture medium in addition to the 0.1% BSA. Thus the ligand demonstrates specificity towards TGF β 1 which is not interfered with by the presence of the horse serum in this assay. The biggest indication that the inhibition of TGF β 1 receptor binding is a specific phenomenon is the fact that not all TGF β 1 ligands block receptor binding, but the ones that do, do so reproducibly. There are no examples of ligands that do not bind to TGF β 1 blocking TGF β 1 receptor binding activity.

[0137] In summary, RNA ligands that can block TGF β 1 receptor binding are a subset of ligands. Binding is necessary but not sufficient for bio-activity. Roughly 50% of the high affinity ligands tested were inhibitors. Of the inhibitors, 30% were good inhibitors (IC₅₀<25 nM).

Example 4

Boundary Analysis

[0138] Truncation experiments were done on a number of TGF β 1 ligands to determine the nucleotides essential for binding. Group A ligands, lib3-13 (SEQ ID NO:12) and lib8-9 (SEQ ID NO:16), were truncated with consistent results. The fragment lib3-13-79 binds to TGF β 1, thus none of the nucleotides 3' to nucleotide 79 in lib3-13 are essential for binding. Similarly when all nucleotides 5' to nucleotide 38 are deleted the remaining fragment, lib3-13-(38-123) can still bind to TGF β 1. The 5' boundary is in agreement with the sequence lib6-23 (SEQ ID NO:21), which has a deletion corresponding to nucleotides 19-36 of lib3-13 (SEQ ID NO:25), and still binds to TGF β 1. Thus, all high affinity binding determinants for Group A clones may lie wholly within the random region and may correspond to a 42 nucleotide fragment, lib3-13-(38-79). Many Group A ligands contain deletions or substitutions within the predicted essential binding domain, in the region corresponding to lib3-13-(72-81). The deletion and substitution in lib4-32 have no effect on its 3' boundary which corresponds to lib3-13 nucleotide 80. Thus, the 3' boundary is probably correct and the alterations in nucleotide sequence 72-81 are ones that do not significantly alter the nucleic acid structure required for binding. Mutations in this region, most notably nucleotide 72 may, however, modify the ability of the ligand to block TGF β 1 receptor binding as noted earlier.

[0139] Boundary analysis of the 3' end of Group B ligand, lib4-12 (SEQ ID NO:28), predicts that nothing beyond nucleotide 72 is required for TGF β 1 binding. When the 5' boundary of lib4-12 was determined, all but the first three nucleotides were required for binding, indicating that the 5' constant region is an essential part of the ligand at least when the boundary of the full length ligand was determined. Assuming that ligand lib3-44 (SEQ ID NO:29) has a similar binding determinant as lib4-12 (SEQ ID NO:28), we can also conclude that nucleotides 37-46 of lib4-12 are not required for binding since these are deleted in lib3-44 and lib3-42 (SEQ ID NO:30).

[0140] The 3' constant region is not necessary for binding in Group C and D ligands. Both ligand types bind without

the 3' nucleotides in the constant region. Lib1-20-3-82, an 82 nucleotide truncated version of lib1-20-3 (SEQ ID NO:32), binds as well as the full length lib1-20-3. Likewise binding and bioactivity of lib2-6-1 is unaffected by the 3' truncation found in lib2-6-1-81 (SEQ ID NO:36).

Example 5

Experimental Procedures

[0141] In the preferred embodiment, a second set of SELEX experiments was performed in search of RNA and DNA ligands with specific high affinity for TGF β 1 from degenerate libraries containing 40 random positions (40N). This Example provides the general procedures followed and incorporated in Example 6.

[0142] A. Materials.

[0143] M-MLV superscript reverse transcriptase was purchased from Gibco BRL (Gaithersburg, Md.). T7 RNA polymerase was purified according to standard procedures at NeXstar Pharmaceuticals, Inc. (Boulder, Colo.). Taq DNA polymerase (Amplitaq) was from Perkin Elmer/Cetus (Richmond, Calif.). T4 polynucleotide kinase, DNA polymerase (Klenow fragment), and alkaline phosphatase were purchased from New England Biolabs, Inc. (Beverly, Mass.). The 2'-amino substituted nucleotide triphosphates amino-UTP and amino-CTP were synthesized according to standard procedures at NeXstar Pharmaceuticals, Inc. (Boulder, Colo.). Other reagents used in this work were of the highest quality obtainable.

[0144] B. Nucleic Acids.

[0145] RNAs were synthesized by in vitro transcription using double-stranded DNA oligonucleotides and T7 RNA polymerase. DNA oligonucleotides (Table 5) were purchased from Operon, Inc. (Alameda, Calif.) and purified by 6% preparative polyacrylamide gel electrophoresis. PCR amplification was performed in 50 mM KCl, 10 mM Tris-HCl (pH 8.6), 2.5 mM MgCl₂, 170 mg/mL BSA, and dNTPs (present at 1 mM each). Taq DNA polymerase was used at 100 units per 0.1 mL reaction, and the 5'- and 3'-primers were present at 1 mM. Transcription was performed in 40 mM NaCl, 10 mM dithiothreitol, 50 mM Tris-acetate (pH 8.0), 8 mM magnesium acetate, 2 mM spermidine, and 2 mM NTP. T7 RNA polymerase was present at 1 unit/mL. The reaction was incubated at 28 degrees for 16 hours and then treated with 20 units of DNase I for an additional 10 min at 37 degrees. The reaction was stopped by the addition of one half volume of loading buffer (93% formamide, 10 mM EDTA, pH 8.0) and heated to 95 degrees for 3 min prior to electrophoresis on a 6% polyacrylamide/8 M urea denaturing gel. The RNA transcript was visualized by UV shadowing and was excised from the gel and eluted into TE buffer (10 mM Tris-acetate pH 8.0, 2 mM EDTA). The RNA transcript was ethanol precipitated, dried under vacuum, and redissolved in distilled H₂O. The concentration of RNA as well as single-stranded DNA was quantified by measuring the A₂₆₀ and assuming that 1 A₂₆₀ unit equaled 40 mg/mL and 33 mg/mL, respectively.

[0146] C. Evolution of High-Affinity Ligands.

[0147] SELEX ligands that bind to TGF β 1 were derived essentially as described in U.S. Pat. No. 5,270,163 (see also Tuerk and Gold (1990) Science 249:505-510) using the

oligonucleotides illustrated in Table 5 (SEQ ID NOS:43-54). The DNA templates contained a 40-nucleotide (40N) variable sequence generated by mixed-nucleotide DNA synthesis, as well as 5'- and 3'-fixed sequences, necessary for PCR amplification of the template. The 5'-fixed sequence of oligonucleotides 40N7 (SEQ ID NO:43) and 40N8 (SEQ ID NO:49) also contained a T7 RNA polymerase promoter. RNA for the first round of RNA SELEX was transcribed from double-stranded DNA templates generated by primer extension on single-stranded DNA templates 40N7 and 40N8 with the Klenow fragment of DNA polymerase I. RNA SELEX consisted of up to 15 rounds of RNA synthesis, binding to target, partitioning of bound and unbound RNA by nitrocellulose filtration, cDNA synthesis, and PCR amplification to regenerate the double-stranded DNA template. Binding to the target by the RNA pool was performed in binding buffer A (120 mM NaCl, 2.5 mM KCl, 0.12 mM MgSO₄, 40 mM HEPES, 20 mM NaH₂PO₄/Na₂HPO₄ pH 7.4, 0.01% HSA) at 37 degrees for at least 10 min prior to filtration. In contrast, the first round of single-stranded DNA SELEX was performed by using the synthetically synthesized oligonucleotides 40D7 and 40D8 directly. SELEX consisted of 25 rounds of binding to target, partitioning of bound and unbound single-stranded DNA by nitrocellulose filtration, PCR amplification to generate a double-stranded DNA population, and preparative polyacrylamide gel electrophoresis to purify single-stranded DNA for the next round of SELEX. Binding of the target to the single-stranded DNA pool was performed in binding buffer B (150 mM NaCl, 10 mM Tris-acetate pH 7.5, 0.001% BSA) at 37 degrees for at least 15 min prior to filtration. Radiolabeling of RNA as well as DNA repertoires was performed by incubation of 5 picomoles nucleic acid, 2 units of T4 polynucleotide kinase, and 6 mL [³²P] ATP (800 Ci/mmol) in a volume of 10 mL at 37 degrees for 30 min. The concentration of nucleic acid at each round of the SELEX experiment varied between 1500 nM and 1 nM while the concentration of the target TGF- β 1 varied between 150 nM and 0.03 nM.

[0148] D. Cloning and Sequencing of Ligands.

[0149] Cloning of the nucleic acid repertoire was performed as described by Tuerk and Gold (1990) Science 249:505-510 using double-stranded DNA that was generated from the RNA repertoire by PCR amplification. PCR-amplified DNA was digested with the restriction enzymes SphI and HindIII and ligated into compatible sites within pGEM. Ligated plasmids were transformed into *E. coli* and plated onto LB agar containing 5-bromo-4-chloro-3-indolyl β -D-galactoside, isopropyl thiogalactoside, and 100 mg/mL ampicillin. Colonies not expressing β -galactosidase were analyzed. Sequencing of DNA was performed as described by Tuerk and Gold (1990) using the dideoxynucleotide procedure of Sanger et al. (1977) *Proc. Natl. Acad. Sci. USA* 74:5463-5467. Plasmids were isolated from *E. coli* by the alkaline lysis miniprep procedure (Manitatis et al. (1982) in *Molecular Cloning: A laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). DNA was incubated in 50 mM Tris-HCl (pH 8.3), 60 mM NaCl, 6 mM magnesium acetate, and 1 mM DTT with 0.4 mM dNTP and 0.2 mM dideoxy-NTP for 20 min at 48 degrees. DNA polymerase was present at 4 units per reaction. The reactions were stopped by the addition of 10 mL of loading buffer and heated to 95 degrees for 3 min prior to gel electrophoresis on a 6% polyacrylamide/8 M urea denaturing gel. G-track sequencing was performed as described and

provided a convenient method to quickly screen the cloned library for ligands of different sequence. Briefly, the G-track sequencing reaction contained 50 mM Tris-HCl (pH 8.3), 60 mM NaCl, 6 mM magnesium acetate, and 1 mM DTT with 0.4 mM dNTP, 0.2 mM dideoxy-GTP, and 4 units of DNA polymerase. The reaction was performed at 48 degrees for 20 min and was stopped by the addition of 10 μ L of loading buffer and heated to 95 degrees for 3 min prior to gel electrophoresis on a 6% polyacrylamide/8 M urea denaturing gel.

Example 6

Binding Analysis, Bioassay Results, and Sequences of a ssDNA Library

[0150] Binding analysis of the 40D7 DNA library for TGF- β 1 is shown in FIG. 1. Binding data obtained from round 19 (triangles) and round 0 (circles) are shown. The experiment was performed by incubating nucleic acid (less than 1 nM) and the indicated concentration of TGF- β 1 in Binding Buffer (150 mM NaCl, 10 mM Tris-acetate pH 8.2, 0.001% BSA) for 15 minutes at 37 degrees in a volume of 0.1 mL. Samples were filtered through nitrocellulose and were immediately followed by 3 mL of TE Buffer (10 mM Tris-acetate pH 8.0, 0.1 mM EDTA). The percentage of radiolabel bound was calculated from the amount of radiolabel retained on the nitrocellulose filter and the total radiolabel added to the binding reaction. The results show that the apparent K_d of the 40D7 library is 1 nM, whereas the starting pool has an apparent K_d of 30 nM. Thus, the 40D7 library shows an increase of about three fold in binding.

[0151] A PAI-luciferase assay to detect TGF- β 1 activity in the presence of the nucleic acid libraries generated in Example 5 was performed as described in Abe et al. (1994) *Analytical Biochem.* 216:276-284. Mink lung epithelial cells containing the PAI-luciferase reporter gene were incubated with TGF- β 1 (10 pM) and oligonucleotides from the DNA libraries or anti-TGF- β antibody (60 μ g/mL). The mink lung epithelial cells were incubated for 18 hours and oligonucleotides were pre-incubated with TGF- β 1 before the assay and read after 8 hours. Addition of oligonucleotides alone (100 nM) to the cell culture did not affect the assay (data not shown). The identity of the oligonucleotide libraries as well as their effect on luciferase activity is indicated in FIG. 2. The ssDNA library 40N7 completely inhibited the activity of TGF- β 1, while the control (an equal concentration of randomized nucleic acid) showed a small stimulation of TGF- β 1 activity.

[0152] Based on the results of the binding analysis and PAI-luciferase assay, DNA ligands from the 40N7 library were sequenced as described in Example 5. The sequences are shown in Table 6 (SEQ ID NOS:55-89). As the DNA 40N7 library showed inhibition in the PAI-luciferase bioassay, it is reasonable to suggest that the individual clones from the library are TGF β 1 binders.

Example 7

Experimental Procedures

[0153] This Example provides the general procedures followed and incorporated in Examples 8-15 for the evolution of nucleic acid ligands to PDGF.

[0154] A. Materials.

[0155] Recombinant human PDGF-AA (Mr=29,000), PDGF-AB (Mr=27,000) and PDGF-BB (Mr=25,000) were purchased from R&D Systems (Minneapolis, Minn.) in lyophilized form, free from carrier protein. All three isoforms were produced in *E. coli* from synthetic genes based on the sequences for the long form of the mature human PDGF A-chain (Betsholtz et al., (1986) *Nature* 320: 695-699) and the naturally occurring mature form of human PDGF B-chain (Johnsson et al., (1984) *EMBO J.* 3: 921-928). Randomized DNA libraries, PCR primers and DNA ligands and 5'-iodo-2'-deoxyuridine-substituted DNA ligands were synthesized by NeXstar Pharmaceuticals, Inc. (Boulder, Colo.) or by Operon Technologies (Alameda, Calif.) using the standard solid phase phosphoramidite method (Sinha et al., (1984) *Nucleic Acids Res.* 12:4539-4557).

[0156] B. Single Stranded DNA (ssDNA) Selex

[0157] Essential features of the SELEX procedure have been described in detail in the SELEX Patent Applications (see also, Tuerk and Gold, *Science* 249:505 (1990); Jellinek et al., *Biochemistry* 33:10450 (1994); Jellinek et al., *Proc. Natl. Acad. Sci.* 90:11227 (1993)), which are incorporated by reference herein. The initial ssDNA library containing a contiguous randomized region of forty nucleotides, flanked by primer annealing regions (Table 7; SEQ ID NO:90) of invariant sequence, was synthesized by the solid phase phosphoramidite method using equal molar mixture of the four phosphoramidites to generate the randomized positions. The ssDNA library was purified by electrophoresis on an 8% polyacrylamide/7 M urea gel. The band that corresponds to the full-length DNA was visualized under UV light, excised from the gel, eluted by the crush and soak method, ethanol precipitated and pelleted by centrifugation. The pellet was dried under vacuum and resuspended in phosphate buffered saline supplemented with 1 mM MgCl₂ (PBSM=10.1 mM Na₂HPO₄, 1.8 mM KH₂PO₄, 137 mM NaCl and 2.7 mM KCl, 1 mM MgCl₂, pH 7.4) buffer. Prior to incubation with the protein, the ssDNA was heated at 90° C. for 2 minutes in PBSM and cooled on ice. The first selection was initiated by incubating approximately 500 pmol (3 \times 10¹⁴ molecules) of 5³²P end-labeled random ssDNA with PDGF-AB in binding buffer (PBSM containing 0.01% human serum albumin (HSA)). The mixture was incubated at 4° C. overnight, followed by a brief (15 min) incubation at 37° C. The DNA bound to PDGF-AB was separated from unbound DNA by electrophoresis on an 8% polyacrylamide gel (1:30 bis-acrylamide:acrylamide) at 4° C. and at 5 V/cm with 89 mM Tris-borate (pH 8.3) containing 2 mM EDTA as the running buffer. The band that corresponds to the PDGF-ssDNA complex, which runs with about half the electrophoretic mobility of the free ssDNA, was visualized by autoradiography, excised from the gel and eluted by the crush and soak method. In subsequent affinity selections, the ssDNA was incubated with PDGF-AB for 15 minutes at 37° C. in binding buffer and the PDGF-bound ssDNA was separated from the unbound DNA by nitrocellulose filtration, as previously described (Green, et al., (1995) *Chemistry and Biology* 2, 683-695). All affinity-selected ssDNA pools were amplified by PCR in which the DNA was subjected to 12-20 rounds of thermal cycling (30 s at 93° C., 10 s at 52° C., 60 s at 72° C.) in 10 mM Tris-Cl (pH 8.4) containing 50 mM KCl, 7.5 mM MgCl₂, 0.05 mg/ml bovine serum albumin, 1

mM deoxynucleoside triphosphates, 5 μ M primers (Table 7) and 0.1 units/ μ l Taq polymerase. The 5' PCR primer was 5' end-labeled with polynucleotide kinase and [γ - 32 P]ATP and the 3' PCR primer was biotinylated at the 5' end using biotin phosphoramidite (Glen Research, Sterling, Va.). Following PCR amplification, streptavidin (Pierce, Rockford, Ill.) was added to the unpurified PCR reaction mixture at a 10-fold molar excess over the biotinylated primer and incubated for 15 min at room temperature. The dsDNA was denatured by adding an equal volume of stop solution (90% formamide, 1% sodium dodecyl sulfate, 0.025% bromophenol blue and xylene cyanol) and incubating for 20 min at room temperature. The radiolabeled strand was separated from the streptavidin-bound biotinylated strand by electrophoresis on 12% polyacrylamide/7M urea gels. The faster migrating radiolabeled (non-biotinylated) ssDNA strand was cut out of the gel and recovered as described above. The amount of ssDNA was estimated from the absorbance at 260 nm using the extinction coefficient of 33 μ g/ml/absorbance unit (Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual*, 2 Ed. 3 vols., Cold Spring Harbor Laboratory Press, Cold Spring Harbor).

[0158] C. Cloning and Sequencing.

[0159] The amplified affinity-enriched pool from SELEX round 12 was purified on a 12% polyacrylamide gel and cloned between HindIII and PstI sites in JM109 strain of *E. coli* (Sambrook, et al., (1989) *Molecular Cloning: A Laboratory Manual*, 2 Ed. 3 vols., Cold Spring Harbor Laboratory Press, Cold Spring Harbor). Individual clones were used to prepare plasmids by alkaline lysis. Plasmids were sequenced at the insert region using the forward sequencing primer and Sequenase 2.0 (Amersham, Arlington Heights, Ill.) according to the manufacturer's protocol.

[0160] D. Determination of the Apparent Equilibrium Dissociation Constants and the Dissociation Rate Constants.

[0161] The binding of ssDNA ligands at low concentrations to varying concentrations of PDGF was determined by the nitrocellulose filter binding method as described (Green et al., (1995) *Chemistry and Biology* 2:683-695). The concentrations of PDGF stock solutions (in PBS) were determined from the absorbance readings at 280 nm using the following ϵ_{280} values calculated from the amino acid sequences (Gill, S. C. and von Hippel, P. H. (1989) *Anal. Biochem.* 182:319-326): 19,500 $M^{-1}cm^{-1}$ for PDGF-AA, 15,700 $M^{-1}cm^{-1}$ for PDGF-AB and 11,800 $M^{-1}cm^{-1}$ for PDGF-BB. ssDNA for all binding experiments were purified by electrophoresis on 8% (>80 nucleotides) or 12% (<40 nucleotides) polyacrylamide/7 M urea gels. All ssDNA ligands were heated at 90° C. in binding buffer at high dilution (\approx 1 nM) for 2 min and cooled on ice prior to further dilution into the protein solution. The binding mixtures were typically incubated for 15 min at 37° C. before partitioning on nitrocellulose filters.

[0162] The binding of DNA ligands (L) to PDGF-AA (P) is adequately described with the bimolecular binding model for which the fraction of bound DNA at equilibrium (q) is given by eq. 1,

$$q = \frac{f/2[L]_t \{ [P]_t + [L]_t + K_d + \sqrt{([P]_t + [L]_t + K_d)^2 - 4[P]_t} \}}{[L]_t} \quad (1)$$

where $[P]_t$ and $[L]_t$ are total protein and total DNA concentrations, K_d is the equilibrium dissociation constant and f is

the efficiency of retention of protein-DNA complexes on nitrocellulose filters (Irvine et al., (1991) *J. Mol. Biol.* 222:739-761; Jellinek et al., (1993) *Proc. Nat'l. Acad. Sci. USA* 90:11227-11231).

[0163] The binding of DNA ligands to PDGF-AB and PDGF-BB is biphasic and can be described by a model in which the DNA ligand is composed of two non-interconverting components (L_1 and L_2) that bind to the protein with different affinities, described by corresponding dissociation constants, K_{d1} and K_{d2} (Jellinek et al., (1993) *Proc. Nat'l. Acad. Sci. USA* 90:11227-11231). In this case, the explicit solution for the fraction of bound DNA (q) is given by eq. 2,

$$q = f \left(\frac{\chi_1 K_{d1}}{1 + K_{d1}[P]} + \frac{\chi_2 K_{d2}}{1 + K_{d2}[P]} \right)^{[P]_t} \quad (2)$$

with

$$[P]_t = \frac{[P]_i}{1 + \frac{\chi_1 K_{d1} [L]_i}{1 + K_{d1}[P]} + \frac{\chi_2 K_{d2} [L]_i}{1 + K_{d2}[P]}}$$

where χ_1 and $\chi_2 (=1-\chi_1)$ are the mole fractions of L_1 and L_2 . The K_d values for the binding of DNA ligands to PDGF were calculated by fitting the data points to eq. 1 (for PDGF-AA) or eq. 2 (for PDGF-AB and PDGF-BB) using the non-linear least squares method.

[0164] The dissociation rate constants (k_{off}) were determined by measuring the amount of 32 P 5'-end labeled minimal ligands (0.17 nM) bound to PDGF-AB (1 nM) as a function of time following the addition of 500-fold excess of unlabeled ligands, using nitrocellulose filter binding as the partitioning method. The k_{off} values were determined by fitting the data points to the first-order rate equation (eq. 3)

$$(q - q_\infty) / (q_0 - q_\infty) = \exp(-k_{off}t) \quad (3)$$

where q, q_0 and q_∞ represent the fractions of DNA bound to PDGF-AB at any time (t), t=0 and t= ∞ , respectively.

[0165] E. Minimal Ligand Determinations.

[0166] To generate a population of 5' end-labeled DNA ligands serially truncated from the 3' end, a primer complementary to the 3' invariant sequence region of a DNA ligand template (truncated primer 5N2, Table 7; SEQ ID NO:92) was radiolabeled at the 5' end with [γ - 32 P]-ATP and T4 polynucleotide kinase, annealed to the template and extended with Sequenase (Amersham, Arlington Heights, Ill.) and a mixture of all four dNTPs and ddNTPs. Following incubation in binding buffer for 15 min at 37° C., the fragments from this population that retain high affinity binding to PDGF-AB were separated from those with weaker affinity by nitrocellulose filter partitioning. Electrophoretic resolution of the fragments on 8% polyacrylamide/7 M urea gels, before and after affinity selection, allows determination of the 3' boundary. To generate a population of 3' end-labeled DNA ligands serially truncated from the 5' end, the DNA ligands were radiolabeled at the 3' end with [α - 32 P]-cordycepin-5'-triphosphate (New England Nuclear, Boston, Mass.) and T4 RNA ligase (Promega, Madison, Wis.), phosphorylated at the 5' end with ATP and T4 polynucleotide kinase, and partially digested with lambda exonuclease (Gibco BRL, Gaithersburg, Md.). Par-

tial digestion of 10 pmols of 3'-labeled ligand was done in 100 μ L volume with 7 mM glycine-KOH (pH 9.4), 2.5 mM $MgCl_2$, 1 μ g/ml BSA, 15 μ g tRNA, and 4 units of lambda exonuclease for 15 min at 37°. The 5' boundary was determined in an analogous manner to that described for the 3' boundary.

[0167] F. Melting Temperature (T_m) Measurements.

[0168] Melting profiles for the minimal DNA ligands were obtained on a Cary Model 1 E spectrophotometer. Oligonucleotides (320-400 nM) were heated to 95° C. in PBS, PBSM or PBS with 1 mM EDTA and cooled to room temperature prior to the melting profile determination. Melting profiles were generated by heating the samples at the rate of 1° C./min from 15-95° C. and recording the absorbance every 0.1° C. The first derivative of the data points was calculated using the plotting program KaleidaGraph (Synergy Software, Reading, Pa.). The first derivative values were smoothed using a 55 point smoothing function by averaging each point with 27 data points on each side. The peak of the smoothed first derivative curves was used to estimate the T_m values.

[0169] G. Crosslinking of 5-iodo-2'-deoxyuridine-substituted DNA Ligands to PDGF-AB.

[0170] DNA ligands containing single or multiple substitutions of 5'-iodo-2'-deoxyuridine for thymidine were synthesized using the solid phase phosphoramidite method. To test for the ability to crosslink, trace amounts of 5³²P end-labeled ligands were incubated with PDGF-AB (100 nM) in binding buffer at 37° C. for 15 min prior to irradiation. The binding mixture was transferred to a 1 cm path length cuvette thermostated at 37° C. and irradiated at 308 nm for 25-400 s at 20 Hz using a XeCl charged Lumonics Model EX748 excimer laser. The cuvette was positioned 24 cm beyond the focal point of a convergent lens, with the energy at the focal point measuring 175 mJoules/pulse. Following irradiation, aliquots were mixed with an equal volume of formamide loading buffer containing 0.1% SDS and incubated at 95° for 5 min prior to resolution of the crosslinked PDGF/ligand complex from the free ligand on 8% polyacrylamide/7 M urea gels.

[0171] To identify the protein site of crosslinking for ligand 20 t-14, binding and irradiation were done on a larger scale. PDGF-AB and 5³²P end-labeled ligand, each at 1 μ M in PBSM, were incubated and irradiated (300 s) as described above in two 1 ml reaction vessels. The reaction mixtures were combined, ethanol precipitated and resuspended in 0.3 ml of Tris-HCl buffer (100 mM, pH 8.5). The PDGF-AB/ligand crosslinked complex was digested with 0.17 μ g/ μ l of modified trypsin (Boehringer Mannheim) for 20 hours at 37° C. The digest mixture was extracted with phenol/chloroform, chloroform and then ethanol precipitated. The pellet was resuspended in water and an equal volume of formamide loading buffer with 5% (v/v) β -mercaptoethanol (no SDS), incubated at 95° C. for 5 min, and resolved on a 40 cm 8% polyacrylamide/7 M urea gel. The crosslinked tryptic-peptide/ligand that migrated as two closely spaced bands about 1.5 cm above the free ligand band was excised from the gel and eluted by the crush and soak method and ethanol precipitated. The dried crosslinked peptide (about 160 pmoles based on the specific activity) was sequenced by Edman degradation (Midwest Analytical, Inc., St. Louis, Mo.).

[0172] H. Receptor Binding Assay.

[0173] The binding of ¹²⁵I-PDGF-AA and ¹²⁵I-PDGF-BB to porcine aortic endothelial (PAE) cells transfected with PDGF α - or β -receptors were performed as described (Heldin et al., (1988) *EMBO J.* 7:1387-1394). Different concentrations of DNA ligands were added to the cell culture (1.5 cm²) in 0.2 ml of phosphate buffered saline supplemented with 1 mg bovine serum albumin per ml together with ¹²⁵I-PDGF-AA (2 ng, 100,000 cpm) or ¹²⁵I-PDGF-BB (2 ng, 100,000 cpm). After incubation at 4° C. for 90 min, the cell cultures were washed and cell associated radioactivity determined in a γ -counter (Heldin et al., (1988) *EMBO J.* 7:1387-1394).

[0174] I. [³H]thymidine Incorporation Assay.

[0175] The incorporation of [³H]thymidine into PAE cells expressing PDGF β -receptor in response to 20 ng/ml of PDGF-BB or 10% fetal calf serum and in the presence of different concentrations of DNA ligands was performed as described (Mori et al., (1991) *J. Biol. Chem.* 266:21158-21164). After incubation for 24 h at 37° C., ³H-radioactivity incorporated into DNA was determined using a β -counter.

Example 8

ssDNA Ligands of PDGF

[0176] High affinity DNA ligands to PDGF AB were identified by the SELEX process from a library of $\approx 3 \times 10^{14}$ molecules (500 pmol) of single stranded DNA randomized at forty contiguous positions (Table 7; SEQ ID NO:90). The PDGF-bound DNA was separated from unbound DNA by polyacrylamide gel electrophoresis in the first round and by nitrocellulose filter binding in the subsequent rounds. After 12 rounds of SELEX, the affinity-enriched pool bound to PDGF-AB with an apparent dissociation constant (K_d) of ≈ 50 pM (data not shown). This represented an improvement in affinity of ≈ 700 -fold compared to the initial randomized DNA library. This affinity-enriched pool was used to generate a cloning library from which 39 isolates were sequenced. Thirty-two of these ligands were found to have unique sequences (Table 8; SEQ ID NOS:93-124). Ligands that were subjected to the minimal sequence determination are marked with an asterisk (*) next to the clone number. The clone numbers that were found to retain high affinity binding as minimal ligands are italicized. All ligands shown in Table 8 were screened for their ability to bind to PDGF AB using the nitrocellulose filter binding method. To identify the best ligands from this group, we determined their relative affinities for PDGF-AB by measuring the fraction of 5³²P end-labeled ligands bound to PDGF-AB over a range of protein concentrations. For the ligands that bound to PDGF-AB with high affinity, the affinity toward PDGF-BB and PDGF-AA was also examined: in all cases, the affinity of ligands for PDGF-AB and PDGF-BB was comparable while the affinity for PDGF-AA was considerably lower (data not shown).

[0177] Twenty-one of the thirty-two unique ligands can be grouped into a sequence family shown in Table 9. The sequences of the initially randomized region (uppercase letters) are aligned according to the consensus three-way helix junction motif. Nucleotides in the sequence-invariant region (lowercase letters) are only shown where they participate in the predicted secondary structure. Several ligands

were “disconnected” (equality symbol) in order to show their relatedness to the consensus motif through circular permutation. The nucleotides predicted to participate in base pairing are indicated with underline inverted arrows, with the arrow heads pointing toward the helix junction. The sequences are divided into two groups, A and B, based on the first single stranded nucleotide (from the 5' end) at the helix junction (A or G, between helices II and III). Mismatches in the helical regions are shown with dots under the corresponding letters (G-T and T-G base pairs were allowed). In places where single nucleotide bulges occur, the mismatched nucleotide is shown above the rest of the sequence between its neighbors.

[0178] This classification is based in part on sequence homology among these ligands, but in greater part on the basis of a shared secondary structure motif: a three-way helix junction with a three nucleotide loop at the branch point (FIG. 3). These ligands were subdivided into two groups; for ligands in group A, the loop at the branch point has an invariant sequence AGC and in group B, that sequence is G(T/G)(C/T). The proposed consensus secondary structure motif is supported by base-pairing covariation at non-conserved nucleotides in the helices (Table 10). Since the three-way junctions are encoded in continuous DNA strands, two of the helices end in loops at the distal end from the junction. These loops are highly variable, both in length and in sequence. Furthermore, through circular permutation of the consensus motif, the loops occur in all three helices, although they are most frequent in helices II and III. Together these observations suggest that the regions distal from the helix junction are not important for high affinity binding to PDGF-AB. The highly conserved nucleotides are indeed found near the helix junction (Table 9, FIG. 3).

Example 9

Boundary Analysis

[0179] The minimal sequence necessary for high affinity binding was determined for the six best ligands to PDGF-AB. In general, the information about the 3' and 5' minimal sequence boundaries can be obtained by partially fragmenting the nucleic acid ligand and then selecting for the fragments that retain high affinity for the target. With RNA ligands, the fragments can be conveniently generated by mild alkaline hydrolysis (Tuerk et al., (1990) *J. Mol. Biol.* 213:749-761; Jellinek et al., (1994) *Biochemistry* 33:10450-10456; Jellinek et al., (1995) *Biochemistry* 34:11363-11372; Green et al., (1995) *J. Mol. Biol.* 247:60-68). Since DNA is more resistant to base, an alternative method of generating fragments is needed for DNA. To determine the 3' boundary, a population of ligand fragments serially truncated at the 3' end was generated by extending the 5' end-labeled primer annealed to the 3' invariant sequence of a DNA ligand using the dideoxy sequencing method. This population was affinity-selected by nitrocellulose filtration and the shortest fragments (truncated from the 3' end) that retain high affinity binding for PDGF-AB were identified by polyacrylamide gel electrophoresis. The 5' boundary was determined in an analogous manner except that a population of 3' end-labeled ligand fragments serially truncated at the 5' end was generated by limited digestion with lambda exonuclease. The minimal ligand is then defined as the sequence between the two boundaries. It is important to keep in mind that, while

the information derived from these experiments is useful, the suggested boundaries are by no means absolute since the boundaries are examined one terminus at a time. The untruncated (radiolabeled) termini can augment, reduce or have no effect on binding (Jellinek et al., (1994) *Biochemistry* 33:10450-10456).

[0180] Of the six minimal ligands for which the boundaries were determined experimentally, two (20 t (SEQ ID NO:172) and 41 t (SEQ ID NO:174); truncated versions of ligands 20 and 41) bound with affinities comparable (within a factor of 2) to their full-length analogs and four had considerably lower affinities. The two minimal ligands that retained high affinity binding to PDGF, 20 t and 41 t, contain the predicted three-way helix junction secondary structure motif (FIG. 4). The sequence of the third minimal ligand that binds to PDGF-AB with high affinity, 36 t (SEQ ID NO: 173), was deduced from the knowledge of the consensus motif (FIG. 4). In subsequent experiments, we found that the single-stranded region at the 5' end of ligand 20 t is not important for high affinity binding. Furthermore, the trinucleotide loops on helices II and III in ligand 36 t (GCA and CCA) can be replaced with 20 pentaethylene glycol spacers (infra). These experiments provide further support for the importance of the helix junction region in high affinity binding to PDGF-AB.

[0181] The binding of minimal ligands 20 t, 36 t, and 41 t to varying concentrations of PDGF-AA, PDGF-AB and PDGF-BB is shown in FIGS. 5A, 5B and 5C. In agreement with the binding properties of their full length analogs, the minimal ligands 25 bind to PDGF-AB and PDGF-BB with substantially higher affinity than to PDGF AA (FIGS. 5A, 5B and 5C, Table 11). In fact, their affinity for PDGF-AA is comparable to that of random DNA (data not shown). The binding to PDGF-AA is adequately described with a monophasic binding equation while the binding to PDGF-AB and PDGF-BB is notably biphasic. In previous SELEX experiments, biphasic binding has been found to be a consequence of the existence of separable nucleic acid species that bind to their target protein with different affinities (Jellinek et al., (1995) *Biochemistry* 34:11363-11372, and unpublished results). The identity of the high and the low affinity fractions is at present not known. Since these DNA ligands described here were synthesized chemically, it is possible that the fraction that binds to PDGF-AB and PDGF-BB with lower affinity represents chemically imperfect DNA. Alternatively, the high and the low affinity species may represent stable conformational isomers that bind to the PDGF B-chain with different affinities. In any event, the higher affinity binding component is the most populated ligand species in all cases (FIG. 5). For comparison, a 39-mer DNA ligand that binds to human thrombin with a K_d of 0.5 nM (ligand T39 (SEQ ID NO.:177): 5'-CAGTC-CGTGGTAGGGCAGGTTGGGGTGACTTCGTGGAA [3'T], where [3'T] represents a 3'-3' linked thymidine nucleotide added to reduce 3'-exonuclease degradation) and has a predicted stem-loop structure, binds to PDGF-AB with a K_d of 0.23 μ M (data not shown).

Example 10

Kinetic Stability of PDGF-Nucleic Acid Ligand Complexes

[0182] In order to evaluate the kinetic stability of the PDGF-AB/DNA complexes, the dissociation rates were

determined at 37° C. for the complexes of minimal ligands 20 t, 36 t and 41 t (SEQ ID NOS:172-174) with PDGF-AB by measuring the amount of radiolabeled ligands (0.17 nM) bound to PDGF-AB (1 nM) as a function of time following the addition of a large excess of unlabeled ligands (FIG. 6). At these protein and DNA ligand concentrations, only the high affinity fraction of the DNA ligands binds to PDGF-AB. The following values for the dissociation rate constants were obtained by fitting the data points shown in FIG. 6 to the first-order rate equation: $4.5 \pm 0.2 \times 10^{-3} \text{ s}^{-1}$ ($t_{1/2} = 2.6 \text{ min}$) for ligand 20 t, $3.0 \pm 0.2 \times 10^{-3} \text{ s}^{-1}$ ($t_{1/2} = 3.8 \text{ min}$) for ligand 36 t, and $1.7 \pm 0.1 \times 10^{-3} \text{ s}^{-1}$ ($t_{1/2} = 6.7 \text{ min}$) for ligand 41 t. The association rates calculated for the dissociation constants and dissociation rate constants ($k_{on} = k_{off}/K_d$) are $3.1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ for 20 t, $3.1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ for 36 t and $1.2 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ for 41 t.

Example 11

Thermal Melting Properties

[0183] In order to examine the ability of minimal ligands 20 t, 36 t and 41 t to assume folded structures, their melting temperatures (T_m 's) were determined from the UV absorbance vs. temperature profiles in PBSM or PBSE buffers. At the oligonucleotide concentrations used in these experiments (320-440 nM), only the monomeric species were observed as single bands on non-denaturing polyacrylamide gels (data not shown). Ligands 20 t and 41 t underwent thermal melting that is well described by a two-state (folded and unfolded) model with linearly sloping baselines (Peterseim and Turner (1983) *Biochem.* 22:256-263) with T_m values in PBSM buffer of $43.8 \pm 0.4^\circ \text{ C.}$ and $49.2 \pm 0.5^\circ \text{ C.}$, respectively. In PBSE buffer, similar T_m values were obtained: $44.8 \pm 0.5^\circ \text{ C.}$ for ligand 20 t and $48.0 \pm 0.5^\circ \text{ C.}$ for ligand 41 t. Ligand 36 t exhibited a more complex thermal melting profile in which two distinct transitions were observed. In this case, the data were well described by a three-state model in which the fully folded and the unfolded states are connected through a partially unfolded intermediate results. Using this model, we obtained two T_m values for ligand 36 t: $47.0 \pm 0.9^\circ \text{ C.}$ and $67.1 \pm 3.8^\circ \text{ C.}$ in PBSM buffer and $44.2 \pm 1.7^\circ \text{ C.}$ and $64.3 \pm 4.1^\circ \text{ C.}$ in PBSE buffer.

Example 12

Photo-Crosslinking of Nucleic Acid Ligands and PDGF

[0184] In order to determine the sites on the DNA ligands and PDGF that are in close contact, a series of photo-crosslinking experiments were performed with 5'-iodo-2'-deoxyuridine (IdU)-substituted DNA ligands 20 t, 36 t and 41 t (SEQ ID NOS:172-174). Upon monochromatic excitation at 308 nm, 5-iodo- and 5-bromo-substituted pyrimidine nucleotides populate a reactive triplet state following intersystem crossing from the initial $n \rightarrow \pi^*$ transition. The excited triplet state species then reacts with electron rich amino acid residues (such as Trp, Tyr and His) that are in its close proximity to yield a covalent crosslink. This method has been used extensively in studies of nucleic acid-protein interactions since it allows irradiation with $>300 \text{ nm}$ light which minimizes photodamage (Willis et al., (1994) *Nucleic Acids Res.* 22:4947-4952; Stump, W. T., and Hall, K. B. (1995) *RNA* 1:55-63; Willis et al., (1993) *Science* 262:1255-1257; Jensen et al., (1995) *Proc. Natl. Acad. Sci., U. S. A.*

92:12220-12224). Analogs of ligands 20 t, 36 t and 41 t were synthesized in which all thymidine residues were replaced with IdU residues using the solid phase phosphoramidite method. The affinity of these IdU-substituted ligands for PDGF-AB was somewhat enhanced compared to the unsubstituted ligands and based on the appearance of bands with slower electrophoretic mobility on 8% polyacrylamide/7 M urea gels, all three 5' end-labeled IdU-substituted ligands crosslinked to PDGF-AB upon irradiation at 308 nm (data not shown). The highest crosslinking efficiency was observed with IdU-substituted ligand 20 t. To identify the specific IdU position(s) responsible for the observed crosslinking, seven singly or multiply IdU-substituted analogs of 20 t were tested for their ability to photo-crosslink to PDGF-AB: ligands 20 t-I1 through 20 t-17 (5'-TGGGAGGGCGCGT¹T¹CT¹T¹CGT²GGT³T⁴ACT⁵T⁶T⁶AGT⁷CCCG-3' (SEQ ID NOS:178-184) where the numbers indicate IdU substitutions at indicated thymidine nucleotides for the seven ligands). Of these seven ligands, efficient crosslinking to PDGF-AB was observed only with ligand 20 t-14. The photo-reactive IdU position corresponds to the 3' proximal thymidine in the loop at the helix junction (FIG. 4).

[0185] To identify the crosslinked amino acid residue(s) on PDGF-AB, a mixture of 5' end-labeled 20 t-I4 and PDGF-AB was incubated for 15 min at 37° C. followed by irradiation at 308 nm. The reaction mixture was then digested with modified trypsin and the crosslinked fragments resolved on an 8% polyacrylamide/7 M urea gel. Edman degradation of the peptide fragment recovered from the band that migrated closest to the free DNA band revealed the amino acid sequence KKPIXKK (SEQ ID NO:185), where X indicates a modified amino acid that could not be identified with the 20 derivatized amino acid standards. This peptide sequence, where X is phenylalanine, corresponds to amino acids 80-86 in the PDGF-B chain (Johnsson et al., (1984) *EMBO J.* 3:921-928) which in the crystal structure of PDGF-BB comprises a part of solvent-exposed loop III (Oefner et al., (1992) *EMBO J.* 11:3921-3926). In the PDGF A-chain, this peptide sequence does not occur (Betsholtz et al., (1986) *Nature* 320:695-699). Together, these data establish a point contact between a specific thymidine residue in ligand 20 t and phenylalanine 84 of the PDGF B-chain.

Example 13

Inhibition of PDGF by Nucleic Acid Ligands

[0186] In order to determine whether the DNA ligands to PDGF were able to inhibit the effects of PDGF isoforms on cultured cells, the effects on binding of ^{125}I -labeled PDGF isoforms to PDGF α - and β -receptors stably expressed in porcine aortic endothelial (PAE) cells by transfection was determined. Ligands 20 t, 36 t and 41 t (SEQ ID NOS:172-174) all efficiently inhibited the binding of ^{125}I -PDGF-BB to PDGF α -receptors (FIG. 7) or PDGF β -receptors (data not shown), with half maximal effects around 1 nM of DNA ligand. DNA ligand T39, directed against thrombin and included as a control, showed no effect. None of the ligands was able to inhibit the binding of ^{125}I -PDGF-AA to the PDGF α -receptor (FIG. 7), consistent with the observed specificity of ligands 20 t, 36 t and 41 t for PDGF-BB and PDGF-AB.

[0187] The ability of the DNA ligands to inhibit the mitogenic effects of PDGF-BB on PAE cells expressing

PDGF β -receptors was investigated. As shown in **FIG. 8**, the stimulatory effect of PDGF-BB on [^3H]thymidine incorporation was neutralized by ligands 20 t, 36 t and 41 t. Ligand 36 t exhibited half maximal inhibition at the concentration of 2.5 nM; ligands 41 t was slightly more efficient and 20 t slightly less efficient. The control ligand T39 had no effect. Moreover, none of the ligands inhibited the stimulatory effects of fetal calf serum on [^3H]thymidine incorporation in these cells, showing that the inhibitory effects are specific for PDGF.

Example 14

Post-SELEX Process Nucleotide Substitutions

[0188] The stability of nucleic acids to nucleases is an important consideration in efforts to develop nucleic acid-based therapeutics. Experiments have shown that many, and in some cases most of the nucleotides in SELEX-derived ligands can be substituted with modified nucleotides that resist nuclease digestion, without compromising high affinity binding (Green et al., (1995) *Chemistry and Biology* 2:683-695; Green et al., (1995) *J. Mol. Biol.* 247:60-68). Experiments of this type with the DNA ligands reported here suggest that substitutions with modified nucleotides are tolerated at many positions (**FIG. 9**; SEQ ID NOS:175-176). Specifically, we have examined the substitution of 2'-O-methyl-2'-deoxy- and 2'-fluoro-2'-deoxyribonucleotides for 2'-deoxyribonucleotides in ligand 36 t, by examining the PDGF-AB binding properties of singly or multiply substituted ligand 36 t. The substitution pattern indicated in **FIG. 9** is compatible with high affinity binding to PDGF-AB. Furthermore, this ligand tolerates the substitution of pentaethylene glycol spacers (Glen Research, Sterling, Va.) for the trinucleotide loops at the ends of helices II and III (**FIG. 9**). These DNA ligands therefore represent lead compounds for a novel class of high affinity, specific antagonists of PDGF-AB and PDGF-BB.

Example 15

Experimental Procedure for Evolving 2'-fluoro-2'-deoxyuridine RNA Ligands to PDGF and RNA Sequences Obtained

[0189] A. 2'-fluoro-2'-deoxyuridine RNA SELEX

[0190] SELEX with 2'-fluoro-2'-deoxyuridine RNA targeting PDGF AB was done essentially as described previously (vide supra, and Jellinek et al., (1993, 1994) supra) using the primer template set as shown in Table 12 (SEQ ID NOS:125-127). Briefly, the 2'-fluoro-2'-deoxyuridine RNA for affinity selections was prepared by in vitro transcription from synthetic DNA templates using T7 RNA polymerase (Milligan et al., *Nucl. Acids Res.* 15:8783 (1987)). The conditions for in vitro transcription described in detail previously (Jellinek et al., (1994) supra) were used, except that higher concentration (3 mM) of the 2'-fluoro-2'-deoxyuridine nucleoside triphosphates (2'-F-UTP and 2'-F-CTP) was used compared to ATP and GTP (1 mM). Affinity selections were done by incubating PDGF AB with 2'-fluoro-2'-deoxyuridine RNA for at least 15 min at 37°C. in PBS containing 0.01% human serum albumin. Partitioning of free RNA from protein-bound RNA was done by nitrocellulose filtration as described (Jellinek et al., (1993, 1994) supra). Reverse transcription of the affinity-selected

RNA and amplification by PCR were done as described previously (Jellinek et al., (1994) supra). Nineteen rounds of SELEX were performed, typically selecting between 1-12% of the input RNA. For the first eight rounds of selection, suramin (3-15 μM) was included in the selection buffer to increase the selection pressure. The affinity-enriched pool (round 19) was cloned and sequenced as described (Schneider et al., (1992) supra). Forty-six unique sequences have been identified, and the sequences are shown in Table 13 (SEQ ID NOS:128-170). The unique-sequence ligands were screened for their ability to bind PDGF AB with high affinity. While random 2'-fluoropyrimidine RNA (Table 12) bound to PDGF with a dissociation constant (K_d) of 35 ± 7 nM, many of the affinity-selected ligands bound to PDGF AB with ≈ 100 -fold higher affinities. Among the unique ligands, clones 9 ($K_d=91 \pm 16$ pM), 11 ($K_d=120 \pm 21$ pM), 16 ($K_d=116 \pm 34$ pM), 23 ($K_d=173 \pm 38$ pM), 25 ($K_d=80 \pm 22$ pM), 37 ($K_d=97 \pm 29$ pM), 38 ($K_d=74 \pm 39$ pM), and 40 ($K_d=91 \pm 32$ pM) exhibited the highest affinity for PDGF AB (binding of all of these ligands to PDGF AB is biphasic and the K_d for the higher affinity binding component is given).

Example 16

Experimental Procedures

[0191] This Example provides the general procedures followed and incorporated in Examples 17-19 for the evolution of nucleic acid ligands to hKGF.

[0192] A. Materials and Methods

[0193] Recombinant human Keratinocyte Growth Factor (hKGF) and human Epidermal Growth Factor (hEGF) were purchased from Upstate Biotechnology Inc. (Lake Placid, N.Y.). hAFGF, hbFGF, PDGF-AB, TGF β 1, and anti-KGF neutralizing monoclonal antibody were purchased from R&D Systems (Minneapolis, Minn.). Recombinant rat KGF was purchased from QED Advanced Research Technologies (San Diego, Calif.). Human thrombin was purchased from Enzyme Research Laboratories (South Bend, Ind.). T4 DNA ligase, HpaII methylase, and restriction enzymes were purchased from New England Biolabs (Beverly, Mass.). pCR-Script Amp SK(+) cloning kit was purchased from Stratagene (La Jolla, Calif.). AMV reverse transcriptase was purchased from Life Sciences (St. Petersburg, Fla.). Taq DNA polymerase was purchased from Perkin Elmer (Foster City, Calif.). Ultrapure nucleotide triphosphates were purchased from Pharmacia (Piscataway, N.J.). α - ^{32}P -ATP, γ - ^{32}P -ATP, and 5'- ^{32}P -cytidine 3', 5'-bis (phosphate) (5'- ^{32}P -pCp) were from DuPont NEN Research Products (Boston, Mass.). ^{125}I -labeled KGF was prepared as described before (Bottaro et al., (1990) *J. Biol. Chem.* 265:12767-12770). PC-3 prostatic carcinoma cells were obtained from ATCC (catalog number CRL1435). Balb/MK cells and NIH3T3 transfected cells with the human KGF receptor (NIH3T3/KGFR) were a generous gift from S. Aaronson, Mt. Sinai Medical Center, NY, and have been described elsewhere (Miki et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:246-250; Miki et al., (1991) *Science* 251:72-75; Weissman et al., (1983) *Cell* 32:599-606). T7 RNA polymerase, 2' NH_2 - and 2'-F-modified CTP and UTP were from NeXstar Pharmaceuticals, Inc. (Boulder, Colo.). DNA oligonucleotides were obtained from Operon Technologies, Inc. (Alameda, Calif.). Nitrocellulose/cellulose acetate mixed matrix, 0.45 μm , HA filters were from Millipore (Bedford, Mass.). Calcium and

magnesium containing Dulbecco's Phosphate Buffered Saline (DPBS) was purchased from Life Technologies (Gaithersburg, Md.). Chemicals were at least reagent grade and purchased from commercial sources.

[0194] B. SELEX

[0195] The SELEX procedure has been described in detail in U.S. Pat. No. 5,270,163 (see also Tuerk and Gold (1990) *Science* 249:505-510). A single-stranded DNA (ssDNA) pool was used to generate the double-stranded (dsDNA) template for generating the initial random sequence RNA pool by transcription. The DNA template contained 40 random nucleotides, flanked by 5' and 3' constant regions for primer annealing sites for PCR and cDNA synthesis (Table 14; SEQ ID NOS:186-188). The 5' primer contains the T7 promoter sequence for in vitro transcriptions. The template was PCR amplified following an initial denaturation at 93° C. for 3.5 minutes through 15 cycles of 30 second denaturation at 93° C., 1 minute annealing at 60° C., and 1 minute elongation at 72° C., in 50 mM KCl, 10 mM Tris-HCl, pH9, 0.1% Triton X-100, 3 mM MgCl₂, 0.5 mM of each dATP, dCTP, dGTP, and dTTP, 0.1 units/μl Taq DNA polymerase, and 2.5 nM each of 3G7 and 5G7 primers (Table 14; SEQ ID NOS:187-188). SELEX experiments for hKGF were initiated with a random sequence pool of RNA in which all pyrimidines were 2'-NH₂-modified or 2'-F-modified. Transcription reactions were done with about 5 μM DNA template, 5 units/μl T7 RNA polymerase, 40 mM Tris-HCl (pH8), 12 mM MgCl₂, 5 mM DTT, 1 mM spermidine, 0.002% Triton X-100, 4% PEG 8000, 2-4 mM each 2'OH ATP, 2'OH GTP, 2'NH₂ or 2'F CTP, 2'NH₂ or 2'F UTP, and 0.25 μM a ³²P 2'OH ATP (800 Ci/mmmole). The full length transcripts were gel-purified prior to use. To prepare binding reactions, the RNA molecules were incubated with recombinant hKGF in Dulbecco's Phosphate-Buffered Saline (DPBS) with calcium and magnesium (Life Technologies, Gaithersburg, Md., Cat. No 21300-025) containing 0.01% human serum albumin. Following incubation at room temperature (ranging from 10 minutes to 10 hours) the protein-RNA complexes were partitioned from unbound RNA by filtering through nitrocellulose. Nitrocellulose filter bound RNA was recovered by phenol/urea extraction. The partitioned RNA was reverse transcribed into cDNA by AMV reverse transcriptase at 48° C. for 60 min in 50 mM Tris-HCl pH8.3, 60 mM NaCl, 6 mM Mg(OAc)₂, 10 mM DTT, 50 pmol DNA 3' primer (Table 14), 0.4 mM each of dATP, dCTP, dGTP, and dTTP, and 1 unit/μl AMV RT. The cDNA was PCR amplified and used to initiate the next SELEX cycle.

[0196] C. Nitrocellulose Filter Partitioning

[0197] In order to partition the protein-RNA complexes, the binding reactions were filtered through nitrocellulose/cellulose acetated mixed matrix, 0.45 μm pore size (filter disks, Millipore, Co., Bedford, Mass.). For filtration, the filters were placed onto a vacuum manifold and wetted by aspirating 5 ml of DPBS. The binding reactions were aspirated through the filters, and following a 5 ml wash, the filters were counted in a scintillation counter (Beckmann). Higher wash volumes with DPBS or 0.5 M urea were used as a means to increase selection stringency as shown in Table 15. Gel purified, internally α-³²P-ATP labeled transcripts were incubated with various concentrations of hKGF in DPBS at 37° C. for 10 minutes. Oligonucleotide protein mixtures were filtered through prewetted 0.45 μm pore size HA filters, followed by a 5 ml wash with DPBS. The radioactivity retained on the filter was counted and corrected

for background binding in the absence of protein. Nonlinear least square method was used to fit the data into monophasic or biphasic binding curves and to obtain the equilibrium dissociation constant K_d (Jellinek et al., (1993) *Proc. Natl. Acad. Sci. USA* 90:11227-11231) using the software package Kaleidagraph (Synergy Software, Reading, Pa.). Biphasic binding can be described as the binding of two affinity species that are not in equilibrium.

[0198] D. Cloning and Sequencing

[0199] The RNA recovered from the round 8 filters was reverse transcribed and PCR amplified. Following column purification with QIA-quick spin columns (Qiagen, Inc., Chatsworth, Calif.) and ethanol precipitation, the amplified DNA was methylated with HpaII methylase (New England Biolabs, Beverly, Mass.). The methylated DNA was cloned into the SrfI restriction site of pCR-Script Direct SK(+) plasmid using the pCR-Script Amp SK(+) cloning kit (Stratagene Cloning Systems, La Jolla, Calif.). About 80 clones were sequenced with Sequenase sequencing kit (United States Biochemical Corporation, Cleveland, Ohio). Sequence analysis and secondary structure prediction was done by using previously described computer software (Feng and Doolittle (1987) *J. Mol. Evol.* 25:351-360; Jaeger et al., (1989) *Proc. Natl. Acad. Sci. USA* 86:7706-7710; Jaeger et al., (1990) *Methods Enzymol.* 183:281-306; Zucker (1989) *Science* 244:48-52).

[0200] E. Determination of Minimal Sequences Necessary for Binding

[0201] Oligonucleotide ligands end labeled at the 5' end with γ-³²P-ATP using T4 polynucleotide kinase, or at the 3' end with 5'-³²P-pCp and T4 RNA ligase, were used to establish 3' and 5' boundaries respectively (Fitzwater et al., (1996) *Methods Enzymol.* 267:275-301). After partial alkaline hydrolysis, the radiolabeled oligonucleotide was incubated with 0.1, 0.6, and 3.0 nM hKGF, and the protein bound oligonucleotide was isolated by nitrocellulose filtration. The nitrocellulose retained oligonucleotide truncates were analyzed on a high resolution denaturing polyacrylamide gel. An alkaline hydrolysis ladder and a ladder of radioactively labeled ligands terminated with G-residues, generated by partial RNase T1 digestion, were used as markers to map the 3' and 5' boundaries.

[0202] F. Thermal Denaturation Profiles

[0203] Oligonucleotide melting profiles were obtained with a Cary Model 1 E spectrophotometer. Oligonucleotides were heated to 95° C. in PBS (Sambrook et al., (1989) *Molecular Cloning: A Laboratory Manual, 2nd Ed.*, Cold Spring Harbor, N.Y.) or 10 mM phosphate buffer and cooled to room temperature before recording the melting profile. The melting profiles generated show the change in absorbance at 260 nm as a function of temperature. During recording, the samples were heated at a rate of 1° C. min⁻¹ from 20-95° C.

Example 17

RNA Ligands to hKGF

[0204] A. SELEX

[0205] To generate RNA ligands for hKGF, two parallel SELEX experiments were initiated, one with 2'-NH₂ and the other 2'-F pyrimidine modified RNA molecules randomized at 40 contiguous positions. The SELEX conditions and results for each round are summarized in Table 15. The

starting pool contained 5×10^{14} (500 pmoles) and 2.5×10^{14} (250 pmoles) 2'-NH₂ and 2'-F pyrimidine modified RNA molecules, respectively, and bound to hKGF with an approximate K_D of 30 nM. After 8 rounds of SELEX, the evolved pools bound with a K_D of 0.6 nM. No further improvement in the K_D was observed in the subsequent two rounds. The RNA pools from the 8th round were reverse transcribed, PCR amplified and cloned as described.

[0206] B. RNA Sequences

[0207] In the 2'-NH₂ SELEX, 29 out of 31 clones were unique. In the 2'-F SELEX all 43 clones sequenced were unique. A unique sequence is defined as one that differs from all others by three or more nucleotides. Table 16 lists the sequences (SEQ ID NOS:189-262) of all of the clones sequenced in standard single letter code (Cornish-Bowden, (1985) *Nucleic Acid Res* 13:3021-3030). Computer assisted global and local alignment did not reveal any extensive homologies among the clones, and no obvious families were apparent. The 2'-NH₂ clones are in general purine rich while the 2'-F clones are pyrimidine rich. When the alignment parameters were relaxed, the Feng/Doolittle algorithm grouped the 2'-NH₂ clones in one family and the 2'-F clones in another. Visual inspection of the sequences suggested two and three possible families for the 2'-NH₂ and the 2'-F ligands, respectively. Using conserved predicted secondary structure, 38 2'F ligands could be assigned into two classes (**FIGS. 12A and 12B**). Similarly, 15 2'NH₂ ligands could be assigned into two classes (**FIGS. 12C and 12D**). The two proposed classes for the 2'F ligands can be folded into pseudoknot structures (Wyatt et al., (1993) *The RNA World* 465-496; ten Dam, E. (1992) *Biochemistry* 31: 1665-1676). These structures are very related and in fact they could be circular permutations of a common structure. Loop 3 (L3) of class 1 pseudoknots presents the conserved sequence 5'RRYuy while loop 1 (L1) of class 2 ligands presents the sequence 5'AaYY. Both of these sequences contain the consensus 5'RRYY. Some of the 2'F ligands contain two to three copies of the RRY sequence (**FIGS. 12A and 12B**). Another feature of these structures is the unequal distribution of purines and pyrimidines in stem 1 (S1). One strand of that stem contains almost exclusively purines while the other strand contains pyrimidines.

[0208] Class 1 of the 2'NH₂ ligands includes 8 members that can be folded into stem-loop structures with internal symmetric or asymmetric loops. The stem contains three consecutive GC base pairs. The terminal loops are long and present the conserved sequence 5'GGAA(N)₁₋₁₄YAA(N)₁₋₇RCRR (SEQ ID NO:263). Both sides of the internal asymmetric loops of the class 1 ligands contain the sequence 5'AA. Class 2 includes 7 ligands that can be folded into dumbbells with variable sized loops. One loop contains the conserved sequence 5'YGAY while the other loop contains the conserved sequence 5'GGAA(N)₀₋₄YGA (SEQ ID NO:264). Clones 2N and 54N are circular permutations of the remaining 5 clones.

[0209] C. Affinities

[0210] The dissociation constants of the hKGF ligands were determined by nitrocellulose filter binding and are listed in Table 17. Eight out of 41 2'-F ligands bound biphasically. The remaining of the 2'-F and all the 2'-NH₂ ligands bound monophasically. Under protein excess, biphasic binding suggests that the ligand exists as two affinity

species (presumably isoconformers) that are not in equilibrium. The best 2'-F-modified ligand, K14F, binds biphasically with the high and low affinity dissociation constant at about 0.3-3 pM and 2-10 nM respectively. There is some observed variability in the K_D determinations for the various clones and the random RNA. Despite the experimental variability in the K_D determinations, the high affinity species of K14F have a 1,000-5,000 fold better affinity than the random RNA. Among the monophasic 2'-F-modified ligands, K38F had the best KD of about 0.3 nM. The best 2'-NH₂-modified ligands bound with a KD of 0.4 nM which represent about 75 fold improvement over the random RNA.

[0211] D. Determination of Minimal Sequences Necessary for Binding

[0212] Two 2'F ligands (6F and 14F) (SEQ ID NOS:223 and 231) were studied further to determine the minimal sequences necessary for binding. Sequence boundaries were determined by allowing an alkaline hydrolysis ladder, labeled at the 3' or 5' end, to bind to hKGF. The partial fragments were affinity purified by nitrocellulose filtration and analyzed on high resolution denaturing gels. Boundaries were clearly observed only at the 3' ends for both ligands (**FIG. 13**) and are in agreement with the class I proposed folding as shown in **FIGS. 12A and 12B**. Truncated templates were then used to confirm the boundaries (**FIG. 13**). Three truncates were tested for 6F because a run of 7 consecutive pyrimidines did not allow the precise mapping of the boundary. From these three truncates, one lost its KGF binding activity as shown in **FIG. 13**. A single 14F truncate, designated 14F3'T, was tested. This truncate was two bases longer than the observed boundary in order to extend stem 2 (S2) of the proposed pseudoknot structure. The 14F3'T truncated ligand retained binding activity with affinity similar to the full length ligand. Like the full length ligand, 14F3'T bound KGF biphasically where the high affinity species represented about 20% of the molecules and showed K_d values of about 0.3-3 pM. These high affinity species when partially separated from the low affinity species on the basis of differential affinity to KGF, exhibited binding curves with mid points at 0.3-3 pM and maximum plateaus of about 70% (data not shown). **FIG. 13** shows the predicted folding of the shortest active truncates for 6F and 14F which are 53 and 49 bases long respectively. Both proposed pseudoknot structures contain relatively long stems. The two proposed stems of 6F are separated by a single base forming a non-H-type pseudoknot. The proposed 6F structure resembles the solution structure of a similar pseudoknot motif from a frame-shifting element found in the MMTV RNA (Shen et al., (1995) *J. Mol. Biol.* 247:963-978). The two stems (S1 and S2) of 14F could be drawn as two coaxially stacked helices of 16 base pairs total length (H-type pseudoknot). A similar pseudoknot structure has been proposed before, based on NMR data (Du et al., (1996) *Biochemistry* 35:4187-4198). Given the short length of L1, it is possible that ligand 14F forms a non-H-type pseudoknot where the last GU base pair of S1 is not formed allowing a more flexible helical region and a longer L1. Temperature melting curves of 14F and 14F3'T suggest a remarkable thermostability for this ligand (data not shown). These melting curves appear to be concentration independent and biphasic in 150 mM salt. Biphasic melting curves have been observed before with tRNA (Hilbers et al., (1976) *Biochemistry* 15:1874-1882), and have been attributed to the tertiary folding of the RNA molecule. Multiphasic temperature

transitions have also been proposed for RNA pseudoknots (Du et al., (1996) *Biochemistry* 35:4187-4198). The biphasic curves observed include a low T_m at about 55° C. and a high T_m of greater than 85-90° C. In 10 mM salt the low T_m of 14F is not observed while the high T_m is shifted down to 75-78° C. The melting profile for 14F appears to be flatter than 14F3'T even though the T_m values are the same. The data suggest that the observed thermostability is attributable to just the minimal 49-mer.

[0213] In an effort to identify shorter KGF ligands that retained binding, the binding activity of various deletions of the shortest truncate of ligand 14F, namely 14F3'T were tested. Deletions were tested in all the structural elements of the proposed pseudoknot structure. The results are summarized in Table 23 (SEQ ID NOS:272-304). RNA transcripts containing 2'F pyrimidines and 2'OH purines were obtained by in vitro transcription using synthetic DNA templates. The activity of each ligand is shown by scoring for both the high (H) and low (L) affinity component of the 14F3'T binding curve with +(active) or -(not active). Truncates T35 and T36 represent two complementary halves of 14F3'T molecule and were additionally tested as an equimolar mixture. The structural elements of the proposed pseudoknot structure are separated by () and are indicated by symbols S1 (stem 1), S2 (stem 2), L1 (loop 1) and L3 (loop 3). The proposed pseudoknot structure for 14F3'T is a non-H-type pseudoknot and lacks L2 (loop 2). The complementary sequences forming S1 (S1 and S1') and S2 (S2 and S2') are marked by single and double underlines respectfully. In the tabulated sequences, deleted bases were replaced with periods (.). Any deletion attempt in the stems S1 and S2 of the proposed pseudoknot structure resulted in loss of both the high (H) and low (L) affinity component of the binding curve as observed with the 14F3'T ligand. Deletions in loop 3 (L3), however, were tolerated as long as one copy of the RYYY box remained intact. The shortest ligand that retained activity is T22 which is a 43-mer. In trying to obtain shorter ligands by truncating L3 further a mutant version of T22 (designated T22mu) was used where the last GC base pair of S1 was eliminated by a G to U mutation at position 6. The reasoning for this mutation was to enhance the flexibility of the double stranded region of this ligand by allowing an unpaired base between S1 and S2. Although this mutation did not affect the binding of T22 it did not allow further active truncations in L3.

[0214] E. Specificity of RNA Ligands to hKGF

[0215] The specificity of the K14F ligand was tested by determining its K_D against rat hKGF, and the heparin binding human growth factors, aFGF, bFGF, and PDGF (Table 18). The results suggest that the K14F binds all tested targets like random RNA, except hKGF, and it can discriminate between hKGF and other similar proteins by a factor of 400-40,000.

[0216] The specificity of ligand 14F3'T was tested by determining its K_d against a variety of heparin binding proteins. The results summarized in Table 22 show that ligand 14F3'T can discriminate KGF from all other heparin binding proteins tested by a factor of 1.2×10^4 to 3×10^{10} . Ligand 14F3'T binds only to KGF with high affinity while it binds all other heparin binding proteins tested like random RNA. Binding of 14F3'T to the rat KGF, which is 91% identical to human KGF, is with about a 5-10 fold reduced

affinity. Similar specificity was observed during the inhibition of the KGF induced DNA synthesis of Balb/MK cells. Ligand 14F3'T inhibits rat KGF induced DNA synthesis with a K_i of 1.8 nM which is 20-50 fold higher than the K_i observed with the human KGF. Ligand 14F3'T inhibits the DNA synthesis of Balb/MK cells only if it is the result of KGF but not EGF stimulation (data not shown).

Example 18

Inhibition of hKGF Binding to Cell Surface Receptors

[0217] A. Receptor Binding Assay

[0218] To test the ability of the hKGF ligands to competitively inhibit the binding of hKGF to its cell surface receptor, two cell lines were used. The first cell line, PC-3, is an isolate from a grade IV prostatic adenocarcinoma (ATCC CRL 1435). The second cell line is designated as NIH3T3/FGFR-2 and is a recombinant NIH/3T3 cell line carrying the human hKGF receptor at about $0.5-1 \times 10^6$ high affinity KGF binding sites per cell (Miki et al., (1992) *Proc. Natl. Acad. Sci. USA* 89:246-250).

[0219] PC-3 cells were plated in 24-well plates at about 10^5 cells per well. Following growth for 48-36 hours, the cells were serum starved for 24 hours, washed two times with 500 μ l of cold DPBS, and then incubated with 500 μ l binding buffer (BB1; DPBS, 0.5 mM $MgCl_2$, 0.2% BSA, 0.02% sodium azide) containing various concentrations of ^{125}I -labeled KGF ranging from 0 to 0.8 nM. Following 3-3.5 hour incubation at 4° C., the binding mixes were aspirated and the well-adhered cells were washed two times with 1 ml BB1 and once with 1 ml BB1 supplemented with 0.5M NaCl. The remaining bound labeled hKGF was solubilized in 600 μ l 0.5% SDS/0.1M NaOH and counted in a gamma counter (Beckmann). Nonspecific binding was determined in the presence of 100 fold molar excess of unlabeled hKGF. For competition assays, the labeled hKGF was kept constant at 0.3 nM, and varying concentrations of competitor molecules were included in the binding reactions ranging from 0-1,000 nM. Binding curves were fitted to the equation:

$$[\text{Bound Tracer}] = \frac{([\text{Total Tracer}] * [\text{Receptor}])}{(K_D + [\text{Total Tracer}])}$$

where [Total Tracer] and [Bound Tracer] were fixed and the K_D and [Receptor] were determined by regression analysis using the software Kaleidagraph (Synergy Software, Reading, Pa.).

[0220] NIH3T3/KGFR-2 cells were plated in 24-well plates at about 10^5 cells per well. Following growth overnight, the cells were serum starved for 1-5 hours, washed two times with 500 μ l binding buffer (BB2: serum-free MEM growth medium, 0.1% BSA, 25 mM HEPES, pH 7.4), and then incubated with 250 μ l BB2 containing 1 μ g/ml heparin (from bovine lung, SIGMA, St. Louis, Mo.), ^{125}I -labeled hKGF at 0.03 nM, and varying concentrations of competitor molecules (300 nM-0 nM). Following 1 hour incubation at room temperature, the binding mixes were aspirated, and the wells were washed two times with 250 μ l cold DPBS and once with 250 μ l cold DPBS supplemented with 0.5M NaCl. The bound labeled hKGF was solubilized in 500 μ l 0.5% SDS and counted in a scintillation counter (Beckmann).

[0221] The inhibition constants (K_i) of the RNA ligands were determined by a nonlinear regression analysis of the data.

[0222] In search of KGF receptors on the surface of PC-3 cells, different concentrations of ^{125}I -hKGF were used, ranging from 0.002 to 0.8 nM, in the presence and absence of 100 fold molar excess of unlabeled hKGF, and saturation binding of the tracer on the surface of PC-3 cells was observed. **FIG. 10** shows the plot of the concentration of bound tracer as a function of the total concentration of tracer as well as the Scatchard analysis of the same data. Analysis of the data suggested that there are about 5,000 specific hKGF binding sites per cell with a K_D of 100-200 pM. This K_D is in good agreement with the reported K_D for hKGF of 200 pM (Miki et al., (1992) *Proc Natl Acad Sci USA* 89:246-250).

[0223] PC-3 plasma membrane extracts were found to alter the electrophoretic mobility (gel shift) of radiolabeled hKGF upon native gel electrophoresis (**FIG. 11**). For electrophoretic mobility shift gels, about 3×10^7 PC-3 cells were gently spun and washed with PBS and then lysed by mixing with equal volume of lysis buffer containing 40 mM HEPES, pH 7.4, 150 mM NaCl, 20% glycerol, 2% triton X-100, 0.1% sodium azide, 3 mM MgCl_2 , 3 mM EGTA, 2 μM aprotinin, 2 μM leupeptin, 2 mM PMSF, and 400 μM sodium orthovanadate. Following 15 min incubation on ice the extract was spun at 11,000 g at 4° C. for 30 min to remove debris and nuclei and the supernatant was aliquoted and stored at -70° C. For gel analysis, 25 μl binding reactions were set in DPBS, 0.01% HSA, 2 mM MgCl_2 , containing 3 μl of a 10 fold diluted PC-3 membrane extract in 0.01% HSA, and various concentrations of ^{125}I -labeled hKGF. Following a 10 min incubation at room temperature, 6 \times loading dye was added to achieve 1 \times concentration, and the samples were loaded onto a 5% or 10% native TBE polyacrylamide gel. The gel was prerun at room temperature at 100 Volts. Following loading, the gel was run at 200 Volts for 5 min and then at 100 Volts for 30-60 min at room temperature. The radioactive bands were then visualized by autoradiography. The gel shift of radiolabeled hKGF is not observed in the presence of 100 fold molar excess of unlabeled hKGF (**FIG. 11**), demonstrating a specific interaction between a component found in the PC-3 membrane extracts and hKGF. The estimated KD from the gel shift experiment is about 8 nM.

[0224] In agreement with the competition experiments reported in the literature (Miki et al., *Proc Natl Acad Sci USA* 89:246-250), gel shift competition curves using unlabeled hKGF and bFGF as well as an unrelated small basic protein namely lysozyme were obtained. Table 21 lists the IC50 values obtained in this experiment. In agreement with previous reports, the data presented in Table 21 show that bFGF competes about 20 fold worse than hKGF for binding with the hKGF receptor present in the PC-3 plasma membrane extracts. The interaction observed by the gel shift appears to be a specific interaction for FGF and it is not due to a charge-charge interaction, as lysozyme, another small positively charged molecule, competes for the PC-3 membrane extract:hKGF complex with about 100 fold worse affinity than hKGF alone.

[0225] IC50 values for various RNA ligands obtained with the PC-3 assay are shown in Table 19. A subset of these ligands was tested on the NIH3T3/FGFR-2. Competitive

inhibition constants (K_i) were determined from full competition curves and are summarized in Table 20. In determining the K_i values, it was assumed that 3T3 cells have 500,000 binding sites per cell and PC-3 cells have 5,000 binding sites per cell.

[0226] The data show that several hKGF ligands can competitively inhibit binding of hKGF to its cell surface receptors. Some of these ligands, such as K14F, have potent competitive activities with K_i 's in the low nM range.

[0227] This work not only demonstrates that nucleic acid competitors for hKGF were obtained, but also identifies a new assay for screening hKGF competitors including small molecules, antibodies, and peptides. This new assay includes the use of the prostate carcinoma cell line, PC-3.

[0228] The two cell lines, PC3 and NIH3T3/FGFR-2, give slightly different results (see Table 20). KGF binding to PC-3 cells is more sensitive to inhibition by several ligands and by heparin. Random RNA, however, does not effectively compete for KGF binding on the PC-3 cells. KGF binding to NIH3T3/FGFR-2 is resistant to inhibition by some RNA ligands and heparin. This is because the NIH3T3/KGFR assay is more stringent since it is done in the presence of 1 $\mu\text{g}/\text{ml}$ heparin. The random oligonucleotide competition curve with the NIH3T3/FGFR-2 is completely flat with $K_i > 10^{-4}$ M. Ligands 6F and 14F show the best inhibitory activity with K_i values of 100-200 pM and 2-8 nM in the PC-3 and NIH3T3/FGFR-2 assay respectively. Only two $^{2'}\text{NH}_2$ ligands, 14N and 29N, show good activity with the PC-3 cells (K_i value of 1.4 nM). From these two ligands, only 14N retains its inhibitory activity in the NIH3T3/FGFR-2 assay showing a K_i value of 100 nM. The observed inhibition of the KGF mitogenic activity by these ligands is not due to a nonspecific affect in the proliferative ability of the cell lines because these ligands have no antiproliferative activity on cells induced by EGF instead of KGF (data not shown).

[0229] This work not only demonstrates that nucleic acid competitors for hKGF were obtained, but also identifies a new assay for screening hKGF competitors including small molecules, antibodies, and peptides. This new assay includes the use of the prostate carcinoma cell line, PC-3.

Example 19

Inhibition of the Mitogenic Activity of KGF

[0230] One of the biological effects of KGF is the stimulation of proliferation of epithelial cells (Rubin et al., (1989) *Proc Natl Acad Sci USA* 86:802-806). This proliferative effect of KGF can be measured by the stimulation of ^3H -thymidine incorporation in responding cells after exposure to KGF. Three such cell lines have been described before (Rubin et al., (1989) *Proc Natl Acad Sci USA* 86:802-806). Two cell lines were used to test the anti-mitogenic activity of various ligands. One is 4MBr-5 (ATCC #CCL208), a monkey epithelial, low passage, cell line (Caputo et al., (1979) *In Vitro* 15:222-223) while the second is Balb/MK, a transformed rat keratinocyte cell line (Weissman and Aaronson (1983) *Cell* 32:599-606). 4-MBr5 cells grown in F12K containing 30 ng/ml, hEGF, and 10% FCS,

were trypsinized and resuspended in M199 containing 10 mM HEPES, pH 7.4, and 10% FCS at 1.4×10^5 cells/ml. A 96-well microtiter plate was seeded with 100 μ l of cell suspension and KGF was added at 10 ng/ml (0.5 nM), as well as K14F ligand at various concentrations ranging from 0-1000 nM. Each incubation reaction was set in at least triplicates. Following 24 h incubation at 37° C., 3 H-thymidine was added at 1 μ Ci/well along with unlabeled thymidine at 10 nM. The cells were incubated for additional 24 h, the supernatant was aspirated, and the remaining cells were harvested by lysis in 20 μ l of 0.2 N NaOH. The extent of 3 H-thymidine incorporation was determined by TCA precipitation and filtration through GFC filter disks (Whatman, Hillsboro, Oreg.).

[0231] Balb/MK cells grown in Low Ca^{++} EMEM with 10% FCS (dialyzed and heat inactivated) and 5 ng/ml rhEGF were trypsinized and resuspended in Low Ca^{++} EMEM with 1% FCS (dialyzed and heat inactivated) and 0.5 ng/ml rhEGF and plated on 96 well fibronectin coated culture plates at $4-6 \times 10^4$ cells per well in 100 μ l total volume. Following overnight growth, the medium was replaced with Low Ca^{++} EMEM without FCS or rhEGF and serum starved for about 30 hrs. Human recombinant KGF or EGF was then added at 16 and 49 pM respectively, along with various concentrations of competitors ranging from 0-1000 nM. Following over-night incubation, 3 H-thymidine was added at 0.2 μ Ci/well and incubation continued for an additional 7-8 hrs. The extent of 3 H-thymidine incorporation was determined by TCA precipitation and filtration through GFC filter disks.

[0232] The inhibition constants (K_i) of the oligonucleotide ligands were determined by a nonlinear regression analysis of the data as described before (Gill et al., (1991) *J. Mol. Biol.* 220:307-324).

[0233] The two assays give slightly different results. The 4MBr-5 assay was performed in the presence of fetal calf serum, while the Balb/MK was done following serum starvation. The Balb/MK assay is more sensitive and a prototypic assay for the KGF induced mitogenic activity. Similar to the results obtained with the PC-3 cells, the 4MBr-5 assay showed a good activity for ligand 14F (K_i value of 9.8 nM but incomplete inhibition). In the same assay, the random oligonucleotides showed K_i values of $>1 \mu$ M while a monoclonal neutralizing antibody showed a K_i value of 2.9 nM. It appears that ligand 14F is as good or even better than the monoclonal neutralizing antibody. The competition curves for the neutralizing monoclonal antibody and ligand 14N

plateau at about 20-40%, suggesting that these antagonists do not completely abolish the KGF mitogenic activity. In contrast to the monoclonal antibody, ligand 14F completely blocks the KGF mitogenic activity on the 4MBr-5 cells. In the Balb/MK assay, 14N showed K_i values of about 10 nM (incomplete inhibition) while the random oligonucleotide showed K_i values of about 300 nM. The K_i values for 6F and 14F are 830 and 92 pM, respectively. Similar to the 4MBr-5 assay, ligand 14F appears to be as good if not better than the monoclonal neutralizing antibody which shows a K_i value of 980 pM. The best inhibitory activity was observed with 14F3'T with a K_i value of 34 pM.

Example 20

[0234] Nucleic acid ligands that bind to basic fibroblast growth factor (bFGF) have been derived by the SELEX method as described in U.S. Pat. No. 5,459,015 (see also U.S. Pat. No. 5,270,163 and Tuerk and Gold (1990) *Science* 249:505-510). A 2' NH_2 -modified nucleic acid ligand designated 21A having the sequence 5'-GGGAGACAA-GAAU AACGCUCAAGUAGACUAAUGUGUG-GAAGACAGCGG-GUGGUUCGACAGGAGGCUCACAACAGGC (SEQ ID NO:265) was examined by deletion analysis for the minimal sequence information required for high affinity binding to bFGF. This analysis led to truncated ligand 21A-t (GGU-GUGUGGAAGACAGCGGGUGGuuc (SEQ ID NO:266) where the underlined G's are guanines added to improve efficiency of transcription and lowercase letters are from the constant region.

[0235] In order to increase the stability of ligand 21A-t against degradation by nucleases, short phosphorothioate caps were added to the 5' and the 3' ends. In addition, nine ribopurine positions were identified that can be substituted with 2'-deoxy-2'-O-methylpurines without a loss in binding affinity for bFGF, using the method described in Green et al., *Chem. Biol.* 2:683-695, resulting in the ligand designated as NX-286 (5'-TsTsTsTs mGmGaU rGaUrG aUrGrG mAaArG mAaCrA rGaCmG mGmGaU mGmGaU aUaC TsTsTsTsT-3' (SEQ ID NO:267), where s represents phosphorothioate internucleoside linkage, aU and aC are 2'-deoxy-2'-aminouridine and 2'-deoxy-2'-aminocytidine residues, respectively, mA and mG are 2'-deoxy-2'-O-methyladenosine and guanosine residues, respectively, rA and rG are adenosine and guanosine residues and T is 2'-deoxythymidine). The modified nucleic acid ligand had a K_d of 0.4 nM as measured by electrophoretic mobility shift assay.

TABLE 1

Nucleic Acid Sequences Used in SELEX Experiments described in Examples 1-4		SEQ ID NO.
Starting RNAs:		
64N6 transcript: 5' GGGGGAGAACGCGGAUCC [-64N-] AAGCUUCGCUCUAGAUCUCCUUUAGU GAGGGUUA 3'		1
40N6 transcript: 5' GGGGGAGAACGCGGAUCC [-40N-] AAGCUUCGCUCUAGAUCUCCUUUAGU GAGGGUUA 3'		2

TABLE 1-continued

Nucleic Acid Sequences Used in SELEX Experiments described in Examples 1-4		SEQ ID NO.
randomized lib2-6-1 transcript*:		
5'GGGGGAGAACGCGGAUCC[ugucuccaccgcccgaucuggguuccuggggccccuccauggag gaggggggugguucggaga] AAGCUUCGCUCUAGAUCUCCUUUAGUGAGGGUUA 3'		3
Starting DNA templates:		
Z-54 (64N6):		
5'GGGAGAACGCGGATCC [-64N-] AAGCTTCGCTCTAGA3'		4
Z-55 (40N6):		
5'GGGAGAACGCGGATCC [-40N-] AAGCTTCGCTCTAGA3'		5
D-123(randomized lib2-6-1)*:		
5'GGGGGAGAACGCGGATCC[tgtctccaccgcccgatactggggttcctggggcccctccatggaggagg gggtggttcggaga] AAGCTTCGCTCTAG 3'		6
PCR and cloning primers:		
T7SacBam:		
5'TAATACGACTCACTATAGGGGAGTCTGCGGATCC3'		7
	SacI BamH1	
T7SB2N:		
5'TAATACGACTCACTATAGGGGAGAACGCGGATCC3'		8
	BamH1	
3XH:		
5'TAACCCCTACTAAAGGGAGATCTAGAGCGAAGCTT3'		9
	XbaI HindHIII	
BamH1 cloning site engineered into pGem9zf to clone SELEX experiments 3-9.		
GATTTAGGTGACACTATAGAATATGCATCACTAGTAAGCTTTGCTCTAGA		10
	SP6 promoter XbaI	
GGATCCCGAGCTCCCTATAGTGAGTCGTATTA		11
	BamH1 T7 promoter	
*GAUC or GATC, these bases only gauc or gact 62.5% specified base, 12.5% the other three bases		

[0236]

TABLE 2

RNA SELEX Experiments described in Examples 1-4: template, pyrimidine nucleotides, and round cloned.				
SELEX exp	template*	2'substituted UTP	2'substituted CTP	Round cloned
lib1	64N6	OH	OH	20
lib2	64N6	OH	OH	6
lib3	40N6 + 64N6	F	F	4
lib4	40N6 + 64N6	NH ₂	NH ₂	5
lib5	64N6	NH ₂	NH ₂	13
lib6	64N6	F	F	13

TABLE 2-continued

RNA SELEX Experiments described in Examples 1-4: template, pyrimidine nucleotides, and round cloned.				
SELEX exp	template*	2'substituted UTP	2'substituted CTP	Round cloned
lib7	64N6	F	NH ₂	14
lib8	D-123	OH	OH	6
lib9	64N6	NH ₂	F	5

*Sequences of templates are described in Table 1.

[0237]

TABLE 3

TGFb Binding ligands				
Clone	5' CONSTANT	VARIABLE	3' CONSTANT	SEQ ID NO.
Group A lib3	gggggagaacgcggaucc	[40 or 64N]	aagcuucgcucuagaucuccuuuagugaggguaa	2 or 1
13	GAGCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUACUCGGUGUCGUG			12
3	GAGCAACCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUACUCGGUGUCGUG			13
4	GAGCAACCCCAGGCGCAUAGCUUCCGAGUAGACAGGCGGGAGGGGUGGAUGUGGCUCUACUCGGAGUCGUG			14
lib4 32	G GCAACCCCAGGCGCAUAGCUUCCGAGUAGACAGGCGGGAGGGGUGGAUGUGGCUCUACUCGGAGG			15
lib8 9	GCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUACUCGGCUCGUG			16
lib5 5	GAGCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUACUCGGAGG			17
7	GAGCAAGCCCUGGC AUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUACUCGGUGUCGUG			18
48	G GCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGUGU ACGAGG			19
lib2 6-4	GAGCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGUGUCU CGAG			20
lib6 23	A AGCUUC GAGUAGACAGGAGGGAGGGGUGGAUGUGGAGUCU CGAG			21
4	GAGCAAUCCUAA GCAUAGCUUC GAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCU CGAG			22
lib7 21	GAGCAAUCCCAGGCGCAUAGCUUCCGAGGAGACAGGCGGGAGGGGUGGAUGUGGCUCU CGAG			23
43	GAGCAAUCCCAGGCGCAUAGCUUCCGAGUAGACAGGCGGGAGGGGUGGAUGUGGCUCU CGAG			24
Clone	5' CONSTANT	VARIABLE	3' CONSTANT	SEQ ID NO.
Group B.	gggggagaacgcggaucc	[40 or 64N]	aagcuucgcucuagaucuccuuuagugaggguaa	2 or 1
lib4-12	UGAGAAGGACGUCGGGGUCAACGGGGUGAGGUGCAGCAGAAAGGGCCGGCACCACAUGACGUAA			28
lib3-44	UGAGAAGGACGUCGGGGU	GAGGUGCAGCAGAAAGGGCCGGCACCACAUGACGUAA		29
lib3-42	GGUGGGAAA GUCGGAUU	AUGUGU GUAGAUUU GU GUGCGA		30
Group C.				
lib1-20-3**	UGC UAGACCCAGGAUGCAAAGGGACAUGCAUUAGGGAAACCUAUGUAUAAGAACCGGUCGCAG			32
lib1-20-3H**	UGC UAGACCCAGGAUGCAAAGGGACAUGCAUUAGGGAAACCUAUGUAUAAGAACCGGUCGCAG			33
lib6-30**	UGC UAGACCCAGGAUGCAAAGGGACAUGCAUUAGGGAAACCUAU UAUAGAACCGGUCGCAG			34
Group D.				
lib2-6-1*	UGUCUCCACC GCCGAUACUGGGGUUCCUGGGGCCCUCCAUGCAGGAGGGGGUGGUUCGGAGA			35
lib2-6-1-81*	UGUCUCCACC GCCGAUACUGGGGUUCCUGGGGCCCUCCAUGCAGGAGGGGGUGGUUCGGAG			36
lib8-23*	UGUCUCCACC GCCGAUACUGGGGUUCCUGGGGCCCUCCAUGCAGGAGGGGGUGGUUCGGAGA			37
lib9-10*	UGUCUCCACC GCCGAUACUGGGGUUCCUGGGGCCCUCCAUGCAGGAGGGGGUGGUUCGGAGA			38
ORPHANS.				
clone#				
lib3-45	GGAAAGUCUGGUCUUUGGGGAGUCCGCAUGGCCUUGGCGA			39
lib1-20-5**	AAGAAUGUUCGGCCGCACGAGGUGACAGUGGUGCGGAUACGGACCGAUUGGUUUUGCC			40
lib1-20-12***	GGUCACCCGGCAUUAACA AUGCCGACACUGGGGUUCCUGGGACGGGUGGACUGGACGGAAAG			41
lib2-6-8***	AUAACCGGUCGCAUGGGAGGACAUCUGGGAAAGGACGGGUCGAGAUACUGAGCAGUUCGGC			42
Clone				SEQ ID NO.
Group A Boundary Experiments				
lib3-13 boundaries	5' GCUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUAC	3'		25
lib8-9 boundaries	5' CUUCCGAGUAGACAGGAGGGAGGGGUGGAUGUGGCUCUAC	3'		26

TABLE 3-continued

TGFb Binding ligands		
lib4-32 boundary	GGCAACCCAGGCGCAUAGCUUCCGAGUAGACAGGCGGGAGGGUGGAUGUGGCGUCACG 3'	27
Group B Boundary Experiments		
lib4-12 boundaries	5' UGAGAAGGACGUCGGGGUCAACGGGGUGAGGUGCAGCAGAAAGGCCGGCACCA 3'	31

Legend: The constant region of the ligand is shown in lower case and variable in upper. Sequences have been aligned.

Deletions with respect to the first sequence in each group are shown by gaps, substitutions are in bold type.

*2'NH₂-UTP, 2'F-CTP:

**2'F-UTP, 2'F-CTP.

***2'OH-UTP, 2'OH-CTP

Group A and B bind with either 2'NH₂- or 2'F- pyrimidines.

Ligands bind with either 2'NH₂- or 2'F- pyrimidines unless otherwise indicated

[0238]

TABLE 4

Dissociation and Inhibition Constants				
Group	Ligand	B _{max}	K _d	IC ₅₀
A	lib3-13	0.60	0.9 nM	9.7 nM
		0.38	0.7 nM	42 nM
		0.55	0.9 nM	18 nM
				32 nM
	lib3-3	0.44	1.7 nM	NT
	lib4-32	0.50	0.8 nM	20 nM
				157 nM
	lib5-5	0.37	2.4 nM	49 nM
	lib5-7	0.33	3.4 nM	17 nM
	lib8-9	0.4	1.7 nM	210 nM
	lib8-9*	0.35	2.8 nM	124 nM
	lib5-48	0.32	3.8 nM	not inhibitory
	lib2-6-4	0.20	3.1 nM	not inhibitory
	lib6-23	0.35	3.4 nM	not inhibitory
lib7-21****	0.18	2.4 nM	not inhibitory	
lib7-43****	0.33	3.3 nM	not inhibitory	
B	lib4-12	0.15	0.4 nM	109 nM
		0.08	0.2 nM	108 nM
	lib3-44	0.18	1.3 nM	119 nM
	lib3-42	0.16	0.6 nM	22 nM

TABLE 4-continued

Dissociation and Inhibition Constants				
Group	Ligand	B _{max}	K _d	IC ₅₀
C	lib1-20-3**	0.67	30 nM	not inhibitory
	lib1-20-3-82**	0.46	6.1 nM	not inhibitory
	lib6-30**	0.35	8.8 nM	not inhibitory
D	lib2-6-1*	0.40	14.3 nM	112 nM
				103 nM
	lib2-6-1-81*	0.39	10.7 nM	201 nM
				298 nM
Orphans	lib8-23*	0.48	6.6 nM	not inhibitory
	lib9-10*	0.24	1.1 nM	not inhibitory
	lib3-45	0.08	1.9 nM	not inhibitory
	lib1-20-5**	0.42	46 nM	not inhibitory
Controls	lib1-20-12****	0.34	3.1 nM	NT
	lib1-6-8****	0.12	4.7 nM	NT
	lib5-9		nonbinder	not inhibitory
	random 64N6		nonbinder	not inhibitory

ligands are 2'-NH₂ pyrimidines unless otherwise noted

*2'-NH₂-UTP, 2'-F-CTP,

**2'-F pyrimidines,

***2'-OH pyrimidines,

****2'-F-UTP, 2'-NH₂-CTP

[0239]

TABLE 5

DNA oligonucleotides used in Examples 5 and 6 ^a		
Description	Sequence	SEQ ID NO.
40N7 Template for RNA SELEX	TCGGGCGAGTCGTCTG[40N]CCGCATCGTCCTCCC	43
5N7 5'-primer for PCR	TAATACGACTCACTATAGGGAGGACGATGCGG	44
3N7 3'-primer for PCR	TCGGGCGAGTCGTCTG	45
40D7 Starting material for DNA SELEX	GGGAGGACGATGCGG[40N]CAGACGACTCGCCCGA	46
5D7 5'-primer for PCR	GGGAGGACGATGCGG	47
3D7 3'-primer for PCR	(biotin) ₃ TCGGGCGAGTCGTCTG	48
40N8 Template for RNA SELEX	GCCTGTTGTGAGCCCTCTGTCGAA[40N]TTGAGCGTTTATTCTGTCTCCC	49
5N8 5'-primer for PCR	TAATACGACTCACTATAGGGAGACAAGAATAAACGCTCAA	50
3N8 3'-primer for PCR	GCCTGTTGTGAGCCCTCTGTCGAA	51

TABLE 5-continued

<u>DNA oligonucleotides used in Examples 5 and 6^a</u>		
Description	Sequence	SEQ ID NO.
40D8 Starting material for DNA SELEX	GGGAGACAAGAATAAACGCTCAA[40N]TTCGACAGGAGGCTCACAAACAGGC	52
5D8 5'-primer for PCR	GGGAGACAAGAATAAACGCTCAA	53
3D8 3'-primer for PCR	(biotin) ₃ GCCTGTTGTGAGCCTCCTGTCGAA	54

^aDNA oligonucleotides 40N7 and 40N8 were used to generate the double-stranded DNA template for in vitro transcription. The 3'-primers 3N7 and 3N8 were also used to generate cDNA from the RNA repertoire. Synthetically synthesized DNA oligonucleotides 40D7 and 40D8 were used directly as the starting repertoire for the two single-stranded DNA SELEX experiments. PCR amplification of the selected repertoires used the appropriate 5'- or 3'-primer. The symbol 40N indicated a 40-nucleotide randomized region within the oligonucleotide.

[0240]

TABLE 6

<u>TGFβ1 40N7 DNA Selex Sequence of fifty randomly chosen clones.</u>		
		SEQ ID NO.
	5' GGGAGGACGATGCCG . . . 40N . . . CAGACGACTCGCCGA 3'	46
<u>Group A</u>		
20 (11 clones)	CCAGGGGGGTATGGGGTGGTCTACTTACTTGCCTTT	55
4	CCAGGGGGGTATGGGGTAGTGCTACTTACTTGCCTTT	56
5	CCAGGGGGGTATGGGGTAGTACTACTTACTTACGCTTT	57
8	CCAGGGGGGTATGGGGTATACTACTTACTTACGCTTT	58
13	CCAGGGGGGTATGGGGTAATACTACTTACTTACATCTT	59
16	CCAGGGGGGTATGGGGTAATACTACTTACTTACGCTTT	60
40	CCAGGGGGGTATGGGGTGGTGTACTTACTTGCCTTT	61
48	CCAGGGGGGTATGGGGTGGTCTTCTTACTTGCCTTT	62
<u>Group B</u>		
18	CCAGGGGGGTATGGGGTGGTGTACTTTTCTCCTGCTCTTC	63
19	CCAGGGGGGTATGGGGTGGTTCGTTTTCTTTGCGGCTT	64
32	CCAGGGGGGTGTGGGGTGGTGTACTTTTCTTGTCTTC	65
46	CCAGGGGGGTATGGGGTGGTTTGGTATGTTGCGTCCGT	66
<u>Group C</u>		
12 (3 clones)	CCGGGTGGGTATGGGGTAATACTACTTACTTACGCTTT	67
1	CCGGGGTGGGTAGGGGGTAGTCTACTTACTTACGCTTT	68
3	CCAGGGTGGGTGTGGGGTAGTACTACTTACTTGCCTTT	69
10	CCAGGGTGGGTATGGGGTAGTCTACTTACTTGCCTTT	70
23	CCGGGTGGGTATGGGGTGGTGTACTTACTTGCCTTT	71
34	CCTGGGTGGGTATGGGGTGGTCTACTTACTTGCCTTT	72
<u>Group D</u>		
2	CCACGGGTGGGTGTGGGGTAGTGTCTCACTTTACATCAC	73
6	CCCGGGTGGGTGTGGGGTAGTGTATTATTTACAGCCT	74
25 & 38	CCAGGGTGGGTGTGGGGTGGTGTACTTTTCTGCTCCTT	75
7	CCAGGGTGGGTATGGGGTAGTGTACTTTTAAATGATCTT	76
9	CCCGGGGAGAGCGGTGGGTAGTGTCTATAGTATTTCGTGT	77
11	CCAGGGGGGTATGTTTTAATACTACTTACTTACGCTTT	78
17	CCAGGGAGGGTATGGGGTGGTGTCTTACTTGTTCGCGCT	79
21	CCAGGGTGGCATGGGGTGGTGTGGATTAATCTTCGTCC	80
24	CCAGGGTGGGTGTGGGGTGGTGTCTTATTTACTCGTCGC	81
28 & 30	GGGGCGCTTGAAGAGGTTGCCGTTGGAGTATTCGAGC	82
29	CCAGGTGTGGGGTGGTTGGGTTTCTTTCGTCGCC	83
31	CCAGGGTGGGTATGGGGTTAATTAATCTTCGTCCCA	84
35	GGGGCGCTTGAAGAGGTTGCCGTTGGAGTATTCGAGC	85
36	CCCGGGTGGGTGTGGGGTGGTGTGAATTAATCTTCGTCC	86

TABLE 6-continued

<u>TGFβ1 40N7 DNA Selex Sequence of fifty randomly chosen clones.</u>		
		SEQ ID NO.
41	CCCGGGTGGGTGTGGGGTGGTGTATTATATTTGCGGCCT	87
44 & 45	CCAGGGTCGGTGTGGGTGGTGTACTTTTCCCTGTCCTTC	88
50	GGGGCGCTTGAAGAGGTTGCCGGTTGGAGTATTCGAGC	89

Bold typeface indicates a discrepancy with the most common sequence of that group.

[0241]

TABLE 7

<u>Starting DNA and PCR primers for the ssDNA SELEX experiment</u>	
	SEQ ID NO.
Starting ssDNA: 5'-ATCCGCCTGATTAGCGATACT[-40N-]ACTTGAGCAAATCACCTGCAGGGG-3'	90
PCR Primer 3N2*: 5'-BBBCCCTGCAGGTGATTTTGCTCAAGT-3'	91
PCR Primer 5N2**: 5'-CCGAAGCTTAATACGACTCACTATAGGGATCCGCCTGATTAGCGATACT-3'	92

*Biotin phosphoramidite (e. g., Glen Research, Sterling, VA)
**For rounds 10, 11, and 12, the truncated PCR primer 5N2 (underlined) was used to amplify the template.

[0242]

TABLE 8

<u>Unique Sequences of the ssDNA high affinity ligands to PDGF</u>	
	SEQ ID NO
5'-ATCCGCCTGATTAGCGATACT [40N] ACTTGAGCAAATCACCTGCAGGGG-3'	90
*14 AGGCTTGACAAAGGCACCATGGCTTAGTGGTCCCTAGT	93
*41 CAGGGCACTGCAAGCAATTGTGGTCCCAATGGGCTGAGT	94
6 CCAGGCAGTCATGGTCATTGTTTACAGTCGTGGAGTAGGT	95
23 AGGTGATCCCTGCAAAGGCAGGATAACGTCCTGAGCATC	96
2 ATGTGATCCCTGAGAGGGAGGANACGCTGAGCATC	97
34 CACGTGATCCCATAAAGGGCTGCGCAAATAGCAGAGCATC	98
8 GGTGGACTAGAGGGCAGCAAACGATCCTTGGTTAGCGTCC	99
1 GGTGCGACGAGGCTTACACAAACGTACACGTTTCCCCGC	100
5 TGTCCGAGCAGGGGCGTACGAAAACTTTACAGTTCCCCCG	101
*40 AGTGGAACAGGGCACGGAGAGTCAAACCTTGGTTTCCCCC	102
47 GTGGTAGGGATCGGTGGATGCCTCGTCACTTCTAGTCCC	103
18 GGGCGCCCTAAACAAAGGGTGGTCACTTCTAGTCCCAGGA	104
30 TCCGGGCTCGGGATTCGTGGTCACTTTCAGTCCCGGATATA	105

TABLE 8-continued

<u>Unique Sequences of the ssDNA high affinity ligands to PDGF</u>		SEQ ID NO
*20	ATGGGAGGGCGCGTTCTTCGTGGTTACTTTTAGTCCC	106
35	ACGGGAGGGCACGTTCTTCGTGGTTACTTTTAGTCCC	107
13	GCTCGTAGGGGGGATTCTTCGCCGTTACTTCCAGTCCT	108
16	GAGGCATGTTAACATGAGCATCGTCTCAGATCCTCAGCC	109
*36	CCACAGGCTACGGCAGTAGAGCATCACCATGATCCTGTG	110
50	GCGGGCATGGCACATGAGCATCTCTGATCCCGAATCCTC	111
4	ACCGGGCTACTTCGTAGAGCATCTCTGATCCCGGTGCTCG	112
44	AAAGGGCGAACGTAGGTCGAGGCATCCATTGGATCCCTTC	113
24	ACGGGCTCTGTCACTGTGGCACTAGCAATAGTCCCGTCGC	114
7	GGGCAGACCTTCTGGACGAGCATCACCTATGTGATCCC	115
*26	AGAGGGGAAGTAGGCTGCCTGACTCGAGAGAGTCCCTCCG	116
19	AGGGGTGCGAAACACATAATCCTCGCGGATCCCATCGCT	117
48	GGGGGGCAATGGCGGTACCTCTGGTCCCCTAAATAC	118
46	GCGGCTCAAAGTCTGTACCCGACGACATCTGTGGTC	119
25	TTGGGGTGAATGTCCACGGGTACCTCCGGTCCCAAAGAG	120
31	TCCGCGCAAGTCCCTGGTAAAGGGCAGCCCTAATGGTC	121
12	CAAGTTCCCACAAGACTGGGGCTGTTCAAACCGCTAGTA	122
15	CAAGTAGGGCGGACACACGTCGGGCACCTAAGGTCCA	123
*38	AAAGTCGTGCAGGGTCCCTGGAAGCATCTCCGATCCCAG	124

* Indicates a boundary experiment was performed.
 Italics indicate the clones that were found to retain high affinity binding as minimal ligands.

[0243]

TABLE 9

SEQ ID NO:	Group A	Sequence
97	2	=AGGG---AGGA---TACG-----TCTG-AGC-ATCac3' 5'ATGTGAT-CCCTGCAG=
112	4	ACCGGG---CTAC--TTC-----GTAG-AGC-ATC---TCT----GAT-CCCGGTGCTCG
115	7	tGGG---CGACC-TTCT----GGACG-AGC-ATCAC--CTAT--GTGAT-CCCG
109	16	ctGAGG---CATG--TAA-----CATG-AGC-ATCGT--CTC---ACGAT-CCTCAGCC
110	36	CCACAGG---CTACG-GCA-----CGTAG-AGC-ATCA---CCA----TGAT-CCTGTG
124	38	AAAGTCGTGCAGGG---TCC---CCT-----GGA-AGC-ATC---TCC----GAT-CCCAGact
113	44	AAAGGG---CGAAC-GTA-----GGTCG-AGGCATCC---ATT---GGAT-CCCTTC
111	50	GCGGG---CATG--GCA-----CATG-AGC-ATC---TCT----GAT-CCCGAATCCTC
96	23	=AGC---CAGGATAAC-----GTCTG-AGC-ATCac3' 5'AGGTGATCCCTGCAA=
98	34	GGG---CTGC--GCAAAAATA--GCAG-AGC-ATCac3' 5'CACGTGAT-CCCATAA=

TABLE 9-continued

SEQ ID NO: Group A		
108	13	GCTCGTAGG--GGGCGA-TTCTT----TCGCC-GTT- <u>ACT</u> ---TCC-----AGT-CCTac
93	14	tactAGG-- <u>CTT</u> --GACA-----AAG-GGC- <u>ACCAT</u> --GGCTTAGTGGT-CCTAGTa
123	15	ctCAAGTAGGG--CGGAC-ACAC----GTCCG-GGC-ACC---TAA-----GGT-CCCAactgag
104	18	ctGGG--CGCCCTAAACAA--AGGGTG-GTC- <u>ACT</u> ---TCT-----AGT-CCCAGGA
106	20	ATGGGAGGG--CGCG--TTCTT----CGTG-GTT- <u>ACT</u> ---TTT-----AGT-CCCG
120	25	ctTTGGG--CGTG--AATGTC---CACG-GGT- <u>ACC</u> ---TCC-----GGT-CCCAAAGAG
105	30	TCCGGG--CTCGG-GAT-----TCGTG-GTC- <u>ACT</u> ---TTC-----AGT-CCCGGATATA
121	31	5'TCCGCGCAAGT-CCCTGGTAA= =AGGG--CAG--CCCTAA----CTG-GTC-actgagc3'
107	35	ACGGGAGGG--CACG--TTCTT----CGTG-GTT- <u>ACT</u> ---TTT-----AGT-CCCG
94	41	=GGG--CTGAGTa3' 5'tactCAG-GGC- <u>ACTGCAAGCAATTGTGGT-CCCAAT</u> =
103	47	GTGGGTGGGATCGGGG--ATG-----CCTC-GTC- <u>ACT</u> ---TCT-----AGT-CCcact

[0244]

TABLE 10

Frequency of base pairs in the helical regions of the consensus motif shown in FIG. 3							
Position ^a	Base pair ^b						other
	AT	TA	GC	CG	TG	GT	
I-1	0	0	21	0	0	0	0
I-2	0	0	21	0	0	0	0
I-3	5	0	16	0	0	0	0
I-4	3	5	1	4	1	0	7
I-5	2	3	3	4	0	0	9
II-1	0	1	2	17	0	0	1
II-2	5	5	5	1	0	4	1
II-3	3	4	7	6	0	0	1
II-4	3	0	8	5	0	0	4
III-1	21	0	0	0	0	0	0
III-2	0	10	0	11	0	0	0
III-3	0	7	0	13	1	0	0

^aHelices are numbered with roman numerals as shown in FIG. 3. Individual base pairs are numbered with arabic numerals starting with position 1 at the helix junction and increasing with increased distance from the junction.

^bWe have included the TG and GT base pairs to the Watson-Crick base pairs for this analysis. There is a total of 21 sequences in the set.

[0245]

TABLE 11

Affinities of the minimal DNA ligands to PDGF AA, PDGF AB and PDGF BB			
K _d , nM			
Ligand	PDGF AA ^a	PDGF AB ^b	PDGF BB ^b
20t	47 ± 4	0.147 ± 0.011	0.127 ± 0.031
36t	72 ± 12	0.094 ± 0.011	0.093 ± 0.009
41t	49 ± 8	0.138 ± 0.009	0.129 ± 0.011

^aData points shown in FIG. 5A were fitted to eq 1 (Example 7).

^bData points in FIGS. 5B and 5C were fitted to eq. 2. The dissociation constant (K_d) values shown are for the higher affinity binding component. The mole fraction of DNA that binds to PDGF AB or PDGF BB as the high affinity component ranges between 0.58 to 0.88. The K_d values for the lower affinity interaction range between 13 to 78 nM.

[0246]

TABLE 12

Starting RNA and PCR primers for the 2'-fluoropyrimidine RNA SELEX experiment		SEQ ID NO
Starting 2'-fluoropyrimidine RNA: Starting RNA: 5'-GGGAGACAAGAAUAACGCUCAA[-50 N-] UUCGACAGGAGGCUCACAACAGGC-3'		125

TABLE 12-continued

Starting RNA and PCR primers for the 2'-fluoropyrimidine RNA SELEX experiment	
	SEQ ID NO
PCR Primer 1: 5'-TAATACGACTCACTATAGGGAGACAAGAATAACGCTCAA-3'	126
PCR Primer 2: 5'-GCCTGTTGTGAGCCTCCTGTCGAA-3'	127

[0247]

TABLE 13

Sequences of the 2'-fluoropyrimidine RNA high affinity ligands to PDGF AB.	
	SEQ ID NO.
1 CGGUGGCAUUUCUUCACUCCUUCUCGCUUUCUCGCGUUGGGCNCGA	128
2 CCAACCUUCUGUCGCGUUGCUUUUUGGACGGCACUCAGGCCUCA	129
3 UCGAUCGGUUGUGUGCCGGACAGCCUUAACCAGGGCUGGACCGAGGCC	130
4 CUGAGUAGGGGAGGAAGUUGAAUCAGUUGUGGCGCCUCUCAUUCGC	131
5 CAGCACUUUCGCUUUUCAUCAUUUUUUCUUCACUGUUGGGCGCGGAA	132
6 UCAGUGCUGGCGUCAUGUCUCGAUGGGGAUUUUUCUUCAGCACUUUGCCA	133
7 UCUACUUUCCAUUUCUCUUUUUCUUCACGAGCGGGUUCAGUGAACCA	134
8 CGAUAGUGACUACGAUGACGAAGGCCGCGGGUUGGAUGCCCGCAUUGA	135
10 GUCGAUACUGGCGACUUGCUCCAUUGGCCGAUUAACGAUUCGGUCAG	136
13 GUGCAAACUUAACCCGGGAACCGCGCUUUCGAUCGACUUUCUUUCCA	137
15 AUUCCGCGUUCGCAUUAUUCUGUCUCGGAUUCGGUAGCCAUAGUGCA	138
16 CGAACGAGGAGGGAGUGGCAAGGGAUGGUUGGAUAGGCUCUACGCUCA	139
17 GCGAAACUGGCGACUUGCUCCAUUGGCCGAUUAACGAUUCGGUUCAU	140
18 CGAACGAGGAGGGAGUCGCAAGGGAUGGUUGGAUAGGCUCUACGCUCAA	141
19 CGAGAAGUGACUACGAUGACGAAGGCCGCGGGUUGAAUCCCUCAUUGA	142
20 AAGCAACGAGACCUGACGCCUGAUGUGACUGUGCUUGCACCCGAUUCUG	143
21 GUGAUUCUCAUUCUCAAUGCUUUCACACAACUUUUCCACUUCAGCGUGA	144
22 AAGCAACGAGACUCGACGCCUGAUGUGACUGUGCUUGCACCCGAUUCU	145
23 UCGAUCGGUUGUGUGCCGGACAGCUUUGACCAUGAGCUGGGACCGAGGCC	146
24 NGACNGUGGACCUGACUAAUCGACUGAUCAAAGAUCGCCCCAGAUUGGG	147
26 CACUGCGACUUGCAGAAGCCUUGUGUGGCGUACCCCUUUGGCCUCG	148
27 GGUGGCAUUUCUCAAUUUUCUUCUCGCUUUCUCCGCCGUUGGGCGCG	149
29 CCUGAGUAGGGGGAAAGUUGAAUCAGUUGUGGCGCUCUACUCAUUCGCC	150
30 GUCGAAACUGGCGACUUGCUCCAUUGGCCGAUUAACGAUUCGGUUCA	151
31 GCGAUACUGGCGACUUGCUCCAUUGGCCGAUUAACGAUUCGGUCAG	152

TABLE 13-continued

Sequences of the 2'-fluoropyrimidine RNA high affinity ligands to PDGF AB.		SEQ ID NO.
32	ACGUGGGGCACAGGACCGAGAGUCCUCCGGCAAUAGCCGCUACCCACC	153
33	CACAGCCUNANAGGGGGGAAGUUGAAUCAGUUGUGCGCUCUACUCAUUCGC	154
34	ANGGGNUAUGGUGACUUGCUCCAUUGGCCGAUAUAACGAUUCGGUCAG	155
35	CCUGCGUAGGGGGGAAGUUGAAUCAGUUGUGCGCUCUACUCAUUCGCC	156
39	CGAACGAGGAGGGAGUGGCAAGGGAUGGUUGGAUAGGCUCUACGCUCA	157
41	GUGCAAACUUAACCCGGGAACCGCGUUUCGAUUCGCUUCCNUAUCCA	158
42	CGAACGAGGAGGGAGUGGCAAGGACGGUNNAUAGGCUCUACGCUCA	159
43	UCGGUGUGGCUCAGAAACUGACACGCGUGAGCUUCGCACACAUCUGC	160
44	UAUCGCUUUUCAUCAAUCCACUUUUUCACUCUNUAACUUGGGCGUGCA	161
45	GUGCAAACUUAACCCGGGAACCGCGUUUCGAUCCUGCAUCCUUUUUC	162
46	UCGNUMCGUUGUGUGCCGGCAGCUUUGUCCAGCGUUGGGCCGAGGCC	163
47	AGUACCCAUCUCAUCUUUCCUUUCCUUUCUUAAGGCACAUUGAGGGU	164
49	CCUGAGUAGGGGGGAAGUUGAACAGUUGUGGCCUACUCAUUCNCCA	165
51	CCNNCCUNCUGUGCGCGCUUGUCUUUUUGGACGGGCAACCCAGGGCUC	166
52	CCAACCUNCUGUGCGCGCUUGUCUUUUUGGACGAGCAACUCAAGGCUCGU	167
53	CCAGCGCAGAUCCCGGGCUGAAGUGACUGCCGGCAACGGCCGCUCCA	168
54	UUCGCGAACAACUUUCAUUUUCACUUUUCAUCCAACAGUGAGCAGCA	169
55	UAUCGCUUUUCAUCAAUCCACUCCUUCACUUCUUUAACUUGGGCGUGCA	170

[0248]

TABLE 14

Starting RNAs:	
40N7:	
5'GGGAGGACGAUGCGG[-40N-]CAGACGACUCGCCCGA 3'	(SEQ ID NO:186)
SELEX PCR Primers:	
5G7:	
5' <u>TAATACGACTCACTATAGGGAGGACGATGCGG</u> 3'	(SEQ ID NO:187)
T7 Promoter	
3G7:	
5'TCGGGCGAGTCGTCG 3'	(SEQ ID NO:188)

[0249]

TABLE 15

Conditions and progress of the SELEX against hKGF										
Round	[RNA], M	[KGF], M	net % bound	Signal/noise	PF ^a	Spin ^b	B-Wash ^c (ml)	U-Wash ^d (ml)	SPKD ^e , M	KD ^f , nM
1	1.00E-06	3.00E-07	4.4	11.8			4		5.61E-06	30.0
2	4.00E-06	3.00E-07	1.5	4.2			5		1.58E-05	

TABLE 15-continued

Conditions and progress of the SELEX against hKGF										
	[RNA], M	[KGF], M	net % bound	Signal/noise	PF ^a	Spin ^b	B-Wash ^c (ml)	U-Wash ^d (ml)	SPKD ^e , M	KD ^f , nM
3	1.00E-06	1.00E-07	5.9	20.6			5		8.52E-07	
4	1.00E-06	1.00E-07	14.3	12.8	+		8		3.21E-06	17.0
5	3.00E-07	1.00E-08	2.5	4.5	+		8		7.64E-08	
6	3.70E-08	3.70E-09	0.7	2.6	+	+	15	15	3.73E-07	
7	4.10E-09	4.10E-10	1.1	8.2	+	+	20	20	2.46E-08	0.7
8	4.60E-10	4.60E-11	1.5	8.8	+	+	25	25	2.04E-09	0.3
9	5.10E-11	5.10E-12	0.7	5.9	+	+	25	25	8.76E-10	
10	1.70E-11	1.70E-12	0.3	2.1	+	+	25	25	4.12E-10	
2F SELEX										
1	1.00E-06	3.00E-07	2.9	11.0			4		3.39E-06	30.0
2	4.00E-06	3.00E-07	2.2	9.9			5		9.28E-06	
3	3.00E-06	3.00E-07	5.7	5.7			5		2.15E-06	
4	2.50E-06	3.00E-07	3.9	11.7	+		8		4.98E-06	15.0
5	6.70E-07	3.00E-08	2.3	5.8	+		8		3.64E-06	
6	1.20E-08	1.23E-09	0.3	1.8	+	+	15	15	1.59E-07	
7	1.40E-09	1.40E-10	1.1	11.2	+	+	20	20	6.86E-09	0.6
8	1.50E-10	1.50E-11	0.4	4.8	+	+	25	25	5.36E-10	0.3
9	1.70E-11	1.70E-12	0.2	3.1	+	+	25	25	5.67E-10	
10	1.70E-11	1.70E-12	0.3	3.0	+	+	25	25	1.42E-10	

^aPrefiltered RNA through nitrocellulose to counter select for nitrocellulose binding molecules

^bBrief spinning of the binding reactions

^cVolume of buffer used to wash the captured complexes

^dVolume of 0.5 M urea wash following the buffer wash

^eCalculated single point K_D from the binding data at each round

^fK_D values obtained from binding curves

[0250]

TABLE 16

Sequences of 2'-NH ₂ and 2'-F KGF ligands				
Clone	5' constant	random	3' constant	SEQ ID NO:
2'-NH ₂ ligands:				
1N	GGGAGGACGAUGC	GAAGGACGAUAAAGAGGAAUCGAAACAACAGUGGCGG	CAGACGACUCGCCCGA	189
2N	GGGAGGACGAUGC	GCGGGAAGGUCCGAAGACCGGGCAAAGGAACGAGAUUGCC	CAGACGACUCGCCCGA	190
4N	GGGAGGACGAUGC	GUGGUGAAGAGGUACCGGAAUUGCUAAAGAUACCACGGCC	CAGACGACUCGCCCGA	191
6N	GGGAGGACGAUGC	GCAGGAGCAAUGAACUCAAGUCAAGCCGGUGCACGUGGG	CAGACGACUCGCCCGA	192
10N	GGGAGGACGAUGC	UAGCUGCUGUCAUGCAAGACACUAGAAGAUUAAGAUUGGG	CAGACGACUCGCCCGA	193
11N	GGGAGGACGAUGC	GGGCCGGAUUUGAACCGACGACUUCGGGUUAUGAGCCCGACGU	CAGACGACUCGCCCGA	194
14N	GGGAGGACGAUGC	UCCAGGGAUUGAAGUUGUCGGGGUAGGAACAUAAGGCGGC	CAGACGACUCGCCCGA	195
16N	GGGAGGACGAUGC	AAGUUCUAACAAGUUAGUGGAAGGUUCCACUUGAAUGUA	CAGACGACUCGCCCGA	196
22N	GGGAGGACGAUGC	AUGGAGCUGAAA	CAGACGACUCGCCCGA	197
24N	GGGAGGACGAUGC	GUGGGAAGAUAGCCGGUCGCGCAGUUAUGUACACUCGGC	CAGACGACUCGCCCGA	198
25N	GGGAGGACGAUGC	GAGGGAUUGAGGAAACAACUAGCAGAUAAACGAGCUGGC	CAGACGACUCGCCCGA	199
27N	GGGAGGACGAUGC	AUGGAGCUGAAA	CAGACGACUCGCCCGA	200
28N	GGGAGGACGAUGC	UUGCUUACAUAUGACGCGGUGACUCUCCAGUUCUUGGACA	CAGACGACUCGCCCGA	201
29N	GGGAGGACGAUGC	GAGGGGAGAAUGCAGGAAACAGCGAAUUGCGUGUGGC	CAGACGACUCGCCCGA	202
34N	GGGAGGACGAUGC	GCGGGAAGAGCUAAUGGAAGUGGAAUCAGUCACAGUGCGG	CAGACGACUCGCCCGA	203
35N	GGGAGGACGAUGC	GCUUAGGGAAAUGGUUCUGAGGUGGU	CAGACGACUCGCCCGA	204
36N	GGGAGGACGAUGC	GAAGGGAACAGGAUAAGACAAGUCGAAACAAGCCGAGGUG	CAGACGACUCGCCCGA	205
37N	GGGAGGACGAUGC	AUGGAGCUGAAA	CAGACGACUCGCCCGA	206
42N	GGGAGGACGAUGC	GGAGACGUAAGCGGGAACAUAAGAACCAACAACCGCGGC	CAGACGACUCGCCCGA	207
43N	GGGAGGACGAUGC	GAAGUGGAUAGAACAGUCAGAAAUGUAAGCGUGAGGUG	CAGACGACUCGCCCGA	208
47N	GGGAGGACGAUGC	GAAGGGUAGGAAGGUCAAGAGGAAACAGCGCUUCGGGGUG	CAGACGACUCGCCCGA	209
48N	GGGAGGACGAUGC	GGCAAAGGAAGUUGGAAUCGGGACUAAGUAGUGUGUGGC	CAGACGACUCGCCCGA	210
54N	GGGAGGACGAUGC	AGAACAACAGAGCCUUCUGGUGGUGGGGAAGGAUUUCU	CAGACGACUCGCCCGA	211
55N	GGGAGGACGAUGC	ACACACAAGUGAAGGUACAGACGCGAAUUAACGUGGGUGGG	CAGACGACUCGCCCGA	212
57N	GGGAGGACGAUGC	UCGUGGGGUGGGUGGGGCGAGCUUGGAAUUAAGUAACUGGUAACGGCUGGC	CAGACGACUCGCCCGA	213
59N	GGGAGGACGAUGC	GGUGGGUGGUUACCUUUAUAUAUUAUUAUCUGGCUUUAG	CAGACGACUCGCCCGA	214
60N	GGGAGGACGAUGC	CCCUUAGCUCAGUGGUUAGAG	CAGACGACUCGCCCGA	215
65N	GGGAGGACGAUGC	UAACGUGGAAUAGGGUUAACAACGUGGAAUUAACGUAAGGUGGC	CAGACGACUCGCCCGA	216
69N	GGGAGGACGAUGC	GUAGGAGUAGGACAGACAUAAACAGUGCAACCAUCUGGCG	CAGACGACUCGCCCGA	217

TABLE 16-continued

<u>Sequences of 2'-NH₂ and 2'-F KGF ligands</u>				
Clone	5' constant	random	3' constant	SEQ ID NO:
71N	GGGAGGACGAUGC	AAACGGCGUGGCAAAAGUGAGGGGGUAGGAUGUACCAUGGGU	CAGACGACUCGCCCGA	218
72N	GGGAGGACGAUGC	GAGGGGAAAAGAGACCCAGACGAUUGACGGAAGUACUGGG	CAGACGACUCGCCCGA	219
2'-F ligands:				
2F	GGGAGGACGAUGC	GCAUUCGUCAAUACCUUGUUUUUUAUCCUUUUUCUAGCGGCC	CAGACGACUCGCCCGA	220
3F	GGGAGGACGAUGC	AUCGUAAUCGCCACUACUACUUCUCCGAAACCCGACGUGGC	CAGACGACUCGCCCGA	221
5F	GGGAGGACGAUGC	CGUCCCGAGUCACGCUGUCCUGUAACCUUCUCUGUGCC	CAGACGACUCGCCCGA	222
6F	GGGAGGACGAUGC	GAUCCUUUGUGGGUCUUGUUGACCCUCCUUGUGCCCGCC	CAGACGACUCGCCCGA	223
7F	GGGAGGACGAUGC	CGGUACUCUUCGCCAGCUCUCCAAAGCGCGACCUGUGCC	CAGACGACUCGCCCGA	224
8F	GGGAGGACGAUGC	UUUCGAAUAGGGCCAUUUCUCACUAGCUAUCCUACCCUGCC	CAGACGACUCGCCCGA	225
9F	GGGAGGACGAUGC	AUAAUGGCUAGAACUAGCUCGCAUCUUGGUGUCCGGUGCC	CAGACGACUCGCCCGA	226
10F	GGGAGGACGAUGC	GACCAGAUGGGGAUUUUUCAGCAAUCCUCCCGUGCC	CAGACGACUCGCCCGA	227
11F	GGGAGGACGAUGC	UGAUGGCGACCAGUCAAACCGGUGUUUACUCCCGCCG	CAGACGACUCGCCCGA	228
12F	GGGAGGACGAUGC	GAAUUAAACAGGGCCAGAAUUUCUACUUNNCUCCCGUGACC	CAGACGACUCGCCCGA	229
13F	GGGAGGACGAUGC	CACCUUAGACCUGUCCUCCAAAGCGUGAGUUGCUGUGCC	CAGACGACUCGCCCGA	230
14F	GGGAGGACGAUGC	UGGUCUCCCAAUUUCUAAACUUUCUCCAUUGUAUCUGGGC	CAGACGACUCGCCCGA	231
15F	GGGAGGACGAUGC	UCAUGGUGUCUUUCCACAGCUCUUCCCAUGAUCGCCCGC	CAGACGACUCGCCCGA	232
16F	GGGAGGACGAUGC	GAAUCCAGCGCUUGACUGAUACAACNNUCCCGUGCC	CAGACGACUCGCCCGA	233
19F	GGGAGGACGAUGC	CAA-NNNNNNNCUCUCUCCUGGCGUCCCGCAACCCGCC	CAGACGACUCGCCCGA	234
20F	GGGAGGACGAUGC	AGUAUCCAGCCUGGAUUGCAUAGUCAGUGCUCUCCGUGCC	CAGACGACUCGCCCGA	235
21F	GGGAGGACGAUGC	UCCUAGCAGCGAUUCAUCCCGUUCUCUCAGCGUUGCC	CAGACGACUCGCCCGA	236
22F	GGGAGGACGAUGC	CCUGAAGUACAGCGUCUAAACUCCAAAGCGGACCGCCG	CAGACGACUCGCCCGA	237
23F	GGGAGGACGAUGC	CCCUACCACUUUUUCCUCUACUGUUAUCCUGUGCC	CAGACGACUCGCCCGA	238
24F	GGGAGGACGAUGC	UGGUCUCCUAGAUUCACAGCACUCCAUUGCAUUGGGC	CAGACGACUCGCCCGA	239
26F	GGGAGGACGAUGC	UCAAGCUUAACAGUCUGGCAAUUGGCAUUAUGGCGCC	CAGACGACUCGCCCGA	240
27F	GGGAGGACGAUGC	CaGUCUGGAUCUCUAUUGGAAUUUAGUCCUACAUCUGGCC	CAGACGACUCGCCCGA	241
28F	GGGAGGACGAUGC	GAUUCUUUCGGCAGUGAAAAUAUCCUUGCUUCCCGAGC	CAGACGACUCGCCCGA	242
29F	GGGAGGACGAUGC	GGACUUCAAACUAGUCUUAUUGCCUCGUCUCCUGUGCC	CAGACGACUCGCCCGA	243
31F	GGGAGGACGAUGC	AACGGAGAUGUCCCUCAAMAUUUACCGUCUCCGCUUUGCGCC	CAGACGACUCGCCCGA	244
35F	GGGAGGACGAUGC	CGAAUUAGCUUCUUAUGACUCACGUUCCUUGCCGCC	CAGACGACUCGCCCGA	245
37F	GGGAGGACGAUGC	GCCCCAUCUACUGCAUUCCGAAACGAUUUCCCGACUGUG	CAGACGACUCGCCCGA	246
38F	GGGAGGACGAUGC	NGACUGAUUUUUCUUGNCGAGUGUAUUUCCUGGCGGCC	CAGACGACUCGCCCGA	247
41F	GGGAGGACGAUGC	GGACUUUGACAGGCAUUGAUUUCGACCUGUUCUCCCGUGGC	CAGACGACUCGCCCGA	248
42F	GGGAGGACGAUGC	CGACACAAUAGCCUUUGAUUCCAUAGUUGGUCGCGCGUGCC	CAGACGACUCGCCCGA	249
43F	GGGAGGACGAUGC	UGUAGUUUCCUGUAUGCCAUUUCCUCCUAGCCGACCGC	CAGACGACUCGCCCGA	250
44F	GGGAGGACGAUGC	UCGAGUGUUCUCCUUCGGUAACUAUUNNNNAUUUCGUGCC	CAGACGACUCGCCCGA	251
45F	GGGAGGACGAUGC	GUCGUAUUCAUCUCCUUGUUCUGUUCUUGCACCUGGGC	CAGACGACUCGCCCGA	252
49F	GGGAGGACGAUGC	GGACUUUGACAGGCAUUGAUUUCGACGUGUUCUCCCGUGGC	CAGACGACUCGCCCGA	253
50F	GGGAGGACGAUGC	UGAUCAAUCGGCGUUUACUCUUGCGCUCACCGUGGCC	CAGACGACUCGCCCGA	254
51F	GGGAGGACGAUGC	CAGUCUCCUAGGUUUAUCUUCUGCAGCAUUCGGGGUCC	CAGACGACUCGCCCGA	255
53F	GGGAGGACGAUGC	AUCAAAGCACUCUACUCCCGUGUCGUCUUAUUGGUCUCC	CAGACGACUCGCCCGA	256
54F	GGGAGGACGAUGC	AAGAUCCUCCAAUCUGCUGUGGCUAAUUAUUCUCCCGUCC	CAGACGACUCGCCCGA	257
55F	GGGAGGACGAUGC	UCCGUCAUAACGGCCAUAAACUGCGAAUACUCCUGGGC	CAGACGACUCGCCCGA	258
56F	GGGAGGACGAUGC	GGACAAYAGCGGUGUUCUUAUUNKAUCCUCCGACRUCUCC	CAGACGACUCGCCCGA	259
57F	GGGAGGACGAUGC	UGACUUAUCUGGCUAGUCCAAUACCCGAGCCACCAGCGC	CAGACGACUCGCCCGA	260
58F	GGGAGGACGAUGC	GAACUAAUGGCCGUGAUUAAACAAUGCAGGCUUCCUGCG	CAGACGACUCGCCCGA	261
60F	GGGAGGACGAUGC	UGACAUGGAAUUUUCUACGGGCCCGAUCCUGCCAGCGGUGUG	CAGACGACUCGCCCGA	262

[0251]

TABLE 17

<u>K_d values hKGF ligands</u>					
Clone	K _d in nM		Clone	K _d in nM	
	1	2		1	2
1N	0.51		2F	1.77	
2N	0.77		3F	4.47	
4N	0.75		5F	2.53	
6N	0.71		6F	0.05 (37)	3.25
10N	1.10		7F	3.69	
11N	1.28		8F	2.63	
14N	0.44		9F	0.83	
16N	1.40		10F	0.47	

TABLE 17-continued

<u>K_d values hKGF ligands</u>					
Clone	K _d in nM		Clone	K _d in nM	
	1	2		1	2
22N	5.70		11F	3.74	
24N	1.16		12F	1.38	
25N	0.87		13F	0.03 (28)	
27N	ND		14F	0.006-0.03 (25-44)	
28N	2.54		15F	0.07 (33)	
29N	0.43		16F	0.83 (49)	3.39
34N	0.80		19F	1.6	0.94-2.57

TABLE 17-continued

<u>K_d values hKGF ligands</u>					
Clone	<u>K_d in nM</u>		Clone	<u>K_d in nM</u>	
	1	2		1	2
35N	2.32		20F	2.05	8.70
36N	8.27		21F	ND	44.8
37N	ND		22F	2.75	
42N	0.78		23F	2.52	
43N	0.79		24F	2.02	
47N	1.76		26F	0.23 (43)	2.55
48N	1.34		27F	1.52	
54N	5.35		28F	ND	
55N	1.25		29F	3.24	
57N	35.8		31F	1.0	
59N	22.0		35F	1.1	
60N	7.38		37F	0.46	
65N	26.56		38F	0.33	
69N	15.20		41F	1.44	
71N	3.52		42F	0.9	
72N	7.67		43F	1.13	
random	30		44F	1.32	
			45F	4.7	
			49F	1.0	
			50F	0.12 (21)	2.10
			51F	1.27	
			53F	0.70	
			54F	1.23	
			55F	2.52	
			56F	0.07 (32)	3.00
			57F	1.20	
			58F	2.52	
			60F	2.10	
			random	30	

For biphasic curves, Kd1 is for the high affinity component. Number in parentheses indicate the per cent of the high affinity component.

[0252]

TABLE 18

<u>Binding Specificity of the 2'-F Ligand K14F</u>		
Target	<u>K_D Target/K_D hKGF</u>	Ratio:
human hKGF		1
rat hKGF		1,254
human aFGF		38,650
human bFGF		1,071
human PDGF		432

The ratios shown are averages of at least two determinations

[0253]

TABLE 19

<u>IC₅₀ values from the PC-3 assay</u>	
Competitor	IC50, nM
hKGF	70
Heparin, 5,000	30
40N7F	>1000
K6F	4
K13F	30
K14F	10
K15F	20
K56F	1

TABLE 19-continued

<u>IC₅₀ values from the PC-3 assay</u>	
Competitor	IC50, nM
K10F	30
K37F	20
K38F	0.6
K43F	80
40N7N	>1000
K1N	50
K2N	200
K4N	70
K6N	80
K14N	6
K29N	40
K42N	800
K43N	800

[0254]

TABLE 20

<u>Ki values of hKGF competitors on the PC3 and NIH3T3/FGFR-2 competition assay</u>			
Cell line	Competitor	Ki, nM	R
PC-3	hKGF	7.700	0.95519
	2'F random	930.000	0.99713
	2'NH ₂ random	673.000	0.85357
	Hep5000	6.500	0.99984
	K14F	0.200	0.97735
	K6F	0.160	0.95927
	K38F	0.220	0.99013
	K56F	0.160	0.95927
	K14N	1.400	0.94698
	NIH3T3/FGFR-2	hKGF	0.034
2'F random		>10,000.000	
2'NH ₂ random		>10,000.000	
Hep5000		26.300	0.97856
K14F		2.700	partial comp. 0.99047
K6F		6.800	0.96202
K38F		20.000	0.98659
K56F		27.400	0.97582
K14N		10.600	0.97856
			partial comp.

[0255]

TABLE 21

<u>IC50 values obtained with the gel shift assay</u>	
Competitor	IC50, nM
KGF	70
bFGF	1,500
Lysozyme	10,000

[0256]

TABLE 22

Protein	Binding Specificity of Ligand K14F3'T				
	random RNA		K14F3'T		°DF
	^a K _d 1, nM	^b K _d 2, nM	K _d 1, nM	K _d 2, nM	
hKGF		20.1	0.0008	10.2	1
rKGF		45.3	0.0041	70.0	5
hbFGF	0.0375	10.3		10.0	1.2 × 10 ⁴
haFGF		16,000,000		24,000,000	3 × 10 ¹⁰
hPDGF-AB		22.0		50.0	6.2 × 10 ⁴
hTGFβ1		10.4		98.0	1.2 × 10 ⁵

TABLE 22-continued

Protein	Binding Specificity of Ligand K14F3'T				
	random RNA		K14F3'T		°DF
	^a K _d 1, nM	^b K _d 2, nM	K _d 1, nM	K _d 2, nM	
hEGF		2,000		256	3.2 × 10 ⁵
Thrombin		7,200,000		22,700,000	2.8 × 10 ¹⁰

^aHigh affinity dissociation constant from biphasic binding curves.
^bLow affinity dissociation constant from biphasic binding curves or affinity dissociation constant from monophasic binding curves.
 °Discrimination factor defined as the ratio of the highest affinity K_d of 14F3'T for the corresponding protein over the affinity K_d for hKGF.

[0257]

TABLE 23

	S1	L1	S2	S1'	L3	S2'	H L	SEQ ID NO:
143 'T	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUAAACUUUCU	CCAUCGUAUC		272
T2	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUAAACUUUCU	CCAUCGUA..	- +	273
T3	GGGAGG	AC	..UGCGGUGG	UCUCCC	AAUUCUAAACUUUCU	CCAUCGUA..	--	274
T4	GGGAGG	AC	GAUGCGGUGG	UCUCCC	...CUAAACUUUCU	CCAUCGUAUC	++	275
T5	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA...UUCU	CCAUCGUAUC	++	276
T6	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUAAACU...	CCAUCGUAUC	± +	277
T7	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAU...AACU...	CCAUCGUAUC	--	278
T8	..GAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUAAACUUUCU	CCAUCGUAUC	- +	279
T10	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAU.....UUCU	CCAUCGUAUC	--	280
T11	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA.....	CCAUCGUAUC	--	281
T12	.GGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA...UUCU	CCAUCGUAUC	--	282
T13	.GGAGG	AC	GAUGCGGUGG	UCUCC.	AAUUCUA...UUCU	CCAUCGUAUC	--	283
T14	..GAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA...UUCU	CCAUCGUAUC	--	284
T15	..GAGG	AC	GAUGCGGUGG	UCUC..	AAUUCUA...UUCU	CCAUCGUAUC	--	285
T16	GGGAGG	.C	GAUGCGGUGG	UCUCCC	AAUUCUA...UUCU	CCAUCGUAUC	- ±	286
T18	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA...UCU	CCAUCGUAUC	++	287
T19	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA.....CU	CCAUCGUAUC	--	288
T20	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCUA.....U	CCAUCGUAUC	--	289
T21	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUCU...UUCU	CCAUCGUAUC	++	290
T22	GGGAGG	AC	GAUGCGGUGG	UCUCCC	AAUUC...UUCU	CCAUCGUAUC	++	291
T29	GGG.G.	AC	GAUGCGGUGG	.C.CCC	AAUUCUA...UUCU	CCAUCGUAUC	--	292
T30	GGG...	AC	GAUGCGGUGG	...CCC	AAUUCUA...UUCU	CCAUCGUAUC	--	293
T31G	AC	GAUGCGGUGG	...CCC	AAUUCUA...UUCU	CCAUCGUAUC	--	294
T32GG	AC	GAUGCGGUGG	...CCC	AAUUCUA...UUCU	CCAUCGUAUC	--	295
T33	GAUGCGGUGG	...CCC	AAUUCUA...UUCU	CCAUCGUAUC	--	296
T34	GAUGCGGUGG	UCUCCC	AAUUCUA...UUCU	CCAUCGUAUC	--	297

TABLE 23-continued

T35GGUGG UCUCCC AAUUCUA.....UUCU CCAUCGUAUC	--	298
T36	GGGAGG AC GAUG.....	--	299
T37	GGGAGG AC GAUGCGGUGG UCUCCC AAUUC.....UCU CCAUCGUAUC	--	300
T39	GGGAGG AC GAUGCGGUGG UCUCCC AAUUC.....UUCU .GAUCGUAUC	--	301
T40	GGGAGG AC GAUGCGG.GG UCUCCC AAUUC.....UUCU CC.UCGUAUC	--	302
T41	GGGAGG AC GAUGCGG.G. UCUCCC AAUUC.....UUCU .C.UCGUAUC	--	303
T22mu	GGGAGU AC GAUGCGGUGG UCUCCC AAUUC.....UUCU CCAUCGUAUC	--	304
T35/36 equimolar amounts of T35 and T36			--

[0258]

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 304

(2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGGGGAGAAC GCGGAUCCNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50
 NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNAAGCUUCG CUCUAGAUCU 100
 CCCUUUAGUG AGGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 93 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GGGGGAGAAC GCGGAUCCNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50
 NNNNNNNNAA GCUUCGCUCU AGAUCUCCCU UUAGUGAGGG UUA 93

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 123 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGGGGAGAAC GCGGAUCCUG UCUCACCAGC CGAUACUGGG GUUCCUGGGG 50

CCCCUCAUG GAGGAGGGGG GAGGGGGUGG UUCGGAGAAA GCUUCGCUCU 100

GAAUCUCCU UUAGUGAGGG UUA 123

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 95 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GGGAGAACGC GGATCCNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN AAGCTTCGCT CTAGA 95

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GGGAGAACGC GGATCCNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNNAAGC TTCGCTCTAG A 71

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 95 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GGGGGAGAAC GCGGATCCTG TCTCCACCGC CGATACTGGG GTTCCTGGGG 50

CCCTCCATG GAGGAGGGGG TGGTTCGGAG AAAGCTTCGC TCTAG 95

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TAATACGACT CACTATAGGG GGAGTCTGCG GATCC 35

-continued

(2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TAATACGACT CACTATAGGG GGAGAACGCG GATCC

35

(2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

TAACCCTCAC TAAAGGGAGA TCTAGAGCGA AGCTT

35

(2) INFORMATION FOR SEQ ID NO: 10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 50 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

GATTTAGGTG ACACTATAGA ATATGCATCA CTAGTAAGCT TTGCTCTAGA

50

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GGATCCCGGA GCTCCCTATA GTGAGTCGTA TTA

33

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 125 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

-continued

(D) OTHER INFORMATION: All pyri
modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GGGGGAGAAC GCGGAUCCGA GCAAUCCAG GCGCAUAGCU UCCGAGUAGA 50

CAGGAGGGAG GGGUGGAUGU GGCGUCUACU CGGUGUCGUG AAGCUUCGCU 100

CUAGAUCCU CUUUAGUGAG GGUUA 125

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 125 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All pyri
modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GGGGGAGAAC GCGGAUCCGA GCAACCCAG GCGCAUAGCU UCCGAGUAGA 50

CAGGAGGGAG GGGUGGAUGU GGCGUCUACU CGGUGUCGUG AAGCUUCGCU 100

CUAGAUCCU CUUUAGUGAG GGUUA 125

(2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 125 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All pyri
modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GGGGGAGAAC GCGGAUCCGA GCAACCCAG GCGCAUAGCU UCCGAGUAGA 50

CAGGCGGGAG GGGUGGAUGU GGCGUCUACU CGGAGUCGUG AAGCUUCGCU 100

CUAGAUCCU CUUUAGUGAG GGUUA 125

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 116 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All pyri
modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

GGGGGAGAAC GCGGAUCCGG CAACCCAGG CGCAUAGCUU CCGAGUAGAC 50

-continued

AGGCGGGAGG GGUGGAUGUG GCGUCACGAG GAAGCUUCGC UCUAGAUCUC	100
CCUUUAGUGA GGGUUA	116

(2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 123 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

GGGGGAGAAC GCGGATCCGC AAUCCAGGC GCAUAGCUUC CGAGUAGACA	50
GGAGGGAGGG GUGGAUGUGG CGUCUACUCG GCGUCGUGAA GCUUCGCUCU	100
AGAUCUCCCU UUAGUGAGGG UUA	123

(2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

GGGGGAGAAC GCGGAUCCGA GCAAUCCAG GCGCAUAGCU UCCGAGUAGA	50
CAGGAGGGAG GGGUGGAUGU GGCGUCUGA GGAAGCUUCG CUCUAGAUCU	100
CCCUUAGUG AGGGUUA	117

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 123 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

GGGGGAGAAC GCGGAUCCGA GCAAGCCUG GCAUAGCUUC CGAGUAGACA	50
GGAGGGAGGG GUGGAUGUGG CGUCUACUCG GUGUCGUGAA GCUUCGCUCU	100
AGAUCUCCCU UUAGUGAGGG UUA	123

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 115 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

GGGGGAGAAC GCGGAUCCGG CAAUCCAGG CGCAUAGCUU CCGAGUAGAC	50
AGGAGGGAGG GGUGGAUGUG GUGUACGAGG AAGCUUCGCU CUAGAUCUCC	100
CUUUAGUGAG GGUUA	115

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 116 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GGGGGAGAAC GCGGAUCCGA GCAAUCCAG GCGCAUAGCU UCCGAGUAGA	50
CAGGAGGGAG GGGUGGAUGU GGUGUCUGA GAAGCUUCGC UCUAGAUCUC	100
CCUUUAGUGA GGGUUA	116

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 98 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

GGGGGAGAAC GCGGAUCCAA GCUUCGAGUA GACAGGAGGG AGGGGUGGAU	50
GUGGAGUCUC GAGAAGCUUC GCUCUAGAUC UCCUUUAGU GAGGGUUA	98

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 113 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GGGGGAGAAC GCGGAUCCGA GCAAUCCUAA GCAUAGCUUC GAGUAGACAG	50
GAGGAGGGGG UGGAUGUGGC GUCUCGAGAA GCUUCGCUCU AGAUCUCCU	100

-continued

 UUAGUGAGGG UUA 113

(2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 116 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All U's

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All C's

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

GGGGGAGAAC GCGGAUCCGA GCAAUCCCGG GCGCAUAGCU UCCGAGGAGA 50

CAGGCGGGAG GGGUGGAUGU GCGGUCUCGA GAAGCUUCGC UCUAGAUCUC 100

CCUUUAGUGA GGGUUA 116

(2) INFORMATION FOR SEQ ID NO: 24:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 116 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All U's

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All C's

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GGGGGAGAAC GCGGAUCCGA GCAAUCCCGG GCGCAUAGCU UCCGAGUAGA 50

CAGGCGGGAG GGGUGGAUGU GCGGUCUCGA GAAGCUUCGC UCUAGAUCUC 100

CCUUUAGUGA GGGUUA 116

(2) INFORMATION FOR SEQ ID NO: 25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 42 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

GCUUCCGAGU AGACAGGAGG GAGGGGUGGA UGUGGCGUCU AC 42

(2) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:
 CUUCCGAGUA GACAGGAGGG AGGGGUGGAU GUGGCGUCUA CUC 43

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 78 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:
 GGGGGAGAAC GCGGAUCCGG CAACCCAGG CGCAUAGCUU CCGAGUAGAC 50
 AGGCGGGAGG GGUGGAUGUG GCGUCACG 78

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:
 GGGGGAGAAC GCGGAUCCUG AGAAGGACGU CGGGGUCAAC GGGGUGAGGU 50
 GCAGCAGAAA GGGCCGGCAC CACAUGACGU AAAAGCUUCG CUCUAGAUCU 100
 CCCUUUAGUG AGGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 108 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:
 GGGGGAGAAC GCGGAUCCUG AGAAGGACGU CGGGGUGAGG UGCAGCAGAA 50
 AGGGCCGGCA CCACAUGACG UAAAAGCUUC GCUCUAGAUC UCCUUUAGU 100

-continued

GAGGGUUA

108

(2) INFORMATION FOR SEQ ID NO: 30:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 92 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

- (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

GGGGGAGAAC GCGGAUCCGG UGGGAAAGUC GGAUUAUGUG UGUAGAUUUG 50
 UGUGCGAAAG CUUCGCUCUA GAUCUCCCUU UAGUGAGGGU UA 92

(2) INFORMATION FOR SEQ ID NO: 31:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 69 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

- (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

GGAGAACGCG GAUCCUGAGA AGGACGUCGG GGUCAACGGG GUGAGGUGCA 50
 GCAGAAAGGG CCGGCACCA 69

(2) INFORMATION FOR SEQ ID NO: 32:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

- (D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GGGGGAGAAC GCGGAUCCUG CUAGACCGAG GAUGCAAAGG GACAUGCAUU 50
 AGGGAACCU AUGUAUAAGA ACGCGUCGC AGAAGCUUCG CUCUAGAUCU 100
 CCCUUUAGUG AGGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 33:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 83 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

GGGGGAGAAC GCGGAUCCUG CUAGACCGAG GAUGCAAAGG GACAUGCAUU 50

AGGGAAACCU AUGUAUAAGA ACGCGGUCGC AGA 83

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 116 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

GGGGGAGAAC GCGGAUCCUG CUAGACCGAG GAUGCAAAGG GACAUGCAUU 50

AGGGAAACCU AUUAUAAGAA CGCGGUCGCA GAAGCUUCGC UCUAGAUCUC 100

CCUUUAGUGA GGGUUA 116

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All C's

(ix) FEATURE:
 (D) OTHER INFORMATION: All U's

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

GGGGGAGAAC GCGGAUCCUG UCUCACC GC GAUACUGGG GUUCCUGGGG 50

CCCCUCCAUG CAGGAGGGGG GUGGUUCGGA GAAAGCUUCG CUCUAGAUCU 100

CCCUUAGUG AGGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 36:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 81 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All C's

(ix) FEATURE:
 (D) OTHER INFORMATION: All U's

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

GGGGGAGAAC GCGGAUCCUG UCUCACC CGAUCUGGG GUUCCUGGGG 50

CCCCUCAUG CAGGAGGGG GUGGUUCGGA G 81

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 117 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All C's

(ix) FEATURE:

(D) OTHER INFORMATION: All U's

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

GGGGGAGAAC GCGGAUCCUG UCUCACC CGAUCUGGG GUUCCUGGGG 50

CCGCUCCAUG CAGGAGGGG GUGGUUCGGA GAAAGCUUCG CUCUAGAUCU 100

CCUUUAGUG AGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 38:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 115 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All C's

(ix) FEATURE:

(D) OTHER INFORMATION: All U's

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

GGGGGAGAAC GCGGAUCCUG UCUCACC CGAUCUGGG GUUCCUGGGG 50

CCCCUCAUG CAGGAGGGG GGUUCGAGA AAGCUUCGCU CUAGAUCUC 100

CUUUAGUGAG GGUUA 115

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 92 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All pyri modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

GGGGGAGAAC GCGGAUCCG AAGUCUGGUC UUUGGGAGU CCGCAUGGCC 50

-continued

 CUGGGCAAAG CUUCGCUCUA GAUCUCCUU UAGUGAGGU UA 92

(2) INFORMATION FOR SEQ ID NO: 40:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 111 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyri
 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

GGGGGAGAAC GCGGAUCCAA GAAUGUUCGG CCGCACGAGG UGACAGUGGU 50
 GCGGAUACGG ACCGAUUGGG UUUGCCAAGC UUCGCUCUAG AUCUCCUUU 100
 AGUGAGGGUU A 111

(2) INFORMATION FOR SEQ ID NO: 41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

GGGGGAGAAC GCGGAUCCGG UCACCCGGGC AUAUAACAAU GCCGACACUG 50
 GGGUACCUGG GACGGGUGGG ACUGGACGGA AGAAGCUUCG CUCUAGAUCU 100
 CCCUUUAGUG AGGGUUA 117

(2) INFORMATION FOR SEQ ID NO: 42:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 119 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

GGGGGAGAAC GCGGAUCCAU AACCGCUGC AUGGGAGGGA CAUCCUGGGA 50
 AAGGACGGGU CGAGAUGACC UGAGCAGUUC CGGCAAGCUU CGCUCUAGAU 100
 CUCCUUUAG UGAGGGUUA 119

(2) INFORMATION FOR SEQ ID NO: 43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

-continued

 TCGGGCGAGT CGTCTGNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNNCCGC ATCGTCCTCC C 71

(2) INFORMATION FOR SEQ ID NO: 44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

TAATACGACT CACTATAGGG AGGACCATGC GG 32

(2) INFORMATION FOR SEQ ID NO: 45:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

TCGGGCGAGT CGTCTG 16

(2) INFORMATION FOR SEQ ID NO: 46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

GGGAGGACGA TGCGNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 47:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

GGGAGGACGA TGCGG 15

(2) INFORMATION FOR SEQ ID NO: 48:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

-continued

(ix) FEATURE:
 (D) OTHER INFORMATION: N at positions 1-3 is biotin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:
 NNNTCGGGCG AGTCGTCTG 19

(2) INFORMATION FOR SEQ ID NO: 49:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 87 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:
 GCCTGTTGTG AGCCTCCTGT CGAANNNNNN NNNNNNNNNN NNNNNNNNNN 50
 NNNNNNNNNN NNNNTTGAGC GTTTATTCTT GTCTCCC 87

(2) INFORMATION FOR SEQ ID NO: 50:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 40 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:
 TAATACGACT CACTATAGGG AGACAAGAAT AAACGCTCAA 40

(2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:
 GCCTGTTGTG AGCCTCCTGT CGAA 24

(2) INFORMATION FOR SEQ ID NO: 52:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 87 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:
 GGGAGACAAG AATAAACGCT CAANNNNNNN NNNNNNNNNN NNNNNNNNNN 50
 NNNNNNNNNN NNNTTCGACA GGAGGCTCAC AACAGGC 87

(2) INFORMATION FOR SEQ ID NO: 53:

(i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:
 GGGAGACAAG AATAAACGCT CAA 23

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
 (D) OTHER INFORMATION: N at positions 1-3 is biotin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:
 NNNGCCTGTT GTGAGCCTCC TGTCGAA 27

(2) INFORMATION FOR SEQ ID NO: 55:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:
 GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTGGTGCT ACTTACTTGC 50
 GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:
 GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTAGTGCT ACTTACTTGC 50
 GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

-continued

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTAGTACT ACTTACTTAC 50

GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 70 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTATACTA CTTACTTACG 50

TCTTCAGACG ACTCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 59:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTAATACT ACTTACTTAC 50

ATCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 60:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTAATACT ACTTACTTAC 50

GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 61:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTGGTGTT ACTTACTTGC 50

GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 62:

- (i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTGGTGCT TCTTACTTGC 50

GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 63:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 73 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTGGTGTA CTTTTTCCTG 50

CGTCTTCCAG ACGACTCGCC CGA 73

(2) INFORMATION FOR SEQ ID NO: 64:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 72 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

GGGAGGACGA TCGGCCAGG GGGGGTATGG GGGTGGTTCG TTTTTCTTTG 50

CGGCTTCAGA CGACTCGCC GA 72

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

GGGAGGACGA TCGGCCAGG GGGGGTGTGG GGGTGGTGTA CTTTTCTTTG 50

TCTTCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

-continued

GGGAGGACGA TGCGGCCAGG GGGGGTATGG GGGTGGTTTG GTATGTTGCG 50
TCCGTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 67:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

GGGAGGACGA TGCGGCCGGG GTGGGTATGG GGGTAATACT ACTTACTTAC 50
GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO:68 :

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 72 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

GGGAGGACGA TGCGGCCGGG GGTGGGTAGG GGGTAGTGC TACTTACTTA 50
CGTCTTCAGAC GACTCGCCC GA 72

(2) INFORMATION FOR SEQ ID NO: 69:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

GGGAGGACGA TGCGGCCAGG GTCGGTGTGG GGGTAGTACT ACTTACTTGC 50
GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 70:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

GGGAGGACGA TGCGGCCAGG GTGGGTATGG GGGTAGTGCT ACTTACTTGC 50
GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 71:

-continued

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:
GGGAGGACGA TGCGCCGGG GTGGGTATGG GGGTGGTGCT ACTTACTTGC 50
GTCTTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 72:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:
GGGAGGACGA TGCGCCCTGG GTGGGTATGG GGGTGGTGCT ACTTACTTGC 50
GTCTTCAGACG ACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 73:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:
GGGAGGACGA TGCGGCCACG GGTGGGTGTG GGGTAGTGTG TCTCACTTA 50
CATCACCAGA CGACTCGCCC GA 72

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:
GGGAGGACGA TGCGGCCCGG GGTGGGTGTG GGGTAGTGTA TTATATTTAC 50
AGCCTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 75:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

GGGAGGACGA TCGGCCAGG GTCGGTGTGG GGTGGTGTAC TTTTCCTGT 50
CCTTCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

GGGAGGACGA TCGGCCAGG GTCGGTATGG GGTAGTGTAC TTTTAATGA 50
TCTTCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 77:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

GGGAGGACGA TCGGCCCGG GGGAGAGCGG TGGTAGTGT TCTATAGTAT 50
TCGTGTCAGA CGACTCGCCC GA 72

(2) INFORMATION FOR SEQ ID NO: 78:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

GGGAGGACGA TCGGCCAGG GGGGTATGT TTTAATACT ACTTACTTAC 50
GTCTTCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 79:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

GGGAGGACGA TCGGCCAGG GAGGGTATGG GGTGGTGT TCTAGTTTTG 50
CGGCGTCAGA CGACTCGCCC GA 72

(2) INFORMATION FOR SEQ ID NO: 80:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 72 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

GGGAGGACGA TCGGCCAGG GTGGGCATGG GGTGGTGTG GATTAATCT 50
 TCGTCCCAGA CGACTGCCCG GA 72

(2) INFORMATION FOR SEQ ID NO: 81:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

GGGAGGACGA TCGGCCAGG GTCGGTGTGG GGTGGTGTGTTT TTATTACTC 50
 GTCGCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 82:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

GGGAGGACGA TCGGGGGGC GGCTTGAAG AGGTTGCCGG TTGGAGTATT 50
 CGAGCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 83:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 67 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

GGGAGGACGA TCGGCCAGG TGTGGGTGG TTTGGTGTGTTT CTTCGTCGC 50
 CCAGACGACT CGCCCGA 67

(2) INFORMATION FOR SEQ ID NO: 84:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

GGGAGGACGA TCGGCCAGG GTGGGTATGG GGGTTAATT AATTCTTCGT 50

CCCACAGACG ACTCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 85:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

GGGAGGACGA TCGGGGGGC GGCTTGAAG AGGTGCGCG TTGGAGTATT 50

CGAGCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 86:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 72 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

GGGAGGACGA TCGGCCCGG GGTGGGTGTG GGTGGGTGTG AATTAATCTT 50

TCGTCCAGAC GACTCGCCCG GA 72

(2) INFORMATION FOR SEQ ID NO: 87:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

GGGAGGACGA TCGGCCCGG GGTGGGTGTG GGTGGGTGTA TTATATTTC 50

GGCCTCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 88:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

GGGAGGACGA TCGGCCAGG GTCGGTGTGG GTGGTGTACT TTTTCCTGTC 50

CTTCAGACG ACTCGCCCGA 70

-continued

(2) INFORMATION FOR SEQ ID NO: 89:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

GGGAGGACGA TCGGGGGGC GCCTTGAAG AGTTGCCGG TTGAGTATT 50
 CGAGCCAGAC GACTCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 90:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 86 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

ATCCGCCTGA TTAGCGATAC TNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50
 NNNNNNNNNN NACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 91:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 28 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(D) OTHER INFORMATION: N at positions 1-3 is biotin

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

NNNCCCTGC AGGTGATTTT GCTCAAGT 28

(2) INFORMATION FOR SEQ ID NO: 92:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 49 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

CCGAAGCTTA ATACGACTCA CTATAGGGAT CCGCCTGATT AGCGATACT 49

(2) INFORMATION FOR SEQ ID NO: 93:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 84 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

ATCCGCCTGA TTAGCGATAC TAGGCTTGAC AAAGGCACC ATGGCTTAGT 50

GGTCCTAGTA CTTGAGCAA ATCACCTGCA GGGG 84

(2) INFORMATION FOR SEQ ID NO: 94:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 85 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

ATCCGCCTGA TTAGCGATAC TCAGGGCACT GCAAGCAATT GTGGTCCCAA 50

TGGGCTGAGT ACTTGAGCAA AATCACCTGC AGGGG 85

(2) INFORMATION FOR SEQ ID NO: 95:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

ATCCGCCTGA TTAGCGATAC TCCAGGCAGT CATGGTCATT GTTTACAGTC 50

GTGGAGTAGG TACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 96:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 85 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:

ATCCGCCTGA TTAGCGATAC TAGGTGATCC CTGCAAAGGC AGGATAACGT 50

CCTGAGCATC ACTTGAGCAA AATCACCTGC AGGGG 85

(2) INFORMATION FOR SEQ ID NO: 97:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 83 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

ATCCGCCTGA TTAGCGATAC TATGTGATCC CTGCAGAGGG AGGANACGTC 50

TGAGCATCAC TTGAGCAAAA TCACCTGCAG GGG 83

-continued

(2) INFORMATION FOR SEQ ID NO: 98:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 86 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

ATCCGCCTGA TTAGCGATAC TCACGTGATC CCATAAGGGC TCGCAAAAT 50
 AGCAGAGCAT CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 99:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 86 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

ATCCGCCTGA TTAGCGATAC TGGTGGACTA GAGGGCAGCA AACGATCCTT 50
 GGTTAGCGTC CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 100:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 85 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

ATCCGCCTGA TTAGCGATAC TGGTGCACG AGGCTTACAC AAACGTACAC 50
 GTTTCCCCG ACTTGAGCAA AATCACCTGC AGGGG 85

(2) INFORMATION FOR SEQ ID NO: 101:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 86 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

ATCCGCCTGA TTAGCGATAC TTGTCGGAGC AGGGGCGTAC GAAACTTTA 50
 CAGTTCCCC GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 102:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 86 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

ATCCGCCTGA TTAGCGATAC TAGTGGAACA GGGCACGGAG AGTCAAACCTT 50

TGGTTTCCCC CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 103:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

ATCCGCCTGA TTAGCGATAC TGTGGGTAGG GATCGGTGGA TGCCTCGTCA 50

CTTCTAGTCC CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 104:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

ATCCGCCTGA TTAGCGATAC TGGGCGCCCT AAACAAGGG TGGTCACTTC 50

TAGTCCCAGG AACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 105:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 87 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:

ATCCGCCTGA TTAGCGATAC TTCCGGGCTC GGGATTCGTG GTCACCTTCA 50

GTCCCGGATA TAACTTGAGC AAAATCACCT GCAGGGG 87

(2) INFORMATION FOR SEQ ID NO: 106:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 84 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:

ATCCGCCTGA TTAGCGATAC TATGGGAGGG CGCGTTCTTC GTGGTTACTT 50

TTAGTCCCGA CTTGAGCAA ATCACCTGCA GGGG 84

-continued

(2) INFORMATION FOR SEQ ID NO: 107:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 84 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

ATCCGCCTGA TTAGCGATAC TACGGGAGGG CACGTTCTTC GTGGTTACTT 50

TTAGTCCCGA CTTGAGCAAA ATCACCTGCA GGGG 84

(2) INFORMATION FOR SEQ ID NO: 108:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

ATCCGCCTGA TTAGCGATAC TGCTCGTAGG GGGCGATTCT TTCGCCGTTA 50

CTTCCAGTCC TACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 109:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:

ATCCGCCTGA TTAGCGATAC TGAGGCATGT TAACATGAGC ATCGTCTCAC 50

GATCCTCAGC CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 110:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

ATCCGCCTGA TTAGCGATAC TCCACAGGCT ACGGCACGTA GAGCATCACC 50

ATGATCCTGT GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 111:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

ATCCGCCTGA TTAGCGATAC TCGGGCATG GCACATGAGC ATCTCTGATC 50

CCGCAATCCT CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 112:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

ATCCGCCTGA TTAGCGATAC TACCGGGCTA CTTCTAGAG CATCTCTGAT 50

CCCGGTGCTC GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 113:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

ATCCGCCTGA TTAGCGATAC TAAAGGGCGA ACGTAGGTCG AGGCATCCAT 50

TGGATCCCTT CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 114:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 86 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

ATCCGCCTGA TTAGCGATAC TACGGGCTCT GTCCTGTGG CACTAGCAAT 50

AGTCCCGTCG CACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 115:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 85 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

ATCCGCCTGA TTAGCGATAC TGGGCAGACC TTCTGGACGA GCATCACCTA 50

TGTGATCCCG ACTTGAGCAA AATCACCTGC AGGGG 85

-continued

(2) INFORMATION FOR SEQ ID NO: 116:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

ATCCGCCTGA TTAGCGATAC TAGAGGGGAA GTAGGCTGCC TGACTCGAGA 50
GAGTCCTCCC GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 117:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

ATCCGCCTGA TTAGCGATAC TAGGGGTGCG AAACACATAA TCCTCGCGGA 50
TTCCCATCGC TACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 118:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 83 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

ATCCGCCTGA TTAGCGATAC TGGGGGGGCA ATGGCGGTAC CTCTGGTCCC 50
CTAAATACAC TTGAGCAAAA TCACCTGCAG GGG 83

(2) INFORMATION FOR SEQ ID NO: 119:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 85 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

ATCCGCCTGA TTAGCGATAC TGCGGCTCAA AGTCCTGCTA CCCGCAGCAC 50
ATCTGTGGTC ACTTGAGCAA AATCACCTGC AGGGG 85

(2) INFORMATION FOR SEQ ID NO: 120:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

ATCCGCCTGA TTAGCGATAC TTTGGGCGTG AATGTCCACG GGTACCTCG 50

GTCCCAAAGA GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 121:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 85 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

ATCCGCCTGA TTAGCGATAC TTCCGCGCAA GTCCTGGTA AAGGCAGCC 50

CTAACTGGTC ACTTGAGCAA AATCACCTGC AGGGG 85

(2) INFORMATION FOR SEQ ID NO: 122:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

ATCCGCCTGA TTAGCGATAC TCAAGTCCC CACAAGACTG GGGCTGTCA 50

AACCGCTAGT AACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 123:

ATCCGCCTGA TTAGCGATAC TCAAGTAGGG CGGACACAC GTCGGGCAC 50

CTAAGGTCCC AACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 124:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 86 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 124:

ATCCGCCTGA TTAGCGATAC TAAAGTCGTG CAGGTCCCC TGAAGCATC 50

-continued

TCCGATCCCA GACTTGAGCA AAATCACCTG CAGGGG 86

(2) INFORMATION FOR SEQ ID NO: 125:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 96 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 125:

GGGAGACAAG AAUAACGCUC AANNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNNNNNN NNNNNNNNNN NNUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 126:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 126:

TAATACGACT CACTATAGGG AGACAAGAAT AACGCTCAA 39

(2) INFORMATION FOR SEQ ID NO: 127:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 127:

GCCTGTGTG AGCCTCCTGT CGAA 24

(2) INFORMATION FOR SEQ ID NO: 128:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 93 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:

(D) OTHER INFORMATION: All pyrimidines are 2'-F mod

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 128:

GGGAGACAAG AAUAACGCUC AACGGUGGCA UUUCUUCACU UCCUUCUGC 50

UUUCUCGCGU UGGGCNCGAU UCGACAGGAG GCUCACAACA GGC 93

(2) INFORMATION FOR SEQ ID NO: 129:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 91 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 129:

GGGAGACAAG AAUAACGCUC AACCAACCUU CUGUCGGCGU UGCUUUUUGG 50

ACGGCACUCA GGCUCCAUUC GACAGGAGGC UCACAACAGG C 91

(2) INFORMATION FOR SEQ ID NO:130:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 95 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F mod

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:130::

GGGAGACAAG AAUAACGCUC AAUCGAUCGG UUGUGUGCCG GACAGCCUUA 50

ACCAGGGCUG GGACCGAGGC CUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 131:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 92 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F mod

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 131:

GGGAGACAAG AAUAACGCUC AACUGAGUAG GGGAGGAAGU UGAAUCAGUU 50

GUGGCGCCUC UCAUUCGCUU CGACAGGAGG CUCACAACAG GC 92

(2) INFORMATION FOR SEQ ID NO: 132:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 95 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 132:

GGGAGACAAG AAUAACGCUC AACAGCACUU UCGCUUUUCA UCAUUUUUUC 50

UUUCCACUGU UGGGCGCGGA AUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 133:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 133:

GGGAGACAAG AAUAACGCUC AAUCAGUGCU GCGUCAUGU CUCGAUGGGG 50
 AUUUUUUCUUC AGCACUUUGC CAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 134:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 134:

GGGAGACAAG AAUAACGCUC AAUCUACUUU CCAUUUCUCU UUCUUCUCA 50
 CGAGCGGGUU UCCAGUGAAC CAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 135:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 94 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 135:

GGGAGACAAG AAUAACGCUC AACGAUAGUG ACUACGAUGA CGAAGGCCGC 50
 GGGUUGGAUG CCCGCAUUGA UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 136:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 93 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 136:

GGGAGACAAG AAUAACGCUC AAGUCGAUAC UGGCGACUUG CUCCAUUGGC 50
 CGAUUAACGA UUCGGUCAGU UCGACAGGAG GCUCACAACA GGC 93

-continued

(2) INFORMATION FOR SEQ ID NO: 137:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 95 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 137:

GGGAGACAAG AAUAACGCUC AAGUGCAAAC UUAACCCGGG AACCGCGCGU 50

UUCGAUCGAC UUUCCUUUC AUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 138:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 138:

GGGAGACAAG AAUAACGCUC AAAUCCGCG UCCGAUUAU UCCUGUGCUC 50

GGAAAUCGGU AGCCAUAGUG CAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 139:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 139:

GGGAGACAAG AAUAACGCUC AACGAACGAG GAGGGAGUGG CAAGGGAUGG 50

UUGGAUAGGC UCUACGCUCA UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 140:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 140:

-continued

GGGAGACAAG AAUAACGCUC AAGCGAAACU GCGACUUGC UCCAUGGCC 50

GAUUAACGA UUCGGUUCAU UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 141:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 95 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 141:

GGGAGACAAG AAUAACGCUC AACGAACGAG GAGGGAGUCG CAAGGGAUGG 50

UUGGAUAGGC UCUACGCUCA AUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 142:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 142:

GGGAGACAAG AAUAACGCUC AACGAGAAGU GACUACGAUG ACGAAGGCCG 50

CGGGUUGAAU CCCUCAUGA UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 143:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 95 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 143:

GGGAGACAAG AAUAACGCUC AAAAGCAACG AGACCUGACG CCUGAUGUGA 50

CUGUGCUUGC ACCCGAUUCU GUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 144:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 144:
GGGAGACAAG AAUAACGCUC AAGUGAUUCU CAUUCUCAAU GCUUUCUCAC 50
AACUUUUUCC ACUUCAGCGU GAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 145:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 145:
GGGAGACAAG AAUAACGCUC AAAAGCAACG AGACUCGACG CCUGAUGUGA 50
CUGUGCUUGC ACCCGAUUCU UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 146:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 146:
GGGAGACAAG AAUAACGCUC AAUCGAUCGG UUGUGUGCCG GACAGCUUUG 50
ACCAUGAGCU GGGACCGAGG CCUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 147:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 147:
GGGAGACAAG AAUAACGCUC AANGACGNGU GGACCUGACU AAUCGACUGA 50
UCAAGAUAUC CGCCCAUG GGUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 148:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 148:

GGGAGACAAG AAUAACGCUC AACACUGCGA CUUGCAGAAG CCUUGUGUGG 50

CGGUACCCCC UUUGGCCUCG UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 149:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 149:

GGGAGACAAG AAUAACGCUC AAGGUGGCAU UUCUUCAUUU UCCUUCUCGC 50

UUUCUCGCGC GUUGGGCGCG UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 150:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 150:

GGGAGACAAG AAUAACGCUC AACCUGAGUA GGGGGGAAAG UUGAAUCAGU 50

UGUGGCGCUC UACUCAUUCG CCUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 151:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 151:

GGGAGACAAG AAUAACGCUC AAGUCGAAAC UGGCGACUUG CUCCAUUGGC 50

CGAUUAUACG AUUCGGUUA UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 152:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 94 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 152:

GGGAGACAAG AAUAACGCUC AAGCGAUACU GCGGACUUGC UCCAUUGGCC 50
 GAUAUAACGA UUCGGCUCAG UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 153:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 153:

GGGAGACAAG AAUAACGCUC AAACGUGGGG CACAGGACCG AGAGUCCUC 50
 CGGCAAUAGC CGCUACCCCA CCUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 154:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 98 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 154:

GGGAGACAAG AAUAACGCUC AACACAGCCU NANAGGGGGG AAGUUGAAUC 50
 AGUUGUGGCG CUCUACUCAU UCGCUUCGAC AGGAGGCUCA CAACAGGC 98

(2) INFORMATION FOR SEQ ID NO: 155:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 94 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 155:

GGGAGACAAG AAUAACGCUC AAANGGGNUA UGGUGACUUG CUCCAUUGGC 50
 CGAUAUAACG AUUCGGUCAG UUCGACAGGA GGCUCACAAC AGGC 94

-continued

(2) INFORMATION FOR SEQ ID NO: 156:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 156:

GGGAGACAAG AAUAACGCUC AACCGCGUA GGGNGGGAAG UUGAAUCAGU 50

UGUGGCGCUC UACUCAUUCG CCUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 157:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 157:

GGGAGACAAG AAUAACGCUC AACGAACGAG GAGGGAGUGG CAAGGGAUGG 50

UUGGAUAGGC UCUACGCUCA UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 158:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 97 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 158:

GGGAGACAAG AAUAACGCUC AAGUGCAAAC UUAACCCGGG AACCGCGGU 50

UUCGAUUCGC UUUCCNUAUU CCAUUCGACA GGAGGCUCAC AACAGGC 97

(2) INFORMATION FOR SEQ ID NO: 159:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 93 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

-continued

GGGAGACAAG AAUAACGCUC AACGAACGAG GAGGGAGUGG CAAGGGACGG 50

UNNAUAGGCU CUACGCUCAU UCGACAGGAG GCUCACAACA GGC 93

(2) INFORMATION FOR SEQ ID NO: 160:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 93 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 160:

GGGAGACAAG AAUAACGCUC AAUCGGUGUG GCUCAGAAAC UGACACGCGU 50

GAGCUUCGCA CACAUCUGCU UCGACAGGAG GCUCACAACA GGC 93

(2) INFORMATION FOR SEQ ID NO: 161:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 95 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 161:

GGGAGACAAG AAUAACGCUC AAUAUCGCUU UUCAUCAAU CCACUUUUUC 50

ACUCUNUAAC UUGGGCGUGC AUUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 162:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 96 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 162:

GGGAGACAAG AAUAACGCUC AAGUGCAAAC UUAACCCGGG AACCGCGCGU 50

UUGGAUCCUG CAUCCUUUUU CCUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 163:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 93 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 163:

GGGAGACAAG AAUAACGCUC AAUCGNUCGG UUGUGUGCCG GCAGCUUUGU 50

CCAGCGUUGG GCCGAGGCCU UCGACAGGAG GCUCACAACA GGC 93

(2) INFORMATION FOR SEQ ID NO: 164:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 95 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 164:

GGGAGACAAG AAUAACGCUC AAAGUACCCA UCUCAUCUUU UCCUUUCCUU 50

UCUUC AAGGC ACAUUGAGGG UUCGACAGG AGGCUCACAA CAGGC 95

(2) INFORMATION FOR SEQ ID NO: 165:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 97 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 165:

GGGAGACAAG AAUAACGCUC AACCGAGUA GGGGGGAAG UGAACCAGU 50

UGUGGCNGCC UACUCAUUCN CCAUUCGACA GGAGGCUCAC AACAGGC 97

(2) INFORMATION FOR SEQ ID NO: 166:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 94 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 166:

GGGAGACAAG AAUAACGCUC AACCNCCUN CUGUCGGCGC UUGUCUUUUU 50

GGACGGGCAA CCCAGGCUC UUCGACAGGA GGCUCACAAC AGGC 94

(2) INFORMATION FOR SEQ ID NO: 167:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 96 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 167:

GGGAGACAAG AAUAACGCUC AACCAACCUN CUGUCGGCGC UUGUCUUUUU 50

GGACGAGCAA CUCAAGGCUC GUUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 168:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 93 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 168:

GGGAGACAAG AAUAACGCUC AACCAGCGCA GAUCCCGGC UGAAGUGACU 50

GCCGGCAACG GCCGCUCCAU UCGACAGGAG GCUCACAACA GGC 93

(2) INFORMATION FOR SEQ ID NO: 169:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 169:

GGGAGACAAG AAUAACGCUC AAUCCCGUA ACAACUUUUC AUUUUCACUU 50

UUCAUCCAAC CAGUGAGCAG CAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 170:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-F modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 170:

GGGAGACAAG AAUAACGCUC AAUAUCGCUU UCAUCAAAUU CCACUCCUUC 50

ACUUCUUUAA CUUGGGCGUG CAUUCGACAG GAGGCUCACA ACAGGC 96

(2) INFORMATION FOR SEQ ID NO: 171:

-continued

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: N at positions 1 and 23
base pair to each other and can be 1-4
base pairs long.
- (ix) FEATURE:
(D) OTHER INFORMATION: N at positions 5 and 10
is any base pair.
- (ix) FEATURE:
(D) OTHER INFORMATION: N at positions 6 and 9
is any base pair.
- (ix) FEATURE:
(D) OTHER INFORMATION: N at positions 7 and 8 is
any base pair.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 171:
NGGCNNNNNN GRKYAYRRT CCN 23
- (2) INFORMATION FOR SEQ ID NO: 172:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: Nucleotide 38 is an
inverted orientation T (3'-3' linked)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 172:
TGGGAGGGCG CGTCTTCGT GGTTACTTTT AGTCCCGT 38
- (2) INFORMATION FOR SEQ ID NO: 173:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: Nucleotide 40 is an
inverted orientation T (3'-3' linked)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 173:
CACAGGCTAC GGCACGTAGA GCATCACCAT GATCCTGTGT 40
- (2) INFORMATION FOR SEQ ID NO: 174:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(D) OTHER INFORMATION: Nucleotide 45 is an inverted orientation T (3'-3' linked)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 174:

TACTCAGGGC ACTGCAAGCA ATTGTGGTCC CAATGGGCTG AGTAT

45

(2) INFORMATION FOR SEQ ID NO: 175:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(D) OTHER INFORMATION: C at positions 8, 11, 25 and 26 is 2'-O-methyl-2'-deoxycytidine.

(ix) FEATURE:

(D) OTHER INFORMATION: G at positions 9, 10, 17, 19 and 35 is 2'-O-methyl-2'-deoxyguanosine.

(ix) FEATURE:

(D) OTHER INFORMATION: A at positions 12, 24 and 27 is 2'-O-methyl-2'-deoxyadenosine.

(ix) FEATURE:

(D) OTHER INFORMATION: U at position 34 is 2'-O-methyl-2'-deoxyuridine.

(ix) FEATURE:

(D) OTHER INFORMATION: U at positions 6 and 22 is 2'-fluoro-2'-deoxyuridine.

(ix) FEATURE:

(D) OTHER INFORMATION: C at positions 23, 32 and 33 is 2'-fluoro-2'-deoxycytidine.

(ix) FEATURE:

(D) OTHER INFORMATION: Nucleotide 36 is an inverted orientation T (3'-3' linked).

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 175:

CAGGCUACGG CACGTAGAGC AUCACCATGA TCCUGT

36

(2) INFORMATION FOR SEQ ID NO: 176:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(D) OTHER INFORMATION: C at position 8 is 2'-O-methyl-2'-deoxycytidine.

(ix) FEATURE:

(D) OTHER INFORMATION: G at positions 9, 17 and 31 is 2'-O-methyl-2'-deoxyguanosine.

(ix) FEATURE:

(D) OTHER INFORMATION: A at position 22 is 2'-O-methyl-2'-deoxyadenine.

-continued

- (ix) FEATURE:
(D) OTHER INFORMATION: U at position 30 is 2'-
0-methyl-2'-deoxyuridine.
- (ix) FEATURE:
(D) OTHER INFORMATION: U at positions 6 and
20 is 2'-fluoro-2'-deoxyuridine.
- (ix) FEATURE:
(D) OTHER INFORMATION: C at positions 21, 28
and 29 is 2'-fluoro-2'-deoxycytidine.
- (ix) FEATURE:
(D) OTHER INFORMATION: N at positions 10 and 23
is pentaethylene glycol phosphoramidite
spacer.
- (ix) FEATURE:
(D) OTHER INFORMATION: Nucleotide 32 is an
inverted orientation T (3'-3' linked)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 176:
CAGGCUACGN CGTAGAGCAU CANTGATCCU GT 32
- (2) INFORMATION FOR SEQ ID NO: 177:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: Nucleotide 39 is an
inverted orientation T (3'-3' linked)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 177:
CAGTCCGTGG TAGGGCAGGT TGGGGTGACT TCGTGGAAT 39
- (2) INFORMATION FOR SEQ ID NO: 178:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: T at positions 13, 14,
16 and 17 is substituted with IdU.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 178:
TGGGAGGGCG CGTCTTCGT GGTTACTTTT AGTCCCG 37
- (2) INFORMATION FOR SEQ ID NO: 179:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (ix) FEATURE:
(D) OTHER INFORMATION: T at position 20 is substituted with

-continued

IdU.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 179:

TGGGAGGGCG CGTCTTCGT GGTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 180:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(D) OTHER INFORMATION: T at position 23 is substitu

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 180:

TGGGAGGGCG CGTCTTCGT GGTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 181:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(D) OTHER INFORMATION: T at position 24 is substitu

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 181:

TGGGAGGGCG CGTCTTCGT GGTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 182:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(D) OTHER INFORMATION: T at position 27 is substitu

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 182:

TGGGAGGGCG CGTCTTCGT GGTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 183:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(D) OTHER INFORMATION: T at positions 28-30 is subs

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 183:

-continued

 TGGGAGGGCG CGTTCCTTCGT GGTTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 184:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 37 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 - (D) OTHER INFORMATION: T at position 33 is substituted

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 184:

TGGGAGGGCG CGTTCCTTCGT GGTTACTTTT AGTCCCG 37

(2) INFORMATION FOR SEQ ID NO: 185:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 7 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Peptide

- (ix) FEATURE:
 - (D) OTHER INFORMATION: Xaa at position 5 is a modification that could not be identified.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 185:

Lys Lys Pro Ile Xaa Lys Lys
5

(2) INFORMATION FOR SEQ ID NO: 186:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 71 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 186:

GGGAGGACGA UGCGGNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 50

NNNNNCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 187:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 32 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 187:

TAATACGACT CACTATAGGG AGGACGATGC GG 32

(2) INFORMATION FOR SEQ ID NO: 188:

- (i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 188:

TCGGGCGAGT CGTCTG 16

(2) INFORMATION FOR SEQ ID NO: 189:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 189:

GGGAGGACGA UGCGGGAAG GACGAUAAAG AGGAAUCGAA CAACAAGUGG 50
CUGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 190:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 190:

GGGAGGACGA UGCGGGCGGG AAGGUCCGAA GACCGCGAA AGGAACGAGA 50
UUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO:191:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:191:

GGGAGGACGA UGCGGGUGGU GAAGAGGUAC CGGAAUUGCU AAAGAUACCA 50
GGCCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 192:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs

-continued

(B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:192:

GGGAGGACGA UGCGGGCAGG GAGCAAUGAA CUCAAGUCA GCGGUGCAC 50
 GUGGGCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO:193:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:193:

GGGAGGACGA UGCGGUAGCU GCUGUCAUGC AAGACACUAG AAGAUUAAGA 50
 UGGGGCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO:194:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 74 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:194:

GGGAGGACGA UGCGGGGGCC GGAUUUGAAC CGACGACUUC GGUUAUGAG 50
 CCCGACGUCA GACGACUCGC CCGA 74

(2) INFORMATION FOR SEQ ID NO: 195:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 195:

GGGAGGACGA UGCGGUCCAG GGAUUUGAAGU GUCGGGUAG GAACAUAAAG 50
 GCGGCCAGAC GACUCGCCG A 71

-continued

(2) INFORMATION FOR SEQ ID NO: 196:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 196:

GGGAGGACGA UGCGGAAGUU CUAACAAGUU AGUGGAAGGU UCCACUUGAA 50

UGUACAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 197:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 197:

GGGAGGACGA UGCGGAUGGA GCUGAAAUCA GACGACUCGC CCGA 44

(2) INFORMATION FOR SEQ ID NO: 198:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 198:

GGGAGGACGA UGCGGGUGGG AAGAUGAGCC GGUCGGCAGU AAUGUGACAC 50

UGCGGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 199:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 199:

GGGAGGACGA UGCGGGAGGG AAUGAGGAAA CAACUAGCAG AUAACCGAGC 50

UGGCCAGACG ACUCGCCCGA 70

-continued

(2) INFORMATION FOR SEQ ID NO: 200:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 200:

GGGAGGACGA UGCGGAUGGA GCUGAAAUCA GACGACUCGC CCGA 44

(2) INFORMATION FOR SEQ ID NO: 201:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 201:

GGGAGGACGA UGCGGUUGCU CUACAAUGAC GCGGUGACUC CGCAGUUCUU 50

GGACACAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 202:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 202:

GGGAGGACGA UGCGGAGGG GAGAAGAAUG CAGGAAACAG CGAAAUGCGU 50

GUGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 203:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 203:

GGGAGGACGA UGCGGCGGG AAGAGCUAAU GGAAGUGGAA UCAGUCACAG 50

-continued

UGCGGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 204:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 204:

GGGAGGACGA UGCGGGCUUA GGGAAAUGGU UCUGAGGUGG UCAGACGACU 50

GC CCCGA 57

(2) INFORMATION FOR SEQ ID NO: 205:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 205:

GGGAGGACGA UGCGGGAAGG GAACAGGAUA AGACAAGUCG AACAAAGCCG 50

AGGUGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 206:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 206:

GGGAGGACGA UGCGGAUGGA GCUGAAAUCA GACGACUCGC CCGA 44

(2) INFORMATION FOR SEQ ID NO: 207:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 207:

-continued

GGGAGGACGA UGCGGGGAGA CGUAGACGGG AACAUAGAAC GAACAUCAAC 50

GCGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 208:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 69 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH₂ modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:208:

GGGAGGACGA UGCGGGAAGU GGAUAGAACA GUCAGAAAUG UAAGCGUGAG 50

GUGCAGACGA CUCGCCCGA 69

(2) INFORMATION FOR SEQ ID NO: 209:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH₂ modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 209:

GGGAGGACGA UGCGGGAAGG GUAGGAAGGU CAAGAGGAAA CAGCGCUUCG 50

GGGUGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 210:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH₂ modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 210:

GGGAGGACGA UGCGGGGCAA AGGAAGUUGG AAUCGGGACU AAGUAGUGUG 50

UGGCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 211:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 211:
GGGAGGACGA UGCGGAGAAC CAACAGAGCC CCCUGGUGGU GGGGGAAGGA 50
UUCUCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 212:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 212:
GGGAGGACGA UGCGGACACA CAAGUGAAGG UCAGACGCGA AUUACGUGGG 50
UGGCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 213:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 82 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 213:
GGGAGGACGA UGCGGUCGUG GGGUGGGUGG GGCAGCGUU GAAUAAGUA 50
ACUGGUAACG GCUGGCCAGA CGACUCGCCG GA 82

(2) INFORMATION FOR SEQ ID NO: 214:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 214:
GGGAGGACGA UGCGGGGUGG GUGGUUACCU GUAAUUAUUAU UGAUUCUGGC 50
UUUAGCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO: 215:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 53 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 215:

GGGAGGACGA UGCGGCCCCU UAGCUCAGUG GUUAGAGCAG ACGACUCGCC 50

CGA 53

(2) INFORMATION FOR SEQ ID NO: 216:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 74 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 216:

GGGAGGACGA UGCGGUAACG UGGAUAGGG UAAACAGCU GGAAUAACG 50

UAGGUGGCCA GACGACUCGC CCGA 74

(2) INFORMATION FOR SEQ ID NO: 217:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 217:

GGGAGGACGA UGCGGUAGG GAGUAGGACA GACUAACAG UGCAACCAUC 50

GUGGCCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO: 218:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 73 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 218:

GGGAGGACGA UGCGGAAACG GCGUGGCAAA AGUGAGGGGG UAGGAUGUAC 50

CAUGGGUCAG ACGACUCGCC CGA 73

(2) INFORMATION FOR SEQ ID NO: 219:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH₂ modified

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 219:

GGGAGGACGA UGCGGAGGG GAAAUGAGA CCGACAGAUU GACGGAAGUA 50
 CUGGGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO:220:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 220:

GGGAGGACGA UGCGGGCAUU CGUCAAUACC UGUUUUUUUU CCUUUUCUAG 50
 CGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 221:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 221:

GGGAGGACGA UGCGGAUCGU AAUCGCCACU ACUACUUUCC GAACCCGCAC 50
 GUGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 222:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 222:

GGGAGGACGA UGCGGCGUCC CGAGUCACGC UGUCCUGAUA ACCUUCUCUG 50
 UGCCAGACG ACUCGCCCGA 70

-continued

(2) INFORMATION FOR SEQ ID NO:223:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 223:

GGGAGGACGA UGCGGGAUCC UUUGUGGGCU CUUGUUGACC CCCUCGUUGU 50

CCCCCCCAGA CGACUCGCC GA 72

(2) INFORMATION FOR SEQ ID NO: 224:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 224:

GGGAGGACGA UGCGCGGGU ACUCUUCGCC AGCUCCUCCA AGCGCGACCU 50

GUGCCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO: 225:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 225:

GGGAGGACGA UGCGGUUUCG AAUAGGGCCA UUUCACUA GCUAUCCUAC 50

CCUGCCCAGA CGACUCGCC GA 72

(2) INFORMATION FOR SEQ ID NO: 226:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 226:

-continued

GGGAGGACGA UGCGGAUAAU GGCUAGAACU AGCUCGCAUC UUGGUGUCCG 50

GUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 227:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 227:

GGGAGGACGA UGCGGGACCA GAUGGCGGAU UUUUCAGCAA UCCUCCCCCG 50

CUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 228:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 228:

GGGAGGACGA UGCGGUGAUG GCGACCAGUC AAACCGGUGC UUUUACUCCC 50

CCGCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 229:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 72 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 229:

GGGAGGACGA UGCGGAAAU AACAGGGCCA GAAUUCUCAU CUNNCUCCCC 50

GUGACCCAGA CGACUCGCC GA 72

(2) INFORMATION FOR SEQ ID NO: 230:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 230:
GGGAGGACGA UGCGGCACCU UAGACCUGUC CUCCAAGCGU GAGUUGCUGU 50
GGCCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 231:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: RNA
(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 231:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AAACUUUCUC CAUCGUAUCU 50
GGGCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 232:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: RNA
(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:232:
GGGAGGACGA UGCGGUCAUG GUGUCUUUCC ACAGCUCUUC CCAUGAUCGC 50
CCGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 233:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: RNA
(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 233:
GGGAGGACGA UGCGGGAAUU CCCAGCGCUU GACUGAUACA AACNUUCCCG 50
UGCCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO:234:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 234:

GGGAGGACGA UGCGGCAANN NNNNNCUCUC UCCUGGCGUU CCGCAACCCG 50

CCCCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 235:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 235:

GGGAGGACGA UGCGGAGUUAU UCCAGCCUGG AUUCAUAGUC AGUGCUCUCC 50

GUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 236:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 236:

GGGAGGACGA UGCGGUCCUA GCAGCGAUUC AUCCCCGUUC UCUCAGCGUU 50

GCCCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 237:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:237:

GGGAGGACGA UGCGGCCUGA AGUACAGGCU CUA AACUCCA AGCGCGACCG 50

UCCGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 238:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 68 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 238:

GGGAGGACGA UGCGGCCUA CCACUUUUUC CCUCUACUGU UAUCCUGUCC 50
 CCCAGACGAC UCGCCCGA 68

(2) INFORMATION FOR SEQ ID NO: 239:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 239:

GGGAGGACGA UGCGGUGGUC UCCCUAGAUC UACAGCACUU CCAUCGCAUU 50
 GGGCCAGACG ACUCGCCCGA 70

(2) INFORMATION FOR SEQ ID NO: 240:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 69 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 240:

GGGAGGACGA UGCGGUCAAG CUUAAACAGUC UGGCAAUGGC CAUUAUGGCG 50
 CCCCAGACGA CUCGCCCGA 69

(2) INFORMATION FOR SEQ ID NO: 241:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 72 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 241:

GGGAGGACGA UGCGGCAGUC UGGAUCUCUA UUGGAAUUUA GUCCUCAACU 50
 GUGCCCCAGA CGACUCGCC GA 72

-continued

(2) INFORMATION FOR SEQ ID NO: 242:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 242:

GGGAGGACGA UGCGGGAUUC UUUCGGCAAG UGAAAAUAU CCUUGCUUCC 50

CGAGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 243:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 243:

GGGAGGACGA UGCGGGGACU UCAACUAAGU CCUCAUUUGC CUCGCUCCUC 50

GUGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 244:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 74 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 244:

GGGAGGACGA UGCGGAACGG AGAUGUCCCC UCAAMAUUUA CCGUCUCCGU 50

UUGCGCCCA GACGACUCGC CCGA 74

(2) INFORMATION FOR SEQ ID NO: 245:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:245:

-continued

GGGAGGACGA UGCGGCGAAA UUAGCUUCUU AUGACUCACG UUCCUUGCC 50

GCCCCAGACG ACUCGCCCCG 70

(2) INFORMATION FOR SEQ ID NO: 246:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 246:

GGGAGGACGA UGCGGGCCCG AUCUACUGCA UUACCGAAAC GAUUUCCCCA 50

CUGUGCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 247:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 247:

GGGAGGACGA UGCGGNGACU GAUUUUUCCU UGNCAGUGUA AUUCCUGGC 50

UGCCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 248:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 248:

GGGAGGACGA UGCGGGGACU UUGACAGGCA UGAUUUCGA CCUGUCCCC 50

GUGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 249:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 249:
GGGAGGACGA UGCGGCGACA CAUAGCCUU UGAUCCCAUG AUGGCUCGCC 50
GUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 250:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 250:
GGGAGGACGA UGCGGUGUAG UUUCCUGUA UGCCAUUCUU UCCCAUGCCG 50
CACGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 251:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 251:
GGGAGGACGA UGCGGUCGAG UGUUCUCCUU CGGUAACUUA UNNNAUUUC 50
GUGCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 252:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 252:
GGGAGGACGA UGCGGGUCGU AUUCAUCC UGUUCUGUU UCGUUGCACC 50
UGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 253:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 253:

GGGAGGACGA UGCGGGGACU UUGACAGGCA UGAUUUCGA CGUGUCCCC 50

GUGGCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 254:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 69 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 254:

GGGAGGACGA UGCGGUGAUC AAUCGGCGCU UUACUCUUGC GCUCACCGUG 50

CCCCAGACGA CUCGCCCGA 69

(2) INFORMATION FOR SEQ ID NO: 255:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 255:

GGGAGGACGA UGCGGCAGUC UCCCUAGGUU UCAUCUCUGC AGCAUCCGG 50

GGUNCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 256:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 256:

GGGAGGACGA UGCGGAUCA AAGCACUCAU UCCCGUGCUC GCUUCAUUG 50

UCCCCAGAC GACUCGCCCG A 71

(2) INFORMATION FOR SEQ ID NO: 257:

-continued

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 74 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 257:

GGGAGGACGA UGCGGAAGAU CUCCCAACUG CUGUGGCUAA UAAUUCUCUC 50
CGCGUCCCA GACGACUCGC CCGA 74

(2) INFORMATION FOR SEQ ID NO: 258:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 70 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 258:

GGGAGGACGA UGCGGUCCGU CAUAACGGCC AUAACUGCG AAUACUCCU 50
GGCCAGACG ACUCGCCGA 70

(2) INFORMATION FOR SEQ ID NO: 259:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 73 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 259:

GGGAGGACGA UGCGGGGACA AWYAGCGUG UCUUUCAU UNKAUCCUC 50
GACRUCCAG ACGACUGCC CGA 73

(2) INFORMATION FOR SEQ ID NO: 260:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 260:

GGGAGGACGA UGCGGUGACU AUCUGGUCG AUCCAAUCAC CCGAGCCCAC 50
CGCGCCAGAC GACUCGCCG A 71

-continued

(2) INFORMATION FOR SEQ ID NO: 261:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 261:

GGGAGGACGA UGCGGAACU AAUGGCCGUG AUUAACCAU GCAGGCUUCC 50

UGCGCCAGAC GACUCGCCG A 71

(2) INFORMATION FOR SEQ ID NO: 262:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 73 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 262:

GGGAGGACGA UGCGGUGACA UGGAAUUUUC UACGGGCCCG AUCCUGCCAG 50

CCGUGUGCAG ACGACUGCC CGA 73

(2) INFORMATION FOR SEQ ID NO: 263:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 m

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 263:

GGAANYAANR CRR 13

(2) INFORMATION FOR SEQ ID NO: 264:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-NH2 m

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 264:

GGAANYTGA 9

-continued

(2) INFORMATION FOR SEQ ID NO: 265:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 76 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All C's are 2'-NH₂ 2 cytosine

(ix) FEATURE:
 (D) OTHER INFORMATION: All U's are 2'-NH₂ 2 uracil

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 265:

GGGAGACAAG AAUAACGCUC AAGUAGACUA AUGUGUGGAA GACAGCGGGU 50

GGUUCGACAG GAGGCUCACA ACAGGC 76

(2) INFORMATION FOR SEQ ID NO: 266:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All C's are 2'-NH₂ 2 cytosine

(ix) FEATURE:
 (D) OTHER INFORMATION: All U's are 2'-NH₂

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 266:

GGUGUGUGGA AGACAGCGGG UGGUUC 26

(2) INFORMATION FOR SEQ ID NO: 267:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
 (D) OTHER INFORMATION: All U's and C's are 2'- deox
 and 2'-deoxy-2' aminocytidine residues

(ix) FEATURE:
 (D) OTHER INFORMATION: A at positions 14 and 17 is
 methyladenosine

(ix) FEATURE:
 (D) OTHER INFORMATION: G at positions 5, 6, 22, 23,
 2'-deoxy-2'-O-methylguanosine

(ix) FEATURE:
 (D) OTHER INFORMATION: T at positions 1-4 and 31-35
 deoxythymidine

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:267:

TTTTGGUGUG UGGAAGACAG CGGGUGGUUC TTTT 35

-continued

(2) INFORMATION FOR SEQ ID NO: 268:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 268:

GGGAGGSGSG GUYUCYRRY UYYYSYS

27

(2) INFORMATION FOR SEQ ID NO: 269:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 269:

RGGRRGRAYY RSBSYYYYY BSYBSY

26

(2) INFORMATION FOR SEQ ID NO: 270:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 m

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 270:

GGGAAGGAAY AARCRGACC C

21

(2) INFORMATION FOR SEQ ID NO: 271:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-NH2 m

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 271:

GGAAYGAYGA Y

11

(2) INFORMATION FOR SEQ ID NO: 272:

(i) SEQUENCE CHARACTERISTICS:

-continued

(A) LENGTH: 49 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 272:

GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AAACUUUCUC CAUCGUAUC 49

(2) INFORMATION FOR SEQ ID NO: 273:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 273:

GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AAACUUUCUC CAUCGUA 47

(2) INFORMATION FOR SEQ ID NO: 274:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 274:

GGGAGGACUG CGGUGGUCUC CCAAUUCUAA ACUUUCUCCA UCGUA 45

(2) INFORMATION FOR SEQ ID NO: 275:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 275:

GGGAGGACGA UGCGGUGGUC UCCCCUAAAC UUUCUCCAUC GUAUC 45

(2) INFORMATION FOR SEQ ID NO: 276:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-continued

- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 276:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AUUCUCCAUC GUAUC 45
- (2) INFORMATION FOR SEQ ID NO: 277:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 45 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 277:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AAACUCCAUC GUAUC 45
- (2) INFORMATION FOR SEQ ID NO: 278:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 278:
GGGAGGACGA UGCGGUGGUC UCCCAAUUA CUCCAUCGUA UC 42
- (2) INFORMATION FOR SEQ ID NO: 279:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 279:
GAGGACGAUG CGGUGGUCUC CCAAUUCUAA ACUUUCUCCA UCGUAUC 47
- (2) INFORMATION FOR SEQ ID NO: 280:
- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:

-continued

(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 280:
GGGAGGACGA UGCGGUGGUC UCCCAAUUU CUCCAUCGUA UC 42

(2) INFORMATION FOR SEQ ID NO: 281:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 281:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU ACCAUCGUAU C 41

(2) INFORMATION FOR SEQ ID NO: 282:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 282:
GGAGGACGAU GCGGUGGUCU CCCAAUUCUA UUCUCAUCG UAUC 44

(2) INFORMATION FOR SEQ ID NO: 283:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 283:
GGAGGACGAU GCGGUGGUCU CCAAUUCUAU UCUCAUCGU AUC 43

(2) INFORMATION FOR SEQ ID NO: 284:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 284:

-continued

GAGGACGAUG CGGUGGUCUC CCAAUUCUAU UCUCCAUCGU AUC 43

(2) INFORMATION FOR SEQ ID NO: 285:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 285:

GAGGACGAUG CGGUGGUCUC AAUUCUAUUC UCUCGUAU C 41

(2) INFORMATION FOR SEQ ID NO: 286:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 286:

GGGAGGCGAU GCGGUGGUCU CCAAUUCUA UUCUCCAUUG UAUC 44

(2) INFORMATION FOR SEQ ID NO: 287:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 44 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 287:

GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AUCUCCAUUG UAUC 44

(2) INFORMATION FOR SEQ ID NO: 288:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 288:

GGGAGGACGA UGCGGUGGUC UCCCAAUUCU ACUCUCCAUUG AUC 43

(2) INFORMATION FOR SEQ ID NO: 289:

-continued

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 289:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU AUCCAUCGUA UC 42
- (2) INFORMATION FOR SEQ ID NO: 290:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 44 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 290:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU UUCUCCAUCG UAUC 44
- (2) INFORMATION FOR SEQ ID NO: 291:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 43 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 291:
GGGAGGACGA UGCGGUGGUC UCCCAAUUCU UCUCUCCAUCGU AUC 43
- (2) INFORMATION FOR SEQ ID NO: 292:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 41 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: RNA
- (ix) FEATURE:
 (D) OTHER INFORMATION: All pyrimidines are 2'-fluor
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 292:
GGGGACGAUG CGGUGGCCCC AAUUCUAUUC UCCAUCGU AU C 41
- (2) INFORMATION FOR SEQ ID NO: 293:
- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid

-continued

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 293:
GGGACGAUGC GGUGGCCCAA UUCUAUUCUC CAUCGUAUC 39

(2) INFORMATION FOR SEQ ID NO: 294:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 294:
GACGAUGCGG UGGCCCAAU CUAUUCUCA UCGUAUC 37

(2) INFORMATION FOR SEQ ID NO: 295:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 295:
GGACGAUGCG GUGGCCCAAU UCUAUUCUCC AUCGUAUC 38

(2) INFORMATION FOR SEQ ID NO: 296:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 296:
GAUGCGGUGG CCAAUUCUA UUCUCCAUCG UAUC 34

(2) INFORMATION FOR SEQ ID NO: 297:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 37 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

-continued

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 297:

GAUGC GGUGG UCUC CCAAU CUAUUCUCCA UCGUAUC 37

(2) INFORMATION FOR SEQ ID NO: 298:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 298:

GGUGGUCUCC CAAUUCUAU CUCCAUCGUA UC 32

(2) INFORMATION FOR SEQ ID NO: 299:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 299:

GGGAGGACGA UG 12

(2) INFORMATION FOR SEQ ID NO: 300:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 300:

GGGAGGACGA UGCGGUGGUC UCCCAAUUCU CUCCAUCGUA UC 42

(2) INFORMATION FOR SEQ ID NO: 301:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

-continued

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 301:
GGGAGGACGA UGCGGUGUCU CCCAAUUCUU CUCAUCGUAU C 41

(2) INFORMATION FOR SEQ ID NO: 302:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 302:
GGGAGGACGA UGCGGGGUCU CCCAAUUCUU CUCCUCGUAU C 41

(2) INFORMATION FOR SEQ ID NO: 303:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 39 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 303:
GGGAGGACGA UGCGGGUCUC CCAAUUCUUC UCUCGUUAUC 39

(2) INFORMATION FOR SEQ ID NO: 304:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 43 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(ix) FEATURE:
(D) OTHER INFORMATION: All pyrimidines are 2'-fluor

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 304:
GGGAGUACGA UGCGGUGGUC UCCCAAUUCU UCUCAUCGU AUC 43

1. A method of inhibiting a platelet derived growth factor (PDGF) comprising contacting said PDGF with a nucleic acid ligand of PDGF.

2. The method of claim 1, wherein the nucleic acid ligand of PDGF comprises a ligand selected from the group of sequences consisting of SEQ ID NOS:93-124 and 128-176.

3. A method for treating a PDGF mediated disease or medical condition comprising administering to a patient in

need thereof a pharmaceutically effective amount of a nucleic acid ligand capable of binding to PDGF.

4. The method of claim 3, wherein the nucleic acid ligand of PDGF comprises a ligand selected from the group of sequences consisting of SEQ ID NOS:93-124 and 128-176.

5. The method of claim 3, wherein said PDGF mediated disease or medical condition is a proliferative disease.

6. The method of claim 5, wherein said proliferative disease is a tumor expressing PDGF or PDGF receptors.

7. A method of inhibiting a human keratinocyte growth factor (hKGF) comprising contacting said hKGF with a nucleic acid ligand of hKGF.

8. The method of claim 7, wherein the nucleic acid ligand of hKGF comprises a ligand selected from the group of sequences consisting of SEQ ID NOS: 189-264 and 268-304.

9. A method for treating a hKGF mediated disease or medical condition comprising administering to a patient in

need thereof a pharmaceutically effective amount of a nucleic acid ligand capable of binding to hKGF.

10. The method of claim 9, wherein the nucleic acid ligand of hKGF comprises a ligand selected from the group of sequences consisting of SEQ ID NOS: 189-264 and 268-304.

11. The method of claim 9, wherein said hKGF mediated disease or medical condition is select.

* * * * *