




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 231/00	A2	(11) International Publication Number: WO 00/18741 (43) International Publication Date: 6 April 2000 (06.04.00)
<p>(21) International Application Number: PCT/JP99/05289</p> <p>(22) International Filing Date: 27 September 1999 (27.09.99)</p> <p>(30) Priority Data: PP 6231 30 September 1998 (30.09.98) AU PQ 2243 16 August 1999 (16.08.99) AU</p> <p>(71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-8514 (JP).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): KONISHI, Nobukiyo [JP/JP]; 22-7, Aotanai, Nagaokakyo-shi, Kyoto 617-0811 (JP). NAKAMURA, Katsuya [JP/JP]; 12-1-103, Kamihamuro 2-chome, Takatsuki-shi, Osaka 569-1044 (JP). YAMAMOTO, Hirofumi [JP/JP]; 1-4-22-207, Kashio, Takarazuka-shi, Hyogo 665-0054 (JP). MANABE, Takashi [JP/JP]; 1-2-103, Maruyamadai, Kawanishi-shi, Hyogo 666-0152 (JP).</p> <p>(74) Agent: TAKASHIMA, Hajime; Yuki Building, 3-9, Hiranomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541-0046 (JP).</p>	<p>(81) Designated States: BR, CA, CN, JP, KR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
(54) Title: NOVEL COMPOUNDS		
<div style="text-align: center;">  <p style="text-align: right;">(I)</p> </div>		
<p>(57) Abstract</p> <p>A compound of formula (I), wherein R¹ is halo(lower)alkyl, halogen or cyano, R² is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino, R³ is hydrogen, hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl, R⁴ is hydrogen or halogen, Q is pyrazolyl, A is lower alkylene optionally substituted with oxo or hydroxy, and n is 0, 1 or 2, provided that when R³ is hydrogen, R² is aryl substituted with lower alkenyl or A is lower alkylene substituted with oxo, or its salt, processes for their preparation and pharmaceutical compositions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION
NOVEL COMPOUNDS

Technical Field

This invention relates to novel pyrazole compounds and pharmaceutically acceptable salts thereof which have pharmacological activity, to a process for their production; and to a pharmaceutical composition containing the same.

Background of Art

Some pyrazole derivatives having antiinflammatory and analgesic activities have been known as described, for example, in Canadian Patent 1 130 808, and EP Patent Publication Nos. 248 594, 272 704, 293 220, 418 845 and 554 829, and WO Patent Publication Nos. 95/15315, 95/15316, 95/15317, 95/15318, 96/14302 and 97/15271.

Disclosure of Invention

One object of this invention is to provide novel pyrazole compounds and pharmaceutically acceptable salts thereof which have an inhibiting activity of COX-II.


Another object of this invention is to provide a process for production of the novel pyrazole compounds.

A further object of this invention is to provide a pharmaceutical composition containing, as an active ingredient, the pyrazole compound or a pharmaceutically acceptable salt thereof.

A still further object of this invention is to provide a use of the novel pyrazole compounds and pharmaceutically acceptable salts thereof for manufacturing a medicament for treating or preventing various diseases.

The present invention relates to novel pyrazole compounds and pharmaceutically acceptable salts thereof, which have pharmaceutical activity such as inhibiting activity of cyclooxygenase-2 (hereinafter described as COX-II), to a process for their production, to a pharmaceutical composition containing the same, and to a use thereof.

The object pyrazole derivatives of this invention are new and can be represented by the following general formula (I).



wherein

R¹ is halo(lower)alkyl, halogen or cyano,

R² is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino,

R³ is hydrogen, hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl,

R⁴ is hydrogen or halogen,


Q is pyrazolyl,

A is lower alkylene optionally substituted with oxo or hydroxy, and
n is 0, 1 or 2,


provided that when R³ is hydrogen, R² is aryl substituted with lower alkenyl or A is lower alkyl substituted with oxo.

The object compound (I) can be prepared by one of the following processes 1 - 5.

Process 1





(II)
or its salt





(I-1)
or its salt



Process 2




Process 3




Process 4




Process 5






Reference Processes

(1)



(2)



wherein R¹, R², R³, A, Q and n are each as defined above,
 R is lower alkyl,

R^{3a} is amino optionally substituted with hydroxy or lower alkyl,

R^{3b} is acyloxy,

R^{3c} is lower alkylsulfonyl or lower alkylsulfinyl, and

R^{3d} is hydroxy substituted with acyl.

"Pyrazolyl" includes 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl and 5-pyrazolyl, in which preferable one is 5-pyrazolyl.

In the above and subsequent description of the present specification, suitable examples of the various definitions to be included within the scope of the invention are explained in detail in the following.

The term "lower" is intended to mean a group having 1 to 6 carbon atom(s), unless otherwise provided.

Suitable "lower alkyl" and lower alkyl moiety in the terms "lower(halo)alkyl", "lower alkylthio" and "lower alkylsulfonyl" may be a straight or branched one having 1 to 6 carbon atom(s), such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, tert-pentyl, hexyl, and the like, in which preferable one is C₁-C₄ alkyl.

Suitable "lower alkenyl" includes, for example, isopropenyl, vinyl, propenyl, betenyl, pentenyl, hexenyl, in which preferable one is isopropenyl.

Suitable "lower alkoxy" includes, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, tert-pentyloxy and hexyloxy, in which preferable one is C₁ - C₄ alkoxy.

Suitable "lower(halo)alkyl" includes, for example, trifluoromethyl, difluoromethyl, trichloromethyl, and the like.

Suitable "halogen" includes, for example, fluorine, chlorine, bromide and iodide.

Suitable "alkylene" includes, for example, methylene, dimethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene, propylene, ethylidene, and the like.

Suitable "acyl" and acyl moiety in the term "acyloxy" include acyl

such as aliphatic acyl, aromatic acyl and heterocyclic acyl.

The aliphatic acyl includes saturated or unsaturated, acyclic or cyclic ones, for example, alkanoyl such as lower alkanoyl (e.g., formyl, acetyl, propionyl, butyl, isobutyl, valeryl, isovaleryl, pivaloyl, hexanoyl, etc.), alkylsulfonyl, such as lower alkylsulfonyl (e.g., mesyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, isobutylsulfonyl, pentylsulfonyl, hexylsulfonyl, etc.), carbamoyl, N-alkylcarbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), alkoxycarbonyl such as lower alkoxycarbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, tert-butoxycarbonyl, etc.), alkenyloxycarbonyl such as lower alkenyloxycarbonyl (e.g., vinyloxycarbonyl, allyloxycarbonyl, etc.), alkenoyl such as lower alkenoyl (e.g., acryloyl, methacryloyl, crotonoyl, etc.), cycloalkanecarbonyl such as cyclo(lower)alkanecarbonyl (e.g., cyclopropanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, etc.), and the like.


The aromatic acyl may include C₆ - C₁₀ aryl (e.g., benzoyl, toluoyl, xylyl, etc.), N-(C₆ - C₁₀)arylcarbamoyl (e.g., N-phenylcarbamoyl, N-tolylcarbamoyl, N-naphthylcarbamoyl, etc.), C₆ - C₁₀ arenesulfonyl (e.g., benzenesulfonyl, tosyl, etc.), and the like.

The heterocyclic acyl may include heterocyclic carbonyl (e.g., furoyl, thenoyl, nicotinoyl, isonicotinoyl, thiazolylcarbonyl, thiadiazolylcarbonyl, tetrazolylcarbonyl, etc.), and the like.

Suitable "aryl" includes, for example, an aryl having 6 to 10 carbon atoms, such as phenyl, tolyl, xylyl, cumenyl, mesityl, naphthyl, in which preferable one is phenyl and tolyl.

Each of R¹ and R² may be attached to any position of the pyrazole ring.

The preferred ones of the compound (I) are compounds having the formula (I')



wherein

R^1 is halo(lower)alkyl, halogen or cyano,

R^2 is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino,

R^3 is hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl,

R^4 is hydrogen or halogen,

A is lower alkylene optionally substituted with oxo or hydroxy, and n is 0, 1 or 2,

provided that when R^3 is hydrogen, R^2 is aryl substituted with lower alkenyl or A is lower alkylene substituted with oxo, or its salt.

The most preferred ones of the compound (I) are compounds having the formula (I') wherein

R^1 is trifluoromethyl, difluoromethyl, chlorine or cyano,

R^2 is phenyl, tolyl, or phenyl or tolyl substituted with substituent(s) selected from the group consisting of chlorine, fluorine, bromine and methoxy,

R^3 is hydroxy, acetoxy, ethoxy, amino, methylamino, methylthio or methylsulfonyl,

R^4 is hydrogen,

A is dimethylene, trimethylene, methylene or pentamethylene, and n is 2.

The preferred one is the compound selected from the group consisting of

- (1) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chloro-4-methoxyphenyl)pyrazole,
- (2) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (3) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (4) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (5) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (6) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylthio)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (7) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(5-hydroxypentyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (8) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(4-hydroxybutyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (9) 1-(3-chloro-4-methoxyphenyl)-5-[4-(ethoxycarbonylmethylsulfonyl)phenyl]-3-(trifluoromethyl)pyrazole,
- (10) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methylaminoethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (11) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-aminoethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (12) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (13) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]phenyl}pyrazole,
- (14) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylsulfonyl)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (15) 5-{4-[(acetoxymethyl)sulfonyl]phenyl}-1-(4-fluorophenyl)-3-(trifluoromethyl)pyrazole,
- (16) 1-(4-fluorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-

- (trifluoromethyl)pyrazole,
(17) 5-{4-[(acetoxymethyl)sulfonyl]phenyl}-1-(4-chloro-3-methoxyphenyl)-3-cyanopyrazole,
(18) 1-(4-chloro-3-methoxyphenyl)-5-{4-[2-hydroxyethyl)sulfonyl]phenyl}-3-cyanopyrazole,
(19) 5-[4-(acetoxymethylthio)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole,
(20) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole,
(21) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-methylphenyl)-pyrazole,
(22) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-methylphenyl)-pyrazole,
(23) 3-chloro-1-(4-isopropenylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole,
(24) 5-[(4-acetoxymethylthio)phenyl]-3-chloro-1-(isopropenylphenyl)-pyrazole,
(25) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}pyrazole,
(26) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
(27) 1-(4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
(28) 1-(4-methylphenyl)-5-{4-[(3-acetoxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
(29) 1-(4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
(30) 5-[4-(acetoxymethylthio)phenyl]-3-trifluoromethyl-1-[(3-chloro-4-methyl)phenyl]pyrazole,
(31) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,

- (32) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (33) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (34) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (35) 1-(3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (36) 1-(3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (37) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (38) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (39) 5-{4-[(acetoxymethyl)sulfinyl]-3-fluorophenyl}-1-(3-chloro-4-methoxy-phenyl)-3-(trifluoromethyl)pyrazole,
- (40) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-3-fluorophenyl}-3-(trifluoromethyl)pyrazole,
- (41) 1-(4-chloro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (42) 1-(1-hydroxy-1-methylethyl)-5-[4-(methylthio)phenyl]-3-(trifluoromethyl)pyrazole,
- (43) 1-(4-isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (44) 1-(3-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (45) 1-(3-chlorophenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (46) 1-(4-isopropylphenyl)-5-[4-(acetoxymethylthio)phenyl]-3-(trifluoromethyl)pyrazole,
- (47) 1-(4-isopropylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-

- (trifluoromethyl)pyrazole,
- (48) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-cyano-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (49) 1-(3-cyano-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (50) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-methoxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (51) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(N-hydroxycarbamoyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (52) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(methylcarbamoyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (53) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-oxopropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (54) 5-{4-[(3-acetoxypentyl)sulfonyl]phenyl}-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (55) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[3-(dimethylamino)propyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole hydrochloride,
- (56) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(dimethylamino)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole hydrochloride,
- (57) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(difluoromethyl)pyrazole,
- (58) 1-(3-chloro-4-methoxyphenyl)-3-difluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (59) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(2-hydroxyethoxy)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (60) 5-[4-(acetoxymethylthio)phenyl]-1-(4-bromo-3-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (61) 1-(4-bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (62) 1-(4-bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,

- (63) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-difluoromethyl-1-[4-(methoxyphenyl)pyrazole,
- (64) 1-(4-methoxyphenyl)-3-difluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (65) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-bromophenyl)pyrazole,
- (66) 1-(3-bromophenyl)-3-chloro-5-{4-(3-hydroxypropyl)sulfonyl}phenyl}pyrazole,
- (67) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-fluoro-4-methylphenyl)pyrazole,
- (68) 3-chloro-1-(3-fluoro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (69) 5-[(4-acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluoro-3-methylphenyl)pyrazole,
- (70) 3-chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (71) 3-chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (72) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (73) 1-(3-chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (74) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (75) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}pyrazole hydrochloride,
- (76) 5-[4-(4-acetoxymethylthio)phenyl]-3-chloro-1-(4-chloro-3-methylphenyl)pyrazole,
- (77) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (78) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)-

- sulfonyl]phenyl}pyrazole,
(79) 3-chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole,
(80) 3-chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}pyrazole,
(81) 3-chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-(4-methylphenyl)pyrazole,
(82) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole,
(83) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-methylphenyl)-pyrazole,
(84) 3-chloro-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-1-(3-methylphenyl)pyrazole,
(85) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-phenylpyrazole,
(86) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-phenylpyrazole,
(87) 3-chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-phenylpyrazole,
(88) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole,
(89) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole,
(90) 3-chloro-1-(4-chloro-3-methoxyphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole,
(91) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(3-chlorophenyl)-pyrazole,
(92) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chlorophenyl)-pyrazole,
(93) 3-chloro-1-(3-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
(94) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-chlorophenyl)-pyrazole,

- (95) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chlorophenyl)-pyrazole,
- (96) 3-chloro-1-(4-chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-pyrazole,
- (97) 3-chloro-1-(4-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (98) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-fluorophenyl)-pyrazole,
- (99) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluorophenyl)-pyrazole,
- (100) 3-chloro-1-(4-fluorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (101) 5-(4-methylphenyl)-1-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (102) 5-(4-methylphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole, and
- (103) 5-(4-chloro-3-methoxyphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole.

The more preferred one is the compound selected from the group consisting of

- (1) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (2) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (3) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (4) 1-(4-bromo-3-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (5) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (6) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-

phenyl}-3-(trifluoromethyl)pyrazole,

(7) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}pyrazole,

(8) 3-chloro-1-(4-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]-phenyl}pyrazole,

(9) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,

(10) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,

(11) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole, and

(12) 5-(4-chloro-3-methoxyphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole.

The compounds (I) according to the present invention may contain one or more asymmetric centers, and thus they can exist as enantiomers or diastereoisomers, and the invention includes both mixtures and separate individual isomers.

Suitable salts of the compounds (I) are conventional pharmaceutically acceptable salts and may include a salt with a base or an acid addition salt such as a salt with an inorganic base, for example, an alkali metal salt (e.g. sodium salt, potassium salt, cesium salt, etc.), an alkaline earth metal salt (e.g. calcium salt, magnesium salt etc.), an acid addition salt such as a salt with an inorganic base, for example, an alkali metal salt (e.g. sodium salt, potassium salt, cesium salt, etc.), an alkaline earth metal salt (e.g. calcium salt, magnesium salt etc.), an ammonium salt; a salt with an organic base, for example, an organic amine salt (e.g. triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, etc.), etc.; an inorganic acid addition salt (e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.); an organic carboxylic or sulfonic acid addition salt (e.g. formate, acetate,

trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, etc.); a salt with a basic or acidic amino acid (e.g. arginine, aspartic acid, glutamic acid, etc.); and the like, and the preferable example thereof is an acid addition salt.

The compound (I) according to the present invention can be in the form of a solvate, which was included within the scope of the present invention. The solvate preferably includes a hydrate, an ethanolate, and so on.

Also included in the scope of invention are radiolabelled derivatives of compounds (I) which are suitable for biological studies.

Process 1

The compound (I-1) can be prepared by reacting the compound (II) or its salt with CH_3COONa (sodium acetate) or $(\text{CH}_3\text{CO})_2\text{O}$ (acetic anhydride), and then the oxydation.

(First step)

The reaction is usually carried out at reflux temperature. The solvents to be used are conventional solvents such as water, alcohol [e.g., methanol, ethanol, isopropanol, etc.], alkanolic acid, tetrahydrofuran, dioxane, toluene, methylene chloride, chloroform, N,N-dimethylformamide, or any other organic solvents which do not adversely affect the reaction, or the mixture thereof, preferably, acidic solvent such as alkanolic acid (e.g., acetic acid).

(Oxydation)

The oxidizing agent to be used is MMPAC (magnesium monoperoxyphthalate).

This reaction is usually carried out in a solvent which does not adversely influence the reaction such as acetic acid, dichloromethane, acetone, ethyl acetate, chloroform, water, an alcohol [e.g. methanol, ethanol, etc.], a mixture thereof or the like.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process 2

The compound (I-2) can be prepared by reacting a compound (IV) or its salt with a compound (V) or its salt.

The reaction is usually carried out in a conventional solvent such as water, alcohol [e.g., methanol, ethanol, isopropanol, etc.], alkanolic acid, tetrahydrofuran, dioxane, toluene, methylene chloride, chloroform, N,N-dimethylformamide, or any other organic solvents which do not adversely affect the reaction, or the mixture thereof, preferably, acidic solvent such as alkanolic acid (e.g., acetic acid).

The reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

Process 3

The compound (I-4) or its salt can be prepared from the compound (I-3) or its salt.

Amination can preferably be carried out by, for example, reacting the compound (I-4) with a mesylating reagent such as methanesulfonyl chloride and methanesulfonic anhydride to give a mesylate. The solvents to be used are, for example, dichloromethane, chloroform, tetrahydrofuran, or any other organic solvents which do not adversely affect the reaction.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Then, the mesylate is reacted with an alkylamine (e.g., methylamine, dimethylamine, ethylamine, etc.) or hydroxyamine. The solvents to be used are, for example, acetonitrile, tetrahydrofuran, toluene, or any other organic solvents which do not adversely affect the reaction, or the mixture thereof.

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

Process 4

The compound (I-5) can be prepared from the compound (I-3).

The acylating agent to be used are, for example, a lower alkanolic anhydride (e.g., acetic anhydride, etc.), lower alkanoyl halide (e.g., acetyl chloride, etc.), and the like.

The reaction is usually carried out in a solvent such as dichloromethane, tetrahydrofuran, toluene, a solvent which does not adversely influence the reaction, or a mixture thereof.

The reaction temperature is not critical and the reaction can be carried out under cooling to warming.

Process 5

The compound (I-7) can be prepared by reacting the compound (I-6) with an oxidizing agent.

The suitable oxidizing agent may be hydrogen peroxide, cumene hydroperoxide, tert-butyl hydroperoxide, Jones reagent, peracid [e.g. peracetic acid, perbenzoic acid, m-chloroperbenzoic acid, monopersulfate compound (Oxone®), etc.], chromic acid, potassium permanganate, alkali metal periodate [e.g. sodium periodate, etc.], and the like.

This reaction is usually carried out in a solvent which does not adversely influence the reaction such as acetic acid, dichloromethane, acetone, ethyl acetate, chloroform, water, an alcohol [e.g. methanol, ethanol, etc.], a mixture thereof or the like.

The reaction temperature is not critical, and the reaction is usually carried out under cooling to warming.

Reference Processes

(1) The compound (II') can be prepared by reacting the compound (V) or its salt with a hydrazine derivative (VI) or its salt.

The reaction is usually carried out in a conventional solvent such as water, alcohol [e.g., methanol, ethanol, isopropyl alcohol, etc.], alkanolic acid, tetrahydrofuran, dioxane, toluene, methylene chloride, chloroform, N,N-dimethylformamide, or any other organic solvents which do not adversely affect the reaction, or the mixture thereof,

preferably, acidic solvent such as alkanolic acid (e.g., acetic acid.)

The reaction temperature is not critical and the reaction is usually carried out under cooling to warming.

(2) The compound (IV) can be prepared from the compound (I-5) or its salt in a similar manner to that of Process 2 and below mentioned Preparations.

The compounds obtained by the above processes can be isolated and purified by a conventional method such as pulverization, recrystallization, column chromatography, reprecipitation, or the like.

Suitable salts of the compounds (II) and (V) are, for example, an inorganic acid addition salt (e.g. hydrochloride, hydrobromide, sulfate, phosphate, etc.), an organic carboxylic or sulfonic acid addition salt (e.g. formate, acetate, trifluoroacetate, maleate, tartrate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, etc.), and the like.

The object compound (I) possesses inhibiting activity of COX-II and possesses strong antiinflammatory, analgesic, antithrombotic, anti-cancer activities and so on. The object compound (I) and pharmaceutically acceptable salts thereof, therefore, are useful for the treatment and/or prevention of inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunological diseases, thrombosis, cancer and neurodegenerative diseases in human beings or animals, and more particularly for the treatment and/or prevention of inflammation and pain in joint and muscle [e.g. rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, juvenile arthritis, etc.], inflammatory skin condition [e.g. sunburn, burns, eczema, dermatitis, etc.], inflammatory eye condition [e.g. conjunctivitis, etc.], lung disorder in which inflammation is involved [e.g. asthma, bronchitis, pigeon fancier's disease, farmer's lung, etc.], condition of the gastrointestinal tract associated with inflammation [e.g. aphthous ulcer, Crohn's disease, atopic gastritis, gastritis varioliforme, ulcerative colitis, coeliac disease, regional ileitis, irritable bowel

syndrome, etc.], gingivitis, inflammation, pain and tumescence after operation or injury, pyrexia, pain and other conditions associated with inflammation, particularly those in which lipoxygenase and cyclooxygenase products are a factor, systemic lupus erythematosus, scleroderma, polymyositis, tendinitis, bursitis, periarteritis nodosa, rheumatic fever, Sjogren's syndrome, Behcet disease, thyroiditis, type I diabetes, nephrotic syndrome, aplastic anemia, myasthenia gravis, uveitis contact dermatitis, psoriasis, Kawasaki disease, sarcoidosis, Hodgkin's disease, Alzheimer's disease, and the like. Additionally, the object compound (I) or a salt thereof is expected to be useful as therapeutical and/or preventive agents for cardiovascular or cerebrovascular diseases, the diseases caused by hyperglycemia and hyperlipemia.

The compound (I) of the present invention has much advantage, such as more selective inhibitory activity of COX-II, stronger activity, more suitable half-life, decreased adverse effect, or the like, compared to the known pyrazole compounds shown in the prior arts.

In order to illustrate the usefulness of the object compound (I), the pharmacological test data of the compound (I) are shown in the following.

COX-I and COX-II activity in vitro:

(i) Test Method:

a. Preparation of the recombinant cyclooxygenase (COX)

The human cyclooxygenase COX-I and COX-II were expressed in transfected Chinese hamster ovary (CHO) cells. Monolayer cultures of semi-confluent CHO cells stably expressing COX-I and COX-II were washed twice and scraped into phosphate buffered saline (PBS). The cells were centrifuged at 200 x g for 5 minutes and the cell pellet was sonicated in reaction buffer containing 100 mM Tris-HCl (pH 7.4), 2 μ M hematin and 5 mM tryptophan. Broken cells were centrifuged for 5 minutes at 1700 x g at 4°C and the supernatants were used as crude

enzymes.

Cyclooxygenase activities in the absence or presence of inhibitors were measured by determining the level of prostaglandin E₂ (PGE₂) synthesis from arachidonic acid. Enzymes (1 µg for COX-I and/or 3 µg for COX-II) in a total volume of 200 µl of reaction buffer were incubated in the absence and presence of various concentrations of inhibitors for 5 minutes at 30°C. The reaction was then started by the addition of arachidonic acid to the final concentration of 10 µM. The reaction was terminated by 50 µl of HCl (1N) after incubation at 30°C for 5 minutes.

PGE₂ was extracted with ethyl acetate, concentrated under a stream of nitrogen and analyzed by a radio immunoassay kit (Amersham) according to the manufacture's instructions.

b. Assay for human recombinant COX-I and COX-II activity

COX activity was assayed as PGE₂ formation using radioimmunoassay to detect the prostaglandin release. The appropriate COX enzyme was incubated in 0.1 M Tris-HCl buffer (pH 7.3) containing hematin and tryptophan with the addition of arachidonic acid (10 µM) for 5 minutes at 37°C. Compounds were pre-incubated with the enzyme for 5 minutes prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme was stopped after 5 minutes at 37°C by addition of 20 µl of 1N HCl. PGE₂ formation was measured by radioimmunoassay (Amersham).

(ii) Test Results : COX-I and COX-II activity in vitro :

Test compound (Example No.)	Human COX-II IC ₅₀ (µM)	Human COX-I IC ₅₀ (µM)
5	≤ 1	>100
3	≤ 1	>100
12	≤ 1	>100

The compound (I) and a pharmaceutically acceptable salt thereof, are used as a medicament by intravenous, intracutaneous,

intramuscular, pulmonary, or oral administration, or insufflation to human beings or animals.

A pharmaceutical composition of the present invention is a homogeneous mixture which comprises one of the compounds (I) or pharmaceutically acceptable salts thereof, as an active ingredient, in association with a pharmaceutically acceptable carrier or excipient. The pharmaceutical composition is manufactured by mixing the sufficient amount of the compound (I) or a salt thereof as an active ingredient with a pharmaceutically non-toxic carrier or excipient to give homogeneous mixture. The pharmaceutically non-toxic carriers and excipients may be organic or inorganic and solid or liquid, and can be any of the conventional ones suitable for oral, parenteral or external (topical) administration.

For therapeutic purpose, the pharmaceutical composition of the present invention can be used in a form of a pharmaceutical preparation, for example, in a solid, semisolid, or liquid form. The pharmaceutical preparations may be capsules, tablets, dragees, granules, inhalant, suppositories, solution, lotion, suspension, emulsion, ointment, gel, or the like. If desired, there may be included in these preparations, auxiliary substances, stabilizing agents, wetting or emulsifying agents, buffers and other commonly used additives.

In the pharmaceutical composition the compound (I) or a pharmaceutically acceptable salt thereof is included in an sufficient amount to have the desired effects of aforementioned pharmaceutical activities on the aforesaid diseases in human beings or animals.

While the dosage of therapeutically effective amount of the compound (I) will vary depending upon the age and condition of the patient, an average single dose of about 0.1 mg, 1 mg, 10 mg, 50 mg, 100 mg, 250 mg, 500 mg and 1000 mg of the compound (I) may be effective for treating the above-mentioned diseases. In general, amounts between 0.1 mg/body and about 1,000 mg/body may be administered

per day.

The following Preparations and Examples are given for the purpose of illustrating this invention.

Preparation 1

A mixture containing 1-[4-(methylthio)phenyl]-4,4,4-trifluorobutane-1, 3-dione (8.0 g), (3-chloro-4-methoxy)phenylhydrazine hydrochloride (6.4 g) and acetic acid (45 ml) was stirred at 100-110°C. After 3 hours, the reaction mixture was concentrated in vacuo. The resultant residue was partitioned between ethyl acetate and water. The organic layer was separated, washed with water and dried over magnesium sulfate. Evaporation of the solvent under reduced pressure gave an oil (11.5 g). The obtained product was used for next reaction without purification.

This oil was dissolved in dichloromethane (100 ml), and 3-chlorobenzoic acid (80%, 5.92 g) was added to this solution over a period of 15 minutes at an ice-bath temperature. After being stirred for 1 hour, the reaction mixture was diluted with dichloromethane (50 ml) and then washed with aqueous sodium carbonate solution (40 ml x 2) and brine. The organic layer was dried over $MgSO_4$ and evaporated under reduced pressure. The resulting residue was purified by column chromatography on silica gel eluting with ethyl acetate to give 1-(3-chloro-4-methoxyphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole (8.95 g) in 71% yield.

IR (KBr) : 1505, 1270, 1235, 1160, 1135, 1050 cm^{-1}

NMR ($CDCl_3$, δ) : 2.76(3H,s), 3.93(3H,s), 6.80-7.67(8H,m)

MASS : 415(M+1)

Preparation 2

A mixture of tert-butyl nitrite (2.6 ml) and copper (II) chloride(2.3 g) in acetonitrile (25 ml) was heated at 60-65°C for 30 minutes. After the reaction mixture was cooled to room temperature, 3-amino-1-(3-chloro-4-methoxyphenyl)-5-[4-(methylthio)phenyl]pyrazole (5.0 g) was

added to the mixture, and then the resulting mixture was stirred for 1.5 hours and concentrated in vacuo. The residue was partitioned between aqueous 10% hydrochloric acid solution and ethyl acetate. The organic layer was separated and washed with brine. The evaporation of the solvent followed by column chromatography on silica gel eluting with dichloromethane gave 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-[4-(methylthio)phenyl]pyrazole (3.1 g) in 58% yield. This product was oxidized by using 3-chloroperbenzoic acid, followed by an usual work-up, to give 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole (2.6 g) in 80% yield.

IR (KBr) : 1505, 1375, 1270, 1055 cm^{-1}

NMR (CDCl_3 , δ) : 2.75(3H,s), 3.92(3H,s), 6.49(1H,s), 6.82- 7.65(7H,m)

MASS : 381(M+1)

Preparation 3

A mixture of methyl 4-(4-methylthiophenyl)-2,4-dioxobutanoate (5.0 g) and (4-chloro-3-methoxy)phenylhydrazine hydrochloride (5.2 g) in dioxane (30 ml) and ethanol (30 ml) was stirred at reflux temperature for 3 hours. The reaction mixture was cooled to room temperature and evaporated under reduced pressure. The resulting residue was purified by column chromatography on silica gel eluting with ethyl acetate to give methyl {1-(4-chloro-3-methoxyphenyl)-5-[4-(methylthio)phenyl]pyrazol-3-yl}carboxylate (6.4 g) in 83.0% yield.

NMR (CDCl_3 , δ) : 2.49(3H,s), 3.75(3H,s), 3.98(3H,s), 6.75-7.33(8H,m)

MASS : 389(M+1)

Preparation 4

To methyl {1-(4-chloro-3-methoxyphenyl)-5-[4-(methylthio)phenyl]pyrazol-3-yl}carboxylate (6.3 g) in formamide (35 ml) was added sodium methoxide (920 mg), and the resulting mixture was heated at 100°C under a nitrogen atmosphere. After 40 minutes the reaction mixture was cooled in an ice bath, and the resulting precipitates were collected by filtration and washed with ethyl acetate to give the desired

pyrazole amide (4.18 g). The amide was added to Vilsmeier reagent, prepared from N,N-dimethylformamide (15 ml) and phosphorus oxychloride (1.8 ml), at 5°C and then the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into ice water, and the resulting precipitates were collected by filtration and dried at 50°C for 4 hours to afford the desired pyrazole nitrile (4.1 g). To this nitrile in dichloromethane (40 ml) was added by portions 3-chloroperbenzoic acid (2.84 g) at an ice-bath temperature. After being stirred at the same temperature for 2 hours, the reaction mixture was diluted with dichloromethane (50ml) and washed successively with aqueous sodium carbonate solution and brine, and dried over magnesium sulfate. The evaporation of the solvent gave 1-(4-chloro-3-methoxyphenyl)-3-cyano-5-[4-(methylsulfinyl)phenyl]pyrazole (4.16 g) in 69.1% yield.

IR (KBr) : 2235, 1590, 1285, 1075 cm^{-1}

NMR (CDCl_3 , δ) : 2.76(3H,s), 3.82(3H,s), 6.66-7.68(8H,m)

MASS : 372(M+1)

Preparation 5

1-(4-Fluorophenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 1.

IR (Neat) : 1605, 1245, 1160, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.76(3H), 6.82(1H,s), 7.03-7.67(7H,m)

MASS: 369(M+1)

Preparation 6

To a solution of 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-[4-(acetoxymethylsulfonyl)phenyl]pyrazole (2.0 g) in tetrahydrofuran (12 ml) and methanol (6 ml) was added aqueous 1N NaOH solution (4.8 ml) at an ice-bath temperature. After 15 minutes, the reaction mixture was allowed to warm to room temperature. After 30 minutes the reaction mixture was concentrated under reduced pressure to give sodium 4-[3-

chloro-1-(3-chloro-4-methoxyphenyl)pyrazol-5-yl]benzenesulfinate (1.8 g) as a powder.

IR (KBr) : 3400, 1505, 1370, 1265 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.87(3H,s), 6.76(1H,s), 7.08-7.51(7H,m)

Preparation 7

Sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 1645, 1510, 1235, 1135 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.89(3H,s), 7.15-7.58(8H,m)

Preparation 8

Sodium 4-[1-(4-fluorophenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

Preparation 9

Sodium 4-[1-(4-chloro-3-methoxyphenyl)-3-cyanopyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 2240, 1595, 1490, 1235 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.77(3H,s), 6.79-7.51(8H,m)

Preparation 10

A mixture of 1-[4-(methylthio)phenyl]-4,4,4-trifluoro-butane-1,3-dione (2.5 g) and 3-chlorophenylhydrazine hydrochloride (2.1 g) in acetic acid (20 ml) was stirred at 120°C for 1.5hr. The reaction mixture was cooled to room temperature and evaporated under reduced pressure.

The residue was partitioned between ethyl acetate and water. The organic layer was separated, and washed successively with saturated sodium carbonate solution and brine, dried over magnesium sulfate. The evaporation of the solvent gave 1-(3-chlorophenyl)-5-[4-(methylthio)phenyl]-3-(trifluoromethyl)pyrazole (3.8 g).

Preparation 11.

To a solution of 1-(3-chlorophenyl)-5-[4-(methylthio)phenyl]-3-(trifluoromethyl)pyrazole (3.8 g) in methylene chloride (40 ml) was added m-chloroperbenzoic acid (2.34 g: 70% purity) at 0-5°C with ice-water bath. After being stirred for 1 hour, the reaction mixture was washed with saturated sodium carbonate solution and brine, and dried over magnesium sulfate. The evaporation of the solvent gave 1-(3-chlorophenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole (3.76 g)

Preparation 12

To a solution of 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole (3.05 g) in methanol (75 ml) and tetrahydrofuran (15 ml) was added 1N-NaOH solution (3.9 ml) at room temperature. After being stirred for 30 minutes, the reaction mixture was evaporated under reduced pressure. The residue was triturated with ethanol to give sodium 4-[1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate (1.44 g).

Preparation 13

3-Chloro-1-(4-methylphenyl)-5-[4-(methylthio)phenyl]pyrazole (2.8 g) was dissolved in dichloromethane (30 ml) and cooled with an ice.

Meta-chloroperbenzoic acid (1.69 g) was added portionwise to this solution. After stirred for 1 hour, the reaction mixture was quenched with aqueous sodium thiosulfate solution. The aqueous layer was separated, and the organic layer was washed with aqueous sodium bicarbonate solution and brine, dried over magnesium sulfate. After evaporation of the solvent, the obtained residue was triturated with isopropyl ether to give a powder, 3-chloro-1-(4-methylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole (2.69 g).

NMR (DMSO-d₆, δ); 2.33 (3H, s), 2.75 (3H, s), 6.89 (1H, s), 7.18 (2H, d, J=8.4 Hz), 7.25 (2H, d, J=8.4 Hz),

7.43 (2H, d, J=8.4 Hz), 7.67 (2H, d, J=8.4 Hz).

IR (KBr): 3464, 3429, 3116, 1041, 2993, 2912 cm⁻¹.

Mass m/e : 331 (M++1).

Preparation 14

To a solution of 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-methylphenyl)pyrazole (2.65 g) in tetrahydrofuran (20 ml) was added 1N NaOH (6.54 ml) at room temperature. After stirred for 1 hour, the reaction mixture was concentrated under reduced pressure. The obtained residue was partitioned between ethyl ether (20ml) and water (20ml). The aqueous phase was separated and concentrated under reduced pressure to give a powder, which was washed with isopropyl ether (20 ml) and collected by filtration, giving 4-[3-chloro-1-(4-methylphenyl)pyrazol-5-yl]benzenesulfinate (2.34 g).

NMR (DMSO- d_6 , δ); 2.32 (3H, s), 6.75 (1H, s), 7.13-7.25 (6H, m), 7.48 (2H, d, J=8.1 Hz).

IR (KBr): 3639, 3535, 3396, 3128, 1664, 1604 cm^{-1} .

Mass m/e : 333 (M+).

Preparation 15

Sodium 4-[3-chloro-1-(4-isopropenylphenyl)pyrazol-5-yl]benzenesulfinate was prepared according to a similar manner to that of Preparation 14.

IR (KBr) : 1645, 1565, 1550, 1515, 1375 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.10(3H, s), 5.15(1H, m), 5.50(1H, m), 6.76(1H, s), 7.16-7.58(8H, m)

Preparation 16

1-(4-Methylphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (KBr) : 1510, 1465, 1275, 1235, 1160, 1125 cm^{-1}

Preparation 17

Sodium 4-[1-(4-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 21, Example 22 and Preparation 14.

IR (KBr) : 1645, 1510, 1280, 1235, 1160, 1130 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.34(3H, s), 7.14-7.44(9H, m)

Preparation 18

1-(4-Methoxyphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)-pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (Neat) : 1510, 1465, 1240, 1160, 1130, 1100 cm^{-1}

NMR (CDCl_3 , δ) : 2.75(3H, s), 3.86(3H, s), 6.80-7.63(9H, m).

MASS : 381($M^+ + 1$)

Preparation 19

Sodium 4-[1-(4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (KBr) : 1615, 1510, 1465, 1240, 1165, 1130 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.79(3H, s), 6.98-7.44(9H, m)

Preparation 20

Sodium 4-[1-(3-chloro-4-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 20 and Preparation 12.

IR (KBr) : 3365, 1500, 1270, 1235, 1160, 1130 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.36(3H, s), 7.11-7.53(8H, m)

Preparation 21

1-(4-Fluoro-3-methylphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (Neat) : 1500, 1230, 1160, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.28(3H, s), 2.75(3H, s), 6.81(1H, s), 6.96-7.77(7H, m).

MASS : 383($M^+ + 1$)

Preparation 22

Sodium 4-[1-(4-fluoro-3-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example

19, Example 20 and Preparation 12.

IR (KBr) : 1590, 1575, 1502, 1460, 1415, 1250, 1165 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.25(3H, s), 7.06-7.59(8H, m)

Preparation 23

1-(3-Methylphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (Neat) : 1240, 1160, 1130, 1090 cm^{-1}

NMR (CDCl_3 , δ) : 2.35(3H, s), 2.78(3H, s), 6.81(1H, s), 6.81-7.63(8H, m).

MASS : 365($M^+ + 1$)

Preparation 24

Sodium 4-[1-(3-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (KBr) : 1615, 1465, 1275, 1240, 1160, 1130 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.33(3H, s), 6.99-7.44(9H, m)

Preparation 25

1-(3-Fluoro-4-methylphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (KBr) : 1685, 1505, 1460, 1270, 1245, 1160, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.30(3H, s), 2.76(3H, s), 6.80(1H, s), 6.92-7.66(7H, m).

MASS : 383($M + 1$)

Preparation 26

Sodium 4-[1-(3-fluoro-4-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (KBr) : 1590, 1240, 1165, 1125 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.26(3H, s), 7.02-7.47(8H, m)

Preparation 27

1-(3-Chloro-4-methoxyphenyl)-5-(4-methylsulfinyl-3-fluorophenyl)-3-

(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10.

mp : 117-119°C

IR (KBr) : 1500, 1270, 1225, 1155, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.52(3H, s), 3.93(3H, s), 6.73-7.47(7H, m).

MASS : 417(M^+1)

Preparation 28

Sodium 2-fluoro-4-[1-(4-methoxy-3-chlorophenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 20 and Preparation 12.

NMR (DMSO-d_6 , δ) : 3.90(3H, s), 6.97-7.62(7H, m)

Preparation 29

1-(4-Chloro-3-methylphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (Neat) : 1475, 1405, 1265, 1235, 1160, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.37(3H, s), 2.75(3H, s), 6.81(1H, s), 6.91-7.77(7H, m).

MASS : 399(M^+1)

Preparation 30

Sodium 4-[1-(4-chloro-3-methylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (KBr) : 1610, 1475, 1270, 1240, 1160, 1135 cm^{-1}

NMR (DMSO-d_6 , δ) : 2.34(3H, s), 7.04-7.52(8H, m)

Preparation 31

Sodium 4-[1-(4-isopropenylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 20 and Preparation 12. The obtained compound could not be characterized by its spectroscopic data because of containing some by-products and the crude was used for the next reaction without purification.

Preparation 32

1-(3-Methoxyphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)-pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

IR (Neat) : 1600, 1490, 1465, 1280, 1250, 1215, 1160 cm^{-1}

NMR (CDCl_3 , δ) : 2.75(3H, s), 3.76(3H, s), 6.78-8.13(9H, m).

MASS : 381($\text{M}^+ + 1$)

Preparation 33

Sodium 4-[1-(3-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]-benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (KBr) : 1605, 1490, 1465, 1435, 1280, 1250, 1220, 1165, 1130 cm^{-1}

NMR (DMSO-d_6 , δ) : 3.72(3H, s), 6.82-7.62(9H, m)

Preparation 34

Sodium 4-[1-(4-isopropylphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 20 and Preparation 12.

NMR (DMSO-d_6 , δ) : 1.21(6H, d, $J=7.0\text{Hz}$), 2.87-3.01(1H, m), 7.14-7.44(9H, m)

Preparation 35

1-(3-Cyano-4-methoxyphenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

mp : 148-150°C

IR (KBr) : 2235, 1505, 1285, 1235, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.77(3H, s), 3.98(3H, s), 6.82(1H, s), 6.95-7.69(7H, m).

MASS : 406($\text{M}^+ + 1$)

Preparation 36

Sodium 4-[1-(3-cyano-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 12.

IR (KBr) : 2230, 1645, 1505, 1280 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.95(3H, s), 7.17-7.93(8H, m).

Preparation 37

Sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(difluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 1500, 1450, 1270, 1170 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.88(3H, s), 6.82-7.33(9H, m).

Preparation 38

Sodium 4-[1-(4-bromo-3-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar manner to that of Example 20 and preparation 12.

IR (KBr) : 1590, 1475, 1405, 1255, 1225, 1160 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.75(3H, s), 6.73-7.64(8H, m).

Preparation 39

Sodium 4-[1-(4-methoxyphenyl)-3-(difluoromethyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 3401, 3268, 1616, 1512, 1247, 1033 cm^{-1}

NMR (DMSO- d_6 , δ) : 3.78(3H, s), 6.82-7.44(9H, m).

Preparation 40

1-(3-Bromophenyl)-3-chloro-5-[4-(methylthio)-phenyl]pyrazole was prepared by the treatment of 4-methylthiocinnamionitrile with 3-bromophenylhydrazine hydrochloride followed by dehydrogenation with MnO_2 , and chlorination via the corresponding diazonium salt as described in Preparation 2.

IR (Neat) : 1590, 1480, 1370, 1235, 1095 cm^{-1}

NMR (CDCl_3 , δ) : 2.49(3H, s), 5.77(1H, s), 7.09-7.59(8H, m).

MASS : 379(M^+)

Preparation 41

Sodium 4-[1-(3-bromophenyl)-3-chloropyrazol-5-yl]benzenesulfinate was prepared by a similar manner to that of Preparation 14.

IR (KBr) : 3375, 3140, 1585, 1475, 1370, 1240 cm^{-1}

Preparation 42

Sodium 4-[1-(3-fluoro-4-methylphenyl)-3-chloropyrazol-5-yl]-benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 1590, 1505, 1370 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.24(3H, s), 6.77(1H, s), 6.92-7.68(7H, m).

Preparation 43

Sodium 4-[1-(4-fluoro-3-methylphenyl)-3-chloropyrazol-5-yl]-benzenesulfinate was prepared by a similar procedure to that of Preparation 6.

IR (KBr) : 3375, 1500, 1375, 1230 cm^{-1}

NMR (DMSO- d_6 , δ) : 2.23(3H, s), 6.77(1H, s), 7.03-7.44(7H, m).

Preparation 44

This starting material was prepared by the treatment of 4-methylthiocinnamionitrile and 4-chloro-3-methylphenylhydrazine hydrochloride followed by dehydrogenation with MnO_2 , which was converted to 3-chloro-1-(4-chloro-3-methylphenyl)-5-[4-(methylthio)phenyl]pyrazole by using a similar manner described in Preparation 2.

NMR (CDCl_3 , δ) : 2.38(3H, s), 2.48(3H, s), 6.40(1H, s), 6.90(1H, dd, $J=8.5$ and 2.6Hz), 7.10-7.31(6H, m).

MASS : 349 (M^+)

Preparation 45

Sodium 4-[3-chloro-1-(4-chloro-3-methylphenyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Example 22 and Preparation 14.

Preparation 46

Sodium 4-[3-chloro-1-(3-chloro-4-methylphenyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 13, Example 21, Example 22 and Preparation 14.

Preparation 47

3-Chloro-1-(3-methylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole was obtained from 3-chloro-1-(3-methylphenyl)-5-[4-(methylthio)phenyl]pyrazole in the similar manner that of Preparation 13.

NMR (DMSO- d_6 , δ) : 2.30(3H, s), 2.75(3H, s), 6.90(1H, s), 6.98-7.02(1H, m), 7.22-7.34(3H, m), 7.44(2H, d, J=8.3Hz), 7.67(2H, d, J=8.3Hz).

IR (KBr) : 3120, 3049, 2989, 2910 cm^{-1} .

Mass m/e : 331 ($M^+ + 1$).

Preparation 48

Sodium 4-[3-chloro-1-(3-methylphenyl)pyrazol-5-yl]benzenesulfinate was obtained from 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole in the similar manner that of Preparation 14.

NMR (DMSO- d_6 , δ) ; 2.31(3H, s), 6.77(1H, s), 6.93-6.96(1H, m), 7.17-7.32(5H, m), 7.43(2H, d, J=8.1Hz).

IR (KBr) : 3623, 3396, 3363, 3059, 1668, 1616 cm^{-1} .

Mass m/e : 333 ($M^+ + 1$).

Preparation 49

3-Chloro-1-(3-chlorophenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole was prepared by a similar procedure to that of Preparation 13.

NMR (CDCl_3 , δ) : 2.75(3H, s), 6.51(1H, s), 7.05(1H, dt, J=8.2Hz), 7.28-7.41(3H, m), 7.39(2H, d, J=8Hz), 7.64(2H, d, J=8Hz).

IR (KBr) : 1047 cm^{-1}

Mass(m/z) : 351(M+1)

Preparation 50

Sodium 4-[3-chloro-1-(3-chlorophenyl)pyrazol-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 14.

NMR (DMSO- d_6 , δ) : 6.80(1H, s), 7.13-7.25(4H, m), 7.24-7.47(4H, m),

IR (KBr) : 1621 cm^{-1}

mass (m/z) : 351(M)

Preparation 51

3-Chloro-1-(4-chlorophenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole was prepared by a similar procedure to that of Preparation 13.

NMR (CDCl₃, δ) : 2.76(3H, s), 6.50(1H, s), 7.20(2H, d, J=8.8Hz), 7.33(2H, d, J=9Hz), 7.37(2H, d, J=8Hz), 7.63(2H, d, J=8Hz).

IR (KBr) : 1049 cm⁻¹

mass (m/z) : 351(M+1)

Preparation 52

Sodium 4-[3-chloro-1-(4-chlorophenyl)pyrazole-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 14.

NMR (DMSO-d₆, δ) : 6.79(1H, s), 7.20(2H, d, J=8Hz), 7.29(2H, d, J=9Hz), 7.44(2H, d, J=8Hz), 7.50(2H, d, J=9Hz).

IR (KBr) : 1617 cm⁻¹

mass (m/z) : 351(M)

Preparation 53

Sodium 4-[3-chloro-1-(4-fluorophenyl)pyrazole-5-yl]benzenesulfinate was prepared by a similar procedure to that of Preparation 14.

NMR (DMSO-d₆, δ) : 6.79(1H, s), 7.20(2H, d, J=8Hz), 7.24-7.39(4H, m), 7.45(2H, d, J=8Hz).

IR (KBr) : 1654 cm⁻¹

mass (m/z) : 335(M)

Preparation 54

5-(4-Methylphenyl)-1-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazol was prepared by a similar procedure to that of Preparation 10 and Preparation 11.

Preparation 55

Sodium 4-[5-(4-methylphenyl)-3-(trifluoromethyl)pyrazol-1-yl]benzenesulfinate was prepared by a similar procedure to that of Example 19, Example 20 and Preparation 12.

IR (Neat) : 1645, 1565, 1465, 1235, 1135 cm⁻¹

NMR (DMSO-d₆, d) : 2.29(3H, s), 7.12-7.64(9H, m).

Example 1

A mixture containing 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-[(methylsulfinyl)phenyl]pyrazole (2.5 g), sodium acetate (2.0 g) and

acetic anhydride (30 ml) was stirred at reflux temperature for 4 hours. The reaction mixture was cooled and the insoluble materials were removed by filtration. Toluene (50 ml) was added to the filtrate and then the mixture was concentrated under reduced pressure. To the residue was added toluene (50 ml), and again the reaction mixture was evaporated under reduced pressure. The resulting residue was purified by column chromatography on silica gel eluting with dichloromethane to give 5-[(acetoxymethylsulfinyl)phenyl]-3-chloro-1-(3-chloro-4-methoxyphenyl)-pyrazole as a synthetic intermediate. To the intermediate in a mixed solvent of dichloromethane (20 ml) and methanol (10 ml) was added magnesium monoperoxyphthalate (6.8 g) with stirring at an ice-bath temperature. After the addition was completed, the reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture was diluted with dichloromethane (100 ml) and then washed successively with water, aqueous sodium carbonate solution and brine, and dried over MgSO_4 . The evaporation of the solvent gave a gummy oil, which was crystallized in ethanol and collected by filtration to give 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chloro-4-methoxyphenyl)pyrazole (2.4 g) in 80.3% yield. The analytical sample was obtained by recrystallization from ethanol.

IR (KBr) : 1760, 1505, 1320, 1150, 1060 cm^{-1}

NMR (CDCl_3 , δ) : 2.08(3H,s), 3.92(3H,s), 5.16(2H,s), 6.55(1H,s), 6.84-7.92(7H,m)

MASS : 455(M)

Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{Cl}_2\text{N}_2\text{O}_5\text{S}$: C, 50.12 ; H, 3.54 ; N, 6.15

Found : C, 49.51 ; H, 3.41 ; N, 6.08

mp : 139-140°C

Example 2

5-[4-(Acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole was prepared by a similar procedure

to that of Example 1.

IR (KBr) : 1745, 1505, 1270, 1205, 1155, 1125 cm^{-1}

NMR (CDCl_3 , δ) : 2.09(3H), 3.94(3H,s), 5.16(2H,s), 6.87-7.94(8H,m)

MASS : 489(M+1)

mp : 86 - 88°C

Example 3

A mixture of sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate (200 mg) and 3-bromopropanol (82 mg) in N,N-dimethylformamide (0.5 ml) was heated at 100°C for 1 hour. The reaction mixture was poured into water and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent gave a gummy oil, which was purified by column chromatography on silica gel with eluting a mixed solvent of ethyl acetate and n-hexane (2 : 1) to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (65 mg) as a powder.

IR (Neat) : 3480, 1500, 1275, 1230, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 1.88-2.05(3H,m), 3.23-3.79(4H,m), 3.94(3H,s), 6.84-7.93(8H,m)

MASS : 475(M+1)

Example 4

3-Chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3. Purification by column chromatography on silica gel eluting with ethyl acetate followed by trituration in ethanol gave the desired product as a powder.

IR (KBr) : 3505, 1505, 1310, 1270, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.62(1H,t,J=6.0Hz), 3.35-4.07(4H,m), 3.93(3H,s), 6.53(1H,s), 6.84-7.93(7H,m)

MASS : 427(M+1)

Example 5

1-(3-Chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar manner to that of Example 3. After purification by column chromatography eluting with ethyl acetate, the desired product was obtained as a powder.

IR (KBr) : 3440, 1665, 1500, 1275, 1230, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 2.61(1H,t,J=6.2Hz), 3.36-4.67(4H,m), 3.93(3H,s), 6.74-8.02(8H,m)

MASS : 461(M+1)

Example 6

A mixture of sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoro-methyl)pyrazol-5-yl]benzenesulfinate (300 mg) and 2-chloroethyl methylsulfide (101 mg) in N,N-dimethylformamide (0.75 ml) was heated at 100°C in the presence of catalytic amount of potassium iodide. After 1.5 hours, the reaction mixture was poured into water and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent gave a gummy oil, which was triturated with n-hexane to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylthio)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (274 mg) as a powder.

IR (KBr): 1600, 1500, 1270, 1230, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.10(3H,s), 2.70-4.27(4H,m), 3.94(3H,s), 6.82-7.92(8H, m)

MASS : 491(M+1)

Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{ClF}_3\text{N}_2\text{O}_3\text{S}_2$: C, 48.93 ; H, 3.69 ; N, 5.71

Found : C, 49.18 ; H, 3.68 ; N, 5.60

Example 7

A mixture of sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate (200 mg) and 5-chloropentanol (73 mg) in N,N-dimethylformamide (0.5 ml) was heated at 100°C in the presence of potassium iodide as a catalyst. After 4 hours, the reaction mixture was poured into water and extracted with ethyl

acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent gave the residue, which was purified by column chromatography on silica gel eluting with ethyl acetate to afford 1-(3-chloro-4-methoxyphenyl)-5-{4-[(5-hydroxypentyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (186 mg) in 81.2% yield.

IR (Neat) : 3405, 1500, 1270, 1235, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 1.31-1.85(5H,m), 3.08-3.70(6H,m), 3.94(3H,s), 6.82-7.91 (8H,m)

MASS : 503(M+1)

Example 8

A mixture of sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate (200 mg) and 4-chlorobutanol (64 mg) in N,N-dimethylformamide (0.5 ml) was heated at 100°C in the presence of potassium iodide as a catalyst. Additional 4-chlorobutanol and potassium carbonate were added to the reaction mixture. After 4 hours, the reaction mixture was poured into water and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent gave the residue, which was purified by column chromatography on silica gel eluting with ethyl acetate to afford 1-(3-chloro-4-methoxyphenyl)-5-{4-[(4-hydroxybutyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (42 mg) in 18.8% yield.

IR (Neat) : 3430, 1505, 1275, 1235, 1140 cm^{-1}

MASS : 489(M+1)

Example 9

A mixture of sodium 4-[1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazol-5-yl]benzenesulfinate (300 mg) and ethyl bromoacetate (126 mg) in N,N-dimethylformamide (0.8 ml) was heated at 100°C for 1 hour. The reaction mixture was poured into water and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent gave the residue,

which was purified by column chromatography on silica gel eluting with a mixed solvent of ethyl acetate and n-hexane (1:2) to afford 1-(3-chloro-4-methoxyphenyl)-5-[4-(ethoxycarbonylmethylsulfonyl)phenyl]-3-(trifluoromethyl)pyrazole (200 mg) in 58.1% yield.

IR (KBr) : 1740, 1505, 1330, 1275, 1230, 1160 cm^{-1}

NMR (CDCl_3 , δ) : 1.22(3H,t,J=7.0Hz), 3.92(3H,s), 4.11(2H,s), 4.16(2H,q,J=7.0Hz), 6.85-7.96(8H,m)

MASS : 503(M+1)

Example 10

To a stirred solution of 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (100 mg) and triethylamine (26 mg) in dichloromethane (5 ml) was added dropwise a solution of methanesulfonyl chloride (25 mg) in dichloromethane (0.5 ml) at an ice-bath temperature. After 40 minutes the reaction mixture was evaporated under reduced pressure. The residue was diluted with ethyl acetate and then filtered. The filtrate was evaporated under reduced pressure to afford the desired mesylate. To the mesylate in acetonitrile (5 ml) was added 40% aqueous methylamine solution (42 mg) at room temperature and the resulting mixture was stirred overnight. The reaction mixture was evaporated in vacuo and the residue was partitioned between ethyl acetate and water. The organic layer was separated and washed with brine, and dried over MgSO_4 . The obtained residue was purified by column chromatography on silica gel eluting with a mixed solvent of ethyl acetate and methanol (10 : 1) to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methylaminoethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (72 mg) as a powder.

IR (KBr) : 3405, 1605, 1505, 1315, 1275, 1230, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 2.05(1H,broad s), 2.46(3H,s), 3.07(2H,t,J=6.0Hz), 3.38(2H,t,J=6.0Hz), 3.94(3H,s), 6.85-7.93(8H,m)

MASS : 474(M+1)

Example 11

1-(3-Chloro-4-methoxyphenyl)-5-{4-[(2-aminoethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar manner to that of Example 10.

IR (KBr) : 3465, 1500, 1275, 1230, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 3.03(2H,t,J=6.0Hz), 3.25(2H,t,J=6.0Hz), 3.93(3H,s), 6.85-7.91(8H,m)

Example 12

A mixture of 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (70 mg) and acetic anhydride (0.5 ml) was heated at 150°C with stirring. After 2 hours, the reaction mixture was diluted with toluene (20 ml) and evaporated under reduced pressure. The resulting residue was purified by column chromatography on silica gel eluting with a mixed solvent of ethyl acetate and n-hexane (1 : 1) to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (51 mg) as a powder.

NMR (CDCl_3 , δ) : 1.87(3H,s), 3.48(2H,t,J=6.0Hz), 3.94(3H,s), 4.41(2H,t,J=6.0Hz)

MASS : 503(M+1)

Example 13

3-Chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)-sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 12.

IR (KBr) : 1745, 1505, 1380, 1310, 1245, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 1.87(3H,s), 3.47(2H,t,J=6.0Hz), 3.93(3H,s), 4.41(2H,t,J=6.0Hz), 6.52(1H,s), 6.84-7.91(7H,m)

MASS : 469(M-)

mp : 113 - 114°C

Example 14

To a solution of 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylthio)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (150 mg) in

dichloromethane (5 ml) was added 3-chloroperoxybenzoic acid (70%, 201 mg) at room temperature for 2 days. The reaction mixture was partitioned between dichloromethane and water. The organic layer was separated and washed successively with aqueous sodium carbonate solution and brine, dried over MgSO_4 . The evaporation of the solvent gave a gummy oil, which was purified by column chromatography on silica gel eluting a mixed solvent of dichloromethane and methanol (50 : 1), and then crystallized from ethanol to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylsulfonyl)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (100 mg) in 59% yield.

IR (KBr) : 3430, 1560, 1500, 1375, 1315, 1230, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 3.03(3H,d,10.0Hz), 3.36-4.52(4H,m), 3.94(3H,s), 6.87-7.94(8H,m)

MASS : 523(M+1)

Example 15

5-{4-[(Acetoxymethyl)sulfonyl]phenyl}-1-(4-fluorophenyl)-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 1.

IR (Neat) : 1770, 1605, 1335, 1240, 1200, 1155 cm^{-1}

NMR (CDCl_3 , δ) : 2.08(3H,s), 5.17(2H,s), 6.88(1H,s), 7.00-7.95(7H,m)

MASS : 443(M+1)

Example 16

1-(4-Fluorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

IR (Neat) : 3502, 1605, 1315, 1280, 1245, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 2.56(1H,t,J=6.0Hz), 3.36-4.11(4H,m), 6.87(1H,s), 7.03-7.96(7H,m)

MASS : 415(M+1)

Example 17

5-{4-[(Acetoxymethyl)sulfonyl]phenyl}-1-(4-chloro-3-methoxy-

phenyl)-3-cyanopyrazole was prepared by a similar procedure to that of Example 1.

IR (KBr) : 2240, 1765, 1595, 1490, 1330, 1240, 1195, 1145 cm^{-1}

NMR (CDCl_3 , δ) : 2.08(3H,s), 3.86(3H,s), 5.16(2H,s), 6.60-7.95(8H,m)

MASS : 446(M+1)

Example 18

1-(4-Chloro-3-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-cyanopyrazole was prepared by a similar procedure to that of Example 3.

IR (KBr) : 3500, 2240, 1590, 1485, 1305, 1235, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.52(1H,t,J=6.0Hz), 3.36-4.10(4H,m), 3.85(1H,s), 6.61-7.97(8H,m)

MASS : 418(M+1)

Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{ClN}_3\text{O}_4\text{S}$: C, 54.61 ; H, 3.86 ; N, 10.06

Found : C, 54.36 ; H, 3.80 ; N, 9.98

Example 19

A mixture of 1-(3-chlorophenyl)-5-[4-(methylsulfinyl)phenyl]-3-(trifluoromethyl)pyrazole (3.7 g) and sodium acetate (2.9 g) in acetic anhydride (20 ml) was refluxed for 4 hours. The reaction mixture was cooled to room temperature, and insoluble were removed by filtration and washed with toluene. The filtrates were evaporated under reduced pressure. The residue was partitioned between ethyl acetate and water. The organic layer was separated, and washed with saturated sodium carbonate solution and brine, and dried over magnesium sulfate. The evaporation of the solvent gave 5-[4-(acetoxymethylthio)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole (3.6 g).

Example 20

To a solution of 5-[4-(acetoxymethylthio)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole (3.6 g) in methylene chloride (20 ml) and methanol (13 ml) was added magnesium monoperoxyphthalate (7.3 g) at 0-5°C with ice-water bath. After being stirred at

room temperature for 1 hour, the reaction mixture washed successively with saturated sodium carbonate solution and brine dried over magnesium sulfate. After evaporation of the solvent, the residue was triturated with isopropyl alcohol to give 5-[4-(acetoxymethylsulfonyl)-phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole (3.05 g).

Example 21

To a stirred solution of 3-chloro-1-(4-methylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole (2.65 g) in acetic anhydride (19 ml) was added sodium acetate (2.43 g) portionwise. After stirred for 3 hour at 180°C, the reaction mixture was cooled to room temperature and then filtered. The filter cake was washed with toluene (20 ml). The filtrate and the washing were combined and concentrated under reduced pressure. The resulting residue was partitioned between ethyl acetate (30 ml) and water (30 ml). The organic layer was washed successively with aqueous sodium bicarbonate solution and brine, and dried over magnesium sulfate. After evaporation of the solvent, the obtained residue was purified by column chromatography on silicagel eluting with a mixed solvent of ethyl acetate and n-hexane. The fractions containing a desired product were collected and then concentrated under reduced pressure to give 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-methylphenyl)pyrazole (2.89 g).

NMR (DMSO- d_6 , δ); 2.05 (3H, s), 2.33 (3H, s), 5.53 (2H, s), 6.80 (1H, s), 7.15-7.26 (6H, m), 7.42 (2H, d, J=8.5 HZ).

IR (Neat): 3473, 3134, 3035, 2964, 2927, 2873, 1747, 1601 cm^{-1} .

Mass m/e : 373 (M++1).

Example 22

A stirred solution of 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-methylphenyl)pyrazole (2.81 g) in methanol (10 ml) and dichloromethane (10 ml) was cooled with an ice. Monoperoxyphthalic acid magnesium salt (5.95 g) was added portionwise to this solution. After 1 hour the reaction mixture was poured into a mixture of water (20

ml) and dichloromethane (20 ml). The organic layer was separated and washed successively with aqueous sodium bicarbonate solution and brine, and dried over magnesium sulfate. Evaporation of the solvent followed by triturating the resulting residue with isopropyl ether gave 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-methylphenyl)-pyrazole (2.71 g).

NMR (DMSO- d_6 , δ); 2.04 (3H, s), 2.34 (3H, s), 5.44 (2H, s), 6.98 (1H, s), 7.17 (2H, d, $J=8.5$ Hz), 7.25 (2H, d, $J=8.5$ Hz), 7.53 (2H, d, $J=8.5$ Hz, 1.8 Hz) 7.89 (2H, dt, $J=8.5$ Hz, 1.8 Hz)

IR (KBr): 3126, 3072, 3041, 2993, 2929, 1768 cm^{-1} .

Mass m/e : 405 ($M+1$).

Example 23

To a stirred solution of 3-chloro-1-(4-acetylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole (1.0 g) in tetrahydrofuran (50 ml) was dropwise 1.14 N methyllithium diethyl ether solution (5 ml) at an ice-bath temperature under a nitrogen atmosphere. After the addition was completed, the reaction mixture was allowed to react at room temperature overnight. Aqueous ammonium chloride solution was added to the mixture at an ice-bath temperature. Extraction with ethyl acetate followed by washing with brine and evaporation under reduced pressure gave an oil, which was purified by a column chromatography on silica gel eluting with dichloromethane-methanol (50 : 1) to afford the desired alcohol. This alcohol was dissolved in benzene (10ml), and thionyl chloride (0.3 ml) and pyridine (2 ml) were added. The resulting mixture was heated at 150°C with stirring for 1 hour, and then poured into water. Extraction with ethyl acetate followed by washing with aqueous dilute hydrochloric acid solution and brine, and evaporation under reduced pressure gave a crude product, which was subjected to a column chromatography on silica gel eluting with dichloromethane-n-hexane (3 : 1) to give 3-chloro-1-(4-isopropenylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole (346 mg) in 50% yield.

IR (KBr) : 1600, 1510, 1470, 1430, 1365, 1090 cm^{-1}

NMR (CDCl₃, δ) : 2.14(3H, s), 2.48(3H, s), 5.13(1H, m), 5.40(1H, m), 6.40(1H, s), 7.10-7.45(8H, m).

MASS : 341(M⁺+1)

Example 24

5-[[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(isopropenylphenyl)-pyrazole was prepared according to a similar manner to that of Preparation 13 and Example 21.

IR (Neat) : 1745, 1515, 1375, 1210 cm⁻¹

NMR (CDCl₃, δ) : 2.11(3H, s), 2.14(3H, s), 5.13(1H, m), 5.40(1H, m), 5.44(2H, s), 6.43(1H, s), 7.16-7.46(8H, m).

MASS : 399(M⁺+1)

Example 25

3-Chloro-1-(4-isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-phenyl}pyrazole was prepared by a similar procedure to that of Example 3.
mp : 158-161°C

IR (KBr) : 1510, 1375, 1305, 1150, 1110 cm⁻¹

NMR (CDCl₃, δ) : 2.15(3H, s), 3.21(3H, s), 3.39(1H, t, J=6.0Hz), 3.76(2H, t, J=6.0Hz), 5.15(1H, m), , 5.40(1H, m), 6.53(1H, s), 7.17-7.89(8H, m).

MASS : 417(M⁺+1)

Anal. Calcd for C₂₁H₂₁ClN₂O₃S 1/3H₂O : C, 59.64; H, 5.16; N, 6.62.

Found : C, 59.72; H, 4.98; N, 6.60.

Example 26

3-Chloro-1-(4-isopropenylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 122-125°C

IR (KBr) : 3465, 3435, 1510, 1370, 1305, 1275, 1235, 1130 cm⁻¹

NMR (CDCl₃, δ) : 2.15(3H, s), 3.35-4.19(4H, m), 5.16(1H, m), 5.41(1H, m), 6.51-7.91(9H, m).

MASS : 403(M⁺⁺+1)

Anal Calcd for C₂₀H₂₁ClN₂O₃S : C, 58.74; H, 4.85; N, 6.85.

Found : C, 58.87; H, 4.69; N, 6.80.

Example 27

1-(4-Methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 102-104°C

IR (KBr) : 3480, 1510, 1465, 1280, 1235, 1155, 1120 cm^{-1}

NMR (CDCl_3 , δ) : 2.40(3H, s), 2.63(1H, t, J=6.0Hz), 3.34-4.19(4H, m), 6.82-7.91(9H, m).

MASS : 411($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}$: C, 55.61; H, 4.17; N, 6.82.

Found : C, 55.06 ; H, 4.09 ; N, 6.71.

Example 28

1-(4-Methylphenyl)-5-{4-[(3-acetoxypropyl)sulfonyl] phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 95-96°C

IR (KBr) : 1735, 1285, 1235, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 2.03-2.16(2H, m), 2.08(3H, s), 2.40(3H, s), 3.14-3.22(2H, m), 4.13(2H, t, J=6.0Hz), 6.85(1H, s), 7.13-7.89(8H, m).

MASS : 467($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{F}_3\text{N}_2\text{O}_4\text{S}$: C, 56.56; H, 4.54; N, 6.00

Found : C, 56.73; H, 4.56; N, 5.94.

Example 29

1-(4-Methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 96-98°C

IR (KBr) : 3545, 3480, 1605, 1525, 1315, 1270, 1235, 1165, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.61(1H, t, J=6.0Hz), 3.37(2H, t, J=6.0Hz), 3.84(3H, s), 4.03(2H, q, J=6.0Hz), 6.85-7.91(9H, m).

MASS : 427(M⁺+1)

Anal. Calcd for C₁₉H₁₇F₃N₂O₄S : C, 53.52; H, 4.02; N, 6.57.

Found : C, 53.28 ; H, 3.99 ; N, 6.47.

Example 30

5-[4-(Acetoxymethylthio)phenyl]-3-trifluoromethyl-1-(3-chloro-4-methylphenyl)pyrazole was prepared by a similar method to that of Preparation 10, Preparation 11 and Example 19.

IR (KBr) : 1745, 1500, 1465, 1230, 1210, 1160, 1135 cm⁻¹

NMR (CDCl₃, δ) : 2.11(3H, s), 2.40(3H, s), 5.45(2H, s), 6.74(1H, s), 6.99-7.44(7H, m).

MASS : 441(M⁺+1)

Example 31

1-(3-Chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

IR (KBr) : 3495, 1500, 1315, 1275, 1235, 1135 cm⁻¹

NMR (CDCl₃, δ) : 2.41(3H, s), 2.59(1H, t, J=6.0Hz), 3.38(2H, t, J=6.0Hz), 4.04(2H, q, J=6.0Hz), 6.85(1H, s), 6.97-7.95(7H, m).

MASS : 445(M⁺+1)

Example 32

1-(3-Chloro-4-methylphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 12 (S9803378).

mp : 105-108°C

IR (KBr) : 1740, 1500, 1465, 1395, 1325, 1235, 1150 cm⁻¹

NMR (CDCl₃, δ) : 1.87(3H, s), 2.41(3H, s), 3.48(2H, t, J=6.0Hz), 4.18(2H, t, J=6.0Hz), 6.82-7.93(8H, m).

MASS : 487(M⁺+1)

Anal. Calcd for C₂₁H₁₈ClF₃N₂O₄S : C, 51.80 ; H, 3.73 ; N, 5.75.

Found : C, 51.94 ; H, 3.70 ; N, 5.67.

Example 33

1-(4-Fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 80-83°C

IR (KBr) : 3480, 1500, 1310, 1280, 1230, 1115 cm⁻¹

NMR (CDCl₃, δ) : 2.28(3H, s), 2.58(1H, t, J=6.0Hz), 3.37(2H, t, J=6.0Hz), 4.03(2H, q, J=6.0Hz), 6.85(1H, s), 6.97-7.94(7H, m).

MASS : 429(M⁺+1)

Anal. Calcd for C₁₉H₁₆F₄N₂O₃S : C, 53.27 ; H, 3.77 ; N, 6.54.

Found : C, 53.03 ; H, 3.70 ; N, 6.53.

Example 34

1-(4-Fluoro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 93-94°C

IR (KBr) : 1500, 1400, 1310, 1275, 1225, 1160 cm⁻¹

NMR (CDCl₃, δ) : 2.28(3H, s), 3.21(3H, s), 3.39(2H, t, J=6.0Hz), 3.76(2H, t, J=6.0Hz), 6.85(1H, s), 6.96-7.91(7H, m).

MASS : 443(M⁺+1) Anal. Calcd for C₂₀H₁₈F₄N₂O₃S : C, 54.30; H, 4.10; N, 6.33.

Found : C, 53.62 ; H, 3.93 ; N, 6.28.

Example 35

1-(3-Methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 89-90°C

IR (KBr) : 1605, 1465, 1310, 1280, 1160, 1135 cm⁻¹

NMR (CDCl₃, δ) : 2.36(3H, s), 3.20(3H, s), 3.39 (2H, t, J=6.0Hz), 3.75(2H, t, J=6.0Hz), 6.85(1H, s), 6.95-7.90(8H, m).

MASS : 425(M⁺+1)

Anal. Calcd for C₂₀H₁₉F₃N₂O₃S 1/3H₂O : C, 55.81 ; H, 4.61 ; N, 6.51.

Found : C, 55.89 ; H, 4.46 ; N, 6.63.

Example 36

1-(3-Methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 92-93°C

IR (KBr) : 3475, 1510, 1465, 1310, 1280, 1240, 1160, 1115 cm^{-1}

NMR (CDCl_3 , δ) : 2.37(3H, s), 2.60(1H, t, J=6.0Hz), 3.34-3.39(2H, m), 3.99-4.07(2H, m), 6.86(1H, s), 6.93-7.92(8H, m).

MASS : 411($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}$: C, 55.61 ; H, 4.17; N, 6.82.

Found : C, 55.32 ; H, 4.16 ; N, 6.69.

Example 37

1-(3-Fluoro-4-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 136-137°C

IR (KBr) : 1595, 1505, 1460, 1400, 1310, 1280, 1245, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.30(3H, s), 3.21(3H, s), 3.40(2H, t, J=6.0Hz), 3.77(2H, t, J=6.0Hz), 6.84(1H, s), 6.90-7.92(7H, m).

MASS : 443($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{F}_4\text{N}_2\text{O}_3\text{S}$: C, 54.30 ; H, 4.10 ; N, 6.33.

Found : C, 54.38 ; H, 4.03 ; N, 6.31.

Example 38

1-(3-Fluoro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 100-101°C

IR (KBr) : 3480, 1595, 1470, 1415, 1310, 1280, 1245, 1160, 1120 cm^{-1}

NMR (CDCl_3 , δ) : 2.31(3H, s), 2.60(2H, t, J=6.0Hz), 3.38(2H, t, J=6.0Hz), 4.04(2H, q, J=6.0Hz), 6.85(1H, s), , 6.91-7.95(7H, m).

MASS : 429($\text{M}^+ + 1$)

Anal. Calcd for $C_{19}H_{16}F_4N_2O_3S$: C, 3.27 ; H, 3.77 ; N, 6.54.

Found : C, 53.32 ; H, 3.80 ; N, 6.44.

Example 39

5-{ 4-[(Acetoxymethyl)sulfinyl]-3-fluorophenyl}-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of preparation 11 and Example 19.

IR (Neat) : 1750, 1500, 1270, 1220, 1150, 1135 cm^{-1}

NMR ($CDCl_3$, δ) : 2.10(3H, s), 3.94(3H, s), 5.41(2H, s), 6.76-7.51(7H, m).

MASS : 475(M^+)

Example 40

1-(3-Chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-3-fluorophenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

IR (KBr) : 3435, 1615, 1505, 1270, 1130 cm^{-1}

NMR ($CDCl_3$, δ) : 2.43(1H, t, $J=6.0Hz$), 3.52-4.25(4H, m), 3.95(3H, s), 6.83-7.937(7H, m).

MASS : 479(M^++1)

Example 41

1-(4-Chloro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 132-133°C

IR (KBr) : 1475, 1405, 1305, 1275, 1240, 1160, 1140 cm^{-1}

NMR ($CDCl_3$, δ) : 2.38(3H, s), 3.21(3H, s), 3.40(2H, t, $J=6.0Hz$), 3.77(2H, t, $J=6.0Hz$), 6.85(1H, s), 6.90-7.92(7H, m).

MASS : 459(M^++1)

Anal. Calcd for $C_{20}H_{18}ClF_3N_2O_3S$: C, 52.35; H, 3.95; N, 6.10.

Found : C, 51.62 ; H, 3.84 ; N, 6.04.

Example 42

1-(4-Isopropenylphenyl)-5-[4-(acetoxymethylthio)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of

Preparation 11 and Example 19.

IR (Neat) : 1750, 1460, 1375, 1230, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 2.12(3H, s), 2.15(3H, s), 5.15(1H, m), 5.40(1H, m),
5.44(2H, s), 6.75(1H, s), 7.17-7.49(8H, m).

MASS : 455(M^{+23})

Example 43

1-(4-Isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 141-143°C

IR (KBr) : 1510, 1305, 1280, 1240, 1140 cm^{-1}

NMR (CDCl_3 , δ) : 2.15(3H, s), 3.21(3H, s), 3.40(2H, t, $J=6.0\text{Hz}$), 3.76(2H, t, $J=6.0\text{Hz}$), 5.17(1H, m), 5.41(1H, m), 6.85(1H, s), 7.22-7.91(8H, m).

MASS : 451(M^{+1})

Example 44

1-(3-Methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 115-116°C

IR (KBr) : 3415, 1600, 1315, 1250, 1220, 1170, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.60(2H, t, $J=6.0\text{Hz}$), 3.37(2H, t, $J=6.0\text{Hz}$), 3.78(3H, s),
4.03(2H, q, $J=6.0\text{Hz}$), 6.76-7.93(9H, m).

MASS : 427(M^{+1})

Anal. Calcd for $\text{C}_{19}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_4\text{S}$: C, 53.52 ; H, 4.02 ; N, 6.57.

Found : C, 53.32 ; H, 4.08 ; N, 6.54.

Example 45

1-(3-Chlorophenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 59-61°C

IR (KBr) : 1590, 1475, 1440, 1400, 1370, 1280, 1240, 1145 cm^{-1}

NMR (CDCl₃, δ) : 3.21(3H, s), 3.40(2H, t, J=6.0Hz), 3.76(2H, t, J=6.0Hz), 6.86(1H, s), 7.06-7.94(8H, m).

MASS : 445(M⁺+1)

Anal. Calcd for C₁₉H₁₆ClF₃N₂O₃S : C, 51.30 ; H, 3.63 ; N, 6.30.

Found : C, 51.14 ; H, 3.49 ; N, 6.26.

Example 46

1-(4-Isopropylphenyl)-5-[4-(acetoxymethylthio)phenyl]-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Preparation 10 and Example 19.

IR (Neat) : 1750, 1510, 1465, 1375, 1230, 1135 cm⁻¹

NMR (CDCl₃, δ) : 1.25(6H, d, J=7.0Hz), 2.87-3.00(1H, m), 5.44(2H, s), 6.74(1H, s), 7.16-7.40(8H, m).

MASS : 457(M⁺+23)

Example 47

1-(4-Isopropylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 147-150°C

IR (KBr) : 1510, 1465, 1305, 1280, 1240, 1140 cm⁻¹

NMR (CDCl₃, δ) : 1.25(6H, d, J=7.0Hz), 2.84-2.98(1H, m), 3.20(3H, s), 3.39(2H, t, J=6.0Hz), 3.76(2H, t, J=6.0Hz), 6.84-7.90(9H, m).

MASS : 453(M⁺+1)

Example 48

5-[4-(Acetoxymethylsulfonyl)phenyl]-1-(3-cyano-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 19 and Example 20.

IR (KBr) : 2230, 1750, 1505, 1290, 1225, 1155, 1140 cm⁻¹

NMR (CDCl₃, δ) : 2.17(3H, s), 3.98(3H, s), 5.00(2H, ABq, J=10.0Hz), 6.84(1H, s), 6.96-7.73(7H, m).

Example 49

1-(3-Cyano-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-

phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

mp : 142-143°C

IR (KBr) : 3510, 2230, 1505, 1285, 1235, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.57(1H, t, J=6.0Hz), 3.99(3H, s), 4.03(2H, t, J=6.0Hz), 6.87(1H, s), 6.98-7.98(7H, m).

MASS : 474(M^+ +23)

Anal. Calcd for $\text{C}_{20}\text{H}_{16}\text{F}_3\text{N}_3\text{O}_4\text{S}$: C, 53.21 ; H, 3.57; N, 9.31.

Found : C, 52.68 ; H, 3.56 ; N, 9.14.

Example 50

To a stirred suspension of NaH (about 60%, 12 mg) in N, N-dimethylformamide (0.3 ml) was added a solution of the 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole (147 mg) in N, N-dimethylformamide (0.4 ml) at an ice bath temperature. After 15 minutes methyl iodide (49mg) in N, N-dimethylformamide (0.3 ml) was added to this mixture at the same temperature. After 15 minutes, the reaction mixture was allowed to react at room temperature for 30 minutes. The mixture was diluted with water and then neutralized by addition of aqueous HCl dilute solution, and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of solvent under reduced pressure gave a gummy oil, which was subjected to a column chromatography on silica gel eluting with a mixed solvent of ethyl acetate and n-hexane (2 : 1). Fractions containing the desired product were collected and concentrated under reduced pressure to give 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-methoxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (73 mg).

IR (Neat) : 1505, 1270, 1235, 1145 cm^{-1}

NMR (CDCl_3 , δ) : 1.95-2.04(2H, m), 3.18-3.47(4H, m), 3.28(3H, s), 3.94(3H, s), 6.84-7.96(8H, m).

Example 51

A mixture of 1-(3-chloro-4-methoxyphenyl)-5-{4-[(ethoxycarbonyl)-

methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (200 mg), hydroxyamine hydrochloric acid (30 mg) and sodium bicarbonate (55 mg) in ethanol (5 ml) was refluxed with stirring for 24 hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was diluted with water and extracted with ethyl acetate. The extract was washed with brine and dried over MgSO_4 . The evaporation of the solvent followed by column chromatography on silica gel eluting with ethyl acetate gave 1-(3-chloro-4-methoxyphenyl)-5-{4-[(N-hydroxycarbamoyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (48 mg).

IR (KBr) : 1740, 1505, 1315, 1275, 1230, 1160 cm^{-1}

NMR (CDCl_3 , δ) : 3.91(3H, s), 4.07(2H, s), 6.86-8.21(8H, m).

MASS : 490(M+1)

Example 52

A mixture of 1-(3-chloro-4-methoxyphenyl)-5-{4-[(ethoxycarbonyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (250 mg) and 40% aqueous methylamine solution (0.5 ml) in ethanol (5 ml) was heated at reflux temperature for 8 hours. The reaction mixture was concentrated under reduced pressure, and the resulting residue was purified by column chromatography on silica gel using a mixed solvent of dichloromethane and methanol (20 : 1) as an eluant, giving 1-(3-chloro-4-methoxyphenyl)-5-{4-[(methylcarbonyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole (123 mg).

IR (KBr) : 3380, 1670, 1505, 1270, 1230, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.86(3H, d, J=5.0Hz), 3.93(3H, s), 4.01(2H, s), 6.58(1H, br s), 6.85-7.90(8H, m).

MASS : 488(M+1)

Example 53

1-(3-Chloro-4-methoxyphenyl)-5-{4-[(2-oxopropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 7.

IR (KBr) : 1725, 1505, 1325, 1270, 1235, 1160 cm^{-1}

NMR (CDCl₃, δ) : 2.42(3H, s), 3.94(3H, s), 4.17(2H, s), 6.86-7.89(8H, m).

MASS : 473(M+1)

Example 54

5-{4-[(3-Acetoxypropyl)sulfonyl]phenyl}-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3 and Example 12.

IR (KBr) : 1740, 1505, 1260, 1235, 1135 cm⁻¹

NMR (CDCl₃, δ) : 2.05-2.15(2H, m), 2.09(3H, s), 3.15-3.23(2H, m), 3.94(3H, s), 4.13(2H, t, J=6.0Hz), 6.85-7.94(8H, m).

MASS : 539(M+23) API-ES

Anal. Calcd for C₂₂H₂₀ClF₃N₂O₅S : C, 51.12 ; H, 3.90 ; N, 5.42.

Found : C, 50.78 ; H, 3.87 ; N, 5.42.

Example 55

1-(3-Chloro-4-methoxyphenyl)-5-{4-[[3-(dimethylamino)propyl]-sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole hydrochloride was prepared by a similar procedure to that of Example 10.

mp : 213-215°C

IR (KBr) : 2615, 1505, 1280, 1240, 1160 cm⁻¹

NMR (DMSO-d₆, δ) : 1.95-2.06(2H, m), 2.70(6H, s), 3.07-3.18(2H, m), 3.46-3.54(2H, m), 4.01(3H, s), 7.20-7.97(8H, m)

MASS : 502(M⁺+1, free)

Anal. Calcd for C₂₂H₂₃ClF₃N₃O₃S · HCl : C, 49.08; H, 4.49 ; N, 7.80.

Found : C, 48.83 ; H, 4.46 ; N, 7.69.

Example 56

1-(3-Chloro-4-methoxyphenyl)-5-{4-[[2-(dimethylamino)ethyl]sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole hydrochloride was prepared by a similar procedure to that of Example 10.

mp : 211-212°C

IR (KBr) : 2560, 2505, 2420, 1505, 1265, 1230, 1155 cm⁻¹

NMR (DMSO-d₆, δ) 2.74(6H, s), 3.34-3.40(2H, m), 3.91(3H, s), 3.91-4.00(2H, m), 7.20-7.99(8H, m)

MASS : 488(M⁺+1, free)

Anal. Calcd for C₂₁H₂₁ClF₃N₃O₃S · HCl : C, 48.10 ; H, 4.23 ; N, 8.01.

Found : C, 47.50 ; H, 4.20 ; N, 7.84.

Example 57

5-[4-(Acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(difluoromethyl)pyrazole was prepared according to a similar manner to that of Preparation 10, Preparation 11, Example 19 and Example 20.

IR (KBr) : 1765, 1500, 1330, 1265, 1230, 1150 cm⁻¹

NMR (CDCl₃, δ) : 2.09(3H, s), 3.94(3H, s), 5.16(2H, s), 6.50-7.93(9H, m).

MASS : 471(M⁺+1)

Example 58

1-(3-Chloro-4-methoxyphenyl)-3-difluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 120-122°C

IR (KBr) : 3465, 3440, 1505, 1455, 1410, 1300, 1290, 1265, 1145 cm⁻¹

NMR (CDCl₃, δ) : 1.63(1H, t, J=6.0Hz), 1.93-2.07(2H, m), 3.23-3.31(2H, m), 3.76(2H, q, J=6.0Hz), 3.94(3H, s), 6.52-7.92(9H, m).

MASS : 457(M⁺)

Anal. Calcd for C₂₀H₁₉ClF₂N₂O₄S : C, 52.58 ; H, 4.19 ; N, 6.13.

Found : C, 52.66 ; H, 4.26 ; N, 5.99.

Example 59

1-(3-Chloro-4-methoxyphenyl)-5-{4-[[2-(2-hydroxyethoxy)ethyl]-sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 7.

IR (Neat) : 3495, 1505, 1315, 1270, 1235, 1140 cm⁻¹

NMR (CDCl₃, δ) : 2.89-4.30(9H, m), 3.94(3H, s), 6.81-7.94(8H, m).

MASS : 505(M⁺+1)

Example 60

The key intermediate, 1-(4-bromo-3-methoxyphenyl)-5-(4-methylthiophenyl)-3-(trifluoromethyl)pyrazole, was prepared by refluxing a

mixture of 1, 1, 1-trifluoro-4-[4-(methylthio)phenyl]-2, 4-dioxobutane and 4-bromo-3-methoxyphenylhydrazine hydrochloride in acetic acid, which was converted to the desired product, 5-[4-(acetoxymethylthio)phenyl]-1-(4-bromo-3-methoxyphenyl)-3-(trifluoromethyl)pyrazole, according to a similar manner to that of Preparation 11 and Example 19.

IR (Neat) : 1750, 1595, 1475, 1410, 1255, 1220 cm^{-1}

NMR (CDCl_3 , δ): 2.12(3H, s), 3.81(3H, s), 5.45(2H, s), 6.67-7.52(8H, m).

MASS : 501(M^+)

Example 61

1-(4-Bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 123-124°C

IR (KBr) : 3505, 1595, 1405, 1280, 1225, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.55(1H, t, J=6.0Hz), 3.39(2H, t, J=6.0Hz), 3.84(3H, s), 4.07(2H, q, J=6.0Hz), 6.59-7.97(8H, m).

MASS : 505($\text{M}^+ + 2$)

Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{BrF}_3\text{N}_2\text{O}_4\text{S}$: C, 45.16 ; H, 3.19 ; N, 5.54.

Found : C, 45.15 ; H, 3.18 ; N, 5.43.

Example 62

1-(4-Bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 61-65°C

IR (KBr) : 3430, 1595, 1475, 1405, 1310, 1255, 1225, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 1.62(1H, t, J=6.0Hz), 1.95-2.08(2H, s), 3.23-3.31(2H, m), 3.77(2H, q, J=6.0Hz), 3.91(3H, s), 6.59-7.94(8H, m).

MASS : 519(M^+)

Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{BrF}_3\text{N}_2\text{O}_4\text{S}$: C, 46.25 ; H, 3.49 ; N, 5.39.

Found : C, 46.85 ; H, 3.89 ; N, 5.05.

Example 63

The key intermediate, 1-(4-methoxyphenyl)-5-(4-methylthiophenyl)-3-difluoromethylpyrazole, was prepared by refluxing a mixture of 1, 1-difluoro-4-(4-methylthiophenyl)-2, 4-dioxobutane and 4-methoxyphenylhydrazine hydrochloride in acetic acid, which was converted to the desired product, 5-[4-(acetoxymethylsulfonyl)phenyl]-3-difluoromethyl-1-[4-(methoxyphenyl)pyrazole, according to a similar manner to that of Preparation 11, Example 19 and Example 20.

IR (KBr) : 1755, 1605, 1515, 1325, 1245, 1160 cm^{-1}

NMR (CDCl_3 , δ) : 3.84(3H, s), 5.15(2H, s), 6.51-7.89(9H, m).

MASS : 437($\text{M}^+ + 1$)

Example 64

1-(4-Methoxyphenyl)-3-difluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 80-81 $^{\circ}\text{C}$

IR (KBr) : 3470, 1515, 1305, 1240, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.64(1H, t, $J=6.0\text{Hz}$), 3.37(2H, t, $J=6.0\text{Hz}$), 3.84(3H, s), 4.03(2H, q, $J=6.0\text{Hz}$), 6.51-7.90(9H, m).

MASS : 409($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{19}\text{H}_{18}\text{F}_2\text{N}_2\text{O}_4\text{S}$: C, 55.88 ; H, 4.44 ; N, 6.86.

Found : C, 55.28; H, 4.37; N, 6.72.

Example 65

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-bromophenyl)pyrazole was prepared by a similar manner to that of Preparation 13, Example 19 and Example 20.

IR (KBr) : 1760, 1585, 1480, 1370, 1330, 1205 cm^{-1}

NMR (CDCl_3 , δ) : 2.09(3H, s), 5.16(2H, s), 6.56(1H, s), 7.01-7.93(8H, m).

MASS : 469(M^+)

Example 66

1-(3-Bromophenyl)-3-chloro-5-{4-(3-hydroxypropyl)sulfonyl}phenyl}pyrazole was prepared by a similar manner to that of Example 3.

IR (Neat) : 3435, 1585, 1480, 1370, 1310, 1145 cm^{-1}

NMR (CDCl_3 , δ) : 1.63(1H, br s), 1.93-2.07(2H, m), 3.23-3.31(2H, m), 3.50-3.76(2H, m), 6.54(1H, s), 7.05-7.92(8H, m).

MASS : 519($\text{M}^+ + 2$)

Example 67

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-fluoro-4-methylphenyl)pyrazole was prepared by a similar procedure to that of Preparation 13, Example 21 and Example 22.

IR (KBr) : 1770, 1595, 1505, 1435, 1370, 1320, 1195 cm^{-1}

NMR (CDCl_3 , δ) : 2.09(3H, s), 2.29(3H, s), 5.30(2H, s), 6.54(1H, s), 6.83-7.92(7H, m).

MASS : 423($\text{M}^+ + 1$)

Example 68

3-Chloro-1-(3-fluoro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 126-128°C

IR (KBr) : 1590, 1510, 1370, 1300, 1270, 1240, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 1.64(1H, t, $J=6.0\text{Hz}$), 1.94-2.08(2H, m), 3.23-3.31(2H, m), 3.75(2H, q, $J=6.0\text{Hz}$), 6.52(1H, s), 6.84-7.91(7H, m).

MASS : 409($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{19}\text{H}_{18}\text{ClFN}_2\text{O}_3\text{S}$: C, 55.81 ; H, 4.44 ; N, 6.85.

Found : C, 56.00 ; H, 4.48 ; N, 6.55.

Example 69

5-[(4-Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluoro-3-methylphenyl)pyrazole was prepared by a similar manner to that of Preparation 13, Example 21 and Example 22.

IR (KBr) : 3135, 1765, 1595, 1500, 1375, 1335, 1240, 1200 cm^{-1}

NMR (CDCl_3 , δ) : 2.07(3H, s), 2.26(3H, s), 5.15(2H, s), 6.55(1H, s), 6.90-7.90(7H, m).

MASS : 423($\text{M}^+ + 1$)

Example 70

3-Chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 115-117°C

IR (KBr) : 3465, 1500, 1370, 1310, 1280, 1230, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 2.26(3H, s), 2.60(1H, t, J=6.0Hz), 3.37(2H, t, J=6.0Hz), 4.03(2H, t, J=6.0Hz), 6.54(1H, s), 6.91-7.91(7H, m).

MASS : 395($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{ClFN}_2\text{O}_3\text{S}$: C, 54.75 ; H, 4.09 ; N, 7.09.

Found : C, 54.51 ; H, 4.01 ; N, 6.98.

Example 71

3-Chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 130-132°C

IR (KBr) : 3505, 3405, 1500, 1375, 1305, 1270, 1230, 1130 cm^{-1}

NMR (CDCl_3 , δ) : 1.63(1H, t, J=6.0Hz), 1.93-2.07(2H, m), 2.26(3H, s), 3.22-3.29(2H, m), 3.76(2H, q, J=6.0Hz), 6.53(1H, s), 6.91-7.89(7H, m).

MASS : 409($\text{M}^+ + 1$)

Anal. Calcd for $\text{C}_{19}\text{H}_{18}\text{ClFN}_2\text{O}_3\text{S}$: C, 55.81 ; H, 4.44 ; N, 6.85.

Found : C, 55.49 ; H, 4.39 ; N, 6.69.

Example 72

3-Chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.

mp : 132-134°C

IR (KBr) : 3405, 1505, 1370, 1300, 1270, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 1.64(1H, t, J=5.0Hz), 1.93-2.06(2H, m), 3.22-3.30(2H, m), 3.75(2H, q, J=6.0Hz), 3.93(3H, s), 6.52(1H, s), 6.84-7.92(7H, m).

MASS : 441(M^+)

Anal. Calcd for $\text{C}_{19}\text{H}_{18}\text{Cl}_2\text{N}_2\text{O}_4\text{S}$: C, 51.71 ; H, 4.11 ; N, 6.35.

Found : C, 51.68 ; H, 4.17 ; N, 6.11.

Example 73

1-(3-Chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar procedure to that of Example 3.

IR (Neat) : 3460, 1590, 1475, 1280, 1235, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 3.38(2H, t, J=6.0Hz), 3.78(1H, t, J=6.0Hz), 4.01-4.19(2H, m), 6.84-7.96(9H, m).

MASS : 431($\text{M}^+ + 1$)

Example 74

3-Chloro-1-(4-isopropenylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole was prepared by a similar manner to that of Example 3.
mp : 143-145°C

IR (KBr) : 3405, 1595, 1375, 1235, 1135 cm^{-1}

NMR (CDCl_3 , δ) : 0.79-2.05(6H, m), 3.22-3.78(6H, m), 6.54(1H, s), 7.18-7.90(8H, m).

MASS : 417($\text{M}^+ + 1$)

Example 75

3-Chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 153-154°C

IR (KBr) : 1600, 1505, 1300, 1270, 1150 cm^{-1}

NMR (CDCl_3 , δ) : 3.21(3H, s), 3.39(2H, t, J=6.0Hz), 3.75(2H, t, J=6.0Hz), 3.92(3H, s), 6.49(1H, s), 6.83-7.90(7H, m)

MASS : 441(M^+)

Anal. Calcd for $\text{C}_{19}\text{H}_{18}\text{Cl}_2\text{N}_2\text{O}_4\text{S}$: C, 51.71 ; H, 4.11 ; N, 6.35.

Found : C, 51.14 ; H, 3.95 ; N, 6.22.

Example 76

5-[4-(4-Acetoxyethylthio)phenyl]-3-chloro-1-(4-chloro-3-methylphenyl)pyrazole was prepared by a similar procedure to that of

Preparation 13 and Example 21.

NMR (CDCl₃, δ) : 2.11(3H, s), 2.35(3H, s), 5.44(2H, s), 6.42(1H, s), 6.89(1H, dd, J=8.3 and 2.4Hz), 7.13-7.41(6H, m).

Example 77

3-Chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 101-103°C

NMR (CDCl₃, δ) : 2.36(3H, s), 2.60(1H, t, J=6.4Hz), 3.37(2H, m), 4.02(2H, m), 6.54(1H, s), 6.87(1H, dd, J=8.6 and 2.4Hz), 7.27(2H, m), 7.43(2H, d, J=8.6Hz), 7.90(2H, d, J=8.6Hz).

MASS : 411(M⁺)

Anal. Calcd for C₁₈H₁₆Cl₂N₂O₃S : C, 52.56 ; H, 3.92 ; N, 6.81.

Found : C, 52.52 ; H, 3.85 ; N, 6.75.

Example 78

3-Chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 113-115°C

NMR (CDCl₃, δ) : 1.62(1H, t, J=5.2Hz), 1.97(2H, m), 2.35(3H, s), 3.25(2H, m), 3.75(2H, m), 6.53(1H, s), 6.87(1H, dd, J=8.6 and 2.4Hz), 7.28(2H, m), 7.41(2H, d, J=8.4Hz), 7.89(2H, d, J=8.4Hz).

MASS : 425(M⁺)

Example 79

3-Chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 105-106°C

NMR (CDCl₃, δ) : 1.67(1H, t, J=5.2Hz), 1.99(2H, m), 2.39(3H, s), 3.27(2H, m), 3.75(2H, q, J=5.7Hz), 6.53(1H, s), 6.94(1H, dd, J=8.0 and 2.2Hz), 7.18(1H, d, J=8.0Hz), 7.35(1H, d, J=2.2Hz), 7.41(2H, d, J=8.6Hz), 7.89(2H,

d, J=8.6Hz).

MASS : 425(M⁺)

Anal. Calcd for C₁₉H₁₈Cl₂N₂O₃S : C, 53.65 ; H, 4.27 ; N, 6.59.

Found : C, 53.60 ; H, 4.21 ; N, 6.51.

Example 80

3-Chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}pyrazole was prepared by a similar procedure to that of Example 3.

mp : 108-110°C

NMR (CDCl₃, δ) : 2.39(3H, s), 2.62(1H, t, J=6.4Hz), 3.37(2H, m), 4.04(2H, m), 6.53(1H, s), 6.94(1H, dd, J=8.0 and 2.2Hz), 7.23(1H, d, J=8.0Hz), 7.34(1H, d, J=2.2Hz), 7.42(2H, d, J=8.6Hz), 7.90(2H, d, J=8.6Hz).

MASS : 413(M⁺+2)

Anal. Calcd for C₁₈H₁₆Cl₂N₂O₃S : C, 52.56; H, 3.92; N, 6.81.

Found : C, 51.92 ; H, 3.85 ; N, 6.62.

Example 81

3-Chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-(4-methylphenyl)pyrazole was obtained from 4-[3-chloro-1-(4-methylphenyl)pyrazol-5-yl]benzenesulfinate in the similar manner that of Example 3.

NMR (DMSO-d₆, δ) : 1.60-1.75(2H, m), 2.34(3H, s), 3.28-3.45(4H, m), 4.64(1H, t, J=5.3Hz), 6.96(1H, s), 7.18(2H, d, J=8.5Hz), 7.25(2H, d, J=8.5Hz), 7.51(2H, d, J=8.5Hz), 7.87(2H, d, J=8.5Hz).

IR (KBr) : 3512, 3400, 3128, 2964, 2924, 2881, 1601 cm⁻¹.

Mass m/e : 391 (M⁺+1).

Example 82

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole was obtained from 3-chloro-1-(3-methylphenyl)-5-[4-(methylsulfinyl)phenyl]pyrazole in a similar manner that of Example 21.

NMR (DMSO-d₆, δ) : 2.04(3H, s), 2.31(3H, s), 5.53(2H, s), 6.82(1H, s), 6.97-7.01(1H, m), 7.20-7.34(5H, m), 7.42(2H, d, J=8.4Hz).

IR (KBr) : 3654, 3118, 3047, 2958, 2924, 2868, 1738, 1603 cm^{-1} .

Mass m/e : 373 ($M^+ + 1$).

Example 83

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-methylphenyl)-pyrazole was obtained from 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole in the similar manner that of Example 22.

NMR (DMSO- d_6 , δ) : 2.03(3H, s), 2.31(3H, s), 5.44(2H, s), 6.98-7.02(1H, m), 7.00(1H, s), 7.21-7.34(3H, m), 7.54(2H, d, $J=8.4\text{Hz}$), 7.90(2H, d, $J=8.4\text{Hz}$).

IR (KBr) : 3516, 3134, 3082, 3008, 2945, 1765 cm^{-1} .

Mass m/e : 405 ($M^+ + 1$).

Example 84

3-Chloro-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-1-(3-methylphenyl)-pyrazole was obtained from sodium 4-[3-chloro-1-(3-methylphenyl)-pyrazol-5-yl]benzenesulfinate in the similar manner that of Example 3.

NMR (DMSO- d_6 , δ) : 2.30(3H, s), 3.48(2H, t, $J=6.1\text{Hz}$), 3.62-3.71(2H, m), 4.88(1H, t, $J=5.3\text{Hz}$), 6.97(1H, s), 6.95-7.03(1H, m), 7.20-7.34(3H, m), 7.49(2H, d, $J=8.4\text{Hz}$), 7.88(2H, d, $J=8.4\text{Hz}$).

IR (KBr) : 3464, 3406, 3132, 3055, 3960, 2916, 1601 cm^{-1} .

Mass m/e : 377 ($M^+ + 1$).

Example 85

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-phenylpyrazole was prepared by a similar method to that of Example 21.
yellow oil.

IR (Neat) : 1751, 1738, 1598, 1490 cm^{-1} .

NMR (DMSO- d_6 , δ) : 2.06(3H, s), 5.53(2H, s), 6.83(1H, s), 7.20-7.50(9H, m).

MS (m/z) : 359(M^+).

Example 86

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-phenylpyrazole was prepared by a similar method to that of Example 22.

white crystals.

mp : 139-142°C.

IR (KBr) : 1758, 1594, 1492, 1434 cm^{-1} .

NMR (DMSO- d_6 , δ) : 2.03(3H, s), 5.43 (2H, s), 7.01(1H, s), 7.20-7.50(5H, m)
7.53(2H, dd, J=7, 2Hz), 7.89(2H, dd, J=7, 2Hz) ,

MS(m/z) : 391(M^{+1}).

Example 87

3-Chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-phenylpyrazole was prepared by the two step procedure. In step one, the starting material was hydrolyzed by a similar method to that of Preparation 14, and followed by alkylation by a similar method to that of Example 3.

white crystals.

mp : 110-112°C

IR (KBr) : 3500, 3490, 3400, 1596, 1492 cm^{-1} .

NMR (DMSO- d_6 , δ) : 1.50-1.80(2H, m), 3.30-3.50(4H, m), 4.65(1H, t, J=5Hz),
6.99(1H, s), 7.25-7.60(7H, m), 7.87(2H, d, J=8Hz)

MS(m/z) : 377(M^{+1}).

Anal. Calcd. for $C_{18}H_{17}ClN_2O_3S$: C, 57.37 ; H, 4.55 ; N, 7.43.

Found (1/6 H_2O) : C, 56.92 ; H, 4.60 ; N, 7.37.

Example 88

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole was prepared by a similar method to that of Example 21.

white crystals.

mp : 120-124°C

IR (KBr) : 1737, 1594, 1488 cm^{-1} .

NMR (DMSO- d_6 , δ) : 2.04(3H, s), 3.75(3H, s), 5.53(2H, s), 6.79(1H, dd, J=8,
2Hz), 6.85(1H, s), 7.15(1H, s), 7.10-7.50(5H, m).

MS(m/z) : 423(M^{+1}), 425(M^{+3}).

Example 89

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole was prepared by a similar method to that of Example 22.

white crystals.

mp : 124-125°C

IR (KBr) : 1766, 1594, 1490, 1459 cm^{-1} .

NMR (DMSO- d_6 , δ) : 2.04(3H, s), 3.75(3H, s), 5.44(2H, s), 6.79(1H, dd, J=8, 2Hz), 7.03(1H, s), 7.15(1H, d, J=2Hz), 7.46(1H, d, J=8Hz), 7.57(2H, d, J=8Hz), 7.92(2H, d, J=8Hz).

MS(m/z) : 455(M^+), 457(M^+)

Example 90

3-Chloro-1-(4-chloro-3-methoxyphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole was prepared by the two step procedure. In step one, the starting material was hydrolyzed by similar method as that described for Preparation 14, and followed by alkylation by similar method to that described for Example 3.

white crystals.

mp ; 113-114°C

IR (KBr) : 3519, 3465, 3403, 1594, 1488 cm^{-1} .

NMR (DMSO- d_6 , δ) : 1.50-1.80(2H, m), 3.30-3.50(4H, m), 3.73(3H, s), 4.65(1H, t, J=5Hz), 6.79(1H, dd, J=8, 2Hz), 7.01(1H, s), 7.15(1H, d, J=2Hz), 7.47(1H, d, J=8Hz), 7.56(2H, d, J=8Hz), 7.91(2H, d, J=8Hz).

MS(m/z) : 441(M^+), 443(M^+).

Anal. Calcd for $C_{18}H_{17}ClN_2O_3S$: C, 51.71 ; H, 4.11 ; N, 6.35.

Found : C, 51.57 ; H, 4.06 ; N, 6.26.

Example 91

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(3-chlorophenyl)-pyrazole was prepared by a similar procedure to that of Example 21.

NMR ($CDCl_3$, δ) : 2.11(3H, s), 5.44(2H, s), 6.44(1H, s), 7.06 (1H, pd, t, J=8Hz), 7.17(2H, d, J=8Hz), 7.19-7.43(3H, m), 7.40(2H, d, J=8Hz).

IR (Neat) : 1747 cm^{-1}

mass (m/z) : 393(M+1)

Example 92

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chlorophenyl)-pyrazole was prepared by a similar procedure to that of Example 22.

NMR ($CDCl_3$, δ) : 2.10(3H, s), 5.16(2H, s), 6.56(1H, s), 7.01(1H, dt, J=8.1Hz),

7.26-7.47(3H, m), 7.44(2H, d, J=8Hz), 7.91(2H, d, J=8Hz).

IR (KBr) : 1766,1328, 1151 cm^{-1}

mass (m/z) : 425(M+1)

Example 93

3-Chloro-1-(3-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-pyrazole was prepared a similar procedure to that of Example 3.

NMR (CDCl_3 , δ) : 1.72(1H, t, J=5Hz), 1.92-2.07(2H, m), 3.22-3.31(2H, m), 3.75(2H, q, J=6Hz), 6.55(1H, s), 7.04(1H, dt, J=8.2Hz), 7.23-7.45(1H, m), 7.42(2H, d, J=9Hz), 7.90(2H, d, J=9Hz).

IR (KBr) : 3405, 1305 cm^{-1}

mass (m/z) : 411(M+1)

mp : 108-110°C

Example 94

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(4-chlorophenyl)pyrazole was prepared by a similar procedure to that of Example 21.

NMR (CDCl_3 , δ) : 2.11(3H, s), 5.44(2H, s), 6.43(1H, s), 7.15(2H, d, J=9Hz), 7.21(2H, d, J=9Hz), 7.32(1H, d, J=9Hz), 7.39(2H, d, J=9Hz).

IR (Neat) : 1745 cm^{-1}

mass (m/z) : 393(M+1)

Example 95

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chlorophenyl)-pyrazole was prepared by a similar procedure to that of Example 22.

NMR (CDCl_3 , δ) : 2.08(3H, s), 5.16(2H, s), 6.56(1H, s), 7.18(2H, d, J=9Hz), 7.35(2H, d, J=9Hz), 7.42(2H, d, J=9Hz), 7.90(2H, d, J=9Hz).

IR (KBr) : 1758, 1330,1149 cm^{-1}

mass (m/z) : 425(M+1)

Example 96

3-Chloro-1-(4-chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-pyrazole was prepared by a similar procedure to that of Example 3.

NMR (CDCl_3 , δ) : 2.61(1H, t, J=6Hz), 3.35-3.40(2H, m), 4.00-4.09(2H, m), 6.54(1H, s), 7.19(2H, d, J=9Hz), 7.35(2H, d, J=9Hz), 7.42(2H, d, J=9Hz),

7.91(2H, d, J=9Hz), 7.35(2H, d, J=8.9Hz), 7.42(2H, d, J=8.6Hz), 7.91(2H, d, J=8.6Hz).

IR (KBr) : 3469, 1309 cm^{-1}

mass (m/z) : 397(M+1)

mp : 150-151°C

Example 97

3-Chloro-1-(4-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-pyrazole was prepared by a similar procedure to that of Example 3

NMR (CDCl_3 , δ) : 1.67(1H, t, J=5Hz), 1.94-2.08(2H, m), 3.22-3.31(2H, m), 3.77(2H, q, J=6Hz), 6.54(1H, s), 7.19(2H, d, J=9Hz), 7.35(2H, d, J=9Hz), 7.41(2H, d, J=9Hz), 7.89(2H, d, J=9Hz).

IR (KBr) : 3403, 1305 cm^{-1}

mass (m/z) : 411(M+1)

mp : 156-158°C

Example 98

5-[4-(Acetoxymethylthio)phenyl]-3-chloro-1-(4-fluorophenyl)pyrazole was prepared by a similar procedure to that of Example 21.

NMR (CDCl_3 , δ) : 2.11(3H, s), 5.43(2H, s), 6.43(1H, s), 7.04(2H, t, J=9Hz), 7.14(2H, d, J=9Hz), 7.22-7.29(2H, m), 7.38(2H, d, J=9Hz).

IR (Neat) : 1747 cm^{-1}

mass (m/z) : 377(M+1)

Example 99

5-[4-(Acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluorophenyl)-pyrazole was prepared by a similar procedure to that of Example 22.

NMR (CDCl_3 , δ) : 2.08(3H, s), 5.15(2H, s), 6.56(1H, s), 7.08(2H, t, J=9Hz), 7.19-7.27(2H, m), 7.41(2H, d, J=9Hz), 7.89(2H, d, J=9Hz).

IR (KBr) : 1770, 1317, 1157 cm^{-1}

mass (m/z) : 409(M+1)

Example 100

3-Chloro-1-(4-fluorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-pyrazole was prepared by a similar procedure to that of Example 3.

NMR (CDCl₃, δ) : 1.63(1H, t, J=5Hz), 1.94-2.08(2H, m), 3.26(2H, t, J=8Hz), 2.77(q, J=5Hz), 6.54(1H, s), 7.07(2H, t, J=9Hz), 7.20-7.28(2H, m), 7.40(2H, d, J=8Hz), 7.88(2H, d, J=8Hz).

IR (KBr) : 3403, 1305 cm⁻¹

mass (m/z) : 395(M+1)

mp : 125-126°C

Example 101

5-(4-Methylphenyl)-1-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar manner to that of Example 3.

IR (Neat) : 1595, 1465, 1320, 1270, 1240, 1140 cm⁻¹

NMR (CDCl₃, δ) : 2.38(3H, s), 3.22(3H, s), 3.39(2H, t, J=6.0Hz), 3.77(2H, t, J=6.0Hz), 6.75(1H, s), 7.08-7.92(9H, m).

MASS : 425(M⁺+1)

Example 102

5-(4-Methylphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar manner to that of Example 3.

IR (Neat) : 3475, 1465, 1275, 1235, 1135 cm⁻¹

Example 103

5-(4-Chloro-3-methoxyphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole was prepared by a similar manner to that of Example 3.


IR (Neat) : 3480, 1665, 1475, 1315, 1280, 1240, 1140 cm⁻¹

NMR (CDCl₃, δ) : 2.38(3H, s), 2.59 (1H, t, J=6.0Hz), 3.38(2H, t, J=6.0Hz), 4.04(2H, q, J=6.0Hz), 6.86(1H, s), 6.90-8.02(7H, m).

MASS : 445(M⁺+1)

CLAIMS

1. A compound of the formula (I):



wherein

R^1 is halo(lower)alkyl, halogen or cyano,

R^2 is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino,

R^3 is hydrogen, hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl,


R^4 is hydrogen or halogen,

Q is pyrazolyl,

A is lower alkylene optionally substituted with oxo or hydroxy, and n is 0, 1 or 2,

provided that when R^3 is hydrogen, R^2 is aryl substituted with lower alkenyl or A is lower alkylene substituted with oxo, or its salt.

2. A compound of the formula (I')



wherein

R¹ is halo(lower)alkyl, halogen or cyano,

R² is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino,

R³ is hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl,

R⁴ is hydrogen or halogen,

A is lower alkylene optionally substituted with oxo or hydroxy, and n is 0, 1 or 2,

provided that when R³ is hydrogen, R² is aryl substituted with lower alkenyl or A is lower alkylene substituted with oxo, or its salt.

3. The compound according to claim 2, wherein

R¹ is trifluoromethyl, difluoromethyl, chlorine or cyano,

R² is phenyl, tolyl, or phenyl or tolyl substituted with substituent(s) selected from the group consisting of chlorine, fluorine, bromine and methoxy,

R³ is hydroxy, acetoxy, ethoxy, amino, methylamino, methylthio or methylsulfonyl,

R⁴ is hydrogen,

A is dimethylene, trimethylene, methylene or pentamethylene, and n is 2.

4. The compound according to claim 3, which is a compound selected from the group consisting of

(1) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chloro-4-methoxyphenyl)pyrazole,

(2) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,

(3) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]-

- phenyl}-3-(trifluoromethyl)pyrazole,
- (4) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}pyrazole,
- (5) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (6) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylthio)ethyl]sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (7) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(5-hydroxypentyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (8) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(4-hydroxybutyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (9) 1-(3-chloro-4-methoxyphenyl)-5-[4-(ethoxycarbonylmethylsulfonyl)-phenyl]-3-(trifluoromethyl)pyrazole,
- (10) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methylaminoethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (11) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-aminoethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (12) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (13) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-acetoxyethyl)-sulfonyl]phenyl}pyrazole,
- (14) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(methylsulfonyl)ethyl]-sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (15) 5-{4-[(acetoxymethyl)sulfonyl]phenyl}-1-(4-fluorophenyl)-3-(trifluoromethyl)pyrazole,
- (16) 1-(4-fluorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (17) 5-{4-[(acetoxymethyl)sulfonyl]phenyl}-1-(4-chloro-3-methoxyphenyl)-3-cyanopyrazole,
- (18) 1-(4-chloro-3-methoxyphenyl)-5-{4-[2-hydroxyethyl)sulfonyl]-phenyl}-3-cyanopyrazole,

- (19) 5-[4-(acetoxymethylthio)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole,
- (20) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chlorophenyl)-3-(trifluoromethyl)pyrazole,
- (21) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-methylphenyl)-pyrazole,
- (22) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-methylphenyl)-pyrazole,
- (23) 3-chloro-1-(4-isopropenyl-phenyl)-5-[4-(methylsulfinyl)-phenyl]pyrazole,
- (24) 5-[[4-(acetoxymethylthio)phenyl]-3-chloro-1-(isopropenylphenyl)-pyrazole,
- (25) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-phenyl}pyrazole,
- (26) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}pyrazole,
- (27) 1-(4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (28) 1-(4-methylphenyl)-5-{4-[(3-acetoxypropyl)sulfonyl] phenyl}-3-(trifluoromethyl)pyrazole,
- (29) 1-(4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (30) 5-[4-(acetoxymethylthio)phenyl]-3-trifluoromethyl-1-[(3-chloro-4-methyl)phenyl]pyrazole,
- (31) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (32) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-acetoxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (33) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-phenyl}-3-(trifluoromethyl)pyrazole,
- (34) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]-

- phenyl}-3-(trifluoromethyl)pyrazole,
- (35) 1-(3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (36) 1-(3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (37) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (38) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (39) 5-{4-[(acetoxymethyl)sulfinyl]-3-fluorophenyl}-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (40) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]-3-fluorophenyl}-3-(trifluoromethyl)pyrazole,
- (41) 1-(4-chloro-3-methylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (42) 1-(1-hydroxy-1-methylethyl)-5-[4-(methylthio)phenyl]-3-(trifluoromethyl)pyrazole,
- (43) 1-(4-isopropenylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (44) 1-(3-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (45) 1-(3-chlorophenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (46) 1-(4-isopropylphenyl)-5-[4-(acetoxymethylthio)phenyl]-3-(trifluoromethyl)pyrazole,
- (47) 1-(4-isopropylphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (48) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-cyano-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (49) 1-(3-cyano-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,

- (50) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-methoxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (51) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(N-hydroxycarbamoyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (52) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(methylcarbamoyl)methylsulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (53) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-oxopropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (54) 5-{4-[(3-acetoxypropyl)sulfonyl]phenyl}-1-(3-chloro-4-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (55) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[3-(dimethylamino)propyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole hydrochloride,
- (56) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(dimethylamino)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole hydrochloride,
- (57) 5-[4-(acetoxymethylsulfonyl)phenyl]-1-(3-chloro-4-methoxyphenyl)-3-(difluoromethyl)pyrazole,
- (58) 1-(3-chloro-4-methoxyphenyl)-3-difluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (59) 1-(3-chloro-4-methoxyphenyl)-5-{4-[[2-(2-hydroxyethoxy)ethyl]sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (60) 5-[4-(acetoxymethylthio)phenyl]-1-(4-bromo-3-methoxyphenyl)-3-(trifluoromethyl)pyrazole,
- (61) 1-(4-bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (62) 1-(4-bromo-3-methoxyphenyl)-3-trifluoromethyl-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (63) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-difluoromethyl-1-[4-(methoxyphenyl)pyrazole,
- (64) 1-(4-methoxyphenyl)-3-difluoromethyl-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (65) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-bromophenyl)-

- pyrazole,
- (66) 1-(3-bromophenyl)-3-chloro-5-{4-(3-hydroxypropyl)sulfonyl}-phenyl}pyrazole,
- (67) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-fluoro-4-methylphenyl)pyrazole,
- (68) 3-chloro-1-(3-fluoro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (69) 5-[(4-acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluoro-3-methylphenyl)pyrazole,
- (70) 3-chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (71) 3-chloro-1-(4-fluoro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (72) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (73) 1-(3-chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (74) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (75) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-methoxyethyl)sulfonyl]phenyl}pyrazole hydrochloride,
- (76) 5-[4-(4-acetoxymethylthio)phenyl]-3-chloro-1-(4-chloro-3-methylphenyl)pyrazole,
- (77) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (78) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (79) 3-chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (80) 3-chloro-1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,

- (81) 3-chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-(4-methylphenyl)pyrazole,
- (82) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole,
- (83) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-methylphenyl)pyrazole,
- (84) 3-chloro-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-1-(3-methylphenyl)pyrazole,
- (85) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-phenylpyrazole,
- (86) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-phenylpyrazole,
- (87) 3-chloro-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-1-phenylpyrazole,
- (88) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole,
- (89) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chloro-3-methoxyphenyl)pyrazole,
- (90) 3-chloro-1-(4-chloro-3-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (91) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(3-chlorophenyl)pyrazole,
- (92) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(3-chlorophenyl)pyrazole,
- (93) 3-chloro-1-(3-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
- (94) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-chlorophenyl)pyrazole,
- (95) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-chlorophenyl)pyrazole,
- (96) 3-chloro-1-(4-chlorophenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (97) 3-chloro-1-(4-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]-

- phenyl}pyrazole,
(98) 5-[4-(acetoxymethylthio)phenyl]-3-chloro-1-(4-fluorophenyl)-pyrazole,
(99) 5-[4-(acetoxymethylsulfonyl)phenyl]-3-chloro-1-(4-fluorophenyl)-pyrazole,
(100) 3-chloro-1-(4-fluorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,
(101) 5-(4-methylphenyl)-1-{4-[(2-methoxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
(102) 5-(4-methylphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole, and
(103) 5-(4-chloro-3-methoxyphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole.

5. The compound according to claim 4, which is a compound selected from the group consisting of

- (1) 1-(3-chloro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (2) 1-(3-fluoro-4-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (3) 1-(4-fluoro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (4) 1-(4-bromo-3-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (5) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (6) 1-(3-chloro-4-methoxyphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole,
- (7) 3-chloro-1-(4-isopropenylphenyl)-5-{4-[(2-hydroxyethyl)sulfonyl]phenyl}pyrazole,
- (8) 3-chloro-1-(4-chlorophenyl)-5-{4-[(3-hydroxypropyl)sulfonyl]phenyl}pyrazole,

phenyl}pyrazole,


(9) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(2-hydroxyethyl)-sulfonyl]phenyl}pyrazole,

(10) 3-chloro-1-(3-chloro-4-methoxyphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole,

(11) 3-chloro-1-(4-chloro-3-methylphenyl)-5-{4-[(3-hydroxypropyl)-sulfonyl]phenyl}pyrazole, and

(12) 5-(4-chloro-3-methoxyphenyl)-1-{4-[(2-hydroxyethyl)sulfonyl]phenyl}-3-(trifluoromethyl)pyrazole.

6. A process for preparing a compound of the formula (I):



wherein

R¹ is halo(lower)alkyl, halogen or cyano,

R² is aryl optionally substituted with substituent(s) selected from the group consisting of halogen, lower alkoxy, cyano, nitro, lower alkenyl, amino, acyl and hydroxyimino,

R³ is hydrogen, hydroxy, acyloxy, lower alkoxy optionally substituted with lower alkoxy, amino optionally substituted with hydroxy or lower alkyl, lower alkylthio, or lower alkylsulfonyl,

R⁴ is hydrogen or halogen,

Q is pyrazolyl,


A is lower alkylene optionally substituted with oxo or hydroxy, and

n is 0, 1 or 2,

provided that when R³ is hydrogen, R² is aryl substituted with lower

alkenyl or A is lower alkylene substituted with oxo, or its salt,
which comprises,


(1) reacting a compound of the formula (II)



(II)

or its salt


wherein R^1 , R^2 , R^4 , Q and n are as defined above,
with CH_3COONa or $(\text{CH}_3\text{CO})_2\text{O}$
to give a compound of the formula (I-1)



(I-1)


or its salt

wherein R^1 , R^2 , R^4 , Q and n are as defined above,
(2) reacting a compound of the formula (IV):




(IV)

wherein R^1 , R^2 , R^4 and Q are as defined above,
 with a compound (V): R^3 -A-Hal wherein A and R^3 are as defined above, or
 its salt,
 to give a compound (I-2):




or its salt

wherein R^1 , R^2 , R^3 , R^4 , A and Q are as defined above,
 (3) subjecting a compound of the formula (I-3):




wherein R^1 , R^2 , R^4 , A, Q and n are as defined above,
 to amination to give a compound of the formula (I-4)




or its salt

wherein R^{3a} is amino optionally substituted with hydroxy or lower alkyl, and R¹, R², R⁴, A, Q and n are as defined above,


(4) subjecting a compound of the formula (I-3)




wherein R¹, R², R⁴, A, Q and n are as defined above, to acylation to give a compound of the formula (I-5)



wherein R^{3b} is acyl, and R¹, R², R⁴, A, Q and n are as defined above, or (5) subjecting a compound of the formula (I-6)



wherein R is lower alkyl, and R¹, R², R⁴, A, Q and n are as defined above, to oxydation to give a compound of the formula (I-7)



or its salt

wherein R^{3c} is lower alkylsulfonyl, and R¹, R², R⁴, A, Q and n are as defined above.

7. A pharmaceutical composition comprising the compound of claim 1, as an active ingredient, in association with a pharmaceutically non-toxic carrier or excipient.
8. A compound of claim 1 for use as a medicament.
9. A COX-II inhibiting agent comprising the compound of claim 1.
10. A method for the treatment and/or prevention of inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or neurodegenerative diseases which comprises administering an effective amount of the compound of claim 1 to human beings or animals.
11. Use of the compound of claim 1 for the manufacture of a medicament for treatment and/or prevention of inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or neurodegenerative diseases

in human beings or animals.

12. Use of the compound of claim 1 for the manufacture of a medicament for treatment and/or prevention of inflammatory conditions, various pains, collagen diseases, autoimmune diseases, various immunity diseases, analgesic, thrombosis, cancer or neurodegenerative diseases in human beings or animals.