



HU000027932T2

(19) **HU**(11) Lajstromszám: **E 027 932**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 11 799673**(51) Int. Cl.: **A61K 39/04** (2006.01)(22) A bejelentés napja: **2011. 12. 14.**

(86) A nemzetközi (PCT) bejelentési szám:

(96) Az európai bejelentés bejelentési száma:
EP 20110799673**PCT/EP 11/072816**(97) Az európai bejelentés közzétételi adatai:
EP 2651437 A1 **2012. 06. 21.**

(87) A nemzetközi közzétételi szám:

WO 12080369(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2651437 B1 **2016. 02. 10.**

(30) Elsőbbségi adatok: 422723 P 2010. 12. 14. US	(73) Jogosult(ak): GlaxoSmithKline Biologicals S.A., 1330 Rixensart (BE)
(72) Feltaláló(k): GODART, Stephane Andre Georges, B-1330 Rixensart (BE) LAANAN, Amina, B-1330 Rixensart (BE) LEMOINE, Dominique Ingrid, B-1330 Rixensart (BE)	(74) Képviselő: Danubia Szabadalmi és Jogi Iroda Kft., Budapest

(54) **Antigén mycobacterium-készítmény**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 651 437 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
10.02.2016 Bulletin 2016/06

(21) Application number: **11799673.6**

(22) Date of filing: **14.12.2011**

(51) Int Cl.:
A61K 39/04 (2006.01)

(86) International application number:
PCT/EP2011/072816

(87) International publication number:
WO 2012/080369 (21.06.2012 Gazette 2012/25)

(54) **MYCOBACTERIUM ANTIGENIC COMPOSITION**

ANTIGENE MYCOBACTERIUM-ZUSAMMENSETZUNG

COMPOSITION ANTIGÉNIQUE DE MYCOBACTERIUM

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR
Designated Extension States:
BA ME

(30) Priority: **14.12.2010 US 422723 P**

(43) Date of publication of application:
23.10.2013 Bulletin 2013/43

(60) Divisional application:
15201196.1

(73) Proprietor: **GlaxoSmithKline Biologicals S.A.**
1330 Rixensart (BE)

(72) Inventors:
• **GODART, Stephane Andre Georges**
B-1330 Rixensart (BE)
• **LAANAN, Amina**
B-1330 Rixensart (BE)

• **LEMOINE, Dominique Ingrid**
B-1330 Rixensart (BE)

(74) Representative: **West, Heloise**
GlaxoSmithKline
Global Patents (CN925.1)
980 Great West Road
Brentford, Middlesex TW8 9GS (GB)

(56) References cited:
WO-A1-2010/142685 **WO-A1-2010/142686**
WO-A2-2006/117240 **WO-A2-2009/088255**
US-A1- 2009 291 062 **US-B1- 6 869 607**

• **LEROUX-ROELS ISABEL ET AL: "Evaluation of the Safety and Immunogenicity of Two Antigen Concentrations of the Mtb72F/AS02(A) Candidate Tuberculosis Vaccine in Purified Protein Derivative-Negative Adults", CLINICAL AND VACCINE IMMUNOLOGY, vol. 17, no. 11, November 2010 (2010-11), pages 1763-1771, XP002669941,**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 651 437 B1

Description

Field of the Invention

5 [0001] The present invention relates to immunogenic compositions comprising an M72 related antigen and having a low ionic strength. The present invention also relates to such immunogenic compositions which further comprise one or more immunostimulants.

Background of the Invention

10 [0002] Tuberculosis (TB) is a chronic infectious disease caused by infection with *Mycobacterium tuberculosis* and other *Mycobacterium* species. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world. More than 2 billion people are believed to be infected with TB bacilli, with about 9.4 million new cases of TB and 1.7 million deaths each year. 10% of those infected with TB bacilli will develop active TB, each person with active TB infecting an average of 10 to 15 others per year. While annual incidence rates have peaked globally, the number of deaths and cases is still rising due to population growth (World Health Organisation Tuberculosis Facts 2010).

15 [0003] The protein antigens Mtb72f and M72 (described, for example, in international patent application WO2006/117240) or fragments or derivatives thereof are protein antigens of potential benefit for the treatment or prevention of tuberculosis.

20 [0004] The formulation of protein antigens is extremely important in order to ensure immunogenicity is maintained. Immunostimulants are sometimes used to improve the immune response raised to any given antigen. However, the inclusion of adjuvants into an immunogenic composition increases the complexity of preparation of the components as well as the complexity of distribution and formulation of the composition. The preparation of each of the adjuvant components as well as the antigenic component must be considered by formulators. In particular, the compatibility of the antigenic component with the adjuvant component should be considered. This is particularly the case where lyophilised antigens or antigenic preparations are intended to be reconstituted with an adjuvant preparation. In such a circumstance, it is important that the buffer of the adjuvant preparation is suitable for the antigen and that immunogenicity or solubility of the antigen is not affected by the adjuvant.

25 [0005] WO2006/117240 discloses methods of preventing reactivation of M. tuberculosis infections using a pharmaceutical composition comprising M72 or Mtb72f fusion proteins.

Summary of the invention

35 [0006] The present inventors have identified for the first time that M72 related antigens are particularly sensitive to the presence of salts. Without being limited by theory, it is believed M72 related antigens are detrimentally impacted by a phenomenon known as "salting out" which may be defined as the precipitation of a protein from its solution by interaction with salts, such as sodium chloride. The present inventors have found that these antigens aggregate and precipitate at a concentration of sodium chloride as low as 150 mM. Consequently, the stability of immunogenic compositions comprising M72 related antigens can surprisingly be improved by a reduction in the concentration of sodium chloride.

40 [0007] Accordingly, the present invention provides an immunogenic composition comprising an M72 related antigen, wherein the M72 related antigen comprises a sequence having at least 90% identity to SEQ ID No: 1, wherein the conductivity of the composition is 5 mS/cm or lower and the pH of the composition is in the range 7.0 to 9.0.

Brief description of the drawings

- 45 [0008]
- Figure 1. QS21 lytic activity curve
 - Figure 2. Percentage of each 3D-MPL congener in the different ASA formulations
 - 50 Figure 3. DLS of immunogenic compositions with varied pH and NaCl concentrations after storage
 - Figure 4. Nephelometry of immunogenic compositions with varied pH and NaCl concentrations after storage
 - Figure 5. Antigenic stability of immunogenic compositions with varied pH and NaCl concentrations following after storage
 - Figures 6a-6d. SEC-HPLC analysis of immunogenic compositions with varied pH and NaCl concentrations after storage
 - 55 Figure 7. Antigenicity of immunogenic compositions with varied pH and NaCl concentrations after storage
 - Figure 8. Conductivity of NaCl standard solutions
 - Figure 9. Induction of CD4 T cell responses in mice using immunogenic compositions of the invention

- Figure 10. Induction of CD8 T cell responses in mice using immunogenic compositions of the invention
 Figure 11. Nephelometry of immunogenic compositions with varied pH and NaCl concentrations after storage
 Figure 12. DLS of immunogenic compositions with varied pH and NaCl concentrations after storage
 Figure 13. Antigenicity of immunogenic compositions with varied NaCl concentrations after storage

5

Brief description of sequence identifiers

[0009]

- 10 SEQ ID No: 1 Amino acid sequence for the M72 protein
 SEQ ID No: 2 Nucleotide sequence encoding the M72 protein
 SEQ ID No: 3 Amino acid sequence for the M72 protein with two N-terminal His residues
 SEQ ID No: 4 Nucleotide sequence encoding the M72 protein with two N-terminal His residues
 SEQ ID No: 5 Amino acid sequence for the Mtb72f protein
 15 SEQ ID No: 6 Nucleotide sequence encoding the Mtb72f protein
 SEQ ID No: 7 Amino acid sequence for the Mtb72f protein with six N-terminal His residues
 SEQ ID No: 8 Nucleotide sequence encoding the Mtb72f protein with six N-terminal His residues
 SEQ ID No: 9 Nucleotide sequence for CpG Oligo 1 (CpG 1826)
 SEQ ID No: 10 Nucleotide sequence for CpG Oligo 2 (CpG 1758)
 20 SEQ ID No: 11 Nucleotide sequence for CpG Oligo 3
 SEQ ID No: 12 Nucleotide sequence for CpG Oligo 4 (CpG 2006)
 SEQ ID No: 13 Nucleotide sequence for CpG Oligo 5 (CpG 1686)

Detailed description of the invention

25

[0010] In a first aspect, the present invention provides an immunogenic composition comprising an M72 related antigen, wherein the M72 related antigen comprises a sequence having at least 90% identity to SEQ ID No: 1, the pH of the composition is in the range 7.0 to 9.0 and wherein the conductivity of the composition is 5 mS/cm or lower, 4 mS/cm or lower, or 3 mS/cm or lower. In a particular embodiment the conductivity of the immunogenic composition is 2.5 mS/cm or lower, such as 2.25 mS/cm or lower, or 2.0 mS/cm or lower. In a further specific embodiment the conductivity of the immunogenic composition is 1.5 to 2.5 mS/cm.

30

[0011] In some embodiments, the present invention provides an immunogenic composition wherein the concentration of salts in said composition is 40 mM or lower. In a particular embodiment the concentration of salts in said composition is 35 mM or lower, such as 30 mM or lower, or 25 mM or lower. In a further specific embodiment the concentration of salts in said composition is 20 to 40 mM, such as 25 to 35 mM.

35

[0012] In some embodiments the concentration of sodium chloride is 40 mM or lower, 30 mM or lower, 20 mM or lower or 15 mM or lower. In a particular embodiment the concentration of sodium chloride in the immunogenic composition is 10 mM or lower, such as 7.5 mM or lower. Suitably the concentration of sodium chloride in the immunogenic composition or is at or below 5 mM. In a further specific embodiment, the immunogenic composition is essentially free of sodium chloride. By essentially free is meant that the concentration of sodium chloride is at or very near to zero mM (such as 3 mM or less, 2 mM or less or 1 mM or less).

40

[0013] Suitably, the concentration of CaCl₂ in the immunogenic compositions will be 30 mM or lower, 20 mM or lower, 15 mM or lower or 10 mM or lower.

[0014] Suitably, the concentration of MgSO₄ in the immunogenic compositions will be 60 mM or lower, 40 mM or lower, 30 mM or lower, 20 mM or lower or 10 mM or lower.

45

[0015] Suitably, the total concentration of NH₄⁺, Mg²⁺ and Ca²⁺ ions in the immunogenic compositions will be 60 mM or lower, 40 mM or lower, 30 mM or lower, 20 mM or lower or 10 mM or lower.

[0016] The immunogenic compositions of the invention will be aqueous preparations.

[0017] The conductivity of an immunogenic composition of the invention can be measured using techniques known in the art, for example using a dedicated conductivity meter or other instrument with the capability to measure conductivity. One suitable instrument is the Zetasizer Nano ZS from Malvern Instruments (UK).

50

[0018] The skilled person can readily test for the concentration of both sodium (Na⁺) and chloride (Cl⁻) ions using known techniques and kits. For example, sodium can be determined using a kit such as the Sodium Enzymatic Assay Kit (Catalogue Number: BQ011EAEL) from Biosupply. Chloride can be determined using a kit such as Chloride Enzymatic Assay Kit (Catalogue Number: BQ006EAEL) from Biosupply.

55

[0019] Tuberculosis (TB) is a chronic infectious disease caused by infection with *Mycobacterium tuberculosis* and other *Mycobacterium* species. It is a major disease in developing countries, as well as an increasing problem in developed areas of the world. More than 2 billion people are believed to be infected with TB bacilli, with about 9.4 million new cases

of TB and 1.7 million deaths each year. 10% of those infected with TB bacilli will develop active TB, each person with active TB infecting an average of 10 to 15 others per year. While annual incidence rates have peaked globally, the number of deaths and cases is still rising due to population growth (World Health Organisation *Tuberculosis Facts* 2010).

[0020] *Mycobacterium tuberculosis* infects individuals through the respiratory route. Alveolar macrophages engulf the bacterium, but it is able to survive and proliferate by inhibiting phagosome fusion with acidic lysosomes. A complex immune response involving CD4+ and CD8+ T cells ensues, ultimately resulting in the formation of a granuloma. Central to the success of *Mycobacterium tuberculosis* as a pathogen is the fact that the isolated, but not eradicated, bacterium may persist for long periods, leaving an individual vulnerable to the later development of active TB.

[0021] Fewer than 5% of infected individuals develop active TB in the first years after infection. The granuloma can persist for decades and is believed to contain live *Mycobacterium tuberculosis* in a state of dormancy, deprived of oxygen and nutrients. However, recently it has been suggested that the majority of the bacteria in the dormancy state are located in non-macrophage cell types spread throughout the body (Locht et al, Expert Opin. Biol. Ther. 2007 7(11):1665-1677). The development of active TB occurs when the balance between the host's natural immunity and the pathogen changes, for example as a result of an immunosuppressive event (Anderson P Trends in Microbiology 2007 15(1):7-13; Ehlers S Infection 2009 37(2):87-95).

[0022] A dynamic hypothesis describing the balance between latent TB and active TB has also been proposed (Cardana P-J Inflammation & Allergy - Drug Targets 2006 6:27-39; Cardana P-J Infection 2009 37(2):80-86).

[0023] Although an infection may be asymptomatic for a considerable period of time, the active disease is most commonly manifested as an acute inflammation of the lungs, resulting in tiredness, weight loss, fever and a persistent cough. If untreated, serious complications and death typically result.

[0024] Tuberculosis can generally be controlled using extended antibiotic therapy, although such treatment is not sufficient to prevent the spread of the disease. Actively infected individuals may be largely asymptomatic, but contagious, for some time. In addition, although compliance with the treatment regimen is critical, patient behaviour is difficult to monitor. Some patients do not complete the course of treatment, which can lead to ineffective treatment and the development of drug resistance.

[0025] Multidrug-resistant TB (MDR-TB) is a form which fails to respond to first line medications. 3.3% of all TB cases are MDR-TB, with an estimated 440,000 new MDR-TB cases occurring each year. Extensively drug-resistant TB (XDR-TB) occurs when resistance to second line medications develops on top of resistance to first line medications. The virtually untreatable XDR-TB has been confirmed in 58 countries (World Health Organisation *Tuberculosis Facts* 2010).

[0026] Even if a full course of antibiotic treatment is completed, infection with *M. tuberculosis* may not be eradicated from the infected individual and may remain as a latent infection that can be reactivated. In order to control the spread of tuberculosis, an effective vaccination programme and accurate early diagnosis of the disease are of utmost importance.

[0027] Currently, vaccination with live bacteria is the most widely used method for inducing protective immunity. The most common *Mycobacterium* employed for this purpose is *Bacillus Calmette-Guerin* (BCG), an avirulent strain of *M. bovis* which was first developed over 60 years ago. However, the safety and efficacy of BCG is a source of controversy - while protecting against severe disease manifestation in children, BCG does not prevent the establishment of latent TB or the reactivation of pulmonary disease in adult life. Additionally, some countries, such as the United States, do not vaccinate the general public with this agent.

[0028] Several of the proteins which are strongly expressed during the early stages of *Mycobacterium* infection have been shown to provide protective efficacy in animal vaccination models. However, vaccination with antigens which are highly expressed during the early stages of infection may not provide an optimal immune response for dealing with later stages of infection. Adequate control during latent infection may require T cells which are specific for the particular antigens which are expressed at that time. Post-exposure vaccines which directly target the dormant persistent bacteria may aid in protecting against TB reactivation, thereby enhancing TB control, or even enabling clearance of the infection. A vaccine targeting latent TB could therefore significantly and economically reduce global TB infection rates.

[0029] Subunit vaccines based on late stage antigens could also be utilised in combination with early stage antigens to provide a multiphase vaccine. Alternatively, early and/or late stage antigens could be used to complement and improve BCG vaccination (either by boosting the BCG response or through the development of advanced recombinant BCG strains).

[0030] The protein antigens Mtb72f and M72 are protein antigens of potential benefit for the treatment or prevention of tuberculosis. Mtb72f has been shown to provide protection in a number of animal models (see, for example: Brandt et al Infect. Immun. 2004 72(11):6622-6632; Skeiky et al J. Immunol. 2004 172:7618-7628; Tsenova et al Infect. Immun. 2006 74(4):2392-2401; Reed et al PNAS 2009 106(7):2301-2306). Mtb72f has also been the subject of clinical investigations (Von Eschen et al 2009 Human Vaccines 5(7):475-482). M72 is an improved antigen which incorporates a single serine to alanine mutation relative to Mtb72f, resulting in improved stability characteristics. M72 related antigens have also been shown to be of value in a latent TB model (international patent application WO2006/117240).

[0031] As used herein the term 'M72 related antigen' refers to the M72 protein provided in SEQ ID No: 1 or an immunogenic derivative thereof. As used herein the term "derivative" refers to an antigen that is modified relative to the

reference sequence. Immunogenic derivatives are sufficiently similar to the reference sequence to retain the immunogenic properties of the reference sequence and remain capable of allowing an immune response to be raised against the reference sequence. A derivative may, for example, comprise a modified version of the reference sequence or alternatively may consist of a modified version of the reference sequence.

5 **[0032]** The M72 related antigen may for example contain fewer than 1500 amino acid residues, such as fewer than 1200 amino acid residues, in particular less than 1000 amino acid residues, especially fewer than 800 amino acid residues.

[0033] T cell epitopes are short contiguous stretches of amino acids which are recognised by T cells (e.g. CD4+ or CD8+ T cells). Identification of T cell epitopes may be achieved through epitope mapping experiments which are known to the person skilled in the art (see, for example, Paul, *Fundamental Immunology*, 3rd ed., 243-247 (1993); Beißbarth et al *Bioinformatics* 2005 21(Suppl. 1):i29-i37). In a diverse out-bred population, such as humans, different HLA types mean that particular epitopes may not be recognised by all members of the population. As a result of the crucial involvement of the T cell response in tuberculosis, to maximise the level of recognition and scale of immune response, an immunogenic derivative of M72 is desirably one which contains the majority (or suitably all) T cell epitopes intact.

15 **[0034]** The skilled person will recognise that individual substitutions, deletions or additions to the M72 protein which alters, adds or deletes a single amino acid or a small percentage of amino acids is an "immunogenic derivative" where the alteration(s) results in the substitution of an amino acid with a functionally similar amino acid or the substitution/deletion/addition of residues which do not substantially impact the immunogenic function.

[0035] Conservative substitution tables providing functionally similar amino acids are well known in the art. In general, such conservative substitutions will fall within one of the amino-acid groupings specified below, though in some circumstances other substitutions may be possible without substantially affecting the immunogenic properties of the antigen. The following eight groups each contain amino acids that are typically conservative substitutions for one another:

- 1) Alanine (A), Glycine (G);
 - 2) Aspartic acid (D), Glutamic acid (E);
 - 25 3) Asparagine (N), Glutamine (Q);
 - 4) Arginine (R), Lysine (K);
 - 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
 - 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
 - 7) Serine (S), Threonine (T); and
 - 30 8) Cysteine (C), Methionine (M)
- (see, e.g., Creighton, *Proteins* 1984).

[0036] Suitably such substitutions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

35 **[0037]** Immunogenic derivatives may also include those wherein additional amino acids are inserted compared to the reference sequence. Suitably such insertions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen. One example of insertions includes a short stretch of histidine residues (e.g. 2-6 residues) to aid expression and/or purification of the antigen in question.

[0038] Immunogenic derivatives include those wherein amino acids have been deleted compared to the reference sequence. Suitably such deletions do not occur in the region of an epitope, and do not therefore have a significant impact on the immunogenic properties of the antigen.

[0039] The skilled person will recognise that a particular immunogenic derivative may comprise substitutions, deletions and additions (or any combination thereof).

45 **[0040]** The terms "identical" or percentage "identity," in the context of two or more polypeptide sequences, refer to two or more sequences or sub-sequences that are the same or have a specified percentage of amino acid residues that are the same (i.e., 70% identity, optionally 75%, 80%, 85%, 90%, 95%, 98% or 99% identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. This definition also refers to the compliment of a test sequence. Optionally, the identity exists over a region that is at least 500 amino acids in length, such as at least 600 amino acids or at least 700 amino acids. Suitably, the comparison is performed over a window corresponding to the entire length of the reference sequence (as opposed to the derivative sequence).

[0041] For sequence comparison, one sequence acts as the reference sequence, to which the test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percentage sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

55 **[0042]** A "comparison window", as used herein, refers to a segment in which a sequence may be compared to a

reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerised implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel et al., eds. 1995 supplement)).

[0043] One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7.0 (Devereaux et al., *Nuc. Acids Res.* 12:387-395 (1984)).

[0044] Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* 25:3389-3402 (1997) and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (website at www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al.*, *supra*). These initial neighbourhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

[0045] The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Natl. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

[0046] In any event, immunogenic derivatives of a polypeptide sequence will have essentially the same activity as the reference sequence. By essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in an *in vitro* restimulation assay of PBMC or whole blood with specific antigens (e.g. restimulation for a period of between several hours to up to two weeks, such as up to one day, 1 day to 1 week or 1 to 2 weeks) that measures the activation of the cells via lymphoproliferation, production of cytokines in the supernatant of culture (measured by ELISA, CBA etc) or characterisation of T and B cell responses by intra and extracellular staining (e.g. using antibodies specific to immune markers, such as CD3, CD4, CD8, IL2, TNF-alpha, IFN-gamma, CD40L, CD69 etc) followed by analysis with a flowcytometer. Suitably, by essentially the same activity is meant at least 50%, suitably at least 75% and especially at least 90% activity of the reference sequence in a T cell proliferation and/or IFN-gamma production assay.

5 [0047] Particular derivatives of the M72 protein include those with additional His residues at the N-terminus (e.g. two His residues, as provided in SEQ ID No: 3; or a polyhistadine tag of five or particularly six His residues, which may be used for nickel affinity purification). Mtb72f (SEQ ID No: 5) which contains the original serine residue that has been mutated in M72, is a further derivative of M72, as are Mtb72f proteins with additional His residues at the N-terminus (e.g. two His residues; or a polyhistadine tag of five or particularly six His residues, which may be used for nickel affinity purification).

[0048] Suitably an M72 related antigen will comprise, such as consist of, a sequence having at least 95%, for example at least 99% to SEQ ID No. 1. Optionally, an M72 related antigen will comprise, such as consist of, a sequence having at least 98% identity to SEQ ID No. 1.

10 [0049] Typical M72 related antigens will comprise, such as consist of, an immunogenic derivative of SEQ ID No: 1 or 3 having a small number of deletions insertions and/or substitutions. Examples are those having deletions of up to 5 residues at 0-5 locations, insertions of up to 5 residues at 0-5 five locations and substitutions of up to 20 residues.

[0050] Other immunogenic derivatives of M72 are those comprising, such as consisting of, a fragment of SEQ ID No: 1 or 3 which is at least 500 amino acids in length, such as at least 600 amino acids in length or at least 700 amino acids in length.

15 [0051] M72 related antigens may be prepared by methods previously described (WO2006/117240), those provided in the Examples, or methods analogous thereto.

[0052] The immunogenic compositions may comprise one or more further antigenic components. Such additional antigenic components need not themselves be sensitive to the presence of salts in the composition.

20 [0053] Additional antigenic components may be intended to strengthen or complement the immune responses solicited by the M72 related antigen in the field of tuberculosis prevention and therapy or additional antigens could be associated with other pathogens and are intended for administration with the M72 related antigen for reasons of convenience. Where a number of antigenic components are present within the formulation, these may be provided in the form of individual polypeptides or fusion proteins. In some circumstances additional antigenic components may be provided as a polynucleotide (or polynucleotides).

25 [0054] It is well known that for parenteral administration solutions should have a pharmaceutically acceptable osmolality to avoid cell distortion or lysis. A pharmaceutically acceptable osmolality will generally mean that solutions will have an osmolality which is approximately isotonic or mildly hypertonic. Suitably the immunogenic compositions of the present invention will have an osmolality in the range of 250 to 750 mOsm/kg, for example, the osmolality may be in the range of 250 to 550 mOsm/kg, such as in the range of 280 to 500 mOsm/kg.

30 [0055] Osmolality may be measured according to techniques known in the art, such as by the use of a commercially available osmometer, for example the Advanced® Model 2020 available from Advanced Instruments Inc. (USA).

[0056] An "isotonicity agent" is a compound that is physiologically tolerated and imparts a suitable tonicity to a formulation to prevent the net flow of water across cell membranes that are in contact with the formulation.

35 [0057] Generally, sodium chloride (NaCl) is used as a tonicity agent. The present inventors have shown for the first time that that M72 related antigens are particularly sensitive to "salting out", a process whereby the proteins in solution aggregate or coagulate when in solutions containing high concentrations of salt. Consequently, alternative means are provided for ensuring the immunogenic compositions of the invention have a pharmaceutically acceptable osmolality.

40 [0058] In a particular embodiment there are provided immunogenic compositions further comprising a non-ionic tonicity agent. A non-ionic tonicity agent for use in an immunogenic composition will itself need to be pharmaceutically acceptable, e.g. suitable for use in humans, as well as being compatible with the M72 related antigen and further compatible with other components such as the immunostimulant(s).

45 [0059] In one embodiment of the present invention, suitable non-ionic tonicity agents are polyols, sugars (in particular sucrose, fructose, dextrose or glucose) or amino acids such as glycine. In one embodiment the polyol is a sugar alcohol, especially a C3-6 sugar alcohol. Exemplary sugar alcohols include glycerol, erythritol, threitol, arabitol, xylitol, ribitol, sorbitol, mannitol, dulcitol and iditol. In a specific example of this embodiment, a suitable non-ionic tonicity agent is sorbitol. The skilled person will recognise that an appropriate osmolality may be attained through the use of a mixture of different tonicity agents. In a particular embodiment of the invention the non-ionic tonicity agent in the compositions of the invention incorporates sucrose and/or sorbitol.

50 [0060] In one embodiment, a suitable concentration of polyol within the immunogenic composition is between about 2.5 and about 15% (w/v), in particular between about 2.5 and about 10% (w/v) for example between about 3 and about 7% (w/v), such as between about 4 and about 6% (w/v). In a specific example of this embodiment, the polyol is sorbitol.

55 [0061] In another embodiment, the immunogenic composition comprises sucrose and sorbitol. In such circumstances the immunogenic composition may suitably contain between about 2.5 and about 15% (w/v) of sucrose and between about 2.5 and about 15% (w/v) of sorbitol, in particular between about 2.5 and about 10% (w/v) of sucrose and between about 2.5 and about 10% (w/v) of sorbitol, for example, between about 3 and about 7% (w/v) of sucrose and between about 3 and about 7% (w/v) of sorbitol, such as between about 4 and about 6% (w/v) of sucrose and between about 4 and about 6% (w/v) of sorbitol.

[0062] The pH of the immunogenic compositions should be suitable for parenteral administration. Suitably the pH will be in the range 7.25 to 8.75, such as 7.5 to 8.5, in particular pH 7.75 to 8.25. A pH of about 8.0 is of particular interest.

[0063] The pH may be controlled by the use of buffers, including for example Tris or phosphate buffers.

[0064] In a particular embodiment of the invention, the immunogenic composition comprises one or more immunostimulants.

[0065] In one embodiment, the immunostimulant may be a saponin. A particularly suitable saponin for use in the present invention is Quil A and its derivatives. Quil A is a saponin preparation isolated from the South American tree *Quillaja saponaria* Molina and was first described by Dalsgaard et al. in 1974 ("Saponin adjuvants", Archiv. für die gesamte Virusforschung, Vol. 44, Springer Verlag, Berlin, p243-254) to have adjuvant activity. Purified fractions of Quil A have been isolated by HPLC which retain adjuvant activity without the toxicity associated with Quil A (WO88/09336), for example QS7 and QS21 (also known as QA7 and QA21). QS21 is a natural saponin derived from the bark of *Quillaja saponaria* Molina, which induces CD8+ cytotoxic T cells (CTLs), Th1 cells and a predominant IgG2a antibody response. QS21 is a preferred saponin in the context of the present invention.

[0066] In a suitable form of the present invention, the saponin adjuvant within the immunogenic composition is a derivative of *saponaria* Molina Quil A, in particular an immunologically active fraction of Quil A, such as QS17 or QS21, suitably QS21.

[0067] Desirably, QS21 is provided in a less reactogenic composition where it is quenched with an exogenous sterol, such as cholesterol for example. Several particular forms of less reactogenic compositions wherein QS21 is quenched with cholesterol exist. In a specific embodiment, the saponin/sterol is in the form of a liposome structure (such as described in WO96/33739, Example 1). In this embodiment the liposomes suitably contain a neutral lipid, for example phosphatidylcholine, which is suitably non-crystalline at room temperature, for example egg yolk phosphatidylcholine, dioleoyl phosphatidylcholine (DOPC) or dilauryl phosphatidylcholine. The liposomes may also contain a charged lipid which increases the stability of the liposome-QS21 structure for liposomes composed of saturated lipids. In these cases the amount of charged lipid is suitably 1-20% w/w, such as 5-10%. The ratio of sterol to phospholipid is 1-50% (mol/mol), suitably 20-25%.

[0068] Suitable sterols include β -sitosterol, stigmasterol, ergosterol, ergocalciferol and cholesterol. In one particular embodiment, the immunogenic composition comprises cholesterol as sterol. These sterols are well known in the art, for example cholesterol is disclosed in the Merck Index, 11th Edn., page 341, as a naturally occurring sterol found in animal fat.

[0069] Where the active saponin fraction is QS21, the ratio of QS21:sterol will typically be in the order of 1:100 to 1:1 (w/w), suitably between 1:10 to 1:1 (w/w), and especially 1:5 to 1:1 (w/w). Suitably excess sterol is present, the ratio of QS21:sterol being at least 1:2 (w/w). In one embodiment, the ratio of QS21:sterol is 1:5 (w/w). The sterol is suitably cholesterol.

[0070] In another embodiment, the immunogenic composition comprises an immunostimulant which is a Toll-like receptor 4 (TLR4) agonist. By "TLR agonist" it is meant a component which is capable of causing a signaling response through a TLR signaling pathway, either as a direct ligand or indirectly through generation of endogenous or exogenous ligand (Sabroe et al, J Immunol 2003 p1630-5). A TLR4 agonist is capable of causing a signaling response through a TLR-4 signaling pathway. A suitable example of a TLR4 agonist is a lipopolysaccharide, suitably a non-toxic derivative of lipid A, particularly monophosphoryl lipid A or more particularly 3-de-O-acylated monophosphoryl lipid A (3D-MPL).

[0071] 3D-MPL is sold under the name MPL by GlaxoSmithKline Biologicals N.A. and is referred throughout the document as MPL or 3D-MPL see, for example, US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094. 3D-MPL primarily promotes CD4+ T cell responses with an IFN-gamma (Th1) phenotype. 3D-MPL can be produced according to the methods disclosed in GB2220211A. Chemically it is a mixture of 3-de-O-acylated monophosphoryl lipid A with 3, 4, 5 or 6 acylated chains. In the compositions of the present invention small particle 3D-MPL may be used to prepare the immunogenic composition. Small particle 3D-MPL has a particle size such that it may be sterile-filtered through a 0.22 μ m filter. Such preparations are described in WO94/21292. Suitably, powdered 3D-MPL is used to prepare the immunogenic compositions of the present invention.

[0072] Other TLR4 agonists which can be used are alkyl glucosaminide phosphates (AGPs) such as those disclosed in WO98/50399 or US Patent No. 6,303,347 (processes for preparation of AGPs are also disclosed), suitably RC527 or RC529 or pharmaceutically acceptable salts of AGPs as disclosed in US Patent No. 6,764,840. Some AGPs are TLR4 agonists, and some are TLR4 antagonists.

[0073] Other suitable TLR4 agonists are as described in WO2003/011223 and in WO2003/099195, such as compound I, compound II and compound III disclosed on pages 4-5 of WO2003/011223 or on pages 3-4 of WO2003/099195 and in particular those compounds disclosed in WO2003/011223 as ER803022, ER803058, ER803732, ER804053, ER804057m ER804058, ER804059, ER804442, ER804680 and ER804764. For example, one suitable TLR-4 agonist is ER804057.

[0074] In a particular embodiment, the immunogenic composition comprises both a saponin and a TLR4 agonist. In a specific example, the immunogenic composition comprises QS21 and 3D-MPL.

[0075] A TLR-4 agonist, such as a lipopolysaccharide, such as 3D-MPL, can be used at amounts between 1 and 100

ug per human dose of the immunogenic composition. 3D-MPL may be used at a level of about 50 ug, for example between 40 to 60 ug, suitably between 45 to 55 ug or between 49 and 51 ug or 50 ug. In a further embodiment, the human dose of the immunogenic composition comprises 3D-MPL at a level of about 25 ug, for example between 20 to 30 ug, suitable between 21 to 29 ug or between 22 to 28 ug or between 23 and 27 ug or between 24 and 26 ug, or 25 ug.

5 **[0076]** A saponin, such as QS21, can be used at amounts between 1 and 100 ug per human dose of the immunogenic composition. QS21 may be used at a level of about 50 ug, for example between 40 to 60 ug, suitably between 45 to 55 ug or between 49 and 51 ug or 50 ug. In a further embodiment, the human dose of the immunogenic composition comprises QS21 at a level of about 25 ug, for example between 20 to 30 ug, suitable between 21 to 29 ug or between 22 to 28 ug or between 23 and 27 ug or between 24 and 26 ug, or 25 ug.

10 **[0077]** Where both TLR4 agonist and saponin are present in the immunogenic composition, then the weight ratio of TLR4 agonist to saponin is suitably between 1:5 to 5:1, suitably between 1:2 to 2:1, such as about 1:1. For example, where 3D-MPL is present at an amount of 50 ug or 25 ug, then suitably QS21 may also be present at an amount of 50 ug or 25 ug, respectively, per human dose of the immunogenic composition. Certain immunogenic compositions of the present invention comprise QS21 and 3D-MPL, at an amount of between 1 and 100 ug of each per human dose, such as at an amount of between 10 and 75 ug of each per human dose. Immunogenic compositions of the present invention may suitably comprise QS21 and 3D-MPL, at an amount of between 15 and 35 ug of each per human dose, such as at an amount of between 20 and 30 ug of each per human dose.

15 **[0078]** In one embodiment, the immunostimulant is a TLR9 agonist, for example as set out in WO2008/142133. In a specific example, said TLR9 agonist is an immunostimulatory oligonucleotide, in particular an oligonucleotide containing an unmethylated CpG motif. Such oligonucleotides are well known and are described, for example, in WO96/02555, WO99/33488 and US5,865,462. Suitable TLR9 agonists for use in the immunogenic compositions described herein are CpG containing oligonucleotides, optionally containing two or more dinucleotide CpG motifs separated by at least three, suitably at least six or more nucleotides. A CpG motif is a cytosine nucleotide followed by a guanine nucleotide.

20 **[0079]** In one embodiment the internucleotide bond in the oligonucleotide is phosphorodithioate, or possibly a phosphorothioate bond, although phosphodiester and other internucleotide bonds could also be used, including oligonucleotides with mixed internucleotide linkages. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US5,666,153, US5,278,302 and WO95/26204. Oligonucleotides comprising different internucleotide linkages are contemplated, e.g. mixed phosphorothioate phosphodiesters. Other internucleotide bonds which stabilise the oligonucleotide may be used.

25 **[0080]** Examples of CpG oligonucleotides suitable for inclusion in the immunogenic compositions described herein have the following sequences. In one embodiment, these sequences contain phosphorothioate modified internucleotide linkages.

OLIGO 1 (SEQ ID No: 9): TCC ATG ACG TTC CTG ACG TT (CpG 1826)

30 OLIGO 2 (SEQ ID No: 10): TCT CCC AGC GTG CGC CAT (CpG 1758)

OLIGO 3 (SEQ ID No: 11): ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG

OLIGO 4 (SEQ ID No: 12): TCG TCG TTT TGT CGT TTT GTC GTT (CpG 2006)

OLIGO 5 (SEQ ID No: 13): TCC ATG ACG TTC CTG ATG CT (CpG 1668)

35 **[0081]** Alternative CpG oligonucleotides may comprise the sequences above in that they have inconsequential deletions or additions thereto.

[0082] In one embodiment the immunostimulant is a tocol. Tocols are well known in the art and are described in EP0382271. In a particular embodiment, the tocol is alpha-tocopherol or a derivative thereof such as alpha-tocopherol succinate (also known as vitamin E succinate).

40 **[0083]** The immunogenic compositions according the invention may be used in medicine, in particular for the prophylaxis, treatment or amelioration of infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*. The immunogenic compositions will generally be provided for administration to humans, though they may also be of value in veterinary medicine such as for administration to bovines.

45 **[0084]** There is provided the use of an immunogenic composition according the invention in the manufacture of a medicament, in particular a medicament for the prophylaxis, treatment or amelioration of infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*.

[0085] The immunogenic composition may be provided for the purpose of:

- treating active tuberculosis;
- 50 - prophylaxis of active tuberculosis, such as by administering to a subject who is uninfected, or alternatively a subject who has latent infection;
- treating latent tuberculosis;
- prophylaxis of latent tuberculosis, such as by administering to a subject who is uninfected; or

- preventing or delaying reactivation of tuberculosis, especially the delay of TB reactivation, for example by a period of months, years or even indefinitely.

[0086] The term "active infection" refers to an infection, e.g. infection by *M. tuberculosis*, with manifested disease symptoms and/or lesions, suitably with manifested disease symptoms.

[0087] The terms "inactive infection", "dormant infection" or "latent infection" refer to an infection, e.g. infection by *M. tuberculosis*, without manifested disease symptoms and/or lesions, suitably without manifested disease symptoms. A subject with latent infection will suitably be one which tests positive for infection, e.g. by PPD or T cell based assays, but which has not demonstrated the disease symptoms and/or lesions which are associated with an active infection.

[0088] The term "primary tuberculosis" refers to clinical illness, e.g., manifestation of disease symptoms, directly following infection, e.g. infection by *M. tuberculosis*. See, Harrison's Principles of Internal Medicine, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

[0089] The terms "secondary tuberculosis" or "postprimary tuberculosis" refer to the reactivation of a dormant, inactive or latent infection, e.g. infection by *M. tuberculosis*. See, Harrison's Principles of Internal Medicine, Chapter 150, pp. 953-966 (16th ed., Braunwald, et al., eds., 2005).

[0090] The term "tuberculosis reactivation" refers to the later manifestation of disease symptoms in an individual that tests positive for infection (e.g. in a tuberculin skin test, suitably in an *in vitro* T cell based assay) test but does not have apparent disease symptoms. Suitably the individual will not have been re-exposed to infection. The positive diagnostic test indicates that the individual is infected, however, the individual may or may not have previously manifested active disease symptoms that had been treated sufficiently to bring the tuberculosis into an inactive or latent state.

[0091] Suitably an immunogenic composition is administered to a subject who is uninfected or who has a latent infection by mycobacteria, such as infection by *Mycobacterium tuberculosis*.

[0092] The volume of immunogenic composition administered may vary depending upon a number of other factors, such as the specific delivery route, e.g. intramuscular, subcutaneous or intradermal. Typically, the volume administered in a single injection (the unit dose) for a human will be in the range of 50 ul to 1 ml, such as 100 ul to 750 ul, especially 400 to 600 ul, for example about 500 ul.

[0093] The quantity of M72 related antigen contained within a single dose is dependent upon clinical needs but a single human dose will typically be in the range of 1 to 100 ug, such as 5 to 50 ug, for example 5 to 20 ug. A single human dose may contain about 10 ug of M72 related antigen.

[0094] Suitably, compositions of the invention will be stable, in which is meant that during storage at 25 °C for a period of 24 hours antigenicity as measured by the techniques described herein remains at least 80% of the antigenicity before storage. Desirably, antigenicity will remain at least 85%, such as at least 90% and in particular at least 95% after storage at 25 °C for a period of 24 hours. For compositions of particular interest, at least 80% of the antigenicity of the composition, such as at least 85%, at least 90% and especially at least 95% remains after storage at 30 °C for a period of 24 hours.

[0095] The present invention will now be further described by means of the following non-limiting examples.

EXAMPLES

Example 1: Preparation of adjuvant composition ASA (sorbitol)

[0096] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21 in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

[0097] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A in organic solvent was dried down under vacuum. An aqueous solution (phosphate buffered saline [100 mM NaCl, 50 mM Phosphate pH 6.1]) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposome size was reduced to around 90 nm ± 10 nm measured by DLS. Liposomes were then sterile filtered.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0098] Na₂/K Phosphate buffer 100 mM pH 6.1 when diluted 10 times was added to water for injection to reach a 10 mM phosphate buffer concentration in the final formulation. A 30% (w/v) sorbitol solution in water for injection (WFI) was then added to reach a concentration of 4.7% in the final formulation - this was stirred for 15 to 45 minutes at room

EP 2 651 437 B1

temperature.

[0099] Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 100 ug/ml of 3D-MPL in the final formulation.

[0100] The mixture was subsequently stirred for 15 to 45 minutes at room temperature.

5

Step 2: QS21 addition

[0101] Using a peristaltic pump, QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 100 ug/ml concentration in the final formulation. The mix was stirred for 15 to 45 minutes.

10

[0102] Final ASA(sorbitol) formulation contained 2 mg DOPC, 500 ug cholesterol, 100 ug 3D-MPL/ml and 100 ug QS21/ml, 4.7% sorbitol and 5 mM sodium chloride and 10 mM phosphate.

Step 3: pH was checked to be 6.1 ± 0.1

15

Step 4: Sterile filtration

[0103] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at $+2^{\circ}\text{C}$ to $+8^{\circ}\text{C}$.

20

[0104] The adjuvant composition obtained, which comprised 3-de-O-acylated MPL and QS21 in a liposomal formulation and containing sorbitol as a tonicity agent (designated ASA (sorbitol)), was then stored at 4°C .

Example 2: Preparation of adjuvant composition ASA (150 mM NaCl)

25

[0105] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21 in a liposomal formulation using sodium chloride as a tonicity agent.

A. Method of preparation of liposomes:

30

[0106] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A (3D-MPL) in organic solvent was dried down under vacuum. Phosphate buffered saline (100 mM NaCl, 50 mM Phosphate pH 6.1) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around $90 \text{ nm} \pm 10 \text{ nm}$ measured by DLS. Liposomes were then sterile filtered on 0.22 um PES membrane.

35

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

40

[0107] Na_2/K Phosphate buffer 100 mM pH 6.45 when diluted 10 times and NaCl 1.5 M were added to water for injection to reach respectively 10 mM phosphate and NaCl 150 mM concentrations in the final formulation. This mixture was stirred for 5 minutes at room temperature. Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 100 ug/ml of 3D-MPL in the final formulation. The mixture was subsequently stirred for 5 to 15 minutes at room temperature.

45

Step 2: QS21 addition

[0108] QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 100 ug/ml concentration in the final formulation. The mix was stirred at room temperature.

50

Step 3: pH was checked so as to be 6.1 ± 0.1 .

Step 4: Sterile filtration

55

[0109] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

EP 2 651 437 B1

Step 5: Storage at +2°C to +8°C

[0110] Final composition of ASA(150 mM NaCl) was 2 mg DOPC, 500 ug cholesterol, 100 ug 3-de-O-acylated MPL, 100 ug QS21 per 1 ml, with 10 mM phosphate and 150 mM NaCl.

5

Example 3: QS21 Lytic Activity

[0111] QS21 is known to lyse red blood cells (RBC). The ASA (sorbitol) adjuvant composition prepared as in Example 1 was tested to ensure that QS21 lytic activity was quenched in the same way as was seen with the equivalent adjuvant composition comprising 150 mM NaCl (ASA (150 mM NaCl)).

10

[0112] QS21 lytic activity was measured by a haemolysis assay using chicken Red Blood cells (RBC). RBC were centrifuged at 550 g at 4°C. Supernatant was discarded. The pellet was carefully resuspended in PBS buffer to reach the initial volume and the same operation was repeated until supernatant was no longer red (generally 3 times). The pellet was stored at 4°C for 3 to 4 days maximum if not used directly (and washed again the day it is used) or was diluted around 10 times in buffer if used the same day.

15

[0113] A QS21 dose range curve was prepared in ASA buffer (in salt or in sorbitol buffer following the ASA sample tested) extemporaneously and the adjuvant samples (containing a 50 ug or 90 ug equivalent of QS21 meaning the equivalent of 500 ul or 900 ul ASA) were prepared. Final volume was adjusted to 900 ul in standards and samples with adequate buffer (containing or not sorbitol as a function of the buffer of the sample tested). Due to its opalescence, ASA interferes with optical density (OD). ASA "blanks" were thus prepared and their OD was subtracted from the OD of ASA tested samples. Those blanks corresponded to the same ASA volume as the volume tested in samples, but adjusted to 1 ml with buffer. No RBC were added to these blanks. Standards and samples were then incubated with RBC (100 ul of diluted RBC added to 900 ul of standards and samples) for 30 minutes at room temperature (RT). Samples were then centrifuged 5 minutes at 900 g. Optical density at 540 nm was measured after centrifugation.

20

25

[0114] Determination of lytic activity was carried out by a limit test.

1. Limit of detection (LOD) was defined as the lowest concentration of QS21 leading to an OD:

- Higher than the base level (OD>0.1)
- Around three times higher than OD's buffer (the "0 ug" QS21)
- In the ascendant part of the curve
- Determined for each test.

30

2. QS21 lytic activity was held to be positive in the adjuvant samples if the OD for the adjuvant sample was greater than the OD_{LOD}.

35

Example QS21 curve

ug QS21	OD	QS21 quenched
0	0.029	NA
0.5	0.052	< LOD
0.6	0.073	< LOD
0.7	0.091	< LOD
0.8	0.096	< LOD
0.9	0.12	> 98.2%
1	0.195	> 98%
1.1	0.212	> 97.8%
1.2	0.348	> 97.6%
1.3	0.479	> 97.4%
1.4	0.612	> 97.2%
1.5	0.669	> 97%
2	1.139	> 96%
2.5	1.294	> 95%
3	1.391	> 94%
5	1.416	> 90%

40

45

50

55

EP 2 651 437 B1

(continued)

ug QS21	OD	QS21 quenched
Adjuvant *	0.03	> 98.2 %
*50ug QS21 equivalent tested. 150 mM sodium chloride buffer.		

[0115] The above data is shown graphically in Figure 1.

[0116] The Limit of Detection in this assay is at 0.9 ug QS21, and OD of 0.12

[0117] The QS21 quenching in an adjuvant composition comprising 150 mM sodium chloride was estimated to be more than 98.2% for the equivalent of 50 ug QS21 tested. In the case of an equivalent of 90 ug tested, conclusion is more than 99%.

[0118] QS21 quenching was then compared with an equivalent adjuvant composition comprising sorbitol and only 5 mM sodium chloride. Data were generated after storage of the ASA at 4°C or after accelerated stability (7 days at 37°C). For the ASA in sorbitol, the QS21 standard curve was realised in a sorbitol containing buffer.

Sample	Timepoint	LOD	QS21 quenched
Adjuvant composition (ASA) 150 mM NaCl	T0	< 1.4	> 97.2%
	7 days 37°C	< 0.9	> 98.2%
Adjuvant Composition (ASA) sorbitol, 5 mM NaCl	T0	< 2	> 97.8%
	7 days 37°C	<1	> 96%
	11 months 4°C	<2	> 97.8%*
	Equivalent of 50 ug QS21 tested except * equivalent of 90 ug QS21 tested.		

[0119] It was concluded that QS21 was adequately quenched in a low sodium chloride buffer.

Example 4: MPL congeners.

[0120] Chemically, 3D-MPL is a mixture of 3-de-O-acylated monophosphoryl lipid A with mainly 4, 5 or 6 acylated chains. Each separate 3D-MPL molecule is called a congener. It is important that the congener composition remain constant, with no shift between the proportion of congeners. It is also important that any buffer used enables the congener composition to be the same as in the concentrated liposomes used to make the adjuvant composition.

[0121] As shown on Figure 2, the congener composition was examined in 3D-MPL concentrated liposomes (Conc. Liposomes LIP07-217, first column of Figure 2), an adjuvant composition comprising 3D-MPL liposomes and QS21 in a 150 mM NaCl buffer (Adjuvant 150 mM NaCl, or ASA (150 mM NaCl), second column), and an adjuvant composition comprising 3D-MPL liposomes and QS21 in a sorbitol and 5 mM NaCl buffer (Adjuvant Sorbitol, or ASA (sorbitol), columns 3-7).

[0122] The congener composition was also examined in two lots of ASA (sorbitol) adjuvant at day 0 and 7 days after preparation and maintenance at 37 °C to ensure that there was no evolution over time (see final four columns of Figure 2).

[0123] Relative distribution of tetra-, penta- and hexa-acylated congeners of MPL in concentrated liposomes or ASA (sorbitol) samples was determined by IP-HPLC-Fluo detection (ARD). Both standards and samples were derivatised with dansylhydrazine, which introduces a Fluo-active chromophore on the disaccharide backbone. The derivatised samples were analysed on a C18 reverse phase column using tetrabutylammonium hydroxide (TBAOH) as an ion pair reagent. Congeners containing the same numbers of fatty acyl groups were eluted in distinct groups (tetraacyl, pentaacyl, and hexaacyl). Distribution of congeners is deduced by comparing the peak area of each group to the total peak area of all MPL congeners.

[0124] Figure 2 shows the percentage of each congener. No significant difference in congener composition was found between adjuvant buffers, and the congener composition was consistent over time in the sorbitol buffer.

Example 5: Preparation of adjuvant composition ASA (sorbitol - 2)

[0125] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 1, in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

EP 2 651 437 B1

The adjuvant was prepared by 1:1 dilution of ASA(sorbitol), prepared according to Example 1, with a solution containing 10 mM phosphate, 5 mM NaCl, 4.7% sorbitol at pH 6.1.

Final ASA(sorbitol - 2) formulation contained 1 mg DOPC, 250 ug cholesterol, 50 ug 3D-MPL/ml and 50 ug QS21/ml, 4.7% sorbitol, 5 mM sodium chloride and 10 mM phosphate.

Example 6: Preparation of adjuvant composition ASA (sorbitol - 3)

[0126] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 1, in a liposomal formulation using sorbitol as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

[0127] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A in organic solvent was dried down under vacuum. An aqueous solution (phosphate buffered saline [100 mM NaCl, 50 mM Phosphate pH 6.1]) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around 90 nm \pm 10 nm measured by DLS. Liposomes were then sterile filtered.

B. ASA formulation:

Step 1: Dilution of concentrated liposomes

[0128] Na₂/K Phosphate buffer 100 mM pH 6.1 when diluted 10 times was added to water for injection to reach a 10 mM phosphate buffer concentration in the final formulation. A 30% (w/v) sorbitol solution in water for injection (WFI) was then added to reach a concentration of 4.7% in the final formulation - this was stirred for 15 to 45 minutes at room temperature.

[0129] Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 50 ug/ml of 3D-MPL in the final formulation.

[0130] The mixture was subsequently stirred for 15 to 45 minutes at room temperature.

Step 2: QS21 addition

[0131] Using a peristaltic pump, QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 50 ug/ml concentration in the final formulation. The mix was stirred for 15 minutes.

[0132] Final ASA formulation contained 1 mg DOPC, 250 ug cholesterol, 50 ug 3D-MPL/ml and 50 ug QS21/ml, 4.7% sorbitol and 2.5 mM sodium chloride, 10 mM phosphate.

Step 3: pH was checked to be 6.1 \pm 0.1

Step 4: Sterile filtration

[0133] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C.

[0134] The adjuvant composition obtained, which comprised 3-de-O-acylated MPL and QS21 in a liposomal formulation and containing sorbitol as a tonicity agent (designated ASA (sorbitol-3)), was then stored at 4°C.

Example 7: Preparation of adjuvant composition ASA (150 mM NaCl - 2)

[0135] An adjuvant composition was prepared which comprised 3-de-O-acylated monophosphoryl lipid A and QS21, at a reduced level relative to Example 2, in a liposomal formulation using sodium chloride as a tonicity agent. This was prepared as follows:

A. Method of preparation of liposomes:

5 [0136] A mixture of lipid (DOPC), cholesterol and 3-de-O-acylated monophosphoryl lipid A (3D-MPL) in organic solvent was dried down under vacuum. Phosphate buffered saline (100 mM NaCl, 50 mM Phosphate pH 6.1) was then added and the vessel agitated until all the lipid was in suspension. This suspension was then prehomogenised with high shear mixer and then high pressure homogenised until the liposomes size was reduced to around 90 nm \pm 10 nm measured by DLS. Liposomes were then sterile filtered on 0.22 μ m PES membrane.

B. ASA formulation:Step 1: Dilution of concentrated liposomes

10 [0137] Na₂/K Phosphate buffer 100mM pH 6.45 when diluted 10 times and NaCl 1.5 M were added to water for injection to reach respectively 10 mM phosphate and NaCl 150 mM concentrations in the final formulation. This mixture was stirred for 5 minutes at room temperature. Concentrated liposomes (made of DOPC, cholesterol and 3D-MPL at 40 mg/ml, 10 mg/ml and 2 mg/ml respectively) were then added to the mix to reach a concentration of 50 μ g/ml of 3D-MPL in the final formulation. The mixture was subsequently stirred for 5 to 15 minutes at room temperature.

Step 2: QS21 addition

20 [0138] QS21 bulk stock was added to the diluted liposomes under magnetic stirring to reach a 50 μ g/ml concentration in the final formulation. The mix was stirred at room temperature.

25 Step 3: pH was checked so as to be 6.1 \pm 0.1.

Step 4: Sterile filtration

[0139] Sterile filtration was performed using a polyethersulfone (PES) filter from PALL Corporation.

Step 5: Storage at +2°C to +8°C

30 [0140] Final composition of ASA(150 mM NaCl - 2) was 1 mg DOPC, 250 μ g cholesterol, 50 μ g 3-de-O-acylated MPL, 50 μ g QS21 per 1 ml, 10 mM phosphate and 150 mM NaCl.

Example 8: Preparation of Protein Antigens**M72 with two N-terminal His residues (SEQ ID No: 3)**Construction of the M72 expression Vector

40 [0141] A plasmid coding for the amino acid sequence of Mtb72f with an additional 6-His tag at the N-terminus was generated by the sequential linkage in tandem of the open reading frames (ORFs) encoding the C terminal fragment of Mtb32a to the full length ORF of Mtb39a followed at the C terminus with the N terminal portion of Mtb32a. This was accomplished by using sequence-specific oligonucleotides containing unique restriction sites (EcoRI and EcoRV) and devoid of the stop codons at the C terminal ends (in the case of the C terminal fragment of Mtb32a and Mtb39a) for polymerase chain reaction (PCR) of genomic DNA from the *M. tuberculosis* strain H37Rv. Using this vector as template, a mutation of Ser706 to Ala was performed by site-directed mutagenesis. The proper orientation of inserts as well as the mutation Ser706Ala was verified by DNA sequencing.

45 [0142] In order to obtain the vector coding for M72, which just has 2 His residues at the N terminus, four His were deleted making use of a commercial site-directed mutagenesis system. After sequence verification, the M72 coding sequence was excised from the plasmid by enzymatic reaction, gel purified and ligated into a pET vector. The recombinant plasmid was then sequence verified. This plasmid codes for M72 under the control of a T7 promoter. Expression of T7 RNA polymerase is driven from a genomic integrant in the expression host and is induced using a lac operon-based system (lacI) and an IPTG chemical induction signal. The expression plasmid is provided with kanamycin resistance.

55 [0143] The plasmid coding for the M72 fusion protein under the control of a T7 promoter was transformed into the HMS174 (DE3) strain of *E.coli*, using an electroporation method. The coding sequence of the M72 insert and the flanking regions were sequenced on both strands and were found to be identical to the sequence determined from the original plasmid construct.

EP 2 651 437 B1

Fermentation

[0144] A vial of pelleted working seed was thawed at room temperature. A pre-dilution was prepared by mixing the working seed with 4.9 ml of pre-culture medium. 1 ml of the pre-dilution was used to inoculate the liquid pre-culture which consists of 400 ml of pre-culture medium supplemented with 50 mg/l kanamycin sulfate and 10 g/l glucose.

Pre-culture medium composition

Ingredient	Concentration		
KH ₂ PO ₄	14.83 g/l		
K ₂ HPO ₄	1.65 g/l		
(NH ₄) ₂ SO ₄	5.82 g/l		
Yeast extract	6.21 g/l		
Glycerol 87% (w/w)	14.54 ml/l		
Metal and salt solution (1): - FeCl ₃ 6H ₂ O - MgSO ₄ 7H ₂ O - Micro element solution (2): - ZnSO ₄ 7H ₂ O - MnSO ₄ H ₂ O - CuSO ₄ 5H ₂ O - CoCl ₂ 6H ₂ O - H ₃ BO ₃ - Na ₂ MoO ₄ 2H ₂ O - HCl 4N	9.7 ml/l	3.3 g/l 58 g/l 116 ml/l	7.65 g/l 5.28 g/l 1.1 g/l 1.1 g/l 0.3 g/l 2.64 g/l 6.2 ml/l
Biotine and CaCl ₂ solution (2): - Biotine - CaCl ₂ 2 H ₂ O	0.97 ml/l	0.05 g/l 61.7 g/l	
pH of the medium is adjusted to 6.5 with NaOH (25%) solution The medium is filtered through 0.22 um			
(1) pH adjusted to 1.50 with HCl (37%) solution; the solution is filtered through 0.22 um (2) The solution is filtered through 0.22 um			

[0145] The pre-culture was incubated in a 2 litre shake flask at 30°C under agitation (200 RPM) until the OD_{650nm} reached a value between 2 and 4 (approximate incubation time: 16 hours). At that stage, a 72 litre (total volume) fermenter containing 45 litres of culture medium supplemented with 34 mg/l kanamycin sulfate was inoculated with 52 ml liquid pre-culture.

Culture medium composition

Ingredient	Concentration		
MgSO ₄ 7H ₂ O	0.63 g/l		
FeCl ₃ 6H ₂ O Micro element solution (1): ZnSO ₄ 7H ₂ O - MnSO ₄ H ₂ O - CuSO ₄ 5H ₂ O - CoCl ₂ 6H ₂ O - H ₃ BO ₃ - Na ₂ MoO ₄ 2H ₂ O - HCl 4N	0.056 g/l 1.91 ml/l		7.65 g/l 5.28 g/l 1.1 g/l 1.1 g/l 0.3 g/l 2.64 g/l 6.2 ml/l

EP 2 651 437 B1

(continued)

Ingredient	Concentration	
HCl 37%	0.40 mL/L	
Yeast extract	35 g/L	
(NH ₄) ₂ SO ₄	2.10 g/l	
KH ₂ PO ₄	18.70 g/l	
Sodium glutamate	2.5 g/l	
Glycerol 87%	0.276 ml/l	
Glucose	20 g/l	
Biotine solution (2):	0.22 ml/l	
- Biotine		1 g/l
CaCl ₂ 2 H ₂ O	0.21 g/l	
The solution is filtered through 0.22 um		
(1) The solution is filtered through 0.22 um		
(2) pH adjusted to 11.0 with NaOH (25%) solution; the solution is filtered through 0.22 um		

[0146] During the growth phase, pH was maintained at 6.8 ± 0.2 by periodic addition of 25% (v/v) NH₄OH and 25% (v/v) H₃PO₄. After incubation for 16 hours at 30°C, fed-batch was started with feed medium.

Feed medium composition

Ingredient	Concentration	
MgSO ₄ 7H ₂ O	1.98 g/l	
Ingredient	Concentration	
FeCl ₃ 6H ₂ O	0.178 g/l	
Micro element solution (1):	6.02 ml/l	
- ZnSO ₄ 7H ₂ O		7.65 g/l
- MnSO ₄ H ₂ O		5.28 g/l
- CuSO ₄ 5H ₂ O		1.1 g/l
- CoCl ₂ 6H ₂ O		1.1 g/l
- H ₃ BO ₃		0.3 g/l
- Na ₂ MoO ₄ 2H ₂ O		2.64 g/l
- HCl 4N		6.2 ml/l
HCl 37%	1.24 ml/l	
Sodium glutamate	5 g/l	
Yeast extract	40 g/l	
Glycerol 87%	590 ml/l	
Biotine solution (2):	2 ml/l	
- Biotine		1 g/l
CaCl ₂ 2 H ₂ O	0.66 g/l	
The solution is filtered through 0.22 um		
(1) The solution is filtered through 0.22 um		
(2) pH adjusted to 11.0 with NaOH (25%) solution; the solution is filtered through 0.22 um		

[0147] The temperature was maintained at 30°C for a further 2 hours, then raised to 37°C until the end of fermentation.

EP 2 651 437 B1

The air flow was constantly set to 75 l/min and the dissolved oxygen kept at 17% saturation by feedback control of the agitation and pressure. Small quantities of antifoam solution were added on demand automatically. By the time the OD_{650nm} reached a value of 50 (± 5), 1 mM Isopropylbeta-D-thiogalactopyranoside (IPTG) was added in order to induce the expression of M72. Fermentation ended after 5 hours from the time point induction was started. The cell culture was cooled down to 15°C under slight agitation and centrifuged (at 4°C) to obtain cell pellets which were thereafter stored at -20°C in aliquots.

Isolation of Inclusion Bodies

[0148] The cell pellets collected from the harvest were thawed at room temperature and disrupted in lysis buffer (10 mM Tris, 50 mM NaCl, pH 8.0) with a high pressure homogenizer. Thereafter the cell lysate was centrifuged and the resulting cell pellets (or inclusion bodies, IBs) were washed with wash buffer containing urea, Tris and NaCl. The IBs were solubilised with solubilisation buffer containing 8 M urea and filtered through a 0.2 μ m membrane. This filtered solution was first purified by anion exchange chromatography using a Q Sepharose Fast Flow (QSFF) column. The elution of M72 takes place with a 6 M urea, 20 mM bis-Tris propane, 90 mM NaCl, pH 7.0 solution.

[0149] M72 collected was further purified by Hydroxyapatite chromatography (HA), from which it is eluted with a 6 M urea, 20 mM bis-Tris propane, 250 mM NaCl, pH 7.0 solution. The collected fraction was concentrated with a 30 kDa membrane cassette and diafiltered against 20 mM Tris, pH 7.5. M72 was then sterilised through a 0.22 μ m filter. The purified bulk was then aliquoted and stored at -70°C.

Example 9: Investigation of "salting out" in compositions comprising M72 with different salt concentrations at pH 6.1, 7.5 and 8.5.

[0150] The impact of sodium chloride concentration and pH on M72 antigen stability, as assessed by size and antigenicity, was investigated.

Method

[0151] Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was diluted to a concentration of 100 μ g/ml in three different buffers (10mM phosphate buffer at pH 6.1, 20 mM Tris buffer at pH 7.5 and 20 mM Tris buffer at pH 8.5) containing final sodium chloride concentrations of 0, 50, 150, 300 and 450 mM.

[0152] Samples were analysed immediately (TO), stored overnight at 4°C before analysis (TO O/N) or stored at 25°C for 24 hours before analysis (T24h25°C).

[0153] DLS was performed using a Malvern Zetasizer Nano ZS from Malvern Instruments (UK). The instrument was operated using a laser wavelength of 633 nm and power of 4 mW. Scattered light was detected at 173° at a temperature of 22°C. The Z-average diameter (Zav) and polydispersity index (pl) are calculated by the instrument software.

[0154] Nephelometry was performed using a Nepheloskan® Ascent, available from Thermo Fischer Scientific. Analysis was performed in UV transparent Costar® micro-plates available from Corning Inc (USA).

[0155] Antigenicity was quantified by a sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0156] The findings of this experiment are presented in Figures 3 to 5.

[0157] The results demonstrate for the first time that the stability of solutions containing an M72 related antigen is sensitive to both pH and sodium chloride concentration. The impact of sodium chloride on antigen size and antigenicity is all the more notable as the pH is lower.

[0158] Antigen size and antigenicity are not stable at pH 6.1 even in the absence of sodium chloride. The addition of 50 mM sodium chloride at pH 6.1 led to a size increase from 35 nm (0 mM sodium chloride at TO) up to 58 nm (TO) or 79 nm after 24 hours at 25°C.

[0159] Antigen size and antigenicity are relatively stable over 24 hours at 25°C at pH 7.5 or 8.5, particularly in the absence of sodium chloride or at a sodium chloride concentration of 50 mM. Nevertheless, increasing the concentration of sodium chloride to 150 mM or greater results in a clear increase in antigen size and reduction in antigenicity.

Example 10: Prevention of "salting out" in compositions comprising M72, immunostimulants and using sorbitol as a tonicity agent

[0160] In order to compare the stability of immunogenic compositions containing 150 mM NaCl with compositions using sorbitol as a tonicity agent, a number of samples were monitored using SEC-HPLC and ELISA.

Method

[0161] Three different lyophilisation cakes were prepared, such that when combined with the appropriate adjuvant formulations from Examples 5 and 7 the desired pH would be obtained:

(a) M72 with two N-terminal His residues - target pH 8.5 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 50 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

(b) M72 with two N-terminal His residues - target pH 8.0 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 20 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

(c) M72 with two N-terminal His residues - target pH 7.5 in reconstituted vaccine

15.75% (w/v) sucrose solution (prepared in water for injection) was added to water for injection to reach a sucrose concentration of 6.3%. 3% (w/v) Tween80 solution (prepared in water for injection) was then added to reach a concentration of 0.025%. Tris-HCl buffer 1 M pH 8.8 was then added to reach a 12.5 mM Tris buffer concentration. The mixture was magnetically stirred for 5 minutes at room temperature. Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3, as prepared in Example 8) was then added to reach a protein concentration of 25 ug/ml. The mixture was magnetically stirred for 10 minutes at room temperature. The pH was checked and found to be 8.8.

0.5 ml of the mixture obtained was filled in 3 ml glass vials then freeze dried.

[0162] The lyophilisation cakes described above were reconstituted with 625 ul of the adjuvant solutions prepared in Examples 5 and 7. Upon reconstitution with adjuvant solution, the following immunogenic compositions were obtained:

(i) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.5

- 10 ug antigen (20 ug/ml)
- 5% w/v sucrose
- 40 mM Tris
- 0.02% w/v Tween80
- 500 ug DOPC
- 125 ug cholesterol
- 25 ug 3D-MPL
- 25 ug QS21
- 150 mM NaCl
- 10 mM phosphate
- pH 8.5

EP 2 651 437 B1

(ii) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.0

5 10 ug antigen (20 ug/ml)
5% w/v sucrose
16 mM Tris
0.02% w/v Tween80
500 ug DOPC
125 ug cholesterol
10 25 ug 3D-MPL
25 ug QS21
150 mM NaCl
10 mM phosphate
pH 8.0

15 (iii) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 7.5

10 ug antigen (20 ug/ml)
5% w/v sucrose
12.5 mM Tris
20 0.02% w/v Tween80
500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
25 150 mM NaCl
10 mM phosphate
pH 7.5

30 (iv) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.5

10 ug antigen (20 ug/ml)
5% w/v sucrose
40 mM Tris
0.02% w/v Tween80
35 500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
5 mM NaCl
40 4.7% w/v sorbitol
10 mM phosphate
pH 8.5

45 (v) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.0

10 ug antigen (20 ug/ml)
5% w/v sucrose
16 mM Tris
0.02% w/v Tween80
50 500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
5 mM NaCl
55 4.7% w/v sorbitol
10 mM phosphate
pH 8.0

EP 2 651 437 B1

(vi) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 7.5

10 ug antigen (20 ug/ml)
5% w/v sucrose
12.5 mM Tris
0.02% w/v Tween80
500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
5 mM NaCl
4.7% w/v sorbitol
10 mM phosphate
pH 7.5

Sample Analysis

[0163] The reconstituted immunogenic compositions described above were characterised after storage at 25°C or 30°C (T0, T6h and T24h).

[0164] SEC-HPLC analysis was performed by injection on a TOSOH TSK-Gel5000Pwxl (ID 7.8 mm x 30 cm) equilibrated in 20 mM Tris buffer pH 8.5, detection by UV at 210 nm and flow rate 0.5 ml/min.

[0165] Antigenicity was quantified by a sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0166] The results are shown in Figures 6a-6d and 7.

[0167] SEC-HPLC profiles are stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at each pH (i.e. pH 7.5, 8.0 and 8.5). This may be contrasted with the SEC-HPLC profiles for immunogenic compositions containing 150 mM NaCl, which show clear changes between the initial profile obtained and those following storage at 25°C or 30°C. This evolution becomes more intense when the pH of the 150 mM NaCl composition is lowered.

[0168] The same conclusions can be drawn in terms of antigenicity, with recoveries remaining largely stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at each pH (i.e. pH 7.5, 8.0 and 8.5) up to 24h at 30°C.

Example 11: Conductivity determination for immunogenic compositions of the invention.

[0169] The conductivity of a range of immunogenic compositions according to the present invention was measured and compared to the conductivity of control sodium chloride solutions and with an immunogenic composition containing a conventional quantity of sodium chloride.

Method

[0170] A range of standards having sodium chloride concentrations of 0, 75, 100, 150, 250 and 300 mM were prepared from a stock solution of 1500 mM sodium chloride by dilution in water for injection.

[0171] Immunogenic compositions were prepared using M72 with two N-terminal His residues according to the procedures provided in Example 8. To investigate the contribution from the antigen itself and any residual materials in the purified bulk, placebo lyophilisation cakes were also prepared by excluding the antigen component.

[0172] Using a Malvern Zetasizer Nano and 1.5 ml of each sample in folded capillary cells, a voltage of 30 to 150 V (determined automatically by the instrument) was applied and the conductivity determined.

Results

[0173]

EP 2 651 437 B1

Conductivity of sodium chloride standard solutions

Sodium chloride concentration mM	Conductivity mS/cm
0	0.0
75	8.2
100	10.7
150	15.6
250	23.9
300	30.0

[0174] A standard curve, based on this data, is provided in Figure 8.

Conductivity of test solutions

Description	Sodium chloride concentration mM	Conductivity mS/cm	Equivalent sodium chloride concentration mM
ASA(sorbitol - 2)	5	1.46	9
Placebo pH 8.0 / ASA(sorbitol - 2)	5	1.95	14
M72 pH 8.0 / ASA(sorbitol - 2)	5	1.96	14
Placebo pH 8.5 / ASA(sorbitol - 2)	5	2.36	18
M72 pH 8.5 / ASA(sorbitol - 2)	5	2.28	17
ASA(150 mM NaCl - 2)	150	16	159
Placebo pH 8.5 / ASA(150 mM NaCl - 2)	150	14.8	147
M72 pH 8.5 / ASA(150 mM NaCl - 2)	150	15.3	152

[0175] As can be seen from the data above, the conductivity of solutions which utilise 150 mM NaCl is significantly greater than that of solutions which make minimal use of NaCl.

[0176] The impact of the antigen and any components in the purified bulk is minimal, as placebo preparations have comparable conductivity to their M72 related antigen containing counterparts.

Example 12: Immunogenicity testing of immunogenic compositions of the invention.

[0177] The aim of the this Example was to determine whether or not formulation changes to reduce the quantity of salt in immunogenic compositions of the invention, with a view to improving protein stability, had an impact on *in vivo* immunogenicity.

Method

[0178] Four immunogenic compositions were evaluated:

1. M72 with two N-terminal His residues pH 8.5/ASA (150 mM NaCl - 2)
2. M72 with two N-terminal His residues pH 8.5/ASA (sorbitol - 2)
3. M72 with two N-terminal His residues pH 8/ASA (sorbitol - 2)

EP 2 651 437 B1

4. M72 with two N-terminal His residues pH 7.5/ASA (sorbitol - 2)

[0179] The immunogenicity of these antigen containing compositions was evaluated in C57BL/6 mice. For each of the four compositions, 30 C57BL/6 mice were injected 3 times intramuscularly, on days 0, 14 and 28 with 1 ug of antigen in 50 ul of adjuvant solution (prepared by the procedure provided in Example 10). The elicited M72 specific T cell responses (both CD4 & CD8) were measured 6 days post last immunisation (6dPIII).

[0180] For the determination of M72-specific cellular responses, peripheral blood lymphocytes from 30 mice/group were collected and pooled (six pools of five mice/group). A red blood cells lysis was performed before plating the cells *in vitro*. The cells were restimulated *in vitro* with a pool of overlapping peptides (15-mer peptides with an 11 amino acid overlap, at 1 ug/ml/peptide) covering the M72 sequence (without the N-terminal His residues). Cells remaining in medium (no peptide stimulation) were used to determine the background responses. Two hours after the co-culture with the peptide pool, brefeldin A was added to the wells (to inhibit cytokine excretion) and the cells were stored overnight at 4 °C. The cells were subsequently stained for the following markers: CD4, CD8, IL-2, IFN-gamma and TNF-alpha.

Results

[0181] Each datapoint in Figures 9 and 10 represents the background subtracted M72-specific CD4 or CD8 T cell response, respectively, of a pool of peripheral blood lymphocytes from five mice six days after the third immunisation. The response is expressed as the percentage of CD4 T cells producing IFN-gamma and/or IL-2 and/or TNF-alpha in response to simulation with the M72 peptide pool. The bar represents the median of the responses for each group.

[0182] The results in Figure 9 and 10 show that comparable CD4 and CD8 T cell responses are induced following three immunisations with each of the test formulations. Consequently, it may be concluded that a reduction in the quantity of salts present in the immunogenic compositions of the present invention does not lead to a compromise in the induced T cell responses.

Example 13: Investigation of "salting out" in compositions comprising M72 with CaCl₂ or MgSO₄ at pH 6.1 and 8.0.

[0183] To investigate impact of other salts on M72 antigen stability, solutions were prepared with a range of concentrations of CaCl₂ or MgSO₄ and at different pH levels. Visual inspection was used as a readout of stability.

Method

[0184] Purified bulk antigen (M72 with two N-terminal His residues, SEQ ID No: 3) was diluted to a concentration of 100 ug/ml in two different buffers (10mM succinate buffer at pH 6.1 and 10 mM Tris buffer at pH 8.0) containing specified quantities of salts (0 mM; 150 mM or 300 mM NaCl; 40 mM, 80 mM or 160 mM CaCl₂; 87.5 mM, 175 mM or 430 mM MgSO₄).

[0185] Samples were analysed directly after preparation.

[0186] Using a Mettler Toledo conductivity meter and 6 ml of each sample in an unsiliconised glass vial, the conductivity was determined.

Results

[0187]

Group	Salt	Buffer	pH (theoretical)	Conductivity (mS/cm) (measured)	pH (measured)	Visual Observation
A	0 mM	Succinate 10mM	6.1	1.1	6.3	Clear
B	NaCl 150 mM	Succinate 10mM	6.1	13.4	6.1	Clear
C	NaCl 300 mM	Succinate 10mM	6.1	20.0	6.1	Clear
D	CaCl ₂ 40 mM	Succinate 10mM	6.1	8.0	6.1	Opalescent
E	CaCl ₂ 80 mM	Succinate 10mM	6.1	11.2	5.8	Opalescent + large particles

EP 2 651 437 B1

(continued)

	<u>Group</u>	<u>Salt</u>	<u>Buffer</u>	<u>pH (theoretical)</u>	<u>Conductivity (mS/cm) (measured)</u>	<u>pH (measured)</u>	<u>Visual Observation</u>
5	F	CaCl ₂ 160 mM	Succinate 10mM	6.1	20.2	5.8	Opalescent + large particles
	G	MgSO ₄ 87.5 mM	Succinate 10mM	6.1	7.7	6.1	Opalescent
10	H	MgSO ₄ 175 mM	Succinate 10mM	6.1	12.4	5.9	Opalescent+ very large particles
15	I	MgSO ₄ 430 mM	Succinate 10mM	6.1	20.4	5.9	Opalescent+ very large particles
	J	0 mM	Tris 10mM	8.0	0.463	8.0	Clear
20	K	NaCl 150 mM	Tris 10mM	8.0	12.13	8.0	Clear
	L	NaCl 300 mM	Tris 10mM	8.0	21.1	8.0	Clear
25	M	CaCl ₂ 40 mM	Tris 10mM	8.0	6.7	8.1	Large particles
	N	CaCl ₂ 80 mM	Tris 10mM	8.0	10.8	8.0	Opalescent+ large particles
30	O	CaCl ₂ 160 mM	Tris 10mM	8.0	19.7	8.0	Opalescent+ large particles
	P	MgSO ₄ 87.5 mM	Tris 10mM	8.0	7.5	8.0	Large particles
35	Q	MgSO ₄ 175 mM	Tris 10mM	8.0	10.9	8.2	Opalescent+ very large particles
40	R	MgSO ₄ 430 mM	Tris 10mM	8.0	21.7	8.1	Opalescent+ very large particles

[0188] The results demonstrate that solutions containing an M72 related antigen can be sensitive to salts other than sodium chloride. The impact of CaCl₂ or MgSO₄ appears to be more pronounced than for sodium chloride at comparable concentrations or conductivity.

Example 14: Investigation of "salting out" in compositions comprising Mtb72f with different salt concentrations at pH 6.1, 7.5 and 8.5.

[0189] The impact of sodium chloride concentration and pH on Mtb72f antigen stability, as assessed by size, was investigated.

Method

[0190] Purified bulk antigen (Mtb72f with 6 his residues, SEQ ID No: 7) was diluted to a concentration of 100 ug/ml in three different buffers (10mM phosphate buffer at pH 6.1, 20 mM Tris buffer at pH 7.5 and 20 mM Tris buffer at pH 8.5) containing final sodium chloride concentrations of 0, 150 and 450 mM.

[0191] Samples were stored for 24 hours at 4°C or 25°C before analysis.

[0192] Nephelometry was performed using a Nepheloskan® Ascent, available from Thermo Fischer Scientific. Analysis was performed in UV transparent Costar® micro-plates available from Corning Inc (USA).

[0193] DLS was performed using a Dynapro Plate Reader from Wyatt Instruments. The instrument was operated using a laser wavelength of 830 nm and power of 50 mW. Scattered light was detected at 150° at a temperature of 22°C. The mean hydrodynamic diameter and polydispersity index (pl) are calculated by the instrument software.

Results

[0194] The findings of this experiment are presented in Figures 11 and 12.

[0195] Both DLS and nephelometry demonstrate a general trend that Mtb72f is sensitive to salt concentration and pH, in a similar manner to M72 as shown in previous examples. Consequently, the benefits of the present invention apply to M72 related antigens and not just to the M72 sequence itself.

[0196] In the case of a number of DLS samples, instrumentation was unable to determine a specific particle size (shown as NV in Figure 12).

Example 15: Prevention of "salting out" in compositions comprising M72, immunostimulants and using sorbitol as a tonicity agent

[0197] In order to compare the stability of immunogenic compositions containing 150 mM NaCl with compositions using sorbitol as a tonicity agent, samples were monitored using an alternative ELISA.

Method

[0198] Lyophilisation cake was prepared as described in Example 10 (specifically method (a)) such that when combined with the appropriate adjuvant formulations from Example 7 a pH of 8.5 would be obtained.

[0199] The lyophilisation cake describe above were reconstituted with 625 ul of the adjuvant solutions prepared in Example 7. Upon reconstitution with adjuvant solution, the following immunogenic compositions were obtained:

(i) M72 with two N-terminal His residues - ASA(150 mM NaCl - 2) pH 8.5

10 ug antigen (20 ug/ml)
5% w/v sucrose
40 mM Tris
0.02% w/v Tween80
500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
150 mM NaCl
10 mM phosphate
pH 8.5

(ii) M72 with two N-terminal His residues - ASA(sorbitol - 2) pH 8.5

10 ug antigen (20 ug/ml)
5% w/v sucrose
40 mM Tris
0.02% w/v Tween80
500 ug DOPC
125 ug cholesterol
25 ug 3D-MPL
25 ug QS21
5 mM NaCl
4.7% w/v sorbitol
10 mM phosphate
pH 8.5

[0200] The reconstituted immunogenic compositions described above were characterised after storage at 30°C (T24h)

and compared with an extemporaneously prepared sample (TO).

[0201] Antigenicity was quantified by an indirect sandwich ELISA in which the antigen is captured by a M72-specific rabbit polyclonal antibody and subsequently revealed by a M72(Mtb39)-specific mouse monoclonal antibody. Briefly, the plate is coated with anti-M72 rabbit polyclonal antibody at the dilution of 1/8000 in Dulbecco's Phosphate Buffered Saline overnight at 4°C and after four washes the plates were blocked for 1h at 37°C with saturation buffer (PBS, 0.1 % Tween 20, 1% BSA). After the washing step, protein standard (M72 purified bulk: 1950 ug/ml), internal control (M72: 1768 µg/ml) and samples are loaded in wells from the first column of the plate at a concentration of approximately 0.25µg/ml and then a 2-fold serial dilution is performed in the saturation buffer (PBS, 0.025% Tween 20) from well 1 to 12 and incubated 1h30 at 37°C. After the washing step, the immune complex is then incubated 1h at 37°C with anti-M72 mouse monoclonal antibody at a dilution of 1/1000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, a biotinylated rabbit anti-mouse polyclonal antibody was added at a dilution of 1/1000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, the signal was amplified by adding Streptavidin-Horseradish Peroxidase diluted 1/4000 in saturation buffer (PBS, 0.025% Tween 20). After four washes, the signal was revealed by ortho phenylene diamine dihydrochlorid (OPDA) for 15min at RT and the reaction is stopped by addition of HCl 1M. The coloration is proportional to the quantity of bound anti-M72 antibody, and is measured at 490 nm and 620 nm. All washing steps were performed using PBS, 0.025% Tween 20.

[0202] All measured values are presented relative to the expected antigenicity based on the purified bulk protein used to prepare the tested formulations.

Results

[0203] The results are shown in Figure 13. Diamonds indicating the specific measurements for each of the three test samples, with a line indicating the average value.

[0204] Antigen recovery is largely stable after reconstitution in low salt compositions using sorbitol as a tonicity agent at pH 8.5 up to 24h at 30°C. Recovery in ASA(sorbitol-2) was 83.5% after 24 hours (TO 87.1%, meaning 95.9% of the relative antigenicity was maintained), whereas recovery in ASA(NaCl-2) was 54.5% after 24 hours (TO 81.0%, meaning only 67.3% of the relative antigenicity was maintained after storage).

[0205] In summary, Examples 9, 10 and 15 demonstrate for the first time the detrimental impact resulting from pH and NaCl concentration on the stability of immunogenic compositions containing an M72 related antigen. Example 13 extends this work to show that other salts may also have a detrimental impact on the stability of immunogenic compositions containing an M72 related antigen, with Example 14 demonstrating that the effect is also applicable to M72 related sequences.

[0206] Reformulation of the immunogenic compositions with a non-ionic tonicity agent addresses the antigen stability problems. Additionally, Examples 3, 4 and 12 demonstrate the removal of substantially all NaCl from the immunogenic formulation and its replacement with sorbitol as a tonicity agent does not have a detrimental impact on the stimulation of T cell responses.

[0207] Stability of immunogenic compositions is key and may be particularly challenging when in isolated locations where refrigeration may not be readily accessible. By reducing the presence of salts in the immunogenic compositions, the present inventors have been able to reduce the extent of changes observed when the immunogenic compositions are stored.

SEQUENCE LISTING

[0208]

<110> GlaxoSmithKline Biologicals sa
Godart, Stephane
Laanan, Amina
Lemoine, Dominique

<120> Novel compositions

<130> VB64442

<150> US61/422723

<151> 2010-12-14

<160> 13

EP 2 651 437 B1

<170> PatentIn version 3.5

<210> 1

<211> 723

5 <212> PRT

<213> Artificial Sequence

<220>

10 <223> M72 fusion protein

<400> 1

15 Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
1 5 10 15

Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
20 20 25 30

Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
25 35 40 45

Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
50 55 60

Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
30 65 70 75 80

Val Ile Thr Ala Val Asp Gly Ala Pro Ile Asn Ser Ala Thr Ala Met
85 90 95

Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
35 100 105 110

Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
40 115 120 125

Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
45 130 135 140

50

55

EP 2 651 437 B1

	Glu	Ile	Asn	Ser	Ala	Arg	Met	Tyr	Ala	Gly	Pro	Gly	Ser	Ala	Ser	Leu
	145					150					155					160
5	Val	Ala	Ala	Ala	Gln	Met	Trp	Asp	Ser	Val	Ala	Ser	Asp	Leu	Phe	Ser
					165					170					175	
10	Ala	Ala	Ser	Ala	Phe	Gln	Ser	Val	Val	Trp	Gly	Leu	Thr	Val	Gly	Ser
				180					185					190		
15	Trp	Ile	Gly	Ser	Ser	Ala	Gly	Leu	Met	Val	Ala	Ala	Ala	Ser	Pro	Tyr
			195					200						205		
20	Val	Ala	Trp	Met	Ser	Val	Thr	Ala	Gly	Gln	Ala	Glu	Leu	Thr	Ala	Ala
		210					215					220				
25	Gln	Val	Arg	Val	Ala	Ala	Ala	Ala	Tyr	Glu	Thr	Ala	Tyr	Gly	Leu	Thr
	225				230						235					240
30	Val	Pro	Pro	Pro	Val	Ile	Ala	Glu	Asn	Arg	Ala	Glu	Leu	Met	Ile	Leu
				245						250					255	
35	Ile	Ala	Thr	Asn	Leu	Leu	Gly	Gln	Asn	Thr	Pro	Ala	Ile	Ala	Val	Asn
				260					265					270		
40	Glu	Ala	Glu	Tyr	Gly	Glu	Met	Trp	Ala	Gln	Asp	Ala	Ala	Ala	Met	Phe
			275					280					285			
45	Gly	Tyr	Ala	Ala	Ala	Thr	Ala	Thr	Ala	Thr	Ala	Thr	Leu	Leu	Pro	Phe
		290					295					300				
50	Glu	Glu	Ala	Pro	Glu	Met	Thr	Ser	Ala	Gly	Gly	Leu	Leu	Glu	Gln	Ala
	305					310					315					320
55	Ala	Ala	Val	Glu	Glu	Ala	Ser	Asp	Thr	Ala	Ala	Ala	Asn	Gln	Leu	Met
					325					330					335	
60	Asn	Asn	Val	Pro	Gln	Ala	Leu	Gln	Gln	Leu	Ala	Gln	Pro	Thr	Gln	Gly
				340					345						350	
65	Thr	Thr	Pro	Ser	Ser	Lys	Leu	Gly	Gly	Leu	Trp	Lys	Thr	Val	Ser	Pro
			355					360						365		
70	His	Arg	Ser	Pro	Ile	Ser	Asn	Met	Val	Ser	Met	Ala	Asn	Asn	His	Met
		370					375					380				
75	Ser	Met	Thr	Asn	Ser	Gly	Val	Ser	Met	Thr	Asn	Thr	Leu	Ser	Ser	Met
	385					390					395					400

EP 2 651 437 B1

Leu Lys Gly Phe Ala Pro Ala Ala Ala Gln Ala Val Gln Thr Ala
 405 410 415
 5 Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
 420 425 430
 10 Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala
 435 440 445
 Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala Asn Gln
 450 455 460
 15 Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu Thr Ser
 465 470 475 480
 20 Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro Val Gly
 485 490 495
 25 Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu Arg Val
 500 505 510
 Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly Asp Ile
 515 520 525
 30 Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro Ala Leu
 530 535 540
 35 Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln Val Val
 545 550 555 560
 40 Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala Gly Thr
 565 570 575
 Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn His Val
 580 585 590
 45 Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser Gly Gln
 595 600 605
 50 Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp Val Ala
 610 615 620
 Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala Ile Gly
 625 630 635 640
 55 Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn Ser Gly

EP 2 651 437 B1

645

650

655

5 Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val Ala Leu
660 665 670

10 Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu Glu Thr
675 680 685

15 Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ala
690 695 700

20 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

25 Ala Ala Ser

<210> 2

<211> 2172

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Coding sequence for M72 fusion

35 <400> 2

40

45

50

55

60

65

EP 2 651 437 B1

atgacggccg cgtccgataa cttccagctg tcccagggtg ggcagggatt cgccattccg 60
 atcgggcagg cgatggcgat cgcgggccag atccgatcgg gtgggggggtc acccaccggt 120
 5 catatcgggc ctaccgcctt cctcggcttg ggtgtgtcgc acaacaacgg caacggcgca 180
 cgagtccaac gcgtggtcgg gagcgcctcc gcggcaagtc tcggcatctc caccggcgac 240
 gtgatcaccg cggtcgacgg cgtccgatc aactcggcca ccgcgatggc ggacgcgctt 300
 10 aacgggcatc atcccgggta cgtcatctcg gtgacctggc aaaccaagtc gggcggcacg 360
 cgtacagggg acgtgacatt ggccgagggg cccccggccg aattcatggt ggatttcggg 420
 15 gcgttaccac cggagatcaa ctccgcgagg atgtacgccg gcccgggttc ggctcgcgtg 480
 gtggccgcgg ctcagatgtg ggacagcgtg gcgagtgacc tgttttcggc cgcgtcggcg 540
 tttcagtcgg tggctcgggg tctgacggtg ggtcgtgga taggttcgtc ggcgggtctg 600
 20 atggtggcgg cggcctcgcc gtatgtggcg tggatgagcg tcaccgcggg gcagggcgag 660
 ctgaccgccg cccaggtccg ggttgctgcg gcggcctacg agacggcgta tgggctgacg 720
 gtgccccgcg cggatgatcg cgagaaccgt gctgaactga tgattctgat agcgaccaac 780
 25 ctcttggggc aaaacacccc ggcgatcgcg gtcaacgagg ccgaatacgg cgagatgtgg 840
 gcccaagacg ccgccgcgat gtttgctac gccgcggcga cggcgacggc gacggcgacg 900
 30 ttgctgccgt tcgaggaggc gccggagatg accagcgcgg gtgggctcct cgagcaggcc 960

 35

 40

 45

 50

 55

EP 2 651 437 B1

gccgcggtcg aggaggcctc cgacaccgcc gcggcgaacc agttgatgaa caatgtgcc 1020
 caggcgctgc aacagctggc ccagcccacg cagggcacca cgccttcttc caagctgggt 1080
 5 ggcctgtgga agacggtctc gccgcatcgg tcgccgatca gcaacatggt gtogatggcc 1140
 aacaaccaca tgtcgatgac caactcgggt gtgtcgatga ccaacacctt gagctogatg 1200
 ttgaagggct ttgctccggc ggcgcccgcc caggccgtgc aaaccgcggc gcaaaacggg 1260
 10 gtccgggcga tgagctcgct gggcagctcg ctgggttctt cgggtctggg cgggtggggtg 1320
 gccgccaaact tgggtcgggc ggcctcggtc ggttcgttgt cggtgccgca ggcctgggccc 1380
 gcggccaacc aggcagtcac cccggcggcg cgggcgctgc cgctgaccag cctgaccagc 1440
 15 gccgcggaaa gagggcccgg gcagatgctg ggcgggctgc cgggtggggca gatgggccc 1500
 agggccgggtg gtgggctcag tgggtgtgctg cgtgttccgc cgcgacccta tgtgatgccg 1560
 20 cattctccgg cagccggcga tatcgcctcg ccggccttgt cgcaggaccg gttcgcgcgac 1620
 ttccccgcgc tgcccctcga cccgtccgcg atggtcggcc aagtggggcc acaggtggtc 1680
 aacatcaaca ccaaactggg ctacaacaac gccgtgggcg ccgggaccgg catcgtcatc 1740
 25 gatcccaacg gtgtcgtgct gaccaacaac cacgtgatcg cgggcgccac cgacatcaat 1800
 gcggtcagcg tcggctccgg ccaaactac ggcgtcgatg tggtcgggta tgaccgcacc 1860
 30 caggatgtcg cggtgctgca gctgcgcggc gccggtggcc tgccgtcggc ggcgatcggc 1920
 ggcggcgtcg cggttggtga gccgcctcgc gcgatgggca acagcgggtg gcagggcggga 1980
 acgccccgtg cggtgccctgg caggggtggc gcgctcggcc aaaccgtgca ggcgtcggat 2040
 35 tcgctgaccg gtgccgaaga gacattgaac gggttgatcc agttcgatgc cgcgatccag 2100
 cccggtgatg cgggcgggccc cgtcgtcaac ggcctaggac aggtggtcgg tatgaacacg 2160
 gccgcgtcct ag 2172
 40
 <210> 3
 <211> 725
 <212> PRT
 <213> Artificial Sequence
 45
 <220>
 <223> M72 fusion with 2 additional his residues
 50
 <400> 3
 55

EP 2 651 437 B1

Met His His Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly
1 5 10 15

5 Gln Gly Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln
 20 25 30

10 Ile Arg Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala

15

20

25

30

35

40

45

50

55

EP 2 651 437 B1

	35					40					45					
5	Phe	Leu	Gly	Leu	Gly	Val	Val	Asp	Asn	Asn	Gly	Asn	Gly	Ala	Arg	Val
	50						55					60				
10	Gln	Arg	Val	Val	Gly	Ser	Ala	Pro	Ala	Ala	Ser	Leu	Gly	Ile	Ser	Thr
	65					70					75					80
15	Gly	Asp	Val	Ile	Thr	Ala	Val	Asp	Gly	Ala	Pro	Ile	Asn	Ser	Ala	Thr
					85					90					95	
20	Ala	Met	Ala	Asp	Ala	Leu	Asn	Gly	His	His	Pro	Gly	Asp	Val	Ile	Ser
				100					105					110		
25	Val	Thr	Trp	Gln	Thr	Lys	Ser	Gly	Gly	Thr	Arg	Thr	Gly	Asn	Val	Thr
			115					120					125			
30	Leu	Ala	Glu	Gly	Pro	Pro	Ala	Glu	Phe	Met	Val	Asp	Phe	Gly	Ala	Leu
		130					135					140				
35	Pro	Pro	Glu	Ile	Asn	Ser	Ala	Arg	Met	Tyr	Ala	Gly	Pro	Gly	Ser	Ala
	145					150					155					160
40	Ser	Leu	Val	Ala	Ala	Ala	Gln	Met	Trp	Asp	Ser	Val	Ala	Ser	Asp	Leu
					165					170					175	
45	Phe	Ser	Ala	Ala	Ser	Ala	Phe	Gln	Ser	Val	Val	Trp	Gly	Leu	Thr	Val
				180					185					190		
50	Gly	Ser	Trp	Ile	Gly	Ser	Ser	Ala	Gly	Leu	Met	Val	Ala	Ala	Ala	Ser
			195					200					205			
55	Pro	Tyr	Val	Ala	Trp	Met	Ser	Val	Thr	Ala	Gly	Gln	Ala	Glu	Leu	Thr
		210					215					220				
60	Ala	Ala	Gln	Val	Arg	Val	Ala	Ala	Ala	Ala	Tyr	Glu	Thr	Ala	Tyr	Gly
	225					230					235					240
65	Leu	Thr	Val	Pro	Pro	Pro	Val	Ile	Ala	Glu	Asn	Arg	Ala	Glu	Leu	Met
					245					250					255	
70	Ile	Leu	Ile	Ala	Thr	Asn	Leu	Leu	Gly	Gln	Asn	Thr	Pro	Ala	Ile	Ala
				260					265					270		
75	Val	Asn	Glu	Ala	Glu	Tyr	Gly	Glu	Met	Trp	Ala	Gln	Asp	Ala	Ala	Ala
			275					280					285			

EP 2 651 437 B1

Met Phe Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu
 290 295 300

5 Pro Phe Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu
 305 310 315 320

10 Gln Ala Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln
 325 330 335

15 Leu Met Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr
 340 345 350

20 Gln Gly Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val
 355 360 365

25 Ser Pro His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn
 370 375 380

30 His Met Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser
 385 390 395 400

35 Ser Met Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala Gln Ala Val Gln
 405 410 415

40 Thr Ala Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser
 420 425 430 435

45 Leu Gly Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg
 435 440 445

50 Ala Ala Ser Val Gly Ser Leu Ser Val Pro Gln Ala Trp Ala Ala Ala
 450 455 460

55 Asn Gln Ala Val Thr Pro Ala Ala Arg Ala Leu Pro Leu Thr Ser Leu
 465 470 475 480

60 Thr Ser Ala Ala Glu Arg Gly Pro Gly Gln Met Leu Gly Gly Leu Pro
 485 490 495

65 Val Gly Gln Met Gly Ala Arg Ala Gly Gly Gly Leu Ser Gly Val Leu
 500 505 510

70 Arg Val Pro Pro Arg Pro Tyr Val Met Pro His Ser Pro Ala Ala Gly
 515 520 525

75 Asp Ile Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe Ala Asp Phe Pro
 530 535 540

EP 2 651 437 B1

Ala Leu Pro Leu Asp Pro Ser Ala Met Val Ala Gln Val Gly Pro Gln
 545 550 555 560

5 Val Val Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn Ala Val Gly Ala
 565 570 575

10 Gly Thr Gly Ile Val Ile Asp Pro Asn Gly Val Val Leu Thr Asn Asn
 580 585 590

15 His Val Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe Ser Val Gly Ser
 595 600 605

20 Gly Gln Thr Tyr Gly Val Asp Val Val Gly Tyr Asp Arg Thr Gln Asp
 610 615 620

25 Val Ala Val Leu Gln Leu Arg Gly Ala Gly Gly Leu Pro Ser Ala Ala
 625 630 635

Ile Gly Gly Gly Val Ala Val Gly Glu Pro Val Val Ala Met Gly Asn
 645 650 655

30 Ser Gly Gly Gln Gly Gly Thr Pro Arg Ala Val Pro Gly Arg Val Val
 660 665 670

Ala Leu Gly Gln Thr Val Gln Ala Ser Asp Ser Leu Thr Gly Ala Glu
 675 680 685

35 Glu Thr Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly
 690 695 700

40 Asp Ala Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met
 705 710 715 720

Asn Thr Ala Ala Ser
 725

45 <210> 4
 <211> 2178
 <212> DNA
 <213> Artificial Sequence

50 <220>
 <223> Coding sequence for M72 fusion with 2 additional his residues

<400> 4

55

EP 2 651 437 B1

atgcatcaca cggccgcgtc cgataacttc cagctgtccc agggtagggca gggattcgcc 60
attccgatcg ggcagggcat ggcgatcgcg ggccagatcc gatcgggtgg ggggtcaccc 120
5 accgttcata tcgggcctac cgccttcctc ggcttgggtg ttgtcgacaa caacggcaac 180

10

15

20

25

30

35

40

45

50

55

EP 2 651 437 B1

ggcgcacgag tccaacgcgt ggtcgggagc gctccggcgg caagtctcgg catctccacc 240
 ggcgacgtga tcaccgcggg cgacggcgct ccgatcaact cggccaccgc gatggcggac 300
 5 ggccttaacg ggcacatcc cgggtgacgtc atctcgggga cctggcaaac caagtccggc 360
 ggcacgcgta cagggaacgt gacattggcc gagggacccc cggccgaatt catggtggat 420
 ttcggggcgt taccaccgga gatcaactcc gcgaggatgt acgccggccc gggttcggcc 480
 10 tcgctggtgg ccgcggctca gatgtgggac agcgtggcga gtgacctggt ttcggccgcg 540
 tcggcgtttc agtcgggtgg ctggggctcg acggtggggg cgtggatagg ttcgctcggc 600
 ggtctgatgg tggcggcgcc ctccgctat gtggcgtgga tgagcgtcac cgcggggcag 660
 15 gccgagctga ccgccgcca ggtccgggtt gctgcggcgg cctacgagac ggcgatggg 720
 ctgacggtgc ccccgccggt gatcgcgag aaccgtgctg aactgatgat tctgatagcg 780
 20 accaacctct tggggcaaaa caccggcg atcgcggtca acgaggccga atacggcgag 840
 atgtgggccc aagacccgc cgcgatggtt ggctacgccg cggcgacggc gacggcgacg 900
 gcgacgttgc tgccgttcga ggaggcgccg gagatgacca gcgcgggtgg gctcctcag 960
 25 caggccgccc cggtcgagga ggcctccgac accgccgcgg cgaaccagtt gatgaacaat 1020
 gtgccccagg cgctgcaaca gctggcccag cccacgcagg gcaccacgcc ttcttccaag 1080
 ctgggtggcc tgtggaagac ggtctcgccg catcggtcgc cgatcagcaa catggtgtcg 1140
 30 atggccaaca accacatgct gatgaccaac tcgggtgtgt cgatgaccaa caccttgagc 1200
 tcgatgttga agggctttgc tccggcgccg gccgcccagg ccgtgcaaac cgcggcgcaa 1260
 aacggggtcc gggcgatgag ctccgctggc agctcgtggt gttcttcggg tctggcggt 1320
 35 ggggtggccg ccaacttggg tcggggcgcc tcggtcgggt cgttgtcggg gccgcaggcc 1380
 tgggccgcgg ccaaccaggc agtcaccccg gcggcgccgg cgtgcccgt gaccagcctg 1440
 40 accagcggcc cggaaagagg gccggggcag atgctggggc ggctgccggg ggggcagatg 1500
 ggcgccaggg ccggtggtgg gctcagtggg gtgctgcgtg ttccgccgcg accctatgtg 1560
 atgccgcatt ctccggcagc cggcgatata gcccccgg ccttgtcgca ggaccggttc 1620
 45 gccgacttcc ccgcgctgcc cctcgacccg tccgcgatgg tcgcccaggt ggggcccacag 1680
 gtggtcaaca tcaacaccaa actgggctac aacaacgccg tgggcgcccg gaccggcatc 1740
 gtcacgatc ccaacggtgt cgtgctgacc aacaaccacg tgatcgcggg gccaccgac 1800
 50 atcaatgcgt tcagcgtcgg ctccggccaa acctacggcg tcgatgtggt cgggtatgac 1860
 cgcaccagg atgtcgggt gctgcagctg cgcggtgccg gtggcctgcc gtcggcgccg 1920
 55 atcgggtggc gcgtcgggt tgggtagccc gtcgtcgcga tgggcaacag cgggtggcag 1980
 ggcggaacgc ccgctcgggt gcctggcagg gtggtcgcgc tcggccaaac cgtgcaggcg 2040

EP 2 651 437 B1

	tcggattcgc tgaccggtgc cgaagagaca ttgaacgggt tgatccagtt cgatgccgcg	2100
	atccagcccg gtgatgcggg cgggcccgtc gtcaacggcc taggacaggt ggtcggtatg	2160
5	aacacggccg cgtcctag	2178

<210> 5

<211> 723

<212> PRT

10 <213> Artificial Sequence

<220>

<223> Mtb72f fusion

15 <400> 5

20

25

30

35

40

45

50

55

EP 2 651 437 B1

1 Met Thr Ala Ala Ser Asp Asn Phe Gln Leu Ser Gln Gly Gly Gln Gly
 5 Phe Ala Ile Pro Ile Gly Gln Ala Met Ala Ile Ala Gly Gln Ile Arg
 10 Ser Gly Gly Gly Ser Pro Thr Val His Ile Gly Pro Thr Ala Phe Leu
 15 Gly Leu Gly Val Val Asp Asn Asn Gly Asn Gly Ala Arg Val Gln Arg
 20 Val Val Gly Ser Ala Pro Ala Ala Ser Leu Gly Ile Ser Thr Gly Asp
 25 Ala Asp Ala Leu Asn Gly His His Pro Gly Asp Val Ile Ser Val Thr
 30 Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr Gly Asn Val Thr Leu Ala
 35 Glu Gly Pro Pro Ala Glu Phe Met Val Asp Phe Gly Ala Leu Pro Pro
 40 Val Ala Ala Ala Gln Met Trp Asp Ser Val Ala Ser Asp Leu Phe Ser
 45 Ala Ala Ser Ala Phe Gln Ser Val Val Trp Gly Leu Thr Val Gly Ser
 50
 55

EP 2 651 437 B1

Trp Ile Gly Ser Ser Ala Gly Leu Met Val Ala Ala Ala Ser Pro Tyr
195 200 205

5 Val Ala Trp Met Ser Val Thr Ala Gly Gln Ala Glu Leu Thr Ala Ala
210 215 220

10 Gln Val Arg Val Ala Ala Ala Tyr Glu Thr Ala Tyr Gly Leu Thr
225 230 235 240

15 Val Pro Pro Pro Val Ile Ala Glu Asn Arg Ala Glu Leu Met Ile Leu
245 250 255

Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr Pro Ala Ile Ala Val Asn
260 265 270

20 Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln Asp Ala Ala Ala Met Phe
275 280 285

25 Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr Ala Thr Leu Leu Pro Phe
290 295 300

30 Glu Glu Ala Pro Glu Met Thr Ser Ala Gly Gly Leu Leu Glu Gln Ala
305 310 315 320

Ala Ala Val Glu Glu Ala Ser Asp Thr Ala Ala Ala Asn Gln Leu Met
325 330 335

35 Asn Asn Val Pro Gln Ala Leu Gln Gln Leu Ala Gln Pro Thr Gln Gly
340 345 350

40 Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu Trp Lys Thr Val Ser Pro
355 360 365

45 His Arg Ser Pro Ile Ser Asn Met Val Ser Met Ala Asn Asn His Met
370 375 380

Ser Met Thr Asn Ser Gly Val Ser Met Thr Asn Thr Leu Ser Ser Met
385 390 395 400

50 Leu Lys Gly Phe Ala Pro Ala Ala Ala Gln Ala Val Gln Thr Ala
405 410 415

Ala Gln Asn Gly Val Arg Ala Met Ser Ser Leu Gly Ser Ser Leu Gly
420 425 430

55 Ser Ser Gly Leu Gly Gly Gly Val Ala Ala Asn Leu Gly Arg Ala Ala

EP 2 651 437 B1

		435				440						445				
5	Ser	Val	Gly	Ser	Leu	Ser	Val	Pro	Gln	Ala	Trp	Ala	Ala	Ala	Asn	Gln
		450					455					460				
10	Ala	Val	Thr	Pro	Ala	Ala	Arg	Ala	Leu	Pro	Leu	Thr	Ser	Leu	Thr	Ser
		465					470				475					480
15	Ala	Ala	Glu	Arg	Gly	Pro	Gly	Gln	Met	Leu	Gly	Gly	Leu	Pro	Val	Gly
					485					490						495
20	Gln	Met	Gly	Ala	Arg	Ala	Gly	Gly	Gly	Leu	Ser	Gly	Val	Leu	Arg	Val
				500					505					510		
25	Pro	Pro	Arg	Pro	Tyr	Val	Met	Pro	His	Ser	Pro	Ala	Ala	Gly	Asp	Ile
			515					520						525		
30	Ala	Pro	Pro	Ala	Leu	Ser	Gln	Asp	Arg	Phe	Ala	Asp	Phe	Pro	Ala	Leu
		530					535					540				
35	Pro	Leu	Asp	Pro	Ser	Ala	Met	Val	Ala	Gln	Val	Gly	Pro	Gln	Val	Val
		545				550					555					560
40	Asn	Ile	Asn	Thr	Lys	Leu	Gly	Tyr	Asn	Asn	Ala	Val	Gly	Ala	Gly	Thr
					565					570					575	
45	Gly	Ile	Val	Ile	Asp	Pro	Asn	Gly	Val	Val	Leu	Thr	Asn	Asn	His	Val
				580				585						590		
50	Ile	Ala	Gly	Ala	Thr	Asp	Ile	Asn	Ala	Phe	Ser	Val	Gly	Ser	Gly	Gln
			595					600					605			
55	Thr	Tyr	Gly	Val	Asp	Val	Val	Gly	Tyr	Asp	Arg	Thr	Gln	Asp	Val	Ala
		610					615					620				
60	Val	Leu	Gln	Leu	Arg	Gly	Ala	Gly	Gly	Leu	Pro	Ser	Ala	Ala	Ile	Gly
		625				630					635					640
65	Gly	Gly	Val	Ala	Val	Gly	Glu	Pro	Val	Val	Ala	Met	Gly	Asn	Ser	Gly
					645					650					655	
70	Gly	Gln	Gly	Gly	Thr	Pro	Arg	Ala	Val	Pro	Gly	Arg	Val	Val	Ala	Leu
				660					665					670		
75	Gly	Gln	Thr	Val	Gln	Ala	Ser	Asp	Ser	Leu	Thr	Gly	Ala	Glu	Glu	Thr
			675					680					685			

EP 2 651 437 B1

Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala Ile Gln Pro Gly Asp Ser
690 695 700

5 Gly Gly Pro Val Val Asn Gly Leu Gly Gln Val Val Gly Met Asn Thr
705 710 715 720

Ala Ala Ser

10

<210> 6
<211> 2172
<212> DNA
<213> Artificial Sequence

15

<220>
<223> Coding sequence for Mtb72f fusion

20

<400> 6

25

30

35

40

45

50

55

EP 2 651 437 B1

atgacggccg cgtccgataa cttccagctg tcccaggggtg ggcagggatt cgccattccg 60
 atcggggcagg cgatggcgat cgcgggccag atccgatcgg gtgggggggtc acccaccgtt 120
 5 catatcgggc ctaccgcctt cctcggcttg ggtgttgtcg acaacaacgg caacggcgca 180
 cgagtccaac gcgtggtcgg gagcgtccg gcggcaagtc tcggcatctc caccggcgac 240
 gtgatcaccg cggtcgacgg cgctccgatc aactcggcca ccgcgatggc ggacgcgctt 300
 10 aacgggcatac atcccggatga cgtcatctcg gtgacctggc aaaccaagtc gggcggcacg 360
 cgtacagggga acgtgacatt ggccgagggga cccccggccg aattcatggg ggatttcggg 420
 15 gcgttaccac cggagatcaa ctccgcgagg atgtacgccg gcccgggttc ggctctgctg 480
 gtggcccgcg ctcagatgtg ggacagcgtg gcgagtgacc tgttttcggc cgcgtcggcg 540
 tttcagtcgg tggctctggg tctgacggtg gggctgtgga taggttcgctc ggcgggtctg 600
 20 atggtggcgg cggcctcgcc gtatgtggcg tggatgagcg tcaccgcggg gcaggccgag 660
 ctgaccgccg cccaggtccg ggttgctgcg gcggcctacg agacggcgta tgggctgacg 720
 gtgccccgcg cggatgatcg cgagaaccgt gctgaactga tgattctgat agcgaccaac 780
 25 ctcttggggc aaaacacccc ggcgatcgcg gtcaacgagg ccgaatacgg cgagatgtgg 840
 gcccaagacg ccgcccgcat gtttggtctac gccgcggcga cggcgacggc gacggcgacg 900
 30 ttgctgccgt tcgaggaggc gccggagatg accagcgcgg gtgggctcct cgagcaggcc 960
 gccgcggctg aggaggcctc cgacaccgcc gcggcgaacc agttgatgaa caatgtgcc 1020
 caggcgctgc aacagctggc ccagcccacg cagggcacca cgccttcttc caagctgggt 1080
 35 ggcctgtgga agacggctct gccgcacggt tcgccgatca gcaacatggg gtcgatggcc 1140
 aacaaccaca tgtcgatgac caactcgggt gtgtcgatga ccaacacctt gagctcgatg 1200
 40 ttgaagggct ttgctccggc ggcggccgcc caggccgtgc aaaccgcggc gcaaaacggg 1260

40

45

50

55

EP 2 651 437 B1

gtccgggcca tgagctcgct gggcagctcg ctgggttctt cgggtctggg cgggtggggtg 1320
 gccgccaact tgggtcgggc ggcctcggtc ggttcgttgt cggtgccgca ggcctgggccc 1380
 5 gccggccaacc aggcagtcac cccggcggcg cgggcgctgc cgctgaccag cctgaccagc 1440
 gccgcggaaa gagggcccgg gcagatgctg ggcgggctgc cgggtggggca gatgggcgcc 1500
 agggccgggtg gtgggctcag tgggtgtgctg cgtgttccgc cgcgacccta tgtgatgccg 1560
 10 cattctccgg cagccggcga tatcgccccg ccggccttgt cgcaggaccg gttcgccgac 1620
 ttccccgcgc tgcccctcga cccgtccgcg atggtcggcc aagtggggcc acaggtggtc 1680
 aacatcaaca ccaaactggg ctacaacaac gccgtgggcg ccgggaccgg catcgtcatc 1740
 gatcccaacg gtgtcgtgct gaccaacaac cacgtgatcg cgggcgccac cgacatcaat 1800
 gcgttcagcg tcggctccgg ccaaactac ggcgtcgtatg tggtcgggta tgaccgcacc 1860
 20 caggatgtcg cggtgctgca gctgcgcggt gccggtggcc tgccgctcggc ggcgatcggg 1920
 ggcggcgtcg cggttggtga gcccgtcgtc gcgatgggca acagcgggtg gcagggcgga 1980
 acgccccgtg cgggtgcctgg cagggtggtc gcgctcggcc aaaccgtgca ggcgtcggat 2040
 25 tcgctgaccg gtgccgaaga gacattgaac gggttgatcc agttcgtatc cgcgatccag 2100
 cccggtgatt cgggcggggc cgctcgtcaac ggcctaggac aggtggtcgg tatgaacacg 2160
 30 gccgcgtcct ag 2172

<210> 7

<211> 729

<212> PRT

35 <213> Artificial Sequence

<220>

<223> Mtb72f fusion with 6 additional his residues

40 <400> 7

Met His His His His His His Thr Ala Ala Ser Asp Asn Phe Gln Leu
 1 5 10 15

45 Ser Gln Gly Gly Gln Gly Phe Ala Ile Pro Ile Gly Gln Ala Met Ala
 20 25 30

50 Ile Ala Gly Gln Ile Arg Ser Gly Gly Gly Ser Pro Thr Val His Ile
 35 40 45

55 Gly Pro Thr Ala Phe Leu Gly Leu Gly Val Val Asp Asn Asn Gly Asn
 50 55 60

Gly Ala Arg Val Gln Arg Val Val Gly Ser Ala Pro Ala Ala Ser Leu
 65 70 75 80

EP 2 651 437 B1

Gly Ile Ser Thr Gly Asp Val Ile Thr Ala Val Asp Gly Ala Pro Ile
 85 90 95
 5 Asn Ser Ala Thr Ala Met Ala Asp Ala Leu Asn Gly His His Pro Gly
 100 105 110
 10 Asp Val Ile Ser Val Thr Trp Gln Thr Lys Ser Gly Gly Thr Arg Thr
 115 120 125
 15 Gly Asn Val Thr Leu Ala Glu Gly Pro Pro Ala Glu Phe Met Val Asp
 130 135 140
 20 Phe Gly Ala Leu Pro Pro Glu Ile Asn Ser Ala Arg Met Tyr Ala Gly
 145 150 155 160
 25 Pro Gly Ser Ala Ser Leu Val Ala Ala Ala Gln Met Trp Asp Ser Val
 165 170 175
 30 Ala Ser Asp Leu Phe Ser Ala Ala Ser Ala Phe Gln Ser Val Val Trp
 180 185 190
 35 Gly Leu Thr Val Gly Ser Trp Ile Gly Ser Ser Ala Gly Leu Met Val
 195 200 205
 40 Ala Ala Ala Ser Pro Tyr Val Ala Trp Met Ser Val Thr Ala Gly Gln
 210 215 220
 45 Ala Glu Leu Thr Ala Ala Gln Val Arg Val Ala Ala Ala Tyr Glu
 225 230 235 240
 50 Thr Ala Tyr Gly Leu Thr Val Pro Pro Pro Val Ile Ala Glu Asn Arg
 245 250 255
 55 Ala Glu Leu Met Ile Leu Ile Ala Thr Asn Leu Leu Gly Gln Asn Thr
 260 265 270
 60 Pro Ala Ile Ala Val Asn Glu Ala Glu Tyr Gly Glu Met Trp Ala Gln
 275 280 285
 65 Asp Ala Ala Ala Met Phe Gly Tyr Ala Ala Ala Thr Ala Thr Ala Thr
 290 295 300
 70 Ala Thr Leu Leu Pro Phe Glu Glu Ala Pro Glu Met Thr Ser Ala Gly
 305 310 315 320
 75 Gly Leu Leu Glu Gln Ala Ala Ala Val Glu Glu Ala Ser Asp Thr Ala
 325 330 335

EP 2 651 437 B1

Ala Ala Asn Gln Leu Met Asn Asn Val Pro Gln Ala Leu Gln Gln Leu
340 345 350

5 Ala Gln Pro Thr Gln Gly Thr Thr Pro Ser Ser Lys Leu Gly Gly Leu
355 360 365

10 Trp Lys Thr Val Ser Pro His Arg Ser Pro Ile Ser Asn Met Val Ser
370 375 380

15 Met Ala Asn Asn His Met Ser Met Thr Asn Ser Gly Val Ser Met Thr
385 390 395 400

20 Asn Thr Leu Ser Ser Met Leu Lys Gly Phe Ala Pro Ala Ala Ala Ala
405 410 415

25 Gln Ala Val Gln Thr Ala Ala Gln Asn Gly Val Arg Ala Met Ser Ser
420 425 430 435

30 Leu Gly Ser Ser Leu Gly Ser Ser Gly Leu Gly Gly Gly Val Ala Ala
435 440 445

35 Asn Leu Gly Arg Ala Ala Ser Val Gly Ser Leu Ser Val Pro Gln Ala
450 455 460

40 Trp Ala Ala Ala Asn Gln Ala Val Thr Pro Ala Ala Arg Ala Leu Pro
465 470 475 480

45 Leu Thr Ser Leu Thr Ser Ala Ala Glu Arg Gly Pro Gly Gln Met Leu
485 490 495

50 Gly Gly Leu Pro Val Gly Gln Met Gly Ala Arg Ala Gly Gly Gly Leu
500 505 510 515

55 Ser Gly Val Leu Arg Val Pro Pro Arg Pro Tyr Val Met Pro His Ser
515 520 525

60 Pro Ala Ala Gly Asp Ile Ala Pro Pro Ala Leu Ser Gln Asp Arg Phe
530 535 540

65 Ala Asp Phe Pro Ala Leu Pro Leu Asp Pro Ser Ala Met Val Ala Gln
545 550 555 560

70 Val Gly Pro Gln Val Val Asn Ile Asn Thr Lys Leu Gly Tyr Asn Asn
565 570 575

75 Ala Val Gly Ala Gly Thr Gly Ile Val Ile Asp Pro Asn Gly Val Val
580 585 590

EP 2 651 437 B1

Leu Thr Asn Asn His Val Ile Ala Gly Ala Thr Asp Ile Asn Ala Phe
 595 600 605

5 Ser Val Gly Ser Gly Gln Thr Tyr Gly Val Asp Val Val Gly Tyr Asp
 610 615 620

10 Arg Thr Gln Asp Val Ala Val Leu Gln Leu Arg Gly Ala Gly Gly Leu
 625 630 635 640

15 Pro Ser Ala Ala Ile Gly Gly Gly Val Ala Val Gly Glu Pro Val Val
 645 650 655

Ala Met Gly Asn Ser Gly Gly Gln Gly Gly Thr Pro Arg Ala Val Pro
 660 665 670

20 Gly Arg Val Val Ala Leu Gly Gln Thr Val Gln Ala Ser Asp Ser Leu
 675 680 685

25 Thr Gly Ala Glu Glu Thr Leu Asn Gly Leu Ile Gln Phe Asp Ala Ala
 690 695 700

Ile Gln Pro Gly Asp Ser Gly Gly Pro Val Val Asn Gly Leu Gly Gln
 705 710 715 720

30 Val Val Gly Met Asn Thr Ala Ala Ser
 725

<210> 8

35 <211> 2190

<212> DNA

<213> Artificial Sequence

<220>

40 <223> Coding sequence for M72 fusion with 6 additional his residues

<400> 8

45 atgcatcacc atcaccatca cacggccgcg tccgataact tccagctgtc ccaggggtggg 60
 cagggattcg ccattccgat cgggcaggcg atggcgatcg cgggccagat ccgatcgggt 120
 ggggggtcac ccaccgttca tatcgggcct accgccttcc tcggcttggg tgttgtcgac 180
 50 aacaacggca acggcgcacg agtccaacgc gtggtcggga gcgctccggc ggcaagtctc 240
 ggcattctca ccggcgacgt gatcaccgcg gtcgacggcg ctccgatcaa ctcgccacc 300
 gcgatggcgg acgcgcttaa cgggcatcat cccggtgacg tcatctcggg gacctggcaa 360
 55 accaagtctg gcggcacgcg tacaggggaac gtgacattgg ccgagggacc cccggccgaa 420
 ttcattggtg atttcggggc gttaccaccg gagatcaact ccgagggat gtacgccggc 480

EP 2 651 437 B1

ccgggttcgg cctcgcctggt ggccgcggct cagatgtggg acagcgtggc gagtgacctg 540
 ttttcggccg cgtcggcggtt tcagtcggtg gtctggggtc tgacgggtggg gtcgtggata 600
 5 gggttcgtcgg cgggtctgat ggtggcggcg gcctcgccgt atgtggcgtg gatgagcgtc 660
 accgcggggc aggccgagct gaccgccgcc caggtccggg ttgctgcggc ggccctacgag 720
 acggcgtatg ggctgacggt gccccgcggc gtgatcgccg agaaccgtgc tgaactgatg 780
 10 attctgatag cgaccaacct cttggggcaa aacaccccg cgatcgcggt caacgaggcc 840
 gaatacggcg agatgtgggc ccaagacgcc gccgcgatgt ttggctacgc cgcggcgacg 900
 15 gcgacggcga cggcgacggt gctgccgttc gaggaggcgc cggagatgac cagcgcgggt 960
 gggctcctcg agcaggccgc cgcggtcgag gaggcctccg acaccgccgc ggcgaaccag 1020
 ttgatgaaca atgtgcccc a ggcgctgcaa cagctggccc agcccacgca gggcaccacg 1080
 20 ccttcttcca agctgggtgg cctgtggaag acggtctcgc cgcacggtc gccgatcagc 1140
 aacatggtgt cgatggccaa caaccacatg tcgatgacca actcgggtgt gtcgatgacc 1200
 aacaccttga gctcgatggt gaagggcttt gctccggcgc cggccgccc a ggcctgcaa 1260
 25 accgcggcgc aaaacgggggt ccgggcgatg agctcgtcgg gcagctcgtt gggttcttcg 1320
 ggtctgggcg gtgggggtggc cgccaacttg ggtcgggcgg cctcggtcgg ttcggtgtcg 1380
 gtgccgcagg cctgggcccgc ggccaaccag gcagtcaccc cggcggcgcg ggcgctgccg 1440
 30 ctgaccagcc tgaccagcgc cgcggaaaga gggcccgggc agatgctggg cgggctgccg 1500
 gtggggcaga tgggcgccag ggccggtggt gggctcagtg gtgtgctgcg tgttccgccg 1560
 35 cgaccctatg tgatgccgca ttctccggca gccggcgata tcgcccgcgc ggccctgtcg 1620
 caggaccggt tcgccgactt ccccgcgctg ccctcgcacc cgtccgcgat ggtcgcccaa 1680
 gtggggccac aggtggtcaa catcaacacc aaactgggct acaacaacgc cgtgggcgcc 1740
 40 gggaccggca tcgtcatcga tcccacgggt gtcgtgctga ccaacaacca cgtgatcgcg 1800
 ggcgccaccg acatcaatgc gttcagcgtc ggctccggcc aaacctacgg cgtcogatgtg 1860
 45 gtcgggtatg accgcaccca ggatgtcgcg gtgctgcagc tgccgcggtg cgggtggcctg 1920
 ccgtcggcgg cgatcgggtg cggcgtcgcg gttggtgagc ccgtcgtcgc gatgggcaac 1980
 agcgggtggc agggcggaac gccccgtcgc gtgcctggca ggggtggtcgc gctcggccaa 2040
 50 accgtgcagg cgtcggattc gctgaccggt gccgaagaga cattgaacgg gttgatccag 2100
 ttcgatgccg cgatccagcc cgggtattcg ggcgggcccg tcgtcaacgg cctaggacag 2160
 55 gtggtcggta tgaacacggc cgcgtcctag 2190
 <210> 9
 <211> 20
 <212> DNA

EP 2 651 437 B1

<213> Artificial Sequence

<220>

<223> Cpg Oligo 1 - CpG 1826

5

<400> 9

tccatgacgt tcctgacgtt 20

<210> 10

10

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

15

<223> CpG Oligo 2 - CpG 1758

<400> 10

tctcccagcg tgcgccat 18

20

<210> 11

<211> 30

<212> DNA

<213> Artificial Sequence

25

<220>

<223> CpG Oligo 3

<400> 11

accgatgacg tcgccggtga cggcaccacg 30

30

<210> 12

<211> 24

<212> DNA

<213> Artificial Sequence

35

<220>

<223> CpG Oligo 4 - CpG 2006

<400> 12

40

tcgctgtttt gtcgtttgt cggt 24

<210> 13

<211> 20

<212> DNA

45

<213> Artificial Sequence

<220>

<223> CpG Oligo 5 - CpG 1686

50

<400> 13

tccatgacgt tcctgatgct 20

Claims

55

1. An immunogenic composition comprising an M72 related antigen, wherein the M72 related antigen comprises a sequence having at least 90% identity to SEQ ID No: 1, the pH of said composition is in the range 7.0 to 9.0, and the conductivity of the composition is 5 mS/cm or lower.

EP 2 651 437 B1

2. The immunogenic composition according to claim 1, wherein the conductivity of the composition is 4 mS/cm or lower.
3. The immunogenic composition according to claim 2, wherein the conductivity of the composition is 3 mS/cm or lower.
- 5 4. The immunogenic composition according to any one of claims 1 to 3, wherein the concentration of salts in said composition is 40 mM or lower.
5. The immunogenic composition according to any one of claims 1 to 4, further comprising a non-ionic tonicity agent.
- 10 6. The immunogenic composition according to claim 5, wherein the non-ionic tonicity agent is a polyol.
7. The immunogenic composition according to claim 6, wherein the polyol is sorbitol and wherein the concentration of sorbitol is between about 4 and about 6% (w/v).
- 15 8. The immunogenic composition according to any one of claims 1 to 7, wherein the concentration of sucrose is between about 4 and about 6% (w/v).
9. The immunogenic composition according to any one of claims 1 to 8, further comprising one or more immunostimulants.
- 20 10. The immunogenic composition according to claim 9, wherein the immunogenic composition comprises QS21.
11. The immunogenic composition according to either claim 9 or 10, wherein the immunogenic composition comprises 3-de-O-acylated monophosphoryl lipid A.
- 25 12. The immunogenic composition according to any one of claims 1 to 11, wherein the osmolality is 250 to 750 mOsm/kg.
13. The immunogenic composition according to any one of claims 1 to 12, wherein the composition is provided as a unit dose of between 50 μ l and 1 ml.
- 30 14. The immunogenic composition according to claim 13, wherein the unit dose contains 5 to 50 μ g of M72 related protein.
15. The immunogenic composition according to claim 13 or 14, wherein the unit dose is a human dose and contains between 1 and 100 μ g of 3D-MPL.
- 35 16. The immunogenic composition according to any one of claims 13 to 15, wherein the unit dose is a human dose and contains between 1 and 100 μ g of QS21.
17. The immunogenic composition according to any one of claims 1 to 16, wherein the pH of said composition is in the range 7.5 to 8.5.
- 40 18. The immunogenic composition according to any one of claims 1 to 17, wherein the M72 related antigen comprises the amino acid sequence of SEQ ID No: 3.
- 45 19. The immunogenic composition according to any one of claims 1 to 17, wherein the M72 related antigen consists of a sequence having at least 90% identity to SEQ ID No: 1.
20. The immunogenic composition according to claim 19, wherein the M72 related antigen consists of the amino acid sequence of SEQ ID No: 3.
- 50 21. The immunogenic composition according to any one of claims 1 to 20, for use in medicine.
22. The immunogenic composition according to any one of claims 1 to 21, for use in a method for the prophylaxis, treatment or amelioration of infection by mycobacteria.
- 55 23. The immunogenic composition for use according to either claim 21 or 22, for administration to a human.

Patentansprüche

- 5 1. Immunogene Zusammensetzung, umfassen ein M72-verwandtes Antigen, wobei das M72-verwandte Antigen eine Sequenz mit wenigstens 90 % Identität zu SEQ ID NO: 1 umfasst, wobei der pH dieser Zusammensetzung im Bereich von 7,0 bis 9,0 liegt und die Leitfähigkeit der Zusammensetzung 5 mS/cm oder geringer ist.
2. Immunogene Zusammensetzung gemäß Anspruch 1, wobei die Leitfähigkeit der Zusammensetzung 4 mS/cm oder geringer ist.
- 10 3. Immunogene Zusammensetzung gemäß Anspruch 2, wobei die Leitfähigkeit der Zusammensetzung 3 mS/cm oder geringer ist.
4. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 3, wobei die Konzentration der Salze in dieser Zusammensetzung 40 mM oder geringer ist.
- 15 5. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 4, ferner umfassend ein nicht-ionisches Tonizitätsmittel.
6. Immunogene Zusammensetzung gemäß Anspruch 5, wobei das nicht-ionische Tonizitätsmittel ein Polyol ist.
- 20 7. Immunogene Zusammensetzung gemäß Anspruch 6, wobei das Polyol Sorbitol ist und wobei die Sorbitol-Konzentration zwischen etwa 4 und etwa 6% (G/V) liegt.
8. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 7, wobei die Sukrose-Konzentration zwischen etwa 4 und etwa 6 % (G/V) liegt.
- 25 9. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 8, ferner umfassend ein oder mehrere Immunstimulanzien.
- 30 10. Immunogene Zusammensetzung gemäß Anspruch 9, wobei die immunogene Zusammensetzung QS21 umfasst.
11. Immunogene Zusammensetzung gemäß entweder Anspruch 9 oder 10, wobei die immunogene Zusammensetzung 3-De-O-acyliertes Monophosphoryllipid A umfasst.
- 35 12. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 11, wobei die Osmolalität 250 bis 750 mOsm/kg ist.
13. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 12, wobei die Zusammensetzung als eine Einheitsdosis zwischen 50 µl und 1 ml bereitgestellt wird.
- 40 14. Immunogene Zusammensetzung gemäß Anspruch 13, wobei die Einheitsdosis 5 bis 50 µg M72-verwandtes Protein enthält.
15. Immunogene Zusammensetzung gemäß Anspruch 13 oder 14, wobei die Einheitsdosis eine humane Dosis ist und zwischen 1 und 100 µg 3D-MPL enthält.
- 45 16. Immunogene Zusammensetzung gemäß einem der Ansprüche 13 bis 15, wobei die Einheitsdosis eine humane Dosis ist und zwischen 1 und 100 µg QS21 enthält.
- 50 17. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 16, wobei der pH dieser Zusammensetzung im Bereich von 7,5 bis 8,5 ist.
18. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 17, wobei das M72-verwandte Antigen die Aminosäuresequenz SEQ ID NO: 3 umfasst.
- 55 19. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 17, wobei das M72-verwandte Antigen aus einer Sequenz mit wenigstens 90 % Identität zu SEQ ID NO: 1 besteht.

EP 2 651 437 B1

20. Immunogene Zusammensetzung gemäß Anspruch 19, wobei das M72-verwandte Antigen aus der Aminosäuresequenz SEQ ID NO: 3 besteht.
21. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 20 zur medizinischen Anwendung.
22. Immunogene Zusammensetzung gemäß einem der Ansprüche 1 bis 21 zur Anwendung bei einem Verfahren zur Prophylaxe, Behandlung oder Verbesserung einer Infektion durch Mykobakterien.
23. Immunogene Zusammensetzung zur Anwendung gemäß entweder Anspruch 21 oder 22 zur Verabreichung an einen Menschen.

Revendications

1. Composition immunogène comprenant un antigène apparenté à M72, où l'antigène apparenté à M72 comprend une séquence présentant au moins 90 % d'identité avec SEQ ID NO : 1, le pH de ladite composition se situe dans la plage de 7,0 à 9,0, et la conductivité de la composition est de 5 mS/cm ou inférieure.
2. Composition immunogène selon la revendication 1, la conductivité de la composition étant de 4 mS/cm ou inférieure.
3. Composition immunogène selon la revendication 2, la conductivité de la composition étant de 3 mS/cm ou inférieure.
4. Composition immunogène selon l'une quelconque des revendications 1 à 3, la concentration des sels dans ladite composition étant de 40 mM ou inférieure.
5. Composition immunogène selon l'une quelconque des revendications 1 à 4, comprenant en outre un agent de tonicité non ionique.
6. Composition immunogène selon la revendication 5, dans laquelle l'agent de tonicité non ionique est un polyol.
7. Composition immunogène selon la revendication 6, dans laquelle le polyol est le sorbitol et dans laquelle la concentration de sorbitol se situe entre environ 4 et environ 6 % (p/v).
8. Composition immunogène selon l'une quelconque des revendications 1 à 7, dans laquelle la concentration de saccharose se situe entre environ 4 et environ 6 % (p/v).
9. Composition immunogène selon l'une quelconque des revendications 1 à 8, comprenant en outre un ou plusieurs immunostimulants.
10. Composition immunogène selon la revendication 9, la composition immunogène comprenant du QS21.
11. Composition immunogène selon la revendication soit 9 soit 10, la composition immunogène comprenant du monophosphoryl-lipide A 3-dés-O-acylé.
12. Composition immunogène selon l'une quelconque des revendications 1 à 11, dans laquelle l'osmolalité est de 250 à 750 mOsm/kg.
13. Composition immunogène selon l'une quelconque des revendications 1 à 12, la composition étant fournie sous la forme d'une dose unitaire d'entre 50 µl et 1 ml.
14. Composition immunogène selon la revendication 13, la dose unitaire contenant 5 à 50 µg de protéine apparentée à M72.
15. Composition immunogène selon la revendication 13 ou 14, la dose unitaire étant une dose humaine et contenant entre 1 et 100 µg de 3D-MPL.
16. Composition immunogène selon l'une quelconque des revendications 13 à 15, la dose unitaire étant une dose humaine et contenant entre 1 et 100 µg de QS21.

EP 2 651 437 B1

17. Composition immunogène selon l'une quelconque des revendications 1 à 16, le pH de ladite composition se situant dans la plage de 7,5 à 8,5.
- 5 18. Composition immunogène selon l'une quelconque des revendications 1 à 17, dans laquelle l'antigène apparenté à M72 comprend la séquence d'acides aminés de SEQ ID NO : 3.
19. Composition immunogène selon l'une quelconque des revendications 1 à 17, dans laquelle l'antigène apparenté à M72 est constitué d'une séquence présentant au moins 90 % d'identité avec SEQ ID NO : 1.
- 10 20. Composition immunogène selon la revendication 19, dans laquelle l'antigène apparenté à M72 est constitué de la séquence d'acides aminés de SEQ ID NO : 3.
21. Composition immunogène selon l'une quelconque des revendications 1 à 20, pour une utilisation en médecine.
- 15 22. Composition immunogène selon l'une quelconque des revendications 1 à 21, pour une utilisation dans un procédé de prophylaxie, de traitement ou d'amélioration d'une infection par une mycobactérie.
- 20 23. Composition immunogène pour une utilisation selon la revendication soit 21 soit 22, pour une administration à un être humain.
- 25
- 30
- 35
- 40
- 45
- 50
- 55

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6a






Figure 6b

Figure 6c




Figure 6d




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2006117240 A [0003] [0005] [0030] [0051]
- WO 8809336 A [0065]
- WO 9633739 A [0067]
- US 4436727 A [0071]
- US 4877611 A [0071]
- US 4866034 A [0071]
- US 4912094 A [0071]
- GB 2220211 A [0071]
- WO 9421292 A [0071]
- WO 9850399 A [0072]
- US 6303347 B [0072]
- US 6764840 B [0072]
- WO 2003011223 A [0073]
- WO 2003099195 A [0073]
- WO 803022 A [0073]
- WO 803058 A [0073]
- WO 803732 A [0073]
- WO 804053 A [0073]
- WO 804057 A [0073]
- WO 804058 A [0073]
- WO 804059 A [0073]
- WO 804442 A [0073]
- WO 804680 A [0073]
- WO 804764 A [0073]
- WO 2008142133 A [0078]
- WO 9602555 A [0078]
- WO 9933488 A [0078]
- US 5865462 A [0078]
- US 5666153 A [0079]
- US 5278302 A [0079]
- WO 9526204 A [0079]
- EP 0382271 A [0082]
- US 61422723 B [0208]

Non-patent literature cited in the description

- World Health Organisation. *Tuberculosis Facts*, 2010 [0002] [0025]
- LOCHT et al. *Expert Opin. Biol. Ther.*, 2007, vol. 7 (11), 1665-1677 [0021]
- ANDERSON P. *Trends in Microbiology*, 2007, vol. 15 (1), 7-13 [0021]
- EHLERS S. *Infection*, 2009, vol. 37 (2), 87-95 [0021]
- CARDANA P-J. *Inflammation & Allergy - Drug Targets*, 2006, vol. 6, 27-39 [0022]
- CARDANA P-J. *Infection*, 2009, vol. 37 (2), 80-86 [0022]
- BRANDT et al. *Infect. Immun.*, 2004, vol. 72 (11), 6622-6632 [0030]
- SKEIKY et al. *J. Immunol.*, 2004, vol. 172, 7618-7628 [0030]
- TSENOVA et al. *Infect. Immun.*, 2006, vol. 74 (4), 2392-2401 [0030]
- REED et al. *PNAS*, 2009, vol. 106 (7), 2301-2306 [0030]
- VON ESCHEN et al. *Human Vaccines*, 2009, vol. 5 (7), 475-482 [0030]
- PAUL. *Fundamental Immunology*. 1993, 243-247 [0033]
- BEIßBARTH et al. *Bioinformatics*, 2005, vol. 21, i29-i37 [0033]
- CREIGHTON. *Proteins*, 1984 [0035]
- SMITH ; WATERMAN. *Adv. Appl. Math.*, 1981, vol. 2, 482 [0042]
- NEEDLEMAN ; WUNSCH. *J. Mol. Biol.*, 1970, vol. 48, 443 [0042]
- PEARSON ; LIPMAN. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 2444 [0042]
- Current Protocols in Molecular Biology. 1995 [0042]
- FENG ; DOOLITTLE. *J. Mol. Evol.*, 1987, vol. 35, 351-360 [0043]
- HIGGINS ; SHARP. *CABIOS*, 1989, vol. 5, 151-153 [0043]
- DEVEREAUX et al. *Nuc. Acids Res.*, 1984, vol. 12, 387-395 [0043]
- ALTSCHUL et al. *Nuc. Acids Res.*, 1977, vol. 25, 3389-3402 [0044]
- ALTSCHUL et al. *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0044]
- HENIKOFF ; HENIKOFF. *Proc. Natl. Acad. Sci. USA*, 1989, vol. 89, 10915 [0044]
- KARLIN ; ALTSCHUL. *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-5787 [0045]
- Archiv. für die gesamte Virusforschung. DALSGAARD et al. Saponin adjuvants. Springer Verlag, 1974, vol. 44, 243-254 [0065]
- SABROE et al. *J Immunol*, 2003, 1630-5 [0070]
- Harrison's Principles of Internal Medicine. 2005, 953-966 [0088] [0089]

ANTIGÉN MYCOBACTERIUM-KÉSZÍTMÉNY

Szabadalmi igénypontok

1. Immunogén készítmény, amely M72-eredetű antigént („M72 related antigen”) tartalmaz, ahol az M72-eredetű antigén a SEQ ID NO: 1 szerinti szekvenciával legalább 90%-ban azonos szekvenciát tartalmaz, a készítmény pH-ja a 7,0-9,0 tartományban van, és a készítmény konduktivitása 5 mS/cm vagy alacsonyabb.
2. Az 1. igénypont szerinti immunogén készítmény, ahol a készítmény konduktivitása 4 mS/cm vagy alacsonyabb.
3. A 2. igénypont szerinti immunogén készítmény, ahol a készítmény konduktivitása 3 mS/cm vagy alacsonyabb.
4. Az 1-3. igénypontok bármelyike szerinti immunogén készítmény, ahol a készítményben a sók koncentrációja 40 mM vagy alacsonyabb.
5. Az 1-4. igénypontok bármelyike szerinti immunogén készítmény, amely tartalmaz továbbá nem ionos, tonicitásra ható ágenszt.
6. Az 5. igénypont szerinti immunogén készítmény, ahol a nem ionos, tonicitásra ható ágens poliol.
7. A 6. igénypont szerinti immunogén készítmény, ahol a poliol szorbitol, és amelyben a szorbitolkoncentráció mintegy 4-6 vegyes%.
8. Az 1-7. igénypontok bármelyike szerinti immunogén készítmény, amelyben a szacharózkoncentráció mintegy 4-6 vegyes%.
9. Az 1-8. igénypontok bármelyike szerinti immunogén készítmény, amely tartalmaz továbbá egy vagy több immunstimulánst.
10. A 9. igénypont szerinti immunogén készítmény, ahol az immunogén készítmény QS21-et tartalmaz.
11. A 9. vagy 10. igénypontok bármelyike szerinti immunogén készítmény, ahol az immunogén készítmény 3-de-O-acilezett monofoszforil-lipid-A-t tartalmaz.
12. Az 1-11. igénypontok bármelyike szerinti immunogén készítmény, amelyben az ozmolalitás 250-750 mOsm/kg.
13. Az 1-12. igénypontok bármelyike szerinti immunogén készítmény, amely készítmény 50 µl és 1 ml közötti egységdózisban van kiszerve.

14. A 13. igénypont szerinti immunogén készítmény, amelyben az egységdózis 5-50 µg M72-eredetű fehérjét tartalmaz.

15. A 13. vagy 14. igénypont szerinti immunogén készítmény, ahol az egységdózis humán dózis és 1-100 µg 3D-MPL-t tartalmaz.

16. A 13-15. igénypontok bármelyike szerinti immunogén készítmény, ahol az egységdózis humán dózis és 1-100 µg QS21-et tartalmaz.

17. Az 1-16. igénypontok bármelyike szerinti immunogén készítmény, ahol a készítmény pH-ja 7,5-8,5.

18. Az 1-17. igénypontok bármelyike szerinti immunogén készítmény, ahol az M72-eredetű antigén SEQ ID NO. 3 szerinti aminosav-szekvenciát tartalmaz.

19. Az 1-17. igénypontok bármelyike szerinti immunogén készítmény, ahol az M72-eredetű antigén a SEQ ID NO. 1 szerinti szekvenciával legalább 90%-ban azonos szekvenciából áll.

20. A 19. igénypont szerinti immunogén készítmény, ahol az M72-eredetű antigén SEQ ID NO. 3 szerinti aminosavszekvenciából áll.

21. Az 1-20. igénypontok bármelyike szerinti immunogén készítmény gyógyászatban történő alkalmazásra.

22. Az 1-21. igénypontok bármelyike szerinti immunogén készítmény mycobaktériumok általi fertőzés megelőzésében, kezelésében vagy enyhítésében történő alkalmazásra.

23. A 21. vagy 22. igénypont szerinti immunogén készítmény az ott meghatározott alkalmazásra, embernek történő beadásra.