



(86) Date de dépôt PCT/PCT Filing Date: 1990/12/05
 (87) Date publication PCT/PCT Publication Date: 1991/06/13
 (45) Date de délivrance/Issue Date: 2001/08/14
 (85) Entrée phase nationale/National Entry: 1991/08/01
 (86) N° demande PCT/PCT Application No.: JP 90/01581
 (87) N° publication PCT/PCT Publication No.: WO 91/08183
 (30) Priorités/Priorities: 1989/12/05 (1-314253) JP;
 1989/12/05 (1-314252) JP; 1989/12/19 (1-327338) JP;
 1990/03/01 (2-47232) JP; 1990/03/08 (2-54839) JP;
 1990/07/31 (2-201305) JP

(51) Cl.Int.⁵/Int.Cl.⁵ C07C 17/28, C07C 19/08

(72) Inventeurs/Inventors:
 Takei, Ryutaro, JP;
 Kawasaki, Toru, JP;
 Yanase, Koichi, JP;
 Tanuma, Toshihiro, JP;
 Okamoto, Hidekazu, JP;
 Ohnishi, Keiichi, JP

(73) Propriétaire/Owner:
 ASAHI GLASS COMPANY LTD., JP

(74) Agent: SMART & BIGGAR

(54) Titre : METHODE DE PRODUCTION DE DICHLOROPENTAFLUOROPROPANES
 (54) Title: METHOD FOR PRODUCING A DICHLOROPENTAFLUOROPROPANES

(57) **Abrégé/Abstract:**

A method for producing a dichloropentafluoropropane, which comprises reacting dichlorofluoromethane (R21) with tetrafluoroethylene (4F) in the presence of a Lewis acid catalyst for addition reaction to obtain dichloropentafluoropropane, wherein a halide containing at least one element selected from the group consisting of Sb, Nb, Ta, B, Ga, In, Zr, Hf and Ti, or AlBr₃, or AlI₃, is used as the Lewis acid.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 19/08, 17/26	A1	(11) International Publication Number: WO 91/08183 (43) International Publication Date: 13 June 1991 (13.06.91)																		
(21) International Application Number: PCT/JP90/01581 (22) International Filing Date: 5 December 1990 (05.12.90) (30) Priority data: <table border="0" style="width: 100%;"> <tr> <td style="width: 30%;">1/314252</td> <td style="width: 40%;">5 December 1989 (05.12.89)</td> <td style="width: 30%;">JP</td> </tr> <tr> <td>1/314253</td> <td>5 December 1989 (05.12.89)</td> <td>JP</td> </tr> <tr> <td>1/327338</td> <td>19 December 1989 (19.12.89)</td> <td>JP</td> </tr> <tr> <td>2/47232</td> <td>1 March 1990 (01.03.90)</td> <td>JP</td> </tr> <tr> <td>2/54839</td> <td>8 March 1990 (08.03.90)</td> <td>JP</td> </tr> <tr> <td>2/201305</td> <td>31 July 1990 (31.07.90)</td> <td>JP</td> </tr> </table> (71) Applicant (for all designated States except US): ASAHI GLASS COMPANY LTD. [JP/JP]; 1-2, Marunouchi 2-chome, Chiyoda-ku, Tokyo 100 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only) : OHNISHI, Keiichi [JP/JP]; 62, Kuritaya, Kanagawa-ku, Yokohama-shi, Kanagawa 221 (JP). OKAMOTO, Hidekazu [JP/JP]; 2-59-1, Tsurugamine, Asahi-ku, Yokohama-shi, Kanagawa 241 (JP). TANUMA, Toshihiro [JP/JP]; 2-24-31, Konan, Konan-ku, Yokohama-shi, Kanagawa 233 (JP). YANASE, Koichi [JP/JP]; 5232-2, Goi, Ichihara-shi, Chiba 290 (JP). KAWASAKI, Toru [JP/JP]; 1846-1, Goi, Ichihara-shi, Chiba 290 (JP). TAKEI, Ryutaro [JP/JP]; 2-20-10, Hon-cho, Nakano-ku, Tokyo 164 (JP).		1/314252	5 December 1989 (05.12.89)	JP	1/314253	5 December 1989 (05.12.89)	JP	1/327338	19 December 1989 (19.12.89)	JP	2/47232	1 March 1990 (01.03.90)	JP	2/54839	8 March 1990 (08.03.90)	JP	2/201305	31 July 1990 (31.07.90)	JP	(74) Agents: YAMAMOTO, Ryoza et al.; Torimoto Kogyo Bldg., 38, Kanda-Higashimatsushitacho, Chiyoda-ku, Tokyo 101 (JP). (81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), KR, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i>
1/314252	5 December 1989 (05.12.89)	JP																		
1/314253	5 December 1989 (05.12.89)	JP																		
1/327338	19 December 1989 (19.12.89)	JP																		
2/47232	1 March 1990 (01.03.90)	JP																		
2/54839	8 March 1990 (08.03.90)	JP																		
2/201305	31 July 1990 (31.07.90)	JP																		
(54) Title: METHOD FOR PRODUCING DICHLOROPENTAFLUOROPROPANES																				
(57) Abstract <p>A method for producing a dichloropentafluoropropane, which comprises reacting dichlorofluoromethane (R21) with tetrafluoroethylene (4F) in the presence of a Lewis acid-catalyst for addition reaction to obtain dichloropentafluoropropane, wherein a halide containing at least one element selected from the group consisting of Sb, Nb, Ta, B, Ga, In, Zr, Hf and Ti, or AlBr₃, or AlI₃, is used as the Lewis acid.</p>																				

- 1 -

DESCRIPTION

TITLE OF THE INVENTION


5 METHOD FOR PRODUCING DICHLOROPENTAFLUOROPROPANES

TECHNICAL FIELD

The present invention relates to a method for producing dichloropentafluoropropanes (R225s).

10 Hydrochlorofluoropropanes are expected to be useful as foaming agents, cooling media or cleaning agents like conventional chlorofluorocarbons.

As a method for producing the dichloropentafluoropropanes (R225s), it is known to synthesize 3,3-dichloro-1,1,1,2,2-pentafluoropropane 15 (R225ca) and 1,3-dichloro-1,1,2,2,3-pentafluoropropane (R225cb) as shown below by adding tetrafluoroethylene to dichlorofluoromethane in the presence of aluminum chloride:



(O. Paleta et al., Collect. Czech. Chem. Commun., 36, 1867 (1971)). However, in this reaction, dichlorofluoromethane is disproportionated as shown by

71416-5

2

the following formula,



and by-product chloroform (R20) which is hardly separable by a
 5 usual method such as distillation is formed in a large amount.
 Thus, this method has a disadvantage that a multi-step
 purification process is required to obtain a product in high
 purity.

The present invention provides a method for producing
 10 a dichloropentafluoropropane, which comprises reacting
 dichlorofluoromethane (R21) with tetrafluoroethylene (4F) in
 the presence of a Lewis acid catalyst for addition reaction to
 obtain dichloropentafluoropropane, wherein said Lewis acid is a
 halide containing at least one element selected from the group
 15 consisting of Sb, Nb, Ta, Zr, Hf and Ti, or a fluoride of the
 formula $\text{MX}\alpha\text{F}\beta$, wherein M is one of atoms selected from Group
 IVa, Group Va and Group IIIb, or a mixture of such atoms, X is
 one of Cl, Br and I atoms, or a mixture of such atoms, α is a
 real number of $0 < \alpha < 5$, and β is a real number of $0 < \beta \leq 3.5$
 20 in the case of Group IVa, a real number of $0 < \beta \leq 4.5$ in the
 case of Group Va, and a real number of $0 < \beta \leq 2.5$ in the case
 of Group IIIb, provided that $\alpha + \beta$ is an integer of from 3 to
 5.

The halide containing at least one element selected from the group consisting of Sb, Nb, Ta, B, Ga, In, Zr, Hf and Ti includes chlorides such as $SbCl_5$, $NbCl_5$, $TaCl_5$, BCl_3 , $GaCl_2$, $GaCl_3$, $ZrCl_4$, $HfCl_4$, $InCl_3$ and $TiCl_4$, and partially fluorinated compounds of such chlorides; fluorides such as SbF_5 , NbF_5 , TaF_5 and BF_3 , and partially chlorinated compounds of such fluorides; bromides and iodides such as $TaBr_5$, $AlBr_3$, AlI_3 , BBr_3 , BI_3 , $GaBr_3$, GaI_3 , $HfBr_4$, HfI_4 , $InBr_3$, InI_3 and $TiBr_4$, and partially chlorinated or fluorinated compounds of such bromides or iodides. The partially fluorinated chlorides and partially chlorinated fluorides include, for example, $TiCl_2F_2$, $TiClF_3$ and $ZrCl_2F_2$.

The fluoride of the formula $MX_\alpha F_\beta$ can readily be prepared by treating a halide of the formula:



wherein M is one of atoms selected from the group consisting of Group IVa, Group Va and Group IIIb, or a mixture of such atoms, X is one of Cl, Br and I atoms, or a mixture of such atoms, α is $\alpha = 4$ in the case of Group IVa, $\alpha = 5$ in the case of Group Va, and $\alpha = 3$ in the case of Group IIIb, for example, a chloride such as BCl_3 , $AlCl_3$, $GaCl_3$, $InCl_3$, $TiCl_4$, $ZrCl_4$, $HfCl_4$, $NbCl_5$ or $TaCl_5$, or a bromide or iodide such as $GaBr_3$, GaI_3 , $InBr_3$, InI_3 , $TaBr_5$, $AlBr_3$, AlI_3 , BBr_3 , BI_3 , $TiBr_4$, TiI_4 , $ZrBr_4$, ZrI_4 , $HfBr_4$, HfI_4 , $AlBr_3$ or AlI_3 , with a suitable fluorinating

agent, for example, a chlorofluorocarbon (CFC) such as trichlorofluoromethane (R11), dichlorodifluoromethane (R12) or trichlorotrifluoromethane (R113), a hydrochlorofluorocarbon (HCFC) such as
5 dichlorofluoromethane (R21) or chlorodifluoromethane (furon 22), hydrogen fluoride or fluorine gas.

The treating conditions vary depending upon the halide and the fluorine source. However, it is usual to employ at least an equimolar amount of the fluorine
10 source to the halide.

The reaction temperature is usually from -50 to 200°C, preferably from -20 to 100°C, when chlorofluoromethane or hydrochlorofluoromethane is employed, and usually from -20 to 200°C, preferably from
15 0 to 150°C, when other fluorine source such as hydrogen fluoride is employed.

The reaction time is usually from 10 minutes to 2 weeks, preferably from 1 hour to 1 day, when chlorofluoromethane or hydrochlorofluoromethane is
20 employed, and usually from 30 minutes to two weeks, preferably from one hour to one day, when other fluorine source such as hydrogen fluoride is employed.

The content of fluorine atom contained in the fluoride of the formula MX_aF_b is preferably selected to be
25 within a proper range in order to increase the yield of the dichloropentafluoropropanes while suppressing the production of chloroform. The range may vary depending

- 5 -

upon the particle size when the fluoride is solid.

However, it is usually $0 < \beta \leq 3.5$, preferably $1 < \beta \leq 3$,
in the case of Group IVa, usually $0 < \beta \leq 4.5$, preferably
 $1 \leq \beta \leq 4$, in the case of Group Va, and usually $0 < \beta \leq$
5 2.5 , preferably $0.01 \leq \beta \leq 2$, more preferably $0.1 \leq \beta \leq$
2, in the case of Group IIIb.


The addition reaction of R21 to tetrafluoroethylene
in the presence of a Lewis acid catalyst can be conducted
in an inert solvent such as
10 perfluorobutyltetrahydrofuran. However, in order to
facilitate the purification, it is usually preferred to
conduct the reaction in the absence of a solvent.

The amount of the catalyst varies depending upon the
type of the catalyst used. However, it is usually from
15 0.01 to 50% by weight, preferably from 0.1 to 10% by
weight, relative to the starting material. The reaction
is conducted usually within a temperature range of from
-80 to 200°C, preferably from -20 to 100°C. The reaction
pressure is usually from 0 to 30 kg/cm²·G, preferably
20 from 0 to 15 kg/cm²·G.


The amount of tetrafluoroethylene to be added is
usually preferably at least equimolar to R21 to increase
the conversion of R21.

If the addition reaction of R21 to
25 tetrafluoroethylene (4F) in the presence of a Lewis acid
catalyst is conducted under such reaction condition that
the molar ratio of 4F to R21 is at least equimolar,

preferably in an excess amount, the disproportionation reaction of R21 is substantially suppressed, whereby as shown by the following formula:



3,3-dichloro-1,1,1,2,2-pentafluoropropane (R225ca) and 1,3-dichloro-1,1,2,2,3-pentafluoropropane (R225cb) can be obtained in good yield, and chloroform produced in a small amount as a by-product by the disproportionation, will react with tetrafluoroethylene present in an excess amount to give 1,3,3-trichloro-1,1,2,2-tetrafluoropropane (R224ca) as shown by the following formula:



R224ca has a boiling point different from R225 and can easily be separated by distillation.

The amount of tetrafluoroethylene to be added is preferably at least equimolar to R21, although it may depend also on the amount of the Lewis acid. Preferably, the molar ratio of 4F to R21 is $1.01 \leq 4F/R21 \leq 10$, particularly $1.01 \leq 4F/R21 \leq 5$, more preferably $1.1 \leq 4F/R21 \leq 3$. The manner of supplying tetrafluoroethylene (4F) and dichlorofluoromethane (R21) is not particularly limited so long as the reaction condition is eventually

such that 4F is present in an equimolar amount, preferably in an excess amount, to R21. However, in order to suppress the disproportionation reaction of R21, it is preferred to supply them continuously to the reactor while maintaining the molar ratio in supply of 4F to R21 within a range of $1 \leq 4F/R21$, preferably $1.01 \leq 4F/R21 \leq 10$, particularly $1.01 \leq 4F/R21 \leq 5$, more preferably $1.1 \leq 4F/R21 \leq 3$, and to withdraw the reaction product containing R225 from the reactor continuously.

10 The excess amount of 4F can be recovered after the reaction. Therefore, 4F can be used in large excess at a level of 10 mol times or more, but it is not an economical operation.

The Lewis acid catalyst may be supplied preliminarily to the reactor, or it may be supplied continuously to the reactor together with 4F and R21 and withdrawn continuously from the reactor together with the reaction product. In the latter case, the recovered Lewis acid catalyst can be recycled.

20 If R21 remains in the reaction product, disproportionation takes place to form by-products such as chloroform. Therefore, in order to improve the selectivity for the desired reaction, it is always preferred to control the concentration of R21 to the minimum level. To minimize the concentration of R21 in the reactor, it is preferred to supply the starting

25

materials R21 and 4F (and the catalyst) continuously to the reactor and to withdraw the reaction product continuously, so that the concentrations of the respective components can be maintained at constant levels, and if 4F is used in excess, it is possible to suppress the concentration of R21 to the minimum level. Thus, such an operation is preferred.

The reactor to be used for the continuous operation may be any so-called continuous reactor and may be of either continuous stirred tank reactor or plug flow reactor.

Further, it is necessary to feed solvent for the reaction at the initiation of the reaction. However, in the case of a continuous operation, as the reaction continuously proceeds, the solvent for reaction will gradually be replaced by the reaction product. Therefore, the solvent for reaction is not particularly restricted, so long as it does not adversely affect the main reaction.

As for the solvent at the initiation of the reaction, it is preferred to employ PFC such as perfluorooctane or perfluorobutyltetrahydrofuran, CFC such as 1,1,1-trichloropentafluoropropane (R215cb), 1,1,3-trichloropentafluoropropane (R215ca) or 1,1,1,3-tetrachlorotetrafluoropropane (R214cb), or HCFC such as 3,3-dichloro-1,1,1,2,2-pentafluoropropane (R225ca), 1,3-dichloro-1,1,2,2,3-pentafluoropropane (R225cb) or 1,3,3-

trichloro-1,1,2,2-tetrafluoropropane (R224ca). However, it is possible to conduct the reaction without solvent. The reaction is conducted usually within a temperature range of from -80 to 200°C, preferably from -20 to 100°C.

5 The reaction pressure is usually from 0 to 20 kg/cm², preferably from 0 to 10 kg/cm².

In the case of a continuous operation, the resident time of the reaction solution is usually from 0.1 minute to 24 hours, preferably from 1 minute to 10 hours, 10 although it depends on the reaction temperature and the type of the Lewis acid catalyst used.

The amount of the Lewis acid catalyst is usually from 0.1 to 50 mol%, preferably from 0.1 to 10 mol%, relative to R21. The catalyst to be used for this reaction is not 15 particularly restricted so long as it is a Lewis acid. However, it is preferred to employ a halide containing at least one element selected from the group consisting of Al, Sb, Nb, Ta, B, Ga, In, Zr, Hf and Ti. For example, it is possible to employ chlorides such as GaCl₂, GaCl₃, 20 ZrCl₄, BCl₃, AlCl₃, HfCl₄, InCl₃ and TiCl₄, or partially fluorinated compounds thereof, or bromides and iodides such as GaBr₃, GaI₃, HfBr₄, HfI₄, InBr₃, InI₃, and TiBr₄, or partially chlorinated or fluorinated compounds thereof, such as TiCl₂F₂, TiClF₃, and ZrCl₂F₂.

25 Further, chlorides such as SbCl₅, NbCl₅, TaCl₅, BCl₃ and partially fluorinated compounds thereof, fluorides such as SbF₅, NbF₅, TaF₅, and BF₃ and partially

chlorinated compounds thereof, and bromides and iodides such as $TaBr_5$, $AlBr_3$, AlI_3 , BBr_3 and BI_3 and partially chlorinated or fluorinated compounds thereof, may be used.

5 Now, the present invention will be described in further detail with reference to Examples. However, it should be understood that the present invention is by no means restricted by such specific Examples.

Example 1

10 Into a 200 ml Hastelloy C autoclave, 2 g of antimony pentachloride was added and cooled to $-78^\circ C$. Then, the autoclave was deaerated under reduced pressure, and 155 g (1.5 mol) of R21 was added thereto. After charging 20 g of tetrafluoroethylene, the temperature was raised to
15 $30^\circ C$, and tetrafluoroethylene was continuously added with stirring. Twelve hours later, the supply of tetrafluoroethylene was stopped, and stirring was continued for further 4 hours. The total amount of tetrafluoroethylene charged was 150 g. The pressure was
20 returned to normal pressure, and then the reaction solution was washed with water, and about 220 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 1.

Table 1

Reaction product	Molar ratio (%)
R225	81
Chloroform	1
Other components	18

5

Example 2

The reaction was conducted in the same manner as in
10 Exampel 1 except that 2 g of niobium pentachloride was
used instead of antimony pentachloride, whereby 190 g of
the crude reaction product was recovered. The results of
the analyses by means of gas chromatography and NMR are
shown in Table 2.

15

Table 2

Reaction product	Molar ratio (%)
R225	85
Chloroform	2
Other components	13

20

Example 3

The reaction was conducted in the same manner in
Example 1 except that 2 g of tantalum pentachloride was
used instead of antimony pentachloride, whereby 190 g of
25 the crude reaction product was recovered. The results of
the analyses by means of gas chromatography and NMR are
shown in Table 3.

Table 3

Reaction product	Molar ratio (%)
R225	84
Chloroform	2
Other components	14

5

Example 4

10

The reaction was conducted in the same manner as in Example 1 except that 2 g of aluminum bromide was used instead of antimony pentachloride and the reaction temperature was changed to -10°C , whereby 240 g of the crude reaction product was recovered. The results of the

15 analyses by means of gas chromatography and NMR are shown in Table 4.

Table 4

Reaction product	Molar ratio (%)
R225	77
Chloroform	1
Other components	22

20

Example 5

The reaction was conducted in the same manner as in Example 1 except that 2 g of aluminum iodide was used instead of antimony pentachloride and the reaction temperature was changed to -10°C , whereby 240 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 5.

Table 5

Reaction product	Molar ratio (%)
R225	83
Chloroform	2
Other components	15

15 Example 6

Into a 200 ml Hastelloy C autoclave, 2 g (0.009 mol) of zirconium(IV) chloride was added and cooled to -78°C . Then, the autoclave was deaerated and 155 g (1.5 mol) of R21 was added thereto. After charging 20 g of tetrafluoroethylene, the temperature was raised to 10°C , and tetrafluoroethylene was continuously added while maintaining the reaction temperature within a range of from 10 to 20°C . Twelve hours later, the supply of tetrafluoroethylene was stopped, and stirring was continued for further 4 hours. The total amount of tetrafluoroethylene charged was 150 g. The pressure was returned to normal pressure, and then the reaction

solution was washed with water, and 260 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 6.

Table 6

Reaction product	Molar ratio (%)
R225ca	39
R225cb	42
Chloroform	1
R224ca (CClF ₂ CF ₂ CHCl ₂)	15
Other components	3

Example 7

The reaction was conducted in the same manner as in Example 6 except that 2 g of titanium tetrachloride was used instead of zirconium(IV) chloride, whereby 190 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 7.

Table 7

Reaction product	Molar ratio (%)
R225ca	37
R225cb	43
Chloroform	5
R224ca (CClF ₂ CF ₂ CHCl ₂)	10
Other components	5

Example 8

The reaction was conducted in the same manner as in Example 6 except that 2 g of hafnium trichloride was used instead of zirconium(IV) chloride, whereby 190 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 8.

Table 8

Reaction product	Molar ratio (%)
R225ca	40
R225cb	30
Chloroform	3
R224ca (CClF ₂ CF ₂ CHCl ₂)	16
Other components	11

Example 9

The reaction was conducted in the same manner as in Example 6 except that 2 g of hafnium tetrachloride was used instead of zirconium(IV) chloride and the reaction temperature was changed to from 10 to 20°C, whereby 240 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 9.

Table 9

Reaction product	Molar ratio (%)
R225ca	38
R225cb	41
Chloroform	2
R224ca (CClF ₂ CF ₂ CHCl ₂)	11
Other components	8

10 Example 10

The reaction was conducted in the same manner as in Example 6 except that 2 g of zirconium dichloride difluoride was used instead of zirconium(IV) chloride, whereby about 260 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 10.

Table 10

Reaction product	Molar ratio (%)
R225ca	36
R225cb	54
Chloroform	1
R224ca (CClF ₂ CF ₂ CHCl ₂)	6
Other components	3

25 Example 11

The reaction was conducted in the same manner as in Example 6 except that 2 g of titanium dichloride

difluoride was used instead of zirconium(IV) chloride, whereby about 260 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 11.

5

Table 11

10

Reaction product	Molar ratio (%)
R225ca	37
R225cb	53
Chloroform	1
R224ca (CClF ₂ CF ₂ CHCl ₂)	7
Other components	2

15

Comparative Example 1-I

Into a 200 ml Hastelloy C autoclave, 3 g (0.02 mol) of anhydrous aluminum chloride was added and cooled to -78°C. Then, the autoclave was deaerated, and 155 g (1.5 mol) of R21 was added thereto. The temperature was raised to 10°C, and tetrafluoroethylene was continuously added while maintaining the reaction temperature within a range of from 10 to 20°C. Twelve hours later, the supply of tetrafluoroethylene was stopped, and stirring was continued for further 4 hours. The total amount of tetrafluoroethylene charged was 120 g. The pressure was returned to normal pressure, and then the reaction solution was washed with water, and about 200 g of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 1-I.

Table 1-I

Reaction product	Molar ratio (%)
R225ca	27
R225cb	18
Chloroform	20
R224ca ($\text{CClF}_2\text{CF}_2\text{CHCl}_2$)	10
Other components	25

Preparation Example 1

Into a 2 ℓ three necked round bottom flask equipped with a reflux condenser, 200 g (1.5 mol) of anhydrous aluminum chloride and 2000 g (14.6 mol) of trichlorofluoromethane (R11) were charged under a nitrogen stream and gently stirred at 0°C for 12 hours. After leaving the reaction mixture to stand still for 1 hour, the supernatant was removed, followed by drying under reduced pressure to obtain $AlCl_{3-\beta}F_{\beta}$. The fluorine content β of this product was 0.1.

Preparation Example 2

Into a 2 ℓ three necked round bottom flask equipped with a reflux condenser, 200 g (0.86 mol) of anhydrous zirconium(IV) chloride and 2000 g (14.6 mol) of trichlorofluoromethane (R11) were charged under a nitrogen stream and stirred at 0°C for 2 hours. After leaving the mixture to stand still for 1 hour, the supernatant was removed, followed by drying under reduced pressure to obtain $ZrCl_{4-\beta}F_{\beta}$. The fluorine content β of this product was 1.6.

Preparation Example 3

Into a 2 ℓ three necked round bottom flask equipped with a reflux condenser, 200 g (1.1 mol) of anhydrous titanium tetrachloride and 2000 g (19.4 mol) of dichlorofluoromethane (R21) were charged under a nitrogen stream and stirred at 0°C for 12 hours. After leaving the mixture to stand still for 1 hour, the supernatant

was removed, followed by drying under reduced pressure to obtain $TiCl_{4-\beta}F_{\beta}$. The fluorine content β of this product was 2.3.

Preparation Example 4

5 Into a 1 ℓ Hastelloy C autoclave, 200 g (0,6 mol) of anhydrous tantalum pentachloride was charged, and the autoclave was deaerated under reduced pressure. Then, 500 g (25 mol) of anhydrous hydrogen fluoride was added thereto, and the mixture was stirred at 50°C for 5 hours.
10 Then, hydrogen fluoride and hydrogen chloride were removed under reduced pressure to obtain $TaCl_{5-\beta}F_{\beta}$. The fluorine content β of this product was 4.

Example 12

15 Into a 10 ℓ Hastelloy C autoclave, 0.1 kg of aluminum fluoride chloride prepared in Preparation Example 1 was charged, and the autoclave was deaerated under reduced pressure. Then, 3 kg (14.8 mol) of R225ca ($CF_3CF_2CHCl_2$) was added thereto. The autoclave was cooled to -10°C. Then, while maintaining the reaction temperature within a
20 range of from 0 to 5°C, tetrafluoroethylene and R21 were continuously added at the rates of 850 g/hr and 670 g/hr, respectively. Six hours later, the supply of tetrafluoroethylene and R21 was stopped, and stirring was continued for further 4 hours while maintaining the
25 reaction temperature within a range of from 5 to 10°C. The pressure was returned to normal pressure, and then the reaction solution was subjected to filtration,

whereby about 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and ^{19}F - NMR are shown in Table 12. The crude reaction product was purified by distillation to
 5 obtain 9.8 kg of R225 (dichloropentafluoropropane).

Table 12

Composition of the reaction solution	Molar ratio (%)
R225ca ($\text{CF}_3\text{CF}_2\text{CHCl}_2$)	71.5
R225cb ($\text{CClF}_2\text{CF}_2\text{CHClF}$)	23
Chloroform (CHCl_3)	0
R224ca ($\text{CClF}_2\text{CF}_2\text{CHCl}_2$)	2
Other components	3.5

20 Example 13

The reaction was conducted in the same manner as in Preparation Example 1 except that 2000 g (19.4 mol) of dichlorofluoromethane (R21) was used instead of trichlorofluoromethane (R11), to obtain aluminum fluoride
 25 chloride (β : 0.01). Then, using 0.1 kg of this product, the reaction was conducted in the same manner as in Example 12, whereby 10.8 kg of the crude reaction product

was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 13. The crude reaction product was purified by distillation to obtain 9.6 kg of R225 (dichloropentafluoropropane).

5 Example 14

The reaction was conducted in the same manner as in Example 12 except that 0.1 kg of zirconium fluoride chloride obtained in Preparation Example 2 was used, whereby 10.8 kg of the crude reaction product was
10 recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 13. The reaction product was purified by distillation to obtain 10.2 kg of R225 (dichloropentafluoropropane).

Example 15

15 The reaction was conducted in the same manner as in Preparation Example 2 except that 2000 g (19.4 mol) of dichlorofluoromethane (R21) was used instead of trichlorofluoromethane (R11), whereby zirconium fluoride chloride was prepared (β : 1.8). Then, using 0.1 kg of
20 this product, the reaction was conducted in the same manner as in Example 12, whereby 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 13. The crude reaction product was purified by
25 distillation to obtain 10.1 kg of R225 (dichloropentafluoropropane).

Example 16

The reaction was conducted in the same manner as in Example 12 except that 0.2 kg of tantalum fluoride chloride prepared in Preparation Example 4 was used, whereby 9.5 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 13. The crude reaction product was purified by distillation to obtain 8.5 kg of R225 (dichloropentafluoropropane).

10

Table 13

15

Example Nos.	13	14	15	16
Composition of the reaction products (%)				
R225ca	72	58	58	65
R225cb	22	40	41	23
Chloroform	0	0	0	1
R224ca	2	1.5	0.5	6.5
Others	4	0.5	0.5	4.5

20 Example 17

The reaction was conducted in the same manner as in Example 12 except that 0.1 kg of titanium fluoride chloride prepared in Preparation Example 3 was used, whereby 10.5 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 14. The crude reaction product was purified by distillation to obtain

25

9.5 kg of R225 (dichloropentafluoropropane).

Example 18

The reaction was conducted in the same manner as in Preparation Example 3 except that 2000 g (14.6 mol) of trichlorofluoromethane (R11) was used instead of dichlorofluoromethane (R21) and stirring was conducted at 10°C for one week, whereby titanium fluoride chloride was prepared (β : 2.2). Then, using 0.1 kg of this product, the reaction was conducted in the same manner as in Example 12, whereby 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 14. The crude reaction product was purified by distillation to obtain 9.6 kg of R225 (dichloropentafluoropropane).

Example 19

The reaction was conducted in the same manner as in Preparation Example 1 except that 200 g of hafnium tetrachloride was used instead of zirconium(IV) chloride, whereby hafnium fluoride chloride (β : 1.5) was prepared. Then, using 0.1 kg of this product, the reaction was conducted in the same manner as in Example 12, whereby 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 14. The crude reaction product was purified by distillation to obtain 9.9 kg of R225 (dichloropentafluoropropane).

Example 20

The reaction was conducted in the same manner as in Preparation Example 2 except that 200 g of hafnium tetrachloride was used instead of zirconium(IV) chloride, 5 2000 g (19.4 mol) of dichlorofluoromethane (R21) was used instead of trichlorofluoromethane (R11), whereby hafnium fluoride chloride (β : 1.7) was prepared. Then, using 0.1 kg of this product, the reaction was conducted in the same manner as in Example 12, whereby 10.6 kg of the 10 crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 14. The crude reaction product was purified by distillation to obtain 9.8 kg of R225 (dichloropentafluoropropane).

15

Table 14

Example Nos.	17	18	19	20
Composition of the reaction solution (%)				
R225ca	60	61	60	59
R225cb	37	36	38	39
Chloroform	0	0	0	0
R224ca	2.5	2.5	1.5	1.5
Others	0.5	0.5	0.5	0.5

25

Example 21

A 10 ℓ Hastelloy C autoclave was deaerated under

reduced pressure after 100 g of aluminum fluoride chloride prepared in Preparation Example 1 was charged. And then, 10 kg of 1,1,1-trichloropentafluoropropane (R215cb) was charged as an initial solvent. The
5 autoclave was cooled to -10°C . Then, while maintaining the reaction temperature within a range of from 0 to 5°C , tetrafluoroethylene, dichlorofluoromethane and the aluminum fluoride chloride prepared in Preparation
10 Example 1 were continuously added at the rates of 1300 g/hr, 1030 g/hr and 20 g/hr, respectively. The reaction was conducted with continuously discharging the reaction mixture which is the same amount as the charged one. From the results of the analyses by means of gas chromatography and ^{19}F -NMR, it was found that after 20
15 hours of the reaction, the initial solvent R215cb was not present in the reaction mixture. The composition of the reaction products after 30 hours is shown in Table 15. 10.7 kg of the crude reaction products recovered since 20 hours later was purified by distillation to obtain 9.2 kg
20 (yield: 86%) of dichloropentafluoropropane (R225).

Example 22

The reaction was conducted in the same manner as in Example 21 except that zirconium fluoride chloride prepared in Preparation Example 2 was used instead of
25 aluminum fluoride chloride. The composition of the reaction products after 30 hours of the reaction is shown in Table 15. 10.7 kg of a crude reaction products

- 27 -

recovered since 20 hours later was purified by distillation to obtain 9.2 kg (yield: 86%) of dichloropentafluoropropane (R225).

Example 23

5 The reaction was conducted in the same manner as in Example 21 except that 40 g/hr of titanium fluoride chloride prepared in Preparation Example 3 was supplied instead of aluminum fluoride chloride. The composition of the reaction products after 30 hours of the reaction
10 is shown in Table 15. 10.7 kg of the crude reaction products recovered since 20 hours later was purified by distillation to obtain 9.2 kg (yield: 86%) of dichloropentafluoropropane (R225).

Example 24

15 The reaction was conducted in the same manner as in Example 21 except that hafnium fluoride chloride used in Example 19 was employed instead of aluminum fluoride chloride. The composition of the reaction products after 30 hours of the reaction is shown in Table 15. 10.7 kg
20 of the crude reaction products recovered since 20 hours later was purified by distillation to obtain 9.2 kg (yield: 86%) of dichloropentafluoropropane (R225).

Example 25

25 The reaction was conducted in the same manner as in Example 21 except that 40 g/hr of tantalum fluoride chloride prepared in Preparation Example 4 was used instead of aluminum fluoride chloride. The composition

1

of the reaction products after 30 hours is shown in Table 15. 10.7 kg of the crude reaction product recovered after 20 hours was purified by distillation to obtain 9 kg (yield: 84%) of dichloropentafluoropropane (R225).

5

Table 15

Example Nos.	21	22	23	24	25
Composition of the reaction solution (%)					
R225ca	58	40	39	40	60
R225cb	38	58	57	58	35
Chloroform	0	0	0	0	0
R224ca	2	1	3	1	3
Others	2	1	1	1	2

10

15

Example 26

The reaction was conducted in the same manner as in Example 12 except that 0.1 kg of anhydrous aluminum chloride was used instead of aluminum fluoride chloride, whereby 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 16. The crude reaction product was purified by distillation to obtain 9.1 kg of R225 (dichloropentafluoropropane).

20

25

Example 27

The reaction was conducted in the same manner as in Example 14 except that anhydrous zirconium(IV) chloride was used instead of zirconium(IV) fluoride chloride, whereby 10.7 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 16. The crude reaction product was purified by distillation to obtain 9 kg of R225 (dichloropentafluoropropane).

10 Example 28

The reaction was conducted in the same manner as in Example 17 except that 0.1 kg of anhydrous titanium chloride was used instead of titanium fluoride chloride, whereby 10.5 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 16. The crude reaction product was purified by distillation to obtain 8.5 kg of R225 (dichloropentafluoropropane).

Example 29

20 The reaction was conducted in the same manner as in Example 19 except that 0.1 kg of anhydrous hafnium tetrachloride was used instead of hafnium fluoride chloride, whereby 10.5 kg of a crude reaction products was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 16. The crude reaction product was purified by distillation to obtain 8.9 kg of R225 (dichloropentafluoropropane).

Example 30

The reaction was conducted in the same manner as in Example 16 except that 0.1 kg of anhydrous tantalum pentachloride was used instead of tantalum fluoride chloride, whereby 9 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 16. The crude reaction product was purified by distillation to obtain 7.5 kg of R225 (dichloropentafluoropropane).

10

Table 16

Example Nos.	26	27	28	29	30
Composition of the reaction solution (%)					
R225ca	71	58	59	59	64
R225cb	22	40	37	38	22
Chloroform	0.5	0.5	1	0.5	2
R224ca	2	1	2.5	1.5	8
Others	4.5	0.5	0.5	1	4

20

Example 31

The reaction was conducted in the same manner as in Example 12 except that 3 kg (12.6 mol) of R215cb ($\text{CF}_3\text{CF}_2\text{CCl}_3$) was used instead of R225ca as the solvent for reaction and anhydrous aluminum chloride was used, whereby 10.5 kg of the crude reaction product was

25

recovered. The results of the analyses by means of gas chromatography and ^{19}F -NMR are shown in Table 17.

Table 17

Composition of the reaction solution	Molar ratio (%)
R225	
$\text{CF}_3\text{CF}_2\text{CHCl}_2$ (R225ca)	45
$\text{CClF}_2\text{CF}_2\text{CHClF}$ (R225cb)	23
R224	
$\text{CClF}_2\text{CF}_2\text{CHCl}_2$ (R224ca)	2
CHCl_3 (R20)	0
R215	
$\text{CF}_3\text{CF}_2\text{CCl}_3$ (R215cb)	24
Other components	6

The crude reaction solution was purified by distillation to obtain 6.6 kg (yield: 84%) of R225 (dichloropentafluoropropane).

Example 32

Into a 10 l Hastelloy C autoclave, 0.2 kg (1.5 mol) of anhydrous aluminum chloride was charged, and the autoclave was deaerated under reduced pressure. Then, autoclave was cooled to -10°C , and 5 kg (48.6 mol) of R21 (CHCl_2F) was added thereto. Then, while maintaining the reaction temperature within a range of from 0 to 5°C , tetrafluoroethylene was continuously added at the rate of

800 g/hr. The upper limit of the reaction pressure was 5 kg/cm², and the reaction was continued while purging the gas as the case required. Five hours later, the reaction temperature was raised to 20°C, and the reaction was continued with further supply of tetrafluoroethylene. When chloroform in the system was consumed, the reaction was stopped, and the pressure was returned to normal pressure. Then, the reaction solution was subjected to filtration, whereby 7.8 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and ¹⁹F-NMR are shown in Table 18.

Table 18

Composition of the reaction solution	Molar ratio (%)
R225	
CF ₃ CF ₂ CHCl ₂ (R225ca)	37
CClF ₂ CF ₂ CHClF (R225cb)	10
R224	
CClF ₂ CF ₂ CHCl ₂ (R224ca)	36
Other components	17

The crude reaction product was purified by distillation to obtain 3.5 kg (yield: 35%) of R225 (dichloropentafluoropropane).

Example 33

A 10 ℓ Hastelloy C autoclave was deaerated under reduced pressure. Then, 10 kg of R215cb (CF₃CF₂CCl₃) was

- 33 -

charged as the initial solvent.

The autoclave was cooled to -15°C . Then, while maintaining the reaction temperature within a range of from -10°C to -5°C , tetrafluoroethylene, R21 and aluminum chloride were continuously added at the rates of 1300 g/hr, 1030 g/hr and 20 g/hr (1.5 mol% relative to R21, same applied hereinafter), respectively. The reaction was conducted with continuously withdrawing the reaction mixture which is the same amount as the charged one.

From the results of the analyses by means of gas chromatography and ^{19}F -NMR, it was found that after 20 hours of the reaction, the initial solvent R215cb was not present in the reaction mixture. The composition of the reaction solution at that time is shown in Table 19.

15

Table 19

20

Composition of the reaction solution	Molar ratio (%)
R225	
$\text{CF}_3\text{CF}_2\text{CHCl}_2$ (R225ca)	57
$\text{CClF}_2\text{CF}_2\text{CHClF}$ (R225cb)	37
R224	
$\text{CClF}_2\text{CF}_2\text{CHCl}_2$ (R224ca)	3
CHCl_3 (R20)	0
Other components	3

25

The reaction mixture which is the same amount as the charged one was continuously withdrawn, and 10.7 kg of the crude reaction product thereby obtained was purified

by distillation to obtain 9.1 kg (yield: 85%) of R225 (dichloropentafluoropropane).

Example 34

The reaction was conducted in the same manner as in Example 33 except that the supplying rates of tetrafluoroethylene, R21 and aluminum chloride were doubled by twice, whereby 10.5 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and ^{19}F -NMR are shown in Table 20.

Table 20

Composition of the reaction solution	Molar ratio (%)
R225	
$\text{CF}_3\text{CF}_2\text{CHCl}_2$ (R225ca)	55
$\text{CClF}_2\text{CF}_2\text{CHClF}$ (R225cb)	36
R224	
$\text{CClF}_2\text{CF}_2\text{CHCl}_2$ (R224ca)	5
CHCl_3 (R20)	0
Other components	4

The crude reaction product was purified by distillation to obtain 8.8 kg (yield: 84%) of R225 (dichloropentafluoropropane).

Example 35

The reaction was conducted in the same manner as in Example 33 except that the supplying rate of aluminum

chloride was changed to 67 g/hr (5.5 mol%), whereby 9.8 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and ^{19}F -NMR are shown in Table 21.

5

Table 21

Composition of the reaction solution	Molar ratio (%)
R225	
$\text{CF}_3\text{CF}_2\text{CHCl}_2$ (R225ca)	53
$\text{CClF}_2\text{CF}_2\text{CHClF}$ (R225cb)	37
R224	
$\text{CClF}_2\text{CF}_2\text{CHCl}_2$ (R224ca)	5
CHCl_3 (R20)	1
Other components	4

10

15

The crude reaction product was purified by distillation to obtain 8.0 kg (yield: 82%) of R225 (dichloropentafluoropropane).

Example 36

20

The reaction was conducted in the same manner as in Example 26 except that the reaction temperature was changed in the range of 0 to 5°C, whereby 8.5 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and ^{19}F -NMR are shown in Table 22.

25

Table 22

Composition of the reaction solution	Molar ratio (%)
R225 $\text{CF}_3\text{CF}_2\text{CHCl}_2$ (R225ca) $\text{CClF}_2\text{CF}_2\text{CHClF}$ (R225cb)	52 36
R224 $\text{CClF}_2\text{CF}_2\text{CHCl}_2$ (R224ca)	5
CHCl_3 (R20)	0
Other components	7

The crude reaction product was purified by distillation to obtain 6.8 kg (yield: 80%) of R225 (dichloropentafluoropropane).

15 Example 37

The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to zirconium(IV) chloride, and the supplying rate of the catalyst was changed to 20 g/hr (0.9 mol%), whereby 11 kg of the crude reaction solution was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 23.

Table 23

	Composition of the reaction solution	Molar ratio (%)
5	R225ca	39
	R225cb	58
	Chloroform	0
	R224 (CCl ₂ F ₂ CF ₂ CHCl ₂)	2
10	Other components	1

The crude reaction solution was purified by distillation to obtain 9.5 kg (yield: 86%) of R225 (dichloropentafluoropropane).

15 Example 38

The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to titanium tetrachloride, and the supplying rate of the catalyst was changed to 20 g/hr (1.1 mol%), whereby 11 kg
20 of the crude reaction solution was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 24.

Table 24

5	Composition of the reaction solution	Molar ratio (%)
	R225ca	38
	R225cb	56
	Chloroform	0
	R224ca	4
10	Other components	2

The crude reaction solution was purified by distillation to obtain 9.2 kg (yield: 84%) of R225 (dichloropentafluoropropane).

15 Example 39

The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to zirconium dichloride difluoride, and the supplying rate of the catalyst was changed to 20 g/hr (1 mol%), whereby 20 11 kg of the crude reaction product was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 25.

Table 25

Composition of the reaction solution	Molar ratio (%)
R225ca	39
R225cb	58
Chloroform	0
R224ca (CClF ₂ CF ₂ CHCl ₂)	2
Other components	1

The crude reaction product was purified by distillation to obtain 9.5 kg (yield: 86%) of R225 (dichloropentafluoropropane).

Example 40

The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to titanium dichloride difluoride, and the supplying rate of the catalyst was changed to 20 g/hr (1.3 mol%), whereby 11 kg of the crude reaction solution was recovered. The results of the analyses by means of gas chromatography and NMR are shown in Table 26.

Table 26

Composition of the reaction solution	Molar ratio (%)
R225ca	38
R225cb	56
Chloroform	0
R224ca	4
Other components	2

5
10 The crude reaction solution was purified by distillation to obtain 9.2 kg (yield: 84%) of R225 (dichloropentafluoropropane).

Example 41

15 The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to gallium trichloride, and the supplying rate of the catalyst was changed to 20 g/hr (1.1 mol%), whereby 11 kg of a crude reaction solution was recovered. The results of the analyses by means of gas chromatography and NMR
20 are shown in Table 27.

Table 27

Composition of the reaction solution	Molar ratio (%)
R225ca	40
R225cb	50
Chloroform	1
R224ca	7
Other components	2

25

The crude reaction solution was purified by distillation to obtain 9 kg (yield: 81%) of R225 (dichloropentafluoropropane).

Example 42

5 The reaction was conducted in the same manner as in Example 33 except that the catalyst was changed to hafnium tetrachloride, and the supplying rate of the catalyst was changed to 20 g/hr (0.6 mol%), whereby 11 kg of the crude reaction solution was recovered. The
10 results of the analyses by means of gas chromatography and NMR are shown in Table 28.

Table 28

Composition of the reaction solution	Molar ratio (%)
R225ca	38
R225cb	57
Chloroform	0
R224ca	3
Other components	2

15 The crude reaction solution was purified by distillation to obtain 9.5 kg (yield: 86%) of R225 (dichloropentafluoropropane).

As shown by the Examples, according to the present
5 invention, dichloropentafluoropropane (R225), a high-purity product of which used to be hardly available, can be produced in good yield.


71416-5

42

CLAIMS:

1. A method for producing a dichloropentafluoropropane, which comprises reacting dichlorofluoromethane (R21) with tetrafluoroethylene (4F) in the presence of a Lewis acid
 5 catalyst for addition reaction to obtain dichloropentafluoropropane, wherein said Lewis acid is a halide containing at least one element selected from the group consisting of Sb, Nb, Ta, Zr, Hf and Ti, or a fluoride of the formula

10



wherein M is one of atoms selected from Group IVa, Group Va and Group IIIb, or a mixture of such atoms, X is one of Cl, Br and I atoms, or a mixture of such atoms, α is a real number of $0 < \alpha < 5$, and β is a real number of $0 < \beta \leq 3.5$ in the case of
 15 Group IVa, a real number of $0 < \beta \leq 4.5$ in the case of Group Va, and a real number of $0 < \beta \leq 2.5$ in the case of Group IIIb, provided that $\alpha + \beta$ is an integer of from 3 to 5.

2. The method according to Claim 1, wherein the fluoride is a compound obtained by fluorinating a halogen compound of
 20 the formula



wherein M is one of atoms selected from Group IVa, Group Va and Group IIIb, or a mixture of such atoms, X is one of Cl, Br, and I atoms, or a mixture of such atoms, and α is $\alpha = 4$ in the case
 25 of Group IVa, $\alpha = 5$ in the case of Group Va, and $\alpha = 3$ in the case of Group IIIb, with a fluorinating agent.

3. The method according to Claim 2, wherein a chlorofluorocarbon (CFC) or hydrochlorofluorocarbon (HCFC) is used as the fluorinating agent.

71416-5

43

4. The method according to Claim 1, wherein the reaction is conducted under the reaction condition such that the molar ratio of 4F to R21 is $1 \leq 4F/R21$.

5. The method according to Claim 4, wherein the molar ratio of 4F to R21 is $1.01 \leq 4F/R21 \leq 10$.

6. The method according to Claim 4, wherein 4F and R21 are continuously supplied to a reactor in a molar ratio of 4F to R21 being $1 \leq 4F/R21$, and the reaction product containing the dichloropentafluoropropane is continuously withdrawn from the reactor.

7. The method according to Claim 4, wherein the Lewis acid catalyst, 4F and R21 are continuously supplied to a reactor in a molar ratio of 4F to R21 being $1 \leq 4F/R21$, and the reaction product containing the dichloropentafluoropropane and the Lewis acid catalyst are continuously withdrawn from the reactor.

SMART & BIGGAR

OTTAWA, CANADA

PATENT AGENTS