

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2015/111060 A1

(43) International Publication Date
30 July 2015 (30.07.2015)

(51) International Patent Classification:

A61Q 13/00 (2006.01) *A61K 8/19* (2006.01)
A61K 8/11 (2006.01) *A61K 8/25* (2006.01)
A61K 8/06 (2006.01) *A61K 8/29* (2006.01)
A61K 8/81 (2006.01) *A61K 8/26* (2006.01)
A61K 8/73 (2006.01) *A61Q 15/00* (2006.01)

(21) International Application Number:

PCT/IL2015/050088

(22) International Filing Date:

22 January 2015 (22.01.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/930,539 23 January 2014 (23.01.2014) US

(71) Applicant: MAORI S.C. LTD. [IL/IL]; 108 Herzl, Apt. 401, 6655419 Tel Aviv (IL).

(72) Inventor: SHAVIT, Shoval; Herzl 108 apt. 401, 6655419 Tel Aviv (IL).

(74) Agents: GEYRA, Assaf et al.; S. Friedman & Co., P.O. Box 33123, 6133101 Tel Aviv (IL).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2015/111060 A1

(54) Title: SCENTED BODY COMPOSITIONS

(57) Abstract: A fragrance releasing complex with extended fragrance retention capability including an acrylic polymer, to be applied a human body surface is provided. The fragrance releasing complex can be applied by using a kit which includes means for applying the fragrance releasing complex to the body surface.

SCENTED BODY COMPOSITIONS

BACKGROUND OF THE INVENTION

[001] Since early human history, scents and fragrances have been used to modify body odor. They were obtained in raw form such as resins, gums or essential oils and derived from natural sources, such as the bark, roots, leaves and fruit of indigenous plants and trees. These raw materials were then diluted with water or other solvents and applied to the skin. In modern times, the chemical components responsible for the odor properties of these raw materials were isolated and identified. Current day perfumery engages in combining various fragrance materials to obtain novel fragrance compositions with specific "characteristics".

[002] In light of the advancements in chemical technology many fragrances are no longer derived from natural sources but are synthesized as highly pure fragrance raw materials (FRM). Furthermore, fragrances have been categorized into three "note" types based on their relative volatility; base, having the most long lasting aroma; middle having a medium volatility, and top notes being the most volatile. If used correctly the different note combinations can produce a "balanced fragrance" composition i.e. one which diffuses in a manner having an aesthetic pleasing effect.

[003] In addition, fragrances have been grouped according to the odor they produce, by using both broad and specific descriptions. For example, "floral" is a term used for odors associated with flowers while the term "lilac" is more specific. Additional examples of descriptive terms include "rose", "floral", "green", "citrus", "spicy", "honey", and "musk".

[004] Due to an uneven evaporation rate of the different components, the initial fragrance may be quite different from the aroma perceived several hours later. Several methods have been commonly employed to address this issue. One method is to "load up" on

the perfume initially and rely on the natural evaporation rate to diminish the fragrance so as to reach a suitable level several hours later when the desired effect is required. Another method is to continually reapply small amounts of the perfume to the skin at short time intervals. However neither of these solutions overcomes the problem of the diminishing level of top and middle notes over time. In fact, base notes which are present over an extended period in light of their low volatility, begin to accumulate with each renewed appliance of perfume, with the possible outcome of overwhelming the other fragrance notes and negating the original fragrance balance.

[005] Acrylic acid (prop-2-enoic acid) is an organic compound with the formula $\text{CH}_2=\text{CHCO}_2\text{H}$. It consists of a vinyl group connected directly to a carboxylic acid terminus and is the simplest unsaturated carboxylic acid. It is a colorless liquid with an acrid or tart smell. It is miscible with water, alcohols, ethers, and chloroform. It is used extensively in different forms and more than one billion kilograms are produced annually. Acrylic acid is produced from propene which is a byproduct of ethylene and gasoline production.

[006] Acrylate polymers belong to a group of polymers which are commonly referred to as plastics. Some of their notable characteristics include transparency, resistance to breakage, and elasticity. They are also generally known as acrylics or polyacrylates. Acrylate polymers are formed from Acrylate monomers which are based on the structure of acrylic acid or are derivatives of acrylic acid, such as methyl methacrylate in which one vinyl hydrogen and the carboxylic acid hydrogen are both replaced by methyl groups, and acrylonitrile in which the carboxylic acid group is replaced by the related nitrile group.

[007] Acrylic paint is a paint containing a pigment suspension in an acrylic polymer emulsion. It can be diluted with water, but becomes water-resistant when dry. Depending on the degree to which the paint is diluted with water, or modified with acrylic gels, media, or

pastes, the finished acrylic painting can have unique characteristics not attainable with other media.

SUMMARY OF THE INVENTION

[008] In one embodiment, this invention provides a composition and/or a fragrance-releasing complex comprising: (1) a fragrance; and (2) a fragrance retention composition, the fragrance retention composition comprises an acrylic polymer wherein the fragrance-releasing complex is adapted to extend fragrance retention. In one embodiment, this invention provides a composition comprising: (1) a fragrance; and (2) an acrylic polymer. In one embodiment, this invention provides a composition comprising: (1) a fragrance; and (2) an acrylic polymer, wherein the fragrance is embedded within the acrylic polymer.

[009] In a further embodiment, this invention provides a fragrance-releasing complex, comprising from 50 to 80% by weight polymer.

[010] In a further embodiment, this invention provides a composition and/or a fragrance-releasing complex, wherein the acrylic polymer is within an emulsion. In a further embodiment, this invention provides a fragrance-releasing complex, wherein the emulsion is an acrylic paint.

[011] In a further embodiment, this invention provides a composition and/or a fragrance-releasing complex, wherein the fragrance comprises: an alcohol, an ether, a nitrile, an aldehyde, an ester, a ketone, a lactone, a thiol, an amine, a schiff-base, a terpene, a cyclic alkene, a cyclic oxide, an oxime, an essential oil, an aromatic species, or any combination thereof.

[012] In a further embodiment, this invention provides a composition and/or a fragrance-releasing complex, wherein the fragrance comprises: i) a base note fragrance; ii) a middle

note fragrance; iii) a top note fragrance; iv) a balance carrier; v) a fixative, or any combination thereof.

[013] In a further embodiment, this invention provides a composition and/or a fragrance-releasing complex, wherein the fragrance comprises a solvent suitable for human topical
5 administration.

[014] In a further embodiment, this invention provides a composition and/or a fragrance-releasing complex, wherein the fragrance comprises a volatile fragrance.

[015] In a further embodiment, this invention provides a composition or a fragrance-releasing complex, retaining a perceptible fragrance for 15 to 40 hours.

10 [016] In a further embodiment, this invention provides a method for conferring, enhancing, improving or modifying the odor properties of a body surface, comprising contacting or treating the body surface with the fragrance-releasing complex.

[017] In a further embodiment, this invention provides a method for decorating the body surface.

15 [018] In a further embodiment, this invention provides a kit for applying a fragrance to a subject, comprising the composition and/or fragrance-releasing complex and means for topically applying the fragrance-releasing complex to a body surface of a subject.

[019] In a further embodiment, this invention provides a kit wherein applying a fragrance to a subject is further applying a body decoration.

20 [020] In a further embodiment, this invention provides a kit wherein the composition and/or the fragrance-releasing complex comprise a pigment.

DETAILED DESCRIPTION OF THE INVENTION

[021] In one embodiment, the present invention provides a composition comprising: a fragrance; an acrylic polymer, a thickener, an emulsifier, an aqueous solution, and a pigment. In another embodiment, the composition is a visible fragrance-releasing complex. In another embodiment, visible is visible light. In another embodiment, a composition of the invention is a scented body composition. In another embodiment, a composition of the invention is a topical scented body composition. In another embodiment, a composition of the invention is a topical scented colored body composition and/or decoration.

[022] In another embodiment, an aqueous solution is water, distilled water, or purified water. In another embodiment, an aqueous solution is a buffer. In another embodiment, an aqueous solution is an isotonic buffer.

[023] In another embodiment, an emulsifier is a nonionic emulsifier. In another embodiment, an emulsifier is any nonionic emulsifier known to one of average skill in the art. In another embodiment, an emulsifier is a combination of emulsifiers or a combination of nonionic emulsifiers.

[024] In another embodiment, a composition as described herein is stable after it was applied to the skin or a body surface and is dried. In another embodiment, a stable composition is a composition not affected by cracks and/or peelings. In another embodiment, a stable composition is a composition which releases noticeable amount of fragrance for at least 8 hours. In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance for at least 10 hours. In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance for at least 12 hours. In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance for at least 15 hours. In another embodiment, a stable

composition is a composition which releases a noticeable amount of fragrance for at least 20 hours. In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance for at least 24 hours. In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance for 8 to 36 hours.

[025] In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance within 2 to 120 seconds from applying it to a skin surface (immediate fragrance release). In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance within 1 to 15 seconds from applying it to a skin surface (immediate fragrance release). In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance within 5 to 30 seconds from applying it to a skin surface (immediate fragrance release). In another embodiment, a stable composition is a composition which releases a noticeable amount of fragrance within 10 to 40 seconds from applying it to a skin surface (immediate fragrance release).

[026] In another embodiment, an emulsifier is Glyceryl Stearate. In another embodiment, an emulsifier is a Polysorbate. In another embodiment, an emulsifier is Polysorbate 80. In another embodiment, an emulsifier is Polysorbate 20. In another embodiment, an emulsifier is Cetareth-20.

[027] In another embodiment, a thickener is a cosmetic thickener. In another embodiment, a thickener comprises cellulose or a derivative thereof. In another embodiment, a thickener is an aqueous thickener. In another embodiment, a thickener comprises clay. In another embodiment, a thickener comprises a gum. In another embodiment, a thickener comprises a carbomer. In another embodiment, a thickener comprises a polyethylene glycol. In another embodiment, a thickener comprises a non-aqueous thickener. In another embodiment, a thickener is an organic thickener. In another embodiment, a thickener is an organoclay.

[028] In another embodiment, a thickener is a clay based thickener having a density of at least 1 g/cm³. In another embodiment, a thickener is a clay based thickener having a density of 1 to 2.5 g/cm³. In another embodiment, a thickener is a clay based thickener having a density of 1.2 to 2.0 g/cm³. In another embodiment, a thickener is a clay based thickener having a density of 1 to 1.8 g/cm³. In another embodiment, a thickener is a clay based thickener having a density of 1.4 to 2.0 g/cm³. In another embodiment, a thickener is a clay based thickener having a density of about 1.6 g/cm³.

[029] In another embodiment, the invention provides a composition and/or a fragrance-releasing complex comprising: at least one fragrance and a fragrance retention composition.

10 In one embodiment, this invention provides a composition comprising: (1) a fragrance; and (2) an acrylic polymer. In one embodiment, this invention provides a composition comprising: (1) a fragrance; and (2) an acrylic polymer, wherein the fragrance is embedded within the acrylic polymer. In one embodiment, the fragrance retention composition comprises an acrylic polymer. In another embodiment, the fragrance-releasing complex is adapted to extend

15 fragrance retention within the acrylic polymer. In another embodiment, the acrylic polymer engulfs, encapsulates, and/or retains a fragrance. In another embodiment, the acrylic polymer provides a solid or a semi-solid vehicle to a fragment. In another embodiment, the acrylic polymer provides a gel, a semi-gel, or a viscous vehicle to a fragment. In another embodiment, the acrylic polymer is in the form of a pattern that adheres to a body surface. In

20 another embodiment, the fragrance-releasing complex is non-irritable, non-allergenic, non-toxic or any combination thereof to a body surface such as skin. In another embodiment, the acrylic polymer has a pH value of 7 to 9. In another embodiment, the acrylic polymer has a pH value of 7.5 to 8.5.

[030] In another embodiment, a composition as described herein forms a film once applied

25 to a surface. In another embodiment, the quality of film correlates to the stability of the

formulation on a surface. In another embodiment, the quality of film correlates to the stability of the formulation on the skin.

[031] In one embodiment, the term "comprises" includes or can be replaced with the term "consists".

5 [032] In one embodiment, the term "about", as used herein, means within 10% of the recited numerical value; In another embodiment, the term "about" as used herein, means within 5% of the recited numerical value.

[033] In one embodiment, the term "fragrance-releasing complex" is synonymous with the term "composition" as used herein.

10 [034] In another embodiment, a composition comprising: a fragrance; an acrylic polymer, a thickener, an emulsifier, an aqueous solution, and a pigment is used for skin or a body surface decoration. In another embodiment, a composition as described herein is a visible fragrance-releasing complex. In another embodiment, a composition as described herein is active for at least 10 hours. In another embodiment, a composition as described herein is active for at least
15 12 hours. In another embodiment, a composition as described herein is active for at least 15 hours. In another embodiment, a composition as described herein is active for at least 18 hours. In another embodiment, a composition as described herein is active for at least 20 hours. In another embodiment, a composition as described herein is active for 10 to 40 hours. In another embodiment, a composition as described herein is active for 10 to 30 hours. In
20 another embodiment, a composition as described herein is active for 10 to 25 hours.

[035] In another embodiment, active is visible upon placement on a body surface. In another embodiment, active is air exposed. In another embodiment, active is air exposed but not liquid exposed. In another embodiment, active is a noticeable amount of fragrance for at least 5 hours. In another embodiment, active is a noticeable amount of fragrance for at least 10 hours.

In another embodiment, active is a noticeable amount of fragrance for at least 15 hours. In another embodiment, active is a noticeable amount of fragrance for at least 25 hours. In another embodiment, active is a noticeable amount of fragrance for 5 to 40 hours. In another embodiment, active is a noticeable amount of fragrance for 10 to 40 hours. In another embodiment, active is a noticeable amount of fragrance for 15 to 40 hours. In another embodiment, active is a noticeable amount of fragrance for 15 to 35 hours.

[036] In another embodiment, noticeable scent or fragrance is when at least one healthy adult individual can detect or smell a fragrance from a composition placed at a distance of 5 cm from his or hers nose. In another embodiment, “noticeable” is smellable (capable of being smelled) amount of fragrance noticed by healthy and normal human nose of three adult female individuals from a distance of 5 cm from the composition as described herein. In another embodiment, unnoticeable scent or fragrance is when at least one healthy adult female individual of three adult female individuals cannot detect or smell a fragrance from a composition placed at a distance of 5 cm from her nose.

[037] In another embodiment, the composition as described herein comprises a fragrance in an amount of from 2 to 40% by weight of the total composition. In another embodiment, the composition as described herein comprises a fragrance in an amount of from 5 to 30% by weight of the total composition. In another embodiment, the composition as described herein comprises a fragrance in an amount of from 5 to 20% by weight of the total composition. In another embodiment, the composition as described herein comprises a fragrance in an amount of from 5 to 15% by weight of the total composition. In another embodiment, the composition as described herein comprises a fragrance in an amount of from 7 to 15% by weight of the total composition.

[038] In another embodiment, a recitation of weight percent as described herein is weight percent from/of the total composition.

[039] In another embodiment, the composition as described herein comprises from 0.2 to 8% by weight thickener. In another embodiment, the composition as described herein comprises from 0.2 to 5% by weight thickener. In another embodiment, the composition as described herein comprises from 0.2 to 4% by weight thickener. In another embodiment, the composition as described herein comprises from 0.5 to 3% by weight thickener. In another embodiment, the composition as described herein comprises from 0.5 to 2.5% by weight thickener.

[040] In another embodiment, the composition as described herein comprises from 0.2 to 8% by weight emulsifier. In another embodiment, the composition as described herein comprises from 0.2 to 5% by weight emulsifier. In another embodiment, the composition as described herein comprises from 0.2 to 4% by weight emulsifier. In another embodiment, the composition as described herein comprises from 0.5 to 3% by weight emulsifier. In another embodiment, the composition as described herein comprises from 0.5 to 2.5% by weight emulsifier.

[041] In another embodiment, the composition as described herein comprises from 2 to 30% by weight aqueous solution. In another embodiment, the composition as described herein comprises from 5 to 15% by weight aqueous solution. In another embodiment, the composition as described herein comprises from 8 to 12% by weight aqueous solution. In another embodiment, the composition as described herein comprises from 8 to 25% by weight aqueous solution. In another embodiment, the composition as described herein comprises from 15 to 28% by weight aqueous solution. In another embodiment, the composition as described herein comprises from 10 to 30% by weight aqueous solution.

[042] In another embodiment, the composition as described herein comprises from 2 to 50% by weight pigment. In another embodiment, the composition as described herein comprises from 5 to 30% by weight pigment. In another embodiment, the composition as described

herein comprises from 5 to 20% by weight pigment. In another embodiment, the composition as described herein comprises from 15 to 30% by weight pigment. In another embodiment, the composition as described herein comprises from 7 to 15% by weight pigment. In another embodiment, the composition as described herein comprises from 25 to 40% by weight pigment. In another embodiment, the composition as described herein comprises from 25 to 35% by weight pigment.

[043] In another embodiment, the composition as described herein comprises: from 30 to 75% by weight acrylic polymer, from 10 to 40% by weight pigment, from 0.4 to 4% by weight thickener, from 0.5 to 2.5% by weight emulsifier, from 8 to 42% by weight aqueous solution, and from 7 to 15% by weight fragrance.

[044] In another embodiment, the composition as described herein comprises: from 40 to 80% by weight acrylic polymer, from 1 to 20% by weight pigment, from 0.1 to 2% by weight thickener, from 0.5 to 2.0% by weight emulsifier, from 3 to 20% by weight aqueous solution, and from 1 to 10% by weight fragrance.

[045] In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 5 to 50 seconds after exposure to air at a temperature of 15 to 30°C. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 5 to 50 seconds after it is applied on a body surface at an ambient temperature of 15 to 30°C. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 5 to 50 seconds after it is applied onto the skin at an ambient temperature of 15 to 30°C. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 5 to 50 seconds after it is applied onto a cloth at an ambient temperature of 15 to 30°C. In another embodiment, a stabilized composition is a composition that does not stain another surface upon contact with another surface. In another embodiment,

a stabilized composition is a solid composition. In another embodiment, a stabilized composition is a composition that sticks to a body surface.

[046] In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 5 to 30 seconds. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 10 to 40 seconds. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 10 to 20 seconds. In another embodiment, the composition as described herein solidifies, dries, and/or stabilizes within 10 to 30 seconds.

[047] In another embodiment, each physical measure described herein is obtained under standard conditions for temperature and pressure (STP) or STAP if not otherwise stated.

[048] In another embodiment, a thickener of the invention has a density of 0.5 to 2.5 g/cm³ at 20°C. In another embodiment, a thickener of the invention has a density of 0.7 to 2.2 g/cm³ at 20°C. In another embodiment, a thickener of the invention has a density of 0.7 to 1.8 g/cm³ at 20°C. In another embodiment, a thickener of the invention has a density of 0.9 to 1.6 g/cm³ at 20°C.

[049] In another embodiment, an acrylic polymer has a viscosity of 1500 to 12000 mPas. In another embodiment, an acrylic polymer has a viscosity of 2500 to 10000 mPas. In another embodiment, an acrylic polymer has a viscosity of 2500 to 7500 mPas. In another embodiment, an acrylic polymer has a viscosity of 2000 to 4000 mPas. In another embodiment, an acrylic polymer has a viscosity of 3000 to 6000 mPas.

[050] In another embodiment, an acrylic polymer of the invention has a density of 0.5 to 2.5 g/cm³ at 20°C. In another embodiment, an acrylic polymer of the invention has a density of 0.7 to 2.2 g/cm³ at 20°C. In another embodiment, an acrylic polymer of the invention has a

density of 0.7 to 1.8 g/cm³ at 20°C. In another embodiment, an acrylic polymer of the invention has a density of 0.8 to 1.5 g/cm³ at 20°C.

[051] In another embodiment, an acrylic polymer has an ultimate tensile strength of 7 to 12 N/mm². In another embodiment, an acrylic polymer has an ultimate tensile strength of 8 to 12 N/mm². In another embodiment, an acrylic polymer has an ultimate tensile strength of 8 to 10 N/mm². In another embodiment, an acrylic polymer has an ultimate tensile strength of 8.5 to 9.5 N/mm².

[052] In another embodiment, an acrylic polymer has an elongation at break measure of 200 to 800%. In another embodiment, an acrylic polymer has an elongation at break measure of 300 to 800%. In another embodiment, an acrylic polymer has an elongation at break measure of 300 to 700%. In another embodiment, an acrylic polymer has an elongation at break measure of 400 to 600%. In another embodiment, an acrylic polymer has an elongation at break measure of 450 to 550%.

[053] In another embodiment, an acrylic polymer has an E-modulus of 0.5 to 5 N/mm². In another embodiment, an acrylic polymer has an E-modulus of 1 to 4 N/mm². In another embodiment, an acrylic polymer has an E-modulus of 1 to 3 N/mm². In another embodiment, an acrylic polymer has an E-modulus of 1.5 to 2.5 N/mm².

[054] In another embodiment, an acrylic polymer of the invention comprises from 30 to 75% solid content. In another embodiment, an acrylic polymer of the invention comprises from 40 to 75% solid content. In another embodiment, an acrylic polymer of the invention comprises from 40 to 60% solid content. In another embodiment, an acrylic polymer of the invention comprises from 45 to 55% solid content.

[055] In another embodiment, a pigment of the invention has a density of 1 to 10 g/cm³ at 20°C. In another embodiment, a pigment of the invention has a density of 1.5 to 5 g/cm³ at

20°C. In another embodiment, a pigment of the invention has a density of 1 to 3 g/cm³ at 20°C. In another embodiment, a pigment of the invention has a density of 2 to 4 g/cm³ at 20°C.

[056] In another embodiment, a pigment of the invention comprises aluminum. In another
5 embodiment, a pigment of the invention comprises titanium dioxide. In another embodiment, a pigment of the invention comprises silicon dioxide.

[057] In one embodiment, the fragrance-releasing complex comprises a retention
composition. In another embodiment, the fragrance-releasing complex comprises a
combination of retention compositions. In one embodiment, the fragrance-releasing complex
10 comprises a fragrance. In another embodiment, the fragrance-releasing complex comprises a
combination of fragrances. In another embodiment, the present invention provides a
composition and/or a fragrance-releasing complex comprising one or more fragrances and one
or more fragrance retention compositions. In another embodiment, the fragrance-releasing
complex comprises an admixture or an accord.

[058] In another embodiment, the fragrance retention composition comprises an acrylic
15 polymer or a combination of acrylic polymers. In another embodiment, the acrylic polymer, is
comprises starch or modified starch. In another embodiment, the acrylic polymer, is a:
homopolymer, copolymer or multipolymer of one or more monomers including, but not
limited to, acrylic acid, methacrylic acid, esters of acrylic acid and methacrylic acid (acrylates
20 and methacrylates), acrylamides, acrylonitriles and derivatives and any combinations
thereof. In another embodiment, the acrylic polymer comprises acrylate monomers, such as
but not limited to: Methacrylate, Methyl acrylate, Ethyl acrylate, 2-Chloroethyl vinyl ether, 2-
Ethylhexyl acrylate, Hydroxyethyl methacrylate, Butyl acrylate, Butyl methacrylate,
trimethylolpropane triacrylate (TMPTA) and any combination thereof. In another

embodiment, the acrylic polymer is a copolymer-dispersion of acrylic and methacrylic acid esters.

[059] In another embodiment, an acrylic polymer is an acrylate polymer or copolymer. In another embodiment, the fragrance retention composition comprises solid polymers such as PMMA (polymethyl methacrylate). In another embodiment, the fragrance retention composition comprises silica. In another embodiment, the fragrance retention composition comprises methyl methacrylate. In another embodiment, the fragrance retention composition comprises acrylated beads. In another embodiment, the fragrance retention composition comprises microparticles comprising polymers of the invention. In another embodiment, the fragrance retention composition comprises nanoparticles comprising polymers of the invention. In another embodiment, the acrylic polymer is a copolymer. In another embodiment, the acrylic polymer is a dispersion of acrylic and methacrylic acid esters.

[060] In another embodiment, the fragrance retention composition is nonporous. In another embodiment, the fragrance retention composition comprises a composite product such as but not limited to: PMMA 2MUSI. In another embodiment, the fragrance retention composition is a film-forming agent. In another embodiment, the fragrance retention composition comprises a wax, a resin, a gum, cellulose, polyquaternium and any combination thereof. In another embodiment, the fragrance retention composition comprises any cosmetic polymer and/or copolymer.

[061] In one embodiment, the fragrance retention composition has the capacity to affect the characteristics of the fragrance composition, such as but not limited to, limiting evaporation rate and intensity, in a manner that can be perceived by an observer or user thereof, over time, as compared to the same perception in the absence of the fragrance retention composition. In another embodiment, the fragrance retention composition is non-irritable, non-allergenic, non-toxic or any combination thereof to a body surface such as skin. In another embodiment, the

fragrance retention composition thickens upon exposure to air. In another embodiment, the fragrance retention composition dries upon exposure to air. In another embodiment, the fragrance retention composition solidifies upon exposure to air.

[062] In one embodiment, the present invention provides a fragrance-releasing complex, wherein the weight ratio of the fragrance to the acrylic polymer is at least 1:90. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is about 1:50. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is about 1:30. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:175 to 1:3. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:150 to 1:3. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:150 to 1:4. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:135 to 1:4. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:135 to 1:30. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:200 to 1:10. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:120 to 1:30. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:500 to 1:100. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:100 to 1:10. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:90 to 1:15. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:90 to 1:10. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:75 to 1:8. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:75 to 1:9. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:60 to 1:9. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:60 to 1:10. In another embodiment, the weight ratio of the fragrance

to the acrylic polymer is from about 1:45 to 1:10. In another embodiment, the weight ratio of the fragrance to the acrylic polymer is from about 1:45 to 1:15.

[063] In one embodiment, the acrylic polymer is within an emulsion. In another embodiment, an emulsion is oil in water emulsion. In another embodiment, an emulsion is water in oil emulsion. In another embodiment, the acrylic polymer is within dispersion. In another embodiment, the acrylic polymer is within a suspension.

[064] In one embodiment, the emulsion comprises polyethyl acrylate, polymethyl acrylate, a copolymer of ethyl acrylate, and/or methyl methacrylate. In one embodiment, the emulsion comprises an ethylenically unsaturated compound such as, but not limited to: styrene. In another embodiment, the emulsion comprises a higher alkyl methacrylate.

In one embodiment, the emulsion is an acrylic paint. In one embodiment, the term “acrylic paint” refers to water based paint. In another embodiment the term acrylic paint refers to latex paint. In another embodiment the term acrylic paint refers to emulsion paint. In another embodiment the term acrylic paint refers to water soluble paint.

[065] In one embodiment, the fragrance retention composition comprises a pigment. In another embodiment, the fragrance retention composition comprises a dispersing agent. In another embodiment, the fragrance retention composition comprises a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises an inorganic filler. In another embodiment, the fragrance retention composition comprises a binding agent. In another embodiment, the fragrance retention composition comprises a resin. In another embodiment, the fragrance retention composition comprises a rheological agent. In another embodiment, the fragrance retention composition comprises an anti-foaming agent.

[066] In one embodiment, the pigment is an organic pigment. In another embodiment, the pigment is an inorganic pigment. In one embodiment, the pigment is a combination of

pigments. In one embodiment, the pigment is a combination of inorganic and organic pigments. In another embodiment, the pigment is a colorant. In another embodiment, the pigment comprises a combination of colorants. In another embodiment, the pigment is a dye. In another embodiment, the pigment comprises a combination of dyes. In one embodiment, the pigment comprises a colored component. In another embodiment, the pigment comprises a combination of colored components. In another embodiment, the pigment is an ink. In another embodiment, the pigment comprises a combination of inks.

[067] In another embodiment, the pigment comprises: an anthraquinone, a phthalocyanine, a pyrroline, a triphenodioxazine, a methine, a benzodifuranone, a coumarin, an indoaniline, a benzenoid, a xanthene, a phenazine, a solvent soluble sulphur dye, a quinophthalone, a pyridone, an aminopyrazole, a pyrrolidine, a styrylic and any combination thereof. In another embodiment, the pigment comprises an azoic, such as, but not limited to, a monoazo, a disazo, a trisazo and any combination thereof. In another embodiment, the pigment comprises an azoic containing a heterocyclic group.

[068] In one embodiment, the fragrance retention composition comprises, about 30% to 95% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 30% to 90% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 40% to 90% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 40% to 80% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 50% to 80% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 50% to 70% by weight of a pigment. In one embodiment, the fragrance retention composition comprises, about 60% to 70% by weight of a pigment.

[069] In one embodiment, the fragrance retention composition comprises, about 0.5% to 30% by weight of a dispersing agent. In one embodiment, the fragrance retention composition

comprises, about 0.5% to 25% by weight of a dispersing agent. In one embodiment, the fragrance retention composition comprises, about 1.5% to 25% by weight of a dispersing agent. In one embodiment, the fragrance retention composition comprises, about 3% to 20% by weight of a dispersing agent. In one embodiment, the fragrance retention composition comprises, about 5% to 20% by weight of a dispersing agent. In one embodiment, the fragrance retention composition comprises, about 5% to 10% by weight of a dispersing agent.

[070] In another embodiment, the fragrance retention composition comprises, about 2% to 50% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 2% to 40% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 5% to 40% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 5% to 35% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 5% to 30% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 10% to 20% by weight of a solubilization accelerating agent. In another embodiment, the fragrance retention composition comprises, about 15% to 20% by weight of a solubilization accelerating agent.

[071] In another embodiment, the fragrance retention composition comprises, about 1% to 50% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 1% to 40% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 3% to 40% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 5% to 40% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 5% to 35% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 5% to 30% by weight of

inorganic filler. In another embodiment, the fragrance retention composition comprises, about 10% to 20% by weight of inorganic filler. In another embodiment, the fragrance retention composition comprises, about 15% to 20% by weight of inorganic filler.

[072] In another embodiment, the fragrance retention composition comprises, about 1 to 10% by weight of a binding agent. In another embodiment, the fragrance retention composition comprises, about 1 to 8% by weight of a binding agent. In another embodiment, the fragrance retention composition comprises, about 3 to 8% by weight of a binding agent. In another embodiment, the fragrance retention composition comprises, about 2 to 6% by weight of a binding agent. In another embodiment, the fragrance retention composition comprises, about 4 to 6% by weight of a binding agent.

[073] In another embodiment, the fragrance retention composition comprises, about 1% to 50% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 1% to 40% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 3% to 40% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 5% to 40% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 5% to 35% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 5% to 30% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 10% to 20% by weight of a resin. In another embodiment, the fragrance retention composition comprises, about 15% to 20% by weight of a resin.

[074] In one embodiment, the fragrance retention composition comprises, about 0.5% to 20% by weight of a rheological agent. In one embodiment, the fragrance retention composition comprises, about 0.5% to 17% by weight of a rheological agent. In one embodiment, the fragrance retention composition comprises, about 1.5% to 17% by weight of

a rheological agent. In one embodiment, the fragrance retention composition comprises, about 1.5% to 15% by weight of a rheological agent. In one embodiment, the fragrance retention composition comprises, about 3% to 15% by weight of a rheological agent. In one embodiment, the fragrance retention composition comprises, about 3% to 10% by weight of a
5 rheological agent. In one embodiment, the fragrance retention composition comprises, about 5% to 10% by weight of a rheological agent. In one embodiment, the fragrance retention composition comprises, about 5% to 7% by weight of a rheological agent.

[075] In another embodiment, the fragrance retention composition comprises, about 0.5 to 10% by weight of an anti-foaming agent. In another embodiment, the fragrance retention
10 composition comprises, about 0.5 to 8% by weight of an anti-foaming agent. In another embodiment, the fragrance retention composition comprises, about 1.5 to 8% by weight of an anti-foaming agent. In another embodiment, the fragrance retention composition comprises, about 1.5 to 6% by weight of an anti-foaming agent. In another embodiment, the fragrance retention composition comprises, about 3 to 6% by weight of an anti-foaming agent. In
15 another embodiment, the fragrance retention composition comprises, about 3 to 4.5% by weight of an anti-foaming agent.

[076] In one embodiment, the fragrance comprises: an alcohol, an ether, a nitrile, an aldehyde, an ester, a ketone, a lactone, a thiol, an amine, a schiff-base, a terpene, a cyclic alkene, a cyclic oxide, an oxime, an essential oil, an aromatic species, or any combination
20 thereof.

[077] In another embodiment the fragrance comprises a natural product, such as, but not limited to, an absolute, a resinoid, a resin and a concrete and any combination thereof.

[078] In another embodiment the fragrance comprises a saturated compound, an unsaturated compound, an aliphatic compound, a carbocyclic compound and a heterocyclic compound and any combination thereof.

[079] In another embodiment the fragrance comprises a component that is selected from the following: alicyclic ester, aliphatic ester, cyclic ester, aromatic ester, alicyclic ether, cyclic ether, aromatic ether, primary alcohols, tertiary alcohols, aromatic alcohols, cyclic alcohols, aliphatic aldehyde, cyclic aldehydes, aliphatic primary aldehyde, aromatic aldehydes, tertiary aldehydes, aliphatic ketones, aromatic ketones, cyclic ketones, macrocyclic ketones, aliphatic nitrile, aromatic nitrile, cyclic nitrile, 1,2-benzopyrone, Methyl 2-((1-(2,4-dimethyl-3-cyclohexenyl)methylidene)amino)-1-benzenecarboxylate, Methyl 2-((2-methylpentylidene)amino)-1-benzenecarboxylate, dimethyl-2-Methylene bicyclo (3,1,1) Heptane (6,6-), para-mentha-1,4(8)-diene, and any combination thereof.

[080] In another embodiment the fragrance comprises a component that is selected from the following: an ethyl 2,6,6-trimethyl-1,3-cyclohexadiene-1-carboxylate, cis-beta-gamma-hexenyl acetate, 2-methylbuten-2-ol-4-acetate; glycolic acid, 2-pentyloxy:allyl ester, methyl-2-nonenoate, cis-beta-gamma-Hexenyl salicylate, 4-methyl-pentan-2-ol 2-butenate, hexyl 2-butenate, acetyl diisoamylene, 3-methylene-7-methyl-1-octen-7-yl acetate, methyl 2-nonenoate, citronellyl acetate, isobutyl angelate, tricyclo decenyl acetate, tricyclo decenyl propionate, 7-Acetyl, 1,2,3,4,5,6,7,8-octahydro-1,1,6,7-tetramethyl naphthalene, givescone, allyl cyclohexyloxy acetate, 4(3)-(4-Methyl-3-pentenyl)-3-cyclohexenyl-methyl acetate, 2-cyclopentyl-cyclopentyl 2-butenate, datilat, 6-Butyl-3,6-dihydro-2,4-dimethyl-2-hydro-pyran, 3,6-dihydro-4,6-dimethyl-2-phenyl-2hydro-pyran, 9-(Methyloxy)tricydo[5,2,1,0^{2,6}]dec-3-ene, beta gamma hexenol, 3,7-dimethyl-6-octen-1-ol, cis-3-hexenol, 9-Decen-1-ol, geraniol, 2,6-dimethyl-7-octen-2-ol, linalool, dihydro myrcenol, 3,7-Dimethyl-1,6-octadien-3-ol, 4-Methyl-3-decen-5-ol, amyl vinyl carbinol, 3-Phenyl-2-

propen-1-ol, 2-Methyl-4-phenyl-1-pentanol, 2-Ethyl-4-(2,2,3-trimethyl-3-cyclopenten-1-yl)-
 2-buten-1-ol, 4,7-Methano-1H-inden-5-ol, 3a,4,5,6,7,7a-Hexahydro-2(or 3), 4-dimethyl,
 dimethyl cyclohexene methanol, 5-(2,2,3-Trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol, 1-
 Para-Menthen-4-ol, terpineol, ambrinol L20, 3,7-Dimethyl-6-octen-1-al, 2-Methyl-4(2,6,6-
 5 trimethyl-1-cyclohexenyl)-2-butenal, 1 -Methyl-4-(4-methylpentyl)-3-
 cyclohexenecarbaldehyde, 2,7-Dimethyloct-5-en-4-one 50% in Iso Propyl Myristate,
 Hexahydro Tetramethyl Methanonaphtalenone, 4-Penten-1-one, 1-(5,5-dimethyl-1-
 cyclohexen-1-yl)-, 5,8-Methano-2H-1-benzopyran; 6-ethylideneoctahydro-, 2-Buten-1-one; 1-
 (2,6,6-trimethyl-2-cyclohexen-1-yl)-, Diethyl dimethylcyclohex-2-en-1-one, 2-Buten-1-one,
 10 1-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, 2,5,10-Trimethyl-2,5,9-cyclododecatrien-1-yl methyl
 ketone, Methyl cedr-8-enyl ketone, Oxacyclohexadecen-2-one, 3,7-dimethyl-2,6-
 octadienenitrile, 3,12-tridecadiene nitrile, iris nitrile, ozonil, teamonyl, 5-Phenyl-3-methyl-
 penten-2-acid-nitrile, 2-Benzyl-2-methyl-3-butenenitril, rose nitrile, 1,2-benzopyrone, Methyl
 2-((-1-(2, 4-dimethyl-3-cyclohexenyl)methylidene)amino)-1-benzenecarboxylate, Methyl 2-
 15 ((-2-methylpentylidene)amino)-1-benzenecarboxylate, Dimethyl-2-Methylene Bicyclo(3,1,1)
 Heptane (6,6-); Para-Mentha-1 ,4(8)-diene and any combination thereof.

[081] In one embodiment, the fragrance comprises a base note fragrance. In one
 embodiment, the fragrance comprises a middle note fragrance. In one embodiment, the
 fragrance comprises a top note fragrance. In one embodiment, the fragrance comprises a
 20 balance carrier. In one embodiment, the term "balance carrier" is synonymous with the term
 "carrier" as used herein. In one embodiment, the fragrance comprises a carrier. In one
 embodiment, the fragrance comprises a fixative. In one embodiment, the term "fragrance" is
 synonymous with perfume. In another embodiment, the term "fragrance" means scent. In
 another embodiment, the term "fragrance" means aroma. In one embodiment, the term
 25 "fragrance", includes but is not limited to, conventional fragrances known in the art, including

but not limited to, U.S. Pat. Nos.: 4,145,184, Brain and Cummins, issued Mar. 20, 1979; 4,209,417, Whyte, issued June 24, 1980; 4,515,705, Moeddel, issued May 7, 1985; and 4,152,272, Young, issued May 1, 1979, all of the patents being incorporated herein by reference.

5 [082] In one embodiment, the top note fragrance, the middle note fragrance and the base note fragrance are scents that can be sensed with respect to the time after the application of a perfume.

[083] In one embodiment, the top note fragrance has an evaporation coefficient (perfume notes) in the range of about 1 to 14. In one embodiment, the middle note fragrance has an
10 evaporation coefficient (perfume notes) in the range of about 15 to 60. In one embodiment, the base note fragrance has an evaporation coefficient (perfume notes) in the range of about 61 to 100.

[084] In one embodiment, the carrier comprises a component selected from the following: water, C1-C4 monohydric alcohol, C2-C6 polyhydric alcohol, propylene carbonate, liquid
15 polyalkylene glycol and any combinations thereof.

[085] In one embodiment, the fixative comprises a component selected from the following: Ambergris, sandalwood, musk, vetiver, orris root, and bergamot orange and any combinations thereof.

[086] In one embodiment, the fragrance is selected according to the
20 fragrance odor characteristics and other fragrance characteristics such as, but not limited to: viscosity, specific gravity, refractive index, saponification value and flash point or any combination thereof. In another embodiment the fragrance is selected according to fragrance odor characteristics obtained when combined with other components. In one embodiment, fragrance odor characteristics include a pleasant odor and/or clean odor such as, but not

limited to: lavender, violet, rose, jasmin, pine, woody, floral, fruity, lemon, lime, apple, peach, raspberry, strawberry, banana, plum, apricot, vanilla, pear, eucalyptus, aromatic, aldehydic, tutti frutti, oriental, sweet, amber, Paola, Muguet, Citron (lime) ella and any combination thereof. In another embodiment the fragrance is selected according to fragrance
5 characteristics obtained when combined with other components.

[087] In another embodiment, fragrance odor characteristics and other fragrance characteristics include, but are not limited to, those given in "Perfume and Flavor Chemicals (Aroma Chemicals)," Steffen Arctander, published by the author, 1969. incorporated herein by reference.

10 [088] In another embodiment the fragrance comprises a component that is selected from the following: anethole, benzaldehyde, decyl aldehyde, benzyl acetate, benzyl alcohol, benzyl formate, benzyl propionate, iso-bornyl acetate, camphene, cis-citral (neral), citronellal, citronellol, citronellyl acetate, paracymene, decanal, dihydrolinalool, dihydromyrcenol, methyl benzyl carbinyl acetate, dimethyl benzyl carbinyl acetate, dimethyl phenyl carbinol,
15 eucalyptol, helional, geranial, geraniol, geranyl acetate, geranyl nitrile, cis-3-hexenyl acetate, dihydrocitronellal, d-limonene, linalool, linalool oxide, tetra-hydro linalool, alpha-methyl ionone, methyl nonyl acetaldehyde, methyl phenyl carbinyl acetate, laevo-menthyl acetate, menthone, iso-menthone, myrcene, myrcenyl acetate, myrcenol, nerol, neryl acetate, nonyl acetate, phenyl ethyl alcohol, phenyl acetaldehyde, alpha-pinene, beta-pinene, gamma-
20 terpinene, terpineol, alpha-terpineol, beta-terpineol, terpinyl acetate, vertenex (para-tertiary-butyl cyclohexyl acetate), gamma-methyl ionone, undecalactone, undecylenic aldehyde, alpha-damascone, beta-damascone, amyl acetate, lemon oil, orange oil and any combination thereof.

[089] In another embodiment the fragrance comprises a component that is selected from the
25 following: hexyl cinnamic aldehyde, alpha-amylcinnamic aldehyde, p-anis aldehyde,

cinnamic aldehyde, cuminic aldehyde, p-t-butyl-alpha-methyldihydrocinnamaldehyde, 4-hydroxy-3-methoxycinnamaldehyde, 2-phenyl-3-(2-furyl)prop-2-enal, vanillin isobutyrate, ethyl vanillin acetate, vanillin acetate, cyclamen aldehyde, heptanal, lauryl aldehyde, nonanal, octanal, phenyl propyl aldehyde, vanillin, salicyl aldehyde, cytral, 2,4-dihydroxy-3-methylbenzaldehyde, 2-hydroxy-4-methylbenzaldehyde, 5-methyl salicylic aldehydes, 4-nitrobenzaldehyde, o-nitrobenzaldehyde, 5-ethyl-2-thiophenecarbaldehyde, 5-methyl-2-thiophenecarboxaldehyde, 2-thiophenecarb aldehyde, as aronaldehyde, 5-(hydroxymethyl)-2-furaldehyde, 2-benzofurancarboxaldehyde, 2-benzofurancarboxaldehyde, 4-ethoxy-3-methoxy benz aldehyde, Protocatechualdehyde, Heliotropine, 2,3,4-trimethoxybenz aldehyde, 3,4,5-trimethoxybenzaldehyde, 2,8-dithianon-4-carboxaldehyde, Sorbinaldehyde, 2,4-heptadienal, 2,4-decadienal, 2,4-nonadienal, (E,E)-2,4-octadien-1-al, 2,4-octadienal, 2,4-dodecadienal, 4-undecadienal, 2,4-tridecadien-1-al, 2-trans-4-cis-7-cis-tridecatrienal, piperonylidene propionaldehyde, 2-methyl-3-(2-furyl)acrolein, 2,4-pentadienal, 2-furfurylidene butyraldehyde, 3-(2-furyl)acrolein, Pyruvaldehyde, Ethanediol, menthol, 3-buten-2-one, 3-methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl), 4-(2,6,6-trimethylcyclohexen-1-en-1-yl)but-3-en-2-one, 3-buten-2-one, 4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-, (E)-, menthyl lactate, isomenthyl acetate, isomenthyl propionate, isomenthyl isobutyrate, camphor, p-menthane, cresol, tetrahydromyrcenol, citronellol, citronellyl derivatives, geranyl derivatives, linalyl acetate, mugetanol, eugenol, jasmal, pinanol, cedrene, beta pinene, cineole, nonadienol, ethylhexanal, octanol acetate, methyl furfural, terpinene, thujene, amylacetate, benzylacetate, dihydrocumarin, 5 dihydromyrcenyl acetate, Isoamylacetate, para-cymene, triethyl acetate, para-cresol, ethyl acetate, benzyl-benzoate, isopropyl myristate, methyl abietate, Ethanol, Isopropanol, diethyl sebacate, Glycerol, propylene glycol, 1,2-butylene glycol, dipropylene glycol, 2-methyl-2,4-pentanediol, diethylene glycol monoethyl ether, diethyl phthalate, hexyl salicylate, triethyl citrate, benzyl salicylate and any

combination thereof. In another embodiment, the fragrance comprises the odor characteristics of the acrylic polymer.

[090] In another embodiment, the fragrance includes, but is not limited to those disclosed in "Perfume and Flavor Chemicals (Aroma Chemicals)," Steffen Arctander, published by the author, 1969. incorporated herein by reference.

[091] In one embodiment, the fragrance releasing complex comprises a solvent suitable for human topical administration. In another embodiment the solvent comprises a component selected from the following: dipropylene glycol, diethyl phthalate, isopropyl myristate, benzyl benzoate, 2-(2-ethoxyethoxy)-1-ethanol or ethyl citrate, propylene glycol, triacetin, triethyl citrate, benzyl alcohol, ethanol, vegetable oil, terpene and any combination thereof.

[092] In one embodiment, the fragrance releasing complex comprises a volatile fragrance.

[093] In one embodiment, a volatile fragrance has a boiling point of less than about 475°C to 525°C. In another embodiment a volatile fragrance has a boiling point of less than about 475°C to 515°C. In another embodiment a volatile fragrance has a boiling point of less than about 485°C to 515°C. In another embodiment a volatile fragrance has a boiling point of less than about 495°C to 505°C. In another embodiment a volatile fragrance has a boiling point of less than about 500°C.

[094] In one embodiment, a volatile fragrance is a highly volatile fragrance. In one embodiment, highly volatile fragrance means having a boiling point of about 230°C to 250°C or lower. In another embodiment highly volatile fragrance means having a boiling point of about 230°C to 245°C or lower. In another embodiment highly volatile fragrance means having a boiling point of about 235°C to 245°C or lower. In another embodiment highly volatile fragrance means having a boiling point of about 250°C or lower.

[095] In another embodiment the highly volatile fragrance comprises a component selected from the following: anethole, benzaldehyde, benzyl acetate, benzyl alcohol, benzyl formate, iso-bornyl acetate, camphene, cis-citral (neral), citronellal, citronellol, citronellyl acetate, para-cymene, decanal, dihydrolinalool, dihydromyrcenol, dimethyl phenyl carbinol, 5 eucalyptol, geranial, geraniol, geranyl acetate, geranyl nitrile, cis-3-hexenyl acetate, hydroxycitronellal, d-limonene, linalool, linalool oxide, linalyl acetate, linalyl propionate, methyl anthranilate, alpha-methyl ionone, methyl nonyl acetaldehyde, methyl phenyl carbinyl acetate, laevo-menthyl acetate, menthone, iso-menthone, myrcene, myrcenyl acetate, myrcenol, nerol, neryl acetate, nonyl acetate, phenyl ethyl alcohol, alpha-pinene, beta-pinene, 10 gamma-terpinene, alpha-terpineol, beta-terpineol, terpinyl acetate, and vertenex (para-tertiary-butyl cyclohexyl acetate) and any combination thereof.

[096] In one embodiment, a volatile fragrance is a moderately volatile fragrance. In one embodiment, a moderately volatile fragrance means having a boiling point of about 250°C to 300°C. In another embodiment a moderately volatile fragrance means having a boiling point 15 of about 250°C to 300°C. In another embodiment a moderately volatile fragrance means having a boiling point of about 250°C to 290°C. In another embodiment a moderately volatile fragrance means having a boiling point of about 260°C to 290°C. In another embodiment a moderately volatile fragrance means having a boiling point of about 260°C to 280°C.

[097] In another embodiment, the moderately volatile fragrance comprises a component 20 selected from the following: amyl cinnamic aldehyde, iso-amyl salicylate, beta-caryophyllene, cedrene, cinnamic alcohol, coumarin, dimethyl benzyl carbinyl acetate, ethyl vanillin, eugenol, iso-eugenol, flor acetate, heliotropine, 3-cis-hexenyl salicylate, hexyl salicylate, lilial (para- tertiarybutyl-alpha-methyl hydrocinnamic aldehyde), gamma-methyl ionone, nerolidol, patchouli alcohol, phenyl hexanol, beta-selinene, trichloromethyl phenyl carbinyl acetate, 25 triethyl citrate, vanillin, veratraldehyde and any combination thereof.

[098] In one embodiment, a volatile fragrance is a less volatile fragrance. In one embodiment, a less volatile fragrance means having a boiling point of about 300°C to 500°C. In another embodiment a less volatile fragrance means having a boiling point of about 300°C to 450°C. In another embodiment a less volatile fragrance means having a boiling point of about 350°C to 450°C.

[099] In another embodiment, the less volatile fragrance comprises a component selected from the following: benzophenone, benzyl salicylate, ethylene brassiate, galaxolide (1,3,4,6,7,8-hexahydro-4,6,6,7,8,8-hexamethyl-cyclopenta-gama-2-benzopyran), hexyl cinnamic aldehyde, lylal (4-(4-hydroxy-4-methyl pentyl)-3-cyclohexene-10-carboxaldehyde), methyl cedrylone, methyl dihydro jasmonate, methyl-beta-naphthyl ketone, musk indanone, musk ketone, musk tibetene, phenylethyl phenyl acetate and any combination thereof.

[0100] In one embodiment, the volatile fragrance is encapsulated or entrapped within the acrylic polymer. In one embodiment, the volatile fragrance is partially encapsulated within the acrylic polymer. In another embodiment the volatile fragrance is fully encapsulated within the acrylic polymer. In one embodiment, the volatile fragrance is partially entrapped within the acrylic polymer. In another embodiment the volatile fragrance is fully entrapped within the acrylic polymer.

[0101] In another embodiment, “retains a perceptible fragrance” is retains a perceptible fragrance after applied to a body surface. In another embodiment, “retains a perceptible fragrance” is smellable or detected by smell from a distance of 5 cm as described herein. In one embodiment, the fragrance-releasing complex retains a perceptible fragrance for at least 7 days. In another embodiment, the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 24 hours to 18 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex

for 24 hours to 16 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 3 to 16 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 3 to 14 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 5 to 14 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 5 to 12 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 7 to 10 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 7 days. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 10 days.

[0102] In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 10 to 50 hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 10 to 40 hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 15 to 50 hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 10 to 30

hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 15 to 30 hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 18 to 28 hours. In another embodiment, the composition and/or the fragrance-releasing complex retains a perceptible fragrance attributable to the composition and/or the fragrance-releasing complex for 20 to 30 hours.

[0103] In one embodiment, the present invention provides a method for conferring, enhancing, improving or modifying the odor properties of a body surface, contacting or treating the body surface with the composition and/or the fragrance-releasing complex as described herein.

[0104] In one embodiment, the composition comprises an alcohol such as ethanol. In another embodiment the composition comprises a perfume concentrate. In another embodiment the composition comprises an eaux de parfum. In another embodiment, the composition comprises an eaux de toilette. In another embodiment the composition comprises a cologne. In another embodiment, the composition comprises a body splash.

[0105] In one embodiment, a fragrance means, a compound of current use in perfumery, which is used essentially for its ability to smell pleasantly and to be capable of imparting a pleasant odor to the products into which it is incorporated, or to the surfaces, such as, but not limited to, skin or hair, to which it is applied, on its own or in admixture with other components.

[0106] In another embodiment a fragrance is a composition capable to impart or modify the odor of a surface such as a body surface.

[0107] In another embodiment a fragrance is a composition that has the ability to mask a malodor of a surface.

[0108] In one embodiment, the present invention provides a method for decorating a body surface. In another embodiment, the body surface is skin. In one embodiment, the present invention provides a kit for applying a fragrance to a subject, comprising a fragrance-releasing complex; and means for topically applying the composition and/or the fragrance-releasing complex to a body surface of a subject.

[0109] In one embodiment, the kit comprises a stencil having a decorative shape therein. In another embodiment, the kit comprises a transferable sheet with a decorative shape thereon. In another embodiment, the kit comprises a cutout in the shape of the body decoration. In another embodiment, the kit comprises an image or pattern to be placed on any part of the human body.

[0110] In another embodiment, the kit comprises decorative element comprising a substrate, a decorative pattern thereon. In another embodiment, the substrate comprises a plastic film. In another embodiment, the plastic film comprises a polymer selected from the following: polyethylene, polypropylene, polyester, polyvinyl chloride and plasticized polyvinyl chloride; a metalized plastic film; and a metal foil.

[0111] In another embodiment, the plastic film has a thickness of about 0.01 mm to 0.5 mm. In another embodiment, the plastic film has a thickness of about 0.01 mm to 0.45 mm. In another embodiment, the plastic film has a thickness of about 0.05 mm to 0.45 mm. In another embodiment, the plastic film has a thickness of about 0.05 mm to 0.4 mm. In another embodiment, the plastic film has a thickness of about 0.1 mm to 0.4 mm. In another embodiment, the plastic film has a thickness of about 0.1 mm to 0.35 mm. In another embodiment, the plastic film has a thickness of about 0.15 mm to 0.35 mm. In another

embodiment, the plastic film has a thickness of about 0.15 mm to 0.3 mm. In another embodiment, the plastic film has a thickness of about 0.2 mm to 0.25 mm.

[0112] In another embodiment, the kit comprises an adhesive sheet. In another embodiment, the adhesive sheet comprises an adhesive component selected from the following: acrylic emulsion, a solvent-based acrylic and an acrylic hot melt adhesive.

[0113] In another embodiment, the kit comprises a decorative template that covers one or more regions of the body area. In another embodiment, the decorative template limits application of the composition and/or the fragrance-releasing complex to those regions of the body area not covered by the decorative template. In another embodiment, the body area is selected from the group comprising skin, nails, teeth, and any combination thereof. In another embodiment, the kit is for home use. In one embodiment, the present invention provides a kit for decorating a body surface with a composition or a complex as described herein. In one embodiment, the present invention provides a kit wherein the composition and/or the fragrance-releasing complex further comprise a pigment.

EXAMPLES

[0114] The experiments conducted with the below formulations/compositions involved applying 6 ml of each composition to the arm of an adult female. The fragrance duration experiments involved applying 6 ml of the below compositions to a designated glass surface.

[0115] All fragrance related tests were performed by one of the inventors who smelled the compositions in a close room at 25°C by placing her nose 5 cm from the composition.

[0116] Fragrance duration tests were performed every 4 hours until 20 hours and every hour from 20 hours to 30 hours.

[0117] The intensity of immediate Fragrance release was measured 1 minute after the composition was applied to an adult skin. The fragrance intensity was given a score from 1 to 10, wherein 1 is minimal fragrance and 10 is maximal fragrance.

[0118] The stability of the compositions was examined 30 minutes after each composition was applied to an adult female in a close room at 25°C.

[0119] The perfume used was a mixture of natural essential oils with aromatic chemicals. The perfume was insoluble in water, and soluble in alcohol. The perfume's refractive index is 1.4830 to 1.4970 and its gravity is 0.920 to 1.060.

[0120] The emulsifier used for these experiments was POLYSORBATE 80 (POE 20 sorbitan monoleate, SABO).

[0121] The pigments used were: 1. IRID-100, Mica coated with metallic oxides/tin dioxides, (manufactured by Pritty); 2. Aluminum Powder; 3. Bronze Powder; 4. Copper Powder; and mixtures of 2-4. Pigments 2-4 were purchased from ECKART GmbH. Additional pigments No effect to the pigment type was recorded.

15

[0122] Several organic polymers including modified starch polymers, calcium carbonate containing polymers, cellulose and its derivatives polymers, and mixture polymers were tested with the ingredients of the compositions of the invention. All the above tested polymers cause dispersion insufficiencies, instability, fragrance inhibition properties, skin incompatibility and other deleterious effects.

[0123] The effective polymer that was utilized is ALBERDINGK® AC 548 (manufactured by Alberdingk Boley GmbH). The polymer is aqueous, fine disperse, medium viscous, APEO-free copolymer-dispersion of acrylic and methacrylic acid esters and without plasticizers. It includes from 49-51% solids and has a pH value of 7.5-8.5. It has viscosity of

3000 – 6000 mPas, density of 1.07 g/cm³, ultimate tensile strength of 9 N/mm², elongation at break of 500%, and E-Modulus of 2 N/mm².

EXAMPLES

Example 1: 12 selected formulations of the invention

5 [0124] 12 different formulas of the invention were applied to the skin around the arm of an adult user. The water based copolymer-dispersion of acrylic and methacrylic acid esters was previously selected as it provided short drying/stabilization period and sufficient elasticity. Likewise, a clay thickener was chosen as it stabilized the emulsion, optimized its rheological behavior, its homogeneity, enhanced drying and increased elasticity.

10 [0125] 3 different clay thickeners were tested and the most dense clay thickener proved to provide superior in-vivo results. As can be clearly viewed from table 1, the concentration of the 603 thickener has direct impact on drying time.

[0126] A drying period of 15 to 30 seconds is optimal. Therefore the formulations having an optimal drying period were further developed.

15 [0127] Interestingly, the smell and the intensity of the fragrance did not change within the various tested formulation and seemed to be solely dependent on the perfume used and the concentration.

[0128] The results provided in table 1 clearly indicate that lack of a thickener and/or insufficient polymer concentration result in a peeling effect of the composition within up to 5
20 minutes. The thickener influenced the quality of film on skin and drying time.

[0129] As stated above clay based thickeners were previously selected due to their superior stability, and rheological properties, also, size of particles, the morphology, diffusion in solution.

[0130] Thickener 600 is CLiQFLOW 600 has solids content of ca. 100 %, specific gravity of: ca. 2.5, and particle size of: 98% < 53 microns (CLiQ SwissTech (Deutschland) GmbH).

[0131] Thickener 602 is CLiQFLOW 602 has active content of : 100 % and density (20°C) of: 1.0 g/cm³ (CLiQ SwissTech (Deutschland) GmbH).

5 [0132] Thickener 603 is CLiQFLOW 603 CLiQFLOW 603 has active content of: 100 % and density (20°C) of: 1.6 g/cm³ (CLiQ SwissTech (Deutschland) GmbH). The 603 includes bentonite clay and is modified with cellulose. Thus rheological properties, size of particles, the diffusion within solution are important factors for a successful product of the invention. thickener having a density of at least 1.0 g/cm³ was required to maintain the integrity of the present composition after it was applied to the skin. It should be noted that only the combination of sufficient amount of polymer together with a clay dense thickener result in sufficient, immediate fragrance release, and skin stability upon drying.

10

[0133] "CF 603 "

Table 1

	Formula #	Type of thickener	The intensity of immediate Fragrance release (1-10)	Quality of the film 30 minutes after it was applied to the skin	Film formation
By weight percent	1				
40%	Polymer		7	Few short cracks	yes
30%	Pigment				
2%	Thickener	603			
1%	Emulsifier				
20%	Water				
7%	Fragrance				
	2				
50%	Polymer		9	Very few short cracks	yes
20%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
20%	Water				
8%	Fragrance				

	3				
63%	Polymer		7	No Cracks	yes
10%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
20%	Water				
5%	Fragrance				
	4				
71%	Polymer		7	No Cracks	yes
10%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
10%	Water				
8%	Fragrance				
	5				
60%	Polymer		8	Small cracks	yes
20%	Pigment				
0%	Thickener				
2%	Emulsifier				
10%	Water				
8%	Fragrance				
	6				
60%	Polymer		8	Small cracks	yes
15%	Pigment				
0%	Thickener				
1%	Emulsifier				
15%	Water				
8%	Fragrance				
	7				
60%	Polymer		9	No Cracks	yes
20%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	8				
30%	Polymer		10	No film	no
30%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
8%	Water				
30%	Fragrance				
	9				
55%	Polymer		8	Many cracks and peelings	yes
30%	Pigment				
5%	Thickener	602			

1%	Emulsifier				
8%	Water				
10%	Fragrance				
	10				
55%	Polymer		8	Many cracks	yes
30%	Pigment				
2%	Thickener	602			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	11				
55%	Polymer		9	Many cracks and peelings	yes
30%	Pigment				
1%	Thickener	600			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	12				
20%	Polymer		8	Many cracks and peelings	yes
30%	Pigment				
1%	Thickener	600			
1%	Emulsifier				
40%	Water				
8%	Fragrance				

[0134] In summary, it seems that the actual presence of a thickener and particularly thickener 603 having an elevated density of 1.6 g/cm³ and at least 50% polymer confer proper stability together with fragrance retention and release as evidenced in formulations: 3-6 and 7 which were chosen for additional testing and assessments. Interestingly insufficient amount of polymer inhibited film formation.

Example 2: Formulation drying time and stability

[0135] Obviously optimal drying time is crucial to the success of a formulation of the invention. A very short drying time such as 15 seconds and less may render a kit of the

invention unusable as the formulation will dry within the applicator and prior to applying it to the skin. A long drying time will render a composition of the invention as a stain hazard.

[0136] Likewise, it is essential that a stable smell or fragrance will remain during at least 12 hours from applying the formulation to a body surface.

- 5 [0137] Table 2 summarizes the impact of the amount and type of thickener and the amount of polymer on the drying time and stability (fragrance release duration and physical integrity) of the 12 formulations presented in table 2.

Table 2

By weight percent	Formula #	Type of thickener	Duration of continuous Fragrance release (hours)	Film Quality of film	Reaching film skin stability from the time of application (seconds) at room temperature
	1				
40%	Polymer		23	Few cracks	15
30%	Pigment				
2%	Thickener	603			
1%	Emulsifier				
20%	Water				
7%	Fragrance				
	2				
50%	Polymer		24	Minor peeling	19
20%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
20%	Water				
8%	Fragrance				
	3				
63%	Polymer		20	Stable	19
10%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
20%	Water				
5%	Fragrance				
	4				
71%	Polymer		25	Stable	16

10%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
10%	Water				
8%	Fragrance				
	5				
60%	Polymer		24	Single crack	40
20%	Pigment				
0%	Thickener	603			
2%	Emulsifier				
10%	Water				
8%	Fragrance				
	6				
60%	Polymer		24	Single crack	38
15%	Pigment				
0%	Thickener	603			
1%	Emulsifier				
15%	Water				
8%	Fragrance				
	7				
60%	Polymer		30	Stable	20
20%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	8				
30%	Polymer		34	Many cracks and peelings	30
30%	Pigment				
1%	Thickener	603			
1%	Emulsifier				
8%	Water				
30%	Fragrance				
	9				
55%	Polymer		30	Many cracks and peelings	25
30%	Pigment				
5%	Thickener	602			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	10				
55%	Polymer		30	Many cracks and peelings	35
30%	Pigment				
2%	Thickener	602			

1%	Emulsifier				
8%	Water				
10%	Fragrance				
	11				
55%	Polymer		30	Many cracks and peelings	30
30%	Pigment				
1%	Thickener	600			
1%	Emulsifier				
8%	Water				
10%	Fragrance				
	12				
20%	Polymer		24	Many cracks and peelings	40
30%	Pigment				
1%	Thickener	600			
1%	Emulsifier				
40%	Water				
8%	Fragrance				

[0138] Interestingly, table 2 unexpectedly further demonstrates that a stable composition should include at least 50% and even 60% polymer and a clay based thickener having a high density as provided in example 1. With respect to the duration of continuous fragrance release (hours), all formulation were effective in maintain a solid fragrance for at least 15 hours as required. Interestingly, the amount of fragrance or water did not directly correlate with: (a) the duration of fragrance release and/or the intensity of immediate fragrance release; and (b) the physical stability of the dried composition – the skin decoration.. Thus, the 603 thickener, at least 5% fragrance, and over 50% polymer render the present composition effective in providing a novel smell releasing body decoration system.

CLAIMS**What is claimed is:**

1. A composition comprising: a fragrance; an acrylic polymer, a thickener, an emulsifier, an aqueous solution, and a pigment.
2. The composition of claim 1, wherein said composition is a visible fragrance-releasing
5 complex.
3. The composition of claim 1, wherein a noticeable amount of fragrance is released for at least 15 hours.
4. The composition of claim 1, comprising from 50 to 80% by weight acrylic polymer.
5. The composition of claim 1, wherein said fragrance is in an amount of from ~~5 to 30%~~
10 1%-15% by weight.
6. The composition of claim 1, wherein said thickener is in an amount of from 0.2 to 2%
by weight.
7. The composition of claim 1, wherein said emulsifier is in an amount of from 0.2 to 2%
by weight.
- 15 8. The composition of claim 1, wherein said aqueous solution is in an amount of from 2
to 20% by weight.
9. The composition of claim 1, wherein said pigment is in an amount of from 5 to 40%
by weight.
10. The composition of claim 1, wherein said aqueous solution is water.
- 20 11. The composition of claim 1, wherein said emulsifier is a nonionic emulsifier.
12. The composition of claim 1, wherein said acrylic polymer is within an emulsion.

13. The composition of claim 1, wherein said fragrance comprises: an alcohol, an ether, a nitrile, an aldehyde, an ester, a ketone, a lactone, a thiol, an amine, a schiff-base, a terpene, a cyclic alkene, a cyclic oxide, an oxime, an essential oil, an aromatic species, or any combination thereof.
- 5 14. The composition of claim 1, wherein said fragrance comprises: i) a base note fragrance; ii) a middle note fragrance; iii) a top note fragrance; iv) a balance carrier; v) a fixative, or any combination thereof.
15. The composition of claim 1, wherein said fragrance comprises a solvent suitable for human topical administration.
- 10 16. The composition of claim 1, wherein said fragrance comprises a volatile fragrance.
17. The composition of claim 15, wherein said ~~volatile~~ fragrance is encapsulated or entrapped within said acrylic polymer.
18. The composition of claim 1, retaining a perceptible fragrance for at least seven days.
- 15 19. The composition of claim 1, wherein said thickener has a density of 0.7 to 2 g/cm³ at 20°C.
20. The composition of claim 1, wherein said acrylic polymer has a viscosity of 1500 to 12000 mPas.
21. The composition of claim 1, wherein said acrylic polymer has a density of 0.5 to 2
20 g/cm³ at 20°C.
22. A method for conferring, enhancing, improving or modifying the odor properties of a body surface, comprising contacting or treating said body surface with said composition of any one of claims 1 to 17.
23. The method of claim 22, further comprising decorating said body surface.

24. A kit for applying a fragrance to a subject, comprising the composition of any one of claims 1 to 17; and means for topically applying said composition to a body surface of said subject.
- 5 25. The kit of claim 24, wherein said applying said composition to said subject is further applying a body decoration.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2015/050088

A. CLASSIFICATION OF SUBJECT MATTER

See extra sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC (2015.01) A61Q 13/00, A61K 8/11, A61K 8/06, A61K 8/81, A61K 8/73, A61K 8/19, A61K 8/25, A61K 8/29, A61K 8/26, A61Q 15/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

fragrance, acrylic polymer, thickner, emulsifier, pigment, extended, release, coated

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Databases consulted: PATENTSCOPE, USPTO, THOMSON INNOVATION, Eponline, Esp@cenet, Google Patents, Google Scholar, PatBase

Search terms used: fragrance, acrylic polymer, thickner, emulsifier, pigment, extended, release, coated

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 8921303 B1 LULL MICHAEL et al 30 Dec 2014 (2014/12/30) abstract, column 11 line 31-36, column 13 line 26-30	1-3
X	abstract, column 11 line 31-36, column 13 line 26-30	4-25
Y	WO 2005070371 A1 KULKARNI RUPALI et al 04 Aug 2005 (2005/08/04) page 1 line 30-34)	1-25
Y	US 6143353 A1 OSHLACK BENJAMIN et al 07 Nov 2000 (2000/11/07) abstract	1-25
Y	US 2006287205 A1 POPPLEWELL LEWIS et al 06 Apr 2006 (2006/04/06) abstract, claim 20	1-25

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

05 May 2015

Date of mailing of the international search report

06 May 2015

Name and mailing address of the ISA:

Israel Patent Office

Technology Park, Bldg.5, Malcha, Jerusalem, 9695101, Israel

Facsimile No. 972-2-5651616

Authorized officer

SHERAN Tally

Telephone No. 972-2-5651730

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No. PCT/IL2015/050088
--

Patent document cited search report	Publication date	Patent family member(s)	Publication Date
US 8921303 B1	30 Dec 2014	US 8921303 B1	30 Dec 2014
WO 2005070371 A1	04 Aug 2005	NONE	
US 6143353 A1	07 Nov 2000	NONE	
US 2006287205 A1	06 Apr 2006	US 2006287205 A1	21 Dec 2006
		US 7294612 B2	13 Nov 2007
		BR 0303858 A	08 Sep 2004
		CN 1497009 A	19 May 2004
		CN 1316002 C	16 May 2007
		DE 60325153 D1	22 Jan 2009
		EP 1407754 A1	14 Apr 2004
		EP 1407754 B1	10 Dec 2008
		EP 1407754 B2	30 Nov 2011
		ES 2316699 T3	16 Apr 2009
		ES 2316699 T5	21 Mar 2012
		MX PA03009246 A	11 Apr 2005
		US 2004142828 A1	22 Jul 2004
		US 7119057 B2	10 Oct 2006
		US 2004071746 A1	15 Apr 2004
		US 7585824 B2	08 Sep 2009
		US 2004071742 A1	15 Apr 2004

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2015/050088

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (2015.01) A61Q 13/00, A61K 8/11, A61K 8/06, A61K 8/81, A61K 8/73, A61K 8/19, A61K 8/25, A61K 8/29, A61K 8/26, A61Q 15/00