



(11) **EP 3 330 352 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
06.06.2018 Bulletin 2018/23

(51) Int Cl.:
C11D 3/386 (2006.01)

(21) Application number: **17204768.0**

(22) Date of filing: **30.11.2017**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**
Designated Extension States:
BA ME
Designated Validation States:
MA MD

(71) Applicant: **The Procter & Gamble Company**
Cincinnati, OH 45202 (US)

(72) Inventor: **LANT, Neil Joseph**
Newcastle upon Tyne, NE12 9TS (GB)

(74) Representative: **Peet, Jillian Wendy**
Procter & Gamble Technical Centres Limited
Whitley Road
Longbenton
Newcastle upon Tyne
NE12 9TS (GB)

(30) Priority: **02.12.2016 EP 16202073**

(54) **CLEANING COMPOSITIONS INCLUDING ENZYMES AND ALKOXYLATED PHENOL**

(57) Cleaning compositions that include a galactanase enzyme and alkoxyated phenol, in particular an alkoxyated polyaryl phenol and/or an alkoxyated polyalkyl phenol. Methods of making and using such cleaning compositions. Use of a galactanase enzyme and an alkoxyated phenol.

EP 3 330 352 A1

Description

REFERENCE TO A SEQUENCE LISTING

5 **[0001]** This application contains a Sequence Listing in computer readable form, which is incorporated herein by reference.

FIELD OF THE INVENTION

10 **[0002]** The present invention relates to cleaning compositions that include a galactanase enzyme and alkoxyated phenol, in particular an alkoxyated polyaryl phenol and/or an alkoxyated polyalkyl phenol. The present invention also relates to methods of making and using such cleaning compositions. The present invention also relates to the use of a galactanase and an alkoxyated phenol.

15 BACKGROUND OF THE INVENTION

[0003] The laundry detergent formulator is constantly aiming to improve the performance of detergent compositions. Many detergent components may provide stain removal benefits, but some stains are particularly resistant to removal. Even when a soil is lifted from a fabric, the lifted material may redeposit onto the fabric during the wash cycle, impacting

20 whiteness and/or color contrast between white and colored regions on the same fabric.

[0004] There is a need for improved cleaning compositions that provide whiteness and/or color contrast benefits.

SUMMARY OF THE INVENTION

25 **[0005]** The present invention relates to a cleaning composition that includes: an endo-beta-1,6-galactanase enzyme; and an alkoxyated phenol compound that is selected from the group consisting of an alkoxyated polyaryl phenol compound, an alkoxyated polyalkyl phenol compound, and mixtures thereof.

[0006] The present invention also relates to a method of cleaning a surface, preferably a textile, where the method includes mixing the cleaning composition as described herein with water to form an aqueous liquor and contacting a

30 surface, preferably a textile, with the aqueous liquor in a laundering step.

[0007] The present invention also relates to the use of an endo-beta-1,6-galactanase enzyme and an alkoxyated polyaryl phenol and/or an alkoxyated polyalkyl phenol in a cleaning composition to enhance the stain-removal, whiteness, and/or for malodor reduction.

35 DETAILED DESCRIPTION OF THE INVENTION

[0008] The present invention relates to cleaning compositions that include a galactanase enzyme and an alkoxyated (polyaryl/polyalkyl) phenol. Without wishing to be bound by theory, it is believed that the galactanase enzymes are active on soils including soils of microbial origin such as extracellular DNA materials that adhere to, and cause other soils to

40 adhere to, fabric. It is further believed that the alkoxyated phenols described herein work synergistically with the galactanase enzymes to assist in the removal of extracellular DNA-bound soils and prevent redeposition of the reaction products arising from the action of the galactanase enzymes. In doing so, it is believed that the activity of the enzyme on the textile surface is enhanced by reduced interference or inhibition from redeposited soils.

[0009] The components of the compositions and processes of the present invention are described in more detail below.

45 **[0010]** As used herein, the articles "a" and "an" when used in a claim, are understood to mean one or more of what is claimed or described. As used herein, the terms "include," "includes," and "including" are meant to be non-limiting. The compositions of the present invention can comprise, consist essentially of, or consist of, the components of the present invention.

[0011] The terms "substantially free of" or "substantially free from" may be used herein. This means that the indicated material is at the very minimum not deliberately added to the composition to form part of it, or, preferably, is not present at analytically detectable levels. It is meant to include compositions whereby the indicated material is present only as an impurity in one of the other materials deliberately included. The indicated material may be present, if at all, at a level of less than 1%, or less than 0.1 %, or less than 0.01 %, or even 0%, by weight of the composition.

50 **[0012]** Unless otherwise noted, all component or composition levels are in reference to the active portion of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources of such components or compositions.

55 **[0013]** All temperatures herein are in degrees Celsius (°C) unless otherwise indicated. Unless otherwise specified, all measurements herein are conducted at 20°C and under the atmospheric pressure.

[0014] In all embodiments of the present invention, all percentages are by weight of the total composition, unless specifically stated otherwise. All ratios are weight ratios, unless specifically stated otherwise.

[0015] It should be understood that every maximum numerical limitation given throughout this specification includes every lower numerical limitation, as if such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

[0016] As used herein, the term "alkoxy" is intended to include C1-C8 alkoxy and C1-C8 alkoxy derivatives of polyols having repeating units such as butylene oxide, glycidol oxide, ethylene oxide or propylene oxide.

[0017] As used herein, unless otherwise specified, the terms "alkyl" and "alkyl capped" are intended to include C1-C18 alkyl groups, or even C1-C6 alkyl groups.

[0018] As used herein, unless otherwise specified, the term "aryl" is intended to include C3-12 aryl groups.

[0019] As used herein, unless otherwise specified, the term "arylalkyl" and "alkaryl" are equivalent and are each intended to include groups comprising an alkyl moiety bound to an aromatic moiety, typically having C1-C18 alkyl groups and, in one aspect, C1-C6 alkyl groups.

[0020] The terms "ethylene oxide," "propylene oxide" and "butylene oxide" may be shown herein by their typical designation of "EO," "PO" and "BO," respectively.

[0021] As used herein, the term "cleaning and/or treatment composition" includes, unless otherwise indicated, granular, powder, liquid, gel, paste, unit dose, bar form and/or flake type washing agents and/or fabric treatment compositions.

[0022] As used herein, "cellulosic substrates" are intended to include any substrate which comprises cellulose, either 100% by weight cellulose or at least 20% by weight, or at least 30% by weight or at least 40% or at least 50% by weight or even at least 60% by weight cellulose. Cellulose may be found in wood, cotton, linen, jute, and hemp. Cellulosic substrates may be in the form of powders, fibers, pulp and articles formed from powders, fibers and pulp. Cellulosic fibers, include, without limitation, cotton, rayon (regenerated cellulose), acetate (cellulose acetate), triacetate (cellulose triacetate), and mixtures thereof. Typically cellulosic substrates comprise cotton. Articles formed from cellulosic fibers include textile articles such as fabrics. Articles formed from pulp include paper.

[0023] As used herein, the term "maximum extinction coefficient" is intended to describe the molar extinction coefficient at the wavelength of maximum absorption (also referred to herein as the maximum wavelength), in the range of 400 nanometers to 750 nanometers.

[0024] As used herein "average molecular weight" is reported as a weight average molecular weight, as determined by its molecular weight distribution; as a consequence of their manufacturing process, polymers disclosed herein may contain a distribution of repeating units in their polymeric moiety.

[0025] As used herein the term "variant" refers to a polypeptide that contains an amino acid sequence that differs from a wild type or reference sequence. A variant polypeptide can differ from the wild type or reference sequence due to a deletion, insertion, or substitution of a nucleotide(s) relative to said reference or wild type nucleotide sequence. The reference or wild type sequence can be a full-length native polypeptide sequence or any other fragment of a full-length polypeptide sequence. A polypeptide variant generally has at least about 70% amino acid sequence identity with the reference sequence, but may include 75% amino acid sequence identity within the reference sequence, 80% amino acid sequence identity within the reference sequence, 85% amino acid sequence identity with the reference sequence, 86% amino acid sequence identity with the reference sequence, 87% amino acid sequence identity with the reference sequence, 88% amino acid sequence identity with the reference sequence, 89% amino acid sequence identity with the reference sequence, 90% amino acid sequence identity with the reference sequence, 91% amino acid sequence identity with the reference sequence, 92% amino acid sequence identity with the reference sequence, 93% amino acid sequence identity with the reference sequence, 94% amino acid sequence identity with the reference sequence, 95% amino acid sequence identity with the reference sequence, 96% amino acid sequence identity with the reference sequence, 97% amino acid sequence identity with the reference sequence, 98% amino acid sequence identity with the reference sequence, 98.5% amino acid sequence identity with the reference sequence or 99% amino acid sequence identity with the reference sequence.

[0026] As used herein, the term "solid" includes granular, powder, bar and tablet product forms.

[0027] As used herein, the term "fluid" includes liquid, gel, paste, and gas product forms.

Cleaning Composition

[0028] The present disclosure relates to cleaning and/or treatment compositions. The cleaning composition may be selected from the group of light duty liquid detergents compositions, heavy duty liquid detergent compositions, solid, for example particulate/powder or "dry" cleaning compositions, hard surface cleaning compositions, detergent gels commonly used for laundry, bleaching compositions, laundry additives, fabric enhancer compositions, shampoos, body

washes, other personal care compositions, and mixtures thereof. The cleaning composition may be a hard surface cleaning composition (such as a dishwashing composition) or a laundry composition (such as a heavy duty liquid or solid detergent composition).

[0029] The cleaning compositions may be in any suitable form. The composition can be selected from a liquid, solid, or combination thereof. As used herein, "liquid" includes free-flowing liquids, as well as pastes, gels, foams and mousses. Non-limiting examples of liquids include light duty and heavy duty liquid detergent compositions, fabric enhancers, detergent gels commonly used for laundry, bleach and laundry additives. Gases, e.g., suspended bubbles, or solids, e.g. particles, may be included within the liquids. A "solid" as used herein includes, but is not limited to, powders, agglomerates, and mixtures thereof. Non-limiting examples of solids include: granules, micro-capsules, beads, noodles, and pearlised balls. Solid compositions may provide a technical benefit including, but not limited to, through-the-wash benefits, pre-treatment benefits, and/or aesthetic effects.

[0030] The cleaning composition may be in the form of a unitized dose article, such as a tablet or in the form of a pouch. Such pouches typically include a water-soluble film, such as a polyvinyl alcohol water-soluble film, that at least partially encapsulates a composition. Suitable films are available from MonoSol, LLC (Indiana, USA). The composition can be encapsulated in a single or multi-compartment pouch. A multi-compartment pouch may have at least two, at least three, or at least four compartments. A multi-compartmented pouch may include compartments that are side-by-side and/or superposed. The composition contained in the pouch may be liquid, solid (such as powders), or combinations thereof.

Galactanase Enzyme

[0031] The endo-beta-1,6-galactanase enzyme is an extracellular polymer-degrading enzyme. The term "endo-beta-1,6-galactanase" or "a polypeptide having endo-beta-1,6-galactanase activity" means a endo-beta-1,6-galactanase activity (EC 3.2.1.164) that catalyzes the hydrolytic cleavage of 1,6-3-D-galactooligosaccharides with a degree of polymerization (DP) higher than 3, and their acidic derivatives with 4-O-methylglucosyluronate or glucosyluronate groups at the non-reducing terminals.

[0032] For purposes of the present disclosure, endo-beta-1,6-galactanase activity is determined according to the procedure described in WO 2015185689 in Assay I. Suitable examples from class EC 3.2.1.164 are described in WO 2015185689, such as the mature polypeptide SEQ ID NO: 2 described therein. Preferably the galactanase enzyme is selected from Glycoside Hydrolase (GH) Family 30.

[0033] Preferably, the endo-beta-1,6-galactanase comprises a microbial enzyme. The endo-beta-1,6-galactanase may be fungal or bacterial in origin. Bacterial endo-beta-1,6-galactanase may be most preferred. Fungal endo-beta-1,6-galactanase may be most preferred.

[0034] A bacterial endo-beta-1,6-galactanase is obtainable from *Streptomyces*, for example *Streptomyces davawensis*. A preferred endo-beta-1,6-galactanase is obtainable from *Streptomyces davawensis* JCM 4913 defined in SEQ ID NO: 1 herein, or a variant thereof, for example having at least 40% or 50% or 60% or 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identity thereto.

[0035] Other bacterial endo-beta-1,6-galactanase include those encoded by the DNA sequences of *Streptomyces avermitilis* MA-4680 with amino acid sequence defined in SEQ ID NO: 2 herein, or a variant thereof, for example having at least 40% or 50% or 60% or 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identity thereto.

[0036] A fungal endo-beta-1,6-galactanase is obtainable from *Trichoderma*, for example *Trichoderma harzianum*. A preferred endo-beta-1,6-galactanase is obtainable from *Trichoderma harzianum* defined in SEQ ID NO: 3 herein, or a variant thereof, for example having at least 40% or 50% or 60% or 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto.

[0037] Other fungal endo-beta-1,6-galactanases include those encoded by the DNA sequences of *Ceratocystis fimbriata* f. sp. Platani, *Muscodor strobilii* WG-2009a, *Oculimacula yallundae*, *Trichoderma viride* GD36A, *Thermomyces stellatus*, *Myceliophthora thermophila*.

[0038] Preferably the galactanase has an amino acid sequence having at least 60%, or at least 80%, or at least 90% or at least 95% identity with the amino acid sequence shown in SEQ ID NO:1, SEQ ID NO:2 or SEQ ID NO:3. Preferably the galactanase is an isolated galactanase.

[0039] Preferably the galactanase enzyme is present in a laundering aqueous liquor in an amount of from 0.01ppm to 1000 ppm of the galactanase enzyme, or from 0.05 or from 0.1ppm to 750 or 500ppm.

[0040] The galactanase or composition comprising galactanase may also give rise to biofilm-disrupting effects.

Alkoxyated phenol compound


[0041] The cleaning compositions of the present invention include an alkoxyated phenol compound. The alkoxyated phenol compound may be selected from the group consisting of an alkoxyated polyaryl phenol compound, an alkoxyated

polyalkyl phenol compound, and mixtures thereof. The alkoxyated phenol compound may be an alkoxyated polyaryl phenol compound. The alkoxyated phenol compound may be an alkoxyated polyalkyl phenol compound.

[0042] The alkoxyated phenol compound may be present in the cleaning composition at a level of from about 0.2% to about 10%, or from about 0.5% to about 5%, by weight of the cleaning composition.

[0043] The alkoxyated phenol compound preferably has a weight average molecular weight between 280 and 2880.


[0044] The alkoxyated phenol compound may have the following structure:



wherein each R_1 is independently selected from linear or branched C3-C15 alkyl groups and aryl groups, X is selected from ethoxy or propoxy groups, n is from 2 to 70, T is selected from H, SO_3^- , COO^- and PO_3^{2-} , preferably H and SO_3^- . Each R_1 may be the same.


[0045] The alkoxyated polyaryl or alkoxyated polyalkyl phenol compound is preferably selected from groups (i) to (v):

(i) Uncharged alkoxyated tristyrylphenols of the following structure:



wherein n is selected from 2 to 70, more preferably n is selected from 10 to 54, most preferably n = 16 or 20;


(ii) Anionic alkoxyated tristyrylphenols of the following structure



wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- , wherein n is selected from 2 to 54;

(iii) Uncharged alkoxyated tri(n-butyl)phenols of the following structure:

5




10

wherein n is selected from 2 to 50;

(iv) Anionic alkoxyated tri(n-butyl)phenols of the following structure:

15



20

25

wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- , wherein n is selected from 6 to 50; and

(v) mixtures thereof.

30

[0046] Such compounds are available from industrial suppliers, for example Solvay under the Soprophor trade name, from Clariant under the Emulsogen trade name, Aoki Oil Industrial Co. under the Blaunon trade name, from Stepan under the Makon trade name, and from TOTO Chemical Industry Co. under the Sorpol trade name.

35 Adjuncts

[0047] The cleaning compositions described herein may optionally include other adjunct components, for example selected from surfactants, fabric shading dyes, fabric care benefit agent; additional enzyme; deposition aid; rheology modifier; builder; chelant; bleach; bleaching agent; bleach precursor; bleach booster; bleach activator, bleach catalyst; perfume and/or perfume microcapsules; perfume loaded zeolite; starch encapsulated accord; polyglycerol esters; whitening agent; pearlescent agent; enzyme stabilizing systems; scavenging agents including fixing agents for anionic dyes, complexing agents for anionic surfactants, and mixtures thereof; optical brighteners or fluoescers; polymer including but not limited to soil release polymer and/or soil suspension polymer; dispersants; antifoam agents; non-aqueous solvent; fatty acid; suds suppressors, e.g., silicone suds suppressors; cationic starches; scum dispersants; substantive dyes; colorants; opacifier; antioxidant; hydrotropes such as toluenesulfonates, cumenesulfonates and naphthalenesulfonates; color speckles; colored beads, spheres or extrudates; clay softening agents; anti-bacterial agents. Additionally or alternatively, the compositions may comprise surfactants, and/or solvent systems. Quaternary ammonium compounds may be present, particularly in fabric enhancer compositions, such as fabric softeners, and comprise quaternary ammonium cations that are positively charged polyatomic ions of the structure NR_4^+ , where R is an alkyl group or an aryl group.

50

Additional Enzymes

[0048] Preferably the composition of the invention comprises additional enzymes, for example selected from lipases, amylases, proteases, nucleases, pectate lyases, cellulases, cutinases, and mixtures thereof. The cleaning compositions preferably comprise one or more additional enzymes from the group selected from nucleases. The cleaning compositions preferably comprises one or more additional enzymes selected from the group amylases, lipases, proteases, pectate lyases, cellulases, cutinases, and mixtures thereof. Preferably, the cleaning compositions comprises one or more addi-

55

tional enzymes selected from amylases and proteases and mixtures thereof. Preferably the cleaning compositions comprise one or more additional enzymes selected from lipases. The compositions may also comprise hemicellulases, peroxidases, xylanases, pectinases, keratinases, reductases, oxidases, phenoloxidases, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, lac-
 5 case and mixtures thereof. When present in the composition, the aforementioned additional enzymes may be present at levels from about 0.00001% to about 2%, from about 0.0001 % to about 1% or even from about 0.001 % to about 0.5% enzyme protein by weight of the composition. Preferably the or each additional enzyme is present in the laundering aqueous liquor in an amount of from 0.01ppm to 1000 ppm of the active enzyme protein, or from 0.05 or from 0.1ppm to 750 or 500ppm.

Nucleases

[0049] Preferably the composition additionally comprises a nuclease enzyme. The nuclease enzyme is an enzyme capable of cleaving the phosphodiester bonds between the nucleotide subunits of nucleic acids. Suitable nuclease
 15 enzymes may be deoxyribonuclease or ribonuclease enzyme or a functional fragment thereof. By functional fragment or part is meant the portion of the nuclease enzyme that catalyzes the cleavage of phosphodiester linkages in the DNA backbone and so is a region of said nuclease protein that retains catalytic activity. Thus it includes truncated, but functional versions, of the enzyme and/or variants and/or derivatives and/or homologues whose functionality is maintained.

[0050] Preferably the nuclease enzyme is a deoxyribonuclease, preferably selected from any of the classes E.C. 3.1.21.x, where x=1, 2, 3, 4, 5, 6, 7, 8 or 9, E.C. 3.1.22.y where y=1, 2, 4 or 5, E.C. 3.1.30.z where z= 1 or 2, E.C. 3.1.31.1 and mixtures thereof. Nuclease enzymes from class E.C. 3.1.21.x and especially where x=1 are particularly preferred. Nucleases in class E.C. 3.1.22.y cleave at the 5' hydroxyl to liberate 3' phosphomonoesters. Enzymes in class E.C. 3.1.30.z may be preferred as they act on both DNA and RNA and liberate 5'-phosphomonoesters. Suitable examples from class E.C. 3.1.31.2 are described in US2012/0135498A, such as SEQ ID NO:3 therein. Such enzymes are commercially available as DENARASE® enzyme from c-LECTA. Nuclease enzymes from class E.C. 3.1.31.1 produce 3'phosphomonoesters.

[0051] Preferably, the nuclease enzyme comprises a microbial enzyme. The nuclease enzyme may be fungal or bacterial in origin. Bacterial nucleases may be most preferred. Fungal nucleases may be most preferred.

[0052] The microbial nuclease is obtainable from *Bacillus*, such as a *Bacillus licheniformis* or *Bacillus subtilis* bacterial nucleases. A preferred nuclease is obtainable from *Bacillus licheniformis*, preferably from strain EI-34-6. A preferred deoxyribonuclease is a variant of *Bacillus licheniformis*, from strain EI-34-6 nucB deoxyribonuclease defined in SEQ ID NO:4 herein, or variant thereof, for example having at least 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto. Other suitable nucleases are defined in SEQ ID NO: 5 herein, or variant thereof, for example having at least 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto.
 35 Other suitable nucleases are defined in SEQ ID NO: 6 herein, or variant thereof, for example having at least 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto.

[0053] A fungal nuclease is obtainable from *Aspergillus*, for example *Aspergillus oryzae*. A preferred nuclease is obtainable from *Aspergillus oryzae* defined in SEQ ID NO: 7 herein, or variant thereof, for example having at least 60% or 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto.

[0054] Another suitable fungal nuclease is obtainable from *Trichoderma*, for example *Trichoderma harzianum*. A preferred nuclease is obtainable from *Trichoderma harzianum* defined in SEQ ID NO: 8 herein, or variant thereof, for example having at least 60% or 70% or 75% or 80% or 85% or 90% or 95%, 96%, 97%, 98%, 99% or 100% identical thereto.

[0055] Other fungal nucleases include those encoded by the DNA sequences of *Aspergillus oryzae* RIB40, *Aspergillus oryzae* 3.042, *Aspergillus flavus* NRRL3357, *Aspergillus parasiticus* SU-1, *Aspergillus nomius* NRRL13137, *Trichoderma reesei* QM6a, *Trichoderma virens* Gv29-8, *Oidiodendron maius* Zn, *Metarhizium guizhouense* ARSEF 977, *Metarhizium majus* ARSEF 297, *Metarhizium robertsii* ARSEF 23, *Metarhizium acridum* CQMa 102, *Metarhizium brunneum* ARSEF 3297, *Metarhizium anisopliae*, *Colletotrichum fiorinae* PJ7, *Colletotrichum sublineola*, *Trichoderma atroviride* IMI 206040, *Tolyposcladium ophioglossoides* CBS 100239, *Beauveria bassiana* ARSEF 2860, *Colletotrichum higginsianum*, *Hirsutella minnesotensis* 3608, *Scedosporium apiospermum*, *Phaeoconiella chlamydozpora*, *Fusarium verticillioides* 7600, *Fusarium oxysporum* f. sp. cubense race 4, *Colletotrichum graminicola* M1.001, *Fusarium oxysporum* FOSC 3-a, *Fusarium avenaceum*, *Fusarium langsethiae*, *Grosmannia clavigera* kw1407, *Claviceps purpurea* 20.1, *Verticillium longisporum*, *Fusarium oxysporum* f. sp. cubense race 1, *Magnaporthe oryzae* 70-15, *Beauveria bassiana* D1-5, *Fusarium pseudograminearum* CS3096, *Neonectria ditissima*, *Magnaportheopsis poae* ATCC 64411, *Cordyceps militaris* CM01, *Marssonina brunnea* f. sp. 'multigermtubi' MB_m1, *Diaporthe ampelina*, *Metarhizium album* ARSEF 1941, *Colletotrichum gloeosporioides* Nara gc5, *Madurella mycetomatis*, *Metarhizium brunneum* ARSEF 3297, *Verticillium alfalfae* VaMs.102, *Gaeumannomyces graminis* var. tritici R3-111a-1, *Nectria haematococca* mpVI 77-13-4, *Verticillium longisporum*, *Verticillium dahliae* VdLs.17, *Torrubiella hemipterigena*, *Verticillium longisporum*, *Verticillium dahliae* VdLs.17, *Botrytis cinerea* B05.10, *Chaetomium globosum* CBS 148.51, *Metarhizium anisopliae*, *Stemphylium lycopersici*, *Sclerotinia bore-*

alis F-4157, *Metarhizium robertsii* ARSEF 23, *Myceliophthora thermophila* ATCC 42464, *Phaeosphaeria nodorum* SN15, *Phialophora attae*, *Ustilagoidea virens*, *Diplodia seriata*, *Ophiostoma piceae* UAMH 11346, *Pseudogymnoascus pan-norum* VKM F-4515 (FW-2607), *Bipolaris oryzae* ATCC 44560, *Metarhizium guizhouense* ARSEF 977, *Chaetomium thermophilum* var. *thermophilum* DSM 1495, *Pestalotiopsis fici* W106-1, *Bipolaris zeicola* 26-R-13, *Setosphaeria turcica* Et28A, *Arthroderma otae* CBS 113480 and *Pyrenophora tritici-repentis* Pt-1C-BFP.

[0056] Preferably the nuclease is an isolated nuclease.

[0057] Preferably the nuclease enzyme is present in the laundering aqueous liquor in an amount of from 0.01ppm to 1000 ppm of the nuclease enzyme, or from 0.05 or from 0.1ppm to 750 or 500ppm.

Acetylglucosaminidases.

[0058] Preferably the composition comprises an acetylglucosaminidase enzyme, preferably a β -N-acetylglucosaminidase enzyme from E.C. 3.2.1.52, preferably an enzyme having at least 70%, or at least 75% or at least 80% or at least 85% or at least 90% or at least 95% or at least 96% or at least 97% or at least 98% or at least 99% or at least or 100% identity to SEQ ID NO: 9.

Mannanases

[0059] Preferably the composition comprises a mannanase enzyme. The term "mannanase" means a polypeptide having mannan endo-1,4- beta-mannosidase activity (EC 3.2.1.78) from the glycoside hydrolase family 26 that catalyzes the hydrolysis of 1 ,4-3-D-mannosidic linkages in mannans, galactomannans and glucomannans. Alternative names of mannan endo-1,4-beta-mannosidase are 1,4-3-D-mannan mannanohydrolase; endo-1,4-3-mannanase; endo- β -1,4-mannase; β -mannanase B; 3-1,4-mannan 4-mannanohydrolase; endo-3-mannanase; and β -D-mannanase. Preferred mannanases are members of the glycoside hydrolase family 26.

[0060] For purposes of the present disclosure, mannanase activity may be determined using the Reducing End Assay as described in the experimental section of WO 2015040159. Suitable examples from class EC 3.2.1.78 are described in WO 2015040159, such as the mature polypeptide SEQ ID NO: 2 described therein.

[0061] Preferred mannanases are variants having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 81 %, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91 %, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide SEQ ID NO: 10 from *Ascobolus stictioideus*; Preferred mannanases are variants having at least 81 %, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91 %, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide SEQ ID NO: 11 from *Chaetomium virescens*.

[0062] Preferred mannanases are variants having at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91 %, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide SEQ ID NO: 12 from *Preussia aemulans*.

[0063] Preferred mannanases are variants having at least at least 65%, at least 66%, at least 67%, at least 68%, at least 69%, at least 70%, at least 71%, at least 72%, at least 73%, at least 74%, at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91 %, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide SEQ ID NO: 13 from *Yunnania penicillata*.

[0064] Preferred mannanases are variants having at least at least 75%, at least 76%, at least 77%, at least 78%, at least 79%, at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91 %, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide SEQ ID NO: 14 from *Myrothecium roridum*. Preferably the mannanase is an isolated mannanase.

[0065] Preferably the mannanase enzyme is present in the cleaning compositions in an amount from 0.001 to 1 wt% based on active protein in the composition, or from 0.005 to 0.5 wt% or from 0.01 to 0.25 wt%. Preferably the mannanase enzyme is present in the laundering aqueous liquor in an amount of from 0.01ppm to 1000 ppm of the mannanase enzyme, or from 0.05 or from 0.1 ppm to 750 or 500ppm. The compositions of the invention comprising both galactanase and mannanase may be particularly effective against sticky soils and for improved cleaning. It is believed the two enzymes function together in a complementary way.

Further Glycosyl Hydrolases

[0066] The composition may comprise a glycosyl hydrolase selected from GH family 39 and GH family 114 and mixtures thereof, for example as described in co-pending applications having applicants reference numbers CM4645FM and CM4646 FM, respectively.

Proteases.

[0067] Preferably the composition comprises one or more proteases. Suitable proteases include metalloproteases and serine proteases, including neutral or alkaline microbial serine proteases, such as subtilisins (EC 3.4.21.62). Suitable proteases include those of animal, vegetable or microbial origin. In one aspect, such suitable protease may be of microbial origin. The suitable proteases include chemically or genetically modified mutants of the aforementioned suitable proteases. In one aspect, the suitable protease may be a serine protease, such as an alkaline microbial protease or/and a trypsin-type protease. Examples of suitable neutral or alkaline proteases include:

(a) subtilisins (EC 3.4.21.62), preferably those derived from *Bacillus* sp., such as *B. lentus*, *B. alkalophilus*, *B. subtilis*, *B. amyloliquefaciens*, *B. pumilus* and *B. gibsonii* and *B. akibaii* described in WO2004067737, WO2015091989, WO2015091990, WO2015024739, WO2015143360, US 6,312,936 B1, US 5,679,630, US 4,760,025, US7,262,042 and WO09/021867, DE102006022216A1, DE102006022224A1, WO2015089447, WO2015089441, WO2016066756, WO2016066757, WO2016069557, WO2016069563, WO2016069569.

(b) trypsin-type or chymotrypsin-type proteases, such as trypsin (e.g., of porcine or bovine origin), including the *Fusarium* protease described in WO 89/06270 and the chymotrypsin proteases derived from *Cellulomonas* described in WO 05/052161 and WO 05/052146.

(c) metalloproteases, preferably those derived from *Bacillus amyloliquefaciens* described in WO 07/044993A2; from *Bacillus*, *Brevibacillus*, *Thermoactinomyces*, *Geobacillus*, *Paenibacillus*, *Lysinibacillus* or *Streptomyces* spp. Described in WO2014194032, WO2014194054 and WO2014194117; from *Kribella alluminosa* described in WO2015193488; and from *Streptomyces* and *Lysobacter* described in WO2016075078.

(d) protease having at least 90% identity to the subtilase from *Bacillus* sp. TY145, NCIMB 40339, described in WO92/17577 (Novozymes A/S), including the variants of this *Bacillus* sp TY145 subtilase described in WO2015024739, and WO2016066757.

[0068] Preferred proteases include those derived from *Bacillus gibsonii* or *Bacillus Lentus*.

[0069] Suitable commercially available protease enzymes include those sold under the trade names Alcalase®, Savinase®, Primase®, Durazym®, Polarzyme®, Kannase®, Liquanase®, Liquanase Ultra®, Savinase Ultra®, Ovozyme®, Neutrase®, Everlase® and Esperase® by Novozymes A/S (Denmark), those sold under the tradename Maxatase®, Maxacal®, Maxapem®, Properase®, Purafect®, Purafect Prime®, Purafect Ox®, FN3®, FN4®, Excellase® and Purafect OXP® by Genencor International, those sold under the tradename Opticlean® and Optimase® by Solvay Enzymes, those available from Henkel/ Kemira, namely BLAP (sequence shown in Figure 29 of US 5,352,604 with the following mutations S99D + S101 R + S103A + V104I + G159S, hereinafter referred to as BLAP), BLAP R (BLAP with S3T + V4I + V199M + V205I + L217D), BLAP X (BLAP with S3T + V4I + V205I) and BLAP F49 (BLAP with S3T + V4I + A194P + V199M + V205I + L217D) - all from Henkel/Kemira; and KAP (*Bacillus alkalophilus* subtilisin with mutations A230V + S256G + S259N) from Kao, or as disclosed in WO2009/149144, WO2009/149145, WO2010/56653, WO2010/56640, WO2011/07211 US2011/0237487, WO2011/140316, WO2012/151480, EP2510092, EP2566960 OR EP2705145.

Amylases

[0070] Preferably the composition may comprise an amylase. Suitable alpha-amylases include those of bacterial or fungal origin. Chemically or genetically modified mutants (variants) are included. A preferred alkaline alpha-amylase is derived from a strain of *Bacillus*, such as *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Bacillus stearothermophilus*, *Bacillus subtilis*, or other *Bacillus* sp., such as *Bacillus* sp. NCIB 12289, NCIB 12512, NCIB 12513, DSM 9375 (USP 7,153,818) DSM 12368, DSMZ no. 12649, KSM AP1378 (WO 97/00324), KSM K36 or KSM K38 (EP 1,022,334). Preferred amylases include:

(a) the variants described in WO 94/02597, WO 94/18314, WO96/23874 and WO 97/43424, especially the variants with substitutions in one or more of the following positions versus the enzyme listed as SEQ ID No. 2 in WO 96/23874: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444.

(b) the variants described in USP 5,856,164 and WO99/23211, WO 96/23873, WO00/60060 and WO 06/002643, especially the variants with one or more substitutions in the following positions versus the AA560 enzyme listed as

SEQ ID No. 12 in WO 06/002643:

26, 30, 33, 82, 37, 106, 118, 128, 133, 149, 150, 160, 178, 182, 186, 193, 203, 214, 231, 256, 257, 258, 269,
270, 272, 283, 295, 296, 298, 299, 303, 304, 305, 311, 314, 315, 318, 319, 339, 345, 361, 378, 383, 419, 421,
437, 441, 444, 445, 446, 447, 450, 461, 471, 482, 484, preferably that also contain the deletions of D183* and
G184*.

(c) variants exhibiting at least 90% identity with SEQ ID No. 4 in WO06/002643, the wild-type enzyme from *Bacillus* SP722, especially variants with deletions in the 183 and 184 positions and variants described in WO 00/60060, which is incorporated herein by reference.

(d) variants exhibiting at least 95% identity with the wild-type enzyme from *Bacillus* sp.707 (SEQ ID NO:7 in US 6,093, 562), especially those comprising one or more of the following mutations M202, M208, S255, R172, and/or M261. Preferably said amylase comprises one or more of M202L, M202V, M202S, M202T, M202I, M202Q, M202W, S255N and/or R172Q. Particularly preferred are those comprising the M202L or M202T mutations.

(e) variants described in WO 09/149130, preferably those exhibiting at least 90% identity with SEQ ID NO: 1 or SEQ ID NO:2 in WO 09/149130, the wild-type enzyme from *Geobacillus Stearophermophilus* or a truncated version thereof;

(f) variants as described in EP2540825 and EP2357220, EP2534233; (g) variants as described in WO2009100102 and WO2010115028.

[0071] Suitable commercially available alpha-amylases include DURAMYL®, LIQUEZYME®, TERMAMYL®, TERMAMYL ULTRA®, NATALASE®, SUPRAMYL®, STAINZYME®, STAINZYME PLUS®, FUNGAMYL® and BAN® (Novozymes A/S, Bagsvaerd, Denmark), KEMZYM® AT 9000 Biozym Biotech Trading GmbH Wehlstrasse 27b A-1200 Wien Austria, RAPIDASE®, PURASTAR®, ENZYSIZE®, OPTISIZE HT PLUS®, POWERASE® and PURASTAR OX-AM® (Genencor International Inc., Palo Alto, California) and KAM® (Kao, 14-10 Nihonbashi Kayabacho, 1-chome, Chuo-ku Tokyo 103-8210, Japan). In one aspect, suitable amylases include NATALASE®, STAINZYME® and STAINZYME PLUS® and mixtures thereof.

Lipases

[0072] Preferably the composition comprises one or more lipases, including "first cycle lipases" such as those described in U.S. Patent 6,939,702 B1 and US PA 2009/0217464. Preferred lipases are first-wash lipases. In one embodiment of the invention the composition comprises a first wash lipase. First wash lipases includes a lipase which is a polypeptide having an amino acid sequence which: (a) has at least 90% identity with the wild-type lipase derived from *Humicola lanuginosa* strain DSM 4109; (b) compared to said wild-type lipase, comprises a substitution of an electrically neutral or negatively charged amino acid at the surface of the three-dimensional structure within 15Å of E1 or Q249 with a positively charged amino acid; and (c) comprises a peptide addition at the C-terminal; and/or (d) comprises a peptide addition at the N-terminal and/or (e) meets the following limitations: i) comprises a negative amino acid in position E210 of said wild-type lipase; ii) comprises a negatively charged amino acid in the region corresponding to positions 90-101 of said wild-type lipase; and iii) comprises a neutral or negative amino acid at a position corresponding to N94 or said wild-type lipase and/or has a negative or neutral net electric charge in the region corresponding to positions 90-101 of said wild-type lipase. Preferred are variants of the wild-type lipase from *Thermomyces lanuginosus* comprising one or more of the T231R and N233R mutations. The wild-type sequence is the 269 amino acids (amino acids 23 - 291) of the Swissprot accession number Swiss-Prot 059952 (derived from *Thermomyces lanuginosus* (*Humicola lanuginosa*)). Preferred lipases include those sold under the tradenames Lipex® and Lipolex® and Lipoclean®. Other suitable lipases include those described in European Patent Application No. 12001034.3 or EP2623586.

Endoglucanases

[0073] Other preferred enzymes include microbial-derived endoglucanases exhibiting endo-beta-1,4-glucanase activity (E.C. 3.2.1.4), including a bacterial polypeptide endogenous to a member of the genus *Bacillus* which has a sequence of at least 90%, 94%, 97% and even 99% identity to the amino acid sequence SEQ ID NO:2 in US7,141,403B2) and mixtures thereof. Suitable endoglucanases are sold under the tradenames Celluclean® and Whitezyme® (Novozymes A/S, Bagsvaerd, Denmark).

Pectate Lyases

[0074] Other preferred enzymes include pectate lyases sold under the tradenames Pectawash®, Pectaway®, Xpect®

and mannanases sold under the tradenames Mannaway® (all from Novozymes A/S, Bagsvaerd, Denmark), and Purabrite® (Genencor International Inc., Palo Alto, California).

Cleaning Cellulase

5

[0075] The cleaning composition described herein may additionally comprise a cleaning cellulase. The cellulase may be an endoglucanase. The cellulase may have endo beta 1,4-glucanase activity and a structure which does not comprise a class A Carbohydrate Binding Module (CBM). A class A CBM is defined according to A. B. Boraston et al. Biochemical Journal 2004, Volume 382 (part 3) pages 769-781. In particular, the cellulase does not comprise a class A CBM from families 1, 2a, 3, 5 and 10.

10

[0076] The cellulase may be a glycosyl hydrolase having enzymatic activity towards amorphous cellulose substrates, wherein the glycosyl hydrolase is selected from GH families 5, 7, 12, 16, 44 or 74. Preferably, the cellulase is a glycosyl hydrolase selected from GH family 5. A preferred cellulase is Celluclean, supplied by Novozymes. This preferred cellulase is described in more detail in WO2002/099091. The glycosyl hydrolase (GH) family definition is described in more detail in Biochem J. 1991, v280, 309-316. Another preferred cellulase is a glycosyl hydrolase having enzymatic activity towards both xyloglucan and amorphous cellulose substrates, wherein the glycosyl hydrolase is selected from GH families 5, 12, 44 or 74. Preferably, the glycosyl hydrolase selected from GH family 44.

15

[0077] For purposes of the present invention, the degree of identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends in Genetics 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows: (Identical Residues x 100)/(Length of Alignment - Total Number of Gaps in Alignment).

20

[0078] Suitable cleaning cellulase glycosyl hydrolases are selected from the group consisting of: GH family 44 glycosyl hydrolases from *Paenibacillus polyxyrna* (wild-type) such as XYG1006 described in WO 01/062903 or are variants thereof; GH family 12 glycosyl hydrolases from *Bacillus licheniformis* (wild-type) such as Seq. No. ID: 1 described in WO 99/02663 or are variants thereof; GH family 5 glycosyl hydrolases from *Bacillus agaradhaerens* (wild type) or variants thereof; GH family 5 glycosyl hydrolases from *Paenibacillus* (wild type) such as XYG1034 and XYG 1022 described in WO 01/064853 or variants thereof; GH family 74 glycosyl hydrolases from *Jonesia sp.* (wild type) such as XYG1020 described in WO 2002/077242 or variants thereof; and GH family 74 glycosyl hydrolases from *Trichoderma Reesei* (wild type), such as the enzyme described in more detail in Sequence ID no. 2 of WO03/089598, or variants thereof.

25

[0079] Preferred glycosyl hydrolases are selected from the group consisting of: GH family 44 glycosyl hydrolases from *Paenibacillus polyxyrna* (wild-type) such as XYG1006 or are variants thereof.

30

[0080] Typically, the cellulase modifies the fabric surface during the laundering process so as to improve the removal of soils adhered to the fabric after the laundering process during wearing and usage of the fabric, in subsequent wash cycles. Preferably, the cellulase modifies the fabric surface during the laundering process so as to improve the removal of soils adhered to the fabric after the laundering process during wearing and usage of the fabric, in the subsequent two, or even three wash cycles.

35

[0081] Typically, the cellulase is used at a concentration of 0.005ppm to 1.0ppm in the aqueous liquor during the first laundering process. Preferably, the cellulase is used at a concentration of 0.02ppm to 0.5ppm in the aqueous liquor during the first laundering process.

40

Surfactant system

45

[0082] The cleaning composition may comprise a surfactant system. The cleaning composition may comprise from about 1% to about 80%, or from 1% to about 60%, preferably from about 5% to about 50% more preferably from about 8% to about 40%, by weight of the cleaning composition, of a surfactant system.

[0083] Surfactants suitable for use in the surfactant system may be derived from natural and/or renewable sources.

50

[0084] The surfactant system may comprise an anionic surfactant, more preferably an anionic surfactant selected from the group consisting of, alkyl benzene sulfonate, alkyl sulfate, alkyl alkoxy sulfate, especially alkyl ethoxy sulfate, paraffin sulfonate and mixtures thereof, alkyl benzene sulfonates are particularly preferred. The surfactant system may further comprise a surfactant selected from the group consisting of nonionic surfactant, cationic surfactant, amphoteric surfactant, zwitterionic surfactant, and mixtures thereof. The surfactant system preferably comprises a nonionic surfactant, for example an ethoxylated nonionic surfactant. The surfactant system may comprise an amphoteric surfactant, for example an amine oxide surfactant, such as an alkyl dimethyl amine oxide. The surfactant system may comprise a zwitterionic surfactant, such as a betaine.

55

[0085] The most preferred surfactant system for the detergent composition of the present invention comprises from

1% to 40%, preferably 6% to 35%, more preferably 8% to 30% weight of the total composition of an anionic surfactant, preferably comprising an alkyl benzene sulphonate. The preferred surfactant system may optionally in addition comprise an alkyl alkoxy sulfate surfactant, more preferably an alkyl ethoxy sulfate, optionally combined with 0.5% to 15%, preferably from 1% to 12%, more preferably from 2% to 10% by weight of the composition of amphoteric and/or zwitterionic surfactant, more preferably an amphoteric and even more preferably an amine oxide surfactant, especially an alkyl dimethyl amine oxide.

[0086] Preferably the composition further comprises a nonionic surfactant, especially an alcohol alkoxyate in particular an alcohol ethoxyate nonionic surfactant. Most preferably the surfactant system comprises an anionic and a nonionic surfactant, preferably the weight ratio of the anionic to nonionic surfactant is from 25:1 to 1:2.

Anionic surfactant

[0087] Anionic surfactants may be in salt form or acid form, typically in the form of a water-soluble sodium, potassium, ammonium, magnesium or mono-, di- or tri- C2-C3 alkanolammonium salt, with the sodium cation being the usual one chosen.

Sulfonate Surfactant

[0088] Suitable anionic sulfonate surfactants for use herein include water-soluble salts of C8-C18 alkyl or hydroxyalkyl sulfonates; C11-C18 alkyl benzene sulfonates (LAS), modified alkylbenzene sulfonate (MLAS) as discussed in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO 00/23549, and WO 00/23548; methyl ester sulfonate (MES); and alpha-olefin sulfonate (AOS). Those also include the paraffin sulfonates may be monosulfonates and/or disulfonates, obtained by sulfonating paraffins of 10 to 20 carbon atoms. The sulfonate surfactant may also include the alkyl glyceryl sulfonate surfactants.

Sulfated anionic surfactant

[0089] Preferably the sulfated anionic surfactant is alkoxyated, more preferably, an alkoxyated branched sulfated anionic surfactant having an alkoxylation degree of from about 0.2 to about 4, even more preferably from about 0.3 to about 3, even more preferably from about 0.4 to about 1.5 and especially from about 0.4 to about 1. Preferably, the alkoxy group is ethoxy. When the sulfated anionic surfactant is a mixture of sulfated anionic surfactants, the alkoxylation degree is the weight average alkoxylation degree of all the components of the mixture (weight average alkoxylation degree). In the weight average alkoxylation degree calculation the weight of sulfated anionic surfactant components not having alkoxyated groups should also be included.

$$\text{Weight average alkoxylation degree} = (x_1 * \text{alkoxylation degree of surfactant 1} + x_2 * \text{alkoxylation degree of surfactant 2} + \dots) / (x_1 + x_2 + \dots)$$

wherein x_1, x_2, \dots are the weights in grams of each sulfated anionic surfactant of the mixture and alkoxylation degree is the number of alkoxy groups in each sulfated anionic surfactant.

[0090] Preferably, the branching group is an alkyl. Typically, the alkyl is selected from methyl, ethyl, propyl, butyl, pentyl, cyclic alkyl groups and mixtures thereof. Single or multiple alkyl branches could be present on the main hydrocarbyl chain of the starting alcohol(s) used to produce the sulfated anionic surfactant used in the detergent of the invention. Most preferably the branched sulfated anionic surfactant is selected from alkyl sulfates, alkyl ethoxy sulfates, and mixtures thereof.

[0091] The branched sulfated anionic surfactant can be a single anionic surfactant or a mixture of anionic surfactants. In the case of a single surfactant the percentage of branching refers to the weight percentage of the hydrocarbyl chains that are branched in the original alcohol from which the surfactant is derived.

[0092] In the case of a surfactant mixture the percentage of branching is the weight average and it is defined according to the following formula:

$$\text{Weight average of branching (\%)} = [(x_1 * \text{wt\% branched alcohol 1 in alcohol 1} + x_2 * \text{wt\% branched alcohol 2 in alcohol 2} + \dots) / (x_1 + x_2 + \dots)] * 100$$

wherein x1, x2, ... are the weight in grams of each alcohol in the total alcohol mixture of the alcohols which were used as starting material for the anionic surfactant for the detergent of the invention. In the weight average branching degree calculation the weight of anionic surfactant components not having branched groups should also be included.

[0093] Suitable sulfate surfactants for use herein include water-soluble salts of C8-C18 alkyl or hydroxyalkyl, sulfate and/or ether sulfate. Suitable counterions include alkali metal cation or ammonium or substituted ammonium, but preferably sodium.

[0094] The sulfate surfactants may be selected from C8-C18 primary, branched chain and random alkyl sulfates (AS); C8-C18 secondary (2,3) alkyl sulfates; C8-C18 alkyl alkoxy sulfates (AExS) wherein preferably x is from 1-30 in which the alkoxy group could be selected from ethoxy, propoxy, butoxy or even higher alkoxy groups and mixtures thereof.

[0095] Alkyl sulfates and alkyl alkoxy sulfates are commercially available with a variety of chain lengths, ethoxylation and branching degrees. Commercially available sulfates include, those based on Neodol alcohols ex the Shell company, Lial - Isalchem and Safol ex the Sasol company, natural alcohols ex The Procter & Gamble Chemicals company.

[0096] Preferred alkyl sulfates are those in which the anionic surfactant is an alkyl ethoxy sulfate with a degree of ethoxylation of from about 0.2 to about 3, more preferably from about 0.3 to about 2, even more preferably from about 0.4 to about 1.5, and especially from about 0.4 to about 1. They are also preferred anionic surfactant having a level of branching of from about 5% to about 40%, even more preferably from about 10% to 35% and especially from about 20% to 30%.

Nonionic surfactant

[0097] Preferably the surfactant system comprises a nonionic surfactant, in an amount of from 0.1% to 40%, preferably 0.2% to 20%, most preferably 0.5% to 10% by weight of the composition. Suitable nonionic surfactants include the condensation products of aliphatic alcohols with from 1 to 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 8 to 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from 10 to 18 carbon atoms, preferably from 10 to 15 carbon atoms with from 2 to 18 moles, preferably 2 to 15, more preferably 5-12 of ethylene oxide per mole of alcohol. Highly preferred nonionic surfactants are the condensation products of guerbet alcohols with from 2 to 18 moles, preferably 2 to 15, more preferably 5-12 of ethylene oxide per mole of alcohol.

[0098] Other suitable non-ionic surfactants for use herein include fatty alcohol polyglycol ethers, alkylpolyglucosides and fatty acid glucamides.


Amphoteric surfactant

[0099] The surfactant system may include amphoteric surfactant, such as amine oxide. Preferred amine oxides are alkyl dimethyl amine oxide or alkyl amido propyl dimethyl amine oxide, more preferably alkyl dimethyl amine oxide and especially coco dimethyl amino oxide. Amine oxide may have a linear or mid-branched alkyl moiety. Typical linear amine oxides include water-soluble amine oxides containing one R1 C8-18 alkyl moiety and 2 R2 and R3 moieties selected from the group consisting of C1-3 alkyl groups and C1-3 hydroxyalkyl groups. Preferably amine oxide is characterized by the formula $R1 - N(R2)(R3)$ wherein R1 is a C8-18 alkyl and R2 and R3 are selected from the group consisting of methyl, ethyl, propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl and 3-hydroxypropyl. The linear amine oxide surfactants in particular may include linear C10-C18 alkyl dimethyl amine oxides and linear C8-C12 alkoxy ethyl dihydroxy ethyl amine oxides. Preferred amine oxides include linear C10, linear C10-C12, and linear C12-C14 alkyl dimethyl amine oxides. As used herein "mid-branched" means that the amine oxide has one alkyl moiety having n1 carbon atoms with one alkyl branch on the alkyl moiety having n2 carbon atoms. The alkyl branch is located on the α carbon from the nitrogen on the alkyl moiety. This type of branching for the amine oxide is also known in the art as an internal amine oxide. The total sum of n1 and n2 is from 10 to 24 carbon atoms, preferably from 12 to 20, and more preferably from 10 to 16. The number of carbon atoms for the one alkyl moiety (n1) should be approximately the same number of carbon atoms as the one alkyl branch (n2) such that the one alkyl moiety and the one alkyl branch are symmetric. As used herein "symmetric" means that $|n1 - n2|$ is less than or equal to 5, preferably 4, most preferably from 0 to 4 carbon atoms in at least 50 wt%, more preferably at least 75 wt% to 100 wt% of the mid-branched amine oxides for use herein.

[0100] The amine oxide may further comprise two moieties, independently selected from a C1-3 alkyl, a C1-3 hydroxyalkyl group, or a polyethylene oxide group containing an average of from about 1 to about 3 ethylene oxide groups. Preferably the two moieties are selected from a C1-3 alkyl, more preferably both are selected as a C 1 alkyl.

Zwitterionic surfactant

[0101] Other suitable surfactants include betaines, such as alkyl betaines, alkylamidobetaine, amidazoliniumbetaine, sulfobetaine (INCI Sultaines) as well as the Phosphobetaine and preferably meets formula (I):



wherein

R¹ is a saturated or unsaturated C6-22 alkyl residue, preferably C8-18 alkyl residue, in particular a saturated C10-16 alkyl residue, for example a saturated C12-14 alkyl residue;

X is NH, NR⁴ with C1-4 Alkyl residue R⁴, O or S,

n a number from 1 to 10, preferably 2 to 5, in particular 3,

x 0 or 1, preferably 1,





R², R³ are independently a C1-4 alkyl residue, potentially hydroxy substituted such as a hydroxyethyl, preferably a methyl.

m a number from 1 to 4, in particular 1, 2 or 3,

y 0 or 1 and

Y is COO, SO₃, OPO(OR⁵)O or P(O)(OR⁵)O, whereby R⁵ is a hydrogen atom H or a C1-4 alkyl residue.

[0102] Preferred betaines are the alkyl betaines of the formula (Ia), the alkyl amido propyl betaine of the formula (Ib), the Sulfo betaines of the formula (Ic) and the Amido sulfobetaine of the formula (Id);



in which R¹ as the same meaning as in formula I. Particularly preferred betaines are the Carbobetaine [wherein Y = COO⁻], in particular the Carbobetaine of the formula (Ia) and (Ib), more preferred are the Alkylamidobetaine of the formula (Ib).

[0103] Examples of suitable betaines and sulfobetaine are the following [designated in accordance with INCI]: Almondamidopropyl of betaines, Apricotamidopropyl betaines, Avocamidopropyl of betaines, Babassamidopropyl of betaines, Behenam idopropyl betaines, Behenyl of betaines, betaines, Canolamidopropyl betaines, Capryl/Capram idopropyl betaines, Carnitine, Cetyl of betaines, Cocamidoethyl of betaines, Cocamidopropyl betaines, Cocamidopropyl Hydroxysultaine, Coco betaines, Coco Hydroxysultaine, Coco/Oleamidopropyl betaines, Coco Sultaine, Decyl of betaines, Dihydroxyethyl Oleyl Glycinate, Dihydroxyethyl Soy Glycinate, Dihydroxyethyl Stearyl Glycinate, Dihydroxyethyl Tallow Glycinate, Dimethicone Propyl of PG-betaines, Erucamidopropyl Hydroxysultaine, Hydrogenated Tallow of betaines, Isostearamidopropyl betaines, Lauramidopropyl betaines, Lauryl of betaines, Lauryl Hydroxysultaine, Lauryl Sultaine, Milkamidopropyl betaines, Minkamidopropyl of betaines, Myristamidopropyl betaines, Myristyl of betaines, Oleamidopropyl betaines, Oleamidopropyl Hydroxysultaine, Oleyl of betaines, Olivamidopropyl of betaines, Palmamidopropyl betaines, Palm itamidopropyl betaines, Palmitoyl Carnitine, Palm Kernelamidopropyl betaines, Polytetrafluoroethylene Acetoxypromyl of betaines, Ricinoleamidopropyl betaines, Sesamidopropyl betaines, Soyamidopropyl betaines, Stearamidopropyl betaines, Stearyl of betaines, Tallowamidopropyl betaines, Tallowamidopropyl Hydroxysultaine, Tallow of betaines, Tallow Dihydroxyethyl of betaines, Undecylenamidopropyl betaines and Wheat Germamidopropyl betaines. A preferred betaine is, for example, Cocoamidopropylbetaine.

Fatty Acid

[0104] Especially when in liquid form, preferably, the detergent composition comprises between 1.5% and 20%, more preferably between 2% and 15%, even more preferably between 3% and 10%, most preferably between 4% and 8% by weight of the liquid detergent composition of soap, preferably a fatty acid salt, more preferably an amine neutralized fatty acid salt, wherein preferably the amine is an alkanolamine more preferably selected from monoethanolamine, diethanolamine, triethanolamine or a mixture thereof, more preferably monoethanolamine.

Perfume

[0105] Preferred compositions of the invention comprise perfume. Typically the composition comprises a perfume that comprises one or more perfume raw materials, selected from the group as described in WO08/87497. However, any

perfume useful in a detergent may be used. A preferred method of incorporating perfume into the compositions of the invention is via an encapsulated perfume particle comprising either a water-soluble hydroxylic compound or melamine-formaldehyde or modified polyvinyl alcohol. In one aspect the encapsulate comprises (a) an at least partially water-soluble solid matrix comprising one or more water-soluble hydroxylic compounds, preferably starch; and (b) a perfume oil encapsulated by the solid matrix. In a further aspect the perfume may be pre-complexed with a polyamine, preferably a polyethylenimine so as to form a Schiff base.


Polymers

[0106] The detergent composition may comprise one or more polymers for example for cleaning and/or care. Examples are optionally modified carboxymethylcellulose, poly (ethylene glycol), poly(vinyl alcohol), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid co-polymers and carboxylate polymers.

[0107] Suitable carboxylate polymers include maleate/acrylate random copolymer or polyacrylate homopolymer. The carboxylate polymer may be a polyacrylate homopolymer having a molecular weight of from 4,000 Da to 9,000 Da, or from 6,000 Da to 9,000 Da. Other suitable carboxylate polymers are co-polymers of maleic acid and acrylic acid, and may have a molecular weight in the range of from 4,000 Da to 90,000 Da.


[0108] Other suitable carboxylate polymers are co-polymers comprising: (i) from 50 to less than 98 wt% structural units derived from one or more monomers comprising carboxyl groups; (ii) from 1 to less than 49 wt% structural units derived from one or more monomers comprising sulfonate moieties; and (iii) from 1 to 49 wt% structural units derived from one or more types of monomers selected from ether bond-containing monomers represented by formulas (I) and (II):

formula (I):



wherein in formula (I), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5 provided X represents a number 1-5 when R is a single bond, and R_1 is a hydrogen atom or C1 to C20 organic group;

formula (II)



in formula (II), R_0 represents a hydrogen atom or CH_3 group, R represents a CH_2 group, CH_2CH_2 group or single bond, X represents a number 0-5, and R_1 is a hydrogen atom or C1 to C20 organic group.

[0109] The composition may comprise one or more amphiphilic cleaning polymers such as the compound having the

following general structure: bis((C₂H₅O)(C₂H₄O)_n)(CH₃)-N⁺-C_xH_{2x}-N⁺-(CH₃)-bis((C₂H₅O)(C₂H₄O)_n), wherein n = from 20 to 30, and x = from 3 to 8, or sulphated or sulphonated variants thereof. In one aspect, this polymer is sulphated or sulphonated to provide a zwitterionic soil suspension polymer.

[0110] The composition preferably comprises amphiphilic alkoxyated grease cleaning polymers which have balanced hydrophilic and properties such that they remove grease particles from fabrics and surfaces. Preferred amphiphilic alkoxyated grease cleaning polymers comprise a core structure and a plurality of alkoxyate groups attached to that core structure. These may comprise alkoxyated polyalkylenimines, preferably having an inner polyethylene oxide block and an outer polypropylene oxide block. Typically these may be incorporated into the compositions of the invention in amounts of from 0.005 to 10 wt%, generally from 0.5 to 8 wt%.

[0111] Alkoxyated polycarboxylates such as those prepared from polyacrylates are useful herein to provide additional grease removal performance. Such materials are described in WO 91/08281 and PCT 90/01815. Chemically, these materials comprise polyacrylates having one ethoxy side-chain per every 7-8 acrylate units. The side-chains are of the formula -(CH₂CH₂O)_m(CH₂)_nCH₃ wherein m is 2-3 and n is 6-12. The side-chains are ester-linked to the polyacrylate "backbone" to provide a "comb" polymer type structure. The molecular weight can vary, but is typically in the range of about 2000 to about 50,000. Such alkoxyated polycarboxylates can comprise from about 0.05% to about 10%, by weight, of the compositions herein.

[0112] The composition may comprise polyethylene glycol polymers and these may be particularly preferred in compositions comprising mixed surfactant systems. Suitable polyethylene glycol polymers include random graft co-polymers comprising: (i) hydrophilic backbone comprising polyethylene glycol; and (ii) side chain(s) selected from the group consisting of: C4-C25 alkyl group, polypropylene, polybutylene, vinyl ester of a saturated C1-C6 mono-carboxylic acid, C1-C6 alkyl ester of acrylic or methacrylic acid, and mixtures thereof. Suitable polyethylene glycol polymers have a polyethylene glycol backbone with random grafted polyvinyl acetate side chains. The average molecular weight of the polyethylene glycol backbone can be in the range of from 2,000 Da to 20,000 Da, or from 4,000 Da to 8,000 Da. The molecular weight ratio of the polyethylene glycol backbone to the polyvinyl acetate side chains can be in the range of from 1:1 to 1:5, or from 1:1.2 to 1:2. The average number of graft sites per ethylene oxide units can be less than 1, or less than 0.8, the average number of graft sites per ethylene oxide units can be in the range of from 0.5 to 0.9, or the average number of graft sites per ethylene oxide units can be in the range of from 0.1 to 0.5, or from 0.2 to 0.4. A suitable polyethylene glycol polymer is Sokalan HP22.

[0113] Typically these polymers when present are each incorporated into the compositions of the invention in amounts from 0.005 to 10 wt%, more usually from 0.05 to 8 wt%.

[0114] Preferably the composition comprises one or more carboxylate polymer, such as a maleate/acrylate random copolymer or polyacrylate homopolymer. In one aspect, the carboxylate polymer is a polyacrylate homopolymer having a molecular weight of from 4,000 Da to 9,000 Da, or from 6,000 Da to 9,000 Da. Typically these are incorporated into the compositions of the invention in amounts from 0.005 to 10 wt%, or from 0.05 to 8 wt%.

[0115] Preferably the composition comprises one or more soil release polymers.

[0116] Suitable soil release polymers are polyester soil release polymers such as Repel-o-tex polymers, including Repel-o-tex SF, SF-2 and SRP6 supplied by Rhodia. Other suitable soil release polymers include Texcare polymers, including Texcare SRA100, SRA300, SRN100, SRN170, SRN240, SRN260, SRN300 and SRN325 supplied by Clariant. Other suitable soil release polymers are Marloquest polymers, such as Marloquest SL supplied by Sasol.

[0117] Preferably the composition comprises one or more cellulosic polymer, including those selected from alkyl cellulose, alkyl alkoxyalkyl cellulose, carboxyalkyl cellulose, alkyl carboxyalkyl cellulose. Preferred cellulosic polymers are selected from the group comprising carboxymethyl cellulose, methyl cellulose, methyl hydroxyethyl cellulose, methyl carboxymethyl cellulose, and mixtures thereof. In one aspect, the carboxymethyl cellulose has a degree of carboxymethyl substitution from 0.5 to 0.9 and a molecular weight from 100,000 Da to 300,000 Da.

[0118] The composition preferably comprises a cationically-modified polysaccharide polymer. Preferably, the cationic polysaccharide polymer is selected from cationically modified hydroxyethyl cellulose, cationically modified hydroxypropyl cellulose, cationically and hydrophobically modified hydroxyethyl cellulose, cationically and hydrophobically modified hydroxypropyl cellulose, or a mixture thereof, more preferably cationically modified hydroxyethyl cellulose, cationically and hydrophobically modified hydroxyethyl cellulose, or a mixture thereof.


Amines

[0119] The cleaning compositions described herein may contain an amine. The cleaning compositions may include from about 0.1% to about 10%, or from about 0.2% to about 5%, or from about 0.5% to about 4%, or from about 0.1 % to about 4%, or from about 0.1 % to about 2%, by weight of the composition, of an amine. The amine can be subjected to protonation depending on the pH of the cleaning medium in which it is used. Non-limiting examples of amines include, but are not limited to, etheramines, cyclic amines, polyamines, oligoamines (e.g., triamines, diamines, pentamines, tetraamines), or combinations thereof. The compositions described herein may comprise an amine selected from the

group consisting of oligoamines, etheramines, cyclic amines, and combinations thereof. In some aspects, the amine is not an alkanolamine. In some aspects, the amine is not a polyalkyleneimine. Examples of suitable oligoamines include tetraethylenepentamine, triethylenetetraamine, diethylenetriamine, and mixtures thereof. Etheramines and cyclic amines may be particularly preferred.

Fabric Shading Dye

[0120] The composition may comprise a fabric shading agent. Suitable fabric shading agents include dyes, dye-clay conjugates, and pigments. Suitable dyes include small molecule dyes and polymeric dyes. Suitable small molecule dyes include small molecule dyes selected from the group consisting of dyes falling into the Colour Index (C.I.) classifications of Direct Blue, Direct Red, Direct Violet, Acid Blue, Acid Red, Acid Violet, Basic Blue, Basic Violet and Basic Red, or mixtures thereof. Preferred dyes include alkoxyated azothiophenes, Solvent Violet 13, Acid Violet 50 and Direct Violet 9. Particularly preferred dyes are polymeric dyes, particularly comprising polyalkoxy, most preferably polyethoxy groups, for example:



wherein the index values x and y are independently selected from 1 to 10.

Dye Transfer Inhibitors

[0121] Suitable dye transfer inhibitors include polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone, polyvinylloxazolidone, polyvinylimidazole and mixtures thereof. Preferred are poly(vinyl pyrrolidone), poly(vinylpyridine betaine), poly(vinylpyridine N-oxide), poly(vinyl pyrrolidone-vinyl imidazole) and mixtures thereof. Suitable commercially available dye transfer inhibitors include PVP-K15 and K30 (Ashland), Sokalan® HP165, HP50, HP53, HP59, HP56K, HP56, HP66 (BASF), Chromabond® S-400, S403E and S-100 (Ashland).

Chelant

[0122] The composition may comprise chelant for example selected from phosphonic, sulphonic, succinic and acetic chelants or mixtures thereof. Suitable examples include HEDP, DTPA, EDTA, MGDA, GLDA, EDDS and 4,5-dihydroxy-1,3-benzenedisulfonic acids and salts thereof.

Encapsulated Benefit Agent

[0123] The composition may further comprise an encapsulated benefit agent. The encapsulated benefit may comprise a shell surrounding a core. The core may comprise a benefit agent. The benefit agent may comprise perfume raw materials.

[0124] The shell may comprise a material selected from the group consisting of aminoplast copolymer, an acrylic, an acrylate, and mixtures thereof. The aminoplast copolymer may be melamine-formaldehyde, urea-formaldehyde, cross-linked melamine formaldehyde, or mixtures thereof.

[0125] The shell may be coated with one or more materials, such as a polymer, that aids in the deposition and/or retention of the perfume microcapsule on the site that is treated with the composition disclosed herein. The polymer may be a cationic polymer selected from the group consisting of polysaccharides, cationically modified starch, cationically modified guar, polysiloxanes, poly diallyl dimethyl ammonium halides, copolymers of poly diallyl dimethyl ammonium chloride and vinyl pyrrolidone, acrylamides, imidazoles, imidazolium halides, imidazolium halides, poly vinyl amine, copolymers of poly vinyl amine and N-vinyl formamide, and mixtures thereof.

[0126] The core may comprise a benefit agent. Suitable benefit agents include a material selected from the group consisting of perfume raw materials, silicone oils, waxes, hydrocarbons, higher fatty acids, essential oils, lipids, skin coolants, vitamins, sunscreens, antioxidants, glycerine, catalysts, bleach particles, silicon dioxide particles, malodor reducing agents, odor-controlling materials, chelating agents, antistatic agents, softening agents, insect and moth repelling agents, colorants, antioxidants, chelants, bodying agents, drape and form control agents, smoothness agents, wrinkle control agents, sanitization agents, disinfecting agents, germ control agents, mold control agents, mildew control

agents, antiviral agents, drying agents, stain resistance agents, soil release agents, fabric refreshing agents and freshness extending agents, chlorine bleach odor control agents, dye fixatives, dye transfer inhibitors, color maintenance agents, optical brighteners, color restoration/rejuvenation agents, anti-fading agents, whiteness enhancers, anti-abrasion agents, wear resistance agents, fabric integrity agents, anti-wear agents, anti-pilling agents, defoamers, anti-foaming agents, UV protection agents, sun fade inhibitors, anti-allergenic agents, enzymes, water proofing agents, fabric comfort agents, shrinkage resistance agents, stretch resistance agents, stretch recovery agents, skin care agents, glycerin, and natural actives, antibacterial actives, antiperspirant actives, cationic polymers, dyes and mixtures thereof. The benefit agent may comprise perfume raw materials.

[0127] The composition may comprise, based on total composition weight, from about 0.01% to about 10%, or from about 0.1% to about 5%, or from about 0.2% to about 1%, of encapsulated benefit agent. The encapsulated benefit agent may be friable and/or have a mean particle size of from about 10 microns to about 500 microns or from about 20 microns to about 200 microns.

[0128] Suitable encapsulated benefit agents may be obtained from Encapsys, LLC, of Appleton, Wisconsin USA.

[0129] Formaldehyde scavengers may also be used in or with such encapsulated benefit agents.

[0130] In a further preferred aspect of the invention, the composition is preferably liquid and comprises particulate benefit agents such as the encapsulated benefit agents mentioned above. The combination of the galactanase enzyme in addition to the plant fiber and particulate benefit agent has been found to provide the additional benefit of enhanced deposition of the particulate benefit agent. Thus, the present invention also provides a method of enhancing deposition of a particulate benefit agent comprising contacting a textile with an aqueous liquor comprising a composition defined herein, comprising a galactanase enzyme and a plant fiber and in addition a particulate benefit agent in a textile treatment step, preferably a laundering step, and optionally rinsing and drying the textile. In a preferred method, the aqueous liquor is an aqueous wash liquor. In a preferred method, the particulate benefit agent comprises an encapsulated perfume particle, most preferably comprising a shell which comprises a material selected from the group consisting of polymers or copolymers comprising acrylic acid and/or acrylates, and mixtures thereof.

Methods of Making the Composition

Methods of Making the Composition

[0131] The present invention relates to methods of making the compositions described herein. The compositions of the invention may be solid (for example granules or tablets) or liquid form. Preferably the compositions are in liquid form. They may be made by any process chosen by the formulator, including by a batch process, a continuous loop process, or combinations thereof.

[0132] When in the form of a liquid, the compositions of the invention may be aqueous (typically above 2 wt% or even above 5 or 10 wt% total water, up to 90 or up to 80wt% or 70 wt% total water) or non-aqueous (typically below 2 wt% total water content). Typically the compositions of the invention will be in the form of an aqueous solution or uniform dispersion or suspension of optical brightener, DTI and optional additional adjunct materials, some of which may normally be in solid form, that have been combined with the normally liquid components of the composition, such as the liquid alcohol ethoxylate nonionic, the aqueous liquid carrier, and any other normally liquid optional ingredients. Such a solution, dispersion or suspension will be acceptably phase stable. When in the form of a liquid, the detergents of the invention preferably have viscosity from 1 to 1500 centipoises (1-1500 mPa*s), more preferably from 100 to 1000 centipoises (100-1000 mPa*s), and most preferably from 200 to 500 centipoises (200-500 mPa*s) at 20s-1 and 21°C. Viscosity can be determined by conventional methods. Viscosity may be measured using an AR 550 rheometer from TA instruments using a plate steel spindle at 40 mm diameter and a gap size of 500 μm. The high shear viscosity at 20s-1 and low shear viscosity at 0.05-1 can be obtained from a logarithmic shear rate sweep from 0.1-1 to 25-1 in 3 minutes time at 21C. The preferred rheology described therein may be achieved using internal existing structuring with detergent ingredients or by employing an external rheology modifier. More preferably the detergents, such as detergent liquid compositions have a high shear rate viscosity of from about 100 centipoise to 1500 centipoise, more preferably from 100 to 1000 cps. Unit Dose detergents, such as detergent liquid compositions have high shear rate viscosity of from 400 to 1000cps. Detergents such as laundry softening compositions typically have high shear rate viscosity of from 10 to 1000, more preferably from 10 to 800 cps, most preferably from 10 to 500 cps. Hand dishwashing compositions have high shear rate viscosity of from 300 to 4000 cps, more preferably 300 to 1000 cps.

[0133] The cleaning and/or treatment compositions in the form of a liquid herein can be prepared by combining the components thereof in any convenient order and by mixing, e.g., agitating, the resulting component combination to form a phase stable liquid detergent composition. In a process for preparing such compositions, a liquid matrix is formed containing at least a major proportion, or even substantially all, of the liquid components, e.g., nonionic surfactant, the non-surface active liquid carriers and other optional liquid components, with the liquid components being thoroughly admixed by imparting shear agitation to this liquid combination. For example, rapid stirring with a mechanical stirrer may

usefully be employed. While shear agitation is maintained, substantially all of any anionic surfactants and the solid form ingredients can be added. Agitation of the mixture is continued, and if necessary, can be increased at this point to form a solution or a uniform dispersion of insoluble solid phase particulates within the liquid phase. After some or all of the solid-form materials have been added to this agitated mixture, particles of any enzyme material to be included, e.g., enzyme granulates, are incorporated. As a variation of the composition preparation procedure hereinbefore described, one or more of the solid components may be added to the agitated mixture as a solution or slurry of particles premixed with a minor portion of one or more of the liquid components. After addition of all of the composition components, agitation of the mixture is continued for a period of time sufficient to form compositions having the requisite viscosity and phase stability characteristics. Frequently this will involve agitation for a period of from about 30 to 60 minutes.

[0134] The adjunct ingredients in the compositions of this invention may be incorporated into the composition as the product of the synthesis generating such components, either with or without an intermediate purification step. Where there is no purification step, commonly the mixture used will comprise the desired component or mixtures thereof (and percentages given herein relate to the weight percent of the component itself unless otherwise specified) and in addition unreacted starting materials and impurities formed from side reactions and/or incomplete reaction. For example, for an ethoxylated or substituted component, the mixture will likely comprise different degrees of ethoxylation/substitution.

Method of Use

[0135] The present invention relates to methods of using the cleaning compositions of the present invention to clean a surface, such as a textile. In general, the method includes mixing the cleaning composition as described herein with water to form an aqueous liquor and contacting a surface, preferably a textile, with the aqueous liquor in a laundering step. The target surface may include a greasy soil such as a body soil. The compositions herein, typically prepared as hereinbefore described, can be used to form aqueous washing/treatment solutions for use in the laundering/treatment of fabrics and/or hard surfaces. Generally, an effective amount of such a composition is added to water, for example in a conventional fabric automatic washing machine, to form such aqueous liquor laundering solutions. The aqueous liquor so formed is then contacted, typically under agitation, with the fabrics to be laundered/treated therewith. An effective amount of the cleaning composition herein added to water to form aqueous liquors for washing can comprise amounts sufficient to form from about 500 to 25,000 ppm, or from 500 to 15,000 ppm of composition in aqueous liquor, or from about 1,000 to 3,000 ppm of the cleaning compositions herein will be provided in aqueous liquor.

[0136] Typically, the aqueous liquor is formed by contacting the detergent with wash water in such an amount so that the concentration of the cleaning composition in the aqueous liquor is from above 0.1 g/l to 5g/l, or from 1g/l, and to 4.5g/l, or to 4.0g/l, or to 3.5g/l, or to 3.0g/l, or to 2.5g/l, or even to 2.0g/l, or even to 1.5g/l. The method of laundering fabric or textile may be carried out in a top-loading or front-loading automatic washing machine, or can be used in a hand-wash laundry application. In these applications, the aqueous liquor formed and concentration of laundry detergent composition in the aqueous liquor is that of the main wash cycle. Any input of water during any optional rinsing step(s) is not included when determining the volume of the aqueous liquor.

[0137] The aqueous liquor may comprise 40 litres or less of water, or 30 litres or less, or 20 litres or less, or 10 litres or less, or 8 litres or less, or even 6 litres or less of water. The wash liquor may comprise from above 0 to 15 litres, or from 2 litres, and to 12 litres, or even to 8 litres of water. Typically from 0.01kg to 2kg of fabric per litre of aqueous liquor is dosed into said aqueous liquor. Typically from 0.01kg, or from 0.05kg, or from 0.07kg, or from 0.10kg, or from 0.15kg, or from 0.20kg, or from 0.25kg fabric per litre of aqueous liquor is dosed into said aqueous liquor. Optionally, 50g or less, or 45g or less, or 40g or less, or 35g or less, or 30g or less, or 25g or less, or 20g or less, or even 15g or less, or even 10g or less of the composition is contacted to water to form the aqueous liquor. Such compositions are typically employed at concentrations of from about 500 ppm to about 15,000 ppm in solution. When the wash solvent is water, the water temperature typically ranges from about 5 °C to about 90 °C and, when the situs comprises a fabric, the water to fabric ratio is typically from about 1:1 to about 30:1. Typically the aqueous liquor comprising the detergent of the invention has a pH of from 3 to 11.5.

[0138] In one aspect, such method comprises the steps of optionally washing and/or rinsing said surface or fabric, contacting said surface or fabric with any composition disclosed in this specification then optionally washing and/or rinsing said surface or fabric is disclosed, with an optional drying step.

[0139] Drying of such surfaces or fabrics may be accomplished by any one of the common means employed either in domestic or industrial settings: machine drying or open-air drying. The fabric may comprise any fabric capable of being laundered in normal consumer or institutional use conditions, and the invention is particularly suitable for synthetic textiles such as polyester and nylon and especially for treatment of mixed fabrics and/or fibres comprising synthetic and cellulosic fabrics and/or fibres. As examples of synthetic fabrics are polyester, nylon, these may be present in mixtures with cellulosic fibres, for example, polycotton fabrics. The solution typically has a pH of from 7 to 11, more usually 8 to 10.5. The compositions are typically employed at concentrations from 500 ppm to 5,000 ppm in solution. The water temperatures typically range from about 5 °C to about 90 °C. The water to fabric ratio is typically from about 1:1 to about

30:1.

Use of an Alkoxylated Phenol

5 **[0140]** The present invention further relates to a use of an alkoxylated polyaryl phenol and/or an alkoxylated polyalkyl phenol in a cleaning composition to enhance the stain-removal, whiteness, and/or malodor-reducing benefits of a galactanase enzyme.

EXAMPLES

10 **[0141]** The following are illustrative examples of cleaning compositions according to the present invention and are not intended to be limiting.

Examples 1-7

15 **[0142]** Heavy Duty Liquid laundry detergent compositions.

Ingredients	1	2	3	4	5	6	7
	% weight						
AE _{1.8} S	6.77	5.16	1.36	1.30	-	-	-
AE ₃ S	-	-	-	-	0.45	-	-
LAS	0.86	2.06	2.72	0.68	0.95	1.56	3.55
HSAS	1.85	2.63	1.02	-	-	-	-
AE9	6.32	9.85	10.20	7.92			
AE8							35.45
AE7					8.40	12.44	
C ₁₂₋₁₄ dimethyl Amine Oxide	0.30	0.73	0.23	0.37	-	-	-
C ₁₂₋₁₈ Fatty Acid	0.80	1.90	0.60	0.99	1.20	-	15.00
Citric Acid	2.50	3.96	1.88	1.98	0.90	2.50	0.60
Optical Brightener 1	1.00	0.80	0.10	0.30	0.05	0.50	0.001
Optical Brightener 3	0.001	0.05	0.01	0.20	0.50	-	1.00
Sodium formate	1.60	0.09	1.20	0.04	1.60	1.20	0.20
DTI 1	0.32	0.05	-	0.60	0.10	0.60	0.01
DTI 2	0.32	0.10	0.60	0.60	0.05	0.40	0.20
Sodium hydroxide	2.30	3.80	1.70	1.90	1.70	2.50	2.30
Monoethanolamine	1.40	1.49	1.00	0.70	-	-	-
Diethylene glycol	5.50	-	4.10	-	-	-	-
Chelant 1	0.15	0.15	0.11	0.07	0.50	0.11	0.80
4-formyl-phenylboronic acid	-	-	-	-	0.05	0.02	0.01
Sodium tetraborate	1.43	1.50	1.10	0.75	-	1.07	-
Ethanol	1.54	1.77	1.15	0.89	-	3.00	7.00
Polymer 1	0.10	-	-	-	-	-	2.00
Polymer 2	0.30	0.33	0.23	0.17	-	-	-
Polymer 3	-	-	-	-	-	-	0.80
Polymer 4	0.80	0.81	0.60	0.40	1.00	1.00	-

EP 3 330 352 A1

(continued)

Ingredients	1	2	3	4	5	6	7
	% weight						
1,2-Propanediol	-	6.60	-	3.30	0.50	2.00	8.00
Structurant	0.10	-	-	-	-	-	0.10
Perfume	1.60	1.10	1.00	0.80	0.90	1.50	1.60
Perfume encapsulate	0.10	0.05	0.01	0.02	0.10	0.05	0.10
Protease	0.80	0.60	0.70	0.90	0.70	0.60	1.50
Mannanase	0.07	0.05	-	0.06	0.04	0.045	-
Amylase 1	0.30	-	0.30	0.10	-	0.40	0.10
Amylase 2	-	0.20	0.10	0.15	0.07	-	0.10
Xyloglucannase	0.20	0.10	-	-	0.05	0.05	0.20
Lipase	0.40	0.20	0.30	0.10	0.20	-	-
Polishing enzyme	-	0.04	-	-	-	0.004	-
Galactanase	0.05	0.03	0.01	0.03	0.03	0.003	0.003
Dispersin B	-	-	-	0.05	0.03	0.001	0.001
Acid Violet 50	0.05	-	-	-	-	-	0.005
Direct Violet 9	-	-	-	-	-	0.05	-
Violet DD	-	0.035	0.02	0.037	0.04	-	-
Alkoxylated polyaryl/polyalkyl phenol	0.8	1.0	1.2	0.9	0.5	2.2	3.1
Water, dyes & minors	Balance						
pH	8.2						

Based on total cleaning and/or treatment composition weight. Enzyme levels are reported as raw material.

Examples 8 to 18: Unit Dose Compositions.

[0143] These examples provide various formulations for unit dose laundry detergents. Compositions 8 to 12 comprise a single unit dose compartment. The film used to encapsulate the compositions is a polyvinyl-alcohol-based film.

Ingredients	8	9	10	11	12
	% weight				
LAS	19.09	16.76	8.59	6.56	3.44
AE3S	1.91	0.74	0.18	0.46	0.07
AE7	14.00	17.50	26.33	28.08	31.59
Citric Acid	0.6	0.6	0.6	0.6	0.6
C12-15 Fatty Acid	14.8	14.8	14.8	14.8	14.8
Polymer 3	4.0	4.0	4.0	4.0	4.0
Chelant 2	1.2	1.2	1.2	1.2	1.2
Optical Brightener 1	0.20	0.25	0.01	0.01	0.50
Optical Brightener 2	0.20	-	0.25	0.03	0.01
Optical Brightener 3	0.18	0.09	0.30	0.01	-

EP 3 330 352 A1

(continued)

5

10

15

20

25

30

35

<u>Ingredients</u>	8	9	10	11	12
	% weight				
DTI 1	0.10	-	0.20	0.01	0.05
DTI 2	-	0.10	0.20	0.25	0.05
Glycerol	6.1	6.1	6.1	6.1	6.1
Monoethanol amine	8.0	8.0	8.0	8.0	8.0
Tri-isopropanol amine	-	-	2.0	-	-
Tri-ethanol amine	-	2.0	-	-	-
Cumene sulfonate	-	-	-	-	2.0
Protease	0.80	0.60	0.07	1.00	1.50
Mannanase	0.07	0.05	0.05	-	-
Amylase 1	0.20	0.11	0.30	0.50	0.05
Amylase 2	0.11	0.20	0.10	-	0.50
Polishing enzyme	0.005	0.05	-	-	-
Galactanase	0.005	0.05	0.005	0.010	0.005
Dispersin B	0.010	0.05	-	0.005	-
Cyclohexyl dimethanol	-	-	-	2.0	-
Acid violet 50	0.03	0.02			
Violet DD			0.01	0.05	0.02
Structurant	0.14	0.14	0.14	0.14	0.14
Perfume	1.9	1.9	1.9	1.9	1.9
Alkoxylated polyaryl/polyalkyl phenol	1.0	1.7	1.9	0.9	0.8
Water and miscellaneous	To 100%				
pH	7.5-8.2				

[0144] Based on total cleaning and/or treatment composition weight. Enzyme levels are reported as raw material.

40

[0145] In the following examples the unit dose has three compartments, but similar compositions can be made with two, four or five compartments. The film used to encapsulate the compartments is polyvinyl alcohol.

45

50

55

<u>Base compositions Ingredients</u>	13	14	15	16
	% weight			
HLAS	26.82	16.35	7.50	3.34
AE7	17.88	16.35	22.50	30.06
Citric Acid	0.5	0.7	0.6	0.5
C12-15 Fatty acid	16.4	6.0	11.0	13.0
Polymer 1	2.9	0.1	-	-
Polymer 3	1.1	5.1	2.5	4.2
Cationic cellulose polymer	-	-	0.3	0.5
Polymer 6	-	1.5	0.3	0.2
Chelant 2	1.1	2.0	0.6	1.5

EP 3 330 352 A1

(continued)

5

10

15

20

25

30

<u>Base compositions Ingredients</u>	13	14	15	16
	% weight			
Optical Brightener 1	0.20	0.25	0.01	0.005
Optical Brightener 3	0.18	0.09	0.30	0.005
DTI 1	0.1	-	0.2	-
DTI 2	-	0.1	0.2	-
Glycerol	5.3	5.0	5.0	4.2
Monoethanolamine	10.0	8.1	8.4	7.6
Polyethylene glycol	-	-	2.5	3.0
Potassium sulfite	0.2	0.3	0.5	0.7
Protease	0.80	0.60	0.40	0.80
Amylase 1	0.20	0.20	0.200	0.30
Polishing enzyme	-	-	0.005	0.005
Galactanase	0.05	0.010	0.005	0.005
Dispersin B	-	0.010	0.010	0.010
MgCl ₂	0.2	0.2	0.1	0.3
Structurant	0.2	0.1	0.2	0.2
Acid Violet 50	0.04	0.03	0.05	0.03
Perfume / encapsulates	0.10	0.30	0.01	0.05
Alkoxylated polyaryl/polyalkyl phenol	1.1	1.8	0.9	1.2
Solvents and misc.	To 100%			
pH	7.0-8.2			

35

40

45

50

<u>Finishing compositions</u>	17			18		
Compartment	A	B	C	A	B	C
Volume of each compartment	40 ml	5 ml	5 ml	40 ml	5 ml	5 ml
<u>Ingredients</u>	Active material in Wt. %					
Perfume	1.6	1.6	1.6	1.6	1.6	1.6
Violet DD	0	0.006	0	0	0.004	-
TiO ₂	-	-	0.1	-	-	0.1
Sodium Sulfite	0.4	0.4	0.4	0.3	0.3	0.3
Polymer 5	-	-	-	2	-	-
Hydrogenated castor oil	0.14	0.14	0.14	0.14	0.14	0.14
Base Composition 13, 14, 15 or 16	Add to 100%					

[0146] Based on total cleaning and/or treatment composition weight, enzyme levels are reported as raw material.

55

Examples 19 to 24

[0147] Granular laundry detergent compositions for hand washing or washing machines, typically top-loading washing

EP 3 330 352 A1

machines.

Ingredient	19	20	21	22	23	24
	% weight					
LAS	11.33	10.81	7.04	4.20	3.92	2.29
Quaternary ammonium	0.70	0.20	1.00	0.60	-	-
AE3S	0.51	0.49	0.32	-	0.08	0.10
AE7	8.36	11.50	12.54	11.20	16.00	21.51
Sodium Tripolyphosphate	5.0	-	4.0	9.0	2.0	-
Zeolite A	-	1.0	-	1.0	4.0	1.0
Sodium silicate 1.6R	7.0	5.0	2.0	3.0	3.0	5.0
Sodium carbonate	20.0	17.0	23.0	14.0	14.0	16.0
Polyacrylate MW 4500	1.0	0.6	1.0	1.0	1.5	1.0
Polymer 6	0.1	0.2	-	-	0.1	-
Carboxymethyl cellulose	1.0	0.3	1.0	1.0	1.0	1.0
Acid Violet 50	0.05	-	0.02	-	0.04	-
Violet DD	-	0.03	-	0.03	-	0.03
Protease 2	0.10	0.10	0.10	0.10	-	0.10
Amylase	0.03	-	0.03	0.03	0.03	0.03
Lipase	0.03	0.07	0.30	0.10	0.07	0.40
Polishing enzyme	0.002	-	0.05	-	0.02	-
Galactanase	0.001	0.001	0.01	0.05	0.002	0.02
Dispersin B	0.001	0.001	0.05	-	0.001	-
Optical Brightener 1	0.200	0.001	0.300	0.650	0.050	0.001
Optical Brightener 2	0.060	-	0.650	0.180	0.200	0.060
Optical Brightener 3	0.100	0.060	0.050	-	0.030	0.300
Chelant 1	0.60	0.80	0.60	0.25	0.60	0.60
DTI 1	0.32	0.15	0.15	-	0.10	0.10
DTI 2	0.32	0.15	0.30	0.30	0.10	0.20
Sodium Percarbonate	-	5.2	0.1	-	-	-
Sodium Perborate	4.4	-	3.85	2.09	0.78	3.63
Nonanoyloxybenzenesulfonate	1.9	0.0	1.66	0.0	0.33	0.75
Tetraacetylenediamine	0.58	1.2	0.51	0.0	0.015	0.28
Photobleach	0.0030	0.0	0.0012	0.0030	0.0021	-
S-ACMC	0.1	0.0	0.0	0.0	0.06	0.0
Alkoxylated polyaryl/ polyalkyl phenol	1.9	1.1	2.0	1.8	2.2	1.2
Sulfate/Moisture	Balance					

Examples 25-30

[0148] Granular laundry detergent compositions typically for front-loading automatic washing machines.

EP 3 330 352 A1

	<u>Ingredient</u>	25	26	27	28	29	30
		% weight					
5	LAS	6.08	5.05	4.27	3.24	2.30	1.09
	AE3S	-	0.90	0.21	0.18	-	0.06
	AS	0.34	-	-	-	-	-
10	AE7	4.28	5.95	6.72	7.98	9.20	10.35
	Quaternary ammonium	0.5	-	-	0.3	-	-
	Crystalline layered silicate	4.1	-	4.8	-	-	-
	Zeolite A	5.0	-	2.0	-	2.0	2.0
15	Citric acid	3.0	4.0	3.0	4.0	2.5	3.0
	Sodium carbonate	11.0	17.0	12.0	15.0	18.0	18.0
	Sodium silicate 2R	0.08	-	0.11	-	-	-
20	Optical Brightener 1	-	0.25	0.05	0.01	0.10	0.02
	Optical Brightener 2	-	-	0.25	0.20	0.01	0.08
	Optical Brightener 3	-	0.06	0.04	0.15	-	0.05
	DTI 1	0.08	-	0.04	-	0.10	0.01
25	DTI 2	0.08	-	0.04	0.10	0.10	0.02
	Soil release agent	0.75	0.72	0.71	0.72	-	-
	Acrylic /maleic acid copolymer	1.1	3.7	1.0	3.7	2.6	3.8
30	Carboxymethyl cellulose	0.2	1.4	0.2	1.4	1.0	0.5
	Protease 3	0.20	0.20	0.30	0.15	0.12	0.13
	Amylase 3	0.20	0.15	0.20	0.30	0.15	0.15
	Lipase	0.05	0.15	0.10	-	-	-
35	Amylase 2	0.03	0.07	-	-	0.05	0.05
	Cellulase 2	-	-	-	-	0.10	0.10
	Polishing enzyme	0.003	0.005	0.020	-	-	-
40	Galactanase	0.002	0.010	0.020	0.020	0.010	0.003
	Dispersin B	0.002	0.010	0.020	0.020	-	0.002
	Tetraacetylenehtylenediamine	3.6	4.0	3.6	4.0	2.2	1.4
45	Sodium percarbonate	13.0	13.2	13.0	13.2	16.0	14.0
	Chelant 3	-	0.2	-	0.2	-	0.2
	Chelant 2	0.2	-	0.2	-	0.2	0.2
	MgSO ₄	-	0.42	-	0.42	-	0.4
50	Perfume	0.5	0.6	0.5	0.6	0.6	0.6
	Suds suppressor agglomerate	0.05	0.10	0.05	0.10	0.06	0.05
	Soap	0.45	0.45	0.45	0.45	-	-
55	Acid Violet 50	0.04	-	0.05	-	0.04	-
	Violet DD	-	0.04	-	0.05	-	0.04
	S-ACMC	0.01	0.01	-	0.01	-	-

EP 3 330 352 A1

(continued)

<u>Ingredient</u>	25	26	27	28	29	30
	% weight					
Direct Violet 9 (active)	-	-	0.0001	0.0001	-	-
Alkoxyated polyaryl/ polyalkyl phenol	1.1	1.1	0.81	1.6	2.3	0.6
Sulfate/ Water & Miscellaneous	Balance					

Examples 31-37: Heavy Duty Liquid laundry detergent compositions.

[0149]

<u>Ingredients</u>	31	32	33	34	35	36	37
	% weight						
AE _{1,8} S	6.77	5.16	1.36	1.30	-	-	-
AE ₃ S	-	-	-	-	0.45	-	-
LAS	0.86	2.06	2.72	0.68	0.95	1.56	3.55
HSAS	1.85	2.63	1.02	-	-	-	-
AE9	6.32	9.85	10.20	7.92			
AE8							35.45
AE7					8.40	12.44	
C ₁₂₋₁₄ dimethyl Amine Oxide	0.30	0.73	0.23	0.37	-	-	-
C ₁₂₋₁₈ Fatty Acid	0.80	1.90	0.60	0.99	1.20	-	15.00
Citric Acid	2.50	3.96	1.88	1.98	0.90	2.50	0.60
Optical Brightener 1	1.00	0.80	0.10	0.30	0.05	0.50	0.001
Optical Brightener 3	0.001	0.05	0.01	0.20	0.50	-	1.00
Sodium formate	1.60	0.09	1.20	0.04	1.60	1.20	0.20
DTI 1	0.32	0.05	-	0.60	0.10	0.60	0.01
DTI 2	0.32	0.10	0.60	0.60	0.05	0.40	0.20
Sodium hydroxide	2.30	3.80	1.70	1.90	1.70	2.50	2.30
Monoethanolamine	1.40	1.49	1.00	0.70	-	-	-
Diethylene glycol	5.50	-	4.10	-	-	-	-
Chelant 1	0.15	0.15	0.11	0.07	0.50	0.11	0.80
4-formyl-phenylboronic acid	-	-	-	-	0.05	0.02	0.01
Sodium tetraborate	1.43	1.50	1.10	0.75	-	1.07	-
Ethanol	1.54	1.77	1.15	0.89	-	3.00	7.00
Polymer 1	0.10	-	-	-	-	-	2.00
Polymer 2	0.30	0.33	0.23	0.17	-	-	-
Polymer 3	-	-	-	-	-	-	0.80
Polymer 4	0.80	0.81	0.60	0.40	1.00	1.00	-
1,2-Propanediol	-	6.60	-	3.30	0.50	2.00	8.00
Structurant	0.10	-	-	-	-	-	0.10

EP 3 330 352 A1

(continued)

Ingredients	31	32	33	34	35	36	37
	% weight						
Perfume	1.60	1.10	1.00	0.80	0.90	1.50	1.60
Perfume encapsulate	0.10	0.05	0.01	0.02	0.10	0.05	0.10
Protease	0.80	0.60	0.70	0.90	0.70	0.60	1.50
Galactanase of any of SEQ ID Nos: 1-3	0.07	0.05	0.045	0.06	0.04	0.045	0.10
Amylase 1	0.30	-	0.30	0.10	-	0.40	0.10
Amylase 2	-	0.20	0.10	0.15	0.07	-	0.10
Xyloglucanase	0.20	0.10	-	-	0.05	0.05	0.20
Lipase	0.40	0.20	0.30	0.10	0.20	-	-
Polishing enzyme	-	0.04	-	-	-	0.004	-
Nuclease	0.05	0.03	0.01	0.03	0.03	0.003	0.003
Dispersin B	-	-	-	0.05	0.03	0.001	0.001
Acid Violet 50	0.05	-	-	-	-	-	0.005
Direct Violet 9	-	-	-	-	-	0.05	-
Violet DD	-	0.035	0.02	0.037	0.04	-	-
Water insoluble plant fiber	0.2	0.1	0.3	0.25	1.2	1.5	0.25
Dye control agent	-	0.3	-	0.5	-	0.3	-
Alkoxylated polyaryl/ polyalkyl phenol	0.8	1.1	1.2	1.5	0.85	1.1	3.1
Water, dyes & minors	Balance						
pH	8.2						

[0150] Based on total cleaning and/or treatment composition weight. Unless indicated otherwise, enzyme levels are reported as raw material.

AE1.8S	is C ₁₂₋₁₅ alkyl ethoxy sulfate with an average degree of ethoxylation of 1.8
AE3S	is C ₁₂₋₁₅ alkyl ethoxy sulfate with an av degree of ethoxylation of 3
AE7	is C ₁₂₋₁₃ alcohol ethoxylate, with an average degree of ethoxylation of 7
AE8	is C ₁₂₋₁₃ alcohol ethoxylate, with an average degree of ethoxylation of 8
AE9	is C ₁₂₋₁₃ alcohol ethoxylate, with an average degree of ethoxylation of 9
Alkoxylated polyaryl/ polyalkyl phenol	is alkoxylated polyaryl/polyalkyl phenol in accordance with the invention, for example Emulsogen® TS160, Hostapal® BV conc., Sapogenat® T110 or Sapogenat® T139, all from Clariant
Amylase 1	is Stainzyme®, 15 mg active/g
Amylase 2	is Natalase®, 29 mg active/g
Amylase 3	is Stainzyme Plus®, 20 mg active/g,
AS	is C ₁₂₋₁₄ alkylsulfate
Cellulase 2	is Celluclean™, 15.6 mg active/g
Xyloglucanase	is Whitezyme®, 20mg active/g
Chelant 1	is diethylene triamine pentaacetic acid
Chelant 2	is 1-hydroxyethane 1,1-diphosphonic acid
Chelant 3	is sodium salt of ethylenediamine-N,N'-disuccinic acid, (S,S) isomer (EDDS)
Dispersin B	is a glycoside hydrolase, reported as 1000mg active/g
DTI 1	is poly(4-vinylpyridine-1-oxide) (such as Chromabond S-403E®),
DTI 2	is poly(1-vinylpyrrolidone-co-1-vinylimidazole) (such as Sokalan HP56®).
Dye control agent	Dye control agent in accordance with the invention, for example Suparex®

EP 3 330 352 A1

	HSAS	O.IN (M1), Nylofixan® P (M2), Nylofixan® PM (M3), or Nylofixan® HF (M4)
	LAS	is mid-branched alkyl sulfate as disclosed in US 6,020,303 and US6,060,443
5	Galactanase	is linear alkylbenzenesulfonate having an average aliphatic carbon chain length C ₉ -C ₁₅ (HLAS is acid form).
	Lipase	is SEQ ID NO: 1, 2 and/or 3, as active protein.
	Mannanase	is Lipex®, 18 mg active/g
	Optical Brightener 1	is Mannaway®, 25 mg active/g
10	Optical Brightener 2	is disodium 4,4'-bis[[4-anilino-6-morpholino-s-triazin-2-yl]-amino]-2,2'-stilbenedisulfonate
	Optical Brightener 3	is disodium 4,4'-bis-(2-sulfostyryl)biphenyl (sodium salt)
	Perfume encapsulate	is Optiblanc SPL10® from 3V Sigma
	Photobleach	is a core-shell melamine formaldehyde perfume microcapsules.
	Polishing enzyme	is a sulfonated zinc phthalocyanine
15	Polyetheramine	is Para-nitrobenzyl esterase, reported as 1000mg active/g
	Polymer 1	as described in present disclosure.
	Polymer 2	is bis((C ₂ H ₅ O)(C ₂ H ₄ O) _n)(CH ₃)-N ⁺ -C _x H _{2x} -N ⁺ -(CH ₃)-bis((C ₂ H ₅ O)(C ₂ H ₄ O) _n),
	Polymer 3	wherein n = 20-30, x = 3 to 8 or sulphated or sulfonated variants thereof
20	Polymer 4	is ethoxylated (EO ₁₅) tetraethylene pentamine
	Polymer 5	is ethoxylated polyethylenimine
	Polymer 6	is ethoxylated hexamethylene diamine
	Protease	is Acusol 305, provided by Rohm&Haas
25	Protease 2	is a polyethylene glycol polymer grafted with vinyl acetate side chains, provided by BASF.
	Protease 3	is Purafect Prime®, 40.6 mg active/g
	Quaternary ammonium	is Savinase®, 32.89 mg active/g
	S-ACMC	is Purafect®, 84 mg active/g
	Soil release agent	is C ₁₂₋₁₄ Dimethylhydroxyethyl ammonium chloride
30	Structurant	is Reactive Blue 19 Azo-CM-Cellulose provided by Megazyme
	Violet DD	is Repel-o-tex® SF2
		is Hydrogenated Castor Oil
		is a thiophene azo dye provided by Milliken

[0151] The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm."

40

45

50

55

EP 3 330 352 A1

SEQUENCE LISTING

<110> P&G

5 <120> CLEANING COMPOSITIONS INCLUDING ENZYME AND BLEACH

<130> CM04649FM

<160> 14

10 <170> PatentIn version 3.5

<210> 1

<211> 463

<212> PRT

15 <213> Streptomyces davawensis

<400> 1

20 Asp Ala Thr Ile Val Ile Asn Pro Gly Thr Arg Tyr Gly Thr Trp Glu
1 5 10 15

Gly Trp Gly Thr Ser Leu Ala Trp Trp Gly Asn Val Phe Gly Thr Arg
20 25 30

25 Asp Asp Phe Ala Asp Leu Phe Phe Thr Thr Lys Ser Val Thr Tyr Asn
35 40 45

30 Gly Thr Ser Leu Pro Gly Leu Gly Leu Asn Ile Ala Arg Tyr Asn Leu
50 55 60

35 Gly Ala Cys Ser Trp Asn Ala Val Asn Gly Glu Thr Met Val Lys Ser
65 70 75 80

Pro Asn Ile Pro Ala Phe Lys Gln Ile Glu Gly Phe Trp Gln Asp Trp
85 90 95

40 Asn Asn Glu Asp Pro Thr Ser Ser Ala Trp Asp Trp Thr Ala Asp Ala
100 105 110

45 Thr Gln Arg Ala Met Leu Val Lys Ala Thr Gln Arg Gly Ala Val Thr
115 120 125

Glu Leu Phe Ala Asn Ser Pro Met Trp Trp Met Cys Tyr Asn His Asn
130 135 140

50 Pro Ser Gly Ala Ala Asp Gly Gly Asn Asn Leu Gln Thr Trp Asn Tyr
145 150 155 160

55 Arg Gln His Ala Ser His Leu Ala Ala Val Ala Leu Tyr Ala Arg Thr
165 170 175

EP 3 330 352 A1

Asn Trp Gly Val Asn Phe Ala Thr Val Asp Pro Phe Asn Glu Pro Ala
180 185 190

5 Ser Ser Trp Trp Thr Ala Ser Gly Thr Gln Glu Gly Cys His Leu Asp
195 200 205

10 Pro Ala Val Gln Ala Ala Val Leu Pro Tyr Met Arg Ser Glu Leu Asp
210 215 220

Lys Arg Gly Leu Thr Gly Val Arg Ile Ser Ala Ser Asp Glu Thr Asn
225 230 235 240

15 Tyr Asp Thr Ala Arg Ser Thr Trp Ser Ser Phe Gly Ser Ala Thr Lys
245 250 255

20 Ala Leu Val Ser Gln Val Asn Val His Gly Tyr Gln Gly Thr Gly Gly
260 265 270

Arg Arg Asp Leu Leu Tyr Thr Asp Val Val Thr Thr Ser Gly Lys Lys
275 280 285

25 Leu Trp Asn Ser Glu Thr Gly Asp Ser Asp Gly Thr Gly Leu Ser Met
290 295 300

30 Ala Arg Asn Leu Cys Tyr Asp Phe Arg Trp Leu His Pro Thr Ala Trp
305 310 315 320

35 Cys Tyr Trp Gln Val Met Asp Pro Ser Thr Gly Trp Ala Met Ile Ala
325 330 335

Tyr Asp Ala Asn Thr Leu Gln Pro Thr Thr Val Gln Pro Lys Tyr Tyr
340 345 350

40 Val Met Ala Gln Phe Ser Arg His Ile Arg Pro Gly Met Thr Ile Leu
355 360 365

45 Asp Thr Gly Val Ser Phe Ala Ala Ala Tyr Asp Ala Ser Ala Arg
370 375 380

Arg Leu Val Leu Val Ala Val Asn Thr Ser Thr Ser Pro Gln Thr Phe
385 390 395 400

50 Thr Phe Asp Leu Ser Arg Phe Thr Thr Val Thr Gly Gly Ser Gly Gly
405 410 415

55 Leu Val Pro Arg Trp Asn Thr Val Thr Gly Gly Gly Asp Met Tyr Arg
420 425 430

EP 3 330 352 A1

Ala Tyr Thr Asn Thr Tyr Val Thr Gly Lys Ser Val Ser Ala Thr Phe
435 440 445

5 Ala Ala Gly Ser Val Gln Thr Leu Gln Val Asp Gly Val Thr Thr
450 455 460

<210> 2
<211> 464
<212> PRT
<213> Streptomyces avermitilis

<400> 2

15 Asp Ala Thr Ile Ala Val Asn Pro Ser Thr Thr Tyr Gly Lys Trp Glu
1 5 10 15

20 Gly Trp Gly Thr Ser Leu Ala Trp Trp Ala Asn Val Phe Gly Ala Arg
20 25 30

25 Asp Asp Phe Ala Asp Leu Phe Phe Thr Thr Lys Ser Val Thr Tyr Asn
35 40 45

30 Gly Arg Thr Leu Pro Gly Leu Gly Leu Asn Ile Ala Arg Tyr Asn Leu
50 55 60

35 Ala Asn Ile Pro Ala Phe Lys Gln Ile Glu Gly Tyr Trp Gln Asp Trp
85 90 95

40 Asn Asn Glu Asp Pro Thr Ser Ser Ala Trp Lys Trp Thr Ala Asp Ala
100 105 110

45 Ala Gln Arg Thr Met Leu Val Lys Ala Thr Ala Arg Gly Ala Thr Thr
115 120 125

50 Glu Leu Phe Ala Asn Ser Pro Met Trp Trp Met Cys Leu Asn His Asn
130 135 140

55 Pro Ser Gly Ala Ser Gly Gly Gly Asn Asn Leu Gln Ser Trp Asn Tyr
145 150 155 160

Arg Gln His Ala Ser His Leu Ala Ala Val Ala Leu Tyr Ala Lys Ser
165 170 175

Asn Trp Gly Val Asn Phe Ala Thr Val Asp Pro Phe Asn Glu Pro Ser
180 185 190

EP 3 330 352 A1

Ser Ser Trp Trp Thr Ala Thr Gly Thr Gln Glu Gly Cys His Met Asp
 195 200 205

5
 Ala Ser Val Gln Ala Ala Val Leu Pro Tyr Leu Arg Ser Glu Leu Asp
 210 215 220

10
 Arg Arg Gly Leu Thr Gly Thr Lys Ile Ser Ala Ser Asp Glu Thr Ser
 225 230 235 240

15
 Tyr Asp Leu Ala Arg Thr Thr Trp Gly Ser Phe Gly Ser Ser Thr Lys
 245 250 255

20
 Ala Leu Val Asn Arg Val Asn Val His Gly Tyr Gln Gly Ser Gly Gly
 260 265 270

25
 Arg Arg Asp Leu Leu Tyr Thr Asp Val Val Thr Thr Ala Gly Lys Ala
 275 280 285

30
 Leu Trp Asn Ser Glu Thr Gly Asp Ser Asp Gly Thr Gly Leu Thr Leu
 290 295 300

35
 Ala Ser Asn Leu Cys Leu Asp Phe Arg Trp Leu His Pro Thr Ala Trp
 305 310 315 320

40
 Val Tyr Trp Gln Val Met Asp Pro Ser Ser Gly Trp Ala Met Ile Ala
 325 330 335

45
 Tyr Asp Ala Ser Thr Leu Gln Pro Gly Ala Val Gln Thr Lys Tyr Tyr
 340 345 350

50
 Val Met Ala Gln Phe Ser Arg His Ile Arg Ala Gly Met Thr Ile Val
 355 360 365

55
 Asp Thr Gly Val Gly Tyr Ala Ala Ala Ala Tyr Asp Ala Thr Ala Arg
 370 375 380

60
 Arg Leu Val Ile Val Ala Val Asn Thr Ser Thr Ser Ala Gln Thr Leu
 385 390 395 400

65
 Thr Phe Asp Leu Ser Arg Phe Ser Thr Val Thr Gly Gly Thr Gly Gly
 405 410 415

70
 Leu Val Arg Arg Trp Asn Thr Val Thr Gly Gly Gly Gly Asp Leu Tyr
 420 425 430

75
 Ala Ala His Ser Asp Thr Tyr Leu Ser Gly Lys Ser Leu Ser Val Pro
 435 440 445

EP 3 330 352 A1

Phe Ala Ala Gly Ala Val Gln Thr Leu Glu Val Asp Gly Val Thr Val
 450 455 460

5 <210> 3
 <211> 458
 <212> PRT
 <213> Trichoderma harzianum

10 <400> 3

Asp Thr Thr Leu Ser Ile Asp Pro Thr Ser Asn Trp Gly Thr Trp Glu
 1 5 10 15

15 Gly Trp Gly Val Ser Leu Ala Trp Trp Ala Lys Ala Phe Gly Asn Arg
 20 25 30

20 Asp Asp Leu Ala Asn Val Phe Phe Thr Arg Asn Asn Gln Val Ile Asn
 35 40 45

Gly Gln Asn Leu Pro Gly Leu Gly Phe Asn Ile Ala Arg Tyr Asn Ala
 50 55 60

25 Gly Ala Cys Ser Thr Asn Thr Tyr Asn Gly Ser Ser Met Val Val Ser
 65 70 75 80

30 Ser Ser Ile Lys Pro Ser Arg Gln Val Asp Gly Tyr Trp Leu Asp Trp
 85 90 95

35 Ala Ser Thr Asp Pro Ala Ser Ser Ser Trp Asn Trp Asn Val Asp Ala
 100 105 110

Asn Gln Arg Ala Met Leu Gln Lys Ala Lys Ala Asn Gly Ala Asn Ile
 115 120 125

40 Phe Glu Leu Phe Ser Asn Ser Pro Met Trp Trp Met Cys Leu Asn His
 130 135 140

45 Asn Pro Ser Gly Ser Gly Ser Ser Asp Asn Leu Gln Ser Trp Asn Tyr
 145 150 155 160

Gln Asn His Ala Val Tyr Leu Ala Asn Ile Ala Gln His Ala Gln Gln
 165 170 175

50 Asn Trp Gly Ile Gln Phe Gln Ser Val Glu Ala Phe Asn Glu Pro Ser
 180 185 190

55 Ser Gly Trp Gly Pro Thr Gly Thr Gln Glu Gly Cys His Phe Ala Val
 195 200 205

EP 3 330 352 A1

Ser Thr Met Ala Thr Val Ile Gly Tyr Leu Asn Thr Glu Leu Ala Gln
 210 215 220

5
 Arg Gly Leu Ser Ser Phe Ile Ser Ala Ser Asp Glu Thr Ser Tyr Asp
 225 230 235 240

10
 Leu Ala Ile Ser Thr Trp Gln Gly Leu Gly Ser Ser Ala Gln Asn Ala
 245 250 255

15
 Val Lys Arg Val Asn Val His Gly Tyr Gln Gly Gly Gly Gly Arg Arg
 260 265 270

20
 Asp Thr Leu Tyr Ser Leu Val Ser Gln Ala Gly Lys Arg Leu Trp Asn
 275 280 285

25
 Ser Glu Tyr Gly Asp Ala Asp Ala Ser Gly Lys Ser Met Tyr Thr Asn
 290 295 300

30
 Leu Leu Leu Asp Phe Thr Trp Leu His Pro Thr Ala Trp Val Tyr Trp
 305 310 315 320

35
 Gln Ala Ile Asp Gly Ser Gly Trp Gly Leu Ile Val Gly Asp Asn Asp
 325 330 335

40
 Gln Leu Thr Leu Ser Ser Ala Ser Thr Lys Tyr Phe Val Leu Ala Gln
 340 345 350

45
 Leu Thr Arg His Ile Arg Pro Gly Met Gln Ile Leu Thr Thr Pro Asp
 355 360 365

50
 Gly Asn Thr Val Ala Ala Tyr Asp Ser Gly Ser Gln Lys Leu Val Ile
 370 375 380

55
 Val Ala Ala Asn Trp Gly Ser Ala Gln Thr Ile Thr Phe Asp Leu Thr
 385 390 395 400

60
 Arg Ala Lys Thr Ala Gly Ser Asn Gly Ala Thr Val Pro Arg Trp Ser
 405 410 415

65
 Thr Gln Thr Ser Gly Gly Asp Gln Tyr Lys Ser Tyr Ser Asp Thr Lys
 420 425 430

70
 Ile Asn Asn Gly Lys Phe Ser Val Ser Phe Ser Thr Gly Gln Val Gln
 435 440 445

75
 Thr Phe Glu Ile Ser Gly Val Val Leu Lys

EP 3 330 352 A1

450

455

5

<210> 4
 <211> 109
 <212> PRT
 <213> Bacillus licheniformis

<400> 4

10

Ala Arg Tyr Asp Asp Val Leu Tyr Phe Pro Ala Ser Arg Tyr Pro Glu
 1 5 10 15

15

Thr Gly Ala His Ile Ser Asp Ala Ile Lys Ala Gly His Ala Asp Val
 20 25 30

20

Cys Thr Ile Glu Arg Ser Gly Ala Asp Lys Arg Arg Gln Glu Ser Leu
 35 40 45

25

Lys Gly Ile Pro Thr Lys Pro Gly Phe Asp Arg Asp Glu Trp Pro Met
 50 55 60

30

Ala Met Cys Glu Glu Gly Gly Lys Gly Ala Ser Val Arg Tyr Val Ser
 65 70 75 80

35

Ser Ser Asp Asn Arg Gly Ala Gly Ser Trp Val Gly Asn Arg Leu Asn
 85 90 95

40

Gly Tyr Ala Asp Gly Thr Arg Ile Leu Phe Ile Val Gln
 100 105

45

<210> 5
 <211> 109
 <212> PRT
 <213> Bacillus subtilis

<400> 5

Ala Ser Ser Tyr Asp Lys Val Leu Tyr Phe Pro Leu Ser Arg Tyr Pro
 1 5 10 15

50

Glu Thr Gly Ser His Ile Arg Asp Ala Ile Ala Glu Gly His Pro Asp
 20 25 30

55

Ile Cys Thr Ile Asp Asp Gly Ala Asp Lys Arg Arg Glu Glu Ser Leu
 35 40 45

Lys Gly Ile Pro Thr Lys Pro Gly Tyr Asp Arg Asp Glu Trp Pro Met
 50 55 60

60

Ala Val Cys Glu Glu Gly Gly Ala Gly Ala Asp Val Arg Tyr Val Thr
 65 70 75 80

EP 3 330 352 A1

Pro Ser Asp Asn Arg Gly Ala Gly Ser Trp Val Gly Asn Gln Met Ser
 85 90 95

5 Ser Tyr Pro Asp Gly Thr Arg Val Leu Phe Ile Val Gln
 100 105

10 <210> 6
 <211> 109
 <212> PRT
 <213> Bacillus licheniformis
 <400> 6

15 Ala Arg Tyr Asp Asp Ile Leu Tyr Phe Pro Ala Ser Arg Tyr Pro Glu
 1 5 10 15

20 Thr Gly Ala His Ile Ser Asp Ala Ile Lys Ala Gly His Ser Asp Val
 20 25 30

25 Cys Thr Ile Glu Arg Ser Gly Ala Asp Lys Arg Arg Gln Glu Ser Leu
 35 40 45

Lys Gly Ile Pro Thr Lys Pro Gly Phe Asp Arg Asp Glu Trp Pro Met
 50 55 60

30 Ala Met Cys Glu Glu Gly Gly Lys Gly Ala Ser Val Arg Tyr Val Ser
 65 70 75 80

35 Ser Ser Asp Asn Arg Gly Ala Gly Ser Trp Val Gly Asn Arg Leu Ser
 85 90 95

Gly Phe Ala Asp Gly Thr Arg Ile Leu Phe Ile Val Gln
 100 105

40 <210> 7
 <211> 204
 <212> PRT
 <213> Aspergillus oryzae
 45 <400> 7

50 Lys Thr Gly Ser Gly Asp Ser Gln Ser Asp Pro Ile Lys Ala Asp Leu
 1 5 10 15

Glu Val Lys Gly Gln Ser Ala Leu Pro Phe Asp Val Asp Cys Trp Ala
 20 25 30

55 Ile Leu Cys Lys Gly Ala Pro Asn Val Leu Gln Arg Val Asn Glu Lys
 35 40 45

EP 3 330 352 A1

Thr Lys Asn Ser Asn Arg Asp Arg Ser Gly Ala Asn Lys Gly Pro Phe
 50 55 60
 5 Lys Asp Pro Gln Lys Trp Gly Ile Lys Ala Leu Pro Pro Lys Asn Pro
 65 70 75 80
 10 Ser Trp Ser Ala Gln Asp Phe Lys Ser Pro Glu Glu Tyr Ala Phe Ala
 85 90 95
 15 Ser Ser Leu Gln Gly Gly Thr Asn Ala Ile Leu Ala Pro Val Asn Leu
 100 105 110
 20 Ala Ser Gln Asn Ser Gln Gly Gly Val Leu Asn Gly Phe Tyr Ser Ala
 115 120 125
 25 Thr Trp Phe Gln Ile Thr Lys Phe Thr Gly Ala Ala Gly Pro Tyr Cys
 145 150 155 160
 30 Lys Ala Leu Gly Ser Asn Asp Lys Ser Val Cys Asp Lys Asn Lys Asn
 165 170 175
 35 Ile Ala Gly Asp Trp Gly Phe Asp Pro Ala Lys Trp Ala Tyr Gln Tyr
 180 185 190
 40 <210> 8
 <211> 188
 <212> PRT
 <213> Trichoderma harzianum
 <400> 8
 45 Ala Pro Ala Pro Met Pro Thr Pro Pro Gly Ile Pro Thr Glu Ser Ser
 1 5 10 15
 50 Ala Arg Thr Gln Leu Ala Gly Leu Thr Val Ala Val Ala Gly Ser Gly
 20 25 30
 55 Thr Gly Tyr Ser Arg Asp Leu Phe Pro Thr Trp Asp Ala Ile Ser Gly
 35 40 45
 60 Asn Cys Asn Ala Arg Glu Tyr Val Leu Lys Arg Asp Gly Glu Gly Val
 50 55 60

EP 3 330 352 A1

Gln Val Asn Asn Ala Cys Glu Ser Gln Ser Gly Thr Trp Ile Ser Pro
 65 70 75 80
 5 Tyr Asp Asn Ala Ser Phe Thr Asn Ala Ser Ser Leu Asp Ile Asp His
 85 90 95
 10 Met Val Pro Leu Lys Asn Ala Trp Ile Ser Gly Ala Ser Ser Trp Thr
 100 105 110
 15 Thr Ala Gln Arg Glu Ala Leu Ala Asn Asp Val Ser Arg Pro Gln Leu
 115 120 125
 20 Trp Ala Val Ser Ala Ser Ala Asn Arg Ser Lys Gly Asp Arg Ser Pro
 130 135 140
 25 Asp Gln Trp Lys Pro Pro Leu Thr Ser Phe Tyr Cys Thr Tyr Ala Lys
 145 150 155 160
 30 Ser Trp Ile Asp Val Lys Ser Phe Tyr Lys Leu Thr Ile Thr Ser Ala
 165 170 175
 35 Glu Lys Thr Ala Leu Ser Ser Met Leu Asp Thr Cys
 180 185
 40 <210> 9
 <211> 361
 <212> PRT
 <213> *Aggregatibacter actinomycetemcomitans*
 45 <400> 9
 50 Asn Cys Cys Val Lys Gly Asn Ser Ile Tyr Pro Gln Lys Thr Ser Thr
 1 5 10 15
 55 Lys Gln Thr Gly Leu Met Leu Asp Ile Ala Arg His Phe Tyr Ser Pro
 20 25 30
 60 Glu Val Ile Lys Ser Phe Ile Asp Thr Ile Ser Leu Ser Gly Gly Asn
 35 40 45
 65 Phe Leu His Leu His Phe Ser Asp His Glu Asn Tyr Ala Ile Glu Ser
 50 55 60
 70 His Leu Leu Asn Gln Arg Ala Glu Asn Ala Val Gln Gly Lys Asp Gly
 65 70 75 80
 75 Ile Tyr Ile Asn Pro Tyr Thr Gly Lys Pro Phe Leu Ser Tyr Arg Gln
 85 90 95

EP 3 330 352 A1

Leu Asp Asp Ile Lys Ala Tyr Ala Lys Ala Lys Gly Ile Glu Leu Ile
 100 105 110
 5 Pro Glu Leu Asp Ser Pro Asn His Met Thr Ala Ile Phe Lys Leu Val
 115 120 125
 10 Gln Lys Asp Arg Gly Val Lys Tyr Leu Gln Gly Leu Lys Ser Arg Gln
 130 135 140
 Val Asp Asp Glu Ile Asp Ile Thr Asn Ala Asp Ser Ile Thr Phe Met
 145 150 155 160
 15 Gln Ser Leu Met Ser Glu Val Ile Asp Ile Phe Gly Asp Thr Ser Gln
 165 170 175
 20 His Phe His Ile Gly Gly Asp Glu Phe Gly Tyr Ser Val Glu Ser Asn
 180 185 190
 His Glu Phe Ile Thr Tyr Ala Asn Lys Leu Ser Tyr Phe Leu Glu Lys
 195 200 205
 25 Lys Gly Leu Lys Thr Arg Met Trp Asn Asp Gly Leu Ile Lys Asn Thr
 210 215 220
 30 Phe Glu Gln Ile Asn Pro Asn Ile Glu Ile Thr Tyr Trp Ser Tyr Asp
 225 230 235 240
 Gly Asp Thr Gln Asp Lys Asn Glu Ala Ala Glu Arg Arg Asp Met Arg
 245 250 255
 35 Val Ser Leu Pro Glu Leu Leu Ala Lys Gly Phe Thr Val Leu Asn Tyr
 260 265 270
 40 Asn Ser Tyr Tyr Leu Tyr Ile Val Pro Lys Ala Ser Pro Thr Phe Ser
 275 280 285
 45 Gln Asp Ala Ala Phe Ala Ala Lys Asp Val Ile Lys Asn Trp Asp Leu
 290 295 300
 Gly Val Trp Asp Gly Arg Asn Thr Lys Asn Arg Val Gln Asn Thr His
 305 310 315 320
 50 Glu Ile Ala Gly Ala Ala Leu Ser Ile Trp Gly Glu Asp Ala Lys Ala
 325 330 335
 55 Leu Lys Asp Glu Thr Ile Gln Lys Asn Thr Lys Ser Leu Leu Glu Ala
 340 345 350

EP 3 330 352 A1

Val Ile His Lys Thr Asn Gly Asp Glu
 355 360

5 <210> 10
 <211> 541
 <212> PRT
 <213> *Ascobolus stictoides*

10 <400> 10

Gln Thr Tyr Thr Leu Glu Ala Glu Ala Gly Thr Leu Thr Gly Val Thr
 1 5 10 15

15 Val Met Asn Glu Ile Ala Gly Phe Ser Gly Thr Gly Tyr Val Gly Gly
 20 25 30

20 Trp Asp Glu Asp Ala Asp Thr Val Ser Leu Thr Phe Thr Ser Asp Ala
 35 40 45

25 Thr Lys Leu Tyr Asp Val Lys Ile Arg Tyr Ser Gly Pro Tyr Gly Ser
 50 55 60

Lys Tyr Thr Arg Ile Ser Tyr Asn Gly Ala Thr Gly Gly Asp Ile Ser
 65 70 75 80

30 Leu Pro Glu Thr Thr Glu Trp Ala Thr Val Asn Ala Gly Gln Ala Leu
 85 90 95

35 Leu Asn Ala Gly Ser Asn Thr Ile Lys Leu His Asn Asn Trp Gly Trp
 100 105 110

Tyr Leu Ile Asp Ala Val Ile Leu Thr Pro Ser Val Pro Arg Pro Pro
 115 120 125

40 His Gln Val Thr Asp Ala Leu Val Asn Thr Asn Ser Asn Ala Val Thr
 130 135 140

45 Lys Gln Leu Met Lys Phe Leu Val Ser Lys Tyr His Lys Ala Tyr Ile
 145 150 155 160

50 Thr Gly Gln Gln Glu Leu His Ala His Gln Trp Val Glu Lys Asn Val
 165 170 175

Gly Lys Ser Pro Ala Ile Leu Gly Leu Asp Phe Met Asp Tyr Ser Pro
 180 185 190

55 Ser Arg Val Glu Phe Gly Thr Thr Ser Gln Ala Val Glu Gln Ala Ile
 195 200 205

EP 3 330 352 A1

Asp Phe Asp Lys Arg Gly Gly Ile Val Thr Phe Ala Trp His Trp Asn
 210 215 220
 5
 Ala Pro Ser Gly Leu Ile Asn Thr Pro Gly Ser Glu Trp Trp Arg Gly
 225 230 235 240
 10
 Phe Tyr Thr Glu His Thr Thr Phe Asp Val Ala Ala Ala Leu Gln Asn
 245 250 255
 15
 Thr Thr Asn Ala Asn Tyr Asn Leu Leu Ile Arg Asp Ile Asp Ala Ile
 260 265 270
 20
 Ala Val Gln Leu Lys Arg Leu Gln Thr Ala Gly Val Pro Val Leu Trp
 275 280 285
 25
 Arg Pro Leu His Glu Ala Glu Gly Gly Trp Phe Trp Trp Gly Ala Lys
 290 295 300
 30
 Gly Pro Glu Pro Ala Lys Lys Leu Tyr Lys Ile Leu Tyr Asp Arg Leu
 305 310 315 320
 35
 Thr Asn Tyr His Lys Leu Asn Asn Leu Ile Trp Val Trp Asn Ser Val
 325 330 335
 40
 Ala Lys Asp Trp Tyr Pro Gly Asp Glu Ile Val Asp Val Leu Ser Phe
 340 345 350
 45
 Asp Ser Tyr Pro Ala Gln Pro Gly Asp His Gly Pro Val Ser Ala Gln
 355 360 365
 50
 Tyr Asn Ala Leu Val Glu Leu Gly Lys Asp Lys Lys Leu Ile Ala Ala
 370 375 380
 55
 Thr Glu Val Gly Thr Ile Pro Asp Pro Asp Leu Met Gln Leu Tyr Glu
 385 390 395 400
 Ser Tyr Trp Ser Phe Phe Val Thr Trp Glu Gly Glu Phe Ile Glu Asn
 405 410 415
 Gly Val His Asn Ser Leu Glu Phe Leu Lys Lys Leu Tyr Asn Asn Ser
 420 425 430
 Phe Val Leu Asn Leu Asp Thr Ile Gln Gly Trp Lys Asn Gly Ala Gly
 435 440 445
 Ser Ser Thr Thr Thr Val Lys Ser Thr Thr Thr Thr Pro Thr Thr Thr

EP 3 330 352 A1

	450		455		460														
5	Ile	Lys	Ser	Thr	Thr	Thr	Thr	Pro	Val	Thr	Thr	Pro	Thr	Thr	Val	Lys			
	465						470					475				480			
	Thr	Thr	Thr	Thr	Pro	Thr	Thr	Thr	Ala	Thr	Thr	Val	Lys	Ser	Thr	Thr			
10					485					490					495				
	Thr	Thr	Ala	Gly	Pro	Thr	Pro	Thr	Ala	Val	Ala	Gly	Arg	Trp	Gln	Gln			
				500					505					510					
15	Cys	Gly	Gly	Ile	Gly	Phe	Thr	Gly	Pro	Thr	Thr	Cys	Glu	Ala	Gly	Thr			
			515					520					525						
	Thr	Cys	Asn	Val	Leu	Asn	Pro	Tyr	Tyr	Ser	Gln	Cys	Leu						
20		530					535					540							
	<210>	11																	
	<211>	526																	
	<212>	PRT																	
25	<213>	Chaetomium virescens																	
	<400>	11																	
	Pro	Arg	Asp	Pro	Gly	Ala	Thr	Ala	Arg	Thr	Phe	Glu	Ala	Glu	Asp	Ala			
30	1				5					10					15				
	Thr	Leu	Ala	Gly	Thr	Asn	Val	Asp	Thr	Ala	Leu	Ser	Gly	Phe	Thr	Gly			
				20					25					30					
35	Thr	Gly	Tyr	Val	Thr	Gly	Phe	Asp	Gln	Ala	Ala	Asp	Lys	Val	Thr	Phe			
			35					40					45						
	Thr	Val	Asp	Ser	Ala	Ser	Thr	Glu	Leu	Tyr	Asp	Leu	Ser	Ile	Arg	Val			
40		50					55					60							
	Ala	Ala	Ile	Tyr	Gly	Asp	Lys	Arg	Thr	Ser	Val	Val	Leu	Asn	Gly	Gly			
	65					70					75					80			
45	Ala	Ser	Ser	Glu	Val	Tyr	Phe	Pro	Ala	Gly	Glu	Thr	Trp	Thr	Asn	Val			
					85					90					95				
	Ala	Ala	Gly	Gln	Leu	Leu	Leu	Asn	Gln	Gly	Ser	Asn	Thr	Ile	Asp	Ile			
50				100					105					110					
	Val	Ser	Asn	Trp	Gly	Trp	Tyr	Leu	Ile	Asp	Ser	Ile	Thr	Leu	Thr	Pro			
			115					120					125						
55	Ser	Thr	Pro	Arg	Pro	Ala	His	Gln	Ile	Asn	Glu	Ala	Pro	Val	Asn	Ala			

EP 3 330 352 A1

	130		135		140												
5	Ala 145	Ala	Asp	Lys	Asn	Ala 150	Lys	Ala	Leu	Tyr	Ser 155	Tyr	Leu	Arg	Ser	Ile 160	
10	Tyr	Gly	Lys	Lys	Ile 165	Leu	Ser	Gly	Gln	Gln 170	Glu	Leu	Ser	Leu	Ser	Asn 175	
15	Trp	Ile	Ala	Gln 180	Gln	Thr	Gly	Lys	Thr 185	Pro	Ala	Leu	Val	Ser 190	Val	Asp	
20	Leu	Met	Asp 195	Tyr	Ser	Pro	Ser	Arg 200	Val	Glu	Arg	Gly	Thr 205	Val	Gly	Thr	
25	Ala	Val 210	Glu	Glu	Ala	Ile	Gln 215	His	His	Asn	Arg	Gly 220	Gly	Ile	Val	Ser	
30	Val 225	Leu	Trp	His	Trp	Asn 230	Ala	Pro	Thr	Gly	Leu 235	Tyr	Asp	Thr	Glu	Glu 240	
35	His	Arg	Trp	Trp	Ser 245	Gly	Phe	Tyr	Thr	Ser 250	Ala	Thr	Asp	Phe	Asp 255	Val	
40	Ala	Ala	Ala	Leu 260	Ser	Ser	Thr	Thr	Asn 265	Ala	Asn	Tyr	Thr	Leu 270	Leu	Ile	
45	Arg	Asp	Ile 275	Asp	Ala	Ile	Ala	Val 280	Gln	Leu	Lys	Arg	Leu 285	Gln	Ser	Ala	
50	Gly	Val 290	Pro	Val	Leu	Phe	Arg 295	Pro	Leu	His	Glu	Ala 300	Glu	Gly	Gly	Trp	
55	Phe 305	Trp	Trp	Gly	Ala	Lys 310	Gly	Pro	Glu	Pro	Ala 315	Lys	Lys	Leu	Trp	Gly 320	
60	Ile	Leu	Tyr	Asp	Arg 325	Val	Thr	Asn	His 330	His	Gln	Ile	Asn	Asn	Leu 335	Leu	
65	Trp	Val	Trp	Asn 340	Ser	Ile	Leu	Pro	Glu 345	Trp	Tyr	Pro	Gly	Asp 350	Ala	Thr	
70	Val	Asp	Ile 355	Leu	Ser	Ala	Asp 360	Val	Tyr	Ala	Gln	Gly	Asn 365	Gly	Pro	Met	
75	Ser	Thr 370	Gln	Tyr	Asn	Gln	Leu 375	Ile	Glu	Leu	Gly	Lys 380	Asp	Lys	Lys	Met	

EP 3 330 352 A1

Ile Ala Ala Ala Glu Val Gly Ala Ala Pro Leu Pro Asp Leu Leu Gln
 385 390 395 400

5 Ala Tyr Glu Ala His Trp Leu Trp Phe Thr Val Trp Gly Asp Ser Phe
 405 410 415

10 Ile Asn Asn Ala Asp Trp Asn Ser Leu Asp Thr Leu Lys Lys Val Tyr
 420 425 430

15 Thr Ser Asp Tyr Val Leu Thr Leu Asp Glu Ile Gln Gly Trp Gln Gly
 435 440 445

20 Ser Thr Pro Ser Ala Thr Thr Thr Ser Ser Thr Thr Thr Pro Ser Ala
 450 455 460

25 Thr Thr Thr Thr Thr Thr Pro Ser Thr Thr Ala Thr Thr Ala Thr Pro
 465 470 475 480

30 Ser Ala Thr Thr Thr Ala Ser Pro Val Thr Tyr Ala Glu His Trp Gly
 485 490 495

35 Gln Cys Ala Gly Lys Gly Trp Thr Gly Pro Thr Thr Cys Arg Pro Pro
 500 505 510

40 Tyr Thr Cys Lys Tyr Gln Asn Asp Trp Tyr Ser Gln Cys Leu
 515 520 525

<210> 12
 <211> 452
 <212> PRT
 <213> *Preussia aemulans*

45 <400> 12

50 Gln Thr Val Ile Tyr Gln Ala Glu Gln Ala Lys Leu Ser Gly Val Thr
 1 5 10 15

55 Val Glu Phe Ser Ile Ile Lys Gln Val Val Gly Thr Gly Tyr Val Glu
 20 25 30

Gly Phe Asp Glu Ser Thr Asp Ser Ile Thr Phe Thr Val Glu Ser Thr
 35 40 45

60 Thr Ala Ala Leu Tyr Asp Leu Ala Leu Thr Tyr Asn Gly Pro Tyr Gly
 50 55 60

65 Asp Lys Tyr Thr Asn Val Val Leu Asn Asn Ala Ala Gly Ser Gln Val
 65 70 75 80

EP 3 330 352 A1

Ser Leu Pro Ala Thr Thr Ala Trp Thr Thr Val Pro Ala Gly Gln Val
 85 90 95
 5 Leu Leu Asn Ala Gly Ala Asn Thr Ile Gln Ile Gln Asn Asn Trp Gly
 100 105 110
 Trp Tyr Leu Val Asp Ser Ile Ser Leu Lys Pro Ala Ala Thr Arg Gly
 115 120 125
 10 Ala His Gln Ile Thr Thr Lys Pro Val Asn Lys Asn Ala Asn Ser Asp
 130 135 140
 15 Ala Lys Ala Leu Leu Lys Tyr Leu Gly Ser Ile Tyr Gly Lys Lys Ile
 145 150 155 160
 20 Leu Ser Gly Gln Gln Asp Leu Ser Ser Leu Asp Trp Val Thr Lys Asn
 165 170 175
 Val Gly Lys Thr Pro Ala Val Leu Gly Leu Asp Thr Met Asp Tyr Ser
 180 185 190
 25 Glu Ser Arg Lys Ser Arg Gly Ala Val Ser Thr Asp Val Asp Lys Ala
 195 200 205
 30 Ile Ala Phe Ala Lys Lys Gly Gly Ile Val Thr Phe Cys Trp His Trp
 210 215 220
 Gly Ala Pro Thr Gly Leu Phe Asp Ser Ala Ala Gln Pro Trp Tyr Arg
 225 230 235 240
 35 Gly Phe Tyr Thr Asp Ala Thr Asp Phe Asn Ile Glu Thr Ala Leu Lys
 245 250 255
 40 Asp Thr Thr Asn Ala Asn Tyr Thr Leu Leu Met Lys Asp Ile Asp Thr
 260 265 270
 45 Ile Ala Val Gln Leu Lys Lys Leu Gln Asp Ala Gly Val Pro Val Ile
 275 280 285
 Trp Arg Pro Leu His Glu Ala Glu Gly Gly Trp Phe Trp Trp Gly Ala
 290 295 300
 50 Lys Gly Pro Glu Pro Ala Lys Lys Leu Trp Lys Ile Met Tyr Asp Arg
 305 310 315 320
 55 Leu Thr Asn Gln His Gly Leu Asn Asn Leu Val Trp Thr Trp Asn Ser
 325 330 335

EP 3 330 352 A1

Val Ala Pro Asn Trp Tyr Pro Gly Asp Asp Thr Val Asp Ile Val Ser
340 345 350

5 Ala Asp Thr Tyr Ser Gln Gly Asp His Gly Pro Ile Ser Ala Thr Tyr
355 360 365

10 Asn Asn Leu Leu Ala Leu Thr Asn Asp Thr Lys Ile Ile Ala Ala Ala
370 375 380

15 Glu Ile Gly Ser Val Met Glu Pro Ala Gln Leu Gln Ala Tyr Gln Ala
385 390 395 400

20 Asp Trp Val Tyr Phe Cys Val Trp Ser Gly Glu Phe Ile Asp Gly Gly
405 410 415

25 Val Trp Asn Ser Leu Asp Phe Leu Lys Lys Val Tyr Asn Asp Pro Tyr
420 425 430

30 Val Leu Thr Leu Asp Glu Ile Gln Gly Trp Lys Thr Ala Arg Gly Lys
435 440 445

35 Pro Arg Val Ser
450

<210> 13
<211> 312
<212> PRT
<213> Yunnania penicillata

40 Ala Pro Ser Thr Thr Pro Val Asn Glu Lys Ala Thr Asp Ala Ala Lys
1 5 10 15

45 Asn Leu Leu Ser Tyr Leu Val Glu Gln Ala Ala Asn Gly Val Thr Leu
20 25 30

50 Ser Gly Gln Gln Asp Leu Glu Ser Ala Gln Trp Val Ser Asp Asn Val
35 40 45

Gly Lys Trp Pro Ala Ile Leu Gly Ile Asp Phe Met Asp Tyr Ser Pro
50 55 60

Ser Arg Val Glu Tyr Gly Ala Val Gly Ser Thr Val Pro Asp Ala Ile
65 70 75 80

55 Ser Tyr Asp Ser Asp Gly Gly Ile Val Thr Phe Cys Trp His Trp Gly
85 90 95

EP 3 330 352 A1

Ser Pro Ser Gly Thr Tyr Asn Thr Thr Asp Gln Pro Trp Trp Ser Asn
100 105 110

5 Phe Tyr Thr Glu Ala Thr Ala Phe Asp Ile Ala Ala Ala Met Asp Asp
115 120 125

10 Pro Asp Ser Ala Asp Tyr Asn Leu Leu Val Arg Asp Ile Asp Ala Ile
130 135 140

15 Ser Glu Leu Leu Leu Gln Leu Gln Asp Leu Asp Ile Pro Ile Leu Trp
145 150 155 160

Arg Pro Leu His Glu Ala Glu Gly Gly Trp Phe Trp Trp Gly Ala Lys
165 170 175

20 Gly Pro Glu Ala Cys Ile Ala Leu Tyr Arg Leu Met Phe Asp Arg Met
180 185 190

25 Thr Asn His His Gly Leu Asn Asn Leu Leu Trp Val Trp Asn Ser Val
195 200 205

Asp Pro Ser Trp Tyr Pro Gly Asn Asp Val Val Asp Ile Val Ser Ala
210 215 220

30 Asp Ile Tyr Ala Asp Ala Gly Asp His Ser Pro Gln Glu Glu Thr Phe
225 230 235 240

35 Ala Ser Leu Gln Ser Leu Thr Gly Asp Thr Lys Leu Val Ala Leu Gly
245 250 255

Glu Val Gly Asn Ile Pro Asp Pro Ala Ser Thr Gly Gly Val Ala Asp
260 265 270

40 Trp Ala Tyr Trp Val Thr Trp Asn Gly Asp Phe Ile Lys Gly Glu Asp
275 280 285

45 Tyr Asn Pro Leu Glu Tyr Lys Lys Glu Val Phe Ser Ala Glu Asn Ile
290 295 300

Ile Thr Arg Asp Glu Val Asp Val
305 310

50

<210> 14
<211> 327
<212> PRT
55 <213> Myrothecium roridum

<400> 14

EP 3 330 352 A1

Gly Thr Ile Glu Asn Arg Gln Trp Leu Thr Tyr Asn Pro Val Asp Ser
 1 5 10 15
 5 Ala Ala Thr Thr Glu Ala Arg Ala Leu Leu Arg Tyr Ile Gln Ser Gln
 20 25 30
 Tyr Gly Trp Arg Tyr Leu Ser Gly Gln Gln Glu Arg Ala Glu Val Gln
 10 35 40 45
 Trp Leu Lys Ser Asn Ile Gly Lys Thr Pro Ala Ile Gln Gly Ser Asp
 50 55 60
 15 Leu Ile Asp Tyr Ser Pro Ser Arg Val Ser Tyr Gly Ala Thr Ser Thr
 65 70 75 80
 Ala Val Glu Asp Ala Ile Ala Phe Asp Arg Gln Gly Gly Ile Val Thr
 20 85 90 95
 Phe Thr Trp His Trp Asn Ala Pro Asn Cys Leu Tyr Asn Ser Ala Asp
 100 105 110
 25 Gln Pro Trp Tyr Phe Gly Phe Tyr Thr Lys Ala Thr Cys Phe Asn Ile
 115 120 125
 30 Gln Ala Ala Leu Ala Gln Gly Ser Asn Gly Ala Asp Tyr Lys Leu Leu
 130 135 140
 Ile Arg Asp Ile Asp Ala Ile Ala Val Gln Leu Lys Arg Leu Arg Asp
 145 150 155 160
 35 Ala Lys Val Pro Ile Leu Phe Arg Pro Leu His Glu Pro Asp Gly Ala
 165 170 175
 40 Trp Phe Trp Trp Gly Ala Lys Gly Ser Gly Pro Phe Lys Gln Leu Trp
 180 185 190
 45 Asp Ile Leu Tyr Asp Arg Leu Thr Lys Tyr His Gly Leu His Asn Met
 195 200 205
 50 Leu Trp Val Cys Asn Thr Glu Lys Ser Asp Trp Tyr Pro Gly Asn Asn
 210 215 220
 Lys Cys Asp Ile Ala Thr Thr Asp Val Tyr Val Asn Ala Gly Asp His
 225 230 235 240
 55 Ser Val Gln Lys Ser His Trp Asp Ala Leu Tyr Gly Val Ser Gly Gly
 245 250 255

EP 3 330 352 A1

Gln Arg Ile Leu Ala Leu Gly Glu Val Gly Val Ile Pro Asp Pro Glu
260 265 270

5 Arg Gln Ala Ser Glu Asn Val Pro Trp Ala Tyr Trp Met Thr Trp Asn
275 280 285

10 Gly Tyr Phe Ile Arg Asp Gly Asn Tyr Asn Ser Arg Asn Phe Leu Gln
290 295 300

15 Ser Thr Phe Ser Asn Ala Arg Val Val Thr Leu Asp Gly Thr Ser Pro
305 310 315 320

Leu Gly Asn Trp Lys Ser Ser
325

20 **Claims**

1. A cleaning composition comprising:

25 a endo-beta-1,6-galactanase enzyme; and
an alkoxylated phenol compound that is selected from the group consisting of an alkoxylated polyaryl phenol compound, an alkoxylated polyalkyl phenol compound, and mixtures thereof.

30 2. A cleaning composition according to claim 1, wherein the enzyme has an amino acid sequence having at least 60%, or at least 80%, or at least 90% or at least 95% identity with the amino acid sequence shown in SEQ ID NO:1, SEQ ID NO:2 or SEQ ID NO:3.


35 3. A cleaning composition according to claims 1 and 2, wherein the galactanase enzyme is selected from Glycoside Hydrolase Family 30.

4. A cleaning composition according to any preceding claim, wherein the galactanase enzyme is obtainable from *Streptomyces davawensis*, *Trichoderma harzianum*, *Streptomyces avermitilis*, or a mixture thereof.

40 5. A cleaning composition according to any preceding claim, wherein the composition further comprises a β -N-acetylglucosaminidase enzyme from E.C. 3.2.1.52, preferably an enzyme having at least 70% identity to SEQ ID NO:9.

45 6. A cleaning composition according to any preceding claim, wherein the alkoxylated phenol compound has a structure according to Formula I:

Formula I



wherein


each R_1 is independently selected from linear or branched C3-C15 alkyl groups and C3-C15 aryl groups,
 X is selected from ethoxy or propoxy groups,
 n is from 2 to 70, and
 T is selected from H, SO_3^- , COO^- and PO_3^{2-} , preferably H and SO_3^- .

7. A cleaning composition according to claim 6, wherein each R_1 is the same.

8. A cleaning composition according to any preceding claim, wherein the alkoxyated phenol is selected from the group consisting of:

(i) an uncharged alkoxyated tristyrylphenols according to Formula II:


Formula II



wherein n is selected from 2 to 70, more preferably n is selected from 10 to 54, most preferably n = 16 or 20;

(ii) an anionic alkoxyated tristyrylphenol according to Formula III:


Formula III



wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- ; and wherein n is selected from 2 to 54;

(iii) an uncharged alkoxyated tri(n-butyl)phenol according to Formula IV:


Formula IV



wherein n is selected from 2 to 50;

(iv) an anionic alkoxyated tri(n-butyl)phenol of the following structure according to Formula V:

Formula V




wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- , and wherein n is selected from 6 to 50; and

(v) mixtures thereof.

9. A cleaning composition according to claim 8, wherein the alkoxyated phenol comprises:

(i) an uncharged alkoxyated tristyrylphenols according to Formula II:

Formula II




wherein n is selected from 2 to 70, more preferably n is selected from 10 to 54, most preferably n = 16 or 20.

10. A cleaning composition according to claim 8, wherein the alkoxyated phenol comprises:

(ii) an anionic alkoxyated tristyrylphenol according to Formula III:

Formula III




wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- , and wherein n is selected from 2 to 54.

11. A cleaning composition according to claim 8, wherein the alkoxyated phenol comprises:

(iii) an uncharged alkoxyated tri(n-butyl)phenol according to Formula IV:

Formula IV




wherein n is selected from 2 to 50.

12. A cleaning composition according to claim 8, wherein the alkoxyated phenol comprises:

(iv) an anionic alkoxyated tri(n-butyl)phenol of the following structure according to Formula V:

Formula V



wherein R is selected from SO_3^- , COO^- and PO_3^{2-} , preferably selected from SO_3^- and COO^- , and wherein n is selected from 6 to 50.

13. A cleaning composition according to any preceding claim, wherein the alkoxyated phenol compound is an alkoxyated polyaryl phenol compound or an alkoxyated polyalkyl phenol compound.

14. A method of cleaning a surface, preferably a textile, comprising mixing the cleaning composition according to any preceding claim with water to form an aqueous liquor and contacting a surface, preferably a textile, with the aqueous liquor in a laundering step.

15. The use of an endo-beta-1,6-galactanase enzyme and an alkoxyated polyaryl phenol and/or an alkoxyated polyalkyl phenol in a cleaning composition to enhance the stain-removal, whiteness, and/or malodor-reducing benefits.



EUROPEAN SEARCH REPORT

Application Number
EP 17 20 4768

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y,D	WO 2015/185689 A1 (NOVOZYMES AS [DK]) 10 December 2015 (2015-12-10) * page 38, line 31 - page 54, line 24; claims; examples; sequence 2 * -----	1-15	INV. C11D3/386
Y	WO 2016/110379 A1 (UNILEVER PLC [GB]; UNILEVER NV [NL]; CONOPCO INC DBA UNILEVER [US]) 14 July 2016 (2016-07-14) * claims; examples * -----	1-15	
			TECHNICAL FIELDS SEARCHED (IPC)
			C11D
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 11 January 2018	Examiner Vernier, Frédéric
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

1
EPO FORM 1503 03 82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 17 20 4768

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

11-01-2018

10	Patent document cited in search report	Publication date	Patent family member(s)	Publication date
15	WO 2015185689 A1	10-12-2015	CN 106414698 A EP 3152290 A1 WO 2015185689 A1	15-02-2017 12-04-2017 10-12-2015
20	WO 2016110379 A1	14-07-2016	NONE	

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2015185689 A [0032]
- US 20120135498 A [0050]
- WO 2015040159 A [0060]
- WO 2004067737 A [0067]
- WO 2015091989 A [0067]
- WO 2015091990 A [0067]
- WO 2015024739 A [0067]
- WO 2015143360 A [0067]
- US 6312936 B1 [0067]
- US 5679630 A [0067]
- US 4760025 A [0067]
- US 7262042 B [0067]
- WO 09021867 A [0067]
- DE 102006022216 A1 [0067]
- DE 102006022224 A1 [0067]
- WO 2015089447 A [0067]
- WO 2015089441 A [0067]
- WO 2016066756 A [0067]
- WO 2016066757 A [0067]
- WO 2016069557 A [0067]
- WO 2016069563 A [0067]
- WO 2016069569 A [0067]
- WO 8906270 A [0067]
- WO 05052161 A [0067]
- WO 05052146 A [0067]
- WO 07044993 A2 [0067]
- WO 2014194032 A [0067]
- WO 2014194054 A [0067]
- WO 2014194117 A [0067]
- WO 2015193488 A [0067]
- WO 2016075078 A [0067]
- WO 9217577 A [0067]
- US 5352604 A [0069]
- WO 2009149144 A [0069]
- WO 2009149145 A [0069]
- WO 201056653 A [0069]
- WO 201056640 A [0069]
- WO 201107211 A [0069]
- US 20110237487 A [0069]
- WO 2011140316 A [0069]
- WO 2012151480 A [0069]
- EP 2510092 A [0069]
- EP 2566960 A [0069]
- EP 2705145 A [0069]
- US 7153818 B [0070]
- WO 9700324 A [0070]
- EP 1022334 A [0070]
- WO 9402597 A [0070]
- WO 9418314 A [0070]
- WO 9623874 A [0070]
- WO 9743424 A [0070]
- US 5856164 A [0070]
- WO 9923211 A [0070]
- WO 9623873 A [0070]
- WO 0060060 A [0070]
- WO 06002643 A [0070]
- US 6093562 A [0070]
- WO 09149130 A [0070]
- EP 2540825 A [0070]
- EP 2357220 A [0070]
- EP 2534233 A [0070]
- WO 2009100102 A [0070]
- WO 2010115028 A [0070]
- US 6939702 B1 [0072]
- US 20090217464 A [0072]
- EP 12001034 A [0072]
- EP 2623586 A [0072]
- US 7141403 B2 [0073]
- WO 2002099091 A [0076]
- WO 01062903 A [0078]
- WO 9902663 A [0078]
- WO 01064853 A [0078]
- WO 2002077242 A [0078]
- WO 03089598 A [0078]
- WO 9905243 A [0088]
- WO 9905242 A [0088]
- WO 9905244 A [0088]
- WO 9905082 A [0088]
- WO 9905084 A [0088]
- WO 9905241 A [0088]
- WO 9907656 A [0088]
- WO 0023549 A [0088]
- WO 0023548 A [0088]
- WO 0887497 A [0105]
- WO 9108281 A [0111]
- WO 9001815 A [0111]
- US 6020303 A [0150]
- US 6060443 A [0150]

Non-patent literature cited in the description

- A. B. BORASTON et al. *Biochemical Journal*, 2004, vol. 382, 769-781 [0075]

EP 3 330 352 A1

- *Biochem J.*, 1991, vol. 280, 309-316 [0076]
- **NEEDLEMAN ; WUNSCH.** *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0077]
- **RICE et al.** EMBOSS: The European Molecular Biology Open Software Suite. *Trends in Genetics*, 2000, vol. 16, 276-277 [0077]