

**(12) STANDARD PATENT APPLICATION (11) Application No. AU 2013224661 A1**  
**(19) AUSTRALIAN PATENT OFFICE**

(54) Title  
**Agricultural adjuvant compositions, pesticide compositions, and methods for using such compositions**

(51) International Patent Classification(s)  
**A01N 25/00** (2006.01)

(21) Application No: **2013224661** (22) Date of Filing: **2013.09.04**

(43) Publication Date: **2013.10.03**

(43) Publication Journal Date: **2013.10.03**

(62) Divisional of:  
**2006315513**

(71) Applicant(s)  
**Rhodia Inc.**

(72) Inventor(s)  
**Smith, Francis George;Pazhianur, Rajesh;Modaressi, Hedieh**

(74) Agent / Attorney  
**Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000**

AGRICULTURAL ADJUVANT COMPOSITIONS, PESTICIDE COMPOSITIONS, AND  
METHODS FOR USING SUCH COMPOSITIONS

**Abstract**

An adjuvant composition that contains, based on 100 parts by weight of the adjuvant composition (a) one or more betaine surfactant compounds, and (b) one or more surfactant compounds selected from, alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols is useful as a component of pesticide compositions.

**AGRICULTURAL ADJUVANT COMPOSITIONS, PESTICIDE  
COMPOSITIONS, AND METHODS FOR USING SUCH COMPOSITIONS**

5 Field of the Invention

This invention relates to agricultural adjuvant compositions, pesticide compositions, and methods for using such compositions.

10 Background of the Invention

Many agricultural pesticides, including insecticides, fungicides, herbicides, miticides, and plant growth regulators, are applied in the form of a liquid composition. In addition to the pesticide and a solvent, such  
15 liquid compositions typically include one or more adjuvant compounds intended to improve one or more properties of the liquid composition, such as for example, storage stability, ease of handling, pesticide efficacy against target organisms.

20 There is a continuing interest in pesticide compositions that exhibit improved properties.

Summary of the Invention

25 In a first aspect, the present invention is directed to an adjuvant composition comprising,

- (a) one or more betaine surfactant compounds, and
- (b) one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates,  
30 alkoxyated fatty acids, and alkoxyated alcohols.

In a second aspect, the present invention is directed to a pesticide composition, comprising:

- (a) one or more betaine surfactant compounds,  
(b) one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols, and  
5 (c) an effective amount of a pesticide.

In a third aspect, the present invention is directed to a method for treating a target plant, comprising applying the above described pesticide composition to such plant.

10

#### Detailed Description of Invention and Preferred Embodiments

As used herein, the term "alkyl" means a saturated straight chain, branched chain, or cyclic hydrocarbon radical, such as for example,  
15 methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, t-butyl, pentyl, n-hexyl, and cyclohexyl.

As used herein, the term "alkoxy" means an oxy radical that is substituted with an alkyl group, such as for example, methoxy, ethoxy,  
20 propoxy, isopropoxy, and butoxy. As used herein in reference to an organic compound, the term "alkoxyated" means that the compound comprises one or more alkoxy or, more typically, poly(alkyleneoxy) moieties, such as, for example a poly(ethyleneoxy), poly(propyleneoxy), or poly(ethyleneoxypropyleneoxy) moiety and the term "ethoxyated"  
25 means that the compound comprises at least one ethoxy or poly(ethyleneoxy) moiety. As used herein in reference to a poly(alkyleneoxy) moiety, the notation "(n)", wherein n is an integer, indicates the number of alkyleneoxy monomeric units in the poly(alkyleneoxy) moiety. For example such as "ethoxyated (15) tridecyl  
30 alcohol" means a tridecyl alcohol ethoxyated with 15 moles of ethyleneoxy units per mole of tridecyl alcohol.

As used herein, the term "alkenyl" means an unsaturated straight chain, branched chain, or cyclic hydrocarbon radical that contains one or more carbon-carbon double bonds, such as, for example, ethenyl, 1-propenyl, and 2-propenyl.

5

As used herein, the term "aryl" means a monovalent unsaturated hydrocarbon radical containing one or more six-membered carbon rings in which the unsaturation may be represented by three conjugated double bonds, which may be substituted one or more of carbons of the ring with hydroxy, alkyl, alkenyl, halo, haloalkyl, or amino, such as, for example, phenoxy, phenyl, methylphenyl, dimethylphenyl, trimethylphenyl, chlorophenyl, trichloromethylphenyl, aminophenyl, and tristyrylphenyl.

10

As used herein, the term "arylene" means a divalent unsaturated hydrocarbon radical containing one or more six-membered carbon rings in which the unsaturation may be represented by three conjugated double bonds, which may be substituted one or more of carbons of the ring with hydroxy, alkyl, alkenyl, halo, haloalkyl, or amino, such as, for example, phenylene, methylphenylene, trimethylphenylene, aminophenylene, and tristyrylphenylene.

15  
20

As used herein, the term "aralkyl" means an alkyl group substituted with one or more aryl groups, such as, for example, phenylmethyl, phenylethyl, and triphenylmethyl.

25

As used herein, the term "aralkenyl" means an alkenyl group substituted with an aryl group, such as, for example, phenylethenyl, and phenylpropenyl.

30

As used herein, the term "aryloxy" means an oxygen radical substituted with an aryl group, such as, for example, phenoxy, methylphenoxy, and trimethylphenoxy.

As used herein, the terminology "(C<sub>n</sub>-C<sub>m</sub>)" in reference to an organic group, wherein n and m are each integers, indicates that the group may contain from n carbon atoms to m carbon atoms per group.

5 As used herein, the term "agronomically acceptable salts" refers to salts prepared from agronomically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Typical agronomically acceptable salts the compound referred to herein  
 10 comprise an anion derived from the compound, for example, by deprotonation of a hydroxy or hydroxyalkyl substituent, and one or more positively charged counterions. Suitable positively charged counterions include inorganic cations and organic cations, such as for example, sodium cations, potassium cations, calcium cations, magnesium cations, isopropylamine cations, ammonium cations, and tetraalkylammonium  
 15 cations.

Betaine surfactant compounds are generally known compounds. In one embodiment, the betaine surfactant compound comprises one or more compounds according to formula (I):



wherein:


R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> are each independently alkyl, hydroxyalkyl, alkylamidoalkyl, or -CH<sub>2</sub>COOH, or an agronomically acceptable salt thereof.

25 In one embodiment, R<sup>1</sup> and R<sup>2</sup> are each independently (C<sub>1</sub>-C<sub>6</sub>)alkyl, more typically methyl, and R<sup>3</sup> is (C<sub>1</sub>-C<sub>30</sub>)alkyl, more typically (C<sub>8</sub>-C<sub>18</sub>)alkyl.

In one embodiment,  $R^1$  and  $R^2$  are each independently ( $C_1$ - $C_6$ )alkyl, more typically methyl, and  $R^3$  is alkylamidoalkyl, more typically ( $C_1$ - $C_{30}$ )alkylamido( $C_1$ - $C_6$ )alkyl.

5 Suitable betaines include, for example, decyl dimethyl betaine, lauryl dimethyl betaine, coco dimethyl betaine, stearyl dimethyl betaine, cocoamidopropyl betaine, and mixtures thereof.

Alkyl ether sulfate surfactant compounds are generally known  
10 compounds and include agronomically acceptable salts of alkoxyated alkyl sulfates. In one embodiment, the adjuvant composition comprises one or more alkyl ether sulfate surfactant compounds according to formula (II):



15 wherein:

each  $R^4$  is independently H or methyl,

$R^5$  is alkyl,

$m$  is an integer of from 1 to 100, more typically from 1 to 50, and

$X^+$  is an agronomically acceptable cation.

20

In one embodiment,  $R^5$  is ( $C_1$ - $C_{30}$ )alkyl, more typically ( $C_8$ - $C_{18}$ )alkyl.


In one embodiment, each  $R^4$  is H and is from 1 to about 10.

25

Suitable alkyl ether sulfate surfactant compounds, include, for example, sodium ( $C_8$ - $C_{10}$ )alkyl ether sulfate and sodium lauryl ether sulfate.

Sulfonate surfactant compounds are generally known compounds and include agronomically acceptable salts of mono-sulfonic acids, agronomically acceptable salts of di-sulfonic acids, and mixtures thereof.

In one embodiment, the adjuvant composition comprises one or more sulfonate surfactant compounds according to formula (III) or (IV):



wherein:

- 10  $\text{R}^6$  is aryl, aryloxy, or alkenyl,  
 $\text{R}^7$  and  $\text{R}^{7'}$  are each independently arylene, and  
 $\text{X}^+$  is an agronomically acceptable cation.


In one embodiment,  $\text{R}^6$  is phenyl, (C<sub>8</sub>-C<sub>18</sub>)alkylphenyl, or (C<sub>8</sub>-  
 15 C<sub>18</sub>)alkylphenoxy.

In one embodiment,  $\text{R}^7$  and  $\text{R}^{7'}$  are each (C<sub>8</sub>-C<sub>18</sub>)alkylphenylene.

Suitable sulfonate surfactant compounds include, for example,  
 20 calcium dodecylbenzene sulfonate, calcium octadecylphenyl sulfonate,  
 sodium tridecyl benzene sulfonate, isopropylamine dodecyl benzene  
 sulfonate, isopropylamine tridecyl benzene sulfonate, ammonium tridecyl  
 phenyl sulfonate, sodium (C<sub>8</sub>-C<sub>18</sub>)alkylphenoxy sulfonate, sodium xylene  
 sulfonate, sodium (C<sub>14</sub>-C<sub>16</sub>)alpha olefin sulfonate, disodium  
 25 alkyldiphenyloxide disulfonates such as, for example, DOWFAX 2A1  
 (Dow Chemical Company), and mixtures thereof.





Sulfosuccinate surfactant compounds are generally known compounds and include agronomically acceptable salts of mono-esters of sulfosuccinic acid, agronomically acceptable salts of di-esters of sulfosuccinic acid, each of which may, optionally, be alkoxyated, as well as mixtures thereof. In one embodiment, the adjuvant composition comprises one or more sulfosuccinate surfactant compounds according to formula (V):



10 wherein:

A and A' are each independently  $-O^-X^+$ , or




each  $R^8$  is independently H or methyl,

15 each  $R^9$  is independently H, alkyl, aryl, or alkylamidoalkyl,


each  $n$  is independently 0 or an integer of from 1 to about 100,  
more typically from 1 to 50, and

each  $X^+$  is an agronomically acceptable cation.

20 In one embodiment, one of A and A' is  $-O^-X^+$ , and the other of A  
and A' is




5 In one embodiment, A and A' are each



10 Suitable sulfosuccinate surfactant compounds include, for example, disodium mono-octylsulfosuccinate, sodium dioctylsulfosuccinate, disodium lauryl sulfosuccinate, disodium laureth sulfosuccinate, disodium laurimide (MEA) sulfosuccinate, disodium mono-alkylphenyl ether sulfosuccinate, and mixtures thereof.

15 Alkyl ether carboxylate surfactant compounds are generally known compounds and include agronomically acceptable salts of alkoxyated carboxylic acids. In one embodiment, the adjuvant composition comprises one or more alkyl ether carboxylate surfactant compounds according to formula (VI):



wherein:

each R<sup>10</sup> is independently H or methyl,

R<sup>11</sup> is alkyl or alkenyl,

25 p is 0 or an integer of from 1 to 100, more typically an integer of from 1 to 50, and

$X^+$  is an agronomically acceptable cation.

In one embodiment,  $R^{11}$  is  $(C_8-C_{18})$ alkyl.

5 In one embodiment, each  $R^{10}$  is H and p is an integer of from 1 to about 10.

Suitable alkyl ether carboxylate surfactant compounds include, for example, sodium laureth-13 carboxylate.

10


In one embodiment, the adjuvant composition comprises one or more alkoxyated fatty acid surfactant compounds. The fatty acid portion of such alkoxyated fatty acid surfactant compounds is derived from a saturated or unsaturated mono- or di- fatty acids, typically  $(C_6-C_{30})$ fatty acids, such as, for example, lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, linoleic acid, behenic acid, erucic acid, or a mixture thereof, including vegetable oils, such as, for example, tall oil, rapeseed oil, canola oil, soy oil, coconut oil, castor oil, corn oil, olive oil, sunflower oil, cottonseed oil, palm oil, peanut oil, sesame oil, safflower oil, linseed oil, flax seed oil, palm kernel oil, and mixtures thereof. These acids are alkoxyated with from 2 to 20 moles, more typically from 5 to 20 moles of a  $(C_2-C_4)$ alkylene oxide, more typically, ethylene oxide.

15

20

Alkoxyated alcohol surfactant compounds are generally known compounds. In one embodiment, the adjuvant composition comprises one or more alkoxyated alcohol surfactant compounds according to formula (VII):

25



(VII)

wherein

30 each  $R^{12}$  is independently H or methyl,  
 $R^{13}$  is alkyl or alkenyl,

q is an integer of from 1 to 100, more typically from 1 to 50, or an agronomically acceptable salt thereof.

In one embodiment, R<sup>13</sup> is (C<sub>6</sub>-C<sub>22</sub>)alkyl.

5

In one embodiment, each R<sup>12</sup> is H and q is an integer of from 1 to about 30.

Suitable alkoxyated alcohol surfactant compounds include, for example, ethoxylated (15) tridecyl alcohol, ethoxylated (7) lauryl alcohol, ethoxylated (20) oleyl alcohol, ethoxylated (15) stearyl alcohol, and mixtures thereof.

10

The adjuvant composition comprises a non-zero amount of the one or more betaine surfactant compounds and a non-zero amount of one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols. As used herein, "non-zero amount" means an amount greater than 0.

15

20

In one embodiment, the adjuvant composition comprises, based on 100 parts by weight ("pbw") of the adjuvant composition:

- (a) greater or equal to than about 0.1 pbw, more typically from about 1 pbw to about 99 pbw, even more typically from about 1 pbw to about 50 pbw, and still more typically from about 5 pbw to about 25 pbw, of one or more betaine surfactant compounds, and
- (b) greater or equal to than about 0.1 pbw, more typically from about 0.001 pbw to about 0.1 pbw, even more typically from about 0.005 pbw to about 0.095 pbw, and still more typically from about 0.02 pbw to about 0.08 pbw, of one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

25

30

In one embodiment of the adjuvant composition, the one or more surfactants comprise at least one alkyl ether sulfate surfactant and may optionally further comprise one or more surfactants selected from  
5 sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

In one embodiment of the adjuvant composition, the one or more surfactants comprise at least one sulfonate surfactant and may optionally  
10 further comprise one or more surfactants selected from alkyl ether sulfates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

In one embodiment of the adjuvant composition, the one or more  
15 surfactants comprise at least one sulfosuccinate surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

In one embodiment of the adjuvant composition, the one or more  
20 surfactants comprise at least one alkyl ether carboxylate surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkoxyated fatty acids, and alkoxyated alcohols

25  
In one embodiment of the adjuvant composition, the one or more surfactants comprise at least one alkoxyated fatty acid surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates,  
30 and alkoxyated alcohols

In one embodiment of the adjuvant composition, the one or more surfactants comprise at least one alkoxyated alcohol surfactant and may

optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, and alkoxyated fatty acids.

5           In one embodiment, the adjuvant composition comprises, per 1 pbw betaine component of such adjuvant composition, from about 0.1 to about 10, more typically from about 1 to about 8, and even more typically from about 2 to about 5 pbw, of one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether  
10 carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

Suitable pesticides are biologically active compounds used to control agricultural pests and include, for example, herbicides, plant growth regulators, crop dessicants, fungicides, bacteriocides,  
15 bacteriostats, insecticides, and insect repellants. Suitable pesticides include, for example, triazine herbicides such as metribuzin, hexaxinone, or atrazine; sulfonylurea herbicides such as chlorsulfuron; uracils such as lenacil, bromacil, or terbacil; urea herbicides such as linuron, diuron, siduron, or neburon; acetanilide herbicides such as alachlor, or  
20 metolachlor; thiocarbamate herbicides such as benthocarb, triallate; oxadiazolone herbicides such as oxadiazon; phenoxyacetic acids such as 2,4-D; diphenyl ether herbicides such as fluazifop, acifluorfen, bifenox, or oxyfluorfen; dinitro aniline herbicides such as trifluralin; organophosphonate herbicides such as glyphosate salts and esters;  
25 dihalobenzonitrile herbicides such as bromoxynil, or ioxynil, dipyridilium herbicides such as paraquat. Suitable fungicides include, for example, nitrilo oxime fungicides such as cymoxanil; imidazole fungicides such as benomyl, carbendazim, or thiophanate-methyl; triazole fungicides such as triadimefon; sulfenamide fungicides, such as captan; dithio-carbamate  
30 fungicides such as maneb, mancozeb, or thiram; chloronated aromatic fungicides such as chloroneb; dichloro aniline fungicides such as iprodione, strobilurin fungicides such as kresoxim-methyl, trifloxystrobin or azoxystrobin; chlorothalonil; copper salt fungicides such as copper

oxychloride; sulfur; phenylamides; and acylamino fungicides such as metalaxyl or mefenoxam. Suitable insecticides, include, for example, carbamate insecticides, such as methomyl, carbaryl, carbofuran, or aldicarb; organo thiophosphate insecticides such as EPN, isofenphos, 5 isoxathion, chlorpyrifos, or chlormephos; organophosphate insecticides such as terbufos, monocrotophos, or terachlorvinphos; perchlorinated organic insecticides such as methoxychlor; synthetic pyrethroid insecticides such as fenvalerate, abamectin or emamectin benzoate, neonicotinoide insecticides such as thiamethoxam or imidacloprid; 10 pyrethroid insecticides such as lambda-cyhalothrin, cypermethrin or bifenthrin, and oxadiazine insecticides such as indoxacarb, imidachlopryd, or fipronil. Suitable miticides include, for example, propynyl sulfite miticides such as propargite; triazapentadiene miticides such as amitraz; chlorinated aromatic miticides such as chlorobenzilate, 15 or tetradifan; and dinitrophenol miticides such as binapacryl. Suitable nematicides include carbamate nematicides, such as oxamyl.

Pesticide compounds are, in general, referred herein to by the names assigned by the International Organization for Standardization 20 (ISO). ISO common names may be cross-referenced to International Union of Pure and Applied Chemistry ("IUPAC") and Chemical Abstracts Service ("CAS") names through a number of sources such as, for example, the *Compendium of Pesticide Common Names*, which is available on-line at <http://www.hclrss.demon.co.uk/index.html>.

25

In one embodiment, the pesticide comprises one or more compounds selected from group consisting of herbicides, plant growth regulators, crop dessicants, fungicides, bacteriocides, bacteriostats, insecticides, miticides, nematocides, and insect repellants.

30

In one embodiment, the pesticide comprises one or more compounds selected from the group consisting of glufosinate, glyphosate, water soluble glufosinate salts, water soluble glyphosate salts, and

mixtures thereof, including, for example sodium, potassium, isopropyl amine, and ammonium salts.

In one embodiment, the pesticide comprises one or more  
5 compounds selected from the group consisting of the potassium salt of glyphosate, the sodium salt of glyphosate, the isopropyl amine salt of glyphosate, the ammonium salt of glyphosate.

Herbicidal compositions containing glyphosate or derivatives  
10 thereof are used to control a very wide variety of plants worldwide and can, when applied to the target plant in a herbicidally effective amount, reportedly control one or more target plant species of one or more of the following genera: Abutilon, Amaranthus, Artemisia, Asclepias, Avena, Axonopus, Borreria, Brachiaria, Brassica, Bromus, Chenopodium,  
15 Cirsium, Commelina, Convolvulus, Cynodon, Cyperus, Digitaria, Echinochloa, Eleusine, Elymus, Equisetum, Erodium, Helianthus, Imperata, Ipomoea, Kochia, Lolium, Malva, Oryza, Ottochloa, Panicum, Paspalum, Phalaris, Phragmites, Polygonum, Portulaca, Pteridium, Pueraria, Rubus, Salsola, Setaria, Sida, Sinapis, Sorghum, Triticum,  
20 Typha, Ulex, Xanthium and Zea, including annual broadleaf species such as, for example, velvetleaf (*Abutilon theophrasti*), pigweed (*Amaranthus* spp.), buttonweed (*Borreria* spp.), oilseed rape, canola, indian mustard, etc. (*Brassica* spp.), commelina (*Commelina* spp.), filaree (*Erodium* spp.), sunflower (*Helianthus* spp.), morningglory (*Ipomoea* spp.), kochia (*Kochia scoparia*), mallow (*Malva* spp.), wild buckwheat, smartweed, etc.  
25 (*Polygonum* spp.), purslane (*Portulaca* spp.), russian thistle (*Salsola* spp.), sida (*Sida* spp.), wild mustard (*Sinapis arvensis*) and cocklebur (*Xanthium* spp.), annual narrowleaf species such as for example, wild oat (*Avena fatua*), carpetgrass (*Axonopus* spp.), downy brome (*Bromus tectorum*), crabgrass (*Digitaria* spp.), barnyardgrass (*Echinochloa crus-galli*), goosegrass (*Eleusine indica*), annual ryegrass (*Lolium multiflorum*), rice (*Oryza sativa*), ottochloa (*Ottochloa nodosa*), bahiagrass (*Paspalum notatum*), canarygrass (*Phalaris* spp.), foxtail (*Setaria* spp.), wheat



(*Triticum aestivum*) and corn (*Zea mays*), perennial broadleaf species such as, for example, mugwort (*Artemisia* spp.), milkweed (*Asclepias* spp.), canada thistle (*Cirsium arvense*), field bindweed (*Convolvulus arvensis*) and kudzu (*Pueraria* spp.), perennial narrowleaf species such as for example, brachiaria (*Brachiaria* spp.), bermudagrass (*Cynodon dactylon*), yellow nutsedge (*Cyperus esculentus*), purple nutsedge (*C. rotundus*), quackgrass (*Elymus repens*), lalang (*Imperata cylindrica*), perennial ryegrass (*Lolium perenne*), guineagrass (*Panicum maximum*), dallisgrass (*Paspalum dilatatum*), reed (*Phragmites* spp.), johnsongrass (*Sorghum halepense*) and cattail (*Typha* spp.), and other perennial species such as, for example, horsetail (*Equisetum* spp.), bracken (*Pteridium aquilinum*), blackberry (*Rubus* spp.), and gorse (*Ulex europaeus*).

15 As used herein, the terminology "effective amount" in reference to the relative amount of a pesticide in a pesticide composition means the relative amount of pesticide that is effective to control the growth of a target organism, for example, a target plant, fungus, bacterium, or insect, when the pesticide composition is applied to the organism at a given application rate.

20 In one embodiment, the pesticide is glyphosate herbicide and the pesticide composition is an herbicide composition that comprises a herbicidally effective amount of glyphosate.

25 As used herein, the terminology "an herbicidally effective amount" in reference to the relative amount of herbicide in an herbicidal composition means the relative amount that is effective to control growth of a target plant when the herbicidal composition is spray applied to the target plant at a given application rate.

30 The adjuvant composition comprises a non-zero amount of the one or more betaine surfactant compounds and a non-zero amount of one or

more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

- 5           In one embodiment, the pesticide composition comprises, based on 100 pbw of the adjuvant composition:
- (a)   greater than or equal to about 0.001 part by weight, more typically from about 0.001 pbw to about 0.1 pbw, even more typically from about 0.001 pbw to about 0.05 pbw, and still more typically from about 0.005 pbw to about 0.025 pbw, of one or more betaine surfactant compounds,
- 10           (b)   greater than or equal to about 0.001 part by weight, more typically from about 0.001 pbw to about 0.1 pbw, even more typically from about 0.005 pbw to about 0.095 pbw, and still more typically from about 0.02 pbw to about 0.08 pbw, of one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols, and
- 15           (c)   an effective amount of a pesticide.

20           In one embodiment of the pesticide composition, the one or more surfactants comprise at least one alkyl ether sulfate surfactant and may optionally further comprise one or more surfactants selected from sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

25

          In one embodiment of the adjuvant composition, the one or more surfactants comprise at least one sulfonate surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

30

In one embodiment of the pesticide composition, the one or more surfactants comprise at least one sulfosuccinate surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

In one embodiment of the pesticide composition, the one or more surfactants comprise at least one alkyl ether carboxylate surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkoxyated fatty acids, and alkoxyated alcohols

In one embodiment of the pesticide composition, the one or more surfactants comprise at least one alkoxyated fatty acid surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, and alkoxyated alcohols

In one embodiment of the pesticide composition, the one or more surfactants comprise at least one alkoxyated alcohol surfactant and may optionally further comprise one or more surfactants selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, and alkoxyated fatty acids.

In one embodiment, the pesticide composition comprises, based on 100 pbw of the composition, from about 10 pbw to about 90 pbw, more typically from about 30 to about 60 pbw, glyphosate.

The adjuvant and pesticide compositions of the present invention may each, optionally, further comprise one or more agronomically acceptable solvent. Suitable solvents include, for example, water, and organic solvents, such as for example, alkylated aromatic solvents, such

as toluene or alkylated naphthalenes and mineral oil fractions, such as paraffinic hydrocarbons.

5 In one embodiment, the adjuvant composition of the present invention is an aqueous composition and thus comprises water.

In one embodiment, the pesticide composition of the present invention is an aqueous composition and thus comprises water.

10 In one embodiment, the adjuvant composition of the present invention further comprises, based on 100 pbw of such composition, up to about 25 pbw an organic solvent.

15 In one embodiment, the pesticide composition further comprises a fertilizer. Such fertilizers can provide the primary nutrients of nitrogen, phosphorus and/or potassium such as urea ammonium nitrate (30-0-0), 10-34-0, secondary nutrients sulfur, calcium, magnesium such as ammonium thiosulfate 12-0-0-26S, micronutrient fertilizers containing zinc, iron, molybdenum, copper, boron, chlorine, magnesium, for example  
20 0-0-1 3%-S; 3%-Zn; 2%-Fe; 2%-Mn and mixtures thereof. In one embodiment, the pesticide composition comprises from about 85 to about 99 pbw, more typically from about 90 to about 99 pbw, and even more typically from about 93 to about 99 pbw, of a mixture of fertilizer and water.

25

In one embodiment, the pesticide composition of the present invention further comprises one or more water conditioners, such as for example, chelating agents, such as ethylenediamine tetraacetic acid, complexing agents such as ammonium sulfate, and pH adjusting agents,  
30 such as citric acid and polyacrylic acid.

In one embodiment, the pesticide composition of the present invention comprises, based on 100 pbw of such composition, from about 0.1 to about 3 pbw, more typically from about 0.7 to about 2.5 pbw, of one or more water conditioners, typically ammonium sulfate.

5

The pesticide composition of the present invention may, optionally, further comprise other ingredients, such as one or more additional surfactants, one or more thickeners, such as polysaccharide thickeners, and polyacrylamide thickeners, as well as antifoams, spreaders, and drift control agents.

10

The adjuvant composition of the present invention is made by combining and mixing the components of such composition.

15

The pesticide composition of the present invention is made by mixing the ingredients together. In one embodiment, the pesticide composition is made by combining and mixing the adjuvant composition of the present invention, a pesticide compound, and water. Alternatively, the pesticide composition is made by combining and mixing the separate components of the adjuvant composition, a pesticide, and water.

20

In one embodiment, the pesticide composition is spray applied to foliage of a target plant at a rate of from about 0.5 pint/acre to about 3 pints/acre, more typically from about 0.5 pint/acre to about 2.5 pints/acre.

25

#### EXAMPLES 1 -10 AND COMPARATIVE EXAMPLES C1-C4

The pesticide compositions of Examples 1 - 10 and Comparative Examples C1-C4 were aqueous solutions made by combining the ingredients in the relative amounts (based on 100 pbw of the respective pesticide composition) set forth below and mixing:

30

EX #	Ingredients (pbw/ 100 pbw pesticide composition)
1	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 3.5 pbw 30% aqueous solution of sodium lauryl ether sulfate, ethoxylated with 1-3 moles ethylene oxide per mole 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 57 pbw water
2	2.25 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 2.5 pbw tall oil fatty acid ethoxylated with 10 moles of ethylene oxide per mole 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 57.25 pbw water
3	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 4.95 pbw tall oil fatty acid ethoxylated with 10 moles of ethylene oxide per mole 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 55.55 pbw water
4	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 2.5 pbw tall oil fatty acid ethoxylated with 10 moles of ethylene oxide per mole 1.12 pbw 45% aqueous solution of sodium alkylphenoxy benzene sulfonate 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 56.88 pbw water
5	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 2.8 pbw 56% aqueous solution of ammonium (C <sub>8</sub> -C <sub>10</sub> )alkyl ether sulfate, ethoxylated with 3 moles ethylene oxide per mole 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 57.7 pbw water
6	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 4 pbw sodium dioctyl sulfosuccinate 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 56.5 pbw water
7	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 3.75 pbw sodium laureth-13 carboxylate 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 56.75 pbw water
8	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 2 pbw disodium laurimide (MEA) sulfosuccinate 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 58.5 pbw water

EX #	Ingredients (pbw/ 100 pbw pesticide composition)
9	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 5 pbw ethoxylated (20 moles ethylene oxide per mole) oleyl alcohol 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 55.5 pbw water
10	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 2 pbw ammonium xylene sulfonate 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 58.5 pbw water
C1	38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 62 pbw water
C2	3 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 59 pbw water
C3	1.5 pbw 30% aqueous solution of (C <sub>12</sub> -C <sub>14</sub> )alkyl dimethyl betaine 38 pbw glyphosate isopropyl amine salt (MON0139, Monsanto) 60.5 pbw water
C4	Roundup <sup>®</sup> Ultramax

The efficacy of the compositions in controlling plant growth was tested by applying the compositions to the following plant species: Green Foxtail ("GRF"), Giant Foxtail ("GF"), Shattercane ("SC"), Barnyard Grass ("BG"), Redroot Pigweed ("PW"), Common Waterhemp ("WH"), Common Purslane ("CP"), Kochia ("KO"), Velvetleaf ("VL"), Common Lambsquarter ("LQ"), Ivy Morning Glory ("MG"), at a rate of 0.5 pint per acre and at a rate of 1.0 pint per acre. Results for the 0.5 pint per acre application rate at 28 days post application are given in TABLE I below as percent control of plant growth for various plant species. Results for the 1.0 pint per acre application rate are given below in TABLE II as percent control of plant growth for various plant species.

TABLE I - 28 Days, 0.5 pint/acre

Example #	% Control of Plant Growth										
	GRF	GF	SC	BG	PW	WH	CP	KO	VL	LQ	MG
1	100	100	84	77	100	100	100	91	78	85	91
2	100	100	91	83	100	100	73	72	68	88	91
3	100	100	73	63	100	100	95	73	73	73	78
4	100	100	79	62	100	100	88	71	42	78	91
5	100	98	58	63	100	100	73	58	78	63	88
6	100	99	58	68	100	100	78	63	30	58	86
7	100	100	88	86	100	100	68	73	62	90	78
8	100	100	73	72	100	100	78	58	28	73	81
9	100	100	91	92	100	100	83	73	83	87	83
10	100	100	81	74	100	100	83	63	73	81	71
C1	100	97	38	38	100	100	22	48	22	18	62
C2	100	100	78	73	100	100	73	73	48	82	63
C3	100	100	73	68	100	100	58	82	63	73	80
C4	100	100	73	82	100	100	78	92	72	85	48

TABLE II - 28 Days, 1.0 pint/acre

Example #	% Control of Plant Growth										
	GRF	GF	SC	BG	PW	WH	CP	KO	VL	LQ	MG
1	100	100	99	100	100	100	100	100	100	97	93
2	100	100	100	100	100	100	94	100	85	100	90
3	100	100	95	88	100	100	100	94	89	94	86
4	100	100	94	78	100	100	100	90	91	94	96
5	100	100	68	68	100	100	100	83	94	72	88
6	100	100	84	78	100	100	96	83	83	83	92
7	100	100	94	88	100	100	96	94	83	91	94
8	100	100	88	88	100	100	100	94	82	92	91
9	100	100	100	100	100	100	100	93	93	100	88
10	100	100	100	100	100	100	100	100	93	100	73
C1	100	98	48	48	100	100	72	78	70	12	75
C2	100	100	88	91	100	100	85	98	73	91	78
C3	100	100	86	88	100	100	100	88	98	85	78
C4	100	100	86	91	100	100	94	100	88	94	71



Claims:

1. An adjuvant composition comprising,
  - (a) one or more betaine surfactant compounds, and
  - 5 (b) one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acid, and alkoxyated alcohols.
  
- 10 2. The composition of claim 1, wherein the composition comprises, based on 100 parts by weight of the composition, greater than or equal to about 0.1 part by weight of the one or more betaine surfactant compounds and greater than or equal to about 0.1 part by weight of one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and  
15 alkoxyated alcohols.
  
3. The composition of claim 1, wherein one or more surfactant compounds comprises an alkyl ether sulfate surfactant.
  
- 20 4. The composition of claim 1, wherein one or more surfactant compounds comprise a sulfonate surfactant.
  
5. The composition of claim 1, wherein one or more surfactant compounds comprise a sulfosuccinate surfactant.  
25
  
6. The composition of claim 1, wherein one or more surfactant compounds comprise an alkyl ether carboxylate surfactant
  
7. The composition of claim 1, wherein one or more surfactant  
30 compounds comprise an alkoxyated fatty acid surfactant.
  
8. The composition of claim 1, wherein one or more surfactant compounds comprise an alkoxyated alcohol surfactant.

9. The composition of claim 1, wherein composition comprises, per 1 part by weight betaine component of such composition, from about 0.1 to about 10 parts by weight of the one or more surfactant compounds  
5 selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.
10. A pesticide composition, comprising:  
(a) one or more betaine surfactant compounds,  
10 (b) one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acid ethoxylates, and alkoxyated alcohols, and  
(c) an effective amount of a pesticide.
- 15 11. The composition of claim 10, wherein the composition comprises, based on 100 parts by weight of the composition, greater than or equal to about 0.001 part by weight of the one or more betaine surfactant compounds and greater than or equal to about 0.001 part by weight of the one or more surfactant compounds selected from alkyl ether sulfates,  
20 sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acid ethoxylates, and alkoxyated alcohols.
- 25 12. The composition of claim 10, wherein the pesticide compound comprises one or more compounds selected from herbicides, plant growth regulators, crop dessicants, fungicides, bacteriocides, bacteriostats, insecticides, miticides, nematocides, and insect repellants.
- 30 13. The composition of claim 10, wherein the pesticide compound comprises one or more compounds selected from glufosinate, glyphosate, water soluble glufosinate salts, and water soluble glyphosate salts.
14. The composition of claim 10, wherein the pesticide compound comprises one or more compounds selected from the potassium salt of

glyphosate, the sodium salt of glyphosate, the isopropyl amine salt of glyphosate, and the ammonium salt of glyphosate.

5 15. The composition of claim 10, wherein one or more surfactant compounds comprise an alkyl ether sulfate surfactant.

16. The composition of claim 10, wherein one or more surfactant compounds comprise a sulfonate surfactant.

10 17. The composition of claim 10, wherein one or more surfactant compounds comprise a sulfosuccinate surfactant.

15 18. The composition of claim 10, wherein one or more surfactant compounds comprise an alkyl ether carboxylate surfactant.

19. The composition of claim 10, wherein one or more surfactant compounds comprise an alkoxyated fatty acid surfactant.

20 20. The composition of claim 10, wherein one or more surfactant compounds comprise an alkoxyated alcohol surfactant.

25 21. The composition of claim 10, wherein composition comprises, per 1 part by weight betaine component of such composition, from about 0.1 to about 10 part by weight of the one or more surfactant compounds selected from alkyl ether sulfates, sulfonates, sulfosuccinates, alkyl ether carboxylates, alkoxyated fatty acids, and alkoxyated alcohols.

30 22. A method for treating a target organism to control the growth of the organism, comprising applying to the organism a pesticide composition according to claim 10.

23. The method of claim 22, wherein the method comprises treating a target plant to control the growth of the plant and the pesticide is selected from herbicides and plant growth regulators.

5 24. The method of claim 23, wherein the pesticide comprises one or more compounds selected from glufosinate, glyphosate, water soluble glufosinate salts, and water soluble glyphosate salts.

10 25. The method of claim 23, wherein the pesticide comprises one or more compounds selected from the potassium salt of glyphosate, the sodium salt of glyphosate, the isopropyl amine salt of glyphosate, and the ammonium salt of glyphosate.