



US 20030077696A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0077696 A1**

**Thim et al.** (43) **Pub. Date: Apr. 24, 2003**

(54) **HUMAN SPAMOLYTIC POLYPEPTIDE IN GLYCOSYLATED FORM**

(21) Appl. No.: **09/528,644**

(22) Filed: **Mar. 20, 2000**

(76) Inventors: **Lars Thim**, Gentofte (DK); **Kjeld Norris**, Hellerup (DK); **Fanny Norris**, Hellerup (DK); **Soren Bjorn**, Lyngby (DK); **Mogens Christensen**, Gentofte (DK); **Per Franklin Nielsen**, Varlose (DK)

(30) **Foreign Application Priority Data**

Jan. 20, 1994 (J)..... PCT/DK94/00037  
Jan. 21, 1993 (DK)..... 0068/93

**Publication Classification**

(51) **Int. Cl.<sup>7</sup>** ..... **C07K 17/00**; A01N 37/18;  
C07K 14/00; C07K 1/00;  
C12P 21/06; A61K 38/00  
(52) **U.S. Cl.** ..... **435/69.1**; 514/2; 530/350

(57) **ABSTRACT**








A human spasmodic polypeptide (HSP) having the amino acid SEQ ID NO:1, characterized by being in a glycosylated form.


Correspondence Address:

**Steve T Zelson Esq**  
**Novo Nordisk of North America Inc**  
**405 Lexington Avenue**  
**Suite 6400**  
**New York, NY 10017 (US)**

(\* ) Notice: This is a publication of a continued prosecution application (CPA) filed under 37 CFR 1.53(d).

## Trefoil family of peptides

| Peptide                          |       | Number of trefoil domains   | Species      |
|----------------------------------|-------|---|--------------|
| Intestinal Trefoil Factor        | (ITF) |    | Rat<br>Human |
| Breast cancer associated peptide | (pS2) |    | Human        |
| Spasmolytic polypeptide          | (hSP) |    | Human        |
|                                  | (PSP) |     | Porcine      |
|                                  | (mSP) |      | Mouse        |
| Spasmolysins                     | (xP1) |  | Frog         |
|                                  | (xP4) |   | Frog         |



ECORI

GAATTCATTCAAGAATAGTTCAAACAAGAAGATTACAAACTATCAATTCATACACAAT

ATAAACGACCAAAGAATGAAGGCTGTTTTCTTGGTTTTGTCTTGATCGGATTCTGCTG

M K A V F L V L S L I G F C W

GGCCCAACCAGTCACTGGCGATGAATCATCTGTTGAGATTCCGGAAGAGTCTCTGATCAT

A Q P V T G D E S S V E I P E E S L I I

CGCTGAAAACACCACTTTGGCTAACGTCGCCATGGCTGAGAGATTGGAGAAGAGAgagaa

A E N T T L A N V A M A E R L E K R E K

Dde I 1 →

accctccccctgccagtgtccagggcTGAGCCCCCATAACAGGACGAACTGCGGCTTCCC

P S P C Q C S R L S P H N R T N C G F P

TGGAATCACCAGTGACCAGTGTTTTGACAATGGATGCTGTTTCGACTCCAGTGTCACTGG

G I T S D Q C F D N G C C F D S S V T G

AvaII ← 2

Ggtccccctggtgtttccacccccctcccaagcaagagtcggatcagtgcgatcatggaggt

V P W C F H P L P K Q E S D Q C V M E V

→ 4

Dde I

CTCAGACAGAAGAAACTGTGGCTACCCGGGCATCAGCCCCGAGGAATGCGCCTCTCGGAA

S D R R N C G Y P G I S P E E C A S R K

← 5 Nco I

GTGCTGCTTCTCCAACCTTCATCTTTGAAGTGccatggtgcttcttcccgaactctgtgga

C C F S N F I F E V P W C F F P N S V E

Xba I

agactgccattactaagtCTAGA

D C H Y

Fig. 3

# hSP expression vector




Fig. 4




Fig. 5




Fig. 6




Fig. 7







Fig. 8



## HUMAN SPASMOLYTIC POLYPEPTIDE IN GLYCOSYLATED FORM

### FIELD OF INVENTION

[0001] The present invention relates to human spasmolytic polypeptide in glycosylated form, variants of human and porcine spasmolytic polypeptides and a method of producing spasmolytic polypeptides in glycosylated form.

### BACKGROUND OF THE INVENTION

[0002] Human spasmolytic polypeptide (HSP) belongs to a family of peptides containing one or more characteristic trefoil domains [1]. The trefoil domain is made up of a sequence of 38 or 39 amino acid residues in which 6 cysteine residues are linked in the configuration 1-5, 2-4 and 3-6 thus forming a characteristic trefoil structure [1]. The trefoil family of peptides consists of rat intestinal trefoil factor, ITF [2], human breast cancer associated peptide, pS2 [3,4,5], porcine, human and murine spasmolytic polypeptide (PSP, HSP, MSP) [6,7,8] and frog spasmolysins (xP1, xP2 and xP4) [8,10,11] all containing 1, 2 or 4 trefoil domains (**FIG. 1**).

[0003] The physiological function of the trefoil peptides is poorly understood, and so far only PSP has been studied in any detail. In the porcine pancreas, PSP is found in the acinar cells and to be secreted in large amounts (50-100 mg/ml) into the pancreatic juice upon stimulation with pancreozymin or secretin [12,13,14]. PSP is resistant to digestion by intestinal proteases in the gastrointestinal tract [12], and specific binding of PSP to rat intestinal mucosa cells and membrane preparations from these cells has been demonstrated [15,16]. In the porcine gastrointestinal tract, specific receptor-like binding to Paneth cells in the duodenum has been found [17]. These results suggest a unique intraluminal function of the peptide. A pharmacological screening has indicated that PSP has spasmolytic and gastric acid secretion inhibitory effects [18], and studies on mammalian cells have indicated a growth factor-like activity of PSP [19].

[0004] The DNA sequence and derived amino acid sequence of the human counterpart of porcine SP is shown in [8]. Unlike PSP, human SP (**FIG. 2**), has been found to be expressed in the stomach, but not in the pancreas to any greater extent [8]. An increased expression of HSP and pS2 has been reported to be associated with peptic ulcers and mucosal injury in inflammatory bowel disease [20,21] indicating a possible healing function of these peptides.

[0005] Only very limited amounts of HSP can be prepared by extraction of human tissue. An object of study resulting in the present invention was therefore to prepare recombinant HSP in sufficient amounts for physiological and biochemical studies of the peptide.

### SUMMARY OF THE INVENTION

[0006] It has surprisingly been found that when recombinant HSP is produced in certain host organisms, a proportion of it is produced in glycosylated form by posttranslational modifications. The glycosylated form of HSP has not, to applicant's best knowledge, been described previously.

[0007] Accordingly, the present invention relates to human spasmolytic polypeptide (HSP) which has the amino acid sequence

(SEQ ID NO:1)

Glu Lys Pro Ser Pro Cys Gln Cys Ser Arg Leu Ser  
 Pro His Asn Arg Thr Asn Cys Gly Phe Pro Gly Ile  
 Thr Ser Asp Gln Cys Phe Asp Asn Gly Cys Cys Phe  
 Asp Ser Ser Val Thr Gly Val Pro Trp Cys Phe His  
 Pro Leu Pro Lys Gln Glu Ser Asp Gln Cys Val Met  
 Glu Val Ser Asp Arg Arg Asn Cys Gly Tyr Pro Gly  
 Ile Ser Pro Glu Glu Cys Ala Ser Arg Lys Cys Cys  
 Phe Ser Asn Phe Ile Phe Glu Val Pro Trp Cys Phe  
 Phe Pro Asn Ser Val Glu Asp Cys His Tyr

[0008] or a functionally equivalent homologue thereof, characterized by being in glycosylated form.

[0009] In the present context, the term "functionally equivalent" is intended to indicate that the homologous polypeptide has a biological activity (e.g. spasmolytic effect) corresponding to that of native HSP. The term "homologue" is intended to indicate a polypeptide encoded by DNA which hybridizes to the same probe as the DNA coding for HSP under conditions of high or low stringency (e.g. as described in Sambrook et. al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989). More specifically, the term is intended to refer to a DNA sequence which is at least 60% homologous to the sequence encoding HSP with the amino acid sequence shown above. The term is intended to include modifications of the DNA sequence such as nucleotide substitutions which do not give rise to another amino acid sequence of the polypeptide, but which correspond to the codon usage of the host organism into which the DNA construct is introduced or nucleotide substitutions which do give rise to a different amino acid sequence and therefore, possibly, a different protein structure which might give rise to a mutant polypeptide with different properties than the native enzyme. Other examples of possible modifications are insertion of one or more codons into the sequence, addition of one or more codons at either end of the sequence, or deletion of one or more codons at either end or within the sequence. The term "glycosylated" is intended to indicate that a carbohydrate moiety is present at one or more sites of the protein molecule.

[0010] It is at present contemplated that glycosylation of HSP may give rise to differences in the biological activity of the protein, for instance with respect to stability towards proteolytic enzymes in the gastrointestinal tract, solubility at gastric and/or intestinal pH compared to non-glycosylated HSP, antigenicity, half-life, tertiary structure, and targeting to receptors on appropriate cells.

[0011] In another aspect, the present invention relates to a variant of a spasmolytic polypeptide (SP) which is a fragment of human spasmolytic polypeptide (HSP) or porcine spasmolytic polypeptide (PSP) comprising at least one trefoil domain.

[0012] The variant SP may be provided in both glycosylated and non-glycosylated form. It is at present contemplated that such a variant may be advantageous to use

instead of full-length SP because of a higher specific biological activity, increased solubility and stability, longer half-life, easier way of production, or the like.

[0013] It is assumed that other spasmolytic polypeptides than HSP will, if provided with a glycosylation site, also be expressed in predominantly glycosylated form. In a further aspect, the present invention therefore relates to a method of preparing a spasmolytic polypeptide in at least 60% glycosylated form, wherein a host cell transformed with a DNA fragment encoding a spasmolytic polypeptide and capable of providing glycosylation of said spasmolytic polypeptide is cultured under conditions permitting production of said spasmolytic polypeptide and recovering the resulting spasmolytic polypeptide from the culture.

#### DETAILED DESCRIPTION OF THE INVENTION

[0014] It has been found that, at least when recombinant HSP is produced in yeast, the proportion of it that is provided in glycosylated form is in N-glycosylated form. It has further been found that glycosylation takes place at Asn15 of the sequence shown above. In preferred embodiments of glycosylated HSP, the glycosylated side chain contains at least one hexose unit. In particular, the glycosylated side chain may contain at least one mannose unit, preferably at least five mannose units, most preferably at least ten mannose units. In one preferred embodiment of glycosylated HSP of the invention, the glycosylated side chain contains 13-17 mannose units. In other preferred embodiments, the glycosylated HSP is in addition glycosylated with at least one unit of N-acetyl glucosamine (GlcNAc). In the currently preferred embodiment, the glycosylated HSP is glycosylated at Asn15 with (GlcNAc)<sub>2</sub>(Man)<sub>10-15</sub>.

[0015] It is further contemplated to produce homologues of HSP which are provided with one or more additional glycosylation sites. Thus, the present invention also relates to HSP homologues, wherein Lys2 is replaced by Asn, Gln7 is replaced by Asn, Arg10 is replaced by Asn, Gly 20 is replaced by Thr or Ser, Gly23 is replaced by Asn, Ile 24 is replaced by Asn, Phe 36 is replaced by Asn, Asp 37 is replaced by Asn, Ser39 is replaced by Asn, Gln53 is replaced by Asn, Glu61 is replaced by Asn, Asp64 is replaced by Asn, Arg66 is replaced by Thr or Ser, Gly69 is replaced by Thr or Ser, Gly72 is replaced by Asn, Ile 89 is replaced by Thr or Ser, Pro98 is replaced by Asn or Val101 is replaced by Thr or Ser, or a combination of two or more of these substitutions. In a currently preferred embodiment of such an HSP homologue, Asp64 is replaced by Asn, and Arg66 is replaced by Thr or Ser.

[0016] It is of course understood that HSP homologues of the invention may be glycosylated in the same manner at one or more of these sites as described above for glycosylation at Asn15.

[0017] It is assumed that the trefoil structure common among spasmolytic polypeptides is important for the function of HSP and PSP. The variant human or porcine SP comprising a fragment of the full-length polypeptide should therefore include at least three disulfide bonds to provide this structure. Consequently, the variant may comprise at least a sequence of amino acids from position 8 to 46 or from position 58 to 95, each of which sequences defines a trefoil domain of HSP and PSP.

[0018] As indicated above, the SP variant of the invention may be provided in non-glycosylated form. This may, for instance, be accomplished by substituting Asn15 by another amino acid, e.g. Asp or Glu, or by substituting Thr17 by another amino acid except Ser, e.g. Ala. It is more likely, however, that one or more additional glycosylation sites will be introduced into this domain, for instance by replacing Arg10 by Asn, Gly 20 by Thr or Ser, Gly23 by Asn, Ile 24 by Asn, Phe 36 by Asn, Asp 37 by Asn, or Ser39 by Asn, or a combination of two or more of these substitutions.

[0019] On the other hand, it may be desirable to provide the trefoil domain from position 58 to 95 with a glycosylation site lacking in this domain in native HSP and PSP. Thus, Glu61 may be replaced by Asn, Asp64 by Asn, Arg66 by Thr or Ser, Gly69 by Thr or Ser, or Gly72 is replaced by Asn, or a combination of two or more of these substitutions. In a currently preferred embodiment of the variant, Asp64 is replaced by Asn, and Arg66 is replaced by Thr or Ser.

[0020] It is of course understood that variants of the invention may be glycosylated in the same manner at one or more of these sites as described above for glycosylation at Asn15 in full-length HSP.

[0021] A DNA sequence encoding HSP may suitably be isolated from a human genomic DNA library by PCR (polymerase chain reaction) cloning using primers based on the published cDNA sequence [8]. Alternatively, the DNA sequence may be prepared synthetically by established standard methods, e.g. the phosphoamidite method described by S. L. Beaucage and M. H. Caruthers, *Tetrahedron Letters* 22, 1981, pp. 1859-1869, or the method described by Matthes et al., *EMBO Journal* 3, 1984, pp. 801-805. According to the phosphoamidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in suitable vectors. The cDNA sequence shown in [8] may be used as the basis of oligonucleotide synthesis.

[0022] Alternatively, it is possible to use cDNA coding for HSP obtained by screening a human cDNA library with oligonucleotide probes in accordance with well-known procedures.

[0023] Furthermore, the DNA sequence may be of mixed synthetic and genomic, mixed synthetic and cDNA or mixed genomic and cDNA origin prepared by ligating fragments of genomic, synthetic or cDNA origin (as appropriate), the fragments corresponding to various parts of the entire DNA sequence, in accordance with standard techniques.

[0024] The SP variant of the invention may be encoded by a fragment of the full-length DNA sequence, prepared by one of the methods indicated above, or by suitably truncating the full-length sequence.

[0025] The DNA sequence encoding HSP or an SP variant of the invention may then be inserted in a suitable expression vector. The recombinant expression vector may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the

host cell genome and replicated together with the chromosome(s) into which it has been integrated.

[0026] In the vector, the DNA sequence encoding HSP or an SP variant of the invention should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA encoding the inhibitor of the invention in mammalian cells are the SV 40 promoter (Subramani et al., *Mol. Cell Biol.* 1, 1981, pp. 854-864), the MT-1 (metallothionein gene) promoter (Palmiter et al., *Science* 222, 1983, pp. 809-814) or the adenovirus 2 major late promoter. Suitable promoters for use in yeast host cells include promoters from yeast glycolytic genes (Hitzeman et al., *J. Biol. Chem.* 255, 1980, pp. 12073-12080; Alber and Kawasaki, *J. Mol. Appl. Gen.* 1, 1982, pp. 419-434) or alcohol dehydrogenase genes (Young et al., in *Genetic Engineering of Microorganisms for Chemicals* (Hollaender et al., eds.), Plenum Press, New York, 1982), or the TPI1 (U.S. Pat. No. 4,599,311) or ADH2-4c (Russell et al., *Nature* 304, 1983, pp. 652-654) promoters. Suitable promoters for use in filamentous fungus host cells are, for instance, the ADH3 promoter (McKnight et al., *The EMBO J.* 4, 1985, pp. 2093-2099) or the *tpiA* promoter.

[0027] The DNA sequence encoding HSP or an SP variant may also be operably connected to a suitable terminator, such as the human growth hormone terminator (Palmiter et al., op. cit.) or (for fungal hosts) the TPI1 (Alber and Kawasaki, op. cit.) or ADH3 (McKnight et al., op. cit.) promoters. The vector may further comprise elements such as polyadenylation signals (e.g. from SV 40 or the adenovirus 5' Elb region), transcriptional enhancer sequences (e.g. the SV 40 enhancer) and translational enhancer sequences (e.g. the ones encoding adenovirus VA RNAs).

[0028] The recombinant expression vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. An example of such a sequence (when the host cell is a mammalian cell) is the SV 40 origin of replication, or (when the host cell is a yeast cell) the yeast plasmid 2  $\mu$  replication genes REP 1-3 and origin of replication. The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or one which confers resistance to a drug, e.g. neomycin, hygromycin or methotrexate, or the *Schizosaccharomyces pombe* TPI gene (described by P. R. Russell, *Gene* 40, 1985, pp. 125-130).

[0029] The procedures used to ligate the DNA sequences coding for HSP or the SP variant, the promoter and the terminator, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y., 1989).

[0030] The host cell into which the expression vector is introduced may be any cell which is capable of producing the inhibitor of the invention and is preferably a eukaryotic cell, such as a mammalian, yeast or fungal cell.

[0031] The yeast organism used as the host cell may be any yeast organism which, on cultivation, produces large

quantities of the inhibitor of the invention. Examples of suitable yeast organisms are strains of the yeast species *Saccharomyces cerevisiae*, *Saccharomyces kluyveri*, *Schizosaccharomyces pombe* or *Saccharomyces uvarum*. The transformation of yeast cells may for instance be effected by protoplast formation followed by transformation in a manner known per se.

[0032] Examples of suitable mammalian cell lines are the COS (ATCC CRL 1650), BHK (ATCC CRL 1632, ATCC CCL 10) or CHO (ATCC CCL 61) cell lines. Methods of transfecting mammalian cells and expressing DNA sequences introduced in the cells are described in e.g. Kaufman and Sharp, *J. Mol. Biol.* 159, 1982, pp. 601-621; Southern and Berg, *J. Mol. Appl. Genet.* 1, 1982, pp. 327-341; Loyer et al., *Proc. Natl. Acad. Sci. USA* 79, 1982, pp. 422-426; Wigler et al., *Cell* 14, 1978, p. 725; Corsaro and Pearson, *Somatic Cell Genetics* 7, 1981, p. 603; Graham and van der Eb, *Virology* 52, 1973, p. 456; and Neumann et al., *EMBO J.* 1, 1982, pp. 841-845.

[0033] Alternatively, fungal cells may be used as host cells. Examples of suitable fungal cells are cells of filamentous fungi, e.g. *Aspergillus* spp. or *Neurospora* spp., in particular strains of *Aspergillus oryzae* or *Aspergillus niger*. The use of *Aspergillus* spp. for the expression of proteins is described in, e.g., EP 238 023.

[0034] According to the present method, yeast cells are currently preferred for producing HSP and other SPs (such as those shown in FIG. 1), as they have surprisingly been found to produce SP in a high yield and in at least 60% glycosylated form. For instance, about two thirds of the HSP produced by yeast may be recovered in glycosylated form.

[0035] The medium used to cultivate the cells may be any conventional medium suitable for growing mammalian cells or fungal (including yeast) cells, depending on the choice of host cell. The spasmolytic polypeptide will be secreted by the host cells to the growth medium and may be recovered therefrom by conventional procedures including separating the cells from the medium by centrifugation or filtration, precipitating the proteinaceous components of the supernatant or filtrate by means of a salt, e.g. ammonium sulfate, purification by a variety of chromatographic procedures, e.g. ion exchange chromatography or affinity chromatography, or the like.

[0036] The present invention also relates to a pharmaceutical composition comprising HSP or a variant spasmolytic polypeptide of the invention together with a pharmaceutically acceptable carrier or excipient. In the composition of the invention, the variant may be formulated by any of the established methods of formulating pharmaceutical compositions, e.g. as described in *Remington's Pharmaceutical Sciences*, 1985. The composition may typically be in a form suited for oral or rectal administration and may, as such, be formulated as tablets or suppositories.

[0037] HSP or an SP variant of the invention is contemplated to be useful for the prophylaxis or treatment of gastrointestinal disorders. More specifically, it is contemplated for the treatment of gastric or peptic ulcers, inflammatory bowel disease, Crohn's disease or injury to the intestinal tract caused by radiation therapy, bacterial or other infections, etc.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0038] The invention is further illustrated in the following example with reference to the appended drawings in which

[0039] FIG. 1 shows the trefoil family of peptides. Intestinal trefoil factor (ITF) contains one trefoil domain [2], as does the breast cancer associated pS2 peptide [3,4]. The spasmolytic polypeptides from man, pig and mouse contain two trefoil domains [1,8]. Spasmodolysins from *Xenopus laevis* contain one or four trefoil domains [10]. Recently, a member of the frog trefoil family containing two domains has been described [11].

[0040] FIG. 2 shows the proposed structure of human spasmolytic polypeptide, HSP. The primary amino acid sequence is taken from Tomasetto et al. [8], and the disulfide bonds are placed in homology to PSP [1].

[0041] FIG. 3 shows the nucleotide sequence (SEQ ID NO:2) and corresponding amino acid sequence (SEQ ID NO: 3) of the 563 bp EcoRI-XbaI fragment encoding the leader-HSP fusion protein. The Kex 2 processing site is indicated by a vertical arrow. The leader and the PCR cloned parts of the HSP gene are shown in capital letters, while the synthetic parts are shown in small letters. The underlined sequences correspond to the PCR primers with horizontal arrows indicating the direction. Restriction sites relevant for the construction are shown.

[0042] FIG. 4 shows the *S. cerevisiae* plasmid for the expression and secretion of HSP. TPI-prom. and TPI-term. are *S. cerevisiae* triosephosphate isomerase transcription promoter and terminator sequences, respectively. POT is a selective marker, the *Schizosaccharomyces pombe* triosephosphate isomerase gene. Only restriction sites relevant for the construction of the plasmid have been indicated.

[0043] FIG. 5 shows reversed-phase HPLC on a Vydac 214TP54 column of yeast fermentation broth. The two peaks corresponding to r-HSP and glycosylated r-HSP are indicated. The dashed line shows the concentration of acetonitrile in the eluting solvent.

[0044] FIG. 6 shows ion exchange chromatography on a Fast Flow S column of concentrated yeast supernatant. The amount of r-HSP and glycosylated r-HSP were determined by the use of the HPLC system shown in FIG. 5. The bars indicate the fractions pooled for further purification of r-HSP and glycosylated r-HSP. The dashed line shows the concentration of NaCl in the eluting solvent. For details, see Material and Methods.

[0045] FIG. 7 shows the final purification of r-HSP (A) and glycosylated r-HSP (B) on a preparative reversed-phase HPLC Vydac 214TP1022 column. The bars indicate the fractions pooled for lyophilization. The dashed lines show the concentration of acetonitrile in the eluting solvent. For details, see Material and Methods.

[0046] FIG. 8 shows reversed-phase HPLC on a Vydac 214TP54 column of purified, glycosylated r-HSP (A) and r-HSP (B). The dashed lines show the concentration of acetonitrile in the eluting solvent.

[0047] FIG. 9 shows mass spectra of purified r-HSP (A and B) and glycosylated r-HSP (C and D). FIG. A and FIG. C show the original mass spectrum of r-HSP and glycosylated r-HSP, respectively. FIG. B and FIG. D show the

reconstructed mass spectrum for r-HSP and glycosylated r-HSP on the basis of FIG. A and FIG. C.

## EXAMPLE

## Material and Methods

## [0048] General Methods

[0049] Standard DNA techniques were used as previously described [29]. Synthetic oligonucleotides were prepared on an automatic DNA synthesizer (380B, Applied Biosystems) using commercially available reagents. DNA sequence determinations were performed by the dideoxy chain-termination technique [30]. Polymerase chain reactions (PCR) were performed on a DNA Thermal Cycler (Perkin Elmer Cetus) using a commercial kit (GeneAmp, Perkin Elmer Cetus).

## [0050] PCR Cloning of HSP

[0051] The first trefoil domain of HSP was isolated by a PCR reaction in which 1  $\mu$ g human genomic DNA (Clontech, Palo Alto, Calif., USA) was used as a template. The reaction mixture contained 100 pmole each of the forward primer 1 (GGCTGAGCCCCATAACAG) (SEQ ID NO:4) and reverse primer 2 (TGGAAACACCAGGGGAC) (SEQ ID NO:5) (FIG. 3) and was carried out in a 100  $\mu$ l volume. The cycle was: 94° C. for 1 min, 50° C. for 1 min, and 72° C. for 1 min. After 30 cycles a final cycle was performed in which the 72° C. step was maintained for 10 min. The PCR product, a 115 bp fragment, was isolated by electrophoresis on a 2% agarose gel.

[0052] The 115 bp PCR fragment was digested with DdeI and then ligated to a 31 bp duplex formed from the oligonucleotides (GAGAAACCCTCCCCCTGCCAGTGCTC-CAGGC) (SEQ ID NO:6) and (TCAGCCTGGAGCACTGGCAGGGGGAGGGTTTCTC). The ligation product was amplified by PCR using forward primer 3 (GCTGAGAGAT-TGGAGAAGAGAGAGAAACCCTCCCCCT) (SEQ ID NO:7) and reverse primer 2. The 3' part of primer 3 is identical to the N-terminal encoding part of the HSP gene and the 5' part of primer 3 is identical to the C-terminal encoding part of the hybrid leader gene (FIG. 3). In-frame fusion of the hybrid leader gene and the first trefoil domain from HSP was obtained by overlay extension PCR [31]. The product was digested with EcoRI and AvaII and isolated as a 360 bp DNA fragment.

[0053] The second trefoil domain of HSP was PCR-cloned from human genomic DNA as described for the first domain by replacing primers 1 and 2 with forward primer 4 (TGCGTCATGGAGGTCTC) (SEQ ID NO:8) and reverse primer 5 (AGCACCATGGCACTTCAAAG) (SEQ ID NO:9) (FIG. 3). Reverse primer 5 introduces a NcoI site as a silent mutation. The PCR product, a 115 bp fragment, was isolated and digested with DdeI and NcoI resulting in a 91 bp fragment. To this fragment were ligated two synthetic duplexes. The first, encoding the amino acid sequence between the two trefoil domains, consisted of the oligonucleotides (GTCCCCTGGTGTTCACCCCCCTC-CCAAAGCAAGAGTCGGATCAGTGCGT-CATGGAGGTC) (SEQ ID NO:10) and (TGAGACCTCCATGACGCACTGATC-CGACTCTGCTTTGGGAGGGGGTGGAAA-CACCAGGG) (SEQ ID NO:11). The second, a 46 bp

NcoI-XbaI fragment encoding the C-terminal part of HSP, consisted of the oligonucleotides (CATGGTGCTTCTCCCGAACTCTGTGGAAGACTGCCATTACTAAGT) (SEQ ID NO:12) and (CTAGACTTAGTAATGGCAGTCTTCCACAGAGTTCCGGGAAGAAGCAC) (SEQ ID NO:13). After AvaII digestion a 195 bp AvaII-XbaI fragment was isolated.

**[0054]** A DNA construct encoding the hybrid leader fused in-frame to the entire HSP gene was obtained by ligation of the 360 bp EcoRI-AvaII fragment and the 195 bp AvaII-XbaI fragment described above to the 2.7 kb EcoRI-XbaI fragment from vector pTZ19R [32]. This construct was then transformed into *E. coli* strain MT-172 ( $r^-$ ,  $m^+$ ) by selection for resistance to ampicillin. DNA sequencing of the resulting plasmid, KFN-1843, showed that the correct construction had been obtained.

**[0055]** Construction of the HSP Secreting Yeast Strain

**[0056]** Plasmid KFN-1843 described above was digested with EcoRI and XbaI. The resulting 558 bp fragment was isolated and ligated to the 9.3 kb NcoI-XbaI fragment and the 1.6 kb NcoI-EcoRI fragment both from the yeast expression vector pMT-636. Plasmid pMT-636 is derived from the *S. cerevisiae*-*E. coli* shuttle vector CPOT [25,33] by deletion of the 0.4 kb HpaI-NruI fragment from the Leu-2 gene. The ligation mixture was transformed into *E. coli* strain MT-172, and the HSP expression plasmid, KFN-1847, was isolated (**FIG. 4**). Plasmid pKFN-1847 was transformed into *S. cerevisiae* strain MT-663 by selection for growth on glucose as the sole carbon source. One transformant, KFN-1852, was selected for fermentation.

**[0057]** Fermentation

**[0058]** The transformant described above was cultivated at 30° C. for 3 days in yeast peptone dextrose (YPD) medium [40] supplied with additional yeast extract (60 g/l). An OD 650 nm value of 52 was reached at the end of the fermentation.

**[0059]** Purification of r-HSP

**[0060]** The concentration of r-HSP in the yeast fermentation broth and fractions obtained during the purification was measured by analytical HPLC. Aliquots (usually 50-200  $\mu$ l) were injected onto a Vydac 214TP54 reverse-phase C4 HPLC column (0.46x25 cm) equilibrated at 30° C. at a flow rate of 1.5 ml/min with 0.1% (v/v) TFA in 5% (v/v) acetonitrile. The concentration of acetonitrile in the eluting solvent was raised to 65% (v/v) over 30 min. Absorbance was measured at 280 nm. The peaks eluting at 15.6 min. and 16.1 min. (**FIG. 5**) was found by mass spectrometry analysis to represent glycosylated r-HSP and unglycosylated r-HSP, respectively. The peptides were quantified using a calibrated PSP sample as standard as both peptides contain two Trp and two Tyr out of 106 amino acid residues.

**[0061]** From a 10 litre fermentor, 8 litres of fermentation broth was isolated by centrifugation at 3,000 rpm for 10 min. The supernatant was concentrated to 0.9 litre using an Amicon ultrafiltration unit (RA 2000) equipped with an Amicon spiral ultrafiltration cartridge type S1Y3, MW cut-off 3,000 (Product No. 540620). The pH was adjusted to 1.7 and the conductivity in the resulting concentrated sample was measured to 4.7 mS.

**[0062]** The sample was pumped onto a Fast Flow S-Sepharose (Pharmacia) column (5x11 cm) with a flow rate of 40 ml/h. Previous to the application, the column was

equilibrated in 50 mM formic acid buffer, pH=3.7. After application of the sample, the column was washed with 500 ml of 50 mM formic acid buffer, pH=3.7. The peptides were eluted from the column by a linear gradient between 1.5 litres of 50 mM formic acid buffer, pH=3.7 and 1.5 litres of 50 mM formic acid buffer, pH=3.7 containing 0.6 M NaCl. Fractions of 10 ml were collected at a flow rate of 40 ml/h and the absorbance was measured at 280 nm. Fractions were assayed for the content of r-HSP and glycosylated r-HSP in the HPLC-system previously described. The elution profile is shown in **FIG. 6**. Fractions corresponding to r-HSP (fract. Nos. 107-128) and glycosylated r-HSP (fract. Nos. 78-95), respectively, were pooled.

**[0063]** Glycosylated r-HSP and r-HSP were further purified by preparative HPLC chromatography. Pooled fractions (approx. 200 ml) were pumped onto a Vydac 214TP1022 C4 column (2.2x25 cm) equilibrated in 0.1% (v/v) TFA. The column was washed with 100 ml of 0.1% (v/v) TFA in 10% (v/v) MeCN. The peptides were eluted at 25° C. and at a flow rate of 5 ml/min with a linear gradient (650 ml) formed from MeCN/H<sub>2</sub>O/TFA (10.0:89.9:0.1 v/v/v) and MeCN/H<sub>2</sub>O/TFA (60.0:39.9:0.1 v/v/v). UV-absorption was monitored at 280 nm, and fractions corresponding to 10 ml were collected and analysed for the content of r-HSP or glycosylated r-HSP. **FIG. 7** shows the preparative HPLC purification of r-HSP (**FIG. 7A**) and glycosylated r-HSP (**FIG. 7B**). Fractions corresponding to the bars were pooled, and the volume reduced to 30% by vacuum centrifugation. From the two resulting pools, r-HSP and glycosylated r-HSP were isolated by lyophilization.

**[0064]** Characterization of r-HSP and Glycosylated r-HSP

**[0065]** Amino acid composition analysis were carried out by hydrolysis of 50  $\mu$ g peptide with 6M HCl for 24 h at 110° C. as previously described [6]; no correction for loss during hydrolysis was carried out. Amino acid sequence analysis was determined by automated Edman degradation using an Applied Biosystems Model 470A gas-phase sequencer [22]. Carbohydrate composition analysis was carried out by hydrolysis of 50  $\mu$ g peptide with 2M HCl for 1 h, 2 h and 4 h at 100° C. and monosaccharides were separated on a CarboPac PAI (Dionex, Sunnyvale, Calif.) column (4x250 mm) eluted with 14 mM NaOH. The monosaccharides were detected by pulsed amperometric detection (Dionex PAD-detector). The amount of monosaccharides was corrected to zero time of hydrolysis and calculated as nmol of monosaccharide per nmol of peptide.

**[0066]** Mass spectrometry analysis was performed using an API III LC/MS/MS system (Sciex, Thornhill, Ontario, Canada). The triple quadrupole instrument has a mass-to-charge ( $m/z$ ) range of 2400 and is fitted with a pneumatically assisted electrospray (also referred to as ion-spray) interface [23,24]. Sample introduction was done by a syringe infusion pump (Sage Instruments, Cambridge, Mass.) through a fused capillary (75  $\mu$ m i.d.) with a liquid flow-rate set at 0.5-1  $\mu$ l/min. The instrument  $m/z$  scale was calibrated with the singly-charged ammonium adduct ions of poly(propylene glycols) (PPG's) under unit resolution.

**[0067]** The accuracy of mass measurements was generally better than 0.02%.

## Results

**[0068]** Expression and Purification

**[0069]** DNA fragments encoding the two trefoil domains of HSP were isolated by PCR from human genomic DNA

using primers based on the published cDNA sequence [8]. The full length HSP gene was obtained from the PCR cloned fragments by addition of synthetic DNA fragments. The HSP gene was fused in-frame to a hybrid yeast leader sequence by overlay extension PCR [31] (FIG. 3). The hybrid leader is based on the mouse salivary amylase signal peptide [34] and the *S. kluyveri*  $\alpha$  mating factor leader sequence [35] and is further modified near the Kex 2 cleavage site for efficient processing [36, 41].

[0070] The yeast expression plasmid pKFN-1847 contains the leader-HSP gene inserted between the *S. cerevisiae* triose phosphate isomerase promoter and terminator [37]. The expression vector (FIG. 4) also contains the *Schizosaccharomyces pombe* TPI gene (POT) [38].

[0071] The plasmid was transformed into the yeast strain MT-663, carrying a deletion in the TPI gene, by selecting for growth on glucose.

[0072] The expression level of r-HSP in the present yeast system is approx. 120 mg/l. As can be seen from FIG. 5, the yeast supernatant contains two forms of r-HSP; one eluting at  $R_t=15.6$  min. and one eluting at  $R_t=16.1$  min. These two forms were purified separately, and by using the analytical HPLC-system (FIG. 5), these two forms can be quantified individually during the different steps of the purification.

[0073] After the initial concentration of the yeast supernatant by ultrafiltration, the first purification step was cationic exchange chromatography on a Fast Flow S column. FIG. 6 shows the elution profile from the column including the amount of r-HSP and glycosylated r-HSP determined in the fractions. A complete separation of the two forms of r-HSP was obtained in this step.

[0074] The fractions from the Fast Flows S column were pooled as indicated in FIG. 6, and the two peptides were further purified by preparative HPLC (FIG. 7). The r-HSP and glycosylated r-HSP were recovered from the fractions indicated in FIG. 7A and FIG. 7B by vacuum centrifugation and lyophilization. The purification is summarized in Table 1. The overall yield of r-HSP and glycosylated r-HSP from 8 litres of fermentation broth was 160 mg and 219 mg corresponding to 50% and 34%, respectively.

[0075] Characterization of r-HSP and Glycosylated r-HSP

[0076] FIG. 8 shows the purity of r-HSP and glycosylated r-HSP as analysed by analytical HPLC. From these results none of the peptides looks completely pure. However, upon rechromatography of material eluting in the minor as well as the major peak, similar chromatograms were obtained for both peptides (results not shown). This seems to indicate that the double peak observed for both r-HSP and glycosylated r-HSP reflects an atypical behaviour of these peptides on reverse phase columns rather than impurities in the preparations.

[0077] Table 2 shows the amino acid sequencing results obtained on r-HSP and glycosylated r-HSP. The average repetitive yield was 94.4% (r-HSP) and 94.6% (glycosylated r-HSP), respectively. In both cases the first 40 residues of the two peptides were confirmed by the sequence analysis. In the glycosylated HSP, no PTH-a.a. was found in Edman degradation cycle No. 15. The HSP sequence from residue 15-17 (Asn-Arg-Thr) corresponds to a consensus sequence for N-glycosylation of Asn-15.

[0078] The carbohydrate composition analysis of glycosylated r-HSP showed the presence of 12.8 nmol mannose (Man) and 1.6 nmol of N-acetyl glucoseamine (GlcNAc) per nmol of r-HSP. By peptide mapping of r-HSP and glycosylated r-HSP in combination with mass spectrometry and sequencing analysis (results not shown), no other residue besides Asn-15 of the glycosylated r-HSP was found to be modified, i.e. no O-glycosylation was found.

[0079] In FIG. 9, the electro-spray mass spectrometry (ESMS) analysis is shown for r-HSP and glycosylated r-HSP. FIGS. 9A and 9C are original mass spectra displaying characteristic series of multiply charged protonated ions always observed in ESMS spectra of proteins. FIGS. 9B and 9D are the corresponding computer reconstructed mass spectra from which the molecular weight of individual components may be read directly. As can be seen from FIG. 9B, the MW found for r-HSP is  $11961.5 \pm 2$  which is in very good agreement with a calculated mass of 11961.3. FIG. 9D shows the reconstructed ion spray mass spectrum of the glycosylated r-HSP. From the sequence analysis and the carbohydrate composition analysis, it is known that only Asn-15 is glycosylated and that only two monosaccharide residues, mannose and N-acetyl glucoseamine, occur in the glycosylated form of r-HSP. From these results in combination with the mass spectrometry data, it is possible to deduce the different glycosylated forms of r-HSP (Table 3).

[0080] Molecular weights corresponding to two series of carbohydrate side chains can be deduced from the combination of carbohydrate composition data and ISMS-data, namely (GlcNAc)<sub>2</sub>(Hex)<sub>10-15</sub> and (Hex)<sub>13-17</sub> (Table 3). As mannose is the only hexose in the glycosylated r-HSP, and as Asn-15 is the only glycosylated residue, it seems reasonable to conclude that the structure of the glycosylation site is Asn-(GlcNAc)<sub>2</sub>(Man)<sub>10-15</sub>. The observed Asn-(Hex)<sub>13-17</sub> forms are thus most likely to arise from fragmentation in the mass spectrometer, by which the two GlcNAc residues lose an acetyl group and are converted into two hexoses.

[0081] The structure of Asn-(GlcNAc)<sub>2</sub>(Man)<sub>10-15</sub> has previously been reported as high mannose type of N-glycosylation for other peptides and proteins expressed in yeast [26].

TABLE 1

| STEP                        | Purification of r-hSP and glycosylated r-hSP from yeast supernatant |             |                       |           |                       |
|-----------------------------|---|-------------|-----------------------|-----------|-----------------------|
|                             | Vol-<br>[ml]  | Amount [mg] |                       | Yield [%] |                       |
|                             |   | r-hSP       | glycosylated<br>r-hSP | r-hSP     | glycosylated<br>r-hSP |
| Yeast supernatant           | 8000  | 320         | 640                   | 100       | 100                   |
| Ultrafiltration             | 900   | 207         | 405                   | 65        | 63                    |
| Ion exchange chromatography | Pool 1  | 160         | 275                   |           | 43                    |
|                             | Pool 2  | 220         | 182                   | 57        |                       |
| Prep HPLC                   | Pool 1  | 54          | 219                   |           | 34                    |
|                             | Pool 2  | 80          | 160                   | 50        |                       |



[0082]

TABLE 2

| Amino acid sequence analysis of r-hSP and glycosylated r-hSP |          |              |                    |
|--|----------|--------------|--------------------|
| Cycle No.  | PTH-a.a. | Yield (pmol) |                    |
|  |          | r-hSP        | glycosylated r-hSP |
| 1  | Glu      | 4304         | 8853               |
| 2  | Lys      | 6925         | 8292               |
| 3  | Pro      | 6027         | 12837              |
| 4  | Ser      | 2890         | 5602               |
| 5  | Pro      | 4336         | 8802               |
| 6  | (Cys)    | ND           | ND                 |
| 7  | Gln      | 3388         | 5689               |
| 8  | (Cys)    | ND           | ND                 |
| 9  | Ser      | 1279         | 2417               |
| 10   | Arg      | 1876         | 2523               |
| 11   | Leu      | 2277         | 4290               |
| 12   | Ser      | 877          | 1790               |
| 13   | Pro      | 1545         | 2963               |
| 14   | His      | 517          | 574                |
| 15   | Asn      | 1202         | 0*                 |
| 16   | Arg      | 959          | 1471               |
| 17   | Thr      | 978          | 2172               |
| 18   | Asn      | 1066         | 1509               |
| 19   | (Cys)    | ND           | ND                 |
| 20   | Gly      | 836          | 1857               |
| 21   | Phe      | 993          | 1958               |
| 22   | Pro      | 843          | 1839               |
| 23   | Gly      | 785          | 2049               |
| 24   | Ile      | 640          | 1400               |
| 25   | Thr      | 589          | 1454               |
| 26   | Ser      | 274          | 621                |
| 27   | Asp      | 581          | 1391               |
| 28   | Gln      | 445          | 952                |
| 29   | (Cys)    | ND           | ND                 |
| 30   | Phe      | 623          | 1562               |
| 31   | Asp      | 483          | 1210               |
| 32   | Asn      | 369          | 823                |
| 33   | Gly      | 359          | 885                |
| 34   | (Cys)    | ND           | ND                 |
| 35   | (Cys)    | ND           | ND                 |
| 36   | Phe      | 422          | 1094               |
| 37   | Asp      | 268          | 783                |
| 38   | Ser      | 127          | 324                |
| 39   | Ser      | 145          | 394                |
| 40   | Val      | 298          | 827                |

ND: Not determined

\*No trace of PTH-Asn or PTH-Asp was seen in cycle No. 15 of glycosylated r-hSP.

[0083]

TABLE 3

| Mass analysis of glycosylated r-hSP |               |                            |
|-------------------------------------|---------------|----------------------------|
| Structure                           | Calculated MW | MW found by ESMS (FIG. 9D) |
| hSP + 2 GlcNAc + 10 Man             | 13989.1       | 13989.5                    |
| hSP + 2 GlcNAc + 11 Man             | 14151.2       | 14151.0                    |
| hSP + 2 GlcNAc + 12 Man             | 14313.4       | 14313.5                    |
| hSP + 2 GlcNAc + 13 Man             | 14475.5       | 14475.0                    |
| hSP + 2 GlcNAc + 14 Man             | 14639.7       | 14639.5                    |
| hSP + 2 GlcNAc + 15 Man             | 14799.8       | 14801.5                    |
| hSP + 13 Man                        | 14069.1       | 14072.0                    |
| hSP + 14 Man                        | 14231.3       | 14232.5                    |
| hSP + 15 Man                        | 14393.4       | 14393.0                    |
| hSP + 16 Man                        | 14555.5       | 14557.5                    |
| hSP + 17 Man                        | 14717.7       | 14720.0                    |

## REFERENCES

- [0084] [1] Thim, L. (1989) FEBS Lett. 250, 85-90.
- [0085] [2] Suemori, S., Lynch-Devaney, K. and Podolsky, D. K. (1991) Proc. Natl. Acad. Sci. USA 88, 11017-11021.
- [0086] [3] Jakowlew, S. B., Breathnach, R., Jeltsch, J. M., Masiakowski, P. and Chambon, P. (1984) Nucleic Acids Res. 12, 1861-2878.
- [0087] [4] Prud'homme, J. -F., Fridlansky, F., Le Cunff, M., Atger, M., Mercier-Bodart, C., Pichon, M. -F. and Milgrom, E. (1985) DNA 4, 11-21.
- [0088] [5] Rio, M. C., Bellocq, J. P., Daniel, J. Y., Tomasetto, C., Lathe, R., Chenard, M. P., Batzenschlager, A. and Chambon, P. (1988) Science 241, 705-708.
- [0089] [6] Thim, L., Thomsen, J., Christensen, M. and Jørgensen, K. H. (1985) Biochim. Biophys. Acta 827, 410-418.
- [0090] [7] Rose, K., Savoy, L. -A., Thim, L., Christensen, M. and Jørgensen, K. H. (1989) Biochim. Biophys. Acta 998, 297-300.
- [0091] [8] Tomasetto, C., Rio, M. -C., Gautier, C., Wolf, C., Hareuveni, M., Chambon, P. and Lathe, R. (1990) EMBO J. 9, 407-414.
- [0092] [9] Hoffmann, W. (1988) J. Biol. Chem. 263, 7686-7690.
- [0093] [10] Hauser, F. and Hoffmann, W. (1991) J. Biol. Chem. 266, 21306-21309.
- [0094] [11] Hauser, F., Roeben, C. and Hoffmann, W. (1992) J. Biol. Chem. 267, 14451-14455.
- [0095] [12] Jørgensen, K. H., Thim, L. and Jacobsen, H. E. (1982) Regul. Peptides 3, 207-219.
- [0096] [13] Thim, L., Jørgensen, K. H. and Jørgensen, K. D. (1982) Regul. Peptides 3, 221-230.
- [0097] [14] Rasmussen, T. N., Raabjerg, L., Poulsen, S. S., Thim, L. and Holst, J. J. (1992) Histochemistry 98, 113-119.
- [0098] [15] Frandsen, E. K., Jørgensen, K. H. and Thim, L. (1986) Regul. Peptides 16, 291-297.
- [0099] [16] Frandsen, E. K. (1988) Regul. Peptides 20, 45-52.
- [0100] [17] Rasmussen, T. N., Raabjerg, L., Poulsen, S. S., Thim, L. and Holst, J. J. (1992) Am. J. Physiol. (in press)
- [0101] [18] Jørgensen, K. D., Diamant, B., Jørgensen, K. H. and Thim, L. (1982) Regul. Peptides 3, 231-243.
- [0102] [19] Hoosein, N. M., Thim, L., Jørgensen, K. H. and Brattain, M. G. (1989) FEBS Lett. 247, 303-306.
- [0103] [20] Wright, N. A., Poulsom, R., Stamp, G. W., Hall, P. A., Jeffery, R. E., Longcroft, J. M., Rio, M. -C., Tomasetto, C. and Chambon, P. (1990) J. Pathol. 162, 279-284.

- [0104] [21] Rio, M. -C., Chenard, M. -P., Wolf, C., Marcellin, L., Tomasetto, C., Lathe, R., Bellocq, J. P. and Chambon, P. (1991) *Gastroenterology* 100, 375-379.
- [0105] [22] Thim, L., Hansen, M. T. and Soerensen, A. R. (1987) *FEBS Lett.* 212, 307-312.
- [0106] [23] Bruins, A. P., Covey, T. R. and Henion, J. D. (1987) *Anal. Chem.* 59, 2642-2646.
- [0107] [24] Covey, T. R., Bonner, R. F., Shushan, B. I. and Henion, J. D. (1988) *Rapid Commun. Mass Spectrom.* 2, 249-256.
- [0108] [25] Thim, L., Hansen, M. T., Norris, K., Hoegh, I., Boel, E., Forstrom, J., Ammerer, G. and Fiil, N. P. (1986) *Proc. Natl. Acad. Sci. USA* 83, 6766-6770.
- [0109] [26] Poulter, L. and Burlingame, A. L. (1990) in: *Methods in Enzymology* (McCloskey, J. A., ed.) 193, 661-689. Academic Press, Inc., San Diego, Calif.
- [0110] [27] Gajhede, M., Thim, L., Jørgensen, K. H. and Melberg, S. G. (1992) *Proteins: Structure, Function, and Genetics* 13, 364-368.
- [0111] [28] Carr, M. D. (1992) *Biochemistry* 31, 1998-2004.
- [0112] [29] Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- [0113] [30] Sanger, F., Micklen, S., and Coulson, A. R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
- [0114] [31] Horton, R. M., Hunt, H. D., Ho, S. N., Pullen, J. K., and Pease, L. R. (1989) *Gene* 77, 61-68.
- [0115] [32] Mead, D. A., Szczesna-Skorupa, E. and Kemper, B. (1986) *Prot. Engin.* 1, 67-74.
- [0116] [33] Kawasaki, G. (1984) 12th International Conference on Yeast Genetics and Molecular Biology, Sep. 17-24, 1984, Edinburgh, Scotland, Abstr. P15.
- [0117] [34] Hagenbüchle, O., Tosi, M., Schibler, U., Bovey, R., Wellauer, P. K., and Young, R. A. (1981) *Nature* 289, 643-646.
- [0118] [35] Egel-Mitani, M. and Hansen, M. T. (1987) *Nucl. Acids Res.* 15, 6303-6304.
- [0119] [36] Christiansen, L. and Norris, K., personal communication.
- [0120] [37] Alber, T. and Kawasaki, G. (1982) *J. Mol. Appl. Genet.* 1, 419-434.
- [0121] [38] Russell, P. R. (1985) *Gene* 40, 125-130.
- [0122] [39] Jeltsch, J. M., Roberts, M., Schatz, C., Garnier, J. M., Brown, A. M. C., and Chambon, P. (1987) *Nucl. Acids. Res.* 15, 1401-1414.
- [0123] [40] Sherman, F., Fink, G. R. and Hicks, J. B. (1981) *Methods in Yeast Genetics*. Cold Spring Harbor Laboratory Press, New York.
- [0124] [41] Norris, K., Norris, F. and Bjoern, S. E. (1990) International Patent Application WO 90/10075.

---

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 14

<210> SEQ ID NO 1

<211> LENGTH: 106

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 1

Glu Lys Pro Ser Pro Cys Gln Cys Ser Arg Leu Ser Pro His Asn Arg  
1 5 10 15

Thr Asn Cys Gly Phe Pro Gly Ile Thr Ser Asp Gln Cys Phe Asp Asn  
20 25 30

Gly Cys Cys Phe Asp Ser Ser Val Thr Gly Val Pro Trp Cys Phe His  
35 40 45

Pro Leu Pro Lys Gln Glu Ser Asp Gln Cys Val Met Glu Val Ser Asp  
50 55 60

Arg Arg Asn Cys Gly Tyr Pro Gly Ile Ser Pro Glu Glu Cys Ala Ser  
65 70 75 80

Arg Lys Cys Cys Phe Ser Asn Phe Ile Phe Glu Val Pro Trp Cys Phe  
85 90 95

Phe Pro Asn Ser Val Glu Asp Cys His Tyr  
100 105

-continued

---

```

<210> SEQ ID NO 2
<211> LENGTH: 563
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (77)...(553)
<221> NAME/KEY: sig_peptide
<222> LOCATION: (77)...(235)
<221> NAME/KEY: mat_peptide
<222> LOCATION: (236)...(553)

<400> SEQUENCE: 2

gaattccatt caagaatagt tcaaacaaga agattacaaa ctatcaattt catacacaat      60
ataaacgacc aaaaga atg aag gct gtt ttc ttg gtt ttg tcc ttg atc gga      112
      Met Lys Ala Val Phe Leu Val Leu Ser Leu Ile Gly
                -50                -45

ttc tgc tgg gcc caa cca gtc act ggc gat gaa tca tct gtt gag att      160
Phe Cys Trp Ala Gln Pro Val Thr Gly Asp Glu Ser Ser Val Glu Ile
      -40                -35                -30

ccg gaa gag tct ctg atc atc gct gaa aac acc act ttg gct aac gtc      208
Pro Glu Glu Ser Leu Ile Ile Ala Glu Asn Thr Thr Leu Ala Asn Val
      -25                -20                -15                -10

gcc atg gct gag aga ttg gag aag aga gag aaa ccc tcc ccc tgc cag      256
Ala Met Ala Glu Arg Leu Glu Lys Arg Glu Lys Pro Ser Pro Cys Gln
                -5                1                5

tgc tcc agg ctg agc ccc cat aac agg acg aac tgc ggc ttc cct gga      304
Cys Ser Arg Leu Ser Pro His Asn Arg Thr Asn Cys Gly Phe Pro Gly
      10                15                20

atc acc agt gac cag tgt ttt gac aat gga tgc tgt ttc gac tcc agt      352
Ile Thr Ser Asp Gln Cys Phe Asp Asn Gly Cys Cys Phe Asp Ser Ser
      25                30                35

gtc act ggg gtc ccc tgg tgt ttc cac ccc ctc cca aag caa gag tcg      400
Val Thr Gly Val Pro Trp Cys Phe His Pro Leu Pro Lys Gln Glu Ser
      40                45                50                55

gat cag tgc gtc atg gag gtc tca gac aga aga aac tgt ggc tac ccg      448
Asp Gln Cys Val Met Glu Val Ser Asp Arg Arg Asn Cys Gly Tyr Pro
                60                65                70

ggc atc agc ccc gag gaa tgc gcc tct cgg aag tgc tgc ttc tcc aac      496
Gly Ile Ser Pro Glu Glu Cys Ala Ser Arg Lys Cys Cys Phe Ser Asn
                75                80                85

ttc atc ttt gaa gtg cca tgg tgc ttc ttc ccg aac tct gtg gaa gac      544
Phe Ile Phe Glu Val Pro Trp Cys Phe Phe Pro Asn Ser Val Glu Asp
                90                95                100

tgc cat tac taagtctaga      563
Cys His Tyr
      105

<210> SEQ ID NO 3
<211> LENGTH: 159
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: SIGNAL
<222> LOCATION: (1)...(53)

<400> SEQUENCE: 3

Met Lys Ala Val Phe Leu Val Leu Ser Leu Ile Gly Phe Cys Trp Ala
      -50                -45                -40

Gln Pro Val Thr Gly Asp Glu Ser Ser Val Glu Ile Pro Glu Glu Ser
      -35                -30                -25

```

-continued

Leu Ile Ile Ala Glu Asn Thr Thr Leu Ala Asn Val Ala Met Ala Glu  
 -20 -15 -10

Arg Leu Glu Lys Arg Glu Lys Pro Ser Pro Cys Gln Cys Ser Arg Leu  
 -5 1 5 10

Ser Pro His Asn Arg Thr Asn Cys Gly Phe Pro Gly Ile Thr Ser Asp  
 15 20 25

Gln Cys Phe Asp Asn Gly Cys Cys Phe Asp Ser Ser Val Thr Gly Val  
 30 35 40

Pro Trp Cys Phe His Pro Leu Pro Lys Gln Glu Ser Asp Gln Cys Val  
 45 50 55

Met Glu Val Ser Asp Arg Arg Asn Cys Gly Tyr Pro Gly Ile Ser Pro  
 60 65 70 75

Glu Glu Cys Ala Ser Arg Lys Cys Cys Phe Ser Asn Phe Ile Phe Glu  
 80 85 90

Val Pro Trp Cys Phe Phe Pro Asn Ser Val Glu Asp Cys His Tyr  
 95 100 105

<210> SEQ ID NO 4  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Synthetic

<400> SEQUENCE: 4

ggctgagccc ccataacag 19

<210> SEQ ID NO 5  
 <211> LENGTH: 17  
 <212> TYPE: DNA  
 <213> ORGANISM: Synthetic

<400> SEQUENCE: 5

tggaaacacc aggggac 17

<210> SEQ ID NO 6  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Synthetic

<400> SEQUENCE: 6

gagaaaccct cccctgcca gtgctccagg c 31

<210> SEQ ID NO 7  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Synthetic

<400> SEQUENCE: 7

tcagcctgga gcactggcag ggggagggtt tctc 34

<210> SEQ ID NO 8  
 <211> LENGTH: 37  
 <212> TYPE: DNA  
 <213> ORGANISM: Synthetic

<400> SEQUENCE: 8

gctgagagat tggagaagag agagaaacc tccccct 37

-continued

---

```

<210> SEQ ID NO 9
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 9

tgcgtcatgg aggtctc                                     17

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 10

agcaccatgg cacttcaaag                                   20

<210> SEQ ID NO 11
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 11

gtcccctggt gttccacc cctcccaaag caagatcgg atcagtgcgt catggagtc   60

<210> SEQ ID NO 12
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 12

tgagacctcc atgacgcact gatccgactc ttgctttggg aggggggtgga aacaccaggg   60

<210> SEQ ID NO 13
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 13

catggtgctt cttcccgaac tctgtggaag actgccatta ctaagt               46

<210> SEQ ID NO 14
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Synthetic

<400> SEQUENCE: 14

ctagacttag taatggcagt cttccacaga gttcgggaag aagcac               46
    
```

---

1. Human spasmolytic polypeptide (HSP) which has the amino acid sequence

(SEQ ID NO:1)

```

Glu Lys Pro Ser Pro Cys Gln Cys Ser Arg Leu Ser
Pro His Asn Arg Thr Asn Cys Gly Phe Pro Gly Ile
Thr Ser Asp Gln Cys Phe Asp Asn Gly Cys Cys Phe
Asp Ser Ser Val Thr Gly Val Pro Trp Cys Phe His
    
```

-continued

```

Pro Leu Pro Lys Gln Glu Ser Asp Gln Cys Val Met
Glu Val Ser Asp Arg Arg Asn Cys Gly Tyr Pro Gly
Ile Ser Pro Glu Glu Cys Ala Ser Arg Lys Cys Cys
Phe Ser Asn Phe Ile Phe Glu Val Pro Trp Cys Phe
Phe Pro Asn Ser Val Glu Asp Cys His Tyr
    
```

or a functionally equivalent homologue thereof, characterized by being in glycosylated form.

2. HSP according to claim 1, characterized by being in N-glycosylated form.

3. HSP according to claim 1 or 2, which is glycosylated at Asn15.

4. HSP according to claim 3, in which the glycosylated side chain contains at least one hexose unit.

5. HSP according to claim 4, in which the glycosylated side chain contains at least one mannose unit, preferably at least five mannose units, most preferably at least ten mannose units.

6. HSP according to claim 5, in which the glycosylated side chain contains 13-17 mannose units.

7. HSP according to claim 4, which in addition is glycosylated with at least one unit of N-acetyl glucosamine (GlcNAc).

8. HSP according to claim 7, which is glycosylated with  $(\text{GlcNAc})_2(\text{Man})_{10-15}$ .

9. HSP according to claim 1, wherein Lys2 is replaced by Asn, Gln7 is replaced by Asn, Arg10 is replaced by Asn, Gly 20 is replaced by Thr or Ser, Gly23 is replaced by Asn, Ile 24 is replaced by Asn, Phe 36 is replaced by Asn, Asp 37 is replaced by Asn, Ser39 is replaced by Asn, Gln53 is replaced by Asn, Glu61 is replaced by Asn, Asp64 is replaced by Asn, Arg66 is replaced by Thr or Ser, Gly69 is replaced by Thr or Ser, Gly72 is replaced by Asn, Ile 89 is replaced by Thr or Ser, Pro98 is replaced by Asn or Val101 is replaced by Thr or Ser, or a combination of two or more of these substitutions.

10. HSP according to claim 9, wherein Asp64 is replaced by Asn, and wherein Arg66 is replaced by Thr or Ser.

11. HSP according to claim 9, in which the glycosylated Asn contains at least one hexose unit.

12. HSP according to claim 11, which is glycosylated with at least one mannose unit, preferably at least five mannose units, most preferably at least ten mannose units.

13. HSP according to claim 12, which is glycosylated with 13-17 mannose units.

14. HSP according to claim 11, which in addition is glycosylated with at least one unit of N-acetyl glucosamine (GlcNAc).

15. HSP according to claim 14, which is glycosylated with  $(\text{GlcNAc})_2(\text{Man})_{10-15}$ .

16. A variant of a spasmolytic polypeptide which is a fragment of human spasmolytic polypeptide (HSP) or porcine spasmolytic polypeptide (PSP) comprising at least one trefoil domain.

17. A variant according to claim 16, which comprises at least a sequence of amino acids from position 8 to 46 or from position 58 to 95.

18. A variant according to claim 16 or 17, wherein Asn15 is substituted by another amino acid, e.g. Asp or Glu.

19. A variant according to claim 16 or 17, wherein Thr17 is substituted by another amino acid except Ser, e.g. Ala.

20. A variant according to claim 16 or 17, wherein Arg10 is replaced by Asn, Gly 20 is replaced by Thr or Ser, Gly23 is replaced by Asn, Ile 24 is replaced by Asn, Phe 36 is replaced by Asn, Asp 37 is replaced by Asn, Ser39 is replaced by Asn, Glu61 is replaced by Asn, Asp64 is replaced by Asn, Arg66 is replaced by Thr or Ser, Gly69 is replaced by Thr or Ser, or Gly72 is replaced by Asn, or a combination of two or more of these substitutions.

21. A method of preparing a spasmolytic polypeptide in at least 60% glycosylated form, wherein a host cell transformed with a DNA fragment encoding a spasmolytic polypeptide and capable of providing glycosylation of said spasmolytic polypeptide is cultured under conditions permitting production of said spasmolytic polypeptide and recovering the resulting spasmolytic polypeptide from the culture.

22. A method according to claim 21, wherein the host cell is a fungal cell such as a yeast cell, e.g. a strain of *Saccharomyces cerevisiae*, or a filamentous fungus cell, e.g. a strain of *Aspergillus*, such as *Aspergillus oryzae*.

23. A pharmaceutical composition comprising HSP according to any of claims 1-15 together with a pharmaceutically acceptable carrier or excipient.

24. A pharmaceutical composition comprising a variant spasmolytic polypeptide according to any of claims 16-20 together with a pharmaceutically acceptable carrier or excipient.

25. Use of HSP according to any of claims 1-15 for the manufacture of a medicament for the prophylaxis or treatment of gastrointestinal disorders.

26. Use of a variant spasmolytic polypeptide according to any of claims 16-20 for the manufacture of a medicament for the prophylaxis or treatment of gastrointestinal disorders.

\* \* \* \* \*