

(19) **DANMARK**

(10) **DK/EP 2921485 T3**



Patent- og
Varemærkestyrelsen

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **C 07 D 261/08 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-11-12**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-07-18**
- (86) Europæisk ansøgning nr.: **15154130.7**
- (86) Europæisk indleveringsdag: **2012-12-21**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-09-23**
- (30) Prioritet: **2011-12-27 EP 11195764**
- (62) Stamansøgningsnr: **12806489.6**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Bayer CropScience AG, Alfred-Nobel-Straße 50, 40789 Monheim, Tyskland**
- (72) Opfinder: **TSUCHIYA, Tomoki, 16 rue Jean Marie Leclair, 69009 Lyon, Frankrig
HOFFMANN, Sebastian, Zypressenweg 12, 41470 Neuss, Tyskland
SEITZ, Thomas, Rietherbach 10b, 40764 Langenfeld, Tyskland
HILLEBRAND, Stefan, Lothringer Str. 22, 41462 Neuss, Tyskland
Schmidt, Jan Peter, 130 Prospector Court, 95630 Folsom, USA
Cristau Pierre, 4 Rue Marietton, 69009 Lyon, Frankrig
Wasnaire, Pierre, Christophstraße 55, 49225 Düsseldorf, Tyskland
BENTING, Jürgen, Amselstraße 7, 42799 Leichlingen, Tyskland**
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **ISOXAZOLDERIVATER**
- (56) Fremdragne publikationer:
US-A- 5 814 627
JONATHAN D. TOKER ET AL: "Exploring the Scope of the 29G12 Antibody Catalyzed 1,3-Dipolar Cycloaddition Reaction", THE JOURNAL OF ORGANIC CHEMISTRY, Bd. 70, Nr. 20, 1. September 2005 (2005-09-01), Seiten 7810-7815, XP055195563, ISSN: 0022-3263, DOI: 10.1021/jo050410b
ALAM ATHAR ET AL: "Synthesis and bio-evaluation of human macrophage migration inhibitory factor inhibitor to develop anti-inflammatory agent", BIOORGANIC & MEDICINAL CHEMISTRY, PERGAMON, GB, Bd. 19, Nr. 24, 24. Oktober 2011 (2011-10-24), Seiten 7365-7373, XP028597161, ISSN: 0968-0896, DOI: 10.1016/J.BMC.2011.10.056
V. SINGH ET AL: "Interesting Results of Catalytic Hydrogenation of 3-(2-Nitrophenyl)isoxazoles and 3-(Nitrophenyl)-4,5-dihydroisoxazoles", SYNTHESIS, Bd. 2006, Nr. 12, 1. Juni 2006 (2006-06-01), Seiten 1995-2004, XP055195562, ISSN: 0039-7881, DOI: 10.1055/s-2006-942388
ZAGOZDA ET AL: "Optically active nitrile oxides: synthesis and 1,3-dipolar cycloaddition reactions", TETRAHEDRON ASYMMETRY, PERGAMON PRESS LTD, OXFORD, GB, Bd. 18, Nr. 12, 4. Juli 2007 (2007-07-04),

Fortsættes ...

Seiten 1457-1464, XP022157444, ISSN: 0957-4166, DOI: 10.1016/J.TETASY.2007.05.021

MOLINA C T ET AL: "SYNTHESIS OF NEW 4,5-DIHYDROISOXAZOLES WITH POTENTIAL ANTI-INFLAMMATORY ACTIVITY", HETEROCYCLIC COMMUNICATIONS, FREUND PUBLISHING HOUSE, TEL AVIV, IL, Bd. 9, Nr. 5, 1. Januar 2003 (2003-01-01), Seiten 535-538, XP009044208, ISSN: 0793-0283

Description

The invention relates to compounds suitable for preparation of heteroarylpiperidine derivatives, and to compositions for
5 controlling phytopathogenic fungi in and/or on plants or in and/or on seed of plants containing said heteroarylpiperidine derivatives.

The synthesis of aryl-substituted isoxazolines by 1,3-dipolar
10 cycloaddition is already known (J. D. Toker et al., J. Org. Chem. 70 (2005), 7810-7815; A. Alam et al., Bioorg. Med. Chem. 19 (2011), 7365-7373).

Furthermore, it is already known that particular
15 heterocyclically substituted thiazoles can be used as fungicidal crop protection compositions (see WO 07/014290, WO 08/013925, WO 08/013622, WO 08/091594, WO 08/091580, WO 09/055514, WO 09/094407, WO 09/094445, WO 09/132785, WO 10/037479, WO 10/065579, WO 11/076510, WO 11/018415, WO
20 11/018401, WO 11/076699, WO 11/146182, WO 12/055837, WO 12/025557, WO 12/082580). However, specifically at relatively low application rates, the fungicidal efficacy of these compounds is not always adequate.


25 Since the ecological and economical demands made on modern crop protection agents are increasing constantly, for example with respect to activity spectrum, toxicity, selectivity, application rate, formation of residues and favourable manufacture, and there can furthermore be problems, for
30 example, with resistances, there is a constant need to develop novel crop protection compositions, in particular fungicides, which, at least in some areas, have advantages over the known ones.

35 It has now been found that, surprisingly, the present heteroarylpiperidine and -piperazine derivatives achieve at least some aspects of the objects mentioned and are suitable for use as crop protection compositions, especially

fungicides.

The invention provides compounds of the formulae **(VIIIa)** and **(VIIIb)**

5



in which the radical definitions have the following meanings:

10 R⁵ represents hydrogen,

L² represents a direct bond,

R¹ represents phenyl which may contain 1, 2 or 3
15 substituents, where the substituents independently of one another are selected at least once from Z⁴ and optionally from the list below:

fluorine, chlorine, bromine, iodine, cyano, nitro, hydroxy,
20 amino, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl, 1,2-dimethylethyl, ethenyl, ethynyl, trifluoromethyl, difluoromethyl, trichloromethyl, dichloromethyl, cyclopropyl, methoxy, ethoxy, n-propoxy, 1-methylethoxy, 1,1-dimethylethoxy, methylcarbonyl, ethylcarbonyl,
25 methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, 1-methyl-ethoxycarbonyl, 1,1-dimethylethoxycarbonyl, 1-methylcarbonyloxy, methylthio, ethylthio, methylsulphonyl or -L³R³,

30 L³ represents a direct bond,

R³ represents hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₈-cycloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-

alkoxycarbonyl or C₁-C₄-haloalkoxycarbonyl,

Z³ represents a phenyl radical which may contain up to two substituents, where the substituents independently of one another are selected from the list below:

chlorine, bromine, iodine, fluorine, cyano, nitro, hydroxy, amino, -SH, methyl, ethyl, n-propyl, 1-methylethyl, 1,1-dimethylethyl, ethenyl, propen-2-yl, ethynyl, propyn-2-yl,
 10 trifluoromethyl, difluoromethyl, methoxymethyl, methylcarbonyl, ethylcarbonyl, trifluoromethylcarbonyl, methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, 1-methylethoxycarbonyl, 1,1-dimethylethoxycarbonyl, methoxy, ethoxy, n-propoxy, 1-methylethoxy, 1,1-dimethylethoxy,
 15 trifluoromethoxy, ethenyloxy, 2-propenyloxy, ethynyloxy, 2-propynyloxy, methylthio, ethylthio, trifluoromethylthio, methylsulphonyl, ethylsulphonyl, propylthionyl, 1-methylethylthio, trifluoromethylsulphonyl, methylamino, ethylamino, n-propylamino, 1-methylethylamino, 1,1-dimethylethylamino or dimethylamino or

Z³ represents naphthalenyl,

R¹³ and R¹⁴ are the same or different and independently represent hydrogen, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl or 1,1-dimethylethyl,

Z⁴ represents -formyl, methoxymethoxy, 2-methoxyethoxy, allyloxy, 2-fluoroprop-2-en-1-yloxy, 2-chloroprop-2-en-1-yloxy, 3-chloroprop-2-en-1-yloxy, 2-bromoprop-2-en-1-yloxy, 2-methylprop-2-en-1-yloxy, 3,3-dichloroprop-2-en-1-yloxy, 3,3-dichloro-2-fluoroprop-2-en-1-yloxy, but-2-en-1-yloxy, but-3-en-2-yloxy, but-3-en-1-yloxy, 3-chlorobut-2-en-1-yloxy, 3-methylbut-2-en-1-yloxy, 4,4,4-trifluorobut-2-en-1-yloxy, prop-2-yn-1-yloxy, 3-chloroprop-2-yn-1-yloxy, 3-bromoprop-2-yn-1-yloxy, but-2-yn-1-yloxy, pent-2-yn-1-yloxy, 2-fluoro-2-methylpropanoyloxy, 3,3,3-trifluoropropanoyloxy, cyclopropylcarbonyloxy, cyclohexylcarbonyloxy, (1-

chlorocyclopropyl)carbonyloxy, but-2-enoyloxy, acryloyloxy,
 cyanomethoxy, methylsulphonyloxy, ethylsulphonyloxy,
 trifluoromethylsulphonyloxy, cyclopropylsulphonyloxy, 2-
 methoxyethoxymethyl, allyloxymethyl, prop-2-yn-1-yloxymethyl,
 5 methylsulphonylmethyl, methylcarbonylaminoethyl,
 methylsulphonylaminoethyl, $-C(=NOR^7)R^8$,
 dimethylaminosulphonyl, ethylaminosulphonyl,
 trimethylsilylethynyl, diethylaminosulphonyl,
 methylaminosulphonyl, trimethylsilyloxy, trimethylsilylprop-2-
 10 yn-1-yloxy, trifluoromethylamino, dimethylaminocarbonylamino,
 $-C(=O)OH$, $-NHC(=O)H$, $-C(=O)NH_2$,
 $-C(=S)NR^{13}R^{14}$, 1,1-dimethylethylcarbonylamino,
 chloromethylcarbonylamino, trifluoromethylcarbonylamino, 1,1-
 dimethylethoxycarbonylamino, ethylcarbonylamino, 1-
 15 methylethoxycarbonylamino, trifluoromethylcarbonylamino,
 methylcarbonylamino, methoxycarbonylamino,
 ethoxycarbonylamino, isopropoxycarbonylamino,
 1-methylethylcarbonylamino, methylsulphonylamino or
 phenylsulphonylamino, 3-bromoprop-2-en-1-yloxy or $-L^4Z^3$,


20 L^4 represents $-C(=O)O-$ or $-OCH_2C\equiv C-$,

R^7 represents hydrogen, methyl, ethyl, n-propyl,
 1-methylethyl, n-butyl, 1,1-dimethylethyl or 2-methylpropyl,

25 R^8 represents hydrogen, methyl, ethyl, n-propyl,
 1-methylethyl, n-butyl, 1,1-dimethylethyl or 2-methylpropyl,

and salts thereof.

30 The compounds of the formulae **(VIIIa)** and **(VIIIb)** are suitable
 for preparation of the compounds of the formula **(I)**



(I)

in which L^2 and R^1 are each defined as in the formulae **(VIIIa)** and **(VIIIb)**, and where the remaining radical definitions have the following meanings:

5

R^{A1} is hydrogen, halogen, cyano, amino, -CHO, -C(=O)OH, -C(=O)NH₂, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, cycloalkyl, halocycloalkyl, alkylcycloalkyl, cycloalkylalkyl, halocycloalkylalkyl, cycloalkenyl, halocycloalkenyl, alkoxyalkyl, alkylthioalkyl, alkylsulphinylalkyl, alkylsulphonylalkyl, alkylaminoalkyl, dialkylaminoalkyl, haloalkylaminoalkyl, alkylcarbonyl, haloalkylcarbonyl, cycloalkylcarbonyl, alkoxycarbonyl, cycloalkoxycarbonyl, cycloalkylalkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkoxy, haloalkoxy, cycloalkoxy, halocycloalkoxy, alkenyloxy, haloalkenyloxy, alkynyloxy, haloalkynyloxy, alkoxyalkoxy, alkylcarbonyloxy, haloalkylcarbonyloxy, alkylthio, haloalkylthio, cycloalkylthio, alkylamino, dialkylamino, haloalkylamino, halodialkylamino, cycloalkylamino, alkylcarbonylamino, haloalkylcarbonylamino, alkylsulphonylamino or haloalkylsulphonylamino,

R^{A2} is hydrogen, halogen, cyano, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, or

R^{A2} is an unsubstituted or substituted phenyl, an optionally benzofused, substituted 5- or 6-membered heterocyclyl, where the substituents are each independently selected from the following list: hydroxyl, cyano, halogen, alkyl, haloalkyl, alkoxy, haloalkoxy, or

R^{A1} and R^{A2} , together with the carbon atom to which they are bonded, form a three- to seven-membered saturated or partially unsaturated ring optionally containing one, two, three or four heteroatoms from the group of oxygen, nitrogen, silicon or sulphur, where optionally one, two or three carbon ring members are selected from C(=O) and C(=S) and the sulphur ring

35

members are selected from $S(=O)_s(=NR^{A3})_f$, and the silicon ring members are selected from $SiR^{A4}R^{A5}$, where the ring may be unsubstituted or substituted, where the substituents are each independently selected from R^{A6} ,

5

R^{A3} is hydrogen, cyano, alkyl, haloalkyl, cycloalkyl, halocycloalkyl, alkoxy, haloalkoxy, alkylamino, dialkylamino, haloalkylamino or phenyl,

10 R^{A4} and R^{A5} are the same or different and are each independently alkyl, alkenyl, alkynyl, cycloalkyl, halocycloalkyl, cycloalkylalkyl, alkylcycloalkyl, alkylcycloalkylalkyl, haloalkyl, alkoxy or haloalkoxy,

15 s is 0, 1 or 2, and

f is 0, 1 or 2, and

$s+f$ is 0, 1 or 2,

20

R^{A6} is halogen, cyano, alkyl, haloalkyl, alkoxy or haloalkoxy on the carbon ring members and cyano, alkyl or alkoxy on the nitrogen ring members,

25 L^1 is oxygen, sulphur, $-N(R^{L1})-$, $-C(R^{L2})_2-$, $-OC(R^{L2})_2-$, $-SC(R^{L2})_2-$, $-N(R^{L1})-C(R^{L2})_2-$,

30 where the bond pointing to the left is bonded to the nitrogen atom of formula I and the bond pointing to the right is bonded to the nitrogen atom of formula I,

R^{L1} is hydrogen, cyano, alkyl, haloalkyl, alkoxyalkyl, alkylthioalkyl, alkylcarbonyl, haloalkylcarbonyl, alkoxyalkyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylsulphonyl or haloalkylsulphonyl, or the two R^{L1} and R^{A2} radicals together with the carbon atom to which they are bonded form a five- to seven-membered partially unsaturated ring optionally containing one, two or three heteroatoms from

the group of oxygen, nitrogen and sulphur, where the ring may be unsubstituted or substituted, where the substituents are each independently selected from R^{A6},

5 R^{L2} is hydrogen, alkyl or haloalkyl,

R^{B1} is hydrogen, halogen, cyano, hydroxy, formyl, alkyl, alkenyl, alkynyl, haloalkyl, haloalkenyl, haloalkynyl, alkoxyalkyl, alkylthioalkyl, alkylsulphinylalkyl,
10 alkylsulphonylalkyl, alkylcarbonyl, haloalkylcarbonyl, alkoxy, haloalkoxy, alkylthio, haloalkylthio, alkylsulphinyl, haloalkylsulphinyl, alkylsulphonyl, haloalkylsulphonyl, alkylcarbonyloxy, haloalkylcarbonyloxy, alkoxy, haloalkoxy, alkylaminocarbonyloxy, dialkylaminocarbonyloxy, or
15

R^{B1} is a phenyl radical, naphthalenyl radical or a 5- or 6-membered heteroaryl radical, each of which may contain 0, 1, 2 or 3 substituents, where the substituents are each
20 independently selected from the following list: hydrogen, halogen, alkyl, haloalkyl, cycloalkyl, alkoxy, haloalkoxy, alkylcarbonyloxy, haloalkylcarbonyloxy,

R^{B2} is hydrogen, alkyl or haloalkyl,
25 or the two R^{B1} und R^{B2} radicals together with the carbon atom to which they are bonded form a three- to six-membered saturated ring,

Y is sulphur or oxygen,
30


X represents carbon or nitrogen,

R² represents hydrogen, alkyl, alkenyl, haloalkyl, alkoxy, halogen, cyano or hydroxy,
35

R¹⁰ is the same or different and independently represents hydrogen, alkyl, alkenyl, haloalkyl, alkoxy, halogen, cyano or hydroxy,

p represents 0, 1 or 2,

G represents 5-membered heteroaryl which is substituted by
5 Q and may otherwise be unsubstituted or substituted,



where the bond identified by "*" is bonded directly to G or L²,
and where the bond identified by "#" is bonded directly
10 to L² or G or where the bond identified by "*" is bonded
directly to L², and at the same time the bond identified by "#" is bonded directly to G,

R⁵ represents hydrogen,

15

and salts of the compounds of the formula **(I)**.

Heteroarylpiperidine and -piperazine derivatives of the
formula **(I)** and the salts, metal complexes and N-oxides
20 thereof are very suitable for controlling phytopathogenic
harmful fungi. The aforementioned compounds exhibit, in
particular, potent fungicidal activity and can be used in crop
protection, in the domestic and hygiene sector and in the
protection of materials.

25

The compounds of the formula **(I)** may be present either in pure
form or as mixtures of different possible isomeric forms,
especially of stereoisomers, such as E and Z, threo and
erythro, and also optical isomers, such as R and S isomers or
30 atropisomers, and, if appropriate, also of tautomers. Both the
E and Z isomers, and the threo and erythro isomers, and also
the optical isomers, any desired mixtures of these isomers,
and the possible tautomeric forms are claimed.

35 The radical definitions of the compounds of the formula **(I)**
preferably, more preferably and most preferably have the

following definitions:

5 R^{A1} is preferably hydrogen, cyano, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, C₁-C₄-haloalkyl, C₂-C₄-haloalkenyl, C₂-C₄-haloalkynyl, C₂-C₄-alkoxyalkyl, C₂-C₄-alkylthioalkyl, C₁-C₃-alkylcarbonyl, C₁-C₃-haloalkylcarbonyl, C₁-C₃-alkoxycarbonyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₂-C₄-alkenyloxy, C₂-C₄-haloalkenyloxy, C₂-C₄-alkynyloxy, C₂-C₄-haloalkynyloxy, C₂-C₄-alkoxyalkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkylthio, C₁-C₄-alkylamino, C₂-C₄-dialkylamino, C₁-C₄-haloalkylamino, C₂-C₄-halodialkylamino, C₃-C₆-cycloalkyl and more preferably hydrogen, methyl, ethyl, propan-2-yl, *t*-butyl, difluoromethyl, trifluoromethyl, methoxymethyl, ethoxymethyl, methoxycarbonyl, ethoxycarbonyl or cyclopropyl,

15

R^{A2} is preferably hydrogen, C₁-C₃-alkyl, C₁-C₃-haloalkyl, C₁-C₃-alkoxy, or

20 R^{A2} is preferably an unsubstituted or substituted phenyl, an optionally benzofused, substituted 5- or 6-membered heterocyclyl, where the substituents are each independently selected from the following list: hydroxyl, cyano, halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy,

25 R^{A2} is more preferably methyl, ethyl, propyl, isopropyl, butyl, *tert*-butyl, *iso*-butyl, 1,3-benzodioxolyl or an unsubstituted or substituted phenyl, where the substituents are each independently selected from the following list: hydroxyl, cyano, fluorine, chlorine, bromine, methyl, difluoromethyl, trifluoromethyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy,

35 R^{A1} and R^{A2} preferably form, together with the carbon atom to which they are bonded, a three- to six-membered saturated or partially unsaturated ring optionally containing one, two, three or four heteroatoms from the group of oxygen, nitrogen and sulphur, where one carbon ring member is optionally selected from C(=O) and C(=S), where the ring may contain no,

one, two or three substituents, where the substituents are each independently selected from R^{A6} ,

5 R^{A6} is preferably halogen, cyano, C_1 - C_2 -alkyl, C_1 - C_2 -haloalkyl, C_1 - C_2 -alkoxy or C_1 - C_2 -haloalkoxy on the carbon ring members, and cyano, C_1 - C_2 -alkyl or C_1 - C_2 -alkoxy on the nitrogen ring members,

L^1 is preferably oxygen, sulphur, $-N(R^{L1})-$,

10 R^{L1} is preferably hydrogen, C_1 - C_2 -alkyl, C_1 - C_2 -haloalkyl, $-C(=O)CH_3$, $-C(=O)CF_3$, $C(=O)OCH_3$, or the two R^{L1} and R^{A2} radicals together with the carbon atom to which they are bonded form a five- to seven-membered partially unsaturated ring optionally containing one, two or three heteroatoms from the group of
15 oxygen, nitrogen and sulphur, where the ring may be unsubstituted or substituted, where the substituents are each independently selected from R^{A6} ,

R^{B1} is preferably hydrogen, cyano, hydroxyl, C_1 - C_3 -alkyl, C_2 -
20 C_3 -alkenyl, C_2 - C_3 -alkynyl, C_1 - C_3 -haloalkyl, C_2 - C_3 -haloalkenyl, C_2 - C_3 -haloalkynyl, C_2 - C_3 -alkylcarbonyl, C_2 - C_3 -haloalkylcarbonyl, C_1 - C_3 -alkoxy, C_1 - C_3 -haloalkoxy, C_1 - C_3 -alkylthio, C_1 - C_3 -haloalkylthio, C_1 - C_2 -alkylcarbonyloxy, C_1 - C_2 -haloalkylcarbonyloxy, or

25 R^{B1} is preferably a phenyl radical, naphthalenyl radical or a 5- or 6-membered heteroaryl radical, each of which may contain 0, 1, 2 or 3 substituents, where the substituents are each independently selected from the following list:

30 hydrogen, fluorine, chlorine, bromine, C_1 - C_3 -alkyl, C_1 - C_3 -haloalkyl, C_1 - C_3 -alkoxy, C_1 - C_3 -haloalkoxy,

R^{B2} is preferably hydrogen or C_1 - C_2 -alkyl, and more preferably hydrogen,

35 Y is preferably sulphur or oxygen, and more preferably oxygen,

X is preferably carbon or nitrogen, and more preferably carbon,


R² is preferably hydrogen, C₁-C₄-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, halogen, cyano or hydroxy, and more preferably hydrogen, fluorine, methoxy or hydroxy, and most preferably hydrogen,

R¹⁰ is preferably the same or different and is independently hydrogen, C₁-C₄-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy, halogen, cyano or hydroxy, and more preferably hydrogen, fluorine, methoxy or hydroxy,

p is preferably 0 to 1, and more preferably 0,

15

G is preferably




20 where the bond identified by "v" is bonded directly to X and where the bond identified by "w" is bonded directly to Q,

G is more preferably G¹, G² or G³, and most preferably G¹,

25 R^{G1} is preferably hydrogen, C₁-C₃-alkyl or halogen and more preferably hydrogen,

Q is more preferably



where the bond identified by "x" is bonded directly to G, and
 5 where the bond identified by "y" is bonded directly to L²,

R⁵ is hydrogen,

R³ is hydrogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₈-
 10 cycloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-haloalkylcarbonyl, C₁-C₄-
 alkoxy carbonyl or C₁-C₄-haloalkoxy carbonyl,

L² is a direct bond,

15 R¹ is phenyl containing 1, 2 or 3 substituents, where the
 substituents are independently selected at least once from Z⁴
 and optionally from the following list: fluorine, chlorine,
 bromine, iodine, cyano, nitro, hydroxy, amino, methyl, ethyl,
 n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl, 1,2-
 20 dimethylethyl, ethenyl, ethynyl, trifluoromethyl,
 difluoromethyl, trichloromethyl, dichloromethyl, cyclopropyl,
 methoxy, ethoxy, n-propoxy, 1-methylethoxy, 1,1-
 dimethylethoxy, methylcarbonyl, ethylcarbonyl,
 methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, 1-
 25 methylethoxycarbonyl, 1,1-dimethylethoxycarbonyl, 1-
 methylcarbonyloxy, methylthio, ethylthio, methylsulphonyl or -
 L³R³, and most preferably phenyl containing 1, 2 or 3
 substituents, where the substituents are independently
 selected from the following list:

30

formyl, methoxymethoxy, 2-methoxyethoxy, allyloxy, 2-
 fluoroprop-2-en-1-yloxy, 2-chloroprop-2-en-1-yloxy, 3-
 chloroprop-2-en-1-yloxy, 2-bromoprop-2-en-1-yloxy, 2-

methylprop-2-en-1-yloxy, 3,3-dichloroprop-2-en-1-yloxy, 3,3-
 dichloro-2-fluoroprop-2-en-1-yloxy, but-2-en-1-yloxy, but-3-
 en-2-yloxy, but-3-en-1-yloxy, 3-chlorobut-2-en-1-yloxy, 3-
 methylbut-2-en-1-yloxy, 4,4,4-trifluorobut-2-en-1-yloxy, prop-
 5 2-yn-1-yloxy, 3-chloroprop-2-yn-1-yloxy, 3-bromoprop-2-yn-1-
 yloxy, but-2-yn-1-yloxy, pent-2-yn-1-yloxy, 2-fluoro-2-
 methylpropanoyloxy, 3,3,3-trifluoropropanoyloxy,
 cyclopropylcarbonyloxy, cyclohexylcarbonyloxy, (1-
 chlorocyclopropyl)carbonyloxy, but-2-enoyloxy, acryloyloxy,
 10 benzoyloxy, 2-fluorobenzoyloxy, 3-fluorobenzoyloxy, 4-
 fluorobenzoyloxy, cyanomethoxy, methylsulphonyloxy,
 ethylsulphonyloxy, trifluoromethylsulphonyloxy,
 cyclopropylsulphonyloxy, 2-methoxyethoxymethyl,
 allyloxymethyl, prop-2-yn-1-yloxymethyl,
 15 methylsulphonylmethyl, methylcarbonylaminomethyl,
 methylsulphonylaminomethyl, $-C(=NOH)H$, $-C(=NOCH_3)H$, $-$
 $C(=NOCH_2CH_3)H$, $-C(=NOCH(CH_3)CH_3)H$, $-C(=NOH)CH_3$, $-C(=NOCH_3)CH_3$, $-$
 $C(=NOCH_2CH_3)CH_3$, $-C(=NOCH(CH_3)CH_3)CH_3$, dimethylaminosulphonyl,
 $C(=O)NH_2$, ethylaminosulphonyl, trimethylsilylethynyl,
 20 diethylaminosulphonyl, methylaminosulphonyl,
 trimethylsilyloxy, trimethylsilylprop-2-yn-1-yloxy,
 trifluoromethylamino, dimethylaminocarbonylamino, $-C(=O)OH$,
 1,1-dimethylethylcarbonylamino, chloromethylcarbonylamino,
 trifluoromethylcarbonylamino, 1,1-dimethylethoxycarbonylamino,
 25 ethylcarbonylamino, 1-methylethoxycarbonylamino,
 trifluoromethylcarbonylamino, methylcarbonylamino,
 methoxycarbonylamino, ethoxycarbonylamino, iso-
 propoxycarbonylamino, 1-methylethylcarbonylamino,
 methylsulphonylamino or phenylsulphonylamino, 3-bromoprop-2-
 30 en-1-yloxy,

and additional substituents are optionally selected from the
 following list: fluorine, chlorine, methyl, trifluoromethyl,
 methoxy,

35

L^3 is a direct bond,

Z^3 is a phenyl radical, which may contain up to two

substituents, where the substituents are each independently selected from the following list:

chlorine, bromine, iodine, fluorine, cyano, nitro, hydroxyl,
 5 amino, -SH, methyl, ethyl, *n*-propyl, 1-methylethyl, 1,1-
 dimethylethyl, ethenyl, propen-2-yl, ethynyl, propyn-2-yl,
 trifluoromethyl, difluoromethyl, methoxymethyl,
 methylcarbonyl, ethylcarbonyl, trifluoromethylcarbonyl,
 methoxycarbonyl, ethoxycarbonyl, *n*-propoxycarbonyl, 1-
 10 methylethoxycarbonyl, 1,1-dimethylethoxycarbonyl, methoxy,
 ethoxy, *n*-propoxy, 1-methylethoxy, 1,1-dimethylethoxy,
 trifluoromethoxy, ethenyloxy, 2-propenyloxy, ethynyloxy, 2-
 propynyloxy, methylthio, ethylthio, trifluoromethylthio,
 methylsulphonyl, ethylsulphonyl, propylthionyl, 1-
 15 methylethylthio, trifluoromethylsulphonyl, methylamino,
 ethylamino, *n*-propylamino, 1-methylethylamino, 1,1-
 dimethylethylamino or dimethylamino, or

Z³ is naphthalenyl,
 20

R¹³ and R¹⁴ are the same or different and are each independently hydrogen, methyl, ethyl, *n*-propyl, 1-methylethyl, *n*-butyl or 1,1-dimethylethyl,

Z⁴ is - formyl, methoxymethoxy, 2-methoxyethoxy, allyloxy,
 25 2-fluoroprop-2-en-1-yloxy, 2-chloroprop-2-en-1-yloxy, 3-
 chloroprop-2-en-1-yloxy, 2-bromoprop-2-en-1-yloxy, 2-
 methylprop-2-en-1-yloxy, 3,3-dichloroprop-2-en-1-yloxy, 3,3-
 dichloro-2-fluoroprop-2-en-1-yloxy, but-2-en-1-yloxy, but-3-
 30 en-2-yloxy, but-3-en-1-yloxy, 3-chlorobut-2-en-1-yloxy 3-
 methylbut-2-en-1-yloxy, 4,4,4-trifluorobut-2-en-1-yloxy, prop-
 2-yn-1-yloxy, 3-chloroprop-2-yn-1-yloxy, 3-bromoprop-2-yn-1-
 yloxy, but-2-yn-1-yloxy, pent-2-yn-1-yloxy, 2-fluoro-2-
 methylpropanoyloxy, 3,3,3-trifluoropropanoyloxy,
 35 cyclopropylcarbonyloxy, cyclohexylcarbonyloxy, (1-
 chlorocyclopropyl)carbonyloxy, but-2-enoyloxy, acryloyloxy,
 cyanomethoxy, methylsulphonyloxy, ethylsulphonyloxy,
 trifluoromethylsulphonyloxy, cyclopropylsulphonyloxy, 2-

methoxyethoxymethyl, allyloxymethyl, prop-2-yn-1-yloxymethyl,
 methylsulphonylmethyl, methylcarbonylaminoethyl,
 methylsulphonylaminoethyl, $-C(=NOR^7)R^8$,
 dimethylaminosulphonyl, ethylaminosulphonyl,
 5 trimethylsilylethynyl, diethylaminosulphonyl,
 methylaminosulphonyl, trimethylsilyloxy, trimethylsilylprop-2-
 yn-1-yloxy, trifluoromethylamino, dimethylaminocarbonylamino,
 $-C(=O)OH$, $-NHC(=O)H$, $-C(=O)NH_2$, $-C(=S)NR^{13}R^{14}$ 1,1-
 dimethylethylcarbonylamino, chloromethylcarbonylamino,
 10 trifluoromethylcarbonylamino, 1,1-dimethylethoxycarbonylamino,
 ethylcarbonylamino, 1-methylethoxycarbonylamino,
 trifluoromethylcarbonylamino, methylcarbonylamino,
 methoxycarbonylamino, ethoxycarbonylamino, iso-
 propoxycarbonylamino, 1-methylethylcarbonylamino,
 15 methylsulphonylamino or phenylsulphonylamino, 3-bromoprop-2-
 en-1-yloxy, or $-L^4Z^3$,

L^4 is $-OCH_2C\equiv C-$ or $-C(=O)O-$,

20 R^7 is hydrogen, methyl, ethyl, *n*-propyl, 1-methylethyl, *n*-
 butyl, 1,1-dimethylethyl or 2-methylpropyl,

R^8 is hydrogen, methyl, ethyl, *n*-propyl, 1-methylethyl, *n*-
 butyl, 1,1-dimethylethyl or 2-methylpropyl.

25

The radical definitions of the radical definitions above and
 specified below of the formula **(I)** apply to the end products
 of the formula **(I)**, and also equally to all intermediates (see
 also below under "Elucidations of the processes and
 30 intermediates").

The radical definitions and elucidations listed above and
 below, in general terms or in areas of preference, can be
 combined with one another as desired, i.e. including
 35 combinations between the particular areas and areas of
 preference. They apply both to the end products and
 correspondingly to precursors and intermediates. Moreover,
 individual definitions may not apply.

Preference is given to those compounds of the formula **(I)** in which all radicals have the abovementioned preferred definitions.

5

Particular preference is given to those compounds of the formula **(I)** in which all radicals have the abovementioned more preferred definitions.

10 Very particular preference is given to those compounds of the formula **(I)** in which all radicals have the abovementioned most preferred definitions.

Preference is additionally given to compounds of the formula **(I)** and agrochemically active salts, metal complexes and N-oxides thereof, in which:

15

R^{A1} is methyl, trifluoromethyl or cyclopropyl,

20 R^{A2} is methyl or propan-2-yl,

R^{A2} is 1,3-benzodioxol-5-yl, 4-ethoxyphenyl, 3-fluorophenyl, 3,4-dimethylphenyl, 3-(trifluoromethoxy)phenyl, 3,4-dimethylphenyl or 4-ethoxyphenyl,

25

L^1 is oxygen,

R^{B1} and R^{B2} are each hydrogen,

30 Y is oxygen;

G is G^1 ;

R^{G1} is hydrogen;

35

Q is Q^{24-3} or Q is Q^{11-1} ;

R^5 is hydrogen or R^5 is methyl;

L² is a direct bond;

R¹ is 2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl or

5

R¹ is 2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl or

10 R¹ is 2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,3-difluoro-4-formylphenyl or

R¹ is 2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl or

15

R¹ is 2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

20 R¹ is 2,4-difluoro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,4-difluoro-3-formylphenyl or

R¹ is 2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl or

25

R¹ is 2,5-dichloro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl or

30 R¹ is 2,5-dichloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2,5-difluoro-3-[(methylsulphonyl)oxy]phenyl or

35

R¹ is 2,5-difluoro-3-formylphenyl or

R¹ is 2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2,5-difluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,5-difluoro-4-formylphenyl or

5

R¹ is 2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2,6-dichloro-3-[(methylsulphonyl)oxy]phenyl or

10 R¹ is 2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2,6-dichloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

15

R¹ is 2,6-difluoro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,6-difluoro-3-formylphenyl or

20 R¹ is 2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2,6-difluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2,6-difluoro-4-formylphenyl or

25

R¹ is 2-(allyloxy)-3,4-dichlorophenyl or

R¹ is 2-(allyloxy)-3,4-difluorophenyl or

30 R¹ is 2-(allyloxy)-3,5-dichlorophenyl or

R¹ is 2-(allyloxy)-3,5-difluorophenyl or

R¹ is 2-(allyloxy)-3,6-dichlorophenyl or

35

R¹ is 2-(allyloxy)-3,6-difluorophenyl or

R¹ is 2-(allyloxy)-3-chlorophenyl or

R¹ is 2-(allyloxy)-3-fluorophenyl or

R¹ is 2-(allyloxy)-3-methylphenyl or

5

R¹ is 2-(allyloxy)-4,5-dichlorophenyl or

R¹ is 2-(allyloxy)-4,5-difluorophenyl or

10 R¹ is 2-(allyloxy)-4,6-dichlorophenyl or

R¹ is 2-(allyloxy)-4,6-difluorophenyl or

R¹ is 2-(allyloxy)-4-chlorophenyl or

15

R¹ is 2-(allyloxy)-4-fluorophenyl or

R¹ is 2-(allyloxy)-4-methylphenyl or

20 R¹ is 2-(allyloxy)-5,6-dichlorophenyl or

R¹ is 2-(allyloxy)-5,6-difluorophenyl or

R¹ is 2-(allyloxy)-5-chlorophenyl or

25

R¹ is 2-(allyloxy)-5-fluorophenyl or

R¹ is 2-(allyloxy)-5-methylphenyl or

30 R¹ is 2-(allyloxy)-6-chlorophenyl or

R¹ is 2-(allyloxy)-6-fluorophenyl or

R¹ is 2-(allyloxy)-6-methylphenyl or

35

R¹ is 2-(allyloxy)phenyl or

R¹ is 2-(cyanomethoxy)-3,4-dichlorophenyl or

- R¹ is 2-(cyanomethoxy)-3,4-difluorophenyl or
- 5 R¹ is 2-(cyanomethoxy)-3,5-dichlorophenyl or
- R¹ is 2-(cyanomethoxy)-3,5-difluorophenyl or
- R¹ is 2-(cyanomethoxy)-3,6-dichlorophenyl or
- 10 R¹ is 2-(cyanomethoxy)-3,6-difluorophenyl or
- R¹ is 2-(cyanomethoxy)-3-chlorophenyl or
- R¹ is 2-(cyanomethoxy)-3-fluorophenyl or
- 15 R¹ is 2-(cyanomethoxy)-3-methylphenyl or
- R¹ is 2-(cyanomethoxy)-4,5-dichlorophenyl or
- 20 R¹ is 2-(cyanomethoxy)-4,5-difluorophenyl or
- R¹ is 2-(cyanomethoxy)-4,6-dichlorophenyl or
- R¹ is 2-(cyanomethoxy)-4,6-difluorophenyl or
- 25 R¹ is 2-(cyanomethoxy)-4-chlorophenyl or
- R¹ is 2-(cyanomethoxy)-4-fluorophenyl or
- 30 R¹ is 2-(cyanomethoxy)-4-methylphenyl or
- R¹ is 2-(cyanomethoxy)-5,6-dichlorophenyl or
- R¹ is 2-(cyanomethoxy)-5,6-difluorophenyl or
- 35 R¹ is 2-(cyanomethoxy)-5-chlorophenyl or
- R¹ is 2-(cyanomethoxy)-5-fluorophenyl or

R¹ is 2-(cyanomethoxy)-5-methylphenyl or

R¹ is 2-(cyanomethoxy)-6-chlorophenyl or

5

R¹ is 2-(cyanomethoxy)-6-fluorophenyl or

R¹ is 2-(cyanomethoxy)-6-methylphenyl or

10 R¹ is 2-(cyanomethoxy)phenyl or

R¹ is 2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl or

15

R¹ is 2-chloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2-chloro-3-[(methylsulphonyl)oxy]phenyl or

20 R¹ is 2-chloro-3-formylphenyl or

R¹ is 2-chloro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2-chloro-4-[(methylsulphonyl)oxy]phenyl or

25

R¹ is 2-chloro-4-formylphenyl or

R¹ is 2-fluoro-3-(prop-2-yn-1-yloxy)phenyl or

30 R¹ is 2-fluoro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 2-fluoro-3-formylphenyl or

R¹ is 2-fluoro-4-(prop-2-yn-1-yloxy)phenyl or

35

R¹ is 2-fluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 2-fluoro-4-formylphenyl or

R¹ is 2-formyl-3-methylphenyl or

R¹ is 2-formyl-4-methylphenyl or

5

R¹ is 2-formyl-5-methylphenyl or

R¹ is 2-formyl-6-methylphenyl or

10 R¹ is 2-formylphenyl or

R¹ is 2-methyl-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2-methyl-3-[(methylsulphonyl)oxy]phenyl or

15

R¹ is 2-methyl-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 2-methyl-4-[(methylsulphonyl)oxy]phenyl or

20 R¹ is 2-[(hydroxyimino)methyl]-3,4-difluorophenyl or

R¹ is 2-[(hydroxyimino)methyl]-3,5-difluorophenyl or

R¹ is 2-[(hydroxyimino)methyl]-3,6-difluorophenyl or

25

R¹ is 2-[(hydroxyimino)methyl]-3-chlorophenyl or

R¹ is 2-[(hydroxyimino)methyl]-3-fluorophenyl or

30 R¹ is 2-[(hydroxyimino)methyl]-3-methylphenyl or

R¹ is 2-[(hydroxyimino)methyl]-4,5-difluorophenyl or

R¹ is 2-[(hydroxyimino)methyl]-4,6-difluorophenyl or

35

R¹ is 2-[(hydroxyimino)methyl]-4-chlorophenyl or

R¹ is 2-[(hydroxyimino)methyl]-4-fluorophenyl or

- R¹ is 2-[(hydroxyimino)methyl]-4-methylphenyl or
- 5 R¹ is 2-[(hydroxyimino)methyl]-5,6-difluorophenyl or
- R¹ is 2-[(hydroxyimino)methyl]-5-chlorophenyl or
- R¹ is 2-[(hydroxyimino)methyl]-5-fluorophenyl or
- 10 R¹ is 2-[(hydroxyimino)methyl]-5-methylphenyl or
- R¹ is 2-[(hydroxyimino)methyl]-6-chlorophenyl or
- R¹ is 2-[(hydroxyimino)methyl]-6-fluorophenyl or
- 15 R¹ is 2-[(hydroxyimino)methyl]-6-methylphenyl or
- R¹ is 2-[(hydroxyimino)methyl]phenyl or
- 20 R¹ is 2-[(methoxyimino)methyl]-3,4-difluorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-3,5-difluorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-3,6-difluorophenyl or
- 25 R¹ is 2-[(methoxyimino)methyl]-3-chlorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-3-fluorophenyl or
- 30 R¹ is 2-[(methoxyimino)methyl]-3-methylphenyl or
- R¹ is 2-[(methoxyimino)methyl]-4,5-difluorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-4,6-difluorophenyl or
- 35 R¹ is 2-[(methoxyimino)methyl]-4-chlorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-4-fluorophenyl or

- R¹ is 2-[(methoxyimino)methyl]-4-methylphenyl or
- 5 R¹ is 2-[(methoxyimino)methyl]-5,6-difluorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-5-chlorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-5-fluorophenyl or
- 10 R¹ is 2-[(methoxyimino)methyl]-5-methylphenyl or
- R¹ is 2-[(methoxyimino)methyl]-6-chlorophenyl or
- R¹ is 2-[(methoxyimino)methyl]-6-fluorophenyl or
- 15 R¹ is 2-[(methoxyimino)methyl]-6-methylphenyl or
- R¹ is 2-[(methoxyimino)methyl]phenyl or
- 20 R¹ is 2-[(methylsulphonyl)oxy]phenyl or
- R¹ is 2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl or
- R¹ is 3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl or
- 25 R¹ is 3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl or
- R¹ is 3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl or
- 30 R¹ is 3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl or
- R¹ is 3,4-difluoro-2-formylphenyl or
- R¹ is 3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl or
- 35 R¹ is 3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl or
- R¹ is 3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl or

5

R¹ is 3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,5-difluoro-2-formylphenyl or

10 R¹ is 3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,5-difluoro-4-formylphenyl or

15

R¹ is 3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl or

20 R¹ is 3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl or

25

R¹ is 3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,6-difluoro-2-formylphenyl or

30 R¹ is 3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3,6-difluoro-4-formylphenyl or

35

R¹ is 3-(allyloxy)-2,4-dichlorophenyl or

R¹ is 3-(allyloxy)-2,4-difluorophenyl or

R¹ is 3-(allyloxy)-2,5-dichlorophenyl or

R¹ is 3-(allyloxy)-2,5-difluorophenyl or

5

R¹ is 3-(allyloxy)-2,6-dichlorophenyl or

R¹ is 3-(allyloxy)-2,6-difluorophenyl or

10 R¹ is 3-(allyloxy)-2-chlorophenyl or

R¹ is 3-(allyloxy)-2-fluorophenyl or

R¹ is 3-(allyloxy)-2-methylphenyl or

15

R¹ is 3-(allyloxy)-4,5-dichlorophenyl or

R¹ is 3-(allyloxy)-4,5-difluorophenyl or

20 R¹ is 3-(allyloxy)-4,6-dichlorophenyl or

R¹ is 3-(allyloxy)-4,6-difluorophenyl or

R¹ is 3-(allyloxy)-4-chlorophenyl or

25

R¹ is 3-(allyloxy)-4-fluorophenyl or

R¹ is 3-(allyloxy)-4-methylphenyl or

30 R¹ is 3-(allyloxy)-5,6-dichlorophenyl or

R¹ is 3-(allyloxy)-5,6-difluorophenyl or

R¹ is 3-(allyloxy)-5-chlorophenyl or

35

R¹ is 3-(allyloxy)-5-fluorophenyl or

R¹ is 3-(allyloxy)-5-methylphenyl or

R¹ is 3-(allyloxy)-6-chlorophenyl or

R¹ is 3-(allyloxy)-6-fluorophenyl or

5

R¹ is 3-(allyloxy)-6-methylphenyl or

R¹ is 3-(allyloxy)phenyl or

10 R¹ is 3-(cyanomethoxy)-2,4-dichlorophenyl or

R¹ is 3-(cyanomethoxy)-2,4-difluorophenyl or

R¹ is 3-(cyanomethoxy)-2,5-dichlorophenyl or

15

R¹ is 3-(cyanomethoxy)-2,5-difluorophenyl or

R¹ is 3-(cyanomethoxy)-2,6-dichlorophenyl or

20 R¹ is 3-(cyanomethoxy)-2,6-difluorophenyl or

R¹ is 3-(cyanomethoxy)-2-chlorophenyl or

R¹ is 3-(cyanomethoxy)-2-fluorophenyl or

25

R¹ is 3-(cyanomethoxy)-2-methylphenyl or

R¹ is 3-(cyanomethoxy)-4,5-dichlorophenyl or

30 R¹ is 3-(cyanomethoxy)-4,5-difluorophenyl or

R¹ is 3-(cyanomethoxy)-4,6-dichlorophenyl or

R¹ is 3-(cyanomethoxy)-4,6-difluorophenyl or

35

R¹ is 3-(cyanomethoxy)-4-chlorophenyl or

R¹ is 3-(cyanomethoxy)-4-fluorophenyl or

R¹ is 3-(cyanomethoxy)-4-methylphenyl or

R¹ is 3-(cyanomethoxy)-5,6-dichlorophenyl or

5

R¹ is 3-(cyanomethoxy)-5,6-difluorophenyl or

R¹ is 3-(cyanomethoxy)-5-chlorophenyl or

10 R¹ is 3-(cyanomethoxy)-5-fluorophenyl or

R¹ is 3-(cyanomethoxy)-5-methylphenyl or

R¹ is 3-(cyanomethoxy)-6-chlorophenyl or

15

R¹ is 3-(cyanomethoxy)-6-fluorophenyl or

R¹ is 3-(cyanomethoxy)-6-methylphenyl or

20 R¹ is 3-(cyanomethoxy)phenyl or

R¹ is 3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3-chloro-2-(prop-2-yn-1-yloxy)phenyl or

25

R¹ is 3-chloro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-chloro-2-formylphenyl or

30 R¹ is 3-chloro-4-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3-chloro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-chloro-4-formylphenyl or

35

R¹ is 3-fluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3-fluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-fluoro-2-formylphenyl or

R¹ is 3-fluoro-4-(prop-2-yn-1-yloxy)phenyl or

5

R¹ is 3-fluoro-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-fluoro-4-formylphenyl or

10 R¹ is 3-formyl-2-methylphenyl or

R¹ is 3-formyl-4-methylphenyl or

R¹ is 3-formyl-5-methylphenyl or

15

R¹ is 3-formyl-6-methylphenyl or

R¹ is 3-formylphenyl or

20 R¹ is 3-methyl-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 3-methyl-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-methyl-4-(prop-2-yn-1-yloxy)phenyl or

25

R¹ is 3-methyl-4-[(methylsulphonyl)oxy]phenyl or

R¹ is 3-[(hydroxyimino)methyl]-2,4-difluorophenyl or

30 R¹ is 3-[(hydroxyimino)methyl]-2,5-difluorophenyl or

R¹ is 3-[(hydroxyimino)methyl]-2,6-difluorophenyl or

R¹ is 3-[(hydroxyimino)methyl]-2-chlorophenyl or

35

R¹ is 3-[(hydroxyimino)methyl]-2-fluorophenyl or

R¹ is 3-[(hydroxyimino)methyl]-2-methylphenyl or

- R¹ is 3-[(hydroxyimino)methyl]-4,5-difluorophenyl or
- 5 R¹ is 3-[(hydroxyimino)methyl]-4,6-difluorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-4-chlorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-4-fluorophenyl or
- 10 R¹ is 3-[(hydroxyimino)methyl]-4-methylphenyl or
- R¹ is 3-[(hydroxyimino)methyl]-5,6-difluorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-5-chlorophenyl or
- 15 R¹ is 3-[(hydroxyimino)methyl]-5-fluorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-5-methylphenyl or
- 20 R¹ is 3-[(hydroxyimino)methyl]-6-chlorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-6-fluorophenyl or
- R¹ is 3-[(hydroxyimino)methyl]-6-methylphenyl or
- 25 R¹ is 3-[(hydroxyimino)methyl]phenyl or
- R¹ is 3-[(methoxyimino)methyl]-2,4-difluorophenyl or
- 30 R¹ is 3-[(methoxyimino)methyl]-2,5-difluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-2,6-difluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-2-chlorophenyl or
- 35 R¹ is 3-[(methoxyimino)methyl]-2-fluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-2-methylphenyl or

- R¹ is 3-[(methoxyimino)methyl]-4,5-difluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-4,6-difluorophenyl or
5
- R¹ is 3-[(methoxyimino)methyl]-4-chlorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-4-fluorophenyl or
- 10 R¹ is 3-[(methoxyimino)methyl]-4-methylphenyl or
- R¹ is 3-[(methoxyimino)methyl]-5,6-difluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-5-chlorophenyl or
15
- R¹ is 3-[(methoxyimino)methyl]-5-fluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-5-methylphenyl or
- 20 R¹ is 3-[(methoxyimino)methyl]-6-chlorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-6-fluorophenyl or
- R¹ is 3-[(methoxyimino)methyl]-6-methylphenyl or
25
- R¹ is 3-[(methoxyimino)methyl]phenyl or
- R¹ is 3-[(methylsulphonyl)oxy]phenyl or
- 30 R¹ is 4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl or
- R¹ is 4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl or
- R¹ is 4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl or
35
- R¹ is 4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl or
- R¹ is 4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 4,5-difluoro-2-formylphenyl or

5

R¹ is 4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl or

10 R¹ is 4,5-difluoro-3-formylphenyl or

R¹ is 4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl or

15

R¹ is 4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl or

20 R¹ is 4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 4,6-difluoro-2-formylphenyl or

25

R¹ is 4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl or

30 R¹ is 4,6-difluoro-3-formylphenyl or

R¹ is 4-(allyloxy)-2,3-dichlorophenyl or

R¹ is 4-(allyloxy)-2,3-difluorophenyl or

35

R¹ is 4-(allyloxy)-2,5-dichlorophenyl or

R¹ is 4-(allyloxy)-2,5-difluorophenyl or

R¹ is 4-(allyloxy)-2,6-dichlorophenyl or

R¹ is 4-(allyloxy)-2,6-difluorophenyl or

5

R¹ is 4-(allyloxy)-2-chlorophenyl or

R¹ is 4-(allyloxy)-2-fluorophenyl or

10 R¹ is 4-(allyloxy)-2-methylphenyl or

R¹ is 4-(allyloxy)-3,5-dichlorophenyl or

R¹ is 4-(allyloxy)-3,5-difluorophenyl or

15

R¹ is 4-(allyloxy)-3,6-dichlorophenyl or

R¹ is 4-(allyloxy)-3,6-difluorophenyl or

20 R¹ is 4-(allyloxy)-3-chlorophenyl or

R¹ is 4-(allyloxy)-3-fluorophenyl or

R¹ is 4-(allyloxy)-3-methylphenyl or

25

R¹ is 4-(allyloxy)phenyl or

R¹ is 4-(cyanomethoxy)-2,3-dichlorophenyl or

30 R¹ is 4-(cyanomethoxy)-2,3-difluorophenyl or

R¹ is 4-(cyanomethoxy)-2,5-dichlorophenyl or

R¹ is 4-(cyanomethoxy)-2,5-difluorophenyl or

35

R¹ is 4-(cyanomethoxy)-2,6-dichlorophenyl or

R¹ is 4-(cyanomethoxy)-2,6-difluorophenyl or

R¹ is 4-(cyanomethoxy)-2-chlorophenyl or

R¹ is 4-(cyanomethoxy)-2-fluorophenyl or

5

R¹ is 4-(cyanomethoxy)-2-methylphenyl or

R¹ is 4-(cyanomethoxy)-3,5-dichlorophenyl or

10 R¹ is 4-(cyanomethoxy)-3,5-difluorophenyl or

R¹ is 4-(cyanomethoxy)-3,6-dichlorophenyl or

R¹ is 4-(cyanomethoxy)-3,6-difluorophenyl or

15

R¹ is 4-(cyanomethoxy)-3-chlorophenyl or

R¹ is 4-(cyanomethoxy)-3-fluorophenyl or

20 R¹ is 4-(cyanomethoxy)-3-methylphenyl or

R¹ is 4-(cyanomethoxy)phenyl or

R¹ is 4-(prop-2-yn-1-yloxy)phenyl or

25

R¹ is 4-chloro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4-chloro-2-[(methylsulphonyl)oxy]phenyl or

30 R¹ is 4-chloro-2-formylphenyl or

R¹ is 4-chloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4-chloro-3-[(methylsulphonyl)oxy]phenyl or

35

R¹ is 4-chloro-3-formylphenyl or

R¹ is 4-fluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4-fluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 4-fluoro-2-formylphenyl or

5

R¹ is 4-fluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4-fluoro-3-[(methylsulphonyl)oxy]phenyl or

10 R¹ is 4-fluoro-3-formylphenyl or

R¹ is 4-formyl-2-methylphenyl or

R¹ is 4-formyl-3-methylphenyl or

15

R¹ is 4-formylphenyl or

R¹ is 4-methyl-2-(prop-2-yn-1-yloxy)phenyl or

20 R¹ is 4-methyl-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 4-methyl-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 4-methyl-3-[(methylsulphonyl)oxy]phenyl or

25

R¹ is 4-[(hydroxyimino)methyl]-2,3-difluorophenyl or

R¹ is 4-[(hydroxyimino)methyl]-2,5-difluorophenyl or

30 R¹ is 4-[(hydroxyimino)methyl]-2,6-difluorophenyl or

R¹ is 4-[(hydroxyimino)methyl]-2-chlorophenyl or

R¹ is 4-[(hydroxyimino)methyl]-2-fluorophenyl or

35

R¹ is 4-[(hydroxyimino)methyl]-2-methylphenyl or

R¹ is 4-[(hydroxyimino)methyl]-3,5-difluorophenyl or

- R¹ is 4-[(hydroxyimino)methyl]-3,6-difluorophenyl or
- R¹ is 4-[(hydroxyimino)methyl]-3-chlorophenyl or
5
- R¹ is 4-[(hydroxyimino)methyl]-3-fluorophenyl or
- R¹ is 4-[(hydroxyimino)methyl]-3-methylphenyl or
- 10 R¹ is 4-[(hydroxyimino)methyl]phenyl or
- R¹ is 4-[(methoxyimino)methyl]-2,3-difluorophenyl or
- R¹ is 4-[(methoxyimino)methyl]-2,5-difluorophenyl or
15
- R¹ is 4-[(methoxyimino)methyl]-2,6-difluorophenyl or
- R¹ is 4-[(methoxyimino)methyl]-2-chlorophenyl or
- 20 R¹ is 4-[(methoxyimino)methyl]-2-fluorophenyl or
- R¹ is 4-[(methoxyimino)methyl]-2-methylphenyl or
- R¹ is 4-[(methoxyimino)methyl]-3,5-difluorophenyl or
25
- R¹ is 4-[(methoxyimino)methyl]-3,6-difluorophenyl or
- R¹ is 4-[(methoxyimino)methyl]-3-chlorophenyl or
- 30 R¹ is 4-[(methoxyimino)methyl]-3-fluorophenyl or
- R¹ is 4-[(methoxyimino)methyl]-3-methylphenyl or
- R¹ is 4-[(methoxyimino)methyl]phenyl or
35
- R¹ is 4-[(methylsulphonyl)oxy]phenyl or
- R¹ is 5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl or

5

R¹ is 5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl or

10 R¹ is 5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 5,6-difluoro-2-formylphenyl or

R¹ is 5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl or

15

R¹ is 5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 5,6-difluoro-3-formylphenyl or

20 R¹ is 5-chloro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-chloro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 5-chloro-2-formylphenyl or

25

R¹ is 5-chloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-chloro-3-[(methylsulphonyl)oxy]phenyl or

30 R¹ is 5-chloro-3-formylphenyl or

R¹ is 5-fluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-fluoro-2-[(methylsulphonyl)oxy]phenyl or

35

R¹ is 5-fluoro-2-formylphenyl or

R¹ is 5-fluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-fluoro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 5-fluoro-3-formylphenyl or

5

R¹ is 5-methyl-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-methyl-2-[(methylsulphonyl)oxy]phenyl or

10 R¹ is 5-methyl-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 5-methyl-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 6-chloro-2-(prop-2-yn-1-yloxy)phenyl or

15

R¹ is 6-chloro-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 6-chloro-2-formylphenyl or

20 R¹ is 6-chloro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 6-chloro-3-[(methylsulphonyl)oxy]phenyl or

R¹ is 6-chloro-3-formylphenyl or

25

R¹ is 6-fluoro-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 6-fluoro-2-[(methylsulphonyl)oxy]phenyl or

30 R¹ is 6-fluoro-2-formylphenyl or

R¹ is 6-fluoro-3-(prop-2-yn-1-yloxy)phenyl or

R¹ is 6-fluoro-3-[(methylsulphonyl)oxy]phenyl or

35

R¹ is 6-fluoro-3-formylphenyl or

R¹ is 6-methyl-2-(prop-2-yn-1-yloxy)phenyl or

R¹ is 6-methyl-2-[(methylsulphonyl)oxy]phenyl or

R¹ is 6-methyl-3-(prop-2-yn-1-yloxy)phenyl or

5

R¹ is 6-methyl-3-[(methylsulphonyl)oxy]phenyl steht.

The radical definitions specified above can be combined with one another as desired. Moreover, individual definitions may not apply.

10

According to the type of substituents defined above, the compounds of the formula **(I)** have acidic or basic properties and can form salts, possibly also internal salts or adducts, with inorganic or organic acids or with bases or with metal ions. If the compounds of the formula **(I)** bear amino, alkylamino or other groups which induce basic properties, these compounds can be reacted with acids to give salts, or they are obtained directly as salts by the synthesis. If the compounds of the formula **(I)** bear hydroxyl, carboxyl or other groups which induce acidic properties, these compounds can be reacted with bases to give salts. Suitable bases are, for example, hydroxides, carbonates, hydrogencarbonates of the alkali metals and alkaline earth metals, especially those of sodium, potassium, magnesium and calcium, and also ammonia, primary, secondary and tertiary amines having C₁-C₄-alkyl groups, mono-, di- and trialkanolamines of C₁-C₄-alkanols, choline and chlorocholine.

20

25

The salts thus obtainable likewise have fungicidal properties.

30

Examples of inorganic acids are hydrohalic acids, such as hydrogen fluoride, hydrogen chloride, hydrogen bromide and hydrogen iodide, sulphuric acid, phosphoric acid and nitric acid, and acidic salts, such as NaHSO₄ and KHSO₄. Useful organic acids include, for example, formic acid, carbonic acid and alkanolic acids such as acetic acid, trifluoroacetic acid, trichloroacetic acid and propionic acid, and also glycolic

35

acid, thiocyanic acid, lactic acid, succinic acid, citric acid, benzoic acid, cinnamic acid, oxalic acid, saturated or mono- or diunsaturated C₆-C₂₀ fatty acids, alkylsulphuric monoesters, alkylsulphonic acids (sulphonic acids having
5 straight-chain or branched alkyl radicals having 1 to 20 carbon atoms), arylsulphonic acids or aryldisulphonic acids (aromatic radicals, such as phenyl and naphthyl, which bear one or two sulphonic acid groups), alkylphosphonic acids (phosphonic acids having straight-chain or branched alkyl
10 radicals having 1 to 20 carbon atoms), arylphosphonic acids or aryldiphosphonic acids (aromatic radicals, such as phenyl and naphthyl, which bear one or two phosphonic acid radicals), where the alkyl and aryl radicals may bear further substituents, for example p-toluenesulphonic acid, salicylic
15 acid, p-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, etc.

Useful metal ions are especially the ions of the elements of the second main group, especially calcium and magnesium, of
20 the third and fourth main group, especially aluminium, tin and lead, and also of the first to eighth transition groups, especially chromium, manganese, iron, cobalt, nickel, copper, zinc and others. Particular preference is given to the metal ions of the elements of the fourth period. Here, the metals
25 can be present in the various valencies that they can assume.

Optionally substituted groups may be mono- or polysubstituted, where the substituents in the case of polysubstitutions may be identical or different.
30

In the definitions of the symbols given in the above formulae, collective terms were used, which are generally representative of the following substituents:

35 **Halogen:** fluorine, chlorine, bromine and iodine and preferably fluorine, chlorine, bromine and more preferably fluorine, chlorine.

Alkyl: saturated, straight-chain or branched hydrocarbyl radicals having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms, for example (but not limited to) C₁-C₆-alkyl such as methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl. This definition also applies to alkyl as part of a composite substituent, for example cycloalkylalkyl, hydroxyalkyl etc., unless defined elsewhere like, for example, alkylthio, alkylsufinyl, alkylsulphonyl, haloalkyl or haloalkylthio. If the alkyl is at the end of a composite substituent, as, for example, in alkylcycloalkyl, the part of the composite substituent at the start, for example the cycloalkyl, may be mono- or polysubstituted identically or differently and independently by alkyl. The same also applies to composite substituents in which other radicals, for example alkenyl, alkynyl, hydroxyl, halogen, formyl etc., are at the end.

25

Alkenyl: unsaturated, straight-chain or branched hydrocarbyl radicals having 2 to 8, preferably 2 to 6, carbon atoms and one double bond in any position, for example (but not limited to) C₂-C₆-alkenyl such as ethenyl, 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-1-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-

35

pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 5 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, 2,2-dimethyl-3-butenyl, 2,3-10 dimethyl-1-butenyl, 2,3-dimethyl-2-butenyl, 2,3-dimethyl-3-butenyl, 3,3-dimethyl-1-butenyl, 3,3-dimethyl-2-butenyl, 1-ethyl-1-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl, 2-ethyl-1-butenyl, 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-propenyl, 1-ethyl-1-methyl-2-propenyl, 1-ethyl-2-15 methyl-1-propenyl and 1-ethyl-2-methyl-2-propenyl. This definition also applies to alkenyl as part of a composite substituent, for example haloalkenyl etc., unless defined elsewhere.

20 **Alkynyl:** straight-chain or branched hydrocarbyl groups having 2 to 8, preferably 2 to 6, carbon atoms and one triple bond in any position, for example (but not limited to) C₂-C₆-alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 1-methyl-2-propynyl, 1-pentylnyl, 2-pentylnyl, 25 pentylnyl, 4-pentylnyl, 1-methyl-2-butylnyl, 1-methyl-3-butylnyl, 2-methyl-3-butylnyl, 3-methyl-1-butylnyl, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentylnyl, 1-methyl-3-pentylnyl, 1-methyl-4-pentylnyl, 2-methyl-3-pentylnyl, 2-methyl-30 4-pentylnyl, 3-methyl-1-pentylnyl, 3-methyl-4-pentylnyl, 4-methyl-1-pentylnyl, 4-methyl-2-pentylnyl, 1,1-dimethyl-2-butylnyl, 1,1-dimethyl-3-butylnyl, 1,2-dimethyl-3-butylnyl, 2,2-dimethyl-3-butylnyl, 3,3-dimethyl-1-butylnyl, 1-ethyl-2-butylnyl, 1-ethyl-3-butylnyl, 2-ethyl-3-butylnyl and 1-ethyl-1-methyl-2-35 propynyl. This definition also applies to alkynyl as part of a composite substituent, for example haloalkynyl etc., unless defined elsewhere.

Alkoxy: saturated, straight-chain or branched alkoxy radicals having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms, for example (but not limited to) C₁-C₆-alkoxy such as methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy, 1,1-dimethylethoxy, pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy. This definition also applies to alkoxy as part of a composite substituent, for example haloalkoxy, alkynylalkoxy, etc., unless defined elsewhere.

Alkylthio: saturated, straight-chain or branched alkylthio radicals having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms, for example (but not limited to) C₁-C₆-alkylthio such as methylthio, ethylthio, propylthio, 1-methylethylthio, butylthio, 1-methylpropylthio, 2-methylpropylthio, 1,1-dimethylethylthio, pentylthio, 1-methylbutylthio, 2-methylbutylthio, 3-methylbutylthio, 2,2-dimethylpropylthio, 1-ethylpropylthio, hexylthio, 1,1-dimethylpropylthio, 1,2-dimethylpropylthio, 1-methylpentylthio, 2-methylpentylthio, 3-methylpentylthio, 4-methylpentylthio, 1,1-dimethylbutylthio, 1,2-dimethylbutylthio, 1,3-dimethylbutylthio, 2,2-dimethylbutylthio, 2,3-dimethylbutylthio, 3,3-dimethylbutylthio, 1-ethylbutylthio, 2-ethylbutylthio, 1,1,2-trimethylpropylthio, 1,2,2-trimethylpropylthio, 1-ethyl-1-methylpropylthio and 1-ethyl-2-methylpropylthio. This definition also applies to alkylthio as part of a composite substituent, for example haloalkylthio etc., unless defined elsewhere.

Alkoxy carbonyl: an alkoxy group which has 1 to 6, preferably 1

to 3, carbon atoms (as specified above) and is bonded to the skeleton via a carbonyl group (-CO-). This definition also applies to alkoxy carbonyl as part of a composite substituent, for example cycloalkylalkoxy carbonyl etc., unless defined
5 elsewhere.

Alkylsulphinyl: saturated, straight-chain or branched alkylsulphinyl radicals having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms, for example (but not
10 limited to) C₁-C₆-alkylsulphinyl such as methylsulphinyl, ethylsulphinyl, propylsulphinyl, 1-methylethylsulphinyl, butylsulphinyl, 1-methylpropylsulphinyl, 2-methylpropylsulphinyl, 1,1-dimethylethylsulphinyl, pentylsulphinyl, 1-methylbutylsulphinyl, 2-methylbutylsulphinyl, 3-methylbutylsulphinyl, 2,2-dimethylpropylsulphinyl, 1-ethylpropylsulphinyl, hexylsulphinyl, 1,1-dimethylpropylsulphinyl, 1,2-dimethylpropylsulphinyl, 1-methylpentylsulphinyl, 2-methylpentylsulphinyl, 3-methylpentylsulphinyl, 4-methylpentylsulphinyl, 1,1-dimethylbutylsulphinyl, 1,2-dimethylbutylsulphinyl, 1,3-dimethylbutylsulphinyl, 2,2-dimethylbutylsulphinyl, 2,3-dimethylbutylsulphinyl, 3,3-dimethylbutylsulphinyl, 1-ethylbutylsulphinyl, 2-ethylbutylsulphinyl, 1,1,2-trimethylpropylsulphinyl, 1,2,2-trimethylpropylsulphinyl, 1-ethyl-1-methylpropylsulphinyl and 1-ethyl-2-methylpropylsulphinyl. This definition also applies to alkylsulphinyl as part of a composite substituent, for example haloalkylsulphinyl etc., unless defined elsewhere.

Alkylsulphonyl: saturated, straight-chain or branched alkylsulphonyl radicals having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms, for example (but not limited to) C₁-C₆-alkylsulphonyl such as methylsulphonyl, ethylsulphonyl, propylsulphonyl, 1-methylethylsulphonyl, butylsulphonyl, 1-methylpropylsulphonyl, 2-methylpropylsulphonyl, 1,1-dimethylethylsulphonyl, pentylsulphonyl, 1-methylbutylsulphonyl, 2-methylbutylsulphonyl, 3-methylbutylsulphonyl, 2,2-

dimethylpropylsulphonyl, 1-ethylpropylsulphonyl,
 hexylsulphonyl, 1,1-dimethylpropylsulphonyl, 1,2-
 dimethylpropylsulphonyl, 1-methylpentylsulphonyl, 2-
 methylpentylsulphonyl, 3-methylpentylsulphonyl, 4-
 5 methylpentylsulphonyl, 1,1-dimethylbutylsulphonyl, 1,2-
 dimethylbutylsulphonyl, 1,3-dimethylbutylsulphonyl, 2,2-
 dimethylbutylsulphonyl, 2,3-dimethylbutylsulphonyl, 3,3-
 dimethylbutylsulphonyl, 1-ethylbutylsulphonyl, 2-
 ethylbutylsulphonyl, 1,1,2-trimethylpropylsulphonyl, 1,2,2-
 10 trimethylpropylsulphonyl, 1-ethyl-1-methylpropylsulphonyl and
 1-ethyl-2-methylpropylsulphonyl. This definition also applies
 to alkylsulphonyl as part of a composite substituent, for
 example alkylsulphonylalkyl etc., unless defined elsewhere.

15 **Cycloalkyl:** monocyclic, saturated hydrocarbyl groups having 3
 to 10, preferably 3 to 8 and more preferably 3 to 6 carbon
 ring members, for example (but not limited to) cyclopropyl,
 cyclopentyl and cyclohexyl. This definition also applies to
 cycloalkyl as part of a composite substituent, for example
 20 cycloalkylalkyl etc., unless defined elsewhere.

Cycloalkenyl: monocyclic, partially unsaturated hydrocarbyl
 groups having 3 to 10, preferably 3 to 8 and more preferably 3
 to 6 carbon ring members, for example (but not limited to)
 25 cyclopropenyl, cyclopentenyl and cyclohexenyl. This definition
 also applies to cycloalkenyl as part of a composite
 substituent, for example cycloalkenylalkyl etc., unless
 defined elsewhere.

30 **Cycloalkoxy:** monocyclic, saturated cycloalkyloxy radicals
 having 3 to 10, preferably 3 to 8 and more preferably 3 to 6
 carbon ring members, for example (but not limited to)
 cyclopropyloxy, cyclopentyloxy and cyclohexyloxy. This
 definition also applies to cycloalkoxy as part of a composite
 35 substituent, for example cycloalkoxyalkyl etc., unless defined
 elsewhere.

Haloalkyl: straight-chain or branched alkyl groups having 1 to

8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms (as specified above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as specified above, for example (but not limited to) C₁-C₃-haloalkyl such as

5 chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-

10 trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl and 1,1,1-trifluoroprop-2-yl. This definition also applies to haloalkyl as part of a composite substituent, for example haloalkylaminoalkyl etc.,

15 unless defined elsewhere.

Haloalkenyl and haloalkynyl are defined analogously to haloalkyl except that, instead of alkyl groups, alkenyl and alkynyl groups are present as part of the substituent.

20

Haloalkoxy: straight-chain or branched alkoxy groups having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms (as specified above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as

25 specified above, for example (but not limited to) C₁-C₃-haloalkoxy such as chloromethoxy, bromomethoxy, dichloromethoxy, trichloromethoxy, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chlorofluoromethoxy, dichlorofluoromethoxy, chlorodifluoromethoxy, 1-chloroethoxy,

30 1-bromoethoxy, 1-fluoroethoxy, 2-fluoroethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-2-fluoroethoxy, 2-chloro-2-difluoroethoxy, 2,2-dichloro-2-fluoroethoxy, 2,2,2-trichloroethoxy, pentafluoroethoxy and

35 haloalkoxy as part of a composite substituent, for example haloalkoxyalkyl etc., unless defined elsewhere.

Haloalkylthio: straight-chain or branched alkylthio groups

having 1 to 8, preferably 1 to 6 and more preferably 1 to 3 carbon atoms (as specified above), where some or all of the hydrogen atoms in these groups may be replaced by halogen atoms as specified above, for example (but not limited to) C₁-
 5 C₃-haloalkylthio such as chloromethylthio, bromomethylthio, dichloromethylthio, trichloromethylthio, fluoromethylthio, difluoromethylthio, trifluoromethylthio, chlorofluoromethylthio, dichlorofluoromethylthio, chlorodifluoromethylthio, 1-chloroethylthio, 1-bromoethylthio,
 10 1-fluoroethylthio, 2-fluoroethylthio, 2,2-difluoroethylthio, 2,2,2-trifluoroethylthio, 2-chloro-2-fluoroethylthio, 2-chloro-2-difluoroethylthio, 2,2-dichloro-2-fluoroethylthio, 2,2,2-trichloroethylthio, pentafluoroethylthio and 1,1,1-trifluoroprop-2-ylthio. This definition also applies to
 15 haloalkylthio as part of a composite substituent, for example haloalkylthioalkyl etc., unless defined elsewhere.

Heteroaryl: 5 or 6-membered, fully unsaturated monocyclic ring system containing one to four heteroatoms from the group of
 20 oxygen, nitrogen and sulphur; if the ring contains more than one oxygen atom, they are not directly adjacent;

5-membered heteroaryl containing one to four nitrogen atoms or one to three nitrogen atoms and one sulphur or oxygen atom: 5-
 25 membered heteroaryl groups which, in addition to carbon atoms, may contain one to four nitrogen atoms or one to three nitrogen atoms and one sulphur or oxygen atom as ring members, for example (but not limited thereto) 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 3-isoxazolyl, 4-
 30 isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-
 35 5-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl and 1,3,4-triazol-2-yl;

nitrogen-bonded 5-membered heteroaryl containing one to four

nitrogen atoms, or benzofused nitrogen-bonded 5-membered heteroaryl containing one to three nitrogen atoms: 5-membered heteroaryl groups which, in addition to carbon atoms, may contain one to four nitrogen atoms or one to three nitrogen atoms as ring members and in which two adjacent carbon ring members or one nitrogen and one adjacent carbon ring member may be bridged by a buta-1,3-diene-1,4-diyl group in which one or two carbon atoms may be replaced by nitrogen atoms, where these rings are attached to the skeleton via one of the nitrogen ring members, for example (but not limited to) 1-pyrrolyl, 1-pyrazolyl, 1,2,4-triazol-1-yl, 1-imidazolyl, 1,2,3-triazol-1-yl and 1,3,4-triazol-1-yl;

6-membered heteroaryl which contains one to four nitrogen atoms: 6-membered heteroaryl groups which, in addition to carbon atoms, may contain, respectively, one to three and one to four nitrogen atoms as ring members, for example (but not limited thereto) 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl, 1,2,4-triazin-3-yl and 1,2,4,5-tetrazin-3-yl;

benzofused 5-membered heteroaryl containing one to three nitrogen atoms or one nitrogen atom and one oxygen or sulphur atom: for example (but not limited to) indol-1-yl, indol-2-yl, indol-3-yl, indol-4-yl, indol-5-yl, indol-6-yl, indol-7-yl, benzimidazol-1-yl, benzimidazol-2-yl, benzimidazol-4-yl, benzimidazol-5-yl, indazol-1-yl, indazol-3-yl, indazol-4-yl, indazol-5-yl, indazol-6-yl, indazol-7-yl, indazol-2-yl, 1-benzofuran-2-yl, 1-benzofuran-3-yl, 1-benzofuran-4-yl, 1-benzofuran-5-yl, 1-benzofuran-6-yl, 1-benzofuran-7-yl, 1-benzothiophen-2-yl, 1-benzothiophen-3-yl, 1-benzothiophen-4-yl, 1-benzothiophen-5-yl, 1-benzothiophen-6-yl, 1-benzothiophen-7-yl, 1,3-benzothiazol-2-yl, 1,3-benzothiazol-4-yl, 1,3-benzothiazol-5-yl, 1,3-benzothiazol-6-yl, 1,3-benzothiazol-7-yl, 1,3-benzoxazol-2-yl, 1,3-benzoxazol-4-yl, 1,3-benzoxazol-5-yl, 1,3-benzoxazol-6-yl and 1,3-benzoxazol-7-yl;

benzofused 6-membered heteroaryl which contains one to three nitrogen atoms: for example (but not limited to) quinolin-2-yl, quinolin-3-yl, quinolin-4-yl, quinolin-5-yl, quinolin-6-yl, quinolin-7-yl, quinolin-8-yl, isoquinolin-1-yl, isoquinolin-3-yl, isoquinolin-4-yl, isoquinolin-5-yl, isoquinolin-6-yl, isoquinolin-7-yl and isoquinolin-8-yl.

This definition also applies to heteroaryl as part of a composite substituent, for example heteroarylalkyl etc., unless defined elsewhere.

Heterocyclyl: three- to fifteen-membered, preferably three- to nine-membered, saturated or partially unsaturated heterocycle containing one to four heteroatoms from the group of oxygen, nitrogen and sulphur: mono, bi- or tricyclic heterocycles which contain, in addition to carbon ring members, one to three nitrogen atoms and/or one oxygen or sulphur atom or one or two oxygen and/or sulphur atoms; if the ring contains more than one oxygen atom, they are not directly adjacent; for example (but not limited to) oxiranyl, aziridinyl, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothienyl, 3-tetrahydrothienyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 3-isoxazolidinyl, 4-isoxazolidinyl, 5-isoxazolidinyl, 3-isothiazolidinyl, 4-isothiazolidinyl, 5-isothiazolidinyl, 3-pyrazolidinyl, 4-pyrazolidinyl, 5-pyrazolidinyl, 2-oxazolidinyl, 4-oxazolidinyl, 5-oxazolidinyl, 2-thiazolidinyl, 4-thiazolidinyl, 5-thiazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 1,2,4-oxadiazolidin-3-yl, 1,2,4-oxadiazolidin-5-yl, 1,2,4-thiadiazolidin-3-yl, 1,2,4-thiadiazolidin-5-yl, 1,2,4-triazolidin-3-yl, 1,3,4-oxadiazolidin-2-yl, 1,3,4-thiadiazolidin-2-yl, 1,3,4-triazolidin-2-yl, 2,3-dihydrofur-2-yl, 2,3-dihydrofur-3-yl, 2,4-dihydrofur-2-yl, 2,4-dihydrofur-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,4-dihydrothien-2-yl, 2,4-dihydrothien-3-yl, 2-pyrrolin-2-yl, 2-pyrrolin-3-yl, 3-pyrrolin-2-yl, 3-pyrrolin-3-yl, 2-isoxazolin-3-yl, 3-isoxazolin-3-yl, 4-isoxazolin-3-yl, 2-isoxazolin-4-yl, 3-isoxazolin-4-yl, 4-isoxazolin-4-yl, 2-isoxazolin-5-yl, 3-

isoxazolin-5-yl, 4-isoxazolin-5-yl, 2-isothiazolin-3-yl, 3-isothiazolin-3-yl, 4-isothiazolin-3-yl, 2-isothiazolin-4-yl, 3-isothiazolin-4-yl, 4-isothiazolin-4-yl, 2-isothiazolin-5-yl, 3-isothiazolin-5-yl, 4-isothiazolin-5-yl, 2,3-dihydropyrazol-1-yl, 2,3-dihydropyrazol-2-yl, 2,3-dihydropyrazol-3-yl, 2,3-dihydropyrazol-4-yl, 2,3-dihydropyrazol-5-yl, 3,4-dihydropyrazol-1-yl, 3,4-dihydropyrazol-3-yl, 3,4-dihydropyrazol-4-yl, 3,4-dihydropyrazol-5-yl, 4,5-dihydropyrazol-1-yl, 4,5-dihydropyrazol-3-yl, 4,5-dihydropyrazol-4-yl, 4,5-dihydropyrazol-5-yl, 2,3-dihydrooxazol-2-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrooxazol-4-yl, 2,3-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 3,4-dihydrooxazol-5-yl, 3,4-dihydrooxazol-2-yl, 3,4-dihydrooxazol-3-yl, 3,4-dihydrooxazol-4-yl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1,3-dioxan-5-yl, 2-tetrahydropyranyl, 4-tetrahydropyranyl, 2-tetrahydrothienyl, 3-hexahydropyridazinyl, 4-hexahydropyridazinyl, 2-hexahydropyrimidinyl, 4-hexahydropyrimidinyl, 5-hexahydropyrimidinyl, 2-piperazinyl, 1,3,5-hexahydrotriazin-2-yl and 1,2,4-hexahydrotriazin-3-yl. This definition also applies to heterocyclyl as part of a composite substituent, for example heterocyclylalkyl etc., unless defined elsewhere.

25 **Leaving group:** S_N1 or S_N2 leaving group, for example chlorine, bromine, iodine, alkylsulphonates (-OSO₂-alkyl, e.g. -OSO₂CH₃, -OSO₂CF₃) or arylsulphonates (-OSO₂-aryl, e.g. -OSO₂Ph, -OSO₂PhMe).

30 Not included are combinations which are against natural laws and which the person skilled in the art would therefore exclude based on his/her expert knowledge. Ring structures having three or more adjacent oxygen atoms, for example, are excluded.

35

Elucidation of the preparation processes and intermediates

The heteroarylpiperidine and -piperazine derivatives of the

formula **(I)** can be prepared in different ways. First of all, the possible processes are shown schematically below. Unless indicated otherwise, the radicals given have the meanings given above.

5

The processes for preparing compounds of the formula **(I)** are optionally performed using one or more reaction auxiliaries.

Useful reaction auxiliaries are, as appropriate, inorganic or organic bases or acid acceptors. These preferably include 10 alkali metal or alkaline earth metal acetates, amides, carbonates, hydrogencarbonates, hydrides, hydroxides or alkoxides, for example sodium acetate, potassium acetate or calcium acetate, lithium amide, sodium amide, potassium amide or calcium amide, sodium carbonate, potassium carbonate or 15 calcium carbonate, sodium hydrogencarbonate, potassium hydrogencarbonate or calcium hydrogencarbonate, lithium hydride, sodium hydride, potassium hydride or calcium hydride, lithium hydroxide, sodium hydroxide, potassium hydroxide or calcium hydroxide, sodium methoxide, ethoxide, n- or i- 20 propoxide, n-, i-, s- or t-butoxide or potassium methoxide, ethoxide, n- or i-propoxide, n-, i-, s- or t-butoxide; and also basic organic nitrogen compounds, for example trimethylamine, triethylamine, tripropylamine, tributylamine, ethyldiisopropylamine, N,N-dimethylcyclohexylamine, 25 dicyclohexylamine, ethyldicyclohexylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine, pyridine, 2-methyl-, 3-methyl-, 4-methyl-, 2,4-dimethyl-, 2,6-dimethyl-, 3,4-dimethyl- and 3,5-dimethylpyridine, 5-ethyl-2-methylpyridine, 30 4-dimethylaminopyridine, N-methylpiperidine, 1,4-diazabicyclo[2.2.2]-octane (DABCO), 1,5-diazabicyclo[4.3.0]-non-5-ene (DBN) or 1,8-diazabicyclo[5.4.0]-undec-7-ene (DBU).

Useful reaction auxiliaries are, as appropriate, inorganic or 35 organic acids. These preferably include inorganic acids, for example hydrogen fluoride, hydrogen chloride, hydrogen bromide and hydrogen iodide, sulphuric acid, phosphoric acid and nitric acid, and acidic salts such as NaHSO₄ and KHSO₄, or

organic acids, for example, formic acid, carbonic acid and alkanolic acids such as acetic acid, trifluoroacetic acid, trichloroacetic acid and propionic acid, and also glycolic acid, thiocyanic acid, lactic acid, succinic acid, citric acid, benzoic acid, cinnamic acid, oxalic acid, saturated or mono- or diunsaturated C₆-C₂₀ fatty acids, alkylsulphuric monoesters, alkylsulphonic acids (sulphonic acids having straight-chain or branched alkyl radicals having 1 to 20 carbon atoms), arylsulphonic acids or aryldisulphonic acids (aromatic radicals, such as phenyl and naphthyl, which bear one or two sulphonic acid groups), alkylphosphonic acids (phosphonic acids having straight-chain or branched alkyl radicals having 1 to 20 carbon atoms), arylphosphonic acids or aryldiphosphonic acids (aromatic radicals, such as phenyl and naphthyl, which bear one or two phosphonic acid radicals), where the alkyl and aryl radicals may bear further substituents, for example p-toluenesulphonic acid, salicylic acid, p-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, etc.

20

The processes are optionally performed using one or more diluents. Useful diluents are virtually all inert organic solvents. These preferably include aliphatic and aromatic, optionally halogenated hydrocarbons, such as pentane, hexane, heptane, cyclohexane, petroleum ether, benzene, ligroin, benzene, toluene, xylene, methylene chloride, ethylene chloride, chloroform, carbon tetrachloride, chlorobenzene and o-dichlorobenzene, ethers such as diethyl ether and dibutyl ether, glycol dimethyl ether and diglycol dimethyl ether, tetrahydrofuran and dioxane, ketones such as acetone, methyl ethyl ketone, methyl isopropyl ketone and methyl isobutyl ketone, esters, such as methyl acetate and ethyl acetate, nitriles, for example acetonitrile and propionitrile, amides, for example dimethylformamide, dimethylacetamide and N-methylpyrrolidone, and also dimethyl sulphoxide, tetramethylenesulphone and hexamethylphosphoramide and DMPU.

35

In the processes, the reaction temperatures can be varied

within a relatively wide range. In general, the temperatures employed are between 0°C and 250°C, preferably temperatures between 10°C and 185°C.


- 5 The reaction time varies as a function of the scale of the reaction and of the reaction temperature, but is generally between a few minutes and 48 hours.

The processes are generally performed under standard pressure. However, it is also possible to work under elevated or reduced pressure.

For performance of the processes, the starting materials required in each case are generally used in approximately equimolar amounts. However, it is also possible to use one of the components used in each case in a relatively large excess.

Process A

20 Scheme 1: Process A



in which the symbols R^{A1} , R^{A2} , R^{B1} , R^{B2} , R^{10} , p , R^2 , X , G , Q , L^1 , L^2 and R^1 each have the general definitions specified above.

The amides **(Ia)** obtained in the performance of process A (Scheme 1) can be converted by means of methods described in the literature to the corresponding thioamides **(Ib)** (e.g. *Bioorganic & Medicinal Chemistry Letters*, **2009**, 19(2), 462-468). This involves reacting the compounds of the formula **(Ia)** typically with phosphorus pentasulphide or 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane 2,4-disulphide (Lawesson's reagent) (see Scheme 7, Process F).

Process A is preferably carried out using one or more diluents. The preferred solvents are toluene, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane.


5

After the reaction has ended, the compounds **(Ib)** are separated from the reaction mixture by one of the customary separation techniques. If necessary, the compounds are purified by recrystallization or chromatography.

10

Process B

Scheme 2: Process B



15


in which the symbols R^{A1} , R^{A2} , R^{B1} , R^{B2} , Y , R^{10} , p , R^2 , X , G , Q , L^1 , L^2 and R^1 each have the general definitions specified above and W^1 and W^2 are functional groups suitable for the formation of the desired heterocycle **Q**

20


In general, it is possible to prepare compounds of the formula **(I)** from corresponding compounds **(XXIII)** and **(XXIV)** with suitable functional groups W^1 and W^2 **(I)** (see Scheme 2, process B). The possible functional groups for W^1 and W^2 are, for example, aldehydes, ketones, esters, carboxylic acids, amides, thioamides, nitriles, alcohols, thiols, hydrazines, oximes, amidines, amide oximes, olefins, acetylenes, halides, alkyl halides, methanesulphonates, trifluoromethanesulphonates, boronic acids, boronates, dialkyl acetal, ketoximes, etc., which can form the desired heterocycle **Q** under suitable reaction conditions. There are numerous literature methods for the preparation of heterocycles (see WO2008/013622; *Comprehensive Heterocyclic Chemistry Vol. 4-6*, A. R. Katritzky

30


and C. W. Rees editors, Pergamon Press, New York, 1984; *Comprehensive Heterocyclic Chemistry II*, Vol 2-4, A. R. Katritzky, C. W. Rees and E. F. Scriven editors, Pergamon Press, New York, 1996; *The Chemistry of Heterocyclic Compounds*, E. C. Taylor, editor, Wiley, New York; *Rodd's Chemistry of Carbon Compounds*, Vol. 2-4, Elsevier, New York; *Synthesis*, **1982**, 6, 508-509; *Tetrahedron*, **2000**, 56, 1057-1094).



(XXXIa)



(XXXIb)



(XXXIc)

10

The substances of the formula **(I)** prepared with the aid of the inventive compounds **(XIII)** have potent microbicidal activity and can be used for control of unwanted microorganisms, such as fungi and bacteria, in crop protection and in the protection of materials.

15

The heteroarylpiperidine and -piperazine derivatives of the formula **(I)** have very good fungicidal properties and can be used in crop protection, for example for control of Plasmodiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

20

Bactericides can be used in crop protection, for example, for control of Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomyetaceae.

25

The fungicidal compositions prepared with the aid of the inventive compounds **(XIII)** can be used for curative or protective control of phytopathogenic fungi. The compositions for controlling phytopathogenic fungi in crop protection comprise an effective but non-phytotoxic amount of the active ingredients. An "effective but non-phytotoxic amount" means an amount of the composition which is sufficient to control the fungal disease of the plant in a satisfactory manner or to eradicate the fungal disease completely, and which, at the

30

35

same time, does not cause any significant symptoms of phytotoxicity. In general, this application rate may vary within a relatively wide range. It depends on several factors, for example on the fungus to be controlled, the plant, the climatic conditions and the ingredients of the fungicidal compositions.

All plants and plant parts can be treated. Plants are understood here to mean all plants and plant populations, such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants may be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and including the plant cultivars which are protectable and non-protectable by plant breeders' rights. Plant parts are understood to mean all parts and organs of plants above and below the ground, such as shoot, leaf, flower and root, examples of which include leaves, needles, stalks, stems, flowers, fruit bodies, fruits and seeds, and also roots, tubers and rhizomes. The plant parts also include harvested material and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, slips and seeds.

Plants which can be treated include the following: cotton, flax, grapevine, fruit, vegetables, such as *Rosaceae* sp. (for example pome fruits such as apples and pears, but also stone fruits such as apricots, cherries, almonds and peaches, and soft fruits such as strawberries), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for example banana trees and plantations), *Rubiaceae* sp. (for example coffee), *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for example lemons, oranges and grapefruit); *Solanaceae* sp. (for example tomatoes), *Liliaceae* sp., *Asteraceae* sp. (for example lettuce), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp. (for

example cucumber), *Alliaceae sp.* (for example leek, onion),
Papilionaceae sp. (for example peas); major crop plants, such
 as *Gramineae sp.* (for example maize, turf, cereals such as
 5 wheat, rye, rice, barley, oats, millet and triticale),
Asteraceae sp. (for example sunflower), *Brassicaceae sp.* (for
 example white cabbage, red cabbage, broccoli, cauliflower,
 Brussels sprouts, pak choi, kohlrabi, radishes, and oilseed
 rape, mustard, horseradish and cress), *Fabaceae sp.* (for
 example bean, peanuts), *Papilionaceae sp.* (for example soya
 10 bean), *Solanaceae sp.* (for example potatoes), *Chenopodiaceae*
sp. (for example sugar beet, fodder beet, swiss chard,
 beetroot); useful plants and ornamental plants for gardens and
 wooded areas; and genetically modified varieties of each of
 these plants.

15

Non-limiting examples of pathogens of fungal diseases which
 can be treated include:

and stem blight (*Diaporthe phaseolorum*), stem canker
 20 (*Diaporthe phaseolorum var. caulivora*), phytophthora rot
 (*Phytophthora megasperma*), brown stem rot (*Phialophora*
gregata), pythium rot (*Pythium aphanidermatum*, *Pythium*
irregulare, *Pythium debaryanum*, *Pythium myriotylum*, *Pythium*
ultimum), rhizoctonia root rot, stem decay, and damping-off
 25 (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia*
sclerotiorum), sclerotinia southern blight (*Sclerotinia*
rolfsii), thielaviopsis root rot (*Thielaviopsis basicola*).

The active ingredients prepared with the aid of the inventive
 30 compounds also have very good fortifying action in plants.
 They are therefore suitable for mobilizing the plant's own
 defences against attack by undesirable microorganisms.

Plant-fortifying (resistance-inducing) substances are
 35 understood to mean, in the present context, those substances
 which are capable of stimulating the defence system of plants
 in such a way that the treated plants, when subsequently
 inoculated with undesirable microorganisms, develop a high

degree of resistance to these microorganisms.

In the present case, undesirable microorganisms are understood to mean phytopathogenic fungi and bacteria. The substances
5 prepared with the aid of the inventive compounds can thus be used to protect plants for a certain period after the treatment against attack by the pathogens mentioned. The period for which protection is provided generally extends over 1 to 10 days, preferably 1 to 7 days, after the treatment of
10 the plants with the active ingredients.

The fact that the active ingredients are well tolerated by plants at the concentrations required for controlling plant diseases allows the treatment of above-ground parts of plants,
15 of propagation stock and seeds, and of the soil.

The active ingredients prepared with the aid of the inventive compounds can be used particularly successfully to control diseases in viticulture and potato, fruit and vegetable
20 growing, for example against powdery mildew fungi, Oomycetes, for example Phytophthora, Plasmopara, Pseudoperonospora and Pythium species.

The active ingredients prepared with the aid of the inventive
25 compounds are also suitable for enhancing harvest yield. In addition, they have low toxicity and are well tolerated by plants.

If appropriate, the compounds prepared with the aid of the
30 inventive compounds can, at certain concentrations or application rates, also be used as herbicides, safeners, growth regulators or agents to improve plant properties, or as microbicides, for example as fungicides, antimycotics, bactericides, viricides (including agents against viroids) or
35 as agents against MLO (mycoplasma-like organisms) and RLO (rickettsia-like organisms). If appropriate, they can also be used as insecticides. If appropriate, they can also be used as intermediates or precursors for the synthesis of other active

ingredients.

The active ingredients prepared with the aid of the inventive compounds, when they are well tolerated by plants, have
5 favourable homeotherm toxicity and are well tolerated by the environment, are suitable for protecting plants and plant organs, for enhancing harvest yields, for improving the quality of the harvested material in agriculture, in horticulture, in animal husbandry, in forests, in gardens and
10 leisure facilities, in the protection of stored products and of materials, and in the hygiene sector. They can preferably be used as crop protection agents. They are effective against normally sensitive and resistant species and against all or some stages of development.

15

The treatment of the plants and plant parts with the active ingredients or compositions is effected directly or by action on their surroundings, habitat or storage space by the customary treatment methods, for example by dipping, spraying,
20 atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, watering (drenching), drip irrigating and, in the case of propagation material, especially in the case of seeds, also by dry seed treatment, wet seed treatment, slurry treatment, incrustation,
25 coating with one or more coats, etc. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject the active ingredient preparation/the active ingredient itself into the soil.

30 In addition, in the protection of materials, the active ingredients or compositions prepared with the aid of the inventive compounds can be used for protection of industrial materials against attack and destruction by unwanted microorganisms, for example fungi.

35

Industrial materials in the present context are understood to mean inanimate materials which have been prepared for use in industry. For example, industrial materials which are to be

protected by active ingredients prepared with the aid of the inventive compounds from microbial alteration or destruction may be adhesives, sizes, paper and board, textiles, leather, wood, paints and plastic articles, cooling lubricants and other materials which can be infected with or destroyed by microorganisms. The range of materials to be protected also includes parts of production plants, for example cooling water circuits, which may be impaired by the proliferation of microorganisms. Industrial materials within the scope of the present invention preferably include adhesives, sizes, paper and cardboard, leather, wood, paints, cooling lubricants and heat transfer fluids, more preferably wood. The inventive active ingredients or compositions may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

Microorganisms capable of degrading or altering the industrial materials include, for example, bacteria, fungi, yeasts, algae and slime organisms. The active ingredients preferably act against fungi, especially moulds, wood-discoloring and wood-destroying fungi (Basidiomycetes), and against slime organisms and algae. Examples include microorganisms of the following genera: *Alternaria*, such as *Alternaria tenuis*; *Aspergillus*, such as *Aspergillus niger*; *Chaetomium*, such as *Chaetomium globosum*; *Coniophora*, such as *Coniophora puetana*; *Lentinus*, such as *Lentinus tigrinus*; *Penicillium*, such as *Penicillium glaucum*; *Polyporus*, such as *Polyporus versicolor*; *Aureobasidium*, such as *Aureobasidium pullulans*; *Sclerophoma*, such as *Sclerophoma pityophila*; *Trichoderma*, such as *Trichoderma viride*; *Escherichia*, such as *Escherichia coli*; *Pseudomonas*, such as *Pseudomonas aeruginosa*; *Staphylococcus*, such as *Staphylococcus aureus*.

The active ingredients prepared with the aid of the inventive compounds can also be used, as such or in formulations thereof, in a mixture with known fungicides, bactericides, acaricides, nematocides or insecticides, in order thus to broaden, for example, the activity spectrum or to prevent

development of resistance.

Useful mixing partners include, for example, known fungicides,
insecticides, acaricides, nematicides or else bactericides
5 (see also Pesticide Manual, 14th ed.).

A mixture with other known active ingredients, such as
herbicides, or with fertilizers and growth regulators,
safeners and/or semiochemicals, is also possible.

10

Application is accomplished in a customary manner appropriate
for the use forms.

The active ingredients or compositions prepared with the aid
15 of the inventive compounds are also suitable for the treatment
of seed. A large part of the damage to crop plants caused by
harmful organisms is triggered by the infection of the seed
during storage or after sowing both during and after
germination of the plant. This phase is particularly critical
20 since the roots and shoots of the growing plant are
particularly sensitive, and even small damage may result in
the death of the plant. Accordingly, there is great interest
in protecting the seed and the germinating plant by using
appropriate compositions.

25

The control of phytopathogenic harmful fungi by treating the
seed of plants has been known for a long time and is the
subject of constant improvements. However, the treatment of
seed entails a series of problems which cannot always be
30 solved in a satisfactory manner. For instance, it is desirable
to develop methods for protecting the seed and the germinating
plant, which dispense with, or at least significantly reduce,
the additional deployment of crop protection compositions
after planting or after emergence of the plants. It is also
35 desirable to optimize the amount of the active ingredient used
so as to provide the best possible protection for the seed and
the germinating plant from attack by phytopathogenic fungi,
but without damaging the plant itself by the active ingredient

used. In particular, methods for the treatment of seed should also take into consideration the intrinsic fungicidal properties of transgenic plants in order to achieve optimum protection of the seed and the germinating plant with a minimum of crop protection compositions being employed.

Animal pests and/or phytopathogenic harmful fungi which damage plants post-emergence are controlled primarily by the treatment of the soil and of the exposed plant parts with crop protection compositions. Owing to the concerns regarding a possible influence of the crop protection compositions on the environment and the health of humans and animals, there are efforts to reduce the amount of active ingredients deployed.

One of the advantages of the present invention is that, because of the particular systemic properties of the compositions prepared with the aid of the inventive compounds, the treatment of the seed with these compositions not only protects the seed itself, but also the resulting plants after emergence, from animal pests and/or phytopathogenic harmful fungi. In this way, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

It is likewise considered to be advantageous that the active ingredients or compositions prepared with the aid of the inventive compounds can be used especially also for transgenic seed, in which case the plant which grows from this seed is capable of expressing a protein which acts against pests. The treatment of such seed with the active ingredients or compositions prepared with the aid of the inventive compounds, merely through the expression of the protein, for example an insecticidal protein, can result in control of certain pests. Surprisingly, a further synergistic effect can be observed in this case, which additionally increases the effectiveness for protection against attack by pests.

The compositions prepared with the aid of the inventive

compounds are suitable for protection of seed of any plant variety which is used in agriculture, in the greenhouse, in forests or in horticulture. More particularly, the seed is that of cereals (such as wheat, barley, rye, millet and oats),
5 maize, cotton, soya, rice, potatoes, sunflower, bean, coffee, beet (e.g. sugar beet and fodder beet), peanut, vegetables (such as tomato, cucumber, onions and lettuce), lawns and ornamental plants. Of particular significance is the treatment of the seed of cereals (such as wheat, barley, rye and oats),
10 maize and rice.

As also described below, the treatment of transgenic seed with the active ingredients or compositions prepared with the aid of the inventive compounds is of particular significance. This
15 refers to the seed of plants containing at least one heterologous gene which allows the expression of a polypeptide or protein having insecticidal properties. The heterologous gene in transgenic seed may originate, for example, from microorganisms of the species *Bacillus*, *Rhizobium*,
20 *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. This heterologous gene preferably originates from *Bacillus* sp., in which case the gene product is effective against the European corn borer and/or the Western corn rootworm. Particularly preferably, the heterologous gene
25 originates from *Bacillus thuringiensis*.

The composition prepared with the aid of the inventive compounds can be applied to the seed either alone or in a suitable formulation. Preferably, the seed is treated in a
30 state in which it is sufficiently stable for no damage to occur in the course of treatment. In general, the seed can be treated at any time between harvest and sowing. It is customary to use seed which has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh
35 of the fruits. For example, it is possible to use seed which has been harvested, cleaned and dried down to a moisture content of less than 15% by weight. Alternatively, it is also possible to use seed which, after drying, for example, has

been treated with water and then dried again.

When treating the seed, it generally has to be ensured that the amount of the composition prepared with the aid of the inventive compounds applied to the seed and/or the amount of further additives is selected such that the germination of the seed is not impaired, or that the resulting plant is not damaged. This must be ensured particularly in the case of active ingredients which can exhibit phytotoxic effects at certain application rates.

The compositions prepared with the aid of the inventive compounds can be applied directly, i.e. without containing any other components and without having been diluted. In general, it is preferable to apply the compositions to the seed in the form of a suitable formulation. Suitable formulations and methods for seed treatment are known to those skilled in the art and are described, for example, in the following documents: US 4,272,417, US 4,245,432, US 4,808,430, US 5,876,739, US 2003/0176428, WO 2002/080675, WO 2002/028186.

The active ingredients prepared with the aid of the inventive compounds can be converted to the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating compositions for seed, and also ULV formulations.

These formulations are prepared in a known manner, by mixing the active ingredients or active ingredient combinations with customary additives, for example customary extenders and solvents or diluents, dyes, wetting agents, dispersants, emulsifiers, antifoams, preservatives, secondary thickeners, adhesives, gibberellins, and also water.

Useful dyes which may be present in the seed dressing formulations are all dyes which are customary for such purposes. It is possible to use either pigments, which are sparingly soluble in water, or dyes, which are soluble in

water. Examples include the dyes known by the names Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

5 Useful wetting agents which may be present in the seed dressing formulations are all substances which promote wetting and which are conventionally used for the formulation of active agrochemical ingredients. Usable with preference are alkylnaphthalenesulphonates, such as diisopropyl- or diisobutyl-naphthalenesulphonates.

10

Useful dispersants and/or emulsifiers which may be present in the seed dressing formulations are all nonionic, anionic and cationic dispersants conventionally used for the formulation of active agrochemical ingredients. Usable with preference are
15 nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Useful nonionic dispersants include especially ethylene oxide/propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ether, and the phosphated or sulphated derivatives thereof.
20 Suitable anionic dispersants are especially lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.

Antifoams which may be present in the seed dressing
25 formulations are all foam-inhibiting substances conventionally used for the formulation of active agrochemical ingredients. Silicone antifoams and magnesium stearate can be used with preference.

30 Preservatives which may be present in the seed dressing formulations are all substances usable for such purposes in agrochemical compositions. Examples include dichlorophene and benzyl alcohol hemiformal.

35 Secondary thickeners which may be present in the seed dressing formulations are all substances usable for such purposes in agrochemical compositions. Preferred examples include cellulose derivatives, acrylic acid derivatives, xanthan,

modified clays and finely divided silica.

Adhesives which may be present in the seed dressing formulations are all customary binders usable in seed dressing products. Preferred examples include polyvinylpyrrolidone, 5 polyvinyl acetate, polyvinyl alcohol and tylose.

The gibberellins which may be present in the seed dressing formulations may preferably be gibberellins A1, A3 10 (= gibberellic acid), A4 and A7; particular preference is given to using gibberellic acid. The gibberellins are known (cf. R. Wegler "Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel" [Chemistry of Crop Protection Agents and Pesticides], vol. 2, Springer Verlag, 1970, p. 401- 15 412).

The seed dressing formulations can be used to treat a wide variety of different kinds of seed either directly or after prior dilution with water. For instance, the concentrates or 20 the preparations obtainable therefrom by dilution with water can be used to dress the seed of cereals, such as wheat, barley, rye, oats, and triticale, and also the seed of maize, rice, oilseed rape, peas, beans, cotton, sunflowers, and beets, or else a wide variety of different vegetable seed. The 25 seed dressing formulations, or the dilute preparations thereof, can also be used to dress seed of transgenic plants. In this case, additional synergistic effects may also occur in interaction with the substances formed by expression.

30 For treatment of seed with the seed dressing formulations, or the preparations prepared therefrom by adding water, all mixing units usable customarily for the seed dressing are useful. Specifically, the procedure in the seed dressing is to place the seed into a mixer, to add the particular desired 35 amount of seed dressing formulations, either as such or after prior dilution with water, and to mix everything until the formulation is distributed homogeneously on the seed. If appropriate, this is followed by a drying operation.

The application rate of the seed dressing formulations can be varied within a relatively wide range. It is guided by the particular content of the active ingredients in the formulations and by the seed. The application rates of active ingredient combination are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 15 g per kilogram of seed.

10 In addition, the compounds of the formula **(I)** prepared with the aid of the inventive compounds also have very good antimycotic effects. They have a very broad antimycotic activity spectrum, especially against dermatophytes and yeasts, moulds and diphasic fungi (for example against *Candida* species, such as *Candida albicans*, *Candida glabrata*), and *Epidermophyton floccosum*, *Aspergillus* species, such as *Aspergillus niger* and *Aspergillus fumigatus*, *Trichophyton* species, such as *Trichophyton mentagrophytes*, *Microsporon* species such as *Microsporon canis* and *audouinii*. The
15 enumeration of these fungi by no means constitutes a restriction of the mycotic spectrum covered, and is merely of illustrative character.

The active ingredients of the formula **(I)** can therefore be
25 used both in medical and in non-medical applications.

The active ingredients can be used as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, suspensions, wettable powders, pastes, soluble powders, dusts and granules. Application is
30 accomplished in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading-on and the like. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject
35 the active ingredient preparation/the active ingredient itself into the soil. It is also possible to treat the seed of the plants.

When using the active ingredients as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. The application rate of the active ingredients is

5

- in the case of treatment of plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 50 to 300 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates such as rockwool or perlite are used);

10

- in the case of seed treatment: from 2 to 200 g per 100 kg of seed, preferably from 3 to 150 g per 100 kg of seed, more preferably from 2.5 to 25 g per 100 kg of seed, even more preferably from 2.5 to 12.5 g per 100 kg of seed;

15

- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

20

These application rates are merely by way of example and are not limiting for the purposes of the invention.

25

The active ingredients are used in the veterinary sector and in animal husbandry in a known manner, by enteral administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boluses, the feed-through process and suppositories, by parenteral administration, for example by injection (intramuscular, subcutaneous, intravenous, intraperitoneal and the like), implants, by nasal administration, by dermal use in the form, for example, of dipping or bathing, spraying, pouring on and spotting on, washing and powdering, and also with the aid of molded articles containing the active ingredient, such as collars, ear marks, tail marks, limb bands, halters, marking devices and the like.

30

35

When used for livestock, poultry, domestic animals and the

like, the active ingredients of the formula (I) can be used as formulations (for example powders, emulsions, flowables) comprising the active ingredients in an amount of 1 to 80% by weight, either directly or after 100 to 10 000-fold dilution, 5 or they may be used as a chemical bath.

The ready-to-use compositions may optionally also comprise other insecticides, and optionally also one or more fungicides.

10

With respect to possible additional mixing partners, reference is made to the insecticides and fungicides mentioned above.

At the same time, the compounds of the formula (I) prepared 15 with the aid of the inventive compounds can be used for protection of objects which come into contact with saltwater or brackish water, especially hulls, screens, nets, buildings, moorings and signalling systems, against fouling.

20 In addition, the compounds of the formula (I) can be used as anti-fouling compositions, alone or in combinations with other active ingredients.

At certain application rates, the active ingredient 25 combinations may also have a fortifying effect on plants. Accordingly, they are suitable for mobilizing the defence system of the plant against attack by unwanted phytopathogenic fungi and/or microorganisms and/or viruses. This may, if appropriate, be one of the reasons for the enhanced activity 30 of the combinations, for example against fungi. Plant-fortifying (resistance-inducing) substances shall be understood to mean, in the present context, also those substances or combinations of substances which are capable of stimulating the defence system of plants in such a way that, 35 when subsequently inoculated with unwanted phytopathogenic fungi and/or microorganisms and/or viruses, the plants treated display a substantial degree of resistance to these unwanted phytopathogenic fungi and/or microorganisms and/or viruses. In

the present case, unwanted phytopathogenic fungi and/or microorganisms and/or viruses are understood to mean phytopathogenic fungi, bacteria and viruses. Thus, the substances prepared with the aid of the inventive compounds
5 **(XIII)** can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period within which protection is achieved generally extends for from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active
10 ingredients.

Plants and plant varieties which are preferably treated include all plants which have genetic material which imparts particularly advantageous, useful traits to these plants
15 (whether obtained by breeding and/or biotechnological means).

Plants and plant varieties which are also preferably treated are resistant against one or more biotic stress factors, i.e. said plants have a better defence against animal and microbial
20 pests, such as against nematodes, insects, mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

Plants and plant varieties which may also be treated are those plants which are resistant to one or more abiotic stress
25 factors. Abiotic stress conditions may include, for example, drought, cold and hot conditions, osmotic stress, waterlogging, increased soil salinity, increased exposure to minerals, ozone conditions, strong light conditions, limited availability of nitrogen nutrients, limited availability of
30 phosphorus nutrients or shade avoidance.

Plants and plant varieties which can likewise be treated are those plants which are characterized by increased yield
35 properties. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination

efficiency and accelerated maturation. Yield can also be affected by improved plant architecture (under stress and non-stress conditions), including early flowering, flowering control for hybrid seed production, seedling vigour, plant
5 size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as
10 carbohydrate content, protein content, oil content and composition, nutritional value, reduction in anti-nutritional compounds, improved processibility and better storage stability.

15 Plants that may be treated are hybrid plants that already express the characteristics of heterosis, or hybrid effect, which results in generally higher yield, vigour, health and resistance towards biotic and abiotic stress factors. Such plants are typically produced by crossing an inbred male-
20 sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). The hybrid seed is typically harvested from the male-sterile plants and sold to growers. Male-sterile plants can sometimes (e.g. in corn) be produced by detasseling (i.e. the mechanical removal of the
25 male reproductive organs or male flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants, it is typically beneficial to ensure that male fertility in hybrid
30 plants, which contain the genetic determinants responsible for male sterility, is fully restored. This can be accomplished by ensuring that the male parents have appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic
35 determinants responsible for male sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described for Brassica species. However, genetic

determinants for male sterility can also be located in the nuclear genome. Male-sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 89/10396 in which, for example, a ribonuclease such as a barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar.

10

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

15

Herbicide-tolerant plants are, for example, glyphosate-tolerant plants, i.e. plants which have been made tolerant to the herbicide glyphosate or salts thereof. For example, glyphosate-tolerant plants can be obtained by transforming the plant with a gene which encodes the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are the AroA gene (mutant CT7) of the bacterium *Salmonella typhimurium*, the CP4 gene of the bacterium *Agrobacterium sp.*, the genes encoding a petunia EPSPS, a tomato EPSPS, or an Eleusine EPSPS. It can also be a mutated EPSPS. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate oxidoreductase enzyme. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate acetyl transferase enzyme. Glyphosate-tolerant plants can also be obtained by selecting plants with naturally-occurring mutations of the above-mentioned genes.

20

25

30

35

Other herbicide-resistant plants are for example plants that have been made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or

glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition. One such efficient detoxifying enzyme is, for example, an enzyme
5 encoding a phosphinothricin acetyltransferase (for example the bar or pat protein from *Streptomyces* species). Plants expressing an exogenous phosphinothricin acetyltransferase have been described.

10 Further herbicide-tolerant plants are also plants that have been made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvate dioxygenase (HPPD). Hydroxyphenylpyruvate dioxygenases are enzymes that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is
15 converted to homogentisate. Plants tolerant to HPPD inhibitors can be transformed with a gene encoding a naturally occurring resistant HPPD enzyme, or a gene encoding a mutated HPPD enzyme. Tolerance to HPPD inhibitors can also be obtained by transforming plants with genes encoding certain enzymes
20 enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD inhibitor. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an enzyme prephenate dehydrogenase in addition to a gene encoding an HPPD-tolerant enzyme.

25 Further herbicide-resistant plants are plants that have been made tolerant to acetolactate synthase (ALS) inhibitors. The known ALS inhibitors include, for example, sulphonylurea, imidazolinone, triazolopyrimidines, pyrimidinyl
30 oxy(thio)benzoates and/or sulphonylaminocarbonyltriazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxy acid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides. The production of sulphonylurea-tolerant plants and
35 imidazolinone-tolerant plants has been described in the international publication WO 1996/033270. Further sulphonylurea- and imidazolinone-tolerant plants have also been described, for example in WO 2007/024782.

Other plants tolerant to imidazolinone and/or sulphonylurea can be obtained by induced mutagenesis, by selection in cell cultures in the presence of the herbicide or by mutation breeding.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which can also be treated are insect-resistant transgenic plants, i.e. plants which have been made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation which imparts such insect resistance.

In the present context, the term "insect-resistant transgenic plant" includes any plant containing at least one transgene comprising a coding sequence which encodes the following:

1) an insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins compiled online at: http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/, or insecticidal portions thereof, for example proteins of the Cry protein classes Cry1Ab, Cry1Ac, Cry1F, Cry2Ab, Cry3Ae or Cry3Bb or insecticidal portions thereof; or

2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein as *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cy34 and Cy35 crystal proteins; or

3) a hybrid insecticidal protein comprising parts of two different insecticidal crystal proteins from *Bacillus thuringiensis*, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2) above, for example the Cry1A.105 protein produced by maize event MON98034 (WO 2007/027777); or

- 4) a protein of any one of points 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes induced in the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in maize events MON863 or MON88017, or the Cry3A protein in maize event MIR604; or
- 5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal proteins (VIP) listed at: http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html, for example proteins from the VIP3Aa protein class; or
- 6) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B. cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins;
- 7) a hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the proteins in 2) above; or
- 8) a protein of any one of points 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes induced in the encoding DNA during cloning or transformation (while still encoding an insecticidal protein), such as the VIP3Aa protein in cotton event COT102.

Of course, insect-resistant transgenic plants, as used herein,

also include any plant comprising a combination of genes encoding the proteins of any one of the abovementioned classes 1 to 8. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the abovementioned classes 1 to 8, to expand the range of target insect species affected or to delay insect resistance development to the plants, by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated are tolerant to abiotic stress factors. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress-tolerant plants include the following:

a. plants which contain a transgene capable of reducing the expression and/or the activity of the poly(ADP-ribose)polymerase (PARP) gene in the plant cells or plants;

b. plants which contain a stress tolerance-enhancing transgene capable of reducing the expression and/or the activity of the PARG-encoding genes of the plants or plant cells;

c. plants which contain a stress tolerance-enhancing transgene coding for a plant-functional enzyme of the nicotinamide adenine dinucleotide salvage biosynthesis pathway, including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenyltransferase, nicotinamide adenine dinucleotide synthetase or nicotinamide phosphoribosyltransferase.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated

show altered quantity, quality and/or storage stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as:

5 1) Transgenic plants which synthesize a modified starch which is altered with respect to its chemophysical traits, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the distribution of the side chains, the viscosity behaviour, the
10 gel resistance, the grain size and/or grain morphology of the starch in comparison to the synthesized starch in wild-type plant cells or plants, such that this modified starch is better suited for certain applications.

15 2) Transgenic plants which synthesize non-starch carbohydrate polymers or which synthesize non-starch carbohydrate polymers with altered properties in comparison to wild-type plants without genetic modification. Examples are plants which produce polyfructose, especially of the inulin
20 and levan type, plants which produce alpha-1,4-glucans, plants which produce alpha-1,6-branched alpha-1,4-glucans, and plants producing alternan.

3) Transgenic plants which produce hyaluronan.
25

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated are plants, such as cotton plants, with altered fibre characteristics. Such plants can be obtained by genetic
30 transformation, or by selection of plants containing a mutation imparting such altered fibre characteristics, and include:

a) plants, such as cotton plants, which contain an altered
35 form of cellulose synthase genes;

b) plants, such as cotton plants, which contain an altered form of rsw2 or rsw3 homologous nucleic acids;


c) plants, such as cotton plants, with an increased expression of sucrose phosphate synthase;

5 d) plants, such as cotton plants, with an increased expression of sucrose synthase;


e) plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fibre cell is altered, for example through downregulation of fibre-selective
10 β -1,3-glucanase;

f) plants, such as cotton plants, which have fibres with altered reactivity, for example through the expression of the
15 N-acetylglucosaminetransferase gene including nodC and chitin synthase genes.


Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated
20 are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be:



(XXXIa)



(XXXIb)



(XXXIc)

25 The substances of the formula **(I)** prepared with the aid of the inventive compounds **(XIII)** have potent microbicidal activity and can be used for control of unwanted microorganisms, such as fungi and bacteria, in crop protection and in the protection of materials.

30

The heteroarylpiperidine and -piperazine derivatives of the formula **(I)** have very good fungicidal properties and can be used in crop protection, for example for control of Plasmodiophoromycetes, Oomycetes, Chytridiomycetes,
35 Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

Bactericides can be used in crop protection, for example, for control of Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomycetaceae.

5 diseases caused by powdery mildew pathogens, for example Blumeria species, for example Blumeria graminis; Podosphaera species, for example Podosphaera leucotricha; Sphaerotheca species, for example Sphaerotheca fuliginea; Uncinula species, for example Uncinula necator;

10

diseases caused by rust disease pathogens, for example Gymnosporangium species, for example Gymnosporangium sabinae; Hemileia species, for example Hemileia vastatrix; Phakopsora species, for example Phakopsora pachyrhizi or Phakopsora
15 meibomiae; Puccinia species, for example Puccinia recondita, Puccinia graminis oder Puccinia striiformis; Uromyces species, for example Uromyces appendiculatus;

diseases caused by pathogens from the group of the Oomycetes,
20 for example Albugo species, for example Albugo candida; Bremia species, for example Bremia lactucae; Peronospora species, for example Peronospora pisi or P. brassicae; Phytophthora species, for example Phytophthora infestans; Plasmopara species, for example Plasmopara viticola; Pseudoperonospora
25 species, for example Pseudoperonospora humuli or Pseudoperonospora cubensis; Pythium species, for example Pythium ultimum;

leaf blotch diseases and leaf wilt diseases caused, for
30 example, by Alternaria species, for example Alternaria solani; Cercospora species, for example Cercospora beticola; Cladosporium species, for example Cladosporium cucumerinum; Cochliobolus species, for example Cochliobolus sativus (conidial form: Drechslera, syn: Helminthosporium) or
35 Cochliobolus miyabeanus; Colletotrichum species, for example Colletotrichum lindemuthanium; Cycloconium species, for example Cycloconium oleaginum; Diaporthe species, for example Diaporthe citri; Elsinoe species, for example Elsinoe

fawcettii; Gloeosporium species, for example Gloeosporium
laeticolor; Glomerella species, for example Glomerella
cingulata; Guignardia species, for example Guignardia
bidwelli; Leptosphaeria species, for example Leptosphaeria
5 maculans; Magnaporthe species, for example Magnaporthe grisea;
Microdochium species, for example Microdochium nivale;
Mycosphaerella species, for example Mycosphaerella
graminicola, Mycosphaerella arachidicola or Mycosphaerella
fijiensis; Phaeosphaeria species, for example Phaeosphaeria
10 nodorum; Pyrenophora species, for example Pyrenophora teres or
Pyrenophora tritici repentis; Ramularia species, for example
Ramularia collo-cygni or Ramularia areola; Rhynchosporium
species, for example Rhynchosporium secalis; Septoria species,
for example Septoria apii or Septoria lycopersici;
15 Stagonospora species, for example Stagonospora nodorum;
Typhula species, for example Typhula incarnata; Venturia
species, for example Venturia inaequalis;

root and stem diseases caused, for example, by Corticium
20 species, for example Corticium graminearum; Fusarium species,
for example Fusarium oxysporum; Gaeumannomyces species, for
example Gaeumannomyces graminis; Plasmodiophora species, for
example Plasmodiophora brassicae; Rhizoctonia species, for
example Rhizoctonia solani; Sarocladium species, for example
25 Sarocladium oryzae; Sclerotium species, for example Sclerotium
oryzae; Tapesia species, for example Tapesia acuformis;
Thielaviopsis species, for example Thielaviopsis basicola;

ear and panicle diseases (including corn cobs) caused, for
30 example, by Alternaria species, for example Alternaria spp.;
Aspergillus species, for example Aspergillus flavus;
Cladosporium species, for example Cladosporium
cladosporioides; Claviceps species, for example Claviceps
purpurea; Fusarium species, for example Fusarium culmorum;
35 Gibberella species, for example Gibberella zeae; Monographella
species, for example Monographella nivalis; Stagnospora
species, for example Stagnospora nodorum;

diseases caused by smut fungi, for example *Sphacelotheca* species, for example *Sphacelotheca reiliana*; *Tilletia* species, for example *Tilletia caries* or *Tilletia controversa*; *Urocystis* species, for example *Urocystis occulta*; *Ustilago* species, for example *Ustilago nuda*;

fruit rot caused, for example, by *Aspergillus* species, for example *Aspergillus flavus*; *Botrytis* species, for example *Botrytis cinerea*; *Penicillium* species, for example *Penicillium expansum* or *Penicillium purpurogenum*; *Rhizopus* species, for example *Rhizopus stolonifer*; *Sclerotinia* species, for example *Sclerotinia sclerotiorum*; *Verticilium* species, for example *Verticilium alboatrum*;

seed- and soil-borne rot and wilt diseases, and also diseases of seedlings, caused, for example, by *Alternaria* species, for example *Alternaria brassicicola*; *Aphanomyces* species, for example *Aphanomyces euteiches*; *Ascochyta* species, for example *Ascochyta lentis*; *Aspergillus* species, for example *Aspergillus flavus*; *Cladosporium* species, for example *Cladosporium herbarum*; *Cochliobolus* species, for example *Cochliobolus sativus* (conidial form: *Drechslera*, *Bipolaris* Syn: *Helminthosporium*); *Colletotrichum* species, for example *Colletotrichum coccodes*; *Fusarium* species, for example *Fusarium culmorum*; *Gibberella* species, for example *Gibberella zeae*; *Macrophomina* species, for example *Macrophomina phaseolina*; *Microdochium* species, for example *Microdochium nivale*; *Monographella* species, for example *Monographella nivalis*; *Penicillium* species, for example *Penicillium expansum*; *Phoma* species, for example *Phoma lingam*; *Phomopsis* species, for example *Phomopsis sojae*; *Phytophthora* species, for example *Phytophthora cactorum*; *Pyrenophora* species, for example *Pyrenophora graminea*; *Pyricularia* species, for example *Pyricularia oryzae*; *Pythium* species, for example *Pythium ultimum*; *Rhizoctonia* species, for example *Rhizoctonia solani*; *Rhizopus* species, for example *Rhizopus oryzae*; *Sclerotium* species, for example *Sclerotium rolfsii*; *Septoria* species, for example *Septoria nodorum*; *Typhula* species, for example *Typhula*

incarnata; *Verticillium* species, for example *Verticillium dahliae*;

5 cancers, galls and witches' broom caused, for example, by *Nectria* species, for example *Nectria galligena*;

wilt diseases caused, for example, by *Monilinia* species, for example *Monilinia laxa*;

10 deformations of leaves, flowers and fruits caused, for example, by *Exobasidium* species, for example *Exobasidium vexans*; *Taphrina* species, for example *Taphrina deformans*;

15 degenerative diseases in woody plants, caused, for example, by *Esca* species, for example *Phaeomoniella chlamydospora*, *Phaeoacremonium aleophilum* or *Fomitiporia mediterranea*; *Ganoderma* species, for example *Ganoderma boninense*;

20 diseases of flowers and seeds caused, for example, by *Botrytis* species, for example *Botrytis cinerea*;

diseases of plant tubers caused, for example, by *Rhizoctonia* species, for example *Rhizoctonia solani*; *Helminthosporium* species, for example *Helminthosporium solani*;

25 diseases caused by bacterial pathogens, for example *Xanthomonas* species, for example *Xanthomonas campestris* pv. *oryzae*; *Pseudomonas* species, for example *Pseudomonas syringae* pv. *lachrymans*; *Erwinia* species, for example *Erwinia amylovora*.

Preference is given to controlling the following diseases of soya beans:

35 Fungal diseases on leaves, stems, pods and seeds caused, for example, by *Alternaria* leaf spot (*Alternaria spec. atrans tenuissima*), Anthracnose (*Colletotrichum gloeosporoides dematium* var. *truncatum*), brown spot (*Septoria glycines*),

cercospora leaf spot and blight (*Cercospora kikuchii*),
 choanephora leaf blight (*Choanephora infundibulifera trispora*
 (Syn.)), dactuliophora leaf spot (*Dactuliophora glycines*),
 downy mildew (*Peronospora manshurica*), drechslera blight
 5 (*Drechslera glycini*), frog-eye leaf spot (*Cercospora sojina*),
 leptosphaerulina leaf spot (*Leptosphaerulina trifolii*),
 phyllosticta leaf spot (*Phyllosticta sojaecola*), pod and stem
 blight (*Phomopsis sojae*), powdery mildew (*Microsphaera*
diffusa), pyrenochaeta leaf spot (*Pyrenochaeta glycines*),
 10 rhizoctonia aerial, foliage, and web blight (*Rhizoctonia*
solani), rust (*Phakopsora pachyrhizi*, *Phakopsora meibomia*),
 scab (*Sphaceloma glycines*), stemphylium leaf blight
 (*Stemphylium botryosum*), target spot (*Corynespora cassiicola*).

15 Fungal diseases on roots and the stem base caused, for
 example, by black root rot (*Calonectria crotalariae*), charcoal
 rot (*Macrophomina phaseolina*), fusarium blight or wilt, root
 rot, and pod and collar rot (*Fusarium oxysporum*, *Fusarium*
orthoceras, *Fusarium semitectum*, *Fusarium equiseti*),
 20 mycoleptodiscus root rot (*Mycoleptodiscus terrestris*),
 neocosmospora (*Neocosmospora vasinfecta*), pod
 and stem blight (*Diaporthe phaseolorum*), stem canker
 (*Diaporthe phaseolorum* var. *caulivora*), phytophthora rot
 25 (*Phytophthora megasperma*), brown stem rot (*Phialophora*
gregata), pythium rot (*Pythium aphanidermatum*, *Pythium*
irregulare, *Pythium debaryanum*, *Pythium myriotylum*, *Pythium*
ultimum), rhizoctonia root rot, stem decay, and damping-off
 (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia*
 30 *sclerotiorum*), sclerotinia southern blight (*Sclerotinia*
rolfsii), thielaviopsis root rot (*Thielaviopsis basicola*).

The active ingredients prepared with the aid of the inventive
 compounds also have very good fortifying action in plants.
 35 They are therefore suitable for mobilizing the plant's own
 defences against attack by undesirable microorganisms.

Plant-fortifying (resistance-inducing) substances are

understood to mean, in the present context, those substances which are capable of stimulating the defence system of plants in such a way that the treated plants, when subsequently inoculated with undesirable microorganisms, develop a high
5 degree of resistance to these microorganisms.

In the present case, undesirable microorganisms are understood to mean phytopathogenic fungi and bacteria. The substances prepared with the aid of the inventive compounds can thus be
10 used to protect plants for a certain period after the treatment against attack by the pathogens mentioned. The period for which protection is provided generally extends over 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active ingredients.

15 The fact that the active ingredients are well tolerated by plants at the concentrations required for controlling plant diseases allows the treatment of above-ground parts of plants, of propagation stock and seeds, and of the soil.

20 The active ingredients prepared with the aid of the inventive compounds can be used particularly successfully to control diseases in viticulture and potato, fruit and vegetable growing, for example against powdery mildew fungi, Oomycetes,
25 for example Phytophthora, Plasmopara, Pseudoperonospora and Pythium species.

The active ingredients prepared with the aid of the inventive compounds are also suitable for enhancing harvest yield. In
30 addition, they have low toxicity and are well tolerated by plants.

If appropriate, the compounds prepared with the aid of the inventive compounds can, at certain concentrations or
35 application rates, also be used as herbicides, safeners, growth regulators or agents to improve plant properties, or as microbicides, for example as fungicides, antimycotics, bactericides, viricides (including agents against viroids) or

as agents against MLO (mycoplasma-like organisms) and RLO (rickettsia-like organisms). If appropriate, they can also be used as insecticides. If appropriate, they can also be used as intermediates or precursors for the synthesis of other active ingredients.

The active ingredients prepared with the aid of the inventive compounds, when they are well tolerated by plants, have favourable homeotherm toxicity and are well tolerated by the environment, are suitable for protecting plants and plant organs, for enhancing harvest yields, for improving the quality of the harvested material in agriculture, in horticulture, in animal husbandry, in forests, in gardens and leisure facilities, in the protection of stored products and of materials, and in the hygiene sector. They can preferably be used as crop protection agents. They are effective against normally sensitive and resistant species and against all or some stages of development.

The treatment of the plants and plant parts with the active ingredients or compositions is effected directly or by action on their surroundings, habitat or storage space by the customary treatment methods, for example by dipping, spraying, atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, watering (drenching), drip irrigating and, in the case of propagation material, especially in the case of seeds, also by dry seed treatment, wet seed treatment, slurry treatment, incrustation, coating with one or more coats, etc. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject the active ingredient preparation/the active ingredient itself into the soil.

In addition, in the protection of materials, the active ingredients or compositions prepared with the aid of the inventive compounds can be used for protection of industrial materials against attack and destruction by unwanted microorganisms, for example fungi.

Industrial materials in the present context are understood to mean inanimate materials which have been prepared for use in industry. For example, industrial materials which are to be protected by active ingredients prepared with the aid of the inventive compounds from microbial alteration or destruction may be adhesives, sizes, paper and board, textiles, leather, wood, paints and plastic articles, cooling lubricants and other materials which can be infected with or destroyed by microorganisms. The range of materials to be protected also includes parts of production plants, for example cooling water circuits, which may be impaired by the proliferation of microorganisms. Industrial materials within the scope of the present invention preferably include adhesives, sizes, paper and cardboard, leather, wood, paints, cooling lubricants and heat transfer fluids, more preferably wood. The inventive active ingredients or compositions may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

Microorganisms capable of degrading or altering the industrial materials include, for example, bacteria, fungi, yeasts, algae and slime organisms. The active ingredients preferably act against fungi, especially moulds, wood-discoloring and wood-destroying fungi (Basidiomycetes), and against slime organisms and algae. Examples include microorganisms of the following genera: *Alternaria*, such as *Alternaria tenuis*; *Aspergillus*, such as *Aspergillus niger*; *Chaetomium*, such as *Chaetomium globosum*; *Coniophora*, such as *Coniophora puetana*; *Lentinus*, such as *Lentinus tigrinus*; *Penicillium*, such as *Penicillium glaucum*; *Polyporus*, such as *Polyporus versicolor*; *Aureobasidium*, such as *Aureobasidium pullulans*; *Sclerophoma*, such as *Sclerophoma pityophila*; *Trichoderma*, such as *Trichoderma viride*; *Escherichia*, such as *Escherichia coli*; *Pseudomonas*, such as *Pseudomonas aeruginosa*; *Staphylococcus*, such as *Staphylococcus aureus*.

The active ingredients prepared with the aid of the inventive

compounds can also be used, as such or in formulations thereof, in a mixture with known fungicides, bactericides, acaricides, nematocides or insecticides, in order thus to broaden, for example, the activity spectrum or to prevent
5 development of resistance.

Useful mixing partners include, for example, known fungicides, insecticides, acaricides, nematocides or else bactericides (see also Pesticide Manual, 14th ed.).
10

A mixture with other known active ingredients, such as herbicides, or with fertilizers and growth regulators, safeners and/or semiochemicals, is also possible.

15 Application is accomplished in a customary manner appropriate for the use forms.

The active ingredients or compositions prepared with the aid of the inventive compounds are also suitable for the treatment
20 of seed. A large part of the damage to crop plants caused by harmful organisms is triggered by the infection of the seed during storage or after sowing both during and after germination of the plant. This phase is particularly critical since the roots and shoots of the growing plant are
25 particularly sensitive, and even small damage may result in the death of the plant. Accordingly, there is great interest in protecting the seed and the germinating plant by using appropriate compositions.

30 The control of phytopathogenic harmful fungi by treating the seed of plants has been known for a long time and is the subject of constant improvements. However, the treatment of seed entails a series of problems which cannot always be solved in a satisfactory manner. For instance, it is desirable
35 to develop methods for protecting the seed and the germinating plant, which dispense with, or at least significantly reduce, the additional deployment of crop protection compositions after planting or after emergence of the plants. It is also

desirable to optimize the amount of the active ingredient used so as to provide the best possible protection for the seed and the germinating plant from attack by phytopathogenic fungi, but without damaging the plant itself by the active ingredient used. In particular, methods for the treatment of seed should also take into consideration the intrinsic fungicidal properties of transgenic plants in order to achieve optimum protection of the seed and the germinating plant with a minimum of crop protection compositions being employed.

10

Animal pests and/or phytopathogenic harmful fungi which damage plants post-emergence are controlled primarily by the treatment of the soil and of the exposed plant parts with crop protection compositions. Owing to the concerns regarding a possible influence of the crop protection compositions on the environment and the health of humans and animals, there are efforts to reduce the amount of active ingredients deployed.

15

One of the advantages of the present invention is that, because of the particular systemic properties of the compositions prepared with the aid of the inventive compounds, the treatment of the seed with these compositions not only protects the seed itself, but also the resulting plants after emergence, from animal pests and/or phytopathogenic harmful fungi. In this way, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

20

25

It is likewise considered to be advantageous that the active ingredients or compositions prepared with the aid of the inventive compounds can be used especially also for transgenic seed, in which case the plant which grows from this seed is capable of expressing a protein which acts against pests. The treatment of such seed with the active ingredients or compositions prepared with the aid of the inventive compounds, merely through the expression of the protein, for example an insecticidal protein, can result in control of certain pests. Surprisingly, a further synergistic effect can be observed in

30

35

this case, which additionally increases the effectiveness for protection against attack by pests.

5 The compositions prepared with the aid of the inventive compounds are suitable for protection of seed of any plant variety which is used in agriculture, in the greenhouse, in forests or in horticulture. More particularly, the seed is that of cereals (such as wheat, barley, rye, millet and oats), maize, cotton, soya, rice, potatoes, sunflower, bean, coffee,
10 beet (e.g. sugar beet and fodder beet), peanut, vegetables (such as tomato, cucumber, onions and lettuce), lawns and ornamental plants. Of particular significance is the treatment of the seed of cereals (such as wheat, barley, rye and oats), maize and rice.

15

As also described below, the treatment of transgenic seed with the active ingredients or compositions prepared with the aid of the inventive compounds is of particular significance. This refers to the seed of plants containing at least one
20 heterologous gene which allows the expression of a polypeptide or protein having insecticidal properties. The heterologous gene in transgenic seed may originate, for example, from microorganisms of the species Bacillus, Rhizobium, Pseudomonas, Serratia, Trichoderma, Clavibacter, Glomus or
25 Gliocladium. This heterologous gene preferably originates from Bacillus sp., in which case the gene product is effective against the European corn borer and/or the Western corn rootworm. Particularly preferably, the heterologous gene originates from Bacillus thuringiensis.

30

The composition prepared with the aid of the inventive compounds can be applied to the seed either alone or in a suitable formulation. Preferably, the seed is treated in a state in which it is sufficiently stable for no damage to
35 occur in the course of treatment. In general, the seed can be treated at any time between harvest and sowing. It is customary to use seed which has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh

of the fruits. For example, it is possible to use seed which has been harvested, cleaned and dried down to a moisture content of less than 15% by weight. Alternatively, it is also possible to use seed which, after drying, for example, has been treated with water and then dried again.

When treating the seed, it generally has to be ensured that the amount of the composition prepared with the aid of the inventive compounds applied to the seed and/or the amount of further additives is selected such that the germination of the seed is not impaired, or that the resulting plant is not damaged. This must be ensured particularly in the case of active ingredients which can exhibit phytotoxic effects at certain application rates.

The compositions prepared with the aid of the inventive compounds can be applied directly, i.e. without containing any other components and without having been diluted. In general, it is preferable to apply the compositions to the seed in the form of a suitable formulation. Suitable formulations and methods for seed treatment are known to those skilled in the art and are described, for example, in the following documents: US 4,272,417, US 4,245,432, US 4,808,430, US 5,876,739, US 2003/0176428, WO 2002/080675, WO 2002/028186.

The active ingredients prepared with the aid of the inventive compounds can be converted to the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating compositions for seed, and also ULV formulations.

These formulations are prepared in a known manner, by mixing the active ingredients or active ingredient combinations with customary additives, for example customary extenders and solvents or diluents, dyes, wetting agents, dispersants, emulsifiers, antifoams, preservatives, secondary thickeners, adhesives, gibberellins, and also water.

Useful dyes which may be present in the seed dressing formulations are all dyes which are customary for such purposes. It is possible to use either pigments, which are sparingly soluble in water, or dyes, which are soluble in water. Examples include the dyes known by the names Rhodamine B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

Useful wetting agents which may be present in the seed dressing formulations are all substances which promote wetting and which are conventionally used for the formulation of active agrochemical ingredients. Usable with preference are alkylnaphthalenesulphonates, such as diisopropyl- or diisobutyl-naphthalenesulphonates.

Useful dispersants and/or emulsifiers which may be present in the seed dressing formulations are all nonionic, anionic and cationic dispersants conventionally used for the formulation of active agrochemical ingredients. Usable with preference are nonionic or anionic dispersants or mixtures of nonionic or anionic dispersants. Useful nonionic dispersants include especially ethylene oxide/propylene oxide block polymers, alkylphenol polyglycol ethers and tristyrylphenol polyglycol ether, and the phosphated or sulphated derivatives thereof. Suitable anionic dispersants are especially lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.

Antifoams which may be present in the seed dressing formulations are all foam-inhibiting substances conventionally used for the formulation of active agrochemical ingredients. Silicone antifoams and magnesium stearate can be used with preference.

Preservatives which may be present in the seed dressing formulations are all substances usable for such purposes in agrochemical compositions. Examples include dichlorophene and benzyl alcohol hemiformal.

Secondary thickeners which may be present in the seed dressing formulations are all substances usable for such purposes in agrochemical compositions. Preferred examples include cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica.

Adhesives which may be present in the seed dressing formulations are all customary binders usable in seed dressing products. Preferred examples include polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose.

The gibberellins which may be present in the seed dressing formulations may preferably be gibberellins A1, A3 (= gibberellic acid), A4 and A7; particular preference is given to using gibberellic acid. The gibberellins are known (cf. R. Wegler "Chemie der Pflanzenschutz- und Schädlingsbekämpfungsmittel" [Chemistry of Crop Protection Agents and Pesticides], vol. 2, Springer Verlag, 1970, p. 401-412).

The seed dressing formulations can be used to treat a wide variety of different kinds of seed either directly or after prior dilution with water. For instance, the concentrates or the preparations obtainable therefrom by dilution with water can be used to dress the seed of cereals, such as wheat, barley, rye, oats, and triticale, and also the seed of maize, rice, oilseed rape, peas, beans, cotton, sunflowers, and beets, or else a wide variety of different vegetable seed. The seed dressing formulations, or the dilute preparations thereof, can also be used to dress seed of transgenic plants. In this case, additional synergistic effects may also occur in interaction with the substances formed by expression.

For treatment of seed with the seed dressing formulations, or the preparations prepared therefrom by adding water, all mixing units usable customarily for the seed dressing are useful. Specifically, the procedure in the seed dressing is to place the seed into a mixer, to add the particular desired

amount of seed dressing formulations, either as such or after prior dilution with water, and to mix everything until the formulation is distributed homogeneously on the seed. If appropriate, this is followed by a drying operation.

5

The application rate of the seed dressing formulations can be varied within a relatively wide range. It is guided by the particular content of the active ingredients in the formulations and by the seed. The application rates of active ingredient combination are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 15 g per kilogram of seed.

In addition, the compounds of the formula **(I)** prepared with the aid of the inventive compounds also have very good antimycotic effects. They have a very broad antimycotic activity spectrum, especially against dermatophytes and yeasts, moulds and diphasic fungi (for example against *Candida* species, such as *Candida albicans*, *Candida glabrata*), and *Epidermophyton floccosum*, *Aspergillus* species, such as *Aspergillus niger* and *Aspergillus fumigatus*, *Trichophyton* species, such as *Trichophyton mentagrophytes*, *Microsporon* species such as *Microsporon canis* and *audouinii*. The enumeration of these fungi by no means constitutes a restriction of the mycotic spectrum covered, and is merely of illustrative character.

The active ingredients of the formula **(I)** can therefore be used both in medical and in non-medical applications.

30

The active ingredients can be used as such, in the form of their formulations or the use forms prepared therefrom, such as ready-to-use solutions, suspensions, wettable powders, pastes, soluble powders, dusts and granules. Application is accomplished in a customary manner, for example by watering, spraying, atomizing, broadcasting, dusting, foaming, spreading-on and the like. It is also possible to deploy the active ingredients by the ultra-low volume method or to inject

35

the active ingredient preparation/the active ingredient itself into the soil. It is also possible to treat the seed of the plants.

- 5 When using the active ingredients as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. The application rate of the active ingredients is
- 10 • in the case of treatment of plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 50 to 300 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates
- 15 such as rockwool or perlite are used);
- in the case of seed treatment: from 2 to 200 g per 100 kg of seed, preferably from 3 to 150 g per 100 kg of seed, more preferably from 2.5 to 25 g per 100 kg of seed, even more
- 20 preferably from 2.5 to 12.5 g per 100 kg of seed;
- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

25 These application rates are merely by way of example and are not limiting for the purposes of the invention.

The active ingredients are used in the veterinary sector and in animal husbandry in a known manner, by enteral

30 administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boluses, the feed-through process and suppositories, by parenteral administration, for example by injection (intramuscular, subcutaneous, intravenous, intraperitoneal and the like), implants, by nasal

35 administration, by dermal use in the form, for example, of dipping or bathing, spraying, pouring on and spotting on, washing and powdering, and also with the aid of molded articles containing the active ingredient, such as collars,

ear marks, tail marks, limb bands, halters, marking devices and the like.

5 When used for livestock, poultry, domestic animals and the like, the active ingredients of the formula (I) can be used as formulations (for example powders, emulsions, flowables) comprising the active ingredients in an amount of 1 to 80% by weight, either directly or after 100 to 10 000-fold dilution, or they may be used as a chemical bath.

10

The ready-to-use compositions may optionally also comprise other insecticides, and optionally also one or more fungicides.

15 With respect to possible additional mixing partners, reference is made to the insecticides and fungicides mentioned above.

20 At the same time, the compounds of the formula (I) prepared with the aid of the inventive compounds can be used for protection of objects which come into contact with saltwater or brackish water, especially hulls, screens, nets, buildings, moorings and signalling systems, against fouling.

25 In addition, the compounds of the formula (I) can be used as anti-fouling compositions, alone or in combinations with other active ingredients.

30 At certain application rates, the active ingredient combinations may also have a fortifying effect on plants. Accordingly, they are suitable for mobilizing the defence system of the plant against attack by unwanted phytopathogenic fungi and/or microorganisms and/or viruses. This may, if appropriate, be one of the reasons for the enhanced activity of the combinations, for example against fungi. Plant-
35 fortifying (resistance-inducing) substances shall be understood to mean, in the present context, also those substances or combinations of substances which are capable of stimulating the defence system of plants in such a way that,

when subsequently inoculated with unwanted phytopathogenic fungi and/or microorganisms and/or viruses, the plants treated display a substantial degree of resistance to these unwanted phytopathogenic fungi and/or microorganisms and/or viruses. In
5 the present case, unwanted phytopathogenic fungi and/or microorganisms and/or viruses are understood to mean phytopathogenic fungi, bacteria and viruses. Thus, the substances prepared with the aid of the inventive compounds **(XIII)** can be employed for protecting plants against attack by
10 the abovementioned pathogens within a certain period of time after the treatment. The period within which protection is achieved generally extends for from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active ingredients.

15 Plants and plant varieties which are preferably treated include all plants which have genetic material which imparts particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

20 Plants and plant varieties which are also preferably treated are resistant against one or more biotic stress factors, i.e. said plants have a better defence against animal and microbial pests, such as against nematodes, insects, mites,
25 phytopathogenic fungi, bacteria, viruses and/or viroids.

Plants and plant varieties which may also be treated are those plants which are resistant to one or more abiotic stress factors. Abiotic stress conditions may include, for example,
30 drought, cold and hot conditions, osmotic stress, waterlogging, increased soil salinity, increased exposure to minerals, ozone conditions, strong light conditions, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients or shade avoidance.

35 Plants and plant varieties which can likewise be treated are those plants which are characterized by increased yield properties. Increased yield in said plants can be the result

of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can also be affected by improved plant architecture (under stress and non-stress conditions), including early flowering, flowering control for hybrid seed production, seedling vigour, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content, protein content, oil content and composition, nutritional value, reduction in anti-nutritional compounds, improved processibility and better storage stability.

Plants that may be treated are hybrid plants that already express the characteristics of heterosis, or hybrid effect, which results in generally higher yield, vigour, health and resistance towards biotic and abiotic stress factors. Such plants are typically produced by crossing an inbred male-sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). The hybrid seed is typically harvested from the male-sterile plants and sold to growers. Male-sterile plants can sometimes (e.g. in corn) be produced by detasseling (i.e. the mechanical removal of the male reproductive organs or male flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants, it is typically beneficial to ensure that male fertility in hybrid plants, which contain the genetic determinants responsible for male sterility, is fully restored. This can be accomplished by ensuring that the male parents have appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic

determinants responsible for male sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described for Brassica species. However, genetic
5 determinants for male sterility can also be located in the nuclear genome. Male-sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 89/10396 in which, for example, a ribonuclease
10 such as a barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar.

15 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a
20 mutation imparting such herbicide tolerance.

Herbicide-tolerant plants are, for example, glyphosate-tolerant plants, i.e. plants which have been made tolerant to the herbicide glyphosate or salts thereof. For example,
25 glyphosate-tolerant plants can be obtained by transforming the plant with a gene which encodes the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are the AroA gene (mutant CT7) of the bacterium *Salmonella typhimurium*, the CP4 gene of the
30 bacterium *Agrobacterium sp.*, the genes encoding a petunia EPSPS, a tomato EPSPS, or an Eleusine EPSPS. It can also be a mutated EPSPS. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate oxidoreductase enzyme. Glyphosate-tolerant plants can also be obtained by
35 expressing a gene that encodes a glyphosate acetyl transferase enzyme. Glyphosate-tolerant plants can also be obtained by selecting plants with naturally-occurring mutations of the above-mentioned genes.

Other herbicide-resistant plants are for example plants that have been made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition. One such efficient detoxifying enzyme is, for example, an enzyme encoding a phosphinothricin acetyltransferase (for example the bar or pat protein from Streptomyces species). Plants expressing an exogenous phosphinothricin acetyltransferase have been described.

Further herbicide-tolerant plants are also plants that have been made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvate dioxygenase (HPPD). Hydroxyphenylpyruvate dioxygenases are enzymes that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is converted to homogentisate. Plants tolerant to HPPD inhibitors can be transformed with a gene encoding a naturally occurring resistant HPPD enzyme, or a gene encoding a mutated HPPD enzyme. Tolerance to HPPD inhibitors can also be obtained by transforming plants with genes encoding certain enzymes enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD inhibitor. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an enzyme prephenate dehydrogenase in addition to a gene encoding an HPPD-tolerant enzyme.

Further herbicide-resistant plants are plants that have been made tolerant to acetolactate synthase (ALS) inhibitors. The known ALS inhibitors include, for example, sulphonylurea, imidazolinone, triazolopyrimidines, pyrimidinyl oxy(thio)benzoates and/or sulphonylaminocarbonyl triazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxy acid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides. The production of sulphonylurea-tolerant plants and

imidazolinone-tolerant plants has been described in the international publication WO 1996/033270. Further sulphonylurea- and imidazolinone-tolerant plants have also been described, for example in WO 2007/024782.

5

Other plants tolerant to imidazolinone and/or sulphonylurea can be obtained by induced mutagenesis, by selection in cell cultures in the presence of the herbicide or by mutation breeding.

10

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which can also be treated are insect-resistant transgenic plants, i.e. plants which have been made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation which imparts such insect resistance.

15

In the present context, the term "insect-resistant transgenic plant" includes any plant containing at least one transgene comprising a coding sequence which encodes the following:

20

1) an insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins compiled online at: http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/, or insecticidal portions thereof, for example proteins of the Cry protein classes Cry1Ab, Cry1Ac, Cry1F, Cry2Ab, Cry3Ae or Cry3Bb or insecticidal portions thereof; or

25

30

2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein as *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cy34 and Cy35 crystal proteins; or

35

3) a hybrid insecticidal protein comprising parts of two different insecticidal crystal proteins from *Bacillus*

thuringiensis, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2) above, for example the Cry1A.105 protein produced by maize event MON98034 (WO 2007/027777); or

5

4) a protein of any one of points 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target
10 insect species affected, and/or because of changes induced in the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in maize events MON863 or MON88017, or the Cry3A protein in maize event MIR604; or

15 5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal proteins (VIP) listed
at:
http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.htm
20 1, for example proteins from the VIP3Aa protein class; or

6) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B.*
25 *cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins;

7) a hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or
30 *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the proteins in 2) above; or

8) a protein of any one of points 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by
35 another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes induced in the encoding DNA during cloning or transformation (while still

encoding an insecticidal protein), such as the VIP3Aa protein in cotton event COT102.

Of course, insect-resistant transgenic plants, as used herein, also include any plant comprising a combination of genes encoding the proteins of any one of the abovementioned classes 1 to 8. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the abovementioned classes 1 to 8, to expand the range of target insect species affected or to delay insect resistance development to the plants, by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated are tolerant to abiotic stress factors. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress-tolerant plants include the following:

a. plants which contain a transgene capable of reducing the expression and/or the activity of the poly(ADP-ribose)polymerase (PARP) gene in the plant cells or plants;

b. plants which contain a stress tolerance-enhancing transgene capable of reducing the expression and/or the activity of the PARG-encoding genes of the plants or plant cells;

c. plants which contain a stress tolerance-enhancing transgene coding for a plant-functional enzyme of the nicotinamide adenine dinucleotide salvage biosynthesis pathway, including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenyltransferase, nicotinamide adenine dinucleotide

synthetase or nicotinamide phosphoribosyltransferase.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated
5 show altered quantity, quality and/or storage stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as:

1) Transgenic plants which synthesize a modified starch
10 which is altered with respect to its chemophysical traits, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the distribution of the side chains, the viscosity behaviour, the gel resistance, the grain size and/or grain morphology of the
15 starch in comparison to the synthesized starch in wild-type plant cells or plants, such that this modified starch is better suited for certain applications.

2) Transgenic plants which synthesize non-starch
20 carbohydrate polymers or which synthesize non-starch carbohydrate polymers with altered properties in comparison to wild-type plants without genetic modification. Examples are plants which produce polyfructose, especially of the inulin and levan type, plants which produce alpha-1,4-glucans, plants
25 which produce alpha-1,6-branched alpha-1,4-glucans, and plants producing alternan.

3) Transgenic plants which produce hyaluronan.

30 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated are plants, such as cotton plants, with altered fibre characteristics. Such plants can be obtained by genetic transformation, or by selection of plants containing a
35 mutation imparting such altered fibre characteristics, and include:

a) plants, such as cotton plants, which contain an altered

form of cellulose synthase genes;

b) plants, such as cotton plants, which contain an altered form of rsw2 or rsw3 homologous nucleic acids;

5

c) plants, such as cotton plants, with an increased expression of sucrose phosphate synthase;

10

d) plants, such as cotton plants, with an increased expression of sucrose synthase;

15

e) plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fibre cell is altered, for example through downregulation of fibre-selective β -1,3-glucanase;

20

f) plants, such as cotton plants, which have fibres with altered reactivity, for example through the expression of the N-acetylglucosaminetransferase gene including nodC and chitin synthase genes.

25

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such altered oil characteristics, and include:

30

a) plants, such as oilseed rape plants, which produce oil having a high oleic acid content;

35

b) plants, such as oilseed rape plants, which produce oil having a low linolenic acid content;

c) plants, such as oilseed rape plants, which produce oil having a low level of saturated fatty acids.

Particularly useful transgenic plants which may be treated are plants which comprise one or more genes which encode one or more toxins, and are the transgenic plants which are sold under the following trade names: YIELD GARD® (for example
5 corn, cotton, soybeans), KnockOut® (for example corn), BiteGard® (for example corn), BT-Xtra® (for example corn), StarLink® (for example corn), Bollgard® (cotton), Nucotn® (cotton), Nucotn 33B® (cotton), NatureGard® (for example corn), Protecta® and NewLeaf® (potato). Examples of herbicide-
10 tolerant plants which should be mentioned are corn varieties, cotton varieties and soybean varieties which are available under the following trade names: Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya beans), Liberty Link® (tolerance to phosphinothricin, for example oilseed
15 rape), IMI® (tolerance to imidazolinone) and SCS® (tolerance to sulphonylurea, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned include the varieties sold under the name Clearfield (for example maize).

20

Particularly useful transgenic plants which may be treated are plants containing transformation events, or a combination of transformation events, and that are listed for example in the databases for various national or regional regulatory agencies
25 (see for example http://gmoinfo.jrc.it/gmp_browse.aspx and <http://www.agbios.com/dbase.php>).

The plants listed can be treated in a particularly advantageous manner with the compounds of the general formula
30 **(I)** and/or the active ingredient mixtures. The preferred ranges stated above for the active ingredients or mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the compounds or mixtures specifically mentioned in the present
35 text.

The active ingredients or compositions prepared with the aid of the inventive compounds can thus be used to protect plants

from attack by the pathogens mentioned for a certain period of time after treatment. The period for which protection is provided extends generally for 1 to 28 days, preferably for 1 to 14 days, particularly preferably for 1 to 10 days, very particularly preferably for 1 to 7 days after the treatment of the plants with the active ingredients, or for up to 200 days after a seed treatment.

The preparation and the use of the active ingredients of the formula (I) is illustrated by the examples which follow. However, the invention is not limited to these examples.

Preparation examples

General notes: Unless stated otherwise, all chromatographic purification and separation steps are carried out on silica gel and using a solvent gradient from 0:100 ethyl acetate/cyclohexane to 100:0 ethyl acetate/cyclohexane.

Preparation of compounds of the formula (I)

2-{3-[2-(1-[(Propan-2-ylideneamino)oxy]acetyl)piperidin-4-yl]-1,3-thiazol-4-yl]-1,2-oxazol-5-yl}benzaldehyde (I-8)

To a solution of *tert*-butyl 4-{4-[5-(2-formylphenyl)-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidine-1-carboxylate (2.20 g) in 1,4-dioxane was added dropwise, at 0°C, a 4 molar solution of hydrogen chloride (12 ml) in 1,4-dioxane. The reaction mixture was stirred at 0°C and then gradually warmed to room temperature. After stirring overnight, the solvent and excess hydrogen chloride were removed. This gave 4-{4-[5-(2-formylphenyl)-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidinium chloride (**XIIIa-99**, 2.0 g).

To a solution of [(propan-2-ylideneamino)oxy]acetic acid (185 mg) in dichloromethane (10 ml) were added, at 0°C, oxalyl chloride (168 µl) and one drop of *N,N*-dimethylformamide. The reaction mixture was stirred at room temperature for 120

minutes. Thereafter, the solvent and the excess reagent were removed under reduced pressure. The solid residue was dissolved again in dichloromethane and added dropwise at 0°C to a solution of 4-{4-[5-(2-formylphenyl)-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidinium chloride (484 mg) and triethylamine (357 µl) in dichloromethane (10 ml). The reaction mixture was stirred at room temperature for 1 h. Then concentrated sodium hydrogencarbonate solution was added thereto, and the aqueous phase was removed and extracted with ethyl acetate. The combined organic phases were dried over sodium sulphate and concentrated. Purification by column chromatography gave 2-{3-[2-(1-[(propan-2-ylideneamino)oxy]acetyl)piperidin-4-yl]-1,3-thiazol-4-yl]-1,2-oxazol-5-yl}benzaldehyde (150 mg).

15

2-{3-[2-(1-[(Propan-2-ylideneamino)oxy]acetyl)piperidin-4-yl]-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulphonate (I-12)

To a solution of acetone oxime (9.3 mg) in N,N-dimethylformamide (0.28 ml) at room temperature was added 3Å molecular sieve, and the mixture was stirred at this temperature for 2 hours. Thereafter, 2-(3-{2-[1-(chloroacetyl)piperidin-4-yl]-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulphonate (50 mg) and caesium carbonate (64 mg) were added thereto, and the mixture was stirred at room temperature for 18 hours. Subsequently, the mixture was filtered and extracted twice with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. Purification by column chromatography gave 2-{3-[2-(1-[(propan-2-ylideneamino)oxy]acetyl)piperidin-4-yl]-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulphonate (5 mg, 10%).

35 **Preparation of compounds of the formula (IV)**

[(Propan-2-ylideneamino)oxy]acetic acid (IV-1)

A mixture of (aminoxy)acetic acid hemihydrochloride (2.51 g) and acetone (6.0 g) was stirred at room temperature for 64 hours. Subsequently, dichloromethane (10 ml) was added to the mixture. Thereafter, the solvent and the excess reagent were removed under reduced pressure. This gave [(propan-2-ylideneamino)oxy]acetic acid (3.1 g), which was converted further without further purification.

{{[1-(4-Fluorophenyl)ethylidene]amino}oxy}acetic acid (IV-2)

10

Step 1:

A mixture of 1-(4-fluorophenyl)ethanone oxime (8.00 g) and caesium carbonate (20.4 g) in acetonitrile was stirred at 20°C for 30 minutes. Then ethyl bromoacetate (12.2 g) and potassium iodide (8.7 g) were added thereto and the mixture was stirred at 82°C for 3 hours. Then the reaction mixture was filtered. The solvent was removed from the filtrate under reduced pressure. Purification by column chromatography gave ethyl {{[1-(4-fluorophenyl)ethylidene]amino}oxy}acetate (8.7 g).

20

Step 2:

To a solution of ethyl {{[1-(4-fluorophenyl)ethylidene]amino}oxy}acetate (8.7 g) in a mixture of 50 ml of tetrahydrofuran and 10 ml of water was added, at 20°C, lithium hydroxide monohydrate (2.3 g) and the mixture was stirred at this temperature for 18 hours. Subsequently, the mixture was stirred into ice-cold 10% hydrochloric acid and extracted twice with ethyl acetate (50 ml each time). The combined organic phases were dried over sodium sulphate and concentrated. Purification by column chromatography on silica gel with a solvent gradient from 0:100 methanol/dichloromethane to 60:0 methanol/dichloromethane gave {{[1-(4-fluorophenyl)ethylidene]amino}oxy}acetic acid (2.9 g).

30

35

Preparation of compounds of the formula (X)

2-(3-{2-[1-(N,N-Dimethylglycyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl methanesulphonate (X-1)

To 4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidinium chloride (500 mg) in dimethylformamide (6 ml) under argon were added N,N-dimethylglycine (122 mg), diisopropylethylamine (582 mg) and O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU, 542 mg). The reaction mixture was stirred at room temperature for 18 hours. Then ice-cold sodium hydrogencarbonate solution was added thereto, the mixture was filtered, and the aqueous phase was removed and extracted with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. This gave 2-(3-{2-[1-(N,N-dimethylglycyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl methanesulphonate (310 mg, 55%).

LogP (pH2.7): 1.48

2-(3-{2-[1-({[tert-butyl(dimethyl)silyl]oxy}acetyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl methanesulphonate (X-2)

To 4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidinium chloride (500 mg) in dimethylformamide (6 ml) under argon were added {[tert-butyl(dimethyl)silyl]oxy}acetic acid (225 mg), diisopropylethylamine (582 mg) and O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU, 542 mg). The reaction mixture was stirred at room temperature for 18 hours. Then ice-cold sodium hydrogencarbonate solution was added thereto, the mixture was filtered, and the aqueous phase was removed and extracted with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. Purification by column chromatography gave 2-(3-{2-[1-({[tert-butyl(dimethyl)silyl]oxy}acetyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl

methanesulphonate (150 mg, 22%).

LogP (pH2.7): 4.14

5 **2-{4-[4-(5-{2-[(Methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}-2-oxoethyl acetate (X-3)**

To 4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidinium chloride (500 mg)
10 in dichloromethane (6 ml) under argon were added 2-chloro-2-oxoethyl acetate (154 mg) and triethylamine (342 mg). The reaction mixture was stirred at room temperature for 18 hours. Then water was added thereto, and it was filtered, dried and
15 concentrated. Purification by column chromatography gave 2-{4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}-2-oxoethyl acetate (170 mg, 30%).

20 LogP (pH2.7): 2.18

2-(3-{2-[1-(Chloroacetyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl methanesulphonate (Xc-a-142)

25

To a solution of chloroacetyl chloride (22 mg) was added, at 0°C, a solution of 4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidinium chloride (87 mg) and triethylamine (41 mg) in dichloromethane
30 (1 ml). The reaction mixture was stirred at 0°C for 15 minutes, and at room temperature for a further 18 hours. Then water was added thereto, and the aqueous phase was removed and extracted with dichloromethane. The combined organic phases were dried over magnesium sulphate and concentrated.
35 Purification by column chromatography gave 2-(3-{2-[1-(chloroacetyl)piperidin-4-yl]-1,3-thiazol-4-yl}-4,5-dihydro-1,2-oxazol-5-yl)phenyl methanesulphonate (60 mg, 60%).

LogP (pH2.7): 2.42

¹H NMR (250 MHz, CDCl₃): δ ppm : 1.7-2.0 (m, 2H), 2.15-2.35 (m, 2H), 2.80-2.95 (m, 1H), 3.05-3.20 (m, 1H), 3.30 (s, 3H), 3.27-
5 3.38 (m, 1H), 3.39-3.50 (dd, 1H), 3.85-3.97 (dd, 1H), 3.90-
4.10 (m, 1H), 4.20 (s, 2H), 4.55-4.66 (m, 1H), 5.98-6.06 (dd, 1H),
7.30-7.42 (m, 3H), 7.55-7.62 (m, 1H), 7.62 (s, 1H)

Preparation of compounds of the formula (XXVIIa)

10

2-{3-[2-(1-glycoloylpiperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulphonate (XXVIIa-142)

To 4-[4-(5-{2-[(methylsulphonyl)oxy]phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidinium chloride (110 mg) in dimethylformamide (6 ml) were added, under argon, glycolic acid (19 mg), diisopropylethylamine (32 mg) and O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU, 159 mg). Then diisopropylethylamine
20 (64 mg) was added once again to the reaction mixture. The reaction mixture was stirred at room temperature for 1 h. Then ice-cold sodium hydrogencarbonate solution was added thereto, the mixture was filtered, and the aqueous phase was removed and extracted with dichloromethane. The combined organic
25 phases were dried over magnesium sulphate and concentrated. Purification by column chromatography gave 2-{3-[2-(1-glycoloylpiperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulphonate (15 mg, 12%).

30 LogP (pH2.7): 1.88

¹H NMR (500 MHz, CDCl₃): δ ppm : 1.7-1.9 (m, 2H), 2.15-2.25 (m, 2H), 2.88-3.00 (m, 1H), 3.10-3.20 (m, 1H), 3.27 (s, 3H), 3.27-
35 3.38 (m, 1H), 3.39-3.47 (dd, 1H), 3.62 (m, 1H), 3.89-3.97 (dd, 1H),
4.20 (s, 2H), 4.60-4.66 (m, 1H), 5.98-6.06 (dd, 1H),
7.30-7.40 (m, 3H), 7.55-7.60 (m, 1H), 7.61 (s, 1H)

Preparation of compounds of the formula (XVI)

***tert*-Butyl 4-[4-(5-{2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidine-1-carboxylate (XVI-143)**

5

To a solution of 2-[3-(chloroacetyl)-4,5-dihydro-1,2-oxazol-5-yl]-5-(trifluoromethyl)phenyl methanesulphonate (200 mg) and *tert*-butyl 4-carbamothioylpiperidine-1-carboxylate (108 mg) in tetrahydrofuran (2 ml) at room temperature was added
10 tetrabutylammonium bromide. The reaction mixture was stirred at room temperature for 12 hours. Then water was added thereto, and the aqueous phase was removed and extracted with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. Purification by column
15 chromatography gave *tert*-butyl 4-[4-(5-{2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl}-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidine-1-carboxylate (168 mg, 56%).

20 LogP (pH2.7): 4.36

Preparation of compounds of the formula (VIIa)

2-[3-(Chloroacetyl)-4,5-dihydro-1,2-oxazol-5-yl]-5-(trifluoromethyl)phenyl methanesulphonate (VIIa-a-143)

25

To a solution of 5-(trifluoromethyl)-2-vinylphenyl methanesulphonate (1.05 g) in acetonitrile (10 ml) were added sodium hydrogencarbonate (2.55 g) and 3-chloro-N-hydroxy-2-oxopropanimidoyl chloride (0.60 g) at room temperature under
30 argon. The reaction mixture was stirred at room temperature for one hour. The solids were filtered off with suction and the filtrate was concentrated under reduced pressure. The residue was stirred with heptane to obtain 2-[3-(chloroacetyl)-4,5-dihydro-1,2-oxazol-5-yl]-5-(trifluoromethyl)phenyl methanesulphonate (1.33 g, 86% pure,
35 75%).

LogP (pH2.7): 3.25

Preparation of compounds of the formula (VIII)

5 **1-[5-(2-{[prop-2-yn-1-yl]oxy}phenyl)-4,5-dihydro-1,2-oxazol-3-yl]ethanone (VIIIa-a-81)**

Step 1:

10 To a solution of 3,3-dimethoxybutan-2-one (1.00 g) in ethanol (10 ml) was added dropwise hydroxylamine (50% in water, 0.23 ml) at room temperature. The reaction mixture was stirred at 50°C for 4 hours. Then water was added thereto, and the aqueous phase was removed and extracted with ethyl acetate.
15 The combined organic phases were dried over magnesium sulphate and concentrated. This gave 3,3-dimethoxybutan-2-one oxime (800 mg, 72%).

Step 2:

20 To a solution of 3,3-dimethoxybutan-2-one oxime (270 mg) in tetrahydrofuran (2.7 ml) was added dropwise, at 0°C under argon, n-butyllithium (2M in tetrahydrofuran, 1.83 ml). After stirring for a further 5 minutes, a solution of 2-{{3-(trimethylsilyl)prop-2-yn-1-yl}oxy}benzaldehyde (232 mg) in
25 tetrahydrofuran (1 ml) was added dropwise to the reaction mixture, which was stirred for a further 1 hour. Subsequently, concentrated ammonium chloride solution was added to the reaction mixture, and the aqueous phase was removed and
30 extracted with ethyl acetate. The combined organic phases were dried over magnesium sulphate and concentrated. Purification by column chromatography gave 1-hydroxyl-4,4-dimethoxy-1-(2-{{3-(trimethylsilyl)prop-2-yn-1-yl}oxy}phenyl)pentan-3-one oxime (482 mg, 69%).

35


LogP (pH2.7): 3.19

Step 3:

A solution of hydrochloric acid (4M in dioxane, 3.80 ml) was added to 1-hydroxyl-4,4-dimethoxy-1-(2-{{[3-(trimethylsilyl)prop-2-yn-1-yl]oxy}phenyl)pentan-3-one oxime. After stirring for a further 15 minutes, concentrated sodium hydrogencarbonate solution was added to the reaction mixture, and the aqueous phase was removed and extracted with dichloromethane. The combined organic phases were dried over magnesium sulphate and concentrated. This gave 1-[5-(2-{{[prop-2-yn-1-yl]oxy}phenyl)-4,5-dihydro-1,2-oxazol-3-yl]ethanone (315 mg, 99%).

LogP (pH2.7): 4.41

15 Compound examples



The structural elements G and Q listed in Table 1 are defined as follows:



For all compounds listed in Table 1, p=0 and L² = direct bond.


25

Table 1:

Ex.	R ^{A1}	R ^{A2}	L ¹	R ^{B1}	R ^{B2}	Y	X	R ²	R ¹	Log P
I-1	CH ₃	1,3-benzodio	O	H	H	O	C	H	2-fluoro-6-(prop-2-yn-	3.29 [a];

		xol-5-yl							1- yloxy)phenyl	3.28 [b]
I-2	propa n-2- yl	4- ethoxyph enyl	O	H	H	O	C	H	2-fluoro-6- (prop-2-yn- 1- yloxy)phenyl	4.19 [a]; 4.21 [b]
I-3	CH ₃	3- fluoroph enyl	O	H	H	O	C	H	2-fluoro-6- (prop-2-yn- 1- yloxy)phenyl	3.48 [a]; 3.51 [b]
I-4	CH ₃	3,4- dimethyl phenyl	O	H	H	O	C	H	2-fluoro-6- (prop-2-yn- 1- yloxy)phenyl	3.93 [a]; 3.94 [b]
I-5	CH ₃	3- (trifluo romethox y)phenyl	O	H	H	O	C	H	2-fluoro-6- (prop-2-yn- 1- yloxy)phenyl	4.01 [a]; 4.01 [b]
I-6	CH ₃	3,4- dimethyl phenyl	O	H	H	O	C	H	5-fluoro-2- (prop-2-yn- 1- yloxy)phenyl	4.18 [a]; 4.19 [b]
I-7	propa n-2- yl	4- ethoxyph enyl	O	H	H	O	C	H	5-fluoro-2- (prop-2-yn- 1- yloxy)phenyl	4.42 [a]; 4.42 [b]
I-8	CH ₃	CH ₃	O	H	H	O	C	H	2- formylphenyl	2.45 [a]
I-9	CH ₃	trifluor omethyl	O	H	H	O	C	H	2-fluoro-6- [(methylsulph honyl)oxy]ph enyl	2.91 [a]
I-10	CH ₃	trifluor omethyl	O	H	H	O	C	H	2- [(methylsulph honyl)oxy]ph enyl	2.94 [a]
I-11	CH ₃	CH ₃	O	H	H	O	C	H	2-(prop-2- yn-1-	2.75 [a]

										yl oxy) phenyl	
I-12	CH ₃	CH ₃	O	H	H	O	C	H		2-[(methylsulphonyl)oxy]phenyl	2.36 [a]
I-13	CH ₃	CH ₃	O	H	H	O	C	H		2-fluoro-6-[(methylsulphonyl)oxy]phenyl	2.34 [a]
I-14	CH ₃	1,1-dimethyl ethyl	O	H	H	O	C	H		2-[(methylsulphonyl)oxy]phenyl	3.42 [a]

Table 2:**(XIIIa)**

5

Ex.	R
XIIIa-1	2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-2	2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl
XIIIa-3	2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-4	2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIa-5	2,3-difluoro-4-formylphenyl
XIIIa-6	2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-7	2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-8	2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-9	2,4-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-10	2,4-difluoro-3-formylphenyl
XIIIa-11	2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-12	2,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-13	2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-14	2,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
XIIIa-15	2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-16	2,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-17	2,5-difluoro-3-formylphenyl

XIIIIa-18	2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-19	2,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-20	2,5-difluoro-4-formylphenyl
XIIIIa-21	2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-22	2,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-23	2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-24	2,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-25	2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-26	2,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-27	2,6-difluoro-3-formylphenyl
XIIIIa-28	2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-29	2,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-30	2,6-difluoro-4-formylphenyl
XIIIIa-31	2-(allyloxy)-3,4-dichlorophenyl
XIIIIa-32	2-(allyloxy)-3,4-difluorophenyl
XIIIIa-33	2-(allyloxy)-3,5-dichlorophenyl
XIIIIa-34	2-(allyloxy)-3,5-difluorophenyl
XIIIIa-35	2-(allyloxy)-3,6-dichlorophenyl
XIIIIa-36	2-(allyloxy)-3,6-difluorophenyl
XIIIIa-37	2-(allyloxy)-3-chlorophenyl
XIIIIa-38	2-(allyloxy)-3-fluorophenyl
XIIIIa-39	2-(allyloxy)-3-methylphenyl
XIIIIa-40	2-(allyloxy)-4,5-dichlorophenyl
XIIIIa-41	2-(allyloxy)-4,5-difluorophenyl
XIIIIa-42	2-(allyloxy)-4,6-dichlorophenyl
XIIIIa-43	2-(allyloxy)-4,6-difluorophenyl
XIIIIa-44	2-(allyloxy)-4-chlorophenyl
XIIIIa-45	2-(allyloxy)-4-fluorophenyl
XIIIIa-46	2-(allyloxy)-4-methylphenyl
XIIIIa-47	2-(allyloxy)-5,6-dichlorophenyl
XIIIIa-48	2-(allyloxy)-5,6-difluorophenyl
XIIIIa-49	2-(allyloxy)-5-chlorophenyl
XIIIIa-50	2-(allyloxy)-5-fluorophenyl
XIIIIa-51	2-(allyloxy)-5-methylphenyl
XIIIIa-52	2-(allyloxy)-6-chlorophenyl
XIIIIa-53	2-(allyloxy)-6-fluorophenyl
XIIIIa-54	2-(allyloxy)-6-methylphenyl
XIIIIa-55	2-(allyloxy)phenyl

XIIIIa-56	2-(cyanomethoxy)-3,4-dichlorophenyl
XIIIIa-57	2-(cyanomethoxy)-3,4-difluorophenyl
XIIIIa-58	2-(cyanomethoxy)-3,5-dichlorophenyl
XIIIIa-59	2-(cyanomethoxy)-3,5-difluorophenyl
XIIIIa-60	2-(cyanomethoxy)-3,6-dichlorophenyl
XIIIIa-61	2-(cyanomethoxy)-3,6-difluorophenyl
XIIIIa-62	2-(cyanomethoxy)-3-chlorophenyl
XIIIIa-63	2-(cyanomethoxy)-3-fluorophenyl
XIIIIa-64	2-(cyanomethoxy)-3-methylphenyl
XIIIIa-65	2-(cyanomethoxy)-4,5-dichlorophenyl
XIIIIa-66	2-(cyanomethoxy)-4,5-difluorophenyl
XIIIIa-67	2-(cyanomethoxy)-4,6-dichlorophenyl
XIIIIa-68	2-(cyanomethoxy)-4,6-difluorophenyl
XIIIIa-69	2-(cyanomethoxy)-4-chlorophenyl
XIIIIa-70	2-(cyanomethoxy)-4-fluorophenyl
XIIIIa-71	2-(cyanomethoxy)-4-methylphenyl
XIIIIa-72	2-(cyanomethoxy)-5,6-dichlorophenyl
XIIIIa-73	2-(cyanomethoxy)-5,6-difluorophenyl
XIIIIa-74	2-(cyanomethoxy)-5-chlorophenyl
XIIIIa-75	2-(cyanomethoxy)-5-fluorophenyl
XIIIIa-76	2-(cyanomethoxy)-5-methylphenyl
XIIIIa-77	2-(cyanomethoxy)-6-chlorophenyl
XIIIIa-78	2-(cyanomethoxy)-6-fluorophenyl
XIIIIa-79	2-(cyanomethoxy)-6-methylphenyl
XIIIIa-80	2-(cyanomethoxy)phenyl
XIIIIa-81	2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-82	2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl
XIIIIa-83	2-chloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-84	2-chloro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-85	2-chloro-3-formylphenyl
XIIIIa-86	2-chloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-87	2-chloro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-88	2-chloro-4-formylphenyl
XIIIIa-89	2-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-90	2-fluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-91	2-fluoro-3-formylphenyl
XIIIIa-92	2-fluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-93	2-fluoro-4-[(methylsulphonyl)oxy]phenyl

XIIIIa-94	2-fluoro-4-formylphenyl
XIIIIa-95	2-formyl-3-methylphenyl
XIIIIa-96	2-formyl-4-methylphenyl
XIIIIa-97	2-formyl-5-methylphenyl
XIIIIa-98	2-formyl-6-methylphenyl
XIIIIa-99	2-formylphenyl
XIIIIa-100	2-methyl-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-101	2-methyl-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-102	2-methyl-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-103	2-methyl-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-104	2-[(hydroxyimino)methyl]-3,4-difluorophenyl
XIIIIa-105	2-[(hydroxyimino)methyl]-3,5-difluorophenyl
XIIIIa-106	2-[(hydroxyimino)methyl]-3,6-difluorophenyl
XIIIIa-107	2-[(hydroxyimino)methyl]-3-chlorophenyl
XIIIIa-108	2-[(hydroxyimino)methyl]-3-fluorophenyl
XIIIIa-109	2-[(hydroxyimino)methyl]-3-methylphenyl
XIIIIa-110	2-[(hydroxyimino)methyl]-4,5-difluorophenyl
XIIIIa-111	2-[(hydroxyimino)methyl]-4,6-difluorophenyl
XIIIIa-112	2-[(hydroxyimino)methyl]-4-chlorophenyl
XIIIIa-113	2-[(hydroxyimino)methyl]-4-fluorophenyl
XIIIIa-114	2-[(hydroxyimino)methyl]-4-methylphenyl
XIIIIa-115	2-[(hydroxyimino)methyl]-5,6-difluorophenyl
XIIIIa-116	2-[(hydroxyimino)methyl]-5-chlorophenyl
XIIIIa-117	2-[(hydroxyimino)methyl]-5-fluorophenyl
XIIIIa-118	2-[(hydroxyimino)methyl]-5-methylphenyl
XIIIIa-119	2-[(hydroxyimino)methyl]-6-chlorophenyl
XIIIIa-120	2-[(hydroxyimino)methyl]-6-fluorophenyl
XIIIIa-121	2-[(hydroxyimino)methyl]-6-methylphenyl
XIIIIa-122	2-[(hydroxyimino)methyl]phenyl
XIIIIa-123	2-[(methoxyimino)methyl]-3,4-difluorophenyl
XIIIIa-124	2-[(methoxyimino)methyl]-3,5-difluorophenyl
XIIIIa-125	2-[(methoxyimino)methyl]-3,6-difluorophenyl
XIIIIa-126	2-[(methoxyimino)methyl]-3-chlorophenyl
XIIIIa-127	2-[(methoxyimino)methyl]-3-fluorophenyl
XIIIIa-128	2-[(methoxyimino)methyl]-3-methylphenyl
XIIIIa-129	2-[(methoxyimino)methyl]-4,5-difluorophenyl
XIIIIa-130	2-[(methoxyimino)methyl]-4,6-difluorophenyl
XIIIIa-131	2-[(methoxyimino)methyl]-4-chlorophenyl

XIIIIa-132	2-[(methoxyimino)methyl]-4-fluorophenyl
XIIIIa-133	2-[(methoxyimino)methyl]-4-methylphenyl
XIIIIa-134	2-[(methoxyimino)methyl]-5,6-difluorophenyl
XIIIIa-135	2-[(methoxyimino)methyl]-5-chlorophenyl
XIIIIa-136	2-[(methoxyimino)methyl]-5-fluorophenyl
XIIIIa-137	2-[(methoxyimino)methyl]-5-methylphenyl
XIIIIa-138	2-[(methoxyimino)methyl]-6-chlorophenyl
XIIIIa-139	2-[(methoxyimino)methyl]-6-fluorophenyl
XIIIIa-140	2-[(methoxyimino)methyl]-6-methylphenyl
XIIIIa-141	2-[(methoxyimino)methyl]phenyl
XIIIIa-142	2-[(methylsulphonyl)oxy]phenyl
XIIIIa-143	2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl
XIIIIa-144	3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-145	3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-146	3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-147	3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-148	3,4-difluoro-2-formylphenyl
XIIIIa-149	3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-150	3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-151	3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-152	3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-153	3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-154	3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-155	3,5-difluoro-2-formylphenyl
XIIIIa-156	3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-157	3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-158	3,5-difluoro-4-formylphenyl
XIIIIa-159	3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-160	3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-161	3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-162	3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-163	3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-164	3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-165	3,6-difluoro-2-formylphenyl
XIIIIa-166	3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-167	3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIIa-168	3,6-difluoro-4-formylphenyl
XIIIIa-169	3-(allyloxy)-2,4-dichlorophenyl

XIIIa-170	3-(allyloxy)-2,4-difluorophenyl
XIIIa-171	3-(allyloxy)-2,5-dichlorophenyl
XIIIa-172	3-(allyloxy)-2,5-difluorophenyl
XIIIa-173	3-(allyloxy)-2,6-dichlorophenyl
XIIIa-174	3-(allyloxy)-2,6-difluorophenyl
XIIIa-175	3-(allyloxy)-2-chlorophenyl
XIIIa-176	3-(allyloxy)-2-fluorophenyl
XIIIa-177	3-(allyloxy)-2-methylphenyl
XIIIa-178	3-(allyloxy)-4,5-dichlorophenyl
XIIIa-179	3-(allyloxy)-4,5-difluorophenyl
XIIIa-180	3-(allyloxy)-4,6-dichlorophenyl
XIIIa-181	3-(allyloxy)-4,6-difluorophenyl
XIIIa-182	3-(allyloxy)-4-chlorophenyl
XIIIa-183	3-(allyloxy)-4-fluorophenyl
XIIIa-184	3-(allyloxy)-4-methylphenyl
XIIIa-185	3-(allyloxy)-5,6-dichlorophenyl
XIIIa-186	3-(allyloxy)-5,6-difluorophenyl
XIIIa-187	3-(allyloxy)-5-chlorophenyl
XIIIa-188	3-(allyloxy)-5-fluorophenyl
XIIIa-189	3-(allyloxy)-5-methylphenyl
XIIIa-190	3-(allyloxy)-6-chlorophenyl
XIIIa-191	3-(allyloxy)-6-fluorophenyl
XIIIa-192	3-(allyloxy)-6-methylphenyl
XIIIa-193	3-(allyloxy)phenyl
XIIIa-194	3-(cyanomethoxy)-2,4-dichlorophenyl
XIIIa-195	3-(cyanomethoxy)-2,4-difluorophenyl
XIIIa-196	3-(cyanomethoxy)-2,5-dichlorophenyl
XIIIa-197	3-(cyanomethoxy)-2,5-difluorophenyl
XIIIa-198	3-(cyanomethoxy)-2,6-dichlorophenyl
XIIIa-199	3-(cyanomethoxy)-2,6-difluorophenyl
XIIIa-200	3-(cyanomethoxy)-2-chlorophenyl
XIIIa-201	3-(cyanomethoxy)-2-fluorophenyl
XIIIa-202	3-(cyanomethoxy)-2-methylphenyl
XIIIa-203	3-(cyanomethoxy)-4,5-dichlorophenyl
XIIIa-204	3-(cyanomethoxy)-4,5-difluorophenyl
XIIIa-205	3-(cyanomethoxy)-4,6-dichlorophenyl
XIIIa-206	3-(cyanomethoxy)-4,6-difluorophenyl
XIIIa-207	3-(cyanomethoxy)-4-chlorophenyl

XIIIa-208	3-(cyanomethoxy)-4-fluorophenyl
XIIIa-209	3-(cyanomethoxy)-4-methylphenyl
XIIIa-210	3-(cyanomethoxy)-5,6-dichlorophenyl
XIIIa-211	3-(cyanomethoxy)-5,6-difluorophenyl
XIIIa-212	3-(cyanomethoxy)-5-chlorophenyl
XIIIa-213	3-(cyanomethoxy)-5-fluorophenyl
XIIIa-214	3-(cyanomethoxy)-5-methylphenyl
XIIIa-215	3-(cyanomethoxy)-6-chlorophenyl
XIIIa-216	3-(cyanomethoxy)-6-fluorophenyl
XIIIa-217	3-(cyanomethoxy)-6-methylphenyl
XIIIa-218	3-(cyanomethoxy)phenyl
XIIIa-219	3-(prop-2-yn-1-yloxy)phenyl
XIIIa-220	3-chloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-221	3-chloro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-222	3-chloro-2-formylphenyl
XIIIa-223	3-chloro-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-224	3-chloro-4-[(methylsulphonyl)oxy]phenyl
XIIIa-225	3-chloro-4-formylphenyl
XIIIa-226	3-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-227	3-fluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-228	3-fluoro-2-formylphenyl
XIIIa-229	3-fluoro-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-230	3-fluoro-4-[(methylsulphonyl)oxy]phenyl
XIIIa-231	3-fluoro-4-formylphenyl
XIIIa-232	3-formyl-2-methylphenyl
XIIIa-233	3-formyl-4-methylphenyl
XIIIa-234	3-formyl-5-methylphenyl
XIIIa-235	3-formyl-6-methylphenyl
XIIIa-236	3-formylphenyl
XIIIa-237	3-methyl-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-238	3-methyl-2-[(methylsulphonyl)oxy]phenyl
XIIIa-239	3-methyl-4-(prop-2-yn-1-yloxy)phenyl
XIIIa-240	3-methyl-4-[(methylsulphonyl)oxy]phenyl
XIIIa-241	3-[(hydroxyimino)methyl]-2,4-difluorophenyl
XIIIa-242	3-[(hydroxyimino)methyl]-2,5-difluorophenyl
XIIIa-243	3-[(hydroxyimino)methyl]-2,6-difluorophenyl
XIIIa-244	3-[(hydroxyimino)methyl]-2-chlorophenyl
XIIIa-245	3-[(hydroxyimino)methyl]-2-fluorophenyl


XIIIa-246	3-[(hydroxyimino)methyl]-2-methylphenyl
XIIIa-247	3-[(hydroxyimino)methyl]-4,5-difluorophenyl
XIIIa-248	3-[(hydroxyimino)methyl]-4,6-difluorophenyl
XIIIa-249	3-[(hydroxyimino)methyl]-4-chlorophenyl
XIIIa-250	3-[(hydroxyimino)methyl]-4-fluorophenyl
XIIIa-251	3-[(hydroxyimino)methyl]-4-methylphenyl
XIIIa-252	3-[(hydroxyimino)methyl]-5,6-difluorophenyl
XIIIa-253	3-[(hydroxyimino)methyl]-5-chlorophenyl
XIIIa-254	3-[(hydroxyimino)methyl]-5-fluorophenyl
XIIIa-255	3-[(hydroxyimino)methyl]-5-methylphenyl
XIIIa-256	3-[(hydroxyimino)methyl]-6-chlorophenyl
XIIIa-257	3-[(hydroxyimino)methyl]-6-fluorophenyl
XIIIa-258	3-[(hydroxyimino)methyl]-6-methylphenyl
XIIIa-259	3-[(hydroxyimino)methyl]phenyl
XIIIa-260	3-[(methoxyimino)methyl]-2,4-difluorophenyl
XIIIa-261	3-[(methoxyimino)methyl]-2,5-difluorophenyl
XIIIa-262	3-[(methoxyimino)methyl]-2,6-difluorophenyl
XIIIa-263	3-[(methoxyimino)methyl]-2-chlorophenyl
XIIIa-264	3-[(methoxyimino)methyl]-2-fluorophenyl
XIIIa-265	3-[(methoxyimino)methyl]-2-methylphenyl
XIIIa-266	3-[(methoxyimino)methyl]-4,5-difluorophenyl
XIIIa-267	3-[(methoxyimino)methyl]-4,6-difluorophenyl
XIIIa-268	3-[(methoxyimino)methyl]-4-chlorophenyl
XIIIa-269	3-[(methoxyimino)methyl]-4-fluorophenyl
XIIIa-270	3-[(methoxyimino)methyl]-4-methylphenyl
XIIIa-271	3-[(methoxyimino)methyl]-5,6-difluorophenyl
XIIIa-272	3-[(methoxyimino)methyl]-5-chlorophenyl
XIIIa-273	3-[(methoxyimino)methyl]-5-fluorophenyl
XIIIa-274	3-[(methoxyimino)methyl]-5-methylphenyl
XIIIa-275	3-[(methoxyimino)methyl]-6-chlorophenyl
XIIIa-276	3-[(methoxyimino)methyl]-6-fluorophenyl
XIIIa-277	3-[(methoxyimino)methyl]-6-methylphenyl
XIIIa-278	3-[(methoxyimino)methyl]phenyl
XIIIa-279	3-[(methylsulphonyl)oxy]phenyl
XIIIa-280	4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-281	4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-282	4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-283	4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl

XIIIa-284	4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-285	4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-286	4,5-difluoro-2-formylphenyl
XIIIa-287	4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-288	4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-289	4,5-difluoro-3-formylphenyl
XIIIa-290	4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-291	4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-292	4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-293	4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-294	4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-295	4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-296	4,6-difluoro-2-formylphenyl
XIIIa-297	4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-298	4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-299	4,6-difluoro-3-formylphenyl
XIIIa-300	4-(allyloxy)-2,3-dichlorophenyl
XIIIa-301	4-(allyloxy)-2,3-difluorophenyl
XIIIa-302	4-(allyloxy)-2,5-dichlorophenyl
XIIIa-303	4-(allyloxy)-2,5-difluorophenyl
XIIIa-304	4-(allyloxy)-2,6-dichlorophenyl
XIIIa-305	4-(allyloxy)-2,6-difluorophenyl
XIIIa-306	4-(allyloxy)-2-chlorophenyl
XIIIa-307	4-(allyloxy)-2-fluorophenyl
XIIIa-308	4-(allyloxy)-2-methylphenyl
XIIIa-309	4-(allyloxy)-3,5-dichlorophenyl
XIIIa-310	4-(allyloxy)-3,5-difluorophenyl
XIIIa-311	4-(allyloxy)-3,6-dichlorophenyl
XIIIa-312	4-(allyloxy)-3,6-difluorophenyl
XIIIa-313	4-(allyloxy)-3-chlorophenyl
XIIIa-314	4-(allyloxy)-3-fluorophenyl
XIIIa-315	4-(allyloxy)-3-methylphenyl
XIIIa-316	4-(allyloxy)phenyl
XIIIa-317	4-(cyanomethoxy)-2,3-dichlorophenyl
XIIIa-318	4-(cyanomethoxy)-2,3-difluorophenyl
XIIIa-319	4-(cyanomethoxy)-2,5-dichlorophenyl
XIIIa-320	4-(cyanomethoxy)-2,5-difluorophenyl
XIIIa-321	4-(cyanomethoxy)-2,6-dichlorophenyl

XIIIIa-322	4-(cyanomethoxy)-2,6-difluorophenyl
XIIIIa-323	4-(cyanomethoxy)-2-chlorophenyl
XIIIIa-324	4-(cyanomethoxy)-2-fluorophenyl
XIIIIa-325	4-(cyanomethoxy)-2-methylphenyl
XIIIIa-326	4-(cyanomethoxy)-3,5-dichlorophenyl
XIIIIa-327	4-(cyanomethoxy)-3,5-difluorophenyl
XIIIIa-328	4-(cyanomethoxy)-3,6-dichlorophenyl
XIIIIa-329	4-(cyanomethoxy)-3,6-difluorophenyl
XIIIIa-330	4-(cyanomethoxy)-3-chlorophenyl
XIIIIa-331	4-(cyanomethoxy)-3-fluorophenyl
XIIIIa-332	4-(cyanomethoxy)-3-methylphenyl
XIIIIa-333	4-(cyanomethoxy)phenyl
XIIIIa-334	4-(prop-2-yn-1-yloxy)phenyl
XIIIIa-335	4-chloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-336	4-chloro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-337	4-chloro-2-formylphenyl
XIIIIa-338	4-chloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-339	4-chloro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-340	4-chloro-3-formylphenyl
XIIIIa-341	4-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-342	4-fluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-343	4-fluoro-2-formylphenyl
XIIIIa-344	4-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-345	4-fluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-346	4-fluoro-3-formylphenyl
XIIIIa-347	4-formyl-2-methylphenyl
XIIIIa-348	4-formyl-3-methylphenyl
XIIIIa-349	4-formylphenyl
XIIIIa-350	4-methyl-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-351	4-methyl-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-352	4-methyl-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-353	4-methyl-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-354	4-[(hydroxyimino)methyl]-2,3-difluorophenyl
XIIIIa-355	4-[(hydroxyimino)methyl]-2,5-difluorophenyl
XIIIIa-356	4-[(hydroxyimino)methyl]-2,6-difluorophenyl
XIIIIa-357	4-[(hydroxyimino)methyl]-2-chlorophenyl
XIIIIa-358	4-[(hydroxyimino)methyl]-2-fluorophenyl
XIIIIa-359	4-[(hydroxyimino)methyl]-2-methylphenyl

XIIIa-360	4-[(hydroxyimino)methyl]-3,5-difluorophenyl
XIIIa-361	4-[(hydroxyimino)methyl]-3,6-difluorophenyl
XIIIa-362	4-[(hydroxyimino)methyl]-3-chlorophenyl
XIIIa-363	4-[(hydroxyimino)methyl]-3-fluorophenyl
XIIIa-364	4-[(hydroxyimino)methyl]-3-methylphenyl
XIIIa-365	4-[(hydroxyimino)methyl]phenyl
XIIIa-366	4-[(methoxyimino)methyl]-2,3-difluorophenyl
XIIIa-367	4-[(methoxyimino)methyl]-2,5-difluorophenyl
XIIIa-368	4-[(methoxyimino)methyl]-2,6-difluorophenyl
XIIIa-369	4-[(methoxyimino)methyl]-2-chlorophenyl
XIIIa-370	4-[(methoxyimino)methyl]-2-fluorophenyl
XIIIa-371	4-[(methoxyimino)methyl]-2-methylphenyl
XIIIa-372	4-[(methoxyimino)methyl]-3,5-difluorophenyl
XIIIa-373	4-[(methoxyimino)methyl]-3,6-difluorophenyl
XIIIa-374	4-[(methoxyimino)methyl]-3-chlorophenyl
XIIIa-375	4-[(methoxyimino)methyl]-3-fluorophenyl
XIIIa-376	4-[(methoxyimino)methyl]-3-methylphenyl
XIIIa-377	4-[(methoxyimino)methyl]phenyl
XIIIa-378	4-[(methylsulphonyl)oxy]phenyl
XIIIa-379	5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-380	5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-381	5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-382	5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-383	5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-384	5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-385	5,6-difluoro-2-formylphenyl
XIIIa-386	5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-387	5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-388	5,6-difluoro-3-formylphenyl
XIIIa-389	5-chloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-390	5-chloro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-391	5-chloro-2-formylphenyl
XIIIa-392	5-chloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIa-393	5-chloro-3-[(methylsulphonyl)oxy]phenyl
XIIIa-394	5-chloro-3-formylphenyl
XIIIa-395	5-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIa-396	5-fluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIa-397	5-fluoro-2-formylphenyl

XIIIIa-398	5-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-399	5-fluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-400	5-fluoro-3-formylphenyl
XIIIIa-401	5-methyl-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-402	5-methyl-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-403	5-methyl-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-404	5-methyl-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-405	6-chloro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-406	6-chloro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-407	6-chloro-2-formylphenyl
XIIIIa-408	6-chloro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-409	6-chloro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-410	6-chloro-3-formylphenyl
XIIIIa-411	6-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-412	6-fluoro-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-413	6-fluoro-2-formylphenyl
XIIIIa-414	6-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-415	6-fluoro-3-[(methylsulphonyl)oxy]phenyl
XIIIIa-416	6-fluoro-3-formylphenyl
XIIIIa-417	6-methyl-2-(prop-2-yn-1-yloxy)phenyl
XIIIIa-418	6-methyl-2-[(methylsulphonyl)oxy]phenyl
XIIIIa-419	6-methyl-3-(prop-2-yn-1-yloxy)phenyl
XIIIIa-420	6-methyl-3-[(methylsulphonyl)oxy]phenyl

Table 3:

5

Ex.	R ¹
XXVIIa-1	2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-2	2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-3	2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-4	2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-5	2,3-difluoro-4-formylphenyl
XXVIIa-6	2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-7	2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl

XXVIIa-8	2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-9	2,4-difluoro-3-[(methylysulphonyl)oxy]phenyl
XXVIIa-10	2,4-difluoro-3-formylphenyl
XXVIIa-11	2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-12	2,5-dichloro-3-[(methylysulphonyl)oxy]phenyl
XXVIIa-13	2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-14	2,5-dichloro-4-[(methylysulphonyl)oxy]phenyl
XXVIIa-15	2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-16	2,5-difluoro-3-[(methylysulphonyl)oxy]phenyl
XXVIIa-17	2,5-difluoro-3-formylphenyl
XXVIIa-18	2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-19	2,5-difluoro-4-[(methylysulphonyl)oxy]phenyl
XXVIIa-20	2,5-difluoro-4-formylphenyl
XXVIIa-21	2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-22	2,6-dichloro-3-[(methylysulphonyl)oxy]phenyl
XXVIIa-23	2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-24	2,6-dichloro-4-[(methylysulphonyl)oxy]phenyl
XXVIIa-25	2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-26	2,6-difluoro-3-[(methylysulphonyl)oxy]phenyl
XXVIIa-27	2,6-difluoro-3-formylphenyl
XXVIIa-28	2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-29	2,6-difluoro-4-[(methylysulphonyl)oxy]phenyl
XXVIIa-30	2,6-difluoro-4-formylphenyl
XXVIIa-31	2-(allyloxy)-3,4-dichlorophenyl
XXVIIa-32	2-(allyloxy)-3,4-difluorophenyl
XXVIIa-33	2-(allyloxy)-3,5-dichlorophenyl
XXVIIa-34	2-(allyloxy)-3,5-difluorophenyl
XXVIIa-35	2-(allyloxy)-3,6-dichlorophenyl
XXVIIa-36	2-(allyloxy)-3,6-difluorophenyl
XXVIIa-37	2-(allyloxy)-3-chlorophenyl
XXVIIa-38	2-(allyloxy)-3-fluorophenyl
XXVIIa-39	2-(allyloxy)-3-methylphenyl
XXVIIa-40	2-(allyloxy)-4,5-dichlorophenyl
XXVIIa-41	2-(allyloxy)-4,5-difluorophenyl
XXVIIa-42	2-(allyloxy)-4,6-dichlorophenyl
XXVIIa-43	2-(allyloxy)-4,6-difluorophenyl
XXVIIa-44	2-(allyloxy)-4-chlorophenyl
XXVIIa-45	2-(allyloxy)-4-fluorophenyl

XXVIIa-46	2-(allyloxy)-4-methylphenyl
XXVIIa-47	2-(allyloxy)-5,6-dichlorophenyl
XXVIIa-48	2-(allyloxy)-5,6-difluorophenyl
XXVIIa-49	2-(allyloxy)-5-chlorophenyl
XXVIIa-50	2-(allyloxy)-5-fluorophenyl
XXVIIa-51	2-(allyloxy)-5-methylphenyl
XXVIIa-52	2-(allyloxy)-6-chlorophenyl
XXVIIa-53	2-(allyloxy)-6-fluorophenyl
XXVIIa-54	2-(allyloxy)-6-methylphenyl
XXVIIa-55	2-(allyloxy)phenyl
XXVIIa-56	2-(cyanomethoxy)-3,4-dichlorophenyl
XXVIIa-57	2-(cyanomethoxy)-3,4-difluorophenyl
XXVIIa-58	2-(cyanomethoxy)-3,5-dichlorophenyl
XXVIIa-59	2-(cyanomethoxy)-3,5-difluorophenyl
XXVIIa-60	2-(cyanomethoxy)-3,6-dichlorophenyl
XXVIIa-61	2-(cyanomethoxy)-3,6-difluorophenyl
XXVIIa-62	2-(cyanomethoxy)-3-chlorophenyl
XXVIIa-63	2-(cyanomethoxy)-3-fluorophenyl
XXVIIa-64	2-(cyanomethoxy)-3-methylphenyl
XXVIIa-65	2-(cyanomethoxy)-4,5-dichlorophenyl
XXVIIa-66	2-(cyanomethoxy)-4,5-difluorophenyl
XXVIIa-67	2-(cyanomethoxy)-4,6-dichlorophenyl
XXVIIa-68	2-(cyanomethoxy)-4,6-difluorophenyl
XXVIIa-69	2-(cyanomethoxy)-4-chlorophenyl
XXVIIa-70	2-(cyanomethoxy)-4-fluorophenyl
XXVIIa-71	2-(cyanomethoxy)-4-methylphenyl
XXVIIa-72	2-(cyanomethoxy)-5,6-dichlorophenyl
XXVIIa-73	2-(cyanomethoxy)-5,6-difluorophenyl
XXVIIa-74	2-(cyanomethoxy)-5-chlorophenyl
XXVIIa-75	2-(cyanomethoxy)-5-fluorophenyl
XXVIIa-76	2-(cyanomethoxy)-5-methylphenyl
XXVIIa-77	2-(cyanomethoxy)-6-chlorophenyl
XXVIIa-78	2-(cyanomethoxy)-6-fluorophenyl
XXVIIa-79	2-(cyanomethoxy)-6-methylphenyl
XXVIIa-80	2-(cyanomethoxy)phenyl
XXVIIa-81	2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-82	2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl
XXVIIa-83	2-chloro-3-(prop-2-yn-1-yloxy)phenyl

XXVIIa-84	2-chloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-85	2-chloro-3-formylphenyl
XXVIIa-86	2-chloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-87	2-chloro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-88	2-chloro-4-formylphenyl
XXVIIa-89	2-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-90	2-fluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-91	2-fluoro-3-formylphenyl
XXVIIa-92	2-fluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-93	2-fluoro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-94	2-fluoro-4-formylphenyl
XXVIIa-95	2-formyl-3-methylphenyl
XXVIIa-96	2-formyl-4-methylphenyl
XXVIIa-97	2-formyl-5-methylphenyl
XXVIIa-98	2-formyl-6-methylphenyl
XXVIIa-99	2-formylphenyl
XXVIIa-100	2-methyl-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-101	2-methyl-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-102	2-methyl-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-103	2-methyl-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-104	2-[(hydroxyimino)methyl]-3,4-difluorophenyl
XXVIIa-105	2-[(hydroxyimino)methyl]-3,5-difluorophenyl
XXVIIa-106	2-[(hydroxyimino)methyl]-3,6-difluorophenyl
XXVIIa-107	2-[(hydroxyimino)methyl]-3-chlorophenyl
XXVIIa-108	2-[(hydroxyimino)methyl]-3-fluorophenyl
XXVIIa-109	2-[(hydroxyimino)methyl]-3-methylphenyl
XXVIIa-110	2-[(hydroxyimino)methyl]-4,5-difluorophenyl
XXVIIa-111	2-[(hydroxyimino)methyl]-4,6-difluorophenyl
XXVIIa-112	2-[(hydroxyimino)methyl]-4-chlorophenyl
XXVIIa-113	2-[(hydroxyimino)methyl]-4-fluorophenyl
XXVIIa-114	2-[(hydroxyimino)methyl]-4-methylphenyl
XXVIIa-115	2-[(hydroxyimino)methyl]-5,6-difluorophenyl
XXVIIa-116	2-[(hydroxyimino)methyl]-5-chlorophenyl
XXVIIa-117	2-[(hydroxyimino)methyl]-5-fluorophenyl
XXVIIa-118	2-[(hydroxyimino)methyl]-5-methylphenyl
XXVIIa-119	2-[(hydroxyimino)methyl]-6-chlorophenyl
XXVIIa-120	2-[(hydroxyimino)methyl]-6-fluorophenyl
XXVIIa-121	2-[(hydroxyimino)methyl]-6-methylphenyl

XXVIIa-122	2-[(hydroxyimino)methyl]phenyl
XXVIIa-123	2-[(methoxyimino)methyl]-3,4-difluorophenyl
XXVIIa-124	2-[(methoxyimino)methyl]-3,5-difluorophenyl
XXVIIa-125	2-[(methoxyimino)methyl]-3,6-difluorophenyl
XXVIIa-126	2-[(methoxyimino)methyl]-3-chlorophenyl
XXVIIa-127	2-[(methoxyimino)methyl]-3-fluorophenyl
XXVIIa-128	2-[(methoxyimino)methyl]-3-methylphenyl
XXVIIa-129	2-[(methoxyimino)methyl]-4,5-difluorophenyl
XXVIIa-130	2-[(methoxyimino)methyl]-4,6-difluorophenyl
XXVIIa-131	2-[(methoxyimino)methyl]-4-chlorophenyl
XXVIIa-132	2-[(methoxyimino)methyl]-4-fluorophenyl
XXVIIa-133	2-[(methoxyimino)methyl]-4-methylphenyl
XXVIIa-134	2-[(methoxyimino)methyl]-5,6-difluorophenyl
XXVIIa-135	2-[(methoxyimino)methyl]-5-chlorophenyl
XXVIIa-136	2-[(methoxyimino)methyl]-5-fluorophenyl
XXVIIa-137	2-[(methoxyimino)methyl]-5-methylphenyl
XXVIIa-138	2-[(methoxyimino)methyl]-6-chlorophenyl
XXVIIa-139	2-[(methoxyimino)methyl]-6-fluorophenyl
XXVIIa-140	2-[(methoxyimino)methyl]-6-methylphenyl
XXVIIa-141	2-[(methoxyimino)methyl]phenyl
XXVIIa-142	2-[(methylsulphonyl)oxy]phenyl
XXVIIa-143	2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl
XXVIIa-144	3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-145	3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-146	3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-147	3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-148	3,4-difluoro-2-formylphenyl
XXVIIa-149	3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-150	3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-151	3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-152	3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-153	3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-154	3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-155	3,5-difluoro-2-formylphenyl
XXVIIa-156	3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-157	3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-158	3,5-difluoro-4-formylphenyl
XXVIIa-159	3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl

XXVIIa-160	3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-161	3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-162	3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-163	3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-164	3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-165	3,6-difluoro-2-formylphenyl
XXVIIa-166	3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-167	3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-168	3,6-difluoro-4-formylphenyl
XXVIIa-169	3-(allyloxy)-2,4-dichlorophenyl
XXVIIa-170	3-(allyloxy)-2,4-difluorophenyl
XXVIIa-171	3-(allyloxy)-2,5-dichlorophenyl
XXVIIa-172	3-(allyloxy)-2,5-difluorophenyl
XXVIIa-173	3-(allyloxy)-2,6-dichlorophenyl
XXVIIa-174	3-(allyloxy)-2,6-difluorophenyl
XXVIIa-175	3-(allyloxy)-2-chlorophenyl
XXVIIa-176	3-(allyloxy)-2-fluorophenyl
XXVIIa-177	3-(allyloxy)-2-methylphenyl
XXVIIa-178	3-(allyloxy)-4,5-dichlorophenyl
XXVIIa-179	3-(allyloxy)-4,5-difluorophenyl
XXVIIa-180	3-(allyloxy)-4,6-dichlorophenyl
XXVIIa-181	3-(allyloxy)-4,6-difluorophenyl
XXVIIa-182	3-(allyloxy)-4-chlorophenyl
XXVIIa-183	3-(allyloxy)-4-fluorophenyl
XXVIIa-184	3-(allyloxy)-4-methylphenyl
XXVIIa-185	3-(allyloxy)-5,6-dichlorophenyl
XXVIIa-186	3-(allyloxy)-5,6-difluorophenyl
XXVIIa-187	3-(allyloxy)-5-chlorophenyl
XXVIIa-188	3-(allyloxy)-5-fluorophenyl
XXVIIa-189	3-(allyloxy)-5-methylphenyl
XXVIIa-190	3-(allyloxy)-6-chlorophenyl
XXVIIa-191	3-(allyloxy)-6-fluorophenyl
XXVIIa-192	3-(allyloxy)-6-methylphenyl
XXVIIa-193	3-(allyloxy)phenyl
XXVIIa-194	3-(cyanomethoxy)-2,4-dichlorophenyl
XXVIIa-195	3-(cyanomethoxy)-2,4-difluorophenyl
XXVIIa-196	3-(cyanomethoxy)-2,5-dichlorophenyl
XXVIIa-197	3-(cyanomethoxy)-2,5-difluorophenyl

XXVIIa-198	3-(cyanomethoxy)-2,6-dichlorophenyl
XXVIIa-199	3-(cyanomethoxy)-2,6-difluorophenyl
XXVIIa-200	3-(cyanomethoxy)-2-chlorophenyl
XXVIIa-201	3-(cyanomethoxy)-2-fluorophenyl
XXVIIa-202	3-(cyanomethoxy)-2-methylphenyl
XXVIIa-203	3-(cyanomethoxy)-4,5-dichlorophenyl
XXVIIa-204	3-(cyanomethoxy)-4,5-difluorophenyl
XXVIIa-205	3-(cyanomethoxy)-4,6-dichlorophenyl
XXVIIa-206	3-(cyanomethoxy)-4,6-difluorophenyl
XXVIIa-207	3-(cyanomethoxy)-4-chlorophenyl
XXVIIa-208	3-(cyanomethoxy)-4-fluorophenyl
XXVIIa-209	3-(cyanomethoxy)-4-methylphenyl
XXVIIa-210	3-(cyanomethoxy)-5,6-dichlorophenyl
XXVIIa-211	3-(cyanomethoxy)-5,6-difluorophenyl
XXVIIa-212	3-(cyanomethoxy)-5-chlorophenyl
XXVIIa-213	3-(cyanomethoxy)-5-fluorophenyl
XXVIIa-214	3-(cyanomethoxy)-5-methylphenyl
XXVIIa-215	3-(cyanomethoxy)-6-chlorophenyl
XXVIIa-216	3-(cyanomethoxy)-6-fluorophenyl
XXVIIa-217	3-(cyanomethoxy)-6-methylphenyl
XXVIIa-218	3-(cyanomethoxy)phenyl
XXVIIa-219	3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-220	3-chloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-221	3-chloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-222	3-chloro-2-formylphenyl
XXVIIa-223	3-chloro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-224	3-chloro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-225	3-chloro-4-formylphenyl
XXVIIa-226	3-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-227	3-fluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-228	3-fluoro-2-formylphenyl
XXVIIa-229	3-fluoro-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-230	3-fluoro-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-231	3-fluoro-4-formylphenyl
XXVIIa-232	3-formyl-2-methylphenyl
XXVIIa-233	3-formyl-4-methylphenyl
XXVIIa-234	3-formyl-5-methylphenyl
XXVIIa-235	3-formyl-6-methylphenyl

XXVIIa-236	3-formylphenyl
XXVIIa-237	3-methyl-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-238	3-methyl-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-239	3-methyl-4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-240	3-methyl-4-[(methylsulphonyl)oxy]phenyl
XXVIIa-241	3-[(hydroxyimino)methyl]-2,4-difluorophenyl
XXVIIa-242	3-[(hydroxyimino)methyl]-2,5-difluorophenyl
XXVIIa-243	3-[(hydroxyimino)methyl]-2,6-difluorophenyl
XXVIIa-244	3-[(hydroxyimino)methyl]-2-chlorophenyl
XXVIIa-245	3-[(hydroxyimino)methyl]-2-fluorophenyl
XXVIIa-246	3-[(hydroxyimino)methyl]-2-methylphenyl
XXVIIa-247	3-[(hydroxyimino)methyl]-4,5-difluorophenyl
XXVIIa-248	3-[(hydroxyimino)methyl]-4,6-difluorophenyl
XXVIIa-249	3-[(hydroxyimino)methyl]-4-chlorophenyl
XXVIIa-250	3-[(hydroxyimino)methyl]-4-fluorophenyl
XXVIIa-251	3-[(hydroxyimino)methyl]-4-methylphenyl
XXVIIa-252	3-[(hydroxyimino)methyl]-5,6-difluorophenyl
XXVIIa-253	3-[(hydroxyimino)methyl]-5-chlorophenyl
XXVIIa-254	3-[(hydroxyimino)methyl]-5-fluorophenyl
XXVIIa-255	3-[(hydroxyimino)methyl]-5-methylphenyl
XXVIIa-256	3-[(hydroxyimino)methyl]-6-chlorophenyl
XXVIIa-257	3-[(hydroxyimino)methyl]-6-fluorophenyl
XXVIIa-258	3-[(hydroxyimino)methyl]-6-methylphenyl
XXVIIa-259	3-[(hydroxyimino)methyl]phenyl
XXVIIa-260	3-[(methoxyimino)methyl]-2,4-difluorophenyl
XXVIIa-261	3-[(methoxyimino)methyl]-2,5-difluorophenyl
XXVIIa-262	3-[(methoxyimino)methyl]-2,6-difluorophenyl
XXVIIa-263	3-[(methoxyimino)methyl]-2-chlorophenyl
XXVIIa-264	3-[(methoxyimino)methyl]-2-fluorophenyl
XXVIIa-265	3-[(methoxyimino)methyl]-2-methylphenyl
XXVIIa-266	3-[(methoxyimino)methyl]-4,5-difluorophenyl
XXVIIa-267	3-[(methoxyimino)methyl]-4,6-difluorophenyl
XXVIIa-268	3-[(methoxyimino)methyl]-4-chlorophenyl
XXVIIa-269	3-[(methoxyimino)methyl]-4-fluorophenyl
XXVIIa-270	3-[(methoxyimino)methyl]-4-methylphenyl
XXVIIa-271	3-[(methoxyimino)methyl]-5,6-difluorophenyl
XXVIIa-272	3-[(methoxyimino)methyl]-5-chlorophenyl
XXVIIa-273	3-[(methoxyimino)methyl]-5-fluorophenyl


XXVIIa-274	3-[(methoxyimino)methyl]-5-methylphenyl
XXVIIa-275	3-[(methoxyimino)methyl]-6-chlorophenyl
XXVIIa-276	3-[(methoxyimino)methyl]-6-fluorophenyl
XXVIIa-277	3-[(methoxyimino)methyl]-6-methylphenyl
XXVIIa-278	3-[(methoxyimino)methyl]phenyl
XXVIIa-279	3-[(methylsulphonyl)oxy]phenyl
XXVIIa-280	4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-281	4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-282	4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-283	4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-284	4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-285	4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-286	4,5-difluoro-2-formylphenyl
XXVIIa-287	4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-288	4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-289	4,5-difluoro-3-formylphenyl
XXVIIa-290	4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-291	4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-292	4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-293	4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-294	4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-295	4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-296	4,6-difluoro-2-formylphenyl
XXVIIa-297	4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-298	4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-299	4,6-difluoro-3-formylphenyl
XXVIIa-300	4-(allyloxy)-2,3-dichlorophenyl
XXVIIa-301	4-(allyloxy)-2,3-difluorophenyl
XXVIIa-302	4-(allyloxy)-2,5-dichlorophenyl
XXVIIa-303	4-(allyloxy)-2,5-difluorophenyl
XXVIIa-304	4-(allyloxy)-2,6-dichlorophenyl
XXVIIa-305	4-(allyloxy)-2,6-difluorophenyl
XXVIIa-306	4-(allyloxy)-2-chlorophenyl
XXVIIa-307	4-(allyloxy)-2-fluorophenyl
XXVIIa-308	4-(allyloxy)-2-methylphenyl
XXVIIa-309	4-(allyloxy)-3,5-dichlorophenyl
XXVIIa-310	4-(allyloxy)-3,5-difluorophenyl
XXVIIa-311	4-(allyloxy)-3,6-dichlorophenyl

XXVIIa-312	4-(allyloxy)-3,6-difluorophenyl
XXVIIa-313	4-(allyloxy)-3-chlorophenyl
XXVIIa-314	4-(allyloxy)-3-fluorophenyl
XXVIIa-315	4-(allyloxy)-3-methylphenyl
XXVIIa-316	4-(allyloxy)phenyl
XXVIIa-317	4-(cyanomethoxy)-2,3-dichlorophenyl
XXVIIa-318	4-(cyanomethoxy)-2,3-difluorophenyl
XXVIIa-319	4-(cyanomethoxy)-2,5-dichlorophenyl
XXVIIa-320	4-(cyanomethoxy)-2,5-difluorophenyl
XXVIIa-321	4-(cyanomethoxy)-2,6-dichlorophenyl
XXVIIa-322	4-(cyanomethoxy)-2,6-difluorophenyl
XXVIIa-323	4-(cyanomethoxy)-2-chlorophenyl
XXVIIa-324	4-(cyanomethoxy)-2-fluorophenyl
XXVIIa-325	4-(cyanomethoxy)-2-methylphenyl
XXVIIa-326	4-(cyanomethoxy)-3,5-dichlorophenyl
XXVIIa-327	4-(cyanomethoxy)-3,5-difluorophenyl
XXVIIa-328	4-(cyanomethoxy)-3,6-dichlorophenyl
XXVIIa-329	4-(cyanomethoxy)-3,6-difluorophenyl
XXVIIa-330	4-(cyanomethoxy)-3-chlorophenyl
XXVIIa-331	4-(cyanomethoxy)-3-fluorophenyl
XXVIIa-332	4-(cyanomethoxy)-3-methylphenyl
XXVIIa-333	4-(cyanomethoxy)phenyl
XXVIIa-334	4-(prop-2-yn-1-yloxy)phenyl
XXVIIa-335	4-chloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-336	4-chloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-337	4-chloro-2-formylphenyl
XXVIIa-338	4-chloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-339	4-chloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-340	4-chloro-3-formylphenyl
XXVIIa-341	4-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-342	4-fluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-343	4-fluoro-2-formylphenyl
XXVIIa-344	4-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-345	4-fluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-346	4-fluoro-3-formylphenyl
XXVIIa-347	4-formyl-2-methylphenyl
XXVIIa-348	4-formyl-3-methylphenyl
XXVIIa-349	4-formylphenyl

XXVIIa-350	4-methyl-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-351	4-methyl-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-352	4-methyl-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-353	4-methyl-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-354	4-[(hydroxyimino)methyl]-2,3-difluorophenyl
XXVIIa-355	4-[(hydroxyimino)methyl]-2,5-difluorophenyl
XXVIIa-356	4-[(hydroxyimino)methyl]-2,6-difluorophenyl
XXVIIa-357	4-[(hydroxyimino)methyl]-2-chlorophenyl
XXVIIa-358	4-[(hydroxyimino)methyl]-2-fluorophenyl
XXVIIa-359	4-[(hydroxyimino)methyl]-2-methylphenyl
XXVIIa-360	4-[(hydroxyimino)methyl]-3,5-difluorophenyl
XXVIIa-361	4-[(hydroxyimino)methyl]-3,6-difluorophenyl
XXVIIa-362	4-[(hydroxyimino)methyl]-3-chlorophenyl
XXVIIa-363	4-[(hydroxyimino)methyl]-3-fluorophenyl
XXVIIa-364	4-[(hydroxyimino)methyl]-3-methylphenyl
XXVIIa-365	4-[(hydroxyimino)methyl]phenyl
XXVIIa-366	4-[(methoxyimino)methyl]-2,3-difluorophenyl
XXVIIa-367	4-[(methoxyimino)methyl]-2,5-difluorophenyl
XXVIIa-368	4-[(methoxyimino)methyl]-2,6-difluorophenyl
XXVIIa-369	4-[(methoxyimino)methyl]-2-chlorophenyl
XXVIIa-370	4-[(methoxyimino)methyl]-2-fluorophenyl
XXVIIa-371	4-[(methoxyimino)methyl]-2-methylphenyl
XXVIIa-372	4-[(methoxyimino)methyl]-3,5-difluorophenyl
XXVIIa-373	4-[(methoxyimino)methyl]-3,6-difluorophenyl
XXVIIa-374	4-[(methoxyimino)methyl]-3-chlorophenyl
XXVIIa-375	4-[(methoxyimino)methyl]-3-fluorophenyl
XXVIIa-376	4-[(methoxyimino)methyl]-3-methylphenyl
XXVIIa-377	4-[(methoxyimino)methyl]phenyl
XXVIIa-378	4-[(methylsulphonyl)oxy]phenyl
XXVIIa-379	5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-380	5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-381	5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-382	5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-383	5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-384	5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-385	5,6-difluoro-2-formylphenyl
XXVIIa-386	5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-387	5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl

XXVIIa-388	5,6-difluoro-3-formylphenyl
XXVIIa-389	5-chloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-390	5-chloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-391	5-chloro-2-formylphenyl
XXVIIa-392	5-chloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-393	5-chloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-394	5-chloro-3-formylphenyl
XXVIIa-395	5-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-396	5-fluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-397	5-fluoro-2-formylphenyl
XXVIIa-398	5-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-399	5-fluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-400	5-fluoro-3-formylphenyl
XXVIIa-401	5-methyl-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-402	5-methyl-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-403	5-methyl-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-404	5-methyl-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-405	6-chloro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-406	6-chloro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-407	6-chloro-2-formylphenyl
XXVIIa-408	6-chloro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-409	6-chloro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-410	6-chloro-3-formylphenyl
XXVIIa-411	6-fluoro-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-412	6-fluoro-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-413	6-fluoro-2-formylphenyl
XXVIIa-414	6-fluoro-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-415	6-fluoro-3-[(methylsulphonyl)oxy]phenyl
XXVIIa-416	6-fluoro-3-formylphenyl
XXVIIa-417	6-methyl-2-(prop-2-yn-1-yloxy)phenyl
XXVIIa-418	6-methyl-2-[(methylsulphonyl)oxy]phenyl
XXVIIa-419	6-methyl-3-(prop-2-yn-1-yloxy)phenyl
XXVIIa-420	6-methyl-3-[(methylsulphonyl)oxy]phenyl

Table 4:



Ex.	R ¹
Xc-a-1	2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-2	2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-3	2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-4	2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-5	2,3-difluoro-4-formylphenyl
Xc-a-6	2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-7	2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-8	2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-9	2,4-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-10	2,4-difluoro-3-formylphenyl
Xc-a-11	2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-12	2,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-13	2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-14	2,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-15	2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-16	2,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-17	2,5-difluoro-3-formylphenyl
Xc-a-18	2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-19	2,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-20	2,5-difluoro-4-formylphenyl
Xc-a-21	2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-22	2,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-23	2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-24	2,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-25	2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-26	2,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-27	2,6-difluoro-3-formylphenyl
Xc-a-28	2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-29	2,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-30	2,6-difluoro-4-formylphenyl
Xc-a-31	2-(allyloxy)-3,4-dichlorophenyl
Xc-a-32	2-(allyloxy)-3,4-difluorophenyl
Xc-a-33	2-(allyloxy)-3,5-dichlorophenyl

Xc-a-34	2-(allyloxy)-3,5-difluorophenyl
Xc-a-35	2-(allyloxy)-3,6-dichlorophenyl
Xc-a-36	2-(allyloxy)-3,6-difluorophenyl
Xc-a-37	2-(allyloxy)-3-chlorophenyl
Xc-a-38	2-(allyloxy)-3-fluorophenyl
Xc-a-39	2-(allyloxy)-3-methylphenyl
Xc-a-40	2-(allyloxy)-4,5-dichlorophenyl
Xc-a-41	2-(allyloxy)-4,5-difluorophenyl
Xc-a-42	2-(allyloxy)-4,6-dichlorophenyl
Xc-a-43	2-(allyloxy)-4,6-difluorophenyl
Xc-a-44	2-(allyloxy)-4-chlorophenyl
Xc-a-45	2-(allyloxy)-4-fluorophenyl
Xc-a-46	2-(allyloxy)-4-methylphenyl
Xc-a-47	2-(allyloxy)-5,6-dichlorophenyl
Xc-a-48	2-(allyloxy)-5,6-difluorophenyl
Xc-a-49	2-(allyloxy)-5-chlorophenyl
Xc-a-50	2-(allyloxy)-5-fluorophenyl
Xc-a-51	2-(allyloxy)-5-methylphenyl
Xc-a-52	2-(allyloxy)-6-chlorophenyl
Xc-a-53	2-(allyloxy)-6-fluorophenyl
Xc-a-54	2-(allyloxy)-6-methylphenyl
Xc-a-55	2-(allyloxy)phenyl
Xc-a-56	2-(cyanomethoxy)-3,4-dichlorophenyl
Xc-a-57	2-(cyanomethoxy)-3,4-difluorophenyl
Xc-a-58	2-(cyanomethoxy)-3,5-dichlorophenyl
Xc-a-59	2-(cyanomethoxy)-3,5-difluorophenyl
Xc-a-60	2-(cyanomethoxy)-3,6-dichlorophenyl
Xc-a-61	2-(cyanomethoxy)-3,6-difluorophenyl
Xc-a-62	2-(cyanomethoxy)-3-chlorophenyl
Xc-a-63	2-(cyanomethoxy)-3-fluorophenyl
Xc-a-64	2-(cyanomethoxy)-3-methylphenyl
Xc-a-65	2-(cyanomethoxy)-4,5-dichlorophenyl
Xc-a-66	2-(cyanomethoxy)-4,5-difluorophenyl
Xc-a-67	2-(cyanomethoxy)-4,6-dichlorophenyl
Xc-a-68	2-(cyanomethoxy)-4,6-difluorophenyl
Xc-a-69	2-(cyanomethoxy)-4-chlorophenyl
Xc-a-70	2-(cyanomethoxy)-4-fluorophenyl
Xc-a-71	2-(cyanomethoxy)-4-methylphenyl

Xc-a-72	2-(cyanomethoxy)-5,6-dichlorophenyl
Xc-a-73	2-(cyanomethoxy)-5,6-difluorophenyl
Xc-a-74	2-(cyanomethoxy)-5-chlorophenyl
Xc-a-75	2-(cyanomethoxy)-5-fluorophenyl
Xc-a-76	2-(cyanomethoxy)-5-methylphenyl
Xc-a-77	2-(cyanomethoxy)-6-chlorophenyl
Xc-a-78	2-(cyanomethoxy)-6-fluorophenyl
Xc-a-79	2-(cyanomethoxy)-6-methylphenyl
Xc-a-80	2-(cyanomethoxy)phenyl
Xc-a-81	2-(prop-2-yn-1-yloxy)phenyl
Xc-a-82	2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl
Xc-a-83	2-chloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-84	2-chloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-85	2-chloro-3-formylphenyl
Xc-a-86	2-chloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-87	2-chloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-88	2-chloro-4-formylphenyl
Xc-a-89	2-fluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-90	2-fluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-91	2-fluoro-3-formylphenyl
Xc-a-92	2-fluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-93	2-fluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-94	2-fluoro-4-formylphenyl
Xc-a-95	2-formyl-3-methylphenyl
Xc-a-96	2-formyl-4-methylphenyl
Xc-a-97	2-formyl-5-methylphenyl
Xc-a-98	2-formyl-6-methylphenyl
Xc-a-99	2-formylphenyl
Xc-a-100	2-methyl-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-101	2-methyl-3-[(methylsulphonyl)oxy]phenyl
Xc-a-102	2-methyl-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-103	2-methyl-4-[(methylsulphonyl)oxy]phenyl
Xc-a-104	2-[(hydroxyimino)methyl]-3,4-difluorophenyl
Xc-a-105	2-[(hydroxyimino)methyl]-3,5-difluorophenyl
Xc-a-106	2-[(hydroxyimino)methyl]-3,6-difluorophenyl
Xc-a-107	2-[(hydroxyimino)methyl]-3-chlorophenyl
Xc-a-108	2-[(hydroxyimino)methyl]-3-fluorophenyl
Xc-a-109	2-[(hydroxyimino)methyl]-3-methylphenyl

Xc-a-110	2-[(hydroxyimino)methyl]-4,5-difluorophenyl
Xc-a-111	2-[(hydroxyimino)methyl]-4,6-difluorophenyl
Xc-a-112	2-[(hydroxyimino)methyl]-4-chlorophenyl
Xc-a-113	2-[(hydroxyimino)methyl]-4-fluorophenyl
Xc-a-114	2-[(hydroxyimino)methyl]-4-methylphenyl
Xc-a-115	2-[(hydroxyimino)methyl]-5,6-difluorophenyl
Xc-a-116	2-[(hydroxyimino)methyl]-5-chlorophenyl
Xc-a-117	2-[(hydroxyimino)methyl]-5-fluorophenyl
Xc-a-118	2-[(hydroxyimino)methyl]-5-methylphenyl
Xc-a-119	2-[(hydroxyimino)methyl]-6-chlorophenyl
Xc-a-120	2-[(hydroxyimino)methyl]-6-fluorophenyl
Xc-a-121	2-[(hydroxyimino)methyl]-6-methylphenyl
Xc-a-122	2-[(hydroxyimino)methyl]phenyl
Xc-a-123	2-[(methoxyimino)methyl]-3,4-difluorophenyl
Xc-a-124	2-[(methoxyimino)methyl]-3,5-difluorophenyl
Xc-a-125	2-[(methoxyimino)methyl]-3,6-difluorophenyl
Xc-a-126	2-[(methoxyimino)methyl]-3-chlorophenyl
Xc-a-127	2-[(methoxyimino)methyl]-3-fluorophenyl
Xc-a-128	2-[(methoxyimino)methyl]-3-methylphenyl
Xc-a-129	2-[(methoxyimino)methyl]-4,5-difluorophenyl
Xc-a-130	2-[(methoxyimino)methyl]-4,6-difluorophenyl
Xc-a-131	2-[(methoxyimino)methyl]-4-chlorophenyl
Xc-a-132	2-[(methoxyimino)methyl]-4-fluorophenyl
Xc-a-133	2-[(methoxyimino)methyl]-4-methylphenyl
Xc-a-134	2-[(methoxyimino)methyl]-5,6-difluorophenyl
Xc-a-135	2-[(methoxyimino)methyl]-5-chlorophenyl
Xc-a-136	2-[(methoxyimino)methyl]-5-fluorophenyl
Xc-a-137	2-[(methoxyimino)methyl]-5-methylphenyl
Xc-a-138	2-[(methoxyimino)methyl]-6-chlorophenyl
Xc-a-139	2-[(methoxyimino)methyl]-6-fluorophenyl
Xc-a-140	2-[(methoxyimino)methyl]-6-methylphenyl
Xc-a-141	2-[(methoxyimino)methyl]phenyl
Xc-a-142	2-[(methylsulphonyl)oxy]phenyl
Xc-a-143	2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl
Xc-a-144	3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-145	3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-146	3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-147	3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl

Xc-a-148	3,4-difluoro-2-formylphenyl
Xc-a-149	3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-150	3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-151	3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-152	3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-153	3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-154	3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-155	3,5-difluoro-2-formylphenyl
Xc-a-156	3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-157	3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-158	3,5-difluoro-4-formylphenyl
Xc-a-159	3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-160	3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-161	3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-162	3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-163	3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-164	3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-165	3,6-difluoro-2-formylphenyl
Xc-a-166	3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-167	3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-168	3,6-difluoro-4-formylphenyl
Xc-a-169	3-(allyloxy)-2,4-dichlorophenyl
Xc-a-170	3-(allyloxy)-2,4-difluorophenyl
Xc-a-171	3-(allyloxy)-2,5-dichlorophenyl
Xc-a-172	3-(allyloxy)-2,5-difluorophenyl
Xc-a-173	3-(allyloxy)-2,6-dichlorophenyl
Xc-a-174	3-(allyloxy)-2,6-difluorophenyl
Xc-a-175	3-(allyloxy)-2-chlorophenyl
Xc-a-176	3-(allyloxy)-2-fluorophenyl
Xc-a-177	3-(allyloxy)-2-methylphenyl
Xc-a-178	3-(allyloxy)-4,5-dichlorophenyl
Xc-a-179	3-(allyloxy)-4,5-difluorophenyl
Xc-a-180	3-(allyloxy)-4,6-dichlorophenyl
Xc-a-181	3-(allyloxy)-4,6-difluorophenyl
Xc-a-182	3-(allyloxy)-4-chlorophenyl
Xc-a-183	3-(allyloxy)-4-fluorophenyl
Xc-a-184	3-(allyloxy)-4-methylphenyl
Xc-a-185	3-(allyloxy)-5,6-dichlorophenyl

Xc-a-186	3-(allyloxy)-5,6-difluorophenyl
Xc-a-187	3-(allyloxy)-5-chlorophenyl
Xc-a-188	3-(allyloxy)-5-fluorophenyl
Xc-a-189	3-(allyloxy)-5-methylphenyl
Xc-a-190	3-(allyloxy)-6-chlorophenyl
Xc-a-191	3-(allyloxy)-6-fluorophenyl
Xc-a-192	3-(allyloxy)-6-methylphenyl
Xc-a-193	3-(allyloxy)phenyl
Xc-a-194	3-(cyanomethoxy)-2,4-dichlorophenyl
Xc-a-195	3-(cyanomethoxy)-2,4-difluorophenyl
Xc-a-196	3-(cyanomethoxy)-2,5-dichlorophenyl
Xc-a-197	3-(cyanomethoxy)-2,5-difluorophenyl
Xc-a-198	3-(cyanomethoxy)-2,6-dichlorophenyl
Xc-a-199	3-(cyanomethoxy)-2,6-difluorophenyl
Xc-a-200	3-(cyanomethoxy)-2-chlorophenyl
Xc-a-201	3-(cyanomethoxy)-2-fluorophenyl
Xc-a-202	3-(cyanomethoxy)-2-methylphenyl
Xc-a-203	3-(cyanomethoxy)-4,5-dichlorophenyl
Xc-a-204	3-(cyanomethoxy)-4,5-difluorophenyl
Xc-a-205	3-(cyanomethoxy)-4,6-dichlorophenyl
Xc-a-206	3-(cyanomethoxy)-4,6-difluorophenyl
Xc-a-207	3-(cyanomethoxy)-4-chlorophenyl
Xc-a-208	3-(cyanomethoxy)-4-fluorophenyl
Xc-a-209	3-(cyanomethoxy)-4-methylphenyl
Xc-a-210	3-(cyanomethoxy)-5,6-dichlorophenyl
Xc-a-211	3-(cyanomethoxy)-5,6-difluorophenyl
Xc-a-212	3-(cyanomethoxy)-5-chlorophenyl
Xc-a-213	3-(cyanomethoxy)-5-fluorophenyl
Xc-a-214	3-(cyanomethoxy)-5-methylphenyl
Xc-a-215	3-(cyanomethoxy)-6-chlorophenyl
Xc-a-216	3-(cyanomethoxy)-6-fluorophenyl
Xc-a-217	3-(cyanomethoxy)-6-methylphenyl
Xc-a-218	3-(cyanomethoxy)phenyl
Xc-a-219	3-(prop-2-yn-1-yloxy)phenyl
Xc-a-220	3-chloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-221	3-chloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-222	3-chloro-2-formylphenyl
Xc-a-223	3-chloro-4-(prop-2-yn-1-yloxy)phenyl

Xc-a-224	3-chloro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-225	3-chloro-4-formylphenyl
Xc-a-226	3-fluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-227	3-fluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-228	3-fluoro-2-formylphenyl
Xc-a-229	3-fluoro-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-230	3-fluoro-4-[(methylsulphonyl)oxy]phenyl
Xc-a-231	3-fluoro-4-formylphenyl
Xc-a-232	3-formyl-2-methylphenyl
Xc-a-233	3-formyl-4-methylphenyl
Xc-a-234	3-formyl-5-methylphenyl
Xc-a-235	3-formyl-6-methylphenyl
Xc-a-236	3-formylphenyl
Xc-a-237	3-methyl-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-238	3-methyl-2-[(methylsulphonyl)oxy]phenyl
Xc-a-239	3-methyl-4-(prop-2-yn-1-yloxy)phenyl
Xc-a-240	3-methyl-4-[(methylsulphonyl)oxy]phenyl
Xc-a-241	3-[(hydroxyimino)methyl]-2,4-difluorophenyl
Xc-a-242	3-[(hydroxyimino)methyl]-2,5-difluorophenyl
Xc-a-243	3-[(hydroxyimino)methyl]-2,6-difluorophenyl
Xc-a-244	3-[(hydroxyimino)methyl]-2-chlorophenyl
Xc-a-245	3-[(hydroxyimino)methyl]-2-fluorophenyl
Xc-a-246	3-[(hydroxyimino)methyl]-2-methylphenyl
Xc-a-247	3-[(hydroxyimino)methyl]-4,5-difluorophenyl
Xc-a-248	3-[(hydroxyimino)methyl]-4,6-difluorophenyl
Xc-a-249	3-[(hydroxyimino)methyl]-4-chlorophenyl
Xc-a-250	3-[(hydroxyimino)methyl]-4-fluorophenyl
Xc-a-251	3-[(hydroxyimino)methyl]-4-methylphenyl
Xc-a-252	3-[(hydroxyimino)methyl]-5,6-difluorophenyl
Xc-a-253	3-[(hydroxyimino)methyl]-5-chlorophenyl
Xc-a-254	3-[(hydroxyimino)methyl]-5-fluorophenyl
Xc-a-255	3-[(hydroxyimino)methyl]-5-methylphenyl
Xc-a-256	3-[(hydroxyimino)methyl]-6-chlorophenyl
Xc-a-257	3-[(hydroxyimino)methyl]-6-fluorophenyl
Xc-a-258	3-[(hydroxyimino)methyl]-6-methylphenyl
Xc-a-259	3-[(hydroxyimino)methyl]phenyl
Xc-a-260	3-[(methoxyimino)methyl]-2,4-difluorophenyl
Xc-a-261	3-[(methoxyimino)methyl]-2,5-difluorophenyl


Xc-a-262	3-[(methoxyimino)methyl]-2,6-difluorophenyl
Xc-a-263	3-[(methoxyimino)methyl]-2-chlorophenyl
Xc-a-264	3-[(methoxyimino)methyl]-2-fluorophenyl
Xc-a-265	3-[(methoxyimino)methyl]-2-methylphenyl
Xc-a-266	3-[(methoxyimino)methyl]-4,5-difluorophenyl
Xc-a-267	3-[(methoxyimino)methyl]-4,6-difluorophenyl
Xc-a-268	3-[(methoxyimino)methyl]-4-chlorophenyl
Xc-a-269	3-[(methoxyimino)methyl]-4-fluorophenyl
Xc-a-270	3-[(methoxyimino)methyl]-4-methylphenyl
Xc-a-271	3-[(methoxyimino)methyl]-5,6-difluorophenyl
Xc-a-272	3-[(methoxyimino)methyl]-5-chlorophenyl
Xc-a-273	3-[(methoxyimino)methyl]-5-fluorophenyl
Xc-a-274	3-[(methoxyimino)methyl]-5-methylphenyl
Xc-a-275	3-[(methoxyimino)methyl]-6-chlorophenyl
Xc-a-276	3-[(methoxyimino)methyl]-6-fluorophenyl
Xc-a-277	3-[(methoxyimino)methyl]-6-methylphenyl
Xc-a-278	3-[(methoxyimino)methyl]phenyl
Xc-a-279	3-[(methylsulphonyl)oxy]phenyl
Xc-a-280	4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-281	4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-282	4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-283	4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-284	4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-285	4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-286	4,5-difluoro-2-formylphenyl
Xc-a-287	4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-288	4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-289	4,5-difluoro-3-formylphenyl
Xc-a-290	4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-291	4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-292	4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-293	4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-294	4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-295	4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-296	4,6-difluoro-2-formylphenyl
Xc-a-297	4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-298	4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-299	4,6-difluoro-3-formylphenyl

Xc-a-300	4-(allyloxy)-2,3-dichlorophenyl
Xc-a-301	4-(allyloxy)-2,3-difluorophenyl
Xc-a-302	4-(allyloxy)-2,5-dichlorophenyl
Xc-a-303	4-(allyloxy)-2,5-difluorophenyl
Xc-a-304	4-(allyloxy)-2,6-dichlorophenyl
Xc-a-305	4-(allyloxy)-2,6-difluorophenyl
Xc-a-306	4-(allyloxy)-2-chlorophenyl
Xc-a-307	4-(allyloxy)-2-fluorophenyl
Xc-a-308	4-(allyloxy)-2-methylphenyl
Xc-a-309	4-(allyloxy)-3,5-dichlorophenyl
Xc-a-310	4-(allyloxy)-3,5-difluorophenyl
Xc-a-311	4-(allyloxy)-3,6-dichlorophenyl
Xc-a-312	4-(allyloxy)-3,6-difluorophenyl
Xc-a-313	4-(allyloxy)-3-chlorophenyl
Xc-a-314	4-(allyloxy)-3-fluorophenyl
Xc-a-315	4-(allyloxy)-3-methylphenyl
Xc-a-316	4-(allyloxy)phenyl
Xc-a-317	4-(cyanomethoxy)-2,3-dichlorophenyl
Xc-a-318	4-(cyanomethoxy)-2,3-difluorophenyl
Xc-a-319	4-(cyanomethoxy)-2,5-dichlorophenyl
Xc-a-320	4-(cyanomethoxy)-2,5-difluorophenyl
Xc-a-321	4-(cyanomethoxy)-2,6-dichlorophenyl
Xc-a-322	4-(cyanomethoxy)-2,6-difluorophenyl
Xc-a-323	4-(cyanomethoxy)-2-chlorophenyl
Xc-a-324	4-(cyanomethoxy)-2-fluorophenyl
Xc-a-325	4-(cyanomethoxy)-2-methylphenyl
Xc-a-326	4-(cyanomethoxy)-3,5-dichlorophenyl
Xc-a-327	4-(cyanomethoxy)-3,5-difluorophenyl
Xc-a-328	4-(cyanomethoxy)-3,6-dichlorophenyl
Xc-a-329	4-(cyanomethoxy)-3,6-difluorophenyl
Xc-a-330	4-(cyanomethoxy)-3-chlorophenyl
Xc-a-331	4-(cyanomethoxy)-3-fluorophenyl
Xc-a-332	4-(cyanomethoxy)-3-methylphenyl
Xc-a-333	4-(cyanomethoxy)phenyl
Xc-a-334	4-(prop-2-yn-1-yloxy)phenyl
Xc-a-335	4-chloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-336	4-chloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-337	4-chloro-2-formylphenyl

Xc-a-338	4-chloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-339	4-chloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-340	4-chloro-3-formylphenyl
Xc-a-341	4-fluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-342	4-fluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-343	4-fluoro-2-formylphenyl
Xc-a-344	4-fluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-345	4-fluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-346	4-fluoro-3-formylphenyl
Xc-a-347	4-formyl-2-methylphenyl
Xc-a-348	4-formyl-3-methylphenyl
Xc-a-349	4-formylphenyl
Xc-a-350	4-methyl-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-351	4-methyl-2-[(methylsulphonyl)oxy]phenyl
Xc-a-352	4-methyl-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-353	4-methyl-3-[(methylsulphonyl)oxy]phenyl
Xc-a-354	4-[(hydroxyimino)methyl]-2,3-difluorophenyl
Xc-a-355	4-[(hydroxyimino)methyl]-2,5-difluorophenyl
Xc-a-356	4-[(hydroxyimino)methyl]-2,6-difluorophenyl
Xc-a-357	4-[(hydroxyimino)methyl]-2-chlorophenyl
Xc-a-358	4-[(hydroxyimino)methyl]-2-fluorophenyl
Xc-a-359	4-[(hydroxyimino)methyl]-2-methylphenyl
Xc-a-360	4-[(hydroxyimino)methyl]-3,5-difluorophenyl
Xc-a-361	4-[(hydroxyimino)methyl]-3,6-difluorophenyl
Xc-a-362	4-[(hydroxyimino)methyl]-3-chlorophenyl
Xc-a-363	4-[(hydroxyimino)methyl]-3-fluorophenyl
Xc-a-364	4-[(hydroxyimino)methyl]-3-methylphenyl
Xc-a-365	4-[(hydroxyimino)methyl]phenyl
Xc-a-366	4-[(methoxyimino)methyl]-2,3-difluorophenyl
Xc-a-367	4-[(methoxyimino)methyl]-2,5-difluorophenyl
Xc-a-368	4-[(methoxyimino)methyl]-2,6-difluorophenyl
Xc-a-369	4-[(methoxyimino)methyl]-2-chlorophenyl
Xc-a-370	4-[(methoxyimino)methyl]-2-fluorophenyl
Xc-a-371	4-[(methoxyimino)methyl]-2-methylphenyl
Xc-a-372	4-[(methoxyimino)methyl]-3,5-difluorophenyl
Xc-a-373	4-[(methoxyimino)methyl]-3,6-difluorophenyl
Xc-a-374	4-[(methoxyimino)methyl]-3-chlorophenyl
Xc-a-375	4-[(methoxyimino)methyl]-3-fluorophenyl

Xc-a-376	4-[(methoxyimino)methyl]-3-methylphenyl
Xc-a-377	4-[(methoxyimino)methyl]phenyl
Xc-a-378	4-[(methylsulphonyl)oxy]phenyl
Xc-a-379	5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-380	5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-381	5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-382	5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-383	5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-384	5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-385	5,6-difluoro-2-formylphenyl
Xc-a-386	5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-387	5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-388	5,6-difluoro-3-formylphenyl
Xc-a-389	5-chloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-390	5-chloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-391	5-chloro-2-formylphenyl
Xc-a-392	5-chloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-393	5-chloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-394	5-chloro-3-formylphenyl
Xc-a-395	5-fluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-396	5-fluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-397	5-fluoro-2-formylphenyl
Xc-a-398	5-fluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-399	5-fluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-400	5-fluoro-3-formylphenyl
Xc-a-401	5-methyl-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-402	5-methyl-2-[(methylsulphonyl)oxy]phenyl
Xc-a-403	5-methyl-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-404	5-methyl-3-[(methylsulphonyl)oxy]phenyl
Xc-a-405	6-chloro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-406	6-chloro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-407	6-chloro-2-formylphenyl
Xc-a-408	6-chloro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-409	6-chloro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-410	6-chloro-3-formylphenyl
Xc-a-411	6-fluoro-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-412	6-fluoro-2-[(methylsulphonyl)oxy]phenyl
Xc-a-413	6-fluoro-2-formylphenyl

Xc-a-414	6-fluoro-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-415	6-fluoro-3-[(methylsulphonyl)oxy]phenyl
Xc-a-416	6-fluoro-3-formylphenyl
Xc-a-417	6-methyl-2-(prop-2-yn-1-yloxy)phenyl
Xc-a-418	6-methyl-2-[(methylsulphonyl)oxy]phenyl
Xc-a-419	6-methyl-3-(prop-2-yn-1-yloxy)phenyl
Xc-a-420	6-methyl-3-[(methylsulphonyl)oxy]phenyl

Table 5:**(VIIa-a)**

5

Ex.	R ¹
VIIa-a-1	2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-2	2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-3	2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-4	2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-5	2,3-difluoro-4-formylphenyl
VIIa-a-6	2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-7	2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-8	2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-9	2,4-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-10	2,4-difluoro-3-formylphenyl
VIIa-a-11	2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-12	2,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-13	2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-14	2,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-15	2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-16	2,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-17	2,5-difluoro-3-formylphenyl
VIIa-a-18	2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-19	2,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-20	2,5-difluoro-4-formylphenyl
VIIa-a-21	2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-22	2,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-23	2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl

VIIa-a-24	2,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-25	2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-26	2,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-27	2,6-difluoro-3-formylphenyl
VIIa-a-28	2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-29	2,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-30	2,6-difluoro-4-formylphenyl
VIIa-a-31	2-(allyloxy)-3,4-dichlorophenyl
VIIa-a-32	2-(allyloxy)-3,4-difluorophenyl
VIIa-a-33	2-(allyloxy)-3,5-dichlorophenyl
VIIa-a-34	2-(allyloxy)-3,5-difluorophenyl
VIIa-a-35	2-(allyloxy)-3,6-dichlorophenyl
VIIa-a-36	2-(allyloxy)-3,6-difluorophenyl
VIIa-a-37	2-(allyloxy)-3-chlorophenyl
VIIa-a-38	2-(allyloxy)-3-fluorophenyl
VIIa-a-39	2-(allyloxy)-3-methylphenyl
VIIa-a-40	2-(allyloxy)-4,5-dichlorophenyl
VIIa-a-41	2-(allyloxy)-4,5-difluorophenyl
VIIa-a-42	2-(allyloxy)-4,6-dichlorophenyl
VIIa-a-43	2-(allyloxy)-4,6-difluorophenyl
VIIa-a-44	2-(allyloxy)-4-chlorophenyl
VIIa-a-45	2-(allyloxy)-4-fluorophenyl
VIIa-a-46	2-(allyloxy)-4-methylphenyl
VIIa-a-47	2-(allyloxy)-5,6-dichlorophenyl
VIIa-a-48	2-(allyloxy)-5,6-difluorophenyl
VIIa-a-49	2-(allyloxy)-5-chlorophenyl
VIIa-a-50	2-(allyloxy)-5-fluorophenyl
VIIa-a-51	2-(allyloxy)-5-methylphenyl
VIIa-a-52	2-(allyloxy)-6-chlorophenyl
VIIa-a-53	2-(allyloxy)-6-fluorophenyl
VIIa-a-54	2-(allyloxy)-6-methylphenyl
VIIa-a-55	2-(allyloxy)phenyl
VIIa-a-56	2-(cyanomethoxy)-3,4-dichlorophenyl
VIIa-a-57	2-(cyanomethoxy)-3,4-difluorophenyl
VIIa-a-58	2-(cyanomethoxy)-3,5-dichlorophenyl
VIIa-a-59	2-(cyanomethoxy)-3,5-difluorophenyl
VIIa-a-60	2-(cyanomethoxy)-3,6-dichlorophenyl
VIIa-a-61	2-(cyanomethoxy)-3,6-difluorophenyl

VIIa-a-62	2-(cyanomethoxy)-3-chlorophenyl
VIIa-a-63	2-(cyanomethoxy)-3-fluorophenyl
VIIa-a-64	2-(cyanomethoxy)-3-methylphenyl
VIIa-a-65	2-(cyanomethoxy)-4,5-dichlorophenyl
VIIa-a-66	2-(cyanomethoxy)-4,5-difluorophenyl
VIIa-a-67	2-(cyanomethoxy)-4,6-dichlorophenyl
VIIa-a-68	2-(cyanomethoxy)-4,6-difluorophenyl
VIIa-a-69	2-(cyanomethoxy)-4-chlorophenyl
VIIa-a-70	2-(cyanomethoxy)-4-fluorophenyl
VIIa-a-71	2-(cyanomethoxy)-4-methylphenyl
VIIa-a-72	2-(cyanomethoxy)-5,6-dichlorophenyl
VIIa-a-73	2-(cyanomethoxy)-5,6-difluorophenyl
VIIa-a-74	2-(cyanomethoxy)-5-chlorophenyl
VIIa-a-75	2-(cyanomethoxy)-5-fluorophenyl
VIIa-a-76	2-(cyanomethoxy)-5-methylphenyl
VIIa-a-77	2-(cyanomethoxy)-6-chlorophenyl
VIIa-a-78	2-(cyanomethoxy)-6-fluorophenyl
VIIa-a-79	2-(cyanomethoxy)-6-methylphenyl
VIIa-a-80	2-(cyanomethoxy)phenyl
VIIa-a-81	2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-82	2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl
VIIa-a-83	2-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-84	2-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-85	2-chloro-3-formylphenyl
VIIa-a-86	2-chloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-87	2-chloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-88	2-chloro-4-formylphenyl
VIIa-a-89	2-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-90	2-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-91	2-fluoro-3-formylphenyl
VIIa-a-92	2-fluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-93	2-fluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-94	2-fluoro-4-formylphenyl
VIIa-a-95	2-formyl-3-methylphenyl
VIIa-a-96	2-formyl-4-methylphenyl
VIIa-a-97	2-formyl-5-methylphenyl
VIIa-a-98	2-formyl-6-methylphenyl
VIIa-a-99	2-formylphenyl

VIIa-a-100	2-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-101	2-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-102	2-methyl-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-103	2-methyl-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-104	2-[(hydroxyimino)methyl]-3,4-difluorophenyl
VIIa-a-105	2-[(hydroxyimino)methyl]-3,5-difluorophenyl
VIIa-a-106	2-[(hydroxyimino)methyl]-3,6-difluorophenyl
VIIa-a-107	2-[(hydroxyimino)methyl]-3-chlorophenyl
VIIa-a-08	2-[(hydroxyimino)methyl]-3-fluorophenyl
VIIa-a-109	2-[(hydroxyimino)methyl]-3-methylphenyl
VIIa-a-110	2-[(hydroxyimino)methyl]-4,5-difluorophenyl
VIIa-a-111	2-[(hydroxyimino)methyl]-4,6-difluorophenyl
VIIa-a-112	2-[(hydroxyimino)methyl]-4-chlorophenyl
VIIa-a-113	2-[(hydroxyimino)methyl]-4-fluorophenyl
VIIa-a-114	2-[(hydroxyimino)methyl]-4-methylphenyl
VIIa-a-115	2-[(hydroxyimino)methyl]-5,6-difluorophenyl
VIIa-a-116	2-[(hydroxyimino)methyl]-5-chlorophenyl
VIIa-a-117	2-[(hydroxyimino)methyl]-5-fluorophenyl
VIIa-a-118	2-[(hydroxyimino)methyl]-5-methylphenyl
VIIa-a-119	2-[(hydroxyimino)methyl]-6-chlorophenyl
VIIa-a-120	2-[(hydroxyimino)methyl]-6-fluorophenyl
VIIa-a-121	2-[(hydroxyimino)methyl]-6-methylphenyl
VIIa-a-122	2-[(hydroxyimino)methyl]phenyl
VIIa-a-123	2-[(methoxyimino)methyl]-3,4-difluorophenyl
VIIa-a-124	2-[(methoxyimino)methyl]-3,5-difluorophenyl
VIIa-a-125	2-[(methoxyimino)methyl]-3,6-difluorophenyl
VIIa-a-126	2-[(methoxyimino)methyl]-3-chlorophenyl
VIIa-a-127	2-[(methoxyimino)methyl]-3-fluorophenyl
VIIa-a-128	2-[(methoxyimino)methyl]-3-methylphenyl
VIIa-a-129	2-[(methoxyimino)methyl]-4,5-difluorophenyl
VIIa-a-130	2-[(methoxyimino)methyl]-4,6-difluorophenyl
VIIa-a-131	2-[(methoxyimino)methyl]-4-chlorophenyl
VIIa-a-132	2-[(methoxyimino)methyl]-4-fluorophenyl
VIIa-a-133	2-[(methoxyimino)methyl]-4-methylphenyl
VIIa-a-134	2-[(methoxyimino)methyl]-5,6-difluorophenyl
VIIa-a-135	2-[(methoxyimino)methyl]-5-chlorophenyl
VIIa-a-136	2-[(methoxyimino)methyl]-5-fluorophenyl
VIIa-a-137	2-[(methoxyimino)methyl]-5-methylphenyl

VIIa-a-138	2-[(methoxyimino)methyl]-6-chlorophenyl
VIIa-a-139	2-[(methoxyimino)methyl]-6-fluorophenyl
VIIa-a-140	2-[(methoxyimino)methyl]-6-methylphenyl
VIIa-a-141	2-[(methoxyimino)methyl]phenyl
VIIa-a-142	2-[(methylsulphonyl)oxy]phenyl
VIIa-a-143	2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl
VIIa-a-144	3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-145	3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-146	3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-147	3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-148	3,4-difluoro-2-formylphenyl
VIIa-a-149	3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-150	3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-151	3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-152	3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-153	3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-154	3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-155	3,5-difluoro-2-formylphenyl
VIIa-a-156	3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-157	3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-158	3,5-difluoro-4-formylphenyl
VIIa-a-159	3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-160	3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-161	3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-162	3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-163	3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-164	3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-165	3,6-difluoro-2-formylphenyl
VIIa-a-166	3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-167	3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-168	3,6-difluoro-4-formylphenyl
VIIa-a-169	3-(allyloxy)-2,4-dichlorophenyl
VIIa-a-170	3-(allyloxy)-2,4-difluorophenyl
VIIa-a-171	3-(allyloxy)-2,5-dichlorophenyl
VIIa-a-172	3-(allyloxy)-2,5-difluorophenyl
VIIa-a-173	3-(allyloxy)-2,6-dichlorophenyl
VIIa-a-174	3-(allyloxy)-2,6-difluorophenyl
VIIa-a-175	3-(allyloxy)-2-chlorophenyl

VIIa-a-176	3-(allyloxy)-2-fluorophenyl
VIIa-a-177	3-(allyloxy)-2-methylphenyl
VIIa-a-178	3-(allyloxy)-4,5-dichlorophenyl
VIIa-a-179	3-(allyloxy)-4,5-difluorophenyl
VIIa-a-180	3-(allyloxy)-4,6-dichlorophenyl
VIIa-a-181	3-(allyloxy)-4,6-difluorophenyl
VIIa-a-182	3-(allyloxy)-4-chlorophenyl
VIIa-a-183	3-(allyloxy)-4-fluorophenyl
VIIa-a-184	3-(allyloxy)-4-methylphenyl
VIIa-a-185	3-(allyloxy)-5,6-dichlorophenyl
VIIa-a-186	3-(allyloxy)-5,6-difluorophenyl
VIIa-a-187	3-(allyloxy)-5-chlorophenyl
VIIa-a-188	3-(allyloxy)-5-fluorophenyl
VIIa-a-189	3-(allyloxy)-5-methylphenyl
VIIa-a-190	3-(allyloxy)-6-chlorophenyl
VIIa-a-191	3-(allyloxy)-6-fluorophenyl
VIIa-a-192	3-(allyloxy)-6-methylphenyl
VIIa-a-193	3-(allyloxy)phenyl
VIIa-a-194	3-(cyanomethoxy)-2,4-dichlorophenyl
VIIa-a-195	3-(cyanomethoxy)-2,4-difluorophenyl
VIIa-a-196	3-(cyanomethoxy)-2,5-dichlorophenyl
VIIa-a-197	3-(cyanomethoxy)-2,5-difluorophenyl
VIIa-a-198	3-(cyanomethoxy)-2,6-dichlorophenyl
VIIa-a-199	3-(cyanomethoxy)-2,6-difluorophenyl
VIIa-a-200	3-(cyanomethoxy)-2-chlorophenyl
VIIa-a-201	3-(cyanomethoxy)-2-fluorophenyl
VIIa-a-202	3-(cyanomethoxy)-2-methylphenyl
VIIa-a-203	3-(cyanomethoxy)-4,5-dichlorophenyl
VIIa-a-204	3-(cyanomethoxy)-4,5-difluorophenyl
VIIa-a-205	3-(cyanomethoxy)-4,6-dichlorophenyl
VIIa-a-206	3-(cyanomethoxy)-4,6-difluorophenyl
VIIa-a-207	3-(cyanomethoxy)-4-chlorophenyl
VIIa-a-208	3-(cyanomethoxy)-4-fluorophenyl
VIIa-a-209	3-(cyanomethoxy)-4-methylphenyl
VIIa-a-210	3-(cyanomethoxy)-5,6-dichlorophenyl
VIIa-a-211	3-(cyanomethoxy)-5,6-difluorophenyl
VIIa-a-212	3-(cyanomethoxy)-5-chlorophenyl
VIIa-a-213	3-(cyanomethoxy)-5-fluorophenyl

VIIa-a-214	3-(cyanomethoxy)-5-methylphenyl
VIIa-a-215	3-(cyanomethoxy)-6-chlorophenyl
VIIa-a-216	3-(cyanomethoxy)-6-fluorophenyl
VIIa-a-217	3-(cyanomethoxy)-6-methylphenyl
VIIa-a-218	3-(cyanomethoxy)phenyl
VIIa-a-219	3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-220	3-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-221	3-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-222	3-chloro-2-formylphenyl
VIIa-a-223	3-chloro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-224	3-chloro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-225	3-chloro-4-formylphenyl
VIIa-a-226	3-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-227	3-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-228	3-fluoro-2-formylphenyl
VIIa-a-229	3-fluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-230	3-fluoro-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-231	3-fluoro-4-formylphenyl
VIIa-a-232	3-formyl-2-methylphenyl
VIIa-a-233	3-formyl-4-methylphenyl
VIIa-a-234	3-formyl-5-methylphenyl
VIIa-a-235	3-formyl-6-methylphenyl
VIIa-a-236	3-formylphenyl
VIIa-a-237	3-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-238	3-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-239	3-methyl-4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-240	3-methyl-4-[(methylsulphonyl)oxy]phenyl
VIIa-a-241	3-[(hydroxyimino)methyl]-2,4-difluorophenyl
VIIa-a-242	3-[(hydroxyimino)methyl]-2,5-difluorophenyl
VIIa-a-243	3-[(hydroxyimino)methyl]-2,6-difluorophenyl
VIIa-a-244	3-[(hydroxyimino)methyl]-2-chlorophenyl
VIIa-a-245	3-[(hydroxyimino)methyl]-2-fluorophenyl
VIIa-a-246	3-[(hydroxyimino)methyl]-2-methylphenyl
VIIa-a-247	3-[(hydroxyimino)methyl]-4,5-difluorophenyl
VIIa-a-248	3-[(hydroxyimino)methyl]-4,6-difluorophenyl
VIIa-a-249	3-[(hydroxyimino)methyl]-4-chlorophenyl
VIIa-a-250	3-[(hydroxyimino)methyl]-4-fluorophenyl
VIIa-a-251	3-[(hydroxyimino)methyl]-4-methylphenyl


VIIa-a-252	3-[(hydroxyimino)methyl]-5,6-difluorophenyl
VIIa-a-253	3-[(hydroxyimino)methyl]-5-chlorophenyl
VIIa-a-254	3-[(hydroxyimino)methyl]-5-fluorophenyl
VIIa-a-255	3-[(hydroxyimino)methyl]-5-methylphenyl
VIIa-a-256	3-[(hydroxyimino)methyl]-6-chlorophenyl
VIIa-a-257	3-[(hydroxyimino)methyl]-6-fluorophenyl
VIIa-a-258	3-[(hydroxyimino)methyl]-6-methylphenyl
VIIa-a-259	3-[(hydroxyimino)methyl]phenyl
VIIa-a-260	3-[(methoxyimino)methyl]-2,4-difluorophenyl
VIIa-a-261	3-[(methoxyimino)methyl]-2,5-difluorophenyl
VIIa-a-262	3-[(methoxyimino)methyl]-2,6-difluorophenyl
VIIa-a-263	3-[(methoxyimino)methyl]-2-chlorophenyl
VIIa-a-264	3-[(methoxyimino)methyl]-2-fluorophenyl
VIIa-a-265	3-[(methoxyimino)methyl]-2-methylphenyl
VIIa-a-266	3-[(methoxyimino)methyl]-4,5-difluorophenyl
VIIa-a-267	3-[(methoxyimino)methyl]-4,6-difluorophenyl
VIIa-a-268	3-[(methoxyimino)methyl]-4-chlorophenyl
VIIa-a-269	3-[(methoxyimino)methyl]-4-fluorophenyl
VIIa-a-270	3-[(methoxyimino)methyl]-4-methylphenyl
VIIa-a-271	3-[(methoxyimino)methyl]-5,6-difluorophenyl
VIIa-a-272	3-[(methoxyimino)methyl]-5-chlorophenyl
VIIa-a-273	3-[(methoxyimino)methyl]-5-fluorophenyl
VIIa-a-274	3-[(methoxyimino)methyl]-5-methylphenyl
VIIa-a-275	3-[(methoxyimino)methyl]-6-chlorophenyl
VIIa-a-276	3-[(methoxyimino)methyl]-6-fluorophenyl
VIIa-a-277	3-[(methoxyimino)methyl]-6-methylphenyl
VIIa-a-278	3-[(methoxyimino)methyl]phenyl
VIIa-a-279	3-[(methylsulphonyl)oxy]phenyl
VIIa-a-280	4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-281	4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-282	4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-283	4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-284	4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-285	4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-286	4,5-difluoro-2-formylphenyl
VIIa-a-287	4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-288	4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-289	4,5-difluoro-3-formylphenyl

VIIa-a-290	4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-291	4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-292	4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-293	4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-294	4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-295	4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-296	4,6-difluoro-2-formylphenyl
VIIa-a-297	4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-298	4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-299	4,6-difluoro-3-formylphenyl
VIIa-a-300	4-(allyloxy)-2,3-dichlorophenyl
VIIa-a-301	4-(allyloxy)-2,3-difluorophenyl
VIIa-a-302	4-(allyloxy)-2,5-dichlorophenyl
VIIa-a-303	4-(allyloxy)-2,5-difluorophenyl
VIIa-a-304	4-(allyloxy)-2,6-dichlorophenyl
VIIa-a-305	4-(allyloxy)-2,6-difluorophenyl
VIIa-a-306	4-(allyloxy)-2-chlorophenyl
VIIa-a-307	4-(allyloxy)-2-fluorophenyl
VIIa-a-308	4-(allyloxy)-2-methylphenyl
VIIa-a-309	4-(allyloxy)-3,5-dichlorophenyl
VIIa-a-310	4-(allyloxy)-3,5-difluorophenyl
VIIa-a-311	4-(allyloxy)-3,6-dichlorophenyl
VIIa-a-312	4-(allyloxy)-3,6-difluorophenyl
VIIa-a-313	4-(allyloxy)-3-chlorophenyl
VIIa-a-314	4-(allyloxy)-3-fluorophenyl
VIIa-a-315	4-(allyloxy)-3-methylphenyl
VIIa-a-316	4-(allyloxy)phenyl
VIIa-a-317	4-(cyanomethoxy)-2,3-dichlorophenyl
VIIa-a-318	4-(cyanomethoxy)-2,3-difluorophenyl
VIIa-a-319	4-(cyanomethoxy)-2,5-dichlorophenyl
VIIa-a-320	4-(cyanomethoxy)-2,5-difluorophenyl
VIIa-a-321	4-(cyanomethoxy)-2,6-dichlorophenyl
VIIa-a-322	4-(cyanomethoxy)-2,6-difluorophenyl
VIIa-a-323	4-(cyanomethoxy)-2-chlorophenyl
VIIa-a-324	4-(cyanomethoxy)-2-fluorophenyl
VIIa-a-325	4-(cyanomethoxy)-2-methylphenyl
VIIa-a-326	4-(cyanomethoxy)-3,5-dichlorophenyl
VIIa-a-327	4-(cyanomethoxy)-3,5-difluorophenyl

VIIa-a-328	4-(cyanomethoxy)-3,6-dichlorophenyl
VIIa-a-329	4-(cyanomethoxy)-3,6-difluorophenyl
VIIa-a-330	4-(cyanomethoxy)-3-chlorophenyl
VIIa-a-331	4-(cyanomethoxy)-3-fluorophenyl
VIIa-a-332	4-(cyanomethoxy)-3-methylphenyl
VIIa-a-333	4-(cyanomethoxy)phenyl
VIIa-a-334	4-(prop-2-yn-1-yloxy)phenyl
VIIa-a-335	4-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-336	4-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-337	4-chloro-2-formylphenyl
VIIa-a-338	4-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-339	4-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-340	4-chloro-3-formylphenyl
VIIa-a-341	4-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-342	4-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-343	4-fluoro-2-formylphenyl
VIIa-a-344	4-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-345	4-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-346	4-fluoro-3-formylphenyl
VIIa-a-347	4-formyl-2-methylphenyl
VIIa-a-348	4-formyl-3-methylphenyl
VIIa-a-349	4-formylphenyl
VIIa-a-350	4-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-351	4-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-352	4-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-353	4-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-354	4-[(hydroxyimino)methyl]-2,3-difluorophenyl
VIIa-a-355	4-[(hydroxyimino)methyl]-2,5-difluorophenyl
VIIa-a-356	4-[(hydroxyimino)methyl]-2,6-difluorophenyl
VIIa-a-357	4-[(hydroxyimino)methyl]-2-chlorophenyl
VIIa-a-358	4-[(hydroxyimino)methyl]-2-fluorophenyl
VIIa-a-359	4-[(hydroxyimino)methyl]-2-methylphenyl
VIIa-a-360	4-[(hydroxyimino)methyl]-3,5-difluorophenyl
VIIa-a-361	4-[(hydroxyimino)methyl]-3,6-difluorophenyl
VIIa-a-362	4-[(hydroxyimino)methyl]-3-chlorophenyl
VIIa-a-363	4-[(hydroxyimino)methyl]-3-fluorophenyl
VIIa-a-364	4-[(hydroxyimino)methyl]-3-methylphenyl
VIIa-a-365	4-[(hydroxyimino)methyl]phenyl

VIIa-a-366	4-[(methoxyimino)methyl]-2,3-difluorophenyl
VIIa-a-367	4-[(methoxyimino)methyl]-2,5-difluorophenyl
VIIa-a-368	4-[(methoxyimino)methyl]-2,6-difluorophenyl
VIIa-a-369	4-[(methoxyimino)methyl]-2-chlorophenyl
VIIa-a-370	4-[(methoxyimino)methyl]-2-fluorophenyl
VIIa-a-371	4-[(methoxyimino)methyl]-2-methylphenyl
VIIa-a-372	4-[(methoxyimino)methyl]-3,5-difluorophenyl
VIIa-a-373	4-[(methoxyimino)methyl]-3,6-difluorophenyl
VIIa-a-374	4-[(methoxyimino)methyl]-3-chlorophenyl
VIIa-a-375	4-[(methoxyimino)methyl]-3-fluorophenyl
VIIa-a-376	4-[(methoxyimino)methyl]-3-methylphenyl
VIIa-a-377	4-[(methoxyimino)methyl]phenyl
VIIa-a-378	4-[(methylsulphonyl)oxy]phenyl
VIIa-a-379	5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-380	5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-381	5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-382	5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-383	5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-384	5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-385	5,6-difluoro-2-formylphenyl
VIIa-a-386	5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-387	5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-388	5,6-difluoro-3-formylphenyl
VIIa-a-389	5-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-390	5-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-391	5-chloro-2-formylphenyl
VIIa-a-392	5-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-393	5-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-394	5-chloro-3-formylphenyl
VIIa-a-395	5-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-396	5-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-397	5-fluoro-2-formylphenyl
VIIa-a-398	5-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-399	5-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-400	5-fluoro-3-formylphenyl
VIIa-a-401	5-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-402	5-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-403	5-methyl-3-(prop-2-yn-1-yloxy)phenyl

VIIa-a-404	5-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-405	6-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-406	6-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-407	6-chloro-2-formylphenyl
VIIa-a-408	6-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-409	6-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-410	6-chloro-3-formylphenyl
VIIa-a-411	6-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-412	6-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-413	6-fluoro-2-formylphenyl
VIIa-a-414	6-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-415	6-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIa-a-416	6-fluoro-3-formylphenyl
VIIa-a-417	6-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIa-a-418	6-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIa-a-419	6-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIa-a-420	6-methyl-3-[(methylsulphonyl)oxy]phenyl

Table 6:**(VIIIa-a)**

5

Ex.	R ¹
VIIIa-a-1	2,3-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-2	2,3-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-3	2,3-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-4	2,3-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-5	2,3-difluoro-4-formylphenyl
VIIIa-a-6	2,4-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-7	2,4-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-8	2,4-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-9	2,4-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-10	2,4-difluoro-3-formylphenyl
VIIIa-a-11	2,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-12	2,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-13	2,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl

VIIIa-a-14	2,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-15	2,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-16	2,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-17	2,5-difluoro-3-formylphenyl
VIIIa-a-18	2,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-19	2,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-20	2,5-difluoro-4-formylphenyl
VIIIa-a-21	2,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-22	2,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-23	2,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-24	2,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-25	2,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-26	2,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-27	2,6-difluoro-3-formylphenyl
VIIIa-a-28	2,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-29	2,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-30	2,6-difluoro-4-formylphenyl
VIIIa-a-31	2-(allyloxy)-3,4-dichlorophenyl
VIIIa-a-32	2-(allyloxy)-3,4-difluorophenyl
VIIIa-a-33	2-(allyloxy)-3,5-dichlorophenyl
VIIIa-a-34	2-(allyloxy)-3,5-difluorophenyl
VIIIa-a-35	2-(allyloxy)-3,6-dichlorophenyl
VIIIa-a-36	2-(allyloxy)-3,6-difluorophenyl
VIIIa-a-37	2-(allyloxy)-3-chlorophenyl
VIIIa-a-38	2-(allyloxy)-3-fluorophenyl
VIIIa-a-39	2-(allyloxy)-3-methylphenyl
VIIIa-a-40	2-(allyloxy)-4,5-dichlorophenyl
VIIIa-a-41	2-(allyloxy)-4,5-difluorophenyl
VIIIa-a-42	2-(allyloxy)-4,6-dichlorophenyl
VIIIa-a-43	2-(allyloxy)-4,6-difluorophenyl
VIIIa-a-44	2-(allyloxy)-4-chlorophenyl
VIIIa-a-45	2-(allyloxy)-4-fluorophenyl
VIIIa-a-46	2-(allyloxy)-4-methylphenyl
VIIIa-a-47	2-(allyloxy)-5,6-dichlorophenyl
VIIIa-a-48	2-(allyloxy)-5,6-difluorophenyl
VIIIa-a-49	2-(allyloxy)-5-chlorophenyl
VIIIa-a-50	2-(allyloxy)-5-fluorophenyl
VIIIa-a-51	2-(allyloxy)-5-methylphenyl

VIIIa-a-52	2-(allyloxy)-6-chlorophenyl
VIIIa-a-53	2-(allyloxy)-6-fluorophenyl
VIIIa-a-54	2-(allyloxy)-6-methylphenyl
VIIIa-a-55	2-(allyloxy)phenyl
VIIIa-a-56	2-(cyanomethoxy)-3,4-dichlorophenyl
VIIIa-a-57	2-(cyanomethoxy)-3,4-difluorophenyl
VIIIa-a-58	2-(cyanomethoxy)-3,5-dichlorophenyl
VIIIa-a-59	2-(cyanomethoxy)-3,5-difluorophenyl
VIIIa-a-60	2-(cyanomethoxy)-3,6-dichlorophenyl
VIIIa-a-61	2-(cyanomethoxy)-3,6-difluorophenyl
VIIIa-a-62	2-(cyanomethoxy)-3-chlorophenyl
VIIIa-a-63	2-(cyanomethoxy)-3-fluorophenyl
VIIIa-a-64	2-(cyanomethoxy)-3-methylphenyl
VIIIa-a-65	2-(cyanomethoxy)-4,5-dichlorophenyl
VIIIa-a-66	2-(cyanomethoxy)-4,5-difluorophenyl
VIIIa-a-67	2-(cyanomethoxy)-4,6-dichlorophenyl
VIIIa-a-68	2-(cyanomethoxy)-4,6-difluorophenyl
VIIIa-a-69	2-(cyanomethoxy)-4-chlorophenyl
VIIIa-a-70	2-(cyanomethoxy)-4-fluorophenyl
VIIIa-a-71	2-(cyanomethoxy)-4-methylphenyl
VIIIa-a-72	2-(cyanomethoxy)-5,6-dichlorophenyl
VIIIa-a-73	2-(cyanomethoxy)-5,6-difluorophenyl
VIIIa-a-74	2-(cyanomethoxy)-5-chlorophenyl
VIIIa-a-75	2-(cyanomethoxy)-5-fluorophenyl
VIIIa-a-76	2-(cyanomethoxy)-5-methylphenyl
VIIIa-a-77	2-(cyanomethoxy)-6-chlorophenyl
VIIIa-a-78	2-(cyanomethoxy)-6-fluorophenyl
VIIIa-a-79	2-(cyanomethoxy)-6-methylphenyl
VIIIa-a-80	2-(cyanomethoxy)phenyl
VIIIa-a-81	2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-82	2-(prop-2-yn-1-yloxy)-4-(trifluoromethyl)phenyl
VIIIa-a-83	2-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-84	2-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-85	2-chloro-3-formylphenyl
VIIIa-a-86	2-chloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-87	2-chloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-88	2-chloro-4-formylphenyl
VIIIa-a-89	2-fluoro-3-(prop-2-yn-1-yloxy)phenyl

VIIIa-a-90	2-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-91	2-fluoro-3-formylphenyl
VIIIa-a-92	2-fluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-93	2-fluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-94	2-fluoro-4-formylphenyl
VIIIa-a-95	2-formyl-3-methylphenyl
VIIIa-a-96	2-formyl-4-methylphenyl
VIIIa-a-97	2-formyl-5-methylphenyl
VIIIa-a-98	2-formyl-6-methylphenyl
VIIIa-a-99	2-formylphenyl
VIIIa-a-100	2-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-101	2-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-102	2-methyl-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-103	2-methyl-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-104	2-[(hydroxyimino)methyl]-3,4-difluorophenyl
VIIIa-a-105	2-[(hydroxyimino)methyl]-3,5-difluorophenyl
VIIIa-a-106	2-[(hydroxyimino)methyl]-3,6-difluorophenyl
VIIIa-a-107	2-[(hydroxyimino)methyl]-3-chlorophenyl
VIIIa-a-108	2-[(hydroxyimino)methyl]-3-fluorophenyl
VIIIa-a-109	2-[(hydroxyimino)methyl]-3-methylphenyl
VIIIa-a-110	2-[(hydroxyimino)methyl]-4,5-difluorophenyl
VIIIa-a-111	2-[(hydroxyimino)methyl]-4,6-difluorophenyl
VIIIa-a-112	2-[(hydroxyimino)methyl]-4-chlorophenyl
VIIIa-a-113	2-[(hydroxyimino)methyl]-4-fluorophenyl
VIIIa-a-114	2-[(hydroxyimino)methyl]-4-methylphenyl
VIIIa-a-115	2-[(hydroxyimino)methyl]-5,6-difluorophenyl
VIIIa-a-116	2-[(hydroxyimino)methyl]-5-chlorophenyl
VIIIa-a-117	2-[(hydroxyimino)methyl]-5-fluorophenyl
VIIIa-a-118	2-[(hydroxyimino)methyl]-5-methylphenyl
VIIIa-a-119	2-[(hydroxyimino)methyl]-6-chlorophenyl
VIIIa-a-120	2-[(hydroxyimino)methyl]-6-fluorophenyl
VIIIa-a-121	2-[(hydroxyimino)methyl]-6-methylphenyl
VIIIa-a-122	2-[(hydroxyimino)methyl]phenyl
VIIIa-a-123	2-[(methoxyimino)methyl]-3,4-difluorophenyl
VIIIa-a-124	2-[(methoxyimino)methyl]-3,5-difluorophenyl
VIIIa-a-125	2-[(methoxyimino)methyl]-3,6-difluorophenyl
VIIIa-a-126	2-[(methoxyimino)methyl]-3-chlorophenyl
VIIIa-a-127	2-[(methoxyimino)methyl]-3-fluorophenyl

VIIIa-a-128	2-[(methoxyimino)methyl]-3-methylphenyl
VIIIa-a-129	2-[(methoxyimino)methyl]-4,5-difluorophenyl
VIIIa-a-130	2-[(methoxyimino)methyl]-4,6-difluorophenyl
VIIIa-a-131	2-[(methoxyimino)methyl]-4-chlorophenyl
VIIIa-a-132	2-[(methoxyimino)methyl]-4-fluorophenyl
VIIIa-a-133	2-[(methoxyimino)methyl]-4-methylphenyl
VIIIa-a-134	2-[(methoxyimino)methyl]-5,6-difluorophenyl
VIIIa-a-135	2-[(methoxyimino)methyl]-5-chlorophenyl
VIIIa-a-136	2-[(methoxyimino)methyl]-5-fluorophenyl
VIIIa-a-137	2-[(methoxyimino)methyl]-5-methylphenyl
VIIIa-a-138	2-[(methoxyimino)methyl]-6-chlorophenyl
VIIIa-a-139	2-[(methoxyimino)methyl]-6-fluorophenyl
VIIIa-a-140	2-[(methoxyimino)methyl]-6-methylphenyl
VIIIa-a-141	2-[(methoxyimino)methyl]phenyl
VIIIa-a-142	2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-143	2-[(methylsulphonyl)oxy]-4-(trifluoromethyl)phenyl
VIIIa-a-144	3,4-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-145	3,4-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-146	3,4-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-147	3,4-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-148	3,4-difluoro-2-formylphenyl
VIIIa-a-149	3,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-150	3,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-151	3,5-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-152	3,5-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-153	3,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-154	3,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-155	3,5-difluoro-2-formylphenyl
VIIIa-a-156	3,5-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-157	3,5-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-158	3,5-difluoro-4-formylphenyl
VIIIa-a-159	3,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-160	3,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-161	3,6-dichloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-162	3,6-dichloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-163	3,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-164	3,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-165	3,6-difluoro-2-formylphenyl

VIIIa-a-166	3,6-difluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-167	3,6-difluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-168	3,6-difluoro-4-formylphenyl
VIIIa-a-169	3-(allyloxy)-2,4-dichlorophenyl
VIIIa-a-170	3-(allyloxy)-2,4-difluorophenyl
VIIIa-a-171	3-(allyloxy)-2,5-dichlorophenyl
VIIIa-a-172	3-(allyloxy)-2,5-difluorophenyl
VIIIa-a-173	3-(allyloxy)-2,6-dichlorophenyl
VIIIa-a-174	3-(allyloxy)-2,6-difluorophenyl
VIIIa-a-175	3-(allyloxy)-2-chlorophenyl
VIIIa-a-176	3-(allyloxy)-2-fluorophenyl
VIIIa-a-177	3-(allyloxy)-2-methylphenyl
VIIIa-a-178	3-(allyloxy)-4,5-dichlorophenyl
VIIIa-a-179	3-(allyloxy)-4,5-difluorophenyl
VIIIa-a-180	3-(allyloxy)-4,6-dichlorophenyl
VIIIa-a-181	3-(allyloxy)-4,6-difluorophenyl
VIIIa-a-182	3-(allyloxy)-4-chlorophenyl
VIIIa-a-183	3-(allyloxy)-4-fluorophenyl
VIIIa-a-184	3-(allyloxy)-4-methylphenyl
VIIIa-a-185	3-(allyloxy)-5,6-dichlorophenyl
VIIIa-a-186	3-(allyloxy)-5,6-difluorophenyl
VIIIa-a-187	3-(allyloxy)-5-chlorophenyl
VIIIa-a-188	3-(allyloxy)-5-fluorophenyl
VIIIa-a-189	3-(allyloxy)-5-methylphenyl
VIIIa-a-190	3-(allyloxy)-6-chlorophenyl
VIIIa-a-191	3-(allyloxy)-6-fluorophenyl
VIIIa-a-192	3-(allyloxy)-6-methylphenyl
VIIIa-a-193	3-(allyloxy)phenyl
VIIIa-a-194	3-(cyanomethoxy)-2,4-dichlorophenyl
VIIIa-a-195	3-(cyanomethoxy)-2,4-difluorophenyl
VIIIa-a-196	3-(cyanomethoxy)-2,5-dichlorophenyl
VIIIa-a-197	3-(cyanomethoxy)-2,5-difluorophenyl
VIIIa-a-198	3-(cyanomethoxy)-2,6-dichlorophenyl
VIIIa-a-199	3-(cyanomethoxy)-2,6-difluorophenyl
VIIIa-a-200	3-(cyanomethoxy)-2-chlorophenyl
VIIIa-a-201	3-(cyanomethoxy)-2-fluorophenyl
VIIIa-a-202	3-(cyanomethoxy)-2-methylphenyl
VIIIa-a-203	3-(cyanomethoxy)-4,5-dichlorophenyl

VIIIa-a-204	3-(cyanomethoxy)-4,5-difluorophenyl
VIIIa-a-205	3-(cyanomethoxy)-4,6-dichlorophenyl
VIIIa-a-206	3-(cyanomethoxy)-4,6-difluorophenyl
VIIIa-a-207	3-(cyanomethoxy)-4-chlorophenyl
VIIIa-a-208	3-(cyanomethoxy)-4-fluorophenyl
VIIIa-a-209	3-(cyanomethoxy)-4-methylphenyl
VIIIa-a-210	3-(cyanomethoxy)-5,6-dichlorophenyl
VIIIa-a-211	3-(cyanomethoxy)-5,6-difluorophenyl
VIIIa-a-212	3-(cyanomethoxy)-5-chlorophenyl
VIIIa-a-213	3-(cyanomethoxy)-5-fluorophenyl
VIIIa-a-214	3-(cyanomethoxy)-5-methylphenyl
VIIIa-a-215	3-(cyanomethoxy)-6-chlorophenyl
VIIIa-a-216	3-(cyanomethoxy)-6-fluorophenyl
VIIIa-a-217	3-(cyanomethoxy)-6-methylphenyl
VIIIa-a-218	3-(cyanomethoxy)phenyl
VIIIa-a-219	3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-220	3-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-221	3-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-222	3-chloro-2-formylphenyl
VIIIa-a-223	3-chloro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-224	3-chloro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-225	3-chloro-4-formylphenyl
VIIIa-a-226	3-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-227	3-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-228	3-fluoro-2-formylphenyl
VIIIa-a-229	3-fluoro-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-230	3-fluoro-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-231	3-fluoro-4-formylphenyl
VIIIa-a-232	3-formyl-2-methylphenyl
VIIIa-a-233	3-formyl-4-methylphenyl
VIIIa-a-234	3-formyl-5-methylphenyl
VIIIa-a-235	3-formyl-6-methylphenyl
VIIIa-a-236	3-formylphenyl
VIIIa-a-237	3-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-238	3-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-239	3-methyl-4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-240	3-methyl-4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-241	3-[(hydroxyimino)methyl]-2,4-difluorophenyl

VIIIa-a-242	3-[(hydroxyimino)methyl]-2,5-difluorophenyl
VIIIa-a-243	3-[(hydroxyimino)methyl]-2,6-difluorophenyl
VIIIa-a-244	3-[(hydroxyimino)methyl]-2-chlorophenyl
VIIIa-a-245	3-[(hydroxyimino)methyl]-2-fluorophenyl
VIIIa-a-246	3-[(hydroxyimino)methyl]-2-methylphenyl
VIIIa-a-247	3-[(hydroxyimino)methyl]-4,5-difluorophenyl
VIIIa-a-248	3-[(hydroxyimino)methyl]-4,6-difluorophenyl
VIIIa-a-249	3-[(hydroxyimino)methyl]-4-chlorophenyl
VIIIa-a-250	3-[(hydroxyimino)methyl]-4-fluorophenyl
VIIIa-a-251	3-[(hydroxyimino)methyl]-4-methylphenyl
VIIIa-a-252	3-[(hydroxyimino)methyl]-5,6-difluorophenyl
VIIIa-a-253	3-[(hydroxyimino)methyl]-5-chlorophenyl
VIIIa-a-254	3-[(hydroxyimino)methyl]-5-fluorophenyl
VIIIa-a-255	3-[(hydroxyimino)methyl]-5-methylphenyl
VIIIa-a-256	3-[(hydroxyimino)methyl]-6-chlorophenyl
VIIIa-a-257	3-[(hydroxyimino)methyl]-6-fluorophenyl
VIIIa-a-258	3-[(hydroxyimino)methyl]-6-methylphenyl
VIIIa-a-259	3-[(hydroxyimino)methyl]phenyl
VIIIa-a-260	3-[(methoxyimino)methyl]-2,4-difluorophenyl
VIIIa-a-261	3-[(methoxyimino)methyl]-2,5-difluorophenyl
VIIIa-a-262	3-[(methoxyimino)methyl]-2,6-difluorophenyl
VIIIa-a-263	3-[(methoxyimino)methyl]-2-chlorophenyl
VIIIa-a-264	3-[(methoxyimino)methyl]-2-fluorophenyl
VIIIa-a-265	3-[(methoxyimino)methyl]-2-methylphenyl
VIIIa-a-266	3-[(methoxyimino)methyl]-4,5-difluorophenyl
VIIIa-a-267	3-[(methoxyimino)methyl]-4,6-difluorophenyl
VIIIa-a-268	3-[(methoxyimino)methyl]-4-chlorophenyl
VIIIa-a-269	3-[(methoxyimino)methyl]-4-fluorophenyl
VIIIa-a-270	3-[(methoxyimino)methyl]-4-methylphenyl
VIIIa-a-271	3-[(methoxyimino)methyl]-5,6-difluorophenyl
VIIIa-a-272	3-[(methoxyimino)methyl]-5-chlorophenyl
VIIIa-a-273	3-[(methoxyimino)methyl]-5-fluorophenyl
VIIIa-a-274	3-[(methoxyimino)methyl]-5-methylphenyl
VIIIa-a-275	3-[(methoxyimino)methyl]-6-chlorophenyl
VIIIa-a-276	3-[(methoxyimino)methyl]-6-fluorophenyl
VIIIa-a-277	3-[(methoxyimino)methyl]-6-methylphenyl
VIIIa-a-278	3-[(methoxyimino)methyl]phenyl
VIIIa-a-279	3-[(methylsulphonyl)oxy]phenyl

VIIIa-a-280	4,5-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-281	4,5-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-282	4,5-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-283	4,5-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-284	4,5-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-285	4,5-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-286	4,5-difluoro-2-formylphenyl
VIIIa-a-287	4,5-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-288	4,5-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-289	4,5-difluoro-3-formylphenyl
VIIIa-a-290	4,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-291	4,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-292	4,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-293	4,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-294	4,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-295	4,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-296	4,6-difluoro-2-formylphenyl
VIIIa-a-297	4,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-298	4,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-299	4,6-difluoro-3-formylphenyl
VIIIa-a-300	4-(allyloxy)-2,3-dichlorophenyl
VIIIa-a-301	4-(allyloxy)-2,3-difluorophenyl
VIIIa-a-302	4-(allyloxy)-2,5-dichlorophenyl
VIIIa-a-303	4-(allyloxy)-2,5-difluorophenyl
VIIIa-a-304	4-(allyloxy)-2,6-dichlorophenyl
VIIIa-a-305	4-(allyloxy)-2,6-difluorophenyl
VIIIa-a-306	4-(allyloxy)-2-chlorophenyl
VIIIa-a-307	4-(allyloxy)-2-fluorophenyl
VIIIa-a-308	4-(allyloxy)-2-methylphenyl
VIIIa-a-309	4-(allyloxy)-3,5-dichlorophenyl
VIIIa-a-310	4-(allyloxy)-3,5-difluorophenyl
VIIIa-a-311	4-(allyloxy)-3,6-dichlorophenyl
VIIIa-a-312	4-(allyloxy)-3,6-difluorophenyl
VIIIa-a-313	4-(allyloxy)-3-chlorophenyl
VIIIa-a-314	4-(allyloxy)-3-fluorophenyl
VIIIa-a-315	4-(allyloxy)-3-methylphenyl
VIIIa-a-316	4-(allyloxy)phenyl
VIIIa-a-317	4-(cyanomethoxy)-2,3-dichlorophenyl

VIIIa-a-318	4-(cyanomethoxy)-2,3-difluorophenyl
VIIIa-a-319	4-(cyanomethoxy)-2,5-dichlorophenyl
VIIIa-a-320	4-(cyanomethoxy)-2,5-difluorophenyl
VIIIa-a-321	4-(cyanomethoxy)-2,6-dichlorophenyl
VIIIa-a-322	4-(cyanomethoxy)-2,6-difluorophenyl
VIIIa-a-323	4-(cyanomethoxy)-2-chlorophenyl
VIIIa-a-324	4-(cyanomethoxy)-2-fluorophenyl
VIIIa-a-325	4-(cyanomethoxy)-2-methylphenyl
VIIIa-a-326	4-(cyanomethoxy)-3,5-dichlorophenyl
VIIIa-a-327	4-(cyanomethoxy)-3,5-difluorophenyl
VIIIa-a-328	4-(cyanomethoxy)-3,6-dichlorophenyl
VIIIa-a-329	4-(cyanomethoxy)-3,6-difluorophenyl
VIIIa-a-330	4-(cyanomethoxy)-3-chlorophenyl
VIIIa-a-331	4-(cyanomethoxy)-3-fluorophenyl
VIIIa-a-332	4-(cyanomethoxy)-3-methylphenyl
VIIIa-a-333	4-(cyanomethoxy)phenyl
VIIIa-a-334	4-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-335	4-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-336	4-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-337	4-chloro-2-formylphenyl
VIIIa-a-338	4-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-339	4-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-340	4-chloro-3-formylphenyl
VIIIa-a-341	4-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-342	4-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-343	4-fluoro-2-formylphenyl
VIIIa-a-344	4-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-345	4-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-346	4-fluoro-3-formylphenyl
VIIIa-a-347	4-formyl-2-methylphenyl
VIIIa-a-348	4-formyl-3-methylphenyl
VIIIa-a-349	4-formylphenyl
VIIIa-a-350	4-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-351	4-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-352	4-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-353	4-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-354	4-[(hydroxyimino)methyl]-2,3-difluorophenyl
VIIIa-a-355	4-[(hydroxyimino)methyl]-2,5-difluorophenyl

VIIIa-a-356	4-[(hydroxyimino)methyl]-2,6-difluorophenyl
VIIIa-a-357	4-[(hydroxyimino)methyl]-2-chlorophenyl
VIIIa-a-358	4-[(hydroxyimino)methyl]-2-fluorophenyl
VIIIa-a-359	4-[(hydroxyimino)methyl]-2-methylphenyl
VIIIa-a-360	4-[(hydroxyimino)methyl]-3,5-difluorophenyl
VIIIa-a-361	4-[(hydroxyimino)methyl]-3,6-difluorophenyl
VIIIa-a-362	4-[(hydroxyimino)methyl]-3-chlorophenyl
VIIIa-a-363	4-[(hydroxyimino)methyl]-3-fluorophenyl
VIIIa-a-364	4-[(hydroxyimino)methyl]-3-methylphenyl
VIIIa-a-365	4-[(hydroxyimino)methyl]phenyl
VIIIa-a-366	4-[(methoxyimino)methyl]-2,3-difluorophenyl
VIIIa-a-367	4-[(methoxyimino)methyl]-2,5-difluorophenyl
VIIIa-a-368	4-[(methoxyimino)methyl]-2,6-difluorophenyl
VIIIa-a-369	4-[(methoxyimino)methyl]-2-chlorophenyl
VIIIa-a-370	4-[(methoxyimino)methyl]-2-fluorophenyl
VIIIa-a-371	4-[(methoxyimino)methyl]-2-methylphenyl
VIIIa-a-372	4-[(methoxyimino)methyl]-3,5-difluorophenyl
VIIIa-a-373	4-[(methoxyimino)methyl]-3,6-difluorophenyl
VIIIa-a-374	4-[(methoxyimino)methyl]-3-chlorophenyl
VIIIa-a-375	4-[(methoxyimino)methyl]-3-fluorophenyl
VIIIa-a-376	4-[(methoxyimino)methyl]-3-methylphenyl
VIIIa-a-377	4-[(methoxyimino)methyl]phenyl
VIIIa-a-378	4-[(methylsulphonyl)oxy]phenyl
VIIIa-a-379	5,6-dichloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-380	5,6-dichloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-381	5,6-dichloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-382	5,6-dichloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-383	5,6-difluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-384	5,6-difluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-385	5,6-difluoro-2-formylphenyl
VIIIa-a-386	5,6-difluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-387	5,6-difluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-388	5,6-difluoro-3-formylphenyl
VIIIa-a-389	5-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-390	5-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-391	5-chloro-2-formylphenyl
VIIIa-a-392	5-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-393	5-chloro-3-[(methylsulphonyl)oxy]phenyl

VIIIa-a-394	5-chloro-3-formylphenyl
VIIIa-a-395	5-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-396	5-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-397	5-fluoro-2-formylphenyl
VIIIa-a-398	5-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-399	5-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-400	5-fluoro-3-formylphenyl
VIIIa-a-401	5-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-402	5-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-403	5-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-404	5-methyl-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-405	6-chloro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-406	6-chloro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-407	6-chloro-2-formylphenyl
VIIIa-a-408	6-chloro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-409	6-chloro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-410	6-chloro-3-formylphenyl
VIIIa-a-411	6-fluoro-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-412	6-fluoro-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-413	6-fluoro-2-formylphenyl
VIIIa-a-414	6-fluoro-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-415	6-fluoro-3-[(methylsulphonyl)oxy]phenyl
VIIIa-a-416	6-fluoro-3-formylphenyl
VIIIa-a-417	6-methyl-2-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-418	6-methyl-2-[(methylsulphonyl)oxy]phenyl
VIIIa-a-419	6-methyl-3-(prop-2-yn-1-yloxy)phenyl
VIIIa-a-420	6-methyl-3-[(methylsulphonyl)oxy]phenyl

The logP values were measured according to EEC directive 79/831 Annex V.A8 by HPLC (High Performance Liquid Chromatography) on reversed-phase columns (C 18), using the methods below:

(a) The LC-MS determination in the acidic range is effected at pH 2.7 with 0.1% aqueous formic acid and acetonitrile (contains 0.1% formic acid) as eluents; linear gradient from 10% acetonitrile to 95% acetonitrile.

[b] The LC-MS determination in the neutral range is effected at pH 7.8 using 0.001 molar aqueous ammonium hydrogencarbonate solution and acetonitrile as eluents; linear gradient from 10% acetonitrile to 95% acetonitrile.

5

The calibration is effected with unbranched alkan-2-ones (having 3 to 16 carbon atoms) with known logP values (logP values determined on the basis of the retention times by linear interpolation between two successive alkanones).

10

The lambda-max values were determined in the maxima of the chromatographic signals using the UV spectra from 200 nm to 400 nm.

15 **NMR data of selected examples**

NMR peak list method

The ¹H NMR data of selected examples are stated in the form of ¹H NMR peak lists. For each signal peak, first the δ -value in ppm and then the signal intensity in round brackets are listed. The δ value - signal intensity number pairs for different signal peaks are listed with separation from one another by semicolons.

25

The peak list for one example therefore takes the form of:

δ_1 (intensity₁); δ_2 (intensity₂);.....; δ_i (intensity_i);.....; δ_n (intensity_n)

30

Ex. I-1, solvent: DMSO-d₆, spectrometer: 399.95 MHz

7.9545 (3.60); 7.4134 (0.43); 7.3965 (0.44); 7.1822 (0.83);
 7.1782 (1.01); 7.1516 (0.33); 7.1314 (0.39); 7.0088 (0.67);
 35 6.9876 (0.60); 6.9104 (0.58); 6.8890 (0.39); 6.8848 (0.41);
 6.0411 (1.73); 6.0305 (0.76); 4.8750 (0.67); 4.8476 (0.96);
 4.8430 (1.60); 4.8377 (1.51); 3.5424 (0.37); 3.5368 (0.73);
 3.5312 (0.35); 3.3287 (11.63); 2.8904 (16.00); 2.7310 (13.61);

2.6890 (0.37); 2.5110 (4.13); 2.5066 (8.51); 2.5021 (11.46);
 2.4976 (8.43); 2.4933 (4.16); 2.1951 (4.61); 2.0941 (0.33);
 2.0668 (0.35); -0.0002 (1.94)

5 **Ex. I-2, solvent: DMSO-d₆, spectrometer: 399.95 MHz**

8.7704 (0.40); 7.9674 (7.50); 7.9568 (14.86); 7.4320 (2.09);
 7.4148 (2.86); 7.4110 (4.55); 7.3941 (4.71); 7.3901 (3.04);
 7.3731 (2.52); 7.3581 (4.38); 7.3361 (5.35); 7.3250 (10.03);
 10 7.3031 (10.99); 7.0071 (6.19); 6.9860 (5.62); 6.9292 (5.16);
 6.9241 (2.51); 6.9113 (10.58); 6.9073 (11.20); 6.8896 (8.88);
 6.8594 (2.99); 6.0694 (0.97); 6.0558 (1.97); 6.0472 (1.41);
 6.0337 (2.54); 6.0256 (2.36); 6.0168 (1.28); 6.0031 (1.97);
 4.8896 (0.32); 4.8834 (0.74); 4.8772 (0.76); 4.8656 (1.59);
 15 4.8492 (4.59); 4.8436 (12.06); 4.8381 (14.93); 4.8323 (10.43);
 4.8048 (0.62); 4.7986 (0.99); 4.7923 (0.83); 4.7709 (3.30);
 4.7542 (0.83); 4.7321 (1.79); 4.6953 (1.81); 4.6238 (1.78);
 4.5884 (0.76); 4.4207 (1.52); 4.4000 (1.80); 4.3934 (1.73);
 4.0553 (0.87); 4.0376 (3.47); 4.0330 (2.34); 4.0245 (7.36);
 20 4.0200 (5.20); 4.0159 (5.42); 4.0073 (7.33); 3.9985 (5.23);
 3.9898 (3.31); 3.9811 (2.74); 3.9477 (1.45); 3.8563 (0.63);
 3.8236 (0.70); 3.8067 (0.86); 3.7756 (1.65); 3.7647 (1.16);
 3.7337 (2.17); 3.7023 (1.19); 3.5447 (2.02); 3.5394 (5.22);
 3.5351 (8.28); 3.5292 (3.52); 3.5185 (1.34); 3.4974 (2.51);
 25 3.4824 (3.33); 3.4765 (2.53); 3.4648 (4.49); 3.4535 (2.24);
 3.4470 (3.58); 3.4292 (1.81); 3.4119 (0.33); 3.4007 (0.64);
 3.3917 (1.11); 3.3816 (0.83); 3.3719 (1.35); 3.3629 (2.44);
 3.3537 (1.84); 3.3270 (73.42); 3.3064 (0.72); 3.2965 (0.67);
 3.2873 (0.41); 3.2150 (0.97); 3.1832 (1.71); 3.1539 (1.00);
 30 3.1259 (0.53); 3.0951 (0.78); 3.0660 (0.44); 2.9458 (0.50);
 2.8589 (0.43); 2.8419 (1.16); 2.8248 (1.83); 2.8184 (1.23);
 2.8078 (2.05); 2.7794 (2.09); 2.7501 (1.64); 2.7210 (0.51);
 2.6754 (0.60); 2.6709 (0.81); 2.6663 (0.59); 2.5411 (0.32);
 2.5241 (2.86); 2.5107 (45.32); 2.5063 (87.61); 2.5018
 35 (114.13); 2.4973 (83.01); 2.4929 (40.26); 2.3373 (0.35);
 2.3330 (0.64); 2.3286 (0.84); 2.3241 (0.63); 2.0941 (2.96);
 2.0679 (3.61); 1.9892 (9.12); 1.9618 (0.66); 1.7451 (0.50);
 1.7164 (1.15); 1.6926 (1.08); 1.6868 (1.06); 1.6630 (0.48);

1.5840 (0.92); 1.5537 (1.64); 1.5224 (1.55); 1.4918 (0.83);
 1.3970 (7.54); 1.3356 (7.58); 1.3299 (6.61); 1.3183 (14.57);
 1.3126 (11.93); 1.3009 (7.37); 1.2952 (5.52); 1.2490 (0.42);
 1.1921 (2.98); 1.1745 (12.90); 1.1567 (16.00); 1.1374 (7.43);
 5 1.0126 (11.42); 0.9956 (11.25); 0.0079 (1.26); -0.0002
 (27.88); -0.0085 (1.05)

Ex. I-3, solvent: DMSO-d₆, spectrometer: 399.95 MHz

10 7.9542 (6.35); 7.5019 (0.97); 7.4823 (2.04); 7.4638 (0.97);
 7.4485 (2.11); 7.4441 (2.24); 7.4361 (1.00); 7.4282 (2.01);
 7.4200 (1.99); 7.4158 (2.15); 7.4097 (0.92); 7.3982 (1.43);
 7.3944 (0.97); 7.3773 (0.73); 7.2380 (0.71); 7.2200 (0.68);
 7.0109 (2.14); 6.9897 (1.93); 6.9118 (1.09); 6.8863 (1.32);
 15 6.8649 (1.02); 6.0571 (0.93); 6.0347 (1.14); 6.0267 (1.10);
 6.0041 (0.97); 4.9751 (0.54); 4.9393 (2.21); 4.8987 (2.15);
 4.8847 (0.37); 4.8619 (0.69); 4.8500 (3.18); 4.8452 (4.90);
 4.8404 (3.23); 4.4223 (0.59); 4.3898 (0.64); 4.0557 (1.11);
 4.0379 (3.38); 4.0201 (3.44); 4.0023 (1.19); 3.9716 (0.57);
 20 3.9378 (0.63); 3.7681 (0.61); 3.7375 (0.70); 3.7262 (0.86);
 3.6955 (0.76); 3.5451 (1.35); 3.5393 (2.76); 3.5337 (1.30);
 3.4808 (0.60); 3.4595 (0.60); 3.4396 (0.46); 3.4174 (0.45);
 3.3913 (0.47); 3.3817 (0.33); 3.3721 (0.55); 3.3626 (0.96);
 3.3535 (0.61); 3.3283 (18.68); 3.2282 (0.43); 3.1978 (0.77);
 25 3.1681 (0.44); 2.8255 (0.39); 2.8204 (0.41); 2.7906 (0.75);
 2.7627 (0.42); 2.5111 (12.72); 2.5069 (24.81); 2.5024 (32.41);
 2.4980 (24.08); 2.2501 (16.00); 2.0863 (1.06); 2.0698 (1.09);
 1.9895 (14.31); 1.7192 (0.49); 1.6894 (0.48); 1.5510 (0.51);
 1.5202 (0.48); 1.1924 (3.91); 1.1745 (7.74); 1.1568 (3.84); -
 30 0.0002 (5.50)

Ex. I-4, solvent: DMSO-d₆, spectrometer: 399.95 MHz

7.9541 (4.56); 7.4349 (0.46); 7.4130 (2.11); 7.3972 (1.13);
 35 7.3933 (0.72); 7.3762 (0.52); 7.3518 (0.76); 7.3325 (0.92);
 7.1471 (0.94); 7.1284 (0.78); 7.0095 (1.47); 6.9883 (1.33);
 6.9105 (0.74); 6.8852 (0.91); 6.8637 (0.69); 6.0562 (0.54);
 6.0339 (0.68); 6.0262 (0.65); 6.0034 (0.56); 5.7593 (2.88);

4.9251 (0.42); 4.8892 (1.54); 4.8423 (4.60); 4.8048 (0.46);
 4.4214 (0.40); 4.3913 (0.45); 4.0553 (0.49); 4.0375 (1.49);
 4.0197 (1.55); 4.0019 (0.81); 3.9619 (0.44); 3.7305 (0.47);
 3.5436 (0.89); 3.5378 (1.80); 3.5321 (0.84); 3.4860 (0.65);
 5 3.4635 (0.65); 3.4431 (0.50); 3.4211 (0.49); 3.3685 (0.39);
 3.3592 (0.66); 3.3498 (0.42); 3.3270 (13.45); 3.1941 (0.53);
 2.8901 (1.43); 2.7854 (0.52); 2.7310 (1.19); 2.5105 (11.52);
 2.5063 (22.30); 2.5018 (29.06); 2.4973 (21.43); 2.4931
 (10.72); 2.2184 (10.46); 2.2101 (16.00); 2.0946 (0.80); 2.0695
 10 (0.81); 1.9892 (6.42); 1.7172 (0.34); 1.6895 (0.33); 1.5506
 (0.35); 1.5203 (0.34); 1.1921 (1.74); 1.1743 (3.47); 1.1565
 (1.71); -0.0002 (6.69)

Ex. I-5, solvent: DMSO-d₆, spectrometer: 399.95 MHz

15
 7.9523 (6.32); 7.6893 (1.32); 7.6698 (1.71); 7.5807 (2.39);
 7.5613 (1.28); 7.5412 (2.22); 7.5213 (1.15); 7.4361 (0.74);
 7.4188 (1.57); 7.4151 (2.00); 7.3983 (2.20); 7.3774 (1.00);
 7.0114 (2.15); 6.9902 (1.94); 6.9109 (1.08); 6.8854 (1.35);
 20 6.8640 (1.02); 6.0567 (0.92); 6.0343 (1.13); 6.0264 (1.10);
 6.0037 (0.96); 5.7598 (1.88); 4.9935 (0.58); 4.9577 (2.20);
 4.9145 (2.12); 4.8908 (0.35); 4.8797 (0.65); 4.8505 (3.16);
 4.8457 (4.91); 4.8408 (3.18); 4.4224 (0.61); 4.3902 (0.67);
 4.0560 (0.79); 4.0382 (2.41); 4.0204 (2.45); 4.0026 (0.85);
 25 3.9687 (0.59); 3.9353 (0.66); 3.7654 (0.64); 3.7334 (0.72);
 3.7233 (0.90); 3.6918 (0.79); 3.5457 (1.30); 3.5399 (2.71);
 3.5342 (1.28); 3.4790 (0.55); 3.4564 (0.55); 3.4410 (0.42);
 3.4139 (0.41); 3.3912 (0.48); 3.3814 (0.36); 3.3717 (0.57);
 3.3623 (0.97); 3.3532 (0.61); 3.3429 (0.45); 3.3286 (11.09);
 30 3.2288 (0.44); 3.1983 (0.80); 3.1689 (0.44); 2.8263 (0.47);
 2.8214 (0.43); 2.7907 (0.79); 2.7622 (0.44); 2.5251 (0.45);
 2.5073 (19.73); 2.5028 (25.92); 2.4984 (19.39); 2.2689
 (16.00); 2.1042 (1.05); 2.0709 (1.17); 1.9897 (10.36); 1.7171
 (0.52); 1.6895 (0.50); 1.5492 (0.53); 1.5205 (0.50); 1.3969
 35 (2.38); 1.1926 (2.89); 1.1748 (5.76); 1.1570 (2.84); -0.0002
 (6.56); -0.0084 (0.35)

Ex. I-6, solvent: DMSO-d₆, spectrometer: 399.95 MHz

8.7705 (0.42); 8.0099 (3.69); 7.4036 (1.92); 7.3536 (0.94);
 7.3499 (0.82); 7.3340 (1.13); 7.3301 (1.00); 7.1772 (2.14);
 7.1651 (2.47); 7.1611 (2.22); 7.1538 (1.50); 7.1340 (1.17);
 5 7.0888 (0.77); 7.0667 (0.89); 5.8531 (0.60); 5.8355 (0.71);
 5.8255 (0.74); 5.8078 (0.62); 4.9178 (0.40); 4.8774 (4.12);
 4.8717 (4.06); 4.8408 (1.59); 4.8165 (0.33); 4.8050 (0.44);
 4.4103 (0.40); 4.3795 (0.46); 4.0558 (0.43); 4.0380 (1.31);
 4.0202 (1.34); 4.0023 (0.51); 3.9802 (0.41); 3.9468 (0.46);
 10 3.9011 (0.54); 3.8735 (0.56); 3.8579 (0.65); 3.8304 (0.54);
 3.6078 (0.75); 3.6020 (1.51); 3.5968 (0.75); 3.3502 (0.40);
 3.3408 (0.75); 3.3282 (13.10); 3.3124 (0.41); 3.2678 (0.69);
 3.2503 (0.67); 3.2245 (0.70); 3.2073 (0.72); 3.1835 (0.56);
 3.1533 (0.35); 2.7724 (0.55); 2.7438 (0.35); 2.5246 (0.64);
 15 2.5112 (11.67); 2.5070 (22.74); 2.5025 (29.67); 2.4982
 (21.87); 2.2281 (3.23); 2.2182 (16.00); 2.2074 (13.09); 2.1996
 (2.30); 2.1902 (0.67); 2.0916 (0.70); 2.0820 (0.72); 2.0499
 (0.83); 1.9897 (5.58); 1.9092 (1.42); 1.7042 (0.33); 1.6968
 (0.34); 1.6732 (0.34); 1.5307 (0.33); 1.5202 (0.35); 1.5001
 20 (0.34); 1.4907 (0.34); 1.3969 (4.19); 1.1925 (1.54); 1.1747
 (3.02); 1.1569 (1.51); -0.0002 (6.53)

Ex. I-7, solvent: DMSO-d₆, spectrometer: 399.95 MHz

25 12.9922 (0.38); 8.7715 (0.37); 8.6617 (1.17); 8.0222 (3.57);
 8.0128 (9.56); 7.9937 (0.35); 7.3508 (1.87); 7.3231 (7.08);
 7.3012 (7.21); 7.1746 (6.71); 7.1621 (7.00); 7.1585 (6.15);
 7.0890 (2.56); 7.0861 (2.56); 7.0636 (2.71); 6.9204 (3.70);
 6.9137 (6.26); 6.8981 (4.17); 6.8921 (5.48); 5.8648 (0.63);
 30 5.8525 (1.65); 5.8357 (2.26); 5.8251 (1.92); 5.8076 (1.59);
 4.8718 (11.07); 4.8662 (10.08); 4.8259 (3.15); 4.8014 (0.46);
 4.7948 (0.36); 4.7674 (3.06); 4.7309 (1.39); 4.6919 (1.12);
 4.6139 (1.01); 4.5791 (0.47); 4.4126 (1.20); 4.4035 (1.23);
 4.3825 (1.37); 4.0563 (0.82); 4.0470 (1.50); 4.0386 (2.60);
 35 4.0305 (4.26); 4.0211 (3.54); 4.0130 (4.45); 4.0056 (3.60);
 3.9955 (1.88); 3.9881 (2.69); 3.9706 (1.68); 3.9259 (1.56);
 3.9044 (1.84); 3.8977 (1.24); 3.8820 (1.30); 3.8765 (1.96);
 3.8611 (2.11); 3.8552 (1.16); 3.8334 (1.85); 3.8136 (0.48);

3.6065 (2.67); 3.6008 (5.18); 3.5953 (2.56); 3.5802 (0.39);
 3.4978 (0.71); 3.4801 (1.85); 3.4623 (2.60); 3.4445 (2.00);
 3.4340 (0.43); 3.4268 (0.82); 3.3819 (0.51); 3.3731 (0.82);
 3.3635 (0.69); 3.3535 (1.10); 3.3442 (2.07); 3.3304 (20.08);
 5 3.3156 (1.38); 3.3057 (1.23); 3.2835 (1.13); 3.2674 (2.56);
 3.2501 (1.95); 3.2403 (0.91); 3.2242 (2.21); 3.2069 (2.25);
 3.1709 (1.36); 3.1419 (0.82); 3.1111 (0.45); 3.0754 (0.71);
 3.0483 (0.44); 3.0138 (0.53); 2.9472 (0.39); 2.8372 (0.51);
 2.8205 (0.92); 2.8016 (1.01); 2.7648 (1.65); 2.7342 (1.22);
 10 2.7036 (0.40); 2.6771 (0.33); 2.6724 (0.45); 2.5080 (36.63);
 2.5036 (46.90); 2.4992 (34.39); 2.3305 (0.33); 2.0721 (2.16);
 2.0460 (2.77); 1.9904 (5.45); 1.9584 (0.71); 1.9316 (0.42);
 1.7294 (0.39); 1.7003 (0.88); 1.6773 (0.85); 1.6485 (0.39);
 1.5890 (0.35); 1.5595 (0.97); 1.5289 (1.45); 1.4981 (1.32);
 15 1.4631 (0.69); 1.4041 (11.07); 1.3961 (3.74); 1.3397 (7.11);
 1.3224 (16.00); 1.3052 (11.20); 1.2880 (2.43); 1.2491 (0.35);
 1.2342 (0.35); 1.1932 (1.83); 1.1753 (10.41); 1.1572 (12.78);
 1.1348 (7.30); 1.0097 (5.45); 0.9929 (5.28); 0.0079 (0.57); -
 0.0002 (12.39); -0.0085 (0.54)

20

Ex. I-8, solvent: CD₃CN, spectrometer: 399.95 MHz

10.1384 (4.87); 7.9635 (1.51); 7.9450 (1.44); 7.6833 (2.36);
 7.6811 (2.86); 7.6714 (3.22); 7.6684 (1.83); 7.6487 (5.99);
 25 7.5976 (0.76); 7.5861 (1.11); 7.5783 (0.82); 7.5664 (0.76);
 7.5568 (0.48); 6.4608 (1.02); 6.4443 (1.03); 6.4329 (1.06);
 6.4162 (1.04); 4.6285 (2.33); 4.6116 (2.40); 4.4785 (0.38);
 4.4512 (0.39); 4.3267 (0.42); 4.0990 (1.23); 4.0864 (0.94);
 4.0686 (2.80); 4.0554 (1.49); 4.0507 (2.75); 4.0329 (0.94);
 30 4.0275 (1.36); 3.9830 (0.37); 3.9498 (0.40); 3.3104 (0.47);
 3.2908 (0.48); 3.2811 (0.95); 3.2712 (0.51); 3.2524 (0.50);
 3.1672 (1.48); 3.1505 (1.74); 3.1236 (1.53); 3.1069 (1.36);
 2.7614 (0.48); 2.1913 (0.46); 2.1898 (0.51); 2.1692 (146.43);
 2.1651 (275.77); 2.1639 (218.61); 2.1469 (1.17); 2.1323
 35 (0.61); 2.1199 (0.97); 2.1139 (1.36); 2.1076 (1.38); 2.1015
 (1.04); 2.0954 (0.76); 2.0812 (0.94); 1.9721 (12.09); 1.9645
 (14.96); 1.9584 (2.42); 1.9526 (40.42); 1.9464 (79.16); 1.9402
 (115.91); 1.9340 (78.66); 1.9279 (39.79); 1.9150 (0.58);

1.8888 (0.58); 1.8536 (15.08); 1.8373 (0.55); 1.8175 (0.82);
 1.8083 (16.00); 1.7748 (0.55); 1.7686 (0.77); 1.7623 (0.56);
 1.7565 (0.44); 1.7305 (0.37); 1.6986 (0.38); 1.6154 (0.35);
 1.5913 (0.33); 1.5812 (0.32); 1.3349 (0.33); 1.2218 (3.22);
 5 1.2040 (6.45); 1.1861 (3.15); -0.0002 (1.13)

Ex. I-9, solvent: DMSO-d6, spectrometer: 399.95 MHz

8.0327 (6.22); 7.5906 (0.54); 7.5747 (0.64); 7.5696 (1.15);
 10 7.5538 (1.19); 7.5487 (0.76); 7.5329 (0.70); 7.3481 (0.81);
 7.3300 (2.48); 7.3238 (1.11); 7.3092 (1.66); 7.3025 (0.75);
 6.0465 (0.62); 6.0230 (0.78); 6.0166 (0.74); 5.9930 (0.65);
 5.0844 (0.38); 5.0466 (1.73); 5.0071 (1.67); 4.9696 (0.39);
 4.3987 (0.40); 4.3659 (0.43); 4.0380 (0.51); 4.0202 (0.52);
 15 3.8983 (0.43); 3.8942 (0.40); 3.8679 (0.49); 3.8636 (0.49);
 3.8550 (0.59); 3.8506 (0.60); 3.8242 (0.85); 3.8205 (0.87);
 3.7880 (0.43); 3.5470 (16.00); 3.5275 (0.76); 3.5040 (0.72);
 3.4826 (0.57); 3.4598 (0.56); 3.3900 (0.34); 3.3702 (0.45);
 3.3613 (0.73); 3.3519 (0.42); 3.3245 (11.87); 3.1663 (0.53);
 20 2.7902 (0.52); 2.5245 (0.41); 2.5197 (0.70); 2.5112 (8.64);
 2.5067 (17.31); 2.5021 (22.66); 2.4974 (16.20); 2.4929 (7.58);
 2.0999 (0.80); 2.0945 (0.84); 2.0629 (15.57); 1.9891 (2.31);
 1.7285 (0.35); 1.6980 (0.32); 1.5563 (0.33); 1.5271 (0.32);
 1.2496 (0.35); 1.1924 (0.64); 1.1746 (1.29); 1.1568 (0.62);
 25 0.0080 (0.52); -0.0002 (14.82); -0.0086 (0.44)

Ex. I-10, solvent: DMSO-d6, spectrometer: 399.95 MHz

8.0469 (5.74); 7.5142 (0.80); 7.5100 (0.92); 7.4931 (1.33);
 30 7.4682 (0.81); 7.4637 (0.99); 7.4529 (2.98); 7.4486 (3.63);
 7.4345 (0.58); 7.4232 (1.09); 7.4170 (0.76); 7.4070 (0.65);
 7.4046 (0.79); 7.4015 (0.59); 7.3979 (0.57); 7.3893 (0.39);
 7.3827 (0.36); 6.0059 (0.79); 5.9865 (0.93); 5.9780 (0.90);
 5.9586 (0.81); 5.0803 (0.36); 5.0426 (1.65); 5.0030 (1.60);
 35 4.9651 (0.36); 4.3930 (0.37); 4.3602 (0.40); 4.0381 (0.64);
 4.0203 (0.65); 3.9914 (0.81); 3.9634 (0.95); 3.9479 (1.06);
 3.9200 (0.87); 3.8135 (0.36); 3.7798 (0.40); 3.5526 (16.00);
 3.3767 (0.36); 3.3561 (1.28); 3.3483 (0.78); 3.3367 (1.27);

3.3240 (8.73); 3.3126 (1.08); 3.2932 (0.91); 3.1569 (0.50);
 2.7799 (0.48); 2.5247 (0.40); 2.5199 (0.61); 2.5113 (8.66);
 2.5067 (17.70); 2.5021 (23.44); 2.4975 (16.86); 2.4930 (7.97);
 2.0767 (0.90); 2.0611 (14.17); 2.0522 (1.34); 1.9892 (2.84);
 5 1.2495 (0.33); 1.1927 (0.79); 1.1749 (1.58); 1.1571 (0.78);
 0.0080 (0.43); -0.0002 (14.15); -0.0085 (0.45)

Ex. I-11, solvent: DMSO-d6, spectrometer: 399.95 MHz

10 7.9982 (5.94); 7.3449 (0.52); 7.3407 (0.70); 7.3199 (1.87);
 7.3060 (0.69); 7.3018 (1.61); 7.2985 (1.84); 7.1486 (1.75);
 7.1285 (1.48); 7.0156 (0.96); 6.9981 (1.67); 6.9795 (0.77);
 5.8806 (0.85); 5.8628 (0.97); 5.8529 (0.94); 5.8350 (0.86);
 5.7577 (1.13); 4.8781 (4.66); 4.8722 (4.67); 4.6666 (2.44);
 15 4.6483 (2.43); 4.5951 (0.67); 4.3878 (0.43); 4.3549 (0.47);
 4.0557 (0.38); 4.0379 (1.14); 4.0201 (1.16); 4.0023 (0.39);
 3.9284 (0.41); 3.8900 (1.26); 3.8621 (1.05); 3.8470 (1.15);
 3.8192 (0.96); 3.5879 (1.22); 3.5821 (2.71); 3.5762 (1.21);
 3.3571 (0.39); 3.3375 (0.56); 3.3240 (19.15); 3.3094 (0.40);
 20 3.2993 (0.51); 3.2588 (1.16); 3.2409 (1.12); 3.2158 (1.04);
 3.1981 (1.00); 3.1408 (0.57); 2.7458 (0.55); 2.5243 (0.43);
 2.5194 (0.75); 2.5108 (10.95); 2.5064 (22.13); 2.5018 (29.24);
 2.4973 (21.30); 2.4928 (10.34); 2.0607 (0.89); 2.0308 (1.05);
 1.9891 (5.09); 1.8288 (15.58); 1.8235 (1.85); 1.8196 (1.01);
 25 1.7932 (16.00); 1.7823 (0.95); 1.6664 (0.46); 1.6433 (0.37);
 1.6320 (0.39); 1.5163 (0.36); 1.5084 (0.38); 1.4872 (0.37);
 1.4792 (0.35); 1.1924 (1.33); 1.1746 (2.66); 1.1568 (1.31); -
 0.0002 (2.35)

30 Ex. I-12, solvent: DMSO-d6, spectrometer: 399.95 MHz

8.0416 (5.69); 7.5142 (0.85); 7.5102 (0.98); 7.4930 (1.44);
 7.4727 (0.34); 7.4683 (0.84); 7.4638 (1.03); 7.4530 (3.23);
 7.4486 (3.86); 7.4344 (0.68); 7.4234 (1.18); 7.4173 (0.86);
 35 7.4048 (0.89); 7.4018 (0.70); 7.3983 (0.66); 7.3896 (0.45);
 7.3830 (0.41); 6.0049 (0.84); 5.9855 (1.01); 5.9772 (0.97);
 5.9577 (0.88); 5.7581 (0.70); 4.6691 (2.29); 4.6515 (2.34);
 4.3911 (0.40); 4.3593 (0.44); 4.0558 (0.32); 4.0380 (0.93);

4.0202 (0.95); 4.0024 (0.35); 3.9909 (0.84); 3.9629 (1.00);
 3.9474 (1.24); 3.9323 (0.43); 3.9196 (1.13); 3.8993 (0.45);
 3.5533 (16.00); 3.3691 (0.43); 3.3555 (1.19); 3.3501 (0.58);
 3.3360 (1.53); 3.3254 (14.63); 3.3120 (1.37); 3.2927 (0.98);
 5 3.1461 (0.53); 2.7512 (0.53); 2.5247 (0.41); 2.5198 (0.67);
 2.5113 (7.79); 2.5068 (15.48); 2.5023 (20.20); 2.4977 (14.58);
 2.4932 (6.96); 2.0677 (0.85); 2.0372 (0.99); 1.9893 (4.12);
 1.8296 (14.61); 1.8187 (0.70); 1.7936 (15.22); 1.7815 (0.68);
 1.6721 (0.35); 1.6676 (0.35); 1.6494 (0.33); 1.6417 (0.33);
 10 1.5159 (0.35); 1.4932 (0.33); 1.1925 (1.10); 1.1747 (2.18);
 1.1569 (1.07); -0.0002 (1.61)

Ex. I-13, solvent: DMSO-d6, spectrometer: 399.95 MHz

15 8.0271 (6.25); 7.5903 (0.53); 7.5745 (0.62); 7.5694 (1.14);
 7.5535 (1.18); 7.5484 (0.76); 7.5326 (0.69); 7.3484 (0.80);
 7.3297 (2.54); 7.3243 (1.19); 7.3088 (1.66); 7.3029 (0.80);
 6.0454 (0.63); 6.0220 (0.78); 6.0155 (0.75); 5.9920 (0.66);
 4.6725 (2.29); 4.6557 (2.27); 4.3964 (0.39); 4.3649 (0.42);
 20 4.0379 (0.66); 4.0201 (0.66); 3.9404 (0.38); 3.8978 (0.69);
 3.8939 (0.59); 3.8672 (0.49); 3.8630 (0.49); 3.8543 (0.58);
 3.8499 (0.59); 3.8238 (0.49); 3.8197 (0.51); 3.5471 (16.00);
 3.5264 (0.76); 3.5028 (0.73); 3.4818 (0.56); 3.4586 (0.56);
 3.3817 (0.36); 3.3623 (0.42); 3.3528 (0.77); 3.3433 (0.46);
 25 3.3250 (15.50); 3.1551 (0.51); 2.7608 (0.50); 2.5245 (0.34);
 2.5197 (0.56); 2.5111 (7.41); 2.5066 (14.97); 2.5020 (19.76);
 2.4974 (14.25); 2.4929 (6.74); 2.0859 (0.82); 2.0555 (0.94);
 1.9891 (2.95); 1.8325 (15.49); 1.7955 (15.54); 1.6908 (0.33);
 1.6698 (0.34); 1.6620 (0.32); 1.5326 (0.33); 1.1924 (0.80);
 30 1.1745 (1.59); 1.1568 (0.78); -0.0002 (1.73)

Ex. I-14, solvent: CDCl3, spectrometer: 250.13 MHz

7.6093 (1.70); 7.5932 (0.43); 7.5684 (0.47); 7.5599 (0.47);
 35 7.4024 (0.76); 7.3874 (1.25); 7.3561 (0.38); 7.3447 (0.46);
 7.2658 (5.88); 7.2635 (5.81); 6.0595 (0.34); 6.0271 (0.40);
 6.0145 (0.39); 5.9817 (0.38); 5.3038 (3.98); 5.3015 (3.99);
 4.7093 (1.76); 4.7048 (1.79); 3.9822 (0.33); 3.9378 (0.34);

3.9131 (0.46); 3.8680 (0.42); 3.4797 (0.45); 3.4464 (0.43);
 3.4092 (0.35); 3.3769 (0.35); 3.2957 (0.33); 3.2640 (5.71);
 3.2621 (5.51); 2.8063 (0.39); 2.1921 (0.35); 2.1471 (0.46);
 1.8761 (0.37); 1.8514 (5.47); 1.8494 (5.40); 1.7818 (0.42);
 5 1.7665 (0.44); 1.7310 (0.37); 1.7150 (0.36); 1.6003 (5.45);
 1.2560 (0.32); 1.0915 (16.00); 1.0898 (15.63); 0.0020 (3.70);
 -0.0002 (3.83)

Ex. XVI-81, solvent: DMSO-d6, spectrometer: 399.95 MHz

10
 8.0023 (2.21); 7.3274 (0.63); 7.3094 (0.52); 7.3064 (0.61);
 7.1559 (0.55); 7.1361 (0.48); 7.0063 (0.54); 4.8848 (1.59);
 4.8789 (1.58); 4.0457 (0.33); 4.0280 (0.46); 3.8681 (0.37);
 3.8529 (0.41); 3.8252 (0.34); 3.5917 (0.42); 3.5858 (0.91);
 15 3.5799 (0.39); 3.3325 (21.59); 3.2674 (0.44); 3.2496 (0.49);
 3.2243 (0.41); 3.2066 (0.40); 2.5323 (0.43); 2.5275 (0.67);
 2.5189 (7.55); 2.5144 (15.19); 2.5098 (20.02); 2.5052 (14.38);
 2.5006 (6.78); 2.0156 (0.36); 1.9968 (1.09); 1.5688 (0.37);
 1.5590 (0.33); 1.5375 (0.32); 1.4130 (16.00); 1.4055 (4.84);
 20 1.3983 (0.54); 1.1828 (0.57)

Ex. XVI-96, solvent: DMSO-d6, spectrometer: 399.95 MHz

10.1331 (1.80); 7.9900 (2.32); 7.8225 (0.80); 7.5122 (0.57);
 25 7.5092 (0.58); 7.4848 (0.99); 7.4649 (0.42); 6.3801 (0.33);
 6.3686 (0.33); 4.0440 (0.39); 4.0162 (0.67); 4.0004 (0.66);
 3.9728 (0.57); 3.3355 (5.02); 3.2206 (0.32); 3.1791 (0.44);
 3.1628 (0.41); 3.1354 (0.38); 3.1192 (0.38); 2.5121 (5.31);
 2.5077 (10.74); 2.5032 (14.32); 2.4987 (10.76); 2.4943 (5.47);
 30 2.3956 (3.63); 2.0202 (0.40); 1.9900 (0.60); 1.5485 (0.36);
 1.5391 (0.41); 1.5179 (0.35); 1.5082 (0.34); 1.4022 (16.00);
 1.3969 (13.31); 1.3823 (0.80); -0.0002 (6.28)

Ex. XVI-99, solvent: DMSO-d6, spectrometer: 399.95 MHz

35
 10.1791 (1.59); 8.0298 (0.44); 8.0257 (0.50); 8.0094 (0.57);
 8.0067 (0.52); 7.9879 (2.03); 7.7141 (0.51); 7.6953 (0.37);
 7.6920 (0.34); 7.6213 (0.41); 7.6054 (0.91); 7.5869 (0.61);

4.0770 (0.34); 4.0490 (0.43); 4.0400 (0.56); 4.0332 (0.54);
 4.0222 (0.72); 4.0050 (0.74); 3.9820 (0.36); 3.3146 (76.20);
 3.2228 (0.38); 3.2155 (0.57); 3.1993 (0.48); 3.1719 (0.40);
 3.1557 (0.39); 2.5109 (4.05); 2.5066 (7.25); 2.5021 (9.26);
 5 2.4977 (6.41); 2.4933 (3.09); 2.0288 (0.39); 1.9877 (2.29);
 1.5561 (0.38); 1.5457 (0.37); 1.5242 (0.34); 1.4203 (0.57);
 1.4095 (3.12); 1.4028 (16.00); 1.3837 (0.49); 1.1936 (0.57);
 1.1758 (1.11); 1.1580 (0.55); -0.0002 (0.49)

10 **Ex. XVI-142, solvent: DMSO-d6, spectrometer: 399.95 MHz**

8.0379 (2.21); 7.5135 (0.38); 7.5098 (0.43); 7.4923 (0.63);
 7.4673 (0.37); 7.4629 (0.44); 7.4520 (1.34); 7.4477 (1.52);
 7.4226 (0.47); 7.4165 (0.33); 7.4040 (0.37); 6.0031 (0.35);
 15 5.9837 (0.42); 5.9753 (0.40); 5.9558 (0.36); 4.0252 (0.33);
 3.9887 (0.66); 3.9606 (0.44); 3.9452 (0.47); 3.9173 (0.39);
 3.5512 (6.33); 3.3554 (0.46); 3.3357 (0.63); 3.3237 (9.19);
 3.3121 (0.51); 3.2926 (0.41); 3.2460 (0.35); 2.5108 (5.80);
 2.5066 (10.93); 2.5021 (14.10); 2.4976 (10.36); 2.4933 (5.17);
 20 2.0453 (0.42); 2.0403 (0.42); 2.0134 (0.49); 1.5667 (0.40);
 1.5584 (0.44); 1.5366 (0.40); 1.5274 (0.37); 1.4061 (16.00);
 1.0435 (0.44); 1.0283 (0.43)

Ex. XVI-405, solvent: DMSO-d6, spectrometer: 399.95 MHz

25
 7.9403 (2.52); 7.4067 (0.50); 7.3861 (0.88); 7.3654 (0.70);
 7.1577 (0.74); 7.1497 (0.90); 7.1364 (0.57); 7.1297 (0.75);
 7.1275 (0.53); 6.2254 (0.35); 6.2008 (0.46); 6.1946 (0.42);
 6.1699 (0.36); 4.7922 (0.79); 4.7859 (1.02); 4.7840 (0.98);
 30 4.7776 (0.78); 4.0380 (1.03); 4.0202 (0.99); 4.0024 (0.55);
 3.7230 (0.34); 3.7118 (0.48); 3.6810 (0.39); 3.5723 (0.44);
 3.5476 (0.44); 3.4430 (0.48); 3.4371 (1.11); 3.4312 (0.46);
 3.3224 (6.80); 2.5112 (3.77); 2.5066 (7.61); 2.5020 (10.08);
 2.4974 (7.28); 2.4929 (3.45); 2.0660 (0.34); 2.0609 (0.36);
 35 2.0341 (0.41); 2.0288 (0.39); 1.9887 (3.52); 1.5770 (0.32);
 1.4097 (16.00); 1.1930 (0.97); 1.1752 (1.92); 1.1573 (0.95); -
 0.0002 (0.69)

Ex. XVI-411, solvent: DMSO-d6, spectrometer: 399.95 MHz

7.9551 (2.65); 7.4104 (0.46); 7.3936 (0.47); 7.0086 (0.64);
6.9874 (0.58); 6.9077 (0.32); 6.8862 (0.35); 6.8829 (0.37);
5 6.0427 (0.34); 6.0346 (0.32); 4.8486 (0.94); 4.8431 (1.62);
4.8375 (0.91); 3.5414 (0.53); 3.5355 (1.20); 3.5296 (0.51);
3.5158 (0.33); 3.4932 (0.33); 3.3217 (3.50); 3.2549 (0.33);
2.8906 (0.76); 2.7311 (0.48); 2.5109 (3.82); 2.5064 (7.66);
2.5018 (10.06); 2.4972 (7.17); 2.4926 (3.38); 2.0629 (0.34);
10 2.0576 (0.35); 2.0311 (0.41); 2.0258 (0.40); 1.5840 (0.32);
1.5741 (0.35); 1.5532 (0.33); 1.4094 (16.00); 1.3974 (4.87); -
0.0002 (7.08)

Ex. XVI-412, solvent: DMSO-d6, spectrometer: 399.95 MHz

15
8.0235 (2.59); 7.5685 (0.48); 7.5526 (0.49); 7.3468 (0.35);
7.3290 (0.99); 7.3225 (0.47); 7.3083 (0.67); 6.0203 (0.33);
4.0380 (0.97); 4.0202 (0.94); 4.0024 (0.54); 3.5455 (6.73);
3.5261 (0.33); 3.3235 (9.89); 3.2588 (0.34); 2.5195 (0.38);
20 2.5111 (4.57); 2.5066 (9.08); 2.5020 (11.93); 2.4975 (8.59);
2.4930 (4.09); 2.0632 (0.36); 2.0576 (0.37); 2.0310 (0.42);
1.9889 (3.21); 1.5730 (0.32); 1.4085 (16.00); 1.1928 (0.84);
1.1750 (1.67); 1.1572 (0.82); -0.0002 (1.02)

25 Ex. XIIIa-81, solvent: DMSO-d6, spectrometer: 250.13 MHz

8.0476 (3.47); 7.3672 (0.36); 7.3353 (1.41); 7.3054 (1.33);
7.1686 (0.81); 7.1370 (0.63); 7.0414 (0.47); 7.0114 (0.78);
6.9820 (0.33); 5.9127 (0.37); 5.8843 (0.48); 5.8686 (0.45);
30 5.8401 (0.38); 4.8933 (2.26); 4.8838 (2.22); 3.9297 (0.40);
3.8852 (0.47); 3.8609 (0.55); 3.8164 (0.46); 3.6534 (0.50);
3.6266 (0.34); 3.6146 (0.78); 3.6053 (1.72); 3.5958 (0.80);
3.5776 (16.00); 3.4196 (0.66); 3.4064 (0.47); 3.3943 (0.67);
3.3770 (0.53); 3.3618 (0.48); 3.3296 (0.55); 3.2946 (0.60);
35 3.2661 (0.51); 3.2261 (0.44); 3.1973 (0.43); 3.0480 (0.41);
3.0062 (0.39); 2.5269 (2.31); 2.5197 (4.97); 2.5123 (6.87);
2.5050 (4.89); 2.4977 (2.22); 2.2226 (0.38); 2.1762 (0.54);
1.9494 (0.44); 1.9198 (0.45); 1.9077 (0.37)

Ex. XIIIa-99, solvent: DMSO-d6, spectrometer: 399.95 MHz

10.1852 (0.84); 8.0314 (1.21); 7.6096 (0.56); 7.5908 (0.35);
5 4.9198 (0.83); 3.5685 (16.00); 2.5153 (1.65); 2.5110 (3.07);
2.5066 (4.02); 2.5022 (2.88); 2.4979 (1.45)

Ex. XIIIa-142, solvent: DMSO-d6, spectrometer: 399.95 MHz

10 8.0794 (1.10); 7.4568 (0.52); 7.4530 (0.70); 3.5680 (16.00);
3.5561 (3.16); 3.3620 (0.33); 3.3187 (0.33); 2.5122 (3.27);
2.5077 (6.64); 2.5031 (8.90); 2.4985 (6.58); 2.4940 (3.22);
1.5955 (0.36)

15 Ex. XIIIa-236, solvent: DMSO-d6, spectrometer: 399.95 MHz

10.0393 (1.14); 10.0353 (1.07); 8.0737 (1.13); 8.0695 (1.10);
7.9323 (0.83); 7.9014 (0.45); 7.8983 (0.42); 7.8826 (0.51);
7.7509 (0.36); 7.7316 (0.53); 7.6643 (0.37); 7.6455 (0.55);
20 6.1538 (0.93); 5.8819 (0.37); 3.5694 (16.00); 3.5671 (12.49);
3.5649 (15.32); 3.4501 (0.42); 3.4466 (0.44); 3.4306 (0.45);
3.4269 (0.47); 3.4123 (0.43); 3.4073 (0.49); 3.3878 (0.48);
3.3347 (0.59); 3.3038 (0.70); 3.0320 (0.55); 3.0039 (0.53);
2.5163 (4.08); 2.5125 (5.39); 2.5084 (4.95); 2.2135 (0.49);
25 2.1809 (0.67); 2.0048 (0.54); 1.9753 (0.48); -0.0002 (0.36); -
0.0047 (0.34)

Ex. XIIIa-349, solvent: DMSO-d6, spectrometer: 399.95 MHz

30 10.0194 (2.71); 8.0738 (3.28); 7.9545 (1.88); 7.9378 (0.78);
7.9339 (2.11); 7.6260 (1.88); 7.6055 (1.65); 5.8958 (0.45);
5.8766 (0.52); 5.8683 (0.50); 5.8490 (0.45); 3.9953 (0.51);
3.9675 (0.57); 3.9521 (0.66); 3.9244 (0.56); 3.5689 (16.00);
3.4221 (0.79); 3.4095 (0.55); 3.4028 (0.83); 3.3915 (0.34);
35 3.3792 (0.76); 3.3597 (0.82); 3.3475 (0.61); 3.3157 (0.71);
3.0334 (0.58); 3.0066 (0.56); 2.5679 (0.44); 2.5157 (4.27);
2.5113 (8.22); 2.5068 (10.52); 2.5022 (7.51); 2.4978 (3.56);
2.2079 (0.52); 2.1829 (0.68); 2.1792 (0.69); 1.9731 (0.55);

1.9474 (0.49); 1.3565 (0.95); 0.0079 (0.43); -0.0002 (10.08);
-0.0085 (0.34)

Ex. XIIIa-405, solvent: DMSO-d6, spectrometer: 399.95 MHz

5
7.9866 (2.24); 7.4120 (0.45); 7.3913 (0.91); 7.3707 (0.66);
7.1635 (0.77); 7.1531 (0.89); 7.1425 (0.65); 7.1331 (0.74);
6.2352 (0.36); 6.2106 (0.47); 6.2044 (0.43); 6.1797 (0.37);
4.7980 (0.90); 4.7920 (1.02); 4.7883 (1.01); 4.7821 (0.87);
10 3.7339 (0.37); 3.7229 (0.50); 3.6921 (0.42); 3.5774 (0.70);
3.5680 (16.00); 3.5532 (0.52); 3.5356 (0.34); 3.5111 (0.34);
3.4624 (0.59); 3.4567 (1.14); 3.4508 (0.58); 3.4175 (0.36);
3.3598 (0.44); 3.3279 (0.54); 3.0421 (0.44); 3.0155 (0.43);
2.5130 (3.61); 2.5087 (7.01); 2.5042 (9.07); 2.4996 (6.63);
15 2.4953 (3.26); 2.2274 (0.40); 2.1984 (0.54); 1.9769 (0.40);
1.9498 (0.38); -0.0002 (5.24)

Ex. XIIIa-411, solvent: DMSO-d6, spectrometer: 399.95 MHz

20 8.0015 (2.35); 7.4157 (0.58); 7.3985 (0.55); 7.0128 (0.89);
6.9917 (0.82); 6.9107 (0.47); 6.8864 (0.61); 6.8637 (0.43);
6.0736 (0.39); 6.0514 (0.48); 6.0436 (0.47); 6.0210 (0.41);
5.7614 (0.38); 4.8476 (2.03); 3.7794 (0.35); 3.7680 (0.43);
3.7375 (0.39); 3.5680 (16.00); 3.5655 (11.38); 3.5537 (1.28);
25 3.5212 (0.47); 3.4989 (0.47); 3.4787 (0.37); 3.4564 (0.44);
3.4173 (0.45); 3.3537 (0.58); 3.3226 (0.71); 3.0381 (0.58);
3.0107 (0.56); 2.8913 (0.38); 2.7306 (0.35); 2.5089 (10.65);
2.5049 (14.30); 2.2234 (0.55); 2.1921 (0.72); 1.9766 (0.58);
1.9481 (0.51); -0.0002 (3.54); -0.0027 (2.58)

30

Ex. XIIIa-412, solvent: DMSO-d6, spectrometer: 399.95 MHz

8.0656 (0.97); 7.3355 (0.35); 3.5681 (16.00); 3.5526 (2.63);
2.5128 (2.19); 2.5085 (4.36); 2.5039 (5.79); 2.4994 (4.34);
35 2.4951 (2.19)

Ex. VIIa-a-143, solvent: DMSO-d6, spectrometer: 399.95 MHz

13.7691 (1.28); 8.0269 (0.33); 7.8403 (1.07); 7.8195 (1.73);
 7.8042 (2.66); 7.7584 (2.21); 7.7381 (1.42); 6.1631 (0.88);
 6.1492 (0.57); 6.1423 (1.04); 6.1335 (0.98); 6.1126 (0.91);
 6.1064 (0.46); 5.6296 (0.44); 5.6015 (0.41); 5.0141 (1.08);
 5 4.9727 (4.57); 4.9653 (2.84); 4.9504 (4.46); 4.9090 (1.05);
 4.2764 (1.13); 3.8091 (1.02); 3.7793 (1.13); 3.7648 (1.26);
 3.7351 (1.12); 3.6493 (16.00); 3.5693 (3.62); 3.3743 (5.39);
 3.1903 (1.20); 3.1695 (1.19); 3.1460 (1.06); 3.1253 (1.05);
 2.5114 (7.75); 2.5070 (1.02); 2.5027 (7.52)

10

Ex. VIIa-a-81, solvent: DMSO-d6, spectrometer: 399.95 MHz

7.3037 (0.63); 7.2995 (0.72); 7.2809 (1.24); 7.2646 (0.85);
 7.2604 (1.00); 7.2462 (1.33); 7.2423 (1.23); 7.2273 (1.52);
 15 7.2234 (1.28); 7.0579 (1.80); 7.0379 (1.52); 6.9489 (0.97);
 6.9473 (0.98); 6.9303 (1.69); 6.9287 (1.66); 6.9116 (0.81);
 6.9100 (0.78); 5.8459 (0.94); 5.8251 (1.09); 5.8160 (1.08);
 5.7952 (0.98); 4.7535 (4.77); 4.7476 (4.83); 3.5303 (1.35);
 3.5244 (2.79); 3.5186 (1.36); 3.4687 (1.12); 3.4387 (1.19);
 20 3.4251 (1.39); 3.3952 (1.25); 3.2600 (4.66); 2.9524 (1.32);
 2.9315 (1.30); 2.9089 (1.14); 2.8880 (1.13); 2.4336 (5.79);
 2.4292 (7.64); 2.4248 (5.66); 2.3714 (16.00)

The intensity of sharp signals correlates with the height of
 25 the signals in a printed example of an NMR spectrum in cm and
 shows the true ratios of the signal intensities. In the case
 of broad signals, several peaks or the middle of the signal
 and their relative intensities may be shown in comparison to
 the most intense signal in the spectrum.

30

For calibration of the chemical shift of the ¹H NMR spectra we
 use tetramethylsilane and/or the chemical shift of the
 solvent, particularly in the case of spectra measured in DMSO.
 Therefore, the tetramethylsilane peak may but need not occur
 35 in the NMR peak lists.

The lists of the ¹H NMR peaks are similar to the conventional
¹H NMR printouts and thus usually contain all peaks listed in

conventional NMR interpretations.

In addition, like conventional ¹H NMR printouts, they may show solvent signals, signals of stereoisomers of the target
5 compounds, which likewise form part of the subject-matter of the invention, and/or peaks of impurities.

In the reporting of compound signals in the delta range of solvents and/or water, our lists of ¹H NMR peaks show the
10 usual solvent peaks, for example peaks of DMSO in DMSO-d₆ and the peak of water, which usually have a high intensity on average.

The peaks of stereoisomers of the target compounds and/or
15 peaks of impurities usually have a lower intensity on average than the peaks of the target compounds (for example with a purity of >90%).

Such stereoisomers and/or impurities may be typical of the
20 particular preparation process. Their peaks can thus help to identify reproduction of our preparation process with reference to "by-product fingerprints".

An expert calculating the peaks of the target compounds by
25 known methods (MestreC, ACD simulation, but also with empirically evaluated expected values) can, if required, isolate the peaks of the target compounds, optionally using additional intensity filters. This isolation would be similar to the relevant peak picking in conventional ¹H NMR
30 interpretation.

Further details of ¹H NMR peak lists can be found in Research Disclosure Database Number 564025.

35 Use Examples

Phytophthora test (tomato) / protective

Solvents: 49 parts by weight of N,N-dimethylformamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable active ingredient formulation, 1 part by weight of active ingredient is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.
- 10 To test for protective efficacy, young tomato plants are sprayed with the active ingredient formulation at the stated application rate. 1 day after the treatment, the plants are inoculated with a spore suspension of *Phytophthora infestans*, and then left to stand at 100% rel. humidity and 22°C for 24
- 15 h. Subsequently, the plants are placed in a climate chamber at approx. 96% relative air humidity and a temperature of approx. 20°C.

Evaluation follows 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an

20 efficacy of 100% means that no infection is observed.


In this test, the following inventive compounds show, at an active ingredient concentration of 100 ppm, an efficacy of 70%

25 or more:

Ex.	Eff. %
I-1	94
I-2	89
I-3	83
I-4	78
I-8	89

Patentkrav

1. Forbindelser med formel (VIIIa) og (VIIIb)



5 hvor

R³ står for hydrogen,

L² står for en direkte binding,

R¹ står for phenyl, som indeholder 1, 2 eller 3 substituent, idet substituenterne uafhængigt af hinanden er udvalgt i det
 10 mindste en gang af Z⁴ og i givet fald fra følgende liste:
 fluor, chlor, brom, iod, cyano, nitro, hydroxy, amino, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl, 1,2-dimethylethyl, ethenyl, ethinyl, trifluormethyl, difluormethyl, trichlormethyl, dichlormethyl, cyclopropyl,
 15 methoxy, ethoxy, n-propoxy, 1-methylethoxy, 1,1-dimethylethoxy, methylcarbonyl, ethylcarbonyl, methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, 1-methylethoxycarbonyl, 1,1-dimethylethoxycarbonyl, 1-methylcarbonyloxy, methylthio, ethylthio, methylsulfonyl eller
 20 -L³R³,

L³ står for en direkte binding,

R³ står for hydrogen, C₁-C₆-alkyl, C₁-C₆-halogenalkyl, C₃-C₈-cycloalkyl, C₁-C₄-alkylcarbonyl, C₁-C₄-halogenalkylcarbonyl, C₁-C₄-alkoxycarbonyl eller C₁-C₄-halogenalkoxycarbonyl,

25 Z³ står for en phenylgruppe, som kan indeholde op til to substituent, idet substituenterne uafhængigt af hinanden er udvalgt af følgende liste: chlor, brom, iod, fluor, cyano, nitro, hydroxy, amino, -SH, methyl, ethyl, n-propyl, 1-methylethyl, 1,1-dimethylethyl, ethenyl, propen-2-yl, ethinyl,
 30 propin-2-yl, trifluormethyl, difluormethyl, methoxymethyl, methylcarbonyl, ethylcarbonyl, trifluormethylcarbonyl, methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, 1-methylethoxycarbonyl, 1,1-dimethylethoxycarbonyl, methoxy, ethoxy, n-propoxy, 1-methylethoxy, 1,1-dimethylethoxy,
 35 trifluormethoxy, ethenyloxy, 2-propenyloxy, ethinyloxy, 2-propinyloxy, methylthio, ethylthio, trifluormethylthio,

methylsulfonyl, ethylsulfonyl, propylthionyl, 1-
 methylethylthio, trifluormethylsulfonyl, methylamino,
 ethylamino, n-propylamino, 1-methylethylamino, 1,1-
 dimethylethylamino eller dimethylamino, eller
 5 Z³ står for naphthalenyl,
 R¹³ og R¹⁴ står ens eller forskelligt uafhængigt af hinanden for
 hydrogen, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl
 eller 1,1-dimethylethyl,
 Z⁴ står for -formyl, methoxymethoxy, 2-methoxyethoxy, allyloxy,
 10 2-fluorprop-2-en-1-yloxy, 2-chlorprop-2-en-1-yloxy, 3-
 chlorprop-2-en-1-yloxy, 2-bromprop-2-en-1-yloxy, 2-methylprop-
 2-en-1-yloxy, 3,3-dichlorprop-2-en-1-yloxy, 3,3-dichlor-2-
 fluorprop-2-en-1-yloxy, but-2-en-1-yloxy, but-3-en-2-yloxy,
 but-3-en-1-yloxy, 3-chlorbut-2-en-1-yloxy 3-methylbut-2-en-1-
 15 yloxy, 4,4,4-trifluorbut-2-en-1-yloxy, prop-2-in-1-yloxy, 3-
 chlorprop-2-in-1-yloxy, 3-bromprop-2-in-1-yloxy, but-2-in-1-
 yloxy, pent-2-in-1-yloxy, 2-fluor-2-methylpropanoyloxy, 3,3,3-
 trifluorpropanoyloxy, cyclopropylcarbonyloxy,
 cyclohexylcarbonyloxy, (1-chlorcyclopropyl)-carbonyloxy, but-
 20 2-enoyloxy, acryloyloxy, cyanomethoxy, methylsulfonyloxy,
 ethylsulfonyloxy, trifluormethylsulfonyloxy,
 cyclopropylsulfonyloxy, 2-methoxyethoxymethyl, allyloxymethyl,
 Prop-2-in-1-yloxymethyl, methylsulfonylmethyl,
 methylcarbonylaminomethyl, methylsulfonylaminomethyl, -
 25 C(=NOR⁷)R⁸, dimethylaminosulfonyl, ethylaminosulfonyl,
 trimethylsilylethynyl, diethylaminosulfonyl,
 methylaminosulfonyl, trimethylsilyloxy, trimethylsilylprop-2-
 in-1-yloxy, trifluormethylamino, dimethylaminocarbonylamino, -
 C(=O)OH, -NHC(=O)H, -C(=O)NH₂, -C(=S)NR¹³R¹⁴ 1,1-
 30 dimethylethylcarbonylamino, chlormethylcarbonylamino,
 Trifluormethylcarbonylamino, 1,1-dimethylethoxycarbonylamino,
 ethylcarbonylamino, 1-methylethoxycarbonylamino,
 trifluormethylcarbonylamino, methylcarbonylamino,
 methoxycarbonylamino, ethoxycarbonylamino, iso-
 35 propoxycarbonylamino, 1-methylethylcarbonylamino,
 methylsulfonylamino eller phenylsulfonylamino, 3-bromprop-2-
 en-1-yloxy eller -L⁴Z³,
 L⁴ står for -C(=O)O- eller -OCH₂C≡C-,

R⁷ står for hydrogen, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl eller 2-methylpropyl,

R⁸ står for hydrogen, methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl eller 2-methylpropyl samt salte

5 heraf.

2. Forbindelse ifølge krav 1, idet R¹ står for 6-chlor-2-[(methylsulfonyl)oxy]phenyl.