

AUSTRALIA

NOTICE OF ENTITLEMENT
(To be filed before acceptance)

I/We Unilever PLC
of Unilever House, Blackfriars, London EC4P 4BQ, England
being the applicant(s) and nominated person(s) in respect of this Application, No.
14680/92

- (a) I/We have entitlement to the grant of the patent from the actual inventor(s)
listed on the Patent Request Form as follows:

The applicant is the assignee of the employer of the actual inventors namely
Unilever UK Central Resources Limited. Unilever UK Central Resources
Limited was entitled to have assigned to it a patent granted to any of the
actual inventors in respect of the invention.

- (b) The person nominated for the grant of the patent is the assignee from the
applicants of the application listed in the declaration under Article 8 of the
PCT.
- (c) The basic application listed in the declaration made under Article 8 of the PCT
is the first application made in a Convention country in respect of the
invention.

Signed for and on behalf of
Unilever PLC by
Unilever Australia Limited



.....
B.F. JONES, Company Secretary

26 October, 1995
Date



AU9214680

(12) PATENT ABRIDGMENT (11) Document No. AU-B-14680/92
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 669478

- (54) Title
PLANT PROMOTER INVOLVED IN CONTROLLING LIPID BIOSYNTHESIS IN SEEDS
- International Patent Classification(s)
- (51)⁵ C12N 015/82 A01H 005/10 C11B 001/00 C12N 015/29
- (21) Application No. : 14680/92 (22) Application Date : 08.04.92
- (87) PCT Publication Number : WO92/18634
- (30) Priority Data
- (31) Number (32) Date (33) Country
91303098 09.04.91 EP EUROPEAN PATENT OFFICE (EPO)
- (43) Publication Date : 17.11.92
- (44) Publication Date of Accepted Application : 13.06.96
- (71) Applicant(s)
UNILEVER PLC
- (72) Inventor(s)
JACQUELINE DE SILVA; RICHARD SAFFORD; STEPHEN GLYN HUGHES
- (74) Attorney or Agent
UNILEVER AUSTRALIA LIMITED , B F Jones, Private Mailbag 2, EPPING NSW 2121
- (56) Prior Art Documents
DE SILVA ET AL PLANT MOL.BIOL.VOL.14,1990. P537-548
- (57) Claim

1. A recombinant DNA construct containing a promoter that is capable of acting as a seed-specific plant promoter, said promoter controlling expression of a heterologous gene placed under control of said promoter in concert with the fatty acid or lipid biosynthesis in a plant cell, such that the polypeptide resulting from the expression of the heterologous gene can play a role in the fatty acid or lipid biosynthesis in the plant cell, and wherein said promoter comprises at least the 291 bp polynucleotide of clone ACP05 given in the specification, i.e.

1

```

AGATCTGATT GGTAAGATAT GGGTACTGTT TGGTTTATAT GTTTGGACTA 50
TTCAGTCACT ATGGCCCCCA TAAATTTTAA TTCGGCTGGT ATGTCTCGGT 100
TAAGACCGGT TTGACATGGT TCATTTTCAGT TCAATTATGT GAATCTGGCA 150
CGTGATATGT TTACCTTCAC ACGAACATTA GTAATGATGG GCTAATTTAA 200
GACTTAACAG CCTAGAAAGG CCCATCTTAT TACGTAACGA CATCGTTTAG 250
AGTGCACCAA GCTTATAAAT GACGACGAGC TACCTCGGGG C 291

```

OPI DATE 17/11/92 APPLN. ID 14680/92
AOJP DATE 27/12/92 PCT NUMBER PCT/GB92/00627



AU9214680

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12N 15/82, A01H 5/10 C11B 1/00, C12P 21/02 // C12N 15/55, 9/16, 15/29</p>	<p>AI</p>	<p>(11) International Publication Number: WO 92/18634 (43) International Publication Date: 29 October 1992 (29.10.92)</p>
<p>(21) International Application Number: PCT/GB92/00627 (22) International Filing Date: 8 April 1992 (08.04.92) (30) Priority data: 91303098.7 9 April 1991 (09.04.91) EP (34) Countries for which the regional or international application was filed: GB et al. (71) Applicant (for AU CA GB only): UNILEVER PLC [G3/ GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU CA GB US): UNILEVER N.V. [NL/NL]; Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).</p> <p style="text-align: center; font-size: 2em; font-weight: bold;">069478</p>		<p>(72) Inventors; and (75) Inventors/Applicants (for US only): DE SILVA, Jacqueline [GB/GB]; 10 Tithe Barn, Felmersham, Bedford MK43 7JF (GB). SAFFORD, Richard [GB/GB]; 10 Furness Close, Bedford MK41 8RN (GB). HUGHES, Stephen, Glyn [GB/GB]; 70 Ashdon Road, Saffron Walden, Cambridgeshire CB10 2AL (GB). (74) Agent: KEITH W. NASH & CO.; Pearl Assurance House, 90-92 Regent Street, Cambridge CB2 1DP (GB). (81) Designated States: AT (European patent), AU, BE (Euro- pean patent), CA, CH (European patent), DE (Euro- pean patent), DK (European patent), ES (European pa- tent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (Euro- pean patent), MC (European patent), NL (European pa- tent), SE (European patent), US.</p> <p>Published With international search report.</p>
<p>(54) Title: PLANT PROMOTER INVOLVED IN CONTROLLING LIPID BIOSYNTHESIS IN SEEDS</p>		
<p>(57) Abstract</p> <p>A new seed-specific plant promoter is provided, capable of expressing a gene placed under control of said promoter before or during fatty acid or lipid biosynthesis in plant cells. In nature it occurs in the acyl carrier protein (ACP) gene. This opens the possibility of modifying the fatty acid synthesis in plants, which may result in changing the triacylglycerol composition of oil-containing seeds. Another option is the production of a desired protein in plants, either to improve the nutritional value of the seeds, or for the production of specific proteins that can be isolated from the fruits of plants.</p>		

TITLE: PLANT PROMOTER INVOLVED IN CONTROLLING LIPID
BIOSYNTHESIS IN SEEDS

INTRODUCTION

5

Technical Field

The invention relates to transforming plant cells for modifying the seed-specific production of fatty acids resulting in a changed fatty acid composition of seed oils. In particular the invention provides a new promoter isolated from a seed-specific acyl carrier protein (ACP) gene present in *Brassica napus* (oil seed rape).

15

Background and Prior Art

During the last decade methods have been developed for transforming plants by introducing genes into plants which on expression give new or improved properties to the resulting transformed plants. One of these methods is the use of the bacterium *Agrobacterium tumefaciens* for introducing the desired gene into the chromosome of the plant to be transformed. Many articles have been published on this technique. For a general introduction reference is made to Chapter 13 (Genetic Engineering of Plants by Using Crown Gall Plasmids) on pages 164-175 of the book 'Recombinant DNA, A Short Course' by James D. Watson, John Tooze and David T. Kurtz, published by Scientific American Books in 1983 and distributed by W.H. Freeman and Company, New York, U.S.A.

30

According to European patent specification EP-A2-0255378 (CALGENE, INC.), published on 3 Feb. 1988 with claimed priority date of 31 July 1986, a so-called transcriptional initiation region of the napin gene is identified and isolated from plant cells, and used to prepare expression cassettes which may then be inserted into plant cells for seed specific transcription. It is stated in that patent specification that the method may

35

be applied in conjunction with modifying fatty acid production in seed tissue.

From that EP-A-0255378 the following passages are quoted:

5 on page 3, lines 6-9:

" Transcriptional initiation regions of particular interest are those associated with the *Brassica napus* or *campestris* napin genes, acyl carrier proteins, genes that express from about day 7 to day 40 in seed, particularly having maximum expression from about day 10 to about day 20, where the expressed gene is not found in leaves, while the expressed product is found in seed in high abundance. "

15

on page 4, lines 11-22:

" The constructs may be used to modify the fatty acid composition in seeds, that is changing the ratio and/or amounts of the various fatty acids, as to length, unsaturation, or the like. These results can be achieved by providing for reduction of expression of one or more endogenous products, particularly enzymes or cofactors, by producing a transcription product which is complementary to the transcription product of a native gene, so as to inhibit the maturation and/or expression of the transcription product, or providing for expression of a gene, either endogenous or exogenous, associated with fatty acid synthesis. Expression products associated with fatty acid synthesis include acyl carrier protein, thioesterase, acetyl transacylase, acetyl-coA carboxylase[m], ketoacyl-synthases, malonyl transacylase, stearyl-ACP desaturase, and other desaturase enzymes. "

35 on page 4, lines 57-64:

" Expression cassettes of particular interest include transcriptional regions from napin genes, particularly *Brassica napin* genes, more particularly

Brassica napus or *Brassica campestris* genes, regulating structural genes associated with lipid production, particularly fatty acid production, including acyl carrier proteins, which may be
5 endogenous or exogenous to the particular plant, such as spinach acyl carrier protein, *Brassica* acyl carrier protein, acyl carrier protein, either *napus* or *campestris*, *Cuphea* acyl carrier protein, acetyl transacylase, malonyl transacylase, β -ketoacyl
10 synthases I and II, thioesterase, particularly thioesterase II, from plant, mammalian, or bacterial sources, for example rat thioesterase II, acyl ACP, or phospholipid acyl desaturases. "

It should be noted that the time periods indicated in
15 the passage quoted from page 3, lines 6-9, sometimes mentioned as Days After Flowering (DAF) is not always a precise unit, because for the same plants it can differ depending on the location and conditions of growth. Thus, DAF should not be used as an absolute but as a
20 comparative parameter. Therefore, when comparing results of this nature from experiments done at different times or different locations, one has to be particularly careful in drawing conclusions on the basis of differences in DAF values.

25 Furthermore, with experiments in tobacco the rate of development of individual seed pods even on a single plant may be variable, i.e. the earliest pods developing fastest, in which case it is not possible with tobacco to use DAF for determining accurately the developmental
30 stage of a seed. In such a situation seeds have to be staged using morphological characteristics. This is expanded later on in this specification (see Example 1.c.5 and Example 3, experiment 2).

Example I of EP-A-0255378 discloses a construct
35 comprising the structural gene encoding spinach leaf acyl carrier protein under control of the napin promoter from *B. napus*. No evidence was given that the protein was formed: by means of Northern blots only the presence

of the corresponding mRNA was made plausible in embryos but not in leaves indicating seed-specific expression of the spinach leaf ACP gene.

Example II discloses construction of a *B. campestris* napin promoter cassette. It was only suggested that a gene involved in the fatty acid synthesis might be inserted into that cassette. No experimental details on such insertion or expression of such gene were given.

According to Example III:

10 " Other seed-specific promoters may be isolated from genes encoding proteins involved in seed triacyl-glycerol synthesis, such as acyl carrier protein from *Brassica* seeds. Immature seed[s] were collected from *Brassica campestris* cv."R-500," a self-compatible
15 variety of turnip rape. Whole seeds were collected at stages corresponding approximately to 14 to 28 days after flowering. preparation of a cDNA bank To probe the cDNA bank, the oligonucleotide was synthesized This synthetic DNA
20 molecule will hybridize at low stringencies to DNA or RNA sequences coding for the amino acid sequence (ala-ala-lys-pro-glu-thr-val-glu-lys-val). This amino acid sequence has been reported for ACP isolated from seeds of *Brassica napus* (A.R. Slabas et al., 7th International Symposium of the Structure and Function of
25 Plant lipids, University of California, Davis, CA, 1986); ACP from *B. campestris* seed is highly homologous. DNA sequence analysis of two DNA clones showing obvious hybridization to the oligonucleotide probe indicated that one, designated pCGN1Bcs, indeed
30 coded for an ACP-precursor protein by the considerable homology of the encoded amino acid sequence with ACP proteins described from *Brassica napus* (A.R. Slabas et al., 198[6] *supra*). Similarly
35 to Example II, the ACP cDNA clone can be used to isolate a genomic clone from which an expression cassette can be fashioned in a manner directly analogous to the *B. campestris* napin cassette. "

Under the next heading Other Examples it was stated:

" Ninety-six clones from the 14-28 day post-anthesis *B. campestris* seed cDNA library (described in the previous example) were screened Other seed-specific genes may also serve as useful sources of promoters. Without knowing their specific functions, yet other cDNA clones can be classified as to their level of expression in seed tissues, their timing of expression (i.e., when post-anthesis they are expressed) Clones fitting the criteria necessary for expressing genes relating to fatty acid synthesis or other seed functions can be used to screen a genomic library for genomic clones which contain the 5' and 3' regulatory regions necessary for expression. The non-coding regulatory regions can be manipulated to make a tissue-specific expression cassette in the general manner described for the napin genes in previous examples.

One example of a cDNA clone is EA9. It is highly expressed in seeds and not leaves from *B. campestris*. Northern blot analysis of mRNA isolated from day 14 seed, and day 21 and 28 post-anthesis embryos using a 700 bp *EcoRI* fragment of EA9 as a probe shows that EA9 is highly expressed at day 14 and expressed at a much lower level at day 21 and day 28. The partial sequence provided here for clone EA9 (Figure 3) can be used to synthesize a probe which will identify a unique class of *Brassica* seed-specific promoters!"

In related European patent specification EP-A2-0255377 (CALGENE, INC.), also published on 3 Feb. 1988 with claimed priority date of 31 July 1986, the DNA sequences of structural genes encoding ACP of spinach and *Brassica campestris* are provided, which can be used for production of ACPs as an end product or may enhance seed oil production in plant seed. Also described are napin promoters of *B. napus* and *B. campestris* substantially limiting expression of the ACP genes to seed tis-

sue, which promoters are the same as described in EP-A2-0255378.

5 However, according to M.A. Post-Beittenmiller *et al.*
in *The Plant Cell* 1 (1989) 889-899 no significant
alterations in leaf lipid biosynthesis were detected by
lipid analysis, neither in level nor in composition,
when tobacco was transformed with a chimaeric gene
10 consisting of the tobacco ribulose-1,5-biphosphate
carboxylase promoter and transit peptide and the
sequence encoding mature spinach ACP-I. They further
showed that the mature spinach ACP-I gene was expressed
at higher levels than the endogenous tobacco ACPs as
shown by protein immunoblots. Thus this later work
15 shows that increased production of ACP in plant tissue
need not necessarily result in an altered fatty acid
composition.

 This finding is in agreement with statements made by
20 V.C. Knauf in "The application of genetic engineering
to oilseed crops" published in *TIBTECH* 5 (Feb 1987) 40-
47, in which he mentioned many possibilities why "a
'typical' project" for altering the fatty acid composi-
tion of rape seed might fail due to the complexity of
25 lipid biosynthesis in plant tissues.

 In a paper entitled "Plastid-localised seed acyl-
carrier protein of *Brassica napus* is encoded by a
distinct, nuclear multigene family" R. Safford *et al.*
30 provide in *Eur. J. Biochem.* 174 (1988) 287-295 the first
insight into the origin, structure and expression of
genes co-ordinating fatty acid biosynthesis in oil-
bearing seeds. It reveals seed ACP to be localised
within plastid bodies and to be encoded in nuclear DNA,
35 being synthesised as a precursor containing an N-termi-
nal extension sequence which presumably directs import
of the protein into the plastids. Analysis of several
cDNA clones revealed sequence heterogeneity and thus

evidence for an ACP multigene family. Further experiments showed that at least some of these genes were not expressed in leaf tissue. In Fig. 8A of this publication not more than 56 nucleotides of the 5'-non-coding regions upstream of the ACP start codon are given. The promoters could not be determined because the work was done with cDNA clones from mRNA as starting material, which contains only transcribed sequences, instead of chromosomal DNA, the latter containing transcription regulating sequences including the promoter region.

In a subsequent paper of J. de Silva *et al.* in Plant Molecular Biology 14 (1990) 537-548 the same group described the isolation and sequence analysis of two genomic clones encoding seed-expressed acyl carrier protein genes from *Brassica napus*. The latter paper discloses that the transcription start site is situated 69 bp upstream of the (ATG) start-codon of the structural gene.

Thus, the 5' regions shown in both papers are less than 70 nucleotides long and describe the DNA sequences downstream of the transcription start site. Therefore, they neither comprise a promoter region nor a regulatory region conferring seed-specific temporal regulation of gene transcription.

Summarizing the prior art, EP-A-0255378 describes the isolation and use of napin promoters of *B. napus* and *B. campestris* and suggests how other promoters like the ACP promoters can be isolated, EP-A-0255377 describes the same napin promoters, and the R. Safford *et al.* (1988) and the J. de Silva *et al.* (1990) publications describe only less than 70 nucleotides of the 5' region preceding the structural gene encoding the seed-specific ACP of *B. napus*, so that neither of these publications discloses the nucleotide sequence of an ACP promoter.

SUMMARY OF THE INVENTION

The invention provides a novel seed-specific ACP promoter isolated from *Brassica napus* and DNA constructs which can be employed in manipulating plant cells to provide for seed-specific transcription.

According to one embodiment of the invention, a desired gene encoding a protein active in the biosynthetic pathway for fatty acid production is placed under control of the novel ACP promoter and introduced into plant genomes to provide for seed-specific transcription, whereby the gene may be homologous or heterologous to the plant genome. The constructs provide for modulation of endogenous products or of their production, as well as for production of heterologous products. In order to be capable of influencing the fatty acid biosynthesis the protein to be produced must be targeted to the correct intracellular site of the cell, i.e. the plastids. This requires that a chimaeric DNA construct be made, in which the desired gene is linked to a suitable transit sequence such that the resulting chimaeric protein will be targeted to the plastid, thus enabling release of the mature form of the protein corresponding to the desired gene within the plastid. Some examples of proteins active in the biosynthetic pathway for fatty acid production are acetyl-coA carboxylases, acetyl transacylases, ACPs, desaturases, elongases, enoyl-reductases, β -keto-reductases, ketoacyl-synthases, malonyl transacylases, and thioesterases.

According to another embodiment of the invention a gene encoding a desired protein is placed under control of the novel ACP promoter. Such a desired protein can be any protein the production of which in plants and, optionally its subsequent isolation, is desirable. As examples of such proteins can be mentioned

- (1) enzymes to be used in food processing, e.g. guar α -galactosidase, thaumatin, chymosin,
- (2) proteins that can inhibit the formation of anti-nutritive factors,

- (3) pharmaceutically active proteins, e.g. blood factors, interferon, hormones, human serum albumin, and
- 5 (4) plant proteins with a more desirable amino acid composition, e.g. one with a higher lysine content than occurring in the non-transformed plant.

Depending on the influence of the desired protein on the cell metabolism it can be allowed to reside in the cytoplasm, or it can be targeted to the plastids, in which case the gene encoding the desired protein should be linked to a target sequence as described above. Alternatively it can be targeted to other organelles within the cell, for which other targeting sequences are required. Several other targeting sequences are described before; see for example G. Van den Broek et al. in *Nature* 313 (31 jan 1985) 358-363 on targeting of a foreign protein to chloroplasts, and a review on protein targeting by R.J. Ellis & C. Robinson in *Advances in Botanical Research* 14 (1987/8) 1-24 published by Academic Press Ltd. (ISBN 0-12-005914-2) showing targeting to the chloroplasts, the mitochondria and the nucleus of plants.

25 DESCRIPTION OF THE SPECIFIC EMBODIMENTS

The present invention is based upon carrying out research on the factors which may influence the expression of a protein involved in the fatty acid synthesis in the seeds of plants like *Brassica napus*. This research was needed in a project directed to changing the fatty acid profile of seed oils such as rape. In order to change the fatty acid composition of a storage lipid in plants or to induce production of a desired polypeptide or protein in seeds of a plant by means of genetic engineering, the expression of the relevant genes has to be controlled in the following manner:

- 35
- i) The expression of the desired gene should be confined to seed tissue only, at the correct

developmental stage and at an appropriate level.

- ii) Specifically in the case of modifying the fatty acid profile of the seed lipids:
 - 5 a) Transport of the resulting protein into the fatty acid synthesizing sub-compartment of the cell (so-called plastid) in a biologically active form, and
 - 10 b) Transportation of synthesized fatty acids out of the plastid and their subsequent conversion to triacylglycerides.

One way to achieve step i) is to utilize a promoter region isolated from an endogenous gene involved in the fatty acid biosynthesis. Such a promoter would activate gene expression before or at the stage that fatty acid biosynthesis in the seed occurs.

To obtain controlled expression of a desired gene in plants, e.g. rape seed, the structural gene can, for example, be placed under control of a promoter region isolated from an endogenous rape seed lipid biosynthetic gene. Although some seed-specific promoters were already known as described above, there have been no reports describing the isolation and functional characterisation of such regulatory DNA sequences of genes involved in the biosynthesis of seed lipids in plants.

An example of one such gene is the gene encoding ACP, which is a key component of the fatty acid biosynthetic machinery of plants, serving as a component of the fatty acid synthetase (FAS) and is also involved in desaturation and acyl transfer reactions (P.K. Stumpf *et al.* Fatty acid biosynthesis in higher plants. In: Fatty Acid Metabolism and Its Regulation. Elsevier Press, Amsterdam, Numa S. (ed), (1984) 155-179).

In the course of the research resulting in the

present invention a promoter was selected belonging to the chromosomal ACP05 gene described by J. de Silva et al. (1990) *supra*, which corresponds to the ACP-encoding 29C08 cDNA described earlier by R. Safford et al. (1988) *supra*. The latter publication showed the isolation and characterisation of cDNA clones encoding rape embryo ACP and that ACP is synthesized as a precursor containing an N-terminal transit sequence. The latter can be used as a 'tool' to transport products of foreign genes into fatty acid synthesising plastids.

Functional analysis of the promoter element present in the 5' upstream region of the ACP gene was carried out using the β -glucuronidase (GUS) reporter gene and tobacco transformation.

A 1.4 kb 5' upstream fragment of rape ACP05 gene (AP1 promoter) was fused to the β -glucuronidase (GUS) reporter gene and transferred, via *Agrobacterium* infection, into tobacco. Analysis of leaf and seed tissue from 15 transgenic tobacco plants showed the level of GUS expression to increase through seed development to a value that was, on average, approximately 100x higher than that observed in leaf. Analysis of control plants transformed with constructs containing the GUS gene linked to a constitutive plant promoter (CaMV 35S) showed similar levels of GUS expression in leaf and all stages of seed development. These results demonstrate that the isolated AP1 promoter sequence functions to control expression of the GUS gene in transgenic tobacco in the desired seed specific and developmental manner.

In a comparative study with CaMV 35S and AP1 promoters, the level of GUS expression obtained with the AP1 promoter at the stage of maximum seed lipid synthesis was equivalent to that obtained with the powerful plant promoter CaMV 35S. Therefore, the AP1 promoter can be considered as a 'strong', seed-specific promoter. The CaMV 35S promoter itself cannot be used for this

purpose, because it is a constitutive promoter that produces the protein throughout the plant, which is undesirable.

5 In order to define more precisely the essential part of the rape ACP05 promoter, including any specific, developmental regulation sequence(s), a deletion analysis was performed. Chimaeric gene constructs containing 1.4 kb, 0.92 kb and 0.29 kb, respectively, of the region
10 upstream from the transcription initiation start site of the ACP05 gene fused to the β -glucuronidase (GUS) reporter gene were transferred into tobacco. For each construct 10 transformed plants were analysed for mode of GUS expression. No significant differences were observed
15 in either the level or tissue distribution of GUS activity in plants transformed with the 'deleted' versions of the ACP promoter compared to the 'original' 1.4 kb construct. It was therefore concluded that the DNA sequences which determine the level of transcription and seed specificity of the ACP05 gene reside within the
20 0.29 kb region immediately upstream of the start site of transcription of the gene, which start site was described in the J. de Silva et al. (1990) publication *supra*.

25 This DNA fragment was sequenced and the resulting 291 bp DNA sequence is

```

1
AGATCTGATT GGTAAGATAT GGGTACTGTT TGGTTTATAT GTTTTGACTA 50
30 TTCAGTCACT ATGGCCCCCA TAAATTTTAA TTCGGCTGGT ATGTCTCGGT 100
TAAGACCGGT TTGACATGGT TCATTTTCAGT TCAATTATGT GAATCTGGCA 150
CGTGATATGT TTACCTTCAC ACGAACATTA GTAATGATGG GCTAATTTAA 200
GACTTAACAG CCTAGAAAGG CCCATCTTAT TACGTAACGA CATCGTTTAA 250
AGTGCACCAA GCTTATAAAT GACGACGAGC TACCTCGGGG C 291.
35

```

To compare the temporal regulation of the ACP05 promoter in relation to other seed-specific plant promoters it was determined if the respective promoter ele-

- ments from the lipid biosynthetic gene (ACP05) and from two storage protein genes are indeed capable of conferring differential gene expression during seed development. This was investigated by transferring AP1-GUS, napin-GUS and cruciferin-GUS chimaeric genes into tobacco and monitoring their expression at various stages throughout seed development, whereas similar experiments with expression of these combinations in rape are in progress.
- 10 The 1.4 kb 5' upstream fragment of the ACP05 gene, a 1.1 kb upstream fragment of a rape napin gene and a 2 kb upstream fragment of a rape cruciferin gene were fused to the β -glucuronidase (GUS) reporter gene and transferred into tobacco via *Agrobacterium* infection.
- 15 GUS activity was assayed in seed from two stages of development (mid-mature and mature) and in leaf tissue. All three rape promoters were found to regulate GUS expression in a seed-specific manner, but they differed in their mode of temporal regulation during seed development. Thus with AP1-GUS transformed plants, GUS activity was maximal in mid-mature seeds, whilst in plants transformed with the two storage protein promoter constructs, GUS activity was maximal in mature seed (see Fig. 12).
- 20
- 25 The maximal level of GUS expression obtained in tobacco seeds with both the ACP and cruciferin promoters was similar to that obtained with the powerful constitutive cauliflower mosaic virus 35S promoter (average of 10 plants/construct), and approximately 3x that obtained with the napin promoter.
- 30

A second study was carried out to determine more precisely the nature of the temporal differences conferred on GUS expression during seed development by a lipid biosynthetic gene promoter (AP1) and a storage protein gene promoter (cruciferin). Seed from the highest expressing AP1GUS and CRUGUS plants of the first study were used to propagate new plants. Flowers were

tagged on opening and 7-20 days after flowering seed pods were collected and the GUS activity of seed extracts determined. In plants transformed with the ACP promoter-GUS fusion, activity peaked at 11-12 DAF corresponding to the maximum rate of lipid synthesis in the seed. In plants transformed with the cruciferin promoter-GUS fusion, activity peaked at 16-19 DAF, which corresponded to the stage in seed development when protein content was rapidly increasing but the rate of lipid synthesis was decreasing (see Fig. 14).

The results of these findings have important implications for a programme aimed at modifying rapeseed oil composition via genetic engineering. In order to be successful a transferred gene, encoding a protein that may perturb fatty acid biosynthesis, should be expressed coincidentally with the lipid synthetic phase of seed development. The present results clearly demonstrate that in order to achieve that objective, the expression of the transferred gene would have to be controlled by a seed lipid biosynthetic gene promoter, e.g. of an ACP gene. Fusion of the gene to a seed storage protein gene promoter, e.g. of a napin gene, would result in maximal expression of the transferred gene after the most active phase of lipid synthesis and would, therefore, not likely result in a significant perturbation of the fatty acid profile.

Thus the present invention provides a new seed lipid biosynthetic gene promoter, and its use in expressing a gene at such stage in the seed development that the protein or polypeptide formed can influence the biosynthetic formation of fatty acids and lipidic esters thereof. Although the invention is illustrated on the basis of a nature-identical plant promoter isolated from a particular *Brassica napus* species, it will be clear to a skilled person that other nature-identical seed-specific promoters that are also capable of expressing genes

in concert with the fatty acid or lipid biosynthesis can be isolated using the teachings of the present specification. The expression "in concert with" used in this specification means that the gene is expressed at such place and at such time that the protein resulting from the expression can play a role in the fatty acid or lipid biosynthesis in the plant cell. Specifically said promoter has the ability to express a gene placed under control of said promoter at a stage before or during fatty acid or lipid biosynthesis.

Moreover, by using known techniques for DNA modification, other DNA sequences can be prepared that can be tested for their ability to promote gene expression in a seed-specific and temporal fashion.

Thus in a broader sense the expression "promoter that is capable of acting as a seed-specific plant promoter" covers both nature-identical promoters and modifications thereof that are also active as seed-specific plant promoters, as well as otherwise designed promoters being active as seed-specific plant promoters.

Therefore, the present invention relates to a recombinant DNA construct containing a promoter that is capable of acting as a seed-specific plant promoter, said promoter being also capable of expressing a gene placed under control of said promoter in concert with the fatty acid or lipid biosynthesis in a plant cell.

The expression "recombinant DNA construct" is used in this specification to exclude similar DNA sequences in their natural environment. It indicates that human intervention is used to prepare the construct, which can then be incorporated into plants and stably inherited in their progeny. For example, the nature-identical promoter can be combined with a structural gene different from the gene it controls in nature. Or it can be combined with an enhancer to increase the level of transcription of its natural gene. The level of production of the protein encoded by its natural

structural gene can be reduced by combining the promoter with anti-sense DNA, or by combining with a truncated structural gene. These methods are described in the prior art. Of course, one can also apply these methods
5 to a heterologous gene.

Or it can be combined with antisense DNA to reduce the level of production of the protein encoded by its natural structural gene.

10 More specifically the promoter in said DNA construct comprises at least the 291 bp polynucleotide of clone ACP05 given above. This DNA sequence was determined from the 1 kb *Pst*I-*Bgl*III 5' upstream fragment of the rape ACP05 gene given in Fig. 1. The larger 1.4 kb *Bam*HI-
15 *Bgl*III 5' upstream fragment of the genomic rape ACP05 gene given in Fig. 1, called the AP1 promoter, was taken up in plasmid pAP1GUS present in *E. coli* JM101- /pAP1GUS (NCIMB 40396).

20 Another embodiment of the invention is the use of a DNA construct containing a seed-specific promoter according to the invention for transforming plant cells, preferably for modifying the seed-specific biosynthesis of fatty acids. Preferably the plant cells are subsequently
25 grown to whole plants in which the modified biosynthesis of fatty acids occurs specifically in the seeds. A practical embodiment of such use is a process of transforming plant cells, in which a DNA construct containing a seed-specific promoter according to the inven-
30 tion is introduced into a transformable plant cell in such a way, that after growing the resulting transformed plant cell to whole plants the structural gene forming part of said gene controlled by the introduced seed-specific and temporally regulating plant promoter is
35 expressed in concert with the fatty acid or lipid biosynthesis in the seeds of the plants, thereby producing the protein corresponding to said structural gene. In one preferred way of carrying out such process said

structural gene encodes a protein required for the seed-specific biosynthesis of fatty acids or the corresponding lipids.

Such process can result in a method for modifying the formation of vegetable seed oils, which method comprises
5 growing a plant cell via plantlet to a plant bearing seed and harvesting the resulting seed containing a vegetable oil with modified composition, whereby the cells of said plant cell or plant or seed comprise a DNA
10 construct according to the invention, in particular, if said DNA construct comprises a gene encoding a protein active in the biosynthetic pathway for fatty acid production or lipid formation, and the protein or proteins introduced in this way can modify the biosynthetic
15 pathway.

If a promoter essentially consisting of a seed-specific ACP promoter, preferably one originating from *Brassica napus* and a structural gene encoding ACP is used, the latter should differ from the wild-type gene.
20 This embodiment may result in seeds comprising a DNA construct according to the invention, but if said DNA construct contains a seed-specific plant promoter homologous to the seed, said DNA construct should be present in the genome of said seed at a site other than the
25 natural site for said promoter.

Another embodiment of the invention relates to a seed, preferably of the *Brassica* family, wherein said DNA construct also contains a DNA sequence of interest encoding an exogenous protein, whereby the DNA sequence
30 of interest is under control of the seed-specific plant promoter. However, for practical purpose the exogenous protein should be present in addition to the DNA construct. This latter embodiment is thus mainly directed to a process for producing a desired protein in plant
35 cells, preferably seed cells, which comprises expressing a structural gene encoding said protein, said plant cells containing a recombinant DNA construct according to the invention comprising said structural gene, the

production of said protein. For some applications, e.g. animal feedstuff or for human consumption, it is sufficient if the seeds contain the desired protein, because the seeds can be used as such. For other applications it is desirable that such process is followed by
 5 isolation of said protein from the plant cells.

The invention is illustrated by the following Examples without being limited thereto.

10

Mostly standard methods were used as described in Maniatis, T., Fritsch, E.F., & Sambrook, J.; Molecular Cloning; Cold Spring Harbor Laboratory Publ. (1982). If modifications were used, they are described below.
 15 The following restriction sites are mentioned in this specification:

<i>Bam</i> HI	G↓GATCC	<i>Kpn</i> I	GGTAC↓C
<i>Bgl</i> III	A↓GATCT	<i>Pst</i> I	CTGCA↓G
<i>Eco</i> RI	G↓AATTC	<i>Sal</i> I	G↓TCGAC
20 <i>Hae</i> III	GG↓CC	<i>Sau</i> 3A	↓GATC
<i>Hind</i> III	A↓AGCTT	<i>Sst</i> I	GAGCT↓C

Example 1 - Isolation of *B. napus* ACP promoter

25

a) Isolation of ACP genomic clones

Nuclear DNA was isolated from leaves of field grown *B. napus* plants. Leaves were homogenised in 0.6 M sucrose, 50 mM Tris-HCl pH 8.0, 10 mM MgCl₂, 10 mM β-mercapto-
 30 ethanol, the nuclei pelleted, washed twice in the same homogenisation buffer also containing 1.2% Triton X-100 and lysed in 50 mM Tris-HCl pH 8.0, 20 mM EDTA, and 1% Sarkosyl. DNA was purified by phenol extraction/CsCl centrifugation, partially digested with *Sau*3A and frac-
 35 tionated by sucrose density gradient centrifugation. 15-20 kb DNA was ligated to *Bam*HI-digested lambda EMBL 4 arms previously purified by electroelution from agarose. The ligation mixture was packaged using Gigapack Plus

extracts (ex Stratagene) and propagated in *E. coli* K803. The resultant library was screened with a ^{32}P -labelled RNA probe derived from ACP cDNA clone 29C08 (R. Safford *et al.* (1988) *supra*), cloned into a SP6 vector (ex
5 Amersham), and positive plaques obtained. One of these, designated ACP05, was purified, the DNA was isolated and subcloned into pTZ18R (ex United States Biochemical Corp.) and characterised by restriction mapping and DNA sequencing.

10

b) Structural analysis of ACP05

ACP05 DNA was digested with a range of restriction enzymes and the cleavage sites mapped (Fig. 1). Restricted DNA was Southern blotted to identify DNA fragments with
15 homology to the ACP RNA probe described above. A 1.15 kb *HindIII* fragment, a 5 kb *BamHI* fragment and a 1 kb *SaII* fragment were found to hybridize (see Fig. 1). Overlapping restriction fragments were sub-cloned into M13 vectors and DNA sequenced using a modified bacteriophage
20 T7 DNA polymerase (ex United States Biochemical Corp.). Universal and synthetic oligonucleotide primers were used to obtain a total of about 2233 bp of sequence (Fig. 3). Using dot matrix analysis homology was found
25 between the DNA sequences of ACP05 and rape embryo cDNA clone 29C08, see (Fig. 2), thus confirming that the cloned genomic fragment encoded an ACP gene. Alignment of the nucleotide sequences of genomic clone ACP05 and cDNA clone 29C08 identified 3 intervening sequences (in-
30 trons) within the ACP gene corresponding to nucleotides 1048-1317, 1426-1501, and 1625-1726 of the genomic sequence given in Fig. 3. Complete homology (100%) was found between the exons and the seed expressed cDNA sequence, providing evidence that the ACP05 gene is expressed in the seed.

35

To determine the transcription start site of ACP05 and hence also to define the start of the upstream regulatory sequences of the gene, RNase protection studies

were carried out (see J. de Silva *et al.* (1990) *supra*). A 1 kb *Pst*I-*Sal*I fragment of genomic clone ACP05, spanning the predicted transcription start site of the gene, was ligated into the SP65 transcription vector (ex
5 Amersham) and used as a template to produce full length ³²P-labelled antisense ACP RNA. The antisense probe (about 200,000 dpm) was hybridised to *B. napus* embryo poly A⁺ RNA, isolated according to (R. Safford *et al.* (1988) *supra*), and to an RNA transcript of ACP cDNA
10 clone 29C08 in 50% formamide, 40 mM PIPES pH 6.4, 0.4 M NaCl, 1 mM EDTA at 45°C overnight followed by treatment with RNase A (40 mg/l) and RNase TI (2 mg/l) for 30 minutes at 30°C (dpm = desintegrations per minute, PIPES
= piperazine-N,N'-bis[2-ethanesulfonic acid] or 1,4-
15 piperazine diethanesulfonic acid ex Sigma). RNase activity was destroyed by treatment with proteinase K (125 mg/l) and SDS (0.5%) for 30 minutes at 37°C. Protected RNA was recovered by phenol/chloroform extraction and ethanol precipitation and analysed on a
20 6% acrylamide/urea sequencing gel. The major fragment protected by embryo poly A⁺ RNA was found to be 12 bases longer than the major fragment protected by the cDNA derived transcript.

25 This result therefore identifies the start of transcription of ACP05 as 12 bases upstream from the 5' end of ACP cDNA 29C08, at the first adenine within the sequence GGCATCA, and defines the length of the 5' non-coding sequence as 69 nucleotides (Fig. 3).

30 c) Functional analysis of ACP05 promoter

To evaluate the ability of the 5' upstream region (promoter) of ACP05 to confer seed specific and temporal regulation of gene expression in plants, a transcriptional fusion was made between a 1.4 kb 5' upstream
35 fragment of ACP05 and the reporter gene β -glucuronidase (GUS). The chimaeric gene (AP1GUS) was transferred into tobacco and expression of GUS activity was monitored in leaf and seed tissue of the resultant transgenic plants.

c.1) Construction of pAP1GUS

A 2.5 kb *Bam*HI-*Sst*I restriction fragment of genomic clone ACP05, containing approximately 1 kb of the transcriptional unit of the gene together with 1.5 kb of the 5' upstream region of the gene, was cloned into pTZ18R (see Example 1.a) to form pTZ5BS. This recombinant plasmid was linearised by digestion with *Bam*HI and partially digested with *Bgl*III to produce restriction fragments of 4.9, 3.8, 3.5, 1.4, 1.1 and 0.3 kb in length. The 1.4 kb *Bam*HI-*Bgl*III fragment, containing the promoter region of the gene, was recovered and ligated into *Bam*HI-linearised, phosphatased pTAK vector DNA (Fig. 4) to form pAP1GUS (pTAK is a binary plant transformation vector ex Clontech Labs. Inc., containing a GUS marker gene between the T-DNA border sequences which define the region of DNA capable of transfer to the plant chromosome following agroinfection of damaged plant tissue; in addition to the GUS gene, the T-DNA contains the bacterial neomycin phosphotransferase, NPTII, gene which confers resistance to the antibiotic kanamycin, thus allowing for selection of transformed plant cells). The mix containing pAP1GUS was used to transform commercially available *E. coli* JM101 and recombinant clones were screened to confirm insertion of a single promoter fragment in the correct orientation.

c.2) Transformation of *Agrobacterium* with pAP1GUS

The recombinant pAP1GUS plasmid was mobilised, in a tripartate mating, from *E. coli* JM101 to *Agrobacterium tumefaciens* ACH5/pLBA4404 (see A. Hoekema *et al.*; Nature 303 (1983) 179-181) using commercially available *E. coli* HB101 carrying the helper plasmid pRK2013 (Holsters *et al.*; Mol. Gen. Genet. 163 (1978) 181-187). An overnight culture of the recipient *Agrobacterium* strain and exponential cultures of the donor and helper *E. coli* strains were grown. Of each culture 2 ml was centrifuged and the cells resuspended in 1 ml 10 mM MgSO₄. Equal amounts of

the 3 cell suspensions were mixed, spread on L-agar plates and incubated overnight at 28°C. Resultant bacterial lawn was suspended in 10 mM MgSO₄ and plated onto L-agar containing 50 mg/l rifampicin + 50 mg/l kanamycin. Rifampicin-resistant, kanamycin-resistant colonies were selected, plasmid DNA isolated, transformed back into *E. coli* and characterised by restriction analysis to verify the presence of intact copies of pAP1GUS.

10 c.3) Transformation and regeneration of tobacco

This was performed essentially as described by Horsch *et al.*; Science 227 (1985) 1229-1231. Leaf discs of *Nicotiana tabacum* (var. SR1) with a diameter of 0.5 cm were incubated for 10 min with an overnight culture of *A. tumefaciens* ACH5 containing pLBA4404/pAP1GUS. After blotting dry, discs were placed on *Nicotiana plumbaginifolia* feeder plates, prepared by plating 2 ml of haploid *N. plumbaginifolia* suspension culture (Barfield *et al.*; Plant Cell Reports 4 (1985) 104-107) onto petri dishes containing 20 ml of shoot-inducing medium [0.9% agar, MS salts, 3% sucrose, 0.02 mg/l indole acetic acid (IAA), 1 mg/ml benzylaminopurine (BAP)]. MS-media were described by Murishige, T. & Skoog, F. in Physiol. Plant 15 (1962) 473-497.

25 After 3 days in culture, discs were transferred to shoot-inducing medium containing 500 mg/l cefotaxime and 100 mg/l kanamycin. Shoots regenerating on selective media were excised and placed on minus hormone media (MS salts, 3% sucrose, 0.9% agar) containing 500 mg/l cefotaxime. Once roots had become established, shoots were again excised and placed on minus-hormone medium containing 100 mg/l kanamycin. Plants that rooted on selective media were transferred to soil and grown at 25°C under a 16 hour photoperiod (=AP1GUS plants).

35

c.4) Southern analysis of transformed plants

DNA was isolated from leaves of regenerated AP1GUS plants (Dellaporta; Plant Mol. Biol. Reporter 1 (1983)

19-21), restricted with *Pst*I and *Eco*RI, transferred to nitrocellulose and hybridised to a ³²P-labelled ACP RNA probe. A 3.0 kb *Pst*I-*Eco*RI hybridising fragment was indicative of integration of an intact ACP promoter-GUS cassette into the plant genome, and 15 plants whose DNA digests showed this fragment were chosen for further analysis. Scanning laser densitometry of autoradiographs showed the number of copies of the AP1GUS gene in transformed plants to vary from 1 to 4.

10

c.5) Analysis of transformed plants for GUS enzyme activity

GUS activity was monitored in leaf tissue and in seeds at various stages of development. Under the particular growth conditions used the rate of development of individual seed pods on a single tobacco plant was found to be variable, with the earliest pods frequently developing the fastest. As such it was not possible to use DAF to accurately determine the developmental stage of a seed. For this reason, seeds were staged using morphological characteristics. Five stages of seed development were identified on the basis of seed size and pigmentation:

1. 0.4-0.5mm long and no pigment (white)
2. 0.5-0.6mm long and light brown
3. 0.6-0.8mm long and pigmented with a hard seed coat
4. 0.6-0.8mm long and highly pigmented (brown)
5. Desiccated (which is called mature seed)

30

The nature of the relationship between DAF and the various stages of seed development are shown in Fig. 5.

15 AP1GUS transformed plants were analysed for GUS activity along with 4 control plants which were transformed with pcTAK, a construct containing the GUS gene linked to the constitutive plant promoter, cauliflower mosaic virus (CaMV) 35S - this construct was kindly provided by

35

Richard Jefferson, Plant Breeding Institute, Cambridge.

Extracts from leaf and seed stages 1, 2 and 3 of the AP1GUS and pCTAK transformed plants were assayed for GUS activity by incubation with methyl umbelliferyl glucuronide (MUG) and measurement of the fluorescence of released methyl umbelliferone (MU). Plant extracts were prepared by grinding tissue on ice in GUS extraction buffer [50 mM sodium phosphate pH 7.5, 0.1% Triton X-100, 1 mM EDTA, 10 mM dithiothreitol (DTT), 1 mM phenylmethylsulphonyl fluoride (PMSF)] in the presence of acid washed sand. Insoluble material was removed by centrifugation for 5 minutes in a bench top centrifuge. Of the soluble extract 0.05 ml was incubated with 1 ml of assay buffer [=GUS extraction buffer containing 1 mM MUG] at 37°C and 0.25 ml aliquots were removed at 0, 5, 15 and 30 minutes into 0.75 ml 0.2M Na₂CO₃. MU fluorescence was measured on a Baird Nova 1 spectrofluorimeter set at excitation wavelength 365nm and emission wavelength 455 nm.

Consistent patterns of GUS activity were obtained in individual transformed plants in terms of tissue specificity and developmental regulation, although the absolute levels of expression varied considerably from plant to plant (Fig. 6). The variable levels of gene expression obtained within the transgenic population is a common phenomenon and is thought to be due to position effects resulting from random integration of the transferred DNA (Jones et al.; EMBO J. 4 (1985) 2411-2418). There was no correlation between level of GUS expression and copy number of inserted genes.

In pAP1GUS transformed plants, the level of GUS activity (pmol MU/minute/mg fresh wt.) in leaf tissue was very low (average value = 7.2), compared to seed tissue where it was found to increase through development from an average of 68.7 in stage 1 to 301 in stage 3. In con-

trast, in plants transformed with pcTAK, the level of GUS expression was higher in leaf (average value = 237) than in seed tissue where the average value dropped from 188 at stage 1 to 56 at stage 3 (see Fig. 6).

5

The relative level of GUS activity in seed tissue compared to leaf was calculated for each of the seed stages of the AP1GUS and cTAK transformed plants. This provides data on the tissue specificity/developmental regulation of expression that is independent of plant to plant differences in absolute levels of GUS activity. The average values obtained from 13 AP1GUS plants and 4 cTAK plants are presented in the form of a histogram (Fig. 7). In AP1GUS transformed plants the level of GUS activity increases during seed development to reach a maximum value that is, on average, 100-fold higher than that observed in leaf tissue. In contrast, in cTAK plants, the level of GUS activity in all 3 seeds stages of seed development is similar to that found in leaf tissue and thus the average seed/leaf values approximate to 1. This data therefore demonstrates that the isolated AP1 promoter functions to control gene expression in a seed specific, developmentally regulated fashion.

25 Furthermore, comparison of GUS expression in stage 3 seeds of AP1GUS and cTAK transformed plants shows a higher average level of activity in the AP1GUS plants. Since the CaMV 35S promoter present in the cTAK construct is widely acknowledged as a powerful constitutive plant promoter, it follows that the AP1 promoter that has been isolated is itself a powerful promoter for
30 expressing genes in this particular tissue.

35 Example 2 - Deletion analysis of ACP promoter fragment

The 1.4 kb ACP promoter fragment described in Example 1 was subjected to deletion analysis to define more

precisely the location of sequences which are able to confer tissue specific, developmental regulation of gene expression.

5 Chimaeric gene constructs containing 1.4 kb (pAP1GUS), 0.93 kb (pAP3GUS) and 0.29 kb (pAP2GUS) respectively of the 5' upstream region of the acyl carrier protein gene ACP05 fused to the β -glucuronidase (GUS) reporter gene were transferred into tobacco and the resultant trans-
10 formed plants assayed for expression of GUS activity.

a) Construction of pAP1GUS

See Example 1.

15

b) Construction of pAP2GUS (see Fig. 8)

pTZ5BS DNA (see Example 1, item c.1) was digested to completion with *Bgl*III, producing restriction fragments
20 of 3.5, 1.1 and 0.3 kb in length. The 0.3 kb fragment was ligated to *Bam*HI linearised pTAK and the mix used to transform *E. coli* RRI.

Recombinant clones were screened by digestion with *Pst*I + *Sst*I and *Hind*III to confirm insertion of a single
25 promoter fragment in the correct orientation. The resultant plasmid AP2GUS contains 0.29 kb of the ACP promoter linked to the GUS gene.

c) Construction of AP3GUS (see Fig. 9)

30 pTZ5BS DNA was digested to completion with *Pst*I, self-ligated and the mix used to transform *E. coli* RRI. Recombinant clones were screened for recircularised large *Pst*I fragment by digestion with *Hind*III yielding a plasmid indicated with pTZ5PS. DNA of this latter plasmid
35 was digested with *Hind*III, the 1 kb promoter fragment recovered, ligated into *Hind*III digested AP2GUS DNA and transformed into *E. coli* RRI. Recombinant clones were screened by digestion with *Pst*I + *Bgl*III for the presence

of the 1 kb promoter fragment instead of the 0.29 kb AP2 promoter fragment. The resultant plasmid (pAP3GUS) contains the GUS gene linked to and under control of the the first 924 bp of the ACP promoter immediately up-
5 stream of the transcriptional start site of the structural ACP gene.

d) Transformation of *Agrobacterium*

The vectors pAP1GUS, pAP2GUS, pAP3GUS were transferred
10 into *A. tumefaciens* ACH5/pLBA4404, along with control vectors pCTAK and pTAK (a promoterless GUS gene construct) using a direct DNA uptake protocol (An *et al.*; Binary vectors; In: Plant Molecular Biology Manual (edited by Galvin and Schilperoort) A3 (1988) 1-19).
15 Transformed colonies were selected, plasmid DNA isolated, transformed back into *E. coli* and resultant plasmids subjected to restriction analysis to confirm the presence of intact copies of the respective genes.

20 e) Tobacco transformation

As described in Example 1.

f) Southern analysis of transformed plants

Regenerated plants were confirmed to contain inserted
25 copies of intact chimaeric genes via Southern analysis as described in Example 1.

g) Analysis of transgenic plants for GUS activity

Extracts from leaf and seed stages 3 and 5 of transgenic
30 plants were assayed for GUS activity (10 plants per construct group). In this experiment GUS activity was expressed as a function of DNA, which represents a more constant cell parameter than either fresh weight or protein concentration, both of which increase dramati-
35 cally during the cell expansion phase of embryo development. During initial DNA estimation of leaf extracts it was found that Triton X-100, present in the GUS extraction buffer (see Example 1), interfered with measurement

of DNA concentration using the Hoechst method. SDS was able to replace Triton X-100 without interference and was used in subsequent extractions.

5 As described in Example 1, the absolute level of GUS expression (pMol MU/minute/ μ g DNA) in individual plants was found to vary considerably. However, consistent tissue patterns of GUS expression were obtained in individual plants from each construct group. The mode
10 of GUS expression in plants transformed with the 2 deleted ACP promoter constructs (AP2GUS and AP3GUS) showed no difference to that obtained in plants transformed with the 1.4 kb promoter (AP1GUS). Thus plants transformed with each of the 3 ACP promoter constructs
15 had higher levels of expression in seed than leaf, with maximum activity in stage 3 seeds. By contrast, plants transformed with pCTAK showed similar levels of expression in seed and leaf tissues. Fig. 10 shows the values obtained from averaging the GUS activity of the
20 10 plants in each construct group. In the 3 groups transformed with the ACP promoter constructs, GUS activity is highest in stage 3 seed > mature seed > leaf (the level of activity in leaf was no higher than the background level obtained in plants transformed with the promoterless pTAK construct), thus demonstrating that
25 both deleted versions of the ACP promoter still contain the elements responsible for tissue specific and developmental regulation. In regard to levels of GUS activity, the 0.29 kb ACP promoter construct (AP2GUS),
30 in fact showed slightly higher average levels of activity than the 1.4 kb promoter, which in turn was higher than the 924 bp AP3 promoter values.

In conclusion, it has been shown that the elements
35 conferring both tissue specific, developmental regulation and also level of expression of gene ACP05 reside within 0.29 kb of the transcription start site of the gene.

Example 3 - Comparison of ACP, napin and cruciferin
promoters from rape

5

In oil seed rape, synthesis of the various storage products during seed development is differentially regulated. Thus storage lipid is synthesized first, followed by the two storage proteins, napin and cruciferin. It might be that this regulation is mediated by specific groups of seed promoters differentially activating the genes responsible for the synthesis of the various storage products during seed development.

15 The purpose of the study described in this Example was to determine if the respective rape promoter elements from a lipid biosynthetic gene (ACP) and from 2 storage protein genes, (napin and cruciferin) are indeed capable of conferring differential gene expression during seed development. This was investigated by transferring chimaeric ACP-GUS, napin-GUS and cruciferin-GUS genes into tobacco and monitoring their expression at various stages during seed development.

25 Construction of chimaeric genes

The construction of pAP1GUS, the transcriptional fusion between the 1.4 kb 5' upstream region of ACP05 and the GUS gene was described in Example 1. Constructs containing rape storage protein promoters fused to the GUS gene, cloned into the plant transformation vector pTAK, were obtained from Dr. A. Ryan at Durham University. The latter are respective transcriptional fusions (Fig. 11) between a 1.1 kb napin 5' upstream fragment and the GUS gene (pNAPGUS) and a 2.0 kb cruciferin 5' upstream fragment and the GUS gene (pCRUGUS).

Vectors were transferred into *A. tumefaciens* ACH5-
/pLBA4404 as described in Example 2 and these were used

to transform tobacco, along with pTAK and pcTAK controls, as described in Example 1. Ten plants were regenerated for each construct.

5 Southern analysis

Regenerated plants were confirmed to contain inserted copies of intact chimaeric genes via Southern analysis as described in Example 1.

10 Analysis of transgenic plants for GUS activity

Experiment 1

GUS activity was assayed in leaf and seed stages 3 and 5 of the transgenic plants as described in Example 2.

15 Consistent patterns of GUS activity in the tissues examined were obtained for individual plants in each construct group, although absolute values varied (as noted previously in Examples 1 and 2).

20 In plants transformed with either napin or cruciferin constructs the level of GUS activity in leaves was low (Fig. 12) and maximum activity was observed in stage 5 seed. With napin transformants, GUS activity in stage 3 seeds was, on average, 75% of maximum, whilst with
25 cruciferin transformants only 30% of maximum activity was observed in stage 3 seeds.
With AP1GUS transformed plants, maximum GUS activity was observed in stage 3 seeds, and this was, on average, 2.5 fold higher than in stage 5 seed.

30

These results demonstrate an earlier activation of the ACP promoter during seed development compared to the 2 storage protein gene promoters. To be able to determine the nature of this differential activation more accurately,
35 a second experiment (see below) was carried out employing a detailed seed staging GUS analysis.

Comparison of the average values of GUS activities ob-

tained from each of the construct groups allows a quantitative evaluation of the relative promoter strengths. In AP1GUS transformed plants the maximum level of GUS activity, in stage 3 seed was similar to that obtained in stage 5 seed of pCRUGUS transformed plants (also similar to the maximum level obtained in pCTAK transformed plants in leaf tissue). This value was approximately 3 fold higher than the maximum observed in pNAPGUS transformed plants, in stage 5 seed.

10

Experiment 2

Seed from the highest GUS expressing transgenic AP1GUS and CRUGUS plants was germinated on kanamycin and the resultant FI plants were used to carry out a detailed analysis of GUS expression during seed development.

15

In earlier studies (see Example 1), where plants were grown in 5 inch pots and all of the flowers pollinated, a variable rate of pod development was observed and hence morphological characteristics were used to stage seed development. In order to be able to determine more accurately the nature of the observed differential gene activation conferred by the ACP and storage protein gene promoters, an alternative system of flower tagging was utilised, which did enable DAF to be used as a meaningful marker for seed development. Thus, at the start of flowering, tobacco plants were transferred to 7.5 inch diameter compost pots. Each day at a pre-set time, the number of new flowers with anthers open was scored, a single flower was tagged and the remaining flowers removed. Tagging was carried out for 14 days (day 0 to day 13) and on day 20, all 14 pods were harvested, representing 7-20 DAF. Seeds were collected and assayed for protein and lipid content. Protein estimation was carried out by homogenising seeds in 0.1% SDS, 1 M NaCl, 50 mM sodium phosphate pH 7.5, 1 mM EDTA, 10 mM dithiothreitol, centrifuging and assaying the supernatant using a protein reagent (ex Bio-Rad). Total fatty acid content

20

25

30

35

was determined by extraction of seeds in chloroform-
/methanol (2:1) and GLC analysis of fatty acid methyl
esters produced by refluxing the extracted lipids with
methanol:toluene:conc. sulphuric acid (20:10:1). The
5 data obtained on the synthesis of lipid and protein
during seed development is shown in Fig. 13. Fatty acid
content is seen to increase sharply between days 9 and
13, whilst the major phase of protein synthesis occurs
during days 17 to 20. The sigmoidal patterns of accumu-
10 lation observed for the 2 storage products shows that,
under these established growth conditions, flower tag-
ging can be used as a meaningful developmental marker.

'Tagged' seeds were extracted and analysed for GUS acti-
15 vity as described in Example 1. Fig. 14 shows that in
AP1GUS transformed plants, GUS activity (average of 2
plants) commenced at 9 DAF, reached 50% of maximum at
10 DAF and peaked at 11-12 DAF, corresponding to the
most active phase of lipid synthesis. By 14 DAF, activi-
20 ty had fallen to 20% of maximum and it remained at this
level until 20 DAF (stage 5).

In CRUGUS transformed plants GUS activity was 1.3% of
maximum at 10 DAF and only 7% of maximum at 11 DAF, the
25 phase of most active lipid synthesis. By 15 DAF 50% of
maximum activity was reached and activity peaked between
16 and 19 DAF, corresponding to the most active phase of
protein synthesis.

Superimposition of GUS activities on fatty acid and
30 protein accumulation during tobacco seed development
(Fig. 15) shows GUS expression driven by the ACP promo-
ter to be maximum coincident with the most active phase
of lipid biosynthesis, whilst cruciferin promoter
driven GUS expression to peak several days later in
35 concert with the major phase of storage protein syn-
thesis.

These findings are of crucial importance to any pro-

gramme aimed at modifying storage lipid composition by genetic engineering. In order to perturb the process of lipid biosynthesis, the transferred gene must be under the control of a seed lipid biosynthetic promoter.

- 5 Linkage of the transferred gene to a storage protein gene promoter would lead to expression of the gene after the bulk of the storage lipid had been synthesized within the seed.

10

Example 4 - AP1-controlled MCH production in rape

This example demonstrates the functionality of the AP1 promoter in the homologous plant i.e. oil seed rape.

- 15 The AP1 promoter is shown to temporally regulate the expression of a foreign gene during seed development in oil seed rape. The foreign gene used encodes the medium chain s-acyl fatty acid synthetase thioester hydrolase (MCH) from rat. MCH is an enzyme which is induced in
20 the rat mammary gland during lactation, whereupon it causes premature chain termination of fatty acid synthesis resulting in the synthesis of medium chain fatty acids for milk production (Libertini and Smith, J. Biol. Chem. 253, (1978) 1393-1401).

25

- cdNA encoding rat MCH has been isolated and sequenced (R. Safford *et al.*, Biochem. 26 (1987) 1358-1364). In order to target the MCH gene product to the correct intracellular site of fatty acid synthesis within the
30 cotyledon cells, namely the plastid organelle, it was necessary to link the MCH cdNA to a plastid targeting sequence. This sequence was isolated from the rape ACP cdNA (R. Safford *et al.* (1988) *supra*). The resultant ACPMCH chimaeric gene was fused to the rape AP1 promoter
35 and the final construct, called pAP1A2M, was transferred, via *Agrobacterium* infection, into oil seed rape.

- a) Construction of pAP1A2M

Historically, pAP1A2M was constructed via a very circuitous route, being the final product of a number of exploratory constructs. Since all the component parts of the construct have either been described already in this patent or have been published in the literature, we will, for simplicity, only provide a description of the final construct plus the DNA sequence (see Fig. 16 and the legends to Fig. 16).

10 The construct comprises the following elements

i) ACP promoter

The 1.4 kb *Bam*H1-*Bgl*III sequence from clone ACP05 (see example 1: Construction of pAP1GUS) which comprises a sequenced 975 bp *Pst*I-*Bgl*III fragment (polynucleotide 7-981 of Fig. 3) of the non-coding part of the gene) plus a further 0.4 kb *Bam*H1-*Pst*I 5' fragment (not sequenced).

ii) ACP transit sequence

20 A 183 bp *Sau*3A-*Hae*III fragment of ACP genomic clone 05E01 (R. Safford *et al.* (1988) *supra*). A *Bgl*III linker was attached to the 3' end of the fragment to permit fusion with the MCH structural gene.

25 iii) MCH structural gene

A fragment representing nucleotides 320-1179 of MCH cDNA clone 43H09 (from the GAG codon immediately downstream of the ATG initiation codon to 71 nucleotides downstream of the translation stop codon), whereby the last two nucleotides AG were enlarged to a *Bgl*III linker to facilitate cloning into the plant vector.

35 In order to preserve the natural cleavage site (C↓A) of the ACP molecule, the junction between the ACP transit sequence and the MCH structural gene was modified by site directed mutagenesis. This involved deletion of a 9 base sequence corresponding to the *Bgl*III linker and the ATG initiation codon of MCH. Thus the final pAP1A2M

construct encoded a fusion protein consisting of the ACP transit sequence plus the first two amino acids of the mature ACP protein (Ala-Ala) followed by the MCH protein lacking the initiating ATG.

5

b) Transformation of *Agrobacterium tumefaciens*

A binary vector was constructed by transferring pAP1A2M into *A. tumefaciens* pGV3850 (Zambryski *et al.*, EMBO 2 (1983) 2143-2150) using a direct DNA uptake procedure (An *et al.* (1988) *supra*). From the resultant *Agrobacterium* colonies, DNA was extracted and transformed into *E. coli*, from which it was re-isolated enabling correct gene insertion to be verified.

15 c) Transformation of *Brassica napus* cv. Westar

Stem segments were cut and transformed with *A. tumefaciens* containing the binary vector pGV3850:pAP1A2M. The procedure used was that of Fry *et al.*, Plant Cell Reports 6 (1987) 321-325 with the following modifications:

- 20 i. Kanamycin selection was at 20 μ g/ml and was delayed until 2 weeks after infection,
- ii. Carbenicillin was replaced by cefotaxime: 500 μ g/ml,
- 25 iii. Arginine was omitted from the regeneration media,
- iv. 0.8% agar was replaced by 1% agarose,
- v. a 2-3 day pretreatment of the stem segments on standard shooting media was carried out prior to infection, and
- 30 vi. a *N. plumbaginifolia* cell line (Barfield *et al.* (1985) *supra*) was used as feeder layer.

Shoots staying green on selective media after 2 transfers were tested for the presence of nopaline (Otten and Schilperoort, B.B.A. 527 (1978) 497-500). Positive shoots were transferred to soil, potted on into 5" pots and transferred to growth rooms operating a 16 hour day (22 $^{\circ}$ C) and 8 hour night (18 $^{\circ}$ C) cycle.

d) Analysis of transformed plant tissue

DNA was extracted from leaf tissue, restricted and digests subject to Southern blot analysis, using a *KpnI*-
5 *BglII* MCH fragment, to confirm presence of inserted MCH genes.

Seeds were harvested from Southern positive plants at 5 specified developmental stages and analysed for expres-
10 sion of MCH protein using Western blotting. The 5 stages and their relationship to DAF is as follows:

Stage 1	< 15 DAF	Stage 4	25-30 DAF
Stage 2	15-20 DAF	Stage 5	30-35 DAF
Stage 3	20-25 DAF		

15 Seeds were homogenised in Laemmli sample buffer (Laemmli, Nature 227 (1970) 680-685) (1:1 v/v) using sand as an abrasive. Extracts were boiled for 5 min, microfuged and supernatants removed for analysis. 10mg equivalent fresh weight of extracts were electrophoresed on 10%
20 SDS-PAGE, blotted onto nitrocellulose and blots reacted with rat- α -MCH antibodies as described in R. Safford *et al.* (1987) *supra*.

In the resultant autoradiograph (Fig. 17) the seed ex-
25 tracts show a single cross-reactive band which co-migrates with purified MCH protein. This indicates that the ACP transit sequence of the chimaeric ACP-MCH protein has been processed, presumably upon import of MCH into plastids. The autoradiograph shows MCH expression to be
30 regulated in a temporal fashion during rape seed development. MCH expression is barely detectable in stages 1 and 2, but a dramatic increase is observed during stage 3, just prior to the onset of storage lipid deposition in oil seed rape (see Fig. 18). This result therefore
35 demonstrates that the AP1 promoter functions to express genes specifically in concert with the storage lipid synthetic phase of seed development in oil seed rape.

A culture of *E. coli* JM101/pAP1GUS was deposited under the Budapest Treaty on 22 March 1991 at the National Collection of Industrial and Marine Bacteria (Aberdeen) and obtained deposit number NCIMB 40396.

5

In agreement with Rule 28 (4) EPC the availability of a sample to a third person shall be effected only by the issue of a sample to an nominated expert.

Legends to the Figures

Figure 1

Restriction map of ACP05 genomic clone, obtained by
5 digesting ACP05 DNA with restriction enzymes BamHI,
*Pst*I, *Bgl*III, *Hind*III, *Sal*I and *Sst*I and mapping of the
restriction sites.

Figure 2

10 Dot matrix analysis of homology between ACP05 (x axis)
and ACP cDNA 29C08 (y axis) using DNA Star Dotplot
software. Blocks of 10 nucleotide sequences are compared
and regions sharing 100% homology identified by a dot.

15 Figure 3 (1/3 - 3/3)

This shows the determined nucleotide sequence (2233 bp)
of the ACP05 genomic clone provided with the deduced
amino acid sequence above the corresponding DNA sequence
and the restriction enzyme sites indicated in Fig. 1
20 given in bold type letters and indicated below the DNA
sequence.

Nucleotides 1-6 (*Sal*I site) are the remainder of the M13
cloning vector described in Example 1.b, whereas
polynucleotide 7-2233 originates from the ACP05 clone.

25 Nucleotides 7-12 form the *Pst*I restriction site CTGCAG.
The AP2 promoter (0.29 kb) is the polynucleotide 640-
930, thus starting with the *Bgl*III restriction site
AGATCT and ending just before the start site of
transcription, the first A in GGGCATCACG.

30 The introns are polynucleotides 1048-1317, 1426-1501 and
1625-1726.

Polynucleotides 1000-1047 and 1318-1422 encode the
transit peptide -51 to -1 (Met Ser Thr ... Val Ser Cys).

Nucleotides 1423-1425, 1502-1624 and 1727-1849 encode
35 the mature ACP 1 to 83 (Ala Ala Lys ... Ala Lys Lys).

Nucleotides 1850-1852 form the stop codon TGA.

Figure 4

Construction of plant transformation vector pAP1GUS.

Figure 5

- 5 Tobacco seed development. Relationship between days after flowering (DAF) = days post anthesis (dpa) and morphological stage of tobacco seeds.

Figure 6

- 10 GUS (β -glucuronidase) activity in individual AP1GUS transgenic tobacco plants. GUS activities (pmol MU/minute/mg fresh weight) were measured in seed stages 1, 2 and 3 and leaf of plants transformed with pAP1GUS or pcTAK (control) vectors.

15

Figure 7

- GUS activity in AP1GUS transgenic tobacco plants. Working out the seed/leaf values for individual plants obtained from 13 AP1GUS and 4 cTAK (control) transgenic plants and then averaging the figures obtained, the relative GUS activities (seed/leaf) for stages 1, 2 and 3 were calculated.

Figure 8

- 25 Construction of plant transformation vector pAP2GUS.

Figure 9

Construction of plant transformation vector pAP3GUS.

30 Figure 10

- GUS activity in AP1GUS, AP2GUS and AP3GUS transgenic tobacco plants. Average values (from 10 plants) of GUS activity (pmol MU/minute/ μ g DNA) of leaf, stage 3 seed and mature seed from AP1GUS, AP2GUS, AP3GUS plants and control TAK and cTAK plants.

35

Figure 11

Transcriptional fusions between napin and cruciferin

promoter sequences and the GUS structural gene, yielding plasmids pNAPGUS and pCRUGUS, respectively. The triangles at both sides indicate the left and right T-DNA borders.

5

Figure 12

GUS activity in AP1GUS, NAPGUS, CRUGUS, cTAK and TAK transgenic tobacco plants. Average values (from 10 plants) of GUS activity (pmol MU/minute/ μ g DNA) of leaf, stage 3 seed and mature seed.

10

Figure 13

Accumulation of lipid and protein during development of tobacco seed. Values (as % of fresh weight) are plotted as a percentage of the maximum recorded measurement.

15

Figure 14

GUS activity (pmol MU/minute/mg fresh weight) through seed development of AP1GUS and CRUGUS transgenic tobacco plants.

20

Figure 15

GUS activity measurements through seed development of AP1GUS and CRUGUS transgenic tobacco plants superimposed upon accumulation of lipid and protein during seed development (i.e. Figures 13 and 14 combined).

25

Figure 16 (1/4 - 4/4)

This shows the nucleotide sequence of a chimaeric construct containing the 1.4 kb AP1 promoter of which only the sequenced part is shown, i.e. the about 970 bp *Pst*I-*Bgl*III fragment, an ACP transit sequence, almost the complete MCH gene, and part of the 3'-non-coding sequence of the MCH gene. The nucleotide sequence is provided with the deduced amino acid sequence above the corresponding DNA sequence and some relevant restriction enzyme sites are given in bold type letters and indicated below the DNA sequence.

30
35

Thus polynucleotide 1-987 is identical to polynucleotide 7-993 of the DNA sequence of the ACP05 gene given in Fig. 3.

Polynucleotide 988-1149 originates from various
5 chimaeric constructs and comprises the transit sequence of ACP cDNA clone 05E01 plus the first codon of the ACP structural gene (GCG encoding Ala). This ACP-originating part was connected to the MCH structural gene in which
its ATG start codon was replaced by GCA encoding Ala
10 Thus polynucleotide 1153-1938 encodes polypeptide 3-264 in Fig. 16 being identical to polypeptide 2-263 of the MCH protein. Codon 1939-1941 is the stop codon of the MCH structural gene. Polynucleotide 1942-2010 is part of the 3'-non-coding region of the MCH gene. Nucleotides
15 2011-2015 originate from the *Bgl*III site added for facilitating cloning of the DNA sequence.

Figure 17

This autoradiograph shows the MCH expression by means of
20 Western blot analysis with rape seed extracts at stages 1-5 of rape transformed with the pAP1A2M plasmid. It shows that MCH expression is barely detectable in stages 1 and 2, but a dramatic increase is observed during stage 3, just prior to the onset of storage lipid
25 deposition in oil seed rape (see Fig. 18).

The left hand site of the autoradiograph shows a
molecular weight marker, pure MCH, an extract from
plants transformed with pTAK as a control, and seed of
tobacco transformed with the pAP1A2M plasmid.

30

Figure 18

Lipid accumulation during rape seed development.
The onset of storage lipid deposition in oil seed rape
starts about 14 DAF, reaches a reasonable value between
35 19 and 27 DAF corresponding to stage 3 seed and increases dramatically after 29 DAF.

* * * * *

LITERATURE mentioned in the specification

European patent specification EP-A2-0255377 (CALGENE,
INC.), published on 3 Feb. 1988 with claimed priority
5 date of 31 July 1986,

European patent specification EP-A2-0255378 (CALGENE,
INC.), published on 3 Feb. 1988 with claimed priority
10 date of 31 July 1986,

An *et al.*; Binary vectors; In: Plant Molecular Biology
Manual (edited by Galvin and Schilperoort) A3 (1988)
15 1-19

Barfield *et al.*; Plant Cell Reports 4 (1985) 104-107

Dellaporta; Plant Mol. Biol. Reporter 1 (1983) 19-21
20

R.J. Ellis & C. Robinson; In: Advances in Botanical
Research 14 (1987/8) 1-24; published by Academic Press
Ltd. (ISBN 0-12-005914-2)

25 Fry *et al.*, Plant Cell Reports 6 (1987) 321-325

A. Hoekema *et al.*; Nature 303 (1983) 179-181

Holsters *et al.*; Mol. Gen. Genet. 163 (1978) 181-187
30

Horsch *et al.*; Science 227 (1985) 1229-1231

Jones *et al.*; EMBO J. 4 (1985) 2411-2418

35 V.C. Knauf; TIBTECH 5 (Feb 1987) 40-47; The application
of genetic engineering to oilseed crops

Laemmli, Nature 227 (1970) 680-685

- Libertini and Smith; J. Biol. Chem. 253 (1978) 13931401;
- 5 T. Maniatis, E.F. Fritsch, & J. Sambrook; Molecular Cloning; Cold Spring Harbor Laboratory Publ. (1982).
- T. Murishige & F. Skoog; Physiol. Plant 15 (1962) 473-497
- 10 Otten and Schilperoort, Biochem. Biophys. Acta 527 (1978) 497-500
- M.A. Post-Beittenmiller *et al.*; The Plant Cell 1 (1989) 889-899
- 15 R. Safford *et al.*; Biochem. 26 (1987) 1358-1364;
- R. Safford *et al.*; Eur. J. Biochem. 174 (1988) 287-295; Plastid-localised seed acyl-carrier protein of *Brassica napus* is encoded by a distinct, nuclear multigene family
- 20 J. de Silva *et al.*; Plant Molecular Biology 14 (1990) 537-548; The isolation and sequence analysis of two seed-expressed acyl carrier protein genes from *Brassica napus*
- 25 A.R. Slabas *et al.*; 7th International Symposium of the Structure and Function of Plant lipids, University of California, Davis, CA, (1986);
- 30 P.K. Stumpf *et al.*; Fatty acid biosynthesis in higher plants; In: Fatty Acid Metabolism and Its Regulation; Elsevier Press, Amsterdam, Numa S. (ed), (1984) 155-179
- 35 G. Van den Broek *et al.*; Nature 313 (1985) 358-363;
- J.D. Watson, J. Tooze & D.T. Kurtz; Chapter 13 (Genetic Engineering of Plants by Using Crown Gall Plasmids) on

pages 164-175; In: Recombinant DNA, A Short Course,
publ. Scientific American Books (1983) distr. W.H.
Freeman and Company, New York, U.S.A.

- 5 Zambryski *et al.*; EMBO 2 (1983) 2143-2150

* * * * *

CLAIMS

1. A recombinant DNA construct containing a promoter that is capable of acting as a seed-specific plant promoter, said promoter controlling expression of a heterologous gene placed under control of said promoter in concert with the fatty acid or lipid biosynthesis in a plant cell, such that the polypeptide resulting from the expression of the heterologous gene can play a role in the fatty acid or lipid biosynthesis in the plant cell, and wherein said promoter comprises at least the 291 bp polynucleotide of clone ACP05 given in the specification, i.e.

1

```

AGATCTGATT GGTAAGATAT GGGTACTGTT TGGTTTATAT GTTTTGACTA 50
TTCAGTCACT ATGGCCCCCA TAAATTTTAA TTCGGCTGGT ATGTCTCGGT 100
TAAGACCGGT TTGACATGGT TCATTTTCAGT TCAATTATGT GAATCTGGCA 150
CGTGATATGT TTACCTTCAC ACGAACATTA GTAATGATGG GCTAATTTAA 200
GACTTAACAG CCTAGAAAGG CCCATCTTAT TACGTAACGA CATCGTTTATAG 250
AGTGCACCAA GCTTATAAAT GACGACGAGC TACCTCGGGG C          291

```

2. A DNA construct according to claim 1, comprising at least the 1 kb Pst-BglII 5' upstream fragment of the rape ACP05 gene given in Fig. 1.

3. A DNA construct according to claim 1, comprising at least the 1.4 kb BamHI-BglII 5' upstream fragment of the rape ACP05 gene given in Fig. 1.

4. A DNA construct according to claim 1, comprising a seed-specific plant promoter present in plasmid pAP1GUS present in *E. coli* JM101/pAP1GUS (NCIMB 40396).

5. Use of a DNA construct containing a seed-specific promoter as described in any one of claims 1-4 for transforming plant cells, preferably for modifying the seed-specific biosynthesis of fatty acids.

6. Use according to claim 5, whereby the plant cells are subsequently grown to whole plants in which modified biosynthesis of fatty acids occurs specifically in the seeds.



7. A process of transforming plant cells, in which a DNA construct containing a seed-specific promoter as described in any one of claims 1-4 is introduced into a transformable plant cell in such a way, that after growing the resulting transformed plant cell to whole plants a structural gene forming part of said gene controlled by the introduced seed-specific and temporally regulating plant promoter is expressed in concert with the fatty acid or lipid biosynthesis in the seeds of the plants, thereby producing the protein corresponding to said structural gene, so as to allow the protein to play a role in the fatty acid or lipid biosynthesis in the plant cell.
8. A method for modifying the formation of vegetable seed oils, which comprises growing a plant cell via plantlet to a plant bearing seed and harvesting the resulting seed containing a vegetable oil with modified composition, whereby the cells of said plant cell or plant or seed comprise a DNA construct according to any one of claims 1-4.
9. A method according to claim 8, in which said DNA construct comprises a gene under control of said promoter, which gene encodes a protein which modifies the biosynthetic pathway for fatty acid production or lipid formation.
10. A method according to claim 8 or 9, in which said DNA construct comprises a promoter essentially consisting of a seed-specific acyl carrier protein (ACP) promoter, preferably one originating from *Brassica napus*, operably linked to a structural gene encoding ACP, which gene differs in any way from the wild-type gene.
11. A seed comprising a DNA construct according to any one of claims 1-4, provided that if said DNA construct contains a seed-specific plant promoter homologous to the seed, said DNA construct being present in the genome of said seed at a site other than the natural site for said promoter.
12. A seed according to claim 11, wherein said DNA construct also contains a DNA sequence of interest encoding an exogenous protein, whereby the DNA sequence of interest is under control of the seed-specific plant promoter.



13. A seed according to claim 12, wherein said seed is of the *Brassica* family.

14. A process for producing a desired protein in plant cells, preferably seed cells, which comprises expressing a structural gene encoding said protein, said plant cells containing a recombinant DNA construct according to any one of claims 1-4 comprising said structural gene, the production of said protein optionally being followed by isolating said protein from the plant cells.

DATED 16 April 1986
Signed for and on behalf of UNILEVER PLC
by Unilever Australia Limited

.....
B. F. JONES, Company Secretary.






Fig. 1

2/23




Fig. 2

001				GTCGAC	006
				<i>SalI</i>	
CTGCAGCCAG	AAGGATAAAG	AAATTTTGGA	CGCCTGAAGA	AGAGGCAGTT	056
<i>PstI</i>					
CTGAGGGAAG	GAGTAAAAGA	GTATGTCTCC	TTAACTCTAC	TATCAAGTTT	106
CAAGAAGCTG	AGCTTGGCTC	TACCTTGATA	TGTTTATTGC	TGTTGTGCAG	156
GTATGGTAAA	TCATGGAAAG	AGATAAAGAA	TGCAAACCCT	GAAGTATTCG	206
CAGAGAGGAC	TGAGGTGAGA	GAGCATGTCA	CTTTTGTGTT	ACTCATCTGA	256
ATTATCTTAT	ATGCGAATTG	TGAGTGGTAC	TAAAAAAGGT	TGTAACCTTTT	306
GGTAGGTTGA	TTTGAAGGAT	AAATGGAGGA	ACTTGGTTCG	GTAGCCGTAA	356
CAAGTTTTTG	GGAATCTCTT	GGTTTTTAAA	TTGCTATGGA	GTTTTTTTTT	406
GCCTGCGTGA	CAACATATCA	TCAGCTGTTG	AGAAGGAAGA	TGGTATTAGA	456
AAGGGTCTTT	CTTTCACATT	TTGTGTTGTG	GACAAATATT	AAAGTCAAAT	506
GTGGCACATG	GATTTTAATT	CGGCCGGTAT	GGTTTGGTTA	AGACTGGTTT	556
AACATGTATA	ATTAGTCTTT	GTTTTATTG	GCTCAGCGGT	TTGTTGGTGT	606
TGGTTAGGAA	CTTAGGCTTG	TCTCTTCTG	ATAAGATCTG	ATTGGTAAGA	656
			<i>BglII</i>		
TATGGGTACT	GTTTGGTTTA	TATGTTTTGA	CTATTCAGTC	ACTATGGCCC	706
CCATAAATTT	TAATTCGGCT	GGTATGTCTC	GGTTAAGACC	GGTTTGACAT	756
GGTTCATTTT	AGTTCAATTA	TGTGAATCTG	GCACGTGATA	TGTTTACCTT	806
CACACGAACA	TTAGTAATGA	TGGGCTAATT	TAAGACTTAA	CAGCCTAGAA	856
AGGCCCATCT	TATTACGTAA	CGACATCGTT	TAGAGTGCAC	CAAGCTTATA	906
			<i>HindIII</i>		
AATGACGACG	AGCTACCTCG	GGGCATCAGC	CTCTTTGTAC	ACTCCGCCAT	956

Fig. 3 (1/3)

5/23

	15						20					25	
Ile	Val	Lys	Lys	Gln	Leu	Ser	Leu	Lys	Asp	Asp	Gln	Asn	
ATC	GTC	AAG	AAG	CAG	CTA	TCA	CTC	AAA	GAC	GAT	CAA	AAC	1573
				30						35			
Val	Val	Ala	Glu	Thr	Lys	Phe	Ala	Asp	Leu	Gly	Ala	Asp	
GTC	GTT	GCG	GAA	ACC	AAA	TTT	GCT	GAT	CTT	GGA	GCA	GAT	1612
	40		42										
Ser	Leu	Asp	Thr	1625									
TCT	CTC	GAC	ACT		GT	AATTCACCAA	ATGAATCACT	CTCTATGTGA	1656				
ATTAAACAAC	TTGTGTAGTT	TTTTTTTTTT	TTTTTTTTTAA	TACTGATTAG	1706								
						43		45					
				1726	Val	Glu	Ile	Val	Met	Gly	Leu		
ATTGAGTGTT	TTGCATGCAG	GTT	GAG	ATA	GTG	ATG	GGT	TTA	1747				
	50				55					60			
Glu	Glu	Glu	Phe	His	Ile	Glu	Met	Ala	Glu	Glu	Lys	Ala	
GAG	GAA	GAG	TTT	CAT	ATC	GAA	ATG	GCT	GAA	GAA	AAA	GCA	1786
		65								70		75	
Gln	Lys	Ile	Thr	Thr	Val	Glu	Glu	Ala	Ala	Glu	Leu	Ile	
CAG	AAG	ATC	ACA	ACG	GTG	GAG	GAA	GCT	GCT	GAG	CTC	ATT	1825
				80		83							
Asp	Glu	Leu	Val	Gln	Ala	Lys	Lys	***					
GAT	GAG	CTC	GTG	CAA	GCC	AAG	AAG	TGACTTT	TAGTATTAAG	1866			
AGAAGAACCA	AAGGCTTTGT	TGTTTTCATA	ATCTTTCTGT	CATTTTCTTT	1916								
TATTATGATG	TCAAGTCAAG	CGACTCTTTG	CTAGTAATCT	GTATGCCATG	1966								
GATCTCTCTC	TCTATTTGTC	GACTGAAAAC	TTTTGGGTTA	CACATGAAAG	2016								
		<i>Sall</i>		<i>HindIII</i>									
CTTTTTCTTT	TTCTAAAATC	CAAATGAAA	GAGTTGTATT	AACAGATACA	2066								
TAAGTGAAAG	AGTAGTCCCT	AAGATGACAC	TAGCTTCATT	TATAACAAT	2116								
CCTATCACAT	TGTATATACA	GGTTATGATT	TATTCCAAT	CAGCGTCAA	2166								
GARTCCAGCA	TCTTTCATCT	CTGAATAGTA	GACATTCTCC	AAGTTTAGAT	2216								
CTTCCTCCTC	GATCAAA				2233								

Fig. 3 (3/3)

6/23




Fig. 4




Fig. 5

8/23

AP1GUS	LEAF	SEED STAGE 1	SEED STAGE 2	SEED STAGE 3
5	3.4	10.0	187.2	551.2
6	5.0	68.5	99.2	249.6
10	2.4	8.6	5.6	38.7
12	0.6	0.2	0.5	1.3
14	14.3	1.2	2.0	165.6
16	25.4	209.2	424.0	264.0
18	18.4	320.0	348.0	248.0
19	8.8	144.0	200.0	284.0
25	7.8	79.2	136.4	516.0
26	2.1	41.2	137.6	426.0
27	1.8	0.3	138.4	448.0
28	0.6	0.1	52.4	300.0
29	3.3	10.9	22.0	416.0
AVE	7.22	68.72	134.87	300.65
cTAK	LEAF	SEED STAGE 1	SEED STAGE 2	SEED STAGE 3
3A	14.8	15.1	32.2	5.1
3B	9.6	2.6	7.0	24.4
21	24.8	12.4	5.1	18.4
23	900.0	720.0	-	176.0
AVE	237.3	187.5		56.0

Fig. 6

SUBSTITUTE SHEET

9/23




Fig. 7

10/23




Fig. 8

11/23




Fig. 9




Fig. 10

SUBSTITUTE SHEET





Fig. 11




Fig. 12

15/23




Fig. 13

16/23




Fig. 14

17/23




Fig. 15

18/23

CTGCAGCCAG AAGGATAAAG AAATTTTGGG CGCCTGAAGA AGAGGCAGTT 50
*Pst*I
 CTGAGGGAAG GAGTAAAAGA GTATGTCTCC TTAAGTCTAC TATCAAGTTT 100
 CAAGAAGCTG AGCTTGGCTC TACCTTGATA TGTTTATTGC TGTTGTGCAG 150
 GTATGGTAAA TCATGGAAAG AGATAAAGAA TGCAAACCCT GAAGTATTCG 200
 CAGAGAGGAC TGAGGTGAGA GAGCATGTCA CTTTGTGTGTT ACTCATCTGA 250
 ATTATCTTAT ATGCGAATTG TGAGTGGTAC TAAAAAAGGT TGTAAGTCTT 300
 GGTAGGTTGA TTTGAAGGAT AAATGGAGGA ACTTGGTTCG GTAGCCGTAA 350
 CAAGTTTTTG GGAATCTCTT GGGTTTTAAA TTGCTATGGA GTTTTTTTTT 400
 GCCTCCGTGA CAACATATCA TCAGCTGTTG AGAAGGAAGA TGGTATTAGA 450
 AAGGGTCTTT CTTTCACATT TTGTGTGTG GACAAATATT AAAGTCAAAT 500
 GTGGCACATG GATTTTAATT CGGCCGGTAT GGTTTGTTA AGACTGGTTT 550
 AACATGTATA ATTAGTCTTT GTTTTATTTG GCTCAGCGGT TTGTGGTGT 600
 TGGTTAGGAA CTTAGGCTTG TCTCTTCTG ATAAGATCTG ATTGGTAAGA 650
 TATGGGTACT GTTTGGTTTA TATGTTTTGA CTATTCAGTC ACTATGGCCC 700
*Bgl*III
 CCATAAATTT TAATTCGGCT GGTATGTCTC GGTTAAGACC GGTTTGACAT 750
 GGTTCAATTC AGTTCAATTA TGTGAATCTG GCACGTGATA TGTTTACCTT 800
 CACACGAACA TTAGTAATGA TGGGCTAATT TAAGACTTAA CAGCCTAGAA 850
 AGGCCCATCT TATTACGTAA CGACATCGTT TAGAGTGCAC CAAGCTTATA 900
*Hind*III
 AATGACGACG AGCTACCTCG GGCATCAGC CTCTTTGTAC ACTCCGCCAT 950

Fig. 16 (1/4)

19/23

CTCTCTCTCC TTCGAGCACA GATCTCTCTC GTGAATAACG AAA												-51 Met	996			
<i>BglIII</i>																
-50	-45						-40									
Ala	Thr	Thr	Phe	Ser	Ala	Ser	Val	Ser	Met	Gln	Ala	Thr				
GCG	ACC	ACT	TTC	AGC	GCT	TCA	GTC	TCC	ATG	CAA	GCT	ACC	1035			
	-35			-30				-25								
Ser	Leu	Val	Thr	Thr	Thr	Arg	Ile	Ser	Phe	Gln	Lys	Pro				
TCT	CTG	GTC	ACA	ACA	ACG	AGG	ATT	AGT	TTC	CAA	AAG	CCA	1074			
	-20				-15											
Val	Leu	Val	Ser	Asn	His	Gly	Arg	Thr	Asn	Leu	Ser	Phe				
GTT	TTG	GTT	TCC	AAC	CAT	GGA	AGG	ACT	AAT	CTC	TCC	TTC	1113			
	-10			-5				-1					1			
Asn	Leu	Ser	Arg	Thr	Arg	Leu	Ser	Ile	Ser	Cys	Ala	Ala				
AAC	CTA	AGC	CGC	ACT	CGC	CTT	TCA	ATC	TCT	TGC	GCG	GCA	1152			
	5			10						15						
Glu	Thr	Ala	Val	Asn	Ala	Lys	Ser	Pro	Arg	Asn	Glu	Lys				
GAG	ACA	GCA	GTC	AAT	GCT	AAG	AGT	CCC	AGG	AAT	GAA	AAG	1191			
	20			25												
Val	Leu	Asn	Cys	Leu	Tyr	Gln	Asn	Pro	Asp	A	Val	Phe				
GTT	TTG	AAC	TGT	TTG	TAT	CAA	AAT	CCT	GAT	GCA	GTT	TTC	1230			
	30			35						40						
Lys	Leu	Ile	Cys	Phe	Pro	Trp	Ala	Gly	Gly	Gly	Ser	Ile				
AAG	CTG	ATC	TGC	TTC	CCT	TGG	GCA	GGA	GGC	GGC	TCC	ATC	1269			
	45				50											
His	Phe	Ala	Lys	Trp	Gly	Gln	Lys	Ile	Asn	Asp	Ser	Leu				
CAT	TTT	GCC	AAG	TGG	GGC	CAA	AAG	ATT	AAC	GAC	TCT	CTG	1308			
	55			60						65						
Glu	Val	His	Ala	Val	Arg	Leu	Ala	Gly	Arg	Glu	Thr	Arg				
GAA	GTG	CAT	GCT	GTA	AGA	CTG	GCT	GGA	AGA	GAA	ACC	CGA	1347			

Fig. 16 (2/4)

20/23

Leu	Gly	Glu	Pro	Pne	Ala	Asn	Asp	Ile	Tyr	Gln	Ile	Ala	
CTT	GGA	GAA	CCT	TTC	GCA	AAT	GAC	ATC	TAC	CAG	ATA	GCT	1386
				85					90				
Asp	Glu	Ile	Val	Thr	Ala	Leu	Leu	Pro	Ile	Ile	Gln	Asp	
GAT	GAA	ATC	GTG	ACC	GCC	CTG	TTG	CCC	ATC	ATT	CAG	GAT	1425
		95					100				105		
Lys	Ala	Phe	Ala	Phe	Phe	Gly	His	Ser	Phe	Gly	Ser	Tyr	
AAA	GCT	TTT	GCG	TTT	TTT	GGC	CAC	AGT	TTT	GGA	TCC	TAC	1464
			110					115					
Thr	Ala	Leu	Ile	Thr	Ala	Leu	Leu	Leu	Lys	Glu	Lys	Tyr	
ACT	GCT	CTT	ATT	ACT	GCT	CTG	CTC	CTA	AAG	GAG	AAA	TAC	1503
						120							
Lys	Met	Glu	Pro	Leu	His	Ile	Phe	Val	Ser	Gly	Ala	Ser	
AAA	ATG	GAG	CCG	CTG	CAT	ATT	TTT	GTA	TCC	GGT	GCA	TCC	1542
			135					140				145	
Ala	Pro	His	Ser	Thr	Ser	Arg	Pro	Gln	Val	Pro	Asp	Leu	
GCC	CCT	CAC	TCA	ACA	TCC	CGG	CCT	CAA	GTT	CCT	GAT	CTT	1581
				150					155				
Asn	Glu	Leu	Thr	Glu	Glu	Gln	Val	Arg	His	His	Leu	Leu	
AAC	GAA	TTG	ACA	GAA	GAA	CAA	GTC	AGA	CAT	CAC	CTT	CTG	1620
						160			165			170	
Asp	Phe	Gly	Gly	Thr	Pro	Lys	His	Leu	Ile	Glu	Asp	Gln	
GAT	TTC	GGA	GGC	ACG	CCC	AAG	CAT	CTC	ATA	GAA	GAC	CAG	1659
				175					180				
Asp	Val	Leu	Arg	Met	Phe	Ile	Pro	Leu	Leu	Lys	Ala	Asp	
GAT	GTT	CTG	AGG	ATG	TTC	ATT	CCT	TTG	CTG	AAG	GCA	GAT	1698
						185							
Ala	Gly	Val	Val	Lys	Lys	Phe	Ile	Phe	Asp	Lys	Pro	Ser	
GCT	GGC	GTT	GTG	AAA	AAA	TTC	ATC	TTT	GAC	AAG	CCC	TCC	1737

Fig. 16 (3/4)

SUBSTITUTE SHEET

21/23

Lys	Ala	Leu	Leu	Ser	Leu	Asp	Ile	Thr	Gly	Phe	Leu	Gly	
AAA	GCT	CTT	CTC	TCT	CTG	GAC	ATA	ACG	GGC	TTC	CTT	GGA	1776
Ser	Glu	Asp	Thr	Ile	Lys	Asp	Ile	Glu	Gly	Trp	Gln	Asp	
TCT	GAA	GAT	ACA	ATA	AAG	GAC	ATA	GAA	GGC	TGG	CAA	GAC	1815
Leu	Thr	Ser	Gly	Lys	Phe	Asp	Val	His	Met	Leu	Pro	Gly	
CTA	ACC	AGT	GGG	AAG	TTT	GAT	GTC	CAC	ATG	CTG	CCA	GGC	1854
Asp	His	Phe	Tyr	Leu	Met	Lys	Pro	Asp	Asn	Glu	Asn	Phe	
GAC	CAC	TTT	TAT	CTG	ATG	AAG	CCC	GAC	AAC	GAG	AAC	TTT	1893
Ile	Lys	Asn	Tyr	Ile	Ala	Lys	Cys	Leu	Glu	Leu	Ser	Ser	
ATC	AAG	AAC	TAC	ATA	GCC	AAG	TGC	TTG	GAA	CTC	TCG	TCA	1932
Leu	Thr	***											
CTC	ACT	TGA	CTACTTTTA	GATGAGCTTT	CTTTGGGGCT								1970
GTGGATATGC	AGACGGTTCA	AAAGCTGCTC	CTCTGGGTCC	AGATC									2015
				BgIII									

Fig. 16 (4/4)

22/23




Fig. 17

23/23





Fig. 18

INTERNATIONAL SEARCH REPORT

PCT/GB 92/00627

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/82; A01H5/10; C11B1/00; C12P21/02 //C12N15/55//C12N9/16//C12N15/29		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N ; C11B ; A01H ; C12P	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ^o	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	PLANT MOLECULAR BIOLOGY. vol. 14, 1990, DORDRECHT, THE NETHERLANDS. pages 537 - 548; DE SILVA, J., ET AL.: 'The isolation and sequence analysis of two seed-expressed acyl carrier protein genes from Brassica napus' cited in the application see the whole document ---	1-4
O,X	J. EXP. BOT. vol. 41, 1990, SUPPLEMENT, P5-1 DE SILVA, J.: 'Isolation of an acyl carrier protein gene from oil seed rape and functional analysis of the 5' upstream sequence' * abstract P5.02 *	1-6,8, 13-16
O,Y	---	7-12
	-/--	
<p>^o Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
13 JULY 1992	24. 07. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	MADDOX A. D. 	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with Indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	TIBTECH vol. 5, no. 2, February 1987, pages 40 - 47; KNAUF, V. C.: 'The application of genetic engineering to oilseed crops' see page 44, column 2, line 28 - page 45, column 2, line 46 ---	7-12
X	BIOTECHNOLOGY. vol. 7, no. 9, September 1989, NEW YORK, USA. pages 929 - 932; VANDERKERCKHOVE, J., ET AL: 'Enkephalins produced in transgenic plants using modified 2S seed storage proteins' * page 930 left col. last para * ---	16
P,X	WO,A,9 113 980 (CALGENE) 19 September 1991 see example 4 ---	1,6
P,X	WO,A,9 113 972 (CALGENE) 19 September 1991 see page 13, line 29 - page 14, line 10 see page 20, line 14 - page 21 see page 41 - page 42 see page 51 - page 60 see page 73, line 18 - page 75, line 3 see page 77, line 30 - page 78, line 15 ---	1,6-10, 13-16
P,X	WO,A,9 116 421 (CALGENE) 31 October 1991 see page 85 - page 86 ---	1
P,X	WO,A,9 203 564 (CALGENE) 5 March 1992 see page 84 - page 87 ---	1,6-12
A	BIOTECHNOLOGY. vol. 6, no. 10, October 1988, NEW YORK, USA. pages 1219 - 1221; BAYLEY, S. A., ET AL.: 'Metabolic consequences of expression of the medium chain hydrolase gene of the rat in mouse NIH 3T3 cells' see the whole document ---	7-12
A	EP,A,0 255 377 (CALGENE) 3 February 1988 see page 12, line 9 - page 26, line 27 ---	8,9
A	EP,A,0 255 378 (CALGENE) 3 February 1988 see page 3, line 6 - line 9 see page 4, line 5 - line 65; example 3 ---	1-12
	-/--	

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. GB 9200627
SA 58153**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 13/07/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9113980	19-09-91	EP-A- 0472712	04-03-92
WO-A-9113972	19-09-91	EP-A- 0472722	04-03-92
WO-A-9116421	31-10-91	EP-A- 0480024	15-04-92
WO-A-9203564	05-03-92	None	
EP-A-0255377	03-02-88	AU-B- 609724	09-05-91
		AU-A- 7630387	04-02-88
		US-A- 5110728	05-05-92
		AU-B- 612326	11-07-91
		AU-A- 7630287	04-02-88
		EP-A- 0255378	03-02-88
		JP-A- 63119680	24-05-88
		JP-A- 63112987	18-05-88
EP-A-0255378	03-02-88	AU-B- 612326	11-07-91
		AU-A- 7630287	04-02-88
		AU-B- 609724	09-05-91
		AU-A- 7630387	04-02-88
		EP-A- 0255377	03-02-88
		JP-A- 63119680	24-05-88
		JP-A- 63112987	18-05-88
		US-A- 5110728	05-05-92
NL-A-9002130	16-04-92	None	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
O,A	J. CELL. BIOCHEM. SUPPL. vol. 14E, 1990, MEETING APRIL 16-22, 1990 page 266; KNAUF, V. C., ET AL.: 'Reprogramming levels of fatty acid synthesis enzymes in developing embryos of rapeseed' * abstract R 018 *	1-12
E	--- NL,A,9 002 130 (STICHTING TECHNISCHE WETENSCHAPPEN UTRECHT) 16 April 1992 see the whole document ---	1,8,9,16