

(19) **DANMARK**

(10) **DK/EP 2623599 T3**



(12) **Oversættelse af  
europæisk patentskrift**

Patent- og  
Varemærkestyrelsen

- (51) Int.Cl.: **C 12 N 15/11 (2006.01)**                      **A 61 K 31/7088 (2006.01)**                      **A 61 K 31/7115 (2006.01)**  
**A 61 K 31/712 (2006.01)**                      **A 61 P 3/06 (2006.01)**                      **A 61 P 31/14 (2006.01)**  
**C 12 N 15/113 (2010.01)**
- (45) Oversættelsen bekendtgjort den: **2019-04-08**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-01-02**
- (86) Europæisk ansøgning nr.: **12191739.7**
- (86) Europæisk indleveringsdag: **2008-10-03**
- (87) Den europæiske ansøgnings publiceringsdag: **2013-08-07**
- (30) Prioritet:            **2007-10-04 US 977497 P**                                              **2007-10-11 US 979217 P**  
                                 **2008-02-12 US 28062**                                              **2008-07-17 EP 08104780**
- (62) Stamansøgningsnr: **08801378.4**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **ROCHE INNOVATION CENTER COPENHAGEN A/S, Fremtidsvej 3, 2970 Hørsholm, Danmark**
- (72) Opfinder: **Obad, Susanna, Nicoloviusgatan 10 A, 217 57 Malmø, Sverige**  
**Kauppinen, Sakari, Norskekrogen 12, 2765 Smørum, Danmark**  
**Elmén, Joacim, Sickla Kanalgata 19 A, 120 67 Stockholm, Sverige**  
**Lindow, Morten, Thomas Koppels Allé 22C, 2450 København SV, Danmark**  
**Heidenblad, Markus, Mantalskroken 9, 226 47 Lund, Sverige**
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **Mikromirer**
- (56) Fremdragne publikationer:  
**WO-A2-2007/112753**  
**US-A1- 2007 065 840**  
**ELMÉN J ET AL: "LNA-mediated microRNA silencing in non-human primates", NATURE: INTERNATIONAL WEEKLY JOURNAL OF SCIENCE, NATURE PUBLISHING GROUP, UNITED KINGDOM, vol. 452, 17 April 2008 (2008-04-17), pages 896-900, XP002533567, ISSN: 0028-0836, DOI: 10.1038/NATURE06783 -& Elmén J. et al.: "LNA-mediated microRNA silencing in non-human primates", , 17 April 2008 (2008-04-17), XP002699495, Retrieved from the Internet: URL:http://www.nature.com/nature/journal/v 452/n7189/extref/nature06783-s2.pdf [retrieved on 2013-06-25]**  
**FABANI M M ET AL: "miR-122 targeting with LNA/2'-O-methyl oligonucleotide mixmers, peptide nucleic acids (PNA), and PNA-peptide conjugates", RNA, COLD SPRING HARBOR LABORATORY PRESS, US, vol. 14, 1 February 2008 (2008-02-01), pages 336-346, XP002533356, ISSN: 1355-8382, DOI: 10.1261/RNA.844108**  
**ELMÉN J ET AL: "Antagonism of microRNA-122 in mice by sistemically administered LNA-antimiR leads to up-regulation of a large set of predicted target mRNAs in the liver", NUCLEIC ACIDS RESEARCH, OXFORD**

Fortsættes ...

UNIVERSITY PRESS, SURREY, GB, vol. 36, no. 4, 23 December 2007 (2007-12-23), pages 1153-1162, XP002533568, ISSN: 0305-1048, DOI: 10.1093/NAR/GKM1113

OBAD S ET AL: "Targeting of cancer-associated microRNAs using short LNA-antimiR oligonucleotides", EUROPEAN JOURNAL OF CANCER. SUPPLEMENT, PERGAMON, OXFORD, GB, vol. 6, no. 9 (549), 1 July 2008 (2008-07-01), page 142, XP002533357, ISSN: 1359-6349, DOI: 10.1016/S1359-6349(08)71721-X

ELAYADI A ET AL: "Implications of High-Affinity Hybridization by locked Nucleic Acid Oligomers for Inhibition of human Telomerase", BIOCHEMISTRY, AMERICAN CHEMICAL SOCIETY, US, vol. 41, no. 31, 1 January 2002 (2002-01-01), pages 9973-9981, XP002990849, ISSN: 0006-2960, DOI: 10.1021/BI025907J

WIENHOLDS E ET AL: "MicroRNA expression in zebrafish embryonic development", SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, US, vol. 309, no. 5732, 8 July 2005 (2005-07-08), pages 310-311, XP002386061, ISSN: 0036-8075, DOI: 10.1126/SCIENCE.1114519

WIENHOLDS E ET AL: "MicroRNA expression in zebrafish embryonic development", SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, US, vol. 309, no. 5732, 8 July 2005 (2005-07-08), pages S1-S34, XP002388752, ISSN: 0036-8075, DOI: 10.1126/SCIENCE.1114519

CHAN J A ET AL: "MicroRNA-21 is an antiapoptotic factor in human glioblastoma cells", CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 65, no. 14, 15 July 2005 (2005-07-15), pages 6029-6033, XP002457479, ISSN: 0008-5472, DOI: 10.1158/0008-5472.CAN-05-0137

**Description****FIELD OF THE INVENTION**

5 **[0001]** The present invention relates to very short oligonucleotides which target and inhibit microRNAs *in vivo*, and their use in medicaments and pharmaceutical compositions.

**BACKGROUND OF THE INVENTION**

10 **[0002]** MicroRNAs (miRNAs) are an abundant class of short endogenous RNAs that act as post-transcriptional regulators of gene expression by base-pairing with their target mRNAs. They are processed from longer (ca 70-80 nt) hairpin-like precursors termed pre-miRNAs by the RNase III enzyme Dicer. MicroRNAs assemble in ribonucleoprotein complexes termed miRNPs and recognize their target sites by antisense complementarity thereby mediating down-regulation of their target genes. Near-perfect or perfect complementarity between the miRNA and its target site results in target mRNA cleavage, whereas limited complementarity between the microRNA and the target site results in translational inhibition of the target gene.

15 **[0003]** A summary of the role of microRNAs in human diseases, and the inhibition of microRNAs using single stranded oligonucleotides is provided by WO2007/112754 and WO2007/112753, WO2008046911, provides microRNA sequences which are associated with cancer. Numerous microRNAs have been associated with disease phenotypes and it is therefore desirable to provide substances capable of modulating the availability of microRNAs *in vivo*. WO2007/112754 and WO2007/112753 disclose short single stranded oligonucleotides which are considered to form a strong duplex with their target miRNA. SEQ ID NOs 1 - 45 are examples of anti microRNA oligonucleotides as disclosed in WO2007/112754 and WO2007/112753.

**SUMMARY OF THE INVENTION**

25 **[0004]** The present invention is based upon the discovery that the use of very short oligonucleotides which target microRNAs and which have a high proportion of nucleotide analogue nucleotides, such as LNA nucleotides, are highly effective in alleviating the repression of RNAs, such as an mRNA, by the targeted microRNAs *in vivo*.

30 **[0005]** The present invention provides for an oligomer of 7-12 nucleobases in length, for use in reducing the effective amount of a microRNA target in a cell; wherein all of the nucleobase units of the contiguous nucleobase sequence are LNA nucleobase units, wherein the contiguous nucleobase sequence is complementary to the corresponding sequence of at least two miRNA sequences, and wherein at least 75% of the internucleoside linkages present between the nucleobase units of the contiguous nucleobase sequence are phosphorothioate internucleoside linkages.

35 **[0006]** The present invention provides for an oligomer of 7-12 nucleobases in length, for use in reducing the effective amount of a microRNA target in a cell; wherein all of the nucleobase units of the contiguous nucleobase sequence are LNA nucleobase units, wherein the contiguous nucleobase sequence is complementary to the corresponding sequence of at least two miRNA sequences, and wherein at least 75% of the internucleoside linkages present between the nucleobase units of the contiguous nucleobase sequence are phosphorothioate internucleoside linkages, for use as a medicament.

40 **[0007]** The oligomer of the invention may in some embodiments be between 7-10 nucleotides in length which comprises a contiguous nucleotide sequence of a total of between 7-10 nucleotides, such as 7, 8, 9, or 10, nucleotide units, wherein the nucleotide sequence is complementary to a corresponding nucleotide sequence found in mammalian or viral microRNA.

45 **[0008]** The present invention provides oligomers according to the invention as a medicament.

**[0009]** The present invention provides pharmaceutical compositions comprising the oligomer of the invention and a pharmaceutically acceptable diluent, carrier, salt or adjuvant.

**[0010]** The invention provides for a conjugate comprising an oligomer according to the invention, conjugated to at least one non-nucleotide or polynucleotide entity, such as a sterol, such as cholesterol.

50 **[0011]** The invention provides for the use of an oligomer or a conjugate according to the invention, for the manufacture of a medicament for the treatment of a disease or medical disorder associated with the presence or over-expression of a microRNA, such as one or more of the microRNAs referred to herein.

**[0012]** The invention provides for the treatment of a disease or medical disorder associated with the presence or overexpression of the microRNA, comprising the step of administering a composition (such as the pharmaceutical composition) comprising an oligomer or conjugate according to the invention to a patient suffering from or likely to suffer from said disease or medical disorder.

55 **[0013]** The invention provides for an *in vitro* method for reducing the effective amount of a microRNA target in a cell, comprising administering the oligomer of the invention, or a composition (such as a pharmaceutical composition) com-

prising the oligomer or conjugate according to the invention to the cell.

**[0014]** The invention provides for an *in vitro* method for reducing the effective amount of a microRNA target in a cell, comprising administering the oligomer or conjugate or pharmaceutical composition according to the invention to the cell.

**[0015]** The invention provides for an *in vitro* method for de-repression of a target mRNA (or one or more RNAs) in a cell, comprising administering an oligomer or conjugate according to the invention, or a composition comprising said oligomer or conjugate, to said cell.

**[0016]** The invention provides for the use of an oligomer or a conjugate according to the invention, for inhibiting the microRNA in a cell which comprises said microRNA, such as a human cell. The use may be *in vivo* or *in vitro*.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0017]**

**Figure 1.** Schematic presentation of the miR-21, miR-155 and miR-122 8-mer LNA-antimiRs, indicating the targeting positions with the fully LNA-modified and phosphorothiolated LNA-antimiR. Preferred hybridisation positions for 7mer, 8mer, 9mer and 10mer LNA oligonucleotides on the mature microRNA are also indicated.

**Figure 2.** Assessment of miR-21 antagonism by SEQ ID #3205 and SEQ ID #3204 LNA-antimiRs in MCF-7 cells using a luciferase sensor assay. MCF-7 cells were co-transfected with luciferase sensor plasmids containing a perfect match target site for miR-21 or a mismatch target site (.mm2) and LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean of renilla/firefly ratios for three separate experiments (bars = s.e.m), were all have been normalized against 0 nM psiCHECK2 (=control).

**Figure 3.** Assessment of miR-21 antagonism by SEQ ID #3205 and SEQ ID #3204 LNA-antimiRs in HeLa cells using a luciferase sensor assay. HeLa cells were co-transfected with luciferase sensor plasmids containing a perfect match target site for miR-21 (mir-21) or a mismatch target site (mm2) and LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean of renilla/firefly ratios for three separate experiments (bars = s.e.m), were all have been normalized against 0 nM psiCHECK2 (=control).

**Figure 4.** Assessment of miR-155 antagonism by SEQ ID #3206 and SEQ ID #3207 LNA-antimiRs in LPS-treated mouse RAW cells using a luciferase sensor assay. RAW cells were co-transfected with miR-155 and the different LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean of renilla/firefly, were all have been normalized against 0 nM psiCHECK2.

**Figure 5.** Assessment of miR-122 antagonism by SEQ ID #3208 and SEQ ID #4 LNA-antimiRs in HuH-7 cells using a luciferase sensor assay. HuH-7 cells were co-transfected with a miR-122 luciferase sensor containing a perfect match miR-122 target site and the different LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean of renilla/firefly ratios for three separate experiments (bars = s.e.m), where all have been normalized against 0 nM psiCHECK2 (=control).

**Figure 6.** Schematic presentation of the miR-21 luciferase reporter constructs.

**Figure 7.** Assessment of miR-21 antagonism by an 8-mer LNA-antimiR (SEQ ID #3205) versus a 15-mer LNA-antimiR (SEQ ID #3204) in PC3 cells using a luciferase reporter assay. PC3 cells were co-transfected with luciferase reporter plasmids containing a perfect match target site for miR-21 or a mismatch target site and LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) of three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs. LNA nucleotides are indicated by ovals, and DNA residues are indicated by bars.

**Figure 8.** Specificity assessment of miR-21 antagonism by an 8-mer LNA-antimiR in HeLa cells using a luciferase reporter assay. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match or a mismatched target site for miR-21 and LNA-antimiRs (SEQ ID #3205) or an 8-mer LNA mismatch control oligo (SEQ ID #3218) at different concentrations. After 24 hours, cells were harvested and luciferase activity was measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the Renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs. Mismatches are indicated by filled ovals.

**Figure 9.** Assessment of the shortest possible length of a fully LNA-modified LNA-antimiR that mediates effective antagonism of miR-21. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match or a mismatch target site for miR-21 and the LNA-antimiRs at different concentrations (SEQ ID #3209 =6-mer and SEQ ID #3210=7-mer). After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 10.** Length assessment of fully LNA-substituted LNA-antimiRs antagonizing miR-21. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match or a mismatch target site for miR-21 and LNA-antimiRs at different concentrations (SEQ ID #3211 =9-mer, SEQ ID #3212=10-mer, SEQ ID #3213=12-mer and SEQ ID #3214=14-mer). After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 11.** Determination of the most optimal position for an 8-mer LNA-antimiR within the miR target recognition sequence. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match or a mismatch target site for miR-21 and the LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 12.** Validation of interaction of the Pcd4-3'-UTR and miR-21 by the 8-mer SEQ ID #3205 LNA-antimiR. HeLa cells were co-transfected with a luciferase reporter plasmid containing part of the 3'UTR of Pcd4 gene and LNA-antimiRs at different concentrations (SEQ ID #3205 = 8 mer, perfect match; SEQ ID #3218 = 8 mer, mismatch; SEQ ID #3204 = 15 mer, LNA/DNA mix; SEQ ID #3220 = 15 mer, gapmer). After 24 hours, cells were harvested and luciferase activity measured. Shown are renilla/firefly ratios that have been normalized against 0 nM. Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 13.** Comparison of an 8-mer LNA-antimiR (SEQ ID #3207) with a 15-mer LNA-antimiR (SEQ ID #3206) in antagonizing miR-155 in mouse RAW cells. Mouse RAW cells were co-transfected with luciferase reporter plasmids containing a perfect match for miR-155 and the different LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) of three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without miR-155 target site (=control). Shown is also a schematic presentation of the miR-155 sequence and the design and position of the LNA-antimiRs.

**Figure 14.** Assessment of c/EBP $\beta$  Assessment of c/EBP $\beta$  LNA-antimiR (SEQ ID #3207) with a 15-mer LNA-antimiR (SEQ ID #3206) in antagonizing miR-155 in mouse RAW cells. Mouse RAW cells were co-transfected with luciferase reporter plasmids containing a perfect match for miR-155 and the different LNA-antimiRs at different concentrations. After 20 hours, cells were harvested and western blot analysis of protein extracts from RAW cells was performed. The different isoforms of c/EBP $\beta$  are indicated, and the ratios calculated on c/EBP $\beta$  LIP and beta-tubulin are shown below.

**Figure 15.** Antagonism of miR-106b by a fully LNA-modified 8-mer (SEQ ID #3221) LNA-antimiR or by a 15-mer mixmer (SEQ ID #3228) antimiR. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match for miR-106b and the different LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values of four replicates where the renilla/firefly ratios have been normalized against 0 nM empty vector without miRNA target site (=control). Shown is also a schematic presentation of the miR-106b sequence and the design and position of the LNA-antimiRs.

**Figure 16.** Antagonism of miR-19b by a fully LNA-modified 8-mer (SEQ ID #3222) LNA-antimiR and a 15-mer (SEQ ID #3229) mixmer antimiR. HeLa cells were co-transfected with luciferase reporter plasmids containing a perfect match for miR-19a and the two LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values of four replicate experiments, where the renilla/firefly ratios have been normalized against 0 nM empty vector without a miR-19a target site (=control). Shown is also a schematic presentation of the miR-19a sequence and the design and position of the LNA-antimiRs.

**Figure 17.** Schematic presentation showing the mature human miR-221 and miR-222 sequences. Shown in the square is the seed sequence (7-mer) that is conserved in both miRNA sequences.

**Figure 18.** Targeting of a microRNA family using short, fully LNA-substituted LNA-antimiR. PC3 cells were co-transfected with luciferase reporter plasmids for miR-221 and miR-222 separately or together and with the different LNA-antimiRs at varying concentrations. When co-transfecting with the LNA-antimiRs (15-mers) SEQ ID #3223 (against miR-221) and SEQ ID #3224 (against miR-222), the total concentration was 2 nM (1 nM each), while transfecting the cells with SEQ ID #3225 (7-mer) the concentrations were 0, 1, 5, 10 or 25 nM. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) of three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without a miRNA target site (=control). Shown is also a schematic presentation of the miR-221/222 sequence and the design and position of the LNA-antimiRs.

**Figure 19.** Assessment of p27 protein levels as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR. PC3 cells were transfected with the 7-mer LNA-antimiR SEQ ID #3225 targeting both miR-221 and miR-222 at varying concentrations. After 24 hours, cells were harvested and protein levels were measured on a western blot. Shown are the ratios of p27/tubulin.

**Figure 20.** Assessment of miR-21 antagonism by an 8-mer LNA-antimiR (SEQ ID #3205) versus a 15-mer LNA-antimiR (SEQ ID #3204) and an 8-mer with 2 mismatches (SEQ ID #3218) in HepG2 cells using a luciferase reporter assay.

HepG2 cells were co-transfected with luciferase reporter plasmid containing a perfect match target site for miR-21 and LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) of three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 21.** Validation of interaction of the Pdc4 3'UTR and miR-21 by the 8-mer SEQ ID #3205 LNA-antimiR versus the 15-mer (SEQ ID #3204) and an 8-mer with two mismatches (SEQ ID #3218).

Huh-7 cells were co-transfected with a luciferase reporter plasmid containing part of the 3'UTR of Pdc4 gene, pre-miR-21 (10 nM) and LNA-antimiRs at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) of three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=control). Shown is also a schematic presentation of the miR-21 sequence and the design and position of the LNA-antimiRs.

**Figure 22.** Antagonism of miR-21 by SEQ ID #3205 leads to increased levels of Pdc4 protein levels.

HeLa cells were transfected with 5 nM LNA-antimiR SEQ ID #3205 (perfect match), or SEQ ID #3219 LNA scrambled (8mer) or SEQ ID #3218 (8-mer mismatch). Cells were harvested after 24 hours and subjected to Western blot with Pdc4 antibody.

**Figure 23.** ALT and AST levels in mice treated with SEQ ID #3205 (perfect match) or SEQ ID #3218 (mismatch control). Mice were sacrificed after 14 days and after receiving 25 mg/kg every other day.

**Figure 24.** Assessment of PU.1 protein levels as a functional readout for miR-155 antagonism by short LNA-antimiR (SEQ ID #3207).

THP-1 cells were co-transfected with pre- miR-155 (5 nmol) and different LNA oligonucleotides (5 nM) and 100 ng/ml LPS was added. After 24 hours, cells were harvested and western blot analysis of protein extracts from the THP-1 cells was performed. PU.1 and tubulin are indicated.

**Figure 25.** Assessment of p27 protein levels as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR.

PC3 cells were transfected with the 7-mer LNA-antimiR SEQ ID #3225 targeting both miR-221 and miR-222 and a LNA scrambled control at 5 and 25 nM. After 24 hours, cells were harvested and protein levels were measured on a western blot. Shown are the ratios of p27/tubulin.

**Figure 26.** Knock-down of miR-221/222 by the 7-mer SEQ ID #3225 (perfect match) LNA-antimiR reduces colony formation in soft agar in PC3 cells.

PC3 cells were transfected with 25 nM of the 7-mer LNA-antimiR SEQ ID #3225 targeting both miR-221 and miR-222 or a 7-mer scrambled control ((SEQ ID #3231). After 24 hours, cells were harvested and seeded on soft agar. After 12 days, colonies were counted. One experiment has been done in triplicate.

**Figure 27.** Overview of the human let-7 family, and of tested antagonists.

(upper) The sequences represent the mature miRNA for each member and the box depicts nucleotides 2-16, the positions typically antagonized by LNA-antimiRs. Columns to the right show the number of nucleotide differences compared to let-7a, within the seed (S; position 2-8), extended seed (ES; position 2-9), and the remaining sequence typically targeted by LNA-antimiRs (NE; position 9-16), respectively. Nucleotides with inverted colors are altered compared to let-7a. (lower) Summary of tested antagonists against the let-7 family, including information on design, length and perfectly complementary targets. All compounds are fully phosphorothiolated.

**Figure 28.** Assessment of let-7 antagonism by six different LNA-antimiRs in Huh-7 cells using a luciferase sensor assay.

Huh-7 cells were co-transfected with luciferase sensor plasmids containing a partial HMGA2 3'UTR (with four let-7 binding sites), with or without let-7a precursor (grey and black bars, respectively), and with 6 different LNA-antimiRs at increasing concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean of renilla/firefly ratios for duplicate measurements and standard deviations for each assay. Within each LNA-antimiR group all ratios have been normalized to the average of wells containing no let-7a precursor (black bars).

**Figure 29.** Luciferase results from Huh-7 cells transfected with the HMGA2 3'UTR sensor plasmid, LNA-antimiRs SEQ ID #3226 (left) and SEQ ID #3227 (right), and pre-miRs for let-7a (A), let-7d (B), let-7e (C), and let-7i (D). Grey bars indicate the target de-repression after pre-mis inclusion, whereas black control bars represent the equivalent level without pre-miR addition. Each ratio is based on quadruplicate measurements and have been normalized against the average of wells containing no precursor (black bars) within each treatment group.

**Figure 30.** Luciferase results from HeLa cells transfected with the HMGA2 3'UTR sensor plasmid or control vector, and the LNA-antimiR SEQ ID #3227 at various concentrations. Each ratio is based on quadruplicate measurements normalized against untreated (0 nM) empty control vector (psi-CHECK-2; grey bars).

**Figure 31.** Assessment of miR-21 antagonism by 8mer (#3205) in HCT116 cells using a luciferase sensor assay. HCT116 cells were co-transfected with luciferase sensor plasmids containing a perfect match target site for miR-21 (grey bars) and LNA-antimiR and control oligonucleotides at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown is one typical example of two where the renilla/firefly ratios have been normalized against 0 nM empty vector (=black bars).

**Figure 32.** Silencing of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation in soft agar in PC3 cells. PC3 cells were transfected with 25 nM of the 8-mer LNA-antimiR #3205 targeting miR-21. After 24 hours, cells were harvested and seeded on soft agar. After 12 days, colonies were counted. Shown is the mean of three separate experiments, each performed in triplicate, and normalised against 0 nM control (i.e. transfection but with no LNA).  $p=0.01898$  for #3205.

**Figure 33.** Knock-down of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation in soft agar in HepG2 cells. HepG2 cells were transfected with 25 nM of the 8-mer LNA-antimiR #3205 targeting miR-21. After 24 hours, cells were harvested and seeded on soft agar. After 17 days, colonies were counted. Shown is the mean of three replicates from one experiment (bars=SEM).

**Figure 34.** Wound closure in the invasive human prostate cell line PC3 after treatment with #3205. (A) PC3 cells were transfected at day 3 with LNA-antimiR and control oligonucleotides at 25 nM, #3205 (8mer, perfect match) and #3219 (8mer, mismatch) and the following day a scratch was made. Pictures were taken after 24 hours in order to monitor the migration. (B) The area in each timepoint has been measured with the software program Image J and normalized against respective 0 h time-point.

**Figure 35.** Length assessment of fully LNA-substituted LNA-antimiRs antagonizing miR-155.

RAW cells were co-transfected with luciferase reporter plasmids containing a perfect match target site for miR-155 and with LNA-antimiR oligonucleotides at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=mock). Shown is also a schematic presentation of the miR sequence and the design and position of the LNA-antimiRs.

**Figure 36.** Binding of 5'-FAM labeled LNA-antimiR-21 (#3205) to mouse plasma protein.

(A) % unbound LNA-antimiR-21 compound as a function of oligonucleotide concentration in mouse plasma. (B) Concentration of unbound LNA-antimiR-21 compound #3205 as a function of #3205 concentration in mouse plasma.

**Figure 37.** Quantification Ras protein levels by Western blot analysis.

A. Gel image showing Ras and Tubulin (internal standard) protein in treated (anti-let-7; 8-mer) vs. untreated (saline) lung and kidney samples. B. Quantifications of Ras protein levels in the lung and kidney, respectively, of LNA-antimiR-treated mice (black bars), normalized against equivalent saline controls (grey bars), using tubulin as equal-loading control.

B. Silencing of miR-21 by #3205 leads to increased levels of Pcd4 protein levels *in vivo*.

C. Mice were injected with saline or 25 mg/kg LNA-antimiR (#3205) over 14 days every other day, with a total of 5 doses. Mice were sacrificed and protein was isolated from kidney and subjected to Western blot analysis with Pcd4 antibody. A. Gel image showing Pcd4 and Gapdh (internal standard) protein in treated (antimiR-21; 8-mer) vs. untreated (saline) kidney samples (M1, mouse 1; M2, mouse 2). B. Quantification of Pcd4 protein levels in kidneys of LNA-antimiR-treated mice (dark grey bars), normalized against the average of equivalent saline controls (light grey bars), using Gapdh as loading control.

## DETAILED DESCRIPTION OF THE INVENTION

**[0018]** Short oligonucleotides which incorporate LNA are known from the *in vitro* reagents area, (see for example WO2005/098029 and WO 2006/069584). However the molecules designed for diagnostic or reagent use are very different in design than those for *in vivo* or pharmaceutical use. For example, the terminal nucleotides of the reagent oligos are typically not LNA, but DNA, and the internucleoside linkages are typically other than phosphorothioate, the preferred linkage for use in the oligonucleotides of the present invention. The invention therefore provides for a novel class of oligonucleotides (referred to herein as oligomers) per se.

**[0019]** The following embodiments refer to certain embodiments of the oligomer of the invention, which may be used in a pharmaceutical composition. Aspects which refer to the oligomer may also refer to the contiguous nucleotide sequence, and vice versa.

### The Oligomer

**[0020]** The oligomer of the invention is a single stranded oligonucleotide wherein LNA, form the entire contiguous nucleotide sequence of the oligonucleotide. The nucleotide sequence of the oligomer consists of a contiguous nucleotide

sequence.

**[0021]** The term "oligonucleotide" (or simply "oligo"), which is used interchangeably with the term "oligomer" refers, in the context of the present invention, to a molecule formed by covalent linkage of two or more nucleotides. When used in the context of the oligonucleotide of the invention (also referred to the single stranded oligonucleotide), the term "oligonucleotide" may have, in one embodiment, for example have between 7 - 10 nucleotides, such as in individual embodiments, 7, 8, 9, or 10.

**[0022]** The term 'nucleotide' refers to nucleotides, such as DNA and RNA, and nucleotide analogues. It should be recognised that, in some aspects, the term nucleobase may also be used to refer to a nucleotide which may be either naturally occurring or non-naturally occurring - in this respect the term nucleobase and nucleotide may be used interchangeably herein.

**[0023]** In some embodiments, the contiguous nucleotide sequence consists of 7 nucleotide LNA units. In some embodiments, the contiguous nucleotide sequence consists of 8 nucleotide LNA units. In some embodiments, the contiguous nucleotide sequence consists of 9 nucleotide LNA units.

**[0024]** As described below, the contiguous nucleotide sequence consist only of LNA units (including linkage groups, such as phosphorothioate linkages),

**[0025]** The oligomer may consist of the contiguous nucleotide sequence.

**[0026]** In a specially preferred embodiment, all the nucleotide analogues are LNA. In a further All the nucleotides of the oligomer are LNA. In a further preferred embodiment, all nucleotides of the oligomer are LNA and all internucleoside linkage groups are phosphothioate.

**[0027]** Herein, the term "nitrogenous base" is intended to cover purines and pyrimidines, such as the DNA nucleobases A, C, T and G, the RNA nucleobases A, C, U and G, as well as non-DNA/RNA nucleobases, such as 5-methylcytosine (MeC), isocytosine, pseudoisocytosine, 5-bromouracil, 5-propynyluracil, 5-propynyl-6-fluoroluracil, 5-methylthiazoleuracil, 6-aminopurine, 2-aminopurine, inosine, 2,6-diaminopurine, 7-propyne-7-deazaadenine, 7-propyne-7-deazaguanine and 2-chloro-6-aminopurine, in particular MeC. It will be understood that the actual selection of the non-DNA/RNA nucleobase will depend on the corresponding (or matching) nucleotide present in the microRNA strand which the oligonucleotide is intended to target. For example, in case the corresponding nucleotide is G it will normally be necessary to select a non-DNA/RNA nucleobase which is capable of establishing hydrogen bonds to G. In this specific case, where the corresponding nucleotide is G, a typical example of a preferred non-DNA/RNA nucleobase is MeC.

**[0028]** It should be recognised that the term in 'one embodiment' should not necessarily be limited to refer to one specific embodiment, but may refer to a feature which may be present in 'some embodiments', or even as a generic feature of the invention. Likewise, the use of the term 'some embodiments' may be used to describe a feature of one specific embodiment, or a collection of embodiments, or even as a generic feature of the invention.

**[0029]** The terms "corresponding to" and "corresponds to" refer to the comparison between the nucleotide sequence of the oligomer or contiguous nucleotide sequence (a first sequence) and the equivalent contiguous nucleotide sequence of a further sequence selected from either i) a sub-sequence of the reverse complement of the microRNA nucleic acid target (such as a microRNA target selected from SEQ ID 40 - SEQ ID 976, and/or ii) the sequence of nucleotides provided herein such as the group consisting of SEQ ID NO 977 - 1913, or SEQ ID NO 1914-2850, or SEQ ID NO 2851 - 3787. Nucleotide analogues are compared directly to their equivalent or corresponding nucleotides. A first sequence which corresponds to a further sequence under i) or ii) typically is identical to that sequence over the length of the first sequence (such as the contiguous nucleotide sequence).

**[0030]** When referring to the length of a nucleotide molecule as referred to herein, the length corresponds to the number of monomer units, *i.e.* nucleotides, irrespective as to whether those monomer units are nucleotides or nucleotide analogues. With respect to nucleotides or nucleobases, the terms monomer and unit are used interchangeably herein.

**[0031]** It should be understood that when the term "about" is used in the context of specific values or ranges of values, the disclosure should be read as to include the specific value or range referred to.

**[0032]** As used herein, "hybridisation" means hydrogen bonding, which may be Watson-Crick, Hoogsteen, reversed Hoogsteen hydrogen bonding, etc., between complementary nucleoside or nucleotide bases. The four nucleobases commonly found in DNA are G, A, T and C of which G pairs with C, and A pairs with T. In RNA T is replaced with uracil (U), which then pairs with A. The chemical groups in the nucleobases that participate in standard duplex formation constitute the Watson-Crick face. Hoogsteen showed a couple of years later that the purine nucleobases (G and A) in addition to their Watson-Crick face have a Hoogsteen face that can be recognised from the outside of a duplex, and used to bind pyrimidine oligonucleotides via hydrogen bonding, thereby forming a triple helix structure.

**[0033]** In the context of the present invention "complementary" refers to the capacity for precise pairing between two nucleotides sequences with one another. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the corresponding position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The DNA or RNA strand are considered complementary to each other when a sufficient number of nucleotides in the oligonucleotide can form hydrogen bonds with corresponding nucleotides in the target DNA or RNA to enable the formation of a stable complex. To be



stable *in vitro* or *in vivo* the sequence of an oligonucleotide need not be 100% complementary to its target microRNA. The terms "complementary" and "specifically hybridisable" thus imply that the oligonucleotide binds sufficiently strong and specific to the target molecule to provide the desired interference with the normal function of the target whilst leaving the function of non-target RNAs unaffected. However, in one preferred embodiment the term complementary shall mean 100% complementary or fully complementary.

**[0034]** In a preferred example the oligonucleotide of the invention is 100% complementary to a miRNA sequence, such as a human microRNA sequence, or one of the microRNA sequences referred to herein.

**[0035]** In a preferred example, the oligonucleotide of the invention comprises a contiguous sequence, which is 100% complementary to the seed region of the human microRNA sequence.

**[0036]** Preferably, the term "microRNA" or "miRNA", in the context of the present invention, means an RNA oligonucleotide consisting of between 18 to 25 nucleotides in length. In functional terms miRNAs are typically regulatory endogenous RNA molecules.

**[0037]** The terms "target microRNA" or "target miRNA" refer to a microRNA with a biological role in human disease, e.g. an upregulated, oncogenic miRNA or a tumor suppressor miRNA in cancer, thereby being a target for therapeutic intervention of the disease in question.

**[0038]** The terms "target gene" or "target mRNA" refer to regulatory mRNA targets of microRNAs, in which said "target gene" or "target mRNA" is regulated post-transcriptionally by the microRNA based on near-perfect or perfect complementarity between the miRNA and its target site resulting in target mRNA cleavage; or limited complementarity, often conferred to complementarity between the so-called seed sequence (nucleotides 2-7 of the miRNA) and the target site resulting in translational inhibition of the target mRNA.

**[0039]** In the context of the present invention the oligonucleotide is single stranded, this refers to the situation where the oligonucleotide is in the absence of a complementary oligonucleotide - i.e. it is not a double stranded oligonucleotide complex, such as an siRNA. In one embodiment, the composition according to the invention does not comprise a further oligonucleotide which has a region of complementarity with the oligomer of 5 or more, such as 6, 7, 8, 9, or 10 consecutive nucleotides, such as eight or more.

### Length

**[0040]** Surprisingly we have found that such short 'antimiRs' provide an improved specific inhibition of microRNAs *in vivo*, whilst retaining remarkable specificity for the microRNA target. A further benefit has been found to be the ability to inhibit several microRNAs simultaneously due to the conservation of homologous short sequences between microRNA species - such as the seed regions as described herein. According to the present invention, it has been found that it is particularly advantageous to have short oligonucleotides of 7, 8, 9, 10 nucleotides, such as 7, 8 or 9 nucleotides.

### Sequences

**[0041]** The contiguous nucleotide sequence is complementary (such as 100% complementary - i.e. perfectly complementary) to a corresponding region of a mammalian, human or viral microRNA (miRNA) sequence, preferably a human or viral miRNA sequence.

**[0042]** The microRNA sequence may suitably be a mature microRNA. In some embodiments the microRNA may be a microRNA precursor.

**[0043]** The human microRNA sequence may be selected from SEQ ID No 1 - 558 as disclosed in WO2008/046911. As described in WO2008/046911, these microRNAs are associated with cancer.

**[0044]** The viral microRNA sequence may, in some embodiments, be selected from the group consisting of Herpes simplex virus 1, Kaposi sarcoma-associated herpesvirus, Epstein Barr virus and Human cytomegalovirus.

**[0045]** In one embodiment, the contiguous nucleotide sequence is complementary (such as 100% complementary) to a corresponding region of a miRNA sequence selected from the group of miRNAs listed in table 1. Table 1 provides 7mer, 8mer and 9mer oligomers which target human and viral microRNAs published in miRBase (Release 12.0 - <http://microrna.sanger.ac.uk/sequences/>).

**[0046]** In some embodiments, the oligomers according to the invention may consist of or comprise a contiguous nucleotide sequence which is complementary to a corresponding microRNA sequence selected from the group consisting of miR-1, miR-10b, miR-17-3p, miR-18, miR-19a, miR-19b, miR-20, miR-21, miR-34a, miR-93, miR-106a, miR-106b, miR-122, miR-133, miR-134, miR-138, miR-155, miR-192, miR-194, miR-221, miR-222, miR-375.

**[0047]** Therefore, in one embodiment, the miRNA (*i.e.* target miRNA) is selected from the group consisting of miR-1, miR-10b, miR-17-3p, miR-18, miR-19a, miR-19b, miR-20, miR-21, miR-34a, miR-93, miR-106a, miR-106b, miR-122, miR-133, miR-134, miR-138, miR-155, miR-192, miR-194, miR-221, miR-222, and miR-375.

**[0048]** In one embodiment, the miRNA target is a member of the miR 17 - 92 cluster, such as miR 17, miR 106a, miR 106b, miR 18, miR 19a, miR 19b/1, miR 19b/2, miR20/93, miR92/1, miR92/2 and miR25.

[0049] In some embodiments the contiguous nucleotide sequence is complementary to a corresponding region of a microRNA (miRNA) sequence selected from the group consisting of miR-21, miR-155, miR-221, miR-222, and miR-122.

[0050] In some embodiments said miRNA is selected from the group consisting of miR-1, miR-10, miR-29, miR-125b, miR-126, miR-133, miR-141, miR-143, miR-200b, miR-206, miR-208, miR-302, miR-372, miR-373, miR-375, and miR-520c/e.

[0051] In some embodiments the contiguous nucleotide sequence is complementary to a corresponding region of a microRNA (miRNA) sequence present in the miR 17 - 92 cluster, such as a microRNA selected from the group consisting of miR-17-5p, miR-20a/b, miR-93, miR-106a/b, miR-18a/b, miR-19a/b, miR-25, miR-92a, , miR-363.

[0052] In one embodiment, the miRNA (*i.e* target miRNA) is miR-21, such as hsa-miR-21. In one embodiment, the miRNA (*i.e* target miRNA) is miR-122, such as hsa-miR-122. In one embodiment, the miRNA (*i.e* target miRNA) is miR-19b, such as hsa-miR-19b. In one embodiment, the miRNA (*i.e* target miRNA) is miR-155, such as hsa-miR-155. In one embodiment, the miRNA (*i.e* target miRNA) is miR-375, such as hsa-miR-375. In one embodiment, the miRNA (*i.e* target miRNA) is miR-375, such as hsa-miR-106b.

[0053] Suitably, the contiguous nucleotide sequence may be complementary to a corresponding region of the microRNA, such as a hsa-miR selected from the group consisting of 19b, 21, 122, 155 and 375.

### ***The Seed Region and Seedmers***

[0054] The inventors have found that carefully designed short single stranded oligonucleotides comprising or consisting of nucleotide analogues, such as high affinity nucleotide analogues such as locked nucleic acid (LNA) units, show significant silencing of microRNAs, resulting in reduced microRNA levels. It was found that tight binding of said oligonucleotides to the so-called seed sequence, typically nucleotides 2 to 8 or 2 to 7, counting from the 5' end, of the target microRNAs was important. Nucleotide 1 of the target microRNAs is a non-pairing base and is most likely hidden in a binding pocket in the Ago 2 protein. Whilst not wishing to be bound to a specific theory, the present inventors consider that by selecting the seed region sequences, particularly with oligonucleotides that comprise LNA in the region which is complementary to the seed region, the duplex between miRNA and oligonucleotide is particularly effective in targeting miRNAs, avoiding off target effects, and possibly providing a further feature which prevents RISC directed miRNA function.

[0055] The inventors have found that microRNA silencing is even more enhanced when LNA-modified single stranded oligonucleotides do not contain a nucleotide at the 3' end corresponding to this non-paired nucleotide 1. It was further found that at least two LNA units in the 3' end of the oligonucleotides according to the present invention made said oligonucleotides highly nuclease resistant.

[0056] In one embodiment, the first or second 3' nucleotide of the oligomer corresponds to the second 5' nucleotide of the microRNA sequence.

[0057] In one embodiment, nucleotide units 1 to 6 (inclusive) of the oligomer as measured from the 3' end the region of the oligomer are complementary to the microRNA seed region sequence.

[0058] In one embodiment, nucleotide units 1 to 7 (inclusive) of the oligomer as measured from the 3' end the region of the oligomer are complementary to the microRNA seed region sequence.

[0059] In one embodiment, nucleotide units 2 to 7 (inclusive) of the oligomer as measured from the 3' end the region of the oligomer are complementary to the microRNA seed region sequence.

[0060] In one embodiment, the oligomer comprises at least one LNA unit, in a position which is within the region complementary to the miRNA seed region. The oligomer may, in one embodiment comprise at between one and 6 or between 1 and 7 nucleotide analogue units, such as between 1 and 6 and 1 and 7 LNA units, in a position which is within the region complementary to the miRNA seed region.

[0061] In one embodiment, the contiguous nucleotide sequence consists of or comprises a sequence which is complementary (such as 100% complementary) to the seed sequence of said microRNA.

[0062] In one embodiment, the contiguous nucleotide sequence consists of or comprises a sequence selected from any one of the seedmer sequences listed in table 1.

[0063] In one embodiment, the 3' nucleotide of the seedmer forms the 3' most nucleotide of the contiguous nucleotide sequence, wherein the contiguous nucleotide sequence may, optionally, comprise one or two further nucleotide 5' to the seedmer sequence.

[0064] In one embodiment, the oligomer does not comprise a nucleotide which corresponds to the first nucleotide present in the microRNA sequence counted from the 5' end.

[0065] In one embodiment, the oligonucleotide according to the invention does not comprise a nucleotide at the 3' end that corresponds to the first 5' end nucleotide of the target microRNA.

### ***Nucleotide Analogues***

[0066] According to the present invention, it has been found that it is particularly advantageous to have short oligo-

nucleotides of 7, 8, 9, 10 nucleotides, such as 7, 8 or 9 nucleotides, wherein 100% of the nucleotide units of the oligomer are Locked Nucleic Acid (LNA) nucleotide unit.

**[0067]** In some embodiments, the oligonucleotide of the invention is 7, 8 or 9 nucleotides long, and comprises a contiguous nucleotide sequence which is complementary to a seed region of a human or viral microRNA, and wherein 100% of the nucleotides are Locked Nucleic Acid (LNA) nucleotide units.

**[0068]** In such oligomers, at least 75% of the internucleoside linkage groups are phosphorothioate linkages.

**[0069]** All of the nucleotide units of the contiguous nucleotide sequence are LNA nucleotide units.

**[0070]** In one embodiment, the contiguous nucleotide sequence comprises or consists of 7, 8, 9 or 10 contiguous, LNA nucleotide units.

**[0071]** In a further preferred embodiment, the oligonucleotide of the invention is 7, 8 or 9 nucleotides long, and comprises a contiguous nucleotide sequence which is complementary to a seed region of a human or viral microRNA, and wherein all of the nucleotides are LNA, and wherein at least 80%, such as 85%, such as 90%, such as 95%, such as 100% of the internucleotide bonds are phosphorothioate bonds. It will be recognised that the contiguous nucleotide sequence of the oligomer (a seedmer) may extend beyond the seed region.

**[0072]** In some embodiments, the oligonucleotide of the invention is 7 nucleotides long, which are all LNA.

**[0073]** High affinity nucleotide analogues are nucleotide analogues which result in oligonucleotides which has a higher thermal duplex stability with a complementary RNA nucleotide than the binding affinity of an equivalent DNA nucleotide. This may be determined by measuring the  $T_m$ .

**[0074]** In one embodiment, the oligomer has at least 7 LNA units, such as at least 8 LNA units, such as at least 9 LNA units, such as 10 LNA.

**[0075]** In one embodiment wherein at least one of the LNA units, is either cytosine or guanine, such as between 1 - 10 of the of the LNA units, is either cytosine or guanine, such as 2, 3, 4, 5, 6, 7, 8, or 9 of the of the LNA units, is either cytosine or guanine.

**[0076]** In one embodiment at least two of the LNA units are either cytosine or guanine. In one embodiment at least three of the LNA units are either cytosine or guanine. In one embodiment at least four of the LNA units are either cytosine or guanine. In one embodiment at least five of the LNA units are either cytosine or guanine. In one embodiment at least six of the nucleotide LNA units are either cytosine or guanine. In one embodiment at least seven of the LNA units are either cytosine or guanine. In one embodiment at least eight of the LNA units are either cytosine or guanine.

**[0077]** In a preferred embodiment the LNA units have a higher thermal duplex stability for a complementary RNA nucleotide than the binding affinity of an equivalent DNA nucleotide to said complementary RNA nucleotide.


**[0078]** In one embodiment, the LNA units confer enhanced serum stability to the single stranded oligonucleotide.

**[0079]** Whilst the specific SEQ IDs in the sequence listing and table 1 refer to oligomers of LNA monomers with phosphorothioate (PS) backbone, it will be recognised that the invention also encompasses the use of other linkages, as long as 75% of linkages are phosphorothioate. As such, the sequence of nucleotides (bases) shown in the sequence listings may be of LNA such as LNA/PS, LNA or may be oligomers containing alternative linkage chemistry, whilst retaining the same base sequence (A, T, C or G) as long as 75% of linkages are phosphorothioate.

## LNA

**[0080]** When used in the present context, the terms "LNA unit", "LNA monomer", "LNA residue", "locked nucleic acid unit", "locked nucleic acid monomer" or "locked nucleic acid residue", refer to a bicyclic nucleoside analogue. LNA units are described in *inter alia* WO 99/14226, WO 00/56746, WO 00/56748, WO 01/25248, WO 02/28875, WO 03/006475 and WO 03/095467. The LNA unit may also be defined with respect to its chemical formula. Thus, an "LNA unit", as used herein, has the chemical structure shown in Scheme 1 below:


Scheme 1



wherein

X is selected from the group consisting of O, S and  $\text{NR}^{\text{H}}$ , where  $\text{R}^{\text{H}}$  is H or  $\text{C}_{1-4}$ -alkyl; Y is  $(-\text{CH}_2)_r$ , where r is an integer of 1-4; and B is a nitrogenous base.


[0081] In a preferred embodiment of the invention,  $r$  is 1 or 2, in particular 1, *i.e.* a preferred LNA unit has the chemical structure shown in Scheme 2 below:



wherein X and B are as defined above.


15 [0082] In an interesting embodiment, the LNA units incorporated in the oligonucleotides of the invention are independently selected from the group consisting of thio-LNA units, amino-LNA units and oxy-LNA units.

[0083] Thus, the thio-LNA unit may have the chemical structure shown in Scheme 3 below:



wherein B is as defined above.


[0084] Preferably, the thio-LNA unit is in its beta-D-form, *i.e.* having the structure shown in 3A above. likewise, the amino-LNA unit may have the chemical structure shown in Scheme 4 below:



wherein B and  $R^H$  are as defined above.

[0085] Preferably, the amino-LNA unit is in its beta-D-form, *i.e.* having the structure shown in 4A above.

[0086] The oxy-LNA unit may have the chemical structure shown in Scheme 5 below:



wherein B is as defined above.

**[0087]** Preferably, the oxy-LNA unit is in its beta-D-form, i.e. having the structure shown in **5A** above. As indicated above, B is a nitrogenous base which may be of natural or non-natural origin. Specific examples of nitrogenous bases include adenine (A), cytosine (C), 5-methylcytosine (<sup>Me</sup>C), isocytosine, pseudoisocytosine, guanine (G), thymine (T), uracil (U), 5-bromouracil, 5-propynyluracil, 5-propyne-6, 5-methylthiazoleuracil, 6-aminopurine, 2-aminopurine, inosine, 2,6-diaminopurine, 7-propyne-7-deazaadenine, 7-propyne-7-deazaguanine and 2-chloro-6-aminopurine.

**[0088]** The term "thio-LNA unit" refers to an LNA unit in which X in Scheme 1 is S. A thio-LNA unit can be in both the beta-D form and in the alpha-L form. Generally, the beta-D form of the thio-LNA unit is preferred. The beta-D-form and alpha-L-form of a thio-LNA unit are shown in Scheme 3 as compounds **3A** and **3B**, respectively.

**[0089]** The term "amino-LNA unit" refers to an LNA unit in which X in Scheme 1 is NH or NR<sup>H</sup>, where R<sup>H</sup> is hydrogen or C<sub>1-4</sub>-alkyl. An amino-LNA unit can be in both the beta-D form and in the alpha-L form. Generally, the beta-D form of the amino-LNA unit is preferred. The beta-D-form and alpha-L-form of an amino-LNA unit are shown in Scheme 4 as compounds **4A** and **4B**, respectively.

**[0090]** The term "oxy-LNA unit" refers to an LNA unit in which X in Scheme 1 is O. An Oxy-LNA unit can be in both the beta-D form and in the alpha-L form. Generally, the beta-D form of the oxy-LNA unit is preferred. The beta-D form and the alpha-L form of an oxy-LNA unit are shown in Scheme 5 as compounds **5A** and **5B**, respectively.

**[0091]** In the present context, the term "C<sub>1-6</sub>-alkyl" is intended to mean a linear or branched saturated hydrocarbon chain wherein the longest chains has from one to six carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl and hexyl. A branched hydrocarbon chain is intended to mean a C<sub>1-6</sub>-alkyl substituted at any carbon with a hydrocarbon chain.

**[0092]** In the present context, the term "C<sub>1-4</sub>-alkyl" is intended to mean a linear or branched saturated hydrocarbon chain wherein the longest chains has from one to four carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl. A branched hydrocarbon chain is intended to mean a C<sub>1-4</sub>-alkyl substituted at any carbon with a hydrocarbon chain.

**[0093]** When used herein the term "C<sub>1-6</sub>-alkoxy" is intended to mean C<sub>1-6</sub>-alkyl-oxy, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentoxy, isopentoxy, neopentoxy and hexoxy.

**[0094]** In the present context, the term "C<sub>2-6</sub>-alkenyl" is intended to mean a linear or branched hydrocarbon group having from two to six carbon atoms and containing one or more double bonds. Illustrative examples of C<sub>2-6</sub>-alkenyl groups include allyl, homo-allyl, vinyl, crotyl, butenyl, butadienyl, pentenyl, pentadienyl, hexenyl and hexadienyl. The position of the unsaturation (the double bond) may be at any position along the carbon chain.

**[0095]** In the present context the term "C<sub>2-6</sub>-alkynyl" is intended to mean linear or branched hydrocarbon groups containing from two to six carbon atoms and containing one or more triple bonds. Illustrative examples of C<sub>2-6</sub>-alkynyl groups include acetylene, propynyl, butynyl, pentynyl and hexynyl. The position of unsaturation (the triple bond) may be at any position along the carbon chain. More than one bond may be unsaturated such that the "C<sub>2-6</sub>-alkynyl" is a di-yne or enedi-yne as is known to the person skilled in the art.

**[0096]** When referring to substituting a DNA unit by its corresponding LNA unit in the context of the present invention, the term "corresponding LNA unit" is intended to mean that the DNA unit has been replaced by an LNA unit containing the same nitrogenous base as the DNA unit that it has replaced, e.g. the corresponding LNA unit of a DNA unit containing the nitrogenous base A also contains the nitrogenous base A. The exception is that when a DNA unit contains the base C, the corresponding LNA unit may contain the base C or the base <sup>Me</sup>C, preferably <sup>Me</sup>C.

**[0097]** Herein, the term "non-LNA unit" refers to a nucleoside different from an LNA-unit, i.e. the term "non-LNA unit" includes a DNA unit as well as an RNA unit. A preferred non-LNA unit is a DNA unit.

**[0098]** The terms "unit", "residue" and "monomer" are used interchangeably herein.

**[0099]** The term "at least one" encompasses an integer larger than or equal to 1, such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 and so forth.

**[0100]** The terms "a" and "an" as used about a nucleotide, an agent, an LNA unit, etc., is intended to mean one or more. In particular, the expression "a component (such as a nucleotide, an agent, an LNA unit, or the like) selected from the group consisting of ..." is intended to mean that one or more of the cited components may be selected. Thus, expressions like "a component selected from the group consisting of A, B and C" is intended to include all combinations of A, B and C, i.e. A, B, C, A+B, A+C, B+C and A+B+C.

### **Internucleoside Linkages**

**[0101]** The term "internucleoside linkage group" is intended to mean a group capable of covalently coupling together two nucleotides, such as between DNA units, between DNA units and nucleotide analogues, between two non-LNA units, between a non-LNA unit and an LNA unit, and between two LNA units, etc. Examples include phosphate, phosphodiester groups and phosphorothioate groups.

**[0102]** In some embodiments, at least one of, the internucleoside linkage in the oligomer is phosphodiester. However

for *in vivo* use, phosphorothioate linkages may be preferred.

**[0103]** Typical internucleoside linkage groups in oligonucleotides are phosphate groups, but these may be replaced by internucleoside linkage groups differing from phosphate. In a further interesting embodiment of the invention, the oligonucleotide of the invention is modified in its internucleoside linkage group structure, i.e. the modified oligonucleotide comprises an internucleoside linkage group which differs from phosphate. Accordingly, in a preferred embodiment, the oligonucleotide according to the present invention comprises at least one internucleoside linkage group which differs from phosphate.

**[0104]** Specific examples of internucleoside linkage groups which differ from phosphate (-O-P(O)<sub>2</sub>-O-) include -O-P(O,S)-O-, -O-P(S)<sub>2</sub>-O-, -S-P(O)<sub>2</sub>-O-, -S-P(O,S)-O-, -S-P(S)<sub>2</sub>-O-, -O-P(O)<sub>2</sub>-S-, -O-P(O,S)-S-, -S-P(O)<sub>2</sub>-S-, -O-PO(R<sup>H</sup>)-O-, O-PO(OCH<sub>3</sub>)-O-, -O-PO(NR<sup>H</sup>)-O-, -O-PO(OCH<sub>2</sub>CH<sub>2</sub>S-R)-O-, -O-PO(BH<sub>3</sub>)-O-, -O-PO(NHR<sup>H</sup>)-O-, -O-P(O)<sub>2</sub>-NR<sup>H</sup>-, -NR<sup>H</sup>-P(O)<sub>2</sub>-O-, -NR<sup>H</sup>-CO-O-, -NR<sup>H</sup>-CO-NR<sup>H</sup>-, -O-CO-O-, -O-CO-NR<sup>H</sup>-, -NR<sup>H</sup>-CO-CH<sub>2</sub>-, -O-CH<sub>2</sub>-CO-NR<sup>H</sup>-, -O-CH<sub>2</sub>-CH<sub>2</sub>-NR<sup>H</sup>-, -CO-NR<sup>H</sup>-CH<sub>2</sub>-, -CH<sub>2</sub>-NR<sup>H</sup>-CO-, -O-CH<sub>2</sub>-CH<sub>2</sub>-S-, -S-CH<sub>2</sub>-CH<sub>2</sub>-O-, -S-CH<sub>2</sub>-CH<sub>2</sub>-S-, -CH<sub>2</sub>-SO<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CO-NR<sup>H</sup>-, -O-CH<sub>2</sub>-CH<sub>2</sub>-NR<sup>H</sup>-CO-, -CH<sub>2</sub>-NCH<sub>3</sub>-O-CH<sub>2</sub>-, where R<sup>H</sup> is hydrogen or C<sub>1-4</sub>-alkyl.

**[0105]** When the internucleoside linkage group is modified, the internucleoside linkage group is preferably a phosphorothioate group (-O-P(O,S)-O-). In a preferred embodiment, all internucleoside linkage groups of the oligonucleotides according to the present invention are phosphorothioate.

**[0106]** The internucleoside linkage may be selected from the group consisting of: -O-P(O)<sub>2</sub>-O-, -O-P(O,S)-O-, -O-P(S)<sub>2</sub>-O-, -S-P(O)<sub>2</sub>-O-, -S-P(O,S)-O-, -S-P(S)<sub>2</sub>-O-, -O-P(O)<sub>2</sub>-S-, -O-P(O,S)-S-, -S-P(O)<sub>2</sub>-S-, -O-PO(R<sup>H</sup>)-O-, O-PO(OCH<sub>3</sub>)-O-, -O-PO(NR<sup>H</sup>)-O-, -O-PO(OCH<sub>2</sub>CH<sub>2</sub>S-R)-O-, -O-PO(BH<sub>3</sub>)-O-, -O-PO(NHR<sup>H</sup>)-O-, -O-P(O)<sub>2</sub>-NR<sup>H</sup>-, -NR<sup>H</sup>-P(O)<sub>2</sub>-O-, -NR<sup>H</sup>-CO-O-, -NR<sup>H</sup>-CO-NR<sup>H</sup>-, and/or the internucleoside linkage may be selected from the group consisting of: -O-CO-O-, -O-CO-NR<sup>H</sup>-, -NR<sup>H</sup>-CO-CH<sub>2</sub>-, -O-CH<sub>2</sub>-CO-NR<sup>H</sup>-, -O-CH<sub>2</sub>-CH<sub>2</sub>-NR<sup>H</sup>-, -CO-NR<sup>H</sup>-CH<sub>2</sub>-, -CH<sub>2</sub>-NR<sup>H</sup>-CO-, -O-CH<sub>2</sub>-CH<sub>2</sub>-S-, -S-CH<sub>2</sub>-CH<sub>2</sub>-O-, -S-CH<sub>2</sub>-CH<sub>2</sub>-S-, -CH<sub>2</sub>-SO<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CO-NR<sup>H</sup>-, -O-CH<sub>2</sub>-CH<sub>2</sub>-NR<sup>H</sup>-CO-, -CH<sub>2</sub>-NCH<sub>3</sub>-O-CH<sub>2</sub>-, where R<sup>H</sup> is selected from hydrogen and C<sub>1-4</sub>-alkyl. Suitably, in some embodiments, sulphur (S) containing internucleoside linkages as provided above may be preferred. The internucleoside linkages may be independently selected, or all be the same, such as phosphorothioate linkages.

**[0107]** In one embodiment, 80% or 85% or 90% or 95% or all of the internucleoside linkages present between the nucleotide units of the contiguous nucleotide sequence are phosphorothioate internucleoside linkages.

### ***Micromir oligonucleotides targeting more than one microRNA***

**[0108]** In one embodiment, the contiguous nucleotide sequence is complementary to the corresponding sequence of at least two miRNA sequences such as 2, 3, 4, 5, 6, 7, 8, 9, or 10 miRNA sequence,. The use of a single universal base may allow a single oligomer of the invention to target two independent microRNAs which either one or both have a single mismatch in the region which corresponds to oligomer at the position where the universal nucleotide is positioned.

**[0109]** In one embodiment, the contiguous nucleotide sequence consists of or comprises a sequence which is complementary to the sequence of at least two miRNA seed region sequences such as 2, 3, 4, 5, 6, 7, 8, 9, or 10 miRNA seed region sequences.

**[0110]** In one embodiment, the contiguous nucleotide sequence is complementary to the corresponding region of both miR-221 and miR-222.

**[0111]** In one embodiment, the contiguous nucleotide sequence is complementary to the corresponding region of more than one member of the miR-17-92 cluster - such as two or more or all of miR-17-5p, miR-20a/b, miR-93, miR-106a/b; or two or more or all of miR-25, miR-92a and miR-363.

**[0112]** In one embodiment, the contiguous nucleotide sequence consists of or comprises a sequence that is complementary to 5'GCTACAT3'.

### ***Pharmaceutical Composition and Medical Application***

**[0113]** The invention provides for a pharmaceutical composition comprising the oligomer according to the invention, and a pharmaceutically acceptable diluent, carrier, salt or adjuvant.

**[0114]** The invention further provides for the use of an oligonucleotide according to the invention, such as those which may form part of the pharmaceutical composition, for the manufacture of a medicament for the treatment of a disease or medical disorder associated with the presence or over-expression (upregulation) of the microRNA.

**[0115]** The invention further provides for a method for the treatment of a disease or medical disorder associated with the presence or over-expression of the microRNA, comprising the step of administering a composition (such as the pharmaceutical composition) according to the invention to a person in need of treatment.

**[0116]** The invention further provides for a method for reducing the effective amount of a miRNA in a cell or an organism, comprising administering a composition (such as the pharmaceutical composition) according to the invention or a oligomer

according to the invention to the cell or the organism. Reducing the effective amount in this context refers to the reduction of functional miRNA present in the cell or organism. It is recognised that the preferred oligonucleotides according to the invention may not always significantly reduce the actual amount of miRNA in the cell or organism as they typically form very stable duplexes with their miRNA targets. The reduction of the effective amount of the miRNA in a cell may, in one

embodiment, be measured by detecting the level of de-repression of the miRNA's target in the cell.  
**[0117]** The invention further provides for a method for de-repression of a target mRNA of a miRNA in a cell or an organism, comprising administering a composition (such as the pharmaceutical composition) or an oligomer according to the invention to the cell or the organism.

**[0118]** The invention further provides for the use of an oligomer of the invention, for the manufacture of a medicament for the treatment of a disease or medical disorder associated with the presence or over-expression of the microRNA.

**[0119]** In one embodiment the medical condition (or disease) is hepatitis C (HCV), and the miRNA is miR-122.

**[0120]** In one embodiment, the pharmaceutical composition according to the invention is for use in the treatment of a medical disorder or disease selected from the group consisting of: hepatitis C virus infection and hypercholesterolemia and related disorders, and cancers.

**[0121]** In one embodiment the medical disorder or disease is a CNS disease, such as a CNS disease where one or more microRNAs are known to be indicated.

**[0122]** In the context of hypercholesterolemia related disorders refers to diseases such as atherosclerosis or hyperlipidemia. Further examples of related diseases also include different types of HDL/LDL cholesterol imbalance; dyslipidemias, e.g., familial combined hyperlipidemia (FCHL), acquired hyperlipidemia, statin-resistant hypercholesterolemia; coronary artery disease (CAD) coronary heart disease (CHD), atherosclerosis.

**[0123]** In one embodiment, the pharmaceutical composition according to the invention further comprises a second independent active ingredient that is an inhibitor of the VLDL assembly pathway, such as an ApoB inhibitor, or an MTP inhibitor (such as those disclosed in US 60/977,497).

**[0124]** The disclosure provides a method for the treatment of a disease or medical disorder associated with the presence or over-expression of the microRNA, comprising the step of administering a composition (such as the pharmaceutical composition) comprising an oligomer of the invention to a person in need of treatment.

**[0125]** The disclosure provides a method for reducing the effective amount of a miRNA target (i.e. 'available' miRNA) in a cell or an organism, comprising administering a composition (such as the pharmaceutical composition) comprising an oligomer of the invention to the cell or the organism.

**[0126]** It should be recognised that "reducing the effective amount" of one or more microRNAs in a cell or organism, refers to the inhibition of the microRNA function in the cell or organism. The cell is preferably a mammalian cell or a human cell which expresses the microRNA or microRNAs.

**[0127]** The disclosure provides a method for de-repression of a target mRNA of a miRNA in a cell or an organism, comprising an oligomer of the invention (or a composition comprising said oligonucleotide) to the cell or the organism.

**[0128]** As mentioned above, microRNAs are related to a number of diseases. Hence, a fourth aspect of the invention relates to the use of an oligonucleotide as defined herein for the manufacture of a medicament for the treatment of a disease associated with the expression of microRNAs selected from the group consisting of spinal muscular atrophy, Tourette's syndrome, hepatitis C, fragile X mental retardation, DiGeorge syndrome and cancer, such as in non limiting example, chronic lymphocytic leukemia, breast cancer, lung cancer and colon cancer, in particular cancer.

#### **Methods of Synthesis**

**[0129]** The disclosure provides a method for the synthesis of an oligomer of the invention targeted against a human microRNA, such as an oligomer described herein, said method comprising the steps of:

- a. Optionally selecting a first nucleotide, counting from the 3' end, which is a LNA nucleotide.
- b. Optionally selecting a second nucleotide, counting from the 3' end, which is a LNA nucleotide.
- c. Selecting a region of the oligomer which corresponds to the miRNA seed region, wherein said region is as defined herein.
- d. Selecting a seventh and optionally an eight nucleotides defined herein.
- e. Optionally selecting one or two further 5' terminal of the oligomer is as defined herein; wherein the synthesis is performed by sequential synthesis of the regions defined in steps a - e, wherein said synthesis may be performed in either the 3'-5' (a to f) or 5' - 3' (e to a) direction, and wherein said oligomer is complementary to a sequence of the miRNA target.

**[0130]** The invention further provides for a method for the preparation of an oligomer (such as an oligomer according to the invention), said method comprising the steps of a) comparing the sequences of two or more miRNA sequences to identify two or more miRNA sequences which comprise a common contiguous nucleotide sequence of at least 7

nucleotides in length, such as 7, 8, 9 or 10 nucleotides in length (i.e. a sequence found in both non-identical miRNAs),  
 b) preparing an oligomer sequence which consists or comprises of a contiguous nucleotide sequence which is complementary to said common contiguous nucleotide sequence, wherein said oligomer is, as according to the invention. In a preferred example, the common contiguous nucleotide sequence consists or comprises of the seed region  
 5 of each of said two or more miRNA sequences (which comprise a common contiguous nucleotide sequence of at least 6 nucleotides in length). In one embodiment, the seed regions of the two or more miRNAs are identical. Suitably the oligomer consists or comprises a seedmer sequence of 7 or 8 nucleotides in length which comprises of a sequence which is complementary to said two or more miRNAs. This method may be used in conjunction with step c of the above method.

10 **[0131]** The method for the synthesis of the oligomer according to the invention may be performed using standard solid phase oligonucleotide synthesis.

**[0132]** In one embodiment, the method for the synthesis of an oligomer targeted against a human microRNA, is performed in the 3' to 5' direction a - e.

15 **[0133]** A further aspect of the invention is a method to reduce the levels of target microRNA by contacting the target microRNA to an oligonucleotide as defined herein, wherein the oligonucleotide (i) is complementary to the target microRNA sequence (ii) does not contain a nucleotide at the 3' end that corresponds to the first 5' end nucleotide of the target microRNA.

### 20 *Duplex stability and $T_m$*

**[0134]** In one embodiment, the oligomer of the invention is capable of forming a duplex with a complementary single stranded RNA nucleic acid molecule (typically of about the same length of said single stranded oligonucleotide) with phosphodiester internucleoside linkages, wherein the duplex has a  $T_m$  of between 30°C and 70°C or 80°C, such as between 30°C and 60°C or 70°C, or between 30°C and 50°C or 60°C. In one embodiment the  $T_m$  is at least 40°C.  
 25  $T_m$  may be determined by determining the  $T_m$  of the oligomer and a complementary RNA target in the following buffer conditions: 100mM NaCl, 0.1mM EDTA, 10mM Na-phosphate, pH 7.0 (see examples for a detailed protocol). A high affinity analogue may be defined as an analogue which, when used in the oligomer of the invention, results in an increase in the  $T_m$  of the oligomer as compared to an identical oligomer which contains only DNA bases.

### 30 *Conjugates*

**[0135]** In one embodiment, said oligomer is conjugated with one or more non-nucleotide (or polynucleotide) compounds.

35 **[0136]** In the context the term "conjugate" is intended to indicate a heterogenous molecule formed by the covalent attachment ("conjugation") of the oligomer as described herein to one or more non-nucleotide, or non-polynucleotide moieties. Examples of non-nucleotide or non-polynucleotide moieties include macromolecular agents such as proteins, fatty acid chains, sugar residues, glycoproteins, polymers, or combinations thereof. Typically proteins may be antibodies for a target protein. Typical polymers may be polyethylene glycol.

40 **[0137]** Therefore, in various embodiments, the oligomer of the invention may comprise both a polynucleotide region which typically consists of a contiguous sequence of nucleotides, and a further non-nucleotide region. When referring to the oligomer of the invention consisting of a contiguous nucleotide sequence, the compound may comprise non-nucleotide components, such as a conjugate component.

**[0138]** In various embodiments of the invention the oligomeric compound is linked to ligands/conjugates, which may be used, e.g. to increase the cellular uptake of oligomeric compounds. WO2007/031091 provides suitable ligands and conjugates.

45 **[0139]** The invention also provides for a conjugate comprising the compound according to the invention as herein described, and at least one non-nucleotide or non-polynucleotide moiety covalently attached to said compound. Therefore, in various embodiments where the compound of the invention consists of a specified nucleic acid or nucleotide sequence, as herein disclosed, the compound may also comprise at least one non-nucleotide or non-polynucleotide moiety (e.g. not comprising one or more nucleotides or nucleotide analogues) covalently attached to said compound.

50 **[0140]** Conjugation (to a conjugate moiety) may enhance the activity, cellular distribution or cellular uptake of the oligomer of the invention. Such moieties include, but are not limited to, antibodies, polypeptides, lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g. Hexyl-s-tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipids, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-o-hexadecyl-rac-glycero-3-h-phosphonate, a polyamine or a polyethylene glycol chain, an adamantane acetic acid, a palmitoyl moiety, an octadecylamine or hexylamino-carbonyl-oxcholesterol moiety.



55 **[0141]** The oligomers of the invention may also be conjugated to active drug substances, for example, aspirin, ibuprofen, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.



[0142] In certain embodiments the conjugated moiety is a sterol, such as cholesterol.

[0143] In various embodiments, the conjugated moiety comprises or consists of a positively charged polymer, such as a positively charged peptides of, for example between 1 -50, such as 2 - 20 such as 3 - 10 amino acid residues in length, and/or polyalkylene oxide such as polyethylglycol(PEG) or polypropylene glycol - see WO 2008/034123. Suitably the positively charged polymer, such as a polyalkylene oxide may be attached to the oligomer of the invention via a linker such as the releasable inker described in WO 2008/034123.

[0144] By way of example, the following conjugate moieties may be used in the conjugates of the invention:



### Activated oligomers

20 [0145] The term "activated oligomer," as used herein, refers to an oligomer of the invention that is covalently linked (i.e., functionalized) to at least one functional moiety that permits covalent linkage of the oligomer to one or more conjugated moieties, i.e., moieties that are not themselves nucleic acids or monomers, to form the conjugates herein described. Typically, a functional moiety will comprise a chemical group that is capable of covalently bonding to the oligomer via, e.g., a 3'-hydroxyl group or the exocyclic NH<sub>2</sub> group of the adenine base, a spacer that is preferably hydrophilic and a terminal group that is capable of binding to a conjugated moiety (e.g., an amino, sulfhydryl or hydroxyl group). In some embodiments, this terminal group is not protected, e.g., is an NH<sub>2</sub> group. In other embodiments, the terminal group is protected, for example, by any suitable protecting group such as those described in "Protective Groups in Organic Synthesis" by Theodora W Greene and Peter G M Wuts, 3rd edition (John Wiley & Sons, 1999). Examples of suitable hydroxyl protecting groups include esters such as acetate ester, aralkyl groups such as benzyl, diphenylmethyl, or triphenylmethyl, and tetrahydropyranyl. Examples of suitable amino protecting groups include benzyl, alpha-methylbenzyl, diphenylmethyl, triphenylmethyl, benzyloxycarbonyl, tert-butoxycarbonyl, and acyl groups such as trichloroacetyl or trifluoroacetyl. In some embodiments, the functional moiety is self-cleaving. In other embodiments, the functional moiety is biodegradable. See e.g., U.S. Patent No. 7,087,229.

35 [0146] In some embodiments, oligomers of the invention are functionalized at the 5' end in order to allow covalent attachment of the conjugated moiety to the 5' end of the oligomer. In other embodiments, oligomers of the invention can be functionalized at the 3' end. In still other embodiments, oligomers of the invention can be functionalized along the backbone or on the heterocyclic base moiety. In yet other embodiments, oligomers of the invention can be functionalized at more than one position independently selected from the 5' end, the 3' end, the backbone and the base.

40 [0147] In some embodiments, activated oligomers of the invention are synthesized by incorporating during the synthesis one or more monomers that is covalently attached to a functional moiety. In other embodiments, activated oligomers of the invention are synthesized with monomers that have not been functionalized, and the oligomer is functionalized upon completion of synthesis. In some embodiments, the oligomers are functionalized with a hindered ester containing an aminoalkyl linker, wherein the alkyl portion has the formula (CH<sub>2</sub>)<sub>w</sub>, wherein w is an integer ranging from 1 to 10, preferably about 6, wherein the alkyl portion of the alkylamino group can be straight chain or branched chain, and wherein the functional group is attached to the oligomer via an ester group (-O-C(O)-(CH<sub>2</sub>)<sub>w</sub>NH).

45 [0148] In other embodiments, the oligomers are functionalized with a hindered ester containing a (CH<sub>2</sub>)<sub>w</sub>-sulfhydryl (SH) linker, wherein w is an integer ranging from 1 to 10, preferably about 6, wherein the alkyl portion of the alkylamino group can be straight chain or branched chain, and wherein the functional group attached to the oligomer via an ester group (-O-C(O)-(CH<sub>2</sub>)<sub>w</sub>SH).

50 [0149] In some embodiments, sulfhydryl-activated oligonucleotides are conjugated with polymer moieties such as polyethylene glycol or peptides (via formation of a disulfide bond).

[0150] Activated oligomers containing hindered esters as described above can be synthesized by any method known in the art, and in particular by methods disclosed in PCT Publication No. WO 2008/034122 and the examples therein.

55 [0151] In still other embodiments, the oligomers of the invention are functionalized by introducing sulfhydryl, amino or hydroxyl groups into the oligomer by means of a functionalizing reagent substantially as described in U.S. Patent Nos. 4,962,029 and 4,914,210, i.e., a substantially linear reagent having a phosphoramidite at one end linked through a hydrophilic spacer chain to the opposing end which comprises a protected or unprotected sulfhydryl, amino or hydroxyl group. Such reagents primarily react with hydroxyl groups of the oligomer. In some embodiments, such activated oli-

omers have a functionalizing reagent coupled to a 5'-hydroxyl group of the oligomer. In other embodiments, the activated oligomers have a functionalizing reagent coupled to a 3'-hydroxyl group. In still other embodiments, the activated oligomers of the invention have a functionalizing reagent coupled to a hydroxyl group on the backbone of the oligomer. In yet further embodiments, the oligomer of the invention is functionalized with more than one of the functionalizing reagents as described in U.S. Patent Nos. 4,962,029 and 4,914,210.

**[0152]** Methods of synthesizing such functionalizing reagents and incorporating them into monomers or oligomers are disclosed in U.S. Patent Nos. 4,962,029 and 4,914,210.

**[0153]** In some embodiments, the 5'-terminus of a solid-phase bound oligomer is functionalized with a dienyl phosphoramidite derivative, followed by conjugation of the deprotected oligomer with, e.g., an amino acid or peptide via a Diels-Alder cycloaddition reaction.

**[0154]** In various embodiments, the incorporation of monomers containing 2'-sugar modifications, such as a 2'-carbamate substituted sugar or a 2'-(O-pentyl-N-phthalimido)-deoxyribose sugar into the oligomer facilitates covalent attachment of conjugated moieties to the sugars of the oligomer. In other embodiments, an oligomer with an amino-containing linker at the 2'-position of one or more monomers is prepared using a reagent such as, for example, 5'-dimethoxytrityl-2'-O-(e-phthalimidylaminopentyl)-2'-deoxyadenosine-3'-- N,N-diisopropyl-cyanoethoxy phosphoramidite. See, e.g., Manoharan, et al., *Tetrahedron Letters*, 1991, 34, 7171.

**[0155]** In still further embodiments, the oligomers of the invention may have amine-containing functional moieties on the nucleotide, including on the N6 purine amino groups, on the exocyclic N2 of guanine, or on the N4 or 5 positions of cytosine. In various embodiments, such functionalization may be achieved by using a commercial reagent that is already functionalized in the oligomer synthesis.

**[0156]** Some functional moieties are commercially available, for example, heterobifunctional and homobifunctional linking moieties are available from the Pierce Co. (Rockford, Ill.). Other commercially available linking groups are 5'-Amino-Modifier C6 and 3'-Amino-Modifier reagents, both available from Glen Research Corporation (Sterling, Va.). 5'-Amino-Modifier C6 is also available from ABI (Applied Biosystems Inc., Foster City, Calif.) as Aminolink-2, and 3'-Amino-Modifier is also available from Clontech Laboratories Inc. (Palo Alto, Calif.).

#### ***Therapy and pharmaceutical compositions - formulation and administration***

**[0157]** As explained initially, the oligonucleotides of the invention will constitute suitable drugs with improved properties. The design of a potent and safe drug requires the fine-tuning of various parameters such as affinity/specificity, stability in biological fluids, cellular uptake, mode of action, pharmacokinetic properties and toxicity.

**[0158]** Accordingly, in a further aspect the present invention relates to a pharmaceutical composition comprising an oligonucleotide according to the invention and a pharmaceutically acceptable diluent, carrier or adjuvant. Preferably said carrier is saline or buffered saline.

**[0159]** In a still further aspect the present invention relates to an oligonucleotide according to the present invention for use as a medicament.

**[0160]** As will be understood, dosing is dependent on severity and responsiveness of the disease state to be treated, and the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Optimum dosages may vary depending on the relative potency of individual oligonucleotides. Generally it can be estimated based on EC50s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 µg to 1 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 10 years or by continuous infusion for hours up to several months. The repetition rates for dosing can be estimated based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state.

**[0161]** As indicated above, the invention also relates to a pharmaceutical composition, which comprises at least one oligonucleotide of the invention as an active ingredient. It should be understood that the pharmaceutical composition according to the invention optionally comprises a pharmaceutical carrier, and that the pharmaceutical composition optionally comprises further compounds, such as chemotherapeutic compounds, anti-inflammatory compounds, antiviral compounds and/or immuno-modulating compounds.

**[0162]** The oligonucleotides of the invention can be used "as is" or in form of a variety of pharmaceutically acceptable salts. As used herein, the term "pharmaceutically acceptable salts" refers to salts that retain the desired biological activity of the herein-identified oligonucleotides and exhibit minimal undesired toxicological effects. Non-limiting examples of such salts can be formed with organic amino acid and base addition salts formed with metal cations such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium, sodium, potassium, and the like, or with a cation formed from ammonia, *N,N*-dibenzylethylene-diamine, D-glucosamine, tetraethylammonium, or ethylenediamine.

**[0163]** In one embodiment of the invention, the oligonucleotide may be in the form of a pro-drug. Oligonucleotides are

by virtue negatively charged ions. Due to the lipophilic nature of cell membranes the cellular uptake of oligonucleotides are reduced compared to neutral or lipophilic equivalents. This polarity "hindrance" can be avoided by using the pro-drug approach (see e.g. Crooke, R. M. (1998) in Crooke, S. T. Antisense research and Application. Springer-Verlag, Berlin, Germany, vol. 131, pp. 103-140).

**[0164]** Pharmaceutically acceptable binding agents and adjuvants may comprise part of the formulated drug.

**[0165]** Examples of delivery methods for delivery of the therapeutic agents described herein, as well as details of pharmaceutical formulations, salts, may be well described elsewhere for example in US provisional application 60/838,710 and 60/788,995, and Danish applications, PA 2006 00615.

**[0166]** Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids. Delivery of drug to tumour tissue may be enhanced by carrier-mediated delivery including, but not limited to, cationic liposomes, cyclodextrins, porphyrin derivatives, branched chain dendrimers, polyethylenimine polymers, nanoparticles and microspheres (Dass CR. J Pharm Pharmacol 2002; 54(1):3-27). The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product. The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels and suppositories. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers. The compounds of the invention may also be conjugated to active drug substances, for example, aspirin, ibuprofen, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

**[0167]** In another embodiment, compositions of the invention may contain one or more oligonucleotide compounds, targeted to a first microRNA and one or more additional oligonucleotide compounds targeted to a second microRNA target. Two or more combined compounds may be used together or sequentially.

**[0168]** The compounds disclosed herein are useful for a number of therapeutic applications as indicated above. In general, therapeutic methods of the invention include administration of a therapeutically effective amount of an oligonucleotide to a mammal, particularly a human. In a certain embodiment, the present invention provides pharmaceutical compositions containing (a) one or more compounds of the invention, and (b) one or more chemotherapeutic agents. When used with the compounds of the invention, such chemotherapeutic agents may be used individually, sequentially, or in combination with one or more other such chemotherapeutic agents or in combination with radiotherapy. All chemotherapeutic agents known to a person skilled in the art are here incorporated as combination treatments with compound according to the invention. Other active agents, such as anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, antiviral drugs, and immuno-modulating drugs may also be combined in compositions of the invention. Two or more combined compounds may be used together or sequentially.

**[0169]** Examples of therapeutic indications which may be treated by the pharmaceutical compositions of the invention:

microRNA	Possible medical indications
miR-1	Cardiac arrhythmia
miR-21	Glioblastoma, breast cancer, hepatocellular carcinoma, colorectal cancer, sensitization of gliomas to cytotoxic drugs, cardiac hypertrophy
miR-21, miR-200b and miR-141	Response to chemotherapy and regulation of cholangiocarcinoma growth
miR-122	hypercholesterolemia, hepatitis C infection, hemochromatosis
miR-19b	lymphoma and other tumour types
miR-26a	Osteoblast differentiation of human stem cells
miR-155	lymphoma, pancreatic tumor development, breast and lung cancer
miR-203	Psoriasis
miR-375	diabetes, metabolic disorders, glucose-induced insulin secretion from pancreatic endocrine cells

(continued)

microRNA	Possible medical indications
miR-181	myoblast differentiation, auto immune disorders
miR-10b	Breast cancer cell invasion and metastasis
miR-125b-1	Breast, lung, ovarian and cervical cancer
miR-221 and 222	Prostate carcinoma, human thyroid papillary car, human hepatocellular carcinoma
miRNA-372 and - 373	testicular germ cell tumors.
miR-142	B-cell leukemia
miR-17 - 19b cluster	B-cell lymphomas, lung cancer, hepatocellular carcinoma

**[0170]** Tumor suppressor gene tropomyosin 1 (TPM1) mRNA has been indicated as a target of miR-21. Myotrophin (mtpn) mRNA has been indicated as a target of miR 375.

**[0171]** In an even further aspect, the present invention relates to the use of an oligonucleotide according to the invention for the manufacture of a medicament for the treatment of a disease selected from the group consisting of: atherosclerosis, hypercholesterolemia and hyperlipidemia; cancer, glioblastoma, breast cancer, lymphoma, lung cancer; diabetes, metabolic disorders; myoblast differentiation; immune disorders.

**[0172]** The invention further refers to oligonucleotides according to the invention for the use in the treatment of from a disease selected from the group consisting of: atherosclerosis, hypercholesterolemia and hyperlipidemia; cancer, glioblastoma, breast cancer, lymphoma, lung cancer; diabetes, metabolic disorders; myoblast differentiation; immune disorders.

**[0173]** The disclosure provides a method of treating a subject suffering from a disease or condition selected from from the group consisting of: atherosclerosis, hypercholesterolemia and hyperlipidemia; cancer, glioblastoma, breast cancer, lymphoma, lung cancer; diabetes, metabolic disorders; myoblast differentiation; immune disorders, the method comprising the step of administering an oligonucleotide or pharmaceutical composition of the invention to the subject in need thereof.

**[0174]** The invention further provides for a kit comprising a pharmaceutical composition according to the invention, and a second independent active ingredient that is an inhibitor of the VLDL assembly pathway, such as an ApoB inhibitor, or an MTP inhibitor.

### **Cancer**

**[0175]** In an even further aspect, the present invention relates to the use of an oligonucleotide according to the invention for the manufacture of a medicament for the treatment of cancer. In another aspect, the present invention concerns a method for treatment of, or prophylaxis against, cancer, said method comprising administering an oligonucleotide of the invention or a pharmaceutical composition of the invention to a patient in need thereof.

**[0176]** Such cancers may include lymphoreticular neoplasia, lymphoblastic leukemia, brain tumors, gastric tumors, plasmacytomas, multiple myeloma, leukemia, connective tissue tumors, lymphomas, and solid tumors.

**[0177]** In the use of a compound of the invention for the manufacture of a medicament for the treatment of cancer, said cancer may suitably be in the form of a solid tumor. Analogously, in the method for treating cancer disclosed herein said cancer may suitably be in the form of a solid tumor.

**[0178]** Furthermore, said cancer is also suitably a carcinoma. The carcinoma is typically selected from the group consisting of malignant melanoma, basal cell carcinoma, ovarian carcinoma, breast carcinoma, non-small cell lung cancer, renal cell carcinoma, bladder carcinoma, recurrent superficial bladder cancer, stomach carcinoma, prostatic carcinoma, pancreatic carcinoma, lung carcinoma, cervical carcinoma, cervical dysplasia, laryngeal papillomatosis, colon carcinoma, colorectal carcinoma and carcinoid tumors. More typically, said carcinoma is selected from the group consisting of malignant melanoma, non-small cell lung cancer, breast carcinoma, colon carcinoma and renal cell carcinoma. The malignant melanoma is typically selected from the group consisting of superficial spreading melanoma, nodular melanoma, lentigo maligna melanoma, acral melanoma, amelanotic melanoma and desmoplastic melanoma.

**[0179]** Alternatively, the cancer may suitably be a sarcoma. The sarcoma is typically in the form selected from the group consisting of osteosarcoma, Ewing's sarcoma, chondrosarcoma, malignant fibrous histiocytoma, fibrosarcoma and Kaposi's sarcoma.

**[0180]** Alternatively, the cancer may suitably be a glioma.

**[0181]** A further embodiment is directed to the use of an oligonucleotide according to the invention for the manufacture

of a medicament for the treatment of cancer, wherein said medicament further comprises a chemotherapeutic agent selected from the group consisting of adrenocorticosteroids, such as prednisone, dexamethasone or decadron; altretamine (hexalen, hexamethylmelamine (HMM)); amifostine (ethyol); aminoglutethimide (cytadren); amsacrine (M-AMSA); anastrozole (arimidex); androgens, such as testosterone; asparaginase (elspar); bacillus calmette-gurin; bicalutamide (casodex); bleomycin (blenoxane); busulfan (myleran); carboplatin (paraplatin); carmustine (BCNU, BiCNU); chlorambucil (leukeran); chlorodeoxyadenosine (2-CDA, cladribine, leustatin); cisplatin (platinol); cytosine arabinoside (cytarabine); dacarbazine (DTIC); dactinomycin (actinomycin-D, cosmegen); daunorubicin (cerubidine); docetaxel (taxotere); doxorubicin (adriomycin); epirubicin; estramustine (emcyt); estrogens, such as diethylstilbestrol (DES); etoposide (VP-16, VePesid, etopophos); fludarabine (fludara); flutamide (eulexin); 5-FUDR (floxuridine); 5-fluorouracil (5-FU); gemcitabine (gemzar); goserelin (zodalex); herceptin (trastuzumab); hydroxyurea (hydrea); idarubicin (idamycin); ifosfamide; IL-2 (proleukin, aldesleukin); interferon alpha (intron A, roferon A); irinotecan (campptosar); leuprolide (lupron); levamisole (ergamisole); lomustine (CCNU); mechlorathamine (mustargen, nitrogen mustard); melphalan (alkeran); mercaptopurine (purinethol, 6-MP); methotrexate (mexate); mitomycin-C (mutamucin); mitoxantrone (novantrone); octreotide (sandostatin); pentostatin (2-deoxycoformycin, nipent); plicamycin (mithramycin, mithracin); prorocarbazine (matulane); streptozocin; tamoxifen (nolvadex); taxol (paclitaxel); teniposide (vumon, VM-26); thiotepa; topotecan (hycamtin); tretinoin (vesanoid, all-trans retinoic acid); vinblastine (valban); vincristine (oncovin) and vinorelbine (navelbine). Suitably, the further chemotherapeutic agent is selected from taxanes such as Taxol, Paclitaxel or Docetaxel.

**[0182]** Similarly, the invention is further directed to the use of an oligonucleotide according to the invention for the manufacture of a medicament for the treatment of cancer, wherein said treatment further comprises the administration of a further chemotherapeutic agent selected from the group consisting of adrenocorticosteroids, such as prednisone, dexamethasone or decadron; altretamine (hexalen, hexamethylmelamine (HMM)); amifostine (ethyol); aminoglutethimide (cytadren); amsacrine (M-AMSA); anastrozole (arimidex); androgens, such as testosterone; asparaginase (elspar); bacillus calmette-gurin; bicalutamide (casodex); bleomycin (blenoxane); busulfan (myleran); carboplatin (paraplatin); carmustine (BCNU, BiCNU); chlorambucil (leukeran); chlorodeoxyadenosine (2-CDA, cladribine, leustatin); cisplatin (platinol); cytosine arabinoside (cytarabine); dacarbazine (DTIC); dactinomycin (actinomycin-D, cosmegen); daunorubicin (cerubidine); docetaxel (taxotere); doxorubicin (adriomycin); epirubicin; estramustine (emcyt); estrogens, such as diethylstilbestrol (DES); etoposide (VP-16, VePesid, etopophos); fludarabine (fludara); flutamide (eulexin); 5-FUDR (floxuridine); 5-fluorouracil (5-FU); gemcitabine (gemzar); goserelin (zodalex); herceptin (trastuzumab); hydroxyurea (hydrea); idarubicin (idamycin); ifosfamide; IL-2 (proleukin, aldesleukin); interferon alpha (intron A, roferon A); irinotecan (campptosar); leuprolide (lupron); levamisole (ergamisole); lomustine (CCNU); mechlorathamine (mustargen, nitrogen mustard); melphalan (alkeran); mercaptopurine (purinethol, 6-MP); methotrexate (mexate); mitomycin-C (mutamucin); mitoxantrone (novantrone); octreotide (sandostatin); pentostatin (2-deoxycoformycin, nipent); plicamycin (mithramycin, mithracin); prorocarbazine (matulane); streptozocin; tamoxifen (nolvadex); taxol (paclitaxel); teniposide (vumon, VM-26); thiotepa; topotecan (hycamtin); tretinoin (vesanoid, all-trans retinoic acid); vinblastine (valban); vincristine (oncovin) and vinorelbine (navelbine). Suitably, said treatment further comprises the administration of a further chemotherapeutic agent selected from taxanes, such as Taxol, Paclitaxel or Docetaxel.

**[0183]** Alternatively stated, the disclosure provides a method for treating cancer, said method comprising administering an oligonucleotide of the invention or a pharmaceutical composition according to the invention to a patient in need thereof and further comprising the administration of a further chemotherapeutic agent. Said further administration may be such that the further chemotherapeutic agent is conjugated to the compound of the invention, is present in the pharmaceutical composition, or is administered in a separate formulation.

### ***Infectious diseases***

**[0184]** It is contemplated that the compounds of the invention may be broadly applicable to a broad range of infectious diseases, such as diphtheria, tetanus, pertussis, polio, hepatitis B, hepatitis C, hemophilus influenza, measles, mumps, and rubella.

**[0185]** Hsa-miR122 is indicated in hepatitis C infection and as such oligonucleotides according to the invention which target miR-122 may be used to treat Hepatitis C infection.

**[0186]** Accordingly, in yet another aspect the present invention relates the use of an oligonucleotide according to the invention for the manufacture of a medicament for the treatment of an infectious disease, as well as to a method for treating an infectious disease, said method comprising administering an oligonucleotide according to the invention or a pharmaceutical composition according to the invention to a patient in need thereof.

**[0187]** In a preferred embodiment, the invention provides for a combination treatment providing an anti miR-122 oligomer in combination with an inhibitor of VLDL assembly, such as an inhibitor of apoB, or of MTP.

**Inflammatory diseases**

**[0188]** The inflammatory response is an essential mechanism of defense of the organism against the attack of infectious agents, and it is also implicated in the pathogenesis of many acute and chronic diseases, including autoimmune disorders. In spite of being needed to fight pathogens, the effects of an inflammatory burst can be devastating. It is therefore often necessary to restrict the symptomatology of inflammation with the use of anti-inflammatory drugs. Inflammation is a complex process normally triggered by tissue injury that includes activation of a large array of enzymes, the increase in vascular permeability and extravasation of blood fluids, cell migration and release of chemical mediators, all aimed to both destroy and repair the injured tissue.

**[0189]** In yet another aspect, the present invention relates to the use of an oligonucleotide according to the invention for the manufacture of a medicament for the treatment of an inflammatory disease, as well as to a method for treating an inflammatory disease, said method comprising administering an oligonucleotide according to the invention or a pharmaceutical composition according to the invention to a patient in need thereof.

**[0190]** In one preferred embodiment of the invention, the inflammatory disease is a rheumatic disease and/or a connective tissue diseases, such as rheumatoid arthritis, systemic lupus erythematosus (SLE) or Lupus, scleroderma, polymyositis, inflammatory bowel disease, dermatomyositis, ulcerative colitis, Crohn's disease, vasculitis, psoriatic arthritis, exfoliative psoriatic dermatitis, pemphigus vulgaris and Sjorgren's syndrome, in particular inflammatory bowel disease and Crohn's disease.

**[0191]** Alternatively, the inflammatory disease may be a non-rheumatic inflammation, like bursitis, synovitis, capsulitis, tendinitis and/or other inflammatory lesions of traumatic and/or sportive origin.

**Metabolic diseases**

**[0192]** A metabolic disease is a disorder caused by the accumulation of chemicals produced naturally in the body. These diseases are usually serious, some even life threatening. Others may slow physical development or cause mental retardation. Most infants with these disorders, at first, show no obvious signs of disease. Proper screening at birth can often discover these problems. With early diagnosis and treatment, metabolic diseases can often be managed effectively.

**[0193]** In yet another aspect, the present invention relates to the use of an oligonucleotide according to the invention or a conjugate thereof for the manufacture of a medicament for the treatment of a metabolic disease, as well as to a method for treating a metabolic disease, said method comprising administering an oligonucleotide according to the invention or a conjugate thereof, or a pharmaceutical composition according to the invention to a patient in need thereof.

**[0194]** In one preferred embodiment of the invention, the metabolic disease is selected from the group consisting of Amyloidosis, Biotinidase, OMIM (Online Mendelian Inheritance in Man), Crigler Najjar Syndrome, Diabetes, Fabry Support & Information Group, Fatty acid Oxidation Disorders, Galactosemia, Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency, Glutaric aciduria, International Organization of Glutaric Acidemia, Glutaric Acidemia Type I, Glutaric Acidemia, Type II, Glutaric Acidemia Type I, Glutaric Acidemia Type-II, F-HYPDRR - Familial Hypophosphatemia, Vitamin D Resistant Rickets, Krabbe Disease, Long chain 3 hydroxyacyl CoA dehydrogenase deficiency (LCHAD), Mannosidosis Group, Maple Syrup Urine Disease, Mitochondrial disorders, Mucopolysaccharidosis Syndromes: Niemann Pick, Organic acidemias, PKU, Pompe disease, Porphyria, Metabolic Syndrome, Hyperlipidemia and inherited lipid disorders, Trimethylaminuria: the fish malodor syndrome, and Urea cycle disorders.

**Liver disorders**

**[0195]** In yet another aspect, the present invention relates to the use of an oligonucleotide according to the invention or a conjugate thereof for the manufacture of a medicament for the treatment of a liver disorder, as well as to a method for treating a liver disorder, said method comprising administering an oligonucleotide according to the invention or a conjugate thereof, or a pharmaceutical composition according to the invention to a patient in need thereof.

**[0196]** In one preferred embodiment of the invention, the liver disorder is selected from the group consisting of Biliary Atresia, Alagille Syndrome, Alpha-1 Antitrypsin, Tyrosinemia, Neonatal Hepatitis, and Wilson Disease.

**Other uses**

**[0197]** The oligonucleotides of the present invention can be utilized for as research reagents for diagnostics, therapeutics and prophylaxis. In research, the oligonucleotide may be used to specifically inhibit the synthesis of target genes in cells and experimental animals thereby facilitating functional analysis of the target or an appraisal of its usefulness as a target for therapeutic intervention. In diagnostics the oligonucleotides may be used to detect and quantitate target expression in cell and tissues by Northern blotting, *in-situ* hybridisation or similar techniques. For therapeutics, an animal or a human, suspected of having a disease or disorder, which can be treated by modulating the expression of target is

treated by administering the oligonucleotide compounds in accordance with this invention. Further provided are methods of treating an animal particular mouse and rat and treating a human, suspected of having or being prone to a disease or condition, associated with expression of target by administering a therapeutically or prophylactically effective amount of one or more of the oligonucleotide compounds or compositions of the invention.

5

#### ***Therapeutic use of oligonucleotides targeting miR-122a***

**[0198]** We have demonstrated that a LNA-antimiR, targeting miR-122a reduces plasma cholesterol levels. Therefore, another aspect of the invention is use of the above described oligonucleotides targeting miR-122a as medicine.

10 **[0199]** Still another aspect of the invention is use of the above described oligonucleotides targeting miR-122a for the preparation of a medicament for treatment of increased plasma cholesterol levels (or hypercholesterolemia and related disorders). The skilled man will appreciate that increased plasma cholesterol levels is undesirable as it increases the risk of various conditions, e.g. atherosclerosis.

15 **[0200]** Still another aspect of the invention is use of the above described oligonucleotides targeting miR-122a for upregulating the mRNA levels of Nrdg3, Aldo A, Bckdk or CD320.

#### **EMBODIMENTS**

20 **[0201]** The following embodiments of the present invention may be used in combination with the other embodiments described herein.

25 1. A pharmaceutical composition comprising an oligomer of 7 - 12 nucleotides in length, wherein said oligomer comprises a total of 7 - 12 nucleotides, such as 7, 8, 9, 10, 11 or 12 nucleotide units, where all nucleotide units are LNA units, and wherein at least 75% of the internucleoside linkages present between the nucleobase units of the contiguous nucleobase sequence are phosphorothioate internucleoside linkages, wherein the oligomer is for use in reducing the effective amount of a microRNA target in a cell; and a pharmaceutically acceptable diluent, carrier, salt or adjuvent.

30 2. The pharmaceutical composition according to embodiment 1, wherein the contiguous nucleotide sequence is complementary to a corresponding region of a mammalian, human or viral microRNA (miRNA) sequence.

3. The pharmaceutical composition according to embodiment 2, wherein the contiguous nucleotide sequence is complementary to a corresponding region of a miRNA sequence selected from the group of miRNAs listed in any one of tables 3, 4 or 5.

4. The pharmaceutical composition according to embodiment 2 or 3, wherein the contiguous nucleotide sequence consists of or comprises a sequence which is complementary to the seed sequence of said microRNA.

35 5. The pharmaceutical composition according to any one of embodiments 2 - 4, wherein the contiguous nucleotide sequence consists of or comprises a sequence selected from any one of the sequences listed in table 3 or 4.

6. The pharmaceutical composition according to embodiment 4 or 5, wherein the 3' nucleobase of the seedmer forms the 3' most nucleobase of the contiguous nucleotide sequence, wherein the contiguous nucleotide sequence may, optionally, comprise one or two further 5' nucleobases.

40 7. The pharmaceutical composition according to any one of embodiments 1-6, wherein said contiguous nucleotide sequence does not comprise a nucleotide which corresponds to the first nucleotide present in the micro RNA sequence counted from the 5' end.

8. The pharmaceutical composition according to any one of embodiments 1-7, wherein the contiguous nucleotide sequence is complementary to a corresponding nucleotide sequence present in a miRNA selected from those shown in table 3 or 4 or 5.

9. The pharmaceutical composition according to embodiment 8, wherein said miRNA is selected from the group consisting of miR-1, miR-10b, miR-17-3p, miR-18, miR-19a, miR-19b, miR-20, miR-21, miR-34a, miR-93, miR-106a, miR-106b, miR-122, miR-133, miR-134, miR-138, miR-155, miR-192, miR-194, miR-221, miR-222, and miR-375.

50 14. The pharmaceutical composition according to any one of embodiments 1 - 13, wherein the contiguous nucleotide sequence consists of 7, 8, 9 or 10 contiguous, LNA nucleobase units.

15. The pharmaceutical composition according to any one of embodiments 1-14, wherein the oligomer consist of 7, 8, 9 or 10 contiguous LNA.

18. The pharmaceutical composition according to any one of embodiments 1 - 9 wherein 80% or 85% or 90% or 95% or all of the internucleoside linkages present between the nucleobase units of the contiguous nucleotide sequence are phosphorothioate internucleoside linkages.

55 19. The pharmaceutical composition according to any one of embodiments 1 - 18, wherein said oligomer is conjugated with one or more non-nucleobase compounds.

20. The pharmaceutical composition according to any one of embodiments 1 - 19, wherein the contiguous nucleotide

sequence is complementary to the corresponding sequence of at least two miRNA sequences such as 2, 3, 4, 5, 6, 7, 8, 9, or 10 miRNA sequences.

21. The pharmaceutical composition according to any one of embodiments 1 - 20, wherein the contiguous nucleotide sequence consists or comprises of a sequence which is complementary to the sequence of at least two miRNA seed region sequences such as 2, 3, 4, 5, 6, 7, 8, 9, or 10 miRNA seed region sequences.

22. The pharmaceutical composition according to any one of embodiments 20 or 21, wherein the contiguous nucleotide sequence is complementary to the corresponding region of both miR-221 and miR-222.

23. The pharmaceutical composition according to embodiment 22, wherein the contiguous nucleotide sequence consists or comprises of a sequence that is complementary to 5'GCUACAU3'.

24. The pharmaceutical composition according to any one of embodiments 1 - 23, wherein the oligomer is constituted as a prodrug.

25. The pharmaceutical composition according to any one of embodiments 1 - 24, wherein the contiguous nucleotide sequence is complementary to a corresponding region of has-miR-122.

26. The pharmaceutical composition according to embodiment 25, for use in the treatment of a medical disorder or disease selected from the group consisting of: hepatitis C virus infection and hypercholesterolemia and related disorders.

27. The pharmaceutical composition according to embodiment 25 or 26, wherein the composition further comprises a second independent active ingredient that is an inhibitor of the VLDL assembly pathway, such as an ApoB inhibitor, or an MTP inhibitor.

28. A kit comprising a pharmaceutical composition according to embodiment 25 or 26, and a second independent active ingredient that is an inhibitor of the VLDL assembly pathway, such as an ApoB inhibitor, or an MTP inhibitor.

29. The pharmaceutical composition of the invention for use in the treatment of a disease or medical disorder associated with the presence or overexpression of a microRNA.

30. An oligomer, as defined according to anyone of embodiments 1 - 25.

31. A conjugate comprising the oligomer according to embodiment 30, and at least one non-nucleobase compounds.

32. The use of an oligomer or a conjugate as defined in any one of embodiments 30 - 31, for the manufacture of a medicament for the treatment of a disease or medical disorder associated with the presence or over-expression of the microRNA.

33. An *in vitro* a method for reducing the amount, or effective amount, of a miRNA in a cell, comprising administering an oligomer, a conjugate or a pharmaceutical composition, according to any one of the preceding embodiments to the cell which is expressing said miRNA so as to reduce the amount, or effective amount of the miRNA in the cell.

34. An *in vitro* a method for de-repression of a mRNA whose expression is repressed by a miRNA in a cell comprising administering an oligomer, a conjugate or a pharmaceutical composition, according to any one of the preceding embodiments to the cell to the cell which expressed both said mRNA and said miRNA, in order to de-repress the expression of the mRNA.

## EXAMPLES

**[0202]** LNA Monomer and oligonucleotide synthesis were performed using the methodology referred to in Examples 1 and 2 of WO2007/112754. The stability of LNA oligonucleotides in human or rat plasma is performed using the methodology referred to in Example 4 of WO2007/112754. The treatment of *in vitro* cells with LNA anti-miR antisense oligonucleotide (targeting miR-122) is performed using the methodology referred to in Example 6 of WO2007/112754. The analysis of Oligonucleotide Inhibition of miR expression by microRNA specific quantitative PCR in both an *in vitro* and *in vivo* model is performed using the methodology referred to in Example 7 of WO2007/112754. The assessment of LNA antimir knock-down specificity using miRNA microarray expression profiling is performed using the methodology referred to in Example 8 of WO2007/112754. The detection of microRNAs by *in situ* hybridization is performed using the methodology referred to in Example 9 of WO2007/112754. The isolation and analysis of mRNA expression (total RNA isolation and cDNA synthesis for mRNA analysis) in both an *in vitro* and *in vivo* model is performed using the methodology referred to in Example 10 of WO2007/112754. *In vivo* Experiments using Oligomers of the invention targeting microRNA-122. and subsequent analysis are performed using the methods disclosed in Examples 11 - 27 of WO2007/112754.

### Example 1: Design of the LNA anti-miR oligonucleotides and melting temperatures

#### [0203]



**Table 2** - Oligomers used in the examples and figures. The SEQ# is an identifier used throughout the examples and figures - the SEQ ID NO which is used in the sequence listing is also provided.

Example/Figure SEQ#	SEQ ID NO	Compound Sequence	Comment
5 #3204	1	TcAGtCTGaTaAgCT	
#3205	2	GATAAGCT	
#3206	3	TcAcAATtaGCAItA	
10 #3207	4	TAGCATTa	
#4	5	CcAttGTcaCaCtCC	
#3208	6	CACACTCC	
#3209	7	TAAGCT	
15 #3210	8	ATAAGCT	
#3211	9	TGATAAGCT	
#3212	10	CTGATAAGCT	
20 #3213	11	GTCTGATAAGCT	
#2114	12	CAGTCTGATAAGCT	
#3215	13	TCTGATAA	
#3216	14	ATCAGTCT	
25 #3217	15	TCAACATC	
#3218/#3230	16	<u>GGTAAACT</u>	Underline = mismatch
#3219	17	<u>CGTAATGA</u>	Underline = mismatch
30 #3220	18	TCAgctgataaGCTa	5' fluorescent label (FAM)
#3221	19	AGCACTTT	
#3222	20	ATTTGCAC	
35 #3223	21	AgCagACaaTgTaGC	5' fluorescent label (FAM)
#3224	22	GtAgcCAgaTgTaGC	5' fluorescent label (FAM)
40 #3225	23	ATGTAGC	
#3226	24	ACaAcCTacTaCcTC	
#3227	25	ACTACCTC	
#3228	26	CaCtgTCagCaCtTT	
45 #3229	27	TgCatAGatTtGcAC	
#3231	28	GTAGACT	
#3232	29	TACCTC	
50 #3233	30	CTACCTC	
#3234	31	TNCTACCTC	N = universal base.
#3235	32	TNCTACCTC	N = universal base.
#3236	33	GCaAcCTacTaCcTC	
55 #3237	34	ACaAcCTccTaCcTC	
#3238	35	ACaAaCTacTaCcTC	

(continued)

Example/Figure SEQ#	SEQ ID NO	Compound Sequence	Comment
#3239	36	CTACCTC	
#3240	37	CTAACTC	
#3241	38	TTAGCATTA	
#3242	39	CGATTAGCATTA	
#3243	977	CACGATTAGCATTA	
#3244	978	GCATTA	
#3245	979	AGCATTA	
#3246	980	ATTAGCATTA	
Capital and lower case letters denote LNA and DNA, respectively. LNA cytosines are preferably methyl cytosine/5'methyl-cytosine* All internucleoside linkages are preferably phosphorothioate* All LNA may, for example, be beta-D-oxy LNA* *Used in the specific examples.			

**Example 2: In vitro model: Cell culture**

**[0204]** The effect of LNA oligonucleotides on target nucleic acid expression (amount) can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. Target can be expressed endogenously or by transient or stable transfection of a nucleic acid encoding said nucleic acid.

**[0205]** The expression level of target nucleic acid can be routinely determined using, for example, Northern blot analysis (including microRNA northern), Quantitative PCR (including microRNA qPCR), Ribonuclease protection assays. The following cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen.

**[0206]** Cells were cultured in the appropriate medium as described below and maintained at 37°C at 95-98% humidity and 5% CO<sub>2</sub>. Cells were routinely passaged 2-3 times weekly.

15PC3: The human prostate cancer cell line 15PC3 was kindly donated by Dr. F. Baas, Neurozintuigen Laboratory, AMC, The Netherlands and was cultured in DMEM (Sigma) + 10% fetal bovine serum (FBS) + Glutamax I + gentamicin.

PC3: The human prostate cancer cell line PC3 was purchased from ATCC and was cultured in F12 Coon's with glutamine (Gibco) + 10% FBS + gentamicin.

518A2: The human melanoma cancer cell line 518A2 was kindly donated by Dr. B. Jansen, Section of experimental Oncology, Molecular Pharmacology, Department of Clinical Pharmacology, University of Vienna and was cultured in DMEM (Sigma) + 10% fetal bovine serum (FBS) + Glutamax I + gentamicin.

HeLa: The cervical carcinoma cell line HeLa was cultured in MEM (Sigma) containing 10% fetal bovine serum gentamicin at 37°C, 95% humidity and 5% CO<sub>2</sub>.

MPC-11: The murine multiple myeloma cell line MPC-11 was purchased from ATCC and maintained in DMEM with 4mM Glutamax+ 10% Horse Serum.

DU-145: The human prostate cancer cell line DU-145 was purchased from ATCC and maintained in RPMI with Glutamax + 10% FBS.

RCC-4 +/- VHL: The human renal cancer cell line RCC4 stably transfected with plasmid expressing VHL or empty plasmid was purchased from ECACC and maintained according to manufacturers instructions.

786-0: The human renal cell carcinoma cell line 786-0 was purchased from ATCC and maintained according to manufacturers instructions

HUVEC: The human umbilical vein endothelial cell line HUVEC was purchased from Camcrex and maintained in EGM-2 medium.

K562: The human chronic myelogenous leukaemia cell line K562 was purchased from ECACC and maintained in RPMI with Glutamax + 10% FBS. U87MG: The human glioblastoma cell line U87MG was purchased from ATCC and maintained according to the manufacturers instructions.

B16: The murine melanoma cell line B16 was purchased from ATCC and maintained according to the manufacturers

instructions.

LNCap: The human prostate cancer cell line LNCap was purchased from ATCC and maintained in RPMI with Glutamax + 10% FBS

Huh-7: Human liver, epithelial like cultivated in Eagles MEM with 10 % FBS, 2mM Glutamax I, 1x non-essential amino acids, Gentamicin 25 µg/ml

L428: (Deutsche Sammlung für Mikroorganismen (DSM, Braunschweig, Germany)): Human B cell lymphoma maintained in RPMI 1640 supplemented with 10% FCS, L-glutamine and antibiotics.

L1236: (Deutsche Sammlung für Mikroorganismen (DSM, Braunschweig, Germany)): Human B cell lymphoma maintained in RPMI 1640 supplemented with 10% FCS, L-glutamine and antibiotics.

### Example 3: Design of a LNA anti-miR library for all human microRNA sequences in miRBase microRNA database.

**[0207]** The miRBase version used was version 12, as reported in Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A., Enright, A.J. 2006. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 34: D140-4, and available via <http://microrna.sanger.ac.uk/sequences/index.shtml>.

**[0208]** **Table 1** shows 7, 8 and 9mer nucleotide sequences comprising the seedmer sequence of micro RNA's according to the miRBase micro RNA database. The seedmer sequence comprises the reverse complement of the microRNA seed region. In some embodiments the oligomer of the invention has a contiguous nucleotide sequence selected from the 7mer, 8mer or 9mer sequences. With respect to the 7mer, 8mer and 9mer sequences, in some embodiments, all the internucleoside linkages are phosphorothioate. The 7mer, 8mer and 9mer nucleotide sequences may consist of sequence of nucleotide analogues as described herein, such as LNA nucleotide analogues. LNA cytosines may be methyl-cytosine (5'-methyl-cytosine). In some embodiments, the LNA is beta-D-oxy-LNA.

**Table 3** provides a list of microRNAs grouped into those which can be targeted by the same seedmer oligomers, such as the 7, 8 or 9mers provided herein (see table 1).

<b>Table 3</b>
hsa-let-7a*, hsa-let-7f-1*
hsa-let-7a, hsa-let-7b, hsa-let-7c, hsa-let-7d, hsa-let-7f, hsa-miR-98, hsa-let-7g, hsa-let-7i
hsa-miR-1, hsa-miR-206
hsa-miR-103, hsa-miR-107
hsa-miR-10a, hsa-miR-10b
hsa-miR-125b, hsa-miR-125a-5p
hsa-miR-129*, hsa-miR-129-3p
hsa-miR-130a, hsa-miR-301a, hsa-miR-130b, hsa-miR-454, hsa-miR-301b
hsa-miR-133a, hsa-miR-133b
hsa-miR-135a, hsa-miR-135b
hsa-miR-141, hsa-miR-200a
hsa-miR-146a, hsa-miR-146b-5p
hsa-miR-152, hsa-miR-148b
hsa-miR-154*, hsa-miR-487a
hsa-miR-15a, hsa-miR-16, hsa-miR-15b, hsa-miR-195, hsa-miR-497
hsa-miR-17, hsa-miR-20a, hsa-miR-93, hsa-miR-106a, hsa-miR-106b, hsa-miR-20b, hsa-miR-526b*
hsa-miR-181a, hsa-miR-181c
hsa-miR-181b, hsa-miR-181d
hsa-miR-18a, hsa-miR-18b
hsa-miR-190, hsa-miR-190b
hsa-miR-192, hsa-miR-215

(continued)

<b>Table 3</b>	
5	hsa-miR-196a, hsa-miR-196b
	hsa-miR-199a-3p, hsa-miR-199b-3p
	hsa-miR-199a-5p, hsa-miR-199b-5p
	hsa-miR-19a*, hsa-miR-19b-1*, hsa-miR-19b-2*
10	hsa-miR-19a, hsa-miR-19b
	hsa-miR-200b, hsa-miR-200c
	hsa-miR-204, hsa-miR-211
15	hsa-miR-208a, hsa-miR-208b
	hsa-miR-212, hsa-miR-132
	hsa-miR-23a*, hsa-miR-23b*
	hsa-miR-23a, hsa-miR-23b, hsa-miR-130a*
20	hsa-miR-24-1*, hsa-miR-24-2*
	hsa-miR-25, hsa-miR-92a, hsa-miR-367, hsa-miR-92b
	hsa-miR-26a, hsa-miR-26b
25	hsa-miR-26a-1*, hsa-miR-26a-2*
	hsa-miR-27a, hsa-miR-27b
	hsa-miR-29a, hsa-miR-29b, hsa-miR-29c
30	hsa-miR-302a, hsa-miR-302b, hsa-miR-302c, hsa-miR-302d, hsa-miR-373, hsa-miR-520e, hsa-miR-520a-3p, hsa-miR-520b, hsa-miR-520c-3p, hsa-miR-520d-3p
	hsa-miR-302b*, hsa-miR-302d*
	hsa-miR-30a*, hsa-miR-30d*, hsa-miR-30e*
35	hsa-miR-30a, hsa-miR-30c, hsa-miR-30d, hsa-miR-30b, hsa-miR-30e
	hsa-miR-330-5p, hsa-miR-326
	hsa-miR-34a, hsa-miR-34c-5p, hsa-miR-449a, hsa-miR-449b
	hsa-miR-362-3p, hsa-miR-329
40	hsa-miR-374a, hsa-miR-374b
	hsa-miR-376a, hsa-miR-376b
	hsa-miR-378, hsa-miR-422a
45	hsa-miR-379*, hsa-miR-411*
	hsa-miR-381, hsa-miR-300
	hsa-miR-509-5p, hsa-miR-509-3-5p
	hsa-miR-515-5p, hsa-miR-519e*
50	hsa-miR-516b*, hsa-miR-516a-3p
	hsa-miR-517a, hsa-miR-517c
	hsa-miR-518a-5p, hsa-miR-527
55	hsa-miR-518f, hsa-miR-518b, hsa-miR-518c, hsa-miR-518a-3p, hsa-miR-518d-3p
	hsa-miR-519c-3p, hsa-miR-519b-3p, hsa-miR-519a
	hsa-miR-519c-5p, hsa-miR-519b-5p, hsa-miR-523*, hsa-miR-518f*, hsa-miR-526a, hsa-miR-520c

(continued)

Table 3	
5p, hsa-miR-518e*, hsa-miR-518d-5p, hsa-miR-522*, hsa-miR-519a*	
hsa-miR-519e, hsa-miR-33b*	
hsa-miR-520a-5p, hsa-miR-525-5p	
hsa-miR-520g, hsa-miR-520h	
hsa-miR-524-5p, hsa-miR-520d-5p	
hsa-miR-525-3p, hsa-miR-524-3p	
hsa-miR-548b-5p, hsa-miR-548a-5p, hsa-miR-548c-5p, hsa-miR-548d-5p	
hsa-miR-7-1*, hsa-miR-7-2*	
hsa-miR-99a, hsa-miR-100, hsa-miR-99b	

We have constructed an 8-mer LNA-antimiR against miR-21, miR-155 and miR-122 (designated here as micromiR) that is fully LNA modified and phosphorothiolated (see figure 1 and Table 6). Our results from repeated experiments in MCF-7, HeLa, Raw and Huh-7 cells using a luciferase sensor plasmid for miR-21, miR-155 and miR-122 demonstrate that the fully LNA-modified short LNA-antimiRs are highly potent in antagonizing microRNAs.

Table 4. LNA_antimiR & MicromiR sequences and predicted $T_m$ s			
SEQ ID #	microRNA	sequence	$T_m$ (°C)
3204	miR-21	T c A G t C T G a T a A g C T	73
3205		G A T A A G C T	33
3206	miR-155	T c A c A A T t a G C A t T A	63
3207		T A G C A T T A	45
4	miR-122	C c A t t G T c a C a C t C C	73
3208		C A C A C T C C	62
Capital letters are LNA units, such as beta-D-oxy LNA. Lower case letters are DNA units. Intemudeoside linkages are preferably phosphorothioate. LNA cytosines are all preferably methylated/5-methyl cytosine.			

The melting temperatures can be assessed towards the mature microRNA sequence, using a synthetic microRNA oligonucleotide (typically consisting of RNA nucleotides with a phosphodiester backbone). Typically measured  $T_m$ s are higher than predicted  $T_m$ s when using LNA oligomers against the RNA target.

#### Example 4: Assessment of miR-21 antagonism by SEQ ID #3205 LNA-antimiR in MCF-7 cells using a luciferase sensor assay.

**[0209]** In order to assess the efficiency of a fully LNA-modified 8-mer LNA-antimiR (SEQ ID #3205) oligonucleotide in targeting and antagonizing miR-21, luciferase sensor constructs were made containing a perfect match target site for the mature miR-21 and as control, a target site with two mutations in the seed (Fig. 6). In order to monitor microRNA-21 inhibition, the breast carcinoma cell line MCF-7 was transfected with the different luciferase constructs together with the miR-21 antagonist SEQ ID #3205 at varying concentrations in comparison with a 15-mer LNA-antimiR SEQ ID #3204 against miR-21. After 24 hours, luciferase activity was measured.

**[0210] Results:** As seen in Figure 2, the new fully LNA-modified 8-mer LNA-antimiR (SEQ ID #3205) shows two-fold higher potency compared to SEQ ID #3204, as shown by de-repression of the luciferase activity. By contrast, the control miR-21 sensor construct with two mismatches in the miR-21 seed did not show any de-repression of the firefly luciferase activity, thereby demonstrating the specificity of the perfect match miR-21 sensor in monitoring miR-21 activity in cells. The de-repression of luciferase activity by the 8-mer LNA-antimiR is clearly dose-dependent, which is not seen with SEQ ID #3204. Moreover, the new 8-mer is also much more potent at lower doses than SEQ ID #3204.

**[0211]** To conclude, the 8-mer LNA-antimiR (SEQ ID #3205) shows significantly improved potency in inhibition of miR-

21 *in vitro* compared to the 15-mer LNA-antimiR SEQ ID #3204 targeting miR-21.

#### Materials and Methods:

5 [0212] Cell line: The breast carcinoma cell line MCF-7 was purchased from ATCC (#HTB-22™). MCF-7 cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

10 [0213] Transfection: 400.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, MCF-7 cells were transfected with 0.8 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector (SDS Promega) together with 1 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

15 [0214] Luciferase assay: The cells were washed with PBS and harvested with cell scraper, after which cells were centrifugated for 5 min at 10.000 rpm. The supernatant was discarded and 50 µl 1 x Passive Lysis Buffer (Promega) was added to the cell pellet, after which cells were put on ice for 30 min. The lysed cells were spinned at 10.000 rpm for 30 min after which 20 µl were transferred to a 96 well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### 20 Example 5: Assessment of miR-21 antagonism by SEQ ID #3205 LNA-antimiR in HeLa cells using a luciferase sensor assay.

[0215] To further assess the efficiency of the fully LNA-modified 8-mer LNA-antimiR SEQ ID #3205 in targeting miR-21, the cervix carcinoma cell line HeLa was also transfected with the previously described miR-21 luciferase sensor constructs alongside SEQ ID #3205 at varying concentrations as described in the above section (Figure 3).

25 [0216] **Results**: The SEQ ID #3205 shows complete de-repression of the miR-21 luciferase sensor construct in HeLa cells already at 5 nM compared to SEQ ID #3204, which did not show complete de-repression until the highest dose (50 nM). In addition, antagonism of miR-21 by the 8-mer SEQ ID #3205 LNA-antimiR is dose-dependent. To demonstrate the specificity of the miR-21 luciferase sensor assay, a mismatched miR-21 target site (2 mismatches in seed) was also transfected into HeLa cells, but did not show any de-repression of the firefly luciferase activity.

30 [0217] *To conclude*, the fully LNA-modified SEQ ID #3205 shows significantly improved potency in inhibition of miR-21 *in vitro*, in both MCF-7 and HeLa cells compared to the 15-mer LNA-antimiR SEQ ID #3204.

#### Materials and Methods:

35 [0218] Cell line: The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

40 [0219] Transfection: 60.000 cells were seeded per well in a 24 well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,7 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

45 [0220] Luciferase assay: The cells were washed with PBS and 100 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24 well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96 well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### Example 6: Assessment of miR-155 antagonism by SEQ ID #3207 LNA-antimiR in mouse RAW cells using a luciferase sensor assay.

50 [0221] To ask whether a fully LNA-modified 8-mer LNA-antimiR can effectively antagonize miR-155, a perfect match target site for miR-155 was cloned into the same luciferase vector (psiCHECK2) and transfected into the mouse leukaemic monocyte macrophage RAW cell line. Because the endogenous levels of miR-155 are low in the RAW cell line, the cells were treated with 100 ng/ml LPS for 24 hours in order to induce miR-155 accumulation.

55 [0222] **Results**: Luciferase measurements showed that the fully LNA-modified 8-mer LNA-antimiR SEQ ID #3207 targeting miR-155 was similarly effective in antagonizing miR-155 compared to the 15-mer LNA-antimiR SEQ ID #3206 (Figure 4). Both LNA-antimirs showed a >50% de-repression of the miR-155 luciferase sensor at 0.25 nM concentration and inhibited miR-155 in a dose-dependent manner.

[0223] **Conclusion**: These data further support the results from antagonizing miR-21, as shown in examples 1 and

2, demonstrating that a fully thiolated 8-mer LNA-antimiR is highly potent in microRNA targeting.

#### Materials and Methods:

5 **[0224] Cell line:** The mouse leukaemic monocyte macrophage RAW 264.7 was purchased from ATCC (TIB-71). RAW cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 4 mM Glutamax and 25 ug/ml Gentamicin.

10 **[0225] Transfection:** 500.000 cells were seeded per well in a 6 well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, MCF-7 cells were transfected with 0.3 ug miR-155 or empty psiCHECK2 vector together with 10 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. In order to induce miR-155 accumulation, LPS (100 ng/ml) was added to the RAW cells after the 4 hour incubation with the transfection complexes. After another 24 hours, cells were harvested for luciferase measurements.

15 **[0226] Luciferase assay:** The cells were washed with PBS and harvested with cell scraper, after which cells were centrifugated for 5 min at 2.500 rpm. The supernatant were discarded and 50 µl 1 x Passive Lysis Buffer (Promega) was added to the cell pellet, after which cells were put on ice for 30 min. The lysed cells were spinned at 10.000 rpm for 30 min after which 20 µl were transferred to a 96 well plate and luciferase measurments were performed according to manufacturer's instructions (Promega).

#### 20 **Example 7: Assessment of miR-122 antagonism by SEQ ID #3208 LNA-antimiR in HuH-7 cells using a luciferase sensor assay.**

**[0227]** The potency of the fully modified 8-mer LNA-antimiR SEQ ID #3208 against miR-122 was assessed in the human hepatoma cell line HuH-7. The HuH-7 cells were transfected with luciferase sensor construct containing a perfect match miR-122 target site. After 24 hours luciferase measurements were performed (Figure 5).

25 **[0228] Results:** The fully LNA-modified 8-mer LNA-antimiR SEQ ID #3208 is more potent than the 15-mer LNA-antimiR SEQ ID #4 at low concentration, as shown by de-repression of the miR-122 luciferase sensor. Both LNA-antimiRs inhibit miR-122 in a dose-dependet manner (Figure 5). **Conclusion:** The fully LNA-modified 8-mer LNA-antimiR SEQ ID #3208 targeting miR-122 shows improved potency in inhibition of miR-122 *in vitro*.

#### 30 **Materials and Methods:**

**[0229] Cell line:** The human hepatoma cell line HuH-7 was a kind gift from R. Bartenschlager, Heidelberg. Huh-7 cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

35 **[0230] Transfection:** 8.000 cells were seeded per well in a 96 well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, HuH-7 cells were transfected with 57 ng miR-122 or empty psiCHECK2 vector together with 1 µl Lipofectamine2000 (Invitrogen). After 24 hours, cells were harvested for luciferase measurements.

40 **[0231] Luciferase assay:** 50 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 96 well plate was put on an orbital shaker for 30 min. To each well the Dual-luciferase Reporter assay system (Promega) was added and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### 45 **Example 8. Assessment of miR-21 antagonism by comparing an 8-mer (SEQ ID #3205) versus a 15-mer (SEQ ID #3204) LNA-antimiR in human prostate carcinoma cells (PC3).**

**[0232]** We have previously shown (patent application 1051), that an 8-mer LNA-antimiR that is fully LNA-modified and phosphorothiolated is able to completely de-repress the miR-21 luciferase reporter levels in the human cervix carcinoma cell line HeLa and partly de-repress the miR-21 luciferase reporter levels in the human breast carcinoma cell line MCF-7. We next extended this screening approach to the human prostate cancer cell line PC3. To assess the efficiency of the different LNA-antimiR oligonucleotides against miR-21, luciferase reporter constructs were generated in which a perfect match target site for the mature miR-21 and a target site with two mismatches in the seed were cloned in the 3'UTR of Renilla luciferase gene (Figure 7). In order to monitor miR-21 inhibition, PC3 cells were transfected with the different luciferase constructs together with the miR-21 antagonist SEQ ID #3205 (8-mer) and for comparison with the 15-mer LNA-antimiR perfect match SEQ ID #3204 at varying concentrations. After 24 hours, luciferase activity was measured.

55 **[0233] Results:** The luciferase reporter experiments showed a dose-dependent de-repression of the luciferase miR-21 reporter activity with the 15-mer LNA-antimiR against miR-21 (SEQ ID #3204). However, complete de-repression of the luciferase reporter was not obtained even at the highest concentrations (Figure 7). In contrast, the cells that were

transfected with the 8-mer fully LNA substituted LNA-antimiR showed complete de-repression already at 1 nM, indicating significantly improved potency compared to the 15-mer LNA-antimiR. The luciferase control reporter harboring a mismatch target site for miR-21 was not affected by either LNA-antimiR, demonstrating high specificity of both LNA-antimiRs.

**[0234] Conclusion:** The micromer is far more potent than the 15-mer LNA-antimiR in targeting miR-21 and has so far shown to be most potent in prostate carcinoma cells.

#### Materials and Methods:

**[0235] Cell line:** The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714). PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

**[0236] Transfection:** 100.000 cells were seeded per well in a 12-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with 0.3 µg miR-21 or empty psiCHECK2 vector together with 1,2 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0237] Luciferase assay:** The cells were washed with PBS and 250 µl 1 x Passive Lysis Buffer (Promega) was added to the wells. The plates were placed on a shaker for 30 min., after which the cell lysates were transferred to eppendorf tubes. The cell lysate was centrifugated for 10 min at 2.500 rpm after which 20 µl were transferred to a 96 well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### Example 9. Specificity assessment of miR-21 antagonism by an 8-mer LNA-antimiR

**[0238]** To investigate the specificity of our short LNA-antimiR targeting miR-21, we designed an 8-mer mismatch control LNA-antimiR (SEQ ID #3218) containing 2 mismatches in the seed recognition sequence (see Figure 8). The luciferase reporter constructs described in example 1 were transfected into the human cervix carcinoma cell line HeLa together with the LNA mismatch control oligo SEQ ID #3218 and its efficacy was compared with the 8-mer LNA-antimiR (SEQ ID #3205) targeting miR-21. After 24 hours, luciferase activity was measured. **Results:** As shown in Figure 8, transfection of the fully LNA-modified 8-mer LNA-antimiR in HeLa cells resulted in complete de-repression of the luciferase miR-21 reporter already at 5 nM. In contrast, when the cells were transfected with the 8-mer LNA mismatch control oligo, combined with the results obtained with the control miR-21 luciferase reporter having two mismatches in the miR-21 seed, these data demonstrate high specificity of the fully LNA-substituted 8-mer LNA-antimiR in HeLa cells.

**[0239]** Analysis of the miRBase microRNA sequence database showed that the miR-21 recognition sequence, of the LNA-antimiR SEQ ID #3205 is unique for microRNA-21. However, when decreasing the micromer length to 7 nt, it is not specific for only miR-21, since ath-miR-844, mmu-miR-590-3p and has-miR-590-3p are also targeted.

**[0240]** Conclusion: Exchanging two nucleotide positions within the 8-mer LNA-antimiR with two mismatching nucleotides completely abolished the antagonizing activity of the LNA-antimiR for miR-21.

#### Materials and Methods:

**[0241] Cell line:** The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0242] Transfection:** 60.000 cells were seeded per well in a 24-well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,7 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0243] Luciferase assay:** The cells were washed with PBS and 100 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates were put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96-well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### Example 10. Assessment of the shortest possible length of a fully LNA-modified LNA-antimiR that mediates effective antagonism of miR-21.

**[0244]** To further investigate the LNA-antimiR length requirements, we designed a 7-mer and a 6-mer LNA-antimiR targeting miR-21, both fully LNA-modified and phosphorothiolated oligonucleotides. The miR-21 luciferase reporter



constructs were transfected into HeLa cells along with the LNA-antimiRs at varying concentrations. Luciferase measurements were performed after 24 hours.

**[0245] Results:** As seen in Figure 9, the 7-mer LNA-antimiR mediates de-repression of the miR-21 luciferase reporter plasmid, but at lower potency compared to the 8-mer LNA-antimiR (SEQ ID #3205). Nevertheless, a dose-dependent trend can still be observed. By contrast, the 6-mer LNA-antimiR did not show any inhibitory activity.

**[0246] Conclusion:** To conclude, the shortest possible length of an LNA-antimiR which is able to mediate miR-21 inhibition is 7 nucleotides. However, the 7-mer LNA-antimiR is less potent compared to the 8-mer LNA-antimiR for miR-21.

#### Materials and Methods:

**[0247] Cell line:** The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0248] Transfection:** 60.000 cells were seeded per well in a 24 well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,7 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0249] Luciferase assay:** The cells were washed with PBS and 100 µl 1x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96-well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### Example 11. Length assessment of fully LNA-substituted LNA-antimiRs antagonizing miR-21

**[0250]** Next, we investigated the effect of increasing the length from a 9-mer to a 14-mer fully LNA substituted LNA-antimiRs on antagonizing miR-21 in HeLa cells. The resulting LNA-antimiRs were transfected into HeLa cells together with the miR-21 luciferase reporter constructs (Figure 10). Luciferase measurements were performed after 24 hours.

**[0251] Results:** The 9-mer LNA-antimiR SEQ ID #3211 (9-mer) showed dose-dependent de-repression of the miR-21 luciferase reporter which did not reach complete de-repression, as demonstrated for the 7-mer LNA-antimiR (SEQ ID #3210). Increasing the length to 10-mer to 14-mer (SEQ ID #3212, SEQ ID #3213 and SEQ ID #3214) decreased the potency as shown by less efficient de-repression of the miR-21 reporter.

**[0252] Conclusion:** As shown in Figure 10, the longest fully LNA-modified and phosphorothiolated LNA-antimiR which is still able to mediate miR-21 inhibition is a 9-mer LNA-antimiR SEQ ID #3211. However, it is clearly less efficient than the 7-mer and 8-mer LNA-antimiRs.

**[0253] Materials and Methods: Cell line:** The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0254] Transfection:** 60.000 cells were seeded per well in a 24-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 control vector without target site together with 0,7 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0255] Luciferase assay:** The cells were washed with PBS and 100 µl 1x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates were put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96-well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

#### Example 12. Determination of the most optimal position for an 8-mer LNA-antimiR within the miR target recognition sequence.

**[0256]** Our experiments have shown that the most potent fully LNA-modified phosphorothiolated LNA-antimiR is 8 nucleotides in length. To assess the most optimal position for an 8-mer LNA-antimiR within the miR target recognition sequence, we designed four different fully LNA-modified 8-mer LNA-antimiRs tiled across the mature miR-21 sequence as shown in Figure 11. The different LNA-antimiRs were co-transfected together with the miR-21 luciferase reporter constructs into HeLa cells. Luciferase measurements were performed after 24 hours.

**[0257] Results:** The only LNA-antimiR that mediated efficient silencing of miR-21 as measured by the luciferase reporter was SEQ ID #3205, which targets the seed region of miR-21. Neither SEQ ID #3215 which was designed to

cover the 3' end of the seed (50% seed targeting) did not show any effect, nor did the other two LNA-antimiRs SEQ ID #3216 or SEQ ID #3217, which were positioned to target the central region and the 3' end of the mature miR-21, respectively. **Conclusion:** The only 8-mer LNA-antimiR mediating potent silencing of miR-21 is the one targeting the seed of the miR-21.

5

#### Materials and Methods:

**[0258] Cell line:** The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0259] Transfection:** 60.000 cells were seeded per well in a 24-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,7 µl Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0260] Luciferase assay:** The cells were washed with PBS and 100 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96 well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

20

#### Example 13. Validation of interaction of the miR-21 target site in the Pcd4-3'-UTR and miR-21 using the 8-mer SEQ ID #3205 LNA-antimiR.

**[0261]** The tumour suppressor protein Pcd4 inhibits TPA-induced neoplastic transformation, tumour promotion and progression. Pcd4 has also been shown to be upregulated in apoptosis in response to different inducers. Furthermore, downregulation of Pcd4 in lung and colorectal cancer has also been associated with a poor patient prognosis. Recently, Asangani *et al* and Frankel *et al* showed that the Pcd4-3'-UTR contains a conserved target site for miR-21, and transfecting cells with an anti-miR-21, resulted in an increase in Pcd4 protein. We therefore constructed a luciferase reporter plasmid, harboring 313 nt of the 3'UTR region of Pcd4 encompassing the aforementioned miR-21 target site, which was co-transfected together with different LNA-antimiRs into HeLa cells. The different LNA-antimiRs were; SEQ ID #3205 (8-mer, perfect match) or SEQ ID #3218 (8-mer, mismatch). Luciferase measurements were performed after 24 hours.

25

30

**[0262] Results:** As shown in Figure 12, in cells transfected with the Pcd4 3'UTR luciferase reporter and SEQ ID #3205, an increase in luciferase activity was observed, indicating interaction between the Pcd4 3'UTR and miR-21. However, transfecting the cells with the mismatch compound, SEQ ID #3218, no change in luciferase activity was observed, which was expected since the compound does not antagonize miR-21. When comparing the 8-mer LNA-antimiR against two longer designed LNA-antimiRs, the short fully LNA-modified and phosphorothiolated LNA-antimiR was significantly more potent, confirming previous luciferase assay data. **Conclusion:** These data conclude that SEQ ID #3205, which antagonizes miR-21, can regulate the interaction between Pcd4 3'UTR and miR-21.

40

#### Materials and Methods:

**[0263] Cell line:** The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

45

**[0264] Transfection:** 60.000 cells were seeded per well in a 24-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 0.2 ug Pcd4-3'UTR/psiCHECK2 or empty psiCHECK2 vector together with 0,7 µl Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Varying concentrations of the LNA-antimiR oligonucleotides were also transfected. After 24 hours, cells were harvested for luciferase measurements.

50

**[0265] Luciferase assay:** The cells were washed with PBS and 100 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to an eppendorf tube and spinned at 10.000 rpm for 30 min after which 10 µl were transferred to a 96 well plate and luciferase measurements were performed according to manufacturer's instructions (Promega).

55

**Example 14. Comparison of an 8-mer LNA-antimiR (SEQ ID #3207) with a 15-mer LNA-antimiR (SEQ ID #3206) in antagonizing miR-155 in mouse RAW cells.**

[0266] To ask whether our approach of using short LNA-antimiRs could be adapted to targeting other miRNAs we designed a fully LNA-modified 8-mer LNA-antimiR against microRNA-155. A perfect match target site for miR-155 was cloned into the 3'UTR of the luciferase gene in the reporter plasmid psiCHECK2 and transfected into the mouse RAW macrophage cell line together with an 8-mer or a 15-mer LNA-antimiR. Because the endogenous levels of miR-155 are low in the RAW cell line, the cells were treated with 100 ng/ml LPS for 24 hours in order to induce miR-155 accumulation. After 24 hours, luciferase analysis was performed.

[0267] **Results:** Luciferase measurements showed that the fully LNA-modified 8-mer LNA-antimiR SEQ ID #3207 targeting miR-155 was similarly effective in antagonizing miR-155 compared to the 15-mer LNA-antimiR SEQ ID #3206 (Figure 13). Both LNA-antimiRs showed a >50% de-repression of the miR-155 luciferase sensor at 0.25 nM concentration and inhibited miR-155 in a dose-dependent manner.

[0268] Analysis of the miRBase microRNA sequence database showed that the miR-155 recognition sequence, of the LNA-antimiR SEQ ID #3207 is unique for microRNA-155. However, when decreasing the LNA-antimiR length to 7 nt, it is not specific for only miR-155, mdv1-miR-M4 and kshv-miR-K12-11 is also targeted.

[0269] **Conclusion:** A fully LNA-modified and phosphorothiolated 8-mer LNA-antimiR is equally potent compared with a 15-mer LNA-antimiR of a mixed LNA/DNA design in antagonizing miR-155. Thus, our approach of using short LNA-antimiRs can be readily adapted to targeting of other miRNAs

**Materials and Methods:**

[0270] **Cell line:** The mouse macrophage RAW 264.7 cell line was purchased from ATCC (TIB-71). RAW cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 4 mM Glutamax and 25 ug/ml Gentamicin.

[0271] **Transfection:** 500,000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, RAW 264.7 cells were transfected with 0.3 ug miR-155 perfect match/psiCHECK2 or empty psiCHECK2 vector together with 10  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. In order to induce miR-155 accumulation, LPS (100 ng/ml) was added to the RAW cells after the 4 hour incubation with the transfection complexes. After another 24 hours, cells were harvested for luciferase measurements.

[0272] **Luciferase assay:** The cells were washed with PBS and harvested with cell scraper, after which cells were spinned for 5 min at 2,500 rpm. The supernatant was discarded and 50  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to the cell pellet, after which cells were put on ice for 30 min. The lysed cells were spinned at 10,000 rpm for 30 min after which 20  $\mu$ l were transferred to a 96-well plate and luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 15. Assessment of c/EBP $\beta$  protein levels as a functional readout for miR-155 antagonism by short LNA-antimiR (SEQ ID #3207).**

[0273] As a functional readout for miR-155 antagonism by short LNA-antimiR (SEQ ID #3207) we determined the protein levels of a novel miR-155 target, c/EBP $\beta$ . The mouse macrophage RAW cell line was transfected together with either an 8-mer (SEQ ID #3207) or a 15-mer (SEQ ID #3206) LNA-antimiR in the absence or presence of pre-miR-155. As mismatch controls for the 15-mer, SEQ ID #4 was used, which targets miR-122 and for the 8-mer SEQ ID #3205 was used, which targets miR-21. These two control miRNAs do not regulate c/EBP $\beta$  expression levels. LPS was used to induce miR-155 accumulation and cells were harvested after 16 hours with LPS. c/EBP $\beta$  has three isoforms; LIP, LAP and LAP\* that were detected by Western blot analysis and the same membranes were re-probed with beta-tubulin as loading control. **Results:** Ratios were calculated for c/EBP $\beta$  LIP and beta-tubulin as indicated in Figure 14. RAW cells that were transfected with the 15-mer LNA-antimiR and no pre-miR-155 all showed equal c/EBP $\beta$  LIP/beta-tubulin ratios, due to inhibition of miR-155 increases the c/EBP $\beta$  LIP levels (Figure 14, left panel). By comparison, transfection of pre-miR-155 in RAW cells resulted in decreased c/EBP $\beta$  LIP levels as expected, if c/EBP $\beta$  was a miR-155 target, as shown in lanes with protein extracts from RAW cells treated with no LNA or a mismatch. However, protein extracts from RAW cells transfected with LNA-antimiR against miR-155, showed an increase of c/EBP $\beta$  LIP levels. The same experiments were also carried out with the 8-mer LNA-antimiR-155 (SEQ ID #3207) and as shown in Figure 14 (right panel) comparable results to those with the 15-mer LNA-antimiR SEQ ID #3206 were obtained.

[0274] **Conclusion:** Antagonism of miR-155 using either an 8-mer or a 15-mer LNA-antimiR leads to de-repression of the direct target c/EBP $\beta$ .

**Materials and Methods:**

[0275] Cell line: The mouse macrophage RAW 264.7 cell line was purchased from ATCC (TIB-71). RAW cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 4 mM Glutamax and 25 ug/ml Gentamicin.

[0276] Transfection: 500,000 cells were seeded per well in a 6-well plate the day before transfection in order to achieve 50% confluency the next day. On the day of transfection, RAW 264.7 cells were transfected with 5 nmol pre-miR-155 (Ambion) and/or 5 nM LNA-antimiR together with 10  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. In order to induce miR-155 accumulation, LPS (100 ng/ml) was added to the RAW cells after the 4 hour incubation with the transfection complexes. After 16 hours, cells were harvested for protein extraction and western blot analysis. Western blot: Cells were washed with PBS, trypsinated, transferred to eppendorf tubes and 250  $\mu$ l lysis buffer (1xRIPA) was added. The cell lysate was placed on ice for 20 min and spinned at 10,000 rpm for 10 minutes. The protein concentration was measured with Coomassie Plus according to the manufacturer's instructions and 80 ug was loaded onto a 4-12% BIS-TRIS gel. The membrane was incubated overnight at 4°C with the primary monoclonal mouse antibody C/EBP  $\beta$  (Santa Cruz) with a 1:100 concentration, Immunoreactive bands were visualized with ECL Plus (Amersham).

**Example 16. Antagonism of miR-106b by a fully LNA-modified 8-mer (SEQ ID #3221) LNA-antimiR**

[0277] To confirm that our approach of using short LNA-antimiRs could be adapted to targeting of other miRNAs we designed a fully LNA-modified 8-mer LNA-antimiR against microRNA-106b. A perfect match target site for miR-106b was cloned into the 3'UTR of the luciferase gene in the vector (psiCHECK2) and transfected into the human cervix carcinoma HeLa cell line together with a short LNA-antimiR (SEQ ID #3221) or with a 15-mer LNA-antimiR (SEQ ID #3228) at varying concentrations. Luciferase measurements were performed after 24 hours.

[0278] **Results:** Transfection of the 8-mer LNA-antimiR SEQ ID #3221 against miR-106b resulted in dose-dependent inhibition of miR-106b as shown by de-repression of the luciferase reporter, which was completely de-repressed at 1 nM LNA-antimiR concentration (Figure 15). Comparable results were obtained using the 15-mer LNA-antimiR SEQ ID #3228 demonstrating that an 8-mer LNA-antimiR is similarly potent to a 15-mer.

[0279] **Conclusion:** Targeting of miR-106b in HeLa cells shows that an 8-mer fully LNA-modified and phosphorotiolated LNA-antimiR is equally potent compared with a 15-mer LNA/DNA mixmer LNA-antimiR.

**Materials and Methods:**

[0280] Cell line: The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

[0281] Transfection: 5,200 cells were seeded per well in a 96-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 57 ng miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,14  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

[0282] Luciferase assay: The cells were washed with PBS and 30  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to eppendorf tubes and spinned at 10,000 rpm for 30 min after which luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 17. Antagonism of miR-19a by a fully LNA-modified 8-mer (SEQ ID #3222) LNA-antimiR**

[0283] To further confirm that our approach of using short LNA-antimiRs can be readily adapted to targeting of other miRNAs we designed a fully LNA-modified 8-mer LNA-antimiR against microRNA-19a. A perfect match target site for miR-19a was cloned in the 3'UTR of the luciferase gene in the psiCHECK2 vector. The reporter plasmid was transfected into the human cervix carcinoma HeLa cell line together with a short LNA-antimiR (SEQ ID #3222) or with a 15-mer LNA-antimiR (SEQ ID #3229) targeting miR-19a at varying concentrations. Luciferase measurements were performed after 24 hours.

[0284] **Results:** As shown in Figure 16, transfection of the 15-mer LNA-antimiR SEQ ID #3229 into HeLa efficiently antagonizes miR-19a as demonstrated by complete de-repression at 1 nM LNA-antimiR concentration. By comparison, transfection of the 8-mer LNA-antimiR SEQ ID #3222 resulted in effective miR-19a antagonism already at 0.5 nM concentration, indicating that this 8-mer LNA-antimiR is at least equally potent compared with a 15-mer LNA-antimiR in HeLa cells.

**[0285] Conclusion:** Targeting of miR-19a in HeLa cells shows that an 8-mer fully LNA-modified and phosphorothiolated LNA-antimiR is at least equally potent compared with a 15-mer LNA/DNA mixmer LNA-antimiR.

**[0286] Materials and Methods:** *Cell line:* The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0287] Transfection:** 5.200 cells were seeded per well in a 96-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 57 ng miR-21 perfect match/psiCHECK2, miR-21.mm2/psiCHECK2 or empty psiCHECK2 vector together with 0,14 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0288] Luciferase assay:** The cells were washed with PBS and 30 µl 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 24-well plates was put on an orbital shaker for 30 min. The cells were collected and transferred to eppendorf tubes and spinned at 10.000 rpm for 30 min after which luciferase measurements were performed according to the manufacturer's instructions (Promega).

### Example 18. Targeting of a microRNA family using short, fully LNA-substituted LNA-antimiR.

**[0289]** Next, we investigated whether it is possible to target a microRNA family using a single short 7-mer LNA-antimiR complementary to the seed sequence that is common for all family members (see Figure 17). In this experiment, we focused on miR-221 and miR-222 that are overexpressed in solid tumors of the colon, pancreas, prostate and stomach. It has also been shown that miR-221 and miR-222 are the most significantly upregulated microRNAs in glioblastoma multiforme. Furthermore, overexpression of miR-221 and miR-222 may contribute to the growth and progression of prostate carcinoma, at least in part by blocking the tumor suppressor protein p27. A perfect match target site for both miR-221 and miR-222, respectively, was cloned into the 3'UTR of the luciferase gene resulting in two reporter constructs. These constructs were then transfected either separate or combined into the prostate carcinoma cell line, PC3. In addition to the 7-mer, targeting both miR-221 and miR-222, we also co-transfected a 15-mer LNA-antimiR (15mer) targeting either miR-221 (SEQ ID #3223) or miR-222 (SEQ ID #3224), each transfected separately or together (see Figure 18 left). **Results:** As shown in Figure 18, transfection of PC3 cells with the LNA-antimiR SEQ ID #3223 against miR-221 resulted in efficient inhibition of miR-221 at 1 nM LNA-antimiR concentration. An inhibitory effect is also observed when using the luciferase reporter plasmid for miR-222 as well as when co-transfecting both luciferase reporters for miR-221 and miR-222 simultaneously into PC3 cells. This inhibitory effect is most likely due to the shared seed sequence between miR-221 and miR-222. Similarly, transfection of PC3 cells with the LNA-antimiR SEQ ID #3224 against miR-222 resulted in efficient inhibition of miR-222 at 1 nM LNA-antimiR concentration as shown by complete de-repression of the luciferase reporter for miR-222. An inhibitory effect is also observed when using the luciferase reporter plasmid for miR-222 as well as when co-transfecting both luciferase reporters for miR-221 and miR-222 simultaneously into PC3 cells. Co-transfection of both LNA-antimiR compounds SEQ ID #3223 and SEQ ID #3224 against miR-221 and miR-222, respectively, (see Figure 18 left), resulted in effective inhibition of both miRNAs as shown by complete de-repression of the luciferase reporter plasmids both when separately transfected and when co-transfected into PC3 cells. Interestingly, transfection of a single fully LNA-modified 7-mer LNA-antimiR (SEQ ID #3225) targeting the seed sequence of miR-221 and miR-222 into PC3 cells resulted in efficient, dose-dependent antagonism of miR-221 and miR-222 simultaneously as shown by complete de-repression of the luciferase reporter plasmids both when separately transfected and when co-transfected into PC3 cells. This demonstrates that a single, short LNA-substituted LNA-antimiR can effectively target seed sequences thereby antagonizing entire microRNA families simultaneously. Analysis of the miRBase microRNA sequence database showed that the miR-221/222 seed recognition sequence, of the LNA-antimiR SEQ ID #3225 is unique for both miRNAs.

**[0290] Conclusion:** Our results demonstrate that LNA enables design and synthesis of short fully LNA-substituted LNA-antimiR oligonucleotides that can effectively target microRNA seed sequences thereby antagonizing entire microRNA families simultaneously.

### Materials and Methods:

**[0291] Cell line:** The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714) PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

**[0292] Transfection:** 100.000 cells were seeded per well in a 12-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with 0.3 ug of luciferase reporter plasmid for miR-221 or for miR-222 or with empty psiCHECK2 vector without miRNA target site as control together with 1,2 µl Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

**[0293]** Luciferase assay: The cells were washed with PBS and 250  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to the wells. The plates were placed on a shaker for 30 min., after which the cell lysates was transferred to eppendorf tubes. The cell lysate was spinned for 10 min at 2.500 rpm after which 20  $\mu$ l were transferred to a 96-well plate and luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 19. Assessment of p27 protein levels as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR.**

**[0294]** Previous work has shown (le Sage et al. 2007, Galardi et al. 2007) that miR-221 and miR-222 post-transcriptionally regulate the expression of the tumour suppressor gene p27, which is involved in cell cycle regulation. In these studies, down-regulation of miR-221 and miR-222 was shown to increase expression levels of p27. Thus, as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR we determined the protein levels of p27 after transfection of the LNA-antimiR SEQ ID #3225 into PC3 cells in comparison with an 8-mer LNA mismatch control. After 24 hours the cells were harvested for western blot analysis (Figure 19).

**[0295] Results:** As shown in Figure 19, transfection of the 7-mer LNA-antimiR SEQ ID #3225 targeting the seed sequence in miR-221 and miR-222 resulted in dose-dependent increase of the p27 protein levels compared to either untransfected or LNA mismatch control transfected PC3 cells. These results clearly demonstrate that the 7-mer LNA-antimiR is able to effectively antagonize the miR-221/222 family leading to de-repression of the direct target p27 at the protein level. **Conclusion:** A fully LNA-modified 7-mer LNA-antimiR targeting the seed sequence in the miR-221/222 family effectively antagonized both miRNAs leading to de-repression of the direct target p27 at the protein level.

**Materials and Methods:**

**[0296]** Cell line: The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714) PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

**[0297]** Transfection: 250.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with LNA-antimiRs at varying concentrations with Lipofectamine2000. Cells were harvested after 24 hours for protein extraction and western blot analysis.

**[0298]** Western blot: Cells were washed with PBS, trypsinated, transferred to eppendorf tubes and 250  $\mu$ l lysis buffer (1xRIPA) was added. The cell lysate was placed on ice for 20 min, then spinned at 10.000 rpm for 10 minutes. The protein concentration was measured with Coomassie Plus according to the manufacturer's instructions and 100 ug was loaded onto a 4-12% BIS-TRIS gel. The membrane was incubated overnight at 4°C with the primary monoclonal mouse antibody p27 (BD Biosciences) at a 1:1000 dilution. Immunoreactive bands were visualized with ECL Plus (Amersham).

**Example 20. Duplex melting temperatures ( $T_m$ ) of the LNA-antimiRs.**

**[0299]** As shown in Table 5,  $T_m$  values increase with increasing the length of short fully modified LNA-antimiRs (see  $T_m$  values for SEQ ID #3205, SEQ ID #3209-3214 in Table 7). Most optimal inhibitory effect was achieved with the 8-mer LNA-antimiR SEQ ID #3205 against miR-21, whereas the very low  $T_m$  of the 6-mer SEQ ID #3209 is most likely not sufficient to mediate antagonism of the miR-21 target. On the other hand, increasing the length beyond a 10-mer (SEQ ID #3212) significantly increases the  $T_m$ , while simultaneously decreasing the inhibitory activity as measured using the luciferase miR-21 reporter, which is most likely due to high propensity of the fully modified 12- and 14-mer LNA-antimiRs to form homodimers. The experiments using a sliding window of fully LNA-modified 8-mer LNA-antimirs across the mir-21 recognition sequence clearly demonstrate that in addition to adequate  $T_m$  value of the LNA-antimiR, the seed region is most critical for miRNA function and, thus, the most optimal region to be targeted by an LNA-antimiR.

**Table 5:**  $T_m$  values for miR-21 LNA-antimiRs, measured against a complementary RNA oligonucleotide

SEQ ID #	microRN A	Length (bp)	Sequence	Measured $T_m$ (RNA) °C
3205	miR-21	8	5'- GATAAGCT -3'	64,0
3209	miR-21	6	5'- TAAGCT -3'	32,0
3210	miR-21	7	5'- ATAAGCT -3'	45,0
3211	miR-21	9	5'- TGATAAGCT -3'	65,0
3212	miR-21	10	5'- CTGATAAGCT -3'	63,0

(continued)

SEQ ID #	microRN A	Length (bp)	Sequence	Measured $T_m$ (RNA) °C
3213	miR-21	12	5'- GTCTGATAAGCT -3'	86,8
3214	miR-21	14	5'- CAGTCTGATAAGCT -3'	89,9
3215	miR-21	8	5'- TCTGATAA - 3'	56,0
3216	miR-21	8	5'- ATCAGTCT - 3	72,0
3217	miR-21	8	5'- TCAACATC - 3	48,0

**[0300] Conclusion:** The  $T_m$  values along with experimental data obtained with luciferase reporters show that potent antagonism by LNA-antimiR is not only dependent on  $T_m$  but also depends on the positioning of the LNA-antimiR within the microRNA recognition sequence.

#### Materials and Methods:

**[0301]  $T_m$  measurements:** The oligonucleotide:miR-21 RNA duplexes were diluted to 3  $\mu$ M in 500  $\mu$ l RNase free H<sub>2</sub>O and mixed with 500  $\mu$ l 2x  $T_m$ -buffer (200 mM NaCl, 0.2 mM EDTA, 20 mM Na-phosphate, pH 7,0). The solution was heated to 95°C for 3 min and then allowed to anneal in RT for 30 min. The duplex melting temperatures ( $T_m$ ) were measured on a Lambda 40 UV/VIS Spectrophotometer equipped with a Peltier temperature programmer PTP6 using PE Templab software (Perkin Elmer). The temperature was ramped up from 20°C to 95°C and then down to 25°C, recording absorption at 260 nm. First derivative and the local maximums of both the melting and annealing were used to assess the duplex melting temperatures.

#### Example 21. Assessment of miR-21 antagonism by comparing an 8-mer (SEQ ID #3205) versus a 15-mer (SEQ ID #3204) LNA-antimiR in human hepatocytic cell line HepG2.

**[0302]** We have previously shown in this application, that an 8-mer LNA-antimiR that is fully LNA-modified and phosphorothiolated effectively antagonizes miR-21 in the human cervix carcinoma cell line HeLa, the human breast carcinoma cell line MCF-7 and the human prostate cancer cell line PC3. We extended this screening approach to the human hepatocellular cancer cell line HepG2. To assess the efficiency of the 8-mer LNA-antimiR oligonucleotide against miR-21, luciferase reporter constructs were generated in which a perfect match target site for the mature miR-21 was cloned into the 3'UTR of the Renilla luciferase gene. In order to monitor miR-21 inhibition, HepG2 cells were transfected with the luciferase constructs together with the miR-21 antagonist SEQ ID #3205 (8-mer) and for comparison of specificity with the 8-mer LNA-antimiR mismatch (SEQ ID #3218) and for comparison of potency together with the 15-mer (SEQ ID #3204) at varying concentrations. After 24 hours, luciferase activity was measured.

**[0303] Results:** The luciferase reporter experiments showed a dose-dependent de-repression of the luciferase miR-21 reporter activity with the 15-mer LNA-antimiR against miR-21 (SEQ ID #3204). However, complete de-repression of the luciferase reporter was not obtained, not even at the higher concentrations (Figure 20). In contrast, the cells that were transfected with the 8-mer fully LNA modified LNA-antimiR (SEQ ID #3205) showed complete de-repression already at 5 nM, indicating significantly improved potency compared to the 15-mer LNA-antimiR. Comparing the specificity of the 8-mer perfect match and the 8-mer mismatch, the mismatch LNA-antimiR (SEQ ID #3218) did not show any de-repression at all, demonstrating high specificity of the LNA-antimiR compound against miR-21.

**[0304] Conclusion:** The 8-mer (SEQ ID #3205) is more potent than the 15-mer LNA-antimiR in targeting miR-21 and antagonism of miR-21 by SEQ ID #3205 is specific.

#### Materials and Methods:

**[0305] Cell line:** The human hepatocytic HepG2 cell line was purchased from ECACC (#85011430). HepG2 cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

**[0306] Transfection:** 650.000 cells were seeded per well in a 6-well plate and reverse transfection were performed. HepG2 cells were transfected with 0.6  $\mu$ g miR-21 or empty psiCHECK2 vector together with 2,55  $\mu$ l Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. After 24 hours, cells were harvested for luciferase measurements.

**[0307] Luciferase assay:** The cells were washed with PBS and 300  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to the wells. The plates were placed on a shaker for 30 min., after which the cell lysates were transferred to eppendorf

tubes. The cell lysate was centrifugated for 10 min at 2.500 rpm after which 50  $\mu$ l were transferred to a 96 well plate and luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 22. Validation of interaction of the miR-21 target site in the Pdc4 3'UTR and miR-21 using the 8-mer SEQ ID #3205 LNA-antimiR in human hepatocellular cell line Huh-7.**

**[0308]** The tumour suppressor protein Pdc4 inhibits tumour promotion and progression. Furthermore, downregulation of Pdc4 in lung and colorectal cancer has also been associated with poor patient prognosis. Recently, Asangani et al (Oncogene 2007) and Frankel et al (J Biol Chem 2008) showed that the Pdc4 3'UTR contains a conserved target site for miR-21, and transfecting cells with an anti-miR-21, resulted in an increase in Pdc4 protein. We therefore constructed a luciferase reporter plasmid, harboring 313 nt of the 3'UTR region of Pdc4 encompassing the aforementioned miR-21 target site, which was co-transfected together with different LNA-antimiRs and pre-miR-21 (10 nM) into Huh-7 cells. The different LNA-antimiRs were; SEQ ID #3205 (8-mer, perfect match), SEQ ID #3218 (8-mer, mismatch) and SEQ ID #3204 (15-mer, DNA/LNA mixmer). Luciferase measurements were performed after 24 hours. **Results:** As shown in Figure 21, cells transfected with the Pdc4 3'UTR luciferase reporter and SEQ ID #3205, an increase in luciferase activity was observed, indicating interaction between the Pdc4 3'UTR and miR-21. However, transfecting the cells with the mismatch compound, SEQ ID #3218, no change in luciferase activity was observed, which was expected since the compound does not antagonize miR-21. When comparing the 8-mer LNA-antimiR against the 15-mer LNA-antimiR (SEQ ID #3204), the short fully LNA-modified and phosphorothiolated LNA-antimiR was significantly more potent, confirming previous data.

**Materials and Methods:**

**[0309]** Cell line: The human hepatoma cell line Huh-7 was a kind gift from R. Bartinschlager (Dept Mol Virology, University of Heidelberg). Huh-7 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25  $\mu$ g/ml Gentamicin.

**[0310]** Transfection: 11.000 cells were seeded per well in a 96-well plate the day before transfection in order to achieve 50-70% confluency the next day. On the day of transfection, Huh-7 cells were transfected with 20 ng Pdc4 3'UTR/psiCHECK2 or empty psiCHECK2 vector together with 10 nM pre-miR-21 (Ambion) and 0,14  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Varying concentrations of the LNA-antimiR oligonucleotides were also transfected. After 24 hours, cells were harvested for luciferase measurements.

**[0311]** Luciferase assay: Cells were washed and 30  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to each well, after which the 96-well plates was put on an orbital shaker. After 30 min., 50  $\mu$ l luciferase substrate dissolved in Luciferase Assay Buffer II (Dual-Luciferase Reporter Assay System from Promega, Cat# E1910) was added to the wells with lysated cells and luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 23. Assessment of Pdc4 protein levels as a functional readout for miR-21 antagonism by the 8-mer LNA-antimiR (SEQ ID #3205).**

**[0312]** In addition, we also transfected HeLa cells with SEQ ID #3205 (perfect match), SEQ ID #3218 (mismatch), SEQ ID #3219 (scrambled) and analyzed Pdc4 protein levels after 24 hours with Western blot (Figure 22). As shown, in the protein extracts from cells where SEQ ID #3205 had been added, the Pdc4 protein levels increase, due to antagonism of miR-21 by SEQ ID #3205 in contrast to the two control LNA oligonucleotides.

**[0313]** **Conclusion:** Antagonism of miR-21 using an 8-mer (SEQ ID #3205) leads to derepression of the direct target Pdc4 antagonism of miR-21

**Materials and Methods:**

**[0314]** Cell line: The human cervix carcinoma cell line HeLa was purchased from ECACC (#93021013). HeLa cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25  $\mu$ g/ml Gentamicin.

**[0315]** Transfection: 200.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50-70% confluency the next day. On the day of transfection, HeLa cells were transfected with 5 nM LNA oligonucleotides and 2,5  $\mu$ g/ml Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. After 24 hours, cells were harvested for Western blot analysis.

**[0316]** Western blot: Cells were washed with PBS, trypsinated, transferred to eppendorf tubes and 50  $\mu$ l lysis buffer (1xR1PA) was added. The cell lysate was placed on ice for 20 min and spinned at 10.000 rpm for 10 minutes. Equal amounts (15  $\mu$ l cell lysate) were loaded onto a 4-12% BIS-TRIS gel. The proteins were transferred to a nitrocellulose



membrane using iBlot (Invitrogen) according to manufacturers instructions. The membrane was incubated overnight at 4°C with the primary affinity purified rabbit serum antibody Pdc4 (Rockland) with a 1:2000 concentration. As control, anti- beta tubulin antibodies (Thermo Scientific) were used at a 1:5000 dilution. Immunoreactive bands were visualized with ECL Plus (Amersham).

**Example 24. Assessment of potential hepatotoxicity of the 8-mer perfect match LNA-antimiR SEQ ID #3205 and the LNA mismatch control SEQ ID #3218.**

[0317] Each compound was injected into female NMRI mice, at doses of 25 mg/kg, 5 mg/kg and 1 mg/kg, every other day for 2 weeks. The animals were sacrificed and serum was collected from whole blood for ALT and AST analyses. As seen in Figure 23, the ALT and AST levels were not elevated for SEQ ID #3205 compared to saline or SEQ ID #3218 (mismatch control). However, one mouse showed increased levels (marked red), since the serum samples were contaminated with red blood cells, which contain 6-8 times higher levels of ALT and AST compared to plasma. The mice that received 5 mg/kg and 1 mg/kg were also analyzed for ALT and AST levels and showed no changes compared to saline treated control animals (data not shown).

**Materials and Methods:**

Experimental design:

[0318]

Gr. no.	Animal IDno.	No. of mice	Compound Dose level per day	Conc. at dose vol. 10 ml/ kg	Adm. Route	Dosing
1	1 -10	10	NaCl 0.9%	-	i.v	0, 2, 4, 7, 9
2	11-15	5	SEQ ID # 3205 25mg/kg	2.5mg/ml	i.v	0, 2, 4, 7, 9
3	16-20	5	SEQ ID #3205	0.5mg/ml	i.v	0, 2, 4, 7, 9
4	21-25	5	SEQ ID # 3205 1mg/kg	0.1m/ml	i.v	0, 2, 4, 7, 9
5	26-30	5	SEQ ID # 3230 25mg/kg	2.5mg/ml	i.v	0, 2, 4, 7, 9
6	31-35	5	SEQ ID # 3230 5mg/kg	0.5mg/ml	i.v	0, 2, 4, 7, 9

[0319] Sacrifice: The animals was sacrificed by cervical dislocation.

[0320] Sampling of serum for ALT/AST; The animals were anaesthetised with 70% CO<sub>2</sub>-30% O<sub>2</sub> before collection of retro orbital sinus blood. The blood was collected into S-monovette Serum-Gel vials. The serum samples were harvested and stored from each individual mouse. The blood samples were stored at room temperature for two hours and thereafter centrifuged 10 min, 3000 rpm, at room temp. The serum fractions were harvested into Eppendorf tubes on wet ice.

[0321] ALT and AST measurements; ALT and AST measurements were performed in 96-well plates using ALT and AST reagents from ABX Pentra (A11A01627 - ALT, A11A01629 - AST) according to the manufacturer's instructions. In short, serum samples were diluted 2.5 fold with H<sub>2</sub>O and each sample was assayed in duplicate. After addition of 50 µl diluted sample or standard (multical from ABX Pentra - A11A01652) to each well, 200 µl of 37 °C AST or ALT reagent mix was added to each well. Kinetic measurements were performed for 5 min with an interval of 30s at 340 nm and 37 °C.

**Example 25. Assessment of PU.1 protein levels as a functional readout for miR-155 antagonism by short LNA-antimiR (SEQ ID #3207).**

[0322] We have previously shown that the 8-mer (SEQ ID #3207) antagonizing miR-155 leads to derepression of the miR-155 target c/EBPbeta in the mouse macrophage RAW cells. To further verify the potency of SEQ ID #3207 we

determined the protein levels of another miR-155 target, PU.1 As a functional readout for miR-155 antagonism by short LNA-antimiR (SEQ ID #3207) we performed Western blot. The antagonism was verified in the human monocytic THP-1 cell line which was transfected together with either an 8-mer (SEQ ID #3207) perfect match or a 8-mer control LNA in the absence or presence of pre-miR-155. LPS was used to induce miR-155 accumulation and cells were harvested after 24 hours.

**[0323] Results:** THP-1 cells that were transfected with pre-miR-155 shows a decrease in PU.1 levels (Figure 24). Transfecting the cells with the fully LNA-modified and phosphorothiolated SEQ ID #3207 effectively antagonizes miR-155, leading to unaltered levels of PU.1 protein. By comparison, transfecting the cells with an 8-mer LNA control, PU.1 levels decreased, indicating that antagonism of miR-155 by SEQ ID #3207 LNA-antimiR is specific.

**[0324] Conclusion:** Antagonism of miR-155 using an 8-mer leads to de-repression of the direct target PU.1 in human THP-1 cells.

#### Materials and Methods:

**[0325] Cell line:** The human monocytic THP-1 cell line was purchased from ECACC (#88081201). THP-1 cells were cultured in RPMI with L-glutamine, supplemented with 10% fetal bovine serum.

**[0326] Transfection:** 200.000 cells were seeded per well in a 12-well plate the day before. On the day of transfection, THP-1 cells were transfected with 5 nmol pre-miR-155 (Ambion) and/or 5 nM LNA-antimiR together with Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. LPS (100 ng/ml) was added to the cells after the 4 hour incubation with the transfection complexes. After 24 hours, cells were harvested for protein extraction and western blot analysis.

**[0327] Western blot:** Cells were washed with PBS, trypsinated, transferred to eppendorf tubes and 50 µl lysis buffer (1xRIPA) was added. The cell lysate was placed on ice for 20 min and spinned at 10.000 rpm for 10 minutes. Equal amounts (15 µl cell lysate) were loaded onto a 4-12% BIS-TRIS gel. The proteins were transferred to a nitrocellulose membrane using iBlot (Invitrogen) according to manufacturers instructions The membrane was incubated overnight at 4°C with the rabbit monoclonal PU.1 antibody (Cell Signaling) with a 1:2000 concentration. As equal loading, Tubulin (Thermo Scientific) was used at a 1:5000 dilution. Immunoreactive bands were visualized with ECL Plus (Amersham).

#### Example 26. Assessment of p27 protein levels as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR.

**[0328]** Previous work has shown (le Sage et al. 2007, Galardi et al. 2007) that miR-221 and miR-222 post-transcriptionally regulate the expression of the tumour suppressor gene p27, which is involved in cell cycle regulation. In these studies, down-regulation of miR-221 and miR-222 was shown to increase expression levels of p27. Thus, as a functional readout for antagonism of the miR-221/222 family by the 7-mer SEQ ID #3225 LNA-antimiR we determined the protein levels of p27 after transfection of the LNA-antimiR SEQ ID #3225 into PC3 cells.

**[0329] Results:** As shown in Figure 25, transfection of the 7-mer LNA-antimiR SEQ ID #3225 targeting the seed sequence of miR-221 and miR-222 resulted in dose-dependent increase of the p27 protein levels compared to either untransfected or our LNA scrambled control transfected PC3 cells. These results clearly demonstrate that the 7-mer LNA-antimiR is able to effectively antagonize the miR-221/222 family leading to de-repression of the direct target p27 at the protein level at concentrations as low as 5 nM.

**[0330] Conclusion:** A fully LNA-modified 7-mer LNA-antimiR targeting the seed sequence in the miR-221/222 family at 5 nM can effectively antagonize both miRNAs leading to de-repression of the direct target p27 at protein level.

#### Materials and Methods:

**[0331] Cell line:** The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714). PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

**[0332] Transfection:** 250.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with LNA-oligonucleotides at varying concentrations (see Figure 25) with Lipofectamine2000. Cells were harvested after 24 hours for protein extraction and western blot analysis.

**[0333] Western blot:** Cells were washed with PBS, trypsinated, transferred to eppendorf tubes and 50 µl lysis buffer (1xRIPA) was added. The cell lysate was placed on ice for 20 min, then spinned at 10.000 rpm for 10 minutes. Equal amounts (15 µl cell lysate) were loaded onto a 4-12% BIS-TRIS gel. The proteins were transferred to a nitrocellulose membrane using iBlot (Invitrogen) according to manufacturers instructions. The membrane was incubated overnight at 4°C with the primary monoclonal mouse antibody p27 (BD Biosciences) at a 1:1000 dilution. As loading control, Tubulin (Thermo Scientific) was used at a 1:5000 dilution. Immunoreactive bands were visualized with ECL Plus (Amersham).

**Example 27. Knock-down of miR-221/222 by the 7-mer SEQ ID #3225 LNA-antimiR reduces colony formation of PC3 cells**

[0334] A hallmark of cellular transformation is the ability for tumour cells to grow in an anchorage-independent way in semisolid medium. We have therefore performed soft agar assay which is a phenotypic assay that is relevant for cancer, given that it measures the decrease of tumour cells. We transfected SEQ ID #3225 (perfect match) and SEQ ID #3231 (scrambled) into PC3 cells, and after 24 hours plated cells in soft agar. Colonies were counted after 12 days. We show in Figure 26 that inhibition of miR-221 and miR-222 by SEQ ID #3225 can reduce the amount of colonies growing in soft agar compared to the scrambled control LNA-antimiR, indicating decrease of tumour cells.

[0335] **Conclusion:** The 7-mer (SEQ ID #3225) targeting the miR-221/222 family reduces the number of colonies in soft agar, indicating proliferation arrest of PC3 cells.

**Materials and Methods:**

[0336] Cell line: The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714). PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

[0337] Transfection: 250.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with 25 nM of different LNA oligonucleotides with Lipofectamine2000.

[0338] Clonogenic growth in soft agar:  $2.5 \times 10^3$  PC3 cells were seeded in 0.35% agar on the top of a base layer containing 0.5% agar. Cells were plated 24 hours after transfection. Plates were incubated in at 37°C, 5%CO<sub>2</sub> in a humidified incubator for 12 days and stained with 0.005% crystal violet for 1 h, after which cells were counted. The assay was performed in triplicate.

**Example 28: Assessment of let-7 antagonism by 6-9-mer LNA-antimiRs in Huh-7 cells transfected with let-7a precursor miRNA, and a luciferase sensor assay.**

[0339] In order to assess the efficiency of fully LNA-modified 6-9-mer oligonucleotides in targeting and antagonizing the let-7 family of miRNAs, a luciferase sensor construct was made, containing some 800 bp of the HMGA2 3'UTR. The sequence cloned into the vector contains four out of seven functional let-7 binding sites (sites 2-5), as previously demonstrated by Mayr et al. (Science, 2007) and Lee and Dutta (Genes Dev, 2007). In order to monitor let-7 inhibition, the hepatocellular carcinoma cell line Huh-7 (with low to non-existing levels of endogenous let-7) was transfected with the luciferase sensor construct, with let-7a precursor miRNA, and with the 6-9 mer let-7 antagonists SEQ ID #3232, -3233, -3227, -3234, -3235; see Figure 27) at increasing concentrations. The 6-9-mer LNA-antimiRs were compared with SEQ ID #3226, a 15-mer against let-7a as a positive control. After 24 hours, luciferase activity was measured. **Results:** As seen in Figure 28, the fully LNA-modified 8- and 9-mer LNA-antimiRs (SEQ ID #3227, SEQ ID #3234, and SEQ ID #3235) show similar potencies in de-repressing the let-7 targets in the luciferase sensor assay, as the positive control 15-mer SEQ ID #3226. Full target de-repression for these highly potent compounds is achieved already at 1-5 nM, whereas the 7-mer SEQ ID #3233 needs to be present at slightly higher concentrations (10 nM) to generate the same effect. However, the 6-mer SEQ ID #3232 shows no effect even at as high concentrations as 50 nM. The de-repression of luciferase activity by the 7-9- and the 15-mer LNA-antimiRs is dose-dependent, which is particularly clear in the case of the slightly less potent SEQ ID #3233.

[0340] **Conclusion:** To conclude, the 8-9-mer LNA-antimiRs (SEQ ID #3227, SEQ ID #3234, and SEQ ID #3235) show equal antagonist potencies in inhibition of let-7a *in vitro* compared to the 15-mer LNA-antimiR SEQ ID #3226 targeting let-7a. A potent effect, albeit at slightly higher concentrations is also seen for the 7-mer SEQ ID #3233, whereas a 6-mer has no effect at tested concentrations.

**Materials and Methods:**

[0341] Cell line: The hepatocellular carcinoma cell line Huh-7 was a kind gift from R. Bartinschlager (Dept Mol Virology, University of Heidelberg). Huh-7 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

[0342] Transfection: 8,000 cells were seeded per well in a 96-well plate the day before transfection in order to receive 60-80% confluency the next day. On the day of transfection, Huh-7 cells in each well were transfected with 20 ng HMGA2 3'UTR/psiCHECK2 plasmid, let-7a precursor miRNA (Dharmacon; 10 nM end-concentration), LNA-antimiRs SEQ ID #3232, -3233, -3227, - 3234, -3235, -3226; 0-50 nM end concentrations) together with 0.17 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

[0343] Luciferase assay: Growth media was discarded and 30 µl 1x Passive Lysis Buffer (Promega) was added to

each well. After 15-30 minutes of incubation on an orbital shaker, renilla and firefly luciferase measurements were performed according to manufacturers instructions (Promega).

**Example 29: Assessment of entire let-7 family antagonism by 8-, and 15-mer LNA-antimiRs in Huh-7 cells transfected with a luciferase sensor assay.**

**[0344]** In order to assess the efficiency of a fully LNA-modified 8-mer oligonucleotide in antagonizing the entire let-7 family of miRNAs, the same luciferase sensor construct as described in the previous example was used. Again, Huh-7 cells (with low to non-existing levels of endogenous let-7) were transfected with the sensor construct, with one of the family-representative let-7a, let-7d, let-7e, or let-7i precursors, and with the antagonist SEQ ID #3227 at increasing concentrations. The 8-mer LNA-antimiR was compared to SEQ ID #3226, a 15-mer against let-7a as a positive and potent control. After 24 hours, luciferase activity was measured.

**[0345] Results:** As seen in Figure 29 the fully LNA-modified 8-mer LNA-antimiRs (SEQ ID #3227) show similar potencies in de-repressing the various let-7 targets in the luciferase sensor assay, as the positive control 15-mer SEQ ID #3226. Nearly full target de-repression for the 8-mer is achieved already at 0.5-1 nM, except in the case with let-7e premiR (Fig. 29C), to which only 7 out of 8 nucleotides of SEQ ID #3227 hybridizes to the target. However, despite the terminal mismatch in this case, SEQ ID #3227 generates full target de-repression at 5 nM. The positive control 15-mer shows potent antagonism of all precursors and gives nearly full de-repression at 0.5 nM. The de-repression of luciferase activity by both the 8- and the 15-mer LNA-antimiRs is clearly dose-dependent, as seen in all four panels (Fig 29A-D).

**[0346] Conclusion:** *To conclude*, the 8-mer LNA-antimiR (SEQ ID #3227), is a potent antagonist against four representative let-7 family members *in vitro*, and thus likely against the entire family. Compared to a 15-mer positive control antagonist, SEQ ID #3226, the 8-mer is equally potent for three of four targets, and slightly less potent for the fourth target, let-7e, explained by a terminal mismatch in this case.

**Materials and Methods:**

**[0347] Cell line:** The hepatocellular carcinoma cell line Huh-7 was a kind gift from R. Bartinschlager (Dept Mol Virology, University of Heidelberg). Huh-7 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0348] Transfection:** 8,000 cells were seeded per well in a 96-well plate the day before transfection in order to receive 60-80% confluency the next day. On the day of transfection, Huh-7 cells in each well were transfected with 20 ng HMGA2 3'UTR/psiCHECK2 plasmid, with let-7a, -7d, -7e, or -7i precursor miRNA (Dharmacon; 10 nM end-concentration), and with LNA-antimiRs SEQ ID #3227 and SEQ ID #3226; 0-50 nM end concentrations) together with 0.17 µl Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

**[0349] Luciferase assay:** Growth medium was discarded and 30 µl 1x Passive Lysis Buffer (Promega) was added to each well. After 15-30 minutes of incubation on an orbital shaker, renilla and firefly luciferase measurements were performed according to manufacturer's instructions (Promega).

**Example 30. Assessment of endogenous let-7 antagonism by SEQ ID #3227, an 8-mer LNA-antimiRs, in HeLa cells transfected with a luciferase sensor assay.**

**[0350]** In order to determine the efficiency of a fully LNA-modified 8-mer oligonucleotide in targeting and antagonizing endogenous let-7, the same luciferase sensor construct as described in previous two examples, was co-transfected with SEQ ID #3227 into the cervical cancer cell line HeLa (that expresses moderate to high levels of let-7 as determined by Q-PCR; data not shown). Empty psiCHECK-2 vector was included as a negative control.

**[0351] Results:** As seen in Figure 30, the fully LNA-modified 8-mer LNA-antimiR SEQ ID #3227 shows potent antagonism of endogenous let-7, and gives full target de-repression at concentrations of 5-10 nM. The de-repression of luciferase activity is dose-dependent, starting around 1 nM and reaching a plateau at approximately 10 nM.

**[0352] Conclusion:** *To conclude*, the 8-mer LNA-antimiR (SEQ ID #3227), is a potent antagonist against also endogenous let-7 *in vitro*, and thus provides definite evidence that entire miRNA families can be successfully targeted by short and fully LNA-modified antagonists.

**Materials and Methods:**

**[0353] Cell line:** The cervical cancer cell line HeLa was purchased from ATCC (#CCL-2™). HeLa cells were cultured in Eagle's MEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax, 1x NEAA and 25 ug/ml Gentamicin.

**[0354] Transfection:** 8,000 cells were seeded per well in a 96-well plate the day before transfection in order to receive

50-70% confluency the next day. On the day of transfection, HeLa cells in each well were co-transfected with 20 ng HMGA2 3'UTR/psiCHECK2 plasmid or psiCHECK-2 (empty vector), and with LNA-antimiR SEQ ID #3227 (0-50 nM, end concentrations) together with 0.17  $\mu$ l Lipofectamine2000 (Invitrogen) according to manufacturer's instructions. After 24 hours, cells were harvested for luciferase measurements.

5 **[0355] Luciferase assay:** Growth media was discarded and 30  $\mu$ l 1x Passive Lysis Buffer (Promega) was added to each well. After 15-30 minutes of incubation on an orbital shaker, renilla and firefly luciferase measurements were performed according to manufacturer's instructions (Promega).

10 **Example 31. Assessment of miR-21 antagonism by an 8-mer LNA-antimiR-21 (#3205) versus an 8-mer (#3219) scrambled control LNA in the human colon carcinoma cell line HCT116.**

15 **[0356]** We have previously shown in this application, that an 8-mer LNA-antimiR that is fully LNA-modified and phosphorothiolated effectively antagonizes miR-21 in the human cervix carcinoma cell line HeLa, the human breast carcinoma cell line MCF-7, the human prostate cancer cell line PC3 and human hepatocellular carcinoma HepG2 cell line. We extended this screening approach to the human colon carcinoma cell line HCT116. To assess the efficiency of the 8-mer LNA-antimiR oligonucleotide against miR-21, luciferase reporter constructs were generated in which a perfect match target site for the mature miR-21 was cloned into the 3'UTR of the Renilla luciferase gene. In order to monitor miR-21 inhibition, HCT116 cells were transfected with the luciferase constructs together with the miR-21 antagonist #3205 (8-mer) and for comparison of specificity with the 8-mer LNA scrambled control (#3219). After 24 hours, luciferase activity was measured.

20 **[0357] Results:**The luciferase reporter experiments showed a dose-dependent de-repression of the luciferase miR-21 reporter activity with the 8-mer LNA-antimiR against miR-21 (#3205) and complete de-repression was obtained at 5 nM (Figure 31). When comparing the specificity of the 8-mer perfect match and the 8-mer scrambled control, the scrambled control LNA-antimiR (#3219) did not show any de-repression at all, demonstrating high specificity of the LNA-antimiR compound against miR-21.

25 **[0358] Conclusion:** The 8-mer (#3205) is potent in targeting miR-21 and antagonism of miR-21 by #3205 is specific.

**Materials and Methods:**

30 **[0359] Cell line:** The human colon carcinoma HCT116 cell line was purchased from ATCC (CCL-247). HCT116 cells were cultured in RPMI medium, supplemented with 10% fetal bovine serum, and 25  $\mu$ g/ml Gentamicin.

35 **[0360] Transfection:** 110,000 cells were seeded per well in a 12-well plate and transfection was performed. HCT116 cells were transfected with 0.3  $\mu$ g miR-21 luciferase sensor plasmid or empty psiCHECK2 vector together with 1.2  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instruction. Transfected were varying concentration of LNA-antimiR and control oligonucleotides. After 24 hours, cells were harvested for luciferase measurements.

40 **[0361] Luciferase assay:** The cells were washed with PBS and 250  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to the wells. The plates were placed on a shaker for 30 min., after which the cell lysates were transferred to eppendorf tubes. The cell lysate was centrifugated for 10 min at 2,500 rpm after which 50  $\mu$ l were transferred to a 96 well plate and luciferase measurements were performed according to the manufacturer's instructions (Promega).

**Example 32. Knock-down of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation of PC3 cells.**

45 **[0362]** A hallmark of cellular transformation is the ability for tumour cells to grow in an anchorage-independent way in semisolid medium. We therefore performed soft agar assay which is a phenotypic assay that is relevant for cancer, given that it measures the decrease of tumour cells. We transfected #3205 (perfect match LNA-antimiR-21) and #3219 (LNA scrambled control) into PC3 cells, and after 24 hours plated cells in soft agar. Colonies were counted after 12 days. We show in Figure 32 that inhibition of miR-21 by #3205 can reduce the amount of colonies growing in soft agar compared to the scrambled control LNA treated or untreated control (transfected, but with no LNA), demonstrating decrease of tumour cells.

50 **[0363] Conclusion:** The 8-mer (#3205) targeting the miR-21 family reduces the number of colonies in soft agar, demonstrating proliferation arrest of PC3 cells.

**Materials and Methods:**

55 **[0364] Cell line:** The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714). PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25  $\mu$ g/ml Gentamicin.

**[0365] Transfection:** 250,000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, PC3 cells were transfected with 25 nM of different LNA

oligonucleotides with Lipofectamine2000.

[0366] Clonogenic growth in soft agar:  $2.5 \times 10^3$  PC3 cells were seeded in 0.35% agar on the top of a base layer containing 0.5% agar. Cells were plated 24 hours after transfection. Plates were incubated in at 37°C, 5% CO<sub>2</sub> in a humidified incubator for 12 days and stained with 0.005% crystal violet for 1 h, after which cells were counted. The assay was performed in triplicate.

[0367] **Example 33. Silencing of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation of HepG2 cells.** miR-21 is overexpressed in the human hepatocellular carcinoma cell line HepG2 and we have previously shown that we are able to regulate the luciferase activity of a miR-21 sensor plasmid with #3205 in these cells. HepG2 cells were transfected with #3205 and #3219 (scrambled 8-mer), and after 24 hours plated into soft agar. Colonies were counted after 17 days with a microscope.

[0368] **Results:** We show in Figure 33 that inhibition of miR-21 by #3205 can reduce the amount of colonies growing in soft agar, showing that proliferation arrest has occurred. In addition, our scrambled 8-mer control, #3219, had no significant effect on the number of colonies. **Conclusion:** The 8-mer (#3205) targeting the miR-21 reduces the number of colonies in soft agar, indicating proliferation arrest of HepG2 cells.

#### Materials and Methods:

[0369] Cell line: The human hepatocytic HepG2 cell line was purchased from ECACC (#85011430). HepG2 cells were cultured in EMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

[0370] Transfection: 650.000 cells were seeded per well in a 6-well plate and reverse transfection was performed. HepG2 cells were transfected with 0.6 µg miR-21 luciferase sensor plasmid or empty psiCHECK2 vector together with 2,55 µl Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected were also LNA-antimiR and control oligonucleotides as varying concentrations. After 24 hours, the cells were harvested for luciferase measurements.

[0371] Clonogenic growth in soft agar:  $2.0 \times 10^3$  HepG2 cells were seeded in 0.35% agar on the top of a base layer containing 0.5% agar. Cells were plated 24 hours after transfection. Plates were incubated in at 37°C, 5% CO<sub>2</sub> in a humidified incubator for 17 days and stained with 0.005% crystal violet for 1 h, after which cells were counted. The assay was performed in triplicate.

#### Example 34. Silencing of miR-21 by the 8-mer #3205 LNA-antimiR inhibits cell migration in PC3 cells.

[0372] Cell migration can be monitored by performing a wound healing assay (=scratch assay) where a "scratch" is made in a cell monolayer, and images are captured at the beginning and at regular intervals during cell migration. By comparing the images, quantification of the migration rate of the cells can be determined. This was done in the human prostate cancer cell line PC3. Cells were seeded, and on day 3 the cells were transfected, and the next day, when 100% confluency was reached, a scratch (=wound) was made. When the scratch was made, pictures were taken in order to document the initial wound. Afterwards the area of the wound closure is measured at different time points with the free software program Image J. As shown in Figure 34A, PC3 cells had been treated with 25 nM #3205 (perfect match, miR-21), the control #3219 or left untransfected. Pictures were taken after 24 hours, and the area was calculated for the wound closure at respective time-point. The wound closure for the untransfected cells and for the control, #3219, was faster as compared to our LNA-antimiR against miR-21, #3205, indicating that #3205 inhibits miR-21 and prevents the cells from migrating (Figure 34B).

[0373] **Conclusion:** The 8-mer (#3205) targeting miR-21 inhibits the cell migration of PC3 cells compared to untransfected and control transfected cells.

#### Materials and Methods:

[0374] Cell line: The human prostate carcinoma PC3 cell line was purchased from ECACC (#90112714). PC3 cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 2 mM Glutamax and 25 ug/ml Gentamicin.

[0375] Scratch assay: 150.000 cells were seeded per well in a 6-well plate three days before transfection in order to receive 100% confluency the next day. At 24 hours after transfection, a scratch was made in the cell monolayer with a 200 µl tip. Pictures were taken at 0 h and after 24 hours by using a digital camera coupled to a microscope. The software program Image J was used to determine wound closure.

#### Example 35. Length assessment of fully LNA-substituted LNA-antimiRs antagonizing miR-155.

[0376] We have previously shown a length assessment for miR-21 regarding fully LNA-substituted LNA-antimiRs, and showed that the most potent LNA-antimiRs are 7-, 8- or 9 nt in length. The same experiment was repeated with miR-

155. A perfect match target site for miR-155 was cloned into the 3'UTR of the luciferase gene in the reporter plasmid psiCHECK2 and transfected into the mouse RAW macrophage cell line together with fully LNA-substituted LNA-antimiRs of different lengths. Because the endogenous levels of miR-155 are low in the RAW cell line, the cells were treated with 100 ng/ml LPS for 24 hours in order to induce miR-155 accumulation. After 24 hours, luciferase analysis was performed.

5 [0377] **Results:** As shown in Figure 35, the most potent LNA-antimiRs are #3207(8 nt) and #3241 (9 nt), reaching almost a 80% de-repression at only 0.25 nM LNA concentration. The 6-mer (#3244) shows no significant de-repression. Increasing the length to 12-mer to 14-mer (#3242 and #3243) decreased the potency as shown by less efficient de-repression of the miR-155 reporter.

10 [0378] **Conclusion:** The most potent fully LNA-substituted LNA-antimiRs targeting miR-155 were an 8- and 9-mer (#3207and #3241).

#### Materials and Methods:

15 [0379] **Cell line:** The mouse macrophage RAW 264.7 cell line was purchased from ATCC (TIB-71). RAW cells were cultured in DMEM medium, supplemented with 10% fetal bovine serum, 4 mM Glutamax and 25 ug/ml Gentamicin.

20 [0380] **Transfection:** 500.000 cells were seeded per well in a 6-well plate the day before transfection in order to receive 50% confluency the next day. On the day of transfection, RAW 264.7 cells were transfected with 0.3 ug miR-155 perfect match/psiCHECK2 or empty psiCHECK2 vector together with 10  $\mu$ l Lipofectamine2000 (Invitrogen) according to the manufacturer's instructions. Transfected was also varying concentrations of LNA-antimiRs. In order to induce miR-155 accumulation, LPS (100 ng/ml) was added to the RAW cells after the 4 hour incubation with the transfection complexes. After another 24 hours, cells were harvested for luciferase measurements.

25 [0381] **Luciferase assay:** The cells were washed with PBS and harvested with cell scraper, after which cells were spinned for 5 min at 2.500 rpm. The supernatant was discarded and 50  $\mu$ l 1 x Passive Lysis Buffer (Promega) was added to the cell pellet, after which cells were put on ice for 30 min. The lysed cells were spinned at 10.000 rpm for 30 min after which 20  $\mu$ l were transferred to a 96-well plate and luciferase measurements were performed according to the manufacturer's instructions (Promega).

#### Example 36. Plasma protein binding for the fully LNA-substituted 8-mer #3205 targeting miR-21 (LNA-antimiR-21).

30 [0382] The plasma proteins are not saturated with #3205 at the plasma concentrations in the experiment shown in Figure 36A. In a wide range of #3205 concentrations in the plasma the protein binding is around 95% of the #3205 LNA-antimiR-21 in Figure 36B. At #3205 concentrations 50.1  $\mu$ M (174  $\mu$ g/mL) the binding capacity of plasma proteins for FAM-labeled #3205 has not been saturated.

35 [0383] **Materials and Methods:** Mouse plasma (100  $\mu$ L) was spiked with FAM-labeled #3205 to 0.167, 1.67, 5.01, 10.02, 16.7, 25.05 and 50.1  $\mu$ M concentrations. The solutions were incubated at 37°C for 30 minutes. The solutions were transferred to a Microcon Ultracel YM-30 filter (regenerated cellulose 30.000 MWCO). The filters were spun for 20 minutes at 2000g and at room temperature in a microcentrifuge. The filtrate was diluted 5, 10 and 20 times and 100 $\mu$ L samples were transferred to a microtiter plate (Polystyrene Black NUNC-237108). The fluorescence was detected using a FLUOstar Optima elisa reader with excitation 458 nm and emission 520 nm. The amount of unbound FAM-labeled #3205 was calculated from a standard curve derived from filtrated plasma spiked with FAM-labeled #3205 at 12 different (0.45 -1000 nM) concentrations. The numbers were corrected with the recovery number established from filtration experiments with #3205 concentrations 0.167, 1.67, 5.01, 10.02, 16.7, 25.05 and 50.1  $\mu$ M in filtrated plasma. The recovery of FAM-labeled #3205 was 86%.

#### 45 Example 37. Quantitative whole body autoradiography study in female pigmented mice after single intravenous administration of <sup>35</sup>S-labelled #3205 LNA-antimiR-21.

[0384] In order to determine the biodistribution of a short fully LNA-modified LNA-antimiR (#3205, 8-mer) a whole body tissue distribution of radioactively labeled compound was done in mice. <sup>35</sup>S-labelled #3205 was dosed to mice with a single intravenous administration and mice were sacrificed at different time-points, ranging from 5 min to 21 days.

55 **Table 6(i).** Individual tissue concentrations ( $\mu$ g #3205/g tissue) after a single intravenous administration of <sup>35</sup>S-labelled #3205 in female pigmented mice. The figures are mean values of three measurements for each tissue and ratio. The coefficient of variation (CV) is generally about 10%.

Tissue	Max. Conc. of oligo $\mu$ g #3205/g tissue	Time of max conc. hours	T <sub>1/2</sub> hours
Adrenal gl.	13,6	0,083	374

(continued)

	Tissue	Max. Conc. of oligo $\mu\text{g}$ #3205/g tissue	Time of max conc. hours	T $\frac{1}{2}$ hours
5	Bile	4	1	
	Bone marrow	7,2	0,083	411
	Brain	0,4	0,083	
10	Brown fat	8,8	0,083	
	Gastric muc.	10,1	0,083	
	Heart blood	26,2	0,083	10.3
	Kidney ctx.	58,7	24	104
15	Liver	11,8	0,083	588
		10,7	24	
	Lung	13,2	0,083	289
20	Lymph node	5	0,083	262
		2,4	48	
	lymph	18,8	4	
		20,8	168	
25	Myocardium	8,1	0,083	662
	Ovary	13	0,083	198
	Pancreas	5	0,083	
30	Pituitary gl.	6,7	0,083	
	Salivary gl.	8,6	0,083	405
		5,5	168	
	skel. Muscle	4,8	0,083	
35	Skin pig.	5,4	0,25	
	Spleen	9,8	0,083	564
	Thymus	3,8	0,083	185
40	Thyroid gl.	10,9	0,083	592
	Urine	328,9	0,083	
	Uterus	9,6	0,25	177
45	Uvea of the eye	13,6	0,083	
	LOQ	0,045	0,083	
		0,033	24	
		0,03	168	

50

**Table 6(ii)** Tissue to liver ratios after single intravenous administration of  $^{35}\text{S}$ -labelled #3205 in female pigmented mice.

$^{35}\text{S}$ -#3205										
55	Animal no	10	11	12	13	14	15	16	17	18
	Surv. Time (h)	0,083	0,25	1h	4h	24h	48h	96h	168	504



(continued)

<sup>35</sup> S-#3205										
Organ										
5	Adrenal gl	liver	liver	liver	liver	liver	liver	liver	liver	liver
	Bile	1,15	1,08	0,52	0,27	0,24	0,26	0,23	0,18	0,17
10	Bone marrow	0,03	0,11	0,55	0,10	0,03	0,07	0,04	0,03	0,04
	Brain	0,61	0,81	0,55	0,45	0,40	0,48	0,43	0,42	0,34
	Brown fat	0,03	0,03	0,01	0,00	0,00	0,00	0,00	0,00	0,00
15	Gastric muc	0,75	0,57	0,29	0,12	0,07	0,12	0,08	0,10	0,07
	Heart blood	0,86	0,71	0,31	0,22	0,10	0,21	0,15	0,16	0,12
	Kidney ctx	2,23	1,91	0,74	0,11	0,01	0,00	0,00	0,00	0,00
	Liver	2,87	3,94	6,45	6,95	5,51	6,68	3,92	2,24	0,40
20	Lung	1,00	1,00	1,00	1,00	1,00	1,00	1,00	1,00	1,00
	Lymph node	1,12	0,97	0,63	0,09	0,04	0,04	0,03	0,02	0,02
	Lymph	0,43	0,30	0,25	0,19	0,11	0,32	0,20	0,17	0,12
	Myocardium	0,82	1,09	1,78	2,78	1,03	2,05	1,62	3,17	1,89
25	Ovary	0,69	0,63	0,30	0,13	0,10	0,15	0,09	0,11	0,12
	Pancreas	1,10	1,40	0,61	0,31	0,27	0,28	0,21	0,21	0,08
	Pituitary gland	0,42	0,37	0,22	0,18	0,12	0,17	0,12	0,15	0,11
	Salivary gland	0,57	0,54	0,28	0,11	0,15	0,16	0,12	0,10	0,08
30	Skel. muscle	0,73	0,81	0,38	0,25	0,25	0,42	0,23	0,85	0,24
	Skin, pigm.	0,40	0,28	0,14	0,04	0,02	0,04	0,03	0,03	0,03
	Spleen	0,34	0,69	0,65	0,36	0,20	0,26	0,20	0,19	0,13
35	Thymus	0,83	0,86	0,44	0,32	0,24	0,34	0,35	0,29	0,31
	Thyroid gland	0,32	0,31	0,14	0,07	0,09	0,08	0,05	0,04	0,02
	Urine	0,9	1,2	0,43	0,28	0,25	0,34	0,19	0,26	0,25
40	Uterus	27,96	39,48	9,90	5,44	0,24	0,39	0,12	0,15	0,03
	Uvea of the eye	0,56	1,23	0,65	0,30	0,30	0,07	0,27	0,16	0,08

45 **[0385] Conclusions:** #3205 shows blood clearance of radioactivity with elimination half-lives of 8-10 hours. High levels of radioactivity were registered in the kidney cortex, lymph, liver, bone marrow, spleen, ovary and uterus. The highest level of radioactivity was registered in the kidney cortex showing five times higher levels than that of the liver for #3205. A strong retention of radioactivity was noticed in the kidney cortex, lymph, liver, bone marrow and spleen for #3205 LNA-antimiR-21.

50 **Materials and Methods:**

55 **[0386] Dose administration:** All mice were weighed before administration. Nine female mice were given 10 mg/kg of <sup>35</sup>S-#3205 intravenously in a tail vein. The volume given to each animal was 10 mL/kg of the test formulation. The specific activity 75.7 µCi/mg. Individual mice were killed 5 min, 15 min, 1 hour, 4 hours, 24 hours, 2 days, 4 days, 7 days and 21 days after administration of #3205. Whole body autoradiography: The mice were anaesthetized by sevoflurane, and then immediately immersed in heptane, cooled with dry ice to -80°C, ABR-SOP-0130. The frozen carcasses were embedded in a gel of aqueous carboxymethyl cellulose (CMC), frozen in ethanol, cooled with dry ice (-80°C) and sectioned sagittally for whole body autoradiography, according to the standard method, ABR-SOP-0131. From each animal 20 µm

sections were cut at different levels with a cryomicrotome (Leica CM 3600) at a temperature of about -20°C. The obtained sections were caught on tape (Minnesota Mining and Manufacturing Co., No. 810) and numbered consecutively with radioactive ink. After being freeze-dried at -20°C for about 24 hours, selected sections were covered with a thin layer of mylar foil, and put on imaging plates (Fuji, Japan). Exposure took place in light tight cassettes in a lead shielding box at -20°C, to protect the image plates from environmental radiation. After exposure the imaging plates were scanned at a pixel size of 50 µm and analyzed by radioluminography using a bioimaging analysis system (Bas 2500, Fuji, Japan), and described in ABR-SOP-0214. A water-soluble standard test solution of <sup>35</sup>S radioactivity was mixed with whole blood and used for production of a calibration scale, ABR-SOP-0251. However, the different blood standards were dissolved in 500 µL Soluene-35. 4.5 mL Ultima Gold was then added to the dissolved samples. As <sup>35</sup>S and <sup>14</sup>C have very similar energy spectra, a standard <sup>14</sup>C-programme (Packard 2200CA) was used when the radioactivity for the different blood samples was settled.

**[0387] Pharmacokinetic calculations:** The <sup>35</sup>S radioactivity measured in whole blood and tissues was expressed as nCi/g tissue and recalculated to nmol equiv/g tissue for the pharmacokinetic evaluation. The pharmacokinetic parameters C<sub>max</sub>, t<sub>1/2</sub> and AUC were determined for the whole blood and tissues by non-compartmental analysis using WinNonlin Professional (Pharsight Corporation, Mountain View, CA, USA). After intravenous administration, the concentration was extrapolated back to zero and expressed as (C<sub>0</sub>). The elimination rate constant λ was estimated by linear regression analysis of the terminal slope of the logarithmic plasma concentration-time curve. The elimination half-life, t<sub>1/2</sub>, was calculated using the equation, t<sub>1/2</sub> = ln2/λ. The last three time-points above LOQ were used in the elimination half-life calculations, if not stated otherwise.

#### **Example 38. Assessment of let-7 inhibition *in vivo* by an 8-mer LNA-antimiR, as determined through Ras protein quantification in mouse lung and kidney**

**[0388]** In order to investigate the possibility to antagonize the abundantly expressed let-7 family *in vivo*, mice were intravenously (i.v.) injected with an 8-mer LNA-antimiR antagonist or with saline. To measure treatment effect, proteins were isolated from lungs and kidneys. Because the Ras family of proteins (N-Ras, K-Ras, and H-Ras), in particular N-Ras and K-Ras, has previously been shown to be regulated (repressed) by the let-7 family by Johnson et al. (Cell, 2005), the aim was to analyze whether these let-7 targets could be de-repressed *in vivo*.

**[0389] Results:** As seen in Figure 37, the 8-mer LNA-antimiR potently de-repressed Ras protein levels in the kidneys of treated mice, normalized against saline controls. The up-regulation in this organ was more than 3-fold, showing a clear *in vivo* effect. In the lungs, however, only a minimal (1.2-fold) Ras de-repression was observed (Fig 1B), suggesting that insufficient amounts of LNA-antimiR has entered this organ in order to inhibit its massive amounts of let-7, as previously described by Johnson et al. (Cancer Research, 2007).

**[0390] Conclusion:** The 8-mer LNA-antimiR shows a clear effect in regulating target let-7 miRNA *in vivo*, as evaluated based on Ras protein levels in treated vs. control mice. Whereas the effect seems to be smaller in lungs, Ras levels in the kidney show a substantial up-regulation upon antimiRs-treatment.

**[0391] Materials and Methods: Animals and dosing:** C57BL/6 female mice were treated with 10 mg/kg LNA-antimiR or saline for three consecutive days (0, 1, and 2) and sacrificed on day 4. Tissue samples from lungs and kidneys were snapfrozen and stored at -80°C until further processing.

**[0392] Western blot analysis:** Lung and kidney proteins from saline and LNA-antimiR-treated mice were separated on NuPAGE Bis Tris 4-12% (Invitrogen), using 100 µg per sample. The proteins were transferred to a nitrocellulose membrane using iBlot (Invitrogen) according to the manufacturer's instructions. Blocking, antibody dilution and detection was performed according to the manufacturer's specifications. For Ras detection, a primary rabbit-anti Ras antibody (SC-3339, Santa Cruz Biotechnology) and a secondary HRP-conjugated swine-anti-rabbit antibody (P0399, Dako) was used, and for tubulin detection, a primary tubulin alpha (MS-581-P1, Neomarkers) and a secondary HRP-conjugated goat-anti-mouse antibody (P0447, Dako) was used.

#### **Example 40. *In vivo* efficacy assessment of the 8-mer LNA-antimiR (#3205) in targeting miR-21, as determined by Pdcd4 protein up-regulation in mouse kidney.**

**[0393]** We have shown that an 8-mer LNA-antimiR that is fully LNA-modified antagonizes miR-21 and has the ability to regulate the protein levels of the miR-21 target Pdcd4 *in vitro*. We therefore injected the LNA-antimiR into mice to determine the effects of the LNA-antimiR *in vivo*. The mice received 25 mg/kg of #3205 by i.v. injection every other day for 14 days (a total of 5 doses). The mice were sacrificed on day 14, the kidney was removed, and protein was isolated. In order to determine target regulation, Western blot analysis was performed.

**[0394] Results:** As shown in Figure 38, treating mice with #3205 showed significantly increased Pdcd4 protein levels as compared to the saline control. While the normalized Pdcd4 versus Gapdh ratio was consistent in both saline samples, the protein up-regulation in the two LNA-antimiR-treated (#32059 mice were measured to 3.3- and 6.3-fold, respectively,

demonstrating an *in vivo* pharmacological effect of the #3205 8-mer LNA-antimiR.

**[0395]** Conclusion: The fully LNA-modified 8-mer LNA-antimiR #3205 antagonizes miR-21 *in vivo*, as demonstrated through its ability to de-repress (up-regulate) mouse kidney levels of Pdc4, a validated miR-21 target.

5 **Materials and Methods:**

**[0396]** *Animals and dosing:* C57BL/6 female mice with average of 20 g body weight at first dosing were used in all experiments and received regular chow diet (Altromin no 1324, Brogaarden, Gentofte, Denmark). Substances were formulated in physiological saline (0.9% NaCl). The animals were dozed with LNA-antimiR or saline (0.9% NaCl), receiving an injection of 25 mg/kg every other day for 14 days, a total of 5 doses. Animals were sacrificed on day 14.

10 **[0397]** *Western blot analysis:* 80  $\mu$ g kidney tissue from saline or LNA-treated mice was separated on NuPAGE Bis Tris 4-12% (Invitrogen). The proteins were transferred to a nitrocellulose membrane using iBlot (Invitrogen) according to the manufacturer's instructions. The membrane was incubated with Pdc4 antibody (Bethyl Laboratories), followed by HRP-conjugated swine-anti-rabbit antibody (Dako). As equal loading control, GAPDH (Abcam) was used, followed by HRP-conjugated swine-anti-mouse antibody. The membranes were visualized by chemiluminescence (ECL, Amer-  
15 sham).

20

25

30

35

40

45

50

55

Table 1

microRNA	MicroRNA Sequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
ebv-miR-BART1-3p	UAGCACCGCUAUCCACUAUGUC	40	AGCGGTGCT	977	GCGGTGCT	1914	CGGTGCT	2851
ebv-miR-BART1-5p	UCUUAGUGGAAGUGACGUGCUGUG	41	TCCACTAAG	978	CCACTAAG	1915	CACTAAG	2852
ebv-miR-BART10	UACAUAAACCAUGGAGUUGGCUGU	42	TGTTATGT	979	GGTTATGT	1916	GTTATGT	2853
ebv-miR-BART10*	GCCACCUCUUUGGUUCUGUACA	43	AAGAGGTGG	980	AGAGGTGG	1917	GAGGTGG	2854
ebv-miR-BART11-3p	ACGCACACCAGGCUGACUGCC	44	TGGTGTGCG	981	GGTGTGCG	1918	GTGTGCG	2855
ebv-miR-HART11-5p	UCAGACAGUUUGGUGCGCUAGUUG	45	AACTGTCTG	982	ACTGTCTG	1919	CTGTCTG	2856
ebv-miR-BART12	UCCUGUGGUUUUGGUGUGGUU	46	CACCACAGG	983	ACCACAGG	1920	CCACAGG	2857
ebv-miR-BART13	UGUAACUUGCCAGGACGGCUGA	47	GCAAGTTAC	984	CAAGTTAC	1921	AAGTTAC	2858
ebv-miR-BART13*	AACCGGCUCGUGGCUCGUACAG	48	CGAGCCGGT	985	GAGCCGGT	1922	AGCCGGT	2859
ebv-miR-BART14	UAAUUGCUGCAGUAGAGGGAU	49	GCAGCATTT	986	CAGCATTT	1923	AGCATTT	2860
ebv-miR-BART14*	UACCCUACGCUGCCGAAUUACA	50	GCGTAGGGT	987	CGTAGGGT	1924	GTAGGGT	2861
ebv-miR-BART15	GUCAGUGUUUUUGUUCCUUGA	51	AACCACTGA	988	ACCACTGA	1925	CCACTGA	2862
ebv-miR-BART16	UUAGAUAGAGUGGGUGUGUCUCU	52	CTCTATCTA	989	TCTATCTA	1926	CTATCTA	2863
ebv-miR-BART17-3p	UGUAUGCCUGGUGUCCCCUUGU	53	CAGGCATAC	990	AGGCATAC	1927	GGCATAC	2864
ebv-miR-BART17-5p	UAAGAGGACGCAGGCAUACAAG	54	CGTCCTCTT	991	GTCCCTTT	1928	TCCTCTT	2865
ebv-miR-BART18-3p	UAUCGGAAGUUUGGCUCUUCGUC	55	ACTTCCGAT	992	CTTCCGAT	1929	TTCCGAT	2866
ebv-miR-BART18-5p	UCAAGUUCGCACUCCUUAUACA	56	GCGAACTTG	993	CGAACTTG	1930	GAACTTG	2867
ebv-miR-BART19-3p	UUUUUUUGCUUGGAAUGCU	57	GCAAACAAA	994	CAAACAAA	1931	AAACAAA	2868
ebv-miR-BART19-5p	ACAUUCCCCGCAACAUACAUG	58	CGGGAAATG	995	GGGAAATG	1932	GGAATG	2869
ebv-miR-BART2-3p	AAGGAGCGAUUUGGAGAAAAUAAA	59	ATCGCTCCT	996	TCGCTCCT	1933	CGCTCCT	2870
ebv-miR-BART2-5p	UAUUUCUGCAUUCGCCCUUGC	60	GCAGAAAAT	997	CAGAAAAT	1934	AGAAAAT	2871
ebv-miR-BART20-3p	CAUGAAGGCACAGCCUUGUACC	61	TGCCTTCAT	998	GCCCTCAT	1935	CCTTCAT	2872
ebv-miR-BART20-5p	UAGCAGGCAUGUCUCAUCC	62	ATGCCTGCT	999	TGCCTGCT	1936	GCCTGCT	2873
ebv-miR-BART3	CGCACCAUAGUCACCAGGUGU	63	TAGTGGTGC	1000	AGTGGTGC	1937	GTGGTGC	2874

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
ebv-miR-BART3*	ACCUAGUGUUAGUGUUGUGCU	64	AACACTAGG	1001	ACACTAGG	1938	CACTAGG	2875
ebv-miR-BART4	GACCUGAUGCUCUGGGUGUGCU	65	GCATCAGGT	1002	CATCAGGT	1939	ATCAGGT	2876
ebv-miR-BART5	CAAGGUGAAUAGCUGCCCAUCG	66	ATTCACCTT	1003	TTCACCTT	1940	TCACCTT	2877
ebv-miR-BART6-3p	CGGGGAUCGGACUAGCCUUAGA	67	CCGATCCCC	1004	CGATCCCC	1941	GATCCCC	2878
ebv-miR-BART6-5p	UAAAGUUUGUCCAAUCCAUAGG	68	ACCAACCTT	1005	CCAACCTT	1942	CAACCTT	2879
ebv-miR-BART7	CAUCAUAGUCCAGUGUCCAGGG	69	GACTATGAT	1006	ACTATGAT	1943	CTATGAT	2880
ebv-miR-BART7*	CCUGGACCUUGACUAUGAAACA	70	AAGGTCCAG	1007	AGGTCCAG	1944	GGTCCAG	2881
ebv-miR-BART8	UACGGUUUCCUAGAUUGUACAG	71	GGAACCCGT	1008	GAACCCGT	1945	AAACCCGT	2882
ebv-miR-BART8*	GUCACAUCUAGGGGUCGUAGA	72	AGATTGTGA	1009	GATTGTGA	1946	ATTGTGA	2883
ebv-miR-BART9	UAACACUUAUGGGUCCCGUAGU	73	TGAAGTGTT	1010	GAAGTGTT	1947	AAGTGTT	2884
ebv-miR-BART9*	UACUGGACCCUGAAUUGGAAAC	74	GGGTCCAGT	1011	GGTCCAGT	1948	GTCCAGT	2885
ebv-miR-BHRF1-1	UAACCUGAUCAGCCCCGGAGUU	75	GATCAGGTT	1012	ATCAGGTT	1949	TCAGGTT	2886
ebv-miR-BHRF1-2	UAUCUUUUGCGGCAGAAUUUGA	76	GCAAAAGAT	1013	CAAAAGAT	1950	AAAAGAT	2887
ebv-miR-BHRF1-2*	AAAUUCUGUUGCAGCAGAUAGC	77	AACAGAATT	1014	ACAGAATT	1951	CAGAATT	2888
ebv-miR-BHRF1-3	UAACGGGAAGUGUGUAAGCACA	78	CTTCCCGTT	1015	TTCCCGTT	1952	TCCCGTT	2889
hcmv-miR-UL112	AAGUGACGGUGAGAUCCAGGCU	79	ACCGTCACT	1016	CCGTCACT	1953	CGTCACT	2890
hcmv-miR-UL148D	UCGUCCUCCCCUUCUUCACCCG	80	GGGAGGACG	1017	GGAGGACG	1954	GAGGACG	2891
hcmv-miR-UL22A	UAACUAGCCUUCUCCGUGAGA	81	AGGCTAGTT	1018	GGCTAGTT	1955	GCTAGTT	2892
hcmv-miR-UL22A*	UCACCAGAAUGCUAGUUUGUAG	82	ATTCTGGTG	1019	TTCTGGTG	1956	TCTGGTG	2893
hcmv-miR-UL36	UCGUUGAAGACACCCUGGAAAGA	83	TCTCAACG	1020	CTTCAACG	1957	TTCAACG	2894
hcmv-miR-UL36*	UUUCCAGGUGUUUCAACGUGC	84	CACCTGGAA	1021	ACCTGGAA	1958	CCTGGAA	2895
hcmv-miR-UL70-3p	GGGGAUGGCUUGCGCGCGCGG	85	GCCCATCCC	1022	CCCATCCC	1959	CCATCCC	2896
hcmv-miR-UL70-5p	UGCGUCUCGGCCUUCGUCCAGA	86	CCGAGACGC	1023	CGAGACGC	1960	GAGACGC	2897

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hcmv-miR-US25-1	AACCGCUCAGUGGUCGGACC	87	CTGAGCGGT	1024	TGAGCGGT	1961	GAGCGGT	2898
hcmv-miR-US25-1*	UCCGAACGCUAGGUCGGUUCUC	88	AGCGTTCGG	1025	GCGTTCGG	1962	CGTTCGG	2899
hcmv-miR-US25-2-3p	AUCCACUUGGAGAGCUCGCCGGG	89	CCAAGTGGA	1026	CAAGTGGA	1963	AAGTGGA	2900
hcmv-miR-US25-2-5p	AGCGGUCUGUUCAGGUGGAUGA	90	ACAGACCGC	1027	CAGACCGC	1964	AGACCGC	2901
hcmv-miR-US33-3p	UCACGGUCCGAGCACAUCCA	91	CGGACCGTG	1028	GGACCGTG	1965	GACCGTG	2902
hcmv-miR-US33-5p	GAUUGUCCCCGGACCGUGGGCG	92	GGGCACAAT	1029	GGCACAAT	1966	GCACAAT	2903
hcmv-miR-US4	CGACAUUGGACGUGCAGGGGGAU	93	GTCCATGTC	1030	TCCATGTC	1967	CCATGTC	2904
hcmv-miR-US5-1	UGACAAGCCUGACGAGAGCGU	94	AGGCTTGTC	1031	GGCTTGTC	1968	GCTTGTC	2905
hcmv-miR-US5-2	UUUAUGAUGGUGGACGAUGUC	95	CCTATCATA	1032	CTATCATA	1969	TATCATA	2906
hsa-let-7a	UGAGGUAGUAGGUUGUAUAGUU	96	TACTACCTC	1033	ACTACCTC	1970	CTACCTC	2907
hsa-let-7a*	CUAUACAUCUACUGUCUUUC	97	GATTGTATA	1034	ATTGTATA	1971	TTGTATA	2908
hsa-let-7b	UGAGGUAGUAGGUUGUGUGUU	98	TACTACCTC	1035	ACTACCTC	1972	CTACCTC	2909
hsa-let-7b*	CUAUACAACCUACUGCCUUCCC	99	GGTTGTATA	1036	GTTGTATA	1973	TTGTATA	2910
hsa-let-7c	UGAGGUAGUAGGUUGUAUGUU	100	TACTACCTC	1037	ACTACCTC	1974	CTACCTC	2911
hsa-let-7c*	UAGAGUUACACCCUUGGGAGUUA	101	TGTAACCTC	1038	GTAACCTC	1975	TAACCTC	2912
hsa-let-7d	AGAGGUAGUAGGUUGCAUAGUU	102	TACTACCTC	1039	ACTACCTC	1976	CTACCTC	2913
hsa-let-7d*	CUAUACGACCUCGUCUUUCU	103	GGTCCGTATA	1040	GTCGTATA	1977	TCGTATA	2914
hsa-let-7e	UGAGGUAGGAGGUUGUAUAGUU	104	TCCTACCTC	1041	CCTACCTC	1978	CTACCTC	2915
hsa-let-7e*	CUAUACGGCCUCCUAGCUUUCC	105	GGCCGTATA	1042	GCCGTATA	1979	CCGTATA	2916
hsa-let-7f	UGAGGUAGUAGUUGUAUAGUU	106	TACTACCTC	1043	ACTACCTC	1980	CTACCTC	2917
hsa-let-7f-1*	CUAUACAACUUAUUGCCUUCCC	107	GATTGTATA	1044	ATTGTATA	1981	TTGTATA	2918
hsa-let-7f-2*	CUAUACAGUCUACUGUCUUUCC	108	GACTGTATA	1045	ACTGTATA	1982	CTGTATA	2919

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-let-7g	UGAGGUAGUAGUUUGUACAGUU	109	TACTACCTC	1046	ACTACCTC	1983	CTACCTC	2920
hsa-let-7g*	CUGUACAGGCCACUGCCUUGC	110	GCCTGTACA	1047	CCTGTACA	1984	CTGTACA	2921
hsa-let-7i	UGAGGUAGUAGUUUGUGCUGUU	111	TACTACCTC	1048	ACTACCTC	1985	CTACCTC	2922
hsa-let-7i*	CUGCGAAGCUACUGCCUUGC	112	GCTTGCGCA	1049	CTTGCGCA	1986	TTGCGCA	2923
hsa-miR-1	UGGAUUGUAAAGAAGUAUGUAAU	113	TTACATTCC	1050	TACATTCC	1987	ACATTCC	2924
hsa-miR-100	AACCCGUAGAUCCGAACUUGUG	114	TCTACGGGT	1051	CTACGGGT	1988	TACGGGT	2925
hsa-miR-100*	CAAGCUUGUAUCUAUAGGUAG	115	TACAAGCTT	1052	ACAAGCTT	1989	CAAGCTT	2926
hsa-miR-101	UACAGUACUGUGUAUACUGAA	116	CAGTACTGT	1053	AGTACTGT	1990	GACTGT	2927
hsa-miR-101*	CAGUUUACACAGUGCUGAUGCU	117	GTGATAACT	1054	TGATAACT	1991	GATAACT	2928
hsa-miR-103	AGCAGCAUUGUACAGGGCUAUGA	118	CAATGCTGC	1055	AATGCTGC	1992	ATGCTGC	2929
hsa-miR-103-as	UCAUAGCCCGUACAAUGCUGCU	119	AGGGCTATG	1056	GGGCTATG	1993	GGCTATG	2930
hsa-miR-105	UCAAAUGCUCAGACUCCUGUGGU	120	GAGCAATTTG	1057	AGCAATTTG	1994	GCATTTG	2931
hsa-miR-105*	ACGGAUGUUUGAGCAUGUGCUA	121	AAACATCCG	1058	AACATCCG	1995	ACATCCG	2932
hsa-miR-106a	AAAAGUGCUUACAGUGCAGGUAG	122	AAGCACTTT	1059	AGCACTTT	1996	GCACTTT	2933
hsa-miR-106a*	CUGCAAUGUAAGCACUUCUUC	123	TACATTGCA	1060	ACATTGCA	1997	CATTGCA	2934
hsa-miR-106b*	UAAAUGUCUGACAGUGCAGAU	124	CAGCACTTT	1061	AGCACTTT	1998	GCACTTT	2935
hsa-miR-106b*	CCGCACUGUGGGUACUUGCUGC	125	CACAGTGCG	1062	ACAGTGCG	1999	CAGTGCG	2936
hsa-miR-107	AGCAGCAUUGUACAGGGCUAUC	126	CAATGCTGC	1063	AATGCTGC	2000	ATGCTGC	2937
hsa-miR-10a	UACCCUGUAGAUCCGAAUUUGUG	127	CTACAGGGT	1064	TACAGGGT	2001	ACAGGGT	2938
hsa-miR-10a*	CAAAUUCGUUUCUAGGGGAUA	128	TACGAATTT	1065	ACGAATTT	2002	CGAATTT	2939
hsa-miR-10b	UACCCUGUAGAACCGAAUUUGUG	129	CTACAGGGT	1066	TACAGGGT	2003	ACAGGGT	2940
hsa-miR-10b*	ACAGAUUCGAUUCUAGGGGAU	130	TCGAATCTG	1067	CGAATCTG	2004	GAATCTG	2941
hsa-miR-1178	UUGCUCACUGUUUCUCCCUAG	131	CAGTGAGCA	1068	AGTGAGCA	2005	GTGAGCA	2942

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1179	AAGCAUUCUUUCAUUUGGUUGG	132	AAGAATGCT	1069	AGAATGCT	2006	GAATGCT	2943
hsa-miR-1180	UUUCCGGCUCGCGUGGGUGUGU	133	GAGCCGGAA	1070	AGCCGGAA	2007	GCCGGAA	2944
hsa-miR-1181	CCGUCGCCGCCACCCGAGCCG	134	GCGCGACG	1071	CGCGACG	2008	GCGGACG	2945
hsa-miR-1182	GAGGUCUUUGGGGGAUGUGAC	135	CAAGACCCCT	1072	AAGACCCCT	2009	AGACCCCT	2946
hsa-miR-1183	CACUGAGGUGAUGGUGAGAGUGG GCA	136	ACCTACAGT	1073	CCTACAGT	2010	CTACAGT	2947
hsa-miR-1184	CCUGCAGGACUUUGAUGGCUUCC	137	TCGCTGCAG	1074	CGCTGCAG	2011	GCTGCAG	2948
hsa-miR-1185	AGAGGAUACCCUUUGUAUGUU	138	GGTATCCTC	1075	GTATCCTC	2012	TATCCTC	2949
hsa-miR-1197	UAGGACACAUGGUCUACUUCU	139	ATGTGTCTT	1076	TGTGTCTT	2013	GTGTCTT	2950
hsa-miR-1200	CUCCUGAGCCAUUCUGAGCCUC	140	GGCTCAGGA	1077	GCTCAGGA	2014	CTCAGGA	2951
hsa-miR-1201	AGCCUGAUUAAACACAUGCUCUGA	141	TAATCAGGC	1078	AATCAGGC	2015	ATCAGGC	2952
hsa-miR-1202	GUGCCAGCUGCAGUGGGGAG	142	CAGCTGGCA	1079	AGCTGGCA	2016	GCTGGCA	2953
hsa-miR-1203	CCCGGAGCCAGGAUGCAGCUC	143	TGGCTCCGG	1080	GGCTCCGG	2017	GCTCCGG	2954
hsa-miR-1204	UCGUGGCCUGGUCUCCAUUUAU	144	CAGGCCACG	1081	AGGCCACG	2018	GGCCACG	2955
hsa-miR-1205	UCUGCAGGGUUUGCUUUUGAG	145	ACCCTGCAG	1082	CCCTGCAG	2019	CCTGCAG	2956
hsa-miR-1206	UGUUCAUGAUGAUGUUUAAGC	146	TACATGAAC	1083	ACATGAAC	2020	CATGAAC	2957
hsa-miR-1207-3p	UCAGCUGGCCUCCAUUUUC	147	GGCCAGCTG	1084	GCCAGCTG	2021	CCAGCTG	2958
hsa-miR-1207-5p	UGGCAGGGAGGCUGGGAGGGG	148	CTCCCTGCC	1085	TCCCTGCC	2022	CCCTGCC	2959
hsa-miR-1208	UCACUGUUCAGACAGCGGGA	149	TGAACAGTG	1086	GAACAGTG	2023	AACAGTG	2960
hsa-miR-122	UGGAGUGUGACAAUGGUGUUUG	150	TCACACTCC	1087	CACACTCC	2024	ACACTCC	2961
hsa-miR-122*	AACGCCAUUUAUCACACUAAAUA	151	TAATGGCGT	1088	AATGGCGT	2025	ATGGCGT	2962
hsa-miR-1224-3p	CCCCACCUCCUCUCUCCUCAG	152	GGAGGTGGG	1089	GAGGTGGG	2026	AGGTGGG	2963
hsa-miR-1224-5p	GUGAGGACUCGGGAGGUGG	153	GAGTCCCTCA	1090	AGTCCCTCA	2027	GTCCTCA	2964
hsa-miR-1225-3p	UGAGCCCCCUGUGCCGCCCCCCAG	154	CAGGGGCTC	1091	AGGGGCTC	2028	GGGGCTC	2965



(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1225-5p	GUGGGUACGGCCCGAGUGGGGG	155	CCGTACCCA	1092	CGTACCCA	2029	GTACCCA	2966
hsa-miR-1226	UACACAGCCUCUGUUCUUCCUAG	156	GGGCTGGTG	1093	GGCTGGTG	2030	GCTGGTG	2967
hsa-miR-1226*	GUGAGGGCAUGCAGGCCUCUGGAUG GGG	157	ATGCCCTCA	1094	TGCCCTCA	2031	GCCCTCA	2968
hsa-miR-1227	CGUGCCACCCUUUJCCCCAG	158	GGGTGGCAC	1095	GGTGGCAC	2032	GTGGCAC	2969
hsa-miR-1228	UCACACCUCCUCGCCCCCCC	159	GCAGGTGTG	1096	CAGGTGTG	2033	AGGTGTG	2970
hsa-miR-1228*	GUGGGGGGGGCAGGUGUGUG	160	CCCCGCCCA	1097	CCCCGCCCA	2034	CCGCCCA	2971
hsa-miR-1229	CUCUCACCACUGCCCCUCCCCACAG	161	GTGGTGAGA	1098	TGGTGAGA	2035	GGTGAGA	2972
hsa-miR-1231	GUGUCUGGGCGGACAGCUGC	162	GCCCAGACA	1099	CCCAGACA	2036	CCAGACA	2973
hsa-miR-1233	UGAGCCUCUGUCCUCCCGCAG	163	ACAGGGCTC	1100	CAGGGCTC	2037	AGGGCTC	2974
hsa-miR-1234	UCGGCCUGACACCCACCCAC	164	GTCAGGCCG	1101	TCAGGCCG	2038	CAGGCCG	2975
hsa-miR-1236	CCUCUCCCCUUGUCUCUCCAG	165	GGGGAAGAG	1102	GGGAAGAG	2039	GGAAGAG	2976
hsa-miR-1237	UCCUUCUGCUCUCCUCCCCCAG	166	AGCAGAAGG	1103	GCAGAAGG	2040	CAGAAGG	2977
hsa-miR-1238	CUUCCUCGUCUGUCUGCCCC	167	GACGAGGAA	1104	ACGAGGAA	2041	CGAGGAA	2978
hsa-miR-124	UAAGGCACGGGUGAAUGCC	168	GCGTGCCTT	1105	CGTGCCTT	2042	GTGCCTT	2979
hsa-miR-124*	CGUGUUCACACGGGACCUUGAU	169	TGTGAACAC	1106	GTGAACAC	2043	TGAACAC	2980
hsa-miR-1243	AACUGGAUCAUUUUAAGGAGUG	170	TGATCCAGT	1107	GATCCAGT	2044	ATCCAGT	2981
nsa-miR-1244	AAGUAGUUGUUUUGUAUGAGAUUG UU	171	CCAACTACT	1108	CAACTACT	2045	AACTACT	2982
hsa-miR-1245	AAGUGAUCUAAAAGCCUACAU	172	TAGATCACT	1109	AGATCACT	2046	GATCACT	2983
nsa-miR-1246	AAUGGAUUUUUGGAGCAGG	173	AAAATCCAT	1110	AAATCCAT	2047	AATCCAT	2984
hsa-miR-1247	ACCCGUCCGUUCGUCCCCCGGA	174	CGGGACGGG	1111	GGGACGGG	2048	GGACGGG	2985
hsa-miR-1248	ACCUUCUUGUAUAAGCACUGUGCU AAA	175	ACAAGAAGG	1112	CAAGAAGG	2049	AAGAAGG	2986

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1249	ACGCCUUUCCCCCCCUCUUCUA	176	GGAAGGGCG	1113	GAAGGGCG	2050	AAGGGCG	2987
hsa-miR-1250	ACGGUGCUGGAUGUGGCCUUU	177	CCAGCACCG	1114	CAGCACCG	2051	AGCACCG	2988
hsa-miR-1251	ACUCUAGCUGCCAAAGGCGCU	178	CAGCTAGAG	1115	AGCTAGAG	2052	GCTAGAG	2989
hsa-miR-1252	AGAGGAAUUUGAAUUCAUUUA	179	ATTTCCTTC	1116	TTTCCTTC	2053	TTCCTTC	2990
hsa-miR-1253	AGAGAAAGAAGAUACAGCCUGCA	180	CTTCTTCTC	1117	TTCTTCTC	2054	TCTTCTC	2991
hsa-miR-1254	AGCCUGGAAGCUGGAGCCUGCAGU	181	CTTCCAGGC	1118	TTCCAGGC	2055	TCCAGGC	2992
hsa-miR-1255a	AGGAUGAGCAAAAGAAAGUAGAUU	182	TGCTCATCC	1119	GCTCATCC	2056	CTCATCC	2993
hsa-miR-1255b	CGGAUGAGCAAAAGAAAGUGGUU	183	TGCTCATCC	1120	GCTCATCC	2057	CTCATCC	2994
hsa-miR-1256	AGGCAUUGACUUCACUAGCU	184	GTC AATGCC	1121	TCAATGCC	2058	CAATGCC	2995
hsa-miR-1257	AGUGAAUGAUGGGUUCUGACC	185	ATCATTAC	1122	TCATTAC	2059	CATTAC	2996
hsa-miR-1258	AGUUAGGAUUAGGUCGUGGAA	186	AATCCTAAC	1123	ATCCTAAC	2060	TCCTAAC	2997
hsa-miR-1259	AUAUAUGAUGACUUAGCUUUU	187	CATCATATA	1124	ATCATATA	2061	TCATATA	2998
hsa-miR-125a-3p	ACAGGUGAGGUUCUUGGGAGCC	188	CCTCACCTG	1125	CTCACCTG	2062	TCACCTG	2999
hsa-miR-125a-5p	UCCUGAGACCCUUUAACCCUGUGA	189	GTCTCAGGG	1126	TCTCAGGG	2063	CTCAGGG	3000
hsa-miR-125b	UCCUGAGACCCUUAACUUGUGA	190	GTCTCAGGG	1127	TCTCAGGG	2064	CTCAGGG	3001
hsa-miR-125b-1*	ACGGUUAGGCUCUUGGGAGCU	191	CCTAACCCG	1128	CTAACCCG	2065	TAACCCG	3002
hsa-miR-125b-2*	UCACAAGUCAGGCUCUUGGGAC	192	TGACTTGTG	1129	GACTTGTG	2066	ACTTGTG	3003
hsa-miR-126	UCGUACCGUGAGUAAUUGCG	193	CACGGTACG	1130	ACGGTACG	2067	CGGTACG	3004
hsa-miR-126*	CAUUUUACUUUUGGUACGCG	194	AGTAATAAT	1131	GTAATAAT	2068	TAATAAT	3005
hsa-miR-1260	AUCCCACCUCUGCCACCA	195	GAGGTGGGA	1132	AGGTGGGA	2069	GGTGGGA	3006
hsa-miR-1261	AUGGAUAAGCCUUUGGCUU	196	CCTTATCCA	1133	CTTATCCA	2070	TTATCCA	3007
hsa-miR-1262	AUGGGUAAUUUUGUAGAAGGAU	197	ATTCACCCA	1134	TTCACCCA	2071	TCACCCA	3008
hsa-miR-1263	AUGGUACCCUGGCAUACUGAGU	198	AGGGTACCA	1135	GGGTACCA	2072	GGTACCA	3009

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1264	CAAGUCUUUUUAGAGCACCUGUU	199	ATAAGACTT	1136	TAAGACTT	2073	AAGACTT	3010
hsa-miR-1265	CAGGAUGUGGUCUAAGUGUUUGUU	200	CCACATCCT	1137	CACATCCT	2074	ACATCCT	3011
hsa-miR-1266	CCUCAGGGCUGUAGAACAGGGGCU	201	AGCCCTGAG	1138	GCCCTGAG	2075	CCCTGAG	3012
hsa-miR-1267	CCUGUUGAAGUGUAAUCCCCA	202	CTTCAACAG	1139	TTCAACAG	2076	TCAACAG	3013
hsa-miR-1268	CGGGCGUGUGGUGGGGG	203	ACCACGCC	1140	CCACGCC	2077	CACGCC	3014
hsa-miR-1269	CUGGACUGAGCCGUGCUACUGG	204	CTCAGTCCA	1141	TCAGTCCA	2078	CAGTCCA	3015
hsa-miR-127-3p	UCGGAUCCGUCUGAGCUUGGCU	205	ACGGATCCG	1142	CGGATCCG	2079	GGATCCG	3016
hsa-miR-127-5p	CUGAAGCUCAGAGGGCUCUGAU	206	TGAGCTTCA	1143	GAGCTTCA	2080	AGCTTCA	3017
hsa-miR-1270	CUGGAGAUUGGAAGAGCUGUGU	207	ATATCTCCA	1144	TATCTCCA	2081	ATCTCCA	3018
hsa-miR-1271	CUUGGCACCUAGCAAGCACUCA	208	AGGTGCCAA	1145	GGTGCCAA	2082	GTGCCAA	3019
hsa-miR-1272	GAUGAUGAGGGCAGCAAUUUCUGA AA	209	CATCATCAT	1146	ATCATCAT	2083	TCATCAT	3020
hsa-miR-1273	GGGGACAAAAGCAAGACUCUUUCU U	210	TTTGTGCC	1147	TTGTGCC	2084	TGTGCC	3021
hsa-miR-1274a	GUCCUGUUUCAGGCGCCA	211	GAACAGGGA	1148	AACAGGGA	2085	ACAGGGA	3022
hsa-miR-1274b	UCCUGUUCGGGCGCCA	212	CGAACAGGG	1149	GAACAGGG	2086	AACAGGG	3023
hsa-miR-1275	GUGGGGAGAGGCUGUC	213	TCTCCCCA	1150	CTCCCCA	2087	TCCCCA	3024
hsa-miR-1276	UAAAGAGCCUGUGGAGACA	214	GGGCTCTT	1151	GGCTCTT	2088	GCTCTT	3025
hsa-miR-1277	UACGUAGAUUUUUUUUUUU	215	TATCTACGT	1152	ATCTACGT	2089	TCTACGT	3026
hsa-miR-1278	UAGUACUGGCAUAUCAUCUUAU	216	CACAGTACT	1153	ACAGTACT	2090	CAGTACT	3027
hsa-miR-1279	UCAUUUGCUUUUUUUU	217	AGCAATATG	1154	GCAATATG	2091	CAATATG	3028
hsa-miR-128	UCACAGUGAACGGGUCUCUUU	218	TTCACTGTG	1155	TCACTGTG	2092	CACTGTG	3029
hsa-miR-1280	UCCACCCGUCGCCACCC	219	AGCGGTGGG	1156	GCGGTGGG	2093	CGGTGGG	3030
hsa-miR-1281	UCGCCUCCUCUCUCC	220	GAGGAGGCG	1157	AGGAGGCG	2094	GGAGGCG	3031

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1282	UCGUUUGCCUUUUUCUGCUU	221	AGGCAAACG	1158	GGCAAACG	2095	GCAAACG	3032
hsa-miR-1283	UCUACAAAGGAAAGCGCUUUCU	222	CCTTTGTAG	1159	CTTTGTAG	2096	TTTGTAG	3033
hsa-miR-1284	UCUAUACAGACCCUGGCUUUUC	223	TCTGTATAG	1160	CTGTATAG	2097	TGTATAG	3034
hsa-miR-1285	UCUGGGCAACAAGUGAGACCU	224	GTTGCCCAG	1161	TTGCCCAG	2098	TGCCCAG	3035
hsa-miR-1286	UGCAGGACCAAGAUGAGCCCU	225	TGGTCCTGC	1162	GGTCCTGC	2099	GTCCTGC	3036
hsa-miR-1287	UGCUGGAUCAGUGGUUCGAGUC	226	TGATCCAGC	1163	GATCCAGC	2100	ATCCAGC	3037
hsa-miR-1288	UGGACUGCCCUGAUCUGGAGA	227	GGCAGTCC	1164	GGCAGTCC	2101	GCAGTCC	3038
hsa-miR-1289	UGGAGUCCAGGAAUCUGCAUUUU	228	CTGGACTCC	1165	TGGACTCC	2102	GGACTCC	3039
hsa-miR-129*	AAGCCCUUACCCCAAAAAGUAU	229	GTAAGGGCT	1166	TAAGGGCT	2103	AAGGGCT	3040
hsa-miR-129-3p	AAGCCCUUACCCCAAAAAGCAU	230	GTAAGGGCT	1167	TAAGGGCT	2104	AAGGGCT	3041
hsa-miR-129-5p	CUUUUUGCGGUCUGGCUUUGC	231	CCGCAAAA	1168	CGCAAAA	2105	GCAAAA	3042
hsa-miR-1290	UGGAUUUUUGGAUCAGGGA	232	CAAAAATCC	1169	AAAAATCC	2106	AAAATCC	3043
hsa-miR-1291	UGGCCCUGACUGAAGACCAGCAGU	233	GTCAGGGCC	1170	TCAGGGCC	2107	CAGGGCC	3044
hsa-miR-1292	UGGGAACGGGUUCCGGCAGACGCU G	234	CCCGTTCCC	1171	CCGTCCC	2108	CGTTCCC	3045
hsa-miR-1293	UGGGUGGUCUGGAGAUUUUGUC	235	AGACCACCC	1172	GACCACCC	2109	ACCACCC	3046
hsa-miR-1294	UGUGAGGUUGGCAUUGUUGUCU	236	CAACCTCAC	1173	AACCTCAC	2110	ACCTCAC	3047
hsa-miR-1295	UUAGGCCGCAGAUUGGGUGA	237	TGCGGCCTA	1174	GCGGCCTA	2111	CGGCCTA	3048
hsa-miR-1296	UUAGGCCCUUGGCUCCAUCUCC	238	AGGGCCCTA	1175	GGGCCCTA	2112	GGCCCTA	3049
hsa-miR-1297	UUCAAGUAUUUCAGGUG	239	ATTACTTGA	1176	TTACTTGA	2113	TACTTGA	3050
hsa-miR-1298	UUCAUUCGGCUGUCCAGAUGUA	240	GCCGAATGA	1177	CCGAATGA	2114	CGAATGA	3051
hsa-miR-1299	UUCUGGAAUUCUGUGAGGGA	241	AATCCAGA	1178	ATCCAGA	2115	TTCCAGA	3052
hsa-miR-1300	UUGAGAAGGAGGCUGCUG	242	TCCTTCTCA	1179	CCTTCTCA	2116	CTTCTCA	3053
hsa-miR-1301	UUGCAGCUGCCUGGGAGUGACUUC	243	GCAGCTGCA	1180	CAGCTGCA	2117	AGCTGCA	3054

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1302	UUGGGACAUACUUAUGCUAAA	244	TATGTCCCA	1181	ATGTCCCA	2118	TGTCCCA	3055
hsa-miR-1303	UUUAGAGACGGGUCUUGCUCU	245	CGTCTCTAA	1182	GTCTCTAA	2119	TCTCTAA	3056
hsa-miR-1304	UUUGAGGCUACAGUGAGAUGUG	246	TAGCCTCAA	1183	AGCCTCAA	2120	GCCTCAA	3057
hsa-miR-1305	UUUUAACUCUAAUUGGGAGAGA	247	GAGTTGAAA	1184	AGTTGAAA	2121	GTTGAAA	3058
hsa-miR-1306	ACGUUGGCUCUGGUGGUG	248	GAGCCAACG	1185	AGCCAACG	2122	GCCAACG	3059
hsa-miR-1307	ACUCGGGUGGGCGUCGGUGGUG	249	CACGCCGAG	1186	ACGCCGAG	2123	CGCCGAG	3060
hsa-miR-1308	GCAUGGGUGGUUCAGUGG	250	CCACCCATG	1187	CACCCATG	2124	ACCCATG	3061
hsa-miR-130a*	CAGUGCAAUGUUAAAAAGGGCAU	251	CATTGCACT	1188	ATTGCACT	2125	TTGCACT	3062
hsa-miR-130b	UUCACAUUGUGCUACUGUCUGC	252	ACAATGTGA	1189	CAATGTGA	2126	AATGTGA	3063
hsa-miR-130b*	CAGUGCAAUGAUGAAAGGGCAU	253	CATTGCACT	1190	ATTGCACT	2127	TTGCACT	3064
hsa-miR-130b*	ACUUUUCCCUUGUCACUAC	254	GGGAAAGAG	1191	GGAAAGAG	2128	GAAAGAG	3065
hsa-miR-132	UAAAGUCUACAGCCAUUGGUCG	255	TAGACTGTT	1192	AGACTGTT	2129	GACTGTT	3066
hsa-miR-132*	ACCGUGGCUUUUGAUUGUUACU	256	AAGCCACGG	1193	AGCCACGG	2130	GCCACGG	3067
hsa-miR-1321	CAGGGAGGUGAAUGUGAU	257	CACCTCCCT	1194	ACCTCCCT	2131	CCTCCCT	3068
hsa-miR-1322	GAUGAUGCUGCUGAUGCUG	258	CAGCATCAT	1195	AGCATCAT	2132	GCATCAT	3069
hsa-miR-1323	UCAAAAACUGAGGGCAUUUUUCU	259	TCAGTTTTG	1196	CAGTTTTG	2133	AGTTTTG	3070
hsa-miR-1324	CCAGACAGAAUUCUAUGCACUUCU	260	TTCTGTCTG	1197	TCTGTCTG	2134	CTGTCTG	3071
hsa-miR-133a	UUUGGUCCCUUCAACCAGCUG	261	GGGACCAA	1198	GGGACCAA	2135	GGACCAA	3072
hsa-miR-133b	UUUGGUCCCCUUCAACCAGCUA	262	GGGACCAA	1199	GGGACCAA	2136	GGACCAA	3073
hsa-miR-134	UGUGACUGGUUGACCAGAGGGG	263	ACCAGTCAC	1200	CCAGTCAC	2137	CAGTCAC	3074
hsa-miR-135a	UAUGGCUUUUUUAUCCUAUGUGA	264	AAAAGCCAT	1201	AAAGCCAT	2138	AAGCCAT	3075
hsa-miR-135a*	UAUAGGGAUUGGAGCCGUGGCG	265	AATCCCTAT	1202	ATCCCTAT	2139	TCCCTAT	3076
hsa-miR-135b	UAUGGCUUUUCAUCCUAUGUGA	266	AAAAGCCAT	1203	AAAGCCAT	2140	AAGCCAT	3077

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-135b*	AUGUAGGGCUAAAAAGCCAUUGGG	267	AGCCCTACA	1204	GCCCTACA	2141	CCCTACA	3078
hsa-miR-136	ACUCCAUUUGUUUUUGAUGAUGGA	268	CAAAATGGAG	1205	AAATGGAG	2142	AATGGAG	3079
hsa-miR-136*	CAUCAUCGUCUCAAAUGAGUCU	269	GACGATGAT	1206	ACGATGAT	2143	CGATGAT	3080
hsa-miR-137	UUUUUGCUUAAGAAUACGGCUAG	270	TAAGCAATA	1207	AAGCAATA	2144	AGCAATA	3081
hsa-miR-138	AGCUGGUGUUUGAAUCAGGCCG	271	AACACCAGC	1208	ACACCAGC	2145	CACCAGC	3082
hsa-miR-138-1*	GCUACUUCACAAACACCAGGGCC	272	GTGAAGTAG	1209	TGAAGTAG	2146	GAAGTAG	3083
hsa-miR-138-2*	GCUUUUUCACGACACCAGGGUU	273	GTGAAATAG	1210	TGAAATAG	2147	GAAATAG	3084
hsa-miR-139-3p	GGAGACGGGCCCGUUUGGAGU	274	CCGGTCTC	1211	CGGTCTC	2148	GCGTCTC	3085
hsa-miR-139-5p	UCUACAGUGCACGGUGUCUCCAG	275	GCACTGTAG	1212	CACTGTAG	2149	ACTGTAG	3086
hsa-miR-140-3p	UACCACAGGGUAGAACCACGG	276	CCCTGTGGT	1213	CCTGTGGT	2150	CTGTGGT	3087
hsa-miR-140-5p	CAGUGUUUUACCCUAGUUAG	277	AAAACCACT	1214	AAACCACT	2151	AACCACT	3088
hsa-miR-141	UAACACUGUCUGGUAAAAGAUUG	278	GACAGTGTT	1215	ACAGTGTT	2152	CAGTGTT	3089
hsa-miR-141*	CAUCUCCAGUACAGUUGGA	279	CTGGAAGAT	1216	TGGAAGAT	2153	GGAAGAT	3090
hsa-miR-142-3p	UGUAGUUUUCCUACUUUUGGA	280	AAACACTAC	1217	AACACTAC	2154	ACACTAC	3091
hsa-miR-142-5p	CAUAAAAGUAGAAAGCACUACU	281	CTACTTTAT	1218	TACTTTAT	2155	ACTTTAT	3092
hsa-miR-143	UGAGAUGAAGCACUGUAGCUC	282	CTTCATCTC	1219	TTCATCTC	2156	TCATCTC	3093
hsa-miR-143*	GGUGCAGUGCUGCAUCUCUGGU	283	GCACTGCAC	1220	CACTGCAC	2157	ACTGCAC	3094
hsa-miR-144	UACAGUAUGAUGAUGUACU	284	CTATACTGT	1221	TATACTGT	2158	ATACTGT	3095
hsa-miR-144*	GGAAUUAUCAUUAUACUGUAAG	285	GATGATATC	1222	ATGATATC	2159	TGATATC	3096
hsa-miR-145	GUCCAGUUUUCCCAGGAAUCCCU	286	AAAACCTGA	1223	AAACTGGA	2160	AACTGGA	3097
hsa-miR-145*	GGAUUCCUGGAAAUAUCUGUUCU	287	CCAGGAATC	1224	CAGGAATC	2161	AGGAATC	3098
hsa-miR-1468	CUCGGUUUGCCUGUUUUGCUG	288	GCAAACGGA	1225	CAAACGGA	2162	AAACGGA	3099
hsa-miR-1469	CUCGGCGGGGGCGGGGCUCC	289	CCGGCCCGA	1226	CGGCCCGA	2163	GCGCCGA	3100

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-146a	UGAGAACUGAAUUCCAUGGGUU	290	TCAGTTCTC	1227	CAGTTCTC	2164	AGTTCTC	3101
hsa-miR-146a*	CCUCUGAAUUUCAGUUCUUCAG	291	ATTCAGAG	1228	TTTCAGAG	2165	TTCAGAG	3102
hsa-miR-146b-3p	UGCCUCGDGGACUCAGUUCUGG	292	CCACAGGGC	1229	CACAGGGC	2166	ACAGGGC	3103
hsa-MiR-146b-5p	UGAGAACUGAAUUCCAUAGGCU	293	TCAGTTCTC	1230	CAGTTCTC	2167	AGTTCTC	3104
hsa-miR-147	GUGUGGAAAUGCUUCUGC	294	TCCACACA	1231	TCCACACA	2168	CCACACA	3105
hsa-miR-1470	GCCUCCGCCCGUGCACC	295	GGCGAGGG	1232	GCGGAGGG	2169	CGGAGGG	3106
hsa-miR-1471	GCCCGGUGUGGAGCCAGGUGU	296	ACACGGGG	1233	CACGCGGG	2170	ACGCGGG	3107
hsa-miR-147b	GUGUGGAAAUGCUUCUGUA	297	TCCGCACA	1234	TCCGCACA	2171	CCGCACA	3108
hsa-miR-148a	UCAGUGCACUACAGAAUUUGU	298	AGTGCACTG	1235	GTGCACTG	2172	TGCACTG	3109
hsa-miR-148a*	AAAGUUCUGAGACACUCGACU	299	TCAGAACTT	1236	CAGAACTT	2173	AGAACTT	3110
hsa-miR-148b	UCAGUGCAUCACAGAAUUUGU	300	GATGCACTG	1237	ATGCACTG	2174	TGCACTG	3111
hsa-miR-148b*	AAGUUCUGUUUAACACUCAGGC	301	AACAGAACT	1238	ACAGAACT	2175	CAGAACT	3112
hsa-miR-149	UCUGGCUCGUGUCUUCACUCCC	302	CGGAGCCAG	1239	GGAGCCAG	2176	GAGCCAG	3113
hsa-miR-149*	AGGGAGGACGGGGGUGUGC	303	GTCCCTCCC	1240	TCCCTCCC	2177	CCCTCCC	3114
hsa-miR-150	UCUCCCAACCCUUUGUACCCAGUG	304	GGTTGGGAG	1241	GTTGGGAG	2178	TTGGGAG	3115
hsa-miR-150*	CUGGUACAGGCCUUGGGGACAG	305	CCTGTACCA	1242	CTGTACCA	2179	TGTACCA	3116
hsa-miR-151-3p	CUAGACUGAAGCUCUUGAGG	306	TTCAGTCTA	1243	TCAGTCTA	2180	CAGTCTA	3117
hsa-miR-151-5p	UCGAGGAGCUCACAGUCUAGU	307	AGCTCCTCG	1244	GCTCCTCG	2181	CTCCTCG	3118
hsa-miR-152	UCAGUGCAUGACAGAAUUUGG	308	CATGCACTG	1245	ATGCACTG	2182	TGCACTG	3119
hsa-miR-153	UUGCAUAGUCACAAAAGUGAUC	309	GACTATGCA	1246	ACTATGCA	2183	CTATGCA	3120
hsa-miR-1537	AAAACCGUCUAGUUACAGUUUGU	310	AGACGGTTT	1247	GACGGTTT	2184	ACGGTTT	3121
hsa-miR-1538	CGGCCGGGUCUGUCUUGUCCU	311	GCCCCGGCC	1248	CCCCGGCC	2185	CCGGGCC	3122
hsa-miR-1539	UCCUGCGGUCUCCAGAUGCC	312	ACGGCCAGG	1249	CGGCAGG	2186	GCGCAGG	3123

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-154	UAGGUUAUCCGGUUGCCUUCG	313	GGATAACCT	1250	GATAACCT	2187	ATAACCT	3124
hsa-miR-154*	AAUCAUACACGGUUGACCUAUU	314	GTGTATGAT	1251	TGTATGAT	2188	GTATGAT	3125
hsa-miR-155	UUAUUGCUAUUCGUGAUAGGGGU	315	TTAGCATT	1252	TAGCATT	2189	AGCATT	3126
hsa-miR-155*	CUCCUACAUUAGCAUUAACA	316	TATGTAGGA	1253	ATGTAGGA	2190	TGTAGGA	3127
hsa-miR-15a	UAGCAGCACAUAAUGUUUGUG	317	TGTGCTGCT	1254	GTGCTGCT	2191	TGCTGCT	3128
hsa-miR-15a*	CAGGCCAUUUGUCGCCUCA	318	ATATGGCCT	1255	TATGGCCT	2192	ATGGCCT	3129
hsa-miR-15b	UAGCAGCACAUCAUGGUUUAACA	319	TGTGCTGCT	1256	GTGCTGCT	2193	TGCTGCT	3130
hsa-miR-15b*	CGAAUCAUUUUUGCUCUCUA	320	TAATGATTC	1257	AATGATTC	2194	ATGATTC	3131
hsa-miR-16	UAGCAGCACGUAAAUUUGGCG	321	CGTGCTGCT	1258	GTGCTGCT	2195	TGCTGCT	3132
hsa-miR-16-1*	CCAGUAUUUAAACUGUCGUCUGA	322	TTAATACTG	1259	TAATACTG	2196	AATACTG	3133
hsa-miR-16-2*	CCAAUUUACUGUCGUCUUUA	323	GTAATATTG	1260	TAATATTG	2197	AATATTG	3134
hsa-miR-17	CAAGUGCUUACAGUCGAGGUAG	324	AAGCACTTT	1261	AGCACTTT	2198	GCACTTT	3135
hsa-miR-17*	ACUGCAGUGAAGGCACUUGUAG	325	TCACTGCAG	1262	CACTGCAG	2199	ACTGCAG	3136
hsa-miR-181a	AACAUUCAACGCUGUCGGUGAGU	326	GTTGAATGT	1263	TTGAATGT	2200	TGAATGT	3137
hsa-miR-181a*	ACCAUCGACCGUUUGAUUGUACC	327	GGTCGATGG	1264	GTCGATGG	2201	TCGATGG	3138
hsa-miR-181a-2*	ACCACUGACCGUUUGACUUGUACC	328	GGTCAGTGG	1265	GTCAGTGG	2202	TCAGTGG	3139
hsa-miR-181b	AACAUUCAUUGCUGUCGGUGGGU	329	AATGAATGT	1266	ATGAATGT	2203	TGAATGT	3140
hsa-miR-181c	AACAUUCAACCGUCGUGAGU	330	GTTGAATGT	1267	TTGAATGT	2204	TGAATGT	3141
hsa-miR-181c*	AACCAUCGACCGUUUGAGUGGAC	331	GTCGATGCT	1268	TCGATGCT	2205	CGATGCT	3142
hsa-miR-181d	AACAUUCAUUGUUGCUGGGUGGU	332	AATGAATGT	1269	ATGAATGT	2206	TGAATGT	3143
hsa-miR-182	UUUGGCAAUGGUAGAACUCACACU	333	CATTGCCAA	1270	ATTGCCAA	2207	TTGCCAA	3144
hsa-miR-182*	UGGUUCUAGAGUUGCCAACUA	334	TCTAGAACC	1271	CTAGAACC	2208	TAGAACC	3145
hsa-miR-1825	UCCAGUGCCCUCCUCUCC	335	GGGCACTGG	1272	GGCACTGG	2209	GCACCTGG	3146



(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-1826	AUUGAUCaucGACACUUCGAAACGGC AAU	336	GATGATCAA	1273	ATGATCAA	2210	TGATCAA	3147
hsa-miR-1827	UGAGGCAGUAGAUUGAAU	337	TACTGCCTC	1274	ACTGCCTC	2211	CTGCCTC	3148
hsa-miR-183	UAUGGCACUGGUAGAAUUCACU	338	CAGTGCCAT	1275	AGTGCCAT	2212	GTGCCAT	3149
hsa-miR-183*	GUGAAUACCGAAGGCCAUAA	339	GGTAATTCA	1276	GTAATTCA	2213	TAATTCA	3150
hsa-miR-184	UGGACGGAGAACUGAUUAGGGU	340	TCTCCGTCC	1277	CTCCGTCC	2214	TCCGTCC	3151
hsa-miR-185	UGGAGAGAAAAGGCAGUUCUGA	341	TTTCTCTCC	1278	TTCTCTCC	2215	TCTCTCC	3152
hsa-miR-185*	AGGGCUGGCUUUCCUCUGGUC	342	GCCAGCCCC	1279	CCAGCCCC	2216	CAGCCCC	3153
hsa-miR-186	CAAAGAAUUCUCCUUUUGGGCU	343	GAATTCITT	1280	AATTCITT	2217	ATTCITT	3154
hsa-miR-186*	GCCCAAAGGUGAAUUUUUUGGG	344	ACCTTTGGG	1281	CCTTTGGG	2218	CTTTGGG	3155
hsa-miR-187	UCGUGUCUUUGUUGCAGCCGG	345	CAAGACACG	1282	AAGACACG	2219	AGACACG	3156
hsa-miR-187*	GGCUACAACACAGGACCCGGGC	346	TGTTGTAGC	1283	GTTGTAGC	2220	TTGTAGC	3157
hsa-miR-188-3p	CUCCCAUCAGCAGGGUUUGCA	347	CATGTGGGA	1284	ATGTGGGA	2221	TGTGGGA	3158
hsa-miR-188-5p	CAUCCCUUGCAUGGUGGAGGG	348	GCAAGGGAT	1285	CAAGGGAT	2222	AAGGGAT	3159
hsa-miR-18a	UAAGGUGCAUCUAGUGCAGAUAG	349	ATGCACCTT	1286	TGCACCTT	2223	GCACCTT	3160
hsa-miR-18a*	ACUGCCCUAAGUGCUCCUUCUGG	350	TTAGGGCAG	1287	TAGGGCAG	2224	AGGGCAG	3161
hsa-miR-18b	UAAGGUGCAUCUAGUGCAGUUAG	351	ATGCACCTT	1288	TGCACCTT	2225	GCACCTT	3162
hsa-miR-18b*	UGCCCUAAAUGCCCCUUCUGGC	352	ATTTAGGGC	1289	TTTAGGGC	2226	TTAGGGC	3163
hsa-miR-190	UGAUUGUUUGAUUUAUUAGGU	353	AAACATATC	1290	AACATATC	2227	ACATATC	3164
hsa-miR-1908	CGGCGGGACGGCGAUUUGGUC	354	GTCCCCGCC	1291	TCCCCGCC	2228	CCCCGCC	3165
hsa-miR-1909	CGCAGGGCCGGGUGUCUCACCCG	355	GGCCCCCTGC	1292	GCCCCCTGC	2229	CCCCTGC	3166
hsa-miR-1909*	UGAGUGCCGGUGCCUCCCCUG	356	CCGGCACTC	1293	CGGCACTC	2230	GGCACTC	3167
hsa-miR-190b	UGAUUGUUUGAUUUAUUUGGUU	357	AAACATATC	1294	AACATATC	2231	ACATATC	3168
hsa-miR-191	CAACGGAAUCCCAAAAAGCAGCUG	358	GATTCGGTT	1295	ATTCGGTT	2232	TTCCGTT	3169

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-191*	GCUGGCUUUGGAUUUGUCUCCC	359	CAAGCGCAG	1296	AAGCGCAG	2233	AGCGCAG	3170
hsa-miR-1910	CCAGUCCUGUGCCUGCCGCCU	360	ACAGGACTG	1297	CAGGACTG	2234	AGGACTG	3171
hsa-miR-1911	UGAGUACCGCCAUUGUCUUGGG	361	GCGTACTC	1298	CGGTACTC	2235	GGTACTC	3172
hsa-miR-1911*	CACCAGGCAUUGUGUCUCC	362	ATGCCTGGT	1299	TGCCTGGT	2236	GCCTGGT	3173
hsa-miR-1912	UACCCAGAGCAUGCAGUGUGAA	363	GCTCTGGGT	1300	CTCTGGGT	2237	TCTGGGT	3174
hsa-miR-1913	UCUGCCCCUCCGUCUGGCCA	364	AGGGGCAG	1301	GGGGGCAG	2238	GGGGCAG	3175
hsa-miR-1914	CCUUGUCCCCGGCCACUUCUG	365	GGCACACAGG	1302	GGCACACAGG	2239	GCACACAGG	3176
hsa-miR-1914*	GGAGGGUCCCCGCACUGGGAGG	366	GGACCCCTC	1303	GACCCCTC	2240	ACCCCTC	3177
hsa-miR-1915	CCCCAGGGCGACGGCGGG	367	CGCCCTGGG	1304	GCCCTGGG	2241	CCCTGGG	3178
hsa-miR-1915*	ACCUUGCCUUGCUGCCCGGGCC	368	AAGGCAAGG	1305	AGGCAAGG	2242	GGCAAGG	3179
hsa-miR-192	CUGACCUAUGAAUUGACAGCC	363	CATAGGTCA	1306	ATAGGTCA	2243	TAGGTCA	3180
hsa-MIR-192*	CUGCCAAUCCAUAGGUCACAG	370	GAATTGGCA	1307	AATTGGCA	2244	ATTGGCA	3181
hsa-miR-193a-3p	AACUGGCCUACAAAGUCCCAGU	371	TAGGCCAGT	1308	AGGCCAGT	2245	GGCCAGT	3182
hsa-miR-193a-5p	UGGGUCUUUGCGGGCGAGAUGA	372	CAAAGACCC	1309	AAAGACCC	2246	AAGACCC	3183
hsa-miR-193b	AACUGGCCUCAAAGUCCCAGU	373	AGGGCCAGT	1310	GGGGCCAGT	2247	GGCCAGT	3184
hsa-miR-193b*	CGGGUUUUUGAGGGCGAGAUGA	374	CAAAACCCC	1311	AAAACCCC	2248	AAACCCC	3185
hsa-miR-194	UGUAACAGCAACUCCAUUGGGA	375	TGCTGTTAC	1312	GCTGTTAC	2249	CTGTTAC	3186
hsa-miR-194*	CCAGUGGGCUGCUGUUUUCUG	376	GCCCCACTG	1313	CCCCACTG	2250	CCCACTG	3187
hsa-miR-195	UAGCAGCACAGAAAUUUGGC	377	TGTGCTGCT	1314	GTGCTGCT	2251	TGCTGCT	3188
hsa-miR-195*	CCAAUJUUGGCUGUCUCUCC	378	CCAATATTG	1315	CAATATTG	2252	AATATTG	3189
hsa-miR-196a	UAGGUAGUUUCAUUGUUGGG	379	AAACTACCT	1316	AACTACCT	2253	ACTACCT	3190
hsa-miR-196a*	CGGCAACAAGAAACUCCUGAG	380	CTTGTGCCC	1317	TTGTGCCC	2254	TGTTGCC	3191
hsa-miR-196b	UAGGUAGUUUCCUGUUGGG	381	AAACTACCT	1318	AACTACCT	2255	ACTACCT	3192

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-197	UUCACCACCUUCUCCACCCAGC	382	AGGTGGTGA	1319	GGTGGTGA	2256	GTGGTGA	3193
hsa-miR-198	GGUCCAGAGGGGAGAUAGGUUC	383	CCTCTGGAC	1320	CTCTGGAC	2257	TCTGGAC	3194
hsa-miR-199a-5p	CCCAGUGUUUCAGACUACCUUGUUC	384	GAACACTGG	1321	AACACTGG	2258	ACACTGG	3195
hsa-miR-199b-3p	ACAGUAGUCUGCACAUUGGUUA	385	AGACTACTG	1322	GACTACTG	2259	ACTACTG	3196
hsa-miR-199b-5p	CCCAGUGUUUAGACUAUCUGUUC	386	AAACACTGG	1323	AACACTGG	2260	ACACTGG	3197
hsa-miR-19a	UGUGCAAUUCUUGCAAAAACUGA	387	GATTTGCAC	1324	ATTTGCAC	2261	TTTGCAC	3198
hsa-miR-19a*	AGUUUUUGCAUAGUUGCACUACA	388	ATGCAAAAAC	1325	TGCAAAAAC	2262	GCAAAAAC	3199
hsa-miR-19b	UGUGCAAUUCUUGCAAAAACUGA	389	GATTTGCAC	1326	ATTTGCAC	2263	TTTGCAC	3200
hsa-miR-19b-1*	AGUUUUUGCAGGUUUGCAUCCAGC	390	CTGCAAAAAC	1327	TGCAAAAAC	2264	GCAAAAAC	3201
hsa-miR-19b-2*	AGUUUUUGCAGGUUUGCAUUUCA	391	CTGCAAAAAC	1328	TGCAAAAAC	2265	GCAAAAAC	3202
hsa-miR-200a	UAACACUGUCUGGUAAACGAUGU	392	GACAGTGTT	1329	ACAGTGTT	2266	CAGTGTT	3203
hsa-miR-200a*	CAUCUUACCCGGACAGUGCUGGA	393	CGGTAAGAT	1330	GGTAAGAT	2267	GTAAGAT	3204
hsa-miR-200b	UAAUACUGCCUGGUAAUUGAUGA	394	GGCAGTATT	1331	GCAGTATT	2268	CAGTATT	3205
hsa-miR-200b*	CAUCUUACUGCCGGUAAUUGAUGA	395	CAGTAAGAT	1332	AGTAAGAT	2269	GTAAGAT	3206
hsa-miR-200c	UAAUACUGCCGGUAAUUGAUGA	396	GGCAGTATT	1333	GCAGTATT	2270	CAGTATT	3207
hsa-miR-200c*	CGUCUUACCCAGCAGUGUUUUGG	397	GGGTAAGAC	1334	GGTAAGAC	2271	GTAAGAC	3208
hsa-miR-202	AGAGGUUAGGGCAUUGGGAA	398	CTATACCTC	1335	TATACCTC	2272	ATACCTC	3209
hsa-miR-202*	UUCCUUGCAUUAUCUUCUUG	399	TGCATAGGA	1336	GCATAGGA	2273	CATAGGA	3210
hsa-miR-203	GUGAAAUGUUUAGGACCACUAG	400	AACATTTCA	1337	ACATTTCA	2274	CATTTCA	3211
hsa-miR-204	UUCCUUUGUCAUCCUUAUGCCU	401	ACAAAGGGA	1338	CAAAGGGA	2275	AAAGGGA	3212
hsa-miR-205	UCCUUCUUCUCCACCGGAGUCUG	402	GAATGAAGG	1339	AATGAAGG	2276	ATGAAGG	3213
hsa-miR-206	UGGAAUGUAAGGAAAGUGUGUGG	403	TTACATTCC	1340	TACATTCC	2277	ACATTCC	3214
hsa-miR-208a	AUAAGACGAGCAAAAAGCUUGU	404	CTCGTCTTA	1341	TCGTCTTA	2278	CGTCTTA	3215

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-208b	AUAAAGACGAACAAAAGGUUUUGU	405	TTCGTCTTA	1342	TCGTCTTA	2279	CGTCTTA	3216
hsa-miR-20a	UAAAAGUGCUUUAUAGUGCAGGUAG	406	AAGCACTTT	1343	AGCACTTT	2280	GCACTTT	3217
hsa-miR-20a*	ACUGCAUUUAUGAGCACUUAAAG	407	ATAATGCAG	1344	TAATGCAG	2281	AATGCAG	3218
hsa-miR-20b	CAAAUGUCUCAUAGUGCAGGUAG	408	GAGCACTTT	1345	AGCACTTT	2282	GCACTTT	3219
hsa-miR-20b*	ACUGUAGUAUGGGCACUUCAG	409	ATACTACAG	1346	TACTACAG	2283	ACTACAG	3220
hsa-miR-21	UAGCUUAUCAGACUGAUUGA	410	TGATAAGCT	1347	GATAAGCT	2284	ATAAGCT	3221
hsa-miR-21*	CAACACCAGUCGAUGGGCUGU	411	ACTGGTGT	1348	CTGGTGT	2285	TGGTGT	3222
hsa-miR-210	CUGUGCGUGUGACAGCGGCUGA	412	ACACGCACA	1349	CACGCACA	2286	ACGCACA	3223
hsa-miR-211	UUCUUUUUGUCAUCCUUCGCCU	413	ACAAAGGGA	1350	CAAAGGGA	2287	AAAGGGA	3224
hsa-miR-212	UAACAGUCUCCAGUCACGGCC	414	GAGACTGTT	1351	AGACTGTT	2288	GACTGTT	3225
hsa-miR-214	ACAGCAGGCACAGACAGGCAGU	415	TGCCTGCTG	1352	GCCTGCTG	2289	CCTGCTG	3226
hsa-miR-214*	UGCCUGUCUACACUUGCUGUGC	416	TAGACAGGC	1353	AGACAGGC	2290	GACAGGC	3227
hsa-miR-215	AUGACCUAUGAAUUUGACAGAC	417	CATAGGTCA	1354	ATAGGTCA	2291	TAGGTCA	3228
hsa-miR-216a	UAAUCUCAGCUGGCAACUGUGA	418	GCTGAGATT	1355	CTGAGATT	2292	TGAGATT	3229
hsa-miR-216b	AAAUCUCUGCAGGCAAAUUGUGA	419	GCAGAGATT	1356	CAGAGATT	2293	AGAGATT	3230
hsa-miR-217	UACUGCAUCAGGAACUGAUUGGA	420	TGATGCAGT	1357	GATGCAGT	2294	ATGCAGT	3231
hsa-miR-218	UUUGUCUUGAUCUAACCAUGU	421	TCAAGCACA	1358	CAAGCACA	2295	AAGCACA	3232
hsa-miR-218-1*	AUGGUUCCGUCUAAAGCACC AUGG	422	ACGGAACCA	1359	CGGAACCA	2296	GGAACCA	3233
hsa-miR-218-2*	CAUGGUUCUGUCAAGCACCCGCG	423	CAGAACCAT	1360	AGAACCAT	2297	GAACCAT	3234
hsa-miR-219-1-3p	AGAGUUGAGUCUGGACGUCGCCG	424	ACTCAACTC	1361	CTCAACTC	2298	TCAACTC	3235
hsa-miR-219-2-3p	AGAAUUGUGGCUGGACAUUCUGU	425	CCACAATTC	1362	CACAATTC	2299	ACAATTC	3236
hsa-miR-219-Sp	UGAUUGUCCAAACGCAAUUCU	426	TGGACAATC	1363	GGACAATC	2300	GACAATC	3237
hsa-miR-22	AAGCUGCCAGUUUGAAGAACUGU	427	CTGGCAGCT	1364	TGGCAGCT	2301	GGCAGCT	3238

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-22*	AGUUCUUCAGUGGCAAGCUUUA	428	CTGAAGAAC	1365	TGAAGAAC	2302	GAAGAAC	3239
hsa-miR-220a	CCACACCGUAUCUGACACUUU	429	TACGGTGTG	1366	ACGGTGTG	2303	CGGTGTG	3240
hsa-miR-220b	CCACCACCGUGUCGACACUU	430	ACGGTGGTG	1367	CGGTGGTG	2304	GGTGGTG	3241
hsa-miR-220c	ACACAGGGCUGUUGAAGACU	431	AGCCCTGTG	1368	GCCCTGTG	2305	CCCTGTG	3242
hsa-miR-221	AGCUACAUUGUCUGCGGUGUUUC	432	CAATGTAGC	1369	AATGTAGC	2306	ATGTAGC	3243
hsa-miR-221*	ACCUGGCAUACAAGUAGAUUU	433	TATGCCAGG	1370	ATGCCAGG	2307	TGCCAGG	3244
hsa-miR-222	AGCUACAUCUGGCUACUUGGU	434	AGATGTAGC	1371	GATGTAGC	2308	ATGTAGC	3245
hsa-miR-222*	CUCAGUAGCCAGUGUAGAUCU	435	GGCTACTGA	1372	GCTACTGA	2309	CTACTGA	3246
hsa-miR-223	UGUCAGUUUGUCAAAUACCCCA	436	CAAACTGAC	1373	AAACTGAC	2310	AACTGAC	3247
hsa-miR-223*	CGUGAUUUUGACAAGCUGAGUU	437	CAAAATACAC	1374	AAATACAC	2311	AATACAC	3248
hsa-miR-224	CAAGUCACUAGUGUUCGGUU	438	TAGTGACTT	1375	AGTGACTT	2312	GTGACTT	3249
hsa-miR-23a	AUCACAUUGCCAGGGAUUUUC	439	GCAATGTGA	1376	CAATGTGA	2313	AATGTGA	3250
hsa-miR-23a*	GGGUUCCUGGGGUAUGGGAUUU	440	CAGGAACCC	1377	AGGAACCC	2314	GGAACCC	3251
hsa-miR-23b	AUCACAUUGCCAGGGAUUACC	441	GCAATGTGA	1378	CAATGTGA	2315	AATGTGA	3252
hsa-miR-23b*	UGGGUCCUGGGCAUGCUGAUUU	442	CAGGAACCC	1379	AGGAACCC	2316	GGAACCC	3253
hsa-miR-24	UGGCUCAGUUCAGCAGGAACAG	443	AACTGAGCC	1380	ACTGAGCC	2317	CTGAGCC	3254
hsa-miR-24-1*	UGCCUACUGAGCUGAUUACAGU	444	TCAGTAGGC	1381	CAGTAGGC	2318	AGTAGGC	3255
hsa-miR-24-2*	UGCCUACUGAGCUGAAACACAG	445	TCAGTAGGC	1382	CAGTAGGC	2319	AGTAGGC	3256
hsa-miR-25	CAUUGCACUUGUCUCGGUCUGA	446	AAGTGCAAT	1383	AGTGCAAT	2320	GTGCAAT	3257
hsa-miR-25*	AGGCGGAGACUUGGGCAAUUG	447	GTCTCCGCC	1384	TCTCCGCC	2321	CTCCGCC	3258
hsa-miR-26a	UUCAAGUAAUCCAGGAUAGGCU	448	ATTACTTGA	1385	TTACTTGA	2322	TACTTGA	3259
hsa-miR-26a-1*	CCUUAUUCUUGGUUACUUGCACG	449	CAAGAATAG	1386	AAGAATAG	2323	AGAATAG	3260
hsa-MIR-26a-2*	CCUUAUUCUUGAUUACUUGUUUC	450	CAAGAATAG	1387	AAGAATAG	2324	AGAATAG	3261

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-26b	UUCAAGUAAUUCAGGAUAGGU	451	ATTACTTGA	1388	TTACTTGA	2325	TACTTGA	3262
hsa-miR-26b*	CCUGUUCUCCAUUACUUGGCUC	452	GGAGAACAG	1389	GAGAACAG	2326	AGAACAG	3263
hsa-miR-27a	UUCACAGUGGCUAAGUCCGC	453	CCACTGTGA	1390	CACTGTGA	2327	ACTGTGA	3264
hsa-miR-27a*	AGGGCUUAGCUGCUUUGAGCA	454	GCTAAGCCC	1391	CTAAGCCC	2328	TAAGCCC	3265
hsa-miR-27b	UUCACAGUGGCUAAGUUCUGC	455	CCACTGTGA	1392	CACTGTGA	2329	ACTGTGA	3266
hsa-miR-27b*	AGAGCUUAGCUGAUUGGUAAC	456	GCTAAGCTC	1393	CTAAGCTC	2330	TAAGCTC	3267
hsa-miR-28-3p	CACUAGAUUGAGCUCUCCUGGA	457	CAATCTAGT	1394	AATCTAGT	2331	ATCTAGT	3268
hsa-miR-28-5p	AAGGAGCUCACAGUCUUAUUGAG	458	TGAGCTCCT	1395	GAGCTCCT	2332	AGCTCCT	3269
hsa-miR-296-3p	GAGGGUUGGGUGGAGGCUCUCC	459	CCCAACCCCT	1396	CCAACCCCT	2333	CAACCCCT	3270
hsa-miR-296-5p	AGGGCCCCCCCUCAAUCCUGU	460	GGGGGCCCC	1397	GGGGGCCCC	2334	GGGGCCC	3271
hsa-miR-297	AUGUAUGUGCAUGUGCAUG	461	ACACATACA	1398	CACATACA	2335	ACATACA	3272
hsa-miR-298	AGCAGAAGCAGGGAGGUUCUCCCA	462	TGCTTCTGC	1399	GCTTCTGC	2336	CTTCTGC	3273
hsa-miR-299-3p	UAUGUGGGAUGGUAAAACCGCUU	463	ATCCACAT	1400	TCCACAT	2337	CCACAT	3274
hsa-miR-299-Sp	UGGUUUACCGUCCCAUACAU	464	CGGTAAACC	1401	GGTAAACC	2338	GTAACC	3275
hsa-miR-29a	UAGCACCAUCUGAAAUCGGUUA	465	GATGGTGCT	1402	ATGGTGCT	2339	TGGTGCT	3276
hsa-miR-29a*	ACUGAUUUUUUUUGGUGUUCAG	466	AGAAATCAG	1403	GAAATCAG	2340	AAATCAG	3277
hsa-miR-29b	UAGCACCAUUUGAAAUCAGUGUU	467	AATGGTGCT	1404	ATGGTGCT	2341	TGGTGCT	3278
hsa-miR-29b-1*	GCUGGUUUCAUUGGUGGUUUAGA	468	TGAAACCAG	1405	GAAACCAG	2342	AAACCAG	3279
hsa-miR-29b-2*	CUGGUUUACAUUGGUGGUUUAG	469	GTGAAACCA	1406	TGAAACCA	2343	GAAACCA	3280
hsa-miR-29c	UAGCACCAUUUGAAAUCGGUUA	470	AATGGTGCT	1407	ATGGTGCT	2344	TGGTGCT	3281
hsa-miR-29c*	UGACCGAUUUUCUCCUGGUUUUC	471	AAATCGGTC	1408	AATCGGTC	2345	ATCGGTC	3282
hsa-miR-300	UAUACAAGGGCAGACUCUCUCU	472	CCCTTGAT	1409	CCTTGAT	2346	CTTGAT	3283
hsa-miR-301a	CAGUGCAAUAGUUAUUUGUCAAAGC	473	TATTGCACT	1410	ATTGCACT	2347	TTGCACT	3284

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-301b	CAGUGCAAUGAUUUUGUCAAAAGC	474	CATTGCACT	1411	ATTGCACT	2348	TTGCACT	3285
hsa-miR-302a	UAAUGUCUCCAUUUUUGGUGA	475	GAAGCACTT	1412	AAGCACTT	2349	AGCACTT	3286
hsa-miR-302a*	ACUAAAACGUGGAUGUACUUGCU	476	ACGTTTAAG	1413	CGTTTAAG	2350	GTTTAAG	3287
hsa-miR-302b	UAAUGUCUCCAUUUUUGUAG	477	GAAGCACTT	1414	AAGCACTT	2351	AGCACTT	3288
hsa-miR-302b*	ACUUUAACAUGGAAGUCUUC	478	ATGTTAAAG	1415	TGTTAAAG	2352	GTTAAAG	3289
hsa-miR-302c	UAAUGUCUCCAUUUUUGAGUGG	479	GAAGCACTT	1416	AAGCACTT	2353	AGCACTT	3290
hsa-miR-302c*	UUUAACAUGGGGUACCUUGCUG	480	CCATGTTAA	1417	CATGTTAA	2354	ATGTTAA	3291
hsa-miR-302d	UAAUGUCUCCAUUUUUGAGUGU	481	GAAGCACTT	1418	AAGCACTT	2355	AGCACTT	3292
hsa-miR-302d*	ACUUUAACAUGGAGGCACUUGC	482	ATGTTAAAG	1419	TGTTAAAG	2356	GTTAAAG	3293
hsa-miR-302e	UAAUGUCUCCAUUGCUU	483	GAAGCACTT	1420	AAGCACTT	2357	AGCACTT	3294
hsa-miR-302f	UAAUGUCUCCAUUGUUU	484	GAAGCAATT	1421	AAGCAATT	2358	AGCAATT	3295
hsa-miR-30a	UGUAAACAUCUCCAGCUGGAAG	485	GATGTTTAC	1422	ATGTTTAC	2359	TGTTTAC	3296
hsa-miR-30a*	CUUUCAGUCGGAUUUUUGCAGC	486	CGACTGAAA	1423	GACTGAAA	2360	ACTGAAA	3297
hsa-miR-30b	UGUAAACAUCUCCAGCUGCUGCU	487	GATGTTTAC	1424	ATGTTTAC	2361	TGTTTAC	3298
hsa-miR-30b*	CUGGGAGGUGGAUGUUUACUUC	488	CACCTCCCA	1425	ACCTCCCA	2362	CCTCCCA	3299
hsa-miR-30c	UGUAAACAUCUCCAGCUGCAGC	489	GATGTTTAC	1426	ATGTTTAC	2363	TGTTTAC	3300
hsa-miR-30c-1*	CUGGGAGGUGGUUUUACUUC	490	CCTCTCCCA	1427	CTCTCCCA	2364	TCTCCCA	3301
hsa-miR-30c-2*	CUGGGAGAAGGUCUUUUACUCU	491	CTTCTCCCA	1428	TTCTCCCA	2365	TCTCCCA	3302
hsa-miR-30d	UGUAAACAUCUCCAGCUGGAAG	492	GATGTTTAC	1429	ATGTTTAC	2366	TGTTTAC	3303
hsa-miR-30d*	CUUUCAGUCAGAUUUUUGCUGC	493	TGACTGAAA	1430	GACTGAAA	2367	ACTGAAA	3304
hsa-miR-30e	UGUAAACAUCUCCAGCUGGAAG	494	GATGTTTAC	1431	ATGTTTAC	2368	TGTTTAC	3305
hsa-miR-30e*	CUUUCAGUCGGAUGUUUACAGC	495	CGACTGAAA	1432	GACTGAAA	2369	ACTGAAA	3306
hsa-miR-31	AGGCAAGAUGCUGGCAUAGCU	496	CATCTTGCC	1433	ATCTTGCC	2370	TCTTGCC	3307

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-31*	UGCUAUGCCACAUAUUGCCAU	497	TGGCATAGC	1434	GGCATAGC	2371	GCATAGC	3308
hsa-miR-32	UAUUGCACAUUACUAAGUUGCA	498	ATGTGCAAT	1435	TGTGCAAT	2372	GTGCAAT	3309
hsa-miR-32*	CAUUUAGUGUGUGUAUUUU	499	CACTAAATT	1436	ACTAAATT	2373	CTAAATT	3310
hsa-miR-320a	AAAAGCUGGUUGAGAGGGCGA	500	CCCAGCTTT	1437	CCAGCTTT	2374	CAGCTTT	3311
hsa-miR-320b	AAAAGCUGGUUGAGAGGGCAA	501	CCCAGCTTT	1438	CCAGCTTT	2375	CAGCTTT	3312
hsa-miR-320c	AAAAGCUGGUUGAGAGGGGU	502	CCCAGCTTT	1439	CCAGCTTT	2376	CAGCTTT	3313
hsa-miR-320d	AAAAGCUGGUUGAGAGGGA	503	CCCAGCTTT	1440	CCAGCTTT	2377	CAGCTTT	3314
hsa-miR-323-3p	CACAUUACACGGUCGACCUCU	504	GTGTAATGT	1441	TGTAATGT	2378	GTAATGT	3315
hsa-miR-323-5p	AGGUGGUCGUGGGCGGUUCGC	505	CGGACCACC	1442	GGACCACC	2379	GACCACC	3316
hsa-miR-324-3p	ACUGCCCCAGGUGCUCUGG	506	CTGGGCAG	1443	TGGGCAG	2380	GGGCAG	3317
hsa-miR-324-5p	CGAUCCCAGGCAUUGGUGU	507	AGGGATGC	1444	GGGATGC	2381	GGATGC	3318
hsa-miR-325	CCUAGUAGGUGUCCAGUAAGUGU	508	ACCTACTAG	1445	CCTACTAG	2382	CTACTAG	3319
hsa-miR-326	CCUCUGGGCCCUCCUCCAG	509	GGCCCAGAG	1446	GCCCAGAG	2383	CCCAGAG	3320
hsa-miR-328	CUGGCCUCUCUGCCCCUCCGU	510	AGAGGGCCA	1447	GAGGGCCA	2384	AGGGCCA	3321
hsa-miR-329	AACACACCUUGUUAAACCUCUUU	511	CAGGTGTGT	1448	AGGTGTGT	2385	GGTGTGT	3322
hsa-miR-330-3p	GCAAAGCACACGGCCUGCAGAGA	512	TGTGCTTTG	1449	GTGCTTTG	2386	TGCTTTG	3323
hsa-miR-330-5p	UCUCUGGGCCUGUGUCUUAGGC	513	GGCCCAGAG	1450	GCCCAGAG	2387	CCCAGAG	3324
hsa-miR-331-3p	GCCCCUGGGCCUAUCCUAGAA	514	GCCCCAGGG	1451	CCCAGGG	2388	CCAGGG	3325
hsa-miR-331-5p	CUAGGUAGUCCAGGGAUCC	515	CCATACCTA	1452	CATACCTA	2389	ATACCTA	3326
hsa-miR-335	UCAAGAGCAUAACGAAAAAUGU	516	TTGCTCTTG	1453	TGCTCTTG	2390	GCTCTTG	3327
hsa-miR-335*	UUUUUCAUUUUUCCUCCGACC	517	TAATGAAAA	1454	AATGAAAA	2391	ATGAAAA	3328
hsa-miR-337-3p	CUCCUAUAUGAUGCCUUUCUUC	518	CATATAGGA	1455	ATATAGGA	2392	TATAGGA	3329
hsa-miR-337-5p	GAACGGCUUCAUACAGGAGUU	519	GAAGCCGTT	1456	AAGCCGTT	2393	AGCCGTT	3330



(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-338-3p	UCCAGCAUCAGUAUUUUGUUG	520	TGATGCTGG	1457	GATGCTGG	2394	ATGCTGG	3331
hsa-miR-338-5p	AACAUAUCCUGGUGCUGAGUG	521	GGATATTGT	1458	GATATTGT	2395	ATATTGT	3332
hsa-miR-339-3p	UGAGCGCCUCGACGACAGAGCCCG	522	GAGCGCTC	1459	AGGCGCTC	2396	GGCGCTC	3333
hsa-miR-339-5p	UCCUGUCCUCCAGGAGCUCACG	523	AGGACAGGG	1460	GGACAGGG	2397	GACAGGG	3334
hsa-miR-33a	GUGCAUUGUAGUUGCAUUGCA	524	TACAATGCA	1461	ACAATGCA	2398	CAATGCA	3335
hsa-miR-33a*	CAUUGUUCCACAGUGCAUCAC	525	GGAACATT	1462	GAAACATT	2399	AAACATT	3336
hsa-miR-33b	GUGCAUUGCUGUUGCAUUGC	526	AGCAATGCA	1463	GCAATGCA	2400	CAATGCA	3337
hsa-miR-33b*	CAGUGCCUCGGCAGUGCAGCCC	527	CGAGGCACT	1464	GAGGCACT	2401	AGGCACT	3338
hsa-miR-340	UUUAAAAGCAAUGAGACUGAUU	528	TGCTTTATA	1465	GCTTTATA	2402	CTTTATA	3339
hsa-miR-340*	UCCGUCUCAGUUACUUUUAJAGC	529	CTGAGACGG	1466	TGAGACGG	2403	GAGACGG	3340
hsa-miR-342-3p	UCUCACACAGAAUUCGACCCGU	530	CTGTGTGAG	1467	TGTGTGAG	2404	GTGTGAG	3341
hsa-miR-342-5p	AGGGGUGCUAUCUGUAGUUGA	531	TAGCACCCC	1468	AGCACCCC	2405	GCACCCC	3342
hsa-miR-345	GCUGACUCCUAGUCCAGGGCUC	532	AGGAGTCAG	1469	GGAGTCAG	2406	GAGTCAG	3343
hsa-miR-346	UGUCUGCCCGCAUGCCUGCCUCU	533	CGGGCAGAC	1470	GGGCAGAC	2407	GGCAGAC	3344
hsa-miR-34a	UGGCAGUGUCUAGCUGGUUGU	534	GACACTGCC	1471	ACACTGCC	2408	CACTGCC	3345
hsa-miR-34a*	CAAUCAGCAAGUAUACUGCCCU	535	TTGCTGATT	1472	TGCTGATT	2409	GCTGATT	3346
hsa-miR-34b	CAAUCACUAAUCCACUGCCAU	536	TTAGTGATT	1473	TAGTGATT	2410	AGTGATT	3347
hsa-miR-34b*	UAGGCAGUGUCAUJAGCUGAUUG	537	ACACTGCCT	1474	CACTGCCT	2411	ACTGCCT	3348
hsa-miR-34c-3p	AAUCACUAAACCACACGCCCAGG	538	GTTAGTGAT	1475	TTAGTGAT	2412	TAGTGAT	3349
hsa-miR-34c-5p	AGGCAGUGUAGUJAGCUGAUUGC	539	TACACTGCC	1476	ACACTGCC	2413	CACTGCC	3350
hsa-miR-361-3p	UCCCCAGGUGUAGUUCUGAUUU	540	ACCTGGGGG	1477	CCTGGGGG	2414	CTGGGGG	3351
hsa-miR-361-5p	UUUUCAGAAUUCUCCAGGGGUAC	541	ATTCTGATA	1478	TTCTGATA	2415	TCTGATA	3352
hsa-miR-362-3p	AACACACCUAUUCAAGGAUUCA	542	TAGGTGTGT	1479	AGGTGTGT	2416	GGTGTGT	3353

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-362-5p	AAUCCUUGGAACCUAGGUGUGAGU	543	TCCAAGGAT	1480	CCAAGGAT	2417	CAAGGAT	3354
hsa-miR-363	AAUUGCACGGUAUCCAUUCUGUA	544	CCGTGCAAT	1481	CGTGCAAT	2418	GTGCAAT	3355
hsa-miR-363*	CGGGUGGAUCACGAUGCAAUUU	545	GATCCACCC	1482	ATCCACCC	2419	TCCACCC	3356
hsa-miR-365	UAAUGCCCCUAAAAUCCUUAU	546	AGGGGCATT	1483	GGGGCATT	2420	GGGCATT	3357
hsa-miR-367	AAUUGCACUUUAGCAAUGGUGA	547	AAGTGCAAT	1484	AGTGCAAT	2421	GTGCAAT	3358
hsa-miR-367*	ACUGUUGCUAUUUUGCAACUCU	548	TAGCAACAG	1485	AGCAACAG	2422	GCAACAG	3359
hsa-miR-369-3p	AAUAAUACAUGGUUGAUUCUUU	549	ATGTATTAT	1486	TGTATTAT	2423	GTATTAT	3360
hsa-miR-369-5p	AGAUGCACCGUGUUAUUAUUCGC	550	CGGTCGATC	1487	GGTCGATC	2424	GTCGATC	3361
hsa-miR-370	GCCUCUGGGGUGGAACCCUGGU	551	CCCAGCAGG	1488	CCAGCAGG	2425	CAGCAGG	3362
hsa-miR-371-3p	AAGUCCGCCCAUCUUUUGAGUGU	552	GGCGGCACT	1489	GCGGCACT	2426	CGGCACT	3363
hsa-miR-371-5p	ACUCAACUGUGGGGGCACU	553	CAGTTTGAG	1490	AGTTTGAG	2427	GTTTGAG	3364
hsa-miR-372	AAAGUGCUCGACAUUUGAGCGU	554	GCAGCACTT	1491	CAGCACTT	2428	AGCACTT	3365
hsa-miR-373	GAAGUCUUCGAUUUUGGGGUGU	555	GAAGCACTT	1492	AAGCACTT	2429	AGCACTT	3366
hsa-miR-373*	ACUCAAAUUGGGGGCGCUUCC	556	CATTTTGAG	1493	ATTTTGAG	2430	TTTTGAG	3367
hsa-miR-374a	UUUAAUACAACCUUGAUUAGUG	557	TGTATTATA	1494	GTATTATA	2431	TATTATA	3368
hsa-miR-374a*	CUUAUCAGAUUGUAUUUGUAAUU	558	ATCTGATAA	1495	TCTGATAA	2432	CTGATAA	3369
hsa-miR-374b	AUAUAAUACAACCUUGCUAAGUG	559	TGTATTATA	1496	GTATTATA	2433	TATTATA	3370
hsa-miR-374b*	CUUAGCAGGUUGUAUUUAUCAUU	560	ACCTGCTAA	1497	CCTGCTAA	2434	CTGCTAA	3371
hsa-miR-375	UUUGUUCGUUCGGCCUCGCGUGA	561	AACGAACAA	1498	ACGAACAA	2435	CGAACAA	3372
hsa-miR-376a	AUCAUAGAGGAAAAUCCACGU	562	CCTCTATGA	1499	CTCTATGA	2436	TCTATGA	3373
hsa-miR-376a*	GUAGAUUCUCCUUUAUGAGUA	563	GAGAATCTA	1500	AGAATCTA	2437	GAATCTA	3374
hsa-miR-376b	AUCAUAGAGGAAAAUCCAUUUU	564	CCTCTATGA	1501	CTCTATGA	2438	TCTATGA	3375
hsa-miR-376c	AACAUAAGAGGAAAAUCCACGU	565	CCTCTATGT	1502	CTCTATGT	2439	TCTATGT	3376

(continued)

microRNA	MicroRNASequences	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-377	AUCACACAAAGGCAACUUUUUGU	566	TTTGTGTGA	1503	TTGTGTGA	2440	TGTGTGA	3377
hsa-miR-377*	AGAGGUUGCCCUUGGUGAAUUC	567	GGCAACCTC	1504	GCAACCTC	2441	CAACCTC	3378
hsa-miR-378	ACUGGACUUGGAGUCAGAAAGG	568	CAAGTCCAG	1505	AAGTCCAG	2442	AGTCCAG	3379
hsa-miR-378*	CUCCUGACUCCAGGUCCUGUGU	569	GAGTCAGGA	1506	AGTCAGGA	2443	GTCAGGA	3380
hsa-miR-379	UGGUAGACUAUGGAACGUAGG	570	TAGTCTACC	1507	AGTCTACC	2444	GTCTACC	3381
hsa-miR-379*	UAUGUAACAUGGUCCACUAACU	571	ATGTTACAT	1508	TGTTACAT	2445	GTTACAT	3382
hsa-miR-380	UAUGUAAUUGGUCCACAUCUU	572	ATATTACAT	1509	TATTACAT	2446	ATTACAT	3383
hsa-miR-380*	UGGUUGACCAUAGAACAUGCGC	573	TGGTCAACC	1510	GGTCAACC	2447	GTC AACC	3384
hsa-miR-381	UAUACAAGGGCAAGCUCUCUGU	574	CCCTTGAT	1511	CCTTGAT	2448	CTTGAT	3385
hsa-miR-382	GAAGUUGUUCGUGGUGGAUUCG	575	GAACAACCT	1512	AACAACCT	2449	ACAACCT	3386
hsa-miR-383	AGAUCAGAAGGUGAUUGUGGCU	576	CTTCTGATC	1513	TTCTGATC	2450	TCTGATC	3387
hsa-miR-384	AUUCUAGAAUUUUUCAUA	577	TTCTAGGAA	1514	TCTAGGAA	2451	CTAGGAA	3388
hsa-miR-409-3p	GAAUGUUGCUCGGUGAACCCCU	578	AGCAACATT	1515	GCAACATT	2452	CAACATT	3389
hsa-miR-409-5p	AGGUUACCCGAGCAACUUUGCAU	579	CGGTAACC	1516	GGTAACC	2453	GGTAACC	3390
hsa-miR-410	AAUUAACACAGAUAGCCUGU	580	GTGTTATAT	1517	TGTTATAT	2454	GTTATAT	3391
hsa-miR-411	UAGUAGACCGUAUAGCGUACG	581	CGGTCTACT	1518	GGTCTACT	2455	GTCTACT	3392
hsa-miR-411*	UAUGUAACACGGUCCACUAACC	582	GTGTTACAT	1519	TGTTACAT	2456	GTTACAT	3393
hsa-miR-412	ACUUCACCUUGUCCACUAGCCGU	583	CAGGTGAAG	1520	AGGTGAAG	2457	GGTGAAG	3394
hsa-miR-421	AUCAACAGACAUUAAUUUGGCGC	584	GTCTGTTGA	1521	TCTGTTGA	2458	CTGTTGA	3395
hsa-miR-422a	ACUGGACUUAGGGUCAGAAAGC	585	TAAGTCCAG	1522	AAGTCCAG	2459	AGTCCAG	3396
hsa-miR-423-3p	AGCUCGGUCUGAGGCCCCUCAGU	586	AGACCGAGC	1523	GACCGAGC	2460	ACCGAGC	3397
hsa-miR-423-5p	UGAGGGGCAGAGAGCGGAGACUUU	587	CTGCCCTC	1524	TGCCCTC	2461	GCCCTC	3398
hsa-miR-424	CAGCAGCAUUUCAUGUUUUGAA	588	ATTGCTGCT	1525	TTGCTGCT	2462	TGCTGCT	3399

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-424*	CAAAAACGUGAGGCGCUGCUAU	589	TCACGTTTT	1526	CACGTTTT	2463	ACGTTTT	3400
hsa-miR-425	AAUGACACGAUCACUCCCGUUGA	590	TCGTGTTCAT	1527	CGTGTTCAT	2464	GTGTTCAT	3401
hsa-miR-425*	AUCGGAAUGUCGUGUCCGCCCC	591	CATCCCGA	1528	ATCCCGA	2465	TTCCCGA	3402
hsa-miR-429	UAAUACUGUCUGGUAAAACCGU	592	GACAGTATT	1529	ACAGTATT	2466	CAGTATT	3403
hsa-miR-431	UGUCUUGCAGGCCCGUCAUGCA	593	CTGCAAGAC	1530	TGCAAGAC	2467	GCAAGAC	3404
hsa-miR-431*	CAGGUCGUCUUGCAGGGCUUCU	594	AGACGACCT	1531	GACGACCT	2468	ACGACCT	3405
hsa-miR-432	UCUUGGAGUAGGUCAUUGGGUGG	595	TACTCCAAG	1532	ACTCCAAG	2469	CTCCAAG	3406
hsa-miR-432*	CUGGAUGGCCUCCUCCAUGUCU	596	AGCCATCCA	1533	GCCATCCA	2470	CCATCCA	3407
hsa-miR-433	AUCAUGAUGGGCUCUCCGGUGU	597	CCATCATGA	1534	CATCATGA	2471	ATCATGA	3408
hsa-miR-448	UUGCAUUGAAGGAUGUCCCAU	598	ACATATGCA	1535	CATATGCA	2472	ATATGCA	3409
hsa-miR-449a	UGGCAGUGUAUUGUUAGCUGGU	599	TACACTGCC	1536	ACACTGCC	2473	CACTGCC	3410
hsa-miR-449b	AGGCAGUGUAUUGUUAGCUGGC	600	TACACTGCC	1537	ACACTGCC	2474	CACTGCC	3411
hsa-miR-450a	UUUUGCGAUGUUUCCUAAUUAU	601	CATGCAAA	1538	ATGCAAA	2475	TCGCAAA	3412
hsa-miR-450b-3p	UUGGGAUCAUUUUGCAUCCAU	602	ATGATCCCA	1539	TGATCCCA	2476	GATCCCA	3413
hsa-miR-450b-5p	UUUUGCAAUAUGUUUCCUGAAUA	603	TATTGCAAA	1540	ATTGCAAA	2477	TTGCAAA	3414
hsa-miR-451	AAACCGUUACCAUUACUGAGUU	604	GTAACGGTT	1541	TAACGGTT	2478	AACGGTT	3415
hsa-miR-452	AACUGUUUGCAGAGGAAACUGA	605	GCAAAACAGT	1542	CAAAACAGT	2479	AAACAGT	3416
hsa-miR-452*	CUCAUCUGCAAAGAAGUAAGUG	606	TGCAGATGA	1543	GCAGATGA	2480	CAGATGA	3417
hsa-miR-453	AGGUUGUCCGUGGUGAGUUUGCA	607	CGGACAACC	1544	GGACAACC	2481	GACAACC	3418
hsa-miR-454	UAGUGCAAUUUUGCUUUAJAGGGU	608	TATTGCACT	1545	ATTGCACT	2482	TTGCACT	3419
hsa-miR-454*	ACCCUAUCAUUUUGUCUCUCUGC	609	TTGATAGGG	1546	TGATAGGG	2483	GATAGGG	3420
hsa-miR-455-3p	GCAGUCCAUGGGCAUUAJACAC	610	CATGGACTG	1547	ATGGACTG	2484	TGGACTG	3421
hsa-miR-455-5p	UAUGUGCCUUUUGGACUACAUCG	611	AAGGCACAT	1548	AGGCACAT	2485	GGCACAT	3422

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-483-3p	UCACUCCUCUCUCCUCCGUCUU	612	AGAGGAGTG	1549	GAGGAGTG	2486	AGGAGTG	3423
hsa-miR-483-5p	AAGACGGGAGGAAAAGAGGGAG	613	CTCCCGTCT	1550	TCCCGTCT	2487	CCCGTCT	3424
hsa-miR-484	UCAGGCUACAGUCCUCCUCCGGAU	614	CTGAGCCTG	1551	TGAGCCTG	2488	GAGCCTG	3425
hsa-miR-485-3p	GUAUACACGGCUCUCCUCUCU	615	CGTGATGA	1552	GTGTATGA	2489	TGTATGA	3426
hsa-miR-485-5p	AGAGGCUCCGUGAUGAAUUC	616	GCCAGCCTC	1553	CCAGCCTC	2490	CAGCCTC	3427
hsa-miR-486-3p	CGGGCAGCUCAGUACAGGAU	617	AGTGCCCC	1554	GCTGCCCC	2491	CTGCCCC	3428
hsa-miR-486-5p	UCCUGUACUGAGCUGCCCCGAG	618	CAGTACAGG	1555	AGTACAGG	2492	GTACAGG	3429
hsa-miR-487a	AAUCAUACAGGGACAUCCAGUU	619	CTGTATGAT	1556	TGTATGAT	2493	GTATGAT	3430
hsa-miR-487b	AAUCGUACAGGGUCAUCCACUU	620	CTGTACGAT	1557	TGTACGAT	2494	GTACGAT	3431
hsa-miR-488	UUGAAAGGCUAUUUUUGGUC	621	AGCCTTTCA	1558	GCCTTTCA	2495	CCTTTCA	3432
hsa-miR-488*	CCCAGAUAAUGGCACUCUCAA	622	ATTATCTGG	1559	TTATCTGG	2496	TATCTGG	3433
hsa-miR-489	GUGACAUACAUUACGGCAGC	623	GTGATGTCA	1560	TGATGTCA	2497	GATGTCA	3434
hsa-miR-490-3p	CAACCUGGAGGACUCCAUUGCUG	624	CTCCAGGTT	1561	TCCAGGTT	2498	CCAGGTT	3435
hsa-miR-490-5p	CCAUGGAUCUCCAGGUGGGU	625	AGATCCATG	1562	GATCCATG	2499	ATCCATG	3436
hsa-miR-491-3p	CUUAUGCAAGAUUCCUUCUAC	626	CTTGATATA	1563	TTGCATAA	2500	TGCATAA	3437
hsa-miR-491-5p	AGUGGGGAACCCUUCUCCAUAGAGG	627	GTTCCCCAC	1564	TTCCCCAC	2501	TCCCCAC	3438
hsa-miR-492	AGGACCUCCGGGACAAAGAUUCUU	628	CGCAGGTCC	1565	GCAGGTCC	2502	CAGGTCC	3439
hsa-miR-493	UGAAGGUCUACUGUGCCAGG	629	TAGACCTTC	1566	AGACCTTC	2503	GACCTTC	3440
hsa-miR-493*	UUGUACAUUGGUAGGCUUUCUU	630	CCATGTACA	1567	CATGTACA	2504	ATGTACA	3441
hsa-miR-494	UGAAACAUAACACGGGAAACCUC	631	GTATGTTTC	1568	TATGTTTC	2505	ATGTTTC	3442
hsa-miR-495	AAACAAACAUGGUGCACUUCUU	632	ATGTTTGT	1569	TGTTTGT	2506	GTTTGT	3443
hsa-miR-496	UGAGUAUUACAUGGCCAAUCUC	633	GTAATACTC	1570	TAATACTC	2507	AATACTC	3444
hsa-miR-497	CAGCAGCACACUGUGGUUUUGU	634	TGTGCTGCT	1571	GTGCTGCT	2508	TGCTGCT	3445

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-497*	CAAACCCACACUGUGGUGUAGA	635	GTGTGGTTT	1572	TGTGGTTT	2509	GTGGTTT	3446
hsa-miR-498	UUUCAAGCCAGGGGGUUUUUC	636	TGGCTTGAA	1573	GGCTTGAA	2510	GCTTGAA	3447
hsa-miR-499-3p	AACAUCACAGCAAGUCUGUCU	637	CTGTGATGT	1574	TGTGATGT	2511	GTGATGT	3448
hsa-miR-499-5p	UUAGACUUGCAGUGAUUUU	638	CAAGTCTTA	1575	AAGTCTTA	2512	AGTCTTA	3449
hsa-miR-500	UAAUCCUUGCACCCUGGGUGAGA	639	GCAAGGATT	1576	CAAGGATT	2513	AAGGATT 3450	
hsa-miR-500*	AUGCACCUGGGCAAGGAUUCUG	640	CCAGGTGCA	1577	CAGGTGCA	2514	AGGTGCA	3451
hsa-miR-501-3p	AAUGCACCCGGGCAAGGAUUCU	641	CGGGTGCAT	1578	GGGTGCAT	2515	GGTGCAT	3452
hsa-miR-501-5p	AAUCCUUUGUCCUUGGGUGAGA	642	ACAAAGGAT	1579	CAAAGGAT	2516	AAAGGAT	3453
hsa-miR-502-3p	AAUGCACCUGGGCAAGGAUUCU	643	CAGGTGCAT	1580	AGGTGCAT	2517	GGTGCAT	3454
hsa-miR-502-5p	AUCCUUGCUAUCUGGGUCUA	644	TAGCAAGGA	1581	AGCAAGGA	2518	GCAAGGA	3455
hsa-miR-503	UAGCAGCGGGAACAGUUCUCGAG	645	CCCGTGCT	1582	CCGCTGCT	2519	CGCTGCT	3456
hsa-miR-504	AGACCCUGGUCUGCACUCUAUC	646	ACCAGGTC	1583	CCAGGTC	2520	CAGGTC	3457
hsa-miR-505	CGUCAACACUUGCUGGUUCCU	647	AGTGTGAC	1584	GTGTGAC	2521	TGTTGAC	3458
hsa-miR-505*	GGGAGCCAGGAAGUAUUGAUGU	648	CCTGGCTCC	1585	CTGGCTCC	2522	TGGCTCC	3459
hsa-miR-506	UAAAGCACCCUUCUGAGUAGA	649	GGGTGCCTT	1586	GGTGCCTT	2523	GTGCCTT	3460
hsa-miR-507	UUUUGCACCCUUUGGAGUGAA	650	AGGTGCAAA	1587	GGTGCAAA	2524	GTGCAAA	3461
hsa-miR-508-3p	UGAUUGUAGCCUUUUGGAGUAGA	651	GCTACAATC	1588	CTACAATC	2525	TACAATC	3462
hsa-miR-508-5p	UACUCCAGAGGGGUCACUCAUG	652	CTCTGGAGT	1589	TCTGGAGT	2526	CTGGAGT	3463
hsa-miR-509-3-5p	UACUGCAGACGUGGCAAUCAUG	653	GTCTGCAGT	1590	TCTGCAGT	2527	CTGCAGT	3464
hsa-miR-509-3p	UGAUUGGUACGUCUGGGUAG	654	GTACCAATC	1591	TACCAATC	2528	ACCAATC	3465
hsa-miR-509-5p	UACUGCAGACAGUGGCAAUCA	655	GTCTGCAGT	1592	TCTGCAGT	2529	CTGCAGT	3466
hsa-miR-510	UACUCAGGAGAGUGGCAAUCA	656	CTCCTGAGT	1593	TCCTGAGT	2530	CCTGAGT	3467
hsa-miR-511	GUGUCUUUUGCUCUGCAGUCA	657	CAAAAGACA	1594	AAAAGACA	2531	AAAGACA	3468

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-512-3p	AAGUGCUGUCAUAGCUGAGGUC	658	GACAGCACT	1595	ACAGCACT	2532	CAGCACT	3469
hsa-miR-512-5p	CACUCAGCCUUGAGGGCACUUUC	659	AGGCTGAGT	1596	GGCTGAGT	2533	GCTGAGT	3470
hsa-miR-513a-3p	UAAUUUACCCUUUCUGAGAAGG	660	GTGAAATTT	1597	TGAAATTT	2534	GAAATTT	3471
hsa-miR-513a-5p	UUCACAGGGAGGUGUCAU	661	TCCTGTGA	1598	CCCTGTGA	2535	CCTGTGA	3472
hsa-miR-513b	UUCACAAGGAGGUGUCAUUUUAU	662	TCCTTGTGA	1599	CCTTGTGA	2536	CTTGTGA	3473
hsa-miR-513c	UUCUCAAGGAGGUGUCGUUUUAU	663	TCCTTGTGA	1600	CCTTGTGA	2537	CTTGTGA	3474
hsa-miR-514	AUUGACACUUCUGUGAGUAGA	664	AAGTGCAA	1601	AGTGCAA	2538	GTGTCAA	3475
hsa-miR-515-3p	GAGUGCCUUUUUUGGAGCGUU	665	GAAGGCACT	1602	AAGGCACT	2539	AGGCACT	3476
hsa-miR-515-5p	UUCUCCAAAAGAAAGCACUUUCUG	666	TTTTGGAGA	1603	TTTGGAGA	2540	TTGGAGA	3477
hsa-miR-516a-3p	UGCUUCCUUUCAGAGGGU	667	AAAGAAAGC	1604	AAGAAAGC	2541	AGAAAGC	3478
hsa-miR-516a-5p	UUCUCGAGGAAAGAACACUUUC	668	TCCTCGAGA	1605	CCTCGAGA	2542	CTCGAGA	3479
hsa-miR-516b	AUCUGGAGGUAAGAAAGCACUUU	669	ACCTCCAGA	1606	CCTCCAGA	2543	CTCCAGA	3480
hsa-miR-517*	CCUCUAGAUGGAAAGCACUGUCU	670	CATCTAGAG	1607	ATCTAGAG	2544	TCTAGAG	3481
hsa-miR-517a	AUCGUGCAUCCUUUJAGAGUGU	671	GATGCACGA	1608	ATGCACGA	2545	TGCACGA	3482
hsa-miR-517b	UCGUGCAUCCUUUJAGAGUGUU	672	GGATGCACG	1609	GATGCACG	2546	ATGCACG	3483
hsa-miR-517c	AUCGUGCAUCCUUUJAGAGUGU	673	GATGCACGA	1610	ATGCACGA	2547	TGCACGA	3484
hsa-miR-518a-3p	GAAAGCGCUUCCUUUGCUGGA	674	AAGCGCTTT	1611	AGCGCTTT	2548	GCGCTTT	3485
hsa-miR-518b	CAAAAGCGCUUCCUUUJAGAGGU	675	GAGCGCTTT	1612	AGCGCTTT	2549	GCGCTTT	3486
hsa-miR-518c	CAAAAGCGCUUUCUUUJAGAGUGU	676	AAGCGCTTT	1613	AGCGCTTT	2550	GCGCTTT	3487
hsa-miR-518c*	UCUCUGGAGGGAAGCACUUUCUG	677	CCTCCAGAG	1614	CTCCAGAG	2551	TCCAGAG	3488
hsa-miR-518d-3p	CAAAAGCGCUUCCUUUJGGAGC	678	AAGCGCTTT	1615	AGCGCTTT	2552	GCGCTTT	3489
hsa-miR-518d-5p	CUCUAGAGGGAAGCACUUUCUG	679	CCCTCTAGA	1616	CCTCTAGA	2553	CTCTAGA	3490
hsa-miR-518e	AAAGCGCUUCCUUUCAGAGUG	680	GAAGCGCTT	1617	AAGCGCTT	2554	AGCGCTT	3491

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-518f	GAAAGCGCUUCUCUUUAGAGGG	681	AAGCGCTTT	1618	AGCGCTTT	2555	GCGCTTT	3492
hsa-miR-518f*	CUCUAGAGGGAAGCACUUUCUC	682	CCCTCTAGA	1619	CCTCTAGA	2556	CTCTAGA	3493
hsa-miR-519a	AAAGUGCAUCCUUUUUAGAGUGU	683	GATGCACTT	1620	ATGCACTT	2557	TGCACTT	3494
hsa-miR-519a*	CUCUAGAGGGAAGCGCUUUUCUG	684	CCCTCTAGA	1621	CCTCTAGA	2558	CTCTAGA	3495
hsa-miR-519b-3p	AAAGUGCAUCCUUUUUAGAGGUU	685	GATGCACTT	1622	ATGCACTT	2559	TGCACTT	3496
hsa-miR-519c-3p	AAAGUGCAUCCUUUUUAGAGGAU	686	GATGCACTT	1623	ATGCACTT	2560	TGCACTT	3497
hsa-miR-519d	CAAAGUGCCUCCUUUUUAGAGUG	687	AGGCACTTT	1624	GGCACTTT	2561	GCACCTT	3498
hsa-miR-519e	AAGUGCCUCCUUUUUAGAGUGUU	688	GGAGGCACT	1625	GAGGCACT	2562	AGGCACT	3499
hsa-miR-519e*	UUCUCCAAAAGGGAGCACUUUC	689	TTTTGGAGA	1626	TTTGGAGA	2563	TTGGAGA	3500
hsa-miR-520a-3p	AAAGUGCUUCCUUUUGGACUGU	690	GAAGCACTT	1627	AAGCACTT	2564	AGCACTT	3501
hsa-miR-520a-5p	CUCCAGAGGGAAGUACUUUCU	691	CCCTCTGGA	1628	CCTCTGGA	2565	CTCTGGA	3502
hsa-miR-520b	AAAGUGCUUCCUUUUUAGAGGG	692	GAAGCACTT	1629	AAGCACTT	2566	AGCACTT	3503
hsa-miR-520c-3p	AAAGUGCUUCCUUUUUAGAGGGU	693	GAAGCACTT	1630	AAGCACTT	2567	AGCACTT	3504
hsa-miR-520d-3p	AAAGUGCUUCCUUUUGGUGGGU	694	GAAGCACTT	1631	AAGCACTT	2568	AGCACTT	3505
hsa-miR-520d-5p	CUACAAAGGGGAAGCCUUUC	695	CCCTTTGTA	1632	CCTTTGTA	2569	CTTTGTA	3506
hsa-miR-520e	AAAGUGCUUCCUUUUUAGAGGG	696	GAAGCACTT	1633	AAGCACTT	2570	AGCACTT	3507
hsa-miR-520f	AAGUGCUUCCUUUUUAGAGGGUU	697	GGAAGCACT	1634	GAAGCACT	2571	AAGCACT	3508
hsa-miR-520g	ACAAAGUGCUUCCUUUAGAGUGU	698	AGCACTTTG	1635	GCACTTTG	2572	CACTTTG	3509
hsa-miR-520h	ACAAAGUGCUUCCUUUAGAGU	699	AGCACTTTG	1636	GCACTTTG	2573	CACTTTG	3510
hsa-miR-521	AACGCACUCCUUUUUAGAGUGU	700	GAAGTCCGT	1637	AAGTCCGT	2574	AGTCCGT	3511
hsa-miR-522	AAAAUGGUUCCUUUUUAGAGUGU	701	GAACCATTT	1638	AACCATTT	2575	ACCATTT	3512
hsa-miR-523	GAACGCGCUUCCUUUAGAGGGU	702	AAGCGCGTT	1639	AGCGCGTT	2576	GCGCGTT	3513
hsa-miR-524-3p	GAAGGCGCUUCCUUUUGGAGU	703	AAGCGCCTT	1640	AGCGCCTT	2577	GCGCCTT	3514



5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-524-5p	CUACAAAGGGGAAGCACUUUCUC	704	CCCTTTGTA	1641	CCTTTGTGTA	2578	CTTTGTA	3515
hsa-miR-525-3p	GAAGGGCUUCCUUUAGAGCG	705	AAGCGCCTT	1642	AGCGCCTT	2579	GCGCCTT	3516
hsa-miR-525-5p	CUCCAGAGGGAUGCACUUUCU	706	CCCTCTGGA	1643	CCTCTGGA	2580	CTCTGGA	3517
hsa-miR-526b	CUCUUGAGGGAAGCACUUUCUGU	707	CCCTCAAGA	1644	CCTCAAGA	2581	CTCAAGA	3518
hsa-miR-526b*	GAAAGUCUUCCUUUJAGAGGC	708	AAGCACTTT	1645	AGCACTTT	2582	GCACCTT	3519
hsa-miR-527	CUGCAAAGGGGAAGCCUUUC	709	CCCTTTGCA	1646	CCTTTGCA	2583	CTTTGCA	3520
hsa-miR-532-3p	CCUCCACACCCAAAGCCUUGCA	710	GTGTGGGAG	1647	TGTGGGAG	2584	GTGGGAG	3521
hsa-miR-532-5p	CAUGCCUUGAGUGUAGGACCGU	711	TCAAAGCAT	1648	CAAGGCAT	2585	AAGGCAT	3522
hsa-miR-539	GGAGAAUUUCCUUUGGUGUGU	712	TAATTTCTC	1649	AATTTCTC	2586	ATTTCTC	3523
hsa-miR-541	UGGUGGCACAGAAUCUGGACU	713	GTGCCACC	1650	TGCCACC	2587	GCCACC	3524
hsa-miR-541*	AAAGAUUCUGCUGUGCGGCCAC U	714	AGAATCCTT	1651	GAATCCTT	2588	AATCCTT	3525
hsa-miR-542-3p	UGUGACAGAUUGAUAAUCUGAAA	715	ATCTGTCAC	1652	TCTGTCAC	2589	CTGTCAC	3526
hsa-miR-542-5p	UCGGGGAUCAUCAUGUCACGAGA	716	TGATCCCCG	1653	GATCCCCG	2590	ATCCCCG	3527
hsa-miR-543	AAACAUUCGGGUGCACUUCUU	717	GCGAATGTT	1654	CGAATGTT	2591	GAATGTT	3528
hsa-miR-544	AUUCUGCAUUUUUAGCAAGUUC	718	AATGCAGAA	1655	ATGCAGAA	2592	TGCAGAA	3529
hsa-miR-545	UCAGCAAACAUUUUUUGUGUGC	719	TGTTTGCTG	1656	GTTTGCTG	2593	TTTGCTG	3530
hsa-miR-545*	UCAGUAAAUGUUUUUAGAUGA	720	CATTTACTG	1657	ATTTACTG	2594	TTTACTG	3531
hsa-miR-548a-3p	CAAAACUGGCAUUACUUUUGC	721	GCCAGTTTT	1658	CCAGTTTT	2595	CAGTTTT	3532
hsa-miR-548a-5p	AAAAGUAAUUGCGAGUUUJACC	722	AATTACTTT	1659	ATTACTTT	2596	TTACTTT	3533
hsa-miR-548b-3p	CAAGAACCUCAGUUUGCUUUUGU	723	GAGGTTCTT	1660	AGGTTCTT	2597	GGTTCTT	3534
hsa-miR-548b-5p	AAAAGUAAUUGGUUUUGGCC	724	AATTACTTT	1661	ATTACTTT	2598	TTACTTT	3535
hsa-miR-548c-3p	CAAAAAUCUCAAUUACUUUUGC	725	GAGATTTTT	1662	AGATTTTT	2599	GATTTTT	3536
hsa-miR-548c-5p	AAAAGUAAUUGCGGUUUUUGCC	726	AATTACTTT	1663	ATTACTTT	2600	TTACTTT	3537

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-548d-3p	CAAAAACCCACAGUUUUUUUUGC	727	GTGGTTTTT	1664	TGGTTTTT	2601	GGTTTTT	3538
hsa-miR-548d-5p	AAAAAGUAAUUGUGUUUUUUGCC	728	AATTACTTT	1665	ATTACTTT	2602	TTACTTT	3539
hsa-miR-548e	AAAAACUGAGACUACUUUUUGCA	729	CTCAGTTTT	1666	TCAGTTTT	2603	CAGTTTT	3540
hsa-miR-548f	AAAAACUGOAAUUACUUUUU	730	TACAGTTTT	1667	ACAGTTTT	2604	CAGTTTT	3541
hsa-miR-548g	AAAAACUGAAUUACUUUUUGUAC	731	TTACAGTTT	1668	TACAGTTT	2605	ACAGTTT	3542
hsa-miR-548h	AAAAAGUAAUCGGGUUUUUUGUC	732	GATTACTTT	1669	ATTACTTT	2606	TTACTTT	3543
hsa-miR-548i	AAAAAGUAAUUGCGGAUUUUUGCC	733	AATTACTTT	1670	ATTACTTT	2607	TTACTTT	3544
hsa-miR-548j	AAAAAGUAAUUGCGGUUUUUUGGU	734	AATTACTTT	1671	ATTACTTT	2608	TTACTTT	3545
hsa-miR-548k	AAAAAGUACUUGCGGAUUUUUGCU	735	AAGTACTTT	1672	AGTACTTT	2609	GTACTTT	3546
hsa-miR-548l	AAAAAGUAAUUGCGGUUUUUUGUC	736	AAACTACTTT	1673	AATACTTT	2610	ATACTTT	3547
hsa-miR-548m	CAAGGUAAUUGUGUUUUUUG	737	AATACCTTT	1674	ATACCTTT	2611	TACCTTT	3548
hsa-miR-548n	CAAAAGUAAUUGUGGAUUUUUGU	738	ATTACTTTT	1675	TTACTTTT	2612	TACTTTT	3549
hsa-miR-548o	CCAAAACUGCAGUUUACUUUUUGC	739	GCAGTTTTG	1676	CAGTTTTG	2613	AGTTTTG	3550
hsa-miR-548p	UAGCAAAAACUCGAGUUUACUUU	740	GTTTTTGCT	1677	TTTTTGCT	2614	TTTTGCT	3551
hsa-miR-549	UGACAACUAUGGAUGAGCUCU	741	ATAGTTGTC	1678	TAGTTGTC	2615	AGTTGTC	3552
hsa-miR-550	AGUGCCUGAGGGAGUAAGAGCCC	742	CTCAGGCAC	1679	TCAGGCAC	2616	CAGGCAC	3553
hsa-miR-550*	UGUCUUACUCCCCUCAGGCACAU	743	GAGTAAGAC	1680	AGTAAGAC	2617	GTAAGAC	3554
hsa-miR-551a	GCGACCCACUUUGUUUCCA	744	AGTGGTCTG	1681	GTGGTCTG	2618	TGGTCTG	3555
hsa-miR-551b	GCGACCCAUACUUUGUUUUCAG	745	TATGGTCTG	1682	ATGGTCTG	2619	TGGTCTG	3556
hsa-miR-551b*	GAAUCAAGCGUGGGUGAGACC	746	GCTTGATTT	1683	CTTGATTT	2620	TTGATTT	3557
hsa-miR-552	AACAGGUGACUGGUUAGACAA	747	GTCACCTGT	1684	TCACCTGT	2621	CACCTGT	3558
hsa-miR-553	AAAACGGUGAGAUUUUGUUUU	748	TCACCGTTT	1685	CACCGTTT	2622	ACCGTTT	3559
hsa-miR-554	GCUAGUCCUGACUCAGCCAGU	4	CAGGACTAG	1686	AGGACTAG	2623	GGACTAG	3560

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-555	AGGUAAGCUGAACCCUCUGAU	750	AGCTTACCC	1687	GCTTACCC	2624	CTTACCC	3561
hsa-miR-556-3p	AUAUUACCAUUAGCUCUCAUCUUU	751	ATGGTAATA	1688	TGGTAATA	2625	GGTAATA	3562
hsa-miR-556-5p	GAUGAGCUCAUUGUAUAUAGAG	752	TGAGCTCAT	1689	GAGCTCAT	2626	AGCTCAT	3563
hsa-miR-557	GUUUGCACGGUGGGCCUUGUCU	753	CCGTGCAAA	1690	CGTGCAAA	2627	GTGCAAA	3564
hsa-miR-558	UGAGCUCUGUACCAAAU	754'	CAGCAGCTC	1691	AGCAGCTC	2628	GCAGCTC	3565
hsa-miR-559	UAAAGUAAUUAUGCACCACAAA	755	ATTTACTTT	1692	TTTACTTT	2629	TTACTTT	3566
hsa-miR-561	CAAAGUUUAAAGAUCCUUGAAGU	756	TAAACTTT	1693	TAAACTTT	2630	AAACTTT	3567
hsa-miR-562	AAAGUAGCUGUACCAUUUGC	757	CAGCTACTT	1694	AGCTACTT	2631	GCTACTT	3568
hsa-miR-563	AGGUUGACAUACGUUUUCCC	758	ATGTC AAC	1695	TGTC AAC	2632	GTC AAC	3569
hsa-miR-564	AGGCACGGUGUCAGCAGGC	759	CACCGTGCC	1696	ACCGTGCC	2633	CCGTGCC	3570
hsa-miR-566	GGCGCCUGUGAUCCCAAC	760	ACAGGCGCC	1697	CAGGCGCC	2634	AGGCGCC	3571
hsa-miR-567	AGUAUGUUUUCCAGGACAGAAC	761	AGAACATAC	1698	GAACATAC	2635	AACATAC	3572
hsa-miR-568	AUGUAUAAUUGUAUACACAC	762	ATTTATACA	1699	TTTATACA	2636	TTATACA	3573
hsa-miR-569	AGUUAUAGAAUCCUGGAAAGU	763	TTCATTAAC	1700	TCATTAAC	2637	CATTAAC	3574
hsa-miR-570	CGAAAACAGCAAUUACCUUUGC	764	GCTGTTTTTC	1701	CTGTTTTTC	2638	TGTTTTTC	3575
hsa-miR-571	UGAGUUGGCCAUUCUGAGUGAG	765	GGCCAACTC	1702	GCCAACTC	2639	CCA ACTC	3576
hsa-miR-572	GUCCGCUCCGGUGGCCCA	766	CCGAGCGGA	1703	CGAGCGGA	2640	GAGCGGA	3577
hsa-miR-573	CUGAAGUGAUGUGUAACUGAUCAG	767	ATCACTTCA	1704	TCACTTCA	2641	CACTTCA	3578
hsa-miR-574-3p	CACGCUCAUGCACACACCCACA	768	CATGAGCGT	1705	ATGAGCGT	2642	TGAGCGT	3579
hsa-miR-574-5p	UGAGUGUGUGUGUGAGUGUGU	769	CACACACTC	1706	ACACACTC	2643	CACACTC	3580
hsa-miR-575	GAGCCAGUUGGACAGGAGC	770	CAACTGGCT	1707	AACTGGCT	2644	ACTGGCT	3581
hsa-miR-576-3p	AAGAUGUGGAAAAUUUGGAAUC	771	TCCACATCT	1708	CCACATCT	2645	CACATCT	3582
hsa-miR-576-5p	AUUCUAAUUUUCUCCACCGUCUUU	772	AAATTAGAA	1709	AATTAGAA	2646	ATTAGAA	3583

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-577	UAGAUAAAAUUAUUGGUACCUG	773	ATTTTATCT	1710	TTTTTATCT	2647	TTTTATCT	3584
hsa-miR-578	CUUCUUGUGUCUCUAGGAUUGU	774	GCACAAGAA	1711	CACAAGAA	2648	ACAAGAA	3585
hsa-miR-579	UUCAUUUGGUUAAAACCGGAUU	775	ACCAATGA	1712	CCAAATGA	2649	CAAAATGA	3586
hsa-miR-580	UUGAGAUGAUGAAUCUUAUAGG	776	TCATTCTCA	1713	CATTCTCA	2650	ATTCTCA	3587
hsa-miR-581	UCUUGUUCUCUAGAUCAGU	777	GAACACAAG	1714	AACACAAG	2651	ACACAAG	3588
hsa-miR-582-3p	UAACUGGUUGAACAAACUGAACC	778	CAACCAGTT	1715	AACCAGTT	2652	ACCAGTT	3589
hsa-miR-582-5p	UUACAGUUGUUCAAACCAGUUAUCU	779	ACAACGTGA	1716	CAACTGTA	2653	AACTGTA	3590
hsa-miR-583	CAAAGAGGAAGGUCCCAUUAC	780	TTCCTCTTT	1717	TCCTCTTT	2654	CCTCTTT	3591
hsa-miR-584	UUUUGGUUUGCCUGGGACUGAG	781	CAAAACATA	1718	AAACCATA	2655	AACCATA	3592
hsa-miR-585	UGGGCUAUCUGUAUCUA	782	GATACGCC	1719	ATACGCC	2656	TACGCC	3593
hsa-miR-586	UAUGCAUUGUAUUUUUAGGUCC	783	ACAATGCAT	1720	CAATGCAT	2657	AATGCAT	3594
hsa-miR-587	UUUCCAUGGUGAUGAGUCAC	784	CCTATGGAA	1721	CTATGGAA	2658	TATGGAA	3595
hsa-miR-588	UUGGCCACAUGGGUUAAGAAC	785	TTGTGGCCA	1722	TGTGGCCA	2659	GTGGCCA	3596
hsa-miR-589	UGAGAACCACGUCUCUCUGAG	786	GTGGTTCTC	1723	TGGTTCTC	2660	GGTTCTC	3597
hsa-miR-589*	UCAGAACAUAUUGCCGGUUCGCCAGA	787	TTTGTCTG	1724	TTGTTCTG	2661	TGTTCTG	3598
hsa-miR-590-3p	UAAUUUAUGUAUAAGCUAGU	788	CATAAAATT	1725	ATAAAATT	2662	TAAAAATT	3599
hsa-miR-590-5p	GAGCUUAUUCAUAAAAAGUCAG	789	GAATAAGCT	1726	AATAAGCT	2663	ATAAGCT	3600
hsa-miR-591	AGACCAUGGUUCUCAUUGU	790	CCCATGGTC	1727	CCATGGTC	2664	CATGGTC	3601
hsa-miR-592	UUUGUCAUUUAUGCGAUGAUGU	791	ATTGACACA	1728	TTGACACA	2665	TGACACA	3602
hsa-miR-593	UGUCUCUGUGGGUUUCU	792	AGCAGAGAC	1729	GCAGAGAC	2666	CAGAGAC	3603
hsa-miR-593*	AGGCACCAGCCAGGCAUUGCUCAG C	793	GCTGGTGCC	1730	CTGGTGCC	2667	TGGTGCC	3604
hsa-miR-595	GAAGUGUGCCCGUGGUGUGUCU	794	GGCACACTT	1731	GCACACTT	2668	CACACTT	3605
hsa-miR-596	AAGCCUCCCCGGCUCCUCGGG	795	GGCAGGCT	1732	GGCAGGCT	2669	GCAGGCT	3606

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-597	UGUGUCACUCGAUGACCACUGU	796	GAGTGACAC	1733	AGTGACAC	2670	GTGACAC	3607
hsa-miR-598	UACGUCaucGUUGUCAUCGUGA	797	CGATGACGT	1734	GATGACGT	2671	ATGACGT	3608
hsa-miR-599	GUUGUGUCAGUUUAUCAAAAC	798	CTGACACAA	1735	TGACACAA	2672	GACACAA	3609
hsa-miR-600	ACUUACAGACAAGAGCCUUGCUC	799	GTCTGTAAG	1736	TCTGTAAG	2673	CTGTAAG	3610
hsa-miR-601	UGGUCUAGGAUUGUUGGAGGAG	800	TCCTAGACC	1737	CCTAGACC	2674	CTAGACC	3611
hsa-miR-602	GACACGGCGACAGCUGCGGCC	801	CGCCCGTGT	1738	GCCCGTGT	2675	CCCGTGT	3612
hsa-miR-603	CACACACUGCAAUUACUUUJGC	802	GCAGTGTGT	1739	CAGTGTGT	2676	AGTGTGT	3613
hsa-miR-604	AGGCUGCGGAAUUCAGGAC	803	TCCGCAGCC	1740	CCGCAGCC	2677	CGCAGCC	3614
hsa-miR-605	UAAAUCCCAUGGUCUUCUCCU	804	ATGGGATTT	1741	TGGGATTT	2678	GGGATTT	3615
hsa-miR-606	AAACUACUGAAAAUCAAAAGAU	805	TCAGTAGTT	1742	CAGTAGTT	2679	AGTAGTT	3616
hsa-miR-607	GUUCAAUCCAGAUUAUAAAC	806	GGATTTGAA	1743	GATTTGAA	2680	ATTTGAA	3617
hsa-miR-608	AGGGUGGUGUUGGGACAGCUCCU GU	807	CACCACCCC	1744	ACCACCCC	2681	CCACCCC	3618
hsa-miR-609	AGGGUUUUUCUCUCAUCUCU	808	GAACACCCC	1745	AAACACCCC	2682	AACACCCC	3619
hsa-miR-610	UGAGCUAAAUGUGUCUGGGA	809	ATTTAGCTC	1746	TTTAGCTC	2683	TTAGCTC	3620
hsa-miR-611	GCGAGGACCCUCCGGGUCUGAC	810	GGTCCCTCG	1747	GGTCCCTCG	2684	GTCCTCG	3621
hsa-miR-612	GCUGGCAGGGCUUCUGAGCUCCU U	811	CCTGCCCAG	1748	CTGCCCAG	2685	TGCCCAG	3622
hsa-miR-613	AGGAUUGUUCCUUUUUGCC	812	GAACATTCC	1749	AACATTCC	2686	ACATTCC	3623
hsa-miR-614	GAACGCCUUGUUCUUGCCAGGUGG	813	ACAGGCGTT	1750	CAGGCGTT	2687	AGGCGTT	3624
hsa-miR-615-3p	UCCGAGCCUGGGUCUCCUUCU	814	CAGGCTCGG	1751	AGGCTCGG	2688	GGCTCGG	3625
hsa-miR-615-5p	GGGGUCCCCGGUGUCUGGAUC	815	GGGACCCCC	1752	GGGACCCC	2689	GGACCCC	3626
hsa-miR-616	AGUCAUUGGAGGUUUGAGCAG	816	TCCAATGAC	1753	CCAATGAC	2690	CAATGAC	3627
hsa-miR-616*	ACUCAAACCCUUCAGUGACUU	817	GGTTTGGAG	1754	GTTTGGAG	2691	TTTTGGAG	3628

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-617	AGACUCCCAUUUGAAGGUGGC	818	TGGGAAGTC	1755	GGGAAGTC	2692	GGAAGTC	3629
hsa-miR-618	AAACUCUACUUGUCUUCUGAGU	819	AGTAGAGTT	1756	GTAGAGTT	2693	TAGAGTT	3630
hsa-miR-619	GACCUAGACAUUUUGUCCCCAGU	820	TGTCCAGGT	1757	GTCAGGT	2694	TCCAGGT	3631
hsa-miR-620	AUGGAGUAUAUAGAAAU	821	CTATCTCCA	1758	TATCTCCA	2695	ATCTCCA	3632
hsa-miR-621	GGCUAGCAACAGCGCUUACCU	822	GTTGCTAGC	1759	TTGCTAGC	2696	TGCTAGC	3633
hsa-miR-622	ACAGUCUGCUGAGGUUGGAGC	823	AGCAGACTG	1760	GCAGACTG	2697	CAGACTG	3634
hsa-miR-623	AUCCCUUGCAGGGGCUUUGGGU	824	TGCAAGGGA	1761	GCAAGGGA	2698	CAAGGGA	3635
hsa-miR-624	CACAAGGUUUUGUAUUACCU	825	ATACCTTGT	1762	TACCTTGT	2699	ACCTTGT	3636
hsa-miR-624*	UAGUACCAGUACCUUUGUUA	826	ACTGGTACT	1763	CTGGTACT	2700	TGGTACT	3637
hsa-miR-625	AGGGGAAAGUUUAUAGUCC	827	CTTCCCCC	1764	TTTCCCCC	2701	TTCCCCC	3638
hsa-miR-625*	GACUUAAGAACUUUCCCCUCA	828	TTCTATAGT	1765	TCTATAGT	2702	CTATAGT	3639
hsa-miR-626	AGCUGUCUGAAAAUGUCUU	829	TCAGACAGC	1766	CAGACAGC	2703	AGACAGC	3640
hsa-miR-627	GUGAGUCUCUAAGAAAAGAGGA	830	AGAGACTCA	1767	GAGACTCA	2704	AGACTCA	3641
hsa-miR-628-3p	UCUAGUAAGAGUGGCAGUCGA	831	TCTTACTAG	1768	CTTACTAG	2705	TTACTAG	3642
hsa-miR-628-5p	AUGCUGACAUUUUAUCUAGAGG	832	ATGTCAGCA	1769	TGTCAGCA	2706	GTCAGCA	3643
hsa-miR-629	UGGGUUUACGUUGGGAGAAACU	833	CGTAAACCC	1770	GTAACCCC	2707	TAAACCC	3644
hsa-miR-629*	GUUCUCCCAACGUAAAGCCAGC	834	TTGGGAGAA	1771	TGGGAGAA	2708	GGGAGAA	3645
hsa-miR-630	AGUUAUCUGUACCAGGGAAGGU	835	ACAGAATAC	1772	CAGAATAC	2709	AGAATAC	3646
hsa-miR-631	AGACCUGGCCCCAGACCUCAGC	836	GGCCAGGTC	1773	GCCAGGTC	2710	CCAGGTC	3647
hsa-miR-632	GUGUCUGCUUCCUGUGGGA	837	AAGCAGACA	1774	AGCAGACA	2711	GCAGACA	3648
hsa-miR-633	CUAAUAGUAUCUACCACAAUAAA	838	ATACTATTA	1775	TACTATTA	2712	ACTATTA	3649
hsa-miR-634	AACCAGCACCCCAACUUUGGAC	839	GGTGCTGGT	1776	GTGCTGGT	2713	TGCTGGT	3650
hsa-miR-635	ACUUGGGCACUGAAAACAAUUGUCC	840	GTGCCCAAG	1777	TGCCCAAG	2714	GCCCAAG	3651

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-636	UGUGCUUUCUCUCGUCUCCGCCCGCA	841	AGCAAGCAC	1778	GCAAGCAC	2715	CAAGCAC	3652
hsa-miR-637	ACUGGGGGCUUUCGGGCUUCUGCG	842	AGCCCCCAG	1779	GCCCCCAG	2716	CCCCCAG	3653
hsa-miR-638	AGGAUCGCCGGGGGUGGGCGGC	843	CGGATCCC	1780	GCGATCCC	2717	CGATCCC	3654
hsa-miR-639	AUCGCUCCGGUUCGAGCGCUGU	844	CCGACGGA	1781	CGCAGCGA	2718	GCAGCGA	3655
hsa-miR-640	AUGAUCCAGGAACCCUCCUCU	845	CCTGGATCA	1782	CTGGATCA	2719	TGGATCA	3656
hsa-miR-641	AAAGACAUAGGAUAGAGUACCCUC	846	CTATGTCTT	1783	TATGTCTT	2720	ATGTCTT	3657
hsa-miR-642	GUCCUCUCCAAAUGUGUCUUG	847	GGAGAGGGA	1784	GAGAGGGA	2721	AGAGGGA	3658
hsa-miR-643	ACUUGUAUCUAGCUCAGGUAG	848	GCATACAAG	1785	CATACAAG	2722	ATACAAG	3659
hsa-miR-644	AGUGUGCUUUUCUAGAGC	849	AAGCCACAC	1786	AGCCACAC	2723	GCCACAC	3660
hsa-miR-645	UCUAGGCUUGUACUCUGA	850	CCAGCCTAG	1787	CAGCCTAG	2724	AGCCTAG	3661
hsa-miR-646	AAGCAGCUCUCCUCUGAGGC	851	GCAGCTGCT	1788	CAGCTGCT	2725	AGCTGCT	3662
hsa-miR-647	GUGGCUCCAGCUCUCCUUC	852	GTGCAGCCA	1789	TGCAGCCA	2726	GCAGCCA	3663
hsa-miR-648	AAGUGCAGGGCACUCUGU	853	CTGCACACT	1790	TGCACACT	2727	GCACACT	3664
hsa-miR-649	AAACCUUGUUGUUCUAAAGAGUC	854	ACACAGGTT	1791	CACAGGTT	2728	ACAGGTT	3665
hsa-miR-650	AGGAGGCAGCCUCUCAGGAC	855	GCTGCCCTCC	1792	CTGCCCTCC	2729	TGCCCTCC	3666
hsa-miR-651	UUUAGGAUAAGCUUAGACUUUUG	856	TTATCCTAA	1793	TATCCTAA	2730	ATCCTAA	3667
hsa-miR-652	AAUGGCACACUAGGUUGUG	857	TGGCGCCAT	1794	GGCGCCAT	2731	GCGCCAT	3668
hsa-miR-653	GUGUUGAAACAUCUCUACUG	858	GTTTCAACA	1795	TTTCAACA	2732	TTCAACA	3669
hsa-miR-654-3p	UAUGUCUGCUGACCACUACCCUU	859	AGCAGACAT	1796	GCAGACAT	2733	CAGACAT	3670
hsa-miR-654-5p	UGGUGGCCCGCAGAACUAGUGC	860	CGGCCACC	1797	GGCCCACC	2734	GCCCACC	3671
hsa-miR-655	AUAUAUAUGGUUAACCCUUCUU	861	CATGTATTA	1798	ATGTATTA	2735	TGTATTA	3672
hsa-miR-656	AAUAUAUAACAGUCAACCCUCU	862	GTATAATAT	1799	TATAATAT	2736	ATAATAT	3673

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

**Table 1**

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-657	GGCAGGUUCUCACCCUCUCUAGG	863	AGAACCTGC	1800	GAACCTGC	2737	AACCTGC	3674
hsa-miR-658	GGCGGAGGGAAGUAGGUCCGUUG GU	864	TCCCTCCGC	1801	CCCTCCGC	2738	CCTCCGC	3675
hsa-miR-659	CUUGGUUCAGGGAGGUCCCA	865	CTGAACCAA	1802	TGAACCAA	2739	GAACCAA	3676
hsa-miR-660	UACCCAUUGCAUUCGGAGUUG	866	GCAATGGGT	1803	CAATGGGT	2740	AATGGGT	3677
hsa-miR-661	UGCCUGGGUCUCUGGCCUGGGCG U	867	GACCCAGGC	1804	ACCCAGGC	2741	CCCAGGC	3678
hsa-miR-662	UCCACGUUGUGGCCCCAGCAG	868	CAACGTGGG	1805	AACGTGGG	2742	ACGTGGG	3679
hsa-miR-663	AGCGGGGGCCCGGGACCCG	869	CGCCCCGCC	1806	GCCCCGCC	2743	CCCCGCC	3680
hsa-miR-663b	GGUGGGCCCGCCGUGCCUGAGG	870	CCGGGCCAC	1807	CGGGCCAC	2744	GGGCCAC	3681
hsa-miR-664	UAUUAUUUAUCCCCAGCCUACA	871	TAAATGAAT	1808	AAATGAAT	2745	AATGAAT	3682
hsa-miR-664*	ACUGGCUAGGGAAAUAUUGGAU	872	CCTAGCCAG	1809	CTAGCCAG	2746	TAGCCAG	3683
hsa-miR-665	ACCAGGAGCUGAGGCCCCU	873	GCCTCCTGG	1810	CCTCCTGG	2747	CTCCTGG	3684
hsa-miR-668	UGUCACUCGGCUCGGCCACUAC	874	CCGAGTGAC	1811	CGAGTGAC	2748	GAGTGAC	3685
hsa-miR-671-3p	UCCGGUUCUCAGGGCUCACC	875	GAGAACC GG	1812	AGAACC GG	2749	GAACC GG	3686
hsa-miR-671-5p	AGGAAGCCCUUGAGGGGCUGGAG	876	AGGGCTTCC	1813	GGGCTTCC	2750	GGCTTCC	3687
hsa-miR-675	UGGUGGGAGAGGGCCACAGUG	877	CTCCGCACC	1814	TCCGCACC	2751	CCGCACC	3688
hsa-miR-675b	CUGUAUGCCUCACCCGCUCA	878	GGGCATACA	1815	GGCATACA	2752	GCATACA	3689
hsa-miR-7	UGGAAGACUAGUAUUUUGUUGU	879	TAGTCTTCC	1816	AGTCTTCC	2753	GTCTTCC	3690
hsa-miR-7-1*	CAACAAUACACAGUCGCCAU	880	TGATTTGTT	1817	GATTTGTT	2754	ATTTGTT	3691
hsa-miR-7-2*	CAACAAUCCCAGUCUACCUAA	881	GGATTTGTT	1818	GATTTGTT	2755	ATTTGTT	3692
hsa-miR-708	AAGGAGCUUACAAUCUAGCUGGG	882	TAAGCTCCT	1819	AAGCTCCT	2756	AGCTCCT	3693
hsa-miR-708*	CAACUAGACUGAGCUUCUAG	883	AGTCTAGTT	1820	GTCTAGTT	2757	TCTAGTT	3694
hsa-miR-720	UCUCGCGGGGCCUCCA	884	CCCAGCGAG	1821	CCAGCGAG	2758	CAGCGAG	3695



5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-744	UGCGGGGCUAGGGCUAACAGCA	885	TAGCCCCGC	1822	AGCCCCGC	2759	GCCCCGC	3696
hsa-miR-744*	CUGUUGCCACUAACCCUCAACCU	886	GTGGCAACA	1823	TGGCAACA	2760	GGCAACA	3697
hsa-miR-758	UUUGAGCCUGGUCACUAACC	887	AGTCAACA	1824	GGTCAACA	2761	GTCACAA	3698
hsa-miR-760	CGGCUUGGUCUGGGGA	888	CCCAGGCC	1825	CCAGAGCC	2762	CAGAGCC	3699
hsa-miR-765	UGGAGGAGGAAGGUGAUG	889	TTCTCTCC	1826	TCTCTCC	2763	CTCCTCC	3700
hsa-miR-766	ACUCCAGCCCCACAGCCUCAGC	890	GGGCTGGAG	1827	GGCTGGAG	2764	GCTGGAG	3701
hsa-miR-767-3p	UCUGCUAUACCCCAUGGUUCU	891	TATGAGCAG	1828	ATGAGCAG	2765	TGAGCAG	3702
hsa-miR-767-5p	UGCACCAUGGUUGUCUGAGCAUG	892	CCATGGTGC	1829	CATGGTGC	2766	ATGGTGC	3703
hsa-miR-769-3p	CUGGGAUCUCCGGGUCUUGGUU	893	GAGATCCCA	1830	AGATCCCA	2767	GATCCCA	3704
hsa-miR-769-5p	UGAGACCUCUGGGUUCUGAGCU	894	AGAGTCTC	1831	GAGTCTC	2768	AGGTCTC	3705
hsa-miR-770-5p	UCCAGUACCACGUCAGGGCCA	895	TGGTACTGG	1832	GGTACTGG	2769	GTACTGG	3706
hsa-miR-802	CAGUAACAAAGAUUAUCCUUGU	896	TTTGTACT	1833	TTGTACT	2770	TGTTACT	3707
hsa-miR-873	GCAGGAACUUGUGAGUCUCU	897	AAGTTCCTG	1834	AGTTCCTG	2771	GTTTCCTG	3708
hsa-miR-874	CUGCCUUGCCCGAGGGACCGA	898	GCCAGGGCA	1835	CCAGGGCA	2772	CAGGGCA	3709
hsa-miR-875-3p	CCUGGAAACACUGAGGUUGUG	899	TGTTTCCAG	1836	GTTTCCAG	2773	TTTTCCAG	3710
hsa-miR-875-5p	UAUACCUACAGUUUUUAUCAGGUG	900	CTGAGGTAT	1837	TGAGGTAT	2774	GAGGTAT	3711
hsa-miR-876-3p	UGGUGUUUACAAAGUAUUUCA	901	TAAACCACC	1838	AAACCACC	2775	AACCACC	3712
hsa-miR-876-5p	UGGAUUUUUGUGAAUCACCA	902	AAGAAATCC	1839	AGAAATCC	2776	GAAATCC	3713
hsa-miR-877	GUAGAGGAGAUUGCGCAGGG	903	TCTCCTCTA	1840	CTCCTCTA	2777	TCCTCTA	3714
hsa-miR-877*	UCCUCUUCUCCUCCUCCUCCAG	904	GAGAAGAGG	1841	AGAAGAGG	2778	GAAGAGG	3715
hsa-miR-885-3p	AGGCAGCGGGUGUAGUGGAUA	905	CCCGCTGCC	1842	CCGCTGCC	2779	CGCTGCC	3716
hsa-miR-885-5p	UCCAUUACACUACCCUCCUCUCU	906	GTGTAATGG	1843	TGTAATGG	2780	GTAATGG	3717
hsa-miR-886-3p	CGCGGGUUCUUUACUGACCCUU	907	AGCACCCGC	1844	GCACCCGC	2781	CACCCGC	3718

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-886-5p	CGGGUCGGAGUAGUCUCAAGCGG	908	CTCCGACCC	1845	TCCGACCC	2782	CCGACCC	3719
hsa-miR-887	GUGAACGGCGCCCAUCCCGAGG	909	GCCCCTTCA	1846	CCCGTTCA	2783	CCGTTCA	3720
hsa-miR-888	UACUCAAAAAGCUGUCAGUCA	910	TTTTTGAGT	1847	TTTTGAGT	2784	TTTGAGT	3721
hsa-miR-888*	GACUGACACCUCUUUGGGUGAA	911	GGTGTCAGT	1848	GTGTCAGT	2785	TGTCAGT	3722
hsa-miR-889	UUAAUUCGGACAACCAUUGU	912	CCGATATTA	1849	CGATATTA	2786	GATATTA	3723
hsa-miR-890	UACUUGGAAAGGCAUCAGUUG	913	TTTCCAAGT	1850	TTCCAAGT	2787	TCCAAGT	3724
hsa-miR-891a	UGCAACGAACCUGAGCCACUGA	914	GTTCTGTC	1851	TTCGTTGC	2788	TCGTTGC	3725
hsa-miR-891b	UGCAACUUAACCUGAGUCAUUGA	915	GTAAGTTGC	1852	TAAGTTGC	2789	AAGTTGC	3726
hsa-miR-892a	CACUGUGUCCUUUCUGCGUAG	916	GGACACAGT	1853	GACACAGT	2790	ACACAGT	3727
hsa-miR-892b	CACUGGCCUUUCUGGGUAGA	917	GGAGCCAGT	1854	GAGCCAGT	2791	AGCCAGT	3728
hsa-miR-9	UCUUUGGUUAUCUAGCUGUAUGA	918	TAACCAAAG	1855	AACCAAAG	2792	ACCAAAG	3729
hsa-miR-9*	AUAAAAGCUAGUAACCGAAAGU	919	CTAGCTTTA	1856	TAGCTTTA	2793	AGCTTTA	3730
hsa-miR-920	GGGAGCUGUGGAAGCAGUA	920	ACAGTCCC	1857	CAGTCCC	2794	AGTCCC	3731
hsa-miR-921	CUAGUGAGGGACACAGAACCAGGAUU C	921	CCCTCACTA	1858	CCTCACTA	2795	CTCACTA	3732
hsa-miR-922	GCAGCAGAGAAUAGGACUACGUC	922	TCTCTGCTG	1859	CTCTGCTG	2796	TCTGCTG	3733
hsa-miR-923	GUCAGCGGAGGAAAAGAAACU	923	CTCCGCTGA	1860	TCCGCTGA	2797	CCGCTGA	3734
hsa-miR-924	AGAGUCUUGUGAUGUCUUGC	924	ACAAGACTC	1861	CAAGACTC	2798	AAGACTC	3735
hsa-miR-92a	UAUUGCACUUGUCCCGCCUGU	925	AAGTGCAAT	1862	AGTGCAAT	2799	GTGCAAT	3736
hsa-miR-92a-1*	AGGUUGGGAUCGGUUGCAUUGCU	926	ATCCCAACC	1863	TCCCAACC	2800	CCCAACC	3737
hsa-miR-92a-2*	GGGUGGGAAUUGUUGCAUJAC	927	ATCCCCACC	1864	TCCCCACC	2801	CCCCACC	3738
hsa-miR-92b	UAUUGCACUCUCCCGCCUCC	928	GAGTGCAAT	1865	AGTGCAAT	2802	GTGCAAT	3739
hsa-miR-92b*	AGGACGGGACGGGUGCAGUG	929	TCCCGTCCC	1866	CCCGTCCC	2803	CCGTCCC	3740
hsa-miR-93	CAAAGUCGUGUUCGUGCAGGUAG	930	CAGCACTTT	1867	AGCACTTT	2804	GCACTTT	3741

(continued)

microRNA	MicroRNASequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsa-miR-93*	ACUGCUGAGCUAGCACUUCUCCG	931	GCTCAGCAG	1868	CTCAGCAG	2805	TCAGCAG	3742
hsa-miR-933	UGUGCGCAGGGAGACCUUCUCCC	932	CCTGCGCAC	1869	CTGCGCAC	2806	TGCGCAC	3743
hsa-miR-934	UGUCUACUACUGGAGACACUGG	933	GTAGTAGAC	1870	TAGTAGAC	2807	AGTAGAC	3744
hsa-miR-935	CCAGUUACCGCUUCGCGUACCGC	934	CGGTAAC TG	1871	GGTAACTG	2808	GTAAC TG	3745
hsa-miR-936	ACAGUAGAGGGAGGAUUCGCAG	935	CCTCTACTG	1872	CTCTACTG	2809	TCTACTG	3746
hsa-miR-937	AUCCGGCUCUGACUCUCUGCC	936	GAGCGCGGA	1873	AGCGCGGA	2810	GCGCGGA	3747
hsa-miR-938	UGCCCUUAAAAGGUGAACCCAGU	937	TTTAAGGGC	1874	TTAAGGGC	2811	TAAGGGC	3748
hsa-miR-939	UGGGGAGCUGAGGCUCUGGGGGU G	938	CAGCTCCCC	1875	AGCTCCCC	2812	GCTCCCC	3749
hsa-miR-940	AAGGCAGGGCCCCCGCUCUCCC	939	GCCCTGCCT	1876	CCCTGCCT	2813	CCTGCCT	3750
hsa-miR-941	CACCCGGCUGUGGCACAUUGUG	940	CAGCCGGGT	1877	AGCCGGGT	2814	GCCGGGT	3751
hsa-miR-942	UCUUCUCUGUUUUGGCCAUUGUG	941	ACAGAGAAG	1878	CAGAGAAG	2815	AGAGAAG	3752
hsa-miR-943	CUGACUGUUGCCGUCCUCCAG	942	CAACAGTCA	1879	AACAGTCA	2816	ACAGTCA	3753
hsa-miR-944	AAAUUUUUGUACAUCGGAUGAG	943	ACAATAATT	1880	CAATAATT	2817	AATAATT	3754
hsa-miR-95	UUCAACGGGUUUUUUUUGAGCA	944	ACCCGTTGA	1881	CCCGTTGA	2818	CCGTTGA	3755
hsa-miR-96	UUUGGCACUAGCACAUUUUUUGCU	945	TAGTGCCAA	1882	AGTGCCAA	2819	GTGCCAA	3756
hsa-miR-96*	AAUCAUGUGCAGUGCCAAUAUG	946	GCACATGAT	1883	CACATGAT	2820	ACATGAT	3757
hsa-miR-98	UGAGGUAGUAAGUUGUAUUUGUU	947	TACTACCTC	1884	ACTACCTC	2821	CTACCTC	3758
hsa-miR-99a	AACCCGUAGAUCCGAUCUUGUG	948	TCTACGGGT	1885	CTACGGGT	2822	TACGGGT	3759
hsa-miR-99a*	CAAGCUCGCUUCUAUGGGUCUG	949	AGCGAGCTT	1886	GCGAGCTT	2823	CGAGCTT	3760
hsa-miR-99b	CACCCGUAGAACCAGCCUUGCG	950	TCTACGGGT	1887	CTACGGGT	2824	TACGGGT	3761
hsa-miR-99b*	CAAGCUCGUGUCUGUGGGUCCG	951	CACGAGCTT	1888	ACGAGCTT	2825	CGAGCTT	3762
hsv1-miR-H1	UGGAAGGACGGGAAGUGGAAG	952	CGTCCCTCC	1889	GTCCTTCC	2826	TCCTTCC	3763
hsv1-miR-H2-3p	CCUGAGCCAGGGACGAGUGCGGACU	953	CTGGCTCAG	1890	TGGCTCAG	2827	GGCTCAG	3764

(continued)

microRNA	MicroRNA Sequence	SEQ ID NO	9-mer	SEQ ID NO	8-mer	SEQ ID NO	7-mer	SEQ ID NO
hsv1-miR-H2-5p	UCGCACGGCGCCCGGCACAGACU	954	GCGCGTGCGG	1891	CGCGTGCGG	2828	GCGTGCGG	3765
hsv1-miR-H3	CUGGGACUGUGCGGUUGGGA	955	ACAGTCCCA	1892	CAGTCCCA	2829	AGTCCCA	3766
hsv1-miR-H4-3p	CUUGCCUGUCUAAACUCGCUAGU	956	GACAGGCAA	1893	ACAGGCAA	2830	CAGGCAA	3767
hsv1-miR-H4-5p	GGUAGAGUUUGACAGGCAAGCA	957	AAACTCTAC	1894	AACTCTAC	2831	ACTCTAC	3768
hsv1-miR-H5	GUCAGAGAUCCAAACCCUCCGG	958	GATCTCTGA	1895	ATCTCTGA	2832	TCTCTGA	3769
hsv1-miR-H6	CACUCCCCGUCCUCCAUCCC	959	ACGGGAAGT	1896	CGGGAAGT	2833	GGGAAGT	3770
kshv-miR-K12-1	AUUACAGGAAACUGGGUGUAAGC	960	TTCCCTGTAA	1897	TCCTGTAA	2834	CCTGTAA	3771
kshv-miR-K12-10a	UAGUGUUUUCUCCCCCGAGUGGC	961	GACAACACT	1898	ACAACACT	2835	CAACACT	3772
kshv-miR-K12-10b	UGGUGUUUUCUCCCCCGAGUGGC	962	GACAACACC	1899	ACAACACC	2836	CAACACC	3773
kshv-miR-K12-11	UUAAUGCUUAGCCUGUGUCCGA	963	TAAGCATTAA	1900	AAGCATTAA	2837	AGCATTAA	3774
kshv-miR-K12-12	ACCAGGCCACCAUUCUCCUCCG	964	GTGGCCTGG	1901	TGGCCTGG	2838	GGCCTGG	3775
kshv-miR-K12-2	AACUGUAGUCCGGGUCGAUCUG	965	GACTACAGT	1902	ACTACAGT	2839	CTACAGT	3776
kshv-miR-K12-3	UCACAUUCUGAGGACGGCAGCCGA	966	CAGAAATGTG	1903	AGAATGTG	2840	GAATGTG	3777
kshv-miR-K12-3*	UCGCGGUACACAGAAUUGACA	967	GTGACCCGG	1904	TGACCCGG	2841	GACCCGG	3778
kshv-miR-K12-4-3p	UAGAAUACUGAGGCCUAGCUGA	968	CAGTATTCT	1905	AGTATTCT	2842	GTATTCT	3779
kshv-miR-K12-4-5p	AGCUAAACCCGACGACUCUAGG	969	CGGTTTAGC	1906	GGTTTAGC	2843	GTTTAGC	3780
kshv-miR-K12-5	UAGGAUGCCUGGAACUUGCCGG	970	AGGCATCCT	1907	GGCATCCT	2844	GCATCCT	3781
kshv-miR-K12-6-3p	UGAUGGUUUUCGGCUGUUGAG	971	AAAACCATC	1908	AAACCATC	2845	AACCATC	3782
kshv-miR-K12-6-5p	CCAGCAGCACCUAAUCCAUCCGG	972	GTGCTGCTG	1909	TGCTGCTG	2846	GCTGCTG	3783
kshv-miR-K12-7	UGAUCCCAUGUUGCUGGGCGCU	973	CATGGGATC	1910	ATGGGATC	2847	TGGGATC	3784
kshv-miR-K12-8	UAGGCGGACUCGAGAGAGCAGC	974	GTCGCGCCT	1911	TCGCGCCT	2848	CGCGCCT	3785
kshv-miR-K12-9	CUGGGUUAACGCAGCUCGCUAA	975	GTATACCCA	1912	TATACCCA	2849	ATACCCA	3786
kshv-miR-K12-9*	ACCCAGCUCGCGUAAACCCCGCU	976	GCAGCTGGG	1913	CAGCTGGG	2850	AGCTGGG	3787

**[0398]** Figure 32. Silencing of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation in soft agar in PC3 cells. PC3 cells were transfected with 25 nM of the 8-mer LNA-antimiR #3205 targeting miR-21. After 24 hours, cells were harvested and seeded on soft agar. After 12 days, colonies were counted. Shown is the mean of three separate experiments, each performed in triplicate, and normalised against 0 nM control (i.e. transfection but with no LNA).  
 5 p=0.01898 for #3205.

**[0399]** Figure 33. Knock-down of miR-21 by the 8-mer #3205 LNA-antimiR reduces colony formation in soft agar in HepG2 cells. HepG2 cells were transfected with 25 nM of the 8-mer LNA-antimiR #3205 targeting miR-21. After 24 hours, cells were harvested and seeded on soft agar. After 17 days, colonies were counted. Shown is the mean of three replicates from one experiment (bars=SEM).

**[0400]** **Figure 34.** Wound closure in the invasive human prostate cell line PC3 after treatment with #3205. (A) PC3 cells were transfected at day 3 with LNA-antimiR and control oligonucleotides at 25 nM, #3205 (8mer, perfect match) and #3219 (8mer, mismatch) and the following day a scratch was made. Pictures were taken after 24 hours in order to monitor the migration. (B) The area in each timepoint has been measured with the software program Image J and normalized against respective 0 h time-point.

**[0401]** **Figure 35.** Length assessment of fully LNA-substituted LNA-antimiRs antagonizing miR-155. RAW cells were co-transfected with luciferase reporter plasmids containing a perfect match target site for miR-155 and with LNA-antimiR oligonucleotides at different concentrations. After 24 hours, cells were harvested and luciferase activity measured. Shown are the mean values (bars=s.e.m) for three independent experiments where the renilla/firefly ratios have been normalized against 0 nM empty vector without target site (=mock). Shown is also a schematic presentation of the miR sequence and the design and position of the LNA-antimiRs.

**[0402]** **Figure 36.** Binding of 5'-FAM labeled LNA-antimiR-21 (#3205) to mouse plasma protein. (A) % unbound LNA-antimiR-21 compound as a function of oligonucleotide concentration in mouse plasma. (B) Concentration of unbound LNA-antimiR-21 compound #3205 as a function of #3205 concentration in mouse plasma.

**[0403]** **Figure 37.** Quantification Ras protein levels by Western blot analysis.

A. Gel image showing Ras and Tubulin (internal standard) protein in treated (anti-let-7; 8-mer) vs. untreated (saline) lung and kidney samples. B. Quantifications of Ras protein levels in the lung and kidney, respectively, of LNA-antimiR-treated mice (black bars), normalized against equivalent saline controls (grey bars), using tubulin as equal-loading control.

B. Silencing of miR-21 by #3205 leads to increased levels of Pdc4 protein levels *in vivo*.

**[0404]** **Conclusion:** The 8-mer LNA-antimiR shows a clear effect in regulating target let-7 miRNA *in vivo*, as evaluated based on Ras protein levels in treated vs. control mice. Whereas the effect seems to be smaller in lungs, Ras levels in the kidney show a substantial up-regulation upon antimiRs-treatment.

**[0405]** **Materials and Methods:** *Animals and dosing:* C57BL/6 female mice were treated with 10 mg/kg LNA-antimiR or saline for three consecutive days (0, 1, and 2) and sacrificed on day 4. Tissue samples from lungs and kidneys were snapfrozen and stored at -80°C until further processing.

**[0406]** **Western blot analysis:** Lung and kidney proteins from saline and LNA-antimiR-treated mice were separated on NuPAGE Bis Tris 4-12% (Invitrogen), using 100 µg per sample. The proteins were transferred to a nitrocellulose membrane using iBlot (Invitrogen) according to the manufacturer's instructions. Blocking, antibody dilution and detection was performed according to the manufacturer's specifications. For Ras detection, a primary rabbit-anti Ras antibody (SC-3339, Santa Cruz Biotechnology) and a secondary HRP-conjugated swine-anti-rabbit antibody (P0399, Dako) was used, and for tubulin detection, a primary tubulin alpha (MS-581-P1, Neomarkers) and a secondary HRP-conjugated goat-anti-mouse antibody (P0447, Dako) was used.

**Example 40. *In vivo* efficacy assessment of the 8-mer LNA-antimiR (#3205) in targeting miR-21, as determined by Pdc4 protein up-regulation in mouse kidney.**

**[0407]** We have shown that an 8-mer LNA-antimiR that is fully LNA-modified antagonizes miR-21 and has the ability to regulate the protein levels of the miR-21 target Pdc4 *in vitro*. We therefore injected the LNA-antimiR into mice to determine the effects of the LNA-antimiR *in vivo*. The mice received 25 mg/kg of #3205 by i.v. injection every other day for 14 days (a total of 5 doses). The mice were sacrificed on day 14, the kidney was removed, and protein was isolated. In order to determine target regulation, Western blot analysis was performed.

**[0408]** **Results:** As shown in Figure 37, treating mice with #3205 showed significantly increased Pdc4 protein levels as compared to the saline control. While the normalized Pdc4 versus Gapdh ratio was consistent in both saline samples, the protein up-regulation in the two LNA-antimiR-treated (#32059 mice were measured to 3.3- and 6.3-fold, respectively, demonstrating an *in vivo* pharmacological effect of the #3205 8-mer LNA-antimiR.

**[0409]** **Conclusion:** The fully LNA-modified 8-mer LNA-antimiR #3205 antagonizes miR-21 *in vivo*, as demonstrated

through its ability to de-repress (up-regulate) mouse kidney levels of Pdc4, a validated miR-21 target.

#### Materials and Methods:

5 **[0410]** *Animals and dosing:* C57BL/6 female mice with average of 20 g body weight at first dosing were used in all experiments and received regular chow diet (Altromin no 1324, Brogaarden, Gentofte, Denmark). Substances were formulated in physiological saline (0.9% NaCl). The

#### SEQUENCE LISTING

10

#### **[0411]**

<110> Santaris Pharma A/S

15

<120> Micromirs

<130> 1051WO

20

<150> US 60/977497

<151> 2007-10-04

<150> US 60/979217

<151> 2007-10-11

25

<150> US 61/028062

<151> 2008-02-12

<150> EP08104780.5

<151> 2008-07-17

30

<160> 980

<170> PatentIn version 3.5

35

<210> 1

<211> 15

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Seedmer, optionally fully LNA phosphorothioate

<220>

<221> misc\_feature

45

<222> (1)..(15)

<223> Phosphorothioate linkages, LNAs at positions 1, 3, 4, 6, 7, 8, 10, 12, 14 & 15 otherwise DNAs.

<400> 1

tcagtctgat aagct 15

50

<210> 2

<211> 8

<212> DNA

<213> Artificial Sequence

55

<220>

<223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 5  
 <400> 2  
 gataagct 8  
 <210> 3  
 10 <211> 15  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 15 <223> LNA Oligomer  
 <220>  
 <221> misc\_feature  
 <222> (1)..(15)  
 20 <223> Phosphorothioate linkages, LNAs at positions 1, 3, 5, 6, 7, 10, 11, 12, 14 & 15 otherwise DNAnts.  
 <400> 3  
 tcacaattag catta 15  
 25 <210> 4  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence  
 30 <220>  
 <223> LNA Oligomer  
 <220>  
 <221> misc\_feature  
 35 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 <400> 4  
 40 tagcatta 8  
 <210> 5  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence  
 45 <220>  
 <223> LNA Oligomer  
 <220>  
 <221> misc\_feature  
 50 <222> (1)..(15)  
 <223> Phosphorothioate linkages, LNAs at positions 1, 3, 6, 7, 8, 10, 12, 14 & 15 otherwise DNAnts.  
 <400> 5  
 55 ccattgtcac actcc 15  
 <210> 6  
 <211> 8

<212> DNA  
 <213> Artificial Sequence

5

<220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages

10

<400> 6  
 cacactcc 8

15

<210> 7  
 <211> 6  
 <212> DNA  
 <213> Artificial Sequence

20

<220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(6)  
 <223> Fully LNA & Phosphorothioate linkages

25

<400> 7  
 taagct 6

30

<210> 8  
 <211> 7  
 <212> DNA  
 <213> Artificial Sequence

35

<220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(7)  
 <223> Fully LNA & Phosphorothioate linkages

40

<400> 8  
 ataagct 7

45

<210> 9  
 <211> 9  
 <212> DNA  
 <213> Artificial Sequence

50

<220>  
 <223> LNA Oligomer

55

<220>  
 <221> misc\_feature  
 <222> (1)..(9)  
 <223> Fully LNA & Phosphorothioate linkages



<400> 9  
 tgataagct 9

5  
 <210> 10  
 <211> 10  
 <212> DNA  
 <213> Artificial Sequence

10  
 <220>  
 <223> LNA Oligomer

15  
 <220>  
 <221> misc\_feature  
 <222> (1) .. (10)  
 <223> Fully LNA & Phosphorothioate linkages

20  
 <400> 10  
 ctgataagct 10

25  
 <210> 11  
 <211> 12  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> LNA Oligomer

35  
 <220>  
 <221> misc\_feature  
 <222> (1)..(12)  
 <223> Fully LNA & Phosphorothioate linkages

40  
 <400> 11  
 gtctgataag ct 12

45  
 <210> 12  
 <211> 14  
 <212> DNA  
 <213> Artificial Sequence

50  
 <220>  
 <223> LNA Oligomer

55  
 <220>  
 <221> misc\_feature  
 <222> (1)..(14)  
 <223> Fully LNA & Phosphorothioate linkages

<400> 12  
 cagtctgata agct 14

<210> 13  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 5  
 <400> 13  
 tctgataa 8  
 <210> 14  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence  
 10  
 <220>  
 <223> LNA Oligomer  
 15  
 <220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 20  
 <400> 14  
 atcagtct 8  
 25  
 <210> 15  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> LNA Oligomer  
 <220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 35  
 <400> 15  
 tcaacatc 8  
 40  
 <210> 16  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence  
 45  
 <220>  
 <223> LNA Oligomer  
 <220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Fully LNA & Phosphorothioate linkages  
 50  
 <400> 16  
 ggtaaact 8  
 55  
 <210> 17  
 <211> 8

<212> DNA  
 <213> Artificial Sequence

<220>  
 5 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 10 <223> Fully LNA & Phosphorothioate linkages

<400> 17  
 cgtaatga 8

15 <210> 18  
 <211> 16  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 25 <222> (1)..(16)  
 <223> LNA gapmer 3LNA- 9DNA-3LNA-1DNA, 5' FAM label, phosphorothioate linkages.

<400> 18  
 tcagtctgat aagcta 16

30 <210> 19  
 <211> 8  
 <212> DNA  
 <213> Artificial Sequence

35 <220>  
 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 40 <222> (1)..(8)  
 <223> Full LNA & Phosphorothioate linkages

<400> 19  
 45 agcacttt 8

<210> 20  
 <211> 8  
 <212> DNA  
 50 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer

55 <220>  
 <221> misc\_feature  
 <222> (1)..(8)  
 <223> Full LNA & Phosphorothioate linkages

<400> 20  
attgcac 8

5 <210> 21  
<211> 15  
<212> DNA  
<213> Artificial Sequence

10 <220>  
<223> LNA Oligomer

<400> 21  
agcagacaat gtagc 15

15 <210> 22  
<211> 15  
<212> DNA  
<213> Artificial Sequence

20 <220>  
<223> LNA Oligomer

<400> 22  
gtagccagat gtagc 15

25 <210> 23  
<211> 7  
<212> DNA  
<213> Artificial Sequence

30 <220>  
<223> LNA Oligomer

35 <220>  
<221> misc\_feature  
<222> (1)..(7)  
<223> Full LNA & Phosphorothioate linkages

40 <400> 23  
atgtagc 7

<210> 24  
<211> 15  
<212> DNA

45 <213> Artificial Sequence

<220>  
<223> LNA Oligomer

50 <400> 24  
acaacctact acctc 15

<210> 25  
<211> 8  
<212> DNA

55 <213> Artificial Sequence

<220>

<223> LNA Oligomer  
  
 <220>  
 <221> misc\_feature  
 5 <222> (1)..(8)  
 <223> Full LNA & Phosphorothioate linkages  
  
 <400> 25  
 10 actacctc 8  
  
 <210> 26  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence  
 15  
  
 <220>  
 <223> LNA Oligomer  
  
 <400> 26  
 20 cactgtcagc acttt 15  
  
 <210> 27  
 <211> 15  
 <212> DNA  
 25 <213> Artificial Sequence  
  
 <220>  
 <223> LNA Oligomer  
  
 <400> 27  
 30 tgcatagatt tgcac 15  
  
 <210> 28  
 <211> 7  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> LNA Oligomer  
 40  
  
 <220>  
 <221> misc\_feature  
 <222> (1)..(7)  
 <223> Full LNA & Phosphorothioate linkages  
 45  
  
 <400> 28  
 gtagact 7  
  
 <210> 29  
 50 <211> 6  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 55 <223> LNA Oligomer  
  
 <220>  
 <221> misc\_feature

<222> (1)..(6)  
 <223> Full LNA & Phosphorothioate linkages

5 <400> 29  
 tacctc 6

<210> 30  
 <211> 7  
 <212> DNA  
 10 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer

15 <220>  
 <221> misc\_feature  
 <222> (1)..(7)  
 <223> Full LNA & Phosphorothioate linkages

20 <400> 30  
 ctacctc 7

<210> 31  
 <211> 9  
 25 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer

30 <220>  
 <221> misc\_feature  
 <222> (1)..(9)  
 <223> Full LNA & Phosphorothioate linkages, N = Universal base

35 <400> 31  
 tntacctc 9

<210> 32  
 <211> 9  
 40 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer

45 <220>  
 <221> misc\_feature  
 <222> (1) .. (9)  
 50 <223> Full LNA & Phosphorothioate linkages, N = Universal base

<400> 32  
 tntacctc 9

55 <210> 33  
 <211> 15  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> LNA Oligomer  
  
 <400> 33  
 5 gcaacctact acctc 15  
  
 <210> 34  
 <211> 15  
 <212> DNA  
 10 <213> Artificial Sequence  
  
 <220>  
 <223> LNA Oligomer  
  
 15 <400> 34  
 acaacctcct acctc 15  
  
 <210> 35  
 <211> 15  
 20 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> LNA Oligomer  
  
 25 <400> 35  
 acaaactact acctc 15  
  
 <210> 36  
 <211> 7  
 30 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> LNA Oligomer  
  
 35 <220>  
 <221> misc\_feature  
 <222> (1)..(7)  
 40 <223> Full LNA & Phosphorothioate linkages,  
  
 <400> 36  
 ctacctc 7  
  
 45 <210> 37  
 <211> 7  
 <212> DNA  
 <213> Artificial Sequence  
  
 50 <220>  
 <223> LNA Oligomer  
  
 <220>  
 <221> misc\_feature  
 55 <222> (1)..(7)  
 <223> Full LNA & Phosphorothioate linkages,  
  
 <400> 37

ctaactc 7

<210> 38  
 <211> 9  
 5 <212> DNA  
 <213> Artificial Sequence

<220>  
 10 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(9)  
 15 <223> Full LNA & Phosphorothioate linkages,

<400> 38  
 ttagcatta 9

<210> 39  
 20 <211> 9  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 25 <223> LNA Oligomer

<220>  
 <221> misc\_feature  
 <222> (1)..(9)  
 30 <223> Full LNA & Phosphorothioate linkages,

<400> 39  
 ttagcatta 9

35 <210> 40  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

40 <400> 40  
 uagcaccgcu auccacuaug uc 22

<210> 41  
 <211> 24  
 45 <212> RNA  
 <213> Epstein Barr virus

<400> 41  
 ucuuagugga agugacgugc ugug 24  
 50

<210> 42  
 <211> 23  
 <212> RNA  
 <213> Epstein Barr virus  
 55

<400> 42  
 uacauaacca uggaguuggc ugu 23



<210> 43  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 5  
 <400> 43  
 gccaccucuu ugguucugua ca 22  
 <210> 44  
 <211> 21  
 <212> RNA  
 <213> Epstein Barr virus  
 10  
 <400> 44  
 acgcacacca ggcugacugc c 21  
 15  
 <210> 45  
 <211> 24  
 <212> RNA  
 <213> Epstein Barr virus  
 20  
 <400> 45  
 ucagacaguu uggugcgua guug 24  
 25  
 <210> 46  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 30  
 <400> 46  
 uccuguggug uuuggugugg uu 22  
 35  
 <210> 47  
 <211> 23  
 <212> RNA  
 <213> Epstein Barr virus  
 40  
 <400> 47  
 uguaacuugc cagggacggc uga 23  
 45  
 <210> 48  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 50  
 <400> 48  
 aaccggcucg uggcucguac ag 22  
 55  
 <210> 49  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 <400> 49  
 uaaaugcugc aguaguaggg au 22  
 <210> 50  
 <211> 22

<212> RNA  
 <213> Epstein Barr virus

<400> 50  
 5 uaccuacgc ugccgauua ca 22

<210> 51  
 <211> 22  
 <212> RNA  
 10 <213> Epstein Barr virus

<400> 51  
 gucagugguu uuguuuccu ga 22

15 <210> 52  
 <211> 24  
 <212> RNA  
 <213> Epstein Barr virus

20 <400> 52  
 uuagauagag ugggugugug cucu 24

25 <210> 53  
 <211> 23  
 <212> RNA  
 <213> Epstein Barr virus

30 <400> 53  
 uguaugccug gugucccuu agu 23

35 <210> 54  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

<400> 54  
 uaagaggacg caggcauaca ag 22

40 <210> 55  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

45 <400> 55  
 uaucggaagu uugggcuucg uc 22

<210> 56  
 <211> 22  
 <212> RNA  
 50 <213> Epstein Barr virus

<400> 56  
 ucaaguucgc acuuccuaua ca 22

55 <210> 57  
 <211> 21  
 <212> RNA  
 <213> Epstein Barr virus

<400> 57  
 uuuuguuugc uugggaaugc u 21

5  
 <210> 58  
 <211> 23  
 <212> RNA  
 <213> Epstein Barr virus

10  
 <400> 58  
 acauuccccg caaacaugac aug 23

15  
 <210> 59  
 <211> 24  
 <212> RNA  
 <213> Epstein Barr virus

20  
 <400> 59  
 aaggagcgau uuggagaaaa uaaa 24

25  
 <210> 60  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

30  
 <400> 60  
 uauuuucugc auucgccuu gc 22

35  
 <210> 61  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

40  
 <400> 61  
 caugaaggca cagccuguua cc 22

45  
 <210> 62  
 <211> 21  
 <212> RNA  
 <213> Epstein Barr virus

50  
 <400> 62  
 uagcaggcau gucucauuc c 21

55  
 <210> 63  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

<400> 63  
 cgcaccacua gucaccaggu gu 22

<210> 64  
 <211> 21  
 <212> RNA  
 <213> Epstein Barr virus

<400> 64  
 accuaguguu aguguugugc u 21

<210> 65  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 5  
 <400> 65  
 gaccugaucg ugcuggugug cu 22  
 <210> 66  
 <211> 24  
 <212> RNA  
 <213> Epstein Barr virus  
 10  
 <400> 66  
 caaggugaau auagcugccc aucg 24  
 <210> 67  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 20  
 <400> 67  
 cggggaucgg acuagccuua ga 22  
 <210> 68  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 25  
 <400> 68  
 uaagguuggu ccaauccaau gg 22  
 <210> 69  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 35  
 <400> 69  
 caucuauguc caguguccag gg 22  
 <210> 70  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 40  
 <400> 70  
 ccuggaccuu gacuaugaaa ca 22  
 <210> 71  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus  
 45  
 <400> 71  
 uacgguuucc uagauuguac ag 22  
 <210> 72  
 <211> 23

<212> RNA  
 <213> Epstein Barr virus

5 <400> 72  
 gucacaauuc auggggucgu aga 23

10 <210> 73  
 <211> 23  
 <212> RNA  
 <213> Epstein Barr virus

15 <400> 73  
 uaacacuuc auggguccgu agu 23

20 <210> 74  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

25 <400> 74  
 uacuggaccc ugaauuggaa ac 22

30 <210> 75  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

35 <400> 75  
 uaaccugauc agccccggag uu 22

40 <210> 76  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

45 <400> 76  
 uaucuuuugc ggcagaaau ga 22

50 <210> 77  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

55 <400> 77  
 aaauucuguu gcagcagaua gc 22

60 <210> 78  
 <211> 22  
 <212> RNA  
 <213> Epstein Barr virus

65 <400> 78  
 uaacgggaag uguguaagca ca 22

70 <210> 79  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus

<400> 79  
 aagugacggu gagauccagg cu 22

5 <210> 80  
 <211> 21  
 <212> RNA  
 <213> Human cytomegalovirus

10 <400> 80  
 ucguccuccc cuucucacc g 21

15 <210> 81  
 <211> 20  
 <212> RNA  
 <213> Human cytomegalovirus

20 <400> 81  
 uaacuagccu ucccgugaga 20

25 <210> 82  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus

30 <400> 82  
 ucaccagaau gcuaguugu ag 22

35 <210> 83  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus

40 <400> 83  
 ucguugaaga caccuggaaa ga 22

45 <210> 84  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus

50 <400> 84  
 uuuccaggug uuuucaacgu gc 22

55 <210> 85  
 <211> 20  
 <212> RNA  
 <213> Human cytomegalovirus

<400> 85  
 ggggaugggc uggcgcgcg 20

<210> 86  
 <211> 21  
 <212> RNA  
 <213> Human cytomegalovirus

<400> 86  
 ugcgucucgg ccucguccag a 21

<210> 87  
 <211> 21  
 <212> RNA  
 <213> Human cytomegalovirus  
 5  
 <400> 87  
 aaccgcucag uggcucggac c 21  
 <210> 88  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus  
 10  
 <400> 88  
 uccgaacgcu aggucgguuc uc 22  
 <210> 89  
 <211> 23  
 <212> RNA  
 <213> Human cytomegalovirus  
 20  
 <400> 89  
 auccacuugg agagcucccg cgg 23  
 <210> 90  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus  
 25  
 <400> 90  
 agcggucugu ucagguggau ga 22  
 <210> 91  
 <211> 20  
 <212> RNA  
 <213> Human cytomegalovirus  
 35  
 <400> 91  
 ucacgguccg agcacaucca 20  
 <210> 92  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus  
 40  
 <400> 92  
 gauugugccc ggaccguggg cg 22  
 <210> 93  
 <211> 22  
 <212> RNA  
 <213> Human cytomegalovirus  
 45  
 <400> 93  
 cgacauggac gugcaggggg au 22  
 <210> 94  
 <211> 21

<212> RNA  
 <213> Human cytomegalovirus

<400> 94  
 5 ugacaagccu gacgagagcg u 21

<210> 95  
 <211> 22  
 <212> RNA  
 10 <213> Human cytomegalovirus

<400> 95  
 uuaugauagg ugugacgaug uc 22

15 <210> 96  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 96  
 ugagguagua gguuguauag uu 22

<210> 97  
 <211> 21  
 25 <212> RNA  
 <213> Homo sapiens

<400> 97  
 30 cuauacaauc uacugucuuu c 21

<210> 98  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35

<400> 98  
 ugagguagua gguugugugg uu 22

<210> 99  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40

<400> 99  
 45 cuauacaacc uacugccuuc cc 22

<210> 100  
 <211> 22  
 <212> RNA  
 50 <213> Homo sapiens

<400> 100  
 ugagguagua gguuguauagg uu 22

55 <210> 101  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens



<400> 101  
uagaguuaca cccugggagu ua 22

5 <210> 102  
<211> 22  
<212> RNA  
<213> Homo sapiens

10 <400> 102  
agagguagua gguugcauag uu 22

15 <210> 103  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 103  
cuauacgacc ugcugccuuu cu 22

25 <210> 104  
<211> 22  
<212> RNA  
<213> Homo sapiens

30 <400> 104  
ugagguagga gguuguauag uu 22

35 <210> 105  
<211> 22  
<212> RNA  
<213> Homo sapiens

40 <400> 105  
cuauacggcc uccuagccuuu cc 22

45 <210> 106  
<211> 22  
<212> RNA  
<213> Homo sapiens

50 <400> 106  
ugagguagua gauuguauag uu 22

55 <210> 107  
<211> 22  
<212> RNA  
<213> Homo sapiens

60 <400> 107  
cuauacauc uauugccuuc cc 22

65 <210> 108  
<211> 22  
<212> RNA  
<213> Homo sapiens

70 <400> 108  
cuauacaguc uacuguccuuu cc 22

<210> 109  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 109  
 ugagguagua guuuguacag uu 22  
 <210> 110  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 110  
 cuguacaggc cacugccuug c 21  
 15  
 <210> 111  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 111  
 ugagguagua guuugugcug uu 22  
 25  
 <210> 112  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 112  
 cugcgcaagc uacugccuug cu 22  
 35  
 <210> 113  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 113  
 uggaauguaa agaaguaugu au 22  
 45  
 <210> 114  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 114  
 aaccguaga uccgaacuug ug 22  
 55  
 <210> 115  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 <400> 115  
 caagcuugua ucuauaggua ug 22  
 <210> 116  
 <211> 21

<212> RNA  
<213> Homo sapiens

<400> 116  
5 uacaguacug ugauaacuga a 21

<210> 117  
<211> 22  
<212> RNA  
10 <213> Homo sapiens

<400> 117  
caguuauacac agugcugaug cu 22

<210> 118  
<211> 23  
<212> RNA  
15 <213> Homo sapiens

<400> 118  
20 agcagcauug uacagggcua uga 23

<210> 119  
<211> 23  
<212> RNA  
25 <213> Homo sapiens

<400> 119  
ucauagcccu guacaaugcu gcu 23  
30

<210> 120  
<211> 23  
<212> RNA  
<213> Homo sapiens  
35

<400> 120  
ucaaauugcuc agacuccugu ggu 23

<210> 121  
<211> 22  
<212> RNA  
40 <213> Homo sapiens

<400> 121  
45 acggauguuu gagcaugugc ua 22

<210> 122  
<211> 23  
<212> RNA  
50 <213> Homo sapiens

<400> 122  
aaaagugcuu acagugcagg uag 23

<210> 123  
<211> 22  
<212> RNA  
55 <213> Homo sapiens

<400> 123  
 cugcaaugua agcacuucuu ac 22

5  
 <210> 124  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 124  
 uaaagugcug acagugcaga u 21

15  
 <210> 125  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 125  
 ccgcacugug gguacuugcu gc 22

25  
 <210> 126  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 126  
 agcagcauug uacagggcua uca 23

35  
 <210> 127  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 127  
 uaccuguag aaccgaauuu gug 23

45  
 <210> 128  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 128  
 caaaauucgua ucuaggggaa ua 22

55  
 <210> 129  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 129  
 uaccuguag aaccgaauuu gug 23

<210> 130  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 130  
 acagauucga uucuagggga au 22

<210> 131  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 131  
 uugcucacug uucuuccua g 21  
 10  
 <210> 132  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 15  
 <400> 132  
 aagcauucuu ucauugguug g 21  
 20  
 <210> 133  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 <400> 133  
 uuuccggcuc gcgugggugu gu 22  
 25  
 <210> 134  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 134  
 ccgucgccgc cacccgagcc g 21  
 35  
 <210> 135  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 <400> 135  
 gagggucuug ggagggaugu gac 23  
 40  
 <210> 136  
 <211> 27  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 136  
 cacuguaggu gauggugaga gugggca 27  
 50  
 <210> 137  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 55  
 <400> 137  
 ccugcagcga cuugauggcu ucc 23  
 <210> 138  
 <211> 21

<212> RNA  
 <213> Homo sapiens

5 <400> 138  
 agaggauacc cuuuguaugu u 21

10 <210> 139  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

15 <400> 139  
 uaggacacau ggucuacuuc u 21

20 <210> 140  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 140  
 cuccugagcc auucugagcc uc 22

30 <210> 141  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

35 <400> 141  
 agccugauua aacacaugcu cuga 24

40 <210> 142  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

45 <400> 142  
 gugccagcug caguggggga g 21

50 <210> 143  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

55 <400> 143  
 cccggagcca ggaugcagcu c 21

<210> 144  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 144  
 ucguggccug guccauua u 21

<210> 145  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 145  
 ucugcagggg uugcuuugag 20

5  
 <210> 146  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 146  
 uguucaugua gauguuuag c 21

15  
 <210> 147  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 147  
 ucagcuggcc cucuuuc 18

25  
 <210> 148  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 148  
 uggcagggag gcuggaggg g 21

35  
 <210> 149  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 149  
 ucacuguuca gacaggcggg 20

45  
 <210> 150  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 150  
 uggaguguga caaugguguu ug 22

55  
 <210> 151  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 151  
 aacgccauua ucacacuaaa ua 22

<210> 152  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 152  
 ccccaccucc ucucuccua g 21

<210> 153  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 153  
 gugaggacuc gggaggugg 19  
 10  
 <210> 154  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 15  
 <400> 154  
 ugagccccug ugccgcccc ag 22  
 20  
 <210> 155  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 <400> 155  
 guggguacgg cccagugggg gg 22  
 25  
 <210> 156  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 156  
 ucaccagccc uguguucccu ag 22  
 35  
 <210> 157  
 <211> 26  
 <212> RNA  
 <213> Homo sapiens  
 <400> 157  
 gugagggcau gcaggccugg augggg 26  
 40  
 <210> 158  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 158  
 cgugccaccc uuuucccag 20  
 50  
 <210> 159  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 55  
 <400> 159  
 ucacaccugc cugcccccc 20  
 <210> 160  
 <211> 21



<212> RNA  
 <213> Homo sapiens

5 <400> 160  
 gugggcgggg gcaggugugu g 21

10 <210> 161  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

15 <400> 161  
 cucucaccac ugccuccca cag 23

20 <210> 162  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

25 <400> 162  
 gugucugggc ggacagcugc 20

30 <210> 163  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

35 <400> 163  
 ugagcccugu ccucccgag 20

40 <210> 164  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 164  
 ucggccugac cacccacccc ac 22

50 <210> 165  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 165  
 ccucuucucc uugucucucc ag 22

<210> 166  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 166  
 uccuucugcu ccgucccca g 21

<210> 167  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 167  
 cuuccucguc ugucugcccc 20

5  
 <210> 168  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 168  
 uaaggcacgc ggugaaugcc 20

15  
 <210> 169  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 169  
 cguguucaca gcggaccuug au 22

25  
 <210> 170  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 170  
 aacuggauca auuauaggag ug 22

35  
 <210> 171  
 <211> 26  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 171  
 aaguaguugg uuuguaugag augguu 26

45  
 <210> 172  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 172  
 aagugaucua aaggccuaca u 21

55  
 <210> 173  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 173  
 aauggauuuu uggagcagg 19

<210> 174  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 174  
 acccgucccg uucgucccg ga 22

<210> 175  
 <211> 27  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 175  
 accuucuugu auaagcacug ugcuaaa 27  
 <210> 176  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 176  
 acgcccuucc ccccuucuu ca 22  
 <210> 177  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 177  
 acggugcugg auguggccuu u 21  
 <210> 178  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 178  
 acucuagcug ccaaaggcgc u 21  
 <210> 179  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 179  
 agaaggaaau ugaauucauu ua 22  
 <210> 180  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 180  
 agagaagaag aucagccugc a 21  
 <210> 181  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 181  
 agccuggaag cuggagccug cagu 24  
 <210> 182  
 <211> 23

<212> RNA  
 <213> Homo sapiens

5 <400> 182  
 aggaugagca aagaaaguag auu 23

10 <210> 183  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 183  
 cggaugagca aagaaagugg uu 22

20 <210> 184  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 184  
 aggcauugac uucucacuag cu 22

30 <210> 185  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

35 <400> 185  
 agugaaugau ggguuucugac c 21

40 <210> 186  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

45 <400> 186  
 aguuaggauu aggucgugga a 21

50 <210> 187  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

55 <400> 187  
 auauaugaug acuuagcuuu u 21

<210> 188  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 188  
 acaggugagg uucuugggag cc 22

<210> 189  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

<400> 189  
 ucccugagac ccuuuaccu guga 24

5  
 <210> 190  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 190  
 ucccugagac ccuaacuugu ga 22

15  
 <210> 191  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 191  
 acggguuagg cucuugggag cu 22

25  
 <210> 192  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 192  
 ucacaaguca ggcucuuggg ac 22

35  
 <210> 193  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 193  
 ucguaccgug aguaauaaug cg 22

45  
 <210> 194  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 194  
 cauuaauacu uuugguacgc g 21

55  
 <210> 195  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens

<400> 195  
 auccaccuc ugccacca 18

<210> 196  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 196  
 auggauaagg cuuuggcuu 19

<210> 197  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 197  
 augggugaau uuguagaagg au 22  
 <210> 198  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 198  
 augguacccu ggcauacuga gu 22  
 <210> 199  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 199  
 caagucuuau uugagcaccu guu 23  
 <210> 200  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 200  
 caggaugugg ucaaguguug uu 22  
 <210> 201  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 201  
 ccucagggcu guagaacagg gcu 23  
 <210> 202  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 202  
 ccuguugaag uguaaucccc a 21  
 <210> 203  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 203  
 cgggcguggu gguggggg 18  
 <210> 204  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 204  
 cuggacugag ccgugcuacu gg 22

10 <210> 205  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 205  
 ucggauccgu cugagcuugg cu 22

20 <210> 206  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 206  
 cugaagcuca gagggcucug au 22

30 <210> 207  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

35 <400> 207  
 cuggagauau ggaagagcug ugu 23

40 <210> 208  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 208  
 cuuggcaccu agcaagcacu ca 22

50 <210> 209  
 <211> 26  
 <212> RNA  
 <213> Homo sapiens

55 <400> 209  
 gaugaugaug gcagcaaaau cugaaa 26

60 <210> 210  
 <211> 25  
 <212> RNA  
 <213> Homo sapiens

65 <400> 210  
 gggcgacaaa gcaagacucu uucuu 25

70 <210> 211  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens

<400> 211  
 gucccuguuc aggcgcca 18

5  
 <210> 212  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 212  
 ucccuguucg ggcgcca 17

15  
 <210> 213  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 213  
 gugggggaga ggcuguc 17

25  
 <210> 214  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 214  
 uaaagagccc uguggagaca 20

35  
 <210> 215  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 215  
 uacguagaua uauauguauu uu 22

45  
 <210> 216  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 216  
 uaguacugug cauaucaucu au 22

55  
 <210> 217  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens

<400> 217  
 ucatauugcu ucuuucu 17

<210> 218  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 218  
 ucacagugaa ccggucucu u 21



<210> 219  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 219  
 ucccaccgcu gccaccc 17  
 <210> 220  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 220  
 ucgccuccuc cucucc 17  
 <210> 221  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 221  
 ucguuugccu uuuucugcu 20  
 <210> 222  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 222  
 ucuacaaagg aaagcgcuu cu 22  
 <210> 223  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 223  
 ucuauacaga cccuggcuu uc 22  
 <210> 224  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 224  
 ucugggcaac aaagugagac cu 22  
 <210> 225  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 225  
 ucgaggacca agaugagccc u 21  
 <210> 226  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 226  
 ugcuggauca gugguucgag uc 22

10 <210> 227  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

15 <400> 227  
 uggacugccc ugaucuggag a 21

20 <210> 228  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

25 <400> 228  
 uggaguccag gaaucugcau uuu 23

30 <210> 229  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 229  
 aagcccuuac ccaaaaaagu au 22

40 <210> 230  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 230  
 aagcccuuac ccaaaaaagc au 22

50 <210> 231  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

55 <400> 231  
 cuuuuugcgg ucugggcuug c 21

<210> 232  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 232  
 uggauuuuug gaucagga 19

<210> 233  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

<400> 233  
 uggcccugac ugaagaccag cagu 24

5  
 <210> 234  
 <211> 25  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 234  
 ugggaacggg uuccggcaga cgcug 25

15  
 <210> 235  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 235  
 uggguggucu ggagauuugu gc 22

25  
 <210> 236  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 236  
 ugugagguug gcauuguugu cu 22

35  
 <210> 237  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 237  
 uuaggccgca gaucugggug a 21

45  
 <210> 238  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 238  
 uuagggcccu ggcuccaucu cc 22

55  
 <210> 239  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens

<400> 239  
 uucaaguaau ucaggug 17

<210> 240  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 240  
 uucauucggc uguccagaug ua 22

<210> 241  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 241  
 uucuggaauu cugugagagg ga 22  
 <210> 242  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 242  
 uugagaagga ggcugcug 18  
 <210> 243  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 243  
 uugcagcugc cuggaguga cuuc 24  
 <210> 244  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 244  
 uuggacaua cuuauacuaa a 21  
 <210> 245  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 245  
 uuuagagacg gggucuugcu cu 22  
 <210> 246  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 246  
 uuugaggcua cagugagaug ug 22  
 <210> 247  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 247  
 uuuucaacuc uauugggaga ga 22  
 <210> 248  
 <211> 18

<212> RNA  
 <213> Homo sapiens

5 <400> 248  
 acguuggcuc ugguggug 18

10 <210> 249  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 249  
 acucggcgug gcgucggucg ug 22

20 <210> 250  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens

25 <400> 250  
 gcaugggugg uucagugg 18

30 <210> 251  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 251  
 cagugcaaug uuaaaagggc au 22

40 <210> 252  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 252  
 uucacauugu gcuacugucu gc 22

50 <210> 253  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 253  
 cagugcaaug augaaagggc au 22

60 <210> 254  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

65 <400> 254  
 acucuuucc uguugcacua c 21

70 <210> 255  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 255		
	uaacagucua cagccauggu cg	22	
5	<210> 256		
	<211> 22		
	<212> RNA		
	<213> Homo sapiens		
10	<400> 256		
	accguggcuu ucgauuguua cu	22	
15	<210> 257		
	<211> 18		
	<212> RNA		
	<213> Homo sapiens		
20	<400> 257		
	cagggaggug aaugugau	18	
25	<210> 258		
	<211> 19		
	<212> RNA		
	<213> Homo sapiens		
30	<400> 258		
	gaugaugcug cugaugcug	19	
35	<210> 259		
	<211> 22		
	<212> RNA		
	<213> Homo sapiens		
40	<400> 259		
	ucaaaacuga ggggcacuuuu cu	22	
45	<210> 260		
	<211> 24		
	<212> RNA		
	<213> Homo sapiens		
50	<400> 260		
	ccagacagaa uucuaugcac uuuc	24	
55	<210> 261		
	<211> 22		
	<212> RNA		
	<213> Homo sapiens		
60	<400> 261		
	uuuggucucc uucaaccagc ug	22	
65	<210> 262		
	<211> 22		
	<212> RNA		
	<213> Homo sapiens		
70	<400> 262		
	uuuggucucc uucaaccagc ua	22	

<210> 263  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 263  
 ugugacuggu ugaccagagg gg 22  
 <210> 264  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 264  
 uauggcuuuu uauuccuaug uga 23  
 <210> 265  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 265  
 uauagggauu ggagccgugg cg 22  
 <210> 266  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 266  
 uauggcuuuu cauuccuaug uga 23  
 <210> 267  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 267  
 auguagggcu aaaagccaug gg 22  
 <210> 268  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 268  
 acuccauuug uuuugaugau gga 23  
 <210> 269  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 269  
 caucaucguc ucaaaugagu cu 22  
 <210> 270  
 <211> 23

<212> RNA  
 <213> Homo sapiens

5 <400> 270  
 uuauugcuua agaauacgcg uag 23

10 <210> 271  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

15 <400> 271  
 agcugguguu gugaacagg ccg 23

20 <210> 272  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 272  
 gcuacuucac aacaccaggg cc 22

30 <210> 273  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 273  
 gcuauuucac gacaccaggg uu 22

40 <210> 274  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 274  
 ggagacgagg ccuguugga gu 22

50 <210> 275  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 275  
 ucuacagugc acgugucucc ag 22

<210> 276  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 276  
 uaccacaggg uagaaccacg g 21

<210> 277  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens



<400> 277  
 cagugguuuu acccuauggu ag 22

5  
 <210> 278  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 278  
 uaacacuguc ugguaaagau gg 22

15  
 <210> 279  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 279  
 caucuuccag uacaguguug ga 22

25  
 <210> 280  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 280  
 uguaguuuu ccuacuuuau gga 23

35  
 <210> 281  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 281  
 cauaaaguag aaagcacuac u 21

45  
 <210> 282  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 282  
 ugagaugaag cacuguagcu c 21

55  
 <210> 283  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 283  
 ggugcagugc ugcaucucug gu 22

<210> 284  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 284  
 uacagauag augauguacu 20

<210> 285  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 285  
 ggauaucauc auauacugua ag 22  
 <210> 286  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 286  
 guccaguuuu cccaggauc ccu 23  
 <210> 287  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 287  
 ggauuccugg aaauacuguu cu 22  
 <210> 288  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 288  
 cuccguuugc cuguuucgu g 21  
 <210> 289  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 289  
 cucggcgcgg ggcgcgggcu cc 22  
 <210> 290  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 290  
 ugagaacuga auuccauggg uu 22  
 <210> 291  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 291  
 ccucugaaau ucaguucuuc ag 22  
 <210> 292  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 292  
 ugcccugugg acucaguucu gg 22

10 <210> 293  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 293  
 ugagaacuga auuccauagg cu 22

20 <210> 294  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

25 <400> 294  
 guguguggaa augcuucugc 20

30 <210> 295  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

35 <400> 295  
 gccuccgcc cgugcacccc g 21

40 <210> 296  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 296  
 gcccgugugu ggagccaggu gu 22

50 <210> 297  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 297  
 gugugcggaa augcuucugc ua 22

<210> 298  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 298  
 ucagugcacu acagaacuuu gu 22

<210> 299  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 299	
	aaaguucuga gacacuccga cu	22
5	<210> 300 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 300	
	ucagugcauc acagaacuuu gu	22
15	<210> 301 <211> 22 <212> RNA <213> Homo sapiens	
20	<400> 301	
	aaguucuguu auacacucag gc	22
25	<210> 302 <211> 23 <212> RNA <213> Homo sapiens	
30	<400> 302	
	ucuggcuccg ugucuucacu ccc	23
35	<210> 303 <211> 21 <212> RNA <213> Homo sapiens	
40	<400> 303	
	agggaggac gggggcugug c	21
45	<210> 304 <211> 22 <212> RNA <213> Homo sapiens	
50	<400> 304	
	ucuccaacc cuuguaccag ug	22
55	<210> 305 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 305	
	cugguacagg ccugggggac ag	22
	<210> 306 <211> 21 <212> RNA <213> Homo sapiens	
	<400> 306	
	cuagacugaa gcuccuugag g	21

<210> 307  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 307  
 ucgaggagcu cacagucuag u 21  
 <210> 308  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 308  
 ucagugcaug acagaacuug g 21  
 <210> 309  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 309  
 uugcauaguc acaaaguga uc 22  
 <210> 310  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 310  
 aaaaccgucu aguuacaguu gu 22  
 <210> 311  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 311  
 cggcccgggc ugcugcuguu ccu 23  
 <210> 312  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 312  
 uccugcgcu cccagaugcc c 21  
 <210> 313  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 313  
 uagguuaucc guguugccuu cg 22  
 <210> 314  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 314  
 aaucauacac gguugaccua uu 22

10 <210> 315  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

15 <400> 315  
 uuaaugcuaa ucgugauagg ggu 23

20 <210> 316  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 316  
 cuccuacaua uuagcauuaa ca 22

30 <210> 317  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 317  
 uagcagcaca uaaugguuug ug 22

40 <210> 318  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 318  
 caggccauau ugugcugccu ca 22

50 <210> 319  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 319  
 uagcagcaca ucaugguuuu ca 22

<210> 320  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 320  
 cgaaucauua uuugcugcuc ua 22

<210> 321  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 321 uagcagcacg uaaauauugg cg	22
5	<210> 322 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 322 ccaguauuaa cugugcugcu ga	22
15	<210> 323 <211> 22 <212> RNA <213> Homo sapiens	
20	<400> 323 ccaauuuuac ugugcugcuu ua	22
25	<210> 324 <211> 23 <212> RNA <213> Homo sapiens	
30	<400> 324 caaagugcuu acagucgagg uag	23
35	<210> 325 <211> 22 <212> RNA <213> Homo sapiens	
40	<400> 325 acugcaguga aggcacuugu ag	22
45	<210> 326 <211> 23 <212> RNA <213> Homo sapiens	
50	<400> 326 aacaaucaac gcugucggug agu	23
55	<210> 327 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 327 accaucgacc guugauugua cc	22
	<210> 328 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 328 accacugacc guugacugua cc	22

<210> 329  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 329  
 aacaucauu gcugucggug ggu 23  
 <210> 330  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 330  
 aacaucaac cugucgguga gu 22  
 <210> 331  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 331  
 aaccaucgac cguugagugg ac 22  
 <210> 332  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 332  
 aacaucauu guugucggug ggu 23  
 <210> 333  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 333  
 uuuggcaaug guagaacuca cacu 24  
 <210> 334  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 334  
 ugguucuaga cuugccaacu a 21  
 <210> 335  
 <211> 18  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 335  
 uccagugccc uccucucc 18  
 <210> 336  
 <211> 27



<212> RNA  
 <213> Homo sapiens  
  
 <400> 336  
 5 auugaucauc gacacuucga acgcaau 27  
  
 <210> 337  
 <211> 18  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 337  
 ugaggcagua gauugaau 18  
  
 15 <210> 338  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 338  
 uauggcacug guagaauuca cu 22  
  
 <210> 339  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 339  
 gugaauuacc gaaggccau aa 22  
 30  
  
 <210> 340  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
  
 <400> 340  
 uggacggaga acugauaagg gu 22  
  
 40 <210> 341  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 <400> 341  
 45 uggagagaaa ggcaguuccu ga 22  
  
 <210> 342  
 <211> 22  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 342  
 aggggcuggc uuuccucugg uc 22  
  
 55 <210> 343  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 343 caaagaauc uccuuuggg cu	22
5	<210> 344 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 344 gcccaaaggu gaauuuuug gg	22
15	<210> 345 <211> 22 <212> RNA <213> Homo sapiens	
20	<400> 345 ucgugucuug uguugcagcc gg	22
25	<210> 346 <211> 22 <212> RNA <213> Homo sapiens	
30	<400> 346 ggcuacaaca caggaccgg gc	22
35	<210> 347 <211> 21 <212> RNA <213> Homo sapiens	
40	<400> 347 cuccacaug cagguuugc a	21
45	<210> 348 <211> 21 <212> RNA <213> Homo sapiens	
50	<400> 348 caucccuugc augguggagg g	21
55	<210> 349 <211> 23 <212> RNA <213> Homo sapiens	
	<400> 349 uaaggugcau cuagucaga uag	23
	<210> 350 <211> 23 <212> RNA <213> Homo sapiens	
	<400> 350 acugccuaa gugcuccuuc ugg	23

<210> 351  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 351  
 uaaggugcau cuagugcagu uag 23  
 <210> 352  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 352  
 ugcccuaaaau gcccucug gc 22  
 <210> 353  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 353  
 ugauauguuu gauauuuag gu 22  
 <210> 354  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 354  
 cggcggggac ggcgauuggu c 21  
 <210> 355  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 355  
 cgcaggggcc gggugcucac cg 22  
 <210> 356  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 356  
 ugagugccgg ugccugccu g 21  
 <210> 357  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 357  
 ugauauguuu gauauugggu u 21  
 <210> 358  
 <211> 23

<212> RNA  
 <213> Homo sapiens

<400> 358  
 5 caacggaauc ccaaaagcag cug 23

<210> 359  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens

<400> 359  
 gcugcgcug gauuucgucc cc 22

<210> 360  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 360  
 20 ccaguccugu gccugccgc u 21

<210> 361  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 361  
 30 ugaguaccgc caugucuguu ggg 23

<210> 362  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 362  
 35 caccaggcau uguggucucc 20

<210> 363  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 363  
 45 uaccagagc augcagugug aa 22

<210> 364  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 364  
 ucugccccc ccgucgucg ca 22

<210> 365  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 365  
 cccugugccc ggcccacuuc ug 22

5  
 <210> 366  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 366  
 ggaggggucc cgcacuggga gg 22

15  
 <210> 367  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 367  
 ccccagggcg acgcggcggg 20

25  
 <210> 368  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 368  
 accuugccuu gcugcccggg cc 22

35  
 <210> 369  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 369  
 cugaccuaug aaugacagc c 21

45  
 <210> 370  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 370  
 cugccaauuc cauaggucac ag 22

55  
 <210> 371  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 371  
 aacuggccua caaagucca gu 22

<210> 372  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 372  
 ugggucuug cgggcgagau ga 22

<210> 373  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 373  
 aacuggcccu caaagucccg cu 22  
 <210> 374  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 374  
 cgggguuuug agggcgagau ga 22  
 <210> 375  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 375  
 uguaacagca acuccaugug ga 22  
 <210> 376  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 376  
 ccaguggggc ugcuguuau c 22  
 <210> 377  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 377  
 uagcagcaca gaaauuugg c 21  
 <210> 378  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 378  
 ccaauuugg cugucugcu cc 22  
 <210> 379  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 379  
 uagguaguuu cauguuguug gg 22  
 <210> 380  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 380  
 cggcaacaag aaacugccug ag 22

10 <210> 381  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 381  
 uagguaguuu ccuguuguug gg 22

20 <210> 382  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 382  
 uucaccaccu ucuccacca gc 22

30 <210> 383  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 383  
 gguccagagg ggagauaggu uc 22

40 <210> 384  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

45 <400> 384  
 cccaguguuc agacuaccug uuc 23

50 <210> 385  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 385  
 acaguagucu gcacauuggu ua 22

60 <210> 386  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

65 <400> 386  
 cccaguguuu agacuaucug uuc 23

70 <210> 387  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

	<400> 387	
	ugugcaaauc uaugcaaaac uga	23
5	<210> 388 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 388	
	aguuuugcau aguugcacua ca	22
15	<210> 389 <211> 23 <212> RNA <213> Homo sapiens	
20	<400> 389	
	ugugcaaauc caugcaaaac uga	23
25	<210> 390 <211> 23 <212> RNA <213> Homo sapiens	
30	<400> 390	
	aguuuugcag guuugcaucc age	23
35	<210> 391 <211> 22 <212> RNA <213> Homo sapiens	
40	<400> 391	
	aguuuugcag guuugcauuu ca	22
45	<210> 392 <211> 22 <212> RNA <213> Homo sapiens	
50	<400> 392	
	uaacacuguc ugguaacgau gu	22
55	<210> 393 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 393	
	caucuaccg gacagugcug ga	22
	<210> 394 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 394	
	uaauacugcc ugguaaugau ga	22



<210> 395  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 395  
 caucuacug ggcagcauug ga 22  
 <210> 396  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 396  
 uaauacugcc ggguaaugau gga 23  
 <210> 397  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 397  
 cgucuuacc agcaguguuu gg 22  
 <210> 398  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 398  
 agagguauag ggcaugggaa 20  
 <210> 399  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 399  
 uuccuaugca uauacuucu ug 22  
 <210> 400  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 400  
 gugaauguu uaggaccacu ag 22  
 <210> 401  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 401  
 uucccuugu cauccaugc cu 22  
 <210> 402  
 <211> 22

<212> RNA  
 <213> Homo sapiens  
  
 <400> 402  
 5 uccuucauuc caccggaguc ug 22  
  
 <210> 403  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 403  
 uggaauguaa ggaagugugu gg 22  
  
 <210> 404  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 404  
 auaagacgag caaaaagcuu gu 22  
  
 <210> 405  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 405  
 30 auaagacgaa caaaagguuu gu 22  
  
 <210> 406  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 406  
 uaaagugcuu auagugcagg uag 23  
  
 <210> 407  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 <400> 407  
 45 acugcauuau gagcacuuaa ag 22  
  
 <210> 408  
 <211> 23  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 408  
 caaagugcuc auagugcagg uag 23  
  
 <210> 409  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 409 acuguaguau gggcacuucc ag	22
5	<210> 410 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 410 uagcuuauca gacugauguu ga	22
15	<210> 411 <211> 21 <212> RNA <213> Homo sapiens	
20	<400> 411 caacaccagu cgaugggcug u	21
25	<210> 412 <211> 22 <212> RNA <213> Homo sapiens	
30	<400> 412 cugugcgugu gacagcggcu ga	22
35	<210> 413 <211> 22 <212> RNA <213> Homo sapiens	
40	<400> 413 uucccuugu cauccuucgc cu	22
45	<210> 414 <211> 21 <212> RNA <213> Homo sapiens	
50	<400> 414 uaacagucuc cagucacggc c	21
55	<210> 415 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 415 acagcaggca cagacaggca gu	22
	<210> 416 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 416 ugccugucua cacuugcugu gc	22

<210> 417  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 417  
 augaccuaug aauugacaga c 21  
 <210> 418  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 418  
 uaaucucagc uggcaacugu ga 22  
 <210> 419  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 419  
 aaaucucugc aggcaaau gu ga 22  
 <210> 420  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 420  
 uacugcauca ggaacugauu gga 23  
 <210> 421  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 421  
 uugugcuuga ucuaccaug u 21  
 <210> 422  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 422  
 augguuccgu caagaccau gg 22  
 <210> 423  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 423  
 caugguucug ucaagcaccg cg 22  
 <210> 424  
 <211> 22

<212> RNA  
 <213> Homo sapiens

5 <400> 424  
 agaguugagu cuggacgucc cg 22

10 <210> 425  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 425  
 agaaauuggg cuggacaucu gu 22

20 <210> 426  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

25 <400> 426  
 ugauugucca aacgcaauuc u 21

30 <210> 427  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 427  
 aagcugccag uugaagaacu gu 22

40 <210> 428  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 428  
 aguucuucag uggcaagcuu ua 22

50 <210> 429  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

55 <400> 429  
 ccacaccgua ucugacacu u 21

<210> 430  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 430  
 ccaccaccgu gucugacacu u 21

<210> 431  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

	<400> 431 acacagggcu guugugaaga cu	22
5	<210> 432 <211> 23 <212> RNA <213> Homo sapiens	
10	<400> 432 agcuacauug ucugcugggu uuc	23
15	<210> 433 <211> 22 <212> RNA <213> Homo sapiens	
20	<400> 433 accuggcaua caauguagau uu	22
25	<210> 434 <211> 21 <212> RNA <213> Homo sapiens	
30	<400> 434 agcuacaucu ggcuacuggg u	21
35	<210> 435 <211> 22 <212> RNA <213> Homo sapiens	
40	<400> 435 cucaguagcc aguguagauc cu	22
45	<210> 436 <211> 22 <212> RNA <213> Homo sapiens	
50	<400> 436 ugucaguuuug ucaaauaccc ca	22
55	<210> 437 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 437 cguguauuug acaagcugag uu	22
	<210> 438 <211> 21 <212> RNA <213> Homo sapiens	
	<400> 438 caagucacua gugguuccgu u	21

<210> 439  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 439  
 aucacauugc cagggauuuc c 21  
  
 <210> 440  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 440  
 gggguuccug gggauuggau uu 22  
  
 <210> 441  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 441  
 aucacauugc cagggauuac c 21  
  
 <210> 442  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 442  
 uggguuccug gcaugcugau uu 22  
  
 <210> 443  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 443  
 uggcucaguu cagcaggaac ag 22  
  
 <210> 444  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 444  
 ugccuacuga gcugauauca gu 22  
  
 <210> 445  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 445  
 ugccuacuga gcugaaacac ag 22  
  
 <210> 446  
 <211> 22

<212> RNA  
 <213> Homo sapiens  
  
 <400> 446  
 5   cauugcacuu gucucggucu ga       22  
  
 <210> 447  
 <211> 21  
 <212> RNA  
 10   <213> Homo sapiens  
  
 <400> 447  
      aggcggagac uugggcaauu g       21  
  
 15   <210> 448  
      <211> 22  
      <212> RNA  
      <213> Homo sapiens  
  
 20   <400> 448  
      uucaaguaau ccaggauagg cu       22  
  
      <210> 449  
      <211> 22  
 25   <212> RNA  
      <213> Homo sapiens  
  
      <400> 449  
      ccuauucuug guuacuugca cg       22  
 30  
      <210> 450  
      <211> 22  
      <212> RNA  
      <213> Homo sapiens  
 35  
      <400> 450  
      ccuauucuug auuacuuguu uc       22  
  
      <210> 451  
      <211> 21  
      <212> RNA  
      <213> Homo sapiens  
 40  
  
      <400> 451  
      uucaaguaau ucaggauagg u       21  
 45  
  
      <210> 452  
      <211> 22  
      <212> RNA  
 50   <213> Homo sapiens  
  
      <400> 452  
      ccuguucucc auuacuuggc uc       22  
  
 55   <210> 453  
      <211> 21  
      <212> RNA  
      <213> Homo sapiens



<400> 453  
 uucacagugg cuaaguuccg c 21

5  
 <210> 454  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 454  
 agggcuuagc ugcuuugag ca 22

15  
 <210> 455  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 455  
 uucacagugg cuaaguucug c 21

25  
 <210> 456  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 456  
 agagcuuagc uguuugguga ac 22

35  
 <210> 457  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 457  
 cacuagauug ugagcuccug ga 22

45  
 <210> 458  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 458  
 aaggagcuca cagucuauug ag 22

55  
 <210> 459  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 459  
 gagguuggg uggaggcucu cc 22

<210> 460  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 460  
 agggccccc cucauuccug u 21

<210> 461  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 461  
 auguaugugu gcaugugcau g 21  
 <210> 462  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 462  
 agcagaagca gggagguucu ccca 24  
 <210> 463  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 463  
 uaugugggau gguaaacgc uu 22  
 <210> 464  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 464  
 ugguuuaccg ucccacauac au 22  
 <210> 465  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 465  
 uagcaccauc ugaaaucggu ua 22  
 <210> 466  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 466  
 acugauuucu uuugguguuc ag 22  
 <210> 467  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 467  
 uagcaccauu ugaaucagu guu 23  
 <210> 468  
 <211> 24

<212> RNA  
 <213> Homo sapiens

5 <400> 468  
 gcugguuuca uauggugguu uaga 24

10 <210> 469  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 469  
 cugguuucac augguggcuu ag 22

20 <210> 470  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

25 <400> 470  
 uagcaccauu ugaaucggu ua 22

30 <210> 471  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 471  
 ugaccgauuu cuccuggugu uc 22

40 <210> 472  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 472  
 uauacaaggg cagacucucu cu 22

50 <210> 473  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

55 <400> 473  
 cagugcaaua guauugucua age 23

<210> 474  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 474  
 cagugcaaug auauugucua age 23

<210> 475  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 475  
 uaagugcuuc cauguuuugg uga 23

5  
 <210> 476  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 476  
 acuuaaacgu ggauguacuu gcu 23

15  
 <210> 477  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 477  
 uaagugcuuc cauguuuuag uag 23

25  
 <210> 478  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 478  
 acuuuaacau ggaagugcuu uc 22

35  
 <210> 479  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 479  
 uaagugcuuc cauguuucag ugg 23

45  
 <210> 480  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 480  
 uuuuaacaugg gguuaccugc ug 22

55  
 <210> 481  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 481  
 uaagugcuuc cauguuugag ugu 23

<210> 482  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 482  
 acuuuaacau ggaggcacuu gc 22

<210> 483  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 483  
 uaagugcuuc caugcuu 17  
 <210> 484  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 484  
 uaaugcuuc cauguuu 17  
 15  
 <210> 485  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 485  
 uguaaacauc cugcacugga ag 22  
 25  
 <210> 486  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 486  
 cuuucagucg gauguuugca gc 22  
 35  
 <210> 487  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 487  
 uguaaacauc cuacacucag cu 22  
 45  
 <210> 488  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 488  
 cugggaggug gauguuacu uc 22  
 55  
 <210> 489  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 <400> 489  
 uguaaacauc cuacacucuc age 23  
 <210> 490  
 <211> 22

<212> RNA  
 <213> Homo sapiens

<400> 490  
 5 cugggagagg guuguuuacu cc 22

<210> 491  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens

<400> 491  
 cugggagaag gcuguuuacu cu 22

15 <210> 492  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 492  
 uguaaacauc cccgacugga ag 22

<210> 493  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens

<400> 493  
 30 cuuucaguca gauguuugcu gc 22

<210> 494  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35

<400> 494  
 uguaaacauc cuugacugga ag 22

<210> 495  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40

<400> 495  
 45 cuuucagucg gauguuuaca gc 22

<210> 496  
 <211> 21  
 <212> RNA  
 50 <213> Homo sapiens

<400> 496  
 55 aggcaagaug cuggcauagc u 21

<210> 497  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 497  
 ugcuaugcca acauauugcc au 22

5 <210> 498  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10 <400> 498  
 uauugcacau uacuaaguug ca 22

15 <210> 499  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 499  
 caauuuagug ugugugauau uu 22

25 <210> 500  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30 <400> 500  
 aaaagcuggg uugagagggc ga 22

35 <210> 501  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40 <400> 501  
 aaaagcuggg uugagagggc aa 22

45 <210> 502  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

50 <400> 502  
 aaaagcuggg uugagagggg u 20

55 <210> 503  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 503  
 aaaagcuggg uugagagga 19

<210> 504  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 504  
 cacauuacac ggucgaccuc u 21

<210> 505  
<211> 22  
<212> RNA  
<213> Homo sapiens  
5  
<400> 505  
aggugguccg uggcgcuuc gc 22  
10  
<210> 506  
<211> 20  
<212> RNA  
<213> Homo sapiens  
15  
<400> 506  
acugccccag gugcugcugg 20  
20  
<210> 507  
<211> 23  
<212> RNA  
<213> Homo sapiens  
25  
<400> 507  
cgcauccccu agggcauugg ugu 23  
30  
<210> 508  
<211> 23  
<212> RNA  
<213> Homo sapiens  
35  
<400> 508  
ccuaguaggu guccaguaag ugu 23  
40  
<210> 509  
<211> 20  
<212> RNA  
<213> Homo sapiens  
45  
<400> 509  
ccucugggcc cuuccuccag 20  
50  
<210> 510  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 510  
cuggcccucu cugcccuucc gu 22  
60  
<210> 511  
<211> 22  
<212> RNA  
<213> Homo sapiens  
65  
<400> 511  
aacacaccug guuaaccucu uu 22  
70  
<210> 512  
<211> 23



<212> RNA  
 <213> Homo sapiens

<400> 512  
 5 gcaaagcaca cggccugcag aga 23

<210> 513  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens

<400> 513  
 ucucugggcc ugugucuag gc 22

15 <210> 514  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20 <400> 514  
 gcccugggc cuauccuaga a 21

<210> 515  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens

<400> 515  
 30 cuagguaugg uccagggau cc 22

<210> 516  
 <211> 23  
 <212> RNA  
 35 <213> Homo sapiens

<400> 516  
 ucaagagcaa uaacgaaaaa ugu 23

40 <210> 517  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 517  
 45 uuuuucuuu uugcuccuga cc 22

<210> 518  
 <211> 22  
 <212> RNA  
 50 <213> Homo sapiens

<400> 518  
 cuccuauaug augccuuucu uc 22

55 <210> 519  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 519  
 gaacggcuuc auacaggagu u 21

5  
 <210> 520  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 520  
 uccagcauca gugauuuugu ug 22

15  
 <210> 521  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 521  
 aacaauauucc uggugcugag ug 22

25  
 <210> 522  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 522  
 ugagcgccuc gacgacagag ccg 23

35  
 <210> 523  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 523  
 ucccuguccu ccaggagcuc acg 23

45  
 <210> 524  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 524  
 gugcauugua guugcauugc a 21

55  
 <210> 525  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 525  
 caauguuucc acagugcauc ac 22

<210> 526  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 526  
 gugcauugcu guugcauugc 20

<210> 527  
<211> 22  
<212> RNA  
<213> Homo sapiens  
5  
<400> 527  
cagugccucg gcagucgagc cc 22  
10  
<210> 528  
<211> 22  
<212> RNA  
<213> Homo sapiens  
15  
<400> 528  
uuauaaagca augagacuga uu 22  
20  
<210> 529  
<211> 22  
<212> RNA  
<213> Homo sapiens  
25  
<400> 529  
uccgucucag uuacuuuaua gc 22  
30  
<210> 530  
<211> 23  
<212> RNA  
<213> Homo sapiens  
35  
<400> 530  
ucucacacag aaaucgcacc cgu 23  
40  
<210> 531  
<211> 21  
<212> RNA  
<213> Homo sapiens  
45  
<400> 531  
aggggugcua ucugugauug a 21  
50  
<210> 532  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 532  
gcugacuccu aguccagggc uc 22  
60  
<210> 533  
<211> 23  
<212> RNA  
<213> Homo sapiens  
65  
<400> 533  
ugucugcccg caugccugcc ucu 23  
70  
<210> 534  
<211> 22

<212> RNA  
<213> Homo sapiens

<400> 534  
5 uggcaguguc uuagcugguu gu 22

<210> 535  
<211> 22  
<212> RNA  
10 <213> Homo sapiens

<400> 535  
caaucagcaa guauacugcc cu 22

<210> 536  
<211> 22  
<212> RNA  
15 <213> Homo sapiens

<400> 536  
20 caaucacuaa cuccacugcc au 22

<210> 537  
<211> 23  
25 <212> RNA  
<213> Homo sapiens

<400> 537  
30 uaggcagugu cauuagcuga uug 23

<210> 538  
<211> 22  
<212> RNA  
35 <213> Homo sapiens

<400> 538  
aaucacuaac cacacggcca gg 22

<210> 539  
<211> 23  
40 <212> RNA  
<213> Homo sapiens

<400> 539  
45 aggcagugua guuagcugau ugc 23

<210> 540  
<211> 23  
<212> RNA  
50 <213> Homo sapiens

<400> 540  
ucccccaggu gugauucuga uuu 23

<210> 541  
55 <211> 22  
<212> RNA  
<213> Homo sapiens

<400> 541  
uuaucagaau cuccaggggu ac 22

5 <210> 542  
<211> 22  
<212> RNA  
<213> Homo sapiens

10 <400> 542  
aacacaccua uucaaggauu ca 22

15 <210> 543  
<211> 24  
<212> RNA  
<213> Homo sapiens

20 <400> 543  
aauccuugga accuaggugu gagu 24

25 <210> 544  
<211> 22  
<212> RNA  
<213> Homo sapiens

30 <400> 544  
aaugcacgg uauccaucug ua 22

35 <210> 545  
<211> 22  
<212> RNA  
<213> Homo sapiens

40 <400> 545  
cggguggauc acgaugcaau uu 22

45 <210> 546  
<211> 22  
<212> RNA  
<213> Homo sapiens

50 <400> 546  
uaaugcccu aaaaauccu au 22

55 <210> 547  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 547  
aaugcacuu uagcaauggu ga 22

<210> 548  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 548  
acuguucua auaugcaacu cu 22

<210> 549  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 549  
 aaauaacau gguugaucuu u 21  
 <210> 550  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 550  
 agaucgaccg uguuauauuc gc 22  
 <210> 551  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 551  
 gccugcuggg guggaaccug gu 22  
 <210> 552  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 552  
 aagugccgcc aucuuuugag ugu 23  
 <210> 553  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 553  
 acucuaacug ugggggcacu 20  
 <210> 554  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 554  
 aaagugcugc gacuuugag cgu 23  
 <210> 555  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 555  
 gaagugcuuc gauuuugggg ugu 23  
 <210> 556  
 <211> 22

<212> RNA  
 <213> Homo sapiens  
  
 <400> 556  
 5 acucaaaaug ggggcgcuuu cc 22  
  
 <210> 557  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 557  
 uuauauaca accugauaag ug 22  
  
 15 <210> 558  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 558  
 cuuaucagau uguauuguaa uu 22  
  
 <210> 559  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 559  
 auauauaca accugcuaag ug 22  
 30  
  
 <210> 560  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
  
 <400> 560  
 cuuagcaggu uguauuauca uu 22  
  
 <210> 561  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
  
 <400> 561  
 45 uuuguucguu cggcucgcgu ga 22  
  
 <210> 562  
 <211> 21  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 562  
 aucauagagg aaaauccacg u 21  
  
 55 <210> 563  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 563  
 guagauucuc cuucuaugag ua 22

5  
 <210> 564  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 564  
 aucauagagg aaaaucraig uu 22

15  
 <210> 565  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 565  
 aacauagagg aaauccacg u 21

25  
 <210> 566  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 566  
 aucacacaaa ggcaacuuuu gu 22

35  
 <210> 567  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 567  
 agagguugcc cuuggugaau uc 22

45  
 <210> 568  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 568  
 acuggacuug gagucagaag g 21

55  
 <210> 569  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 569  
 cuccugacuc cagguccugu gu 22

<210> 570  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 570  
 ugguagacua uggaacguag g 21



<210> 571  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 571  
 uauguaacau gguccacuaa cu 22  
 <210> 572  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 572  
 uauguaauau gguccacauc uu 22  
 <210> 573  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 573  
 uggugacca uagaacaugc gc 22  
 <210> 574  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 574  
 uauacaaggg caagcucucu gu 22  
 <210> 575  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 575  
 gaaguuguuc gugguggauu cg 22  
 40  
 <210> 576  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 576  
 agaucagaag gugauugugg cu 22  
 <210> 577  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 577  
 auuccuagaa auugucaua 20  
 55  
 <210> 578  
 <211> 22

<212> RNA  
 <213> Homo sapiens

<400> 578  
 5 gaauguugcu cggugaacct cu 22

<210> 579  
 <211> 23  
 <212> RNA  
 10 <213> Homo sapiens

<400> 579  
 agguuacctg agcaacuuug cau 23

15 <210> 580  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20 <400> 580  
 aauauaacac agauggccug u 21

<210> 581  
 <211> 21  
 25 <212> RNA  
 <213> Homo sapiens

<400> 581  
 uaguagacctg uauagcguac g 21  
 30

<210> 582  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35

<400> 582  
 uauguaacac gguccacuaa cc 22

<210> 583  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 40

<400> 583  
 45 acuucacctg guccacuagc cgu 23

<210> 584  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 50

<400> 584  
 aucaacagac auuaauuggg cgc 23

55 <210> 585  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 585  
 acuggacuua gggucagaag gc 22

5  
 <210> 586  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 586  
 agcucggucu gaggccccuc agu 23

15  
 <210> 587  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 587  
 ugaggggcag agagcgagac uuu 23

25  
 <210> 588  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 588  
 cagcagcaau ucauguuuug aa 22

35  
 <210> 589  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 589  
 caaaacguga ggcgcugcua u 21

45  
 <210> 590  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 590  
 aaugacacga ucacuccgu uga 23

55  
 <210> 591  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 591  
 aucgggaaug ucguguccgc cc 22

<210> 592  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 592  
 uaaucuguc ugguaaaacc gu 22

<210> 593  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 593  
 ugucuugcag gccgucaugc a 21  
 <210> 594  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 594  
 caggucgucu ugcagggcuu cu 22  
 <210> 595  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 595  
 ucuuggagua ggucauuggg ugg 23  
 <210> 596  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 596  
 cuggauggcu ccuccauguc u 21  
 <210> 597  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 597  
 aucaugaugg gcuccucggu gu 22  
 <210> 598  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 598  
 uugcauangu aggauguccc au 22  
 <210> 599  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 599  
 uggcagugua uuguuagcug gu 22  
 <210> 600  
 <211> 22

<212> RNA  
<213> Homo sapiens

<400> 600  
5 aggcagugua uuguuagcug gc 22

<210> 601  
<211> 22  
<212> RNA  
10 <213> Homo sapiens

<400> 601  
uuuugcgaug uguuccuaau au 22

15 <210> 602  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 602  
uugggaucan uuugcaucca ua 22

<210> 603  
<211> 22  
25 <212> RNA  
<213> Homo sapiens

<400> 603  
uuuugcaaua uguuccugaa ua 22

30 <210> 604  
<211> 22  
<212> RNA  
<213> Homo sapiens

35 <400> 604  
aaaccguuac cauucugag uu 22

<210> 605  
40 <211> 22  
<212> RNA  
<213> Homo sapiens

<400> 605  
45 aacuguuugc agaggaaacu ga 22

<210> 606  
<211> 22  
<212> RNA  
50 <213> Homo sapiens

<400> 606  
cucaucugca aagaaguaag ug 22

55 <210> 607  
<211> 23  
<212> RNA  
<213> Homo sapiens

<400> 607  
 agguuguccg uggugaguuc gca 23

5  
 <210> 608  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 608  
 uagugcaaua uugcuauag ggu 23

15  
 <210> 609  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 609  
 acccuaucaa uauugucucu gc 22

25  
 <210> 610  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 610  
 gcaguccaug ggcauuaca c 21

35  
 <210> 611  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 611  
 uaugugccuu uggacuacau cg 22

45  
 <210> 612  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 612  
 ucacuccucu ccuccgucu u 21

55  
 <210> 613  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 613  
 aagacgggag gaaagaagg ag 22

<210> 614  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 614  
 ucaggcucag uccccuccg au 22

<210> 615  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 615  
 gucauacacg gcucuccucu cu 22  
 <210> 616  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 616  
 agaggcuggc cgugaugaau uc 22  
 <210> 617  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 617  
 cggggcagcu caguacagga u 21  
 <210> 618  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 618  
 uccuguacug agcugccccg ag 22  
 <210> 619  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 619  
 aaucauacag ggacauccag uu 22  
 <210> 620  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 620  
 aaucguacag ggauccac uu 22  
 <210> 621  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 621  
 uugaaaggcu auuucuuggu c 21  
 <210> 622  
 <211> 21

<212> RNA  
 <213> Homo sapiens

<400> 622  
 5 cccagauaau ggcacucuca a 21

<210> 623  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens

<400> 623  
 gugacaucac auauacggca gc 22

15 <210> 624  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 624  
 caaccuggag gacuccaugc ug 22

<210> 625  
 <211> 20  
 25 <212> RNA  
 <213> Homo sapiens

<400> 625  
 30 ccauggaucu ccaggugggu 20

<210> 626  
 <211> 22  
 <212> RNA  
 35 <213> Homo sapiens

<400> 626  
 cuuaugcaag auucccuucu ac 22

<210> 627  
 <211> 22  
 <212> RNA  
 40 <213> Homo sapiens

<400> 627  
 45 agugggaac ccuuccauga gg 22

<210> 628  
 <211> 23  
 <212> RNA  
 50 <213> Homo sapiens

<400> 628  
 aggaccugcg ggacaagauu cuu 23

55 <210> 629  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens



<400> 629  
 ugaaggucua cugugugcca gg 22

5  
 <210> 630  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 630  
 uuguacaugg uaggcuuua uu 22

15  
 <210> 631  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 631  
 ugaacaauac acgggaaacc uc 22

25  
 <210> 632  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 632  
 aaacaacau ggugcacuuc uu 22

35  
 <210> 633  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 633  
 ugaguauuac auggccaau uc 22

45  
 <210> 634  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 634  
 cagcagcaca cugugguuug u 21

55  
 <210> 635  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 635  
 caaacacac ugugguguua ga 22

<210> 636  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 636  
 uuucaagcca gggggcguuu uuc 23

<210> 637  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 637  
 aacaucacag caagucugug cu 22  
 10  
 <210> 638  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 15  
 <400> 638  
 uuaagacuug cagugauguu u 21  
 20  
 <210> 639  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 639  
 uaaucuuugc uaccugggug aga 23  
 30  
 <210> 640  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 640  
 augcaccugg gcaaggauuc ug 22  
 40  
 <210> 641  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 641  
 aaugcaccgg ggcaaggauu cu 22  
 50  
 <210> 642  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 55  
 <400> 642  
 aaucuuugu cccuggguga ga 22  
 <210> 643  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 <400> 643  
 aaugcaccug ggcaaggauu ca 22  
 <210> 644  
 <211> 21

<212> RNA  
<213> Homo sapiens

<400> 644  
5 auccuugcua ucugggugcu a 21

<210> 645  
<211> 23  
<212> RNA  
10 <213> Homo sapiens

<400> 645  
uagcagcggg aacaguucug cag 23

15 <210> 646  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 646  
agaccucggu cugcacucua uc 22

<210> 647  
<211> 22  
25 <212> RNA  
<213> Homo sapiens

<400> 647  
30 cgucaacacu ugcugguuuc cu 22

<210> 648  
<211> 22  
<212> RNA  
35 <213> Homo sapiens

<400> 648  
gggagccagg aaguauugau gu 22

40 <210> 649  
<211> 21  
<212> RNA  
<213> Homo sapiens

<400> 649  
45 uaaggcacc uucugaguag a 21

<210> 650  
<211> 21  
<212> RNA  
50 <213> Homo sapiens

<400> 650  
uuuugaccu uuuggaguga a 21

55 <210> 651  
<211> 23  
<212> RNA  
<213> Homo sapiens

<400> 651  
ugauuguagc cuuuggagu aga 23

5  
<210> 652  
<211> 23  
<212> RNA  
<213> Homo sapiens

10  
<400> 652  
uacuccagag ggcgucacuc aug 23

15  
<210> 653  
<211> 22  
<212> RNA  
<213> Homo sapiens

20  
<400> 653  
uacugcagac guggcaauca ug 22

25  
<210> 654  
<211> 22  
<212> RNA  
<213> Homo sapiens

30  
<400> 654  
ugauugguac gucugugggu ag 22

35  
<210> 655  
<211> 21  
<212> RNA  
<213> Homo sapiens

40  
<400> 655  
uacugcagac aguggcaauc a 21

45  
<210> 656  
<211> 22  
<212> RNA  
<213> Homo sapiens

50  
<400> 656  
uacucaggag aguggcaauc ac 22

55  
<210> 657  
<211> 21  
<212> RNA  
<213> Homo sapiens

60  
<400> 657  
gugucuuug cucugcaguc a 21

65  
<210> 658  
<211> 22  
<212> RNA  
<213> Homo sapiens

70  
<400> 658  
aagucguc auagcugagg uc 22

<210> 659  
<211> 23  
<212> RNA  
<213> Homo sapiens  
5  
<400> 659  
cacucagccu ugagggcacu uuc 23  
10  
<210> 660  
<211> 23  
<212> RNA  
<213> Homo sapiens  
15  
<400> 660  
uaaaauucac cuuucugaga agg 23  
20  
<210> 661  
<211> 18  
<212> RNA  
<213> Homo sapiens  
<400> 661  
uucacaagga ggugucau 18  
25  
<210> 662  
<211> 22  
<212> RNA  
<213> Homo sapiens  
30  
<400> 662  
uucacaagga ggugucuuu au 22  
35  
<210> 663  
<211> 22  
<212> RNA  
<213> Homo sapiens  
<400> 663  
uucucaagga ggugucguuu au 22  
40  
<210> 664  
<211> 21  
<212> RNA  
<213> Homo sapiens  
45  
<400> 664  
auugacacuu cugugaguag a 21  
50  
<210> 665  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 665  
gagugccuuc uuuuggagcg uu 22  
<210> 666  
<211> 24

<212> RNA  
 <213> Homo sapiens  
  
 <400> 666  
 5 uucuccaaaa gaaagcacuu ucug 24  
  
 <210> 667  
 <211> 18  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 667  
 ugcuuccuuu cagaggu 18  
  
 15 <210> 668  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 668  
 uucucgagga aagaagcacu uuc 23  
  
 <210> 669  
 <211> 22  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 669  
 aucuggaggu aagaagcacu uu 22  
 30  
  
 <210> 670  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
  
 <400> 670  
 ccucuagaug gaagcacugu cu 22  
  
 <210> 671  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
  
 <400> 671  
 aucgugcauc ccuuuagagu gu 22  
 45  
  
 <210> 672  
 <211> 22  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 672  
 ucgugcaucc cuuuagagug uu 22  
  
 55 <210> 673  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 673  
 aucgugcauc cuuuuagagu gu 22

5  
 <210> 674  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 674  
 gaaagcgcuu cccuuugcug ga 22

15  
 <210> 675  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 675  
 caaagcguc cccuuuagag gu 22

25  
 <210> 676  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 676  
 caaagcgcuu cucuuuagag ugu 23

35  
 <210> 677  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 677  
 ucucuggagg gaagcacuuu cug 23

45  
 <210> 678  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 678  
 caaagcgcuu cccuuuggag c 21

55  
 <210> 679  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 679  
 cucuagaggg aagcacuuuc ug 22

<210> 680  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 680  
 aaagcgcuuc ccuucagagu g 21

<210> 681  
<211> 21  
<212> RNA  
<213> Homo sapiens  
5  
<400> 681  
gaaagcgcuu cucuuuagag g 21  
10  
<210> 682  
<211> 22  
<212> RNA  
<213> Homo sapiens  
15  
<400> 682  
cucuagaggg aagcacuuuc uc 22  
20  
<210> 683  
<211> 22  
<212> RNA  
<213> Homo sapiens  
25  
<400> 683  
aaagugcauc cuuuuagagu gu 22  
30  
<210> 684  
<211> 22  
<212> RNA  
<213> Homo sapiens  
35  
<400> 684  
cucuagaggg aagcgcuuuc ug 22  
40  
<210> 685  
<211> 22  
<212> RNA  
<213> Homo sapiens  
45  
<400> 685  
aaagugcauc cuuuuagagg uu 22  
50  
<210> 686  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 686  
aaagugcauc uuuuuagagg au 22  
60  
<210> 687  
<211> 22  
<212> RNA  
<213> Homo sapiens  
65  
<400> 687  
caaagugccu cccuuuagag ug 22  
70  
<210> 688  
<211> 22



<212> RNA  
 <213> Homo sapiens  
  
 <400> 688  
 5 aagugccucc uuuuagagug uu 22  
  
 <210> 689  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 689  
 uucuccaaaa gggagcacuu uc 22  
  
 15 <210> 690  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 690  
 aaagugcuuc ccuuuggacu gu 22  
  
 <210> 691  
 <211> 21  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 691  
 30 cuccagaggg aaguacuuuc u 21  
  
 <210> 692  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 692  
 aaagugcuuc cuuuuagagg g 21  
  
 <210> 693  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
  
 40  
 <400> 693  
 45 aaagugcuuc cuuuuagagg gu 22  
  
 <210> 694  
 <211> 22  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 694  
 aaagugcuuc ucuuuggugg gu 22  
  
 55 <210> 695  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 695  
 cuacaaaggg aagccuuuc 20

5 <210> 696  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

10 <400> 696  
 aaagugcuuc cuuuuugagg g 21

15 <210> 697  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 697  
 aagugcuucc uuuuagaggg uu 22

25 <210> 698  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

30 <400> 698  
 acaaagugcu uccuuuaga gugu 24

35 <210> 699  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40 <400> 699  
 acaaagugcu uccuuuaga gu 22

45 <210> 700  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50 <400> 700  
 aacgcacuuc ccuuuagagu gu 22

55 <210> 701  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 701  
 aaaaugguuc ccuuuagagu gu 22

<210> 702  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 702  
 gaacgcguu ccuauagag ggu 23

<210> 703  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 703  
 gaaggcgcuu cccuuuggag u 21  
 <210> 704  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 704  
 cuacaaaggg aagcacuuuc uc 22  
 15  
 <210> 705  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 705  
 gaaggcgcuu cccuuuagag cg 22  
 25  
 <210> 706  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 706  
 cuccagaggg augcacuuuc u 21  
 35  
 <210> 707  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 707  
 cucuugaggg aagcacuuuc ugu 23  
 45  
 <210> 708  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 708  
 gaaagugcuu ccuuuagag gc 22  
 55  
 <210> 709  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 <400> 709  
 cugcaaaggg aagccuuuc 20  
 <210> 710  
 <211> 22

<212> RNA  
<213> Homo sapiens

<400> 710  
5 ccucccacac ccaagguug ca 22

<210> 711  
<211> 22  
<212> RNA  
10 <213> Homo sapiens

<400> 711  
caugccuuga guguaggacc gu 22

15 <210> 712  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 712  
ggagaaaua uccuuggugu gu 22

<210> 713  
<211> 22  
25 <212> RNA  
<213> Homo sapiens

<400> 713  
uggugggcac agaaucugga cu 22

30 <210> 714  
<211> 25  
<212> RNA  
<213> Homo sapiens

35 <400> 714  
aaaggauucu gcugucgguc ccacu 25

<210> 715  
40 <211> 22  
<212> RNA  
<213> Homo sapiens

<400> 715  
45 ugugacagau ugauaacuga aa 22

<210> 716  
<211> 23  
<212> RNA  
50 <213> Homo sapiens

<400> 716  
ucggggauca ucaugucacg aga 23

55 <210> 717  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 717  
aaacauucgc ggugcacuuc uu 22

5 <210> 718  
<211> 22  
<212> RNA  
<213> Homo sapiens

10 <400> 718  
auucugcauu uuuagcaagu uc 22

15 <210> 719  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 719  
ucagcaaaca uuuauugugu gc 22

25 <210> 720  
<211> 22  
<212> RNA  
<213> Homo sapiens

30 <400> 720  
ucaguaaaug uuuuuagau ga 22

35 <210> 721  
<211> 22  
<212> RNA  
<213> Homo sapiens

40 <400> 721  
caaaacuggc aauuacuuuu gc 22

45 <210> 722  
<211> 22  
<212> RNA  
<213> Homo sapiens

50 <400> 722  
aaaaguaauu gcgaguuuua cc 22

55 <210> 723  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 723  
caagaaccuc aguugcuuuu gu 22

<210> 724  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 724  
aaaaguaauu gugguuuugg cc 22

<210> 725  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 725  
 caaaaaucuc aauuacuuuu gc 22  
 <210> 726  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 726  
 aaaaguaauu gcgguuuuug cc 22  
 15  
 <210> 727  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 727  
 caaaaaccac aguuucuuuu gc 22  
 25  
 <210> 728  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 728  
 aaaaguaauu gugguuuuug cc 22  
 35  
 <210> 729  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 729  
 aaaaacugag acuacuuuuug ca 22  
 45  
 <210> 730  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 730  
 aaaaacugua auuacuuuu 19  
 55  
 <210> 731  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 <400> 731  
 aaaacuguaa uuacuuuuugu ac 22  
 <210> 732  
 <211> 22

<212> RNA  
<213> Homo sapiens

<400> 732  
5 aaaaguaauc gcgguuuuug uc 22

<210> 733  
<211> 22  
<212> RNA  
10 <213> Homo sapiens

<400> 733  
aaaaguaauu gcggauuuug cc 22

15 <210> 734  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 734  
aaaaguaauu gcggucuuug gu 22

<210> 735  
<211> 22  
25 <212> RNA  
<213> Homo sapiens

<400> 735  
30 aaaaguacuu gcggauuuug cu 22

<210> 736  
<211> 22  
<212> RNA  
35 <213> Homo sapiens

<400> 736  
aaaaguauuu gcggguuuug uc 22

<210> 737  
40 <211> 21  
<212> RNA  
<213> Homo sapiens

<400> 737  
45 caaagguauu ugugguuuuu g 21

<210> 738  
<211> 22  
<212> RNA  
50 <213> Homo sapiens

<400> 738  
caaaaguaau uguggauuuu gu 22

55 <210> 739  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 739  
ccaaaacugc aguuacuuuu gc 22

5  
<210> 740  
<211> 22  
<212> RNA  
<213> Homo sapiens

10  
<400> 740  
uagcaaaaac ugcaguuacu uu 22

15  
<210> 741  
<211> 21  
<212> RNA  
<213> Homo sapiens

20  
<400> 741  
ugacaacuau ggaugagcuc u 21

25  
<210> 742  
<211> 23  
<212> RNA  
<213> Homo sapiens

30  
<400> 742  
agugccugag ggaguaagag ccc 23

35  
<210> 743  
<211> 22  
<212> RNA  
<213> Homo sapiens

40  
<400> 743  
ugucuuacuc ccucaggcac au 22

45  
<210> 744  
<211> 21  
<212> RNA  
<213> Homo sapiens

50  
<400> 744  
gcgaccacac cuugguuucc a 21

55  
<210> 745  
<211> 21  
<212> RNA  
<213> Homo sapiens

60  
<400> 745  
gcgaccacua cuugguuuca g 21

65  
<210> 746  
<211> 22  
<212> RNA  
<213> Homo sapiens

70  
<400> 746  
gaaaucaagc guggugaga cc 22



<210> 747  
<211> 21  
<212> RNA  
<213> Homo sapiens  
5  
<400> 747  
aacaggugac ugguuagaca a 21  
10  
<210> 748  
<211> 21  
<212> RNA  
<213> Homo sapiens  
15  
<400> 748  
aaaacgguga gauuuuguuu u 21  
20  
<210> 749  
<211> 21  
<212> RNA  
<213> Homo sapiens  
25  
<400> 749  
gcuaguccug acucagccag u 21  
30  
<210> 750  
<211> 21  
<212> RNA  
<213> Homo sapiens  
35  
<400> 750  
aggguaagcu gaaccucuga u 21  
40  
<210> 751  
<211> 22  
<212> RNA  
<213> Homo sapiens  
45  
<400> 751  
auauuaccau uagcucaucu uu 22  
50  
<210> 752  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 752  
gaugagcuca uuguauaug ag 22  
60  
<210> 753  
<211> 23  
<212> RNA  
<213> Homo sapiens  
65  
<400> 753  
guuugcacgg gugggccuug ucu 23  
70  
<210> 754  
<211> 19

<212> RNA  
<213> Homo sapiens

<400> 754  
5 ugagcugcug uaccaaaa 19

<210> 755  
<211> 21  
<212> RNA  
10 <213> Homo sapiens

<400> 755  
uaaaguaaa augcaccaa a 21

15 <210> 756  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 756  
caaaguuaa gaucuuuga gu 22

<210> 757  
<211> 20  
25 <212> RNA  
<213> Homo sapiens

<400> 757  
aaaguagcug uaccuuugc 20  
30

<210> 758  
<211> 19  
<212> RNA  
35 <213> Homo sapiens

<400> 758  
agguugacau acguuucc 19

40 <210> 759  
<211> 19  
<212> RNA  
<213> Homo sapiens

45 <400> 759  
aggcacggug ucagcaggc 19

<210> 760  
<211> 19  
<212> RNA  
50 <213> Homo sapiens

<400> 760  
ggcgccugu gauccaac 19

55 <210> 761  
<211> 23  
<212> RNA  
<213> Homo sapiens

<400> 761  
 aguauguucu uccaggacag aac 23

5  
 <210> 762  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 762  
 auguauaaau guauacacac 20

15  
 <210> 763  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 763  
 aguuuaugaa uccuggaaag u 21

25  
 <210> 764  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 764  
 cgaaaacagc aaauaccuuu gc 22

35  
 <210> 765  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 765  
 ugaguuggcc aucugaguga g 21

45  
 <210> 766  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 766  
 guccgcucgg cgguggcca 20

55  
 <210> 767  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

<400> 767  
 cugaagugau guguaacuga ucag 24

<210> 768  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 768  
 cacgcucaug cacacacca ca 22

<210> 769  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 769  
 ugagugugug ugugugagug ugu 23  
 <210> 770  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 770  
 gagccaguug gacaggagc 19  
 <210> 771  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 771  
 aagaugugga aaaauuggaa uc 22  
 <210> 772  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 772  
 auucuaauuu cuccagguu uu 22  
 <210> 773  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 773  
 uagauaaaau auugguaccu g 21  
 <210> 774  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 774  
 cuucuugugc ucuaggauug u 21  
 <210> 775  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 50  
 <400> 775  
 uucauuuggu auaaacgagc auu 23  
 <210> 776  
 <211> 22

<212> RNA  
<213> Homo sapiens

<400> 776  
5 uugagaauga ugaaucauaa gg 22

<210> 777  
<211> 21  
<212> RNA  
10 <213> Homo sapiens

<400> 777  
ucuuguguuc ucuagaucag u 21

15 <210> 778  
<211> 22  
<212> RNA  
<213> Homo sapiens

20 <400> 778  
uaacugguug aacaacugaa cc 22

<210> 779  
<211> 23  
25 <212> RNA  
<213> Homo sapiens

<400> 779  
uuacaguugu ucaaccaguu acu 23

30 <210> 780  
<211> 21  
<212> RNA  
<213> Homo sapiens

35 <400> 780  
caaagaggaa gguccauua c 21

<210> 781  
40 <211> 22  
<212> RNA  
<213> Homo sapiens

<400> 781  
45 uuaugguuug ccugggacug ag 22

<210> 782  
<211> 19  
<212> RNA  
50 <213> Homo sapiens

<400> 782  
ugggcguauc uguaugcua 19

55 <210> 783  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 783  
uaugcauugu auuuuuaggu cc 22

5  
<210> 784  
<211> 21  
<212> RNA  
<213> Homo sapiens

10  
<400> 784  
uuuccauagg ugaugaguca c 21

15  
<210> 785  
<211> 21  
<212> RNA  
<213> Homo sapiens

20  
<400> 785  
uuggccacaa ugguuagaa c 21

25  
<210> 786  
<211> 22  
<212> RNA  
<213> Homo sapiens

30  
<400> 786  
ugagaaccac gucugcucug ag 22

35  
<210> 787  
<211> 24  
<212> RNA  
<213> Homo sapiens

40  
<400> 787  
ucagaacaaa ugccguuucc caga 24

45  
<210> 788  
<211> 21  
<212> RNA  
<213> Homo sapiens

50  
<400> 788  
uaauuuuug uauaagcuag u 21

55  
<210> 789  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 789  
gagcuauuc auaaaagugc ag 22

<210> 790  
<211> 20  
<212> RNA  
<213> Homo sapiens

<400> 790  
agaccauggg uucucauugu 20

<210> 791  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 791  
 uugugucaau augcgau gau gu 22  
 <210> 792  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 792  
 ugucucugcu gggguucu 19  
 <210> 793  
 <211> 25  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 793  
 aggcaccagc caggcauugc ucagc 25  
 <210> 794  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 794  
 gaagugugcc gugguguguc u 21  
 <210> 795  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 795  
 aagccugccc ggcuccucgg g 21  
 <210> 796  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 796  
 ugugucacuc gaugaccacu gu 22  
 <210> 797  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 797  
 uacgucaucg uugucaucgu ca 22  
 <210> 798  
 <211> 20

<212> RNA  
 <213> Homo sapiens

<400> 798  
 5 guugugucag uuuaucaaac 20

<210> 799  
 <211> 23  
 <212> RNA  
 10 <213> Homo sapiens

<400> 799  
 acuuacagac aagagccuug cuc 23

<210> 800  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 800  
 20 uggucuagga uuguuggagg ag 22

<210> 801  
 <211> 23  
 <212> RNA  
 25 <213> Homo sapiens

<400> 801  
 gacacgggcg acagcugcgg ccc 23

<210> 802  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 802  
 35 cacacacugc aauuacuuuu gc 22

<210> 803  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 803  
 45 aggcugcgga auucaggac 19

<210> 804  
 <211> 23  
 <212> RNA  
 50 <213> Homo sapiens

<400> 804  
 uaaaucccau ggugccuucu ccu 23

<210> 805  
 <211> 21  
 <212> RNA  
 55 <213> Homo sapiens



<400> 805  
aaacuacuga aaaucaaaga u 21

5  
<210> 806  
<211> 21  
<212> RNA  
<213> Homo sapiens

10  
<400> 806  
guucaaaucc agaucuauaa c 21

15  
<210> 807  
<211> 25  
<212> RNA  
<213> Homo sapiens

20  
<400> 807  
agggguggug uugggacagc uccgu 25

25  
<210> 808  
<211> 20  
<212> RNA  
<213> Homo sapiens

30  
<400> 808  
aggguguuuc ucucaucucu 20

35  
<210> 809  
<211> 21  
<212> RNA  
<213> Homo sapiens

40  
<400> 809  
ugagcuaaaau gugugcuggg a 21

45  
<210> 810  
<211> 23  
<212> RNA  
<213> Homo sapiens

50  
<400> 810  
gcgaggaccc cucggggucu gac 23

55  
<210> 811  
<211> 25  
<212> RNA  
<213> Homo sapiens

<400> 811  
gcugggcagg gcuucugagc uccuu 25

<210> 812  
<211> 20  
<212> RNA  
<213> Homo sapiens

<400> 812  
aggaauguuc cuucuuugcc 20

<210> 813  
<211> 23  
<212> RNA  
<213> Homo sapiens  
5  
<400> 813  
gaacgccugu ucuugccagg ugg 23  
10  
<210> 814  
<211> 22  
<212> RNA  
<213> Homo sapiens  
15  
<400> 814  
uccgagccug ggucuccuc uu 22  
20  
<210> 815  
<211> 22  
<212> RNA  
<213> Homo sapiens  
25  
<400> 815  
ggggguccccc ggugcucgga uc 22  
30  
<210> 816  
<211> 22  
<212> RNA  
<213> Homo sapiens  
35  
<400> 816  
agucauugga ggguuugagc ag 22  
40  
<210> 817  
<211> 22  
<212> RNA  
<213> Homo sapiens  
45  
<400> 817  
acucaaaacc cuucagugac uu 22  
50  
<210> 818  
<211> 22  
<212> RNA  
<213> Homo sapiens  
55  
<400> 818  
agacuuccca uuugaaggug gc 22  
60  
<210> 819  
<211> 23  
<212> RNA  
<213> Homo sapiens  
65  
<400> 819  
aaacucuacu uguccuucug agu 23  
70  
<210> 820  
<211> 24

<212> RNA  
<213> Homo sapiens

<400> 820  
5 gaccuggaca uguuugugcc cagu 24

<210> 821  
<211> 20  
<212> RNA  
10 <213> Homo sapiens

<400> 821  
auggagauag auauagaaau 20

15 <210> 822  
<211> 21  
<212> RNA  
<213> Homo sapiens

20 <400> 822  
ggcuagcaac agcgcuuacc u 21

<210> 823  
<211> 21  
25 <212> RNA  
<213> Homo sapiens

<400> 823  
30 acagucugcu gagguuggag c 21

<210> 824  
<211> 23  
<212> RNA  
35 <213> Homo sapiens

<400> 824  
aucccuugca ggggcuguug ggu 23

<210> 825  
40 <211> 21  
<212> RNA  
<213> Homo sapiens

<400> 825  
45 cacaagguau uggauuacc u 21

<210> 826  
<211> 22  
<212> RNA  
50 <213> Homo sapiens

<400> 826  
uaguaccagu accuuguguu ca 22

55 <210> 827  
<211> 21  
<212> RNA  
<213> Homo sapiens

<400> 827  
aggggaaag uucuauaguc c 21

5  
<210> 828  
<211> 22  
<212> RNA  
<213> Homo sapiens

10  
<400> 828  
gacuauagaa cuuuccccu ca 22

15  
<210> 829  
<211> 19  
<212> RNA  
<213> Homo sapiens

20  
<400> 829  
agcugucuga aaaugucuu 19

25  
<210> 830  
<211> 22  
<212> RNA  
<213> Homo sapiens

30  
<400> 830  
gugagucucu aagaaaagag ga 22

35  
<210> 831  
<211> 21  
<212> RNA  
<213> Homo sapiens

40  
<400> 831  
ucuaguaaga guggcagucg a 21

45  
<210> 832  
<211> 22  
<212> RNA  
<213> Homo sapiens

50  
<400> 832  
augcugacau auuuacuaga gg 22

55  
<210> 833  
<211> 21  
<212> RNA  
<213> Homo sapiens

<400> 833  
uggguuuacg uugggagaac u 21

<210> 834  
<211> 22  
<212> RNA  
<213> Homo sapiens

<400> 834  
guucucccaa cguaagccca gc 22

<210> 835  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 835  
 aguauucugu accaggaag gu 22  
 10  
 <210> 836  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 15  
 <400> 836  
 agaccuggcc cagaccucag c 21  
 20  
 <210> 837  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 837  
 gugucugcuu ccugugga 19  
 30  
 <210> 838  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 838  
 cuaauaguau cuaccacaau aaa 23  
 40  
 <210> 839  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 839  
 aaccagcacc ccaacuuugg ac 22  
 50  
 <210> 840  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 55  
 <400> 840  
 acuuggcac ugaacaaug ucc 23  
 60  
 <210> 841  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 65  
 <400> 841  
 ugugcuugcu cgucccgccc gca 23  
 70  
 <210> 842  
 <211> 24

<212> RNA  
 <213> Homo sapiens

5 <400> 842  
 acugggggcu uucgggcucu gcgu 24

10 <210> 843  
 <211> 25  
 <212> RNA  
 <213> Homo sapiens

15 <400> 843  
 agggaucgcg ggcggguggc ggccu 25

20 <210> 844  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

25 <400> 844  
 aucgcugcgg uugcgagcgc ugu 23

30 <210> 845  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

35 <400> 845  
 augauccagg aaccugccuc u 21

40 <210> 846  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

45 <400> 846  
 aaagacauag gauagaguca ccuc 24

50 <210> 847  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

55 <400> 847  
 guccucucc aaaugugucu ug 22

<210> 848  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 848  
 acuuguaugc uagcucaggu ag 22

<210> 849  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

<400> 849  
 aguguggcuu ucuuagagc 19

5  
 <210> 850  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 850  
 ucuaggcugg uacugcuga 19

15  
 <210> 851  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 851  
 aagcagcugc cucugaggc 19

25  
 <210> 852  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 852  
 guggcugcac ucacuuccu c 21

35  
 <210> 853  
 <211> 19  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 853  
 aagugugcag ggcacuggu 19

45  
 <210> 854  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 854  
 aaaccugugu uguucaagag uc 22

55  
 <210> 855  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 855  
 aggaggcagc gcucucagga c 21

<210> 856  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 856  
 uuuaggauaa gcuugacuuu ug 22

<210> 857  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 857  
 aauggcgcca cuagguugu g 21  
 <210> 858  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 858  
 guguugaaac aaucucuacu g 21  
 <210> 859  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 859  
 uaugucugcu gaccaucacc uu 22  
 <210> 860  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 860  
 uggugggccg cagaacaugu gc 22  
 <210> 861  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 861  
 auaauacaug guuaaccucu uu 22  
 <210> 862  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 862  
 aaauuuauac agucaaccuc u 21  
 <210> 863  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 863  
 ggcagguucu caccucucu agg 23  
 <210> 864  
 <211> 25



<212> RNA  
 <213> Homo sapiens

5 <400> 864  
 gccggaggga aguagguccg uuggu 25

<210> 865  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens

<400> 865  
 cuugguucag ggaggguccc ca 22

15 <210> 866  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20 <400> 866  
 uacccauugc auaucggagu ug 22

25 <210> 867  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

30 <400> 867  
 ugccuggguc ucuggccugc gcgu 24

35 <210> 868  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

<400> 868  
 ucccacguug uggcccagca g 21

40 <210> 869  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

45 <400> 869  
 aggcggggcg cgcgggacc gc 22

50 <210> 870  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 870  
 gguggcccgg ccgugccuga gg 22

55 <210> 871  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 871  
 uauucauuua ucccagccu aca 23

5  
 <210> 872  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 872  
 acuggcuagg gaaaugauu ggau 24

15  
 <210> 873  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 873  
 accaggaggc ugaggcccu 20

25  
 <210> 874  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 874  
 ugucacucgg cucggccac uac 23

35  
 <210> 875  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 875  
 uccgguucuc agggcuccac c 21

45  
 <210> 876  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 876  
 aggaagcccu ggaggggcug gag 23

55  
 <210> 877  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 877  
 uggugcgag agggcccaca gug 23

<210> 878  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

<400> 878  
 cugaugccc ucaccgcuca 20

<210> 879  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 879  
 uggaagacua gugauuuugu ugu 23  
 <210> 880  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 880  
 caacaaaucagucugcca ua 22  
 <210> 881  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 881  
 caacaaauccagucuaccu aa 22  
 <210> 882  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 882  
 aaggagcuua caaucuagcu ggg 23  
 <210> 883  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 883  
 caacuagacu gugagcuucu ag 22  
 <210> 884  
 <211> 17  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 884  
 ucucgcuggg gccucca 17  
 <210> 885  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 45  
 <400> 885  
 ugcggggcua gggcuaacag ca 22  
 <210> 886  
 <211> 22

<212> RNA  
 <213> Homo sapiens  
  
 <400> 886  
 5 cuguugccac uaaccucaac cu 22  
  
 <210> 887  
 <211> 22  
 <212> RNA  
 10 <213> Homo sapiens  
  
 <400> 887  
 uuugugaccu gguccacuaa cc 22  
  
 15 <210> 888  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
  
 20 <400> 888  
 cggcucuggg ucugugggga 20  
  
 <210> 889  
 <211> 21  
 25 <212> RNA  
 <213> Homo sapiens  
  
 <400> 889  
 30 uggaggagaa ggaaggugau g 21  
  
 <210> 890  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 890  
 acuccagccc cacagccuca gc 22  
  
 <210> 891  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
  
 <400> 891  
 45 ucugcucuaa cccaugguu ucu 23  
  
 <210> 892  
 <211> 23  
 <212> RNA  
 50 <213> Homo sapiens  
  
 <400> 892  
 ugaccaugg uugucugagc aug 23  
  
 55 <210> 893  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

	<400> 893 cugggaucuc cggggucuug guu	23
5	<210> 894 <211> 22 <212> RNA <213> Homo sapiens	
10	<400> 894 ugagaccucu gguucugag cu	22
15	<210> 895 <211> 23 <212> RNA <213> Homo sapiens	
20	<400> 895 uccaguacca cgugucaggg cca	23
25	<210> 896 <211> 23 <212> RNA <213> Homo sapiens	
30	<400> 896 caguaacaaa gauucauccu ugu	23
35	<210> 897 <211> 21 <212> RNA <213> Homo sapiens	
40	<400> 897 gcaggaacuu gugagucucc u	21
45	<210> 898 <211> 22 <212> RNA <213> Homo sapiens	
50	<400> 898 cugcccuggc ccgagggacc ga	22
55	<210> 899 <211> 21 <212> RNA <213> Homo sapiens	
	<400> 899 ccuggaaaca cugagguugu g	21
	<210> 900 <211> 22 <212> RNA <213> Homo sapiens	
	<400> 900 uauaccucag uuuuaucaagg ug	22

<210> 901  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 901  
 uggugguuua caaaguaauu ca 22  
 <210> 902  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 902  
 uggauuucuu ugugaaucac ca 22  
 <210> 903  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 903  
 guagaggaga uggcgcaggg 20  
 <210> 904  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 904  
 uccucuucuc ccuccuccca g 21  
 <210> 905  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 905  
 aggcagcggg guguagugga ua 22  
 <210> 906  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 906  
 uccauuacac uaccugccu cu 22  
 <210> 907  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 907  
 cgcgggugcu uacugacccu u 21  
 <210> 908  
 <211> 23

<212> RNA  
 <213> Homo sapiens

5 <400> 908  
 cgggucggag uuagcucaag cgg 23

10 <210> 909  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

15 <400> 909  
 gugaacgggc gccaucccga gg 22

20 <210> 910  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

25 <400> 910  
 uacucaaaaa gcugucaguc a 21

30 <210> 911  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

35 <400> 911  
 gacugacacc ucuuugggug aa 22

40 <210> 912  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

45 <400> 912  
 uuaauaucgg acaaccuug u 21

50 <210> 913  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

55 <400> 913  
 uacuuggaaa ggcaucaguu g 21

<210> 914  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 914  
 ugcaacgaac cugagccacu ga 22

<210> 915  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 915  
 ugcaacuuac cugagucauu ga 22

5  
 <210> 916  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 916  
 cacugugucc uuucugcgua g 21

15  
 <210> 917  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 917  
 cacuggcucc uuucugggua ga 22

25  
 <210> 918  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 918  
 ucuuugguuu ucuagcugua uga 23

35  
 <210> 919  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 919  
 auaaagcuag auaaccgaaa gu 22

45  
 <210> 920  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 920  
 ggggagcugu ggaagcagua 20

55  
 <210> 921  
 <211> 25  
 <212> RNA  
 <213> Homo sapiens

<400> 921  
 cuagugaggg acagaaccag gauuc 25

<210> 922  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

<400> 922  
 gcagcagaga auaggacuac guc 23



<210> 923  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 923  
 gucagcggag gaaaagaaac u 21  
 <210> 924  
 <211> 20  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 924  
 agagucuugu gaugucuugc 20  
 <210> 925  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 925  
 uauugcacuu gucccggccu gu 22  
 <210> 926  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 926  
 agguugggau cgguugcaau gcu 23  
 <210> 927  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 927  
 ggguggggau uuguugcauu ac 22  
 <210> 928  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 928  
 uauugcacuc gucccggccu cc 22  
 <210> 929  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 929  
 agggacggga cgcgugcag ug 22  
 <210> 930  
 <211> 23

<212> RNA  
 <213> Homo sapiens

5      <400> 930  
       caaagugcug uucgucagg uag      23

10     <210> 931  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

15     <400> 931  
       acugcugagc uagcacuucc cg      22

20     <210> 932  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

25     <400> 932  
       ugugcgagg gagaccucuc cc      22

30     <210> 933  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

35     <400> 933  
       ugucuacuac uggagacacu gg      22

40     <210> 934  
       <211> 23  
       <212> RNA  
       <213> Homo sapiens

45     <400> 934  
       ccaguuaccg cuuccgcuac cgc      23

50     <210> 935  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

55     <400> 935  
       acaguagagg gaggaaucgc ag      22

60     <210> 936  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

65     <400> 936  
       auccg'gcuc ugacucucug cc      22

70     <210> 937  
       <211> 22  
       <212> RNA  
       <213> Homo sapiens

<400> 937  
 ugcccuuaaa ggugaaccca gu 22

5  
 <210> 938  
 <211> 24  
 <212> RNA  
 <213> Homo sapiens

10  
 <400> 938  
 uggggagcug aggcucuggg ggug 24

15  
 <210> 939  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

20  
 <400> 939  
 aaggcagggc cccgcuccc c 21

25  
 <210> 940  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens

30  
 <400> 940  
 caccggcug ugucacaug ugc 23

35  
 <210> 941  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

40  
 <400> 941  
 ucuucucugu uuuggccaug ug 22

45  
 <210> 942  
 <211> 21  
 <212> RNA  
 <213> Homo sapiens

50  
 <400> 942  
 cugacuguug ccguccuca g 21

55  
 <210> 943  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 943  
 aaauuauugu acaucggaug ag 22

<210> 944  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens

<400> 944  
 uucaacgggu auuuauugag ca 22

<210> 945  
 <211> 23  
 <212> RNA  
 <213> Homo sapiens  
 5  
 <400> 945  
 uuuggcacua gcacauuuu gcu 23

<210> 946  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 10  
 <400> 946  
 aaucaugugc agugccaaua ug 22

<210> 947  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 20  
 <400> 947  
 ugagguagua aguuguauug uu 22

<210> 948  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 25  
 <400> 948  
 aaccguaga uccgaucuug ug 22

<210> 949  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 30  
 <400> 949  
 caagcucgcu ucuauuggguc ug 22

<210> 950  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 35  
 <400> 950  
 caccguaga accgaccuug cg 22

<210> 951  
 <211> 22  
 <212> RNA  
 <213> Homo sapiens  
 40  
 <400> 951  
 caagcucgug ucuguggguc cg 22

<210> 952  
 <211> 21

<212> RNA  
 <213> Herpes simplex virus 1

5 <400> 952  
 uggaaggacg ggaaguggaa g 21

10 <210> 953  
 <211> 24  
 <212> RNA  
 <213> Herpes simplex virus 1

15 <400> 953  
 ccugagccag ggacgagugc gacu 24

20 <210> 954  
 <211> 22  
 <212> RNA  
 <213> Herpes simplex virus 1

25 <400> 954  
 ugcacgcgc ccggcacaga cu 22

30 <210> 955  
 <211> 20  
 <212> RNA  
 <213> Herpes simplex virus 1

35 <400> 955  
 cugggacugu gcgguuggga 20

40 <210> 956  
 <211> 22  
 <212> RNA  
 <213> Herpes simplex virus 1

45 <400> 956  
 cuugccuguc uaacucgcu gu 22

50 <210> 957  
 <211> 22  
 <212> RNA  
 <213> Herpes simplex virus 1

55 <400> 957  
 gguagaguuu gacaggcaag ca 22

<210> 958  
 <211> 22  
 <212> RNA  
 <213> Herpes simplex virus 1

<400> 958  
 gucagagauc caaacccucc gg 22

<210> 959  
 <211> 21  
 <212> RNA  
 <213> Herpes simplex virus 1

<400> 959  
 cacuucccggu ccuuccaucc c 21

5  
 <210> 960  
 <211> 23  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

10  
 <400> 960  
 auuacaggaa acugggugua agc 23

15  
 <210> 961  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

20  
 <400> 961  
 uaguguuguc cccccgagug gc 22

25  
 <210> 962  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

30  
 <400> 962  
 ugguguuguc cccccgagug gc 22

35  
 <210> 963  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

40  
 <400> 963  
 uuaaugcuua gccugugucc ga 22

45  
 <210> 964  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

50  
 <400> 964  
 accaggccac cauuccucuc cg 22

55  
 <210> 965  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

<400> 965  
 aacuguaguc cgggucgauc ug 22

<210> 966  
 <211> 23  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

<400> 966  
 ucacauucug aggacggcag cga 23

<210> 967  
 <211> 21  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 5  
 <400> 967  
 ucgcgguac agaaugugac a 21  
 <210> 968  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 10  
 <400> 968  
 uagaauacug aggccuagcu ga 22  
 <210> 969  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 20  
 <400> 969  
 agcuaaacg caguacucua gg 22  
 <210> 970  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 25  
 <400> 970  
 uaggaugccu ggaacuugcc gg 22  
 <210> 971  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 30  
 <400> 971  
 ugauguuuu cgggcuguug ag 22  
 <210> 972  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 35  
 <400> 972  
 ccagcagcac cuaaucauc gg 22  
 <210> 973  
 <211> 21  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus  
 40  
 <400> 973  
 ugaucoccaug uugcuggcgc u 21  
 <210> 974  
 <211> 22

<212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

5 <400> 974  
 uaggcgcgac ugagagagca cg 22

<210> 975  
 <211> 22  
 <212> RNA  
 10 <213> Kaposi's sarcoma-associated herpesvirus

<400> 975  
 cuggguauac gcagcugcgu aa 22

15 <210> 976  
 <211> 22  
 <212> RNA  
 <213> Kaposi's sarcoma-associated herpesvirus

20 <400> 976  
 acccagcugc guaaaccccg cu 22

25 <210> 977  
 <211> 14  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> LNA Oligomer

<220>  
 <221> misc-feature  
 <222> (1)..(14)  
 <223> Full LNA & Phosphorothioate linkages,  
 35

<400> 977  
 cacgattagc atta 14

40 <210> 978  
 <211> 6  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <223> LNA Oligomer

<220>  
 <221> misc-feature  
 <222> (1)..(6)  
 50 <223> Full LNA & Phosphorothioate linkages,

<400> 978  
 gcatta 6

55 <210> 979  
 <211> 7  
 <212> DNA  
 <213> Artificial Sequence



<220>  
<223> LNA Oligomer

5 <220>  
<221> misc-feature  
<222> (1)..(7)  
<223> Full LNA & Phosphorothioate linkages,

10 <400> 979  
agcatta 7

<210> 980  
<211> 10  
<212> DNA  
15 <213> Artificial Sequence

<220>  
<223> LNA Oligomer

20 <220>  
<221> misc-feature  
<222> (1)..(10)  
<223> Full LNA & Phosphorothioate linkages,

25 <400> 980  
attagcatta 10

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

## Patent documents cited in the description

- WO 2007112754 A [0003] [0202]
- WO 2007112753 A [0003]
- WO 2008046911 A [0003] [0043]
- WO 2005098029 A [0018]
- WO 2006069584 A [0018]
- WO 9914226 A [0080]
- WO 0056746 A [0080]
- WO 0056748 A [0080]
- WO 0125248 A [0080]
- WO 0228875 A [0080]
- WO 03006475 A [0080]
- WO 03095467 A [0080]
- US 60977497 B [0123] [0411]
- WO 2007031091 A [0138]
- WO 2008034123 A [0143]
- US 7087229 B [0145]
- WO 2008034122 A [0150]
- US 4962029 A [0151] [0152]
- US 4914210 A [0151] [0152]
- US 60838710 B [0165]
- US 60788995 B [0165]
- DK 200600615 [0165]
- WO 20071112754 A [0202]
- US 60979217 B [0411]
- US 61028062 B [0411]
- EP 08104780 A [0411]

## Non-patent literature cited in the description

- THEODORA W GREENE ; PETER G M WUTS. Protective Groups in Organic Synthesis. John Wiley & Sons, 1999 [0145]
- MANOHARAN et al. *Tetrahedron Letters*, 1991, vol. 34, 7171 [0154]
- CROOKE, R. M. Antisense research and Application. Springer-Verlag, 1998, vol. 131, 103-140 [0163]
- DASS CR. *J Pharm Pharmacol*, 2002, vol. 54 (1), 3-27 [0166]
- DR. B. JANSEN. Section of experimental Oncology, Molecular Pharmacology. Department of Clinical Pharmacology, University of Vienna [0206]
- GRIFFITHS-JONES, S. ; GROCOCK, R.J. ; VAN DONGEN, S. ; BATEMAN, A. ; ENRIGHT, A.J. miR-Base: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.*, 2006, vol. 34, D140-4 [0207]
- ASANGANI et al. *Oncogene*, 2007 [0308]
- FRANKEL et al. *J Biol Chem*, 2008 [0308]
- R. BARTINSCHLAGER. Dept Mol Virology. University of Heidelberg [0309] [0341] [0347]
- MAYR et al. *Science*, 2007 [0339]
- LEE ; DUTTA. *Genes Dev*, 2007 [0339]
- JOHNSON et al. *Cell*, 2005 [0388]
- JOHNSON et al. *Cancer Research*, 2007 [0389]

## Patentkrav

1. Oligomer af 7-12 nukleobasers længde til anvendelse til reducere af den effektive mængde af et mikroRNA-mål i en  
5 celle; hvor alle nukleobaseenhederne i den sammenhængende nukleobasesekvens er LNA-nukleobaseenheder, hvor den sammenhængende nukleobasesekvens er komplementær til den tilsvarende sekvens af mindst to miRNA-sekvenser, og hvor mindst 75 % af internukleosidkoblingerne, der er til stede  
10 mellem nukleobaseenhederne i den sammenhængende nukleobasesekvens, er phosphorothioat-internukleosidkoblinger.
2. Oligomer af 7-12 nukleobasers længde til anvendelse til reducere af den effektive mængde af et mikroRNA-mål i en  
15 celle; hvor alle nukleobaseenhederne i den sammenhængende nukleobasesekvens er LNA-nukleobaseenheder, hvor den sammenhængende nukleobasesekvens er komplementær til den tilsvarende sekvens af mindst to miRNA-sekvenser, og hvor mindst 75 % af internukleosidkoblingerne, der er til stede  
20 mellem nukleobaseenhederne i den sammenhængende nukleobasesekvens, er phosphorothioat-internukleosidkoblinger, til anvendelse som et medikament.
3. Oligomer eller anvendelse ifølge krav 1 eller 2, hvor  
25 oligomeren er til reducere af mængden af mindst to mikroRNA-mål.
4. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 3, hvor den sammenhængende nukleobasesekvens er  
30 komplementær til en mammalia-afledt, human eller viral mikroRNA (miRNA)-sekvens.
5. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 4, hvor den sammenhængende nukleobasesekvens er  
35 100 % komplementær til den mammalia-afledte, humane eller virale mikroRNA (miRNA)-sekvens.
6. Oligomer eller anvendelse ifølge et hvilket som helst af

kravene 1 til 5, hvor den sammenhængende nukleobasesekvens består af eller omfatter en sekvens, der er 100 % komplementær til den mammalia-afledte, humane eller virale mikroRNA (miRNA)-podeseqvens.

5

7. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 6, hvor den sammenhængende nukleobasesekvens består af eller omfatter en sekvens, der er komplementær til sekvensen af mindst to miRNA-poderegionsekvenser.

10

8. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor den sammenhængende nukleobasesekvens består af eller omfatter en sekvens, der er 100 % komplementær til sekvensen af mindst to miRNA-poderegionsekvenser.

15

9. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor sekvensen, der er komplementær til podeseqvensen af de mindst to mikroRNA-sekvenser, er mindst 6 eller mindst 7 nukleobaser lang og er 100 % komplementær til podeseqvensen af det mindst to mikroRNA-sekvenser.

20

10. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 9, hvor alle internukleosidkoblingerne, der er til stede mellem nukleobaseenhederne i den sammenhængende nukleobasesekvens, er phosphorothioat-internukleosidkoblinger.

25

11. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 10, hvor alle de bicykliske nukleosidanalogenheder har formelen:




30

hvor X er vagt fra gruppen, der består af O, S og  $\text{NR}^{\text{H}}$ , hvor  $\text{R}^{\text{H}}$  er H eller  $\text{C}_{1-4}$ -alkyl; Y er  $(-\text{CH}_2)_r$ , hvor r er et helt tal på 1-2; og B er en nitrogenholdig base.

35

12. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 10, hvor alle de bicykliske

nukleosidanalogenheder har formelen:



hvor B er en nitrogenholdig base.

- 5 13. Oligomer eller anvendelse ifølge et hvilket som helst af kravene 1 til 12, hvor oligomeren er 7 sammenhængende nukleobaser lang.
14. Oligomer eller anvendelse ifølge et hvilket som helst af  
10 kravene 1 til 12, hvor oligomeren er 8 sammenhængende nukleobaser lang.
15. Oligomer eller anvendelse ifølge et hvilket som helst af  
15 kravene 1 til 12, hvor oligomeren er 9 sammenhængende nukleobaser lang.
16. Oligomer eller anvendelse ifølge et hvilket som helst af  
kravene 1 til 12, hvor oligomeren er 10 sammenhængende nukleobaser lang.  
20
17. Konjugat, der omfatter oligomeren ifølge et hvilket som helst af kravene 1 til 16 og mindst én ikke-nukleotiddel, der er kovalent bundet dertil.
- 25 18. Oligomer ifølge et hvilket som helst af kravene 1 til 16 eller konjugat ifølge krav 17 til anvendelse som et medikament.
19. Farmaceutisk sammensætning, der omfatter oligomeren  
30 ifølge et hvilket som helst af kravene 1 til 16 eller konjugatet ifølge krav 17 og et farmaceutisk acceptabelt fortyndingsmiddel, bæremiddel, salt eller adjuvans.
20. *In vitro*-fremgangsmåde til reducere af den effektive  
35 mængde af et mikroRNA-mål i en celle, som omfatter administrering af en oligomer ifølge et hvilket som helst af

kravene 1 til 16 eller et konjugat ifølge krav 17 til cellen eller organismen.

21. *In vitro*-fremgangsmåde til derepression af et mål-mRNA i  
5 en celle, som omfatter administrering af en oligomer ifølge et hvilket som helst af kravene 1 til 16 eller et konjugat ifølge krav 17 til cellen eller organismen.

22. Fremgangsmåde ifølge krav 20 eller 21, hvor  
10 fremgangsmåden er til reducere af mængden af mindst to mikroRNA-mål i cellen.

# DRAWINGS




Figure 1




Figure 2




Figure 3




Figure 4






Figure 5






Figure 6



**Figure 7**



**Figure 8**




Figure 9




Figure 10






Figure 13



Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20








Figure 21



SEQ ID #3205      5'- GATAAGCT - 3'

SEQ ID #3219      5'- CGTAATGA - 3'

SEQ ID #3218      5'- GGTAAACT - 3'

Figure 22




Figure 23





Figure 24



SEQ ID # 3225      5'- ATGTAGC -3'  
 LNA control      5'- GTAGACT -3'

Figure 25




Figure 26

Name	Mature miRNA sequence	S (2-8)	ES (2-9)	NE (9-16)	Total (2-16)
let-7a	UGAGGUAGUAGGUUGUAUAGUU	N/A	N/A	N/A	N/A
let-7b	UGAGGUAGUAGGUUGUAUAGUU	0	0	0	0
let-7c	UGAGGUAGUAGGUUGUAUAGUU	0	0	0	0
let-7d	AGAGGUAGUAGGUUGUAUAGUU	0	0	1	1
let-7e	UGAGGUAGUAGGUUGUAUAGUU	0	1	0	1
let-7f	UGAGGUAGUAGGUUGUAUAGUU	0	0	1	1
let-7g	UGAGGUAGUAGGUUGUAUAGUU	0	0	1	1
let-7i	UGAGGUAGUAGGUUGUAUAGUU	0	0	1	1
miR-98	UGAGGUAGUAGGUUGUAUAGUU	0	0	1	1

Number	Compound	Sequence (5' to 3') <sup>a</sup>	Length (nt)	Complementary target(s)
1.	SEQ ID #3226	A-C-a-A-c-C-T-a-c-T-a- C-c-T-C	15	let-7a/b/c
2.	SEQ ID #3236	G-C-a-A-c-C-T-a-c-T-a- C-c-T-C	15	let-7d
3.	SEQ ID #3237	A-C-a-A-c-C-T-c-c-T-a- C-c-T-C	15	let-7e
4.	SEQ ID #3238	A-C-a-A-a-C-T-a-c-T-a- C-c-T-C	15	let-7g/i
5.	SEQ ID #3239	C-T-A-C- C-T-C	7	all members
6.	SEQ ID #3240	C-T-A-A- C-T-C	7	none
7.	SEQ ID #3227	A-C-T-A-C- C-T-C	8	all except let-7e
8.	SEQ ID #3232	T-A-C- C-T-C	6	all members
9.	SEQ ID #3234	T-N <sub>1</sub> -C-T-A-C- C-T-C	9	all members <sup>b</sup>
10.	SEQ ID #3235	T-N <sub>2</sub> -C-T-A-C- C-T-C	9	all members <sup>b</sup>

<sup>a</sup>Capital and lower case letters denote LNA and DNA, respectively.

<sup>b</sup>Both 9-mers theoretically target all members since they contain 2 different universally hybridizing chemistries at their 2<sup>nd</sup> position.

Figure 27




Figure 28




Figure 29




Figure 30





Figure 31.




Figure 32.






Figure 33.





(A)



(B)

Figure 34 A and B







Figure 35.




A)




B)

Figure 36.


**A**



**B**



**A**



**B**






Figure 38.


**A**



**B**



**A**



**B**




Figure 37.