


(72) STEINER, GERD, DE
(72) HOGER, THOMAS, DE
(72) UNGER, LILIANE, DE
(72) TESCHENDORF, HANS-JURGEN, DE
(72) JUCHELKA, FRIEDER, DE
(71) BASF AKTIENGESELLSCHAFT, DE
(51) Int.Cl.⁷ C07D 403/06, A61K 31/4184
(30) 1998/08/12 (198 36 404.0) DE

(54) DERIVES A SUBSTITUTION N D'AZABICYCLOHEPTANE,
PREPARATION ET UTILISATION DESDITS DERIVES
(54) N-SUBSTITUTED AZABICYCLOHEPTANE DERIVATIVES,
PRODUCTION AND USE THEREOF



(57) The invention relates to compounds of formula (1), wherein R¹, R² and R³ have the meaning cited in the description. The novel substances are suitable for use in disease control.




PCT WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales Büro
INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)

<p>(51) Internationale Patentklassifikation ⁷ : C07D 403/06, A61K 31/4184</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 00/09499 (43) Internationales Veröffentlichungsdatum: 24. Februar 2000 (24.02.00)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP99/05166 (22) Internationales Anmeldedatum: 20. Juli 1999 (20.07.99) (30) Prioritätsdaten: 198 36 404.0 12. August 1998 (12.08.98) DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): STEINER, Gerd [DE/DE]; Oberer Waldweg 1, D-67281 Kirchheim (DE). HÖGER, Thomas [DE/DE]; Rathenaustrasse 12, D-68535 Edingen-Neckarhausen (DE). UNGER, Liliane [DE/DE]; Wollstrasse 129, D-67065 Ludwigshafen (DE). TESCHENDORF, Hans-Jürgen [DE/DE]; Georg-Nuss-Strasse 5, D-67373 Dudenhofen (DE). JUCHELKA, Frieder [DE/DE]; Tuerkisweg 22, D-69181 Leimen (DE). (74) Gemeinsamer Vertreter: BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE).</p>	<p>(81) Bestimmungsstaaten: AL, AU, BG, BR, BY, CA, CN, CZ, GE, HR, HU, ID, IL, IN, JP, KR, KZ, LT, LV, MK, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, ZA, eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Veröffentlicht Mit internationalem Recherchenbericht.</p>	

(54) Title: N-SUBSTITUTED AZABICYCLOHEPTANE DERIVATIVES, PRODUCTION AND USE THEREOF

(54) Bezeichnung: N-SUBSTITUIERTE AZABICYCLOHEPTAN-DERIVATE, IHRE HERSTELLUNG UND VERWENDUNG



(57) Abstract

The invention relates to compounds of formula (1), wherein R¹, R² and R³ have the meaning cited in the description. The novel substances are suitable for use in disease control.

(57) Zusammenfassung

Es werden Verbindungen der Formel (I) beschrieben, worin R¹, R² und R³ die in der Beschreibung angegebene Bedeutung haben. Die neuen Substanzen eignen sich zur Bekämpfung von Krankheiten.

N-SUBSTITUTED AZABICYCLOHEPTANE DERIVATIVES,
PRODUCTION AND USE THEREOF

The invention relates to novel N-substituted azabicycloheptane derivatives, their preparation and use for controlling diseases.

Exo-6-phenyl-3-azabicyclo[3.2.0]heptane derivatives have interesting properties as potential neuroleptics (WO 94/00458, 10 WO 95/15312). Of particular importance in this connection are the observed high affinities for D₄ and 5-HT₂ receptors.

The most interesting substance from the above class of compounds with high D₄/5-HT_{2A} affinity and good selectivity versus D₂ is 15 (+)-(1S,5R,6S)-exo-3-[2-[6-(4-fluorophenyl)-3-azbicyclo[3.2.0]-heptan-3-yl]ethyl]-1H,3H-quinazoline-2,4-dione (= substance A), which is a potential neuroleptic. However, there is an upper limit to dosage of substance A due to prolongations which occur in the QT interval in the cardiac [sic] ECG.


20

We have now found substances with better properties.

The invention relates to N-substituted 3-azabicyclo[3.2.0]-heptane derivatives of the formula I

25

30



in which

35

- R¹ is fluorine or chlorine,
R² is C₁-C₃-alkyl or cyclopropyl, and
R³ is hydrogen, fluorine or chlorine,

40 and the salts thereof with physiologically tolerated acids.


1a

Preferred compounds are those in which

R¹ is chlorine, preferably in the p position,

R² is methyl or cyclopropyl and

R³ is hydrogen.



0050/49277


2

The following compounds are to be mentioned as particularly preferred:

(+)-(1S,5R,6S)-exo-1-[2-[6-(4-chlorophenyl)-3-azabicyclo[3.2.0]-
 5 heptan-3-yl]ethyl]-3-methyl-2H-1,3-dihydrobenzimidazol-2-one,
 (+)-(1S,5R,6S)-exo-1-[2-[6-(4-chlorophenyl)-3-azabicyclo[3.2.0]-
 heptan-3-yl]ethyl]-3-ethyl-2H-1,3-dihydrobenzimidazol-2-one, and
 (+)-(1S,5R,6S)-exo-1-[2-[6-(4-fluorophenyl)-3-azabicyclo[3.2.0]-
 10 heptan-3-yl]ethyl]-3-cyclopropyl-2H-1,3-dihydrobenzimidazol-
 2-one.

The compounds of the formula I according to the invention can be prepared by reacting a compound of the formula II


15



II,

20

in which R^2 and R^3 have the abovementioned meanings, and in Nu is a nucleofugic leaving group, with a 3-azabicyclo[3.2.0]heptane
 25 derivative of the formula III as (+)-(1S,5R,exo-6S) enantiomer



III,

30

in which R^1 has the abovementioned meaning, and converting the compound obtained in this way where appropriate into the acid addition salt with a physiologically tolerated acid.

35

Suitable and preferred nucleofugic leaving groups for Nu are halogen atoms, in particular bromine or chlorine.

The reaction expediently takes place in the presence of an inert
 40 base such as triethylamine or potassium carbonate as acid acceptor in an inert solvent such as a cyclic saturated ether, in particular tetrahydrofuran or dioxane, or a benzenoid hydrocarbon such as toluene or xylene.

45

0050/49277

3

The reaction generally takes place at temperatures from 20 to 150°C, in particular from 80 to 140°C, and is generally complete within 1 to 10 hours.

- 5 The compounds of the formula I according to the invention can be either recrystallized by recrystallization [sic] from conventional organic solvents, preferably from a lower alcohol such as ethanol, or purified by column chromatography.
- 10 The free 3-azabicyclo[3.2.0]heptane derivatives of the formula I can be converted in a conventional way into the acid addition salt with a pharmacologically suitable acid, preferably by treating a solution with one equivalent of the appropriate acid. Examples of pharmaceutically suitable acids are hydrochloric
- 15 acid, phosphoric acid, sulfuric acid, methanesulfonic acid, sulfamic acid, maleic acid, fumaric acid, oxalic acid, tartaric acid or citric acid.

The compounds according to the invention have valuable

20 pharmacological properties. They can be used as neuroleptics (in particular atypical), antidepressants, sedatives, hypnotics, CNS protective agents or agents for treating cocaine dependency. It is possible for several of the types of action mentioned to occur in combination in a compound according to the invention.

25

The substances are characterized in particular by a very high and selective affinity for the dopamine D₄ and serotonin 2A receptors.

The prolongations of the QT interval measured using the model of

30 the guinea pig papillary muscle are negligibly small. This means that the novel substances are well tolerated even at high dosages.

The invention accordingly also relates to a therapeutic

35 composition which has a content of a compound of the formula I or its pharmacologically suitable acid addition salt as active ingredient in addition to conventional carriers and diluents, and to the use of novel compounds for controlling diseases.

40 The compounds according to the invention can be administered in a conventional way orally or parenterally, intravenously or intramuscularly.

The dosage depends on the age, condition and weight of the

45 patient and on the mode of administration. The daily dose of active ingredient is usually between about 1 and 100 mg/kg of

0050/49277

4

body weight on oral administration and between 0.1 and 10 mg/kg of body weight on parenteral administration.

The novel compounds can be used in conventional solid or liquid
5 pharmaceutical forms, e.g. as uncoated or (film-coated) tablets, capsules, powders, granules, suppositories, solutions, ointments, creams or sprays. These are produced in a conventional way. The active ingredients can for this purpose be processed with conventional pharmaceutical aids such as tablet binders, bulking
10 agents, preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, release-slowing agents, antioxidants and/or propellant gases (cf. H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1978). The administration forms obtained in this way
15 normally contain the active ingredient in an amount of from 1 to 99% by weight.

The substances of the formula II and III required as starting materials for synthesizing the compounds according to the
20 invention are known (WO 94/00458; Heterocycles 40 (1), 319-330 (1995); J. Heterocyclic Chem. 18, 85-89 (1981)) or can be synthesized from analogous starting materials using the preparation methods described in the literature.

25 The following examples serve to illustrate the invention:

A Preparation of the starting materials

a) 1-(α -Phenylvinyl)-2H-1,3-dihydrobenzimidazol-2-one

30

21.6 g (200 mM [sic]) of o-phenylenediamine and 37 ml (214 mM [sic]) of ethyl benzoylacetate in 75 ml of 4-tert-butyltoluene with the addition of a spatula tip of p-toluenesulfonic acid were refluxed with a water trap at
35 a bath temperature of 200°C under nitrogen for 1 h, and the liberated water was separated off. After cooling, 80 ml of acetonitrile were added to the reaction mixture and, after stirring in an ice bath, the solid was filtered off with suction and washed with cold
40 acetonitrile. 39.5 g (84%) of product were isolated, melting point 167-169°C.

1-(α -Phenylvinyl)-6-chloro-2H-1,3-dihydrobenzimidazol-2-one can be prepared in an analogous manner (starting material:
45 4-chloro-1,2-diaminobenzene).

0050/49277

5

- b) 1-(α -Phenylvinyl)-3-methyl-2H-1,3-dihydrobenzimidazol-2-one

5 40 ml of 10% strength sodium hydroxide solution, 0.2 g of benzyltriethylammonium chloride and 1.76 ml (18.5 mM [sic]) of dimethyl sulfate were added to 3.5 g (14.8 mM [sic]) of 1-(α -phenylvinyl)-2H-1,3-dihydrobenzimidazol-2-one in 70 ml of toluene, and the mixture was stirred at 10 60°C for 2 h. The toluene phase was then concentrated, and 3.65 g (99%) of product were isolated as an oil with sufficient purity for the next reaction.

The following were prepared in an analogous manner:

15 1-(α -phenylvinyl)-3-ethyl-2H-1,3-dihydrobenzimidazol-2-one (alkylating agent: diethyl sulfate) and 1-(α -phenylvinyl)-3-n-propyl-2H-1,3-dihydrobenzimidazol-2-one (alkylating agent 1-bromopropane), 1-(α -phenyl-20 vinyl)-3-methyl-6-chloro-2H-1,3-dihydrobenzimidazol-2-one (starting material: 1-(α -phenylvinyl)-6-chloro-2H-1,3-dihydrobenzimidazol-2-one) and 1-(2-chloroethyl-3-methyl-5-chloro-2H-1,3-dihydrobenzimidazol-2-one [sic] (starting material: 1-(2-chloroethyl-5-chloro-2H-1,3-dihydrobenzimidazol-2-one [sic])).

- 25 c) 1-Methyl-2H-1,3-dihydrobenzimidazol-2-one

3.65 g (14.6 mM [sic]) 1-(α -phenylvinyl)-3-methyl-2H-1,3-dihydrobenzimidazol-2-one were dissolved in 30 ml of ethanol and, after addition of 60 ml of 10% strength 30 hydrochloric acid and of 10 ml of concentrated hydrochloric acid, stirred at 80°C for 2 h. The mixture was then concentrated, and ice was added to the remaining aqueous solution. The precipitated solids were stirred 35 while cooling in an ice bath, filtered off with suction and washed with water. 1.7 g (79%) of product were isolated.

The following were prepared analogously: 1-(2-chloro-ethyl)-5-chloro-2H-1,3-dihydrobenzimidazol-2-one 40 (starting material: 1-(2-chloroethyl)-3-(α -phenylvinyl)-5-chloro-2H-1,3-dihydrobenzimidazol-2-one) and 1-methyl-5-chloro-2H-1,3-dihydrobenzimidazol-2-one (starting material: 1-(α -phenylvinyl)-3-methyl-6-chloro-2H-1,3-dihydrobenzimidazol-2-one). 45

0050/49277

6

- d) 1-(2-Chloroethyl)-3-methyl-2H-1,3-dihydrobenzimidazol-2-one

5 1.7 g (11.5 mM [sic]) of 1-methyl-2H-1,3-dihydrobenzimidazol-2-one in 40 ml of acetonitrile were mixed with 1.6 g (11.5 mM [sic]) of finely powdered potassium carbonate and 2.9 ml (35 mM [sic]) of 1-bromo-2-chloroethane and refluxed for 14 h. After cooling, the solids were filtered off with suction,
10 washing with acetonitrile, and then the filtrate was concentrated. 2.3 g (95%) of product were isolated as an oil which slowly crystallized, melting point 87-89°C.

15 The following were prepared analogously: 1-(2-chloroethyl)-3-(α -phenylvinyl)-5-chloro-2H-1,3-dihydrobenzimidazol-2-one (starting material: 1-(α -phenylvinyl)-6-chloro-2H-1,3-dihydrobenzimidazol-2-one) and 1-(2-chloroethyl)-3-methyl-6-chloro-2H-1,3-dihydrobenzimidazol-2-one (starting material: 1-methyl-5-chloro-2H-1,3-dihydrobenzimidazol-2-one).
20

- e) (+)-(1S,5R,6S)-exo-6-(4-Chlorophenyl)-3-azabicyclo-[3.2.0]heptane

25 The (+) enantiomer was isolated by the method described in Heterocycles 40 (1), 326 (1995).

- f) 1-(α -Phenylvinyl)-3-cyclopropyl-2H-1,3-dihydrobenzimidazol-2-one

30 1.35 g (45 mM [sic]) of sodium hydride (80 percent) were added in portions to 10.0 g (42.4 mM [sic]) of 1-(α -phenylvinyl)-2H-1,3-dihydrobenzimidazol-2-one in 80 ml of dimethylformamide with thorough stirring, and
35 the reaction mixture was then stirred for 2 h. Subsequently 8.0 ml (100 mM [sic]) of cyclopropyl bromide were added, and the reaction mixture was transferred into a 0.3 l stirred autoclave which was then heated at 200°C for 10 h. After cooling and concentration in a rotary
40 evaporator, the residue was partitioned between methylene chloride and water, acidifying with 10 percent hydrochloric acid. The aqueous phase was extracted once more with methylene chloride. After drying and concentration of the organic phase, 12.4 g of crude
45 product were isolated and were purified by column chromatography (silica gel, mobile phase methylene chloride). Yield 3.6 g (31%) of adequate purity.

0050/49277

7

B Preparation of the final products

Example 1

5 (+)-(1S,5R,6S)-exo-1-[2-[6-(4-Chlorophenyl)-3-azabicyclo-
[3.2.0]heptan-3-yl]ethyl]-3-methyl-2H-1,3-dihydrobenz-
imidazol-2-one tartrate x 2H₂O

10 2.5 g (12.1 mM [sic]) of (+)-(1S,5R,6S)-exo-6-(4-
chlorophenyl)-3-azabicyclo[3.2.0]heptane in 60 ml of xylene
were mixed with 2.55 g (12.1 mM [sic]) of
1-(2-chloroethyl)-3-methyl-2H-1,3-dihydrobenzimidazol-2-one
and 1.7 g (12.1 mM [sic]) of finely powdered potassium
15 carbonate and refluxed for 20 h. The mixture was then
concentrated in a rotary evaporator, and the residue was
partitioned between water and methyl tert-butyl ether at
pH 10. The aqueous phase was extracted once more with methyl
tert-butyl ether, and then the combined organic phases were
concentrated. The crude product was purified by column
20 chromatography (silica gel, mobile phase methylene
chloride/methanol 97/3. 3.3 g (71%) of product were isolated
as an oil which was dissolved in 200 ml of ether and
converted with 1.4 g (9.3 mM [sic]) of tartaric acid,
dissolved in ethanol, into the tartrate (melting point 107 to
109°C). $[\alpha]_D = + 55.4^\circ$ (EtOH)

25

Elemental analysis C₂₂H₂₄N₃OCl x C₄H₆O₆ x 2H₂O

calculated	C 54.97	H 6.03	N 7.39	O 25.34	Cl 6.24
found	C 54.9	H 5.8	N 7.1	O 25.5	Cl 6.2

30 The following were prepared in an analogous manner:

2. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-Chlorophenyl)-3-azabicyclo-
[3.2.0]heptan-3-yl]ethyl]-3-ethyl-2H-1,3-dihydrobenzimidazol-
2-one x HCl, melting point 174 to 176°C, $[\alpha]_D = + 67.7^\circ$ (EtOH)

35

3. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-Chlorophenyl)-3-azabicyclo-
[3.2.0]heptan-3-yl]ethyl]-3-n-propyl-2H-1,3-dihydrobenz-
imidazol-2-one x HCl, melting point 178 to 180°C, $[\alpha]_D =$
+66.4° (EtOH)

40

4. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-chlorophenyl)-3-azabicyclo-
[3.2.0]heptan-3-yl]ethyl]-3-methyl-5-chloro-2H-1,3-dihydro-
benzimidazol-2-one x HCl, melting point 101 to 103°C, $[\alpha]_D =$
+92.3° (EtOH)

45

0050/49277

8

5. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-chlorophenyl)-3-azabicyclo-[3.2.0]heptan-3-yl]ethyl]-3-methyl-6-chloro-2H-1,3-dihydro-benzimidazol-2-one x HCl, melting point 230 to 232°C, $[\alpha]_D = +77.2^\circ$ (EtOH)

5

6. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-chlorophenyl)-3-azabicyclo-[3.2.0]heptan-3-yl]ethyl]-3-cyclopropyl-2H-1,3-dihydro-benzimidazol-2-one x HCl, melting point 109 to 111°C, $[\alpha]_D = +73.5^\circ$ (EtOH)

10

7. (+)-(1S,5R,6S)-exo-1-[2-[6-(4-fluorophenyl)-3-azabicyclo-[3.2.0]heptan-3-yl]ethyl]-3-cyclopropyl-2H-1,3-dihydro-benzimidazol-2-one x HCl, melting point 121 to 123°C, $[\alpha]_D = +64.5^\circ$ (EtOH)

15

20

25

30

35

40


45

0050/49277

9

We claim:

1. An N-substituted 3-azabicyclo[3.2.0]heptane derivative of the
5 formula I



15 in which

R¹ is fluorine or chlorine,
R² is C₁-C₃-alkyl or cyclopropyl, and
R³ is hydrogen, fluorine or chlorine,

20

and the salts thereof with physiologically tolerated acids.

2. A compound of the formula I as claimed in claim 1 for use in
25 controlling diseases.

30

35

40

45

