



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A01H 1/00, 5/00, C12N 15/00 C12N 9/02, C12P 7/64, 1/02 C12P 1/04, 21/06, C07H 15/12 C07H 17/00</p>	A1	<p>(11) International Publication Number: WO 93/06712</p> <p>(43) International Publication Date: 15 April 1993 (15.04.93)</p>						
<p>(21) International Application Number: PCT/US92/08746</p> <p>(22) International Filing Date: 13 October 1992 (13.10.92)</p> <p>(30) Priority data:</p> <table border="0"> <tr> <td>774,475</td> <td>10 October 1991 (10.10.91)</td> <td>US</td> </tr> <tr> <td>817,919</td> <td>8 January 1992 (08.01.92)</td> <td>US</td> </tr> </table> <p>(60) Parent Application or Grant (63) Related by Continuation US 817,919 (CIP) Filed on 8 January 1992 (08.01.92)</p> <p>(71) Applicant (for all designated States except US): RHONE-POULENC AGROCHIMIE [FR/FR]; 14-20, rue Pierre-Baizet, B.P. 9163, F-69263 Lyon Cédex 09 (FR).</p>		774,475	10 October 1991 (10.10.91)	US	817,919	8 January 1992 (08.01.92)	US	<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : THOMAS, Terry [US/US]; 3004 Normand, College Station, TX 77845 (US). REDDY, Avutu, S. [IN/US]; 3902 E. 29th Street, #G11, Bryan, TX 77802 (US). NUCCIO, Michael [US/US]; P.O. Box 553, College Station, TX 77841 (US). FREYSINET, Georges [FR/FR]; 21, rue de Nervieux, F-Saint-Cyr-au-Mont-d'Or (FR).</p> <p>(74) Agents: SCOTT, Anthony, C. et al.; Scully, Scott, Murphy & Presser, 400 Garden City Plaza, Garden City, NY 11530 (US).</p> <p>(81) Designated States: AU, BG, BR, CA, CS, HU, JP, KR, PL, RO, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).</p> <p>Published <i>With international search report.</i></p>
774,475	10 October 1991 (10.10.91)	US						
817,919	8 January 1992 (08.01.92)	US						
<p>(54) Title: PRODUCTION OF GAMMA LINOLENIC ACID BY A Δ6-DESATURASE</p>								
<p>(57) Abstract</p> <p>Linoleic acid is converted into γ-linolenic acid by the enzyme Δ6-desaturase. The present invention is directed to an isolated nucleic acid comprising the Δ6-desaturase gene. More particularly, the isolated nucleic acid comprises the promoter, coding region and termination regions of the Δ6-desaturase gene. The present invention provides recombinant constructions comprising the Δ6-desaturase coding region in functional combination with heterologous regulatory sequences. The nucleic acids and recombinant constructions of the instant invention are useful in the production of GLA in transgenic organisms.</p>								

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SK	Slovak Republic
CI	Côte d'Ivoire	LK	Sri Lanka	SN	Senegal
CM	Cameroon	LU	Luxembourg	SU	Soviet Union
CS	Czechoslovakia	MC	Monaco	TD	Chad
CZ	Czech Republic	MG	Madagascar	TG	Togo
DE	Germany	ML	Mali	UA	Ukraine
DK	Denmark	MN	Mongolia	US	United States of America
ES	Spain			VN	Viet Nam
FI	Finland				

PRODUCTION OF GAMMA LINOLENIC ACID
BY A Δ 6-DESATURASE

1

Linoleic acid (18:2) (LA) is transformed into
gamma linolenic acid (18:3) (GLA) by the enzyme Δ 6-
5 desaturase. When this enzyme, or the nucleic acid
encoding it, is transferred into LA-producing cells, GLA
is produced. The present invention provides a nucleic
acid comprising the Δ 6-desaturase gene. More
specifically, the nucleic acid comprises the promoter,
10 coding region and termination regions of the Δ 6-
desaturase gene. The present invention is further
directed to recombinant constructions comprising a Δ 6-
desaturase coding region in functional combination with
heterologous regulatory sequences. The nucleic acids
15 and recombinant constructions of the instant invention
are useful in the production of GLA in transgenic
organisms.

Unsaturated fatty acids such as linoleic
($C_{18}\Delta^{9,12}$) and α -linolenic ($C_{18}\Delta^{9,12,15}$) acids are
20 essential dietary constituents that cannot be
synthesized by vertebrates since vertebrate cells can
introduce double bonds at the Δ^9 position of fatty acids
but cannot introduce additional double bonds between the
 Δ^9 double bond and the methyl-terminus of the fatty acid
25 chain. Because they are precursors of other products,
linoleic and α -linolenic acids are essential fatty
acids, and are usually obtained from plant sources.
Linoleic acid can be converted by mammals into γ -
linolenic acid (GLA, $C_{18}\Delta^{6,9,12}$) which can in turn be
30 converted to arachidonic acid (20:4), a critically
important fatty acid since it is an essential precursor
of most prostaglandins.

35

-2-

1 The dietary provision of linoleic acid, by virtue
of its resulting conversion to GLA and arachidonic acid,
satisfies the dietary need for GLA and arachidonic acid.
However, a relationship has been demonstrated between
5 consumption of saturated fats and health risks such as
hypercholesterolemia, atherosclerosis and other chemical
disorders which correlate with susceptibility to
coronary disease, while the consumption of unsaturated
fats has been associated with decreased blood
10 cholesterol concentration and reduced risk of
atherosclerosis. The therapeutic benefits of dietary
GLA may result from GLA being a precursor to arachidonic
acid and thus subsequently contributing to prostaglandin
synthesis. Accordingly, consumption of the more
15 unsaturated GLA, rather than linoleic acid, has
potential health benefits. However, GLA is not present
in virtually any commercially grown crop plant.

Linoleic acid is converted into GLA by the enzyme
 $\Delta 6$ -desaturase. $\Delta 6$ -desaturase, an enzyme of about 359
20 amino acids, has a membrane-bound domain and an active
site for desaturation of fatty acids. When this enzyme
is transferred into cells which endogenously produce
linoleic acid but not GLA, GLA is produced. The present
invention, by providing the gene encoding $\Delta 6$ -desaturase,
25 allows the production of transgenic organisms which
contain functional $\Delta 6$ -desaturase and which produce GLA.
In addition to allowing production of large amounts of
GLA, the present invention provides new dietary sources
of GLA.

30 The present invention is directed to an isolated
 $\Delta 6$ -desaturase gene. Specifically, the isolated gene

35

1 comprises the $\Delta 6$ -desaturase promoter, coding region, and
termination region.

The present invention is further directed to
expression vectors comprising the $\Delta 6$ -desaturase
5 promoter, coding region and termination region.

The present invention is also directed to
expression vectors comprising a $\Delta 6$ -desaturase coding
region in functional combination with heterologous
regulatory regions, i.e. elements not derived from the
10 $\Delta 6$ -desaturase gene.

Cells and organisms comprising the vectors of the
present invention, and progeny of such organisms, are
also provided by the present invention.

The present invention further provides isolated
15 bacterial $\Delta 6$ -desaturase and is still further directed to
an isolated nucleic acid encoding bacterial $\Delta 6$ -
desaturase.

The present invention further provides a method
for producing plants with increased gamma linolenic acid
20 (GLA) content which comprises transforming a plant cell
with an isolated nucleic acid of the present invention
and regenerating a plant with increased GLA content from
said plant cell.

A method for producing chilling tolerant plants
25 is also provided by the present invention.

Fig. 1 depicts the hydropathy profiles of the
deduced amino acid sequences of Synechocystis $\Delta 6$ -
desaturase (Panel A) and $\Delta 12$ -desaturase (Panel B).
Putative membrane spanning regions are indicated by
30 solid bars. Hydrophobic index was calculated for a
window size of 19 amino acid residues [Kyte, et al.
(1982) J. Molec. Biol. 157].

1 Fig. 2 provides gas liquid chromatography
profiles of wild type (Panel A) and transgenic (Panel B)
Anabaena.

5 Fig. 3 is a diagram of maps of cosmid cSy75,
cSy13 and cSy7 with overlapping regions and subclones.
The origins of subclones of cSy75, cSy75-3.5 and cSy7
are indicated by the dashed diagonal lines. Restriction
sites that have been inactivated are in parentheses.

10 Fig. 4 provides gas liquid chromatography
profiles of wild type (Panel A) and transgenic (Panel B)
tobacco.

 The present invention provides an isolated
nucleic acid encoding $\Delta 6$ -desaturase. To identify a
nucleic acid encoding $\Delta 6$ -desaturase, DNA is isolated
15 from an organism which produces GLA. Said organism can
be, for example, an animal cell, certain fungi (e.g.
Mortierella), certain bacteria (e.g. Synechocystis) or
certain plants (borage, Oenothera, currants). The
isolation of genomic DNA can be accomplished by a
20 variety of methods well-known to one of ordinary skill
in the art, as exemplified by Sambrook et al. (1989) in
Molecular Cloning: A Laboratory Manual, Cold Spring
Harbor, NY. The isolated DNA is fragmented by physical
methods or enzymatic digestion and cloned into an
25 appropriate vector, e.g. a bacteriophage or cosmid
vector, by any of a variety of well-known methods which
can be found in references such as Sambrook et al.
(1989). Expression vectors containing the DNA of the
present invention are specifically contemplated herein.
30 DNA encoding $\Delta 6$ -desaturase can be identified by gain of
function analysis. The vector containing fragmented DNA
is transferred, for example by infection,

-5-

1 transconjugation, transfection, into a host organism
that produces linoleic acid but not GLA. As used
herein, "transformation" refers generally to the
incorporation of foreign DNA into a host cell. Methods
5 for introducing recombinant DNA into a host organism are
known to one of ordinary skill in the art and can be
found, for example, in Sambrook et al. (1989).
Production of GLA by these organisms (i.e., gain of
function) is assayed, for example by gas chromatography
10 or other methods known to the ordinarily skilled
artisan. Organisms which are induced to produce GLA,
i.e. have gained function by the introduction of the
vector, are identified as expressing DNA encoding $\Delta 6$ -
desaturase, and said DNA is recovered from the
15 organisms. The recovered DNA can again be fragmented,
cloned with expression vectors, and functionally
assessed by the above procedures to define with more
particularity the DNA encoding $\Delta 6$ -desaturase.

As an example of the present invention, random
20 DNA is isolated from the cyanobacteria Synechocystis
Pasteur Culture Collection (PCC) 6803, American Type
Culture Collection (ATCC) 27184, cloned into a cosmid
vector, and introduced by transconjugation into the GLA-
deficient cyanobacterium Anabaena strain PCC 7120, ATCC
25 27893. Production of GLA from Anabaena linoleic acid is
monitored by gas chromatography and the corresponding
DNA fragment is isolated.

The isolated DNA is sequenced by methods well-
known to one of ordinary skill in the art as found, for
30 example, in Sambrook et al. (1989).

In accordance with the present invention, a DNA
comprising a $\Delta 6$ -desaturase gene has been isolated. More

35

1 particularly, a 3.588 kilobase (kb) DNA comprising a $\Delta 6$ -
desaturase gene has been isolated from the cyanobacteria
2 Synechocystis. The nucleotide sequence of the 3.588 kb
DNA was determined and is shown in SEQ ID NO:1. Open
5 reading frames defining potential coding regions are
present from nucleotide 317 to 1507 and from nucleotide
2002 to 3081. To define the nucleotides responsible for
encoding $\Delta 6$ -desaturase, the 3.588 kb fragment that
confers $\Delta 6$ -desaturase activity is cleaved into two
10 subfragments, each of which contains only one open
reading frame. Fragment ORF1 contains nucleotides 1
through 1704, while fragment ORF2 contains nucleotides
1705 through 3588. Each fragment is subcloned in both
forward and reverse orientations into a conjugal
15 expression vector (AM542, Wolk et al. [1984] Proc. Natl.
Acad. Sci. USA 81, 1561) that contains a cyanobacterial
carboxylase promoter. The resulting constructs (i.e.
ORF1(F), ORF1(R), ORF2(F) and ORF2(R)) are conjugated to
wild-type Anabaena PCC 7120 by standard methods (see,
20 for example, Wolk et al. (1984) Proc. Natl. Acad. Sci.
USA 81, 1561). Conjugated cells of Anabaena are
identified as Neo^R green colonies on a brown background
of dying non-conjugated cells after two weeks of growth
on selective media (standard mineral media BG11N +
25 containing 30 μ g/ml of neomycin according to Rippka et
al., (1979) J. Gen Microbiol. 111, 1). The green
colonies are selected and grown in selective liquid
media (BG11N + with 15 μ g/ml neomycin). Lipids are
extracted by standard methods (e.g. Dahmer et al.,
30 (1989) Journal of American Oil Chemical Society 66, 543)
from the resulting transconjugants containing the
forward and reverse oriented ORF1 and ORF2 constructs.

1 For comparison, lipids are also extracted from wild-type
cultures of Anabaena and Synechocystis. The fatty acid
methyl esters are analyzed by gas liquid chromatography
(GLC), for example with a Tracor-560 gas liquid
5 chromatograph equipped with a hydrogen flame ionization
detector and a capillary column. The results of GLC
analysis are shown in Table 1.

Table 1: Occurrence of C18 fatty acids in wild-type and
transgenic cyanobacteria

10

SOURCE	18:0	18:1	18:2	γ 18:3	α 18:3	18:4
Anabaena (wild type)	+	+	+	-	+	-
Anabaena + ORF1(F)	+	+	+	-	+	-
15 Anabaena + ORF1(R)	+	+	+	-	+	-
Anabaena + ORF2(F)	+	+	+	+	+	+
Anabaena + ORF2(R)	+	+	+	-	+	-
Synechocystis (wild type)	+	+	+	+	-	-

20

As assessed by GLC analysis, GLA deficient
Anabaena gain the function of GLA production when the
construct containing ORF2 in forward orientation is
introduced by transconjugation. Transconjugants
25 containing constructs with ORF2 in reverse orientation
to the carboxylase promoter, or ORF1 in either
orientation, show no GLA production. This analysis
demonstrates that the single open reading frame (ORF2)
within the 1884 bp fragment encodes Δ 6-desaturase. The
30 1884 bp fragment is shown as SEQ ID NO:3. This is
substantiated by the overall similarity of the
hydropathy profiles between Δ 6-desaturase and Δ 12-

35

1 desaturase [Wada et al. (1990) Nature 347] as shown in
Fig. 1 as (A) and (B), respectively.

Isolated nucleic acids encoding Δ^6 -desaturase can
be identified from other GLA-producing organisms by the
5 gain of function analysis described above, or by nucleic
acid hybridization techniques using the isolated nucleic
acid which encodes Anabaena Δ^6 -desaturase as a
hybridization probe. Both genomic and cDNA cloning
methods are known to the skilled artisan and are
10 contemplated by the present invention. The
hybridization probe can comprise the entire DNA sequence
disclosed as SEQ. ID NO:1, or a restriction fragment or
other DNA fragment thereof, including an oligonucleotide
probe. Methods for cloning homologous genes by cross-
15 hybridization are known to the ordinarily skilled
artisan and can be found, for example, in Sambrook
(1989) and Beltz et al. (1983) Methods in Enzymology
100, 266.

Transgenic organisms which gain the function of
20 GLA production by introduction of DNA encoding Δ^6 -
desaturase also gain the function of octadecatetraenoic
acid ($18:4\Delta^{6,9,12,15}$) production. Octadecatetraenoic
acid is present normally in fish oils and in some plant
species of the Boraginaceae family (Craig et al. [1964]
25 J. Amer. Oil Chem. Soc. 41, 209-211; Gross et al. [1976]
Can. J. Plant Sci. 56, 659-664). In the transgenic
organisms of the present invention, octadecatetraenoic
acid results from further desaturation of α -linolenic
acid by Δ^6 -desaturase or desaturation of GLA by Δ^{15} -
30 desaturase.

The 359 amino acids encoded by ORF2, i.e. the
open reading frame encoding Δ^6 -desaturase, are shown as

1 SEQ. ID NO:2. The present invention further
contemplates other nucleotide sequences which encode the
amino acids of SEQ ID NO:2. It is within the ken of the
ordinarily skilled artisan to identify such sequences
5 which result, for example, from the degeneracy of the
genetic code. Furthermore, one of ordinary skill in the
art can determine, by the gain of function analysis
described hereinabove, smaller subfragments of the 1884
bp fragment containing ORF2 which encode $\Delta 6$ -desaturase.

10 The present invention contemplates any such
polypeptide fragment of $\Delta 6$ -desaturase and the nucleic
acids therefor which retain activity for converting LA
to GLA.

In another aspect of the present invention, a
15 vector containing the 1884 bp fragment or a smaller
fragment containing the promoter, coding sequence and
termination region of the $\Delta 6$ -desaturase gene is
transferred into an organism, for example,
cyanobacteria, in which the $\Delta 6$ -desaturase promoter and
20 termination regions are functional. Accordingly,
organisms producing recombinant $\Delta 6$ -desaturase are
provided by this invention. Yet another aspect of this
invention provides isolated $\Delta 6$ -desaturase, which can be
purified from the recombinant organisms by standard
25 methods of protein purification. (For example, see
Ausubel et al. [1987] Current Protocols in Molecular
Biology, Green Publishing Associates, New York).

Vectors containing DNA encoding $\Delta 6$ -desaturase are
also provided by the present invention. It will be
30 apparent to one of ordinary skill in the art that
appropriate vectors can be constructed to direct the
expression of the $\Delta 6$ -desaturase coding sequence in a

-10-

1 variety of organisms. Replicable expression vectors are
particularly preferred. Replicable expression vectors
as described herein are DNA or RNA molecules engineered
for controlled expression of a desired gene, i.e. the
5 $\Delta 6$ -desaturase gene. Preferably the vectors are
plasmids, bacteriophages, cosmids or viruses. Shuttle
vectors, e.g. as described by Wolk et al. (1984) Proc.
Natl. Acad. Sci. USA, 1561-1565 and Bustos et al. (1991)
J. Bacteriol. 174, 7525-7533, are also contemplated in
10 accordance with the present invention. Sambrook et al.
(1989), Goeddel, ed. (1990) Methods in Enzymology 185
Academic Press, and Perbal (1988) A Practical Guide to
Molecular Cloning, John Wiley and Sons, Inc., provide
15 detailed reviews of vectors into which a nucleic acid
encoding the present $\Delta 6$ -desaturase can be inserted and
expressed. Such vectors also contain nucleic acid
sequences which can effect expression of nucleic acids
encoding $\Delta 6$ -desaturase. Sequence elements capable of
20 effecting expression of a gene product include
promoters, enhancer elements, upstream activating
sequences, transcription termination signals and
polyadenylation sites. Both constitutive and tissue
specific promoters are contemplated. For transformation
of plant cells, the cauliflower mosaic virus (CaMV) 35S
25 promoter and promoters which are regulated during plant
seed maturation are of particular interest. All such
promoter and transcriptional regulatory elements, singly
or in combination, are contemplated for use in the
present replicable expression vectors and are known to
30 one of ordinary skill in the art. The CaMV 35S promoter
is described, for example, by Restrepo et al. (1990)

35

1 Plant Cell 2, 987. Genetically engineered and mutated
regulatory sequences are also contemplated.

The ordinarily skilled artisan can determine
vectors and regulatory elements suitable for expression
5 in a particular host cell. For example, a vector
comprising the promoter from the gene encoding the
carboxylase of Anabaena operably linked to the coding
region of $\Delta 6$ -desaturase and further operably linked to a
termination signal from Synechocystis is appropriate for
10 expression of $\Delta 6$ -desaturase in cyanobacteria. "Operably
linked" in this context means that the promoter and
terminator sequences effectively function to regulate
transcription. As a further example, a vector
appropriate for expression of $\Delta 6$ -desaturase in
15 transgenic plants can comprise a seed-specific promoter
sequence derived from helianthinin, napin, or glycin
operably linked to the $\Delta 6$ -desaturase coding region and
further operably linked to a seed termination signal or
the nopaline synthase termination signal.

20 In particular, the helianthinin regulatory
elements disclosed in applicant's copending U.S.
Application Serial No. 682,354, filed April 8, 1991 and
incorporated herein by reference, are contemplated as
promoter elements to direct the expression of the $\Delta 6$ -
25 desaturase of the present invention.

Modifications of the nucleotide sequences or
regulatory elements disclosed herein which maintain the
functions contemplated herein are within the scope of
this invention. Such modifications include insertions,
30 substitutions and deletions, and specifically
substitutions which reflect the degeneracy of the
genetic code.

-12-

1 Standard techniques for the construction of such
hybrid vectors are well-known to those of ordinary skill
in the art and can be found in references such as
Sambrook et al. (1989), or any of the myriad of
5 laboratory manuals on recombinant DNA technology that
are widely available. A variety of strategies are
available for ligating fragments of DNA, the choice of
which depends on the nature of the termini of the DNA
fragments. It is further contemplated in accordance
10 with the present invention to include in the hybrid
vectors other nucleotide sequence elements which
facilitate cloning, expression or processing, for
example sequences encoding signal peptides, a sequence
encoding KDEL, which is required for retention of
15 proteins in the endoplasmic reticulum or sequences
encoding transit peptides which direct $\Delta 6$ -desaturase to
the chloroplast. Such sequences are known to one of
ordinary skill in the art. An optimized transit peptide
is described, for example, by Van den Broeck et al.
20 (1985) Nature 313, 358. Prokaryotic and eukaryotic
signal sequences are disclosed, for example, by
Michaelis et al. (1982) Ann. Rev. Microbiol. 36, 425.

A further aspect of the instant invention
provides organisms other than cyanobacteria which
25 contain the DNA encoding the $\Delta 6$ -desaturase of the
present invention. The transgenic organisms
contemplated in accordance with the present invention
include bacteria, cyanobacteria, fungi, and plants and
animals. The isolated DNA of the present invention can
30 be introduced into the host by methods known in the art,
for example infection, transfection, transformation or
transconjugation. Techniques for transferring the DNA

35

1 of the present invention into such organisms are widely
known and provided in references such as Sambrook et al.
(1989).

5 A variety of plant transformation methods are
known. The $\Delta 6$ -desaturase gene can be introduced into
plants by a leaf disk transformation-regeneration
procedure as described by Horsch et al. (1985) Science
227, 1229. Other methods of transformation, such as
protoplast culture (Horsch et al. (1984) Science 223,
10 496; DeBlock et al. (1984) EMBO J. 2, 2143; Barton et
al. (1983) Cell 32, 1033) can also be used and are
within the scope of this invention. In a preferred
embodiment plants are transformed with Agrobacterium-
derived vectors. However, other methods are available
15 to insert the $\Delta 6$ -desaturase gene of the present
invention into plant cells. Such alternative methods
include biolistic approaches (Klein et al. (1987) Nature
327, 70), electroporation, chemically-induced DNA
uptake, and use of viruses or pollen as vectors.

20 When necessary for the transformation method, the
 $\Delta 6$ -desaturase gene of the present invention can be
inserted into a plant transformation vector, e.g. the
binary vector described by Bevan (1984) Nucleic Acids
Res. 12, 8111. Plant transformation vectors can be
25 derived by modifying the natural gene transfer system of
Agrobacterium tumefaciens. The natural system comprises
large Ti (tumor-inducing)-plasmids containing a large
segment, known as T-DNA, which is transferred to
transformed plants. Another segment of the Ti plasmid,
30 the vir region, is responsible for T-DNA transfer. The
T-DNA region is bordered by terminal repeats. In the
modified binary vectors the tumor-inducing genes have

35

-14-

1 been deleted and the functions of the vir region are
utilized to transfer foreign DNA bordered by the T-DNA
border sequences. The T-region also contains a
selectable marker for antibiotic resistance, and a
5 multiple cloning site for inserting sequences for
transfer. Such engineered strains are known as
"disarmed" A. tumefaciens strains, and allow the
efficient transformation of sequences bordered by the T-
region into the nuclear genomes of plants.

10 Surface-sterilized leaf disks are inoculated with
the "disarmed" foreign DNA-containing A. tumefaciens,
cultured for two days, and then transferred to
antibiotic-containing medium. Transformed shoots are
selected after rooting in medium containing the
15 appropriate antibiotic, transferred to soil and
regenerated.

Another aspect of the present invention provides
transgenic plants or progeny of these plants containing
the isolated DNA of the invention. Both
20 monocotyledenous and dicotyledenous plants are
contemplated. Plant cells are transformed with the
isolated DNA encoding $\Delta 6$ -desaturase by any of the plant
transformation methods described above. The transformed
plant cell, usually in a callus culture or leaf disk, is
25 regenerated into a complete transgenic plant by methods
well-known to one of ordinary skill in the art (e.g.
Horsch et al. (1985) Science 227, 1129). In a preferred
embodiment, the transgenic plant is sunflower, oil seed
rape, maize, tobacco, peanut or soybean. Since progeny
30 of transformed plants inherit the DNA encoding $\Delta 6$ -
desaturase, seeds or cuttings from transformed plants
are used to maintain the transgenic plant line.

35

1 The present invention further provides a method
for providing transgenic plants with an increased
content of GLA. This method includes introducing DNA
encoding $\Delta 6$ -desaturase into plant cells which lack or
5 have low levels of GLA but contain LA, and regenerating
plants with increased GLA content from the transgenic
cells. In particular, commercially grown crop plants
are contemplated as the transgenic organism, including,
but not limited to, sunflower, soybean, oil seed rape,
10 maize, peanut and tobacco.

 The present invention further provides a method
for providing transgenic organisms which contain GLA.
This method comprises introducing DNA encoding $\Delta 6$ -
desaturase into an organism which lacks or has low
15 levels of GLA, but contains LA. In another embodiment,
the method comprises introducing one or more expression
vectors which comprise DNA encoding $\Delta 12$ -desaturase and
 $\Delta 6$ -desaturase into organisms which are deficient in both
GLA and LA. Accordingly, organisms deficient in both LA
20 and GLA are induced to produce LA by the expression of
 $\Delta 12$ -desaturase, and GLA is then generated due to the
expression of $\Delta 6$ -desaturase. Expression vectors
comprising DNA encoding $\Delta 12$ -desaturase, or $\Delta 12$ -
desaturase and $\Delta 6$ -desaturase, can be constructed by
25 methods of recombinant technology known to one of
ordinary skill in the art (Sambrook et al., 1989) and
the published sequence of $\Delta 12$ -desaturase (Wada et al
[1990] Nature (London) 347, 200-203. In addition, it
has been discovered in accordance with the present
30 invention that nucleotides 2002-3081 of SEQ. ID NO:1
encode cyanobacterial $\Delta 12$ -desaturase. Accordingly, this
sequence can be used to construct the subject expression

-16-

1 vectors. In particular, commercially grown crop plants
are contemplated as the transgenic organism, including,
but not limited to, sunflower, soybean, oil seed rape,
maize, peanut and tobacco.

5 The present invention is further directed to a
method of inducing chilling tolerance in plants.
Chilling sensitivity may be due to phase transition of
lipids in cell membranes. Phase transition temperature
depends upon the degree of unsaturation of fatty acids
10 in membrane lipids, and thus increasing the degree of
unsaturation, for example by introducing $\Delta 6$ -desaturase
to convert LA to GLA, can induce or improve chilling
resistance. Accordingly, the present method comprises
introducing DNA encoding $\Delta 6$ -desaturase into a plant
15 cell, and regenerating a plant with improved chilling
resistance from said transformed plant cell. In a
preferred embodiment, the plant is a sunflower, soybean,
oil seed rape, maize, peanut or tobacco plant.

The following examples further illustrate the
20 present invention.

25

30

35

1

EXAMPLE 1

Strains and Culture Conditions

Synechocystis (PCC 6803, ATCC 27184), Anabaena (PCC 7120, ATCC 27893) and Synechococcus (PCC 7942, ATCC 33912) were grown photoautotrophically at 30°C in BG11N+ medium (Rippka et al. [1979] J. Gen. Microbiol. 111, 1-61) under illumination of incandescent lamps ($60\mu\text{E}\cdot\text{m}^{-2}\cdot\text{S}^{-1}$). Cosmids and plasmids were selected and propagated in Escherichia coli strain DH5 α on LB medium supplemented with antibiotics at standard concentrations as described by Maniatis et al. (1982) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring, New York.

15

20

25

30

35

1

EXAMPLE 2

Construction of Synechocystis Cosmid Genomic Library

Total genomic DNA from Synechocystis (PCC 6803) was partially digested with Sau3A and fractionated on a sucrose gradient (Ausubel et al. [1987] Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley Interscience, New York). Fractions containing 30 to 40 kb DNA fragments were selected and ligated into the dephosphorylated BamHI site of the cosmid vector, pDUCA7 (Buikema et al. [1991] J. Bacteriol. 173, 1879-1885). The ligated DNA was packaged in vitro as described by Ausubel et al. (1987), and packaged phage were propagated in E. coli DH5 α containing the AvaI and Eco4711 methylase helper plasmid, pRL528 as described by Buikema et al. (1991). A total of 1152 colonies were isolated randomly and maintained individually in twelve 96-well microtiter plates.

20

25

30

35

1

EXAMPLE 3

Gain-of-Function Expression of GLA in Anabaena

Anabaena (PCC 7120), a filamentous cyanobacterium, is deficient in GLA but contains significant amounts of linoleic acid, the precursor for GLA (Figure 2; Table 2). The Synechocystis cosmid library described in Example 2 was conjugated into Anabaena (PCC 7120) to identify transconjugants that produce GLA. Anabaena cells were grown to mid-log phase in BG11N+ liquid medium and resuspended in the same medium to a final concentration of approximately 2×10^8 cells per ml. A mid-log phase culture of E. coli RP4 (Burkardt et al. [1979] J. Gen. Microbiol. 114, 341-348) grown in LB containing ampicillin was washed and resuspended in fresh LB medium. Anabaena and RP4 were then mixed and spread evenly on BG11N+ plates containing 5% LB. The cosmid genomic library was replica plated onto LB plates containing 50 $\mu\text{g/ml}$ kanamycin and 17.5 $\mu\text{g/ml}$ chloramphenicol and was subsequently patched onto BG11N+ plates containing Anabaena and RP4. After 24 hours of incubation at 30°C, 30 $\mu\text{g/ml}$ of neomycin was underlaid; and incubation at 30°C was continued until transconjugants appeared.

Individual transconjugants were isolated after conjugation and grown in 2 ml BG11N+ liquid medium with 15 $\mu\text{g/ml}$ neomycin. Fatty acid methyl esters were prepared from wild type cultures and cultures containing pools of ten transconjugants as follows. Wild type and transgenic cyanobacterial cultures were harvested by centrifugation and washed twice with distilled water. Fatty acid methyl esters were extracted from these cultures as described by Dahmer et al. (1989) J. Amer.

35

1 Oil. Chem. Soc. 66, 543-548 and were analyzed by Gas
Liquid Chromatography (GLC) using a Tracor-560 equipped
with a hydrogen flame ionization detector and capillary
5 Associates Inc., IL). Retention times and co-
chromatography of standards (obtained from Sigma
Chemical Co.) were used for identification of fatty
acids. The average fatty acid composition was
determined as the ratio of peak area of each C18 fatty
10 acid normalized to an internal standard.

Representative GLC profiles are shown in Fig. 2.
C18 fatty acid methyl esters are shown. Peaks were
identified by comparing the elution times with known
standards of fatty acid methyl esters and were confirmed
15 by gas chromatography-mass spectrometry. Panel A
depicts GLC analysis of fatty acids of wild type
Anabaena. The arrow indicates the migration time of
GLA. Panel B is a GLC profile of fatty acids of
transconjugants of Anabaena with pAM542+1.8F. Two GLA
20 producing pools (of 25 pools representing 250
transconjugants) were identified that produced GLA.
Individual transconjugants of each GLA positive pool
were analyzed for GLA production; two independent
transconjugants, AS13 and AS75, one from each pool, were
25 identified which expressed significant levels of GLA and
which contained cosmids, cSy13 and cSy75, respectively
(Figure 3). The cosmids overlap in a region
approximately 7.5 kb in length. A 3.5 kb NheI fragment
of cSy75 was recloned in the vector pDUCA7 and
30 transferred to Anabaena resulting in gain-of-function
expression of GLA (Table 2).

-21-

1 Two NheI/Hind III subfragments (1.8 and 1.7 kb)
of the 3.5 kb Nhe I fragment of cSy75-3.5 were subcloned
into "pBLUESCRIPT" (Stratagene) (Figure 3) for
sequencing. Standard molecular biology techniques were
5 performed as described by Maniatis et al. (1982) and
Ausubel et al. (1987). Dideoxy sequencing (Sanger et al.
[1977] Proc. Natl. Acad. Sci. USA 74, 5463-5467) of
pBS1.8 was performed with "SEQUENASE" (United States
Biochemical) on both strands by using specific
10 oligonucleotide primers synthesized by the Advanced DNA
Technologies Laboratory (Biology Department, Texas A & M
University). DNA sequence analysis was done with the
GCG (Madison, WI) software as described by Devereux et
al. (1984) Nucleic Acids Res. 12, 387-395.

15 Both NheI/HindIII subfragments were transferred
into a conjugal expression vector, AM542, in both
forward and reverse orientations with respect to a
cyanobacterial carboxylase promoter and were introduced
into Anabaena by conjugation. Transconjugants
20 containing the 1.8 kb fragment in the forward
orientation (AM542-1.8F) produced significant quantities
of GLA and octadecatetraenoic acid (Figure 2; Table 2).
Transconjugants containing other constructs, either
reverse oriented 1.8 kb fragment or forward and reverse
25 oriented 1.7 kb fragment, did not produce detectable
levels of GLA (Table 2).

Figure 2 compares the C18 fatty acid profile of
an extract from wild type Anabaena (Figure 2A) with that
of transgenic Anabaena containing the 1.8 kb fragment of
30 cSy75-3.5 in the forward orientation (Figure 2B). GLC
analysis of fatty acid methyl esters from AM542-1.8F
revealed a peak with a retention time identical to that

35

-22-

1 of authentic GLA standard. Analysis of this peak by gas
chromatography-mass spectrometry (GC-MS) confirmed that
it had the same mass fragmentation pattern as a GLA
reference sample. Transgenic Anabaena with altered
5 levels of polyunsaturated fatty acids were similar to
wild type in growth rate and morphology.

10

15

20

25

30

35

1

Table 2
Composition of C18 Fatty Acids in
Wild Type and Trasgenic Cyanobacteria

5

Strain	Fatty acid (%)					
	18:0	18:1	18:2	18:3 (α)	18:3 (γ)	18:4
Wild type						
Synechocystis (sp.PCC6803)	13.6	4.5	54.5	-	27.3	-
Anabaena (sp.PCC7120)	2.9	24.8	37.1	35.2	-	-
Synechococcus (Sp.PCC7942)	20.6	79.4	-	-	-	-
Anabaena Transconjugants						
cSy75	3.8	24.4	22.3	9.1	27.9	12.5
cSy75-3.5	4.3	27.6	18.1	3.2	40.4	6.4
pAM542-1.8F	4.2	13.9	12.1	19.1	25.4	25.4
pAM542-1.8R	7.7	23.1	38.4	30.8	-	-
pAM542-1.7F	2.8	27.8	36.1	33.3	-	-
pAM542-1.7R	2.8	25.4	42.3	29.6	-	-
Synechococcus Transformants						
pAM854	27.8	72.2	-	-	-	-
pAM854-Δ ¹²	4.0	43.2	46.0	-	-	-
pAM854-Δ ⁶	18.2	81.8	-	-	-	-
pAM854-Δ ⁶ & Δ ¹²	42.7	25.3	19.5	-	16.5	-

10

15

20

25

30

18:0, stearic acid; 18:1, oleic acid; 18:2, linoleic acid; 18:3(α), α-linolenic acid; 18:3(γ), γ-linolenic acid; 18:4, octadecatetraenoic acid

35

1

EXAMPLE 4

Transformation of Synechococcus
with $\Delta 6$ and $\Delta 12$ Desaturase Genes

5 A third cosmid, cSy7, which contains a $\Delta 12$ -
desaturase gene, was isolated by screening the
Synechocystis genomic library with a oligonucleotide
synthesized from the published Synechocystis $\Delta 12$ -
desaturase gene sequence (Wada et al. [1990] Nature
10 (London) 347, 200-203). A 1.7 kb AvaI fragment from
this cosmid containing the $\Delta 12$ -desaturase gene was
identified and used as a probe to demonstrate that cSy13
not only contains a $\Delta 6$ -desaturase gene but also a $\Delta 12$ -
desaturase gene (Figure 3). Genomic Southern blot
analysis further showed that both the $\Delta 6$ -and $\Delta 12$ -
15 desaturase genes are unique in the Synechocystis genome
so that both functional genes involved in C18 fatty acid
desaturation are linked closely in the Synechocystis
genome.

20 The unicellular cyanobacterium Synechococcus (PCC
7942) is deficient in both linoleic acid and GLA(3).
The $\Delta 12$ and $\Delta 6$ -desaturase genes were cloned individually
and together into pAM854 (Bustos et al. [1991] J.
Bacteriol. 174, 7525-7533), a shuttle vector that
contains sequences necessary for the integration of
25 foreign DNA into the genome of Synechococcus (Golden et
al. [1987] Methods in Enzymol. 153, 215-231).
Synechococcus was transformed with these gene constructs
and colonies were selected. Fatty acid methyl esters
were extracted from transgenic Synechococcus and
30 analyzed by GLC.

Table 2 shows that the principal fatty acids of
wild type Synechococcus are stearic acid (18:0) and

35

SUBSTITUTE SHEET

-25-

1 oleic acid (18:1). Synechococcus transformed with
pAM854- Δ 12 expressed linoleic acid (18:2) in addition to
the principal fatty acids. Transformants with pAM854- Δ 6
and Δ 12 produced both linoleate and GLA (Table 1).
5 These results indicated that Synechococcus containing
both Δ 12- and Δ 6-desaturase genes has gained the
capability of introducing a second double bond at the
 Δ 12 position and a third double bond at the Δ 6 position
of C18 fatty acids. However, no changes in fatty acid
10 composition was observed in the transformant containing
pAM854- Δ 6, indicating that in the absence of substrate
synthesized by the Δ 12 desaturase, the Δ 6-desaturase is
inactive. This experiment further confirms that the 1.8
kb NheI/HindIII fragment (Figure 3) contains both coding
15 and promoter regions of the Synechocystis Δ 6-desaturase
gene. Transgenic Synechococcus with altered levels of
polyunsaturated fatty acids were similar to wild type in
growth rate and morphology.

20

25

30

35

1

EXAMPLE 5**Nucleotide Sequence of $\Delta 6$ -Desaturase**

The nucleotide sequence of the 1.8 kb fragment of cSy75-3.5 including the functional $\Delta 6$ -desaturase gene was determined. An open reading frame encoding a polypeptide of 359 amino acids was identified (Figure 4). A Kyte-Doolittle hydrophathy analysis (Kyte et al. [1982] J. Mol. Biol. 157, 105-132) identified two regions of hydrophobic amino acids that could represent transmembrane domains (Figure 1A); furthermore, the hydrophathic profile of the $\Delta 6$ -desaturase is similar to that of the $\Delta 12$ -desaturase gene (Figure 1B; Wada et al.) and $\Delta 9$ -desaturases (Thiede et al. [1986] J. Biol. Chem. 261, 13230-13235). However, the sequence similarity between the Synechocystis $\Delta 6$ - and $\Delta 12$ -desaturases is less than 40% at the nucleotide level and approximately 18% at the amino acid level.

20

25

30

35

1

EXAMPLE 6

Transfer of Cyanobacterial Δ^6 -Desaturase into Tobacco

The cyanobacterial Δ^6 -desaturase gene was mobilized into a plant expression vector and transferred to tobacco using Agrobacterium mediated gene transfer techniques. To ensure that the transferred desaturase is appropriately expressed in leaves and developing seeds and that the desaturase gene product is targeted to the endoplasmic reticulum or the chloroplast, various expression cassettes with Synechocystis Δ^6 -desaturase open reading frame (ORF) were constructed. Components of these cassettes include: (i) a 35S promoter or seed specific promoter derived from the sunflower helianthinin gene to drive Δ^6 -desaturase gene expression in all plant tissues or only in developing seeds respectively, (ii) a putative signal peptide either from carrot extensin gene or sunflower helianthinin gene to target newly synthesized Δ^6 -desaturase into the ER, (iii) an ER lumen retention signal sequence (KDEL) at the COOH-terminal of the Δ^6 -desaturase ORF, and (iv) an optimized transit peptide to target Δ^6 desaturase into the chloroplast. The 35S promoter is a derivative of pRTL2 described by Restrepo et al. (1990). The optimized transit peptide sequence is described by Van de Broeck et al. (1985). The carrot extensin signal peptide is described by Chen et al. (1985) EMBO J. 9, 2145.

Transgenic tobacco plants were produced containing a chimeric cyanobacterial desaturase gene, comprised of the Synechocystis Δ^6 desaturase gene fused to an endoplasmic reticulum retention sequence (KDEL) and extensin signal peptide driven by the CaMV 35S promoter. PCR amplifications of transgenic tobacco genomic DNA indicate that the Δ^6 desaturase gene was incorporated into the tobacco genome. Fatty acid methyl esters of leaves of these transgenic tobacco plants were

-28-

1 extracted and analyzed by Gas Liquid Chromatography
(GLC). These transgenic tobacco accumulated significant
amounts of GLA (Figure 4). Figure 4 shows fatty acid
methyl esters as determined by GLC. Peaks were
5 identified by comparing the elution times with known
standards of fatty acid methyl ester. Accordingly,
cyanobacterial genes involved in fatty acid metabolism
can be used to generate transgenic plants with altered
fatty acid compositions.

10

15

20

25

30

35

SUBSTITUTE SHEET

-29-

1

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Thomas, Terry L.
Reddy, Avutu S.
Nuccio, Michael
Freyssinet, Georges L.

(ii) TITLE OF INVENTION: PRODUCTION OF GAMMA LINOLENIC
ACID BY A DELTA 6-DESATURASE

10

(iii) NUMBER OF SEQUENCES: 3

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Scully, Scott, Murphy & Presser
(B) STREET: 400 Garden City Plaza
(C) CITY: Garden City
(D) STATE: New York
(E) COUNTRY: United States
(F) ZIP: 11530

15

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

20

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: To be assigned
(B) FILING DATE: 08-JAN-1992
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

25

(A) NAME: McNulty, William E.
(B) REGISTRATION NUMBER: 22,606
(C) REFERENCE/DOCKET NUMBER: 8383Z

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (516) 742-4343
(B) TELEFAX: (516) 742-4366
(C) TELEX: 230 901 SANS UR

30

35

1

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 3588 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 2002..3081

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10 GCTAGCCACC AGTGACGATG CCTTGAATTT GGCCATTCTG ACCCAGGCC GTATTCTGAA 60
 TCCCCGCATT CGCATTGTTA ATCGTTTGTT CAACCATGCC CTGGGTAAAC GTTTAGACAC 120
 CACCTTGCCA GACCACGTTA GTTTGAGTGT TTCGCCCTG GCGGCCCGA TTTTTCCTT 180
 TGCGGCTTTG GGCAATCAGG CGATCGGCA ATTGCGTTTG TTTGACCAGA CTGGGCCCAT 240
 TCAGGAAATT GTCATTCACC AAGACCATCC CTGGCTCAAT TTACCCCTGG CGGATTTATG 300
 15 GGATGATCCG AGCCGAATGT TGATCTATTA CCTACCGGCC CACAGTGAAA CGGATTTAGT 360
 AGGCGCAGTG GTGAATAATT TAACGTTGCA ATCTGGGGAC CATTTAATAG TGGGACAAAA 420
 ACCCCAACCC AAGACCAAAC GGCATCGCC TTGGCGCAA TTTTCCAAAC TGATTACCAA 480
 CCTGCGGGAG TATCAGCGGT ATGTCCAACA GGTGATATGG GTGGTGTGT TTTTATTGTT 540
 GATGATTTT CTGGCCACCT TCATCTACGT TTCCATTGAT CAACATATTG CCCCAGTGGA 600
 20 CGCGTTGTAT TTTTCCGTGG GCATGATTAC CGGGGCCGGT GGCAAGGAAG AGGTGGCCGA 660
 AAAGTCCCC GATATCATCA AAGTATTCAC AGTGGTGATG ATGATCGCCG GGGCGGGGT 720
 GATTGGTATT TGTTATGCC TACTGAATGA TTTCATCCTT GGCAGTCGCT TTAGTCAGTT 780
 TTTGGATGCG GCCAAGTTAC CCGATCGCCA TCACATCATC ATTTGTGGC TGGGGGAGT 840
 25 GAGCATGGCC ATTATGAAG AGTTAATCA CCAGGGCCAT GAAATGTGG TAATCGAAAA 900
 GGATACAGAT AATCGTTTCT TGCATACGGC CCGCTCCCTG GGGGTGCCCC TAATTGTGGA 960

30

35

1 GGATGCCCGC CTAGAAAGAA CGTTGGCCTG CGCCAATATC AACCGAGCCG AAGCCATTGT 1020
 GGTGGCCACC AGCGACGACA CCGTTAACTT GGAAATTGGC CTAAGTGCCA AGGCGATCGC 1080
 CCCTAGCCTG CCAGTGGTGT TGCCTTGCCA GGATGCCCAG TTTAGCCTGT CCCTGCAGGA 1140
 AGTATTTGAA TTTGAAACGG TGCTTTGTCC GCGGGAATTG GCCACCTATT CCTTTGCGGC 1200
 5 GCGGGCCCTG GGGGGCAAAA TTTTGGGCAA CGGCATGACC GATGATTTGC TGTGGGTAGC 1260
 CCTAGCCACC TTAATCACTC CTAACCATCC CTTTGCCGAC CAATTGGTTA AAATTGCAGC 1320
 CCAAAAGTCT GATTTTCGTTT CCTCTATCT AGAACGGGGT GGCAAAACCA TCCATAGCTG 1380
 GGAATTATTG GGTACCCATC TCGACTCTGG AGACGTGTTG TATTTAACCA TGCCCCCCAC 1440
 10 TGCCCTAGAG CAACTTTGGC GATCGCCCCG TGCCACTGCT GATCCTCTGG ACTCTTTTTT 1500
 GGTTTAGCAT GGGGGGATGG AACTCTTGAC TCGGCCCAAT GGTGATCAAG AAAGAACGCT 1560
 TTGTCTATGT TTAGTATTTT TAAGTTAACC AACAGCAGAG GATAACTTCC AAAAGAAATT 1620
 AAGCTCAAAA AGTAGCAAAA TAAGTTTAAAT TCATAACTGA GTTTTACTGC TAAACAGCGG 1680
 TGCAAAAAAG TCAGATAAAA TAAAAGCTTC ACTTCGGTTT TATATTGTGA CCATGGTTCC 1740
 15 CAGGCATCTG CTCTAGGGAG TTTTTCGGCT GCCTTTAGAG AGTATTTTCT CCAAGTCGGC 1800
 TAACTCCCCC ATTTTITAGGC AAAATCATAT ACAGACTATC CCAATATTGC CAGAGCTTTG 1860
 ATGACTCACT GTAGAAGGCA GACTAAAATT CTAGCAATGG ACTCCCAGTT GGAATAAATT 1920
 TTTAGTCTCC CCCGGCGCTG GAGTTTTTTT GTAGTTAATG GCGGTATAAT GTGAAAGTTT 1980
 TTTATCTATT TAAATTTATA A ATG CTA ACA GCG GAA AGA ATT AAA TTT ACC 2031
 20 Met Leu Thr Ala Glu Arg Ile Lys Phe Thr
 1 5 10
 CAG AAA CGG GGG TTT CGT CGG GTA CTA AAC CAA CGG GTG GAT GCC TAC 2079
 Gln Lys Arg Gly Phe Arg Arg Val Leu Asn Gln Arg Val Asp Ala Tyr
 15 20 25
 TTT GCC GAG CAT GGC CTG ACC CAA AGG GAT AAT CCC TCC ATG TAT CTG 2127
 Phe Ala Glu His Gly Leu Thr Gln Arg Asp Asn Pro Ser Met Tyr Leu
 30 35 40
 25
 30
 35

1	AAA ACC CTG ATT ATT GTG CTC TGG TTG TTT TCC GCT TGG GCC TTT GTG	2175
	Lys Thr Leu Ile Ile Val Leu Trp Leu Phe Ser Ala Trp Ala Phe Val	
	45 50 55	
	CTT TTT GCT CCA GTT ATT TTT CCG GTG CGC CTA CTG GGT TGT ATG GTT	2223
	Leu Phe Ala Pro Val Ile Phe Pro Val Arg Leu Leu Gly Cys Met Val	
	60 65 70	
5	TTG GCG ATC GCC TTG GCG GCC TTT TCC TTC AAT GTC GGC CAC GAT GCC	2271
	Leu Ala Ile Ala Leu Ala Ala Phe Ser Phe Asn Val Gly His Asp Ala	
	75 80 85 90	
	AAC CAC AAT GCC TAT TCC TCC AAT CCC CAC ATC AAC CGG GTT CTG GGC	2319
	Asn His Asn Ala Tyr Ser Ser Asn Pro His Ile Asn Arg Val Leu Gly	
	95 100 105	
10	ATG ACC TAC GAT TTT GTC GGG TTA TCT AGT TTT CTT TGG CGC TAT CGC	2367
	Met Thr Tyr Asp Phe Val Gly Leu Ser Ser Phe Leu Trp Arg Tyr Arg	
	110 115 120	
	CAC AAC TAT TTG CAC CAC ACC TAC ACC AAT ATT CTT GGC CAT GAC GTG	2415
	His Asn Tyr Leu His His Thr Tyr Thr Asn Ile Leu Gly His Asp Val	
	125 130 135	
	GAA ATC CAT GGA GAT GGC GCA GTA CGT ATG AGT CCT GAA CAA GAA CAT	2463
	Glu Ile His Gly Asp Gly Ala Val Arg Met Ser Pro Glu Gln Glu His	
	140 145 150	
15	GTT GGT ATT TAT CGT TTC CAG CAA TTT TAT ATT TGG GGT TTA TAT CTT	2511
	Val Gly Ile Tyr Arg Phe Gln Gln Phe Tyr Ile Trp Gly Leu Tyr Leu	
	155 160 165 170	
	TTC ATT CCC TTT TAT TGG TTT CTC TAC GAT GTC TAC CTA GTG CTT AAT	2559
	Phe Ile Pro Phe Tyr Trp Phe Leu Tyr Asp Val Tyr Leu Val Leu Asn	
	175 180 185	
20	AAA GGC AAA TAT CAC GAC CAT AAA ATT CCT CCT TTC CAG CCC CTA GAA	2607
	Lys Gly Lys Tyr His Asp His Lys Ile Pro Pro Phe Gln Pro Leu Glu	
	190 195 200	
	TTA GCT AGT TTG CTA GGG ATT AAG CTA TTA TGG CTC GGC TAC GTT TTC	2655
	Leu Ala Ser Leu Leu Gly Ile Lys Leu Leu Trp Leu Gly Tyr Val Phe	
	205 210 215	
	GGC TTA CCT CTG GCT CTG GGC TTT TCC ATT CCT GAA GTA TTA ATT GGT	2703
25	Gly Leu Pro Leu Ala Leu Gly Phe Ser Ile Pro Glu Val Leu Ile Gly	
	220 225 230	

30

35

1	GCT TCG GTA ACC TAT ATG ACC TAT GGC ATC GTG GTT TGC ACC ATC TTT Ala Ser Val Thr Tyr Met Thr Tyr Gly Ile Val Val Cys Thr Ile Phe 235 240 245 250	2751
	ATG CTG GCC CAT GTG TTG GAA TCA ACT GAA TTT CTC ACC CCC GAT GGT Met Leu Ala His Val Leu Glu Ser Thr Glu Phe Leu Thr Pro Asp Gly 255 260 265	2799
5	GAA TCC GGT GCC ATT GAT GAC GAG TGG GCT ATT TGC CAA ATT CGT ACC Glu Ser Gly Ala Ile Asp Asp Glu Trp Ala Ile Cys Gln Ile Arg Thr 270 275 280	2847
	ACG GCC AAT TTT GCC ACC AAT AAT CCC TTT TGG AAC TGG TTT TGT GGC Thr Ala Asn Phe Ala Thr Asn Asn Pro Phe Trp Asn Trp Phe Cys Gly 285 290 295	2895
10	GGT TTA AAT CAC CAA GTT ACC CAC CAT CTT TTC CCC AAT ATT TGT CAT Gly Leu Asn His Gln Val Thr His His Leu Phe Pro Asn Ile Cys His 300 305 310	2943
	ATT CAC TAT CCC CAA TTG GAA AAT ATT ATT AAG GAT GTT TGC CAA GAG Ile His Tyr Pro Gln Leu Glu Asn Ile Ile Lys Asp Val Cys Gln Glu 315 320 325 330	2991
15	TTT GGT GTG GAA TAT AAA GTT TAT CCC ACC TTC AAA GCG GCG ATC GCC Phe Gly Val Glu Tyr Lys Val Tyr Pro Thr Phe Lys Ala Ala Ile Ala 335 340 345	3039
	TCT AAC TAT CGC TGG CTA GAG GCC ATG GGC AAA GCA TCG TGACATTGCC Ser Asn Tyr Arg Trp Leu Glu Ala Met Gly Lys Ala Ser 350 355 360	3088
	TTGGGATTGA AGCAAAATGG CAAAATCCCT CGTAAATCTA TGATCGAAGC CTTTCTGTTG	3148
20	CCCGCCGACC AAATCCCCGA TGCTGACCAA AGGTTGATGT TGGCATTGCT CCAAACCCAC	3208
	TTTGAGGGGG TTCATTGGCC GCAGTTTCAA GCTGACCTAG GAGGCAAAGA TTGGGTGATT	3268
	TTGCTCAAAT CCGCTGGGAT ATTGAAAGGC TTCACCACCT TTGGTTTCTA CCCTGCTCAA	3328
	TGGGAAGGAC AAACCGTCAG AATTGTTTAT TCTGGTGACA CCATCACCGA CCCATCCATG	3388
	TGGTCTAACC CAGCCCTGGC CAAGGCTTGG AÇCAAGGCCA TGCAAATCTT CCACGAGGCT	3448
25	AGGCCAGAAA AATTATATTG GCTCCTGATT TCTTCCGGCT ATCGCACCTA CCGATTTTTG	3508
	AGCATTTTTG CCAAGGAATT CTATCCCCAC TATCTCCATC CCACTCCCCC GCCTGTACAA	3568
30		
35		

1 AATTTTATCC ATCAGCTAGC 3588

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 359 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Phe	Thr	Gln	Lys	Arg	Gly	Phe	Arg
	1				5					10					15	
10	Arg	Val	Leu	Asn	Gln	Arg	Val	Asp	Ala	Tyr	Phe	Ala	Glu	His	Gly	Leu
			20						25					30		
	Thr	Gln	Arg	Asp	Asn	Pro	Ser	Met	Tyr	Leu	Lys	Thr	Leu	Ile	Ile	Val
			35					40					45			
	Leu	Trp	Leu	Phe	Ser	Ala	Trp	Ala	Phe	Val	Leu	Phe	Ala	Pro	Val	Ile
		50					55				60					
15	Phe	Pro	Val	Arg	Leu	Leu	Gly	Cys	Met	Val	Leu	Ala	Ile	Ala	Leu	Ala
	65				70						75				80	
	Ala	Phe	Ser	Phe	Asn	Val	Gly	His	Asp	Ala	Asn	His	Asn	Ala	Tyr	Ser
				85					90					95		
	Ser	Asn	Pro	His	Ile	Asn	Arg	Val	Leu	Gly	Met	Thr	Tyr	Asp	Phe	Val
				100					105					110		
20	Gly	Leu	Ser	Ser	Phe	Leu	Trp	Arg	Tyr	Arg	His	Asn	Tyr	Leu	His	His
			115					120					125			
	Thr	Tyr	Thr	Asn	Ile	Leu	Gly	His	Asp	Val	Glu	Ile	His	Gly	Asp	Gly
		130					135					140				
	Ala	Val	Arg	Met	Ser	Pro	Glu	Gln	Glu	His	Val	Gly	Ile	Tyr	Arg	Phe
	145					150					155					160
	Gln	Gln	Phe	Tyr	Ile	Trp	Gly	Leu	Tyr	Leu	Phe	Ile	Pro	Phe	Tyr	Trp
25					165					170					175	

30

35

1 Phe Leu Tyr Asp Val Tyr Leu Val Leu Asn Lys Gly Lys Tyr His Asp
 180 185 190
 His Lys Ile Pro Pro Phe Gln Pro Leu Glu Leu Ala Ser Leu Leu Gly
 195 200 205
 5 Ile Lys Leu Leu Trp Leu Gly Tyr Val Phe Gly Leu Pro Leu Ala Leu
 210 215 220
 Gly Phe Ser Ile Pro Glu Val Leu Ile Gly Ala Ser Val Thr Tyr Met
 225 230 235 240
 Thr Tyr Gly Ile Val Val Cys Thr Ile Phe Met Leu Ala His Val Leu
 245 250 255
 10 Glu Ser Thr Glu Phe Leu Thr Pro Asp Gly Glu Ser Gly Ala Ile Asp
 260 265 270
 Asp Glu Trp Ala Ile Cys Gln Ile Arg Thr Thr Ala Asn Phe Ala Thr
 275 280 285
 Asn Asn Pro Phe Trp Asn Trp Phe Cys Gly Gly Leu Asn His Gln Val
 290 295 300
 15 Thr His His Leu Phe Pro Asn Ile Cys His Ile His Tyr Pro Gln Leu
 305 310 315 320
 Glu Asn Ile Ile Lys Asp Val Cys Gln Glu Phe Gly Val Glu Tyr Lys
 325 330 335
 Val Tyr Pro Thr Phe Lys Ala Ala Ile Ala Ser Asn Tyr Arg Trp Leu
 340 345 350
 Glu Ala Met Gly Lys Ala Ser
 355

20

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1884 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

30

35

1	AGCTTCACTT CGGTTTTATA TTGTGACCAT GGTTCCCAGG CATCTGCTCT AGGGAGTTTT	60
	TCCGCTGCCT TTAGAGAGTA TTTTCTCCAA GTCGGCTAAC TCCCCCATTT TTAGGCAAAA	120
	TCATATACAG ACTATCCCAA TATTGCCAGA GCTTTGATGA CTCACTGTAG AAGGCAGACT	180
5	AAAATICTAG CAATGGACTC CCAGTTGGAA TAAATTTTTA GTCTCCCCCG GCGCTGGAGT	240
	TTTTTTGTAG TTAATGGCGG TATAATGTGA AAGTTTTTTA TCTATTTAAA TTTATAAATG	300
	CTAACAGCGG AAAGAATTAA ATTTACCCAG AAACGGGGGT TTCGTCGGGT ACTAAACCAA	360
	CGGGTGGATG CCTACTTTGC CGAGCATGGC CTGACCCAAA GGGATAATCC CTCCATGTAT	420
	CTGAAAACCC TGATTATTGT GCTCTGGTTG TTTTCCGCTT GGGCCTTTGT GCTTTTTGCT	480
10	CCAGTTATTT TTCCGGTGGC CCTACTGGGT TGTATGGTTT TGGCGATCGC CTTGGCGGCC	540
	TTTTCCCTCA ATGTCGGCCA CGATGCCAAC CACAATGCCT ATTCCTCCAA TCCCCACATC	600
	AACCGGGTTC TGGGCATGAC CTACGATTTT GTCGGGTTAT CTAGTTTTCT TTGGCGCTAT	660
	CGCCACAACCT ATTTGCACCA CACCTACACC AATATTCTTG GCCATGACGT GGAAATCCAT	720
	GGAGATGGCG CAGTACGTAT GAGTCCTGAA CAAGAACATG TTGGTATTTA TCGTTTCCAG	780
15	CAATTTTATA TTTGGGGTTT ATATCTTTTC ATTCCTTTT ATTTGGTTTCT CTACGATGTC	840
	TACCTAGTGC TTAATAAAGG CAAATATCAC GACCATAAAA TTCCTCCTTT CCAGCCCCTA	900
	GAATTAGCTA GTTTGCTAGG GATTAAGCTA TTATGGCTCG GCTACGTTTT CGGCTTACCT	960
	CTGGCTCTGG GCTTTTCCAT TCCTGAAGTA TTAATTGGTG CTTCGGTAAC CTATATGACC	1020
	TATGGCATCG TGGTTTGCAC CATCTTTATG CTGGCCCATG TGTGGAATC AACTGAATTT	1080
20	CTCACCCCG ATGGTGAATC CGGTGCCATT GATGACGAGT GGGCTATTTG CCAAATTCGT	1140
	ACCACGGCCA ATTTTGCAC CAATAATCCC TTTTGGAACT GTTTTTGTGG CGGTTTAAAT	1200
	CACCAAGTTA CCCACCATCT TTTCCCAAT ATTTGTCATA TTCACTATCC CCAATTGGAA	1260
	AATATTATTA AGGATGTTTG CCAAGAGTTT GGTGTGGAAT ATAAAGTTTA TCCCACCTTC	1320
25	AAAGCGGCGA TCGCCTCTAA CTATCGCTGG CTAGAGGCCA TGGGCAAAGC ATCGTGACAT	1380
	TGCCTTGGGA TTGAAGCAA ATGGCAAAT CCCTCGTAAA TCTATGATCG AAGCCTTTCT	1440

30

35

-37-

1 GTTGCCCGCC GACCAAATCC CCGATGCTGA CCAAAGGTTG ATGTTGGCAT TGCTCCAAAC 1500
CCACTTTGAG GGGGTTTCATT GGCCGCAGTT TCAAGCTGAC CTAGGAGGCA AAGATTGGGT 1560
GATTTTGCTC AAATCCGCTG GGATATTGAA AGGCTTCACC ACCTTTGGTT TCTACCCTGC 1620
TCAATGGGAA GGACAAACCG TCAGAATTGT TTATTCTGGT GACACCATCA CCGACCCATC 1680
5 CATGTGGTCT AACCCAGCCC TGGCCAAGGC TTGGACCAAG GCCATGCAAA TTCTCCACGA 1740
GGCTAGGCCA GAAAAATTAT ATTGGCTCCT GATTTCTTCC GGCTATCGCA CCTACCGATT 1800
TTTGAGCATT TTTGCCAAGG AATTCTATCC CCACTATCTC CATCCCACTC CCCC GCCTGT 1860
ACAAAATTTT ATCCATCAGC TAGC 1884

10

15

20

25

30

35

1 WHAT IS CLAIMED:

1. An isolated nucleic acid encoding bacterial $\Delta 6$ -desaturase.
2. The nucleic acid of Claim 1 comprising the
5 nucleotides of SEQ. ID NO:3.
3. An isolated nucleic acid that codes for the amino acid sequence encoded by the nucleic acid of Claim 1.
4. The isolated nucleic acid of any one of Claims 1-3 wherein said nucleic acid is contained in a vector.
- 10 5. The isolated nucleic acid of Claim 4 operably linked to a promoter and/or a termination signal capable of effecting expression of the gene product of said isolated nucleic acid.
6. The isolated nucleic acid of Claim 5 wherein said
15 promoter is a $\Delta 6$ -desaturase promoter, an Anabaena carboxylase promoter, a helianthinin promoter, a glycin promoter, a napin promoter, or a helianthinin tissue-specific promoter.
7. The isolated nucleic acid of Claim 5 wherein said
20 termination signal is a Synechocystis termination signal, a nopaline synthase termination signal, or a seed termination signal.
8. The isolated nucleic acid of any one of Claims 1-7 wherein said isolated nucleic acid is contained within a transgenic organism.
- 25 9. The isolated nucleic acid of Claim 8 wherein said transgenic organism is a bacterium, a fungus, a plant cell or an animal.
10. A plant or progeny of said plant which has been regenerated from the transgenic plant cell of Claim 9.
- 30 11. The plant of Claim 10 wherein said plant is a sunflower, soybean, maize, tobacco, peanut or oil seed rape plant.

1 12. A method of producing a plant with increased
gamma linolenic acid (GLA) content which comprises:

(a) transforming a plant cell with the isolated
nucleic acid of any one of Claims 1-7; and

5 (b) regenerating a plant with increased GLA content
from said plant cell.

13. The method of Claim 12 wherein said plant is a
sunflower, soybean, maize, tobacco, peanut or oil seed rape
plant.

10 14. A method of inducing production of gamma
linolenic acid (GLA) in an organism deficient or lacking in
GLA with comprises transforming said organism with the
isolated nucleic acid of any one of Claims 1-7.

15 15. A method of inducing production of gamma
linolenic acid (GLA) in an organism deficient or lacking in
GLA and linoleic acid (LA) which comprises transforming said
organism with an isolated nucleic acid encoding bacterial $\Delta 6$ -
desaturase and an isolated nucleic acid encoding $\Delta 12$ -
desaturase.

20 16. A method of inducing production of gamma
linolenic acid (GLA) in an organism deficient or lacking in
GLA and linoleic acid (LA) which comprises transforming said
organism with at least one expression vector comprising an
isolated nucleic acid encoding bacterial $\Delta 6$ -desaturase and an
25 isolated nucleic acid encoding $\Delta 12$ -desaturase.

17. The method of any one of Claims 15 or 16 wherein
said isolated nucleic acid encoding $\Delta 6$ -desaturase comprises
nucleotides 317 to 1507 of SEQ. ID NO:1.

30 18. A method of inducing production of
octadecatetraenoic acid in an organism deficient or lacking
in gamma linolenic acid with comprises transforming said
organism with isolated nucleic acid of any one of Claims 1-7.

1 19. The method of Claim 18 wherein said organism is a
bacterium, a fungus, a plant or an animal.

 20. A method of use of the isolated nucleic acid of
any one of Claims 1-7 to produce a plant with improved
5 chilling resistance which comprises:

 a) transforming a plant cell with the isolated
nucleic acid of any one of Claims 1-7; and

 b) regenerating said plant with improved chilling
resistance from said transformed plant cell.

10 21. The method of Claim 20 wherein said plant is a
sunflower, soybean, maize, tobacco, peanut or oil seed rape
plant.

 22. Isolated bacterial $\Delta 6$ -desaturase.

 23. The isolated bacterial $\Delta 6$ -desaturase of Claim 22
15 which has an amino acid sequence of SEQ ID NO:2.

20

25

30

35




FIG. IA






FIG. IB






FIG. 4A




FIG. 4B

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US92/08746

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : Please See Extra Sheet.

US CL : 800/205; 435/172.3, 189, 69.1, 320.1, 134, 170, 171; 536/27

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 800/205; 435/172.3, 189, 69.1, 320.1, 134, 170, 171; 536/27;
935/9, 30, 6, 24, 29, 38

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN/BIOSIS, CA; APS

search terms: linolenic, desaturase, delta-6, gene, DNA, cDNA,
purif?, cyanobacteri?,


C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Nature, Volume 347, issued 13 September 1990, H. Wada et al., "Enhancement of Chilling Tolerance of a Cyanobacterium by Genetic Manipulation of Fatty Acid Desaturation", pages 200-203, especially pages 201-203.	1-23
Y	Biochemical Journal, Volume 240, issued 1986, S. Stymne et al., "Biosynthesis of γ -Linolenic Acid in Cotyledons and Microsomal Preparations of the Developing Seeds of Common Borage (<i>Borago officinalis</i>)", pages 385-392, especially pages 385 and 392.	1-23
Y	EP, A, 0,255, 378 (Kridl et al.) 3 February 1988, see entire document, especially columns 3-5 and 7-11.	1-23

Further documents are listed in the continuation of Box C. See patent family annex.

<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be part of particular relevance</p> <p>"E" earlier document published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Date of the actual completion of the international search 03 DECEMBER 1992	Date of mailing of the international search report 13 JAN 1993
--------------------------------------------------------------------------------------	--------------------------------------------------------------------------

Name and mailing address of the ISA/ Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer  CHARLES C. P. RORIES, PH.D.
Facsimile No. NOT APPLICABLE	Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US92/08746

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (5):

A01H 1/00, 5/00; C12N 15/00, 9/02; C12P 7/64, 1/02, 1/04, 21/06; C07H 15/12, 17/00