

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
16 December 2010 (16.12.2010)

PCT

(10) International Publication Number  
**WO 2010/142141 A1**

(51) International Patent Classification:  
*A61K 38/18* (2006.01) *A61P 9/04* (2006.01)

(21) International Application Number:  
PCT/CN2010/000845

(22) International Filing Date:  
12 June 2010 (12.06.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
200910057390.5 9 June 2009 (09.06.2009) CN  
61/236,849 25 August 2009 (25.08.2009) US

(71) Applicant (for all designated States except US): **ZEN-SUN (SHANGHAI) SCIENCE & TECHNOLOGY LIMITED** [CN/CN]; No.68 Ju Li Road, Zhangjiang Hi-Tech Park, Pudong, Shanghai 201203 (CN).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ZHOU, Mingdong** [AU/CN]; No.68 Ju Li Road, Zhangjiang Hi-Tech Park, Pudong, Shanghai 201203 (CN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,

KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

**Published:**

- with international search report (Art. 21(3))
- the filing date of the international application is within two months from the date of expiration of the priority period (Rule 26bis.3)
- with sequence listing part of description (Rule 5.2(a))



**WO 2010/142141 A1**

(54) Title: NEUREGULIN BASED METHODS FOR TREATING HEART FAILURE

(57) Abstract: The present invention features methods of treating patients with chronic heart failure by administering a neuregulin polypeptide within a dosage range which is both effective and safe.

# NEUREGULIN BASED METHODS FOR TREATING HEART FAILURE

## 1. FIELD OF THE INVENTION

[0001] The present invention relates to methods for preventing, treating or delaying heart failure in a mammal using a neuregulin protein. Particularly, the present invention provides the dosage range of neuregulin protein for preventing, treating or delaying heart failure in the mammal.

## 2. BACKGROUND OF THE INVENTION

[0002] Heart failure affects approximately five million Americans, and more than 550,000 new patients are diagnosed with the condition each year. Current drug therapy for heart failure is primarily directed to angiotensin-converting enzyme (ACE) inhibitors, which are vasodilators that cause blood vessels to expand, lowering blood pressure and reducing the heart's workload. While the percent reduction in mortality has been significant, the actual reduction in mortality with ACE inhibitors has averaged only 3%-4%, and there are several potential side effects. Additional limitations are associated with other options for preventing or treating heart failure. For example, heart transplantation is clearly more expensive and invasive than drug treatment, and it is further limited by the availability of donor hearts. Use of mechanical devices, such as biventricular pacemakers, is similarly invasive and expensive. Thus, there has been a need for new therapies given the deficiencies in current therapies.

[0003] One promising new therapy involves administration of neuregulin (hereinafter referred to as "NRG") to a patient suffering from or at risk of developing

heart failure. NRGs, a family of EGF-like growth factors, comprises a family of structurally related growth and differentiation factors that include NRG1, NRG2, NRG3 and NRG4 and isoforms thereof, are involved in an array of biological responses: stimulation of breast cancer cell differentiation and secretion of milk proteins; induction of neural crest cell differentiation to Schwann cells; stimulation of skeletal muscle cell synthesis of acetylcholine receptors; and, promotion of myocardial cell survival and DNA synthesis. In vivo studies of neuregulin gene-targeted homozygous mouse embryos with severe defects in ventricular trabeculae formation and dorsal root ganglia development indicate that neuregulin is essential for heart and neural development.

**[0004]** NRGs bind to the EGF receptor family, which comprises EGFR, ErbB2, ErbB3 and ErbB4, each of which plays an important role in multiple cellular functions, including cell growth, differentiation and survival. They are protein tyrosine kinase receptors, consisting of an extracellular ligand-binding domain, transmembrane kinase domain and cytoplasmic tyrosine kinase domain. After NRG bind to the extracellular domain of ErbB3 or ErbB4, it induces a conformational change that leads to heterodimer formation between ErbB3, ErbB4 and ErbB2 or homodimer formation between ErbB4 itself, which results in phosphorylation of the receptor's C-terminal domain inside the cell membrane. The phosphorylated intracellular domain then binds additional signal proteins inside the cell, activating the corresponding downstream AKT or ERK signaling pathway, and inducing a series of cell reactions, such as stimulation or depression of cell proliferation, cell differentiation, cell apoptosis, cell migration or cell adhesion. Among these receptors, mainly ErbB2 and ErbB4 are expressed in the heart.

**[0005]** It has been shown that the EGF-like domains of NRG-1, ranging in size from 50 to 64-amino acids, are sufficient to bind to and activate these receptors. Previous studies have shown that neuregulin-1 $\beta$  (NRG-1 $\beta$ ) can bind directly to ErbB3 and ErbB4 with high affinity. The orphan receptor, ErbB2, can form heterodimer with ErbB3 and ErbB4 with higher affinity than ErbB3 or ErbB4 homodimers. Research in

neural development has indicated that the formation of the sympathetic nervous system requires an intact NRG-1 $\beta$ , ErbB2 and ErbB3 signaling system. Targeted disruption of the NRG-1 $\beta$  or ErbB2 or ErbB4 led to embryonic lethality due to cardiac development defects. Recent studies also highlighted the roles of NRG-1 $\beta$ , ErbB2 and ErbB4 in the cardiovascular development as well as in the maintenance of adult normal heart function. NRG-1 $\beta$  has been shown to enhance sarcomere organization in adult cardiomyocytes. The administration of a recombinant NRG-1 $\beta$  EGF-like domain significantly improves or protects against deterioration in myocardial performance in distinct animal models of heart failure. More importantly, NRG-1 $\beta$  significantly prolongs survival of heart failure animals. These effects make NRG-1 $\beta$  promising as a broad spectrum therapeutic or lead compound for heart failure due to a variety of common diseases. However, there is still a need for more information about the tolerance and effective dosage of NRG, which can be used in a clinical setting for prevention, treating or delaying of heart failure.

### **3. SUMMARY OF THE INVENTION**

[0006] The present invention is based on the discovery that NRG enhances cardiac muscle cell differentiation and organization of sarcomeric and cytoskeleton structure, as well as cell adhesion. The present invention is also based on the discovery that NRG significantly improves or protects against deterioration in myocardial performance in distinct animal models of heart failure and prolongs survival of heart failure animals. Neuregulin, neuregulin polypeptide, neuregulin derivatives, or compounds which mimic the activities of neuregulins, fall within the scope of the methods of the present invention.

[0007] In a first aspect of the invention, the tolerance dosage range of NRG protein when used in mammals (e.g., humans) is provided. Said tolerance dosage range means the largest quantity of NRG that may be administrated into the mammal (e.g., human) without harm. In one embodiment, the tolerance dosage range does not

exceed 2.4  $\mu\text{g}/\text{kg}/\text{day}$ , e.g., 2.4  $\mu\text{g}/\text{kg}/\text{day}$ , 1.2  $\mu\text{g}/\text{kg}/\text{day}$ , 0.6  $\mu\text{g}/\text{kg}/\text{day}$  or 0.3  $\mu\text{g}/\text{kg}/\text{day}$ . In one embodiment, the tolerance dosage range does not exceed 48 U/kg/day, e.g., 48 U/kg/day, 24 U/kg/day, 12 U/kg/day or 6 U/kg/day. In another embodiment, the tolerance dosage range does not exceed 0.34 nmol/kg/day, e.g., 0.34 nmol/kg/day, 0.17 nmol/kg/day, 0.08 nmol/kg/day or 0.04 nmol/kg/day.

**[0008]** In a second aspect of the invention, the effective dosage range of NRG protein for preventing, treating or delaying heart failure in mammals (e.g., humans) is provided. Said effective dosage range means the quantity of NRG that will produce one or more beneficial effects when administered into the mammal (e.g., human). The beneficial effects could be improvement of heart function of patients with heart failure, or prevention of the deterioration of heart function of patients with heart failure, or delaying the deterioration of heart function of patients with heart failure. In one embodiment, the effective dosage range is 0.3-2.4 $\mu\text{g}/\text{kg}/\text{day}$ , e.g., 2.4  $\mu\text{g}/\text{kg}/\text{day}$ , 1.2  $\mu\text{g}/\text{kg}/\text{day}$ , 0.6  $\mu\text{g}/\text{kg}/\text{day}$  or 0.3  $\mu\text{g}/\text{kg}/\text{day}$ . In one embodiment, the effective dosage range is 6-48 U/kg/day, e.g., 48 U/kg/day, 24 U/kg/day, 12 U/kg/day or 6 U/kg/day. In another embodiment, the effective dosage range is 0.04-0.34 nmol/kg/day, e.g., 0.34 nmol/kg/day, 0.17 nmol/kg/day, 0.08 nmol/kg/day or 0.04 nmol/kg/day.

**[0009]** In a third aspect of the invention, a method is provided for preventing, treating or delaying heart failure in mammals (e.g., humans), the method comprising administering a NRG protein within the tolerance dosage range into the mammal (e.g., human) in need thereof. In one embodiment, the NRG protein within the tolerance dosage range is administered once a day. In another embodiment, the NRG protein within the tolerance dosage range is administered several times per day. In one embodiment, the NRG protein within the tolerance dosage range is administered for one day. In another embodiment, the NRG protein within the tolerance dosage range is administered for several days.

**[0010]** In a fourth aspect of the invention, a method is provided for preventing, treating or delaying heart failure in mammals (e.g., humans), the method comprising administering a NRG protein within the effective dosage range into the mammal (e.g.,

human) in need thereof. In one embodiment, the NRG protein within the effective dosage range is administered once a day. In another embodiment, the NRG protein within the effective dosage range is administered several times per day. In one embodiment, the NRG protein within the effective dosage range is administered for one day. In another embodiment, the NRG protein within the effective dosage range is administered for several days.

[0011] In a fifth aspect of the invention, a pharmaceutical composition for preventing, treating or delaying heart failure is provided, the composition comprises NRG protein with the amount within the tolerance dosage range.

[0012] In a sixth aspect of the invention, a pharmaceutical composition for preventing, treating or delaying heart failure is provided, the composition comprises NRG protein with the amount within the effective dosage range.

[0013] In a seventh aspect of the invention, a kit for preventing, treating or delaying heart failure is provided, the kit comprises one dose of or several doses of aforesaid pharmaceutical composition for preventing, treating or delaying heart failure.

#### **4. DETAILED DESCRIPTION OF THE INVENTION**

[0014] Although any methods similar or equivalent to those described herein can be used in the practice of the present invention, the preferred methods and materials are now described.

[0015] The present invention provides methods for preventing, treating or delaying heart failure in mammals (e.g., humans) in need thereof using a NRG protein within the tolerance and/or effective dosage range. Preferably, the mammals (e.g., humans) are subjects or patients suffering from or at risk of developing heart failure.

[0016] For clarity of disclosure, and not by way of limitation, the detailed description of the invention hereinafter is divided into the subsections that follow. All

publications mentioned herein are incorporated by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

#### **A. DEFINITIONS**

[0017] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published applications and other publications referred to herein are incorporated by reference in their entirety. If a definition set forth in this section is contrary to or otherwise inconsistent with a definition set forth in the patents, applications, published applications and other publications that are herein incorporated by reference, the definition set forth in this section prevails over the definition that is incorporated herein by reference.

[0018] As used herein, the singular forms “a”, “an”, and “the” mean “at least one” or “one or more” unless the context clearly dictates otherwise.

[0019] As used herein, “neuregulin” or “NRG” used in the present invention refers to proteins or peptides that can bind and activate ErbB2, ErbB3, ErbB4 or combinations thereof, including but not limited to all neuregulin isoforms, neuregulin EGF domain alone, polypeptides comprising neuregulin EGF-like domain, neuregulin mutants or derivatives, and any kind of neuregulin-like gene products that also activate the above receptors as described in detail below. Neuregulin also includes NRG-1, NRG-2, NRG-3 and NRG-4 proteins, peptides, fragments and compounds that mimic the activities of neuregulin. Neuregulin used in the present invention can activate the above ErbB receptors and modulate their biological reactions, e.g., stimulate acetylcholine receptor synthesis in skeletal muscle cell; and/or improve cardiocyte differentiation, survival and DNA synthesis. Neuregulin also includes those variants with conservative amino acid substitutions that do not substantially alter their biological activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that,

in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al., Molecular Biology of the Gene, 4<sup>th</sup> Edition, 1987, The Bejacmin/Cummings Pub:co., p.224). In preferred embodiments, neuregulin used in the present invention binds to and activates ErbB2/ ErbB4 or ErbB2/ ErbB3 heterodimers, for example, but not for the purpose of restriction, peptides including the 177-237 residues of NRG-1  $\beta$ 2 isoform containing the amino acid sequence: SHLVKCAEKEKTFCVNGGECFMVKDLSNPSRYLCKC PNEFTGDRCQNYVMASFYKAEELYQ (SEQ ID NO:1).

**[0020]** Neuregulin can be administered by any route according to the judgment of those skill in the art, including but not limited to orally, inhalationally, parenterally (e.g., intravenously, intramuscularly, subcutaneously, or intradermally). In certain embodiments, neuregulin is administered orally. In certain embodiments, neuregulin is administered intravenously. In certain embodiments, neuregulin is administered subcutaneously. In preferred embodiments, neuregulin is extendedly released to the mammal intravenously or subcutaneously.

**[0021]** Extended release of neuregulin provides continuous therapeutic level of neuregulin over a period of time. In some embodiments, neuregulin is released over a period of 1 hour, 2 hours, 4 hours, 8 hours, 10 hours, 12 hours, 14 hours, 16 hours, 20 hours, 24 hours or longer. In some embodiments, neuregulin is released over a period of 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days or longer. In another embodiments, neuregulin is released over a period of 1 week, 2 weeks, 3 weeks, 4 weeks or longer. In another embodiments, neuregulin is released over a period of 1 month, 2 months, 4 months, 8 months, 12 months or longer. In another embodiments, neuregulin is released over a period of 1 year, 2 years, 3 years, 4 years or longer. In another embodiments, neuregulin is released 4 hours each day for consecutive 3 days, 4 hours each day for consecutive 5 days, 4 hours each day for consecutive 7 days, 4 hours each day for consecutive 10 days, 6 hours each day for consecutive 3 days, 6 hours each day for consecutive 5 days, 6 hours each day for consecutive 7 days, 6 hours each day for consecutive 10 days, 8 hours each day for



consecutive 3 days, 8 hours each day for consecutive 5 days, 8 hours each day for consecutive 7 days, 8 hours each day for consecutive 10 days.

**[0022]** As used herein, “epidermal growth factor-like domain” or “EGF-like domain” refers to a polypeptide motif encoded by the neuregulin gene that binds to and activates ErbB2, ErbB3, ErbB4, or combinations thereof, and bears a structural similarity to the EGF receptor-binding domain as disclosed in WO 00/64400, Holmes et al., *Science*, 256:1205-1210 (1992); US Patent Nos. 5,530,109 and 5,716,930; Hijazi et al., *Int. J. Oncol.*, 13:1061-1067 (1998); Chang et al., *Nature*, 387:509-512 (1997); Carraway et al., *Nature*, 387:512-516 (1997); Higashiyama et al., *J. Biochem.*, 122:675-680 (1997); and WO 97/09425, the contents of which are all incorporated herein by reference. In certain embodiments, EGF-like domain binds to and activates ErbB2/ErbB4 or ErbB2/ErbB3 heterodimers. In certain embodiments, EGF-like domain comprises the amino acid sequence of the receptor binding domain of NRG-1. In some embodiments, EGF-like domain comprises the amino acid sequence corresponding to amino acid residues 177-226, 177-237, or 177-240 of NRG-1. In certain embodiments, EGF-like domain comprises the amino acid sequence of the receptor binding domain of NRG-2. In certain embodiments, EGF-like domain comprises the amino acid sequence of the receptor binding domain of NRG-3. In certain embodiments, EGF-like domain comprises the amino acid sequence of the receptor binding domain of NRG-4. In certain embodiments, EGF-like domain comprises the amino acid sequence of Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro (SEQ ID NO:2), as described in US Patent No. 5,834,229.

**[0023]** As used herein, “tolerance dosage” refers to the largest quantity of NRG that may be administered into a mammal without harm. In certain embodiments, the harm may be determined by the incidence and severity of clinical adverse events, abnormalities of vital signs including blood pressure, heart rate, breath and body temperature, new electrocardiographic changes and physical or laboratory abnormalities. In certain embodiments, tolerance dosage refers to the largest quantity

of NRG that may be administrated into a human without serious adverse events. In one specific embodiment, the serious adverse events is fatal or life threatening. In another specific embodiment, the serious adverse events results in in-patient hospitalisation or prolonged existing hospitalisation. In another specific embodiment, the serious adverse events results in a persistent or significant disability or incapacity. In another specific embodiment, the serious adverse events results in myocardial infarction, progression to unstable angina or other increase in myocardial ischaemia. In another specific embodiment, the serious adverse events results in the occurrence of a new sustained arrhythmia, e.g., atrial flutter or fibrillation, ventricular tachycardia or fibrillation, complete heart block. In another specific embodiment, the serious adverse events results in congenital anomaly. In other embodiments, the serious adverse events may not be immediately life threatening or result in death or hospitalisation but may jeopardise the subject or require intervention to prevent one of the outcomes listed above.

[0024] As used herein, “effective dosage” refers to the quantity of NRG that will produce one or more beneficial effects when administered into a mammal. In certain embodiments, the beneficial effects may be determined by changes of cardiac function including but not limited by LVEF, LVEDV and LVESV measured by MRI. In certain embodiments, the beneficial effects may be determined by the acute haemodynamics changes, including but not limited by cardiac output, left and right heart filling pressures, pulmonary and systemic blood pressures, and systemic and pulmonary vascular resistance. In certain embodiments, the beneficial effects may be determined by Neuro-hormonal and immunological markers in blood, e.g., noradrenaline, aldosterone, endothelin-1, NT-proBNP, terminal propeptide of type III procollagen (PIIINP), hsCRP, TNF and IL6. In certain embodiments, the beneficial effects may be determined by Six-minute Walk Test. In certain embodiments, the beneficial effects may be determined by New York Heart Association Classification of CHF patients. In certain embodiments, the beneficial effects may be determined by the quality of life (QOL, Minnesota, 1986). In certain embodiments, the beneficial

effects may be determined by the frequency of re-hospitalization of the patient. In certain embodiments, the beneficial effects may be determined by the survival rate or mortality of the patients.

**[0025]** As used herein, “treat”, “treatment” and “treating” refer to any manner in which the symptoms of a condition, disorder or disease are ameliorated or otherwise beneficially altered. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease. Treatment also encompasses any pharmaceutical use of the compositions herein.

**[0026]** As used herein, “left ventricle ejection fraction” or “LVEF” means the portion of blood that is pumped out of a filled ventricle as the result of a heartbeat. It may be defined by the following formula:  $(LVEDV-LVESV) / LVEDV$ .

**[0027]** As used herein, “LVEDV” means left ventricular end-diastolic volume.

**[0028]** As used herein, “LVESV” means left ventricular end-systolic volume.

**[0029]** As used herein, “fractional shortening” or “FS” means a ratio of the change in the diameter of the left ventricle between the contracted and relaxed states. It may be defined by the following formula:  $(LV \text{ end diastolic diameter} - LV \text{ end systolic diameter}) / LV \text{ end diastolic diameter}$ .

**[0030]** As used herein, “heart failure” means an abnormality of cardiac function where the heart does not pump blood at the rate needed for the requirements of metabolizing tissues. Heart failure includes a wide range of disease states such as congestive heart failure, myocardial infarction, tachyarrhythmia, familial hypertrophic cardiomyopathy, ischemic heart disease, idiopathic dilated cardiomyopathy, myocarditis and the like. The heart failure can be caused by any number of factors, including, without limitation, ischemic, congenital, rheumatic, viral, toxic or idiopathic forms. Chronic cardiac hypertrophy is a significantly diseased state which is a precursor to congestive heart failure and cardiac arrest.

**[0031]** As used herein, “protein” is synonymous with “polypeptide” or “peptide” unless the context clearly dictates otherwise.

**[0032]** As used herein, “activity unit” or “EU” or “U” means the quantity of standard product that can induce 50% maximal reaction. In other words, to determine the activity unit for a given active agent, the EC50 must be measured. For example, if the EC50 for a batch of product was 0.1 µg, then that would be one unit. Further, if 1 µg of that products is being used, then 10 EU (1/0.1) is being used. The EC50 can be determined by any method known in the art, including the method employed by the inventor. This determination of the activity unit is important for quality control of genetically engineered products and clinically used drugs, permits product from different pharmaceuticals and/or different batch numbers to be quantified with uniform criteria.

**[0033]** The following is an exemplary, rapid, sensitive, high flux and quantitative method for determination of biological activity of NRG-1 through combining NRG with cell surface ErbB3/ErbB4 molecule and indirect mediation of ErbB2 phosphorylation (See e.g., Michael D. Sadick et al., 1996, Analytical Biochemistry, 235:207-214 and WO03/099300).

**[0034]** Briefly, the assay, termed a kinase receptor activation enzyme-linked immunosorbant assay (KIRA-ELISA), consists of two separate microtiter plates, one for cell culture, ligand stimulation, and cell lysis/receptor solubilization and the other plate for receptor capture and phosphotyrosine ELISA. The assay was developed for analysis of NRG-induced ErbB2 activation and utilizes the stimulation of intact receptor on the adherent breast carcinoma cell line, MCF-7. Membrane proteins are solubilized via Triton X-100 lysis and the receptor is captured in ELISA wells coated with ErbB2-specific antibodies with no cross-reaction to ErbB3 or ErbB4. The degree of receptor phosphorylation is then quantified by antiphosphotyrosine ELISA. A reproducible standard curve is generated with a EC50 of approximately 360pM for heregulin beta 1 (177-244). When identical samples of HRG beta 1 (177-244) are analyzed by both the KIRA-ELISA and quantitative antiphosphotyrosine Western Blot

analysis, the results correlate very closely with one another. The assay described in this report is able to specifically quantify tyrosine phosphorylation of ErbB2 that results from the interaction of HRG with ErbB3 and/or ErbB4.

[0035] Since most of the genetically engineered medicines are proteins and polypeptides, their activity can be determined by their amino acid sequences or the activity center formed by their spatial structure. Activity titer of protein and polypeptide is not consistent with their absolute quality, therefore cannot be determined with weight unit as that of chemical drugs. However, biological activity of genetically engineered medicines is generally consistent with their pharmacodynamics and titer determination system established through given biological activity can determine its titer unit. Therefore, biological activity determination can be part of a process of titering the substance with biological activity and is an important component of quality control of genetically engineered product. It is important to determine biological activity criteria for quality control of genetically engineered product and clinically used drugs.

[0036] Quantity of standard product that can induce 50% maximal reaction is defined as an activity unit (1 EU). Accordingly, product from different pharmaceuticals and of different batch numbers can be quantitated with uniform criteria.

## **B. EXAMPLES**

[0037] The invention is illustrated by the following examples which are not intended to be limiting in any way.

[0038] **Example 1: A phase I, open-label, non-comparative, single-dose study to evaluate the tolerance of recombinant human Neuregulin-1 for injection in healthy volunteers**

[0039] To determine the safety and tolerance of single dose of recombinant human

Neuregulin-1 for injection in healthy subjects, the phase I clinical trial (Protocol ID: KW-70112) was carried out in Peking University First Hospital in China, 28 healthy volunteers were enrolled and were randomized into six dosage groups according to their enrolment order, among which the gender ratio was about 1:1.

**[0040]** Investigational product:

**[0041]** Specification: Neucardin<sup>TM</sup>, a 61-amino acid polypeptide comprising the EGF-like domain of Neuregulin-1  $\beta$ 2 isoform and having the amino acid sequence set forth in SEQ ID NO:1, with the molecular weight of 7054 Dal ( $1\mu\text{g}=0.14\text{nmol}$ ).  $250\mu\text{g}$  (5000 EU) /vial ( $1\mu\text{g}=20$  EU).

**[0042]** Preparation: For injection.

**[0043]** Mode of administration: Slow bolus (20ml/10min, infusion pump control).

**[0044]** Storage: in safe place, with limited access and protected from light, at 3-8 °C.

**[0045]** Dosage groups:

**[0046]** Twenty eight volunteers were divided into six groups with escalating doses equivalent to 0.2 $\mu\text{g}$ , 0.4 $\mu\text{g}$ , 0.8 $\mu\text{g}$ , 1.2 $\mu\text{g}$ , 1.6 $\mu\text{g}$  and 2.4 $\mu\text{g}$  (refer to Table 1 for details). Started from 0.2 $\mu\text{g}/\text{kg}$ , under precondition of confirmation of the safety and tolerability of the former dosage group, the study escalated to next dose.

**Table 1. Dosage groups and subjects in each group**

Group	Dosage	Samples
A	0.2 $\mu\text{g}/\text{kg}$ (4 EU/kg)	4
B	0.4 $\mu\text{g}/\text{kg}$ (8 EU/kg)	4
C	0.8 $\mu\text{g}/\text{kg}$ (16 EU/kg)	4
D	1.2 $\mu\text{g}/\text{kg}$ (24 EU/kg)	5
E	1.6 $\mu\text{g}/\text{kg}$ (32 EU/kg)	6
F	2.4 $\mu\text{g}/\text{kg}$ (48 EU/kg)	5

**[0047]** Study Procedures**[0048]** 1) Screening period:

- Obtain the written informed consent form;
- Verify the inclusion/exclusion criteria;
- Demographics
- Collect the medical history;
- Physical examination;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Haematology, urinalysis;
- Blood biochemistry;
- Serum cardiac markers;
- ECG;
- Ultrasonic cardiogram;
- Blood coagulation test: APTT, PT;
- Serology test (including HBsAg, Anti-HCV, Anti-HIV assay);
- Chest X-ray;
- Urine HCG test (female subject, if necessary);
- Record adverse event and serious adverse event;
- Record concomitant medications.

**[0049]** 2) Baseline: The time from the night before scheduled study drug administration to initiation of study drug administration.

- Physical examination;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Specific antibody assay;
- Re-verify the inclusion/ exclusion criteria;
- Record adverse event and serious adverse event;
- Record concomitant medication.

**[0050]** 3) The day study drug was administrated

- Study drug administration ;
- ECG;
- Echocardiography;
- Record adverse event and serious adverse event;
- Record concomitant medication (each day).

**[0051]** 4) The 1st day and 7th day after cessation of drug administration

- Physical examination;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Specific antibody assay;
- Haematology, urinalysis;
- Blood biochemistry;
- Serum cardiac markers;
- ECG;
- Ultrasonic cardiogram (7th day after cessation of drug administration);
- Blood coagulation test: APTT, PT;
- Urine HCG test (female subject, if necessary);
- Record adverse events and serious adverse events;
- Record concomitant medications.

**[0052]** 5) The 10th day after cessation of drug administration

**[0053]** As to any abnormal item in the 7th day test results after cessation of drug administration, this item was re-tested on the 10th days after dose. After that, it was re-tested with a certain interval until the result became normal.

**[0054]** Results and Analysis

**[0055]** Totally 28 healthy subjects were enrolled into 6 dosage groups of 0.2, 0.4, 0.8, 1.2, 1.6, 2.4 $\mu$ g/kg, the number of subjects who completed study in each dosage



groups was 4, 4, 4, 5, 6, 5 respectively (refer to Table 1).

[0056] Adverse events (AE) were monitored from the time informed consent was signed until the end-of-treatment visit. Sixteen subjects (57.1%) experienced at least one adverse event during the course of the study, all adverse events were mild. None of subject experienced moderate or severe adverse events during the study.

[0057] During the study, the most frequently reported adverse events were listed in Table 2, adverse events by severity were listed in Table 3.

**Table 2. Listing of adverse events**

Group	Subject	AE	Severity	Outcome
0.2µg/kg	402	Abnormal ECG	Mild	Recovery
0.8µg/kg	101	Nausea	Mild	Recovery
	102	Nausea; Lower limb fatigue	Mild	Recovery
	103	Nausea; Fatigue; Loose stools; Light headness	Mild	Recovery
	104	Nausea	Mild	Recovery
1.2µg/kg	601	Abnormal ECG	Mild	Recovery
	603	Hematology abnormality	Mild	Recovery
	605	Abnormal ECG	Mild	Recovery
1.6µg/kg	201	Poor appetite; Hematology abnormality	Mild	Recovery
	202	ECG and Urinary abnormality	Mild	Recovery
	204	Nausea; Abnormal ECG	Mild	Recovery
	206	Nausea; Poor appetite	Mild	Recovery
2.4µg/kg	302	Abnormal ECG	Mild	Recovery
	303	Common cold	Mild	Recovery
	304	Abnormal ECG	Mild	Recovery
	305	Nausea; Light headness; Abnormal ECG	Mild	Recovery

**Table 3. Summary of adverse events by severity**

	0.2µg/kg (N=4) n (%)	0.4µg/kg (N=4) n (%)	0.8µg/kg (N=4) n (%)	1.2µg/kg (N=5) n (%)	1.6µg/kg (N=6) n (%)	2.4µg/kg (N=5) n (%)	Total (N=28) n (%)
Subjects experienced at least one adverse events	1 (25.0)	0 (0.0)	4 (100.0)	3 (60.0)	4 (66.7)	4 (80.0)	16 (57.1)
Subjects classified by worst adverse events							
Mild	1 (25.0)	0 (0.0)	4 (100.0)	3 (60.0)	4 (66.7)	4 (80.0)	16 (57.1)
Moderate	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Subjects experienced at least one SAE	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Subjects experienced at least one adverse events leading to premature termination	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Death	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

**[0058]** The study drug related adverse events were ECG abnormality (8 subjects, 28.6%) and gastrointestinal disorders (7 subjects, 25.0%). ECG abnormalities included slight ST-T segment depression, T-wave flattening or minimal T-wave inversion. During study drug administration period and observation period, none of the subjects experienced precordial discomfort, choking sensation, chest pain, dyspnea; blood pressure was normal at 2 hour and 24 hour after dosing; no abnormality was found in blood biochemistry, electrolytes, serum cardiac markers and echocardiography.

**[0059]** All subjects of each dosage were not changed in physical examination parameters from baseline. Serum antibody assay showed negative results in all subjects (refer to Table 4).

**Table 4. Serum anti-NRG-1 antibody of each group**

Group	First sampling		Second sampling		Third sampling	
	1:50	1:100	1:50	1:100	1:50	1:100
0.2µg/kg	0.18±0.05	0.12±0.02	0.18±0.06	0.13±0.02	0.17±0.06	0.14±0.02
0.4µg/kg	0.12±0.01	0.10±0.01	0.12±0.06	0.10±0.02	0.11±0.01	0.10±0.01
0.8µg/kg	0.16±0.05	0.15±0.02	0.18±0.03	0.15±0.04	0.18±0.05	0.16±0.05

1.2µg/kg	0.12±0.02	0.10±0.02	0.12±0.03	0.10±0.02	0.11±0.02	0.10±0.02
1.6µg/kg	0.21±0.12	0.17±0.08	0.22±0.10	0.17±0.07	0.22±0.10	0.18±0.06
2.4µg/kg	0.15±0.04	0.15±0.03	0.17±0.04	0.13±0.03	0.14±0.03	0.14±0.03

[0060] The results of the phase I clinical trial showed that in the dosage of not exceeding 2.4 µg/kg/day, the drug-related adverse events are mainly mild ECG abnormality and gastrointestinal disorders, no moderate or severe adverse events were observed. So, dosage not exceeding 2.4µg/kg/day when administered with a single dose is tolerable.

**[0061] Example 2: A phase I, open-label, non-comparative, multi-dose study to evaluate the tolerance of recombinant human Neuregulin-1 for injection in healthy volunteers**

[0062] To evaluate the safety and tolerance of multi-dose of recombinant human Neuregulin-1 for injection in healthy subjects for 5 consecutive days, the phase I clinical trial (Protocol ID: KW-70112) was carried out in Peking University First Hospital in China, 32 healthy volunteers were enrolled and were randomized into 4 dosage groups.

[0063] Investigational product:

[0064] Specification: Neucardin™, a 61-amino acid polypeptide comprising the EGF-like domain of Neuregulin-1 β2 isoform and having the amino acid sequence set forth in SEQ ID NO:1, with the molecular weight of 7054 Dal (1µg=0.14nmol). 250µg (5000 EU) /vial (1µg=20 EU).

[0065] Preparation: For injection.

[0066] Mode of administration: Slow bolus (20ml/10min, infusion pump control).

[0067] Storage: in safe place, with limited access and protected from light, at 3-8 °C.

[0068] Dosage groups:

[0069] Thirty two volunteers were divided into four groups with escalating doses equivalent to 0.2µg, 0.4µg, 0.8µg and 1.2µg (refer to Table 5 for details). Started from 0.2µg/kg, under precondition of confirmation of the safety and tolerability of the former dosage group, the study escalated to next dose.

**Table 5. Dosage groups and subjects in each group**

Group	Dosage	Samples
I (5 days)	0.2µg/kg (4EU)	8
II (5 days )	0.4µg/kg (8EU)	8
III (5 days )	0.8µg/kg (16EU)	8
IV (5 days )	1.2µg/kg (24EU)	8

[0070] Study Procedures

[0071] 1) Screening period:

- Obtain the written informed consent form;
- Verify the inclusion/exclusion criteria;
- Demographics;
- Collect the medical history;
- Physical examination;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Haematology, urinalysis;
- Blood biochemistry;
- Serum cardiac markers;
- ECG;
- Ultrasonic cardiogram;
- Blood coagulation test: APTT, PT;
- Serology test (including HBsAg, Anti-HCV, Anti-HIV assay);
- Chest X-ray;
- Urine HCG test (female subject, if necessary);
- Record adverse event and serious adverse event;

- Record concomitant medications.
- [0072] 2) Baseline: The time from the night before scheduled study drug administration to initiation of study drug administration.
- Physical examination;
  - Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
  - Re-verify the inclusion/ exclusion criteria;
  - Record adverse event and serious adverse event;
  - Record concomitant medication.
- [0073] 3) 1<sup>st</sup> day to 5<sup>th</sup> day of drug administration
- Study drug administration;
  - ECG;
  - Record adverse event and serious adverse event (each day);
  - Record concomitant medication (each day).
- [0074] 4) The 1st day and 7th day after cessation of drug administration
- Physical examination;
  - Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
  - Haematology, urinalysis;
  - Blood biochemistry;
  - Serum cardiac markers;
  - ECG;
  - Ultrasonic cardiogram (7th day after cessation of drug administration);
  - Blood coagulation test: APTT, PT;
  - Urine HCG test (female subject, if necessary);
  - Record adverse events and serious adverse events;
  - Record concomitant medications.
- [0075] 5) The 14th day after cessation of drug administration

[0076] As to any abnormal item in the 7th day test results after cessation of drug administration, this item was re-tested on the 10th days after dose. After that, it was re-tested with a certain interval until the result became normal.

[0077] Results and analysis

[0078] A total of 32 healthy subjects were enrolled and randomized into 4 different dosage groups of 0.2, 0.4, 0.8 and 1.2 µg/kg, with 8 subjects in each group and the gender ratio of 1:1. One subject in 1.2 µg/kg dosage group discontinued the study because of adverse event, all other subjects completed the study.

[0079] Adverse events were monitored from the time informed consent was signed until the end-of-treatment visit. Twenty-three subjects (71.9%) experienced at least one adverse event during the course of the study, all adverse events were mild. None of subject experienced severe or serious adverse events during the study (refer to Table 6).

**Table 6. Adverse events by severity**

	0.2µg/kg (N=8) n(%)	0.4µg/kg (N=8) n(%)	0.8µg/kg (N=8) n(%)	1.2µg/kg (N=8) n(%)	Total (N=32) n(%)
Subjects experienced at least one adverse events	4 (50.0)	5 (62.5)	7 (87.5)	7 (87.5)	23(71.9)
Subjects classified by worst adverse events					
Mild	4 (50.0)	5 (62.5)	7 (87.5)	7 (87.5)	23(71.9)
Moderate	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Severe	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Subjects experienced at least one SAE	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Subjects experienced at least one adverse events leading to premature termination	0 (0.0)	0 (0.0)	0 (0.0)	1 (12.5)	1 (3.1)
Death	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

[0080] The frequently reported study drug related adverse events included ECG abnormalities (14 cases, 40.63%) and gastrointestinal disorders (9 cases, 28.13%).

ECG abnormalities included slight ST-T segment depression, T-wave flattening or minimal T-wave inversion. During study drug administration period and observation period, none of the subjects with abnormal ECG experienced precordial discomfort, choking sensation, chest pain, and dyspnea. No treatment was taken in the whole process of study, almost all ECG abnormalities return to normal within 48 hours after 5<sup>th</sup> study drug administration.

[0081] Gastrointestinal disorders related to study drug were stomach discomfort, nausea, vomit and loose stools (refer to Table 7).

**Table 7. Frequently reported study drug related adverse events**

MedDRA system organ class/ MedDRA preferred term	0.2µg/kg (N=8) n(%)	0.4µg/kg (N=8) n(%)	0.8µg/kg (N=8) n(%)	1.2µg/kg (N=8) n(%)	Total (N=32) n(%)
Subjects experienced at least one adverse event	2 (25.0)	4 (50.0)	7 (87.5)	7 (87.5)	20 (62.5)
Skin and hypodermis disorders	0 (0.0)	0 (0.0)	2 (25.0)	0 (0.0)	2 (6.3)
Rash	0 (0.0)	0 (0.0)	2 (25.0)	0 (0.0)	2 (6.3)
Nervous system disorder	0 (0.0)	0 (0.0)	1 (12.5)	0 (0.0)	1 (3.1)
Headache	0 (0.0)	0 (0.0)	1 (12.5)	0 (0.0)	1 (3.1)
Gastrointestinal disorder	0 (0.0)	1 (12.5)	4 (50.0)	4 (50.0)	9 (28.1)
Nausea	0 (0.0)	0 (0.0)	4 (50.0)	2 (25.0)	6 (18.8)
Vomit	0 (0.0)	0 (0.0)	0 (0.0)	2 (25.0)	2 (6.3)
Stomach discomfort	0 (0.0)	1 (12.5)	0 (0.0)	0 (0.0)	1 (3.1)
Loose stool	0 (0.0)	0 (0.0)	0 (0.0)	1 (12.5)	1 (3.1)
Cardiac disorder	2 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (6.3)
Ventricular arrhythmia	2 (25.0)	0 (0.0)	0 (0.0)	0 (0.0)	2 (6.3)
ECG abnormalities	0 (0.0)	3 (37.5)	5 (62.5)	6 (75.0)	14 (43.8)
T wave inversion	0 (0.0)	2 (25.0)	2 (25.0)	3 (37.5)	7 (21.9)
T wave depression	0 (0.0)	2 (25.0)	5 (62.5)	4 (50.0)	11 (34.4)
T wave abnormalities	0 (0.0)	0 (0.0)	0 (0.0)	1 (12.5)	1 (3.1)

[0082] The results of the phase I clinical trial showed that in the dosage of not

exceeding 1.2 µg/kg/day, the drug-related adverse events are mainly mild ECG abnormality and gastrointestinal disorders, no moderate or severe adverse events were observed. The AEs experienced in this multi-dose trial were similar with that of single-dose trial, supporting that the incidence and intensity of AEs will not increase when the study drug was administered for multiple days.

**[0083] Example 3: An open-label, single-center, parallel group study to evaluate the efficacy and safety of recombinant human neuregulin-1 for injection in heart failure patients on standard therapy**

[0084] To evaluate the efficacy and safety of recombinant human neuregulin-1 for injection on chronic heart failure and to investigate the effective dosage range of recombinant human neuregulin-1 for injection for treating chronic heart failure, the phase II, open-label, single-center, parallel group, standard treatment based study (Protocol ID: HREC 06/035) was carried out in St Vincent's Hospital in Australia. Fifteen patients with chronic heart failure were enrolled and randomized into 3 groups.

[0085] Investigational product:

Specification: Neucardin™, a 61-amino acid polypeptide comprising the EGF-like domain of Neuregulin-1 β2 isoform and having the amino acid sequence set forth in SEQ ID NO:1, with the molecular weight of 7054 Dal (1µg=0.14nmol). 250µg (5000 EU)/vial (1µg=20 EU).

[0086] Preparation: For injection.

[0087] Mode of administration: Intravenously drip.

[0088] Storage: in safe place, with limited access and protected from light, at 3-8 °C.

[0089] Dosage groups:

[0090] Fifteen patients with chronic heart failure (LVEF≤40%, NYHA class II/III, established on ACEI/ARA and beta blocker with stable doses of these medications for



the preceding 3 months) were enrolled and randomized into 3 dosage groups with 5 patients in each group (refer to Table 8).

**Table 8. Dosage regimen of phase II trial in Australia**

Dosage group	Sample	Dosage (1 <sup>st</sup> day)	Dosage (2 <sup>nd</sup> -11 <sup>th</sup> day)
A	5	1.2µg/kg/6hrs	2.4µg/kg/12hrs
B	5	1.2µg/kg/6hrs	1.2µg/kg/12hrs
C	5	1.2µg/kg/6hrs	0.6µg/kg/12hrs

**[0091]** Study Procedures

**[0092]** 1) Screening and baseline:

- Obtain the written informed consent form;
- Collect the medical history;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Physical examination;
- Verify the inclusion/exclusion criteria;
- ECG;
- FBC;
- Neurohormonal, immunological marker (NT-proBNP、PIIINP、NA、Aldos、Endothelin、hsCRP、TNF-a、IL-6);
- Ultrasonic cardiogram;
- MRI.

**[0093]** 2) 1<sup>st</sup> day of drug administration

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Study drug administration;
- ECG;
- RH Catheter;

- NT-proBNP.

3) 2<sup>nd</sup> to 11<sup>th</sup> day of drug administration

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Study drug administration;
- ECG.

[0094] 4) 12<sup>th</sup> day of the study

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- ECG;
- RH Catheter;
- FBC;
- Neurohormonal, immunological marker (NT-proBNP, PIIINP, NA, Aldos, Endothelin, hsCRP, TNF-a, IL-6);
- Ultrasonic cardiogram;
- MRI.

[0095] 5) 30<sup>th</sup> day of the study

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- ECG;
- FBC;
- Neurohormonal, immunological marker (NT-proBNP, PIIINP, NA, Aldos, Endothelin, hsCRP, TNF-a, IL-6);
- Ultrasonic cardiogram;
- MRI.

[0096] 6) 60<sup>th</sup> day of the study

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);

- ECG;
- FBC;
- Neurohormonal, immunological marker (NT-proBNP、PIIINP、NA、Aldos、Endothelin、hsCRP、TNF-a、IL-6);

**[0097]** 7) 90<sup>th</sup> day of the study

- Collect the medical history;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Physical examination;
- ECG;
- FBC;
- RH Catheter;
- Neurohormonal, immunological marker (NT-proBNP、PIIINP、NA、Aldos、Endothelin、hsCRP、TNF-a、IL-6);
- Ultrasonic cardiogram;
- MRI.

**[0098]** Results and analysis

**[0099]** The primary efficacy endpoint is evaluating the left ventricular function and remodeling (as determined by change in LVEF, LVEDV and LVESV measured by MRI). Hemodynamics parameters and neurohormonal and immunological markers act as secondary efficacy endpoints.

**[0100]** As a total count, 13 subjects (13/15, 86.7%) received the test drug by IV infusion for 11 consecutive days, one subject (1/15, 6.7%) received the test drug by IV infusion for 10 days, one subject (1/15, 6.7%) received the test drug by IV infusion for 7 days. No subject withdrew.

**[0101]** In Group A (1.2 $\mu$ g/kg/day\* 1 day + 2.4 $\mu$ g/kg/day\* 10 days), 4 subjects (4/5, 80.0%) received the test drug by IV infusion for 11 consecutive days, one subject (1/5, 20.0%) received the test drug by IV infusion for 7 days. In Group B

(1.2µg/kg/day\* 1 day + 1.2µg/kg/day\* 10 days), 5 subjects (5/5, 100.0%) received the test drug by IV infusion for 11 consecutive days. In Group C (1.2µg/kg/day\* 1 day + 0.6µg/kg/day\* 10 days), 4 subjects (4/5, 80.0%) received the test drug by IV infusion for 11 consecutive days, one subject (1/5, 20%) received the test drug by IV infusion for 10 days.

[0102] In group A, after the administration of Neucardin<sup>TM</sup>, at Day 12, the LVEF (%) was improved by 5.4% compared to baseline; the percentage increase was 16.0%; at 1 Month, the LVEF (%) was improved by 3.3% compared to baseline, the percentage of increase was 8.8%; at 3 Months, the LVEF (%) was still improved by 2.6% compared to baseline, the percentage of increase was 6.1%.

[0103] In group B, at Day 12, the LVEF (%) was improved by 3.8% compared to baseline; the percentage increase was 11.8%; at 1 Month, the LVEF (%) was improved by 3.4% compared to baseline, the percentage of increase was 9.7%; at 3 Months, the improvement of LVEF (%) was increased to 4.8% compared to baseline, the percentage of increase was 15.7%.

[0104] In group C, at Day 12, the LVEF (%) was improved by 2.2% compared to baseline; the percentage increase was 8.4%; at 1 Month, the LVEF (%) was improved by 4.6% compared to baseline, the percentage of increase was 15.4%; and the improvement of LVEF (%) sustained to 3 Months by 4.4% compared to baseline, the percentage of increase was 15.5%.

[0105] And for all 15 subjects who completed the study (combined), the improvement of LVEF (%) was 3.8%, 3.8% and 3.9% at day 12, 1 month, 3 month, compared to baseline, respectively (refer to Table 9).

[0106] For all 15 subjects who completed the study (combined) and for each individual dosage group, there was no significant change compared to baseline in LVEDV (ml) at any time point.

[0107] Concerning LVESV (ml), in group A, at Day 12, the LVESV (ml) was decreased by 8.4 ml compared to baseline, the percentage decrease was 4.9%; At 1 Month, the LVESV (ml) was decreased by 14.3 ml compared to baseline, the percentage decrease was 8.8%; At 3 month, the LVESV (ml) was decreased by 12.0

ml compared to baseline, the percentage decrease was 8.3%.

**Table 9. Measured values (Mean±SD) of LVEF (%)**

Dosage group	Time points	Measured LVEF value (%)	Absolute change of LVEF (%)	Percent change of LVEF compared to baseline (%)
Group A (1.2mcg/kg/day*1 day + 2.4mcg/kg/day*10 days)	Baseline (n=5)	33.2±9.7		
	Day 12 (n=5)	38.6±11.6	5.4±2.9	16.0±8.2
	1 Month (n=4)	37.5±13.1	3.3±2.8	8.8±6.3
	3 Months (n=5)	35.8±13.1	2.6±4.0	6.1±11.0
Group B (1.2mcg/kg/day*1 day +1.2mcg/kg/day*10days)	Baseline (n=5)	30.8±8.0		
	Day 12 (n=5)	34.6±10.0	3.8±2.6	11.8±5.9
	1 Month (n=5)	34.2±10.9	3.4±4.1	9.7±12.4
	3 Months (n=5)	35.6±9.4	4.8±3.4	15.7±11.2
Group C (1.2mcg/kg/day*1 day + 0.6mcg/kg/day*10 days)	Baseline (n=5)	32.6±7.3		
	Day 12 (n=5)	34.8±5.4	2.2±3.7	8.4±11.9
	1 Month (n=5)	37.2±6.8	4.6±3.4	15.4±11.5
	3 Months (n=5)	37.0±5.2	4.4±3.0	15.5±12.9
All subjects	Baseline (n=15)	32.2±7.9		
	Day 12 (n=15)	36.0±8.9	3.8±3.2	12.1±8.9
	1 Month (n=14)	36.2±9.6	3.8±3.3	11.5±10.3
	3 Months (n=15)	36.1±9.1	3.9±3.4	12.4±11.8

**[0108]** In group B, at Day 12, the LVESV (ml) was decreased by 1.0 ml compared to baseline, the percentage decrease was 2.5%; At 1 Month, the LVESV (ml) was decreased by 3.8 ml compared to baseline, the percentage decrease was 2.5%; At 3 month, the LVESV (ml) was decreased by 12.6 ml compared to baseline, the percentage decrease was 6.3%.

**[0109]** In group C, at Day 12, the LVESV (ml) was decreased by 11.4 ml compared to baseline, the percentage decrease was 8.6%; At 1 Month, the LVESV (ml) was decreased by 8.0 ml compared to baseline, the percentage decrease was 5.1%; At 3 month, the LVESV (ml) was decreased by 10.8 ml compared to baseline, the percentage decrease was 7.1%.

**[0110]** And for all 15 subjects who completed the study (combined), the decrease of LVESV (ml) was 6.9, 8.3 and 11.8 ml at day 12, 1 month, 3 month, compared to baseline, respectively (refer to Table 10).

**Table 10. Measured values (Mean±SD) of LVESV (ml)**

Dosage groups	Time points	Measured value of LVESV (ml)	Absolutely change of LVESV (ml)	Percent change of LVESV compared to baseline (%)
Group A (1.2mcg/kg/day*1 day + 2.4mcg/kg/day*10days)	Baseline (n=5)	194.2±74.0		
	Day 12 (n=5)	185.8±73.2	-8.4±10.4	-4.9±6.3
	1 Month (n=4)	172.8±81.5	-14.3±14.9	-8.8±9.8
	3 Months (n=5)	182.2±83.4	-12.0±17.6	-8.3±11.4
Group B (1.2mcg/kg/day*1 day + 1.2mcg/kg/day*10 days)	Baseline (n=5)	179.6±34.7		
	Day 12 (n=5)	178.6±60.5	-1.0±29.0	-2.5±16.6
	1 Month (n=5)	175.8±40.1	-3.8±14.5	-2.5±8.3
	3 Months (n=5)	167.0±26.0	-12.6±15.6	-6.3±8.3
Group C (1.2mcg/kg/day*1 day + 0.6mcg/kg/day*10 days)	Baseline (n=5)	182.4±70.5		
	Day 12 (n=5)	171.0±81.6	-11.4±22.2	-8.6±14.2
	1 Month (n=5)	174.4±73.4	-8.0±9.2	-5.1±6.9
	3 Months (n=5)	171.6±74.4	-10.8±8.9	-7.1±6.6
All subjects	Baseline (n=15)	185.4±58.1		
	Day 12 (n=15)	178.5±67.2	-6.9±20.8	-5.3±12.4
	1 Month (n=14)	174.4±60.7	-8.3±12.7	-5.2±8.1
	3 Months (n=15)	173.6±61.7	-11.8±13.5	-7.2±8.4

[0111] A 6-hour infusion of Neucardin<sup>TM</sup> acutely and significantly influenced several hemodynamic parameters when monitored over 24 hours. As shown in Table 11, the acute influence of Neucardin<sup>TM</sup> on hemodynamics was quite positive.

**Table 11. Summary of significant changes in hemodynamics parameters during & after a 6-hour infusion of Neucardin<sup>TM</sup>**

Hemodynamics parameters	Time points when the parameters significantly increased compared to baseline	Time points when the parameters significantly decreased compared to baseline
Heart rate (HR , beat/min)	1 Hour, 2 Hours, 4 Hours & 6 Hours	
Systolic Blood Pressure (SBP , mmHg)	6 Hours	
Diastolic Blood Pressure (DBP , mmHg)		24 Hours
Mean Blood Pressure (MBP , mmHg)		24 Hours

Right Atrial Pressure (RA , mmHg)		1 Hour, 2 Hours, 4 Hours, 6 Hours, 12 Hours & 24 Hours
Pulmonary Artery Systolic Pressure (PASP, mmHg)	30 minutes, 1 Hour	
Pulmonary Artery Diastolic Pressure (PADP , mmHg)		2 Hours, 12 Hours & 24 Hours
Mean Pulmonary Arterial Pressure (MPAP, mmHg)		2 Hours, 12 Hours & 24 Hours
Pulmonary Arterial Wedge Pressure (PAWP , mmHg)		30 minutes, 1 Hour, 2 Hours, 4 Hours, 6 Hours, 12 Hours & 24 Hours
Transpulmonary Pressure Gradient (TPG , mmHg)	30 minutes, 1 Hour, 2 Hours, 4 Hours, 6 Hours & 24 Hours	
Pulmonary Vascular Resistance (PVR , dynes. sec.cm <sup>-5</sup> )		
Systemic Vascular Resistance (SVR , dynes. sec.cm <sup>-5</sup> )		30 minutes, 2 Hours, 4 Hours, 6 Hours & 24 Hours
Cardiac Output (CO, L/min)	30 minutes, 1 Hour, 2 Hours, 4 Hours, 6 Hours & 24 Hours	
Cardiac Index (CI , L/min/m <sup>2</sup> )	30 minutes, 1 Hour, 2 Hours, 4 Hours, 6 Hours & 24 Hours	
Stroke Volume (SV , ml)	30 minutes, 2 Hours, 4 Hours, 12 Hours & 24 Hours	
Stroke Volume Index (SVI , ml/m <sup>2</sup> )	30 minutes, 2 Hours, 4 Hours, 12 Hours & 24 Hours	

[0112] The long term significant changes in hemodynamics for the study subjects are summarized in the following table, and details are listed below (Table 12).

**Table 12. Summary of significant changes in long-term hemodynamic parameters**

Hemodynamics parameters	Time points when the parameters significantly increased compared to baseline	Time points when the parameters significantly decreased compared to baseline
SBP (mmHg)	Day 12, 3 Months	
DBP (mmHg)	Day 12, 3 Months	
MBP (mmHg)	Day 12, 3 Months	
TPG (mmHg)	Day 12, 3 Months	
PAWP (mmHg)		3 Months

[0113] Concerning the neurohormonal and immunological biomarkers, the

NT-proBNP was increased after the 6-hour infusion of Neucardin<sup>TM</sup> on the first day, and did not return to baseline at 24 hours from the start of the administration, and after the end of medication, it decreased and returned to baseline at the end of the study.

[0114] No significant change was observed in hs-CRP.

[0115] Regarding the safety evaluation, 10 subjects (10/15, 66.7%) experienced at least one adverse event (AE). The most commonly reported individual AEs were nausea (20%) , lethargy (20%) and chest pain (20%), without a clear causal relationship to dose. Approximately 67% of subjects had ECG abnormalities during or following study drug dosing, including: T wave inversions or flattening, ST-T segment changes, premature ventricular contractions, non-sustained runs of supraventricular or ventricular tachycardia or atrial flutter/fibrillation. As ECG abnormalities are common in CHF subjects and there was no placebo group for comparison, it is difficult to assess causality or relationship to drug in this study.

[0116] One subject in Group A (2.4 µg/kg/day) and one subject in Group C (0.6 µg/kg/day) experienced serious adverse event(s) (SAEs).

[0117] There were no clinical significant abnormalities identified concerning to vital signs, physical examination, laboratory parameters.

[0118] According to the results of the phase II trial, recombinant human neuregulin-1 is believed to be effective and safe to treat patients with chronic heart failure in the dosage range between 0.6 to 2.4 µg/kg/day (12-48EU/kg/day or 0.08-0.34nmol/kg/day).

[0119] **Example 4: A randomized, double-blinded, multi-center, placebo controlled study to evaluate the efficacy and safety of recombinant human neuregulin 1 in patients with chronic heart failure based on standard treatment**

[0120] To evaluate the efficacy and safety of recombinant human neuregulin-1 for injection on chronic congestive heart failure and to investigate the effective dosage



range of recombinant human neuregulin-1 for injection for treating chronic congestive heart failure, the phase II, double-blinded, multi-center, placebo controlled, standard treatment based study (Protocol ID: ZS-01-206) was carried out in multiple clinical centers in China. Sixty four patients diagnosed as chronic heart failure originating from one or more factors including dilated cardiomyopathy, hypertension, viral myocarditis, myocardial infarction, and alcoholic cardiomyopathy, were enrolled and randomized into 4 groups.

**[0121]** Investigational product:

**[0122]** Specification: Neucardin<sup>TM</sup>, a 61-amino acid polypeptide comprising the EGF-like domain of Neuregulin-1  $\beta$ 2 isoform and having the amino acid sequence set forth in SEQ ID NO:1, with the molecular weight of 7054 Dal ( $1\mu\text{g}=0.14\text{nmol}$ ). 250 $\mu\text{g}$  (5000 EU)/vial ( $1\mu\text{g}=20$  EU).

**[0123]** Preparation: For injection.

**[0124]** Mode of administration: Intravenously drip.

**[0125]** Storage: in safe place, with limited access and protected from light, at 3-8 °C.

**[0126]** Placebo:

**[0127]** Specification: Excipient for Neucardin<sup>TM</sup>. 250 $\mu\text{g}$ /vial and without active recombinant human neuregulin-1 protein.

**[0128]** Dosage groups:

**[0129]** Sixty four patients with chronic heart failure (LVEF $\leq$ 40%, NYHA class II/III, established on ACEI/ARA, beta blocker, diuretic or digoxin with stable doses of these medications for 1 month) were designed to be enrolled and randomized into 4 dosage groups with 16 patients in each group (refer to Table 13).

**Table 13. Dosage groups and regimen**

Dosage	0µg/kg/day	0.3µg/kg/day	0.6µg/kg/day	1.2µg/kg/day
Administration	Intravenous infusion			
Volume	50ml			
Course	10 hours per day, for consecutive 10 days			

**[0130]** Study Procedures

**[0131]** 1) Screening and baseline:

- Obtain the written informed consent form;
- Demography;
- Collect the medical history;
- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Physical examination;
- Blood/Urine routine, blood biochemistry;
- Blood coagulation test: APTT, PT;
- NT-proBNP;
- ECG;
- NYHA classification;
- Ultrasonic cardiogram;
- Chest radiography;
- MRI;
- Tissue Doppler (breast, liver, gallbladder, kidney, adrenal gland, pancreas, spleen, ovary, uterus, prostate);
- Urine pregnancy test (women of child-bearing age);
- Dyspnea evaluation at rest stage;
- Quality of life;
- Six minute walking test;
- Verify the inclusion/exclusion criteria;

- Urine volume for 24 hours;
- Record adverse events and serious adverse events;
- Record concomitant medications.

**[0132]** 2) 1<sup>st</sup> to 10<sup>th</sup> day of drug administration

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- ECG;
- Study drug administration;
- Urine volume for 24 hours;
- Record adverse events and serious adverse events;
- Record concomitant medications.

**[0133]** 3) 11<sup>th</sup> -13<sup>th</sup> day of the study

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Physical examination;
- ECG;
- Blood/Urine routine, blood biochemistry;
- Blood coagulation test: APTT, PT;
- NT-proBNP;
- Urine pregnancy test (women of child-bearing age);
- Quality of life;
- Six minute walking test;
- MRI;
- NYHA classification;
- Dyspnea evaluation at rest stage;
- Urine volume for 24 hours;
- Record adverse events and serious adverse events;
- Record concomitant medications.

**[0134]** 4) 30<sup>th</sup> and 90<sup>th</sup> day of the study

- Vital signs (supine blood pressure, pulse, respiratory rate, temperature);
- Physical examination;
- Blood/Urine routine, blood biochemistry;
- Blood coagulation test: APTT, PT;
- NT-proBNP;
- ECG;
- Chest radiography;
- Six minute walking test;
- MRI;
- NYHA classification;
- Dyspnea evaluation at rest stage;
- Quality of life;
- Record adverse events and serious adverse events;
- Record concomitant medications.

**[0135]** Results and analysis

**[0136]** Forty patients completed the trial. The primary efficacy endpoint is heart function parameters (LVEF, LVEDV and LVESV) at day 30. Concerning LVEF, the mean value of LVEF is decreased from 21.54% at baseline to 20.93% at day 30 in the group of placebo. While in the group of 0.3 µg/kg/day, the mean value of LVEF is slightly improved from 25.08% at baseline to 26.61%, the percentage increase is 4.88%. And in the group of 0.6 µg/kg/day, the mean value of LVEF is significantly increased from 23.03% at baseline to 28.01% at day 30, the percentage increase is 27.11% (refer to Table 14).

**[0137]** Concerning LVEDV, the mean value of LVEDV is increased from 392.16 ml at baseline to 413.35 ml at day 30 in the group of placebo, with the percentage increase of 5.93%. While in the group of 0.3 µg/kg/day, the mean value of LVEDV slightly decreased from 333.62 ml at baseline to 323.22 ml at day 30, with the percentage decrease 3.28%. In the group of 0.6 µg/kg/day and 1.2µg/kg/day, the mean

value of LVEDV significantly decreased from 408.51 ml and 397.04 ml at baseline to 386.89 ml and 374.46 ml at day 30, respectively, both with the percentage decrease more than 5% (refer to Table 15).

**Table 14. LVEF value before and after treatment**

	Group A (Placebo) n=10	Group B 0.3 $\mu$ g/kg/day n=11	Group C 0.6 $\mu$ g/kg/day n=11	Group D 1.2 $\mu$ g/kg/day n=8
Baseline (%)	21.54 $\pm$ 4.73	25.08 $\pm$ 7.64	23.03 $\pm$ 10.23	22.64 $\pm$ 4.24
Day 30 (%)	20.93 $\pm$ 8.53	26.61 $\pm$ 9.68	28.01 $\pm$ 11.27	22.01 $\pm$ 7.21
$\Delta$ LVEF (%)	-0.61 $\pm$ 8.22	1.53 $\pm$ 3.81	4.98 $\pm$ 5.37	-0.63 $\pm$ 4.34
Changed percentage (%)	-1.39 $\pm$ 33.34	4.88 $\pm$ 17.07	27.11 $\pm$ 31.12	-4.01 $\pm$ 19.42

**Table 15. LVEDV value before and after treatment**

	Group A (Placebo) n=10	Group B 0.3 $\mu$ g/kg/day n=11	Group C 0.6 $\mu$ g/kg/day n=11	Group D 1.2 $\mu$ g/kg/day n=8
Baseline (ml)	392.16 $\pm$ 107.38	333.62 $\pm$ 106.14	408.51 $\pm$ 142.85	397.04 $\pm$ 86.94
Day 30 (ml)	413.35 $\pm$ 143.98	323.22 $\pm$ 111.86	386.89 $\pm$ 150.14	374.46 $\pm$ 84.47
$\Delta$ LVEDV (ml)	21.20 $\pm$ 108.13	-10.40 $\pm$ 33.19	-21.63 $\pm$ 42.72	-22.58 $\pm$ 37.72
Changed percentage (%)	5.93 $\pm$ 31.48	-3.28 $\pm$ 11.34	-5.64 $\pm$ 10.03	-5.41 $\pm$ 8.53

[0138] The result of LVESV is similar with that of LVEDV, in the group of placebo, the mean value of LVESV increased from 310.54 ml at baseline to 335.78 ml at day 30, with the percentage increase 9.45%. While in each group of recombinant neuregulin treatment, the mean value of LVESV decreased. For the group of 0.6  $\mu$ g/kg/day as an example, the mean value of LVESV decreased from 325.02 ml at baseline to 291.71 ml at day 30, with the percentage decrease 11.58% (refer to Table 16).

**Table 16. LVESV value before and after treatment**

	Group A (Placebo) n=10	Group B 0.3µg/kg/day n=11	Group C 0.6µg/kg/day n=11	Group D 1.2µg/kg/day n=8
Baseline (ml)	310.54±97.54	254.85±100.86	325.02±142.70	308.46±74.37
Day 30 (ml)	335.78±145.37	244.01±110.30	291.71±147.66	292.22±69.20
△LVESV (ml)	25.24±119.19	-10.84±25.10	-33.31±44.54	-16.24±20.89
Changed percentage (%)	9.45±48.04	-5.33±12.89	-11.58±12.74	-5.01±6.77

[0139] NT-proBNP is an important independent prognostic factor for survival rate of patients with chronic heart failure. From the result, the value of NT-proBNP in the group of placebo and 1.2 µg/kg/day at day 30 and day 90 increased significantly compared to that of baseline. While in the group of 0.3µg/kg/day and 0.6 µg/kg/day, it shows a trend of decrease (refer to Table 17), suggesting a better prognosis.

**Table 17. Change of NT-proBNP at day 30 and day 90**

	Group A (Placebo) n=10	Group B 0.3µg/kg/day n=11	Group C 0.6µg/kg/day n=11	Group D 1.2µg/kg/day n=8
Baseline (fmol/ml)	1013±634.5	1732±1638.4	1403±1755.2	859±460.7
Day 30 (fmol/ml)	811±436.6	1631±1222.1	1021±1045.9	822±349.4
Changed Percentage (%)	6.54±74.29	25.68±72.09	-4.27±41.37	49.42±118.19
Day 90 (fmol/ml)	869±440.9	1663±1664.1	1112±1418.5	847±316.2
Changed Percentage (%)	26.79±102.61	-3.5±60.07	8.66±74.83	45.99±97.45

[0140] There were adverse events in each group, particularly, in the group of 1.2 µg/kg/day, 100% patients in the group experienced at least on adverse event. The most commonly reported individual AEs were nausea, poor appetite and headache. There was no serious adverse events reported (refer to Table 18). Most reported adverse

events relieved without treatment and the rest relieved after the cease of study drug administration or by symptomatic treatment and without sequelae.

**[0141]** There were no clinical significant abnormalities identified concerning to vital signs and physical examination.

**[0142]** From the results of efficacy and safety, we can find that in the dosage of 0.3-1.2 $\mu$ g/kg/day (6-24 EU/kg/day or 0.04-0.17nmol/kg/day), recombinant human neuregulin-1 is effective and safe for the treatment of chronic heart failure.

**Table 18. Incidence of AE and SAE in each group**

	Dosage group	With	Without	Incidence (%)
AE	Group A (n=11)	6	5	54.5
	Group B (n=11)	7	4	63.64
	Group C (n=12)	8	4	66.67
	Group D (n=10)	10	0	100
SAE	Group A (n=11)	0	11	0
	Group B (n=11)	0	11	0
	Group C (n=12)	0	12	0
	Group D (n=10)	0	10	0

**[0143]** From the results of these clinical trials, we can make a conclusion that in the dosage of 0.3-2.4  $\mu$ g/kg/day (6-48 EU/kg/day or 0.04-0.34 nmol/kg/day), recombinant human neuregulin-1 is tolerant and effective for the treatment of chronic heart failure.

What is claimed is:

1. A method for preventing, treating or delaying heart failure in mammals in need thereof, which method comprises administering to the mammal in need thereof a neuregulin 1 protein in an amount:
  - a) from about 0.3  $\mu\text{g}$  to about 2.4  $\mu\text{g}/\text{kg}/\text{day}$ ;
  - b) from about 6 EU to about 48 EU /kg /day; or
  - c) from about 0.04 nmol to about 0.34 nmol /kg /day.
2. The method of claim 1, wherein the neuregulin 1 protein is neuregulin 1  $\beta$ 2 isoform.
3. The method of claim 1, wherein the neuregulin 1 protein comprises the amino acid sequence set forth in SEQ ID NO:1.
4. The method of claim 1, wherein the neuregulin 1 protein consists of the amino acid sequence set forth in SEQ ID NO:1.
5. The method of claim 1, wherein the neuregulin 1 protein is administered to the mammal intravenously.
6. The method of claim 1, wherein the neuregulin 1 protein is administered to the mammal for one day or multiple days.
7. The method of claim 6, wherein the neuregulin 1 protein is administered to the mammal for at least 10 days.
8. The method of claim 7, wherein the neuregulin 1 protein is administered to the mammal for from 10 minutes to 12 hours per day.
9. The method of claim 8, wherein the neuregulin 1 protein is administered to the mammal 10 hours per day for 10 days.
10. The method of claim 1, wherein the mammal has or is suspected of having heart failure.
11. The method of claim 10, wherein the heart failure is caused by one or more



ischaemic, congenital, rheumatic, idiopathic, viral or toxic factors.

12. The method of claim 1, where in the mammal is human.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2010/000845

## A. CLASSIFICATION OF SUBJECT MATTER

See extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CNPAT, CNKI, WPI, EPODOC, NCBI, ISI Web of Knowledge: neuregulin 1, NRG-1, heart failure, sequence search on SEQ ID NO:1

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CN1276381A (QIU L) 13 Dec. 2000 (13.12.2000) See claim 5, lines 15-28 in page 7 and lines 20-28 in page 8 of description.	1-12
X	CN101394861A (ZENSUN SHANGHAI SCI & TECH LTD) 25 March 2009 (25.03.2009), See claims 4, 94, pages 10, 20 in description, and SEQ ID NO: 1.	1-12
X	WO2003099300A1 (ZENSUN SHANGHAI SCI & TECH LTD) 04 Dec. 2003 (04.12.2003), See claims 22, 23, 29, 31, 57, 60, pages 4, 14, 27-28 and examples 1, 2 in description, and SEQ ID NO: 4.	1-12

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
“A” document defining the general state of the art which is not considered to be of particular relevance	“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
“E” earlier application or patent but published on or after the international filing date	“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
“L” document which may throw doubts on priority claim (S) or which is cited to establish the publication date of another citation or other special reason (as specified)	“&” document member of the same patent family
“O” document referring to an oral disclosure, use, exhibition or other means	
“P” document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  
08 Sept. 2010 (08.09.2010)Date of mailing of the international search report  
**16 Sep. 2010 (16.09.2010)**Name and mailing address of the ISA/CN  
The State Intellectual Property Office, the P.R.China  
6 Xitucheng Rd., Jimen Bridge, Haidian District, Beijing, China  
100088  
Facsimile No. 86-10-62019451Authorized officer  
**TIAN, Tian**  
Telephone No. (86-10)62411042

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2010/000845

### Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: 1-12  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 1-12 relate to methods for treating live human/animal bodies. However, the search has been carried out on the basis of the reasonable expectation that the subject matter of claims 1-12 is amended as follows: use of neuregulin 1 protein for manufacturing the drug for preventing, treating or delaying heart failure in mammals in need thereof.
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**International application No.  
PCT/CN2010/000845

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO2009033373A1 (ZENSUN SHANGHAI SCI&TECHNOLOGY LTD) 19 March 2009 (19.03.2009), See claims 14, 15, pages 6, 14 in description and SEQ ID NO: 1.	1-12
X	US7226907B1 (ZENSUN SHANGHAI SCI & TECHNOLOGY LTD) 05 June 2007 (05.06.2007), See claims 24-26, columns 4, 8, 10, 16 in description, and SEQ ID NO: 2.	1-12

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.  
PCT/CN2010/000845

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
CN 1276381 A	13.12.2000	CN 1138785 C	18.02.2004
CN 101394861 A	25.03.2009	WO 2007076701 A1	12.07.2007
		US 2007190127 A1	16.08.2007
		EP 1981525 A1	22.10.2008
		AU 2006332340 A1	12.07.2007
		CA 2634974 A1	12.07.2007
		JP 2009522209T	11.06.2009
		WO 03099300 A1	04.12.2003
WO 2009033373 A1	19.03.2009	AU 2002336030 A1	12.12.2003
		AU 2003238336 A1	12.12.2003
		EP 1553960 A1	20.07.2005
		JP 2005532332 T	27.10.2005
		US 2006019888 A1	26.01.2006
		AU 2003238336 B2	16.07.2009
		WO 03099320 A1	04.12.2003
		AU 2002304965 A1	12.12.2003
		CN 1655804 A	17.08.2005
		CN 1935254 A	28.03.2007
		CN 101007027 A	01.08.2007
		CN 101491670 A	29.07.2009
		CN 100522997 C	05.08.2009
		US 2009156488 A1	18.06.2009
US 7226907 B1	05.06.2007	WO 0037095 A1	29.06.2000
		AU 2422400 A	12.07.2000
		EP 1158998 A1	05.12.2001
		US 2006194734 A1	31.08.2006
		US 2006199767 A1	07.09.2006
		US 2007129296 A1	07.06.2007
		US 2007264254 A1	15.11.2007
		EP 1158998 B1	19.11.2008

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.  
PCT/CN2010/000845

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
		DE 69939950 E	02.01.2009
		EP 2027869 A2	25.02.2009
		EP 2027869 A3	04.03.2009
		ES 2318909 T3	01.05.2009
		US 7612164 B2	03.11.2009

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2010/000845

Continuation of:

## **CLASSIFICATION OF SUBJECT MATTER**

A61K38/18 (2006.01) i

A61P9/04 (2006.01) i