




(86) **Date de dépôt PCT/PCT Filing Date:** 2009/11/20
 (87) **Date publication PCT/PCT Publication Date:** 2010/06/03
 (45) **Date de délivrance/Issue Date:** 2016/08/16
 (85) **Entrée phase nationale/National Entry:** 2011/05/24
 (86) **N° demande PCT/PCT Application No.:** JP 2009/070071
 (87) **N° publication PCT/PCT Publication No.:** 2010/061938
 (30) **Priorité/Priority:** 2008/11/25 (JP2008-299271)

(51) **Cl.Int./Int.Cl. A01N 37/38** (2006.01),
A01N 37/46 (2006.01), **A01N 57/14** (2006.01),
A01P 3/00 (2006.01), **A23B 9/26** (2006.01)
 (72) **Inventeurs/Inventors:**
 TAKAISHI, MASANAO, JP;
 SOMA, MASATO, JP
 (73) **Propriétaire/Owner:**
 SUMITOMO CHEMICAL COMPANY, LIMITED, JP
 (74) **Agent:** FETHERSTONHAUGH & CO.

(54) **Titre : COMPOSITION ET METHODE DE CONTROLE DE MALADIES DES VEGETAUX RENFERMANT DE LA
 MANDESTROBINE ET DU TOLCLOFOS-METHYLE**
 (54) **Title: COMPOSITION AND METHOD FOR CONTROLLING PLANT DISEASES COMPRISING MANDESTROBIN AND
 TOLCLOFOS-METHYL**




(57) **Abrégé/Abstract:**

There is provided a composition and method for controlling plant fungal diseases. The composition comprises a compound of formula (1), together with at least one compound which is tolclofos-methyl, metalaxyl, and mfenoxam. Compound (1) is as follows: (see formula 1) wherein X¹ represents a methyl group, a difluoromethyl group or an ethyl group; X² represents a methoxy group or a methylamino group; and X³ represents a phenyl group, a 2-methylphenyl group or a 2,5-dimethylphenyl group. Compound (1) may be mandestrobin.

25711-877

Abstract

There is provided a composition and method for controlling plant fungal diseases. The composition comprises a compound of formula (1), together with at least one compound which is tolclfos-methyl, metalaxyl, and mfenoxam. Compound (1) is as follows:



wherein X¹ represents a methyl group, a difluoromethyl group or an ethyl group; X² represents a methoxy group or a methylamino group; and X³ represents a phenyl group, a 2-methylphenyl group or a 2,5-dimethylphenyl group. Compound (1) may be mandestrobin.

25711-877

1

DESCRIPTION

COMPOSITION AND METHOD FOR CONTROLLING PLANT DISEASES COMPRISING
MANDESTROBIN AND TOLCLOFOS-METHYL

TECHNICAL FIELD

[0001]

The present invention relates to a
composition for controlling plant diseases and a method
5 for controlling plant diseases.

BACKGROUND ART

[0002]

α -Substituted phenylacetic acid compounds,
tolclofos-methyl, metalaxyl and mefenoxam are
10 conventionally known as active ingredients of agents
for controlling plant diseases (see, for example,
Patent Document 1 and Non-Patent Document 1).
Nevertheless, there is a continuing need for more
highly active agents for controlling plant diseases.

15 [0003]

Patent Document 1: WO 95/27,693

Non-Patent Document 1: "The Pesticide Manual - 14th
edition" published by BCPC, ISBN: 1901396142

DISCLOSURE OF THE INVENTION

20 PROBLEMS TO BE SOLVED BY THE INVENTION

[0004]

An object of the present invention is to provide a composition for controlling plant diseases and a method for controlling plant diseases, having excellent control effect for plant diseases, and so on.

5 MEANS FOR SOLVING THE PROBLEMS


[0005]

The present invention provides a composition for controlling plant diseases and a method for controlling plant diseases, having an improved control
 10 effect for plant diseases by combining a compound represented by the following formula (1) with a specific germicidally active compound.

Specifically, the present invention takes the following constitutions.

15 [1] A composition for controlling plant diseases comprising, as active ingredients, a compound represented by formula (1):

[Formula 1]



wherein X¹ represents a methyl group, a difluoromethyl
 20 group or an ethyl group; X² represents a methoxy group or a methylamino group; and X³ represents a phenyl group, a 2-methylphenyl group or a 2,5-dimethylphenyl

group;

and at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam;

5 [2] The composition according to [1], which has a weight ratio of the compound represented by formula (1) to the at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam falling within the range of from 0.0125:1 to
10 500:1;

[3] A seed treatment agent comprising the compound represented by formula (1) of [1] and at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam as active
15 ingredients;

[4] A plant seed treated with effective amounts of the compound represented by formula (1) of [1] and at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and
20 mefenoxam;

[5] A method for controlling plant diseases which comprises applying effective amounts of the compound represented by formula (1) of [1] and at least one compound selected from the group consisting of
25 tolclofos-methyl, metalaxyl and mefenoxam to a plant or a locus where a plant is allowed to grow; and


[6] Combined use for controlling plant diseases of the acid compound represented by formula

25711-877

4

(1) of [1] and at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam; and so on.

The application further discloses a composition for
5 controlling plant fungal diseases comprising, a compound represented by formula (1b):



and tolclofos-methyl.

Still further, the application discloses a seed
10 treatment agent for controlling plant fungal diseases comprising the compound represented by formula (1b) as defined above and tolclofos-methyl.

Still further, the application discloses use of the
15 compound represented by formula (1b) as defined above and tolclofos-methyl for controlling plant fungal diseases in a plant seed.

Still further, the application discloses a method for
controlling plant fungal diseases which comprises applying
effective amounts of the compound represented by formula (1b) as
20 defined above and tolclofos-methyl to a plant or a locus where a plant is allowed to grow.

25711-877

4a

Still further, the application discloses combined use of the compound represented by formula (1b) as defined above and tolclofos-methyl for controlling plant fungal diseases.

ADVANTAGE OF THE INVENTION

5 [0006]

The composition according to the present invention exhibits an excellent control effect for plant diseases.

BEST MODE FOR CARRYING OUT THE INVENTION

[0007]

10 The compound represented by formula (1) for use in the composition for controlling plant diseases according to the present invention is described.

Examples of the compound represented by formula (1) includes the following compounds.

15 [0008]

A compound in which X^1 is a methyl group, a difluoromethyl group or an ethyl group in formula (1);

a compound in which X^1 is a methyl group in formula (1);

20 a compound in which X^2 is a methoxy group or a methylamino group in formula (1);

a compound in which X^1 is a methyl group and X^2 is a methoxy group in formula (1);

25 a compound in which X^1 is a methyl group and X^2 is methylamino group in formula (1);

[0009]

a compound in which X^3 is a phenyl group, a 2-methylphenyl group or a 2,5-dimethylphenyl group in formula (1);

5 a compound in which X^3 is a phenyl group or a 2,5-dimethylphenyl group in formula (1);

a compound in which X^1 is a methyl group, X^2 is a methoxy group, and X^3 is a 2,5-dimethylphenyl group in formula (1);

10 a compound in which X^1 is a methyl group, X^2 is methylamino group, and X^3 is a phenyl group in formula (1); and

a compound in which X^1 is a methyl group, X^2 is methylamino group, and X^3 is a 2,5-dimethylphenyl
15 group in formula (1).

[0010]

Next, specific examples of the compound represented by formula (1) are shown.

In the compound represented by formula (1),
20 X^1 , X^2 , X^3 are one of the combinations of substituents shown in Table 1.

[0011]

[Table 1]

X ¹	X ²	X ³
CH ₃	OCH ₃	Ph
CH ₃	OCH ₃	2-CH ₃ Ph
CH ₃	OCH ₃	2, 5-(CH ₃) ₂ Ph
CH ₃	NHCH ₃	Ph
CH ₃	NHCH ₃	2-CH ₃ Ph
CH ₃	NHCH ₃	2, 5-(CH ₃) ₂ Ph
CHF ₂	OCH ₃	Ph
CHF ₂	OCH ₃	2-CH ₃ Ph
CHF ₂	OCH ₃	2, 5-(CH ₃) ₂ Ph
CHF ₂	NHCH ₃	Ph
CHF ₂	NHCH ₃	2-CH ₃ Ph
CHF ₂	NHCH ₃	2, 5-(CH ₃) ₂ Ph
C ₂ H ₅	OCH ₃	Ph
C ₂ H ₅	OCH ₃	2-CH ₃ Ph
C ₂ H ₅	OCH ₃	2, 5-(CH ₃) ₂ Ph
C ₂ H ₅	NHCH ₃	Ph
C ₂ H ₅	NHCH ₃	2-CH ₃ Ph
C ₂ H ₅	NHCH ₃	2, 5-(CH ₃) ₂ Ph

[0012]

The compound represented by formula (1) may have isomers such as stereoisomers such as optical isomers based on an asymmetric carbon atoms and tautomers, and any isomer can be contained and used solely or in a mixture of any isomer ratio in the

present invention.

[0013]

The compound represented by formula (1) may be in a form of a solvate (for example, hydrate) and it
5 can be used in a form of a solvate in the present invention.

[0014]

The compound represented by formula (1) may be in a form of a crystal form and/or an amorphous form
10 and it can be used in any form in the present invention.

[0015]

The compound represented by formula (1) is a compound described in W095/27,693 pamphlet. These
15 compounds can be synthesized, for example, by a method described in the pamphlet.

[0016]

Next, the germicidally active compound for use in the composition for controlling plant diseases
20 according to the present invention in combination with the compound represented by formula (1) is at least one compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam. All of the germicidally active compounds included in this group
25 are well-known compounds, such as those described in "The Pesticide Manual - 14th edition" published by BCPC, ISBN:1901396142, pp.1043, 678 and 679. These compounds can be obtained from commercial agents or

prepared using well-known methods.

[0017]

In the composition for controlling plant diseases according to the present invention, the weight ratio of the compound represented by formula (1) to the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam is typically in the range of 0.0125:1 to 500:1, preferably 0.025:1 to 100:1. In addition, when used as a dusting powder, the range of 0.025:1 to 40:1 is more preferable, and when used as a seed treatment agent, the range of 0.25:1 to 100:1 is more preferable.

[0018]

The composition for controlling plant diseases according to the present invention may be a simple mixture of the compound represented by formula (1) and the compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam. Alternatively, the composition for controlling plant diseases is typically produced by mixing the compound represented by formula (1) and the compound selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam with an inert carrier, and adding to the mixture a surfactant and other adjuvants as needed so that the mixture can be formulated into an oil agent, an emulsion, a flowable agent, a wettable powder, a granulated wettable powder, a powder agent, a granule agent and so on. The composition for

controlling the compound(s) mentioned above can be used as a seed treatment agent of the present invention as it is or added with other inert ingredients.

In the composition for controlling plant
5 diseases according to the present invention, the total amount of the compound represented by formula (1) and the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam is typically in the range of 0.1 to 99% by weight, preferably 0.2 to
10 90% by weight.

[0019]

Examples of the solid carrier used in formulation include fine powders or granules such as minerals such as kaolin clay, attapulgite clay,
15 bentonite, montmorillonite, acid white clay, pyrophyllite, talc, diatomaceous earth and calcite; natural organic materials such as corn rachis powder and walnut husk powder; synthetic organic materials such as urea; salts such as calcium carbonate and
20 ammonium sulfate; synthetic inorganic materials such as synthetic hydrated silicon oxide; and as a liquid carrier, aromatic hydrocarbons such as xylene, alkylbenzene and methylnaphthalene; alcohols such as 2-propanol, ethyleneglycol, propylene glycol, and
25 ethylene glycol monoethyl ether; ketones such as acetone, cyclohexanone and isophorone; vegetable oil such as soybean oil and cotton seed oil; petroleum aliphatic hydrocarbons, esters, dimethylsulfoxide,

25711-877

10

acetonitrile and water.

Examples of the surfactant include anionic surfactants such as alkyl sulfate ester salts, alkylaryl sulfonate salts, dialkyl sulfosuccinate salts, polyoxyethylene alkylaryl ether phosphate ester salts, lignosulfonate salts and naphthalene sulfonate formaldehyde polycondensates; and nonionic surfactants such as polyoxyethylene alkyl aryl ethers, polyoxyethylene alkylpolyoxypropylene block copolymers and sorbitan fatty acid esters and cationic surfactants such as alkyltrimethylammonium salts.

Examples of the other formulation auxiliary agents include water-soluble polymers such as polyvinyl alcohol and polyvinylpyrrolidone, polysaccharides such as Arabic gum, alginic acid and the salt thereof, CMC (carboxymethyl-cellulose), Xanthan gum, inorganic materials such as aluminum magnesium silicate and alumina sol, preservatives, coloring agents and stabilization agents such as PAP (acid phosphate isopropyl) and BHT.

[0020]

The composition for controlling plant diseases according to the present invention is effective for the following plant diseases and the

25711-877

11

Diseases of rice: blast (*Magnaporthe grisea*), *Helminthosporium* leaf spot (*Cochliobolus miyabeanus*), sheath blight (*Rhizoctonia solani*), and bakanae disease (*Gibberella fujikuroi*).

Diseases of wheat: powdery mildew (*Erysiphe*
 5 *graminis*), *Fusarium* head blight (*Fusarium graminearum*, *F.*
avenacerum, *F. culmorum*, *Microdochium nivale*), rust (*Puccinia*
striiformis, *P. graminis*, *P. recondita*), pink snow mold
 (*Micronectriella nivale*), *Typhula* snow blight (*Typhula* sp.),
 loose smut (*Ustilago tritici*), bunt (*Tilletia caries*), eyespot
 10 (*Pseudocercosporella herpotrichoides*), leaf blotch
 (*Mycosphaerella graminicola*), glume blotch (*Stagonospora*
nodorum), and yellow spot (*Pyrenophora tritici-repentis*).

Diseases of barley: powdery mildew (*Erysiphe*
graminis), *Fusarium* head blight (*Fusarium graminearum*, *F.*
 15 *avenacerum*, *F. culmorum*, *Microdochium nivale*), rust (*Puccinia*
striiformis, *P. graminis*, *P. hordei*), loose smut (*Ustilago*
nuda), scald (*Rhynchosporium secalis*), net blotch (*Pyrenophora*
teres), spot blotch (*Cochliobolus sativus*), leaf stripe
 (*Pyrenophora graminea*), and *Rhizoctonia* damping-off
 20 (*Rhizoctonia solani*).

Diseases of corn: smut (*Ustilago maydis*), brown spot
 (*Cochliobolus heterostrophus*), copper spot (*Gloeocercospora*
sorghii), southern rust (*Puccinia polysora*), gray leaf spot
 (*Cercospora zea-maydis*), and *Rhizoctonia* damping-off
 25 (*Rhizoctonia solani*).

[0021]

Diseases of citrus: melanose (*Diaporthe citri*), scab
 (*Elsinoe fawcetti*), penicillium rot (*Penicillium digitatum*, *P.*

25711-877

12

italicum), and brown rot (*Phytophthora parasitica*, *Phytophthora citrophthora*).

Diseases of apple: blossom blight (*Monilinia mali*), canker (*Valsa ceratosperma*), powdery mildew (*Podosphaera leucotricha*), *Alternaria* leaf spot (*Alternaria alternata* apple pathotype), scab (*Venturia inaequalis*), bitter rot (*Colletotrichum acutatum*), crown rot (*Phytophthora cactorum*), blotch (*Diplocarpon mali*), and ring rot (*Botryosphaeria berengeriana*).

10 Diseases of pear: scab (*Venturia nashicola*, *V. pirina*), black spot (*Alternaria alternata* Japanese pear pathotype), rust (*Gymnosporangium haraeaeum*), and phytophthora fruit rot (*Phytophthora cactorum*);

15 Diseases of peach: brown rot (*Monilinia fructicola*), scab (*Cladosporium carpophilum*), and phomopsis rot (*Phomopsis* sp.).

Diseases of grape: anthracnose (*Elsinoe ampelina*), ripe rot (*Glomerella cingulata*), powdery mildew (*Uncinula necator*), rust (*Phakopsora ampelopsidis*), black rot (*Guignardia bidwellii*), and downy mildew (*Plasmopara viticola*).

Diseases of Japanese persimmon: anthracnose (*Gloeosporium kaki*), and leaf spot (*Cercospora kaki*, *Mycosphaerella nawae*).

25 Diseases of gourd: anthracnose (*Colletotrichum lagenarium*), powdery mildew (*Sphaerotheca fuliginea*), gummy stem blight (*Mycosphaerella melonis*), *Fusarium* wilt (*Fusarium oxysporum*), downy mildew (*Pseudoperonospora cubensis*),

25711-877

13

Phytophthora rot (*Phytophthora* sp.), and damping-off (*Pythium* sp.);

Diseases of tomato: early blight (*Alternaria solani*), leaf mold (*Cladosporium fulvum*), and late blight (*Phytophthora*
5 *infestans*).

Diseases of eggplant: brown spot (*Phomopsis vexans*), and powdery mildew (*Erysiphe cichoracearum*).

Diseases of cruciferous vegetables: *Alternaria* leaf spot (*Alternaria japonica*), white spot (*Cercospora*
10 *brassicae*), clubroot (*Plasmodiophora brassicae*), and downy mildew (*Peronospora parasitica*).

Diseases of welsh onion: rust (*Puccinia allii*), and downy mildew (*Peronospora destructor*).

[0022]

15 Diseases of soybean: purple seed stain (*Cercospora kikuchii*), sphaceloma scab (*Elsinoe glycines*), pod and stem blight (*Diaporthe phaseolorum* var. *sojae*), septoria brown spot (*Septoria glycines*), frog-eye leaf spot (*Cercospora sojae*),
20 rust (*Phakopsora pachyrhizi*), brown stem rot (*Phytophthora sojae*), and *Rhizoctonia* damping-off (*Rhizoctonia solani*).

Diseases of kidney bean: anthracnose (*Colletotrichum lindemthianum*). Diseases of peanut: leaf spot (*Cercospora personata*), brown leaf spot (*Cercospora arachidicola*) and southern blight (*Sclerotium rolfsii*).

25 Diseases of garden pea: powdery mildew (*Erysiphe pisi*), and root rot (*Fusarium solani* f. sp. *pisi*).

25711-877

14

Diseases of potato: early blight (*Alternaria solani*), late blight (*Phytophthora infestans*), pink rot (*Phytophthora erythroseptica*), powdery scab (*Spongospora subterranean* f. sp. *subterranea*) and black scurf (*Rhizoctonia solani*).

5 Diseases of strawberry: powdery mildew (*Sphaerotheca humuli*), and anthracnose (*Glomerella cingulata*).

Diseases of tea: net blister blight (*Exobasidium reticulatum*), white scab (*Elsinoe leucospila*), gray blight (*Pestalotiopsis* sp.), and anthracnose (*Colletotrichum theae-sinensis*).

Diseases of tobacco: brown spot (*Alternaria longipes*), powdery mildew (*Erysiphe cichoracearum*), anthracnose (*Colletotrichum tabacum*), downy mildew (*Peronospora tabacina*), and black shank (*Phytophthora nicotianae*).

15 Diseases of rapeseed: sclerotinia rot (*Sclerotinia sclerotiorum*), and *Rhizoctonia* damping-off (*Rhizoctonia solani*).

Diseases of cotton: *Rhizoctonia* damping-off (*Rhizoctonia solani*).

20 Diseases of sugar beet: *Cercospora* leaf spot (*Cercospora beticola*), leaf blight (*Thanatephorus cucumeris*), Root rot (*Thanatephorus cucumeris*), and *Aphanomyces* root rot (*Aphanomyces cochlioides*).

Diseases of rose: black spot (*Diplocarpon rosae*),
25 powdery mildew (*Sphaerotheca pannosa*), and downy mildew (*Peronospora sparsa*).

25711-877

15

Diseases of chrysanthemum and asteraceous plants: downy mildew (*Bremia lactucae*), leaf blight (*Septoria chrysanthemi-indici*), and white rust (*Puccinia horiana*).

5 Diseases of various groups: diseases caused by *Pythium* spp. (*Pythium aphanidermatum*, *Pythium debarianum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium ultimum*), gray mold (*Botrytis cinerea*), and *Sclerotinia* rot (*Sclerotinia sclerotiorum*).

10 Disease of Japanese radish: *Alternaria* leaf spot (*Alternaria brassicicola*).

Diseases of turfgrass: dollar spot (*Sclerotinia homeocarpa*), and brown patch and large patch (*Rhizoctonia solani*).

15 Disease of banana: sigatoka (*Mycosphaerella fijiensis*, *Mycosphaerella musicola*).

Disease of sunflower: downy mildew (*Plasmopara halstedii*).

20 Seed diseases or diseases in the early stages of the growth of various plants caused by bacteria of *Aspergillus* genus, *Penicillium* genus, *Fusarium* genus, *Gibberella* genus, *Trichoderma* genus, *Thielaviopsis* genus, *Rhizopus* genus, *Mucor* genus, *Corticium* genus, *Phoma* genus, *Rhizoctonia* genus and *Diplodia* genus.

Viral diseases of various plants mediated by

25711-877

16

Polymixa genus or the *Olpidium* genus and so on.

[0023]

plant diseases can be controlled by applying effective amounts of the compound represented by formula (1) and the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam to the plant pathogens or a place where the plant pathogens inhabit or a place (plant, soil) where the plant pathogens may inhabit.

10 Plant diseases can be controlled by applying effective amounts of the compound represented by formula (1) and the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam to a plant or a place where a plant is allowed to grow.

15 As a plant which is the object of application, stalk and leaves of the plant, seed of the plant, bulbs of the plant can be included. Here, the bulb means a bulb, corm, rhizoma, stem tuber, root tuber and rhizophore.

20 When the application is conducted to plant diseases, a plant or the soil where the plant is allowed to grow, the compound represented by formula (1) and the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam

25 may be separately applied for the same period, but they are typically applied as a composition for controlling plant diseases of the present invention from the viewpoint of simplicity of the application.

The controlling method of the present invention includes treatment of stalk and leaves of a plant, treatment of the place where the plant is allowed to grow such as the soil, treatment of the seeds such as seed sterilization/seed coating and treatment of the bulb such as potato sets.

As the treatment of stalk and leaves of a plant in the control method of the present invention, specifically, for example, application onto the surface of the plant such as spraying to the stalk and leaves and spraying to the trunk can be included.

As the treatment of the soil in the control method of the present invention, for example, spraying onto the soil, admixing with the soil, perfusion of an agent liquid into the soil (irrigation of an agent liquid, injection into the soil, dripping of an agent liquid) can be included and the examples of the place to be treated include a planting hole, a furrow, peripheral of the planting hole, peripheral of the planting furrow, the entire surface of the growing area, the parts between the soil and the plant, area between roots, area beneath the trunk, main furrow, growing soil, box for raising seedlings, tray for raising seedlings, seedbed. The treatment can be performed before dissemination, at the time of dissemination, immediately after the dissemination, during the raising period of seedlings, before settled planting, at the time of settled planting and growing

time after settled planting. In the soil treatment mentioned above, the active ingredients may be applied to the plant at the same time, or solid manure such as paste manure containing the active ingredients may be applied to the soil. The active ingredients may be mixed in irrigating liquid, and, for example, may be injected to irrigating facilities (irrigating tube, irrigating pipe, sprinkler, etc.), mixed into the flooding liquid between furrows, or mixed into a water culture medium. Alternatively, the irrigating liquid and the active ingredients may be mixed beforehand and, for example, used for treatment by an appropriate irrigating method including the irrigating method mentioned above and the other methods such as sprinkling and flooding.

Treatment of a seed in the control method of the present invention is, for example, a method for treating a seed, a bulb or the like to be protected from plant diseases with a composition for controlling plant diseases of the present invention and specific examples thereof include a spraying treatment in which a suspension of the composition for controlling plant diseases of the present invention is atomized and sprayed on the seed surface or the bulb surface; smearing treatment in which a wettable powder, an emulsion, a flowable agent or the like of the composition for controlling plant diseases of the present invention as it is or added with a small amount

of water is applied on the seed surface or the bulb surface; immersing treatment in which the seed is immersed in a solution of the composition for controlling plant diseases of the present invention for a certain period of time; film coating treatment and pellet coating treatment.

[0024]

When a plant or the soil for growing a plant is treated with a compound represented by formula (1) and compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam, the amount for the treatment may be changed depending on the kind of the plant to be treated, the kind and the occurring frequency of the diseases to be controlled, formulation form, treatment period, climatic condition and so on but the total amount of the compound represented by formula (1) and the compound(s) selected from the group consisting of tolclofos-methyl, metalaxyl and mefenoxam (hereinbelow referred to as the amount of the active ingredients) per 10,000m² is typically 1 to 5000 g and preferably 2 to 200 g.

The emulsion, wettable powder, flowable agent or the like is typically diluted with water, and then sprinkled for treatment. In this case, the concentration of the active ingredients is typically in the range of 0.0001 to 3% by weight and preferably 0.0005 to 1% by weight. The powder agent, granule agent or the like is typically used for treatment

without dilution.

In the treatment of seeds, the amount of the applied active ingredients is typically in the range of 0.001 to 20 g, preferably 0.01 to 5 g per 1 kg of
5 seeds.

[0025]

The control method of the present invention can be used in agricultural lands such as fields, paddy fields, lawns and orchards or in non-agricultural
10 lands.

The present invention can be used to control diseases in agricultural lands for cultivating the following "plant" and the like without adversely affecting the plant and so on.

15 Examples of the crops are as follows:

crops: corn, rice, wheat, barley, rye, oat, sorghum, cotton, soybean, peanut, buckwheat, beet, rapeseed, sunflower, sugar cane, tobacco, etc.;

vegetables: solanaceous vegetables (eggplant,
20 tomato, pimento, pepper, potato, etc.), cucurbitaceous vegetables (cucumber, pumpkin, zucchini, water melon, melon, squash, etc.), cruciferous vegetables (Japanese radish, white turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, leaf mustard, broccoli, cauliflower,
25 etc.), asteraceous vegetables (burdock, crown daisy, artichoke, lettuce, etc.), liliaceous vegetables (green onion, onion, garlic, and asparagus), ammiaceous vegetables (carrot, parsley, celery, parsnip, etc.),

25711-877

21

chenopodiaceous vegetables (spinach, Swiss chard, etc.),
 lamiaceous vegetables (*Perilla frutescens*, mint, basil, etc.),
 strawberry, sweet potato, *Dioscorea japonica*, colocasia, etc.,

flowers,

5 foliage plants,

turf grasses,

fruits: pomaceous fruits (apple, pear, Japanese pear,
 Chinese quince, quince, etc.), stone fleshy fruits (peach, plum,
 nectarine, *Prunus mume*, cherry fruit, apricot, prune, etc.),
 10 citrus fruits (*Citrus unshiu*, orange, lemon, lime, grapefruit,
 etc.), nuts (chestnuts, walnuts, hazelnuts, almond, pistachio,
 cashew nuts, macadamia nuts, etc.), berries (blueberry,
 cranberry, blackberry, raspberry, etc.), grape, kaki fruit,
 olive, Japanese plum, banana, coffee, date palm, coconuts, etc.,

15 trees other than fruit trees; tea, mulberry, flowering
 plant, roadside trees (ash, birch, dogwood, *Eucalyptus*, *Ginkgo*
biloba, lilac, maple, *Quercus*, poplar, Judas tree, *Liquidambar*
formosana, plane tree, zelkova, Japanese arborvitae, fir wood,
 hemlock, juniper, *Pinus*, *Picea*, and *Taxus cuspidate*), etc.

20 [0026]

The aforementioned "plants" include plants, to which
 resistance to HPPD inhibitors such as isoxaflutole, ALS
 inhibitors such as imazethapyr or thifensulfuron-methyl, EPSP
 synthetase inhibitors such

as glyphosate, glutamine synthetase inhibitors such as the glufosinate, acetyl-CoA carboxylase inhibitors such as sethoxydim, PPO inhibitors such as flumioxazin, and herbicides such as bromoxynil, dicamba, 2,4-D, etc. has
5 been conferred by a classical breeding method or genetic engineering technique.

Examples of a "plant" on which resistance has been conferred by a classical breeding method include rape, wheat, sunflower and rice resistant to
10 imidazolinone ALS inhibitory herbicides such as imazethapyr, which are already commercially available under a product name of Clearfield (registered trademark). Similarly, there is soy bean on which resistance to sulfonylurea ALS inhibitory herbicides
15 such as thifensulfuron-methyl has been conferred by a classical breeding method, which is already commercially available under a product name of STS soy bean. Similarly, examples on which resistance to acetyl-CoA carboxylase inhibitors such as trione oxime
20 or aryloxy phenoxypropionic acid herbicides has been conferred by a classical breeding method include SR corn. The plant on which resistance to acetyl-CoA carboxylase inhibitors has been conferred is described in Proceedings of the National Academy of Sciences of
25 the United States of America (Proc. Natl. Acad. Sci. USA), vol. 87, pp. 7175-7179 (1990). A variation of acetyl-CoA carboxylase resistant to an acetyl-CoA carboxylase inhibitor is reported in Weed Science, vol.

53, pp. 728-746 (2005) and a plant resistant to acetyl-CoA carboxylase inhibitors can be generated by introducing a gene of such an acetyl-CoA carboxylase variation into a plant by genetically engineering technology, or by introducing a variation conferring resistance into a plant acetyl-CoA carboxylase. Furthermore, plants resistant to acetyl-CoA carboxylase inhibitors or ALS inhibitors or the like can be generated by introducing a site-directed amino acid substitution variation into an acetyl-CoA carboxylase gene or the ALS gene of the plant by introduction a nucleic acid into which has been introduced a base substitution variation represented Chimeraplasty Technique (Gura T. 1999. Repairing the Genome's Spelling Mistakes. Science 285: 316-318) into a plant cell.

[0027]

Examples of a plant on which resistance has been conferred by genetic engineering technology include corn, soy bean, cotton, rape, sugar beet resistant to glyphosate, which is already commercially available under a product name of RoundupReady (registered trademark), AgrisureGT, etc. Similarly, there are corn, soy bean, cotton and rape which are made resistant to glufosinate by genetic engineering technology, a kind, which is already commercially available under a product name of LibertyLink (registered trademark). A cotton made resistant to

25711-877

24

bromoxynil by genetic engineering technology is already commercially available under a product name of BXN likewise.

[0028]

5 The aforementioned "plants" include genetically engineered crops produced using such genetic engineering techniques, which, for example, are able to synthesize selective toxins as known in genus *Bacillus*.

10 Examples of toxins expressed in such genetically engineered crops include: insecticidal proteins derived from *Bacillus cereus* or *Bacillus popilliae*; δ -endotoxins such as Cry1Ab, Cry1Ac, Cry1F, Cry1Fa2, Cry2Ab, Cry3A, Cry3Bb1 or Cry9C, derived from
15 *Bacillus thuringiensis*; insecticidal proteins such as VIP1, VIP2, VIP3, or VIP3A; insecticidal proteins derived from nematodes; toxins generated by animals, such as scorpion toxin, spider toxin, bee toxin, or insect-specific neurotoxins; mold fungi toxins; plant
20 lectin; agglutinin; protease inhibitors such as a trypsin inhibitor, a serine protease inhibitor, patatin, cystatin, or a papain inhibitor; ribosome-inactivating proteins (RIP) such as lycine, corn-RIP, abrin, luffin, saporin, or briodin; steroid-
25 metabolizing enzymes such as 3-hydroxysteroid oxidase, ecdysteroid-UDP-glucosyl transferase, or cholesterol oxidase; an ecdysone inhibitor; HMG-COA reductase; ion channel inhibitors such as a sodium channel inhibitor

or calcium channel inhibitor; juvenile hormone esterase; a diuretic hormone receptor; stilbene synthase; bibenzyl synthase; chitinase; and glucanase.

Moreover, toxins expressed in such

5 genetically engineered crops also include: hybrid toxins of δ -endotoxin proteins such as Cry1Ab, Cry1Ac, Cry1F, Cry1Fa2, Cry2Ab, Cry3A, Cry3Bb1, Cry9C, Cry34Ab or Cry35Ab and insecticidal proteins such as VIP1, VIP2, VIP3 or VIP3A; partially deleted toxins; and

10 modified toxins. Such hybrid toxins are produced from a new combination of the different domains of such proteins, using a genetic engineering technique. As a partially deleted toxin, Cry1Ab comprising a deletion of a portion of an amino acid sequence has been known.

15 A modified toxin is produced by substitution of one or multiple amino acids of natural toxins.

Examples of such toxins and genetically engineered plants capable of synthesizing such toxins are described in EP-A-0 374 753, WO 93/07278, WO

20 95/34656, EP-A-0 427 529, EP-A-451 878, WO 03/052073, etc.

Toxins contained in such genetically engineered plants are able to confer resistance particularly to insect pests belonging to Coleoptera,

25 Hemiptera, Diptera, Lepidoptera and Nematodes, to the plants.

[0029]

Furthermore, genetically engineered plants,

which comprise one or multiple insecticidal pest-resistant genes and which express one or multiple toxins, have already been known, and some of such genetically engineered plants have already been on the market. Examples of such genetically engineered plants include YieldGard (registered trademark) (a corn variety for expressing Cry1Ab toxin), YieldGard Rootworm (registered trademark) (a corn variety for expressing Cry3Bb1 toxin), YieldGard Plus (registered trademark) (a corn variety for expressing Cry1Ab and Cry3Bb1 toxins), Herculex I (registered trademark) (a corn variety for expressing phosphinotricine N-acetyl transferase (PAT) so as to confer resistance to Cry1Fa2 toxin and glufosinate), NuCOTN33B (registered trademark) (a cotton variety for expressing Cry1Ac toxin), Bollgard I (registered trademark) (a cotton variety for expressing Cry1Ac toxin), Bollgard II (registered trademark) (a cotton variety for expressing Cry1Ac and Cry2Ab toxins), VIPCOT (registered trademark) (a cotton variety for expressing VIP toxin), NewLeaf (registered trademark) (a potato variety for expressing Cry3A toxin), NatureGard (registered trademark) Agrisure (registered trademark) GT Advantage (GA21 glyphosate-resistant trait), Agrisure (registered trademark) CB Advantage (Bt11 corn borer (CB) trait), and Protecta (registered trademark).

[0030]

The aforementioned "plants" also include

crops produced using a genetic engineering technique, which have ability to generate antipathogenic substances having selective action.

A PR protein and the like have been known as such antipathogenic substances (PRPs, EP-A-0 392 225). Such antipathogenic substances and genetically engineered crops that generate them are described in EP-A-0 392 225, WO 95/33818, EP-A-0 353 191, etc.

Examples of such antipathogenic substances expressed in genetically engineered crops include: ion channel inhibitors such as a sodium channel inhibitor or a calcium channel inhibitor (KP1, KP4 and KP6 toxins, etc., which are produced by viruses, have been known); stilbene synthase; bibenzyl synthase; chitinase; glucanase; a PR protein; and antipathogenic substances generated by microorganisms, such as a peptide antibiotic, an antibiotic having a hetero ring, a protein factor associated with resistance to plant diseases (which is called a plant disease-resistant gene and is described in WO 03/000906). These antipathogenic substances and genetically engineered plants producing such substances are described in EP-A-0392225, WO95/33818, EP-A-0353191, etc.

[0031]

The "plant" mentioned above includes plants on which advantageous characters such as characters improved in oil stuff ingredients or characters having reinforced amino acid content have been conferred by

25711-877

28

genetically engineering technology. Examples thereof include VISTIVE (registered trademark) (low linolenic soy bean having reduced linolenic content) or high-lysine (high-oil) corn (corn with increased lysine or oil content).

5 [0032]

Furthermore, stack varieties are also included in which a plurality of advantageous characters such as the classic herbicide characters mentioned above or herbicide tolerance genes, insecticidal harmful insect resistance genes,
 10 antipathogenic substance producing genes, characters improved in oil stuff ingredients or characters having reinforced amino acid content are combined.

[0033]

Examples of the diseases on which a high control
 15 effect is expected among the above include Rhizoctonia damping-off (*Rhizoctonia solani*) of wheat, corn, rice, soybean, cotton, rapeseed, sugar beet and turfgrass, damping-off and root rot of wheat, barley, corn, rice, sorghum, soybean, cotton, rapeseed, sugar beet and turfgrass caused by *Pythium* spp. (*Pythium*
 20 *aphanidermatum*, *Pythium debarianum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium ultimum*), seed diseases and diseases in the early stages of the growth of wheat, corn, cotton, soybean, rapeseed and turfgrass caused by *Fusarium* spp., pink snow mold (*Microdochium nivale*), *Rhizoctonia* damping-off (*Rhizoctonia*
 25 *solani*), *Fusarium* head blight (*Fusarium graminearum*, *F. avenacerum*, *F. culmorum*, *Microdochium nivale*) and eyespot (*Pseudocercospora herpotrichoides*) of wheat, diseases of citrus: melanose (*Diaporthe citri*) and scab (*Elsinoe fawcetti*),

25711-877

29

purple seed stain (*Cercospora kikuchii*), rust (*Phakopsora pachyrhizi*) and brown stem rot (*Phytophthora sojae*) of soybean, black shank (*Phytophthora nicotianae*) of tobacco, *Rhizoctonia* damping-off (*Rhizoctonia solani*) of cotton, *Rhizoctonia* damping-off (*Rhizoctonia solani*) and sclerotinia rot (*Sclerotinia sclerotiorum*) of rapeseed, anthracnose (*Elsinoe ampelina*), ripe rot (*Glomerella cingulata*), powdery mildew (*Uncinula necator*), black rot (*Guignardia bidwellii*) and gray mold (*Botrytis cinerea*) of grape, dollar spot (*Sclerotinia homeocarpa*) and brown patch (*Rhizoctonia solani*) of turfgrass, scab (*Venturia nashicola*, *V. pirina*) of pear, blossom blight (*Monilinia mali*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*), blotch (*Diplocarpon mali*) and ring rot (*Botryosphaeria berengeriana*) of apple, brown rot (*Monilinia fructicola*) and phomopsis rot (*Phomopsis* sp.) of peach, early leaf spot (*Cercospora arachidicola*) of peanut, gray blight (*Pestalotiopsis* sp.) and anthracnose (*Colletotrichum theae-sinensis*) of tea, *Cercospora* leaf spot (*Cercospora beticola*), leaf blight (*Thanatephorus cucumeris*), root rot (*Thanatephorus cucumeris*) and *Aphanomyces* root rot (*Aphanomyces cochlioides*) of sugar beet, sigatoka (*Mycosphaerella fijiensis*, *Mycosphaerella musicola*) of banana, blast (*Magnaporthe grisea*) and bakanae disease (*Gibberella fujikuroi*) of rice, *Rhizoctonia* damping-off (*Rhizoctonia solani*) of gourd, downy mildew (*Plasmopara halstedii*) of sunflower, late blight (*Phytophthora infestans*) and black scarf (*Rhizoctonia solani*) of potato, dollar spot (*Sclerotinia homeocarpa*), and brown patch and large patch (*Rhizoctonia solani*) of turfgrass, gray mold (*Botrytis cinerea*) and *Sclerotinia* rot (*Sclerotinia sclerotiorum*) of the other crops.

25711-877

30

[0034]

Examples of the diseases on which a particularly high control effect is expected among the above include *Rhizoctonia* damping-off (*Rhizoctonia solani*) of wheat, corn, rice, soybean, 5 cotton, rapeseed, sugar beet and turfgrass, damping-off and root rot of wheat, barley, corn, rice, sorghum, soybean, cotton, rapeseed, sugar beet and turfgrass caused by *Pythium* spp. (*Pythium aphanidermatum*, *Pythium debarianum*, *Pythium* 10 *graminicola*, *Pythium irregulare*, *Pythium ultimum*), seed diseases and diseases in the early stages of the growth of wheat, corn, cotton, soybean, rapeseed and turfgrass caused by *Fusarium* spp., brown stem rot (*Phytophthora sojae*) of soybean, black shank (*Phytophthora nicotianae*) of tobacco, downy mildew (*Plasmopara halstedii*) of sunflower, late blight (*Phytophthora infestans*) 15 and black scarf of potato, dollar spot (*Sclerotinia homeocarpa*), and brown patch and large patch (*Rhizoctonia solani*) of turfgrass, *Aphanomyces* root rot (*Aphanomyces cochlioides*) of sugar beet, gray mold (*Botrytis cinerea*) and *Sclerotinia* rot (*Sclerotinia sclerotiorum*) of the other crops.

20 EXAMPLES

[0035]


In the following, the present invention will be more specifically described by way of formulation examples, seed treatment formulation examples, and test examples. However, the 25 present invention is not limited to the following examples. In the following examples, the part represents part by weight unless otherwise noted in particular.

25711-877

31


[0036]

The compound (1a) is a compound represented by formula (1) wherein X^1 is a methyl group, X^2 is a methylamino group, and X^3 is a 2,5-dimethylphenyl group and the compound has an R type steric structure according to Cahn-Ingold-Prelog order rule, and represented by the following formula (1a).



[0037]

The compound (1b) is a compound represented by formula (1) wherein X^1 is a methyl group, X^2 is a methylamino group, and X^3 is a 2,5-dimethylphenyl group and the compound is an racemic body and represented by the following formula (1b).



25711-877

32

[0038]

Formulation example 1

2.5 Parts of the compound (1a) or the compound (1b),
1.25 parts of tolcllofos-methyl, 14 parts of polyoxyethylene
5 styrylphenyl ether, 6 parts of calcium dodecyl benzene sulfonate
and 76.25 parts of xylene are fully mixed, so as to obtain
respective emulsions.

[0039]

Formulation example 2

5 Parts of the compound (1a) or the compound (1b), 5 parts of metalaxyl, 35 parts of a mixture of 5 white carbon and a polyoxyethylene alkyl ether sulfate ammonium salt (weight ratio 1:1) and 55 parts of water are mixed, and the mixture is subjected to fine grinding according to a wet grinding method, so as to obtain respective flowable agents.

10 [0040]

Formulation example 3

5 Parts of the compound (1a) or the compound (1b), 10 parts of mefenoxam, 1.5 parts of sorbitan trioleate and 28.5 parts of an aqueous solution 15 containing 2 parts of polyvinyl alcohol are mixed, and the mixture is subjected to fine grinding according to a wet grinding method. Thereafter, 45 parts of an aqueous solution containing 0.05 part of Xanthan gum and 0.1 part of aluminum magnesium silicate is added to 20 the resultant mixture, and 10 parts of propylene glycol is further added thereto. The obtained mixture is blended by stirring, so as to obtain respective flowables.

[0041]

25 Formulation example 4

5 Parts of the compound (1a) or the compound (1b), 20 parts of tolclofos-methyl, 1.5 parts of sorbitan trioleate and 23.5 parts of an aqueous

25711-877

34

solution containing 2 parts of polyvinyl alcohol are mixed, and the mixture is subjected to fine grinding according to a wet grinding method. Thereafter, 40 parts of an aqueous solution containing 0.05 part of Xanthan gum and 0.1 part of aluminum magnesium silicate is added to the resultant mixture, and 10 parts of propylene glycol is further added thereto. The obtained mixture is blended by stirring, so as to obtain respective flowable formulations.

10 [0042]

Formulation example 5

40 Parts of the compound (1a) or the compound (1b), 5 parts of metalaxyl, 5 parts of propylene glycol (manufactured by Nacalai Tesque), 5 parts of SoprophorTMFLK (manufactured by Rhodia Nikka), 0.2 parts of an anti-form C emulsion (manufactured by Dow Corning), 0.3 parts of proxelTM GXL (manufactured by Arch Chemicals) and 49.5 parts of ion-exchange water are mixed so as to obtain a bulk slurry. 150 parts of glass beads (diameter = 1 mm) are put into 100 parts of the slurry, and the slurry is ground for 2 hours while being cooled with a cooling water. After ground, the resultant is filtered to remove the glass beads and respective flowables were obtained.

25 [0043]

Formulation example 6

50 Parts of the compound (1a) or the compound (1b), 0.5 part of mefenoxam, 38.5 parts of NN kaolin

clay (manufactured by Takehara Chemical Industrial), 10 parts of MorwetD425 and 1 part of MorwerEFW (manufactured by Akzo Nobel Corp.) are mixed to obtain an AI premix. This premix was ground with a jet mill so as to obtain respective powders.

[0044]

Formulation example 7

1 Part of the compound (1a) or the compound (1b), 4 parts of tolclofos-methyl, 1 part of synthetic hydrated silicon oxide, 2 parts of calcium lignin sulfonate, 30 parts of bentonite and 62 parts of kaolin clay are fully ground and mixed, and the resultant mixture is added with water and fully kneaded, and then subjected to granulation and drying so as to obtain respective granules.

[0045]

Formulation example 8

1 Part of the compound (1a) or the compound (1b), 40 parts of metalaxyl, 3 parts of calcium lignin sulfonate, 2 parts of sodium lauryl sulfate and 54 parts of synthetic hydrated silicon oxide are fully ground and mixed so as to obtain respective wettable powders.

[0046]

Formulation example 9

1 Part of the compound (1a) or the compound (1b), 2 parts of mefenoxam, 85 parts of kaolin clay and 12 parts of talc are fully ground and mixed so as to

obtain respective powders.

[0047]

Formulation example 10

2 Parts of the compound (1a) or the compound
5 (1b), 0.25 part of tolclafos-methyl, 14 parts of
polyoxyethylene styrylphenyl ether, 6 parts of calcium
dodecyl benzene sulfonate and 77.75 parts of xylene are
fully mixed, so as to obtain respective emulsions.

[0048]

10 Formulation example 11

10 Parts of the compound (1a) or the compound
(1b), 2.5 parts of metalaxyl, 1.5 parts of sorbitan
trioleate, 30 parts of an aqueous solution containing 2
parts of polyvinyl alcohol are subjected to fine
15 grinding according to a wet grinding method.

Thereafter, 46 parts of an aqueous solution containing
0.05 part of Xanthan gum and 0.1 part of aluminum
magnesium silicate is added to the ground solution, and
10 parts of propylene glycol is further added thereto.
20 The obtained mixture is blended by stirring, so as to
obtain respective flowables.

[0049]

Formulation example 12

1 Part of the compound (1a) or the compound
25 (1b), 20 parts of mefenoxam, 1 part of synthetic
hydrated silicon oxide, 2 parts of calcium lignin
sulfonate, 30 parts of bentonite and 47 parts of kaolin
clay are ground and mixed, and the resultant mixture is

added with water and fully kneaded, and then subjected
granulation and drying so as to obtain respective
granules.

[0050]

5 Formulation example 13

40 Parts of the compound (1a) or the compound
(1b), 1 part of tolclufos-methyl, 3 parts of calcium
lignin sulfonate, 2 parts of sodium lauryl sulfate and
54 parts of synthetic hydrated silicon oxide are fully
10 ground and mixed so as to obtain respective wettable
powders.

[0051]

Formulation example 14

2.5 Parts of the compound (1a) or the
15 compound (1b), 1.25 parts of metalaxyl, 14 parts of
polyoxyethylene styrylphenyl ether, 6 parts of calcium
dodecyl benzene sulfonate and 76.25 parts of xylene are
fully mixed, so as to obtain respective emulsions.

[0052]

20 Formulation example 15

5 Parts of the compound (1a) or the compound
(1b), 5 parts of mefenoxam, 35 parts of a mixture of
white carbon and a polyoxyethylene alkyl ether sulfate
ammonium salt (weight ratio 1:1) and 55 parts of water
25 are mixed, and the mixture is subjected to fine
grinding according to a wet grinding method, so as to
obtain respective flowable agents.

[0053]

Formulation example 16

5 Parts of the compound (1a) or the compound (1b), 10 parts of tolcllofos-methyl, 1.5 parts of sorbitan trioleate and 28.5 parts of an aqueous solution containing 2 parts of polyvinyl alcohol are mixed, and the mixture is subjected to fine grinding according to a wet grinding method. Thereafter, 45 parts of an aqueous solution containing 0.05 part of Xanthan gum and 0.1 part of aluminum magnesium silicate is added to the resultant mixture, and 10 parts of propylene glycol is further added thereto. The obtained mixture is blended by stirring, so as to obtain respective flowables.

[0054]

15 Formulation example 17

5 Parts of the compound (1a) or the compound (1b), 20 parts of metalaxyl, 1.5 parts of sorbitan trioleate and 28.5 parts of an aqueous solution containing 2 parts of polyvinyl alcohol are mixed, and the mixture is subjected to fine grinding according to a wet grinding method. Thereafter, 45 parts of an aqueous solution containing 0.05 part of Xanthan gum and 0.1 part of aluminum magnesium silicate is added to the resultant mixture, and 10 parts of propylene glycol is further added thereto. The obtained mixture is blended by stirring, so as to obtain respective flowable formulations.

[0055]

Formulation example 18

40 Parts of the compound (1a) or the compound (1b), 5 parts of mefenoxam, 5 parts of propylene glycol (manufactured by Nacalai Tesque), 5 parts of

5 SoprophorFLK (manufactured by Rhodia Nikka), 0.2 parts of an anti-form C emulsion (manufactured by Dow Corning), 0.3 parts of proxel GXL (manufactured by Arch Chemicals) and 49.5 parts of ion-exchange water are mixed so as to obtain a bulk slurry. 150 parts of

10 glass beads (diameter = 1 mm) are put into 100 parts of the slurry, and the slurry is ground for 2 hours while being cooled with a cooling water. After ground, the resultant is filtered to remove the glass beads and respective flowables were obtained.

15 [0056]

Formulation example 19a

50 Parts of the compound (1a) or the compound (1b), 0.5 part of tolclofos-methyl, 38.5 parts of NN kaolin clay (manufactured by Takehara Chemical

20 Industrial), 10 parts of MorwetD425 and 1 part of MorwerEFW (manufactured by Akzo Nobel Corp.) are mixed to obtain an AI premix. This premix was ground with a jet mill so as to obtain respective powders.

[0057]

25 Formulation example 19b

1 Part of the compound (1a) or the compound (1b), 4 parts of metalaxyl, 1 part of synthetic hydrated silicon oxide, 2 parts of calcium lignin

sulfonate, 30 parts of bentonite and 62 parts of kaolin clay are fully ground and mixed, and the resultant mixture is added with water and fully kneaded, and then subjected to granulation and drying so as to obtain
5 respective granules.

[0058]

Formulation example 20

1 Part of the compound (1a) or the compound (1b), 40 parts of mefenoxam, 3 parts of calcium lignin
10 sulfonate, 2 parts of sodium lauryl sulfate and 54 parts of synthetic hydrated silicon oxide are fully ground and mixed so as to obtain respective wettable powders.

[0059]

15 Formulation example 21

1 Part of the compound (1a) or the compound (1b), 2 parts of tolclofos-methyl, 87 parts of kaolin clay and 10 parts of talc are fully ground and mixed so as to obtain respective powders.

20 [0060]

Formulation example 22

2 Parts of the compound (1a) or the compound (1b), 0.25 part of metalaxyl, 14 parts of polyoxyethylene styrylphenyl ether, 6 parts of calcium
25 dodecyl benzene sulfonate and 77.75 parts of xylene are fully mixed, so as to obtain respective emulsions.

[0061]

Formulation example 23

41

10 Parts of the compound (1a) or the compound
(1b), 2.5 parts of mefenoxam, 1.5 parts of sorbitan
trioleate, 30 parts of an aqueous solution containing 2
parts of polyvinyl alcohol are subjected to fine
5 grinding according to a wet grinding method.
Thereafter, 46 parts of an aqueous solution containing
0.05 part of Xanthan gum and 0.1 part of aluminum
magnesium silicate is added to the ground solution, and
10 parts of propylene glycol is further added thereto.
10 The obtained mixture is blended by stirring, so as to
obtain respective flowables.

[0062]

Formulation example 24

1 Part of the compound (1a) or the compound
15 (1b), 20 parts of tolclofos-methyl, 1 part of synthetic
hydrated silicon oxide, 2 parts of calcium lignin
sulfonate, 30 parts of bentonite and 46 parts of kaolin
clay are ground and mixed, and the resultant mixture is
added with water and fully kneaded, and then subjected
20 granulation and drying so as to obtain respective
granules.

[0063]

Formulation example 25

40 Parts of the compound (1a) or the compound
25 (1b), 1 part of metalaxyl, 3 parts of calcium lignin
sulfonate, 2 parts of sodium lauryl sulfate and 54
parts of synthetic hydrated silicon oxide are fully
ground and mixed so as to obtain respective wetttable

powders.

[0064]

Seed treatment example 1

An emulsion prepared as in Formulation
5 example 1 is used for smear treatment in an amount of
500 ml per 100 kg of dried sorghum seeds using a rotary
seed treatment machine (seed dresser, produced by Hans-
Ulrich Hege GmbH) so as to obtain treated seeds.

[0065]

10 Seed treatment example 2

A flowable prepared as in Formulation example
16 is used for smear treatment in an amount of 50 ml
per 10 kg of dried rape seeds using a rotary seed
treatment machine (seed dresser, produced by Hans-
15 Ulrich Hege GmbH) so as to obtain treated seeds.

[0066]

Seed treatment example 3

A flowable prepared as in Formulation example
17 is used for smear treatment in an amount of 40 ml
20 per 10 kg of dried corn seeds using a rotary seed
treatment machine (seed dresser, produced by Hans-
Ulrich Hege GmbH) so as to obtain treated seeds.

[0067]

Seed treatment example 4

25 5 Parts of a flowable agent prepared as in
Formulation example 18, 5 parts of pigment BPD6135
(manufactured by Sun Chemical) and 35 parts of water
are mixed to prepare a mixture. The mixture is used

for smear treatment in an amount of 60 ml per 10 kg of dried rice seeds using a rotary seed treatment machine (seed dresser, produced by Hans-Ulrich Hege GmbH) so as to obtain treated seeds.

5 [0068]

Seed treatment example 5

A powder agent prepared as in Formulation example 19a is used for powder coating treatment in an amount of 50 g per 10 kg of dried corn seeds so as to
10 obtain treated seeds.

[0069]

Seed treatment example 6

An emulsion prepared as in Formulation example 22 is used for smear treatment in an amount of
15 500 ml per 100 kg of dried sugar beet seeds using a rotary seed treatment machine (seed dresser, produced by Hans-Ulrich Hege GmbH) so as to obtain treated seeds.

[0070]

20 Seed treatment example 7

A flowable prepared as in Formulation example 23 is used for smear treatment in an amount of 50 ml per 10 kg of dried soy bean seeds using a rotary seed treatment machine (seed dresser, produced by Hans-
25 Ulrich Hege GmbH) so as to obtain treated seeds.

[0071]

Seed treatment example 8

A granule agent prepared as in Formulation

example 24 is used for smear treatment in an amount of 50 ml per 10 kg of dried wheat seeds using a rotary seed treatment machine (seed dresser, produced by Hans-Ulrich Hege GmbH) so as to obtain treated seeds.

5 [0072]

Seed treatment example 9

5 Parts of a wettable powder prepared as in Formulation example 25, 5 parts of pigment BPD6135 (manufactured by Sun Chemical) and 35 parts of water
10 are mixed and the resultant mixture is used for smear treatment in an amount of 70 ml per 10 kg of potato tuber pieces using a rotary seed treatment machine (seed dresser, produced by Hans-Ulrich Hege GmbH) so as to obtain treated seeds.

15 [0073]

Seed treatment example 10

A wettable powder prepared as in Formulation example 20 is used for powder coating treatment in an amount of 40 g per 10 kg of dried cotton seeds so as to
20 obtain treated seeds.

[0074]

Test Example 1

An acetone solution of compound (1b) and an acetone solution of tolclofos-methyl were mixed so as
25 to prepare mixed liquids containing compound (1b) and tolclofos-methyl in predetermined concentration. These mixed liquids were used for smear treatment of cucumber (Sagamihanjiro) seeds using a rotary seed treatment

25711-877

45

machine (seed dresser, produced by Hans-Ulrich Hege GmbH) so as to obtain treated seeds. The treated seeds were left untouched overnight and then disseminated on the soil which filled a plastic pot and covered with the soil containing *Rhizoctonia solani*, pathogen of cucumber damping-off, which had been cultured on a bran medium. They were allowed to grow in a greenhouse while irrigated and the number of non-budding seeds was checked on the seventh day after the dissemination and the incidence of disease was calculated by Equation 3. The control value was calculated by Equation 1 based on the incidence of disease. The incidence of disease was also checked in the case in which the plants had not been treated with the agent in order to calculate the control value.

As a comparison, a compound (1b) liquid and a tolclofos-methyl liquid respectively in the predetermined concentration were prepared as the above-mentioned acetone solution and subjected to similar tests.

The results are shown in Table 2.

[0075]

"Equation 1"

Control value = $100(A - B)/A$

A: Incidence of disease of plant in untreated area

B: Incidence of disease of plant in treated area

[0076]

Generally, the control value expected for the case where the given two kinds of active ingredient compounds are mixed and used for the treatment, the so-called control value expectation is calculated from the following Colby's calculating equation.

"Equation 2"

$$E = X + Y - (X \times Y) / 100$$

X: Control value (%) when active ingredient compound A is used for treatment in M g per hectare or in M g per 100 kg of seeds

Y: Control value (%) when active ingredient compound B is used for treatment in N g per hectare or in N g per 100 kg of seeds

E: Control value (%) expected for the case where active ingredient compound A in M g per hectare or in M g per 100 kg of seeds and active ingredient compound B in N g per hectare or in N g per 100 kg of seeds are mixed and used for treatment (hereinbelow referred to as "control value expectation")

$$\text{"Synergetic effect"} = (\text{Actual control value}) \times 100 / (\text{Control value expectation})$$

[0077]

"Equation 3"

Incidence of disease = (Number of no budding seeds) \times 100 / (Number of total disseminated seeds)

[0078]

[Table 2]

Compound (1b)	Tolclofos-methyl	Actual control value	Control value expectation	Synergistic effect (%)
1g/100kg-seed	10g/100kg-seed	96	87	110.3
1g/100kg-seed	0g/100kg-seed	30	-	-
0g/100kg-seed	10g/100kg-seed	57	-	-

[0079]

Test Example 2

5 A plastic pot was filled with sandy soil. Turfgrass (bentgrass penncross) was disseminated and allowed to grow in a green house for 3 weeks. A wettable powder of compound (1b) and a wettable powder of tolclofos-methyl (Grancer Wettable Powder

10 manufactured by Nihon Green & Garden Corporation) were respectively diluted with water and then tank-mixed so as to prepare tank-mixed liquids containing compound (1b) and tolclofos-methyl in predetermined

15 concentration. The tank-mixed liquids were subjected to foliage application such that they could be sufficiently adhered to the leaves of the aforementioned turfgrass. Thereafter, the plants were air-dried. After the plants were air-dried, Sclerotinia homeocarpa, pathogen of dollar spot of

20 turfgrass, which had been cultured on a bran medium was sprayed on the plants to be treated from above. Then, the plants were placed in a greenhouse at 19°C to 23°C

under high humidity for 14 days, and thereafter control effect was checked. The diseased area ratio of the pot plants on which the agents had been sprayed was determined as the incidence of disease at the time of 5 checking and the control value calculated by Equation 1 based on the incidence of disease thus determined. In order to calculate the control value, the incidence of disease was also determined in the case in which the plants had not been treated with the agent.

10 As a comparison, the respective wettable powders described above were diluted with water in predetermined concentration so as to prepare a compound (1b) liquid and a tolclofos-methyl liquid respectively and they were subjected to similar tests.

15 The results are shown in Table 3.

[0080]

[Table 3]

Compound (1b)	Tolclofos-methyl	Actual control value (%)	Control value expectation (%)	Synergistic effect (%)
150g/ha	600g/ha	100	86.7	115.4
150g/ha	150g/ha	100	81.5	122.7
150g/ha	37.5g/ha	98.7	71.3	138.6
150g/ha	0g/ha	67.2	-	-
0g/ha	600g/ha	59.3	-	-
0g/ha	150g/ha	43.7	-	-
0g/ha	37.5g/ha	12.4	-	-

25711-877

49

[0081]

Test Example 3

A plastic pot was filled with sandy soil. Turfgrass (bentgrass penncross) was disseminated and allowed to grow in a green house for 3 weeks. A wettable powder of compound (1b) and a wettable powder of tolclofos-methyl (Grancer Wettable Powder manufactured by Nihon Green & Garden Corporation) were respectively diluted with water and then tank-mixed so as to prepare tank-mixed liquids containing compound (1b) and tolclofos-methyl in predetermined concentration. The tank-mixed liquids were subjected to foliage application such that they could be sufficiently adhered to the leaves of the aforementioned turfgrass. Thereafter, the plants were air-dried. After the plants were air-dried, *Rhizoctonia solani*, pathogen of brown patch of turfgrass, which had been cultured on a bran medium was sprayed on the plants to be treated from above. Then, the plants were placed in a greenhouse at 19°C to 23°C under high humidity for 14 days, and thereafter control effect was checked. The diseased area ratio of the pot plants on which the agents had been sprayed was determined as the incidence of disease at the time of checking and the control value was calculated by Equation 1 based on the incidence of disease thus determined. In order to calculate the control value, the incidence of disease was also determined in the

case in which the plants had not been treated with the agent.

As a comparison, the respective wettable powders described above were diluted with water in predetermined concentration so as to prepare a compound (1b) liquid and a tolclofos-methyl liquid respectively and they were subjected to similar tests.

The results are shown in Table 4.

[0082]

10 [Table 4]

Compound (1b)	Tolclofos-methyl	Actual control value	Control value expectation	Synergistic effect (%)
37.5g/ha	37.5g/ha	90	61.5	146.3
9.4g/ha	37.5g/ha	80	51	156.7
37.5g/ha	0g/ha	45	-	-
9.4g/ha	0g/ha	30	-	-
0g/ha	37.5g/ha	30	-	-

[0083]

Test Example 4

An acetone solution of compound (1b) and an acetone solution of metalaxyl were mixed so as to prepare mixed liquids containing compound (1b) and metalaxyl in predetermined concentration. These mixed liquids were adhered on the surface of cucumber (Sagamihanjiro) seeds so as to obtain treated seeds. The treated seeds were left untouched overnight and then disseminated on the soil which filled a plastic

pot and covered with the soil containing Pythium irregulare, pathogen of cucumber damping-off, which had been cultured on a bran medium. They were allowed to grow in a greenhouse while irrigated and the number of 5 non-budding seeds was checked on the seventh day after the dissemination and the incidence of disease was calculated by Equation 3. The control value was calculated by Equation 1 based on the incidence of disease. In order to calculate the control value, the 10 incidence of disease was also determined in the case in which the plants had not been treated with the agent.

As a comparison, a compound (1b) liquid and a metalaxyl liquid respectively in the predetermined concentration were prepared as the above-mentioned 15 acetone solution and subjected to similar tests.

The results are shown in Table 5.

[0084]

[Table 5]

Compound (1b)	Metalaxyl	Actual control value	Control value expectation	Synergistic effect (%)
0.005g/ 100kg-seed	0.0025g/ 100kg-seed	62	37	182
0.005g/ 100kg-seed	0.00125g/ 100kg-seed	62	49	143
0.005g/ 100kg-seed	0g/100kg-seed	25	-	-
0g/100kg-seed	0.0025g/ 100kg-seed	12.5	-	-
0g/100kg-seed	0.00125g/ 100kg-seed	25	-	-

[0085]

Test Example 5

An acetone solution of compound (1b) and an acetone solution of mefenoxam were mixed so as to
5 prepare mixed liquids containing compound (1b) and mefenoxam in predetermined concentration. These mixed liquids were adhered on the surface of cucumber (Sagamihanjiro) seeds so as to obtain treated seeds. The treated seeds were left untouched overnight and
10 then disseminated on the soil which filled a plastic pot and covered with the soil containing *Pythium irregulare*, pathogen of cucumber damping-off, which had been cultured on a bran medium. They were allowed to grow in a greenhouse while irrigated and the number of
15 non-budding seeds was checked on the seventh day after the dissemination and the incidence of disease was calculated by Equation 3. The control value was calculated by Equation 1 based on the incidence of disease. In order to calculate the control value, the
20 incidence of disease was also determined in the case in which the plants had not been treated with the agent.

As a comparison, a compound (1b) liquid and a mefenoxam liquid respectively in the predetermined concentration were prepared as the above-mentioned
25 acetone solution and subjected to similar tests.

The results are shown in Table 6.

[0086]

[Table 6]

Compound (1b)	Mefenoxam	Actual control value	Control value expectation	Synergistic effect (%)
0.005g/ 100kg-seed	0.005g/ 100kg-seed	75	63	120
0.005g/ 100kg-seed	0.00125g/ 100kg-seed	63	44	143
0.005g/ 100kg-seed	0g/100kg-seed	25	-	-
0g/100kg-seed	0.005g/ 100kg-seed	50	-	-
0g/100kg-seed	0.00125g/ 100kg-seed	25	-	-

INDUSTRIAL APPLICABILITY

[0087]


- 5 According to the present invention, a composition for controlling plant diseases having high activity, and a method for effectively controlling plant diseases, and so on can be provided.

25711-877

54

CLAIMS:

1. A composition for controlling plant fungal diseases comprising, a compound represented by formula (1b):



5 and tolclofos-methyl.

2. The composition according to claim 1, which has a weight ratio of the compound represented by formula (1b) to tolclofos-methyl falling within the range of from 0.0125:1 to 500:1.

10 3. A seed treatment agent for controlling plant fungal diseases comprising the compound represented by formula (1b) as defined in claim 1 and tolclofos-methyl.

4. Use of the compound represented by formula (1b) as defined in claim 1 and tolclofos-methyl for controlling plant
15 fungal diseases in a plant seed.

5. A method for controlling plant fungal diseases which comprises applying effective amounts of the compound represented by formula (1b) as defined in claim 1 and tolclofos-methyl to a plant or a locus where a plant is allowed to grow.

25711-877

55

6. Combined use of the compound represented by formula (1b) as defined in claim 1 and tolclofos-methyl for controlling plant fungal diseases.

