

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199892598 B2
(10) Patent No. 756471

(54) Title
Method for producing glufosinates and intermediate products for the same

(51)⁷ International Patent Classification(s)
C07F 009/30 C07F 009/6571
C07F 009/32

(21) Application No: **199892598**

(22) Application Date: **1998.08.08**

(87) WIPO No: **WO99/09039**

(30) Priority Data

(31) Number	(32) Date	(33) Country
19736125	1997.08.20	DE

(43) Publication Date : **1999.03.08**

(43) Publication Journal Date : **1999.05.06**

(44) Accepted Journal Date : **2003.01.16**

(71) Applicant(s)
Hoechst Schering Agrevo GmbH

(72) Inventor(s)
Lothar Willms

(74) Agent/Attorney
**WATERMARK PATENT and TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN
VIC 3122**




AU9892598

<p>(51) Internationale Patentklassifikation ⁶ : C07F 9/30, 9/6571, 9/32</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 99/09039 (43) Internationales Veröffentlichungsdatum: 25. Februar 1999 (25.02.99)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP98/05053 (22) Internationales Anmeldedatum: 8. August 1998 (08.08.98) (30) Prioritätsdaten: 197 36 125.0 20. August 1997 (20.08.97) DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): HOECHST SCHERING AGREVO GMBH [DE/DE]; Mirastrasse 54, D-13509 Berlin (DE). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): WILLMS, Lothar [DE/DE]; Königsteiner Strasse 50, D-65719 Hofheim (DE).</p>	<p>(81) Bestimmungsstaaten: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO Patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Veröffentlicht <i>Mit internationalem Recherchenbericht.</i> <i>Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist; Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i></p>	

(54) Title: METHOD FOR PRODUCING GLUFOSINATES AND INTERMEDIATE PRODUCTS FOR THE SAME

(54) Bezeichnung: VERFAHREN ZUR HERSTELLUNG VON GLUFOSINATE UND ZWISCHENPRODUKTE DAFÜR



(57) Abstract

According to the invention, glufosinates and their 2-methyl analogues are produced in a multi-stage synthesis from methyl phosphor compounds (II) with unsaturated keto compounds (III) via adducts (IV), subsequent reaction in Strecker synthesis conditions and finally, hydrolysis of the aminonitrile (V): Stage 1: (II) + (III) → Adduct (IV); stage 2: Adduct (IV) for example $\text{NH}_3/\text{NaCN}/\text{NH}_4\text{Cl}$ (V); stage 3: Hydrolysis of (V) to glufosinates. Various compounds are identifiable as adducts (IV), depending on the method conditions and substrates.

(57) Zusammenfassung


Glufosinate und das 2-Methylanaloge davon lassen sich in einer Mehrstufensynthese aus Methylphosphorverbindungen (II) mit ungesättigten Ketoverbindungen (III) über Addukte (IV), anschließende Umsetzung unter Bedingungen einer Strecker-Synthese und abschließende Hydrolyse des Aminonitrils (V) herstellen: Stufe 1: (II) + (III) → Addukt IV; Stufe 2: Addukt IV z.B. $\text{NH}_3/\text{NaCN}/\text{NH}_4\text{Cl}$ (V); Stufe 3: Hydrolyse von (V) zu Glufosinate. Als Addukte (IV) können je nach Verfahrensbedingungen und Substraten verschiedene Verbindungen identifiziert werden.

Abstract


Process for preparing glufosinate and phosphorus-containing α -aminonitriles as intermediates

Glufosinate and the 2-methyl analog thereof can be prepared in a multi-step synthesis from methylphosphorus compounds (II) with unsaturated keto compounds (III) via adducts (IV), subsequent reaction under the conditions of a Strecker synthesis and finally hydrolysis of the aminonitrile (V):

Step 1:




Step 2:



Step 3: Hydrolysis of (V) to give glufosinate

Depending on process conditions and substrates, various compounds can be identified as adducts (IV).



Description

PROCESS FOR PREPARING GLUFOSINATE AND INTERMEDIATES THEREFOR


5

The invention relates to the technical field of the processes for preparing biologically active compounds and precursors thereof, preferably of crop protection agents, in particular the herbicide glufosinate, also known as phosphinothricin.

10

Glufosinate (see formula (Ia)) is the common name for the active compound (D,L)-2-amino-4-[hydroxy(methyl)phosphinyl]butanoic acid, which is commercially available as monoammonium salt and is used as foliar herbicide (see DE-A-2717440, US-A-4168963).

15



20



The herbicide can be employed for the non-selective control of weeds in fruit growing and viticulture, in plantation crops, in vegetable growing prior to sowing or transplanting, prior to direct sowing of maize or soya beans, and also on uncultivated land, such as roadsides, industrial terrain and railroad tracks (cf. Z. PflKrankh.PflSchutz, Special Edition IX, 431-440, 1981). Also known is the selective use for controlling weeds in crops of useful plants, such as, inter alia, maize and rapeseed, which have been

25


made resistant by gene technology (cf. EP-A-0242246).

A large number of processes for preparing glufosinate have been disclosed. According to the variant described in EP-A-0011245 (US-A-4521348), phosphorus-containing cyanhydrin derivatives of the formula

30



in which R is a hydrocarbon radical such as alkyl, haloalkyl, cycloalkyl, phenyl or benzyl, with or without substitution, R' is hydrogen, alkyl, phenyl or benzyl and R'' is hydrogen, acyl, trialkylsilyl or alkylsulfonylalkyl, can be converted into aminonitriles, which in turn can be hydrolyzed to give
 5 glufosinate. According to EP-A-0011245, the preparation of the cyanohydrin derivatives is carried out by reaction of a monoalkyl methanephosphonate and an acroleincyanohydrin derivative of the formula




10

in which R' and R'' are as defined above. The described process has the disadvantage that the phosphorus-containing derivative and its precursors have to be provided in the form of esters, whereas in the desired product glufosinate (Ia), the (hydroxy)(methyl)phosphinyl radical is present in
 15 hydrolyzed form.

20

It is an object of the present invention to provide an alternative process to the process described above, said process allowing the number of ester precursors to be reduced and being suitable for preparing glufosinate and related compounds.

The invention provides a process for preparing compounds of the formula (I),



25

in which R* is hydrogen or (C₁-C₄)-alkyl, preferably H or methyl, or salts thereof with acids or bases, which comprises


30

a) (Step 1)

reacting a trivalent methylphosphorus compound of the formula (II) with an unsaturated derivative of the formula (III), if appropriate in the presence of a condensing agent or activator and, if appropriate, alcohols, to give an adduct (IV),



Step 1:




where in the formulae

- 5 R^1 and R^2 independently of one another are halogen, such as, for example, fluorine, chlorine, bromine or iodine, $(\text{C}_1\text{-C}_{18})$ alkoxy with or without substitution, benzyloxy or phenoxy, which may also be substituted, or one of the radicals R^1 and R^2 is hydroxyl, and
- 10 R^* is as defined in formula (I),

b) (Step 2)


the adduct (IV) is, if appropriate after hydrolytic ring opening to aldehydes ($\text{R}^* = \text{H}$) or ketones ($\text{R}^* = \text{alkyl}$) of the formula (IV') or salt thereof,

15



- in which Z is OH, R^1 or R^2 , reacted under the conditions of a Strecker synthesis with ammonia/ammonium chloride and sodium cyanide or
- 20 alternatively with mixtures of ammonia and hydrocyanic acid or with ammonia and a salt of hydrocyanic acid, such as, for example, ammonium cyanide or potassium cyanide, if appropriate in the presence of ammonium chloride, to give the α -aminonitriles of the formula (V) or a salt thereof,

25 Step 2:




where in the formulae (IV') and (V) the radical R^* is as defined in formula (I) and Z is as defined in formula (IV') or is OH, and



c) (Step 3)

the compound of the formula (V) is hydrolyzed under acidic or basic conditions to give the compound of the formula (I) or the salt thereof.


- 5 In the abovementioned formulae and in the formulae used hereinbelow, the radicals alkyl, alkoxy, haloalkyl, haloalkoxy, alkylamino and alkylthio, and also the corresponding unsaturated radicals and/or radicals which are substituted in the carbon skeleton, may in each case be straight chain or branched. Unless specifically indicated, preference for these radicals is given to the lower carbon skeletons, for example those having 1 to 4 carbon atoms and, in the case of unsaturated groups, those having 2 to 4 carbon atoms. Alkyl radicals, also in the composed meanings such as alkoxy, haloalkyl, etc., are, for example, methyl, ethyl, n- or i-propyl, n-, i-, t- or 2-butyl, pentyls, hexyls, such as n-hexyl, i-hexyl and 1,3-dimethylbutyl, 10 heptyls, such as n-heptyl, 1-methylhexyl and 1,4-dimethylpentyl; cycloalkyl is a carbocyclic saturated ring system, for example having 3 to 8 ring atoms, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.; alkenyl, alkynyl and cycloalkenyl radicals have the meaning of the possible unsaturated radicals which correspond to the alkyl or cycloalkyl radicals; 15 alkenyl is, for example, allyl, 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, but-2-en-1-yl, but-3-en-1-yl, methylbut-3-en-1-yl and 1-methylbut-2-en-1-yl; cycloalkenyl is, for example, cyclopentenyl or cyclohexenyl; alkynyl is, for example, propargyl, but-2-yn-1-yl, but-3-yn-1-yl or 1-methylbut-3-yn-1-yl. Alkenyl in the form "(C₃-C₄)-alkenyl" or "(C₃-C₆)-alkenyl" is preferably an 25 alkenyl radical having 3 to 4 and 3 to 6 carbon atoms, respectively, where the double bond is not adjacent to the carbon atom which is attached to the rest of the molecule moiety of the compound (I) ("yl" position). This applies correspondingly to (C₃-C₄)-alkynyl, etc..
- 30 Halogen is, for example, fluorine, chlorine, bromine or iodine. Haloalkyl, -alkenyl and -alkynyl are alkyl, alkenyl and alkynyl, respectively, which are partially or fully substituted by halogen, preferably by fluorine, chlorine and/or bromine, in particular by fluorine or chlorine, for example CF₃, CHF₂, CH₂F, CF₃CF₂, CH₂FCHCl₂, CCl₃, CHCl₂, CH₂CH₂Cl; haloalkoxy is, for example, OCF₃, OCHF₂, OCH₂F, CF₃CF₂O, OCH₂CF₃ and 35 OCH₂CH₂Cl; this applies correspondingly to haloalkenyl and to other halogen-substituted radicals.



If substitutions are defined by "one or more radicals selected from a group of radicals", this includes both the substitution by one or more identical radicals and mono- or polysubstitution by different radicals.

- 5 Substituted radicals, such as substituted hydrocarbon radicals, for example substituted alkyl, alkenyl, alkynyl, aryl, phenyl and benzyl, or substituted heterocyclyl, are, for example, a substituted radical derived from the unsubstituted parent radical, where the substituents are, for example, one or more, preferably 1, 2 or 3, radicals selected from the group consisting of
- 10 halogen, alkoxy, haloalkoxy, alkylthio, hydroxyl, amino, nitro, cyano, azido, alkoxy-carbonyl, alkyl-carbonyl, formyl, carbamoyl, mono- and dialkyl-aminocarbonyl, substituted amino such as acylamino, mono- or dialkyl-amino, and alkylsulfinyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl and, in the case of cyclic radicals, also alkyl and haloalkyl, and unsaturated
- 15 aliphatic radicals corresponding to the abovementioned saturated hydrocarbon-containing radicals, such as alkenyl, alkynyl, alkenyloxy, alkynyloxy etc. Preferred radicals having carbon atoms are those having 1 to 4 carbon atoms, in particular 1 or 2 carbon atoms. Preferred substituents are usually those from the group consisting of halogen, for example
- 20 fluorine and chlorine, (C₁-C₄)-alkyl, preferably methyl or ethyl, (C₁-C₄)-haloalkyl, preferably trifluoromethyl, (C₁-C₄)-alkoxy, preferably methoxy or ethoxy, (C₁-C₄)-haloalkoxy, nitro and cyano. Particular preference is given to the substituents methyl, methoxy and chlorine.
- 25 Phenyl with or without substitution is preferably phenyl which is unsubstituted or mono- or polysubstituted, preferably up to trisubstituted, by identical or different radicals selected from the group consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkyl, (C₁-C₄)-haloalkoxy and nitro, for example o-, m- and p-tolyl, dimethylphenyls, 2-, 3- and
- 30 4-chlorophenyl, 2-, 3- and 4-trifluoro- and -trichlorophenyl, 2,4-, 3,5-, 2,5- and 2,3-dichlorophenyl, o-, m- and p-methoxyphenyl.

35 An acyl radical is the radical of an organic acid, for example the radical of a carboxylic acid, and radicals of acids derived therefrom, such as the thiocarboxylic acid, iminocarboxylic acids with or without N-substitution, or the radical of carbonic acid monoesters, carbaminic acids with or without N-substitution, sulfonic acids, sulfinic acids, phosphonic acids, phosphinic acids. Acyl is, for example, formyl, alkylcarbonyl such as (C₁-C₄)-alkyl-carbonyl, phenylcarbonyl, where the phenyl ring may be substituted, for



example as shown above for phenyl, or alkyloxycarbonyl, phenyloxycarbonyl, benzyloxycarbonyl, alkylsulfonyl, alkylsulfinyl, N-alkyl-1-iminoalkyl and other radicals of organic acids.

- 5 Compounds of the formula (II) are known or can be prepared by known processes, see, for example, J.B. Miles et al. in *Org. Prep. Proc. Int.*, 11 (1), 11 (1979); B.M. Gladshstein et al., *Zh. Obshch. Khim.* 39, 1951 (1969); DAS 1098940 (1959), *Farbf. Bayer, Boetzel et al.*, *J. Fluorine Chem.* 68, 11 (1994); Hoffmann et al., *JACS* 80, 1150 (1958).

10

In the compounds of the formula (II), R^1 and R^2 independently of one another are preferably halogen, such as, for example, fluorine, chlorine, bromine or iodine, (C_1-C_6) alkoxy, (C_1-C_6) haloalkoxy, benzyloxy or phenoxy, where each of the two last-mentioned radicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, alkyl, haloalkyl, alkylthio, nitro, cyano, alkylsulfonyl and haloalkylsulfonyl, preferably in each case having 1 to 6 carbon atoms, in particular 1 to 4 carbon atoms, in the alkyl moiety, or one of the radicals R^1 and R^2 is preferably hydroxyl.

20


Particularly preferably, R^1 and R^2 are each (C_1-C_4) alkoxy.

The compounds of the formula (III) are basic chemicals and therefore also known.

25

The adducts (IV) may have various structures. Intermediates which are possible in some cases are 2-methyl-1,2-oxa-4-phospholenes of the formula (IV*), i.e. the subsequent reactions are consistent with an intermediate of the formula (IV*):

30




In certain cases, the compounds of the formula (IV*) occur as intermediates which cannot be detected, or do not occur as intermediates at all, depending on which activators or condensing agents or reactive additives such as alcohols are employed in the addition/condensation reaction.

35



In a preferred embodiment, preference is given to reacting compounds of the formula (II-1) with a compound of the formula (III) in the presence of anhydrides A_2O , preferably carboxylic anhydrides, and alcohols ROH, to give adducts (IV-1), the latter being semiacetals or a salt thereof,

5



in which

-O-X and -O-Y correspond to the radicals R^1 and R^2 , respectively, if these
10 are radicals of alcohols, i.e.

each of the radicals X and Y independently of one another is H or (C₁-C₁₈)-alkyl which is unsubstituted or substituted, benzyl or phenyl, where
each of the two abovementioned radicals is unsubstituted or substituted,
15 preferably unsubstituted or substituted by one or more radicals selected
from the group consisting of halogen, alkyl, haloalkyl, alkylthio, nitro,
cyano, alkylsulfonyl and haloalkylsulfonyl, preferably in each case having 1
to 6 carbon atoms, in particular 1 to 4 carbon atoms in the alkyl moiety,
and

X and Y are preferably identical radicals, and in particular X, Y and R are
20 identical radicals,

R^* is as defined in formula (I), preferably H,


A is an acyl radical, preferably the acyl radical of a carboxylic acid having 1
to 6 carbon atoms, in particular 1 to 4 carbon atoms,



R is a radical selected from the group of the radicals defined for X and Y,
25 preferably the same radical as X or Y.

Particularly preferably

X, Y and R are in each case identical radicals selected from the group
consisting of (C₁-C₆)alkyl, phenyl or benzyl, in particular (C₁-C₄)alkyl, for
30 example methyl, ethyl, n-, i-propyl, n-, i-, s- or t-butyl.

Correspondingly, the compounds (IV') and (V) are, in the preferred variant
(starting from compounds (II-1)), compounds of the formula (IV'-1) and
(V-1) or salts thereof, respectively,






5


in which X' = H or X and R* and X are as defined above, or salts thereof.

2-Methyl-1,2-oxa-4-phospholenes of the formula (IV*) and the semiacetals
 10 of the formula (IV-1) have hitherto been unknown, as have been the
 aminonitriles of the formula (V-2) (= formula (V) where Z = OH)




15 and they therefore also form part of the subject matter of the present
 invention.

From the 1,2-oxa-4-phospholenes (IV*), only some higher homologs are
 known. Thus, phenyldichlorophosphane (IIa) reacts with α,β -unsaturated
 20 ketones (VI) with addition of acetic anhydride to give the 2-phenyl-2-oxo-
 1,2-oxa-4-phospholenes (VII) (K. Bergesen, Acta Chem. Scand. 19, 1784
 (1965)),




25




in which R^3 and R^4 are hydrogen, methyl or phenyl and R^5 is methyl or phenyl.

Furthermore, it is known that ethyldichlorophosphane IIb reacts with methyl vinyl ketone (VIa) to give 5-methyl-2-ethyl-2-oxo-1,2-oxa-4-phospholene (VIIa) (A.N. Pudovik et al., *Isv. Akad. Nauk. SSSR, Ser. Khim.* (Engl. version) 2543 (1970));

Et = ethyl in formula IIb; Me = methyl in formula VIIa; Ac = acetyl;



Finally, the reaction of 2-thienyldichlorophosphane (IIc) with α,β -unsaturated ketones gives 2-thienyl-2-oxo-1,2-oxa-4-phospholenes (VIIb) (R.Z. Aliev, *Isv. Akad. Nauk., SSSR, Ser. Khim* (Engl. version), 2719 (1973)),



in which R^6 is hydrogen or methyl.

20

Analogous reactions, for example with methyldichlorophosphane which is highly reactive compared to phenyldichlorophosphane (cf. H. Heydt et al., *Methoden der Organischen Chemie XII E2*, p. 29 (1982)) and methanephosphonous acid diesters have hitherto not been described in the literature. Analogous reactions with acrolein (III; $R^2 = \text{H}$) are likewise not known.

25

Because the components (II) and (III) are much more reactive, and because of the complex reaction mixture or the complex course of the reaction in step 1, it is extremely surprising that the process according to the invention can be realized in high yields via the intermediates (adducts

30



IV) to give the α -aminonitrile derivatives (V) or (V-1) and subsequently the compounds (I).

5 In step 1, the process according to the invention is generally carried out by reacting compounds of the formula (II) or (II-1) with unsaturated compounds of the formula (III), preferably in the presence of a condensing agent or activator. Suitable activators/condensing agents are substances which are suitable for promoting or catalyzing the addition of the phosphorus component to the α,β -unsaturated keto compound (III).
10 Suitable condensing agents or activators are carboxylic anhydrides, preferably anhydrides of alkanecarboxylic acids having 1 to 6 carbon atoms, for example acetic anhydride or propionic anhydride.

15 Also suitable are mixtures of the anhydrides with certain proportions of alcohols ROH, where R is as defined above.

The reaction of compounds (II) and (III) can be carried out without solvent or in the presence of an organic solvent, for example in the presence of aliphatic or aromatic hydrocarbons which may be halogenated, such as
20 dichloromethane, toluene, xylene, chlorobenzene, or ethers, such as dioxane, or alcohols, such as ethanol, n-butanol, etc., or mixtures of these exemplary solvents.

25 The phosphorus components of the formula (II) are employed in molar ratios which can deviate considerably from the stoichiometry, preferably in molar ratios of 1:2 to 2:1, but in particular essentially in equimolar amounts, based on the component (III).

30 If the reaction of the components (II) and (III) is carried out in the presence of an anhydride A_2O , such as, for example, acetic anhydride or propionic anhydride, suitable anhydride inputs are usually in the range of from more than 0 to 400 mol%, preferably amounts of from 50 to 150 mol%, based on the starting component (II) or (III), which is employed in the lowest molar amount.

35

If the reaction of the components (II-1) and (III) is carried out in the presence of the anhydride A_2O , for example acetic anhydride, and an alcohol ROH, for example (C₁-C₅)alkanol such as ethanol, preference is given to using 50 to 150 mol% of acetic anhydride and 50 to 200 mol% of



alcohol, in particular in the anhydride:alcohol ratio of 1:1 to 1:1.5, based on the starting component (II) or (III), which is employed in the lowest molar amount.

- 5 The reaction, according to the invention, of compounds (II) and (III) succeeds generally at reaction temperatures between -80°C and $+200^{\circ}\text{C}$, preferably between -10°C and $+60^{\circ}\text{C}$. The duration of the reaction depends in general on the reaction temperature, the size of the batch, the specific reactants, the solvent and the condensing agents/activators and is,
10 for example, in the range of 0.5 - 48 hours (h), preferably 0.5 - 18 h.


Surprisingly, the reaction, according to the invention, of the intermediates (IV) and (IV-1) to give the desired α -aminonitriles (V) and (V-1), respectively, (step 2) can be carried out under conditions which are known
15 analogously to the preparation of aminonitriles from aldehydes or ketones by the type of the "Strecker synthesis" (see textbooks and handbooks of organic chemical synthesis). According to one possible procedure, the reaction solution which contains the crude product (IV) or (IV-1) is added to a solution or suspension comprising an alkali metal cyanide and
20 ammonium chloride in aqueous ammonia solution.

It is also possible to employ mixtures of the abovementioned organic solvents, such as, for example, toluene, xylene, chlorobenzene, dichloromethane, ethanol, butanol etc., for this purpose. Instead of alkali metal cyanides, it is also possible to use alkaline earth metal cyanides or
25 ammonium cyanide, or solutions of hydrocyanic acid in ammonia.

The cyanides or the hydrocyanic acid are employed, for example, in amounts of 80 - 130 mol%, but preferably in essentially equimolar amounts, based on the components of the formula (IV). The amount of
30 ammonia based on the compound (IV) is, for example, between 100 and 800 mol%, preferably from 100 to 400 mol%.

The reactions of the compounds (IV) under the conditions of the Strecker synthesis are carried out, for example, at from -10°C to 100°C , preferably at $0-45^{\circ}\text{C}$.
35

The compounds of the formula (V) or (V-1) are preferably obtained as salts in which the acidic hydrogen atom at the phosphinoyl group is replaced by a cation equivalent, preferably by a cation equivalent such as, for example, Li^+ , Na^+ , K^+ , $(\text{Mg}^{2+})_{1/2}$, $(\text{Ca}^{2+})_{1/2}$, NH_4^+ .




Alternatively, it is possible to initially purify the intermediates (IV) or (IV-1) by distillation or extractive methods and to react them in purified form to give the aminonitriles (V) or (V-1).

5 In a further variant, the intermediates (adducts IV) or (IV-1) are initially hydrolyzed with water to give the aldehydes or ketones of the formula (IV') or (IV-1) and reacted in a further step to give the α -aminonitriles (V) or (V-1).

10 In the formula (IV'), R* is hydrogen or (C₁-C₄)-alkyl. The compound where R* = hydrogen and Z = hydroxyl or salts thereof are novel compounds in the methylphosphinic acid series and therefore also form part of the subject matter of the invention, i.e. the compound of the formula (IV'-2) or salts thereof

15



However, the compound (IV) where R* = methyl and Z = hydroxyl is known (L.D. Quin et al., J. Org. Chem. 39, 686 (1974)).

20

According to step 3 of the process according to the invention, the α -aminonitriles of the formula (V) or (V-1) are useful intermediates which, in analogy with the process conditions known from the literature (Houben - Weyl, Methoden der Organischen Chemie XI/2, p. 305 and p. 371, 1958),

25 can be hydrolyzed both in acidic and in basic medium, to give the biologically active amino acids of the formula (I), in particular glufosinate of the formula (Ia).

30 Compared with known processes for the synthesis of the herbicidal amino acid (Ia), the process according to the invention has a number of advantages, for example, the additional esterification of the phosphorus components of the intermediates (IV), (IV') and (V) of the intermediates is unnecessary. Moreover, the process can optionally be carried out separately for each step, or as a one-pot process over all 3 steps.

35

Thus, the essential PC linkage for building up the amino acid side chain can be carried out in one step, for example with methyldihalophosphanes



or, preferably, methanephosphonous acid diesters (II) or (II-1) and olefins (III), without the complicated conversion of, for example, methyldichlorophosphane to methanephosphonous acid monoesters being necessary. Moreover, in contrast to the process known from EP-A-0011245, the radical addition of the methanephosphonous acid monoesters to acrolein derivatives, which readily leads to by-products, is avoided.

In the process according to the invention it is possible to employ, for example, the readily obtainable olefin components acrolein or methyl vinyl ketone directly, without derivatization being necessary. Furthermore, the α -aminonitriles (V) or (V-1) are obtained in the process according to the invention with a free phosphinic acid or phosphinate grouping, so that in the last step of the synthesis only the nitrile group has to be hydrolyzed to give the free amino acid. Deblocking of the phosphinic ester group to the free phosphinic acid, which is required in the prior art method mentioned above, is thus superfluous.

The examples below illustrate the process, without limiting the possible process conditions. Unless specifically defined otherwise, the amounts stated are based on weight.



Example 1

2-Amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

5 At room temperature and under an atmosphere of inert gas, 7.01 g (0.10 mol) of methyl vinyl ketone are admixed with 10.21 g (0.10 mol) of acetic anhydride. With cooling at at most 25-30°C, 13.61 g (0.10 mol) of diethyl methanephosphonate are subsequently added dropwise. The reaction mixture is stirred at 30°C for approximately 6 hours. At 20-25°C,
10 the mixture is then added dropwise to a solution of 4.41 g (0.09 mol) of sodium cyanide and 9.63 g (0.18 mol) of ammonium chloride in 50 ml of ammonia solution (25% strength). The mixture is stirred at 25°C for another 4 hours and the crude aminonitrile is then rapidly added dropwise without isolation to 200 ml of hydrochloric acid (37% strength). The
15 reaction mixture is subsequently boiled under reflux for approximately 4 hours, while ethanol and acetic acid are distilled off. The mixture is concentrated using a rotary evaporator, a pH of approximately 9 is set using ammonia solution and the desired product is freed of salts by recrystallization from methanol.

20

This gives 19.1 g (corresponding to 94.5% of theory) of 2-amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt.

¹H NMR (D₂O) : 1.56 (d, J = 14Hz, 3H); 1.63 (s, 3H); 1.7-2.3 (m, 4H)


³¹P NMR (D₂O): 54.4.

25

Example 2

2-Amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

At room temperature, 2.10 g (0.03 mol) of methyl vinyl ketone and
30 3.06 g (0.03 mol) of acetic anhydride are dissolved in 20 ml of dichloromethane. At 25-28°C, 3.51 g (0.03 mol) of methyldichlorophosphane are subsequently rapidly added dropwise and the mixture is stirred at approximately 30°C for 3 hours and then added dropwise to a solution of 1.375 g (0.0275 mol) of sodium cyanide and 2.94 g (0.055 mol) of
35 ammonium chloride in 25 ml of ammonia (25% strength). The mixture is stirred at 28-30°C for approximately 4 hours and the two-phase crude aminonitrile solution is added dropwise at 25-30°C to 100 ml of hydrochloric acid (37% strength). The mixture is subsequently heated under reflux for approximately 4 hours and worked-up as under Example 1.



This gives 5.83 g (corresponding to 92% of theory) of 2-amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt.

^1H NMR (D_2O) : 1.57 (d, $J = 14$ Hz, 3H); 1.65 (s, 3H); 1.7-2.3 (m, 14H).

^{31}P NMR : 54.5

5

Example 3

2-Amino-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

At room temperature, 5.61 g (0.10 mol) of freshly distilled acrolein are added to 10.21 g (0.10 mol) of acetic anhydride. At 25-30°C, 13.61 g (0.10 mol) of diethyl methanephosphonate are subsequently added dropwise. The mixture is stirred at 30°C for 2 hours and then, at 25-28°C, added dropwise to a solution of 4.9 g (0.10 mol) of sodium cyanide and 10.7 g (0.20 mol) of ammonium chloride in 50 ml of ammonia (25% strength). After 2 hours at 30°C, the crude aminonitrile is added dropwise to 200 ml of hydrochloric acid (37% strength). The mixture is subsequently heated under reflux for 2 hours, while ethanol and acetic acid are distilled off. The mixture is concentrated using a rotary evaporator, a pH of approximately 9 is set using ammonia solution and the product is purified by crystallization from methanol. This gives 19.4 g (98% of theory) of 2-amino-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt.

10

15

20

^1H NMR (D_2O) : 1.60 (d, 14Hz, 3H); 1.8-2.4 (m, 4H); 4.28 (t, $J = 6$ Hz, 1H).

^{31}P NMR (D_2O) : 55.9.

25


Example 4

2-Amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

At room temperature and under an atmosphere of inert gas, 14.02 g (0.20 mol) of methyl vinyl ketone are admixed with 20.42 g (0.20 mol) of acetic anhydride. With cooling and at at most 25 to 30°C, a mixture of 27.22 g (0.20 mol) of diethyl methanephosphonate and 9.2 g (0.2 mol) of ethanol is subsequently added dropwise. The reaction mixture is stirred at 30°C for approximately 6 hours. At 20 to 25°C, the mixture is then added dropwise to a solution of 8.82 g (0.18 mol) of sodium cyanide and 19.26 g (0.36 mol) of ammonium chloride in 100 ml of ammonia solution (25% strength). The mixture is stirred at 25°C for another 4 hours and the crude aminonitrile is then rapidly added dropwise without isolation to 400 ml of hydrochloric acid (37% strength). The reaction mixture is subsequently boiled under reflux for approximately 4 hours, while ethanol and acetic acid

30

35



are distilled off. The mixture is concentrated using a rotary evaporator, a pH of approximately 9 is set using ammonia solution and the desired product is freed of salts by recrystallization from methanol.

- 5 This gives 38.8 g (corresponding to 96% of theory) of 2-amino-2-methyl-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt (physical data see Ex. 1).

Example 5

- 10 2-Amino-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

At room temperature, 5.61 g (0.10 mol) of freshly distilled acrolein are added to 10.21 g (0.10 mol) of acetic anhydride. At 25 to 30°C, this mixture is subsequently added dropwise to 13.61 g (0.10 mol) of diethyl methane-
15 phosphonate and 4.6 g (0.1 mol) of ethanol. The mixture is stirred at 30°C for 2 hours and then, at 25 to 28°C, added dropwise to a solution of 4.9 g (0.10 mol) of sodium cyanide and 10.7 g (0.20 mol) of ammonium chloride in 50 ml of ammonia (25% strength). After 2 hours at 30°C, the crude aminonitrile is added dropwise to 200 ml of hydrochloric acid (37%
20 strength). The mixture is subsequently heated under reflux for 2 hours, while ethanol and acetic acid are distilled off. The mixture is concentrated using a rotary evaporator, a pH of approximately 9 is set using ammonia solution and the product is purified by crystallization from methanol. This gives 19.6 g (99% of theory) of 2-amino-4-(hydroxymethylphosphinyl)-
25 butyric acid, ammonium salt.

¹H NMR (D₂O): 1.60 (d, 14Hz, 3H); 1,8-2.4 (m, 4H); 4.28 (t, J = 6 Hz, 1H).
³¹P NMR (D₂O): 55.9.

Example 6

- 30 2-Amino-4-(hydroxymethylphosphinyl)butyric acid, ammonium salt

At room temperature, 5.61 g (0.10 mol) of freshly distilled acrolein are added to 10.21 g (0.10 mol) of acetic anhydride. At 25 to 30°C, this mixture is subsequently added dropwise to 16.41 g (0.10 mol) of dibutyl methane-
35 phosphonate and 14.8 g (0.2 mol) of n-butanol. The mixture is stirred at 30°C for 2 hours and then, at 25 to 28°C, added dropwise to a solution of 4.9 g (0.10 mol) of sodium cyanide and 10.7 g (0.20 mol) of ammonium chloride in 50 ml of ammonia (25% strength). After 2 hours at 30°C, the crude aminonitrile is added dropwise to 200 ml of hydrochloric acid (37%




strength). The mixture is subsequently heated under reflux for 2 hours, while ethanol and acetic acid are distilled off. The mixture is concentrated using a rotary evaporator, a pH of approximately 9 is set using ammonia solution and the product is purified by crystallization from methanol. This gives 17.8 g (90% of theory) of 2-amino-4-(hydroxymethylphosphinyl)-butyric acid, ammonium salt.

^1H NMR (D_2O) : 1.60 (d, 14Hz, 3H); 1.8 – 2.4 (m, 4H); 4.28 (t, J = 6 Hz, 1H).


^{31}P NMR (D_2O) : 55.9.

Comprises/comprising and grammatical variations thereof when used in this specification are to be taken to specify the presence of stated features, integers, steps or components or groups thereof, but do not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A process for preparing compounds of the formula I




in which R* is hydrogen or (C₁-C₄)-alkyl, or salts thereof with acids or bases, which comprises

- a) (Step 1)

reacting a trivalent methylphosphorus compound of the formula (II) with an unsaturated derivative of the formula (III) in the presence of a condensing agent or activator and, optionally, alcohols, to give an adduct (IV),

Step 1:





where in the formulae

R¹ and R² independently of one another are (C₁-C₁₆)alkoxy with or without substitution, benzyloxy or phenoxy, which may also be substituted, or one of the radicals R¹ and R² is hydroxyl, and R* is as defined in formula (I),

- b) (Step 2)


the adduct (IV) is, optionally after hydrolytic ring opening to aldehydes (R* = H) or ketones (R* = alkyl) of the formula (IV'),





in which Z is OH, R¹ or R², reacted under the conditions of a Strecker synthesis with ammonia/ammonium chloride and sodium cyanide or alternatively with mixtures of ammonia and hydrocyanic acid or with ammonia and a salt of hydrocyanic acid, optionally in the presence of ammonium chloride, to give the α-aminonitriles of the formula (V) or a salt thereof,

Step 2:



where in the formulae (IV') and (V) the radical R* is as defined in formula (I) and Z is as defined in formula (IV') or is OH, and

c) (Step 3)


the compound of the formula (V) is hydrolyzed under acidic or basic conditions to give the compound of the formula (I).

2. The process as claimed in claim 1, wherein R¹ and R² independently of one another are halogen, (C₁-C₆)alkoxy, (C₁-C₆)haloalkoxy, benzyloxy or phenoxy, where each of the two last-mentioned radicals is unsubstituted or substituted by one or more radicals selected from the group consisting of halogen, alkyl, haloalkyl, alkylthio, nitro, cyano, alkylsulfonyl and haloalkylsulfonyl having in each case 1 to 6 carbon atoms in the alkyl moiety, or one of the radicals R¹ or R² is hydroxyl.

3. The process as claimed in claim 1 or 2, wherein in step 1, as compounds (II), compounds of the formula (II-1) are reacted with a compound of the formula (III) in



the presence of anhydrides A_2O and alcohols ROH to give adducts (IV) of the formula (IV-1),



in which

each of the radicals X and Y independently of one another is H or (C_1-C_{18}) alkyl which is unsubstituted or substituted, benzyl or phenyl, where each of the two abovementioned radicals is unsubstituted or substituted, and A is an acyl radical,

R is (C_1-C_{18}) alkyl, which is unsubstituted or substituted, benzyl or phenyl, where each of the two radicals above is unsubstituted or substituted.


4. The process as claimed in claim 3, wherein

X, Y and R are each (C_1-C_4) alkyl,


A is an acyl radical of an alkanecarboxylic acid having 1 to 6 carbon atoms and

R^* is a hydrogen atom.

5. A process for preparing compounds of the formula (I),




in which R^* is hydrogen or (C_1-C_4) -alkyl, or salts thereof with acids or bases, which comprises hydrolyzing a compound of the formula (V-2) or salts thereof,



in which R* is as defined in formula (I)

under acidic or basic conditions to give the compound of the formula (I) or salts thereof.


6. The process as claimed in claim 5, wherein the compound of the formula (V-2) is prepared from compounds of the formula (IV*),



(IV)

in which R* is as defined in formula (V-2),


optionally after hydrolytic ring opening to aldehydes or ketones of the formula (IV'),



(IV')

in which Z = OH and R* is as defined in formula (V-2), by reaction under the conditions of a Strecker synthesis with ammonia/ammonium chloride and sodium cyanide or alternatively with mixtures of ammonia and hydrocyanic acid or with ammonia and a salt of hydrocyanic acid, optionally in the presence of ammonium chloride, to give the α -aminonitriles of the formula (V-2) or a salt thereof.


7. A process for preparing compounds of the formula (V) or salts thereof,



(V)


in which Z and R* are as defined in step 2 of claim 1, which comprises reacting compounds of the formula (IV-1),






in which

X, A and R are as defined in claim 3 and R* is as defined in formula (V),
optionally after hydrolytic ring opening to aldehydes or ketones of the formula (IV'),



in which Z = OH or OX, X is as defined in formula (IV-1) and R* is as defined in formula (V), by reaction under the conditions of a Strecker synthesis with ammonia/ammonium chloride and sodium cyanide or alternatively with mixtures of ammonia and hydrocyanic acid or with ammonia and a salt of hydrocyanic acid, optionally in the presence of ammonium chloride, to give the α -aminonitriles of the formula (V) or a salt thereof.


8. A compound of the formula (IV-1),




in which

X, A and R are as defined in claim 3 and R* is H or (C₁-C₄)-alkyl, or salts thereof.

9. The process for preparing compounds of the formula (IV-1) as claimed in claim 8, wherein compounds of the formula (II-1) are reacted with a compound of the formula (III) in the presence of anhydrides A₂O and alcohols ROH to give the adducts of the formula






In which

each of the radicals X and Y independently of one another is H or (C₁-C₁₈)alkyl which is unsubstituted or substituted, benzyl or phenyl, where each of the two abovementioned radicals is unsubstituted or substituted, and A is an acyl radical,


R is (C₁-C₁₈)alkyl which is unsubstituted or substituted, benzyl or phenyl, where each of the two radicals above is unsubstituted or substituted.

10. The process as claimed in claim 9, wherein from 50 to 150 mol% of anhydride A₂O and from 50 to 200 mol% of alcohol ROH, based on the starting component (II-1) or (III) which has the lowest molarity, are employed.

11. A process for preparing compounds of the formula (I),



In which R^{*} is hydrogen or (C₁-C₄)-alkyl, or salts thereof with acids or bases, which comprises reacting compounds of the formula (IV-1),




R^* is as defined in formula (I),


each of the radicals X and Y independently of one another is H or (C₁-C₁₈)alkyl which is unsubstituted or substituted, benzyl or phenyl, where each of the two abovementioned radicals is unsubstituted or substituted, and

A is an acyl radical,

R is (C₁-C₁₈)alkyl which is unsubstituted or substituted, benzyl or phenyl, where each of the two abovementioned radicals is unsubstituted or substituted, optionally after hydrolytic ring opening to aldehydes ($R^* = H$) or ketones ($R^* = \text{alkyl}$) of the formula (IV'),



in which Z is OH, or OX, or to salts thereof under the conditions of a Strecker synthesis with ammonia/ammonium chloride and sodium cyanide or alternatively with mixtures of ammonia and hydrocyanic acid or with ammonia and a salt of hydrocyanic acid, optionally in the presence of ammonium chloride, to give the α -aminonitriles of the formula (V) or a salt thereof,



where in the formulae (IV') and (V) the radical R^* is as defined in formula (I) and Z is as defined in formula (IV') or is OH, and

the compound of the formula (V) is hydrolyzed under acidic or basic conditions to give the compound of the formula (I).

