

(19) **DANMARK**

(10) **DK/EP 2498756 T3**



Patent- og
Varemærkestyrelsen

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **A 61 K 9/20 (2006.01)** **A 61 K 31/4709 (2006.01)** **C 07 D 401/12 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2019-11-25**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-09-04**
- (86) Europæisk ansøgning nr.: **10779339.0**
- (86) Europæisk indleveringsdag: **2010-11-02**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-09-19**
- (86) International ansøgning nr.: **IB2010054964**
- (87) Internationalt publikationsnr.: **WO2011055303**
- (30) Prioritet: **2009-11-09 US 259403 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Wyeth LLC, 235 East 42nd Street, New York, NY 10017, USA**
- (72) Opfinder: **ASHRAF, Muhammad, 67 Boulevard, Elmwood Park, New Jersey 07407, USA**
GHOSH, Krishnendu, Pfizer Inc., 401 N. Middletown Road Building 140-4th Fl., Pearl River, New York 10965, USA
GOOLCHARRAN, Chimanlall, Pfizer Global Research&Development, Eastern Point Road, Groton, Connecticut 06340, USA
MAHMUD, Mainuddin, 12 Meredith Court, Oak Ridge, New Jersey 07438, USA
NAGI, Arwinder Singh, 12366 Nelson Road, Moorpark, California 93021, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **TABLETFORMULERINGER AF NERATINIBMALEAT**
- (56) Fremdragne publikationer:
WO-A1-2005/037287
WO-A1-2008/093878
WO-A2-2006/116514
WO-A2-2007/130438
WO-A2-2008/089087
WO-A2-2011/002857

DESCRIPTION

[0001] This application claims the benefit of United States Application No. 61/259,403, filed November 9, 2009.

FIELD OF THE INVENTION

[0002] The present invention relates to oral pharmaceutical formulations of neratinib provided in the form of coated tablets prepared by fluid bed granulation or by wet granulation, and improved methods of making these coated tablets.

BACKGROUND OF THE INVENTION

[0003] Protein kinases are important in the transmission of biochemical signals, which initiate cell replication. Protein kinases are enzymes that catalyze the transfer of a phosphate group from ATP to an amino acid residue, such as tyrosine, serine, threonine, or histidine on a protein. Regulation of these protein kinases is essential for the control of a wide variety of cellular events including proliferation and migration. Specific protein kinases have been implicated in adverse conditions including cancer [Traxler, P. M., Exp. Opin. Ther. Patents, 8, 1599 (1998); Bridges, A. J., Emerging Drugs, 3, 279 (1998)], restenosis [Mattsson, E., Trends Cardiovas. Med. 5, 200 (1995); Shaw, Trends Pharmacol. Sci. 16, 401 (1995)], atherosclerosis [Raines, E. W., Bioessays, 18, 271 (1996)], angiogenesis [Shawver, L. K., Drug Discovery Today, 2, 50 (1997); Folkman, J., Nature Medicine, 1, 27 (1995)] and osteoporosis [Boyce, J. Clin. Invest., 90,1622 (1992)]. Compounds capable of inhibiting the activity of receptor tyrosine kinases are known to be useful in the treatment of cancers, including but not limited to for example, non-small cell lung cancer (NSCLC), breast cancer, polycystic kidney disease, colonic polyps, and stroke in mammals. A specific kinase inhibitor is (E)-N-(4-(3-chloro-4-(pyridin-2-ylmethoxy)phenylamino)-3-cyano-7-ethoxyquinolin-6-yl)-4-(dimethylamino)but-2-enamide, also known as neratinib. Nerartinib is a weak base having low bioavailability and low solubility in both water and alcohol.

[0004] Neratinib maleate particles exhibit very high surface free energy, (work of cohesion = 45.62 mN / m). This property renders the primary particles very cohesive and prone to aggregation as described by B. Janczuk and T.Bialopiotrowicz, "Surface Free-Energy Components of Liquids and Low Energy Solids and Contact Angles," in J. Colloid Interf. Sci. 127 (1989), p.189-204; W.R. Good, "A Comparison of Contact Angle Interpretations," in J. Colloid Interf. Sci. 44 (1973), p. 63; M.D. Lechner (Ed.), Landolt Börnstein, New Series, Vol. IV/16, "Surface Tension of Pure Liquids and Binary Liquid Mixture," Springer Verlag, 1998.; and J.J. Jasper, "The Surface Tension of Pure Liquid Compounds," in J. Phys. Chem. Ref. Data, Vol.1, No. 4, 1972, p. 859. As a consequence of cohesiveness, neratinib maleate powder does

not lend itself easily to pharmaceutical operations such as mixing, flow or fluidization especially when it constitutes a high proportion in a composition. Due to these limitations, it was not possible to develop a neratinib maleate formulation comprising a capsule or tablet of higher strength employing a direct compression or roller compaction processes successfully. A formulation using a conventional wet granulation method led to chemical degradation and stability issues.

SUMMARY OF THE INVENTION

[0005] It is desirable to provide a neratinib maleate formulation, where the surface property of the active ingredient is modified by spraying or otherwise applying a substance, such as a polymer like povidone, of low surface energy (for instance about 38 mN/m) on the surface of neratinib maleate particles.

[0006] The present invention provides pharmaceutically acceptable solid compositions suitable for oral administration comprising the active ingredient neratinib maleate. In certain embodiments, such solid compositions are provided in the form of coated tablets prepared by fluid bed granulation. In some embodiments, the present invention provides a unit dosage form comprising neratinib maleate.

[0007] The present invention provides a pharmaceutically acceptable composition comprising: a granulation comprising intragranular components: (a) 10-70 weight percent neratinib maleate; (b) 15-65 weight percent of one or more fillers; (c) 0-8 or 0.5-8 weight percent of one or more disintegrants; and (d) 0.2-8 weight percent, in certain embodiments 0.2-6 weight percent, of one or more glidants; and (e) 5-15 weight percent of one or more surface modifying agents. The granulation is combined with extragranular components (f) 1-25 or 4-25 weight percent of one or more fillers; (g) 1-8 or 0-8 weight percent of one or more disintegrants and (h) 0.1-3 or 0.5-3 weight percent of one or more lubricants, and then compressed into tablets or dry-filled into capsules.

[0008] The present invention provides a pharmaceutically acceptable composition comprising: a granulation comprising intragranular components (a) 10-70 weight percent of neratinib maleate; (b) 15-65 weight percent of mannitol and microcrystalline cellulose; (c) 0.5-8 weight percent of crospovidone or croscarmellose sodium; and (d) 0.2-8 weight percent, in certain embodiments 0.2-6 weight percent, of colloidal silicon dioxide, and (e) 5-15 weight percent of povidone. The granulation is combined with extragranular components (f) 1-25 or 4-25 weight percent of microcrystalline cellulose; (g) 0-8 or 1-80-8 weight percent of crospovidone or croscarmellose sodium and (h) 0.1-3 or 0.5-3 weight percent of magnesium stearate, and then compressed into tablets or dry-filled into capsules.

[0009] The present invention also provides methods of preparing stable, pharmaceutically acceptable neratinib-maleate formulations for oral administration comprising components described above and herein, which allow for improved processing characteristics while

maintaining acceptable pharmacokinetic properties.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] Figure 1 summarizes mean pharmacokinetic parameters versus time for neratinib maleate following the administration of immediate release tablet formulations of neratinib with different dissolution rates. TR refers to a rapid dissolving tablet, while SR dissolve relatively slowly. The presented data shows plasma concentration levels following administration of a single oral dose (240-mg tablet) in subjects.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

1. Definitions:

[0011] As used herein, an "effective amount" of a compound or pharmaceutically acceptable composition can achieve a desired therapeutic and/or prophylactic effect. In some embodiments, an "effective amount" is at least a minimal amount of a compound, or composition containing a compound, which is sufficient for treating one or more symptoms of a disorder or condition associated with modulation of protein tyrosine kinases. In certain embodiments, an "effective amount" of a compound, or composition containing a compound, is sufficient for treating symptoms associated with, a disease associated with an aberrant tyrosine kinase receptor (e.g. cancer, including malignant and benign tumor growths).

[0012] The term "subject", as used herein, means a mammal and includes human and animal subjects, such as domestic animals (e.g., horses, dogs, cats, etc.).

[0013] The terms "suffer" or "suffering" as used herein refers to one or more conditions that a patient has been diagnosed with, or is suspected to have.

[0014] The terms "treat" or "treating," as used herein, refers to partially or completely alleviating, inhibiting, delaying onset of, preventing, ameliorating and/or relieving a disorder or condition, or one or more symptoms of the disorder or condition.

[0015] "Therapeutically active agent" or "active agent" refers to a substance, including a biologically active substance, that is useful for therapy (e.g., human therapy, veterinary therapy), including prophylactic and therapeutic treatment. Therapeutically active agents include organic molecules that are drug compounds, peptides, proteins, carbohydrates, monosaccharides, oligosaccharides, polysaccharides, nucleoprotein, mucoprotein, lipoprotein, synthetic polypeptide or protein, small molecules linked to a protein, glycoprotein, steroid, nucleic acid, DNA, RNA, nucleotide, nucleoside, oligonucleotides, antisense oligonucleotides,


lipid, hormone, and vitamin. Therapeutically active agents include any substance used as a medicine for treatment, prevention, delay, reduction or amelioration of a disease, condition, or disorder. Among therapeutically active agents useful in formulations are opioid receptor antagonist compounds, opioid analgesic compounds, and the like. Further detailed description of compounds useful as therapeutically active agents is provided below. A therapeutically active agent includes a compound that increases the effect or effectiveness of a second compound, for example, by enhancing potency or reducing adverse effects of a second compound.

[0016] "Unit dosage form" as used herein refers to a physically discrete unit of inventive formulation appropriate for the subject to be treated. It will be understood, however, that the total daily usage of the compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular subject or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of specific active agent employed; specific composition employed; age, body weight, general health, sex and diet of the subject; time of administration, and rate of excretion of the specific active agent employed; duration of the treatment; drugs and/or additional therapies used in combination or coincidental with specific compound(s) employed, and like factors well known in the medical arts.

[0017] In dry granulation (slugging or roller compaction) intragranular materials are blended to prepare slugs or roller compaction. Material is milled and blended with extragranular materials followed by capsule filling or tablet 20 compression. Wet granulation entails blending intragranular materials. Wet granulate the blend with water, with or without a binder, (using high shear, low shear granulators) and dry (using temperatures up to 100° C). Material is milled and blended with extragranular materials followed by capsule filling or tablet compression. See, 25 Handbook of Pharmaceutical Granulation Technology, 1997, Dilip Parikh, Marcel Dekker, Inc. ISBN 0-8247-9882-1, pages 338-368.

2. Pharmaceutically Acceptable Compositions and Formulations:

[0018] In certain embodiments, the present invention provides a pharmaceutically acceptable composition for intravenous administration comprising: neratinib maleate. Neratinib and other 4-amino-3-cyanoquinoline compounds are disclosed in U.S. Pat. Nos. 6,002,008, 6,288,082, 6,297,258, 6,384,051 and 7,399,865. Neratinib has the following chemical structure:



and is isolated as a free base or prepared as a pharmaceutically acceptable salt, such as a

maleate salt. Neratinib is a weak base with an intrinsic low solubility in water.

[0019] In certain embodiments, solid pharmaceutically acceptable compositions of neratinib maleate are provided in the form of tablets prepared by fluid bed granulation. Intragranular particulate components comprising an active ingredient, namely neratinib maleate, one or more fillers, a disintegrant and a glidant are sprayed with, or otherwise wholly or partially covered with, a surface modifying agent, such as povidone, to lower the surface energy of the particles. The fluid bed process is employed to effectively modify the surface behavior of the particulate active ingredient, so that any water instantly dries and does not bring about any polymorphic or chemical change in active during the process. The surface property of the active ingredient is modified by spraying a polymer, for instance povidone, having a low surface energy (for instance of about 38 mN/m) on the surface of the intragranular particles. After the modification of surface properties, the intragranular particles are no longer cohesive, or are significantly less cohesive, and are easily rendered to all pharmaceutical operations. The surface-modified intragranular particles are then further processed, typically by combining with extragranular components typically comprising a filler, a disintegrant and a lubricant, and further processed into dry-filled capsules or tablets for oral administration. The surface modified intragranular components can also be used directly to make dosage forms without combination with extragranular components, for instance in connection with dry-filled capsules.

[0020] In certain embodiments, solid pharmaceutically acceptable compositions of neratinib maleate are provided in the form of tablets prepared by wet granulation. Increasing the glidant and lubrication levels provided an improved neratinib maleate formulation that flowed without aggregation of granules, as compared to a wet granulated neratinib maleate formulation used in clinical trials. The glidant was increased from 0.5% to 2.0 % to improve the flow of the pre-blended material. The picking and sticking issues observed during compression were eliminated by increasing the lubrication level from 0.5% to 3.0%, in certain embodiments 0.5% to 2.0%. In certain embodiments the amount of lubricant such as magnesium stearate needed is as low as 0.2% or even 0.1%. The increase in amount of glidant and lubricant was compensated by a corresponding reduction in the amount of fillers added to the formulation.

[0021] In some cases, the active ingredient comprises a 4-amino-3-cyanoquinoline compound such as neratinib, specially neratinib maleate, or a pharmaceutically acceptable salt thereof. Suitable examples of 4-amino-3-cyanoquinoline compounds are disclosed in U.S. Pat. Nos. 6,002,008, 6,288,082, 6,297,258, 6,384,051 and 7,399,865. According to one embodiment, neratinib maleate is the active ingredient. The active ingredient comprises from about 10 weight % to about 70 weight %, including from 20-50 weight % and about 35 weight % or 41 weight %, based upon total weight of the formulation.

[0022] According to one embodiment, a surface modifying agent is sprayed onto particulate intragranular components before further processing with extragranular components. Suitable surface modifying agents include, but are not limited to for example, povidone, gelatin, starch, hydroxy propyl methyl cellulose and hydroxy propyl cellulose. In one embodiment, povidone is the surface modifying agent. The surface modifying agent comprises from about 1 weight % to

about 15 weight %, including from 3-12 weight % and from 5-10 weight %, based upon total weight of the formulation.

[0023] Suitable fillers (also referred to as "diluent") are known in the art. For example, suitable fillers include but are not limited to starch, dextrin, sucrose, Sorbitol, Sodium Saccharin, Acesulfame potassium, Xylitol, Aspartame, Mannitol, starch, PVP (polyvinyl pyrrolidone), low molecular weight HPC (hydroxypropyl cellulose), microcrystalline cellulose (MCC), low molecular weight HPMC (hydroxypropyl methylcellulose), low molecular weight carboxymethyl cellulose, ethylcellulose, dicalcium phosphate, silicified microcrystalline cellulose, alginates, gelatin, polyethylene oxide, acacia, dextrin, sucrose, magnesium aluminum silicate, and polymethacrylates. Fillers include agents selected from the group consisting of microcrystalline cellulose, starch, lactitol, lactose, a suitable inorganic calcium salt, sucrose, glucose, mannitol, silicic acid, or a combination thereof. The fillers, as an intragranular component, comprise from about 15 weight % to about 65 weight %, based upon total weight of the formulation. In one embodiment, the intragranular filler is a combination of mannitol and microcrystalline cellulose. The fillers, as an extragranular component, comprise from about 4 weight % to about 25 weight %, based upon total weight of the formulation. In one embodiment, the extragranular filler is microcrystalline cellulose.

[0024] Suitable disintegrants are known in the art and include but are not limited to, agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, sodium carbonate, crospovidone (cross-linked PVP), sodium carboxymethyl starch (sodium starch glycolate), cross-linked sodium carboxymethyl cellulose (croscarmellose), pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, sodium starch glycolate, potassium polacrilin, sodium alginate, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum) or a combination thereof. In some embodiments, a disintegrant is crospovidone. The disintegrant, as an intragranular component, comprises from about 0 or 0.5 weight % to about 8 weight %, including from 0 or 0.5-6 weight % and from 0 or 0.5-50-5 weight %, based upon total weight of the formulation. The disintegrant, as an extragranular component, comprises from about 0 or 1 weight % to about 8 weight %, based upon total weight of the formulation.

[0025] A glidant is used as an intragranular component of the formulation. Suitable glidants include, without limitation, colloidal silicon dioxide, talc, magnesium carbonate, calcium silicate, fumed silicon dioxide, and combinations thereof. In some embodiments, the glidant is colloidal silicon dioxide. The amount of glidants used is 0.2-8 weight percent, or 0.2-5 weight percent, including 0.5-2 weight %, based on the total weight of the formulation.

[0026] A lubricant is used as an extragranular component of the formulation. Suitable lubricants or glidants include for example stearates, sodium stearyl fumarate, magnesium salts and magnesium stearate. In some embodiments, the lubricant is magnesium stearate. The amount of lubricants used is 0.2-4 weight percent, in certain embodiments 0.5-3 weight %, based on the total weight of the formulation.

[0027] Provided compositions may be formulated into a unit dosage form. Such formulations

are well known to one of ordinary skill in the art. In certain embodiments, the present invention provides a formulation comprising a solid dosage form as a tablet. In other embodiments, the present invention provides a solution for oral administration. In some embodiments, a unit dosage form contains 5, 10, 20, 25, 40, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 225 mg, 250 mg, 275 mg, 300 mg, 325 mg, 350 mg, 375 mg, 400 mg, 425 mg, 450 mg, 475 mg, or 500 mg, 525 mg, 550 mg, 575 mg, 600 mg, 625 mg, 650 mg, 675 mg, 700 mg, 725 mg, 750 mg, 775 mg, 800 mg, 825 mg, 850 mg, 875 mg, 900 mg, 925 mg, 950 mg, 975 mg, 1000 mg, 1025 mg, 1050 mg, 1075 mg, 1100 mg, 1125 mg, 1150 mg, 1175 mg, 1200 mg, 1225 mg, 1250 mg, 1275 mg, 1300 mg, 1325 mg, 1350 mg, 1375 mg, 1400 mg, 1425 mg, 1450 mg, 1475 mg, 1500 mg of neratinib. In some embodiments, a unit dosage form contains between 5 mg and 500 mg, inclusive, or between 10 mg and 450 mg, inclusive, of neratinib. In some embodiments, a unit dosage form contains 40 mg, 80 mg, 100 mg, 120 mg, 240 mg, 360 mg, or 480 mg. In some embodiments, a unit dosage form contains more than 500 mg of neratinib.

[0028] In some embodiments, satisfactory results are obtained when the compounds of the invention are administered at a daily dosage of from about 0.5 to about 1000 mg/kg of body weight, optionally given in divided doses two to four times a day, or in sustained release form. The total daily dosage is projected to be from about 1 to 1000 mg, preferably from about 2 to 500 mg. Dosage forms suitable for internal use comprise from about 0.5 to 1000 mg of the active compound in intimate admixture with a solid or liquid pharmaceutically acceptable carrier. This dosage regimen may be adjusted to provide the optimal therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

[0029] For the treatment of cancer, the inventive formulations of this invention can be administered in combination with other anti-tumor substances or with radiation therapy. These other substances or radiation treatments can be given at the same or at different times as the compounds of this invention. These combined therapies may effect synergy and result in improved efficacy. For example, the compounds of this invention can be used in combination with mitotic inhibitors such as taxol or vinblastine, alkylating agents such as cisplatin or cyclophosphamide, anti-metabolites such as 5-fluorouracil or hydroxyurea, DNA intercalators such as adriamycin or bleomycin, topoisomerase inhibitors such as etoposide or camptothecin, antiangiogenic agents such as angiostatin, and antiestrogens such as tamoxifen.

[0030] Based on the results disclosed for neratinib and other 4-amino-3-cyanoquinoline compounds in U.S. Pat. No. 6,297,258, the invented formulations are useful antineoplastic agents of significant efficacy, which are useful in treating, inhibiting the growth of, or eradicating neoplasms. In particular, the compounds of this invention are useful in treating, inhibiting the growth of, or eradicating neoplasms that express the receptor protein produced by the erbB2 (Her2) oncogene.

3. Combined Administration:

[0031] In certain embodiments, inventive compositions, and formulations thereof, may be administered alone to treat one or more disorders as described herein, or alternatively may be administered in combination with (whether simultaneously or sequentially) one or more other active agents useful to treat one or more disorders as described herein. Thus, an inventive composition, or formulation thereof, can be administered concurrently with, prior to, or subsequent to, one or more active agents.

[0032] In certain described aspects, compositions include one or more other active agents in addition to neratinib that is not neratinib. In some described aspects, formulations comprise both another anticancer compound and neratinib.

[0033] The amount of additional active agent(s) present in combination compositions will typically be no more than the amount that would normally be administered in a composition comprising that active agent as the only therapeutic agent. In certain described aspects, the amount of additional active agent will range from about 50% to 100% of the amount normally present in a composition comprising that compound as the only therapeutic agent.

4. Uses and Kits of Inventive Compositions:

[0034] Provided compositions, and formulations thereof, are also useful in treatment of conditions including cancers.

[0035] In still further embodiments, veterinary applications (e.g., treatment of domestic animals, e.g. horse, dogs, cats, etc.) of use of inventive compositions, and formulations thereof, are provided. Thus, use of provided formulations in veterinary applications analogous to those discussed above for human subjects is contemplated.

[0036] It will also be appreciated that inventive compositions, and formulations thereof, can be employed in combination therapies, that is, an inventive composition, or formulation thereof, can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. Particular combination therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that therapies employed may achieve a desired effect for the same disorder (for example, a formulation may be administered concurrently with another compound used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic compounds which are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[0037] In other embodiments, inventive compositions, and formulations thereof, and unit dose forms are useful in preparation of medicaments, including, but not limited to medicaments

useful in the treatment of cancer.

[0038] Still further encompassed by the invention are pharmaceutical packs and/or kits comprising inventive compositions, and formulations thereof, and a container (e.g., a foil or plastic package, or other suitable container). Optionally instructions for use are additionally provided in such kits.

[0039] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

[0040] All features of each of the aspects of the invention apply to all other aspects *mutatis mutandis*.

EXAMPLES

Example 1. Preparation of Coated Tablets of a Neratinib Maleate Formulation by Fluid Bed Wet Granulation Process

[0041] A pharmaceutically acceptable formulation of neratinib maleate is prepared: a granulation comprising intragranular components (a) 10-70 weight percent of neratinib maleate; (b) 15-65 weight percent of mannitol and microcrystalline cellulose; (c) 0.5-8 weight percent of crospovidone or croscarmellose sodium; (d) 0.2-8 weight percent of colloidal silicon dioxide, and (e) 5-15 weight percent of povidone. The granulation is combined with extragranular components (f) 4-25 weight percent of microcrystalline cellulose; (g) 1-8 weight percent of crospovidone and (h) 0.5-3 weight percent of magnesium stearate, and then compressed into tablets or dry-filled into capsules. This and certain preferred ranges of materials are shown below in Table 1.

[0042] The formulation was prepared according to the following procedure:

1. Mixed neratinib maleate, mannitol, microcrystalline cellulose and crospovidone and silicon dioxide. Any diffusive or convective mixer may be used.

Table 1

Component	Weight %	Weight % range	Weight % range
Intragranular Components			
HKI-272 Maleate, anhydrous	41	20-50	10-70
Mannitol			15-65
Microcrystalline Cellulose			5-35
Crospovidone/croscarmellose sodium	0.5-5.00	2-6	0.5-8

Component	Weight %	Weight % range	Weight % range
Intragranular Components			
Povidone	5-10	3-12	1-15
Colloidal Silicon Dioxide	0.5-2.00	0.2-3.0	0.2-5
Extragranular Components			
Microcrystalline Cellulose	4-25	4-25	4-25
Crospovidone	1-8	1-8	1-8
Magnesium Stearate	0.5-3.0	0.2-3.5	0.2-4.0
Total (Core Tablet)	1.1.1	1.1.2	1.1.3
Film Coat:			
Opadry®			

- Dissolved povidone in purified water.
- Fluidized the powder blend in step 1 and sprayed it with solution prepared in step 2 in a suitable fluid bed granulator and dryer.
- Dried the granulation.
- Milled the granulation.
- Added microcrystalline cellulose and crospovidone to the granulation in step 5 and mixed.
- Added magnesium stearate to the mixture in step 6 and mixed.
- Compressed the blend in step 7 into tablets of desired strength.
- Applied film-coat to compressed tablets using Opadry II of desired color.
- Alternatively the powder blend can be filled in capsules shells.

Example 2. Unit dosage forms of an Exemplary Neratinib Maleate Formulation

[0043] Using the fluid bed process described in Example 1, different unit dosages of neratinib maleate were prepared from an exemplary formulation, as summarized in Table 2.

Table 2

Ingredient	Function	Granulation wt/wt (%)	40 mg mg/tablet	80 mg mg/tablet	240 mg mg/tablet
Intragranular Components					
HKI-272 Maleate ^a	Active	35.00	40.00	80.00	240.00
Mannitol (Pearlitol 200 SD)	Diluent	38.94	44.50	89.01	267.02
Macrocrystalline (Avicel PH 101)	Diluent	10.56	12.07	24.14	72.41
Crospovidone	Disintegrant	3.00	3.43	6.86	20.57
Povidone USP/K-	Surface	5.00	5.71	11.43	34.29

Ingredient	Function	Granulation wt/wt (%)	40 mg mg/tablet	80 mg mg/tablet	240 mg mg/tablet
Intragranular Components					
25	Modifying Agent				
Colloidal SiO ₂	Glidant	2.00	2.29	4.57	13.71
Extragranular Components					
Avicel PH 101	Diluent	1.50	1.71	3.43	10.29
Crospovidone	Disintegrant	2.00	2.29	4.57	13.71
Mg Stearate	Lubricant	2.00	2.29	4.57	13.71
Total Weight		100.00	114.29	228.57	685.71
Film Coating:					
Opadry II (85F15443) Red	Film Coat	-	3.429 (3%)	-	-
Opadry II (85F92177) Yellow	Film Coat	-	-	6.86 (3%)	-
Opadry II (85F94211) Pink	Film Coat	-	-	-	20.57 (3%)
Total Tablet Wt		-	117.714	235.43	706.28
^a Weight percent as HKI-272 free base					

Example 3. Coated Tablets of Targeted Release Neratinib Maleate Manufactured by Spraying Povidone on Intragranular Components in a Fluid Bed.

[0044] An exemplary targeted release (TR) neratinib maleate formulation is summarized in Table 3.

Example 4. Coated Tablets of Slow Release Neratinib Maleate Manufactured by Spraying Povidone on Intragranular Components in a Fluid Bed.

[0045] An exemplary slow release (SR) neratinib maleate formulation is summarized in Tables 4A and 4B.

Table 3

Ingredient	% Wt/Wt	40 mg Tablet (mg)	240 mg Tablet (mg)	Function
Intragranular Components				
HKI-272 Maleate, anhydrous	35.00	40.00	240.00	Active
Mannitol	38.25	43.79	262.72	Filler
Microcrystalline Cellulose	12.75	14.50	86.99	Filler
Crospovidone	3.00	3.43	20.57	Disintegrant
Povidone	5.00	5.71	34.29	Surface Modifying Agent
Colloidal Silicon Dioxide	0.50	0.57	3.43	Glidant
Extragranular Components				
Microcrystalline Cellulose	3.00	3.43	20.57	Filler
Crospovidone	2.00	2.29	13.71	Disintegrant
Magnesium Stearate	0.50	0.57	3.43	Lubricant
Total (Core Tablet)	100.00	114.29	685.71	
Film Coat:				
Opadry®	-	3.43	20.57	Film Coat

Table 4A

Ingredient Name	Function	& Wt/Wt	mg / tablet 240mg Tablet
Intragranular Components			
HKI-272 Maleate	Active	35.00	240.00
Mannitol USP	Filler	41.81	286.69
Microcrystalline Cellulose NF	Filler	9.19	63.02
Crospovidone NF	Disintegrant	0	
Colloidal Silicon Dioxide, NF	Glidant	0.50	3.43
Povidone USP/K-25	Binder	10.00	68.57
Purified Water, EP/BP/USP	Solvent	-	
Extragranular Components			
Microcrystalline Cellulose NF	Filler	-	
Crospovidone NF	Disintegrant	0.50	3.43
Magnesium Stearate NF/Ph/EU,	Lubricant	3.00	20.57

Extragranular Components			
(Vegetable grade)			

Table 4B

Ingredient	Function	Granulation wt/wt (%)	40 mg mg/tablet	80 mg mg/tablet	240 mg mg/tablet
Intragranular Components					
HKI-272 Maleate Mannitol (Pearlitol 200 SD)	Active	35.00	40.00	80.00	240.00
Mannitol (Pearlitol 200 SD)	Diluent	38.94	44.50	89.01	267.02
Macrocrystalline (Avicel PH 101)	Diluent	10.56	12.07	24.14	72.41
Croscarmellose Sodium	Disintegrant	3.00	3.43	6.86	20.57
Povidone USP/K-25	Surface Modifying Agent	5.00	5.71	11.43	34.29
Colloidal SiO ₂	Glidant	2.00	2.29	4.57	13.71
Extragranular Components					
Avicel PH 101	Diluent	1.50	1.71	3.43	10.29
Croscarmellose Sodium	Disintegrant	2.00	2.29	4.57	13.71
Mg Stearate	Lubricant	2.00	2.29	4.57	13.71
Total Weight		100.00	114.29	228.57	685.71
Film Coating:					
Opadry II (85F15443) Red	Film Coat	-	3.429 (3%)	-	-
Opadry II (85F92177) Yellow	Film Coat	-	-	6.86 (3%)	-
Opadry II (85F94211) Pink	Film Coat	-	-	-	20.57 (3%)
Total Tablet Wt		-	117.714	235.43	706.28

Example 5: Drug Release Data

[0046] Drug release data are summarized for neratinib maleate formulations in Examples 3

and 4, as summarized in Table 5. Dissolution of tablets was carried out employing 900ml of 0.1N HCl as dissolution medium in USP dissolution apparatus 2, and a paddle speed of 50 ± 1 rpm at $37 \pm 0.5^\circ\text{C}$. Samples were taken at specific time points and were analyzed by UV spectrometer at 266 nm.

[0047] Mean pharmacokinetic parameters for neratinib maleate in targeted release and slow release formulations following administration of a single oral dose (240-mg tablet) in subjects were evaluated and summarized in Table 6. The mean concentration versus time profiles for targeted release and slow release formulations are summarized in Figure 1.

Table 6. Summary of Mean Pharmacokinetic Parameters for Neratinib Maleate Formulations Following Single Oral Dose (240-mg tablet) in Healthy Subjects Under Fed Conditions

Mean \pm SD (CV%)	240 mg TR	240 mg SR
[Geometric Mean]	(n=26)	(n=26)
C_{\max} (ng/mL)	81.38 ± 27.49 (34)	82.21 ± 29.73 (36)
	[77.13]	[77.12]
t_{\max} (hr)	5.00	6.00
	(3.00, 12.00)	(3.00, 12.00)
AUC	1432 ± 450 (31)	1474 ± 467 (32)
	[1358]	[1393]

Table 5

Time (minutes)	% Dissolved	
	TR Tablet Formulation	SR Tablet Formulation
15	47	24
30	83	43
45	99	64
60	99	82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61259403B](#) [0001]
- [US6002008A](#) [0018] [0021]
- [US6288082B](#) [0018] [0021]
- [US6297258B](#) [0018] [0021] [0030]
- [US6384051B](#) [0018] [0021]
- [US7399865B](#) [0018] [0021]

Non-patent literature cited in the description

- **TRAXLER, P. M.**Exp. Opin. Ther. Patents, 1998, vol. 8, 1599- [0003]
- **BRIDGES, A. J.**Emerging Drugs, 1998, vol. 3, 279- [0003]
- **MATTSSON, E.**Trends Cardiovas. Med., 1995, vol. 5, 200- [0003]
- **SHAW**Trends Pharmacol. Sci., 1995, vol. 16, 401- [0003]
- **RAINES, E. W.**Bioessays, 1996, vol. 18, 271- [0003]
- **SHAWVER, L. K.**Drug Discovery Today, 1997, vol. 2, 50- [0003]
- **FOLKMAN, J.**Nature Medicine, 1995, vol. 1, 27- [0003]
- **BOYCE**J. Clin. Invest., 1992, vol. 90, 1622- [0003]
- **B. JANCZUKT.BIALOPIOTROWICZ**Surface Free-Energy Components of Liquids and Low Energy Solids and Contact AnglesJ. Colloid Interf. Sci., 1989, vol. 127, 189-204 [0004]
- **W.R. GOODA** Comparison of Contact Angle InterpretationsJ. Colloid Interf. Sci., 1973, vol. 44, 63- [0004]
- **Surface Tension of Pure Liquids and Binary Liquid Mixture**Landolt Börnstein, New SeriesSpringer Verlag19980000vol. IV/16, [0004]
- **J.J. JASPER**The Surface Tension of Pure Liquid CompoundsJ. Phys. Chem. Ref. Data, 1972, vol. 1, 4859- [0004]
- **DILIP PARIKH**Handbook of Pharmaceutical Granulation TechnologyMarcel Dekker, Inc.19970000vol. 25, 338-368 [0017]

Patentkrav

1. Farmaceutisk acceptabel sammensætning omfattende:
en granulering omfattende intragranulære komponenter:

- (a) 10-70 vægtprocent neratinibmaleat;
- 5 (b) 15-65 vægtprocent af et eller flere fyldstoffer;
- (c) 0-8 vægtprocent af et eller flere disintegrationsmidler;
- (d) 0,2-8 vægtprocent af et eller flere glidemidler; og
- (e) 5-15 vægtprocent af et eller flere overflademodificerende midler;

hvor granuleringen er kombineret med extragranulære komponenter:

- 10 (f) 1-25 vægtprocent af et eller flere fyldstoffer;
- (g) 0-8 vægtprocent af et eller flere disintegrationsmidler; og
- (h) 0,1 -3 vægtprocent af et eller flere smøremidler.

2. Farmaceutisk acceptabel sammensætning omfattende 10-70 vægtprocent af
15 neratinibmaleat og 5-15 vægtprocent af povidon.

3. Farmaceutisk acceptabel sammensætning omfattende intragranulære
komponenter: 10-70 vægtprocent af neratinibmaleat og 5-15 vægtprocent af
povidon; og extragranulære komponenter: mikrokrystallinsk cellulose,
20 crospovidon og magnesiumstearat.

4. Farmaceutisk acceptabel sammensætning omfattende intragranulære
komponenter:

- (a) 10-70 vægtprocent af neratinibmaleat;
- 25 (b) 15-65 vægtprocent af mannitol, mikrokrystallinsk cellulose eller en
kombination af begge;
- (c) 0,5-8 vægtprocent crospovidon, croscarmellosenatrium eller en
kombination af begge;
- (d) 0,2-8 vægtprocent kolloid siliciumdioxid; og
- 30 (e) 5-15 vægtprocent povidon;

og extragranulære komponenter:

- (f) 1-25 vægtprocent af mikrokrystallinsk cellulose;
- (g) 1-8 vægtprocent af crospovidon; og
- (h) 0,5-3 vægtprocent af magnesiumstearat.

- 5.** Tablet eller tørfyldt kapsel omfattende sammensætningen ifølge krav 4.
- 6.** Sammensætning ifølge krav 4, hvor mængden af neratinibmaleat er omkring 41 vægtprocent; eller hvor mængden af neratinibmaleat er omkring 35
5 vægtprocent.
- 7.** Sammensætning ifølge krav 4, hvor mængden af mannitol og mikrokrystallinsk cellulose, som en intragranulær komponent, er omkring 50 vægtprocent.
- 10 **8.** Sammensætning ifølge krav 4, hvor mængden af crospovidon, som en intragranulær komponent, er omkring 3 vægtprocent; og/eller hvor mængden af povidon, som en intragranulær komponent, er omkring 5 vægtprocent.
- 9.** Sammensætning ifølge et hvilket som helst af kravene 4 og 6-8, i form af en
15 oral dosis på 40 mg, 80 mg eller 240 mg.
- 10.** Sammensætning ifølge krav 4, hvor sammensætningen omfatter
(b) mannitol og mikrokrystallinsk cellulose og
(c) crospovidon.
20
- 11.** Sammensætning ifølge krav 4, hvor sammensætningen omfatter som intragranulære komponenter:
(a) 35,00 vægtprocent af neratinibmaleat som neratinibfri base;
(b) 38,94 vægtprocent af mannitol;
25 (c) 10,56 vægtprocent af mikrokrystallinsk cellulose;
(d) 3,00 vægtprocent af crospovidon;
(e) 2,00 vægtprocent af kolloid siliciumdioxid; og
(f) 5,00 vægtprocent af povidon;
og som extragranulære komponenter:
30 (g) 1,50 vægtprocent af mikrokrystallinsk cellulose;
(h) 2,00 vægtprocent af crospovidon; og
(i) 2,00 vægtprocent af magnesiumstearat; og hvor sammensætningen yderligere omfatter en filmcoating.

12. Fremgangsmåde til at fremstille en farmaceutisk acceptabel sammensætning omfattende: at sprøjte 5-15 vægtprocent af et eller flere overflademodificerende midler på intragranulære komponenter af en granulering, idet nævnte intragranulære komponenter omfatter:

- 5 (a) 10-70 vægtprocent neratinibmaleat;
 - (b) 15-65 vægtprocent af et eller flere fyldstoffer;
 - (c) 0-8 vægtprocent af et eller flere disintegrationsmidler;
 - (d) 0,2-8 vægtprocent af et eller flere glidemidler; og
 - (e) 5-15 vægtprocent af et eller flere overflademodificerende stoffer;
- 10 hvor nævnte coatede intragranulære komponenter er kombineret med extragranulære komponenter af granuleringen, idet nævnte extragranulære komponenter omfatter:
- (f) 1-25 vægtprocent af et eller flere fyldstoffer;
 - (g) 0-8 vægtprocent af et eller flere disintegrationsmidler; og
- 15 (h) 0,1-3 vægtprocent af et eller flere smøremidler.

13. Fremgangsmåde ifølge krav 12, hvor det overflademodificerende middel er povidon.

20 **14.** Farmaceutisk acceptabel sammensætning ifølge et hvilket som helst af kravene 1 til 4 og 6 til 11 eller tabletten eller den tørfyldte kapsel ifølge krav 5 til anvendelse i en fremgangsmåde til behandling af cancer.

15. Sammensætning ifølge krav 11, hvor mængden af filmcoatingen er omkring 3
25 vægtprocent.

16. Farmaceutisk sammensætning ifølge krav 15, hvor filmcoatingen er rød, gul eller lyserød.

DRAWINGS

FIG. 1

