

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 November 2002 (28.11.2002)

PCT

(10) International Publication Number  
WO 02/094229 A1

- (51) International Patent Classification<sup>7</sup>: A61K 9/72, 31/49, 31/423, 31/27, 31/135
- (21) International Application Number: PCT/US02/15459
- (22) International Filing Date: 15 May 2002 (15.05.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/294,203 24 May 2001 (24.05.2001) US  
60/317,479 5 September 2001 (05.09.2001) US
- (71) Applicant: ALEXZA MOLECULAR DELIVERY CORPORATION [US/US]; 1001 E.Meadow Circle, Palo Alto, CA 94303 (US).
- (72) Inventors: RABINOWITZ, Joshua, D.; 750 N. Shoreline Blvd., #50, Mountain View, CA 94043 (US). ZAFFARONI, Alejandro, C.; 1 Faxon Forest, Atherton, CA 94027 (US).
- (74) Agents: ECKMAN, Richard, R. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/094229 A1

(54) Title: DELIVERY OF MUSCLE RELAXANTS THROUGH AN INHALATION ROUTE

(57) Abstract: The present invention relates to the delivery of muscle relaxants through an inhalation route. Specifically, it relates to aerosols containing a muscle relaxant that is used in inhalation therapy. In a composition aspect of the present invention, the aerosol comprises particles comprising at least 5 percent by weight of muscle relaxant. In a method aspect of the present invention, a muscle relaxant is delivered to a mammal through an inhalation route. The method comprises: a) heating a composition, wherein the composition comprises at least 5 percent by weight of a muscle relaxant, to form a vapor; and, b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal. In a kit aspect of the present invention, a kit for delivering a muscle relaxant through an inhalation route to a mammal is provided which comprises: a) a composition comprising at least 5 percent by weight of a muscle relaxant; and, b) a device that forms a muscle relaxant containing aerosol from the composition, for inhalation by the mammal.

## **DELIVERY OF MUSCLE RELAXANTS THROUGH AN INHALATION ROUTE**

**[0001]** This application claims priority to U.S. provisional application Ser. No. 60/294,203 entitled "Thermal Vapor Delivery of Drugs," filed May 24, 2001, Rabinowitz and Zaffaroni, the entire disclosure of which is hereby incorporated by reference. This application further claims priority to U.S. provisional application Ser. No. 60/317,479 entitled "Aerosol Drug Delivery," filed September 5, 2001, Rabinowitz and Zaffaroni, the entire disclosure of which is hereby incorporated by reference.

### **Field of the Invention**

**[0002]** The present invention relates to the delivery of muscle relaxants through an inhalation route. Specifically, it relates to aerosols containing muscle relaxants that are used in inhalation therapy.

### **Background of the Invention**

**[0003]** There are a number of compositions currently marketed as muscle relaxants. The compositions contain at least one active ingredient that provides for observed therapeutic effects. Among the active ingredients given in muscle relaxant compositions are quinine, chlorzoxazone, carisprodol and cyclobenzaprine.

**[0004]** It is desirable to provide a new route of administration for muscle relaxants that rapidly produces peak plasma concentrations of the compound. The provision of such a route is an object of the present invention.

### **Summary of the Invention**

**[0005]** The present invention relates to the delivery of muscle relaxants through an inhalation route. Specifically, it relates to aerosols containing muscle relaxants that are used in inhalation therapy.

**[0006]** In a composition aspect of the present invention, the aerosol comprises particles comprising at least 5 percent by weight of a muscle relaxant. Preferably, the particles comprise at least 10 percent by weight of a muscle relaxant. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent or 99.97 percent by weight of a muscle relaxant.

[0007] Typically, the aerosol has a mass of at least 10  $\mu\text{g}$ . Preferably, the aerosol has a mass of at least 100  $\mu\text{g}$ . More preferably, the aerosol has a mass of at least 200  $\mu\text{g}$ .

[0008] Typically, the particles comprise less than 10 percent by weight of muscle relaxant degradation products. Preferably, the particles comprise less than 5 percent by weight of muscle relaxant degradation products. More preferably, the particles comprise less than 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of muscle relaxant degradation products.

[0009] Typically, the particles comprise less than 90 percent by weight of water. Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

[0010] Typically, at least 50 percent by weight of the aerosol is amorphous in form, wherein crystalline forms make up less than 50 percent by weight of the total aerosol weight, regardless of the nature of individual particles. Preferably, at least 75 percent by weight of the aerosol is amorphous in form. More preferably, at least 90 percent by weight of the aerosol is amorphous in form.

[0011] Typically, the aerosol has an inhalable aerosol particle density greater than  $10^6$  particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than  $10^7$  particles/mL or  $10^8$  particles/mL.

[0012] Typically, the aerosol particles have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

[0013] Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 3.0. Preferably, the geometric standard deviation is less than 2.5. More preferably, the geometric standard deviation is less than 2.1.

[0014] Typically, the aerosol is formed by heating a composition containing a muscle relaxant to form a vapor and subsequently allowing the vapor to condense into an aerosol.

[0015] In another composition aspect of the present invention, the aerosol comprises particles comprising at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine. Preferably, the particles comprise at least 10 percent by weight of

quinine, chlorzoxazone, carisprodol or cyclobenzaprine. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent or 99.97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

**[0016]** Typically, the aerosol has a mass of at least 10  $\mu\text{g}$ . Preferably, the aerosol has a mass of at least 100  $\mu\text{g}$ . More preferably, the aerosol has a mass of at least 200  $\mu\text{g}$ .

**[0017]** Typically, the particles comprise less than 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products. Preferably, the particles comprise less than 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products. More preferably, the particles comprise less than 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products.

**[0018]** Typically, the particles comprise less than 90 percent by weight of water. Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

**[0019]** Typically, at least 50 percent by weight of the aerosol is amorphous in form, wherein crystalline forms make up less than 50 percent by weight of the total aerosol weight, regardless of the nature of individual particles. Preferably, at least 75 percent by weight of the aerosol is amorphous in form. More preferably, at least 90 percent by weight of the aerosol is amorphous in form.

**[0020]** Typically, where the aerosol comprises quinine, the aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 500 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 400 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 300 mg/L.

**[0021]** Typically, where the aerosol comprises chlorzoxazone, the aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 400 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 300 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 200 mg/L.

[0022] Typically, where the aerosol comprises carisprodol, the aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 500 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 400 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 200 mg/L.

[0023] Typically, where the aerosol comprises cyclobenzaprine, the aerosol has an inhalable aerosol drug mass density of between 1 mg/L and 20 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 2 mg/L and 15 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 4 mg/L and 10 mg/L.

[0024] Typically, the aerosol has an inhalable aerosol particle density greater than  $10^6$  particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than  $10^7$  particles/mL or  $10^8$  particles/mL.

[0025] Typically, the aerosol particles have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

[0026] Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 3.0. Preferably, the geometric standard deviation is less than 2.5. More preferably, the geometric standard deviation is less than 2.1.

[0027] Typically, the aerosol is formed by heating a composition containing quinine, chlorzoxazone, carisprodol or cyclobenzaprine to form a vapor and subsequently allowing the vapor to condense into an aerosol.

[0028] In a method aspect of the present invention, a muscle relaxant is delivered to a mammal through an inhalation route. The method comprises: a) heating a composition, wherein the composition comprises at least 5 percent by weight of a muscle relaxant, to form a vapor; and, b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal. Preferably, the composition that is heated comprises at least 10 percent by weight of a muscle relaxant. More preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent,

70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of a muscle relaxant.

**[0029]** Typically, the particles comprise at least 5 percent by weight of a muscle relaxant. Preferably, the particles comprise at least 10 percent by weight of a muscle relaxant. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of a muscle relaxant.

**[0030]** Typically, the aerosol has a mass of at least 10  $\mu\text{g}$ . Preferably, the aerosol has a mass of at least 100  $\mu\text{g}$ . More preferably, the aerosol has a mass of at least 200  $\mu\text{g}$ .

**[0031]** Typically, the particles comprise less than 10 percent by weight of muscle relaxant degradation products. Preferably, the particles comprise less than 5 percent by weight of muscle relaxant degradation products. More preferably, the particles comprise 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of muscle relaxant degradation products.

**[0032]** Typically, the particles comprise less than 90 percent by weight of water. Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

**[0033]** Typically, the particles of the delivered condensation aerosol have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

**[0034]** Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 3.0. Preferably, the geometric standard deviation is less than 2.5. More preferably, the geometric standard deviation is less than 2.1.

**[0035]** Typically, the delivered aerosol has an inhalable aerosol particle density greater than  $10^6$  particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than  $10^7$  particles/mL or  $10^8$  particles/mL.

**[0036]** Typically, the rate of inhalable aerosol particle formation of the delivered condensation aerosol is greater than  $10^8$  particles per second. Preferably, the aerosol is

formed at a rate greater than  $10^9$  inhalable particles per second. More preferably, the aerosol is formed at a rate greater than  $10^{10}$  inhalable particles per second.

**[0037]** Typically, the delivered condensation aerosol is formed at a rate greater than 0.5 mg/second. Preferably, the aerosol is formed at a rate greater than 0.75 mg/second. More preferably, the aerosol is formed at a rate greater than 1 mg/second, 1.5 mg/second or 2 mg/second.

**[0038]** Typically, the delivered condensation aerosol results in a peak plasma concentration of a muscle relaxant in the mammal in less than 1 h. Preferably, the peak plasma concentration is reached in less than 0.5 h. More preferably, the peak plasma concentration is reached in less than 0.2, 0.1, 0.05, 0.02, 0.01, or 0.005 h (arterial measurement).

**[0039]** In another method aspect of the present invention, one of quinine, chlorzoxazone, carisprodol or cyclobenzaprine is delivered to a mammal through an inhalation route. The method comprises: a) heating a composition, wherein the composition comprises at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine, to form a vapor; and, b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal. Preferably, the composition that is heated comprises at least 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine. More preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

**[0040]** Typically, the particles comprise at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine. Preferably, the particles comprise at least 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine. More preferably, the particles comprise at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

**[0041]** Typically, the aerosol has a mass of at least 10  $\mu\text{g}$ . Preferably, the aerosol has a mass of at least 100  $\mu\text{g}$ . More preferably, the aerosol has a mass of at least 200  $\mu\text{g}$ .

[0042] Typically, the particles comprise less than 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products. Preferably, the particles comprise less than 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products. More preferably, the particles comprise 2.5, 1, 0.5, 0.1 or 0.03 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products.

[0043] Typically, the particles comprise less than 90 percent by weight of water. Preferably, the particles comprise less than 80 percent by weight of water. More preferably, the particles comprise less than 70 percent, 60 percent, 50 percent, 40 percent, 30 percent, 20 percent, 10 percent, or 5 percent by weight of water.

[0044] Typically, the particles of the delivered condensation aerosol have a mass median aerodynamic diameter of less than 5 microns. Preferably, the particles have a mass median aerodynamic diameter of less than 3 microns. More preferably, the particles have a mass median aerodynamic diameter of less than 2 or 1 micron(s).

[0045] Typically, the geometric standard deviation around the mass median aerodynamic diameter of the aerosol particles is less than 3.0. Preferably, the geometric standard deviation is less than 2.5. More preferably, the geometric standard deviation is less than 2.1.

[0046] Typically, where the aerosol comprises quinine, the delivered aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 500 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 400 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 300 mg/L.

[0047] Typically, where the aerosol comprises chlorzoxazone, the delivered aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 400 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 300 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 200 mg/L.

[0048] Typically, where the aerosol comprises carisprodol, the delivered aerosol has an inhalable aerosol drug mass density of between 10 mg/L and 500 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 20 mg/L and 400 mg/L.



More preferably, the aerosol has an inhalable aerosol drug mass density of between 50 mg/L and 300 mg/L.

**[0049]** Typically, where the aerosol comprises cyclobenzaprine, the delivered aerosol has an inhalable aerosol drug mass density of between 1 mg/L and 20 mg/L. Preferably, the aerosol has an inhalable aerosol drug mass density of between 2 mg/L and 15 mg/L. More preferably, the aerosol has an inhalable aerosol drug mass density of between 4 mg/L and 10 mg/L.

**[0050]** Typically, the delivered aerosol has an inhalable aerosol particle density greater than  $10^6$  particles/mL. Preferably, the aerosol has an inhalable aerosol particle density greater than  $10^7$  particles/mL or  $10^8$  particles/mL.

**[0051]** Typically, the rate of inhalable aerosol particle formation of the delivered condensation aerosol is greater than  $10^8$  particles per second. Preferably, the aerosol is formed at a rate greater than  $10^9$  inhalable particles per second. More preferably, the aerosol is formed at a rate greater than  $10^{10}$  inhalable particles per second.

**[0052]** Typically, the delivered condensation aerosol is formed at a rate greater than 0.5 mg/second. Preferably, the aerosol is formed at a rate greater than 0.75 mg/second. More preferably, the aerosol is formed at a rate greater than 1 mg/second, 1.5 mg/second or 2 mg/second.

**[0053]** Typically, where the condensation aerosol comprises quinine, between 50 mg and 500 mg of quinine are delivered to the mammal in a single inspiration. Preferably, between 100 mg and 450 mg of quinine are delivered to the mammal in a single inspiration. More preferably, between 100 mg and 400 mg of quinine are delivered in a single inspiration.

**[0054]** Typically, where the condensation aerosol comprises chlorzoxazone, between 50 mg and 400 mg of chlorzoxazone are delivered to the mammal in a single inspiration. Preferably, between 100 mg and 350 mg of chlorzoxazone are delivered to the mammal in a single inspiration. More preferably, between 100 mg and 300 mg of chlorzoxazone are delivered in a single inspiration.

**[0055]** Typically, where the condensation aerosol comprises carisprodol, between 70 mg and 500 mg of carisprodol are delivered to the mammal in a single inspiration. Preferably, between 150 mg and 450 mg of carisprodol are delivered to the mammal in a

single inspiration. More preferably, between 150 mg and 400 mg of carisprodol are delivered in a single inspiration.

**[0056]** Typically, where the condensation aerosol comprises cyclobenzaprine, between 2 mg and 25 mg of cyclobenzaprine are delivered to the mammal in a single inspiration. Preferably, between 5 mg and 20 mg of cyclobenzaprine are delivered to the mammal in a single inspiration. More preferably, between 5 mg and 15 mg of cyclobenzaprine are delivered to the mammal in a single inspiration.

**[0057]** Typically, the delivered condensation aerosol results in a peak plasma concentration of quinine, chlorzoxazone, carisprodol or cyclobenzaprine in the mammal in less than 1 h. Preferably, the peak plasma concentration is reached in less than 0.5 h. More preferably, the peak plasma concentration is reached in less than 0.2, 0.1, 0.05, 0.02, 0.01 h, or 0.005 h (arterial measurement).

**[0058]** Typically, the delivered condensation aerosol is used to treat musculoskeletal pain.

**[0059]** In a kit aspect of the present invention, a kit for delivering a muscle relaxant through an inhalation route to a mammal is provided which comprises: a) a composition comprising at least 5 percent by weight of a muscle relaxant; and, b) a device that forms a muscle relaxant aerosol from the composition, for inhalation by the mammal. Preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent, 97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of a muscle relaxant.

**[0060]** Typically, the device contained in the kit comprises: a) an element for heating the muscle relaxant composition to form a vapor; b) an element allowing the vapor to cool to form an aerosol; and, c) an element permitting the mammal to inhale the aerosol.

**[0061]** In another kit aspect of the present invention, a kit for delivering quinine, chlorzoxazone, carisprodol or cyclobenzaprine through an inhalation route to a mammal is provided which comprises: a) a composition comprising at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine; and, b) a device that forms an quinine, chlorzoxazone, carisprodol or cyclobenzaprine aerosol from the composition, for inhalation by the mammal. Preferably, the composition comprises at least 20 percent, 30 percent, 40 percent, 50 percent, 60 percent, 70 percent, 80 percent, 90 percent, 95 percent,

97 percent, 99 percent, 99.5 percent, 99.9 percent or 99.97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

[0062] Typically, the device contained in the kit comprises: a) an element for heating the quinine, chlorzoxazone, carisprodol or cyclobenzaprine composition to form a vapor; b) an element allowing the vapor to cool to form an aerosol; and, c) an element permitting the mammal to inhale the aerosol.

### **Brief Description of the Figure**

[0063] Fig. 1 shows a cross-sectional view of a device used to deliver muscle relaxant aerosols to a mammal through an inhalation route.

### **Detailed Description of the Invention**

#### ***Definitions***

[0064] “Aerodynamic diameter” of a given particle refers to the diameter of a spherical droplet with a density of 1 g/mL (the density of water) that has the same settling velocity as the given particle.

[0065] “Aerosol” refers to a suspension of solid or liquid particles in a gas.

[0066] “Aerosol drug mass density” refers to the mass of muscle relaxant per unit volume of aerosol.

[0067] “Aerosol mass density” refers to the mass of particulate matter per unit volume of aerosol.

[0068] “Aerosol particle density” refers to the number of particles per unit volume of aerosol.

[0069] “Amorphous particle” refers to a particle that does not contain more than 50 percent by weight of a crystalline form. Preferably, the particle does not contain more than 25 percent by weight of a crystalline form. More preferably, the particle does not contain more than 10 percent by weight of a crystalline form.

[0070] “Carisprodol” refers to N-isopropyl-2-methyl-2-propyl-1,3-propanediol-dicarbamate.

[0071] “Carisprodol degradation product” refers to a compound resulting from a chemical modification of carisprodol. The modification, for example, can be the result of a

thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

[0072] “Chlorzoxazone” refers to 5-chloro-2-hydroxy-benzoxazole.

[0073] “Chlorzoxazone degradation product” refers to a compound resulting from a chemical modification of chlorzoxazone. The modification, for example, can be the result of a thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

[0074] “Cyclobenzaprine” refers to 3-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-N,N-dimethyl-1-propanamine.

[0075] “Cyclobenzaprine degradation product” refers to a compound resulting from a chemical modification of cyclobenzaprine. The modification, for example, can be the result of a thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

[0076] “Condensation aerosol” refers to an aerosol formed by vaporization of a substance followed by condensation of the substance into an aerosol.

[0077] “Inhalable aerosol drug mass density” refers to the aerosol drug mass density produced by an inhalation device and delivered into a typical patient tidal volume.

[0078] “Inhalable aerosol mass density” refers to the aerosol mass density produced by an inhalation device and delivered into a typical patient tidal volume.

[0079] “Inhalable aerosol particle density” refers to the aerosol particle density of particles of size between 100 nm and 5 microns produced by an inhalation device and delivered into a typical patient tidal volume.

[0080] “Mass median aerodynamic diameter” or “MMAD” of an aerosol refers to the aerodynamic diameter for which half the particulate mass of the aerosol is contributed by particles with an aerodynamic diameter larger than the MMAD and half by particles with an aerodynamic diameter smaller than the MMAD.

[0081] “Muscle relaxant degradation product” refers to a compound resulting from a chemical modification of a muscle relaxant. The modification, for example, can be the result of a thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

[0082] “Quinine” refers to 6-methoxy- $\alpha$ -(5-vinyl-2-quinuclidinyl)-4-quinoline-methanol.

[0083] “Quinine degradation product” refers to a compound resulting from a chemical modification of quinine. The modification, for example, can be the result of a thermally or photochemically induced reaction. Such reactions include, without limitation, oxidation and hydrolysis.

[0084] “Rate of aerosol formation” refers to the mass of aerosolized particulate matter produced by an inhalation device per unit time.

[0085] “Rate of inhalable aerosol particle formation” refers to the number of particles of size between 100 nm and 5 microns produced by an inhalation device per unit time.

[0086] “Rate of drug aerosol formation” refers to the mass of muscle relaxant produced by an inhalation device per unit time.

[0087] “Settling velocity” refers to the terminal velocity of an aerosol particle undergoing gravitational settling in air.

[0088] “Typical patient tidal volume” refers to 1 L for an adult patient and 15 mL/kg for a pediatric patient.

[0089] “Vapor” refers to a gas, and “vapor phase” refers to a gas phase. The term “thermal vapor” refers to a vapor phase, aerosol, or mixture of aerosol-vapor phases, formed preferably by heating.

#### **Formation of Muscle Relaxant Containing Aerosols**

[0090] Any suitable method is used to form the aerosols of the present invention. A preferred method, however, involves heating a composition comprising a muscle relaxant to form a vapor, followed by cooling of the vapor such that it condenses to provide a muscle relaxant comprising aerosol (condensation aerosol). The composition is heated in one of four forms: as pure active compound (*e.g.*, pure quinine, chlorzoxazone, carisprodol or cyclobenzaprine); as a mixture of active compound and a pharmaceutically acceptable excipient; as a salt form of the pure active compound; and, as a mixture of active compound salt form and a pharmaceutically acceptable excipient.

[0091] Salt forms of muscle relaxants (*e.g.*, quinine, chlorzoxazone, carisprodol or cyclobenzaprine) are either commercially available or are obtained from the corresponding free base using well known methods in the art. A variety of pharmaceutically acceptable

salts are suitable for aerosolization. Such salts include, without limitation, the following: hydrochloric acid, hydrobromic acid, acetic acid, maleic acid, formic acid, and fumaric acid salts.

[0092] Pharmaceutically acceptable excipients may be volatile or nonvolatile. Volatile excipients, when heated, are concurrently volatilized, aerosolized and inhaled with the muscle relaxant. Classes of such excipients are known in the art and include, without limitation, gaseous, supercritical fluid, liquid and solid solvents. The following is a list of exemplary carriers within the classes: water; terpenes, such as menthol; alcohols, such as ethanol, propylene glycol, glycerol and other similar alcohols; dimethylformamide; dimethylacetamide; wax; supercritical carbon dioxide; dry ice; and mixtures thereof.

[0093] Solid supports on which the composition is heated are of a variety of shapes. Examples of such shapes include, without limitation, cylinders of less than 1.0 mm in diameter, boxes of less than 1.0 mm thickness and virtually any shape permeated by small (*e.g.*, less than 1.0 mm-sized) pores. Preferably, solid supports provide a large surface to volume ratio (*e.g.*, greater than 100 per meter) and a large surface to mass ratio (*e.g.*, greater than 1 cm<sup>2</sup> per gram).

[0094] A solid support of one shape can also be transformed into another shape with different properties. For example, a flat sheet of 0.25 mm thickness has a surface to volume ratio of approximately 8,000 per meter. Rolling the sheet into a hollow cylinder of 1 cm diameter produces a support that retains the high surface to mass ratio of the original sheet but has a lower surface to volume ratio (about 400 per meter).

[0095] A number of different materials are used to construct the solid supports. Classes of such materials include, without limitation, metals, inorganic materials, carbonaceous materials and polymers. The following are examples of the material classes: aluminum, silver, gold, stainless steel; copper and tungsten; silica, glass, silicon and alumina; graphite, porous carbons, carbon yarns and carbon felts; polytetrafluoroethylene and polyethylene glycol. Combinations of materials and coated variants of materials are used as well.

[0096] Where aluminum is used as a solid support, aluminum foil is a suitable material. Examples of silica, alumina and silicon based materials include amorphous silica S-5631 (Sigma, St. Louis, MO), BCR171 (an alumina of defined surface area greater than 2 m<sup>2</sup>/g from Aldrich, St. Louis, MO) and a silicon wafer as used in the semiconductor industry.

Carbon yarns and felts are available from American Kynol, Inc., New York, NY.

Chromatography resins such as octadecyl silane chemically bonded to porous silica are exemplary coated variants of silica.

**[0097]** The heating of the muscle relaxant compositions is performed using any suitable method. Examples of methods by which heat can be generated include the following: passage of current through an electrical resistance element; absorption of electromagnetic radiation, such as microwave or laser light; and, exothermic chemical reactions, such as exothermic solvation, hydration of pyrophoric materials and oxidation of combustible materials.

#### **Delivery of Muscle Relaxant Containing Aerosols**

**[0098]** Muscle relaxant containing aerosols of the present invention are delivered to a mammal using an inhalation device. Where the aerosol is a condensation aerosol, the device has at least three elements: an element for heating a muscle relaxant containing composition to form a vapor; an element allowing the vapor to cool, thereby providing a condensation aerosol; and, an element permitting the mammal to inhale the aerosol.

Various suitable heating methods are described above. The element that allows cooling is, in its simplest form, an inert passageway linking the heating means to the inhalation means. The element permitting inhalation is an aerosol exit portal that forms a connection between the cooling element and the mammal's respiratory system.

**[0099]** One device used to deliver the muscle relaxant containing aerosol is described in reference to Fig. 1. Delivery device 100 has a proximal end 102 and a distal end 104, a heating module 106, a power source 108, and a mouthpiece 110. A muscle relaxant composition is deposited on a surface 112 of heating module 106. Upon activation of a user activated switch 114, power source 108 initiates heating of heating module 106 (*e.g.*, through ignition of combustible fuel or passage of current through a resistive heating element). The muscle relaxant composition volatilizes due to the heating of heating module 106 and condenses to form a condensation aerosol prior to reaching the mouthpiece 110 at the proximal end of the device 102. Air flow traveling from the device distal end 104 to the mouthpiece 110 carries the condensation aerosol to the mouthpiece 110, where it is inhaled by the mammal.

[0100] Devices, if desired, contain a variety of components to facilitate the delivery of muscle relaxant containing aerosols. For instance, the device may include any component known in the art to control the timing of drug aerosolization relative to inhalation (*e.g.*, breath-actuation), to provide feedback to patients on the rate and/or volume of inhalation, to prevent excessive use (*i.e.*, “lock-out” feature), to prevent use by unauthorized individuals, and/or to record dosing histories.

#### **Dosage of Muscle Relaxant Containing Aerosols**

[0101] The dosage amount of muscle relaxants in aerosol form is generally no greater than twice the standard dose of the drug give orally. For instance, quinine, chlorzoxazone, carisprodol and cyclobenzaprine are given at strengths of 260 mg to 325 mg, 250 mg, 350 mg, and 10 mg respectively for the treatment of musculoskeletal pain. As aerosols, 50 mg to 500 mg of quinine, 50 mg to 400 mg of chlorzoxazone, 70 mg to 500 mg of carisprodol, and 2 mg to 25 mg of quinine are generally provided per inspiration for the same indication. A typical dosage of a muscle relaxant aerosol is either administered as a single inhalation or as a series of inhalations taken within an hour or less (dosage equals sum of inhaled amounts). Where the drug is administered as a series of inhalations, a different amount may be delivered in each inhalation.

One can determine the appropriate dose of muscle relaxant containing aerosols to treat a particular condition using methods such as animal experiments and a dose-finding (Phase I/II) clinical trial. One animal experiment involves measuring plasma concentrations of drug in an animal after its exposure to the aerosol. Mammals such as dogs or primates are typically used in such studies, since their respiratory systems are similar to that of a human. Initial dose levels for testing in humans is generally less than or equal to the dose in the mammal model that resulted in plasma drug levels associated with a therapeutic effect in humans. Dose escalation in humans is then performed, until either an optimal therapeutic response is obtained or a dose-limiting toxicity is encountered.

#### **Analysis of Muscle Relaxant Containing Aerosols**

[0102] Purity of a muscle relaxant containing aerosol is determined using a number of methods, examples of which are described in Sekine *et al.*, *Journal of Forensic Science* 32:1271-1280 (1987) and Martin *et al.*, *Journal of Analytic Toxicology* 13:158-162 (1989). One method involves forming the aerosol in a device through which a gas flow (*e.g.*, air



flow) is maintained, generally at a rate between 0.4 and 60 L/min. The gas flow carries the aerosol into one or more traps. After isolation from the trap, the aerosol is subjected to an analytical technique, such as gas or liquid chromatography, that permits a determination of composition purity.

**[0103]** A variety of different traps are used for aerosol collection. The following list contains examples of such traps: filters; glass wool; impingers; solvent traps, such as dry ice-cooled ethanol, methanol, acetone and dichloromethane traps at various pH values; syringes that sample the aerosol; empty, low-pressure (*e.g.*, vacuum) containers into which the aerosol is drawn; and, empty containers that fully surround and enclose the aerosol generating device. Where a solid such as glass wool is used, it is typically extracted with a solvent such as ethanol. The solvent extract is subjected to analysis rather than the solid (*i.e.*, glass wool) itself. Where a syringe or container is used, the container is similarly extracted with a solvent.

**[0104]** The gas or liquid chromatograph discussed above contains a detection system (*i.e.*, detector). Such detection systems are well known in the art and include, for example, flame ionization, photon absorption and mass spectrometry detectors. An advantage of a mass spectrometry detector is that it can be used to determine the structure of muscle relaxant degradation products.

**[0105]** Particle size distribution of a muscle relaxant containing aerosol is determined using any suitable method in the art (*e.g.*, cascade impaction). An Andersen Eight Stage Non-viable Cascade Impactor (Andersen Instruments, Smyrna, GA) linked to a furnace tube by a mock throat (USP throat, Andersen Instruments, Smyrna, GA) is one system used for cascade impaction studies.

**[0106]** Inhalable aerosol mass density is determined, for example, by delivering a drug-containing aerosol into a confined chamber via an inhalation device and measuring the mass collected in the chamber. Typically, the aerosol is drawn into the chamber by having a pressure gradient between the device and the chamber, wherein the chamber is at lower pressure than the device. The volume of the chamber should approximate the tidal volume of an inhaling patient.

**[0107]** Inhalable aerosol drug mass density is determined, for example, by delivering a drug-containing aerosol into a confined chamber via an inhalation device and measuring

the amount of active drug compound collected in the chamber. Typically, the aerosol is drawn into the chamber by having a pressure gradient between the device and the chamber, wherein the chamber is at lower pressure than the device. The volume of the chamber should approximate the tidal volume of an inhaling patient. The amount of active drug compound collected in the chamber is determined by extracting the chamber, conducting chromatographic analysis of the extract and comparing the results of the chromatographic analysis to those of a standard containing known amounts of drug.

**[0108]** Inhalable aerosol particle density is determined, for example, by delivering aerosol phase drug into a confined chamber via an inhalation device and measuring the number of particles of given size collected in the chamber. The number of particles of a given size may be directly measured based on the light-scattering properties of the particles. Alternatively, the number of particles of a given size is determined by measuring the mass of particles within the given size range and calculating the number of particles based on the mass as follows: Total number of particles = Sum (from size range 1 to size range N) of number of particles in each size range. Number of particles in a given size range = Mass in the size range/Mass of a typical particle in the size range. Mass of a typical particle in a given size range =  $\pi \cdot D^3 \cdot \rho / 6$ , where D is a typical particle diameter in the size range (generally, the mean boundary MMADs defining the size range) in microns,  $\rho$  is the particle density (in g/mL) and mass is given in units of picograms ( $g^{-12}$ ).

**[0109]** Rate of inhalable aerosol particle formation is determined, for example, by delivering aerosol phase drug into a confined chamber via an inhalation device. The delivery is for a set period of time (*e.g.*, 3 s), and the number of particles of a given size collected in the chamber is determined as outlined above. The rate of particle formation is equal to the number of 100 nm to 5 micron particles collected divided by the duration of the collection time.

**[0110]** Rate of aerosol formation is determined, for example, by delivering aerosol phase drug into a confined chamber via an inhalation device. The delivery is for a set period of time (*e.g.*, 3 s), and the mass of particulate matter collected is determined by weighing the confined chamber before and after the delivery of the particulate matter. The rate of aerosol formation is equal to the increase in mass in the chamber divided by the duration of the collection time. Alternatively, where a change in mass of the delivery

device or component thereof can only occur through release of the aerosol phase particulate matter, the mass of particulate matter may be equated with the mass lost from the device or component during the delivery of the aerosol. In this case, the rate of aerosol formation is equal to the decrease in mass of the device or component during the delivery event divided by the duration of the delivery event.

[0111] Rate of drug aerosol formation is determined, for example, by delivering a muscle relaxant containing aerosol into a confined chamber via an inhalation device over a set period of time (*e.g.*, 3 s). Where the aerosol is pure muscle relaxant, the amount of drug collected in the chamber is measured as described above. The rate of drug aerosol formation is equal to the amount of muscle relaxant collected in the chamber divided by the duration of the collection time. Where the muscle relaxant containing aerosol comprises a pharmaceutically acceptable excipient, multiplying the rate of aerosol formation by the percentage of muscle relaxant in the aerosol provides the rate of drug aerosol formation.

#### **Utility of Muscle Relaxant Containing Aerosols**

[0112] The muscle relaxant containing aerosols of the present invention are typically used for the treatment musculoskeletal pain or restless leg syndrome.

[0113] The following examples are meant to illustrate, rather than limit, the present invention.

[0114] Quinine sulfate and cyclobenzaprine hydrochloride are commercially available from Sigma ([www.sigma-aldrich.com](http://www.sigma-aldrich.com)). Carisprodol is available in tablet form (SOMA®), from which it is isolated using standard procedures known to one of ordinary skill in the art. Chlorzoxazone is available in caplet form (PARAFON FORTE® DSC), from which it is isolated using standard procedures known to one of ordinary skill in the art.

### **EXAMPLE 1**

#### ***General Procedure for Obtaining Free Base of a Compound Salt***

[0115] Approximately 1 g of salt (*e.g.*, mono hydrochloride) is dissolved in deionized water (~30 mL). Three equivalents of sodium hydroxide (1 N NaOH<sub>aq</sub>) is added dropwise to the solution, and the pH is checked to ensure it is basic. The aqueous solution is extracted four times with dichloromethane (~50 mL), and the extracts are combined, dried (Na<sub>2</sub>SO<sub>4</sub>) and filtered. The filtered organic solution is concentrated using a rotary

evaporator to provide the desired free base. If necessary, purification of the free base is performed using standard methods such as chromatography or recrystallization.

## EXAMPLE 2

### *General Procedure for Volatilizing Compounds from Halogen Bulb*

[0116] A solution of drug in approximately 120  $\mu$ L dichloromethane is coated on a 3.5 cm x 7.5 cm piece of aluminum foil (precleaned with acetone). The dichloromethane is allowed to evaporate. The coated foil is wrapped around a 300 watt halogen tube (Feit Electric Company, Pico Rivera, CA), which is inserted into a glass tube sealed at one end with a rubber stopper. Running 60 V of alternating current (driven by line power controlled by a variac) through the bulb for 6 s (quinine) or 90 V for 5 s (chlorzoxazone, carisprodol and cyclobenzaprine) affords thermal vapor (including aerosol), which is collected on the glass tube walls. Reverse-phase HPLC analysis with detection by absorption of 225 nm light is used to determine the purity of the aerosol. (When desired, the system is flushed through with argon prior to volatilization.)

[0117] Quinine, chlorzoxazone, and cyclobenzaprine aerosol were obtained in 100% (0.97 mg), 99.71% (1.55 mg), and 99.03% purity (6.33 mg) respectively using this procedure.

CLAIMS

1. An aerosol for inhalation therapy, wherein the aerosol comprises particles comprising at least 10 percent by weight of a muscle relaxant.
2. An aerosol for inhalation therapy, wherein the aerosol comprises particles comprising at least 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.
3. The aerosol according to Claim 2, wherein the aerosol particles have a mass median aerodynamic diameter of less than 3 microns.
4. The aerosol according to Claim 2, wherein the particles comprise less than 2.5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products.
5. The aerosol according to Claim 2, wherein the aerosol comprises particles comprising at least 90 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.
6. The aerosol according to Claim 5, wherein the aerosol particles have a mass median aerodynamic diameter less than 2 microns.
7. The aerosol according to Claim 6, wherein the aerosol comprises particles comprising at least 97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.
8. A method of delivering quinine, chlorzoxazone, carisprodol or cyclobenzaprine to a mammal through an inhalation route, wherein the route comprises:

a) heating a composition, wherein the composition comprises at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine, to form a vapor; and,

b) allowing the vapor to cool, thereby forming a condensation aerosol comprising particles, which is inhaled by the mammal.

9. The method according to Claim 8, wherein the particles comprise at least 10 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

10. The method according to Claim 8, wherein the aerosol particles have a mass median aerodynamic diameter of less than 3 microns.

11. The method according to Claim 8, wherein the particles comprise less than 2.5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine degradation products.

12. The method according to Claim 10, wherein the aerosol comprises particles comprising at least 90 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

13. The method according to Claim 12, wherein the aerosol particles have a mass median aerodynamic diameter less than 2 microns.

14. The method according to Claim 13, wherein the aerosol comprises particles comprising at least 97 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine.

15. A kit for delivering quinine, chlorzoxazone, carisprodol or cyclobenzaprine through an inhalation route to a mammal, wherein the kit comprises:

a) a composition comprising at least 5 percent by weight of quinine, chlorzoxazone, carisprodol or cyclobenzaprine; and,

b) a device that forms a quinine, chlorzoxazone, carisprodol or cyclobenzaprine aerosol from the composition, for inhalation by the mammal and wherein the device comprises:

a) an element for heating the quinine, chlorzoxazone, carisprodol or cyclobenzaprine composition to form a vapor;

b) an element allowing the vapor to cool to form an aerosol; and,

c) an element permitting the mammal to inhale the aerosol.




Fig. 1



## INTERNATIONAL SEARCH REPORT

Internat Application No

PCT/US 02/15459

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/72 A61K31/49 A61K31/423 A61K31/27 A61K31/135

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 376 386 A (GANDERTON DAVID ET AL) 27 December 1994 (1994-12-27) column 3, line 44 -column 4, line 68 ---	1-7
X	WO 00 27363 A (NANOSYSTEMS) 18 May 2000 (2000-05-18) page 7, line 23 -page 11, line 18 page 25, line 26 ---	1-7
X	WO 01 32144 A (INHALE THERAPEUTIC SYST ;LECHUGA BALLESTEROS DAVID (US); KUO MEI C) 10 May 2001 (2001-05-10) page 2, line 27 -page 3, line 31 page 9, line 22 page 19, line 24 -page 20, line 25 -----	1-7

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

\* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

25 September 2002

Date of mailing of the international search report

09/10/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Giménez Miralles, J

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 02/15459

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: 8-14  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2.  Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.  No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 02 15459

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 8-14 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

Continuation of Box I.1

Claims Nos.: 8-14

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 02/15459

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5376386	A	27-12-1994	AT 98487 T	15-01-1994
			DE 69100792 T2	14-04-1994
			NO 913731 A	21-11-1991
			AU 635616 B2	25-03-1993
			AU 7155991 A	21-08-1991
			CA 2049302 A1	25-07-1991
			DE 69100792 D1	27-01-1994
			EP 0464171 A1	08-01-1992
			WO 9111179 A1	08-08-1991
			GB 2240337 A ,B	31-07-1991
			HU 59821 A2	28-07-1992
			IE 910222 A1	31-07-1991
			JP 3100626 B2	16-10-2000
			JP 4504427 T	06-08-1992
			PT 96567 A	15-10-1991
			US 5254330 A	19-10-1993
WO 0027363	A	18-05-2000	US 2002102294 A1	01-08-2002
			AU 1346900 A	29-05-2000
			EP 1128814 A1	05-09-2001
			WO 0027363 A1	18-05-2000
WO 0132144	A	10-05-2001	AU 4235300 A	14-05-2001
			EP 1223915 A1	24-07-2002
			NO 20021800 A	24-06-2002
			WO 0132144 A1	10-05-2001