

(19) **DANMARK**

(10) **DK/EP 2920208 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 K 16/22 (2006.01)** **C 07 K 16/28 (2006.01)** **C 07 K 16/42 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2020-09-14**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-08-19**
- (86) Europæisk ansøgning nr.: **13798483.7**
- (86) Europæisk indleveringsdag: **2013-11-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-09-23**
- (86) International ansøgning nr.: **US2013069993**
- (87) Internationalt publikationsnr.: **WO2014078475**
- (30) Prioritet: **2012-11-14 US 201261726040 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591, USA**
- (72) Opfinder: **DESHPANDE, Dipali, 62 Eldorado Court, White Plains, New York 10603, USA**
CHEN, Gang, 1600 Fox Tail Lane, Yorktown Heights, New York 10598, USA
BURAKOV, Darya, 10 Ashton Road, Yonkers, New York 10705, USA
FANDL, James, 40 Amanda's Way, La Grangeville, New York 12540, USA
ALDRICH, Thomas, 2360 Bunney Court, Yorktown Heights, New York 10598, USA
KAMAT, Vishal, 129 Liberty Road, Apt. P, Bergenfield, New Jersey 07621, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Rekombinante celleoverflade-capture-proteiner**
- (56) Fremdragne publikationer:
EP-A1- 2 522 724
EP-A2- 0 107 509
WO-A2-02/057423
US-A1- 2009 137 416
US-A1- 2010 331 527
MENG Y G ET AL: "Green fluorescent protein as a second selectable marker for selection of high producing clones from transfected CHO cells", GENE, ELSEVIER, AMSTERDAM, NL, vol. 242, no. 1-2, 1 January 2000 (2000-01-01), pages 201-207, XP004196520, ISSN: 0378-1119, DOI: 10.1016/S0378-1119(99)00524-7
MANZ R ET AL: "Analysis and sorting of live cells according to secreted molecules, relocated to a cell-surface affinity matrix", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, NATIONAL ACADEMY OF SCIENCES, US, vol. 92, no. 6, 14 March 1995 (1995-03-14), pages 1921-1925, XP002007404, ISSN: 0027-8424, DOI: 10.1073/PNAS.92.6.1921
GESTUR VIDARSSON ET AL: "IgG Subclasses and Allotypes: From Structure to Effector Functions", FRONTIERS IN IMMUNOLOGY, vol. 5, 1 October 2014 (2014-10-01), XP055416330, DOI:

Fortsættes ...

10.3389/fimmu.2014.00520

DESCRIPTION

BACKGROUND

Field of the Invention

[0001] The field of this invention is related to recombinant cell surface capture proteins and methods for identifying, isolating and enriching cells that produce secreted proteins that are heterodimers, e.g. bispecific proteins. More specifically, the cell surface capture proteins and methods allow rapid and efficient isolation of high expression recombinant antibody-producing cell lines, including rapid and efficient isolation of specific hybridomas and cells secreting heterodimeric proteins, e.g. bispecific antibodies, thereby enriching the heterodimeric species (bispecific molecule) and preferentially isolating the heterodimeric from the homodimeric species.

[0002] Prior art methods for expressing a gene of interest (GOI) in a host cell are known. Briefly, an expression vector carrying the GOI is introduced into the cell. Following stable integration, standard methods for isolating high expression cells involve collection of cell pools, hand-picking colonies from plates, isolation of single cells by limited dilution, or other methods known in the art. Pools or individual clones are then expanded and screened for production of the protein of interest (POI) by direct measurement of POI activity, by immunological detection of POI, or by other suitable techniques. These procedures are laborious, inefficient, expensive, and the number of clones that can be analyzed is usually limited to a few hundred.

[0003] The large degree of heterogeneity in protein expression by cells following stable integration requires that many individual clones be screened in an effort to identify the rare integration event that results in a stable, high expression production cell line. This requirement calls for methods that enable rapid identification and isolation of cells expressing the highest level of protein production. Moreover, the collection of clone pools or hand-picked colonies risks losing high expression cells, which often grow more slowly, to faster growing low expression cells. Therefore, a need exists for methods that allow rapid screening and isolation of individual cells capable of high level expression of a secreted POI. Where the POI contains more than one subunit, it is necessary to select preferentially for a desired heterodimeric species versus a homodimeric species.

[0004] Incorporation of flow cytometry into methods used for the isolation of stable expression cell lines has improved the capability of screening large numbers of individual clones, however, currently available methods remain inadequate for diverse reasons. Diffusion of the POI between cells of different characteristics was also a problem.

The following documents are also acknowledged:

- WO 02/057423 which is concerned with isolating cells expressing secreted proteins; and
- US2020/0331527 which is concerned with readily isolated bispecific antibodies with native immunoglobulin format.

BRIEF SUMMARY

[0005] A high-throughput screening method for the rapid isolation of those cells that secrete a heterodimeric protein by directly screening for the protein of interest (POI) is disclosed. This also allows for the convenient monitoring of POI expression on a single-cell basis during the manufacturing process. Furthermore, this technology can be directly applied to screening of bispecific antibody-producing cells, or any cell producing a heterodimeric protein. The technology can also be directly applied to screening of cells producing modified T cell receptors, such as, for example, cells that produce soluble forms of T cell receptors.

[0006] Accordingly, the present invention provides recombinant antigen-binding protein that binds a human IgG1-Fc domain, a human IgG2-Fc domain, or a human IgG4-Fc domain, wherein:

1. (a) the recombinant antigen-binding protein is specific for a wild type CH3 domain containing a histidine residue at position 95 according to the IMGT exon numbering system and a tyrosine residue at position 96 according to the IMGT exon numbering system (Fc) over a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*); and
2. (b) the recombinant antigen-binding protein comprises: (i) a ScFv which comprises an HCVR having the amino acid sequence of SEQ ID NO:15 and an LCVR having the amino acid sequence of SEQ ID NO:16; and (ii) a membrane anchor.

[0007] The present invention further provides an isolated polynucleotide comprising a nucleic acid sequence that encodes: (a) the antigen binding protein of the invention; or (b) a polypeptide comprising the amino acid sequence of SEQ ID NO:19. The present invention also provides a nucleic acid vector comprising: (a) a polynucleotide according to the invention; (b) a promoter, which is operably linked to the polynucleotide; and (c) a polyadenylation sequence. The invention further provides a host cell expressing an antigen-binding protein of the invention.

[0008] The invention further provides a method of detecting or isolating a cell that stably expresses a heterodimeric protein comprising the steps of:

1. (a) expressing in a host cell a cell surface capture protein (CSCP) comprising an

- antigen-binding protein of the invention and a heterodimeric protein, wherein (i) the CSCP binds to a first site on the heterodimeric protein to form a CSCP- heterodimeric protein complex inside the host cell, (ii) the CSCP- heterodimeric protein complex is transported through the host cell, and (iii) then displayed on the surface of the host cell;
2. (b) contacting the host cell with a detection molecule, wherein the detection molecule binds to a second site on the heterodimeric protein; and
 3. (c) selecting the host cell which binds the detection molecule, wherein in the heterodimeric protein:
 1. (i) the first site on the heterodimeric protein resides on a first subunit comprising a heavy chain comprising a wild type CH3 domain containing a histidine residue at position 95 according to the IMGT exon numbering system and a tyrosine residue at position 96 according to the IMGT exon numbering system (Fc); and
 2. (ii) the second site on the heterodimeric protein resides on a second subunit comprising a heavy chain comprising a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*);

wherein the heterodimeric protein comprises a bispecific antibody or an Fc-containing fusion protein.

[0009] Also disclosed is a method of detecting and isolating cells that produce a secreted protein of interest (POI), comprising: a) constructing a nucleic acid molecule that encodes a cell surface capture molecule capable of binding a POI; b) transfecting a cell expressing the POI with the nucleic acid molecule of step a); c) detecting the surface-displayed POI by contacting the cells with a detection molecule, where in the detection molecule binds the POI; and d) isolating cells based on the detection molecule.

[0010] The protein of interest is a heterodimeric protein. In various embodiments, the protein of interest includes a ligand, a soluble receptor protein, a growth factor, a fusion protein, an antibody, a bispecific antibody, an Fab, a single chain antibody (ScFv), or a fragment thereof. When the protein of interest is an antibody, the antibody is selected from the group consisting of IgM, IgG, IgA, IgD or IgE, as well as various subtypes or variants of these. In a specific embodiment, the antibody is an anti-DII4 antibody, an anti-ErbB3 antibody, an anti-EGFR antibody, a dual-specific anti-ErbB3/EGFR bispecific antibody, or an anti-IL-6 receptor antibody.

[0011] [Deleted]

[0012] In various embodiments, the protein of interest comprises a variable domain of a T cell receptor. In specific embodiments, the protein of interest is a protein comprising a T cell receptor extracellular domain fused to an Fc (TCR-Fc). In a specific embodiment, the Fc is a human Fc. In various embodiments, the protein comprises a variable domain of a T cell receptor extracellular domain. In various embodiments, the protein comprises a variable domain and a constant region of a T cell receptor extracellular domain.

[0013] The nucleic acid that encodes the protein of interest may be from any source, naturally occurring or constructed through recombinant technology, and may be selected from a DNA library.

[0014] The cell surface capture protein (CSCP) employed comprises an antigen-binding protein of the invention. The capture molecule is a fusion protein comprising an anti-Ig ScFv fused to a transmembrane domain or a GPI linker.

[0015] [Deleted]

[0016] The protein of interest is a heterodimeric protein having a first subunit and a second subunit, the cell surface capture molecule comprises an ScFv capable of binding the first subunit and not the second subunit, or such cell surface capture molecule binds the second subunit and not the first subunit.

[0017] In various embodiments where the protein of interest is an IgG1, IgG2, IgG4, or a bispecific antibody having one CH3 domain comprising a mutation that abrogates binding to protein A and the other CH3 domain capable of binding to protein A; or a fusion protein comprising an Fc region from IgG1, IgG2, IgG4, or an Fc region having one CH3 domain comprising a mutation that abrogates binding to protein A and the other CH3 domain capable of binding to protein A, the cell surface capture molecule comprises an anti-immunoglobulin ScFv, which is an anti-Fc.

[0018] In several instances, the methods of the invention further comprise a membrane anchor that serves to anchor the POI to the cell membrane, exposed to the outside of the cell, and thus functions as a cell surface capture molecule. In specific embodiments, the membrane anchor is a transmembrane anchor or a GPI link. Examples of specific transmembrane anchors include the transmembrane domain of an Fc receptor, such as the transmembrane domain of human FcγRI, an example of which is cited in SEQ ID NO:17. The membrane anchor may be native to the cell, recombinant, or synthetic.

[0019] Also disclosed are instances where the protein of interest comprises a T cell receptor variable region, and the cell surface capture molecule comprises a membrane-associated antigen. In a specific instances, the membrane-associated antigen is a recombinant fusion protein comprising an antigen capable of being recognized by the T cell receptor variable region fused to a membrane anchor wherein the antigen is associated with the cell surface. In a specific instance the recombinant fusion protein comprises an antigen fused to a transmembrane anchor or a GPI link. In another specific instance, the cell surface capture molecule comprises a recombinant fusion protein comprising a membrane anchor and an antigen that is capable of binding to a major histocompatibility (MHC) molecule, including but not limited to, for example, tumor antigens and self-proteins of transformed phenotype.

[0020] In further embodiments, a signal sequence is added to the amino terminus of a POI, such that the protein is transported to the cell surface, and functions as a cell surface capture

molecule. The signal sequence may be native to the cell, recombinant, or synthetic.

[0021] In various embodiments, a blocking molecule which binds the cell surface capture molecule is added to reduce the diffusion of the POI from the expressing cell to a neighboring cell. In another embodiment, the diffusion of the POI from the expressing cell to a neighboring cell and its adherence to that cell is reduced by increasing the viscosity of the media.

[0022] The cell isolated by the methods of the invention may be an antibody-producing cell fused to an immortalized cell. In more specific embodiments, the antibody-producing cell is a B-cell or derivative thereof. A B-cell derivative may be a plasma cell, a hybridoma, a myeloma, or a recombinant cell.

[0023] In addition, the methods of the invention are useful for identification of B-cells and derivatives thereof, or hybridomas that express secreted antibodies of a desired specificity, affinity or isotype. The invention can also be used for isolation of cells that express desired levels of an antibody or antibody fragments.

[0024] Detection of the cells with the displayed POI may be accomplished through the use of any molecule capable of directly or indirectly binding the displayed POI. Such detection molecules may facilitate the detection and/or isolation of the cells displaying the POI. In one embodiment, two molecules that bind each other and are differentially labeled are utilized. The detection and/or isolation may be accomplished through standard techniques known in the art.

[0025] Also disclosed is a method of detecting and isolating cells that produce a secreted protein of interest (POI), comprising: a) transfecting a cell with a nucleic acid that encodes a cell surface capture molecule, wherein the cell surface capture molecule is capable of binding the POI; b) transfecting the cell of a) simultaneously or subsequently with a second nucleic acid that encodes a POI wherein the POI is expressed and secreted; c) detecting the surface-displayed POI by contacting the cell with a detection molecule, which binds the POI; and d) isolating cells based on the detection molecule.

[0026] Also disclosed is a method of detecting and isolating cells that produce a POI, comprising: a) detecting a cell that expresses a cell surface capture molecule in high yield; b) isolating and culturing the cell detected in (a); c) transfecting the cell in (b) with a nucleic acid that encodes a POI wherein such POI is secreted; d) detecting the surface-displayed POI by contacting the cells with a detection molecule which binds the POI; and e) isolating cells based on the detection molecule.

[0027] Also disclosed is a method of detecting and isolating cells that produce high levels of protein of interest (POI), comprising: a) transfecting cells with a nucleic acid that encodes such cell surface capture molecule capable of binding the POI, wherein the cell expresses the POI; b) detecting a cell from (a) that expresses said cell surface capture molecule in high yield; c) isolating and culturing a high yield cell; d) detecting the surface-displayed POI by contacting the cell with a detection molecule binds the POI; and e) isolating the detected cell.

[0028] Also disclosed is a method of detecting and isolating cells that produce high levels of a heterodimeric protein, comprising: (a) transfecting cells with a nucleic acid that encodes a cell surface capture molecule, which is a fusion protein comprising a membrane anchor domain and is capable of binding a first subunit of the heterodimeric protein, wherein the cell expresses the heterodimeric protein; (b) detecting a cell of (a) that expresses the surface capture molecule in high yield; (c) isolating and culturing the cell that expresses the surface capture molecule in high yield; (d) detecting the heterodimeric protein on the surface of the isolated and cultured cell of step (c) with a detection molecule that binds a second subunit of the heterodimeric protein; and (e) isolating the cell detected in step (d) that bears the detected heterodimeric protein on its surface.

[0029] Also disclosed is a method of detecting and isolating cells that produce high levels of an immunoglobulin, comprising: (a) transfecting cells with a nucleic acid that encodes a cell surface capture molecule capable of binding the immunoglobulin, wherein the cell expresses the immunoglobulin; (b) detecting a cell of (a) that expresses the surface capture molecule in high yield; (c) isolating and culturing the cell that expresses the surface capture molecule in high yield; (d) detecting the immunoglobulin on the surface of the isolated and cultured cell of step (c) with a detection molecule that binds the immunoglobulin; and (e) isolating the cell detected in step (d) that bears the detected immunoglobulin on its surface.

[0030] Also disclosed is a method of detecting and isolating cells that produce high levels of a bispecific antibody, comprising: (a) transfecting cells with a nucleic acid that encodes a cell surface capture molecule, which is a fusion protein comprising a membrane anchor domain, such as an ScFv fusion protein and is capable of binding the bispecific antibody, wherein the cell expresses the bispecific antibody; (b) detecting a cell of (a) that expresses the surface capture molecule in high yield; (c) isolating and culturing the cell that expresses the surface capture molecule in high yield; (d) detecting the bispecific antibody on the surface of the isolated and cultured cell of step (c) with a detection molecule that binds the bispecific antibody; and (e) isolating the cell detected in step (d) that bears the detected bispecific antibody on its surface.

[0031] Also disclosed is a method for detecting cells that produce a desired level of an affinity agent that comprises a T-cell receptor (TCR) variable region.

[0032] Also disclosed is a method for detecting cells that produce a desired level of a TCR-Fc is provided, comprising: (a) transfecting cells with a nucleic acid that encodes an Fc receptor capable of binding a TCR-Fc, wherein the cell expresses an antigen recognized by the TCR-Fc; (b) detecting a cell of (a) that expresses the TCR-Fc in high yield; (c) isolating and culturing the cell that expresses the TCR-Fc in high yield; (d) detecting the antigen on the surface of the isolated and cultured cell of step (c) with a detection molecule; and (e) isolating the cell detected in step (d) that bears the detected antigen on its surface.

[0033] In various embodiments, the TCR is selected from a human TCR and a rodent TCR

such as a rat, mouse, or hamster TCR. In a specific embodiment the Fc is a human Fc. In another specific embodiment, the Fc is a human Fc and the Fc receptor is a high affinity human Fc receptor. In a specific embodiment, the high affinity human Fc receptor is a human FcγRI.

[0034] [Deleted].

[0035] [Deleted]

[0036] Also disclosed is a recombinant antigen-binding protein that binds a human IgG1-Fc domain, a human IgG2-Fc domain, or a human IgG4-Fc domain, or any protein that comprises for example an amino acid sequence of SEQ ID NO:26, which encodes a human Fc. In some instances the recombinant antigen-binding protein binds the polypeptide with a K_D of less than about 40 nM as measured in a surface plasmon resonance assay.

[0037] The recombinant antigen-binding protein comprises (i) a ScFv which comprises an HCVR having the amino acid sequence of SEQ ID NO:15 and an LCVR having the amino acid sequence of SEQ ID NO:16; and (ii) a membrane anchor.

[0038] Recombinant antigen-binding proteins, which are antibodies, are useful as detection molecules (DMs).

[0039] In one embodiment, the membrane anchor domain is derived from an Fc receptor, such as the transmembrane domain of the human FcγR1 protein, as represented by SEQ ID NO:17, or SEQ ID NO:21, which contains not only the transmembrane domain, but also the C-terminal cytoplasmic domain (SEQ ID NO:18). In one specific embodiment, the ScFv fusion protein has the amino acid sequence of SEQ ID NO:19. Recombinant antigen-binding proteins, which are ScFv fusion proteins, are useful as CSCPs and as DMs.

[0040] In another aspect, the invention provides a polynucleotide that encodes the antigen-binding protein of the preceding aspect. In one embodiment, such as in the case where the antigen-binding protein is an antibody, the polynucleotide encodes the light chain. Likewise, the polynucleotide may encode the heavy chain. In the case in which the antigen-binding protein is an ScFv fusion protein, the polynucleotide may encode the ScFv-FcγRTM-cyto fusion protein of SEQ ID NO:19. For example, the polynucleotide of SEQ ID NO: 20 encodes SEQ ID NO:19.

[0041] In another aspect, the invention provides a nucleic acid vector that encompasses the polynucleotide of the preceding aspect. In one embodiment, the vector comprises the polynucleotide, which encodes the antigen-binding protein, operably linked to an upstream promoter, and followed by a downstream polyadenylation sequence. The promoter can be any promoter, such as for example a CMV promoter. Thus in one case, the vector may contain the sequence of SEQ ID NO:25. In one embodiment, the vector may contain a nucleic acid sequence that encodes a selectable marker, such as for example neomycin resistance. In one embodiment, the vector may contain a nucleic acid sequence that encodes an energy transfer

protein, such as green fluorescence protein (GFP), or a derivative thereof, such as yellow fluorescence protein (YFP). Thus in one case, the vector may contain the sequence of SEQ ID NO:24.

[0042] The vector may be circular or linear, episomal to a host cell's genome or integrated into the host cell's genome. In some embodiments, the vector is a circular plasmid, which in one specific embodiment has the nucleic acid sequence of SEQ ID NO:23 for the ScFv-FcγR-TM-cyto-encoding polynucleotide, in another specific embodiment comprises the nucleic acid sequence of the antibody heavy chain-encoding polynucleotide, and yet another specific embodiment comprises the nucleic acid sequence of the antibody light chain-encoding polynucleotide. In some embodiments, the vector is a linear construct, which may be integrated into a host cell chromosome. In one specific embodiment, the linear construct has the nucleic acid sequence of SEQ ID NO:22 for the ScFv-FcγR-TM-cyto-encoding polynucleotide. In another specific embodiment, the linear construct comprises the nucleic acid sequence of the antibody heavy chain-encoding polynucleotide. In yet another specific embodiment, the linear construct comprises the nucleic acid sequence of the antibody light chain-encoding polynucleotide.

[0043] The host cell may be any cell, prokaryotic or eukaryotic. However, in one specific embodiment, the host cell is a CHO cell, such as a CHO-K1 cell.

[0044] In another aspect, the invention provides a host cell that expresses the antigen-binding protein of the preceding aspect, and/or contains the polynucleotide or nucleic acid vector of the preceding aspects. In some embodiments, the host cell is a CHO cell. In a specific embodiment, the host cell is a CHO-K1 cell. In one embodiment, host cell is used in the production of a protein of interest, and the antigen-binding protein is used as a cell surface capture protein according to the methods disclosed in this application.

[0045] In one aspect, the invention provides a host cell useful in the production of a protein of interest. The host cell harbors a polynucleotide or nucleic acid vector of a preceding aspect, and produces an antigen-binding protein of a preceding aspect, which serves as a cell surface capture protein. The cell surface capture protein binds to the protein of interest inside the host cell, and is transported through the secretory apparatus of the cell, and is expressed on the surface of the host cell. Thus, in one embodiment, the host cell comprises a cell surface capture protein positioned in the host cell plasma membrane, with the capturing moiety facing outside of the cell. In one embodiment, the cell surface capture molecule is bound to a protein of interest, which is positioned at the plasma membrane and oriented outside of the cell.

[0046] In one embodiment, the host cell produces or is capable of producing an ScFv fusion protein that binds to a protein of interest that contains an Fc domain, which contains a histidine at IMGT position 95 and a tyrosine at IMGT position 96. Examples include IgG1, IgG2, and IgG4 proteins. In one embodiment, the ScFv fusion protein contains amino acid sequences set forth in SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, and SEQ ID NO:31. In one specific embodiment, the ScFv fusion protein comprises the amino acid sequence of SEQ

ID In a specific embodiment, the host cell comprises a cell surface capture protein positioned at the plasma membrane and bound to an IgG1, IgG2 or IgG4, or a bispecific antibody containing at least one heavy chain of an IgG1, IgG2 or IgG4, and which may have a second heavy chain that is of another type or contains one or more amino acid substitutions.

[0047] Also disclosed is a recombinant antigen-binding protein that binds a substituted CH3 polypeptide comprising one or more amino acid substitutions selected from the group consisting of (a) 95R, and (b) 95R and 96F according to the IMGT exon numbering system, or (a') 435R, and (b') 435R and 436F according to the EU numbering system, or any protein that comprises for example an amino acid sequence of SEQ ID NO:42, which encodes a substituted human Fc (also known as Fc*). In some instances, the recombinant antigen-binding protein binds the polypeptide with a K_D of less than about 60 nM as measured in a surface plasmon resonance assay.

[0048] In some instances, the recombinant antigen-binding protein comprises one or more complementarity determining regions (CDRs) of a heavy chain variable region (HCVR) having an amino acid sequence that is at least 95% identical to SEQ ID NO:38, or of a light chain variable region (LCVR) having an amino acid sequence that is at least 95% identical to SEQ ID NO:39. In one case, the protein comprises a heavy chain CDR-1 (HCDR1) having the amino acid sequence of SEQ ID NO:32, an HCDR-2 having the amino acid sequence of SEQ ID NO:33, an HCDR-3 having the amino acid sequence of SEQ ID NO:34, a light chain CDR-1 (LCDR-1) having the amino acid sequence of SEQ ID NO:35, and an LCDR-2 having the amino acid sequence of SEQ ID NO:36. In some cases, the protein comprises an HCVR having an amino acid sequence that is at least 95% identical to SEQ ID NO:38 (some of which are identical to SEQ ID NO:38) and an LCVR having an amino acid sequence that is at least 95% identical to SEQ ID NO:39 (some of which are identical to SEQ ID NO:39).

[0049] In some instances, the recombinant antigen-binding protein is an antibody, which comprises a heavy chain and a light chain. The heavy chain may comprise an amino acid sequence that is at least 95% identical to (or 100% identical to) SEQ ID NO:40. The light chain may comprise an amino acid sequence that is at least 95% identical to (or 100% identical to) SEQ ID NO:41. Recombinant antigen-binding proteins, which are antibodies, are used as detection molecules (DMs).

[0050] In some instances, the recombinant antigen-binding protein is an ScFv fusion protein, which in some cases comprises a heavy chain variable domain with an amino acid sequence that is at least 95% identical to (or identical to) SEQ ID NO:38, a light chain variable domain with an amino acid sequence that is at least 95% identical to (or identical to) SEQ ID NO:39, and a membrane anchor domain. In one instance, the membrane anchor domain is derived from an Fc receptor, such as the transmembrane domain of the human FcγR1 protein, as represented by SEQ ID NO:17, or SEQ ID NO:21, which contains not only the transmembrane domain, but also the C-terminal cytoplasmic domain of SEQ ID NO:19. In one specific instance, the ScFv fusion protein has the amino acid sequence of SEQ ID NO:43. Recombinant antigen-binding proteins, which are ScFv fusion proteins, are useful as CSCPs and as DMs.

[0051] Also disclosed is a polynucleotide that encodes the antigen-binding protein of the preceding aspect. In one embodiment, such as in the case where the antigen-binding protein is an antibody, the polynucleotide encodes the light chain, such as for example the light chain of SEQ ID NO:41. Likewise, the polynucleotide may encode the heavy chain, such as for example, the heavy chain of SEQ ID NO:40. In the case in which the antigen-binding protein is an ScFv fusion protein, the polynucleotide may encode the ScFv-FcγR-TM-cyto fusion protein of SEQ ID NO:43. Representative exemplar polynucleotides include those polynucleotides of SEQ ID NO:49, 50 and 51, respectively.

[0052] Also disclosed is a nucleic acid vector that encompasses the polynucleotide of the preceding aspect. In one instance, the vector comprises the polynucleotide, which encodes the antigen-binding protein, operably linked to an upstream promoter, and followed by a downstream polyadenylation sequence. The promoter can be any promoter, such as for example a CMV promoter. Thus in one case, the vector may contain the sequence of SEQ ID NO:47. In one instance, the vector may contain a nucleic acid sequence that encodes a selectable marker, such as for example neomycin resistance. In one embodiment, the vector may contain a nucleic acid sequence that encodes an energy transfer protein, such as green fluorescence protein (GFP), or a derivative thereof, such as yellow fluorescence protein (YFP). Thus in one case, the vector may contain the sequence of SEQ ID NO:46.

[0053] The vector may be circular or linear, episomal to a host cell's genome or integrated into the host cell's genome. In some instances, the vector is a circular plasmid, which in one specific embodiment has the nucleic acid sequence of SEQ ID NO:44 for the ScFv-FcγR-TM-cyto-encoding polynucleotide, in another specific instance has the nucleic acid sequence of the antibody heavy chain-encoding polynucleotide, and yet another specific embodiment has the nucleic acid sequence of the antibody light chain-encoding polynucleotide. In some instances, the vector is a linear construct, which may be integrated into a host cell chromosome. In one specific embodiment, the linear construct comprises the nucleic acid sequence of SEQ ID NO:51 for the ScFv-FcγR-TM-cyto-encoding polynucleotide. In another specific embodiment, the linear construct comprises the nucleic acid sequence of SEQ ID NO:50 for the antibody heavy chain-encoding polynucleotide. In yet another specific instance, the linear construct comprises the nucleic acid sequence of SEQ ID NO:49 for the antibody light chain-encoding polynucleotide.

[0054] The host cell may be any cell, prokaryotic or eukaryotic. However, in one specific embodiment, the host cell is a CHO cell, such as a CHO-K1 cell.

[0055] Also disclosed is a host cell that expresses the antigen-binding protein of the preceding aspect, and/or contains the polynucleotide or nucleic acid vector of the preceding aspects. In some embodiments, the host cell is a CHO cell. In a specific embodiment, the host cell is a CHO-K1 cell. In one embodiment, host cell is used in the production of a protein of interest, and the antigen-binding protein is used as a cell surface capture protein according to the methods disclosed in this application.

[0056] Also disclosed is a host cell useful in the production of a protein of interest. The host cell harbors a polynucleotide or nucleic acid vector of a preceding aspect, and produces an antigen-binding protein of a preceding aspect, which serves as a cell surface capture protein. The cell surface capture protein binds to the protein of interest inside the host cell, and is transported through the secretory apparatus of the cell, and is expressed on the surface of the host cell. Thus, in one instance, the host cell comprises a cell surface capture protein positioned in the host cell plasma membrane, with the capturing moiety facing outside of the cell. In one instance, the cell surface capture molecule is bound to a protein of interest, which is positioned at the plasma membrane and oriented outside of the cell.

[0057] In one instance, the host cell produces or is capable of producing an ScFv fusion protein that binds to a protein of interest that contains an Fc domain, which contains an arginine at IMGT position 95 and a phenylalanine at IMGT position 96 (Fc*). Examples include IgG3 and substituted CH3 regions of IgG1, IgG2, and IgG4 proteins. In one embodiment, the ScFv fusion protein contains amino acid sequences set forth in SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36 and SEQ ID NO:37. In one specific instance, the ScFv fusion protein comprises the amino acid sequence of SEQ ID NO:43. In a specific instance, the host cell comprises a cell surface capture protein positioned at the plasma membrane and bound to an IgG3 or a substituted IgG1, IgG2 or IgG4, which contain the arginine at IMGT position 95 and phenylalanine at IMGT position 96 ("Fc*"), or a bispecific antibody containing at least one heavy chain of a the Fc* type and the other heavy chain of the IgG1, IgG2 or IgG4 wildtype.

[0058] Also disclosed is a method of detecting, isolating, or enriching for a cell that stably expresses a protein of interest (POI). The method includes the step of expressing in the host cell a cell surface capture protein (CSCP) and a POI. According to this method, the CSCP binds to a "first site" on the POI to form a CSCP-POI complex inside the host cell. This CSCP-POI complex is then transported through the secretory system of the host cell, and is secreted from the cell. Since the CSCP contains a membrane binding domain (e.g., SEQ ID NO:17), the CSCP-POI complex is displayed on the surface of the host cell, with the POI exposed outside of the cell. According to the method, the host cell is then contacted with a detection molecule (DM), which binds to a "second site" on the POI. Those cells that bind the DM are selected for identification, isolation, pooling, and/or enrichment. In one embodiment, the DM-bound host cell is selected by fluorescence activated cell sorting.

[0059] In one instance, the method also includes the step of contacting the cell with a blocking molecule prior to selecting the host cell. The blocking molecule binds to any CSCP that is not bound to the POI. The blocking molecule does not bind to the CSCP-POI complex.

[0060] The POI contains multiple subunits, such as an antibody that comprises two heavy chains and two light chains. The first site on the POI resides on a first subunit, and the second site on the POI resides on a second subunit. The POI is a heterodimeric protein. Where the POI is an antibody, the first site on the POI resides on a first heavy chain, and the second site

on the POI may reside on a second heavy chain. the antibody contains subunits that differ by at least one amino acid, such as an antibody having at least one heavy chain with a wild type CH3 domain and the other heavy chain having at least one amino acid substitution in the CH3 domain. The CSCP comprises an antigen-binding protein as described herein, such as an anti-Ig ScFv fusion protein. Here, the detection molecule (DM) may comprise a labeled recombinant antigen-binding protein as described herein, such as a labeled antigen or anti-Ig antibody or ScFv molecule.

[0061] Where the POI is a bispecific antibody, the first site resides on a heavy chain that has a CH3 domain containing a histidine residue at position 95 according to the IMGT exon numbering system and a tyrosine residue at position 96 according to the IMGT exon numbering system (Fc). Then, the second site may reside on a heavy chain that has a CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*). In this case, the CSCP may be an antigen-binding protein described in a preceding aspect, such as an ScFv fusion protein containing the amino acid sequences of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, and SEQ ID NO:31; which in a specific embodiment comprises SEQ ID NO:19. Here also, the detection molecule (DM) may comprise a labeled recombinant antigen-binding protein described in a preceding aspect, such as an antibody or ScFv molecule containing the amino acid sequences of SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, and SEQ ID NO:37; which in a specific embodiment comprises either SEQ ID NO:40 and SEQ ID NO:41 (anti-Fc* antibody), or SEQ ID NO:43 (ScFv*). Here, the blocking molecule may be an Fc polypeptide (*e.g.*, single chain), such as hFc, or any molecule that can bind to the CSCP without also binding to the DM. In one embodiment, the detection molecule may be a labeled anti-human IgG F(ab')₂.

[0062] [Deleted]

[0063] Also disclosed is a method of detecting or isolating a cell that stably expresses a heterodimeric protein comprising the steps of (a) expressing in a host cell a cell surface capture protein (CSCP) and a heterodimeric protein, wherein (i) the CSCP binds to a first site on the heterodimeric protein to form a CSCP- heterodimeric protein complex inside the host cell, (ii) the CSCP- heterodimeric protein complex is transported through the host cell, and (iii) then displayed on the surface of the host cell; (b) contacting the host cell with a detection molecule, wherein the detection molecule binds to a second site on the heterodimeric protein; and (c) selecting the host cell which binds the detection molecule. In some instances, the heterodimeric protein comprises multiple subunits and the first site on the heterodimeric protein resides on a first subunit, and the second site on the heterodimeric protein resides on a second subunit. In some instances, the cell surface capture molecule comprises an antigen, Protein A, or ScFv capable of binding the first subunit and not the second subunit.

[0064] Also disclosed is a method of producing a bispecific antibody comprising the step of expressing in a host cell a cell surface capture protein ("CSCP"), an antibody light chain, a first antibody heavy chain, which contains a CH3 domain comprising a histidine at IMGT position 95

and a tyrosine at IMGT position 96, and a second antibody heavy chain, which contains a CH3 domain comprising an arginine at IMGT position 95 and a phenylalanine at IMGT position 96. While inside the host cell, the CSCP binds to the first antibody heavy chain but does not bind to the second antibody heavy chain, the second antibody heavy chain binds to the first antibody heavy chain, and the light chains bind to the heavy chains, thus forming a CSCP-Antibody ternary complex. This ternary complex is secreted and presented onto the surface of the host cell. The host cell may be contacted with a blocking molecule, which binds to a CSCP on the cell surface, but only in those situations in which the CSCP is not bound to the antibody-of-interest, i.e., an "empty" CSCP. The host cell is then contacted with a DM that binds to or is capable of binding to the second antibody heavy chain. The host cell that binds the DM is identified, selected, and/or pooled. In some embodiments, the host cells that bind the DM are selected, pooled, cultured and expanded, and then subjected to another round of expression, detection, selection, pooling and expansion. This process may be reiterated multiple times to enrich for the production of high titers of bispecific antibodies.

[0065] In one embodiment, the CSCP employed in the method is an ScFv-fusion protein containing the amino acid sequences of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, and SEQ ID NO:31. In one embodiment, the CSCP comprises the amino acid sequence of SEQ ID NO:31. In one embodiment, the DM employed in the method is a protein containing the amino acid sequences of SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, and SEQ ID NO:37. In one embodiment, the DM is an antibody comprising a heavy chain sequence of SEQ ID NO:40 and a light chain sequence of SEQ ID NO:41. In another embodiment, the DM is an ScFv fusion protein containing the amino acid sequence of SEQ ID NO:43. A label, for example a fluorescent moiety like FITC or Alexa Fluor® 488, may be attached to the DM. Fluorescence activated cell sorting may be used as the detection and selection means.

[0066] [Deleted]

[0067] [Deleted]

[0068] The host cell, which is the product of the iterative selection, pooling and expansion, may be capable of producing, or does produce bispecific antibody at a titer of at least 2 g/L, wherein the bispecific antibody species (Fc/Fc*) represents at least 40% by mass of the total antibody produced by the host cell (Fc/Fc + Fc*/Fc* + Fc/Fc*).

[0069] Other objects and advantages will become apparent from a review of the ensuing detailed description.

DETAILED DESCRIPTION

[0070] [Deleted]

[0071] As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, a reference to "a method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure and so forth.

[0072] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described.

General Description

[0073] The method of the invention provides substantial advantages over current methods for isolation and identification of protein-secreting cells. For example, cells that secrete antibodies may be rapidly and conveniently isolated based on desired specificity, avidity, or isotype. Furthermore, the amount of secreted protein produced may be directly quantified, unlike many methods in the prior art wherein production of secreted protein is indirectly quantified.

[0074] Recently, two additional methods that utilize flow cytometry have been developed for the high throughput isolation of stable high expression cell lines. The first method involves modification of the expression plasmid to include a transcriptional read out for the GOI mRNA. This is most often accomplished by inserting an internal ribosomal entry site (IRES) and a gene whose protein product is easily monitored by flow cytometry, most frequently green fluorescent protein (GFP), between the stop codon of the GOI and the terminal poly A site (Meng et al. (2000) Gene 242:201). The presence of an IRES allows the POI and GFP to be translated from the same mRNA. Therefore, the expression level of the GFP gene is indirectly related to the mRNA level for the GOI. Clones that accumulate the GFP at high levels are isolated by flow cytometry and then screened for POI production. Because this method depends on the coupling of GOI expression to the reporter gene by use of an IRES in a recombinant construction, it is not applicable to the isolation of hybridomas.

[0075] The use of flow cytometry in the isolation of expression clones allows for the rapid analysis of large numbers of clones in a high throughput format. Moreover, use of flow cytometry significantly reduces the direct handling of cells. Unfortunately, the level of GFP production is not a direct measure of the production level of the POI. Various mechanisms may uncouple the production of secreted POI from accumulation of GFP. Differences in production of the POI and the GFP reporter may result from differences in the translation efficiency of the two genes, secretion efficiency of the POI, or stability of the polycistronic mRNA.

[0076] Another method that uses flow cytometry to isolate expression clones involves encapsulation of cells within agarose microdrops (Weaver et al. (1990) Methods Enzymol.

2:234). In this method biotinylated antibodies specific for the POI are bound to the biotinylated agarose through streptavidin such that secreted POI is captured and retained within the microdrop (Gray et al., (1995) J. Immunol. Methods 182:155). The trapped POI is detected by immuno-staining with an antibody specific for the POI. To reduce the encapsulating agarose from absorbing POI secreted from adjacent cells, the cells are placed in a low-permeability medium. Those cells with the highest antibody staining of the POI in the embedding agarose are identified and isolated by flow cytometry. The gel microdrop approach screens cells directly for their ability to secrete POI, rather than indirectly screening for expression of GOI mRNA, but requires the availability of suitable antibodies for trapping and staining the secreted POI and the procedure requires special equipment to generate the agarose gel microdrops. Moreover, some cells may be sensitive to the encapsulation process.

[0077] A variation of this method circumvents the requirement for embedding cells in a matrix by directly binding an antibody, specific for the POI, to the cell surface (Manz et al. (1995) PNAS 92:1921-1925). In this method, non-specific biotinylation of cell surface proteins with biotin-hydroxysuccinimide ester is followed by contact with a streptavidin-conjugated antibody capable of binding the POI. Cells secreting the POI become decorated with the POI which is then detected with an appropriately labeled second antibody. However, diffusion of POI between neighboring cells is problematic, and this method also requires a high viscosity medium to reduce diffusion of POI away from expressing cells. Because these high viscosity media are required for discriminating cells, the cells must be washed and placed in a medium suitable for cell sorting if so desired.

[0078] The problems associated with identification and isolation of high expression recombinant cell lines especially applies to the isolation of hybridomas that express an antibody of interest. However, the identification of useful hybridomas includes several additional problems; they must be screened first for antigen-binding activity, then for immunoglobulin isotype. Moreover, GFP-based methods are not applicable to the identification and isolation of hybridomas because construction of hybridomas does not include a recombinant construct such that expression of the antibody genes can be linked to a transcriptional reporter such as GFP. Hybridoma screening is a slow, laborious endeavor where the number of clones screened is limited by existing technologies.

[0079] The instant invention describes a novel and previously unknown method of identifying and isolating cells that produce secreted proteins. The invention is based on the production of a cell line that expresses a molecule, localized to the cell surface, which binds the POI. The cell surface-displayed POI can then be detected by labeling with various detection molecules. The amount of POI displayed on the cell surface, under specific conditions, is a direct measure of the total amount of POI secreted. POI producers may then be isolated from non-producers, and levels of production or POI characteristics may be differentiated. The advantage of the invention is that it directly quantifies the secreted POI rather than indirectly measuring the mRNA.

[0080] The present invention provides a recombinant antigen-binding protein that binds a

human IgG1-Fc domain, a human IgG2-Fc domain, or a human IgG4-Fc domain, wherein:

1. (a) the recombinant antigen-binding protein is specific for a wild type CH3 domain containing a histidine residue at position 95 according to the IMGT exon numbering system and a tyrosine residue at position 96 according to the IMGT exon numbering system (Fc) over a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*); and
2. (b) the recombinant antigen-binding protein comprises: (i) a ScFv which comprises a heavy chain CDR-1 (HCDR-1) having the amino acid sequence of SEQ ID NO:27, a HCDR-2 having the amino acid sequence of SEQ ID NO:28, a HCDR-3 having the amino acid sequence of SEQ ID NO:29, and a light chain CDR-1 (LCDR-1) having the amino acid sequence of SEQ ID NO:30, and a LCDR-2 having the amino acid sequence of SEQ ID NO:31; and (ii) a membrane anchor.

[0081] The present invention further provides A method of detecting or isolating a cell that stably expresses a heterodimeric protein comprising the steps of:

1. (a) expressing in a host cell a cell surface capture protein (CSCP) comprising an antigen-binding protein of any one of claims 1 to 3 and a heterodimeric protein, wherein (i) the CSCP binds to a first site on the heterodimeric protein to form a CSCP-heterodimeric protein complex inside the host cell, (ii) the CSCP- heterodimeric protein complex is transported through the host cell, and (iii) then displayed on the surface of the host cell;
2. (b) contacting the host cell with a detection molecule, wherein the detection molecule binds to a second site on the heterodimeric protein; and
3. (c) selecting the host cell which binds the detection molecule, wherein in the heterodimeric protein:
 1. (i) the first site on the heterodimeric protein resides on a first subunit comprising a heavy chain comprising a wild type CH3 domain containing a histidine residue at position 95 according to the IMGT exon numbering system and a tyrosine residue at position 96 according to the IMGT exon numbering system (Fc); and
 2. (ii) the second site on the heterodimeric protein resides on a second subunit comprising a heavy chain comprising a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*);

wherein the heterodimeric protein comprises a bispecific antibody or an Fc-containing fusion protein..

[0082] This invention therefore relates to the construction or use of cells that express cell surface capture molecules which bind various secreted POIs in the same cell that produces the

POI. As the cell secretes the POI, these cell surface capture molecules bind it, or complexes of POI and cell surface capture molecules may form intracellularly and then get secreted. Binding may occur in an autocrine manner or while being secreted. The cells that produce the secreted POI may then be identified and isolated. Such identification and isolation may be based on characteristics of the POI, production of the POI or lack thereof, or by specified levels of production. The cell surface capture molecule and/or the POI may be produced by the cell in its native state, or the cell surface capture molecules and/or the POI may be recombinantly produced. Through the construction or use of such a cell, any secreted protein may be captured by the cell surface capture molecule provided there is a corresponding affinity between the two. As explained further, any molecule may be manipulated such that it can be used as a cell surface capture molecule. Therefore, this invention may be utilized to isolate any cell that secretes a protein.

[0083] The surface capture protein (CSCP) comprising an antigen-binding protein and a membrane anchor domain . What is necessary is the ability of the desired protein to be anchored to the cell membrane and exposed to the extracellular space. If the desired cell has a signal sequence then only a membrane anchor, including but not limited to a transmembrane anchor or a GPI linkage signal, need be added to the cell surface capture molecule such that it remains anchored in the cell membrane exposed to the outside of the cell. Furthermore, if the desired protein lacks a signal sequence, a signal sequence may be added to the amino terminus of the desired protein, such that it is transported to the cell surface. A signal sequence and a membrane anchor may be native to the cell, recombinant, or synthetic.

[0084] Cells often secrete a wide variety of proteins, endogenously or following the introduction of recombinant DNA. Any secreted heterodimeric protein may be identified and the cell producing it may be isolated according to the method of this invention. Such secreted proteins include but are not limited to growth factors, growth factor receptors, ligands, soluble receptor components, antibodies, bispecific antibodies, recombinant Trap molecules, Fc-containing fusion proteins, sTCRs, TCR-Fc's, and peptide hormones. Such secreted proteins may or may not be recombinant. That is, the secretion of some proteins of interest from the desired cell may not require the introduction of additional nucleotide sequences. For example, the secretion of antibodies from B-cells or plasma cells is not the result of introduction of recombinant nucleotide sequences into the B-cell or plasma cell. Recombinant secreted proteins may be produced by standard molecular biology techniques well known to the skilled artisan (see e.g., Sambrook, J., E. F. Fritsch And T. Maniatis. *Molecular Cloning: A Laboratory Manual*, Second Edition, Vols 1, 2, and 3, 1989; *Current Protocols in Molecular Biology*, Eds. Ausubel et al., Greene Publ. Assoc., Wiley Interscience, NY). These secreted proteins are useful for many commercial and research purposes. This invention encompasses the production of such secreted proteins through the methodologies of the invention. Detection of the cells with the displayed POI is accomplished through the use of a detection molecule binds to the second site on the heterodimeric protein, wherein the second site resides on a second subunit comprising a heavy chain comprising a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*)a. Such detection

molecules may facilitate the detection and/or isolation of the cells displaying the POI.

[0085] The disclosure is applicable to the isolation of, inter alia, a) ligand-producing cells by using the ligand-specific receptor as the cell surface capture molecule, b) soluble receptor-producing cells by using a surface bound receptor-specific ligand as the cell surface capture molecule, c) antibody-producing cells by using an antibody-binding protein as the cell surface capture molecule, d) sTCR's by using an s-TCR-binding protein (e.g., and antigen recognized by the TCR) as the cell surface capture molecule, e) TCR-Fc's, by using an Fc-binding protein as a cell surface capture molecule, or f) bispecific antibodies that harbor a mutation in one of its CH3 domains that abrogates protein A binding, by using a fusion protein capture molecule that comprises an ScFv domain fused to an FcγR transmembrane and cytoplasmic domain.

[0086] In accordance with the methodology disclosed, a cell is first transfected with a vector containing a nucleotide sequence that encodes a cell surface capture molecule that is capable of binding the secreted POI, under conditions in which such cell surface capture molecule is expressed. Transfected cells which are appropriate producers of such cell surface capture molecules are then detected and isolated, and such cells are cultured. These cells may either naturally produce the POI, or the POI may be recombinantly produced. If the cells naturally produce the POI, they are ready for detection and isolation. If the POI is to be recombinantly produced, then the isolated and cultured cells expressing the specified cell surface capture molecule are transfected with second nucleotide sequence that encodes the secreted POI, under conditions in which the secreted POI is expressed. Upon expression, the secreted POI binds to the cell surface capture molecules and the cells displaying bound POI are detected and isolated.

[0087] If the POI is naturally produced by the cell, the cell will not be transfected with nucleotide sequence encoding the POI. Therefore, this aspect of the invention is applicable to any and all cells producing a POI. In addition, if the cell surface capture molecule is naturally produced by the cell, the cell need not be transfected with nucleotide sequences encoding the cell surface capture molecule. Therefore, this aspect of the invention is applicable to any and all cells producing a cell surface capture molecule.

[0088] A wide variety of host cells may be transfected. These cells may be either of eukaryotic or of prokaryotic origin. The cells will often be immortalized eukaryotic cells, and in particular, mammalian cells, for example monkey kidney cells (COS), Chinese hamster ovary cells (CHO), HeLa cells, baby hamster kidney cells (BHK), human embryonic kidney cells (HEK293), leukocytes, myelomas, cell lines transfected with adenovirus genes, for example, AD5 E1, including but not limited to immortalized human retinal cells transfected with an adenovirus gene, for example, PER.C6™ cells, and embryonic stem cells. The cells may also be non mammalian cells including bacterial, fungi, yeast and insect cells, including, but not limited to, for example *Escherichia coli*, *Bacillus subtilis*, *Aspergillus* species, *Saccharomyces cerevisiae*, and *Pichia pastoris*. All cells may be grown in culture trays medium under appropriate conditions or in a synergistic host. The most desirable cells will be mammalian cells capable of culture.

[0089] The secreted POI bound to the cell surface capture molecule is detected using a detection molecule, wherein the detection molecule binds to a second site on the heterodimeric protein, wherein the second site on the heterodimeric protein resides on a second subunit comprising a heavy chain comprising a substituted CH3 domain containing an arginine residue at position 95 according to the IMGT exon numbering system and a phenylalanine residue at position 96 according to the IMGT exon numbering system (Fc*). Cultures cells displaying the secreted POI may be contacted with the detection molecule wherein such detection molecule(s) may contain a detection label, such as, for example, a chromogenic, fluorogenic, colored, fluorescent, or magnetic label. The label bound to the detection molecule may be detected and the cell isolated using various methods. Most preferably, within a cell population the label will be detected and the cell isolated utilizing flow cytometry. Alternatively, the detection molecule may be used for the direct isolation of cells displaying the POI. This may be accomplished by conjugation of the detection molecule to a culture plate, paramagnetic molecules, or any other particle or solid support. In addition, displayed POI may be detected directly by a property of the detection molecule or the POI.

[0090] [Deleted]

[0091] In any of the embodiments and with regards to isolating expressing cells from non-expressing cells or lesser expressing cells, one of the principal difficulties, when the POI is a secreted protein, is diffusion of POI between neighboring cells. Therefore, it is critical that any system that is designed to capture the secreted POI on the cell surface must prevent the diffusion of the POI from the expressing cell to a neighboring cell and its adherence to that cell. If diffusion is allowed to occur, and neighboring cells become decorated with the secreted POI, then separation of cells based upon the degree of POI decoration will fail to discriminate high expressing cells from cells with low expression levels, and may fail to effectively isolate expressing from non-expressing cells.

[0092] Therefore one embodiment of this invention is to block the diffusion of the secreted POI between neighboring cells. This may be accomplished by the addition of a blocking molecule that binds either the cell surface capture molecule or the POI and prevents the binding of the secreted POI to the cell surface capture molecule. In this aspect, the detection molecules do not bind the blocking molecule. For example, if the cell surface receptor is the hFcγRI and the secreted POI possesses the human IgG Fc fragment, then diffusion of the secreted POI between neighboring cells may be blocked by the addition of exogenous rat IgG to the culture media. Detection of cells displaying secreted POI, and not bound rat IgG, is achieved by use of antibodies specific for human IgG Fc that do not recognize rat IgG. In another embodiment, binding of the secreted POI between neighboring cells is reduced by increasing the viscosity of the media.

[0093] In one embodiment of this invention, the secreted POI is not allowed to accumulate in the media. This may be accomplished by regulating the expression of the secreted POI and/or the cell surface capture molecule such that brief expression of the POI results in sufficient POI

to bind the cell surface capture molecule but insufficient amounts for diffusion. In another embodiment, cells may be removed from the media containing accumulated POI, the POI bound to the cells is stripped off, and POI expression is allowed to continue for a limited period of time such that secreted POI does not accumulate in the media. Proteins may be stripped by methods known in the art, for example, washing cells with low pH buffer.

[0094] According to this invention, those cells in a cell population that bind the most detection molecules also express the most secreted POI. In fact, the more POI that an individual cell secretes, the more POI is displayed on the cell surface. This correlation between the amount of surface-displayed POI and the expression level of the POI in that cell allows one to rapidly identify cells with a desired relative expression level from a population of cells.

[0095] Also disclosed is the use of a DNA library to express secreted protein which may be displayed on the cell surface by the cell surface capture molecule. For example, a library of DNA may also be generated from the coding regions of the antibody variable domains from B-cells isolated from immunized animals. The DNA library may then be expressed in a cell that expresses a cell surface capture molecule specific for antibodies such that clones of desired specificity, isotype, or avidity may be identified and isolated by the method of the invention. In another embodiment, a library of DNA may be generated from the coding regions of T cell receptor variable domains from T-cells, and fused to, for example, an Fc capable of binding to an Fc-binding protein. The DNA library may then be expressed in a cell that expresses an Fc-binding protein such that clones of desired specificity, isotype, or avidity may be identified and isolated as described herein.

[0096] Also disclosed is the creation of transgenic mammals that express a particular cell surface capture molecule in one or more cell types. The cells from such transgenic mammals may then be screened directly for the production of a POI. For example, it may be desirable to express a cell surface capture molecule, specific for antibodies, in plasma cells. Accordingly, plasma cells from immunized mice may be harvested and those cells producing antibodies specific to the desired antigen may be isolated by the method of the invention.

[0097] In a further embodiment of the invention, antibody production is measured through the use of a CHO cell line that expresses the human FcγR1 receptor (FcγRI) which binds the particular antibody or TCR-Fc that is the POI.

[0098] In another aspect of the invention, the protein of interest comprises one or more T cell receptor variable domains or a soluble T cell receptor. The one or more T cell receptor variable domains can be covalently linked to a moiety that can bind a cell surface capture protein. In a specific embodiment, the one or more T cell receptor variable domains are fused to an Fc sequence, e.g., a human Fc sequence, and the cell surface capture protein is an Fc receptor, e.g., an FcγR.

[0099] The general structures of TCR variable domains are known (see, e.g., Lefranc and Lefranc (2001) *The T Cell Receptor FactsBook*, Academic Press; see, e.g., pp. 17-20; see

also, Lefranc et al. (2003) IMGT unique numbering for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains, *Developmental and Comparative Immunology* 27:55-77, and Lefranc et al. (2005) IMGT unique numbering for immunoglobulin and T cell receptor constant domains and Ig superfamily C-like domains, *Developmental and Comparative Immunology* 29:185-203). In one embodiment, a TCR variable domain of a TCR-Fc comprises an N-terminal region having a variable domain of 104-125 amino acids. In another embodiment, the TCR-Fc further comprises a TCR constant region comprising 91-129 amino acids. In another embodiment, the TCR-Fc further comprises a connecting peptide comprising 21-62 amino acids.

[0100] In one embodiment, the Fc sequence is fused directly or through a linker to the TCR variable domain. In another embodiment, the TCR-Fc comprises a TCR variable region and a TCR constant region, and the Fc sequence is fused directly or through a linker to the TCR constant region. In another embodiment, the TCR-Fc comprises a TCR variable region, a TCR constant region, and a connecting peptide, and the Fc sequence is fused directly or through a linker to the connecting peptide.

[0101] The sTCR, TCR-Fc, or fusion protein comprising one or more T cell receptor variable regions can be selected so as to specifically bind an antigen of interest, for example, a substance produced by a tumor cell, for example, tumor cell substance that is capable of producing an immune response in a host. In a specific embodiment, the antigen is an antigen that is present on the surface of a tumor cell (i.e., a tumor antigen), is recognized by a T cell, and that produces an immune response in a host. Tumor antigens include, for example, alphafetoprotein (AFP), carcinoembryonic antigen (CEA), MUC-1, epithelial tumor antigen (ETA), tyrosinase (e.g., for malignant melanoma), melanoma-associated antigen (MAGE), and mutated or abnormal forms of other proteins such as, for example, ras, p53, etc.

[0102] In one embodiment, the POI is a TCR-Fc, and the TCR-Fc comprises a TCR α chain variable region fused to an Fc sequence and a TCR β chain fused to the Fc sequence (each directly or through a linker), wherein the TCR α chain-Fc fusion and the TCR β chain-Fc fusion associate to form an $\alpha\beta$ TCR-Fc. In a specific embodiment, the $\alpha\beta$ TCR-Fc comprises the following two polypeptides: (1) a TCR α chain variable region fused to a TCR α chain constant region fused to an Fc sequence, and (2) a TCR β chain variable region fused to a TCR β chain constant region fused to an Fc sequence.

[0103] In another embodiment, the POI is a TCR-Fc having a TCR α variable region and a TCR β variable region and, optionally, a TCR α constant region and/or a TCR β constant region. In a specific embodiment, the TCR-Fc is encoded by a nucleic acid comprising (5' to 3') a TCR α variable region sequence, optionally followed by a TCR α constant region sequence, a TCR β variable region sequence, optionally followed by a TCR β constant region sequence, optionally a linker, then an Fc sequence. In a specific embodiment, the TCR-Fc is encoded by a nucleic acid comprising (5' to 3') a TCR β variable region sequence, optionally followed by a TCR β constant region sequence, a TCR α variable region sequence, optionally followed by a TCR α constant region sequence, optionally a linker, then an Fc sequence. In various

embodiments, constructs encoding TCR-Fc's are preceded by signal sequences, e.g., secretion signal sequences, to render them secretable.

[0104] In another embodiment, the POI is a TCR-Fc, and the TCR-Fc comprises a TCR-Fc comprising a TCR γ chain fused to an Fc sequence and a TCR δ chain variable region fused to an Fc sequence to form a $\gamma\delta$ TCR-Fc. In a specific embodiment, the $\gamma\delta$ TCR-Fc comprises the following two polypeptides: a TCR γ chain variable region fused to a TCR γ chain constant region fused to an Fc sequence, and (2) a TCR δ chain variable region fused to a TCR δ chain constant region fused to an Fc sequence.

[0105] T cell receptor variable regions can be identified and/or cloned by any method known in the art. The T cell receptor variable regions of the protein of interest are obtainable, for example, by expressing rearranged T cell receptor variable region DNA in a cell, for example, fused to a human Fc sequence. Rearranged T cell receptor variable regions specific for a particular antigen can be obtained by any suitable method known in the art (see references below), for example, by exposing a mouse to an antigen and isolating T cells of the mouse, making hybridomas of the T cells of the mouse, and screening the hybridomas with the antigen of interest to obtain a hybridoma of interest. Rearranged T cell variable regions specific for the antigen of interest can be cloned from the hybridoma(s) of interest. T cell receptor variable regions specific for an antigen can also be identified using phage display technology, for example, as provided in references below. The variable regions can then be cloned and fused, for example, to a human Fc to make a protein of interest that can bind to a cell surface capture molecule that is an Fc γ R.

[0106] Methods for identifying and/or cloning T cell receptor variable regions are described, for example, in US Patent No. 5,635,354 (primers and cloning methods); Genevée et al. (1992) An experimentally validated panel of subfamily-specific oligonucleotide primers (V α 1-w29/V β 1-w24) for the study of human T cell receptor variable V gene segment usage by polymerase chain reaction, *Eur. J. Immunol.* 22:1261-1269 (primers and cloning methods); Gorski et al. (1994) Circulating T Cell Repertoire Complexity in Normal Individuals and Bone Marrow Recipients Analyzed by CDR3 Size Spectratyping, *J. Immunol.* 152:5109-5119 (primers and cloning methods); Johnston, S. et al. (1995) A novel method for sequencing members of multi-gene families, *Nucleic Acids Res.* 23/15:3074-3075 (primers and cloning methods); Pannetier et al. (1995) T-cell repertoire diversity and clonal expansions in normal and clinical samples, *Immunology Today* 16/4:176-181 (cloning methods); Hinz, T. and Kabelitz, D. (2000) Identification of the T-cell receptor alpha variable (TRAV) gene(s) in T-cell malignancies, *J. Immunol. Methods* 246:145-148 (cloning methods); van Dongen et al. (2002) Design and standardization of PCR primers and protocols for detection of clonal immunoglobulin and T-cell receptor gene recombinations in suspect lymphoproliferations: US Patent No. 6,623,957 (cloning methods and primers); Report of the BIOMED-2 Concerted Action BMH4-CT98-3936, *Leukemia* 17:2257-2317 (primers and cloning methods); Hodges et al. (2002) Diagnostic role of tests for T cell receptor (TCR) genes, *J. Clin. Pathol.* 56:1-11 (cloning methods); Moysey, R. et al. (2004) Amplification and one-step expression cloning of human T cell receptor genes, *Anal. Biochem.* 326:284-286 (cloning methods); Fernandes et al. (2005) Simplified Fluorescent

Multiplex PCR Method for Evaluation of the T-Cell Receptor V β -Chain Repertoire, Clin. Diag. Lab. Immunol. 12/4:477-483 (primers and cloning methods); Li, Y. et al. (2005) Directed evolution of human T-cell receptors with picomolar affinities by phage display, Nature Biotech. 23/3:349-354 (primers and cloning methods); Wlodarski et al. (2005) Pathologic clonal cytotoxic T-cell responses: nonrandom nature of the T-cell receptor restriction in large granular lymphocyte leukemia, Blood 106/8:2769-2780 (cloning methods); Wlodarski et al. (2006) Molecular strategies for detection and quantitation of clonal cytotoxic T-cell responses in aplastic anemia and myelodysplastic syndrome, Blood 108/8:2632-2641 (primers and cloning methods); Boria et al. (2008) Primer sets for cloning the human repertoire of T cell Receptor Variable regions, BMC Immunology 9:50 (primers and cloning methods); Richman, S. and Kranz, D. (2007) Display, engineering, and applications of antigen-specific T cell receptors, Biomolecular Engineering 24:361-373 (cloning methods). Examples of sTCRs are provided in, for example, US Patent Nos. 6,080,840 and 7,329,731; and, Laugel, B et al. (2005) Design of Soluble Recombinant T Cell Receptors for Antigen Targeting and T Cell Inhibition, J. Biol. Chem. 280:1882-1892. Fc sequences are disclosed herein; examples of Fc sequences, and their use in fusion proteins, are provided, for example, in US Pat. No. 6,927,044 to Stahl et al.

[0107] In a further disclosure, the cell surface capture molecule is designed to engage and display those proteins of interest that are normally incapable of binding with sufficient affinity or bind with low affinity to an Fc γ R capture molecule. Those proteins of interest include IgG4 and IgG2 molecules. Thus, a modular capture molecule was designed and built based upon an ScFv domain fused to an Fc γ R transmembrane and cytoplasmic domain. The ScFv domain was derived from a high affinity anti-humanFc antibody, and contains a heavy chain variable domain fused to a light chain variable domain. The Fc γ R-TM-cytoplasmic domain was used to enable the proper insertion and orientation in the plasma membrane. The ScFv-Fc γ R-TM-cyto fusion protein is capable of binding IgG4 and other Fc containing molecules, as well as IgG2 and IgG1 subtypes, and those heterodimeric (e.g., bispecific antibodies) comprising at least one wild type CH3 domain, wherein the other CH3 domain may contain an Fc*-type substitution.

[0108] [Deleted]

EXAMPLES

[0109] Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1

[0110] Construction of pTE084. pTE084 was constructed by ligating the 1,436 bp Xba I fragment from pCAE100 that encodes the human FcγRI (hFcγRI; GenBank accession number M21091) into the Xba I site of pRG821. The orientation of hFcγRI in desirable plasmids resulting from the ligation was examined by restriction mapping with Not I, Pst I, Eco RI, and Stu I. pTE084 was designed for the high level expression of hFcγRI, the high affinity cell surface receptor for the Fc domain of human IgG. It contains two independent expression cassettes. One cassette is a hFcγRI gene driven by the CMV-MIE promoter, and the second cassette is the neomycin phosphotransferase II (*npt*) gene, which confers resistance to G418, driven by the SV40 late promoter.

[0111] Construction of a CHO K1 derivative that expresses hFcγRI. CHO K1 cells (4×10^6) were transfected with pTE084 using Lipofectamine™ (Life Technologies; Rockville, MD) following manufacturer's suggestions. The cells were placed in the culture medium (10% fetal bovine serum, 90% Ham's F-12, 2 mM L-glutamine; all reagents were from Life Technologies, Rockville, MD) containing 500 μg/ml G418 (Life Technologies) for 15 days. The cells that survived G418 selection were trypsinized, pooled, and stained with FITC-conjugated human IgG, Fc fragment (FITC-hFc; Jackson ImmunoResearch Laboratories, West Grove, PA). Briefly, the cells grown on 10 cm culture plates were washed once with Dulbecco's phosphate-buffered saline (PBS) without calcium chloride and magnesium chloride (Life Technologies). Three milliliters of 0.25% trypsin (Life Technologies) were added to each plate. The plates were swirled until the cells detached from the plate. Ten milliliters of culture medium was immediately added to each plate of the detached cells. The cells were then collected by centrifugation at 1,000 x g for 4 minutes. After removal of supernatant, the cells were resuspended in 4 ml of 2 μg/ml FITC-hFc diluted in culture medium. The cells were then placed on a platform shaker and stained for one hour at room temperature. To remove unbound FITC-hFc, the cells were washed twice with 20 ml PBS. The degree of FITC-hFc label on the cells was measured by flow cytometry on a MOFLO™ cell sorter (Cytomation; Fort Collins, CO). The FITC-hFc did not stain mock-transfected parental CHO K1 cells but gave rise to a distribution of fluorescence in the G418-resistant, pTE084-transfected pool. The top 1% most fluorescent cells from the selected pool were placed into 96-well plates at 1 cell/well by flow cytometry. Nine days later, 88 cell clones in the 96-well plates were expanded into 24-well plates. After 3 days, the cells in individual wells were washed once with 1 ml PBS, stained with 0.5 ml of 2 μg/ml FITC-hFc for 1 hour, washed twice with 1 ml PBS and examined for cell surface staining under a fluorescent microscope. The thirty three most fluorescent clones were chosen, expanded, then screened by flow cytometry.

[0112] Diffusion of secreted protein between expressing cells and non-expressing cells among cells was blocked by adding IgG: As all cells in a hFcγRI clonal cell line express a cell surface hFcγRI, they all possess the ability to bind IgG or fusion proteins consisting of the Fc domain of IgG. Because hFcγRI binds IgG from a variety of species (van de Winkel and Anderson, 1991), a panel of animal IgGs was tested for the ability to block the binding of a protein containing a human IgG1 (hIgG1) Fc tag (4SC622) to hFcγRI-expressing cells. 4SC622 is a chimeric molecule consisting of IL-2Rγ extracellular domain fused to the hIL-4Rγ extracellular domain

which is then fused to the hlgG1-Fc domain. In this experiment, cultures of RGC1, an hFcyRI-expressing cell line selected from CHO K1 cells that have been stably transfected with pTE084, were incubated with 1 µg/ml 4SC622 for 18 hours in the presence or absence of 1 mg/ml IgG from different species in a 37°C tissue culture incubator.

[0113] Cell surface binding of 4SC622 was determined by flow cytometry after washed cells were stained with phycoerythrin-conjugated mouse IgG1 monoclonal AG184 (PE-AG184) specific for the hIL-2R γ component of 4SC622 (BD Pharmingen; San Diego, CA), following procedures outlined for cell staining with FITC-hFc.

[0114] It was found that hlgG completely blocked 4SC622 from binding to the hFcyR1 expressed on the surface of RGC1. Rat, rabbit and canine-derived IgG also effectively blocked binding whereas bovine and ovine-derived IgG did not block. The ability of exogenously added rat IgG to block the binding of an exogenously added hlgG1 Fc-tagged protein (4SC622) to cell surface hFcyRI suggests that rat IgG can also block transfer between cells expressing a hlgG1 Fc-tagged protein at different levels. To test this, two cell lines that can be distinguished by the presence or absence of the green fluorescent protein (EGFP) were generated from RGC1. Briefly, to mark RGC1 cells with EGFP, 2×10^6 RGC1 cells were co-transfected with 0.5 mg PTE073 which encodes a hygromycin B phosphotransferase gene driven by phosphoglycerate kinase promoter, and 5 mg pRG816-EGFP which encodes EGFP gene driven by CMV-MIE promoter. The transfected cells were selected with 200 µg/ml hygromycin B (Sigma; St. Louis, MO) for two weeks. Green fluorescent cells were isolated by flow cytometry. One EGFP and hFcyRI-expressing clone, RGC2, was used in cell mixing experiments. The other cell line used in these experiments, RGC4, was generated by stable transfection of RGC1 with plasmid pEE14.1-622. pEE14.1-622 is a plasmid in which expression of 4SC622 is driven by the CMV-MIE promoter and includes a glutamine synthetase minigene, which confers resistance to the analog methionine sulfoximine (MSX), and allows for selection of stable integration events. RGC4 cells express hFcyRI on the cell surface and secrete the hlgG1 Fc-tagged protein 4SC622. One plate of mixed cells comprising 50% RGC2 and 50% RGC4 cells was incubated with 1 mg/ml rat IgG for 18 hours prior to staining with PE-AG184 then examined by flow cytometry. EGFP fluorescence of RGC2 cells shows that RGC2 cells also bind exogenously added 4SC622 (1 µg/ml) as indicated by an increase in PE-AG184 fluorescence. RGC4 did not fluoresce in the EGFP gate. Significantly, exogenously added rat IgG did not reduce the percentage of RGC4 cells that stained positive for cell surface 4SC622, suggesting that the binding of 4SC622 to hFcyRI occurred while the proteins were in transit to the cell surface. When RGC2 and RGC4 cells were mixed, the 4SC622 protein secreted from RGC4 cells accumulated in the medium and bound most of the RGC2 cells. However, the addition of 1 mg/ml rat IgG significantly reduced the percentage of RGC2 cells that bound 4SC622, demonstrating that rat IgG blocked the transfer of secreted hlgG1 Fc-tagged protein from expressing cells to non-expressing cells.

Example 2: Cell surface fluorescence correlates with the expression level of 4SC622

[0115] RGC1 cells (4×10^6) were transfected with pEE14.1-622 and a pool of stable transfectants was obtained after selection for 2 weeks in medium comprised of 10% dialyzed fetal bovine serum, 90% glutamine-free Dulbecco's Modified Eagle's Medium (DMEM), 1 x GS supplement, and 25 μ M MSX (All reagents were from JRH Biosciences, Lenexa, KS). Rat IgG was added to the culture medium to 1 mg/ml 18 hours prior to immunostaining. The cells were trypsinized, washed with PBS, and stained with 1.5 μ g/ml of a polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment (Jackson ImmunoResearch Laboratories) for one hour at room temperature following procedures as described for FITC-hFc staining in Example 1. Cell staining was then analyzed by flow cytometry. The distribution of fluorescence suggested that the selected pool contained cells with a wide range of 4SC622 expression levels. Cells in the top 3% (R3 bracket), 7-11% (R5 bracket), and 15-19% (R7 bracket) with respect to their immunofluorescence were sorted into three distinct pools and expanded for 9 days. Average 4SC622 production per cell for the pools was determined by measuring cell numbers and 4SC622 levels in the media after 3 days growth by an immuno-based Pandex assay (Idexx; Westbrook, ME) following the manufacturer's recommendations. In the Pandex assay, fluoricon polystyrene assay particles coated with goat anti-human IgG, g-chain specific antibody (Sigma) were used to capture 4SC622 from the medium, and a FITC-conjugated goat anti-human IgG, Fc specific (Sigma) was used to detect bead-bound 4SC622. Known amounts of purified 4SC622 were included in the assay for calibration. Cells in the top 3%, 7-11%, and 15-19% pool were found to produce 4SC622 at 1.42, 0.36, and 0.22 pg/cell/day, respectively. Thus, there was a correlation between cell surface 4SC622 staining and specific protein production. This result suggests that individual cells that express 4SC622 at high levels may be obtained by isolating cells that were stained brightest by the polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment.

Example 3: Isolation of expression clones in RGC1: IL-4 Trap

[0116] To directly demonstrate the efficiency in generating clonal cell lines with high level secreted protein production by our methodology, clonal 4SC622 producing cell lines were generated from RGC1. RGC1 cells (4×10^6) were transfected with pEE14.1-622, and selected for two weeks with 25 μ M MSX to obtain a pool of stable transfectants. MSX-resistant cells were pooled and incubated with 1 mg/ml human IgG for 18 hours, prior to staining with PE-AG184. Six cells from the top 5% gate, as determined by flow cytometry analysis of cell surface 4SC622 staining, were isolated and expanded. 4SC622 production from the six clonal lines was determined and compared to 4SC622 production from clones obtained by hand-picking selected colonies followed by dilution cloning and amplification. One RGC1-derived clone, RGC4, produced 4SC622 at 12 pg/cell/day. This level is similar to that of the best 4SC622 producer isolated by hand-picking and analyzing 2,700 clones. Thus, compared with hand-picking colonies, the methodology outlined in this invention proves to be far more efficient in the screening and cloning of high producers.

[0117] VEGF Trap. Plasmids pTE080 and pTE081 encode the genes for VEGF Traps, hVEGF-

R1R2 and hVEGF-R1 R3. hVEGF-R1 R2 is a chimeric molecule consisting of the first Ig domain of hVEGFR1 fused to the second Ig domain of hVEGFR2 which is then fused to the hlg1FC domain. hVEGF-R1R3 is a chimeric molecule consisting of the first Ig domain of hVEGFR1 fused to the second Ig domain of hVEGFR3 which is then fused to the hlgG1-Fc domain. In these plasmids, the gene for the VEGF Trap is driven by the CMV-MIE promoter and a glutamine synthetase minigene, which confers resistance to MSX, is expressed for selection of stable integration events. RGC1 cells were transfected with either of these plasmids and grown in medium containing 25 μ M MSX for 2 weeks to select for cells in which the plasmid has stably integrated. MSX-resistant cells were incubated with 0.1 μ g/ml IgG2a and mouse IgG3 for 18 hours prior to staining with 1.5 μ g/ml polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment. Cells were stained for 1 hour then washed twice with PBS prior to flow cytometry. Single cells were sorted into 96-well tissue culture plates from the pool of cells whose fluorescence was among the highest 1%. The cells in individual wells were expanded and their productivities were determined by Pandex assays. RGC-derived clones expressing both hVEGF-R1R2 and hVEGF-R1R3 had higher specific productivities and were isolated by screening fewer clones as compared to the highest-expressing hand-picked MSX-resistant colonies. See Table 1.

Table I SPECIFIC PRODUCTIVITY COMPARISON					
Protein	Transient (μg/ml)	Hand-picked CHO K1 Stable Cell Lines		RGC1-derived Stable Cell Lines	
		Sp. Prod. (pg/cell/day)	# clones screened	Sp. Prod. (pg/cell/day)	# clones screened
4SC622	1.1	12	2700	12	6
hVEGF-R1 R2	33	68	190	77	62
hVEGF-R1 R3	27	5	100	22.6	42

Example 4: Cell surface-bound hlgG1 Fc-tagged protein is internalized by RGC1

[0118] hFcyRI is known to induce internalization of its cell surface-bound ligand. To analyze whether RGC1 cells could internalize cell surface-bound 4SC622, 1 μ g/ml 4SC622 was added to RGC1 cells for 1 hour and then the cells were immediately processed for 4SC622 immunostaining with PE-AG184 and flow cytometry analysis. Ninety-three percent of the cells stained positive for cell surface 4SC622. Alternatively, 1 μ g/ml 4SC622 was added to RGC1 cells for 1 hour, then the cells were washed and incubated in culture medium without 4SC622 with PE-AG184 for 18 hours. Flow cytometry analysis following immunostaining for 4SC622 showed that 9% of the cells retained 4SC622 on the cell surface. To further characterize the loss of surface-bound 4SC622, purified 4SC622 protein was added to the media of RGC1 and parental CHO K1 cells, then levels of 4SC622 in the media were measured over time. 4SC622, added to 2 μ g/ml to the culture media in a 10 cm plate, was significantly lower in RGC1

conditioned medium after 3 days incubation as compared to the CHO K1 control. These results show that the concentration of 4SC622 in the culture medium is reduced by the presence of hFcγRI on the cell surface. The results suggest that the depletion of 4SC622 from the media was the result of hFcγRI-4SC622 complex internalization. This internalization of receptor-ligand complexes may facilitate the effective removal of all 4SC622 from non-expressing cells in the presence of blocking IgG during the 18-hour blocking step.

Example 5: Construction of CHO K1 cell lines with inducible hFcγRI expression

[0119] Flow cytometry-based autologous secretion trap (FASTR™) methods that utilize the hFcγRI allow rapid isolation of high expression clones. However, if hFcγRI mediates turnover of Fc-tagged proteins, then the realized production of the secreted protein by engineered hFcγRI expressing cells would be higher if hFcγRI expression could be inhibited during the production period. To this end, a CHO K1 cell line in which the expression of hFcγRI is induced by tetracycline, or the analog doxycycline, was constructed. In this system, CHO K1 cells were first engineered to express the tetracycline repressor protein (TetR) and hFcγRI was placed under transcriptional control of a promoter whose activity was regulated by TetR. Two tandem TetR operators (TetO) were placed immediately downstream of the CMV-MIE promoter/enhancer in pTE084 to generate pTE158. Transcription of hFcγRI from the CMV-MIE promoter in pTE158 was blocked by TetR in the absence of tetracycline or some other suitable inducer. In the presence of inducer, TetR protein was incapable of binding TetO and transcription of hFcγRI occurred.

[0120] CHO K1 cells were transfected with pcDNA6/TR, a plasmid that confers resistance to blasticidin in which expression of TetR originates from the CMV-MIE promoter (Invitrogen; Carlsbad, CA). After two weeks of selection with 2.5 µg/ml blasticidin (Invitrogen), the stable transfectants were pooled. This pool was then transfected with pTE158, a plasmid that confers resistance to G418 in which the expression of hFcγRI is dependent on a CMV-MIE/TetO hybrid promoter. The cells consecutively transfected with pcDNA6/TR and pTE158 were selected with 400 µg/ml G418 and 2.5 µg/ml blasticidin for 12 days then pooled. The pool was induced for two days by the addition of 1 µg/ml doxycycline then stained with FITC-hFc to identify cells that express hFcγRI. The top 5% of cells expressing hFcγRI were collected as a pool, expanded for 6 days in the absence of doxycycline, and were again stained with FITC-hFc for the presence of hFcγRI. Cells that did not stain for hFcγRI were collected and expanded in culture medium containing 1 µg/ml of doxycycline for three days. The pool was then stained for the presence of hFcγRI and were isolated by flow cytometry. Cells that expressed the highest levels of hFcγRI (top 1%) were sorted onto 96 well plates at one cell per well. These cells presumably contained cell that had low non-induced expression levels of FcγR1 and high inducible levels of FcγR1. After expansion, the induction of hFcγRI by doxycycline in 20 clones was confirmed by immunostaining with FITC-hFc and flow cytometry. One clone was chosen for further characterization and was named RGC10.

[0121] In the absence of doxycycline, RGC10 did not express detectable levels of hFcγRI, whereas high levels of hFcγRI were observed in cells that were induced with 1 μg/ml of doxycycline for three days. The mean fluorescence of RGC10 cells increased by more than 1,000 fold after induction by doxycycline.

Example 6: Isolation of 4SC622-producing cell lines from RGC10

[0122] RGC10 cells were transfected with pEE14.1-622, and MSX-resistant cells were pooled after selection with 25 mM MSX for two weeks. Expression of hFcγRI was induced by the addition of 1 μg/ml of doxycycline to the culture medium for three days. One mg/ml rat IgG was added to the culture medium containing doxycycline 18 hours prior to staining with polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment and analysis by flow cytometry. Cells that expressed the highest levels of 4SC622 (top 1%) were sorted into 96 well plates at 1 cell per well. Without induction of hFcγRI expression by doxycycline, staining with polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment fails to detect cell surface bound 4SC622. Sixty clones were expanded in the absence of doxycycline. The specific productivity of the 13 highest producers was determined by Pandex assay. The specific productivity of clone 1C2 was 17.8 pg/cell/day, significantly better than the 12 pg/cell/day observed for the best 4SC622 cell line previously isolated using the unregulated hFcγRI cell line RGC1.

Example 7: Sp2/0 myeloma cells can be engineered to express a cell surface capture protein

[0123] In this example, the Sp2/0-Ag14 myeloma cell line was engineered to stably express hFcγRI to demonstrate that the autologous secretion trap method was applicable to cell lines other than CHO. The gene for hFcγRI was introduced into the myeloma cell by retroviral infection. The plasmid pLXRN (Clontech; Palo Alto, CA), a retroviral DNA vector wherein a gene of interest may be expressed from the upstream Moloney murine sarcoma virus long terminal repeat (MoMuSV LTR) promoter, was used to generate retrovirus encoding the hFcγRI gene. The 1,363 bp Xho I fragment from pTE084, encoding the human FcγRI gene, was cloned into the Xho I site of pLXRN. A plasmid in which hFcγRI cDNA expression was dependent on the MoMuSV LTR was chosen and named pTE255.

[0124] Pantropic retrovirus for the expression of hFcγRI was generated essentially following the manufacturer's guidelines. The packaging cell line GP-293, a HEK 293-based cell line that stably expresses the viral gag and pol proteins (Clontech; Palo Alto, CA), was co-transfected with 10 mg each of pVSV-G and pTE255. The plasmid pVSV-G allows expression of the viral envelope protein VSV-G that confers broad host range upon the infective particles.

[0125] **Construction of Sp2-hFcγRI-4.** The pantropic hFcγRI retrovirus was used to infect 1×10^7 Sp2/0-Ag14 myeloma cells (American Type Culture Collection; Manassas, VA) at a

multiplicity of about 10 infective particles per cell. Three days after infection, cells were stained for 1 hour then washed twice with PBS prior to analysis by flow cytometry. Those cells expressing hFcγRI, as indicated by bound FITC-hFc, were collected as a pool by flow cytometry. The pool was expanded for 13 days then again stained with FITC-hFc and cells expressing hFcγRI were collected as a pool by flow cytometry. These sorted cells were cultured in 10% fetal bovine serum 90% Dulbecco's Modified Eagle's Medium (DMEM) with 4.5 g/l glucose and 4 mM glutamine for 3 weeks, stained with FITC-hFc, and the cells with mean fluorescence in the top 1% of the population were cloned by single cell sorting. After expansion, 24 clones were examined by flow cytometry for expression of hFcγRI, as described above, and one clone, Sp2-hFcγRI-4, was chosen for additional characterization.

[0126] Isolation of Sp2-hFcγRI-4 cells expressing 4SC622 protein. Sp2-hFcγRI-4 cells (1×10^7) were transfected with pTE209, a plasmid that allows constitutive expression of 4SC622 from the CMV-MIE promoter and confers resistance to hygromycin. The transfected cells were placed in medium containing 10% FCS, 90% D-MEM and 400 μg/ml hygromycin for 14 days. Hygromycin-resistant cells were incubated with 1 mg/ml rabbit IgG for eighteen hours prior to staining with polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment. Cells were stained for 1 hour then washed twice with PBS prior to analysis by flow cytometry. Labeled cells were collected as a pool by flow cytometry then cultured for 5 days and sorted as described above. Cells from the expanded pool that bound the most polyclonal FITC-conjugated anti-human IgG(H+L)F(ab')₂ fragment, top 1% population, were then cloned by single cell sorting. Production of 4SC622 from ten clones was analyzed by ELISA and all 10 clones were found to express 4SC622; clone 5H11 produced 4SC622 at 0.5 pg per cell per day. These data showed that clones secreting 4SC622 were efficiently isolated by the autologous secretion trap method from a heterogeneous pool of cells derived from stable transfection of Sp2-hFcγRI-4 cells with pTE209.

[0127] To confirm that 4SC622 was autologously displayed on the surface of myeloma cells expressing both 4SC622 and hFcγRI, clone 5H11 was incubated with 1 mg/ml rabbit IgG for 18 hours then stained with FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment and found to display cell surface 4SC622. Secreted protein was displayed under conditions in which cross-feeding was blocked by rabbit IgG, demonstrating the autologous display of 4SC622. These data indicated that the autologous secretion trap method described above was not limited to CHO cells and may be extended to myeloma and other cell types as well.

Example 8. Protein G chimeric protein can function as a cell surface capture protein

[0128] To demonstrate the application of the autologous secretion trap method to a cell surface capture protein other than hFcγRI, a cell line expressing Protein G was constructed. Protein G, from the *Streptococcus* strain G148, binds to all human and mouse IgG subclasses, and as such has utility for the isolation of recombinant cells expressing antibodies or IgG Fc fusion proteins. To demonstrate that the Protein G IgG Fc binding domain could be used as a

cell surface capture protein capable of binding to all human and mouse IgG subclasses, we constructed a CHO line expressing a chimeric protein comprised of the Fc binding domain of Protein G fused to the hFcγRI transmembrane and intracellular domain. The Fc binding domain of Protein G contains three homologous repeats of 55 amino acids long (Guss et al., (1986) EMBO 5:1567 and Sjöbring et al., (1991) J. Biol. Chem. 266:399) and each repeat is capable of binding one IgG Fc. To improve the expression of this chimeric protein in CHO cells, we constructed a synthetic DNA in which the signal sequence from the mouse ROR1 gene was fused to the Fc binding domain, amino acids 303 to 497 of Protein G (accession # X06173) (SEQ ID NO:1). This synthetic DNA was generated by a combination of oligonucleotide annealing, gap filling, and PCR amplification. The synthetic DNA was then fused, by PCR, to DNA encoding the transmembrane and intracellular domains, amino acids 279 to 374 (SEQ ID NO:2), of hFcγRI (accession M21091). The resultant DNA encoding the Protein G/hFcγRI chimeric protein was cloned into pTE158 downstream of the CMV-MIE promoter, replacing the gene encoding hFcγRI, to yield the plasmid pTE300.

[0129] A CHO K1 cell line adapted to grow in serum-free medium, RGC14, was transfected with pTE300, and after three days 400 µg/ml G418 was added to the culture medium to select for stable integration of pTE300. Two weeks after the start of selection, the cells were stained with FITC-hFc to identify cells that expressed hFcγRI. These cells were analyzed by flow cytometry and cells expressing hFcγRI were collected as a pool. The cells were expanded for 10 days and the population of cells expressing hFcγRI was again isolated by flow cytometry. The cells were again expanded, stained with FITC-hFc, and single cells expressing high levels of the Protein G/hFcγRI chimeric protein were isolated by flow cytometry. Single cells that stained positive for FITC-hFc binding were sorted into medium composed of 10% fetal bovine serum, 90% Ham's F12, and 400 µg/ml G418. After two weeks incubation, 48 clones were examined for binding to bovine IgG present in the culture medium by staining with FITC-conjugated anti-bovine IgG F(ab')₂ fragment (Jackson ImmunoResearch Laboratories, West Grove, PA). One clone, RGC18 that stained positive with this antibody was chosen for further characterization.

[0130] Isolation of expression clones in RGC18: RGC18 cells (6×10^6) were transfected with pTE209 and selected for integration of the plasmid by growth in 400 µg/ml hygromycin for 18 days. Hygromycin-resistant cells were incubated with 1 mg/ml rabbit IgG for eighteen hours prior to staining with polyclonal FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment. Cells were stained for 1 hour then washed twice with PBS prior to analysis by flow cytometry. The most fluorescent cells (top 5%) were isolated by single cell sorting and expanded for 3 weeks. Ten clones were examined for 4SC622 secretion. All clones tested secreted 4SC622 at high level, and the best clone, RGC19, had a specific productivity of 6.4 pg/cell day. This result demonstrated that 4SC622-expressing cells were efficiently isolated from a heterogeneous pool of cells derived from stable transfection of RGC18 with pTE209 by the autologous secretion trap method. Furthermore, these data clearly demonstrated that a fragment of Protein G could be engineered to include a signal sequence and transmembrane domain, and function as a cell surface capture protein.

[0131] To confirm that 4SC622 was autologously displayed on the surface of RGC19 cells expressing both Protein G/hFcγRI chimeric protein and 4SC622, RGC19 was incubated with 1 mg/ml rabbit IgG for 18 hours then stained with FITC-conjugated anti-human IgG (H+L) F(ab')₂ fragment and analyzed by flow cytometry. RGC19 cells were found to possess cell surface 4SC622 under these conditions in which cross-feeding was blocked by rabbit IgG, suggesting autologous display of 4SC622. Rabbit IgG effectively blocked binding of exogenous 4SC622 protein to RGC18 cells, but did not block display of 4SC622 on the cell surface of cells expressing 4SC622. These data demonstrated that the properties of the Protein G/hFcγRI chimeric protein were similar to those of hFcγRI as a cell surface capture protein, and suggested that the autologous secretion trap method can employ other proteins as cell surface capture proteins.

Example 9: Isolation of antibody-producing cells from RGC10

[0132] To demonstrate the utility of the autologous secretion trap method for the isolation of CHO cell lines that express recombinant antibodies we cloned the DNA encoding variable light and variable heavy genes from the KD5 hybridoma. KD5 is a hybridoma that expresses a monoclonal antibody specific for the human Tie-2 receptor.

[0133] The mouse IgG constant region gene sequences were cloned from 500ng of mouse spleen polyA⁺ RNA (Clontech, Palo Alto, CA). Single stranded cDNA was synthesized using SuperScript First-Strand Synthesis System for RT-PCR, primed with 50ng of random hexamers (Invitrogen Life Technologies, Carlsbad, CA). The mouse kappa light constant DNA sequence (accession # Z37499) was amplified from this cDNA by PCR using the primers 5' mCLK1 (Z37499) (5'-CGGGCTGATG CTGCACCAAC TGTATCCATC TTC-3') (SEQ ID NO:3) and 3' mCLK1(Z37499) (5'-ACACTCTCCC CTGTTGAAGC TCTTGACAAT GGG-3') (SEQ ID NO:4). The mouse IgG2a constant region DNA sequence (accession # AJ294738) was also amplified from this cDNA by PCR using the primers 5' mCH2a(AJ294738) (5'-GCCAAAACAA CAGCCCCATC GGTCTATCCA C-3') (SEQ ID NO:5) and 3' mCH2a(AJ294738) (5'-TCATTTACCC GGAGTCCGGG AGAAGCTCTT AGTCG-3') (SEQ ID NO:6). The PCR products were cloned into pCR2.1-TOPO using TOPO TA Cloning kit (Invitrogen Life Technologies, Carlsbad, CA) and the sequence of the constant regions were verified.

[0134] The KD5 variable region genes were amplified by RT-PCR from KD5 hybridoma mRNA and cloned into pCR2.1-TOPO using the heavy and light chain variable region primer mixes from Amersham-Pharmacia Biotech (Piscataway, NJ). The variable heavy chain gene was PCR amplified using the pCR2.1-TOPO cloned variable region as template with the primers 5' BspMI/KD5VH N- term (5'-GAGAGTACCT GCGTCATGCA GATGTGAAAC TGCAGGAGTC TGGCCCT-3') (SEQ ID NO:7) and 3' BspMI/KD5VH C- term (5'-GAGAGACCTG CGTCAGCTGA GGAGACGGTG ACCGTGGT-3') (SEQ ID NO:8), digested with BspMI and ligated to the Bsal-digested IgG2a constant heavy gene PCR fragment amplified with the primers 5' Bsal/CH2a N- term (5'-GAGAGGGTCT CACAGCCAAA ACAACAGCCC CATCG-3')

(SEQ ID NO:9) and 3' BsaI/ CH2a C- term (5'-GAGAGGGTCT CCGGCCGCTC ATTTACCCGG AGTCCGGG AGAA-3') (SEQ ID NO:10). This fragment was then ligated into the BspMI and NotI sites of pRG882. The resulting plasmid, pTE317, was capable of expressing the KD5 recombinant heavy chain gene, fused to the mROR1 signal sequence, from the CMV-MIE promoter. The variable light chain gene was PCR amplified using the pCR2.1-TOPO cloned variable region as template with the primers 5' BsmBI/KD5VL N- term (5'-GAGAGCGTCT CATGCAGACA TCCAGATGAC CCAGTCTCCA-3') (SEQ ID NO:11) and 3' BsmBI/KD5VL C- term (5'-GAGAGCGTCT CACAGCCCGT TTTATTTCCA GCTTGGTCCC-3') (SEQ ID NO:12), digested with BsmBI and ligated to the BsaI-digested kappa constant light gene PCR fragment amplified with the primers 5' BsaI/CLK N- term (5'-GAGAGGGTCT CAGCTGATGC TGCACCAACT GTATCC-3') (SEQ ID NO:13) and 3' BsaI/CLK C- term (5'-GAGAGGGTCT CAGGCCGCTC AACACTCTCC CCTGTTGAAG CTCTTGAC-3') (SEQ ID NO: 14). This fragment was then ligated into the BspMI and NotI sites of pRG882. The resulting plasmid, pTE316, was capable of expressing the KD5 recombinant light chain gene, fused to the mROR1 signal sequence, from the CMV-MIE promoter.

[0135] The 1450 bp EcoRI-NotI fragment from pTE317, encoding the KD5 heavy chain gene, was cloned into the EcoRI and NotI sites of pRG980, a vector that confers resistance to hygromycin and allows expression of recombinant genes for the UbC promoter, to yield plasmid pTE322. Similarly, the 750 bp EcoRI-NotI fragment from pTE316, encoding the KD5 light chain gene, was cloned into the EcoRI and NotI sites of pRG985, a vector that confers resistance to puromycin and allows expression of recombinant genes for the UbC promoter, to yield plasmid pTE324. RGC10 cells (5×10^6) were transfected with 3 μ g pTE322 and 3 μ g pTE324 and selected for integration of the plasmids by growth in F12 medium supplemented with 10% fetal calf serum with 20 μ g puromycin and 400 μ g/ml hygromycin for 14 days. Expression of hFc γ R1 was induced by the addition of 1 μ g/ml of doxycycline to the culture medium for three days. Double-resistant cells were incubated with 1 mg/ml rabbit IgG for eighteen hours prior to staining with goat polyclonal FITC-conjugated anti-mouse IgG (Fc γ) F (ab')₂ fragment (Jackson ImmunoResearch Laboratories, West Grove, PA). Cells were stained for 1 hour then washed twice with PBS prior to analysis by flow cytometry. The most fluorescent cells (top 5%) were isolated as a pool and expanded for 10 days, after which the protocol was repeated but the top 1% most fluorescent cells were isolated as a pool. This pool was expanded for 10 days then the top 0.1% most fluorescent cells were isolated as single cells into 96-well plates. Clones were analyzed by ELISA for expression of antibody and seven clones were chosen from 53 clones analyzed. The average specific productivity of these clones was 35 pg/cell/day and the best clone expressed the recombinant KD5 monoclonal antibody at 54 pg/cell/day.

Example 10: FASTR™ screens unaffected by CSCP expression level

[0136] To demonstrate that the expression level of the CSCP does not significantly affect the ability to isolate cells expressing an associated sPOI, FASTR™ screens for the same sPOI in

two different host cell lines that each express the same CSCP but at either a high level or a low level were compared.

[0137] The FASTR™ host cell line RGC10 was selected for high-level expression of hFcγRI protein by stable integration of pTE158 and was found to contain 40 hFcγRI integrated gene copies. A new cell line, RS527, that expressed hFcγRI protein at a lower level, was generated from CHO K1 after stable transfection and selection for single copy gene integration. RS527 cells expressed significantly less hFcγRI protein than RGC10 cells as determined by Western blot analysis of whole cell lysates of the FASTR™ cell lines.

[0138] Briefly, RGC10 and RS527 cells were transfected with pTE462, a plasmid capable of expressing a secreted hFc-fusion protein Rc1-hFc and conferring resistance to hygromycin. The transfected cultures were selected with hygromycin for two weeks. The hygromycin-resistant cells were induced with 1 µg/ml doxycycline (Dox) and blocked with rabbit IgG overnight, following the FASTR™ method described herein. The next day, the RGC10/pTE462 and RS527/pTE462 cultures were stained by a FITC-conjugated antibody specific for hFc and then analyzed by flow cytometry. Three cell bins R4, R5, and R6 marking cells with low, medium, and high fluorescence respectively were sorted from each host line and expanded in tissue culture.

[0139] To compare Rc1-hFc protein production level from the six cell bins, six cultures were set up using equal number of cells for each bin. Three days later, conditioned media were collected. The Rc1-hFc protein titers in the conditioned media were determined by ELISA and were plotted against mean fluorescence of the respective cell bins. For both RGC10 and RS527 host lines, there was a similar correlation between mean fluorescence (amount of Rc1-hFc displayed on the cell surface) and sPOI protein production levels of the isolated cell pools. Most significantly, the sPOI titers in the two high fluorescence R6 bins derived from RGC10 and RS527 were similar. These data demonstrate that the expression level of the CSCP in a FASTR™ host cell line does not significantly affect the use of that host to isolate transfected cells based on expression level of a sPOI.

Example 11: Tie2 receptor as a cell surface capture protein

[0140] Cell surface capture proteins (CSCP's) other than FcγR1 can be used in the methods described herein. In this example, the Tie2 receptor functions as a CSCP and is used to isolate cells expressing a Tie-specific ScFv_{C1b}-Fc fusion protein made from the C1b monoclonal antibody that specifically binds the extracellular domain of Tie2 receptor. Although the CSCP for ScFv_{C1b}-Fc can be hFcγRI, this example demonstrates that Tie2 can also be used as the CSCP for ScFv_{C1b}-Fc.

[0141] To construct an inducible Tie2 CSCP cell line, CHO K1 was first stably transfected with the TetR plasmid pcDNA6/TR. The blasticidin-resistant cell pool was then stably transfected

with pTE259, a plasmid that allows inducible expression of a protein comprised of the extracellular domain and transmembrane domain of Tie2. Inducible cell clones were isolated by flow cytometry after staining with an antibody specific for Tie2. The RGC54 clone was chosen to study the feasibility of FASTR™ for the expression of ScFv_{C1b}-Fc.

[0142] RGC54 cells were stably transfected with pTE988, a plasmid capable of expressing the secreted hFc-fusion protein ScFv_{C1b}-Fc and conferring resistance to hygromycin. The transfected culture was selected with hygromycin for two weeks. The hygromycin-resistant cells were induced with Dox and blocked with 1 mg/ml of purified C1b mAb. The C1b monoclonal antibody was the source of the variable regions in ScFv_{C1b}-Fc. The next day, the cell pool was stained by a FITC-conjugated antibody specific for hFc and then analyzed by flow cytometry. Three cell bins R6, R7, and R8 marking cells with high, medium, and low fluorescence respectively were sorted and expanded in tissue culture. Three cultures were set up using an equal number of cells for each bin to determine ScFv_{C1b}-Fc protein production as determined by ELISA. A correlation existed between mean fluorescence (amount of ScFv_{C1b}-Fc binding to Tie2 on the cell surface) and ScFv_{C1b}-Fc protein production levels of the isolated cell pools.

[0143] These data show that CSCP other than hFcγRI can serve as a CSCP, and also suggest that any receptor may be converted into a CSCP by removal of its cytoplasmic domain. These data also demonstrate that an antigen can be made into a CSCP and used for FASTR™ screening cells expressing an antigen-specific antibody-related molecule.

Example 12: Effective FASTR™ screens with CSCP:sPOI pairs having low affinity

[0144] Angiopoetin-1 is a ligand for the Tie2 receptor. A chimeric protein comprising angiopoetin-1 receptor binding domain and hFc (FD1-hFc) binds to Tie2 with an affinity constant of 174 nM as determined by BIAcore™. FD1-hFc and Tie2 were chosen as sPOI and CSCP, respectively, to determine if a minimum affinity between CSCP and sPOI is required for FASTR™ screens.

[0145] In cell decoration experiments, exogenously added FD1-hFc bound specifically to RGC54 cells through Tie2. To determine if the affinity between Tie2 and FD1-hFc is sufficient to allow FASTR™ screening, RGC54 cells were stably transfected with pTE942, a plasmid capable of expressing the secreted hFc-fusion protein FD1-hFc and conferring resistance to hygromycin. The transfected culture was selected with hygromycin for two weeks. The hygromycin-resistant cells were induced with Dox and blocked with 1 mg/ml of purified FD1-mFc comprising mouse IgG1 Fc. The next day, the cell pool was stained by a FITC-conjugated antibody specific for hFc and then analyzed by flow cytometry. Three cell bins R6, R7, and R8 marking cells with high, medium, and low fluorescence, respectively, were collected. Cultures were set up using equal number of cells for each bin to determine FD1-hFc protein production levels in the conditioned media as determined by ELISA. There was a correlation between

mean fluorescence (FD1-Fc binding to cell surface-bound Tie2) and FD1-hFc protein production levels of the isolated cell pools. The bin with the highest fluorescence produced the most FD1-hFc.

[0146] These data demonstrate that a CSCP:sPOI pair with low affinity (174 nM KD) can be used for effective FASTR™ screens. Importantly, the dissociation $t_{1/2}$ for FD1-Fc: Tie2 binding is less than 2 minutes, suggesting that any CSCP:sPOI pair with a measurable affinity can work in FASTR™ screens. In addition, this experiment also shows that a non-FcγRI receptor may be used as the CSCP to isolate cells expressing its ligand.

Example 12: Fusing a transmembrane domain onto an ScFv makes a functional CSCP

[0147] An CSCP can be any cell surface-bound protein that has a measurable affinity to the sPOI. To demonstrate this, a totally synthetic CSCP was constructed by fusing the transmembrane domain from the PDGF receptor to an ScFv containing the variable regions from the murine kappa chain-specific monoclonal antibody HB58. A FASTR™ host was constructed that expresses this chimeric protein (ScFv_{HB58}-TM_{PDGFR}) and was used to isolate cells expressing the angiopoietin-2 FD domain-specific P12 antibody.

[0148] The RS655 cell line, derived from CHO K1, constitutively expresses ScFv_{HB58}-TM_{PDGFR}. Cells expressing ScFv_{HB58}-TM_{PDGFR} can be stained by sequential incubation with P12 mAb, FD2-hFc, and FITC-conjugated anti-hIgG - P12 captured on the cell surface by the HB58 ScFv was detected by its affinity for FD2, which in turn was detected by recognition of the hFc tag. RS656 cells were derived from RS655 cells after stable transfection with a plasmid encoding the gene for eYFP. Nearly 100% of RS656 cells were eYFP-positive, and most (76%) maintained expression of ScFv_{HB58}-TM_{PDGFR} as detected by binding to FD2-hFc.

[0149] RS655 cells were stably transfected with pTE693, a plasmid capable of expressing the heavy and light chains of the P12 antibody, and conferring resistance to puromycin. The transfected culture was selected with puromycin for two weeks to yield a pool of cells that were heterogeneous with regard to P12 mAb expression (RS655/pTE693).

[0150] To determine if ScFv_{HB58}-TM_{PDGFR} could function as a CSCP and facilitate isolation of antibody-producing cells from non-producers, equal numbers of RS656 cells and RS655/pTE693 cells were mixed and co-cultured. When P12 expressed from RS655/pTE693 cells was allowed to diffuse and bind to ScFv_{HB58} on the surface of RS656 cells a large population of yellow cells were also positive for binding FD2-hFc. However, if the ScFv_{HB58} on the surface of RS656 was bound with excess murine IgG, then only non-yellow cells were positive for binding FD2-hFc, demonstrating that expressing cells were effectively separated from non-expressing cells.

[0151] These data demonstrate that an ScFv can be made into a functional CSCP by targeting

it to the cell membrane. The data also show that FASTR™ allows cells expressing a secreted antibody to be detected with the antibody's antigen.

Example 13: A protein of interest comprising a T cell receptor variable region

[0152] A flow cytometry-based autologous secretion trap (FASTR™) method for isolating high expression clones of a cell line that expresses a protein of interest that is a TCR-Fc is prepared in a manner analogous to preparing a cell line that expresses an antibody of interest. High expression clones are identified by screening cells that display on their surface the TCR-Fc of interest bound to hFcγR.

[0153] In these examples, the CHO K1 cell line RGC10, comprising an inducible FcγR1 as a cell surface capture molecule, is employed. RGC10 is made to express recombinant TCR-Fc's by cloning TCR variable regions, in frame, to a human Fc region either directly in frame or with a linker sequence between the TCR variable regions and the human Fc region.

[0154] To make a protein of interest that is a dimer comprising an Fc-linked TCR α variable domain and an Fc-linked TCR β variable domain, RGC10 is transfected with two vectors: a first vector capable of expressing a TCR α variable domain fusion protein with a human Fc sequence, and a second vector capable of expressing a TCR β domain fusion protein with the same human Fc sequence. Each vector includes leader sequence (e.g., a secretion signal sequence) 5' with respect to the TCR variable region, and a selectable marker that is a drug resistance gene. Following each vector transfection, cells containing the vector are selected by an appropriate drug selection. The selection results in an RGC10 cell line having both the first and the second vectors. Cells expressing proteins of interest can be detected by one or more of an antibody to the β variable domain, an antibody to the α variable domain, and an antibody to the Fc domain.

[0155] To make a protein of interest that is a dimer comprising both an α and a β TCR variable domain fused to an Fc, RGC10 is transfected with a single vector encoding a protein of interest that is constructed as follows: a leader sequence (e.g., a secretion signal sequence), followed by a TCR variable β domain fused to a linker, where the linker is, in turn, fused to a TCR variable α domain, which in turn is fused to an Fc sequence. Alternatively, the single vector can be constructed as follows: a leader sequence (e.g., a secretion signal sequence), followed by a TCR variable α domain fused to a linker, where the linker is, in turn, fused to a TCR variable β domain, which in turn is fused to an Fc sequence. Cells expressing proteins of interest can be detected by one or more of an antibody to the β variable domain, an antibody to the α variable domain, and an antibody to the Fc domain.

[0156] To make proteins of interest, as above, which also comprise a TCR α and/or TCR β constant domain, the TCR variable domain (α or β) is fused to a TCR constant domain (e.g., TCR variable domain α is fused to TCR constant domain α, and TCR variable domain β is fused to TCR constant domain β), and the TCR variable + constant domain is fused directly or

through a linker to the Fc domain. Cells expressing proteins of interest can be detected by one or more of an antibody to the β variable domain, an antibody to the α variable domain, and an antibody to the Fc domain.

[0157] Cells expressing desired amounts of the TCR-Fc are isolated using the same procedure as used in isolating 4SC622-producing cell lines described herein, using one or more of an antibody to the α variable domain, an antibody to the β variable domain, an antibody to the α constant domain, and antibody to the β constant domain, and an antibody to the Fc domain. Cells expressing the highest levels of the TCR-Fc are selected as TCR-Fc-producing cell lines.

Example 14: ScFv-based CSCP for the isolation of multiple IgG isotypes and bispecific antibodies

[0158] Genetically modified mice, whose immunoglobulin heavy chain VDJ region and immunoglobulin kappa chain VJ region of their genomes were replaced with the human orthologs (*i.e.*, Velocimmune® mice; see US Pat. No. 7,105,348), were immunized with either an Fc fragment of a human IgG4 protein (hFc, or simply Fc; SEQ ID NO: 26), or a human Δ AdpFc polypeptide containing the dipeptide mutation (H95R, Y96F by IMGT; also known as Fc*; SEQ ID NO: 42). Monoclonal antibodies were obtained from the mice and screened for their ability to bind Fc, Fc* or antibodies comprising Fc and/or Fc*. Three antibodies that were capable of binding Fc (Ab1, Ab2, Ab3) and three that were capable of binding Fc* (Ab4, Ab5, Ab6) were tested for their ability to bind molecules having one of the following formats: Fc/Fc, Fc/Fc* (which can be a bispecific antibody), and Fc*/Fc*.

[0159] Measurements to determine binding affinities and kinetic constants were made on a Biacore 2000 instrument. Antibodies (each of Ab1-Ab8) were captured onto an anti-mouse-Fc sensor surface (Mab capture format), and human Fc (SEQ ID NO 26) homodimers, human Fc* homodimers (SEQ ID NO:42), or Fc/Fc* heterodimers were injected over the surface. Kinetic association (k_a) and dissociation (k_d) rate constants were determined by processing and fitting the data to a 1:1 binding model using Scrubber 2.0 curve fitting software. Binding dissociation equilibrium constants (K_D) and dissociative half-lives ($t_{1/2}$) were calculated from the kinetic rate constants as: K_D (M) = k_d / k_a ; and $t_{1/2}$ (min) = $(\ln 2)/(60 * k_d)$. As shown in Table 2 antibodies were of 3 distinct categories: Fc specific, Fc* specific, and those showing no discrimination between Fc and Fc* (non-specific). The Fc specific antibodies were dependent on amino acids His 95 and/or Tyr 96, since these antibodies do not bind human Fc* with its dipeptide mutation (H95R, Y96F). In contrast the Fc* specific antibodies were dependent on Arg 95 and/or Phe 96, since these antibodies do not bind wild type human Fc.

Example 15: Cell lines producing Ab2 and Ab2-derived ScFv-Fc γ R fusion protein

[0160] The heavy chain and the light chain of the Fc-specific Ab2 were sequenced. To manufacture the recombinant Ab2 antibody, an expression vector plasmid was constructed that encodes the heavy chain and an expression vector plasmid was constructed that encodes the light chain. Both vectors enable expression and secretion of the respective subunits in a CHO cell. To express the antibody, both plasmids were transfected into a CHO-K1 cell and stable transformants were isolated. Expression of the antibody chains was driven by the constitutive CMV promoter.

Table 2: Affinity of Antibodies - Surface Plasmon Resonance Studies

Antibody	POI-Target	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	KD (M)	$t_{1/2}$ (min)	Specificity
Ab1	Fc/Fc	1.07E+05	3.79E-04	3.54E-09	30	Fc
	Fc/Fc*	8.16E+04	3.01 E-04	3.69E-09	38	
	Fc*/Fc*	NB	NB	NB	NB	
Ab2	Fc/Fc	7.86E+04	3.50E-05	4.45E-10	330	Fc
	Fc/Fc*	5.45E+04	1.00-06	1.84E-11	11550	
	Fc*/Fc*	NB	NB	NB	NB	
Ab3	Fc/Fc	1.77E+05	4.08E-02	2.30E-07	0.3	Fc
	Fc/Fc*	4.51 E+04	2.60E-02	5.77E-07	0.4	
	Fc*/Fc*	NB	NB	NB	NB	
Ab4	Fc/Fc	NB	NB	NB	NB	Fc*
	Fc/Fc*	6.00E+03	1.00E-06	2.00E-10	11550	
	Fc*/Fc*	2.22E+04	9.56E-06	4.50E-10	1209	
Ab5	Fc/Fc	NB	NB	NB	NB	Fc*
	Fc/Fc*	3.11 E+05	1.00E-06	3.21 E-12	11550	
	Fc*/Fc*	5.57E+05	1.00E-06	1.79E-12	11550	
Ab6	Fc/Fc	NB	NB	NB	NB	Fc*
	Fc/Fc*	4.48E+05	7.43E-04	1.66E-09	16	
	Fc*/Fc*	8.73E+05	5.93E-04	6.79E-10	19	
Ab7	Fc/Fc	6.02E+05	2.42E-04	4.02E-10	48	Non-specific
	Fc/Fc*	4.90E+05	2.15E-04	4.39E-10	54	
	Fc*/Fc*	4.46E+05	3.20E-02	7.18E-08	0.4	
Ab8	Fc/Fc	2.59E+05	4.88E-04	1.88E-09	24	Non-specific
	Fc/Fc*	1.88E+05	4.02E-04	2.14E-09	29	
	Fc*/Fc*	4.10E+04	3.90E-02	9.60E-07	0.3	

[0161] The heavy chain and light chain sequences were used to develop an anti-Fc ScFv surface capture molecule. To manufacture the nucleic acid encoding the Ab2-derived anti-Fc

ScFv-FcγR surface capture molecule, the Ab2 immunoglobulin heavy chain variable domain (SEQ ID NO:15) and the Ab2 immunoglobulin light chain variable domain (SEQ ID NO:16) amino acid sequences were reverse translated and codon optimized for CHO cell expression. Likewise, the C-terminal portion of human FcγRI was codon optimized for CHO cell expression. The codon optimized nucleotide sequences were amplified via polymerase chain reaction and ligated to form a contiguous nucleic acid sequence (SEQ ID NO:20) that encodes the ScFv-FcγR fusion protein of SEQ ID NO:19.

[0162] The nucleic acid encoding the ScFv-FcγR-TM-cyto fusion protein was inserted into an expression vector using standard PCR and restriction endonuclease cloning techniques. The resultant circular plasmid, exemplified in SEQ ID NO:23, comprises a beta-lactamase-encoding nucleic acid sequence, and two operons. The first operon comprises a nucleic acid sequence encoding yellow fluorescence protein (YFP), a variant of green fluorescent protein, in frame with a neomycin resistance marker, driven by an SV40 promoter (e.g., SEQ ID NO:24). The second operon, which is the "business-end" of the vector for the purposes of this aspect of the invention, comprises a nucleic acid sequence encoding the codon-optimized ScFv-FcγR fusion protein, driven by an hCMV-IE promoter and hCMV intron (e.g., SEQ ID NO:25).

[0163] CHO-K1 cells were transfected with the plasmid of SEQ ID NO:23. Stable integrants, which have integrated the linear construct of SEQ ID NO:22 into their genomes, were isolated.

[0164] The circular plasmid contains two Lox sites flanking the first operon and the second operon, to allow for the integration of those operons as a linear construct into the genome of the host cell. The linear construct spanning from the first Lox site to the second Lox site is exemplified in SEQ ID NO:22 and comprises from 5-prime to 3-prime: SV40 promoter, nucleic acid encoding neomycin-resistance, IRES, nucleic acid encoding eYFP, SV40 polyadenylation sequence, hCMV-IE promoter, hCMV intron, Tet-operator sequence (for controlled expression of the ScFv-FcγR-TM-cyto fusion protein), nucleic acid encoding mROR signal sequence, nucleic acid encoding Ab2 ScFv, nucleic acid encoding the FcγR transmembrane and cytoplasmic portion (SEQ ID NO: 21), and SV40 polyadenylation sequence.

Example 16: ScFv-FcγR-TM-cyto Surface Capture Targets

[0165] CHO-K1 cells containing the integrated sequence of SEQ ID NO:22 were transfected with plasmids that encode antibodies of various subtypes, e.g., IgG1, IgG2, IgG4, an IgG4 bispecific antibody containing one CH3 domain with the 95R/435R - 96F/436F dual substitution while the other CH3 domain is wild-type (IgG4 Fc/Fc*), and an IgG1 bispecific antibody of the IgG1 Fc/Fc* format. The cells were treated with doxycycline to induce production of the capture molecule along with the antibody. After co-expression of the antibody and capture molecule, the cells in some cases were treated with hFc blocking protein, and detection molecule (FITC-labeled anti-hFab). Table 3 summarizes the results, and generally shows that the ScFv-FcγR surface capture fusion protein binds IgG4, IgG2, and IgG1 molecules, while the wildtype FcγR surface capture molecule binds IgG1, but not IgG4 or IgG2.

Table 3: Blocking Molecule Competition Assays

Antibody	Arbitrary FITC Units (with or without hFc blocking molecule) - Mode					hFc displacement?
	No hFc	hFc (1 hr)	hFc (2 hr)	hFc (20 hr)	No coat	
		Capture molecule = ScFv-FcγR-TM-cyto				
		Detection molecule = FITC-anti-hFab				
IgG1 mAb-3	250	120	80	20	10	Yes
IgG4 mAb-4	250	100	55	20	10	Yes
IgG4 mAb-5	250	70	40	20	10	Yes
IgG2 mAb-6	200 ¹	ND	ND	ND	12 ²	Yes
		Capture molecule = hFcγR				
		Detection molecule = FITC-anti-hFab				
IgG1 mAb-3	300	80	30	9	3.5	Yes
IgG4 mAb-4	100	2	2	2	2	No
IgG4 mAb-5	35	5	5	5	5	No
¹ + Dox ² -Dox						

Example 17: Cell Lines Producing Ab6 and Ab6-derived ScFv*-FcγR-TM-cyto

[0166] The heavy chain and the light chain of the Fc*-specific Ab6 were sequenced. The amino acid sequence of the light chain was determined to be SEQ ID NO:41. The amino acid sequence of the heavy chain was determined to be SEQ ID NO:40. To manufacture the recombinant Ab6 antibody, an expression vector plasmid was constructed that encodes the heavy chain and an expression vector plasmid was constructed that encodes the light chain. To express the antibody, both plasmids were transfected into a CHO-K1 cell, stable transformants were isolated, and expression was driven by the constitutive CMV promoter.

[0167] To manufacture the nucleic acid encoding the Ab6-derived anti-Fc*-specific ScFv*-FcγR surface capture molecule, the immunoglobulin heavy chain variable domain of the Ab6 antibody (SEQ ID NO:38) and the immunoglobulin light chain variable domain of Ab6 (SEQ ID NO:39) amino acid sequences were reverse translated and codon optimized for CHO cell

expression. Likewise, the C-terminal portion of human FcγRI (SEQ ID NO: 21) was codon optimized for CHO cell expression. The codon optimized nucleotide sequences were amplified via polymerase chain reaction and ligated to form a contiguous nucleic acid sequence (SEQ ID NO:45) that encodes the anti-Fc* ScFv*-FcγR fusion protein (SEQ ID NO:43).

[0168] The nucleic acid encoding the ScFv*-FcγR-TM-cyto fusion protein was inserted into an expression vector using standard PCR and restriction endonuclease cloning techniques. The resultant circular plasmid, exemplified in SEQ ID NO:44, comprises a beta-lactamase-encoding nucleic acid sequence, and two operons. The first operon comprises a nucleic acid sequence encoding yellow fluorescence protein (YFP), a variant of green fluorescent protein, in frame with a neomycin resistance marker, driven by an SV40 promoter (e.g., SEQ ID NO:46). The second operon, which is the "business-end" of the vector for the purposes of this aspect of the invention, comprises a nucleic acid sequence encoding the codon-optimized anti-Fc* ScFv*-FcγR fusion protein, driven by an hCMV-IE promoter and hCMV intron (e.g., SEQ ID NO:47).

[0169] CHO-K1 cells were transfected with the plasmid of SEQ ID NO:44. Stable integrants, which have integrated the linear construct of SEQ ID NO:48, were isolated.

[0170] The circular plasmid contains two Lox sites flanking the first operon and the second operon, to allow for the integration of those operons as a linear construct into the genome of the host cell. The linear construct spanning from the first Lox site to the second Lox site is exemplified in SEQ ID NO:48 and comprises from 5-prime to 3-prime: SV40 promoter, nucleic acid encoding neomycin-resistance, IRES, nucleic acid encoding eYFP, SV40 polyadenylation sequence, hCMV-IE promoter, hCMV intron, Tet-operator sequence (for controlled expression of the anti-Fc* ScFv*-FcγR fusion protein), nucleic acid encoding mROR signal sequence, nucleic acid encoding the Ab6-derived anti-Fc*-specific ScFv* nucleic acid encoding the FcγR transmembrane and cytoplasmic domain polypeptide (SEQ ID NO: 21), and SV40 polyadenylation sequence.

Example 18: Sorting Bispecific Antibodies

Anti-Fc capture & anti-Fc* detection

[0171] The Ab2-derived anti-Fc-specific ScFv-FcγR surface capture system was tested for its ability to detect and enrich for cells that produce bispecific antibodies. To assess the ability to detect bispecific antibodies, which harbor the 95R/435R-96F/436F substitution in one of the CH3 domains (designated Fc*), various antibodies were expressed in the Ab2-derived anti-Fc-specific ScFv-FcγR surface capture cell line, using hFc as the blocking molecule, and a FITC-labeled Ab6 anti-Fc* antibody (e.g., mAb with HC of SEQ ID NO:40, and LC of SEQ ID NO:41) as the detection molecule. The Ab2-derived anti-Fc-specific ScFv-FcγR surface capture cell line was able to detect and distinguish the bispecific antibody (Fc/Fc*) over any Fc*/Fc* or

Fc/Fc monospecific antibodies using the Fc*-specific Ab6 as the detection molecule (Table 4). The wildtype FcγR surface capture cell line was not able to distinguish between the Fc/Fc*, Fc*/Fc*, and Fc/Fc IgG4 species, since FcγR is unable to bind, or binds at very low affinity to IgG4.

Anti-Fc* capture & anti-Fc detection

[0172] Conversely, the Ab6-derived anti-Fc*-specific ScFv*-FcγR surface capture system was tested for its ability to detect and enrich for cells that produce bispecific antibodies. To assess the ability to detect bispecific antibodies, which harbor the 95R/435R-96F/436F substitution in one of the CH3 domains (designated Fc*), various antibodies were expressed in the Ab6-derived anti-Fc*-specific ScFv*-FcγR surface capture cell line, using hFc as the blocking molecule, and an Alexa 488-labeled Ab2 anti-Fc antibody, which recognizes non-substituted CH3, as the detection molecule. The Ab6-derived anti-Fc*-specific ScFv*-FcγR surface capture cell line was able to detect and distinguish the bispecific antibody (Fc/Fc*) over the Fc*/Fc* or Fc/Fc monospecific antibodies using the Fc-specific Ab2 as the detection molecule (Table 4). The FcγR surface capture cell line was not able to distinguish between the Fc/Fc*, Fc*/Fc*, and Fc/Fc IgG4 species.

Table 4: Detection of Bispecific Antibody - Mean Fluorescence Intensity (MFI)

1 CSCP	2 DM	IgG1			IgG4			Fc/Fc*Specificity
		Fc/Fc*	Fc*/Fc*	Fc/Fc	Fc/Fc*	Fc*/Fc*	Fc/Fc	
FcγR	Ab2	500	ND	350	200	200	200	NO
	Ab6	200	200	200	ND	ND	ND	NO
	Anti-hFc	1800	ND	1000	ND	ND	ND	NO
ScFv-FcγR	Ab6	500	15	15	500	15	15	YES
	Anti-hFc	ND	ND	ND	ND	ND	ND	ND
ScFv*-FcγR	Ab2	150	10	10	ND	ND	ND	YES
	Anti-hFc	200	ND	10	ND	ND	ND	YES

1Cell surface capture protein 2Detection molecule

Example 19: Enrichment of Fc/Fc* Bispecific Antibodies

[0173] To assess the ability of the (Ab2-derived) ScFv-FcγR CSCP/ (Ab6) anti-Fc* DM and the (Ab6-derived) ScFv*-FcγR CSCP/ (Ab2) anti-Fc DM systems to sort and enrich bispecific antibodies, cell lines co-expressing an Fc/Fc* IgG4 monoclonal antibody (IgG4-mAb-2) and the

anti-Fc ScFv-FcγR fusion protein, using hFc as the blocking molecule and the FITC-labeled anti-Fc* (Ab6) antibody as the detection molecule, were subjected to serial fluorescence activated cell sorting and pooling to enrich for production of the Fc/Fc* species. Cells yielding Fc/Fc* from the fifth and sixth series pools were analyzed for total antibody titer and titers of each antibody format: Fc/Fc*, Fc/Fc, and Fc*/Fc*. Since the cells encode both a heavy chain encoding the non-substituted CH3 domain ("Fc", *i.e.*, comprising a histidine at IMGT position 95 and a tyrosine at IMGT position 96) and a heavy chain encoding the substituted CH3 domain ("Fc*", *i.e.*, comprising an arginine at IMGT position 95 and a phenylalanine at IMGT position 96), by purely mathematical Punnett square analysis, the cell is theoretically expected to produce 25% Fc/Fc, 50% Fc/Fc*, and 25% Fc*/Fc*. Biologically, however, one might expect (pre-enrichment) most of the antibody produced to be Fc/Fc.

[0174] As shown in Table 5, cells selected, pooled, and enriched for bispecific antibody production produced as much as 49% Fc/Fc* species, with titers of Fc/Fc* bispecific antibodies of at least about 3.2 g/L.

Table 5: Enrichment of Fc/Fc* bispecific antibody IgG4-mAb-2

pool	Cell line	Fc/Fc*		Fc/Fc		Fc*/Fc*	
		Titer (g/L)	%	Titer (g/L)	%	Titer (g/L)	%
5	1	1.2	28	2.2	50	0.99	23
	2	1.9	49	1.3	32	0.73	19
	3	1.5	47	1.2	40	0.40	13
	4	1.6	37	1.3	31	1.3	32
	5	1.5	48	1.1	35	0.58	18
	6	1.8	47	1.3	33	0.75	20
6	7	2.6	44	2.0	34	1.3	23
	8	3.2	42	2.4	31	2.0	27
	9	2.1	45	1.5	33	1.0	22
	10	2.8	43	2.0	31	1.7	28
	11	2.3	44	1.6	31	1.3	24

SEQUENCE LISTING

[0175]

<110> Regeneron Pharmaceuticals, Inc.

<120> RECOMBINANT CELL SURFACE CAPTURE PROTEINS

<130> 8600-WO

<150> US 61/726,040

<151> 2012-11-14

<160> 51

<170> PatentIn version 3.5

<210> 1

<211> 195

<212> PRT

<213> Streptococcus

<400> 1

Thr Tyr Lys Leu Ile Leu Asn Gly Lys Thr Leu Lys Gly Glu Thr Thr
 1 5 10 15

Thr Glu Ala Val Asp Ala Ala Thr Ala Glu Lys Val Phe Lys Gln Tyr
 20 25 30

Ala Asn Asp Asn Gly Val Asp Gly Glu Trp Thr Tyr Asp Asp Ala Thr
 35 40 45

Lys Thr Phe Thr Val Thr Glu Lys Pro Glu Val Ile Asp Ala Ser Glu
 50 55 60

Leu Thr Pro Ala Val Thr Thr Tyr Lys Leu Val Ile Asn Gly Lys Thr
 65 70 75 80

Leu Lys Gly Glu Thr Thr Thr Glu Ala Val Asp Ala Ala Thr Ala Glu
 85 90 95

Lys Val Phe Lys Gln Tyr Ala Asn Asp Asn Gly Val Asp Gly Glu Trp
 100 105 110

Thr Tyr Asp Asp Ala Thr Lys Thr Phe Thr Val Thr Glu Lys Pro Glu
 115 120 125

Val Ile Asp Ala Ser Glu Leu Thr Pro Ala Val Thr Thr Tyr Lys Leu
 130 135 140

Val Ile Asn Gly Lys Thr Leu Lys Gly Glu Thr Thr Thr Lys Ala Val

145 150 155 160

Asp Ala Glu Thr Ala Glu Lys Ala Phe Lys Gln Tyr Ala Asn Asp Asn
 165 170 175

Gly Val Asp Gly Val Trp Thr Tyr Asp Asp Ala Thr Lys Thr Phe Thr
 180 185 190

Val Thr Glu
 195

<210> 2
 <211> 96
 <212> PRT
 <213> Homo sapiens

<400> 2
 Gln Val Leu Gly Leu Gln Leu Pro Thr Pro Val Trp Phe His Val Leu
 1 5 10 15
 Phe Tyr Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val Leu Trp
 20 25 30
 Val Thr Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp Leu Glu
 35 40 45
 Ile Ser Leu Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser Leu Gln
 50 55 60
 Glu Asp Arg His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln Lys Glu
 65 70 75 80
 Glu Gln Leu Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly Ala Thr
 85 90 95

<210> 3
 <211> 33
 <212> DNA
 <213> artificial sequence

<220>
 <223> synthetic

<400> 3
 cgggctgatg ctgcaccaac tgtatccatc ttc

<210> 4
 <211> 33
 <212> DNA
 <213> artificial sequence

<220>
 <223> synthetic

<400> 4
 acactctccc ctgtgaagc tcttgacaat ggg 33

<210> 5
 <211> 31
 <212> DNA
 <213> artificial sequence

<220>

<223> synthetic

<400> 5

gccaaaacaa cagcccatc ggtctatcca c 31

<210> 6

<211> 35

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 6

tcatttaccg ggagtcggg agaagctctt agtcg 35

<210> 7

<211> 47

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 7

gagagtacct gcgcatgca gatgtgaaac tgcaggagtc tggccct 47

<210> 8

<211> 38

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 8

gagagacctg cgtcagctga ggagacgggtg accgtggt 38

<210> 9

<211> 35

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 9

gagaggggtct cacagccaaa acaacagccc catcg 35

<210> 10

<211> 42

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 10

gagaggggtct ccggccgctc atttaccgag agtccgggag aa 42

<210> 11

<211> 40

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 11

gagagcgtct catgcagaca tccagatgac ccagtctcca 40

<210> 12

<211> 40

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 12

gagagcgtct cacagcccgt ttatttcca gcttgggtccc 40

<210> 13

<211> 36

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 13

gagaggggtct cagctgatgc tgcaccaact gtatcc 36

<210> 14

<211> 48

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 14

gagagggtct caggccgctc aacactctcc cctgttgaag ctcttgac 48

<210> 15

<211> 113

<212> PRT

<213> Homo sapiens

<400> 15

Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Lys Pro Gly Ala Ser Val
1 5 10 15

Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Trp Ile
20 25 30

His Trp Glu Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile Gly Tyr
35 40 45

Ile Asn Pro Asn Thr Gly His Thr Glu Tyr Asn Gln Lys Phe Lys Asp
50 55 60

Lys Ala Thr Leu Thr Ala Asp Arg Ser Ser Thr Ala Tyr Met Gln
65 70 75 80

Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg
85 90 95

Thr Tyr Ser Gly Ser Ser His Phe Asp Tyr Trp Gly Gln Gly Thr Thr
100 105 110

Leu

<210> 16

<211> 112

<212> PRT

<213> Homo sapiens

<400> 16

Ser Asp Ile Val Met Thr Gln Thr Pro Val Ser Leu Pro Val Ser Leu
1 5 10 15

Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His
20 25 30

Asn Asn Gly Asp Thr Phe Leu His Trp Tyr Leu Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val
 50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys
 65 70 75 80

Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln
 85 90 95

Thr Thr Leu Ile Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 100 105 110

<210> 17
 <211> 21
 <212> PRT
 <213> Homo sapiens

<400> 17
 Val Leu Phe Tyr Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val
 1 5 10 15

Leu Trp Val Thr Ile
 20

<210> 18
 <211> 61
 <212> PRT
 <213> Homo sapiens

<400> 18
 Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp Leu Glu Ile Ser Leu
 1 5 10 15

Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser Leu Gln Glu Asp Arg
 20 25 30

His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln Lys Glu Glu Gln Leu
 35 40 45

Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly Ala Thr
 50 55 60

<210> 19
 <211> 334
 <212> PRT
 <213> artificial sequence

<220>
 <223> synthetic

<400> 19
 Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30
 Trp Ile His Trp Glu Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Asn Thr Gly His Thr Glu Tyr Asn Gln Lys Phe
 50 55 60
 Lys Asp Lys Ala Thr Leu Thr Ala Asp Arg Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 Ala Arg Thr Tyr Ser Gly Ser Ser His Phe Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Ile Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 115 120 125
 Ser Gly Gly Gly Gly Ser Asp Ile Val Met Thr Gln Thr Pro Val Ser
 130 135 140
 Leu Pro Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser
 145 150 155 160
 Gln Ser Leu Val His Asn Asn Gly Asp Thr Phe Leu His Trp Tyr Leu
 165 170 175
 Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn
 180 185 190
 Arg Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr
 195 200 205
 Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val
 210 215 220
 Tyr Phe Cys Ser Gln Thr Thr Leu Ile Pro Arg Thr Phe Gly Gly Gly
 225 230 235 240
 Thr Lys Leu Glu Ile Lys Arg Gly Gly Gly Gly Ser Val Leu Phe Tyr
 245 250 255
 Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val Leu Trp Val Thr
 260 265 270
 Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp Leu Glu Ile Ser
 275 280 285
 Leu Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser Leu Gln Glu Asp
 290 295 300

Arg His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln Lys Glu Glu Gln
 305 310 315 320

Leu Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly Ala Thr
 325 330

- <210> 20
- <211> 1005
- <212> DNA
- <213> artificial sequence

- <220>
- <223> synthetic

<400> 20
 caagtacaac tgcaacaaag cggagctgaa ctggccaaac caggcgcttc cgtgaagatg 60
 tcttgtaaag ccagcgggta tacatttact aattactgga ttcactggga gaagcaaaga 120
 cctgaacagg gattggaatg gattggatac attaatceta acaccggaca cacagagtat 180
 aatcaaaaat tcaaggataa ggccaccctc acagccgaca gatcttcttc aaccgcctat 240
 atgcaacttt cttccctcac ttctgaagac tccgcagttt acttttgccg acgaacttat 300
 tctggaagct cccatttcga ctactggggt caaggaacaa cactgatcgt gtctagcggc 360
 ggccggagggt ccggcggggg cggtagcggg ggcggagggt ctgatattgt catgactcaa 420
 acacctgtct ctctgcctgt ttcacttggg gatcaageta gcatttctct ccgctctagt 480
 caatctctcg tccacaacaa cggegatact ttcttgcatt ggtatctgca gaaaccaggt 540
 cagtcacctt aactgcttat atacaaagtc tctaatagat tctcaggggt gccagatcga 600
 ttcagtgggt ctgggtccgg tacagatddd aactcaaga tatccagagt agaagcagaa 660
 gatctggggc tgtatttctg cagtcaaaca acacttattc ctctactttt tggaggcggg 720
 acaaaaactg agatcaagcg tggaggcggg gggagtggtt tgttttatct ggccggttggg 780
 ataatgtttc tcgtaaatac agtactttgg gtaacaataa ggaaggaact gaagagaaag 840
 aaaaaatggg atctggaaat atcattggac agtggacacg aaaaaaaagt cacatcatca 900
 ttgcaagaag accggcactt ggaggaggaa ctgaaatgct aagagcaaaa agaagaacaa 960
 ctgcaagaag gcgtacatag aaaagaacca caggaggcaa catag 1005

- <210> 21
- <211> 82
- <212> PRT
- <213> Homo sapiens

<400> 21
 Val Leu Phe Tyr Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val
 1 5 10 15
 Leu Trp Val Thr Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp
 20 25 30

Leu Glu Ile Ser Leu Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser
 35 40 45

Leu Gln Glu Asp Arg His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln
 50 55 60

Lys Glu Glu Gln Leu Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly
 65 70 75 80

Ala Thr

<210> 22

<211> 5759

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 22

acaacttcgt atagcataca ttatacgaag ttatggtacc aagcctaggc ctccaaaaaa 60
 gcctcctcac tacttctgga atagctcaga ggcagaggcg gcctcggcct ctgcataaat 120
 aaaaaaatt agtcagccat ggggcggaga atgggcggaa ctgggcggag ttaggggcgg 180
 gatgggcgga gttagggcg ggactatggt tgctgactaa ttgagatgca tgctttgcat 240
 acttctgcct gctggggagc ctggggactt tccacacctg gttgctgact aattgagatg 300
 catgctttgc atacttctgc ctgctgggga gcctggggac tttccacacc ggatccacca 360
 tgggttcagc tattgagcag gatgggttgc atgctggtag tccgcgcgca tgggtcgaac 420
 gactgtttgg atacgattgg gcccaacaga ctataggctg ttccgacgct gctgtctttc 480
 gtctttctgc acaaggtcgt ccagttctgt tcgtgaaaac cgacttgtcc ggagccctca 540
 atgagttgca agacgaagct gcacgactga gttggcttgc caccactggt gtcccatgtg 600
 ccgcagtact tgacgtcgtc acagaggctg gtcgcgattg gttgctcctt ggagaagtgc 660
 ccggccaaga tcttctcagt tcccaccttg ccctgcca aaaagtttca ataatggctg 720
 acgctatgag aaggctgcac acccttgacc ctgccacatg tccattcgat caccaagcca 780
 aacaccgaat tgaacgagct agaaccgcga tggaagccgg cctcgttgat caagacgatt 840
 tggatgagga acaccagggt ctgcacccg ctgaactctt cgetcgcctc aaagcacgaa 900
 tgccagacgg agatgacttg gtcgtaacc acggagatgc ctgccttctt aacataatgg 960
 tagagaatgg aagatttagc ggcttcattg attgtggacg acttggagtt gcagatcggg 1020
 accaagatat cgctctcgct accagagata ttgctgaaga attgggcgga gaatgggctg 1080
 atcggtttct cgtactctac ggaattgccg cacctgatte ccaacgcatt gctttttacc 1140
 gtcttctgga tgagttcttc taaacgcgtc cccctctcc ctccccccc cctaacgtta 1200
 ctggccgaag ccgcttgaa taaggccggt gtgcgtttgt ctatatgtta ttttccacca 1260

tattgcogtc	ttttggcaat	gtgagggccc	ggaaacctgg	ccctgtcttc	ttgacgagca	1320
ttectagggg	tctttcccct	ctcgccaaag	gaatgcaagg	tctgttgaat	gtcgtgaagg	1380
aagcagttcc	tctggaagct	tcttgaagac	aaacaacgtc	tgtagegacc	ctttgcagge	1440
agcggaacct	cccacctggc	gacaggtgcc	tctgcggcca	aaagccacgt	gtataagata	1500
cacctgcaaa	ggcggcacia	ccccagtgcc	acgttgtgag	ttggatagtt	gtggaaagag	1560
tcaaatggct	ctcctcaagc	gtattcaaca	aggggctgaa	ggatgccccag	aaggtacccc	1620
attgtatggg	atctgatctg	ggcctcgggt	gcacatgctt	tacatgtggt	tagtcogaggt	1680
taaaaaacgt	ctaggcccc	cgaaccacgg	ggacgtgggt	ttcctttgaa	aaacacgatt	1740
gtcgaatca	ccatggtgag	caagggcgag	gagctgttca	ccgggggtgt	gcccacctctg	1800
gtcagactgg	acggcgacgt	aaacggccac	aagttcagcg	tgtccggcga	gggcgagggc	1860
gatgccacct	acggcaagct	gacctgaag	ttcatctgca	ccaccggcaa	gctgcccgctg	1920
ccctggccca	ccctcgtgac	caccttcggc	tacggcctgc	agtgettcgc	ccgtacccc	1980
gaccacatga	agcagcacga	cttcttcaag	tccgccatgc	ccgaaggcta	cgtccaggag	2040
cgcaccatct	tcttcaagga	cgacggcaac	tacaagacce	gcgccgaggt	gaagttcgag	2100
ggcgacacct	tggatgaaccg	catcgagctg	aagggcatcg	acttcaagga	ggacggcaac	2160
atctggggc	acaagctgga	gtacaactac	aacagccaca	acgtctatat	catggccgac	2220
aagcagaaga	acggcatcaa	ggtgaacttc	aagatccgcc	acaacatcga	ggacggcagc	2280
gtgcagctcg	ccgaccacta	ccagcagaac	acccccatcg	gcgacggccc	cgtgetgctg	2340
cccgacaacc	actacctgag	ctaccagtcc	gccctgagca	aagaccccaa	cgagaagcgc	2400
gatcacatgg	tcctgctgga	gttcgtgacc	gccgccggga	tactctcggg	catggacgag	2460
ctgtacaagt	aatcgggcgc	taatcagcca	taccacattt	gtagaggttt	tacttgettt	2520
aaaaaacctc	ccacacctcc	ccctgaacct	gaaacataaa	atgaatgcaa	ttgttgttgt	2580
taacttgttt	attgcagctt	ataatggtta	caaataaagc	aatagcatca	caaatttcac	2640
aaataaagca	tttttttcac	tgcattctag	ttgtggtttg	tccaaactca	tcaatgtatc	2700
ttatcatgtc	ggcgcgttga	cattgattat	tgactagtta	ttaatagtaa	tcaattacgg	2760
ggtcattagt	tcatagccca	tatatggagt	tccgcgttac	ataacttacg	gtaaattggcc	2820
cgctggctg	accgccaac	gacccccgcc	cattgacgtc	aataatgacg	tatgttccca	2880
tagtaacgcc	aatagggact	ttccattgac	gtcaatgggt	ggagtattta	cggtaaactg	2940
cccacttggc	agtacatcaa	gtgtatcata	tgccaagtac	gccccctatt	gacgtcaatg	3000
acggtaaagt	gcccgcctgg	cattatgcc	agtacatgac	cttatgggac	tttccactt	3060
ggcagtacat	ctacgtatta	gtcategcta	ttaccatggt	gatgcggttt	tggcagtaca	3120
tcaatgggcg	tggatagcgg	tttgactcac	ggggatttcc	aagtctccac	cccattgacg	3180
tcaatgggag	tttgttttgg	caccaaatac	aacgggactt	tccaaatgt	cgtaacaact	3240
ccgccccatt	gacgcaaatg	ggcggtaggc	gtgtacgggt	ggaggtctat	ataagcagag	3300
ctctccctat	cagtgataga	gatctcccta	tcagtgatag	agatcgtcga	cgtttagtga	3360
accgtcagat	cgctggaga	cgccatccac	gctgttttga	cctccataga	agacaccggg	3420

accgatccag	cctccgcgge	cggaacggg	gcattggaac	gcggatccccc	cgtgcccaaga	3480
gtgacgtaag	taccgcctat	agagtetata	ggcccccccc	cttggettct	tatgeatget	3540
atactgtttt	tggcttgggg	tctatacacc	cccgttctct	catgttatag	gtgatgggat	3600
agcttagcct	ataggtgtgg	gttattgacc	attattgacc	actccccctat	tggtgacgat	3660
actttccatt	actaatccat	aacatggctc	tttgccacaa	ctctctttat	tggctatatg	3720
ccaatacact	gtccttcaga	gactgacacg	gactctgtat	ttttacagga	tggggctctca	3780
tttattattt	acaaattcac	atatacaaca	ccaccgtccc	cagtgccccgc	agtttttatt	3840
aaacataacg	tgggatctcc	acgcgaatct	cgggtacgtg	ttccggacat	ggtctcttct	3900
ccggtagcgg	cggagcttct	acatccgagc	cctgctccca	tgctcccagc	gactcatggt	3960
cgctcggcag	ctccttgctc	ctaacagtgg	aggccagact	taggcacagc	acgatgccca	4020
ccaccaccag	tgtgcgcgac	aaggccgtgg	cggtagggta	tgtgtctgaa	aatgagctcg	4080
gggagcgggc	ttgcaccgct	gacgcatttg	gaagacttaa	ggcagcggca	gaagaagatg	4140
caggcagctg	agttgttgtg	ttctgataag	agtcagaggt	aactcccgtt	gcgggtgctgt	4200
taacgggtga	gggcagtgta	gtctgagcag	tactcgttgc	tgccgcgcgc	gccaccagac	4260
ataatagctg	acagactaac	agactgttcc	tttccatggg	tcttttctgc	agtcaccgtc	4320
cttgacacga	agcttatact	cgagctctag	attgggaacc	cgggtctctc	gaattcgaga	4380
tctccaccat	gcacagacct	agacgtcgtg	gaactcgtcc	acctccactg	gcactgctcg	4440
ctgctctcct	cctggctgca	cgtgggtgctg	atgcacaagt	acaactgcaa	caaagcggag	4500
ctgaactggc	caaaccaggc	gcttccgtga	agatgtcttg	taaagccagc	gggtatacat	4560
ttactaatta	ctggattcac	tgggagaagc	aaagacctga	acagggattg	gaatggattg	4620
gatacattaa	tcctaacacc	ggacacacag	agtataatca	aaaattcaag	gataaggcca	4680
ccctcacagc	cgacagatct	tcttcaaccg	cctatatgca	actttcttcc	ctcacttctg	4740
aagactccgc	agtttacttt	tgcgcaagaa	cttattctgg	aagctcccat	ttcgactact	4800
gggggtcaagg	aacaacactg	atcgtgtcta	gcggcgccgg	agggtccggc	gggggcggta	4860
gcgggtggcgg	aggttctgat	attgtcatga	ctcaaacacc	tgtctctctg	cctgtttcac	4920
ttggagatca	agctagcatt	tcctgcgcgt	ctagtcaatc	tctcgtccac	aacaacggcg	4980
atactttctt	gcattgggat	ctgcagaaac	caggtcagtc	acctaaactg	cttatataca	5040
aagtctctaa	tagattctca	ggggtgccag	atcgattcag	tggttctggg	tccggtacag	5100
attttacact	caagatatcc	agagtagaag	cagaagatct	gggcgtgtat	ttctgcagtc	5160
aaacaacact	tattcctcgt	acttttggag	gcgggtacaaa	actggagatc	aagcgtggag	5220
gcggaggggag	tgttttgttt	tatctggccg	ttgggataat	gtttctcgtg	aatacagtac	5280
tttgggtaac	aataaggaag	gaactgaaga	gaaagaaaaa	atgggatctg	gaaatatcat	5340
tggacagtgg	acacgaaaaa	aaagtcacat	catcattgca	agaagaccgg	cacttggagg	5400
aggaactgaa	atgtcaagag	caaaaagaag	aacaactgca	agaaggcgta	catagaaaag	5460
aaccacaggg	agcaacatag	gcggcccgct	atcagccata	ccacatttgt	agaggtttta	5520
cttgctttaa	aaaacctccc	acacctcccc	ctgaacctga	aacataaaat	gaatgcaatt	5580

gttgttgtta acttgtttat tgcagcttat aatggttaca aataaagcaa tagcatcaca 5640
aatttcacaa ataaagcatt tttttcactg cattctagtt gtggtttgtc caaactcatc 5700
aatgtatctt atcatgtcta ccggtataac ttcgtataat gtatactata cgaagttag 5759

<210> 23

<211> 7627

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 23

aagcttatac tcgagctcta gattgggaac ccgggtctct cgaattcgag atctccacca 60
tgcacagacc tagacgtcgt ggaactcgtc cacctccact ggcaactgctc gctgctctcc 120
tcttggtgct acgtggtgct gatgcacaag tacaactgca acaaagcgga gctgaactgg 180
ccaaaccagg cgcttccgtg aagatgtctt gtaaagccag ccgggtataca tttactaatt 240
actggattca ctgggagaag caaagacctg aacagggatt ggaatggatt ggatacatta 300
atcctaacac cggacacaca gagtataatc aaaaattcaa ggataaggcc accctcacag 360
ccgacagatc ttcttcaacc gcctatatgc aactttcttc cctcacttct gaagactccg 420
cagtttactt ttgcgcacga acttattctg gaagctccca tttcgactac tggggctcaag 480
gaacaacact gatcgtgtct agcggcggcg gagggctccg cggggcggt agcgggtggcg 540
gaggttctga tattgtcatg actcaaacac ctgtctctct gcctgtttca cttggagatc 600
aagctagcat ttctgcccgc tctagtcfaat ctctcgtcca caacaacggc gatactttct 660
tgcattggta tctgcagaaa ccaggctcagt cacctaaact gcttatatac aaagtctcta 720
atagattctc aggggtgcca gatcgattca gtggttctgg gtccggtaca gattttacac 780
tcaagatata cagagtagaa gcagaagatc tgggcgtgta tttctgcagt caaacaacac 840
ttattcctcg tacttttgga ggcgggtacaa aactggagat caagcgtgga ggcggagggg 900
gtgttttggt ttatctggcc gttgggataa tgtttctcgt aaatacagta ctttgggtaa 960
caataaggaa ggaactgaag agaagaaaa aatgggatct ggaaatatca ttggacagtg 1020
gacacgaaaa aaaagtcaca tcatcattgc aagaagaccg gcacttggag gaggaactga 1080
aatgtcaaga gcaaaaagaa gaacaactgc aagaaggcgt acatagaaaa gaaccacagg 1140
gagcaacata ggcggccgct aatcagccat accacatttg tagaggtttt acttgcttta 1200
aaaaacctcc cacacctccc cctgaacctg aaacataaaa tgaatgcaat tgttggtggt 1260
aacttgttta ttgcagctta taatggttac aaataaagca atagcatcac aaatttcaca 1320
aataaagcat tttttcact gcattctagt tgtggtttgt ccaaactcat caatgtatct 1380
tatcatgtct accggataaa ctctgtataa tgtatactat acgaagttag ccggtagggc 1440
ccctctcttc atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa aggcgcggtt 1500
gctggcggtt ttccataggc tccgcccccc tgacgagcat cacaaaaatc gacgctcaag 1560

tcagaggtgg	cgaaacecga	caggactata	aagataaccag	gcgtttcccc	ctggaagctc	1620
cctcgtgccc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	1680
ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cggtgtaggt	1740
cgttcgcctc	aagctgggct	gtgtgcacga	acccccggtt	cagccccgacc	gctgcgcctt	1800
atccggtaac	tatcgtcttg	agtccaacc	ggtaagacac	gacttatogc	cactggcagc	1860
agccactggt	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	1920
gtgggtggcct	aactacggct	acactagaag	aacagtattt	ggtatctgcg	ctctgctgaa	1980
gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaaaaaa	ccaccgctgg	2040
tagcgggtgg	ttttttggtt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	2100
agatcctttg	atcttttcta	cggggctctga	cgctcagtgg	aacgaaaact	cacgttaagg	2160
gattttggtc	atgggcgcgc	ctcactctcc	tgcaggcatg	agattatcaa	aaaggatctt	2220
cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	2280
aacttggctc	gacagttacc	aatgcttaat	cagtgaggca	cctatctcag	cgatctgtct	2340
atttcgttca	tccatagttg	cctgactccc	cgctcgttag	ataactacga	tacgggaggg	2400
cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	cggctccaga	2460
tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	agaagtggtc	ctgcaacttt	2520
atccgcctcc	atccagtcta	ttaattggtg	ccgggaagct	agagtaagta	gttcgccagt	2580
taatagtttg	cgcaacgttg	ttgccattgc	tacaggcatc	gtgggtgtcac	gctcgtcgtt	2640
tggtatggct	tcattcagct	ccggttccca	acgatcaagg	cgagttacat	gateccccat	2700
gttgtgcaaa	aaagcgggta	gctccttcgg	tcctccgata	gttgtcagaa	gtaagttggc	2760
cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat	tctcttactg	tcatgccatc	2820
cgtaagatgc	ttttctgtga	ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	2880
gcggcgaccg	agttgctctt	gccccggcgc	aatacgggat	aatactgcgc	cacatagcag	2940
aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	cgaaaactct	caaggatctt	3000
accgctggtg	agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	cttcagcatc	3060
ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	aggcaaatg	ccgcaaaaaa	3120
gggaataagg	gcgacacgga	aatggtgaat	actcactctc	ttcctttttc	aatattattg	3180
aagcatttat	cagggttatt	gtctcatgag	cggatacata	tttgaatgta	tttagaaaaa	3240
taaacaaata	ggggttccgc	gcacatttcc	ccgaaaagtg	ccacctgacg	tcaggtagac	3300
aacttcgtat	agcatacatt	atacgaagtt	atggtaccaa	gcctaggcct	ccaaaaaagc	3360
ctctcacta	cttctggaat	agctcagagg	cagagggcgc	ctcggcctct	gcataaataa	3420
aaaaaattag	tcagccatgg	ggcggagaat	gggcgggaact	gggcggagtt	aggggcggga	3480
tgggcggagt	taggggcggg	actatgggtg	ctgactaatt	gagatgcatg	ctttgcatac	3540
ttctgcctgc	tggggagcct	ggggactttc	cacacctggt	tgctgactaa	ttgagatgca	3600
tgctttgcat	acttctgcct	gctggggagc	ctggggactt	tccacaccgg	atccaccatg	3660
gqttcaqcta	ttgaqcaqqa	tggqttqcat	qctqgtaqtc	ccqccqcatq	qqtcqaacqa	3720

ctgtttggat	acgattgggc	ccaacagact	ataggctggt	ccgacgctgc	tgtctttcgt	3780
ctttctgcac	aaggctgtec	agttctgttc	gtgaaaaccg	acttgtccgg	agccctcaat	3840
gagttgcaag	acgaagetgc	acgactgagt	tggettgcc	ccactggtgt	cccattgtgcc	3900
gcagtacttg	acgtcgtcac	agaggctggt	cgcgattggt	tgtcccttgg	agaagtgcc	3960
ggccaagatc	ttctcagttc	ccaccttgcc	cctgccgaaa	aagtttcaat	aatggctgac	4020
gctatgagaa	ggctgcacac	ccttgaccct	gccacatgtc	cattcgatca	ccaagccaaa	4080
caccgaattg	aacgagctag	aaccgcgatg	gaagccggcc	tcgttgatca	agacgatttg	4140
gatgaggaac	accagggctc	cgcacccgct	gaactcttcg	ctgcctcaa	agcacgaatg	4200
ccagacggag	atgacttggt	cgtaaccac	ggagatgctt	gccttctaa	cataatggta	4260
gagaatggaa	gatttagcgg	cttcattgat	tgtggacgac	ttggagttgc	agatcggtag	4320
caagatatcg	ctctcgtcac	cagagatatt	gctgaagaat	tggcgggaga	atgggctgat	4380
cggtttctcg	tactctacgg	aattgcgcga	cctgattccc	aacgcattgc	ttttaccgt	4440
cttctggatg	agttcttcta	aacgcgtccc	ccctctcctt	cccccccc	taacgttact	4500
ggccgaagcc	gcttggaata	aggccggtgt	gcgtttgtct	atatgttatt	ttccaccata	4560
ttgccgtctt	ttggcaatgt	gagggcccgg	aaacctggcc	ctgtcttctt	gacgagcatt	4620
cctaggggct	ttccctctct	cgccaaagga	atgcaaggtc	tgttgaatgt	cgtgaaggaa	4680
gcagttcctc	tggaagcttc	ttgaagacaa	acaacgtctg	tagcgacctt	ttgcaggcag	4740
cggaaccccc	cacctggcga	caggtgcctc	tgcggccaaa	agccacgtgt	ataagataca	4800
cctgcaaagg	cggcacaacc	ccagtgcac	gttgtgagtt	ggatagttgt	ggaaagagtc	4860
aatggctctt	cctcaagcgt	attcaacaag	gggctgaagg	atgcccagaa	ggtaccccat	4920
tgtatgggat	ctgatctggg	gcctcgggtc	acatgcttta	catgtgttta	gtcgaggtta	4980
aaaaacgtct	aggccccccg	aaccacgggg	acgtggtttt	cctttgaaaa	acacgattgc	5040
tcgaatcacc	atggtgagca	agggcgagga	gctgttcacc	ggggtggtgc	ccatcctggt	5100
cgagctggac	ggcgacgtaa	acggccacaa	gttcagcgtg	tcggcgagg	gcgagggcga	5160
tgccacctac	ggcaagctga	ccctgaagtt	catctgcacc	accggcaagc	tgcccgtgcc	5220
ctggcccacc	ctcgtgacca	ccttcggcta	cggcctgcag	tgettccccc	gctaccccga	5280
ccacatgaag	cagcagcact	tcttcaagtc	cgccatgccc	gaaggctacg	tccaggagcg	5340
caccatcttc	ttcaaggacg	acggcaacta	caagaccgc	gcccaggtga	agttcgaggg	5400
cgacaccctg	gtgaaccgca	tcgagctgaa	gggcatcgac	ttcaaggagg	acggcaacat	5460
cctggggcac	aagctggagt	acaactacaa	cagccacaac	gtctatatca	tggccgacaa	5520
gcagaagaac	ggcatcaagg	tgaacttcaa	gatccgccac	aacatcgagg	acggcagcgt	5580
gcagctcgcc	gaccactacc	agcagaacac	ccccatcggc	gacggccccg	tgetgctgcc	5640
cgacaaccac	tacctgagct	accagtcgc	cctgagcaaa	gaccccaacg	agaagcgcga	5700
tcacatggtc	ctgctggagt	tcgtgaccgc	cgcggggtac	actctcggca	tggacgagct	5760
gtacaagtaa	tcggccgcta	atcagccata	ccacatttgt	agaggtttta	cttgetttaa	5820

aaaacctccc acacctcccc ctgaacctga aacataaaat gaatgcaatt gttggttgta	5880
acttgtttat tgcagcttat aatggttaca aataaagcaa tagcatcaca aatttcacaa	5940
ataaagcatt tttttcactg cattctagtt gtggtttgtc caaactcacc aatgtatctt	6000
atcatgtcgg cgcggtgaca ttgattattg actagttatt aatagtaacc aattacgggg	6060
tcattagttc atagcccata tatggagttc cgcggttacc aacttacggt aatggccccg	6120
cctggctgac cgcaccaacga cccccgcca ttgacgtcaa taatgacgta tgttcccata	6180
gtaacgcca tagggacttt ccattgacgt caatgggtgg agtatttacg gtaaacctgcc	6240
cacttggcag tacatcaagt gtatcatatg ccaagtacgc cccctattga cgtcaatgac	6300
ggtaaattgg ccgcctggca ttatgccag tacatgacct tatgggactt tccacttgg	6360
cagtacatct acgtattagt catcctatt accatggtga tgcggttttg gcagtacacc	6420
aatgggogtg gatagcgggt tgactcaccg ggatttccaa gtctccacc cattgacgtc	6480
aatgggagtt tgttttgga ccaaaatcaa cgggactttc caaaatgtcg taacaactcc	6540
gccccattga cgc aaatggg cggtagcgt gtacgggtgg aggtctatat aagcagagct	6600
ctccctatca gtgatagaga tctccctacc agtgatagag atcgtcagcg tttagtgaac	6660
cgtcagatcg cctggagacg ccattccacc tgttttgacc tccatagaag acaccgggac	6720
cgatccagcc tccgcggccg ggaacgggtgc attggaacgc ggattccccg tgccaagagt	6780
gacgtaagta ccgcctatag agtctatagg cccacccctc tggtctctta tgcattgctat	6840
actgtttttg gcttggggtc tatacaccac cgttctctca tgttataggt gatggtatag	6900
cttagcctat aggtgtgggt tattgacct tattgaccac tccctattg gtgacgatac	6960
ttccattac taatccataa catggctctt tgccacaact ctctttattg gctatatgcc	7020
aatacactgt ccttcagaga ctgacacgga ctctgtatct ttacaggatg gggctctcatt	7080
tattatttac aaattcacat atacaacacc accgtccccca gtgccccgag tttttattaa	7140
acataacgtg ggatctccac gcgaatctcg ggtacgtggt ccggacatgg tctctctctc	7200
ggtagcggcg gacgtctacc atccgagccc tgctcccatg cctccagcga ctcatggctc	7260
ctcggcagct ccttgcctct aacagtggag gccagactta ggcacagcac gatgccacc	7320
accaccagtg tgccgcacaa ggccgtggcg gtaggggatg tgtctgaaaa tgagctcggg	7380
gagcgggctt gcaccgctga cgcatttggg agacttaagg cagcggcaga agaagatgca	7440
ggcagctgag ttggttggtt ctgataagag tcagaggtaa ctcccgttgc ggtgctgta	7500
acgggtggag gcagtgtagt ctgagcagta ctggttgctg ccgcgcgcgc caccagacat	7560
aatagctgac agactaacag actgttccct tccatgggtc tttctgcag tcaccgtcct	7620
tgacacg	7627

<210> 24

<211> 2669

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 24

agcctaggcc	tccaaaaaag	cctcctcact	acttctggaa	tagctcagag	gcagaggcgg	60
cctcggcctc	tgcataaata	aaaaaaatta	gtcagccatg	gggcggagaa	tgggcggaac	120
tgggcggagt	taggggcggg	atgggcggag	ttaggggcgg	gactatgggt	gctgactaat	180
tgagatgcat	gctttgcata	cttctgcctg	ctggggagcc	tggggacttt	ccacacctgg	240
ttgctgacta	attgagatgc	atgctttgca	tacttctgce	tgctggggag	cctggggact	300
ttccacaccg	gatccaccat	gggttcagct	attgagcagg	atgggttgca	tgctggtagt	360
cccgcgcgat	gggtcgaacg	actgtttgga	facgattggg	cccaacagac	tataggetgt	420
tccgaecgtg	ctgtctttcg	tctttctgca	caaggctgtc	cagttctggt	cgtgaaaacc	480
gacttgtccg	gagccctcaa	tgagttgcaa	gacgaagctg	cacgactgag	ttggettgcc	540
accactgggtg	tcccatgtgc	cgcagtaact	gacgtcgtca	cagaggctgg	tcgcgattgg	600
ttgctccttg	gagaagtgcc	cgccaagat	cttctcagtt	cccaccttgc	ccctgccgaa	660
aaagtttcaa	taatggctga	cgctatgaga	aggetgcaca	cccttgacc	tgccacatgt	720
ccattcgatc	accaagccaa	acaccgaatt	gaacgagcta	gaaccgcgat	ggaagccggc	780
ctcgttgatc	aagacgattt	ggatgaggaa	caccagggtc	tcgcaccgc	tgaactcttc	840
gctcgcctca	aagcacgaat	gccagacgga	gatgacttgg	tcgtaacca	cggagatgcc	900
tgccttecta	acataatggt	agagaatgga	agatttagcg	gcttcattga	ttgtggacga	960
cttggagttg	cagatcggta	ccaagatata	gctctcgtca	ccagagatat	tgctgaagaa	1020
ttgggcggag	aatgggctga	tgggtttctc	gtactctacg	gaattgcgc	acctgatccc	1080
caacgcattg	ctttttaccg	tcttctggat	gagttcttct	aaacgcgtcc	cccctctccc	1140
tcccccccc	ctaacgttac	tgcccgaaag	cgcttggaat	aaggccgggtg	tgcgtttgtc	1200
tatatgttat	ttccaccat	attgccgtct	tttggaatg	tgagggcccg	gaaacctggc	1260
cctgtcttct	tgacgagcat	tcctaggggt	ctttccctc	tcgccaaagg	aatgcaaggt	1320
ctgttgaatg	tcgtgaagga	agcagttcct	ctggaagctt	cttgaagaca	aacaacgtct	1380
gtagegaccc	tttgcaggca	gcggaacccc	ccacctggcg	acaggtgcct	ctgcggccaa	1440
aagccacgtg	tataagatac	acctgcaaag	gcggcacaac	cccagtgcca	cgttgtgagt	1500
tggatagttg	tggaaagagt	caaatggctc	tcctcaagcg	tattcaacaa	ggggetgaag	1560
gatgccca	aggtaccca	ttgtatggga	tctgatctgg	ggcctcgggtg	cacatgcttt	1620
acatgtgttt	agtcgaggtt	aaaaaacgtc	taggcccccc	gaaccacggg	gacgtggttt	1680
tcctttgaaa	aacacgattg	ctcgaatcac	catggtgagc	aagggcgagg	agctgttcac	1740
cggggtgggtg	cccatcctgg	tcgagctgga	cggcgacgta	aacggccaca	agttcagcgt	1800
gtccggcgag	ggcgagggcg	atgccaccta	cggcaagctg	acctgaagt	tcctctgcac	1860
caccggcaag	ctgcccgtgc	cctggcccac	cctcgtgacc	accttcggct	acggcctgca	1920
gtgcttcgcc	cgctaccccg	accacatgaa	gcagcagcag	ttcttcaagt	ccgccatgcc	1980
cgaaggctac	gtccaggagc	gcacatctt	cttcaaggac	gacggcaact	acaagacccg	2040

cgccgaggtg aagttcgagg gcgacacccct ggtgaaccgc atcgagctga agggcatcga 2100
 cttcaaggag gacggcaaca tcctggggca caagctggag tacaactaca acagccacaa 2160
 cgtctatata atggccgaca agcagaagaa cggcatcaag gtgaacttca agatccgcca 2220
 caacatcgag gacggcagcg tgcagctcgc cgaccactac cagcagaaca ccccatcgg 2280
 cgacggcccc gtgctgctgc ccgacaacca ctacctgagc taccagtccg ccctgagcaa 2340
 agaccccaac gagaagcgcg atcacatggt cctgctggag ttcgtgaccg ccgcccggat 2400
 cactctcggc atggacgagc tgtacaagta atcgcccgct aatcagccat accacatttg 2460
 tagaggtttt acttgcttta aaaaacctcc cacacctccc cctgaacctg aaacataaaa 2520
 tgaatgcaat tggtgtgtt aacttgctta ttgcagctta taatggttac aaataaagca 2580
 atagcatcac aaatttcaca aataaagcat ttttttact gcattctagt tgtggtttgt 2640
 ccaaactcat caatgtatct tatcatgtc 2669

<210> 25

<211> 3003

<212> DNA

<213> artificial sequence

<220>

<223> synthetic

<400> 25

gttgacattg attattgact agttattaat agtaatcaat tacggggtea ttagttcata 60
 gccatataat ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc 120
 ccaacgacct ccgcccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag 180
 ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
 atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggg aaatggcccg 300
 cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg 360
 tattagtcac cgctattacc atggtgatgc ggttttggca gtacatcaat gggcgtggat 420
 agcggtttga ctcaacggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt 480
 tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
 aaatgggccc taggcgtgta cgggtggagg tctatataag cagagctctc cctatcagtg 600
 atagagatct ccctatcagt gatagagatc gtcgacgttt agtgaaccgt cagatcgctt 660
 ggagacgcca tccacgctgt tttgacctcc atagaagaca ccgggaccga tccagcctcc 720
 gcgcccgga acggtgcatt ggaacgcgga ttccccgtgc caagagtgac gtaagtaccg 780
 cctatagagt ctataggccc acccccttgg cttcttatgc atgctatact gtttttggct 840
 tggggtctat acacccccgc ttctcatgt tataggtgat ggtatagctt agcctatagg 900
 tgtgggttat tgaccattat tgaccactcc cctattggtg acgatacttt ccattactaa 960
 tccataacat ggctctttgc cacaactctc tttattggct atatgccaat aactgtctct 1020
 tcagagactg acacggactc tgtattttta caggatgggg tctcatttat tatttataaa 1080

ttcacatata caacaccacc gtccccagtg cccgcagttt ttattaaaca taacgtggga	1140
tctccacgcg aatctcgggt acgtgttccg gacatggtct cttctccggt agcggcggag	1200
cttctacatc cgagccctgc tcccatgcct ccagcgcactc atggtcgctc ggcagctcct	1260
tgetcetaac agtggaggee agaacttaggc acagcaecat gccaccacc accagtggtc	1320
cgcacaaggc cgtggcggta gggatatgtg ctgaaaatga gctcggggag cgggcttgca	1380
ccgctgacgc atttggaaga ctttaaggcag cggcagaaga agatgcagge agctgagttg	1440
ttgtgttctg ataagagtca gaggtaactc ccgttgccgt gctgttaacg gtggagggca	1500
gtgtagtctg agcagtactc gttgctgccc cgcgcgccac cagacataat agctgacaga	1560
ctaacagact gttcctttcc atgggtcttt tctgcagtca ccgtccttga cacgaagctt	1620
atactcgagc tctagattgg gaaccgggt ctctcgaatt cgagatctcc accatgcaca	1680
gacctagacg tcgtggaact cgtccacctc cactggcact gctcgtctgt ctcctcctgg	1740
ctgcacgtgg tgctgatgca caagtacaac tgcaacaaag cggagctgaa ctggccaaac	1800
caggcgttc cgtgaagatg tcttgtaaag ccagcgggta tacatttact aattactgga	1860
ttcactggga gaagcaaaga cctgaacagg gattggaatg gattggatac attaatccta	1920
acaccggaca cacagagtat aatcaaaaat tcaaggataa ggccaccctc acagccgaca	1980
gatcttcttc aaccgcctat atgcaacttt ctccctcac ttctgaagac tccgcagttt	2040
acttttgccg acgaacttat tctggaagct cccatttcca ctactgggggt caaggaacaa	2100
cactgatcgt gtctagcggc ggcggagggt cgggcggggg cggtagcgggt ggcggaggtt	2160
ctgatattgt catgactcaa acacctgtct ctctgcctgt ttcacttggga gatcaageta	2220
gcatttctcg ccgctctagt caatctctcg tccacaacaa cggcgatact ttcttgcatt	2280
ggtatctgca gaaaccagggt cagtcacctc aactgcttat atacaaagtc tctaatagat	2340
tctcaggggt gccagatcga ttcagtgggt ctgggtccgg tacagatfff aactcaaga	2400
tatccagagt agaagcagaa gatctgggcg tgtatctctg cagtcaaaca acacttattc	2460
ctcgtacttt tggaggcgggt acaaaactgg agatcaagcg tggaggcggga gggagtgttt	2520
tgttttatct ggcggttggg ataatgtttc tcgtaaatac agtactttgg gtaacaataa	2580
ggaaggaact gaagagaaag aaaaaatggg atctggaaat atcattggac agtggacacg	2640
aaaaaaaaagt cacatcatca ttgcaagaag accggcactt ggaggaggaa ctgaaatgtc	2700
aagagcaaaa agaagaacaa ctgcaagaag gcgtacatag aaaagaacca caggagcaa	2760
cataggcggc cgctaatcag ccataccaca tttgtagagg ttttacttgc tttaaaaaac	2820
ctcccacacc tccccctgaa cctgaaacat aaaatgaatg caattgttgt tgttaacttg	2880
tttattgcag cttataatgg ttacaaataa agcaatagca tcacaaatft cacaaataaa	2940
gcattttttt cactgcattc tagttgtgggt ttgtccaaac tcatcaatgt atcttatcat	3000
gtc	3003

<210> 26

<211> 208

<212> PRT

<213> Homo sapiens

<400> 26

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
 1 5 10 15
 Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro
 20 25 30
 Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
 35 40 45
 Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val
 50 55 60
 Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
 65 70 75 80
 Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr
 85 90 95
 Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
 100 105 110
 Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
 115 120 125
 Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 130 135 140
 Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 145 150 155 160
 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser
 165 170 175
 Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 180 185 190
 Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
 195 200 205

<210> 27

<211> 8

<212> PRT

<213> Homo sapiens

<400> 27

Gly Tyr Thr Phe Thr Asn Tyr Trp

1

5

<210> 28

<211> 8

<212> PRT

<213> Homo sapiens

<400> 28

Ile Asn Pro Asn Thr Gly His Thr
1 5

<210> 29

<211> 13

<212> PRT

<213> Homo sapiens

<400> 29

Cys Ala Arg Thr Tyr Ser Gly Ser Ser His Phe Asp Tyr
1 5 10

<210> 30

<211> 10

<212> PRT

<213> Homo sapiens

<400> 30

Ser Leu Val His Asn Asn Gly Asp Thr Phe
1 5 10

<210> 31

<211> 9

<212> PRT

<213> Homo sapiens

<400> 31

Ser Gln Thr Thr Leu Ile Pro Arg Thr
1 5

<210> 32

<211> 8

<212> PRT

<213> Homo sapiens

<400> 32

Gly Phe Thr Phe Ser Asn Ala Trp
1 5

<210> 33

<211> 10

<212> PRT

<213> Homo sapiens

<400> 33

Ile Leu Ser Lys Thr Asp Gly Gly Thr Thr
1 5 10

<210> 34

<211> 13

<212> PRT

<213> Homo sapiens

<400> 34

Thr Thr Ala Asp Phe Trp Ser Ala Tyr Ser Ser Asp Tyr
 1 5 10

<210> 35

<211> 4

<212> PRT

<213> Homo sapiens

<400> 35

Gln Ser Leu Leu
 1

<210> 36

<211> 7

<212> PRT

<213> Homo sapiens

<400> 36

His Ser Asn Gly Tyr Asn Tyr
 1 5

<210> 37

<211> 9

<212> PRT

<213> Homo sapiens

<400> 37

Met Gln Gly Leu Gln Thr Pro Tyr Thr
 1 5

<210> 38

<211> 122

<212> PRT

<213> Homo sapiens

<400> 38

Glu Val Gln Leu Val Glu Ser Gly Gly Ala Ile Val Lys Pro Gly Gly
 1 5 10 15

Ser His Arg Val Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Arg Gly Leu Glu Trp Val
 35 40 45

Gly Arg Ile Leu Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60

Pro Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Met
65 70 75 80

Leu Phe Leu Gln Met Asp Ser Leu Lys Ile Glu Asp Thr Ala Val Tyr
85 90 95

Phe Cys Thr Thr Ala Asp Phe Trp Ser Ala Tyr Ser Ser Asp Tyr Trp
100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 39

<211> 112

<212> PRT

<213> Homo sapiens

<400> 39

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
20 25 30

Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Met Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Gly
85 90 95

Leu Gln Thr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 40

<211> 452

<212> PRT

<213> Homo sapiens

<400> 40

Glu Val Gln Leu Val Glu Ser Gly Gly Ala Ile Val Lys Pro Gly Gly
1 5 10 15

Ser His Arg Val Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Asn Ala
20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Arg Gly Leu Glu Trp Val
35 40 45

Gly Arg Ile Leu Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60
 Pro Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Met
 65 70 75 80
 Leu Phe Leu Gln Met Asp Ser Leu Lys Ile Glu Asp Thr Ala Val Tyr
 85 90 95
 Phe Cys Thr Thr Ala Asp Phe Trp Ser Ala Tyr Ser Ser Asp Tyr Trp
 100 105 110
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Lys Thr Thr Ala Pro
 115 120 125
 Ser Val Tyr Pro Leu Ala Pro Val Cys Gly Asp Thr Thr Gly Ser Ser
 130 135 140
 Val Thr Leu Gly Cys Leu Val Lys Gly Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160
 Leu Thr Trp Asn Ser Gly Ser Leu Ser Ser Gly Val His Thr Phe Pro
 165 170 175
 Ala Val Leu Gln Ser Asp Leu Tyr Thr Leu Ser Ser Ser Val Thr Val
 180 185 190
 Thr Ser Ser Thr Trp Pro Ser Gln Ser Ile Thr Cys Asn Val Ala His
 195 200 205
 Pro Ala Ser Ser Thr Lys Val Asp Lys Lys Ile Glu Pro Arg Gly Pro
 210 215 220
 Thr Ile Lys Pro Cys Pro Pro Cys Lys Cys Pro Ala Pro Asn Leu Leu
 225 230 235 240
 Gly Gly Pro Ser Val Phe Ile Phe Pro Pro Lys Ile Lys Asp Val Leu
 245 250 255
 Met Ile Ser Leu Ser Pro Ile Val Thr Cys Val Val Val Asp Val Ser
 260 265 270
 Glu Asp Asp Pro Asp Val Gln Ile Ser Trp Phe Val Asn Asn Val Glu
 275 280 285
 Val His Thr Ala Gln Thr Gln Thr His Arg Glu Asp Tyr Asn Ser Thr
 290 295 300
 Leu Arg Val Val Ser Ala Leu Pro Ile Gln His Gln Asp Trp Met Ser
 305 310 315 320
 Gly Lys Glu Phe Lys Cys Lys Val Asn Asn Lys Asp Leu Pro Ala Pro
 325 330 335
 -- -- - -- - - - - - - - - - - - - - - - -

Ile Glu Arg Thr Ile Ser Lys Pro Lys Gly Ser Val Arg Ala Pro Gln
340 345 350

Val Tyr Val Leu Pro Pro Pro Glu Glu Glu Met Thr Lys Lys Gln Val
355 360 365

Thr Leu Thr Cys Met Val Thr Asp Phe Met Pro Glu Asp Ile Tyr Val
370 375 380

Glu Trp Thr Asn Asn Gly Lys Thr Glu Leu Asn Tyr Lys Asn Thr Glu
385 390 395 400

Pro Val Leu Asp Ser Asp Gly Ser Tyr Phe Met Tyr Ser Lys Leu Arg
405 410 415

Val Glu Lys Lys Asn Trp Val Glu Arg Asn Ser Tyr Ser Cys Ser Val
420 425 430

Val His Glu Gly Leu His Asn His His Thr Thr Lys Ser Phe Ser Arg
435 440 445

Thr Pro Gly Lys
450

- <210> 41
- <211> 219
- <212> PRT
- <213> Homo sapiens

<400> 41

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
20 25 30

Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Met Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Gly
85 90 95

Leu Gln Thr Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105 110

Arg Ala Asp Ala Ala Pro Thr Val Ser Ile Phe Pro Pro Ser Ser Glu
115 120 125

Gln Leu Thr Ser Gly Gly Ala Ser Val Val Cys Phe Leu Asn Asn Phe
130 135 140

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
 180 185 190

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 195 200 205

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
 210 215 220

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 225 230 235 240

Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 245 250 255

<210> 43

<211> 336

<212> PRT

<213> Artificial sequence

<220>

<223> synthetic

<400> 43

Glu Val Gln Leu Val Glu Ser Gly Gly Ala Ile Val Lys Pro Gly Gly
 1 5 10 15

Ser His Arg Val Ser Cys Glu Ala Ser Gly Phe Thr Phe Ser Asn Ala
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Arg Gly Leu Glu Trp Val
 35 40 45

Gly Arg Ile Leu Ser Lys Thr Asp Gly Gly Thr Thr Asp Tyr Ala Ala
 50 55 60

Pro Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Met
 65 70 75 80

Leu Phe Leu Gln Met Asp Ser Leu Lys Ile Glu Asp Thr Ala Val Tyr
 85 90 95

Phe Cys Thr Thr Ala Asp Phe Trp Ser Ala Tyr Ser Ser Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly
 115 120 125

Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Val Met Thr Gln Ser
 130 135 140

Pro Leu Ser Leu Pro Val Thr Pro Gly Glu Pro Ala Ser Ile Ser Cys
 145 150 155 160

Arg Ser Ser Gln Ser Leu Leu His Ser Asn Gly Tyr Asn Tyr Leu Asp
 165 170 175

Trp Tyr Leu Gln Lys Pro Gly Gln Ser Pro Gln Leu Leu Ile Tyr Leu
 180 185 190

Gly Ser Asn Arg Ala Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly
 195 200 205

Ser Gly Thr Asp Phe Thr Leu Lys Ile Ser Arg Met Glu Ala Glu Asp
 210 215 220

Val Gly Val Tyr Tyr Cys Met Gln Gly Leu Gln Thr Pro Tyr Thr Phe
 225 230 235 240

Gly Gln Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Val Leu
 245 250 255

Phe Tyr Leu Ala Val Gly Ile Met Phe Leu Val Asn Thr Val Leu Trp
 260 265 270

Val Thr Ile Arg Lys Glu Leu Lys Arg Lys Lys Lys Trp Asp Leu Glu
 275 280 285

Ile Ser Leu Asp Ser Gly His Glu Lys Lys Val Thr Ser Ser Leu Gln
 290 295 300

Glu Asp Arg His Leu Glu Glu Glu Leu Lys Cys Gln Glu Gln Lys Glu
 305 310 315 320

Glu Gln Leu Gln Glu Gly Val His Arg Lys Glu Pro Gln Gly Ala Thr
 325 330 335

<210> 44

<211> 7633

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 44

aagcttatac tcgagctcta gattgggaac ccgggtctct cgaattcgag atctccacca 60

tgcacagacc tagacgtcgt ggaactcgtc cacctccact ggcactgctc gctgctctcc 120

tcttggtgct acgtggtgct gatgcagagg tgcagctggt ggagtctggg ggagccatag 180

taaagecggg ggggtcccat agagtctcct gtgaagcctc tggattcact ttcagtaacg 240

cctggatgag ttgggtccgc caggctccag ggagggggct ggagtgggtt ggccgtattt 300

taagcaagac tgatggtggg acgacagact acgctgcacc cgtgaaagac agattcacca 360

tttcaagaga tgattctaaa aatatgttgt ttctgcaaat ggacagcctg aaaatcgagg 420

..... 480

acacagccgt	gtatttctgt	accacggccg	atthtttgag	tgettattct	tctgactact	480
ggggccagg	aaccctggte	accgtctect	caggaggtgg	aggttccggg	ggcgggggct	540
ccggcggagg	tggatcagat	attgtgatga	ctcagtctcc	actctccctg	cccgtcaccc	600
ctggagagcc	ggcctccate	tectgcaggt	ctagtcagag	cctcctgcat	agtaatgggt	660
acaactatth	ggattggtag	ctacagaagc	cagggcagtc	tccacaactc	ctgatctatt	720
tgggttctaa	tggggcctcc	ggggtccctg	acaggttcag	tggcagtggg	tcaggcacag	780
atthtacact	gaaaatcagc	agaatggagg	ctgaggatgt	tggggthtat	tactgcatgc	840
aaggtctaca	aactccgtac	actthtggcc	aggggaccaa	gctggagatc	aaaggaggcg	900
gagggagtgt	thtggthtat	ctggccgttg	ggataatgtt	tctcgtaaat	acagtactth	960
gggtaacaat	aaggaaggaa	ctgaagagaa	agaaaaaatg	ggatctggaa	atatcattgg	1020
acagtggaca	cgaaaaaaa	gtcacatcat	cattgcaaga	agaccggcac	ttggaggagg	1080
aactgaaatg	tcaagagcaa	aaagaagaac	aactgcaaga	aggcgtacat	agaaaagaac	1140
cacagggagc	aacataggcg	gccgctaata	agccatacca	catttgtaga	ggthttactt	1200
gctthaaaa	acctcccaca	cctccccctg	aacctgaaac	ataaaatgaa	tgcaattgtt	1260
gttgthtaact	tgtthtattgc	agcttataat	ggttacaaat	aaagcaatag	catcacaat	1320
thcacaata	aagcaththt	thcactgcat	tctagttgtg	gthtgtccaa	actcatcaat	1380
gtatcttata	atgtctaccg	gtataacttc	gtataatgta	tactatacga	agttagccgg	1440
tagggccct	ctcttcatgt	gagcaaaaag	ccagcaaaaag	gccaggaacc	gtaaaaaggc	1500
cgcgthtctg	gcgththtcc	ataggctccg	ccccctgac	gagcatcaca	aaaatcgacg	1560
ctcaagtcag	aggtggcgaa	acccgacagg	actataaaga	taccaggcgt	thccccctgg	1620
aagctccctc	gtgcgctctc	ctgttccgac	cctgcccgtt	accggatacc	tgteccgctt	1680
tctcccttcg	ggaagcgtgg	cgctthtctca	tagctcacgc	tgtaggtata	tcagttcggg	1740
gtaggtcgtt	cgctccaagc	tgggctgtgt	gcacgaacce	cccgttcagc	ccgaccgctg	1800
cgctttatcc	ggtaactata	gtcttgagtc	caaccgggta	agacacgaact	tatcgccact	1860
ggcagcagcc	actggtaaca	ggattagcag	agcgaggtat	gtaggcgggtg	ctacagagtt	1920
cttgaagtgg	tggcctaact	acggctacac	tagaagaaca	gtatthggta	tctgcgctct	1980
gctgaagcca	gthaccttcg	gaaaaagagt	tggtagctct	tgateccggca	aacaaaccac	2040
cgctggtagc	ggtggththt	thgthtgcaa	gcagcagatt	acgcgcagaa	aaaaaggatc	2100
tcaagaagat	cctthgatct	thtctacggg	gtctgacgct	cagtggaacg	aaaactcacg	2160
thtaagggatt	thggctcatgg	gcgcgcctca	tactcctgca	ggcatgagat	tatcaaaaag	2220
gatcttcacc	tagatcctth	thaaatthaaa	atgaagthth	aatcaatct	aaagtatata	2280
tgagthaaact	tggctctgaca	gthaccaatg	ctthaatcagt	gaggcaccta	tctcagcgat	2340
ctgtctatth	cgthcatcca	tagthgcctg	actccccgtc	gtgtagataa	ctacgatacg	2400
ggagggctta	ccatctggcc	ccagtgcctgc	aatgataccg	cgagaccac	gctcaccggc	2460
tccagatthta	tcagcaataa	accagccagc	cggaagggcc	gagcgcagaa	gtggctcctgc	2520
aactthtatcc	gcctccatcc	agctctatthaa	thgthtgcgg	gaagctagag	thagtagtht	2580

gccagttaat	agtttgcgca	acgttggtgc	cattgctaca	ggcatcgtgg	tgtcacgctc	2640
gtcgtttggg	atggcttcat	tcagctccgg	ttcccaacga	tcaaggcgag	ttacatgatc	2700
ccccatggtg	tgcaaaaaag	cggttagctc	cttcggctct	ccgatcgttg	tcagaagtaa	2760
gttggccgca	gtgttatcac	tcatggttat	ggcagcactg	cataattctc	ttactgtcat	2820
gccatecgta	agatgctttt	ctgtgactgg	tgagtaactca	accaagtcac	tctgagaata	2880
gtgtatgcgg	cgaccgagtt	gctcttgccc	ggcgtcaata	cgggataata	ctgcgccaca	2940
tagcagaact	ttaaaagtgc	tcacattgg	aaaacgttct	tcggggcgaa	aactctcaag	3000
gatcttaccg	ctggtgagat	ccagttegat	gtaaccact	cgtgcacca	actgatcttc	3060
agcatctttt	actttcacca	gcgtttctgg	gtgagcaaaa	acaggaaggc	aaaatgccgc	3120
aaaaaagga	ataagggcga	cacggaaatg	ttgaatactc	atactcttcc	tttttcaata	3180
ttattgaagc	atztatcagg	gttattgtct	catgagcgga	tacatatttg	aatgtattta	3240
gaaaaataaa	caaatagggg	ttccgcgcac	atctccccga	aaagtgccac	ctgacgtcag	3300
gtacacaact	tcgtatagca	tacattatac	gaagttatgg	taccaagcct	aggcctccaa	3360
aaaagcctcc	tcactacttc	tggaatagct	cagaggcaga	ggcggcctcg	gcctctgcat	3420
aaataaaaaa	aattagtcag	ccatggggcg	gagaatgggc	ggaactgggc	ggagttaggg	3480
gcgggatggg	cggagttagg	ggcgggacta	tggttgetga	ctaattgaga	tgcatgcttt	3540
gcatacttct	gcctgctggg	gagcctgggg	actttccaca	cctggttgct	gactaattga	3600
gatgcatgct	ttgcatactt	ctgcctgctg	gggagcctgg	ggactttcca	caccggatcc	3660
accatggggt	cagctattga	gcaggatggg	ttgcattgctg	gtagtcccgc	cgcattgggtc	3720
gaacgactgt	ttggatacga	ttgggccccaa	cagactatag	gctgttccga	cgtgctgtc	3780
tttcgtcttt	ctgcacaagg	tcgtccagtt	ctgttcgtga	aaaccgactt	gtccggagcc	3840
ctcaatgagt	tgcaagacga	agctgcacga	ctgagttggc	ttgccaccac	tggtgtccca	3900
tgtgccgcag	tacttgacgt	cgtcacagag	gctggtcgcg	attggttgct	ccttgagaaa	3960
gtgcccggcc	aagatcttct	cagttcccac	cttgcccctg	ccgaaaaagt	ttcaataatg	4020
gctgacgcta	tgagaaggct	gcacaccctt	gaccctgcca	catgtccatt	cgatcaccaa	4080
gcaaacacc	gaattgaacg	agctagaacc	cgcattggaag	ccggcctcgt	tgatcaagac	4140
gatttggatg	aggaacacca	gggtctcgca	cccgtgaaac	tcttcgctcg	cctcaaagca	4200
cgaatgccag	acggagatga	cttggctgta	accacggag	atgcctgcct	tcctaacata	4260
atggtagaga	atggaagatt	tageggcttc	attgattgtg	gacgacttgg	agttgcagat	4320
cgttaccag	atatcgtctt	cgctaccaga	gatattgctg	aagaattggg	cggagaatgg	4380
gctgatcggg	ttctcgtact	ctacggaatt	gcgcacactg	attcccacgc	cattgctttt	4440
taccgtcttc	tggatgagtt	cttctaaacg	cgccccctct	ctccctcccc	ccccctaac	4500
gttactggcc	gaagccgctt	ggaataaggc	cgggtgtcgt	ttgtctatat	gttattttcc	4560
accatattgc	cgtcttttgg	caatgtgagg	gcccggaaac	ctggccctgt	cttcttgacg	4620
agcattccta	ggggctcttc	ccctctcgcc	aaaggaatgc	aaggctctgt	gaatgtcgtg	4680
aaggaagcag	ttctctgga	agcttcttga	agacaaacaa	ctctctatgc	gaccctttac	4740

agggaagcag	cccccccgga	agcccccgga	agccccccaa	cgcccccgga	gccccccgga	4740
aggcagcggg	acccccccacc	tgggcgacagg	tgccctctgcg	gccaaaagcc	acgtgtataa	4800
gatacacctg	caaagggcggc	acaacccccag	tgcccacgttg	tgagttggat	agttgtggaa	4860
agagtcaaat	ggctctcctc	aagcgtatcc	aacaaggggc	tgaaggatgc	ccagaaggta	4920
ccccattgta	tgggatctga	tctggggcct	cggtgcacat	gctttacatg	tgtttagtcg	4980
aggttaaaaa	acgtctaggc	ccccggaacc	acggggacgt	ggttttcctt	tgaaaaacac	5040
gattgctcga	atcaccatgg	tgagcaaggg	cgaggagctg	ttcaccgggg	tggtgccccat	5100
cctggtegag	ctggacggcg	acgtaaacgg	ccacaagttc	agcgtgtccg	gcgagggcga	5160
gggcgatgcc	acctacggca	agctgaccct	gaagttcatc	tgaccaccg	gcaagctgcc	5220
cgtgccctgg	cccaccctcg	tgaccacctt	cggtacggc	ctgcagtget	tcgcccgcta	5280
ccccgaccac	atgaagcagc	acgacttctt	caagtccgcc	atgcccgaag	gctacgtcca	5340
ggagcgcacc	atctttctca	aggacgacgg	caactacaag	acccgcgccg	aggtgaagtt	5400
cgagggcgac	accctgggtga	accgcatcga	gctgaagggc	atcgacttca	aggaggacgg	5460
caacatcctg	gggcacaagc	tgaggtacaa	ctacaacagc	cacaacgtct	atatcatggc	5520
cgacaagcag	aagaacggca	tcaaggtgaa	cttcaagatc	cgccacaaca	tcgaggacgg	5580
cagcgtgcag	ctcgccgacc	actaccagca	gaacaccccc	atcgggcgacg	gccccgtgct	5640
gctgccccgac	aaccactacc	tgagctacca	gtccgcctg	agcaaagacc	ccaacgagaa	5700
gcgcgatcac	atggctctgc	tgagttcgt	gaccgcgcc	gggatcactc	tcggcatgga	5760
cgagctgtac	aagtaatcgg	ccgctaatca	gccataccac	atttgtagag	gttttacttg	5820
ctttaaaaaa	cctcccacac	ctccccctga	acctgaaaca	taaaatgaat	gcaattgttg	5880
ttgttaactt	gtttattgca	gcttataatg	gttacaaata	aagcaatagc	atcaciaaatt	5940
tcacaaataa	agcatttttt	tcactgcatt	ctagttgtgg	tttgtccaaa	ctcatcaatg	6000
tatcttatca	tgctggcgcg	ttgacattga	ttattgacta	gttattaata	gtaatcaatt	6060
acggggatcat	tagttcatag	cccatatatg	gagttccgcg	ttacataact	tacggtaaatt	6120
ggccccgctg	gctgaccgcc	caacgacccc	cgccccattga	cgtaataat	gacgtatggt	6180
cccatagtaa	cgccaatagg	gactttccat	tgacgtcaat	gggtggagta	tttacggtaa	6240
actgcccact	tggcagtaca	tcaagtgtat	catatgccaa	gtacgcccc	tattgacgtc	6300
aatgacggta	aatggcccgc	ctggcattat	gcccagtaca	tgaccttatg	ggactttcct	6360
acttggcagt	acatctacgt	attagtcatc	gctattacca	tggtgatgcg	gttttggcag	6420
tacatcaatg	ggcgtggata	gcggtttgac	tcacggggat	ttccaagtct	ccaccccatt	6480
gacgtcaatg	ggagtttggt	ttggcaccaa	aatcaacggg	actttccaaa	atgtcgtaac	6540
aactccgccc	cattgacgca	aatgggcggg	aggcgtgtac	gggtggagggt	ctatataagc	6600
agagctctcc	ctatcagtga	tagagatctc	cctatcagtg	atagagatcg	tcgacgttta	6660
gtgaaccgtc	agatcgccctg	gagacgccat	ccacgctggt	ttgacctcca	tagaagacac	6720
cgggaccgat	ccagcctccg	cggccgggaa	cggtgcattg	gaacgcggat	tccccgtgcc	6780
aagagtgcag	taagtaccgc	ctatagagtc	tataggccca	ccccctggc	ttcttatgca	6840

tgctatactg tttttggcct ggggtctata ccccccgct tctcatggt ataggtgatg 6900
 gtatagetta gcctataggt gtgggttatt gaccattatt gaccactecc ctattggtga 6960
 cgatacttcc cattaactaat ccataacatg gctctttgcc acaactctct ttattggcta 7020
 tatgccaata cactgtcctt cagagactga cacggactct gtatTTTTac aggatggggg 7080
 ctcatttatt atttacaat tcacatatac aacaccaccg tccccagtgc ccgcagtttt 7140
 tattaacat aacgtgggat ctccacgca atctcgggta cgtgttcagg acatggtctc 7200
 ttctccggta gcggcgggag ttctacatcc gagccctgct cccatgcctc cagcgactca 7260
 tggtcgctcg gcagctcctt gctcctaaca gtggaggcca gacttaggca cagcacgatg 7320
 cccaccacca ccagtgtgcc gcacaaggcc gtggcggtag ggtatgtgtc tgaaaatgag 7380
 ctccggggagc gggcttgca cgtgacgca tttggaagac ttaaggcagc ggcagaagaa 7440
 gatgcaggca gctgagttgt tgtgttctga taagagtcag aggtaactcc cgttgccggtg 7500
 ctgttaacgg tggagggcag tgtagtctga gcagtactcg ttgctgcgc gcgcgccacc 7560
 agacataata gctgacagac taacagactg ttcctttcca tgggtctttt ctgcagtcac 7620
 cgtccttgac acg 7633

<210> 45

<211> 1011

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 45

gaggtgcagc tgggtggagtc tggggggagcc atagtaaagc cggggggggtc ccatagagtc 60
 tctgtgaag cctctggatt cactttcagt aacgcctgga tgagttgggt ccgccaggct 120
 ccagggaggg ggctggagtg ggttggcctt attttaagca agactgatgg tgggacgaca 180
 gactacgctg caccogtgaa agacagattc accatttcaa gagatgattc taaaaatatg 240
 ttgtttctgc aaatggacag cctgaaaatc gaggacacag ccgtgtattt ctgtaccacg 300
 gccgattttt ggagtgetta ttcttctgac tactggggcc aggaaccct ggtcaccgtc 360
 tctcaggag gtggagggtc cggggggcgg ggctccggcg gaggtggatc agatattgtg 420
 atgactcagt ctccactctc cctgcccgtc acccctggag agccggcctc catctcctgc 480
 aggtctagtc agagcctct gcatagtaat gggtaaacct atttggattg gtacctacag 540
 aagccagggc agtctccaca actcctgac tatttgggtt ctaatcgggc ctccggggtc 600
 cctgacaggt tcagtggcag tggatcaggc acagatttta cactgaaaat cagcagaatg 660
 gaggtcagg atgttggggg ttattactgc atgcaaggtc taaaactcc gtacactttt 720
 ggccagggga ccaagctgga gatcaaagga ggcggaggga gtgttttgtt ttatctggcc 780
 gttgggataa tgtttctcgt aaatacagta ctttgggtaa caataaggaa ggaactgaag 840
 agaaagaaaa aatgggatct ggaaatatca ttggacagtg gacacgaaaa aaaagtcaca 900
 tcatcattcc aagaagaccg ccaattccag gaggaaactga aatgtcaaga gcaaaaagaa 960

ccaccattgc aagaagaccg gcactctggag gagggaactga aatgctcaaga gcaaaaagaa 500
 gaacaactgc aagaaggcgt acatagaaaa gaaccacagg gagcaacata g 1011

- <210> 46
- <211> 2669
- <212> DNA
- <213> Artificial sequence

- <220>
- <223> synthetic

<400> 46
 agcctaggcc tccaaaaaag cctcctcact acttctggaa tagctcagag gcagaggcgg 60
 cctcggcctc tgcataaata aaaaaatta gtcagccatg gggcggagaa tgggcggaac 120
 tgggcggagt taggggcggg atgggcggag ttaggggcgg gactatggtt gctgactaat 180
 tgagatgcat gctttgcata cttctgcctg ctggggagcc tggggacttt ccacacctgg 240
 ttgctgacta attgagatgc atgctttgca tacttctgcc tgctggggag cctggggact 300
 ttccacaccg gatccaccat gggttcagct attgagcagg atgggttgca tgctggtagt 360
 cccgcgcgat gggtcgaacg actgtttggg tacgattggg cccaacagac tataggctgt 420
 tccgacgctg ctgtctttcg ttttctgca caaggtcgtc cagttctggt cgtgaaaacc 480
 gacttgctcc gagccctcaa tgagttgcaa gacgaagctg cacgactgag ttggettgcc 540
 accactggtg tcccatgtgc cgcagtaact gacgtcgtca cagaggctgg tcgcgattgg 600
 ttgctccttg gagaagtgcc cggccaagat cttctcagtt ccacacttgc cctgcccga 660
 aaagtttcaa taatggetga cgtatgaga aggetgcaca ccettgacct tgccacatgt 720
 ccattcgatc accaagccaa acaccgaatt gaacgagcta gaaccgcgat ggaagccggc 780
 ctgcttgatc aagacgattt ggatgaggaa caccagggtc tcgcaccogc tgaactcttc 840
 gctcgcctca aagcacgaat gccagacgga gatgacttgg tcgtaacca cggagatgcc 900
 tgccttccta acataatggt agagaatgga agatttagcg gcttcattga ttgtggacga 960
 cttggagttg cagatcggta ccaagatatc gctctcgcta ccagagatat tgctgaagaa 1020
 ttgggcggag aatgggetga tgggtttctc gtaactctac gaattgcgc acctgattcc 1080
 caacgcattg ctttttaccg tcttctggat gagttcttct aaacgcgtcc cccctctccc 1140
 tcccccccc ctaacgttac tggccgaagc cgcttggaat aaggccgggtg tgcgtttgtc 1200
 tatatgttat tttccaccat attgccgtct tttggcaatg tgagggcccc gaaacctggc 1260
 cctgtcttct tgacgagcat tctaggggtt ctttccctc tcgccaagg aatgcaagg 1320
 ctggtgaatg tcgtgaagga agcagttcct ctggaagctt cttgaagaca aacaacgtct 1380
 gtagegacct tttgcaggca gcggaacccc ccacctggcg acaggtgcct ctgcggccaa 1440
 aagccaogtg tataagatac acctgcaaag gcggeacaac cccagtgcc cgttgtgagt 1500
 tggatagttg tggaaagagt caaatggctc tectcaagcg tattcaaca ggggetgaag 1560
 gatgccca gaaggtaccca ttgtatggga tctgatctgg ggctcgggtg cacatgcttt 1620
 1680

acatgtggtt agtcgagggt aaaaaacgtc taggcccccc gaaccacggg gacgtgggtt 1680
 tcctttgaaa aacacgattg ctggaatcac catggtgagc aagggcgagg agctgttcac 1740
 cggggtggtg cccatcctgg tcgagctgga cggcgacgta aacggccaca agttcagcgt 1800
 gtccggcgag ggcgagggcg atgccaccta cggcaagctg acctgaagt tcatctgcac 1860
 cafcggcaag ctgcccgtgc cctggccccac cctcgtgacc accttcgggt acggcctgca 1920
 gtgcttcgcc cgctaccccg accacatgaa gcagcagcag ttcttcaagt ccgccatgcc 1980
 cgaaggctac gtccaggagc gcaccatctt ctccaaggac gacggcaact acaagaccgg 2040
 cgccgagggt aagttcgagg gcgacaccct ggtgaaccgc atcgagctga agggcctega 2100
 ctccaaggag gacggcaaca tctgggggca caagctggag tacaactaca acagccacaa 2160
 cgtctatata atggccgaca agcagaagaa cggcatcaag gtgaacttca agatccgcca 2220
 caacatcgag gacggcagcg tgcagctcgc cgaccactac cagcagaaca ccccatcgg 2280
 cgacggcccc gtgctgctgc ccgacaacca ctacctgagc taccagtcgg cctgagcaa 2340
 agaccccaac gagaagcgcg atcacatggt cctgctggag ttctgtgacc cgcccggtat 2400
 cactctcggc atggacgagc tgtacaagta atcggccgct aatcagccat accacatttg 2460
 tagaggtttt acttgcttta aaaaacctcc cacacctccc cctgaacctg aaacataaaa 2520
 tgaatgcaat tgttgttggt aacttgctta ttgcagctta taatggttac aaataaagca 2580
 atagcatcac aaatttcaca aataaagcat ttttttact gcattctagt tgtggtttgt 2640
 ccaaactcat caatgtatct tatcatgtc 2669

<210> 47

<211> 2992

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 47

gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
 gccatataat ggagttccgc gttacataac ttacggtaaa tgccccgctt ggctgaccgc 120
 ccaacgacct ccgcccattg acgtcaataa tgacgtatgt tcccatagta acgccaatag 180
 ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccac ttggcagtac 240
 atcaagtgta tcatatgcca agtacgcccc ctattgacgt caatgacggt aaatggcccc 300
 cctggcatta tgcccagtac atgaccttat gggactttcc tacttggcag tacatctacg 360
 tattagtcac cgctattacc atggtgatgc ggttttggca gtacatcaat gggcgtggat 420
 agcggtttga ctacggggga tttccaagtc tccaccccat tgacgtcaat gggagtttgt 480
 tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
 aaatggggcg taggcgtgta cgggtggagg tctatataag cagagctctc cctatcagtg 600
 atagagatct ccctatcagt gatagagatc gtcgacgttt agtgaaccgt cagatcgctt 660

ggagacgcca	tccacgctgt	tttgacctcc	atagaagaca	ccgggaccga	tccagcctcc	720
gcggccggga	acgggtgcatt	ggaacgogga	ttccccgtgc	caagagtgac	gtaagtaccg	780
cctatagagt	ctataggccc	acccccttgg	cttcttatgc	atgctatact	gtttttggct	840
tggggctctat	acacccccgc	ttcctcatgt	tataggtgat	ggtatagctt	agcctatagg	900
tgtgggttat	tgaccattat	tgaccactcc	cctattgggtg	acgatacttt	ccattactaa	960
tccataacat	ggctctttgc	cacaactctc	tttattggct	atatgccaat	acactgtcct	1020
tcagagactg	acacggactc	tgtattttta	caggatgggg	tctcatttat	tatttacaaa	1080
ttcacatata	caacaccacc	gtccccagtg	cccgagttt	ttattaaaca	taacgtggga	1140
tctccacgcg	aatctcgggt	acgtgttccg	gacatggctc	cttctccggt	agcggcggag	1200
cttctacatc	cgagccctgc	tcccatgcct	ccagcagctc	atggtcgctc	ggcagctcct	1260
tgctcctaac	agtggaggcc	agacttaggc	acagcacgat	gccaccacc	accagtgtgc	1320
cgcacaaaggc	cgtggcggta	gggtatgtgt	ctgaaaatga	gctcggggag	cgggcttgca	1380
ccgctgacgc	atltggaaga	cttaaggcag	cggcagaaga	agatgcaggc	agctgagttg	1440
ttgtgttctg	ataagagtca	gaggtaactc	ccgttgcggt	gctgttaacg	gtggagggca	1500
gtgtagtctg	agcagtactc	gttgctgccg	cgcgcgccac	cagacataat	agctgacaga	1560
ctaacagact	gttcccttcc	atgggtcttt	tctgcagaag	cttatactcg	agetctagat	1620
tgggaaccgg	ggtctctcga	attecagatc	tccaccatgc	acagacctag	acgtcgtgga	1680
actcgtccac	ctccactggc	actgctcgct	gctctcctcc	tggtgcacg	tggtgctgat	1740
gcagaggtgc	agctggtgga	gtctggggga	gccatagtaa	agccgggggg	gtcccataga	1800
gtctcctgtg	aagcctctgg	attcactttc	agtaacgcct	ggatgagttg	ggtccgccag	1860
gctccagggg	gggggctgga	gtgggttggc	cgtattttaa	gcaagactga	tggtgggacg	1920
acagactacg	ctgcacccgt	gaaagacaga	ttcaccattt	caagagatga	ttctaaaaat	1980
atgttgtttc	tgcaaatgga	cagcctgaaa	atcgaggaca	cagccgtgta	tttctgtacc	2040
acggccgatt	tttggagtgc	ttattcttct	gactactggg	gccagggaac	cctggtcacc	2100
gtctcctcag	gaggtggagg	ttccgggggc	gggggctccg	gcggaggtgg	atcagatatt	2160
gtgatgactc	agtctccact	ctccctgccc	gtcacccctg	gagagccggc	ctccatctcc	2220
tgcaggtcta	gtcagagcct	cctgcatagt	aatgggtaca	actatlttga	ttggtacct	2280
cagaagccag	ggcagtctcc	acaactcctg	atctatlttg	gttctaactg	ggcctccggg	2340
gtccctgaca	ggttcagtgg	cagtggatca	ggcacagatt	ttacactgaa	aatcagcaga	2400
atggaggtctg	aggatgttgg	ggtttattac	tgcattgcaag	gtctacaaac	tccgtacact	2460
tttggccagg	ggaccaagct	ggagatcaaa	ggaggcggag	ggagtgtttt	gttttatctg	2520
gccgttgga	taatgtttct	cgtaaataca	gtactlttggg	taacaataag	gaaggaactg	2580
aagagaaaga	aaaaatggga	tctggaaata	tcattggaca	gtggacacga	aaaaaaagtc	2640
acatcatcat	tgcaagaaga	ccggcacttg	gaggaggaac	tgaaatgtca	agagcaaaaa	2700
gaagaacaac	tgcaagaagg	cgtacataga	aaagaaccac	aggagcaac	ataggcggcc	2760
gctaatacagc	cataccacat	ttgtagaggt	tttacttgcct	ttaaaaaac	tcccacacct	2820

cccctgaac ctgaaacata aatgaatgc aattggtgtt gttacttgt ttattgcagc 2880
 ttataatggt tacaaataaa gcaatagcat cacaaatttc acaataaag catttttttc 2940
 actgcattct agttgtgggt tgcctaaaact catcaatgta tcttatcatg tc 2992

<210> 48

<211> 5765

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 48

acaacttcgt atagcataca ttatacgaag ttatggtacc aagcctaggc ctccaaaaaa 60
 gcctcctcac tacttctgga atagctcaga ggcagaggcg gcctcggcct ctgcataaat 120
 aaaaaaaatt agtcagccat ggggcggaga atgggcggaa ctgggcggag ttaggggcgg 180
 gatgggcgga gttagggcg ggactatggt tgetgactaa ttgagatgca tgetttgcat 240
 acttctgcct gctggggagc ctggggactt tccacacctg gttgctgact aattgagatg 300
 catgctttgc ataacttctgc ctgctgggga gcctggggac tttccacacc ggatecacca 360
 tgggttcagc tattgagcag gatgggttgc atgctggtag tcccgccgca tgggtcgaac 420
 gactgtttgg atacgattgg gcccaacaga ctataggctg ttccgacgct gctgtctttc 480
 gtctttctgc acaaggtcgt ccagttctgt tegtgaaaac cgacttgtcc ggagccctca 540
 atgagttgca agacgaaget gcacgactga gttggcttgc caccactggt gteccatgtg 600
 ccgcagtaet tgacgtcgtc acagaggctg gtcgcgattg gttgctcctt ggagaagtgc 660
 ccggccaaga tcttctcagt tcccacctg cccctgccga aaaagtttca ataatggtc 720
 acgctatgag aaggctgcac acccttgacc ctgccacatg tccattcgat caccaagcca 780
 aacaccgaat tgaacgaget agaaccgcga tggaagccgg cctcgttgat caagacgatt 840
 tggatgagga acaccagggt ctgcaccccg ctgaactctt cgetcgcctc aaagcacgaa 900
 tgccagacgg agatgacttg gtcgtaaacc acggagatgc ctgccttctt aacataatgg 960
 tagagaatgg aagatttagc ggcttcattg attgtggacg acttgaggtt gcagatcgg 1020
 accaagatat cgtctctcgt accagagata ttgctgaaga attgggcgga gaatgggctg 1080
 ateggtttct cgtactctac ggaattgccc cacctgatte ccaacgcatt gctttttacc 1140
 gtcttctgga tgagttcttc taaacgcgtc cccctctcc ctccccccc cctaacgta 1200
 ctggccgaag ccgcttgga taaggccgg gtgcggttgt ctatatgtta tttccacca 1260
 tattgcggtc ttttggcaat gtgagggccc ggaaacctgg ccctgtcttc ttgacgagca 1320
 ttctagggg tctttcccct ctgcctaaaag gaatgcaagg tctggtgaat gtcgtgaagg 1380
 aagcagttcc tctggaaget tcttgaagac aaacaacgtc tgtagcgacc ctttgcagc 1440
 ageggaacct cccacctggc gacaggtgcc tctgcggcca aaagccaagt gtataagata 1500
 cacctqcaaa ggcggcacia cccaagtccc acqttgtgag ttgqataqtt gtqgaaagag 1560

tcaa	atgg	ctcc	taag	gtatt	caaca	agggg	ctgaa	ggatg	cccag	aaggt	acccc	1620
attg	tatg	gg	ctc	gctg	gg	gctc	gg	gcac	atg	tacat	gtg	1680
taaaa	aac	ctag	gcccc	cga	acc	acg	gg	ga	ctg	ttc	cttt	1740
gctc	gaat	ccat	ggt	ca	aggg	gc	gag	gag	ctg	ccg	ggg	1800
gtc	gag	ac	ggc	gac	aa	ac	gg	cc	ac	aag	ttc	1860
gatg	cc	ac	ct	gac	g	ac	ct	gaa	tt	cat	ctg	1920
cc	ct	gg	cc	ca	cc	ct	gac	ca	ct	ct	gac	1980
gac	ca	at	ga	ag	cag	ca	ca	g	ct	ct	ca	2040
cg	ca	ca	ct	ct	ca	ag	ga	ca	ac	tac	aa	2100
gg	ca	ac	cc	tg	gt	ga	ac	cc	g	at	g	2160
at	ct	g	gg	gc	aca	ag	ct	g	ga	gt	g	2220
aag	caga	aga	ac	gg	ca	tcaa	g	gt	g	aa	ct	2280
gt	g	ca	g	ct	g	ac	ca	ac	cc	ca	ct	2340
cc	g	ca	ac	ac	ta	ct	g	ag	ct	cc	ca	2400
gat	ca	ca	tg	g	ct	g	ct	g	ga	ca	g	2460
ct	g	ta	ca	ag	aa	tc	g	cc	gc	ta	ac	2520
aaaa	ac	ct	cc	cc	ct	ga	ac	ct	g	aa	ca	2580
ta	ac	tt	g	tt	g	tt	g	tt	g	tt	g	2640
aa	ta	aa	g	ca	tt	tt	tt	tt	tt	tt	tt	2700
tt	at	ca	tg	ct	g	tt	g	tt	g	tt	g	2760
gg	ta	ca	tg	ct	g	tt	g	tt	g	tt	g	2820
cg	ct	g	gg	ct	g	ct	g	ct	g	ct	g	2880
tag	ta	ac	gc	ca	aa	ta	gg	g	ac	tt	tt	2940
cc	ca	ct	tg	gc	ca	ta	ca	gt	g	ta	ca	3000
ac	gg	ta	aa	tg	g	cc	gc	ct	g	g	ct	3060
gg	ca	g	ta	ca	ta	ct	g	ta	ca	ta	ct	3120
tca	at	gg	gc	tg	ga	ta	gc	gg	tt	tt	g	3180
tca	at	gg	gc	tg	ga	ta	gc	gg	tt	tt	g	3240
cc	g	cc	cc	ca	at	g	gg	gc	g	g	g	3300
ct	ct	cc	ct	at	ca	g	at	g	at	g	at	3360
ac	gc	ta	gc	at	gc	gg	gc	g	g	g	g	3420
ac	gc	ta	gc	at	gc	gg	gc	g	g	g	g	3480
gt	g	ac	g	ta	gc	ta	gc	gg	gc	g	g	3540
at	ac	tt	g	tt	g	tt	g	tt	g	tt	g	3600
ag	ct	ta	gc	at	gc	gg	gc	g	g	g	g	3660

actttccatt	actaatccat	aacatggctc	tttgccacaa	ctctctttat	tggctatatg	3720
ccaatacact	gtccttcaga	gactgacacg	gactctgtat	ttttacagga	tggggctctca	3780
tttattat	acaaattcac	atatacaaca	ccaccgtccc	cagtgccccg	agtttttatt	3840
aaacataacg	tgggatctcc	acgcgaatct	cgggtacgtg	ttccggacat	ggtctcttct	3900
cggtagcgg	cggagcttct	acatccgagc	cctgctccca	tgctccagc	gactcatggt	3960
cgctcggcag	ctccttgctc	ctaacagtgg	aggccagact	taggcacagc	acgatgccca	4020
ccaccaccag	tgtgcgcgac	aaggccgtgg	cggtagggta	tgtgtctgaa	aatgagctcg	4080
gggagcgggc	ttgcaccgct	gacgcatttg	gaagacttaa	ggcagcggca	gaagaagatg	4140
caggcagctg	agttgttg	ttctgataag	agtcagaggt	aactcccgtt	gcgggtgctgt	4200
taacgggtga	gggcagtgta	gtctgagcag	tactcgttgc	tgccgcgcgc	gccaccagac	4260
ataatagctg	acagaactaac	agactgttcc	tttccatggg	tctttctctc	agtcaccgctc	4320
cttgacacga	agcttatact	cgagctctag	attgggaacc	cgggtctctc	gaattcgaga	4380
tctccaccat	gcacagacct	agacgtcgtg	gaactcgtcc	acctccactg	gcactgctcg	4440
ctgctctcct	cctggctgca	cgtgggtgctg	atgcagaggt	gcagctgggtg	gagtctgggg	4500
gagccatagt	aaagccgggg	gggtcccata	gagtctcctg	tgaagcctct	ggattcactt	4560
tcagtaacgc	ctggatgagt	tgggtccgcc	aggetccagg	gagggggctg	gagtggggtg	4620
gccgtat	aagcaagact	gatgggtgga	cgacagacta	cgtgcaccc	gtgaaagaca	4680
gattcaccat	ttcaagagat	gattctaaaa	atatgttggt	tctgcaaatg	gacagcctga	4740
aaatcgagga	cacagccgtg	tattttctgta	ccacggccga	tttttgagat	gcttattctt	4800
ctgactactg	gggccaggga	acctgggtca	ccgtctctct	aggaggtgga	ggttccgggg	4860
gcgggggctc	cggcggaggt	ggatcagata	ttgtgatgac	tcagtctcca	ctctccctgc	4920
ccgtcacccc	tggagagccg	gcctccatct	cctgcaggte	tagtcagagc	ctcctgcata	4980
gtaatgggta	caactat	gattggtacc	tacagaagcc	agggcagtct	ccacaactcc	5040
tgatctat	gggttcta	cgggcctc	gggtccctga	caggttcagt	ggcagtggat	5100
caggcacaga	ttttacactg	aaaatcagca	gaatggaggc	tgaggatggt	ggggtttatt	5160
actgcatgca	aggtctacaa	actccgtaca	cttttgcca	ggggaccaag	ctggagatca	5220
aaggaggcgg	agggagtgtt	ttgttttata	tggccgttgg	gataatg	ctcgtaaata	5280
cagtactttg	ggtaacaata	aggaaggaa	tgaagagaaa	gaaaaaatgg	gatctggaaa	5340
tatcattgga	cagtggacac	gaaaaaaaa	tcacatc	attgcaagaa	gaccggcact	5400
tggaggagga	actgaaatgt	caagagcaaa	aagaagaaca	actgcaagaa	ggcgtacata	5460
gaaaagaacc	acagggagca	acataggcgg	ccgctaatca	gccataccac	at	5520
gttttacttg	ctttaaaaaa	cctcccacac	ctccccctga	acctgaaaca	taaaatgaat	5580
gcaattggtg	ttgttaactt	gtttattgca	gcttataatg	gttacaata	aagcaatagc	5640
atcaciaaatt	tcaciaaata	agcatttttt	tactgcatt	ctagttgtgg	tttgtccaaa	5700
ctcatcaatg	tatcttatca	tgtctaccgg	tataacttcg	tataatgtat	actatacgaa	5760
gtag						5765

<210> 49
 <211> 660
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 49
 gacatcgtga tgacccagtc tccactctcc ctgcccgtca cccttgaga gccggcctcc 60
 atctcctgca ggtctagtea gagcctcctg catagtaatg ggtacaacta tttggattgg 120
 tacctacaga agccagggca gtctccaca ctctgatct atttgggttc taatcgggcc 180
 tccggggctc ctgacagggt cagtggcagt ggatcaggca cagattttac actgaaaate 240
 agcagaatgg aggctgagga tgttgggggt tattactgca tgcaaggctt acaaaactccg 300
 tacacttttg gccaggggac caagctggag atcaaacgag ctgatgctgc accaactgta 360
 tccatcttcc caccatccag tgagcagtta acatctggag gtgcctcagt cgtgtgcttc 420
 ttgaacaact tctaccccaa agacatcaat gtcaagtgga agattgatgg cagtgaacga 480
 caaaatggcg tctgaacag ttggactgat caggacagca aagacagcac ctacagcatg 540
 agcagcacc ctcacgttgac caaggacgag tatgaacgac ataacagcta tacctgtgag 600
 gccactcaca agacatcaac ttcacccatt gtcaagagct tcaacagggg agagtgttga 660

<210> 50
 <211> 1359
 <212> DNA
 <213> Artificial sequence

<220>
 <223> synthetic

<400> 50
 gaggtgcagc tgggtggagtc tgggggagcc atagtaaagc cggggggggt ccatagagtc 60
 tctgtggaag cctctggatt cactttcagt aacgcctgga tgagttgggt ccgccaggct 120
 ccagggaggg ggctggagtg ggttggccgt attttaagca agactgatgg tgggacgaca 180
 gactacgctg caccctgaa agacagattc accatttcaa gagatgattc taaaaatag 240
 ttgtttctgc aaatggacag cctgaaaatc gaggacacag ccgtgtattt ctgtaccacg 300
 gccgattttt ggagtgetta ttcttctgac tactggggcc agggaaccct ggtcacctgc 360
 tctcagcca aaacaacagc cccatcggtc tateactgg ccctgtgtg tggagataca 420
 actggetcct cggtgactct aggatgcctg gtcaagggtt atttccctga gccagtgacc 480
 ttgacctgga actctggate cctgtccagt ggtgtgcaca ccttcccagc tgtcctgcag 540
 tetgacctct acacctcag cagctcagt actgtaacct cgagcacctg gccagccag 600
 tccatcacct gcaatgtggc ccacccggca agcagcacca aggtggacaa gaaaattgag 660

cccagagggc	ccacaatcaa	gccctgtcct	ccatgcaa	gcccagcaec	taacctcttg	720	
ggtggaccat	ccgtcttcat	cttccctcca	aagatcaagg	atgtactcat	gatctccctg	780	
agccccatag	tcacatgtgt	ggtgggtgat	gtgagcgagg	atgaccacaga	tgtccagatc	840	
agctggtttg	tgaacaacgt	ggaagtacac	acagctcaga	cacaaaccca	tagagaggat	900	
tacaacagta	ctctccgggt	ggtcagtgcc	ctccccatcc	agcaccagga	ctggatgagt	960	
ggcaaggagt	tcaa	aatgcaa	ggtcaacaac	aaagacctcc	cagcgcccat	cgagagaacc	1020
atctcaaaac	ccaaagggtc	agtaagagct	ccacaggtat	atgtcttgcc	tccaccagaa	1080	
gaagagatga	ctaagaaaca	ggtcactctg	acctgcatgg	tcacagactt	catgcctgaa	1140	
gacatttacg	tggagtggac	caacaacggg	aaaacagagc	taaactacaa	gaacactgaa	1200	
ccagtctctg	actctgatgg	ttcttacttc	atgtacagca	agctgagagt	ggaaaagaag	1260	
aactgggtgg	aaagaaatag	ctactcctgt	tcagtgggcc	acgaggggtct	gcacaatcac	1320	
cacacgacta	agagcttctc	cggactccg	ggtaaatga			1359	

<210> 51

<211> 1011

<212> DNA

<213> Artificial sequence

<220>

<223> synthetic

<400> 51

gaggtgcage	tgggtggagtc	tgggggagcc	atagtaaage	cggggggggtc	ccatagagtc	60
tctgtgaag	cctctggatt	cacttctcagt	aacgcctgga	tgagttgggt	ccgccaggct	120
ccagggaggg	ggctggagtg	ggttggccgt	atthtaagca	agactgatgg	tgggacgaca	180
gactacgctg	caccctgaa	agacagattc	accatttcaa	gagatgattc	taaaaatag	240
ttgtttctgc	aatggacag	cctgaaaatc	gaggacacag	ccgtgtatth	ctgtaccacg	300
gccgatttht	ggagtgcctta	ttcttctgac	tactggggcc	aggaaccct	ggtcaccgtc	360
tcctcaggag	gtggaggttc	cgggggccc	ggctccggcg	gaggtggatc	agatattgtg	420
atgactcagt	ctccactctc	cctgcccgtc	accctggag	agccggcctc	catctcctgc	480
aggtctagtc	agagcctcct	gcatagtaat	gggtacaact	atthggattg	gtacctacag	540
aagccagggc	agtctccaca	actcctgatc	tatthgggtt	ctaategggc	ctccggggtc	600
cctgacaggt	tcagtggcag	tggatcaggc	acagatthta	cactgaaaat	cagcagaatg	660
gaggtctgag	atgthgggggt	ttattactgc	atgcaaggtc	tacaaactcc	gtacacttht	720
ggccagggga	ccaagctgga	gatcaaagga	ggcggaggga	gtgthttgtt	ttatctggcc	780
gttgggataa	tgtthctcgt	aaatacagta	ctthgggtaa	caataaggaa	ggaactgaag	840
agaaagaaaa	aatgggatct	ggaaatatca	ttggacagtg	gacacgaaaa	aaaagtcaca	900
tcatcattgc	aagaagaccg	gcacttggag	gaggaactga	aatgtcaaga	gcaaaaagaa	960
gaacaactgc	aagaaggcgt	acatagaaaa	gaaccacagg	gagcaacata	g	1011

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO02057423A [0004]
- US20200331527A [0004]
- US5635354A [0106]
- US6623957B [0106]
- US6080840A [0106]
- US7329731B [0106]
- US6927044B [0106]
- US7105348B [0158]
- US61726040B [0175]

Non-patent literature cited in the description

- MENG et al. Gene, 2000, vol. 242, 201- [0074]
- WEAVER et al. Methods Enzymol., 1990, vol. 2, 234- [0076]
- GRAY et al. J. Immunol. Methods, 1995, vol. 182, 155- [0076]
- MANZ et al. PNAS, 1995, vol. 92, 1921-1925 [0077]
- SAMBROOK, J.E. F. FRITSCHT. MANIATIS Molecular Cloning: A Laboratory Manual 1989 0000 vol. 1-3, [0084]
- Current Protocols in Molecular Biology Greene Publ. Assoc., Wiley Interscience [0084]
- LEFRANC LEFRANC The T Cell Receptor Facts Book Academic Press 2001 0000 [0099]
- LEFRANC et al. IMGT unique numbering for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains Developmental and Comparative Immunology, 2003, vol. 27, 55-77 [0099]
- LEFRANC et al. IMGT unique numbering for immunoglobulin and T cell receptor constant domains and Ig superfamily C-like domains Developmental and Comparative

- Immunology, 2005, vol. 29, 185-203 [0099]
- **GENEVÉE et al.**An experimentally validated panel of subfamily-specific oligonucleotide primers (V α 1-w29/V β 1-w24) for the study of human T cell receptor variable V gene segment usage by polymerase chain reaction *Eur. J. Immunol.*, 1992, vol. 22, 1261-1269 [0106]
 - **GORSKI et al.**Circulating T Cell Repertoire Complexity in Normal Individuals and Bone Marrow Recipients Analyzed by CDR3 Size Spectratyping *J. Immunol.*, 1994, vol. 152, 5109-5119 [0106]
 - **JOHNSTON, S. et al.**A novel method for sequencing members of multi-gene families *Nucleic Acids Res.*, 1995, vol. 23/15, 3074-3075 [0106]
 - **PANNETIER et al.**T-cell repertoire diversity and clonal expansions in normal and clinical samples *Immunology Today*, 1995, vol. 16/4, 176-181 [0106]
 - **HINZ, T.KABELITZ, D.**Identification of the T-cell receptor alpha variable (TRAV) gene(s) in T-cell malignancies *J. Immunol. Methods*, 2000, vol. 246, 145-148 [0106]
 - *Leukemia*, vol. 17, 2257-2317 [0106]
 - **HODGES et al.**Diagnostic role of tests for T cell receptor (TCR) genes *J. Clin. Pathol.*, 2002, vol. 56, 1-11 [0106]
 - **MOYSEY, R. et al.**Amplification and one-step expression cloning of human T cell receptor genes *Anal. Biochem.*, 2004, vol. 326, 284-286 [0106]
 - **FERNANDES et al.**Simplified Fluorescent Multiplex PCR Method for Evaluation of the T-Cell Receptor V β -Chain Repertoire *Clin. Diag. Lab. Immunol.*, 2005, vol. 12/4, 477-483 [0106]
 - **LI, Y. et al.**Directed evolution of human T-cell receptors with picomolar affinities by phage display *Nature Biotech.*, 2005, vol. 23/3, 349-354 [0106]
 - **WLODARSKI et al.**Pathologic clonal cytotoxic T-cell responses: nonrandom nature of the T-cell receptor restriction in large granular lymphocyte leukemia *Blood*, 2005, vol. 106/8, 2769-2780 [0106]
 - **WLODARSKI et al.**Molecular strategies for detection and quantitation of clonal cytotoxic T-cell responses in aplastic anemia and myelodysplastic syndrome *Blood*, 2006, vol. 108/8, 2632-2641 [0106]
 - **BORIA et al.**Primer sets for cloning the human repertoire of T cell Receptor Variable regions *BMC Immunology*, 2008, vol. 9, 50- [0106]
 - **RICHMAN, S.KRANZ, D.**Display, engineering, and applications of antigen-specific T cell receptors *Biomolecular Engineering*, 2007, vol. 24, 361-373 [0106]
 - **LAUGEL, B et al.**Design of Soluble Recombinant T Cell Receptors for Antigen Targeting and T Cell Inhibition *J. Biol. Chem.*, 2005, vol. 280, 1882-1892 [0106]
 - **GUSS et al.** *EMBO*, 1986, vol. 5, 1567- [0128]
 - **SJOBRING et al.** *J. Biol. Chem.*, 1991, vol. 266, 399- [0128]

PATENTKRAV

1. Rekombinant antigenbindende protein, der binder et humant IgG1-Fc-domæne, et humant IgG2-Fc-domæne eller et humant IgG4-Fc-domæne, hvor:
 - 5 (a) det rekombinante antigenbindende protein er specifikt for et vildtype-CH3-domæne indeholdende en histidinrest på position 95 ifølge IMGT-exon-nummereringssystemet og en tyrosinrest på position 96 ifølge IMGT-exon-nummereringssystemet (Fc) over et substitueret CH3-domæne indeholdende en argininrest på position 95 ifølge IMGT-exon-nummereringssystemet og en
10 phenylalaninrest på position 96 ifølge IMGT-exon-nummereringssystemet (Fc*); og
 - (b) det rekombinante antigenbindende protein omfatter: (i) et ScFv, der omfatter et HCVR med aminosyresekvensen ifølge SEQ ID NO: 15 og et LCVR med aminosyresekvensen ifølge SEQ ID NO: 16; og (ii) et membrananker.
2. Rekombinant antigenbindende protein ifølge krav 1, hvor det
15 antigenbindende protein er et ScFv-fusionsprotein omfattende:
 - a. et membranforankringsdomæne omfattende aminosyresekvensen ifølge SEQ ID NO: 17 eller SEQ ID NO:21; eller
 - b. aminosyresekvensen ifølge SEQ ID NO: 19.
3. Isoleret polynukleotid, der omfatter en nukleinsyresekvens, der koder for:
 - 20 a. det antigenbindende protein ifølge krav 1 eller 2; eller
 - b. et polypeptid, der omfatter aminosyresekvensen ifølge SEQ ID NO: 19.
4. Nukleinsyrevektor, der omfatter:
 - (a) et polynukleotid ifølge krav 3;
 - (b) en promoter, der er operativt bundet til polynukleotidet; og
 - 25 (c) en polyadenyleringssekvens.
5. Nukleinsyrevektor ifølge krav 4, der endvidere omfatter en nukleinsyre, der koder for:
 - a. en valgbar markør eller

b. et energioverføringsprotein.

6. Nukleinsyrevektor ifølge kravene 4 eller 5, hvor:

a. promoteren er en CMV-promoter;

b. den valgbare markør tilvejebringer neomycinresistens; eller

5 c. energioverføringsproteinet er et derivat af grønt fluorescensprotein, såsom gult fluorescensprotein ("YFP").

7. Nukleinsyrevektor ifølge et hvilket som helst af kravene 4-6, hvor vektoren er cirkelformet eller lineær.

8. Nukleinsyrevektor ifølge et hvilket som helst af kravene 4-7, hvor vektoren
10 er integreret i en værtscelles genom og/eller værtscellen er en CHO-celle.

9. Værtscelle, der udtrykker et antigenbindende protein ifølge krav 1 eller 2.

10. Værtscelle ifølge krav 9, hvor cellen er en CHO-celle.

11. Fremgangsmåde til detektering eller isolering af en celle, der stabilt udtrykker et heterodimert protein, hvilken fremgangsmåde omfatter følgende trin:

15 (a) ekspression i en værtscelle af et celleoverflade-capture-protein (CSCP), der omfatter et antigenbindende protein ifølge krav 1 eller 2 og et heterodimert protein, hvor (i) CSCP'et binder til et første site på det heterodimere protein for at danne et CSCP-heterodimert proteinkompleks inde i værtscellen, (ii) det CSCP-heterodimere proteinkompleks transporteres gennem værtscellen og (iii) vises derefter på
20 værtscellens overflade;

(b) at bringe værtscellen i kontakt med et detekteringsmolekyle, hvor detekteringsmolekylet binder til et andet site på det heterodimere protein; og

(c) udvælgelse af værtscellen, der binder detekteringsmolekylet, hvor i det heterodimere protein:

25 (i) det første site på det heterodimere protein befinder sig på en første underenhed, der omfatter en tungkæde omfattende et vildtype CH3-domæne indeholdende en histidinrest på position 95 ifølge IMGT-exon-nummereringssystemet og tyrosinrest på position 96 ifølge IMGT-exon-nummereringssystemet (Fc); og

- (ii) det andet site på det heterodimere protein befinder sig på en anden underenhed, der omfatter en tungkæde omfattende et substitueret CH3-domæne indeholdende en argininrest på position 95 ifølge IMGT-exon-nummereringssystemet og en phenylalaninrest på position 96 ifølge IMGT-exon-nummereringssystemet (Fc*);
- 5 hvor det heterodimere protein omfatter et bispecifikt antistof eller et Fc-holdigt fusionsprotein.

12. Fremgangsmåde ifølge krav 10 eller 11, der omfatter trinnet med at bringe cellen i kontakt med et blokeringsmolekyle før udvælgelse af værtscellen på trin (c), hvor blokeringsmolekylet binder til CSCP, der ikke er bundet til det heterodimere protein, men ikke binder til det CSCP-heterodimere proteinkompleks.

10

13. Fremgangsmåde ifølge kravene 10 eller 11, hvor udvælgelsestrinnet (c) udføres ved fluorescensaktiveret celledatering.