



- (51) International Patent Classification:
C07K 14/705 (2006.01) A61K 38/17 (2006.01)
- (21) International Application Number:
PCT/US2016/059838
- (22) International Filing Date:
1 November 2016 (01.11.2016)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/249,836 2 November 2015 (02.11.2015) US
62/373,654 11 August 2016 (11.08.2016) US
- (71) Applicant: FIVE PRIME THERAPEUTICS, INC. [US/US]; Two Corporate Drive, South San Francisco, California 94080 (US).
- (72) Inventors: BRENNAN, Thomas; Two Corporate Drive, South San Francisco, California 94080 (US). BELLOVIN, David; Two Corporate Drive, South San Francisco, California 94080 (US). BUSH, David; Two Corporate Drive, South San Francisco, California 94080 (US). SENNINO,

Barbara; Two Corporate Drive, South San Francisco, California 94080 (US).

(74) Agents: DOHERTY, Elizabeth A. et al.; McNeill Baur PLLC, 125 Cambridge Park Drive, Suite 301, Cambridge, Massachusetts 02140 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,

[Continued on next page]

(54) Title: CD80 EXTRACELLULAR DOMAIN POLYPEPTIDES AND THEIR USE IN CANCER TREATMENT

(57) Abstract: This application relates to CD80 (B7-1) extracellular domain (ECD) polypeptides and CD80-ECD fusion molecules and their use in treatment of cancer, both alone and in combination with other therapeutic agents, such as immune stimulating agents such as PD-1/PD-L1 inhibitors.




Fig. 1A





TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

— *with sequence listing part of description (Rule 5.2(a))*

Published:

— *with international search report (Art. 21(3))*

CD80 EXTRACELLULAR DOMAIN POLYPEPTIDES AND THEIR USE IN CANCER TREATMENT

CROSS-REFERENCE TO RELATED APPLICATIONS

[001] This application claims priority to US Provisional Patent Application Nos. 62/373,654, filed August 11, 2016, and 62/249,836, filed November 2, 2015, both of which are incorporated herein by reference in their entirety.

FIELD

[002] This application relates to CD80 (B7-1) extracellular domain (ECD) polypeptides and CD80-ECD fusion molecules and their use in treatment of cancer, both alone and in combination with other therapeutic agents, such as immune stimulating agents such as PD-1/PD-L1 inhibitors.

BACKGROUND

[003] CD80, also known as B7-1, is one of the B7 family of membrane-bound proteins involved in immune regulation by delivering costimulatory or coinhibitory responses through their ligand binding activities. Other members of the B7 family of proteins include CD86 (B7-2), inducible costimulator ligand (ICOS-L), programmed death-1 ligand (PD-L1; B7-H1), programmed death-2 ligand (PD-L2; B7-H2), B7-H3, and B7-H4. CD80 is a transmembrane protein expressed on the surface of T cells, B cells, dendritic cells and monocytes, and binds to the receptors CD28, CTLA4 (CD152), and PD-L1. CD80 and CD86 and their receptors CTLA4 and CD28 operate as a costimulatory-coinhibitory system, for example, to control T cell activation, expansion, differentiation, and survival. CD80 and CD86 interaction with CD28 results in costimulatory signals that lead, for example, to activation of T cell responses. CD80, in turn, stimulates upregulation of CTLA4, which, upon binding to CD80, acts to suppress the T cell response previously triggered by CD80/CD28 interactions. This feedback loop allows for fine control of immune responses.

[004] CD80 has also been shown to interact with another B7 family member, PD-L1 with similar affinity to CD28, whereas CD86 does not interact with PD-L1. PD-

L1 is one of two ligands for the programmed death-1 (PD-1) protein, which is also involved in T cell regulation. Specifically, expression of PD-1 on T cells may be induced after T cells have been activated, and binding of PD-1 to PD-L1 downregulates T cell activity by promoting T cell inactivation. Many tumor cells express PD-L1 on their surface, potentially leading to PD-1/PD-L1 interactions and the inhibition of T cell responses against the tumor. This observation has led to the development of inhibitors of the PD-1/PD-L1 interaction as cancer therapeutics designed to stimulate natural immune responses against tumors in patients.

[005] Binding of CD80 to PD-L1 may serve as an alternative mechanism to block the PD-1/PD-L1 interaction and prevent inhibition of T cell responses at the site of a tumor. At the same time, however, increased levels of CD80 might also be available to bind to CD28 and to induce CTLA4, thus either inducing or inhibiting T cell responses. Some soluble forms of CD80 may also function to block CTLA4 activation by blocking endogenous CD80 activity. In addition, different soluble CD80 protein forms may have different effects on tumor growth through other interactions between the protein forms and tumor cells whose impact cannot be predicted in advance of testing. How various soluble forms of CD80 actually impact tumor growth *in vivo* has also not previously been directly tested. The present inventors have developed a set of CD80 extracellular domain (ECD) fusion molecules with particularly potent effects on tumor growth in a mouse model, both when administered alone, and when administered in conjunction with a PD-1/PD-L1 inhibitor. Based on the data shown in the working examples below, embodiments herein may provide superb therapeutic effects in cancer treatment.

SUMMARY

[006] In some embodiments, a CD80 extracellular domain (ECD) polypeptide or a CD80 ECD fusion molecule is provided. In some embodiments, the fusion molecule comprises a CD80 ECD and at least one fusion partner, comprising an Fc domain of an immunoglobulin, such as a human IgG1, IgG2, IgG3, or IgG4, albumin, or a polymer such as PEG. In some embodiments, the CD80 ECD or CD80 ECD fusion molecule comprises a human CD80, such as that of SEQ ID NO:5, or a human CD80 ECD from CD80 isoform 2 or isoform 3 (SEQ ID NOs: 3 and 4). In some embodiments, the

fusion molecule comprises an Fc domain, such as an Fc domain comprising a sequence selected from SEQ ID NOs: 9-16. In some embodiments, the fusion molecule comprises a human IgG1 Fc domain, such as one with a wild-type sequence such as that of SEQ ID NO:14, or alternatively, a mutant sequence with L234F, L235E, and P331S amino acid substitutions such as that of SEQ ID NO:12. In some embodiments, the CD80 ECD fusion molecule comprises an amino acid sequence selected from SEQ ID NO: 5, SEQ ID NO: 12, SEQ ID NO:14, SEQ ID NO: 20, and SEQ ID NO: 21. In some embodiments, the CD80 ECD fusion molecule fusion partner is directly attached to the C-terminal amino acid of the CD80 ECD amino acid sequence or to the N-terminal amino acid of the mature CD80 ECD amino acid sequence. In some embodiments, the CD80 ECD fusion molecule may be attached to the CD80 ECD through a linker peptide, such as a GS linker.

[007] In some embodiments, the CD80 ECD fusion molecule has a sialic acid content of 10-60 mol sialic acid (SA)/mol protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some such embodiments, the CD80 ECD fusion molecule is a CD80 ECD Fc fusion, for example, with a wild-type Fc domain such as a human IgG1, IgG2, or IgG4 Fc domain, or, alternatively, an IgG1 Fc domain with substitutions at L234F, L235E, and P331S. In some embodiments, the CD80 ECD fusion molecule comprises an amino acid sequence selected from SEQ ID NO: 5, SEQ ID NO: 12, SEQ ID NO:14, SEQ ID NO: 20, and SEQ ID NO: 21. In some embodiments above, the fusion molecule has a greater percentage tumor growth inhibition in a mouse syngeneic or xenograft model, such as in a CT26 mouse model than a fusion molecule of identical amino acid sequence but a lower SA content. In some embodiments above, where the fusion molecule comprises at least 10 mol SA/mol protein, such as at least 15 mol SA/mol protein, such as at least 20 mol SA/mol protein, the fusion molecule has a greater percentage tumor growth inhibition in a mouse syngeneic or xenograft model, such as in a CT26 mouse model than a fusion molecule of

identical amino acid sequence but having less than 10 mol SA/mol protein or less than 15 mol SA/mol protein or less than 20 mol SA/mol protein, respectively.

[008] Some embodiments herein comprise a CD80 ECD fusion molecule wherein the molecule is capable of at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 90%, such as at least 95%, such as at least 98% tumor cell growth inhibition in at least one mouse syngeneic or xenograft cancer model, such as a CT26 model, over a period of at least ten days, such as at least two weeks, such as over a period of ten days to two weeks or two to three weeks, or at least three weeks.

[009] In some embodiments, mice are given one to three doses of 0.3 to 3 mg/kg, such as 0.3 to 0.6 mg/kg, of the CD80 ECD Fc fusion molecule. In some such embodiments, the CD80 ECD fusion molecule also has a sialic acid content of 10-60 mol sialic acid (SA)/mol protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the CD80 ECD fusion molecule has an Fc as a fusion partner, such as a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the CD80 ECD fusion molecule comprises an amino acid sequence selected from SEQ ID NO: 5, SEQ ID NO: 12, SEQ ID NO:14, SEQ ID NO: 20, and SEQ ID NO: 21. In some embodiments above where the fusion molecule comprises at least 10 mol SA/mol protein, such as at least 15 mol SA/mol protein, such as at least 20 mol SA/mol protein, the molecule has a greater percentage tumor growth inhibition in a mouse syngeneic or xenograft model, such as in a CT26 mouse model, than a fusion molecule of identical amino acid sequence but having less than 10 mol SA/mol protein or less than 15 mol SA/mol protein or less than 20 mol SA/mol protein, respectively. In some embodiments above where the fusion molecule comprises at least 10 mol SA/mol protein, such as at least 15 mol SA/mol protein, such as at least 20 mol SA/mol protein, the molecule has a greater percentage tumor growth inhibition in a mouse syngeneic or xenograft model, such as in a CT26 mouse model, after at least ten days or at least two

weeks or at least three weeks, such as ten days to two weeks or two to three weeks, than an anti-CTLA4 antibody, such as anti-CTLA4 antibody clone 9D9.

[0010] In some of the above embodiments, the CD80 ECD Fc fusion molecule is also capable of inducing complete tumor regression in mice from the syngeneic or xenograft tumor model, such as a CT26 model.

[0011] Also provided herein are compositions comprising a CD80 ECD or CD80 ECD fusion molecule of any of the embodiments described above, and further comprising at least one pharmaceutically acceptable carrier. Some such compositions further comprise at least one additional therapeutic agent.

[0012] In some embodiments, the additional therapeutic agent comprises at least one immune stimulating agent. In some embodiments, the immune stimulating agent comprises a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor. The PD-1/PD-L1 inhibitor may be an antibody, such as an anti-PD-1 antibody or anti-PD-L1 antibody, a peptide or fusion molecule, or a small molecule.

[0013] In some embodiments, the PD-1/PD-L1 inhibitor is an anti-PD-1 antibody, such as an antibody comprising the heavy chain and light chain CDRs, or comprising the heavy and light chain variable regions, or comprising the full amino acid sequence, of an antibody selected from nivolumab, pidilizumab, and pembrolizumab. In some embodiments, the PD-1/PD-L1 inhibitor is an anti-PD-L1 antibody, such as an antibody comprising the heavy chain and light chain CDRs, the heavy and light chain variable regions, or the full amino acid sequence, of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.

[0014] In some embodiments, the PD-1/PD-L1 inhibitor is a PD-1 fusion molecule, such as AMP-224 or a polypeptide such as AUR-012.

[0015] Also included herein are methods of treating cancer in a subject comprising administering to the subject an effective amount of a CD80 ECD or CD80 ECD fusion protein or a composition from among the embodiments described above. In some embodiments, the cancer is a solid tumor. In some embodiments, the cancer is selected from colorectal cancer, breast cancer, gastric cancer, non-small cell lung cancer, melanoma, squamous cell carcinoma of the head and neck, ovarian cancer, pancreatic cancer, renal cell carcinoma, hepatocellular carcinoma, bladder cancer, and endometrial

cancer. In some embodiments, the cancer is recurrent or progressive after a therapy selected from surgery, chemotherapy, radiation therapy, or a combination thereof.

[0016] In some of the methods herein, the CD80 ECD, CD80 ECD fusion molecule, or composition is administered in combination with at least one additional therapeutic agent. In some such embodiments, the additional therapeutic agent may be packaged with the CD80 ECD or CD80 ECD fusion molecule as part of the same composition, e.g. mixed together in one composition or provided in separate containers, vials, or other packages. In some embodiments, the additional therapeutic agent comprises at least one immune stimulating agent. In some embodiments, the immune stimulating agent comprises a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor. The PD-1/PD-L1 inhibitor may be an antibody, such as an anti-PD-1 antibody or anti-PD-L1 antibody, a peptide or fusion molecule, or a small molecule.

[0017] In some embodiments, the PD-1/PD-L1 inhibitor is an anti-PD-1 antibody, such as an antibody comprising the heavy chain and light chain CDRs, or comprising the heavy and light chain variable regions, or comprising the full amino acid sequence, of an antibody selected from nivolumab, pidilizumab, and pembrolizumab. In some embodiments, the PD-1/PD-L1 inhibitor is an anti-PD-L1 antibody, such as an antibody comprising the heavy chain and light chain CDRs, the heavy and light chain variable regions, or the full amino acid sequence, of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.

[0018] In some embodiments, the PD-1/PD-L1 inhibitor is a PD-1 fusion molecule, such as AMP-224 or a polypeptide such as AUR-012.

[0019] In some embodiments of the methods herein, the CD80 ECD or CD80 ECD fusion molecule and the additional therapeutic agent, such as an immune stimulating agent, such as a PD-1/PD-L1 inhibitor, may be administered concurrently or sequentially. In some cases, one or more doses of the PD-1/PD-L1 inhibitor are administered prior to administering the CD80 ECD or CD80 ECD fusion molecule. In some cases, the subject has received a complete course of immune stimulating agent, e.g., PD-1/PD-L1 inhibitor therapy prior to administration of the CD80 ECD or CD80 ECD fusion molecule. In some cases, the CD80 ECD or CD80 ECD fusion molecule is administered during a second course of immune stimulation agent, e.g. PD-1/PD-L1

inhibitor therapy. In some cases, the subject has received at least one, at least two, at least three, or at least four doses of the immune stimulating agent, such as PD-1/PD-L1 inhibitor prior to administration of the CD80 ECD or CD80 ECD fusion molecule. In some cases, at least one dose of the immune stimulating agent, e.g. PD-1/PD-L1 inhibitor is administered concurrently with the CD80 ECD or CD80 ECD fusion molecule.

[0020] In some embodiments, one or more doses of the CD80 ECD or CD80 ECD fusion molecule are administered prior to administering an additional therapeutic agent, such as an immune stimulating agent, such as a PD-1/PD-L1 inhibitor. In some such cases, the subject has received at least one, at least two, at least three, or at least four doses of the CD80 ECD or CD80 ECD fusion molecule prior to administration of an immune stimulating agent, e.g. a PD-1/PD-L1 inhibitor. In some cases, at least one dose of the CD80 ECD or CD80 ECD fusion molecule is administered concurrently with the immune stimulating agent, e.g. PD-1/PD-L1 inhibitor.

[0021] In any of the methods herein, the subject may be resistant to treatment with a PD-1/PD-L1 inhibitor. In some such cases, the subject has previously received PD-1/PD-L1 inhibitor therapy, while in other such cases, the subject has not previously received PD-1/PD-L1 inhibitor therapy but is identified as resistant through other means such as certain phenotypic traits.

[0022] In any of the above methods, the subject may be administered an additional therapeutic agent comprising at least one chemotherapy agent, growth inhibitory agent, anti-angiogenesis agent and/or anti-neoplastic composition, in addition to the CD80 ECD or CD80 ECD fusion molecule.

[0023] In some embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule and an immune stimulating agent, such as a PD-1/PD-L1 inhibitor that is administered to the subject has been shown to reduce or inhibit tumor growth in at least one mouse syngeneic or xenograft cancer model in a synergistic fashion compared to treatment with either the CD80 ECD or fusion molecule or the immune stimulating agent, such as the PD-1/PD-L1 inhibitor, given alone. In some embodiments, the mouse model is a colorectal cancer model with murine colorectal carcinoma CT26 cells. In other embodiments, the model may be an MC38 model or a B16 model.

[0024] In any of the above method embodiments, the CD80 ECD or CD80 ECD fusion molecule administered to the subject may inhibit tumor growth in at least one mouse syngeneic or xenograft cancer model over a period of 1 week, 10 days, 2 weeks, or 3 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, or may inhibit growth of tumors by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% in a patient over a period of one month, two months, three months, six months, or one year. In any of the above method embodiments, administration of the CD80 ECD or CD80 ECD fusion molecule may reduce the volume of at least one tumor in an animal or human subject by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, for example, over a period of one month, two months, three months, six months, or one year. In some such embodiments, the tumor is a solid tumor.

[0025] In any of the above combination therapy method embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with an additional therapeutic agent, such as an immune stimulator, such as a PD-1/PD-L1 inhibitor, administered to the subject may inhibit tumor growth in at least one mouse syngeneic or xenograft cancer model over a period of 1 week, 10 days, 2 weeks, or 3 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, or may inhibit growth of tumors by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% in a patient over a period of one month, two months, three months, six months, or one year. In any of the above combination therapy method embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with an additional therapeutic agent, such as an immune stimulator, such as a PD-1/PD-L1 inhibitor, administered to the subject may reduce the volume of at least one tumor in an animal or human subject by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, for example, over a period of one month, two months, three months, six months, or one year. In some such embodiments, the tumor is a solid tumor.

[0026] In some of the above combination therapy embodiments, the CD80 ECD or CD80 ECD fusion molecule is a CD80 ECD fusion molecule comprising 10-60 mol sialic acid (SA) to mol of CD80 ECD protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some such embodiments, the CD80 ECD fusion molecule comprises an Fc domain as fusion partner, such as a wild-type Fc domain, such as a wild-type human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the CD80 ECD fusion molecule comprises an amino acid sequence selected from SEQ ID NO: 5, SEQ ID NO: 12, SEQ ID NO:14, SEQ ID NO: 20, and SEQ ID NO: 21. In some such embodiments, the CD80 ECD or CD80 ECD fusion molecule is capable of at least 90% reduction of growth of CT26 tumor cells in mice, such as at least 95%, or at least 98%, over a period of at least ten days, such as at least two weeks, such as at least three weeks, such as over a period of ten days to two weeks or two to three weeks. In some embodiments, these results are obtained after mice are given one to three doses of 0.3 to 3 mg/kg, such as 0.3 to 0.6 mg/kg, of the CD80 ECD Fc fusion molecule.

[0027] Also comprised herein is a CD80 ECD fusion molecule comprising a human CD80 ECD polypeptide and a human IgG1 Fc domain, such as a wild-type human IgG1 Fc, wherein the CD80 ECD Fc comprises 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20. In some embodiments, the molecule is capable

of at least 90% reduction, such as at least 95%, or at least 98% of growth of CT26 tumor cells in mice over a period of at least ten days, such as at least two weeks, such as at least three weeks, such as over a period of ten days to two weeks or two to three weeks. In some embodiments, these results are obtained after mice are given one to three doses of 0.3 to 3 mg/kg, such as 0.3 to 0.6 mg/kg, of the CD80 ECD Fc fusion molecule.

[0028] Also comprised are compositions comprising the CD80 ECD IgG1 Fc and further comprising at least one pharmaceutically acceptable carrier. Such compositions may also contain an additional therapeutic agent. In some embodiments, the additional therapeutic agent is at least one immune stimulating agent, such as a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor. In some cases, the PD-1/PD-L1 inhibitor is an antibody, such as an anti-PD-1 antibody, such as nivolumab, pidilizumab, and pembrolizumab. For example, the antibody may have the heavy chain and light chain CDRs or the heavy and light chain variable regions of an antibody selected from nivolumab, pidilizumab, and pembrolizumab. In other embodiments, the PD-1/PD-L1 inhibitor is an anti-PD-L1 antibody. An anti-PD-L1 antibody may have the heavy chain and light chain CDRs of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C, for example, or may comprise the heavy chain and light chain variable regions of BMS-936559, MPDL3280A, MEDI4736, or MSB0010718C. In some embodiments, the anti-PD-1 antibody is selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C. Alternatively, the PD-1/PD-L1 inhibitor may be a PD-1 fusion molecule such as AMP-224 or a polypeptide such as AUR-012.

[0029] This disclosure also encompasses methods of treating cancer in a subject comprising administering to the subject an effective amount of a CD80 ECD IgG1 Fc fusion molecule as described above. In some embodiments, the cancer is a solid tumor, such as a cancer selected from colorectal cancer, breast cancer, gastric cancer, non-small cell lung cancer, melanoma, squamous cell carcinoma of the head and neck, ovarian cancer, pancreatic cancer, renal cell carcinoma, hepatocellular carcinoma, bladder cancer, and endometrial cancer. In some embodiments, the cancer is recurrent or progressive after a therapy selected from surgery, chemotherapy, radiation therapy, or a combination thereof.

[0030] In some of these method embodiments, the CD80 ECD Fc comprises 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21. In some embodiments, the molecule is capable of at least 90% reduction of growth of CT26 tumor cells in mice over a period of two or three weeks following inoculation of mice with tumor cells. In some embodiments, the molecule is capable of at least 95% reduction of growth of CT26 tumor cells, such as at least 98% reduction, in mice over a period of two or three weeks following inoculation of mice with tumor cells. For example, such results may be obtained when the mice are given one to three doses of 0.3 to 3.0 mg/kg, such as 0.3 to 0.6 mg/kg, of the ECD Fc fusion molecule. In some of the method embodiments, the CD80 ECD Fc comprises 10-40 mol SA to mol of CD80 ECD Fc protein, and the CD80 ECD Fc reduces growth of CT26 tumor cells in mice over a period of two or three weeks by a greater degree than a CD80 ECD Fc protein of the same amino acid sequence comprising less than 10 mol SA to mol of CD80 ECD Fc protein. The disclosure herein also comprises methods of enhancing efficacy of a CD80 ECD fusion protein in treating cancer in a subject comprising increasing the level of sialic acid (SA) in the CD80 ECD fusion protein or providing a CD80 ECD fusion protein with an increased SA level and administering the CD80 ECD fusion protein comprising an increased level of SA to the subject. In some such embodiments, the SA level is increased by 5, 10, 20, 30, 40, or 50 mol to mol of CD80 ECD protein. In some of these method embodiments, the CD80 ECD Fc comprises 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA

to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21. In some embodiments, the molecule is capable of at least 90% reduction of growth of CT26 tumor cells in mice over a period of two or three weeks. In some embodiments, the molecule is capable of at least 95%, such as at least 98%, reduction of growth of CT26 tumor cells in mice over a period of at least ten days, such as at least two weeks, such as at least three weeks, such as over a period of ten days to two weeks or two to three weeks. In some embodiments, these results are obtained after mice are given one to three doses of 0.3 to 3 mg/kg, such as 0.3 to 0.6 mg/kg, of the CD80 ECD Fc fusion molecule.

[0031] In some of these method embodiments, the enhanced efficacy is measured as an increase in overall survival, an increase in disease-free survival, or as a greater reduction in the growth of at least one tumor in an animal or human subject. In other words, one or more of these parameters is improved upon administration of the CD80 ECD fusion molecule with higher SA content compared to a CD80 ECD fusion molecule with a lower SA content. In other embodiments, the enhanced efficacy is measured as a greater reduction in tumor growth in a mouse syngeneic or xenograft model such as a CT26 mouse model or as a reduced rate of clearance in an animal or human subject. In some embodiments, the efficacy is measured as a greater reduction in tumor growth of at least one tumor in the subject or a greater reduction in tumor growth in at least one mouse syngeneic or xenograft model, and wherein the tumor growth is further reduced by at least 10%, such as at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% upon administration of the CD80 ECD fusion molecule with increased SA level in comparison to administration of the CD80 ECD fusion molecule without the increased SA level.

[0032] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the claims. The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All references cited

herein, including patent applications and publications, are incorporated herein by reference in their entireties for any purpose.

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] **Figs. 1a-1b** show effects of administering a CD80 ECD Fc fusion molecule compared with a CTLA4 ECD Fc fusion molecule and a saline control to mice implanted with murine colorectal carcinoma cell line CT26 cells. **Fig. 1a** shows tumor volume at up to 21 days post-inoculation of mice with the CT26 cells. As shown in the figure, CTLA4 ECD Fc enhanced tumor growth while CD80 ECD Fc inhibited tumor growth in a statistically significant manner compared to the saline control. *P*-values were calculated using unpaired, two-tailed t-test analyses of the calculated tumor volumes on each day of the study (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). **Fig. 1b** shows individual tumor volumes on Day 19 post-inoculation for the three groups.

[0034] **Figs. 2a-2b** show effects of administering a CD80 ECD Fc fusion molecule or an anti-PD-1 antibody or a combination of the two compared to a saline control to mice implanted with murine colorectal carcinoma cell line CT26 cells. **Fig. 2a** shows tumor volume at up to 14 days post-inoculation. Mice administered with the CD80 ECD Fc and anti-PD-1 combination showed statistically significant reduction in tumor growth compared to either the CD80 ECD Fc ($p < 0.01$ beginning after Day 9) or anti-PD-1 ($p < 0.01$ on Day 14) single therapies. Statistical significance was determined via two-tailed, unpaired t-Test comparing the combination group to the CD80 ECD Fc group. **Fig. 2b** shows individual tumor volumes on Day 14.

[0035] **Figs. 3a-b** show the effect of the Fc fusion polypeptide sequence on effects of a CD80 ECD Fc fusion molecule on tumor growth of the CT26 tumors in mice. Specifically, mice were administered saline control, or a CD80 ECD Fc with a human IgG1 wild-type Fc domain fusion partner (CD80-IgG1 WT), or with a CD80 ECD Fc with a mutant (L234F/L235E/P331S) human IgG1 Fc domain fusion partner (CD80-IgG1 MT). **Fig. 3a** shows changes in tumor volume up to Day 21 post-inoculation. The mutated Fc domain resulted in enhanced anti-tumor activity, which was statistically significant beginning on Day 14 after inoculation ($p < 0.01$). Statistical significance was determined via a two-tailed, unpaired t-Test. **Fig. 3b** shows individual tumor volumes on Day 21 post-inoculation.

[0036] **Figs. 4a-b** show staining of murine tumor cells for presence of CD3+ and CD4+ T cells after exposure to saline control or to CD80 ECD Fc fusion molecules with either wild-type and mutant Fc fusion partners. **Fig. 4a** provides representative images showing CD3+ cells (top images) and corresponding DAPI staining (nuclei, bottom images) in CT26 tumors collected 7 days after injection of Saline, CD80-IgG1 WT or CD80-IgG1 MT. Both CD80-IgG1 WT and CD80-IgG1 MT increased the number of CD3+ cells within the tumors compared to vehicle but the magnitude of the increase was greater after CD80-IgG1 MT. Images were collected using the 10X objective. **Fig. 4b** provides representative images showing CD3+ cells (top row) and CD4+ cells (bottom row) in CT26 tumors collected 7 days after injection of Saline, CD80-IgG1 WT or CD80-IgG1 MT. The images were taken in the same field of view but with different channels. Both CD80-IgG1 WT and CD80-IgG1 MT increased the number of infiltrating CD4+ cells compared to vehicle. The ratio of CD3+ to CD4+ cells was increased with the CD80-IgG1 MT compared to the CD80-IgG1 WT. Images were collected using the 10X objective.

[0037] **Figs. 5a-d** show release of cytokines IFN- γ and TNF- α from T-cells on 96 well tissue culture plates exposed to protein A beads coated with 0.01, 0.1, or 1 μ g/well of a CD80 ECD IgG1 Fc domain fusion molecule (CD80-Fc). **Figs. 5a** and **5c** show that bead-immobilized CD80-Fc alone did not cause significant T-cell activation, as measured by soluble cytokine production. **Figs. 5b** and **5d** show that when a small amount of OKT3-scFv (too low to cause T-cell stimulation on its own) was immobilized along with the CD80-Fc, cytokine release was observed.

[0038] **Fig. 6** shows tumor growth of murine CT26 tumors following treatment with a saline control or either 0.3 or 0.6 mg/kg doses of three different lots of a CD80 ECD Fc fusion molecule having three different sialic acid (SA) contents. Lot A has 5 mol SA/mol protein, lot D has 15 mol SA/mol protein and lot E has 20 mol SA/mol protein. Treatment with CD80 ECD Fc lot E dosed at 0.3 or 0.6 mg/kg resulted in a 93% and 98% inhibition of tumor growth compared to the control ($P < 0.001$). Treatment with CD80 ECD Fc lot D dosed at 0.3 or 0.6 mg/kg resulted in a 93% and 95% inhibition of tumor growth compared to the control ($P < 0.001$). By comparison, treatment with CD80 ECD Fc lot A at 0.3mg/kg did not inhibit tumor growth compared

to the control and when dosed at 0.6mg/kg it only induced 70% inhibition ($P < 0.001$) of tumor growth.

[0039] **Fig. 7** shows tumor growth of CT26 tumors treated with mouse IgG2b at 10mg/kg; murine CD80 ECD-Fc SA 20 mol/mol at 0.3mg/kg; anti-CTLA4 antibody clone 9D9 at 10mg/kg; and anti-CTLA4 antibody clone 9D9 at 1.5mg/kg. Arrows indicate when mice were dosed. The asterisk symbol (*) denotes statistically significant differences between murine CD80 ECD-Fc SA 20 mol/mol at 0.3mg/kg and the other treatments.

[0040] **Fig. 8** shows tumor growth of MC38 tumors treated with mouse IgG2b at 10mg/kg; murine CD80 ECD-Fc SA 20 mol/mol at 3mg/kg; anti-CTLA4 antibody clone 9D9 at 10mg/kg; and anti-CTLA4 antibody clone 9D9 at 1.5mg/kg. Arrows indicate when mice were dosed. The asterisk symbol (*) denotes statistically significant differences between murine CD80 ECD-Fc SA 20 mol/mol at 3mg/kg and the other treatments.

[0041] **Fig. 9** shows tumor growth of B16 tumors treated with mouse IgG2b at 10mg/kg; murine CD80 ECD-Fc SA 20 mol/mol at 3mg/kg; anti-CTLA4 antibody clone 9D9 at 10mg/kg; and anti-CTLA4 antibody clone 9D9 at 1.5mg/kg. Arrows indicate when mice were dosed. The asterisk symbol (*) denotes statistically significant differences between murine CD80 ECD-Fc SA 20 mol/mol at 3mg/kg and the other treatments.

DESCRIPTION OF PARTICULAR EMBODIMENTS

Definitions

[0042] Unless otherwise defined, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular.

[0043] In this application, the use of “or” means “and/or” unless stated otherwise. In the context of a multiple dependent claim, the use of “or” refers back to more than one preceding independent or dependent claim in the alternative only. Also, terms such as “element” or “component” encompass both elements and components

comprising one unit and elements and components that comprise more than one subunit unless specifically stated otherwise.

[0044] Exemplary techniques used in connection with recombinant DNA, oligonucleotide synthesis, tissue culture and transformation (e.g., electroporation, lipofection), enzymatic reactions, and purification techniques are described, e.g., in Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)), among other places.

[0045] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0046] The terms “**nucleic acid molecule**” and “**polynucleotide**” may be used interchangeably, and refer to a polymer of nucleotides. Such polymers of nucleotides may contain natural and/or non-natural nucleotides, and include, but are not limited to, DNA, RNA, and PNA. “**Nucleic acid sequence**” refers to the linear sequence of nucleotides that comprise the nucleic acid molecule or polynucleotide.

[0047] The terms “**polypeptide**” and “**protein**” are used interchangeably to refer to a polymer of amino acid residues, and are not limited to a minimum length. Such polymers of amino acid residues may contain natural or non-natural amino acid residues, and include, but are not limited to, peptides, oligopeptides, dimers, trimers, and multimers of amino acid residues. Both full-length proteins and fragments thereof are encompassed by the definition. The terms also include post-expression modifications of the polypeptide, for example, glycosylation, sialylation, acetylation, phosphorylation, and the like. Furthermore, for purposes of the present invention, a “polypeptide” refers to a protein which includes modifications, such as deletions, additions, and substitutions (generally conservative in nature), to the native sequence, as long as the protein maintains the desired activity. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the proteins or errors due to PCR amplification.

[0048] A “**CD80 extracellular domain**” or “**CD80 ECD**” refers to an extracellular domain polypeptide of CD80, including natural and engineered variants thereof. Nonlimiting examples of CD80 ECDs include SEQ ID NOs:--. A “**CD80 ECD fusion molecule**” refers to a molecule comprising a CD80 ECD and a fusion partner such as an Fc domain, albumin, or PEG. The fusion partner may be covalently

attached, for example, to the N- or C- terminal of the CD80 ECD or at an internal location. Nonlimiting examples of CD80 ECD fusion molecules include SEQ ID NOs:--

[0049] The terms “**programmed cell death protein 1**” and “**PD-1**” refer to an immunoinhibitory receptor belonging to the CD28 family. PD-1 is expressed predominantly on previously activated T cells *in vivo*, and binds to two ligands, PD-L1 and PD-L2. The term "PD-1" as used herein includes human PD-1 (hPD-1), variants, isoforms, and species homologs of hPD-1, and analogs having at least one common epitope with hPD-1. The complete hPD-1 sequence can be found under GenBank Accession No. U64863. In some embodiments, the PD-1 is a human PD-1 having the amino acid sequence of SEQ ID NO: -- (precursor, with signal sequence) or SEQ ID NO: -- (mature, without signal sequence).

[0050] The terms “**programmed cell death 1 ligand 1**” and “**PD-L1**” refer to one of two cell surface glycoprotein ligands for PD-1 (the other being PD-L2) that down regulate T cell activation and cytokine secretion upon binding to PD-1. The term "PD-L1" as used herein includes human PD-L1 (hPD-L1), variants, isoforms, and species homologs of hPD-L1, and analogs having at least one common epitope with hPD-L1. The complete hPD-L1 sequence can be found under GenBank[®] Accession No. Q9NZQ7. In some embodiments, the PD-L1 is a human PD-L1 having the amino acid sequence of SEQ ID NO: -- (precursor, with signal sequence) or SEQ ID NO: -- (mature, without signal sequence).

[0051] The term “**immune stimulating agent**” as used herein refers to a molecule that stimulates the immune system by either acting as an agonist of an immune-stimulatory molecule, including a co-stimulatory molecule, or acting as an antagonist of an immune inhibitory molecule, including a co-inhibitory molecule. An immune stimulating agent may be a biologic, such as an antibody or antibody fragment, other protein, or vaccine, or may be a small molecule drug. An “**immune stimulatory molecule**” includes a receptor or ligand that acts to enhance, stimulate, induce, or otherwise “turn-on” an immune response. Immune stimulatory molecules as defined herein include co-stimulatory molecules. An “**immune inhibitory molecule**” includes a receptor or ligand that acts to reduce, inhibit, suppress, or otherwise “turn-off” an immune response. Immune inhibitory molecules as defined herein include co-inhibitory

molecules. Such immune stimulatory and immune inhibitory molecules may be, for example, receptors or ligands found on immune cells such as T cells, or found on cells involved in innate immunity such as NK cells.

[0052] The term “**PD-1/PD-L1 inhibitor**” refers to a moiety that disrupts the PD-1/PD-L1 signaling pathway. In some embodiments, the inhibitor inhibits the PD-1/PD-L1 signaling pathway by binding to PD-1 and/or PD-L1. In some embodiments, the inhibitor also binds to PD-L2. In some embodiments, a PD-1/PD-L1 inhibitor blocks binding of PD-1 to PD-L1 and/or PD-L2. Nonlimiting exemplary PD-1/PD-L1 inhibitors include antibodies that bind to PD-1; antibodies that bind to PD-L1; fusion proteins, such as AMP-224; and peptides, such as AUR-012.

[0053] The term “**antibody that inhibits PD-1**” refers to an antibody that binds to PD-1 or binds to PD-L1 and thereby inhibits PD-1 and/or PD-L1 signaling. In some embodiments, an antibody that inhibits PD-1 binds to PD-1 and blocks binding of PD-L1 and/or PD-L2 to PD-1. In some embodiments, an antibody that inhibits PD-1 binds to PD-L1 and blocks binding of PD-1 to PD-L1. An antibody that inhibits PD-1 that binds to PD-L1 may be referred to as an anti-PD-L1 antibody. An antibody that inhibits PD-1 that binds to PD-1 may be referred to as an anti-PD-1 antibody.

[0054] With reference to CD80 ECDs and CD80 ECD fusion molecules, the term “**blocks binding of**” a ligand, and grammatical variants thereof, refers to the ability to inhibit an interaction between CD80 and a CD80 ligand, such as CD28, CTLA4, or PD-L1. Such inhibition may occur through any mechanism, including by the CD80 ECDs or CD80 ECD fusion molecules competing for binding with CD80 ligands.

[0055] With reference to anti-PD-1 antibodies and PD-1 fusion molecules or peptides the term “**blocks binding of**” a ligand, such as PD-L1, and grammatical variants thereof, are used to refer to the ability to inhibit the interaction between PD-1 and a PD-1 ligand, such as PD-L1. Such inhibition may occur through any mechanism, including direct interference with ligand binding, e.g., because of overlapping binding sites on PD-1, and/or conformational changes in PD-1 induced by the antibody that alter ligand affinity, etc., or, in the case of a PD-1 fusion molecule or peptide, by competing for binding with a PD-1 ligand.

[0056] “**Affinity**” or “**binding affinity**” refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., a polypeptide)

and its binding partner (*e.g.*, a ligand). In some embodiments, “binding affinity” refers to intrinsic binding affinity, which reflects a 1:1 interaction between members of a binding pair (*e.g.*, polypeptide and ligand). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d).

[0057] The term “**antibody**” as used herein refers to a molecule comprising at least complementarity-determining region (CDR) 1, CDR2, and CDR3 of a heavy chain and at least CDR1, CDR2, and CDR3 of a light chain, wherein the molecule is capable of binding to antigen. The term antibody includes, but is not limited to, fragments that are capable of binding antigen, such as Fv, single-chain Fv (scFv), Fab, Fab’, and (Fab’)₂. The term antibody also includes, but is not limited to, chimeric antibodies, humanized antibodies, and antibodies of various species such as mouse, human, cynomolgus monkey, etc.

[0058] In some embodiments, an antibody comprises a heavy chain variable region and a light chain variable region. In some embodiments, an antibody comprises at least one heavy chain comprising a heavy chain variable region and at least a portion of a heavy chain constant region, and at least one light chain comprising a light chain variable region and at least a portion of a light chain constant region. In some embodiments, an antibody comprises two heavy chains, wherein each heavy chain comprises a heavy chain variable region and at least a portion of a heavy chain constant region, and two light chains, wherein each light chain comprises a light chain variable region and at least a portion of a light chain constant region. As used herein, a single-chain Fv (scFv), or any other antibody that comprises, for example, a single polypeptide chain comprising all six CDRs (three heavy chain CDRs and three light chain CDRs) is considered to have a heavy chain and a light chain. In some such embodiments, the heavy chain is the region of the antibody that comprises the three heavy chain CDRs and the light chain in the region of the antibody that comprises the three light chain CDRs.

[0059] The term “**heavy chain variable region**” refers to a region comprising heavy chain HVR1, framework (FR) 2, HVR2, FR3, and HVR3. In some embodiments, a heavy chain variable region also comprises at least a portion of an FR1 and/or at least a portion of an FR4.

[0060] The term “**heavy chain constant region**” refers to a region comprising at least three heavy chain constant domains, C_{H1}, C_{H2}, and C_{H3}. Nonlimiting exemplary

heavy chain constant regions include γ , δ , and α . Nonlimiting exemplary heavy chain constant regions also include ϵ and μ . Each heavy constant region corresponds to an antibody isotype. For example, an antibody comprising a γ constant region is an IgG antibody, an antibody comprising a δ constant region is an IgD antibody, and an antibody comprising an α constant region is an IgA antibody. Further, an antibody comprising a μ constant region is an IgM antibody, and an antibody comprising an ϵ constant region is an IgE antibody. Certain isotypes can be further subdivided into subclasses. For example, IgG antibodies include, but are not limited to, IgG1 (comprising a γ_1 constant region), IgG2 (comprising a γ_2 constant region), IgG3 (comprising a γ_3 constant region), and IgG4 (comprising a γ_4 constant region) antibodies; IgA antibodies include, but are not limited to, IgA1 (comprising an α_1 constant region) and IgA2 (comprising an α_2 constant region) antibodies; and IgM antibodies include, but are not limited to, IgM1 and IgM2.

[0061] The term “**heavy chain**” refers to a polypeptide comprising at least a heavy chain variable region, with or without a leader sequence. In some embodiments, a heavy chain comprises at least a portion of a heavy chain constant region. The term “full-length heavy chain” refers to a polypeptide comprising a heavy chain variable region and a heavy chain constant region, with or without a leader sequence.

[0062] The term “**light chain variable region**” refers to a region comprising light chain HVR1, framework (FR) 2, HVR2, FR3, and HVR3. In some embodiments, a light chain variable region also comprises an FR1 and/or an FR4.

[0063] The term “**light chain constant region**” refers to a region comprising a light chain constant domain, C_L . Nonlimiting exemplary light chain constant regions include λ and κ .

[0064] The term “**light chain**” refers to a polypeptide comprising at least a light chain variable region, with or without a leader sequence. In some embodiments, a light chain comprises at least a portion of a light chain constant region. The term “full-length light chain” refers to a polypeptide comprising a light chain variable region and a light chain constant region, with or without a leader sequence.

[0065] The term “**hypervariable region**” or “**HVR**” refers to each of the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops (“hypervariable loops”). Generally, native four-chain

antibodies comprise six HVRs; three in the V_H (H1, H2, H3), and three in the V_L (L1, L2, L3). HVRs generally comprise amino acid residues from the hypervariable loops and/or from the “**complementarity determining regions**” (“**CDRs**”), the latter being of highest sequence variability and/or involved in antigen recognition. Exemplary hypervariable loops occur at amino acid residues 26-32 (L1), 50-52 (L2), 91-96 (L3), 26-32 (H1), 53-55 (H2), and 96-101 (H3). (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987).) Exemplary CDRs (CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3) occur at amino acid residues 24-34 of L1, 50-56 of L2, 89-97 of L3, 31-35B of H1, 50-65 of H2, and 95-102 of H3. (Kabat *et al.*, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991)). The terms hypervariable regions (HVRs) and complementarity determining regions (CDRs) both refer to portions of the variable region that form the antigen binding regions.

[0066] A “**chimeric antibody**” as used herein refers to an antibody comprising at least one variable region from a first species (such as mouse, rat, cynomolgus monkey, etc.) and at least one constant region from a second species (such as human, cynomolgus monkey, etc.). In some embodiments, a chimeric antibody comprises at least one mouse variable region and at least one human constant region. In some embodiments, a chimeric antibody comprises at least one cynomolgus variable region and at least one human constant region. In some embodiments, a chimeric antibody comprises at least one rat variable region and at least one mouse constant region. In some embodiments, all of the variable regions of a chimeric antibody are from a first species and all of the constant regions of the chimeric antibody are from a second species.

[0067] A “**humanized antibody**” as used herein refers to an antibody in which at least one amino acid in a framework region of a non-human variable region has been replaced with the corresponding amino acid from a human variable region. In some embodiments, a humanized antibody comprises at least one human constant region or fragment thereof. In some embodiments, a humanized antibody is a Fab, an scFv, a (Fab')₂, etc.

[0068] A “**human antibody**” as used herein refers to antibodies produced in humans, antibodies produced in non-human animals that comprise human immunoglobulin genes, such as XenoMouse®, and antibodies selected using in vitro

methods, such as phage display, wherein the antibody repertoire is based on a human immunoglobulin sequences.

[0069] The term “**leader sequence**” refers to a sequence of amino acid residues located at the N terminus of a polypeptide that facilitates secretion of a polypeptide from a mammalian cell. A leader sequence may be cleaved upon export of the polypeptide from the mammalian cell, forming a mature protein. Leader sequences may be natural or synthetic, and they may be heterologous or homologous to the protein to which they are attached. Nonlimiting exemplary leader sequences also include leader sequences from heterologous proteins. In some embodiments, an antibody lacks a leader sequence. In some embodiments, an antibody comprises at least one leader sequence, which may be selected from native antibody leader sequences and heterologous leader sequences.

[0070] The term “**isolated**” as used herein refers to a molecule that has been separated from at least some of the components with which it is typically found in nature. For example, a polypeptide is referred to as “isolated” when it is separated from at least some of the components of the cell in which it was produced. Where a polypeptide is secreted by a cell after expression, physically separating the supernatant containing the polypeptide from the cell that produced it is considered to be “isolating” the polypeptide. Similarly, a polynucleotide is referred to as “isolated” when it is not part of the larger polynucleotide (such as, for example, genomic DNA or mitochondrial DNA, in the case of a DNA polynucleotide) in which it is typically found in nature, or is separated from at least some of the components of the cell in which it was produced, e.g., in the case of an RNA polynucleotide. Thus, a DNA polynucleotide that is contained in a vector inside a host cell may be referred to as “isolated” so long as that polynucleotide is not found in that vector in nature.

[0071] The term “**reduce**” or “**reduces**” when applied to a parameter such as tumor volume means to lower the level of that parameter in an observable, measurable way. In some embodiments, the reduction may be by at least 10%, such as by at least 20%, at least 30%, at least 40%, or at least 50%. In some embodiments, the reduction may be statistically significant compared to an alternative treatment or control.

[0072] The terms “**subject**” and “**patient**” are used interchangeably herein to refer to a human. In some embodiments, methods of treating other mammals, including, but not limited to, rodents, simians, felines, canines, equines, bovines, porcines, ovines,

caprines, mammalian laboratory animals, mammalian farm animals, mammalian sport animals, and mammalian pets, are also provided.

[0073] The terms “**resistant**” or “**nonresponsive**” when used in the context of treatment with a therapeutic agent, means that the subject shows decreased response or lack of response to a standard dose of the therapeutic agent, relative to the subject’s response to the standard dose of the therapeutic agent in the past, or relative to the expected response of a similar subject with a similar disorder to the standard dose of the therapeutic agent. Thus, in some embodiments, a subject may be resistant to therapeutic agent although the subject has not previously been given the therapeutic agent, or the subject may develop resistance to the therapeutic agent after having responded to the agent on one or more previous occasions.

[0074] The term “**sample**,” as used herein, refers to a composition that is obtained or derived from a subject that contains a cellular and/or other molecular entity that is to be characterized, quantitated, and/or identified, for example based on physical, biochemical, chemical and/or physiological characteristics. An exemplary sample is a tissue sample.

[0075] The term “**tissue sample**” refers to a collection of similar cells obtained from a tissue of a subject. The source of the tissue sample may be solid tissue as from a fresh, frozen and/or preserved organ or tissue sample or biopsy or aspirate; blood or any blood constituents; bodily fluids such as cerebral spinal fluid, amniotic fluid, peritoneal fluid, synovial fluid, or interstitial fluid; cells from any time in gestation or development of the subject. In some embodiments, a tissue sample is a synovial biopsy tissue sample and/or a synovial fluid sample. In some embodiments, a tissue sample is a synovial fluid sample. The tissue sample may also be primary or cultured cells or cell lines. Optionally, the tissue sample is obtained from a disease tissue/organ. The tissue sample may contain compounds that are not naturally intermixed with the tissue in nature such as preservatives, anticoagulants, buffers, fixatives, nutrients, antibiotics, or the like. A “control sample” or “control tissue”, as used herein, refers to a sample, cell, or tissue obtained from a source known, or believed, not to be afflicted with the disease for which the subject is being treated.

[0076] For the purposes herein a “**section**” of a tissue sample means a part or piece of a tissue sample, such as a thin slice of tissue or cells cut from a solid tissue sample.

[0077] The term “**cancer**” is used herein to refer to a group of cells that exhibit abnormally high levels of proliferation and growth. A cancer may be benign (also referred to as a benign tumor), pre-malignant, or malignant. Cancer cells may be solid cancer cells (i.e. “**solid tumors**”) or may be leukemic cancer cells. The term “**cancer growth**” is used herein to refer to proliferation or growth by a cell or cells that comprise a cancer that leads to a corresponding increase in the size or extent of the cancer.

[0078] Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular nonlimiting examples of such cancers include squamous cell cancer, small-cell lung cancer, pituitary cancer, esophageal cancer, astrocytoma, soft tissue sarcoma, non-small cell lung cancer (including squamous cell non-small cell lung cancer), adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, renal cell carcinoma, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, brain cancer, endometrial cancer, testis cancer, cholangiocarcinoma, gallbladder carcinoma, gastric cancer, melanoma, and various types of head and neck cancer (including squamous cell carcinoma of the head and neck).

[0079] “**Treatment**,” as used herein, refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. In certain embodiments, the term “**treatment**” covers any administration or application of a therapeutic for disease in a mammal, including a human, and includes inhibiting or slowing the disease or progression of the disease; partially or fully relieving the disease, for example, by causing regression, or restoring or repairing a lost, missing, or defective function; stimulating an inefficient process; or causing the disease plateau to have reduced severity. The term “**treatment**” also includes reducing the severity of any phenotypic characteristic and/or reducing the incidence, degree, or likelihood of that characteristic. Those in need of

treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

[0080] The term “**efficacy**” as used herein may be determined from one or more parameters such as survival or disease-free survival over a period of time such as 1 year, 5 years, or 10 years, as well as parameters such as the reduction in growth of one or more tumors in a subject. Pharmacokinetic parameters such as bioavailability and underlying parameters such as clearance rate may also impact efficacy. Thus, an “enhanced efficacy” (i.e. an improvement in efficacy) may be due to improved pharmacokinetic parameters as well as improved potency, and may be measured by comparing clearance rates and tumor growth in test animals or in human subjects, as well as parameters such as survival, rate of recurrence, or disease-free survival.

[0081] The term “**effective amount**” or “**therapeutically effective amount**” refers to an amount of a drug effective to treat a disease or disorder in a subject. In certain embodiments, an effective amount refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result. A therapeutically effective amount of a CD80 ECD or CD80 ECD fusion molecule may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the drug to elicit a desired response in the individual. A therapeutically effective amount encompasses an amount in which any toxic or detrimental effects of the drug are outweighed by the therapeutically beneficial effects. In some embodiments, the expression “effective amount” refers to an amount of the drug that is effective for treating the cancer.

[0082] Administration “**in combination with**” one or more further therapeutic agents, such as an immune stimulating agent, includes simultaneous (concurrent) and consecutive (sequential) administration in any order.

[0083] A “**pharmaceutically acceptable carrier**” refers to a non-toxic solid, semisolid, or liquid filler, diluent, encapsulating material, formulation auxiliary, or carrier conventional in the art for use with a therapeutic agent that together comprise a “**pharmaceutical composition**” for administration to a subject. A pharmaceutically acceptable carrier is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. The pharmaceutically acceptable carrier is appropriate for the formulation employed. For example, if the

therapeutic agent is to be administered orally, the carrier may be a gel capsule. If the therapeutic agent is to be administered subcutaneously, the carrier ideally is not irritable to the skin and does not cause injection site reaction.

Exemplary CD80 Extracellular Domain and Extracellular Domain Fusion

Molecules

[0084] CD80 ECD and CD80 ECD fusion molecules are provided herein. CD80 ECDs, for example, may comprise the ECDs of human CD80 isoform 1, isoform 2, and isoform 3 (see SEQ ID NOs: 1-3). In some embodiments, CD80 ECDs may comprise the amino acid sequence of SEQ ID NO:5.

[0085] CD80 ECD fusion molecules may comprise fusion partners such as polymers, polypeptides, lipophilic moieties, and succinyl groups. Exemplary polypeptide fusion partners include, but are not limited to, serum albumin and an IgG Fc domain. Further exemplary polymer fusion partners include, but are not limited to, polyethylene glycol, including polyethylene glycols having branched and/or linear chains. The amino acid sequences of certain exemplary Fc domains are shown in SEQ ID NOs: 9-16 herein.

[0086] In certain embodiments, the CD80 ECD or CD80 ECD fusion molecule lacks a signal peptide. In certain embodiments, the CD80 ECD or CD80 ECD fusion molecule includes at least one signal peptide, which may be selected from a native CD80 signal peptide (SEQ ID NO: 7 or amino acids 1-34 of SEQ ID NO:1) and/or a heterologous signal peptide.

[0087] In the case of a CD80 ECD fusion molecule, the fusion partner may be linked to either the amino-terminus or the carboxy-terminus of the polypeptide. In certain embodiments, the polypeptide and the fusion partner are covalently linked. If the fusion partner is also a polypeptide (“the fusion partner polypeptide”), the polypeptide and the fusion partner polypeptide may be part of a continuous amino acid sequence. In such cases, the polypeptide and the fusion partner polypeptide may be translated as a single polypeptide from a coding sequence that encodes both the polypeptide and the fusion partner polypeptide. In some such cases, the two polypeptides are directly linked in sequence such that the N-terminal of one polypeptide immediately follows the C-terminal of the other with no intervening amino acids. In other cases, a linker peptide sequence is inserted in between the two polypeptides, such as a GS linker sequence. In certain embodiments, a CD80 ECD and the fusion partner are covalently linked through

other means, such as, for example, a chemical linkage other than a peptide bond. In certain embodiments, the polypeptide and the fusion partner are noncovalently linked. In certain such embodiments, they may be linked, for example, using binding pairs. Exemplary binding pairs include, but are not limited to, biotin and avidin or streptavidin, an antibody and its antigen, etc.

[0088] In some embodiments, the CD80 ECD fusion molecule comprises the sequence of SEQ ID NO: 20 or 21.

[0089] CD80 ECD fusion molecules may, depending on how they are produced, have different levels of particular glycosylation modifications. For example, a CD80 ECD fusion molecule may have different concentrations of sialic acid residues in relation to the concentration of the CD80 ECD protein. In some embodiments, a higher sialic acid content may have a longer clearance time in the body and thus an increased overall bioavailability. In some embodiments, the sialic acid content of the CD80 ECD fusion molecule is from 10 to 60 mol sialic acid (SA) to mol protein. In some embodiments, the sialic acid content of the CD80 ECD fusion molecule is from 15 to 60 mol sialic acid (SA) to mol protein. For example, in some embodiments, the SA content is 10-40 mol SA/mol protein, such as 15-30 mol SA/mol protein, such as 15-25 mol SA/mol protein, such as 20-40 mol SA/mol protein, such as 20-30 mol SA/mol protein, such as 30-40 mol SA/mol protein, such as 10, 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the SA content is at least 15 mol SA/mol protein, such as at least 20 mol SA/mol protein, at least 25 mol SA/mol protein, at least 30 mol SA/mol protein, at least 35 mol SA/mol protein, or at least 40 mol SA/mol protein. In some such embodiments, the fusion partner is an Fc domain, such as a human IgG1, IgG2, or IgG4 Fc domain.

[0090] In some embodiments, the SA content of the CD80 ECD fusion molecule is increased or is maintained at a relatively high level in comparison to current CD80 ECD fusion molecules. In some embodiments, an increase in SA content, such as by 5, 10, 15, 20, 30, 40 or 50 mol SA to mol of CD80 ECD protein, may lead to an enhanced efficacy in at least one mouse syngeneic or xenograft tumor model. For example, in some embodiments, tumor growth in a mouse tumor model may be further reduced by at least 5%, 10%, 20%, 30%, 40% 50%, 60%, 70%, 80%, 90%, 95%, or 98% when there is an increase in SA content, such as by 5, 10, 15, 20, 30, 40 or 50 mol SA to mol of CD80 ECD protein.

[0091] For example, in some embodiments, a CD80 ECD Fc fusion molecule, such as a fusion molecule comprising a human IgG1 Fc domain comprising between 10 and 60 mol SA/mol protein is capable of at least 80%, such as at least 90%, such as at least 95%, such as at least 98% tumor cell growth inhibition in at least one mouse syngeneic or xenograft cancer model over a period of at least ten days or at least two weeks or at least three weeks, such as ten days to two weeks or two to three weeks following inoculation with tumor cells. In some such embodiments, the molecule comprises at least 15 mol SA/mol protein, such as at least 20 mol SA/mol protein, or a range from 15-30, 15-25, or 20-30 mol SA/mol protein. In some embodiments, the mouse model is a CT26, MC38, or B16 mouse tumor model. In some embodiments, the mice are given one to three doses of the molecule at 0.3 to 3.0 mg/kg, such as at 0.3 to 0.6 mg/kg, for example over a period of one week, once tumors have reached a minimum volume. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the CD80 ECD fusion molecule comprises the sequence of SEQ ID NO: 20 or 21.

[0092] In some embodiments, the CD80 ECD Fc fusion molecule reduces growth of CT26 tumor cells in mice over a period of at least ten days or at least two weeks or at least three weeks, such as ten days to two weeks or two to three weeks, after inoculation by a greater degree than a CD80 ECD Fc fusion protein with the identical amino acid sequence but a lower level of SA per mol of protein. In some embodiments, the CD80 ECD Fc fusion molecule reduces growth of CT26 tumors in mice over a period of at least ten days or at least two weeks, such as over ten days to two weeks or two to three weeks, after inoculation by a greater degree than an anti-CTLA4 antibody, such as anti-CTLA4 antibody clone 9D9. In some such embodiments, the CD80 ECD Fc molecule is dosed one to three times at 0.3 mg/kg, 0.6 mg/kg, or 3.0 mg/kg while the anti-CTLA4 antibody is dosed the same number of times at 1.5 or 10 mg/kg. In some such embodiments, the model is a CT26, MC38, or B16 murine tumor model.

[0093] Example 6 herein, for example, provides data showing that treatment of a mouse syngeneic tumor model with a CD80 ECD fusion molecule having 15 or 20 mol SA/mol protein resulted in at least 93% inhibition of tumor growth after one dose of 0.3 mg/kg, whereas the same treatment with a molecule having only 5 mol SA/mol protein did not significantly inhibit tumor growth. Similarly, a 0.6 mg/kg dose of the CD80

ECD fusion molecule having 15 or 20 mol SA/mol protein resulted in 95% to 98% inhibition of tumor growth, whereas the same treatment with the molecule with 5 mol SA/mol protein inhibited tumor growth by only 70%. (See Fig. 6.) The degree of inhibition was assessed about three weeks following inoculation with the tumors.

[0094] Further, Example 7 herein shows data on a CD80 ECD Fc fusion molecule (a mouse surrogate) having 20 mol SA/mol protein in three different syngeneic mouse tumor models, the CT26, MC38, and B16 models at 0.3 mg/kg (CT26) or 3.0 mg/kg (MC38 and B16) doses compared to an anti-CTLA4 antibody (clone 9D9) at 1.5 mg/kg and 10 mg/kg doses. Each protein was dosed three times over a 7-day period a few days after inoculation with tumor cells, as depicted in Figs. 7-9 (the arrows showing the days of dosing). In each case, the CD80 ECD Fc was superior in tumor growth inhibition to the anti-CTLA4 antibody over the course of the two to three week study. (Figs. 7-9.) For example, in the CT26 model at day 21 after inoculation with tumor cells, the CD80 ECD Fc fusion molecule showed a 90% reduction in tumor growth compared to 75% or 53% for the two dose levels of anti-CTLA4. In the MC38 model at day 19 after inoculation, the CD80 ECD Fc molecule showed about 80% reduction in tumor growth inhibition compared to only 21% tumor growth for the higher dose of anti-CTLA4 and no tumor growth inhibition for the lower anti-CTLA4 dose. In the B16 model, the CD80 ECD Fc fusion molecule showed 41% tumor growth inhibition on day 13 after inoculation while the anti-CTLA4 antibody did not inhibit tumor growth at either dose level. (See Figs. 7-9.)

[0095] Based on these studies, a CD80 ECD fusion molecule may be capable of a certain percentage of tumor growth inhibition over at least a two week period of time, for example, when about two weeks after the mice have been inoculated with the tumor cells, and also following dosing with the fusion molecule, an average tumor growth inhibition at about the stated percentage is observed in the treated mice. A CD80 ECD fusion molecule may be capable of a certain percentage of tumor growth inhibition over a two to three week period of time, for example, when between two and three weeks after the mice have been inoculated with the tumor cells, and also following dosing with the fusion molecule, an average tumor growth inhibition at about the stated percentage is observed in the treated mice.

[0096] Examples 6 and 7 also show that many mice from the CT26 model treated with CD80 ECD fusion molecule had a complete tumor regression over these two to three week time periods. Moreover, a larger percentage of mice had complete tumor regression with CD80 ECD fusion molecule with higher SA content than with the comparison treatments and a larger percentage of mice had complete tumor regression with CD80 ECD fusion molecule than with an anti-CTLA4 antibody. Thus, in some embodiments, treatment CD80 ECD fusion molecule, such as with 0.3 mg/kg to 0.6 mg/kg CD80 ECD fusion molecule or with 0.3 mg/kg to 3.0 mg/kg CD80 ECD fusion molecule, may result in complete tumor regression in mice in a syngeneic or xenograft model such as CT26, MC38, or B16.

Exemplary Fc Domain Fusion Partners

[0097] In some embodiments, the CD80 ECD fusion molecule has an Fc domain as fusion partner. In some embodiments, the Fc domain is derived from human IgG1, IgG2, IgG3, or IgG4. In some embodiments, the Fc domain has a wild-type sequence, such as a wild-type human IgG1 or IgG2 (e.g. IgG2a) sequence. In other embodiments, the Fc domain is either a natural or engineered variant. In some embodiments, an Fc domain is chosen that has altered interactions of the Fc with one or more Fc gamma receptors. In some embodiments, an Fc domain is chosen that has altered interactions of the Fc with one or more complement factors. In some embodiments, an Fc domain is chosen that has altered interactions of the Fc with one or more Fc gamma receptors and that has altered interactions with one or more complement factors.

[0098] In some embodiments, the Fc domain comprises at least one point mutation as described in WO 2014/144960. In some embodiments, the Fc domain is a human Fc domain with a substitution at one or more of positions E233, L234, L235, P238, D265, N297, A327, P329, or P331 (wherein the numbering of these positions is according to the EU index as in Kabat). In some embodiments, the Fc domain is a human Fc domain with a mutation at L234, L235, and/or P331. In some embodiments, the Fc domain is a human Fc domain with the substitutions L234F, L235E, and P331S. (See, e.g., SEQ ID NO:12.) In some embodiments, the Fc domain has an amino acid substitution at position N297. (See, e.g., SEQ ID NO: 13.) In some embodiments, the Fc domain comprises a C237S mutation. (See, e.g., SEQ ID NO: 9.)

[0099] In some embodiments, a mutated Fc fusion partner causes the CD80 ECD Fc fusion molecule to have altered interactions with one or more Fc gamma receptors compared to those of a CD80 ECD fusion molecule with the same amino acid sequence except for the Fc domain mutations. In some embodiment, the Fc has reduced affinity for Fc gamma receptors such as one or more of FcRN, RI, RIIA, RIIB, and RIII compared to a wild-type Fc domain. In some embodiments, the Fc has reduced affinity for all of FcRN, RI, RIIA, RIIB, and RIII compared to a wild-type Fc domain.

[00100] In some embodiments, a mutated Fc fusion partner causes the CD80 ECD Fc fusion molecule to have altered interactions with at one or more complement factors such as C1, C2, C3, C4, and their cleavage products, such as C4a, C4b, C2a, C2b, C3a, and C3b. In some embodiments, a mutated Fc fusion partner causes the CD80 ECD Fc fusion molecule to have altered interactions with one or more complement factors compared to those of a CD80 ECD fusion molecule with the same amino acid sequence except for the Fc domain mutations.

[00101] In some embodiments the CD80 ECD and the fusion partner, such as an Fc fusion partner, are directly linked such that the N- or C-terminal amino acid of the Fc immediately precedes or follows the N- or C-terminal amino acid of the CD80 ECD sequence. (See, e.g., SEQ ID NOs: 20 and 21.) In other embodiments, the CD80 ECD and fusion partner are joined by a linker molecule, such as by a linker peptide sequence, such as by a GS linker sequence.

Therapeutic Compositions and Methods

Methods of Treating Cancer

[00102] In some embodiments, methods for treating cancer are provided, comprising administering an effective amount of a CD80 ECD or CD80 ECD fusion molecule.

[00103] In some embodiments, the cancer may be benign (also referred to as a benign tumor), pre-malignant, or malignant. In some embodiments, the cancer may comprise solid cancer cells (i.e. "solid tumors") or alternatively, it may comprise leukemic cancer cells. In some embodiments, the CD80 ECD or CD80 ECD fusion molecule is effective to reduce cancer growth in a human or animal subject, or in a mouse syngeneic or xenograft model for the cancer being treated. In some embodiments, the CD80 ECD

or CD80 ECD fusion molecule is effective to reduce tumor volume, such as in a mouse syngeneic or xenograft model for the cancer being treated.

[00104] Examples of particular cancers that may be treated include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular nonlimiting examples of such cancers include but are not limited to squamous cell cancer, small-cell lung cancer, pituitary cancer, esophageal cancer, astrocytoma, soft tissue sarcoma, non-small cell lung cancer (including squamous cell non-small cell lung cancer), adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, renal cell carcinoma, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, brain cancer, endometrial cancer, testis cancer, cholangiocarcinoma, gallbladder carcinoma, gastric cancer, melanoma, and various types of head and neck cancer (including squamous cell carcinoma of the head and neck).

[00105] In any of the above method embodiments, the CD80 ECD or CD80 ECD fusion molecule administered to the subject may inhibit tumor growth in a mouse syngeneic xenograft cancer model over a period of 1 week, 10 days, 2 weeks, or 3 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%. In some embodiments, the CD80 ECD fusion molecule may inhibit tumor growth in a CT26 mouse xenograft tumor model by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% at two weeks or at three weeks post-inoculation. In some such cases, the fusion molecule may be dosed one to three times at 0.3 to 3 mg/kg, such as at 0.3 to 0.6 mg/kg. In any of the above method embodiments, administration of the CD80 ECD or CD80 ECD fusion molecule administered to the subject may reduce the volume of at least one tumor in a human or animal subject by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, for example, over a period of one month, two months, three months, six months, or one year. In some cases, the CD80 ECD Fc fusion molecule may be capable of resulting in complete tumor regression in a mouse tumor model such as a

CT26 model, for example in a significant portion of tested mice, such as at least 40%, or at least 50% of mice.

[00106] In any of these methods, the CD80 ECD or CD80 ECD fusion molecule may be a CD80 ECD Fc comprising 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21.

Combination Treatments with Immune Stimulating Agents Including PD-1/PD-L1 Inhibitors

[00107] In some embodiments, the CD80 ECD or CD80 ECD fusion molecule is administered to treat one of the above cancers in combination with an effective amount of at least one immune stimulating agent. Immune stimulating agents may include, for example, a small molecule drug or a biologic. Examples of biologic immune stimulating agents include, but are not limited to, antibodies, antibody fragments, fragments of receptor or ligand polypeptides, for example that block receptor-ligand binding, vaccines and cytokines.

[00108] In some embodiments, the at least one immune stimulating agent comprises an agonist of an immune stimulatory molecule, including a co-stimulatory molecule, while in some embodiments, the at least one immune stimulating agent comprises an antagonist of an immune inhibitory molecule, including a co-inhibitory molecule. In some embodiments, the at least one immune stimulating agent comprises an agonist of an immune-stimulatory molecule, including a co-stimulatory molecule, found on immune cells, such as T cells. In some embodiments, the at least one immune stimulating agent comprises an antagonist of an immune inhibitory molecule, including a co-inhibitory molecule, found on immune cells, such as T cells. In some embodiments,

the at least one immune stimulating agent comprises an agonist of an immune stimulatory molecule, including a co-stimulatory molecule, found on cells involved in innate immunity, such as NK cells. In some embodiments, the at least one immune stimulating agent comprises an antagonist of an immune inhibitory molecule, including a co-inhibitory molecule, found on cells involved in innate immunity, such as NK cells. In some embodiments, the combination enhances the antigen-specific T cell response in the treated subject and/or enhances the innate immunity response in the subject. In some embodiments, the combination results in an improved anti-tumor response in an animal cancer model, such as a syngeneic or xenograft model, compared to administration of either the CD80 ECD or CD80 ECD fusion molecule or immune stimulating agent alone. In some embodiments, the combination results in a synergistic response in an animal cancer model, such as a syngeneic or xenograft model, compared to administration of either the CD80 ECD or CD80 ECD fusion molecule or immune stimulating agent alone.

[00109] In any of the above combination therapy method embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with the immune stimulating agent, such as a PD-1/PD-L1 inhibitor, that is administered to the subject may inhibit tumor growth in a mouse syngeneic or xenograft cancer model over a period of 1 week, 10 days, 2 weeks, or 3 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%. In any of the above combination therapy method embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with the immune stimulating agent, such as a PD-1/PD-L1 inhibitor, that is administered to the subject may reduce the volume of at least one tumor in the subject or in an animal model by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95%, for example, over a period of one month, two months, three months, six months, or one year.

[00110] In any of the combination therapy methods, the CD80 ECD or CD80 ECD fusion molecule may be a CD80 ECD Fc comprising 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30

mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21.

[00111] In certain embodiments, an immune stimulating agent targets a stimulatory or inhibitory molecule that is a member of the immunoglobulin super family (IgSF). For example, an immune stimulating agent may be an agent that targets (or binds specifically to) another member of the B7 family of polypeptides. An immune stimulating agent may be an agent that targets a member of the TNF family of membrane bound ligands or a co-stimulatory or co-inhibitory receptor binding specifically to a member of the TNF family. Exemplary TNF and TNFR family members that may be targeted by immune stimulating agents include CD40 and CD40L, OX-40, OX-40L, GITR, GITRL, CD70, CD27L, CD30, CD30L, 4-1BBL, CD137 (4-1BB), TRAIL/Apo2-L, TRAILR1/DR4, TRAILR2/DR5, TRAILR3, TRAILR4, OPG, RANK, RANKL, TWEAKR/Fn14, TWEAK, BAFFR, EDAR, XEDAR, TACI, APRIL, BCMA, LT β R, LIGHT, DcR3, HVEM, VEGI/TL1A, TRAMP/DR3, EDAR, EDA1, XEDAR, EDA2, TNFR1, Lymphotoxin α /TNF β , TNFR2, TNF α , LT β R, Lymphotoxin α 1 β 2, FAS, FASL, RELT, DR6, TROY and NGFR.

[00112] In some embodiments, an immune stimulating agent may comprise (i) an antagonist of a protein that inhibits T cell activation (e.g., immune checkpoint inhibitor) such as CTLA4, LAG-3, TIM3, Galectin 9, CEACAM-1, BTLA, CD69, Galectin-1, TIGIT, CD113, GPR56, VISTA, B7-H3, B7-H4, 2B4, CD48, GARP, PD1H, LAIR1, TIM-1, TIM-4, and ILT4 and/or may comprise (ii) an agonist of a protein that stimulates T cell activation such as B7-2, CD28, 4-1BB (CD137), 4-1BBL, ICOS, ICOS-L, OX40, OX40L, GITR, GITRL, CD70, CD27, CD40, CD40L, DR3 and CD28H.

[00113] In some embodiments, an immune stimulating agent may comprise an agent that inhibits or is an antagonist of a cytokine that inhibits T cell activation (e.g., IL-6, IL-10, TGF- β , VEGF, and other immunosuppressive cytokines), and in some embodiments an immune stimulating agent may comprise an agent that is an agonist of a

cytokine, such as IL-2, IL-7, IL-12, IL-15, IL-21 and IFN α (e.g., the cytokine itself) that stimulates T cell activation. In some embodiments, immune stimulating agents may comprise an antagonist of a chemokine, such as CXCR2 (e.g., MK-7123), CXCR4 (e.g. AMD3100), CCR2, or CCR4 (mogamulizumab).

[00114] In some embodiments, immune stimulating agents may include antagonists of inhibitory receptors on NK cells or agonists of activating receptors on NK cells. For example, a CD80 ECD or CD80 ECD fusion molecule can be combined with an antagonist of KIR.

[00115] Immune stimulating agents may also include agents that inhibit TGF- β signaling, agents that enhance tumor antigen presentation, e.g., dendritic cell vaccines, GM-CSF secreting cellular vaccines, CpG oligonucleotides, and imiquimod, or therapies that enhance the immunogenicity of tumor cells (e.g., anthracyclines).

[00116] Immune stimulating agents may also include certain vaccines such as mesothelin-targeting vaccines or attenuated listeria cancer vaccines, such as CRS-207.

[00117] Immune stimulating agents may also comprise agents that deplete or block Treg cells, such as agents that specifically bind to CD25.

[00118] Immune stimulating agents may also comprise agents that inhibit a metabolic enzyme such as indoleamine dioxigenase (IDO), dioxigenase, arginase, or nitric oxide synthetase.

[00119] Immune stimulating agents may also comprise agents that inhibit the formation of adenosine or inhibit the adenosine A2A receptor.

[00120] Immune stimulating agents may also comprise agents that reverse/prevent T cell anergy or exhaustion and agents that trigger an innate immune activation and/or inflammation at a tumor site.

[00121] In some embodiments, immune stimulating agents may comprise a CD40 agonist such as a CD40 agonist antibody. The CD80 ECD or CD80 ECD fusion molecule may also be combined with a combinatorial approach that targets multiple elements of the immune pathway, such as one or more of the following: at least one agent that enhances tumor antigen presentation (e.g., dendritic cell vaccine, GM-CSF secreting cellular vaccines, CpG oligonucleotides, imiquimod); at least one agent that inhibits negative immune regulation e.g., by inhibiting CTLA4 pathway and/or depleting or blocking Treg or other immune suppressing cells; a therapy that stimulates positive

immune regulation, e.g., with agonists that stimulate the CD-137, OX-40 and/or GITR pathway and/or stimulate T cell effector function; at least one agent that increases systemically the frequency of anti-tumor T cells; a therapy that depletes or inhibits Tregs, such as Tregs in the tumor, e.g., using an antagonist of CD25 (e.g., daclizumab) or by ex vivo anti-CD25 bead depletion; at least one agent that impacts the function of suppressor myeloid cells in the tumor; a therapy that enhances immunogenicity of tumor cells (e.g., anthracyclines); adoptive T cell or NK cell transfer including genetically modified cells, e.g., cells modified by chimeric antigen receptors (CAR-T therapy); at least one agent that inhibits a metabolic enzyme such as indoleamine dioxigenase (IDO), dioxigenase, arginase or nitric oxide synthetase; at least one agent that reverses/prevents T cell anergy or exhaustion; a therapy that triggers an innate immune activation and/or inflammation at a tumor site; administration of immune stimulatory cytokines or blocking of immunorepressive cytokines.

[00122] For example, a CD80 ECD or CD80 ECD fusion molecule can be used with one or more agonistic agents that ligate positive costimulatory receptors; one or more antagonists (blocking agents) that attenuate signaling through inhibitory receptors, such as antagonists that overcome distinct immune suppressive pathways within the tumor microenvironment; one or more agents that increase systemically the frequency of anti-tumor immune cells, such as T cells, deplete or inhibit Tregs (e.g., by inhibiting CD25); one or more agents that inhibit metabolic enzymes such as IDO; one or more agents that reverse/prevent T cell anergy or exhaustion; and one or more agents that trigger innate immune activation and/or inflammation at tumor sites.

[00123] In one embodiment, the at least one immune stimulating agent comprises a CTLA4 antagonist, such as an antagonistic CTLA4 antibody. Suitable CTLA4 antibodies include, for example, YERVOY (ipilimumab) or tremelimumab.

[00124] In some embodiments, the at least one immune stimulating agent comprises a LAG-3 antagonist, such as an antagonistic LAG-3 antibody. Suitable LAG-3 antibodies include, for example, BMS-986016 (WO10/19570, WO14/08218), or IMP-731 or IMP-321 (WO08/132601, WO09/44273).

[00125] In some embodiments, the at least one immune stimulating agent comprises a CD137 (4-1BB) agonist, such as an agonistic CD137 antibody. Suitable CD137 antibodies include, for example, urelumab or PF-05082566 (WO12/32433).

[00126] In some embodiments, the at least one immune stimulating agent comprises a GITR agonist, such as an agonistic GITR antibody. Suitable GITR antibodies include, for example, TRX-518 (WO06/105021, WO09/009116), MK-4166 (WO11/028683) or a GITR antibody disclosed in WO2015/031667.

[00127] In some embodiments, the at least one immune stimulating agent comprises an OX40 agonist, such as an agonistic OX40 antibody. Suitable OX40 antibodies include, for example, MEDI-6383, MEDI-6469 or MOXR0916 (RG7888; WO06/029879).

[00128] In some embodiments, the at least one immune stimulating agent comprises a CD27 agonist, such as an agonistic CD27 antibody. Suitable CD27 antibodies include, for example, varlilumab (CDX-1127).

[00129] In some embodiments, the at least one immune stimulating agent comprises MGA271, which targets B7H3 (WO11/109400).

[00130] In some embodiments, the at least one immune stimulating agent comprises a KIR antagonist, such as lirilumab.

[00131] In some embodiments, the at least one immune stimulating agent comprises an IDO antagonist. IDO antagonists include, for example, INCB-024360 (WO2006/122150, WO07/75598, WO08/36653, WO08/36642), indoximod, NLG-919 (WO09/73620, WO09/1156652, WO11/56652, WO12/142237) or F001287.

[00132] In some embodiments, the at least one immune stimulating agent comprises a Toll-like receptor agonist, e.g., a TLR2/4 agonist (e.g., Bacillus Calmette-Guerin); a TLR7 agonist (e.g., Hiltonol or Imiquimod); a TLR7/8 agonist (e.g., Resiquimod); or a TLR9 agonist (e.g., CpG7909).

[00133] In some embodiments, the at least one immune stimulating agent comprises a TGF- β inhibitor, e.g., GC1008, LY2157299, TEW7197 or IMC-TR1.

[00134] In some embodiments, the CD80 ECD or CD80 ECD fusion molecule is administered to treat one of the above cancers in combination with an effective amount of a PD-1/PD-L1 inhibitor.

Exemplary PD-1/PD-L1 Inhibitors

[00135] PD-1/PD-L1 inhibitors include antibodies, fusion proteins, and peptides. A nonlimiting exemplary fusion protein that is a PD-1/PD-L1 inhibitor is AMP-224 (Amplimmune, GlaxoSmithKline). A nonlimiting exemplary peptide that is a

PD-1/PD-L1 inhibitor is AUR-012. Other exemplary PD-1/PD-L1 inhibitors include antibodies that inhibit PD-1, such as anti-PD-1 antibodies and anti-PD-L1 antibodies. Such antibodies may be humanized antibodies, chimeric antibodies, mouse antibodies, and human antibodies.

[00136] In some embodiments, the combination results in an improved anti-tumor response in an animal cancer model, such as a xenograft model, compared to administration of either the CD80 ECD or CD80 ECD fusion molecule or PD-1/PD-L1 inhibitor alone. In some embodiments, the combination results in a synergistic response in an animal cancer model, such as a xenograft model, compared to administration of either the CD80 ECD or CD80 ECD fusion molecule or PD-1/PD-L1 inhibitor alone.

[00137] PD-1 is a key immune checkpoint receptor expressed by activated T and B cells and mediates immunosuppression. PD-1 is a member of the CD28 family of receptors, which includes CD28, CTLA4, ICOS, PD-1, and BTLA. Two cell surface glycoprotein ligands for PD-1 have been identified, Programmed Death Ligand-1 (PD-L1) and Programmed Death Ligand-2 (PD-L2). These ligands are expressed on antigen-presenting cells as well as many human cancers and have been shown to down regulate T cell activation and cytokine secretion upon binding to PD-1. Inhibition of the PD-1/PD-L1 interaction mediates potent antitumor activity in preclinical models.

[00138] Human monoclonal antibodies (HuMAbs) that bind specifically to PD-1 with high affinity have been disclosed in U.S. Patent No. 8,008,449. Other anti-PD-1 mAbs have been described in, for example, U.S. Patent Nos. 6,808,710, 7,488,802, 8,168,757 and 8,354,509, and PCT Publication No. WO 2012/145493. Each of the anti-PD-1 HuMAbs disclosed in U.S. Patent No. 8,008,449: (a) binds to human PD-1 with a K_D of 1×10^{-7} M or less, as determined by surface plasmon resonance using a Biacore biosensor system; (b) does not substantially bind to human CD28, CTLA-4 or ICOS; (c) increases T-cell proliferation in a Mixed Lymphocyte Reaction (MLR) assay; (d) increases interferon- γ production in an MLR assay; (e) increases IL-2 secretion in an MLR assay; (f) binds to human PD-1 and cynomolgus monkey PD-1; (g) inhibits the binding of PD-L1 and/or PD-L2 to PD-1; (h) stimulates antigen-specific memory responses; (i) stimulates antibody responses; and/or (j) inhibits tumor cell growth *in vivo*. Anti-PD-1 antibodies usable in the present invention include antibodies that bind specifically to

human PD-1 and exhibit at least one, at least two, at least three, at least four or at least five of the preceding characteristics (a) through (j).

[00139] In one embodiment, the anti-PD-1 antibody is nivolumab. Nivolumab (also known as "Opdivo®"; formerly designated 5C4, BMS-936558, MDX-1106, or ONO-4538) is a fully human IgG4 (S228P) PD-1 immune checkpoint inhibitor antibody that selectively prevents interaction with PD-1 ligands (PD-L1 and PD-L2), thereby blocking the down-regulation of antitumor T-cell functions (U.S. Patent No. 8,008,449; Wang *et al.*, 2014 *Cancer Immunol Res.* 2(9):846-56).

[00140] In another embodiment, the anti-PD-1 antibody is pembrolizumab. Pembrolizumab (also known as "Keytruda®", lambrolizumab, and MK-3475) is a humanized monoclonal IgG4 antibody directed against human cell surface receptor PD-1 (programmed death-1 or programmed cell death-1). Pembrolizumab is described, for example, in U.S. Patent No. 8,900,587; *see also* the site with the address: "www" dot "cancer" dot "gov" slash "drugdictionary?cdrid=695789" (last accessed: December 14, 2014). Pembrolizumab has been approved by the FDA for the treatment of relapsed or refractory melanoma.

[00141] In other embodiments, the anti-PD-1 antibody is MEDI0608 (formerly AMP-514), which is a monoclonal antibody against the PD-1 receptor. MEDI0608 is described, for example, in US Pat. No. 8,609,089, B2 or at www "dot" cancer "dot" gov "slash" drugdictionary?cdrid=756047 (last accessed December 14, 2014).

[00142] In some embodiments, the anti-PD-1 antibody is Pidilizumab (CT-011), which is a humanized monoclonal antibody. Pidilizumab is described in US Pat. No. 8,686,119 B2 or WO 2013/014668 A1.

[00143] Anti-PD-1 antibodies usable in the disclosed methods also include isolated antibodies that bind specifically to human PD-1 and cross-compete for binding to human PD-1 with nivolumab (see, *e.g.*, U.S. Patent No. 8,008,449; WO 2013/173223). The ability of antibodies to cross-compete for binding to an antigen indicates that these antibodies bind to the same epitope region of the antigen and sterically hinder the binding of other cross-competing antibodies to that particular epitope region. These cross-competing antibodies are expected to have functional properties very similar to those of nivolumab by virtue of their binding to the same epitope region of PD-1. Cross-

competing antibodies can be readily identified based on their ability to cross-compete with nivolumab in standard PD-1 binding assays such as Biacore analysis, ELISA assays or flow cytometry (*see, e.g.*, WO 2013/173223).

[00144] In certain embodiments, the antibodies that cross-compete for binding to human PD-1 with, or bind to the same epitope region of human PD-1 as, nivolumab are monoclonal antibodies. For administration to human subjects, these cross-competing antibodies can be chimeric antibodies, or can be humanized or human antibodies.

[00145] Anti-PD-1 antibodies usable in the methods of the disclosed invention also include antigen-binding portions of the above antibodies. Examples include (i) a Fab fragment, a monovalent fragment consisting of the V_L , V_H , C_L and C_{H1} domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the V_H and C_{H1} domains; and (iv) a Fv fragment consisting of the V_L and V_H domains of a single arm of an antibody.

Administration of CD80 ECDs or CD80 ECD Fusion Proteins in Combination with Immune Stimulating Agents or PD-1/PD-L1 Inhibitors

[00146] In some embodiments, the CD80 ECD or CD80 ECD fusion molecule and the immune stimulating agent or PD-1/PD-L1 inhibitor are administered concurrently. In some embodiments, the CD80 ECD or CD80 ECD fusion molecule and the immune stimulating agent or PD-1/PD-L1 inhibitor are administered sequentially. In some embodiments, at least one, at least two, at least three doses, at least five doses, or at least ten doses of a CD80 ECD or CD80 fusion molecule is administered prior to administration of an immune stimulating agent or PD-1/PD-L1 inhibitor. In some embodiments, at least one, at least two, at least three doses, at least five doses, or at least ten doses of an immune stimulating agent or PD-1/PD-L1 inhibitor is administered prior to administration of a CD80 ECD or CD80 fusion molecule. In some embodiments, the last dose of immune stimulating agent or PD-1/PD-L1 inhibitor is administered at least one, two, three, five, days or ten, or one, two, three, five, twelve, or twenty four weeks prior to the first dose of CD80 ECD or CD80 fusion molecule. In some other embodiment, the last dose of CD80 ECD or CD80 fusion molecule is administered at least one, two, three, five, days or ten, or one, two, three, five, twelve, or

twenty four weeks prior to the first dose of immune stimulating agent or PD-1/PD-L1 inhibitor. In some embodiments, a subject has received, or is receiving, immune stimulating agent or PD-1/PD-L1 inhibitor therapy, and a CD80 ECD or CD80 fusion molecule is added to the therapeutic regimen.

[00147] In some embodiments, the subject is an immune stimulating agent or PD-1/PD-L1 inhibitor inadequate responder (i.e. shows resistance to one or more immune stimulating agents or PD-1/PD-L1 inhibitors). A subject who is a PD-1/PD-L1 inhibitor inadequate responder, for example, may have previously responded to a PD-1/PD-L1 inhibitor, but may have become less responsive to the PD-1/PD-L1 inhibitor, or the subject may have never responded to the PD-1/PD-L1 inhibitor. Inadequate response to an immune stimulating agent or PD-1/PD-L1 inhibitor means that aspects of the condition that would be expected to improve following a standard dose of the PD-1/PD-L1 inhibitor do not improve, and/or improvement only occurs if greater than a standard dose is administered. In some embodiments, an immune stimulating agent or PD-1/PD-L1 inhibitor inadequate responder has experienced, or is experiencing, an inadequate response to the drug after receiving a standard dose for at least two weeks, at least three weeks, at least four weeks, at least six weeks, or at least twelve weeks. A “standard” dose of an immune stimulating agent or PD-1/PD-L1 inhibitor may be determined by a medical professional, and may depend on the subject’s age, weight, healthy history, severity of disease, the frequency of dosing, etc. In some embodiments, an immune stimulating agent or PD-1/PD-L1 inhibitor inadequate responder has experienced, or is experiencing, an inadequate response to an anti-PD-1 antibody and/or an anti-PD-L1 antibody. In some embodiments, a PD-1/PD-L1 inhibitor inadequate responder has experienced, or is experiencing, an inadequate response to AMP-224. In some embodiments, a PD-1/PD-L1 inhibitor inadequate responder has experienced, or is experiencing, an inadequate response to a PD-1/PD-L1 inhibitor selected from nivolumab, pidilizumab, and pembrolizumab.

[00148] In any of the above embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with the PD-1/PD-L1 inhibitor that is administered to the subject may inhibit tumor growth in a mouse syngeneic or xenograft cancer model over a period of 1 week, 10 days, or 2 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least

80%, at least 90%, at least 95%, or at least 98%. In any of the above combination therapy method embodiments, the combination of the CD80 ECD or CD80 ECD fusion molecule with the PD-1/PD-L1 inhibitor that is administered to the subject may reduce the volume of at least one tumor in the subject or in an animal model subject by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, for example, over a period of one month, two months, three months, six months, or one year.

[00149] In any of these combination therapy methods, the CD80 ECD or CD80 ECD fusion molecule may be a CD80 ECD Fc comprising 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments, the Fc domain comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21.

Routes of Administration and Carriers

[00150] In various embodiments, polypeptides and fusion molecules may be administered *in vivo* by various routes, including, but not limited to, oral, intra-arterial, parenteral, intranasal, intravenous, intramuscular, intracardiac, intraventricular, intratracheal, buccal, rectal, intraperitoneal, intradermal, topical, transdermal, and intrathecal, or otherwise by implantation or inhalation. The subject compositions may be formulated into preparations in solid, semi-solid, liquid, or gaseous forms; including, but not limited to, tablets, capsules, powders, granules, ointments, solutions, suppositories, enemas, injections, inhalants, and aerosols. A nucleic acid molecule encoding a polypeptide may be coated onto gold microparticles and delivered intradermally by a particle bombardment device, or "gene gun," as described in the literature (see, e.g., Tang et al., *Nature* 356:152-154 (1992)). The appropriate formulation and route of administration may be selected according to the intended application.

[00151] In various embodiments, polypeptide-comprising compositions are provided in formulations with a wide variety of pharmaceutically acceptable carriers (see, e.g., Gennaro, *Remington: The Science and Practice of Pharmacy with Facts and Comparisons: Drugfacts Plus*, 20th ed. (2003); Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th ed., Lippencott Williams and Wilkins (2004); Kibbe et al., *Handbook of Pharmaceutical Excipients*, 3rd ed., Pharmaceutical Press (2000)). Various pharmaceutically acceptable carriers, which include vehicles, adjuvants, and diluents, are available. Moreover, various pharmaceutically acceptable auxiliary substances, such as pH adjusting and buffering agents, tonicity adjusting agents, stabilizers, wetting agents and the like, are also available. Non-limiting exemplary carriers include saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof.

[00152] In various embodiments, compositions comprising polypeptides and fusion molecules may be formulated for injection, including subcutaneous administration, by dissolving, suspending, or emulsifying them in an aqueous or nonaqueous solvent, such as vegetable or other oils, synthetic aliphatic acid glycerides, esters of higher aliphatic acids, or propylene glycol; and if desired, with conventional additives such as solubilizers, isotonic agents, suspending agents, emulsifying agents, stabilizers and preservatives. In various embodiments, the compositions may be formulated for inhalation, for example, using pressurized acceptable propellants such as dichlorodifluoromethane, propane, nitrogen, and the like. The compositions may also be formulated, in various embodiments, into sustained release microcapsules, such as with biodegradable or non-biodegradable polymers. A non-limiting exemplary biodegradable formulation includes poly lactic acid-glycolic acid polymer. A non-limiting exemplary non-biodegradable formulation includes a polyglycerin fatty acid ester. Certain methods of making such formulations are described, for example, in EP 1 125 584 A1.

[00153] Pharmaceutical packs and kits comprising one or more containers, each containing one or more doses of a polypeptide or combination of polypeptides are also provided. In some embodiments, a unit dosage is provided wherein the unit dosage contains a predetermined amount of a composition comprising a polypeptide or combination of polypeptides, with or without one or more additional agents. In some embodiments, such a unit dosage is supplied in a single-use prefilled syringe for injection. In various embodiments, the composition contained in the unit dosage may comprise

saline, sucrose, or the like; a buffer, such as phosphate, or the like; and/or be formulated within a stable and effective pH range. Alternatively, in some embodiments, the composition may be provided as a lyophilized powder that may be reconstituted upon addition of an appropriate liquid, for example, sterile water. In some embodiments, the composition comprises one or more substances that inhibit protein aggregation, including, but not limited to, sucrose and arginine. In some embodiments, a composition of the invention comprises heparin and/or a proteoglycan.

[00154] Pharmaceutical compositions are administered in an amount effective for treatment or prophylaxis of the specific indication. The therapeutically effective amount is typically dependent on the weight of the subject being treated, his or her physical or health condition, the extensiveness of the condition to be treated, or the age of the subject being treated. In some embodiments, a PD-1/PD-L1 inhibitor, such as an antibody or fusion protein, is administered with the CD80 ECD or CD80 ECD fusion molecule at a dose of 1 to 4 mg/kg. In some embodiments, a PD-1/PD-L1 inhibitor is administered at a dose of 1, 2, 3, or 4 mg/kg.

[00155] Determination of the frequency of administration may be made by persons skilled in the art, such as an attending physician based on considerations of the condition being treated, age of the subject being treated, severity of the condition being treated, general state of health of the subject being treated and the like. In some embodiments, an effective dose of a CD80 ECD or CD80 ECD fusion molecule is administered to a subject one or more times. In various embodiments, an effective dose is administered to the subject once a month, less than once a month, such as, for example, every two months or every three months. In other embodiments, an effective dose is administered more than once a month, such as, for example, every three weeks, every two weeks or every week. In some embodiments, an effective dose is administered once per 1, 2, 3, 4, or 5 weeks. In some embodiments, an effective dose is administered twice or three times per week. An effective dose is administered to the subject at least once. In some embodiments, the effective dose may be administered multiple times, including for periods of at least a month, at least six months, or at least a year.

Additional Combination Therapies

[00156] CD80 ECDs or CD80 ECD fusion molecules may be administered alone, with PD-1/PD-L1 inhibitors, and/or with other modes of treatment. CD80

ECDs or CD80 ECD fusion molecules may be provided before, substantially contemporaneous with, or after other modes of treatment, for example, surgery, chemotherapy, radiation therapy, or the administration of another biologic. In some embodiments, the cancer has recurred or progressed following a therapy selected from surgery, chemotherapy, and radiation therapy, or a combination thereof.

[00157] For treatment of cancer, CD80 ECDs or CD80 ECD fusion molecules may be administered in conjunction with one or more additional anti-cancer agents, such as the chemotherapeutic agent, growth inhibitory agent, anti-angiogenesis agent and/or anti-neoplastic composition. Nonlimiting examples of chemotherapeutic agent, growth inhibitory agent, anti-angiogenesis agent, anti-cancer agent and anti-neoplastic composition that can be used in combination with the antibodies of the present invention are provided in the following definitions.

[00158] In any of the above combination therapy method embodiments, the therapy administered to the subject may inhibit tumor growth in a mouse syngeneic or xenograft cancer model over a period of 1 week, 10 days, or 2 weeks, for example, by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95%. In any of the above combination therapy method embodiments, the therapy administered to the subject may reduce the volume of at least one tumor in the subject or in an animal model subject by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98%, for example, over a period of one month, two months, three months, six months, or one year.

[00159] In any of these further combination therapy methods, the CD80 ECD or CD80 ECD fusion molecule may be a CD80 ECD Fc comprising 10-60 mol SA to mol of CD80 ECD Fc protein, such as 15-60 mol SA/mol protein. In some embodiments, the content is 10-40 mol SA/mol protein, such as 15-40 mol SA/mol protein, such as 20-40 mol SA/mol protein, 20-30 mol SA/mol protein, 15-25 mol SA/mol protein, 15-30 mol SA to mol of protein, or 30-40 mol SA/mol protein. In some embodiments, the SA content is at least 15, such as at least 20, at least 25, at least 30, at least 35, or at least 40 mol SA/mol protein. In some embodiments, the SA content is 15, 20, 25, 30, 35, or 40 mol SA/mol protein. In some embodiments, the Fc domain is a human IgG1, IgG2, or IgG4 Fc domain. In some embodiments the Fc domain

comprises the amino acid sequence of SEQ ID NO:14. In some embodiments, the fusion molecule comprises the amino acid sequence of SEQ ID NO:20 or 21.

[00160] A “**chemotherapeutic agent**” is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include, but are not limited to, alkylating agents such as thiotepa and Cytosan[®] cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (*e.g.*, calicheamicin, especially calicheamicin gamma1I and calicheamicin omega1I (*see, e.g.*, Agnew, *Chem Intl. Ed. Engl.*, 33: 183-186 (1994))); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, Adriamycin[®] doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine,

6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK[®] polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, *e.g.*, Taxol[®] paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), Abraxane[®] Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Illinois), and Taxotere[®] doxetaxel (Rhône-Poulenc Rorer, Antony, France); chloranbucil; Gemzar[®] gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin, oxaliplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; Navelbine[®] vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (Camptosar, CPT-11) (including the treatment regimen of irinotecan with 5-FU and leucovorin); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; combretastatin; leucovorin (LV); oxaliplatin, including the oxaliplatin treatment regimen (FOLFOX); inhibitors of PKC-alpha, Raf, H-Ras, EGFR (*e.g.*, erlotinib (Tarceva[®])) and VEGF-A that reduce cell proliferation and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[00161] Further nonlimiting exemplary chemotherapeutic agents include anti-hormonal agents that act to regulate or inhibit hormone action on cancers such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for

example, tamoxifen (including Nolvadex[®] tamoxifen), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and Fareston[®] toremifene; aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, Megase[®] megestrol acetate, Aromasin[®] exemestane, formestanie, fadrozole, Rivisor[®] vorozole, Femara[®] letrozole, and Arimidex[®] anastrozole; and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those which inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Ralf and H-Ras; ribozymes such as a VEGF expression inhibitor (*e.g.*, Angiozyme[®] ribozyme) and a HER2 expression inhibitor; vaccines such as gene therapy vaccines, for example, Allovectin[®] vaccine, Leuvectin[®] vaccine, and Vaxid[®] vaccine; Proleukin[®] rIL-2; Lurtotecan[®] topoisomerase 1 inhibitor; Abarelix[®] rmRH; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[00162] An “**anti-angiogenesis agent**” or “**angiogenesis inhibitor**” refers to a small molecular weight substance, a polynucleotide (including, *e.g.*, an inhibitory RNA (RNAi or siRNA)), a polypeptide, an isolated protein, a recombinant protein, an antibody, or conjugates or fusion proteins thereof, that inhibits angiogenesis, vasculogenesis, or undesirable vascular permeability, either directly or indirectly. It should be understood that the anti-angiogenesis agent includes those agents that bind and block the angiogenic activity of the angiogenic factor or its receptor. For example, an anti-angiogenesis agent is an antibody or other antagonist to an angiogenic agent, *e.g.*, antibodies to VEGF-A (*e.g.*, bevacizumab (Avastin[®])) or to the VEGF-A receptor (*e.g.*, KDR receptor or Flt-1 receptor), anti-PDGFR inhibitors such as Gleevec[®] (Imatinib Mesylate), small molecules that block VEGF receptor signaling (*e.g.*, PTK787/ZK2284, SU6668, Sutent[®]/SU11248 (sunitinib malate), AMG706, or those described in, *e.g.*, international patent application WO 2004/113304). Anti-angiogenesis agents also include native angiogenesis inhibitors, *e.g.*, angiostatin, endostatin, *etc.* See, *e.g.*, Klagsbrun and D’Amore (1991) *Annu. Rev. Physiol.* 53:217-39; Streit and Detmar (2003) *Oncogene* 22:3172-3179 (*e.g.*, Table 3 listing anti-angiogenic therapy in malignant melanoma); Ferrara & Alitalo (1999) *Nature Medicine* 5(12):1359-1364; Tonini *et al.* (2003) *Oncogene* 22:6549-6556

(*e.g.*, Table 2 listing known anti-angiogenic factors); and, Sato (2003) *Int. J. Clin. Oncol.* 8:200-206 (*e.g.*, Table 1 listing anti-angiogenic agents used in clinical trials).

[00163] A “**growth inhibitory agent**” as used herein refers to a compound or composition that inhibits growth of a cell (such as a cell expressing VEGF) either *in vitro* or *in vivo*. Thus, the growth inhibitory agent may be one that significantly reduces the percentage of cells (such as a cell expressing VEGF) in S phase. Examples of growth inhibitory agents include, but are not limited to, agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), taxanes, and topoisomerase II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C. Further information can be found in Mendelsohn and Israel, eds., *The Molecular Basis of Cancer*, Chapter 1, entitled "Cell cycle regulation, oncogenes, and antineoplastic drugs" by Murakami *et al.* (W.B. Saunders, Philadelphia, 1995), *e.g.*, p. 13. The taxanes (paclitaxel and docetaxel) are anticancer drugs both derived from the yew tree. Docetaxel (Taxotere[®], Rhone-Poulenc Rorer), derived from the European yew, is a semisynthetic analogue of paclitaxel (Taxol[®], Bristol-Myers Squibb). Paclitaxel and docetaxel promote the assembly of microtubules from tubulin dimers and stabilize microtubules by preventing depolymerization, which results in the inhibition of mitosis in cells.

[00164] The term “**anti-neoplastic composition**” refers to a composition useful in treating cancer comprising at least one active therapeutic agent. Examples of therapeutic agents include, but are not limited to, *e.g.*, chemotherapeutic agents, growth inhibitory agents, cytotoxic agents, agents used in radiation therapy, anti-angiogenesis agents, other cancer immunotherapeutic agents aside from PD-1/PD-L1 inhibitors, apoptotic agents, anti-tubulin agents, and other agents to treat cancer, such as anti-HER-2 antibodies, anti-CD20 antibodies, an epidermal growth factor receptor (EGFR) antagonist (*e.g.*, a tyrosine kinase inhibitor), HER1/EGFR inhibitor (*e.g.*, erlotinib (Tarceva[®]), platelet derived growth factor inhibitors (*e.g.*, Gleevec[®] (Imatinib Mesylate)), a COX-2 inhibitor (*e.g.*, celecoxib), interferons, CTLA-4 inhibitors (*e.g.*, anti-CTLA antibody ipilimumab (YERVOY[®])), PD-L2 inhibitors (*e.g.*, anti-PD-L2 antibodies),

TIM3 inhibitors (*e.g.*, anti-TIM3 antibodies), cytokines, antagonists (*e.g.*, neutralizing antibodies) that bind to one or more of the following targets ErbB2, ErbB3, ErbB4, PDGFR-beta, BlyS, APRIL, BCMA, PD-L2, CTLA-4, TIM3, or VEGF receptor(s), TRAIL/Apo2, and other bioactive and organic chemical agents, *etc.* Combinations thereof are also included in the invention.

EXAMPLES

[00165] The examples discussed below are intended to be purely exemplary of the invention and should not be considered to limit the invention in any way. The examples are not intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (for example, amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1: A CD80 ECD Fc Fusion Molecule Reduces Tumor Growth in Mice Implanted with Murine Colorectal Carcinoma Cell Line CT26

[00166] Seven-week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for two weeks before the start of the study. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and Matrigel® at 5×10^6 cells per milliliter.

[00167] Mice were monitored twice-weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: Tumor volume (mm³) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to treatment groups. The mean tumor volume for all

animals enrolled into treatment groups was 175 mm³. Mice were administered Saline or plasmid DNA via RIPPSSM. Plasmid DNA that was administered via RIPPSSM contained the sequence for the extracellular domain (ECD) of murine CD80 or CTLA4 as well as the Fc domain of human IgG2a. Tumors continued to be measured at least twice per week until tumor volume exceeded 10% of animal weight, or approximately 2000 mm³.

[00168] The change in tumor size is shown by graphing mean tumor volume relative to the day upon which animals were inoculated with CT26 cells. (**Fig. 1a.**) RIPPS with mouse CD80 ECD significantly reduced tumor growth compared to Saline control ($p < 0.05$) beginning on Day 11. (**Fig. 1a-b.**) *P*-values were calculated using unpaired, two-tailed t-test analyses of the calculated tumor volumes on each day of the study (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$). Tumor growth inhibition by CD80 ECD was determined to be 78.8% compared to saline control, which was calculated as $100 \times (1 - (\text{mean change in tumor volume for CD80} / \text{mean change in tumor volume for saline}))$. As also shown in **Fig. 1a**, RIPPS with mouse CTLA4-ECD Fc actually enhanced tumor growth compared to the saline control. One explanation for this result is that the CTLA4-ECD Fc construct might have acted as a ligand trap for CD80, preventing CD80 from binding to CD28 and stimulating T cell activity against tumor cells.

Example 2: A CD80 ECD Fc Fusion Molecule in Combination with an Anti-PD-1 Antibody Reduces Tumor Growth in Mice Implanted with Murine Colorectal Carcinoma Cell Line CT26

[00169] Seven-week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for 12 days before the start of the study. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and Matrigel[®] at 5×10^6 cells per milliliter.

[00170] Mice were monitored twice-weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: Tumor volume (mm^3) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to treatment groups. The mean tumor volume for all animals enrolled into treatment groups was approximately 150 mm^3 . Mice were administered plasmid DNA coding for mouse CD80 ECD plus Fc from human IgG2a or Fc alone (negative control) via RIPPSSM. Protein was administered for anti-PD1 (Clone RMP1-14) or Rat IgG2a (Clone 2A3, negative control). Dosing groups were as follows: 1) Fc RIPPSSM plus Rat IgG2a, 2) CD80 RIPPSSM plus Rat IgG2a, 3) Fc RIPPSSM plus anti-PD1, or 4) CD80 RIPPSSM plus anti-PD1. Tumors continued to be measured at least twice per week until tumor volume exceeded 10% of animal weight, or approximately 2000 mm^3 .

[00171] The change in tumor size is shown by graphing mean tumor weight for all groups at the end of the study. (**Fig. 2a.**) RIPPSSM with mouse CD80 ECD or administration of anti-PD1 reduced tumor growth compared to control. (**Fig. 2a-b.**) The combination of CD80 RIPPSSM and anti-PD1 resulted in significantly reduced tumor growth compared to either CD80 ($p < 0.01$ beginning after Day 9) or anti-PD-1 ($p < 0.01$ on Day 14) alone ($p < 0.05$) beginning nine days after treatment initiation. *P*-values were calculated using unpaired, two-tailed t-test analyses of the calculated tumor volumes on each day of measurement.

[00172] Tumor growth inhibition (TGI) by CD80 ECD and anti-PD1 were determined to be 28.7% and 41.5%, respectively, compared to control. TGI by the combination of CD80 ECD and anti-PD1 combination was determined to be 83%. (See **Fig. 2b.**) TGI was calculated using the formula $100 \times (1 - (\text{Mean } \Delta \text{volume Treatment group} / \text{Mean } \Delta \text{volume Saline}))$.

Example 3: Comparison of Activity of CD80 ECD Fc Fusion Molecules with Wild-Type and Mutant Human IgG1 Fc Fusion Polypeptide Sequences

[00173] Seven-week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for two weeks before the start of

the study. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and Matrigel® at 5×10^6 cells per milliliter.

[00174] Mice were monitored twice-weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: Tumor volume (mm^3) = (width (mm) x length (mm))²/2. On Day 5, all tumors were measured, and mice were randomly assigned to treatment groups. The mean tumor volume for all animals enrolled into treatment groups was 175 mm^3 . Mice were administered Saline or plasmid DNA via RIPPSSM. Plasmid DNA that was administered via RIPPSSM contained the sequence for the extracellular domain (ECD) of murine CD80 with the Fc domain of human IgG1. Two clones were administered via RIPPSSM, one with the wild type human IgG1 Fc (CD80-IgG1 WT) and the other with this Fc bearing three mutated amino acids (L234F/L235E/P331S) in order to alter the interaction of Fc with Fc gamma receptors (CD80-IgG1 MT). Tumors continued to be measured at least twice per week until tumor volume exceeded 10% of animal weight, or approximately 2000 mm^3 .

[00175] The change in tumor size is shown by graphing mean tumor volume relative to the day upon which animals were inoculated with CT26 cells. (**Fig. 3a-b.**) RIPPSSM with CD80-IgG1 WT significantly reduced tumor growth compared to Saline control ($p < 0.05$) beginning on Day 14. CD80-IgG MT significantly reduced tumor growth compared to CD80-IgG1 WT ($p < 0.01$) beginning on Day 14. *P*-values were calculated using unpaired, two-tailed t-test analyses of the calculated tumor volumes on each day of the study (* $p < 0.05$, ** $p < 0.01$). Tumor growth inhibition (TGI) by CD80-IgG1 WT was determined to be 69.4% compared to Saline control, compared to TGI for CD80-IgG MT, which was calculated as 98%. (See **Fig. 3b.**) TGI was determined using the formula $100 \times (1 - (\text{Mean } \Delta \text{volume Treatment group} / \text{Mean } \Delta \text{volume Saline}))$.

Example 4: Effects of CD80 ECD Fc Fusion Molecules with Wild-Type and Mutant Human IgG1 Fc Fusion Polypeptide Sequences on Infiltrating T Cells in CT26 Tumors

[00176] A separate *in vivo* study was conducted to analyze the effects of the CD80-IgG1 WT and CD80-IgG1 MT on infiltrating T cells in CT26 tumors at an early stage of the treatment.

[00177] Seven-week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for two weeks before the start of the study. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2 mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and Matrigel® at 5×10^6 cells/mL.

[00178] Mice were monitored twice-weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: Tumor volume (mm³) = (width (mm) x length (mm))²/2. On Day 5, all tumors were measured, and mice were randomly assigned to treatment groups (n= 5 mice per group). The mean tumor volume for all animals enrolled into treatment groups was 72 mm³. Mice were administered saline, a murine extracellular domain (ECD) of CD80 with a wild type human IgG1 Fc (CD80-IgG1 WT), or a murine ECD of CD80 with a mutated human IgG1 Fc (CD80-IgG1 MT, mutated L234F/L235E/P331S) in order to alter the interaction with Fc gamma receptors. Tumors were measured on Days 5 and 11.

[00179] On Day 12, mice were euthanized with CO₂ and perfused with phosphate-buffered saline (PBS), pH 7.4. Briefly, the mouse chest was opened rapidly, and a syringe with 20-gauge needle was used to infuse 40mL of PBS into the aorta via an incision in the left ventricle. Blood and PBS exited through an opening in the right atrium. The tumors were removed and immersed in 10% neutral buffered formalin at 4°C. After 2 hours the tissues were rinsed 3 times in PBS and then transferred in 30%

sucrose in PBS overnight. The next day the tumors were frozen in OCT compound and stored at -80C.

[00180] Cryostat sections were cut at 20- μ m in thickness. Sections were dried on Superfrost[®] Plus slides for 1 to 2 hours. Specimens were permeabilized with PBS containing 0.3% Triton[®] X-100 and incubated in 5% goat normal serum in PBS 0.3% Triton[®] X-100 (blocking solution) for 1 hour at room temperature to block nonspecific antibodies binding. To detect T-cells, sections were incubated with rat-anti-CD4 (GK1.5/ eBiosciences) and rabbit anti-CD3 antibody (SP7/ Thermo Scientific) both diluted 1:500 in blocking solution overnight. Control specimens were incubated in 5% normal serum instead of primary antibody for the same period. After rinsing with PBS containing 0.3% Triton[®] X-100, specimens were incubated for 4 hours at room temperature with Alexa[®] 594-labeled goat anti-rat and Alexa[®] 488-labeled goat anti-rabbit secondary antibodies diluted 1:400 in PBS (Jackson ImmunoResearch). Specimens were rinsed with PBS containing 0.3% Triton[®] X-100, fixed in 1% paraformaldehyde (PFA), rinsed again with PBS, and mounted in Vectashield antifade mounting medium with DAPI (Vector laboratories).

[00181] Specimens were examined with a Zeiss Axiophot[®] 2 plus fluorescence microscope equipped with AxioCam[®] HRc camera. Representative images for each experimental group showing the amount and distribution of the CD3⁺ and CD4⁺ cells within the tumor were collected and are shown in **Figs. 4a and 4b**.

[00182] Treatment with CD80-IgG1 WT or CD80-IgG1 MT increased the number of intratumoral CD3⁺ and CD4⁺ cells compared to saline (**Figs. 4a and 4b**). While the amount of CD4⁺ cells was similar between the CD80-IgG1 WT or CD80-IgG1 MT treated tumors, treatment with CD80-IgG1 MT led to a greater increase of tumor infiltrating CD3⁺ T cells compared to CD80-IgG1 WT (**Fig. 4b**). The ratio of CD3⁺ to CD4⁺ cells was increased with the CD80-IgG1 MT compared to the CD80-IgG1 WT.

Example 5: Cytokine Release Effects of a CD80 ECD Fc Fusion Molecule

Methods

Protein Treatments

[00183] A human CD80 ECD IgG1 Fc fusion molecule (CD80-Fc) was bound to magnetic protein-A beads (Life Technologies) in T-cell proliferation media containing RPMI 1640, 100 IU Penicillin/100 ug/ml Streptomycin, 2 mM L-Glutamine, 100 nM non-essential amino acids, 55 uM 2-mercaptoethanol and 10% ultra low-IgG fetal bovine serum. Binding reactions were carried out in 96 well flat-bottom tissue culture plates at a volume of 100ul per well with a bead concentration of 3 million beads per ml. CD80-Fc was bound to the beads across a series of concentrations: 10, 1, 0.1 ug/ml. An additional set of binding reactions was also performed with the addition of 3 ng/ml OKT3-scFv. Proteins were allowed to bind for 1 hour at room temperature on a rocking platform, following which 100ul of 20 ug/ml (final concentration 10 ug/ml) IgG1 Free-Fc (FPT) was added to each well and allowed to bind for an additional hour in order to block any unoccupied Protein-A binding sites on the beads. The fully loaded and blocked beads were then washed 3 times with PBS using a magnetic 96-well plate stand in order to remove unbound proteins. 100ul of Human Pan T-cells at a concentration of 1×10^6 cells/ml was then added to each well of dry, washed beads. Each condition was tested in triplicate.

Cells

[00184] Human PBMCs were isolated from apheresis-enriched blood (buffy coats) collected from healthy donors ~18hrs prior to isolation using Ficoll® (Biochrom) gradient density centrifugation. Pan T-cells were then isolated from PBMCs using a Human Pan T-cell isolation kit (Miltenyi). T-cells were seeded at a density of 1 million cells/ml in T225 tissue culture flasks in proliferation media (above) supplemented with 8 ng/ml IL-2 and Human T-cell Activator Dynabeads® (Life Tech) 1 bead/cell. Following seeding, cells were fed with fresh IL-2 and continually kept at a concentration of 0.3 million cells/ml by the addition of fresh proliferation media every 2 days. Cells were kept in a 37°C water-jacketed incubator maintained at 5% CO₂. After 6 days of expansion, the activator-beads were removed using a magnetic tube stand and the cells were resuspended at a concentration of 1 million cells/ml in fresh proliferation media without IL-2. 24 hours later the cells were put into assay with Protein-A bead immobilized proteins.

Cytokine Measurements

[00185] Soluble Interferon Gamma (IFN- γ) and Tumor Necrosis Factor Alpha (TNF- α) levels were measured in the supernatants using HTRF-ELISA kits (Cisbio) 24 hours after the cells had been treated with the Protein-A bead immobilized proteins according to the manufacturer's instructions.

Results

[00186] Bead-immobilized CD80-Fc alone did not cause significant human T-cell activation, as measured by soluble cytokine production (**Figs. 5a & c**). However, when a small amount of OKT3-scFv was immobilized along with CD80-Fc, robust CD80-dependent IFN- γ and TNF- α release was observed (**Figs. 5b & d**). The amount of OKT3-scFv used here was too low to cause T-cell stimulation on its own and therefore required the presence of CD80 as a co-stimulatory protein. These results therefore confirm the CD80-Fc used in this assay was indeed biologically active.

[00187] While release of IFN- γ and TNF- α in this assay showed that the CD80-Fc was biologically active, an excessive release of cytokines such as IFN- γ and TNF- α can be harmful. Thus, to address the potential safety of CD80 ECD Fc treatment, these results were compared to earlier published results with TGN1412, a monoclonal anti-CD28 antibody that was shown to be a T-cell "superagonist" and to release excessive and harmful levels of cytokines such as IFN- γ and TNF- α in human subjects.

[00188] Immobilized TGN1412 alone appears to be significantly more potent at inducing cytokine release from human T-cells than human CD80 alone. Findlay et al., *J. Immunological Methods* 352: 1-12 (2010), reported that 1 μ g/well of TGN1412 caused robust TNF α release, ~2000pg/ml, and Vessillier et al., *J. Immunological Methods* 424: 43-52 (2015), reported the same amount of TGN1412 caused robust IFN- γ , ~10000 pg/ml. In our assay, the same amount of immobilized CD80-Fc did not cause significant release of either cytokine. These results suggest that CD80-Fc is at least 1000-fold less potent at inducing cytokine release compared to TGN1412 and therefore poses a significantly lower risk of inducing cytokine storm in humans than TGN1412.

Example 6: Effects of a CD80 ECD Fc Fusion Molecule on CT26 Tumors *In Vivo* with Fc Domains with Different Sialic Acid (SA) Content

[00189] An *in vivo* study was conducted in CT26 tumors to analyze the effects of three different lots of CD80 ECD fused to wild-type human IgG1 Fc having different sialic acid (SA) contents. Specifically, lot E of the CD80 ECD Fc contains 20 mol SA/mol protein, lot D contains 15 mol SA/mol protein, and lot A contains 5 mol SA/mol protein.

[00190] Seven week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for one week before the study was initiated. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and Matrigel® at 5×10^6 cells per milliliter.

[00191] Mice were monitored for tumor growth twice weekly following cell implantation. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: tumor volume (mm³) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to seven treatment groups (n= 10 mice per experimental group). The mean tumor volume for all animals enrolled was 94 mm³. The first group was injected with 200 μ l of PBS (control) intravenously (i.v.) into the tail vein. The second group was injected with CD80 ECD Fc at 20 mol SA/mol protein (lot E) i.v. dosed at 0.3 mg/kg. The third group was injected with CD80 ECD Fc at 20 mol SA/mol protein (lot E) i.v. dosed at 0.6 mg/kg. The fourth group was injected with CD80 ECD Fc at 15 mol SA/mol protein (lot D) i.v. dosed at 0.3 mg/kg. The fifth group was injected with CD80 ECD Fc at 15 mol SA/mol protein (lot D) i.v. dosed at 0.6 mg/kg. The sixth group was injected with CD80 ECD Fc at 5 mol SA/mol protein (lot A) i.v. dosed at 0.3 mg/kg. The seventh group was injected with CD80 ECD Fc at 5 mol

SA/mol protein (lot A) i.v. dosed at 0.6 mg/kg. Tumors were measured on day 10, 14, 16, 18, 22, 24.

[00192] Treatment with CD80 ECD Fc at 20 mol SA/mol protein (lot E) dosed at 0.3 or 0.6 mg/kg resulted in a 93% and 98% inhibition of tumor growth compared to the control ($P < 0.001$). Treatment with CD80 ECD Fc at 15 mol SA/mol protein (lot D) dosed at 0.3 or 0.6 mg/kg resulted in a 93% and 95% inhibition of tumor growth compared to the control ($P < 0.001$). By comparison, treatment with CD80 ECD Fc lot A at 0.3mg/kg (with 5 mol SA/mol protein) did not inhibit tumor growth compared to the control and when dosed at 0.6mg/kg it only induced 70% inhibition ($P < 0.001$) (**Fig 6**).

[00193] The incidence of tumor-free mice was analyzed at day 37. Treatment with CD80 ECD-Fc at 20 mol/mol SA (lot E) dosed at 0.3 or 0.6 mg/kg led to complete tumor regression in 8/10 (80%) or 10/10 (100%) of the mice. Treatment with CD80 ECD-Fc at 15 mol/mol SA (lot D) dosed at 0.3 or 0.6 mg/kg led to complete tumor regression in 9/10 (90%) of the mice. By comparison, treatment with CD80 ECD-Fc lot A dosed at 0.6mg/kg induced tumor regression only in 1/10 (10%) of the mice, as shown in the table below.

Treatment group	Tumor free mice at day 37
Saline	0% (0/10 mice)
CD80 ECD-Fc SA 20 mol/mol (lot E) at 0.3mg/kg 1 dose	80% (8/10 mice)
CD80 ECD-Fc SA 20 mol/mol (lot E) at 0.6mg/kg 1 dose	100% (10/10 mice)
CD80 ECD-Fc SA 15 mol/mol (lot D) at 0.3mg/kg 1 dose	90% (9/10 mice)
CD80 ECD-Fc SA 15 mol/mol (lot D) at 0.6mg/kg 1 dose	90% (9/10 mice)
CD80 ECD-Fc SA 5 mol/mol (lot A) at 0.3mg/kg 1 dose	0% (0/10 mice)
CD80 ECD-Fc SA 5 mol/mol (lot A) at 0.6mg/kg 1 dose	10% (1/10 mice)

Example 7: Effects of a Murine CD80 ECD – Murine Fc Fusion Molecule on Tumor Growth in Three Different Syngeneic Tumor Models

[00194] *In vivo* studies were conducted using a mouse surrogate comprising the extracellular domain (ECD) of murine CD80 linked to the Fc domain of mouse IgG2a wild type (murine CD80 ECD-Fc). The effects of murine CD80 ECD-Fc were compared with those of the anti-CTLA4 antibody clone 9D9 (IgG2b) in three different syngeneic tumor models: the CT26 colon carcinoma, the MC38 colon carcinoma and the B16 melanoma models.

CT26 tumor model

[00195] Seven week old female BALB/c mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for one week before the study was initiated. The murine colorectal carcinoma cell line CT26 was implanted subcutaneously over the right flank of the mice at 1.0×10^6 cells/200 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and matrigel.

[00196] Mice were monitored twice weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: tumor volume (mm³) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to seven treatment groups (n= 15 mice per experimental group). The mean tumor volume for all animals enrolled was 96 mm³. Mice were dosed 3 times, on day 4, 7 and 11. The first group was injected with mouse IgG2b (mIgG2b) i.p. dosed at 10mg/kg (control). The second group was injected with murine CD80 ECD-Fc 20 μ mol/ μ mol SA i.v. dosed at 0.3 mg/kg. The third group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 1.5 mg/kg. The fourth group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 10 mg/kg. Tumors were measured on day 10, 13, 17, 19, 21, 24.

[00197] At day 21 (when all the controls were still in the study), treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 0.3 mg/kg resulted in 90% inhibition of tumor growth compared to the control ($P < 0.001$). Treatment with anti-CTLA4 antibody at 10 mg/kg resulted in 75% inhibition of tumor growth compared to the control ($P < 0.001$). By comparison, treatment with anti-CTLA4 antibody at 1.5 mg/kg only induced 53% inhibition of tumor growth ($P < 0.001$) (**Fig 7**). At day 21, the impact of treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 0.3 mg/kg on tumor growth was significantly greater than anti-CTLA4 antibody dosed at 1.5 mg/kg ($P < 0.001$) or at 10 mg/kg ($P = 0.009$).

[00198] The incidence of tumor-free mice was analyzed at day 37. Treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 0.3 mg/kg led to complete tumor regression in 7/15 (47%) of the mice. Treatment with anti-CTLA4 antibody at 10 mg/kg led to complete tumor regression in 3/15 (20%) of the mice. None of the mice treated with anti-CTLA4 antibody at 1.5 mg/kg had complete tumor regression.

MC38 tumor model

[00199] Seven week old female C57Bl/6 mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for one week before the study was initiated. The murine colorectal carcinoma cell line MC38 was implanted subcutaneously over the right flank of the mice at 0.5×10^6 cells/100 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in RPMI 1640 medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free RPMI 1640 and matrigel.

[00200] Mice were monitored twice weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was measured using calipers and volume was calculated according to the formula: tumor volume (mm^3) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to seven treatment groups (n= 15 mice per experimental group). The mean tumor volume for all animals enrolled was 78 mm^3 .

Mice were dosed 3 times, on day 7, 10 and 14. The first group was injected with mouse IgG2b (mIgG2b) i.p. dosed at 10mg/kg (control). The second group was injected with murine CD80 ECD-Fc 20 mol/mol SA i.v. dosed at 3 mg/kg. The third group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 1.5 mg/kg. The fourth group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 10 mg/kg. Tumors were measured on day 11, 14, 17, and 19.

[00201] At day 19 (when all the controls were still in the study), treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 3 mg/kg resulted in 79% inhibition of tumor growth compared to the control ($P<0.001$). Moreover, murine CD80 ECD-Fc at 20 mol/mol SA had a greater impact on tumor growth compared to anti-CTLA4 antibody ($P<0.001$). Treatment with anti-CTLA4 antibody at 10 mg/kg reduced tumor growth by 21% compared to the control ($P=0.05$) while at 1.5 mg/kg did not significantly affect tumor size (**Fig 8**). At day 21, the impact of treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 3 mg/kg on tumor growth was significantly greater than anti-CTLA4 antibody dosed at 1.5mg/kg ($P<0.001$) or at 10mg/kg ($P=0.009$).

[00202] While a 3 mg/kg dose of CD80 ECD-Fc was used for these experiments, a 0.3 mg/kg dose of CD80 ECD-Fc also reduced tumor cell growth in the MC38 tumor model (data not shown).

B16 tumor model

[00203] Seven week old female C57Bl/6 mice were purchased from Charles River Laboratories (Hollister, CA) and were acclimated for one week before the study was initiated. The murine melanoma cell line B16-F10 was implanted subcutaneously over the right flank of the mice at 0.5×10^6 cells/100 μ l/mouse. Prior to inoculation, the cells were cultured for no more than three passages in DMEM medium supplemented with 10% heat-inactivated Fetal Bovine Serum (FBS), 2mM L-Glutamine. Cells were grown at 37°C in a humidified atmosphere with 5% CO₂. Upon reaching 80-85% confluence, cells were harvested and resuspended in a 1:1 mixture of serum-free DMEM and matrigel.

[00204] Mice were monitored twice weekly following cell implantation for tumor growth. For tumor measurements, the length and width of each tumor was

measured using calipers and volume was calculated according to the formula: tumor volume (mm^3) = (width (mm) x length (mm))²/2. On Day 7, all tumors were measured, and mice were randomly assigned to seven treatment groups (n= 15 mice per experimental group). The mean tumor volume for all animals enrolled was 70 mm^3 . Mice were dosed 3 times, on day 3, 6 and 10. The first group was injected with mouse IgG2b (mIgG2b) dosed i.p. at 10mg/kg (control). The second group was injected with murine CD80 ECD-Fc 20 mol/mol SA i.v. dosed at 3 mg/kg. The third group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 1.5 mg/kg. The fourth group was injected with anti-CTLA4 antibody clone 9D9 (IgG2b) i.p. dosed at 10 mg/kg. Tumors were measured on day 10, 13, 15, 16, 17.

[00205] At day 13 (when all the controls were still in the study) treatment with murine CD80 ECD-Fc at 20 mol/mol SA dosed at 3 mg/kg resulted in 41% inhibition of tumor growth compared to the control ($P < 0.001$). Treatment with anti-CTLA4 antibody at 10 mg/kg or 1.5 mg/kg did not significantly affect tumor growth compared to the control (**Fig 9**).

TABLE OF SEQUENCES

[00206] The table below provides a listing of certain sequences referenced herein.

SEQ. ID. NO.	Description	Sequence
1	Human CD80 precursor (with signal sequence) amino acid sequence	MGHTRRQGTSPSKCPYLNFFQLLVLAGLSHFCSGVIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKKMVLTMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYESCVVLKYEKDAFKREHLAEVTLVSKADFPTPSISDFEIPSTNIRRIICSTSGGFPEPHLSWLENGEELNAINTTVSQDPETELYAVSSKLDNFNMTTNHFSFMCLIKYGHLRVNQTFNWNNTTKQEHFPDNLPSWAITLISVNGIFVICCLTYCFAPRCRERRRRNERLRRESVRPV
2	Mouse CD80 precursor (with signal sequence) amino acid sequence	MACNCQLMQDTPLLKFPCLILLFVLLIRLSQVSSDVDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKVLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLSDRGTYSVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSADTKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPESELYTISSQLDFNTRNHTIKCLIKYGDAHVSEDFTWEKPPEDPPDSKNTLVLFAGFGAVITVVIVVIKCFCKHRSCFRRNEASRETNNSLTFGPEEALAEQTVFL
3	Human CD80 Isoform 2 (without signal sequence)	VIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKKMVLTMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYESCVVLKYEKDAFKREHLAEVTLVSKADFPTPSISDFEIPSTNIRRIICSTSGGFPEPHLSWLENGEELNAINTTVSQDPETELYAVSSKLDNFNMTTNHFSFMCLIKYGHLRVNQTFNWNNTSFAPRCRERRRRNERLRRESVRPV
4	Human CD80 Isoform 3 (without signal sequence)	VIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKKMVLTMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYESCVVLKYEKDAFKREHLAEVTLVSKGFAPRCRERRRRNERLRRESVRPV
5	Human CD80 ECD sequence (without signal sequence)	VIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKKMVLTMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYESCVVLKYEKDAFKREHLAEVTLVSKADFPTPSISDFEIPSTNIRRIICSTSGGFPEPHLSWLENGEELNAINTTVSQDPETELYAVSSKLDNFNMTTNHFSFMCLIKYGHLRVNQTFNWNNTTKQEHFPDN
6	Mouse CD80 ECD sequence (without signal sequence)	VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKVLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLSDRGTYSVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSADTKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPESELYTISSQLDFNTRNHTIKCLIKYGDAHVSEDFTWEKPPEDPPDSKN
7	Human CD80 signal sequence	MGHTRRQGTSPSKCPYLNFFQLLVLAGLSHFCSG
8	Mouse CD80	MACNCQLMQDTPLLKFPCLILLFVLLIRLSQVSSD

	signal sequence	
9	Fc C237S	EPKSSDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWWYVDGVEVH NAKTKPREEQ YNSTYRVVSV LTVLHQDWLN GKEYCKVSN KALPAIEKT ISKAKGQPRE PQVYTLPPSRD ELTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTPP VLDSGDSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHYTQKSLSLSPGK
10	Fc	ERKCCVECPC CPAPPVAGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE DPEVQFNWYV DGVEVHNAKT KPREEQFNST FRVVSVLTVV HQDWLNGKEY KCKVSNKGLP APIEKTISK KGQPREPQVY TLPPSREEMT KNQVSLTCLV KGFYPSDIAV EWESNGQPEN NYKTPPMLD SDGSFFLYSK LTVDKSRWQQ GNVFSCSVMH EALHNHYTQK SLSLSPGK
11	Fc	ESKYGPPCPS CPAPEFLGGP SVFLFPPKPK DTLMISRTPE VTCVVVDVSQ EDPEVQFNWY VDGVEVHNAK TKPREEQFNS TYRVVSVLTV LHQDWLNGKE YKCKVSNKGL PSSIEKTISK AKGQPREPQV YTLPPSQEEM TKNQVSLTCL VKGFYPSDIA VEWESNGQPE NNYKTPPVV DSDGSFFLYS RLTVDKSRWQ EGNVFSCSVM HEALHNHYTQ KSLSLSLGK
12	Human IgG1 Fc L234F, L235E, P331S mutant	EPKSSDKTHTCPPCPAPEFEGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPASIEK TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY SDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDK SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
13	Human IgG1 Fc N297 mutant	EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYGSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE KTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTV KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
14	Fc human IgG1	EPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFY SDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDK SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
15	Fc human IgG3	ELKTPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDT PPPCPRCPEPKSCDTPPPCPRCPAPELLGGPSVFLFPPK PKDTLMISRTPEVTCVVVDVSHEDPEVQFKWYVDGVEVH NAKTKPREEQYNSTFRVVSVLTVLHQDWLNGKEYKCKVS NKALPAIEKTISKTKGQPREPQVYTLPPSREEMTKNQVS LTCLVKGFYPSDIAVEWESSGQPENNYNTTPPMLDSDGS FFLYSKLTVDKSRWQQGNIFSCSVMHEALHNRFTQKSLSL SPGK
16	Fc human IgG4	ESKYGPPCPCPAPEFLGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQF NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTI SKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKS RWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK
17	Mouse CD80 ECD mouse Fc	VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKV VLSVIAGKLVWPEYKNRTLYDNTTYSLLIILGLVLSDRGTY

	IgG2a (Fc portion <u>underlined</u>)	SCVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSA DTKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPES ELYTISSQLDFNTTRNHTIKCLIKYGDAHVSEDFTWKPPPE DPPDSKNEPRGPTIKCPPCKCPAPNLLGGPSVFIFPPKIK <u>DVLMISLSPIVTCVVVDVSEDDPDVQISWVFNVEVHTAQ</u> <u>TQTHREDYNSTLRVVSALPIQHQDWMSGKEFKCKVNNKD</u> <u>LPAPIERTISKPKGSVRAPQVYVLPPEEEMTKKQVTLTC</u> <u>MVTDMPEDIYVEWTNNGKTELNYKNTEPVLDSDGSYFM</u> <u>YSKLRVEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTP</u> <u>GK</u>
18	Mouse CD80 ECD Human Fc IgG1 WT (Fc portion <u>underlined</u>)	VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKV VLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLSDRGT SCVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSA DTKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPES ELYTISSQLDFNTTRNHTIKCLIKYGDAHVSEDFTWKPPPE DPPDSKNEPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPK DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL LVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKLSLSLSP GK
19	Mouse CD80 ECD Fc IgG1 MT (234, 235, 331) (Fc portion <u>underlined</u> ; mutants shown in bold)	VDEQLSKSVKDKVLLPCRYNSPHEDESEDRIYWQKHDKV VLSVIAGKLVWPEYKNRTLYDNTTYSLIILGLVLSDRGT SCVVQKKERGTYEVKHLALVKLSIKADFSTPNITESGNPSA DTKRITCFASGGFPKPRFSWLENGRELPGINTTISQDPES ELYTISSQLDFNTTRNHTIKCLIKYGDAHVSEDFTWKPPPE DPPDSKNEPKSSDKTHTCPPCPAPEFEGGPPSVFLFPPKPK K DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN A KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN K ALP A S I EKTISKAKGQPREPQVYTLPPSRDELTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFF LYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKLSLSLSP PGK
20	Human CD80 ECD Human Fc IgG1 WT (Fc portion <u>underlined</u>)	VIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKMMVL TMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYE CVVLKYEKDAFKREHLAEVTLVSKADFPTPSISDFEIPSTNI RRIICSTSGGFPEPHLSWLENGEELNAINTTVSQDPETELY AVSSKLDNFNMTTNSFSMCLIKYGHLRVNQTFNWNTTKQE HFPDNEPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDT LMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYS KLTVDKSRWQQGNVFSCSVMHEALHNHYTQKLSLSLSPG K
21	Human CD80 ECD Human Fc IgG1 L234F, L235E, P331S MT (Fc portion <u>underlined</u> ; mutants in bold)	VIHVTKEVKEVATLSCGHNVSVVEELAQTRIWQKEKMMVL TMMSGDMNIWPEYKNRTIFDITNNLSIVILALRPSDEGTYE CVVLKYEKDAFKREHLAEVTLVSKADFPTPSISDFEIPSTNI RRIICSTSGGFPEPHLSWLENGEELNAINTTVSQDPETELY AVSSKLDNFNMTTNSFSMCLIKYGHLRVNQTFNWNTTKQE HFPDNEPKSSDKTHTCPPCPAPEFEGGPPSVFLFPPKPKD TL M ISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA L P A S I EKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKLSLSLSP GK
22	human PD-1	MQIPQAPWPV VVAVLQLGWR PGWFLDSPDR

	precursor (with signal sequence) UniProtKB/Swiss-Prot: Q15116.3, 01-OCT-2014	PWNPPTFSPA LLVVTEGDNA TFTCSFSNTS ESFVLNWYRM SPSNQTDKLA AFPEDRSQPG QDCRFRVTQL PNGRDFHMSV VRARRNDSGT YLCGAI SLAP KAQIKESLRA ELRV TERRAE VPTAHPSPSP RPAGQFQTLV VGVVGGLLGS LVLLWV LAV ICSRAARGTI GARRTGQPLK EDPSAVPVFS VDYGELDFQW REKTPEPPVP CVPEQTEYAT IVFPSGMGTS SPARRGSADG PRSAQPLRPE DGHCSWPL
23	human PD-1 (mature, without signal sequence)	PGWFLDSPDR PWNPPTFSPA LLVVTEGDNA TFTCSFSNTS ESFVLNWYRM SPSNQTDKLA AFPEDRSQPG QDCRFRVTQL PNGRDFHMSV VRARRNDSGT YLCGAI SLAP KAQIKESLRA ELRV TERRAE VPTAHPSPSP RPAGQFQTLV VGVVGGLLGS LVLLWV LAV ICSRAARGTI GARRTGQPLK EDPSAVPVFS VDYGELDFQW REKTPEPPVP CVPEQTEYAT IVFPSGMGTS SPARRGSADG PRSAQPLRPE DGHCSWPL
24	human PD-L1 precursor (with signal sequence) UniProtKB/Swiss-Prot: Q9NZQ7.1, 01-OCT-2014	MRIFAVFIFM TYWHLLNAFT VTPKDLYVV EYGSNMTIEC KFPVEKQLDL AALIVYWEME DKNIIQFVHG EEDLKVQHSS YRQRARLLKD QLSLGNAALQ ITDVKLQDAG VYRCMISYGG ADYKRITVKV NAPYNKINQR ILVVDPTSE HELTCQAEGY PKAEVIWTSS DHQVLSGKTT TTNSKREEKL FNVSTSLRIN TTTNEIFYCT FRRLDPEENH TAE LVIPELP LAHPPNER TH LVILGAILLC LGVALTFIFR LRKGRMM DVK KCGIQDTNSK KQSDTHLEET
25	human PD-L1 (mature, without signal sequence)	FT VTPKDLYVV EYGSNMTIEC KFPVEKQLDL AALIVYWEME DKNIIQFVHG EEDLKVQHSS YRQRARLLKD QLSL GNAALQ ITDVKLQDAG VYRCMISYGG ADYKRITVKV NAPYNKINQR ILVVDPTSE HELTCQAEGY PKAEVIWTSS DHQVLSGKTT TTNSKREEKL FNVSTSLRIN TTTNEIFYCT FRRLDPEENH TAE LVIPELP LAHPPNER TH LVILGAILLC LGVALTFIFR LRKGRMM DVK KCGIQDTNSK KQSDTHLEET

What is Claimed is:

1. A CD80 extracellular domain (ECD) fusion molecule comprising a human CD80 ECD polypeptide and a human IgG1 Fc domain wherein the CD80 ECD fusion molecule comprises 10-60 mol SA to mol of CD80 ECD Fc protein, wherein the molecule is capable of at least 80%, such as at least 90%, such as at least 95%, such as at least 98% tumor cell growth inhibition in at least one mouse syngeneic or xenograft cancer model over at least two weeks.
2. The CD80 extracellular domain (ECD) fusion molecule of claim 1, wherein the molecule comprises at least 15 mol SA to mol of protein, such as at least 20 mol SA to mol of protein.
3. The CD80 ECD Fc fusion molecule of claim 1 or 2, wherein the molecule comprises 15-30 mol SA to mol of protein.
4. The CD80 ECD Fc fusion molecule of claim 1 or 2, wherein the molecule comprises 15-25 mol SA to mol of protein.
5. The CD80 ECD Fc fusion molecule of claim 1 or 2, wherein the molecule comprises 20-30 mol SA to mol of protein.
6. The CD80 ECD Fc fusion molecule of any one of claims 1-5, wherein the Fc domain comprises the amino acid sequence of SEQ ID NO:14.
7. The CD80 ECD Fc fusion molecule of any one of claims 1-6, wherein the molecule is capable of at least 80%, such as at least 90%, at least 95%, or at least 98% reduction of growth of CT26 tumor cells in mice over a period of two or three weeks.
8. The CD80 ECD Fc fusion molecule of claim 7, wherein the mice are given one, two, or three doses of 0.3 to 0.6 mg/kg of the ECD Fc fusion molecule.
9. A composition comprising the CD80 ECD fusion molecule of any one of claims 1-8, and further comprising at least one pharmaceutically acceptable carrier.
10. The composition of claim 9, further comprising an additional therapeutic agent.
11. The composition of claim 10, wherein the additional therapeutic agent is at least one immune stimulating agent.
12. The composition of claim 11, wherein the at least one immune stimulating agent comprises a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor.

13. The composition of claim 12, wherein the PD-1/PD-L1 inhibitor is an antibody.
14. The composition of claim 13, wherein the PD-1/PD-L1 inhibitor is an anti-PD-1 antibody.
15. The composition of claim 14, wherein the anti-PD-1 antibody comprises the heavy chain and light chain CDRs of an antibody selected from nivolumab, pidilizumab, and pembrolizumab.
16. The composition of claim 14, wherein the anti-PD-1 antibody comprises the heavy chain and light chain variable regions of an antibody selected from nivolumab, pidilizumab, and pembrolizumab.
17. The composition of claim 14, wherein the anti-PD-1 antibody is selected from nivolumab, pidilizumab, and pembrolizumab.
18. The composition of claim 12, wherein the PD-1/PD-L1 inhibitor is an anti-PD-L1 antibody.
19. The composition of claim 18, wherein the anti-PD-L1 antibody comprises the heavy chain and light chain CDRs of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
20. The composition of claim 18, wherein the anti-PD-L1 antibody comprises the heavy chain and light chain variable regions of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
21. The composition of claim 18, wherein the anti-PD-1 antibody is selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
22. The composition of claim 12, wherein the PD-1/PD-L1 inhibitor is a PD-1 fusion molecule.
23. The composition of claim 22, wherein the fusion molecule is AMP-224.
24. The composition of claim 12, wherein the PD-1/PD-L1 inhibitor is a polypeptide such as AUR-012.
25. A method of treating cancer in a subject comprising administering to the subject an effective amount of the CD80 extracellular domain (ECD) fusion molecule according to any one of claims 1-8 or the composition of claim 9.
26. The method of claim 25, wherein the cancer is a solid tumor.
27. The method of claim 26, wherein the cancer is selected from colorectal cancer, breast cancer, gastric cancer, non-small cell lung cancer, melanoma, squamous cell

carcinoma of the head and neck, ovarian cancer, pancreatic cancer, renal cell carcinoma, hepatocellular carcinoma, bladder cancer, and endometrial cancer.

28. The method of any one of claims 25-27, wherein the cancer is recurrent or progressive after a therapy selected from surgery, chemotherapy, radiation therapy, or a combination thereof.

29. The method of any one of claims 25-28, wherein the CD80 ECD fusion molecule is a CD80 ECD Fc fusion molecule wherein: (a) the CD80 ECD Fc fusion molecule reduces growth of CT26 tumor cells in mice over a period of at least two weeks by a greater degree than a CD80 ECD Fc protein of the same amino acid sequence but comprising less than 10 mol SA to mol of CD80 ECD Fc protein, and/or (b) the CD80 ECD Fc fusion molecule reduces growth of CT26 tumor cells in mice over a period of at least two weeks by a greater degree than an anti-CTLA4 antibody.

30. The method of claim 29, wherein the fusion molecule comprises at least 15 mol SA to mol of protein, such as at least 20 mol SA to mol of protein.

31. The method of claim 29, wherein the fusion molecule comprises 15-40, 15-30, 15-25, or 20-30 mol SA to mol of protein.

32. A method of enhancing efficacy of a CD80 ECD fusion protein in treating cancer in a subject comprising increasing the level of sialic acid (SA) in the CD80 ECD fusion protein by administering the CD80 ECD fusion protein comprising an increased level of SA to the subject.

33. The method of claim 32, wherein the SA level is increased by 5, 10, 20, 30, 40, or 50 mol to mol of CD80 ECD protein.

34. The method of claim 32 or 33, wherein enhanced efficacy is measured as an increase in overall survival, an increase in disease-free survival, or as a greater reduction in growth of at least one tumor in a mouse syngeneic or xenograft model in comparison to administration of the CD80 ECD fusion molecule without the increased SA level.

35. The method of any one of claims 32-34, wherein the enhanced efficacy is measured as a greater reduction in tumor growth in a mouse syngeneic or xenograft model such as a CT26 mouse model or as a reduced rate of clearance in an animal or human subject in comparison to administration of the CD80 ECD fusion molecule without the increased SA level.

36. The method of any one of claims 32-35, wherein the enhanced efficacy is measured as (a) a further reduction in tumor growth of at least one tumor in an animal or human subject in comparison to administration of the CD80 ECD fusion molecule without the increased SA level, and/or (b) a further reduction in tumor growth in a mouse syngeneic or xenograft model in comparison to administration of the CD80 ECD fusion molecule without the increased SA level, and wherein the tumor growth is further reduced by at least 10%, such as at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% upon administration of the CD80 ECD fusion molecule with increased SA level in comparison to administration of the CD80 ECD fusion molecule without the increased SA level.
37. A method of treating cancer in a subject comprising administering to the subject an effective amount of a combination of a CD80 extracellular domain (ECD) or CD80 ECD fusion molecule and a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor.
38. The method of claim 37, wherein the combination of the CD80 ECD or CD80 ECD fusion molecule and the PD-1/PD-L1 inhibitor reduces tumor growth in a mouse syngeneic or xenograft model in a synergistic or in an additive fashion compared to treatment with either the CD80 ECD or CD80 ECD fusion molecule or the PD-1/PD-L1 inhibitor alone.
39. The method of claim 38, wherein the mouse model is a colorectal cancer model comprising murine colorectal carcinoma CT26 cells.
40. The method of claim 37 or 38, wherein the cancer is a solid tumor.
41. The method of any one of claims 37-40, wherein the cancer is selected from colorectal cancer, breast cancer, gastric cancer, non-small cell lung cancer, melanoma, squamous cell carcinoma of the head and neck, ovarian cancer, pancreatic cancer, renal cell carcinoma, hepatocellular carcinoma, bladder cancer, and endometrial cancer.
42. The method of any one of claims 37-41, wherein the cancer is recurrent or progressive after a therapy selected from surgery, chemotherapy, radiation therapy, or a combination thereof.
43. The method of any one of claims 37-42, wherein the combination comprises a CD80 ECD fusion molecule with a fusion partner selected from an IgG Fc domain, albumin, and PEG.

44. The method of claim 43, wherein the fusion partner is an Fc domain, such as an Fc domain comprising one of SEQ ID NOs: 9-16.
45. The method of any one of claims 37-44, wherein the CD80 ECD or CD80 ECD fusion molecule comprises an amino acid sequence selected from: amino acids 35 to end of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5, or comprises the sequence of SEQ ID NO:20 or SEQ ID NO: 21.
46. The method of any one of claims 37-45, wherein the PD-1/PD-L1 inhibitor is an anti-PD-1 antibody.
47. The method of claim 46, wherein the anti-PD-1 antibody comprises the heavy chain and light chain CDRs of an antibody selected from nivolumab, pidilizumab, and pembrolizumab.
48. The method of claim 47, wherein the anti-PD-1 antibody comprises the heavy chain and light chain variable regions of an antibody selected from nivolumab, pidilizumab, and pembrolizumab.
49. The method of claim 48, wherein the anti-PD-1 antibody is selected from nivolumab, pidilizumab, and pembrolizumab.
50. The method of any one of claims 37-49, wherein the PD-1/PD-L1 inhibitor is an anti-PD-L1 antibody.
51. The method of claim 50, wherein the anti-PD-L1 antibody comprises the heavy chain and light chain CDRs of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
52. The method of claim 51, wherein the anti-PD-L1 antibody comprises the heavy chain and light chain variable regions of an antibody selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
53. The method of claim 52, wherein the anti-PD-1 antibody is selected from BMS-936559, MPDL3280A, MEDI4736, and MSB0010718C.
54. The method of any one of claims 37-45, wherein the PD-1/PD-L1 inhibitor is a PD-1 fusion molecule.
55. The method of claim 54, wherein the fusion molecule is AMP-224.
56. The method of any one of claims 37-45, wherein the PD-1/PD-L1 inhibitor is a polypeptide such as AUR-012.

57. The method of any one of claims 37-56, wherein the CD80 ECD or CD80 ECD fusion molecule and the PD-1/PD-L1 inhibitor are administered concurrently or sequentially.
58. The method of claim 57, wherein one or more doses of the PD-1/PD-L1 inhibitor are administered prior to administering the CD80 ECD or CD80 ECD fusion molecule.
59. The method of claim 58, wherein the subject received a complete course of PD-1/PD-L1 inhibitor therapy prior to administration of the CD80 ECD or CD80 ECD fusion molecule.
60. The method of claim 58, wherein the CD80 ECD or CD80 ECD fusion molecule is administered during a second course of PD-1/PD-L1 inhibitor therapy.
61. The method of any one of claims 58-60, wherein the subject received at least one, at least two, at least three, or at least four doses of the PD-1/PD-L1 inhibitor prior to administration of the CD80 ECD or CD80 ECD fusion molecule.
62. The method of any one of claims 58-61, wherein at least one dose of the PD-1/PD-L1 inhibitor is administered concurrently with the CD80 ECD or CD80 ECD fusion molecule.
63. The method of any one of claims 37-62, wherein one or more doses of the CD80 ECD or CD80 ECD fusion molecule are administered prior to administering a PD-1/PD-L1 inhibitor.
64. The method of claim 63, wherein the subject received at least two, at least three, at least three, or at least four doses of the CD80 ECD or CD80 ECD fusion molecule prior to administration of the PD-1/PD-L1 inhibitor.
65. The method of claim 63 or 64, wherein at least one dose of the CD80 ECD or CD80 ECD fusion molecule is administered concurrently with the PD-1/PD-L1 inhibitor.
66. The method of any one of claims 37-65, wherein the subject is resistant to treatment with a PD-1/PD-L1 inhibitor.
67. The method of claim 66, wherein the subject has previously received PD-1/PD-L1 inhibitor therapy.

68. The method of any one of claims 37-67, wherein the subject is administered at least one additional therapeutic agent, such as at least one chemotherapy agent, growth inhibitory agent, anti-angiogenesis agent and/or anti-neoplastic composition.
69. The method of any one of claims 37-68, wherein the CD80 ECD or CD80 ECD fusion molecule is a CD80 ECD fusion molecule comprising 10-60 mol sialic acid (SA) to mol of CD80 ECD protein.
70. The method of claim 69, wherein the CD80 ECD fusion molecule comprises at least 15 mol SA to mol of protein, such as at least 20 mol SA to mol of protein.
71. The method of claim 69, wherein the CD80 ECD fusion molecule comprises 15-30 mol SA to mol of protein.
72. The method of claim 69, wherein the CD80 ECD fusion molecule comprises 15-25 mol SA to mol of protein.
73. The method of claim 72, wherein the CD80 ECD fusion molecule comprises 20-30 mol SA to mol of protein.
74. The method of any one of claims 69-73, wherein the CD80 ECD fusion molecule comprises an Fc domain as fusion partner.
75. The method of claim 74, wherein the fusion partner is a wild-type IgG1 Fc domain.
76. The method of any one of claims 37-75, wherein the CD80 ECD or CD80 ECD fusion molecule is capable of at least 50%, 60%, 70%, 80%, or 90% reduction of growth of tumor cells in at least one mouse syngeneic or xenograft model over a period of two or three weeks.
77. The method of claim 76, wherein the CD80 ECD or CD80 ECD fusion molecule is capable of at least 95% reduction, such as at least 98%, of growth of CT26 tumor cells in mice over a period of two or three weeks.
78. The method of claim 77, wherein mice given one, two, or three doses of 0.3 to 0.6 mg/kg of the CD80 ECD or CD80 ECD fusion molecule.
79. The method of any one of claims 37-78, wherein the CD80 ECD or CD80 ECD fusion molecule is a CD80 ECD Fc fusion molecule according to any one of claims 1-8.
80. A CD80 extracellular domain (ECD) fusion molecule comprising a human CD80 ECD polypeptide and a human IgG1 Fc domain with L234F, L235E, and P331S amino acid substitutions.

81. The CD80 ECD fusion molecule of claim 80, comprising an amino acid sequence selected from SEQ ID NO: 5, SEQ ID NO: 12, and SEQ ID NO: 21.
82. The CD80 ECD fusion molecule of claim 80 or 81, wherein the Fc domain amino acid sequence is directly attached to the C-terminal amino acid of the CD80 ECD amino acid sequence.
83. The CD80 ECD fusion molecule of claim 80 or 81, wherein the Fc domain amino acid sequence is attached to the C-terminal amino acid of the CD80 ECD amino acid sequence through a linker peptide, such as a GS linker.
84. A composition comprising the CD80 ECD fusion molecule of any one of claims 80-83, and further comprising at least one pharmaceutically acceptable carrier.
85. The composition of claim 84, further comprising an additional therapeutic agent, such as at least one immune stimulating agent.
86. The composition of claim 85, wherein the at least one immune stimulating agent comprises a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor, such as a PD-1 fusion molecule (i.e. AMP 224), such as a polypeptide (i.e. AUR-012), or such as an antibody, such as an anti-PD-1 antibody (i.e. nivolumab, pidilizumab, or pembrolizumab) or an anti-PD-L1 antibody (i.e. BMS-936559, MPDL3280A, MEDI4736, or MSB0010718C).
87. A method of treating cancer in a subject comprising administering to the subject an effective amount of the CD80 extracellular domain (ECD) fusion molecule according to any one of claims 80-83 or the composition of any one of claims 84-86.
88. The method of claim 87, wherein the cancer is a solid tumor.
89. The method of claim 88, wherein the cancer is selected from colorectal cancer, breast cancer, gastric cancer, non-small cell lung cancer, melanoma, squamous cell carcinoma of the head and neck, ovarian cancer, pancreatic cancer, renal cell carcinoma, hepatocellular carcinoma, bladder cancer, and endometrial cancer.
90. The method of any one of claims 87-89, wherein the cancer is recurrent or progressive after a therapy selected from surgery, chemotherapy, radiation therapy, or a combination thereof.
91. The method of any one of claims 87-90, wherein the CD80 ECD fusion molecule or composition is administered in combination with at least one additional therapeutic agent, such as an agent according to claim 86.

92. The method of claim 91, wherein the additional therapeutic agent comprises at least one chemotherapy agent, growth inhibitory agent, anti-angiogenesis agent and/or anti-neoplastic composition.




Fig. 1A

Fig. 1B




Fig. 2B




Fig. 2A




Fig. 3B




Fig. 3A




Fig. 4A




Fig. 4B




Fig. 5B




Fig. 5A




Fig. 5C




Fig. 5D




Fig. 6




Fig. 7




Fig. 8




Fig. 9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2016/059838

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-11, 25-36(completely); 12-24, 69-73, 79(partially)

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/059838

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K14/705 A61K38/17
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
C07K A61K
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. T. HAILE ET AL: "Soluble CD80 Restores T Cell Activation and Overcomes Tumor Cell Programmed Death Ligand 1-Mediated Immune Suppression", THE JOURNAL OF IMMUNOLOGY, vol. 191, no. 5, 5 August 2013 (2013-08-05), pages 2829-2836, XP055334763, US ISSN: 0022-1767, DOI: 10.4049/jimmunol.1202777 abstract page 5 - page 6 page 9 ----- -/--	1-31, 69-73,79

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search 13 January 2017	Date of mailing of the international search report 29/03/2017
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Brero, Alessandro
--	---

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/059838

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>S. T. HAILE ET AL: "A Soluble Form of CD80 Enhances Antitumor Immunity by Neutralizing Programmed Death Ligand-1 and Simultaneously Providing Costimulation", CANCER IMMUNOLOGY RESEARCH, vol. 2, no. 7, 2 April 2014 (2014-04-02), pages 610-615, XP055334764, US ISSN: 2326-6066, DOI: 10.1158/2326-6066.CIR-13-0204 abstract page 613 - page 614</p> <p style="text-align: center;">-----</p>	1-9, 25-31
X	<p>OSTRAND-ROSENBERG SUZANNE ET AL: "Novel strategies for inhibiting PD-1 pathway-mediated immune suppression while simultaneously delivering activating signals to tumor-reactive T cells", CANCER IMMUNOLOGY, IMMUNOTHERAPY, SPRINGER, BERLIN/HEIDELBERG, vol. 64, no. 10, 20 March 2015 (2015-03-20), pages 1287-1293, XP035533178, ISSN: 0340-7004, DOI: 10.1007/S00262-015-1677-5 [retrieved on 2015-03-20] abstract figure 2 page 1290</p> <p style="text-align: center;">-----</p>	1-9, 25-31
A	<p>DANIEL M. CZAJKOWSKY ET AL: "Fc-fusion proteins: new developments and future perspectives", EMBO MOLECULAR MEDICINE, vol. 4, no. 10, 26 July 2012 (2012-07-26), pages 1015-1028, XP055160259, ISSN: 1757-4676, DOI: 10.1002/emmm.201201379 page 1016 page 1023</p> <p style="text-align: center;">-----</p>	1-36, 69-73,79
A	<p>RICARDO J. SOLÁ ET AL: "Glycosylation of Therapeutic Proteins", BIODRUGS, vol. 24, no. 1, 1 February 2010 (2010-02-01), pages 9-21, XP055016446, ISSN: 1173-8804, DOI: 10.2165/11530550-000000000-00000 page 6</p> <p style="text-align: center;">-----</p>	1-36, 69-73,79

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-11, 25-36(completely); 12-24, 69-73, 79(partially)

A CD80 extracellular domain (ECD) fusion molecule comprising a human CD80 ECD polypeptide and a human IgG1 Fc domain wherein the CD80 ECD fusion molecule comprises 10-60 mol sialic acid (SA) to mol of CD80 ECD Fc protein, wherein the molecule is capable of at least 80% tumor cell growth inhibition in at least one mouse syngeneic or xenograft cancer model over at least two weeks; and subject-matter relating thereto.

2. claims: 37-68, 74-78(completely); 12-24, 69-73, 79, 86-92(partially)

A method of treating cancer in a subject comprising administering to the subject an effective amount of a combination of a CD80 extracellular domain (ECD) or CD80 ECD fusion molecule and a programmed cell death 1 (PD-1) / programmed cell death ligand 1 (PD-L1) inhibitor; and subject-matter relating thereto.

3. claims: 80-85(completely); 86-92(partially)

A CD80 extracellular domain (ECD) fusion molecule comprising a human CD80 ECD polypeptide and a human IgG1 Fc domain with L234F, L235E, and P331S amino acid substitutions; and subject-matter relating thereto.
