



(51) International Patent Classification:

A61P 9/10 (2006.01) C07K 16/22 (2006.01)  
A61K 39/00 (2006.01) C07K 16/40 (2006.01)

(21) International Application Number:

PCT/US2017/020221

(22) International Filing Date:

1 March 2017 (01.03.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/302,907 3 March 2016 (03.03.2016) US

(71) Applicant: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(72) Inventors: **GUSAROVA, Viktoria**; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **GROMADA, Jesper**; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **MURPHY, Andrew J.**; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(74) Agents: **HEUSCH, Marina** et al.; Swanson & Bratschun, L.L.C., 8210 Southpark Terrace, Littleton, Colorado 80120 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2017/151783 A1

(54) Title: METHODS FOR TREATING PATIENTS WITH HYPERLIPIDEMIA BY ADMINISTERING A PCSK9 INHIBITOR IN COMBINATION WITH AN ANGPTL3 INHIBITOR

(57) Abstract: The present invention provides methods for treating patients suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy. The methods of the invention provide for lowering at least one lipid parameter in the patient by administering a therapeutically effective amount of an antibody or antigen-binding fragment thereof that specifically binds to proprotein convertase subtilisin/kexin type 9 (PCSK9) in combination with a therapeutically effective amount of an antibody that specifically binds to angiopoietin-like protein 3 (ANGPTL3). The combination of an anti-PCSK9 antibody with an anti-ANGPTL3 antibody is useful in treating diseases such as hypercholesterolemia, including familial hypercholesterolemia (FH), both heFH and hoFH, as well as hyperlipidemia, hyperlipoproteinemia and dyslipidemia, including hypertriglyceridemia, chylomicronemia, and to prevent or treat diseases or disorders, for which abnormal lipid metabolism is a risk factor, such as cardiovascular diseases.

## METHODS FOR TREATING PATIENTS WITH HYPERLIPIDEMIA BY ADMINISTERING A PCSK9 INHIBITOR IN COMBINATION WITH AN ANGPTL3 INHIBITOR

### FIELD OF THE INVENTION

[0001] The present invention relates to the field of therapeutic treatments of diseases and disorders, which are associated with elevated levels of lipids and lipoproteins. More specifically, the invention relates to the use of a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor in combination with an inhibitor of angiotensin-like protein 3 (ANGPTL3) to treat patients with hypercholesterolemia and related conditions, who are non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy.

### BACKGROUND

[0002] Hyperlipidemia is a general term that encompasses diseases and disorders characterized by or associated with elevated levels of lipids and/or lipoproteins in the blood. Hyperlipidemias include hypercholesterolemia, hypertriglyceridemia, combined hyperlipidemia, and elevated lipoprotein a (Lp(a)). A particular prevalent form of hyperlipidemia in many populations is hypercholesterolemia.

[0003] Hypercholesterolemia, particularly an increase in low-density lipoprotein (LDL) cholesterol (LDL-C) levels, constitutes a major risk for the development of atherosclerosis and coronary heart disease (CHD) (Sharrett *et al.*, 2001, *Circulation* 104:1108-1113). Low-density lipoprotein cholesterol is identified as the primary target of cholesterol lowering therapy and is accepted as a valid surrogate therapeutic endpoint. Numerous studies have demonstrated that reducing LDL-C levels reduces the risk of CHD with a strong direct relationship between LDL-C levels and CHD events; for each 1 mmol/L (~40 mg/dL) reduction in LDL-C, cardiovascular disease (CVD) mortality and morbidity is lowered by 22%. Greater reductions in LDL-C produce greater reduction in events, and comparative data of intensive versus standard statin treatment suggest that the lower the LDL-C level, the greater the benefit in patients at very high cardiovascular (CV) risk.

[0004] Familial hypercholesterolemia (FH) is an inherited disorder of lipid metabolism that predisposes a person to premature severe cardiovascular disease (CVD) (Kolansky *et al.*, (2008), *Am J Cardiology*, 102(11):1438-1443). FH can be either an autosomal dominant or an autosomal recessive disease that results from mutations in the low density lipoprotein receptor (LDLR), or in at least 3 different genes that code for proteins involved in hepatic clearance of LDL-C can cause FH. Examples of such defects include mutations in the gene coding for the LDL receptor (LDLR) that removes LDL-C from the circulation, and in the gene for apolipoprotein (Apo) B, which is the major protein of the LDL particle. In certain cases of FH, the gene coding for proprotein convertase subtilisin/kexin type 9 (PCSK9), an enzyme involved in degrading the LDLR (gain of function mutation), is mutated. In all cases, FH is characterized by an accumulation of LDL-C in the plasma from birth and subsequent development of tendon

xanthomas, xanthelasma, atheromata, and CVD. FH can be classified as either heterozygous FH (heFH) or homozygous FH (hoFH) depending on whether the individual has a genetic defect in one (heterozygous) or both (homozygous) copies of the implicated gene.

**[0005]** Current LDL-C-lowering medications include statins, cholesterol absorption inhibitors, fibrates, niacin, and bile acid sequestrants. Statins are a commonly prescribed treatment for LDL-C lowering. However, despite the availability of such lipid-lowering therapies, many high-risk patients fail to reach their guideline target LDL-C level (Gitt et al., 2010, Clin Res Cardiol 99(11):723-733). For patients who are still unable to achieve guideline target level for LDL-C, despite available lipid-modifying therapy (LMT), mechanical removal of LDL-C by lipoprotein apheresis (*e.g.*, LDL apheresis) is sometimes prescribed.

**[0006]** However, patients who are not at LDL-C goal despite receiving an optimized LMT regimen, would greatly benefit from alternative LDL-C lowering therapies, or through use of a combination of therapeutic agents, such as the agents and regimens described herein.

#### **BRIEF SUMMARY OF THE INVENTION**

**[0007]** The present invention provides methods, uses, and compositions for treating hyperlipidemia in patients who are non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy. The therapeutic methods of the present invention result in a lowering of serum lipoprotein levels to a normal and acceptable range and as such, may act to reduce the risk of development of atherosclerosis, or coronary heart disease.

**[0008]** In one aspect, the invention provides a method of treating a patient suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy, the method comprising treating the patient with a combination of a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor and an inhibitor of angiotensin-like protein 3 (ANGPTL3).

**[0009]** In one embodiment, the invention provides administering one or more doses of a PCSK9 inhibitor in combination with one or more doses of an ANGPTL3 inhibitor to a patient who is being treated, or has been treated with a standard lipid modifying therapy, but has not responded to such therapy. Administration of a combination of a PCSK9 inhibitor with an ANGPTL3 inhibitor to the patient results in lowering the level of at least one lipoprotein in the serum of the patient and consequently reduces or eliminates the need for treatment with the standard lipid lowering therapy by the patient.

**[0010]** In a related aspect, the methods of the present invention comprise selecting a patient with hypercholesterolemia who is being treated, or has been treated with a standard lipid lowering therapy and who is non-responsive to, inadequately controlled by, or intolerant to, such therapy and administering one or more doses of a PCSK9 antibody in combination with

one or more doses of an ANGPTL3 antibody to the patient, thereby lowering the level of at least one lipoprotein in the serum of the patient and consequently replacing the use of the standard lipid modifying therapy with the combination therapy of a PCSK9 antibody plus an ANGPTL3 antibody to achieve a target lipoprotein level.

**[0011]** Patients who are treated or treatable by the methods of the present invention include, *e.g.*, patients with hypercholesterolemia, including patients with familial hypercholesterolemia (FH). In certain embodiments, the patients who are treated or treatable by the methods of the present invention are patients who are diagnosed with (or otherwise known to have), homozygous FH (hoFH) or heterozygous FH (heFH), or at risk for developing abnormally high lipid and/or lipoprotein levels associated with homozygous FH (hoFH) or heterozygous FH (heFH).

**[0012]** The present invention also provides pharmaceutical compositions comprising a PCSK9 inhibitor and an ANGPTL3 inhibitor for use in treating a patient who is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy, such as a statin. The statin may be selected from the group consisting of atorvastatin (LIPITOR®), pitavastatin (LIVALO®), lovastatin (MEVACOR®), simvastatin (ZOCOR®), pravastatin (PRAVACHOL®) fluvastatin (LESCOL®) and rosuvastatin (CRESTOR®). Other standard lipid lowering agents that may be used in patients suffering from hypercholesterolemia, include, but are not limited to, fibrates, niacin, bile acid sequestrants, ezetimibe (ZETIA®), lomitapide (JUZTAPID™), phytosterols, orlistat (XENICAL®).

**[0013]** Exemplary PCSK9 inhibitors, or ANGPTL3 inhibitors that may be used in the context of the methods of the present invention include, *e.g.*, anti-PCSK9 or anti-ANGPTL3 antibodies, small molecule inhibitors, and scaffold-based, *i.e.* PCSK9-binding molecules, or ANGPTL3-binding molecules.

**[0014]** In certain embodiments, it is envisioned that the use of the combination of the PCSK9 inhibitor with the ANGPTL3 inhibitor may be sufficiently effective at lowering serum lipid and/or lipoprotein levels, such that the dose of the standard lipid modifying therapy may be reduced to eliminate any untoward effects, or it may be eliminated altogether.

**[0015]** In one embodiment, the methods provide for treating a patient in need thereof with an antibody, or an antigen-binding fragment thereof, that binds specifically to PCSK9 in combination with an antibody, or an antigen-binding fragment thereof, that binds specifically to ANGPTL3. In one embodiment, the PCSK9 antibody is administered to the patient at a dose of about 75 mg at a frequency of once every two weeks. In one embodiment, the PCSK9 antibody is administered to the patient at a dose of about 140 mg at a frequency of once every two weeks. In one embodiment, the PCSK9 antibody is administered to the patient at a dose of about 150 mg at a frequency of once every two or four weeks. In one embodiment, the PCSK9 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every

four weeks. In one embodiment, the PCSK9 antibody is administered to the patient at a dose of about 420 mg at a frequency of once every four weeks.

**[0016]** In one embodiment, the PCSK9 antibody is selected from the group consisting of alirocumab, evolocumab, bococizumab, lodelcizumab and ralpancizumab.

**[0017]** In one embodiment, the PCSK9 antibody is alirocumab.

**[0018]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to PCSK9 comprises the complementary determining regions (CDRs) of a heavy chain variable (HCVR) having the amino acid sequence of SEQ ID NO: 12 and the CDRs of a light chain variable region (LCVR) of SEQ ID NO: 17.

**[0019]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to PCSK9 comprises a heavy chain CDR1 (HCDR1) having the amino acid sequence of SEQ ID NO: 13, a HCDR2 having the amino acid sequence of SEQ ID NO: 14, a HCDR3 having the amino acid sequence of SEQ ID NO: 15, a light chain CDR1 (LCDR1) having the amino acid sequence of SEQ ID NO: 18, a LCDR2 having the amino acid sequence of SEQ ID NO: 19, and a LCDR3 having the amino acid sequence of SEQ ID NO: 21.

**[0020]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to PCSK9 comprises a HCVR having the amino acid sequence of SEQ ID NO: 12 and a LCVR having the amino acid sequence of SEQ ID NO: 17.

**[0021]** In one embodiment, the PCSK9 antibody is administered to the patient subcutaneously or intravenously.

**[0022]** In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 150 mg at a frequency of once every week. In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every week. In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 450 mg at a frequency of once every week. In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every two weeks. In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 450 mg at a frequency of once every two weeks. In one embodiment, the ANGPTL3 antibody is administered to the patient at a dose of about 20 mg/kg at a frequency of once every four weeks.

**[0023]** In one embodiment, the ANGPTL3 antibody is evinacumab.

**[0024]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to ANGPTL3 comprises the complementary determining regions (CDRs) of a heavy chain variable (HCVR) having the amino acid sequence of SEQ ID NO: 2 and the CDRs of a light chain variable region (LCVR) of SEQ ID NO: 3.

**[0025]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to ANGPTL3 comprises a heavy chain CDR1 (HCDR1) having the amino acid sequence of SEQ ID NO: 4, a HCDR2 having the amino acid sequence of SEQ ID NO: 5, a

HCDR3 having the amino acid sequence of SEQ ID NO: 6, a light chain CDR1 (LCDR1) having the amino acid sequence of SEQ ID NO: 7, a LCDR2 having the amino acid sequence of SEQ ID NO: 8, and a LCDR3 having the amino acid sequence of SEQ ID NO: 9.

**[0026]** In one embodiment, the antibody, or antigen-binding fragment thereof that binds specifically to ANGPTL3 comprises a HCVR having the amino acid sequence of SEQ ID NO: 2 and a LCVR having the amino acid sequence of SEQ ID NO: 3.

**[0027]** In one embodiment, the ANGPTL3 antibody is administered to the patient subcutaneously or intravenously.

**[0028]** In one embodiment, the administration of the PCSK9 antibody in combination with the ANGPTL3 antibody results in an additive effect on lowering the blood level of LDL-C, non-HDL-C and total cholesterol, but has no effect on blood levels of HDL-C.

**[0029]** In one embodiment, the administration of the PCSK9 antibody in combination with the ANGPTL3 antibody results in a synergistic effect on lowering the blood level of LDL-C, non-HDL-C and total cholesterol, but has no effect on blood levels of HDL-C.

**[0030]** In one embodiment, the administration of the PCSK9 antibody in combination with the ANGPTL3 antibody results in lowering one or more of the following parameters:

- (a) a reduction in serum total cholesterol (TC) level;
- (b) a reduction in serum low-density lipoprotein cholesterol (LDL-C) levels; or
- (c) a reduction in serum non-high density lipoprotein cholesterol (non-HDL-C) levels;

wherein the reduction of (a), (b), and/or (c) are determined relative to the patient's serum TC level, serum LDL-C levels and/or serum non-HDL-C levels prior to, or at the time of initiation of treatment with the combination of the PCSK9 inhibitor and the ANGPTL3 inhibitor.

**[0031]** Other embodiments of the present invention will become apparent from a review of the ensuing detailed description.

#### **BRIEF DESCRIPTION OF THE FIGURES**

**[0032] Figure 1** shows the effect of H4H1276P and H1H316P on LDL-C levels in hyperlipidemic *Ldlr<sup>-/-</sup>* mice when used alone or in combination. The mice were placed on a chow diet and were pre-bled five days before the start of the experiment.

**[0033] Figure 2** shows the effect of H4H1276P and H1H316P on total cholesterol levels in hyperlipidemic *Ldlr<sup>-/-</sup>* mice when used alone or in combination. The mice were placed on a chow diet and were pre-bled five days before the start of the experiment.

**[0034] Figure 3** shows the effect of H4H1276P and H1H316P on HDL-C levels in hyperlipidemic *Ldlr<sup>-/-</sup>* mice when used alone or in combination. The mice were placed on a chow diet and were pre-bled five days before the start of the experiment.

**[0035] Figure 4** shows the effect of H4H1276P and H1H316P on Non HDL-C levels in hyperlipidemic *Ldlr<sup>-/-</sup>* mice when used alone or in combination. The mice were placed on a chow diet and were pre-bled five days before the start of the experiment.

**[0036] Figure 5** shows the effect of H4H1276P and H1H316P on LDL-C levels in hyperlipidemic Ldlr<sup>+/+</sup> mice when used alone or in combination. The mice were placed on a high fat Western diet for 3 weeks prior to treatment and were maintained on this diet throughout the course of the study.

**[0037] Figure 6** shows the effect of H4H1276P and H1H316P on total cholesterol levels in hyperlipidemic Ldlr<sup>+/+</sup> mice when used alone or in combination. The mice were placed on a high fat Western diet for 3 weeks prior to treatment and were maintained on this diet throughout the course of the study.

**[0038] Figure 7** shows the effect of H4H1276P and H1H316P on HDL-C levels in hyperlipidemic Ldlr<sup>+/+</sup> mice when used alone or in combination. The mice were placed on a high fat Western diet for 3 weeks prior to treatment and were maintained on this diet throughout the course of the study.

**[0039] Figure 8** shows the effect of H4H1276P and H1H316P on Non HDL-C levels in hyperlipidemic Ldlr<sup>+/+</sup> mice when used alone or in combination. The mice were placed on a high fat Western diet for 3 weeks prior to treatment and were maintained on this diet throughout the course of the study.

#### DETAILED DESCRIPTION

**[0040]** Before the present invention is described, it is to be understood that this invention is not limited to particular methods and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

**[0041]** Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein, the term "about," when used in reference to a particular recited numerical value, means that the value may vary from the recited value by no more than 1%. For example, as used herein, the expression "about 100" includes 99 and 101 and all values in between (*e.g.*, 99.1, 99.2, 99.3, 99.4, etc.).

**[0042]** Although any methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to describe in their entirety.

#### Methods for Treating Hyperlipidemias

**[0043]** The present invention relates generally to methods and compositions for reducing lipoprotein levels in patients suffering from hypercholesterolemia, who are non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapies (*e.g.* a statin). In

certain embodiments of the invention, treatment with a PCSK9 inhibitor in combination with an ANGPTL3 inhibitor may serve to lower the levels of lipoproteins in these patients to an acceptable range, thereby lowering their risk for development of atherosclerosis, stroke and other cardiovascular diseases. In certain embodiments, the methods described may be used to treat patients suffering from hypercholesterolemia, including heterozygous familial hypercholesterolemia (heFH) and/or homozygous familial hypercholesterolemia (hoFH), in the event that these patients are non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapies.

**[0044]** As used herein, the term "lipoprotein" means a biomolecular particle containing both protein and lipid. Examples of lipoproteins include, *e.g.*, low density lipoprotein (LDL), high-density lipoprotein (HDL), very low density lipoprotein (VLDL), intermediate density lipoprotein (IDL), and lipoprotein (a) (Lp(a)).

**[0045]** The present invention, according to certain embodiments, includes methods for treating patients who are non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapy. As used herein, a particular patient who is "non-responsive to, inadequately controlled by, or intolerant to, standard lipid modifying therapy" is determined by a physician, physician's assistant, diagnostician, or other medical professional on the basis of the level of one or more lipoproteins (*e.g.*, LDL-C and/or non-HDL-C) measured or otherwise detected in the serum of the patient after treatment with the standard lipid modifying agent. The physician, physician's assistant, diagnostician, or other medical professional can also determine if the patient is intolerant to standard lipid modifying therapies based on the side effect profile of the standard lipid modifying therapies, which the patient may experience, including, but not limited to, muscle aches, tenderness or weakness (myalgia), headache, skin flushing, difficulty sleeping, abdominal cramping, bloating, diarrhea, constipation, rash, nausea, or vomiting. A patient who is non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapy may also be determined or influenced by other factors such as the patient's family history, medical background, current therapeutic treatment status, as well as generally accepted or prevailing lipoprotein targets adopted by national medical associations and physicians' groups. For example, in certain contexts, if a patient is undergoing therapy with a standard lipid modifying agent, and exhibits an LDL-C level of greater than or equal to about 70 mg/dL, this indicates that the patient is "non-responsive to, or inadequately controlled by, or intolerant to standard lipid modifying therapy" and may benefit by treatment using the therapies described herein. In other contexts, if a patient is undergoing therapy with a standard lipid modifying agent, and exhibits an LDL-C level of greater than or equal to about 100 mg/dL, this indicates that the patient is "non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapy" and may benefit by treatment using the therapies described herein. In certain contexts, if a patient is undergoing therapy with a standard lipid modifying agent, and exhibits an LDL-C level of greater than or equal to about 150 mg/dL, 200 mg/dL,



250 mg/dL, 300 mg/dL, 400 mg/dL or higher, this indicates that the patient is "non-responsive to, inadequately controlled by, or intolerant to standard lipid modifying therapy" and may benefit by treatment using the therapies described herein. In yet other contexts, whether or not a particular percentage reduction in LDL-C or non-HDL-C level is met, relative to the patient's LDL-C or non-HDL-C level at a particular start point ("baseline") can be used to determine whether the patient has responded to standard lipid modifying therapy or whether that patient is in need of further treatment using the methods and agents of the present invention. For instance, a reduction in LDL-C or non-HDL-C of less than 50% (*e.g.*, less than 40%, less than 35%, less than 30%, less than 25%, etc.) from baseline may signify a need for therapy using the methods and agents of the invention.

**[0046]** The present invention, accordingly, includes methods of treatment comprising administration of one or more doses of a PCSK9 inhibitor combined with one or more doses of an ANGPTL3 inhibitor to a patient, whereby the patient's post-treatment levels of total cholesterol, LDL-C, and/or non-HDL-C are significantly reduced in numbers. For example, the present invention includes therapeutic methods comprising administering one or more doses of a PCSK9 inhibitor and one or more doses of an ANGPTL3 inhibitor to a patient who is undergoing standard lipid modifying therapy, but is non-responsive to such therapy, or is intolerant to such therapy, wherein, after receiving one or more doses of the PCSK9 inhibitor and one or more doses of the ANGPTL3 inhibitor, the patient is able to achieve normal levels of total cholesterol, LDL-C, or non-LDL-C. In certain instances, the patient may be taken off of the standard lipid modifying therapy, or the standard lipid modifying therapy may be continued, but may be administered at lower doses and may be used in combination with the PCSK9 inhibitor and the ANGPTL3 inhibitor, to achieve and/or maintain a particular target lipoprotein level. Alternatively, the patient may be administered the standard lipid modifying therapy at the normal prescribed dose, but the frequency of administration of the lipid modifying therapy may be reduced if the standard lipid modifying therapy is to be administered in conjunction with the combination of the PCSK9 inhibitor and the ANGPTL3 inhibitor. In some instances, the need for treatment with the standard lipid modifying therapy by the patient to achieve and/or maintain a particular target lipoprotein level may be eliminated altogether following administration of one or more doses of the PCSK9 inhibitor in conjunction with the ANGPTL3 inhibitor.

**[0047]** According to certain embodiments, the present invention comprises methods for reducing or eliminating the need for standard lipid modifying therapy, wherein the methods comprise selecting a patient with hyperlipidemia (*e.g.*, hypercholesterolemia) who has been treated with lipid modifying therapy within the last month, the last 2 months, the last 3 months, the last 4 months, the last 5 months, the last 6 months, or for a longer period, and administering one or more doses of a PCSK9 inhibitor in combination with an ANGPTL3 inhibitor to the patient. The methods according to this aspect of the invention result in lowering the level of at least one lipoprotein in the serum of the patient, and consequently allow for a reduction or

elimination of the need for treatment with the standard lipid modifying therapy by the patient. For example, in certain embodiments of the present invention, following administration of one or more doses of a PCSK9 inhibitor in combination with an ANGPTL3 inhibitor, the serum LDL-C level of the patient is reduced to less than a defined level (*e.g.*, less than 100 mg/dL or less than 70 mg/dL), or the total cholesterol is lowered to a defined level (*e.g.* less than 200 mg/dL, or less than 150 mg/dL).

**[0048]** According to certain embodiments, the patient who is treatable by the methods of the present invention has hypercholesterolemia (*e.g.*, a serum LDL-C concentration of greater than or equal to 70 mg/dL, or a serum LDL-C concentration greater than or equal to 100 mg/dL). In certain embodiments, the patient's hypercholesterolemia is inadequately controlled by standard lipid modifying therapy, *e.g.* statin therapy. For example, the present invention includes methods for treating a patient who is non-responsive, inadequately controlled by, or intolerant to, therapy with a standard lipid modifying therapy, such as a statin, or who has hypercholesterolemia that is inadequately controlled by a daily dose of a statin selected from the group consisting of atorvastatin (including atorvastatin + ezetimibe), rosuvastatin, cerivastatin, pitavastatin, fluvastatin, lovastatin, simvastatin (including simvastatin + ezetimibe), pravastatin, and combinations thereof. The present invention also includes methods for reducing cholesterol, LDL-C, or non-LDL-C in a patient who has hypercholesterolemia and who exhibits statin intolerance or who otherwise experiences adverse or undesirable reaction(s) to statin therapy (*e.g.*, skeletal muscle pain, aches, weakness or cramping [*e.g.*, myalgia, myopathy, rhabdomyolysis, etc.]).

### **Patient Selection**

**[0049]** The present invention includes methods and compositions useful for treating patients who are suffering from hyperlipidemia, who are non-responsive to, inadequately controlled by, or intolerant to, therapy with a standard lipid modifying therapy. The patients who are treatable by the methods of the present invention may also exhibit one or more of additional selection criteria. For example, a patient may be selected for treatment with the methods of the present invention if the patient is diagnosed with or identified as being at risk of developing a hypercholesterolemia condition such as, *e.g.*, heterozygous Familial Hypercholesterolemia (heFH), homozygous Familial Hypercholesterolemia (hoFH), Autosomal Dominant Hypercholesterolemia (ADH, *e.g.*, ADH associated with one or more gain-of-function mutations in the PCSK9 gene), autosomal recessive hypercholesterolemia (ARH, *e.g.*, ARH associated with mutations in LDLRAP1), as well as incidences of hypercholesterolemia that are distinct from Familial Hypercholesterolemia (nonFH). Diagnosis of familial hypercholesterolemia (*e.g.*, heFH or hoFH) can be made by genotyping and/or clinical criteria. For patients who are not genotyped, clinical diagnosis may be based on either the Simon Broome criteria with a criteria for definite FH, or the WHO/Dutch Lipid Network criteria with a score > 8 points.

**[0050]** According to certain embodiments, the patient may be selected on the basis of having a history of coronary heart disease (CHD). As used herein a "history of CHD" (or "documented history of CHD") includes one or more of: (i) acute myocardial infarction (MI); (ii) silent MI; (iii) unstable angina; (iv) coronary revascularization procedure (e.g., percutaneous coronary intervention [PCI] or coronary artery bypass graft surgery [CABG]); and/or (v) clinically significant CHD diagnosed by invasive or non-invasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography or nuclear imaging).

**[0051]** According to certain embodiments, the patient may be selected on the basis of having non-coronary heart disease cardiovascular disease ("non-CHD CVD"). As used herein, "non-CHD CVD" includes one or more of: (i) documented previous ischemic stroke with a focal ischemic neurological deficit that persisted more than 24 hours, considered as being of atherothrombotic origin; (ii) peripheral arterial disease; (iii) abdominal aortic aneurysm; (iv) atherosclerotic renal artery stenosis; and/or (v) carotid artery disease (transient ischemic attacks or >50% obstruction of a carotid artery).

**[0052]** According to certain embodiments, the patient may be selected on the basis of having one or more additional risk factors such as, e.g., (i) documented moderate chronic kidney disease (CKD) as defined by  $30 \leq \text{eGFR} < 60 \text{ mL/min/1.73 m}^2$  for 3 months or more; (ii) type 1 or type 2 diabetes mellitus with or without target organ damage (e.g., retinopathy, nephropathy, microalbuminuria); (iii) a calculated 10-year fatal CVD risk SCORE  $\geq 5\%$  (ESC/EAS Guidelines for the management of dyslipidemias, Conroy et al., 2003, Eur. Heart J. 24:987-1003).

**[0053]** According to certain embodiments, the patient may be selected on the basis of having one or more additional risk factors selected from the group consisting of age (e.g., older than 40, 45, 50, 55, 60, 65, 70, 75, or 80 years), race, national origin, gender (male or female), exercise habits (e.g., regular exerciser, non-exerciser), other preexisting medical conditions (e.g., type-II diabetes, high blood pressure, etc.), and current medication status (e.g., currently taking beta blockers, niacin, ezetimibe, fibrates, omega-3 fatty acids, bile acid resins, etc.).

**[0054]** According to certain embodiments of the present invention, the subject who is treatable by the methods of the invention exhibits an elevated level of one or more inflammatory marker. Any marker of systemic inflammation can be utilized for the purposes of the present invention. Suitable inflammatory markers include, without limitation, C-reactive protein, cytokines (e.g., IL-6, IL-8, and/or IL-17), and cellular adhesion molecules (e.g., ICAM-1, ICAM-3, BL-CAM, LFA-2, VCAM-1, NCAM, and PECAM).

**[0055]** According to the present invention, patients may be selected on the basis of a combination of one or more of the foregoing selection criteria or therapeutic characteristics. For example, according to certain embodiments, a patient suitable for treatment with the methods of the present invention, may further be selected on the basis of having heFH or non-FH in combination with: (i) a history of documented CHD, (ii) non-CHD CVD, and/or (iii) diabetes

mellitus with target organ damage; such patients may also be selected on the basis of having a serum LDL-C concentration of greater than or equal to 70 mg/dL.

**[0056]** According to certain other embodiments, a patient suitable for treatment with the methods of the present invention, in addition to having hypercholesterolemia that is not adequately controlled by a daily moderate-dose therapeutic statin regimen, may further be selected on the basis of having heFH or non-FH without CHD, or non-CHD CVD, but having either (i) a calculated 10-year fatal CVD risk SCORE  $\geq 5\%$ ; or (ii) diabetes mellitus without target organ damage; such patients may also be selected on the basis of having a serum LDL-C concentration of greater than or equal to 100 mg/dL.

**[0057]** According to certain embodiments of the present invention, the subject who is treatable by the methods of the invention is a subject who has familial chylomicronemia syndrome (FCS; also known as lipoprotein lipase deficiency).

**[0058]** According to certain embodiments of the present invention, the subject who is treatable by the methods of the invention is a subject who is undergoing, or has recently undergone, lipoprotein apheresis (*e.g.*, within the last six months, within the last 12 weeks, within the last 8 weeks, within the last 6 weeks, within the last 4 weeks, within the last 2 weeks, etc.).

#### **Administration of a PCSK9 Inhibitor plus an ANGPTL3 Inhibitor as Add-On Therapy**

**[0059]** The present invention includes methods of treatment wherein a patient who is undergoing, or has recently undergone, standard lipid modifying therapy (*e.g.* a statin) is administered a PCSK9 inhibitor plus an ANGPTL3 inhibitor according to a particular dosing amount and frequency, and wherein the PCSK9 inhibitor and the ANGPTL3 inhibitor are administered as an add-on to the patient's pre-existing lipid modifying therapy (if applicable), such as an add-on to the patient's pre-existing daily therapeutic statin regimen.

**[0060]** For example, the methods of the present invention include add-on therapeutic regimens wherein the PCSK9 inhibitor and the ANGPTL3 inhibitor are administered as add-on therapy to the same stable daily therapeutic statin regimen (*i.e.*, same dosing amount of statin) that the patient was on prior to receiving the PCSK9 and ANGPTL3 inhibitors. In other embodiments, the PCSK9 and ANGPTL3 inhibitors are administered as add-on therapy to a therapeutic statin regimen comprising a statin in an amount that is more than or less than the dose of statin the patient was on prior to receiving the PCSK9 and ANGPTL3 inhibitors. For example, after starting a therapeutic regimen comprising a PCSK9 inhibitor and an ANGPTL3 inhibitor administered at particular dosing frequencies and amounts, the daily dose of statin administered or prescribed to the patient may (a) stay the same, (b) increase, or (c) decrease (*e.g.*, up-titrate or down-titrate) in comparison to the daily statin dose the patient was taking before starting the PCSK9 and ANGPTL3 inhibitors therapeutic regimen, depending on the therapeutic needs of the patient.

### **Therapeutic Efficacy**

[0061] The methods of the present invention may result in the reduction in serum levels of one or more lipid components selected from the group consisting of total cholesterol, LDL-C, non-HDL-C, ApoB100, VLDL-C, triglycerides, Lp(a) and remnant cholesterol. For example, according to certain embodiments of the present invention, administration of a PCSK9 inhibitor in combination with an ANGPTL3 inhibitor to a suitable subject will result in a mean percent reduction from baseline in serum low density lipoprotein cholesterol (LDL-C) of at least about 25%, 30%, 40%, 50%, 60%, or greater; a mean percent reduction from baseline in ApoB100 of at least about 25%, 30%, 40%, 50%, 60%, or greater; a mean percent reduction from baseline in non-HDL-C of at least about 25%, 30%, 40%, 50%, 60%, or greater; a mean percent reduction from baseline in total cholesterol of at least about 10%, 15%, 20%, 25%, 30%, 35%, or greater; a mean percent reduction from baseline in VLDL-C of at least about 5%, 10%, 15%, 20%, 25%, 30%, or greater; a mean percent reduction from baseline in triglycerides of at least about 5%, 10%, 15%, 20%, 25%, 30%, 35% or greater; and/or a mean percent reduction from baseline in Lp(a) of at least about 5%, 10%, 15%, 20%, 25%, or greater.

### **PCSK9 Inhibitors and ANGPTL3 Inhibitors**

[0062] The methods of the present invention comprise administering to a patient a therapeutic composition comprising a PCSK9 inhibitor and an ANGPTL3 inhibitor.

#### **PCSK9 Inhibitors**

[0063] As used herein, a "PCSK9 inhibitor" is any agent, which binds to or interacts with human PCSK9 and inhibits the normal biological function of PCSK9 *in vitro* or *in vivo*. Non-limiting examples of categories of PCSK9 inhibitors include small molecule PCSK9 antagonists, nucleic acid-based inhibitors of PCSK9 expression or activity (*e.g.*, siRNA or antisense), peptide-based molecules that specifically interact with PCSK9 (*e.g.*, peptibodies), receptor molecules that specifically interact with PCSK9, proteins comprising a ligand-binding portion of an LDL receptor, PCSK9-binding scaffold molecules (*e.g.*, DARPins, HEAT repeat proteins, ARM repeat proteins, tetratricopeptide repeat proteins, fibronectin-based scaffold constructs, and other scaffolds based on naturally occurring repeat proteins, etc., [see, *e.g.*, Boersma and Pluckthun, 2011, *Curr. Opin. Biotechnol.* 22:849-857, and references cited therein]), and anti-PCSK9 aptamers or portions thereof. According to certain embodiments, PCSK9 inhibitors that can be used in the context of the present invention are anti-PCSK9 antibodies or antigen-binding fragments of antibodies that specifically bind human PCSK9.

[0064] The term "human proprotein convertase subtilisin/kexin type 9" or "human PCSK9" or "hPCSK9", as used herein, refers to PCSK9 having the nucleic acid sequence shown in SEQ ID NO:22 and the amino acid sequence of SEQ ID NO:23, or a biologically active fragment thereof.

**ANGPTL3 Inhibitors**

**[0065]** As used herein, an "ANGPTL3 inhibitor" is any agent, which binds to or interacts with human ANGPTL3 and inhibits the normal biological function of ANGPTL3 *in vitro* or *in vivo*. Non-limiting examples of categories of ANGPTL3 inhibitors include small molecule ANGPTL3 antagonists, nucleic acid-based inhibitors of ANGPTL3 expression or activity (*e.g.*, siRNA or antisense), peptide-based molecules that specifically interact with ANGPTL3 (*e.g.*, peptibodies), receptor molecules that specifically interact with ANGPTL3, ANGPTL3-binding scaffold molecules (*e.g.*, DARPin, HEAT repeat proteins, ARM repeat proteins, tetratricopeptide repeat proteins, fibronectin-based scaffold constructs, and other scaffolds based on naturally occurring repeat proteins, etc., [*see, e.g.*, Boersma and Pluckthun, 2011, *Curr. Opin. Biotechnol.* 22:849-857, and references cited therein]), and anti-ANGPTL3 aptamers or portions thereof. According to certain embodiments, ANGPTL3 inhibitors that can be used in the context of the present invention are anti-ANGPTL3 antibodies or antigen-binding fragments of antibodies that specifically bind human ANGPTL3.

**[0066]** The term "human angiotensin-like protein-3" or "human ANGPTL3" or "hANGPTL3", as used herein, refers to ANGPTL3 having the amino acid sequence of SEQ ID NO: 1 (see also NCBI Accession NP\_055310), or a biologically active fragment thereof.

**[0067]** The term "antibody", as used herein, is intended to refer to immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, as well as multimers thereof (*e.g.*, IgM). Each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or V<sub>H</sub>) and a heavy chain constant region. The heavy chain constant region comprises three domains, C<sub>H1</sub>, C<sub>H2</sub> and C<sub>H3</sub>. Each light chain comprises a light chain variable region (abbreviated herein as LCVR or V<sub>L</sub>) and a light chain constant region. The light chain constant region comprises one domain (C<sub>L1</sub>). The V<sub>H</sub> and V<sub>L</sub> regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each V<sub>H</sub> and V<sub>L</sub> is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In different embodiments of the invention, the FRs of the anti-PCSK9 antibody (or antigen-binding portion thereof) may be identical to the human germline sequences, or may be naturally or artificially modified. An amino acid consensus sequence may be defined based on a side-by-side analysis of two or more CDRs.

**[0068]** The term "antibody," as used herein, also includes antigen-binding fragments of full antibody molecules. The terms "antigen-binding portion" of an antibody, "antigen-binding fragment" of an antibody, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. Antigen-binding fragments of an antibody may

be derived, *e.g.*, from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Such DNA is known and/or is readily available from, *e.g.*, commercial sources, DNA libraries (including, *e.g.*, phage-antibody libraries), or can be synthesized. The DNA may be sequenced and manipulated chemically or by using molecular biology techniques, for example, to arrange one or more variable and/or constant domains into a suitable configuration, or to introduce codons, create cysteine residues, modify, add or delete amino acids, etc.

**[0069]** Non-limiting examples of antigen-binding fragments include: (i) Fab fragments; (ii) F(ab')<sub>2</sub> fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (*e.g.*, an isolated complementarity determining region (CDR) such as a CDR3 peptide), or a constrained FR3-CDR3-FR4 peptide. Other engineered molecules, such as domain-specific antibodies, single domain antibodies, domain-deleted antibodies, chimeric antibodies, CDR-grafted antibodies, diabodies, triabodies, tetrabodies, minibodies, nanobodies (*e.g.* monovalent nanobodies, bivalent nanobodies, etc.), small modular immunopharmaceuticals (SMIPs), and shark variable IgNAR domains, are also encompassed within the expression "antigen-binding fragment," as used herein.

**[0070]** An antigen-binding fragment of an antibody will typically comprise at least one variable domain. The variable domain may be of any size or amino acid composition and will generally comprise at least one CDR, which is adjacent to or in frame with one or more framework sequences. In antigen-binding fragments having a V<sub>H</sub> domain associated with a V<sub>L</sub> domain, the V<sub>H</sub> and V<sub>L</sub> domains may be situated relative to one another in any suitable arrangement. For example, the variable region may be dimeric and contain V<sub>H</sub>-V<sub>H</sub>, V<sub>H</sub>-V<sub>L</sub> or V<sub>L</sub>-V<sub>L</sub> dimers. Alternatively, the antigen-binding fragment of an antibody may contain a monomeric V<sub>H</sub> or V<sub>L</sub> domain.

**[0071]** In certain embodiments, an antigen-binding fragment of an antibody may contain at least one variable domain covalently linked to at least one constant domain. Non-limiting, exemplary configurations of variable and constant domains that may be found within an antigen-binding fragment of an antibody of the present invention include: (i) V<sub>H</sub>-C<sub>H</sub>1; (ii) V<sub>H</sub>-C<sub>H</sub>2; (iii) V<sub>H</sub>-C<sub>H</sub>3; (iv) V<sub>H</sub>-C<sub>H</sub>1-C<sub>H</sub>2; (v) V<sub>H</sub>-C<sub>H</sub>1-C<sub>H</sub>2-C<sub>H</sub>3; (vi) V<sub>H</sub>-C<sub>H</sub>2-C<sub>H</sub>3; (vii) V<sub>H</sub>-C<sub>L</sub>; (viii) V<sub>L</sub>-C<sub>H</sub>1; (ix) V<sub>L</sub>-C<sub>H</sub>2; (x) V<sub>L</sub>-C<sub>H</sub>3; (xi) V<sub>L</sub>-C<sub>H</sub>1-C<sub>H</sub>2; (xii) V<sub>L</sub>-C<sub>H</sub>1-C<sub>H</sub>2-C<sub>H</sub>3; (xiii) V<sub>L</sub>-C<sub>H</sub>2-C<sub>H</sub>3; and (xiv) V<sub>L</sub>-C<sub>L</sub>. In any configuration of variable and constant domains, including any of the exemplary configurations listed above, the variable and constant domains may be either directly linked to one another or may be linked by a full or partial hinge or linker region. A hinge region may consist of at least 2 (*e.g.*, 5, 10, 15, 20, 40, 60 or more) amino acids, which result in a flexible or

semi-flexible linkage between adjacent variable and/or constant domains in a single polypeptide molecule. Moreover, an antigen-binding fragment of an antibody of the present invention may comprise a homo-dimer or hetero-dimer (or other multimer) of any of the variable and constant domain configurations listed above in non-covalent association with one another and/or with one or more monomeric V<sub>H</sub> or V<sub>L</sub> domain (e.g., by disulfide bond(s)).

**[0072]** As with full antibody molecules, antigen-binding fragments may be monospecific or multispecific (e.g., bispecific). A multispecific antigen-binding fragment of an antibody will typically comprise at least two different variable domains, wherein each variable domain is capable of specifically binding to a separate antigen or to a different epitope on the same antigen. Any multispecific antibody format, including the exemplary bispecific antibody formats disclosed herein, may be adapted for use in the context of an antigen-binding fragment of an antibody of the present invention using routine techniques available in the art.

**[0073]** The constant region of an antibody is important in the ability of an antibody to fix complement and mediate cell-dependent cytotoxicity. Thus, the isotype of an antibody may be selected on the basis of whether it is desirable for the antibody to mediate cytotoxicity.

**[0074]** The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may nonetheless include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. The term includes antibodies recombinantly produced in a non-human mammal, or in cells of a non-human mammal. The term is not intended to include antibodies isolated from or generated in a human subject.

**[0075]** The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further below), antibodies isolated from a recombinant, combinatorial human antibody library (described further below), antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor et al. (1992) Nucl. Acids Res. 20:6287-6295) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for



human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the  $V_H$  and  $V_L$  regions of the recombinant antibodies are sequences that, while derived from and related to human germline  $V_H$  and  $V_L$  sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

**[0076]** Human antibodies can exist in two forms that are associated with hinge heterogeneity. In one form, an immunoglobulin molecule comprises a stable four chain construct of approximately 150-160 kDa in which the dimers are held together by an interchain heavy chain disulfide bond. In a second form, the dimers are not linked via inter-chain disulfide bonds and a molecule of about 75-80 kDa is formed composed of a covalently coupled light and heavy chain (half-antibody). These forms have been extremely difficult to separate, even after affinity purification.

**[0077]** The frequency of appearance of the second form in various intact IgG isotypes is due to, but not limited to, structural differences associated with the hinge region isotype of the antibody. A single amino acid substitution in the hinge region of the human IgG4 hinge can significantly reduce the appearance of the second form (Angal et al. (1993) *Molecular Immunology* 30:105) to levels typically observed using a human IgG1 hinge. The instant invention encompasses antibodies having one or more mutations in the hinge,  $C_{H2}$  or  $C_{H3}$  region which may be desirable, for example, in production, to improve the yield of the desired antibody form.

**[0078]** An "isolated antibody," as used herein, means an antibody that has been identified and separated and/or recovered from at least one component of its natural environment. For example, an antibody that has been separated or removed from at least one component of an organism, or from a tissue or cell in which the antibody naturally exists or is naturally produced, is an "isolated antibody" for purposes of the present invention. An isolated antibody also includes an antibody *in situ* within a recombinant cell. Isolated antibodies are antibodies that have been subjected to at least one purification or isolation step. According to certain embodiments, an isolated antibody may be substantially free of other cellular material and/or chemicals.

**[0079]** The term "specifically binds," or the like, means that an antibody or antigen-binding fragment thereof forms a complex with an antigen that is relatively stable under physiologic conditions. Methods for determining whether an antibody specifically binds to an antigen are well known in the art and include, for example, equilibrium dialysis, surface plasmon resonance, and the like. For example, an antibody that "specifically binds" PCSK9, or that "specifically binds" ANGPTL3, as used in the context of the present invention, includes antibodies that bind PCSK9, or ANGPTL3, or a portion thereof with a  $K_D$  of less than about 1000 nM, less than about 500 nM, less than about 300 nM, less than about 200 nM, less than about 100 nM, less than about 90 nM, less than about 80 nM, less than about 70 nM, less than about 60 nM, less than about 50 nM, less than about 40 nM, less than about 30 nM, less than about 20 nM, less

than about 10 nM, less than about 5 nM, less than about 4 nM, less than about 3 nM, less than about 2 nM, less than about 1 nM or less than about 0.5 nM, as measured in a surface plasmon resonance assay. An isolated antibody that specifically binds human PCSK9, or human ANGPTL3, however, has cross-reactivity to other antigens, such as PCSK9 molecules, or ANGPTL3 molecules from other (non-human) species.

**[0080]** The anti-PCSK9 and the anti-ANGPTL3 antibodies useful for the methods of the present invention may comprise one or more amino acid substitutions, insertions and/or deletions in the framework and/or CDR regions of the heavy and light chain variable domains as compared to the corresponding germline sequences from which the antibodies were derived. Such mutations can be readily ascertained by comparing the amino acid sequences disclosed herein to germline sequences available from, for example, public antibody sequence databases. The present invention includes methods involving the use of antibodies, and antigen-binding fragments thereof, which are derived from any of the amino acid sequences disclosed herein, wherein one or more amino acids within one or more framework and/or CDR regions are mutated to the corresponding residue(s) of the germline sequence from which the antibody was derived, or to the corresponding residue(s) of another human germline sequence, or to a conservative amino acid substitution of the corresponding germline residue(s) (such sequence changes are referred to herein collectively as "germline mutations"). A person of ordinary skill in the art, starting with the heavy and light chain variable region sequences disclosed herein, can easily produce numerous antibodies and antigen-binding fragments which comprise one or more individual germline mutations or combinations thereof. In certain embodiments, all of the framework and/or CDR residues within the V<sub>H</sub> and/or V<sub>L</sub> domains are mutated back to the residues found in the original germline sequence from which the antibody was derived. In other embodiments, only certain residues are mutated back to the original germline sequence, *e.g.*, only the mutated residues found within the first 8 amino acids of FR1 or within the last 8 amino acids of FR4, or only the mutated residues found within CDR1, CDR2 or CDR3. In other embodiments, one or more of the framework and/or CDR residue(s) are mutated to the corresponding residue(s) of a different germline sequence (*i.e.*, a germline sequence that is different from the germline sequence from which the antibody was originally derived). Furthermore, the antibodies of the present invention may contain any combination of two or more germline mutations within the framework and/or CDR regions, *e.g.*, wherein certain individual residues are mutated to the corresponding residue of a particular germline sequence while certain other residues that differ from the original germline sequence are maintained or are mutated to the corresponding residue of a different germline sequence. Once obtained, antibodies and antigen-binding fragments that contain one or more germline mutations can be easily tested for one or more desired property such as, improved binding specificity, increased binding affinity, improved or enhanced antagonistic or agonistic biological properties (as the case may be), reduced immunogenicity, etc. The use of antibodies and antigen-binding

fragments obtained in this general manner are encompassed within the present invention.

**[0081]** The present invention also includes methods involving the use of anti-PCSK9, and anti-ANGPTL3 antibodies comprising variants of any of the HCVR, LCVR, and/or CDR amino acid sequences disclosed herein having one or more conservative substitutions. For example, the present invention includes the use of anti-PCSK9, and anti-ANGPTL3 antibodies having HCVR, LCVR, and/or CDR amino acid sequences with, *e.g.*, 10 or fewer, 8 or fewer, 6 or fewer, 4 or fewer, etc. conservative amino acid substitutions relative to any of the HCVR, LCVR, and/or CDR amino acid sequences disclosed herein.

**[0082]** The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore™ system (Biacore Life Sciences division of GE Healthcare, Piscataway, NJ).

**[0083]** The term " $K_D$ ", as used herein, is intended to refer to the equilibrium dissociation constant of a particular antibody-antigen interaction.

**[0084]** The term "epitope" refers to an antigenic determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. A single antigen may have more than one epitope. Thus, different antibodies may bind to different areas on an antigen and may have different biological effects. Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. In certain circumstance, an epitope may include moieties of saccharides, phosphoryl groups, or sulfonyl groups on the antigen.

**[0085]** According to certain embodiments, the anti-PCSK9 and anti-ANGPTL3 antibodies used in the methods of the present invention are antibodies with pH-dependent binding characteristics. As used herein, the expression "pH-dependent binding" means that the antibody or antigen-binding fragment thereof exhibits "reduced binding to PCSK9 at acidic pH as compared to neutral pH" (for purposes of the present disclosure, both expressions may be used interchangeably), or that the antibody or antigen-binding fragment thereof exhibits "reduced binding to ANGPTL3 at acidic pH as compared to neutral pH" (for purposes of the present disclosure, both expressions may be used interchangeably). For the example, antibodies "with pH-dependent binding characteristics" includes antibodies and antigen-binding fragments thereof that bind either to PCSK9, or to ANGPTL3 with higher affinity at neutral pH than at acidic pH. In certain embodiments, the antibodies and antigen-binding fragments of the present invention bind PCSK9, or ANGPTL3 with at least 3, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, or more times higher affinity at neutral pH than at acidic pH.

**[0086]** According to this aspect of the invention, the anti-PCSK9 antibodies, or the ANGPTL3

antibodies with pH-dependent binding characteristics may possess one or more amino acid variations relative to the parental anti-PCSK9 antibody, or the parental anti-ANGPTL3 antibody. For example, an anti-PCSK9 antibody, or an anti-ANGPTL3 antibody with pH-dependent binding characteristics may contain one or more histidine substitutions or insertions, *e.g.*, in one or more CDRs of a parental anti-PCSK9, or a parental anti-ANGPTL3 antibody. Thus, according to certain embodiments of the present invention, methods are provided comprising administering an anti-PCSK9 antibody and an anti-ANGPTL3 antibody which comprises CDR amino acid sequences (*e.g.*, heavy and light chain CDRs) which are identical to the CDR amino acid sequences of a parental anti-PCSK9 antibody, or parental ANGPTL3 antibody except for the substitution of one or more amino acids of one or more CDRs of the parental antibody with a histidine residue. The anti-PCSK9 antibodies, or anti-ANGPTL3 antibodies with pH-dependent binding may possess, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or more histidine substitutions, either within a single CDR of a parental antibody or distributed throughout multiple (*e.g.*, 2, 3, 4, 5, or 6) CDRs of a parental anti-PCSK9 antibody, or a parental anti-ANGPTL3 antibody. For example, the present invention includes the use of anti-PCSK9 antibodies and anti-ANGPTL3 antibodies with pH-dependent binding comprising one or more histidine substitutions in HCDR1, one or more histidine substitutions in HCDR2, one or more histidine substitutions in HCDR3, one or more histidine substitutions in LCDR1, one or more histidine substitutions in LCDR2, and/or one or more histidine substitutions in LCDR3, of a parental anti-PCSK9 antibody, or a parental anti-ANGPTL3 antibody.

**[0087]** As used herein, the expression "acidic pH" means a pH of 6.0 or less (*e.g.*, less than about 6.0, less than about 5.5, less than about 5.0, etc.). The expression "acidic pH" includes pH values of about 6.0, 5.95, 5.90, 5.85, 5.8, 5.75, 5.7, 5.65, 5.6, 5.55, 5.5, 5.45, 5.4, 5.35, 5.3, 5.25, 5.2, 5.15, 5.1, 5.05, 5.0, or less. As used herein, the expression "neutral pH" means a pH of about 7.0 to about 7.4. The expression "neutral pH" includes pH values of about 7.0, 7.05, 7.1, 7.15, 7.2, 7.25, 7.3, 7.35, and 7.4.

**[0088]** Non-limiting examples of anti-PCSK9 antibodies that can be used in the context of the present invention include, *e.g.*, alirocumab, evolocumab, bococizumab, lodelcizumab, ralpanzumab, or antigen-binding portions of any of the foregoing antibodies.

**[0089]** A non-limiting example of an anti-ANGPTL3 antibody that can be used in the context of the present invention includes evinacumab.

### **Preparation of Human Antibodies**

**[0090]** Anti-PCSK9 antibodies and anti-ANGPTL3 antibodies can be made according to any method of antibody production/isolation known in the art. For example, antibodies for use in the methods of the present invention may be made by hybridoma technologies, by phage display, by yeast display, etc. Antibodies for use in the methods of the present invention may be, *e.g.*, chimeric antibodies, humanized antibodies, or fully human antibodies.

**[0091]** Methods for generating human antibodies in transgenic mice are known in the art. Any such known methods can be used in the context of the present invention to make human antibodies that specifically bind PCSK9, or ANGPTL3.

**[0092]** For example, using VELOCIMMUNE™ technology (see, for example, US 6,596,541, Regeneron Pharmaceuticals) or any other known method for generating monoclonal antibodies, high affinity chimeric antibodies to PCSK9, or to ANGPTL3 are initially isolated having a human variable region and a mouse constant region. The VELOCIMMUNE® technology involves generation of a transgenic mouse having a genome comprising human heavy and light chain variable regions operably linked to endogenous mouse constant region loci such that the mouse produces an antibody comprising a human variable region and a mouse constant region in response to antigenic stimulation. The DNA encoding the variable regions of the heavy and light chains of the antibody are isolated and operably linked to DNA encoding the human heavy and light chain constant regions. The DNA is then expressed in a cell capable of expressing the fully human antibody.

**[0093]** Generally, a VELOCIMMUNE® mouse is challenged with the antigen of interest, and lymphatic cells (such as B-cells) are recovered from the mice that express antibodies. The lymphatic cells may be fused with a myeloma cell line to prepare immortal hybridoma cell lines, and such hybridoma cell lines are screened and selected to identify hybridoma cell lines that produce antibodies specific to the antigen of interest. DNA encoding the variable regions of the heavy chain and light chain may be isolated and linked to desirable isotypic constant regions of the heavy chain and light chain. Such an antibody protein may be produced in a cell, such as a CHO cell. Alternatively, DNA encoding the antigen-specific chimeric antibodies or the variable domains of the light and heavy chains may be isolated directly from antigen-specific lymphocytes.

**[0094]** Initially, high affinity chimeric antibodies are isolated having a human variable region and a mouse constant region. The antibodies are characterized and selected for desirable characteristics, including affinity, selectivity, epitope, etc., using standard procedures known to those skilled in the art. The mouse constant regions are replaced with a desired human constant region to generate the fully human antibody of the invention, for example wild-type or modified IgG1 or IgG4. While the constant region selected may vary according to specific use, high affinity antigen-binding and target specificity characteristics reside in the variable region.

**[0095]** In general, the antibodies that can be used in the methods of the present invention possess high affinities, as described above, when measured by binding to antigen either immobilized on solid phase or in solution phase. The mouse constant regions are replaced with desired human constant regions to generate the fully human antibodies of the invention. While the constant region selected may vary according to specific use, high affinity antigen-binding and target specificity characteristics reside in the variable region.

[0096] Specific examples of human antibodies or antigen-binding fragments of antibodies that specifically bind PCSK9, which can be used in the context of the methods of the present invention include antibodies or antigen-binding proteins comprising the six CDRs (HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and LCDR3) from the heavy and light chain variable region (HCVR/LCVR) amino acid sequence pair comprising SEQ ID NOs: 12/17.

[0097] In certain embodiments of the present invention, the anti-PCSK9 antibody, or antigen-binding fragment thereof, that can be used in the methods of the present invention comprises heavy and light chain complementarity determining regions (HCDR1-HCDR2-HCDR3/LCDR1-LCDR2-LCDR3) comprising the amino acid sequences of SEQ ID NOs:13, 14, 15, 18, 19 and 21.

[0098] In certain embodiments of the present invention, the anti-PCSK9 antibody, or antigen-binding fragment thereof, that can be used in the methods of the present invention comprises an HCVR having the amino acid sequence of SEQ ID NO:12 and an LCVR having the amino acid sequence of SEQ ID NO:17.

[0099] Specific examples of human antibodies or antigen-binding fragments of antibodies that specifically bind ANGPTL3, which can be used in the context of the methods of the present invention include antibodies or antigen-binding proteins comprising the six CDRs (HCDR1, HCDR2, HCDR3, LCDR1, LCDR2 and LCDR3) from the heavy and light chain variable region (HCVR/LCVR) amino acid sequence pair comprising SEQ ID NOs: 2/3.

[0100] In certain embodiments of the present invention, the anti-ANGPTL3 antibody, or antigen-binding fragment thereof, that can be used in the methods of the present invention comprises heavy and light chain complementarity determining regions (HCDR1-HCDR2-HCDR3/LCDR1-LCDR2-LCDR3) comprising the amino acid sequences of SEQ ID NOs:4, 5, 6, 7, 8 and 9.

[0101] In certain embodiments of the present invention, the anti-ANGPTL3 antibody, or antigen-binding fragment thereof, that can be used in the methods of the present invention comprises an HCVR having the amino acid sequence of SEQ ID NO:2 and an LCVR having the amino acid sequence of SEQ ID NO:3.

### **Pharmaceutical Compositions and Methods of Administration**

[0102] The present invention includes methods, which comprise administering a PCSK9 inhibitor to a patient in combination with an ANGPTL3 inhibitor, wherein the PCSK9 inhibitor and the ANGPTL3 inhibitor are contained within the same, or in different pharmaceutical compositions. The pharmaceutical compositions of the invention are formulated with suitable carriers, excipients, and other agents that provide suitable transfer, delivery, tolerance, and the like. A multitude of appropriate formulations can be found in the formulary known to all pharmaceutical chemists: Remington's Pharmaceutical Sciences, Mack Publishing Company,

Easton, PA. These formulations include, for example, powders, pastes, ointments, jellies, waxes, oils, lipids, lipid (cationic or anionic) containing vesicles (such as LIPOFECTIN™), DNA conjugates, anhydrous absorption pastes, oil-in-water and water-in-oil emulsions, emulsions carbowax (polyethylene glycols of various molecular weights), semi-solid gels, and semi-solid mixtures containing carbowax. See also Powell et al. "Compendium of excipients for parenteral formulations" PDA (1998) J Pharm Sci Technol 52:238-311.

**[0103]** Exemplary pharmaceutical formulations comprising anti-PCSK9 antibodies, and/or ANGPTL3 antibodies that can be used in the context of the present invention include any of the formulations as set forth in US 8,795,669 (describing, *inter alia*, exemplary formulations comprising alirocumab), or in WO2013/166448, or WO2012/168491.

**[0104]** Various delivery systems are known and can be used to administer the pharmaceutical composition of the invention, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the mutant viruses, receptor mediated endocytosis (see, *e.g.*, Wu et al., 1987, J. Biol. Chem. 262:4429-4432). Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The composition may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents.

**[0105]** A pharmaceutical composition of the present invention can be delivered subcutaneously or intravenously with a standard needle and syringe. In addition, with respect to subcutaneous delivery, a pen delivery device readily has applications in delivering a pharmaceutical composition of the present invention. Such a pen delivery device can be reusable or disposable. A reusable pen delivery device generally utilizes a replaceable cartridge that contains a pharmaceutical composition. Once all of the pharmaceutical composition within the cartridge has been administered and the cartridge is empty, the empty cartridge can readily be discarded and replaced with a new cartridge that contains the pharmaceutical composition. The pen delivery device can then be reused. In a disposable pen delivery device, there is no replaceable cartridge. Rather, the disposable pen delivery device comes prefilled with the pharmaceutical composition held in a reservoir within the device. Once the reservoir is emptied of the pharmaceutical composition, the entire device is discarded.

**[0106]** Numerous reusable pen and autoinjector delivery devices have applications in the subcutaneous delivery of a pharmaceutical composition of the present invention. Examples include, but are not limited to AUTOPEN™ (Owen Mumford, Inc., Woodstock, UK), DISETRONIC™ pen (Disetronic Medical Systems, Bergdorf, Switzerland), HUMALOG MIX 75/25™ pen, HUMALOG™ pen, HUMALIN 70/30™ pen (Eli Lilly and Co., Indianapolis, IN), NOVOPEN™ I, II and III (Novo Nordisk, Copenhagen, Denmark), NOVOPEN JUNIOR™ (Novo

Nordisk, Copenhagen, Denmark), BD™ pen (Becton Dickinson, Franklin Lakes, NJ), OPTIPEN™, OPTIPEN PRO™, OPTIPEN STARLET™, and OPTICLIK™ (Sanofi-Aventis, Frankfurt, Germany), to name only a few. Examples of disposable pen delivery devices having applications in subcutaneous delivery of a pharmaceutical composition of the present invention include, but are not limited to the SOLOSTAR™ pen (Sanofi-Aventis), the FLEXPEN™ (Novo Nordisk), and the KWIKPEN™ (Eli Lilly), the SURECLICK™ Autoinjector (Amgen, Thousand Oaks, CA), the PENLET™ (Haselmeier, Stuttgart, Germany), the EPIPEN (Dey, L.P.), and the HUMIRA™ Pen (Abbott Labs, Abbott Park IL), to name only a few.

**[0107]** In certain situations, the pharmaceutical composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:201). In another embodiment, polymeric materials can be used; see, *Medical Applications of Controlled Release*, Langer and Wise (eds.), 1974, CRC Pres., Boca Raton, Florida. In yet another embodiment, a controlled release system can be placed in proximity of the composition's target, thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, 1984, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138). Other controlled release systems are discussed in the review by Langer, 1990, *Science* 249:1527-1533.

**[0108]** The injectable preparations may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by known methods. For example, the injectable preparations may be prepared, *e.g.*, by dissolving, suspending or emulsifying the antibody or its salt described above in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (*e.g.*, ethanol), a polyalcohol (*e.g.*, propylene glycol, polyethylene glycol), a nonionic surfactant [*e.g.*, polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, *e.g.*, sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule.

**[0109]** Advantageously, the pharmaceutical compositions for oral or parenteral use described above are prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc.

### **Dosage**

**[0110]** The amount of a PCSK9 inhibitor (*e.g.*, anti-PCSK9 antibody), or an ANGPTL3 inhibitor (*e.g.*, anti-ANGPTL3 antibody) administered to a subject according to the methods of the



present invention is, generally, a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount of a PCSK9 inhibitor" means a dose of a PCSK9 inhibitor, when administered in combination with an ANGPTL3 inhibitor, results in a detectable reduction (at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or more from baseline) in one or more parameters selected from the group consisting of total cholesterol, LDL-C, ApoB100, non-HDL-C, VLDL-C, triglycerides, Lp(a) and remnant cholesterol, or an amount that reduces or eliminates a patient's need for other therapeutic interventions, such as, lipoprotein apheresis, or that reduces a patient's normalized rate of apheresis.

**[0111]** In the case of an anti-PCSK9 antibody, a therapeutically effective amount can be from about 0.05 mg to about 600 mg, *e.g.*, about 0.05 mg, about 0.1 mg, about 1.0 mg, about 1.5 mg, about 2.0 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, about 110 mg, about 120 mg, about 130 mg, about 140 mg, about 160 mg, about 170 mg, about 180 mg, about 190 mg, about 200 mg, about 210 mg, about 220 mg, about 230 mg, about 240 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 310 mg, about 320 mg, about 330 mg, about 340 mg, about 350 mg, about 360 mg, about 370 mg, about 380 mg, about 390 mg, about 400 mg, about 410 mg, about 420 mg, about 430 mg, about 440 mg, about 450 mg, about 460 mg, about 470 mg, about 480 mg, about 490 mg, about 500 mg, about 510 mg, about 520 mg, about 530 mg, about 540 mg, about 550 mg, about 560 mg, about 570 mg, about 580 mg, about 590 mg, or about 600 mg, of the anti-PCSK9 antibody. According to certain exemplary embodiments of the present invention, a therapeutically effective amount of an anti-PCSK9 antibody is 75 mg, 150 mg or 300 mg (*e.g.*, in the case of alirocumab), or 140 mg or 420 mg (*e.g.*, in the case of evolocumab). Other dosing amounts of PCSK9 inhibitors will be apparent to persons of ordinary skill in the art and are contemplated within the scope of the present invention.

**[0112]** The amount of anti-PCSK9 antibody contained within the individual doses may be expressed in terms of milligrams of antibody per kilogram of patient body weight (*i.e.*, mg/kg). For example, the anti-PCSK9 antibody may be administered to a patient at a dose of about 0.0001 to about 10 mg/kg of patient body weight.

**[0113]** The amount of ANGPTL3 inhibitor (*e.g.*, anti-ANGPTL3 antibody) administered to a subject according to the methods of the present invention is, generally, a therapeutically effective amount. As used herein, the phrase "therapeutically effective amount of an ANGPTL3 inhibitor" means a dose of ANGPTL3 inhibitor, when combined with a PCSK9 inhibitor, results in a detectable reduction (at least about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or more from baseline) in one or more parameters selected from the group consisting of total cholesterol, LDL-C, ApoB100, non-HDL-C, VLDL-C,

triglycerides, Lp(a) and remnant cholesterol, or an amount that prevents or attenuates atherosclerosis in a subject (as described elsewhere herein).

**[0114]** In the case of an anti-ANGPTL3 antibody, a therapeutically effective amount can be from about 0.05 mg to about 600 mg, *e.g.*, about 0.05 mg, about 0.1 mg, about 1.0 mg, about 1.5 mg, about 2.0 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, about 110 mg, about 120 mg, about 130 mg, about 140 mg, about 160 mg, about 170 mg, about 180 mg, about 190 mg, about 200 mg, about 210 mg, about 220 mg, about 230 mg, about 240 mg, about 250 mg, about 260 mg, about 270 mg, about 280 mg, about 290 mg, about 300 mg, about 310 mg, about 320 mg, about 330 mg, about 340 mg, about 350 mg, about 360 mg, about 370 mg, about 380 mg, about 390 mg, about 400 mg, about 410 mg, about 420 mg, about 430 mg, about 440 mg, about 450 mg, about 460 mg, about 470 mg, about 480 mg, about 490 mg, about 500 mg, about 510 mg, about 520 mg, about 530 mg, about 540 mg, about 550 mg, about 560 mg, about 570 mg, about 580 mg, about 590 mg, or about 600 mg, of the anti-ANGPTL3 antibody. Other dosing amounts of ANGPTL3 inhibitors will be apparent to persons of ordinary skill in the art and are contemplated within the scope of the present invention.

**[0115]** The amount of anti-ANGPTL3 antibody contained within the individual doses may be expressed in terms of milligrams of antibody per kilogram of patient body weight (*i.e.*, mg/kg). For example, the anti-ANGPTL3 antibody may be administered to a patient at a dose of about 0.0001 to about 10 mg/kg of patient body weight.

### **Combination Therapies**

**[0116]** As described elsewhere herein, the methods of the present invention may comprise administering a PCSK9 inhibitor in combination with an ANGPTL3 inhibitor to a patient who is non-responsive to, inadequately controlled by, or intolerant to a standard lipid lowering therapy. In certain embodiments, the need for further administration of the lipid lowering therapy may be eliminated altogether. In certain embodiments, the combined use of the PCSK9 inhibitor with the ANGPTL3 inhibitor may be used in combination with ("on top of") the patient's previously prescribed lipid lowering therapy. For example, in the context of lowering at least one lipid/lipoprotein parameter in a patient suffering from hyperlipidemia (*e.g.* hypercholesterolemia), wherein the patient is non-responsive to, inadequately controlled by, or intolerant to a standard lipid lowering therapy, a combination of a PCSK9 inhibitor with an ANGPTL3 inhibitor may be administered to a patient in combination with a stable daily therapeutic statin regimen. Exemplary daily therapeutic statin regimens that a PCSK9 inhibitor plus an ANGPTL3 inhibitor may be administered in combination with in the context of the present invention include, *e.g.*, atorvastatin (10, 20, 40 or 80 mg daily), (atorvastatin/ezetimibe 10/10 or 40/10 mg daily), rosuvastatin (5, 10 or 20 mg daily), cerivastatin (0.4 or 0.8 mg daily), pitavastatin (1, 2 or 4 mg daily), fluvastatin (20, 40 or 80 mg daily), simvastatin (5, 10, 20, 40 or

80 mg daily), simvastatin/ezetimibe (10/10, 20/10, 40/10 or 80/10 mg daily), lovastatin (10, 20, 40 or 80 mg daily), pravastatin (10, 20, 40 or 80 mg daily), and combinations thereof. Other lipid modifying therapies that a PCSK9 inhibitor plus an ANGPTL3 inhibitor may be administered in combination with in the context of the present invention include, *e.g.*, (1) an agent which inhibits cholesterol uptake and or bile acid re-absorption (*e.g.*, ezetimibe); (2) an agent which increase lipoprotein catabolism (such as niacin); and/or (3) activators of the LXR transcription factor that plays a role in cholesterol elimination such as 22-hydroxycholesterol.

**[0117]** Non-limiting examples of anti-PCSK9 antibodies that can be used in the context of the present invention include, *e.g.*, alirocumab, evolocumab, bococizumab, lodalcizumab, ralpancizumab, or antigen-binding portions of any of the foregoing antibodies.

**[0118]** A non-limiting example of an ANGPTL3 antibody to be used in the context of the present invention includes evinacumab.

### **Administration Regimens**

**[0119]** According to certain embodiments of the present invention, multiple doses of a PCSK9 inhibitor (*i.e.*, a pharmaceutical composition comprising a PCSK9 inhibitor) and an ANGPTL3 inhibitor (*i.e.*, a pharmaceutical composition comprising an ANGPTL3 inhibitor) may be administered to a subject over a defined time course (*e.g.*, on top of a daily therapeutic statin regimen or other background lipid modifying therapy). The methods according to this aspect of the invention comprise sequentially administering to a subject multiple doses of a PCSK9 inhibitor and an ANGPTL3 inhibitor. As used herein, "sequentially administering" means that each dose of PCSK9 inhibitor and ANGPTL3 inhibitor is administered to the subject at a different point in time, *e.g.*, on different days separated by a predetermined interval (*e.g.*, hours, days, weeks or months). The present invention includes methods which comprise sequentially administering to the patient a single initial dose of a PCSK9 inhibitor and an ANGPTL3 inhibitor, followed by one or more secondary doses of the PCSK9 inhibitor and ANGPTL3 inhibitor, and optionally followed by one or more tertiary doses of the PCSK9 inhibitor and ANGPTL3 inhibitor.

**[0120]** The terms "initial dose," "secondary doses," and "tertiary doses," refer to the temporal sequence of administration of the individual doses of a pharmaceutical composition comprising a PCSK9 inhibitor and the ANGPTL3 inhibitor. Thus, the "initial dose" is the dose which is administered at the beginning of the treatment regimen (also referred to as the "baseline dose"); the "secondary doses" are the doses which are administered after the initial dose; and the "tertiary doses" are the doses which are administered after the secondary doses. The initial, secondary, and tertiary doses may all contain the same amount of the PCSK9 inhibitor and the ANGPTL3 inhibitor, but generally may differ from one another in terms of frequency of administration. In certain embodiments, however, the amount of PCSK9 inhibitor and the ANGPTL3 inhibitor contained in the initial, secondary and/or tertiary doses varies from one

another (*e.g.*, adjusted up or down as appropriate) during the course of treatment. In certain embodiments, two or more (*e.g.*, 2, 3, 4, or 5) doses are administered at the beginning of the treatment regimen as "loading doses" followed by subsequent doses that are administered on a less frequent basis (*e.g.*, "maintenance doses").

**[0121]** According to exemplary embodiments of the present invention, each secondary and/or tertiary dose is administered 1 to 26 (*e.g.*, 1, 1½, 2, 2½, 3, 3½, 4, 4½, 5, 5½, 6, 6½, 7, 7½, 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12, 12½, 13, 13½, 14, 14½, 15, 15½, 16, 16½, 17, 17½, 18, 18½, 19, 19½, 20, 20½, 21, 21½, 22, 22½, 23, 23½, 24, 24½, 25, 25½, 26, 26½, or more) weeks after the immediately preceding dose. The phrase "the immediately preceding dose," as used herein, means, in a sequence of multiple administrations, the dose of antigen-binding molecule, which is administered to a patient prior to the administration of the very next dose in the sequence with no intervening doses.

**[0122]** The methods according to this aspect of the invention may comprise administering to a patient any number of secondary and/or tertiary doses of a PCSK9 inhibitor and ANGPTL3 inhibitor. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient.

**[0123]** In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 1 to 2, 4, 6, 8 or more weeks after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 1 to 2, 4, 6, 8 or more weeks after the immediately preceding dose. Alternatively, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

### EXAMPLES

**[0124]** The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is

average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

#### **Example 1. Generation of Human Antibodies to Human PCSK9**

[0125] Human anti-PCSK9 antibodies were generated as described in US Patent No. 8,062,640. The exemplary PCSK9 inhibitor used in the following Example is the human anti-PCSK9 antibody designated "H1H316P," also known as "alirocumab", or "PRAULENT®". H1H316P has the following amino acid sequence characteristics: a heavy chain comprising SEQ ID NO:16 and a light chain comprising SEQ ID NO:20; a heavy chain variable region (HCVR) comprising SEQ ID NO:12 and a light chain variable domain (LCVR) comprising SEQ ID NO:17; a heavy chain complementarity determining region 1 (HCDR1) comprising SEQ ID NO:13, a HCDR2 comprising SEQ ID NO:14, a HCDR3 comprising SEQ ID NO:15, a light chain complementarity determining region 1 (LCDR1) comprising SEQ ID NO:18, a LCDR2 comprising SEQ ID NO:19 and a LCDR3 comprising SEQ ID NO:21.

#### **Example 2. Generation of Human Antibodies to Human ANGPTL3**

[0126] Human anti-ANGPTL3 antibodies were generated as described in US Patent No. 9,018,356. The exemplary ANGPTL3 inhibitor used in the following Example is the human anti-ANGPTL3 antibody designated "H4H1276S," also known as "evinacumab." H4H1276S has the following amino acid sequence characteristics: a heavy chain comprising SEQ ID NO:10 and a light chain comprising SEQ ID NO:11; a heavy chain variable region (HCVR) comprising SEQ ID NO:2 and a light chain variable domain (LCVR) comprising SEQ ID NO:3; a heavy chain complementarity determining region 1 (HCDR1) comprising SEQ ID NO:4, a HCDR2 comprising SEQ ID NO:5, a HCDR3 comprising SEQ ID NO:6, a light chain complementarity determining region 1 (LCDR1) comprising SEQ ID NO:7, a LCDR2 comprising SEQ ID NO:8 and a LCDR3 comprising SEQ ID NO:9.

#### **Example 3: *In Vivo* Effect of Treatment with a combination of an Anti-hANGPTL3 antibody and an anti-PCSK9 antibody on circulating lipid levels in hyperlipidemic Ldlr<sup>-/+</sup> mice**

[0127] The effect of anti-hANGPTL3 antibody H4H1276 (evinacumab) alone, anti-PCSK9 antibody H1H316P (alirocumab) alone and both antibodies in combination on serum lipids levels was determined in LDLR<sup>-/+</sup> mice. These mice are hyperlipidemic with majority of their circulating cholesterol found in the form of LDL due to partial deficiency in LDLR, the major receptor for LDL-C uptake.

[0128] In the first study, male LDLR<sup>-/+</sup> mice on chow diet were pre-bled 5 days before the experiment and mice were put into groups of five mice each. The antibodies, H4H1276P, H1H316P, their combination and isotype-matched (hlgG4) control with irrelevant specificity,

were administered at a dose of 10mg/kg each by subcutaneous injection on Day 0 of the study. Mice were bled after 4 hours of fasting on consecutive days after the antibodies injections and serum lipids levels (Total Cholesterol, LDL-C, Non-HDL-C and HDL-C) were determined in the serum by ADVIA® 1800 Chemistry System (Siemens). Averages per group were calculated for each of the time points. Results, expressed as mean  $\pm$  SEM of serum lipids concentration, are shown in Figures 1, 2, 3, and 4.

**[0129]** In the second study, male LDLR<sup>-/-</sup> mice were placed on a high fat Western diet for 3 weeks before injection of the antibodies and the mice were fed this diet through the duration of the study. The rest of the study was conducted under the same protocol as for the first study. Results, expressed as mean  $\pm$  SEM of serum lipids concentration, are shown in Figures 5, 6, 7 and 8.

**[0130]** Levels of circulating antibodies (Serum Ab) for both studies were determined using a standard ELISA assay. Briefly, plates were coated with a goat anti-human Fc antibody (Sigma-Aldrich) to capture Serum Ab. Serum was then added to the plates and captured antibodies were detected by chemiluminescence using a horseradish peroxidase (HRP) conjugated goat anti-human IgG antibody (Sigma-Aldrich). Results, expressed as mean  $\pm$  SEM are shown in Tables 1A and 1B (first study) and Tables 2A and 2B (second study). Control: Mice that received an isotype-matched Control Ab

**Results summary:**

**[0131]** The administration of the combination of H1H316P and H4H1276P as single subcutaneous doses to LDLR<sup>-/-</sup> mice on chow and high fat Western diet lead to a significant reduction in total cholesterol, LDL-C and non-HDL-C and had an additive affect on serum lipid levels when compared to the respective single administration of each antibody.

We claim:

1. A method of treating a patient suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy, the method comprising treating the patient with a combination of a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor and an inhibitor of angiotensin-like protein 3 (ANGPTL3).
2. The method of claim 1, wherein the hypercholesterolemia is heterozygous familial hypercholesterolemia (HeFH) or homozygous familial hypercholesterolemia (HoFH).
3. The method of claim 1, wherein the PCSK9 inhibitor is an antibody, or an antigen-binding fragment thereof, that binds specifically to PCSK9.
4. The method of claim 3, wherein the PCSK9 antibody is administered to the patient at a dose of about 75 mg at a frequency of once every two weeks.
5. The method of claim 3, wherein the PCSK9 antibody is administered to the patient at a dose of about 140 mg at a frequency of once every two weeks.
6. The method of claim 3, wherein the PCSK9 antibody is administered to the patient at a dose of about 150 mg at a frequency of once every two weeks, or once every four weeks.
7. The method of claim 3, wherein the PCSK9 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every four weeks.
8. The method of claim 3, wherein the PCSK9 antibody is administered to the patient at a dose of about 420 mg at a frequency of once every four weeks.
9. The method of any of claims 3-8, wherein the PCSK9 antibody is selected from the group consisting of alirocumab, evolocumab, bococizumab, lodelcizumab and ralpancizumab.
10. The method of any one of claims 3-9, wherein the antibody is alirocumab.

11. The method of any of claims 3-10, wherein the antibody or antigen-binding fragment thereof that binds specifically to PCSK9 comprises the complementary determining regions (CDRs) of a heavy chain variable (HCVR) having the amino acid sequence of SEQ ID NO: 12 and the CDRs of a light chain variable region (LCVR) of SEQ ID NO: 17.

12. The method of any of claims 3-11, wherein the antibody or antigen-binding fragment thereof that binds specifically to PCSK9 comprises a heavy chain CDR1 (HCDR1) having the amino acid sequence of SEQ ID NO: 13, a HCDR2 having the amino acid sequence of SEQ ID NO: 14, a HCDR3 having the amino acid sequence of SEQ ID NO: 15, a light chain CDR1 (LCDR1) having the amino acid sequence of SEQ ID NO: 18, a LCDR2 having the amino acid sequence of SEQ ID NO: 19, and a LCDR3 having the amino acid sequence of SEQ ID NO: 21.

13. The method of any of claims 3-12, wherein the antibody or antigen-binding fragment thereof that binds specifically to PCSK9 comprises a HCVR having the amino acid sequence of SEQ ID NO: 12 and a LCVR having the amino acid sequence of SEQ ID NO: 17.

14. The method of claim 3, wherein the PCSK9 antibody is administered to the patient subcutaneously or intravenously.

15. The method of claim 1, wherein the ANGPTL3 inhibitor is an antibody, or an antigen-binding fragment thereof, that binds specifically to ANGPTL3.

16. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 150 mg at a frequency of once every week.

17. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every week.

18. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 450 mg at a frequency of once every week.

19. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 300 mg at a frequency of once every two weeks.

20. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 450 mg at a frequency of once every two weeks.



21. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient at a dose of about 20 mg/kg at a frequency of once every four weeks.
22. The method of any of claims 15-21, wherein the ANGPTL3 antibody is evinacumab.
23. The method of any of claims 15-22, wherein the antibody or antigen-binding fragment thereof that binds specifically to ANGPTL3 comprises the complementary determining regions (CDRs) of a heavy chain variable (HCVR) having the amino acid sequence of SEQ ID NO: 2 and the CDRs of a light chain variable region (LCVR) of SEQ ID NO: 3.
24. The method of any of claims 15-23, wherein the antibody or antigen-binding fragment thereof that binds specifically to ANGTL3 comprises a heavy chain CDR1 (HCDR1) having the amino acid sequence of SEQ ID NO: 4, a HCDR2 having the amino acid sequence of SEQ ID NO: 5, a HCDR3 having the amino acid sequence of SEQ ID NO: 6, a light chain CDR1 (LCDR1) having the amino acid sequence of SEQ ID NO: 7, a LCDR2 having the amino acid sequence of SEQ ID NO: 8, and a LCDR3 having the amino acid sequence of SEQ ID NO: 9.
25. The method of any of claims 15-24, wherein the antibody or antigen-binding fragment thereof that binds specifically to ANGPTL3 comprises a HCVR having the amino acid sequence of SEQ ID NO: 2 and a LCVR having the amino acid sequence of SEQ ID NO: 3.
26. The method of claim 15, wherein the ANGPTL3 antibody is administered to the patient subcutaneously or intravenously.
27. The method of any one of claims 1-26, wherein the treating with a combination of a PCSK9 inhibitor and an inhibitor of ANGPTL3 results in lowering one or more of the following parameters:
- (a) a reduction in serum total cholesterol (TC) level;
  - (b) a reduction in serum low-density lipoprotein cholesterol (LDL-C) level; and
  - (c) a reduction in serum non-high density lipoprotein cholesterol (non-HDL-C) level
- in the patient, wherein the reduction of (a), (b), and/or (c) is determined relative to the patient's serum TC level, serum LDL-C level, and/or serum non-HDL-C level prior to, or at the time of initiation of, treatment with the combination of the PCSK9 inhibitor and the ANGPTL3 inhibitor.

28. A use of a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor in combination with an inhibitor of angiotensin-like protein 3 (ANGPTL3) in the treatment of a patient suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy.

29. A use of a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor in combination with an inhibitor of angiotensin-like protein 3 (ANGPTL3) in the preparation of a medicament for treating a patient suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy.

30. A pharmaceutical composition for treating a patient suffering from hypercholesterolemia, wherein the patient is non-responsive to, inadequately controlled by, or intolerant to treatment with a standard lipid modifying therapy, wherein the composition comprises a proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor in combination with an inhibitor of angiotensin-like protein 3 (ANGPTL3).




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6






Figure 7




Figure 8

INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/020221

A. CLASSIFICATION OF SUBJECT MATTER  
INV. A61P9/10 A61K39/00 C07K16/22 C07K16/40  
ADD.  
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
Minimum documentation searched (classification system followed by classification symbols)  
A61K A61P C07K  
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2012/174178 A1 (REGENERON PHARMA [US]; SLEEMAN MARK W [AU]; GUSAROVA VIKTORIA [US]; MU) 20 December 2012 (2012-12-20) paragraph [0097]; claim 24; examples 7-9; table 1	1-30
X	TIKKA ANNA ET AL: "The role of ANGPTL3 in controlling lipoprotein metabolism", ENDOCRINE, HUMANA PRESS, INC, US, vol. 52, no. 2, 11 January 2016 (2016-01-11), pages 187-193, XP035952116, ISSN: 1355-008X, DOI: 10.1007/S12020-015-0838-9 [retrieved on 2016-01-11] page 187, right-hand column, lines 15-20; figure 1	1-30

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 10 May 2017	Date of mailing of the international search report 19/05/2017
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Cilensek, Zoran
--	---------------------------------------

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2017/020221

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2016/011256 A1 (SANOFI BIOTECHNOLOGY [FR]; REGENERON PHARMA [US]) 21 January 2016 (2016-01-21) the whole document	1-30
A	----- WO 2014/194168 A2 (REGENERON PHARMA [US]) 4 December 2014 (2014-12-04) the whole document	1-30
A	----- VIKTORIA GUSAROVA ET AL: "ANGPTL3 blockade with a human monoclonal antibody reduces plasma lipids in dyslipidemic mice and monkeys", JOURNAL OF LIPID RESEARCH, vol. 56, no. 7, 29 July 2015 (2015-07-29), pages 1308-1317, XP055363055, US ISSN: 0022-2275, DOI: 10.1194/jlr.M054890 the whole document	1-30
A	----- RADER DANIEL J: "New Therapeutic Approaches to the Treatment of Dyslipidemia", CELL METABOLISM, vol. 23, no. 3, 4 February 2015 (2015-02-04), pages 405-412, XP029457099, ISSN: 1550-4131, DOI: 10.1016/J.CMET.2016.01.005 the whole document	1-30
A	----- Anna Borodovsky ET AL: "Development of Monthly to Quarterly Subcutaneous Administration of RNAi Therapeutics Targeting the Metabolic Disease Genes PCSK9, ApoC3 and ANGPTL3 ALN-PCS Phase I Study Results",  25 November 2014 (2014-11-25), XP055370914, Retrieved from the Internet: URL: <a href="http://www.alnylam.com/web/assets/CardioMetabolic_AHA_Poster_111714.pdf">http://www.alnylam.com/web/assets/CardioMetabolic_AHA_Poster_111714.pdf</a> [retrieved on 2017-05-09] the whole document	1-30
	-----	

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/020221

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012174178 A1	20-12-2012	AR 087329 A1	19-03-2014
		AU 2012271663 A1	16-01-2014
		CA 2838867 A1	20-12-2012
		CN 103732624 A	16-04-2014
		CO 6821892 A2	31-12-2013
		CR 20130621 A	12-02-2014
		DO P2013000299 A	15-04-2014
		EC SP13013085 A	31-01-2014
		EP 2721065 A1	23-04-2014
		GT 201300306 A	09-03-2015
		JP 2014523739 A	18-09-2014
		KR 20140037218 A	26-03-2014
		MA 35190 B1	02-06-2014
		MX 337644 B	14-03-2016
		NZ 619092 A	31-03-2016
		PE 11662014 A1	16-09-2014
		RU 2013155906 A	27-07-2015
		SG 195165 A1	30-12-2013
		TW 201313738 A	01-04-2013
		US 2013171149 A1	04-07-2013
		US 2015197564 A1	16-07-2015
UY 34137 A	03-01-2013		
WO 2012174178 A1	20-12-2012		
ZA 201308879 B	30-07-2014		
WO 2016011256 A1	21-01-2016	AU 2015289613 A1	02-02-2017
		EP 3169353 A1	24-05-2017
		KR 20170029613 A	15-03-2017
		US 2016137745 A1	19-05-2016
		WO 2016011256 A1	21-01-2016
WO 2014194168 A2	04-12-2014	AU 2014274046 A1	19-11-2015
		CA 2913550 A1	04-12-2014
		CN 105473161 A	06-04-2016
		EA 201592302 A1	29-04-2016
		EP 3003361 A2	13-04-2016
		HK 1216841 A1	09-12-2016
		JP 2016520616 A	14-07-2016
		KR 20160013047 A	03-02-2016
		US 2014356371 A1	04-12-2014
		WO 2014194168 A2	04-12-2014