










(86) Date de dépôt PCT/PCT Filing Date: 1996/06/17
 (87) Date publication PCT/PCT Publication Date: 1997/01/23
 (45) Date de délivrance/Issue Date: 2008/01/08
 (85) Entrée phase nationale/National Entry: 1997/12/24
 (86) N° demande PCT/PCT Application No.: EP 1996/002601
 (87) N° publication PCT/PCT Publication No.: 1997/002243
 (30) Priorités/Priorities: 1995/06/30 (DE195 23 850.8);
 1996/01/31 (DE196 03 332.2)

(51) Cl.Int./Int.Cl. *C07D 207/38* (2006.01),
A01N 43/04 (2006.01), *A01N 43/36* (2006.01),
A01N 43/56 (2006.01), *A01N 43/86* (2006.01),
A01N 43/90 (2006.01), *A01N 57/08* (2006.01),
C07C 233/51 (2006.01), *C07C 233/52* (2006.01),
C07C 25/02 (2006.01), *C07C 255/29* (2006.01),
C07C 323/56 (2006.01), ...
 (72) Inventeurs/Inventors:
 FISCHER, REINER, DE;
 BRETSCHNEIDER, THOMAS, DE;
 HAGEMANN, HERMANN, DE;
 LIEB, FOLKER, DE;
 RUTHER, MICHAEL, DE; ...

(54) Titre : CETOENOLS A SUBSTITUTION DIALKYLE-HALOGENURE DE PHENYLE S'UTILISANT COMME
 HERBICIDES ET COMME PESTICIDES

(54) Title: DIALKYL PHENYL HALIDE-SUBSTITUTED KETO-ENOLS FOR USE AS HERBICIDES AND PESTICIDES



(57) Abrégé/Abstract:

The present invention relates to new compounds of the formula (I) (see formula I), in which X represents alkyl, Y represents halogen or alkyl and Z represents halogen or alkyl, with the proviso that one of the radicals Y and Z always represents halogen and


- (51) Cl.Int./Int.Cl. (suite/continued) *C07C 57/58* (2006.01), *C07C 57/76* (2006.01), *C07C 69/65* (2006.01), *C07D 209/54* (2006.01), *C07D 209/96* (2006.01), *C07D 213/36* (2006.01), *C07D 231/36* (2006.01), *C07D 279/06* (2006.01), *C07D 307/60* (2006.01), *C07D 307/94* (2006.01), *C07D 309/08* (2006.01), *C07D 309/32* (2006.01), *C07D 309/40* (2006.01), *C07D 311/56* (2006.01), *C07D 311/94* (2006.01), *C07D 333/32* (2006.01), *C07D 333/78* (2006.01), *C07D 405/04* (2006.01), *C07D 407/04* (2006.01), *C07D 409/04* (2006.01), *C07D 417/04* (2006.01), *C07D 471/04* (2006.01), *C07D 487/04* (2006.01), *C07D 491/052* (2006.01), *C07D 491/107* (2006.01), *C07D 493/04* (2006.01), *C07D 493/10* (2006.01), *C07D 495/10* (2006.01), *C07F 9/547* (2006.01)
- (72) Inventeurs(suite)/Inventors(continued): WIDDIG, ARNO, DE; DAHMEN, PETER, DE; DOLLINGER, MARKUS, DE; ERDELEN, CHRISTOPH, DE; SANTEL, HANS-JOACHIM, DE; WACHENDORFF-NEUMANN, ULRIKE, DE; GRAFF, ALAN, DE; ANDERSCH, WOLFRAM, DE
- (73) Propriétaires(suite)/Owners(continued):BAYER AKTIENGESELLSCHAFT, DE
- (74) Agent: FETHERSTONHAUGH & CO.
- (57) Abrégé(suite)/Abstract(continued):
the other alkyl, Het represents one of the groups (see formula 1), (see formula 2), (see formula 3), (see formula 4), (see formula 5), (see formula 6), in which A, B, D and G have the meanings given in the description, a plurality of processes for their preparation and their use as pesticides and herbicides.

Le A 31 064 - Foreign Countries

- 228 -

Alkyl-dihalogenophenyl-substituted ketoenols**A b s t r a c t**

The present invention relates to new compounds of the formula (I)



5 in which





X represents alkyl,

Y represents halogen or alkyl and

Z represents halogen or alkyl,


with the proviso that one of the radicals Y and Z always represents halogen and the
 10 other alkyl,

Het represents one of the groups




Le A 31 064 - Foreign Countries

- 229 -



(5),



(6),

in which

A, B, D and G have the meanings given in the description,

a plurality of processes for their preparation and their use as pesticides and herbicides.

5

Le A 31 064 - Foreign Countries / Ba/bo/Kr/S-P

- 1 -

~~FILED IN THE OFFICE OF THE
TEXT TRANSLATION~~

Dialkyl-halogenophenyl-substituted ketoenols

The invention relates to new phenyl-substituted cyclic ketoenols, a plurality of processes and intermediates for their preparation and their use as pesticides and herbicides.

5 It has already been disclosed that certain phenyl-substituted cyclic ketoenols are effective as insecticides, acaricides and/or herbicides.

Pharmaceutical properties of 3-acyl-pyrrolidine-2,4-diones have previously been described (S. Suzuki et al. Chem. Pharm. Bull. 15 1120 (1967)). Furthermore, N-phenylpyrrolidine-2,4-diones have been synthesized by R. Schmierer and
10 H. Mildenberger (Liebigs Ann. Chem. 1985, 1095). A biological activity of these compounds has not been described.

EP-A-0 262 399 and GB-A-2 266 888 disclose compounds (3-aryl-pyrrolidine-2,4-diones) of similar structure, of which, however, no herbicidal, insecticidal or acaricidal action has been disclosed. Unsubstituted, bicyclic 3-aryl-pyrrolidine-2,4-dione
15 derivatives (EP-A-355 599 and EP-A-415 211) and substituted monocyclic 3-aryl-pyrrolidine-2,4-dione derivatives (EP-A-377 893 and EP-A-442 077) with herbicidal, insecticidal or acaricidal action are known.

Polycyclic 3-arylpyrrolidine-2,4-dione derivatives (EP-A-442 073) and 1H-arylpyrrolidine-dione derivatives (EP-A-456 063, EP-A-521 334, EP-A-596 298,
20 EP-A-613 884, DE 4 440 594, EP-A-613 885, WO 94/01 997 and WO 95/01 358) are furthermore known.

It is known that certain substituted Δ^3 -dihydrofuran-2-one derivatives have herbicidal properties (cf. DE-A-4 014 420). The synthesis of the tetrone acid derivatives used as starting compounds (e.g. 3-(2-methyl-phenyl)-4-hydroxy-5-(4-fluorophenyl)- Δ^3 -
25 dihydrofuran-2-one) is likewise described in DE-A-4 014 420. Compounds of similar structure without details of an insecticidal and/or acaricidal activity are known from the

Le A 31 064 - Foreign Countries

- 2 -

publication Campbell et al., J. Chem. Soc., Perkin Trans. 1, 1985, (8) 1567-76. Furthermore, 3-aryl- Δ^3 -dihydrofuranone derivatives having herbicidal, acaricidal and insecticidal properties are disclosed in EP-A-528 156, but the action described there is not always adequate. 3-Aryl-4-hydroxy- Δ^3 -dihydrofuranone derivatives are disclosed in
5 EP-A-647 637. Thiotetronic acids are disclosed in WO 95/26345.

The literature further discloses certain 3H-pyrazol-3-one derivatives, for example 1,2-diethyl-1,2-dihydro-5-hydroxy-4-phenyl-3H-pyrazol-3-one or {[5-oxo-1,2-diphenyl-4-(p-sulphophenyl)-3-pyrazolin-3-yl]-oxy} [lacuna] disodium salt or p-(3-hydroxy-5-oxo-1,2-diphenyl-3-pyrazolin-4-yl)-benzenesulphonic acid (cf. J.
10 Heterocycl. Chem., 25(5), 1301-1305, 1988 or J. Heterocycl. Chem., 25(5), 1307-1310, 1988 or Zh. Obshch. Khim., 34(7), 2397-2402, 1964). A biological action of these compounds, however, is not described.

It is furthermore known that the trisodium salt of 4,4',4''-(5-hydroxy-3-oxo-1H-pyrazole-1,2,4(3H)-triy)-tris-benzenesulphonic acid has pharmacological properties
15 (cf. Farmakol. Toksikol. (Moscow), 38(2), 180-186, 1976). Its use in plant protection, however, is not known.

In addition, EP-A-508 126 and WO 92/16 510 describe 4-arylpyrazolidine-3,5-dione derivatives having herbicidal, acaricidal and insecticidal properties.

Certain phenyl-pyrone derivatives unsubstituted in the phenyl ring have already been
20 disclosed (cf. A.M. Chirazi, T. Kappe and E. Ziegler, Arch. Pharm. 309, 558 (1976) and K.-H. Boltze and K. Heidenbluth, Chem. Ber. 91, 2849), a possible utility for these compounds as pesticides not being indicated. Phenyl-pyrone derivatives substituted in the phenyl ring and having herbicidal, acaricidal and insecticidal properties are described in EP-A-588 137.

25 Certain 5-phenyl-1,3-thiazine derivatives unsubstituted in the phenyl ring have already been described (cf. E. Ziegler and E. Steiner, Monatsh. 95, 147 (1964), R. Ketcham, T. Kappe and E. Ziegler, J. Heterocycl. Chem. 10, 223 (1973)), a possible application


Le A 31 064 - Foreign Countries

- 3 -

as pesticides not being indicated for these compounds. 5-Phenyl-1,3-thiazine derivatives substituted in the phenyl ring and having herbicidal, acaricidal and insecticidal action are described in WO 94/14 785.

The activity and spectrum of action of these compounds, however, is not always completely satisfactory, in particular at low application rates and concentrations.

New compounds of the formula (I)



have now been found,

in which


10 X represents alkyl,

Y represents halogen or alkyl and


Z represents halogen or alkyl,

with the proviso that one of the radicals Y and Z always represents halogen and the other alkyl,

15 Het represents one of the groups




(1),




(2),

30517-144


- 4 -




(3),



(4),



(5),



(6),

in which

A represents hydrogen or alkyl, alkenyl, alkoxyalkyl, polyalkoxyalkyl or alkylthioalkyl, each of which is optionally substituted by halogen or cycloalkyl or heterocyclyl, each of which is saturated or unsaturated and optionally substituted, or aryl, arylalkyl or hetaryl, each of which is optionally substituted by halogen, alkyl, halogenoalkyl, alkoxy, halogenoalkoxy, cyano or nitro,

B represents hydrogen, alkyl or alkoxyalkyl, or

A and B, together with the carbon atom to which they are bonded, represent a saturated or unsaturated, optionally substituted carbocycle or heterocycle,


D represents hydrogen or an optionally substituted radical of the series alkyl, alkenyl, alkynyl, alkoxyalkyl, polyalkoxyalkyl, alkylthioalkyl, saturated or unsaturated cycloalkyl, saturated or unsaturated heterocyclyl, arylalkyl, aryl, hetarylalkyl or hetaryl, or

A and D, together with the atoms to which they are bonded, represent a saturated or unsaturated and optionally substituted carbocycle or heterocycle,

Le A 31 064 - Foreign Countries

- 5 -

G, in the case in which Het represents one of the radicals (1), (2), (3), (5) or (6), represents hydrogen (a) or, in the case in which Het represents one of the radicals (1), (2), (3), (4), (5) or (6), represents one of the groups



in which

E represents a metal ion equivalent or an ammonium ion,

L represents oxygen or sulphur,

M represents oxygen or sulphur,

10 R¹ represents alkyl, alkenyl, alkoxyalkyl, alkylthioalkyl or polyalkoxyalkyl, each of which is optionally substituted by halogen, or cycloalkyl or heterocyclyl, each of which is optionally substituted by halogen, alkyl or alkoxy, or phenyl, phenylalkyl, hetaryl, phenoxyalkyl or hetaryloxyalkyl, each of which is optionally substituted,

15 R² represents alkyl, alkenyl, alkoxyalkyl or polyalkoxyalkyl, each of which is optionally substituted by halogen, or cycloalkyl, phenyl or benzyl, each of which is optionally substituted,

20 R³, R⁴ and R⁵ independently of one another represent alkyl, alkoxy, alkylamino, dialkylamino, alkylthio, alkenylthio or cycloalkylthio, each of which is optionally substituted by halogen, or phenyl, phenoxy or phenylthio, each of which is optionally substituted,







30517-144

- 6 -

5 R^6 and R^7 independently of one another represent hydrogen, or alkyl, cycloalkyl, alkenyl, alkoxy, alkoxyalkyl, each of which is optionally substituted by halogen, or phenyl or benzyl, each of which is optionally substituted, or, together with the N atom to which they are bonded, form an optionally substituted cyclic system optionally containing oxygen or sulphur.

10 The compounds of the formula (I) can also be present, depending on the nature of the substituents, as geometric and/or optical isomers or isomer mixtures of differing composition which, if appropriate, can be separated in a customary manner. Both the pure isomers and the isomer mixtures, their preparation and use, and compositions containing these are a subject of the present invention. In the following, for the sake of simplicity, however, compounds of the formula (I) are always referred to, although both the pure compounds and, if appropriate, mixtures of different proportions of isomeric compounds are intended.

15 Including the meanings (1) to (6) of the group Het, the following principle structures (I-1) to (I-6) result:



Le A 31 064 - Foreign Countries


- 7 -

in which


A, B, D, G, X, Y and Z have the meanings given above.

Including the various meanings (a), (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-1-a) to (I-1-g) result if Het represents the group (1)


5 (I-1-a):




(I-1-b):




(I-1-c):




(I-1-d):



(I-1-e):




(I-1-f):



Le A 31 064 - Foreign Countries



- 8 -

(I-1-g):



in which

5 A, B, D, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.

Including the various meanings (a), (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-2-a) to (I-2-g) result if Het is the group (2)




(I-2-a):**(I-2-b):**

10

(I-2-c):**(I-2-d):**

Le A 31 064 - Foreign Countries



- 9 -

(I-2-e):**(I-2-f):****(I-2-g):**

5 in which

A, B, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.





Including the various meanings (a), (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-3-a) to (I-3-g) result if Het represents the group (3)

(I-3-a):**(I-3-b):**


10

Le A 31 064 - Foreign Countries

- 10 -

(I-3-c):**(I-3-d):****(I-3-e):****(I-3-f):**

5

(I-3-g):


in which

A, B, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.

Depending on the position of the substituent G, the compounds of the formula (I-4) can
 10 be present in the two isomeric formulae (I-4)_a and (I-4)_b

Le A 31 064 - Foreign Countries

- 11 -





which is intended to be expressed by the dashed line in the formula (I-4):

The compounds of the formulae (I-4)_a and (I-4)_b can be present both as mixtures and in the form of their pure isomers. Mixtures of the compounds of the formula (I-4)_a and (I-4)_b can optionally be separated by physical methods in a manner known per se, for example by chromatographic methods.

For reasons of better clarity, in the following in each case only one of the possible isomers is shown. This does not exclude the fact that the compounds can optionally be present in the form of the isomer mixtures or in the other isomeric form in each case.


10 Including the various meanings (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-4-b) to (I-4-g) result if Het represents the group (4)

(I-4-b):**(I-4-c):**


Le A 31 064 - Foreign Countries

- 12 -


(I-4-d):




(I-4-e)



(I-4-f):





(I-4-g):



5 in which

A, D, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.

Depending on the position of the substituent G, the compounds of the formula (I-5) can be present in the two isomeric forms of the formulae (I-5)_a and (I-5)_b

(I-5)_a(I-5)_b

10

Le A 31 064 - Foreign Countries





- 13 -

which is intended to be expressed by the dashed line in the formula (I-5).

The compounds of the formulae (I-5)_a and (I-5)_b can be present both as mixtures and in the form of their pure isomers. Mixtures of the compounds of the formulae (I-5)_a and (I-5)_b can optionally be separated by physical methods in a manner known per se, for example by chromatographic methods.




For reasons of better clarity, in the following in each case only one of the possible isomers is shown. This does not exclude the fact that the compounds can optionally be present in the form of the isomer mixtures or in the other isomeric form in each case.

Including the various meanings (a), (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-5-a) to (I-5-g) result if Het represents the group (5)

(I-5-a):**(I-5-b):****(I-5-c):****(I-5-d):**

Le A 31 064 - Foreign Countries



- 14 -

(I-5-e):**(I-5-f):****(I-5-g):**

5 in which

A, D, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.





Including the various meanings (a), (b), (c), (d), (e), (f) and (g) of the group G, the following principle structures (I-6-a) to (I-6-g) result if Het represents the group (6)

(I-6-a):**(I-6-b):**


10

Le A 31 064 - Foreign Countries

- 15 -

(I-6-c):**(I-6-d):****(I-6-e):****(I-6-f):**

5

(I-6-g):

in which


A, E, L, M, X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ have the meanings given above.

It has furthermore been found that the new compounds of the formula (I) are obtained
10 by one of the processes described in the following:

Le A 31 064 - Foreign Countries

- 16 -

(A) Compounds of the formula (I-1-a)




in which

A, B, D, X, Y and Z have the meanings given above,

5 are obtained when

compounds of the formula (II)



in which

A, B, D, X, Y and Z have the meanings given above,

10 and


 R^8 represents alkyl (preferably C_1 - C_6 -alkyl),

are intramolecularly condensed in the presence of a diluent and in the presence of a base.

Le A 31 064 - Foreign Countries

- 17 -

(B) It has additionally been found that compounds of the formula (I-2-a)




in which

A, B, X, Y and Z have the meanings given above,

5 are obtained when

compounds of the formula (III)




in which

A, B, X, Y, Z and R⁸ have the meanings given above,

10 are intramolecularly condensed in the presence of a diluent and in the presence of a base.

(C) It has furthermore been found that compounds of the formula (I-3-a)



in which


Le A 31 064 - Foreign Countries

- 18 -

A, B, X, Y and Z have the meanings given above,

are obtained when

compounds of the formula (IV)




5 in which

A, B, X, Y, Z and R⁸ have the meanings given above and

W represents hydrogen, halogen, alkyl (preferably C₁-C₆-alkyl) or alkoxy (preferably C₁-C₈-alkoxy),

10 are cyclized, if appropriate in the presence of a diluent and in the presence of an acid.

(E) It has furthermore been found that the compounds of the formula (I-5-a)



in which

A, D, X, Y and Z have the meanings given above,

Le A 31 064 - Foreign Countries

- 19 -

are obtained when


compounds of the formula (VIII)



in which

5 A and D have the meanings given above,


or their silyl enol ethers of the formula (VIIIa)



in which

10 A and D have the meanings mentioned above and $\text{R}^{\delta'}$ represents alkyl
(preferably methyl),

are reacted with compounds of the formula (V)



in which

X, Y and Z have the meanings given above and

Le A 31 064 - Foreign Countries


- 20 -

Hal represents halogen (preferably chlorine or bromine),

if appropriate in the presence of a diluent and if appropriate in the presence of an acid acceptor.

(F) It has furthermore been found that the compounds of the formula (I-6-a)

5



in which

A, X, Y and Z have the meanings given above,

are obtained when compounds of the formula (IX)




10

in which

A has the meaning given above,

are reacted with compounds of the formula (V)



in which

Le A 31 064 - Foreign Countries


- 21 -

Hal, X, Y and Z have the meanings given above,

if appropriate in the presence of a diluent and if appropriate in the presence of an acid acceptor.

It has additionally been found

- 5 (G) that the compounds of the formulae (I-1-b) to (I-3-b), (I-5-b) and (I-6-b) shown above, in which A, B, D, R¹, X, Y and Z have the meanings given above, are obtained when compounds of the formulae (I-1-a) to (I-3-a), (I-5-a) and (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, and that compounds of the formula (I-4-b) shown above, in which A, D, R¹, X, Y and Z have the meanings given above, are obtained when compounds of the
- 10 formula (I-4-a)



in which

A, D, X, Y and Z have the meaning given above, in each case are reacted

- 15 α) with acid halides of the formula (X)



in which

R¹ has the meaning given above and

Le A 31 064 - Foreign Countries

- 22 -

Hal represents halogen (in particular chlorine or bromine)

or

β) with carboxylic anhydrides of the formula (XI)




5 in which

R¹ has the meaning given above,

if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent;

10 (H) that the compounds of the formulae (I-1-c) to (I-6-c) shown above, in which A, B, D, R², M, X, Y and Z have the meanings given above and L represents oxygen, are obtained when compounds of the formulae (I-1-a) to (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

15 with chloroformic acid esters or chloroformic acid thioesters of the formula (XII)



in which

R² and M have the meanings given above,

20 if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent;


Le A 31 064 - Foreign Countries

- 23 -

(I) that compounds of the formulae (I-1-c) to (I-6-c) shown above, in which A, B, D, R², M, X, Y and Z have the meanings given above and L represents sulphur, are obtained when compounds of the formulae (I-1-a) to (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

5

α) with chloromonothioformic acid esters or chlorodithioformic acid esters of the formula (XIII)



in which

10 M and R² have the meanings given above,

if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent

or

β) with carbon disulphide and then with compounds of the formula (XIV)

15 $\text{R}^2\text{-Hal}$ (XIV)

in which

R² has the meaning given above and

Hal represents chlorine, bromine or iodine,

20

if appropriate in the presence of a diluent and if appropriate in the presence of a base,

Le A 31 064 - Foreign Countries

- 24 -

(J) that compounds of the formulae (I-1-d) to (I-6-d) shown above, in which A, B, D, R³, X, Y and Z have the meanings given above, are obtained when compounds of the formulae (I-1-a) to (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

5 with sulphonyl chlorides of the formula (XV)



in which

R³ has the meaning given above,

10 if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent,

(K) that compounds of the formulae (I-1-e) to (I-6-e) shown above, in which A, B, D, L, R⁴, R⁵, X, Y and Z have the meanings given above, are obtained when compounds of the formulae (I-1-a) to (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

15 with phosphorus compounds of the formula (XVI)



in which

L, R⁴ and R⁵ have the meanings given above and

Hal represents halogen (in particular chlorine or bromine),


Le A 31 064 - Foreign Countries

- 25 -

if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent,

- (L) that compounds of the formulae (I-1-f) to (I-6-f) shown above, in which A, B, D, E, X, Y and Z have the meanings given above, are obtained when
5 compounds of the formulae (I-1-a) to (I-6-a), in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

with metal compounds or amines of the formulae (XVII) or (XVIII)



in which

- 10 Me represents a mono- or divalent metal (preferably an alkali metal or alkaline earth metal such as lithium, sodium, potassium, magnesium or calcium),

t represents the number 1 or 2 and

- 15 R^{10} , R^{11} , R^{12} independently of one another represent hydrogen or alkyl (preferably C_1 - C_8 -alkyl),

if appropriate in the presence of a diluent,

- (M) that compounds of the formulae (I-1-g) to (I-6-g) shown above, in which A, B, D, L, R^6 , R^7 , X, Y and Z have the meanings given above, are obtained when
20 compounds of the formulae (I-1-a) to (I-6-a) shown above, in which A, B, D, X, Y and Z have the meanings given above, in each case are reacted

- α) with isocyanates or isothiocyanates of the formula (XIX)

Le A 31 064 - Foreign Countries

- 26 -




in which

R^6 and L have the meanings given above,

5 if appropriate in the presence of a diluent and if appropriate in the presence of a catalyst or

β) with carbamoyl chlorides or thiocarbamoyl chlorides of the formula (XX)



in which

L, R^6 and R^7 have the meanings given above,

10 if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent.

It has furthermore been found that the new compounds of the formula (I) have a very good activity as pesticides, preferably as insecticides, acaricides and herbicides.

15 Formula (I) provides a general definition of the compounds according to the invention. Preferred substituents or ranges of the radicals shown in the formulae mentioned above and below are illustrated in the following:

X preferably represents C_1 - C_6 -alkyl.

Y preferably represents halogen or C_1 - C_6 -alkyl.

30517-144


- 27 -

Z preferably represents halogen or C₁-C₆-alkyl.


In this case, one of the substituents Y and Z always represents halogen and the other alkyl.

Het preferably represents one of the groups


5




(1),




(2),




(3),



(4),



(5),



(6),

10

A preferably represents hydrogen, or C₁-C₁₂-alkyl, C₂-C₈-alkenyl, C₁-C₁₀-alkoxy-C₁-C₈-alkyl, poly-C₁-C₈-alkoxy-C₁-C₈-alkyl or C₁-C₁₀-alkylthio-C₁-C₆-alkyl, each of which is optionally substituted by halogen, or C₃-C₈-cycloalkyl which is optionally substituted by halogen, C₁-C₆-alkyl or C₁-C₆-alkoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur, or phenyl, naphthyl, phenyl-C₁-C₆-alkyl, naphthyl-C₁-C₆-alkyl or hetaryl having 5 or 6 ring atoms and one to three heteroatoms from the series oxygen, sulphur and nitrogen, each of which is optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkoxy, cyano or nitro,

15

B preferably represents hydrogen, C₁-C₁₂-alkyl or C₁-C₈-alkoxy-C₁-C₆-alkyl, or

Le A 31 064 - Foreign Countries

- 28 -





- 5 A, B and the carbon atom to which they are bonded preferably represent C₃-C₁₀-cycloalkyl or C₅-C₁₀-cycloalkenyl, in which one methylene group is optionally replaced by oxygen or sulphur and each of which is optionally substituted by C₁-C₈-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₈-halogenoalkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio, halogen or phenyl or
- 10 A, B and the carbon atom to which they are bonded preferably represent C₅-C₆-cycloalkyl which is substituted by an alkylenediyl group optionally containing one or two oxygen and/or sulphur atoms, or by an alkylenedioxy group or by an alkylenedithiyl group which, with the carbon atom to which it is bonded, forms a further five- to eight-membered ring, or
- 15 A, B and the carbon atom to which they are bonded preferably represent C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which two substituents, together with the carbon atoms to which they are bonded, represent C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkanedienediyl, each of which is optionally substituted by C₁-C₆-alkyl, C₁-C₆-alkoxy or halogen and in which one methylene group is optionally replaced by oxygen or sulphur in each case.
- 20 D preferably represents hydrogen, or C₁-C₁₂-alkyl, C₃-C₈-alkenyl, C₃-C₈-alkinyl, C₁-C₁₀-alkoxy-C₂-C₈-alkyl, poly-C₁-C₈-alkoxy-C₂-C₈-alkyl or C₁-C₁₀-alkylthio-C₂-C₈-alkyl, each of which is optionally substituted by halogen, or C₃-C₈-cycloalkyl which is optionally substituted by halogen, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-halogenoalkyl and in which one or two non-directly adjacent methylene groups are optionally replaced by oxygen and/or sulphur, or
- 25 phenyl, hetaryl having 5 to 6 ring atoms and one or two heteroatoms from the series oxygen, sulphur and nitrogen, phenyl-C₁-C₆-alkyl or hetaryl-C₁-C₆-alkyl having 5 to 6 ring atoms and one or two heteroatoms from the series oxygen, sulphur and nitrogen, each of which is optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkoxy, cyano or nitro, or

Le A 31 064 - Foreign Countries

- 29 -

A and D together preferably represent a C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkadienediyl group, in which one methylene group in each case is optionally replaced by oxygen or sulphur and each of which is optionally substituted by halogen, hydroxyl, mercapto or C₁-C₁₀-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₃-C₇-cycloalkyl, phenyl or benzyloxy, each of which is optionally substituted by halogen, or by a further C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkadienediyl group forming a fused ring and in which one methylene group in each case is optionally replaced by oxygen or sulphur and which is optionally substituted by C₁-C₆-alkyl or in which two adjacent substituents, together with the carbon atoms to which they are bonded, optionally form a further saturated or unsaturated carbocycle having 5 or 6 ring atoms, or

A and D together represent a C₃-C₆-alkanediyl or C₃-C₆-alkenediyl group which in each case optionally contains one of the following groups




G preferably represents, in the case in which Het represents one of the radicals

Le A 31 064 - Foreign Countries

- 30 -

(1), (2), (3), (5) or (6), hydrogen (a) or, in the case in which Het represents one of the radicals (1), (2), (3), (4), (5) or (6), one of the groups



in which

5 E represents a metal ion equivalent or an ammonium ion,

L represents oxygen or sulphur and

M represents oxygen or sulphur.

10 R¹ preferably represents C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, C₁-C₈-alkylthio-C₁-C₈-alkyl or poly-C₁-C₈-alkoxy-C₁-C₈-alkyl, each of which is optionally substituted by halogen, or C₃-C₈-cycloalkyl which is optionally substituted by halogen, C₁-C₆-alkyl or C₁-C₆-alkoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur,

15 phenyl which is optionally substituted by halogen, cyano, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl, C₁-C₆-halogenoalkoxy, C₁-C₆-alkylthio or C₁-C₆-alkylsulphonyl,

phenyl-C₁-C₆-alkyl which is optionally substituted by halogen, nitro, cyano, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl or C₁-C₆-halogenoalkoxy,

20 5- or 6-membered hetaryl having one or two heteroatoms from the series oxygen, sulphur and nitrogen and which is optionally substituted by halogen or C₁-C₆-alkyl,

Le A 31 064 - Foreign Countries

- 31 -

phenoxy-C₁-C₆-alkyl which is optionally substituted by halogen or C₁-C₆-alkyl,
or

5 5- or 6-membered hetaryloxy-C₁-C₆-alkyl having one or two heteroatoms from the series oxygen, sulphur and nitrogen and which is optionally substituted by halogen, amino or C₁-C₆-alkyl.

R² preferably represents C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₂-C₈-alkyl or poly-C₁-C₈-alkoxy-C₂-C₈-alkyl, each of which is optionally substituted by halogen,

10 C₃-C₈-cycloalkyl which is optionally substituted by halogen, C₁-C₆-alkyl or C₁-C₆-alkoxy, or

phenyl or benzyl, each of which is optionally substituted by halogen, cyano, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl or C₁-C₆-halogenoalkoxy.

15 R³ preferably represents C₁-C₈-alkyl which is optionally substituted by halogen, or phenyl or benzyl, each of which is optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy, cyano or nitro.

20 R⁴ and R⁵ independently of one another preferably represent C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylamino, di-(C₁-C₈-alkyl)amino, C₁-C₈-alkylthio or C₂-C₈-alkenylthio, each of which is optionally substituted by halogen, or phenyl, phenoxy or phenylthio, each of which is optionally substituted by halogen, nitro, cyano, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkylthio, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl.

25 R⁶ and R⁷ independently of one another preferably represent hydrogen, or C₁-C₈-alkyl, C₃-C₈-cycloalkyl, C₁-C₈-alkoxy, C₃-C₈-alkenyl or C₁-C₈-alkoxy-C₂-C₈-alkyl, each of which is optionally substituted by halogen, or phenyl or benzyl, each of

Le A 31 064 - Foreign Countries

- 32 -

which is optionally substituted by halogen, C₁-C₈-alkyl, C₁-C₈-halogenoalkyl or C₁-C₈-alkoxy or together represent a C₃-C₆-alkylene radical which is optionally substituted by C₁-C₆-alkyl and in which a methylene group is optionally replaced by oxygen or sulphur.

5 R¹³ preferably represents hydrogen or C₁-C₈-alkyl or C₁-C₈-alkoxy, each of which is optionally substituted by halogen, or C₃-C₈-cycloalkyl which is optionally substituted by halogen, C₁-C₄-alkyl or C₁-C₄-alkoxy and in which a methylene group is optionally replaced by oxygen or sulphur, or phenyl, phenyl-C₁-C₄-alkyl or phenyl-C₁-C₄-alkoxy, each of which is optionally substituted by
10 halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy, nitro or cyano.

R¹⁴ preferably represents hydrogen or C₁-C₈-alkyl, or

R¹³ and R¹⁴ together preferably represent C₄-C₆-alkanediyl.

R¹⁵ and R¹⁶ are identical or different and preferably represent C₁-C₆-alkyl, or

15 R¹⁵ and R¹⁶ together preferably represent a C₂-C₄-alkanediyl radical which is optionally substituted by C₁-C₆-alkyl, C₁-C₆-halogenoalkyl or by phenyl which is optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₄-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkoxy, nitro or cyano.

20 R¹⁷ and R¹⁸ independently of one another preferably represent hydrogen, or C₁-C₈-alkyl which is optionally substituted by halogen or phenyl which is optionally substituted by halogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-halogenoalkoxy, nitro or cyano, or

25 R¹⁷ and R¹⁸, together with the carbon atom to which they are bonded, represent a carbonyl group, or C₅-C₇-cycloalkyl which is optionally substituted by C₁-C₄-alkyl or C₁-C₄-alkoxy and in which a methylene group is optionally

30517-144

- 33 -

replaced by oxygen or sulphur.

R^{19} and R^{20} independently of one another preferably represent C_1 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, C_1 - C_{10} -alkoxy, C_1 - C_{10} -alkylamino, C_3 - C_{10} -alkenylamino, di- $(C_1$ - C_{10} -alkyl)amino or di- $(C_3$ - C_{10} -alkenyl)amino.

5 X particularly preferably represents C_1 - C_4 -alkyl.







Y particularly preferably represents fluorine, chlorine, bromine or C_1 - C_4 -alkyl.

Z particularly preferably represents fluorine, chlorine, bromine or C_1 - C_4 -alkyl.

In this case one of the radicals Y and Z always represents halogen and the other alkyl.

Het particularly preferably represents one of the groups

10



Le A 31 064 - Foreign Countries

- 34 -

- A particularly preferably represents hydrogen, or C₁-C₁₀-alkyl, C₂-C₆-alkenyl, C₁-C₈-alkoxy-C₁-C₆-alkyl, poly-C₁-C₆-alkoxy-C₁-C₆-alkyl or C₁-C₈-alkylthio-C₁-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine, or C₃-C₇-cycloalkyl which is optionally substituted by fluorine, chlorine, C₁-C₄-alkyl or C₁-C₄-alkoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur, or phenyl, furanyl, pyridyl, imidazolyl, triazolyl, pyrazolyl, indolyl, thiazolyl, thienyl or phenyl-C₁-C₄-alkyl, each of which is optionally substituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, cyano or nitro.
- B particularly preferably represents hydrogen, C₁-C₁₀-alkyl or C₁-C₆-alkoxy-C₁-C₄-alkyl or
- A, B and the carbon atom to which they are bonded particularly preferably represent C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which in each case a methylene group is optionally replaced by oxygen or sulphur and each of which is optionally substituted by C₁-C₆-alkyl, C₃-C₈-cycloalkyl, C₁-C₃-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, fluorine, chlorine or phenyl, or
- A, B and the carbon atom to which they are bonded particularly preferably represent C₅-C₆-cycloalkyl which is substituted by an alkylenediyl group optionally containing one or two oxygen or sulphur atoms or by an alkylenedioxy group or by an alkylenedithiyl group and which, with the carbon atom to which it is bonded, forms a further 5- to 7-membered ring, or
- A, B and the carbon atom to which they are bonded particularly preferably represent C₃-C₆-cycloalkyl or C₅-C₆-cycloalkenyl, in which two substituents, together with the carbon atoms to which they are bonded, represent C₃-C₅-alkanediyl, C₃-C₅-alkenediyl or butadienediyl, each of which is optionally substituted by C₁-C₅-alkyl, C₁-C₅-alkoxy, fluorine, chlorine or bromine, and in which in each case a methylene group is optionally replaced by oxygen or sulphur.




Le A 31 064 - Foreign Countries

- 35 -

D particularly preferably represents hydrogen, or C₁-C₁₀-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkinyl, C₁-C₈-alkoxy-C₂-C₆-alkyl, poly-C₁-C₆-alkoxy-C₂-C₆-alkyl or C₁-C₈-alkylthio-C₂-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine, C₃-C₇-cycloalkyl which is optionally substituted by fluorine, chlorine, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₂-halogenoalkyl and in which one or two non-directly adjacent methylene groups are optionally replaced by oxygen and/or sulphur, or phenyl, furanyl, imidazolyl, pyridyl, thiazolyl, pyrazolyl, pyrimidyl, pyrrolyl, thienyl, triazolyl or phenyl-C₁-C₄-alkyl, each of which is optionally substituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, cyano or nitro, or


A and D together particularly preferably represent a C₃-C₅-alkanediyl or C₃-C₅-alkenediyl group, in which in each case a carbon atom is optionally replaced by oxygen or sulphur and each of which is optionally substituted by fluorine, chlorine, hydroxyl, mercapto or by C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, phenyl or benzoyloxy, each of which is optionally substituted by fluorine or chlorine, or

which in each case optionally contains one of the following groups:




Le A 31 064 - Foreign Countries

- 36 -




or A and D represent (in the case of the compounds of the formula (I-1)), together with the atoms to which they are bonded, one of the groups AD-1 to AD-27


5




AD-1




AD-2




AD-3




AD-4




AD-5




AD-6



AD-7




AD-8




AD-9


10



AD-10




AD-11




AD-12

Le A 31 064 - Foreign Countries


- 37 -




AD-13




AD-14




AD-15



AD-16




AD-17




AD-18


5




AD-19




AD-20




AD-21




AD-22




AD-23




AD-24



AD-25



AD-26




AD-27

10

G particularly preferably represents, in the case in which Het represents one of the radicals (1), (2), (3), (5) or (6), hydrogen (a) or, in the case in which Het represents one of the radicals (1), (2), (3), (4), (5) or (6), one of the groups

Le A 31 064 - Foreign Countries

- 38 -



in which

E represents a metal ion equivalent or an ammonium ion,

L represents oxygen or sulphur and

5 M represents oxygen or sulphur.

R¹ particularly preferably represents C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl or poly-C₁-C₆-alkoxy-C₁-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine, or C₃-C₇-cycloalkyl which is optionally substituted by fluorine, chlorine, C₁-C₅-alkyl or C₁-C₅-alkoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur,

10 phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₃-halogenoalkyl, C₁-C₃-halogenoalkoxy, C₁-C₄-alkylthio or C₁-C₄-alkylsulphonyl,

15 phenyl-C₁-C₄-alkyl which is optionally substituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₃-halogenoalkyl or C₁-C₃-halogenoalkoxy,

pyrazolyl, thiazolyl, pyridyl, pyrimidyl, furanyl or thienyl, each of which is optionally substituted by fluorine, chlorine, bromine or C₁-C₄-alkyl,

Le A 31 064 - Foreign Countries

- 39 -

phenoxy-C₁-C₅-alkyl which is optionally substituted by fluorine, chlorine, bromine or C₁-C₄-alkyl, or

5 pyridyloxy-C₁-C₅-alkyl, pyrimidyloxy-C₁-C₅-alkyl or thiazolyloxy-C₁-C₅-alkyl, each of which is optionally substituted by fluorine, chlorine, bromine, amino or C₁-C₄-alkyl.

R² particularly preferably represents C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₆-alkoxy-C₂-C₆-alkyl or poly-C₁-C₆-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine,

10 C₃-C₇-cycloalkyl which is optionally substituted by fluorine, chlorine, C₁-C₄-alkyl or C₁-C₄-alkoxy, or

phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl, C₁-C₃-alkoxy, C₁-C₃-halogenoalkyl or C₁-C₃-halogenoalkoxy.

15 R³ particularly preferably represents C₁-C₆-alkyl which is optionally substituted by fluorine or chlorine, or phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, C₁-C₅-alkyl, C₁-C₅-alkoxy, C₁-C₃-halogenoalkyl, C₁-C₃-halogenoalkoxy, cyano or nitro.

20 R⁴ and R⁵ independently of one another particularly preferably represent C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylamino, di-(C₁-C₆-alkyl)amino, C₁-C₆-alkylthio or C₃-C₄-alkenylthio, each of which is optionally substituted by fluorine or chlorine, or phenyl, phenoxy or phenylthio, each of which is optionally substituted by fluorine, chlorine, bromine, nitro, cyano, C₁-C₃-alkoxy, C₁-C₃-halogenoalkoxy, C₁-C₃-alkylthio, C₁-C₃-halogenoalkylthio, C₁-C₃-alkyl or C₁-C₃-halogenoalkyl.

25 R⁶ and R⁷ independently of one another particularly preferably represent hydrogen, or

Le A 31 064 - Foreign Countries

- 40 -

5 C₁-C₆-alkyl, C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₃-C₆-alkenyl or C₁-C₆-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine, or phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, C₁-C₅-halogenoalkyl, C₁-C₅-alkyl or C₁-C₅-alkoxy, or together represent a C₃-C₆-alkylene radical which is optionally substituted by C₁-C₄-alkyl and in which a methylene group is optionally replaced by oxygen or sulphur.

10 R¹³ particularly preferably represents hydrogen or C₁-C₆-alkyl or C₁-C₆-alkoxy, each of which is optionally substituted by fluorine or chlorine, or C₃-C₇-cycloalkyl which is optionally substituted by fluorine, C₁-C₂-alkyl or C₁-C₂-alkoxy and in which a methylene group is optionally replaced by oxygen or sulphur, or phenyl, phenyl-C₁-C₃-alkyl or phenyl-C₁-C₂-alkyloxy, each of which is optionally substituted by fluorine, chlorine, bromine, C₁-C₅-alkyl, C₁-C₅-alkoxy, C₁-C₂-halogenoalkyl, C₁-C₂-halogenoalkoxy, nitro or cyano.

R¹⁴ particularly preferably represents hydrogen or C₁-C₆-alkyl, or

15 R¹³ and R¹⁴ together particularly preferably represent C₄-C₆-alkanediyl.

R¹⁵ and R¹⁶ are identical or different and particularly preferably represent C₁-C₄-alkyl, or

20 R¹⁵ and R¹⁶ together particularly preferably represent a C₂-C₃-alkanediyl radical which is optionally substituted by C₁-C₄-alkyl, C₁-C₄-halogenoalkyl or by phenyl which is optionally substituted by fluorine, chlorine, bromine, C₁-C₄-alkyl, C₁-C₂-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₂-halogenoalkoxy, nitro or cyano.

25 R¹⁷ and R¹⁸ independently of one another particularly preferably represent hydrogen, or C₁-C₈-alkyl which is optionally substituted by fluorine or chlorine, or phenyl which is optionally substituted by fluorine, chlorine, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₂-halogenoalkyl, C₁-C₂-halogenoalkoxy, nitro or cyano, or

30517-144

- 41 -

R¹⁷ and R¹⁸, together with the carbon atom to which they are bonded, particularly preferably represent C₅-C₆-cycloalkyl which is optionally substituted by C₁-C₃-alkyl or C₁-C₃-alkoxy and in which a methylene group is optionally replaced by oxygen or sulphur.

5 R¹⁹ and R²⁰ independently of one another particularly preferably represent C₁-C₆-alkyl, C₂-C₆-alkenyl, C₁-C₆-alkoxy, C₁-C₆-alkylamino, C₃-C₆-alkenylamino, di-(C₁-C₆-alkyl)amino or di-(C₃-C₆-alkenyl)amino.


X very particularly preferably represents methyl, ethyl, n-propyl or iso-propyl.

10 Y very particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, n-propyl or iso-propyl.


Z very particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, n-propyl or iso-propyl.

In this case one of the radicals Y and Z always represents halogen and the other alkyl.


Het very particularly preferably represents one of the groups




(1),



(2),



(3),




(4),


15

30517-144

- 42 -



(5),



(6).

A very particularly preferably represents hydrogen, or C₁-C₈-alkyl, C₂-C₄-alkenyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, poly-C₁-C₄-alkoxy-C₁-C₄-alkyl or C₁-C₆-alkylthio-C₁-C₄-alkyl, each of which is optionally substituted by fluorine or chlorine, or C₃-C₆-cycloalkyl which is optionally substituted by fluorine, chlorine, methyl or methoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur, or phenyl, furanyl, thienyl, pyridyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, n-propyl, iso-propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro.

B very particularly preferably represents hydrogen, C₁-C₈-alkyl or C₁-C₄-alkoxy-C₁-C₂-alkyl, or

A, B and the carbon atom to which they are bonded very particularly preferably represent C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which in each case a methylene group is optionally replaced by oxygen or sulphur and each of which is optionally substituted by methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl, tert-butyl, cyclohexyl, trifluoromethyl, methoxy, ethoxy, n-propoxy, iso-propoxy, butoxy, iso-butoxy, sec-butoxy, tert-butoxy, methylthio, ethylthio, fluorine, chlorine or phenyl, or

20 A, B and the carbon atom to which they are bonded very particularly preferably represent C₅-C₆-cycloalkyl which is substituted by an alkylenediyl group optionally containing an oxygen or sulphur atom or by an alkylenedioxyl group which, with the carbon atom to which it is bonded, forms a further five- or six-membered ring, or

Le A 31 064 - Foreign Countries

- 43 -



A, B and the carbon atom to which they are bonded very particularly preferably represent C₃-C₆-cycloalkyl or C₅-C₆-cycloalkenyl, in which two substituents, together with the carbon atoms to which they are bonded, represent C₃-C₄-alkanediyl, C₃-C₄-alkenediyl or butadienediyl, in which in each case a
5 methylene group is optionally replaced by oxygen or sulphur.

D very particularly preferably represents hydrogen, or C₁-C₈-alkyl, C₃-C₄-alkenyl, C₃-C₄-alkinyl, C₁-C₆-alkoxy-C₂-C₄-alkyl, poly-C₁-C₄-alkoxy-C₂-C₄-alkyl, C₁-C₄-alkylthio-C₂-C₄-alkyl or C₃-C₆-cycloalkyl, each of which is optionally substituted by fluorine or chlorine, and in which one or two non-directly
10 adjacent methylene groups are optionally replaced by oxygen and/or sulphur or phenyl, furanyl, pyridyl, thienyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, n-propyl, iso-propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro,

or

15 A and D together very particularly preferably represent a C₃-C₅-alkanediyl or C₃-C₅-alkenediyl group, in which in each case a methylene group is optionally replaced by oxygen or sulphur and each of which is optionally substituted by fluorine, chlorine, hydroxyl, mercapto or by C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, phenyl or benzyloxy, each of which is
20 optionally substituted by fluorine or chlorine, or


which optionally contains one of the following groups




Le A 31 064 - Foreign Countries

- 44 -


or A and D represent, in the case of the compounds of the formula (I-1), together with the atoms to which they are bonded, one of the following groups:



AD-1




AD-2




AD-4


5




AD-6




AD-8




AD-10




AD-12



AD-14




AD-15




10

AD-17



AD-18




AD-27

G very particularly preferably represents, in the case in which Het represents one of the radicals (1), (2), (3), (5) or (6), hydrogen (a) or, in the case in which Het represents one of the radicals (1), (2), (3), (4), (5) or (6), one of the groups

Le A 31 064 - Foreign Countries

- 45 -



in which

E represents a metal ion equivalent or an ammonium ion,

L represents oxygen or sulphur and

5 M represents oxygen or sulphur.

R¹ very particularly preferably represents C₁-C₁₄-alkyl, C₂-C₁₄-alkenyl, C₁-C₄-alkoxy-C₁-C₆-alkyl, C₁-C₄-alkylthio-C₁-C₆-alkyl, poly-C₁-C₄-alkoxy-C₁-C₄-alkyl, each of which is optionally substituted by fluorine or chlorine, or C₃-C₆-cycloalkyl which is optionally substituted by fluorine, chlorine, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, tert-butyl, methoxy, ethoxy, n-propoxy or iso-propoxy and in which one or two methylene groups are optionally replaced by oxygen and/or sulphur,

10 phenyl which is optionally substituted by fluorine, chlorine, bromine, cyano, nitro, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, methylthio, ethylthio, methylsulphonyl or ethylsulphonyl,

15 benzyl which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy,

furanyl, thienyl or pyridyl, each of which is optionally substituted by fluorine, chlorine, bromine, methyl or ethyl,

Le A 31 064 - Foreign Countries

- 46 -

phenoxy-C₁-C₄-alkyl which is optionally substituted by fluorine, chlorine, methyl or ethyl, or

pyridyloxy-C₁-C₄-alkyl, pyrimidyloxy-C₁-C₄-alkyl or thiazolyloxy-C₁-C₄-alkyl, each of which is optionally substituted by fluorine, chlorine, amino, methyl or ethyl.

5

R² very particularly preferably represents C₁-C₁₄-alkyl, C₂-C₁₄-alkenyl, C₁-C₄-alkoxy-C₂-C₆-alkyl or poly-C₁-C₄-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by fluorine or chlorine,

10

C₃-C₆-cycloalkyl which is optionally substituted by fluorine, chlorine, methyl, ethyl, n-propyl, iso-propyl or methoxy,

or phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, cyano, nitro, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy.

15

R³ very particularly preferably represents methyl, ethyl, propyl, isopropyl, each of which is optionally substituted by fluorine or chlorine, or phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, propyl, iso-propyl, tert-butyl, methoxy, ethoxy, isopropoxy, tert-butoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro.

20

R⁴ and R⁵ independently of one another very particularly preferably represent C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylamino, di-(C₁-C₄-alkyl)amino or C₁-C₄-alkylthio, each of which is optionally substituted by fluorine or chlorine, or phenyl, phenoxy or phenylthio, each of which is optionally substituted by fluorine, chlorine, bromine, nitro, cyano, methyl, methoxy, trifluoromethyl or trifluoromethoxy.

Le A 31 064 - Foreign Countries

- 47 -

R⁶ and R⁷ independently of one another very particularly preferably represent hydrogen, or C₁-C₄-alkyl, C₃-C₆-cycloalkyl, C₁-C₄-alkoxy, C₃-C₄-alkenyl or C₁-C₄-alkoxy-C₂-C₄-alkyl, each of which is optionally substituted by fluorine or chlorine, or phenyl or benzyl, each of which is optionally substituted by fluorine, chlorine, bromine, methyl, methoxy or trifluoromethyl, or together represent a C₅-C₆-alkylene radical which is optionally substituted by methyl or ethyl and in which a methylene group is optionally replaced by oxygen or sulphur.

R¹³ very particularly preferably represents hydrogen or C₁-C₄-alkyl or C₁-C₄-alkoxy, each of which is optionally substituted by fluorine or chlorine, or C₃-C₆-cycloalkyl, or phenyl, phenyl-C₁-C₂-alkyl or benzyloxy, each of which is optionally substituted by fluorine, chlorine, bromine, methyl, ethyl, iso-propyl, tert-butyl, methoxy, ethoxy, iso-propoxy, tert-butoxy, trifluoromethyl, trifluoromethoxy, nitro or cyano.

R¹⁴ very particularly preferably represents hydrogen or C₁-C₄-alkyl or

R¹³ and R¹⁴ together very particularly preferably represent C₄-C₆-alkanediyl.

R¹⁵ and R¹⁶ are identical or different and very particularly preferably represent methyl or ethyl or

R¹⁵ and R¹⁶ together very particularly preferably represent a C₂-C₃-alkanediyl radical which is optionally substituted by methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl or by phenyl which is optionally substituted by fluorine, chlorine, methoxy, trifluoromethyl, trifluoromethoxy, nitro or cyano.

The abovementioned definitions or explanations of radicals mentioned generally or in preferred ranges can be combined with one another as desired, i.e. also between the respective ranges and preferred ranges. They apply correspondingly to the final products and to the precursors and intermediates.

Le A 31 064 - Foreign Countries

- 48 -

Preferred compounds of the formula (I) according to the invention are those in which a combination of the meanings (preferably) mentioned above as preferred is present.

Particularly preferred compounds of the formula (I) according to the invention are those in which a combination of the meanings mentioned above as particularly preferred is present.

5

Very particularly preferred compounds of the formula (I) according to the invention are those in which a combination of the meanings mentioned above as very particularly preferred is present.

Saturated or unsaturated hydrocarbon radicals such as alkyl or alkenyl can, as far as possible, in each case be straight-chain or branched, even in combination with heteroatoms, e.g. in alkoxy.


10

Optionally substituted radicals can be mono- or polysubstituted, it being possible in the case of polysubstitution for the substituents to be identical or different.

Le A 31 064 - Foreign Countries

- 49 -

In addition to the compounds mentioned in the Preparation Examples, the following compounds of the formula (I-1-a) may be specifically mentioned:




**Table 1:**

| A | B | D |
|---------------------------------|-----------------|---|
| CH ₃ | H | H |
| C ₂ H ₅ | H | H |
| C ₃ H ₇ | H | H |
| i-C ₃ H ₇ | H | H |
| C ₄ H ₉ | H | H |
| i-C ₄ H ₉ | H | H |
| s-C ₄ H ₉ | H | H |
| t-C ₄ H ₉ | H | H |
| CH ₃ | CH ₃ | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₃ H ₇ | CH ₃ | H |
| i-C ₃ H ₇ | CH ₃ | H |
| C ₄ H ₉ | CH ₃ | H |
| i-C ₄ H ₉ | CH ₃ | H |
| s-C ₄ H ₉ | CH ₃ | H |

Le A 31 064-Foreign Countries

- 50 -



Continuation of Table 1:

| A | B | D |
|---|--|---|
| t-C ₄ H ₉ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |
| C ₃ H ₇ | C ₃ H ₇ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
| | -(CH ₂) ₂ - | H |
| | -(CH ₂) ₄ - | H |
| | -(CH ₂) ₅ - | H |
| | -(CH ₂) ₆ - | H |
| | -(CH ₂) ₇ - | H |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - | H |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - | H |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | H |

Le A 31 064-Foreign Countries

- 51 -




Continuation of Table 1:

| A | B | D |
|---|--|---|
| | $-(\text{CH}_2)_2-\text{CHOC}_2\text{H}_5-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{CHOC}_3\text{H}_7-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{CHi-OC}_3\text{H}_7-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_2-$ | H |
| | $-\text{CH}_2-(\text{CHCH}_3)_2-(\text{CH}_2)_2-$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---}(\text{CH}_2)_2\text{---CH---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{CH}_2 \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---} \qquad \qquad \text{CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_4 \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---} \qquad \qquad \text{CH---}(\text{CH}_2)_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_3 \end{array}$ | H |
| |  | H |
| |  | H |

Le A 31 064-Foreign Countries

- 52 -




Continuation of Table 1:

| A | D | B |
|-----------------|---|---|
| | -(CH ₂) ₃ - | H |
| | -(CH ₂) ₄ - | H |
| | -CH ₂ -CHCH ₃ -CH ₂ - | H |
| | -CH ₂ -CH ₂ -CHCH ₃ - | H |
| | -CH ₂ -CHCH ₃ -CHCH ₃ - | H |
| | -CH ₂ -S-CH ₂ - | H |
| | -CH ₂ -S-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-CH ₂ - | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---} \\ \qquad \qquad \\ \text{---(CH}_2\text{)}_3\text{---} \end{array}$ | H |
| H | CH ₃ | H |
| H | C ₂ H ₅ | H |
| H | C ₃ H ₇ | H |
| H | i-C ₃ H ₇ | H |
| H |  | H |
| H |  | H |
| H |  | H |
| CH ₃ | CH ₃ | H |
| CH ₃ | C ₂ H ₅ | H |

Le A 31 064-Foreign Countries


- 53 -

Continuation of Table 1:

| A | D | B |
|-------------------------------|--|---|
| CH ₃ | C ₃ H ₇ | H |
| CH ₃ | i-C ₃ H ₇ | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |

Le A 31 064-Foreign Countries

- 54 -




**Table 2:**

| A | B | D |
|---------------------------------|-----------------|---|
| CH ₃ | H | H |
| C ₂ H ₅ | H | H |
| C ₃ H ₇ | H | H |
| i-C ₃ H ₇ | H | H |
| C ₄ H ₉ | H | H |
| i-C ₄ H ₉ | H | H |
| s-C ₄ H ₉ | H | H |
| t-C ₄ H ₉ | H | H |
| CH ₃ | CH ₃ | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₃ H ₇ | CH ₃ | H |
| i-C ₃ H ₇ | CH ₃ | H |
| C ₄ H ₉ | CH ₃ | H |
| i-C ₄ H ₉ | CH ₃ | H |
| s-C ₄ H ₉ | CH ₃ | H |

Le A 31 064-Foreign Countries

- 55 -



Continuation of Table 2:

| A | B | D |
|---|--|---|
| t-C ₄ H ₉ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |
| C ₃ H ₇ | C ₃ H ₇ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
| | -(CH ₂) ₂ - | H |
| | -(CH ₂) ₄ - | H |
| | -(CH ₂) ₅ - | H |
| | -(CH ₂) ₆ - | H |
| | -(CH ₂) ₇ - | H |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - | H |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - | H |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - | H |

Le A 31 064-Foreign Countries

- 56 -




Continuation of Table 2:

| A | B | D |
|---|---|---|
| | $-(\text{CH}_2)_2-\text{CHOCH}_3-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{CHOC}_2\text{H}_5-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{CHOC}_3\text{H}_7-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{CHi-OC}_3\text{H}_7-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_2-$ | H |
| | $-\text{CH}_2-(\text{CHCH}_3)_2-(\text{CH}_2)_2-$ | H |
| | $\begin{array}{c} -\text{CH}_2-\text{CH}-(\text{CH}_2)_2-\text{CH}- \\ \qquad \qquad \\ \text{CH}_2 \end{array}$ | H |
| | $\begin{array}{c} -\text{CH}_2-\text{CH} \qquad \qquad \text{CH}-\text{CH}_2- \\ \qquad \qquad \qquad \\ (\text{CH}_2)_4 \end{array}$ | H |
| | $\begin{array}{c} -\text{CH}_2-\text{CH} \qquad \qquad \text{CH}-(\text{CH}_2)_2- \\ \qquad \qquad \qquad \\ (\text{CH}_2)_3 \end{array}$ | H |
| |  | H |
| |  | H |

Le A 31 064-Foreign Countries

- 57 -




Continuation of Table 2:

| A | D | B |
|-----------------|---|----|
| | -(CH ₂) ₃ - | H |
| | -(CH ₂) ₄ - | H |
| | -CH ₂ -CHCH ₃ -CH ₂ - | H |
| | -CH ₂ -CH ₂ -CHCH ₃ - | H |
| | -CH ₂ -CHCH ₃ -CHCH ₃ - | H |
| | -CH ₂ -S-CH ₂ - | H |
| | -CH ₂ -S-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-CH ₂ - | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---} \\ \qquad \qquad \\ \text{---(CH}_2\text{)}_3\text{---} \end{array}$ | H |
| H | CH ₃ | H' |
| H | C ₂ H ₅ | H |
| H | C ₃ H ₇ | H |
| H | i-C ₃ H ₇ | H |
| H |  | H |
| H |  | H |
| H |  | H |
| CH ₃ | CH ₃ | H |

Le A 31 064-Foreign Countries


- 58 -

Continuation of Table 2:

| A | D | B |
|-------------------------------|--|---|
| CH ₃ | C ₂ H ₅ | H |
| CH ₃ | C ₃ H ₇ | H |
| CH ₃ | i-C ₃ H ₇ | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |

Le A 31 064-Foreign Countries

- 59 -




**Table 3:**

| A | B | D |
|---------------------------------|-----------------|---|
| CH ₃ | H | H |
| C ₂ H ₅ | H | H |
| C ₃ H ₇ | H | H |
| i-C ₃ H ₇ | H | H |
| C ₄ H ₉ | H | H |
| i-C ₄ H ₉ | H | H |
| s-C ₄ H ₉ | H | H |
| t-C ₄ H ₉ | H | H |
| CH ₃ | CH ₃ | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₃ H ₇ | CH ₃ | H |
| i-C ₃ H ₇ | CH ₃ | H |
| C ₄ H ₉ | CH ₃ | H |
| i-C ₄ H ₉ | CH ₃ | H |
| s-C ₄ H ₉ | CH ₃ | H |
| t-C ₄ H ₉ | CH ₃ | H |

Le A 31 064-Foreign Countries

- 60 -



Continuation of Table 3:

| A | B | D |
|---|----------------------------------|---|
| C_2H_5 | C_2H_5 | H |
| C_3H_7 | C_3H_7 | H |
|  | CH_3 | H |
|  | CH_3 | H |
|  | CH_3 | H |
| | $-(CH_2)_2-$ | H |
| | $-(CH_2)_4-$ | H |
| | $-(CH_2)_5-$ | H |
| | $-(CH_2)_6-$ | H |
| | $-(CH_2)_7-$ | H |
| | $-(CH_2)_2-O-(CH_2)_2-$ | H |
| | $-(CH_2)_2-S-(CH_2)_2-$ | H |
| | $-CH_2-CHCH_3-(CH_2)_3-$ | H |
| | $-(CH_2)_2-CHCH_3-(CH_2)_2-$ | H |
| | $-(CH_2)_2-CHC_2H_5-(CH_2)_2-$ | H |
| | $-(CH_2)_2-CHC_3H_7-(CH_2)_2-$ | H |
| | $-(CH_2)_2-CHi-C_3H_7-(CH_2)_2-$ | H |
| | $-(CH_2)_2-CHOCH_3-(CH_2)_2-$ | H |

Le A 31 064-Foreign Countries

- 61 -




Continuation of Table 3:

| A | B | D |
|---|---|---|
| | -(CH ₂) ₂ -CHOC ₂ H ₅ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHOC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHi-OC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -C(CH ₃) ₂ -(CH ₂) ₂ - | H |
| | -CH ₂ -(CHCH ₃) ₂ -(CH ₂) ₂ - | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---(CH}_2\text{)}_2\text{---CH---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{---CH}_2\text{---} \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_4 \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---(CH}_2\text{)}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_3 \end{array}$ | H |
| |  | H |
| |  | H |

Le A 31 064-Foreign Countries

- 62 -




Continuation of Table 3:

| A | D | B |
|---------------|---|---|
| | $-(\text{CH}_2)_3-$ | H |
| | $-(\text{CH}_2)_4-$ | H |
| | $-\text{CH}_2-\text{CHCH}_3-\text{CH}_2-$ | H |
| | $-\text{CH}_2-\text{CH}_2-\text{CHCH}_3-$ | H |
| | $-\text{CH}_2-\text{CHCH}_3-\text{CHCH}_3-$ | H |
| | $-\text{CH}_2-\text{S}-\text{CH}_2-$ | H |
| | $-\text{CH}_2-\text{S}-(\text{CH}_2)_2-$ | H |
| | $-(\text{CH}_2)_2-\text{S}-\text{CH}_2-$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---} \\ \qquad \qquad \\ \text{---(CH}_2\text{)}_3\text{---} \end{array}$ | H |
| H | CH_3 | H |
| H | C_2H_5 | H |
| H | C_3H_7 | H |
| H | $i\text{-C}_3\text{H}_7$ | H |
| H |  | H |
| H |  | H |
| H |  | H |
| CH_3 | CH_3 | H |
| CH_3 | C_2H_5 | H |

Le A 31 064-Foreign Countries


- 63 -

Continuation of Table 3:

| A | D | B |
|-------------------------------|--|---|
| CH ₃ | C ₃ H ₇ | H |
| CH ₃ | i-C ₃ H ₇ | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |

Le A 31 064-Foreign Countries

- 64 -




**Table 4:**

| A | B | D |
|---------------------------------|-----------------|---|
| CH ₃ | H | H |
| C ₂ H ₅ | H | H |
| C ₃ H ₇ | H | H |
| i-C ₃ H ₇ | H | H |
| C ₄ H ₉ | H | H |
| i-C ₄ H ₉ | H | H |
| s-C ₄ H ₉ | H | H |
| t-C ₄ H ₉ | H | H |
| CH ₃ | CH ₃ | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₃ H ₇ | CH ₃ | H |
| i-C ₃ H ₇ | CH ₃ | H |
| C ₄ H ₉ | CH ₃ | H |
| i-C ₄ H ₉ | CH ₃ | H |
| s-C ₄ H ₉ | CH ₃ | H |

Le A 31 064-Foreign Countries

- 65 -



Continuation of Table 4:

| A | B | D |
|---|--|---|
| t-C ₄ H ₉ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |
| C ₃ H ₇ | C ₃ H ₇ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
|  | CH ₃ | H |
| | -(CH ₂) ₂ - | H |
| | -(CH ₂) ₄ - | H |
| | -(CH ₂) ₅ - | H |
| | -(CH ₂) ₆ - | H |
| | -(CH ₂) ₇ - | H |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - | H |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - | H |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - | H |

Le A 31 064-Foreign Countries

- 66 -




Continuation of Table 4:

| A | B | D |
|---|---|---|
| | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHOC ₂ H ₅ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHOC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -CHi-OC ₃ H ₇ -(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -C(CH ₃) ₂ -(CH ₂) ₂ - | H |
| | -CH ₂ -(CHCH ₃) ₂ -(CH ₂) ₂ - | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---(CH}_2\text{)}_2\text{---CH---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{CH}_2 \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_4 \end{array}$ | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---(CH}_2\text{)}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_3 \end{array}$ | H |
| |  | H |
| |  | H |

Le A 31 064-Foreign Countries

- 67 -




Continuation of Table 4:

| A | D | B |
|-----------------|---|---|
| | -(CH ₂) ₃ - | H |
| | -(CH ₂) ₄ - | H |
| | -CH ₂ -CHCH ₃ -CH ₂ - | H |
| | -CH ₂ -CH ₂ -CHCH ₃ - | H |
| | -CH ₂ -CHCH ₃ -CHCH ₃ - | H |
| | -CH ₂ -S-CH ₂ - | H |
| | -CH ₂ -S-(CH ₂) ₂ - | H |
| | -(CH ₂) ₂ -S-CH ₂ - | H |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---} \\ \qquad \qquad \\ \text{---(CH}_2\text{)}_3\text{---} \end{array}$ | H |
| H | CH ₃ | H |
| H | C ₂ H ₅ | H |
| H | C ₃ H ₇ | H |
| H | i-C ₃ H ₇ | H |
| H |  | H |
| H |  | H |
| H |  | H |
| CH ₃ | CH ₃ | H |

Le A 31 064-Foreign Countries

- 68 -


Continuation of Table 4:

| A | D | B |
|-------------------------------|--|---|
| CH ₃ | C ₂ H ₅ | H |
| CH ₃ | C ₃ H ₇ | H |
| CH ₃ | i-C ₃ H ₇ | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| CH ₃ |  | H |
| C ₂ H ₅ | CH ₃ | H |
| C ₂ H ₅ | C ₂ H ₅ | H |

Le A 31 064-Foreign Countries

- 69 -

In addition to the compounds mentioned in the Preparation Examples, the following compounds of the formula (I-2-a) may specifically be mentioned:




**Table 5:**

| A | B |
|---------------------------------|-----------------|
| CH ₃ | H |
| C ₂ H ₅ | H |
| C ₃ H ₇ | H |
| i-C ₃ H ₇ | H |
| C ₄ H ₉ | H |
| i-C ₄ H ₉ | H |
| s-C ₄ H ₉ | H |
| t-C ₄ H ₉ | H |
| CH ₃ | CH ₃ |
| C ₂ H ₅ | CH ₃ |
| C ₃ H ₇ | CH ₃ |
| i-C ₃ H ₇ | CH ₃ |
| C ₄ H ₉ | CH ₃ |
| i-C ₄ H ₉ | CH ₃ |
| s-C ₄ H ₉ | CH ₃ |

Le A 31 064-Foreign Countries

- 70 -



Continuation of Table 5:

| A | B |
|---|--|
| t-C ₄ H ₉ | CH ₃ |
| C ₂ H ₅ | C ₂ H ₅ |
| C ₃ H ₇ | C ₃ H ₇ |
|  | CH ₃ |
|  | CH ₃ |
|  | CH ₃ |
| | -(CH ₂) ₂ - |
| | -(CH ₂) ₄ - |
| | -(CH ₂) ₅ - |
| | -(CH ₂) ₆ - |
| | -(CH ₂) ₇ - |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - |

Le A 31 064-Foreign Countries


- 71 -

Continuation of Table 5:

| A | B |
|---|--|
| | $-(\text{CH}_2)_2\text{-CHOCH}_3\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHOC}_2\text{H}_5\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHOC}_3\text{H}_7\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHi-OC}_3\text{H}_7\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-C(CH}_3)_2\text{-(CH}_2)_2\text{-}$ |
| | $-\text{CH}_2\text{-(CHCH}_3)_2\text{-(CH}_2)_2\text{-}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---(CH}_2)_2\text{---CH---} \\ \qquad \qquad \\ \text{---CH}_2\text{---} \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---CH}_2\text{---} \\ \qquad \qquad \\ \text{---(CH}_2)_4\text{---} \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---(CH}_2)_2\text{---} \\ \qquad \qquad \\ \text{---(CH}_2)_3\text{---} \end{array}$ |
| |  |
| |  |

Le A 31 064-Foreign Countries

- 72 -




**Table 6:**

| A | B |
|---------------------------------|-----------------|
| CH ₃ | H |
| C ₂ H ₅ | H |
| C ₃ H ₇ | H |
| i-C ₃ H ₇ | H |
| C ₄ H ₉ | H |
| i-C ₄ H ₉ | H |
| s-C ₄ H ₉ | H |
| t-C ₄ H ₉ | H |
| CH ₃ | CH ₃ |
| C ₂ H ₅ | CH ₃ |
| C ₃ H ₇ | CH ₃ |
| i-C ₃ H ₇ | CH ₃ |
| C ₄ H ₉ | CH ₃ |
| i-C ₄ H ₉ | CH ₃ |
| s-C ₄ H ₉ | CH ₃ |

Le A 31 064-Foreign Countries

- 73 -



Continuation of Table 6:

| A | B |
|---|--|
| t-C ₄ H ₉ | CH ₃ |
| C ₂ H ₅ | C ₂ H ₅ |
| C ₃ H ₇ | C ₃ H ₇ |
|  | CH ₃ |
|  | CH ₃ |
|  | CH ₃ |
| | -(CH ₂) ₂ - |
| | -(CH ₂) ₄ - |
| | -(CH ₂) ₅ - |
| | -(CH ₂) ₆ - |
| | -(CH ₂) ₇ - |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - |

Le A 31 064-Foreign Countries


- 74 -

Continuation of Table 6:

| A | B |
|---|---|
| | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHOC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHOC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHi-OC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -C(CH ₃) ₂ -(CH ₂) ₂ - |
| | -CH ₂ -(CHCH ₃) ₂ -(CH ₂) ₂ - |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---(CH}_2\text{)}_2\text{---CH---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{CH}_2 \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_4 \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---(CH}_2\text{)}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{(CH}_2\text{)}_3 \end{array}$ |
| |  |
| |  |

Le A 31 064-Foreign Countries

- 75 -




**Table 7:**

| A | B |
|---------------------------------|-----------------|
| CH ₃ | H |
| C ₂ H ₅ | H |
| C ₃ H ₇ | H |
| i-C ₃ H ₇ | H |
| C ₄ H ₉ | H |
| i-C ₄ H ₉ | H |
| s-C ₄ H ₉ | H |
| t-C ₄ H ₉ | H |
| CH ₃ | CH ₃ |
| C ₂ H ₅ | CH ₃ |
| C ₃ H ₇ | CH ₃ |
| i-C ₃ H ₇ | CH ₃ |
| C ₄ H ₉ | CH ₃ |
| i-C ₄ H ₉ | CH ₃ |
| s-C ₄ H ₉ | CH ₃ |

Le A 31 064-Foreign Countries

- 76 -



Continuation of Table 7:

| A | B |
|---|--|
| t-C ₄ H ₉ | CH ₃ |
| C ₂ H ₅ | C ₂ H ₅ |
| C ₃ H ₇ | C ₃ H ₇ |
|  | CH ₃ |
|  | CH ₃ |
|  | CH ₃ |
| | -(CH ₂) ₂ - |
| | -(CH ₂) ₄ - |
| | -(CH ₂) ₅ - |
| | -(CH ₂) ₆ - |
| | -(CH ₂) ₇ - |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - |

Le A 31 064-Foreign Countries


- 77 -

Continuation of Table 7:

| A | B |
|---|---|
| | $-(\text{CH}_2)_2-\text{CHOCH}_3-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{CHOC}_2\text{H}_5-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{CHOC}_3\text{H}_7-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{CHi-OC}_3\text{H}_7-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_2-$ |
| | $-\text{CH}_2-(\text{CHCH}_3)_2-(\text{CH}_2)_2-$ |
| | $\begin{array}{c} -\text{CH}_2-\text{CH}-(\text{CH}_2)_2-\text{CH}- \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{CH}_2 \end{array}$ |
| | $\begin{array}{c} -\text{CH}_2-\text{CH} \qquad \qquad \qquad \text{CH}-\text{CH}_2- \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_4 \end{array}$ |
| | $\begin{array}{c} -\text{CH}_2-\text{CH} \qquad \qquad \qquad \text{CH}-(\text{CH}_2)_2- \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_3 \end{array}$ |
| |  |
| |  |

Le A 31 064-Foreign Countries

- 78 -




**Table 8:**

| A | B |
|---------------------------------|-----------------|
| CH ₃ | H |
| C ₂ H ₅ | H |
| C ₃ H ₇ | H |
| i-C ₃ H ₇ | H |
| C ₄ H ₉ | H |
| i-C ₄ H ₉ | H |
| s-C ₄ H ₉ | H |
| t-C ₄ H ₉ | H |
| CH ₃ | CH ₃ |
| C ₂ H ₅ | CH ₃ |
| C ₃ H ₇ | CH ₃ |
| i-C ₃ H ₇ | CH ₃ |
| C ₄ H ₉ | CH ₃ |
| i-C ₄ H ₉ | CH ₃ |
| s-C ₄ H ₉ | CH ₃ |

Le A 31 064-Foreign Countries

- 79 -



Continuation of Table 8:

| A | B |
|---|--|
| t-C ₄ H ₉ | CH ₃ |
| C ₂ H ₅ | C ₂ H ₅ |
| C ₃ H ₇ | C ₃ H ₇ |
|  | CH ₃ |
|  | CH ₃ |
|  | CH ₃ |
| | -(CH ₂) ₂ - |
| | -(CH ₂) ₄ - |
| | -(CH ₂) ₅ - |
| | -(CH ₂) ₆ - |
| | -(CH ₂) ₇ - |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHi-C ₃ H ₇ -(CH ₂) ₂ - |

Le A 31 064-Foreign Countries

- 80 -

Continuation of Table 8:

| A | B |
|---|--|
| | $-(\text{CH}_2)_2-\text{CHOCH}_3-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{CHOC}_2\text{H}_5-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{CHOC}_3\text{H}_7-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{Chi-OC}_3\text{H}_7-(\text{CH}_2)_2-$ |
| | $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_2-$ |
| | $-\text{CH}_2-(\text{CHCH}_3)_2-(\text{CH}_2)_2-$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---}(\text{CH}_2)_2\text{---CH---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad \text{CH}_2 \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---} \qquad \qquad \text{CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_4 \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---} \qquad \qquad \text{CH---}(\text{CH}_2)_2\text{---} \\ \qquad \qquad \qquad \\ \qquad \qquad \qquad (\text{CH}_2)_3 \end{array}$ |
| |  |
| |  |

In addition to the compounds mentioned in the Preparation Examples, the following compounds of the formula (I-3-a) may specifically be mentioned:









Table 9:

| A | B |
|---------------------------------|-----------------|
| CH ₃ | H |
| C ₂ H ₅ | H |
| C ₃ H ₇ | H |
| i-C ₃ H ₇ | H |
| C ₄ H ₉ | H |
| i-C ₄ H ₉ | H |
| s-C ₄ H ₉ | H |
| t-C ₄ H ₉ | H |
| CH ₃ | CH ₃ |
| C ₂ H ₅ | CH ₃ |
| C ₃ H ₇ | CH ₃ |
| i-C ₃ H ₇ | CH ₃ |
| C ₄ H ₉ | CH ₃ |
| i-C ₄ H ₉ | CH ₃ |
| s-C ₄ H ₉ | CH ₃ |

Continuation of Table 9:

| A | B |
|---|--|
| t-C ₄ H ₉ | CH ₃ |
| C ₂ H ₅ | C ₂ H ₅ |
| C ₃ H ₇ | C ₃ H ₇ |
|  | CH ₃ |
|  | CH ₃ |
|  | CH ₃ |
| | -(CH ₂) ₂ - |
| | -(CH ₂) ₄ - |
| | -(CH ₂) ₅ - |
| | -(CH ₂) ₆ - |
| | -(CH ₂) ₇ - |
| | -(CH ₂) ₂ -O-(CH ₂) ₂ - |
| | -(CH ₂) ₂ -S-(CH ₂) ₂ - |
| | -CH ₂ -CHCH ₃ -(CH ₂) ₃ - |
| | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₂ H ₅ -(CH ₂) ₂ - |
| | -(CH ₂) ₂ -CHC ₃ H ₇ -(CH ₂) ₂ - |

Continuation of Table 9:

| A | B |
|---|---|
| | $-(\text{CH}_2)_2\text{-CHi-C}_3\text{H}_7\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHOCH}_3\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHOC}_2\text{H}_5\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHOC}_3\text{H}_7\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-CHi-OC}_3\text{H}_7\text{-(CH}_2)_2\text{-}$ |
| | $-(\text{CH}_2)_2\text{-C(CH}_3)_2\text{-(CH}_2)_2\text{-}$ |
| | $-\text{CH}_2\text{-(CHCH}_3)_2\text{-(CH}_2)_2\text{-}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---(CH}_2)_2\text{---CH---} \\ \qquad \qquad \qquad \\ \text{---CH}_2\text{---} \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---CH}_2\text{---} \\ \qquad \qquad \qquad \\ \text{---(CH}_2)_4\text{---} \end{array}$ |
| | $\begin{array}{c} \text{---CH}_2\text{---CH---CH---(CH}_2)_2\text{---} \\ \qquad \qquad \qquad \\ \text{---(CH}_2)_3\text{---} \end{array}$ |
| |  |
| |  |

Le A 31 064 - Foreign Countries

- 84 -

Table 10:

A, B as indicated in Table 9

X = CH₃; Y = Cl; Z = CH₃**Table 11:**

A, B as indicated in Table 9

X = CH₃; Y = CH₃; Z = Br**Table 12:**

A, B as indicated in Table 9


X = CH₃; Y = Br; Z = CH₃

In addition to the compounds mentioned in the Preparation Examples, the following compounds of the formula (I-5-a) may specifically be mentioned:

Le A 31 064 - Foreign Countries

- 85 -

Table 13:











| A | D |
|-----------------|---|
| H | CH ₃ |
| H | C(CH ₃) ₃ |
| H | C(CH ₃) ₂ CH ₂ Cl |
| CH ₃ | CH ₃ |
| CH ₃ | CH ₂ CHCH ₃ CH ₂ CH ₃ |
| H | CH=C(CH ₃) ₂ |
| CH ₃ | |
| CH ₃ | |
| CH ₃ | |
| CH ₃ | |

Le A 31 064 - Foreign Countries

- 86 -

Continuation of Table 13:

| A | D |
|---|---|
| CH ₃ |  |
|  | CH ₃ |
| H |  |
| CH ₃ |  |
| CH ₃ |  |
| CH ₃ |  |
| CH ₃ |  |
| H |  |
| CH ₃ | C ₅ H ₉ |
| CH ₃ | C ₃ H ₅ |

Continuation of Table 13:

| A | D |
|-----------------------|------------|
| H | C_3H_4Cl |
| $(CH_2)_3$ | |
| $(CH_2)_4$ | |
| $C(CH_3)_2OC(CH_3)_2$ | |

Table 14:

A and D as indicated in Table 13

$X = CH_3$; $Y = Cl$; $Z = CH_3$

Table 15:

A and D as indicated in Table 13

$X = CH_3$; $Y = CH_3$; $Z = Br$

Table 16:


A and D as indicated in Table 13

$X = CH_3$; $Y = Br$; $Z = CH_3$

In addition to the compounds mentioned in the Preparation Examples, the following compounds of the formula (I-6-a) may specifically be mentioned:

Le A 31 064 - Foreign Countries

- 88 -

Table 17:

5

| |
|-----------------------------------|
| A |
| CH ₃ |
| CH(CH ₃) ₂ |
| |
| |
| |
| |

10

Table 18:

A and D as indicated in Table 17

X = CH₃; Y = Cl; Z = CH₃**Table 19:**

A as indicated in Table 17

15

X = CH₃; Y = CH₃; Z = Br

Le A 31 064 - Foreign Countries


- 89 -

Table 20:


A as indicated in Table 17

X = CH₃; Y = Br; Z = CH₃

If according to process (A) ethyl N-[(2-chloro-4,6-dimethyl)-phenylacetyl]-1-amino-4-ethyl-cyclohexane-carboxylate is used as a starting substance, the course of the process according to the invention can be represented by the following equation:




If according to process (B) ethyl O-[(2-chloro-4,6-dimethyl)phenylacetyl]hydroxyacetate is used, the course of the process according to the invention can be represented by the following equation:




If according to process (C) ethyl 2-[(2-bromo-4,6-dimethyl)phenyl]-4-(4-methoxy)benzylmercapto-4-methyl-3-oxo-valerate is used, the course of the process according to the invention can be represented by the following equation:

Le A 31 064 - Foreign Countries


- 90 -



If, for example, according to process (E) chlorocarbonyl 2-[(2-bromo-4,6-dimethyl)phenyl ketene and acetone are used as starting compounds, the course of the process according to the invention can be represented by the following equation:




If, for example, according to process (F) chlorocarbonyl 2-(4-bromo-2,6-dimethyl)phenyl ketene and thiobenzamide are used as starting compounds, the course of the process according to the invention can be represented by the following equation:




If, according to process (G α) 3-[(2-chloro-4,6-dimethyl)-phenyl]-5,5-dimethylpyrrolidine-2,4-dione and pivaloyl chloride are used as starting substances, the course of the process according to the invention can be represented by the following equation:

Le A 31 064 - Foreign Countries


- 91 -



5 If according to process (G) (variant β) 3-[(4-chloro-2,6-dimethyl)-phenyl]-4-hydroxy-5-phenyl- Δ^3 -dihydrofuran-2-one and acetic anhydride are used as starting compounds, the course of the process according to the invention can be represented by the following equation:




10 If according to process (H) 8-[(4-bromo-2-ethyl-6-methyl)-phenyl]-1,6-diaza-bicyclo-(4,3,0^{1,6})-nonane-7,9-dione and ethoxyethyl chloroformate are used as starting compounds, the course of the process according to the invention can be represented by the following equation:




If according to process (I), (variant α), 3-[(2-chloro-4,6-dimethyl)-phenyl]-4-hydroxy-6-(3-pyridyl)-pyrone and methyl chloromonothioformate are used as starting materials, the course of the reaction can be represented in the following manner:

Le A 31 064 - Foreign Countries


- 92 -



If according to process (I), (variant β), 5-[(2-bromo-4-methyl-6-methyl)-phenyl]-6-hydroxy-2-(4-chlorophenyl)-thiazin-4-one, carbon disulphide and methyl iodide are used as starting components, the course of the reaction can be represented as follows:




If according to process (J) 2-[(2-chloro-4,6-dimethyl)-phenyl]-1,5-trimethylene-pyrrolidine-2,4-dione and methanesulfonyl chloride are used as starting materials, the course of the reaction can be represented by the following equation:




10 If according to process (K) 2-[(2-chloro-6-ethyl-4-methyl)-phenyl]-4-hydroxy-5-methyl-6-(2-pyridyl)-pyrone and 2,2,2-trifluoroethyl chloromethanethio-phosphonate are used as starting materials, the course of the reaction can be represented by the following equation:

Le A 31 064 - Foreign Countries


- 93 -



If according to process (L) 3-[(4-bromo-2,6-diethyl)-phenyl]-5-cyclopropyl-5-methyl-pyrrolidine-2,4-dione and NaOH are used as components, the course of the process according to the invention can be represented by the following equation:




If according to process (M) (variant α) 3-[(2-bromo-4,6-dimethyl)-phenyl]-4-hydroxy-5,5-tetramethylene- Δ^3 -dihydrofuran-2-one and ethyl isocyanate are used as starting materials, the course of the reaction can be represented by the following equation:




10 If according to process (M) (variant β) 3-[(2-chloro-4,6-dimethyl)-phenyl]-5-methyl-pyrrolidine-2,4-dione and dimethylcarbamoyl chloride are used as starting materials, the course of the reaction can be represented by the following equation:

Le A 31 064 - Foreign Countries

- 94 -



The compounds of the formula (II)




in which

- 5 A, B, D, X, Y, Z and R⁸ have the meanings given above,

needed as starting substances in process (A) according to the invention are new.

The acylamino acid esters of the formula (II) are obtained, for example, when amino acid derivatives of the formula (XXI)



- 10 in which

A, B, R⁸ and D have the meanings given above,

are acylated using substituted phenylacetyl halides of the formula (XXII)

Le A 31 064 - Foreign Countries

- 95 -




in which

X, Y and Z have the meanings given above and

Hal represents chlorine or bromine,

5 (Chem. Reviews 52, 237-416 (1953); Bhattacharya, Indian J. Chem. 6, 341-5, 1968)

or when acylamino acids of the formula (XXIII)




in which

A, B, D, X, Y and Z have the meanings given above,

10 are esterified (Chem. Ind. (London) 1568 (1968)).

The compounds of the formula (XXIII)



in which


A, B, D, X, Y and Z have the meanings given above,

Le A 31 064 - Foreign Countries

- 96 -

are new.

The compounds of the formula (XXIII) are obtained when amino acids of the formula (XXIV)



5 in which

A, B and D have the meanings given above,

are acylated using substituted phenylacetyl halides of the formula (XXII)



in which

10 X, Y and Z have the meanings given above and

Hal represents chlorine or bromine,


according to Schotten-Baumann (Organikum, VEB Deutscher Verlag der Wissenschaften, Berlin 1977, p. 505).

The compounds of the formula (XXII) are new.

15 The compounds of the formula (XXII) are obtained, for example, by reacting substituted phenylacetic acids of the formula (XXV)

Le A 31 064 - Foreign Countries

- 97 -




in which

X, Y and Z have the meanings given above,

- with halogenating agents (e.g. thionyl chloride, thionyl bromide, oxalyl chloride, phosgene, phosphorus trichloride, phosphorus tribromide or phosphorus pentachloride), if appropriate in the presence of a diluent (e.g. optionally chlorinated aliphatic or aromatic hydrocarbons such as toluene or methylene chloride) at temperatures from -20°C to 150°C, preferably from -10°C to 100°C.

The compounds of the formula (XXV) are new.

- 10 The compounds of the formula (XXV) are obtained, for example, by hydrolysing substituted phenylacetic acid esters of the formula (XXVI)



in which

X, Y, Z and R⁸ have the meaning give above,

- 15 at temperatures between 0°C and 150°C, preferably between 20°C and 100°C, in the presence of an acid (e.g. of an inorganic acid such as hydrochloric acid) or of a base (e.g. of an alkali metal hydroxide such as sodium or potassium hydroxide) and, if appropriate, of a diluent (e.g. of an aqueous alcohol such as methanol or ethanol).

The compounds of the formula (XXVI) are new.

Le A 31 064 - Foreign Countries

- 98 -

The compounds of the formula (XXVI) are obtained, for example, by reacting substituted 1,1,1-trichloro-2-phenylethanes of the formula (XXVII)



in which

5 X, Y and Z have the meanings given above,

first with alkoxides (e.g. alkali metal alkoxides such as sodium methoxide or sodium ethoxide) in the presence of a diluent (e.g. the alcohol derived from the alkoxide) at temperatures between 0°C and 150°C, preferably between 20°C and 120°C, and then reacting with an acid (preferably an inorganic acid, e.g. sulphuric acid) at temperatures
 10 between -20°C and 150°C, preferably 0°C and 100°C (cf. DE-33 14 249).

The compounds of the formula (XXVII) are new.

The compounds of the formula (XXVII) are obtained, for example, when anilines of the formula (XXVIII)



15 in which

X, Y and Z have the meaning given above,

are reacted with vinylidene chloride ($\text{CH}_2 = \text{CCl}_2$) in the presence of an alkyl nitrite of the formula (XXIX)



(XXIX)

in which


R²¹ represents alkyl, preferably C₁-C₆-alkyl,

in the presence of copper(II) chloride and if appropriate in the presence of a diluent
 5 (e.g. of an aliphatic nitrile such as acetonitrile) at a temperature of -20°C to 80°C,
 preferably 0°C to 60°C (cf. J. Org. Chem. 53 (1988), 3637).


The compounds of the formulae (XXVIII) and (XXIX) are known compounds of
 organic chemistry. Copper(II) chloride and vinylidene chloride are long-known and
 commercially available.

10 The compounds of the formulae (XXI) and (XXIV) are known in some cases and/or
 can be prepared by known processes (see, for example, Compagnon, Miocque Ann.
 Chim. (Paris) [14] 5, p. 11-22, 23-27 (1970)).

The substituted cyclic aminocarboxylic acids of the formula (XXIVa), in which A and
 B form a ring, are in general obtainable by the Bucherer-Bergs synthesis or by the
 15 Strecker synthesis and are in each case obtained here in different isomeric forms. Thus,
 according to the conditions of the Bucherer-Bergs synthesis mainly the isomers (in the
 following designated as β for the sake of simplicity) in which the radicals R and the
 carboxyl group are equatorial are obtained, while according to the conditions of the
 Strecker synthesis mainly the isomers (in the following designated as α for the sake of
 20 simplicity) are obtained in which the amino group and the radicals R are equatorial.



Bucherer-Bergs synthesis
 (β -isomer)




Strecker synthesis
 (α -isomer)

Le A 31 064 - Foreign Countries

- 100 -

(L. Munday, J. Chem. Soc. 4372 (1961); J.T. Eward, C. Jitrangeri, Can. J. Chem. 53, 3339 (1975)).

Furthermore, the starting substances of the formula (II)



5 in which

A, B, D, X, Y, Z and R⁸ have the meanings given above,

used in the above process (A) can be prepared when aminonitriles of the formula (XXX)



10 in which

A, B and D have the meanings given above,

are reacted with substituted phenylacetyl halides of the formula (XXII)




in which

15 X, Y, Z and Hal have the meanings given above,

Le A 31 064 - Foreign Countries

- 101 -

to give compounds of the formula (XXXI)




in which

A, B, D, X, Y and Z have the meanings given above,

5 and these are then subjected to an acidic alcoholysis.

The compounds of the formula (XXXI) are also new.

The compounds of the formula (III)



in which

10 A, B, X, Y, Z and R⁸ have the meanings given above,

needed as starting substances in process (B) according to the invention are new.

They can be prepared in a simple manner by methods known in principle.

The compounds of the formula (III) are obtained, for example, when

2-hydroxycarboxylic acid esters of the formula (XXXII)

Le A 31 064 - Foreign Countries


- 102 -



in which

A, B and R⁸ have the meanings given above,

are acylated using substituted phenylacetyl halides of the formula (XXII)




in which

X, Y, Z and Hal have the meanings given above,

(Chem. Reviews 52, 237-416 (1953)).

Furthermore, compounds of the formula (III) are obtained when

10 substituted phenylacetic acids of the formula (XXV)



in which

X, Y and Z have the meanings given above,

are alkylated using α -halogenocarboxylic acid esters of the formula (XXXIII)

Le A 31 064 - Foreign Countries

- 103 -




in which

A, B and R⁸ have the meanings given above and

Hal represents chlorine or bromine.

- 5 The compounds of the formula (XXXIII) are commercially available.

The compounds of the formula (IV)



in which

A, B, W, X, Y, Z and R⁸ have the meanings given above,


- 10 needed as starting substances in the above process (C) are new.

They can be prepared by methods known in principle.

The compounds of the formula (IV) are obtained, for example, when substituted phenylacetic acid esters of the formula (XXVI)

Le A 31 064 - Foreign Countries


- 104 -



in which

X, Y, R⁸ and Z have the meanings given above,

are acylated using 2-benzylthio-carbonyl halides of the formula (XXXIV)



in which

A, B and W have the meanings given above and

Hal represents halogen (in particular chlorine or bromine),


10 in the presence of strong bases (see, for example, M.S. Chambers, E.J. Thomas, D.J. Williams, J. Chem. Soc. Chem. Commun., (1987), 1228).

The benzylthio-carbonyl halides of the formula (XXXIV) are known in some cases and/or can be prepared by known methods (J. Antibiotics (1983), 26, 1589).

15 The halogenocarbonylketenes of the formula (V) needed as starting substances in process (E) are new. They can be prepared in a simple manner by methods known in principle (cf., for example, Org. Prep. Proced. Int., 7, (4), 155-158, 1975 and DE 1 945 703). The compounds of the formula (V)

Le A 31 064 - Foreign Countries

- 105 -




in which

X, Y and Z have the meanings given above and

Hal represents chlorine or bromine,

5 are obtained when

substituted phenylmalonic acids of the formula (XXXV)



in which

X, Y and Z have the meanings given above,

10 are reacted with acid halides, for example thionyl chloride, phosphorus(V) chloride, phosphorus(III) chloride, oxalyl chloride, phosgene or thionyl bromide, if appropriate in the presence of catalysts, for example diethylformamide, methylsterylformamide or triphenylphosphine and if appropriate in the presence of bases, e.g. pyridine or triethylamine, at a temperature between -20°C and 200°C, preferably between 0°C and



15 150°C.

The substituted phenylmalonic acids of the formula (XXXV) are new. However, they can be prepared by known processes in a simple manner (cf., for example Organikum, VEB Deutscher Verlag der Wissenschaften, Berlin 1977, p. 517 ff).

Le A 31 064 - Foreign Countries

- 106 -

The carbonyl compounds of the formula (VIII) or their silyl enol ethers of the formula (VIIIa)



in which

5 A, D and R^{δ'} have the meanings given above,

needed as starting substances for process (E) according to the invention are compounds which are commercially available, generally known or accessible by known processes.

The preparation of the ketene acid chlorides of the formula (V) needed as starting substances for carrying out process (F) according to the invention has already been described in process (E) according to the invention. The thioamides of the formula (IX)



in which

A has the meaning given above,

needed for carrying out process (F) according to the invention are compounds which are generally known in organic chemistry.


The compounds of the formula (I-4-a) needed as starting substances in process (G) are known and/or can be prepared in a simple manner by known methods (cf. WO 92/16510).

The compounds of the formula (I-4-a) are obtained, for example, when

Le A 31 064 - Foreign Countries

- 107 -

compounds of the formula (V)



in which


X, Y and Z have the meanings given above

5 and

Hal represents halogen (in particular chlorine or bromine),

or

compounds of the formula (VI)



10 in which

 R^8 , X, Y and Z have the meanings given above,

are reacted with hydrazines of the formula (VII)



in which

Le A 31 064 - Foreign Countries


- 108 -

A and D have the meanings given above,

if appropriate in the presence of a diluent,

it being possible to use hydrocarbons, such as toluene and xylene, further ethers, such as dibutyl ether, tetrahydrofuran, dioxane, glycol dimethyl ether and diglycol dimethyl ether, additionally polar solvents, such as dimethyl sulphoxide, sulpholane, dimethylformamide and N-methyl-pyrrolidone, and, only in the case in which compounds of the formula (VI) are employed, alcohols such as methanol, ethanol, propanol, iso-propanol, butanol, iso-butanol and tert-butanol, and if appropriate in the presence of a base, where in the case in which compounds of the formula (V) are employed, inorganic bases, in particular alkali metal or alkaline earth metal carbonates such as sodium carbonate, potassium carbonate or calcium carbonate, and organic bases, for example pyridine or triethylamine, are suitable and in the case where compounds of the formula (VI) are employed, alkali metal and alkaline earth metal oxides, hydroxides and carbonates, such as sodium hydroxide, potassium hydroxide, magnesium oxide, calcium oxide, sodium carbonate, potassium carbonate and calcium carbonate, each of which can also be employed in the presence of phase-transfer catalysts, e.g. triethylbenzylammonium chloride, tetrabutylammonium bromide, Adogen 464 (= methyltrialkyl(C₈-C₁₀)ammonium chloride) or TDA 1 (= tris-(methoxyethoxyethyl)-amine), alkali metals such as sodium or potassium, alkali metal and alkaline earth metal amides and hydrides, such as sodium amide, sodium hydride and calcium hydride, and additionally also alkali metal alkoxides, such as sodium methoxide, sodium ethoxide and potassium tert-butoxide are suitable, at temperatures between -20°C and 250°C, preferably between 0°C and 150°C.

The malonic acid esters of the formula (VI)



Le A 31 064 - Foreign Countries


- 109 -

in which

R⁸, X, Y and Z have the meanings given above, are new.

They can be prepared by generally known methods of organic chemistry (cf., for example, Tetrahedron Lett. 27, 2763 (1986) and Organikum VEB Deutscher Verlag der
5 Wissenschaften, Berlin 1977, p. 587 ff.).

The hydrazines of the formula (VII)



in which

A and D have the meanings given above,

10 are known in some cases and/or can be prepared by methods known from the literature (cf., for example, Liebigs Ann. Chem. 585, 6 (1954); Reaktionen der organischen Synthese [Reactions of Organic Synthesis], C. Ferri, page 212, 513; Georg Thieme Verlag Stuttgart, 1978; Liebigs Ann. Chem. 443, 242 (1925); Chem. Ber. 98, 2551 (1965), EP 508 126).

15 The acid halides of the formula (X), carboxylic anhydrides of the formula (XI), chloroformic acid esters or chloroformic acid thioesters of the formula (XII), chloromonothioformic acid esters or chlorodithioformic acid esters of the formula (XIII), alkyl halides of the formula (XIV), sulphonyl chlorides of the formula (XV), phosphorus compounds of the formula (XVI) and metal hydroxides, metal alkoxides or
20 amines of the formula (XVII) and (XVIII) and isocyanates of the formula (XIX) and carbamoyl chlorides of the formula (XX) additionally needed as starting substances for carrying out processes (G), (H), (I), (J), (K), (L) and (M) according to the invention are generally known compounds of organic or inorganic chemistry.

Le A 31 064 - Foreign Countries

- 110 -

The compounds of the formulae (VII), (VIII), (IX) to (XXI), (XXIV) and (XXXII) to (XXXIV) are moreover disclosed in the patent applications cited at the outset and/or can be prepared by the methods given there.

Process (A) is characterized in that compounds of the formula (II), in which A, B, D,
5 X, Y, Z and R⁸ have the meanings given above, are subjected to an intramolecular condensation in the presence of a diluent and in the presence of a base.

The diluents employed in process (A) according to the invention can be all organic solvents which are inert to the reaction participants. Those preferably utilizable are hydrocarbons, such as toluene and xylene, further ethers, such as dibutyl ether,
10 tetrahydrofuran, dioxane, glycol dimethyl ether and diglycol dimethyl ether, additionally polar solvents, such as dimethyl sulphoxide, sulpholane, dimethylformamide and N-methyl-pyrrolidone, and also alcohols such as methanol, ethanol, propanol, iso-propanol, butanol, iso-butanol and tert-butanol.

Suitable bases (deprotonating agents) employed in carrying out process (A) according
15 to the invention can be all customary proton acceptors. Those preferably utilizable are alkali metal and alkaline earth metal oxides, hydroxides and carbonates, such as sodium hydroxide, potassium hydroxide, magnesium oxide, calcium oxide, sodium carbonate, potassium carbonate and calcium carbonate, each of which can also be employed in the presence of phase-transfer catalysts, e.g. triethylbenzylammonium chloride,
20 tetrabutylammonium bromide, Adogen 464 (= methyltrialkyl(C₈-C₁₀)ammonium chloride) or TDA 1 (= tris-(methoxyethoxyethyl)-amine). Alkali metals such as sodium or potassium can furthermore be used. Alkali metal and alkaline earth metal amides and hydrides, such as sodium amide, sodium hydride and calcium hydride, and additionally also alkali metal alkoxides, such as sodium methoxide, sodium ethoxide and potassium
25 tert-butoxide can further be employed.

When carrying out process (A) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the reaction is carried out at temperatures between 0°C and 250°C, preferably between 50°C and 150°C.

Le A 31 064 - Foreign Countries

- 111 -

Process (A) according to the invention is in general carried out under normal pressure.

When carrying out process (A) according to the invention, the reaction component of the formula (II) and the deprotonating base are in general employed in equimolar to approximately double equimolar amounts. However, it is also possible to use one
5 component or the other in a relatively large excess (up to 3 mol).

Process (B) is characterized in that compounds of the formula (III), in which A, B, X, Y, Z and R⁸ have the meanings given above, are condensed intramolecularly in the presence of a diluent and in the presence of a base.

The diluents employed in process (B) according to the invention can be all organic
10 solvents which are inert to the reaction participants. Those preferably utilizable are hydrocarbons, such as toluene and xylene, further ethers, such as dibutyl ether, tetrahydrofuran, dioxane, glycol dimethyl ether and diglycol dimethyl ether, and additionally polar solvents, such as dimethyl sulphoxide, sulpholane, dimethylformamide and N-methyl-pyrrolidone. Alcohols such as methanol, ethanol,
15 propanol, iso-propanol, butanol, iso-butanol and tert-butanol can furthermore be employed.

The bases (deprotonating agents) employed in carrying out process (B) according to the invention can be all customary proton acceptors. Those preferably utilizable are alkali metal and alkaline earth metal oxides, hydroxides and carbonates, such as sodium
20 hydroxide, potassium hydroxide, magnesium oxide, calcium oxide, sodium carbonate, potassium carbonate and calcium carbonate, each of which can also be employed in the presence of phase-transfer catalysts, e.g. triethylbenzylammonium chloride, tetrabutylammonium bromide, Adogen 464 (= methyltrialkyl(C₈-C₁₀)ammonium chloride) or TDA 1 (= tris-(methoxyethoxyethyl)-amine). Alkali metals such as sodium
25 or potassium can furthermore be used. Alkali metal and alkaline earth metal amides and hydrides, such as sodium amide, sodium hydride and calcium hydride, and additionally also alkali metal alkoxides, such as sodium methoxide, sodium ethoxide and potassium tert-butoxide can additionally also be employed.

Le A 31 064 - Foreign Countries

- 112 -

When carrying out process (B) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the reaction is carried out at temperatures between 0°C and 250°C, preferably between 50°C and 150°C.

Process (B) according to the invention is in general carried out under normal pressure.

- 5 When carrying out process (B) according to the invention, the reaction components of the formula (III) and the deprotonating bases are in general employed in approximately equimolar amounts. However, it is also possible to use one component or the other in a relatively large excess (up to 3 mol).

- 10 Process (C) is characterized in that compounds of the formula (IV), in which A, B, W, X, Y, Z and R⁸ have the meaning given above, are cyclized intramolecularly in the presence of an acid and if appropriate in the presence of a diluent.

- 15 Diluents which can be employed in process (C) according to the invention are all organic solvents which are inert to the reaction participants. Those preferably utilizable are hydrocarbons, such as toluene and xylene, further halogenated hydrocarbons such as dichloromethane, chloroform, ethylene chloride, chlorobenzene, dichlorobenzene, additionally polar solvents, such as dimethyl sulphoxide, sulpholane, dimethylformamide and N-methyl-pyrrolidone. Alcohols such as methanol, ethanol, propanol, iso-propanol, butanol, isobutanol, tert-butanol can furthermore be employed.

The acid employed can optionally also be used as a diluent.

- 20 Acids which can be employed in process (C) according to the invention are all customary inorganic and organic acids, e.g. hydrohalic acids, sulphuric acid, alkyl-, aryl- and haloalkylsulphonic acids; halogenated alkylcarboxylic acids, e.g. trifluoroacetic acid, are used in particular.

- 25 When carrying out process (C) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the reaction is carried out at

Le A 31 064 - Foreign Countries

- 113 -

temperatures between 0°C and 250°C, preferably between 50°C and 150°C.

Process (C) according to the invention is in general carried out under normal pressure.

When carrying out process (C) according to the invention, the reaction components of the formula (IV) and the acid are employed, for example, in equimolar amounts.
5 However, it is optionally also possible to employ the acid in catalytic amounts.

Process (E) according to the invention is characterized in that carbonyl compounds of the formula (VIII) or their silyl enol ethers of the formula (VIIIa) are reacted with ketene acid halides of the formula (V), if appropriate in the presence of a diluent and if appropriate in the presence of an acid acceptor.

10 Diluents which can be employed in process (E) according to the invention are all organic solvents which are inert to the reaction participants. Those preferably utilizable are hydrocarbons, such as o-dichlorobenzene, tetralin, toluene and xylene, further ethers, such as dibutyl ether, glycol dimethyl ether and diglycol dimethyl ether, and additionally polar solvents, such as dimethyl sulphoxide, sulpholane,
15 dimethylformamide or N-methyl-pyrrolidone.

Acid acceptors which can be used when carrying out process (E) according to the invention are all customary acid acceptors.

Those preferably utilizable are tertiary amines, such as triethylamine, pyridine, diazabicyclooctane (DABCO), diazabicycloundecane (DBU), diazabicyclononene
20 (DBN), Hünig base or N,N-dimethyl-aniline.

When carrying out process (E) according to the invention, the reaction temperature can be varied within a relatively wide range. The reaction is expediently carried out at temperatures between 0°C and 250°C, preferably between 50°C and 220°C.

Process (E) according to the invention is preferably carried out under normal pressure.

Le A 31 064 - Foreign Countries

- 114 -

When carrying out process (E) according to the invention, the reaction components of the formulae (VIII) and (V) and, if appropriate, the acid acceptor are in general employed in approximately equimolar amounts. However, it is also possible to use one component or the other in a relatively large excess (up to 5 mol).

- 5 Process (F) according to the invention is characterized in that thioamides of the formula (IX) are reacted with ketene acid halides of the formula (V), if appropriate in the presence of a diluent and if appropriate in the presence of an acid acceptor.

Diluents which can be employed in process variant (F) according to the invention are all inert organic solvents. Those preferably utilizable are hydrocarbons, such as
10 o-dichlorobenzene, tetralin, toluene and xylene, further ethers, such as dibutyl ether, glycol dimethyl ether and diglycol dimethyl ether, and additionally polar solvents, such as dimethyl sulphoxide, sulpholane, dimethylformamide and N-methyl-pyrrolidone.

Acid acceptors which can be used in carrying out process (F) according to the invention are all customary acid acceptors.

- 15 Those preferably utilizable are tertiary amines, such as triethylamine, pyridine, diazabicyclooctane (DABCO), diazabicycloundecane (DBU), diazabicyclononene (DBN), Hünig base and N,N-dimethyl-aniline.

When carrying out process (F) according to the invention, the reaction temperature can be varied within a relatively wide range. The reaction is expediently carried out at
20 temperatures between 0°C and 250°C, preferably between 20°C and 220°C.

Process (F) according to the invention is expediently carried out under normal pressure.

When carrying out process (F) according to the invention, the reaction components of the formulae (IX) and (V) and, if appropriate, the acid acceptors are in general employed in approximately equimolar amounts. However, it is also possible to use one
25 component or the other in a relatively large excess (up to 5 mol).

Le A 31 064 - Foreign Countries

- 115 -

Process (G α) is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are in each case reacted with carboxylic acid halides of the formula (X), if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent.

Diluents which can be employed in process (G α) according to the invention are all solvents inert to the acid halides. Those preferably utilizable are hydrocarbons, such as benzine, benzene, toluene, xylene and tetralin, further halogenohydrocarbons, such as methylene chloride, chloroform, carbon tetrachloride, chlorobenzene and o-dichlorobenzene, additionally ketones, such as acetone and methyl isopropyl ketone, furthermore ethers, such as diethyl ether, tetrahydrofuran and dioxane, moreover carboxylic acid esters, such as ethyl acetate, nitriles such as acetonitrile and also strongly polar solvents, such as dimethylformamide, dimethyl sulphoxide and sulpholane. If the stability to hydrolysis of the acid halide permits, the reaction can also be carried out in the presence of water.

In the reaction by process (G α) according to the invention, suitable acid-binding agents are all customary acid acceptors. Those preferably utilizable are tertiary amines, such as triethylamine, pyridine, diazabicyclooctane (DABCO), diazabicycloundecene (DBU), diazabicyclononene (DBN), Hünig base and N,N-dimethyl-aniline, further alkaline earth metal alkoxides, such as magnesium and calcium oxide, additionally alkali metal and alkaline earth metal carbonates, such as sodium carbonate, potassium carbonate and calcium carbonate and also alkali metal hydroxides such as sodium hydroxide and potassium hydroxide.

In process (G α) according to the invention, the reaction temperature can be varied within a relatively wide range. In general, the reaction is carried out at temperatures between -20°C and +150°C, preferably between 0°C and 100°C.

When carrying out process (G α) according to the invention, the starting substances of the formulae (I-1-a) to (I-6-a) and the carboxylic acid halide of the formula (X) are in general each used in approximately equivalent amounts. However, it is also possible to employ the carboxylic acid halide in a relatively large excess (up to 5 mol). Working-

Le A 31 064 - Foreign Countries

- 116 -

up is carried out according to customary methods.

Process (G β) is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted with carboxylic anhydrides of the formula (XI), if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent.

- 5 Diluents which can be used in process (G β) according to the invention are preferably those diluents which are also suitable when using acid halides. Otherwise, a carboxylic anhydride employed in an excess can also simultaneously function as diluent.

Possible acid-binding agents optionally added in process (G β) are preferably those acid-binding agents which are also suitable when using acid halides.

- 10 The reaction temperature in process (G β) according to the invention can be varied within a relatively wide range. In general, the reaction is carried out at temperatures between -20°C and +150°C, preferably between 0°C and 100°C.

- 15 When carrying out process (G β) according to the invention, the starting substances of the formulae (I-1-a) to (I-6-a) and the carboxylic anhydride of the formula (XI) are in general used in approximately equivalent amounts in each case. However, it is also possible to employ the carboxylic anhydride in a relatively large excess (up to 5 mol). Working-up is carried out according to customary methods.

- 20 In general, a procedure is used in which diluent and carboxylic anhydride present in excess and also the resulting carboxylic acid are removed by distillation or by washing with an organic solvent or with water.

Process (H) is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted with chloroformic acid esters or chloroformic acid thioesters of the formula (XII), if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent.

Le A 31 064 - Foreign Countries

- 117 -

Possible acid-binding agents in process (H) according to the invention are all customary acid acceptors. Those preferably utilizable are tertiary amines, such as triethylamine, pyridine, DABCO, DBU, DBA, Hünig base and N,N-dimethyl-aniline, further alkaline earth metal oxides, such as magnesium and calcium oxide, additionally alkali metal and
5 alkaline earth metal carbonates, such as sodium carbonate, potassium carbonate and calcium carbonate and also alkali metal hydroxides such as sodium hydroxide and potassium hydroxide.

Diluents which can be employed in process (H) according to the invention are all solvents which are inert to the chloroformic acid esters or chloroformic acid thioesters.
10 Those preferably utilizable are hydrocarbons, such as benzene, toluene, xylene and tetralin, further halogenohydrocarbons, such as methylene chloride, chloroform, carbon tetrachloride, chlorobenzene and o-dichlorobenzene, additionally ketones, such as acetone and methyl isopropyl ketone, furthermore ethers, such as diethyl ether, tetrahydrofuran and dioxane, nitriles such as acetonitrile, moreover carboxylic acid
15 esters, such as ethyl acetate, and also strongly polar solvents, such as dimethylformamide, dimethyl sulphoxide and sulpholane.

When carrying out process (H) according to the invention, the reaction temperature can be varied within a relatively wide range. The reaction temperature is in general between -20°C and +100°C, preferably between 0°C and 50°C.

20 Process (H) according to the invention is in general carried out under normal pressure.

When carrying out process (H) according to the invention, the starting substances of the formulae (I-1-a) to (I-6-a) and the appropriate chloroformic acid esters or chloroformic acid thioesters of the formula (XII) are in general each used in approximately equivalent amounts. However, it is also possible to employ one component or the other
25 in a relatively large excess (up to 2 mol). Working-up is carried out according to customary methods. In general, a procedure is used in which salts which are deposited are removed and the reaction mixture which remains is concentrated by stripping off the diluent.

Le A 31 064 - Foreign Countries

- 118 -

Process (I) according to the invention is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted with (I α) compounds of the formula (XIII) in the presence of a diluent and if appropriate in the presence of an acid-binding agent or (I β) carbon disulphide and then with alkyl halides of the formula (XIV), if
5 appropriate in the presence of a diluent and if appropriate in the presence of a base.

In preparation process (I α), about 1 mol of chloromonothioformic acid ester or chlorodithioformic acid ester of the formula (XIII) is reacted at 0 to 120°C, preferably at 20 to 60°C, per mole of starting compound of the formulae (I-1-a) to (I-6-a).

Possible diluents optionally added are all inert polar organic solvents, such as ethers,
10 amides, carboxylic acid esters, nitriles, sulphones, sulphoxides, but also halogenoalkanes.

Dimethyl sulphoxide, ethyl acetate, acetonitrile, tetrahydrofuran, dimethylformamide or methylene chloride is preferably employed.

If, in a preferred embodiment, the enolate salt of the compounds (I-1-a) to (I-6-a) is
15 prepared by addition of strong deprotonating agents, e.g. sodium hydride or potassium tertiary butoxide, the further addition of acid-binding agents can be dispensed with.

If acid-binding agents are employed, customary inorganic or organic bases are suitable; sodium hydroxide, sodium carbonate, potassium carbonate, pyridine and triethylamine may be mentioned by way of example.

20 The reaction can be carried out at normal pressure or at elevated pressure; it is preferably carried out at normal pressure. Working-up takes place according to customary methods.

Le A 31 064 - Foreign Countries

- 119 -

In preparation process (I β), the equimolar amount or an excess of carbon disulphide is in each case added per mole of starting compounds of the formulae (I-1-a) to (I-6-a). The reaction is in this case preferably carried out at temperatures from 0 to 50°C and in particular at 20 to 30°C.

5 Often it is expedient first to prepare the corresponding salt from the compounds of the formulae (I-1-a) to (I-6-a) by addition of a base (e.g. potassium tertiary butoxide or sodium hydride). The compounds (I-1-a) to (I-6-a) are each reacted with carbon disulphide until the formation of the intermediate compound is complete, e.g. after stirring at room temperature for several hours.

10 Bases which can be employed in process (I β) are all customary proton acceptors. Those preferably utilizable are alkali metal hydrides, alkali metal alkoxides, alkali metal or alkaline earth metal carbonates or hydrogen carbonates or nitrogen bases. Those which may be mentioned, for example, are sodium hydride, sodium methoxide, sodium hydroxide, calcium hydroxide, potassium carbonate, sodium hydrogen carbonate,
15 triethylamine, dibenzylamine, diisopropylamine, pyridine, quinoline, diazabicyclooctane (DABCO), diazabicyclononene (DBN) and diazabicycloundecene (DBU).

Diluents which can be used in this process are all customary solvents.

Those preferably utilizable are aromatic hydrocarbons such as benzene or toluene, alcohols such as methanol, ethanol, isopropanol or ethylene glycol, nitriles such as
20 acetonitrile, ethers such as tetrahydrofuran or dioxane, amides such as dimethylformamide or other polar solvents such as dimethyl sulphoxide or sulpholane.

Further reaction with the alkyl halide of the formula (XIV) is preferably carried out at 0 to 70°C and in particular at 20 to 50°C. In this case, at least the equimolar amount of alkyl halide is employed.

25 The reaction is carried out at normal pressure or at elevated pressure, preferably at normal pressure.

Le A 31 064 - Foreign Countries

- 120 -

Working-up is in turn carried out according to customary methods.

Process (J) according to the invention is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted with sulphonyl chlorides of the formula (XV), if appropriate in the presence of a diluent and if appropriate in the presence of
5 an acid-binding agent.

In preparation process (J), about 1 mol of sulphonyl chloride of the formula (XV) is reacted at -20 to 150°C, preferably at 20 to 70°C, per mole of starting compound of the formula (I-1-a) to (I-6-a).

Process (J) is preferably carried out in the presence of a diluent.

10 Possible diluents are all inert polar organic solvents, such as ethers, amides, nitriles, sulphones, sulphoxides or halogenated hydrocarbons such as methylene chloride.

Dimethyl sulphoxide, tetrahydrofuran, dimethylformamide and methylene chloride are preferably employed.

If, in a preferred embodiment, the enolate salt of the compounds (I-1-a) to (I-6-a) is
15 prepared by addition of strong deprotonating agents (e.g. sodium hydride or potassium tertiary butoxide), the further addition of acid-binding agents can be dispensed with.

If acid-binding agents are employed, customary inorganic or organic bases are suitable; those which may be mentioned by way of example are sodium hydroxide, sodium carbonate, potassium carbonate, pyridine and triethylamine.

20 The reaction can be carried out at normal pressure or at elevated pressure, preferably it is carried out at normal pressure. Working-up takes place according to customary methods.

Process (K) according to the invention is characterized in that compounds of the

Le A 31 064 - Foreign Countries

- 121 -

formulae (I-1-a) to (I-6-a) are each reacted with phosphorus compounds of the formula (XVI), if appropriate in the presence of a diluent and if appropriate in the presence of an acid-binding agent.

In preparation process (K), to obtain compounds of the formulae (I-1-e) to (I-6-e), 1 to 5 2, preferably 1 to 1.3, mol of the phosphorus compound of the formula (XVI) are reacted at temperatures between -40°C and 150°C, preferably between -10 and 110°C, relative to 1 mol of the compounds (I-1-a) to (I-6-a).

Process (K) is preferably carried out in the presence of a diluent.

Possible diluents are all inert, polar organic solvents, such as halogenohydrocarbons, 10 carboxylic acid esters, ethers, amides, nitriles, sulphones, sulphoxides etc.

Acetonitrile, dimethyl sulphoxide, tetrahydrofuran, dimethylformamide and methylene chloride are preferably employed.

Possible acid-binding agents optionally added are customary inorganic or organic bases such as hydroxides, carbonates or amines. Those which may be mentioned by way of 15 example are sodium hydroxide, sodium carbonate, potassium carbonate, pyridine and triethylamine.

The reaction can be carried out at normal pressure or at elevated pressure, preferably at normal pressure. Working-up takes place according to customary methods of organic chemistry. The final products are preferably purified by crystallization, chromatographic 20 purification or by so-called "incipient distillation", i.e. removal of the volatile constituents in vacuo.

Process (L) is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted with metal hydroxides or metal alkoxides of the formula (XVII) or amines of the formula (XVIII), if appropriate in the presence of a diluent.

Le A 31 064 - Foreign Countries

- 122 -

Diluents which can be employed in process (L) according to the invention are preferably ethers such as tetrahydrofuran, dioxane, diethyl ether or else alcohols such as methanol, ethanol, isopropanol, but also water. Process (L) according to the invention is in general carried out under normal pressure. The reaction temperature is
5 in general between -20°C and 100°C , preferably between 0°C and 50°C .

Process (M) according to the invention is characterized in that compounds of the formulae (I-1-a) to (I-6-a) are each reacted ($M\alpha$) with compounds of the formula (XIX), if appropriate in the presence of a diluent and if appropriate in the presence of a catalyst, or ($M\beta$) with compounds of the formula (XX), if appropriate in the presence
10 of a diluent and if appropriate in the presence of an acid-binding agent.

In preparation process ($M\alpha$), about 1 mol of isocyanate of the formula (XIX) is reacted at 0 to 100°C , preferably at 20 to 50°C , per mole of starting compound of the formulae (I-1-a) to (I-6-a).

Process ($M\alpha$) is preferably carried out in the presence of a diluent.

15 Possible diluents are all inert organic solvents, such as ethers, amides, nitriles, sulphones or sulphoxides.

Catalysts can optionally be added to accelerate the reaction. The catalysts employed can very advantageously be organotin compounds, e.g. dibutyltin dilaurate.

The reaction is preferably carried out at normal pressure.

20 In preparation process ($M\beta$), about 1 mol of carbamoyl chloride of the formula (XX) is reacted at 0 to 150°C , preferably at 20 to 70°C , per mole of starting compound of the formulae (I-1-a) to (I-6-a).

Possible diluents optionally added are all inert polar organic solvents, such as ethers, carboxylic acid esters, nitriles, amides, sulphones, sulphoxides or halogenated

Le A 31 064 - Foreign Countries

- 123 -

hydrocarbons.

Dimethyl sulphoxide, tetrahydrofuran, dimethylformamide and methylene chloride are preferably employed.

5 If, in a preferred embodiment, the enolate salt of the compound (I-1-a) to (I-6-a) is prepared by addition of strong deprotonating agents (e.g. sodium hydride or potassium tertiary butoxide), the further addition of acid-binding agents can be dispensed with.

If acid-binding agents are employed, customary inorganic or organic bases are suitable; those which may be mentioned by way of example are sodium hydroxide, sodium carbonate, potassium carbonate, triethylamine or pyridine.

10 The reaction can be carried out at normal pressure or at elevated pressure, preferably at normal pressure. Working-up takes place according to customary methods.

The active compounds are suitable for controlling animal pests, preferably arthropods and nematodes, in particular insects and arachnida, which are encountered in agriculture, in forestry, in the protection of stored products and of materials, and in the
15 hygiene field. They are active against normally sensitive and resistant species and against all or some stages of development. The abovementioned pests include:

From the order of the Isopoda, for example, *Oniscus asellus*, *Armadillidium vulgare* and *Porcellio scaber*.

From the order of the Diplopoda, for example, *Blaniulus guttulatus*.

20 From the order of the Chilopoda, for example, *Geophilus carpophagus* and *Scutigera spec.*

From the order of the Symphyla, for example, *Scutigera immaculata*.

From the order of the Thysanura, for example, *Lepisma saccharina*.

From the order of the Collembola, for example, *Onychiurus armatus*.

Le A 31 064 - Foreign Countries

- 124 -

From the order of the Orthoptera, for example, *Blatta orientalis*, *Periplaneta americana*, *Leucophaea maderae*, *Blattella germanica*, *Acheta domesticus*, *Gryllotalpa* spp., *Locusta migratoria migratorioides*, *Melanoplus differentialis* and *Schistocerca gregaria*.

From the order of the Dermaptera, for example, *Forficula auricularia*.

5 From the order of the Isoptera, for example, *Reticulitermes* spp..

From the order of the Anoplura, for example, *Phylloxera vastatrix*, *Pemphigus* spp., *Pediculus humanus corporis*, *Haematopinus* spp. and *Linognathus* spp.

From the order of the Mallophaga, for example, *Trichodectes* spp. and *Damalinea* spp.

10 From the order of the Thysanoptera, for example, *Hercinothrips femoralis* and *Thrips tabaci*.

From the order of the Heteroptera, for example, *Eurygaster* spp., *Dysdercus intermedius*, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus* and *Triatoma* spp.

15 From the order of the Homoptera, for example, *Aleurodes brassicae*, *Bemisia tabaci*, *Trialeurodes vaporariorum*, *Aphis gossypii*, *Brevicoryne brassicae*, *Cryptomyzus ribis*, *Aphis fabae*, *Doralis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*, *Macrosiphum avenae*, *Myzus* spp., *Phorodon humuli*, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelis bilobatus*, *Nephotettix cincticeps*, *Lecanium corni*, *Saissetia oleae*, *Laodelphax striatellus*, *Nilaparvata lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp. and *Psylla* spp.

20 From the order of the Lepidoptera, for example, *Pectinophora gossypiella*, *Bupalus piniarius*, *Cheimatobia brumata*, *Lithocolletis blancardella*, *Hyponomeuta padella*, *Plutella maculipennis*, *Malacosoma neustria*, *Euproctis chrysorrhoea*, *Lymantria* spp., *Bucculatrix thurberiella*, *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp., *Earias insulana*, *Heliothis* spp., *Spodoptera exigua*, *Mamestra brassicae*, *Panolis*
25 *flammea*, *Prodenia litura*, *Spodoptera* spp., *Trichoplusia ni*, *Carpocapsa pomonella*, *Pieris* spp., *Chilo* spp., *Pyrausta nubilalis*, *Ephestia kuehniella*, *Galleria mellonella*, *Tineola bisselliella*, *Tinea pellionella*, *Hofmannophila pseudospretella*, *Cacoecia podana*, *Capua reticulana*, *Choristoneura fumiferana*, *Clysia ambiguella*, *Homona magnanima*

Le A 31 064 - Foreign Countries

- 125 -

and *Tortrix viridana*.

From the order of the Coleoptera, for example, *Anobium punctatum*, *Rhizopertha dominica*, *Bruchidius obtectus*, *Acanthoscelides obtectus*, *Hylotrupes bajulus*, *Agelastica alni*, *Leptinotarsa decemlineata*, *Phaedon cochleariae*, *Diabrotica* spp., *Psylliodes chrysocephala*, *Epilachna varivestis*, *Atomaria* spp., *Oryzaephilus surinamensis*,
 5 *Anthonomus* spp., *Sitophilus* spp., *Otiorrhynchus sulcatus*, *Cosmopolites sordidus*, *Ceuthorrhynchus assimilis*, *Hypera postica*, *Dermestes* spp., *Trogoderma* spp., *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*, *Ptinus* spp., *Niptus hololeucus*, *Gibbium psylloides*, *Tribolium* spp., *Tenebrio molitor*, *Agriotes* spp.,
 10 *Conoderus* spp., *Melolontha melolontha*, *Amphimallon solstitialis* and *Costelytra zealandica*.

From the order of the Hymenoptera, for example, *Diprion* spp., *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis* and *Vespa* spp.

From the order of the Diptera, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp.,
 15 *Drosophila melanogaster*, *Musca* spp., *Fannia* spp., *Calliphora erythrocephala*, *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp., *Gastrophilus* spp., *Hyppobosca* spp., *Stomoxys* spp., *Oestrus* spp., *Hypoderma* spp., *Tabanus* spp., *Tannia* spp., *Bibio hortulanus*, *Oscinella frit*, *Phorbia* spp., *Pegomyia hyoscyami*, *Ceratitis capitata*, *Dacus oleae* and *Tipula paludosa*.

20 From the order of the Siphonaptera, for example, *Xenopsylla cheopis* and *Ceratophyllus* spp.

From the order of the Arachnida, for example, *Scorpio maurus* and *Latrodectus mactans*.

From the order of the Acarina, for example, *Acarus siro*, *Argas* spp., *Ornithodoros* spp.,
 25 *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptura oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes* spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp.

Le A 31 064 - Foreign Countries

- 126 -

and Tetranychus spp.

The active compounds according to the invention are distinguished by a high insecticidal and acaricidal activity.

They can be used to particularly good effect for controlling insects which are injurious
 5 to plants, such as, for example, against the larvae of the mustard beetle (*Phaedon cochleariae*) or against the larvae of the green rice leaf hopper (*Nephotettix cincticeps*) or against the caterpillars of the cabbage moth (*Plutella maculipennis*).

The active compounds according to the invention can furthermore be used as defoliant, desiccant, agents for destroying broad-leaved plants and, especially, as weed-killers.
 10 By weeds, in the broadest sense, there are to be understood all plants which grow in locations where they are undesired. Whether the substances according to the invention act as total or selective herbicides depends essentially on the amount used.

The dosages of the active compounds according to the invention necessary for controlling weeds are between 0.001 and 10 kg/ha, preferably between 0.005 and
 15 5 kg/ha.

The active compounds according to the invention can be used, for example, in connection with the following plants:

Dicotyledon weeds of the genera: Sinapis, Lepidium, Galium, Stellaria, Matricaria, Anthemis, Galinsoga, Chenopodium, Urtica, Senecio, Amaranthus, Portulaca, Xanthium,
 20 Convolvulus, Ipomoea, Polygonum, Sesbania, Ambrosia, Cirsium, Carduus, Sonchus, Solanum, Rorippa, Rotala, Lindernia, Lamium, Veronica, Abutilon, Emex, Datura, Viola, Galeopsis, Papaver, Centaurea, Trifolium, Ranunculus and Taraxacum.

Dicotyledon crops of the genera: Gossypium, Glycine, Beta, Daucus, Phaseolus, Pisum, Solanum, Linum, Ipomoea, Vicia, Nicotiana, Lycopersicon, Arachis, Brassica,
 25 Lactuca, Cucumis and Cucurbita.

Le A 31 064 - Foreign Countries

- 127 -

Monocotyledon weeds of the genera: Echinochloa, Setaria, Panicum, Digitaria, Phleum, Poa, Festuca, Eleusine, Brachiaria, Lolium, Bromus, Avena, Cyperus, Sorghum, Agropyron, Cynodon, Monochoria, Fimbristylis, Sagittaria, Eleocharis, Scirpus, Paspalum, Ischaemum, Sphenoclea, Dactyloctenium, Agrostis, Alopecurus and
5 Apera.

Monocotyledon crops of the genera: Oryza, Zea, Triticum, Hordeum, Avena, Secale, Sorghum, Panicum, Saccharum, Ananas, Asparagus and Allium.

However, the use of the active compounds according to the invention is in no way restricted to these genera, but also extends in the same manner to other plants.

10 The compounds are suitable, depending on the concentration, for the total control of weeds, for example on industrial terrain and rail tracks, and on paths and squares with or without tree plantings. Equally, the compounds can be employed for controlling weeds in perennial cultures, for example afforestations, decorative tree plantings, orchards, vineyards, citrus groves, nut orchards, banana plantations, coffee plantations,
15 tea plantations, rubber plantations, oil palm plantations, cocoa plantations, soft fruit plantings and hopfields, on lawns, turf and pasture-land, and for the selective combating of weeds in annual cultures.

The active compounds according to the invention are very highly suitable for the selective control of monocotyledon weeds in dicotyledon crops pre- and post-
20 emergence. They can be employed to very good effect for the control of grass weeds, for example in cotton or sugar beet.

The active compounds can be converted into the customary formulations, such as solutions, emulsions, wettable powders, suspensions, powders, dusting agents, pastes, soluble powders, granules, suspension-emulsion concentrates, natural and synthetic
25 materials impregnated with active compound, and very fine capsules in polymeric substances.

Le A 31 064 - Foreign Countries

- 128 -

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is liquid solvents and/or solid carriers, optionally with the use of surface-active agents, that is emulsifying agents and/or dispersing agents and/or foam-forming agents.

5 In the case of the use of water as an extender, organic solvents can, for example, also be used as auxiliary solvents. As liquid solvents, there are suitable in the main: aromatics, such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons, such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons, such as cyclohexane or paraffins, for
10 example petroleum fractions, mineral and vegetable oils, alcohols, such as butanol or glycol as well as their ethers and esters, ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents, such as dimethylformamide and dimethyl sulphoxide, as well as water.

As solid carriers there are suitable:

15 for example ammonium salts and ground natural minerals, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as highly disperse silica, alumina and silicates, as solid carriers for granules there are suitable: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, as well as synthetic granules of
20 inorganic and organic meals, and granules of organic material such as sawdust, coconut shells, maize cobs and tobacco stalks; as emulsifying and/or foam-forming agents there are suitable: for example non-ionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates as well as albumen hydrolysis
25 products; as dispersing agents there are suitable: for example lignin-sulphite waste liquors and methylcellulose.

Adhesives such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latexes, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, as well as natural phospholipids, such as cephalins and lecithins, and

Le A 31 064 - Foreign Countries

- 129 -

synthetic phospholipids, can be used in the formulations. Further additives can be mineral and vegetable oils.

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs, such as alizarin dyestuffs, azo
5 dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

The formulations in general contain between 0.1 and 95 per cent by weight of active compound, preferably between 0.5 and 90%.

The active compound according to the invention can be present in its commercially
10 available formulations and in the use forms prepared from these formulations, as a mixture with other active compounds, such as insecticides, attractants, sterilizing agents, acaricides, nematocides, fungicides, growth-regulating substances or herbicides. The insecticides include, for example, phosphates, carbamates, carboxylates, chlorinated hydrocarbons, phenylureas and substances produced by microorganisms.

15 Examples of particularly advantageous mixture components are the following compounds:

Fungicides:

2-aminobutane; 2-anilino-4-methyl-6-cyclopropyl-pyrimidine; 2',6'-dibromo-2-methyl-4'-trifluoromethoxy-4'-trifluoro-methyl-1,3-thiazole-5-carboxanilide; 2,6-dichloro-N-
20 (4-trifluoromethylbenzyl)benzamide; (E)-2-methoxyimino-N-methyl-2-(2-phenoxy-phenyl)-acetamide; 8-hydroxyquinoline sulphate; methyl (E)-2-{2-[6-(2-cyanophenoxy)-pyrimidin-4-yloxy]-phenyl}-3-methoxyacrylate; methyl (E)-methoximino[alpha-(o-tolyloxy)-o-tolyl]acetate; 2-phenylphenol (OPP), aldimorph, ampropylfos, anilazine, azaconazole,
25 benalaxyl, benodanil, benomyl, binapacryl, biphenyl, bitertanol, blasticidin-S, bromuconazole, bupirimate, buthiobate, calcium polysulphide, captafol, captan, carbendazim, carboxin, quinomethionate,

Le A 31 064 - Foreign Countries

- 130 -

- chloroneb, chloropicrin, chlorothalonil, chlozolate, cufraneb, cymoxanil, cypro-
conazole, cyprofuram,
dichlorophen, diclobutrazol, diclofluanid, diclomezin, dicloran, diethofencarb,
difenoconazole, dimethirimol, dimethomorph, diniconazole, dinocap, diphenylamine,
5 dipyrithion, ditalimfos, dithianon, dodine, drazoxolon,
edifenphos, epoxyconazole, ethirimol, etridiazole,
fenarimol, fenbuconazole, fenfuram, fenitropan, fenciclonil, fenpropidin, fenpropimorph,
fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, fluoromide,
fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, fosetyl-aluminum,
10 fthalide, fuberidazole, furalaxyl, furmecyclox,
guazatine,
hexachlorobenzene, hexaconazole, hymexazol,
imazalil, imibenconazole, iminoctadine, iprobenfos (IBP), iprodione, isoprothiolane,
kasugamycin, copper preparations such as: copper hydroxide, copper naphthenate,
15 copper oxychloride, copper sulphate, copper oxide, oxine-copper and Bordeaux mixture,
mancopper, mancozeb, maneb, mepanipyrim, mepronil, metalaxyl, metconazole,
methasulfocarb, methfuroxam, metiram, metsulfovax, myclobutanil,
nickel dimethyldithiocarbamate, nitrothal-isopropyl, nuarimol,
ofurace, oxadixyl, oxamocarb, oxycarboxin,
20 pefurazoate, penconazole, pencycuron, phosdiphen, phthalide, pimaricin, piperalin,
polycarbamate, polyoxin, probenazole, prochloraz, procymidone, propamocarb,
propiconazole, propineb, pyrazophos, pyrifenox, pyrimethanil, pyroquilon,
quintozene (PCNB),
sulfur and sulfur preparations,
25 tebuconazole, tecloftalam, tecnazene, tetraconazole, thiabendazole, thicyofen,
thiophanate-methyl, thiram, tolclophos-methyl, tolylfluanid, triadimefon, triadimenol,
triazoxide, trichlamide, tricyclazole, tridemorph, triflumizole, triforine, triticonazole,
validamycin A, vinclozolin,
zineb, ziram
30 **Bactericides:**
bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin,

Le A 31 064 - Foreign Countries

- 131 -

octhilinone, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper sulphate and other copper preparations.

Insecticides/Acaricides/Nematicides:

abamectin, AC 303 630, acephate, acrinathrin, alanycarb, aldicarb, alphamethrin,
 5 amitraz, avermectin, AZ 60541, azadirachtin, azinphos A, azinphos M, azocyclotin,
 Bacillus thuringiensis, bendiocarb, benfuracarb, bensultap, beta-cyfluthrin, bifenthrin,
 BPMC, brofenprox, bromophos A, bufencarb, buprofezin, butocarboxin, butylpyridaben,
 cadusafos, carbaryl, carbofuran, carbophenothion, carbosulfan, cartap, CGA 157 419,
 CGA 184699, chloethocarb, chlorethoxyfos, chlorfenvinphos, chlorfluazuron,
 10 chlormephos, chlorpyrifos, chlorpyrifos M, cis-resmethrin, clocythrin, clofentezine,
 cyanophos, cycloprothrin, cyfluthrin, cyhalothrin, cyhexatin, cypermethrin, cyromazine,
 deltamethrin, demeton-M, demeton-S, demeton-S-methyl, diafenthiuron, diazinon,
 dichlofenthion, dichlorvos, dicliphos, dicrotophos, diethion, diflubenzuron, dimethoate,
 dimethylvinphos, dioxathion, disulfoton,
 15 edifenphos, emamectin, esfenvalerate, ethiofencarb, ethion, ethofenprox, ethoprophos,
 etrimphos,
 fenamiphos, fenazaquin, fenbutatin oxide, fenitrothion, fenobucarb, fenothiocarb,
 fenoxycarb, fenpropathrin, fenpyrad, fenpyroximate, fenthion, fenvalerate, fipronil,
 fluazinam, flucycloxuron, flucythrinate, flufenoxuron, flufenprox, fluvalinate, fonophos,
 20 formothion, fosthiazate, fubfenprox, furathiocarb,
 HCH, heptenophos, hexaflumuron, hexythiazox,
 imidacloprid, iprobenfos, isazophos, isofenphos, isoprocarb, isoxathion, ivermectin,
 lambda-cyhalothrin, lufenuron,
 malathion, mecarbam, mevinphos, mesulfenphos, metaldehyde, methacrifos,
 25 methamidophos, methidathion, methiocarb, methomyl, metolcarb, milbemectin,
 monocrotophos, moxidectin,
 naled, NC 184, NI 25, nitenpyram,
 omethoate, oxamyl, oxydemethon M, oxydeprofos,
 parathion A, parathion M, permethrin, phenthoate, phorate, phosalone, phosmet,
 30 phosphamidon, phoxim, pirimicarb, pirimiphos M, pirimiphos A, profenofos,
 promecarb, propaphos, propoxur, prothiofos, prothoate, pymetrozin, pyrachlophos,

Le A 31 064 - Foreign Countries

- 132 -

pyridaphenthion, pyresmethrin, pyrethrum, pyridaben, pyrimidifen, pyriproxifen,
 quinalphos,

RH 5992,

salithion, sebufos, silafluofen, sulfotep, sulprofos,

- 5 tebufenozid, tebufenpyrad, tebupirimiphos, teflubenzuron, tefluthrin, temephos, terbam,
 terbufos, tetrachlorvinphos, thiafenox, thiodicarb, thiofanox, thiomethon, thionazin,
 thuringiensin, tralomethrin, triarathen, triazophos, triazuron, trichlorfon, triflumuron,
 trimethacarb,
 vamidothion, XMC, xylylcarb, YI 5301/5302, zetamethrin.

10 **Herbicides:**

- for example anilides such as, for example, diflufenican and propanil; arylcarboxylic
 acids such as, for example, dichloropicolinic acid, dicamba and picloram;
 aryloxyalkanoic acids such as, for example, 2,4-D, 2,4-DB, 2,4-DP, fluroxypyr, MCPA,
 MCPP and triclopyr; aryloxy-phenoxy-alkanoic esters such as, for example, diclofop-
 15 methyl, fenoxaprop-ethyl, fluazifop-butyl, haloxyfop-methyl and quizalofop-ethyl;
 azinones such as, for example, chloridazon and norflurazon; carbamates such as, for
 example, chlorpropham, desmedipham, phenmedipham and propham; chloroacetanilides
 such as, for example, alachlor, acetochlor, butachlor, metazachlor, metolachlor,
 pretilachlor and propachlor; dinitroanilines such as, for example, oryzalin,
 20 pendimethalin and trifluralin; diphenyl ethers such as, for example, acifluorfen, bifenox,
 fluoroglycofen, fomesafen, halosafen, lactofen and oxyfluorfen; ureas such as, for
 example, chlortoluron, diuron, fluometuron, isoproturon, linuron and methabenz-
 thiazuron; hydroxylamines such as, for example, alloxydim, clethodim, cycloxydim,
 sethoxydim and tralkoxydim; imidazolinones such as, for example, imazethapyr, imaza-
 25 methabenz, imazapyr and imazaquin; nitriles such as, for example, bromoxynil,
 dichlobenil and ioxynil; oxyacetamides such as, for example, mefenacet; sulfonylureas
 such as, for example, amidosulfuron, bensulfuron-methyl, chlorimuron-ethyl,
 chlorsulfuron, cinosulfuron, metsulfuron-methyl, nicosulfuron, primisulfuron,
 pyrazosulfuron-ethyl, thifensulfuron-methyl, triasulfuron and tribenuron-methyl;
 30 thiocarbamates such as, for example, butylate, cycloate, di-allate, EPTC, esprocarb,
 molinate, prosulfocarb, thiobencarb and tri-allate; triazines such as, for example,

Le A 31 064 - Foreign Countries

- 133 -

atrazine, cyanazine, simazine, simetryn, terbutryn and terbutylazine; triazinones such as, for example, hexazinone, metamilon and metribuzin; others such as, for example, aminotriazole, benfuresate, bentazone, cinmethylin, clomazone, clopyralid, difenzoquat, dithiopyr, ethofumesate, fluorochloridone, glufosinate, glyphosate, isoxaben, pyridate,
5 quinchlorac, quinmerac, sulphosate and tridiphane.

The active compound according to the invention can furthermore be present in its commercially available formulations and in the use forms prepared from these formulations, as a mixture with synergistic agents. Synergistic agents are compounds which increase the action of the active compounds, without it being necessary for the
10 synergistic agent added to be active itself.

The active compound content of the use forms prepared from the commercially available formulations can vary within wide limits. The active compound concentration of the use forms can be from 0.0000001 to 95% by weight of active compound, preferably between 0.0001 and 1% by weight.

15 The compounds are employed in a customary manner appropriate for the use forms.

When used against hygiene pests and pests of stored products, the active compounds are distinguished by an excellent residual action on wood and clay as well as a good stability to alkali on limed substrates.

The active compounds according to the invention are not only active against plant,
20 hygiene and stored product pests, but also, in the veterinary medicine sector, against animal parasites (ectoparasites), such as scaly ticks, Argasidae, scab mites, Trombidae, flies (stinging and sucking), parasitic fly larvae, lice, hair lice, bird lice and fleas. These parasites include:

From the order of the Anoplurida, for example, Haematopinus spp., Linognathus spp.,
25 Pediculus spp., Phtirus spp., Solenopotes spp..

Le A 31 064 - Foreign Countries

- 134 -

From the order of the Mallophagida and the suborders Amblycerina and Ischnocerina, for example, Trimenopon spp., Menopon spp., Trinoton spp., Bovicola spp., Werneckiella spp., Lepikentron spp., Damalina spp., Trichodectes spp., Felicola spp..

From the order Diptera and the suborders Nematocerina and Brachycerina, for example,
 5 Aedes spp., Anopheles spp., Culex spp., Simulium spp., Eusimulium spp., Phlebotomus
 spp., Lutzomyia spp., Culicoides spp., Chrysops spp., Hybomitra spp., Atylotus spp.,
 Tabanus spp., Haematopota spp., Philipomyia spp., Braula spp., Musca spp., Hydrotaea
 spp., Stomoxys spp., Haematobia spp., Morellia spp., Fannia spp., Glossina spp.,
 Calliphora spp., Lucilia spp., Chrysomyia spp., Wohlfahrtia spp., Sarcophaga spp.,
 10 Oestrus spp., Hypoderma spp., Gasterophilus spp., Hippobosca spp., Lipoptena spp.,
 Melophagus spp..

From the order of the Siphonapterida, for example, Pulex spp., Ctenocephalides spp., Xenopsylla spp., Ceratophyllus spp..

From the order of the Heteropterida, for example, Cimex spp., Triatoma spp., Rhodnius
 15 spp., Panstrongylus spp..

From the order of the Blattarida, for example, Blatta orientalis, Periplaneta americana, Blattella germanica, Supella spp..

From the subclass of the Acaria (Acarida) and the orders of the Meta- and Mesostigmata, for example, Argas spp., Ornithodoros spp., Otabius spp., Ixodes spp.,
 20 Amblyomma spp., Boophilus spp., Dermacentor spp., Haemaphysalis spp., Hyalomma
 spp., Rhipicephalus spp., Dermanyssus spp., Raillietia spp., Pneumonyssus spp.,
 Sternostoma spp., Varroa spp..

From the order of the Actinedida (Prostigmata) and Acaridida (Astigmata), for example,
 Acarapis spp., Cheyletiella spp., Ornithocheyletia spp., Myobia spp., Psorergates spp.,
 25 Demodex spp., Trombicula spp., Listrophorus spp., Acarus spp., Tyrophagus spp.,
 Caloglyphus spp., Hypodectes spp., Pterolichus spp., Psoroptes spp., Chorioptes spp.,

Le A 31 064 - Foreign Countries

- 135 -

Otodectes spp., Sarcoptes spp., Notoedres spp., Knemidocoptes spp., Cytodites spp., Laminosioptes spp..

For example, they exhibit an outstanding activity against *Boophilus microplus* and *Lucilia cuprina*.

5 The active compounds of the formula (I) according to the invention are also suitable for controlling arthropods which infest agricultural productive livestock such as, for example, cattle, sheep, goats, horses, pigs, donkeys, camels, buffalo, rabbits, chickens, turkeys, ducks, geese and bees, other pets, such as, for example, dogs, cats, cage birds and aquarium fish, and also so-called test animals, such as, for example, hamsters,
10 guinea pigs, rats and mice. By controlling these arthropods cases of death and reductions in productivity (for meat, milk, wool, hides, eggs, honey etc) should be diminished, so that more economic and simpler animal husbandry is possible by use of the active compounds according to the invention.

The active compounds according to the invention are used in the veterinary sector in
15 a known manner by enteral administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boli, the feed-through process and suppositories, by parenteral administration, such as, for example, by injections (intramuscular, subcutaneous, intravenous, intraperitoneal etc), implants, by nasal administration, by dermal use in the form, for example, of dipping or bathing, spraying,
20 pouring on and spotting on, washing and powdering, and also with the aid of moulded articles containing the active compound, such as collars, ear marks, tail marks, limb bands, halters, marking devices, etc.

When used for livestock, poultry, pets etc., the active compounds of the formula (I) can be applied as formulations (for example powders, emulsions, fluid compositions) which
25 contain the active compounds in an amount from 1 to 80% by weight, directly or after 100 to 10,000-fold dilution or they can be used as a chemical bath.

It was additionally found that the compounds of the formula I according to the

Le A 31 064 - Foreign Countries

- 136 -

invention exhibit a high insecticidal action against insects which destroy industrial materials.

The following insects may be mentioned by way of example and preferably - but without being limitative:

- 5 Beetles such as
 Hylotrupes bajulus, Chlorophorus pilosis, Anobium punctatum, Xestobium rufovillosum, Ptilinus pecticornis, Dendrobium pertinex, Ernobius mollis, Priobium carpini, Lyctus brunneus, Lyctus africanus, Lyctus planicollis, Lyctus linearis, Lyctus pubescens, Trogoxylon aequale, Minthes rugicollis, Xyleborus spec. Tryptodendron
 10 spec. Apate monachus, Bostrychus capucins, Heterobostrychus brunneus, Sinoxylon spec. Dinoderus minutus.

Hymenopterans such as

Sirex juvencus, Urocerus gigas, Urocerus gigas taignus, Urocerus augur.

Termites such as

- 15 Kaloterme flavicollis, Cryptoterme brevis, Heteroterme indicola, Reticuliterme flavipes, Reticuliterme santonensis, Reticuliterme lucifugus, Mastoterme darwiniensis, Zootermopsis nevadensis, Coptoterme formosanus.

Brushtails

such as Lepisma saccharina.

- 20 Industrial materials in the present connection are to be understood as meaning non-living materials, preferably such as plastics, adhesives, sizes, paper and card, leather, wood and wood-processing products and paints.

Very particularly preferably, the materials to be protected from insect attack are wood and wood-processing products.

Le A 31 064 - Foreign Countries

- 137 -

Wood and wood-processing products which can be protected by the agent according to the invention or mixtures containing the latter are, for example, to be understood as meaning: construction timber, wooden beams, railway sleepers, bridge components, jetties, vehicles made of wood, boxes, palettes, containers, telegraph poles, wood
5 lagging, windows and doors made of wood, plywood, particle board, joiner's work or wood products which, quite generally, are used in building or joinery.

The active compounds can be used as such, in the form of concentrates or generally customary formulations such as powders, granules, solutions, suspensions, emulsions or pastes.

10 The formulations mentioned can be prepared in a manner known per se, e.g. by mixing the active compounds with at least one solvent or diluent, emulsifier, dispersant and/or binder or fixative, water-repellent, if appropriate siccatives and UV stabilizers and if appropriate colorants and pigments as well as further processing aids.

15 The insecticidal compositions or concentrates used for the protection of wood and wood materials contain the active compound according to the invention in a concentration from 0.0001 to 95% by weight, in particular 0.001 to 60% by weight.

The amount of the compositions or concentrates employed is dependent on the nature and the occurrence of the insects and on the medium. In use, the optimum amount to employ can in each case be determined by test series. In general, however, it is
20 sufficient to employ 0.0001 to 20% by weight, preferably 0.001 to 10% by weight, of the active compound, based on the material to be protected.

The solvent and/or diluent used is an organic chemical solvent or solvent mixture and/or an oily or oleaginous poorly volatile organic chemical solvent or solvent mixture and/or a polar organic chemical solvent or solvent mixture and/or water and, if
25 appropriate, an emulsifier and/or wetting agent.

Organic chemical solvents employed are preferably oily or oleaginous solvents having

Le A 31 064 - Foreign Countries

- 138 -

an evaporation number of over 35 and a flash point of above 30°C, preferably above 45°C. Poorly volatile, water-insoluble, oily and oleaginous solvents of this type used are appropriate mineral oils or their aromatic fractions or mineral oil-containing solvent mixtures, preferably white spirit, petroleum and/or alkylbenzene.

- 5 Advantageously, mineral oils having a boiling range from 170 to 220°C, white spirit having a boiling range from 170 to 220°C, spindle oil having a boiling range from 250 to 350°C, petroleum or aromatics of boiling range from 160 to 280°C, turpentine and the like are used.

10 In a preferred embodiment, liquid aliphatic hydrocarbons having a boiling range from 180 to 210°C or high-boiling mixtures of aromatic and aliphatic hydrocarbons having a boiling range from 180 to 220°C and/or spindle oil and/or monochloronaphthalene, preferably α -monochloronaphthalene, are used.

15 The organic poorly volatile oily or oleaginous solvents having an evaporation number of over 35 and a flash point of above 30°C, preferably above 45°C, can be partially replaced by readily or moderately volatile organic chemical solvents, with the proviso that the solvent mixture also has an evaporation number of over 35 and a flash point of above 30°C, preferably above 45°C, and that the insecticide-fungicide mixture is soluble or emulsifiable in this solvent mixture.

20 According to a preferred embodiment, a part of the organic chemical solvent or solvent mixture or an aliphatic polar organic chemical solvent or solvent mixture is replaced. Preferably, aliphatic organic chemical solvents containing hydroxyl and/or ester and/or ether groups, such as, for example, glycol ethers, esters or the like are used.

25 Organic chemical binders used in the context of the present invention are the plastic resins and/or binding drying oils which are water-dilutable and/or soluble or dispersible or emulsifiable in the organic chemical solvents employed, in particular binders consisting of or comprising an acrylate resin, a vinyl resin, e.g. polyvinyl acetate, polyester resin, polycondensation or polyaddition resin, polyurethane resin, alkyd resin

Le A 31 064 - Foreign Countries

- 139 -

or modified alkyd resin, phenolic resin, hydrocarbon resin such as indene-coumarone resin, silicone resin, drying vegetable and/or drying oils and/or physically drying binders based on a natural and/or synthetic resin.

5 The synthetic resin used as a binder can be employed in the form of an emulsion, dispersion or solution. The binders used can also be bitumen or bituminous substances up to 10% by weight. Additionally, dyes, pigments, water-repellent agents, odour corrigents and inhibitors or anticorrosive agents and the like known per se can be employed.

10 According to the invention, at least one alkyd resin or modified alkyd resin and/or a drying vegetable oil is preferably contained in the composition or in the concentrate as organic chemical binder. According to the invention, alkyd resins having an oil content of more than 45% by weight, preferably 50 to 68% by weight, are preferably used.

15 The binder mentioned can be completely or partially replaced by a fixing agent (mixture) or a plasticizer (mixture). These additives are intended to prevent volatilization of the active compounds and crystallization or precipitation. They preferably replace 0.01 to 30% of the binder (based on 100% of the binder employed).

20 The plasticizers originate from the chemical classes of phthalic acid esters such as dibutyl, dioctyl or benzyl butyl phthalate, phosphoric acid esters such as tributyl phosphate, adipic acid esters such as di-(2-ethylhexyl) adipate, stearates such as butyl stearate or amyl stearate, oleates such as butyl oleate, glycerol ethers or higher molecular weight glycol ethers, glycerol esters and p-toluenesulphonic acid esters.

Fixing agents are based chemically on polyvinyl alkyl ethers such as, for example, polyvinyl methyl ether or ketones such as benzophenone, ethylenebenzophenone.

25 A possible solvent or diluent is, in particular, also water, if appropriate as a mixture with one or more of the abovementioned organic chemical solvents or diluents, emulsifiers and dispersants.

Le A 31 064 - Foreign Countries

- 140 -

Particularly effective wood protection is achieved by large-scale impregnation processes, e.g. vacuum, double vacuum or pressure processes.

The ready-to-use compositions can optionally also contain further insecticides and optionally also one or more fungicides.


- 5 Possible additional mixture components are preferably the insecticides and fungicides mentioned in WO 94/29 268. The compounds mentioned in this document are explicit constituents of the present application.

10 Very particularly preferred mixture components can be insecticides, such as chlorpyrifos, phoxim, silafluofin, alphamethrin, cyfluthrin, cypermethrin, deltamethrin, permethrin, imidacloprid, NI-25, flufenoxuron, hexaflumuron and triflumuron, as well as fungicides such as epoxyconazole, hexaconazole, azaconazole, propiconazole, tebuconazole, cyproconazole, metconazole, imazalil, dichlofluanid, tolylfluanid, 3-iodo-2-propinyl butylcarbamate, N-octyl-isothiazolin-3-one and 4,5-dichloro-N-octylisothiazolin-3-one.

- 15 The preparation and the use of the active compounds according to the invention follow from the examples below.

Le A 31 064 - Foreign Countries

- 141 -

Preparation examplesExample (I-1-a-1)

32.6 g of the compound according to Example (II-1), dissolved in 200 ml of absolute
5 toluene, are added dropwise at reflux temperature to 20.42 g (0.181 mol) of potassium
tert-butoxide in 70 ml of absolute tetrahydrofuran (THF) and the mixture is stirred at
this temperature for a further 1.5 hours.


For working-up, it is diluted with water, the phases are separated, the toluene phase is
extracted with water and the combined aqueous phases are acidified with conc. HCl.
10 The product is filtered off with suction, washed and dried and finally stirred in methyl
tert-butyl (MTB) ether/n-hexane, filtered off with suction and dried.

Yield: 20.6 g (68% of theory); m.p.: > 220°C.

The compounds of the formula (I-1-a) shown in Table 21 below were obtained
analogously to Example (I-1-a-1) or according to the general details for preparation.

Le A 31 064 - Foreign Countries

- 142 -

Table 21


| Ex. No. | X | Y | Z | B | A | D | M.p.: °C | Iso- mer |
|----------|-------------------------------|-----------------|-----------------|--|---------------------------------|---|-------------|-------------|
| I-1-a-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | 211 | β |
| I-1-a-3 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 201 | - |
| I-1-a-4 | CH ₃ | Br | CH ₃ | CH ₃ | i-C ₃ H ₇ | H | 183 | - |
| I-1-a-5 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-6 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | 196 | β |
| I-1-a-7 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | >220 | - |
| I-1-a-8 | CH ₃ | CH ₃ | Br | CH ₃ | i-C ₃ H ₇ | H | 194 | - |
| I-1-a-9 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-10 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | >220 | β |

Le A 31 064 - Foreign Countries

- 143 -

Table 21 - Continuation

| Ex. No. | X | Y | Z | B | A | D | M.p.: °C | Iso- mer |
|----------|-------------------------------|-----------------|-------------------------------|---|--|---------------------------------|-------------|-------------|
| I-1-a-11 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | 124 | β |
| I-1-a-12 | CH ₃ | CH ₃ | Br | CH ₃ | CH ₃ | H | >220 | - |
| I-1-a-13 | CH ₃ | CH ₃ | Br | H | -CH ₂ -CHCH ₃ -CHCH ₃ - | | >220 | cis |
| I-1-a-14 | CH ₃ | CH ₃ | Br | H | -CH ₂ -CHCH ₃ -CHCH ₃ - | | >220 | trans |
| I-1-a-15 | CH ₃ | CH ₃ | Br | H | -CH ₂ -S-CH ₂ -CH ₂ - | | >220 | - |
| I-1-a-16 | CH ₃ | CH ₃ | Br | H | -(CH ₂) ₄ - | | >220 | - |
| I-1-a-17 | CH ₃ | CH ₃ | Br | H | H | i-C ₃ H ₇ | >220 | - |
| I-1-a-18 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-19 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | >220 | - |
| I-1-a-20 | C ₂ H ₅ | Br | C ₂ H ₅ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-21 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-22 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-23 | CH ₃ | Cl | CH ₃ | CH ₃ | CH ₃ | H | >220 | - |
| I-1-a-24 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | >220 | β |
| I-1-a-25 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₃ -CHCH ₃ -(CH ₂) ₂ - | | H | 196 | β |
| I-1-a-26 | CH ₃ | CH ₃ | Cl | CH ₃ | CH ₃ | H | >220 | - |
| I-1-a-27 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | >220 | - |
| I-1-a-28 | CH ₃ | CH ₃ | Cl | CH ₃ | i-C ₃ H ₇ | H | >220 | - |

Example (I-1-b-1)

1.9 ml (18 mmol) of isobutyryl chloride, dissolved in 5 ml of absolute methylene chloride, are added dropwise at 0°C to 10°C to 4.37 g of the compound according to
 5 Example (I-1-a-1) in 70 ml of absolute methylene chloride and 2.52 ml (18 mmol) of triethylamine. The mixture is stirred at room temperature until the reaction is complete according to thin-layer chromatographic (TLC) checking.


For working-up, it is washed 2 times with 0.5 N NaOH, dried and evaporated. The crude product is recrystallized from MTB ether/n-hexane.

10 Yield: 1.70 g (32% of theory); m.p.: 208°C.



The examples of the formula (I-1-b) shown in Table 22 below were prepared analogously to Example (I-1-b-1) or according to the general details for preparation.

Le A 31 064 - Foreign Countries

- 145 -

Table 22

(I-1-b)

| Ex. No. | X | Y | Z | B | A | D | R ¹ | M.p.: °C | Iso-mer |
|----------|-----------------|-----------------|-----------------|--|---|---|---|----------|---------|
| I-1-b-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ - | 209 | β |
| I-1-b-3 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 212 | β |
| I-1-b-4 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | H ₃ C ₂ -O-CH ₂ - | >220 | β |
| I-1-b-5 | CH ₃ | Br | CH ₃ | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | CH ₃ - | 191 | β |
| I-1-b-6 | CH ₃ | Br | CH ₃ | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H |  | >220 | β |
| I-1-b-7 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | CH ₃ - | 178 | - |
| I-1-b-8 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 202 | - |
| I-1-b-9 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ - | 181 | β |
| I-1-b-10 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 208 | β |
| I-1-b-11 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₄ H ₉ - | 212 | β |
| I-1-b-12 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | t-C ₄ H ₉ -CH ₂ - | >220 | β |
| I-1-b-13 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | CH ₃ - | 196 | β |
| I-1-b-14 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | i-C ₃ H ₇ - | 207 | β |
| I-1-b-15 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | t-C ₄ H ₉ -CH ₂ - | >220 | β |
| I-1-b-16 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | H ₃ C ₂ -O-CH ₂ - | 216 | β |
| I-1-b-17 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H |  | >220 | β |
| I-1-b-18 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | (CH ₃) ₂ C=CH- | >220 | - |

Le A 31 064 - Foreign Countries


- 146 -

Example 22 - Continuation

| Ex. No. | X | Y | Z | B | A | D | R ¹ | M.p.: °C | Iso-mer |
|----------|-------------------------------|-----------------|-----------------|---|---------------------------------|---|--|----------|---------|
| I-1-b-19 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | C ₂ H ₅ -O-CH ₂ | 214 | β |
| I-1-b-20 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | t-C ₄ H ₉ - | >220 | β |
| I-1-b-21 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | (CH ₃) ₂ C=CH- | 193 | β |
| I-1-b-22 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ - | 222 | β |
| I-1-b-23 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 161 | β |
| I-1-b-24 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₄ H ₉ - | 171 | β |
| I-1-b-25 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | C ₂ H ₅ -O-CH ₂ - | 166 | β |
| I-1-b-26 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 211 | - |
| I-1-b-27 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | i-C ₄ H ₉ - | 205 | - |
| I-1-b-28 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | >220 | β |
| I-1-b-29 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 214 | β |
| I-1-b-30 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | C ₂ H ₅ -O-CH ₂ - | 168 | β |
| I-1-b-31 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | i-C ₄ H ₉ - | 190 | β |
| I-1-b-32 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | C ₂ H ₅ -O-CH ₂ - | 153 | β |
| I-1-b-33 | CH ₃ | CH ₃ | Cl | CH ₃ | CH ₃ | H | i-C ₃ H ₇ - | 182 | - |
| I-1-b-34 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | i-C ₃ H ₇ - | 183 | - |
| I-1-b-35 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | i-C ₄ H ₉ - | >220 | - |
| I-1-b-36 | CH ₃ | CH ₃ | Cl | CH ₃ | i-C ₃ H ₇ | H | i-C ₃ H ₇ - | oil | - |
| I-1-b-37 | CH ₃ | CH ₃ | Cl | CH ₃ | i-C ₃ H ₇ | H | i-C ₄ H ₉ - | oil | - |

Le A 31 064 - Foreign Countries

- 147 -

Example (I-1-c-1)

1.2 ml of ethyl chloroformate in 3 ml of absolute methylene chloride are added dropwise at 0°C to 10°C to 4.37 g of the compound according to Example (I-1-a-1) in
5 70 ml of absolute methylene chloride and 1.7 ml (12 mmol) of triethylamine. The mixture is stirred at room temperature until the reaction is complete according to TLC checking.


For working-up, it is washed 2 times with 0.5 N NaOH, dried and evaporated. The crude product is recrystallized from MTB ether/n-hexane.

10 Yield: 3.60 g (68% of theory); m.p.: > 220°C.

The compounds of the formula (I-1-c) shown in Table 23 below were prepared analogously to Example (I-1-c-1) or according to the general details for preparation.

Le A 31 064 - Foreign Countries


- 148 -

Table 23

| Ex. No. | X | Y | Z | B | A | D | L | M | R ² | M.p.: °C | Iso-mer |
|----------|-------------------------------|-----------------|-----------------|--|------------------------------------|---------------------------------|---|---|--|----------|---------|
| I-1-c-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | C ₂ H ₅ - | 217 | β |
| I-1-c-3 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | O | O | C ₂ H ₅ - | >220 | - |
| I-1-c-4 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | C ₂ H ₅ - | 167 | β |
| I-1-c-5 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | O | O | C ₂ H ₅ - | 193 | β |
| I-1-c-6 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | O | O | C ₂ H ₅ - | 217 | - |
| I-1-c-7 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | i-C ₄ H ₉ - | 201 | β |
| I-1-c-8 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | C ₆ H ₅ - | >220 | β |
| I-1-c-9 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | C ₆ H ₅ -CH ₂ | >220 | β |
| I-1-c-10 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | i-C ₄ H ₉ - | 174 | β |
| I-1-c-11 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | O | O | i-C ₄ H ₉ - | >220 | - |
| I-1-c-12 | CH ₃ | Cl | CH ₃ | CH ₃ | CH ₃ | H | O | O | i-C ₄ H ₉ - | 157 | - |
| I-1-c-13 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | O | O | i-C ₄ H ₉ - | 193 | β |
| I-1-c-14 | CH ₃ | CH ₃ | Cl | CH ₃ | CH ₃ | H | O | O | i-C ₄ H ₉ - | 118 | - |
| I-1-c-15 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | O | O | C ₂ H ₅ - | >220 | - |
| I-1-c-16 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | O | O | i-C ₄ H ₉ - | 205 | - |
| I-1-c-17 | CH ₃ | CH ₃ | Br | H | -(CH ₂) ₄ - | | O | S | i-C ₃ H ₇ - | oil | - |
| I-1-c-18 | CH ₃ | CH ₃ | Br | H | -(CH ₂) ₄ - | | O | S | t-C ₄ H ₉ - | oil | - |
| I-1-c-19 | CH ₃ | CH ₃ | Br | H | H | i-C ₃ H ₇ | O | S | i-C ₃ H ₇ - | 124 | - |
| I-1-c-20 | CH ₃ | CH ₃ | Br | H | H | i-C ₃ H ₇ | O | S | t-C ₄ H ₉ - | 169 | - |

Le A 31 064 - Foreign Countries

- 149 -


Example I-1-d-1

3.64 g of the compound according to Example I-1-a-5 and 1.4 ml of triethylamine in 50 ml of abs. methylene chloride are treated at 0 to 10°C with 0.8 ml of methanesulphonyl chloride in 5 ml of abs. methylene chloride and the mixture is then stirred at room temperature. After reaction is complete (checking by means of thin-layer chromatography (TLC)), the mixture is washed 2 times with 50 ml of 0.5 N NaOH, dried over magnesium sulphate and concentrated, and the residue is recrystallized from MTB ether/n-hexane.

10 Yield 2.90 g (65% of theory), m.p. > 220°C.

Le A 31 064 - Foreign Countries


- 150 -

Example (II-1)

23.1 g of 2,6-dimethyl-4-bromo-phenylacetic acid according to Example (XXV-1) and 17.7 ml (0.24 mol) of thionyl chloride are stirred at 80°C until evolution of gas is complete. Excess thionyl chloride is distilled off at 50°C and the residue is taken up in 100 ml of absolute THF. This solution is added dropwise at 0°C to 10°C to a mixture of 20.9 g of methyl 1-amino-3-methyl-cyclohexanecarboxylate in 200 ml of absolute THF and 30.8 ml (0.22 mol) of triethylamine.

For working-up, the solution is filtered off with suction, washed with THF and evaporated, and the residue is taken up in methylene chloride. The solution is washed with 0.5 N HCl, dried and evaporated, and the residue is recrystallized from MTB ether/n-hexane.

Yield: 32.60 g (80% of theory); m.p.: 137°C.

Example (II-2)

28.8 g of the compound according to Example (XXXI-1) in 170 ml of methylene chloride are added at 30 to 40°C to 42 g (0.428 mol) of conc. sulphuric acid and the mixture is stirred at this temperature for a further 2 hours. 57 ml of absolute methanol are then added dropwise such that a temperature of 40°C is established. After addition is complete, the mixture is stirred at 40 to 70°C for a further 6 hours.

For working-up, it is poured onto ice, extracted with methylene chloride, washed with NaHCO₃ solution, dried and evaporated. The crude product is recrystallized from MTB ether/n-hexane.


Yield: 20.7 g (65% of theory); m.p.: 172°C.

The compounds of the formula (II) shown in Table 24 below were prepared analogously to Examples (II-1) and (II-2) or according to the general details for preparation.


Le A 31 064 - Foreign Countries

- 152 -

Table 24



| Ex. No. | X | Y | Z | B | A | D | R ⁸ | M.p.: °C | Iso-mer |
|---------|-------------------------------|-----------------|-------------------------------|---|--|---------------------------------|-------------------------------|----------|---------|
| II-3 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 168 | β |
| II-4 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | CH ₃ | 162 | - |
| II-5 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 129 | β |
| II-6 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 157 | β |
| II-7 | CH ₃ | CH ₃ | Br | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | CH ₃ | 127 | β |
| II-8 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | CH ₃ | 163 | - |
| II-9 | CH ₃ | CH ₃ | Br | CH ₃ | i-C ₃ H ₇ | H | CH ₃ | 183 | - |
| II-10 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 146 | β |
| II-11 | CH ₃ | CH ₃ | Br | CH ₃ | CH ₃ | H | CH ₃ | 141 | - |
| II-12 | CH ₃ | CH ₃ | Br | H | -CH ₂ -CHCH ₃ -CHCH ₃ - | | C ₂ H ₅ | 46 | trans* |
| II-13 | CH ₃ | CH ₃ | Br | H | -CH ₂ -S-(CH ₂) ₂ - | | C ₂ H ₅ | oil | - |
| II-14 | CH ₃ | CH ₃ | Br | H | H | i-C ₃ H ₇ | C ₂ H ₅ | oil | - |
| II-15 | CH ₃ | CH ₃ | Br | H | -(CH ₂) ₄ - | | C ₂ H ₅ | oil | - |
| II-16 | CH ₃ | CH ₃ | Br | H | -CH ₂ -CHCH ₃ -CHCH ₃ | | C ₂ H ₅ | oil | cis* |
| II-17 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 45 | β |
| II-18 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ - | | H | CH ₃ | 110 | β |
| II-19 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₃ -CHCH ₃ -CH ₂ - | | H | CH ₃ | 40 | β |
| II-20 | CH ₃ | CH ₃ | Cl | CH ₃ | CH ₃ | H | CH ₃ | 134 | - |
| II-21 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ | | H | CH ₃ | 132 | β |
| II-22 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ | | H | CH ₃ | 162 | - |
| II-23 | C ₂ H ₅ | Br | C ₂ H ₅ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ | | H | CH ₃ | 163 | β |
| II-24 | C ₂ H ₅ | Br | C ₂ H ₅ | -(CH ₂) ₂ -O-(CH ₂) ₂ | | H | CH ₃ | 163 | - |
| II-25 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -CHCH ₃ -(CH ₂) ₂ | | H | CH ₃ | 179 | β |
| II-26 | CH ₃ | Cl | CH ₃ | CH ₃ | CH ₃ | H | CH ₃ | 172 | - |
| II-27 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ | | H | CH ₃ | 148 | - |
| II-28 | CH ₃ | CH ₃ | Cl | CH ₃ | i-C ₃ H ₇ | H | CH ₃ | 93 | - |
| II-29 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -CHOCH ₃ -(CH ₂) ₂ | | H | CH ₃ | 138 | β |

Example (XXXI-1)


- 23.1 g of 2,6-dimethyl-4-bromophenylacetic acid according to Example (XXV-1) and 17.7 ml of thionyl chloride are stirred at 80°C until the evolution of gas is complete.
- 5 Excess thionyl chloride is then removed in vacuo at 50°C. The residue is taken up in 100 ml of absolute THF and added dropwise at 0 to 10°C to a mixture of 11.2 g of the amine of the formula $(\text{CH}_3)_2\text{CHC}(\text{CH}_2)(\text{CN})\text{NH}_2$ and 14.4 ml (0.11 mol) of triethylamine in 100 ml of absolute THF. It is then stirred at room temperature for a further hour.
- 10 For working-up, it is filtered off with suction and concentrated, the residue is taken up in methylene chloride, and the solution is washed in 0.5 N HCl, dried and concentrated. The crude product is recrystallized from MTB ether/n-hexane.

Yield: 28.8 g (85% of theory); m.p.: 169°C.

- 15 The compounds of the formula (XXXI) shown in Table 25 below were prepared analogously to Example (XXXI-1).

Le A 31 064 - Foreign Countries


- 154 -

Table 25

| Ex. No. | X | Y | Z | A | B | D | M.p.: °C |
|---------|-------------------------------|-----------------|-------------------------------|---|-----------------|---|----------|
| XXXI-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 206 |
| XXXI-3 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 201 |
| XXXI-4 | CH ₃ | CH ₃ | Br | i-C ₃ H ₇ | CH ₃ | H | 139 |
| XXXI-5 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 158 |
| XXXI-6 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 180 |
| XXXI-7 | CH ₃ | CH ₃ | Cl | i-C ₃ H ₇ | CH ₃ | H | 145 |
| XXXI-8 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 172 |
| XXXI-9 | C ₂ H ₅ | Br | C ₂ H ₅ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | H | 158 |

Le A 31 064 - Foreign Countries

- 155 -

Example (I-2-a-1)


(I-2-a-1)

A solution of 19.8 g (50 mmol) of the compound according to Example (III-1) in 50 ml of DMF are added dropwise at 0 to 10°C to 8.42 g (75 mmol) of potassium tert-butoxide in 50 ml of dimethylformamide (DMF) and the mixture is stirred overnight at room temperature.

For working-up, the reaction mixture is added dropwise to 500 ml of ice-cold 1 N HCl, and the crude product which is precipitated is filtered off with suction, washed with water and dried in a vacuum drying oven. For further purification, the crude product is also boiled with n-hexane/acetone.

Yield: 13.6 g (77% of theory); m.p.: > 250°C.

The compounds of the formula (I-2-a) shown in Table 26 below were prepared analogously to Example (I-2-a-1).

Table 26

(I-2-a)

| Ex. No. | X | Y | Z | A | B | M.p.: °C |
|---------|-----------------|-----------------|-----------------|---|------------------------------------|----------|
| I-2-a-2 | CH ₃ | Br | CH ₃ | | -(CH ₂) ₅ - | >250 |
| I-2-a-3 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₅ - | >240 |

Le A 31 064 - Foreign Countries


- 156 -

Table 26 - Continuation

| Ex. No. | X | Y | Z | A | B | M.p.:°C |
|----------|-------------------------------|-----------------|-----------------|---|---|---------|
| I-2-a-4 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ | | 239-241 |
| I-2-a-5 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₄ - | | 268 |
| I-2-a-6 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₅ - | | 238 |
| I-2-a-7 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ | | 223 |
| I-2-a-8 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₄ - | | 258 |
| I-2-a-9 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₅ - | | 233-234 |
| I-2-a-10 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ | | 212-215 |
| I-2-a-11 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | 240-242 |
| I-2-a-12 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ | | 258-259 |
| I-2-a-13 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | 262-263 |

Le A 31 064 - Foreign Countries

- 157 -

Example (I-2-b-1)

(I-2-b-1)

A solution of 1.57 g (13 mmol) of pivaloyl chloride in 40 ml of methylene chloride is added dropwise with ice-cooling to a mixture of 3.52 g (10 mmol) of the compound according to Example (I-2-a-1) and 1.52 g (15 mmol) of triethylamine in 40 ml of methylene chloride and the mixture is stirred at room temperature for 2 hours.


For working up, it is washed successively with 10% strength citric acid, 1 N NaOH and NaCl solution, dried and evaporated. For further purification, the crude product is also stirred with a little petroleum ether.

10 Yield: 1.95 g (45% of theory); m.p.: 107-109°C.

The compounds of the formula (I-2-b) shown in Table 27 below were prepared analogously to Example (I-2-b-1).

Le A 31 064 - Foreign Countries

- 158 -

Table 27

(I-2-b)

| Ex. No. | X | Y | Z | A | B | R ¹ | M.p.: °C |
|----------|-----------------|-----------------|-----------------|---|---|--|----------|
| I-2-b-2 | CH ₃ | Br | CH ₃ | | -(CH ₂) ₅ - | t-C ₄ H ₉ - | 150-152 |
| I-2-b-3 | CH ₃ | Br | CH ₃ | | -(CH ₂) ₅ - | t-C ₄ H ₉ -CH ₂ - | 158-161 |
| I-2-b-4 | CH ₃ | CH ₃ | Br | | -(CH ₂) ₅ - | t-C ₄ H ₉ -CH ₂ - | 147-150 |
| I-2-b-5 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₅ - | t-C ₄ H ₉ - | oil |
| I-2-b-6 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₅ - | t-C ₄ H ₉ -CH ₂ - | 160-163 |
| I-2-b-7 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₅ - | i-C ₄ H ₉ - | 113 |
| I-2-b-8 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₂ -O-(CH ₂) ₂ - | t-C ₄ H ₉ - | 109-110 |
| I-2-b-9 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₄ - | i-C ₄ H ₉ - | 90-91 |
| I-2-b-10 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₄ - | t-C ₄ H ₉ -CH ₂ - | 106-107 |
| I-2-b-11 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₄ - | i-C ₃ H ₇ - | 84-85 |
| I-2-b-12 | CH ₃ | CH ₃ | Cl | | -(CH ₂) ₄ - | t-C ₄ H ₉ - | 105-106 |
| I-2-b-13 | CH ₃ | Cl | CH ₃ | | -(CH ₂) ₅ - | t-C ₄ H ₉ - | oil |
| I-2-b-14 | CH ₃ | Cl | CH ₃ | | -(CH ₂) ₅ - | t-C ₄ H ₉ -CH ₂ - | oil |
| I-2-b-15 | CH ₃ | Cl | CH ₃ | | -(CH ₂) ₂ -O-(CH ₂) ₂ - | t-C ₄ H ₉ - | 148 |
| I-2-b-16 | CH ₃ | Cl | CH ₃ | | -(CH ₂) ₄ - | i-C ₄ H ₉ - | oil |
| I-2-b-17 | CH ₃ | Cl | CH ₃ | | -(CH ₂) ₄ - | t-C ₄ H ₉ -CH ₂ - | 83 |

Le A 31 064 - Foreign Countries

- 159 -


Table 27 - Continuation

| Ex. No. | X | Y | Z | A | B | R ¹ | M.p.:°C |
|----------|-------------------------------|-----------------|-----------------|---|---|--|---------|
| I-2-b-18 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₄ - | | i-C ₃ H ₇ - | oil |
| I-2-b-19 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₄ - | | t-C ₄ H ₉ - | 102 |
| I-2-b-20 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₅ - | | t-C ₄ H ₉ - | 115-116 |
| I-2-b-21 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₅ - | | t-C ₄ H ₉ -CH ₂ - | 139-140 |
| I-2-b-22 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | t-C ₄ H ₉ -CH ₂ - | 158-159 |
| I-2-b-23 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | t-C ₄ H ₉ - | 148-149 |
| I-2-b-24 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | i-C ₄ H ₉ - | 84-85 |
| I-2-b-25 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | t-C ₄ H ₉ -CH ₂ - | 92-93 |
| I-2-b-26 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | i-C ₃ H ₇ - | 83-84 |
| I-2-b-27 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | t-C ₄ H ₉ - | 106-107 |
| I-2-b-28 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | t-C ₄ H ₉ -CH ₂ - | 126-127 |
| I-2-b-29 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | t-C ₄ H ₉ - | 128-129 |
| I-2-b-30 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | i-C ₄ H ₉ - | 99-100 |
| I-2-b-31 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | t-C ₄ H ₉ -CH ₂ - | 89-90 |
| I-2-b-32 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | i-C ₃ H ₇ - | 83-84 |
| I-2-b-33 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | t-C ₄ H ₉ - | 123-124 |
| I-2-b-34 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | C ₆ H ₅ - | 132-135 |

Le A 31 064 - Foreign Countries

- 160 -


The compounds of the formula I-2-c shown in Table 28 were obtained analogously to Example I-1-c-1.

Table 28

| Ex. No. | X | Y | Z | A | B | L | M | R ² | M.p.:°C |
|---------|-----------------|-----------------|-----------------|---|---|---|---|-----------------------------------|---------|
| I-2-c-1 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | O | O | i-C ₃ H ₇ - | oil |
| I-2-c-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | O | O | i-C ₃ H ₇ - | 119-120 |

Le A 31 064 - Foreign Countries

- 161 -


Example (III-1)

12.15 g (50 mmol) of the compound according to Example (XXV-2) and 11.9 g (100 mmol) of thionyl chloride are stirred in 50 ml of toluene at 80°C until evolution of gas is complete. The mixture is then evaporated to dryness and the crude acid chloride thus obtained is heated under reflux overnight in 50 ml of toluene together with 8.6 g (50 mmol) of ethyl 1-hydroxycyclohexanecarboxylate. The mixture is then concentrated.

Yield: 19.8 g (quantitative); colourless oil.

10 ¹H-NMR: δ = 1.20 (t, 3H); 1.40-1.80 (m, 8H); 2.15 (m, 2H); 2.25 (s, 3H); 2.35 (s, 3H); 3.90 (s, 2H); 4.15 (q, 2H); 6.95 (m, 1H); 7.25 (m, 1H).

The compounds of the formula (III) shown in Table 29 are obtained analogously or according to the general details for preparation.

Table 29


| Ex. No. | A | B | X | Y | Z | R ⁸ | M.p.: °C |
|---------|---|------------------------------------|-----------------|-----------------|-----------------|---------------------------------|----------|
| III-2 | | -(CH ₂) ₅ - | CH ₃ | Br | CH ₃ | C ₂ H ₅ - | oil |
| III-3 | | -(CH ₂) ₅ - | CH ₃ | CH ₃ | Cl | C ₂ H ₅ - | oil |

Le A 31 064 - Foreign Countries


- 162 -

Table 29 - Continuation

| Ex. No. | X | Y | Z | A | B | R ⁸ | M.p.:°C |
|---------|-------------------------------|-----------------|-----------------|---|---|-----------------|---------|
| III-4 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₅ - | | CH ₃ | oil |
| III-5 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | CH ₃ | oil |
| III-6 | CH ₃ | Br | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | CH ₃ | oil |
| III-7 | CH ₃ | CH ₃ | Br | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | CH ₃ | oil |
| III-8 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₂ -O-(CH ₂) ₂ - | | CH ₃ | oil |
| III-9 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | CH ₃ | oil |
| III-10 | CH ₃ | Br | CH ₃ | -(CH ₂) ₄ - | | CH ₃ | oil |
| III-11 | CH ₃ | CH ₃ | Cl | -(CH ₂) ₄ - | | CH ₃ | oil |
| III-12 | CH ₃ | Cl | CH ₃ | -(CH ₂) ₄ - | | CH ₃ | oil |
| III-13 | C ₂ H ₅ | Br | CH ₃ | -(CH ₂) ₅ - | | CH ₃ | oil |

Example I-3-a-1

- 34.0 g (69 mmol) of the compound according to Example (IV-1) are heated under reflux for 3 hours in 70 ml of trifluoroacetic acid and 140 ml of toluene. The trifluoroacetic acid is then removed in vacuo and the residue is treated with 400 ml of water and 120 ml of MTB ether. A pH of 14 is established by addition of NaOH, then the mixture is extracted 2 times with MTB ether. The aqueous phase is acidified with HCl and extracted 3 times with MTB ether. After drying, the organic phase is concentrated. Yield 13.0 g (55% of theory), m.p. 235-238°C.
- 10 The compounds of the formula I-3-a shown in Table 30 were prepared analogously to Example I-3-a-1.


Table 30

I-3-a

| Ex. No. | X | Y | Z | A | B | M.p.:°C |
|---------|-----------------|-----------------|-----------------|------------------------------------|---|---------|
| I-3-a-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | 255-257 |
| I-3-a-3 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | >230 |

Le A 31 064 - Foreign Countries

- 164 -

Example I-3-b-1

A solution of 0.74 ml (0.89 g; 5.72 mmol) of 3-chloro-2,2-dimethylpropionyl chloride
 in 3 ml of methylene chloride is added dropwise with ice-cooling to a mixture of 1.5 g
 5 (4.4 mmol) of the compound according to Example (I-3-a-1), 0.92 ml of triethylamine
 and 20 ml of methylene chloride and the mixture is then stirred at room temperature for
 2 hours.

It is then washed 2 times with 10% strength citric acid and the acidic aqueous phases
 are extracted with methylene chloride. The combined organic phases are washed 2
 10 times with 1N NaOH and the aqueous alkaline phases are extracted with methylene
 chloride. The combined organic phases are dried and concentrated. Yield 1.65 g (81%
 of theory), oil.


¹H-NMR in CDCl₃, ppm

| | | | |
|----|-----|-----------|--|
| | δ = | 1.05 | (t, 3H, CH ₂ CH ₃) |
| 15 | | 1.18 | (s, 6H, C(CH ₃) ₂) |
| | | 1.62 | (s, 3H, CCH ₃) |
| | | 1.95-2.05 | (m, 2H, CH ₂ CH ₃) |
| | | 2.08 | (s, 3H, ArCH ₃) |
| | | 2.10 | (s, 3H, ArCH ₃) |
| 20 | | 3.38 | (s, 2H, CH ₂ Cl) |
| | | 7.20 | (s, 2H, ARH) |

Le A 31 064 - Foreign Countries


- 165 -

The compounds of the formula I-3-b shown in Table 31 below were prepared analogously to Example I-3-b-1.

Table 31

| Ex. No. | X | Y | Z | A | B | R ¹ | M.p.:°C |
|----------|-----------------|-----------------|-----------------|------------------------------------|-----------------|--|---------------|
| I-3-b-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | t-C ₄ H ₉ | 105-107 |
| I-3-b-3 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | Cl-CH ₂ -C(CH ₃) ₂ - | 129-131 |
| I-3-b-4 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | 3-(6-Cl-Pyridyl)- | 171-173 |
| I-3-b-5 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | cyclo-C ₃ H ₅ - | 176-180 |
| I-3-b-6 | CH ₃ | Br | CH ₃ | C ₂ H ₅ | CH ₃ | t-C ₄ H ₉ - | oil |
| I-3-b-7 | CH ₃ | Br | CH ₃ | C ₂ H ₅ | CH ₃ | cyclo-C ₃ H ₅ - | oil |
| I-3-b-8 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | t-C ₄ H ₉ - | 81-85 |
| I-3-b-9 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | 4-Cl-C ₆ H ₄ - | ¹⁾ |
| I-3-b-10 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | ClCH ₂ -C(CH ₃) ₂ - | 112-116 |
| I-3-b-11 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | cyclo-C ₃ H ₅ - | oil |

1) ¹H-NMR, CDCl₃, [ppm], δ = 1.4 to 2.0 (m, 10H), 2.15 (s, 3H), 2.30 (s, 3H), 7.00 (s, 1H), 7.30 (s, 1H), 7.42 (d, 2H), 7.81 (d, 2H).

Example I-3-c-1


A solution of 0.74 ml (5.72 mmol) of isobutyl chloroformate in 3 ml of methylene chloride is added dropwise with ice-cooling to a mixture of 1.5 g (4.4 mmol) of the compound according to Example (I-3-a-1), 0.92 ml of triethylamine and 20 ml of methylene chloride. The mixture is stirred at room temperature for 2 hours and worked up as described in Example I-3-b-1. The residue remaining at the end is stirred with petroleum ether. Yield 2.0 g (100% of theory)

| | | |
|----|---|---|
| | ¹ H-NMR, CDCl ₃ , [ppm], δ = 0.68 | (d, 6H, CH(CH ₃) ₂) |
| 10 | 1.04 | (t, 3H, CH ₂ CH ₃) |
| | 1.5-1.6 | (m, 1H, CH(CH ₃) ₂) |
| | 1.71 | (m, 2H, CCH ₃) |
| | 1.9-2.0 | (m, 2H, CH ₂ CH ₃) |
| | 2.08 | (s, 3H, ArCH ₃) |
| 15 | 2.12 | (s, 3H, ArCH ₃) |
| | 3.61 | (d, 2H, OCH ₂) |
| | 7.12 | (s, 2H, ArH) |


Le A 31 064 - Foreign Countries

- 167 -


The compounds of the formula I-3-c shown in Table 32 were prepared analogously to Example I-3-c-1.

Table 32

| Ex. No. | X | Y | Z | A | B | L | M | R ² | M.p.:°C |
|---------|-----------------|-----------------|-----------------|------------------------------------|-----------------|---|---|--|---------|
| I-3-c-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | O | S | i-C ₃ H ₇ - | oil |
| I-3-c-3 | CH ₃ | Br | CH ₃ | C ₂ H ₅ | CH ₃ | O | O | 4-Cl-C ₆ H ₄ - | oil |
| I-3-c-4 | CH ₃ | Br | CH ₃ | C ₂ H ₅ | CH ₃ | O | S | t-C ₄ H ₉ - | oil |
| I-3-c-5 | CH ₃ | Br | CH ₃ | C ₂ H ₅ | CH ₃ | O | S | C ₆ H ₅ -CH ₂ - | oil |
| I-3-c-6 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | O | S | i-C ₃ H ₇ - | 124-127 |

Example IV-1

A: A mixture of 25.0 g (98 mmol) of the compound of the formula



5 1 drop of DMF and 17.5 g (147 mmol) of thionyl chloride in 70 ml of toluene is stirred at room temperature for 5 minutes and then at 100°C until the evolution of gas is complete. Excess thionyl chloride is removed in vacuo.

10 B: 27.7 g of the compound according to Example (XXVI-1) in 40 ml of THF is added dropwise at 0°C to a mixture of 13.0 g (129 mmol) of diisopropylamine and 71.6 ml (118 mmol) of butyllithium (1.6 M in n-hexane) in 100 ml of THF and the mixture is stirred for 30 minutes. The acid chloride prepared under A, dissolved in 40 ml of THF, is then added dropwise at 0°C and the mixture is stirred at room temperature for 1 hour.


15 350 ml of MTB ether and a few drops of water are added, the mixture is washed 2 times with 10% strength ammonium chloride solution, and the organic phase is dried and concentrated. The crude product is chromatographed on silica gel (eluent cyclohexane/ethyl acetate 20:1 to 5:1). Yield 35.0 g (72% of theory).

Le A 31 064 - Foreign Countries

- 169 -

¹H-NMR, CDCl₃, [ppm]; δ = 0.9 to 1.0 (m, 3H), 1.43 (s, 3H), 1.7 to 2.0 (m, 2H), 2.3 to 2.4 (s, 6H), 3.5 to 3.8 (m, 8H), 6.7 to 7.2 (m, 6H)


The compounds of the formula IV shown in the table below were obtained analogously to Example IV-1 and according to the general description.

Table 33

| Ex. No. | X | Y | Z | A | B | R ⁸ | M.p.:°C |
|---------|-----------------|-----------------|-----------------|------------------------------------|---|-----------------|---------|
| IV-2 | CH ₃ | Br | CH ₃ | -(CH ₂) ₅ - | | CH ₃ | 153 |
| IV-3 | CH ₃ | CH ₃ | Br | -(CH ₂) ₅ - | | CH ₃ | 153 |

Le A 31 064 - Foreign Countries

- 170 -


Example (I-5-a-1)

5.7 g (20 mmol) of 2-(4-bromo-2,6-dimethylphenyl)-chlorocarbonylketene are heated under reflux for 8 h with 2.0 g (20 mmol) of cyclohexanone in 60 ml of xylene. The precipitate which is deposited is separated off, washed with cyclohexane and dried. 5.0 g of product (72% of theory) of melting point 244 to 245°C are obtained.


Le A 31 064 - Foreign Countries

- 171 -

The following compounds of the formula (I-5-a) are obtained analogously or according to the general details for preparation:


Table 34

(I-5a)


| Ex. No. | X | Y | Z | A | D | M.p.[°C]- ¹ H-NMR (CDCl ₃): δ [ppm] |
|----------|-----------------|-----------------|-----------------|------------------------------------|---|---|
| I-5-a-2 | CH ₃ | CH ₃ | Br | CH ₃ | 4-F-Phenyl | oil |
| I-5-a-3 | CH ₃ | Br | CH ₃ | CH ₃ | t-Bu | 219-221 |
| I-5-a-4 | CH ₃ | CH ₃ | Br | CH ₃ | 2-Pyridyl | 262-264 |
| I-5-a-5 | CH ₃ | Br | CH ₃ | CH ₃ | 4-F-Phenyl | 210-211 |
| I-5-a-6 | CH ₃ | Br | CH ₃ | CH ₃ | 2-Pyridyl | 104-106 |
| I-5-a-7 | CH ₃ | Br | CH ₃ | CH ₃ |  | 206-208 |
| I-5-a-8 | CH ₃ | CH ₃ | Br | CH ₃ | t-Bu | 7.37 (1H,s), 7.07 (1H,s), 2.3 (3H,s), 2.18 (3H,s), 2.15 (3H,s), 1.49 (9H,s) |
| I-5-a-9 | CH ₃ | CH ₃ | Br | -(CH ₂) ₄ - | | 250-252 |
| I-5-a-10 | CH ₃ | CH ₃ | Cl | CH ₃ | 2-Pyridyl | 197-199 |
| I-5-a-11 | CH ₃ | CH ₃ | Cl | CH ₃ | 4-F-Phenyl | 188-190 |

Le A 31 064 - Foreign Countries

- 172 -

Example (I-5-b-1)

- 1.9 g (5 mmol) of the compound (I-5-a-6) are initially introduced into 20 ml of ethyl acetate and treated with 0.5 g (5 mmol) of triethylamine, and 0.4 g (5 mmol) of acetyl chloride in 5 ml of ethyl acetate is added dropwise at 0°C. The mixture is stirred at room temperature for 20 h, and the precipitate is separated off, washed twice with 50 ml of half-concentrated sodium chloride solution, dried over sodium sulphate and evaporated in vacuo. The residue is chromatographed on silica gel using toluene/acetone 30:1.
- 10 Yield 1.2 g (56% of theory) of melting point 130 to 132°C.


Example (I-6-a-1)

- 2.8 g (10 mmol) of 2-(4-bromo-2,6-dimethylphenyl)-chlorocarbonyl ketene are warmed to 50°C for 6 h with 1.6 g (10 mmol) of 4-fluorothiobenzamide in 80 ml of toluene. The precipitate is separated off, washed with cyclohexane and dried. 3.0 g (74% of theory) of melting point 275 to 276°C are obtained.
- 15


The following compound of the formula (I-6-a-2) of melting point 235 to 236°C is obtained analogously or according to the general details for preparation.

Le A 31 064 - Foreign Countries

- 173 -



(I-6-a-2)


Example (XXII-1)

(XXII-1)

8 g of the compound according to Example (XXV-1) are heated at 80°C with 8.7 ml
 5 of thionyl chloride until the evolution of gas is complete. Excess thionyl chloride is removed in vacuo and the residue is distilled.

Yield: 87% of theory; m.p.: 69-71°C.

The following compounds of the formula (XXII) are obtained analogously or according to the general details for preparation.

Table 35

(XXII)

Le A 31 064 - Foreign Countries

- 174 -

| Ex. No. | X | Y | Z | B.p. [°C], (mbar) |
|---------|-------------------------------|-----------------|-------------------------------|-------------------|
| XXII-2 | CH ₃ | CH ₃ | Br | 114-116, 0.06 |
| XXII-3 | CH ₃ | Br | C ₂ H ₅ | 120-132, 0.1 |
| XXII-4 | CH ₃ | CH ₃ | Cl | * |
| XXII-5 | CH ₃ | Cl | CH ₃ | * |
| XXII-6 | C ₂ H ₅ | Br | C ₂ H ₅ | 131, 0.15 |


* These acid chlorides were used as crude products for the syntheses of the compounds II, III and XXXI and were not characterized in greater detail.

Example (XXXV-1)


222.4 g (0.865 mol) of the compound according to Example (XXVI-1) and 80.56 g (1.438 mol) of potassium hydroxide in 210 ml of methanol and 105 ml of water are heated under reflux for 5 hours. After cooling, the mixture is concentrated and the residue is dissolved in water. The aqueous phase is washed with ethyl acetate and then with dilute hydrochloric acid. The product which is deposited is filtered off with suction, washed with water and dried.

Yield: 197.5 g (94% of theory); m.p.: 185-187°C.

10 The compounds of the formula (XXV) shown in Table 36 are obtained analogously or according to the general details for preparation.

Table 36

| Ex. No. | X | Y | Z | M.p. [°C] |
|---------|-------------------------------|-----------------|-------------------------------|-----------|
| XXV-2 | CH ₃ | CH ₃ | Br | 174-176 |
| XXV-3 | C ₂ H ₅ | Br | CH ₃ | 122-123 |
| XXV-4 | CH ₃ | CH ₃ | Cl | 166-168 |
| XXV-5 | CH ₃ | Cl | CH ₃ | 178-180 |
| XXV-6 | C ₂ H ₅ | Br | C ₂ H ₅ | 143 |

Example (XXVI-1)


349.3 g (1.044 mol) of the compound according to Example (XXVII-1) (94.57% strength), 475 ml of methanol and 842 ml of 30% strength sodium methoxide solution in methanol are heated under reflux for 5 hours. 126 ml of conc. sulphuric acid are then added at room temperature and the mixture is heated under reflux for 1 hour. The solvent is distilled off, and the residue is treated with water and extracted with methylene chloride. After drying, it is filtered, concentrated and finally distilled.

Yield: 222.4 g (82.9% of theory); b.p._{0.2} 98-100°C.

10 The following compounds of the formula (XXVI) are obtained analogously or according to the general procedures for preparation.

Table 37

| Ex. No. | X | Y | Z | R ⁸ | B.p. [°C], (mbar) |
|---------|-------------------------------|-----------------|-------------------------------|-----------------|-------------------|
| XXVI-2 | CH ₃ | CH ₃ | Br | CH ₃ | 93-94, 0.2 |
| XXVI-3 | CH ₃ | Br | C ₂ H ₅ | CH ₃ | 160-165, 20 |
| XXVI-4 | CH ₃ | CH ₃ | Cl | CH ₃ | oil |
| XXVI-5 | CH ₃ | Cl | CH ₃ | CH ₃ | 152-158, 16 |
| XXVI-6 | C ₂ H ₅ | Br | C ₂ H ₅ | CH ₃ | 90-95, 0.015 |

Example (XXVII-1)

326 g (2.673 mol) of anhydrous copper(II) chloride are added to a solution of 326 g (3.175 mol) of tert-butyl nitrite in 1270 ml of dry acetonitrile. 3130 g (32.27 mol) of 1,1-dichloroethane are added dropwise to the well-cooled mixture, the mixture being kept at below 30°C by means of ice-cooling. A solution of 424 g (2.12 mol) of 4-bromo-2,6-dimethylaniline in 2120 ml of acetonitrile is then added dropwise at below 30°C. The mixture is stirred at room temperature until the evolution of gas (N₂) is complete (about 3 hours). The almost black solution is carefully poured into 9 l of 20% strength hydrochloric acid and extracted several times, altogether with 9 l, of methyl tert-butyl ether. The combined organic phases are washed with 20% strength hydrochloric acid and dried over anhydrous magnesium sulphate. The magnesium sulphate is filtered off and the solution is concentrated. The residual oil is


Le A 31 064 - Foreign Countries

- 177 -

fractionated in a high vacuum.


Yield: 349.3 g (49% of theory); b.p._{0.1} 130-137°C.

The following compounds of the formula (XXVII) were prepared analogously or according to the general details for preparation:

Table 38


(XXVII)

| Ex. No. | X | Y | Z | B.p. [°C], (mbar) |
|---------|-------------------------------|-----------------|-------------------------------|-------------------|
| XXVII-2 | CH ₃ | CH ₃ | Br | 110-115, 0.15 |
| XXVII-3 | CH ₃ | Br | C ₂ H ₅ | oil |
| XXVII-4 | CH ₃ | CH ₃ | Cl | oil |
| XXVII-5 | CH ₃ | Cl | CH ₃ | oil |
| XXVII-6 | C ₂ H ₅ | Br | C ₂ H ₅ | oil |

Example VI-1

7.1 g of NaH (80% strength) were initially introduced into 278 ml of dimethyl carbonate and heated to 80 to 90°C. 39 g of methyl 2-chloro-4,6-dimethylphenylacetate were then added dropwise and the mixture was heated at reflux for 20 h. A further 3.4 g of NaH (80% strength) were added and the mixture was heated at reflux for a further 8 h. The mixture was cooled, any NaH still present was destroyed with a little methanol, and it was poured onto ice. After acidifying with half-concentrated HCl, the organic phase was separated off and the aqueous phase was extracted several times with dichloromethane. The combined organic phase was dried and concentrated. Yield: 35.1 g of a solid having a melting point of 67 to 70°C.

¹H-NMR (CDCl₃): δ 7.12 (s, 1H), 6.94 (s, 1H), 5.36 (2, 1H), 3.78 (s, 6H), 2.31 (s, 3H), 2.28 ppm (s, 3H).

Example XXXV-1


10 g of dimethyl 2-chloro-4,6-dimethylphenylmalonate according to Example (VI-1) were initially introduced and treated successively with 20 ml of methanol and 6.8 g of KOH dissolved in 9.1 ml of water. After an hour, the mixture was diluted with a further 20 ml of solvent (MeOH/water 1:1). The mixture was heated at reflux for 10 h, then cooled and concentrated. The residue which remained was taken up in a little water and washed once with toluene. The aqueous phase was then diluted further with water, added to the same amount of ether and cooled to about -10°C. It was acidified with concentrated HCl (pH 1), and the organic phase was separated off and reextracted

Le A 31 064 - Foreign Countries

- 179 -


a further 1 or 2 times. The combined organic phases were dried and concentrated. The residue was crystallized from toluene and afforded 7.6 g of 2-chloro-4,6-dimethylphenylmalonic acid having a melting point of 174 to 176°C (decomposition).

- 5 ¹H-NMR (CDCl₃) δ: 7.10 (s, 1H), 6.95 (s, 1H), 5.00 (s, 1H), 2.36 (s, 3H), 2.30 ppm (s, 3H).

Example V-1

- 7.6 g of 2-chloro-4,6-dimethylphenylmalonic acid according to Example (XXXV-1)
 10 were suspended in 22 ml of toluene and treated dropwise with 19.5 ml of thionyl chloride. The mixture was heated at 95°C for 9.5 h, cooled and freed from the volatile constituents by passing argon through. The residues of thionyl chloride and the solvent were distilled off at 45°C in a high vacuum. 6.6 g of 2-chloro-4,6-dimethylphenylchlorocarbonylketene were obtained as an oil which was slightly
 15 contaminated with 2-chloro-4,6-dimethylphenylacetyl chloride.

¹H-NMR (CDCl₃) δ: 7.16 (s, 1H), 7.02 (s, 1H), 2.33 (s, 3H), 2.30 ppm, (s, 3H).


Example (VI-2)

- 70 g of methyl 4-bromo-2,6-dimethylphenylacetate were reacted in an analogous
 20 manner with 26.8 g of NaH and 7.39 ml of dimethyl carbonate. After working up, 95.4 g of crude product (86.5% pure) were obtained.

Le A 31 064 - Foreign Countries


- 180 -

¹H-NMR (CDCl₃) δ: 7.22 (s, 2H), 5.00 (s, 4H), 3.75 (s, 6H), 2.33 ppm (s, 6H).

Example (XXXV-2)


85 g of dimethyl 4-bromo-2,6-dimethylphenylmalonate according to Example (VI-2) in
 5 158 ml of methanol were reacted in an analogous manner with 49.6 g of KOH in 66 ml
 of H₂O. After working up, 59.7 g of the malonic acid were obtained. Melting point 164
 to 167°C (decomposition).

¹H-NMR (CDCl₃) δ: 7.20 (s, 2H), 7.00 - 6.00 (OH), 4.83 (s, 1H), 2.30 ppm (s, 6H).

Example (V-2)

59 g of 4-bromo-2,6-dimethylphenylmalonic acid in 143 ml of toluene were reacted in
 an analogous manner with 128 ml of thionyl chloride and 49.5 g were isolated as a
 crude product.

¹H-NMR (CDCl₃) δ: 7.31 (s, 2H), 2.33 ppm (s, 6H).

Example (VI-3)


23 g of methyl 2-bromo-4,6-dimethylphenylacetate were reacted in an analogous
 manner with 9.5 g of NaH (80% strength) and 242 ml of dimethyl carbonate. After

Le A 31 064 - Foreign Countries

- 181 -


working up, 31.2 g of crude product (82% pure) were obtained.

¹H-NMR (CDCl₃) δ: 7.31 (s, 1H), 6.98 (s, 1H), 5.45 (s, 1H), 3.88 (s, 6H), 2.32 (s, 3H), 2.28 ppm (s, 3H).

Example (XXXV-3)

27 g of dimethyl 2-bromo-4,6-dimethylphenylmalonate according to Example (VI-3) in 50 ml of methanol were reacted in an analogous manner with 15.7 g of KOH in 21 ml of water. After working up, 17.4 g of malonic acid were obtained. Melting point 167 to 169°C (decomposition).

10 ¹H-NMR (CDCl₃) δ: 8.20 - 7.00 (OH), 7.26 (s, 1H), 6.98 (s, 1H), 5.07 (s, 1H), 2.35 (s, 3H), 2.28 ppm (s, 3H).

Example (V-3)

15 17 g of 2-bromo-4,6-dimethylphenylmalonic acid according to Example (XXXV-3) in 41 ml of toluene were reacted in an analogous manner with 36.8 ml of thionyl chloride and 15.1 g were isolated as a crude product.

IR: μ = 2130 (ketene)

¹H-NMR (CDCl₃) δ: 7.28 (s, 1H), 7.00 (s, 1H), 2.35 (s, 3H), 2.29 ppm (s, 3H).

Le A 31 064 - Foreign Countries

- 182 -

Use ExamplesExample A

Phaedon larvae test

- Solvent: 7 parts by weight of dimethylformamide
5 Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

- 10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with mustard beetle larvae (*Phaedon cochleariae*), while the leaves are still moist.

After 3 days in each case, the destruction in % is determined. 100% means that all the beetle larvae have been killed; 0% means that none of the beetle larvae have been killed.

- 15 In this test, for example, the compounds according to Preparation Examples (I-2-a-2), (I-2-b-2), (I-2-a-1), (I-2-b-1), (I-2-b-4), (I-1-a-2), (I-1-a-1), (I-1-b-2) and (I-1-b-4) at an exemplary active compound concentration of 0.1% caused a destruction of 100% after 7 days.

Le A 31 064 - Foreign Countries

- 183 -

Example B

Plutella test

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with caterpillars of the diamond-back moth (*Plutella xylostella*) while the leaves are still moist.

After the specified period of time, the destruction in % is determined. 100% means that all the caterpillars have been killed; 0% means that none of the caterpillars have been killed.

15 In this test, for example, the compounds according to Preparation Examples (I-2-b-2), (I-2-b-1), (I-1-b-2), (I-1-b-4), (I-1-c-2), (I-1-a-5) and (I-1-1-6) at an exemplary active compound concentration of 0.1% caused a destruction of 100% after 7 days.

Le A 31 064 - Foreign Countries

- 184 -

Example C

Nephotettix test

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

Rice seedlings (*Oryza sativa*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with the green rice
10 leafhopper (*Nephotettix cincticeps*) while the leaves are still moist.

After the specified period of time, the destruction in % is determined. 100% means that all the leafhoppers have been killed; 0% means that none of the leafhoppers have been killed.

15 In this test, for example, the compounds according to Preparation Examples (I-2-a-2), (I-2-a-1), (I-2-b-1), (I-1-a-2), (I-1-a-1), (I-1-b-2), (I-1-b-3), (I-1-b-4), (I-1-c-2), (I-1-a-5) and (I-1-a-6) at an exemplary active compound concentration of 0.1% caused a destruction of 100% after 6 days.

Le A 31 064 - Foreign Countries

- 185 -

Example D

Myzus test

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) which have been heavily infested with the peach aphid (*Myzus persicae*) are treated by being dipped into the preparation of the active compound of the desired concentration.

After the specified period of time, the destruction in % is determined. 100% means that all the aphids have been killed; 0% means that none of the aphids have been killed.

15 In this test, for example, the compounds according to Preparation Examples (I-2-a-1), (I-2-a-2), (I-2-b-2), (I-1-b-3), (I-1-c-2) and (I-1-a-6) at an exemplary active compound concentration of 0.1% caused a destruction of 100% after 6 days.

Le A 31 064 - Foreign Countries

- 186 -

Example E

Tetranychus test (OP-resistant/dipping treatment)

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and the stated amount of emulsifier, and the concentrate is diluted with water to the desired concentration.

- 10 Bean plants (*Phaseolus vulgaris*) which have been heavily infested with the common spider mite or two-spotted spider mite (*Tetranychus urticae*) in all stages of development are treated by being dipped into the preparation of the active compound of the desired concentration.

After the specified period of time, the destruction in % is determined. 100% means that all the spider mites have been killed; 0% means that none of the spider mites have been killed.

- 15 In this test, for example, the compounds according to Preparation Examples (I-2-a-2), (I-2-b-2), (I-2-b-3), (I-2-a-1) and (I-2-b-4) at an exemplary active compound concentration of 0.1% had an action of 100% after 13 days.

Le A 31 064 - Foreign Countries

- 187 -

Example F

Pre-emergence test

Solvent: 5 parts by weight of acetone

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, the stated amount of emulsifier is added, and the concentrate is diluted with water to the desired concentration.

10 Seeds of the test plants are sown in normal soil and, after 24 hours, watered with the preparation of the active compound. It is expedient to keep constant the amount of water per unit area. The concentration of the active compound in the preparation is of no importance, only the amount of active compound applied per unit area being decisive. After three weeks, the degree of damage to the plants is rated in % damage in comparison to the development of the untreated control. The figures denote:

0% = no action (like untreated control)

15 100% = total destruction

Le A 31 064 - Foreign Countries

- 188 -

Pre-emergence test/greenhouse


| Ex. No. | g/ha | Beta vulgaris | Alopecurus myosuroides | Avena fatua | Setaria viridis | Sinapis arvensis l. |
|---------|------|---------------|------------------------|-------------|-----------------|---------------------|
| I-1-a-2 | 250 | 0 | 100 | 60 | 100 | 90 |
| I-1-b-2 | 250 | 0 | 100 | 100 | 100 | - |
| I-1-b-3 | 250 | 0 | - | 80 | 100 | 95 |
| I-1-b-4 | 250 | 0 | - | 100 | 100 | 95 |
| I-1-c-2 | 250 | 0 | 95 | 80 | 100 | 100 |
| I-1-a-5 | 250 | - | 100 | 100 | 100 | 100 |
| I-1-a-6 | 250 | - | 95 | 95 | 100 | 100 |

30517-144

- 189 -

CLAIMS:

1. A compound of general formula (I):



wherein:


X represents C₁-C₆-alkyl;

Y represents a halogen atom or C₁-C₆-alkyl;

Z represents a halogen atom or C₁-C₆-alkyl,

- 10 wherein one of the substituents Y and Z always represents a halogen atom and the other C₁-C₆-alkyl; and

Het represents one of the groups:



30517-144

- 190 -

wherein:

A represents: (i) H, (ii) C₁-C₁₂-alkyl, C₂-C₈-alkenyl, C₁-C₁₀-alkoxy-C₁-C₈-alkyl, poly-C₁-C₈-alkoxy-C₁-C₈-alkyl or C₁-C₁₀-alkylthio-C₁-C₆-alkyl, each of which is
 5 optionally substituted by a halogen atom, (iii) C₃-C₈-cycloalkyl which is optionally substituted by a halogen atom, C₁-C₆-alkyl or C₁-C₆-alkoxy and in which one or two methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, naphthyl, phenyl-C₁-C₆-alkyl, naphthyl-C₁-C₆-
 10 alkyl or hetaryl having 5 or 6 ring atoms and one to three heteroatoms selected from the group consisting of O, S and N, each of which is optionally substituted by a halogen atom, C₁-C₆-alkyl, C₁-C₆-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkoxy, cyano or nitro,

15 B represents H, C₁-C₁₂-alkyl or C₁-C₈-alkoxy-C₁-C₆-alkyl, or

A, B and the carbon atom to which they are bonded represent: (i) C₃-C₁₀-cycloalkyl or C₅-C₁₀-cycloalkenyl, in which in each case one methylene group is optionally
 20 replaced by O or S and each of which is optionally substituted by C₁-C₈-alkyl, C₃-C₁₀-cycloalkyl, C₁-C₈-halogenoalkyl, C₁-C₈-alkoxy, C₁-C₈-alkylthio, a halogen atom or phenyl, (ii) C₅-C₆-cycloalkyl which is substituted by: (α) an alkylenediyl group optionally containing one or two O, S
 25 or O and S, (β) an alkylenedioxyl group or an alkylenedithioyl group which, in each case with the carbon atom to which it is bonded, forms a further five- to eight-membered ring, or (iii) C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which two substituents, together with the
 30 carbon atoms to which they are bonded, represent C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkanedienediyl, each of which is optionally substituted by C₁-C₆-alkyl, C₁-C₆-alkoxy

30517-144

- 191 -

or a halogen atom and in which in each case one methylene group is optionally replaced by O or S,




D represents: (i) H, (ii) C₁-C₁₂-alkyl, C₃-C₈-alkenyl, C₃-C₈-alkinyl, C₁-C₁₀-alkoxy-C₂-C₈-alkyl, poly-C₁-C₈-alkoxy-C₂-C₈-alkyl or C₁-C₁₀-alkylthio-C₂-C₈-alkyl, each of which is optionally substituted by a halogen atom, (iii) C₃-C₈-cycloalkyl which is optionally substituted by a halogen atom, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-halogenoalkyl and in which one or two non-directly adjacent methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, hetaryl having 5 to 6 ring atoms and one or two heteroatoms selected from the group consisting of O, S and N, phenyl-C₁-C₆-alkyl or hetaryl-C₁-C₆-alkyl having 5 to 6 ring atoms and one or two heteroatoms selected from the group consisting of O, S and N, each of which is optionally substituted by a halogen atom, C₁-C₆-alkyl, C₁-C₆-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkoxy, cyano or nitro, or

A and D together represent: (i) a C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkanedienediyl group, in which in each case one methylene group is optionally replaced by O or S and each of which is optionally substituted by: (α) a halogen atom, hydroxyl or mercapto, (β) C₁-C₁₀-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₃-C₇-cycloalkyl, phenyl or benzyloxy, each of which is optionally substituted by a halogen atom, (γ) a further C₃-C₆-alkanediyl, C₃-C₆-alkenediyl or C₄-C₆-alkanedienediyl group forming a fused ring and in which one methylene group in each case is optionally replaced by O or S and which is optionally substituted by C₁-C₆-alkyl, or (δ) in which two adjacent substituents, together with the carbon atoms to which they are bonded, optionally form a further saturated or unsaturated


30517-144

- 192 -


carbocycle having 5 or 6 ring atoms, or (ii) a C₃-C₆-alkanedyl or C₃-C₆-alkenedyl group which in each case optionally contains one of the following groups:




10



G represents: (a) in the case in which Het
15 represents one of the radicals (1), (2), (3), (5) or (6), H,
or, in the case in which Het represents one of the radicals
(1), (2), (3), (4), (5) or (6), one of the groups:



20



wherein:

E represents a metal ion equivalent or an ammonium ion,

30517-144

- 193 -

L represents O or S,

M represents O or S,

R¹ represents: (i) C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₁-C₈-alkyl, C₁-C₈-alkylthio-C₁-C₈-alkyl or poly-C₁-C₈-alkoxy-C₁-C₈-alkyl, each of which is optionally substituted by a halogen atom, (ii) C₃-C₈-cycloalkyl which is optionally substituted by a halogen atom, C₁-C₆-alkyl or C₁-C₆-alkoxy and in which one or two methylene groups are optionally replaced by O, S or O and S, (iii) phenyl which is optionally substituted by a halogen atom, cyano, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl, C₁-C₆-halogenoalkoxy, C₁-C₆-alkylthio or C₁-C₆-alkylsulphonyl, (iv) phenyl-C₁-C₆-alkyl which is optionally substituted by a halogen atom, nitro, cyano, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl or C₁-C₆-halogenoalkoxy, (v) 5- or 6-membered hetaryl having one or two heteroatoms selected from the group consisting of O, S and N, and which is optionally substituted by a halogen atom or C₁-C₆-alkyl, (vi) phenoxy-C₁-C₆-alkyl which is optionally substituted by a halogen atom or C₁-C₆-alkyl, or (vii) 5- or 6-membered hetaryloxy-C₁-C₆-alkyl having one or two heteroatoms selected from the group consisting of O, S and N, and which is optionally substituted by a halogen atom, amino or C₁-C₆-alkyl,

R² represents: (i) C₁-C₂₀-alkyl, C₂-C₂₀-alkenyl, C₁-C₈-alkoxy-C₂-C₈-alkyl or poly-C₁-C₈-alkoxy-C₂-C₈-alkyl, each of which is optionally substituted by a halogen atom, (ii) C₃-C₈-cycloalkyl which is optionally substituted by a halogen atom, C₁-C₆-alkyl or C₁-C₆-alkoxy, or (iii) phenyl or benzyl, each of which is optionally substituted by a halogen atom, cyano, nitro, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-halogenoalkyl or C₁-C₆-halogenoalkoxy,

30517-144

- 194 -

R³ represents: (i) C₁-C₈-alkyl which is optionally substituted by a halogen atom, or (ii) phenyl or benzyl, each of which is optionally substituted by a halogen atom, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-
5 halogenoalkoxy, cyano or nitro,

R⁴ and R⁵ independently of one another represent:
(i) C₁-C₈-alkyl, C₁-C₈-alkoxy, C₁-C₈-alkylamino, di-(C₁-C₈-alkyl)amino, C₁-C₈-alkylthio or C₂-C₈-alkenylthio, each of which is optionally substituted by a halogen atom, or (ii)
10 phenyl, phenoxy or phenylthio, each of which is optionally substituted by a halogen atom, nitro, cyano, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, C₁-C₄-alkylthio, C₁-C₄-halogenoalkylthio, C₁-C₄-alkyl or C₁-C₄-halogenoalkyl,

R⁶ and R⁷ independently of one another represent:
15 (i) H, (ii) C₁-C₈-alkyl, C₃-C₈-cycloalkyl, C₁-C₈-alkoxy, C₃-C₈-alkenyl or C₁-C₈-alkoxy-C₂-C₈-alkyl, each of which is optionally substituted by a halogen atom, or (iii) phenyl or benzyl, each of which is optionally substituted by a halogen atom, C₁-C₈-alkyl, C₁-C₈-halogenoalkyl or C₁-C₈-alkoxy, or

20 R⁶ and R⁷ together represent a C₃-C₆-alkylene radical which is optionally substituted by C₁-C₆-alkyl and in which a methylene group is optionally replaced by O or S,

R¹³ represents: (i) H, (ii) C₁-C₈-alkyl or C₁-C₈-alkoxy, each of which is optionally substituted by a halogen atom, (iii) C₃-C₈-cycloalkyl which is optionally substituted by a halogen atom, C₁-C₄-alkyl or C₁-C₄-alkoxy and in which a methylene group is optionally replaced by O or S, or (iv) phenyl, phenyl-C₁-C₄-alkyl or phenyl-C₁-C₄-alkoxy, each of which is optionally substituted by a halogen atom, C₁-C₆-
25 alkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkyl, C₁-C₄-
30 halogenoalkoxy, nitro or cyano,

30517-144

- 195 -

R^{14} represents H or C₁-C₈-alkyl, or

R^{13} and R^{14} together represent C₄-C₆-alkanediyl,

R^{15} and R^{16} are identical or different and represent C₁-C₆-alkyl, or

5 R^{15} and R^{16} together represent a C₂-C₄-alkanediyl radical which is optionally substituted by: (α) C₁-C₆-alkyl or C₁-C₆-halogenoalkyl or (β) phenyl which is optionally substituted by a halogen atom, C₁-C₆-alkyl, C₁-C₄-halogenoalkyl, C₁-C₆-alkoxy, C₁-C₄-halogenoalkoxy, nitro or
10 cyano,

R^{17} and R^{18} independently of one another represent:
(i) H, (ii) C₁-C₈-alkyl which is optionally substituted by a halogen atom or (iii) phenyl which is optionally substituted by a halogen atom, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₄-
15 halogenoalkyl, C₁-C₄-halogenoalkoxy, nitro or cyano, or

R^{17} and R^{18} , together with the carbon atom to which they are bonded, represent: (i) a carbonyl group, or (ii) C₅-C₇-cycloalkyl which is optionally substituted by C₁-C₄-alkyl or C₁-C₄-alkoxy and in which a methylene group is
20 optionally replaced by O or S, and

R^{19} and R^{20} independently of one another represent C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₁-C₁₀-alkoxy, C₁-C₁₀-alkylamino, C₃-C₁₀-alkenylamino, di-(C₁-C₁₀-alkyl)-amino or di-(C₃-C₁₀-alkenyl)amino.

25 2. A compound of the general formula (I) according to claim 1, wherein:

X represents C₁-C₄-alkyl;

Y represents F, Cl, Br or C₁-C₄-alkyl;

30517-144

- 196 -

Z represents F, Cl, Br or C₁-C₄-alkyl,

wherein one of the radicals Y and Z always represents F, Cl or Br and the other C₁-C₄-alkyl; and

Het represents one of the groups (1) to (6)
5 defined in claim 1, wherein:

A represents: (i) H, (ii) C₁-C₁₀-alkyl, C₂-C₆-alkenyl, C₁-C₈-alkoxy-C₁-C₆-alkyl, poly-C₁-C₆-alkoxy-C₁-C₆-alkyl or C₁-C₈-alkylthio-C₁-C₆-alkyl, each of which is optionally substituted by F or Cl, (iii) C₃-C₇-cycloalkyl
10 which is optionally substituted by F, Cl, C₁-C₄-alkyl or C₁-C₄-alkoxy and in which one or two methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, furanyl, pyridyl, imidazolyl, triazolyl, pyrazolyl, indolyl, thiazolyl, thienyl or phenyl-C₁-C₄-alkyl, each of which is
15 optionally substituted by F, Cl, Br, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, cyano or nitro,

B represents H, C₁-C₁₀-alkyl or C₁-C₆-alkoxy-C₁-C₄-alkyl, or

20 A, B and the carbon atom to which they are bonded represent: (i) C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which in each case one methylene group is optionally replaced by O or S and each of which is optionally substituted by C₁-C₆-alkyl, C₃-C₈-cycloalkyl, C₁-C₃-
25 halogenoalkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, F, Cl or phenyl, (ii) C₅-C₆-cycloalkyl which is substituted by: (α) an alkylenediyl group optionally containing one or two O or S atoms, or (β) an alkylenedioxyl group or an alkylenedithioyl group which, in each case with the carbon atom to which it
30 is bonded, forms a further 5- to 7-membered ring, or (iii) C₃-C₆-cycloalkyl or C₅-C₆-cycloalkenyl, in which two

30517-144

- 197 -



substituents, together with the carbon atoms to which they are bonded, represent C₃-C₅-alkanediyl, C₃-C₅-alkenediyl or butadienediyl, each of which is optionally substituted by C₁-C₅-alkyl, C₁-C₅-alkoxy, F, Cl or Br, and in which in each
5 case one methylene group is optionally replaced by O or S,

D represents: (i) H, (ii) C₁-C₁₀-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkinyl, C₁-C₈-alkoxy-C₂-C₆-alkyl, poly-C₁-C₆-alkoxy-C₂-C₆-alkyl or C₁-C₈-alkylthio-C₂-C₆-alkyl, each of which is optionally substituted by F or Cl, (iii) C₃-C₇-
10 cycloalkyl which is optionally substituted by F, Cl, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₂-halogenoalkyl and in which one or two non-directly adjacent methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, furanyl, imidazolyl, pyridyl, thiazolyl, pyrazolyl, pyrimidyl,
15 pyrrolyl, thienyl, triazolyl or phenyl-C₁-C₄-alkyl, each of which is optionally substituted by F, Cl, Br, C₁-C₄-alkyl, C₁-C₄-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₄-halogenoalkoxy, cyano or nitro, or



A and D together represent: (i) a C₃-C₅-alkanediyl
20 or C₃-C₅-alkenediyl group, in which in each case one methylene group is optionally replaced by O or S and each of which is optionally substituted by: (α) F, Cl, hydroxyl or mercapto, or (β) C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, phenyl or benzoyloxy, each of which is
25 optionally substituted by F or Cl, or (ii) a C₃-C₅-alkanediyl or C₃-C₅-alkenediyl group which in each case optionally contains one of the following groups:

30517-144

- 198 -




5




10


or A and D represent, in the case of a compound of the general formula (I-1), together with the atoms to which they are bonded, one of the groups AD-1 to AD-27:




AD-1




AD-2




AD-3



AD-4




AD-5




AD-6


20



AD-7




AD-8




AD-9

30517-144


- 199 -



AD-10




AD-11




AD-12


5




AD-13




AD-14




AD-15



AD-16




AD-17




AD-18


10



AD-19




AD-20




AD-21


15



AD-22




AD-23




AD-24


20



AD-25



AD-26



AD-27

30517-144

- 200 -

G is as defined in claim 1,

E, L and M are as defined in claim 1,

R¹ represents: (i) C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkylthio-C₁-C₆-alkyl or poly-
5 C₁-C₆-alkoxy-C₁-C₆-alkyl, each of which is optionally substituted by F or Cl, (ii) C₃-C₇-cycloalkyl which is optionally substituted by F, Cl, C₁-C₅-alkyl or C₁-C₅-alkoxy and in which one or two methylene groups are optionally replaced by O, S or O and S, (iii) phenyl which is
10 optionally substituted by F, Cl, Br, cyano, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₃-halogenoalkyl, C₁-C₃-halogenoalkoxy, C₁-C₄-alkylthio or C₁-C₄-alkylsulphonyl, (iv) phenyl-C₁-C₄-alkyl which is optionally substituted by F, Cl, Br, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₃-halogenoalkyl or C₁-C₃-
15 halogenoalkoxy, (v) pyrazolyl, thiazolyl, pyridyl, pyrimidyl, furanyl or thienyl, each of which is optionally substituted by F, Cl, Br or C₁-C₄-alkyl, (vi) phenoxy-C₁-C₅-alkyl which is optionally substituted by F, Cl, Br or C₁-C₄-alkyl, or (vii) pyridyloxy-C₁-C₅-alkyl, pyrimidyloxy-C₁-C₅-
20 alkyl or thiazolyloxy-C₁-C₅-alkyl, each of which is optionally substituted by F, Cl, Br, amino or C₁-C₄-alkyl,

R² represents: (i) C₁-C₁₆-alkyl, C₂-C₁₆-alkenyl, C₁-C₆-alkoxy-C₂-C₆-alkyl or poly-C₁-C₆-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by F or Cl, (ii) C₃-C₇-
25 cycloalkyl which is optionally substituted by F, Cl, C₁-C₄-alkyl or C₁-C₄-alkoxy, or (iii) phenyl or benzyl, each of which is optionally substituted by F, Cl, Br, cyano, nitro, C₁-C₄-alkyl, C₁-C₃-alkoxy, C₁-C₃-halogenoalkyl or C₁-C₃-halogenoalkoxy,

30 R³ represents: (i) C₁-C₆-alkyl which is optionally substituted by F or Cl, or (ii) phenyl or benzyl, each of which is optionally substituted by F, Cl, Br, C₁-C₅-alkyl,

30517-144

- 201 -

C₁-C₅-alkoxy, C₁-C₃-halogenoalkyl, C₁-C₃-halogenoalkoxy, cyano or nitro,

R⁴ and R⁵ independently of one another represent:

(i) C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylamino, di-(C₁-C₆-alkyl)amino, C₁-C₆-alkylthio or C₃-C₄-alkenylthio, each of which is optionally substituted by F or Cl, or (ii) phenyl, phenoxy or phenylthio, each of which is optionally substituted by F, Cl, Br, nitro, cyano, C₁-C₃-alkoxy, C₁-C₃-halogenoalkoxy, C₁-C₃-alkylthio, C₁-C₃-halogenoalkylthio, C₁-C₃-alkyl or C₁-C₃-halogenoalkyl,

R⁶ and R⁷ independently of one another represent:

(i) H, (ii) C₁-C₆-alkyl, C₃-C₆-cycloalkyl, C₁-C₆-alkoxy, C₃-C₆-alkenyl or C₁-C₆-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by F or Cl, or (iii) phenyl or benzyl, each of which is optionally substituted by F, Cl, Br, C₁-C₅-halogenoalkyl, C₁-C₅-alkyl or C₁-C₅-alkoxy, or

R⁶ and R⁷ together represent a C₃-C₆-alkylene radical which is optionally substituted by C₁-C₄-alkyl and in which a methylene group is optionally replaced by O or S,

R¹³ represents: (i) H, (ii) C₁-C₆-alkyl or C₁-C₆-alkoxy, each of which is optionally substituted by F or Cl, (iii) C₃-C₇-cycloalkyl which is optionally substituted by F, C₁-C₂-alkyl or C₁-C₂-alkoxy and in which a methylene group is optionally replaced by O or S, or (iv) phenyl, phenyl-C₁-C₃-alkyl or phenyl-C₁-C₂-alkoxy, each of which is optionally substituted by F, Cl, Br, C₁-C₅-alkyl, C₁-C₅-alkoxy, C₁-C₂-halogenoalkyl, C₁-C₂-halogenoalkoxy, nitro or cyano,

R¹⁴ represents H or C₁-C₆-alkyl, or

R¹³ and R¹⁴ together represent C₄-C₆-alkanediyl,

30517-144

- 202 -

R¹⁵ and R¹⁶ are identical or different and represent C₁-C₄-alkyl, or

R¹⁵ and R¹⁶ together represent a C₂-C₃-alkanediyl radical which is optionally substituted by: (α) C₁-C₄-alkyl or C₁-C₄-halogenoalkyl or (β) phenyl which is optionally substituted by F, Cl, Br, C₁-C₄-alkyl, C₁-C₂-halogenoalkyl, C₁-C₄-alkoxy, C₁-C₂-halogenoalkoxy, nitro or cyano,

R¹⁷ and R¹⁸ independently of one another represent: (i) H, (ii) C₁-C₈-alkyl which is optionally substituted by F or Cl, or (iii) phenyl which is optionally substituted by F, Cl, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₂-halogenoalkyl, C₁-C₂-halogenoalkoxy, nitro or cyano, or

R¹⁷ and R¹⁸, together with the carbon atom to which they are bonded, represent C₅-C₆-cycloalkyl which is optionally substituted by C₁-C₃-alkyl or C₁-C₃-alkoxy and in which a methylene group is optionally replaced by O or S, and

R¹⁹ and R²⁰ independently of one another represent C₁-C₆-alkyl, C₂-C₆-alkenyl, C₁-C₆-alkoxy, C₁-C₆-alkylamino, C₃-C₆-alkenylamino, di-(C₁-C₆-alkyl)amino or di-(C₃-C₆-alkenyl)amino.

3. A compound of the general formula (I) according to claim 1, wherein:

X represents methyl, ethyl, n-propyl or iso-propyl,

Y represents F, Cl, Br, methyl, ethyl, n-propyl or iso-propyl,

Z represents F, Cl, Br, methyl, ethyl, n-propyl or iso-propyl,

30517-144

- 203 -

wherein one of the radicals Y and Z always represents F, Cl or Br and the other methyl, ethyl, n-propyl or iso-propyl; and

Het represents one of the groups (1) to (6)
5 defined in claim 1,

A represents: (i) H, (ii) C₁-C₈-alkyl, C₂-C₄-alkenyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, poly-C₁-C₄-alkoxy-C₁-C₄-alkyl or C₁-C₆-alkylthio-C₁-C₄-alkyl, each of which is optionally substituted by F or Cl, (iii) C₃-C₆-cycloalkyl
10 which is optionally substituted by F, Cl, methyl or methoxy and in which one or two methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, furanyl, thienyl, pyridyl or benzyl, each of which is optionally substituted by F, Cl, Br, methyl, ethyl, n-propyl, iso-
15 propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro,

B represents H, C₁-C₈-alkyl or C₁-C₄-alkoxy-C₁-C₂-alkyl, or

A, B and the carbon atom to which they are bonded
20 represent: (i) C₃-C₈-cycloalkyl or C₅-C₈-cycloalkenyl, in which in each case one methylene group is optionally replaced by O or S and each of which is optionally substituted by methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl, tert-butyl, cyclohexyl,
25 trifluoromethyl, methoxy, ethoxy, n-propoxy, iso-propoxy, butoxy, iso-butoxy, sec-butoxy, tert-butoxy, methylthio, ethylthio, F, Cl or phenyl, (ii) C₅-C₆-cycloalkyl which is substituted by: (α) an alkylenediyl group optionally containing an O or S atom or (β) an alkylenedioxyl group
30 which, with the carbon atom to which it is bonded, forms a further five- or six-membered ring, or (iii) C₃-C₆-cycloalkyl



30517-144

- 204 -

or C₅-C₆-cycloalkenyl, in which two substituents, together with the carbon atoms to which they are bonded, represent C₃-C₄-alkanediyl, C₃-C₄-alkenediyl or butadienediyl, in which in each case one methylene group is optionally replaced by O
5 or S,

D represents: (i) H, (ii) C₁-C₈-alkyl, C₃-C₄-alkenyl, C₃-C₄-alkinyl, C₁-C₆-alkoxy-C₂-C₄-alkyl, poly-C₁-C₄-alkoxy-C₂-C₄-alkyl or C₁-C₄-alkylthio-C₂-C₄-alkyl, each of which is optionally substituted by F or Cl, (iii) C₃-C₆-
10 cycloalkyl, each of which is optionally substituted by F or Cl, and in which one or two non-directly adjacent methylene groups are optionally replaced by O, S or O and S, or (iv) phenyl, furanyl, pyridyl, thienyl or benzyl, each of which is optionally substituted by F, Cl, Br, methyl, ethyl,
15 n-propyl, iso-propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro, or

A and D together represent: (i) a C₃-C₅-alkanediyl or C₃-C₅-alkenediyl group, in which in each case a methylene group is optionally replaced by O or S and each of which is
20 optionally substituted by: (α) F, Cl, hydroxyl or mercapto or (β) C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, phenyl or benzyloxy, each of which is optionally substituted by F or Cl, or (ii) a C₃-C₅-alkanediyl or C₃-C₅-alkenediyl group which in each case optionally contains one
25 of the following groups:




30517-144


- 205 -

or A and D represent, in the case of a compound of the general formula (I-1), together with the atoms to which they are bonded, one of the following groups:


5




AD-1




AD-2




AD-4



AD-6




AD-8




AD-10


10



AD-12




AD-14




AD-15


15



AD-17



AD-18



AD-27

G is as defined in claim 1,

E, L and M are as defined in claim 1,

R¹ represents: (i) C₁-C₁₄-alkyl, C₂-C₁₄-alkenyl, C₁-C₄-alkoxy-C₁-C₆-alkyl, C₁-C₄-alkylthio-C₁-C₆-alkyl or poly-
 20 C₁-C₄-alkoxy-C₁-C₄-alkyl, each of which is optionally substituted by F or Cl, (ii) C₃-C₆-cycloalkyl which is optionally substituted by F, Cl, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, tert-butyl, methoxy, ethoxy, n-propoxy or iso-propoxy and in which one or two methylene
 25 groups are optionally replaced by O, S or O and S, (iii)

30517-144

- 206 -

phenyl which is optionally substituted by F, Cl, Br, cyano, nitro, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl, trifluoromethoxy, methylthio, ethylthio, methylsulphonyl or ethylsulphonyl, (iv) benzyl which is
5 optionally substituted by F, Cl, Br, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy, (v) furanyl, thienyl or pyridyl, each of which is optionally substituted by F, Cl, Br, methyl or ethyl, (vi) phenoxy-C₁-C₄-alkyl which is optionally
10 substituted by F, Cl, methyl or ethyl, or (vii) pyridyloxy-C₁-C₄-alkyl, pyrimidyloxy-C₁-C₄-alkyl or thiazolyloxy-C₁-C₄-alkyl, each of which is optionally substituted by F, Cl, amino, methyl or ethyl,

R² represents: (i) C₁-C₁₄-alkyl, C₂-C₁₄-alkenyl,
15 C₁-C₄-alkoxy-C₂-C₆-alkyl or poly-C₁-C₄-alkoxy-C₂-C₆-alkyl, each of which is optionally substituted by F or Cl, (ii) C₃-C₆-cycloalkyl which is optionally substituted by F, Cl, methyl, ethyl, n-propyl, iso-propyl or methoxy, or (iii) phenyl or benzyl, each of which is optionally substituted by F, Cl,
20 cyano, nitro, methyl, ethyl, n-propyl, i-propyl, methoxy, ethoxy, trifluoromethyl or trifluoromethoxy,

R³ represents: (i) methyl, ethyl, propyl, or isopropyl, each of which is optionally substituted by F or Cl, or (ii) phenyl or benzyl, each of which is optionally
25 substituted by F, Cl, Br, methyl, ethyl, propyl, iso-propyl, tert-butyl, methoxy, ethoxy, isopropoxy, tert-butoxy, trifluoromethyl, trifluoromethoxy, cyano or nitro,

R⁴ and R⁵ independently of one another represent:
(i) C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylamino, di-(C₁-C₄-
30 alkyl)amino or C₁-C₄-alkylthio, each of which is optionally substituted by F or Cl, or (ii) phenyl, phenoxy or phenylthio, each of which is optionally substituted by F,

30517-144

- 207 -

Cl, Br, nitro, cyano, methyl, methoxy, trifluoromethyl or trifluoromethoxy,

R^6 and R^7 independently of one another represent:

(i) H, (ii) C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl, C_1 - C_4 -alkoxy, C_3 - C_4 -
5 alkenyl or C_1 - C_4 -alkoxy- C_2 - C_4 -alkyl, each of which is optionally substituted by F or Cl, or (iii) phenyl or benzyl, each of which is optionally substituted by F, Cl, Br, methyl, methoxy or trifluoromethyl, or

R^6 and R^7 together represent a C_5 - C_6 -alkylene
10 radical which is optionally substituted by methyl or ethyl and in which a methylene group is optionally replaced by O or S,

R^{13} represents: (i) H, (ii) C_1 - C_4 -alkyl or C_1 - C_4 -
alkoxy, each of which is optionally substituted by F or Cl,
15 (iii) C_3 - C_6 -cycloalkyl, or (iv) phenyl, phenyl- C_1 - C_2 -alkyl or benzyloxy, each of which is optionally substituted by F, Cl, Br, methyl, ethyl, iso-propyl, tert-butyl, methoxy, ethoxy, iso-propoxy, tert-butoxy, trifluoromethyl, trifluoromethoxy, nitro or cyano,

20 R^{14} represents H or C_1 - C_4 -alkyl, or

R^{13} and R^{14} together represent C_4 - C_6 -alkanediyl,

R^{15} and R^{16} are identical or different and represent methyl or ethyl, or

R^{15} and R^{16} together represent a C_2 - C_3 -alkanediyl
25 radical which is optionally substituted by: (α) methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or tert-butyl or (β) phenyl which is optionally substituted by F, Cl, methoxy, trifluoromethyl, trifluoromethoxy, nitro or cyano.


30517-144

- 208 -

4. A process for the preparation of a compound of the general formula (I) as defined in any one of claims 1 to 3, wherein:


(A) a compound of the general formula (I-1-a):

5



wherein A, B, D, X, Y and Z are as defined in any one of
10 claims 1 to 3, is obtained when a compound of general
formula (II):


15



wherein A, B, D, X, Y and Z are as defined in any one of
claims 1 to 3, and R⁸ represents alkyl, is intramolecularly
condensed in the presence of a diluent and in the presence
of a base;

20


(B) a compound of general formula (I-2-a):



30517-144


- 209 -

wherein A, B, X, Y and Z are as defined in any one of claims 1 to 3, is obtained when a compound of general formula (III):




wherein A, B, X, Y and Z are as defined in any one of claims 1 to 3, and R⁸ is as defined in (A), is intramolecularly condensed in the presence of a diluent and in the presence of a base;

(C) a compound of general formula (I-3-a):



wherein A, B, X, Y and Z are as defined in any one of claims 1 to 3, is obtained when a compound of general formula (IV):




30517-144

- 210 -

wherein A, B, X, Y and Z are as defined in any one of claims 1 to 3, R⁸ is as defined in (A) and W represents H, a halogen atom, alkyl or alkoxy, is cyclized, optionally in the presence of a diluent and in the presence of an acid;


5 (E) a compound of general formula (I-5-a):



10 wherein A, D, X, Y and Z are as defined in any one of claims 1 to 3, is obtained when a compound of general formula (VIII):



15 wherein A and D are as defined in any one of claims 1 to 3, is reacted with a compound of formula (V):




20 wherein X, Y and Z are as defined in any one of claims 1 to 3, and Hal represents a halogen atom, optionally in the presence of a diluent and optionally in the presence of an acid acceptor;

(F) a compound of general formula (I-6-a):

30517-144

- 211 -




wherein A, X, Y and Z are as defined in any one of claims 1
5 to 3, is obtained when a compound of general formula (IX):




wherein A is as defined in any one of claims 1 to 3, is
reacted with the compound of the general formula (V) as
10 defined in (E), optionally in the presence of a diluent and
optionally in the presence of an acid acceptor, and

optionally then a compound thus obtained of the
general formulae (I-1-a), (I-2-a), (I-3-a), (I-5-a) or
(I-6-a) from (A), (B), (C), (E) or (F), or a compound of
15 general formula (I-4-a):



wherein A, D, X, Y and Z are as defined in any one of
20 claims 1 to 3, in each case are reacted:

(G) (α) with an acid halide of general formula (X):




30517-144

- 212 -


wherein R^1 is as defined in any one of claims 1 to 3, and Hal is as defined in (E), or

(β) with a carboxylic anhydride of general formula (XI):



wherein R^1 is as defined in any one of claims 1 to 3, optionally in the presence of a diluent and optionally in the presence of an acid-binding agent; or

(H) with a chloroformic acid ester or a
10 chloroformic acid thioester of general formula (XII)



wherein R^2 and M are as defined in any one of claims 1 to 3, optionally in the presence of a diluent and optionally in the presence of an acid-binding agent; or

15 (I) (α) with a chloromonothioformic acid ester or a chlorodithioformic acid ester of general formula (XIII):



wherein M and R^2 are as defined in any one of claims 1 to 3,
20 optionally in the presence of a diluent and optionally in the presence of an acid-binding agent, or

(β) with carbon disulphide and then with a compound of general formula (XIV):



30517-144

- 213 -

wherein R^2 is as defined in any one of claims 1 to 3, and Hal represents Cl, Br or I, optionally in the presence of a diluent and optionally in the presence of a base; or

(J) with a sulphonyl chloride of general
5 formula (XV):





wherein R^3 is as defined in any one of claims 1 to 3, optionally in the presence of a diluent and optionally in the presence of an acid-binding agent; or

10 (K) with a phosphorus compound of general formula (XVI):



15 wherein L, R^4 and R^5 are as defined in any one of claims 1 to 3, and Hal is as defined in (E), optionally in the presence of a diluent and optionally in the presence of an acid-binding agent; or

(L) with a metal compound or an amine of general
20 formula (XVII) or (XVIII):



wherein:

Me represents a mono- or divalent metal,


25 t represents the number 1 or 2, and

30517-144

- 214 -


R^{10} , R^{11} , R^{12} independently of one another represent H or alkyl, optionally in the presence of a diluent; or

(M) (α) with an isocyanate or an isothiocyanate of general formula (XIX):



wherein R^6 and L are as defined in any one of claims 1 to 3, optionally in the presence of a diluent and optionally in the presence of a catalyst, or


(β) with a carbamoyl chloride or a thiocarbamoyl chloride of general formula (XX):



wherein L, R^6 and R^7 are as defined in any one of claims 1 to 3, optionally in the presence of a diluent and optionally in the presence of an acid-binding agent.

5. The compound of the general formula (II) as defined in claim 4.

6. A compound of general formula (XXIII):




wherein A, B, D, X, Y and Z are as defined in any one of claims 1 to 3.

30517-144

- 215 -

7. A compound of general formula (XXXI):



5

wherein A, B, D, X, Y and Z are as defined in any one of claims 1 to 3.

8. The compound of the general formula (III) as defined in claim 4.

10 9. The compound of the general formula (IV) as defined in claim 4

10. The compound of the general formula (V) as defined in claim 4.

11. A pesticidal and weedkilling composition,
15 comprising at least one compound of the general formula (I) as defined in any one of claims 1 to 3, and an extender, a surface-active agent or a mixture thereof.

12. Use of a compound of the general formula (I) as defined in any one of claims 1 to 3, or a composition as
20 defined in claim 11, for the control of pests and weeds.

13. A method for the control of pests and weeds,
wherein a compound of the general formula (I) as defined in any one of claims 1 to 3, or a composition as defined in claim 11, is allowed to act on the pests, the weeds or their
25 habitat.

14. A process for the production of a pesticidal and weedkilling composition, wherein a compound of the general

30517-144

- 216 -

formula (I) as defined in any one of claims 1 to 3 is mixed with an extender, a surface-active agent or a mixture thereof.

15. Use of a compound of the general formula (I) as
5 defined in any one of claims 1 to 3, for the production of a
pesticidal and weedkilling composition.

FETHERSTONHAUGH & CO.

OTTAWA, CANADA

PATENT AGENTS

