



- (51) **International Patent Classification:**
A61K 31/609 (2006.01) *A61P 31/04* (2006.01)
A61K 31/612 (2006.01)
- (21) **International Application Number:** PCT/EP2016/061968
- (22) **International Filing Date:** 27 May 2016 (27.05.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:** 1509326.3 29 May 2015 (29.05.2015) GB
- (71) **Applicant:** ANTIBIOTX APS [DK/DK]; Kogle Allé 5, DK-2970 Hørsholm (DK).
- (72) **Inventors:** SOMMER, Morten, Otto, Alexander,; Parsbergsvej 40, 2830 Virum (DK). TOFT-KEHLER, Rasmus, Vendler,; Wilhelm Marstrands Gade 22, 2100 København Ø (DK). SIMON, Daniel, Jean, Jacques,; Toftekaersvej 170, 2860 Søborg (DK).
- (74) **Agents:** HGF LIMITED et al.; 1 City Walk, Leeds, West Yorkshire LS11 9DX (GB).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** HALOGENATED SALICYLANILIDES FOR TREATING CLOSTRIDIUM INFECTIONS




Figure 1

(57) **Abstract:** The present invention relates to halogenated salicylanilides, or pharmaceutically acceptable salts or esters thereof, for use in the treatment of an infection in a subject caused by Clostridium bacteria, particularly a *C. difficile* infection. The halogenated salicylanilides are expected to be useful in the treatment of *C. difficile* associated diseases including *C. difficile* associated diarrhoea and *C. difficile* associated colitis.



HALOGENATED SALICYLANILIDES FOR TREATING CLOSTRIDIUM INFECTIONS

NOVEL USE

[0001] This invention relates to halogenated salicylanilides for use in the prevention or treatment of infections caused by *Clostridium bacteria*, particularly *Clostridium difficile*.

BACKGROUND OF THE INVENTION

[0002] *Clostridium* is a genus of spore forming Gram-positive bacteria that grow under anaerobic conditions comprising more than 100 species. There are four main species responsible for diseases in humans and other warm-blooded animals: *C. botulinum*, an organism producing a toxin in food or wounds that causes botulism; *C. difficile*, which can cause pseudomembranous colitis, toxic megacolon and antibiotic associated diarrheas; *C. tetani*, which is the causative organism of tetanus; and *C. perfringens*, which can cause enterotoxemia, necrotizing enteritis, and gas gangrene.

[0003] *C. perfringens* is ubiquitous in the environment and is found in soil, dust, raw ingredients such as spices used in food processing, and in the intestines of humans and animals. It produces over 15 different toxins resulting in various enteric conditions. *C. perfringens* infections can also cause gut health problems in broiler flocks with significant negative economic consequences.

[0004] *C. difficile* is an opportunistic gram positive, anaerobic, spore forming bacillus, and causes *Clostridium difficile* infections (CDI) such as antibiotic-associated diarrhoea (CDAD) and colitis which burdens healthcare systems across the globe. In the last decade, rates of *C. difficile* infections have increased dramatically, particularly hospital-acquired infection (nosocomial infection), resulting in increased morbidity, an increased incidence of complications requiring colectomy, and rising mortality.

[0005] It is estimated that 3 to 15% of the normal population is infected with *C. difficile*. However, rates of infection are much higher in hospitalised patients. *C. difficile* colonises the intestine and in many subjects infected the bacteria lives in equilibrium with other gut flora and is asymptomatic. However, if the homeostasis of the normal intestinal flora is disturbed, for example as a result of previous antibiotic use, the use of drugs which alter the gastric pH (for example proton pump inhibitors), or gastrointestinal surgery, symptomatic CDI can arise as a result of the proliferation of the *C. difficile* in the intestine. Toxins produced by the *C. difficile* disrupt the colonic epithelium, leading to an inflammatory response and clinical symptoms varying from mild diarrhoea to severe life-threatening pseudomembranous colitis.

[0006] *C. difficile* bacteria produce toxins, which can cause inflammation and damage to the lining of the lower gastro-intestinal tract, including the colon. There are a number of different strains of *C. difficile*, some of which can cause more serious illnesses than others.

Strain NAP1/027/BI 027 (NAP1/027) produces particularly high levels of toxins and is associated with particularly severe CDI and high levels of mortality.

[0007] *C. difficile* infections are particularly associated with the clinical use of broad spectrum antibiotics, for example clindamycin, cephalosporins and amoxicillin-clavulanic acid). Fluoroquinolone antibiotics have been identified as a particular risk factor for CDI. Antibiotics commonly used to treat a primary infection in a subject (for example a urinary infection, a skin infection or other infection), kill the bacteria that cause the primary infection. However, they may also kill many of the bacteria present in the flora of the GI tract. Because *C. difficile* bacteria are not affected by many commonly used antibiotics this can result in the proliferation of *C. difficile* in the intestine and the presence of high levels of associated toxins resulting in the emergence of symptoms of a CDI.

[0008] *C. difficile* infection is the most common infectious cause of nosocomial diarrhoea in elderly patients, accounting for 15% to 25% of all cases of antibiotic-induced diarrhoea. Patients undergoing total joint arthroplasty are at particular risk of CDI because of the advanced age of the patients, the use of prophylactic antibiotic coverage in the perioperative period, multiple comorbid conditions, and length of hospital stay required for recovery.

[0009] The treatment of *C. difficile* infections depends on the severity of the associated symptoms or disease. Generally asymptomatic infections are not treated. However, if symptoms develop treatments are generally required to reduce the symptoms and prevent the infection from worsening.

[0010] Generally a first step in the treatment of CDI is the cessation of the inciting antibiotic. Treatment with concomitant antibiotics (i.e. antibiotics other than those given to treat *C. difficile* infection) is associated both with significant prolongation of diarrhoea and with an increased risk of recurrent CDI. If concomitant antibiotics are essential for treatment of the primary infection, it is generally prudent, if possible, to use an antibiotic therapy that is less frequently implicated in antibiotic-associated CDI, such as parenteral aminoglycosides, sulfonamides, macrolides, vancomycin, or tetracycline (Läkartidningen, 103(46), 2006).

[0011] *C. difficile* infection, such as CDAD is usually treated with metronidazole or oral vancomycin. A new antibiotic against *C. difficile* has recently been approved, fidaxomicin (OPT-80, PAR-101), a macrocyclic antibiotic. In phase III clinical trials, fidaxomicin was non-inferior to vancomycin in achieving clinical cure of CDAD. Fidaxomicin treatment was also superior to vancomycin in preventing recurrence of CDAD. These results, combined with the ease of administration and a somewhat better safety profile has made fidaxomicin an attractive treatment option for treating CDAD. (Louie, T. J., Miller, M. A., Mulvane, K. M., Weiss, K., Lenten, A. Shoe, Y.K. (2011). Fidaxomicin versus Vancomycin for *Clostridium*

difficile infection. New England Journal of Medicine, 364, 422-431). However, resistance towards metronidazole, vancomycin and fidaxomicin has been observed.

[0012] Rifamycins and derivatives, for example rifampicin and rifaximin, have been successfully used to treat recurrent CDI. However, rapid spontaneous resistance evolution has also been observed with this class of antibiotic and the spread of, for example rifampicin-resistant *C. difficile* in hospitals is an increasing concern.

[0013] Teicoplanin (although not widely available and expensive) is another antibiotic with high reported efficacy against CDI, and limited data suggest that it may be effective in recurrent CDI.

[0014] Patients who have had one CDI are at risk of recurrence of the infection. The rate of recurrent CDI (RCDI) is estimated to be 15% to 30%. Patients with recurrent *C. difficile* infections in hospitals and the community constitute an increasing treatment problem. Whilst most patients with a first infection respond to either metronidazole or oral vancomycin, current therapeutic approaches to recurrent *C. difficile* infections are prone to failure, increasing the risk of antibiotic resistance emerging. Most treatment guidelines recommend prolonged oral vancomycin pulse and/or tapering dosage regimens. However, evidence supporting the effectiveness of such dosage regimens is limited.

[0015] The spores formed by *C. difficile* are thought to be the primary mechanism for the transmission or spread of infection. Additionally spores present in the colon of a patient may be responsible for recurrence of *C. difficile* infections, even after elimination of the bacterial with antibiotic treatment. Fidaxomicin has been shown to inhibit *C. difficile* spore formation (Babakhani et al, S162 CID 2012:55 (suppl 2)). There is however a need for additional agents that can inhibit spore formation and thus minimise the risk of transmission and/or recurrence of *C. difficile* infections.

[0016] US patent, US 8,618,100 discloses chromanyl derivatives described as having antibacterial activity against *Clostridium* bacteria, in particular *Clostridium perfringens*.

[0017] PCT patent application WO2008/039640 discloses the compound 5-[3-((R)(+)-6,8-dibromo-chroman-4-ylamino)-propylamino]-4H-thieno[3,2-b]pyridine-7-one, which is also known as REP3123, and its antibacterial activity against *Clostridium difficile*. In vitro tests of the antibacterial activity of the REP3123 compound demonstrate that said compound is active against bacteria of the *Clostridium* genus however REP3123 also has antibacterial activity against a wide variety of bacteria that are present in the gut.

[0018] US patent number US 8,796,292 discloses that certain 7-substituted-2-(benzylamino)-6-oxapurines have potent activity against the growth of the intestinal anaerobe *C. difficile*, but weak activity against other, intestinal Gram-positive anaerobes. The

compounds are described to be useful in reducing the likelihood of developing or to treat *C. difficile* infections.

[0019] PCT application WO2014135891 describes the rectal administration of compositions comprising fidaxomicin. The compositions are described as useful for the treatment or maintenance of remission of infections such as diarrhea caused by *C. difficile*.

[0020] PCT application WO2012/050826 describes the use of reutericyclin or reutericyclin analogs in order to kill *C. difficile* organisms and thus alleviate the signs and symptoms of *C. difficile* infection.

[0021] There is however a need for new treatments for *C. difficile*.

[0022] Halogenated salicylanilides such as niclosamide, closantel and rafoxanide, are important anthelmintics that are used extensively in the control of *Haemonchus* spp. and *Fasciola* spp. infestation in sheep and cattle, and *Oestrus ovis* in sheep.

[0023] Niclosamide is commercially available in a number of formulations including, but not limited to Bayer73®, Bayer2353®, Bayer25648®, Baylucid®, Baylucide®, Cestocid®, Clonitralid, Dichlosale®, Fenasal®, HL 2447®, lomesan®, lomezan®, Manosil®, Nasemo®, Niclosamid®, Phenasal®, Tredemine®, Sulqui®, Vermitid®, Vermitin® and Yomesan®.

[0024] Niclosamide has been proposed as a possible systemic treatment for chronic lung infections caused by the proteobacterium *Pseudomonas aeruginosa* and the actinobacterium *Mycoplasmum tuberculosis* (F. Imperi et al., *Antimicrobial, Agents and Chemotherapy*, 557(2), 996-1005 (2013)).

[0025] J. Vinsova et al. (*Molecules*, vol. 12, no. 1, pp. 1–12, 2007; *Bioorganic and Medicinal Chemistry Letters*, vol. 19, no. 2, pp. 348–351, 2009; *European Journal of Medicinal Chemistry*, vol. 45, no. 12, pp. 6106–6113, 2010) describe certain antibacterial activity of salicylanilides, however, there is no disclosure of the treatment of CDI.

[0026] Ghazi et al. (*Zentralbl. Mikrobiol.* 141 (1986), 225-232) have tested the antibacterial effect and toxicity of synthesized salicylanilide derivatives against *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

[0027] M. J. Macielag et al. tested for antibacterial activity of closantel and related derivatives against the drug-resistant organisms, methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus faecium* (VREF) (*J. Med. Chem.*, 41(16), 2939-45 (1998)).

[0028] D.J. Hlasta et al. found that closantel had antibacterial activity against drug resistant *S. aureus* and *E. faecium* (*Bioorg. Med. Chem. Letters*, 8(14), 1923-28 (1998)).

[0029] R. Rajamuthiah et al. (PloS One, 2014, 9(2): e89189) identified closantel as a hit in a high throughput liquid screening assay and found anti-staphylococcal activity of closantel against vancomycin-resistant *S. aureus* isolates and other Gram-positive bacteria.

[0030] R. Rajamuthiah et al. (PloS One, 2015, 10(4):e0124595) describe that niclosamide and oxyclozanide have activity against MRSA.

[0031] Pauk et al. Bioorg. & Med. Chem. 23, 6574-6581 (2013), discloses the *in-vitro* anti-microbial activity of certain halogenated salicylanilides and derivatives.

[0032] WO 2008/155535 describes the use of halogenated salicylanilides for the treatment of acne resulting from propioni bacterial infection.

BRIEF SUMMARY OF THE DISCLOSURE

[0033] It has been found that halogenated salicylanilides (for example tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clioxanide, dibromosalan, tribromosalan and niclosamide) are active against *Clostridium* bacteria, particularly *C. difficile* and may be useful in treating and/or preventing or reducing *Clostridium* infection and possible reoccurrence of the infection. Use of halogenated salicylanilides may also reduce the rate of developing antibiotic resistance compared to known antibiotics used for the treatment of *Clostridium* infections.

[0034] In accordance with the present invention, there is provided a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof, for use in the treatment of an infection in a subject caused by *Clostridium* bacteria.

[0035] The infection may be caused by a *Clostridium* bacteria selected from for example, *C. perfringens*, *C. difficile*, *C. botulinum*, *C. tetani*, *C. absonum*, *C. argentinense*, *C. baratii*, *C. bifementans*, *C. beijerinckii*, *C. butyricum*, *C. cadaveris*, *C. carnis*, *C. celatum*, *C. clostridioforme*, *C. cochlearium*, *C. cocleatum*, *C. fallax*, *C. ghonii*, *C. glycolicum*, *C. haemolyticum*, *C. hastiforme*, *C. histolyticum*, *C. indolis*, *C. innocuum*, *C. irregulare*, *C. leptum*, *C. limosum*, *C. malenominatum*, *C. novyi*, *C. oroticum*, *C. oedematiens*, *C. paraputrificum*, *C. piliforme*, *C. putrefasciens*, *C. ramosum*, *C. septicum*, *C. sordelii*, *C. sphenoides*, *C. spiroforme*, *C. sporogenes*, *C. subterminale*, *C. symbiosum*, *C. tertium* or *C. tetani*.

[0036] In one embodiment the bacteria causing the infection is not *C. perfringens*.

[0037] The infection is particularly an infection caused by *C. difficile*.

[0038] Infection by *Clostridium difficile* may result in a *C. difficile* disease in the subject. The *C. difficile* disease may be, for example, diarrhoea, colitis (including pseudomembranous colitis) or toxic megacolon. The *C. difficile* infection may be *C. difficile* associated diarrhoea.

The *C. difficile* infection may be *C. difficile* associated colitis, for example pseudomembranous colitis. The *C. difficile* infection may be *C. difficile* associated bloating. The *C. difficile* infection may be *C. difficile* associated abdominal pain.

[0039] *C. difficile* infection in a subject generally arises as a result of the treatment of an infection with an antibiotic. The use of antibiotics to treat an infection will kill the organism causing the underlying infection. However, the antibiotic may also kill many of the bacteria present in the GI tract. The disruption of the normal gut flora can result in the proliferation of the *C. difficile* and emergence of effects of the infection including diarrhoea and colitis. Accordingly it may be that the halogenated salicylanilide is for use in the treatment of an antibiotic induced *Clostridium* infection, particularly an antibiotic induced *C. difficile* infection.

[0040] It may be that the antibiotic responsible for the antibiotic induced *Clostridium* infection is an antibiotic other than a halogenated salicylanilide. It may be that the antibiotic responsible for inducing the *Clostridium* infection is an antibiotic used to treat a primary infection in the body other than a *Clostridium* infection (e.g. other than a *C. difficile* infection). For example, the primary infection may be a skin infection, a urinary tract infection, a lung infection or a bone infection. The antibiotic responsible for inducing the infection may be a broad spectrum antibiotic which may be active against Gram positive and/or Gram negative organisms. The antibiotic may be selected from clindamycin, a cephalosporin (for example cefotaxime and ceftidime), ampicillin, amoxicillin and a quinolone (for example a fluoroquinolone, optionally ciprofloxacin or levofloxacin). For example, the antibiotic induced *C. difficile* infection may be caused by a fluoroquinolone antibiotic, including but not limited to ciprofloxacin or levofloxacin.

[0041] Although *C. difficile* infections are generally caused by prior use of antibiotics to treat an underlying infection, *C. difficile* infections may also be arise without the prior use of an antibiotic. For example, a reduction in the acidity of the stomach can result in colonization of the normally sterile upper gastrointestinal tract. The use of gastric acid suppressive agents, such as proton pump inhibitors (PPIs) and histamine H₂-receptor antagonists (H₂RAs) may therefore be associated with an increased risk of *C. difficile* colonization and subsequent development of CDAD. PPIs include, but are not limited to, omeprazole (Losec, Prilosec, Zegerid), lansoprazole (Prevacid, Zoton, Inhibitol), esomeprazole (Nexium), pantoprazole (Protonix, Somac, Pantoloc, Pantozol, Zurcal, Pan) and rabeprazole (Rabecid, Aciphex, Pariet, Rabeloc). H₂RAs include, but are not limited to, cimetidine (Tagamet), ranitidine (Zinetac, Zantac), famotidine, (Pepcidine, Pepcid), roxatidine (Roxit) and nizatidine (Tazac, Axid). It may be that the halogenated salicylanilide is for use in the treatment of a *Clostridium* infection (for example a *C. difficile* infection) induced by a gastric acid suppressive agent. Accordingly it may be that the halogenated salicylanilide is for use in

treating a *Clostridium* infection in a subject that is or has been treated with a gastric acid suppressive agent, for example a PPI. Accordingly the halogenated salicylanilide may be for use in treating a *Clostridium* infection in a subject treated with a H2RA.

[0042] *C. difficile* infection associated disease may also arise spontaneously, particularly when the subject is infected with certain stains of *C. difficile*, for example an infection caused by the NAP1/027/B strain which produces high levels of Toxin A, Toxin B and other toxins.

[0043] The halogenated salicylanilide may be used as the first line treatment of a *Clostridium* infection, for example a *C. difficile* infection. By "first-line" treatment is meant the first treatment of the *Clostridium* infection. In the first line treatment the *Clostridium* infection has not been treated with an antibiotic active against the *Clostridium* infection, for example, metronidazole, vancomycin, fidaxomicin or a rifamycin such as rifaximin. Accordingly, it may be that the halogenated salicylanilide is for use in the treatment of a *Clostridium* infection (for example *C. difficile* infection), wherein the infection has not been treated with an antibiotic prior to administration of the halogenated salicylanilide to the subject.

[0044] The halogenated salicylanilide may be used to treat a recurrent *Clostridium* infection (e.g. a *C. difficile* infection), for example a *Clostridium* infection which has recurred following prior treatment of the subject with an antibiotic (or other agent) other than a halogenated salicylanilide. For example, the halogenated salicylanilide may be used to treat a *Clostridium* infection (for example a *C. difficile* infection) which has recurred in a subject following prior treatment of the subject with an antibiotic selected from metronidazole, vancomycin, fidaxomicin and a rifamycin such as rifaximin. Suitably an antibiotic selected from metronidazole, vancomycin and fidaxomicin.

[0045] The halogenated salicylanilide may be used to treat a *Clostridium* infection (for example a *C. difficile* infection) which is refractory (for example non-responsive) to treatment with an antibiotic (or other agent) other than a halogenated salicylanilide. For example, the halogenated salicylanilide may be used to treat a refractory *Clostridium* infection (for example a *C. difficile* infection) in a subject. Accordingly, the halogenated salicylanilide may be for use in the treatment of a *Clostridium* infection (e.g. *C. difficile*) that is refractory to a prior antibiotic treatment other than a halogenated salicylanilide. For example the halogenated salicylanilide may be used to treat a *C. difficile* in a subject, wherein the *C. difficile* is refractory to treatment of the subject with an antibiotic selected from metronidazole, vancomycin, fidaxomicin and a rifamycin such as rifaximin.

[0046] It may be that the halogenated salicylanilide is used to treat a *Clostridium* infection (for example a *C. difficile* infection) which is resistant to an antibiotic agent used to treat the

Clostridium infection. Accordingly there is provided a halogenated salicylanilide, or a pharmaceutically acceptable salt thereof, for use in the treatment of a *Clostridium* infection (for example a *C. difficile* infection) which is resistant to an antibiotic agent other than the halogenated salicylanilide.

[0047] It may be that the *Clostridium* (for example a *C. difficile*) is resistant to an antibiotic agent approved by the US FDA or European Medicines Agency prior to 27 May 2016, preferably an antibiotic approved for use in the treatment of a *Clostridium* infection (for example a *C. difficile* infection). It may be that the *Clostridium* (for example a *C. difficile*) which is resistant to an antibiotic other than the halogenated salicylanilide. It may be that the *Clostridium* (for example a *C. difficile*) which is resistant to an antibiotic selected from a penicillin, a cephalosporin, a carbapenem, a monobactam (for example a β -lactam antibiotic), a fusidane, a fluoroquinolone, a tetracycline, a glycylicycline, phenicol (for example chloramphenicol), a macrolide, a macrocyclic (for example fidaxomicin), a rifamycin, a ketolide, a lincosamide, an oxazolidinone (for example cadazolid), an aminocyclitol, a polymyxin, a glycopeptide, an aminoglycoside, a lipopeptide, an antimycobacterial, a nitroimidazole, bacitracin, mupiricin, a pleuromutilin, a rifamycin, a sulphonamide and trimethoprim.

[0048] It may be that the *Clostridium* (for example a *C. difficile*) is resistant to an antibiotic selected from a nitroimidazole, for example metronidazole; a benzimidazole, for example ridinilazole (SMT19969); a glycopeptide, for example vancomycin; a macrocyclic antibiotic, for example fidaxomicin; an oxazolidinone, for example cadazolid; a lipopeptide, for example surothromycin or daptomycin; a glycylicycline, for example tigecycline; a DNA Minor Groove Binder (for example MGB-BP-3) a glycolipodepsipeptide (for example ramoplanin); CRS3123 (a Methionyl-tRNA synthetase (MetRS) inhibitor, Crestone Inc); and a rifamycin such as rifaximin. For example it may be that the *Clostridium* (for example a *C. difficile*) is resistant to an antibiotic selected from metronidazole, vancomycin, fidaxomicin and a rifamycin (e.g. rifaximin).

[0049] The *C. difficile* infection may be any strain of *C. difficile* for example the *C. difficile* strains shown in Table 1 below, in which the NCTC number is the UK National Collection of Type Culture reference number.

Table 1

NCTC no.	Ribotype and Toxin Status
NCTC 11209 (T)	Type strain, PCR-ribotype 001 reference strain
NCTC 11204	PCR-ribotype 001, toxin A and B positive

NCTC no.	Ribotype and Toxin Status
NCTC 11205	PCR-ribotype 001, toxin A and B positive
NCTC 11207	PCR-ribotype 001, toxin A and B positive
NCTC 11208	PCR-ribotype 001, toxin A and B positive
NCTC 11223	PCR-ribotype 001, toxin A and B positive
NCTC 11382	PCR-ribotype 001, toxin A and B positive
NCTC 12729	PCR-ribotype 002, toxin A and B positive
NCTC 13287	PCR-ribotype 017, toxin A negative, toxin B positive
NCTC 13307	PCR-ribotype 012, toxin A and B positive
NCTC 13404	PCR-ribotype 106 reference strain, toxin A and B positive
NCTC 13366	PCR-ribotype 027 reference strain, toxin A and B positive
T= Type Strain	

[0050] It may be that the *C. difficile* infection is a strain of *C. difficile* which produces high levels of Toxin A (entero toxin) and/or Toxin B (cytotoxin). For example it may be that the the *C. difficile* infection is a strain of *C. difficile* which produces higher levels of Toxin A than NCTC 11209 (T). It may be that the the *C. difficile* infection is a strain of *C. difficile* which produces higher levels of Toxin B than NCTC 11209 (T). It may be that the the *C. difficile* infection is a strain of *C. difficile* which produces higher levels of Toxin A and Toxin B than NCTC 11209 (T).

[0051] The *C. difficile* infection may be the hyper-virulent BI/NAP1 (also known as ribotype 027, NAP1/027/BI or NCTC 13366) strain which shows increased Toxin A (entero toxin) and Toxin B (cytotoxin) production as well as the production of additional novel binary toxins. Accordingly, the halogenated salicylanilide may be for use in the treatment of a *C. difficile* caused by the NAP1/027/BI *C. difficile* strain.

[0052] *C. difficile* is an anaerobe and as such the bacteria itself is generally not the primary mechanism for the transmission of infection, because the bacterial is not viable in aerobic conditions. However, *C. difficile* produces spores which are metabolically dormant and very stable. Spores shed in faecal matter are therefore very difficult to eradicate and may persist in the environment for prolonged periods of time, because they are resistant to heat and common cleaning and sterilisation chemicals. The spores represent the primary mechanism

for the transmission of *C. difficile* infections. Current treatments for *C. difficile*, for example, metronidazole, vancomycin and rifamycins are effective against *C. difficile*. However, these compounds have limited effects on sporulation and may not be effective in preventing transmission of infections by the spores. Fidaxomicin has been shown to be superior to metronidazole, vancomycin, and rifaximin in inhibiting sporulation and as such is currently considered to be the “gold-standard” treatment of *C. difficile* infections because of its potential to also inhibit transmission of infection.

[0053] As illustrated in the Examples below, halogenated salicylanilides (for example rafoxanide) are effective inhibitors of *C. difficile* sporulation. In the study performed in the Examples, rafoxanide was more effective than fidaxomicin in inhibiting sporulation. Accordingly it is expected that the halogenated salicylanilides will provide an effective treatment which both kills the *C. difficile* bacteria and inhibits *C. difficile* sporulation. The halogenated salicylanilides are therefore expected to provide an effective treatment of the initial *C. difficile* infection and also prevent or minimise the risk of transmission or spread of infection, for example spread of infection in a community or hospital environment.

[0054] Accordingly there is provided a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof (for example rafoxanide), for use in preventing or inhibiting sporulation of *C. difficile*.

[0055] Also provided is a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof (for example rafoxanide), for use in preventing or inhibiting transmission or spread of a *C. difficile* infection.

[0056] Also provided is a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof (for example rafoxanide), for use in a method of preventing or inhibiting transmission or spread of a *C. difficile* infection, the method comprising administering the halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof to a subject with a *C. difficile* infection.

[0057] A common problem associated with *C. difficile* infection is the recurrence of the infection following initial antibiotic treatment. Often a patient will respond well to the initial antibiotic treatment and will be symptom free for a period of time. However, in many patients recurrence of the infection is common and is often more severe than the initial infection (Louie TJ, et al. N. Eng. J. Med 2011; 364: 422–31). Mortality rates increase as the frequency of recurrent infection increases. A primary factor in the recurrence of infection is thought to be spores residing in the GI tract, particularly in the colon of a patient who has previously had a *C. difficile* infection. Spores present in the colon of a patient that has been infected with *C. difficile* and can persist there in a dormant state for long periods of time.


Upon activation the spores result in recurrence of the infection. The sporulation inhibitory properties of the halogenated salicylanilides are therefore expected to be beneficial in the prevention or reduction of the recurrence of *C. difficile* infection by reducing or eliminating spore formation in a patient infected with *C. difficile*.

[0058] Accordingly also provided is a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof (for example rafoxanide), for use in a method of preventing or inhibiting recurrence of *C. difficile* infection in a subject with a *C. difficile* infection, the method comprising administering the halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof to the subject.

Halogenated salicylanilides

[0059] Halogenated salicylanilides are also known as 2-hydroxy-N-phenylbenzamides or 2-hydroxybenzanilides. Salicylanilides are weakly acidic phenolic compounds. Halogenated salicylanilides are salicylanilides substituted by at least one halo group. The compounds were originally developed as fungicides for topical use and as antimicrobial agents in soaps. Later these compounds were shown to possess potent antihelmintic activity of which niclosamide, tribromosalan and clixanide were some of the first agents to be used. A wide range of halogenated salicylanilide derivatives are known. Any halogenated salicylanilide possessing antibacterial activity against *Clostridium* may be used in the present invention. For example, the halogenated salicylanilide may be any of the niclosamide analogues described in WO 2008/021088, which are incorporated herein by reference thereto.

[0060] The halogenated salicylanilide may be a halogenated salicylanilide of the formula (I):



(I)

wherein

X is O or S;

R¹ and R² are at each occurrence independently selected from halo;

R³ and R⁴ are at each occurrence independently selected from H, C₁₋₆ alkyl, -OR^{A1}, -NO₂ and -CN;

R⁵ is H or -L¹-R⁷;

R⁶ is H or -C(O)R^{A2};

L¹ is selected from a bond, O, S, or -(CR^{A3}R^B)_o-, wherein o is 1 or 2;

R⁶ is phenyl, unsubstituted or substituted with 1, 2, or 3 groups selected from halo, C₁₋₄ alkyl, -OR^{A4}, -NO₂ and -CN;

R^{A1}, R^{A2}, R^{A3} and R^{A4} are at each occurrence independently selected from H and C₁₋₄ alkyl;


R^B is at each occurrence selected from H, C₁₋₄ alkyl and -CN;

n and p are each independently selected from 0, 1, 2, 3 or 4, with the proviso that n+p is at least 1;

t and v are independently selected from 0, 1 and 2;

or a pharmaceutically acceptable salt, or ester thereof.

[0061] The halogenated salicylanilide of formula (I) may be of the formula (II), or a pharmaceutically acceptable salt, or ester thereof.



(II)

[0062] The following statements in the numbered paragraphs below apply to compounds of the formulae (I) or (II). These statements are independent and interchangeable. In other words, any of the features described in any one of the following statements may (where chemically allowable) be combined with the features described in one or more other statements below. In particular, where a compound is exemplified or illustrated in this specification, any two or more of the statements below which describe a feature of that compound, expressed at any level of generality, may be combined so as to represent subject matter which is contemplated as forming part of the disclosure of this invention in this specification.

[0063] 1. X is O.

- [0064] 2. R^1 and R^2 are at each occurrence independently selected from fluoro, chloro, bromo and iodo.
- [0065] 3. R^1 and R^2 are at each occurrence independently selected from chloro, bromo and iodo.
- [0066] 4. R^1 is chloro.
- [0067] 5. R^1 is bromo.
- [0068] 6. R^1 is iodo.
- [0069] 7. R^2 is chloro.
- [0070] 8. R^2 is bromo.
- [0071] 9. R^2 is iodo.
- [0072] 10. R^3 and R^4 are at each occurrence independently selected from H, C_{1-4} alkyl, $-OR^{A1}$, $-NO_2$ and $-CN$.
- [0073] 11. R^3 and R^4 are at each occurrence independently selected from H, C_{1-4} alkyl, $-OR^{A1}$ and $-NO_2$.
- [0074] 12. R^3 and R^4 are at each occurrence independently selected from H, C_{1-4} alkyl, $-OH$, $-OMe$, $-NO_2$ and $-CN$, for example H, C_{1-4} alkyl, $-OH$ or $-NO_2$.
- [0075] 13. R^5 is H.
- [0076] 14. R^5 is $-L^1-R^7$.
- [0077] 15. L^1 is selected from $-O-$, $-CH_2-$ and $-CH(CN)-$, for example $-O-$ or $-CH(CN)-$.
- [0078] 16. R^7 is phenyl, unsubstituted or substituted with 1, 2, or 3 groups selected from halo, C_{1-4} alkyl and $-CN$.
- [0079] 17. R^7 is phenyl unsubstituted or substituted with 1, 2, or 3 groups (for example 1 or 2 groups) selected from halo.
- [0080] 18. R^7 is unsubstituted phenyl.
- [0081] 19. L^1 is selected from $-O-$ and $-CH(CN)-$; and R^7 is phenyl unsubstituted or substituted with 1, 2, or 3 groups selected from halo.
- [0082] 20. R^6 is H.
- [0083] 21. R^6 is $-C(O)R^{A2}$, for example $-C(O)CH_3$.
- [0084] 22. $t = 0$ or 1 .
- [0085] 23. $t = 0$.

[0086] 24. $v = 0$ or 1.

[0087] 25. $v = 0$.

[0088] 26. o is 1.

[0089] 27. $v = 1$ and R^4 is selected from $-OH$, C_{1-4} alkyl and $-NO_2$.

[0090] 28. A compound of any of formulae (I) or (II), or a pharmaceutically acceptable salt thereof.

[0091] Particular compounds are compounds of formula (I) or formula (II), or a pharmaceutically acceptable salt or ester thereof wherein:

X is O;

R^1 and R^2 are at each occurrence independently selected from halo;

R^3 and R^4 are at each occurrence independently selected from H, C_{1-4} alkyl, $-OR^{A1}$, $-NO_2$ and CN;

R^5 is H or $-L^1-R^7$;

R^6 is H or $-C(O)R^{A2}$;

L^1 is selected from O and $-CH(CN)-$;

R^7 is phenyl unsubstituted or substituted with 1, 2, or 3 groups selected from halo;


R^{A1} and R^{A2} are at each occurrence independently selected from H and C_{1-4} alkyl;

n and p are each independently selected from 0, 1, 2, 3 or 4, with the proviso that $n+p$ is at least 1;


t and v are independently selected from 0, 1 and 2.

or a pharmaceutically acceptable salt, or ester thereof.


[0092] The halogenated salicylanilide may be selected from:




15




rafoxanide




oxyclosanide




resorantel



clioxanide




dibromosalan



tribromosalan

and




niclosamide

;

or a pharmaceutically acceptable salt or ester thereof.


[0093] The halogenated salicylanilide may be:




or a pharmaceutically acceptable salt or ester thereof.

[0094] The halogenated salicylanilide may be a compound selected from Table 1 in WO 2008/021088, or a pharmaceutically acceptable salt thereof.

[0095] It may be that the halogenated salicylanilide, for example the halogenated salicylanilide of the formulae (I) or (II) is not the following compounds:





[0096] The halogenated salicylanilide may be selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clioxanide, dibromosalan, tribromosalan, brotitanide and niclosamide, or a pharmaceutically acceptable salt or ester thereof.

[0097] The halogenated salicylanilide may be selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, dibromosalan, tribromosalan and niclosamide, or a pharmaceutically acceptable salt or ester thereof.

[0098] The halogenated salicylanilide may be selected from the group consisting of clioxanide, closantel, oxyclozanide, rafoxanide, tribromosalan or a pharmaceutically acceptable salt or ester thereof.

[0099] The halogenated salicylanilide may be selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clioxanide, dibromosalan, tribromosalan, brotitanide and niclosamide, or a pharmaceutically acceptable salt thereof.

[00100] The halogenated salicylanilide may be selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clioxanide, dibromosalan, tribromosalan and niclosamide, or a pharmaceutically acceptable salt thereof.

[00101] The halogenated salicylanilide may be selected from the group consisting of niclosamide, clioxanide, closantel, oxyclozanide, rafoxanide and tribromosalan, or a pharmaceutically acceptable salt thereof.

[00102] The halogenated salicylanilide may be selected from the group consisting of clioxanide, closantel, oxyclozanide, rafoxanide and tribromosalan, or a pharmaceutically acceptable salt thereof.

[00103] The halogenated salicylanilide may be selected from the group consisting of rafoxanide, oxyclozanide and clioxanide, or a pharmaceutically acceptable salt thereof.

[00104] The halogenated salicylanilide may be selected from the group consisting of clioxanide, closantel, rafoxanide and tribromosalan, or a pharmaceutically acceptable salt thereof.

[00105] The halogenated salicylanilide may be selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clixanide, dibromosalan, tribromosalan, brotitanide and niclosamide.

[00106] The halogenated salicylanilide may be niclosamide, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is niclosamide or a pharmaceutically acceptable salt thereof, suitably the halogenated salicylanilide is niclosamide.

[00107] The halogenated salicylanilide may be clixanide, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is clixanide or a pharmaceutically acceptable salt thereof, suitably the halogenated salicylanilide is clixanide.

[00108] The halogenated salicylanilide may be closantel, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is closantel or a pharmaceutically acceptable salt thereof, suitably the halogenated salicylanilide is closantel.

[00109] The halogenated salicylanilide may be oxyclozanide, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is oxyclozanide or a pharmaceutically acceptable salt thereof, suitably the halogenated salicylanilide is oxyclozanide.

[00110] The halogenated salicylanilide may be rafoxanide, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is rafoxanide or a pharmaceutically acceptable salt thereof, suitably the halogenated salicylanilide is rafoxanide.

[00111] The halogenated salicylanilide may be tribromosalan, or a pharmaceutically acceptable salt or ester thereof, for example the halogenated salicylanilide is tribromosalan or a pharmaceutically acceptable salt thereof, suitably particularly the halogenated salicylanilide is tribromosalan.

[00112] It is to be understood that any of the halogenated salicylanilides described in this section or elsewhere in the application may be used in any of the treatments described herein.

[00113] The halogenated salicylanilide may be administered to the subject using any suitable route, for example parenterally (for example intravenous, intramuscular or subcutaneous administration), mucosal administration (for example oral or rectal administration). Suitably the halogenated salicylanilide is administered orally or rectally. More particularly the halogenated salicylanilide is administered orally.

[00114] The subject or patient in any of the treatments described is suitably a human or animal, for example a warm-blooded animal. Particularly the subject is a human. The subject may be a human aged 65 years or older. The subject may be an animal, e.g. a mammal. In particular, the halogenated salicylanilide may be for use in the treatment of *Clostridium* infections in commercial animals such as livestock (e.g. cows, sheep, chickens, pigs, geese, ducks, goats, etc.). Alternatively, halogenated salicylanilide can be used to treat companion animals such as cats, dogs, horses, etc. The treatment of animals infected a *C. difficile* with may be particularly effective for preventing spread of infection through animal faecal matter to humans or other animals.

[00115] Also provided is the use a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof for the manufacture of a medicament for the treatment of an infection in a subject caused by *Clostridium* bacteria.

[00116] Also provided is a method of treating an infection caused by *Clostridium* bacteria in a subject, the method comprising administering to said subject a therapeutically effective amount of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof.

[00117] It is to be understood that the use and methods of the above two paragraphs are applicable to any of the infections, halogenated salicylanilides and routes of administration described herein.

[00118] Preferred, suitable, and optional features of any one particular aspect of the present invention are also preferred, suitable, and optional features of any other aspect.

BRIEF DESCRIPTION OF THE DRAWINGS

[00119] Figure 1 shows the development of heat-resistant spore count over time for *Clostridium difficile* 7-6011209 in the presence of rafxanide or fidaxomicin at a concentration of 8-fold above the MIC for rafxanide and > 8-fold above the MIC for fidaxomicin. The control shows the spore count over time in the Clospore medium used in the study.

DETAILED DESCRIPTION

Definitions

[00120] Unless otherwise stated, the following terms used in the specification and claims have the following meanings set out below.

[00121] It is to be appreciated that references to “treating” or “treatment” include prophylaxis as well as the alleviation of established symptoms of a condition. “Treating” or “treatment” of a state, disorder or condition therefore includes: (1) preventing or delaying the appearance of

clinical symptoms of the state, disorder or condition developing in a subject, for example a human, that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition, (2) inhibiting the state, disorder or condition, *i.e.*, arresting, reducing or delaying the development of the disease or a relapse thereof (in case of maintenance treatment) or at least one clinical or subclinical symptom thereof, or (3) relieving or attenuating the disease, *i.e.*, causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms. Accordingly in the context of treating infections caused by a *Clostridium* bacteria includes:

- (i) the prevention of a disease caused by *Clostridium* species, particularly *Clostridium difficile*;
- (ii) the suppression of a disease caused by *Clostridium* species, particularly *Clostridium difficile*;
- (iii) the relief of symptoms of a disease caused by *Clostridium* species, particularly *Clostridium difficile*;
- (iv) the eradication of a non-symptomatic colonization by *Clostridium* species, particularly *Clostridium difficile* from an area on or in the body;
- (v) the eradication of a *Clostridium difficile* symptomatic infection;
- (vi) the eradication a *Clostridium* species, particularly *Clostridium difficile*; from an area of the body affected by another disease that could enable establishment of an infection more readily, than in a non-disease affected area – e.g. in the intestinal tract;
- (vii) the suppression of a disease caused a *Clostridium* infection, particularly *Clostridium difficile*; from an area of the body affected by another noninfectious disease that enables establishment of an infection more readily, than in a non-disease affected area;
- (viii) preventing or reducing the risk of transmission or spread of a *Clostridium* infection, particularly *Clostridium difficile*; or
- (ix) preventing or reducing the risk of recurrence of a *Clostridium* infection, particularly *Clostridium difficile*.

[00122] A “therapeutically effective amount” means the amount of a compound that, when administered to a subject, for example a human, for treating a disease, is sufficient to effect such treatment for the disease. The “therapeutically effective amount” will vary depending on the compound, the disease and its severity and the age, weight, etc., of the subject to be treated.

[00123] Minimum inhibitory concentration (MIC) is the lowest concentration of an antibacterial that will inhibit the visible growth of a microorganism after overnight incubation. Minimum inhibitory concentrations are important in diagnostic laboratories to confirm resistance of microorganisms to an antimicrobial agent and also to monitor the activity of new antimicrobial agents. A MIC is generally regarded as the most basic laboratory measurement of the activity of an antimicrobial agent against an organism.

[00124] The median lethal dose, LD50 (abbreviation for "lethal dose, 50%") of a toxin, radiation, or pathogen is the dose required to kill half the members of a tested population after a specified test duration. LD50 figures are frequently used as a general indicator of a substance's acute toxicity.

[00125] Therapeutic index (therapeutic ratio) is defined as the amount of a therapeutic agent causing the therapeutic effect measured as MIC to the amount that causes death in animal studies measured as LD50.

[00126] The rate of resistance development is quantified as the frequency of spontaneous mutants in a population of bacteria that is able to resist a given concentration of an antibiotic. For example the rate of resistance development may be 10^{-9} if on average 1 cell in 10^9 cells is able to survive a concentration of antibiotic corresponding to 1x MIC incubated at 37°C for 48 hours using the method described in Drago et al. Journal of Antimicrobial Chemotherapy, 2005, 56(2), 353 to 359).

[00127] In microbiology, colony-forming unit (CFU) is an estimate of the number of viable bacteria or fungal cells in a sample. Viable is defined as the ability to multiply via binary fission under the controlled conditions.

[00128] The term "halo" or "halogen" refers to one of the halogens, group 17 of the periodic table. In particular the term refers to fluorine, chlorine, bromine and iodine.

[00129] The term C_m-C_n refers to a group with m to n carbon atoms.

[00130] The term "C₁-C₆ alkyl" refers to a linear or branched hydrocarbon chain containing 1, 2, 3, 4, 5 or 6 carbon atoms, for example methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl and *n*-hexyl. "C₁-C₄ alkyl" similarly refers to such groups containing up to 4 carbon atoms.

[00131] The term "optionally substituted" refers to either groups, structures, or molecules that are substituted and those that are not substituted.

[00132] Where optional substituents are chosen from “one or more” groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

[00133] Where a moiety is substituted, it may be substituted at any point on the moiety where chemically possible and consistent with atomic valency requirements. The moiety may be substituted by one or more substituents, e.g. 1, 2, 3 or 4 substituents; optionally there are 1 or 2 substituents on a group. Where there are two or more substituents, the substituents may be the same or different.

[00134] Substituents are only present at positions where they are chemically possible, the person skilled in the art being able to decide (either experimentally or theoretically) without undue effort which substitutions are chemically possible and which are not.

[00135] Throughout the description and claims of this specification, the words “comprise” and “contain” and variations of them mean “including but not limited to”, and they are not intended to (and do not) exclude other moieties, additives, components, integers or steps. Throughout the description and claims of this specification, the singular encompasses the plural unless the context otherwise requires. In particular, where the indefinite article is used, the specification is to be understood as contemplating plurality as well as singularity, unless the context requires otherwise.

[00136] Features, integers, characteristics, compounds, chemical moieties or groups described in conjunction with a particular aspect, embodiment or example of the invention are to be understood to be applicable to any other aspect, embodiment or example described herein unless incompatible therewith. All of the features disclosed in this specification (including any accompanying claims, abstract and drawings), and/or all of the steps of any method or process so disclosed, may be combined in any combination, except combinations where at least some of such features and/or steps are mutually exclusive. The invention is not restricted to the details of any foregoing embodiments. The invention extends to any novel one, or any novel combination, of the features disclosed in this specification (including any accompanying claims, abstract and drawings), or to any novel one, or any novel combination, of the steps of any method or process so disclosed.

[00137] The reader's attention is directed to all papers and documents which are filed concurrently with or previous to this specification in connection with this application and which are open to public inspection with this specification, and the contents of all such papers and documents are incorporated herein by reference.

[00138] Suitable or preferred features of any compounds of the present invention may also be suitable features of any other aspect.

[00139] The invention contemplates pharmaceutically acceptable salts of the halogenated salicylanilide compounds of the invention. These may include the acid addition and base salts of the compounds. These may be acid addition and base salts of the compounds. Suitable acid addition salts are formed from acids which form non-toxic salts. Suitable base salts are formed from bases which form non-toxic salts.

[00140] Pharmaceutically acceptable salts of the halogenated salicylanilide compounds may be prepared by for example, one or more of the following methods:

- (i) by reacting the compound of the invention with the desired acid or base; or
- (ii) by converting one salt of the compound of the invention to another by reaction with an appropriate acid or base or by means of a suitable ion exchange column.

[00141] These methods are typically carried out in solution. The resulting salt may precipitate out and be collected by filtration or may be recovered by evaporation of the solvent. The degree of ionisation in the resulting salt may vary from completely ionised to almost non-ionised.

[00142] Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture". Where a compound of the invention has two or more stereocentres any combination of (R) and (S) stereoisomers is contemplated. The combination of (R) and (S) stereoisomers may result in a diastereomeric mixture or a single diastereoisomer. The compounds of the invention may be present as a single stereoisomer or may be mixtures of stereoisomers, for example racemic mixtures and other enantiomeric mixtures, and diastereomeric mixtures. Where the mixture is a mixture of enantiomers the enantiomeric

excess may be any of those disclosed above. Where the compound is a single stereoisomer the compounds may still contain other diastereoisomers or enantiomers as impurities. Hence a single stereoisomer does not necessarily have an enantiomeric excess (e.e.) or diastereomeric excess (d.e.) of 100% but could have an e.e. or d.e. of about at least 85%

[00143] The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 2001), for example by synthesis from optically active starting materials or by resolution of a racemic form. Some of the compounds of the invention may have geometric isomeric centres (E- and Z- isomers). It is to be understood that the present invention encompasses all optical, diastereoisomers and geometric isomers and mixtures thereof that possess activity against *Clostridium* bacteria, for example *C. difficile*.

[00144] It is also to be understood that certain compounds of the invention, or salts or esters thereof, may exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms that possess activity against *Clostridium* bacteria, for example *C. difficile*.

[00145] It is also to be understood that the halogenated salicylanilides of the invention may exhibit polymorphism, and that the invention encompasses all such forms that possess activity against *Clostridium* bacteria, for example *C. difficile*.

[00146] It is further to be understood that the halogenated salicylanilide may be used in the form of suitable pharmaceutically-acceptable pro-drug of the compound and that such prodrugs are intended to be encompassed by the invention. Accordingly, halogenated salicylanilide may be administered in the form of a pro-drug, that is a compound that is broken down in the human or animal body to release a compound of the invention. A pro-drug may be used to alter the physical properties and/or the pharmacokinetic properties of a compound of the invention. A pro-drug can be formed when the compound of the invention contains a suitable group or substituent to which a property-modifying group can be attached. Examples of pro-drugs include *in vivo* cleavable ester derivatives that may be formed at a hydroxy group in a compound.

[00147] Accordingly, the present invention includes the halogenated salicylanilides as defined hereinbefore when made available by organic synthesis and when made available

within the human or animal body by way of cleavage of a pro-drug thereof. Accordingly, the present invention includes those halogenated salicylanilide compounds that are produced by organic synthetic means and also such compounds that are produced in the human or animal body by way of metabolism of a precursor compound, that is the halogenated salicylanilide may be a synthetically-produced compound or a metabolically-produced compound.

[00148] A suitable pharmaceutically-acceptable pro-drug of a halogenated salicylanilide compound is one that is based on reasonable medical judgement as being suitable for administration to the human or animal body without undesirable pharmacological activities and without undue toxicity.

[00149] Various forms of pro-drug have been described, for example in the following documents :-

[00150] a) Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);

[00151] b) Design of Pro-drugs, edited by H. Bundgaard, (Elsevier, 1985);

[00152] c) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Pro-drugs", by H. Bundgaard p. 113-191 (1991);

[00153] d) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);

[00154] e) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988);

[00155] f) N. Kakeya, *et al.*, Chem. Pharm. Bull., 32, 692 (1984);

[00156] g) T. Higuchi and V. Stella, "Pro-Drugs as Novel Delivery Systems", A.C.S. Symposium Series, Volume 14; and

[00157] h) E. Roche (editor), "Bioreversible Carriers in Drug Design", Pergamon Press, 1987.

[00158] The halogenated salicylanilide may be used in the form of a prodrug of the compound for example, an *in vivo* cleavable ester thereof. An *in vivo* cleavable ester of a compound may be, for example, a pharmaceutically-acceptable ester which is cleaved in the human or animal body to produce the parent compound.

[00159] A suitable pharmaceutically-acceptable pro-drug of a halogenated salicylanilide that possesses a hydroxy group is, for example, an *in vivo* cleavable ester or ether thereof. An *in vivo* cleavable ester or ether of a compound containing a hydroxy group is, for example, a

pharmaceutically-acceptable ester or ether which is cleaved in the human or animal body to produce the parent hydroxy compound. Suitable pharmaceutically-acceptable ester forming groups for a hydroxy group include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters). Further suitable pharmaceutically-acceptable ester forming groups for a hydroxy group include C₁₋₁₀alkanoyl groups such as acetyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl groups, C₁₋₁₀alkoxycarbonyl groups such as ethoxycarbonyl, *N,N*-(C₁₋₆)₂carbamoyl, 2-dialkylaminoacetyl and 2-carboxyacetyl groups. Examples of ring substituents on the phenylacetyl and benzoyl groups include aminomethyl, *N*-alkylaminomethyl, *N,N*-dialkylaminomethyl, morpholinomethyl, piperazin-1-ylmethyl and 4-(C₁₋₄alkyl)piperazin-1-ylmethyl. Suitable pharmaceutically-acceptable ether forming groups for a hydroxy group include α -acyloxyalkyl groups such as acetoxymethyl and pivaloyloxymethyl groups. Accordingly reference to a "pharmaceutically acceptable ester" of a compound encompasses the esters described above.

Halogenated Salicylanilides

[00160] The halogenated salicylanilide used in the present invention may be any of the halogenated salicylanilides described herein, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable salt thereof, or a pro-drug of any thereof.

[00161] Particular halogenated salicylanilides include the compounds of formulae (I) and (II) or a pharmaceutically acceptable salt thereof as described herein.

[00162] More particularly the halogenated salicylanilide is selected from the group consisting of tetrachlorosalicylanilide, closantel, rafoxanide, oxyclozanide, resorantel, clixanide, dibromosalan, tribromosalan and niclosamide.

Niclosamide

[00163] In a one embodiment of the invention the halogenated salicylanilide is niclosamide or a pharmaceutically acceptable salt thereof. Niclosamide (2',5-dichloro-4'-nitrosalicylanilide) exhibits the following acute toxicity:

LD₅₀, mice, p.o., >5000 mg/kg

LD₅₀, rats, p.o., 5000 mg/kg

LD₅₀, rats, dermal, 2000 mg/kg

LD₅₀, rabbits, p.o., 5000 mg/kg

LD₅₀, cats, p.o., >1000 mg/kg

[00164] Niclosamide thus exhibits low toxicity. The compound is poorly soluble in water and shows low intestinal absorption. Once in the bloodstream niclosamide is quickly cleared via the urinary tract or by enzymatic metabolism in the liver.

Niclosamide Derivatives

[00165] It is believed that a number of niclosamide analogs will act in a manner similar to niclosamide in the treatment of the *Clostridium* infections described herein. Illustrative niclosamide analogs include, but are not limited to closantel (CAS #: 57808-65-8), oxyclozanide (CAS #: 2277-92-1), rafoxanide (CAS #: 22662-39-1), clioxanide (CAS #: 14437-41-3). Other suitable niclosamide analogs include brotianide (CAS #: 23233-88-7), 4'-chloro-3-nitrosalicylanilide, 4'-chloro-5-nitrosalicylanilide, 2'-chloro-5'-methoxy-3-nitrosalicylanilide, 2'-methoxy-3,4'-dinitrosalicylanilide, 2',4'-dimethyl-3-nitrosalicylanilide, 2'-chloro-3,4'-dinitrosalicylanilide, 2'-ethyl-3-nitrosalicylanilide and 2'-bromo-3-nitrosalicylanilide; or a pharmaceutically acceptable salt thereof. Further niclosamide derivatives include those described in WO 2008/021088, particularly those described in Table 1 therein, which are incorporated herein by reference.

[00166] Particular niclosamide analogues include closantel, rafoxanide and oxyclozanide. These compounds are expected to have a suitable toxicity profile for the use described herein.

[00167] Acute toxicity of closantel:

LD₅₀, rats, p.o., 262-342 mg/kg (depending on the study), median 302 mg/kg

LD₅₀, rats, s.c., 67 mg/kg

LD₅₀, mice, p.o., 331 mg/kg

LD₅₀, mice, i.m., 57 mg/kg

[00168] Acute toxicity of rafoxanide:

LD₅₀, rats, p.o., 980->2000 mg/kg (depending on the study), median >1490 mg/kg

LD₅₀, mice, p.o., 232-300 mg/kg (depending on the study), median 266 mg/kg

LD₅₀, rabbits, p.o., 3200 mg/kg

[00169] Acute toxicity of oxyclozanide.

LD₅₀, rats, p.o., 980-3519 mg/kg (depending on the study), median 2250 mg/kg

LD₅₀, mice, p.o., 300 mg/kg


LD₅₀, rabbits, p.o., 3200 mg/kg

Brominated Halogenated Salicylanilides

[00170] In another embodiment the halogenated salicylanilide is a brominated halogenated salicylanilide, for example 4',5-dibromosalicylanilide (also known as dibromsalan); 3,5-dibromosalicylanilide (also known as metabromsalan; and 3,4',5-tribromosalicylanilide (also known as tribromsalan).

Synthesis

[00171] The halogenated salicylanilides described herein are known or can be synthesised using known methods. For example using methods analogous to those described in WO2004/006906. The compounds of the Formula (I) herein may be prepared by coupling an amine of the formula (III) with an acid of formula (IV):



[00172] Necessary starting materials are known or can be prepared using standard procedures of organic chemistry.

Pharmaceutical Compositions

[00173] The halogenated salicylanilide may be administered to the subject in the form of a pharmaceutical composition comprising the halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable excipient.

[00174] Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988.

[00175] The composition may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intraperitoneal

dosing or as a suppository for rectal dosing). Suitably the composition is in a form suitable for oral administration.

[00176] The compositions may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Dosage

[00177] The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the subject treated and the particular route of administration. For example, a formulation in a unit dose form such as a tablet or capsule intended for oral administration to humans will generally contain, for example, from 0.1 mg to 5 mg, for example from 0.5 mg to 5 g, from 0.5 to 1000 mg or from 10 to 500 mg of the halogenated salicylanilide compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition.

[00178] The size of the dose of the halogenated salicylanilide for the treatment of the *Clostridium* infections described herein will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well-known principles of medicine.

[00179] The halogenated salicylanilide will generally be administered in a dose of about 0.001 to about 75 mg/kg, for example from about 0.013 to about 66.7 mg/kg, about 0.5 to about 30 mg/kg or from about 2.5 to about 30 mg/kg. The halogenated salicylanilide may be administered within these dosage ranges to the subject from 1 to 4 times per day. The dosage may be administered by any suitable route, for example parenterally, orally or rectally. A particular route of administration which is generally applicable to all of the uses of the halogenated salicylanilides described herein is the oral administration of the halogenated salicylanilide to the subject.

[00180] The particular dosage regimen used to treat a subject will depend on a number of factors that may readily be determined, such as the severity of the condition and its responsiveness to initial treatment, but will normally involve one or more administrations per day on an ongoing basis. The effective dosage of the pharmaceutical composition of the present invention varies from the formulation, administration pathway, age, weight and gender of a human or animal or with a disease caused by *Clostridium species*, particularly *Clostridium difficile* colonizing or infecting the intestinal tract of a human or animal having a *Clostridium difficile* infection.

Therapeutic Use

[00181] As described hereinbefore the halogenated salicylanilide is used for the treatment of an infection caused by *Clostridium* bacteria, particularly *C. difficile*. The halogenated salicylanilide may act to kill or eradicate the infection from the subject, thus providing a bactericidal effect. Alternatively the halogenated salicylanilide may inhibit growth or replication of the bacteria thus producing a bacteriostatic effect. In the context of the present invention, treatment of a condition encompasses both therapeutic and prophylactic treatment, of either an infectious or a non-infectious condition, in a subject for example a mammal such as a human or animal, but in particular a human. It may involve complete or partial eradication of the condition, removal or amelioration of associated symptoms, arresting subsequent development of the condition, and/or prevention of, or reduction of risk of, subsequent occurrence of the condition.

[00182] Generally the halogenated salicylanilide will be administered to a subject experiencing symptoms of a *Clostridium* infection (for example a *C. difficile* infection). Accordingly, the halogenated salicylanilide may be for use in the treatment of a *C. difficile* associated disease, for example the halogenated salicylanilide may be for use in the treatment of a *C. difficile* associated disease selected from diarrhoea and colitis (including pseudomembranous colitis).

[00183] In an alternative embodiment the halogenated salicylanilide is for use in the treatment of a *C. difficile* in a subject, wherein the subject is asymptomatic. Such uses may be useful to eradicate or inhibit a *C. difficile* infection in a subject that is at risk of developing *C. difficile* associated disease. Such subjects could include, for example subjects which require surgical procedures in which prophylactic antibiotics may be administered (for example certain orthopaedic surgery). By eradicating the *C. difficile* infection prior to administration of further antibiotics, the risk of antibiotic induced diarrhoea may be reduced.

[00184] Subjects who have previously suffered from an antibiotic induced *C. difficile* infection may be at a particular risk of developing a *C. difficile* infection if they are administered antibiotics in the future. Accordingly, in another embodiment the halogenated salicylanilide is for use in the treatment of a subject prior to the administration of an antibiotic other than the halogenated salicylanilide, wherein the patient is asymptomatic of a *C. difficile* infection prior to administration of the halogenated salicylanilide and where the subject has previously suffered from an antibiotic induced a *C. difficile* infection.

[00185] When the *Clostridium* infection is an antibiotic induced *Clostridium* infection (for example a *C. difficile* infection) further administration of the antibiotic causing the induced infection is suitably halted. Alternatively, the dosage of the antibiotic may be reduced or

gradually tapered so as to reduce the risk of exacerbating the antibiotic induced *Clostridium* infection. Accordingly the halogenated salicylanilide may be administered to the subject concurrently with another antibiotic being used to treat a primary infection in the subject. For example it may be that the halogenated salicylanilide is administered to the subject concurrently with an antibiotic being used to treat a primary infection other than a *C. difficile* infection. The antibiotic used to treat the primary infection may, for example, be one or more antibiotics selected from a penicillin, a cephalosporin, a carbapenem, a monobactam (for example a β -lactam antibiotic), a fusidane, a fluoroquinolone, a tetracycline, a glycylicycline, phenicol (for example chloramphenicol), a macrolide, a macrocyclic (for example fidaxomicin), a rifamycin, a ketolide, a lincosamide, an oxazolidinone, an aminocyclitol, a polymyxin, a glycopeptide, an aminoglycoside, a lipopeptide, an antimycobacterial, a nitromidazole, bacitracin, mupiricin, a pleuromutilin, a rifamycin, a sulphonamide and trimethoprim, or a combination of two or more thereof. However, preferably the halogenated salicylanilide is used alone or together with a reduced or tapered dose of the antibiotic(s) responsible for the induced *Clostridium* infection. More preferably the halogenated salicylanilide is administered to the subject in the absence of any other antibiotic.

[00186] In one embodiment the halogenated salicylanilide is administered to the subject concurrently with another therapeutic agent in any of the treatments of the *Clostridium* infections (particularly the *C. difficile*) infections described herein. The other therapeutic agent may be, for example, an antibiotic active against *C. difficile*, a microbiome therapeutic or faecal transplant, or a vaccine or an antibody therapy for e.g. *C. difficile*.

[00187] Accordingly it may be that the halogenated salicylanilide is administered to the subject concurrently with another antibiotic active against a *Clostridium* infection, particularly a *C. difficile* infection, other than the halogenated salicylanilide itself. Examples of such antibiotics include a nitroimidazole, for example metronidazole; a benzimidazole, for example ridinilazole (SMT19969); a glycopeptide, for example vancomycin; a macrocyclic antibiotic, for example fidaxomicin; an oxazolidinone, for example cadazolid; a lipopeptide, for example surothromycin or daptomycin; a glycylicycline, for example tigecycline; a DNA Minor Groove Binder (for example MGB-BP-3) a glycolipodepsipeptide (for example ramoplanin); CRS3123 (a Methionyl-tRNA synthetase (MetRS) inhibitor, Crestone Inc); and a rifamycin such as rifaximin, or a combination of two or more thereof. It may be that the halogenated salicylanilide is administered concurrently with metronidazole (for example concurrently with intravenous metronidazole).

[00188] It may be that the halogenated salicylanilide is administered to the subject concurrently with a vaccine, for example concurrently with a vaccine which induces an immune response to *C. difficile* toxins, for example toxins A and B (e.g. ACAM-CDIFF,

Sanofi, or VLA84 (a fusion protein containing cell binding domains of Toxins A and B, Valneva), or a vaccine which prevents a *C. difficile* infection, for example PF06425090 (Pfizer).

[00189] It may be that the halogenated salicylanilide is administered to the subject concurrently with an antibody therapeutic, for example actoxumab and bezlotoxumab or a combination thereof.

[00190] It may be that the halogenated salicylanilide is administered to the subject concurrently with a faecal transplantation or microbiome therapeutics, for example concurrently with a faecal transplant, spores from a non-toxigenic *C. difficile* strain (e.g. VP-20621); spores from microbiome organisms (e.g. SER-109), a microbiota suspension (e.g. RBX2660), a probiotic (e.g. lactobacillus reuteri) or a β -lactamase, for example SYN-004.

[00191] Reference to administration "concurrently" herein includes the separate, simultaneous or separate administration of the halogenated salicylanilide with the other therapy. The halogenated salicylanilide may be administered to the subject administered by the same or different routes of administration, for example oral, intravenously, subcutaneously, or rectally). The halogenated salicylanilide and the other therapy may be administered as a combined preparation; however, generally they will be administered as separate dosage forms to enable the dose and dosing regimen of each to be tailored accordingly.

[00192] The presence of a *Clostridium* infection (for example a *C. difficile* infection) in a subject may be diagnosed using convention methods, for example infection may be suspected from subject exhibiting symptoms of a *C. difficile* associated disease. Infection may also be diagnosed using known methods for example

- A complete blood count to test for the presence of leukocytosis
- Measurement of albumin levels to check for hypoalbuminemia, which may accompany severe disease.
- Testing for elevated serum lactate levels (≥ 5 mmol/L), which can be a sign of severe disease.
- Stool examination. Stools may be positive for blood in severe colitis. Faecal leukocytes are present in about half of cases.

[00193] Stool assays for *C. difficile*, may also be used including the following:

- Stool culture: This is a sensitive test. However, the results are slow and may lead to a delay in the diagnosis.

- Glutamate dehydrogenase enzyme immunoassay (EIA): This is a very sensitive test and detects the presence of glutamate dehydrogenase produced by *C. difficile*.
- Real-time polymerase chain reaction (PCR) assay: This test is an alternative “gold standard” to stool culture. The assay may be used to detect the *C. difficile* gene toxins.
- EIA for detecting toxins A and B produced by *C. difficile*.

[00194] Imaging studies and procedures may also be used to detect infection in a subject. Suitable methods include abdominal computed tomography (CT) scanning. This method is particularly suitable when pseudomembranous colitis or other complications of CDI are suspected. In subjects with sepsis due to suspected megacolon, abdominal radiography may be performed instead of CT scanning to establish the presence of megacolon.

[00195] The use of halogenated salicylanilides for the treatment of a *Clostridium* infection (for example a *C. difficile* infection) as described herein is expected to provide a wider therapeutic window than conventional treatments such as vancomycin, metronidazole or fidaxomicin. Accordingly, the use of halogenated salicylanilides may result in reduced side effects compared to known *C. difficile* treatments.

[00196] Halogenated salicylanilides (e.g. niclosamide) are poorly absorbed following oral administration. Accordingly, the concentration of the halogenated salicylanilide in the faeces is expected to be high compared to for example oral administration of similar doses of vancomycin or metronidazole. A high local concentration of the halogenated salicylanilide in the GI tract, especially the colon, would be expected to enhance the potency and or efficiency of the antibacterial effect locally in the intestine and colon.

[00197] As illustrated in the examples, a number of the halogenated salicylanilide tested had similar or higher potencies than vancomycin, metronidazole or fidaxomicin and may therefore be expected to provide an antibacterial effect on *Clostridium* infections at a similar or lower dose than conventional treatments such as vancomycin.

Clostridium bacteria (for example *C. difficile*) are expected to exhibit a low frequency of spontaneous mutation in the presence of the halogenated salicylanilides described herein. Therefore, it is expected that the risk of resistance to the halogenated salicylanilide emerging will be low.

EXAMPLES

[00198] In the examples below, the effects of various halogenated salicylanilides are compared to vancomycin (a currently approved compound for the treatment *Clostridium difficile* infections).

Example 1 MIC determinations

[00199] *C. difficile* MICs are determined according to CLSI guideline using microbroth dilution as described in “Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard – Eighth Edition” CLSI, ISBN: 1-56238-789-8”, except isovitalex was used in the place of lysed horse blood.

[00200] The media used for the *C. difficile* MIC tests was Brucella broth supplemented with hemin (5 µg/ml), vitamin K and Isovitalex (according to the manufacturer’s instructions). Inoculated plates were incubated for 44 to 48 hours at 36°C under anaerobic conditions (anaerobic chamber or anaerobic jar with gaspack).

[00201] The *C. difficile* strain ATCC 700057 was used as a reference control during the *C. difficile* MIC determinations. This strain has an expected MIC towards vancomycin of 1 µg/mL.

[00202] The results are shown in Tables 2 and 3.

Table 2: MIC of clinical isolates of *C. difficile* against niclosamide and vancomycin

<i>C. difficile</i> isolate	Niclosamide (ug/mL)	Vancomycin(ug/mL)
7-6011209	0.06	0.5
7-7150288	0.125	1
7-7152701	0.25	2
7-7154992	0.25	1
7-5779928	0.25	2
7-6778909	0.25	0.5
7-6870430	0.25	0.5
7-7154712	0.25	2
7-7104022	0.125	1
7-7153872	0.5	2
7-5085357	0.125	0.5
7-7150997	0.125	0.5
7-6008526	0.25	2
7-7124449	0.125	0.5
7-6854508	0.125	0.5
7-7363761	0.125	0.5
7-7200552	0.25	0.5
7-7150318	0.125	0.5
7-7150628	0.125	1
7-7149204	0.125	1
7-7154712	0.25	2
7-7116411	0.125	2
7-7151551	0.25	2
7-7156197	0.25	2
12055	0.125	0.5

12060	0.125	0.5
12061	0.06	0.5
12062	0.06	0.5
12063	0.06	0.5
12064	0.06	1

[00203] Table 2 illustrates that niclosamide has a lower MIC than vancomycin against the *C. difficile* isolates tested.

Table 3

	<i>C. difficile</i> Strain 7-6011209	<i>C. difficile</i> Strain 12055
Compounds	MIC (µg/ml)	MIC (µg/ml)
Clioquinide	0.031	<0.08
Closantel	0.125	0.125
Oxyclozanide	0.25	0.25
Rafoxanide	<0.08	<0.08
Tribromsalan	0.125	0.25
Vancomycin	0.125	0.25

Example 2

[00204] Table 4 compares the properties of niclosamide with those of vancomycin, metronidazole, fidaxomicin. The data shown in Table 4 was obtained from published data together with data from the examples herein.

Table 4

	Vancomycin	Metronidazole	Fidaxomicin	Nicosamide
Class	Glycopeptide	Nitroimidazole	Macrolide	Halogenated salicylanilide
Dosing	TID, oral	TID, oral or IV	BID, oral	1-4 times daily, oral
MIC90 (µg/mL)	2 ^a	1 ^a	0.5 ^a	0.06-0.25 ^b
Highest MIC observed (µg/mL)	16 ^a	>32 ^a	1	0.5
Spectrum of activity	Gram + Bladder pain, bloating, bloody urine, painful urination, fever, dry mouth, irregular heartbeats, loss of appetite, mood changes, muscle pain or cramps, numbness, rapid weight gain, shortness of breath, tiredness	Gram +/- Abdominal or stomach cramps, dizziness, heartburn, Spinning sensation. Trouble sleeping, congestion, dry mouth	Gram + Nausea, vomiting, abdominal pain, gastrointestinal hemorrhage, anemia, neutropenia	Gram + Nausea, retching, abdominal pain
Side effects^c				

Footnotes:

- a ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, June 2002, p. 1647-1650' & 'CID 2012:55 (Suppl 2) • S143'
- b Data from Tables 3 and 4 herein.
- c Data from product labels.

Example 3: Additional MIC Determinations

5 **[00205]** The MIC of the halogenated salicylanilides clioxanide, closantel, oxyclozanide, rafoxanide and tribromsalan was determined against 24 clinical isolate strains of *C. difficile*. Fidaxomicin, metronidazole and vancomycin were used as comparator compounds in the study. The comparator compounds represent the antibiotics most often used in current treatments of *C. difficile* infections.

10 **[00206]** Stock solutions of the test compounds and comparators were made in DMSO at a concentration of 1 mg/mL.

Microorganisms

[00207] The *C. difficile* strains used are shown in Table 5:

Table 5

Name	MLST	Name	MLST
7-6011209	ST002	7-7363761	ST008
7-7150288	ST003	7-7200552	ST017
7-7154992	ST139	7-7150318	ST059
7-6778909	ST016	7-7150628	ST034
7-6870430	ST001	7-7149204	ST006
7-7104022	ST103	7-7116411	ST005
7-5085357	ST028	7-7150997	ST049
7-7124449	ST009	7-6854508	ST013
12055	-	12063	-
12060	-	12064	-
12061	-	12065	-
12062	-	12066	-

15 **Culture medium**

[00208] *C. difficile* strains were grown on Brucella blood agar + hemin + vitamin K [1], plates were incubated 44 to 48 hours at 36°C under anaerobic conditions. Broth cultures were performed in Brucella bouillon supplemented with Isovitalex [2] according the

instructions from the supplier (BBL). All cultures were performed under anaerobic conditions in an anaerobic chamber [3].

Antibacterial activity

5 [00209] The antibacterial activity of the study compounds was determined using the following protocol.

1. Day 1: bacterial strain is isolated and incubated at 37°C on Brucella blood agar + hemin + vitamin K.
2. Day 2: Inoculate 5 ml of Brucella bouillon supplemented with Isovitalex (BBI) with one isolated colony in 15 ml Falcon tube and incubated overnight at 36°C in anaerobic conditions.
3. Day 3:
 - Dilute the antibiotics in BBI to their highest concentration (8 µg/ml in 2 ml).
 - Make a series of two fold dilutions in deep well 96 well plates.
 - Transfer 150 µl of the antibiotics solution to 96-well plates.
 - After 5-6 hours, the culture was stopped and OD₆₀₀ was measured. The culture is diluted to 10⁶ CFU/ml
 - About 1 µl of this diluted culture is added in all wells in order to have 10³ cells per well.
 - Plates are incubated at 44 to 48 hours at 36°C under anaerobic conditions.
4. Day 5: OD₆₀₀ is measured after incubation.
Inhibition calculated as follows:

25
$$Inhibition = 1 - \frac{OD\ antibiotic - OD\ negative\ control}{OD\ positive\ control - OD\ negative\ control}$$

RESULTS

[00210] The MIC values of the tested compounds against the 24 strains of *C. difficile* are shown in Table 6.

Table 6

Strain	Clioquinide	Closoantel	Oxyclozanide	Rifaxanide	Tribromsalan	Fidaxomycin	Metronidazole	Vancomycin
7-6011209	< 0.008	0.016	< 0.008	< 0.008	< 0.008	< 0.008	0.031	0.031
7-7150288	< 0.008	0.031	< 0.008	< 0.008	< 0.008	< 0.008	0.063	0.5
7-7154992	< 0.008	0.063	< 0.008	0.016	0.063	< 0.008	1	0.5
7-6778909	< 0.008	0.031	< 0.008	< 0.008	< 0.008	< 0.008	0.25	0.5
7-6870430	< 0.008	0.031	< 0.008	< 0.008	0.016	0.016	1	0.5
7-7104022	< 0.008	0.25	< 0.008	< 0.008	1	< 0.008	>8	2
7-5085357	< 0.008	0.016	0.016	0.016	0.016	< 0.008	1	0.5
7-7150997	< 0.008	0.016	< 0.008	< 0.008	0.16	< 0.008	0.031	< 0.008
7-7124449	< 0.008	0.25	0.031	0.031	0.063	< 0.008	2	0.5
7-6854508	< 0.008	0.25	< 0.008	< 0.008	0.031	< 0.008	1	0.5
7-7363761	< 0.008					< 0.008		
7-7200552	< 0.008	0.5	0.25	0.063	0.25	< 0.008	2	1
7-7150318	< 0.008	0.63	0.016	< 0.008	< 0.008	< 0.008	1	1
7-7150628	< 0.008	0.063	0.31	0.31	0.031	< 0.008	0.5	1
7-7149204	< 0.008	0.31	< 0.008	< 0.008	< 0.008	< 0.008	0.5	0.5
7-7116411	0.125	0.25	0.125	0.031	< 0.008	2	>16	2

Strain	Clioquinide	Closantel	Oxyclozanide	Rifaxanide	Tribromsalan	Fidaxomycin	Metronidazole	Vancomycin
12055	< 0.008	0.063	0.016	< 0.008	0.016	< 0.008	1	1
12060	< 0.008	0.125	0.031	0.031	0.063	< 0.008	1	0.5
12061	< 0.008	0.063	0.016	0.016	0.016	< 0.008	1	0.25
12062	< 0.008	0.063	< 0.008	0.016	< 0.008	< 0.008	0.125	0.125
12063	< 0.008	0.031	0.016	< 0.008	0.016	< 0.008	0.25	0.25
12064	< 0.008	0.031	< 0.008	< 0.008	< 0.008	< 0.008	0.031	0.5
12065	< 0.008	0.063	0.031	0.016	0.016	< 0.008	1	0.5
12066	0.063	0.5	0.125	0.063	0.031	< 0.008	1	0.5

[00211] Table 6 shows that the tested halogenated salicylanilides were active against the tested strains. The most active compounds were clioxanide, rafoxanide and oxyclozanide, which compared favourably with the activity of the comparator compound fidaxomicin. All of the tested halogenated salicylanilides generally exhibited lower MIC values than the comparator compounds, metronidazole and vancomycin.

References:

- [1] H.-P. Schau, "J. F. MacFaddin, Media for Isolation - Cultivation - Identification - Maintenance of Medical Bacteria, Volume I. XI + 929 S., 163 Abb., 94 Tab. Baltimore, London 1985. Williams and Wilkins. \$ 90.00. ISBN: 0-683-05316-7," J. Basic Microbiol., vol. 26, no. 4, pp. 240–240, 1986.
- [2] L. Pospíšil, "[Isovitalex--a chemically definable enricher of culture media for Neisseria gonorrhoeae]," Ceskoslovenská Dermatol., vol. 46, no. 1, pp. 23–25, Feb. 1971.
- [3] A. N. Edwards, J. M. Suarez, and S. M. McBride, "Culturing and Maintaining Clostridium difficile in an Anaerobic Environment," J. Vis. Exp. JoVE, no. 79, p. e50787, Sep. 2013.

Example 5: Sporulation Study

[00212] *Clostridium difficile* is a spore forming bacteria that causes severe diarrhea in healthcare settings. The spore is the infective agent, and is implicated in disease transmission and recurrence. Prevention or inhibition of spore formation may therefore minimise the risk of transmission and recurrence of infection, particularly in a hospital environment. Currently the main treatments used for the treatment of *C. difficile* infections are vancomycin, metronidazole, rifaximin and fidaxomicin. It has been shown that fidaxomicin inhibits *C. difficile* sporulation [1].

[00213] The halogenated salicylanilide, rafoxanide was tested assess its ability to inhibit spore formation. Fidaxomicin was used as a comparator in the study.

METHODS

Bacterial strain

[00214] The *C. difficile* strain used in the study was 7-6011209, a clinical isolate from the MLST group ST002.

Antimicrobial agents

[00215] Rafoxanide and fidaxomicin (ex.Sigma-Aldrich) were prepared as 10 mg/mL stock solutions in dimethyl sulfoxide (DMSO). The compounds were diluted further to appropriate concentration in growth media prior to testing for their effect on sporulation.

Culture media and culture conditions

[00216] *C. difficile* strains were grown and cultured in Brucella bouillon supplemented with Isovitalax as described above in Example 3.

[00217] Sporulation was carried out using Clospore medium [2], comprising Special
5 Peptone Mix (Oxoid) 10 g/L, yeast extract 10 g/L, $(\text{NH}_4)_2 \text{SO}_4$ 0.6 g/L, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.12 g/L, $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ 0.08 g/L, K_2CO_3 3.48 g/L, KH_2PO_4 2.6 g/L, pH 7.9 ± 0.1 .

[00218] The germination medium was BHIS medium [3] containing 1 g/L of sodium taurocholate. BHIS medium comprises: Brain Heart Infusion 37 g/L, yeast extract 5 g/L, agar 15 g/L, L-cysteine 0.1% (w/v), glucose 0.5% (w/v) and FeSO_4 0.09% (w/v).

10 [00219] All cultures were performed at 37°C under anaerobic conditions in an anaerobic chamber as described in Example 3.

Sporulation kinetics

[00220] *C. difficile* was grown overnight on blood agar plates. One colony was transferred to 10 mL of Brucella bouillon enriched with Isovitalax and grown overnight. Clospore
15 medium containing the appropriate concentration of the test compound was inoculated at 1 % with the overnight culture. Samples were withdrawn every 24 hours for quantitation of heat-resistant spores (survivors after incubation at 65°C for 20 minutes). Spores were serially diluted in 0.09% NaCl and plated on BHIS agar supplemented with 0.1% sodium taurocholate to grow the spores for quantitation.

20 [00221] Concentrations of the test compounds were normalized to the MIC such that they were at least 8-fold above the MIC of the respective compound (8-fold for rafoxanide, and >8-fold for fidaxomicin).

RESULTS

[00222] The impact of the test compounds on sporulation kinetics is shown in Figure 1.

25 [00223] The negative control sporulates rapidly reached a value of 2×10^5 by 24 hours and approached its maximum count of approximately 10^7 by 48 hours sporulation. Rafoxanide, at 8-fold MIC, suppressed formation of spores throughout the 96 hour study period. The comparator compound fidaxomicin, at >8-fold MIC, suppressed formation of spores for the first 48 hours of the study, however, increased spore formation compared to
30 rafoxanide occurred at the 72 and 96 hour time points.

[00224] The data illustrated by Figure 1 shows that rafoxanide suppresses spore formation more effectively than fidaxomicin at a fixed effect level relative to MIC in this study. These results suggest that rafoxanide may inhibit the shedding of *C. difficile* spores and as such be effective in controlling the spread of infection in, for example, a hospital

environment. The compound may also be useful in minimising the risk of recurrent infections in patients.


References:

- 5 [1] F. Babakhani, L. Bouillaut, P. Sears, C. Sims, A. Gomez, and A. L. Sonenshein, "Fidaxomicin inhibits toxin production in *Clostridium difficile*," *J. Antimicrob. Chemother.*, vol. 68, no. 3, pp. 515–522, Mar. 2013.
- [2] J. Perez, V. S. Springthorpe, and S. A. Sattar, "Clospore: a liquid medium for producing high titers of semi-purified spores of *Clostridium difficile*," *J. AOAC Int.*, vol. 94, no. 2, pp. 618–626, Apr. 2011.
- 10 [3] C. J. Smith, S. M. Markowitz, and F. L. Macrina, "Transferable tetracycline resistance in *Clostridium difficile*," *Antimicrob. Agents Chemother.*, vol. 19, no. 6, pp. 997–1003, Jun. 1981.

[00225] The invention is further illustrated by the following numbered clauses:

- 15 1. A halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof for use in the treatment of an infection in a subject caused by *Clostridium* bacteria.
2. The halogenated salicylanilide for the use of Clause 1 wherein the infection is caused by *Clostridium difficile*.
3. The halogenated salicylanilide for the use of Clause 2, wherein the infection is a
20 *Clostridium difficile* associated disease.
4. The halogenated salicylanilide for the use of Clause 3, wherein *Clostridium difficile* associated disease is diarrhoea, colitis (for example pseudomembranous colitis) or toxic megacolon.
5. The halogenated salicylanilide for the use of any of Clauses 1 to 4, wherein the
25 *Clostridium* infection is an antibiotic induced *Clostridium* infection, wherein the antibiotic is other than a halogenated salicylanilide.
6. The halogenated salicylanilide for the use of Clause 5, wherein the antibiotic other than a halogenated salicylanilide is selected from clindamycin, a cephalosporin (for example cefotaxime and ceftaidime), ampicillin, amoxicillin and a quinolone (for example a
30 fluoroquinolone, optionally ciprofloxacin or levofloxacin).
7. The halogenated salicylanilide for the use of any of Clauses 1 to 6, wherein the *Clostridium* infection has not been treated with an antibiotic prior to administration of the halogenated salicylanilide to the subject.

8. The halogenated salicylanilide for the use of any of Clauses 1 to 6, wherein the subject has a *Clostridium* infection which has recurred following treatment with an antibiotic other than a halogenated salicylanilide.
9. The halogenated salicylanilide for the use of Clause 8, wherein *Clostridium* infection has recurred after being treated with an antibiotic selected from metronidazole, vancomycin and fidaxomicin.
10. The halogenated salicylanilide for the use of any of Clauses 1 to 6, wherein the *Clostridium* infection is refractory to a prior antibiotic treatment other than a halogenated salicylanilide.
11. The halogenated salicylanilide for the use of Clause 10, wherein the prior antibiotic treatment is selected from metronidazole, vancomycin and fidaxomicin.
12. The halogenated salicylanilide for the use of any preceding Clause wherein the infection is caused by the NAP1/027/BI *C. difficile* strain.
13. The halogenated salicylanilide for the use of any preceding Clause, wherein the halogenated salicylanilide is of the formula (I):



wherein

- X is O or S;
- 20 R^1 and R^2 are at each occurrence independently selected from halo;
- R^3 and R^4 are at each occurrence independently selected from H, C_{1-6} alkyl, $-OR^{A1}$, $-NO_2$ and $-CN$;
- R^5 is H or $-L^1-R^7$;
- R^6 is H or $-C(O)R^{A2}$;
- 25 L^1 is selected from a bond, O, S, or $-(CR^{A3}R^B)_o-$, wherein o is 1 or 2;
- R^6 is phenyl, unsubstituted or substituted with 1, 2, or 3 groups selected from halo, C_{1-4} alkyl, $-OR^{A4}$, $-NO_2$ and $-CN$;

R^{A1}, R^{A2}, R^{A3} and R^{A4} are at each occurrence independently selected from H and C₁₋₄ alkyl;

R^B is at each occurrence selected from H, C₁₋₄ alkyl and -CN;

n and p are each independently selected from 0, 1, 2, 3 or 4, with the proviso that n+p is at least 1;

5 t and v are independently selected from 0, 1 and 2;

or a pharmaceutically acceptable salt, or ester thereof

14. The halogenated salicylanilide for the use of Clause 13, wherein X is O.


15. The halogenated salicylanilide for the use of Clause 13 or Clause 14, wherein R⁶ is H.


10 16. The halogenated salicylanilide for the use of any of Clauses 13 to 15, wherein R³ and R⁴ are at each occurrence independently selected from H, C₁₋₄ alkyl, -OR^{A1} and -NO₂.

17. The halogenated salicylanilide for the use of any of Clauses 13 to 16, wherein L¹ is selected from O, -CH₂- and -CH(CN)-.

15 18. The halogenated salicylanilide for the use of any of Clauses 13 to 17, wherein R⁷ is phenyl unsubstituted or substituted with 1, 2 or 3 groups selected from halo.


19. The halogenated salicylanilide for the use of any of Clauses 1 to 12, wherein the halogenated salicylanilide is selected from





oxyclozanide


Clioquanide



Dibromosalan

Tribromosalan

and



Niclosamine

;

or a pharmaceutically acceptable salt or ester thereof.

5 20. The halogenated salicylanilide for the use of any of Clauses 1 to 12, wherein the halogenated salicylanilide is selected from the group consisting of niclosamide, clioquanide, closantel, oxyclozanide, rafoxanide, tribromosalan, or a pharmaceutically acceptable salt or ester thereof.

10 21. Use of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof for the manufacture of a medicament for the treatment of an infection in a subject caused by *Clostridium* bacteria.

22. A method of treating an infection caused by *Clostridium* bacteria in a subject, the method comprising administering to said subject an effective amount of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof.


15 23. The halogenated salicylanilide for the use of any of Clauses 1 to 20, the use of Clause 21 or the method of Clause 22, wherein the halogenated salicylanilide is orally administered to the subject.

24. The halogenated salicylanilide for the use of any of Clauses 1 to 20, the use of Clause 21 or the method of Clause 22, wherein the subject is a human or warm blooded animal, optionally wherein the subject is a human.
25. The halogenated salicylanilide for the use of any of Clauses 1 to 20, the use of
5 Clause 21 or the method of Clause 22, wherein the subject is a human aged 65 years or older

CLAIMS

1. A halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof for use in the treatment of an infection in a subject caused by *Clostridium* bacteria.
- 5 2. The halogenated salicylanilide for the use of claim 1 wherein the infection is caused by *Clostridium difficile*.
3. The halogenated salicylanilide for the use of claim 2, wherein the infection is a *Clostridium difficile* associated disease.
4. The halogenated salicylanilide for the use of claim 3, wherein *Clostridium difficile*
10 associated disease is diarrhoea, colitis (for example pseudomembranous colitis) or toxic megacolon.
5. The halogenated salicylanilide for the use of any of claims 1 to 4, wherein the *Clostridium* infection is an antibiotic induced *Clostridium* infection, wherein the antibiotic which induced the infection is other than the halogenated salicylanilide.
- 15 6. The halogenated salicylanilide for the use of claim 5, wherein the antibiotic which induced the infection is selected from clindamycin, a cephalosporin (for example cefotaxime and ceftaidime), ampicillin, amoxicillin and a quinolone (for example a fluoroquinolone, optionally ciprofloxacin or levofloxacin).
7. The halogenated salicylanilide for the use of any of claims 1 to 4, wherein the
20 *Clostridium* infection is induced by a gastric acid suppressive agent.
8. The halogenated salicylanilide for the use of any of claims 1 to 7, wherein the *Clostridium* is resistant to an antibiotic agent other than the halogenated salicylanilide.
9. The halogenated salicylanilide for the use of claim 8, wherein the *Clostridium* is a
25 *Clostridium difficile* strain that is resistant to an antibiotic agent selected from metronidazole, vancomycin, fidaxomicin and a rifamycin.
10. The halogenated salicylanilide for the use of any of claims 1 to 9, wherein the *Clostridium* infection has not been treated with an antibiotic prior to administration of the halogenated salicylanilide to the subject.
11. The halogenated salicylanilide for the use of any of claims 1 to 9, wherein the
30 subject has a recurrent *Clostridium* infection, for example a *Clostridium* infection which has recurred following prior treatment of the subject with an antibiotic other than the halogenated salicylanilide.

12. The halogenated salicylanilide for the use of claim 11, wherein the *Clostridium* infection has recurred after prior treatment of the subject with an antibiotic, for example an antibiotic selected from metronidazole, vancomycin, fidaxomicin and a rifamycin; optionally wherein the *Clostridium* infection has recurred after being treated with an antibiotic selected from metronidazole, vancomycin, fidaxomicin.
13. A halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof, for use in a method of preventing or inhibiting the transmission or spread of a *C. difficile* infection, the method comprising administering the halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof to a subject with a *C. difficile* infection.
14. A halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof, for use in a method of preventing or inhibiting recurrence of *C. difficile* infection in a subject with a *C. difficile* infection, the method comprising administering the halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof to the subject.
15. A halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof, for use in preventing or inhibiting sporulation of *C. difficile* in a subject infected with *C. difficile*.
16. The halogenated salicylanilide for the use of any of claims 13 to 15, wherein the *C. difficile* is resistant to an antibiotic agent other than the halogenated salicylanilide.
17. The halogenated salicylanilide for the use of claim 16, wherein the *C. difficile* is resistant to an antibiotic agent selected from metronidazole, vancomycin, fidaxomicin and a rifamycin.
18. The halogenated salicylanilide for the use of any preceding claim, wherein the halogenated salicylanilide is orally administered to the subject.
19. The halogenated salicylanilide for the use of any preceding claim, wherein the subject is a human or warm blooded animal, optionally wherein the subject is a human.
20. The halogenated salicylanilide for the use of claim 19, wherein the subject is a human aged 65 years or older.
21. The halogenated salicylanilide for the use of any preceding claim wherein the infection is caused by the NAP1/027/BI *C. difficile* strain.
22. The halogenated salicylanilide for the use of any preceding claim, wherein the halogenated salicylanilide is of the formula (I):



(I)

wherein

X is O or S;

5 R^1 and R^2 are at each occurrence independently selected from halo;

R^3 and R^4 are at each occurrence independently selected from H, C_{1-6} alkyl, $-OR^{A1}$, $-NO_2$ and $-CN$;

R^5 is H or $-L^1-R^7$;

R^6 is H or $-C(O)R^{A2}$;

10 L^1 is selected from a bond, O, S, or $-(CR^{A3}R^B)_o-$, wherein o is 1 or 2;

R^6 is phenyl, unsubstituted or substituted with 1, 2, or 3 groups selected from halo, C_{1-4} alkyl, $-OR^{A4}$, $-NO_2$ and $-CN$;

R^{A1} , R^{A2} , R^{A3} and R^{A4} are at each occurrence independently selected from H and C_{1-4} alkyl;

R^B is at each occurrence selected from H, C_{1-4} alkyl and $-CN$;

15 n and p are each independently selected from 0, 1, 2, 3 or 4, with the proviso that n+p is at least 1;

t and v are independently selected from 0, 1 and 2;

or a pharmaceutically acceptable salt, or ester thereof.

23. The halogenated salicylanilide for the use of claim 22, wherein X is O.


20 24. The halogenated salicylanilide for the use of claim 22 or claim 23, wherein R^6 is H.


25. The halogenated salicylanilide for the use of any of claims 22 to 24, wherein R^3 and R^4 are at each occurrence independently selected from H, C_{1-4} alkyl, $-OR^{A1}$ and $-NO_2$.

26. The halogenated salicylanilide for the use of any of claims 22 to 25, wherein L^1 is selected from O, $-CH_2-$ and $-CH(CN)-$.

27. The halogenated salicylanilide for the use of any of claims 22 to 26, wherein R⁷ is phenyl unsubstituted or substituted with 1, 2 or 3 groups selected from halo.

28. The halogenated salicylanilide for the use of any of claims 1 to 21, wherein the halogenated salicylanilide is selected from:





niclosamide

;

or a pharmaceutically acceptable salt or ester thereof.

29. The halogenated salicylanilide for the use of any of claims 1 to 21, wherein the halogenated salicylanilide is selected from the group consisting of niclosamide, clixoxide,
 5 closantel, oxyclozanide, rafoxanide, tribromosalan, or a pharmaceutically acceptable salt or ester thereof.

30. The halogenated salicylanilide for the use of any of claims 1 to 21, wherein the halogenated salicylanilide is selected from the group consisting of the group consisting of clixoxide, closantel, oxyclozanide, rafoxanide and tribromosalan, or a pharmaceutically
 10 acceptable salt thereof.

31. The halogenated salicylanilide for the use of any of claims 1 to 21, wherein the halogenated salicylanilide is selected from the group consisting of the group consisting of rafoxanide, oxyclozanide and clixoxide, or a pharmaceutically acceptable salt thereof,
 15 optionally wherein the halogenated salicylanilide is rafoxanide, or a pharmaceutically acceptable salt thereof.

32. A method of treating an infection caused by *Clostridium* bacteria in a subject, the method comprising administering to said subject an effective amount of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof.

33. The method of claim 32 wherein the infection is caused by *Clostridium difficile*.

20 34. The method of claim 33, wherein the infection is a *Clostridium difficile* associated disease.

35. The method of claim 34, wherein *Clostridium difficile* associated disease is diarrhoea, colitis (for example pseudomembranous colitis) or toxic megacolon.

25 36. The method of any of claims 32 to 34, wherein the *Clostridium* infection is an antibiotic induced *Clostridium* infection, wherein the antibiotic which induced the infection is other than the halogenated salicylanilide.

37. The method of claim 36, wherein the antibiotic which induced the infection is selected from clindamycin, a cephalosporin (for example cefotaxime and ceftidime),

ampicillin, amoxicillin and a quinolone (for example a fluoroquinolone, optionally ciprofloxacin or levofloxacin).

38. The method of any of claims 32 to 35, wherein the *Clostridium* infection is induced by a gastric acid suppressive agent.

5 39. The method of any of claims 32 to 38, wherein the *Clostridium* is resistant to an antibiotic agent other than the halogenated salicylanilide.

40. The method of claim 39, wherein the *Clostridium* is a *Clostridium difficile* strain that is resistant to an antibiotic agent selected from metronidazole, vancomycin, fidaxomicin and a rifamycin.

10 41. The method of any of claims 32 to 40, wherein the *Clostridium* infection has not been treated with an antibiotic prior to administration of the halogenated salicylanilide to the subject.

42. The method of any of claims 32 to 40, wherein the subject has a recurrent *Clostridium* infection, for example a *Clostridium* infection which has recurred following prior
15 treatment with an antibiotic other than the halogenated salicylanilide.

43. The method of claim 42, wherein the *Clostridium* infection has recurred after being treated with an antibiotic selected from metronidazole, vancomycin, fidaxomicin and a rifamycin.

44. A method for preventing or inhibiting the transmission or spread of a *C. difficile*
20 infection, the method comprising administering an effective amount of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof to a subject with a *C. difficile* infection.

45. A method of preventing or inhibiting recurrence of *C. difficile* infection in a subject with a *C. difficile* infection, the method comprising administering to said subject a
25 halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof.

46. A method of preventing or inhibiting sporulation of *C. difficile* in a subject infected with *C. difficile*, the method comprising administering to said subject an effective amount of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof.

47. The method of any of claims 44 to 46, wherein the *C. difficile* is resistant to an
30 antibiotic agent.

48. The method of claim 47, wherein the *C. difficile* is resistant to an antibiotic agent selected from metronidazole, vancomycin, fidaxomicin and a rifamycin.

49. The method of any of claims 32 to 48, wherein the halogenated salicylanilide is orally administered to the subject.
50. The method of any of claims 32 to 49, wherein the halogenated salicylanilide is administered to the subject concurrently with another therapeutic agent; for example
5 wherein the other therapeutic agent is a *C. difficile* therapy selected from an antibiotic active against *C. difficile*, a microbiome therapeutic, a faecal transplant, a vaccine or an antibody therapy.
51. The method of any of claims 32 to 50, wherein the subject is a human or warm blooded animal, optionally wherein the subject is a human.
- 10 52. The method of claim 51, wherein the subject is a human aged 65 years or older.
53. The method of any of claims 32 to 50, wherein the halogenated salicylanilide is as defined in any of claims 22 to 31.
54. Use of a halogenated salicylanilide, or a pharmaceutically acceptable salt or ester thereof for the manufacture of a medicament for the treatment of an infection in a subject
15 caused by *Clostridium* bacteria, for example *C. difficile*.
55. The use of claim 54, wherein the halogenated salicylanilide is orally administered to the subject.
56. The use of claim 54, wherein the subject is a human or warm blooded animal, optionally wherein the subject is a human; optionally wherein the subject is a human aged
20 65 years or older.




Figure 1

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/061968

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K31/609 A61K31/612 A61P31/04
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
A61K
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PAUK KAREL ET AL: "New derivatives of salicylamides: Preparation and antimicrobial activity against various bacterial species", BIOORGANIC & MEDICINAL CHEMISTRY, vol. 21, no. 21, 24 August 2013 (2013-08-24), pages 6574-6581, XP028732732, ISSN: 0968-0896, DOI: 10.1016/J.BMC.2013.08.029 figure 1; table 1 page 6575, right-hand column, paragraph 3 - page 6576, left-hand column, paragraph 1 page 6578, left-hand column, last paragraph ----- -/--	1-56

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search 29 July 2016	Date of mailing of the international search report 05/08/2016
---	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Gradassi, Giulia
--	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/061968

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 2014/125075 A1 (TECH UNIVERSITÄT BERLIN [DE]; CIRAD [FR]) 21 August 2014 (2014-08-21)</p> <p>page 433, lines 8-12 page 436, paragraphs 4,6 compounds 1,4,12,19,22,23,26,28,32</p> <p>-----</p>	<p>1-4, 7-21, 32-52, 54-56</p>
Y	<p>RAJMOHAN RAJAMUTHIAH ET AL: "Repurposing Salicylanilide Anthelmintic Drugs to Combat Drug Resistant Staphylococcus aureus", PLOS ONE, vol. 10, no. 4, 21 April 2015 (2015-04-21) , page e0124595, XP055221148, DOI: 10.1371/journal.pone.0124595 the whole document</p> <p>-----</p>	<p>1-56</p>
Y	<p>WO 2010/005836 A2 (ACTIVBIOTICS PHARMA LLC [US]; SAYADA CHALOM [LU]) 14 January 2010 (2010-01-14) the whole document</p> <p>-----</p>	<p>1-56</p>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/EP2016/061968

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 2014125075	A1	21-08-2014	CA 2901576 A1	21-08-2014
			CN 105143176 A	09-12-2015
			EP 2956437 A1	23-12-2015
			JP 2016513103 A	12-05-2016
			US 2015376120 A1	31-12-2015
			WO 2014125075 A1	21-08-2014

WO 2010005836	A2	14-01-2010	CA 2730274 A1	14-01-2010
			CN 102143751 A	03-08-2011
			EA 201100154 A1	30-08-2011
			EP 2313099 A2	27-04-2011
			US 2011117154 A1	19-05-2011
			WO 2010005836 A2	14-01-2010
