



(86) **Date de dépôt PCT/PCT Filing Date:** 2006/12/05
 (87) **Date publication PCT/PCT Publication Date:** 2007/06/21
 (45) **Date de délivrance/Issue Date:** 2015/11/03
 (85) **Entrée phase nationale/National Entry:** 2008/06/05
 (86) **N° demande PCT/PCT Application No.:** IB 2006/004076
 (87) **N° publication PCT/PCT Publication No.:** 2007/069090
 (30) **Priorités/Priorities:** 2005/12/06 (US11/294,421);
 2005/12/06 (EP05292590.6)

(51) **Cl.Int./Int.Cl. C07K 14/00** (2006.01),
C07K 7/08 (2006.01), **C12N 15/11** (2006.01),
C12N 15/87 (2006.01), **A61K 38/16** (2006.01),
A61K 38/10 (2006.01), **A61K 48/00** (2006.01),
A61P 35/00 (2006.01)
 (72) **Inventeurs/Inventors:**
 DIVITA, GILLES, FR;
 HEITZ, FREDERIC, FR;
 MORRIS, MAY CATHERINE, FR;
 ALDRIAN-HERRADA, GUDRUN, FR
 (73) **Propriétaires/Owners:**
 CENTRE NATIONAL DE LA RECHERCHE
 SCIENTIFIQUE, FR;

(54) **Titre : PEPTIDES DE PENETRATION CELLULAIRE POUR LA DELIVRANCE INTRACELLULAIRE DE MOLECULES**
 (54) **Title: CELL PENETRATING PEPTIDES FOR INTRACELLULAR DELIVERY OF MOLECULES**

(57) **Abrégé/Abstract:**

The present invention concerns cell-penetrating peptides which comprise an amino acid sequence consisting of GLX₉RALX₉RX₁LX₂RSLX₉X₃X₄X₅X₆X₇X₈ (SEQ ID No: 1), wherein X₁ is A, L or G, X₂ is W or none, X₃ is R or K, X₄ is K, L or S, X₅ is L or K, X₆ is R or W, X₇ is K or S, and X₈ is A, V or Q, and X₉ is W, F or Y. These CPPs can be used as vectors for delivering nucleic acids and/or proteins and/or peptides to cells, in vitro or in vivo.

(73) **Propriétaires(suite)/Owners(continued):**AFFYMETRIX, INC., US

(74) **Agent:** BORDEN LADNER GERVAIS LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2007 (21.06.2007)

PCT

(10) International Publication Number
WO 2007/069090 A3

(51) International Patent Classification:

C07K 14/00 (2006.01) *A61P 35/00* (2006.01)
C07K 7/08 (2006.01) *A61K 38/10* (2006.01)
C12N 15/11 (2006.01) *A61K 38/16* (2006.01)
C12N 15/87 (2006.01) *A61K 48/00* (2006.01)

(21) International Application Number:

PCT/IB2006/004076

(22) International Filing Date:

5 December 2006 (05.12.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

05292590.6 6 December 2005 (06.12.2005) EP
11/294,421 6 December 2005 (06.12.2005) US

(71) Applicants (*for all designated States except US*):
CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE [FR/FR]; 3, rue Michel Ange, F-75016 Paris (FR). **PANOMICS INC.** [US/US]; 6519 Dumbarton Circle, Fremont, CA 94555 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **DIVITA, Gilles** [FR/FR]; 476 rue de la Rave, F-34130 Mauguio (FR). **HEITZ, Frédéric** [FR/FR]; 3 rue Traves, F-34790 Grabels (FR). **MORRIS, May Catherine** [GB/FR]; 476 rue de la Rave, F-34130 Mauguio (FR). **ALDRIAN-HERRADA, Gudrun** [AT/FR]; 1161, route de Vendargues, F-34730 Prades-le-Lez (FR).

(74) Agents: **MARCADE, Véronique** et al.; Cabinet Ores, 36, rue de St Péresbourg, F-75008 Paris (FR).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(88) Date of publication of the international search report:

28 February 2008

(54) Title: CELL PENETRATING PEPTIDES FOR INTRACELLULAR DELIVERY OF MOLECULES

(57) Abstract: The present invention concerns cell-penetrating peptides which comprise an amino acid sequence consisting of GLX₉RALX₉RX₁LX₂RSLX₉X₃X₄X₅X₆X₇X₈ (SEQ ID No: 1), wherein X₁ is A, L or G, X₂ is W or none, X₃ is R or K, X₄ is K, L or S, X₅ is L or K, X₆ is R or W, X₇ is K or S, and X₈ is A, V or Q, and X₉ is W, F or Y. These CPPs can be used as vectors for delivering nucleic acids and/or proteins and/or peptides to cells, in vitro or in vivo.



WO 2007/069090 A3

CELL PENETRATING PEPTIDES FOR INTRACELLULAR DELIVERY OF MOLECULES

The present invention pertains to the field of intracellular delivery of molecules such as proteins or nucleic acids. In particular, the invention relates to a new
5 cell-penetrating peptide (CPP) family, which exhibits high efficiency and low toxicity.

Cellular internalization of large hydrophilic therapeutic agents such as proteins or nucleic acids is still a challenging task because of the presence of the plasma membrane, which constitutes an impermeable barrier for such molecules. In order to circumvent this problem, several methods of carrier-mediated delivery systems have been
10 developed. Among them, much attention has recently been given to the use of peptide-based delivery systems. The use of peptides with cell-penetrating properties has several advantages, which are mainly due to the various modifications that can be done to the peptide sequence. This allows the engineering of carriers addressing different cellular subdomains and/or able to transport various types of cargoes.

15 Many CPPs were designed from sequences of membrane-interacting proteins such as fusion proteins, signal peptides, transmembrane domains and antimicrobial peptides (Morris, Chaloin et al. 2000; Jarver and Langel 2004; El-Andaloussi, Holm et al. 2005). Within these sequences, short sequences called Protein Transduction Domains or PTDs proved to efficiently cross biological membranes without
20 the need of a carrier or of a receptor, and to deliver peptides or proteins into intracellular compartments. Many investigations suggested that the use of PTD-based peptides could be of major importance for therapies against viral diseases or cancers. Among these, the third helix of the homeodomain of *antennapedia* called penetratin (Joliot and Prochiantz 2004), the Tat peptide derived from the transactivating protein Tat of HIV-1 (Wadia and Dowdy
25 2002), transportan (Pooga, Hallbrink et al. 1998) and VP22 (Elliott and O'Hare 1997) were used to improve the cellular uptake of peptides, proteins or oligonucleotides.

A second category of cell-penetrating peptides, designated as amphipathic peptides, has been described. An amphipathic molecule can be defined, in short, as consisting of two domains: a hydrophilic (polar) domain and a hydrophobic (non-
30 polar) domain. For peptides, the amphipathic character can arise either from the primary structure, or from the secondary structure. Primary amphipathic peptides can be defined as the sequential assembly of a domain of hydrophobic residues with a domain of hydrophilic residues. Secondary amphipathic peptides are generated by the conformational state which allows the positioning of the hydrophobic and hydrophilic residues on opposite sides of the
35 molecule.

Other peptides, such as polyarginine-based peptides, calcitonin-derived peptides, and oligomers, have also been proposed as tools for intracellular delivery of therapeutics.

5 Deshayes *et al.* have reviewed the available data concerning the use of the above-mentioned cell-penetrating peptides for delivering molecules into cells (Deshayes, Morris *et al.* 2005). It appears that the current vectors are limited, because of their lack of efficiency and/or their toxicity, and that little is known about the pathway of their cellular uptake, which constitutes a handicap for improving their efficiency.

10 The present invention pertains to a new family of cell-penetrating peptides having particularly advantageous properties.

A first aspect of the present invention is hence a cell-penetrating peptide which comprises an amino acid sequence consisting of GLX₉RALX₉RX₁LX₂RSLX₉X₃X₄X₅X₆X₇X₈ (SEQ ID No: 1), wherein X₁ is A, L or G, X₂ is W or none, X₃ is R or K, X₄ is K, L or S, X₅ is L or K, X₆ is R or W, X₇ is K or S, X₈ is 15 A, V or Q, and X₉ is W, F or Y. According to the invention, X₉ can also be another aromatic chemical group. In a preferred embodiment of the invention, X₉ is W, which means that the cell-penetrating peptide comprises an amino acid sequence consisting of GLWRALWRX₁LX₂RSLWX₃X₄X₅X₆X₇X₈ (SEQ ID No: 25), with X₁ to X₈ as described above. The CPPs according to the invention have a secondary amphipathic structure. In the 20 CPPs according to the invention, the amino acids can be either L- or D-amino acids.

In the present text, a "cell-penetrating peptide", also called a "peptide carrier", is a molecule, the core of which is a peptide. Other chemical groups can however be covalently bound to said peptidic core, in order to improve the overall stability of the molecule, and/or to provide it with additional properties, such as targeting ability. For 25 example, a cell-penetrating peptide according to the invention can further comprise, covalently linked to the C-terminal extremity of the peptidic core of SEQ ID No: 1, one or several groups chosen amongst a cysteamide, a cysteine, a thiol, an amide, a carboxyl, a linear or ramified C₁-C₆ alkyl optionally substituted, a primary or secondary amine, an osidic derivative, a lipid, a phospholipid, a fatty acid, a cholesterol, a poly-ethylene glycol, 30 a nuclear localization signal (NLS), and/or a targeting molecule. Alternatively or additionally, the cell-penetrating peptide of the invention can also comprise, covalently linked to the N-terminal end of the peptidic core of SEQ ID No: 1, one or several chemical entities chosen amongst an acetyl, a fatty acid, a cholesterol, a poly-ethylene glycol, a nuclear localization signal, and/or a targeting molecule. If necessary, for example in the 35 case of N-terminal addition of cholesterol, a peptidic bridge can be used to bind a non-peptidic molecule to the peptidic core of the CPP. An example of such a bridge is -CA_β-.

Although some of the entities which can be covalently bound to the peptide of SEQ ID No: 1 comprise amino acids, the invention does not encompass every

peptide or protein comprising SEQ ID No: 1, whatever the amino acids surrounding the core sequence of SEQ ID No: 1. Indeed, the only additional amino acid sequences that can be bound to a CPP according to the invention are the following:

- 5 (i) a peptidic bridge such as CA_β: the size of such a bridge will never exceed 5 amino acids. Preferably, the bridge consists of 1, 2 or 3 amino acids.
- 10 (ii) a nuclear localization signal (NLS): such signals are well known in the scientific literature. An example of NLS that can be used according to the present invention is PKKKRKV (SEQ ID No: 27). Other NLS which can be used can contain up to 11 amino acid residues, in the case of a bipartite NLS (Morris, Chaloin et al. 2002).
- 15 (iii) targeting peptides: targeting molecules which can advantageously be bound to the CPPs according to the invention, can be of peptidic nature. Examples of targeting peptides are epitopes, antibody fragments (Fab) which specifically target tissues or cell surface components or receptors, and the like. Whatever the molecules bound to the peptidic core of SEQ ID No: 1, CPPs according to the invention must retain their vector efficiency, *i.e.*, their ability (i) to interact with the cargo, and (ii) to deliver said cargo into the cells.
- 20 Therefore, in the context of the present invention, the sequence of a targeting peptide will preferably be no longer than 15 amino acids. A examples of such peptides are the RGD motif, NGR, transferrine, antibody fragments, and any of the peptides listed in Table 1 of the review by Allen (Allen 2002).

25 Apart from the targeting peptides, other targeting molecules can be bound to the CPPs of the invention, in order to obtain a specific delivery of molecules to particular cell types and/or organs. For example, sugars, such as monosaccharides (ex : glucose, galactose, glucosamine ou galactosamine), oligosaccharides, polysaccharides, or their analogues, as well as some oligonucleotides, or some organic molecules such as

30 folate, can be used to that aim. Their targeting activity is due to the fact that they are recognized as ligands by some receptors which are over-expressed at the surface of cells in the zone of interest.

In a particular embodiment of the cell-penetrating peptide according to the invention, the amino acid sequence of the peptidic core is chosen in the group

35 consisting of

- GLWRALWRLLRSLWRLWKA (SEQ ID No: 2);
- GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3);
- GLWRALWRALRSLWKLKRKV (SEQ ID No: 4);

- GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5);
- GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6);
- GLWRALWRLLRSLWLLWKA (SEQ ID No: 7);
- GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8);
- 5 - GLWRALWRALWRSLWKSQRKV (SEQ ID No: 9);
- GLWRALWRALWRSLWKKKRKV (SEQ ID No: 10); and
- GLWRALWRLLRSLWLLWSQ (SEQ ID No: 11).

Preferred cell-penetrating peptides according to the invention are
 CADY-1 (Ac-GLWRALWRLLRSLWLLWKA-Cya) and CADY-2 (Ac-
 10 GLWRALWRALWRSLWKLKWKV-Cya).

Another aspect of the present invention is a complex comprising a cell-
 penetrating peptide as described above, and a cargo selected amongst nucleic acids,
 peptides, proteins, contrast agents, and toxins.

In a particular embodiment of the complexes of the invention, the cargo
 15 is a siRNA selected to silence a target mRNA. In this embodiment, the amino acid
 sequence of (core of) the cell-penetrating peptide preferably is
 GLWRALWRX₁LX₂RSLWX₃X₄X₄X₅KX₆ (SEQ ID No: 26), wherein X₁ is A, L or G, X₂
 is W or none, X₃ is R or K, X₄ is K or L, X₅ is R or W, X₆ is A or V. For example, the
 amino acid sequence of the cell-penetrating peptide is chosen in the group consisting of :

- 20 - GLWRALWRLLRSLWLLWKA (SEQ ID No: 2);
- GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3);
- GLWRALWRALRSLWKLKRKV (SEQ ID No: 4);
- GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5);
- GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6); and
- 25 - GLWRALWRGLRSLWKKKRKV (SEQ ID No: 7).

According to the invention, preferred CPPs for siRNA delivery have an
 amino acid sequence corresponding to SEQ ID Nos 2, 6 or 7.

In another embodiment of the complexes of the invention, the cargo is a
 peptide or a protein. In this embodiment, the amino acid sequence of (core of) the cell-
 30 penetrating peptide is SEQ ID No: 1, as described above, preferably with the proviso that
 X₁ is not G and X₈ is not A. Examples of amino acid sequences of cell-penetrating peptides
 that can be incorporated in complexes with peptides or proteins are the following:

- GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8)
- GLWRALWRALWRSLWKSQRKV (SEQ ID No: 9)
- 35 - GLWRALWRALWRSLWKKKRKV (SEQ ID No: 10)
- GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3)
- GLWRALWRLLRSLWLLWSQ (SEQ ID No: 11)
- GLWRALWRLLRSLWLLWSQPKKKKRKV (SEQ ID No: 12)

According to the invention, preferred CPPs for proteins and peptide delivery have an amino acid sequence corresponding to SEQ ID Nos 8, 9 or 10. Depending on the cargo and the application, the skilled artisan can use the intracellular targeting properties of those CPPs, especially those of SEQ ID Nos 8 and 9, which target the nucleus and the cytoplasm, respectively.

Of course, the cell-penetrating peptides comprised in the complexes according to the invention can be modified as mentioned above, by binding additional molecules to the N-terminal and/or the C-terminal ends of their peptidic core of SEQ ID No: 1. In a preferred embodiment of such complexes, the cell-penetrating peptide comprises an acetyl group covalently linked to its N-terminus, and/or a cysteamide group covalently linked to its C-terminus. Alternatively or additionally, the cell-penetrating peptide further comprises a cholesterol molecule, covalently linked to its C-terminus or its N-terminus. When bound to the N-terminus, the cholesterol molecule can be linked via a bridge as described above, for example a CA_β bridge.

The inventors have investigated various parameters in order to optimize the cargo delivery, and have found that when the size of the complexes is < 500 nm, the mechanism of cellular uptake of said complexes is independent on the endosomal pathway. Hence, the size of the complexes according to the invention is preferably between 50 and 300 nm and, more preferably, between 100 and 200 nm. This corresponds, for siRNA/CADY complexes, to a ratio of 1/20 to 1/25.

According to an advantageous embodiment of the complex according to the invention, at least part of the cell-penetrating peptides are bound to a targeting molecule. As mentioned above, a targeting molecule can be a peptide, a sugar, *etc.* When only part of the CPPs are initially bound to a targeting molecule, this latter can be chosen bigger than the maximal size of targeting molecules which are acceptable when all the CPPs are bound to targeting molecules. The targeting molecule can be either covalently linked to a CPP, or bound to the complex via non-covalent bounds. In some cases, the cargo itself can be a targeting molecule, especially for targeting tumor cells.

A therapeutic composition comprising a complex as described above is, of course, part of the present invention.

In one particular aspect, the invention pertains to the use of a cell-penetrating peptide, or of a complex as described above, for the preparation of a therapeutic composition for use in anticancer therapy. For example, a therapeutic composition according to the invention comprises anti-cyclin B1 siRNA/CADY complexes, and/or p1p27/CADY complexes, and/or pRXL/CADY complexes.

The invention also pertains to a method for delivering a molecule into a cell *in vitro*, comprising a step of putting said cell into contact with a complex comprising said molecule and cell-penetrating peptides as described above.

The invention is further illustrated by the following figures and examples.

LEGENDS TO THE FIGURES

5 **Figure 1** : Schematic representation of CADY cellular uptake mechanism.

Figure 2 : Structure of CADY in the interaction with the membrane.

(A) modelisation

(B) CD experiments - Far UV CD spectra were recorded on a JASCO J-
10 810 spectro-polarimeter (JASCO Corporation, Tokyo, Japan) using a quartz cuvette with a path length of 1 mm. Spectra were corrected from the baseline buffer spectrum and smoothed with the JASCO software. Peptides were dissolved to a fixed concentration of 50 μM in water or in PBS. CD spectra were collected at 23°C and each spectrum represents the average of 4 scans. CADY-1 & CADY-2 adopt a α -helical structure characterized by
15 two minima at 208 nm and 222 nm, respectively.

Figure 3 : Formation of CADY/cargo complexes.

Binding of cargoes to CADY was monitored by steady state fluorescence spectroscopy. The intrinsic fluorescence associated to the Trp residues of CADY was used as sensitive probe. The fluorescence of CADY is centered at 340 nm upon excitation at 290
20 nm. Experiments were performed in phosphate buffer (PBS). A fixed concentration of CADY (0.1 μM) was titrated by increasing concentrations of cargoes.

(A) For the experiments with CADY-1, two different siRNAs (double stranded RNA) targeting cyclin B1 and GAPDH, were used.

(B) For the experiment with CADY-2, a short 17-residues peptide was
25 used.

Figure 4 : Toxicity of CADY peptides.

Cells were incubated in the presence of increasing concentrations (from 0.1 μM to 1 mM) of CADY-1, either alone (A) or in complexes with a cargo siRNA at a ratio of 1/20 (B). Cell viability was estimated using MTT assay, after 12 hrs of incubation.

30 **Figure 5** : CADY-1-mediated delivery of siRNA.

The different cell lines were cultured as described in the ATCC protocol. Cell lines were cultured in Dulbecco's Modified Eagle's Medium supplemented with 2mM glutamine, 1% antibiotics (streptomycin 10000 $\mu\text{g}/\text{ml}$, penicillin, 10000 IU/ ml) and 10% (w/v) fetal calf serum (FCS), at 37°C in a humidified atmosphere containing 5% CO_2 .
35 CADY/siRNA complexes were formed by incubation of siRNA (50 nM) with CADY at a molecular ratio of 1:20 (*i.e.*, one molecule of siRNA for 20 molecules of CADY), in 500 μl of DMEM for 30 min at 37°C. Cells grown to 40 to 60 % confluence were then overlaid

with preformed complexes. After 30 min incubation at 37°C, 1 ml of fresh DMEM supplemented with 10% fetal calf serum was added directly to the cells, without removing the overlay of CADY/siRNA complexes, and cells were returned to the incubator. Control experiments were performed with siRNA (50 nM) transfected with cationic lipids Oligofectamine (Invitrogen, Carlsbad, US), according to the guidelines of the manufacturer. A well established siRNA targeting GAPDH protein (Ambion-CA) was used as cargo. The knockdown of the GAPDH at the protein level was quantified 48 hrs after transfection by Western blot analysis using polyclonal antibody (Ref. ABCAM GAPDH antibody – ab9385 rabbit polyclonal). The percentage of GAPDH knockdown was estimated by comparison with control essays using a mismatch siRNA delivered with CADY at the same molecular ratio. Data are the average of 4 separated experiments.

Figure 6 : Mechanism of cellular uptake of CADY.

(A) impact of both the size of the particle and of the molar ratio CADY/siRNA on the efficiency of the siRNA. CADY/siRNA-GAPDH complexes were formed by incubation of siRNA (50 nM) with increasing molecular ratio of CADY from 5:1 to 50:1. Experiments were performed on Human HS-68 fibroblasts or Hela cell lines. The size of the particle was determined by light scattering and the knock down of GAPDH protein was quantified by Western blot analysis.

(B) role of the endosomal pathway on the efficiency of CADY-mediated siRNA delivery. CADY/siRNA-GAPDH complexes were formed by incubation of a fixed concentration of siRNA (50 nM) with CADY-1 at two molar ratios: 20:1 and 50:1, which correspond to CADY/siRNA-GAPDH particles of 100-200 nm and of 500 nm diameter size, respectively. In order to determine the cellular uptake mechanism of CADY, the transfection experiments were performed in the presence of different inhibitors of the endosomal pathway, as previously described by Dowdy and colleagues (Wadia, Stan et al. 2004). The following inhibitors of the endosomal pathway were used: Heparin (20µg/ml), Nystatin or Filipin (25 µg/ml), Cytochalasin D (5µM). The cells were incubated in the presence of the inhibitor one hour prior transfection; the inhibitor was then maintained in the culture medium for one hour after transfection. Cells were then extensively washed, trypsin was added in order to remove any CADY/siRNA complex associated to the cell membrane. Cells were the cultured for 24 additional hours. Experiments were performed on HS-68 fibroblasts or Hela cell lines. The knock down of GAPDH protein was quantified by Western blot analysis.

Figure 7 : CADY-mediated cyclin-B1 siRNA delivery

The effect of increasing concentrations of siRNA from 2 nM to 100 nM complexed with CADY (ratio 1/20) on cyclin B1 protein levels was analyzed in several cell lines (Hela cells, human fibroblasts (HS 68) and 293 cells). A stock solution of CADY/siRNA formulation was prepared using a siRNA concentration of 100 nM

associated with a CADY-1 at a molar ratio of 1/20. Lower concentrations of formulated siRNA (from 50 nM to 0.5 nM) were obtained by serial dilution of the stock solution in PBS, in order to maintain the siRNA/CADY ratio, and therefore the size of the particle around 200 nm diameter. SiRNA/CADY complexes were overlaid onto asynchronous
5 cultured cells in the presence of FBS (10%).

(A) The effect of Cyclin B1 siRNA was monitored on cell cycle progression by FACS analysis 30 hrs after transduction.

(B) Cyclin B1 protein levels were quantified by Western blotting 30 hrs after transduction. To that aim, cells were collected and proteins were separated on SDS
10 gel (12%), then transferred onto nitro-cellulose for Western blotting. Mouse monoclonal anti-Cyclin B1 antibodies (SC-245) and rabbit polyclonal anti-Cdk2 antibodies (SC-163) for Western blotting were obtained from Santa Cruz Biotechnology Inc., (Santa cruz, CA). Cdk2 protein kinase was used as a control to normalize protein levels.

Figure 8 : potency of CADY/siRNA complex to block cancer cell
15 proliferation.

Experiments were performed on MCF-7 or U2OS cells. Cells were treated on day 1 with different concentrations of siRNA and CADY/siRNA complexes (0.1 to 50 nM). A stock solution of CADY/siRNA formulation was prepared using a siRNA concentration of 100 nM associated with a CADY-1 at a molar ratio of 1:20. Lower
20 concentrations of formulated siRNA (from 50 nM to 0.5 nM) were obtained by serial dilution of the stock solution in PBS, in order to maintain the siRNA/CADY ratio and therefore the size of the particle around 200 nm diameter. Mismatched siRNA and free CADY were used as controls, and Oligofectamine (Invitrogen, USA) was used as lipid-based control delivery agent (manufacturer protocol). The inhibition of cell proliferation
25 was determined after 7 days.

Figure 9 : intratumoral injection of anti-cyclin B1/CADY complexes

The effect of the anti-cyclin B1 siRNA was investigated on a tumor animal model, using Swiss nude mice, injected subcutaneously with 2×10^7 15PC3 cells, A549 cells (human bronchoalveolar carcinoma) or HER2 cells. Different CADY/siRNA
30 complexes were injected directly into the tumor (100 μ l) at day 7 and at day15. siRNA/CADY complexes were obtained as described before at a ratio 1:20. The size of the tumor was measured every 2 days and the level of cyclin B1 protein was determined at Day20 by Western blot analysis. Results correspond to the average of 5 different animals.

Figure 10 : Comparison of CADY versus CADY-S

The efficiencies of CADY-1 and CADY-S vectors for delivering anti-cyclin B1 siRNA were compared on Hela/HS68 cells, using the same assay as described in the legend of figure 7A. The figure shows the results obtained on HS68 fibroblasts with a
35 CADY-1 vector devoid of cholesterol.

Figure 11 : CADY-S-mediated delivery of anti-cyclin B1 siRNA through intravenous injection to block tumor growth *in vivo*

Experiments were performed on two tumor mouse models.

(A) 1×10^6 5PC3 adenocarcinoma cells were injected subcutaneously into the flanks of NCR nude mice (day 1). siRNA/CADY-S complexes were obtained as described before at a ratio 1:20. CADY-S/siRNA complexes were injected intravenously at day 7. The size of the tumor was measured every day. Results correspond to the average of 5 different animals.

(B) 1×10^6 tumor cells expressing luciferase were injected intraperitoneally in the mice (4 to 8 week-old immune competent A/J female mice). Tumors were monitored using *in vivo* Imaging System (IVIS-Xenogen), which quantified total body bioluminescence after injection of D-luciferin. The tumor cells are visualized on three mice, each representative of one group (siRNA alone, CADY/siRNA complexes, and CADY-S/siRNA complexes).

Figure 12 : The different p27-derived peptides were associated with CADY-2 at a molar ratio of 1/20, since it was determined that at this ratio, the average size of the particle is of about 100-200 nm in diameter. The formulations were obtained in phosphate buffer, using a fixed concentration of peptide of 1 μ M. Cell lines were cultured in Dulbecco's Modified Eagle's Medium supplemented with 2mM glutamine, 1% antibiotics (streptomycin 10'000 μ g/ml, penicillin, 10'000 IU/ ml) and 10% (w/v) fetal calf serum (FCS), at 37°C in a humidified atmosphere containing 5% CO₂. Cells were synchronized by serum privation during 40 hrs, and cultured in Dulbecco's Modified Eagle's Medium supplemented with 2mM glutamine, 1% antibiotics. G1 synchronized cells were released upon serum addition, then overlaid with preformed CADY-2/peptide complexes. After 30 min incubation at 37°C, 1 ml of fresh DMEM supplemented with 10% fetal calf serum was added directly to the cells, without removing the overlay of CADY/peptide complexes, and cells were returned to the incubator. The level of cell arrest in G1 was analyzed by FACS 24 hrs after release.

(A) Protocol

(B) Results obtained with Jurkat, WI-38, HS-68 and HeLa cells.

Figure 13 : Effect of p1p27/CADY complexes on tumor growth *in vivo*

The tumor animal model consists of Swiss nude mice, injected subcutaneously with 2×10^7 Human H1299 adenocarcinoma cells. CADY2/p1p27 complex and free CADY-2 and peptides were injected directly into the tumor (100 – 300 μ g) after 1 and 4 weeks. p1p27/CADY complexes were obtained as described before at a ratio 1/20. Controls were performed by injecting PBS, CADY alone, or naked p1p27 instead of the p1p27/CADY-2 complex. The size of the tumor volume was measured with calipers once a week. Results correspond to the average of 5 different animals.

EXAMPLES**Example 1: Peptide synthesis**

Several peptides were synthesized, and selected for their ability to deliver
 5 different kinds of cargoes, especially siRNA (CADY-1 family) and peptides or proteins
 (CADY-2 family).

CADY-1 peptides have been designed so that charges are well distributed
 all over the vector. They have been selected for their ability to deliver an anti-GAPDH
 siRNA into HeLa or HS-68 cultured cells, with an efficiency sufficient to obtain at least a
 10 50% decrease of GAPDH protein level after 24 hrs of incubation with 50nM siRNA. The
 efficiency of expression inhibition was determined using CADY-1 (SEQ ID No: 2) as a
 reference, since this peptide enables a 95% inhibition.

An important parameter for designing CADY-2 peptides is the presence
 of tryptophane residues, and their repartition in the peptide. The CPPs have been selected
 15 using the p1p27 peptide (described below) as test cargo. The delivery efficiency was
 functionally determined by measuring the G1 arrest of synchronized HeLa or HS68 cells,
 after 24 hrs of incubation in the presence of 200 nM p1p27 peptide. The CADY-2
 reference peptide is CADY-2 (SEQ ID No: 8), which leads to 75% of cells arrested in G1.

CADY-1 vectors for siRNA delivery

- 20 CADY-1 : GLWRALWRLLRSLWRLWKA- Cya (SEQ ID No: 2)
 CADY-1b : GLWRALWRALWRSLWKLKRKV- Cya (SEQ ID No: 3)
 CADY-1c : GLWRALWRALRSLWKLKRKV- Cya (SEQ ID No: 4)
 CADY-1d : GLWRALWRGLRSLWKLKRKV- Cya (SEQ ID No: 5)
 CADY-1e : GLWRALWRGLRSLWKKKRKV- Cya (SEQ ID No: 6)
 25 CADY-1S : Cholesterol-GLWRALWRLLRSLWRLWKA (SEQ ID No: 7)
 CADY-1S1 : Cholesterol-CA β -GLWRALWRLLRSLWRLWKA (SEQ ID No: 13)
 CADY-1S2 : GLWRALWRLLRSLWRLWKA-Cholesterol (SEQ ID No: 7)

CADY-2 vector for peptide/protein delivery

- CADY-2 : GLWRALWRALWRSLWKLKWKV- Cya (SEQ ID No: 8)
 30 CADY-2b : GLWRALWRALWRSLWKSQRKV- Cya (SEQ ID No: 9)
 CADY-2c : GLWRALWRALWRSLWKKKRKV- Cya (SEQ ID No: 10)
 CADY-2d/1b : GLWRALWRALWRSLWKLKRKV- Cya (SEQ ID No: 3)
 CADY-2e : GLWRALWRLLRSLWRLWSQ-PKKKRKV- Cya (SEQ ID No: 12)

The N-terminal extremity of these peptides is linked to an acetyl group
 35 when no further indication appears. It can also be linked to a cholesterol molecule (when
 indicated), which is either linked directly to the N-terminus, or linked to the vector through
 a CA β bridge, which changes the relative orientation of the cholesterol and the core of the

vector (wherein the "core of the vector" designates the peptidic part of the vector of SEQ ID No:1).

The C-terminal extremity is linked either to a cholesterol molecule (CADY-S2) or to a cysteamide group (*i.e.*, CO-NH-CH₂-CH₂-SH).

5 Peptide synthesis was carried out at a 0.2 mmol scale, starting from a Fmoc-PAL-PEG-PS resin, using a Fmoc-continuous flow peptide synthesis instrument (PioneerTM, Applied Biosystems, Foster City, CA). The coupling reactions were performed with 0.5 M of HATU (2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) in the presence of 1 M of DIEA (diisopropylethylamine).
10 Recoupling was carried out at positions: 6, 7, 10, 11, 13, 14, 17 and 18. Protecting group removal and final cleavage from the resin were carried out with TFA (trifluoroacetic acid) /EDT (ethanedithiol) /thioanisole/phenol/H₂O (82.5/2.5/5/5/5 %) during 5h30min. The reaction mixture was then filtered through a cotton filter in cold diethyl ether. Since there occurred no peptide precipitation, ether was removed under rotary evaporation to obtain a
15 yellow oil. Neat ether was added to the oil and stored at -20°C overnight to allow the peptide to precipitate. A certain quantity of peptide has already precipitated in the cleavage cocktail (it is unusual for a peptide to precipitate in a near pure solution of TFA). This precipitate (+ the cotton and the resin) was resuspended in CH₃CN (containing 0.08% TFA) and H₂O. After filtration and thorough washes by CH₃CN/TFA and H₂O, the filtrate
20 was lyophilized prior to purification.

The crude peptide was purified by RP-HPLC on a C18 column (Interchrom UP5 WOD/25M Uptisperre 300 5 ODB, 250x21.2mm). Electrospray ionization mass spectrum was in complete agreement with the proposed structure.

Example 2: Structure of CADY when interacting with a membrane

25 The structure of CADY peptides was investigated by spectroscopy. It was demonstrated by circular dichroism that CADY adopts a helical structure in water or in phosphate solution. CADY forms a secondary amphipathic organization. Helical structure of CADY results in a hydrophobic and a hydrophilic faces. Molecular dynamic analysis reveals that CADY forms multimeric structures, involving at least 4 CADY
30 molecules within the membrane (figures 1 and 2). FTIR measurement of transferred CADY/membrane have confirmed that CADY peptides form an helical structure in the membrane.

Example 3: CADY forms stable particles with cargoes through non covalent interactions

35 The inventors have investigated the formation of CADY/cargo complexes and demonstrated that CADY peptides are able to form stable complexes with their cargoes. CADY-1 interact mainly with siRNA (Fig 3A) and CADY-2 mainly with

peptides and proteins (Fig 3B). Binding of cargoes to CADY was monitored by steady state fluorescence spectroscopy, using the intrinsic fluorescence associated to the Trp residue of CADY. For each CADY peptide, a fixed concentration of CADY was titrated by increasing concentration of cargoes. The binding of the cargo induced a marked quenching of fluorescence of both CADY peptides, and the dissociation constants were calculated from data fitting, using a quadratic equation which allows the determination of the ratio CADY/cargo. Dissociation constants and ratios were estimated to 2 nM for CADY-1/siRNA and 19 nM for CADY2/peptide, with a ratio at saturation of 1/10 and 1/20, respectively, whatever the variant of CADY-1 or CADY-2. However, cholesterol modification (*i.e.*, addition) reduced 2 to 10 folds the affinity of CADY-1 for the siRNA. These results suggest that in both cases, more than one peptide interact with the cargo, and that both CADY-1 and CADY-2 are able to form particles with their respective cargoes.

Example 4: CADY forms stable and non-toxic particles

The toxicity of CADY peptides has been investigated on different cell lines. Cells were incubated in the presence of increasing concentrations of CADY (from 0.1 μ M to 1 mM) for 12 hrs and cell viability was estimated using MTT assay. Data reported in Figure 4A demonstrated that CADY-1 is not toxic on the tested cell lines, up to a concentration of 100 μ M, with a EC₅₀ of 0.5 mM.

The different CADY variants have been tested on HS-68 and Hela cells. The results are reported in the Table 1 below.

Peptides	free EC50 (mM)	complexed EC50 (mM)
CADY-1	0.4	>2
CADY-1b	0.5	>2
CADY-1c	0.45	>2
CADY-1d	0.5	>2
CADY-1e	0.34	1.5
CADY-S	0.2	1
CADY-S2	0.25	0.8
CADY-2	0.23	1
CADY-2b	0.28	0.9
CADY-2c	0.34	0.8
CADY-2d	0.38	1.5
CADY-2e	0.43	>2
KALA	0.07	
GALA	0.05	
TAT	0.1	
Lipofectamine	0.2	

Table 1

The inventors demonstrated that association with cargo molecule (ratio 1/20) significantly reduced the toxicity of the carrier, an average EC₅₀ of 2 mM was calculated. The lack of toxicity of CADY/cargo complexes can be directly associated to absence of free carrier peptide in the solution (Fig. 4B). Lipofectamine and other

membranolytic CPP (KALA, GALA, TAT), which exhibit more than 70 % toxicity at concentration of 100 μ M, were used as controls.

Example 5: CADY-mediated delivery of siRNA

The ability of CADY-1 and variants to deliver siRNA was evaluated on both adherent and suspension cell lines, using GAPDH as the target of siRNA.

siRNA targeting the 3'UTR of GAPDH were from the SilencerTM GAPDH siRNA kit (Ambion). Fluorescent labeling of siRNA was performed using either the Fam or the Cy3 SilencerTM labeling kit (Ambion) and modified as described in the manufacturer's protocol. pRL-Luc reporter gene was from Promega and siRNA targeting luciferase sense 5'-CUUACGCUGAGUACUUCGATT-3' (SEQ ID No: 14) and antisense 3'-TTGAAUGCGACUCAUGAAGCU-5' (SEQ ID No: 15) and mismatch sense 5'-CGUACGCGGAAUACUUCGATT-3' (SEQ ID No: 16) and antisense 3'-TTGCAUGCGCCUUAUGAAGCU-5' (SEQ ID No: 17) siRNA were obtained from Genset Oligos.

The results, presented in figure 5, show that CADY-1 is more efficient than the lipofectamine control. The percentage of GAPDH knockdown, estimated by comparison with a control essays using a mismatch siRNA delivered with CADY at the same molecular ratio, is also reported in Table 2 below. Data are the average of 4 separated experiments.

Cell lines	Knockdown
Hela	85%
Jurkat	80%
HepG2	70%
C2C12	70%
MEF	70%
HS-68	80%
CEM-SS	80%
U2OS	75%
MCF7	70%
MT4	65%
HER2	60%
MDA-MB	75%
Balb/c3T3	80%

Table 2

Example 6: Mechanism of cellular uptake of CADY-1

Whatever the CPP, the common major concern in the development of new PTD strategy is to avoid any endosomal pathway and/or to facilitate the escape of the cargoes from the early endosomes in order to limit their degradation. The cellular uptake mechanism of CADY-1 was therefore investigated in detail. The inventors demonstrated that, in contrast to most of the delivery systems described so far for siRNA, CADY-mediated delivery of active siRNA is independent of the endosomal pathway and directly correlated to the size of the particle CADY/siRNA.

Indeed, results reported in Fig 6A revealed that efficiency of CADY is dependent on the size of the particle formed with siRNA. Optimal biological response was obtained for a CADY/siRNA ratio of 1:20 to 1:25, which corresponds to a nano particle (size of 100 - 200 nm). Therefore, calibration of the formulation is an essential parameter in the efficiency of CADY-1.

Results reported in Fig 6B demonstrated that none of the tested inhibitors affected the cellular uptake of CADY and the associated siRNA biological response. In contrast to other CPP, the mechanism of cellular uptake of CADY/siRNA particle is independent of the endosomal pathway. However, the size of the particle and CADY/siRNA molar ratio need to be controlled, so that the diameter of the particles remains below 200 nm and the CADY/siRNA molar ratio remains under 1:25. In contrast, for particles of diameter >500 nm, the mechanism is partially associated to the endosomal pathway.

These data demonstrate that calibration of the formulation is an essential parameter in the efficiency of CADY-1, and that the optimal formulation for CADY/siRNA is: ratio 1:20 to 1:25 and nano particle (size of 100 - 200 nm).

Example 7: CADY-mediated *in vitro* delivery of siRNA targeting Cyclin-B1

Cell cycle progression is driven by sequential activation of essential heterodimeric protein kinase complexes (Cdk/cyclin complexes). Most of the drugs currently designed to target cell cycle progression are directed against the kinase activity of the Cdk subunits. However, as such drugs generally tend to affect other cellular kinases non specifically, the inventors have decided to target the regulatory cyclin subunit instead, as a more appropriate means of improving both the selectivity and the efficiency of cell cycle inhibitors. In particular, cyclin B1 constitutes a key target for cancer therapy, as a component of the essential "Mitosis Promoting Factor", together with protein kinase Cdk1.

A siRNA molecule was hence designed to target cyclin B1. The following siRNA sequence was selected out of 10 different sequences, and the oligonucleotides were obtained from proligo:

Si-Cyclin B sense: 5'-GGCGAAGAUAACAUGGCATT-3' (SEQ ID No: 18)

Si-Cyclin B antisense : 5'-UGCCAUGUUGAUCUUCGCCTT-3' (SEQ ID No: 19)

Mismatch sense: 5'-GGUGAAGAUCAGCAUGGCATT-3' (SEQ ID No: 20)

Mismatch antisense: 5'-UGCCAUGUCGAUCUUCACCTT3' (SEQ ID No: 21)

The potency of this siRNA was investigated using CADY as delivery system. The biological response of CADY-mediated delivery of siRNA targeting cyclin B1 was examined. To this aim, the inventors first assessed whether cyclin B1 protein levels were downregulated by CADY-mediated delivery of siRNA, and to what extent, compared to delivery of siRNA using classical lipid formulation.

As shown in Figure 7A, delivery of siRNA cyclin B1 is associated to a cell cycle arrest in G2, characteristic of a knock down of the cyclin B1 protein. Western blot analysis confirmed that the observed G2-arrest is directly associated to a dramatic knock down of the cyclin B1 protein level (Fig. 7B). Interestingly, the G2 arrest is observed at low concentration of siRNA of 2 nM. This demonstrates the high potency of CADY for siRNA delivery and rapid release into the cytoplasm. Using CADY-mediated delivery can hence enable a significant response with only low concentrations of siRNA, which will limit problems related to unspecific targeting and toxicity.

Example 8: Anti-cyclin-B1 siRNA/CADY inhibits proliferation of cancer cells

The potency of CADY/siRNA complex to block cancer cell proliferation was then investigated.

As shown on figure 8, the anti-cyclin B1 siRNA significantly blocks cancer cell proliferation when associated to CADY. 40 % and 72% of inhibition were obtained with siRNA concentrations of 0.1 nM and 1 nM, respectively, associated to CADY-1, whereas the same concentrations were not efficient when lipofectamine was used as vector. These results are in perfect agreement with those obtained on other cell lines and confirm the high potency of CADY and the possibility of using very low concentration (sub nanomolar) of siRNA.

These data are the first demonstration that siRNA can work at concentrations lower than 1 nM. Such low concentrations of siRNA induced a strong biological response, suggesting that CADY-mediated delivery is highly efficient. This is certainly correlated with the rapid release of the siRNA in the cytoplasm, independently of the endosomal pathway.

Example 9: Anti-cyclin-B1 siRNA/CADY inhibits tumor growth: intratumoral injection

The effect of the anti-cyclin B1 siRNA/CADY complexes was then investigated on a tumor animal model, to determine to what extent the results obtained on the proliferation of cancer cells could be extrapolated to the inhibition of tumor growth *in vivo*.

The results are shown on figure 9. In a control experiment, the inventors observed that 20 days after injection of PBS, CADY or naked siRNA, the tumor size increased by 2,5 fold. In contrast, no growth of the tumor was observed, when injected with CADY/siRNA even with low concentration of siRNA (1 µg).

These results suggested that siRNA specifically inhibits tumor growth and, thanks to the used of CADY as carrier, only a low concentration of siRNA is required for a marked anti-tumoral effect. Moreover, a significant reduction in the tumor size was

observed after a second injection of CADY/siRNA. Western blot analysis showed that the level of cyclin B1 was dramatically reduced.

These results demonstrate that CADY constitutes an excellent tool for *in vivo* delivery of siRNA.

5 **Example 10: Cholesterol modified -CADY-formulation for *in vivo* administration**

The above results demonstrate that CADY-1 constitutes an excellent tool for *in vivo* application of siRNA. However, although CADY-1/siRNA formulation is highly efficient upon intra-tumoral injection, the observed response to siRNA was very limited when the complexes were injected intravenously or via the intra-peritoneal route.

10 A new variant of CADY was hence designed to obtain nano-particles including siRNA, which are stable *in vivo* and administrable through intravenous injections. In that purpose, a cholesterol group was added at the N- terminus (CADY-S) or at the C-terminus (CADY-S2) of CADY. Cholesterol linking at the C-terminus was performed through a disulfide bond with the cysteamide group. Linking at the N-terminus
15 was performed through the activation of the amino group and formation of a disulfide bond.

The ability of both CADY-S and CADY-S2 to deliver siRNA was evaluated using anti-cyclin B1 siRNA. In order to identify the optimal formulation, different molar ratios of CADY were used in complex with 50 nM siRNA. Formulations
20 were ranked based on the G2 arrest associated to knock down of cyclin B1.

The results, presented in figure 10, show that the presence of cholesterol attached to CADY does not affect its efficiency to deliver siRNA in cultured cells. The optimal formulation is similar to that obtained with CADY. The optimal size for the particle is of about 200 nm diameter.

25 The *in vivo* efficiency of CADY-S was then investigated, in order to determine if the cholesterol-modified form of CADY can improve siRNA delivery through intravenous injection and block tumor growth *in vivo*. Experiments were performed on two tumor mouse models, in which 5PC3 or Human H1299 adenocarcinoma cells were injected subcutaneously into the flanks of NCR nude mice.

30 The results of both experiments demonstrate that cholesterol-modified CADY was able to deliver siRNA *in vivo*, and induced a significant biological response by blocking tumor growth (figure 11). The presence of cholesterol improves the stability of the particle after intravenous injection. Moreover, only one injection at low concentration of siRNA (10 µg) of siRNA was required to fully abolish the tumor growth.

35 These results constitute the first *in vivo* therapeutic application of a carrier system/siRNA complex using physiological doses. CADY hence appears as the best vector for systematic *in vivo* delivery of siRNA, since other potent methods (such as

cationic polymer/cholesterol-modified siRNA) required an everyday IV injection of siRNA, at concentrations 10 fold higher than those used in the above-described experiments, to obtain a similar efficiency.

Example 11: CADY-mediated delivery of therapeutic peptides

5 Protein transduction technology described so far requires the attachment of the Protein Transduction Domain (PTD) to the target peptide or protein, which can be achieved either by chemical cross-linking or by cloning and expression of a fusion protein. Conjugate method offers several advantages for rationalization and control of the stoichiometry, as well as for reproducibility and calibration of the PTD-cargo for *in vivo* application. However, this approach is limited from a technical point of view and by a risk of alteration of the biological activity of the cargoes.

CADY vectors, especially CADY-2 vectors, offer an alternative to the covalent PTDs technology, for the delivery of full-length proteins and peptides into mammalian cells.

15 In higher eukaryotes, the cell cycle progression is coordinated by several closely related Ser/Thr protein kinases, resulting from association of a catalytic Cyclin-dependent kinase (CDK) subunit with a regulatory cyclin subunit. CDK/cyclin complexes are regulated by both phosphorylation/dephosphorylation and protein-protein interactions with different partners, including structural CDK inhibitors (CKI). Genetic evidence supports a strong correlation between alterations in the regulation of CDKs and the molecular pathology of cancer. Thus, in the past few years, there has been considerable interest in the development of inhibitors of CDK protein kinases. A greater understanding of the cell cycle and of the inherent complexity of CDK regulation now offers a number of possible routes to their inhibition (Swanton 2004). One of the currently most developed strategies to inhibit CDKs is based on the design of small molecules that target the ATP-binding site, thereby interfering directly with their catalytic activity.

25 The tumor suppressor protein p27 is essential in the control of the G1/S transition. Alteration in p27 regulation results in aberrations in cell cycle progression and development of tumors. p27 protein interacts with the Cdk/cyclin complex in order to maintain the complex in an inactive conformation which controls the rate of the G1 phase. The molecular mechanism by which p27 induces cell cycle arrest was investigated in details, and peptide motifs which block both cell cycle in G1 and cancer cell proliferation have been identified.

30 Based on the X-ray structure of the cdk2/cyclinA/ p27 complex (Jeffrey, Russo et al. 1995), the inventors have selected three peptides:

- p1p27 : RVSNGSPSLERMDARQAEHPKPSACRNL (SEQ ID No: 22)
- pRXL : PS�ERMDAR (SEQ ID No: 23), and

- p2p27 : NRTEENVSDGSPNAGSVEQTPKKPGLR (SEQ ID No: 24) as a negative control.

Peptides were synthesized and purified as described previously (Mery, Granier et al. 1993; Morris, Vidal et al. 1997), then their ability to alter the cell cycle progression was investigated.

As shown on figure 12, the delivery of the peptides p1p27 or pRXL induced a cell cycle arrest in G1 in all tested cell lines. In contrast, no effect was observed with CADY or free p1p27, which suggested that CADY dramatically improves the cellular uptake of the peptide, without any toxicity associated to the carrier. Moreover, the fact that no effect was observed with the control peptide p2p27 demonstrates the high specificity of the peptide. In comparison to results previously published with pRXL using a TAT or Pen covalent based strategy (Nagahara, Vocero-Akbani et al. 1998; Chen, Sharma et al. 1999; Snyder, Meade et al. 2003), the efficiency with CADY strategy is at least 2-fold better.

Example 12: CADY2-mediated *in vivo* delivery of p27 inhibits tumor growth

The inventors then investigated the ability of the p1p27/CADY complexes to block tumor growth *in vivo*.

The results are shown on figure 13.

In control experiments, in which PBS, CADY alone or naked p1p27 were injected, the tumor size had increased by 5-fold, 5 weeks after injection. In contrast, no growth of the tumor was observed, when injected with CADY/p1p27. These results suggest that p1p27 specifically inhibits tumor growth, and that the use of CADY as carrier enables to administer only a low amount of p1p27 for a marked anti-tumoral effect. Moreover, a significant reduction in the tumor size was observed after a second injection of CADY/p1p27. These results show that CADY constitutes an excellent tool for *in vivo* application of therapeutic peptides.

Other experiments, performed with peptides having an anti-viral or an antitumoral activity, as described in (Morris, Robert-Hebmann et al. 1999) or in (Gondeau, Gerbal-Chaloin et al. 2005), demonstrated that CADY-2 peptides improve the delivery of several distinct proteins and peptides into different cell lines in a fully biologically active form, without the need for prior chemical covalent coupling or denaturation step. CADY technology hence constitutes a powerful tool for basic research, and demonstrated to be extremely powerful for studying the role of proteins, and for targeting specific protein/protein interactions *in vitro* as well as *in vivo*.

REFERENCES

- Allen, T. M. (2002). "Ligand-targeted therapeutics in anticancer therapy." Nat Rev Cancer **2**(10): 750-63.
- Chen, Y. N., S. K. Sharma, et al. (1999). "Selective killing of transformed cells by cyclin/cyclin-dependent kinase 2 antagonists." Proc Natl Acad Sci U S A **96**(8): 4325-9.
- 5 Deshayes, S., M. C. Morris, et al. (2005). "Cell-penetrating peptides: tools for intracellular delivery of therapeutics." Cell Mol Life Sci **62**(16): 1839-49.
- El-Andaloussi, S., T. Holm, et al. (2005). "Cell-penetrating peptides: mechanisms and applications." Curr Pharm Des **11**(28): 3597-611.
- 10 Elliott, G. and P. O'Hare (1997). "Intercellular trafficking and protein delivery by a herpesvirus structural protein." Cell **88**(2): 223-33.
- Gondeau, C., S. Gerbal-Chaloin, et al. (2005). "Design of a novel class of peptide inhibitors of cyclin-dependent kinase/cyclin activation." J Biol Chem **280**(14): 13793-800.
- 15 Gratton, J. P., J. Yu, et al. (2003). "Cell-permeable peptides improve cellular uptake and therapeutic gene delivery of replication-deficient viruses in cells and in vivo." Nat Med **9**(3): 357-62.
- Jarver, P. and U. Langel (2004). "The use of cell-penetrating peptides as a tool for gene regulation." Drug Discov Today **9**(9): 395-402.
- 20 Jeffrey, P. D., A. A. Russo, et al. (1995). "Mechanism of CDK activation revealed by the structure of a cyclinA-CDK2 complex." Nature **376**(6538): 313-20.
- Joliot, A. and A. Prochiantz (2004). "Transduction peptides: from technology to physiology." Nat Cell Biol **6**(3): 189-96.
- 25 Mery, J., C. Granier, et al. (1993). "Disulfide linkage to polyacrylic resin for automated Fmoc peptide synthesis. Immunochemical applications of peptide resins and mercaptoamide peptides." Int J Pept Protein Res **42**(1): 44-52.
- Morris, M. C., L. Chaloin, et al. (2000). "Translocating peptides and proteins and their use for gene delivery." Curr Opin Biotechnol **11**(5): 461-6.
- 30 Morris, M. C., L. Chaloin, et al. (2002). "Signal sequence based Cell Penetrating Peptides and their applications for gene delivery." Cell Penetrating peptides: Processes and application.
- Morris, M. C., V. Robert-Hebmann, et al. (1999). "A new potent HIV-1 reverse transcriptase inhibitor. A synthetic peptide derived from the interface subunit domains." J Biol Chem **274**(35): 24941-6.
- 35 Morris, M. C., P. Vidal, et al. (1997). "A new peptide vector for efficient delivery of oligonucleotides into mammalian cells." Nucleic Acids Res **25**(14): 2730-6.

- Nagahara, H., A. M. Vocero-Akbani, et al. (1998). "Transduction of full-length TAT fusion proteins into mammalian cells: TAT-p27Kip1 induces cell migration." Nat Med **4**(12): 1449-52.
- Pooga, M., M. Hallbrink, et al. (1998). "Cell penetration by transportan." Faseb J **12**(1):
5 67-77.
- Schwarze, S. R., A. Ho, et al. (1999). "In vivo protein transduction: delivery of a biologically active protein into the mouse." Science **285**(5433): 1569-72.
- Snyder, E. L., B. R. Meade, et al. (2003). "Anti-cancer protein transduction strategies: reconstitution of p27 tumor suppressor function." J Control Release **91**(1-2): 45-51.
- 10 Swanton, C. (2004). "Cell-cycle targeted therapies." Lancet Oncol **5**(1): 27-36.
- Wadia, J. S. and S. F. Dowdy (2002). "Protein transduction technology." Curr Opin Biotechnol **13**(1): 52-6.
- Wadia, J. S., R. V. Stan, et al. (2004). "Transducible TAT-HA fusogenic peptide enhances escape of TAT-fusion proteins after lipid raft macropinocytosis." Nat Med **10**(3):
15 310-5.

CLAIMS:

1. A cell-penetrating peptide comprising an amino acid sequence consisting of GLX₉RALX₉RX₁LX₂RSLX₉X₃X₄X₅X₆X₇X₈ (SEQ ID No: 1), wherein X₁ is selected from the group consisting of A, L and G; X₂ is W or none; X₃ is R or K; X₄ is selected from the group consisting of K, L and S; X₅ is L or K; X₆ is R or W; X₇ is K or S; X₈ is selected from the group consisting of A, V and Q; and X₉ is selected from the group consisting of W, F and Y.
2. The cell-penetrating peptide of claim 1, which comprises an amino acid sequence consisting of GLWRALWRX₁LX₂RSLWX₃X₄X₅X₆X₇X₈ (SEQ ID No: 25), wherein X₁ is A, L or G; X₂ is W or none; X₃ is R or K; X₄ is K, L or S; X₅ is L or K; X₆ is R or W; X₇ is K or S; and X₈ is A, V or Q.
3. The cell-penetrating peptide of claim 1 or 2, further comprising, covalently linked to the C-terminal end of said amino acid sequence, one or several groups selected from the group consisting of a cysteamide, a cysteine, a thiol, an amide, a carboxyl, a linear or ramified C₁-C₆ alkyl optionally substituted, a primary or secondary amine, an osidic derivative, a lipid, a phospholipid, a fatty acid, a cholesterol, a poly-ethylene glycol, a nuclear localization signal and a targeting molecule.
4. The cell-penetrating peptide of any one of claims 1 to 3, further comprising, covalently linked to the N-terminal end of said amino acid sequence, one or several chemical entities selected from the group consisting of an acetyl, a fatty acid, a cholesterol, a poly-ethylene glycol, a nuclear localization signal and a targeting molecule.
5. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is selected from the group consisting of
 - GLWRALWRLLRSLWRLWKA (SEQ ID No: 2);
 - GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3);

- GLWRALWRALRSLWKLKRKV (SEQ ID No: 4);
- GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5);
- GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6);
- GLWRALWRLLRSLWRLWKA (SEQ ID No: 7);
- GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8);
- GLWRALWRALWRSLWKSRRKV (SEQ ID No: 9);
- GLWRALWRALWRSLWKKKRKV (SEQ ID No: 10); and
- GLWRALWRLLRSLWRLWSQ (SEQ ID No: 11).

6. The cell-penetrating peptide of any one of claims 1 to 4, which is selected from the group consisting of CADY-1 (Ac-GLWRALWRLLRSLWRLWKA-Cya) and CADY-2 (Ac-GLWRALWRALWRSLWKLKWKV-Cya).

7. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRLLRSLWRLWKA (SEQ ID No: 2).

8. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3).

9. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRALRSLWKLKRKV (SEQ ID No: 4).

10. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5).

11. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6).

12. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRLLRSLWRLWKA (SEQ ID No: 7).

13. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8).
14. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRALWRSLWKSRRKV (SEQ ID No: 9).
15. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRALWRSLWKKRRKV (SEQ ID No: 10).
16. The cell-penetrating peptide of any one of claims 1 to 4, wherein the amino acid sequence is GLWRALWRLLRSLWRLLWSQ (SEQ ID No: 11).
17. A complex comprising a cell-penetrating peptide according to any one of claims 1 to 6 and a cargo selected from the group consisting of nucleic acids, peptides, proteins, contrast agents and toxins.
18. The complex of claim 17, wherein said cargo is a siRNA selected to silence a target mRNA.
19. The complex of claim 18, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRX₁LX₂RSLWX₃X₄X₅KX₆ (SEQ ID No: 26), wherein X₁ is selected from the group consisting of A, L and G, X₂ is W or none, X₃ is R or K, X₄ is K or L, X₅ is R or W, X₆ is A or V.
20. The complex of claim 18 or 19, wherein the amino acid sequence of the cell-penetrating peptide is selected from the group consisting of
- GLWRALWRLLRSLWRLLWKA (SEQ ID No: 2);
 - GLWRALWRALWRSLWKLKRRKV (SEQ ID No: 3);
 - GLWRALWRALRSLWKLKRRKV (SEQ ID No: 4);

- GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5);
- GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6); and
- GLWRALWRLLRSLWRLWKA (SEQ ID No: 7).

21. The complex of claim 17, wherein said cargo is a peptide or a protein.

22. The complex of claim 21, wherein the amino acid sequence of the cell-penetrating peptide is SEQ ID No: 1, in which X_1 is A or L and X_8 is V or Q.

23. The complex of claim 21 or 22, wherein the amino acid sequence of the cell-penetrating peptide is selected from the group consisting of

- GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8);
- GLWRALWRALWRSLWKSRRKV (SEQ ID No: 9);
- GLWRALWRALWRSLWKKKRKV (SEQ ID No: 10);
- GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3);
- GLWRALWRLLRSLWRLWSQ (SEQ ID No: 11); and
- GLWRALWRLLRSLWRLWSQPKKKRKV (SEQ ID No: 12).

24. The complex of any one of claims 17 to 23, wherein the cell-penetrating peptide comprises an acetyl group covalently linked to its N-terminus, and/or a cysteamide group covalently linked to its C-terminus.

25. The complex of any one of claims 17 to 24, wherein the cell-penetrating peptide further comprises a cholesterol molecule, covalently linked to its C-terminus or its N-terminus.

26. The complex of any one of claims 17 to 25, wherein the size of the complex is between 50 and 300 nm.

27. The complex of claim 26, wherein the size of the complex is between 100 and 200 nm.
28. The complex of any one of claims 17 to 27, wherein at least part of the cell-penetrating peptides are bound to a targeting molecule.
29. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRLLRSLWRLWKA (SEQ ID No: 2).
30. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRALWRSLWKLKRKV (SEQ ID No: 3).
31. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRALRSLWKLKRKV (SEQ ID No: 4).
32. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRGLRSLWKLKRKV (SEQ ID No: 5).
33. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRGLRSLWKKKRKV (SEQ ID No: 6).
34. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRLLRSLWRLWKA (SEQ ID No: 7).
35. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRALWRSLWKLKWKV (SEQ ID No: 8).
36. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRALWRSLWKSRRKV (SEQ ID No: 9).
37. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRALWRSLWKKKRKV (SEQ ID No: 10).

38. The complex of claim 17, wherein the amino acid sequence of the cell-penetrating peptide is GLWRALWRLLRSLWRLWSQ (SEQ ID No: 11).
39. A composition comprising the complex according to any one of claims 17 to 38 and a diluent.
40. A composition for use in anticancer therapy, comprising the complex according to any one of claims 17 to 38 and a diluent.
41. A method for delivering a molecule into a cell *in vitro*, comprising a step of putting said cell into contact with a complex comprising said molecule and the cell-penetrating peptide according to any one of claims 1 to 16.
42. A use of the cell-penetrating peptide according to any one of claims 1 to 16, or of the complex according to any one of claims 17 to 38, for the preparation of a medicament for use in anticancer therapy.
43. The use of claim 42, wherein said medicament comprises anti-cyclin B1 siRNA/CADY complexes and a diluent.
44. The use of claim 42, wherein said medicament comprises p1p27/CADY complexes and/or pRXL/CADY complexes and a diluent.
45. A use of the cell-penetrating peptide according to any one of claims 1 to 16, or of the complex according to any one of claims 17 to 38 for treating or preventing cancer.
46. A commercial package comprising the cell-penetrating peptide according to any one of claims 1 to 16, or of the complex according to any one of claims 17 to 38 and instructions for use in treating or preventing cancer.

47. A use of the cell-penetrating peptide according to any one of claims 1 to 16 for delivering a nucleic acid, a peptide, a protein, a contrast agent, or a toxin to a cell.




Figure 1

A



B





Figure 2

2 / 9

A



B





Figure 3

3 / 9

A



B




Figure 4

4 / 9




Figure 5

5 / 9




Figure 6

6 / 9




Figure 7




Figure 8

7 / 9




Figure 9





Figure 10

8 / 9



H1299 cells




Figure 11

9 / 9

