

COMMONWEALTH OF AUSTRALIA

Patents Act 1952 - 1969

DECLARATION IN SUPPORT OF A CONVENTION APPLICATION FOR A PATENT

In support of the Convention Application made by CIBA-GEIGY AG for a patent for an invention entitled:

Pesticides

I, Werner Waldegg of CIBA-GEIGY AG, Klybeckstrasse 141, 4002 Basle, Switzerland do solemnly and sincerely declare as follows:

1. I am authorised by the applicant for the patent to make this declaration on its behalf.
2. The basic application~~(s)~~ as defined by Section 141 on the Act was ~~(were)~~ made in Switzerland
on May 17, 1990

by CIBA-GEIGY AG, 4002 Basle, Switzerland


3. Adolf Hubele, Obere Egg 9, 4312 Magden, Switzerland

is ~~(are)~~ the actual inventor~~(s)~~ of the invention and the facts upon which the applicant is entitled to make the application are as follows: The said applicant is the assignee of the actual inventor~~(s)~~.

4. The basic application~~(s)~~ referred to in paragraph 2 of this Declaration was ~~(were)~~ the first application~~(s)~~ made in a Convention country in respect of the invention the subject of the application.

DECLARED at Basle, Switzerland on April 17, 1991

CIBA-GEIGY AG


Werner Waldegg
Single Signature, by special power

To: The Commissioner of Patents

S & F Ref: 161812

FORM 10

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

COMPLETE SPECIFICATION

(ORIGINAL)

FOR OFFICE USE:

Class Int Class

Complete Specification Lodged:
Accepted:
Published:

Priority:

Related Art:

Name and Address
of Applicant: Ciba-Geigy AG
 Klybeckstrasse 141
 4002 Basel
 SWITZERLAND


Actual Inventor(s): Adolf Hubele

Address for Service: Spruson & Ferguson, Patent Attorneys
 Level 33 St Martins Tower, 31 Market Street
 Sydney, New South Wales, 2000, Australia

Complete Specification for the invention entitled:

2-anilino-4-cyanopyrimidine derivatives
Pesticides


The following statement is a full description of this invention, including the best method of performing it known to me/us



2-anilino-4-cyanopyrimidine derivatives
pesticides

Abstract

Compounds of formula



wherein R₁ is hydrogen ~~3-fluorine or 4-fluorine~~ and R₂ is C₁-C₄alkyl, halo- or hydroxy-substituted C₁-C₂alkyl, C₃-C₆-cycloalkyl or C₃-C₆cycloalkyl mono- to tri-substituted by identical or different substituents selected from methyl and halogen, have valuable microbicidal properties. The novel compounds can be used in crop protection for preventing an attack on cultivated plants by phytopathogenic microorganisms and for controlling those pests.




PF/5-18084/A

Pesticides

The present invention relates to novel 2-anilinopyrimidine derivatives of formula I below. It relates also to the preparation of those compounds and to agrochemical compositions that comprise at least one of those compounds as active ingredient. The invention relates also to the preparation of the said compositions and to the use of the compounds or compositions for controlling pests, especially harmful insects and plant-destructive microorganisms, especially fungi.

The compounds according to the invention have the general formula I



wherein R_1 is hydrogen ~~3-fluorine or 4-fluorine~~ and R_2 is C_1 - C_4 alkyl, halo- or hydroxy-substituted C_1 - C_2 alkyl, C_3 - C_6 -cycloalkyl or C_3 - C_6 cycloalkyl mono- to tri-substituted by identical or different substituents selected from methyl and halogen; including their acid addition salts and metal salt complexes.

Depending on the number of carbon atoms indicated, alkyl is to be understood as being, for example, methyl, ethyl, propyl, butyl and their isomers, for example isopropyl, isobutyl, tert-butyl or sec-butyl. Halogen is fluorine,

chlorine, bromine or iodine, preferably chlorine. Depending on the number of carbon atoms indicated, cycloalkyl is, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

The invention relates both to the free compounds of formula I and to their addition salts with inorganic and organic acids as well as to their complexes with metal salts.

Salts according to the invention are especially addition salts with physiologically tolerable inorganic or organic acids, for example hydrohalic acids, for example hydrochloric, hydrobromic or hydriodic acid, sulfuric acid, phosphoric acid, phosphorous acid or nitric acid, or organic acids such as acetic acid, trifluoroacetic acid, trichloroacetic acid, propionic acid, glycolic acid, thiocyanic acid, lactic acid, succinic acid, citric acid, benzoic acid, cinnamic acid, oxalic acid, formic acid, benzenesulfonic acid, p-toluenesulfonic acid, methanesulfonic acid, salicylic acid, p-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid or 1,2-naphthalenedisulfonic acid.

Metal salt complexes of formula I comprise the basic organic molecule and an inorganic or organic metal salt, for example the halides, nitrates, sulfates, phosphates, acetates, trifluoroacetates, trichloroacetates, propionates, tartrates, sulfonates, salicylates, benzoates, etc. of the elements of the second main group, such as calcium and magnesium, and of the third and fourth main groups, such as aluminium, tin or lead, and of the first to eighth subgroups, such as chromium, manganese, iron, cobalt, nickel, copper, zinc, etc..

Preference is given to the subgroup elements of the 4th period. The metals may be in the various valencies associated with them. The metal complexes may be mono- or polynuclear, that is to say they may contain one or more organic molecular components as ligands.

The compounds of formula I are oils, resins or solids that are stable at room temperature and are distinguished by valuable microbicidal properties. They can be used in the agricultural sector or related fields preventively and curatively for controlling plant-destructive microorganisms. The compounds of formula I according to the invention are distinguished at low rates of application not only by excellent fungicidal activity, but also by the fact that they are especially well tolerated by plants.

The following groups of compounds are preferred on account of their marked microbicidal, especially phytofungicidal, activity.

Group 1: Compounds of formula I wherein:

R₁ is hydrogen ~~3-fluorine or 4-fluorine~~; and

R₂ is C₁-C₃alkyl, cyclopropyl or cyclopropyl substituted by methyl or by chlorine.

Group 2: Compounds of formula I wherein:

R₁ is hydrogen; and

R₂ is methyl, ethyl, cyclopropyl or 2-methylcyclopropyl.

Group 3: Compounds of formula I wherein:

R₁ is hydrogen ~~3-fluorine or 4-fluorine~~; and

R₂ is methyl, ethyl or cyclopropyl.

For use as fungicidal active ingredients, the following compounds may be mentioned as being especially preferred:

2-phenylamino-4-cyano-6-methylpyrimidine (comp. no. 1.1),

2-phenylamino-4-cyano-6-ethylpyrimidine (comp. no. 1.3),

2-phenylamino-4-cyano-6-cyclopropylpyrimidine (comp.

no. 1.11) and


~~2-(3-fluorophenylamino)-4-cyano-6-methylpyrimidine (comp. no. 1.16).~~




N-pyrimidinylaniline compounds are already known. For example, in published European patent application 0 224 339 and DD patent specification 151 404, compounds having an N-2-pyrimidinyl structure are described as being effective against plant-destructive fungi. In addition, compounds having an anilinyrimidine structure substituted by cyano groups have become known (see European Patent Application No. 0 337 943). According to the description, however, those compounds are used only as intermediates for the synthesis of herbicidally active urea derivatives. The known N-pyrimidinylaniline derivatives have hitherto been unable to meet the requirements made of them for use as fungicides, especially at low rates of application.

The compounds of formula I are prepared by

1.1 reacting a guanidine salt of formula IIa



or a guanidine compound of formula IIb




with a ketone of formula III




wherein R_3 is an alkyl radical, preferably C_1 - C_4 alkyl, and A^\ominus is an acid anion, in a protic solvent or without a solvent,

at temperatures of from 40° to 160°C, preferably from 60° to 110°C, to form a pyrimidine compound of formula IV




and

1.2 hydrolysing the resulting acetal of formula IV in the presence of an acid, for example a hydrohalic acid or sulfuric acid, in water or aqueous mixtures of inert solvents, for example alcohols or dimethylformamide, at temperatures of from 20° to 100°C, preferably from 30° to 60°C, to form the pyrimidinylaldehyde of formula V




and

1.3 then converting the aldehyde of formula V with a hydroxylamine salt and a base that frees the hydroxylamine, in a protic solvent, preferably in the presence of water, at temperatures of from 10° to 90°C, preferably at from 10°C to 60°C, into the oxime of formula VI



and

1.3.1 subjecting the oxime of formula VI to a reaction in which the elements of water are removed; the oxime, for example, being converted in an intermediate step into a compound of formula VII




(VII),

wherein R' is an acyl radical COR'', a carbamoyl radical CONHR'' or an oxycarbonyl radical COOR'', in which R'' is an alkyl radical, preferably C₁-C₄alkyl, and the compound of formula VII then being decomposed by removal of the acyl radical HOCOR'' in inert solvents at temperatures of from 60° to 120°C, preferably from 60° to 100°C; or

1.4 from the pyrimidinealdehyde of formula V, preparing the oxime of formula VI in an intermediate step by reaction in tertiary bases, for example pyridine and hydroxylamine salt, and forming the compound of formula I directly therefrom in situ by treatment with an acid halide or acid anhydride, for example acetic anhydride, at temperatures of from 40° to 120°C, preferably from 60° to 100°C; or

2. reacting a compound of formula VIII



(VIII)

with a cyanide of formula IX




wherein Hal is fluorine, chlorine, bromine or iodine, preferably chlorine or iodine, and Me is an alkali metal cation, alkaline earth metal cation or heavy metal cation, preferably sodium, potassium or copper, in an aprotic solvent, for example dimethylformamide or dimethyl sulfoxide, at temperatures of from 60° to 160°C, preferably from 80° to 120°C; or


3.1 cyclising urea of formula X



with a diketone of formula III




wherein R₃ is an alkyl radical, preferably C₁-C₄alkyl, in the presence of an acid in an inert solvent at temperatures of from 20° to 140°C, preferably from 20° to 40°C, to form a pyrimidine compound of formula XI



and


3.2 replacing the OH group in the compound of formula XI by halogen using excess $\text{PO}(\text{Hal})_3$ in the presence or absence of an inert solvent at temperatures of from 50° to 110°C , preferably at the reflux temperature of the $\text{PO}(\text{Hal})_3$, to form a compound of formula XII



(XII),

wherein Hal is halogen, preferably chlorine or bromine, and then

3.3 reacting a compound of formula XII with an aniline compound of formula XIII




(XIII)

either

a) in the presence of a proton acceptor, for example in an excess of the aniline compound of formula XIII or of an inorganic base, with or without an inert solvent, or

b) in the presence of an acid in an inert solvent, in each case at temperatures of from 60° to 120°C , preferably from 80° to 100°C , to form a compound of formula IV



and converting the resulting acetal into a compound of formula I in the manner described under 1.2, 1.3 and 1.3.1, the radicals R_1 and R_2 in the above-described processes being as defined under formula I.

In the processes described, suitable salt radicals for the acid anion A^\ominus in the compounds of formula IIa are, for example: carbonate, hydrogen carbonate, nitrate, halide, sulfate and hydrogen sulfate.

Halide is to be understood as meaning fluoride, chloride, bromide or iodide, preferably bromide or chloride.

The acids used are especially inorganic acids, for example hydrohalic acids, for example hydrofluoric acid, hydrochloric acid or hydrobromic acid, as well as sulfuric acid, phosphoric acid or nitric acid, but it is also possible to use suitable organic acids, such as acetic acid, toluenesulfonic acid, etc..

The proton acceptors used are, for example, inorganic or organic bases, for example alkali metal or alkaline earth metal compounds, for example the hydroxides, oxides or carbonates of lithium, sodium, potassium, magnesium, calcium, strontium and barium, or alternatively hydrides, for example sodium hydride. There may be mentioned as organic bases, for example, tertiary amines, such as triethylamine, triethylene-diamine and pyridine.

In the processes described above, according to the reaction


conditions, there may be used in addition to the solvents which have been mentioned, for example, the following:

Halogenated hydrocarbons, for example chlorinated hydrocarbons, such as tetrachloroethane, methylene chloride, chloroform, carbon tetrachloride, chlorobenzene, chlorotoluene, ethers, such as ethyl propyl ether, methyl tert-butyl ether, di-n-butyl ether, diisobutyl ether, diisoamyl ether, diisopropyl ether, diethyl ether, ethylene glycol dimethyl ether, tetrahydrofuran, dioxane, nitrated hydrocarbons, such as nitrobenzene, nitriles, such as acetonitrile, butyronitrile, aliphatic or cycloaliphatic hydrocarbons, such as heptane, hexane, petroleum fractions in a boiling point range of from 70°C to 190°C, cyclohexane, petroleum ether, ligroin, esters, such as ethyl acetate, amides, for example formamide, dimethylformamide; ketones, such as acetone, methyl ethyl ketone; alcohols, especially lower aliphatic alcohols, for example methanol, ethanol, n-propanol, isopropanol and the isomers of the butanols; and water. Mixtures of the mentioned solvents and diluents are also suitable.


Methods of synthesis analogous to the above-described preparation processes have been published in the literature. There may be mentioned as references: A. Kreutzberger and J. Gillessen, *J. Heterocyclic Chem.* 22, 101 (1985); O. Stark, *Ber. Dtsch. Chem. Ges.* 42, 699 (1909); J. Hale, *J. Am. Chem. Soc.* 36, 104 (1914); G.M. Kosolapoff, *J. Org. Chem.* 26, 1895 (1961); St. Angerstein, *Ber. Dtsch. Chem. Ges.* 34, 3956 (1901); G.M. Kosolapoff, *J. Org. Chem.* 26, 1895 (1961); M.P.V. Boarland and J.F.W. McOmie, *J. Chem. Soc.* 1951, 1218; T. Matsukawa and K. Shirakuwa, *J. Pharm. Soc. Japan* 71, 933 (1951); *Chem. Abstr.* 46, 4549 (1952).

The described preparation processes, including all steps thereof, are included in the present invention.

The present invention relates also to the level intermediates of formula V




wherein R₁ is hydrogen ~~3-fluorine or 4-fluorine~~ and R₂ is C₁-C₄alkyl or halo- or hydroxy-substituted C₁-C₂alkyl, and to the intermediates of formula VI



wherein R₁ is hydrogen ~~3-fluorine or 4-fluorine~~ and R₂ is C₁-C₄alkyl, halo- or hydroxy-substituted C₁-C₂alkyl, C₃-C₆-cycloalkyl or C₃-C₆cycloalkyl mono- to tri-substituted by identical or different substituents selected from methyl and halogen.

Surprisingly, it has been found that the compounds of formula I have, for practical field application purposes, a very advantageous biocidal spectrum against phytopathogenic microorganisms, especially fungi. Compounds of formula I have very advantageous curative, preventive and, in particular, systemic properties, and can be used for protecting numerous cultivated plants. With the compounds of formula I it is possible to inhibit or destroy the pests which occur on plants or on parts of plants (fruit, blossoms, leaves, stems, tubers, roots) in different crops of useful plants, while at



the same time the parts of plants which grow later are also protected, for example, from attack by phytopathogenic microorganisms.

The compounds of formula I are effective, for example, against the phytopathogenic fungi belonging to the following classes: Fungi imperfecti (especially Botrytis, also Pyricularia, Helminthosporium, Fusarium, Septoria, Cercospora and Alternaria); Basidiomycetes (e.g. Rhizoctonia, Hemileia, Puccinia). They are also effective against the class of the Ascomycetes (e.g. Venturia and Erysiphe, Podosphaera, Monilinia, Uncinula) and of the Oomycetes (e.g. Phytophthora, Pythium, Plasmopara). They can also be used as dressing agents for protecting seeds (fruit, tubers, grains) and plant cuttings against fungus infections as well as against phytopathogenic fungi which occur in the soil.

The invention also relates to compositions comprising as active ingredient compounds of formula I, especially crop protection compositions, and to their use in the agricultural sector or related fields.

The present invention further embraces the preparation of those compositions, which comprises homogeneously mixing the active ingredient with one or more compounds or groups of compounds described herein. The invention furthermore relates to a method of treating plants, which comprises applying thereto the novel compounds of formula I or the novel compositions.

Target crops to be protected within the scope of the present invention comprise e.g. the following species of plants: cereals (wheat, barley, rye, oats, rice, maize, sorghum and related crops), beet (sugar beet and fodder beet), pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and black-

berries), leguminous plants (beans, lentils, peas, soybeans), oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts), cucumber plants (cucumber, marrows, melons), fibre plants (cotton, flax, hemp, jute), citrus fruit (oranges, lemons, grapefruit, mandarins), vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika), lauraceae (avocados, cinnamon, camphor), or plants such as tobacco, nuts, coffee, sugar cane, tea, pepper, vines, hops, bananas and natural rubber plants, as well as ornamentals.

The compounds of formula I are normally applied in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession, with further compounds. These further compounds can be fertilisers or micronutrient donors or other preparations that influence plant growth. They can also be selective herbicides as well as insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired together with further carriers, surfactants or other application-promoting adjuvants customarily employed in formulation technology.

Suitable carriers and adjuvants can be solid or liquid and correspond to the substances ordinarily employed in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilisers.

A preferred method of applying a compound of formula I, or an agrochemical composition which comprises at least one of said compounds, is foliar application. The number of applications and the rate of application depend on the risk of infestation by the corresponding pathogen. However, the compounds of formula I can also penetrate the plant through the roots via

the soil (systemic action) if the locus of the plant is impregnated with a liquid formulation, or if the compounds are applied in solid form to the soil, e.g. in granular form (soil application). In paddy rice crops, such granules may be applied in metered amounts to the flooded rice field. The compounds of formula I may, however, also be applied to seeds (coating) either by impregnating the seeds with a liquid formulation comprising a compound of formula I, or by coating them with a solid formulation.

The compounds of formula I are used in unmodified form or, preferably, together with the adjuvants conventionally employed in formulation technology, and are for this purpose advantageously formulated in known manner e.g. into emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wetttable powders, soluble powders, dusts, granules, and also encapsulations in e.g. polymer substances. As with the nature of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. Advantageous rates of application for direct treatment of the plants or for soil treatment are normally from 50 g to 5 kg of active ingredient (a.i.) per hectare, preferably from 100 g to 2 kg a.i./ha, most preferably from 200 g to 600 g a.i./ha. The amounts of active ingredient used for seeds that are dressed and then sown are markedly lower, being the equivalent of from 0.1 g a.i./ha to 500 g a.i./ha, i.e. the amounts of active ingredient to be found together with the seed in an area of 1 ha. An amount of from 0.5 g a.i./ha to 100 g a.i./ha is preferred.

The formulations, i.e. the compositions, preparations or mixtures comprising the compound (active ingredient) of formula I and, where appropriate, a solid or liquid adjuvant,

are prepared in known manner, e.g. by homogeneously mixing and/or grinding the active ingredients with extenders, e.g. solvents, solid carriers and, where appropriate, surface-active compounds (surfactants).

Suitable solvents are: aromatic hydrocarbons, preferably the fractions containing 8 to 12 carbon atoms, e.g. xylene mixtures or substituted naphthalenes, phthalates such as dibutyl phthalate or dioctyl phthalate, aliphatic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols and their ethers and esters, such as ethanol, ethylene glycol, ethylene glycol monomethyl or monoethyl ether, ketones such as cyclohexanone, strongly polar solvents such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethylformamide, as well as vegetable oils or epoxidised vegetable oils, such as epoxidised coconut oil or soybean oil; or water.

The solid carriers used, e.g. for dusts and dispersible powders, are normally natural mineral fillers such as calcite, talcum, kaolin, montmorillonite or attapulgite. In order to improve the physical properties it is also possible to add highly dispersed silicic acid or highly dispersed absorbent polymers. Suitable granulated adsorptive carriers are porous types, for example pumice, broken brick, sepiolite or bentonite; and suitable nonsorbent carriers are, for example, calcite or sand. In addition, a great number of pregranulated materials of inorganic nature can be used, e.g. especially dolomite or pulverised plant residues.

Particularly advantageous application-promoting adjuvants which are able to reduce substantially the rate of application are also natural (animal or vegetable) or synthetic phospholipids of the series of the cephalins and lecithins, which can be obtained e.g. from soybeans.

Depending on the nature of the compound of formula I to be formulated, suitable surface-active compounds are non-ionic, cationic and/or anionic surfactants having good emulsifying, dispersing and wetting properties. The term "surfactants" will also be understood as comprising mixtures of surfactants.

Both so-called water-soluble soaps and also water-soluble synthetic surface-active compounds are suitable anionic surfactants.

Suitable soaps are the alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts of higher fatty acids ($C_{10}-C_{22}$), e.g. the sodium or potassium salts of oleic or stearic acid or of natural fatty acid mixtures which can be obtained e.g. from coconut oil or tallow oil. Mention may also be made of fatty acid methyl-laurin salts.

More frequently, however, so-called synthetic surfactants are used, especially alkanesulfonates, fatty alcohol sulfates, sulfonated benzimidazole derivatives or alkylsulfonates.

The fatty alcohol sulfonates or sulfates are usually in the form of alkali metal salts, alkaline earth metal salts or unsubstituted or substituted ammonium salts and contain a C_8-C_{22} alkyl radical, which also includes the alkyl moiety of acyl radicals, e.g. the sodium or calcium salt of ligno-sulfonic acid, of dodecyl sulfate or of a mixture of fatty alcohol sulfates obtained from natural fatty acids. These compounds also comprise the salts of sulfated and sulfonated fatty alcohol/ethylene oxide adducts. The sulfonated benzimidazole derivatives preferably contain 2 sulfonic acid groups and one fatty acid radical containing 8 to 22 carbon atoms. Examples of alkylarylsulfonates are the sodium, calcium or triethanolamine salts of dodecylbenzenesulfonic

acid, dibutyl-naphthalenesulfonic acid, or of a condensate of naphthalenesulfonic acid and formaldehyde.

Also suitable are corresponding phosphates, e.g. salts of the phosphoric acid ester of an adduct of p-nonylphenol with 4 to 14 mol of ethylene oxide.

Non-ionic surfactants are preferably polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, or saturated or unsaturated fatty acids and alkylphenols, said derivatives containing 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon moiety and 6 to 18 carbon atoms in the alkyl moiety of the alkylphenols.

Further suitable non-ionic surfactants are the water-soluble adducts of polyethylene oxide with polypropylene glycol, ethylenediaminopolypropylene glycol and alkylpolypropylene glycol containing 1 to 10 carbon atoms in the alkyl chain, which adducts contain 20 to 250 ethylene glycol ether groups and 10 to 100 propylene glycol ether groups. These compounds usually contain 1 to 5 ethylene glycol units per propylene glycol unit.

Representative examples of non-ionic surfactants are nonylphenolpolyethoxyethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxy-polyethyleneethanol, polyethylene glycol and octylphenoxy-polyethoxyethanol.

Fatty acid esters of polyoxyethylene sorbitan, e.g. polyoxyethylene sorbitan trioleate, are also suitable non-ionic surfactants.

Cationic surfactants are preferably quaternary ammonium salts which contain, as N-substituent, at least one C₈-C₂₂alkyl radical and, as further substituents, unsubstituted or halo-

generated lower alkyl, benzyl or hydroxy-lower alkyl radicals. The salts are preferably in the form of halides, methyl sulfates or ethyl sulfates, e.g. stearyltrimethylammonium chloride or benzyldi(2-chloroethyl)ammonium bromide.

Further surfactants customarily employed in formulation technology are known to the person skilled in the art or can be taken from the relevant specialist literature.

The agrochemical compositions usually comprise 0.1 to 99 %, preferably 0.1 to 95 %, of a compound of formula I, 99.9 to 1 %, preferably 99.9 to 5 %, of a solid or liquid adjuvant, and 0 to 25 %, preferably 0.1 to 25 %, of a surfactant.


Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The compositions may also comprise further auxiliaries such as stabilisers, antifoams, viscosity regulators, binders, tackifiers as well as fertilisers or other active ingredients for obtaining special effects.

The following Examples serve to illustrate the invention in greater detail, but do not limit the invention.


1. Preparation Examples

Example 1.1 Preparation of the intermediate 2-anilino-4-formyldiethylacetal-6-methylpyrimidine



11.8 g (60 mmol) of phenylguanidine hydrogen carbonate and 12 g (64 mmol) of 1-methyl-3-formyldiethylacetal-1,3-propanedione in 80 ml of ethanol are heated under reflux for 4 hours with stirring, the evolution of carbon dioxide subsiding as the reaction progresses. After cooling, the ethanol is largely evaporated off and the residue is taken up in 30 ml of diethyl ether, washed three times with 30 ml of water each time, dried over sodium sulfate and filtered, and the diethyl ether is evaporated off. The dark brown oil that remains is dissolved in 350 ml of petroleum ether (b.p. 40-60°C) at elevated temperature, heated under reflux for 10 minutes in the presence of activated carbon, cooled to approximately 40°C and filtered over Hyflo, and concentrated to a volume of approximately 30 ml. The resulting yellowish crystals are isolated by filtration and melt at 63-65°C.


Example 1.2 Preparation of the intermediate 2-anilino-4-formyl-6-methylpyrimidine



3 g (10.4 mmol) of 2-anilino-4-formyldiethylacetal-6-methylpyrimidine and 1 g of concentrated hydrochloric acid in 30 ml of water are stirred at 45°C for 24 hours and then cooled to 15°C, and the pH is adjusted to 8 with sodium hydrogen carbonate solution. The light-brown powder is isolated by filtration and then washed with water and recrystallised from 70 ml of acetonitrile. The yellow crystals melt at 130-131°C.

Example 1.3 Preparation of 2-phenylamino-4-cyano-6-methyl-


pyrimidine



[comp. no. 1.1]


11.8 g (0.17 mol) of hydroxylamine hydrochloride are added in portions over a period of 5 minutes, with stirring, to 21.3 g (0.1 mol) of 2-anilino-4-formyl-6-methylpyrimidine in 90 ml of pyridine, the temperature rising to 40°C. After stirring at 50°C for half an hour, 40 ml of acetic anhydride are added dropwise over a period of a quarter of an hour, the temperature rising to 70°C. After stirring at 90°C for 3 hours, the reaction mixture is cooled and then concentrated in vacuo, and the dark brown oily residue is poured onto 600 ml of ice-water and adjusted to pH 8 with dilute sodium hydroxide solution. After extracting three times with 150 ml of ethyl acetate each time, the combined extracts are washed twice with 50 ml of water each time, dried over sodium sulfate and filtered, and the solvent is evaporated off. The dark brown oily residue is dissolved in 200 ml of tetrahydrofuran, treated with activated carbon and filtered, and the solvent is evaporated off. The brownish solid that remains is recrystallised twice from diisopropyl ether/methanol. The yellow crystalline powder melts at 125-126°C.

Example 1.4 Preparation of the intermediate 2-anilino-4-oximino-6-cyclopropylpyrimidine



First 3.6 g (52 mmol) of hydroxylamine hydrochloride and then 2.8 g (26.4 mmol) of soda in 15 ml of water are added in portions over a period of a quarter of an hour, at room temperature and with stirring, to 10 g (42 mmol) of 2-anilino-4-formyl-6-cyclopropylpyrimidine in 150 ml of ethanol, and the mixture is then stirred at room temperature for 24 hours. The yellowish suspension is diluted with 50 ml of water, filtered and then washed with 10 ml of water, and the yellow semi-crystalline residue is recrystallised from 120 ml of ethanol. The beige crystals melt at 175-185°C.

Example 1.5 Preparation of 2-phenylamino-4-cyano-6-cyclopropylpyrimidine



[comp. no. 1.11]

3 g (53 mmol) of methyl isocyanate in 10 ml of dioxane are added dropwise at room temperature over a period of 10 minutes, with stirring, to 7.6 g (30 mmol) of 2-anilino-4-oximino-6-cyclopropylpyrimidine and 200 mg of 1,4-diazabicyclo[2.2.2]octane in 60 ml of dioxane. After stirring at 75°C for 16 hours, the mixture is cooled, the solvent is evaporated off and the light-brown oil that remains is purified by column chromatography over silica gel (toluene/-

diethyl ether: 3/2). After the eluant mixture has been evaporated off, the yellow solid is purified by recrystallisation from diisopropyl ether. The yellow crystals melt at 122-125°C.

The following compounds of formula I can be prepared in that manner or in accordance with one of the methods indicated above.







Table 1:




Comp. no.	R ₁	R ₂	Physical constant
1.1	H	-CH ₃	m.p. 125-126°C
1.3	H	-C ₂ H ₅	
1.6	H	-C ₃ H _{7-n}	
1.8	H	-C ₃ H _{7-i}	
1.11	H		m.p. 122-125°C
1.14	H		
1.15	H		
1.18	H		
1.21	H		
1.23	H		
1.27	H		
1.30	H	-C ₄ H _{9i}	
1.31	H	-C ₄ H _{9sec}	

Table 2: Novel Intermediates of Formula V



Comp. no.	R ₁	R ₂	Physical content
2.01	H	CH ₃	m.p. 130-131°C
2.04	H	C ₂ H ₅	
2.07	H	nC ₃ H ₇	
2.10	H	isoC ₃ H ₇	
2.13	H	nC ₄ H ₉	
2.16	H	isoC ₄ H ₉	
2.19	H	secC ₄ H ₉	
2.22	H	tertC ₄ H ₉	

Table 3 Novel Intermediates of Formula VI



Comp. no.	R ₁	R ₂	Physical content
3.1	H	-CH ₃	m.p. 240-241°C
3.3	H	-C ₂ H ₅	
3.6	H	-C ₃ H _{7-n}	
3.8	H	-C ₃ H _{7-i}	
3.11	H		m.p. 175-185°C
3.14	H		
3.15	H		
3.18	H		
3.21	H		
3.23	H		
3.27	H		

Table 3 (continuation)

Comp. no.	R ₁	R ₂	Physical constant
3.30	H	-C ₄ H ₉ i	
3.31	H	-C ₄ H ₉ sec	

2. Formulation Examples for solid active ingredients of formula I (throughout, percentages are by weight)

2.1. Wettable powders

	a)	b)	c)
a compound of Table 1	25 %	50 %	75 %
sodium lignosulfonate	5 %	5 %	-
sodium lauryl sulfate	3 %	-	5 %
sodium diisobutyl-naphthalene-sulfonate	-	6 %	10 %
octylphenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
kaolin	62 %	27 %	-

The active ingredient is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders which can be diluted with water to give suspensions of the desired concentration.

2.2. Emulsifiable concentrate

a compound of Table 1	10 %
octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %
calcium dodecylbenzenesulfonate	3 %
castor oil polyglycol ether (35 mol of ethylene oxide)	4 %
cyclohexanone	34 %
xylene mixture	50 %



Emulsions of any required concentration can be obtained from this concentrate by dilution with water.

2.3. Dusts

	a)	b)
a compound of Table 1	5 %	8 %
talcum	95 %	-
kaolin	-	92 %

Ready-for-use dusts are obtained by mixing the active ingredient with the carrier and grinding the mixture in a suitable mill.

2.4. Extruder granules

a compound of Table 1	10 %
sodium lignosulfonate	2 %
carboxymethylcellulose	1 %
kaolin	87 %

The active ingredient is mixed and ground with the adjuvants, and the mixture is subsequently moistened with water. The mixture is extruded and then dried in a stream of air.

2.5. Coated granules

a compound of Table 1	3 %
polyethylene glycol (mol. wt. 200)	3 %
kaolin	94 %

The finely ground active ingredient is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

2.6. Suspension concentrate

a compound of Table 1	40 %
ethylene glycol	10 %
nonylphenol polyethylene glycol	

ether (15 mol of ethylene oxide)	6 %
sodium lignosulfonate	10 %
carboxymethylcellulose	1 %
37 % aqueous formaldehyde solution	0.2 %
silicone oil in the form of a 75 % aqueous emulsion	0.8 %
water	32 %

The finely ground active ingredient is intimately mixed with the adjuvants, giving a suspension concentrate from which suspensions of any desired concentration can be obtained by dilution with water.

3. Biological Examples

Example 3.1: Action against Venturia inaequalis on apple shoots

Residual-protective action

Apple cuttings with 10-20 cm long fresh shoots are sprayed with a spray mixture (0.006 % active ingredient) prepared from a wettable powder formulation of the test compound. The treated plants are infected 24 hours later with a conidia suspension of the fungus. The plants are then incubated for 5 days at 90-100 % relative humidity and stood in a greenhouse for a further 10 days at 20-24°C. Scab infestation is evaluated 15 days after infection.

Compounds of Table 1 exhibit good activity against Venturia (less than 20 % infestation). On the other hand, Venturia infestation is 100 % on untreated and infected control plants.

Example 3.2: Action against Botrytis cinerea on apples

Residual-protective action

Artificially damaged apples are treated by grafting onto the damaged sites a spray mixture (0.002 % active ingredient)



prepared from a wettable powder formulation of the test compound. The treated fruits are then inoculated with a spore suspension of the fungus and incubated for one week at high humidity and about 20°C. Evaluation is made by counting the rotted damaged sites and deriving the fungicidal activity of the test compound therefrom.

Compounds of Table 1 exhibit good activity against Botrytis (less than 20 % infestation). Thus e.g. compounds nos. 1.1 and 1.11 reduce Botrytis infestation to 0 to 10 %. On the other hand, Botrytis infestation is 100 % on untreated and infected control plants.

Example 3.3: Action against Erysiphe graminis on barley


a) Residual-protective action

Barley plants about 8 cm in height are sprayed with a spray mixture (0.006 % active ingredient) prepared from a wettable powder formulation of the test compound. The treated plants are dusted with conidia of the fungus after 3 to 4 hours. The infected barley plants are stood in a greenhouse at about 22°C. The fungus infestation is evaluated after 10 days.

Compounds of Table 1 exhibit good activity against Erysiphe (less than 20 % infestation). On the other hand, Erysiphe infestation is 100 % on untreated and infected control plants.

Example 3.4: Action against Helminthosporium gramineum

Wheat grains are contaminated with a spore suspension of the fungus and dried. The contaminated grains are dressed with a suspension of the test compound prepared from a wettable powder (600 ppm of active ingredient, based on the weight of the seeds). Two days later the grains are placed in suitable agar dishes and the development of fungus colonies around the grains is assessed after a further four days. The effectiveness of the test compound is evaluated on the basis of the



number and size of the colonies. The compounds of the Table substantially prevent fungus infestation (0 to 10 % fungus infestation).

Example 3.5: Action against Colletotrichum lagenarium on cucumbers


After a cultivation period of two weeks, cucumber plants are sprayed with a spray mixture (concentration 0.002 %) prepared from a wettable powder formulation of the test compound. After two days, the plants are infected with a spore suspension (1.5×10^5 spores/ml) of the fungus and incubated for 36 hours at 23°C and high humidity. Incubation is then continued at normal humidity and about 22-23°C. Evaluation of fungus infestation is made 8 days after infection. Fungus infestation is 100 % on untreated and infected control plants.

Compounds of Table 1 exhibit good activity and inhibit the spread of the disease. Fungus infestation is reduced to 20 % or less.



~~What is claimed is:~~ The claims defining the invention are as follows:

1. A compound of formula I



wherein R_1 is hydrogen ~~3-fluorine or 4-fluorine~~ and R_2 is C_1 - C_4 alkyl, halo- or hydroxy-substituted C_1 - C_2 alkyl, C_3 - C_6 -cycloalkyl or C_3 - C_6 cycloalkyl mono- to tri-substituted by identical or different substituents selected from methyl and halogen; or an acid addition salt or metal salt complex thereof.

2. A compound of formula I according to claim 1 wherein R_1 is hydrogen ~~3-fluorine or 4-fluorine~~ and R_2 is C_1 - C_3 alkyl, cyclopropyl or cyclopropyl substituted by methyl or by chlorine.

3. A compound of formula I according to claim 1 wherein R_1 is hydrogen and R_2 is methyl, ethyl, cyclopropyl or 2-methyl-cyclopropyl.

4. A compound of formula I according to claim 1 wherein R_1 is hydrogen ~~3-fluorine or 4-fluorine~~ and R_2 is methyl, ethyl or cyclopropyl.


5. A compound of formula I according to claim 4 selected from the group:

2-phenylamino-4-cyano-6-methylpyrimidine (comp. no. 1.1),
2-phenylamino-4-cyano-6-ethylpyrimidine (comp. no. 1.3), and
2-phenylamino-4-cyano-6-cyclopropylpyrimidine (comp.
no. 1.11), and




6. A process for the preparation of a compound of formula I, which process comprises

1.1 reacting a guanidine salt of formula IIa




wherein A^{\ominus} is an acid anion
or a guanidine compound of formula IIb



with a ketone of formula III



wherein R_3 is an alkyl radical, in a protic solvent or without a solvent, at temperatures of from 40° to 160°C, to form a pyrimidine compound of formula IV




and

1.2 hydrolysing the resulting acetal of formula IV in the




presence of an acid, in water or aqueous mixtures of inert solvents, at temperatures of from 20° to 100°C, to form the pyrimidinylaldehyde of formula V




and

1.3 then converting the aldehyde of formula V with a hydroxylamine salt and a base that frees the hydroxylamine, in a protic solvent, at temperatures of from 10° to 90°C, into the oxime of formula VI



and

1.3.1 subjecting the oxime of formula VI to a reaction in which the elements of water are removed; the oxime being converted in an intermediate step into a compound of formula VII




wherein R' is an acyl radical COR'', a carbamoyl radical



CONHR" or an oxycarbonyl radical COOR", in which R" is an alkyl radical, and the compound of formula VII then being decomposed by removal of the acyl radical HOCOR" in inert solvents at temperatures of from 60° to 120°C; or

1.4 from the pyrimidinealdehyde of formula V, preparing the oxime of formula VI in an intermediate step by reaction in tertiary bases, and forming the compound of formula I directly therefrom in situ by treatment with an acid halide or acid anhydride at temperatures of from 40° to 120°C; or

2. reacting a compound of formula VIII



(VIII)


with a cyanide of formula IX



(IX),

wherein Hal is fluorine, chlorine, bromine or iodine and Me is an alkali metal cation, alkaline earth metal cation or heavy metal cation, in an aprotic solvent at temperatures of from 60° to 160°C; or

3.1 cyclising urea of formula X




(X)

with a diketone of formula III




wherein R₃ is an alkyl radical, in the presence of an acid in an inert solvent at temperatures of from 20° to 140°C, to form a pyrimidine compound of formula XI



and

3.2 replacing the OH group in the compound of formula XI by halogen using excess PO(Hal)₃ in the presence or absence of an inert solvent at temperatures of from 50° to 110°C to form a compound of formula XII



wherein Hal is halogen, and then


3.3 reacting a compound of formula XII with an aniline compound of formula XIII



either



- a) in the presence of a proton acceptor, with or without an inert solvent, or
 b) in the presence of an acid in an inert solvent, in each case at temperatures of from 60° to 120°C, to form a compound of formula IV



5 and converting the resulting acetal into a compound of formula I in the manner described under 1.2, 1.3 and 1.3.1, the radicals R₁ and R₂ in the above-described processes being as defined under formula I.

7. A process for the preparation of a 2-anilino-4-cyanopyrimidine derivative, substantially as hereinbefore described with reference to any one the Preparation
 10 Examples.

8. A 2-anilino-4-cyanopyrimidine prepared by the process of claim 6 or claim 7.

9. A 2-anilino-4-cyanopyrimidine derivative substantially as hereinbefore described with reference to any one of the Preparation Examples.

10. A composition for protecting plants from attack by microorganisms,
 15 comprising a microbicidally effective amount of at least one compound according to any one of claims 1 to 5, 8 or 9 together with a microbicidally acceptable carrier, diluent, excipient and/or adjuvant.

11. A composition according to claim 10, which comprises at least one carrier.

12. A composition for protecting plants from attack by microorganisms,
 20 substantially as hereinbefore described with reference to the Formulation Examples.

13. A process for the preparation of a composition according to any one of claims 10 to 12 which process comprises homogeneously mixing at least one compound defined according to any one of claims 1 to 5, 8 or 9 with a suitable solid or liquid carrier and/or adjuvant.

25 14. A method of protecting plants against attack by phytopathogenic microorganisms, which comprises applying a phytopathogenically effective amount of a compound according to any one of claims 1 to 5, 8 or 9 or a composition according to any one of claims 10 to 12 to the plant or to the locus thereof.

30 15. A method according to claim 14, wherein the phytopathogenic microorganisms are fungus organisms.

Dated 31 August, 1993

Ciba-Geigy AG

Patent Attorneys for the Applicant/Nominated Person
 SPRUSON & FERGUSON

