

(12) **UK Patent Application** (19) **GB** (11) **2491491** (13) **A**

(43) Date of A Publication

**05.12.2012**

(21) Application No: **1210657.1**

(22) Date of Filing: **19.01.2011**

Date Lodged: **15.06.2012**

(30) Priority Data:  
(31) **61297249** (32) **21.01.2010** (33) **US**  
(31) **61385318** (32) **22.09.2010** (33) **US**

(62) Divided from Application No **1201842.0** under section 15(9) of the Patents Act 1977

(71) Applicant(s):  
**Goodchild Investments PTY LTD.**  
**38 Somers Avenue, Malvern 3144, Victoria, Australia**

**Monash University**  
**(Incorporated in Australia – Victoria)**  
**Wellington Road, Clayton 3168, Victoria, Australia**

(72) Inventor(s):  
**Juliet Marguerite Goodchild**  
**Colin Stanley Goodchild**  
**Benjamin James Boyd**

(74) Agent and/or Address for Service:  
**Dehns**  
**St. Bride's House, 10 Salisbury Square, LONDON,**  
**EC4Y 8JD, United Kingdom**

(51) INT CL:  
**A61K 47/40** (2006.01) **A61K 31/56** (2006.01)  
**A61K 31/57** (2006.01) **A61K 47/48** (2006.01)  
**A61P 23/00** (2006.01)

(56) Documents Cited:  
**EP 0399716 B1** **WO 2004/039426 A1**  
**WO 2001/070234 A1** **WO 1993/017711 A1**  
**JOURNAL OF PARENTERAL SCIENCE &**  
**TECHNOLOGY, Vol. 43(6), 1989, (BREWSTER M. E. ET**  
**AL), 'Development of a Non-Surfactant Formulation**  
**for Alfaxalone Through the Use of Chemically-**  
**Modified Cyclodextrins', pages 262 - 265**  
**JOURNAL OF PHARMACEUTICAL SCIENCES Vol. 84**  
**(10), 1995, (BREWSTER M. E. ET AL), 'Preparation,**  
**Characterization, and Anaesthetic Properties of 2-**  
**hydroxypropyl-beta-cyclodextrin Complexes of**  
**Pregnenolome in Rat and Mouse', pages 1154 - 1159**

(58) Field of Search:  
Other: **WPI, EPODOC, MEDLINE**

(54) Title of the Invention: **Anaesthetic formulation**  
Abstract Title: **Improved anaesthetic formulations comprising neuroactive steroids**

(57) The present invention relates generally to the field of drug delivery systems for neuroactive steroid anaesthetic agents. More particularly, anaesthetic and sedative formulations are provided in the form of host/guest preparations comprising one or more neuroactive steroid anaesthetics and a cyclodextrin. Particular cyclodextrins contemplated include sulfoalkyl ether cyclodextrins and modified forms thereof, especially sulfobutyl ether  $\beta$ -cyclodextrin. The neuroactive steroid may be alphaxolone, alphadolone, allopregnanolone or eltanolone (pregnanolone). The formulation may additionally comprise an antimicrobial agent, a preservative, a buffer and/or a copolymer, particularly hydroxypropyl methylcellulose (HPMC), polyvinyl pyrrolidone (PVP) or carboxymethylcellulose. The composition may be administered by infusion or bolus.

**GB**  
**2491491**  
**A**




Figure 1a




Figure 1b




Figure 1c






Figure 1d



**Figure 1e**



**Figure 1f**




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9






Figure 10




Figure 11




Figure 12

## **ANAESTHETIC FORMULATION**

### **FILING DATA**

5 [0001] This application is associated with and claims priority from United States  
Provisional Patent Application No. 61/297,249, filed on 21 January 2010, entitled  
"Anaesthetic formulation" AND United States Provisional Patent Application No.  
61/385,318 filed on 22 September 2010, entitled "Anaesthetic formulation", the entire  
contents of which, are incorporated herein by reference.

10

### **FIELD**

[0002] The present invention relates generally to the field of drug delivery systems for  
neuroactive steroid anaesthetic agents. More particularly, anaesthetic and sedative  
15 compositions are provided in the form of host/guest preparations comprising one or more  
neuroactive steroid anaesthetics and a cyclodextrin or a modified form thereof.

### **BACKGROUND**

20 [0003] Bibliographic details of references provided in the subject specification are listed at  
the end of the specification.

[0004] Reference to any prior art is not, and should not be taken as an acknowledgment or  
any form of suggestion that this prior art forms part of the common general knowledge in  
25 any country.

[0005] Drug delivery systems aim to provide the required amount of drug systemically or  
to a targeted site for a time and under conditions sufficient to have the desired effect.  
Some drug delivery systems require carrier materials to mitigate particular undesirable  
30 properties of the drugs. One such type of carrier molecule is a cyclodextrin which acts as a  
host for a selected guest molecule.

[0006] Cyclodextrins are cyclic oligosaccharides with hydroxyl groups on their outer surface and a void central cavity which has a lipophilic character. Cyclodextrins are capable of forming inclusion complexes with hydrophobic molecules. The stability of the resulting host/guest complex depends on how readily the guest molecule occupies the central cavity of the host.

[0007] The most common cyclodextrins are  $\alpha$ -,  $\beta$ - and  $\gamma$ - cyclodextrins consisting of 6, 7 and 8  $\alpha$ -1,4-linked glucose units, respectively. Cyclodextrins have relatively low solubility in water and organic solvents and this limits their use in pharmaceutical formulations. For a description of the general chemistry of cyclodextrin, reference can be made to Fromming and Szejtlic (eds), *Cyclodextrins in Pharmacy*, Kluwer: Dordrecht, The Netherlands, 1994; Atwood, Davies, MacNicol and Vogtie (Eds), *Comprehensive Supramolecular Chemistry* Vol 4, Pergamon: Oxford UK, 1996; and Thomason, *Crit Rev Ther Drug Carrier Syst* 14:1, 1997.

[0008] Alphaxalone [Alfaxalone or 3- $\alpha$ -hydroxy-5- $\alpha$ -,pregnan-11,20-dione] is a potent neuroactive steroid anaesthetic currently used in veterinary medicine (Child *et al.*, *British Journal of Anaesthesia* 43:2-13, 1971).

[0009] Alphaxalone was widely used around the world as an intravenous anaesthetic together with alphadolone [Althesin; Alfathesin] in human patients until 1983. Although these anaesthetics have a high therapeutic index, they were nevertheless withdrawn from clinical practice due to occasional, unpredictable yet severe anaphylactoid reactions to a polyethoxylated castor oil excipient (Cremophor EL [Registered Trademark]).

[0010] Currently, a lipid formulation of di-isopropyl phenol (propofol) is the most highly used anaesthetic agent. Propofol, however, can be contraindicated in certain at risk patients due to its lowering effect on blood pressure, the effect it has on reducing cardiac output and it can adversely affect respiratory control. In particular, propofol is formulated in a lipid emulsion which can support microbial growth if contaminated. The formulation cannot, in fact, be sterilized. There have been instances where microbially contaminated

propofol formulations have resulted in patients becoming infected. One other issue with the current propofol formulation is the pain induced following or during intravenous injection. Attempts to re-formulate in a water-based preparation have led to increased injection pain. Propofol also can lead to cardiovascular and respiratory depression and has a low therapeutic index of 5, i.e. only 5 times the normal anaesthetic dose can lead to death. Furthermore, the anaesthetic is incompatible with plastic storage containers and plastic syringes which complicates syringe delivery equipment which is frequently in standard use for intravenous anaesthesia and sedation. The drug can also cause hyperlipidaemia and can induce toxicity when used in a larger dose by infusion. This is particularly problematic in the intensive care setting.

**[0011]** A neuroactive steroid anaesthetic has the potential for being more efficacious with fewer side effects than propofol.

**[0012]** There is a need, therefore, to develop a suitable formulation to enable the use of a neuroactive steroid anaesthetic agent in subjects.

## SUMMARY

[0013] Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to  
5 imply the inclusion of a stated element or integer or method step or group of elements or integers or method steps but not the exclusion of any other element or integer or method step or group of elements or integers or method steps.

[0014] The present invention provides a host/guest complex formulation comprising a  
10 neuroactive steroid anaesthetic and a cyclodextrin or modified form thereof for use in inducing anaesthesia or sedation in mammalian subjects. Generally, the neuroactive steroid anaesthetic is sparingly soluble. The host/guest complex formulation is, therefore, a drug delivery system for a neuroactive steroid anaesthetic. In an embodiment, the cyclodextrin is a modified polyanionic  $\beta$ -cyclodextrin and the neuroactive steroid  
15 anaesthetic is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof. However, all cyclodextrins are contemplated herein including  $\gamma$  and  $\alpha$  cyclodextrins or their modified forms as well as their salts. The term "derivative"  
20 includes deuterated derivatives of the neuroactive steroid anaesthetic. Deuterated derivatives are contemplated for use as improved medicaments. One or more hydrogen atoms may be replaced by deuterium. Modified forms of cyclodextrins include methylated, hydroxyalkylated, branched, alkylated, acylated and anionic cyclodextrins. By "alkylated" includes an alkyl ether derivative as well as an alkyl ether-alkyl ether  
25 cyclodextrin. The agent "alphadolone" includes its salt, alphadolone acetate. Reference to a cyclodextrin or a modified form thereof includes its salts (e.g. a sodium salt).

[0015] Accordingly, one aspect of the present invention is directed to an anaesthetic or sedative composition comprising a neuroactive steroid anaesthetic formulated with a  
30 cyclodextrin or modified form thereof.

[0016] The anaesthetic or sedative formulation of the present invention exhibits features

such as being sterilizable, causes reduced incidence of pain on injection, has a larger therapeutic index relative to propofol (including a therapeutic index greater than 5), is capable of storage in a plastic container and induces a rapid induction of anaesthesia to surgical levels with similar or more rapid awakening time than propofol or Althesin  
5 (alphaxalone and alphadolone).

[0017] Hence, another aspect of the present invention provides an anaesthetic or sedative composition comprising a neuroactive steroid anaesthetic and a cyclodextrin or modified form thereof wherein the anaesthetic and cyclodextrin are formulated to provide an  
10 anaesthetic composition which exhibits a property selected from being sterilizable, exhibiting minimal pain on intravenous injection, having a therapeutic index greater than 5 and is storable in a plastic container. In an embodiment, the formulation has one, two, three or all four of these properties.

[0018] In a related embodiment, the present invention provides an anaesthetic or sedative delivery host/guest composition comprising a cyclodextrin host or a modified form thereof with a neuroactive steroid anaesthetic drug guest, the host/guest composition formulated to be sterilizable, administrable by intravenous injection with minimal pain and to exhibit a therapeutic index of greater than 5. The formulation may also be storable in a plastic  
20 container.

[0019] More particularly, the present invention provides an anaesthetic or sedative composition comprising a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone,  
25 hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof formulated with a cyclodextrin or modified form thereof.

[0020] Even more particularly, the present invention is directed to an anaesthetic or  
30 sedative composition comprising a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and

tetrahydrodeoxycorticosterone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof formulated with a cyclodextrin or a modified form thereof wherein the composition exhibits a property selected from being sterilizable, exhibiting minimal pain on intravenous injection, having a therapeutic index greater than 5 and is storable in a plastic container.

[0021] A particular cyclodextrin useful in the practice of the present invention is a sulfoalkyl ether cyclodextrin such as (7) sulfobutyl ether  $\beta$ -cyclodextrin. This compound can be prepared as described in United States Patent No. 5,376,645. Another useful cyclodextrin is an alkyl ether derivative including a sulfoalkyl ether-alkyl ether cyclodextrin. However, the present invention extends to other cyclodextrin derivatives such as methylated, hydroxyalkylated, branched, acylated and anionic forms. The anaesthetic formulation of the present invention enables injectable administration to mammalian subjects and in particular human patients.

[0022] Another aspect of the present invention provides an anaesthetic or sedative composition comprising a neuroactive steroid selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof formulated with a sulfoalkyl ether cyclodextrin or modified form thereof to generate a sterilizable composition with a therapeutic index of greater than 5.

[0023] In an embodiment, the composition is also storable in a plastic container.

[0024] The formulation may comprise a buffer to maintain pH within a range of from about pH5.5 to pH8. Alternatively, the formulation may not be buffered wherein the pH of the formulation may be from about pH3 to about pH9.5. The formulation may also comprise a preservative, anti-microbial agent and/or an agent which reduces toxicity. In addition, to improve solubility and/or stability, a co-polymer may be included. Examples of suitable co-polymers include hydroxyl propyl methyl cellulose (HPMC), polyvinyl pyrrolidone (PVP), and carboxymethyl cellulose (CMC).



[0025] The present invention further contemplates inducing or maintaining by infusion or intermittent bolus administration, anaesthesia or sedation in a subject, the method comprising administering an anaesthetic-effective amount of a neuroactive steroid  
5 anaesthetic formulated with a cyclodextrin, for a time and under conditions to induce anaesthesia or sedation.

[0026] More particularly, the present invention provides a method of inducing or maintaining by infusion or intermittent bolus administration, anaesthesia or sedation in a  
10 subject, the method comprising administering an anaesthetic-effective amount of a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and pharmacologically acceptable derivatives, salts or pro-drug forms thereof formulated with a cyclodextrin or  
15 modified form thereof for a time and under conditions sufficient to induce anaesthesia or sedation wherein the anaesthetic or sedative formulation exhibits a property selected from being sterilizable, exhibiting minimal pain on intravenous injection and having a therapeutic index greater than 5.

20 [0027] In an embodiment, the formulation is also storable in a plastic container.

[0028] Generally, the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is from about 1:1 to about 1:6, more particularly about 1:1 to about 1:4, even more particularly about 1:1 to 1:3 and still more particularly about 1:2.

25

[0029] These aspects of the present invention extend to inducing or maintaining by infusion or intermittent bolus administration, anaesthesia or sedation or both in subjects.

[0030] The formulation may be packaged for sale with a set of instructions. The  
30 instructions may include a patient management protocol comprising administering to the patient an effective amount of neuroactive steroid anaesthetic such as selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone),

ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof formulated with a cyclodextrin for a time and under conditions sufficient to induce anaesthesia. As indicated above, a suitable cyclodextrin includes a  
5 sulfoalkyl ether dextrin, such as (7) sulfobutyl ether  $\beta$ -cyclodextrin as well as alkyl ether derivatives such as sulfoalkyl-alkyl ether cyclodextrins. Other derivatives include methylated, hydroxyalkylated, branched, alkylated, acylated and anionic cyclodextrins.

[0031] The present invention further contemplates the use of a neuroactive steroid  
10 anaesthetic and a cyclodextrin or modified form thereof, in the manufacture of a medicament to induce anaesthesia in a subject. In a particular embodiment, the neuroactive steroid anaesthetic is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and pharmacologically  
15 acceptable derivatives, salts and pro-drug forms thereof.

[0032] Kits comprising in compartmental form a neuroactive steroid anaesthetic in a first compartment and a cyclodextrin, such as a sulfoalkyl ether cyclodextrin, in a second compartment and optionally excipients and/or co-polymers in a third or other compartment  
20 are also contemplated herein. The kit may be in the form of a modified syringe.

[0033] Labeled forms of the neuroactive steroid anaesthetic are also useful in monitoring and tracking the anaesthetic during sedation or anaesthesia. Kits and apparatus are therefore provided herein to assist in the monitoring of labelled neuroactive steroid  
25 anaesthetics. Labeled derivatives include deuterated, tritiated and other labeled agents.

## BRIEF DESCRIPTION OF THE FIGURES

[0034] Some figures contain color representations or entities. Color photographs are available from the Patentee upon request or from an appropriate Patent Office. A fee may be imposed if obtained from a Patent Office.

[0035] **Figure 1a through 1f** are graphical representations of experiments on male Wistar rats implanted with indwelling internal jugular vein intravenous catheters under halothane anaesthesia and then provided with propofol (a,b), Althesin [alphaxalone and alphadolone acetate] (c,d) or Phaxan<sub>CD</sub> [alphaxalone in a 1:2 molar complexation ratio with (7) sulfobutyl ether  $\beta$ -cyclodextrin] (e,f).

[0036] **Figure 2** is a graphical representation of lethal dosing of two alphaxalone preparations [Phaxan<sub>CD</sub> and Althesin in rats].

15

[0037] **Figure 3** is a graphical representation of a probit plot for lethality dosing of an Althesin preparation in rats.

[0038] **Figure 4** is a graphical representation of sleep time in rats using repeated doses of Phaxan<sub>CD</sub> (alphaxalone in a 1:2 molar complexation ratio with (7) sulfobutyl ether  $\beta$ -cyclodextrin).

20

[0039] **Figure 5** is a graphical representation of righting reflex time response curves in rats using pregnanolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

25

[0040] **Figure 6** is a graphical representation of tail pinch time response curve in rats using pregnanolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

[0041] **Figure 7** is a graphical representation of rotarod test time response curve in rats using pregnanolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

30

[0042] **Figure 8** is a graphical representation of righting reflex time response curves in rats using alphadolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

5 [0043] **Figure 9** is a graphical representation of tail pinch time response curves in rats using alphadolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

[0044] **Figure 10** is a graphical representation of rotarod test time response curves in rats using alphadolone in a (7) sulfobutyl ether  $\beta$ -cyclodextrin.

10 [0045] **Figure 11** is a graphical representation of percentage change in mean systolic blood pressure in rats after injection with propofol, Althesin or Phaxan<sub>CD</sub>.

[0046] **Figure 12** is a graphical representation of percentage change in mean diastolic blood pressure in rats after injection with propofol, Althesin or Phaxan<sub>CD</sub>.

15

[0047] Reference to "Phaxan<sub>CD</sub>" means an alphaxalone preparation with (7) sulfobutyl ether  $\beta$ -cyclodextrin.

## DETAILED DESCRIPTION

[0048] The present invention provides a drug delivery system for a neuroactive steroid anaesthetic. Generally, the neuroactive steroid anaesthetic is sparingly soluble in water.

5 The drug delivery system comprises a host carrier in the form of a cyclodextrin or modified form thereof. Reference to "cyclodextrin" includes in one embodiment an  $\alpha$ -,  $\beta$ - or  $\gamma$ -cyclodextrin or a modified or derivatized form thereof. Reference to "cyclodextrin" in another embodiment includes a sulfoalkyl ether dextrin such as (7) sulfobutyl ether  $\beta$ -cyclodextrin or an alkyl ether derivative thereof such as a sulfobutyl ether-alkyl ether

10 cyclodextrin. Derivatives of cyclodextrins include methylated, hydroxyalkylated, branched, alkylated, acylated and anionic cyclodextrins. By "alkylated" includes an alkyl ether derivative such as an alkyl ether-alkyl ether cyclodextrin. Particular cyclodextrins contemplated herein are shown in Table 7 [Uekama *et al.*, *Chem. Rev.* 98: 2045-2076, 1998] and include  $\beta$ -cyclodextrin sulfobutyl ethers, ethyl ethers,  $\beta$ -cyclodextrin sulfobutyl

15 ethers (flat),  $\gamma$ -cyclodextrin sulfobutyl ethers and  $\alpha$ -cyclodextrin sulfobutyl ethers and their salts (e.g. sodium salts).

[0049] The drug delivery system of the present invention enables a neuroactive steroid anaesthetic to be administered to a subject in a sterilized form. Furthermore, the delivery

20 itself is with less pain compared to the intravenous administration of propofol. The formulation of the present invention additionally has a therapeutic index greater than 5 (meaning that administration of greater than 5 times the anaesthetic dose can lead to death in a test animal). By "greater than 5" means a therapeutic index of between 5 and 200 including 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180,

25 190 and 200 as well as integers or fractions in between. The formulation of the present invention is also storable in a plastic container and is compatible for use in plastic delivery apparatus.

[0050] Accordingly, an aspect of the present invention provides an anaesthetic or sedative

30 delivery host/guest composition comprising a cyclodextrin host or modified form thereof with a neuroactive steroid anaesthetic drug guest, the host/guest composition formulated to be sterilizable, administrable by intravenous injection with minimal pain and to exhibit a

therapeutic index of greater than 5. In an embodiment, the formulation may also be storable in a plastic container. The formulation may exhibit one, two, three or all four of these properties.

5 [0051] By "reduced pain" means compared to a formulation comprising propofol as a reference.

[0052] The formulation is useful for inducing anaesthesia or sedation in mammalian subjects and in particular human subjects.

10

[0053] In an embodiment, the neuroactive steroid is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org 20599, Org 21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof.

15

[0054] An example of a pharmacologically acceptable salt is alphadolone acetate, which is encompassed by the present invention. An example of a derivative of a neuroactive steroid anaesthetic is a deuterated derivative. A "modified" cyclodextrin includes a derivative of a cyclodextrin.

20

[0055] Accordingly, another aspect of the present invention is directed to a drug delivery host/guest composition comprising a cyclodextrin host or modified form thereof with a neuroactive steroid anaesthetic drug guest selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, 25 minaxolone, Org 20599, Org 21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof, the host/guest composition being sterilizable, administrable by intravenous injection with minimal pain, exhibiting a therapeutic index of greater than 5 and/or storable in a plastic container. The formulation can also initiate rapid induction of anaesthesia to surgical levels with similar 30 or more rapid waking time compared to propofol. As indicated above, the formulation may exhibit one, two, three or all of these properties.

[0056] The composition of the present invention may be referred to as a formulation, host/guest composition, drug delivery system, medicament, anaesthetic or sedative as well as more descriptively such as an anaesthetic formulation or sedative formulation.

5 [0057] Another aspect of the present invention provides an anaesthetic or sedative formulation comprising a neuroactive steroid anaesthetic and a cyclodextrin or modified form thereof, the formulation exhibiting properties including being sterilizable, inducing reduced pain on intravenous administration, having a therapeutic index of greater than 5 and/or being storable in a plastic container.

10

[0058] More particularly, the present invention relates to an anaesthetic or sedative formulation comprising a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org 20599, Org 21465 and tetrahydrodeoxycorticosterone and  
15 a pharmacologically acceptable derivative, salt or pro-drug form thereof, the formulation exhibiting properties including being sterilizable, inducing reduced pain on intravenous administration and having a therapeutic index of greater than 5.

[0059] In an embodiment, the formulation is also storable in a plastic container.

20

[0060] The present invention extends to mixtures of two or more neuroactive steroid anaesthetic drugs such as a composition comprising alphaxalone and alphadolone and/or alphadolone acetate or their pharmacologically acceptable derivatives, salts or pro-drug forms.

25

[0061] A "pharmacologically acceptable derivative" is a derivative that still induces anaesthesia whilst not increasing adverse side effects. The term "derivative" includes deuterated derivatives where one or more hydrogen atoms are replaced with deuterium. This can lead to improved efficacy. Furthermore, the anaesthetic agents may be subject to  
30 alkylation, alkoxylation, acetylation and/or phosphorylation to generate other derivatives. Other types of derivatives include deuterated or tritiated or other labeled forms useful for monitoring and tracking the anaesthetic in the body. The terms "derivative" and "modified

form" are used interchangeably herein. Salts of alphadolone include alphadolone acetate. Reference to pro-drugs include transported pro-drugs.

5 [0062] In an embodiment, the cyclodextrin is a  $\beta$ -cyclodextrin or a modified form thereof such as but not limited to a sulfoalkyl ether dextrin. A particularly useful sulfoalkyl ether dextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin. Alkyl ether derivatives are also contemplated such as a sulfoalkyl ether-alkyl ether cyclodextrin. An example of an alkyl ether derivative is a sulfobutyl ether-alkyl ether cyclodextrin. Other cyclodextrins contemplated herein are listed in Table 7 and include methylated, hydroxyalkylated, 10 alkylated, branched, acylated and anionic derivatives.

[0063] Accordingly, an aspect of the present invention provides an anaesthetic or sedative delivery host/guest composition comprising a sulfoalkyl ether dextrin host or modified form thereof with a neuroactive steroid anaesthetic drug guest, the host/guest composition 15 formulated to be sterilizable, administrable by intravenous injection with minimal pain, exhibiting a therapeutic index of greater than 5. In an embodiment, the formulation may also be storable in a plastic container.

20 [0064] Another aspect of the present invention is directed to a drug delivery host/guest composition comprising a sulfoalkyl ether dextrin host or modified form thereof with a neuroactive steroid anaesthetic drug guest selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org 20599, Org 21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof, the host/guest 25 composition to be sterilizable, administrable by intravenous injection with minimal pain and exhibiting a therapeutic index of greater than 5.

[0065] In an embodiment, the composition is also storable in a plastic container.

30 [0066] Another aspect of the present invention provides an anaesthetic or sedative formulation comprising a neuroactive steroid anaesthetic and a sulfoalkyl ether dextrin or modified form thereof, the formulation exhibiting properties including being sterilizable,



inducing reduced pain on intravenous administration, having a therapeutic index of greater than 5 and/or being storable in a plastic container.

5 [0067] As indicated above one particularly useful sulfoalkyl ether dextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin. Of the properties exhibited, in a particular embodiment, the formulation exhibits two or more, three or more or all properties. These properties include imitating rapid induction of anaesthesia to surgical levels with similar or more rapid wakening time such as compared to propofol.

10 [0068] The formulation between the neuroactive steroid and cyclodextrin is generally in a molar ratio of from 1:1 to 1:6 (neuroactive steroid:cyclodextrin), more particularly about 1:1 to 1:4, even more particularly about 1:1 to 1:3 and still more particularly about 1:2. The range 1:1 to 1:6 includes 1:1, 1:1.1, 1:1.2, 1:1.3, 1:1.4, 1:1.5, 1:1.6, 1:1.7, 1:1.8, 1:1.9, 1:2, 1:2.1, 1:2.2, 1:2.3, 1:2.4, 1:2.5, 1:2.6, 1:2.7, 1:2.8, 1:2.9, 1:3, 1:3.1, 1:3.2, 1:3.3, 1:3.4, 15 1:3.5, 1:3.6, 1:3.7, 1:3.8, 1:3.9, 1:4, 1:4.1, 1:4.2, 1:4.3; 1:4.4, 1:4.5, 1:4.6, 1:4.7, 1:4.8, 1:4.9, 1:5, 1:5.1, 1:5.2, 1:5.3, 1:5.4, 1:5.5, 1:5.6, 1:5.7, 1:5.8, 1:5.9 and 1:6.

[0069] Accordingly, the present invention provides a drug delivery host/guest composition comprising a cyclodextrin host or modified form thereof with a neuroactive steroid drug 20 guest, wherein the molar ratio of neuroactive steroid to cyclodextrin is from about 1:1 to about 1:6 and wherein the composition is formulated to be sterilizable, administrable by intravenous injection with minimal pain and exhibiting a therapeutic index of greater than 5. In an embodiment, the formulation may also be storable in a plastic container.

25 [0070] More particularly, the present invention is directed to a drug delivery host/guest composition comprising a cyclodextrin selected from an  $\alpha$ -,  $\beta$ - or  $\gamma$ -cyclodextrin or a modified form thereof including a sulfoalkyl ether dextrin or sulfoalkyl ether-alkyl ether derivative or other derivatives listed in Table 7 and a neuroactive steroid drug guest selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone 30 (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org 20599, Org 21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof, wherein the molar ratio of a neuroactive steroid to cyclodextrin is from

about 1:1 to about 1:6 and wherein the composition is formulated to be sterilizable, administrable by intravenous injection with minimal pain and to exhibit a therapeutic index of greater than 5. In one embodiment, the (7) sulfobutyl ether  $\beta$ -cyclodextrin comprises less than 100ppm of a phosphate and has an absorption of less than 0.5AU due to a drug-degrading enzyme, as determined by UV/v is spectrophotometry at a wave length of 245nm to 270nm for an aqueous solution containing 300mg of the dextrin per ml of solution in a cell having a 1cm path length. In an embodiment, the formulation may also be storable in a plastic container.

10 [0071] The anaesthetic composition of the present invention may in one embodiment comprise a buffer such as a phosphate or tris or citrate phosphate buffer to maintain the pH from about 5.5 to about pH8. This includes pH values of 5.5, 6, 6.5, 7, 7.5 and 8. Alternatively, the composition does not comprise a buffer and the pH being from about pH3 to about pH 9.5 such as pH3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9 or 9.5.

15

[0072] In a further aspect, the formulations of the present invention also include one or more agents such as excipients and/or preservatives, microbial retardants. Other agents may also be included to reduce toxicity. Agents include, for example, EDTA, benzyl alcohol, bisulphites, monoglyceryl ester of lauric acid (Monolaurin), capric acid and/or its soluble alkaline salts or its monoglyceryl ester (Monocaprin), edetate, and capric acid and/or its soluble alkaline salts or its monoglyceryl ester (Monocaprin) and edentate. The formulation may also contain one or more co-polymers to assist in solubility or stability of the anaesthetic agent. Examples include hydroxy propyl methyl cellulose (HPMC), polyvinyl pyrrolidone (PVP) and/or carboxymethyl cellulose (CMC).

25

[0073] Conveniently, the neuroactive steroid anaesthetic is provided at a concentration of from about 0.5 to 100 mg/ml in a saline suspension comprising the cyclodextrin. Such a concentration includes 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and 100mg/ml of drug. As indicated above,

30

the composition is generally formulated so that the molar ratio of neuroactive steroid to cyclodextrin is from about 1:1 to about 1:6, particularly from about 1:1 to 1:4, even more particularly from about 1:1 to 1:3 and most particularly about 1:2.

5 [0074] Reference to any particular neuroactive steroid or their salts includes a racemic mixture of enantiomers of each anaesthetic as well as a single enantiomer of the agent.

[0075] In a particular embodiment, the neuroactive steroid is alphaxalone, alphadolone and/or alphadolone acetate. In one embodiment, alphaxalone is in the formulation at a  
10 concentration of from 1 to 100mg/ml such as 10mg/ml. In another embodiment, alphadolone or alphadolone acetate is present at 0.5 to 50mg/ml such as 3 mg/ml.

[0076] The formulations herein are for *in vivo* delivery meaning that the neuroactive steroid anaesthetic is delivered by intravenous, sub-cutaneous, intraperitoneal, intrathecal,  
15 intramuscular, intravitreal, transdermal, suppository (rectal), pessary (vaginal), inhalation, intranasal and the like. Most effectively, the formulation is an intravenous (iv) formulation.

[0077] Accordingly, another aspect of the present invention provides an injectable  
20 formulation of a neuroactive steroid anaesthetic selected to be sterilizable, administrable by intravenous injection with minimal pain, exhibiting a therapeutic index of greater than 5 and storable in a plastic container formulated with cyclodextrin, such as (7) sulfobutyl ether  $\beta$ -cyclodextrin or an alkyl ether derivative.

25 [0078] The neuroactive steroid anaesthetic may be used alone or in combination with another anaesthetic or sedative or other active agent. In one embodiment, alphaxalone is used with alphadolone or its salt, alphadolone acetate. Hence, reference to "alphadolone" includes alphadolone acetate. The composition may comprise, therefore alphaxalone or alphadolone alone or a combination of alphaxalone and alphadolone or any of their  
30 derivatives, salts or pro-drug forms.

[0079] Hence, in a particular embodiment, the present invention further provides a

composition comprising alphaxalone or a pharmacologically acceptable derivative, salt or pro-drug thereof and/or alphadolone or a pharmacologically acceptable derivative, salt or pro-drug thereof formulated with a sulfoalkyl ether dextrin, such as (7) sulfobutyl ether  $\beta$ -cyclodextrin wherein the molar ratio of alphaxalone and/or alphadolone to dextrin is from  
5 about 1:1 to about 1:6. Reference can conveniently be made to Remington's Pharmaceutical Sciences, Mack Publishing Company, Eaton, USA, 1990 and Rowe's Handbook of Pharmaceutical Excipients, 2009 for formulation methods and reagents.

[0080] The present invention contemplates inducing or maintaining by infusion or  
10 intermittent bolus administration, anaesthesia in a human subject, the method comprising administering an anaesthetic-effective amount of a neuroactive steroid anaesthetic formulated with a cyclodextrin, such as sulfoalkyl ether dextrin, for a time and under conditions to induce anaesthesia.

15 [0081] More particularly, the present invention provides a method of inducing or maintaining by infusion or intermittent bolus administration, anaesthesia in a human subject, the method comprising administering an anaesthetic-effective amount of a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone,  
20 Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof formulated with a cyclodextrin, such as (7) sulfobutyl ether  $\beta$ -cyclodextrin, for a time and under conditions sufficient to induce anaesthesia, wherein the anaesthetic formulation is sterilizable, administrateable by intravenous injection with minimal pain and exhibits a therapeutic index of greater than 5.  
25 Additionally, the formulation can initiate rapid anaesthesia with similar or more rapid wakening time compared with propofol or Althesin (Registered Trademark). The formulation may also be storable in a plastic container.

[0082] The present invention extends to inducing or maintaining by infusion or  
30 intermittent bolus administration, sedation. Hence, another aspect of the present invention provides a method of inducing or maintaining by infusion or intermittent bolus administration, sedation in a subject, the method comprising administering a sedation-

effective amount of a neuroactive steroid anaesthetic formulated with a cyclodextrin such as sulfoalkyl ether dextrin, for example, (7) sulfobutyl ether  $\beta$ -cyclodextrin, for a time and under conditions sufficient to induce sedation.

5 [0083] Reference to "(7) sulfobutyl ether  $\beta$ -cyclodextrin" includes methylated, hydroxyalkylated, branched, alkylated, acylated and anionic derivatives thereof such as a sulfobutyl ether-alkyl ether  $\beta$ -cyclodextrin. Other derives include  $\beta$ -cyclodextrin sulfobutyl ethers, ethyl ethers,  $\beta$ -cyclodextrin sulfobutyl ethers (flat),  $\gamma$ -cyclodextrin sulfobutyl ethers and  $\alpha$ -cyclodextrin sulfobutyl ethers and their salts (e.g. sodium salts).

10

[0084] As indicated above, a particular subject is a human subject.

[0085] The anaesthetic formulation may be packaged for sale with instructions for use. The use includes a patient management protocol comprising administering to the patient an effective amount of neuroactive steroid anaesthetic such as selected from alphaxalone, 15 alphadolone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof formulated with a cyclodextrin such as a sulfoalkyl ether dextrin, for example, (7) sulfobutyl ether  $\beta$ -cyclodextrin, for a time and under conditions sufficient to induce anaesthesia.

20

[0086] The present invention further contemplates the use of a neuroactive steroid anaesthetic and a cyclodextrin, such as a sulfoalkyl ether dextrin, for example, (7) sulfobutyl ether  $\beta$ -cyclodextrin, in the manufacture of a medicament to induce anaesthesia in a subject such as a human subject. In a particular embodiment, the neuroactive steroid 25 anaesthetic is selected from alphaxalone, alphadolone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof. In another embodiment, the anaesthetic is selected from acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof.

30

[0087] In terms of an anaesthetic-effective amount, this is generally around 0.25mg/kg to about 100 mg/kg body weight. A sedative-effective amount is provided in similar or lower

amounts and includes from about 0.05mg/kg to about 10mg/kg body weight.

5 [0088] The present invention further provides a kit. The kit may be in any form including a syringe or modified syringe. A kit may comprise alphaxalone and/or alphadolone or other neuroactive steroid anaesthetic or their derivatives, salts or pro-drug forms in one or more compartments and a sulfoalkyl ether dextrin in a further compartment as well as excipients in subsequent compartments. The contents of the compartments may be admixed prior to use.

10 [0089] In a particular embodiment, the present invention provides a formulation comprising alphaxalone and/or alphadolone and/or pharmacologically acceptable derivatives, salts or pro-drug forms thereof complexed with the sulfobutyl ether cyclodextrin, for use in inducing or maintaining by infusion or intermittent bolus administration, anaesthesia or sedation in human subject.

15 [0090] The anaesthetic forms may be labeled such as deuterated or tritiated forms or by other labels to facilitate monitoring and tracking of the anaesthetics in the body. Kits and apparatus are provided, therefore, to monitor labeled neuroactive steroid anaesthetics.

20 [0091] Whilst the present invention is particularly directed to anaesthetic formulations for use in humans, the formulations may also be used in animals such as for clinical trials or veterinary use. Non-human animals contemplated herein include rats, mice, guinea pigs, hamsters, sheep, pigs, dogs, cats, horses, cows, goats, camels and non-human primates.

25 [0092] Hence, the present invention provides an anaesthetic or sedative composition comprising a neuroactive steroid anaesthetic and a cyclodextrin or modified form thereof wherein the composition has the following properties:

- (i) the neuroactive steroid and cyclodextrin are formulated in a molar ratio of from about 1:1 to about 1:6;
- 30 (ii) the neuroactive steroid is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a

- pharmacologically acceptable derivative, salt or pro-drug form thereof formulated;
- (iii) the cyclodextrin is selected from an  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrin or a modified form thereof;
  - (iv) a buffer is optionally present and when present the pH of the formulation is from about pH5.5 to about pH8.0 and in the absence of buffer, the pH is from about pH3 to about pH9.5;
  - (v) the formulation is sterilizable;
  - (vi) intravenous injection of the formulation induces less pain than from a propofol formulation;
  - (vii) the therapeutic index of the formulation is greater than 5;
  - (viii) the formulation can be stored in a plastic container; and
  - (ix) the formulation can initiate rapid induction of anaesthesia to surgical levels with similar or more rapid waking time compared to propofol.

15 **[0093]** In a particular embodiment, an anaesthetic or sedative formulation is provided comprising a sulfoalkyl ether or sulfoalkyl ether-alkyl ether dextrin, a neuroactive steroid anaesthetic such as alphaxalone or alphadolone and one or more co-polymers such as HPMC, PVP and/or CMC.

20 **[0094]** In a particular embodiment, the neuroactive steroid anaesthetic is formulated with a sulfoalkyl ether dextrin such as (7) sulfobutyl ether  $\beta$ -cyclodextrin.

**[0095]** The present invention further contemplates a method for formulating an anaesthetic or sedative composition, the method generating a host/guest composition comprising a cyclodextrin and a neuroactive steroid. In an embodiment, the cyclodextrin is a sulfoalkyl ether or sulfoalkyl ether-alkyl ether dextrin such as (7) sulfobutyl ether  $\beta$ -cyclodextrin or sulfobutyl ether-alkyl ether  $\beta$ -cyclodextrin. Other cyclodextrins include  $\beta$ -cyclodextrin sulfobutyl ether-ethyl ether,  $\beta$ -cyclodextrin sulfobutyl ether (flat),  $\gamma$ -cyclodextrin sulfobutyl ether,  $\alpha$ -cyclodextrin sulfobutyl ether and their sodium salts.

30

**[0096]** Particular embodiments of the invention are:

1. An anaesthetic or sedative delivery host/guest composition comprising a cyclodextrin host or modified form thereof with a neuroactive steroid anaesthetic drug guest, the host/guest composition formulated to be sterilizable, administrable by intravenous injection with minimal pain and to exhibit a therapeutic index of greater than  
5 5.
2. The composition of Embodiment 1 wherein the neuroactive steroid anaesthetic is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and  
10 tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof.
3. The composition of Embodiment 1 or 2 wherein the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is from about 1:1 to 1:6.  
15
4. The composition of Embodiment 3 wherein the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is from about 1:1 to about 1:4.
5. The composition of Embodiment 4 wherein the molar ratio of neuroactive steroid  
20 anaesthetic to cyclodextrin is about 1:2.
6. The composition of any one of Embodiments 1 to 5 further comprising one or more of an antimicrobial agent, a preservative, a buffer and/or a co-polymer.
- 25 7. The composition of Embodiment 6 wherein the co-polymer is selected from hydroxy propyl methyl cellulose, polyvinyl pyrrolidone and carboxymethyl cellulose.
8. The composition of Embodiment 6 wherein if a buffer is present, the pH is from about pH 5.5 to about pH 8.  
30
9. The composition of any one of Embodiments 1 to 8 wherein the cyclodextrin is a  $\beta$ -cyclodextrin or a modified form thereof.



10. The composition of Embodiment 9 wherein the modified form of  $\beta$ -cyclodextrin is a sulfoalkyl ether cyclodextrin or a methylated, hydroxy alkylated, branched, alkylated, acylated or anionic derivative thereof.

5

11. The composition of Embodiment 10 wherein the sulfoalkyl ether  $\beta$ -cyclodextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin.

12. The composition of Embodiment 10 wherein an alkylated derivative is a sulfoalkyl ether-alkyl ether derivative.

10

13. The composition of any one of Embodiments 1 to 8 wherein the cyclodextrin is selected from  $\beta$ -cyclodextrin sulfobutyl ether-ethyl ether,  $\beta$ -cyclodextrin sulfobutyl ether (flat),  $\gamma$ -cyclodextrin sulfobutyl ether,  $\alpha$ -cyclodextrin sulfobutyl ether and their sodium salts.

15

14. The composition of Embodiment 1 wherein the composition initiates rapid induction of anaesthesia with similar or more rapid awakening time relative to propofol.

15. An anaesthetic or sedative composition comprising a neuroactive steroid selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-drug form thereof formulated with a sulfoalkyl ether dextrin or modified form thereof to generate a sterilizable composition with a therapeutic index of greater than 5.

25

16. The anaesthetic or sedative composition of Embodiment 15 wherein the neuroactive steroid anaesthetic is alphaxalone and/or alphadolone or pharmacologically acceptable derivatives including deuterated derivatives, salts and prodrug forms thereof.

30

17. The anaesthetic or sedative composition of Embodiment 15 or 16 wherein the sulfoalkyl ether dextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin or an alkyl ether derivative thereof.
- 5 18. The anaesthetic or sedative composition of Embodiment 17 wherein the dextrin is sulfobutyl ether-alkyl ether cyclodextrin.
19. The anaesthetic or sedation composition of Embodiment 15 to 16 wherein the cyclodextrin is selected from  $\beta$ -cyclodextrin sulfobutyl ether-ethyl ether,  $\beta$ -cyclodextrin  
10 sulfobutyl ether (flat),  $\gamma$ -cyclodextrin sulfobutyl ether,  $\alpha$ -cyclodextrin sulfobutyl ether and their sodium salts.
20. A method of inducing or maintaining by infusion or intermittent bolus administration, anaesthesia or sedation in a subject, the method comprising administering  
15 an anaesthetic- or sedative-effective amount of a composition of any of Embodiments 1 to 19.
21. The method of Embodiment 20 wherein the neuroactive steroid anaesthetic agent is alphaxalone, alphadolone and/or alphadolone acetate.  
20
22. The method of Embodiment 21 wherein alphaxalone and/or alphadolone is/are used at anaesthetic concentration of 0.25mg/kg to 100 mg/kg body weight.
23. The method of Embodiment 21 wherein the alphaxalone is used at a sedating  
25 concentration of 0.05 mg/kg to 10mg/kg body weight.
24. The method of any one of Embodiments 20 to 23 wherein the subject is a human.
25. Use of a neuroactive steroid anaesthetic and a cyclodextrin or a modified form  
30 thereof in the manufacture of a medicament to induce anaesthesia in a subject wherein the medicament is sterilizable, and has a therapeutic index of greater than 5.

26. Use of Embodiment 25 wherein the neuroactive steroid anaesthetic is selected from alphaxalone, alphadolone, acebrochol, allopregnanolone, eltanolone (pregnanolone), ganaxolone, hydroxydione, minaxolone, Org20599, Org21465 and tetrahydrodeoxycorticosterone and a pharmacologically acceptable derivative, salt or pro-  
5 drug form thereof.

27. Use of Embodiment 25 or 26 wherein the cyclodextrin is sulfoalkyl ether dextrin.

28. Use of Embodiment 27 wherein the sulfoalkyl ether dextrin is (7) sulfobutyl ether  
10  $\beta$ -cyclodextrin or an alkyl ether derivative thereof.

29. Use of Embodiment 28 wherein the dextrin is a sulfobutyl ether-alkyl ether cyclodextrin.

15 30. Use of Embodiment 25 or 26 wherein the cyclodextrin is selected from  $\beta$ -cyclodextrin sulfobutyl ether-ethyl ether,  $\beta$ -cyclodextrin sulfobutyl ether (flat),  $\gamma$ -cyclodextrin sulfobutyl ether,  $\alpha$ -cyclodextrin sulfobutyl ether and their sodium salts.

31. A method of inducing or maintaining by infusion or intermittent bolus  
20 administration, sedation in a human subject, said method comprising administration to said subject a sedating-effective amount of a neuroactive steroid anaesthetic selected from alphaxalone, alphadolone and pharmacologically acceptable derivatives, salts and pro-drug forms thereof formulated with a sulfoalkyl ether dextrin or modified form thereof.

25 32. The method of Embodiment 31 wherein the sulfoalkyl ether dextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin or an alkyl ether derivative thereof.

33. The method of Embodiment 32 wherein the dextrin is a sulfobutyl ether-alkyl ether cyclodextrin.

34. The method of Embodiment 31 wherein the sulfoalkyl ether dextrin is selected from  $\beta$ -cyclodextrin sulfobutyl ether-ethyl ether,  $\beta$ -cyclodextrin sulfobutyl ether (flat),  $\gamma$ -cyclodextrin sulfobutyl ether,  $\alpha$ -cyclodextrin sulfobutyl ether and their sodium salts.

5 35. The use of Embodiment 25 or method of Embodiment 31 wherein the subject is a human.

[0097] The present invention is further described by the following non-limiting Examples. When a neuroactive steroid anaesthetic is formulated with a cyclodextrin, it is referred to  
10 as "neuroactive steroid anaesthetic<sub>CD</sub>". An example is Phaxan<sub>CD</sub>, which comprises alphaxalone formulated with a cyclodextrin which in this case is (7) sulfobutyl ether  $\beta$ -cyclodextrin. Other examples include pregnanolone<sub>CD</sub> and alphadolone<sub>CD</sub>

**EXAMPLE 1**

*Anaesthetic effects of alphaxalone in 30% w/v (7) sulfobutyl ether  $\beta$ -cyclodextrin*

[0098] Alphaxalone was formulated as 6ml clear colorless liquid containing:

- 5 • Alphaxalone 60mg (10mg/ml);
- (7) sulfobutyl ether  $\beta$ -cyclodextrin 1800mg;
- Saline (0.9% w/v) 6ml.

[0099] This is a molar complexation ratio of alphaxalone to (7) sulfobutyl ether  $\beta$ -  
 10 cyclodextrin of 1:4.6. Male Wistar rats (weight [wt] 270-315g) with indwelling jugular  
 intravenous catheters were put in a Perspex restrainer and given the injections with the  
 attached observations upon release from the restrainer shown in Table 1.

**TABLE 1**

15

**Effects of alphaxalone formulation on Wistar rats**

TIME MINUTES	PRC447	PRC 446	PRC 468	PRC 461	PRC 448	PRC 469	PRC 455
	C 5	C 10	C 15	C 20	C 20	C 25	C 100
0	s					S	TP-
1	s			RR-	TP-	S	TP-
2	s	s	S	RR-	TP-	RR-	TP-
3	s	s	S	TP-	TP-	RR-	TP-
4	s	RR-	RR-	TP-	TP-	RR-	TP-
5	REC	RR-	RR-	TP-	TP-	RR-	TP-
6		RR-	RR-	TP-	TP-	RR-	TP-
7		RR+	RR-	TP-	TP-	RR-	TP-
8		s	RR-	TP-	TP-	RR-	TP-
9		s	RR-	TP-	TP-	RR-	TP-
10		s	RR-	TP-	TP-	RR-	TP-
11		s	RR-	TP-	TP+	RR+	TP-
12		s	RR+	TP+	RR-	RR+	TP-
13		s	S	RR-	RR-	RR+	TP-
14		s	S	RR-	S	RR+	TP-

KEY

S	sedated
RR+	returned righting reflex
RR-	lost righting reflex
TP+	normal tail pinch response
TP-	none or minimal response to tail pinch
REC	recovered
	no sedation
C = cyclodextrin formulation [5, 10, 15, 20, 25 and 100 mg/kg]	

[0100] The rats given 25 and 100 mg/kg body weight had recovered by 60 minutes; they  
5 did not die or suffer any adverse effects at these doses. In these experiments, it can be seen  
that intravenous injection of alphaxalone 10mg/ml dissolved in (7) sulfobutyl ether  $\beta$ -  
cyclodextrin caused loss of consciousness at doses above 10mg/kg and there was a dose  
related anaesthetic effect with a wide safety margin; rats that received 25 and 100 mg/kg  
body weight (two times and ten times the anaesthetic dose) did not die indicating a wide  
10 safety margin for the preparation. This is clearly different from a formulation of  
alphaxalone in hydroxypropyl  $\beta$ -cyclodextrin. Such a preparation (AlfaxanCD-RTU)  
made by Jurox Pty, Newcastle NSW Australia has a published LD<sub>50</sub>, the dose of  
alphaxalone that causes death in 50% of rats when given intravenously, of 19mg/kg body  
weight, a figure very significantly below that shown here for alphaxalone formulated in  
15 sulfobutyl ether  $\beta$ -cyclodextrin [Alfaxan CD-RTU Material safety Data Sheet; Jurox Pty,  
Newcastle NSW Australia].

**EXAMPLE 2**

***Pharmacokinetics in the rat***

[0101] Two groups of ten rats with implanted internal jugular intravenous and carotid  
5 intraarterial catheters receive bolus intravenous injections *via* the jugular vein of 10mg/kg  
body weight of (7) sulfobutyl ether  $\beta$ -cyclodextrin formulation of alphaxalone (n=10 rats)  
or a mixture of alphaxalone and alphadolone in CremophorEL (a polyethoxylated Caster  
oil), 1.1 ml/kg (n=10 rats). Blood taken from the carotid artery or tail at a number of time  
intervals after this injection is analyzed for alphaxalone blood levels. These are fitted to a  
10 three compartment pharmacokinetic model and mean  $\pm$  sem for key parameters are  
calculated for both preparations of steroid anaesthetic.

[0102] It is expected that there will be no significant difference between the PK parameters  
calculated for the (7) sulfobutyl ether  $\beta$ -cyclodextrin formulation and the Althesin  
15 (Registered Trademark) formulation. This would indicate that the new formulation is  
expected to behave in a similar fashion to the way Althesin (Registered Trademark) has  
behaved in the past, particularly with respect to doses needed for anaesthesia and sedation  
and also the speed of recovery. A sample table of blood levels that can be expected from  
these experiments appears in Table 2.

20

**TABLE 2**

<b>Variable</b>	<b>Units</b>	
Duration of Anaesthesia	Min	35.0 [9.0]
AUC	Min*mg/L	58.6 [11.7]
$t^{1/2}_{(elim)}$	Min	38.2 [5.6]
Cl <sub>T</sub>	mL/min/kg	38.0 [8.2]
V <sub>c</sub>	L/kg	0.5 [0.3]
V <sub>dss</sub>	L/kg	1.8 [0.6]

### EXAMPLE 3

#### *Anaesthetic effects of Alphaxalone in (7) sulfobutyl ether $\beta$ -cyclodextrin compared with alphaxalone as Althesin (Registered Trademark) and propofol*

- 5 [0103] Male Wistar rats (wt 150-220 g weight) were implanted with indwelling internal jugular vein intravenous catheters under halothane anaesthesia. Twenty four hours later each rat received an intravenous injection from a range of doses of either: propofol (10 mg/ml in 10% w/v Intralipid emulsion; Diprivan [Registered Trademark]); Althesin [Registered Trademark] (alphaxalone 9mg/ml plus alphadolone acetate 3 mg/ml in 20%  
10 w/v CremophorEL); or Phaxan<sub>CD</sub> (alphaxalone 10mg/ml in a 1:2 molar complexation ratio with Captisol (Registered Trademark) - (7) sulfobutyl ether  $\beta$ -cyclodextrin). The following were assessed at regular time intervals after the intravenous injection:
1. righting reflex: scored as: 1 normal; 2 slow; 3 some attempt; 4 none – this was a measure of onset and duration of unconsciousness;
  - 15 2. tail pinch response: scored as: 1 normal; 2 weak; 3 just present; 4 none – this was a measure of onset and duration of surgical anaesthesia; and
  3. time to when the rat was able to walk on the rotarod (a rotating cylinder) measured in seconds: the maximum normal run time is 120 s in non sedated rats – attaining this value was a measure of time taken to attain full recovery from the sedating  
20 effects of the anaesthetic injections.

[0104] Results from groups of ten rats treated with the same anaesthetic and dose were combined for statistical purposes. Rats that attained a score of 4 for loss of righting reflex were deemed to have lost consciousness and those that scored a 4 for loss of tail pinch  
25 response were deemed to be surgically anaesthetized. The number of rats in each group of 10 similarly treated animals that scored 4 were subjected to probit regression analysis using SPSS Statistics 18 to produce graphs of probit value v log dose (probit plot) and also to calculate the estimated dose that caused anaesthesia in 50 and 95% of subjects ( $AD_{50}$  and  $AD_{95}$ , respectively) for unconsciousness (righting reflex measurements) and surgical  
30 anaesthesia (tail pinch responses). The rotarod walking times were also plotted for each dose and treatment. This was used as a measurement of complete recovery. The results



are shown in Figures 1a through 1f, rotarod performance for each anaesthetic, n = 10 rats per dose.

[0105] Table 3 below summarises the results of this series of experiments. It can be seen  
5 that Phaxan<sub>CD</sub> is equipotent with Althesin (Registered Trademark) in causing  
unconsciousness and surgical anaesthesia and both are more potent than propofol in this  
respect. Recovery from unconsciousness caused by Phaxan<sub>CD</sub> is just as fast as with  
propofol. However, recovery from Phaxan<sub>CD</sub> is slightly slower than propofol but faster  
10 than Althesin (Registered Trademark) if the depth of anaesthesia is taken to a surgical  
level. Control experiments revealed that the vehicles given alone intravenously, 20%  
CremophorEL, 10% Intralipid and 13% Captisol had no sedating or anaesthetic effects.

TABLE 3

	Althesin (Registered Trademark)	Phaxan <sup>ca</sup>	propofol
minimum dose causing all 10 rats to lose righting reflex mg/kg	5	5	3.0
ED <sub>50</sub> dose for loss of righting reflex mg/kg	2.55	2.79	4.53
ED <sub>95</sub> dose for loss of righting reflex mg/kg	4.39	4.26	8.4
minimum dose causing all 10 rats to lose tail pinch response mg/kg	15	15	15
ED <sub>50</sub> dose for loss of tail pinch response mg/kg	5.88	5.58	8.4
ED <sub>95</sub> dose for loss of tail pinch response mg/kg	14.05	13.55	18.46
ANOVA with Tukey post hoc			
duration of loss of righting reflex at dose causing all 10 rats to lose righting reflex (minutes) mean =	3.5	3.3	3.5
SD =	2.18	0.84	1.15
not significant ANOVA p = 0.0527			
duration of loss of tail pinch response at dose causing all 10 rats to lose tail pinch response (minutes) mean =	13.45	4.85	4.05
SD =	3.9	2.55	1.28
Althesin vs Phaxan <sup>ca</sup> *** p<0.001; Althesin vs propofol *** p<0.001; Phaxan <sup>ca</sup> vs propofol ns p>0.05			
time (minutes) to overall (complete) recovery of rotarod performance after minimum anaesthetic dose (lost RR) mean =	19.9	17	18
SD =	3.93	2.16	1.89
Althesin vs Phaxan <sup>ca</sup> ns p>0.05; Althesin vs propofol * p<0.05; Phaxan <sup>ca</sup> vs propofol ns p>0.05			
time (minutes) to overall (complete) recovery of rotarod performance after minimum surgical anaesthetic dose (lost tail pinch) mean =	39	32.5	23.1
SD =	3.16	3.54	3.25
Althesin vs Phaxan <sup>ca</sup> *** p<0.001; Althesin vs propofol *** p<0.001; Phaxan <sup>ca</sup> vs propofol *** p<0.001			

[0106] It can be concluded from this set of experiments that:

- Phaxan<sub>CD</sub> is an effective intravenous anaesthetic causing fast onset of general anaesthesia after intravenous injection.
- It is equipotent with Althesin and twice as potent as propofol.

5

#### EXAMPLE 4

##### *Lethal dose finding for alphaxalone anaesthetic preparations*

[0107] This series of experiments was undertaken to determine the LD<sub>50</sub> and LD<sub>95</sub> doses for alphaxalone formulated in CremophorEL [Althesin (Registered Trademark)] and (7) sulfobutyl ether  $\beta$ -cyclodextrin (Phaxan<sub>CD</sub>); i.e., the doses of alphaxalone in Captisol (Registered Trademark) that caused 50% and 95% of subjects to die. Male Wistar rats (wt 150-220 g weight) were implanted with indwelling internal jugular vein intravenous catheters under halothane anaesthesia. Twenty four hours later each rat received an intravenous injection from a range of doses of either: Althesin (Registered Trademark) (alphaxalone 9mg/ml plus alphadolone acetate 3 mg/ml in 20% CremophorEL); or Phaxan<sub>CD</sub> (alphaxalone 10mg/ml in a 1:2 molar complexation ratio with Captisol (Registered Trademark) - (7)sulfobutyl ether  $\beta$ -cyclodextrin). The number of rats that died was recorded for each group of 10 rats given the same dose of drug. Results from groups of ten rats treated with the same anaesthetic and dose were combined for statistical purposes. The graph of the raw data is shown in Figure 2 [% rats died in each dosage group v dose].

[0108] At doses of alphaxalone between 50 and 60mg/kg all the rats in the Althesin (Registered Trademark) groups died whereas none died in the Phaxan<sub>CD</sub> groups that received the same doses of alphaxalone. The lethality values for Phaxan<sub>CD</sub> showed a ceiling at 20%; no more than 20% rats died even if the dose of alphaxalone administered as Phaxan<sub>CD</sub> were increased. The percentages of rats that died in all the different dosage treatment groups were subjected to a probit regression analysis using SPSS Statistics 18 and the probit values were plotted on a graph against the log dose of anaesthetic; this is called a probit plot. This is shown in Figure 3.

30

[0109] The probit plot for Althesin (Registered Trademark) was used to calculate the dose of alphaxalone in this formulation that caused death in 50% and 95% rats; the LD<sub>50</sub> and LD<sub>95</sub> respectively). These values were 43.6 mg/kg [LD<sub>50</sub>] and 51.5 mg/kg [LD<sub>95</sub>]. As the dose of alphaxalone was increased the number of rats that died increased in proportion  
5 when the alphaxalone was given as Althesin (Registered Trademark). By contrast there was a ceiling effect of the lethality of the alphaxalone in (7) sulfobutyl ether  $\beta$ -cyclodextrin (Phaxan<sub>CD</sub>) formulation. The alphaxalone was much less toxic as assessed by lethality compared with alphaxalone formulated with CremophorEL (Althesin [Registered Trademark]). A dose of 52mg/kg alphaxalone as Althesin (Registered Trademark) caused  
10 all 10 rats in that group to die but 64mg/kg alphaxalone caused no deaths in the 10 rats which received that dose of alphaxalone formulated with (7) sulfobutyl ether  $\beta$ -cyclodextrin (Phaxan<sub>CD</sub>). Furthermore, unlike the probit plot for Althesin (Registered Trademark), which showed a direct proportional relationship of increasing lethality with increasing dose, alphaxalone formulated in (7) sulfobutyl ether  $\beta$ -cyclodextrin (Phaxan<sub>CD</sub>)  
15 showed a ceiling effect for lethality; when the dose of alphaxalone in this preparation was increased to 71, 78 and then 84 mg/kg only 20% of rats died in each group. Thus, it was not possible to find the dose of alphaxalone in this formulation that caused death in 50% and 95% rats (the LD<sub>50</sub> and LD<sub>95</sub>, respectively). In any event, both of these values are greater than 84mg/kg which is more than double the equivalent values for Althesin  
20 (Registered Trademark) and four times the value of the LD<sub>50</sub> for alphaxalone formulated in hydroxypropyl  $\beta$ -cyclodextrin manufactured by Jurox [Alfaxan CD-RTU Material Safety Data Sheet. Jurox Pty, Newcastle NSW Australia].

[0110] These results make the therapeutic index (ratio of dose that causes death in 50%  
25 subjects (LD<sub>50</sub>) divided by the dose that causes anaesthesia in 50% subjects (the AD<sub>50</sub>) to be 14.8 for Althesin (Registered Trademark) and >30.2 for alphaxalone formulated in (7) sulfobutyl ether  $\beta$ -cyclodextrin (Captisol [Registered Trademark] Phaxan<sub>CD</sub>). This difference is not due to differences in toxicity of the excipients. Table 4 below shows the results of experiments in 10 rats with indwelling jugular intravenous catheters. Five rats  
30 were given a 20% solution of Cremophor EL intravenously and another five rats were given a solution of (7) sulfobutyl ether  $\beta$ -cyclodextrin, both being administered at a dose and volume equal to that administered in the experiments above at the highest dose of

alphaxalone. Neither excipient caused death in any rat indicating that the difference in the safety/lethality of the two formulations of alphaxalone was not due to dose related toxicity of the excipients.

5

**TABLE 4**

n/10 died	vehicle	dose=ml/kg
0	13% Captisol	9.0
0	20% Cremophor EL	4.6

### EXAMPLE 5

#### *Demonstration of the limitation of alphaxalone toxicity by Captisol [Registered Trademark] (sulfobutyl ether $\beta$ -cyclodextrin)*

5

[0111] Since the potency in causing anaesthesia is the same for a bolus dose of alphaxalone whether given in 20% CremophorEL (Althesin [Registered Trademark]) or in (7) sulfobutyl ether  $\beta$ -cyclodextrin (Phaxan<sub>CD</sub>), then the ceiling effect for toxicity must be related to the (7) sulfobutyl ether  $\beta$ -cyclodextrin when higher doses of alphaxalone in the (7) sulfobutyl ether  $\beta$ -cyclodextrin formulation are administered. Such a property has not been described for intravenous anaesthetics before. Furthermore, this property has not been described before for alphaxalone formulated in other cyclodextrins.

[0112] In order to test whether the low toxicity of Phaxan<sub>CD</sub> was due to the Captisol excipient, twenty rats with indwelling jugular intravenous catheters were divided into two groups of 10 rats each. They were all given intravenous injections of alphaxalone formulated in 20% Cremophor at a dose that had, in previous experiments reported in example 4, caused all rats to die (Althesin [Registered Trademark]; alphaxalone dose 52.5mg/kg iv – this equals 16 times the AD<sub>95</sub> for Althesin (Registered Trademark) at which it is expected a very high proportion or all rats will die). Sixty seconds before the Althesin (Registered Trademark) injection a premedication injection was given:

- group 1 (10 rats) received 5.3mls/kg 0.9% sodium chloride solution 60 seconds before 52.5mg/kg alphaxalone as Althesin (Registered Trademark);
- group 2 (10 rats) received 5.3mls/kg 13% solution of (7) sulfobutyl ether  $\beta$ -cyclodextrin in 0.9% sodium chloride solution 60 seconds before 52.5mg/kg alphaxalone as Althesin (Registered Trademark).

[0113] The number of rats that died in each group was recorded shown in Table 5. All 20 rats were anaesthetized by the injection of 52.5mg/kg alphaxalone as Althesin (Registered Trademark). However, the presence of (7) sulfobutyl ether  $\beta$ -cyclodextrin caused a statistically and clinically significant reduction in the mortality caused by the alphaxalone.

TABLE 5

<b>ALL RATS THAT SURVIVED WERE ANAESTHETISED &gt; 1 HOUR</b>		
premed given 60s before Althesin @ 16x AD95		
premed volume = 5.25 ml/kg	n died	n did not die
SALINE PREMED	8	2
SBECD7 PREMED	2	8

Study Of The Effect of Captisol on the Death Rate of Althesin

Fisher's Exact Test

The two-sided P value is 0.0230, considered significant.  
The row/column association is statistically significant.

[0114] This is a much unexpected result.

## EXAMPLE 6

### *The effect on sleep times of repeated dosing with Phaxan<sub>CD</sub>*

[0115] The mechanism responsible for the (7) sulfobutyl ether  $\beta$ -cyclodextrin in causing  
5 the ceiling effect on alphaxalone toxicity is unknown. It is known that alphaxalone is very  
poorly soluble in water and thus the vast majority of alphaxalone molecules are complexed  
with the cyclodextrin molecules in the ratio of 1:2 (the complexation ratio). It is known  
that some of the alphaxalone will dissociate from the cyclodextrin complex when  
Phaxan<sub>CD</sub> is injected intravenously. The question posed by the unique property of the  
10 alphaxalone sulfobutyl ether cyclodextrin complex exhibiting a ceiling effect for toxicity is  
whether this is caused by limiting the amount of alphaxalone released from the complex or  
whether there occurs a “mopping up” of alphaxalone molecules that might otherwise  
penetrate the brain to cause toxicity, by excess uncomplexed cyclodextrin molecules freed  
up by alphaxalone metabolism by the liver. The effect of the latter would be predicted to  
15 cause a progressive decrease in the level of free alphaxalone as the concentration of  
uncomplexed sulfobutyl ether cyclodextrin increased as a result of:

- liver metabolism of the alphaxalone;
- more doses of Phaxan<sub>CD</sub> being administered thereby making more uncomplexed cyclodextrin available.

20

[0116] If this were the case one would predict tolerance to repeated doses of alphaxalone  
formulated in (7) sulfobutyl ether  $\beta$ -cyclodextrin i.e., repeated injections of Phaxan<sub>CD</sub>  
would cause progressively less sedation and anaesthetic effect.

25 [0117] In order to test this five rats with indwelling jugular intravenous catheters were  
given repeated injections of the minimum dose of alphaxalone in (7) sulfobutyl ether  $\beta$ -  
cyclodextrin ( 5mg/kg Phaxan<sub>CD</sub>) that had caused 10/10 rats to be anaesthetized as judged  
by complete loss of righting reflex. The time was measured for each rat to begin recovery  
of righting reflex (progress from a score of 4 to a score of 3 in the righting reflex as  
30 described in example 2 above) and then another dose of 5mg/kg Phaxan<sub>CD</sub> was given. The  
time to the beginning of recovery of the righting reflex after the second dose of anaesthetic  
was measured and then another dose of 5mg/kg Phaxan<sub>CD</sub> was administered iv and the



process was repeated eight more times. The progressive recovery times for the first to the tenth doses are shown in the histogram (Figure 4) and table 6 below as means (sem) for those five rats. It can be seen that there was a significant progressive increase in recovery times after each of the first four doses. Further doses 5-10 caused no further significant increase and more importantly, no decrease, in sleeping time. These results indicate that the alphaxalone toxicity ceiling effect caused by the (7)sulfobutyl ether  $\beta$ -cyclodextrin is probably due to the controlled release of the alphaxalone from the complex at a rate sufficient to cause anaesthesia but no greater than that, no matter how much of the complex is given. This is a unique balance between the chemical relationship of the alphaxalone and the (7) sulfobutyl ether  $\beta$ -cyclodextrin, the low solubility of the alphaxalone in water environments, including biological fluids, the amount of alphaxalone needed to penetrate the brain to cause anaesthesia and the pharmacokinetics of alphaxalone. The results are shown in Figure 4 and Table 6. None of this was expected nor could have been predicted from prior art.

15

**TABLE 6**

dose number	sleep time (mins)	
	mean	sem
first	0.8	0.14
second	2.5	0.35
third	2.9	0.41
fourth	3.2	0.45
fifth	3.7	0.38
sixth	4.1	0.21
seventh	4.1	0.21
eighth	4.2	0.14
ninth	4.3	0.34
tenth	4.2	0.14

[0118] One way ANOVA [Tukey Kramer post hoc] applied to the data of sleep times after X 10 repeated dosing of 5 rats with Phaxan<sub>CD</sub> 5mg/kg revealed that there was a progressive

statistically significant increase in sleeping time from the first to the second, second to the third and third to the fourth but no increase in sleeping times thereafter. Furthermore it is important to note that there was not a progressive decrease in sleeping times that would be predicted if the mechanism for low toxicity and the ceiling effect of Phaxan<sub>CD</sub> was due to uncomplexed “free” sulfobutyl ether cyclodextrin [Captisol (Registered Trademark)] mopping up alphaxalone from the blood. If that were the case then the amount of free uncomplexed Captisol would increase as more doses were given and as alphaxalone was metabolized by the liver so leading to a progressive increase in free uncomplexed Captisol as the repeated exposure experiment progressed. If the alphaxalone were mopped up by this then one would expect less sleeping time with each successive dose. On the contrary the sleep time increased with each of the first four doses and then remained constant thereafter with each subsequent dose.

**EXAMPLE 7**

***Ceiling toxicity***

[0119] The following assumption is made:

5

A. that intravenous injection of Althesin (Registered Trademark) leads to an instantaneous dispersal of alphaxalone in the plasma but, since anaesthesia is caused by drug injection in one circulation and also alphaxalone is cleared from the blood by the liver on first pass, the level achieved by the mixing in the plasma will only reach 30% of the theoretical maximum assuming instantaneous mixing.

10

[0120] The following is considered:

1. alphaxalone is soluble in water to 0.03mg/ml;
2. alphaxalone is 35% protein bound in plasma;
- 15 3. plasma volume is 31ml/kg in rat (Davies and Morris, *Pharmaceutical Research*, 10(7):1093-95, 1993);
4. both Althesin (Registered Trademark) and Phaxan<sub>CD</sub> have alphaxalone concentrations of 10mg/ml;
5. for induction of anaesthesia Althesin (Registered Trademark) and Phaxan<sub>CD</sub> are
- 20 equipotent; a minimum of 4.3mg/kg alphaxalone by either preparation cause sleep in most (95%) rats;
6. after an initial bolus injection a drug preparation will equilibrate with the plasma volume during the first sleep cycle but later the drug will disperse into the extracellular fluid [ECF] which is = 297 ml/kg (Davies and Morris, 1993*supra*);
- 25 7. Captisol is distributed to ECF and restricted to this space; and
8. Alphaxalone is only released from the Captisol into an aqueous environment if the level of free alphaxalone in the aqueous environment is less than saturation i.e., < 0.03mg/ml [fact 1].

30 [0121] It is proposed herein that:

- I. From A and 5 above, the plasma level of drug needed to cause sleep = dose administered as Althesin/plasma volume =  $4.3/31 = 0.14\text{mg/ml}$ .

- II. Applying A, the plasma concentration is expected to be 30% of this when the blood has circulated to mix the drug effectively = 0.046mg/ml.
- III. From this, the level of free unbound alphaxalone in the plasma associated with anaesthesia induction after a single iv bolus injection = 65% of total (from fact 2)  
5 =  $0.045 \times 0.65 = 0.03\text{mg/ml}$ .
- IV. Proposition III above is exactly the known solubility of alphaxalone in water.
- V. Combining points 5 and 8 with proposition IV, it is proposed herein that the first induction dose of Phaxan<sub>CD</sub> caused anaesthesia by releasing all of the alphaxalone from the complex just achieving the anaesthetic level and free alphaxalone  
10 saturation level.
- VI. When the second dose of anaesthetic was administered in Example 9 the rat was starting to recover from the anaesthetic because some free alphaxalone had been metabolized by the liver, some free alphaxalone had been redistributed to the ECF and some of the Captisol containing alphaxalone had also redistributed to the ECF.  
15 Thus, the free alphaxalone level fell and alphaxalone left the brain so causing awakening. Thus, a further dose was given. Unlike the first dose, there was still alphaxalone in the blood so only some of the alphaxalone was released from the complex to bring the free alphaxalone level back to 0.03mg/ml; the brain is reloaded and sleep follows.
- 20 VII. However, sleep follows for a longer time after the second dose and the third and also the fourth until the ECF is loaded with 0.03mg/ml alphaxalone and alphaxalone/Captisol complex [points 6 and 7]. After that further doses of Captisol merely top up the blood level of free alphaxalone and maintain it until hepatic metabolism has cleared that dose of alphaxalone.
- 25 VIII. Once the level of free alphaxalone reaches 0.03mg/ml, the brain is anaesthetized. The brain will only absorb more alphaxalone if the level of free alphaxalone increases. This is possible with Althesin (Registered Trademark) and death ensues when 15 times the anaesthetic dose of that drug is administered as a bolus leading to a theoretical free alphaxalone level of 0.45mg/ml. In contrast, when that dose of  
30 alphaxalone is given as Phaxan<sub>CD</sub>, in a complex with Captisol (Registered Trademark), the alphaxalone is not released from the complex once the level of free alphaxalone reaches 0.03mg/ml [point 1]. This explains the ceiling effect on

lethality with alphaxalone when formulated in 13% sulfobutyl-7-ether  $\beta$ -cyclodextrin.

- IX. Since no further increase in sleep time occurs after the fifth dose of 5mg/kg Phaxan<sub>CD</sub> and subsequent doses of the same magnitude, then the clearance of alphaxalone must be in equilibrium with the dose administration rate. The equilibrium clearance of alphaxalone is therefore 5mg/kg/4.2min. Since the concentration in plasma is approximately 0.04mg/ml then the plasma clearance rate at equilibrium =  $((5 \div 4.2) \div 0.04) = 30\text{mls/kg/min}$ . This is within known values of hepatic blood flow (Davies and Morris, 1993 *supra*). It is well known that alphaxalone is primarily cleared from plasma by first pass hepatic metabolism.
- X. The latter implies that this particular formulation of alphaxalone in sulfobutyl ether  $\beta$ -cyclodextrin controls the level of free unbound anaesthetic that penetrates the brain up to but not above a level that causes anaesthesia; the Captisol cannot release any more compound above the level at which plasma water is saturated with alphaxalone and thus higher blood levels that would cause toxicity are not achieved.

[0122] This property has not been described for intravenous anaesthetics or intravenous cyclodextrins before. It arises from a unique set of circumstances not described or previously discovered:

1. A unique host:guest interaction. The evidence for this is the fact that the same guest [alphaxalone] formulated in another host cyclodextrin (Alfaxan<sub>CD</sub>-RTU; hydroxypropyl  $\beta$ -cyclodextrin - the Jurox preparation) does not have a ceiling on toxicity with a quoted LD<sub>50</sub> of 19mg/kg iv in rats, a dose which is 75% less than the dose of alphaxalone in the (7) sulfobutyl ether  $\beta$ -cyclodextrin which only causes 20% lethality;
2. The guest is a compound that causes anaesthesia at a free drug level which is equal to its aqueous solubility; and
3. The guest is a compound that has a high therapeutic index so that the free drug level is well below the toxic level.

## EXAMPLE 8

### *Pregnanolone formulation*

[0123] The neuroactive steroid anaesthetic pregnanolone was mixed with 13% w/v  
5 sulfobutyl ether  $\beta$ -cyclodextrin (Captisol [Registered Trademark]) in 0.9% saline to form  
pregnanolone<sub>CD</sub>. The pregnanolone dissolved incompletely at a concentration of 10mg/ml,  
and unlike alphaxalone, it only went into solution after 4 hours of continuous stirring. The  
solution was opalescent. This observation indicates that all neuroactive steroids do not  
interact with (7) sulfobutyl ether  $\beta$ -cyclodextrin in the same way. Fifteen male Wistar rats  
10 (150-200g weight) with surgically-implanted internal jugular intravenous catheters were  
used for these experiments in which they received intravenous pregnanolone<sub>CD</sub> injections:  
2.5 mg/kg (n=5); 5 mg/kg (n=5); and 10mg/kg (n=5).

[0124] They were assessed for anaesthesia by the righting reflex which was scored: 1  
15 normal; 2 slow; 3 some attempt; 4 none. A score of 4 means that a state of  
unconsciousness (anaesthesia) has been achieved. Figure 5 below shows the results from  
this test for the rats in the three groups that received 2.5, 5 and 10 mg/kg pregnanolone.  
Results shown are means of the readings from all 5 rats at each time point after intravenous  
injection of pregnanolone.

20

[0125] The rats were also assessed for surgical anaesthesia using tail pinch responses  
scored: 1 normal; 2 weak; 3 just present; 4 none. A score of 4 indicates surgical  
anaesthesia has been achieved. Figure 6 shows the results from this test for the rats in the  
three groups that received 2.5, 5 and 10 mg/kg pregnanolone<sub>CD</sub>. Results shown are means  
25 of the readings from all 5 rats at each time point after intravenous injection of  
pregnanolone<sub>CD</sub>.

[0126] Time to complete recovery from the sedating effects of the pregnanolone<sub>CD</sub> was  
assessed with the rotarod run time – normal non-sedated rats run on the accelerating  
30 rotating drum is 120 seconds; rats are fully recovered when they can walk on the rotarod  
treadmill for 120 seconds. Figure 7 below shows the results from this test for the rats in the  
three groups that received 2.5, 5 and 10 mg/kg pregnanolone<sub>CD</sub>. Results shown are means

of the readings from all 5 rats at each time point after intravenous injection of pregnanolone<sub>CD</sub>.

### **Conclusions**

- 5 [0127] Pregnanolone<sub>CD</sub> is an intravenous anaesthetic but of long duration. It causes anaesthesia induction immediately after intravenous injection. This effect is dose related and it is possible to cause sufficient CNS depression to lead to surgical anaesthesia.

## EXAMPLE 9

### *Alphadolone formulation*

[0128] The neuroactive steroid anaesthetic alphadolone was mixed with 13% w/v  
5 sulfobutyl ether  $\beta$ -cyclodextrin (Captisol) in 0.9% w/v saline to form alphadolone<sub>CD</sub>. The  
alphadolone dissolved completely at a concentration of 10mg/ml, but unlike alphaxalone, it  
only went into solution after 4 hours of continuous stirring. This observation indicates that  
all neuroactive steroids do not interact with (7) sulfobutyl ether  $\beta$ -cyclodextrin in the same  
way. Fifteen male Wistar rats (150-200g weight) with surgically-implanted internal  
10 jugular intravenous catheters were used for these experiments in which they received  
intravenous alphadolone<sub>CD</sub> injections: 10 mg/kg (n=5); 20 mg/kg (n=5); and 40mg/kg  
(n=5).

[0129] They were assessed for anaesthesia by the righting reflex which was scored: 1  
15 normal; 2 slow; 3 some attempt; 4 none. A score of 4 means that a state of  
unconsciousness (anaesthesia) has been achieved. Figure 8 shows the results from this test  
for the rats in the three groups that received 10, 20 and 40 mg/kg alphadolone<sub>CD</sub>. Results  
shown are means of the readings from all five rats at each time point after intravenous  
injection of alphadolone.

20

[0130] The rats were also assessed for surgical anaesthesia using tail pinch responses  
scored: 1 normal; 2 weak; 3 just present; 4 none. A score of 4 indicates surgical  
anaesthesia has been achieved. Figure 9 shows the results from this test for the rats in the  
three groups that received 10, 20 and 40 mg/kg alphadolone<sub>CD</sub>. Results shown are means  
25 of the readings from all five rats at each time point after intravenous injection of  
alphadolone<sub>CD</sub>.

[0131] Time to complete recovery from the sedating effects of the alphadolone<sub>CD</sub> was  
assessed with the rotarod run time – normal non-sedated rats run on the accelerating  
30 rotating drum is 120 seconds; rats are fully recovered when they can walk on the rotarod  
treadmill for 120 seconds. Figure 10 shows the results from this test for the rats in the three



groups that received 10, 20 and 40 mg/kg alphadolone<sub>CD</sub>. Results shown are means of the readings from all 5 rats at each time point after intravenous injection of alphadolone<sub>CD</sub>.

### **Conclusions**

- 5 [0132] Alphadolone<sub>CD</sub> is an intravenous anaesthetic of short duration. It causes anaesthesia induction immediately after intravenous injection. This effect is dose related and it is possible to cause sufficient CNS depression to lead to surgical anaesthesia.

### EXAMPLE 10

#### *Cardiovascular Effects of Phaxan<sub>CD</sub> compared with Althesin and Propofol*

[0133] Fifteen male Wistar rats (150-200g weight) with surgically-implanted internal jugular intravenous catheters were used for these experiments in which rats in three groups received intravenous injections of equipotent AD<sub>95</sub> anaesthetic doses of either propofol (6.6mg/kg; Diprivan 10mg/ml propofol in 10% Intralipid emulsion), Althesin (3.28 mg/kg alphaxalone; Althesin 9mg/ml alphaxalone plus 3mg/ml alphadolone dissolved in 20% CremophorEL), or Phaxan<sub>CD</sub> (3.23 mg/kg alphaxalone; alphaxalone 10mg/ml dissolved in Captisol [(7) sulfobutyl ether  $\beta$ -cyclodextrin] 13%); n = 5 rats per group. Systolic and diastolic blood pressures were measured before and after these injections. Each measurement was calculated as a percentage change from the pre-anaesthetic levels for that rat. Figures 11 and 12 show the percentage changes against time for each of the cardiovascular parameters in each treatment group.

15

#### **Conclusion**

[0134] Both formulations of alphaxalone (Althesin/20% CremophorEL; Phaxan<sub>CD</sub>/sulfobutyl ether  $\beta$ -cyclodextrin (Captisol [Registered Trademark])) caused less cardiovascular disturbance than an equianaesthetic dose of propofol and in one measure (diastolic blood pressure) the sulfobutyl ether  $\beta$ -cyclodextrin formulation of alphaxalone (Phaxan<sub>CD</sub>) caused less cardiovascular disturbance compared with the CremophorEL preparation (Althesin [Registered Trademark]).

20

### EXAMPLE 11

#### *A phase 1/2a clinical trial*

[0135] In these experiments in human volunteers, propofol and (7) sulfobutyl ether  $\beta$ -  
5 cyclodextrin/alphaxalone formulation are compared in a double blind manner. Each  
volunteer is prepared in a fully equipped anaesthetic room. Dosing with either propofol or  
alphaxalone is determined by a randomization schedule to either propofol or alphaxalone.  
One anaesthetist designated to give the anaesthetic opens an envelope to find which drug  
should be given to this patient. The dose of drug is determined from a calculation schedule  
10 (see below) relying on the response of the previous patient to the last dose of that drug  
used – anaesthesia achieved or not based on a measurement of 50 for the bispectral index  
of the electroencephalogram (BIS value). The patient has an intravenous cannula in the  
right hand for drug administration and another for blood withdrawal for samples for  
measurements of blood levels of drug. That arm and the anaesthetist administering the  
15 anaesthetic has no communication with the test subject or a second anaesthetist who is in  
contact with the subject and who is responsible for general care of the subject as well as  
physiological monitoring. The arm and the drug administering anaesthetist are separated  
by a curtain from the anaesthetist caring for the subject as well as the subject and  
anaesthetic nurse present. The first, drug injecting anaesthetist, only communicates that  
20 the anaesthetic injection is about to start by ringing a bell and the caring anaesthetist only  
communicates with the drug administering anaesthetist to say whether a BIS value of 50 or  
less was achieved after that subject leaves the room at the end of the experiment.  
Measurements and assessments made:

- Subjects weight in kg. This is written on the case record before passing to the  
25 administering anaesthetist.
- The patient is asked by the caring anaesthetist to report pain on injection and a positive  
or no report is noted.
- Presence or absence of abnormal movements is noted by the caring anaesthetist.<sup>4</sup>
- Time from the ringing of the bell indicating anaesthetic injection to the subject  
30 dropping a water-loaded 20 ml syringe held between finger and thumb in the  
outstretched left arm.

- Time from the ringing of the bell indicating anaesthetic injection to loss of verbal contact with the patient and time for that contact to return.
  - Time from the ringing of the bell indicating anaesthetic injection to the subject losing the eyelash reflex and the time at which that reflex returned.
- 5 • BIS value and whether a value of 50 or below is achieved and when and for how long after the intravenous injection.
- Blood pressure, systolic and diastolic, and pulse rate using non invasive methods measured every 1 minute for 5 minutes, every 2.5 minutes for a further 10 minutes and every 5 minutes thereafter.
- 10 • Oxygen saturation of the blood measured with a pulse oximeter probe placed on the ear lobe of the left ear. The subject breathes air unless oxygen saturation levels fall below 93% at which time oxygen is given by face mask and anaesthetic circuit. Breathing is assisted if apnoea occurs and persists for longer than 30 seconds. The occurrence of low oxygen saturations and apnoea are noted.
- 15 • Time to full recovery indicated by normal performance in a digital substitution test
- Blood taken for analysis of alphaxalone blood levels at 0.5, 1.0, 1.5, 2, 5 10, 15, 30 and 60 minutes after anaesthetic injection.
  - Blood taken before the experiment, one hour after the experiment and 24 hours and one week later for:
- 20        Full haematological analysis.  
          Hepatic function tests.  
          Renal function tests.

### **Dose schedule**

- 25 [0136] When the envelope is opened by the administering anaesthetist the randomized instruction will be to give propofol or alphaxalone. If this is the first subject to receive the drug they are given: propofol 2 mg/kg; alphaxalone 0.5 mg/kg.
- The dosing for the next patient to receive that drug is determined by the response of the first subject given that drug.
- 30        If the first subject did not achieve a BIS of 50 or less, then for propofol the dose would be 3 mg/kg and for alphaxalone 0.75 mg/kg.

If the first subject did achieve a BIS of 50 or less, then for propofol the dose would be 1 mg/kg and for alphaxalone 0.25 mg/kg.

- Thereafter the dosing is:

5 a decrease of dose of 25% if all subjects so far treated with that drug achieved a BIS of 50 or,

an increase of dose of 25% if all subjects so far treated with that drug had all not achieved a BIS of 50 or,

10 In the case that there have been some subjects treated with this drug that have achieved BIS values of 50 or less and others with BIS values that have not fallen to 50 or less then either:

15 In the case of the last response being a BIS of 50 or less, then the dose of drug for the next subject to receive this drug will be mid way between the dose for the last subject given that drug and the dose given to the most recent previous subject given that drug and who did not achieve a BIS of 50 or below.

In the case of the last response being a BIS greater than 50, then the dose of drug for the next subject to receive this drug will be mid way between the dose for the last subject given that drug and the dose given to the most recent previous subject given that drug who did achieve a BIS of 50 or below.

20 The latter is repeated for each drug series until the dose range variations have become small and six subjects have achieved a BIS value of 50 or less having received the same dose  $\pm$  10% of drug. These 6 doses achieving anaesthetic levels of BIS are combined to calculate the mean  $\pm$  sem induction dose.

25

[0137] The following results are expected:

- Alphaxalone causes general anaesthesia with BIS values <50 achievable in one arm/brain circulation time.
- The quality of induction is at least as good as with propofol but with the added  
30 advantage of no pain on injection.
- At the “induction dose”, propofol causes greater falls in blood pressure, an increased incidence of apnoea and decreased oxygen saturation than alphaxalone.


- After administration of the “induction dose” the speed and quality of recovery is faster for alphaxalone.
- The pharmacokinetics of alphaxalone after intravenous administration are the same as for the alphaxalone figures in the literature after Althesin (Registered Trademark) administration.

5

[0138] Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to, or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

10

**TABLE 7**  
**Cyclodextrin structures**



compd	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
Hydrophilic Derivatives			
methylated cyclodextrins			
3-mono- <i>O</i> -methylcyclodextrins	H	CH <sub>3</sub>	H
2,6-di- <i>O</i> -methylcyclodextrins	CH <sub>3</sub>	H	CH <sub>3</sub>
2,3,6-tri- <i>O</i> -methylcyclodextrins	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>
randomly methylated cyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or CH <sub>3</sub>		
hydroxyalkylated cyclodextrins			
2-hydroxyethylcyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or CH <sub>2</sub> CH <sub>2</sub> OH		
2-hydroxypropylcyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or CH <sub>2</sub> CH(OH)CH <sub>3</sub>		
3-hydroxypropylcyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH		
2,3-dihydroxypropylcyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or CH <sub>2</sub> CH(OH)CH <sub>2</sub> OH		
branched cyclodextrins			
6- <i>O</i> -glucosylcyclodextrins	H	H	H or glucose
6- <i>O</i> -maltosylcyclodextrins	H	H	H or maltose
6- <i>O</i> -dimaltosylcyclodextrins	H	H	H or (maltose) <sub>2</sub>
Hydrophobic Derivatives			
alkylated cyclodextrins			
2,6-di- <i>O</i> -ethylcyclodextrins	C <sub>2</sub> H <sub>5</sub>	H	C <sub>2</sub> H <sub>5</sub>
2,3,6-tri- <i>O</i> -ethylcyclodextrins	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>
acylated cyclodextrins			
2,3-di- <i>O</i> -hexanoylcyclodextrins	COC <sub>6</sub> H <sub>11</sub>	COC <sub>6</sub> H <sub>11</sub>	H
2,3,6-tri- <i>O</i> -acetylcyclodextrins	COCH <sub>3</sub>	COCH <sub>3</sub>	COCH <sub>3</sub>
2,3,6-tri- <i>O</i> -propanoylcyclodextrins	COC <sub>3</sub> H <sub>7</sub>	COC <sub>3</sub> H <sub>7</sub>	COC <sub>3</sub> H <sub>7</sub>
2,3,6-tri- <i>O</i> -butanoylcyclodextrins	COC <sub>4</sub> H <sub>9</sub>	COC <sub>4</sub> H <sub>9</sub>	COC <sub>4</sub> H <sub>9</sub>
2,3,6-tri- <i>O</i> -valerylcyclodextrins	COC <sub>5</sub> H <sub>11</sub>	COC <sub>5</sub> H <sub>11</sub>	COC <sub>5</sub> H <sub>11</sub>
2,3,6-tri- <i>O</i> -hexanoylcyclodextrins	COC <sub>6</sub> H <sub>13</sub>	COC <sub>6</sub> H <sub>13</sub>	COC <sub>6</sub> H <sub>13</sub>
2,3,6-tri- <i>O</i> -octanoylcyclodextrins	COC <sub>8</sub> H <sub>17</sub>	COC <sub>8</sub> H <sub>17</sub>	COC <sub>8</sub> H <sub>17</sub>
Ionizable Derivatives			
anionic cyclodextrins			
6- <i>O</i> -(carboxymethyl)cyclodextrins	H	H	H or CH <sub>2</sub> COONa
6- <i>O</i> -(carboxymethyl)- <i>O</i> -ethylcyclodextrins	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	H, C <sub>2</sub> H <sub>5</sub> or CH <sub>2</sub> COONa
cyclodextrin sulfates	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or SO <sub>3</sub> Na		
sulfobutylcyclodextrins	R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> = H or (CH <sub>2</sub> ) <sub>4</sub> SO <sub>3</sub> Na		

<sup>a</sup> N = 6, α-CDs; N = 7, β-CDs; N = 8, γ-CDs; N = 9, δ-CDs.

From Uekama *et al.*, 1998 *supra*.

## **BIBLIOGRAPHY**

Atwood, Davies, MacNicol and Vogtie (Eds), *Comprehensive Supramolecular Chemistry*  
Vol 4, Pergamon: Oxford UK, 1996

Child *et al.*, *British Journal of Anaesthesia* 43:2-13, 1971

Davies and Morris, *Pharmaceutical Research*, 10(7):1093-95, 1993

Fromming and Szejtlic (eds), *Cyclodextrins in Pharmacy*, Kluwer: Dordrecht, The  
Netherlands, 1994

Remington's Pharmaceutical Sciences, Mack Publishing Company, Eaton, USA, 1990

Rowe's Handbook of Pharmaceutical Excipients, 2009

Thomason, *Crit Rev Ther Drug Carrier Syst* 14:1, 1997


Uekama *et al.*, *Chem. Rev.* 98: 2045-2076, 1998



**CLAIMS:**

1. An anaesthetic or sedative composition comprising a neuroactive steroid anaesthetic formulated with a sulfoalkyl ether cyclodextrin or an alkylated derivative thereof.
2. The composition of claim 1, wherein the neuroactive steroid anaesthetic is selected from alphadolone, allopregnanolone, eltanolone (pregnanolone) and a pharmacologically acceptable derivative or salt thereof.
3. The composition of claim 2, wherein the pharmacologically acceptable derivative is an alkylated, alkoxyated, acetylated, and/or phosphorylated derivative.
4. The composition of claim 2, wherein the pharmacologically acceptable derivative is a deuterated, tritiated or labelled derivative.
5. The composition of any one of claims 1 to 4, wherein the sulfoalkyl ether cyclodextrin is a sulfoalkyl ether  $\beta$ -cyclodextrin.
6. The composition of any one of claims 1 to 5, wherein the alkylated derivative is an alkyl ether derivative.
7. The composition of any one of claims 1 to 6, wherein the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is from 1:1 to 1:6.
8. The composition of claim 7, wherein the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is from 1:1 to 1:4.
9. The composition of claim 8, wherein the molar ratio of neuroactive steroid anaesthetic to cyclodextrin is 1:2.
10. The composition of any one of claims 1 to 9, further comprising one or more of an antimicrobial agent, a preservative, a buffer and/or a co-polymer.

11. The composition of claim 10 wherein the co-polymer is selected from hydroxy propyl methyl cellulose, polyvinyl pyrrolidone and carboxymethyl cellulose.
12. The composition of claim 11 wherein if a buffer is present, the pH is from pH 5.5 to pH 8.
13. The composition of any one of claims 1 to 12, wherein the sulfoalkyl ether cyclodextrin is a  $\beta$ -cyclodextrin sulfobutyl ether, an  $\alpha$ -cyclodextrin sulfobutyl ether, or a  $\gamma$ -cyclodextrin sulfobutyl ether, preferably a  $\beta$ -cyclodextrin sulfobutyl ether.
14. The composition of claim 13, wherein the sulfoalkyl ether cyclodextrin is (7) sulfobutyl ether  $\beta$ -cyclodextrin.
15. The composition of any one of claims 1 to 14, wherein the alkylated derivative is a sulfoalkyl ether-alkyl ether derivative.
16. The composition of any one of claims 1 to 15, wherein the alkylated derivative is a sulfobutyl ether-alkyl ether derivative.
17. The composition of claim 15 or claim 16, wherein the alkylated derivative is sulfobutyl ether-ethyl ether or its sodium salt thereof.
18. The anaesthetic or sedative composition of any one of claims 1 to 12, wherein the neuroactive steroid anaesthetic is alphaxalone and/or alphadolone or a pharmacologically acceptable derivative including a deuterated, tritiated or labelled derivative or salt thereof.
19. An anaesthetic or sedative composition of any one of claims 1 to 18 for use in inducing or maintaining anaesthesia or sedation in a subject, wherein said composition is for administration by infusion or bolus.
20. The composition of Claim 19 wherein the subject is a human.



**Application No:** GB1210657.1

**Examiner:** Dr S. David Evans

**Claims searched:** 1-20

**Date of search:** 27 September 2012

**Patents Act 1977: Search Report under Section 17**

**Documents considered to be relevant:**

Category	Relevant to claims	Identity of document and passage or figure of particular relevance
X	1, 19 at least	WO 93/17711 A1 (AUSTRALIAN COMMERCIAL RESEARCH) See in particular - abstract, page 1 lines 26-28, page 4 line 20 - page 5 line 12, page 9 lines 21-24, page 37 lines 20-25, page 38 lines 5-18, claim 29
Y	1, 19 at least	WO 01/70234 A1 (JUOX PTY LTD) See entire document, especially the preparation of the host-guest formulations on page 5
Y	1, 19 at least	EP 0399716 B1 (INNOVET INC) See whole document, particularly the Experimental section from pages 12-24
Y	1, 19 at least	JOURNAL OF PARENTERAL SCIENCE & TECHNOLOGY, Vol. 43(6), 1989, (BREWSTER M. E. ET AL), 'Development of a Non-Surfactant Formulation for Alfaxalone Through the Use of Chemically-Modified Cyclodextrins', pages 262 - 265 See whole document
Y	1, 19 at least	JOURNAL OF PHARMACEUTICAL SCIENCES Vol. 84(10), 1995, (BREWSTER M. E. ET AL), 'Preparation, Characterization, and Anaesthetic Properties of 2-hydroxypropyl-beta-cyclodextrin Complexes of Pregnenolone in Rat and Mouse', pages 1154 - 1159 See whole document
Y	1, 19 at least	WO 2004/039426 A1 (TRANSPHARMA MEDICAL LTD) See whole document, especially the examples


**Categories:**

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

**Field of Search:**

Search of GB, EP, WO & US patent documents classified in the following areas of the UKC<sup>X</sup> :

Worldwide search of patent documents classified in the following areas of the IPC



The following online and other databases have been used in the preparation of this search report

**International Classification:**

<b>Subclass</b>	<b>Subgroup</b>	<b>Valid From</b>
A61K	0047/40	01/01/2006
A61K	0031/56	01/01/2006
A61K	0031/57	01/01/2006
A61K	0047/48	01/01/2006
A61P	0023/00	01/01/2006