

667947

Patents Act 1990
Section 29

AUSTRALIA
Patents Act 1990

PATENT REQUEST: STANDARD PATENT/PATENT OF ADDITION

We, being the person identified below as the Applicant, request the grant of a patent to the person identified below as the Nominated Person, for an invention described in the accompanying standard complete specification.

Full application details follow.

[71] Applicant: HOECHST AKTIENGESELLSCHAFT
Address: D-65926 FRANKFURT AM MAIN, FEDERAL REPUBLIC OF GERMANY

[70] Nominated Person: HOECHST AKTIENGESELLSCHAFT
Address: D-65926 FRANKFURT AM MAIN, FEDERAL REPUBLIC OF GERMANY

[54] Invention Title: PROCESS FOR THE PREPARATION OF SUBSTITUTED INDANONES

[72] Name(s) of actual inventor(s): LAURENT WEISSE AND HEINZ STRUTZ

[74] Address for service in Australia: c/o WATERMARK PATENT & TRADEMARK ATTORNEYS, of The Atrium,
290 Burwood Road, Hawthorn, Victoria 3122, Australia Attorney Code: WM

BASIC CONVENTION APPLICATION(S) DETAILS

[31] Application Number	[33] Country	Country Code	[32] Date of Application
P42 30 373.7	GERMANY	DE	11TH SEPTEMBER 1992

Basic Applicant(s): HOECHST AKTIENGESELLSCHAFT
Drawing number recommended to accompany the abstract

By our Patent Attorneys,
WATERMARK PATENT & TRADEMARK ATTORNEYS



Darryl B. Mischewski
Registered Patent Attorney

DATED this 9th day of September 1993.

C 005011 100993

AUSTRALIA

Patents Act 1990

NOTICE OF ENTITLEMENT

We, HOECHST AKTIENGESELLSCHAFT of, D65926 Frankfurt Am Main, Germany, being the applicant in respect of Application No. 46260/93 state the following:-


The Persons nominated for the grant of the patent have entitlement from the actual inventors by virtue of an employment contract.

The persons nominated for the grant of the patent are the applicant of the basic application listed on the patent request form.

The basic application listed on the request form is the first application made in a Convention country in respect of the invention

HOECHST AKTIENGESELLSCHAFT

By our Patent Attorneys,
WATERMARK PATENT & TRADEMARK ATTORNEYS


.....

Carolyn J. Harris
Registered Patent Attorney

6 February 1996




AU9346260


(12) PATENT ABRIDGMENT (11) Document No. AU-B-46260/93
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 667947

- (54) Title
PROCESS FOR THE PREPARATION OF SUBSTITUTED INDANONES
- International Patent Classification(s)
(51)^s C07C 045/46 C07C 049/67 C07C 049/675 C07C 049/697
- (21) Application No. : 46260/93 (22) Application Date : 10.09.93
- (30) Priority Data
- (31) Number (32) Date (33) Country
4230373 11.09.92 DE GERMANY
- (43) Publication Date : 17.03.94
- (44) Publication Date of Accepted Application : 18.04.96
- (71) Applicant(s)
HOECHST AKTIENGESELLSCHAFT
- (72) Inventor(s)
LAURENT WEISSE; HEINZ STRUTZ
- (74) Attorney or Agent
WATERMARK PATENT & TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN VIC 3122
- (57) Claim

1. A process for the preparation of a compound of the formula IV or the isomer thereof of the formula IVa



(IV)



(IVa)


in which

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are identical or different and are hydrogen, (C_1-C_{20}) -alkyl, (C_6-C_{14}) aryl, (C_1-C_{10}) alkoxy, (C_2-C_{10}) alkenyl, (C_7-C_{20}) arylalkyl, (C_7-C_{20}) alkylaryl, (C_6-C_{10}) aryloxy, (C_1-C_{10}) fluoroalkyl, (C_6-C_{10}) haloaryl, (C_2-C_{10}) alkynyl, an $-SiR^8_3$ radical where R^8 is (C_1-C_{10}) alkyl, ~~or are~~ a halogen atom or a heteroaromatic radical having 5 or 6 ring members which may contain one or more heteroatoms; or adjacent radicals R^1-R^4 , together with the atoms


(11) AU-B-46260/93
(10) 667947

-2-


connecting them, form one or more substituted or unsubstituted rings, which comprises reacting a compound of the formula I



with a compound of the formula II



or with a compound of the formula (III)



in which R¹-R⁷ are as defined above,
in liquid, anhydrous hydrogen fluoride and with
boron trifluoride.

AUSTRALIA

Patents Act 1990

667947

ORIGINAL
COMPLETE SPECIFICATION
STANDARD PATENT

Application Number:

Lodged:

Invention Title:

PROCESS FOR THE PREPARATION OF SUBSTITUTED INDANONES

The following statement is a full description of this invention, including the best method of performing it known to us :-

Description

Process for the preparation of substituted indanones

The present invention relates to a technically simple process for the preparation of substituted 1-indanones.

5 Compounds of this type are important intermediates in the preparation of metallocene complexes, since 1-indanones can readily be converted into the corresponding indenenes. Indenenes are used as a ligand system for the synthesis of metallocene complexes (EP-A 336 128).

10 Furthermore, substituted 1-indanones are of industrial importance as scents (EP-A 162 465) and as valuable intermediates in the preparation of pharmaceutical products or other bioactive compounds (EP-A 421 759; J. Med. Chem. 25 (1990) 765).

15 A number of processes for the preparation of substituted 1-indanones are described in the literature.

20 1-Indanones which carry substituents on the six-membered ring can be prepared starting from the correspondingly substituted aromatic compounds by fusing the five-membered ring on in 2- to 6-step syntheses (J. Org. Chem., 55 (1990) 247; Bull. Soc. Chim. Fr. 6 (1969) 1981).

25 Processes for the preparation of 1-indanones which carry substituents on the five-membered ring or on both rings are likewise known (J. Org. Chem. 46 (1981) 3758; J. Org. Chem. 23 (1958) 1441).

30 These methods have the disadvantage that they are generally multistep and give only poor overall yields of the desired products. Many of the syntheses cannot be applied generally, but are restricted to specific derivatives. In others, the starting materials are again not

readily available or are very expensive. Certain substitution patterns on the aromatic ring are likewise impossible to achieve by these methods. The few known one-step syntheses have the disadvantage that they are restricted to specific derivatives and give poor yields, so that technically complex purification operations of the products are necessary. Most of these reactions are carried out with the aid of Friedel-Crafts catalysts, such as, for example, $AlCl_3$, which are employed in excess. These reactions require technically complex work-up steps, which are associated with production of a large amount of salt.

Also known are processes for the preparation of substituted indanones by reacting aromatic compounds, such as xylene or acenaphthene, with aqueous methacrylic acid, crotonic acid or cinnamic acid in a large excess of liquid hydrogen fluoride (J. Am. Chem. Soc. 61 (1939) 1272; J. Am. Chem. Soc. 72 (1950) 3287). The yields are between 62% and 81%. This method has the disadvantage that water which is present or formed significantly reduces the activity of the Friedel-Crafts catalyst. This results in low yields and corrosion problems.


EP 93 106 649.2 describes a process which enables the preparation of substituted 1-indanones in one step from relatively inexpensive starting compounds. In the case of certain starting materials, this method gives two or more isomers, it being virtually impossible to preferentially prepare one of these isomers. In the case of highly deactivated aromatic compounds, this process is in addition very time consuming or unsuitable.

The object was therefore to find a process for the preparation of the abovementioned indanones which avoids the disadvantages known from the prior art.


Surprisingly, it has been found that aromatic compounds of the formula I below react with commercial carboxylic

anhydrides of the formula II or carboxylic acid fluorides of the formula III in liquid hydrogen fluoride and with boron trifluoride rapidly, virtually quantitatively and even under relatively mild conditions to give indanones of the formula IV/IVa. In addition, certain indanones which, according to the prior art, were only accessible by complex syntheses can be prepared in one step.

The present invention therefore relates to a process for the preparation of a compound of the formula IV or the isomer thereof of the formula IVa




(IV)




(IVa)

in which


R¹, R², R³, R⁴, R⁵, R⁶ and R⁷ are identical or different and are hydrogen, (C₁-C₂₀)alkyl, (C₅-C₁₄)aryl, (C₁-C₁₀)alkoxy, (C₂-C₁₀)alkenyl, (C₇-C₂₀)arylalkyl, (C₇-C₂₀)alkylaryl, (C₆-C₁₀)aryloxy, (C₁-C₁₀)fluoroalkyl, (C₆-C₁₀)haloaryl, (C₂-C₁₀)alkynyl, an -SiR⁸₃ radical where R⁸ is (C₁-C₁₀)alkyl, or are a halogen atom or a heteroaromatic radical having 5 or 6 ring members which may contain one or more heteroatoms, or adjacent radicals R¹-R⁴, together with the atoms connecting them, form one or more substituted or unsubstituted rings, which comprises reacting a compound of the formula I



with a compound of the formula II



or with a compound of the formula (III)



in which R¹-R⁷ are as defined above,
in liquid, anhydrous hydrogen fluoride and with boron
trifluoride.

In these formulae, alkyl is straight-chain or branched
alkyl. Halogen is fluorine, chlorine, bromine or iodine,
in particular fluorine or chlorine. Examples of hetero-
aromatic radicals are thienyl, furyl and pyridyl.

- 10 In the formulae IV and IVa, it is preferred that R¹, R²,
R³ and R⁴ are identical or different and are hydrogen,
(C₁-C₁₀)-alkyl, (C₁-C₄)-alkoxy, (C₂-C₆)-alkenyl,
(C₆-C₁₄)-aryl, (C₁-C₆)-fluoroalkyl, (C₆-C₁₄)-aryloxy or a
halogen atom, or the radicals R¹ and R², R² and R³ or R³
15 and R⁴, together with the atoms connecting them, form a
five- or six-membered ring, and R⁵, R⁶ and R⁷ are identi-
cal or different and are hydrogen, (C₁-C₁₀)-alkyl or
(C₆-C₁₄)-aryl.

The rings formed by adjacent radicals R¹-R⁴ can be substituted by substituents as defined for R¹-R⁷, or the preferred meanings mentioned therefor.

5 It is particularly preferred that R¹, R², R³ and R⁴ are identical or different and are hydrogen, (C₁-C₁₀)-alkyl, (C₆-C₁₄)-aryl, (C₁-C₄)-alkoxy or a hydrogen atom, or the radicals R¹ and R², R² and R³ or R³ and R⁴, together with the atoms connecting them, form a five- or six-membered, preferably six-membered, saturated or unsaturated carbocyclic ring, and R⁵, R⁶ and R⁷ are identical or different and are hydrogen, methyl or phenyl.

10

The saturated or unsaturated five- or six-membered (carbocyclic) ring formed by adjacent substituents R¹-R⁴ may additionally carry substituents, preferably (C₁-C₁₀) alkyl.

15

Examples which may be mentioned of compounds of the formulae IV and IVa are:

- 6,7-benzo-2-methylindan-1-one,
4,5-benzo-2-methylindan-1-one,
20 5,7-diisopropyl-2-methylindan-1-one,
4,6-diisopropyl-2-methylindan-1-one,
2,5-dimethylindan-1-one,
2,6-dimethylindan-1-one,
5-isobutyl-2-methylindan-1-one,
25 2,5,7-trimethyl-1-indanone,
2,4,6-trimethyl-1-indanone,
2-methylindan-1-one,
2,4,5,6-tetramethylindan-1-one,
5-phenyl-2-methylindan-1-one,
30 8-methyl-4,5,7,8-tetrahydrocyclopenta [e] acenaphthylen-9-one,
2-methyl-3,9-dihydro-2H-cyclopenta [b] fluoren-1-one,
2-methyl-2,10-dihydro-1H-cyclopenta [a] fluoren-3-one,
16-methyl-6,7,15,16-tetrahydrocyclopenta [a] phenanthren-
35 17-one,

9-methyl-5,6,9,10-tetrahydro-cyclopenta [b]phenanthren-8-one,
5-methoxy-2-methylindan-1-one and
5,6-dimethoxy-2-methylindan-2-one.

5 Depending on the substitution pattern on the aromatic ring, the indanones may be produced in the form of two constitutional isomers of the formulae IV and IVa. Depending on the intended application, these can be further reacted in pure form or as a mixture. In the
10 preparation of metallocene complexes and when the 1-indanones are used as scents, the isomer mixture can be employed.

The indanones IV/IVa are preferably prepared by reacting aromatic compounds of the formula I with anhydrides of
15 the formula II.

The starting compounds of the formula I are commercially available or can be prepared by methods known from the literature.

The carboxylic acid fluorides of the formula III can be
20 prepared from the known carboxylic acid chlorides or carboxylic anhydrides (formula II) in a manner known from the literature by reaction with HF (cf., for example, Advanced Organic Chemistry, 1983, 399).

In the preparation of the compounds IV/IVa, additional
25 solvent can be added to the hydrogen fluoride, but the reaction is preferably carried out in pure, anhydrous hydrogen fluoride.

The molar ratio between the compound of the formula I, the compound II or III and the hydrogen fluoride can vary
30 within broad limits. The I : II or III : HF molar ratio is preferably from 1 : 0.5-2.0 : 5-100, in particular from 1 : 0.9-1.2 : 20-50. This means that the hydrogen fluoride is employed in excess.

If x is the total number of ether, keto, thio or carboxyl groups in the starting compounds, the molar ratio between boron trifluoride and the compound of the formula I is from $(0.5-1.5) x : 1$.

- 5 The reaction temperature is preferably from -30°C to 130°C , in particular from -10°C to 80°C .

The reaction times generally vary between 10 minutes and 24 hours, preferably between 30 minutes and 8 hours.

- 10 Reaction is preferably carried out in the pressure range from 1 to 15 atm.

It is preferred to initially introduce a mixture of the compounds I and II (or I and III) and to meter in the hydrogen fluoride. The reverse sequence of addition is also possible.

- 15 When the reaction is complete, the hydrogen fluoride can be removed by distillation and recovered virtually quantitatively without significant impurities. Recovery of boron trifluoride is also possible in principle.

- 20 The indanones of the formulae IV and IVa can be freed from acid components by washing with Na_2CO_3 , NaHCO_3 or KOH solution and water and dried using conventional desiccants, such as Na_2SO_4 , MgSO_4 or molecular sieves. Since the reactions are generally virtually quantitative, further purification can in most cases be omitted.

- 25 However, filtration through silica gel, aluminum oxide or filtration aids, such as, for example, Celite, is frequently advisable. If necessary, further purification can be carried out by distillation, column chromatography or crystallization. If necessary, the constitutional
30 isomers III and IIIa can be separated from one another by column chromatography on silica gel or aluminum oxide.

The process according to the invention is particularly distinguished by the fact that differently substituted 1-indanones can be obtained in a simple and short synthesis (one-step process) and in virtually quantitative yield. The space-time yields can be considerably improved compared with the prior art by the use of boron trifluoride. A further advantage of this method is the ability to optimize the product selectivity by slightly modifying the reaction conditions (for example temperature). A particular advantage is that the process according to the invention also allows the conversion of electron-deficient aromatic compounds, such as, for example, fluorobenzene, into the corresponding indanone, which was not possible by means of the previous prior art.

The indanones IV/IVa are preferably used for the preparation of metallocenes (cf., for example, EP-A 336 128) or as scents (EP-A 162 465).

The examples below serve to illustrate the invention in greater detail.

Example A

6,7-Benzo-2-methylindan-1-one (1) and 4,5-benzo-2-methylindan-1-one (1a)

14 g of boron trifluoride were added to 12.6 g (98 mmol) of naphthalene, 15.8 g (103 mmol) of methacrylic anhydride and 100 g (5 mol) of hydrogen fluoride in a 250 ml stainless-steel autoclave, and were reacted at -10°C for 1 hour. The reaction mixture was subsequently poured onto 1 kilo of ice, and the solution was neutralized by means of dilute KOH. After the aqueous phase had been repeatedly washed with ethyl acetate, the organic phases were separated off, dried and freed from solvent in vacuo, giving 18.7 g (95% yield) of a mixture of (1) and (1a) with a selectivity of 81 and 19% respectively.

Comparative Example to A

12.6 g (98 mmol) of naphthalene and 15.8 g (103 mmol) of methacrylic anhydride were reacted at 50°C for 18 hours in 100 g (5 mol) of HF. Work-up was carried out analogously to Example A, giving 19 g (97% yield) of product. The selectivity was 58% of compound (1) and 39% of compound (1a).

Example B

5-Fluoro-2-methyl-indan-1-one (2)

10 14 g (206 mmol) of boron trifluoride were added analogously to Example A to 9.6 g (100 mmol) of fluorobenzene, 15.8 g (103 mmol) of methacrylic anhydride and 100 g (5 mol) of HF, and the mixture was stirred at 50°C for 3 hours. Work-up was carried out analogously to Example A, giving 16 g (97% yield) of a virtually colorless liquid. The selectivity to (2) was 82%.

Comparative Example to B

9.6 g (100 mmol) of fluorobenzene, 16 g (104 mmol) of methacrylic anhydride and 100 g (5 mol) of HF were reacted at 70°C for 18 hours. Work-up was carried out analogously to Example A, but no 5-fluoro-2-methylindan-1-one was isolated.

Example C

2-Methyl-5-phenylindan-1-one (3)

25 15.4 g (100 mmol) of biphenyl, 16 g (104 mmol) of methacrylic anhydride, 100 g (5 mol) of HF and 14 g (206 mmol) of BF₃ were reacted at 50°C for 2 hours analogously to Example A and worked up. 22 g (99% yield) of product were isolated. The selectivity to (3) was 94%.

Comparative Example to C

15.4 g (100 mmol) of biphenyl and 16 g (104 mmol) of methacrylic anhydride were reacted with 100 g (5 mol) of HF at 70°C for 60 hours. Work-up carried out analogously to A gave 22.2 g (100% yield) of product. The purity was 92%.

5


5

5


5

CLAIMS: THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A process for the preparation of a compound of the formula IV or the isomer thereof of the formula IVa




(IV)



(IVa)

in which


5 $R^1, R^2, R^3, R^4, R^5, R^6$ and R^7 are identical or different and are hydrogen, (C_1-C_{20}) -alkyl, (C_6-C_{14}) aryl, (C_1-C_{10}) alkoxy, (C_2-C_{10}) alkenyl, (C_7-C_{20}) arylalkyl, (C_7-C_{20}) alkylaryl, (C_6-C_{10}) aryloxy, (C_1-C_{10}) fluoroalkyl, (C_6-C_{10}) haloaryl, (C_2-C_{10}) alkynyl, an $-SiR^8$ radical where R^8 is (C_1-C_{10}) alkyl, ~~or a~~ a halogen atom or a heteroaromatic radical having 5 or 6 ring members which may contain one or more heteroatoms; or adjacent radicals R^1-R^4 , together with the atoms connecting them, form one or more substituted or unsubstituted rings, which comprises reacting a compound of the formula I




(I)

with a compound of the formula II





or with a compound of the formula (III)



in which R¹-R⁷ are as defined above,
in liquid, anhydrous hydrogen fluoride and with
boron trifluoride.

5 2. The process as claimed in claim 1, wherein, in the
formulae IV and IVa, R¹, R², R³ and R⁴ are identical
or different and are hydrogen, (C₁-C₁₀)alkyl,
10 (C₁-C₄)alkoxy, (C₂-C₆)alkenyl, (C₆-C₁₄)aryl,
(C₁-C₆)fluoroalkyl, (C₆-C₁₀)aryloxy or a halogen atom,
or the radicals R¹ and R², R² and R³ or R³ and R⁴,
together with the atoms connecting them, form a
five- or six-membered ring, and R⁵, R⁶ and R⁷ are
identical or different and are hydrogen,
15 (C₁-C₁₀)alkyl or (C₆-C₁₄)aryl.

3. The process as claimed in claim 1 or 2, wherein, in
the formulae IV and IVa, R¹, R², R³ and R⁴ are iden-
tical or different and are hydrogen, (C₁-C₁₀)alkyl,
20 (C₆-C₁₄)aryl, (C₁-C₄)alkoxy or a halogen atom, or the
radicals R¹ and R², R² and R³ or R³ and R⁴, together
with the atoms connecting them, form a six-membered,
saturated or unsaturated carbocyclic ring, and R⁵, R⁶
and R⁷ are identical or different and are hydrogen,
methyl or phenyl.

4. The process as claimed in one or more of claims 1 to



3, wherein the compound I : compound II or compound III : hydrogen fluoride molar ratio is from 1 : 0.5-2.0 : 5-100.

5 5. The process as claimed in one or more of claims 1 to 4, wherein the boron trifluoride: compound I molar ratio is from $(0.5-1.5)x : 1$, in which x is the total number of ether, keto, thio or carboxyl groups in the starting compounds.

10 ~~6. The process as claimed in one or more of claims 1 to 5, wherein a compound of the formula I is reacted with a compound of the formula II or of the formula III.~~

DATED this 9th day of September 1993.

HOECHST AKTIENGESELLSCHAFT


WATERMARK PATENT & TRADEMARK ATTORNEYS
"THE ATRIUM"
290 BURWOOD ROAD
HAWTHORN. VIC. 3122.




Abstract of the Disclosure

Process for the preparation of substituted indanones

1-Indanones of the formula IV or IVa




(IV)




(IVa)

5 in which R¹ to R⁷ are preferably hydrogen or alkyl, or adjacent radicals R¹ to R⁴ form a ring, are obtained in a one-step reaction by reacting a compound I




(I)

with a compound of the formula II



(II)

or with a compound of the formula (III)



in anhydrous hydrogen fluoride and with boron trifluoride.

