



EP 2 244 590 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
16.05.2012 Bulletin 2012/20

(51) Int Cl.:
A23K 1/06 (2006.01) **A23K 1/165 (2006.01)**
A23L 1/015 (2006.01) **A23K 1/18 (2006.01)**
C12N 9/18 (2006.01)

(21) Application number: **08865475.1**

(86) International application number:
PCT/EP2008/067885

(22) Date of filing: **18.12.2008**

(87) International publication number:
WO 2009/080701 (02.07.2009 Gazette 2009/27)

(54) CUTINASE FOR DETOXIFICATION OF FEED PRODUCTS

CUTINASE FÜR DIE ENTGIFTUNG VON FUTTERMITTELPRODUKTEN

CUTINASE UTILISÉE POUR LA DÉTOXIFICATION DE PRODUITS DESTINÉS À L'ALIMENTATION POUR BÉTAIL

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR**

(30) Priority: **20.12.2007 EP 07150205**

(43) Date of publication of application:
03.11.2010 Bulletin 2010/44

(73) Proprietor: **Novozymes A/S**
2880 Bagsvaerd (DK)

(72) Inventors:

• **VIKSOE-NIELSEN, Anders**
DK-3550 Slangerup (DK)

• **SOERENSEN, Birthe, Hauerbach**
DK-2000 Frederiksberg (DK)

(56) References cited:

EP-A- 0 981 953 **WO-A-96/12414**
WO-A-2007/133263 **US-A1- 2003 073 239**

• **MARION BRODHAGEN, NANCY P. KELLER:**
"Signalling pathways connecting mycotoxin production and sporulation" MOLECULAR PLANT PATHOLOGY, vol. 7, no. 4, 2006, pages 285-301, XP002478569

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**REFERENCE TO A SEQUENCE LISTING**

5 [0001] This application contains a Sequence Listing in computer readable form. The computer readable form is incorporated herein by reference.

FIELD OF THE INVENTION

10 [0002] The present invention relates to a method comprising treatment with cutinase for detoxification of feed products contaminated by the mycotoxin zearalenone.

BACKGROUND OF THE INVENTION

15 [0003] Several plant pathogenic and/or post-harvest *Fusarium* species on cereals produce toxic substances of considerable concern to livestock and poultry producers, e.g., deoxynivalenol, T-2 toxin, HT-2 toxin, diacetoxyscirpenol and zearalenone.

20 [0004] Zearalenone is found worldwide in a number of cereal crops, such as maize, barley, oats, wheat, rye, rice, millet and sorghum. Zearalenone production does not seem to occur in significant amounts prior to harvest, but under proper environmental conditions, it is readily produced on corn and small grains in storage.

[0005] When cereal grain is used in ethanol production and the starch is consumed the zearalenone is concentrated in the fermentation by-products, e.g., in the distiller's dried grain. The contents of zearalenone in the fermentation by-products may be increased threefold relative to the cereal grain.

25 [0006] The toxin is heat-stable, and it is not destroyed by long storage, roasting, or by the addition of propionic acid or mold retardants.

[0007] Despite their structural dissimilarity to the steroidal estrogens, zearalenone and several of its derivatives possess estrogenic activity. Zearalenone undergoes a folding such that hydroxyl or potential hydroxyl groups become appropriately orientated to facilitate binding to tissue receptors that normally bind estrogens.

30 [0008] Zearalenone is the primary toxin causing infertility, abortion or other breeding problems, especially in swine. The symptoms are especially severe in prepubertal gilts including enlarged mammae, swelling of uterus and vulva, and atrophy of the ovaries. In severe cases, prolapse of the vulva and rectum may occur. Boars exhibit enlarged mammae and atrophied testes.

35 [0009] Zearalenone is present in the meat from animals feeding on contaminated grain as well as in bread baked from contaminated wheat. While cases of poisoning of humans are rare there is concern about the effect of the long term exposure of humans to such an estrogenic activity.

[0010] Inactivation of mycotoxins, including zearalenone, using epoxidase or lactonase is disclosed in W09612414.

40 [0011] There is a need for further methods of detoxification of animal feed products, e.g., such as fermentation by-products, including distiller's wet and dried grain, contaminated by the mycotoxin zearalenone.

SUMMARY OF THE INVENTION

[0012] The inventors of the present invention have discovered that zearalenone in a feed product can be degraded by treating the feed product with a cutinase. Accordingly, in a first aspect the invention provides a process for degrading zearalenone in a feed product which process comprises treating said feed product with a cutinase.

45 [0013] In a second aspect the invention provides a use of a cutinase for degrading the mycotoxin zearalenone.

Detailed description of the inventionZearalenone

50 [0014] In the context of this invention the term "zearalenone" comprises the mycotoxin zearalenone produced from certain *Fusarium* sp. The IUPAC name is (4S,12E)-15, 17-Dihydroxy-4-methyl-3-oxabicyclo[12.4.0]octadeca-12, 15, 17, 19-tetraene-2, 8-dione. The term "zearalenone" also comprises any derivative of zearalenone which comprises an internal carboxylic ester bond susceptible for modification by a cutinase.

55 [0015] The term "animal" includes all animals, including human beings. Examples of animals are cattle, (including but

not limited to cows and calves); mono-gastric animals, e.g. pigs or swine (including, but not limited to, piglets, growing pigs, and sows); poultry such as turkeys and chicken (including but not limited to broiler chicks, layers); and fish (including but not limited to salmon).

[0016] The term "feed" or "feed product" means any compound, preparation, mixture, or composition suitable for, or intended for intake by an animal.

[0017] The feed product may be a product which apart from an unwanted level of zearalenone is suitable for consumption by an animal. The feed product can also be a product suspected of comprising an unwanted level of zearalenone, and/or a product having an unknown level of zearalenone, including products not comprising a detectable level of zearalenone.

[0018] Preferably the feed product is a grain based product. Preferably the grain based product comprises cereal(s), e.g., one or more of corn, wheat, barley, rye, rice, sorghum and millet. Also preferred are grain based product comprising material derived from one or more of corn, wheat, barley, rye, rice, sorghum and millet. In one embodiment, the feed product

US2003/073239 teaches the use of nucleotide sequences encoding zearalenone esterase polypeptides, isolated from the fungus *Gliocladium roseum*, for detoxifying zearalenone and structurally related mycotoxins. may e.g. be derived solely from cereal(s), and in another embodiment partly from legumes, e.g. from soybean, and partly from cereals. The grain based product may comprise whole or milled grain, e.g., wet or dry milled grain, including grain based product comprising fractions of wet or dry milled grain, e.g., gluten, protein, starch, and/or oil fractions. Also preferred are products comprising a by-product from brewing and/or fermentation processes, e.g., spent grain. Spent grain is the by-products from the production of alcoholic beverages and ethanol fuels. Brewers' spent grain (BSG) is the residue of beer making in breweries, which use malted barley as the major raw material. Distiller's spent grain (DSG) is the product left in distilleries after alcohol is removed by distillation from the fermented grains such as corn, wheat, barley, rice, and rye. Distiller's spent grain is also known as distiller's grain. Wet distiller's grain (WDG) is dried to produce dried distiller's grain (DDG) which is used primarily as animal feed.

Cutinases

[0019] In the context of this invention the term "cutinases" include enzymes comprised by the enzyme classification E.C.3.1.1.74. Preferred are the below mentioned enzymes as well as enzymes with homologous sequence, especially recombinant and/or substantially purified enzymes.

[0020] The cutinase may be derived from a fungus. Particularly, the cutinase may be derived from a strain of *Humicola*, particularly *H. insolens*, more particularly *H. insolens* strain DSM1800 (US 5,827,719) or from a strain of *Fusarium*, e.g. *F. roseum culmorum*, or particularly *F. solani pisi* (WO 90/09446; WO 94/14964, WO 94/03578). The fungal cutinase may also be derived from a strain of *Rhizoctonia*, e.g. *R. solani*, or a strain of *Alternaria*, e.g. *A. brassicicola* (WO 94/03578). The cutinase may also be a variant of a parent cutinase such as those described in WO 00/34450, or WO 01/92502, all of which are hereby incorporated by reference. The cutinase may be the variant of the *Humicola insolens* cutinase comprising the substitutions E6Q, G8D, A14P, N15D, E47K, S48E, R51 P, A88H, A91 H, A130V, E179Q and R189V, which is disclosed at p. 24, line 11 of 10038.204-WO.

[0021] SEQ ID NO:1 is the amino acid sequence of the *Humicola insolens* cutinase (corresponding to the mature part of SEQ ID NO:2 of US 5,827,719, and of SEQ ID NO:1 of WO 01/92502), and SEQ ID NO:2 is the amino acid sequence of the *Fusarium solani pisi* according to Fig. 1D of WO 94/14964.

[0022] The cutinase must be present in the medium to be detoxified in effective amounts. Preferably the cutinase is present in concentrations of 0.01-100 mg enzyme protein pr. kg dry matter, preferably 0.1-10 mg enzyme protein pr. kg dry matter, or more preferably 1-5 mg enzyme protein pr. kg dry matter.

The medium

[0023] In an embodiment the cutinase is degrading the zearalenone in a medium comprising the feed product. The medium is preferably aqueous and may be a liquid, a paste or a slurry. To form a suitable medium water may be added to the feed product. The cutinase be comprised in solid or liquid formulations suitable for application to said medium.

[0024] In a embodiment the cutinase is degrading the zearalenone to an extent whereby the content of zearalenone per kg dry matter feed product is reduced to less than 50%, preferably less than 60%, more preferably less than 70%, and most preferably to less than 80% of the initial amount.

[0025] The detoxification efficiency of the invention depends on e.g. availability of water, pH, temperature and buffer of the medium. For example, the treatment may take place at a pH-value at which the relative activity of the actual cutinase is at least 50, or 60, or 70, or 80 or 90%. Likewise, for example, the treatment may take place at a temperature at which the relative activity of the actual cutinase is at least 50, or 60, or 70, or 80 or 90%. The relative activity is calculated relative to the activity at the pH value where the highest activity is observed.

pH in the medium

[0026] Depending, inter alia, on the characteristics of the cutinase employed, the pH in the medium employed should normally be in the range of 5-11, preferably in the range 6-10, e.g. 6.5-8.5.

5

Temperature in the medium

[0027] Preferably a reaction temperature is applied which is close to the optimum temperature for the cutinase. In numerous embodiments of the invention, temperatures in the range of 10-65°C, more preferably 30-50°C, should be employed.

10

Treatment duration

[0028] The duration of treatment depends, inter alia, on the treatment type, the type of item to be treated, the properties of the medium, e.g. temperature and pH and the type and amounts of enzyme employed.

[0029] The enzymatic reaction is continued until the desired result is achieved, following which it may or may not be stopped by inactivating the enzyme, e.g., by a heat-treatment step.

[0030] For detoxification purposes treatment times in the range of 1 minute to 1 week may be employed. In many cases a treatment time in the range of 6 to 48 hours will be suitable.

20

Identity

[0031] The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity".

[0032] For purposes of the present invention, the degree of identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends in Genetics 16: 276-277), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0033] For purposes of the present invention, the degree of identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 3.0.0 or later. The optional parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

45

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

50

Homologous sequence

[0034] The term "homologous sequence" is defined as a predicted protein that gives an E value (or expectancy score) of less than 0.001 in a tfasty search (Pearson, W.R., 1999, in Bioinformatics Methods and Protocols, S. Misener and S. A. Krawetz, ed., pp. 185-219) with a specified sequence.

55

[0035] The term "homologous sequence" may also be defined as a sequence that has a degree of identity at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or even 100%, to a specified sequence.

EXAMPLES

Example 1

5 [0036]

Enzyme: A recombinantly produced enzyme composition comprising the variant of the cutinase from *Humicola insolance* disclosed at p. 24, line 11 of 10038.204-WO.

Assay: Reactions were performed in 300 microL volumes in eppendorf tubes comprising zearalenone 30 microM, Tris 100 mM and enzyme 0.1 mg EP/mL. In control reactions the enzyme volume was substituted an equivalent amount of H₂O. The reactions were incubate 24 hours at 37°C before being terminated by adding 600 microL of a 100 microM acetonitrile stop solution. Reactions were stored at -20° C until chromatographic analysis.

Chromatographic analysis: Samples were centrifugated and the supernatant analysed for zearalenone by HPLC-DAD as described by Smedsgaard (J. Chromatogr. A, 1997, 760, 264-270). The DAD scanned from 200-600 nm. Separation was done on a Phenomenex (Torrance, CA) Luna C18(2) 10×2 mm ID, 3 micrometer, column 2, using a linear gradient moving form 5% to 100% acetonitrile in 20 min. Residual zearalenone was calculated relative to the control. The results are presented in tables 1.

Table 1. Residual zearalenone after 24 hours incubation with or without a cutinase at pH 7.

20 Enzyme	Residual zearalenone (%)
Control	100
Cutinase	19

25 SEQUENCE LISTING

[0037]

<110> Novozymes A/S

30 <120> CUTINASE FOR DETOXIFICATION OF FEED PRODUCTS

<130> 11339.204-WO

35 <160> 2

<170> PatentIn version 3.5

<210> 1

40 <211> 194

<212> PRT

<213> *Humicola insolens*

<220>

45 <221> mat_peptide

<222> (1)..(194)

<400> 1

50

55

EP 2 244 590 B1

Gln Leu Gly Ala Ile Glu Asn Gly Leu Glu Ser Gly Ser Ala Asn Ala
1 5 10 15

5 Cys Pro Asp Ala Ile Leu Ile Phe Ala Arg Gly Ser Thr Glu Pro Gly
20 25 30

10 Asn Met Gly Ile Thr Val Gly Pro Ala Leu Ala Asn Gly Leu Glu Ser
35 40 45

15 His Ile Arg Asn Ile Trp Ile Gln Gly Val Gly Gly Pro Tyr Asp Ala
50 55 60

20 Ala Leu Ala Thr Asn Phe Leu Pro Arg Gly Thr Ser Gln Ala Asn Ile
65 70 75 80

25 Asp Glu Gly Lys Arg Leu Phe Ala Leu Ala Asn Gln Lys Cys Pro Asn
85 90 95

30 Thr Pro Val Val Ala Gly Gly Tyr Ser Gln Gly Ala Ala Leu Ile Ala
100 105 110

35 Ala Ala Val Ser Glu Leu Ser Gly Ala Val Lys Glu Gln Val Lys Gly
115 120 125

40 Val Ala Leu Phe Gly Tyr Thr Gln Asn Leu Gln Asn Arg Gly Gly Ile
130 135 140

45 Pro Asn Tyr Pro Arg Glu Arg Thr Lys Val Phe Cys Asn Val Gly Asp
145 150 155 160

50 Ala Val Cys Thr Gly Thr Leu Ile Ile Thr Pro Ala His Leu Ser Tyr
165 170 175

55 Thr Ile Glu Ala Arg Gly Glu Ala Ala Arg Phe Leu Arg Asp Arg Ile
180 185 190

Arg Ala

<210> 2
<211> 199
<212> PRT
<213> Fusarium solani pisi

<220>
<221> mat_peptide
<222> (1)..(199)

<400> 2

Gly Arg Thr Thr Arg Asp Asp Leu Ile Asn Gly Asn Ser Ala Ser Cys
 1 5 10 15

5 Ala Asp Val Ile Phe Ile Tyr Ala Arg Gly Ser Thr Glu Thr Gly Asn
 20 25 30

10 Leu Gly Thr Leu Gly Pro Ser Ile Ala Ser Asn Leu Glu Ser Ala Phe
 35 40 45

15 Gly Lys Asp Gly Val Trp Ile Gln Gly Val Gly Gly Ala Tyr Arg Ala
 50 55 60

20 Thr Leu Gly Asp Asn Ala Leu Pro Arg Gly Thr Ser Ser Ala Ala Ile
 65 70 75 80

25 Arg Glu Met Leu Gly Leu Phe Gln Gln Ala Asn Thr Lys Cys Pro Asp
 85 90 95

30 Ala Thr Leu Ile Ala Gly Gly Tyr Ser Gln Gly Ala Ala Leu Ala Ala
 100 105 110

35 Ala Ser Ile Glu Asp Leu Asp Ser Ala Ile Arg Asp Lys Ile Ala Gly
 115 120 125

40 Thr Val Leu Phe Gly Tyr Thr Lys Asn Leu Gln Asn Arg Gly Arg Ile
 130 135 140

45 Pro Asn Tyr Pro Ala Asp Arg Thr Lys Val Phe Cys Asn Thr Gly Asp
 145 150 155 160

50 Leu Val Cys Thr Gly Ser Leu Ile Val Ala Ala Pro His Leu Ala Tyr
 165 170 175

55 Gly Pro Asp Ala Arg Gly Pro Ala Pro Glu Phe Leu Ile Glu Lys Val
 180 185 190

60 Arg Ala Val Arg Gly Ser Ala
 195

Claims

- 50 1. A process for degrading zearalenone in a feed product which process comprises treating said feed product with a cutinase.
- 55 2. The process according to claim 1 wherein the dosage of the cutinase is 0.01-100 mg enzyme protein pr. kg dry matter, preferably 0.1-10 mg enzyme protein pr. kg dry matter, or more preferably 1-5 mg enzyme protein pr. kg dry matter.
- 60 3. The process according to any of claims 1 or 2 wherein the feed product is a grain based feed product.

4. The process according to any of claims 1 to 3 wherein the feed product comprises one or more selected from corn, wheat, barley, rye, rice, sorghum and millet.
5. The process according to any of claims 1 to 4 wherein the feed product is an animal feed composition.
6. The process according to any of claims 1 to 5 wherein the feed product is a by-product from a fermentation process.
7. The process according to any of claims 1 to 6 wherein the feed product comprises brewer's spent grain, distiller's spent grain, distiller's wet grain, and/or distiller's dried grain.
10. The process according to any of claims 1 to 7 wherein the feed product is a swine feed product.
9. The process according to any of claims 1 to 8 wherein the cutinase is a cutinase having the sequence shown in SEQ ID NO:1 or a homologous sequence.
15. The process according to any of claims 1 to 9 wherein the cutinase is a variant of the cutinase shown in SEQ ID NO:1 comprising one or more, including all of the substitutions G8D, N15D, S48E, A88H, N91H, A130V and R189V.
20. The process according to any of claims 1 to 10 wherein the cutinase is a cutinase having the sequence shown in SEQ ID NO:2 or a homologous sequence.
12. A use of a cutinase for degrading the mycotoxin zearalenone.
13. The use according to claim 12 in a grain based feed product.
25. The use according to any of claims 12 to 13, wherein the dosage of the cutinase is 0.01-100 mg enzyme protein pr. kg dry matter, preferably 0.1-10 mg enzyme protein pr. kg dry matter, or more preferably 1-5 mg enzyme protein pr. kg dry matter.

Patentansprüche

1. Verfahren zum Abbauen von Zearalenon in einem Futtermittelprodukt, wobei das Verfahren Behandeln des Futtermittelprodukts mit einer Cutinase umfasst.
2. Verfahren nach Anspruch 1, wobei die Dosierung der Cutinase 0,01-100 mg Enzymprotein pro kg Trockenmasse beträgt, bevorzugt 0,1-10 mg Enzymprotein pro kg Trockenmasse, oder weiter bevorzugt 1-5 mg Enzymprotein pro kg Trockenmasse.
3. Verfahren nach einem beliebigen der Ansprüche 1 oder 2, wobei das Futtermittelprodukt ein auf Getreide basierendes Futtermittelprodukt ist.
4. Verfahren nach einem beliebigen der Ansprüche 1 bis 3, wobei das Futtermittelprodukt eines oder mehrere ausgewählt aus Mais, Weizen, Gerste, Roggen, Reis, Sorghum und Hirse umfasst.
45. 5. Verfahren nach einem beliebigen der Ansprüche 1 bis 4, wobei das Futtermittelprodukt eine Tierfuttermittelzusammensetzung ist.
6. Verfahren nach einem beliebigen der Ansprüche 1 bis 5, wobei das Futtermittelprodukt ein Nebenprodukt aus einem Fermentationsverfahren ist.
7. Verfahren nach einem beliebigen der Ansprüche 1 bis 6, wobei das Futtermittelprodukt Biertreber, Schlempe (distiller's spent grain), flüssige Schlempe (distiller's wet grain) und/oder getrocknete Schlempe (distiller's dried grain) umfasst.
55. 8. Verfahren nach einem beliebigen der Ansprüche 1 bis 7, wobei das Futtermittelprodukt ein Schweinefuttermittelprodukt ist.

9. Verfahren nach einem beliebigen der Ansprüche 1 bis 8, wobei die Cutinase eine Cutinase mit der in SEQ ID NO: 1 gezeigten Sequenz oder einer homologen Sequenz ist.
- 5 10. Verfahren nach einem beliebigen der Ansprüche 1 bis 9, wobei die Cutinase eine Variante der in SEQ ID NO: 1 gezeigten Cutinase ist, umfassend eine oder mehrere, einschließlich alle, der Substitutionen G8D, N15D, S48E, A88H, N91H, A130V und R189V.
- 10 11. Verfahren nach einem beliebigen der Ansprüche 1 bis 10, wobei die Cutinase eine Cutinase mit der in SEQ ID NO: 2 gezeigten Sequenz oder einer homologen Sequenz ist.
- 15 12. Verwendung einer Cutinase zum Abbauen des Mycotoxins Zearalenon.
13. Verwendung nach Anspruch 12 in einem auf Getreide basierenden Futtermittelprodukt.
14. Verwendung nach einem beliebigen der Ansprüche 12 bis 13, wobei die Dosierung der Cutinase 0,01-100 mg Enzymprotein pro kg Trockenmasse beträgt, bevorzugt 0,1-10 mg Enzymprotein pro kg Trockenmasse, oder weiter bevorzugt 1-5 mg Enzymprotein pro kg Trockenmasse.
- 20 **Revendications**
1. Procédé de dégradation de la zéaralénone dans un produit alimentaire pour animaux, ledit procédé comprenant le traitement dudit produit alimentaire pour animaux avec une cutinase.
- 25 2. Procédé selon la revendication 1, dans lequel le dosage de la cutinase est de 0,01 à 100 mg de protéine enzymatique par kg de matière sèche, de préférence de 0,1 à 10 mg de protéine enzymatique par kg de matière sèche, ou de manière encore plus préférée de 1 à 5 mg de protéine enzymatique par kg de matière sèche.
- 30 3. Procédé selon l'une quelconque des revendications 1 ou 2, dans lequel le produit alimentaire pour animaux est un produit alimentaire pour animaux à base de grains.
4. Procédé selon l'une quelconque des revendications 1 à 3, dans lequel le produit alimentaire pour animaux comprend un ou plusieurs choix parmi le maïs, le blé, l'orge, le seigle, le riz, le sorgho et le millet.
- 35 5. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel le produit alimentaire pour animaux est une composition alimentaire pour animaux.
6. Procédé selon l'une quelconque des revendications 1 à 5, dans lequel le produit alimentaire pour animaux est un sous-produit issu d'un processus de fermentation.
- 40 7. Procédé selon l'une quelconque des revendications 1 à 6, dans lequel le produit alimentaire pour animaux comprend des drêches de brasserie, des drêches de distillerie, des drêches de distillerie humides, et/ou des drêches de distillerie séchées.
8. Procédé selon l'une quelconque des revendications 1 à 7, dans lequel le produit alimentaire pour animaux est un produit alimentaire pour les porcs.
- 45 9. Procédé selon l'une quelconque des revendications 1 à 8, dans lequel la cutinase est une cutinase ayant la séquence représentée par SEQ ID NO: 1 ou une séquence homologue.
- 50 10. Procédé selon l'une quelconque des revendications 1 à 9, dans lequel la cutinase est une variante de la cutinase représentée dans SEQ ID NO: 1 comprenant une ou plusieurs, y compris toutes les substitutions G8D, N15D, S48E, A88H, N91H, A130V et R189V.
- 55 11. Procédé selon l'une quelconque des revendications 1 à 10, dans lequel la cutinase est une cutinase ayant la séquence représentée par SEQ ID NO: 2 ou une séquence homologue.
12. Utilisation d'une cutinase pour la dégradation de la mycotoxine zéaralénone.

13. Utilisation selon la revendication 12, dans un produit alimentaire pour animaux à base de grains.
14. Utilisation selon l'une quelconque des revendications 12 ou 13, dans laquelle le dosage de la cutinase est de 0,01
5 à 100 mg de protéine enzymatique par kg de matière sèche, de préférence de 0,1 à 10 mg de protéine enzymatique
par kg de matière sèche, ou de manière encore plus préférée de 1 à 5 mg de protéine enzymatique par kg de
matière sèche.

10

15

20

25

30

35

40

45

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 9612414 A [0010]
- US 2003073239 A [0018]
- US 5827719 A [0020] [0021]
- WO 9009446 A [0020]
- WO 9414964 A [0020] [0021]
- WO 9403578 A [0020]
- WO 0034450 A [0020]
- WO 0192502 A [0020] [0021]

Non-patent literature cited in the description

- NEEDLEMAN ; WUNSCH. *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0032]
- RICE et al. EMBOSS: The European Molecular Biology Open Software Suite. *Trends in Genetics*, 2000, vol. 16, 276-277 [0032]
- PEARSON, W.R. Bioinformatics Methods and Protocols. 1999, 185-219 [0034]
- SMEDSGAARD. *J. Chromatogr. A*, 1997, vol. 760, 264-270 [0036]