



(22) **Date de dépôt/Filing Date:** 2006/03/27
(41) **Mise à la disp. pub./Open to Public Insp.:** 2006/10/12
(62) **Demande originale/Original Application:** 2 603 223
(30) **Priorité/Priority:** 2005/04/06 (DE10 2005 015 677.0)


(51) **Cl.Int./Int.Cl.** **A01N 43/36** (2006.01),
A01N 31/00 (2006.01), **A01N 37/00** (2006.01),
A01N 43/00 (2006.01), **A01N 47/00** (2006.01),
A01N 53/12 (2006.01), **A01N 55/10** (2006.01),
A01N 57/00 (2006.01), **A01N 59/26** (2006.01),
A01P 3/00 (2006.01)

(71) **Demandeur/Applicant:**
BAYER CROPSCIENCE AG, DE

(72) **Inventeurs/Inventors:**
WACHENDORFF-NEUMANN, ULRIKE, DE;
DAHMEN, PETER, DE;
DUNKEL, RALF, FR

(74) **Agent:** FETHERSTONHAUGH & CO.

(54) **Titre : COMBINAISONS D'AGENTS ACTIFS FONGICIDES SYNERGIQUES**
(54) **Title: SYNERGISTIC FUNGICIDAL ACTIVE SUBSTANCE COMBINATIONS**



(57) **Abrégé/Abstract:**


The novel active compound combinations of a carboxamide of the general formula (I) (group 1) (see formula I) in which R, G, R¹ and A have the meanings given in the description and the active compound groups (2) to (23) listed in the description have very good fungicidal properties.



30725-1148D1

Abstract

The novel active compound combinations of a carboxamide of the general formula (I) (group 1)



in which R, G, R¹ and A have the meanings given in the description

and the active compound groups (2) to (23) listed in the description have very good fungicidal properties.

Synergistic fungicidal active substance combinations

This is a first divisional application of Canadian Patent Application No. 2,603,223, filed March 27, 2006. It should be understood that the expression "the present invention" or the like used in this specification encompasses not only the subject matter of this first divisional application but that of the parent application and a second divisional also.

The present invention relates to novel active compound combinations comprising, firstly, known carboxamides and, secondly, further known fungicidally active compounds, which combinations are highly suitable for controlling unwanted phytopathogenic fungi.

10


It is already known that certain carboxamides have fungicidal properties: for example *N*-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide from EP-A 0 737 682. The activity of these compounds is good; however, at low application rates it is sometimes unsatisfactory. Furthermore, it is already known that numerous triazole derivatives, aniline derivatives, dicarboximides and other heterocycles can be used for controlling fungi (cf. EP-A 0 040 345, DE-A 22 01 063, DE-A 23 24 010, Pesticide Manual, 9th Edition (1991), pages 249 and 827, EP-A 0 382 375 and EP-A 0 515 901). However, the activity of these compounds, too, is not always sufficient at low application rates. Furthermore, it is already known that 1-(3,5-dimethylisoxazole-4-sulfonyl)-2-chloro-6,6-difluoro-[1,3]-dioxolo-[4,5*f*]-benzimidazole has fungicidal properties (cf. WO 97/06171). Finally, it is also known that substituted halopyrimidines have fungicidal properties (cf. DE-A1-196 46 407, EP-B-712 396).

15

Also known are various mixtures of alkylthienylcarboxamides. The activity of these mixtures, too, is sometimes unsatisfactory (cf. JP-A 11-292715, JP-A 11-302107, JP-A 11-302108, JP-A 11-302109, JP-A 11-302110, JP-A 11-302111, JP-A 2001-72511, JP-A 2001-72512, JP-A 2001-72513, JP-A 11-322513, JP-A 11-322514, JP-A 2000-53506 and JP-A 2000-53507).

20

This invention now provides novel active compound combinations which have very good fungicidal properties and comprise a carboxamide of the general formula (I) (group 1)




(I), in which

30

- R represents hydrogen or methyl,
- G represents hydrogen, fluorine or methyl,
- R¹ represents hydrogen, halogen, C₁-C₃-alkyl or C₁-C₃-haloalkyl having 1 to 7 fluorine, chlorine and/or bromine atoms,
- A represents one of the radicals A1 or A2 below:

30725-1148D1

- 2 -




- X represents CH or N,
 R² represents methyl, difluoromethyl or trifluoromethyl,
 R³ represents hydrogen or fluorine,
 5 R⁴ represents difluoromethyl or trifluoromethyl,


and at least one active compound selected from groups (2) to (23) below:

Group (2) Strobilurins

- 10 (2-1) fluoxastrobin (known from DE-A 196 02 095) of the formula




- (2-2) (2E)-2-(2-{{[6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl}-2-(methoxyimino)-N-methylethanamide (known from DE-A 196 46 407, EP-B 0 712 396) of the formula



15


- (2-3) trifloxystrobin (known from EP-A 0 460 575) of the formula




- (2-4) (2E)-2-(methoxyimino)-N-methyl-2-(2-{{[1-3-(trifluoromethyl)phenyl]ethylidene}amino]oxy}methyl}phenyl)ethanamide (known from EP-A 0 569 384) of the formula

30725-1148D1


- 3 -




- (2-5) (2E)-2-(methoxyimino)-N-methyl-2-{2-[(E)-({1-[3-(trifluoromethyl)phenyl]ethoxy}imino)-methyl]phenyl}ethanamide (known from EP-A 0 596 254) of the formula



- 5 (2-6) orysastrobin (known from DE-A 195 39 324) of the formula




- (2-7) 5-methoxy-2-methyl-4-(2-(((1E)-1-[3-(trifluoromethyl)phenyl]ethyldiene)amino)oxy)-methyl}phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (known from WO 98/23155) of the formula



10


- (2-8) dimoxystrobin (known from EP-A 0 398 692) of the formula




- (2-9) picoxystrobin (known from EP-A 0 278 595) of the formula

30725-1148D1


- 4 -




(2-10) pyraclostrobin (known from DE-A 44 23 612) of the formula

5 Group (3) Triazoles


(3-1) azaconazole (known from DE-A 25 51 560) of the formula




(3-2) etaconazole (known from DE-A 25 51 560) of the formula



10 (3-3) propiconazole (known from DE-A 25 51 560) of the formula




(3-4) difenoconazole (known from EP-A 0 112 284) of the formula




(3-5) bromuconazole (known from EP-A 0 258 161) of the formula

30725-1148D1


- 5 -



(3-6) cyproconazole (known from DE-A 34 06 993) of the formula




(3-7) hexaconazole (known from DE-A 30 42 303) of the formula




5


(3-8) penconazole (known from DE-A 27 35 872) of the formula



(3-9) myclobutanil (known from EP-A 0 145 294) of the formula




10 (3-10) tetraconazole (known from EP-A 0 234 242) of the formula




30725-1148D1

- 6 -


(3-11) flutriafol (known from EP-A 0 015 756) of the formula




(3-12) epoxiconazole (known from EP-A 0 196 038) of the formula




5 (3-13) flusilazole (known from EP-A 0 068 813) of the formula



(3-14) simeconazole (known from EP-A 0 537 957) of the formula




(3-15) prothioconazole (known from WO 96/16048) of the formula



10


(3-16) fenbuconazole (known from DE-A 37 21 786) of the formula




30725-1148D1

- 7 -


(3-17) tebuconazole (known from EP-A 0 040 345) of the formula




(3-18) ipconazole (known from EP-A 0 329 397) of the formula




5 (3-19) metconazole (known from EP-A 0 329 397) of the formula



(3-20) triticonazole (known from EP-A 0 378 953) of the formula




(3-21) bitertanol (known from DE-A 23 24 010) of the formula




10

(3-22) triadimenol (known from DE-A 23 24 010) of the formula




(3-23) triadimefon (known from DE-A 22 01 063) of the formula




30725-1148D1

- 8 -


(3-24) fluquinconazole (known from EP-A 0 183 458) of the formula




(3-25) quinconazole (known from EP-A 0 183 458) of the formula

5 Group (4) Sulfenamides

(4-1) dichlofluanid (known from DE-A 11 93 498) of the formula




(4-2) tolylfuanid (known from DE-A 11 93 498) of the formula




10

Group (5) Valinamides

(5-1) iprovalicarb (known from DE-A 40 26 966) of the formula



(5-2) benthiavalicarb (known from WO 96/04252) of the formula




15

30725-1148D1


- 9 -

Group (6) Carboxamides


- (6-1) 2-chloro-N-(1,1,3-trimethylindan-4-yl)nicotinamide (known from EP-A 0 256 503) of the formula




- 5 (6-2) boscalid (known from DE-A 195 31 813) of the formula



- (6-3) furametpyr (known from EP-A 0 315 502) of the formula




- (6-4) ethaboxam (known from EP-A 0 639 574) of the formula




10

- (6-5) fenhexamid (known from EP-A 0 339 418) of the formula




- (6-6) carpropamid (known from EP-A 0 341 475) of the formula




30725-1148D1

- 10 -

- (6-7) 2-chloro-4-(2-fluoro-2-methylpropionylamino)-N,N-dimethylbenzamide
(known from EP-A 0 600 629) of the formula




- (6-8) picobenzamid (known from WO 99/42447) of the formula




5

- (6-9) zoxamide (known from EP-A 0 604 019) of the formula




- (6-10) 3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide (known from WO 99/24413) of the formula




10


- (6-11) carboxin (known from US 3,249,499) of the formula



- (6-12) tiadinil (known from US 6,616,054) of the formula



- 15 (6-13) siltiofam (known from WO 96/18631) of the formula




30725-1148D1

- 11 -

Group (7) Dithiocarbamates

(7-1) mancozeb (known from DE-A 12 34 704) having the IUPAC name
Manganese ethylenebis(dithiocarbamate) (polymeric) complex with zinc salt


(7-2) maneb (known from US 2,504,404) of the formula




5

(7-3) metiram (known from DE-A 10 76 434) having the IUPAC name
Zinc ammoniate ethylenebis(dithiocarbamate) – poly(ethylenethiuram disulfide)


(7-4) propineb (known from GB 935 981) of the formula




10 (7-5) thiram (known from US 1,972,961) of the formula



(7-6) zineb (known from DE-A 10 81 446) of the formula



(7-7) ziram (known from US 2,588,428) of the formula




15

Group (8) Acylalanines


(8-1) benalaxyl (known from DE-A 29 03 612) of the formula

30725-1148D1


- 12 -



(8-2) furalaxyl (known from DE-A 25 13 732) of the formula




(8-3) benalaxyl-M of the formula




5

Group (9): Anilinopyrimidines

(9-1) cyprodinil (known from EP-A 0 310 550) of the formula




10 (9-2) pyrimethanil (known from DD 151 404) of the formula

Group (10): Benzimidazoles

(10-1) 6-chloro-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2,2-difluoro-5H-[1,3]dioxolo[4,5-f]-benzimidazole (known from WO 97/06171) of the formula


15




30725-1148D1

- 13 -


(10-2) benomyl (known from US 3,631,176) of the formula




(10-3) carbendazim (known from US 3,010,968) of the formula




5 (10-4) chlorfenazole of the formula



(10-5) fuberidazole (known from DE-A 12 09 799) of the formula




(10-6) thiabendazole (known from US 3,206,468) of the formula




10

Group (11): Carbamates


(11-1) propamocarb (known from US 3,513,241) of the formula



15 (11-2) propamocarb hydrochloride (known from US 3,513,241) of the formula



(11-3) propamocarb-fosetyl of the formula




30725-1148D1


- 14 -

Group (12): Dicarboximides

(12-1) captafol (known from US 3,178,447) of the formula




(12-2) folpet (known from US 2,553,770) of the formula




5


(12-3) iprodione (known from DE-A 21 49 923) of the formula




(12-4) procymidone (known from DE-A 20 12 656) of the formula



10 (12-5) vinclozolin (known from DE-A 22 07 576) of the formula

Group (13): Guanidines

(13-1) dodine (known from GB 11 03 989) of the formula




15

(13-2) guazatine (known from GB 11 14 155)


30725-1148D1

- 15 -


(13-3) iminoctadine triacetate (known from EP-A 0 155 509) of the formula

Group (14): Imidazoles


5 (14-1) cyazofamid (known from EP-A 0 298 196) of the formula



(14-2) prochloraz (known from DE-A 24 29 523) of the formula




(14-3) triazoxide (known from DE-A 28 02 488) of the formula




10

(14-4) pefurazoate (known from EP-A 0 248 086) of the formula

Group (15): Morpholines


15 (15-1) aldimorph (known from DD 140 041) of the formula




30725-1148D1

- 16 -


(15-2) tridemorph (known from GB 988 630) of the formula




(15-3) dodemorph (known from DE-A 25 432 79) of the formula



5 (15-4) fenpropimorph (known from DE-A 26 56 747) of the formula


Group (16): Pyrroles

(16-1) fenciclonil (known from EP-A 0 236 272) of the formula




10

(16-2) pyrrolnitrine (known from JP 65-25876) of the formula

Group (17): Phosphonates

15 (17-1) phosphonic acid (known as Chemikalie) of the formula




30725-1148D1

- 17 -


Group (18): Phenylethanamide (known from WO 96/23793, in each case as E or Z isomer, preferably as E isomer)

(18-1) the compound 2-(2,3-dihydro-1H-inden-5-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide of the formula




5

(18-2) the compound N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)-2-(5,6,7,8-tetrahydronaphthalen-2-yl)acetamide of the formula




(18-3) the compound 2-(4-chlorophenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide of the formula


10



(18-4) the compound 2-(4-bromophenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide of the formula




15 (18-5) the compound 2-(4-methylphenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide of the formula




(18-6) the compound 2-(4-ethylphenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide of the formula

30725-1148D1

- 18 -


Group (19): Fungicides

(19-1) acibenzolar-S-methyl (known from EP-A 0 313 512) of the formula




5


(19-2) edifenphos (known from DE-A 14 93 736) of the formula




(19-3) famoxadone (known from EP-A 0 393 911) of the formula



10 (19-4) oxadixyl (known from DE-A 30 30 026) of the formula




(19-5) spiroxamine (known from DE-A 37 35 555) of the formula




(19-6) dithianon (known from JP-A 44-29464) of the formula

30725-1148D1


- 19 -



(19-7) metrafenone (known from EP-A 0 897 904) of the formula




(19-8) fenamidone (known from EP-A 0 629 616) of the formula




5

(19-9) 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)one (known from WO 99/14202) of the formula




(19-10) probenazole (known from US 3,629,428) of the formula




10

(19-11) isoprothiolane (known from US 3,856,814) of the formula




(19-12) kasugamycin (known from GB 1 094 567) of the formula




30725-1148D1

- 20 -


(19-13) phthalide (known from JP-A 57-55844) of the formula




(19-14) N-({4-[(cyclopropylamino)carbonyl]phenyl}sulfonyl)-2-methoxybenzamide of the formula




5 (19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide (known from WO 01/87822) of the formula




(19-16) quinoxifen (known from EP-A 0 326 330) of the formula



10 (19-17) proquinazid (known from WO 94/26722) of the formula

Group (20): (Thio)urea derivatives

(20-1) pencycuron (known from DE-A 27 32 257) of the formula




15


(20-2) thiophanate-ethyl (known from DE-A 18 06 123) of the formula

30725-1148D1




- 21 -

Group (21): Amides

(21-1) fenoxanil (known from EP-A 0 262 393) of the formula




(21-2) diclocymet (known from JP-A 7-206608) of the formula


Group (22): Triazolopyrimidines10 (22-1) 5-chloro-*N*-[(*IS*)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine (known from US 5,986,135) of the formula(22-2) 5-chloro-*N*-[(*IR*)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine (known from WO 02/38565) of the formula(22-3) 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-*a*]pyrimidine (known from US 5,593,996) of the formula

30725-1148D1


- 22 -




- (22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine (known from DE-A 101 24 208) of the formula




- 5 (22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1S)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine (known from US 5,593,996) of the formula




- Group (23): Iodochromones
- 10 (23-1) 2-butoxy-6-iodo-3-propylbenzopyran-4-one (known from WO 03/014103) of the formula



- (23-2) 2-ethoxy-6-iodo-3-propylbenzopyran-4-one (known from WO 03/014103) of the formula




- (23-3) 6-iodo-2-propoxy-3-propylbenzopyran-4-one (known from WO 03/014103) of the formula




30725-1148D1

- 23 -


(23-4) 2-but-2-ynyloxy-6-iodo-3-propylbenzopyran-4-one (known from WO 03/014103) of the formula




5 (23-5) 6-iodo-2-(1-methylbutoxy)-3-propylbenzopyran-4-one (known from WO 03/014103) of the formula



(23-6) 2-but-3-enyloxy-6-iodobenzopyran-4-one (known from WO 03/014103) of the formula



(23-7) 3-butyl-6-iodo-2-isopropoxybenzopyran-4-one (known from WO 03/014103) of the formula



10

Surprisingly, the fungicidal activity of the active compound combinations according to the invention is substantially higher than the sum of the activities of the individual active compounds. Thus, an unforeseeable true synergistic effect is present, and not just an addition of activities.

15 The formula (I) provides a general definition of the compounds of group (1).

Preference is given to carboxamides of the formula (I) in which

R represents hydrogen or methyl,


G represents hydrogen or methyl,

20 R¹ represents hydrogen, fluorine, chlorine, methyl, ethyl, n-propyl, isopropyl, monofluoromethyl, difluoromethyl, trifluoromethyl, monochloromethyl, dichloromethyl or trichloromethyl,

A represents one of the radicals A1 or A2 below:

30725-1148D1

- 24 -



- X represents CH or N,
 R² represents methyl, difluoromethyl or trifluoromethyl,
 R³ represents hydrogen or fluorine,
 5 R⁴ represents difluoromethyl or trifluoromethyl.

The formula (I) comprises in particular the following preferred mixing partners of group (1):

- (1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide
- 10 (1-2) 1-methyl-3-(trifluoromethyl)-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
- (1-3) 1-methyl-N-[2-(3-methylbutyl)-3-thienyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide
- (1-4) N-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide
- 15 (1-5) 3-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrazole-4-carboxamide
- (1-6) 3-(difluoromethyl)-1-methyl-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
- (1-7) 3-(difluoromethyl)-1-methyl-N-[2-(3-methylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
- (1-8) 3-(difluoromethyl)-N-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrazole-4-carboxamide
- 20 (1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide
- (1-10) 5-fluoro-1,3-dimethyl-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
- (1-11) 5-fluoro-1,3-dimethyl-N-[2-(3-methylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
- (1-12) N-[2-(3,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide
- (1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide
- 25 (1-14) 1-methyl-N-[2-(3-methylbutyl)-3-thienyl]-3-(trifluoromethyl)-1H-pyrrole-4-carboxamide
- (1-15) N-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrrole-4-carboxamide
- (1-16) N-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrrole-4-carboxamide
- (1-17) 4-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrrole-3-carboxamide
- (1-18) 4-(difluoromethyl)-1-methyl-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1H-pyrrole-3-
- 30 carboxamide
- (1-19) 4-(difluoromethyl)-1-methyl-N-[2-(3-methylbutyl)-3-thienyl]-1H-pyrrole-3-carboxamide
- (1-20) 4-(difluoromethyl)-N-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrrole-3-carboxamide
- (1-21) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-fluoro-1,4-dimethyl-1H-pyrrole-3-carboxamide

30725-1148D1

- 25 -

- (1-22) 5-fluoro-1,4-dimethyl-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1H-pyrrole-3-carboxamide
 (1-23) 5-fluoro-1,4-dimethyl-N-[2-(3-methylbutyl)-3-thienyl]-1H-pyrrole-3-carboxamide
 (1-24) N-[2-(3,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,4-dimethyl-1H-pyrrole-3-carboxamide
 (1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 5 (1-26) 2-methyl-4-(trifluoromethyl)-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1,3-thiazole-5-
 carboxamide
 (1-27) 2-methyl-N-[2-(3-methylbutyl)-3-thienyl]-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 (1-28) N-[2-(3,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 (1-29) 4-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-1,3-thiazole-5-carboxamide
 10 (1-30) 4-(difluoromethyl)-2-methyl-N-[2-(1,3,3-trimethylbutyl)-3-thienyl]-1,3-thiazole-5-
 carboxamide
 (1-31) 4-(difluoromethyl)-2-methyl-N-[2-(3-methylbutyl)-3-thienyl]-1,3-thiazole-5-carboxamide
 (1-32) 4-(difluoromethyl)-N-[2-(3,3-dimethylbutyl)-3-thienyl]-2-methyl-1,3-thiazole-5-carboxamide

15 Emphasis is given to active compound combinations according to the invention comprising in addition to the carboxamide (1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide (group 1) one or more, preferably one, mixing partner from groups (2) to (23).

20 Emphasis is given to active compound combinations according to the invention comprising in addition to the carboxamide (1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide (group 1) one or more, preferably one, mixing partner from groups (2) to (23).


25 Emphasis is given to active compound combinations according to the invention comprising in addition to the carboxamide (1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide (group 1) one or more, preferably one, mixing partner from groups (2) to (23).

30 Emphasis is given to active compound combinations according to the invention comprising in addition to the carboxamide (1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide (group 1) one or more, preferably one, mixing partner from groups (2) to (23).


The compound (6-6) carpropamid has three asymmetrically substituted carbon atoms. Accordingly, the compound (6-7) can be present as a mixture of various isomers or else in the form of a single component. Particular preference is given to the compounds (1*S*,3*R*)-2,2-dichloro-*N*-[(1*R*)-1-(4-chlorophenyl)ethyl]-1-ethyl-3-methylcyclopropanecarboxamide of the formula

30725-1148D1

- 26 -



(1*R*,3*S*)-2,2-dichloro-*N*-[(1*R*)-1-(4-chlorophenyl)ethyl]-1-ethyl-3-methylcyclopropanecarboxamide of the formula



5

Preferred mixing partners are the following active compounds of groups (2) to (23):

- (2-1) fluoxastrobin, (2-2) (2*E*)-2-(2-{{[6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl}-2-(methoxyimino)-*N*-methylethanamide, (2-3) trifloxystrobin, (2-4) (2*E*)-2-(methoxyimino)-*N*-methyl-2-(2-{{((1*E*)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)ethanamide, (2-5) (2*E*)-2-(methoxyimino)-*N*-methyl-2-{2-[(*E*)-({1-[3-(trifluoromethyl)phenyl]ethoxy}imino)methyl]phenyl}ethanamide, (2-7) 5-methoxy-2-methyl-4-(2-{{((1*E*)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)-2,4-dihydro-3*H*-1,2,4-triazol-3-one, (2-8) dimoxystrobin, (2-9) picoxystrobin, (2-10) pyraclostrobin, (3-3) propiconazole, (3-4) difenoconazole, (3-6) cyproconazole, (3-7) hexaconazole, (3-8) penconazole, (3-9) myclobutanil, (3-10) tetraconazole, (3-13) flusilazole, (3-15) prothioconazole, (3-16) fenbuconazole, (3-17) tebuconazole, (3-21) bitertanol, (3-22) triadimenol, (3-23) triadimefon, (3-12) epoxiconazole, (3-19) metconazole, (3-24) fluquinconazole, (4-1) dichlofluanid, (4-2) tolylfluanid, (5-1) iprovalicarb, (5-2) bentiavalicarb, (6-2) boscalid, (6-4) ethaboxam, (6-5) fenhexamid, (6-6) carpropamid, (6-7) 2-chloro-4-[(2-fluoro-2-methylpropanoyl)amino]-*N,N*-dimethylbenzamide, (6-8) picobenzamid, (6-9) zoxamide, (6-10) 3,4-dichloro-*N*-(2-cyanophenyl)isothiazole-5-carboxamide, (7-1) mancozeb, (7-2) maneb, (7-4) propineb, (7-5) thiram, (7-6) zineb, (8-1) benalaxyl, (8-2) furalaxyl, (8-3) benalaxyl-M, (9-1) cyprodinil, (9-2) pyrimethanil, (10-1) 6-chloro-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2,2-difluoro-5*H*-[1,3]dioxolo[4,5-*f*]benzimidazole, (10-3) carbendazim, (11-1) propamocarb, (11-2) propamocarb hydrochloride, (11-3) propamocarb-fosetyl, (12-2) folpet, (12-3) iprodione, (12-4) procymidone, (13-1) dodine, (13-2) guazatine, (13-3) iminoctadine triacetate, (14-1) cyazofamid, (14-2) prochloraz, (14-3) triazoxide, (15-4) fenpropimorph, (17-1) phosphonic acid, (19-1) acibenzolar-*S*-methyl, (19-3) famoxadone, (19-4) oxadixyl, (19-5) spiroxamine, (19-8) fenamidone, (19-15) 2-(4-chlorophenyl)-*N*-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide, (19-16) quinoxifen, (19-17) proquinazid, (20-1) pencycuron, (22-1) 5-chloro-*N*-[(1*S*)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine, (22-2) 5-chloro-*N*-[(1*R*)-1,2-dimethylpropyl]-

30725-1148D1

- 27 -

6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine, (22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine, (22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-*N*-[(*1S*)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine, (23-1) 2-butoxy-6-iodo-3-propylbenzopyran-4-one, (23-2) 2-ethoxy-6-iodo-3-propylbenzopyran-4-one, (23-3) 6-iodo-2-propoxy-3-propylbenzopyran-4-one.

Particularly preferred mixing partners are the following active compounds of groups (2) to (23):

(2-1) fluoxastrobin, (2-2) (*2E*)-2-(2-{[6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}-phenyl)-2-(methoxyimino)-*N*-methylethanamide, (2-3) trifloxystrobin, (3-15) prothioconazole, (3-17) tebuconazole, (3-21) bitertanol, (3-22) triadimenol, (3-24) fluquinconazole, (4-1) dichlofluanid, (4-2) tolylfluanid, (5-1) iprovalicarb, (6-5) fenhexamid, (6-6) carpropamid, (6-8) picobenzamid, (7-4) propineb, (8-3) benalaxyl-M, (9-2) pyrimethanil, (10-3) carbendazim, (11-3) propamocarb-fosetyl, (12-3) iprodione, (14-2) prochloraz, (14-3) triazoxide, (19-5) spiroxamine, (19-15) 2-(4-chlorophenyl)-*N*-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide, (22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine, (22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-*N*-[(*1S*)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine.

Hereinbelow, preferred active compound combinations consisting of two groups of active compounds and comprising in each case at least one carboxamide of the formula (I) (group 1) and at least one active compound from the stated group (2) to (23) are described. These combinations are the active compound combinations A to R.

Preference is given to active compound combinations A to R in which the carboxamide of the formula (I) (group 1) is selected from the list below:

(1-1) *N*-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide
 (1-3) 1-methyl-*N*-[2-(3-methylbutyl)-3-thienyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide
 (1-5) 3-(difluoromethyl)-*N*-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrazole-4-carboxamide
 (1-7) 3-(difluoromethyl)-1-methyl-*N*-[2-(3-methylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
 (1-9) *N*-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide
 (1-11) 5-fluoro-1,3-dimethyl-*N*-[2-(3-methylbutyl)-3-thienyl]-1H-pyrazole-4-carboxamide
 (1-13) *N*-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide
 (1-15) *N*-[2-(3,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrrole-4-carboxamide
 (1-17) 4-(difluoromethyl)-*N*-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrrole-3-carboxamide
 (1-19) 4-(difluoromethyl)-1-methyl-*N*-[2-(3-methylbutyl)-3-thienyl]-1H-pyrrole-3-carboxamide
 (1-21) *N*-[2-(1,3-dimethylbutyl)-3-thienyl]-2-fluoro-1,4-dimethyl-1H-pyrrole-3-carboxamide

30725-1148D1

- 28 -

- (1-23) 5-fluoro-1,4-dimethyl-N-[2-(3-methylbutyl)-3-thienyl]-1H-pyrrole-3-carboxamide
 (1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 (1-27) 2-methyl-N-[2-(3-methylbutyl)-3-thienyl]-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 (1-29) 4-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-1,3-thiazole-5-carboxamide
 5 (1-31) 4-(difluoromethyl)-2-methyl-N-[2-(3-methylbutyl)-3-thienyl]-1,3-thiazole-5-carboxamide

Particular preference is given to active compound combinations A to R in which the carboxamide of the formula (I) (group 1) is selected from the list below:

- (1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-
 10 carboxamide
 (1-5) 3-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrazole-4-carboxamide
 (1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide
 (1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide
 (1-17) 4-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-1H-pyrrole-3-carboxamide
 15 (1-21) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-fluoro-1,4-dimethyl-1H-pyrrole-3-carboxamide
 (1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide
 (1-29) 4-(difluoromethyl)-N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-1,3-thiazole-5-carboxamide

- Preference is given to active compound combinations A in which the strobilurin (group 2) is selected
 20 from the list below: (2-1) fluoxastrobin, (2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl}oxy}phenyl)-2-(methoxyimino)-N-methylethanamide, (2-3) trifloxystrobin, (2-4) (2E)-2-(methoxyimino)-N-methyl-2-(2-{{(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino}oxy)methyl}phenyl)ethanamide, (2-5) (2E)-2-(methoxyimino)-N-methyl-2-{2-[(E)-({1-[3-(trifluoromethyl)phenyl]ethoxy}imino)methyl]phenyl}ethanamide, (2-6) oryastrobin, (2-7) 5-methoxy-2-methyl-4-(2-{{(1E)-
 25 1-[3-(trifluoromethyl)phenyl]ethylidene}amino}oxy)methyl}phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, (2-8) dimoxystrobin, (2-9) picoxystrobin, (2-10) pyraclostrobin.

- Particular preference is given to active compound combinations A in which the strobilurin (group 2) is selected from the following list: (2-1) fluoxastrobin, (2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl}oxy}phenyl)-2-(methoxyimino)-N-methylethanamide, (2-3) trifloxystrobin, (2-8) dimoxystrobin, (2-9) picoxystrobin, (2-10) pyraclostrobin.
 30

Emphasis is given to the active compound combinations A listed in table 1 below:

Table 1: Active compound combinations A

No.	Carboxamide of the formula (I)	Strobilurin (group 2)
A-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(2-1) fluoxastrobin
A-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-	(2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphen-

Table 1: Active compound combinations A

No.	Carboxamide of the formula (I)	Strobilurin (group 2)
	3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	oxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl)-2-(methoxyimino)-N-methylethanamide
A-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(2-3) trifloxystrobin
A-4	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(2-8) dimoxystrobin
A-5	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(2-9) picoxystrobin
A-6	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(2-10) pyraclostrobin
A-7	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-1) fluoxastrobin
A-8	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl)-2-(methoxyimino)-N-methylethanamide
A-9	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-3) trifloxystrobin
A-10	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-8) dimoxystrobin
A-11	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-9) picoxystrobin
A-12	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(2-10) pyraclostrobin
A-13	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-1) fluoxastrobin
A-14	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl)-2-(methoxyimino)-N-methylethanamide
A-15	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-3) trifloxystrobin
A-16	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-8) dimoxystrobin
A-17	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-9) picoxystrobin
A-18	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(2-10) pyraclostrobin
A-19	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-1) fluoxastrobin
A-20	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-2) (2E)-2-(2-{{6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]oxy}phenyl)-2-(methoxyimino)-N-methylethanamide
A-21	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-3) trifloxystrobin
A-22	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-8) dimoxystrobin
A-23	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-9) picoxystrobin
A-24	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(2-10) pyraclostrobin

30725-1148D1

- 30 -

Preference is given to active compound combinations B in which the triazole (group 3) is selected from the following list: (3-1) azaconazole, (3-2) etaconazole, (3-3) propiconazole, (3-4) difenoconazole, (3-5) bromuconazole, (3-6) cyproconazole, (3-7) hexaconazole, (3-8) penconazole, (3-9) myclobutanil, (3-10) tetraconazole, (3-11) flutriafol, (3-12) epoxiconazole, (3-13) flusilazole, (3-14) simeconazole, (3-15) prothioconazole, (3-16) fenbuconazole, (3-17) tebuconazole, (3-18) ipconazole, (3-19) metconazole, (3-20) triticonazole, (3-21) bitertanol, (3-22) triadimenol, (3-23) triadimefon, (3-24) fluquinconazole, (3-25) quinconazole.

Particular preference is given to active compound combinations B in which the triazole (group 3) is selected from the following list: (3-3) propiconazole, (3-4) difenoconazole, (3-6) cyproconazole, (3-7) hexaconazole, (3-15) prothioconazole, (3-17) tebuconazole, (3-19) metconazole, (3-21) bitertanol, (3-22) triadimenol, (3-24) fluquinconazole.

Emphasis is given to the active compound combinations B listed in table 2 below:

Table 2: Active compound combinations B

No.	Carboxamide of the formula (I)	Triazole (group 3)
B-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-3) propiconazole
B-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-4) difenoconazole
B-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-6) cyproconazole
B-4	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-7) hexaconazole
B-5	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-15) prothioconazole
B-6	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-17) tebuconazole
B-7	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-19) metconazole
B-8	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-21) bitertanol
B-9	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-22) triadimenol
B-10	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(3-24) fluquinconazole
B-11	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-3) propiconazole
B-12	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-4) difenoconazole
B-13	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-6) cyproconazole
B-14	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-7) hexaconazole
B-15	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-15) prothioconazole
B-16	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-	(3-17) tebuconazole

Table 2: Active compound combinations B

No.	Carboxamide of the formula (I)	Triazole (group 3)
	dimethyl-1H-pyrazole-4-carboxamide	
B-17	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-19) metconazole
B-18	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-21) bitertanol
B-19	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-22) triadimenol
B-20	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(3-24) fluquinconazole
B-21	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-3) propiconazole
B-22	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-4) difenoconazole
B-23	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-6) cyproconazole
B-24	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-7) hexaconazole
B-25	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-15) prothioconazole
B-26	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-17) tebuconazole
B-27	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-19) metconazole
B-28	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-21) bitertanol
B-29	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-22) triadimenol
B-30	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(3-24) fluquinconazole
B-31	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-3) propiconazole
B-32	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-4) difenoconazole
B-33	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-6) cyproconazole
B-34	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-7) hexaconazole
B-35	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-15) prothioconazole
B-36	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-17) tebuconazole
B-37	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-19) metconazole
B-38	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-21) bitertanol
B-39	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-22) triadimenol
B-40	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(3-24) fluquinconazole

30725-1148D1

- 32 -

Preference is given to active compound combinations C in which the sulfenamide (group 4) is selected from the following list: (4-1) dichlofluanid, (4-2) tolylfluanid.

Emphasis is given to the active compound combinations C listed in table 3 below:

Table 3: Active compound combinations C

No.	Carboxamide of the formula (I)	Sulfenamide (group 4)
C-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(4-1) dichlofluanid
C-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(4-2) tolylfluanid
C-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(4-1) dichlofluanid
C-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(4-2) tolylfluanid
C-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(4-1) dichlofluanid
C-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(4-2) tolylfluanid
C-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(4-1) dichlofluanid
C-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(4-2) tolylfluanid

5

Preference is given to active compound combinations D in which the valinamide (group 5) is selected from the following list: (5-1) iprovalicarb, (5-2) benthiavalicarb.

Emphasis is given to the active compound combinations D listed in table 4 below:

Table 4: Active compound combinations D

No.	Carboxamide of the formula (I)	Valinamide (group 5)
D-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(5-1) iprovalicarb
D-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(5-2) benthiavalicarb
D-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(5-1) iprovalicarb
D-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(5-2) benthiavalicarb
D-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(5-1) iprovalicarb
D-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(5-2) benthiavalicarb
D-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(5-1) iprovalicarb
D-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(5-2) benthiavalicarb

10

Preference is given to active compound combinations E in which the carboxamide (group 6) is selected from the list below: (6-1) 2-chloro-N-(1,1,3-trimethylindan-4-yl)nicotinamide, (6-2)

30725-1148D1

- 33 -

boscalid, (6-3) furametpyr, (6-4) ethaboxam, (6-5) fenhexamid, (6-6) carpropamid, (6-7) 2-chloro-4-(2-fluoro-2-methylpropionylamino)-N,N-dimethylbenzamide, (6-8) picobenzamid, (6-9) zoxamide, (6-10) 3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide, (6-11) carboxin, (6-12) tiadinil, (6-13) silthiofam.

5

Particular preference is given to active compound combinations E in which the carboxamide (group 6) is selected from the list below: (6-2) boscalid, (6-4) ethaboxam, (6-5) fenhexamid, (6-6) carpropamid, (6-7) 2-chloro-4-(2-fluoro-2-methylpropionylamino)-N,N-dimethylbenzamide, (6-8) picobenzamid, (6-9) zoxamide, (6-10) 3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide.

10

Very particular preference is given to active compound combinations E in which the carboxamide (group 6) is selected from the following list: (6-2) boscalid, (6-5) fenhexamid, (6-6) carpropamid, (6-8) picobenzamid.

15 Emphasis is given to the active compound combinations E listed in table 5 below:

Table 5: Active compound combinations E

No.	Carboxamide of the formula (I)	Carboxamide (group 6)
E-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(6-2) boscalid
E-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(6-5) fenhexamid
E-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(6-6) carpropamid
E-4	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(6-8) picobenzamid
E-5	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(6-2) boscalid
E-6	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(6-5) fenhexamid
E-7	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(6-6) carpropamid
E-8	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(6-8) picobenzamid
E-9	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(6-2) boscalid
E-10	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(6-5) fenhexamid
E-11	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(6-6) carpropamid
E-12	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(6-8) picobenzamid
E-13	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(6-2) boscalid
E-14	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(6-5) fenhexamid
E-15	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-	(6-6) carpropamid

30725-1148D1

- 34 -

Table 5: Active compound combinations E

No.	Carboxamide of the formula (I)	Carboxamide (group 6)
	(trifluoromethyl)-1,3-thiazole-5-carboxamide	
E-16	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(6-8) picobenzamid

Preference is given to active compound combinations F in which the dithiocarbamate (group 7) is selected from the following list: (7-1) mancozeb, (7-2) maneb, (7-4) propineb, (7-5) thiram, (7-6) zineb.

5

Particular preference is given to active compound combinations F in which the dithiocarbamate (group 7) is selected from the following list: (7-1) mancozeb, (7-4) propineb.

Emphasis is given to the active compound combinations F listed in table 6 below:

Table 6: Active compound combinations F

No.	Carboxamide of the formula (I)	Dithiocarbamate (group 7)
F-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(7-1) mancozeb
F-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(7-4) propineb
F-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(7-1) mancozeb
F-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(7-4) propineb
F-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(7-1) mancozeb
F-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(7-4) propineb
F-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(7-1) mancozeb
F-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(7-4) propineb

10

Preference is given to active compound combinations G in which the acylalanine (group 8) is selected from the following list: (8-1) benalaxyl, (8-2) furalaxyl, (8-5) benalaxyl-M.

Particular preference is given to active compound combinations G in which the acylalanine of the formula (VI) (group 8) is selected from the following list: (8-5) benalaxyl-M.

15

Emphasis is given to the active compound combinations G listed in table 7 below:

Table 7: Active compound combinations G

No.	Carboxamide of the formula (I)	Acylalanine (group 8)
G-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(8-5) benalaxyl-M
G-2	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-	(8-5) benalaxyl-M

30725-1148D1

- 35 -

Table 7: Active compound combinations G

No.	Carboxamide of the formula (I)	Acylalanine (group 8)
	dimethyl-1H-pyrazole-4-carboxamide	
G-3	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(8-5) benalaxyl-M
G-4	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(8-5) benalaxyl-M

Preference is given to active compound combinations H in which the anilinopyrimidine (group 9) is selected from the following list: (9-1) cyprodinil, (9-2) pyrimethanil.

- 5 Emphasis is given to the active compound combinations H listed in table 8 below:

Table 8: Active compound combinations H

No.	Carboxamide of the formula (I)	Anilinopyrimidine (group 9)
H-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(9-1) cyprodinil
H-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(9-2) pyrimethanil
H-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(9-1) cyprodinil
H-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(9-2) pyrimethanil
H-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(9-1) cyprodinil
H-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(9-2) pyrimethanil
H-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(9-1) cyprodinil
H-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(9-2) pyrimethanil

- Preference is given to active compound combinations I in which the benzimidazole (group 10) is selected from the following list: (10-1) 6-chloro-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2,2-difluoro-5H-[1,3]dioxolo-[4,5-f]benzimidazole, (10-2) benomyl, (10-3) carbendazim, (10-4) chlorfenazole, (10-5) fuberidazole, (10-6) thiabendazole.

Particular preference is given to active compound combinations I in which the benzimidazole of the formula (VIII) (group 10) is: (10-3) carbendazim.

- 15 Emphasis is given to the active compound combinations I listed in table 9 below:

Table 9: Active compound combinations I

No.	Carboxamide of the formula (I)	Benzimidazole (group 10)
I-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(10-3) carbendazim
I-2	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(10-3) carbendazim

30725-1148D1

- 36 -

Table 9: Active compound combinations I

No.	Carboxamide of the formula (I)	Benzimidazole (group 10)
I-3	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(10-3) carbendazim
I-4	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(10-3) carbendazim

Preference is given to active compound combinations J in which the carbamate (group 11) is selected from the list below: (11-1) propamocarb, (11-2) propamocarb hydrochloride, (11-3) propamocarb-fosetyl.

5

Emphasis is given to the active compound combinations J listed in table 10 below:

Table 10: Active compound combinations J

Nr.	Carboxamide of the formula (I)	Carbamate (group 11)
J-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(11-1) propamocarb
J-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(11-2) propamocarb hydrochloride
J-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(11-3) propamocarb-fosetyl
J-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(11-1) propamocarb
J-5	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(11-2) propamocarb hydrochloride
J-6	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(11-3) propamocarb-fosetyl
J-7	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(11-1) propamocarb
J-8	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(11-2) propamocarb hydrochloride
J-9	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(11-3) propamocarb-fosetyl
J-10	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(11-1) propamocarb
J-11	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(11-2) propamocarb hydrochloride
J-12	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(11-3) propamocarb-fosetyl

Preference is given to active compound combinations K in which the dicarboximide (group 12) is selected from the list below: (12-2) folpet, (12-3) iprodione.

10

Emphasis is given to the active compound combinations K listed in table 11 below:

Table 11: Active compound combinations K

No.	Carboxamide of the formula (I)	Dicarboximide (group 12)
K-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(12-2) folpet
K-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-	(12-3) iprodione

30725-1148D1

- 37 -

Table 11: Active compound combinations K

No.	Carboxamide of the formula (I)	Dicarboximide (group 12)
	(trifluoromethyl)-1H-pyrazole-4-carboxamide	
K-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(12-2) folpet
K-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(12-3) iprodione
K-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(12-2) folpet
K-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(12-3) iprodione
K-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(12-2) folpet
K-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(12-3) iprodione

Preference is given to active compound combinations L in which the guanidine (group 13) is selected from the list below: (13-1) dodine, (13-2) guazatine.

- 5 Emphasis is given to the active compound combinations L listed in table 12 below:

Table 12: Active compound combinations L

No.	Carboxamide of the formula (I)	Guanidine (group 13)
L-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(13-1) dodine
L-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(13-2) guazatine
L-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(13-1) dodine
L-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(13-2) guazatine
L-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(13-1) dodine
L-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(13-2) guazatine
L-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(13-1) dodine
L-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(13-2) guazatine

Preference is given to active compound combinations M in which the imidazole (group 14) is selected from the list below: (14-2) prochloraz, (14-3) triazoxide.

- 10 Emphasis is given to the active compound combinations M listed in table 13 below:

Table 13: Active compound combinations M

No.	Carboxamide of the formula (I)	Imidazole (group 14)
M-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(14-2) prochloraz
M-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-	(14-3) triazoxide

30725-1148D1

- 38 -

Table 13: Active compound combinations M

No.	Carboxamide of the formula (I)	Imidazole (group 14)
	(trifluoromethyl)-1H-pyrazole-4-carboxamide	
M-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(14-2) prochloraz
M-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(14-3) triazoxide
M-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(14-2) prochloraz
M-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(14-3) triazoxide
M-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(14-2) prochloraz
M-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(14-3) triazoxide

Preference is given to active compound combinations N in which the morpholine (group 15) is selected from the following list: (15-1) aldimorph, (15-2) tridemorph, (15-3) dodemorph, (15-4) fenpropimorph.

5

Particular preference is given to active compound combinations N in which the morpholine (group 15) is selected from the list below: (15-4) fenpropimorph.

Emphasis is given to the active compound combinations N listed in table 14 below:

Table 14: Active compound combinations N

No.	Carboxamide of the formula (I)	Morpholine (group 15)
N-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(15-4) fenpropimorph
N-2	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(15-4) fenpropimorph
N-3	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(15-4) fenpropimorph
N-4	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(15-4) fenpropimorph

10

Preference is given to active compound combinations O in which the fungicide (group 19) is selected from the list below: (19-1) acibenzolar-S-methyl, (19-3) famoxadone, (19-4) oxadixyl, (19-5) spiroxamine, (19-8) fenamidone, (19-14) N-({4-[(cyclopropylamino)carbonyl]phenyl}sulfonyl)-2-methoxybenzamide, (19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]-ethyl}-2-(prop-2-yn-1-yloxy)acetamide.

15

Particular preference is given to active compound combinations O in which the fungicide (group 19) is selected from the list below: (19-5) spiroxamine, (19-14) N-({4-[(cyclopropylamino)carbonyl]-

30725-1148D1

- 39 -

phenyl)sulfonyl)-2-methoxybenzamide, (19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide.

Emphasis is given to the active compound combinations O listed in table 15 below:

Table 15: Active compound combinations O

No.	Carboxamide of the formula (I)	Fungicide (group 19)
O-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(19-5) spiroxamine
O-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(19-14) N-({4-[(cyclopropylamino)-carbonyl]phenyl}sulfonyl)-2-methoxybenzamide
O-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide
O-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(19-5) spiroxamine
O-5	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(19-14) N-({4-[(cyclopropylamino)-carbonyl]phenyl}sulfonyl)-2-methoxybenzamide
O-6	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide
O-7	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(19-5) spiroxamine
O-8	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(19-14) N-({4-[(cyclopropylamino)-carbonyl]phenyl}sulfonyl)-2-methoxybenzamide
O-9	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide
O-10	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(19-5) spiroxamine
O-11	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(19-14) N-({4-[(cyclopropylamino)-carbonyl]phenyl}sulfonyl)-2-methoxybenzamide
O-12	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(19-15) 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide

5

Preference is given to the active compound combinations P in which the (thio)urea derivative (group 20) is selected from the list below: (20-1) pencycuron.

Emphasis is given to the active compound combinations P listed in table 16 below:

Table 16: Active compound combinations P

No.	Carboxamide of the formula (I)	(Thio)urea derivative (group 20)
P-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(20-1) pencycuron
P-2	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(20-1) pencycuron
P-3	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(20-1) pencycuron
P-4	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(20-1) pencycuron

Preference is given to active compound combinations Q in which the triazolopyrimidine (group 22) is selected from the list below: (22-1) 5-chloro-*N*-[(*IS*)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine, (22-2) 5-chloro-*N*-[(*IR*)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine, (22-3) 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-*a*]pyrimidine, (22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-*a*]pyrimidine.

10 Particular preference is given to active compound combinations Q in which the triazolopyrimidine (group 22) is selected from the list below: (22-1) 5-chloro-*N*-[(*IS*)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine, (22-2) 5-chloro-*N*-[(*IR*)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine, (22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-*a*]pyrimidine, (22-5) 5-chloro-6-
15 (2,4,6-trifluorophenyl)-*N*-[(*IS*)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-*a*]pyrimidine-7-amine.

Emphasis is given to the active compound combinations Q listed in table 17 below:

Table 17: Active compound combinations Q

No.	Carboxamide of the formula (I)	Triazolopyrimidine (group 22)
Q-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(22-1) 5-chloro- <i>N</i> -[(<i>IS</i>)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5- <i>a</i>]pyrimidine-7-amine
Q-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(22-2) 5-chloro- <i>N</i> -[(<i>IR</i>)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5- <i>a</i>]pyrimidine-7-amine
Q-3	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5- <i>a</i>]pyrimidine
Q-4	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(22-5) 5-chloro-6-(2,4,6-trifluorophenyl)- <i>N</i> -[(<i>IS</i>)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5- <i>a</i>]pyrimidine-7-amine
Q-5	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(22-1) 5-chloro- <i>N</i> -[(<i>IS</i>)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5- <i>a</i>]pyrimidine-7-amine

30725-1148D1

- 41 -

Table 17: Active compound combinations Q

No.	Carboxamide of the formula (I)	Triazolopyrimidine (group 22)
Q-6	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(22-2) 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-7	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine
Q-8	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1S)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-9	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(22-1) 5-chloro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-10	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(22-2) 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-11	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine
Q-12	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1S)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-13	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(22-1) 5-chloro-N-[(1S)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-14	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(22-2) 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine
Q-15	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(22-4) 5-chloro-6-(2,4,6-trifluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine
Q-16	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(22-5) 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1S)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine

Preference is given to active compound combinations R in which the iodochromone (group 23) is selected from the list below: (23-1) 2-butoxy-6-iodo-3-propylbenzopyran-4-one, (23-2) 2-ethoxy-6-iodo-3-propylbenzopyran-4-one, (23-3) 6-iodo-2-propoxy-3-propylbenzopyran-4-one, (23-4) 2-but-2-ynyloxy-6-iodo-3-propylbenzopyran-4-one, (23-5) 6-iodo-2-(1-methylbutoxy)-3-propylbenzopyran-4-one, (23-6) 2-but-3-enyloxy-6-iodobenzopyran-4-one, (23-7) 3-butyl-6-iodo-2-isopropoxybenzopyran-4-one.

Particular preference is given to active compound combinations T in which the iodochromone (group 23) is selected from the list below: (23-1) 2-butoxy-6-iodo-3-propylbenzopyran-4-one, (23-2) 2-ethoxy-6-iodo-3-propylbenzopyran-4-one.

Emphasis is given to the active compound combinations R listed in table 18 below:

30725-1148D1

- 42 -

Table 18: Active compound combinations R

No.	Carboxamide of the formula (I)	Iodochromone of the formula (XV)
R-1	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(23-1) 2-butoxy-6-iodo-3-propyl-benzopyran-4-one
R-2	(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	(23-2) 2-ethoxy-6-iodo-3-propyl-benzopyran-4-one
R-3	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(23-1) 2-butoxy-6-iodo-3-propyl-benzopyran-4-one
R-4	(1-9) N-[2-(1,3-dimethylbutyl)-3-thienyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide	(23-2) 2-ethoxy-6-iodo-3-propyl-benzopyran-4-one
R-5	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(23-1) 2-butoxy-6-iodo-3-propyl-benzopyran-4-one
R-6	(1-13) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide	(23-2) 2-ethoxy-6-iodo-3-propyl-benzopyran-4-one
R-7	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(23-1) 2-butoxy-6-iodo-3-propyl-benzopyran-4-one
R-8	(1-25) N-[2-(1,3-dimethylbutyl)-3-thienyl]-2-methyl-4-(trifluoromethyl)-1,3-thiazole-5-carboxamide	(23-2) 2-ethoxy-6-iodo-3-propyl-benzopyran-4-one

In addition to an active compound of the formula (I), the active compound combinations according to the invention comprise at least one active compound selected from the compounds of groups (2) to (23). In addition, they may also comprise further fungicidally active additives.

5

The synergistic effect is particularly pronounced when the active compounds in the active compound combinations according to the invention are present in certain weight ratios. However, the weight ratios of the active compounds in the active compound combinations can be varied within a relatively wide range. In general, the combinations according to the invention comprise active compounds of the formula (I) and a mixing partner of one of groups (2) to (23) in the mixing ratios given in an exemplary manner in table 19 below.

10

The mixing ratios are based on weight ratios. The ratio is to be understood as meaning active compound of formula (I):mixing partner.

15

Table 19: Mixing ratios

Mixing partner	preferred mixing ratio	particularly preferred mixing ratio
Group (2): Strobilurins	50:1 to 1:50	10:1 to 1:20
Group (3): Triazoles	50:1 to 1:50	20:1 to 1:20
Group (4): Sulfenamides	1:1 to 1:150	1:1 to 1:100
Group (5): Valinamides	50:1 to 1:50	10:1 to 1:20
Group (6): Carboxamides	50:1 to 1:50	20:1 to 1:20

30725-1148D1

- 43 -

Table 19: Mixing ratios

Mixing partner	preferred mixing ratio	particularly preferred mixing ratio
Group (7): Dithiocarbamates	1:1 to 1:150	1:1 to 1:100
Group (8): Acylalanines	10:1 to 1:150	5:1 to 1:100
Group (9): Anilinopyrimidines	5:1 to 1:50	1:1 to 1:20
Group (10): Benzimidazoles	10:1 to 1:50	5:1 to 1:20
Group (11): Carbamates	1:1 to 1:150	1:1 to 1:100
Group (12): (12-1)/(12-2)	1:1 to 1:150	1:5 to 1:100
Group (12): (12-3)/(12-4)/(12-5)	5:1 to 1:50	1:1 to 1:20
Group (13): Guanidines	100:1 to 1:150	20:1 to 1:100
Group (14): Imidazoles	50:1 to 1:50	10:1 to 1:20
Group (15): Morpholines	50:1 to 1:50	10:1 to 1:20
Group (16): Pyrroles	50:1 to 1:50	10:1 to 1:20
Group (17): Phosphonates	10:1 to 1:150	1:1 to 1:100
Group (18): Phenylethanamides	50:1 to 1:50	10:1 to 1:20
Group (19)	50:1 to 1:150	20:1 to 1:100
Group (20): (Thio)urea derivatives	50:1 to 1:50	10:1 to 1:20
Group (21): Amides	50:1 to 1:50	10:1 to 1:20
Group (22): Triazolopyrimidines	50:1 to 1:50	10:1 to 1:20
Group (23): Iodochromones	50:1 to 1:50	10:1 to 1:20

In each case, the mixing ratio is to be chosen such that a synergistic mixture is obtained. The mixing ratios between the compound of the formula (I) and a compound of one of the groups (2) to (23) may also vary between the individual compounds of a group.

- 5 The active compound combinations according to the invention have very good fungicidal properties and can be used for controlling phytopathogenic fungi and bacteria.

In crop protection, fungicides can be used for controlling Plasmodiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

- 10 In crop protection, bactericides can be used for controlling Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomyetaceae.

Some pathogens causing fungal and bacterial diseases which come under the generic names listed above may be mentioned as examples, but not by way of limitation:

Diseases caused by powdery mildew pathogens, such as, for example,

30725-1148D1

- 44 -

- Blumeria species, such as, for example, *Blumeria graminis*;
 Podosphaera species, such as, for example, *Podosphaera leucotricha*;
 Sphaerotheca species, such as, for example, *Sphaerotheca fuliginea*;
 Uncinula species, such as, for example, *Uncinula necator*;
- 5 Diseases caused by rust disease pathogens, such as, for example,
 Gymnosporangium species, such as, for example, *Gymnosporangium sabinae*
 Hemileia species, such as, for example, *Hemileia vastatrix*;
 Phakopsora species, such as, for example, *Phakopsora pachyrhizi* and *Phakopsora meibomia*;
 Puccinia species, such as, for example, *Puccinia recondita*;
- 10 Uromyces species, such as, for example, *Uromyces appendiculatus*;
- Diseases caused by pathogens from the group of the Oomycetes, such as, for example,
 Bremia species, such as, for example, *Bremia lactucae*;
 Peronospora species, such as, for example, *Peronospora pisi* or *P. brassicae*;
 Phytophthora species, such as, for example *Phytophthora infestans*;
- 15 Plasmopara species, such as, for example, *Plasmopara viticola*;
 Pseudoperonospora species, such as, for example, *Pseudoperonospora humuli* or
Pseudoperonospora cubensis;
 Pythium species, such as, for example, *Pythium ultimum*;
- Leaf blotch diseases and leaf wilt diseases caused, for example, by
- 20 Alternaria species, such as, for example, *Alternaria solani*;
 Cercospora species, such as, for example, *Cercospora beticola*;
 Cladosporium species, such as, for example, *Cladosporium cucumerinum*;
 Cochliobolus species, such as, for example, *Cochliobolus sativus*
 (conidia form: *Drechslera*, Syn: *Helminthosporium*);
- 25 Colletotrichum species, such as, for example, *Colletotrichum lindemuthianum*;
 Cycloconium species, such as, for example, *Cycloconium oleaginum*;
 Diaporthe species, such as, for example, *Diaporthe citri*;
 Elsinoe species, such as, for example, *Elsinoe fawcettii*;
 Gloeosporium species, such as, for example, *Gloeosporium laeticolor*;
- 30 Glomerella species, such as, for example, *Glomerella cingulata*;
 Guignardia species, such as, for example, *Guignardia bidwelli*;
 Leptosphaeria species, such as, for example, *Leptosphaeria maculans*;
 Magnaporthe species, such as, for example, *Magnaporthe grisea*;
 Mycosphaerella species, such as, for example, *Mycosphaerella graminicola*;
- 35 Phaeosphaeria species, such as, for example, *Phaeosphaeria nodorum*;

30725-1148D1

- 45 -

- Pyrenophora species, such as, for example, *Pyrenophora teres*;
- Ramularia species, such as, for example, *Ramularia collo-cygni*;
- Rhynchosporium species, such as, for example, *Rhynchosporium secalis*;
- Septoria species, such as, for example, *Septoria apii*;
- 5 Typhula species, such as, for example, *Typhula incarnata*;
- Venturia species, such as, for example, *Venturia inaequalis*;
- Root and stem diseases caused, for example, by
- Corticium species, such as, for example, *Corticium graminearum*;
- Fusarium species, such as, for example, *Fusarium oxysporum*;
- 10 Gaeumannomyces species, such as, for example, *Gaeumannomyces graminis*;
- Rhizoctonia species, such as, for example *Rhizoctonia solani*;
- Tapesia species, such as, for example, *Tapesia acuformis*;
- Thielaviopsis species, such as, for example, *Thielaviopsis basicola*;
- Ear and panicle diseases (including corn cobs) caused, for example, by
- 15 Alternaria species, such as, for example, *Alternaria* spp.;
- Aspergillus species, such as, for example, *Aspergillus flavus*;
- Cladosporium species, such as, for example, *Cladosporium* spp.;
- Claviceps species, such as, for example, *Claviceps purpurea*;
- Fusarium species, such as, for example, *Fusarium culmorum*;
- 20 Gibberella species, such as, for example, *Gibberella zeae*;
- Monographella species, such as, for example, *Monographella nivalis*;
- Diseases caused by smut fungi, such as, for example,
- Sphacelotheca species, such as, for example, *Sphacelotheca reiliana*;
- Tilletia species, such as, for example, *Tilletia caries*;
- 25 Urocystis species, such as, for example, *Urocystis occulta*;
- Ustilago species, such as, for example, *Ustilago nuda*;
- Fruit rot caused, for example, by
- Aspergillus species, such as, for example, *Aspergillus flavus*;
- Botrytis species, such as, for example, *Botrytis cinerea*;
- 30 Penicillium species, such as, for example, *Penicillium expansum*;
- Sclerotinia species, such as, for example, *Sclerotinia sclerotiorum*;
- Verticillium species, such as, for example, *Verticillium alboatrum*;
- Seed- and soil-borne rot and wilt diseases, and also diseases of seedlings, caused, for example, by

30725-1148D1

- 46 -

- Fusarium species, such as, for example, *Fusarium culmorum*;
 Phytophthora species, such as, for example, *Phytophthora cactorum*;
 Pythium species, such as, for example, *Pythium ultimum*;
 Rhizoctonia species, such as, for example, *Rhizoctonia solani*;
- 5 Sclerotium species, such as, for example, *Sclerotium rolfsii*;
- Cancerous diseases, galls and witches' broom caused, for example, by
 Nectria species, such as, for example, *Nectria galligena*;
- Wilt diseases caused, for example, by
 Monilinia species, such as, for example, *Monilinia laxa*;
- 10 Deformations of leaves, flowers and fruits caused, for example, by
 Taphrina species, such as, for example, *Taphrina deformans*;
- Degenerative diseases of woody plants caused, for example, by
 Esca species, such as, for example, *Phaemoniella clamydospora*;
 Diseases of flowers and seeds caused, for example, by
- 15 Botrytis species, such as, for example, *Botrytis cinerea*;
- Diseases of plant tubers caused, for example, by
 Rhizoctonia species, such as, for example, *Rhizoctonia solani*;
- Diseases caused by bacteriopathogens, such as, for example,
 Xanthomonas species, such as, for example, *Xanthomonas campestris* pv. *oryzae*;
- 20 Pseudomonas species, such as, for example, *Pseudomonas syringae* pv. *lachrymans*;
 Erwinia species, such as, for example, *Erwinia amylovora*.
- Preference is given to controlling the following diseases of soya beans:
 fungal diseases on leaves, stems, pods and seeds caused, for example, by
 alternaria leaf spot (*Alternaria spec. atrans tenuissima*), anthracnose (*Colletotrichum*
 25 *gloeosporoides dematium* var. *truncatum*), brown spot (*Septoria glycines*), cercospora leaf spot and
 blight (*Cercospora kikuchii*), choanephora leaf blight (*Choanephora infundibulifera trispora*
 (Syn.)), dactuliophora leaf spot (*Dactuliophora glycines*), downy mildew (*Peronospora*
manshurica), drechslera blight (*Drechslera glycini*), frog-eye leaf spot (*Cercospora sojina*),
 leptosphaerulina leaf spot (*Leptosphaerulina trifolii*), phyllosticta leaf spot (*Phyllosticta sojaecola*),
 30 powdery mildew (*Microsphaera diffusa*), pyrenochaeta leaf spot (*Pyrenochaeta glycines*),
 rhizoctonia aerial, foliage, and web blight (*Rhizoctonia solani*), rust (*Phakopsora pachyrhizi*), scab
 (*Sphaceloma glycines*), stemphylium leaf blight (*Stemphylium botryosum*), target spot

30725-1148D1

- 47 -

(Corynespora cassiicola)

Fungal diseases on roots and the stem base caused, for example, by black root rot (*Calonectria crotalariae*), charcoal rot (*Macrophomina phaseolina*), fusarium blight or wilt, root rot, and pod and collar rot (*Fusarium oxysporum*, *Fusarium orthoceras*, *Fusarium semitectum*, *Fusarium equiseti*), mycoleptodiscus root rot (*Mycoleptodiscus terrestris*), neocosmospora (*Neocosmopora vasinfecta*), pod and stem blight (*Diaporthe phaseolorum*), stem canker (*Diaporthe phaseolorum* var. *caulivora*), phytophthora rot (*Phytophthora megasperma*), brown stem rot (*Phialophora gregata*), pythium rot (*Pythium aphanidermatum*, *Pythium irregulare*, *Pythium debaryanum*, *Pythium myriotylum*, *Pythium ultimum*), rhizoctonia root rot, stem decay, and damping-off (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia sclerotiorum*), sclerotinia Southern blight (*Sclerotinia rolfsii*), thielaviopsis root rot (*Thielaviopsis basicola*).

The fact that the active compound combinations are well tolerated by plants at the concentrations required for controlling plant diseases permits a treatment of entire plants (above-ground parts of plants and roots), of propagation stock and seed, and of the soil. The active compound combinations according to the invention can be used for foliar application or else as seed dressings.

The fact that the active compounds which can be used are well tolerated by plants at the concentrations required for controlling plant diseases permits a treatment of the seed. Accordingly, the active compounds according to the invention can be used as seed dressings.

A large part of the damage to crop plants which is caused by phytopathogenic fungi occurs as early as when the seed is attacked during storage and after the seed is introduced into the soil, as well as during and immediately after germination of the plants. This phase is particularly critical since the roots and shoots of the growing plant are particularly sensitive and even minor damage can lead to the death of the whole plant. Protecting the seed and the germinating plant by the use of suitable compositions is therefore of particularly great interest.

The control of phytopathogenic fungi which damage plants post-emergence is carried out primarily by treating the soil and the above-ground parts of plants with crop protection agents. Owing to the concerns regarding a possible impact of crop protection agents on the environment and the health of man and animals, there are efforts to reduce the amount of active compounds applied.

The control of phytopathogenic fungi by treating the seed of plants has been known for a long time and is subject-matter of continuous improvements. However, the treatment of seed entails a series of problems which cannot always be solved in a satisfactory manner. Thus, it is desirable to develop methods for protecting the seed and the germinating plant which dispense with the

30725-1148D1

- 48 -

additional application of crop protection agents after sowing or after the emergence of the plants or where additional application is at least considerably reduced. It is furthermore desirable to optimize the amount of active compound employed in such a way as to provide maximum protection for the seed and the germinating plant from attack by phytopathogenic fungi, but without damaging the plant itself by the active compound employed. In particular, methods for the treatment of seed should also take into consideration the intrinsic fungicidal properties of transgenic plants in order to achieve optimum protection of the seed and the germinating plant with a minimum of crop protection agents being employed.

The present invention therefore in particular also relates to a method for the protection of seed and germinating plants from attack by phytopathogenic fungi, by treating the seed with a composition according to the invention.

The invention likewise relates to the use of the compositions according to the invention for the treatment of seed for protecting the seed and the germinating plant from phytopathogenic fungi.

Furthermore, the invention relates to seed which has been treated with a composition according to the invention so as to afford protection from phytopathogenic fungi.

One of the advantages of the present invention is that, because of the particular systemic properties of the compositions according to the invention, treatment of the seed with these compositions not only protects the seed itself, but also the resulting plants after emergence, from phytopathogenic fungi. In this manner, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

Furthermore, it must be considered as advantageous that the mixtures according to the invention can also be employed in particular in transgenic seed.

The compositions according to the invention are suitable for protecting seed of any plant variety which is employed in agriculture, in the greenhouse, in forests or in horticulture. In particular, this takes the form of seed of cereals (such as wheat, barley, rye, millet and oats), corn, cotton, soya beans, rice, potatoes, sunflowers, beans, coffee, beet (for example sugar beet and fodder beet), peanuts, vegetables (such as tomatoes, cucumbers, onions and lettuce), lawn and ornamental plants. The treatment of seed of cereals (such as wheat, barley, rye and oats), corn and rice is of particular importance.

In the context of the present invention, the composition according to the invention is applied to the seed either alone or in a suitable formulation. Preferably, the seed is treated in a state which is stable enough to avoid damage during treatment. In general, the seed may be treated at any point in

30725-1148D1

- 49 -

time between harvest and sowing. The seed usually used has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh of the fruit. Thus, for example, it is possible to use seed which has been harvested, cleaned and dried to a moisture content of below 15% by weight. Alternatively, it is also possible to use seed which, after drying, has, for example, been
5 treated with water and then dried again.

When treating the seed, care must generally be taken that the amount of the composition according to the invention applied to the seed and/or the amount of further additives is chosen in such a way that the germination of the seed is not adversely affected, or that the resulting plant is not damaged. This must be borne in mind in particular in the case of active compounds which may
10 have phytotoxic effects at certain application rates.

The compositions according to the invention can be applied directly, that is to say without comprising further components and without having been diluted. In general, it is preferable to apply the composition to the seed in the form of a suitable formulation. Suitable formulations and methods for the treatment of seed are known to the skilled worker and are described, for example,
15 in the following documents: US 4,272,417 A, US 4,245,432 A, US 4,808,430 A, US 5,876,739 A, US 2003/0176428 A1, WO 2002/080675 A1, WO 2002/028186 A2.

The active compound combinations according to the invention are also suitable for increasing the yield of crops. In addition, they show reduced toxicity and are well tolerated by plants.

According to the invention, it is possible to treat all plants and parts of plants. Plants are to be
20 understood here as meaning all plants and plant populations, such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which can be obtained by conventional breeding and optimization methods or by biotechnological and genetic engineering methods or combinations of these methods, including the transgenic plants and including plant cultivars which can or cannot be protected by plant breeders' certificates. Parts of
25 plants are to be understood as meaning all above-ground and below-ground parts and organs of plants, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stems, trunks, flowers, fruit-bodies, fruits and seeds and also roots, tubers and rhizomes. Parts of plants also include harvested material and vegetative and generative propagation material, for example seedlings, tubers, rhizomes, cuttings and seeds.

30 The treatment of the plants and parts of plants according to the invention with the active compounds is carried out directly or by action on their environment, habitat or storage area according to customary treatment methods, for example by dipping, spraying, evaporating,

30725-1148D1

- 50 -

atomizing, broadcasting, brushing-on and, in the case of propagation material, in particular in the case of seeds, furthermore by one- or multilayer coating.

As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, if appropriate in combination with conventional methods (Genetically Modified Organisms), and parts thereof, are treated. The term "parts" or "parts of plants" or "plant parts" has been explained above.

10 Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible which exceed the effects which were actually to be expected.

The transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are preferably to be treated according to the invention include all plants which, in the genetic modification, received genetic material which imparted particularly advantageous useful properties ("traits") to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defense of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), corn, soya beans, potatoes, cotton, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and

grapes), and particular emphasis is given to corn, soya beans, potatoes, cotton and oilseed rape. Traits that are emphasized are in particular increased defense of the plants against insects, by toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as "Bt plants"). Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulfonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are corn varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example corn, cotton, soya beans), KnockOut® (for example corn), Starlink® (eg corn), Bollgard® (cotton), Nucofn® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are corn varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example corn, cotton, soya beans), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulfonylureas, for example corn). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned also include the varieties sold under the name Clearfield® (for example corn). Of course, these statements also apply to plant cultivars which have these genetic traits or genetic traits still to be developed, and which will be developed and/or marketed in the future.

Depending on their particular physical and/or chemical properties, the active compound combinations according to the invention can be converted into the customary formulations, such as solutions, emulsions, suspensions, powders, dusts, foams, pastes, soluble powders, granules, aerosols, suspoemulsion concentrates, natural and synthetic materials impregnated with active compound and microencapsulations in polymeric substances and in coating compositions for seed, and ULV cold and warm fogging formulations.

These formulations are produced in a known manner, for example by mixing the active compounds or active compound combinations with extenders, that is liquid solvents, liquefied gases under pressure, and/or solid carriers, optionally with the use of surfactants, that is emulsifiers and/or dispersants, and/or foam formers.

If the extender used is water, it is also possible to employ, for example, organic solvents as auxiliary solvents. Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as

30725-1148D1

- 52 -

cyclohexane or paraffins, for example petroleum fractions, mineral and vegetable oils, alcohols such as butanol or glycol and their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulfoxide, or else water.

- 5 Liquefied gaseous extenders or carriers are to be understood as meaning liquids which are gaseous at standard temperature and under atmospheric pressure, for example aerosol propellants such as butane, propane, nitrogen and carbon dioxide.

Suitable solid carriers are: for example ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground
10 synthetic minerals such as finely divided silica, alumina and silicates. Suitable solid carriers for granules are: for example crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, or else synthetic granules of inorganic and organic meals, and granules of organic material such as sawdust, coconut shells, corn cobs and tobacco stalks. Suitable emulsifiers and/or foam formers are: for example nonionic and anionic emulsifiers, such as
15 polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulfonates, alkyl sulfates, arylsulfonates, or else protein hydrolyzates. Suitable dispersants are: for example lignosulfite waste liquors and methylcellulose.

Tackifiers such as carboxymethylcellulose, natural and synthetic polymers in the form of powders, granules or latexes, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, or else natural
20 phospholipids such as cephalins and lecithins and synthetic phospholipids can be used in the formulations. Other possible additives are mineral and vegetable oils.

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper,
25 cobalt, molybdenum and zinc.

The active compound content of the use forms prepared from the commercial formulations may be varied within wide ranges. The concentration of active compound of the use forms for controlling animal pests, such as insects and acarids, may be from 0.0000001 to 95% by weight of active compound and is preferably from 0.0001 to 1% by weight. Application is in a customary manner
30 adapted to the use forms.

The formulations for controlling unwanted phytopathogenic fungi generally comprise between 0.1 and 95% by weight of active compounds, preferably between 0.5 and 90%.

30725-1148D1

- 53 -

The active compound combinations according to the invention can be used as such, in the form of their formulations or as the use forms prepared therefrom, such as ready-to-use solutions, emulsifiable concentrates, emulsions, suspensions, wettable powders, soluble powders, dusts and granules. They are used in a customary manner, for example by watering (drenching), drip irrigation, spraying, atomizing, broadcasting, dusting, foaming, spreading-on, and as a powder for dry seed treatment, a solution for seed treatment, a water-soluble powder for seed treatment, a water-soluble powder for slurry treatment, or by encrusting etc.

The active compound combinations according to the invention can, in commercial formulations and in the use forms prepared from these formulations, be present as a mixture with other active compounds, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators or herbicides.

When using the active compound combinations according to the invention, the application rates can be varied within a relatively wide range, depending on the kind of application. In the treatment of parts of plants, the application rates of active compound combination are generally between 0.1 and 10 000 g/ha, preferably between 10 and 1000 g/ha. In the treatment of seed, the application rates of active compound combination are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 10 g per kilogram of seed. In the treatment of the soil, the application rates of active compound combination are generally between 0.1 and 10 000 g/ha, preferably between 1 and 5000 g/ha.

The active compound combinations can be used as such, in the form of concentrates or in the form of generally customary formulations, such as powders, granules, solutions, suspensions, emulsions or pastes.

The formulations mentioned can be prepared in a manner known per se, for example by mixing the active compounds with at least one solvent or diluent, emulsifier, dispersant and/or binder or fixative, water repellent, if desired desiccants and UV stabilizers, and, if desired, colorants and pigments and other processing auxiliaries.

The good fungicidal action of the active compound combinations according to the invention is demonstrated by the examples below. While the individual active compounds show weaknesses in their fungicidal action, the combinations show an action which exceeds a simple sum of actions.

A synergistic effect in the fungicides is always present when the fungicidal action of the active compound combinations exceeds the total of the action of the active compounds when applied individually.

30725-1148D1

- 54 -

The expected fungicidal action for a given combination of two active compounds can be calculated as follows, according to S.R. Colby ("Calculating Synergistic and Antagonistic Responses of Herbicide Combinations", Weeds 1967, 15, 20-22):

If

5 X is the *efficacy* when employing active compound A at an application rate of m g/ha,

Y is the *efficacy* when employing active compound B at an application rate of n g/ha and

E is the *efficacy* when employing active compounds A and B at application rates of m and n g/ha,

then
$$E = X + Y - \frac{X \times Y}{100}$$

10 Here, the efficacy is determined in %. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

If the actual fungicidal action exceeds the calculated value, the action of the combination is superadditive, i.e. a synergistic effect is present. In this case, the actually observed efficacy must exceed the value calculated using the above formula for the expected efficacy (E).

30725-1148D1

- 55 -

Use examplesExample A**Puccinia Test (wheat)/protective**

Solvent: 50 parts by weight of N,N-dimethylacetamide

5 Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier or a commercial active compound formulation, and the concentrate is diluted with water to the desired concentration.

10 To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Puccinia recondita*. The plants remain in an incubation cabin at 20°C and 100% relative atmospheric humidity for 48 hours.

The plants are then placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of 80% to promote the development of rust pustules.

15 Evaluation is carried out 10 days after the inoculation. 0% means inefficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table A: Puccinia test (wheat) protective

Active compounds	Active compound application rate in g/ha	Efficacy in %	
		found*	calc.**
(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	250	67	
(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide	125	33	
(2-9) picoxystrobin	250	94	
(3-6) cyproconazole	125	94	
(9-1) cyprodinil	125	0	
(3-7) hexaconazole	125	78	
(3-3) propiconazole	125	56	
(1-1) + (2-9) picoxystrobin 1:1 (A-5)	250 + 250	100	98
(1-1) + (3-6) cyproconazole 1:1 (B-3)	125 + 125	100	96
(1-1) + (9-1) cyprodinil 1:1 (H-1)	125 + 125	44	33
(1-1) + (3-7) hexaconazole 1:1 (B-4)	125 + 125	100	85
(1-1) + (3-3) propiconazole 1:1 (B-1)	125 + 125	78	70

* found = activity found

** calc. = activity calculated using Colby's formula

30725-1148D1

- 56 -

Example B**Erysiphe Test (wheat)/protective**

Solvent: 50 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylarylpolyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier or a commercial active compound formulation, and the concentrate is diluted with water to the desired concentration.

To test for protective activity, young plants are sprayed with the preparation of active compound at the stated application rate.

- 10 After the spray coating has dried on, the plants are dusted with spores of *Erysiphe graminis* f.sp. *tritici*.

The plants are placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of about 80% to promote the development of mildew pustules.

Evaluation is carried out 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

15

Table B: Erysiphe test (wheat) protective

Active compound	Active compound application rate in g/ha	Efficacy in %	
		found*	calc.**
(1-1) N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-carboxamide	62.5	0	
(2-9) picoxystrobin	62.5	71	
(3-6) cyproconazole	62.5	57	
(9-1) cyprodinil	62.5	0	
(3-7) hexaconazole	62.5	29	
(3-3) propiconazole	62.5	29	
(1-1) + (2-9) picoxystrobin 1:1 (A-5)	62.5 + 62.5	86	71
(1-1) + (3-6) cyproconazole 1:1 (B-3)	62.5 + 62.5	93	57
(1-1) + (9-1) cyprodinil 1:1 (H-1)	62.5 + 62.5	29	0
(1-1) + (3-7) hexaconazole 1:1 (B-4)	62.5 + 62.5	57	29
(1-1) + (3-3) propiconazole 1:1 (B-1)	62.5 + 62.5	57	29

* found = activity found


** calc. = activity calculated using Colby's formula

30725-1148D1

- 57 -

CLAIMS:

1. A fungicidal active compound combination comprising a carboxamide of the general formula (I):



- 5 wherein:


R represents H or methyl,

G represents H, F or methyl,

R¹ represents H, a halogen atom, C₁-C₃-alkyl or C₁-C₃-haloalkyl having 1 to 7 fluorine, chlorine and/or bromine atoms, and

- 10 A represents the radical A1:

A1



wherein:

R² represents methyl, difluoromethyl or trifluoromethyl, and

- 15 R³ represents H or F, and

at least one active compound selected from group consisting of:

30725-1148D1

- 58 -

fluoxastrobin,

(2*E*)-2-(2-{[6-(3-chloro-2-methylphenoxy)-5-fluoro-4-pyrimidinyl]-oxy}phenyl)-2-(methoxyimino)-*N*-methylethanamide,

trifloxystrobin,

5 (2*E*)-2-(methoxyimino)-*N*-methyl-2-(2-{{{(1*E*)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)ethanamide,

(2*E*)-2-(methoxyimino)-*N*-methyl-2-{2-[(*E*)-({1-[3-(trifluoromethyl)-phenyl]ethoxy}imino)methyl]phenyl}ethanamide,

oryastrobin,

10 5-methoxy-2-methyl-4-(2-{{{(1*E*)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy}methyl}phenyl)-2,4-dihydro-3*H*-1,2,4-triazol-3-one,

dimoxystrobin,

picoxystrobin,

pyraclostrobin,

15 azaconazole,

etaconazole,

propiconazole,

difenoconazole,

bromuconazole,

20 cyproconazole,

hexaconazole,

30725-1148D1

- 59 -

penconazole,

myclobutanil,

tetraconazole,

flutriafol,

5 epoxiconazole,

flusilazole,

simeconazole,

prothioconazole,

fenbuconazole,

10 tebuconazole,

ipconazole,

metconazole,

triticonazole,

bitertanol,

15 triadimenol,

triadimefon,

fluquinconazole,

quinconazole,

dichlofluanid,

20 tolylfluanid,

30725-1148D1

- 60 -

iprovalicarb,

benthiavalicarb,

2-chloro-N-(1,1,3-trimethylindan-4-yl)nicotinamide,

boscalid,

5 furametpyr,

ethaboxam,

fenhexamid,

carpropamid,

2-chloro-4-(2-fluoro-2-methylpropionylamino)-N,N-dimethylbenzamide,

10 picobenzamid,

zoxamide,

3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide,

carboxin,

tiadinil,

15 silthiofam,

mancozeb,

maneb,

metiram,

propineb,

20 thiram,

30725-1148D1

- 61 -

zineb,

ziram,

benalaxyl,

furalaxyl,

5 benalaxyl-M,

cyprodinil,

pyrimethanil,

6-chloro-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2,2-difluoro-5H-
[1,3]dioxolo[4,5-f]benzimidazole,

10 benomyl,

carbendazim,

chlorfenazole,

fuberidazole,

thiabendazole,

15 propamocarb,

propamocarb hydrochloride,

propamocarb-fosetyl,

captafol,

folpet,

20 iprodione,

30725-1148D1

- 62 -

procymidone,

vinclozolin,

dodine,

guazatine,

5 iminoctadine triacetate,

cyazofamid,

prochloraz,

triazoxide,

pefurazoate,

10 aldimorph,

tridemorph,

dodemorph,

fenpropimorph,

fenpiclonil,

15 pyrrolnitrine,

phosphonic acid,

2-(2,3-dihydro-1H-inden-5-yl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)-acetamide,

20 N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)-2-(5,6,7,8-tetrahydro-naphthalen-2-yl)acetamide,

30725-1148D1

- 63 -

2-(4-chlorophenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide,

2-(4-bromophenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide,

5 2-(4-methylphenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide,

2-(4-ethylphenyl)-N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(methoxyimino)acetamide,

acibenzolar-S-methyl,

10 edifenphos,

famoxadone,

oxadixyl,

spiroxamine,

dithianon,

15 metrafenone,

fenamidone,

2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)one,

probenazole,

isoprothiolane,

20 kasugamycin,

phthalide,

30725-1148D1

- 64 -

N-({4-[(cyclopropylamino)carbonyl]phenyl}sulfonyl)-2-methoxybenzamide,

2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide,

quinoxifen,

5 proquinazid,

pencycuron,

thiophanate-ethyl,

fenoxanil,

diclocymet

10 5-chloro-*N*-[(*IS*)-2,2,2-trifluoro-1-methylethyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo-[1,5-a]pyrimidine-7-amine,

5-chloro-*N*-[(*IR*)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine,

15 5-chloro-6-(2-chloro-6-fluorophenyl)-7-(4-methylpiperidin-1-yl)[1,2,4]triazolo[1,5-a]pyrimidine,

5-chloro-6-(2,4,6-trifluoro-phenyl)-7-(4-methylpiperidin-1-yl)[1,2,4] triazolo [1,5-a]pyrimidine,

5-chloro-6-(2,4,6-trifluorophenyl)-*N*-[(*IS*)-1,2,2-trimethylpropyl][1,2,4] triazolo[1,5-a]pyrimidine-7-amine,

20 2-butoxy-6-iodo-3-propylbenzopyran-4-one,

2-ethoxy-6-iodo-3-propylbenzopyran-4-one,

6-iodo-2-propoxy-3-propylbenzopyran-4-one,

30725-1148D1

- 65 -

2-but-2-ynyloxy-6-iodo-3-propylbenzopyran-4-one,

6-iodo-2-(1-methylbutoxy)-3-propylbenzopyran-4-one,

2-but-3-enyloxy-6-iodobenzopyran-4-one, and

3-butyl-6-iodo-2-isopropoxybenzopyran-4-one.

5 2. The combination as claimed in claim 1, wherein:

R, R² and R³ are as defined in claim 1,

G represents H or methyl, and

R¹ represents H, F, Cl, methyl, ethyl, n-propyl, isopropyl, monofluoromethyl, difluoromethyl, trifluoromethyl, monochloromethyl, dichloromethyl or trichloromethyl.

10 3. The combination as claimed in claim 1, wherein the carboxamide of the general formula (I) is N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-4-(trifluoromethyl)-1H-pyrrole-3-carboxamide.

4. Use of the combination as claimed in any one of claims 1 to 3, for controlling unwanted phytopathogenic fungi.

15 5. Use of the combination as claimed in any one of claims 1 to 3, for treating seed.

6. Use of the combination as claimed in any one of claims 1 to 3, for treating transgenic plants.

7. Use of the combination as claimed in any one of claims 1 to 3, for treating seed of transgenic plants.

20 8. A method for controlling unwanted phytopathogenic fungi, comprising applying the combination as claimed in any one of claims 1 to 3, to the unwanted phytopathogenic fungi, a habitat thereof or seed thereof.

30725-1148D1

- 66 -

9. A process for preparing a fungicidal composition, comprising mixing the combination as claimed in any one of claims 1 to 3, with an extender, a surfactant or a mixture thereof.