



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12N 15/75, 15/56, 9/28</p>	A1	<p>(11) International Publication Number: WO 93/10249 (43) International Publication Date: 27 May 1993 (27.05.93)</p>
<p>(21) International Application Number: PCT/DK92/00338 (22) International Filing Date: 13 November 1992 (13.11.92) (30) Priority data: PCT/DK91/00343 14 November 1991 (14.11.91) WO (34) Countries for which the regional or international application was filed: DK et al. (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only) : JØRGENSEN, Steen, Troels [DK/DK]; Prumsvej 5, DK-3450 Allerød (DK). DIDERICHSEN, Børge Krag [DK/DK]; Fuglsangsvej 4, DK-3460 Birkerød (DK).</p>		<p>(74) Common Representative: NOVO NORDISK A/S; Patent Department, Novo Allé, DK-2880 Bagsvaerd (DK). (81) Designated States: AU, BR, CA, CS, FI, HU, JP, KR, NO, PL, RU, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published With international search report.</p>
<p>(54) Title: A BACILLUS PROMOTER DERIVED FROM A VARIANT OF A BACILLUS LICHENIFORMIS X-AMYLASE PROMOTER</p> <p style="text-align: center;">GCATGCGTCC TTCTTTGTGC TTGGAAGCAG AGCCCAATAT TATCCCGAAA CGATAAAACG GATGCTGAAG GAAGGAAACG AAGTCGGCAA CCATTCCTGG GACCCATCCG TTATTGACAA GGCTGTCAAA CGAAAAAGCG TATCAGGAGA TTAACGACAC GCAAGAAATG ATCGAAAAAA TCAGCGGACA CCTGCCTGTA CACTTGCGTC CTCCATACGG CGGGATCAAT GATTCGGTCC GCTCGCTTTC CAATCTGAAG GTTTCATTGT GGGATGTTGA TCCGGAAGAT TGGAAGTACA AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGACGGA AAAATCGTCT TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCTGCTGA AGAGATTATT AAAAAGCTGA AAGCAAAGG CTATCAATTG GTAACGTGAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG GCTATTGAAT AAATGAGTAG AAAGCGCCAT ATCGGCGCTT TTCTTTTGGGA AGAAAATATA GGGAAAATGG TAN¹TTGTTAA AAATTCGGAA TATTTATACA ATATCATN²N³N⁴ N⁵N⁶N⁷N⁸N⁹CATTG AAAGGGGAGG AGAATC (SEQ ID#1)</p> <p>(57) Abstract</p> <p>A <i>Bacillus</i> promoter included in DNA sequence (SEQ ID#1), wherein each of N¹-N⁹ is A, T, C or G with the exception that N²-N⁹ do not together form the sequence ATGTTTCA or GTGTTTCA, or a functional homologue of said sequence.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Mongco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

A BACILLUS PROMOTER DERIVED FROM A VARIANT OF A BACILLUS
LICHENIFORMIS X-AMYLASE PROMOTER

FIELD OF INVENTION

5 The present invention relates to a Bacillus licheniformis promoter, a DNA construct comprising said promoter, a host cell transformed with said DNA construct and a method of producing a protein in Bacillus by means of the promoter.

10 BACKGROUND OF THE INVENTION

Various promoter sequences of the Bacillus licheniformis α -amylase gene have been described previously. Thus, M. Sibakov and I. Palva, Eur. J. Biochem. 145, 1984, pp. 567-572, describe
15 the isolation and determination of the 5' end of the Bacillus licheniformis α -amylase gene, including the promoter sequence; T. Yuuki et al., J. Biochem. 98, 1985, pp. 1147-1156, show the complete nucleotide sequence of the Bacillus licheniformis α -amylase gene, including the promoter sequence; and B.M. Laoide
20 et al., J. Bacteriol. 171(5), 1989, pp. 2435-2442, discuss catabolite repression of the Bacillus licheniformis α -amylase gene from a region around the 5' end of the gene and show the sequence of this region.

25 SUMMARY OF THE INVENTION

The present inventors have surprisingly found that a novel promoter homologous to the previously published promoter sequences gives rise to a dramatically increased yield of a
30 protein when the gene coding for the protein is transcribed from the promoter.

Accordingly, the present invention relates to a Bacillus promoter included in the following DNA sequence

35

GCATGCGTCC TTCTTTGTGC TTGGAAGCAG AGCCCAATAT TATCCCGAAA
CGATAAAACG GATGCTGAAG GAAGGAAACG AAGTCGGCAA CCATTCCTGG

GACCCATCCG TTATTGACAA GGCTGTCAAA CGAAAAAGCG TATCAGGAGA
 TTAACGACAC GCAAGAAATG ATCGAAAAAA TCAGCGGACA CCTGCCTGTA
 CACTTGCGTC CTCCATACGG CGGGATCAAT GATTCCGTCC GCTCGCTTTC
 CAATCTGAAG GTTTCATTGT GGGATGTTGA TCCGGAAGAT TGGAAGTACA
 5 AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGACGGA
 AAAATCGTCT TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCTGCTGA
 AGAGATTATT AAAAAGCTGA AAGCAAAAGG CTATCAATTG GTAACTGTAT
 CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG GCTATTGAAT AAATGAGTAG
 AAAGCGCCAT ATCGGCGCTT TTCTTTTGGGA AGAAAATATA GGGAAAATGG
 10 TAN¹TTGTTAA AAATTCGGAA TATTTATAACA ATATCATN²N³N⁴
 N⁵N⁶N⁷N⁸N⁹CATTG AAAGGGGAGG AGAATC (SEQ ID#1)

wherein each of N¹-N⁹ is A, T, C or G with the exception that
 N²-N⁹ do not together form the sequence ATGTTTCA or GTGTTTCA,
 15
 or a functional homologue of said sequence.

In the the previously published sequences, N¹ is either T (cf.
 T. Yuuki et al., supra) or C (B.M. Laoide et al., supra), while
 20 N²-N⁹ is either ATGTTTCA (T. Yuuki et al., supra, and B.M.
 Laoide et al., supra) or GTGTTTCA (cf. M. Sibakov, supra).
 Several papers discuss catabolite repression of Bacillus genes,
 including the B. licheniformis α -amylase gene. Thus, B.M.
 Laoide et al, supra, and B.M. Laoide and D.J. McConnell, J.
 25 Bacteriol. 171, 1989, pp. 2443-2450, map the cis sequences
 essential for mediation of catabolite repression of amyL in B.
subtilis to a 108 bp region downstream from the promoter and
 upstream from the signal sequence cleavage site. They identify
 30 an inverted repeat sequence, TGTTTCAC-20 bp-ATGAAACA, in this
 region but note that deletion into the left-hand part of this
 sequence either abolished or altered expression without
 affecting catabolite repression. They identify sequences
 homologous to the left-hand part of the amyL inverted repeat
 (5'-A/T T G T N A/T-3') around the transcription initiation
 35 sites in a number of B. subtilis catabolite-repressible genes.

Y. Miwa and Y. Fujita, Nucl. Acids Res. 18, pp. 7049-7053,

limit the cis sequences involved in catabolite repression of the B. subtilis gnt operon to a 11 bp region. Within this 11 bp region is a 8 bp sequence, ATTGAAAG, which the authors claim could be a consensus sequence involved in catabolite repression
5 in the genus Bacillus, as it was found in other catabolite repressible Bacillus genes. Interestingly, in the B. licheniformis α -amylase gene, the consensus sequence shown above immediately follows the left-hand part of the inverted repeat sequence identified by Laoide et al.

10

M.J. Weickert and G.H. Chambliss, Proc. Natl. Acad. Sci. USA 87, pp. 6238-6242, describe site-directed mutagenesis of a catabolite repression operator sequence in B. subtilis from the amyE gene. They observe that hyperproduction and catabolite
15 repression of amylase were both affected by mutations in the same region, and sometimes by the same mutation. They found that the B. subtilis α -amylase catabolite repression operator shares significant homology with sequences in other Bacillus amylase gene regulatory regions and with other catabolite
20 repressed genes. The consensus sequence they identified is located from position +70 to +64 with respect to the B. licheniformis α -amylase transcription initiation site.

At least one group considers the sequence N^2-N^9 (according to
25 the present nomenclature) to form an essential part of the cis sequence required for catabolite repression, while another group points to an immediately adjacent sequence. It is noteworthy that N^2-N^9 form part of an inverted repeat sequence. Modifications of these sequences might well influence the
30 transcription levels obtained from the amyL promoter. It cannot, however, be discounted, that substitutions in other parts of the promoter sequence such as at N^1 , may also influence the transcription levels obtained from the promoter.

35 In the present context, the term "functional homologue" is intended to indicate a promoter sequence with at least 70% sequence identity to the sequence shown above, which sequence,

under comparable conditions, promotes a more efficient transcription of the gene it precedes than the promoter disclosed by T. Yuuki et al., supra, or B. Laoide et al., supra. The transcription efficiency may, for instance, be
5 determined by a direct measurement of the amount of mRNA transcription from the promoter, e.g. by Northern blotting or primer extension, or indirectly by measuring the amount of gene product expressed from the promoter. The term is intended to include derivatives of the promoter sequence shown above, such
10 as insertion of one or more nucleotides into the sequence, addition of one or more nucleotides at either end of the sequence and deletion of one or more nucleotides at either end of or within the sequence, provided that such modifications do not impair the promoter function of the sequence. Fragments of
15 the sequence shown above are included in this definition of a functional homologue.

DETAILED DISCLOSURE OF THE INVENTION

20 The promoter of the invention may be derived from the genome of a suitable Bacillus licheniformis strain by hybridisation using oligonucleotide probes based on the promoter sequence known from T. Yuuki et al., supra, or B. Laoide et al., supra, in accordance with standard techniques (cf. Sambrook et al.,
25 Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, NY, 1989). The known promoter sequence may be modified at one or more sites by site-directed mutagenesis in accordance with well-known procedures. The promoter sequence may also be prepared synthetically by established standard methods, e.g.
30 the phosphoamidite method described by S.L. Beaucage and M.H. Caruthers, Tetrahedron Letters 22, 1981, pp. 1859-1869, or the method described by Matthes et al., EMBO J. 3, 1984, pp. 801-805.

35 Examples of preferred promoters of the invention are those wherein N^1 is C or T; or wherein N^7 is A, G or C; in particular wherein N^1 is C and N^7 is A. Thus, N^2-N^9 together preferably

form the sequence ATGTTACA, while N¹ is preferably C.

An example of a suitable fragment of the promoter sequence shown above has the following DNA sequence

5

CTATCAATTG GTAACGTGAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG
GCTATTGAAT AAATGAGTAG AAAGCGCCAT ATCGGCGCTT TTCTTTTGGG
AGAAAATATA GGGAAAATGG TAN¹TTGTTAA AAATTCGGAA TATTTATACA
ATATCATN²N³N⁴ N⁵N⁶N⁷N⁸N⁹CATTG AAAGGGGAGG AGAATC (SEQ ID#2)

10

wherein N¹-N⁹ has the meaning indicated above.

In a preferred embodiment, the promoter of the invention is derived from a B. licheniformis gene, and in particular it is a variant of a Bacillus licheniformis α -amylase promoter.

In another aspect, the present invention relates to a DNA construct comprising a DNA sequence coding for a protein of interest preceded by a promoter sequence as described above. The protein of interest may advantageously be an enzyme, e.g. α -amylase, cyclodextrin glycosyl transferase or a protease. The DNA construct may advantageously also comprise a sequence coding for a signal peptide to ensure secretion into the culture medium of the protein in question on cultivating a cell transformed with the DNA construct.

According to the invention, the DNA construct may be present on an autonomously replicated expression vector. The vector further comprises a DNA sequence enabling the vector to replicate in the host cell. Examples of such sequences are the origins of replication of plasmids pUC19 (C. Yanisch-Perron et al., Gene 33, 1985, pp. 103-119), pACYC177 (A.C.Y. Chang and S.N. Cohen, J. Bacteriol. 134, 1978, pp. 1141-1156), pUB110 (Gryczan et al. 1978) or pIJ702 (E. Katz et al., J. Gen. Microbiol. 129, 1983, pp. 2703-2714). The vector may also comprise a selectable marker, e.g. a gene whose product confers antibiotic resistance such as ampicillin, chloramphenicol or

tetracyclin resistance, or the dal genes from B. subtilis or B. licheniformis (B. Diderichsen, 1986). The procedures used to ligate the DNA sequence coding for the protein of interest, promoter and origin of replication are well known to persons skilled in the art (cf., for instance, Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, NY, 1989).

Alternatively, the DNA construct may be present on the chromosome of the host cell. This is often an advantage as the DNA construct is more likely to be stably maintained in the host cell. Integration of the DNA construct into the host chromosome may be performed according to conventional methods, e.g. by homologous recombination. It should be noted that the promoter sequence, the DNA sequence encoding the protein of interest and optionally the signal sequence may be introduced into the host cell separately.

The host cell may suitably be a strain of Bacillus, in particular a strain of Bacillus licheniformis, Bacillus lentus, Bacillus brevis, Bacillus stearothermophilus, Bacillus alkalophilus, Bacillus amyloliquefaciens, Bacillus coagulans, Bacillus thuringiensis or Bacillus subtilis.

In a further aspect, the present invention relates to a process for producing a protein in Bacilli comprising culturing a Bacillus host cell transformed with a DNA construct or vector according to the invention under conditions permitting production of said protein, and recovering the resulting protein from the culture.

The medium used to cultivate the cells may be any conventional medium suitable for growing bacteria. The product of the expressed gene is preferably recovered from the culture. Recovery of the product may be done by conventional procedures including separating the cells from the medium by centrifugation or filtration., precipitating the proteinaceous

components of the supernatant or filtrate by means of a salt, e.g. ammonium sulphate, followed, if necessary, by a variety of chromatographic procedures, e.g. ion exchange chromatography, affinity chromatography, or the like.

5

The invention is further described in the following example with reference to the appended drawings, in which the following abbreviations are used:

- 10 "pBR322" indicates pBR322-derived DNA;
"+ori pUB110" indicates the plus origin of replication of pUB110;
"rep" indicates the rep gene of pUB110;
"cat" indicates the chloramphenicol resistance gene of pC194;
15 "cgtA" indicates the Thermoanaerobacter CGTase gene;
"PamyM" indicates the promoter of the B. sterothermophilus maltogenic amylase gene (Diderichsen and Christiansen, 1988);
"bla" indicates the ampicillin resistance gene of pBR322;
"pKK233-2" indicates pKK233-2 derived DNA;
20 "PamyL" indicates the promoter of the B. licheniformis α -amylase gene;
"PamyQ" indicates the promoter of the B. amyloliquefaciens α -amylase gene;
"amyL-cgtA" indicates the fusion gene comprising the signal
25 peptide coding part of the B. licheniformis α -amylase gene and the part of the Thermoanaerobacter CGTase gene coding for the mature enzyme;
"erm" indicates the erythromycin resistance gene of pE194;
"ori pE194" indicates the plus origin of replication and rep
30 gene containing region of pE194; and
"'amyL" indicates a DNA fragment spanning the 3'-end of the B. licheniformis α -amylase gene.

Fig. 1 is a restriction map of plasmid pNV601;

35 Fig. 2 is a restriction map of plasmid pPL1878;

Fig. 3 is a restriction map of plasmid pPL1419;

Fig. 4 is a restriction map of plasmid pPL1489;

- Fig. 5 is a restriction map of plasmid pP11540;
Fig. 6 is a restriction map of plasmid pDN3000;
Fig. 7 is a restriction map of plasmid pP11759;
Fig. 8 is a resrtiction map of plasmid pP11892;
5 Fig. 9 is a restriction map of plasmid pP11796;
Fig. 10 is a restriction map of plasmid pBB37;
Fig. 11 is a restriction map of plasmid pP11385;
Fig. 12 is a restriction map of plasmid pP11893;
Fig. 13 is a restriction map of plasmid pSJ1111;
10 Fig. 14 is a restriction map of plasmid pDN3060;
Fig. 15 is a restriction map of plasmid pSJ1277;
Fig. 16 is a restriction map of plasmid pSJ994;
Fig. 17 is a restriction map of plasmid pSJ1283;
Fig. 18 is a restriction map of plasmid pSJ1342;
15 Fig. 19 is a restriction map of plasmid pSJ1359;
Fig. 20 is a restriction map of plasmid pP11483;
Fig. 21 is a restriction map of plasmid pP11487;
Fig. 22 is a restriction map of plasmid pSJ932;
Fig. 23 is a restriction map of plasmid pSJ948;
20 Fig. 24 is a restriction map of plasmid pSJ980;
Fig. 25 is a restriction map of plasmid pSJ1391;
Fig. 26 is a schematic presentation of the exchange, by
homologous recombination, between the chromosomal α -amylase
gene and the amyL-cgtA fusion gene carried on plasmid pSJ1391;
25 Fig. 27 is a schematic presentation of the in vivo
recombination between the 5' ends of the mature parts of cgtA;
and
Fig. 28 is a restriction map of plasmid pSJ1755.

- 30 The invention is further illustrated in the following examples
which are not in any way intended to limit the scope of the
invention as claimed.

EXAMPLE

General Methods

- 5 The experimental techniques used to construct the plasmids were standard techniques within the field of recombinant DNA technology, cf. T. Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, New York, 1982.
- 10 Restriction endonucleases were purchased from New England Biolabs and Boehringer Mannheim and used as recommended by the manufacturers. T4 DNA ligase was purchased from New England Biolabs and used as recommended by the manufacturer.
- 15 Preparation of vector DNA from all strains was conducted by the method described by Kieser, 1984.

Transformation of E. coli:

- Cells of E. coli were made competent and transformed as
20 described by Mandel and Higa, 1970.

Transformation of B. subtilis:

- Competent cells were prepared and transformed as described by
Yasbin et al., 1975.
- 25

Transformation of B. licheniformis:

- Plasmids were introduced into B. licheniformis by polyethylene glycol-mediated protoplast transformation as described by
Akamatzu, 1984.
- 30
- CGTase-producing colonies of either E. coli, B. subtilis or B. licheniformis were identified by plating transformants on LB agar plates supplemented with 1% soluble starch. After incubation at either 37°C or 30°C overnight, plates were
35 stained by iodine vapour to show hydrolysis zones produced by the action of the CGTase on the starch.

Media

	BPX:	Potato starch	100 g/l
		Barley flour	50 g/l
5		BAN 5000 SKB	0.1 g/l
		Sodium caseinate	10 g/l
		Soy Bean Meal	20 g/l
		Na ₂ HPO ₄ , 12 H ₂ O	9 g/l
		Pluronic	0.1g/l
10			
	LB agar:	Bacto-tryptone	10 g/l
		Bacto yeast extract	5 g/l
		NaCl	10 g/l
		Bacto agar	15 g/l
15		Adjusted to pH 7.5 with NaOH	

1. Cloning of a Thermoanaerobacter sp. CGTase gene into
 20 Bacillus subtilis.

The construction of the E. coli plasmid pNV601 (Fig. 1), carrying the Thermoanaerobacter sp. ATCC 53627 CGTase gene referred to in the following as cgtA, is disclosed in WO 25 89/03421. The B. subtilis plasmid pPL1878 (Fig. 2), containing the cgtA gene, is disclosed in WO 91/09129. It was constructed as follows:

pNV601 was digested partially with Sau3A, then religated and 30 transformed into E. coli SCS1 (frozen competent cells purchased from Stratagene, Ja Jolla, California), selecting for ampicillin resistance (200 µg/ml). One CGTase positive colony was PL1419, containing pPL1419 (Fig. 3). Plasmid pPL1419 was partially digested with Sau3A, and fragments ligated to BglII 35 digested pPL1489 (Fig. 4). One CGTase positive, ampicillin resistant (200 µg/ml) E. coli SCS1 transformant contained pPL1540 (Fig. 5). pPL1489 was derived from plasmid pPK233-2

(purchased from Pharmacia LKB Biotechnology) by insertion of a synthetic DNA linker between the PstI and HindIII sites in pKK233-2. This linker was the PstI-HindIII fragment from pDN3000 (Fig. 6; WO 91/09129, Diderichsen et al., 1990).
5 pPL1540 was digested with HaeII and SphI, and the 2.4 kb fragment containing the cgtA gene was inserted into HaeII + SphI digested plasmid pDN1380 (Diderichsen and Christiansen, 1988). A CGTase positive, chloramphenicol resistant (6 µg/ml) transformant of B. subtilis DN1885 (Diderichsen et al., 1990)
10 contained pPL1878.

2. Construction of an α -amylase/CGTase fusion gene.

15 Cloning of the Bacillus licheniformis α -amylase gene, amyL, resulting in plasmid pDN1981, is described by Jørgensen et al., 1990.

In plasmid pPL1759 (Fig. 7), the PstI-HindIII fragment of
20 pDN1981 has been replaced by the PstI-HindIII multilinker fragment from pDN3000 (Fig. 6). It has retained the amyL promoter and most of the signal peptide coding sequence.

Plasmid pPL1892 (Fig. 8) was constructed by insertion of the
25 cgtA gene excised from pPL1878 on a 2.4 kb SalI-NotI fragment into SalI + NotI digested pPL1759, and transformation of DN1885 to kanamycin resistance (10 µg/ml).

Plasmid pPL1796 (Fig. 9) was constructed by insertion of a 0.5
30 kb SacI-EcoRV fragment from pBB37 (Fig. 10; Jørgensen, P. et al., 1991) into SacI + SmaI digested pPL1385 (Fig. 11; Diderichsen et al., 1990), and transformation of DN1885 to chloramphenicol resistance (6 µg/ml).

35 Plasmid pPL1893 (Fig. 12) was constructed by insertion of the CGTase gene excised from pPL1878 on a 2.4 kb BamHI-NotI fragment into BamHI + NotI digested pPL1796, and transformation

of DN1885 to chloramphenicol resistance (6 µg/ml).

The in vivo genetic engineering technique (Jørgensen et al.,
 5 1990), by which two DNA sequences contained on the same plasmid
 and sharing a homologous region can be fused together by recom-
 bination between the homologous regions in vivo (see Fig. 27)
 was used to construct a fusion between the amyL and the cgtA
 genes, in which the cgtA signal peptide coding sequence had
 10 been precisely replaced by the signal peptide coding sequence
 of the amyL gene.

To this end, the following oligonucleotide linker was syn-
 thesized and ligated into SalI digested pUC19 (Yanish-Perron
 15 et al., 1985), giving pSJ1111 (Fig. 13) upon transformation of
E. coli SJ2 (Diderichsen et al., 1990) and selection for
 ampicillin resistance (200 µg/ml):

20 3' end of amyL signal peptide
 coding region

```

          SalI   BclI                               PstI
5' - TCGACTGATCACTTGCTGCCTCATTCTGCAGCAGCGGCG-
3' -      GACTAGTGAACGACGGAGTAAGACGTCGTCGCCGC-
```

25

5' end of cgtA mature protein
 coding region

```

                                XbaI   SalI
30 GCACCGGATACTTCAGTTTCTCTAGAG      - 3'
   CGTGGCCTATGAAGTCAAAGAGATCTCAGCT - 5' (SEQ ID#3)
```

The pC194 (Horinouchi and Weisblum, 1982) derived
 chloramphenicol resistance gene, cat, was excised from pDN3060
 35 (Fig. 14; WO 91/09129) as a 1.1 kb BamHI-BglII fragment and
 inserted into BclI digested pSJ1111, giving pSJ1277 (Fig. 15)
 upon transformation of E. coli SJ 6 (Diderichsen et al., 1990)

and selection for ampicillin (200 $\mu\text{g/ml}$) and chloramphenicol (6 $\mu\text{g/ml}$) resistance.

5 pSJ994 (Fig. 16) was constructed by ligation of the 0.6 kb NotI-NcoI fragment from pPL1893 to the 5.4 kb NotI-NcoI fragment from pPL1892, and transformation into B. subtilis DN1885, selecting for kanamycin resistance (10 $\mu\text{g/ml}$).

10 pSJ1283 (Fig. 17) was constructed by ligation of the 1.1 kb SalI fragment from pSJ1277 to SalI digested pSJ994, and transformation into DN1885, selecting for kanamycin (10 $\mu\text{g/ml}$) and chloramphenicol (6 $\mu\text{g/ml}$) resistance.

15 pSJ1342 (Fig. 18) was constructed by deletion of the 1.1 kb PstI fragment from pSJ1283, and transformation into DN1885, selecting for kanamycin resistance (10 $\mu\text{g/ml}$).

20 pSJ1359 (Fig. 19) was constructed by the actual in vivo recombination from pSJ1342. There is homology between the start of the mature part of the CGTase gene and part of the synthetic oligonucleotide extending between PstI and SalI on pSJ1342. If the plasmid undergoes a recombination event between these two homologous regions, the unique sites for XbaI, SalI and BamHI will be deleted.

25

A batch of pSJ1342 prepared from host strain DN1885 was thoroughly digested with BamHI, XbaI and SalI, and the digested plasmid was directly (i.e. without ligation) transformed into competent cells of DN1885, selecting for kanamycin resistance
30 (10 $\mu\text{g/ml}$). This procedure strongly enriches for recombined plasmids, as linearized plasmid monomers are unable to transform B. subtilis competent cells (Mottes et al., 1979). Recombined plasmids would not be cleaved by the restriction enzymes, and thus exist as a mixture of monomeric and
35 oligomeric forms well able to transform competent B. subtilis cells. One transformant thus obtained contained pSJ1359. This plasmid contains the origin of replication of pUB110 (Lacey and

Chopra, 1974, Gryczan et al., 1978, McKenzie et al., 1986), the pUB110 Rep protein gene, the kanamycin resistance gene, and the B. licheniformis α -amylase (amyL) promoter and signal peptide coding region perfectly fused to the DNA encoding the mature
5 part of the CGTase from Thermoanaerobacter sp. ATCC 53627.

3. Construction of a chromosomal integration vector.

A 1.4 kb BamHI fragment containing the pUB110 kanamycin
10 resistance gene (kan) was excised from plasmid pDN2904 (WO 91/09129), ligated to BglII digested pDN3000 (Fig. 6), transformed into E. coli SCS1 selecting ampicillin resistance (100 μ g/ml), and pPL1483 (Fig. 20) was recovered from one such transformant.

15

This plasmid was then combined with a Bacillus vector temperature sensitive for replication, plasmid pE194 (Horinouchi and Weisblum, 1982b). pPL1483 was digested with AccI, pE194 digested with ClaI, the two linearized plasmids
20 mixed, ligated, and transformed into B. subtilis DN1885 selecting kanamycin resistance (10 μ g/ml) at 30 °C. One such transformant contained pPL1487 (Fig. 21).

25 A 3'-terminal fragment of the amyL gene was excised from plasmid pDN1528 (Jørgensen, S. et al., 1991) as a 0.7 kb SalI-HindIII fragment, ligated to SalI+HindIII digested pUC19, and transformed to E. coli SJ2, selecting for ampicillin resistance (200 μ g/ml). One such transformant contained pSJ932 (Fig. 22).

30

Plasmid pSJ948 (Fig. 23) was obtained by insertion of a BglII linker into HindII digested pSJ932, once more selecting for ampicillin resistance (200 μ g/ml) upon transformation of SJ2.

35 pSJ980 (Fig. 24) was constructed by ligation of the 5.1 kb HindIII fragment of pPL1487 to HindIII digested pSJ948, selecting for kanamycin resistance (10 μ g/ml) in B. subtilis

DN1885 at 30 °C.

Finally, pSJ1391 (Fig. 25) was constructed by ligation of the
5 4.0 kb BglIII fragment of pSJ1359 to the 5.6 kb BglIII fragment
of pSJ980, selecting for kanamycin resistance (10 µg/ml) in
DN1885 at 30 °C. This plasmid contains, on a vector
temperature-sensitive for replication and conferring resistance
to kanamycin and erythromycin, the promoter and upstream region
10 (about 0.4 kb) from the B. licheniformis α-amylase gene (amyL),
the α-amylase/CGTase fusion gene (amyL-cgtA), and then about
0.7 kb from the 3'-region of the α-amylase gene ('amyL).

15 4. Transfer of the fusion gene to B. licheniformis and
integration in the chromosome.

An α-amylase producing strain of B. licheniformis was
20 transformed with pSJ1391 by the protoplast transformation pro-
cedure (Akamatzu, 1984). One regenerating, kanamycin resistant
colony was isolated, and was found to produce both α-amylase
and CGTase. Production of the two enzymes can be easily
distinguished by separating proteins in the culture supernatant
25 from shake flask cultures in BPX medium (WO 91/09129) on
isoelectric focusing gels (e.g. using the Pharmacia Phast sys-
tem), followed by overlaying with an agarose gel containing
1 % soluble starch and subsequent staining by iodine vapour.
The CGTase activity was detected at pI 4.5, the α-amylase
30 activity at pI 8.

When this transformant was analyzed for its plasmid content,
it turned out that a recombination event between the incoming
plasmid and the chromosome had taken place: A double
35 recombination had exchanged the chromosomal α-amylase (amyL)
gene and the plasmid borne amyL-cgtA fusion gene, so that the
plasmid isolated carried the amyL gene (B. subtilis DN1885

transformed with this plasmid produced α -amylase) whereas the amyL-cgtA fusion gene now resided on the chromosome (Fig. 26).

By propagation in TY medium (WO 91/09129) without kanamycin,
5 strains were isolated that had spontaneously lost their plasmid (SJ1599, SJ1603-1607).

The original B. licheniformis transformant was also subjected to experimental conditions to ensure chromosomal integration and subsequent excision of the plasmid, in order to promote
10 recombination events. The transformant was plated on LB agar (WO 91/09129) with 10 μ g/ml kanamycin at 50 °C, individual colonies restreaked a few times at 50 °C, and each then grown in successive overnight TY cultures at 30 °C without kanamycin
15 to permit plasmid excision and loss. Kana^S isolates from each original 50 °C colony were incubated in BPX shake flasks and production of either α -amylase or CGTase determined by analysis on isoelectric focusing gels as above. The plasmid free strains analyzed all produced either CGTase or α -amylase. CGTase
20 producing isolates are e.g. SJ1561-62, 1580-83, 1586-91 and 1595.

One strain, named SJ1608, appeared to produce CGTase in larger amounts than the others.

25

Southern blot analysis of strains SJ1561, 1562, 1599, 1606 and 1608 confirmed that these strains have the chromosomal amyL gene replaced by the amyL-cgtA gene.

30 The following results were obtained by quantitation of the CGTase activity produced on incubation in BPX shake flasks for 6 days at 37 °C (results from several experiments; the variation within each group of strains was mainly due to the use of different batches of shake flasks):

35

Strain	CGTase activity, arbitrary units
SJ1561-62, 1580-83, 1586-91, 1595, 1599, 1603-07	1 - 7.5
SJ1608	200 - 275

10

5. Promoter analysis.

We have investigated whether the large difference in CGTase production between strain SJ1608 and the other strains containing the amyL-cgTA gene was due to differences in the amyL promoter responsible for the CGTase expression.

The amyL promoter sequence of the B. licheniformis host strain is given in SEQ ID#4.

20

The promoter region from a number of the CGTase producing B. licheniformis strains was amplified from chromosomal DNA by the PCR technique (Saiki et al., 1988), using as primers one oligonucleotide corresponding to pos. 204-233 reading downstream through the amyL promoter, and another oligonucleotide corresponding in sequence to the 5'-end of the DNA encoding the mature CGTase and reading upstream. The sequence of this second oligonucleotide was 5'-CCTGTTGGATTATTACTGGG-3' (SEQ ID#5).

The amplified DNA fragment from each strain was excised from an agarose gel and directly sequenced, using as sequencing primers in the dideoxy method (Sanger et al., 1977) the same oligonucleotides that were used for PCR amplification.

The results of the sequence analysis reveal that one or both of two point mutations in the promoter region are responsible for the large difference in CGTase production observed.

Strains SJ1599 and 1603-06, all low-yielding, have the promoter sequence shown in SEQ ID#4. However, the high-yielding strain SJ1608 contains the promoter sequence shown in SEQ ID#6.

- 5 The differences occur at pos. 553, where SJ1608 contains a C instead of a T, and at pos. 593, where SJ1608 contains a A instead of a T.

10 The sequence of the amyL promoter present on pSJ1359 and pSJ1391 was determined using the PCR amplification and sequencing procedure described above. This showed that both plasmids contain the promoter sequence shown in SEQ ID#1, i.e. identical to the promoter sequence of SJ1608.

15

6. Analysis of the promoter effect on expression of the B. licheniformis α -amylase gene amyL

20 pSJ1755 (Fig. 28) was constructed by ligating the 3.3 kb BglIII-HindIII fragment from pDN1981 (cf. Example 2) to the 4.9 kb BglIII-HindIII fragment from pSJ1391 (Fig. 25), selecting for kanamycin resistance (10 μ g/ml) in DN1885 at 30°C. This plasmid contains the entire amyL gene with the promoter sequence shown
25 in SEQ ID#6 (the promoter found in the high-yielding CGTase strain SJ1608) on a vector which is temperature-sensitive for replication and conferring resistance to kanamycin and erythromycin.

30 The α -amylase-producing B. licheniformis strain from which SJ1608 was derived contained a chloramphenicol resistance gene inserted into the alkaline protease gene, thereby disrupting this gene and making the strain alkaline protease negative. A derivative strain, SJ1707, is identical to SJ1608 except that
35 the chloramphenicol resistance gene was replaced by an approximately 150 bp deletion which also makes the strain alkaline protease negative.

Plasmid pSJ1755 was introduced into strain SJ1707 by protoplast transformation, and replacement of the amyL-cgtA fusion gene by the amyL gene was achieved by integration/excision as described in Example 4.

5

Yields of α -amylase from the transformed strain SJ1707 in which the amyL gene is preceded by the promoter sequence shown in SEQ ID#6 were compared to the yield from the strain from which SJ1608 was derived and in which the amyL gene is preceded by the promoter sequence shown in SEQ ID#4.

10

The results obtained from BPX shake flask cultures incubated for 6 days at 37°C.

15

Promoter sequence	amylase, arbitrary units
SEQ ID#4	1
SEQ ID#6	105

20 It clearly appears from these results that the yield of α -amylase is greatly increased using the promoter sequence shown in SEQ ID#6.

REFERENCES.

- 5 Akamatzu, T., Sekiguchi, J. (1984). An improved method of protoplast regeneration for Bacillus species and its application to protoplast fusion and transformation. *Agric. Biol. Chem.*, 48, 651-655.
- 10 Diderichsen, B., Christiansen, L. (1988). Cloning of a maltogenic alpha-amylase from a Bacillus stearothermophilus. *FEMS Microbiology Letters*, 56, 53-60.
- 15 Jørgensen, P. L., Hansen, C. K., Poulsen, G. B., Diderichsen, B. (1990). In vivo genetic engineering: homologous recombination as a tool for plasmid construction. *Gene* 96, 37-41.
- 20 Diderichsen, B. In: Bacillus Molecular Genetics and Biotechnology Applications, A.T. Ganesan and J.A. Hoch, eds., Academic Press, 1986, pp. 35-46.
- 25 Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990). Cloning of aldB, which encodes α -acetolactate decarboxylase, an exoenzyme from Bacillus brevis. *Journal of Bacteriology*, 172, 4315-4321.
- 30 Jørgensen, P. L., Poulsen, G. B., Diderichsen, B. (1991). Cloning of a chromosomal α -amylase gene from Bacillus stearothermophilus. *FEMS Microbiology Letters*, 77, 271-276.
- 35 Jørgensen, S., Skov, K. W., Diderichsen, B. (1991). Cloning, sequence, and expression of a lipase gene from Pseudomonas cepacia: Lipase production in heterologous hosts requires two Pseudomonas genes. *Journal of Bacteriology*, 173, 559-567.
- Horinouchi, S., Weisblum, B. (1982a). Nucleotide sequence and

functional map of pC194, a plasmid that specifies chloramphenicol resistance. *J. Bacteriol.*, 150, 815-825.

Horinouchi, S., Weisblum, B. (1982b). Nucleotide sequence and functional map of pE194, a plasmid that specifies inducible resistance to macrolide, lincosamide, and streptogramin type B antibiotics. *J. Bacteriol.* 150, 804-814.

Kieser, T. (1984). Factors affecting the isolation of CCC DNA from *Streptomyces lividans* and *Escherichia coli*. *Plasmid* 12, 19-36.

Mandel, A., Higa, A. (1970). *J. Mol. Biol.* 53, 159-162.

Mottes, M., Grandi, G., Sgaramella, V., Canosi, U., Morelli, G., Trautner, T. A. (1979). Different specific activities of monomeric and oligomeric forms of plasmid DNA in transformation of *Bacillus subtilis* and *Escherichia coli*. *Mol. Gen. Genet.*, 174, 281-286.

20

Lacey, R., Chopra, I. (1974). Genetic studies of a multiresistant strain of *Staphylococcus aureus*. *J. Med. Microbiol.*, 7, 285-297.

McKenzie, T., Hoshino, T., Tanaka, T., Sueoka, N. (1986). The nucleotide sequence of pUB110: Some salient features in relation to replication and its regulation. *Plasmid* 15, 93-103.

Sanger, F., Nicklen, S., Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA*, 74, 5463-5467.

Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., Mullis, K. B., Erlich, H. A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239, 487-491.

Gryczan, T., Contente, S., Dubnau, D. (1978). Characterization of Staphylococcus aureus plasmids introduced by transformation into Bacillus subtilis. J. Bacteriol., 134, 318-329.

5 Yanish-Perron, C., Vieira, J., Messing, J. (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13 mp18 and pUC19 vectors. Gene 33, 103-119.

10 Yasbin, R. E., Williams, G. A., Young, F. E. (1975). J. Bacteriol. 121, 296-304.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Novo Nordisk A/S
- (B) STREET: Novo Alle
- (C) CITY: Bagsvaerd
- (E) COUNTRY: Denmark
- (F) POSTAL CODE (ZIP): 2880
- (G) TELEPHONE: +45 4444 8888
- (H) TELEFAX: +45 4449 3256
- (I) TELEX: 37304

(ii) TITLE OF INVENTION: A Bacillus Promoter

(iii) NUMBER OF SEQUENCES: 6

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 616 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Bacillus licheniformis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```

GCATGGGTC TTCTTTGTGC TTGGAAGCAG AGCCCAATAT TATCCCGAAA CGATAAAACG      60
GATGCTGAAG GAAGGAAACG AAGTCGGCAA CCATTCCTGG GACCCATCG TTATTGACAA      120
GGCTGTCAAA CGAAAAGCG TATCAGGAGA TTAACGACAC GCAAGAAATG ATCGAAAAAA      180
TCAGCGGACA CCTGCCTGTA CACTTGGGTC CTCCATACGG CGGGATCAAT GATTTCGTCC      240
GCTCGCTTTC CAATCTGAAG GTTTCATTGT GGGATGTTGA TCCGGAAGAT TGGAAGTACA      300
AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGACGGA AAAATOGTCT      360

```

TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCCTGCTGA AGAGATTATT AAAAAGCTGA 420
 AAGCAAAAGG CTATCAATTG GTAACIGTAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG 480
 GCTATTGAAT AAATGAGTAG AAAGGCCAT ATOGGGCGCTT TTCTTTTGGG AGAAAATATA 540
 GGGAAAATGG TANTTGTTAA AAATTCGGAA TATTTATACA ATATCATNNN NNNNNCATTG 600
 AAAGGGGAGG AGAATC 616

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 186 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Bacillus licheniformis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CTATCAATTG GTAACIGTAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG GCTATTGAAT 60
 AAATGAGTAG AAAGGCCAT ATOGGGCGCTT TTCTTTTGGG AGAAAATATA GGGAAAATGG 120
 TANTTGTTAA AAATTCGGAA TATTTATACA ATATCATNNN NNNNNCATTG AAAGGGGAGG 180
 AGAATC 186

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 66 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: synthetic

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

TCGACTGATC ACTTCCTGCC TCATTCCTGCA GCAGCGGGGG CACCGGATAC TTCAGTTTCT 60
 CTAGAG 66

(2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 616 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Bacillus licheniformis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GCATGCGTCC TTCITTTGTGC TTGGAAGCAG AGCCCAATAT TATCCCGAAA CGATAAAAACG	60
GATGCTGAAG GAAGGAAAAG AAGTCGGCAA CCATTCCTGG GACCCATCCG TTATTGACAA	120
GGCTGTCAAA CGAAAAAGCG TATCAGGAGA TTAACGACAC GCAAGAAATG ATCGAAAAAA	180
TCAGGGACA CCTGCCTGTA CACTTGGGTC CTCATAOCCG OGGGATCAAT GATTTCGTCC	240
GCTCGCTTTC CAATCTGAAG GTTTCATTGT GGGATGTTGA TCOGGAAGAT TGAAGTACA	300
AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGAOCGA AAAATOGTCT	360
TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCTGCTGA AGAGATTATT AAAAAGCTGA	420
AAGCAAAAGG CTATCAATTG GTAACGTGAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG	480
GCTATTGAAT AAATGAGTAG AAAGGCCAT ATCGGCGCTT TTCITTTTGGG AGAAAATATA	540
GGGAAAATGG TATTTGTTAA AAATTCGGAA TATTTATAACA ATATCATATG TTTACATTG	600
AAAGGGGAGG AGAATC	616

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: synthetic

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CCTGTGGAT TATTACTGGG 20

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 616 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (vi) ORIGINAL SOURCE:
 (A) ORGANISM: *Bacillus licheniformis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

GCATGCGTCC TTCTTTGTC TTGGAAGCAG AGCCCAATAT TATCCCGAAA CGATAAAAACG      60
GATGCTGAAG GAAGGAAAACG AAGTOGGCAA CCATTCCCTGG GACCCATCCG TTATTGACAA      120
GGCTGTCAAA CGAAAAAGCG TATCAGGAGA TTAACGACAC GCAAGAAATG ATCGAAAAAA      180
TCAGCGGACA CCTGCCCTGA CACTTGGGTC CICCATAACG GGGATCAAT GATTCOGTCC      240
GCTCGCITTC CAATCTGAAG GTTTCATTGT GGGATGTTGA TCOGGAAGAT TGGAAGTACA      300
AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGACGGA AAAATCGTCT      360
TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCTGCTGA AGAGATTATT AAAAAGCTGA      420
AAGCAAAAGG CTATCAATTG GTAACGTGAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG      480
GCTATTGAAT AAATGAGTAG AAAGOGCCAT ATCGGCGCIT TTCTTTTGGG AGAAAATATA      540
GGGAAAATGG TACTTGTATA AAATTCGGAA TATTTATACA ATATCATATG TTACACATTG      600
AAAGGGGAGG AGAATC

```

CLAIMS

1. A Bacillus promoter included in the following DNA sequence
- 5
- GCATGCGTCC TTCTTTGTGC TTGGAAGCAG AGCCCAATAT TATCCCGAAA
 CGATAAACG GATGCTGAAG GAAGGAAACG AAGTCGGCAA CCATTCCTGG
 GACCCATCCG TTATTGACAA GGCTGTCAAA CGAAAAAGCG TATCAGGAGA
 TTAACGACAC GCAAGAAATG ATCGAAAAAA TCAGCGGACA CCTGCCTGTA
 10 CACTTGCGTC CTCCATACGG CGGGATCAAT GATTCGGTCC GCTCGCTTTC
 CAATCTGAAG GTTTCATTGT GGGATGTTGA TCCGGAAGAT TGGAAGTACA
 AAAATAAGCA AAAGATTGTC AATCATGTCA TGAGCCATGC GGGAGACGGA
 AAAATCGTCT TAATGCACGA TATTTATGCA ACGTTCGCAG ATGCTGCTGA
 AGAGATTATT AAAAAGCTGA AAGCAAAAGG CTATCAATTG GTAAGTGTAT
 15 CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG GCTATTGAAT AAATGAGTAG
 AAAGCGCCAT ATCGGCGCTT TTCTTTTGGG AGAAAATATA GGGAAAATGG
 TAN¹TTGTTAA AAATTCGGAA TATTTATACA ATATCATN²N³N⁴
 N⁵N⁶N⁷N⁸N⁹CATTG AAAGGGGAGG AGAATC (SEQ ID#1)
- 20 wherein each of N¹-N⁹ is A, T, C or G with the exception that
 N²-N⁹ do not together form the sequence ATGTTTCA or GTGTTTCA,
 or a functional homologue of said sequence.
- 25 2. A promoter according to claim 1, wherein N¹ is C or T.
3. A promoter according to claim 1, wherein N⁷ is A, G or C.
4. A promoter according to claim 1, wherein N¹ is C and N⁷ is
 30 A.
5. A promoter according to claim 1, wherein N²-N⁹ together form
 the sequence ATGTTACA.
- 35 6. A promoter according to claim 5, wherein N¹ is C.
7. A promoter according to claim 1, which is included in the

following DNA sequence

CTATCAATTG GTAACTGTAT CTCAGCTTGA AGAAGTGAAG AAGCAGAGAG
 GCTATTGAAT AAATGAGTAG AAAGCGCCAT ATCGGCGCTT TTCTTTTGGGA
 5 AGAAAATATA GGGAAAATGG TAN¹TTGTAA AAATTCGGAA TATTTATACA
 ATATCATN²N³N⁴ N⁵N⁶N⁷N⁸N⁹CATTG AAAGGGGAGG AGAATC (SEQ ID#2)

wherein N¹-N⁹ has the meaning indicated above.

10 8. A promoter according to any of claims 1-7, which is derived from a Bacillus licheniformis gene and in particular it is a variant of a Bacillus licheniformis α -amylase promoter.

15 9. A DNA construct comprising a DNA sequence coding for a protein of interest preceded by a promoter sequence according to any of claims 1-8.

20 10. A DNA construct according to claim 9, wherein the protein of interest is an enzyme such as an α -amylase, cyclodextrin glycosyl transferase or protease.

11. A DNA construct according to claim 9 or 10, which further comprises a DNA sequence coding for a signal peptide.

25 12. A DNA construct according to claim 11, wherein the signal peptide is the B. licheniformis α -amylase signal peptide.

13. A recombinant expression vector comprising a DNA construct according to any of claims 9-12.

30

14. A host cell transformed with a DNA construct according to any of claims 9-12, or with a vector according to claim 13.

15. A host cell according to claim 14, which is a strain of Bacillus, in particular a strain of Bacillus licheniformis, Bacillus lentus, Bacillus brevis, Bacillus stearothermophilus, Bacillus alkalophilus, Bacillus amyloliquefaciens, Bacillus

coagulans Bacillus thuringiensis or Bacillus subtilis.

16. A process for producing a protein in Bacilli comprising
culturing a Bacillus host cell transformed with a DNA construct
5 according to any of claims 9-12, or with a vector according to
claim 13 under conditions permitting production of said
protein, and recovering the resulting protein from the culture.




Fig. 1




Fig. 2

3/28




Fig. 3

4/28




Fig. 4

5/28




Fig. 5

6/28




Fig. 6

7/28




Fig. 7

8/28




Fig. 8

9/28




Fig. 9

10/28




Fig. 10

11/28




Fig. 11

12/28




Fig. 12

13/28




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17




Fig. 18




Fig. 19




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25

26/28




Fig. 26

27/28




Fig. 27




Fig. 28

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 92/00338

A. CLASSIFICATION OF SUBJECT MATTER		
IPC5: C12N 15/75, C12N 15/56, C12N 9/28 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC5: C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
WPI, CA, BIOSIS, EMBL		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Chemical Abstracts, Volume 111, No 1, 3 July 1989 (03.07.89), (Columbus, Ohio, USA), Laoide, Brid M et al, "Bacillus licheniformis alfa-amylase gene, amyL, is subject to promoter-independent catabolite repression in Bacillus subtilis", page 171, THE ABSTRACT No 1695c, J. Bacteriol 1989, 171 (5), 2435-2442 --	1-4,7-16
A	Chemical Abstracts, Volume 102, No 7, 18 February 1985 (18.02.85), (Columbus, Ohio, USA), Sibakov, Mervi et al, "Isolation and the 5'-end nucleotide sequence of Bacillus licheniformis alfa-amylase gene", page 149, THE ABSTRACT No 56980m, Eur. J. Biochem. 1984, 145 (3), 567-572 --	1-16
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance		"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"B" earlier document but published on or after the international filing date		"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
22 February 1993	26 -02- 1993	
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer Yvonne Siösteen Telephone No. +46 8 782 25 00	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 92/00338

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>Chemical Abstracts, Volume 104, No 3, 20 January 1986 (20.01.86), (Columbus, Ohio, USA), Yuuki, Toshifumi et al, "Complete nucleotide sequence of a gene coding for heat- and pH-stable alfa-amylase of Bacillus licheniformis: comparison of the amino acid sequences of three bacterial liquefying alfa-amylases deduced from the DNA..", page 147, THE ABSTRACT No 15859b, J. Biochem. 1985, 98 (5), 1147-1156</p> <p style="text-align: center;">-- -----</p>	1-16