

(19) **DANMARK**



Patent- og  
Varemærkestyrelsen

(10) **DK/EP 2485592 T3**

(12) **Oversættelse af  
europæisk patentskrift**

- 
- (51) Int.Cl.: **A 01 N 43/40 (2006.01)**      **A 01 N 43/653 (2006.01)**      **A 01 N 47/06 (2006.01)**  
**A 01 N 47/38 (2006.01)**      **A 01 P 3/00 (2006.01)**      **C 07 D 405/12 (2006.01)**  
**C 07 D 405/14 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-09-05**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-06-01**
- (86) Europæisk ansøgning nr.: **10822587.1**
- (86) Europæisk indleveringsdag: **2010-10-06**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-08-15**
- (86) International ansøgning nr.: **US2010051598**
- (87) Internationalt publikationsnr.: **WO2011044213**
- (30) Prioritet: **2009-10-07 US 249479 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Dow AgroSciences LLC, 9330 Zionsville Road, Indianapolis, Indiana 46268, USA**
- (72) Opfinder: **OWEN, W. John, 12421 Pebblepointe Pass, Carmel, IN 46033, USA**  
**YAO, Chenglin, 14178 Esprit Drive, Westfield, IN 46074, USA**
- (74) Fuldmægtig i Danmark: **Chas. Hude A/S, H.C. Andersens Boulevard 33, 1780 København V, Danmark**
- (54) Benævnelse: **SYNERGISTISKE FUNGICIDE BLANDINGER TIL SVAMPEBEKÆMPELSE I KORNSORTER**
- (56) Fremdragne publikationer:  
**EP-A1- 1 516 874**  
**US-A1- 2007 060 579**  
**US-A1- 2007 066 629**  
**US-B1- 6 355 660**  
**ANONYMOUS: "Synergistic Fungicidal Compositions of Heterocyclic Aromatic Amides and Triazoles", IP.COM JOURNAL, IP.COM INC., WEST HENRIETTA, NY, US, 20 July 2004 (2004-07-20), XP013020905, ISSN: 1533-0001 'Science For A Better Life.' BAYER CROPSCIENCE, [Online] June 2008, page 28 Retrieved from the Internet: <URL:www.BCS\_Roadshow\_London\_20080620\_Schei tza\_Handout\_final\_web[1].pdf> [retrieved on 2010-11-23]**  
**J. E. PARKER ET AL: 'Mechanism of Binding of Prothioconazole to Mycosphaerella graminicola CYP51 Differs from That of Other Azole Antifungals' APPLIED AND ENVIRONMENTAL MICROBIOLOGY vol. 77, no. 4, 17 December 2010, pages 1460 - 1465, XP055163788 DOI: 10.1128/AEM.01332-10 ISSN: 0099-2240**  
**EUGENE O'SULLIVAN ET AL: 'Fungicide Resistance - an increasing problem' TEAGASC - PUBLICATIONS, [Online] 31 January 2007, pages 1 - 13, XP055163833 Retrieved from the Internet: <URL:http://www.teagasc.ie/publications/2007/20070131/ntc2007paper05.asp> [retrieved on 2015-01-21]**



## DESCRIPTION

### Field of the Invention

[0001] This invention concerns a synergistic fungicidal composition containing (a) a compound of Formula I, II, III, IV or V and (b) at least one fungicide selected from the group consisting of prothioconazole and prochloraz.

### Background of the Invention

[0002] Fungicides are compounds, of natural or synthetic origin, which act to protect plants against damage caused by fungi. Current methods of agriculture rely heavily on the use of fungicides. In fact, some crops cannot be grown usefully without the use of fungicides. Using fungicides allows a grower to increase the yield and the quality of the crop, and consequently, increase the value of the crop. In most situations, the increase in value of the crop is worth at least three times the cost of the use of the fungicide.

[0003] However, no one fungicide is useful in all situations and repeated usage of a single fungicide frequently leads to the development of resistance to that and related fungicides. Consequently, research is being conducted to produce fungicides and combinations of fungicides that are safer, that have better performance, that require lower dosages, that are easier to use, and that cost less.


[0004] Synergism occurs when the activity of two or more compounds exceeds the activities of the compounds when used alone. Synergistic mixtures of picolinamides and a second compound selected from, among others, epoxiconazole are described in an anonymous contribution entitled "Synergistic Fungicidal Compositions of Heterocyclic Aromatic Amides and Triazoles", published on IP.COM on July 20, 2004.


### Summary of the Invention

[0005] It is an object of this invention to provide synergistic compositions comprising fungicidal compounds. It is a further object of this invention to provide processes that use these synergistic compositions. The synergistic compositions are capable of preventing or curing, or both, diseases caused by fungi of the classes *Ascomycetes* and *Basidiomycetes*. In addition, the synergistic compositions have improved efficacy against the *Ascomycete* and *Basidiomycete* pathogens, including leaf blotch and brown rust of wheat. In accordance with this invention, synergistic compositions are provided along with methods for their use.


### Detailed Description of the Invention

[0006] The present invention concerns a synergistic fungicidal mixture comprising an fungicidally effective amount of (a) a compound of Formula I, II, III, IV or V, and (b) at least one fungicide selected from the group consisting of prothioconazole and prochloraz.







II



III



IV



V

**[0007]** Prochloraz is the common name for *N*-propyl-*N*-[2,4,6-trichlorophenoxy]ethyl]imidazole-1-carboxamide. Its fungicidal activity is described in The Pesticide Manual, Fifteenth Edition, 2009. Prochloraz provides control against a wide variety of pathogens at application rates between 400 to 600 g a.i./ha.

**[0008]** Prothioconazole is the common name for 2-[2-(1-chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3*H*-1,2,4-triazole-3-thione. Its fungicidal activity is described in The Pesticide Manual, Fourteenth Edition, 2006. Prothioconazole is used for control of diseases such as eyespot, *Fusarium* ear blight, leaf blotch, rust and powdery mildew by foliar application in wheat, barley and other crops.

[0009] In the composition of this invention, the weight ratio of the compounds of Formula I-V to prothioconazole at which the fungicidal effect is synergistic lies within the range of between about 1:10 and about 10:1.

[0010] The weight ratio of the compounds of Formula I-V to prochloraz at which the fungicidal effect is synergistic lies within the range of between about 1:10 and about 10:1.

[0011] The rate at which the synergistic composition is applied will depend upon the particular type of fungus to be controlled, the degree of control required and the timing and method of application. In general, the composition of the invention can be applied at an application rate of between about 65 grams per hectare (g/ha) and about 2300 g/ha based on the total amount of active ingredients in the composition. Prothioconazole is applied at a rate between about 50 g/ha and about 200 g/ha and the compound of Formula I-V is applied at a rate between about 35 g/ha and about 300 g/ha. Prochloraz is applied at a rate between about 50 g/ha and about 450 g/ha and the compound of Formula I-V is applied at a rate between about 35 g/ha and about 300 g/ha.

[0012] The components of the synergistic mixture of the present invention can be applied either separately or as part of a multipart fungicidal system.

[0013] The synergistic mixture of the present invention can be applied in conjunction with one or more other fungicides to control a wider variety of undesirable diseases. When used in conjunction with other fungicide(s), the presently claimed compounds may be formulated with the other fungicide(s), tank mixed with the other fungicide(s) or applied sequentially with the other fungicide(s). Such other fungicides may include 2-(thiocyanatomethylthio)-benzothiazole, 2-phenylphenol, 8-hydroxyquinoline sulfate, ametoctradin, amisulbrom, antimycin, *Ampelomyces quisqualis*, azaconazole, azoxystrobin, *Bacillus subtilis*, *Bacillus subtilis* strain QST713, benalaxyl, benomyl, bentiavalicarbisopropyl, benzylaminobenzene-sulfonate (BABS) salt, bicarbonates, biphenyl, bismethiazol, bitertanol, bixafen, blastidicin-S, borax, Bordeaux mixture, boscalid, bromuconazole, bupirimate, calcium polysulfide, captafol, captan, carbendazim, carboxin, carpropamid, carvone, chlazafenone, chloroneb, chlorothalonil, chlozolinate, *Coniothyrium minitans*, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, copper sulfate (tribasic), cuprous oxide, cyazofamid, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, diammonium ethylenebis-(dithiocarbamate), dichlofluanid, dichlorophen, diclocymet, diclomezine, dichloran, diethofencarb, difenoconazole, difenzoquat ion, diflumetorim, dimethomorph, dimoxystrobin, diniconazole, diniconazole-M, dinobuton, dinocap, diphenylamine, dithianon, dodemorph, dodemorph acetate, dodine, dodine free base, edifenphos, enestrobin, enestroburin, epoxiconazole, ethaboxam, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenoxanil, fempiclonil, fenpropidin, fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, flumorph, fluopicolide, fluopyram, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, formaldehyde, fosetyl, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, GY-81, hexachlorobenzene, hexaconazole, hymexazol, imazalil, imazalil sulfate, imibenconazole, iminocladine, iminocladine triacetate, iminocladine tris(albesilate), iodocarb, ipconazole, ipfenpyrazolone, iprobenfos, iprodione, iprovalicarb, isoprothiolane, isopyrazam, isotianil, kasugamycin, kasugamycin hydrochloride hydrate, kresoxim-methyl, laminarin, mancozeb, mancozeb, mandipropamid, maneb, mefenoxam, mepanipyrim, mepronil, meptyl-dinocap, mercuric chloride, mercuric oxide, mercurous chloride, metalaxyl, metalaxyl-M, metam, metam-ammonium, metam-potassium, metam-sodium, metconazole, methasulfocarb, methyl iodide, methyl isothiocyanate, metiram, metominostrobin, metrafenone, mildiomicin, myclobutanil, nabam, nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid (fatty acids), oryastrobin, oxadixyl, oxine-copper, oxpoconazole fumarate, oxycarboxin, pefurazoate, penconazole, pencycuron, penflufen, pentachlorophenol, pentachlorophenyl laurate, penthiopyrad, phenylmercury acetate, phosphonic acid, phthalide, picoxystrobin, polyoxin B, polyoxins, polyoxorim, potassium bicarbonate, potassium hydroxyquinoline sulfate, probenazole, prochloraz, procymidone, propamocarb, propamocarb hydrochloride, propiconazole, propineb, proquinazid, prothioconazole, pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyrazophos, pyribencarb, pyributicarb, pyrifenoxy, pyrimethanil, pyriofenone, pyroquilon, quinochloramine, quinoxifen, quintozone, *Reynoutria sachalinensis* extract, sedaxane, silthiofam, simeconazole, sodium 2-phenylphenoxide, sodium bicarbonate, sodium pentachlorophenoxide, spiroxamine, sulfur, SYP-Z048, tar oils, tebuconazole, tebufloquin, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tiadinil, tolclofos-methyl, tolyfluanid, triadimefon, triadimenol, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin, valifenalate, valiphenal, vinclozolin, zineb, ziram, zoxamide, *Candida oleophila*, *Fusarium oxysporum*, *Gliocladium* spp., *Phlebiopsis gigantea*, *Streptomyces griseoviridis*, *Trichoderma* spp., (RS)-N-(3,5-dichlorophenyl)-2-(methoxymethyl)-succinimide, 1,2-dichloropropane, 1,3-dichloro-1,1,3,3-tetrafluoroacetone hydrate, 1-chloro-2,4-dinitronaphthalene, 1-chloro-2-nitropropane, 2-(2-heptadecyl-2-imidazolin-1-yl)ethanol, 2,3-dihydro-5-phenyl-1,4-dithiine 1,1,4,4-tetraoxide, 2-methoxyethylmercury acetate, 2-methoxyethylmercury chloride, 2-methoxyethylmercury silicate, 3-(4-chlorophenyl)-5-methylrhodanine, 4-(2-nitroprop-1-enyl)phenyl thiocyanateme, ampropylfos, anilazine, azithiram, barium polysulfide, Bayer 32394, benodanil, benquinox, bentalurone, benzamacril; benzamacril-isobutyl, benzamorf, binapacryl, bis(methylmercury) sulfate, bis(tributyltin) oxide, buthiobate, cadmium calcium copper zinc chromate sulfate, carbamorph, CECA,

chlorthalozole, chloranilmethan, chlorfenazole, chlorquinox, dimbazole, copper bis(3-phenylsalicylate), copper zinc chromate, cufraneb, cupric hydrazinium sulfate, cuprobam, cyclafuramid, cypendazole, cyprofuram, decafentin, dichlone, dichlozoline, diclobutrazol, dimethirimol, dinocron, dinosulfon, dinoterbon, dipyrithione, ditalimfos, dodicin, drazoxolon, EBP, ESBP, etaconazole, etem, ethirim, fenaminosulf, fenapanil, fenitropan, fluotrimazole, furcarbanil, furconazole, furconazole-cis, furemefenox, furophanate, glyodine, griseofulvin, halacrinat, Hercules 3944, hexylthiofos, ICIA0858, isopamphos, isovaldione, mebenil, mecarbinzid, metazoxolon, methfuroxam, methylmercury dicyandiamide, metsulfovax, milneb, mucochloric anhydride, myclozolin, N-3,5-dichlorophenyl-succinimide, N-3-nitrophenylitaconimide, natamycin, N-ethylmercurio-4-toluenesulfonanilide, nickel bis(dimethyldithiocarbamate), OCH, phenylmercury dimethyldithiocarbamate, phenylmercury nitrate, phosdiphen, prothiocarb; prothiocarb hydrochloride, pyracarbolid, pyridinil, pyroxychlor, pyroxyfur, quinacetol; quinacetol sulfate, quinazamid, quinconazole, rabenzazole, salicylanilide, SSF-109, sultropen, tecoram, thiadifluor, thicyofen, thiochlorfenphim, thiophanate, thioquinox, tioxyimid, triamphos, triarimol, triazbutil, trichlamide, urbacid, zarilamid, and any combinations thereof.

**[0014]** The compositions of the present invention are preferably applied in the form of a formulation comprising a composition of (a) a compound of Formula I and (b) at least one fungicide selected from the group consisting of prothioconazole, and prochloraz, together with a phytologically acceptable carrier.

**[0015]** Concentrated formulations can be dispersed in water, or another liquid, for application, or formulations can be dust-like or granular, which can then be applied without further treatment. The formulations are prepared according to procedures which are conventional in the agricultural chemical art, but which are novel and important because of the presence therein of a synergistic composition.

**[0016]** The formulations that are applied most often are aqueous suspensions or emulsions. Either such water-soluble, water-suspendable, or emulsifiable formulations are solids, usually known as wettable powders, or liquids, usually known as emulsifiable concentrates, aqueous suspensions, or suspension concentrates. The present invention contemplates all vehicles by which the synergistic compositions can be formulated for delivery and use as a fungicide.

**[0017]** As will be readily appreciated, any material to which these synergistic compositions can be added may be used, provided they yield the desired utility without significant interference with the activity of these synergistic compositions as antifungal agents.

**[0018]** Wettable powders, which may be compacted to form water-dispersible granules, comprise an intimate mixture of the synergistic composition, a carrier and agriculturally acceptable surfactants. The concentration of the synergistic composition in the wettable powder is usually from about 10% to about 90% by weight, more preferably about 25% to about 75% by weight, based on the total weight of the formulation. In the preparation of wettable powder formulations, the synergistic composition can be compounded with any of the finely divided solids, such as prophyllite, talc, chalk, gypsum, Fuller's earth, bentonite, attapulgite, starch, casein, gluten, montmorillonite clays, diatomaceous earths, purified silicates or the like. In such operations, the finely divided carrier is ground or mixed with the synergistic composition in a volatile organic solvent. Effective surfactants, comprising from about 0.5% to about 10% by weight of the wettable powder, include sulfonated lignins, naphthalenesulfonates, alkylbenzenesulfonates, alkyl sulfates, and non-ionic surfactants, such as ethylene oxide adducts of alkyl phenols.

**[0019]** Emulsifiable concentrates of the synergistic composition comprise a convenient concentration, such as from about 10% to about 50% by weight, in a suitable liquid, based on the total weight of the emulsifiable concentrate formulation. The components of the synergistic compositions, jointly or separately, are dissolved in a carrier, which is either a water-miscible solvent or a mixture of water-immiscible organic solvents, and emulsifiers. The concentrates may be diluted with water and oil to form spray mixtures in the form of oil-in-water emulsions. Useful organic solvents include aromatics, especially the high-boiling naphthalenic and olefinic portions of petroleum such as heavy aromatic naphtha. Other organic solvents may also be used, such as, for example, terpenic solvents, including rosin derivatives, aliphatic ketones, such as cyclohexanone, and complex alcohols, such as 2-ethoxyethanol.

**[0020]** Emulsifiers which can be advantageously employed herein can be readily determined by those skilled in the art and include various nonionic, anionic, cationic and amphoteric emulsifiers, or a blend of two or more emulsifiers. Examples of nonionic emulsifiers useful in preparing the emulsifiable concentrates include the polyalkylene glycol ethers and condensation products of alkyl and aryl phenols, aliphatic alcohols, aliphatic amines or fatty acids with ethylene oxide, propylene oxides such as the ethoxylated alkyl phenols and carboxylic esters solubilized with the polyol or polyoxyalkylene. Cationic emulsifiers include quaternary ammonium compounds and fatty amine salts. Anionic emulsifiers include the oil-soluble salts (e.g., calcium) of alkylaryl sulfonic acids, oil-soluble salts or sulfated polyglycol ethers and appropriate salts of phosphated polyglycol ether.

**[0021]** Representative organic liquids which can be employed in preparing the emulsifiable concentrates of the present invention

are the aromatic liquids such as xylene, propyl benzene fractions, or mixed naphthalene fractions, mineral oils, substituted aromatic organic liquids such as dioctyl phthalate, kerosene, dialkyl amides of various fatty acids, particularly the dimethyl amides of fatty glycols and glycol derivatives such as the *n*-butyl ether, ethyl ether or methyl ether of diethylene glycol, and the methyl ether of triethylene glycol. Mixtures of two or more organic liquids are also often suitably employed in the preparation of the emulsifiable concentrate. The preferred organic liquids are xylene, and propyl benzene fractions, with xylene being most preferred. The surface-active dispersing agents are usually employed in liquid formulations and in the amount of from 0.1 to 20 percent by weight of the combined weight of the dispersing agent with the synergistic compositions. The formulations can also contain other compatible additives, for example, plant growth regulators and other biologically active compounds used in agriculture.

**[0022]** Aqueous suspensions comprise suspensions of one or more water-insoluble compounds, dispersed in an aqueous vehicle at a concentration in the range from about 5% to about 70% by weight, based on the total weight of the aqueous suspension formulation. Suspensions are prepared by finely grinding the components of the synergistic combination either together or separately, and vigorously mixing the ground material into a vehicle comprised of water and surfactants chosen from the same types discussed above. Other ingredients, such as inorganic salts and synthetic or natural gums, may also be added to increase the density and viscosity of the aqueous vehicle. It is often most effective to grind and mix at the same time by preparing the aqueous mixture and homogenizing it in an implement such as a sand mill, ball mill, or piston-type homogenizer.

**[0023]** The synergistic composition may also be applied as a granular formulation, which is particularly useful for applications to the soil. Granular formulations usually contain from about 0.5% to about 10% by weight of the compounds, based on the total weight of the granular formulation, dispersed in a carrier which consists entirely or in large part of coarsely divided attapulgite, bentonite, diatomite, clay or a similar inexpensive substance. Such formulations are usually prepared by dissolving the synergistic composition in a suitable solvent and applying it to a granular carrier which has been preformed to the appropriate particle size, in the range of from about 0.5 to about 3 mm. Such formulations may also be prepared by making a dough or paste of the carrier and the synergistic composition, and crushing and drying to obtain the desired granular particle.

**[0024]** Dusts containing the synergistic composition are prepared simply by intimately mixing the synergistic composition in powdered form with a suitable dusty agricultural carrier, such as, for example, kaolin clay, ground volcanic rock, and the like. Dusts can suitably contain from about 1% to about 10% by weight of the synergistic composition/carrier combination.

**[0025]** The formulations may contain agriculturally acceptable adjuvant surfactants to enhance deposition, wetting and penetration of the synergistic composition onto the target crop and organism. These adjuvant surfactants may optionally be employed as a component of the formulation or as a tank mix. The amount of adjuvant surfactant will vary from 0.01 percent to 1.0 percent volume/volume (v/v) based on a spray-volume of water, preferably 0.05 to 0.5 percent. Suitable adjuvant surfactants include ethoxylated nonyl phenols, ethoxylated synthetic or natural alcohols, salts of the esters or sulfosuccinic acids, ethoxylated organosilicones, ethoxylated fatty amines and blends of surfactants with mineral or vegetable oils.

**[0026]** The formulations may optionally include combinations that can comprise at least 1% by weight of one or more of the synergistic compositions with another pesticidal compound. Such additional pesticidal compounds may be fungicides, insecticides, nematocides, miticides, arthropodocides, bactericides or combinations thereof that are compatible with the synergistic compositions of the present invention in the medium selected for application, and not antagonistic to the activity of the present compounds. Accordingly, in such embodiments the other pesticidal compound is employed as a supplemental toxicant for the same or for a different pesticidal use. The pesticidal compound and the synergistic composition can generally be mixed together in a weight ratio of from 1:100 to 100:1.

**[0027]** The present invention includes within its scope methods for the control or prevention of fungal attack. These methods comprise applying to the locus of the fungus, or to a locus in which the infestation is to be prevented (for example applying to wheat or barley plants), a fungicidally effective amount of the synergistic composition. The synergistic composition is suitable for treatment of various plants at fungicidal levels, while exhibiting low phytotoxicity. The synergistic composition is useful in a protectant or eradicator fashion. The synergistic composition is applied by any of a variety of known techniques, either as the synergistic composition or as a formulation comprising the synergistic composition. For example, the synergistic compositions may be applied to the roots, seeds or foliage of plants for the control of various fungi, without damaging the commercial value of the plants. The synergistic composition is applied in the form of any of the generally used formulation types, for example, as solutions, dusts, wettable powders, flowable concentrates, or emulsifiable concentrates. These materials are conveniently applied in various known fashions.

**[0028]** The synergistic composition has been found to have significant fungicidal effect, particularly for agricultural use. The synergistic composition is particularly effective for use with agricultural crops and horticultural plants, or with wood, paint, leather

or carpet backing.

[0029] In particular, the synergistic composition is effective in controlling a variety of undesirable fungi that infect useful plant crops. The synergistic composition may be used against a variety of *Ascomycete* and *Basidiomycete* fungi, including for example the following representative fungi species: wheat brown rust (*Puccinia recondita*; Bayer code PUCCRT); stripe rust of wheat (*Puccinia striiformis*; Bayer code PUCST); leaf blotch of wheat (*Mycosphaerella graminicola*; anamorph: *Septoria tritici*; Bayer code SEPTTR); glume blotch of wheat (*Leptosphaeria nodorum*; Bayer code LEPTNO; anamorph: *Stagonospora nodorum*); spot blotch of barley (*Cochliobolus sativum*; Bayer code COCHSA; anamorph: *Helminthosporium sativum*); leaf spot of sugar beets (*Cercospora beticola*; Bayer code CERCB); leaf spot of peanut (*Mycosphaerella arachidis*; Bayer code MYCOAR; anamorph: *Cercospora arachidicola*); cucumber anthracnose (*Glomerella lagenarium*; anamorph: *Colletotrichum lagenarium*; Bayer code COLLLA) and black sigatoka disease of banana (*Mycosphaerella fijiensis*; BAYER code MYCOFI). It will be understood by those in the art that the efficacy of the synergistic compositions for one or more of the foregoing fungi establishes the general utility of the synergistic compositions as fungicides.

[0030] The synergistic compositions have a broad range of efficacy as a fungicide. The exact amount of the synergistic composition to be applied is dependent not only on the relative amounts of the components, but also on the particular action desired, the fungal species to be controlled, and the stage of growth thereof, as well as the part of the plant or other product to be contacted with the synergistic composition. Thus, formulations containing the synergistic composition may not be equally effective at similar concentrations or against the same fungal species.

[0031] The synergistic compositions are effective in use with plants in a disease-inhibiting and phytologically acceptable amount. The term "disease-inhibiting and phytologically acceptable amount" refers to an amount of the synergistic composition that kills or inhibits the plant disease for which control is desired, but is not significantly toxic to the plant. The exact concentration of synergistic composition required varies with the fungal disease to be controlled, the type of formulation employed, the method of application, the particular plant species, climate conditions, and the like.

[0032] The present compositions can be applied to fungi or their locus by the use of conventional ground sprayers, granule applicators, and by other conventional means known to those skilled in the art.

[0033] The following examples are provided to further illustrate the invention.

#### **Examples**

##### **Evaluation of Curative and Protectant Activity of Fungicide Mixtures vs. Leaf Blotch of Wheat (*Mycosphaerella graminicola*; anamorph: *Septoria tritici*; Bayer code: SEPTTR)**

[0034] Wheat plants (variety Yuma) were grown from seed in a greenhouse in plastic pots with a surface area of 27.5 square centimeters (cm<sup>2</sup>) containing 50% mineral soil/50% soil-less Metro mix, with 8-12 seedlings per pot. The plants were employed for testing when first leaf was fully emerged, which typically took 7 to 8 days after planting. Test plants were inoculated with an aqueous spore suspension of *Septoria tritici* either 3 days prior to (3-day curative test) or 1 day after fungicide treatments (1-day protectant test). After inoculation the plants were kept in 100% relative humidity (one day in a dark dew chamber followed by two days in a lighted mist chamber) to permit spores to germinate and infect the leaf. The plants were then transferred to a greenhouse for disease to develop.

##### **Evaluation of Curative Activity of Fungicide Mixtures vs. Wheat Brown Rust (*Puccinia recondita*; Bayer code: PUCCRT)**

[0035] Yuma wheat seedlings were grown as described above, and inoculated with an aqueous spore suspension of *Puccinia recondita* 3 days prior to or 1-day after fungicide treatment. After inoculation, plants were kept in 100% relative humidity for 24 hours in a dark dew room to allow spores to germinate and infect plants. The plants were then transferred to a greenhouse for disease to develop.

[0036] Treatments consisted of fungicides compound I, II, III, IV, V, prothioconazole and prochloraz, either using individually or as



two-way mixture with compounds I-V. Technical grades of materials were dissolved in acetone to make stock solutions, which were then used to perform dilutions in acetone either for each individual fungicide component or for the two-way mixture. Desired fungicide rates were obtained after mixing dilutions with 9 volumes of water containing 110 parts per million (ppm) Triton X-100. Twenty milliliter (mL) fungicide solutions were applied onto 12 pots of plants using an automated booth sprayer, which utilized two 6218-1/4 JAUPM spray nozzles operating at 20 pounds per square inch (psi) set at opposing angles to cover both leaf surfaces. All sprayed plants were allowed to air dry prior to further handling. Control plants were sprayed in the same manner with the solvent blank.

[0037] When disease fully developed on the control plants, infection levels were assessed on treated plants visually and scored on a scale of 0 to 100 percent. Percentage of disease control was then calculated using the ratio of disease on treated plants relative to control plants.

[0038] Colby's equation was used to determine the fungicidal effects expected from the mixtures. (See Colby, S. R. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 1967, 15, 20-22.)

[0039] The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:  
 Expected =  $A + B - (A \times B/100)$

A = observed efficacy of active component A at the same concentration as used in the mixture;

B = observed efficacy of active component B at the same concentration as used in the mixture.

[0040] Representative synergistic interactions are presented in the following Tables 1-12.

%DC Obs= Percent disease control observed

%DC Exp= Percent disease control expected

Synergism factor = %DC Obs/%DC Exp

Table 1: Synergistic interactions of compound I and other fungicides in 3-day curative (3DC) *Septoria tritici* (SEPTTR) tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound I+prothioconazole	1.35+1.8	99	83	1.19
Compound I+prothioconazole	0.45+0.6	69	7	10.50

Table 2: Synergistic interactions of compound I and other fungicides in 1-day protectant (1DP) SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound I+prothioconazole	0.1+1.56	94	70	1.34
Compound I+prothioconazole	0.15+0.2	46	15	3.01
Compound I+prochloraz	0.15+0.45	35	7	5.00

Table 3: Synergistic interactions of compound I and other fungicides in 3DC *Puccinia recondita* (PUCCRT) tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound I+prothioconazole	0.4 + 1.56	100	91	1.10
Compound I+prothioconazole	1.35+1.8	94	39	2.43

Table 4: Synergistic interactions of compound I and other fungicides in 1DP PUCCRT tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound I+prothioconazole	0.15+0.2	78	69	1.14
Compound I+prochloraz	0.15+0.45	93	69	1.35

Table 5: Synergistic interactions of compound II and other fungicides in 1DP SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound II+prochloraz	0.45+5	55	36	1.54

Table 6: Synergistic interactions of compound II and other fungicides 3DC SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound II+prothioconazole	1.35+1.8	95	38	2.49

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound II+prothioconazole	0.45+0.6	38	14	2.65
Compound II+prochloraz	1.35+4.05	95	53	1.81
Compound II+prochloraz	1.5+3	75	63	1.20

Table 7: Synergistic interactions of compound III and other fungicides in 1DP SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound III+prothioconazole	1.35+1.8	98	90	1.09
Compound III+prochloraz	1.35+4.05	98	91	1.07
Compound III+prochloraz	0.3+5	62	39	1.60

Table 8: Synergistic interactions of compound III and other fungicides in 3DC SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound III+prothioconazole	0.45+0.6	88	33	2.69
Compound III+prothioconazole	0.15+0.2	50	42	1.20
Compound III+prochloraz	0.45+1.35	71	42	1.68
Compound III+prochloraz	0.15+0.45	67	22	3.08

Table 9: Synergistic interactions of compound IV and other fungicides in 1DP SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound IV+prothioconazole	4.2+5.6	90	78	1.15

Table 10: Synergistic interactions of compound IV and other fungicides in 3DC SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound IV+prothioconazole	4.2+5.6	97	64	1.52
Compound IV+prochloraz	6+3	72	63	1.15

Table 11: Synergistic interactions of compound V and other fungicides in 1DP SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound V+prothioconazole	0.45+0.6	20	14	1.39
Compound V+prochloraz	1.35+4.05	87	38	2.28
Compound V+prochloraz	0.45+1.35	38	5	7.23

Table 12: Synergistic interactions of compound V and other fungicides in 3DC SEPTTR tests

	Rate ppm	%DC obs	%DC exp	Synergism factor
Compound V+prothioconazole	1.35+1.8	78	52	1.51
Compound V+prochloraz	2+3	56	38	1.45
Compound V+prochloraz	0.15+0.45	50	40	1.26

For all tables, %DC = % Disease Control

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.


### Non-patent literature cited in the description

- Synergistic Fungicidal Compositions of Heterocyclic Aromatic Amides and Triazoles IP.COM, 2004, [0004]
- The Pesticide Manual 2009 0000 [0007]


- The Pesticide Manual 20060000 [0008]
- **COLBY, S. R.** Calculation of the synergistic and antagonistic response of herbicide combinations Weeds, 1967, vol. 15, 20-22 [0038]

## PATENTKRAV


1. Synergistisk fungicid blanding indeholdende en fungicid virksam mængde af en forbindelse med formlen I-V og mindst ét fungicid valgt fra gruppen bestående af prothioconazol og prochloraz.




I




II



III



IV



V

10

2. Blanding ifølge krav 1, i hvilken fungicidet er prothioconazol.
3. Blanding ifølge krav 2, i hvilken fungicidet er forbindelse I.
4. Blanding ifølge krav 1, i hvilken vægtforholdet mellem forbindelse I-V og prothioconazol ligger mellem 1:10 og 10:1.
5. Blanding ifølge krav 1, i hvilken vægtforholdet mellem forbindelse I-V og prochloraz ligger mellem 1:10 og 10:1.

6. Fungicid sammensætning indeholdende en fungicid virksam mængde af den fungicide blanding ifølge krav 1 og et landbrugsmæssigt acceptabelt adjuvans eller bærestof.

25

7.      Anvendelse af blandingen eller sammensætningen ifølge ethvert af de foregående krav til bekæmpelse af svampe, der inficerer planteafgrøder.
8.      Anvendelse ifølge krav 7, ved hvilken svampene tilhører klasserne
- 5   *Ascomycetes* og *Basidiomycetes*.
9.      Anvendelse ifølge krav 8, ved hvilken svampene er valgt fra gruppen bestående af hvedebrunrust (*Puccinia recondita*; PUCCRT); hvedegulrust (*Puccinia striiformis*; PUC CST); hvedegråplet (*Mycosphaerella graminicola*;
- 10   anamorf: *Septoria tritici*; SEPTTR); hvedebrunplet (*Leptosphaeria nodorum*;
- LEPTNO; anamorf: *Stagonospora nodorum*); bygpletsyge (*Cochliobolus sativum*;
- COCHSA; anamorf: *Helminthosporium sativum*); bladpletsyge på sukkerroer (*Cercospora beticola*; CERCBE); jordnøddebladpletsyge (*Mycosphaerella arachidis*; MYCOAR; anamorf: *Cercospora arachidicola*); agurkeanthracnose
- 15   (*Glomerella lagenarium*; anamorf: *Colletotrichum lagenarium*; COLLLA) og black sigatoka-syge hos bananer (*Mycosphaerella fijiensis*; MYCOFI).
10.     Anvendelse ifølge krav 7, ved hvilken sammensætningen tilføres i en mængde på mellem 65 gram pr. hektar (g/ha) og 2300 g/ha, baseret på den totale
- 20   mængde aktive bestanddele i sammensætningen.
11.     Anvendelse ifølge krav 7, ved hvilken prothioconazol tilføres i en mængde på mellem 50 g/ha og 200 g/ha, og forbindelsen med formlen I - V tilføres i en mængde på mellem 35 g/ha og 300 g/ha.
- 25
12.     Anvendelse ifølge krav 7, ved hvilken prochloraz tilføres i en mængde på mellem 50 g/ha and 450 g/ha, og forbindelsen med formlen I - V tilføres i en mængde på mellem 35 g/ha og 300 g/ha.