



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : A61K 39/09, 39/40, C07K 16/12, 14/315</p>	<p>A2</p>	<p>(11) International Publication Number: WO 99/51266</p> <p>(43) International Publication Date: 14 October 1999 (14.10.99)</p>					
<p>(21) International Application Number: PCT/US99/07680</p> <p>(22) International Filing Date: 6 April 1999 (06.04.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/080,878</td> <td>7 April 1998 (07.04.98)</td> <td>US</td> </tr> <tr> <td>60/085,743</td> <td>15 May 1998 (15.05.98)</td> <td>US</td> </tr> </table> <p>(71) Applicant: MEDIMMUNE, INC. [US/US]; 35 West Watkins Mill Road, Gaithersburg, MD 20878 (US).</p> <p>(72) Inventors: WIZEMANN, Theresa, M.; 9 Peach Leaf Court, N. Potomac, MD 20878 (US). KOENIG, Scott; 10901 Ralston Road, Rockville, MD 20852 (US). JOHNSON, Leslie, S.; 13545 Ambassador Drive, Germantown, MD 20874 (US).</p> <p>(74) Agents: OLSTEIN, Elliot, M. et al.; Carella, Byrne, Bain, Gilfillan, Cecchi, Stewart & Olstein, 6 Becker Farm Road, Roseland, NJ 07068 (US).</p>	60/080,878	7 April 1998 (07.04.98)	US	60/085,743	15 May 1998 (15.05.98)	US	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
60/080,878	7 April 1998 (07.04.98)	US					
60/085,743	15 May 1998 (15.05.98)	US					
<p>(54) Title: DERIVATIVES OF PNEUMOCOCCAL CHOLINE BINDING PROTEINS FOR VACCINES</p>							
<p>(57) Abstract</p> <p>The present invention provides bacterial immunogenic agents for administration to humans and non-human animals to stimulate an immune response. It particularly relates to the vaccination of mammalian species with pneumococcal derived polypeptides that include an alpha helix but exclude a choline binding region as a mechanism for stimulating production of antibodies that protect the vaccine recipient against infection by pathogenic bacterial species. In another aspect the invention provides antibodies against such proteins and protein complexes that may be used as diagnostics and/or as protective/treatment agents for pathogenic bacterial species.</p>							

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DERIVATIVES OF PNEUMOCOCCAL CHOLINE BINDING PROTEINS FOR VACCINES

5

This application claims the benefit of U.S. Prov. Appl'n Serial No. 60/085,743, filed May 15, 1998 and U.S. Prov. Appl'n Serial NO 60/080,878, filed April 7, 1998.

10

This invention relates generally to the field of bacterial antigens and their use, for example, as immunogenic agents in humans and animals to stimulate an immune response. More specifically, it relates to the vaccination of mammalian species with a polypeptide comprising an alpha helix-forming polypeptide obtained from a choline binding polypeptide as a mechanism for stimulating production of antibodies that protect the vaccine recipient against infection by pathogenic bacterial species. Further, the invention relates to antibodies and antagonists against such polypeptides useful in diagnosis and passive immune therapy with respect to diagnosing and treating such pneumococcal infections.

25

In a particular aspect, the present invention relates to the prevention and treatment of pneumococcal infections such as infections of the middle ear, nasopharynx, lung and bronchial areas, blood, CSF, and the like, that are caused by pneumococcal bacteria. In this regard, certain types of *Streptococcus pneumoniae* are of particular interest.

S. pneumoniae is a gram positive bacteria which is a major causative agent in invasive infections in animals and humans, such as sepsis, meningitis, otitis media and lobar pneumonia (Tuomanen, et al. NEJM 322:1280-1284 (1995)). As part of the infective process, pneumococci

35

readily bind to non-inflamed human epithelial cells of the upper and lower respiratory tract by binding to eukaryotic carbohydrates in a lectin-like manner (Cundell *et al.*, *Micro. Path.* 17:361-374 (1994)). Conversion to invasive pneumococcal infections for bound bacteria may involve the local generation of inflammatory factors which may activate the epithelial cells to change the number and type of receptors on their surface (Cundell, *et al.*, *Nature*, 377:435-438 (1995)). Apparently, one such receptor, platelet activating factor (PAF) is engaged by the pneumococcal bacteria and within a very short period of time (minutes) from the appearance of PAF, pneumococci exhibit strongly enhanced adherence and invasion of tissue. Certain soluble receptor analogs have been shown to prevent the progression of pneumococcal infections (Idanpaan-Heikkila *et al.*, *J. Inf. Dis.*, 176:704-712 (1997)).

A family of choline binding proteins (CBPs), which are non-covalently bound to phosphorylcholine, are present on the surface of pneumococci and have a non-covalent association with teichoic acid or lipoteichoic acid. An example of such family is choline binding protein A (CbpA), an approximately 75kD weight type of CBP which includes a unique N-terminal domain, a proline rich region, and a C-terminal domain comprised of multiple 20 amino acid repeats responsible for binding to choline. A segment of the N-terminal portion of CbpA protein forms an alpha helix as part of its three-dimensional structure.

30

Accordingly, it is an object of the present invention to provide a polypeptide having broad protection against pneumococcal infections.

35

Definitions

In order to facilitate understanding of the description below and the examples which follow certain

frequently occurring methods and/or terms will be described.

"Plasmids" are designated by a lower case p preceded
5 and/or followed by capital letters and/or numbers. The
starting plasmids herein are either commercially
available, publicly available on an unrestricted basis, or
can be constructed from available plasmids in accord with
published procedures. In addition, equivalent plasmids to
10 those described are known in the art and will be apparent
to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of
the DNA with a restriction enzyme that acts only at
15 certain sequences in the DNA. The various restriction
enzymes used herein are commercially available and their
reaction conditions, cofactors and other requirements were
used as would be known to the ordinarily skilled artisan.

For analytical purposes, typically 1 μg of plasmid or DNA
20 fragment is used with about 2 units of enzyme in about 20
 μl of buffer solution. For the purpose of isolating DNA
fragments for plasmid construction, typically 5 to 50 μg
of DNA are digested with 20 to 250 units of enzyme in a
larger volume. Appropriate buffers and substrate amounts
25 for particular restriction enzymes are specified by the
manufacturer. Incubation times of about 1 hour at 37°C are
ordinarily used, but may vary in accordance with the
supplier's instructions. After digestion the reaction is
electrophoresed directly on a polyacrylamide gel to
30 isolate the desired fragment.

Size separation of the cleaved fragments is
performed using 8 percent polyacrylamide gel described by
Goeddel, D. et al., *Nucleic Acids Res.*, 8:4057 (1980).

35

"Oligonucleotides" refers to either a single
stranded polydeoxynucleotide or two complementary
polydeoxynucleotide strands which may be chemically

synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide
5 will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double stranded nucleic
10 acid fragments (Maniatis, T., et al., Id., p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units to T4 DNA ligase ("ligase") per 0.5 μ g of approximately equimolar amounts of the DNA fragments to be ligated.

15

"HPS portion" as used herein refers to an amino acid sequence as set forth in SEQ ID NO:2 for a choline binding protein ("CBP") of a pneumococcal bacteria that may be located amino terminal with respect to the proline rich
20 portion of the overall amino acid sequence for such CBP.

The terms "identity", "% identity" or "percent identity" as utilized in this application refer to a calculation of differences between two contiguous
25 sequences which have been aligned for "best fit" (to provide the largest number of aligned identical corresponding sequence elements, wherein elements are either nucleotides or amino acids) and all individual differences are considered as individual difference with
30 respect to the identity. In this respect, all individual element gaps (caused by insertions and deletions with respect to an initial sequence ("reference sequence")) over the length of the reference sequence and individual substitutions of different elements (for individual
35 elements of the reference sequence) are considered as individual differences in calculating the total number of differences between two sequences. Individual differences may be compared between two sequences where an initial

sequences (reference sequence) has been varied to obtain a variant sequence (comparative sequence) or where a new sequence (comparative sequence) is simply aligned and compared to such a reference sequence. When two aligned sequences are compared all of the individual gaps in BOTH sequences that are caused by the "best fit" alignment over the length of the reference sequence are considered individual differences for the purposes of identity. If an alignment exists which satisfies the stated minimum identity, then a sequence has the stated minimum identity to the reference sequence. For example, the following is a hypothetical comparison of two sequences having 100 elements each that are aligned for best fit wherein one sequence is regarded as the "reference sequence" and the other as the comparative sequence. All of the individual alignment gaps in both sequences are counted over the length of the reference sequence and added to the number of individual element substitution changes (aligned elements that are different) of the comparative sequence for the total number of element differences. The total number of differences (for example 7 gaps and 3 substitutions) is divided by the total number of elements in the length of the reference sequence (100 elements) for the "percentage difference" (10/100). The resulting percentage difference (10%) is subtracted from 100% identity to provide a "% identity" of 90% identity. For the identity calculation all individual differences in both sequences are considered in the above manner over a discrete comparison length (the length of the reference sequence) of two best fit aligned sequences to determine identity. Thus, no algorithm is necessary for such an identity calculation.

"Isolated" in the context of the present invention with respect to polypeptides and/or polynucleotides means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living organism

is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the co-existing materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment. The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

Summary of the Invention

In one aspect the present invention relates to a vaccine for treating or preventing pneumococcal bacterial infections which utilizes as an immunogen at least one polypeptide truncate of a pneumococcal surface-binding protein, analog, or variant having a highly conserved immunogenic alpha-helical portion (corresponding generally to a "consensus" amino acid sequence as set forth in SEQ ID NO:1) with respect to different types of pneumococcal bacteria, which polypeptide does not include a choline-binding portion. Preferably, the C-terminal choline-binding portion is absent from such polypeptides. More preferred are such polypeptides wherein the HPS amino acid sequence is also absent. Even further preferred are polypeptides wherein the highly conserved immunogenic alpha-helical portion corresponding generally to a "consensus" amino acid sequence as set forth in SEQ ID NO:1 also corresponds generally to the amino acid sequence as set forth in SEQ ID NO:19 (amino acids 1 to 103 of SEQ ID NO:19 are identical to amino acids 1 to 103 of SEQ ID NO:1). Also preferred as vaccines are recombinantly-produced, isolated polypeptides that are missing both an HPS portion and the choline-binding portion.

35

More preferred as vaccines are one or more polypeptide truncates of pneumococcal surface-binding proteins, analogs or variants including a single highly conserved alpha-helix immunogenic portion with respect to

different types of pneumococci, which polypeptides do not include a C-terminal choline-binding portion. Further preferred are isolated recombinantly produced polypeptides having such structure. Also preferred are such polypeptides that do not include either a C-terminal choline-binding portion or a HPS portion.

The present invention further provides a vaccine comprising a polypeptide including an immunogenic portion that is capable of forming an alpha helix, which polypeptide includes a sequence that has at least 85% identity and preferably at least 87% identity to the amino acid sequence of SEQ ID NO:1, wherein the isolated polypeptide does not include a C-terminal choline-binding portion. Further preferred are such polypeptides that comprise a polypeptide sequence that has at least 85% identity and preferably at least 87% identity to an amino acid sequence according SEQ ID NO:19. Preferably, the sequence of the isolated polypeptide includes neither an HPS portion (SEQ ID NO:2) nor a C-terminal choline-binding portion. Further preferred are isolated recombinantly produced polypeptides having such structure. In particular, such polypeptides corresponding to alpha helical structures of different types of *S. pneumoniae* bacteria are contemplated. Particularly preferred are the serotypes 1-5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F and 33F of such *S. pneumoniae* bacteria. Examples of such serotypes of bacteria are readily available from standard ATCC catalogs.

In an additional aspect, the present invention further provides a vaccine against *S. pneumoniae* comprising a synthetic or recombinant polypeptide comprising a plurality of alpha-helical portions, each derived from different naturally occurring *S. pneumoniae* choline-binding polypeptides wherein such alpha-helical portions have at least 85% identity to the amino acid sequence of SEQ ID NO:1, and wherein the isolated

polypeptide does not include a choline-binding portion. Further preferred are those wherein the amino acid sequence for the alpha-helix areas is at least 85% identical to the amino acid sequence of SEQ ID NO:19.

5 Preferably, such synthetic polypeptide includes neither a HPS portion nor a choline-binding portion. Analogs and variants of such chain structure polypeptides wherein such alpha helical portions may be synthetic variant amino acid sequences (or may be a mixture of naturally occurring and

10 variant sequences) are also contemplated and embraced by the present invention. In a preferred aspect, chain vaccines polypeptides having at least ten different alpha helical structures corresponding to *S. pneumoniae* serotypes 1-5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14,

15 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F and 33F are provided. Further preferred are polypeptides including at least fifteen of such alpha-helical structures, more preferred are polypeptides including at least 20 such alpha-helical structures and more preferred are

20 polypeptides including at least one alpha-helical structure corresponding to each of the *S. pneumoniae* serotypes 1-5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F and 33F. Another preferred polypeptide comprises each of the alpha helical

25 structures from the amino acid sequences of SEQ ID NOS:3-18 which correspond to SEQ ID NO:1.

In another aspect, the invention relates to passive immunity vaccines formulated from antibodies against a

30 polypeptide including a highly conserved immunogenic portion with respect to different types of pneumococcal bacteria which portion is capable of forming an alpha-helix having the hereinbefore described identity to the amino acid sequence of SEQ ID NO:1, which polypeptide does

35 not include a C-terminal choline-binding portion, wherein said antibodies will bind to at least one *S. pneumoniae* species. Preferably, if such polypeptide is a truncate of a native pneumococcal surface-binding protein both its HPS portion (where applicable) and its choline-binding portion

are absent from such polypeptide. Such passive immunity vaccines can be utilized to prevent and/or treat pneumococcal infections in immunocompromised patients, patients having an immature immune system (such as young children) or patients who already have an ongoing infection. In this manner, according to a further aspect of the invention, a vaccine can be produced from a synthetic or recombinant polypeptide wherein the polypeptide includes the conserved alpha helical portions of two or more different choline binding polypeptides of *S. pneumoniae*.

This invention also relates generally to the use of an isolated polypeptide having a highly conserved immunogenic portion with respect to different types of pneumococcal bacteria which portion is capable of forming an alpha-helix (corresponding generally to SEQ ID NO:1 or to SEQ ID NO:19) wherein the isolated polypeptide does not include a choline-binding portion, to raise antibodies in non-human mammalian species useful, for example, as diagnostic reagents and vaccines.

In yet another aspect, the present invention relates to the production of a polypeptide including a highly conserved immunogenic portion with respect to different types of pneumococcal bacteria which portion is capable of forming an alpha-helix whose sequence corresponds generally to the amino acid sequence of SEQ ID NO:1 or SEQ ID NO:19, wherein the isolated polypeptide does not include a choline-binding portion. Preferably, such recombinant production is of a truncated native pneumococcal surface-binding polypeptide wherein both the HPS portion (where applicable) and the choline-binding portion are absent.

35

In still another aspect, the present invention provides an isolated choline-binding polypeptide, wherein the non-choline binding region of such polypeptide has at least 90% identity to the corresponding amino acid

sequence portion of a naturally occurring pneumococcal surface-binding protein which is a member selected from the group consisting of SEQ ID NOS:3-18. The invention relates to fragments of such polypeptides which include at least the conserved alpha-helical portion corresponding generally to SEQ ID NO:1, and which has at least 85% identity thereto, wherein the isolated polypeptide preferably is free of a choline binding region.

10 In another aspect the present invention provides an isolated polypeptide comprising an amino acid sequence which has at least 90% identity to one of the amino acid sequences selected from the group consisting of SEQ ID NO:3-18. Preferably, such isolated polypeptide comprises
15 an amino acid sequence which has at least 95% identity, and more preferably 97% identity, to one of the amino acid sequences selected from the group consisting of SEQ ID NO:3-18. The invention further relates to fragments of such polypeptides.

20 In a yet further aspect, the present invention provides a *S. pneumoniae* CBP polypeptide encoded by a polynucleotide that will hybridize under highly stringent conditions to the complement of a polynucleotide encoding
25 a polypeptide having an amino acid selected from the group consisting of SEQ ID NOS:1 and 3-18. Particularly preferred are polypeptides comprising an amino acid sequence segment that is at least 90% identical to the amino acid sequence of SEQ ID NO:1. Further preferred are
30 such polypeptides comprising a contiguous amino acid sequence that has at least 95% identity with respect to the amino acid sequence of SEQ ID NO:1. And, even more preferred are polypeptides comprising an amino acid sequence that has at least 97% identity with respect to
35 the amino acid sequence of SEQ ID NO:1.

In another aspect the present invention provides polynucleotides which encode the hereinabove described polypeptides of the invention. The polynucleotide of the

present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The polynucleotides which encode polypeptides including the amino acid sequences of at least one of SEQ ID NOS:3-18 (or polypeptides that have at least 90% identity to the amino acid sequences of such polypeptides) may be one of the coding sequences shown in SEQ ID NOS:20-35 or may be of a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same polypeptides as the DNA of SEQ ID NOS:20-35.

15

The polynucleotides which encode the polypeptides of SEQ ID NOS:3-18 may include: only the coding sequence for the polypeptide; the coding sequence for the polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the polypeptide. The polypeptides encoded may comprise just a single alpha-helical portion or multiple alpha-helical portion and may independently or collectively include N-terminal sequences 5' of such alpha helical areas and/or sequences corresponding to the "X" structures or proline rich areas (as set forth in Figure 1, for example).

The invention further relates to a polynucleotide comprising a polynucleotide sequence that has at least 95% identity and preferably at least 97% identity to a polynucleotide encoding one of the polypeptides comprising SEQ ID NO:3-18. The invention further relates to fragments of such polynucleotides which include at least the portion of the polynucleotide encoding the polypeptide sequence corresponding to SEQ ID NO:1.

Thus, the term "polynucleotide encoding a polypeptide" encompasses a polynucleotide which includes

only coding sequence for the polypeptide as well as a polynucleotide which includes additional coding and/or non-coding sequence. In particular, the polypeptides may include any or all of the types of structures set forth schematically in Figure 1.

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptides including the amino acid sequences of SEQ ID NOS:3-18. The variants of the polynucleotides may be a naturally occurring allelic variant of the polynucleotides or a non-naturally occurring variant of the polynucleotides. Complements to such coding polynucleotides may be utilized to isolate polynucleotides encoding the same or similar polypeptides. In particular, such procedures are useful to obtain alpha helical coding segments from different serotypes of *S. pneumoniae*, which is especially useful in the production of "chain" polypeptide vaccines containing multiple alpha helical segments.

Thus, the present invention includes polynucleotides encoding polypeptides including the same polypeptides as shown in the Sequence Listing as SEQ ID NOS:3-18 as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptides of SEQ ID NOS:3-18. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

As hereinabove indicated, the polynucleotides may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in the Sequence Listing as SEQ ID NOS:20-35. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

The polynucleotides of the present invention may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptides of the present invention. The marker sequence may be, for example, a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptides fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g. COS-7 cells, is used.

10 The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

The present invention further relates to polynucleotides (hybridization target sequences) which hybridize to the complements of the hereinabove-described sequences if there is at least 70% and preferably 80% identity between the target sequence and the complement of the sequence to which the target sequence hybridizes, preferably at least 85% identity. More preferred are such sequences having at least 90% identity, preferably at least 95% and more preferably at least 97% identity between the target sequence and the sequence of complement of the polynucleotide to which it hybridizes. The invention further relates to the complements to both the target sequence and to the polynucleotide sequence that encodes an amino acid sequence selected from the group consisting of SEQ ID NOS:3 to 18. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the complements of the hereinabove-described polynucleotides as well as to those complements. As herein used, the term "stringent conditions" means hybridization will occur with the complement of a polynucleotide and a corresponding sequence only if there is at least 95% and preferably at least 97% identity between the target sequence and the sequence of complement of the polynucleotide to which it hybridizes. The polynucleotides which hybridize to the complements of the hereinabove described polynucleotides

in a preferred embodiment encode polypeptides which retain an immunogenic portion that will cross-react with an antibody to at least one of the polypeptides having a sequence according to SEQ ID NOS:3-18, or to a polypeptide
5 that includes an amino acid sequence which has at least 85% identity to that of SEQ ID NO:1.

In a still further aspect, the present invention provides for the production of such polypeptides and
10 vaccines as set forth above having a histidine label (or other suitable label) such that the full-length proteins, truncates, analogs or variant discussed above can be isolated due to their label.

15 In another aspect the present invention relates to a method of prophylaxis and/or treatment of diseases that are mediated by pneumococcal bacteria that have surface-binding CBP proteins. In particular, the invention relates to a method for the prophylaxis and/or treatment
20 of infectious diseases that are mediated by *S. pneumoniae* that have a CBP surface-binding protein that forms an alpha helix (comprising a sequence that has at least an 85% identity to the amino acid sequence of SEQ ID NO:1). In a still further preferred aspect, the invention relates
25 to a method for the prophylaxis and/or treatment of such infections in humans.

In still another aspect the present invention relates to a method of using one or more antibodies
30 (monoclonal, polyclonal or sera) to the polypeptides of the invention as described above for the prophylaxis and/or treatment of diseases that are mediated by pneumococcal bacteria that have CBP surface-binding proteins. In particular, the invention relates to a
35 method for the prophylaxis and/or treatment of infectious diseases that are mediated by *S. pneumoniae* CBP proteins which include an alpha helical portion having the hereinbefore described identity to the consensus sequence of SEQ ID NO:1. In a still further preferred aspect, the

invention relates to a method for the prophylaxis and/or treatment of otitis media, nasopharyngeal, bronchial infections, and the like in humans by utilizing antibodies to the alpha-helix containing immunogenic polypeptides of
5 the invention as described above.

Brief Description of Drawings

Figure 1 is a diagram of a pneumococcal CBP protein which shows from the N-terminal to the C-terminal, respectively, (a) a N-terminal sequence, (b) one of a potential alpha-helical forming area conserved segment (R1) that may not be present in some CBP polypeptides, (c) an optional small bridging sequence of amino acids that may bridge two conserved alpha-helical segments (X), (d) a second of a potential alpha-helical forming area consensus sequence (R2) related to the first consensus sequence (which corresponds to SEQ ID NO:1), (e) a proline rich area sequence, (f) a choline binding repeats area, and (e) a C-terminal tail sequence. Where relevant, an optional HPS sequence may naturally occur 5' of the proline rich sequence and 3' of the R1, X, and/or R2 areas.

Figure 2 reports the results for passive immunity protection against 1600 cfu virulent serotype 6B *S. pneumoniae* SP317 (in mice) that was provided by day 31 rabbit antisera to a pneumococcal CBP truncate polypeptide, NR1XR2 (truncate missing both the proline and the choline binding areas, but including two conserved alpha-helical areas R1 and R2). Eighty percent of the mice immunized with the truncate antisera prior to challenge survived the 14 day observation period. By contrast, all mice immunized with a control sera (pre-immune rabbit sera) were dead by day 7.

Figure 3 reports the results for passive immunity protection against 3450 cfu virulent serotype 6B *S. pneumoniae* SP317 (in mice) that was provided by day 52 rabbit antisera to a pneumococcal CBP truncate polypeptide, NR1XR2 (truncate missing both the proline and the choline binding areas, but including two conserved alpha-helical areas R1 and R2). One hundred percent of the mice immunized with the truncate antisera prior to challenge survived the 10 day observation period. By

contrast, ninety percent of the mice immunized with a control sera (pre-immune rabbit sera) were dead at day 10.

Figure 4 reports the results for passive immunity protection against 580 cfu virulent serotype 6B *S. pneumoniae* SPSJ2 (in mice) that was provided by day 31 rabbit antisera to a pneumococcal CBP truncate polypeptide, NR1XR2 (truncate missing both the proline and the choline binding areas, but including two conserved alpha-helical areas R1 and R2). Fifty percent of the mice immunized with the truncate antisera prior to challenge survived the 10 day observation period. By contrast, all mice immunized with a control sera (pre-immune rabbit sera) were dead by day 8.

15

Figure 5 reports the results for active immunity protection against 560 cfu virulent serotype 6B *S. pneumoniae* SPSJ2 (in mice) that was provided by immunization with a pneumococcal CBP truncate polypeptide, NR1X (truncate missing the second conserved alpha-helical area R2, as well as both the proline and the choline binding areas). Eighty percent of the mice actively immunized with the NR1X CBP truncate prior to challenge survived the 14 day observation period. By contrast, all mice immunized with a control (sham mice) of PBS and adjuvant were dead by day 8.

25

Figure 6 reports the results for active immunity protection against 680 cfu virulent serotype 6B *S. pneumoniae* SPSJ2 (in mice) that was provided by immunization with a pneumococcal CBP truncate polypeptide, NR1XR2 (truncate missing both the proline and the choline binding areas, but including two conserved alpha-helical areas R1 and R2). Fifty percent of the mice actively immunized with the NR1XR2 CBP truncate prior to challenge survived the 14 day observation period. By contrast, all mice immunized with a control (SP90) protein and adjuvant were dead by day 9.

30

35

Figure 7 is an alignment report of the amino terminus of CBP polypeptides from various types of *S. pneumoniae* and a consensus sequence is reported at the top of each row (sets of lines) of the comparison. The consensus sequence for the comparison is listed as the "Majority" sequence (SEQ ID NO:36). One letter codes are utilized to represent the sequences which are aligned for a "best fit" comparison wherein dashes in a sequence indicate spacing gaps of the contiguous sequence.

10

Figure 8 shows the sequence pair distances for the amino acid sequences as described for Figure 7 and set forth therein. A Clustal method with identity residue weight table is used. The percent similarity for such a comparison is reported for the amino acid sequences set forth in Figure 7.

15

Figure 9 is an alignment report for a first helical region in the amino acid sequences of CBP polypeptides from various types of *S. pneumoniae* and a consensus sequence is reported at the top of each row (sets of lines) of the comparison. The consensus sequence for the comparison is listed as the "Majority" sequence (SEQ ID NO:38). One letter codes are utilized to represent the sequences which are aligned for a "best fit" comparison wherein dashes in a sequence indicate spacing gaps of the contiguous sequence.

20

25

Figure 10 shows the sequence pair distances for the amino acid sequences as described for Figure 9 and set forth therein. A Clustal method with identity residue weight table is used. The percent similarity for such a comparison is reported for the amino acid sequences set forth in Figure 9.

30

35

Figure 11 is an alignment report for the region X in the amino acid sequences of CBP polypeptides from various types of *S. pneumoniae* and a consensus sequence is reported at the top of each row (sets of lines) of the

comparison. The consensus sequence for the comparison is listed as the "Majority" sequence (SEQ ID NO:37). One letter codes are utilized to represent the sequences which are aligned for a "best fit" comparison wherein dashes in a sequence indicate spacing gaps of the contiguous sequence.

Figure 12 shows the sequence pair distances for the amino acid sequences as described for Figure 11 and set forth therein. A Clustal method with identity residue weight table is used. The percent similarity for such a comparison is reported for the amino acid sequences set forth in Figure 11.

Figure 13 is an alignment report for the second helical region A in the amino acid sequences of CBP polypeptides from various types of *S. pneumoniae* and a consensus sequence is reported at the top of each row (sets of lines) of the comparison. The consensus sequence for the comparison is listed as the "Majority" sequence (SEQ ID NO:1). One letter codes are utilized to represent the sequences which are aligned for a "best fit" comparison wherein dashes in a sequence indicate spacing gaps of the contiguous sequence.

Figure 14 shows the sequence pair distances for the amino acid sequences as described for Figure 13 and set forth therein. A Clustal method with identity residue weight table is used. The percent similarity for such a comparison is reported for the amino acid sequences set forth in Figure 13.

Figure 15 is an alignment report for the second helical region B in the amino acid sequences of CBP polypeptides from various types of *S. pneumoniae* and a consensus sequence is reported at the top of each row (sets of lines) of the comparison. The consensus sequence for the comparison is listed as the "Majority" sequence (SEQ ID NO:19). One letter codes are utilized to

represent the sequences which are aligned for a "best fit" comparison wherein dashes in a sequence indicate spacing gaps of the contiguous sequence.

5 Figure 16 shows the sequence pair distances for the amino acid sequences as described for Figure 15 and set forth therein. A Clustal method with identity residue weight table is used. The percent similarity for such a comparison is reported for the amino acid sequences set
10 forth in Figure 15.

Detailed Description of the Invention

15 In accordance with an aspect of the present invention there is provided a vaccine to produce a protective response against *S. pneumoniae* infections which employs a polypeptide which comprises a member selected from the group consisting of:

20 (a) an amino acid sequence which produces an alpha helical structure and which is at least 85% identical to the amino acid sequence of SEQ ID NO:1 and which is free of a choline binding region, and

25 (b) an isolated truncate of a naturally occurring *S. pneumoniae* polypeptide that comprises an alpha helical portion that has at least 85% identity to the amino acid sequence of SEQ ID NO:1 and is free of a choline binding region,

30 (c) an isolated truncate of a naturally occurring *S. pneumoniae* polypeptide that comprises an alpha helical portion that has at least 90% identity to the amino acid sequence of SEQ ID NO:19 and is free of a choline binding region. In a preferred aspect, such isolated truncate polypeptide is a member selected from the group consisting
35 of SEQ ID NOS:3-18 and said isolated polypeptide is free of a choline binding region and, if relevant, a HPS region; or a fragment thereof which includes at least the alpha helical segment which corresponds to the consensus sequence of SEQ ID NO:1. Particularly preferred are

vaccines which utilize such truncate polypeptides that include at least such alpha helical area or utilize a recombinant immunogen polypeptide comprising at least two of such alpha-helical segments. Such polypeptide may be
5 a recombinant polypeptide containing multiple alpha-helical areas from one or more truncates. Further preferred are recombinant immunogen polypeptides comprising at least two alpha-helical areas corresponding to the alpha helical areas of two or more truncates from
10 different types of pneumococcal bacteria. Such polypeptide may be a recombinant polypeptide containing multiple alpha-helical areas from one or more different types of pneumococcal bacteria.

15 In accordance with the present invention, there is provided an isolated polypeptide comprising a truncated surface-binding polypeptide derived from *S. pneumoniae*, said isolated polypeptide containing an alpha-helical area whose amino acid sequence corresponds generally to the
20 amino acid sequence of SEQ ID NO:1, but free of a choline binding area. Preferably, said isolated polypeptide also omits any naturally occurring repeats of the alpha-helical forming area and omits any HPS amino acid sequence that may be present.

25 It is an object of the present invention to utilize as immunogenic composition for a vaccine (or to produce antibodies for use as a diagnostic or as a passive vaccine) comprising an immunogenic polypeptide comprising
30 a pneumococcal surface-binding polypeptide with an alpha helical portion from which a choline binding region has been omitted. In one embodiment, such truncated proteins (naturally or recombinantly produced, as well as functional analogs) from *S. pneumoniae* bacteria are contemplated.
35 Even more particularly, *S. pneumoniae* polypeptides having a single alpha helical portion that omit any HPS areas that occur and choline binding areas of the native protein are contemplated.

A particularly preferred embodiment of such an immunogenic composition is for use as a vaccine (or as an immunogen for producing antibodies useful for diagnostics or vaccines) wherein the active component of the immunogenic composition is an isolated polypeptide comprising at least one member selected from the group consisting of:

(a) an amino acid sequence which is selected from SEQ ID NOS:3-19,

(b) a polypeptide which has at least 90% identity to (a), preferably at least 95% identity to (a), and even more preferred at least 97% identity to (a), or

(c) a fragment of (a) or (b) wherein such fragment includes at least one alpha helical portion that corresponds to the consensus sequence which is SEQ ID NO:1 and said fragment does not comprise a choline binding region. Preferably, such vaccines utilize a polypeptide that contains neither a choline binding region nor an HPS region that occurs as part of the amino acid sequences in the native proteins.

In another preferred embodiment, there is provided a vaccine which includes at least one isolated polypeptide which includes an amino acid sequence which has at least 85% identity (preferably 87% identity and more preferably at least 90% identity) to SEQ ID NO:1, which isolated polypeptide is free of a choline binding portion and, where applicable, is also preferably free of an HPS portion. The preferred polypeptide may also include one or more of the N-terminal sequences that are located 5' of the alpha helical areas in the polypeptides having an amino acid sequence selected from the group consisting of SEQ ID NOS:3-18, or the like. The polypeptide truncate may also include one or more of the proline regions (region "P" in Figure 1) and/or the spanning region (region "X" in Figure 1).

In another aspect of the invention, such an immunogenic composition may be utilized to produce

antibodies to diagnose pneumococcal infections, or to produce vaccines for prophylaxis and/or treatment of such pneumococcal infections as well as booster vaccines to maintain a high titer of antibodies against the immunogen(s) of the immunogenic composition.

While other antigens have been contemplated to produce antibodies for diagnosis and for the prophylaxis and/or treatment of pneumococcal infections, there is a need for improved or more efficient vaccines. Such vaccines should have an improved or enhanced effect in preventing bacterial infections mediated pneumococci having surface-binding polypeptides. Further, to avoid unnecessary expense and provide broad protection against a range of pneumococcal serotypes there is a need for polypeptides that comprise an immunogenic amino acid sequence corresponding to a portion of pneumococcal surface-binding polypeptides that is a highly conserved portion among various types of pneumococci. Preferably, such polypeptides avoid the inclusion of amino acid sequences corresponding to other portions of the native polypeptides, such as the choline binding region and/or the HPS region.

There is a need for improved antigenic compositions comprising highly conserved portions of polypeptides that bind to the surface of pneumococcal bacteria for stimulating high-titer specific antisera to provide protection against infection by pathogenic pneumococcal bacteria and also for use as diagnostic reagents.

In such respect, truncated polypeptides, functional variant analogs, and recombinantly produced truncated polypeptides of the invention are useful as immunogens for preparing vaccine compositions that stimulate the production of antibodies that can confer immunity against pathogenic species of bacteria. Further, preparation of vaccines containing purified proteins as antigenic ingredients are well known in the art.

Generally, vaccines are prepared as injectables, in the form of aqueous solutions or suspensions. Vaccines in an oil base are also well known such as for inhaling.

5 Solid forms which are dissolved or suspended prior to use may also be formulated. Pharmaceutical carriers are generally added that are compatible with the active ingredients and acceptable for pharmaceutical use. Examples of such carriers include, but are not limited to,
10 water, saline solutions, dextrose, or glycerol. Combinations of carriers may also be used.

Vaccine compositions may further incorporate additional substances to stabilize pH, or to function as
15 adjuvants, wetting agents, or emulsifying agents, which can serve to improve the effectiveness of the vaccine.

Vaccines are generally formulated for parenteral administration and are injected either subcutaneously or
20 intramuscularly. Such vaccines can also be formulated as suppositories or for oral administration, using methods known in the art.

The amount of vaccine sufficient to confer immunity
25 to pathogenic bacteria is determined by methods well known to those skilled in the art. This quantity will be determined based upon the characteristics of the vaccine recipient and the level of immunity required. Typically, the amount of vaccine to be administered will be
30 determined based upon the judgment of a skilled physician.

Where vaccines are administered by subcutaneous or intramuscular injection, a range of 50 to 500 μ g purified protein may be given.

35 The term "patient in need thereof" refers to a human that is infected with, or likely, to be infected with, pathogenic pneumococcal bacteria that produce CbpA, or the like, preferably *S. pneumoniae* bacteria (however a mouse

model can be utilized to simulate such a patient in some circumstances).

5 In addition to use as vaccines, the polypeptides of the present invention can be used as immunogens to stimulate the production of antibodies for use in passive immunotherapy, for use as diagnostic reagents, and for use as reagents in other processes such as affinity chromatography.

10

The polynucleotides encoding the immunogenic polypeptides described above may also have the coding sequence fused in frame to a marker sequence which allows for purification of the polypeptides of the present invention. The marker sequence may be, for example, a hexa-histidine tag supplied by a pQE-9 vector to provide for purification of the mature polypeptides fused to the marker in the case of a bacterial host, or, for example, the marker sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g. COS-7 cells, is used. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I., et al., Cell, 37:767 (1984)).

25

The identification of multiple coil structures of alpha helical amino acid segments in the *S. pneumoniae* polypeptides according to the invention may be determined by the location of proline rich areas with respect to one another. Further the "X" area optionally located between two or more alpha-helical structures can play a part in the formation of a coil within a coil structure. Standard three-dimensional protein modeling may be utilized for determining the relative shape of such structures. An example of a computer program, the Paircoil Scoring Form Program ("PairCoil program"), useful for such three-dimensional protein modelling is described by Berger et al. in the Proc. Natl. Acad. of Sci. (USA), 92:8259-8263 (August 1995). The PairCoil program is described as a computer program that utilizes a mathematical algorithm to

30

35

predict locations of coiled-coil regions in amino acid sequences. A further example of such a computer program is described by Wolf et al., Protein Science 6:1179-1189 (June 1997) which is called the Multicoil Scoring Form
5 Program ("Multicoil program"). The MultiCoil program is based on the PairCoil algorithm and is useful for locating dimeric and trimeric coiled coils .

In a preferred aspect, the invention provides for
10 recombinant production of such polypeptides in a host bacterial cell other than a *S. pneumoniae* species host to avoid the inclusion of native surface-binding polypeptides that have a choline binding region. A preferred host is a species of such bacteria that can be cultured under
15 conditions such that the polypeptide of the invention is secreted from the cell.

The present invention also relates to vectors which include polynucleotides encoding one or more of the
20 polypeptides of the invention that include the highly conserved alpha-helical amino acid sequence in the absence of an area encoding a choline binding amino acid sequence, host cells which are genetically engineered with vectors of the invention and the production of such immunogenic
25 polypeptides by recombinant techniques in an isolated and substantially immunogenically pure form.

Host cells are genetically engineered (transduced or transformed or transfected) with the vectors comprising a
30 polynucleotide encoding a polypeptide comprising the highly conserved alpha-helical region but not having a choline binding region, or the like of this invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the
35 form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the polynucleotides which encode such polypeptides. The

culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

5

Vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host.

15 The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the E. coli. lac or trp, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

35 In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampi-

cillin resistance in E. coli.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the proteins.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as E. coli, Streptomyces, Salmonella typhimurium; fungal cells, such as yeast; insect cells such as Drosophila S2 and Spodoptera Sf9; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; plant cells, etc. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example. Bacterial: pQE70, pQE60, pQE-9 (Qiagen, Inc.), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16a, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired

gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P_R, P_L and TRP. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

10

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., Battey, I., Basic Methods in Molecular Biology, (1986)).

20

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

25

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

30

Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

35

Transcription of the DNA encoding the polypeptides

of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples including the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

10

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of E. coli and S. cerevisiae TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), α -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include E. coli, Bacillus subtilis, Salmonella typhimurium and various species within the genera Pseudomonas,

Streptomyces, and Staphylococcus, although others may also be employed as a matter of choice.

5 As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for
10 example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

15

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and
20 cells are cultured for an additional period.

25

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, a french press, mechanical disruption, or use of cell lysing agents, such
30 methods are well know to those skilled in the art. However, preferred are host cells which secrete the polypeptide of the invention and permit recovery of the polypeptide from the culture media.

35

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing

a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

10

The polypeptides can be recovered and/or purified from recombinant cell cultures by well-known protein recovery and purification methods. Such methodology may include ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. In this respect, chaperones may be used in such a refolding procedure. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

25 The polypeptides that are useful as immunogens in the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. Particularly preferred immunogens are truncated pneumococcal polypeptides that comprise a single highly conserved alpha helical area, but do not comprise a choline binding region or a HPS region. Therefore, antibodies against such polypeptides should bind to other pneumococcal bacterial species (in addition to the *S. pneumoniae* species from

30
35

which such polypeptides were derived) and vaccines against such *S. pneumoniae* should give protection against other pneumococcal bacterial infections.

5 Procedures for the isolation of the individually expressed alpha-helical containing polypeptides may be isolated by recombinant expression/isolation methods that are well-known in the art. Typical examples for such isolation may utilize an antibody to a conserved area of
10 the protein or to a His tag or cleavable leader or tail that is expressing as part of the protein structure.

 The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them
15 can be used as an immunogen to produce antibodies thereto.

 These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab
20 expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

 Antibodies generated against the polypeptides
25 corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner,
30 even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides.

 For preparation of monoclonal antibodies, any
35 technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today

4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

5

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic polypeptide products of this invention.

In order to facilitate understanding of the above description and the examples which follow below, as well as the Figures included herewith, Table 1 below sets forth the bacterial source for the polypeptides of SEQ ID NOS:3-18 and the polynucleotides encoding them (SEQ ID NOS:20-35, respectively). The name and/or type of bacteria is specified and a credit or source is named. The sequences from such types of bacteria are for illustrative purposes only since by utilizing probes and/or primers as described herein other sequences of similar type may be readily obtained by utilizing only routine skill in the art.

25

TABLE 1

SEQ ID NO.	Type Of Pneumococcus	Source Credit or ATCC No.
3	1	ATCC 33400
4	2	SPATCC 11733
5	2	ATCC2 (catalog #6302)
6	4	ATCC4 (catalog #6304)
7	6B	ATCC 6B (catalog #6326)
8	18C	SPATCC 18C (ATCC catalog #10356)

TABLE 1
(Continued)

SEQ ID NO.	Type Of Pneumococcus	Source Credit or ATCC No.
9	4	Norway type 4; Nat'l. Inst. of Public Health, Norway Ingeborg Aagerge
10	noncapsulated	R6X; Rockefeller Univ., Rob Masure (from D39, type 2)
11	6B	SPB 105; Boston Univ., Steve Pelton
12	23F	SPB 328; Boston Univ., Steve Pelton
13	14	SPB 331; Boston Univ., Steve Pelton
14	23F	SPB 365; Boston Univ., Steve Pelton
15	9V	SPR 332; Rockefeller Univ., Rob Masure
16	6B	SPSJ 2p; St. Jude Children's Research Hospital, Pat Flynn (clinical isolate passaged 1x in mice for virulence)
17	14	SPSJ 9; St. Jude Children's Research Hospital, Pat Flynn (clinical isolate - nares, pneumonia)
18	19A	SPSJ 12; St. Jude Children's Research Hospital, Pat Flynn (clinical isolate)

5

The present invention will be further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

10

Example 1Generation of CbpA Truncate Protein Vectors**A. Vector for Full Length CbpA (NR1XR2PC)**

5 A virulent serotype 4 *S. pneumoniae* strain, Norway 4
(obtained from I. Aaberge, National Institute of Public
Health, Oslo, Norway) was used as a source of genomic DNA
template for amplifying the polynucleotide encoding full-
length CbpA. Full length CbpA was amplified with PCR
10 primers SJ533 and SJ537 described below.

The degenerate forward primer SJ533 was designed
based on the CbpA N-terminal sequence XENEG provided by
H.R. Masure (St. Jude Children's Research Hospital,
15 Memphis, TN). The SJ533 primer = 5' GGC GGA TCC ATG
GA(A,G) AA(C,T) GA(A,G) GG 3'. It incorporates both
BamHI and NcoI restriction sites and an ATG start codon.

20 The 3' reverse primer SJ537 = 5' GCC GTC GAC TTA GTT
TAC CCA TTC ACC ATT GGC 3'

This primer incorporates a SalI restriction site for
cloning purposes, and the natural stop codon from CbpA,
and is based on type 4 and R6X sequence generated in-
25 house.

PCR product generated from genomic DNA template with
SJ533 and SJ537 was digested with BamHI and SalI, and
cloned into the pQE30 expression vector (Qiagen, Inc.)
30 digested with BamHI, XbaI, and SmaI. The type R6X
template resulted in full-length vector PMI581 and the
type 4 template DNA resulted in full-length vector
PMI580.

B. Vector for CbpA Truncate Protein (NR1XR2)

The naturally occurring PvuII site in the end on the
second amino repeat (nucleic acid 1228 of Type 4
sequence) was exploited to create a truncated version of

CbpA, containing only the amino terminus of the gene. To create the truncate clone, the full-length clone PMI580 (Type 4) or PMI581 (R6X) was digested with PvuII and XbaI, and the amino terminus along with a portion of the expression vector was isolated by size on an agarose gel.

5 pQE30 was digested with XbaI and SmaI, and the band corresponding to the other half of the vector was also size selected on an agarose gel. The two halves were ligated and clones identified by restriction digest, then
10 expressed. In this instance, the stop codon utilized is in the expression vector, so the protein expressed is larger than the predicted size due to additional amino acids at the 5' and 3' end of the cloning site.

15 C. Vector for CbpA Truncate Protein (NR1X)

A similar strategy was used to express only the first amino repeat of CbpA. Here the naturally occurring XmnI site between the two amino repeats (nucleic acid 856 of Type 4 sequence) was utilized. CbpA full-length clone
20 PMI580 was digested with XmnI and AatII. Expression vector pQE30 was digested with AatII and SmaI. Once again, the two sized fragments were ligated, and clones were screened by restriction digest and expressed.

25 **Example 2**

Expression of CbpA Truncate Protein From Expression Vectors

All proteins are expressed from the vectors
30 described in Examples 1A-1C in the Qia expressionist System (Qiagen) using the *E. coli* expression vector pQE30, and the amino terminus His6 tagged proteins are detected by western analysis using both anti-Histidine antibodies and gene specific antibodies.

35

The expressed CBP truncates were purified as follows.

A single colony was selected from plated bacteria

containing the recombinant plasmid and grown overnight at 37° in 6.0 ml LB buffer with 50 ug/ml Kanamycin and 100 ug/ml Ampicillin. This 6.0 ml culture was added to 1L LB with antibiotics at above concentrations. The culture
5 was shaken at 37°C until $A_{600} = \sim 0.400$. 1M IPTG was added to the 1L culture to give a final concentration of 1mM. The culture was then shaken at 37°C for 3-4 h. The 1L culture was spun 15 min. in 250 ml conical tubes at 4000 rpm in a model J-6B centrifuge. The supernatant was
10 discarded and the pellet stored at -20°C until use.

The 1L pellet was resuspended in 25 ml 50 mM NaH_2PO_4 , 10mM Tris, 6M GuCl , 300mM NaCl , pH 8.0 (Buffer A). This mixture was then rotated at room temperature
15 for 30 minutes. The mixture was then subjected to sonication (VibraCell Sonicator, Sonics and Materials, Inc., Danbury, CT) using the microtip, two times, for 30 sec., at 50% Duty Cycle and with the output setting at 7.

The mixture was spun 5 min. at 10K in a JA20 rotor and
20 the supernatant removed and discarded. The supernatant was loaded on a 10 ml Talon (Clonetech, Palo Alto, CA) resin column attached to a GradiFrac System (Pharmacia Biotech, Upsala, Sweden). The column was equilibrated with 100 ml Buffer A and washed with 200 ml of this
25 buffer. A volume based pH gradient using 100% 50 mM NaH_2PO_4 , 8M Urea, 20mM MES, pH 6.0 (Buffer B) as the final target buffer was run over a total volume of 100 ml. Protein eluted at ~ 30% Buffer B. Eluted peaks were collected and pooled.

30

For refolding, dialysis was carried out with a 2L volume of PBS at room temperature for approximately 3 hr. using dialysis tubing with a molecular weight cutoff of 14,000. The sample was then dialyzed overnight in 2L of
35 PBS at 4°C. Additional buffer exchange was accomplished during the concentration of the protein using Centriprep-30 spin columns by adding PBS to the spun retentate and

respinning. The protein concentration was determined using the BCA protein assay and the purity visualized using a Coomassie stained 4-20% SDS-PAGE gel.

5

Example 3

Passive Protection with Anti-CbpA Truncate NR1XR2 Antisera

10 A. Generation of Rabbit Immune Serum

Rabbit immune serum against CbpA truncate was generated at Covance (Denver, PA). Following collection of preimmune serum, a New Zealand white rabbit (#ME101) was immunized with 250 µg CbpA truncate NR1XR2 (containing both alpha helix I and alpha helix II amino acid N-terminal repeats that are prepared from 483:58) in Complete Freund's Adjuvant. The rabbit was given a boost of 125 µg CbpA truncate in Incomplete Freund's Adjuvant on day 21 and bled on days 31 and 52.

20

B. Passive Protection in Mice

C3H/HeJ mice (5 mice/group) were passively immunized intraperitoneally with 100 µl of a 1:2 dilution of rabbit sera in sterile PBS (preimmune or day 31 immune sera). One hour after administration of serum, mice were challenged with 1600 cfu virulent serotype 6B *S. pneumoniae*, strain SP317 (obtained from H.R. Masure). Mice were monitored for 14 days for survival.

30

Eighty percent of the mice immunized with rabbit immune serum raised against CbpA truncate NR1XR2 protein survived the challenge for 14 days (Figure 2). All mice immunized with preimmune rabbit serum were dead by day 7.

35

C. Passive Protection in Mice (Higher Challenge Dose)

C3H/HeJ mice (10 mice/group) were passively immunized intraperitoneally with 100 μ l of a 1:2 dilution of rabbit sera in sterile PBS (preimmune or day 52 immune sera). One hour after administration of serum, mice were
5 challenged with 3450 cfu virulent serotype 6B *S. pneumoniae*, strain SP317. Mice were monitored for 10 days for survival.

One hundred percent of the mice immunized with
10 rabbit immune serum raised against CbpA truncate NR1XR2 protein survived the challenge for ten days (Figure 3). Ninety percent of the mice immunized with preimmune rabbit serum were dead at day 10.

15 D. *Passive Protection in Mice (Against High Virulence)*

C3H/HeJ mice (10 mice/group) were passively immunized intraperitoneally with 100 μ l of a 1:2 dilution of rabbit sera in sterile PBS (preimmune or day 52 immune
20 sera). One hour after administration of serum, mice were challenged with 580 cfu virulent serotype 6B *S. pneumoniae*, strain SPSJ2 (provided by P. Flynn, St. Jude Children's Research Hospital, Memphis, TN). Mice were monitored for 10 days for survival.

25 Fifty percent of the mice immunized with rabbit immune serum raised against CbpA truncate NR1XR2 protein survived the challenge for 10 days (Figure 4). All of the mice immunized with preimmune rabbit serum were dead
30 at day 8.

These data demonstrate that antibodies specific for CbpA are protective against systemic pneumococcal infection. The data further indicate that the choline-
35 binding region is not necessary for protection, as antibody specific for truncated protein NR1XR2, lacking the choline-binding repeats, was sufficient for protection. In addition, serum directed against

recombinant CbpA protein based on a serotype 4 sequence, was still protective against challenge with two different strains of serotype 6B.

5

Example 4**Active Protection with Anti-CbpA Truncates NR1X and NR1XR2**

A. *Active Protection With NR1X Truncate Vaccination*
10 C3H/HeJ mice (10/group) were immunized intraperitoneally with CbpA truncate protein NR1X (15µg in 50 µl PBS, plus 50 µl Complete Freund's Adjuvant). A group of 10 sham immunized mice received PBS and adjuvant. A second immunization was administered four
15 weeks later, 15 µg protein i.p. with Incomplete Freund's Adjuvant (sham group received PBS plus IFA). Blood was drawn (retro-orbital bleed) at weeks 3, 6, and 9 for analysis of immune response. The ELISA end point anti-CbpA truncate titer of pooled sera from the 10 CbpA
20 immunized mice at 9 weeks was 4,096,000. No antibody was detected in sera from sham immunized mice. Mice were challenged at week 10 with 560 CFU serotype 6B *S. pneumoniae* strain SPSJ2. Mice were monitored for 14 days for survival.

25

Eighty percent of the mice immunized with CbpA truncate protein NR1X survived the challenge for 14 Days (results shown in Figure 5). All sham immunized mice were dead by day 8.

30

B. Active Protection With NR1XR2 Truncate Vaccination

C3H/HeJ mice (10/group) were immunized intraperitoneally with CbpA truncate protein NR1XR2 (15µg
35 in 50 µl PBS, plus 50 µl Complete Freund's Adjuvant). A group of 10 control immunized mice received pneumococcal recombinant protein SP90 and adjuvant. A second

immunization was administered four weeks later, 15µg protein i.p. with Incomplete Freund's Adjuvant. Blood was drawn (retro-orbital bleed) at weeks 3, 6, and 9 for analysis of immune response. The ELISA end point anti-CbpA truncate titer of pooled sera from the 10 CbpA immunized mice at 9 weeks was 4,096,000. Mice were challenged at week 10 with 680 CFU serotype 6B *S. pneumoniae* strain SPSJ2. Mice were monitored for 14 days for survival.

10

Fifty percent of the mice immunized with CbpA truncate protein NR1XR2 survived the challenge for 14 days (results shown in Figure 6). All control immunized mice were dead by day 9.

15

These data demonstrate that immunization with recombinant CbpA truncate proteins elicit production of specific antibodies capable of protecting against systemic pneumococcal infection and death. The data further indicate that the choline-binding region is not necessary for protection, as the immunogens were truncated proteins NR1X and NR1XR2. Additionally, the results suggest that a single amino terminal repeat may be sufficient to elicit a protective response. Cross protection is demonstrated as the recombinant pneumococcal protein was generated based on serotype 4 DNA sequence and protection was observed following challenge with a serotype 6B isolate.

30

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described.

35

WHAT IS CLAIMED IS:

1. A vaccine against bacterial infections comprising an immunogen which is a polypeptide truncate
5 of a pneumococcal surface-binding protein, analog or variant having a highly conserved immunogenic alpha-helical portion with respect to different types of pneumococcal bacteria, which polypeptide does not include a choline-binding portion.
- 10
2. A vaccine according to claim 1, wherein the amino acid sequence of said alpha-helical portion has at least 75 % identity with respect to the amino acid sequence of SEQ ID NO:1.
- 15
3. A vaccine according to claim 1, wherein the amino acid sequence of said alpha-helical portion has at least 85 % identity with respect to the amino acid sequence of SEQ ID NO:1.
- 20
4. A vaccine according to claim 1, wherein the amino acid sequence of said alpha-helical portion has at least 90 % identity with respect to the amino acid sequence of a member consisting of:
- 25 (a) the amino acid sequence of SEQ ID NO:1, and
(b) the amino acid sequence of SEQ ID NO:19.
5. A vaccine according to claim 1, wherein the amino acid sequence of said alpha-helical portion has
30 at least 95 % identity with respect to the amino acid sequence of a member selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:1, and
(b) the amino acid sequence of SEQ ID NO:19.
- 35
6. A vaccine according to claim 1, wherein said vaccine is for preventing or treating otitis media, sepsis, meningitis and lobar pneumonia infections.

7. A vaccine according to claim 6, wherein said vaccine is for invasive infections.

8. A vaccine according to claim 6, wherein
5 said vaccine is for otitis media infections caused by *S. pneumoniae*.

9. A vaccine according to claim 1, wherein
10 said polypeptide truncate comprise an amino acid sequence which has at least 90 % identity with respect to a member selected from the group consisting of the amino acid sequences of each of SEQ ID NOS:3 to 18.

10. A vaccine according to claim 1, wherein
15 said polypeptide truncate comprise an amino acid sequence which has at least 95 % identity with respect to a member selected from the group consisting of the amino acid sequences of each of SEQ ID NOS:3 to 18.

20 11. An antibody raised against an immunogen which is a polypeptide truncate of a pneumococcal surface-binding protein, analog or variant having a highly conserved immunogenic alpha-helical portion with respect to different types of pneumococcal bacteria,
25 which polypeptide does not include a choline-binding portion.

12. An antibody according to claim 11, wherein
30 the amino acid sequence of said alpha-helical portion has at least 85 % identity with respect to the amino acid sequence of SEQ ID NO:1.

13. An antibody according to claim 11, wherein
35 the amino acid sequence of said alpha-helical portion has at least 90 % identity with respect to the amino acid sequence of a member selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:1,
and

(b) the amino acid sequence of SEQ ID NO:19.

14. An antibody according to claim 11, wherein the amino acid sequence of said alpha-helical portion has at least 95 % identity with respect to the amino acid sequence of a member selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:1,
and
10 (b) the amino acid sequence of SEQ ID NO:19.

15. An antibody according to claim 11, wherein said polypeptide truncate comprise an amino acid sequence which has at least 95 % identity with respect to a member selected from the group consisting of the amino acid sequences of each of SEQ ID NOS:3 to 18.

16. An antibody according to claim 11, wherein said antibody is an antibody that will detect *S. pneumoniae* infections.

17. An antibody according to claim 15, wherein said antibody is effective for the prevention and/or treatment of *S. pneumoniae* infections.

18. An antibody according to claim 15, wherein said antibody is effective for the prevention and/or treatment of pneumococcal infections caused by types 1-5, 6A, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19F, 19A, 20, 22F, 23F and 33F *S. pneumoniae* bacteria.

19. A method for preventing and/or treating pneumococcal infections in a host comprising immunizing said host with a member selected from the group consisting of:

- (a) a vaccine according to claim 2, and
(b) at least one antibody raised against an immunogen which is a polypeptide truncate of a pneumococcal surface-binding protein, analog or variant

comprising an amino acid sequence that is has at least 90
% identity to the amino acid sequence of a member
selected from the group consisting of SEQ ID NO:3 to 18,
which polypeptide does not include a choline-binding
5 portion.


20. A polypeptide comprising an amino acid
sequence which has at least 90 % identity with respect to
a member selected from the group consisting of the amino
10 acid sequences of each of SEQ ID NOS:3 to 18.

21. An isolated polynucleotide comprising
polynucleotide sequence having at least 90 % identity to
a member selected from the group consisting of:

15 (a) a polynucleotide coding sequence encoding
a polypeptide comprising a member selected from the group
consisting of the amino acid sequences of each of SEQ ID
NOS:3 to 18, and


(b) and the complement of (a).

Figure 1




(Sheet 1 of 16)

Figure 2




(Sheet 2 of 16)

Figure 3




(Sheet 3 of 16)

Figure 4




(Sheet 4 of 16)

Figure 5



(Sheet 5 of 16)

Figure 6



(Sheet 6 of 16)

Figure 8

		Percent Identity																	
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
1	Nonway4	30.7	31.7	28.6	100.0	19.8	19.0	31.7	33.0	36.3	36.3	37.3	30.2	32.3	31.7	23.3	37.0	1	
2	ATCC33400(1)		63.4	55.1	30.7	39.6	34.7	63.4	75.0	65.3	65.3	64.4	27.9	59.4	63.4	37.6	62.0	2	
3	ATCC11733(2)			78.6	31.7	40.6	36.6	100.0	62.0	60.4	60.4	60.4	30.2	59.4	100.0	46.5	65.0	3	
4	ATCC2				28.6	37.8	36.7	78.6	58.2	63.3	63.3	63.3	24.4	67.7	78.6	38.8	60.2	4	
5	ATCC4					19.8	19.0	31.7	33.0	36.3	36.3	37.3	30.2	32.3	31.7	23.3	37.0	5	
6	ATCC6B						89.5	40.6	41.0	38.2	38.2	37.3	51.2	41.7	40.6	64.1	38.0	6	
7	ATCC18C							36.6	35.0	35.3	35.3	34.3	48.8	38.5	36.6	60.2	34.0	7	
8	R6X(2)								62.0	60.4	60.4	60.4	30.2	59.4	100.0	46.5	65.0	8	
9	SPB105(6B)									70.0	70.0	71.0	27.9	65.6	62.0	42.0	67.0	9	
10	SPB328(23F)										100.0	99.0	27.9	63.5	60.4	45.1	81.0	10	
11	SPB331(14)												99.0	27.9	63.5	60.4	45.1	81.0	11
12	SPB365(23F)													27.9	64.6	60.4	45.1	82.0	12
13	SPB609(6B)														23.3	30.2	66.3	29.1	13
14	SPR332(9V)															59.4	42.7	63.5	14
15	SPSJ2(6B)p																46.5	65.0	15
16	SPSJ9(14)																	41.0	16
17	SPSJ12(19A)																		17


(Sheet 8 of 16)

Figure 10

		Percent Identity														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	
1	▲	80.8	▲	98.1	92.7	100.0	78.8	82.7	79.8	79.8	79.8	76.0	78.8	83.7	91.3	Norway 4
2	▲	▲	86.5	78.8	76.0	80.8	82.7	83.7	78.8	78.8	78.8	76.9	82.7	77.9	79.8	ATCC33400(1)
3	▲	▲	▲	76.0	74.0	77.9	84.6	82.7	79.8	79.8	79.8	78.8	84.6	80.8	77.9	ATCC2
4	▲	▲	▲	▲	90.6	98.1	76.9	81.7	77.9	77.9	77.9	74.0	76.9	81.7	89.4	ATCC4
5	▲	▲	▲	▲	▲	92.7	79.2	83.3	84.4	84.4	84.4	76.0	79.2	81.2	93.8	ATCC6B
6	▲	▲	▲	▲	▲	▲	78.8	82.7	79.8	79.8	79.8	76.0	78.8	83.7	91.3	Norway 14
7	▲	▲	▲	▲	▲	▲	▲	82.7	82.7	82.7	82.7	81.7	100.0	83.7	78.8	R6X(2)
8	▲	▲	▲	▲	▲	▲	▲	▲	86.5	86.5	86.5	80.8	82.7	81.7	85.6	SPB105(6B)
9	▲	▲	▲	▲	▲	▲	▲	▲	▲	100.0	100.0	78.8	82.7	80.8	83.7	SPB328(23F)
10	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	100.0	78.8	82.7	80.8	83.7	SPB331(14)
11	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	78.8	82.7	80.8	83.7	SPB365(23F)
12	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	81.7	91.3	79.8	SPR332(9V)
13	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	83.7	78.8	SPSJ2(6B)passed
14	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	79.8	SPSJ9(14)
15	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	SPSJ12(19A)

(Sheet 10 of 16)

Figure 11



Decoration 'Decoration #1': Box residues that differ from the Consensus.

Decoration 'Decoration #2': Box residues that match the Consensus exactly.



(Sheet 11 of 16)

Figure 12

		Percent Identity																
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16		
1	▲	52.0	76.0	98.0	82.0	100.0	86.0	86.0	82.0	82.0	82.0	84.0	76.0	86.0	76.0	76.0	1	Norway4
2	▲	▲	58.5	52.0	55.2	52.0	43.1	44.8	43.1	44.8	43.1	48.3	49.1	43.1	50.9	46.3	2	ATCC33400(1)
3	▲	▲	▲	76.0	81.1	76.0	71.7	69.8	69.8	69.8	69.8	69.8	69.8	71.7	71.7	69.8	3	ATCC2
4	▲	▲	▲	▲	82.0	98.0	84.0	86.0	80.0	80.0	80.0	84.0	76.0	84.0	76.0	76.0	4	ATCC4
5	▲	▲	▲	▲	▲	82.0	62.7	64.5	64.4	64.4	66.1	67.7	71.7	62.7	71.7	70.4	5	ATCC6B
6	▲	▲	▲	▲	▲	▲	86.0	86.0	82.0	82.0	82.0	84.0	76.0	86.0	76.0	76.0	6	Norway14
7	▲	▲	▲	▲	▲	▲	▲	79.7	81.4	81.4	81.4	78.0	66.0	100.0	66.0	66.7	7	R6X(2)
8	▲	▲	▲	▲	▲	▲	▲	▲	78.0	78.0	78.0	83.9	69.8	79.7	67.9	66.7	8	SPB105(6B)
9	▲	▲	▲	▲	▲	▲	▲	▲	▲	98.3	98.3	78.0	67.9	81.4	67.9	64.8	9	SPB328(23F)
10	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	98.3	79.7	67.9	81.4	69.8	64.8	10	SPB331(14)
11	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	78.0	69.8	81.4	67.9	64.8	11	SPB365(23F)
12	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	69.8	78.0	75.5	66.7	12	SPB609(6B)
13	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	66.0	92.5	88.7	13	SPR332(9V)
14	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	66.0	66.7	14	SPSJ2(6B)passaged
15	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	88.7	15	SPSJ9(14)
16	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	▲	16	SPSJ12(19A)

(Sheet 12 of 16)

Figure 13



Decoration '1': Box residues that differ from the Consensus.

(Sheet 13 of 16)

Figure 14

		Percent Identity																	
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1	Norway 4	88.2	88.3	88.3	90.3	100.0	98.1	96.1	100.0	91.3	91.3	90.3	91.3	90.3	92.2	91.3	94.2	97.1	97.1
2	ATCC33400(1)		84.3	90.2	88.2	86.3	86.3	84.3	88.2	91.2	91.2	86.3	87.3	86.3	91.2	91.2	86.3	88.2	89.2
3	ATCC11733(2)			86.4	88.3	88.3	86.4	88.3	88.3	87.4	86.4	87.4	86.4	86.4	90.3	89.3	84.5	85.4	91.3
4	ATCC2				90.3	88.3	86.4	90.3	93.2	95.1	84.5	85.4	85.4	84.5	93.2	93.2	92.2	90.3	91.3
5	ATCC4					98.1	96.1	100.0	91.3	91.3	90.3	91.3	90.3	90.3	92.2	91.3	94.2	97.1	97.1
6	ATCC6B						96.1	98.1	91.3	89.3	90.3	91.3	90.3	90.3	90.3	91.3	92.2	95.1	95.1
7	ATCC18C-3							96.1	87.4	87.4	86.4	87.4	86.4	86.4	88.3	87.4	92.2	93.2	93.2
8	Norway14								91.3	91.3	90.3	91.3	90.3	90.3	92.2	91.3	94.2	97.1	97.1
9	R6X(2)									92.2	87.4	88.3	87.4	87.4	94.2	100.0	89.3	91.3	92.2
10	SPB105(6B)										85.4	86.4	85.4	85.4	96.1	92.2	91.3	93.2	92.2
11	SPB328(23F)											99.0	100.0	88.3	87.4	86.4	87.4	87.4	93.2
12	SPB331(14)													99.0	89.3	88.3	87.4	88.3	94.2
13	SPB365(23F)														88.3	87.4	86.4	87.4	93.2
14	SPR332(9V)															94.2	91.3	95.1	95.1
15	SPSJ2(6B)passaged																89.3	91.3	92.2
16	SPSJ9(14)																	94.2	93.2
17	SPSJ12(19A)																		94.2
18	Consensus A																		

(Sheet 14 of 16)

Figure 16

		Percent Identity																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
1	89.4	91.2	89.5	91.2	100.0	98.2	96.5	100.0	92.1	92.1	91.2	92.1	91.2	93.0	92.1	94.7	97.4	98.2	Norway4
2		85.8	87.7	89.4	87.6	85.8	89.4	92.0	92.0	86.7	87.6	86.7	86.7	92.0	92.0	87.6	89.4	91.2	ATCC33400(1)
3			87.7	89.5	89.5	87.7	89.5	90.4	90.4	87.7	88.6	87.7	87.7	91.2	90.4	86.0	86.8	91.2	ATCC11733(2)
4				91.2	89.5	87.7	91.2	93.9	86.0	86.8	86.0	86.0	86.0	93.9	93.9	93.0	91.2	91.2	ATCC2
5					98.2	96.5	100.0	92.1	92.1	92.1	91.2	91.2	91.2	93.0	92.1	94.7	97.4	98.2	ATCC4
6						96.5	98.2	92.1	92.1	92.1	91.2	92.1	91.2	91.2	92.1	93.0	95.6	96.5	ATCC6B
7							96.5	88.6	87.7	88.6	87.7	88.6	87.7	89.5	88.6	93.0	93.9	94.7	ATCC18C-3
8								92.1	92.1	92.1	91.2	92.1	91.2	93.0	92.1	94.7	97.4	98.2	Norway14
9									100.0	88.6	89.5	88.6	88.6	94.7	100.0	90.4	92.1	93.9	R6X(2)
10										88.6	89.5	88.6	88.6	94.7	100.0	90.4	92.1	93.9	SPB105(6B)
11											99.1	100.0	89.5	88.6	87.7	88.6	89.5	93.0	SPB328(23F)
12													99.1	90.4	89.5	88.6	89.5	93.9	SPB331(14)
13														89.5	88.6	87.7	88.6	93.0	SPB365(23F)
14															94.7	92.1	95.6	94.7	SPR332(9V)
15																90.4	92.1	93.9	SPSJ2(6B)passaged
16																	94.7	93.0	SPSJ9(14)
17																		95.6	SPSJ12(19A)
18																			Consensus B

(Sheet 16 of 16)

SEQUENCE LISTING

<110> Wizemann, Theresa M.
Koenig, Scott
Johnson, Leslie S

<120> Derivatives of Choline Binding Proteins for Vaccines

<130> 469201-364

<140>

<141>

<150> US 60/085,743

<151> 1998-05-15

<160> 38

<170> MS-Word (DOS Text)

<210> 1

<211> 103

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 1

```

Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu Ala Lys Lys Lys
 1                5                10                15
Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn Thr
                20                25                30
Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val Lys
                35                40                45
Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys Glu Ser Arg Asn
                50                55                60
Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys Val Glu Ser Lys Lys Ala
 65                70                75                80
Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp Arg Lys Lys Ala Glu
                85                90                95
Glu Glu Ala Lys Arg Lys Ala
                100

```

<210> 2

<211> 141

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 2

Glu Ala Lys Arg Lys Ala Glu Glu Ser Glu Lys Lys Ala Ala Glu Ala
 1 5 10 15
 Lys Gln Lys Val Asp Ala Glu Glu Tyr Ala Leu Glu Ala Lys Ile Ala
 20 25 30
 Glu Leu Glu Tyr Glu Val Gln Arg Leu Glu Lys Glu Leu Lys Glu Ile
 35 40 45
 Asp Glu Ser Asp Ser Glu Asp Tyr Leu Lys Glu Gly Leu Arg Ala Pro
 50 55 60
 Leu Gln Ser Lys Leu Asp Thr Lys Lys Ala Lys Leu Ser Lys Leu Glu
 65 70 75 80
 Glu Leu Ser Asp Lys Ile Asp Glu Leu Asp Ala Glu Ile Ala Lys Leu
 85 90 95
 Glu Val Gln Leu Lys Asp Ala Glu Gly Asn Asn Asn Val Glu Ala Tyr
 100 105 110
 Phe Lys Glu Gly Leu Glu Lys Thr Thr Ala Glu Lys Lys Ala Glu Leu
 115 120 125
 Glu Lys Ala Glu Ala Asp Leu Lys Lys Ala Val Asp Glu
 130 135 140

<210> 3

<211> 431

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 3

Thr Glu Lys Glu Val Thr Thr Pro Val Ala Thr Ser Ser Asn Lys Ala
 1 5 10 15
 Asn Lys Ser Gln Thr Glu His Met Lys Ala Ala Glu Gln Val Asp Glu
 20 25 30
 Tyr Ile Asn Lys Met Ile Gln Leu Asp Lys Arg Lys His Thr Gln Asn
 35 40 45
 Leu Ala Leu Asn Ile Lys Leu Ser Ala Ile Lys Thr Lys Tyr Leu Arg
 50 55 60
 Glu Leu Asn Val Leu Glu Glu Lys Ser Lys Lys Glu Glu Leu Thr Ser
 65 70 75 80
 Lys Thr Lys Lys Glu Ile Asp Ala Ala Phe Glu Gln Phe Asn Lys Asp
 85 90 95
 Thr Leu Lys Pro Gly Glu Lys Val Glu Glu Ala Glu Lys Lys Val Glu
 100 105 110
 Glu Ala Glu Lys Lys Ala Lys Asp Gln Lys Glu Glu Asp His Arg Asn
 115 120 125

Tyr Pro Thr Ile Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser
 130 135 140
 Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala
 145 150 155 160
 Lys Gly Ser Arg Asn Glu Glu Lys Ile Lys Lys Ala Lys Ala Glu Val
 165 170 175
 Glu Ser Lys Lys Ala Glu Ala Thr Lys Leu Glu Glu Ile Lys Thr Glu
 180 185 190
 Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Glu Ala Glu Glu
 195 200 205
 Glu Val Lys Asn Lys Leu Lys Lys Arg Thr Lys Arg Gly Ala Phe Gly
 210 215 220
 Glu Pro Ala Thr Pro Asp Lys Lys Glu Asn Asp Ala Lys Ser Ser Asp
 225 230 235 240
 Ser Ser Val Val Lys Lys Ser Ser Lys Pro Ile Leu Lys Ser Glu Lys
 245 250 255
 Lys Val Ala Glu Ala Glu Lys Lys Val Ala Glu Ala Glu Lys Lys Val
 260 265 270
 Ala Glu Ala Glu Lys Lys Ala Lys Asp Gln Lys Glu Glu Asp Arg Arg
 275 280 285
 Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu
 290 295 300
 Ser Asp Val Lys Val Lys Glu Ala Glu Leu Glu Leu Val Lys Glu Glu
 305 310 315 320
 Ala Lys Glu Pro Gln Asn Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys
 325 330 335
 Val Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr
 340 345 350
 Asp Arg Lys Lys Ala Glu Glu Ala Lys Arg Lys Val Ala Glu Glu Asp
 355 360 365
 Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Pro
 370 375 380
 Lys Pro Ala Pro Ala Pro Gln Pro Glu Lys Pro Ala Glu Gln Pro Lys
 385 390 395 400
 Ala Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu Asp Tyr Ala Arg Arg
 405 410 415
 Ser Glu Glu Glu Tyr Asn Pro Leu Asp Leu Thr Ala Pro Ala Lys
 420 425 430

<210> 4

<211> 251

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 4

```

Thr Glu Asn Glu Gly Ser Thr Gln Ala Ala Thr Ser Ser Asn Met Ala
 1           5           10           15
Lys Thr Glu His Arg Lys Ala Ala Lys Gln Val Val Asp Glu Tyr Ile
           20           25           30
Glu Lys Met Leu Arg Glu Ile Gln Leu Asp Arg Arg Lys His Thr Gln
           35           40           45
Asn Val Ala Leu Asn Ile Lys Leu Ser Ala Ile Lys Thr Lys Tyr Leu
           50           55           60
Arg Glu Leu Asn Val Leu Glu Glu Lys Ser Lys Asp Glu Leu Pro Ser
           65           70           75           80
Glu Ile Lys Ala Lys Leu Asp Ala Ala Phe Glu Lys Phe Lys Lys Asp
           85           90           95
Thr Leu Lys Pro Gly Glu Lys Val Ala Glu Ala Lys Lys Lys Val Glu
           100           105           110
Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn
           115           120           125
Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Phe
           130           135           140
Asp Val Lys Val Lys Glu Ala Glu Leu Glu Leu Val Lys Glu Glu Ala
           145           150           155           160
Lys Glu Ser Arg Asn Glu Gly Thr Ile Lys Gln Ala Lys Glu Lys Val
           165           170           175
Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys Thr Asp
           180           185           190
Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp
           195           200           205
Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Thr
           210           215           220
Gln Pro Glu Lys Pro Ala Pro Lys Pro Glu Lys Pro Ala Glu Gln Pro
           225           230           235           240
Lys Ala Glu Lys Thr Asp Asp Gln Gln Ala Glu
           245           250

```

<210> 5

<211> 413

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of

Streptococcus pneumoniae

<400> 5

Thr Glu Lys Glu Val Thr Thr Gln Val Pro Thr Tyr Ser Asn Met Ala
 1 5 10 15
 Lys Thr Glu His Arg Lys Ala Ala Lys Gln Val Val Asp Glu Thr Ile
 20 25 30
 Glu Lys Met Leu Arg Glu Ile Gln Leu Asp Arg Arg Lys His Thr Gln
 35 40 45
 Asn Phe Ala Phe Asn Met Lys Leu Ser Ala Ile Lys Thr Glu Tyr Leu
 50 55 60
 Tyr Gly Leu Lys Glu Lys Ser Glu Ala Glu Leu Pro Ser Glu Val Lys
 65 70 75 80
 Ala Lys Leu Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp Thr Leu Lys
 85 90 95
 Pro Gly Glu Lys Val Ala Glu Ala Lys Lys Lys Val Ala Glu Ala Glu
 100 105 110
 Lys Lys Ala Lys Ala Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr
 115 120 125
 Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp Val Glu
 130 135 140
 Val Lys Lys Ala Glu Leu Glu Leu Leu Lys Glu Glu Ala Lys Thr Arg
 145 150 155 160
 Asn Glu Asp Thr Ile Asn Gln Ala Lys Ala Lys Val Glu Ser Lys Lys
 165 170 175
 Ala Glu Ala Thr Leu Lys Glu Glu Ile Lys Thr Asp Arg Lys Lys Ala
 180 185 190
 Glu Glu Glu Ala Lys Arg Lys Ala Glu Ala Glu Glu Asp Lys Val Lys
 195 200 205
 Asp Lys Leu Lys Arg Arg Thr Lys Arg Ala Val Pro Gly Glu Pro Ala
 210 215 220
 Thr Phe Phe Lys Lys Glu Asn Asp Ala Lys Ser Ser Asp Ser Ser Val
 225 230 235 240
 Gly Glu Glu Thr Leu Pro Ser Pro Ser Leu Lys Ser Gly Lys Lys Val
 245 250 255
 Ala Glu Ala Glu Lys Lys Val Ala Glu Ala Glu Lys Lys Ala Lys Asp
 260 265 270
 Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn Thr Thr Lys Thr
 275 280 285
 Leu Asp Leu Glu Ile Ala Glu Ser Asp Val Lys Val Lys Glu Ala Glu
 290 295 300
 Leu Glu Leu Val Lys Glu Glu Ala Lys Gly Ser Arg Asn Glu Glu Lys
 305 310 315 320

Ile Asn Gln Ala Lys Ala Glu Val Glu Ser Lys Lys Ala Glu Ala Thr
 325 330 335

Arg Leu Glu Lys Thr Lys Thr Asp Arg Lys Lys Ala Glu Glu Glu Ala
 340 345 350

Lys Arg Lys Ala Ala Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu
 355 360 365

Gln Pro Gln Pro Ala Pro Ala Pro Gln Pro Glu Lys Pro Thr Glu Glu
 370 375 380

Pro Glu Asn Pro Ala Pro Ala Pro Lys Pro Glu Lys Pro Ala Glu Gln
 385 390 395 400

Pro Lys Ala Glu Lys Thr Asp Asp Gln Gln Ala Glu Glu
 405 410

<210> 6

<211> 446

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 6

Thr Glu Asn Glu Gly Ala Thr Gln Val Pro Thr Ser Ser Asn Arg Ala
 1 5 10 15

Asn Glu Ser Gln Ala Glu Gln Gly Glu Gln Pro Lys Lys Leu Asp Ser
 20 25 30

Glu Arg Asp Lys Ala Arg Lys Glu Val Glu Glu Tyr Val Lys Lys Ile
 35 40 45

Val Gly Glu Ser Tyr Ala Lys Ser Thr Lys Lys Arg His Thr Ile Thr
 50 55 60

Val Ala Leu Val Asn Glu Leu Asn Asn Ile Lys Asn Glu Tyr Leu Asn
 65 70 75 80

Lys Ile Val Glu Ser Thr Ser Glu Ser Gln Leu Gln Ile Leu Met Met
 85 90 95

Glu Ser Arg Ser Lys Val Asp Glu Ala Val Ser Lys Phe Glu Lys Asp
 100 105 110

Ser Ser Ser Ser Ser Ser Ser Asp Ser Ser Thr Lys Pro Glu Ala Ser
 115 120 125

Asp Thr Ala Lys Pro Asn Lys Pro Thr Glu Pro Gly Glu Lys Val Ala
 130 135 140

Glu Ala Lys Lys Lys Val Glu Glu Val Glu Lys Lys Ala Lys Asp Gln
 145 150 155 160

Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Ile Thr Tyr Lys Thr Leu
 165 170 175

Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu
 180 185 190

Glu Leu Val Lys Val Lys Ala Asn Glu Pro Arg Asp Lys Gln Lys Ile
 195 200 205

Lys Gln Ala Glu Ala Glu Val Glu Ser Lys Gln Ala Glu Ala Thr Arg
 210 215 220

Leu Lys Lys Ile Lys Thr Asp Arg Glu Glu Ala Glu Glu Glu Ala Lys
 225 230 235 240

Arg Arg Ala Asp Ala Lys Glu Gln Gly Lys Pro Lys Gly Arg Pro Lys
 245 250 255

Arg Gly Val Pro Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu Asn Asp
 260 265 270

Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Pro Ser Pro
 275 280 285

Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu
 290 295 300

Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn
 305 310 315 320

Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser
 325 330 335

Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala
 340 345 350

Lys Glu Pro Arg Asn Glu Glu Lys Val Lys Gln Ala Lys Ala Glu Val
 355 360 365

Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp
 370 375 380

Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp
 385 390 395 400

Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Pro
 405 410 415

Lys Thr Glu Lys Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu
 420 425 430

Gln Pro Lys Ala Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu
 435 440 445

<210> 7

<211> 428

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 7

Glu Gly Val Arg Ser Gly Asn Asn Ser Thr Val Thr Ser Ser Gly Gln
 1 5 10 15
 Asp Ile Ser Lys Lys Tyr Ala Asp Glu Val Glu Ser His Leu Gln Ser
 20 25 30
 Ile Leu Lys Asp Val Asn Lys Asn Leu Lys Lys Val Gln His Thr Gln
 35 40 45
 Asn Ala Asp Phe Asn Lys Lys Leu Ser Lys Ile Lys Thr Lys Tyr Leu
 50 55 60
 Tyr Glu Leu Asn Val Leu Glu Glu Lys Ser Glu Ala Glu Leu Thr Ser
 65 70 75 80
 Lys Thr Lys Glu Thr Lys Glu Glu Leu Thr Ala Ala Phe Glu Gln Phe
 85 90 95
 Lys Lys Asp Thr Leu Ser Thr Glu Pro Glu Lys Lys Val Ala Glu Ala
 100 105 110
 Lys Lys Lys Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu
 115 120 125
 Lys Asp Arg Arg Asn Tyr Pro Thr Ile Thr Tyr Lys Thr Leu Glu Leu
 130 135 140
 Glu Ile Ala Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu
 145 150 155 160
 Val Lys Val Lys Ala Asn Glu Pro Arg Asp Glu Glu Lys Ile Lys Gln
 165 170 175
 Ala Glu Ala Lys Val Glu Ser Lys Gln Ala Glu Ala Thr Arg Leu Lys
 180 185 190
 Lys Ile Lys Thr Asp Arg Glu Gln Ala Glu Ala Thr Arg Leu Glu Asn
 195 200 205
 Ile Lys Thr Asp Arg Glu Gln Ala Glu Glu Glu Ala Lys Val Lys Asp
 210 215 220
 Glu Pro Lys Lys Arg Thr Lys Arg Gly Val Leu Gly Glu Pro Ala Thr
 225 230 235 240
 Pro Asp Lys Lys Glu Asn Asp Ala Lys Ser Ser Asp Ser Ser Val Gly
 245 250 255
 Glu Glu Thr Leu Pro Ser Pro Ser Leu Lys Pro Glu Lys Lys Val Ala
 260 265 270
 Glu Ala Glu Lys Lys Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln
 275 280 285
 Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu
 290 295 300
 Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu
 305 310 315 320
 Glu Leu Val Lys Glu Glu Ala Lys Glu Pro Arg Asn Glu Glu Lys Val


```

                325                330                335
Lys Gln Ala Lys Ala Glu Val Glu Ser Lys Gln Ala Glu Ala Thr Arg
                340                345                350
Leu Glu Asn Ile Lys Thr Asp Arg Lys Lys Ala Glu Glu Glu Ala Lys
                355                360                365
Arg Lys Ala Ala Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln
                370                375                380
Pro Gln Pro Ala Pro Ala Pro Gln Pro Glu Lys Pro Ala Pro Lys Asp
                385                390                395                400
Glu Lys Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Gln Pro
                405                410                415
Lys Ala Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu
                420                425

```

<210> 8

<211> 219

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 8

```

Glu Gly Val Arg Ser Gly Asn Asn Ser Thr Val Thr Ser Ser Gly Gln
  1                5                10                15
Asp Ile Ser Lys Lys Tyr Ala Asp Glu Val Glu Ser His Leu Gln Ser
                20                25                30
Ile Leu Lys Asp Val Asn Lys Asn Leu Lys Lys Val Gln His Thr Gln
                35                40                45
Asn Ala Asp Phe Asn Lys Lys Leu Ser Lys Ile Lys Pro Lys Tyr Leu
                50                55                60
Tyr Glu Leu Lys Cys Leu Glu Glu Lys Ser Glu Ala Glu Leu Thr Ser
                65                70                75                80
Lys Pro Lys Asn Lys Arg Arg Val Thr Ala Ala Phe Glu Gln Phe Lys
                85                90                95
Lys Asp Thr Leu Ser Thr Glu Pro Glu Lys Lys Val Ala Glu Ala Lys
                100                105                110
Lys Lys Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Lys
                115                120                125
Asp Arg Arg Asn Tyr Pro Thr Ile Thr Tyr Lys Thr Leu Glu Leu Glu
                130                135                140
Ile Ala Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val
                145                150                155                160
Lys Glu Glu Ala Lys Glu Pro Arg Asn Glu Glu Lys Val Lys Gln Ala

```

165 170 175
 Lys Ala Glu Val Glu Ser Lys Gln Ala Glu Ala Thr Arg Leu Glu Lys
 180 185 190
 Ile Lys Thr Asp Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala
 195 200 205
 Ala Glu Glu Asp Lys Val Lys Glu Lys Pro Ala
 210 215

<210> 9
 <211> 446
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description: Amino acid sequence derived from a cDNA from the genome of
 Streptococcus pneumoniae

<400> 9
 Thr Glu Asn Glu Gly Ala Thr Gln Val Pro Thr Ser Ser Asn Arg Ala
 1 5 10 15
 Asn Glu Ser Gln Ala Glu Gln Gly Glu Gln Pro Lys Lys Leu Asp Ser
 20 25 30
 Glu Arg Asp Lys Ala Arg Lys Glu Val Glu Glu Tyr Val Lys Lys Ile
 35 40 45
 Val Gly Glu Ser Tyr Ala Lys Ser Thr Lys Lys Arg His Thr Ile Thr
 50 55 60
 Val Ala Leu Val Asn Glu Leu Asn Asn Ile Lys Asn Glu Tyr Leu Asn
 65 70 75 80
 Lys Ile Val Glu Ser Thr Ser Glu Ser Gln Leu Gln Ile Leu Met Met
 85 90 95
 Glu Ser Arg Ser Lys Val Asp Glu Ala Val Ser Lys Phe Glu Lys Asp
 100 105 110
 Ser Ser Ser Ser Ser Ser Ser Asp Ser Ser Thr Lys Pro Glu Ala Ser
 115 120 125
 Asp Thr Ala Lys Pro Asn Lys Pro Thr Glu Pro Gly Glu Lys Val Ala
 130 135 140
 Glu Ala Lys Lys Lys Val Glu Glu Ala Glu Lys Lys Ala Lys Asp Gln
 145 150 155 160
 Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Ile Thr Tyr Lys Thr Leu
 165 170 175
 Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu
 180 185 190
 Glu Leu Val Lys Val Lys Ala Asn Glu Pro Arg Asp Glu Gln Lys Ile
 195 200 205
 Lys Gln Ala Glu Ala Glu Val Glu Ser Lys Gln Ala Glu Ala Thr Arg

```

210                215                220
Leu Lys Lys Ile Lys Thr Asp Arg Glu Glu Ala Glu Glu Glu Ala Lys
225                230                235                240
Arg Arg Ala Asp Ala Lys Glu Gln Gly Lys Pro Lys Gly Arg Ala Lys
                245                250                255
Arg Gly Val Pro Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu Asn Asp
                260                265                270
Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Pro Ser Pro
                275                280                285
Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu
                290                295                300
Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn
305                310                315                320
Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser
                325                330                335
Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala
                340                345                350
Lys Glu Pro Arg Asn Glu Glu Lys Val Lys Gln Ala Lys Ala Glu Val
                355                360                365
Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp
                370                375                380
Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp
385                390                395                400
Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Pro
                405                410                415
Lys Ala Glu Lys Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu
                420                425                430
Gln Pro Lys Ala Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu
                435                440                445

```

<210> 10

<211> 414

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 10

```

Thr Glu Asn Glu Gly Ser Thr Gln Ala Ala Thr Ser Ser Asn Met Ala
  1                5                10                15

```

```

Lys Thr Glu His Arg Lys Ala Ala Lys Gln Val Val Asp Glu Tyr Ile
  20                25                30

```

```

Glu Lys Met Leu Arg Glu Ile Gln Leu Asp Arg Arg Lys His Thr Gln

```

	35		40		45														
Asn	Val	Ala	Leu	Asn	Ile	Lys	Leu	Ser	Ala	Ile	Lys	Thr	Lys	Tyr	Leu				
	50					55					60								
Arg	Glu	Leu	Asn	Val	Leu	Glu	Glu	Lys	Ser	Lys	Asp	Glu	Leu	Pro	Ser				
	65				70					75					80				
Glu	Ile	Lys	Ala	Lys	Leu	Asp	Ala	Ala	Phe	Glu	Lys	Phe	Lys	Lys	Asp				
				85					90						95				
Thr	Leu	Lys	Pro	Gly	Glu	Lys	Val	Ala	Glu	Ala	Lys	Lys	Lys	Val	Glu				
			100					105						110					
Glu	Ala	Lys	Lys	Lys	Ala	Glu	Asp	Gln	Lys	Glu	Glu	Asp	Arg	Arg	Asn				
		115					120					125							
Tyr	Pro	Thr	Asn	Thr	Tyr	Lys	Thr	Leu	Glu	Leu	Glu	Ile	Ala	Glu	Phe				
	130					135					140								
Asp	Val	Lys	Val	Lys	Glu	Ala	Glu	Leu	Glu	Leu	Val	Lys	Glu	Glu	Ala				
	145				150					155					160				
Lys	Glu	Ser	Arg	Asn	Glu	Gly	Thr	Ile	Lys	Gln	Ala	Lys	Glu	Lys	Val				
				165					170					175					
Glu	Ser	Lys	Lys	Ala	Glu	Ala	Thr	Arg	Leu	Glu	Asn	Ile	Lys	Thr	Asp				
			180					185						190					
Arg	Lys	Lys	Ala	Glu	Glu	Glu	Ala	Lys	Arg	Lys	Ala	Asp	Ala	Lys	Leu				
		195					200					205							
Lys	Glu	Ala	Asn	Val	Ala	Thr	Ser	Asp	Gln	Gly	Lys	Pro	Lys	Gly	Arg				
	210					215					220								
Ala	Lys	Arg	Gly	Val	Pro	Gly	Glu	Leu	Ala	Thr	Pro	Asp	Lys	Lys	Glu				
	225				230					235					240				
Asn	Asp	Ala	Lys	Ser	Ser	Asp	Ser	Ser	Val	Gly	Glu	Glu	Thr	Leu	Pro				
				245					250					255					
Ser	Ser	Ser	Leu	Lys	Ser	Gly	Lys	Lys	Val	Ala	Glu	Ala	Glu	Lys	Lys				
			260					265						270					
Val	Glu	Glu	Ala	Glu	Lys	Lys	Ala	Lys	Asp	Gln	Lys	Glu	Glu	Asp	Arg				
		275					280						285						
Arg	Asn	Tyr	Pro	Thr	Asn	Thr	Tyr	Lys	Thr	Leu	Asp	Leu	Glu	Ile	Ala				
	290					295					300								
Glu	Ser	Asp	Val	Lys	Val	Lys	Glu	Ala	Glu	Leu	Glu	Leu	Val	Lys	Glu				
	305				310					315				320					
Glu	Ala	Lys	Glu	Pro	Arg	Asp	Glu	Glu	Lys	Ile	Lys	Gln	Ala	Lys	Ala				
				325					330					335					
Lys	Val	Glu	Ser	Lys	Lys	Ala	Glu	Ala	Thr	Arg	Leu	Glu	Asn	Ile	Lys				
			340					345					350						
Thr	Asp	Arg	Asp	Asp	Ala	Glu	Glu	Glu	Ala	Lys	Arg	Lys	Ala	Ala	Glu				
	355						360					365							

Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro
 370 375 380

Ala Thr Gln Pro Glu Lys Pro Ala Pro Lys Pro Glu Lys Pro Ala Glu
 385 390 395 400

Gln Pro Lys Ala Glu Lys Thr Asp Asp Gln Gln Ala Glu Glu
 405 410

<210> 11

<211> 425

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of
 Streptococcus pneumoniae

<400> 11

Thr Glu Lys Glu Val Thr Thr Gln Val Ala Thr Ser Ser Asn Arg Ala
 1 5 10 15

Asn Glu Ser Gln Ala Gly His Arg Lys Ala Ala Glu Gln Phe Asp Glu
 20 25 30

Tyr Ile Lys Thr Met Ile Gln Leu Asp Arg Arg Lys His Thr Gln Asn
 35 40 45

Phe Ala Leu Asn Ile Lys Leu Ser Arg Ile Lys Thr Glu Tyr Leu Arg
 50 55 60

Lys Leu Asn Val Leu Glu Glu Lys Ser Lys Ala Glu Leu Pro Ser Glu
 65 70 75 80

Thr Lys Lys Glu Ile Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp Thr
 85 90 95

Asn Arg Thr Lys Lys Thr Val Ala Glu Ala Glu Lys Lys Val Glu Glu
 100 105 110

Ala Lys Lys Lys Ala Lys Ala Gln Lys Glu Glu Asp His Arg Asn Tyr
 115 120 125

Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp
 130 135 140

Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys
 145 150 155 160

Glu Ser Arg Asp Asp Glu Lys Ile Lys Gln Ala Glu Ala Lys Val Glu
 165 170 175

Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys Thr Asp Arg
 180 185 190

Glu Lys Ala Glu Glu Glu Ala Lys Arg Arg Ala Glu Ala Lys Leu Lys
 195 200 205

Glu Ala Val Glu Lys Asn Val Ala Thr Ser Glu Gln Asp Lys Pro Lys
 210 215 220

Gly Arg Arg Lys Arg Gly Val Pro Gly Glu Gln Ala Thr Pro Asp Lys
 225 230 235 240

Lys Glu Asn Asp Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Ala
 245 250 255

Leu Pro Ser Pro Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu
 260 265 270

Lys Lys Val Ala Glu Ala Glu Lys Lys Ala Lys Ala Gln Lys Glu Glu
 275 280 285

Asp Arg Arg Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu
 290 295 300

Ile Ala Glu Ser Asp Val Lys Val Lys Glu Ser Glu Leu Glu Leu Val
 305 310 315 320

Lys Glu Glu Ala Lys Glu Ser Arg Asn Glu Glu Lys Val Asn Gln Ala
 325 330 335

Lys Ala Lys Val Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys
 340 345 350

Ile Lys Thr Asp Arg Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala
 355 360 365

Ala Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro
 370 375 380

Ala Pro Ala Pro Gln Pro Glu Lys Pro Thr Glu Glu Pro Glu Asn Pro
 385 390 395 400

Ala Pro Ala Pro Lys Pro Glu Lys Pro Ala Glu Gln Pro Lys Ala Glu
 405 410 415

Lys Thr Asp Asp Gln Gln Ala Glu Glu
 420 425

<210> 12
 <211> 426
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description: Amino acid sequence derived from a cDNA from the genome of
 Streptococcus pneumoniae

<400> 12
 Thr Glu Lys Glu Val Thr Thr Gln Val Ala Thr Ser Ser Asn Lys Ala
 1 5 10 15

Asn Lys Ser Gln Thr Glu His Met Lys Ala Ala Lys Gln Val Asp Glu
 20 25 30

Tyr Ile Lys Lys Lys Ile Gln Leu Asp Arg Arg Lys His Thr Gln Asn
 35 40 45

Val Gly Leu Leu Thr Lys Leu Gly Val Ile Lys Thr Glu Tyr Leu His
 50 55 60

Gly Leu Ser Val Ser Lys Lys Lys Ser Glu Ala Glu Leu Pro Ser Glu
 65 70 75 80

Ile Lys Ala Lys Leu Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp Thr
 85 90 95

Leu Pro Thr Glu Pro Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val
 100 105 110

Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Lys Asp Leu Arg
 115 120 125

Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Asp Ile Ala Glu
 130 135 140

Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu
 145 150 155 160

Ala Lys Glu Ser Arg Asp Glu Lys Lys Ile Asn Gln Ala Lys Ala Lys
 165 170 175

Val Glu Asn Lys Lys Ala Glu Ala Thr Arg Leu Lys Asn Ile Lys Thr
 180 185 190

Asp Arg Glu Lys Ala Glu Glu Ala Lys Arg Arg Ala Asp Ala Lys Leu
 195 200 205

Gln Glu Ala Asn Val Ala Thr Ser Glu Gln Asp Lys Ser Lys Arg Arg
 210 215 220

Ala Lys Arg Glu Val Leu Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu
 225 230 235 240

Asn Asp Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Thr
 245 250 255

Ser Pro Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys
 260 265 270

Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg
 275 280 285

Arg Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala
 290 295 300

Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu
 305 310 315 320

Glu Ala Lys Glu Ser Arg Asn Glu Glu Lys Ile Lys Gln Val Lys Ala
 325 330 335

Lys Val Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys
 340 345 350

Thr Asp Arg Lys Lys Ala Glu Glu Glu Glu Ala Lys Arg Arg Ala Ala
 355 360 365

Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala
 370 375 380

Pro Ala Pro Gln Pro Glu Lys Pro Thr Glu Glu Pro Glu Asn Pro Ala
 385 390 395 400

Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Lys Pro Lys Ala
 405 410 415

Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu
 420 425

<210> 13

<211> 425

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of
 Streptococcus pneumoniae

<400> 13

Thr Glu Lys Glu Val Thr Thr Gln Val Ala Thr Ser Ser Asn Lys Ala
 1 5 10 15

Asn Lys Ser Gln Thr Glu His Met Lys Ala Ala Lys Gln Val Asp Glu
 20 25 30

Tyr Ile Lys Lys Lys Leu Gln Leu Asp Arg Arg Lys His Thr Gln Asn
 35 40 45

Val Gly Leu Leu Thr Lys Leu Gly Val Ile Lys Thr Glu Tyr Leu His
 50 55 60

Gly Leu Ser Val Ser Lys Lys Lys Ser Glu Ala Glu Leu Pro Ser Glu
 65 70 75 80

Ile Lys Ala Lys Leu Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp Thr
 85 90 95

Leu Pro Thr Glu Pro Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val
 100 105 110

Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Lys Asp Leu Arg
 115 120 125

Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Asp Ile Ala Glu
 130 135 140

Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu
 145 150 155 160

Ala Lys Glu Ser Arg Asp Glu Lys Lys Ile Asn Gln Ala Lys Ala Lys
 165 170 175

Val Glu Asn Lys Lys Ala Glu Ala Thr Arg Leu Lys Asn Ile Lys Thr
 180 185 190

Asp Arg Glu Lys Ala Glu Glu Ala Lys Arg Arg Ala Asp Ala Lys Leu
 195 200 205

Gln Glu Ala Asn Val Ala Thr Ser Glu Gln Asp Lys Ser Lys Arg Arg
 210 215 220

Ala Lys Arg Glu Val Phe Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu
 225 230 235 240

Asn Asp Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Thr
 245 250 255

Ser Pro Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys
 260 265 270

Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg
 275 280 285

Arg Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala
 290 295 300

Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu
 305 310 315 320

Glu Ala Lys Glu Ser Arg Asn Glu Glu Lys Ile Lys Gln Val Lys Ala
 325 330 335

Lys Val Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys
 340 345 350

Thr Asp Arg Lys Lys Ala Glu Glu Glu Glu Ala Lys Arg Arg Ala Ala
 355 360 365

Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala
 370 375 380

Pro Ala Pro Gln Pro Glu Lys Pro Thr Glu Glu Pro Glu Asn Pro Ala
 385 390 395 400

Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Lys Pro Lys Ala
 405 410 415

Glu Lys Pro Ala Asp Gln Gln Ala Glu
 420 425

<210> 14

<211> 424

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 14

Thr Glu Lys Glu Val Thr Thr Gln Val Ala Thr Ser Ser Asn Arg Ala
 1 5 10 15

Asn Lys Ser Gln Thr Glu His Met Lys Ala Ala Lys Gln Val Asp Glu
 20 25 30

Tyr Ile Lys Lys Lys Leu Gln Leu Asp Arg Arg Lys His Thr Gln Asn
 35 40 45

Val Gly Leu Leu Thr Lys Leu Gly Val Ile Lys Thr Glu Tyr Leu His
 50 55 60

Gly Leu Ser Val Ser Lys Lys Lys Ser Glu Ala Glu Leu Pro Ser Glu
 65 70 75 80

Ile Lys Ala Lys Leu Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp Thr
85 90 95

Leu Pro Thr Glu Pro Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val
100 105 110

Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Lys Asp Leu Arg
115 120 125

Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Asp Ile Ala Glu
130 135 140

Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu
145 150 155 160

Ala Lys Glu Ser Arg Asp Glu Lys Lys Ile Asn Gln Ala Lys Ala Lys
165 170 175

Val Glu Asn Lys Lys Ala Glu Ala Thr Arg Leu Lys Asn Ile Lys Thr
180 185 190

Asp Arg Glu Lys Ala Glu Glu Ala Lys Arg Arg Ala Asp Ala Lys Leu
195 200 205

Gln Glu Ala Asn Val Ala Thr Ser Glu Gln Asp Lys Ser Lys Arg Arg
210 215 220

Ala Lys Arg Glu Val Leu Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu
225 230 235 240

Asn Asp Ala Lys Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Thr
245 250 255

Ser Pro Ser Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys
260 265 270

Val Glu Glu Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg
275 280 285

Arg Asn Tyr Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala
290 295 300

Glu Ser Asp Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu
305 310 315 320

Glu Ala Lys Glu Ser Arg Asn Glu Glu Lys Ile Lys Gln Val Lys Ala
325 330 335

Lys Val Glu Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Asn Ile Lys
340 345 350

Thr Asp Arg Lys Lys Ala Glu Glu Glu Glu Ala Lys Arg Arg Ala Ala
355 360 365

Glu Glu Asp Lys Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala
370 375 380

Pro Ala Pro Gln Pro Glu Lys Pro Thr Glu Glu Pro Glu Asn Pro Ala
385 390 395 400

Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Lys Pro Lys Ala


```

                245                250                255
Leu Lys Pro Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu
                260                265                270
Ala Asp Lys Lys Ala Lys Ala Gln Lys Glu Glu Asp Arg Arg Asn Tyr
                275                280                285
Pro Thr Asn Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp
                290                295                300
Val Lys Val Lys Glu Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys
305                310                315                320
Glu Ser Arg Asn Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys Val Glu
                325                330                335
Ser Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp Arg
                340                345                350
Lys Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp Lys
                355                360                365
Val Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Pro Gln
370                375                380
Pro Glu Lys Pro Ala Glu Glu Pro Glu Asn Pro Val Pro Ala Pro Lys
385                390                395                400
Pro Glu Asn Pro Ala Glu Gln Pro Lys Ala Glu Lys Pro Ala Asp Gln
                405                410                415

Gln Ala Glu

```

```

<210> 16
<211> 414
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Description: Amino acid sequence derived from a cDNA from the genome of
        Streptococcus pneumoniae

```

```

<400> 16
Thr Glu Asn Glu Gly Ser Thr Gln Ala Ala Thr Ser Ser Asn Met Ala
  1                5                10                15
Lys Thr Glu His Arg Lys Ala Ala Lys Gln Val Val Asp Glu Tyr Ile
                20                25                30
Glu Lys Met Leu Arg Glu Ile Gln Leu Asp Arg Arg Lys His Thr Gln
                35                40                45
Asn Val Ala Leu Asn Ile Lys Leu Ser Ala Ile Lys Thr Lys Tyr Leu
                50                55                60
Arg Glu Leu Asn Val Leu Glu Glu Lys Ser Lys Asp Glu Leu Pro Ser
                65                70                75                80
Glu Ile Lys Ala Lys Leu Asp Ala Ala Phe Glu Lys Glu Lys Lys Asp

```


<210> 17

<211> 412

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of *Streptococcus pneumoniae*

<400> 17

Glu Gly Val Arg Ser Glu Asn Asn Pro Thr Val Thr Ser Ser Gly Gln
 1 5 10 15
 Asp Ile Ser Lys Lys Tyr Ala Asp Glu Val Lys Ser His Leu Glu Lys
 20 25 30
 Ile Leu Ser Glu Ile Gln Thr Asn Leu Asp Arg Ser Lys His Ile Lys
 35 40 45
 Thr Val Asn Leu Ile Asn Lys Leu Gln Asp Ile Lys Arg Thr Tyr Leu
 50 55 60
 Tyr Glu Leu Asn Val Leu Glu Asp Lys Ser Lys Ala Glu Leu Pro Ser
 65 70 75 80
 Lys Ile Lys Ala Glu Leu Asp Ala Ala Phe Glu Gln Phe Lys Lys Asp
 85 90 95
 Thr Leu Pro Thr Glu Pro Gly Lys Lys Val Ala Glu Ala Lys Lys Lys
 100 105 110
 Val Glu Glu Ala Glu Lys Lys Ala Lys Ala Gln Lys Glu Glu Asp Tyr
 115 120 125
 Arg Asn Tyr Pro Thr Ile Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala
 130 135 140
 Glu Ser Asp Val Lys Val Lys Glu Ala Glu Leu Glu Leu Val Lys Lys
 145 150 155 160
 Glu Ala Asp Glu Ser Arg Asn Glu Gly Thr Ile Asn Gln Ala Lys Ala
 165 170 175
 Lys Val Glu Ser Glu Gln Ala Glu Ala Thr Arg Leu Lys Lys Ile Lys
 180 185 190
 Thr Asp Arg Glu Lys Ala Glu Glu Glu Ala Lys Arg Arg Ala Asp Ala
 195 200 205
 Lys Glu Gln Asp Glu Ser Lys Arg Arg Lys Ser Arg Val Lys Arg Gly
 210 215 220
 Asp Phe Gly Glu Pro Ala Thr Pro Asp Lys Lys Glu Asn Asp Ala Lys
 225 230 235 240
 Ser Ser Asp Ser Ser Val Gly Glu Glu Thr Leu Pro Ser Pro Ser Leu
 245 250 255
 Lys Pro Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu Ala
 260 265 270

Glu Lys Lys Ala Lys Asp Gln Lys Glu Glu Asp His Arg Asn Tyr Pro
 275 280 285

Thr Ile Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp Val
 290 295 300

Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys Gly
 305 310 315 320

Ser Arg Asn Glu Glu Lys Val Lys Gln Ala Lys Ala Glu Val Glu Ser
 325 330 335

Lys Lys Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp Arg Lys
 340 345 350

Lys Ala Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp Lys Val
 355 360 365

Lys Glu Lys Pro Ala Glu Gln Pro Gln Pro Ala Pro Ala Pro Gln Pro
 370 375 380

Glu Lys Pro Ala Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Gln Pro
 385 390 395 400

Lys Ala Glu Lys Pro Ala Asp Gln Gln Ala Glu Glu
 405 410

<210> 18
 <211> 406
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description: Amino acid sequence derived from a cDNA from the genome of
 Streptococcus pneumoniae

<400> 18
 Thr Glu Asn Glu Gly Thr Thr Gln Ala Pro Thr Ser Ser Asn Arg Gly
 1 5 10 15

Asn Glu Ser Gln Ala Glu His Met Lys Ala Ala Lys Gln Val Asp Glu
 20 25 30

Tyr Ile Glu Lys Met Leu Gln Leu Asp Arg Arg Lys His Thr Gln Asn
 35 40 45

Val Gly Leu Leu Thr Lys Leu Gly Ala Ile Lys Thr Glu Tyr Leu Arg
 50 55 60

Gly Leu Ser Val Ser Lys Glu Lys Ser Thr Ala Glu Leu Pro Ser Glu
 65 70 75 80

Ile Lys Glu Lys Leu Thr Ala Ala Phe Lys Gln Phe Lys Lys Asp Thr
 85 90 95

Leu Lys Pro Glu Lys Lys Val Ala Glu Ala Glu Lys Lys Val Ala Glu
 100 105 110

Ala Lys Lys Lys Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn Tyr
 115 120 125

Pro Thr Ile Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp
 130 135 140
 Val Glu Val Lys Lys Ala Glu Leu Glu Leu Val Lys Val Lys Ala Asn
 145 150 155 160
 Glu Pro Arg Asp Glu Glu Lys Ile Lys Gln Ala Glu Ala Glu Val Glu
 165 170 175
 Ser Lys Lys Ala Glu Ala Thr Arg Leu Lys Lys Ile Lys Thr Asp Arg
 180 185 190
 Glu Lys Ala Glu Glu Glu Ala Lys Arg Arg Val Asp Ala Lys Glu Gln
 195 200 205
 Asp Glu Ser Ser Lys Arg Arg Lys Ser Arg Val Lys Arg Gly Asp Leu
 210 215 220
 Gly Glu Gln Ala Thr Pro Asp Lys Lys Glu Asn Asp Ala Lys Ser Ser
 225 230 235 240
 Asp Ser Ser Val Gly Glu Glu Thr Leu Pro Ser Pro Ser Leu Lys Pro
 245 250 255
 Gly Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu Ala Asp Lys
 260 265 270
 Lys Ala Lys Ala Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn
 275 280 285
 Thr Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val
 290 295 300
 Lys Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys Glu Pro Arg
 305 310 315 320
 Asn Glu Glu Lys Val Lys Gln Ala Lys Ala Glu Val Glu Ser Lys Lys
 325 330 335
 Ala Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp Arg Lys Lys Ala
 340 345 350
 Glu Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp Lys Val Lys Glu
 355 360 365
 Lys Pro Ala Glu Gln Pro Lys Pro Ala Pro Ala Pro Gln Pro Glu Lys
 370 375 380
 Pro Ala Pro Lys Pro Glu Asn Pro Ala Glu Gln Pro Lys Ala Glu Lys
 385 390 395 400
 Pro Ala Asp Gln Gln Ala
 405

<210> 19

<211> 114

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from a cDNA from the genome of

Streptococcus pneumoniae

<400> 19

Lys Lys Val Ala Glu Ala Glu Lys Lys Val Glu Glu Ala Lys Lys Lys
 1 5 10 15

Ala Glu Asp Gln Lys Glu Glu Asp Arg Arg Asn Tyr Pro Thr Asn Thr
 20 25 30

Tyr Lys Thr Leu Glu Leu Glu Ile Ala Glu Ser Asp Val Glu Val Lys
 35 40 45

Lys Ala Glu Leu Glu Leu Val Lys Glu Glu Ala Lys Glu Ser Arg Asn
 50 55 60

Glu Glu Lys Ile Lys Gln Ala Lys Ala Lys Val Glu Ser Lys Lys Ala
 65 70 75 80

Glu Ala Thr Arg Leu Glu Lys Ile Lys Thr Asp Arg Lys Lys Ala Glu
 85 90 95

Glu Glu Ala Lys Arg Lys Ala Ala Glu Glu Asp Lys Val Lys Glu Lys
 100 105 110

Pro Ala

<210> 20

<211> 1295

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: cDNA derived from the genome S. pneumoniae

<400> 20

```

acagagaagg aggtaactac cccagtagcc acttcttcta ataaggcaa taaagtcag      60
acagaacata tgaagctgc tgaacaagtc gatgaatata taaacaaaat gatccaatta    120
gataaaagaa aacataccca aaatctcgcc ttaaacataa agttgagcgc aattaaaacg     180
aagtattttg gtgaattaaa tgttttagaa gagaagtcga aaaaagaaga gttgacgtca    240
aaaacaaaaa aagagataga cgcagctttt gagcagttta acaaagatac attgaaacca    300
ggagaaaagg ttgaagaagc tgagaagaag gttgaagaag ctgagaaaaa agccaaggat    360
caaaaagaag aagatcaccg taactaccca accattactt acaaacgct tgaacttgaa    420
attgctgagt ccgatgtgga agttaaaaaa gcgagcttg aactagtaa agaggaagct    480
aagggatctc gaaacgagga aaaaattaag aaagcaaaag cggaagtga gagtaaaaaa    540
gctgaggcta caaagttaga agaatcaag acagaacgta aaaaagcaga agaagaagct    600
aaacgaaaag cagaagcaga agaagaagtt aaaaataaac taaagaagcg gacaaaacga    660
ggagcttttg gagagccagc aacacctgat aaaaaagaaa atgatgcaa gtcttcagat    720
tctagcgtgg tgaagaaatc ttccaagccc atcctgaaat cagaaaaaaa agtagcagaa    780
gctgagaaga aggttgaga agctgagaag aaggttgag aagctgaga aaaagccaag    840
gatcaaaaag aagaagatcg ccgtaactac ccaaccaata cttaaaaaac gcttgaactt    900
gaaattgctg agtccgatgt gaaagttaaa gaagcggagc ttgaactagt aaaagaggaa    960
gctaaggaac ctcaaacgga ggaaaaaatt aagcaagcaa aagcgaagat tgagagtaaa   1020
aaagctgagg ctacaagggt agaaaaaatc aagacagatc gtaaaaaagc agaagaagct   1080
aaacgaaaag tagcagaaga agataaagtt aaagaaaaac cagctgaaca accacaacca   1140
gctcctgcac caaacaccgc gccggctcct caaccagaaa aaccagctga acaaccaaaa   1200
gcagaaaaac cagctgatca acaagctgaa gaagactatg ctctgatatc agaagaagaa   1260
tataaccggc ttgacttaac agcaccggca aaagc                                     1295
    
```

<210> 21

<211> 755

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: cDNA from Streptococcus pneumoniae

<400> 21

```

acagagaacg aggggaagtac ccaagcagcc acttcttcta atatggcaaa gacagaacat      60
aggaaagctg ctaaacaagt cgtc gatgaa tatatagaaa aaatggtgag ggagattcaa      120
ctagatagaa gaaaacatac ccaaaatgtc gccttaacaa taaagttgag cgcaattaaa      180
acgaagtatt tgcgtgaatt aaatgtttta gaagagaagt cgaaagatga gttgccgtca      240
gaaataaaaag caaagttaga cgcagctttt gagaagttta aaaaagatac attgaaacca      300
ggagaaaaagg tagcagaagc taagaagaag gttgaagaag ctaagaaaaa agccgaggat      360
caaaaagaag aagatcgtcg taactaccca accaatactt acaaaacgct tgaacttgaa      420
attgctgagt tcgatgtgaa agttaaagaa gcggagcttg aactagtaaa agaggaagct      480
aaagaatctc gaaacgaggg cacaattaag caagcaaaag agaaagttga gagtaaaaaa      540
gctgaggcta caaggttaga aaacatcaag acagatcgta aaaaagcaga agaagaagct      600
aaacgaaaaag cagcagaaga agataaagtt aagaaaaaac cagctgaaca accacaacca      660
gcgccgggcta ctcaaccaga aaaaccagct ccaaaaccag agaagccagc tgaacaacca      720
aaagcagaaa aaacagatga tcaacaagct gaaga                                755
    
```

<210> 22
 <211> 1239
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: cDNA from Streptococcus pneumoniae

<400> 22

```

acagagaagg aggtaactac ccaagtaccc acttattcta atatggcaaa gacagaacat      60
aggaaagctg ctaaacaagt cgtc gatgaa tatatagaaa aaatggtgag ggagattcaa      120
ttagatagaa gaaaacatac ccaaaatttc gccttcaaca tgaagttgag cgcaattaaa      180
acggaggtatt tgtatggatt aaaagagaag tcggaagctg agttgccgtc agaagtaaaa      240
gcaaaagttag acgcagcttt tgagcagttt aaaaaagata cattgaaact aggagaaaag      300
gtagcagaag ctgagaagaa gggtgcagaa gctgagaaaa aagccaaggc tcaaaaagaa      360
taagatcgcc gtaactaccc aaccaatact tacaaaacgc ttgaacttga aattgctgag      420
tccgatgtgg aagttaaaaa agcggagctt gaactattga aagaggaagc taaaactcga      480
aacgagtgaca caattaacca agcaaaagcg aaagttgaga gtaaaaaagc tgaggctaca      540
aagttagaag aaatcaagac agatcgtaaa aaagcagaag aagaagctaa acgaaaagca      600
gaagcagaag aagataaagt taaagataaa ctaaagaggc ggacaaaacg agcagttcct      660
ggagagccag caacacctga taaaaaagaa aatgatgcca agtcttcaga ttctagcgta      720
ggtgaagaaa ctcttccaag cccatccctg aatcaggaa aaaaggtagc agaagctgag      780
aagaagggtg cagaagctga gaaaaaagcc aaggatcaaa agaagaaga tcgccgtaac      840
taccacaacca atacttacia aacgcttgac cttgaaattg ctgagtcgga tgtgaaagtt      900
aaagaagcgg agcttgaact agtaaaagag gaagctaagg gatctcgaaa cgaggaaaaa      960
attaaccaag caaaagcggg agttgagagt aaaaaagctg aggctacaag gctagaaaaa     1020
atcaagacag atcgtaaaaa agcagaagaa gaagctaac gaaaagcagc agaagaagat     1080
aaagttaaag aaaaaccagc tgaacaacca caaccagcgc cggctcctca accagaaaaa     1140
ccaactgaag agcctgagaa tccagctcca gctccaaaac cagagaagcc agctgaacaa     1200
ccaaaagcag aaaaaacaga tgatcaacaa gctgaagaa                                1239
    
```

<210> 23
 <211> 1338
 <212> DNA
 <213> Artificial Sequence

<220>
 <213> Description: cDNA from Streptococcus pneumoniae

<400> 23

acagagaacg	agggagctac	ccaagtaccc	acttcttcta	atagggcaaa	tgaaagtcat	60
gcagaacaag	gagaacaacc	taaaaaactc	gattcagaac	gagataaggc	aaggaaagag	120
gtcgaggaat	atgtaaaaaa	aatagtgggt	gagagctatg	caaaatcaac	taaaaagcga	180
catacaatta	ctgtagctct	agttaacgag	ttgaacaaca	ttaagaacga	gtatttgaat	240
aaaatagttg	aatcaacctc	agaaagccaa	ctacagatac	tgatgatgga	gagtcgatca	300
aaagttagtg	aagctgtgtc	taagtttgaa	aaggactcat	cttcttcgtc	aagttcagac	360
tcttccacta	aaccggaagc	ttcagatata	gcgaagccaa	acaagccgac	agaaccagga	420
gaaaaggtag	cagaagctaa	gaagaagggt	gaagaagttg	agaaaaaagc	caaggatcaa	480
aaagaagaag	atcgtcgtaa	ctaccaaac	aattacttac	aaacgcttga	acttgaaatt	540
gctgagtccg	atgtggaagt	taaaaaagcg	gagcttgaac	tagtaaaagt	gaaagctaac	600
gaacctcgag	acaagcaaaa	aattaagcaa	gcagaagcgg	aagttgagag	taaacaagct	660
gaggctacaa	ggttaaaaaa	aatcaagaca	gatcgtgaag	aagcagaaga	agaagctaaa	720
cgaaagagcag	atgctaaaga	gcaaggtaaa	ccaaaggggc	ggccaaaacg	aggagtctct	780
ggagagctag	caacacctga	taaaaaagaa	aatgatgcca	agtcttcaga	ttctagcgtg	840
ggtgaagaaa	ctcttccaag	cccattcctg	aaaccagaaa	aaaaggtagc	agaagctgag	900
aagaagttg	aagaagctaa	gaaaaaagcc	gaggatcaaa	aagaagaaga	tcgccgtaac	960
tacccaacca	tacttataca	aacgcttgaa	cttgaattg	ctgagtcgga	tgtggaagt	1020
aaaaaagcgg	agcttgaact	agtaaaagag	gaagctaagg	aacctcgaaa	cgaggaaaaa	1080
gttaagcaag	caaaagcggg	agttgagagt	aaaaaagctg	aggctacaag	gttagaaaaa	1140
atcaagacag	atcgtaaaaa	agcagaagaa	gaagctaaac	gaaaagcagc	agaagaagat	1200
aaagttaaag	aaaaaccagc	tgaacaacca	caaccagcgc	cggctccaaa	aacagaaaaa	1260
ccagctccag	ctccaaaacc	agagaatcca	gctgaacaac	caaaagcaga	aaaaccagct	1320
gatcaacaag	ctgaagaa					1338

<210> 24

<211> 1284

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: cDNA prepared from Streptococcus pneumoniae genome

<400> 24

gaagggggtta	gaagtgggaa	taactccacg	gttacatcta	gtgggcaaga	tatatcgaag	60
aagtatgctg	atgaagtcga	gtcgcatact	caaagtatat	tgaaggatgt	caataaaaaat	120
ttgaagaaaag	ttcaacatac	ccaaaatgcc	gacttcaaca	aaaagttgag	caaaattaaa	180
acgaagtatt	tgtatgaatt	aaatgtttta	gaagagaagt	cggaagctga	gttgacgtca	240
aaaacaaaag	aaacaaaaga	agagtttaac	gcagcttttg	agcagtttaa	aaaagataca	300
ttatcaacag	aaccagaaaa	aaaggtagca	gaagctaaga	agaaggttga	agaagctaag	360
aaaaaagccg	aggatcaaaa	agaaaaagat	cgccgtaact	acccaacat	tacttataaaa	420
acgcttgaac	ttgaaattgc	tgagtccgat	gtggaagtta	aaaaagcggg	gcttgaacta	480
gtaaaagtga	aagctaacga	acctcgagac	gaggaaaaaa	ttaagcaagc	agaagcgaaa	540
gttgagagta	aacaagctga	ggctacaagg	ttaaaaaaaa	tcaagacaga	tcgtgaacaa	600
gctgaggcta	caaggttaga	aaacatcaag	acagatcgtg	aacaagcaga	agaagaagct	660
aaagttaaag	atgaaccaa	gaagcggaca	aaacgaggag	ttcttgagga	gccagcaaca	720
cctgataaaa	aagaaaatga	tgcaaggtct	tcagattcta	gcgtaggtga	agaaactctt	780
ccaagcccat	ccctgaaacc	agaaaaaaag	gttgcagaag	ctgagaagaa	ggttgaagaa	840
gctaagaaaa	aagccgagga	tcaaaaagaa	gaagatcgtc	gtaactacc	aaccaatact	900
tacaaaacgc	ttgaacttga	aattgctgag	tccgatgtgg	aagttaaaaa	agcggagctt	960
gaactagtaa	aaaggaagc	taaggaacct	cgaaacgagg	aaaaagttaa	gcaagcaaaa	1020
gcggaagtgg	agagtaaca	agctgaggct	acaaggttag	aaaacatcaa	gacagatcgt	1080
aaaaaagcag	aagaagaagc	taaacgaaaa	gcagcagaag	aagataaagt	taaagaaaaa	1140
ccagctgaac	aaccacaacc	agcgcggct	cctcaaccag	aaaaaccagc	tccaaaacca	1200
gaaaaaccag	ctccagctcc	aaaaccagag	aatccagctg	aacaaccaa	agcagaaaaa	1260
ccagctgatc	aacaagctga	agaa				1284

<210> 25

<211> 658

<212> DNA
 <213> Artificial Sequence

<220>
 <213> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 25
 gaagggggtta gaagtgggaa taactccacg gttacatcta gtgggcaaga tatatcgaag 60
 aagtatgctg atgaagtcca gtcgcatcta caaagtatat tgaaggatgt caataaaaaat 120
 ttgaaaaaag ttcaacatac ccaaaatgcc gacttcaaca aaaagttgag caaaattaaa 180
 ccgaagtatt tgtatgaatt aaagtgttta gaagagaagt cggaaagctga gttgacgtca 240
 aaaccaaaaga acaaaagaag agttaccgca gcttttgagc agtttaaaaa agatacatta 300
 tcaacagaac cagaaaaaaa ggtagcagaa gctaagaaga aggttgaaga agctaagaaa 360
 aaagccgagg atcaaaaaga aaaagatcgc cgtaactacc caaccattac ttacaaaacg 420
 cttgaacttg aaattgctga gtccgatgtg gaagttaaaa aagcggagct tgaactagta 480
 aaagaggaag ctaaggaacc tcgaaacgag gaaaaagtta agcaagcaaa agcgggaagtt 540
 gagagtaaac aagctgaggc tacaaggtta gaaaaaatca agacagatcg taaaaaagca 600
 gaagaagaag ctaaacgaaa agcagcagaa gaagataaag ttaaagaaaa accagctg 658

<210> 26
 <211> 1338
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 26
 acagagaacg agggagctac ccaagtaccc acttcttcta atagggcaaa tgaaagtcag 60
 gcagaacaag gagaacaacc taaaaaactc gattcagaac gagataaggc aaggaaagag 120
 gtcgaggaat atgtaaaaaa aatagtgggt gagagctatg caaaatcaac taaaaagcga 180
 catacaatta ctgtagctct agttaacgag ttgaacaaca ttaagaacga gtatttgaat 240
 aaaatagttg aatcaacctc agaaagccaa ctacagatac tgatgatgga gagtcatca 300
 aaagtagatg aagctgtgtc taagtttgaa aaggactcat cttcttcgtc aagttcagac 360
 tcttccacta aaccggaagc ttcagatata gcgaagccaa acaagccgac agaaccagga 420
 gaaaaggtag cagaagctaa gaagaagggt gaagaagctg agaaaaaagc caaggatcaa 480
 aaagaagaag atcgtcgtaa ctaccacaacc attacttaca aaacgcttga acttgaatt 540
 gctgagtcg atgtggaagt taaaaaagcg gagcttgaac tagtaaaagt gaaagctaac 600
 gaacctcgag acgagcaaaa aattaagcaa gcagaagcgg aagttgagag taaacaagct 660
 gaggctacaa ggttaaaaaa aatcaagaca gatcgtgaag aagcagaaga agaagctaaa 720
 cgaagagcag atgctaaaaga gcaaggtaaa ccaaaggggc gggcaaaacg aggagttcct 780
 ggagagctag caacacctga taaaaaagaa aatgatgcga agtcttcaga ttctagcgta 840
 ggtgaagaaa ctcttccaag cccatccctg aaaccagaaa aaaaggtagc agaagctgag 900
 aagaaggttg aagaagctaa gaaaaaagcc gaggatcaaa aagaagaaga tcgccgtaac 960
 taccacaacca atacttaca aacgcttgaa cttgaaattg ctgagtcgga tgtggaagtt 1020
 aaaaaagcgg agcttgaact agtaaaagag gaagctaagg aacctcgaaa cgaggaaaaa 1080
 gttaagcaag caaaagcggg agttgagagt aaaaaagctg aggctacaag gttagaaaaa 1140
 atcaagacag atcgtaaaaa agcagaagaa gaagctaaac gaaaagcagc agaagaagat 1200
 aaagttaaag aaaaaccagc tgaacaacca caaccagcgc cggctccaaa agcagaaaaa 1260
 ccagctccag ctccaaaacc agagaatcca gctgaacaac caaaagcaga aaaaccagct 1320
 gatcaacaag ctgaagaa 1338

<210> 27
 <211> 1242
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: cDNA derived from genome of Streptococcus pneumoniae

```

<400> 27
acagaaaaacg aaggaagtac ccaagcagcc acttcttcta atatggcaaa gacagaacat      60
aggaaagctg ctaaacaagt cgtcogatgaa tatatagaaa aaatggtgag ggagattcaa      120
ctagatagaa gaaaacatac ccaaaatgtc gccttaaaca taaagttgag cgcaattaaa      180
acgaagtatt tgcgtgaatt aaatgtttta gaagagaagt cgaaagatga gttgccgtca      240
gaaataaaag caaagttaga cgcagctttt gagaagttta aaaaagatac attgaaacca      300
ggagaaaaggg tagcagaagc taagaagaag gttgaagaag ctaagaaaaa agccgaggat      360
caaaaagaag aagatcgtcg taactacca accaatactt acaaaacgct tgaacttgaa      420
attgctgagt tcgatgtgaa agttaaagaa gcgagcttg aactagtaaa agaggaagct      480
aaagaatctc gaaacgaggg cacaattaag caagcaaaag agaaagttga gagtaaaaaa      540
gctgaggcta caaggttaga aaacatcaag acagatcgta aaaaagcaga agaagaagct      600
aaacgaaaag cagatgctaa gttgaaggaa gctaattgtag cgacttcaga tcaaggtaaa      660
ccaagggggc gggcaaaacg aggagttcct ggagagctag caacacctga taaaaaagaa      720
aatgatgcga agtcttcaga ttctagcgta ggtgaagaaa ctcttccaag ctcacccctg      780
aatcaggaa aaaaggtagc agaagctgag aagaaggttg aagaagctga gaaaaaagcc      840
aaggatcaaa aagaagaaga tcgccgtaac tacccaacca atacttaca aacgcttgac      900
cttgaaattg ctgagtcgga tgtgaaagtt aaagaagcgg agcttgaact agtaaaagag      960
gaagctaagg aacctcgaga cgaggaaaaa attaagcaag caaaagcga agttgagagt     1020
aaaaaagctg aggctacaag gttagaaaa acatcaagcag atcgtaaaaa agcaacaaga     1080
gaagctaaac gaaaagcagc agaagaagat aaagttaaag aaaaaccagc tgaacaacca     1140
caaccagcgc cggctactca accagaaaa ccagctccaa aaccagagaa gccagctgaa     1200
caaccaaag cagaaaaaac agatgatcaa caagctgaag aa                               1242

```

<210> 28

<211> 1275

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 28

```

acagagaagg aggtaactac ccaagtagcc acttcttcta atagggcaaa tgaaagtcag      60
gcaggacata ggaaagctgc tgaacaattc gatgaatata taaaaacaat gatccaatta      120
gatagaagaa aacataccca aaatttcgcc ttaaacataa agttgagcag aattaaaacg      180
gagtattttgc gtaaattaaa tgttttagaa gagaagtcga aagctgagtt gccgtcagaa      240
acaaaaaaag agatagacgc agcttttgag cagtttaaaa aagataccaa cagaacccaa      300
aaaacggtag cagaagctga gaagaaggtt gaagaagcta agaaaaaagc caaggctcaa      360
aaagaagaag atcaccgtaa ctaccaacc aatacttaca aaacgcttga acttgaaatt      420
gctgagtcgg atgtggaagt taaaaaagcg gagcttgaac tagtaaaaga ggaagctaag      480
gaatctcgag acgatgaaaa aattaagcaa gcagaagcga aagttgagag taaaaaagct      540
gaggctacaa ggttagaaaa catcaagaca gatcgtgaaa aagcagaaga agaagctaaa      600
cgaagagcag aagctaagtt gaaggaagct gttgaaaaga atgtagcgac ttcagagcaa      660
gataaaccaa aggggaggag aaaacgagga gttcctggag agcaagcaac acctgataaa      720
aaagaaaatg atgcaagtc ttcagattct agcgtaggtg aagaagctct tccaagccca      780
tcctgaaac cagaaaaaaa ggttgcagaa gctgagaaga aggttgcaga agctgagaaa      840
aaagccaagg ctcaaaaaga agaagatcgc cgtaactacc caaccaatac ttacaaaacg      900
cttgaacttg aaattgctga gtccgatgtg aaagttaaag aagcggagct tgaactagta      960
aaagaggaag ctaaggaatc tcgaaacgag gaaaaagtta atcaagcaaa agcgaaagtt     1020
gagagtaaaa aagctgaggg tacaaggtta gaaaaaatca agacagatcg taaaaaagca     1080
gaagaagaag ctaaacgaaa agcagcagaa gaagataaag ttaaagaaaa accagctgaa     1140
caaccacaac cagcgcgggc tcctcaacca gaaaaaccaa ctgaagagcc tgagaatcca     1200
gctcccgcac caaaaccaga gaagccagct gaacaaccaa aagcagaaaa aacagatgat     1260
caacaagctg aagaa

```

<210> 29

<211> 1278

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 29
 acagagaagg aggtaactac ccaagtagcc acttcttcta ataaggcaaa taaaagtcag 60
 acagaacata tgaaagctgc taacaagtc gatgaatata taaaaaaaaa gctccaatta 120
 gatagaagaa aacataccca aaatgtcggc ttactcacia agttgggctg aattaaaacg 180
 gagtatttgc atggattaag tgtttcaaaa aagaagtcgg aagctgagtt gccgtcagaa 240
 ataaaagcaa agttagacgc agcttttgag cagtttaaaa aagatacatt accaacagaa 300
 ccaggaaaaa aggtagcaga agctgagaag aaggttgaag aagctaagaa aaaagccgag 360
 gatcaaaaag aaaaagatct ccgtaactac ccaaccaata cttacaaaac gcttgaactt 420
 gacattgctg agtccgatgt ggaagttaaa aaagcggagc ttgaactagt aaaagaggaa 480
 gctaaggaat ctcgagacga gaaaaaaatt aatcaagcaa aagcgaagt tgagaataaa 540
 aaagctgagg ctacaagggt aaaaaacatc aagacagatc gtgaaaaagc agaagaagct 600
 aaacgaagag cagatgctaa gttgcaggaa gctaattgtag cgacttcaga gcaagataaa 660
 tcaaagaggc gggcaaaacg agaagttctt ggagagctag caacacctga taaaaaagaa 720
 aatgatgcga agtcttcaga ttctagcgtg ggtgaagaaa ctcttacaag cccatccctg 780
 aaaccagaaa aaaaggtagc agaagctgag aagaaggttg aagaagctaa gaaaaaagcc 840
 gaggatcaaaa aagaagaaga tcgtcgtaac tacccaacca atacttacia aacgcttgaa 900
 cttgaaattg ctgagtcgca tgtggaagtt aaaaaagcgg agcttgaact agtaaaagag 960
 gaagctaagg aatctcgaaa cgaggaaaaa attaagcaag taaaagcgaa agttgagagt 1020
 aaaaaagctg aggctacaag gctagaaaac atcaagacag atcgtaaaaa agcagaagaa 1080
 gaagaagcta aacgaagagc agcagaagaa gataaagtta aagaaaaacc agctgaacaa 1140
 ccacaaccag cgccggctcc tcaaccagaa aaaccaactg aagagcctga gaatccagct 1200
 ccagctccag ctccaaaacc agagaatcca gctgaaaaac caaagcaga aaagccagct 1260
 gatcaacaag ctgaagaa 1278

<210> 30
 <211> 1276
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 30
 acagagaagg aggtaactac ccaagtagcc acttcttcta ataaggcaaa taaaagtcag 60
 acagaacata tgaaagctgc taacaagtc gatgaatata taaaaaaaaa gctccaatta 120
 gatagaagaa aacataccca aaatgtcggc ttactcacia agttgggctg aattaaaacg 180
 gagtatttgc atggattaag tgtttcaaaa aagaagtcgg aagctgagtt gccgtcagaa 240
 ataaaagcaa agttagacgc agcttttgag cagtttaaaa aagatacatt accaacagaa 300
 ccaggaaaaa aggtagcaga agctgagaag aaggttgaag aagctaagaa aaaagccgag 360
 gatcaaaaag aaaaagatct ccgtaactac ccaaccaata cttacaaaac gcttgaactt 420
 gacattgctg agtccgatgt ggaagttaaa aaagcggagc ttgaactagt aaaagaggaa 480
 gctaaggaat ctcgagacga gaaaaaaatt aatcaagcaa aagcgaagt tgagaataaa 540
 aaagctgagg ctacaagggt aaaaaacatc aagacagatc gtgaaaaagc agaagaagct 600
 aaacgaagag cagatgctaa gttgcaggaa gctaattgtag cgacttcaga gcaagataaa 660
 tcaaagaggc gggcaaaacg agaagttttt ggagagctag caacacctga taaaaaagaa 720
 aatgatgcga agtcttcaga ttctagcgtg ggtgaagaaa ctcttacaag cccatccctg 780
 aaaccagaaa aaaaggtagc agaagctgag aagaaggttg aagaagctaa gaaaaaagcc 840
 gaggatcaaaa aagaagaaga tcgtcgtaac tacccaacca atacttacia aacgcttgaa 900
 cttgaaattg ctgagtcgca tgtggaagtt aaaaaagcgg agcttgaact agtaaaagag 960
 gaagctaagg aatctcgaaa cgaggaaaaa attaagcaag taaaagcgaa agttgagagt 1020
 aaaaaagctg aggctacaag gctagaaaac atcaagacag atcgtaaaaa agcagaagaa 1080
 gaagaagcta aacgaagagc agcagaagaa gataaagtta aagaaaaacc agctgaacaa 1140
 ccacaaccag cgccggctcc tcaaccagaa aaaccaactg aagagcctga gaatccagct 1200
 ccagctccag ctccaaaacc agagaatcca gctgaaaaac caaagcaga aaagccagct 1260
 gatcaacaag ctgaag 1278

<210> 31
 <211> 1272
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description: cDNA derived from genome of Streptococcus pneumoniae

<400> 31

```

acagagaagg aggtaactac ccaagtagcc acttcttcta atagggc aaa taaaagtcag      60
acagaacata tgaagctgc taaacaagtc gatgaatata taaaaaaaaa gctccaatta     120
gatagaagaa aacataccca aaatgtcggc ttactcacia agttggggcgt aattaaacg      180
gagtatttgc atggattaag tgtttcaaaa aagaagtcgg aagctgagtt gccgtcagaa     240
ataaaaagcaa agttagacgc agcttttgag cagtttaaaa aagatacatt accaacagaa     300
ccaggtaaaa aggtagcaga agctgagaag aaggttgaag aagctaagaa aaaagccgag     360
gatcaaaaag aaaaagatct ccgtaactac ccaaccaata cttacaaaac gcttgaactt     420
gacattgctg agtccgatgt ggaagttaa aaagcggagc ttgaactagt aaaagaggaa     480
gctaaggaat ctcgagacga gaaaaaatt aatcaagcaa aagcgaaggt tgagaataaa     540
aaagctgagg ctacaagggt aaaaaacatc aagacagatc gtgaaaaagc agaagaagct     600
aaacgaagag cagatgctaa gttgcaggaa gctaattgtag cgacttcaga gcaagataaa     660
tcaaaagagg gggcaaaaac agaagttctt ggagagctag caacacctga taaaaaagaa     720
aatgatgcga agtcttcaga ttctagcgtg ggtgaagaaa ctcttacaag cccatccctg     780
aaaccagaaa aagaagtagc agaagctgag aagaaggttg aagaagctaa gaaaaagcc     840
gaggatcaaa aagaagaaga tcgtcgtaac tacccaacca atacttaca aacgcttgaa     900
cttgaaattg ctgagtcgga tgtggaagtt aaaaaagcgg agcttgaact agtaaaaagag     960
gaagctaagg aatctcgaaa cgaggaaaaa attaagcaag taaaagcgaa agttgagagt    1020
aaaaaagctg aggctacaag gctagaaaac atcaagacag atcgtaaaaa agcagaagaa    1080
gaagaagcta aacgaagagc agcagaagaa gataaagtta aagaaaaacc agctgaacaa    1140
ccacaaccag cgccggctcc tcaaccagaa aaaccaactg aagagcctga gaatccagct    1200
ccagctccag ctccaaaacc agagaatcca gctgaaaaac caaaagcaga aaagccagct    1260
gatcaacaag ct                                     1272

```

<210> 32

<211> 1258

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: Coding strand of cDNA derived from genome of Streptococcus pneumoniae

<400> 32

```

acagagaacg agagaactac ccaagtagcc acttcttcta ataggggaaa gccagaacgt      60
aggaaagctg ctgaacaatt cgatgaatat ataaacaaaa tgatccaatt agataaaaaga     120
aaacataccc aaaatttagc cttcaacata cagttgagca gaattaaac ggagtatttg      180
aatggattaa aagagaagtc ggaagctgag ttgccgtcaa aaataaaaagc agagttagac     240
gcagctttta agcagtttaa aaaagataca ttaccaacag aaccagaaaa aaaagtagca     300
gaagctgaga agaaggttga agaagctgag aagaaggtag cagaagctaa gaaaaaagcc     360
aaggctcaaa aagaagaaga tcaccgtaac tacccaacca ttacttaca aacgcttgac     420
cttgaaattg ctgagttcga tgtgaaagtt aaagaagcgg agcttgaact agtaaaaaag     480
gaagctgacg aatctcgaaa cgagggcaca attaaccaag caaaagcgaa agttgagagt     540
gaaaaagctg aggctacaag gttaaaaaaa atcaagacag atcgtgaaaa agcagaagaa     600
gaagaagcta aacgaagagc agatgctaaa gagcaagatg aatcaaagag gcgaaagagt     660
cggggaaaaa caggagctct tggagagcaa gcaacacctg ataaaaaaga aaatgatgag     720
aagtcttcag attctagcgt aggtgaagaa actcttcaa gccatccct gaaccagga     780
aaaaaggtag cagaaggtg gaagaaggtt gaagaagctg ataaaaaagc caaggctcaa     840
aaagaagaag atcgccgtaa ctaccaacc aatacttaca aaacgcttga acttgaaatt     900
gctgagtcg atgtgaaagt taaagaagcg gagcttgaac tagtaaaaaga ggaagctaag     960
gaatctcgaa acgaggaaaa aattaagcaa gcaaaagcga aagttgagag taaaaaagct    1020
gaggctacaa ggtagaaaa aatcaagaca gatcgtaaaa aagcagaaga agaagctaaa    1080
cgaaaagcag cagaagaaga taaagttaaa gaaaaaccag ctgaacaacc acaaccagcg    1140
ccggctcctc aaccagaaaa accagctgaa gagcctgaga atccagttcc agctccaaaa    1200
ccagagaatc cagctgaaca accaaaagca gaaaaaccag ctgatcaaca agctgaag     1258

```

<210> 33
 <211> 1242
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: Coding strand of cDNA derived from genome of Streptococcus pneumoniae

<400> 33
 acagagaacg agggaagtac ccaagcagcc acttcttcta atatggcaaa gacagaacat 60
 aggaaagctg ctaaacaagt cgtc gatgaa tatatagaaa aaatggtgag ggagattcaa 120
 ctagatagaa gaaaacatac ccaaaatgtc gccttaaaca taaagttgag cgcaattaaa 180
 acgaagtatt tgcgtgaatt taatgtttta gaagagaagt cgaaggatga gttgccgtca 240
 gaaataaaag caaagttaga cgcagctttt gagaagttta aaaaagatac attgaaacca 300
 ggagaaaagg tagcagaagc taagaagaag gttgaagaag ctaagaaaaa agccgaggat 360
 caaaaagaag aagatcgtcg taactaccca accaatactt acaaaacgct tgaacttgaa 420
 attgctgagt tcgatgtgaa agttaaagaa gcggagcttg aactagttaa agaggaagct 480
 aaagaatctc gaaacgaggg cacaattaag caagcaaaag agaaagttga gagtaaaaaa 540
 gctgaggcta caaggttaga aaacatcaag acagatcgta aaaaagcaga agaagaagct 600
 aaacgaaaag cagatgctaa gttgaaggaa gctaagttag cgacttcaga tcaaggtaaa 660
 ccaaaggggc gggcaaaacg aggagttcct ggagagctag caacacctga taaaaagaa 720
 aatgatgcga agtcttcaga ttctagcgta ggtgaagaaa ctcttccaag ctcatccctg 780
 aatcaggaa aaaaggtagc agaagctgag aagaaggttg aagaagctga gaaaaagcc 840
 aaggatcaaa aagaagaaga tcgccgtaac tacccaacca atacttaca aacgcttgac 900
 cttgaaattg ctgagtcgga tgtgaaagtt aaagaagcgg agcttgaact agtaaaagag 960
 gaagctaagg aacctcgaga cgaggaaaaa attaagcaag caaaagcga agttgagagt 1020
 aaaaagctg aggtacaag gttagaaaac atcaagacag atcgtaaaaa agcagaagaa 1080
 gaagctaaac gaaaagcagc agaagaagat aaagttaaag aaaaaccagc tgaacaacca 1140
 caaccagcgc cggctactca accagaaaaa ccagctccaa aaccagagaa gccagctgaa 1200
 caaccaaaag cagaaaaaac agatgatcaa caagctgaag aa 1242

<210> 34
 <211> 1236
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description: Coding strand from cDNA derived from genome of Streptococcus pneumoniae

<400> 34
 gaaggggtta gaagtgagaa taaccccacg gttacatcta gtgggcaaga tatatcgaag 60
 aagtatgctg atgaagtcaa gtcacatcta gaaaaaatat tgagtgagat ccaacaacat 120
 ttagatagaa gtaaacatat caaaactgta aatctaatta acaaattgca agacattaag 180
 agaacgtatt tgtatgaatt aaatgtttta gaagataagt cgaaagctga gttgccgtca 240
 aaaataaaag cagagttaga cgcagctttt gagcagttta aaaaagatac attaccaaca 300
 gaaccaggaa aaaaggtagc agaagctaa gagaaggttg aagaagctga gaaaaagcc 360
 aaggctcaaa aagaagaaga ttaccgtaac tacccaacca ttacttaca aacgcttgaa 420
 cttgaaattg ctgagtcgga tgtgaaagtt aaagaagcgg agcttgaact agtaaaaaag 480
 gaagctagctg aatctcgaaa cgagggcaca attaaccaag caaaagcga agttgagagt 540
 gaacaagcagc aggtacaag gttaaaaaaa atcaagacag atcgtgaaaa agcagaagaa 600
 gaagctaaac gaagagcaga tgctaaagag caagatgaat caaagaggcg aaagagtcgg 660
 gtaaaacgag gagattttgg agagccagca acacctgata aaaaagaaaa tgatgcgaag 720
 tcttcagatt ctagcgtagg tgaagaaact cttccaagcc catccctgaa accaggaaaa 780
 aaggtagcag aagctgagaa gaaggttgaa gaagctgaga aaaaagccaa ggatcaaaaa 840
 gaagaagatc accgtaacta cccaaccatt acttcaaaa cgcttgaact tgaattgct 900
 gagtccgatg tggaagttaa aaaagcggag cttgaaactag taaaagagga agctaagggg 960
 tctcgaacg aggaaaaagt taagcaagca aaagcgggag ttgagagtaa aaaagctgag 1020

gctacaaggt	tagaaaaaat	caagacagat	cgtaaaaaag	cagaagaaga	agctaaacga	1080
aaagcagcag	aagaagataa	agttaaagaa	aaaccagctg	aacaaccaca	accagcgccg	1140
gctcctcaac	cagaaaaacc	agctccagct	ccaaaaccag	agaatccagc	tgaacaacca	1200
aaagcagaaa	aaccagctga	tcaacaagct	gaagaa			1236

<210> 35

<211> 1218

<212> DNA

<213> Artificial Sequence

<220>

<223> Description: Coding strand from cDNA derived from genome of Streptococcus pneumoniae

<400> 35

acagagaacg	agggaactac	ccaagcacc	acttcttcta	ataggggaaa	tgaaagtcag	60
gcagaacata	tgaaagctgc	taaacaagtc	gatgaatata	tagaaaaaat	gctccaatta	120
gatagaagaa	aacataccca	aaatgtcggc	ttactcacia	agttgggagc	aattaaaacg	180
gagtatttgc	gtggattaag	tgtttcaaaa	gagaagtcga	cagctgagtt	gccgtcagaa	240
ataaaagaaa	agttaaccgc	agcttttaag	cagtttaaaa	aagatacatt	gaaaccagaa	300
aaaaaggtag	cagaagctga	gaagaaggta	gcagaagcta	agaaaaaagc	cgaggatcaa	360
aaagaagaag	atcgtcgtaa	ctaccaacc	attacttaca	aaacgcttga	acttgaaatt	420
gctgagtcg	atgtggaagt	taaaaaagcg	gagcttgaac	tagtaaaagt	gaaagctaac	480
gaacctcgag	acgaggaaaa	aattaagcaa	gcagaagcgg	aagttgagag	taaaaaagct	540
gagggtacaa	ggttaaaaaa	aatcaagaca	gatcgtgaaa	aagcagaaga	agaagctaaa	600
cgaagagtag	atgctaaaga	gcaagatgaa	tcatacaaga	ggcgaagag	tcgggtaaaa	660
cgaggagatc	ttggagagca	agcaacacct	gataaaaaag	aaaatgatgc	gaagtcttca	720
gattctagcg	taggtgaaga	aactcttcca	agcccatccc	tgaaaccagg	aaaaaaggta	780
gcagaagctg	agaagaaggt	tgaagaagct	gataaaaaag	ccaaggctca	aaaagaagaa	840
gatcgccgta	actaccaaac	caatacttac	aaaacgcttg	aacttgaaat	tgctgagctc	900
gatgtggaag	ttaaaaaagc	ggagcttgaa	ctagtaaaag	aggaagctaa	ggaacctcga	960
aacgaggaaa	aagttaagca	agcaaaaagc	gaagttgaga	gtaaaaaagc	tgaggctaca	1020
aggttagaaa	aatcaagac	agatcgtaaa	aaagcagaag	aagaagctaa	acgaaaagca	1080
gcagaagaag	ataaagttaa	agaaaaacca	gctgaacaac	caaaaccagc	gccggctcct	1140
caaccagaaa	aaccagctcc	aaaaccagag	aatccagctg	aacaaccaa	agcagaaaaa	1200
ccagctgatc	aacaagct					1218

<210> 36

<211> 102

<212> PRT

<213> Artificial Sequence

<220>

<223> Description: Amino acid sequence derived from consensus cDNA sequence

<400> 36

Thr	Glu	Asn	Glu	Gly	Thr	Thr	Gln	Val	Ala	Thr	Ser	Ser	Asn	Arg	Ala
1				5					10					15	
Asn	Gln	Thr	Glu	His	Arg	Lys	Ala	Ala	Lys	Gln	Val	Val	Asp	Glu	Tyr
			20					25					30		
Ile	Lys	Lys	Met	Leu	Glu	Gln	Leu	Asp	Arg	Arg	Lys	His	Thr	Gln	Asn
		35					40					45			
Val	Ala	Leu	Asn	Ile	Lys	Leu	Ser	Ala	Ile	Lys	Thr	Glu	Tyr	Leu	Arg
	50					55				60					
Glu	Leu	Asn	Val	Leu	Glu	Glu	Lys	Ser	Lys	Ala	Glu	Leu	Pro	Ser	Glu

