

(19) **DANMARK**

(10) **DK/EP 3810283 T3**



Patent- og
Varemærkestyrelsen

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **A 61 P 35/00 (2006.01)** **A 61 K 31/437 (2006.01)** **C 07 D 471/04 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2023-08-21**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-06-14**
- (86) Europæisk ansøgning nr.: **19745335.0**
- (86) Europæisk indleveringsdag: **2019-06-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-04-28**
- (86) International ansøgning nr.: **US2019037492**
- (87) Internationalt publikationsnr.: **WO2019245974**
- (30) Prioritet: **2018-06-21 US 201862687930 P** **2018-08-20 US 201862719896 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **F. Hoffmann-La Roche AG, Grenzacherstrasse 124, 4070 Basel, Schweiz**
- (72) Opfinder: **CHUNG, Cheol Keun, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA**
XU, Jie, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
IDING, Hans, c/o F. Hoffmann-La Roche AG, Grenzacherstrasse 124, 4070 Basel, Schweiz
CLAGG, Kyle, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
DALZIEL, Michael, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
FETTES, Alec, c/o F. Hoffmann-La Roche AG, Grenzacherstrasse 124, 4070 Basel, Schweiz
GOSSELIN, Francis, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
LIM, Ngap-Kie, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
MCCLORY, Andrew, -, deceased, USA
ZHANG, Haiming, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
CHAKRAVARTY, Paroma, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
NAGAPUDI, Kartik, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
ROBINSON, Sarah, c/o Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Strandvejen 70, 2900 Hellerup, Danmark**
- (54) Benævnelse: **Faste former af tartratsalt af 3-((1R,3R)-1-(2,6-difluor-4-((1-(3-fluorpropyl)azetid-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2h-pyrido[3,4-b]indol-2-yl)-2,2-difluorpropan-1-ol, proces til fremstilling deraf og fremgangsmåder til anvendelse deraf til cancerbehandling**
- (56) Fremdragne publikationer:
WO-A1-2016/097072
WO-A1-2017/216280
SHEWALE ET AL: "Pharmaceutical cocrystals: design, development and characterisation", AMERICAN JOURNAL OF PHARMTECH RESEARCH, vol. 5, no. 4, 1 January 2015 (2015-01-01), pages 90-107,

Fortsættes ...

DESCRIPTION

SEQUENCE LISTING

[0001] This application incorporates by reference a Sequence Listing submitted with this application as text file entitled P34807-WO_SL.txt created on June 12, 2019 and having a size of 16.4 kilobytes.

FIELD OF THE INVENTION

[0002] Provided herein are solid forms of 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol, and methods of their use in the treatment of cancer. Further described herein are processes for preparing fused tricyclic compounds comprising a substituted phenyl or pyridinyl moiety.

BACKGROUND

[0003] Fused tricyclic compounds comprising a substituted phenyl or pyridinyl moiety within the scope of the present disclosure are useful as estrogen receptor ("ER") targeting agents.

[0004] The ER is a ligand-activated transcriptional regulatory protein that mediates induction of a variety of biological effects through its interaction with endogenous estrogens. Endogenous estrogens include 17β (beta)-estradiol and estrones. ER has been found to have two isoforms, ER- α (alpha) and ER- β (beta). Estrogens and estrogen receptors are implicated in a number of diseases or conditions, such as breast cancer, lung cancer, ovarian cancer, colon cancer, prostate cancer, endometrial cancer, uterine cancer, as well as others diseases or conditions. ER- α targeting agents have particular activity in the setting of metastatic disease and acquired resistance. ER- α targeting agents are disclosed in U.S. Publication Number 2016/0175289.

[0005] Useful processes for preparing fused tricyclic compounds comprising a substituted phenyl or pyridinyl moiety are disclosed in U.S. Publication Number 2016/0175289. However, there is a need for improved processes for preparing ER- α targeting agents.

[0006] There exists significant complexity surrounding the identification and selection of a solid form of a pharmaceutical compound. The differences in solid forms of such compounds affects both physical and chemical properties and may alter the processing, stability, bioavailability, formulation, and storage of pharmaceutical compounds. There is no reliable predictability of the solid form and its usefulness as a crystalline solid or amorphous solid. Crystalline solids may be considered useful, for example, for physical or chemical stability whereas amorphous

solids may be considered useful, for example, for enhanced dissolution and increased bioavailability.

[0007] Mixtures of crystalline materials arises from polymorphism. It is not possible to predict, *a priori*, if crystalline forms of a compound exist, much less whether crystalline forms can be prepared or isolated. Jones et al., 2006, Pharmaceutical Cocrystals: An Emerging Approach to Physical Property Enhancement," MRS Bulletin 37:875-879 (At present it is not generally possible to computationally predict the number of observable polymorphs of even the simplest molecules). The number of possible solid forms results in differing chemical and physical properties for a pharmaceutical compound and can greatly affect development, stability, and marketing of the product.

[0008] The estrogen receptor ("ER") is a ligand-activated transcriptional regulatory protein that mediates induction of a variety of biological effects through its interaction with endogenous estrogens. Endogenous estrogens include 17β (beta)-estradiol and estrones. ER has been found to have two isoforms, ER- α (alpha) and ER- β (beta). Estrogens and estrogen receptors are implicated in a number of diseases or conditions, such as breast cancer, lung cancer, ovarian cancer, colon cancer, prostate cancer, endometrial cancer, uterine cancer, as well as others diseases or conditions. There is a need for new ER- α targeting agents that have activity in the setting of metastatic disease and acquired resistance. Accordingly, there remains a need for cancer therapies having particular solid forms.

SUMMARY

[0009] Provided herein are solutions to the problems above and other problems in the art.

[0010] In one aspect provided herein is a compound, Compound B, having the name 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol tartrate, as described herein.

[0011] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern comprising peaks at 19.32, 20.26, 21.63, 23.28, or $24.81 \pm 0.1^\circ 2\theta$ ($\pm 0.1^\circ 2\theta$).

[0012] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern comprising peaks at 11.49, 12.54, 19.16, 19.42, or $24.67 \pm 0.1^\circ 2\theta$ ($\pm 0.1^\circ 2\theta$).

[0013] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern substantially as shown in FIG. 10 or FIG. 14.

[0014] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern comprising peaks at 11.31, 15.70, 16.54, 19.10, or $22.76 \pm 0.1^\circ 2\theta$.

[0015] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern comprising peaks at 12.52, 15.90, 19.66, 20.65, or $24.99 \pm 0.1^\circ 2\theta$.

[0016] In another aspect provided herein is a crystal form of Compound B having an X-ray powder diffraction pattern comprising peaks at 11.46, 12.51, 19.29, 19.42, or $20.23 \pm 0.1^\circ 2\theta$.

[0017] Also disclosed herein is an amorphous solid comprising Compound A.

[0018] Further provided herein are pharmaceutical compositions comprising Compound B or a crystal salt thereof. Such compounds and pharmaceutical compositions can be used in methods of treating cancer as set forth herein.

[0019] Also disclosed herein is a process for preparing a compound of formula (IV) or a salt thereof as set forth herein. The process comprises (1) reacting a reaction mixture comprising a compound of formula (I) as described herein, an organic solvent and thionyl chloride to form a compound of formula (IIa) as described herein and (2) reacting a reaction mixture comprising the compound of formula (IIa), a catalyst, an oxidant and a solvent to form a compound of formula (II) as described herein. The process further comprises reacting a reaction mixture comprising the compound of formula (II) and a compound of formula (III) as described herein in an organic solvent to form a compound of formula (IV) or a salt thereof as described herein.

[0020] Also disclosed herein is a process for preparing a compound of formula (VIII) or a pharmaceutically acceptable salt thereof as described herein. The process comprises reacting a reaction mixture comprising a compound of formula (IV) as described herein, a compound of formula (V) as described herein or a compound of formula (X) as described herein, and an organic solvent to form a compound of formula (VI) as described herein. The process further comprises reacting a reaction mixture comprising the compound of formula (VI), an organic solvent, and a compound of formula (VII) as described herein or a salt thereof to form a compound of formula (VIII) or a salt thereof.

[0021] Also disclosed herein is a process for preparing a compound of formula (VIII) or a pharmaceutically acceptable salt thereof as described herein. The process comprises reacting a reaction mixture comprising a compound of formula (IX) as described herein or a compound of formula (X) as described herein, a compound of formula (IV) as described herein and an organic solvent to form the compound of formula (VIII) or a salt thereof as described herein.

[0022] Also disclosed herein is a process for preparing a compound of formula (IX) or a salt thereof as described herein. The process comprises reacting a reaction mixture comprising a compound of formula (X) as described herein, a compound of formula (VII), or a salt thereof, as described herein an organic solvent and a catalyst to form a compound of formula (XI) or a salt thereof as described herein.

[0023] Also disclosed herein is a process for preparing a compound of formula (III) or a salt

thereof as described herein. The process comprises reacting a reaction mixture comprising a compound of formula (XII) as described herein, a compound B and an organic solvent to form the compound of formula (XIII) as described herein.

[0024] Also disclosed herein is a compound of formula (XVI) as described herein.

[0025] Also disclosed herein is a process for preparing a compound having formula (XX) is provided, where the process comprises contacting a compound of formula (XXI) as described herein with a protein transaminase to form a compound of formula (3). The compound of formula (3) is contacted with a compound of formula (II) as described herein to form compound formula (XX).

BRIEF DESCRIPTION OF THE FIGURES

[0026]

FIG. 1 depicts the XRPD pattern for Compound B Form A.

FIG. 2 depicts the TGA and DSC for Compound B Form A.

FIG. 3 depicts a PLM image for Compound B Form A.

FIG. 4 depicts the XRPD pattern for Compound B Form B.

FIG. 5 depicts the TGA and DSC for Compound B Form B.

FIG. 6 depicts the ^{13}C SSNMR for Compound B Form B.

FIG. 7 depicts the ^{19}F SSNMR for Compound B Form B.

FIG. 8 depicts the water sorption/desorption plot for Compound B Form B.

FIG. 9a depicts the SEM image; FIG. 9b depicts the PLM image for Compound B Form B; FIG. 9c depicts the particle size distribution (PSD) for Compound B Form B.

FIG. 10 depicts a comparative XRPD pattern for Compound B Form C against Form A and Form B of Compound B. Form C was found to be a mixture of Form A and Form B.

FIG. 11 depicts the TGA and DSC for Compound B Form C.

FIG. 12 depicts the XRPD pattern for Compound B Form D.

FIG. 13 depicts the TGA and DSC for Compound B Form D.

FIG. 14 depicts the XRPD pattern for Compound B Form E.

FIG. 15 depicts the TGA and DSC for Compound B Form E.

FIG. 16 depicts the XRPD pattern for Compound B Form F.

FIG. 17 depicts the DVS plot for Compound B Form F.

FIG. 18 depicts the DSC for Compound B Form F.

FIG. 19 depicts the XRPD pattern for Compound B Form G.

FIG. 20 depicts the TGA and DSC for Compound B Form G.

FIG. 21 depicts the XRPD pattern overlay for Compound B Form A, Form B, Form C, Form D, Form F, and Form G.

FIG. 22 depicts the XRPD for the compressibility properties for Compound B Form B.

FIG. 23 depicts the ¹⁹F SSNMR for the compressibility properties for Compound B Form B.

FIG. 24 depicts the DSC for the compressibility properties for Compound B Form B.

FIG. 25 depicts phase transformation pathways for Compound B Forms A, B, D, and F.

FIG. 26 depicts phase transformation pathways to obtain Compound B Form B from Form F.

FIG. 27a depicts the XRPD pattern for Compound C Form 1; FIG. 27b depicts the XRPD pattern for Compound C Form 2.

FIG. 28 depicts the TGA and DSC for Compound C Form 1.

FIG. 29 depicts a PLM image for Compound C Form 1.

FIG. 30 depicts the XRPD pattern for Compound D Form M.

FIG. 31 depicts the TGA and DSC for Compound D Form M.

FIG. 32 depicts the XRPD pattern for the amorphous form of Compound A.

FIG. 33 depicts the TGA and DSC for the amorphous form of Compound A.

FIG. 34 depicts cell viability of ER+ breast cancer cell line for Compound B compared to GDC-0810 and GDC-0927.

FIG. 35 depicts effect on tumor volume of Compound B at 0.1 mg/kg and 1 mg/kg compared to GDC-0927 at 100 mg/kg.

FIG. 36a depicts CT scanning and FIG. 36b depicts FES-PET scanning of a breast cancer patient treated with Compound B.

FIG. 37a depicts CT scanning and FIG. 37b depicts FES-PET scanning of a second breast cancer patient treated with Compound B.

DETAILED DESCRIPTION

[0027] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the invention belongs. See, e.g., Singleton et al., *DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY* 2nd ed., J. Wiley & Sons (New York, NY 1994); Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, Cold Springs Harbor Press (Cold Springs Harbor, NY 1989). Any methods, devices and materials similar or equivalent to those described herein can be used in the practice of this invention.

[0028] The following definitions are provided to facilitate understanding of certain terms used frequently herein and are not meant to limit the scope of the present disclosure.

[0029] As used herein, and unless otherwise specified, the terms "about" and "approximately," when referring to doses, amounts, or weight percents of ingredients of a composition or a dosage form, mean a dose, amount, or weight percent that is recognized by one of ordinary skill in the art to provide a pharmacological effect equivalent to that obtained from the specified dose, amount, or weight percent. The equivalent dose, amount, or weight percent can be within 30%, 20%, 15%, 10%, 5%, 1%, or less of the specified dose, amount, or weight percent.

[0030] As used herein, and unless otherwise specified, the terms "about" and "approximately," when referring to a numeric value or range of values used for characterization of a particular solid form described herein (e.g., XRPD peak values) indicate that the value or range of values may deviate from a given value to an extent deemed reasonable to one of ordinary skill in the art while still describing the solid form. In one embodiment, the value of an XRPD peak position may vary by up to $\pm 0.1^\circ 2\theta$ (or ± 0.05 degree 2θ) while still describing the particular XRPD peak.


[0031] As used herein, and unless otherwise specified, a crystalline that is "pure," *i.e.*, substantially free of other crystalline or amorphous solids or other chemical compounds, and contains less than about 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.05%, or 0.01% of one or more other solid forms on a weight basis. The detection of other solid forms can be accomplished by, for example, diffraction analysis, thermal analysis, elemental combustion analysis and/or spectroscopic analysis. The detection of other chemical compounds can be accomplished by, for example, mass spectrometry analysis, spectroscopic analysis, thermal analysis, elemental combustion analysis and/or chromatographic analysis.

[0032] Unless otherwise specified, the terms "solvate" and "solvated," as used herein, refer to a solid form of a substance which contains solvent. The terms "hydrate" and "hydrated" refer to a solvate wherein the solvent is water. The terms "solvate" and "solvated," as used herein, can also refer to a solvate of a salt, cocrystal, or molecular complex. The terms "hydrate" and

"hydrated," as used herein, can also refer to a hydrate of a salt, cocrystal, or molecular complex.

[0033] The term "pharmaceutically acceptable," refers to a diluent, excipient, or carrier in a formulation compatible with the other ingredient(s) of the formulation and not deleterious to the recipient thereof.

[0034] Compound A refers to the compound having the structure:



and having the name 3-(((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol, including a pharmaceutically acceptable salt thereof. Compound A can be a tartaric acid salt as described herein (e.g. Compound B). Compound A can be a fumaric acid salt as described herein (e.g. Compound C). Compound A can be a malonate salt as described herein (e.g. Compound D).

[0035] The term "solid form" refers to a physical form which is not predominantly in a liquid or a gaseous state. A solid form may be a crystalline form or a mixture thereof. In certain embodiments, a solid form may be a liquid crystal. In certain embodiments, the solid form of Compound A is Form A, Form B, Form C, Form D, Form E, Form F, Form G, Form 1, or Form 2, an amorphous solid, or a mixture thereof. In one embodiment, the solid form of Compound A is a tartrate salt. In another embodiment, the solid form of Compound A is a fumarate salt or a mixture thereof. A solid form may be a crystal form as defined herein.

[0036] The term "crystal form" or "crystalline form" refers to a solid form that is crystalline. In certain embodiments, a crystal form of a compound described herein may be substantially free of amorphous solids and/or other crystal forms. In certain embodiments, a crystal form of a compound described herein may contain less than about 1%, less than about 2%, less than about 3%, less than about 4%, less than about 5%, less than about 6%, less than about 7%, less than about 8%, less than about 9%, less than about 10%, less than about 15%, less than about 20%, less than about 25%, less than about 30%, less than about 35%, less than about 40%, less than about 45%, or less than about 50% by weight of one or more amorphous solids and/or other crystal forms. In certain embodiments, a crystal form described herein is pure. In certain embodiments, a crystal form of a compound described herein may be about 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, or 90% pure.

[0037] The term "amorphous" or "amorphous solid" refers to a solid form that not substantially crystalline as determined by X-ray diffraction. In particular, the term "amorphous solid" describes a disordered solid form, i.e., a solid form lacking long range crystalline order. In

certain embodiments, an amorphous solid of a compound described herein may be substantially free of other amorphous solids and/or crystal forms. In certain embodiments, an amorphous solid may be pure. In certain embodiments, an amorphous solid of a compound described herein may be about 99%, 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, or 90% pure.

[0038] "Treating" as used herein, means an alleviation, in whole or in part, of a disorder, disease or condition, or one or more of the symptoms associated with a disorder, disease, or condition, or slowing or halting of further progression or worsening of those symptoms, or alleviating or eradicating the cause(s) of the disorder, disease, or condition itself. In one embodiment, the disorder is a cancer.

[0039] The term "effective amount" or "therapeutically effective amount" refers to an amount of a compound described herein capable of treating or preventing a disorder, disease or condition, or symptoms thereof, disclosed herein.

[0040] "Patient" or "subject" is defined herein to include animals, such as mammals, including, but not limited to, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, monkeys, chickens, turkeys, quails, or guinea pigs and the like, in one embodiment a mammal, in another embodiment a human. In one embodiment, a subject is a human having or at risk for cancer.

[0041] As used herein, the terms "moiety" and "substituent" refer to an atom or group of chemically bonded atoms that is attached to another atom or molecule by one or more chemical bonds thereby forming part of a molecule.

[0042] As used herein, the term "alkyl" refers to an aliphatic straight-chain or branched-chain saturated hydrocarbon moiety having 1 to 20 carbon atoms. In particular embodiments the alkyl has 1 to 10 carbon atoms. In particular embodiments the alkyl has 1 to 6 carbon atoms. Alkyl groups may be optionally substituted independently with one or more substituents described herein.

[0043] As used herein, the term "substituted" refers to the replacement of at least one of hydrogen atom of a compound or moiety with another substituent or moiety. Examples of such substituents include, without limitation, halogen, -OH, -CN, oxo, alkoxy, alkyl, alkylene, aryl, heteroaryl, haloalkyl, haloalkoxy, cycloalkyl and heterocycle. For example, the term "haloalkyl" refers to the fact that one or more hydrogen atoms of an alkyl (as defined below) is replaced by one or more halogen atoms (e.g., trifluoromethyl, difluoromethyl, fluoromethyl, chloromethyl, etc.). In one embodiment, substituted as used herein can refer to replacement of at least one hydrogen atom of a compound or moiety described herein with halogen or alkyl.

[0044] As used herein, the term "alkylene" as used herein refers to a linear or branched saturated divalent hydrocarbon radical of one to twelve carbon atoms, and in another aspect one to six carbon atoms, wherein the alkylene radical may be optionally substituted

independently with one or more substituents described herein. Examples include, but are not limited to, methylene, ethylene, propylene, 2-methylpropylene, pentylene, and the like.

[0045] As used herein, the term "alkoxy" refers to a group of the formula -O-R', wherein R' is an alkyl group. Alkoxy groups may be optionally substituted independently with one or more substituents described herein. Examples of alkoxy moieties include methoxy, ethoxy, isopropoxy, and tert-butoxy.

[0046] As used herein, the term "aryl" refers to a cyclic aromatic hydrocarbon moiety having a mono-, bi- or tricyclic aromatic ring of 5 to 16 carbon ring atoms. Bicyclic aryl ring systems include fused bicyclics having two fused five-membered aryl rings (denoted as 5-5), having a five-membered aryl ring and a fused six-membered aryl ring (denoted as 5-6 and as 6-5), and having two fused six-membered aryl rings (denoted as 6-6). The aryl group can be optionally substituted as defined herein. Examples of aryl moieties include, but are not limited to, phenyl, naphthyl, phenanthryl, fluorenyl, indenyl, pentalenyl, azulenyl, and the like. The term "aryl" also includes partially hydrogenated derivatives of the cyclic aromatic hydrocarbon moiety provided that at least one ring of the cyclic aromatic hydrocarbon moiety is aromatic, each being optionally substituted.

[0047] As used herein, the term "heteroaryl" refers to an aromatic heterocyclic mono-, bi- or tricyclic ring system of 5 to 16 ring atoms, comprising 1, 2, 3 or 4 heteroatoms selected from N, O and S, the remaining ring atoms being carbon. In some embodiments, monocyclic heteroaryl rings may be 5-6 membered. Bicyclic heteroaryl ring systems include fused bicyclics having two fused five-membered heteroaryl rings (denoted as 5-5), having a five-membered heteroaryl ring and a fused six-membered heteroaryl ring (denoted as 5-6 and 6-5), and having two fused six-membered heteroaryl rings (denoted as 6-6). The heteroaryl group can be optionally substituted as defined herein. Examples of heteroaryl moieties include pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyridinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyrimidinyl, triazinyl, isoxazolyl, benzofuranyl, isothiazolyl, benzothienyl, benzothiophenyl, indolyl, aza-indolyl, isoindolyl, isobenzofuranyl, benzimidazolyl, benzoxazolyl, benzoisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzooxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, pyrrolopyridazinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, thienopyridazinyl, thienopyrimidinyl, thienopyrazinyl, furopyridazinyl, furopyrimidinyl, and furopyrazinyl.


[0048] As used herein, the terms "halo", "halogen" and "halide", which may be used interchangeably, refer to a substituent fluorine, chlorine, bromine, or iodine.

[0049] As used herein, the term "haloalkyl" refers to an alkyl group wherein one or more of the hydrogen atoms of the alkyl group has been replaced by the same or different halogen atoms, particularly fluorine and/or chlorine atoms. Examples of haloalkyl include monofluoro-, difluoro- or trifluoro-methyl, -ethyl or -propyl, for example 3,3,3-trifluoropropyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, fluoromethyl, difluoromethyl or trifluoromethyl.

[0050] As used herein, the term "hydroxyalkyl" refers to an alkyl group wherein one or more of the hydrogen atoms of the alkyl group have been replaced by a hydroxyl moiety. Examples include alcohols and diols

[0051] As used herein, the term "heteroalkyl" refers to a straight- or branched-chain alkyl as defined herein having from 2 to 14 carbons, from 2 to 10 carbons, or from 2 to 6 carbons in the chain, one or more of which has been replaced by a heteroatom selected from S, O, P and N. Non-limiting examples of heteroalkyls include alkyl ethers, secondary and tertiary alkyl amines, amides, and alkyl sulfides.

[0052] As used herein, the term "cycloalkyl" means a saturated or partially unsaturated carbocyclic moiety having mono-, bi- (including bridged bicyclic) or tricyclic rings and 3 to 10 carbon atoms in the ring. The cycloalkyl moiety can optionally be substituted with one or more substituents. In particular embodiments cycloalkyl contains from 3 to 8 carbon atoms (i.e., (C₃-C₈)cycloalkyl). In other particular embodiments cycloalkyl contains from 3 to 6 carbon atoms (i.e., (C₃-C₆)cycloalkyl). Examples of cycloalkyl moieties include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and partially unsaturated (cycloalkenyl) derivatives thereof (e.g. cyclopentenyl, cyclohexenyl, and cycloheptenyl), bicyclo[3.1.0]hexanyl, bicyclo[3.1.0]hexenyl, bicyclo[3.1.1]heptanyl, and bicyclo[3.1.1]heptenyl. The cycloalkyl moiety can be attached in a "spirocycloalkyl" fashion such as "spirocyclopropyl":



[0053] As used herein, the terms "heterocycle" or "heterocyclyl" refer to a 4, 5, 6 and 7-membered monocyclic, 7, 8, 9 and 10-membered bicyclic (including bridged bicyclic) or 10, 11, 12, 13, 14 and 15-membered bicyclic heterocyclic moiety that is saturated or partially unsaturated, and has one or more (e.g., 1, 2, 3 or 4 heteroatoms selected from oxygen, nitrogen and sulfur in the ring with the remaining ring atoms being carbon. In some embodiments, the heterocycle is a heterocycloalkyl. In particular embodiments heterocycle or heterocyclyl refers to a 4, 5, 6 or 7-membered heterocycle. When used in reference to a ring atom of a heterocycle, a nitrogen or sulfur may also be in an oxidized form, and a nitrogen may be substituted with one or more (C₁-C₆)alkyl or groups. The heterocycle can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. Any of the heterocycle ring atoms can be optionally substituted with one or more substituents described herein. Examples of such saturated or partially unsaturated heterocycles include, without limitation, tetrahydrofuranyl, tetrahydrothienyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, pyrrolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl. The term the term heterocycle also includes groups in which a heterocycle is fused to one or more aryl, heteroaryl, or cycloalkyl rings, such as indolinyl, 3H-indolyl, chromanyl, azabicyclo[2.2.1]heptanyl, azabicyclo[3.1.0]hexanyl, azabicyclo[3.1.1]heptanyl,

octahydroindolyl, or tetrahydroquinolinyl.

[0054] Unless otherwise indicated, the term "hydrogen" or "hydro" refers to the moiety of a hydrogen atom (-H) and not H₂.

[0055] As used herein, the term "organic solvent" refers to any non-aqueous polar aprotic solvent, polar protic solvent, and non-polar solvent.

[0056] As used herein, the term "polar organic solvent" refers to both polar aprotic solvents and polar protic solvents, excluding water.

[0057] As used herein, the term "polar aprotic solvent" refers to any polar solvent not having a proton-donating ability. Examples include, without any limitation, 2-methyltetrahydrofuran, tetrahydrofuran, ethyl acetate, propyl acetate (e.g., isopropyl acetate), acetone, dimethylsulfoxide, N,N-dimethylformamide, acetonitrile, N,N-dimethylacetamide, N-methylpyrrolidone, hexamethylphosphoramide, and propylene carbonate.

[0058] As used herein, the term "polar protic solvent" refers to any polar solvent having a proton-donating ability. Examples include, without limitation, methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, formic acid, nitromethane and acetic acid. An organic polar protic solvent excludes any effective amount of water.

[0059] As used herein, the term "non-polar solvent" refers to solvents that contain bonds between atoms with similar electronegativities, such as carbon and hydrogen, such that the electric charge on the molecule is evenly distributed. Non-polar solvents are characterized as having a low dielectric constant. Examples include, without limitation, pentane, hexane, heptane, cyclopentane, methyl *tert*-butyl ether (MTBE), diethyl ether, toluene, benzene, 1,4-dioxane, carbon tetrachloride, chloroform and dichloromethane (DCM). In some embodiments, the non-polar solvent has a dielectric constant of less than 2, examples of which include, without limitation, pentane, hexane and heptane. As compared to other non-polar solvents, DCM exhibits some degree of polarity at the bond level (i.e., between carbon and chlorine), but only a small degree of polarity at the molecular level due to symmetry-based cancellation of polarity.

[0060] As used herein, the term "anti-solvent" refers to a solvent in which the referenced compound is poorly soluble and which induces precipitation or crystallization of said compound from solution.

[0061] As used herein, the term "acid catalyst" refers to an acid catalyst such as, but not limited to, a Brønsted acid, a Lewis acid or a Brønsted-Lowry catalyst. Non-limiting examples of acid catalysts include acetic acid, glacial acetic acid, trifluoroacetic acid, benzoic acid, pivalic acid, diphenyl phosphoric acid, triflic acid, formic acid, tartaric acid, fumaric acid, malonic acid, salicylic acid, *p*-toluene sulfonic acid, sulfuric acid, hydrochloric acid, phosphoric acid, methanesulfonic acid, camphor sulfonic acid, naphthalene sulfonic acid, clay-based

montmorillonite K-10 and resin based amberlyst, and combinations thereof.

[0062] As used herein, the term "amine protecting group" refers to any known protecting group that that blocks or protects the functionality of amines. Amine protecting groups within the scope of the disclosure include, without limitation, 1-chloroethyl carbamate (ACD); 4-methoxybenzenesulfonamide; acetamide (Ac); benzylamine (Bn); benzyloxy carbamate (CBz); formamide; methyl carbamate; trifluoroacetamide; tert-butoxy carbamate (Boc); p-methoxybenzyl carbonyl (MeOZ); 9-fluorenylmethoxycarbonyl (FMOC); bezoyl (Bz); p-methoxybenzyl (PMB); 3,4-dimethoxybenzyl (DMPM); p-methoxyphenyl (PMP); Tosyl (Ts); and trichloroethyl chloroformate (Troc). For a description of amine protecting groups and their use, see P. G. M. Wuts and T. W. Greene, *Greene's Protective Groups in Organic Synthesis* 4th edition, Wiley-Interscience, New York, 2006

[0063] As used herein, the term "aldehyde protecting group" refers to any known substituent attached to an aldehyde group that blocks or protects the carbonyl group of the aldehyde functionality. Suitable protecting groups of the aldehyde functionality include, but are not limited to (a) cyclic acetals and ketals, (b) cyclic mono or di-thio acetals or ketals or other derivatives such as imines, hydrazones, cyanohydrin, oximes or semicarbazones, for example, dialkyl or diaryl acetals or 1,3 dithiane, (c) cyclic imines such as substituted methylene derivatives or N,N'-dimethylimidazolidine. Some non-limiting examples of aldehyde protecting groups include 1,3-dithiane, 1,3-dithiolane, diethyl acetal, dimethyl acetal, ethylene glycol acetal, neopentyl glycol acetal, trimethylsilyl cyanohydrin, and trialkyl orthoformates such as triethyl orthoformate. For a description of aldehyde protecting groups and their use, see, Wuts and Greene.

[0064] As used herein, "leaving group" refers to an atom or a group of atoms that is displaced in a chemical reaction as stable species. Suitable leaving groups are well known in the art, e.g., see, *March's Advanced Organic Chemistry*, 5th Ed., Ed.: Smith, M. B. and March, J., John Wiley & Sons, New York: 2001 and T. W. Greene, *Protective Groups in Organic Synthesis*, John Wiley & Sons, New York, 1991. Such leaving groups include, but are not limited to, halogen, alkoxy, sulphonyloxy, optionally substituted alkylsulphonyl, optionally substituted alkenylsulphonyl, optionally substituted arylsulphonyl, and diazonium moieties. Examples of some leaving groups include chloro, iodo, bromo, fluoro, methanesulfonyl (mesyl), tosyl, triflate, nitrophenylsulfonyl (nosyl), and bromo-phenylsulfonyl (brosyl).

[0065] "Transition metal catalysts" within the scope of the disclosure include, without limitation, palladium, platinum, gold, ruthenium, rhodium, and iridium catalysts. Non-limiting examples of suitable catalysts include: (2-Biphenyl)di-tert-butylphosphine gold(I) chloride ("JohnPhos"), 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl gold(I) chloride ("XPhos AuCl"), 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl gold(I) bis(trifluoromethanesulfonyl)imide ("XPhos AuNTf₂"), Chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2-aminoethyl)phenyl]palladium(II) ("XPhos Palladacycle"), Chloro(2-dicyclohexylphosphino-2',6'-dimethoxy-1,1'-biphenyl)[2-(2-aminoethyl)phenyl]palladium(II)-methyl-t-butyl ether adduct ("SPhos Palladacycle"), t-BuXPhos palladium(II) phenethylamine chloride ("tBuXPhos Pd G1"),

Chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) ("Xphos Pd G2"), Chloro(2-dicyclohexylphosphino-2',6'-dimethoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) ("SPhos Pd G2"), Chloro(2-dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) ("RuPhos Pd G2"), Chloro[(2-dicyclohexylphosphino-2',6'-bis(N,N-dimethylamino)-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)]palladium(II) ("CPhos-Pd-G2"), [(2-Dicyclohexylphosphino-2',6'-bis(N,N-dimethylamino)-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)] palladium(II) methanesulfonate ("CPhos-Pd-G3"), [(2-Di-tert-butylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)] palladium(II) methanesulfonate ("tBuXPhos-Pd-G3"), (2-Dicyclohexylphosphino-2',6'-diisopropoxy-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate ("RuPhos-Pd-G3"), (2-Dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate ("XPhos-Pd-G3"), [(2-Di-cyclohexylphosphino-3,6-dimethoxy-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate ("BrettPhos-Pd-G3"), [(2-{Bis[3,5-bis(trifluoromethyl)phenyl]phosphine-3,6-dimethoxy-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate ("JackiePhos-Pd-G3"), tert-butyl BrettPhos-Pd-G3, [tert-butyl BrettPhos-Pd (allyl)]OTf, and combinations thereof.

[0066] As used herein, "inorganic acids" refer to acids such as, but not limited to, hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, sulfamic acid, nitric acid, boric acid, phosphoric acid, and combinations thereof.

[0067] As used herein, "organic acids" refer to acids such as, but not limited to: acetic acid; trifluoroacetic acid; phenylacetic acid; propionic acid; stearic acid; lactic acid; ascorbic acid; maleic acid; hydroxymaleic acid; isethionic acid; succinic acid; valeric acid; fumaric acid; malonic acid; pyruvic acid; oxalic acid; glycolic acid; salicylic acid; oleic acid; palmitic acid; lauric acid; a pyranosidyl acid, such as glucuronic acid or galacturonic acid; an alpha-hydroxy acid, such as mandelic acid, citric acid, or tartaric acid; cysteine sulfinic acid; an amino acid, such as aspartic acid, glutaric acid or glutamic acid; an aromatic acid, such as benzoic acid, 2-acetoxybenzoic acid, naphthoic acid, or cinnamic acid; a sulfonic acid, such as laurylsulfonic acid, p-toluenesulfonic acid, methanesulfonic acid, benzenesulfonic acid or ethanesulfonic acid; cysteine sulfonic acid; and combinations thereof.

[0068] As used herein, "inorganic bases" refer to bases such as, but not limited to, sodium hydroxide, potassium hydroxide, lithium hydroxide, ammonium hydroxide, magnesium hydroxide, sodium carbonate, potassium carbonate, and combinations thereof.

[0069] As used herein, the term "organic base" refers to an organic compound containing one or more nitrogen atoms, and which acts as a base. Examples of organic bases include, but are not limited to, tertiary amine bases. Examples of organic bases include, but are not limited to, 1,8-Diazabicyclo[5.4.0]undec-7-ene ("DBU"), N-methyl-morpholine (NMM), diisopropylethylamine (DIPEA), triethylamine (TEA), a t-butoxide (e.g., sodium, potassium, calcium or magnesium *tert*-butoxide).

[0070] Compounds of the present disclosure may present in a salt form which encompasses pharmaceutically acceptable salts and non-pharmaceutically acceptable salts. As used herein, the term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. In addition to pharmaceutically acceptable salts, the compounds of the present disclosure may be in the form of non-pharmaceutically acceptable salts which can be useful as an intermediate for isolating or purifying said compounds.

[0071] Exemplary acid salts of the compounds of the present disclosure include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate "mesylate", ethanesulfonate, benzenesulfonate, p-toluenesulfonate, and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. A pharmaceutically acceptable salt may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counter ion. The counter ion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically acceptable salt may have more than one charged atom in its structure. Instances where multiple charged atoms are part of the pharmaceutically acceptable salt can have multiple counter ions. Hence, a pharmaceutically acceptable salt can have one or more charged atoms and/or one or more counter ion.

[0072] Exemplary base salts of the compounds of the present disclosure include, but are not limited to, inorganic salts formed from sodium, potassium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, and aluminum cations. Organic salts formed from cations including primary, secondary, and tertiary amines; substituted amines including naturally occurring substituted amines; cyclic amines; basic ion exchange resins; isopropylamine; trimethylamine; diethylamine; trimethylamine; tripropylamine; ethanolamine; 2-diethylaminoethanol; trimethylamine; dicyclohexylamine; lysine; arginine; histidine; caffeine; procaine; hydrabamine; choline; betaine; ethylenediamine; glucosamine; methylglucamine; theobromine; purines; piperazine; piperidine; N-ethylpiperidine; and polyamine resins.

[0073] The compounds of the present disclosure can also be solvated, i.e. hydrated. The solvation can be effected in the course of the manufacturing process or can take place i.e. as a consequence of hygroscopic properties of an initially anhydrous compounds. As herein, "solvate" refers to an association or complex of one or more solvent molecules and a compound of the invention. Non-limiting examples of solvents that form solvates include, but are not limited to, water, isopropanol, ethanol, methanol, DMSO, ethyl acetate (EtOAc), acetic acid (AcOH), and ethanolamine.

[0074] Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers." Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers." Diastereomers are stereoisomers with opposite configuration at one or more chiral centers

which are not enantiomers. Stereoisomers bearing one or more asymmetric centers that are non-superimposable mirror images of each other are termed "enantiomers." When a compound has an asymmetric center, for example, if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center or centers and is described by the R- and S-sequencing rules of Cahn, Ingold and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-) isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture". In certain embodiments the compound is enriched by at least about 90% by weight with a single diastereomer or enantiomer. In other embodiments the compound is enriched by at least about 95%, 98%, or 99% by weight with a single diastereomer or enantiomer.

[0075] Certain compounds of the present disclosure possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, regioisomers and individual isomers (e.g., separate enantiomers) are all intended to be encompassed within the scope of the present disclosure.

[0076] The compounds of the invention may contain asymmetric or chiral centers, and therefore exist in different stereoisomeric forms. It is intended that all stereoisomeric forms of the compounds of the invention, including but not limited to, diastereomers, enantiomers and atropisomers, as well as mixtures thereof such as racemic mixtures, form part of the present invention. In some instances, the stereochemistry has not been determined or has been provisionally assigned. Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane polarized light. In describing an optically active compound, the prefixes D and L, or R and S, are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with (-) or l meaning that the compound is levorotatory. A compound prefixed with (+) or d is dextrorotatory. For a given chemical structure, these stereoisomers are identical except that they are mirror images of one another. A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species, devoid of optical activity. Enantiomers may be separated from a racemic mixture by a chiral separation method, such as supercritical fluid chromatography (SFC). Assignment of configuration at chiral centers in separated enantiomers may be tentative while stereochemistry is definitively established, such as from x-ray crystallographic data.

[0077] As used herein, "essentially" refers to at least 90%, at least 95%, at least 98% or at least 99%.


[0078] In the description herein, if there is a discrepancy between a depicted structure and a

name given to that structure, then the depicted structure controls. Additionally, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold wedged, or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it. In some cases, however, where more than one chiral center exists, the structures and names may be represented as single enantiomers to help describe the relative stereochemistry.


[0079] Unless otherwise indicated, the terms "a compound of the formula" or "a compound of formula" or "compounds of the formula" or "compounds of formula" refer to any compound selected from the genus of compounds as defined by the formula (including any pharmaceutically acceptable salt of any such compound if not otherwise noted).

[0080] The patentable scope of the invention is defined by the claims. Any subject-matter falling outside the scope of the claims is provided for information purposes only. The references to methods of treatment in the subsequent paragraphs of this description are to be interpreted as references to the compounds, pharmaceutical compositions and medicaments of the present invention for use in a method for treatment of the human body by therapy.

[0081] In one aspect provided herein is a crystalline tartrate salt having the structure:




[0082] Also disclosed is a process for preparing a compound of formula (IV), or a salt thereof:




Formula (IV)

where the process comprises the steps:

1. (a) reacting a reaction mixture comprising a compound of formula (I), an organic solvent and thionyl chloride to form a compound of formula (IIa) according to step 1 below, and reacting a reaction mixture comprising the compound of formula (IIa), a catalyst, an oxidant and a solvent to form a compound of formula (II) according to step 2 below






wherein

each of R^{1a} and R^{1b} is independently hydrogen, halogen, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, $-CN$, C_{3-6} cycloalkyl, or C_{3-6} spirocycloalkyl, and

n is an integer of 2 or 3; and

2. (b) reacting a reaction mixture comprising the compound of formula (II) and a compound of formula (III) in an organic solvent to form a compound of formula (IV) or a salt thereof according to step 3 below



wherein B is substituted or unsubstituted indolyl, benzofuranyl, benzothiophenyl, indazolyl, aza-indolyl, benzimidazolyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, pyrrolopyridazinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, thienopyridazinyl, thienopyrimidinyl, thienopyrazinyl, furopyridazinyl, fuopyrimidinyl, or fuopyrazinyl,

each of R^{2a} and R^{2b} is independently hydrogen, halogen, $-OH$, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl, $-CN$, C_{3-6} cycloalkyl, or C_{3-6} spirocycloalkyl,

R^{3a} and R^{3b} are independently hydrogen, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} hydroxyalkyl, $-CN$, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, phenyl, C_{3-6} heteroaryl, or C_{3-6} spirocycloalkyl, and

the asterisk represents a chiral center when R^{3a} and R^{3b} are different.

[0083] In one embodiment, B is substituted indolyl, benzofuranyl, benzothiophenyl, indazolyl, aza-indolyl, benzimidazolyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, pyrrolopyridazinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, thienopyridazinyl, thienopyrimidinyl, thienopyrazinyl, furopyridazinyl, fuopyrimidinyl, or fuopyrazinyl.

[0084] In another embodiment, B is unsubstituted indolyl, benzofuranyl, benzothiophenyl,


indazolyl, aza-indolyl, benzimidazolyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, pyrrolopyridazinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, thienopyridazinyl, thienopyrimidinyl, thienopyrazinyl, furopyridazinyl, furopyrimidinyl, or furopyrazinyl.

[0085] In one embodiment, B is a substituted or unsubstituted indolyl, benzofuranyl, or benzothiophenyl. In another embodiment, B is a indolyl, benzofuranyl, or benzothiophenyl substituted with one or more halogen or C₁₋₃ alkyl as described herein. In still another embodiment, B is a substituted or unsubstituted pyrrolopyridazinyl, pyrrolopyrimidinyl, or pyrrolopyrazinyl. In another embodiment, B is a pyrrolopyridazinyl, pyrrolopyrimidinyl, or pyrrolopyrazinyl substituted with one or more halogen or C₁₋₃ alkyl as described herein. In yet another embodiment, B is a substituted or unsubstituted indolyl. In one preferred embodiment, B is unsubstituted indolyl. In one embodiment, B is substituted indolyl (e.g. substituted with one or more halogen or C₁₋₃ alkyl as described herein). In another preferred embodiment, B is substituted indolyl comprising substitution with at least one moiety selected from the group consisting of methyl, Cl, and F. In still another embodiment, B is a benzofuranyl or a substituted benzofuranyl comprising substitution with at least one moiety selected from the group comprising methyl, Cl, and F.

[0086] B may suitably be substituted with one or two substituents independently selected from the group consisting of fluorine, chlorine, C₁₋₃ alkyl, C₁₋₃ haloalkyl, -CN, -OH, C₁₋₃ alkoxy and C₁₋₃ hydroxyalkyl. In one embodiment, B is indolyl substituted with halogen (e.g. For Cl).

[0087] In one embodiment, R^{1a} and R^{1b} are each independently hydrogen, halogen, C₁₋₃ alkyl, C₁₋₃ haloalkyl, C₁₋₃ alkoxy, -CN, or C₃₋₆ cycloalkyl. In another embodiment, R^{1a} and R^{1b} are each independently hydrogen, -F, -Cl, -OH, -CN, -CH₃, -CF₃, -CHF₂, -CH₂F, or spirocyclopropyl. In one embodiment, R^{1a} and R^{1b} are independently F or hydrogen. In a preferred embodiment, R^{1a} and R^{1b} are each independently hydrogen, -F, or -CH₃. In another preferred embodiment, R^{1a} and R^{1b} are each independently hydrogen, -F, or cyclopropyl. In one embodiment, n is 3.



[0088] In one embodiment:



is of the formula




[0089] R^{2a} and R^{2b} are each independently hydrogen, halogen, -OH, C₁₋₃ alkyl, C₁₋₃ haloalkyl,




including stereoisomers thereof.

[0096] In a particular embodiment, formula (II) is:




(2)

[0097] In some particular embodiments, the compound of formula (II) is:




including stereoisomers thereof.

[0098] In one embodiment, the compound of formula (III) is:




including stereoisomers thereof, where X is -NH-, -N-C₁-C₃ unsubstituted alkyl, -O- or -S-.

[0099] In a particular embodiment, the compound of formula (III) is:




[0100] In another particular embodiment, the compound of formula (III) is:




(3)

[0101] In one embodiment, the compound of formula (IV) is:




or salt thereof, including stereoisomers thereof; and wherein the asterisk denotes a chiral center.


[0102] In a particular embodiment, formula (IV) is:




[0103] In another particular embodiment, formula (IV) is:



[0104] In a particular embodiment, the compound of formula (I) is:




the compound of formula (II) is




(2)

the compound of formula (III) is



(3)

and the compound of formula (IV) is



(4)

[0105] In step 1, a reaction mixture comprising a compound of formula (I), an organic solvent and thionyl chloride is reacted to form a compound of formula (IIa). In one embodiment, the compound of formula (I) is compound 1. In one embodiment, the organic solvent is a non-polar solvent or a polar solvent. In one embodiment, the solvent is non-polar. Non-limiting examples of suitable non-polar solvents include pentane, cyclopentane, hexane, cyclohexane, benzene, toluene, 1,4-dioxane, chloroform, diethyl ether, dichloromethane ("DCM"), and combinations thereof. In one embodiment, the solvent is DCM. In one embodiment, the concentration of formula (I) in the solvent may suitably be about 25 g/L, about 50 g/L, about 100 g/L, about 150 g/L, about 200 g/L, about 250 g/L, and up to a concentration approaching saturation at the reaction temperature, and ranges constructed from those concentrations, such as from about 100 g/L to about 250 g/L. The equivalent ratio of thionyl chloride to the compound of formula (I) is suitably about 1:1, about 1.1:1, about 1.2:1, about 1.3:1, about 1.5:1 or about 2:1, and ranges constructed from those ratios, such as from about 1.1:1 to about 1.5:1. In one embodiment, the reaction temperature is below the reaction mixture reflux temperature. In one embodiment, the reaction is run at reflux. For instance, where the solvent is DCM, the reaction temperature may suitably be from about 25°C to about 40°C. The reaction time is not narrowly limited and the reaction is typically continued until the conversion of formula (I) to formula (IIa) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC).

[0106] Upon reaction completion, the reaction mixture may be quenched. In some such embodiments, the reaction mixture may be quenched with cold water. In such embodiments, the phases may be separated into an aqueous phase and organic phase comprising the compound of formula (IIa). The aqueous phase may be extracted one or more times with organic solvent to recover additional compound formula (IIa).

[0107] In step 2, a reaction mixture comprising the compound of formula (IIa), a catalyst, an oxidant and a solvent is reacted to form a compound of formula (II). In one embodiment, the organic phase or combined organic phases from step 1, comprising formula (IIa), is used as the source of formula (IIa) for step 2. In one embodiment, the catalyst is a redox active metal

catalyst. Non-limiting examples of suitable catalysts include NiCl_2 , RuCl_3 , CoCl_2 , FeCl_3 , FeCl_2 , and MnCl_2 . Non-limiting examples of suitable oxidants include NaIO_4 , NaOCl , and Oxone. Suitable organic solvents include non-polar and polar solvents as discussed elsewhere herein. In general, the oxidant is in equivalent excess to compound formula (IIa), for instance, the ratio of oxidant to compound formula (IIa) may be 1.1:1, 1.5:1, 2:1, 2.5:1, 3:1, 3.5:1, 4:1, 4.5:1 or 5:1. The step 2 reaction mixture may further comprise water. In such embodiments, the volume ratio of water to organic solvent used in the step 1 reaction mixture may be about 9:1, about 5:1, about 3:1, about 2:1, about 1:1, about 1:2, about 1:3, about 1:5 or about 1:9, and ranges constructed therefrom, such as from about 2:1 to about 1:2. The step 2 reaction temperature may suitably be about 25°C , about 15°C , about 5°C , about 0°C , about -5°C or about -10°C , and ranges constructed therefrom, such as from about -10°C to about 10°C . In one embodiment, the organic phase(s) comprising formula (IIa), catalyst, and water are combined and cooled to a reaction temperature. The oxidant is then added over a period of time while maintaining the temperature around the reaction temperature.

[0108] Upon reaction completion, the step 2 reaction mixture may be separated into an aqueous phase and organic phase comprising the compound of formula (II) in solution. In some optional embodiments, the reaction mixture may be filtered, such as through a filter aid (e.g., celite) prior to phase separation. The aqueous phase may be extracted one or more times with organic solvent to recover additional compound formula (II).

[0109] In another embodiment, the step 2 organic phase(s) may be worked up by methods known to those skilled in the art. For instance, the organic phases may be washed with a base, such as with an aqueous solution of Na_2SO_3 . The organic phases may further optionally be dried, such as with a brine solution and/or by the addition of a solid drying agent such as CaCl_2 , MgSO_4 or Na_2SO_4 . Solid desiccants may suitably be removed by filtration. In one embodiment, the compound formula (II) solution may be used for subsequent reaction. In one embodiment, compound formula (II) may be isolated from the solution by methods known in the art such as by distillation, concentration, precipitation (such as by addition of an anti-solvent or pH adjustment) and/or crystallization. In some such embodiments, the organic phase(s) may be concentrated by distillation or stripping to reduce the volume, such as by at least 25%, 50%, 100% or more. Compound formula (II) may then be precipitated/crystallized from solution by addition of an anti-solvent followed by optional further concentration. In one embodiment, the anti-solvent is a C_{4-8} nonionic solvent such as pentane, hexane or heptane. Compound formula (II) solids may be collected by methods known in the art such as filtration or centrifugation. The solids may be dried, such as under partial vacuum, to yield solid compound formula (II). The yield to compound formula (II) from compound formula (I) for steps 1 and 2 is at least 60%, at least 70%, or at least 75%. In one embodiment, the compound of formula (II) is compound 2.

[0110] In step 3, a reaction mixture comprising the compound of formula (II), a compound of formula (III), and an organic solvent is reacted to form a compound of formula (IV). In one embodiment, the organic solvent is a polar aprotic solvent. Non-limiting examples of suitable

solvents include tetrahydrofuran, ethyl acetate, acetone, dimethylformamide, acetonitrile ("ACN"), dimethyl sulfoxide, nitromethane and propylene carbonate. In one embodiment, the solvent is ACN. The mole ratio of compound formula (II) to compound formula (III) is suitably about 1:1, about 1.1:1, about 1.2:1, about 1.3:1, about 1.4:1, about 1.5:1, or greater, and ranges constructed from those ratios, such as between 1:1 and 1.3:1. The concentration of compound formula (II) in the solvent is suitably about 10 g/L, about 25 g/L, about 50 g/L, about 75 g/L, about 100 g/L, about 125 g/L, about 150 g/L, and up to a concentration approaching saturation at the reaction temperature, and ranges constructed from those concentrations, such as from about 50 g/L to about 150 g/L. The acid catalyst may be an acid catalyst as described elsewhere herein. In some embodiments, the acid catalyst is sulfuric acid, *p*-toluene sulfonic acid (*p*-TsOH), or methansulfonic acid, or combinations thereof. In one embodiment, the acid catalyst is *p*-toluene sulfonic acid. The equivalent ratio of acid catalyst to compound formula (II) is suitably about 0.75:1, about 0.9:1, about 1:1, about 1.05:1, about 1.1:1, about 1.2:1, about 1.3:1, about 1.4:1, about 1.5:1, or greater, and ranges constructed therefrom, such as from about 1:1 to about 1.2:1. In one embodiment, the compound of formula (III) is compound 3.


[0111] In some step 3 embodiments, compound formula (II), compound formula (III), the organic solvent and a base are combined to form an admixture. The base may suitably be a moderate base, non-limiting examples of which include potassium *tert*-butoxide, trimethylamine, sodium bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate, sodium hydroxide, ammonium hydroxide, and combinations thereof. The admixture may be heated with agitation to a reaction temperature, typically a temperature of from 2°C to about 30°C below the reflux temperature up to the reflux temperature, and held for a time sufficient to essentially complete the formation of a reaction product comprising compound formula (III). In the case of ACN solvent, the reaction temperature is suitably about 65°C, about 70°C, about 75°C, or about 80°C. The reaction product mixture may then be cooled, such as to less than 50°C, than 40°C, or than 30°C, and optionally filtered to remove solid impurities. The solids may be optionally washed with the solvent to recover additional reaction product. Acid (e.g., *p*-TsOH) and water are then added. The volume ratio of organic solvent to water may be 25:1, 15:1, 10:1, 5:1, 2:1 or 1:1, and ranges constructed therefrom, such as from about 15:1 to about 5:1. The admixture may be heated with agitation to a reaction temperature, typically at a temperature of from 2°C to about 20°C below the reflux temperature to the reflux temperature, and held for a time sufficient to essentially complete the formation of compound formula (IV) such as determined by chromatography (e.g., TLC, GC or HPLC). Upon reaction completion, the reaction mixture may be quenched, such as with cold water (e.g., less than 10°C or less than 5°C). The pH of the quenched reaction mixture may then be adjusted with a base to greater than 7, such as about pH 8, about pH 9, about pH 10 or about pH 11. In one embodiment, the base is an aqueous base such as sodium bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate, sodium hydroxide, or ammonium hydroxide.

[0112] Upon reaction completion, the step 3 reaction mixture may be separated into an aqueous phase and organic phase comprising the compound of formula (IV) in solution. In

some optional embodiments, the reaction mixture may be filtered, such as through a filter aid (e.g., celite) prior to phase separation. The aqueous phase may be extracted one or more times with organic solvent to recover additional compound formula (IV). In one embodiment, the solvent is aprotic. In a particular embodiment, the extracting solvent is suitably isopropyl acetate ("*i*-PrOAc").


[0113] In some embodiments, the step 3 organic phase(s) may be worked up by, for instance, washing the organic phases with water. The organic phases may optionally be dried, such as with a brine solution and/or by the addition of a solid drying agent such as CaCl₂, MgSO₄ or Na₂SO₄. Solid desiccants may suitably be removed by filtration, and the collected desiccant may optionally be washed with solvent to recover compound formula (IV) therefrom. In such embodiments, the organic phase(s) may be concentrated by distillation under partial vacuum or stripping to form compound formula (IV) residue. The compound formula (IV) residue may then be dissolved in organic solvent at a temperature below the reflux temperature. Anti-solvent, such as a non-polar organic solvent as described elsewhere herein, may then be added to the compound formula (IV) solution while cooling, such as to less than about 10°C, to precipitate/crystallize compound formula (IV) from solution. Compound formula (IV) solids may be collected by methods known in the art such as filtration or centrifugation, and optionally washed with anti-solvent. The solids may be dried, such as under partial vacuum, to yield solid compound formula (IV). The yield to compound formula (IV) from compound formula (II) for step 3 is at least 80%, at least 85%, at least 90%, at least 95%, at least 96% or at least 97%. Compound formula (IV) purity is at least 95%, at least 98%, or at least 99%.


[0114] One aspect of the disclosure is directed to a process for preparing a compound of formula (VIII) or a salt thereof:



where the process comprises the steps of:

1. (a) reacting a reaction mixture comprising a compound of formula (IV), a compound of formula (V) or a compound of formula (X), and an organic solvent to form a compound of formula (VI) according to step 1 below





wherein

B is substituted or unsubstituted indolyl, benzofuranyl, benzothiophenyl, aza-indolyl, indazolyl, benzimidazolyl, pyrrolopyridinyl, furopyridinyl, thienopyridinyl, pyrrolopyridazinyl, pyrrolopyrimidinyl, pyrrolopyrazinyl, thienopyridazinyl, thienopyrimidinyl, thienopyrazinyl, furopyridazinyl, furopyrimidinyl, or furopyrazinyl;

each of R^{1a} and R^{1b} is independently hydrogen, fluorine, chlorine, -OH, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl, and -CN, C_{3-6} cycloalkyl, or C_{3-6} spirocycloalkyl,

n is an integer of 2 or 3,

each of R^{2a} and R^{2b} is independently hydrogen, halogen, -OH, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, C_{1-3} hydroxyalkyl, -CN, C_{3-6} cycloalkyl, or C_{3-6} spirocycloalkyl,

R^{3a} and R^{3b} are independently hydrogen, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} alkoxy, -CN, C_{3-6} cycloalkyl, C_{3-6} heterocycloalkyl, phenyl, C_{3-6} heteroaryl, or C_{3-6} spirocycloalkyl,

J is phenyl or pyridinyl;

each R^4 is independently hydrogen, halogen or C_{1-3} alkyl,

s is an integer from 0 to 2,

LG is a leaving group,

LG and CHO are located in the para position with respect to each other on J on the compound of formula (V),


PG is an aldehyde protecting group,

LG and CH-PG are located in the para position with respect to each other on J on the compound formula (X), and

each asterisk independently represents a chiral center wherein the carbon bearing R^{3a} and R^{3b} is a chiral center when R^{3a} and R^{3b} are different; and

2. (b) reacting a reaction mixture comprising the compound of formula (VI), an organic solvent, and a compound of formula (VII) or a salt thereof to form a compound of formula (VIII) or a salt thereof according to step 2 below





wherein

G is C₁₋₃ alkyl,

p is 0 or 1,

E is substituted or unsubstituted azetidinyll or pyrrolidinyl,

each R⁵ is independently hydrogen, halogen, -OH, -CN, C₁₋₅ alkoxy, or C₁₋₅ hydroxyalkyl,

v is an integer from 1 to 5,

R⁶ is halogen or -CN; and

R¹⁰ is hydrogen or C₁₋₃ alkyl.

[0115] B, R^{1a}, R^{1b}, n, R^{2a}, R^{2b}, R^{3a}, R^{3b} and the asterisk (*) are as defined herein.

[0116] In one preferred embodiment, J is phenyl. In another embodiment, J is pyridinyl.

[0117] In one embodiment, each R⁴ is independently hydrogen or halogen. In a preferred embodiment, each R⁴ is fluorine. In one embodiment, s is 1 or 2. In one embodiment, s is 2. In one preferred embodiment, each R⁴ is fluorine and s is 2.

[0118] In one embodiment, G is methylene or ethylene.

[0119] In one embodiment, p is 0.

[0120] In one embodiment, each R⁵ is independently hydrogen, halogen, -OH, or -CN. In one preferred embodiment, each R⁵ is hydrogen. In one embodiment, v is 2. In another embodiment, v is 3. In another embodiment, v is 5. In a preferred embodiment, each R⁵ is hydrogen and v is 3.


[0121] In one preferred embodiment, R⁶ is halogen. In one embodiment, R⁶ is F. In another

embodiment, R⁶ is -CN.


[0122] In one embodiment, R¹⁰ is hydrogen or methyl. In a preferred embodiment, R¹⁰ is hydrogen

[0123] In one embodiment, E is azetidiny. In another embodiment, E is pyrrolidinyl.


[0124] In one embodiment, E has the following structure:



[0125] In one embodiment, E is azetidiny of the following structure:




[0126] In one embodiment, E is of the following structure:




where R⁵ is H, v is 2 or 3, and R⁶ is halogen.






[0127] In one embodiment, E is azetidiny of the following structure:




[0128] In one embodiment, formula (VIII) is an acid salt. Such acid salt can be a pharmaceutically acceptable salt. In some such embodiments, formula (VIII) is a salt of a pharmaceutically acceptable acid. In some particular embodiments, formula (VIII) is a salt of a pharmaceutically acceptable organic acid. In a preferred embodiment, formula (VIII) is a pharmaceutically acceptable salt of tartaric acid. In one embodiment, the compound of formula (VIII) is Compound A as described herein. In another embodiment, the compound of formula (VIII) is Compound A (tartrate salt) of Compound A described herein. In another embodiment, formula (VIII) is a pharmaceutically acceptable salt of fumaric acid. In still another embodiment, the compound of formula (VIII) is Compound B (fumarate salt) of Compound A as described herein.

[0129] In one embodiment, formula (VIII) is of any one of the following structures, or a pharmaceutically acceptable salt thereof:










or a pharmaceutically acceptable salt thereof and including stereoisomers thereof.

[0130] In one embodiment, formula (VIII) is of the following structure, or a pharmaceutically acceptable salt thereof:






[0131] In one embodiment, formula (VIII) is of the following structure:




[0132] In one embodiment, formula (VIII) is of the following structure:





[0133] Step 1 of the process to synthesize compounds of formula (VIII) comprises reacting a reaction mixture comprising a compound of formula (IV), a compound of formula (V) or a compound of formula (X), and an organic solvent to form a compound of formula (VI) as set forth herein. In one embodiment, LG is bromine. In a preferred embodiment, when reacting with compounds of formula (IV) and formula (V), LG and CHO are located on J in the para position with respect to each other. In a preferred embodiment, when reacting compounds of formula (IV) and formula (X), LG and CH-PG are located in the para position with respect to each other on J.

[0134] The compound of formula (IV) is as described herein.

[0135] In one embodiment, the compound of formula (V) is of any one of the following compounds, or a salt thereof:










or a salt thereof, including stereoisomers thereof.

[0138] In step 2 of the synthesis of compounds of formula (VIII) set forth herein, the compound of formula (VII) can be any one of the following structures:




or



or a salt thereof.


[0139] In one embodiment, the salt is an ethane-disulfonate (e.g. a salt of ethane-1,2-disulfonate).

[0140] In one embodiment, the compound of formula (VII) has structure



[0141] Compound (7) can be prepared according to the examples provided herein, such as, for example, Example 4 or Example 4a.

[0142] In one embodiment, Compound 7 is prepared according to the scheme below:



[0143] In step 1, a reaction mixture comprising a compound of formula (IV), a compound of formula (V) or a compound of formula (X), and an organic solvent are reacted to form a compound of formula (VI). In one embodiment, the organic solvent is a polar protic solvent, a non-polar solvent, a polar aprotic solvent, or a combination thereof, as described elsewhere

herein. In some non-limiting embodiments, the solvent is toluene. In one embodiment, the solvent is acetonitrile, methyl ethyl ketone, or methyltetrahydrofuran. In one embodiment, the step 1 reaction mixture further comprises an acid catalyst as described elsewhere herein. In some such non-limiting embodiments, the acid catalyst is acetic acid. The mole ratio of compound (IV) to compound (V) or compound (X) is from about 0.95: 1 to about 1.05: 1, stoichiometric amounts, or, in some embodiments, compound (V) or compound (X) is in slight molar excess. The acid catalyst is generally present in stoichiometric excess, such as in an equivalent ratio to compound formula (IV) of about 1.1:1, about 1.2:1, about 1.3:1, about 1.4:1, about 1.5:1, about 1.6:1, about 1.7:1, about 1.8:1, about 2:1, or greater, and ranges constructed therefrom, such as from about 1.2:1 to about 1.8:1.

[0144] In some step 1 embodiments, the reaction mixture may be heated with agitation to a reaction temperature, of from 2°C to about 30°C below the reflux temperature to the reflux temperature, and held for a time sufficient to essentially complete the reaction to compound formula (VI) such as determined by chromatography (e.g., TLC, GC or HPLC). In the case of toluene solvent, the reaction temperature is suitably about 65°C, about 70°C, about 75°C, or about 80°C. The reaction product mixture may then be cooled and optionally diluted with additional solvent. The reaction product mixture may then be quenched with a base, such as an aqueous solution of a base as described elsewhere herein. A step 1 reaction product mixture organic phase comprising the compound of formula (VI) may then be isolated and, in some embodiments, worked up by methods known in the art such as washing with water and/or a brine solution as described elsewhere herein followed by product isolation as a solid. In one embodiment, the reaction product mixture may be treated with activated charcoal, followed by filtration and optional washing of the activated charcoal filter cake with solvent. As described elsewhere herein: (i) the organic phase(s) containing compound formula (VI) may be concentrated by distillation or stripping to reduce the volume, such as by at least 25%, 50%, 100% or more; and (ii) compound formula (IV) may then be precipitated/crystallized from solution by addition of an anti-solvent followed by optional further concentration.

[0145] In some other step 1 embodiments, the step 1 reaction mixture is heated at reflux for a time to essentially complete the reaction such as determined by chromatography (e.g., TLC, GC or HPLC). The reaction mixture may then be cooled and pH-adjusted with a base, such as an aqueous solution of a base, to a pH at which compound formula (VI) precipitates from solution.

[0146] In any of the various step 1 embodiments, compound formula (VI) solids may be collected by methods known in the art such as filtration or centrifugation. The solids may be dried, such as under partial vacuum, to yield solid compound formula (VI). The yield to compound formula (VI) from compound formula (V) for steps 1 and 2 is at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95%.

[0147] In step 2, a reaction mixture comprising a compound of formula (VI), a compound of formula (VII) and an organic solvent are reacted to form a compound of formula (VIII). In some embodiments, the organic solvent is a polar aprotic solvent as described elsewhere herein. In

some non-limiting embodiments, the solvent is ACN. The step 2 reaction mixture may further comprise a base, such as an organic base. Non-limiting examples of organic bases include DBU, NMM, DIPEA, and TEA. In some such embodiments, the base is DBU. The step 2 reaction mixture may further comprise a catalyst, such as a transition metal catalyst. In some embodiments the catalyst is a Pd catalyst. The mole ratio of compound (VII) to compound (VI) is about 0.95: 1, about 1:1, about 1.05:1, about 1.1:1, about 1.2:1, about 1.3:1, about 1.4:1 or about 1.5:1, and ranges constructed therefrom, such as from about 1:1 to about 1.4:1. The base is generally present in stoichiometric excess, such as in an equivalent ratio to compound formula (VI) of about 1.1:1, about 2: 1, about 3:1, about 4:1, about 5:1, about 6:1, about 7: 1, or about 8:1, and ranges constructed therefrom, such as from about 3: 1 to about 7:1. The reaction mixture may be heated with agitation to a reaction temperature, typically a temperature of from 2°C to about 30°C below the reflux temperature up to the reflux temperature, and held for a time sufficient to essentially complete the reaction such as determined by chromatography (e.g., TLC, GC or HPLC). In the case of ACN solvent, the reaction temperature is suitably about 65°C, about 70°C, about 75°C, or about 80°C.


[0148] After completion the reaction product mixture may suitably be cooled and optionally diluted with an organic solvent. In one embodiment, the reaction product mixture is diluted with a non-polar solvent as described elsewhere herein. One non-limiting example of a suitable non-polar solvent is MTBE. The step 2 reaction product mixture may be worked up by methods known to those skilled in the art including water wash and brine wash. In some such non-limiting embodiments, the work-up may include washing with an aqueous solution of ammonium chloride, brine and water. In some embodiments, the step 2 reaction product mixture may be contacted with a metal scavenger known in the art, such as for instance and without limitation, SiliaMetS Thiol. The reaction product mixture may then be filtered to remove solids prior to isolation of compound formula (VIII) therefrom.

[0149] In some embodiments, the step 2 reaction product mixture may be concentrated, such as by vacuum distillation or stripping, and diluted with an organic solvent, such as an alcohol (e.g., ethanol), such as in a solvent exchange step. An acid may then be added to the diluted solution of compound formula (VIII) followed by cooling to crystallize compound formula (VIII) as an acid salt. In some particular embodiments, the acid is tartaric acid and compound formula (VIII) is the tartaric salt. In some such embodiments, the acid is (2*R*-3*R*)-tartaric acid (L-(+)-tartaric acid). In another aspect, the acid is (2*S*-3*S*)-tartaric acid (D-(-)-tartaric acid). In some such embodiments, the solvent comprises an organic solvent. The crystalline material may be collected by centrifugation or filtration, optionally washed with solvent, and optionally dried.

[0150] In some other embodiments, compound formula (VIII) may be isolated from the step 2 reaction product mixture using methods described elsewhere herein including: (i) distillation, concentration, precipitation (such as by addition of an anti-solvent or pH adjustment) and/or crystallization; (ii) solids collection by centrifugation or filtration; (iii) optional washing of the collected solids; (iv) and drying.


[0151] The step 2 yield of compound formula (VIII) either as a free base or acid salt is at least 80%, at least 85%, or at least 90%.

[0152] Another aspect of the disclosure is directed to a process for preparing a compound of formula (VIII) or a salt thereof, wherein compound formula (VIII) is as described elsewhere herein. The process for preparing formula (VIII) according to this embodiment comprises reaction step 1 as depicted below:



[0153] Each of B, R^{1a}, R^{1b}, n, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R⁴, s, J, R⁵, v, R⁶, R¹⁰, G, p, E, PG and the asterisk are as described elsewhere herein. The CHO moiety and the nitrogen atom linking J and G are located in the para position with respect to each other on J. The CH-PG moiety and the nitrogen atom linking J and G are located in the para position with respect to each other on J.

[0154] In one embodiment, the compound of formula (IX) is:



or a salt thereof.

[0155] In one preferred embodiment, the compound of formula (IX) is compound (8a). In some embodiments, formula (XI) corresponds to any of the above formula (IX) structures, or a salt thereof, but where the aldehyde (-CHO) is a protected moiety of the structure -CH-PG where PG is an aldehyde protecting group as defined elsewhere herein.


[0156] In step 1, a reaction mixture comprising a compound of formula (IX) or of formula (XI),

a compound of formula (IV) and an organic solvent is reacted to form a compound of formula (VIII), or a salt thereof. In one embodiment, the organic solvent is a polar solvent, or is a polar protic solvent. The step 1 reaction mixture further comprises an acid catalyst as described elsewhere herein. In a particular embodiment, the acid catalyst is tartaric acid or fumaric acid. In one embodiment, the acid catalyst is tartaric acid. In another embodiment, the acid catalyst is fumaric acid. Non-limiting examples of suitable solvents include n-butanol, isopropyl alcohol, n-propanol, i-propanol, ethanol, methanol, and combinations thereof. In some particular embodiments, the solvent is ethanol. In one embodiment, the concentration of formula (IX) in the solvent may suitably be about 25 g/L, about 50 g/L, about 100 g/L, about 150 g/L, about 200 g/L, about 250 g/L, and up to a concentration approaching saturation at the reaction temperature, and ranges constructed from those concentrations, such as from about 100 g/L to about 250 g/L. The mole ratio of formula (IX) or of formula (XI) to formula (IV) is suitably about 0.25:1, about 0.3:1, about 0.4:1, about 0.5:1, about 0.6:1, about 0.7:1, about 0.8:1, about 0.9:1, about 0.95:1, about 1:1, about 1.05:1, about 1.1:1 or about 1.2:1, and ranges constructed from those ratios, such as from about 0.95:1 to about 1.05:1. In one embodiment, formula (IX) or formula (XI) and formula (IV) are present in approximately stoichiometric amounts. In one embodiment, the reaction temperature is below the reaction mixture reflux temperature. In some other embodiments, the reaction is run at reflux. For instance, where the solvent is ethanol, the reaction temperature may suitably be from about 50°C to about 75°C. The reaction time is not narrowly limited and the reaction is typically continued until the conversion of formula (IX) or formula (XI) to formula (VIII) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC).


[0157] In one embodiment, formula (VIII) may be formed as a salt of an acid. Suitable acids include inorganic acids and organic acids as described elsewhere herein. In one embodiment, the acid is an organic acid. In a preferred embodiment, the acid is tartaric acid. In another embodiment, the acid is fumaric acid. In one embodiment, formula (VIII) free base may be dissolved in a suitable solvent, such as a polar protic solvent (e.g., an alcohol such as methanol or ethanol) at an elevated temperature followed by addition of an acid. In general, the acid is in stoichiometric excess as compared to compound formula (VIII). In some such embodiments, the solution temperature and/or concentration of formula (VIII) is adjusted to keep the concentration below saturation and thereby avoid formula (VIII) precipitation and/or crystallization. Following acid addition, the solution may optionally be seeded with a crystalline formula (VIII) salt of the acid. In any of the various embodiments, the solution is cooled with stirring to form crystalline formula (VIII) salt. The salt may then be collected by methods known in the art, such as by filtration or centrifugation. In one embodiment, the salt is collected by filtration. The collected formula (VIII) salt may be optionally washed, such as with the dissolution solvent, and then dried, such as under partial vacuum.

[0158] In a particular embodiment, the step 1 reaction mixture as set forth above for synthesis of a compound of formula (VIII) comprises tartaric acid as the acid catalyst, crystalline formula (VIII) tartaric acid, and an alcoholic solvent (e.g. ethanol); the step 1 reaction product mixture is diluted with the alcoholic solvent; and the resultant slurry is cooled with stirring to form crystalline formula (VIII) tartaric acid. In another embodiment, the step 1

reaction mixture comprises fumaric acid as the acid catalyst, crystalline formula (VIII) fumaric acid, and a solvent. The step 1 yield of compound formula (VIII) as a tartaric acid salt is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95%. In such embodiments, the step 1 reaction scheme is as follows:



[0159] Still another aspect of the disclosure is directed to a process for preparing a compound of formula (IX) or a salt thereof. The process for preparing compound formula (IX) comprises two reaction steps as depicted below:



[0160] R⁴, s , LG, PG, R⁵, v , R⁶, R¹⁰, G, p , J, and E are as described elsewhere herein.

[0161] Aldehyde protecting groups are defined herein and non-limiting examples include 1,3-dithiane, 1,3-dithiolane, diethyl acetal, dimethyl acetal, ethylene glycol acetal, neopentyl glycol acetal, trimethylsilyl cyanohydrin, and triethyl orthoformate.

[0162] In step 1, a reaction mixture comprising a compound of formula (X), a compound of formula (VII) or a salt thereof, an organic solvent and a catalyst is reacted to form a compound of formula (XI). In one embodiment, the solvent is a non-polar solvent as described elsewhere herein. Non-limiting examples of suitable solvents include pentane, hexane, heptane, cyclopentane, MTBE, diethyl ether, toluene, 2-methyl tetrahydrofuran (2-MeTHF), benzene, 1,4-dioxane, carbon tetrachloride, chloroform, dichloromethane, and combinations thereof. In some examples, the solvent is toluene. In one embodiment, the catalyst is a transition metal catalyst as described herein. In one embodiment, non-limiting examples of transition metal catalysts include palladium, platinum, gold, ruthenium, rhodium, and iridium catalysts. Non-limiting examples of suitable catalysts include JohnPhos, XPhos AuCl, XPhos AuNTf₂, XPhos Palladacycle, SPhos Palladacycle, tBuXPhos Pd G1, Xphos Pd G2, SPhos Pd G2, RuPhos Pd G2, CPhos-Pd-G2, CPhos-Pd-G3, tBuXPhos-Pd-G3, RuPhos-Pd-G3, XPhos-Pd-G3, BrettPhos-Pd-G3, JackiePhos-Pd-G3, tert-butyl BrettPhos-Pd-G3, [tert-butyl BrettPhos-Pd (allyl)]OTf, and combinations thereof. In some embodiments the catalyst is BrettPhos-Pd-G3. In some embodiments, the step 1 reaction mixture further comprises an organic base as describe elsewhere herein. In some such embodiments, the base is a tert-butoxide such as sodium or potassium *tert*-butoxide. The mole ratio of the compound of formula (VII) to the compound of formula (X) is suitably about 1:1, about 1.1:1, about 1.2:1, about 1.3:1, about 1.5:1 or about 2:1, and ranges constructed from those ratios, such as from about 1.1:1 to about 1.5:1.


[0163] In some step 1 embodiments, the reaction mixture may be heated with agitation to a reaction temperature, of from 2°C to about 30°C below the reflux temperature to the reflux temperature, and held for a time sufficient to essentially complete the reaction to compound formula (XI) such as determined by chromatography (e.g., TLC, GC or HPLC). In the case of toluene solvent, the reaction temperature is suitably about 50°C, about 55°C, about 60°C, about 65°C, about 70°C, about 75°C, or about 80°C. The reaction time is not narrowly limited and the reaction is typically continued until the conversion of formulae (VII) and (X) to formula (XI) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC). After the reaction is complete the reaction product mixture may be suitably quenched. In some embodiments, the step 1 reaction may be quenched with water. Where the reaction is quenched with water, the organic phase comprising the compound of formula (XI) in solution may be isolated and optionally washed at least once with water. In some embodiments, the step 1 reaction product mixture may be contacted with a metal scavenger known in the art, such as for instance and without limitation, SiliaMetS Thiol. The reaction product mixture may then be filtered to remove solids.

[0164] In step 2, the compound of formula (XI) is deprotected to form compound formula (IX) by combining a solution of compound (XI) in an organic solvent (e.g., toluene) with an acid and water. The acid is generally present in equivalent excess, such an equivalent ratio of acid to compound (XI) of 1.01:1, 1.05:1, 1.1:1, 1.15:1, 1.2:1, or greater. In one embodiment, the deprotection temperature is not narrowly critical and may suitably be room temperature. After deprotection, the organic phase and the aqueous phase (comprising the compound of formula (IX)) are separated. The organic phase may optionally be washed with water. The aqueous phase(s) may be treated with a base, such as an inorganic base (e.g., NaOH or KOH),

combined with compound formula (IX) seed crystals. The base may suitably be added in equivalent excess. A slurry of crystalline compound formula (IX) forms with optional cooling. Compound formula (IX) solids may be collected by methods known in the art such as filtration or centrifugation. The solids may be dried, such as under partial vacuum, to yield solid compound formula (IX). The yield to compound formula (IX) from compound formula (XI) for steps 1 and 2 is at least 65%, at least 70%, at least 75%, at least 80% or, at least 85%.


[0165] One aspect of the disclosure is directed to a process for preparing a compound of formula (III) or a salt thereof. The process for preparing compound formula (III) comprises two reaction steps as depicted below:

- (1) reacting a reaction mixture comprising a compound of formula (XII), a compound B and an organic solvent to form the compound of formula (XIII) according to step 1 below



where PG is an amine protecting group; and


- (2) deprotecting the compound of formula (XIII) to form the compound of formula (III) according to step 2 below



where R^{2a}, R^{2b}, R^{3a}, R^{3b}, B, and the asterisk are as described elsewhere herein.


[0166] Non-limiting examples of amine protecting groups include ACD, Ac, Bn, CBz, trifluoroacetamide, Boc, MeOZ, Fmoc, Bz, PMB, DMPM, PMP, Ts and Troc. In one embodiment, PG is Boc.

[0167] In one embodiment, the compound of formula (III) is of the structure



or a salt thereof as described herein.

[0168] In some other embodiments, the compound of formula (III) is of the structure



(3)

or a salt thereof as described elsewhere herein.

[0169] In step 1, a reaction mixture comprising a compound of formula (XII), B and an organic solvent is reacted to form a compound of formula (XIII). In one embodiment, the organic solvent is a non-polar solvent or a polar solvent. In one embodiment, the solvent is non-polar. Non-limiting examples of suitable solvents include pentane, cyclopentane, hexane, cyclohexane, benzene, toluene, 1,4-dioxane, chloroform, diethyl ether, DCM, and combinations thereof. In one embodiment, the solvent is DCM. In some embodiments, the concentration of B in the solvent may suitably be about 10 g/L, about 25 g/L, about 50 g/L, about 75 g/L, about 100 g/L, about 125 g/L, about 150 g/L, about 175 g/L, or about 200 g/L, and up to a concentration approaching saturation at the reaction temperature, and ranges constructed from those concentrations, such as from about 25 g/L to about 125 g/L. The equivalent ratio of B to compound formula (XII) is about 0.75:1, about 0.9:1, about 1:1, about 1.25:1, about 1.5:1, about 1.75:1 or about 2:1, and ranges thereof, such as from about 1.25:1 to about 1.75:1. The step 1 reaction mixture may further comprise a suitable alkylating agent. Non-limiting examples of alkylating reagents include alkyl lithium (e.g., methyl lithium) such as and organomagnesium halide compounds, such as methyl magnesium chloride (e.g., in THF). The equivalent ratio of B to the alkylating reagent is suitably about 0.75:1, about 0.8:1, about 0.85:1, about 0.9:1, about 0.95:1, about 1:1, about 1.05:1, about 1.1:1, about 1.15:2, about 1.2:1, about 1.3:1, about 1.4:1, about 1.5:1 or greater, and ranges constructed therefrom, such as from about 1.1 to about 1.3:1. In one embodiment, the reaction mixture further comprises an alkylation catalyst. In some such embodiments, the catalyst is a transition metal catalyst. In some such embodiments the transition metal is copper. In a particular embodiment, the catalyst is suitably a transition metal halide, such as copper (I) halide (e.g., CuCl). In one embodiment, the reaction temperature is about 25°C, about 20°C, about 15°C, about 10°C, about 5°C, about 0°C, about -5°C, about -10°C, about -15°C, about -20°C, about -25°C, about -30°C, about -35°C, about -40°C, about -45°C, or about -50°C, and ranges constructed therefrom, such as from about -20°C to about 0°C. In one embodiment, the reaction time is not narrowly limited and the reaction is typically continued until the conversion of B and compound formula (XII) to compound formula (XIII) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC).

[0170] After reaction completion, the reaction may be quenched, such as for instance by the addition of aqueous acid. In one embodiment, the acid may be an organic acid as described elsewhere herein, one non-limiting example of which is citric acid. The organic phase comprising the compound of formula (XIII) may then be worked up to dry the compound. In some such embodiments, the quenched reaction product mixture may be separated into an aqueous phase and an organic phase comprising the compound of formula (XIII). The aqueous phase may be washed one or more times with an organic solvent, such as the solvent

used to form the reaction mixture (e.g., two washes, each with one volume of the solvent as compared to the reaction mixture volume). The organic phases may be combined and washed one or more times with brine (e.g., two brine washes, each with one volume of brine as compared to the reaction mixture volume). The washed organic phases may then be combined with stirring with activated carbon and with a solid drying agent (e.g., Na₂SO₄). Any activated carbon and solid drying agent may be removed by filtration or centrifugation. Any collected solids may then be optionally washed with additional solvent to recover compound formula (XIII) therefrom.

[0171] In one embodiment, the solution of compound formula (XIII) may be used as the starting material for step 2. In one embodiment, solid compound formula (XIII) may be prepared. In such embodiments, the collected solution of compound formula (XIII) in organic solvent may be concentrated under partial vacuum to form crude compound formula (XIII). Alternatively, solid compound formula (XIII) may be precipitated/crystallized from solution by the addition of an anti-solvent, such as a non-polar solvent (e.g., 2 volumes of heptane as compared to the solvent volume in the reaction mixture). The solids may be collected by filtration or centrifugation and the collected solids may be washed with anti-solvent. The solids may be dried under partial vacuum, at a temperature of less than 40°C to provide finished compound formula (XIII). The compound formula (XIII) yield based on compound (XII) is at typically least 50%, at least 55%, at least 60%, or at least 65%. The purity by HPLC (area percent) is at least 90%, at least 95%, at least 98% or at least 99%.

[0172] In some step 1 embodiments, compound B, a metal catalyst and an alkylating agent are combined in a first volume of the solvent at the reaction temperature indicated above. The volume of solvent is suitably from about 30% to about 80% of the total volume of solvent used for step 1. Thereafter, a solution of compound formula (XII) in the remainder of the solvent is added at the reaction temperature over a time period to form the reaction mixture. The reaction mixture is then held a temperature for a time to essentially complete the formation of compound formula (XIII), followed by quenching and work-up to a solution of compound formula (XIII) or dried compound formula (XIII).

[0173] In step 2, compound formula (XIII) is deprotected to form compound formula (III). In any of the various embodiments, a solution of compound formula (XIII) is suitably deprotected by the addition of an acid. In one embodiment, solid compound formula (XIII) is dissolved in a polar protic solvent as described elsewhere herein (such as methanol, ethanol or i-propanol). The concentration of compound formula (XIII) in the solvent is about 25 g/L, about 50 g/L, about 75 g/L, about 100 g/L, about 125 g/L, about 150 g/L, about 175 g/L, or about 200 g/L, and ranges constructed therefrom, such as from about 50 g/L to about 150 g/L. Acid addition temperature is not narrowly critical.

[0174] In some step 2 embodiments, the acid is an inorganic acid as described elsewhere herein. A non-limiting example of a suitable inorganic acid is HCl. In one embodiment, the equivalent ratio of acid to compound formula (XIII) is about 1.5:1, about 2.5:1, about 5:1, about 7.5:1, about 10:1, about 12.5:1 or about 15:1 and ranges constructed therefrom, such as from

about 5:1 to about 15: 1. After acid addition, the solution is held at temperature with stirring until the deprotection of compound formula (XIII) to form compound formula (III) is essentially complete. In one embodiment, a solvent exchange from the polar protic solvent to a non-polar solvent (as described elsewhere herein) or a polar aprotic solvent (as described elsewhere herein) may be done. In some such embodiments, the solution of compound formula (III) may be concentrated under partial vacuum and extracted with the non-polar or polar aprotic solvent. In some such embodiments, the extraction solvent is DCM. After extraction, the pH of the aqueous phase may be adjusted to strongly basic (i.e., greater than pH 11) with a suitable base, such as aqueous sodium hydroxide. After basification, the aqueous phase may then be further extracted at least once with the extraction solvent. The organic phase(s) may then be dried, such as with brine and/or over a solid desiccant followed by filtration to remove any solids. Collected solids may be optionally washed to recover additional compound formula (III). Solid compound may be isolated from the solution in organic solvent by methods known in the art. For instance, the solution may be concentrated under partial vacuum to dryness. Alternatively, the solution may be concentrated followed by the addition of an anti-solvent to form solid compound (III) that may be collected by filtration or centrifugation, washed, and dried. The compound formula (III) yield based on compound (XIII) is at typically least 85%, at least 90%, at least 95%, or at least 97%. The purity by HPLC (area percent) is at least 90%, at least 95%, or at least 96%.

[0175] In some other step 2 embodiments, the acid is an organic acid as described elsewhere herein. Non-limiting examples of suitable organic acids include sulfonic acids and camphorsulfonic acid (CSA) (e.g., L(-)-CSA). In the case of CSA, the equivalent ratio of CSA to compound formula (XIII) is about 1.5:1, about 2:1 about 2.5:1, about 5:1, about 7.5:1, or about 10:1, and ranges constructed therefrom, such as from about 2: 1 to about 4:1. After acid addition, the solution may be heated and held at elevated temperature (e.g., about 35°C to about 60°C) with stirring to complete deprotection and form the acid salt of compound formula (XIII). The acid salt solution/suspension may then be cooled, such as to less than about 5°C, to form a suspension of the acid salt of compound formula (XIII). The salt may be collected by filtration or centrifugation and optionally washed with solvent. The salt may then be dried under partial vacuum to yield the solid salt of compound of compound formula (XIII), such as the L(-)-CSA salt thereof. The compound formula (III) salt yield based on compound formula (XIII) is at typically least 85%, at least 90%, at least 92% or at least 95%. The purity by HPLC (area percent) is at least 90%, at least 95%, at least 96%, or at least 97%. The compound formula (XIII) salt may be dissolved in water and pH-adjusted to greater than 13, such as about 14, with a strong base thereby forming a suspension comprising solid compound formula (III) free base. The solid material may be collected by filtration or centrifugation and optionally washed with chilled water. The solids may then be dried under partial vacuum to yield dried compound formula (III) free base. The compound formula (III) yield based on compound (XIII) is at typically at least 80%, at least 85%, or at least 90%. The purity by HPLC (area percent) is at least 90%, at least 95%, at least 96%, or at least 97%.

[0176] In one embodiment, a compound of formula (VIII) as prepared herein is further recrystallized. In one embodiment the recrystallization comprises recrystallizing the compound


of formula (VIII) in a 2-step process. The process comprises heating a slurry comprising a compound of formula (VIII) in a mixture of methanol/ethanol, distilling with methanol, and cooling the mixture. In one embodiment, the mixture of methanol/ethanol is a 95:5, 90:10, 85:15, or 80:20 mixture of methanol/ethanol. In another embodiment, the mixture of ethanol/methanol is a 90:10 mixture of methanol/ethanol. The mixture can be heated at a temperature of > about 50 °C, for example, about 55 °C, 60 °C, or 65 °C. The cooling can be down to about room temperature. In one embodiment, the cooling is to about 20 °C, 25 °C, or 30 °C. The solution can be filtered and dried.

[0177] In another embodiment, the recrystallization comprises recrystallizing the compound of formula (VIII) from MTBE. A base, such as NaOH or KOH is added to the slurry comprising the compound of formula (VIII) in MTBE. The mixture is stirred at, for example, 15 °C, 20 °C, 25 °C or 30 °C, optionally filtered and distilled with ethanol.

[0178] In some embodiments of the process for preparing a compound of formula (III) or a salt thereof, the process further comprises preparing the compound of formula (XII) or a salt thereof.


[0179] The process for preparing the compound of formula (XII) comprises reaction steps 3a and 3b depicted below:

- (1) reacting a reaction mixture comprising a compound of formula (XIV), thionyl chloride, and an organic solvent to form a compound of formula (XV) according to step 3a below



and

- (2) reacting a reaction mixture comprising the compound of formula (XV), a catalyst, an oxidizing agent, and an organic solvent to form the compound of formula (XII) according to step 3b below



[0180] R^{2a}, R^{2b}, R^{3a}, R^{3b}, and the nitrogen protecting group PG are as described herein.

[0181] In step 3a, a reaction mixture comprising a compound of formula (XIV), thionyl chloride, and an organic solvent is reacted to form a compound of formula (XV). In one embodiment, the organic solvent is a non-polar solvent or a polar solvent as described elsewhere herein. In one embodiment, the solvent is non-polar as described elsewhere herein. Non-limiting examples of suitable solvents include pentane, cyclopentane, hexane, cyclohexane, benzene, toluene, 1,4-dioxane, chloroform, diethyl ether, DCM, and combinations thereof. In some particular embodiments, the solvent is DCM. In one embodiment, the concentration of the compound of formula (XIV) in the solvent may suitably be about 10 g/L, about 25 g/L, about 50 g/L, about 75 g/L, about 100 g/L, about 125 g/L, about 150 g/L, about 175 g/L, or about 200 g/L, and up to a concentration approaching saturation at the reaction temperature, and ranges constructed from those concentrations, such as from about 25 g/L to about 125 g/L. The equivalent ratio of thionyl chloride to compound formula (XIV) is about 1: 1, about 1.25:1, about 1.5:1, about 1.75:1, about 2:1, about 2.25: 1, or about 2.5:1, and ranges thereof, such as from about 1.25: 1 to about 1.75:1. In one embodiment, the reaction temperature is about 10°C, about 5°C, about 0°C, about -5°C, or about -10°C, and ranges constructed therefrom, such as from about -5°C to about 5°C. In one embodiment, the reaction time is not narrowly limited and the reaction is typically continued until the conversion of compound formula (XIV) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC). The step 3a reaction mixture may further comprise a base, a non-limiting example of which is imidazole. In such embodiments, the equivalent ratio of thiolation reagent to thionyl chloride may be about 1:1, about 2:1, about 3:1 or about 4:1. The step 3a reaction mixture may further comprise a base, such as an organic base as described elsewhere herein. In some such embodiments, the base may be TEA. In such embodiments, the equivalent ratio of the base to compound formula (XIV) is about 1.25:1, about 1.5:1, about 1.75:1, about 2:1, about 2.25:1, about 2.5:1, about 3:1, about 3.5:1 or about 4:1, and ranges constructed therefrom, such as from about 1.5:1 to about 2.5:1.

[0182] In some particular step 3a embodiments, a solution of the base (e.g., imidazole) in the solvent is formed to which the thionyl chloride is added with stirring while maintaining the reaction temperature. Compound formula (XIV) in the solvent may then be added with stirring while maintaining the reaction temperature, followed by addition of the base with stirring while maintaining the reaction temperature. The reaction mixture is then maintained at the reaction temperature with stirring until the conversion of compound formula (XIV) to form a reaction product mixture comprising the compound of formula (XV) is essentially complete.

[0183] In any of the various step 3a embodiments, the reaction product mixture may be worked up by methods known to those skilled in the art. For instance, the step 3a reaction mixture may be quenched with chilled water (e.g., 0.25 to 2 volumes of water per volume of organic solvent in the step 3a reaction product mixture). An organic phase comprising the compound of formula (XV) in solution may be isolated and the isolated aqueous phase may be extracted with organic solvent to recover additional compound formula (XV). The organic phases may be combined and washed with an aqueous acid solution, and aqueous base solution, and brine. A non-limiting example of an aqueous acid solution is a solution of a weak


acid, such as a citric acid. A non-limiting example of a base solution is a solution of a weak base, such as sodium bicarbonate.


[0184] In step 3b, a reaction mixture comprising formula (XV) in solution in the organic solvent is combined with a catalyst and an oxidizing agent, and reacted to form a reaction product mixture comprising the compound of formula (XII). In one embodiment, the catalyst is a redox active metal catalyst as described elsewhere herein. Non-limiting examples of suitable catalysts include NiCl₂, RuCl₃, CoCl₂, FeCl₃, FeCl₂, and MnCl₂. Non-limiting examples of suitable oxidants include NaIO₄, NaOCl, and Oxone. In some embodiments the reaction mixture further comprises water. The volume ratio of water to organic solvent is suitably about 0.25:1, about 0.5:1, about 0.75:1, about 1:1, about 1.25:1, about 1.50:1, about 1.75:1, about 2:1 or about 2.5:1, and ranges constructed therefrom, such as from about 0.5:1 to about 1.5:1. The reaction temperature is suitably from about 5°C to about 50°C. The reaction mixture is maintained at the reaction temperature with stirring until the conversion of compound formula (XV) to form a reaction product mixture comprising the compound of formula (XII) is essentially complete such as determined by chromatography (e.g., TLC, GC or HPLC).

[0185] The step 3b reaction product mixture may be worked up by methods known to those skilled in the art. For instance, the organic phase comprising the compound of formula (XII) in solution may be isolated and the aqueous phase may be optionally filtered and then extracted with organic solvent to recover compound formula (XII). The organic phase(s) may optionally be washed with a reducing agent (e.g., sodium thiosulfate) and with brine. The organic phase(s) may be dried with a solid desiccant (e.g., sodium sulfate) followed by filtration to remove solids and optional washing thereof with organic solvent. Solid compound formula (XII) from solution in organic solvent by methods known in the art. For instance, the solution may be concentrated under partial vacuum to dryness. Alternatively, the solution may be concentrated followed by the addition of an anti-solvent to form solid compound (XII) that may be collected by filtration or centrifugation, washed, and dried. The compound formula (XII) yield based on compound (XIV) is at typically least 85%, at least 90%, or at least 92%. The purity by HPLC (area percent) is at least 95%, at least 98%, or at least 99%.


[0186] Chiral primary amines can be produced using a stereospecific transaminase either via an asymmetric transamination of a prochiral ketone or kinetic resolution of a racemic amine. The transamination is a reaction equilibrium consisting of a pair of amines and ketones (donors and acceptors) and the reaction conditions applied shift the equilibrium to the chiral target amine. In one such aspect, the amine donor is alanine or 2-propylamine. In one embodiment, the amine acceptor is pyruvate or acetone.

[0187] Thus, further provided herein is a process for preparing a compound of formula (XX) or a salt thereof, wherein conversion of the chiral tryptamine comprises using enzymatic transformation:






[0188] The process for preparing compound formula (XX) comprises contacting a compound of formula (XXI)



with a protein transaminase to form a compound of formula (3):



[0189] The process further comprises the presence of one or more amine donors. In one embodiment the amine donor is alanine or iso-propyl amine.


[0190] The contacting described above can be performed in mixed aqueous organic solvent systems. For example, the transaminase reaction can be performed in aqueous buffer with an organic co-solvent such as cyclo-hexane, methyl-cyclo-hexane, iso-octane, DMSO, acetonitrile, or acetone. Such co-solvents can be present at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, or 50% v/v.

[0191] The mixed organic / aqueous solvent system (micro-aqueous reaction system) can, in certain instances, comprise one or more organic solvents comprising a small amount of aqueous buffer. In one embodiment, the contacting is performed in TBME, dibutyl ether, CPME, toluene, ethyl acetate, butyl acetate, iso-propyl acetate, butyl butyrate, ethyl butyrate or iso-butyl acetate comprising less than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1% v/v aqueous buffer. When performed in organic solvent, the transaminase can be immobilized to a solid support.


[0192] In another embodiment, the conversion of compound (XXI) to compound (3) is above at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%. In another aspect, the conversion is above 90%. In another embodiment, the conversion is above 95%.

[0193] The reductive amination completed by the transaminase can yield the compound of formula (3) in at least 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% enantiomeric excess (EE). In certain instances, the transaminase converts the compound of formula (XXI) to (3) at an EE greater than 99%.

[0194] In one embodiment, the asymmetric transamination of the compound of formula (XXI) is performed using the scheme below:



[0195] The compound of formula (3) is contacted with a compound of formula (II) as described herein:




(II)

to form compound formula (XX).

[0196] R^{1a}, R^{1b} and n are as described elsewhere herein. In one embodiment, R^{1a} and R^{1b} are independently fluorine.


[0197] In one embodiment, compound formula (XX) is of the structure:



[0198] The protein transaminase can be selected from Table 6. In one such aspect, the transaminase is (R)-selective. In one embodiment, the transaminase is TA-P2-A01. In one embodiment, the protein transaminase is selected from a (S)-enantioselective transaminase of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:4.

[0199] Further provided herein are methods of enzymatically synthesizing a compound of formula (3) through kinetic resolution of a racemic amine in the presence of an amine acceptor and a transaminase. The reaction can be performed in conditions as described herein, for example, in mixed aqueous/organic solvent systems as described herein. In one such aspect, the reaction is performed in aqueous buffer comprising acetonitrile. Such kinetic resolution can yield the compound of formula (3) in an EE of at least 99%, requiring above 50% conversion rate. Kinetic resolution comprises use of an (S)-selective transaminase as described herein for performing the kinetic resolution. In one embodiment, the transaminase is selected from Table 6.

[0200] One embodiment of the disclosure is directed to a compound of formula (XVI):



(XVI)

R^4 , s , R^{10} , G , and p are as defined herein.

each of R^{7a} , R^{7b} , R^{8a} and R^{8b} is independently hydrogen, halogen, C_{1-3} alkyl, C_{1-3} haloalkyl, C_{1-3} hydroxyalkyl, or $-CN$. In one embodiment, each of R^{7a} , R^{7b} , R^{8a} and R^{8b} is independently hydrogen, fluorine, $-CH_3$, or $-CN$. In one embodiment, each of R^{7a} , R^{7b} , R^{8a} and R^{8b} is hydrogen.

y is an integer of 1 or 2, x is an integer of 1 or 2, and the total of x and y is 2 or 3.

M is C_{1-5} alkyl;


r is 0 or 1; and


R^9 is halogen or $-CN$.

[0201] In one embodiment, M is $-CH_2CH_2CH_2-$, p is 1, and R^9 is fluorine.


[0202] In one embodiment, M is $-CH_2CH_2CH_2-$, p is 1, and R^9 is $-CN$.

[0203] In one embodiment, the compound of formula (XVI) is:







[0204] In a particular embodiment, the compound of formula (XVI) is:



[0205] In another aspect provided herein is a compound having the structure:



[0206] In another aspect provided herein is a process to make a compound having formula:






(B)


wherein the process is performed as set forth in Scheme A below and in accordance with the embodiments and embodiments provided herein.

Scheme A:



BrettPhos Pd G3 = [(2-di-cyclohexylphosphino-3,6-dimethoxy-2',4',6'- triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate


[0207] In still another embodiment, of the disclosure is a process to make a compound having formula:




(B)

wherein the process is performed as set forth in Scheme B below and in accordance with the embodiments and embodiments provided herein.


Scheme B:






[0208] The processes for synthesis of Compound B above (e.g. Scheme A and B) can further comprise recrystallization according to Scheme C or D below:

Scheme C:




Scheme D:




[0209] In one embodiment, compounds (3) and (4) as described herein and provided in Scheme A and Scheme B, and optionally Scheme C or D, above are synthesized according to Scheme E below and in accordance with the embodiments and embodiments provided herein.

Scheme E:




[0210] Further provided herein is a process for preparing a compound of formula (XXIII) or a salt thereof:





[0211] B, R^{1a}, R^{1b}, n, R^{2a}, R^{2b}, R^{3a}, R^{3b}, J, R⁴, s, G, R⁵, v, R⁶, E, and the asterisks are as defined herein.

[0212] In one embodiment, the compound of formula (XXIII) is an acid salt. In such embodiments, the compound of formula (VIII) is a salt of an acid. In a preferred embodiment, the compound of formula (XXIII) is a salt of tartaric acid. In some embodiments, the compound of formula (XXIII) is a salt of fumaric acid.

[0213] In one embodiment, the compound of formula (XXIII) is of any one of the following structures, or a tartaric acid salt thereof:










[0214] In another embodiment, the compounds above are a fumaric acid salt.

[0215] In one embodiment, the compound of formula (XXIII) is of the following structure, or a pharmaceutically acceptable salt thereof:



[0216] The process for preparing the compound of formula (XXIII) comprises two reaction steps as depicted below:







[0217] LG is as defined herein.

[0218] The variables of formulae (IV), (V) and (VI) are as described herein.

[0219] In one embodiment, the compound of formula (XXIV) is of any one of the following structures, or a salt thereof:




[0220] One embodiment of the disclosure is directed to a process for preparing a compound of formula (XXIII) or a salt thereof, wherein the compound of formula (XXIII) is as described herein. The process for preparing the compound of formula (XXIII) according to this embodiment comprises reaction step 1 as depicted below:



[0221] Each of B, R^{1a}, R^{1b}, n, R^{2a}, R^{2b}, R^{3a}, R^{3b}, R⁴, s, J, R⁵, v, R⁶, G, p, E and the asterisk are as described elsewhere herein. The CHO moiety and the nitrogen atom linking J and G are

located in the para position with respect to each other on J.


[0222] In some embodiments, compound formula (XXIV) is:



or a salt thereof


[0223] One embodiment of the disclosure is directed to a process for preparing a compound of formula (XXVII) or a salt thereof.

[0224] The process for preparing compound formula (XXVII) comprises two reaction steps as depicted below:



[0225] R^4 , s , LG, R^5 , v , R^6 , G, p, E and PG are as described herein.

[0226] Further provided herein are solid forms, formulations comprising such solid forms, and methods of using such solid forms of Compound A:




(A)


and having the name 3-(((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-

ol, including a pharmaceutically acceptable salt thereof.


[0227] In one embodiment provided herein are solid forms of Compound A. Compound A can be a freebase as described herein existing in an amorphous solid form. In another embodiment, Compound A is a crystalline solid as described herein. In still another embodiment, Compound A is a crystalline tartrate salt having the structure:



[0228] In another aspect provided herein are crystalline solid forms of Compound A as a fumarate salt having the structure:



[0229] In still another aspect provided herein are solid forms of Compound A malonate salts having the structure:



[0230] Solid forms described herein can be crystalline. In another embodiment, the solid form is a single-component solid form. Solid forms described herein can be solvates, hydrates, anhydrides, or salts as set forth herein. In one embodiment, solid forms described herein comprise tartrate salts. In another embodiment, solid forms described herein comprise anhydrides of Compound B. In another embodiment, solid forms described herein comprise

fumarate salts. In still another embodiment, solid forms described herein comprise phosphate salts or other salts.

[0231] While not intending to be bound by any particular theory, solid forms can be characterized by physical properties such as, for example, stability, solubility and dissolution rate, density, compressibility, hardness, morphology, cleavage, stickiness, solubility, water uptake, electrical properties, thermal behavior, solid-state reactivity, physical stability, and chemical stability) affecting particular processes (e.g., yield, filtration, washing, drying, milling, mixing, tableting, flowability, dissolution, formulation, and lyophilization) which make certain solid forms suitable for the manufacture of a solid dosage form. Such properties can be determined using particular analytical chemical techniques, including solid-state analytical techniques (e.g., X-ray diffraction, microscopy, spectroscopy and thermal analysis), as described herein.

[0232] The solid forms described herein, including salt forms, crystalline forms, and amorphous solids can be characterized by a number of methods including, for example, single crystal X-ray diffraction, X-ray powder diffraction (XRPD), microscopy (e.g., scanning electron microscopy (SEM)), thermal analysis (e.g., differential scanning calorimetry (DSC), dynamic vapor sorption (DVS), thermal gravimetric analysis (TGA), and hot-stage microscopy), spectroscopy (e.g., infrared, Raman, and solid-state nuclear magnetic resonance), ultra-high performance liquid chromatography (UHPLC), proton nuclear magnetic resonance (spectrum.

[0233] Techniques for characterizing crystal forms and amorphous solids include, for example, thermal gravimetric analysis (TGA), differential scanning calorimetry (DSC), X-ray powder diffractometry (XRPD), single-crystal X-ray diffractometry, vibrational spectroscopy, e.g., infrared (IR) and Raman spectroscopy, solid-state and solution nuclear magnetic resonance (NMR) spectroscopy (including ^1H NMR and ^19F NMR), scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility studies, and dissolution studies.

[0234] The purity of the solid forms provided herein can be determined by standard analytical methods, such as thin layer chromatography (TLC), gel electrophoresis, gas chromatography, ultra-high performance liquid chromatography (UHPLC), and mass spectrometry (MS).

Compound B, Form A:

[0235] In certain embodiments, provided herein is a solid form of Compound B designated as Form A. Form A is a crystalline solid form of Compound B. In one embodiment, Form A is an acetone solvate of Compound B. In one embodiment, Form A is a crystalline acetone solvate tartrate salt of Compound A.

[0236] In another embodiment, Form A of Compound B is obtained by slurring in acetone

followed by evaporation. Form A can be prepared according to the methods and examples described herein.

[0237] In one embodiment, a solid form provided herein, e.g., Form A, is a tartrate salt of Compound A (i.e. Compound B), and is crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD of a solid form provided herein, e.g., Form A, is substantially as shown in FIG. 1. In another embodiment, a solid form provided herein, e.g., Form A, has one or more characteristic XRPD peaks at approximately 4.64, 8.26, 9.28, 11.18, 11.49, 11.96, 12.54, 13.77, 14.22, 14.61, 15.09, 15.56, 16.01, 17.35, 18.55, 18.84, 19.32, 19.82, 20.26, 21.34, 21.63, 21.92, 22.52, 22.97, 23.28, 23.54, 23.94, 24.81, or 25.96 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 1 and as found in Table 16 herein. In still another embodiment, a solid form provided herein, e.g., Form A, has at least 3, at least 5, at least 7, or at least 10 XRPD peaks at approximately 4.64, 8.26, 9.28, 11.18, 11.49, 11.96, 12.54, 13.77, 14.22, 14.61, 15.09, 15.56, 16.01, 17.35, 18.55, 18.84, 19.32, 19.82, 20.26, 21.34, 21.63, 21.92, 22.52, 22.97, 23.28, 23.54, 23.94, 24.81, or 25.96 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 1 and as found in Table 16 herein. In yet another embodiment, a solid form described herein, e.g., Form A, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or all of the characteristic XRPD peaks as set forth in Table 16.

[0238] In still another embodiment, a solid form provided herein, e.g., Form A, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 12.54, 14.61, 16.01, 19.32, 20.26, 21.63, 23.28, 23.54, 23.94, or 24.81 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 1 and as found in Table 16 herein. In another embodiment, a solid form provided herein, e.g., Form A, has one, two, three, four, or five characteristic XRPD peaks at approximately 19.32, 20.26, 21.63, 23.28, or 24.81 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 1 and as found in Table 16 herein. In another embodiment, a solid form provided herein, e.g., Form A, has one, two, three, four, or five characteristic XRPD peaks at approximately 19.32, 20.26, 21.63, 23.28, or 24.81 $\pm 0.05^\circ 2\theta$, as depicted in, for example, FIG. 1 and as found in Table 16 herein.

[0239] In one embodiment described herein, is a solid form, e.g., Form A, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 2. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 7.2% of the total mass of the sample before approximately 125 $^\circ\text{C}$.

[0240] In another embodiment described herein, is a solid form, e.g., Form A, having a DSC thermogram substantially as depicted in FIG. 2 comprising desolvation event at about 124 $^\circ\text{C}$ and a melting temperature of having an onset temperature of about 164 $^\circ\text{C}$ and a peak maximum temperature of about 171 $^\circ\text{C}$.

[0241] In another embodiment described herein, is a solid form, e.g., Form A, having a Polarized Light Microscopy image as depicted in FIG. 3.

[0242] In still another embodiment, Form A is pure. In certain embodiments, pure Form A is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, the purity of Form A is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form B:

[0243] In certain embodiments, provided herein is a solid form of Compound B designated as Form B. Form B is a crystalline solid form of Compound B. In one embodiment, Form B is an anhydrate of Compound B. In another embodiment, Form B is an anhydrate tartrate salt of Compound A.

[0244] In one embodiment, Form B of Compound B is obtained by slurring Compound B in ethyl acetate at RT for about 24 hours. In another embodiment, Form B of Compound B is obtained by slurring Compound B in acetone:water (e.g. 90:10, 95:5, 96:4, 97:3, 99:1 v/v) at about 50 °C for about 6 hours. In still another embodiment, Form B of Compound B is obtained by slurring Compound B in ethanol.

[0245] In certain instances, Form B of Compound B is obtained by slurring Compound B in a solvent system comprising $\geq 95\%$ acetone (e.g. $\geq 95:5$ acetone:water). Form B of Compound B is then isolated from the slurry by, for example, centrifugation or filtration. In another embodiment, Form B of Compound B is obtained from either Form A or Form F as described herein. In one embodiment, Form A of Compound B is reslurried in ethanol (e.g. 100% ethanol) for 10, 12, 16, or 24 hours (e.g. overnight) to obtain Form B. In one embodiment, Form F is converted to Form D as described herein in the presence of water. The mixture can then be slurried in neat ethanol (at for example about 50 °C) or 95:5 or 97:3 acetone:water (v/v) to form Form B. Mixtures can optionally be seeded with Form B crystals.

[0246] Form B can be prepared according to the methods and examples described herein. Thus, provided herein is a method of preparing Form B where the method comprises slurring Compound B in a solvent system comprising acetone or aqueous mixtures of acetone water (e.g., 50%, 90%, 95%, 96%, 97%, 98% and 99% acetone v/v). In one embodiment, the solvent system for crystalizing Form B comprises $\geq 95\%$ acetone. In one embodiment, the solvent system for Form B comprises 96:4 acetone:water. The mixtures can be slurried at RT for about 120 hrs. The mixtures can be filtered and analyzed as described herein (e.g., XRPD). In one embodiment, Form B is prepared according to the methods and/or examples set forth herein.

[0247] In one embodiment, a solid form provided herein, e.g., Form B, is a tartrate salt of Compound A, and is crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD of a solid form provided herein, e.g., Form B, is substantially as shown in FIG. 4. In another embodiment, a solid form provided herein, e.g.,

Form B, has one or more characteristic XRPD peaks at approximately 7.68, 11.49, 12.54, 14.24, 15.30, 15.55, 16.01, 16.63, 17.37, 18.24, 19.16, 19.42, 19.89, 20.24, 21.81, 22.52, 22.99, 23.25, 23.57, 24.67, 25.07, 25.91 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 4 and as found in Table 17 herein. In still another embodiment, a solid form provided herein, e.g., Form B, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 7.68, 11.49, 12.54, 14.24, 15.30, 15.55, 16.01, 16.63, 17.37, 18.24, 19.16, 19.42, 19.89, 20.24, 21.81, 22.52, 22.99, 23.25, 23.57, 24.67, 25.07, 25.91 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 4 and as found in Table 17 herein. In yet another embodiment, a solid form described herein, e.g., Form B, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or all of the characteristic XRPD peaks as set forth in Table 17.

[0248] In still another embodiment, a solid form provided herein, e.g., Form B, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 11.49, 12.54, 15.30, 15.55, 19.16, 19.42, 20.24, 23.25, 24.67, or 25.91 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 4. In still another embodiment, a solid form provided herein, e.g., Form B, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.49, 12.54, 19.16, 19.42, or 24.67 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 4. In still another embodiment, a solid form provided herein, e.g., Form B, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.49, 12.54, 19.16, 19.42, or 24.67 $\pm 0.05^\circ 2\theta$, as depicted in, for example, FIG. 4.

[0249] In one embodiment described herein, is a solid form, e.g., Form B, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 5. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 3.5% of the total mass of the sample.

[0250] In another embodiment described herein, is a solid form, e.g., Form B, having a DSC thermogram substantially as depicted in FIG. 5 comprising an endothermic event with an onset temperature of about 171 $^\circ\text{C}$ and a peak maximum temperature of about 179 $^\circ\text{C}$.

[0251] In another embodiment described herein, is a solid form, e.g., Form B, having a ^{13}C and ^{19}F NMR spectrum substantially as depicted in FIG. 6 and FIG. 7, respectively.

[0252] In another embodiment described herein, is a solid form, e.g., Form B, having a water sorption-desorption profile as depicted in FIG. 8. The solid form, e.g., Form B of Compound B, absorbs about 1.2% w/w moisture up to 90% relative humidity (RH) at about 25 $^\circ\text{C}$.

[0253] In another embodiment described herein, is a solid form, e.g., Form B, having a Scanning Electron Microscope (SEM) image and PLM image as depicted in FIG. 9a and FIG. 9b, respectively. The sample comprises of dense spherical aggregates.

[0254] In another embodiment described herein, is a solid form, e.g., Form B, having a particle size distribution (PSD) as depicted in FIG. 9c.

[0255] In still another embodiment, a solid form, e.g., Form B, remains substantially unchanged following compression as described herein. FIG. 22, FIG. 23, and FIG. 24, respectively show XRPD, ¹⁹F SSNMR, and DSC of Form B of Compound B and compare the compound before and after compression as described herein.

[0256] In still another embodiment, Form B is substantially pure. In certain embodiments, the pure Form B is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, the substantially pure Form B is substantially free of Form A, Form D, or Form F. In certain embodiments, the purity of Form B is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form C:

[0257] In certain embodiments, provided herein is a solid form of Compound B designated as Form C. Form C is a crystalline solid form of Compound B. In one embodiment, Form C is a THF solvate of Compound B.

[0258] In one embodiment, Form C of Compound B is obtained by slurring Compound B in THF. The mixture can then be filtered. Form C can be prepared according to the methods and examples described herein.

[0259] In one embodiment, a solid form provided herein, e.g., Form C, is tartrate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD is substantially as shown in FIG. 10.

[0260] In one embodiment described herein, is a solid form, e.g., Form C, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 11. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 6.8% of the total mass of the sample.

[0261] In another embodiment described herein, is a solid form, e.g., Form C, having a DSC thermogram substantially as depicted in FIG. 11 comprising an endothermic event with an onset temperature of about 118 °C and a peak maximum temperature of about 125 °C.

[0262] In still another embodiment, Form C is pure. In certain embodiments, the purity of Form C is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form D:

[0263] In certain embodiments, provided herein is a solid form of Compound B designated as Form D. Form D is a crystalline solid form of Compound B. In one embodiment, Form D is a hydrate of Compound B. In another embodiment, Form D is a monohydrate of Compound B.

[0264] In one embodiment, Form D of Compound B is obtained by slurring Compound B in 100% ethanol for about 48 hours. The mixture can then be filtered.

[0265] In one embodiment, a solid form provided herein, e.g., Form D, is a tartrate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In another embodiment, Form D is a hydrate of Compound B. In one embodiment, the XRPD of a solid form provided herein, e.g., Form D, is substantially as shown in FIG. 12. In another embodiment, a solid form provided herein, e.g., Form D, has one or more characteristic XRPD peaks at approximately 7.32, 10.99, 11.31, 12.18, 13.23, 13.48, 14.11, 14.66, 15.14, 15.70, 16.03, 16.21, 16.54, 17.24, 17.63, 18.11, 18.34, 19.10, 20.20, 20.58, 21.16, 21.47, 21.89, 22.76, 23.33, or $23.56 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 12 and as found in Table 19. In still another embodiment, a solid form provided herein, e.g., Form D, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 7.32, 10.99, 11.31, 12.18, 13.23, 13.48, 14.11, 14.66, 15.14, 15.70, 16.03, 16.21, 16.54, 17.24, 17.63, 18.11, 18.34, 19.10, 20.20, 20.58, 21.16, 21.47, 21.89, 22.76, 23.33, or $23.56 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 12 and as found in Table 19. In yet another embodiment, a solid form described herein has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or all of the characteristic XRPD peaks as set forth in Table 19.

[0266] In still another embodiment, a solid form provided herein, e.g., Form D, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 11.31, 15.70, 16.54, 19.10, 20.58, 21.16, 21.47, 21.89, 22.76, or $23.33 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 12. In still another embodiment, a solid form provided herein, e.g., Form D, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.31, 15.70, 16.54, 19.10, or $22.76 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 12. In still another embodiment, a solid form provided herein, e.g., Form D, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.31, 15.70, 16.54, 19.10, or $22.76 \pm 0.05^\circ 2\theta$, as depicted in, for example, FIG. 12.

[0267] In one embodiment described herein, is a solid form, e.g., Form D, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 13. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 1.4% of the total mass of the sample before approximately 150 °C.

[0268] In another embodiment described herein, is a solid form, e.g., Form D, having a DSC thermogram substantially as depicted in FIG. 13 comprising an endothermic event having an

onset temperature of about 55 °C and a peak maximum temperature of about 82 °C followed by a second endothermic event with an onset temperature of about 165 °C and a peak maximum temperature of about 172 °C.

[0269] In still another embodiment, Form D is pure. In certain embodiments, Form D is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, Form D is substantially free of Form A, Form B, or Form F. In certain embodiments, the purity of Form D is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form E:

[0270] In certain embodiments, provided herein is a solid form of Compound B designated as Form E. Form E is a solid form of Compound B. In one embodiment, Form E is a DMSO solvate of Compound B.

[0271] In one embodiment, Form E of Compound B is obtained by slurring Compound B in DMSO and adding IPAc, for about 24 hours. The mixture can then be filtered. Form E can be prepared according to the methods and examples described herein.

[0272] In one embodiment, a solid form provided herein, e.g., Form E, is a tartrate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD is substantially as shown in FIG. 14.

[0273] In one embodiment described herein, is a solid form, e.g., Form E, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 15. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 8.3% of the total mass of the sample.

[0274] In another embodiment described herein, is a solid form, e.g., Form E, having a DSC thermogram substantially as depicted in FIG. 15 comprising a first endotherm with an onset temperature of about 126 °C and a peak maximum temperature of about 134 °C and a second endothermic event with an onset temperature of about 143 °C and a peak maximum temperature of about 147 °C.

[0275] In still another embodiment, Form E is pure. In certain embodiments, Form E is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, the purity of Form E is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form F:

[0276] In certain embodiments, provided herein is a solid form of Compound B designated as Form F. Form F is a crystalline solid form of Compound B. In one embodiment, Form F is an anhydrate of Compound B. In another embodiment, Form F is an anhydrate tartrate salt of Compound A.

[0277] In one embodiment, Form F of Compound B is obtained by slurring Compound B in 100% ethanol at RT or at 50 °C for about 8, 10, 12, 15, 20, or 25 hours (e.g. overnight). In another embodiment, Form F can be obtained by slurring Compound B in ethanol/water (e.g. 65:35 v/v). The slurry of Compound B to obtain Form F can optionally include seeding with Form B described herein. The slurry can be filtered to obtain Form F. Form F can also be obtained by slurring at RT in 100% DI water, 1:1 acetone/ water or 100% acetone. Slurries of Form F in neat solvents can be maintained at RT. In one embodiment, Form F can be obtained from 1:1 acetone: water mixture agitated at 5 °C. In still another embodiment, Form F can be obtained by slurring Compound B in 95:5 acetone:water and 97:3 acetone: water mixtures at 50 °C for about 2 hours and cooling to RT. Form F can be prepared according to the methods and examples described herein.

[0278] In one embodiment, a solid form provided herein, e.g., Form F, is a tartrate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD of a solid form provided herein, e.g., Form F, is substantially as shown in FIG. 16. In another embodiment, a solid form provided herein, e.g., Form F, has one or more characteristic XRPD peaks at approximately 3.92, 10.54, 11.72, 12.52, 14.22, 15.40, 15.54, 15.90, 16.48, 16.84, 17.29, 18.26, 18.47, 19.39, 19.66, 20.00, 20.50, 20.65, 21.16, 21.28, 21.95, 22.97, 23.49, 23.70, 23.94, 24.31, 24.67, or 24.99 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 16 and as found in Table 20 herein. In still another embodiment, a solid form provided herein, e.g., Form F, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 3.92, 10.54, 11.72, 12.52, 14.22, 15.40, 15.54, 15.90, 16.48, 16.84, 17.29, 18.26, 18.47, 19.39, 19.66, 20.00, 20.50, 20.65, 21.16, 21.28, 21.95, 22.97, 23.49, 23.70, 23.94, 24.31, 24.67, or 24.99 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 16 and as found in Table 20 herein. In yet another embodiment, a solid form described herein, e.g., Form F, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, twenty, twenty-five, or all of the characteristic XRPD peaks as set forth in Table 20.

[0279] In still another embodiment, a solid form provided herein, e.g., Form F, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 12.52, 15.90, 19.66, 20.65, or 24.99 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 16. In still another embodiment, a solid form provided herein, e.g., Form F, has one, two, three, four, or five characteristic XRPD peaks at approximately 12.52, 15.90, 19.66, 20.65, or 24.99 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 16. In still another embodiment, a solid form provided herein, e.g., Form F, has one, two, three, four, or five characteristic XRPD peaks at approximately 12.52, 15.90, 19.66, 20.65, or 24.99 $\pm 0.05^\circ 2\theta$, as depicted in, for example, FIG. 16.

[0280] In one embodiment described herein, is a solid form, e.g., Form F, having a DVS isotherm plot corresponding substantially to the representative DVS isotherm plot as depicted in FIG. 17.

[0281] In another embodiment described herein, is a solid form, e.g., Form F, having a DSC thermogram substantially as depicted in FIG. 18 comprising an endothermic event with an onset temperature of about 162 °C and a peak maximum temperature of about 167 °C.

[0282] In still another embodiment, Form F is pure. In certain embodiments, pure Form F is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, the purity of the Form F is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound B, Form G

[0283] In certain embodiments, provided herein is a solid form of Compound B designated as Form G. Form G is a solid form of Compound B. In one embodiment, Form G is a methanol solvate of Compound B. In another embodiment, Form F is a methanol solvate tartrate salt of Compound A.

[0284] In one embodiment, Form G of Compound B is obtained by slurring Compound B in methanol and slowly evaporating the methanol. The mixture can be filtered. Form G can be prepared according to the methods and examples described herein.

[0285] In one embodiment, a solid form provided herein, e.g., Form G, is a tartrate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD of a solid form provided herein, e.g., Form G, is substantially as shown in FIG. 19. In another embodiment, a solid form provided herein, e.g., Form G, has one or more characteristic XRPD peaks at approximately 7.65, 11.46, 12.51, 15.27, 15.51, 16.00, 17.34, 18.21, 19.11, 19.29, 19.42, 19.84, 20.23, 21.31, 21.57, 21.79, 22.49, 22.97, 23.22, 24.65, 25.04, or 25.88 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 19 and as found in Table 21 herein. In still another embodiment, a solid form provided herein, e.g., Form G, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 7.65, 11.46, 12.51, 15.27, 15.51, 16.00, 17.34, 18.21, 19.11, 19.29, 19.42, 19.84, 20.23, 21.31, 21.57, 21.79, 22.49, 22.97, 23.22, 24.65, 25.04, or 25.88 $\pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 19 and as found in Table 21 herein. In yet another embodiment, a solid form described herein, e.g., Form G, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, twenty, or all of the characteristic XRPD peaks as set forth in Table 21.

[0286] In still another embodiment, a solid form provided herein, e.g., Form G, has one, two,

three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 11.46, 12.51, 15.27, 16.00, 19.29, 19.42, 20.23, 22.49, 22.97, or $24.65 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 19. In still another embodiment, a solid form provided herein, e.g., Form G, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.46, 12.51, 19.29, 19.42, or $20.23 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 19. In still another embodiment, a solid form provided herein, e.g., Form G, has one, two, three, four, or five characteristic XRPD peaks at approximately 11.46, 12.51, 19.29, 19.42, or $20.23 \pm 0.05^\circ 2\theta$, as depicted in, for example, FIG. 19.

[0287] In one embodiment described herein, is a solid form, e.g., Form G, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 20. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 2% of the total mass of the sample.

[0288] In another embodiment described herein, is a solid form, e.g., Form G, having a DSC thermogram substantially as depicted in FIG. 20 comprising an endothermic event with an onset temperature of about 173°C and a peak maximum temperature of about 178°C .

[0289] In still another embodiment, Form G is pure. In certain embodiments, pure Form G is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, pure Form G is substantially free of Form B, Form D, or Form E. In certain embodiments, the purity of the Form G is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Compound C, Form 1

[0290] In certain embodiments, provided herein is a solid form of Compound C designated as Form 1. Form 1 is a crystalline solid form of Compound C.

[0291] In one embodiment, Form 1 of Compound C is obtained by slurring Compound C in isoamyl alcohol/water at about a 3:1 ratio for about 1.5 hours at about 55°C . The liquid of the mixture is then evaporated under flow of nitrogen and reduced pressure. In another embodiment, Form 1 of compound C is obtained by slurring Compound C in ethanol/heptane at a ratio of about 3:8 at RT. In still another embodiment, Form 1 of compound C is obtained by slurring Compound C in ethanol/heptane at a ratio of about 1: 1 at RT. Form 1 can be prepared according to the methods and examples described herein.

[0292] In one embodiment, a solid form provided herein, e.g., Form 1, is fumarate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD is substantially as shown in FIG. 27a. In another embodiment, a solid form provided herein, e.g., Form 1, has one or more characteristic XRPD peaks at approximately 7.58, 10.59, 11.44, 11.84, 12.5, 14.44, 15.45,

15.78, 16.09, 17.55, 18.92, 19.69, 19.86, 20.23, 21.35, 22.04, 23.16, 23.89, 24.23, 24.67, 25.23, or $25.93 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27a and as found in Table 36 herein. In still another embodiment, a solid form provided herein, e.g., Form 1, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 7.58, 10.59, 11.44, 11.84, 12.5, 14.44, 15.45, 15.78, 16.09, 17.55, 18.92, 19.69, 19.86, 20.23, 21.35, 22.04, 23.16, 23.89, 24.23, 24.67, 25.23, or $25.93 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27a and as found in Table 36 herein. In yet another embodiment, a solid form described herein, e.g., Form 1, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, twenty, or all of the characteristic XRPD peaks as set forth in Table 36.

[0293] In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 10.59, 15.45, 15.78, 16.09, 18.92, 19.69, 19.86, 21.35, 23.16, or $24.23 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27a. In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, or five characteristic XRPD peaks at approximately 16.09, 18.92, 19.69, 19.86, or $23.16 \pm 0.1^\circ 2\theta$ as depicted in, for example, FIG. 27a. In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, or five characteristic XRPD peaks at approximately 16.09, 18.92, 19.69, 19.86, or $23.16 \pm 0.05^\circ 2\theta$ as depicted in, for example, FIG. 27a.

[0294] In one embodiment described herein, is a solid form, e.g., Form 1, having a TGA thermograph corresponding substantially to the representative TGA thermogram as depicted in FIG. 28. In certain embodiments, the crystalline form exhibits a TGA thermogram comprising a total mass loss of approximately 2% of the total mass of the sample.

[0295] In another embodiment described herein, is a solid form, e.g., Form 1, having a DSC thermogram substantially as depicted in FIG. 28, comprising an endothermic event with an onset temperature of about 167°C and a peak maximum temperature of about 172°C .

[0296] In still another embodiment, Form 1 is pure. In certain embodiments, pure Form 1 is substantially free of other solid forms, e.g., amorphous solid. In certain embodiments, the purity of the Form 1 is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

[0297] In another embodiment described herein, is a solid form, e.g., Form 1, having a PLM image as depicted in FIG. 29.

Compound C, Form 2

[0298] In certain embodiments, provided herein is a solid form of Compound C designated as Form 2. Form 2 is a crystalline solid form of Compound C.

[0299] In one embodiment, a solid form provided herein, e.g., Form 2, is fumarate salt of Compound A, and is substantially crystalline, as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD is substantially as shown in FIG. 27b. In another embodiment, a solid form provided herein, e.g., Form 2, has one or more characteristic XRPD peaks at approximately 11.52, 11.87, 15.55, 16.04, 16.51, 17.32, 18.36, 19.00, 19.43, 19.87, 20.24, 21.35, 22.03, 23.23, 23.91, 25.43, or $26.03 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27b and as found in Table 37 herein. In still another embodiment, a solid form provided herein, e.g., Form 2, has at least 3, at least 5, at least 7, or at least 10 characteristic XRPD peaks at approximately 11.52, 11.87, 15.55, 16.04, 16.51, 17.32, 18.36, 19.00, 19.43, 19.87, 20.24, 21.35, 22.03, 23.23, 23.91, 25.43, or $26.03 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27b and as found in Table 37 herein. In yet another embodiment, a solid form described herein, e.g., Form 1, has one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or all of the characteristic XRPD peaks as set forth in Table 37 herein.

[0300] In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, five, six, seven, eight, nine, or ten characteristic XRPD peaks at approximately 11.87, 15.55, 16.04, 16.51, 17.32, 19.43, 19.87, 20.24, 23.23, or $23.91 \pm 0.1^\circ 2\theta$, as depicted in, for example, FIG. 27b. In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, or five characteristic XRPD peaks at approximately 15.55, 19.43, 19.87, 20.24, or $23.23 \pm 0.1^\circ 2\theta$ as depicted in, for example, FIG. 27b. In still another embodiment, a solid form provided herein, e.g., Form 1, has one, two, three, four, or five characteristic XRPD peaks at approximately 15.55, 19.43, 19.87, 20.24, or $23.23 \pm 0.05^\circ 2\theta$ as depicted in, for example, FIG. 27b.

Amorphous Form of Freebase Compound A

[0301] In certain embodiments, provided herein is an amorphous solid form of freebase Compound A.

[0302] In one embodiment, is an amorphous solid form of freebase Compound A as indicated by X-ray powder diffraction pattern (XRPD) measurements. In one embodiment, the XRPD is substantially as shown in FIG. 32.

[0303] In one embodiment described herein, is an amorphous solid form of freebase Compound A having a TGA thermograph corresponding substantially to the representative TGA and DSC thermogram as depicted in FIG. 33. In certain embodiments, the amorphous form exhibits a TGA thermogram comprising a total mass loss of approximately 9.3% of the total mass of the sample.

[0304] In still another embodiment, the amorphous solid form of Compound A is pure. In

certain embodiments, pure amorphous solid form of Compound A is substantially free of other solid forms, e.g., crystalline solids as described herein. In certain embodiments, the purity is no less than about 95%, no less than about 96%, no less than about 97%, no less than about 98%, no less than about 98.5%, no less than about 99%, no less than about 99.5%, or no less than about 99.9%.

Pharmaceutical Compositions

[0305] The compounds described herein can be administered, for example, orally, intramuscularly, subcutaneously, intravenously, intradermally, percutaneously, intraarterially, intraperitoneally, intralesionally, intracranially, intraarticularly, intraprostatically, intrapleurally, intratracheally, intrathecally, intranasally, intravaginally, intrarectally, topically, intratumorally, peritoneally, subconjunctivally, intravesicularly, mucosally, intrapericardially, intraumbilically, intraocularly, intraorbitally, intravitreally (e.g., by intravitreal injection), by eye drop, topically, transdermally, parenterally, by inhalation, by injection, by implantation, by infusion, by continuous infusion, by localized perfusion bathing target cells directly, by catheter, by lavage, in creams, or in lipid compositions. Compounds described herein can be formulated in pharmaceutical compositions as provided herein suitable for oral administration. In another embodiment, a compound described herein can be administered intramuscularly.

[0306] In one embodiment, compounds described herein are administered as pharmaceutical compositions capable of being administered to a subject orally or parenterally. Pharmaceutical compositions of the compounds described herein can be prepared as oral dosage forms such as, for example, capsules, microcapsules, tablets (coated and non-coated tablets), granules, powders, pills, or suppositories. The compounds described herein can be formulated for topical or parenteral use where the compound is dissolved or otherwise suspended in a solution suitable for injections, suspensions, syrups, creams, ointments, gels, sprays, solutions and emulsions.

[0307] Pharmaceutical compositions described herein include one or more pharmaceutically acceptable excipients such as, but not limited to: sucrose, starch, mannitol, sorbitol, lactose, glucose, cellulose, talc, calcium phosphate or calcium carbonate, cellulose, methylcellulose, hydroxymethylcellulose, hydroxypropylcellulose, carboxymethylcellulose, hydroxypropylstarch, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol (PEG), starch, sodium bicarbonate, calcium citrate, magnesium stearate, sodium lauryl sulfate, sodium benzoate, sodium bisulfite, methylparaben, propylparaben, citric acid, sodium citrate or acetic acid, polyvinyl pyrrolidone, aluminum stearate), water, and cocoa butter. Uses as, for example, diluents, binders, lubricants and disintegrators of such excipients is well known in the art.

[0308] The pharmaceutical compositions described herein include an effective amount of the compound described herein (e.g. Compound A, Compound B, Compound C, Compound D, or a solid form thereof). The dose of the compound described herein can be a measure of a specific amount of the compound (e.g. a standard dose amount) or can be measured as a

function of, for example, a patient's body weight. In one embodiment, a compound described herein is administered in an amount equivalent to about 0.1, 0.5, 0.75, 1, 2, 3, 4, 5, 10, 15, 20, 30, 50, 75, 100, 200, or 250 mg/kg. In another embodiment, a compound described herein is administered in an amount of about 0.1 mg/kg to about 1 mg/kg; about 0.5 mg/kg to about 2 mg/kg; about 1 mg/kg to about 5 mg/kg; about 3 mg/kg to about 10 mg/kg; about 8 mg/kg to about 15 mg/kg; or about 15 mg/kg to about 30 mg/kg. In still another embodiment, a compound described herein is administered in an amount less than about 100 mg/kg, less than about 50 mg/kg, less than about 30 mg/kg, less than about 10 mg/kg, or less than about 1 mg/kg.

[0309] In one embodiment, a compound described herein is administered at an amount of about 1, 5, 10, 20, 25, 30, 50, 60, 75, 90, 100, 120, 150, or 250 mg. In another embodiment, a compound described herein is administered in an amount of about 10 mg. In still another embodiment, a compound described herein is administered in an amount of about 30 mg. In still another embodiment, a compound described herein is administered in an amount of about 90 mg. In one embodiment, the pharmaceutical composition comprising the compound described herein is administered in an amount prescribed above once per day (QD). The compound can be Compound B, or a solid form thereof (e.g. Form A, Form B, Form C, Form D, Form E, Form F, or Form G). In another embodiment, the compound is a solid form of Compound B (e.g. Form B, Form D, or Form F). In one embodiment, the compound is Compound C or Compound C Form 1 or Form 2. In another embodiment, the compound is Compound D or a solid form described herein.


[0310] In another embodiment, a compound described herein is administered at an amount of about 1 mg to about 10 mg; about 10 mg to about 30 mg; about 10 mg to about 90 mg; about 30 mg to about 90 mg; or about 90 mg to about 250 mg. In one embodiment, the compound administered is Compound B at an amount of about 1, 10, 30, 50, 90, 100, or 150 mg. The doses of a compound described herein can be provided as a single dose (e.g. a single tablet or capsule of the given dosage amount) or can be provided as multiple doses given over a period of time (e.g. 2 or more tablets or capsules equating to the dosage amount). The compound can be Compound B, or a solid form thereof (e.g. Form A, Form B, Form C, Form D, Form E, Form F, or Form G). In another embodiment, the compound is a solid form of Compound B (e.g. Form B, Form D, or Form F). In one embodiment, the compound is Compound C or Compound C Form 1 or Form 2. In another embodiment, the compound is Compound D or a solid form described herein.

[0311] Pharmaceutical compositions described herein can be administered once daily (QD); twice daily (BID), thrice daily (TID), every other day (Q2D), every three days (Q3D), or once a week. Further, doses of pharmaceutical compositions provided herein comprising a compound described herein can be administered before food (ac), after food (pc), or with food. In one embodiment, a compound described herein is administered QD for a treatment period (a period of time where the drug is administered to a patient described herein) followed by a rest period (a period of time where the drug is not administered to a patient described herein). Rest periods may include administration of anti-cancer agents other than a compound described


herein. In one embodiment, a compound described herein is formulated for oral administration as provided herein and is administered QD for 20-28 days followed by a 3-10 day rest period. In another embodiment, the compound is administered QD with no rest period.

[0312] Preferably a compound described herein is formulated for oral administration. Oral administration can promote patient compliance in taking the compound (e.g. formulated as a pharmaceutical composition), thereby increasing compliance and efficacy. Oral pharmaceutical compositions comprising a compound described herein include, but are not limited to, tablets (e.g. coated, non-coated and chewable) and capsules (e.g. hard gelatin capsules, soft gelatin capsules, enteric coated capsules, and sustained release capsules). Tablets can be prepared by direct compression, by wet granulation, or by dry granulation. Oral pharmaceutical compositions comprising a compound described herein can be formulated as understood in the art for delayed or prolonged release. In one embodiment, Compound B or a solid form described herein (e.g. Form B, Form D, or Form F) is formulated as a tablet or a capsule for oral administration in an amount set forth herein.


[0313] Further provided herein are compounds having the formulae:




(M1)



(M2)



(M3)



(M4)

[0314] Compounds M1, M2, M3, and M4 can be considered metabolites and/or degradants of Compound B, including solid forms described herein. In certain instances, such compounds can be found in compositions described herein where such compositions have been stored for

a given period of time at a relative humidity (RH) of about 50%, 55%, 60%, 65%, 70%, 75%, 80% or more. Such compounds can also be found at elevated temperatures of about 30 °C, 35 °C, 40 °C, 45 °C, or about 50 °C. In one embodiment, compound M1, M2, M3, or M4 is found in composition described herein where the composition comprises less than about 30 mg of Compound B or a solid form thereof. In certain instances, such compounds are found in compositions where the composition comprises an uncoated tablet as described herein.

[0315] In certain embodiments, compositions described herein comprising Compound B or a solid form of Compound B described herein (e.g. Form A, Form B, Form C, Form D, Form E, Form F, or Form G) comprise less than 0.01%, 0.02%, 0.03%, 0.04, 0.05, 0.1, 0.15, 0.75, 0.2, 0.225, 0.25, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1% w/w of one or more of M1, M2, M3, or M4. In one embodiment, a composition described herein comprises less than about 0.5% w/w of one or more of M1, M2, M3, or M4.

[0316] Methods of Treating Cancer: the references to methods of treatment in the following paragraphs are to be interpreted as references to the compound, the solid form or the pharmaceutical composition thereof for use in a method for treatment of the human or animal body.

[0317] The compounds and solid forms described herein can be administered in an effective amount (e.g. an amount as described herein) for treating cancer. It is to be understood that the methods described herein also include treatment with a pharmaceutical composition as described herein comprising a compound (e.g. Compound B or a solid form thereof) described herein and one or more pharmaceutically acceptable excipients.

[0318] In one embodiment provided herein is a method of treating cancer by administering an effective amount of Compound B as described herein to a patient having cancer. In one embodiment, Compound B is a solid form as described herein (e.g. Form A, Form B, Form C, Form D, Form E, Form F, or Form G).

[0319] In another aspect provided herein Compound B, Form A can be administered as described herein to treat a patient having a cancer as set forth herein.

[0320] In another aspect provided herein Compound B, Form B can be administered as described herein to treat a patient having a cancer as set forth herein.

[0321] In another aspect provided herein Compound B, Form C can be administered as described herein to treat a patient having a cancer as set forth herein.

[0322] In another aspect provided herein Compound B, Form D can be administered as described herein to treat a patient having a cancer as set forth herein.

[0323] In another aspect provided herein Compound B, Form E can be administered as described herein to treat a patient having a cancer as set forth herein.

[0324] In another aspect provided herein Compound B, Form F can be administered as described herein to treat a patient having a cancer as set forth herein.

[0325] In another aspect provided herein Compound B, Form G can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0326] In another aspect provided herein an amorphous non-crystalline form of Compound A or Compound B can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0327] In another aspect provided herein is a method of treating lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer by administering an effective amount of Compound B or a solid form as described herein to a patient having said cancer. In one embodiment, the cancer is ovarian cancer or endometrial cancer. In one embodiment, the cancer is breast cancer. In one embodiment of such methods, Compound B is a solid form as described herein (e.g. Form A, Form B, Form C, Form D, Form E, Form F, or Form G).

[0328] In a further aspect, Compound B, Form A can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0329] In a further aspect, Compound B, Form B can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0330] In a further aspect, Compound B, Form C can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0331] In a further aspect, Compound B, Form D can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0332] In a further aspect, Compound B, Form E can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0333] In a further aspect, Compound B, Form F can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0334] In a further aspect, Compound B, Form G can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0335] In a further aspect, an amorphous non-crystalline form of Compound A or Compound B can be administered as described herein to treat a patient having lung cancer, ovarian cancer, endometrial cancer, prostate cancer, uterine cancer, or breast cancer as set forth herein.

[0336] Further provided herein are methods of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form of Compound B as described herein. In one embodiment is a method of treating breast cancer in such a patient by administering an effective amount of a solid form of Compound B as described herein. In one embodiment, the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form B as described herein. In one embodiment, the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of pure Compound B (e.g. substantially free of another solid form described herein, e.g. substantially free of Form D and/or Form F). In another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form D as described herein. In another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form F as described herein. In still another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form A as described herein. In still another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form C as described herein. In still another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form E as described herein. In still another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of Compound B, Form G, as described herein. In still another aspect the method comprises treating breast cancer in a patient having breast cancer by administering to the patient an effective amount of an amorphous non-crystalline form of Compound B as described herein.

[0337] The compounds described herein (e.g. Compound B or a solid form thereof as described herein) can be used in the manufacture of a medicament for use in treating breast cancer as described herein.

[0338] The methods of treating breast cancer provided herein comprise treatment where the breast cancer can be hormone receptor positive breast cancer (e.g. ER+ breast cancer), HER2-positive breast cancer, HER2-negative breast cancer, or triple negative breast cancer (TNBC).

[0339] In one embodiment, the breast cancer is HER2-negative breast cancer. HER2-negative

breast cancer can be defined herein as, for example, a HER2 IHC score of 0 or 1+, or an IHC score of 2+ accompanied by a negative fluorescence, chromogenic, or silver *in situ* hybridization test indicating the absence of HER2-gene amplification, or a HER2/CEP17 ratio of < 2.0, or local clinical guidelines. In one embodiment, the breast cancer is ER+/HER2- breast cancer. The breast cancer can be stage 0, I, II, III, or IV as understood in the art.

[0340] In another embodiment, the breast cancer is locally advanced or metastatic breast cancer (mBC).

[0341] In one embodiment, Compound B or a solid form thereof (e.g. Form B, Form D, or Form F) can be administered as a component of adjuvant therapy. In another embodiment, Compound B or a solid form thereof (e.g. Form B, Form D, or Form F) can be administered as a component of neoadjuvant therapy.

[0342] Breast cancer patients described herein may be premenopausal before treatment with a compound or solid form as described herein. Breast cancer patients described herein may be postmenopausal before treatment with a compound or solid form as described herein.

[0343] The methods provided herein include administering an effective amount of Compound B or a solid form as described herein to the patient at an amount as set forth herein. The effective amount can be, for example, an amount of about 10 mg, 30 mg, 50 mg, 90 mg, 100 mg, 125 mg, or 250 mg. In one embodiment of the methods provided herein, Compound B or a solid form described herein is administered orally. In one embodiment, Compound B or a solid form thereof is administered as a tablet (e.g. a coated or non-coated tablet). In another embodiment, Compound B or a solid form thereof is administered as a capsule. Thus, provided herein are compositions suitable for administration to a breast cancer patient where such compositions comprise an amount of Compound B or a solid form described herein of about 10 mg, 30 mg, 50 mg, 90 mg, 100 mg, 125 mg, or 250 mg in a tablet or capsule as set forth herein. When administered in accordance with the methods provided herein, Compound B or a solid form thereof can be pure as described herein.

[0344] Patients of the methods described above may have had previous treatment with one or more anti-cancer agents or radiation therapy. For example, in one embodiment, a patient may have been previously treated (e.g. with a 1L, 2L, 3L or more line therapy) with doxorubicin, pegylated liposomal doxorubicin, epirubicin, paclitaxel, albumin-bound paclitaxel, docetaxel, 5-fluorouracil, cyclophosphamide, cisplatin, carboplatin, vinorelbine, capecitabine, gemcitabine, ixabepilone, eribulin, olaparib, methotrexate, anastrozole, exemestane, toremifene, letrozole, tamoxifen, 4-hydroxy tamoxifen, raloxifene, droloxifene, trioxifene, keoxifene, ftutamide, nilutamide, bicalutamide, lapatinib, vinblastine, goserelin, leuprolide, pegfilgrastim, filgrastim, or venetoclax.

[0345] In another embodiment, a patient may have been previously treated (e.g. with a 1L, 2L, 3L or more line therapy) with an AKT inhibitor, a CDK4/6 inhibitor, a PARP inhibitor, or an aromatase inhibitor. In one embodiment, the AKT inhibitor is ipatasertib (GDC-0068). In one

embodiment, the CDK4/6 inhibitor is abemaciclib, ribociclib, or palbociclib. In certain instances, a patient may have been previously treated with: (1) abemaciclib, ribociclib, or palbociclib; (2) ipatasertib; (3) everolimus or fulvestrant; (4) -trastuzumab emtansine, trastuzumab, pertuzumab, or atezolizumab; or (5) alemtuzumab, bevacizumab, cetuximab, panitumumab, rituximab, tositumomab, or a combination thereof. Patients described herein may have had surgery prior to treatment with Compound B or the solid form thereof.

[0346] In another embodiment, a patient herein may be refractory to one or more anti-cancer therapies. For example, a patient herein may be refractory to aromatase inhibitors. In another example, a patient herein may be refractory to a selective estrogen receptor degrader (SERD) such as, for example, fulvestrant. In still another example, a patient may be refractory to one or more endocrine therapies such as, clomifene, toremifene, raloxifene, anordrin, bazedoxifene, broparexol, cyclofenil, lasofoxifene, ormeloxifene, acolbifene, elacestrant, brilanestrant, clomifenoxide, droloxifene, etacstil, or ospemifene. In another embodiment, a patient may be refractory to abemaciclib, anastrozole, exemestane, fulvestrant, goserelin, letrozole, leuprorelin, megestrol, palbociclib, tamoxifen, or toremifene. In another example, a patient may be refractory to treatment with trastuzumab emtansine, trastuzumab, pertuzumab, atezolizumab, pembrolizumab, durvalumab, avelumab, or nivolumab.

[0347] The compounds described herein can also be used in methods comprising inhibiting ER α in a patient. Such methods comprise administering an amount of a compound described herein (e.g. Compound A or Compound B, including solid forms thereof as described herein) to the patient.

Combination Therapies

[0348] The compounds and solid forms described herein can be administered in combination with one or more anti-cancer agents. Administration "in combination" as set forth herein includes sequential administration (in any order) of a compound described herein and one or more anti-cancer therapies as well as simultaneous administration. Accordingly, provided herein are methods of treating breast cancer in a patient having breast cancer, such methods comprising administering Compound B or a solid form as described herein in combination with one or more additional anti-cancer therapies. In one embodiment, the anti-cancer therapy comprises doxorubicin, pegylated liposomal doxorubicin, epirubicin, paclitaxel, albumin-bound paclitaxel, docetaxel, 5-fluorouracil, cyclophosphamide, cisplatin, carboplatin, vinorelbine, capecitabine, gemcitabine, ixabepilone, eribulin, olaparib, methotrexate, anastrozole, exemestane, toremifene, letrozole, tamoxifen, 4-hydroxy tamoxifen, raloxifene, droloxifene, trioxifene, keoxifene, ftutamide, nilutamide, bicalutamide, lapatinib, vinblastine, goserelin, leuprolide, pegfilgrastim, filgrastim, or venetoclax.

[0349] In one embodiment provided herein is a method of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form as described herein in combination with doxorubicin, pegylated liposomal doxorubicin, epirubicin,

paclitaxel, albumin-bound paclitaxel, docetaxel, 5-fluorouracil, cyclophosphamide, cisplatin, carboplatin, vinorelbine, capecitabine, gemcitabine, ixabepilone, eribulin, olaparib, methotrexate, anastrozole, exemestane, toremifene, letrozole, tamoxifen, 4-hydroxy tamoxifen, raloxifene, droloxifene, trioxifene, keoxifene, ftutamide, nilutamide, bicalutamide, lapatinib, vinblastine, goserelin, leuprolide, pegfilgrastim, filgrastim, or venetoclax.

[0350] In another aspect provided herein is a method of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form as described herein in combination with paclitaxel, albumin-bound paclitaxel, methotrexate, anastrozole, exemestane, toremifene, letrozole, tamoxifen, 4-hydroxy tamoxifen, raloxifene, droloxifene, trioxifene, keoxifene, or venetoclax. In still another aspect provided herein is a method of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form as described herein in combination with fulvestrant, paclitaxel, albumin-bound paclitaxel, clomifene, toremifene, raloxifene, anordrin, bazedoxifene, broparestrol, cyclofenil, lasofoxifene, ormeloxifene, acolbifene, elacestrant, brilanestrant, clomifenoxide, droloxifene, etacstil, or ospemifene.

[0351] In still another aspect provided herein is a method of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form as described herein in combination with a CDK4/6 inhibitor, a PARP inhibitor, or an aromatase inhibitor.

[0352] In a further aspect provided herein is a method of treating breast cancer in a patient having breast cancer as described herein where the method comprises administering an effective amount of Compound B or a solid form as described herein in combination with a CDK4/6 inhibitor where the CDK4/6 inhibitor is abemaciclib, ribociclib, or palbociclib. In a preferred embodiment, the method comprises administering Compound B or a solid form as described herein in combination with palbociclib. In still another embodiment, the method comprises administering Compound B or a solid form as described herein in combination with abemaciclib or ribociclib. In another aspect provided herein is a kit comprising (i) Compound B or a solid form thereof in a unit dosage form; (ii) a CDK4/6 inhibitor (e.g. palbociclib) in a second unit dosage form; and a container containing each dosage form.

[0353] The dose of abemaciclib may be 50 mg to 500 mg daily, or 150 mg to 450 mg daily and the dosing can be daily in 28 day cycles or less than 28 days per 28 day cycles such as 21 days per 28 day cycle or 14 days per 28 day cycle or 7 days per 28 day cycles. In one embodiment" abemaciclib is dosed once daily or preferably on a bid schedule where dosing is oral. In the case of bid dosing, the doses can be separated by 4 hours, 8 hours or 12 hours. In certain embodiments, abemaciclib is dosed at 150 mg orally bid where each dose is administered about 12 hr apart. In certain embodiments, the dose of abemaciclib is administered in accordance with a package insert.

[0354] The dose of ribociclib may be 200 mg to 1,000 mg daily; or 250 mg to 750 mg daily and the dosuig can be daily in 28 day cycles or less than 28 days per 28 day cycles such as 21

days per 28 day cycle or 14 days per 28 day cycle or 7 days per 28 day cycles. In one embodiment" ribociclib is dosed once daily where dosing is oral. In certain embodiments, the dose of ribociclib is administered in accordance with a package insert.

[0355] The dose of palbociclib may be 25 mg to 250 mg daily or 50 mg to 125 mg daily or from 75 mg to 125 mg daily or 75 mg daily to 100 mg daily or 125 mg daily. The dosing can be daily in 28 day cycles or less than 28 days per 28 day cycles such as 21 days per 28 day cycle or 14 days per 28 day cycle or 7 days per 28 day cycles. In one embodiment" palbociclib is dosed once daily where dosing is oral. In certain embodiments, the dose of palbociclib is administered in accordance with a package insert.

[0356] In another aspect provided herein the methods described herein comprise administering an effective amount of Compound B or a solid form as described herein in combination with an aromatase inhibitor (AI), where the AI is letrozole, anastrozole, exemestane, or testolactone.

[0357] In yet another aspect provided herein is a method of treating breast cancer in a patient having breast cancer by administering an effective amount of Compound B or a solid form as described herein in combination with a cancer immunotherapy (e.g. an antibody). In one embodiment, Compound B or a solid form as described herein is administered in combination with trastuzumab emtansine, trastuzumab, pertuzumab, atezolizumab, pembrolizumab, durvalumab, avelumab, or nivolumab, or a combination thereof. In one embodiment, Compound B or a solid form as described herein is administered in combination with a cancer immunotherapy comprising PD-1 or PD-L1 inhibitor, where the cancer immunotherapy is atezolizumab, pembrolizumab, or nivolumab.

[0358] Administration of a compound described herein (e.g. Compound B or a solid form as described herein) results in patients having adverse effects (AEs) characterized as Grade 2 or lower. In one embodiment, a patient administered Compound B or a solid form as described herein has Grade 2 or lower AEs.

Examples:

[0359] The following Examples are presented by way of illustration, and any example falling outside the scope of the appended claims is provided for information purposes only.

[0360] Synthesis of Compounds described herein. All reagents and solvents were purchased from commercial suppliers and used with no additional purification. Anhydrous solvent (dichloromethane) was utilized. Commercially available solvents were not further purified.

[0361] All reactions were carried out in screw-cap vials equipped with a Teflon septa under a nitrogen atmosphere.

[0362] Flash column chromatography was performed using a teledyne Isco CombiFlash^(R) Rf instrument with pre-packed RediSepRf Gold silica cartridges.

[0363] Unless otherwise indicated, reported yields are for isolated material and are corrected for residual solvents.

[0364] Compounds were characterized by one or more of ¹H NMR, ¹³C NMR, melting point and HRMS, and HPLC analysis (e.g., for confirmation of purity).

[0365] ¹H, and ¹³C Nuclear Magnetic Resonance Spectra were recorded on a Bruker 400 MHz instrument at ambient temperature. All ¹H NMR spectra were measured in parts per million (ppm) relative to residual chloroform signal (7.26 ppm) or dimethyl sulfoxide (2.50 ppm) in the deuterated solvent unless otherwise stated. Data for ¹H NMR are reported as follows: chemical shift, multiplicity (br = broad signal, overlap = overlapping, s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, m = multiplet), coupling constants and integration. All ¹³C NMR spectra are reported in ppm relative to deuteriochloroform (77.06 ppm) or deuterated dimethyl sulfoxide (39.53 ppm), and were obtained with complete ¹H decoupling unless otherwise stated. HPLC analyses were performed on an Agilent 1260 Infinity HPLC system with a UV detector at 220 nm using an Ace Super C18 column. Melting points were obtained using a Buchi B-540 Melting Point Apparatus and are uncorrected. High resolution mass spectrometry (HRMS) data was acquired on a Thermo Scientific Orbitrap Fusion mass spectrometer.


Examples 1-3: Indole Alkylation.

[0366] Indole alkylation was done by the sequence of Boc protection (Example 1), sulfamide formation (Example 2), and indole alkylation (Example 3)

Example 1: Boc Protection


Example 1: Boc protection general reaction scheme.

[0367] Boc protection was done according to the following general reaction scheme where R_A and R_B correspond to the various functional groups in the following Example 1 protection reactions, and wherein the asterisks represent a chiral center:



Example 1A: Preparation of tert-butyl (S)-(2-hydroxy-1-(4-methoxyphenyl)ethyl)carbamate:


[0368]




[0369] General reaction scheme 1 above was performed as follows. To a slurry of (S)-2-amino-2-(4-methoxyphenyl)ethan-1-ol hydrochloride (1.04 g, 5.10 mmol, 100 mol%) in THF (4.4 mL) was added Boc_2O (1.21 mL, 5.61 mmol, 110 mol%), NaHCO_3 (451 mg, 5.10 mmol, 100 mol%), and water (4.4 mL) at rt. The solution was stirred at rt for 18 h, and extracted with *i*PrOAc (20 mL x 2). The organic layer was washed with saturated brine (20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure to provide the product without further purification. Example 1A yield is reported are corrected based on residual solvent from ^1H NMR. The reaction yielded tert-butyl (S)-(2-hydroxy-1-(4-methoxyphenyl)ethyl)carbamate (1.36 g, 100% yield) as a white solid. mp: 139.0-139.9 °C; FTIR (neat, cm^{-1}) 3370, 2984, 2837, 1681, 1613, 1512, 1461; ^1H NMR (400 MHz, CDCl_3): δ 7.24-7.20 (m, 2H), 6.92-6.86 (m, 2H), 5.10 (d, $J = 7.2$ Hz, 1H), 4.72 (br, 1H), 3.82 (t, $J = 5.6$ Hz, 2H), 3.80 (s, 3H), 2.35 (br, 1H), 1.43 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 159.1, 156.2, 131.6, 127.7, 114.2, 80.0, 66.8, 56.4, 55.3, 28.4.

Example 1B: Preparation of tert-butyl (R)-(1-cyclopropyl-2-hydroxyethyl)carbamate:


[0370]



[0371] The general reaction scheme as per Example 1A above was performed with (R)-2-amino-2-cyclopropylethan-1-ol (1.16 g, 11.5 mmol, 100 mol%) to yield tert-butyl (R)-(1-cyclopropyl-2-hydroxyethyl)carbamate (2.31 g, 100% yield) as a white solid. mp: 70.0-70.8 °C; FTIR (neat, cm^{-1}) 3358, 2974, 2937, 1682, 1521, 1366; ^1H NMR (400 MHz, CDCl_3): δ 4.80 (br, 1H), 3.80 (ddd, $J = 10.8, 6.8, 3.2$ Hz, 1H), 3.67 (ddd, $J = 10.8, 6.0, 4.8$ Hz, 1H), 2.94 (dtd, $J = 9.6, 6.4, 3.2$ Hz, 1H), 2.81 (br, 1H), 1.45 (s, 9H), 0.85 (dtt, $J = 9.6, 8.0, 4.8$ Hz, 1H), 0.60-0.47 (m, 2H), 0.44-0.25 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3): δ 156.6, 79.7, 66.3, 57.9, 28.4, 13.0, 3.3, 2.9.


Example 1C: Preparation of tert-butyl ((1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl)carbamate:**[0372]**

[0373] The general reaction scheme as per Example 1A above was performed with (1*R*,2*S*)-1-amino-2,3-dihydro-1*H*-inden-2-ol (5.15 g, 34.5 mmol, 100 mol%) to yield tert-butyl ((1*R*,2*S*)-2-hydroxy-2,3-dihydro-1*H*-inden-1-yl)carbamate (8.61 g, 100% yield) as a white solid. mp: 67.3-68.4 °C; FTIR (neat, cm⁻¹) 3428, 3350, 2983, 2933, 1688, 1509; ¹H NMR (400 MHz, CDCl₃): δ 7.31-7.26 (m, 1H), 7.26-7.18 (m, 3H), 5.17 (br, 1H), 5.05 (br, 1H), 4.57 (ddd, *J* = 7.2, 4.8, 2.0 Hz, 1H), 3.12 (dd, *J* = 16.8, 5.2 Hz, 1H), 2.91 (dd, *J* = 16.8, 2.4 Hz, 1H), 2.31 (d, *J* = 4.8 Hz, 1H), 1.50 (s, 9H); ¹³C NMR (100 MHz, CDCl₃): δ 156.3, 140.9, 139.9, 128.2, 127.1, 125.3, 124.5, 79.9, 73.6, 58.9, 39.4, 28.4.

Example 1D: Preparation of tert-butyl (S)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate:**[0374]**

[0375] The general reaction scheme as per Example 1A above was performed with (S)-2-amino-2-(3-fluorophenyl)ethan-1-ol (1.36 g, 8.73 mmol, 100 mol%) to yield tert-butyl (S)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate (2.23 g, 100% yield) as a white solid. mp: 106.5-107.9 °C; FTIR (neat, cm⁻¹) 3251, 3059, 2977, 2901, 1671, 1587, 1543; ¹H NMR (400 MHz, CDCl₃): δ 7.33 (ddd, *J* = 7.6, 7.6, 6.0 Hz, 1H), 7.12-7.07 (m, 1H), 7.05-6.95 (m, 2H), 5.24 (br, 1H), 4.77 (br, 1H), 3.93-3.77 (m, 2H), 2.02 (br, 1H), 1.44 (s, 9H); ¹³C NMR (100 MHz, CDCl₃): δ 163.0 (d, ¹*J*_{CF} = 246 Hz), 156.0, 142.5, 130.2 (d, ³*J*_{CF} = 9 Hz), 122.2 (d, ⁴*J*_{CF} = 3 Hz), 114.5 (d, ²*J*_{CF} = 21 Hz), 113.6 (d, ²*J*_{CF} = 21 Hz), 80.2, 66.2, 56.3, 28.3; ¹⁹F NMR (CDCl₃, 376 MHz): δ -112.4.


Example 1E: Preparation of (S)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate**[0376]**



[0377] The general reaction scheme as per Example 1A above was performed with (*S*)-2-amino-2-(3-(trifluoromethyl)phenyl)ethan-1-ol hydrochloride (1.00 g, 4.12 mmol, 100 mol%) to yield (*S*)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate (1.26 g, 100% yield) as a white solid. mp: 50.3-52.5 °C; FTIR (neat, cm^{-1}) 3368, 3254, 2979, 2939, 1691, 1510, 1453, 1333; ^1H NMR (400 MHz, CDCl_3): δ 7.59-7.54 (m, 2H), 7.53-7.45 (m, 2H), 5.31 (d, $J = 6.4$ Hz, 1H), 4.83 (br, 1H), 3.92 (ddd, $J = 11.2, 6.8, 4.0$ Hz, 1H), 3.88-3.79 (m, 1H), 1.94 (br, 1H), 1.43 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 156.1, 141.1, 130.9 (q, $^2J_{\text{CF}} = 32$ Hz), 130.1, 129.0, 124.3 (q, $^3J_{\text{CF}} = 4$ Hz), 124.1 (q, $^1J_{\text{CF}} = 270$ Hz), 123.4 (q, $^3J_{\text{CF}} = 4$ Hz), 80.3, 65.8, 56.3, 28.2; ^{19}F NMR (CDCl_3 , 376 MHz): δ -62.6. Example 2: Sulfamidate Formation


Example 2: Sulfamidate formation general reaction scheme

[0378] Sulfamidate formation was done according to the following general reaction scheme where R_A and R_B correspond to the various functional groups in the following Example 2 reactions, and wherein the asterisks represent a chiral center:



Example 2A: Preparation of tert-butyl (*R*)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:

[0379]




[0380] General reaction scheme 2 above was performed as follows. To a cold (-40 °C) solution of SOCl_2 (10.9 mL, 149 mmol, 250 mol%) in CH_2Cl_2 (60.0 mL) was added a solution of *tert*-butyl (*R*)-(1-hydroxy-3-phenylpropan-2-yl)carbamate (15.0 g, 59.7 mmol, 100 mol%) in CH_2Cl_2 (60.0 mL) over 60 min at -40 °C. Pyridine (25.3 mL, 313 mmol, 525 mol%) was then added to the reaction mixture over 30 min at -40 °C. The reaction mixture was stirred at -40 °C for 2 h,

solvent swapped to $\text{CH}_2\text{Cl}_2/i\text{PrOAc}$ (1:1) mixture, filtered. The filtrate was washed with saturated brine solution (20 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was dissolved in CH_3CN (60.0 mL) at 0 °C. NaIO_4 (14.0 g, 65.7 mmol, 110 mol%), RuCl_3 (61.9 mg, 0.298 mmol, 0.5 mol%), and water (60.0 mL) were added into the reaction mixture at 0 °C and stirred for 15 min. The reaction mixture was then warmed to room temperature and stirred at room temperature for 2 h, extracted with $i\text{PrOAc}$ (20 mL), washed with saturated NaHCO_3 solution (15 mL), saturated brine solution (15 mL), dried (Na_2SO_4), filtered, purified by chromatography on SiO_2 . Specific gradient used for each sample is included in the characterization data. All the yields reported are corrected based on residual solvent from ^1H NMR. Reaction 2A yielded *tert*-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (18.7 g, 56% yield) as a white solid. Column Gradient: 0 to 5% CH_3OH in CH_2Cl_2 . mp: 134.4-135.0 °C; FTIR (neat, cm^{-1}) 3261, 2979, 2903, 1712, 1673, 1540; ^1H NMR (400 MHz, CDCl_3): δ 7.38-7.20 (m, 5H), 4.49-4.40 (m, 2H), 4.35-4.28 (m, 1H), 3.37 (dd, $J = 14.0, 4.0$ Hz, 1H), 2.98-2.87 (m, 1H), 1.56 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 148.5, 135.2, 129.5, 129.1, 127.5, 85.6, 68.8, 58.6, 37.9, 28.0.

Example 2B: Preparation of *tert*-butyl (S)-4-phenyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0381]




[0382] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (S)-(2-hydroxy-1-phenylethyl)carbamate (10.0 g, 42.1 mmol, 100 mol%) to yield compound *tert*-butyl (S)-4-phenyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (5.23 g, 42% yield) as a white solid. Column Gradient: 0 to 5% CH_3OH in CH_2Cl_2 . mp: 144.3-145.0 °C; FTIR (neat, cm^{-1}) 2976, 1722, 1458, 1377; ^1H NMR (400 MHz, CDCl_3): δ 7.44-7.35 (m, 5H), 5.28 (dd, $J = 6.4, 4.0$ Hz, 1H), 4.87 (dd, $J = 9.2, 6.4$ Hz, 1H), 4.39 (dd, $J = 9.2, 4.4$ Hz, 1H), 1.42 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 148.3, 137.0, 129.2, 129.1, 126.2, 85.5, 71.8, 60.8, 27.8.

Example 2C: Preparation of *tert*-butyl (S)-4-(4-methoxyphenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:

[0383]






[0384] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (S)-(2-hydroxy-1-(4-methoxyphenyl)ethyl) carbamate (1.45 g, 5.42 mmol, 100 mol%) to yield compound *tert*-butyl (S)-4-(4-methoxyphenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (1.05 g, 59% yield) as a white solid. Column Gradient: 0 to 5% CH₃OH in CH₂Cl₂. mp: 151.6-153.0 °C; FTIR (neat, cm⁻¹) 2979, 2933, 2838, 1721, 1636, 1510, 1457; ¹H NMR (400 MHz, CDCl₃): δ 7.37-7.32 (m, 2H), 6.95-6.90 (m, 2H), 5.24 (dd, *J* = 6.8, 4.4 Hz, 1H), 4.84 (dd, *J* = 9.2, 6.8 Hz, 1H), 4.39 (dd, *J* = 9.2, 4.4 Hz, 1H), 3.82 (s, 3H), 1.44 (s, 9H); ¹³C NMR (100 MHz, CDCl₃): δ 160.2, 148.3, 128.9, 127.7, 114.6, 85.5, 72.0, 60.5, 55.4, 27.9.

Example 2D: Preparation of *tert*-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0385]



[0386] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (R)-(1-hydroxypropan-2-yl)carbamate (5.00 g, 28.5 mmol, 100 mol%) to yield compound *tert*-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (3.85 g, 57% yield) as a white solid. Column Gradient: 0 to 5% CH₃OH in CH₂Cl₂. FTIR (neat, cm⁻¹) 3245, 2982, 1719, 1402, 1329; ¹H NMR (400 MHz, CDCl₃): δ 4.66 (dd, *J* = 9.2, 6.0 Hz, 1H), 4.41 (qdd, *J* = 6.4, 6.0, 2.8 Hz, 1H), 4.19 (dd, *J* = 9.2, 2.8 Hz, 1H), 1.54 (s, 9H), 1.50 (d, *J* = 6.4 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃): δ 148.5, 85.4, 71.4, 53.8, 28.0, 18.3.

Example 2E: Preparation of *tert*-butyl (R)-4-isopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:

[0387]




[0388] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (R)-(1-hydroxy-3-methylbutan-2-yl)carbamate (5.00 g, 24.6 mmol, 100 mol%) to yield

compound tert-butyl (R)-4-isopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (3.93 g, 60% yield) as a white solid. Column Gradient: 0 to 5% CH₃OH in CH₂Cl₂. mp: 104.8-105.8 °C; ¹H NMR (400 MHz, CDCl₃): δ 4.55 (dd, *J* = 9.6, 6.4 Hz, 1H), 4.38 (dd, *J* = 9.6, 2.0 Hz, 1H), 4.17 (ddd, *J* = 6.4, 5.2, 1.6 Hz, 1H), 2.24 (qqd, *J* = 6.8, 6.8, 5.2 Hz, 1H), 1.53 (s, 9H), 1.00 (d, *J* = 6.8 Hz, 3 H), 0.95 (d, *J* = 6.8 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃): δ 149.1, 85.3, 67.0, 62.0, 30.0, 27.9, 18.0, 16.4.

Example 2F: Preparation of tert-butyl (R)-4-cyclopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0389]



[0390] The general reaction scheme as per Example 2A above was performed with tert-butyl (R)-(1-cyclopropyl-2-hydroxyethyl)carbamate (2.31 g, 11.5 mmol, 100 mol%) to yield tert-butyl (R)-4-cyclopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (1.36 g, 45% yield) as a white solid. Column Gradient: 0 to 5% CH₃OH in CH₂Cl₂. mp: 52.7-55.7 °C; FTIR (neat, cm⁻¹) 2977, 1734, 1460, 1363; ¹H NMR (400 MHz, CDCl₃): δ 4.64 (dd, *J* = 9.2, 6.0 Hz, 1H), 4.40 (dd, *J* = 8.8, 2.0 Hz, 1H), 3.77 (ddd, *J* = 9.2, 6.0, 2.0 Hz, 1H), 1.54 (s, 9H), 1.35-1.23 (m, 1 H), 0.74-0.65 (m, 2H), 0.63-0.54 (m, 1H), 0.29-0.20 (m, 1H); ¹³C NMR (100 MHz, CDCl₃): δ 148.9, 85.3, 71.1, 61.6, 27.9, 14.3, 4.4, 1.7.

Example 2G: Preparation of tert-butyl (3aR,8aS)-8,8a-dihydroindeno[1,2-d][1,2,3]oxathiazole-3(3aH)-carboxylate 2,2-dioxide:

[0391]




[0392] The general reaction scheme as per Example 2A above was performed with tert-butyl ((1R,2S)-2-hydroxy-2,3-dihydro-1H-inden-1-yl)carbamate (9.12 g, 36.6 mmol, 100 mol%) to yield tert-butyl (3aR,8aS)-8,8a-dihydroindeno[1,2-d][1,2,3]oxathiazole-3(3aH)-carboxylate 2,2-dioxide (7.50 g, 66% yield) as a white solid. Column Gradient: 0 to 5% CH₃OH in CH₂Cl₂. mp: 134.2-135.0 °C; FTIR (neat, cm⁻¹) 2988, 2937, 1732, 1462, 1375; ¹H NMR (400 MHz, CDCl₃): δ 7.62-7.57 (m, 1H), 7.39-7.24 (m, 3H), 5.71 (d, *J* = 5.6 Hz, 1H), 5.50 (dt, *J* = 6.0, 3.2 Hz, 1 H),

3.38 (d, $J = 3.2$ Hz, 1 H), 1.62 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 149.6, 138.4, 137.9, 129.9, 128.4, 126.2, 125.2, 85.7, 82.2, 65.0, 36.5, 28.0.

Example 2H: Preparation of tert-butyl (S)-5-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0393]



[0394] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (S)-(2-hydroxypropyl)carbamate (6.25 g, 35.7 mmol, 100 mol%) to yield compound *tert*-butyl (S)-5-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (6.10 g, 72% yield) as a white solid. Column Gradient: 0 to 5% CH_3OH in CH_2Cl_2 . mp: 116.9-118.2 °C; FTIR (neat, cm^{-1}) 3370, 2956, 2938, 2837, 1681, 1512, 1461, 1366; ^1H NMR (400 MHz, CDCl_3): δ 5.00-4.90 (m, 1H), 4.06 (dd, $J = 9.6, 5.6$ Hz, 1H), 3.63 (dd, $J = 9.6, 9.2$ Hz, 1H), 1.56 (d, $J = 6.4$ Hz, 3H), 1.53 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 148.6, 85.3, 76.2, 51.7, 27.9, 18.0.

Example 21: Preparation of tert-butyl (S)-4-(3-fluorophenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0395]



[0396] The general reaction scheme as per Example 2A above was performed with *tert*-butyl (S)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate (1.45 g, 5.68 mmol, 100 mol%) to yield *tert*-butyl (S)-4-(3-fluorophenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (0.702 g, 39% yield) as a white solid. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 112.9-114.3 °C; FTIR (neat, cm^{-1}) 2976, 1722, 1636, 1594, 1458; ^1H NMR (400 MHz, CDCl_3): δ 7.44-7.36 (m, 1H), 7.24-7.18 (m, 1H), 7.17-7.11 (m, 1H), 7.11-7.05 (m, 1H), 5.28 (dd, $J = 6.8, 3.6$ Hz, 1H), 4.88 (dd, $J = 9.2, 6.8$ Hz, 1H), 4.39 (dd, $J = 9.2, 3.6$ Hz, 1H), 1.47 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 163.2 (d, $^1J_{\text{CF}} = 246$ Hz), 148.2, 139.6 (d, $^3J_{\text{CF}} = 7$ Hz), 131.1 (d, $^3J_{\text{CF}} = 8$ Hz), 121.8 (d, $^4J_{\text{CF}} = 3$ Hz), 116.2 (d, $^2J_{\text{CF}} = 21$ Hz), 113.4 (d, $^2J_{\text{CF}} = 22$ Hz), 86.0, 71.6, 60.2 (d, $^4J_{\text{CF}} = 3$ Hz), 27.9; ^{19}F NMR (CDCl_3 , 376 MHz): δ -111.0.

Example 2J: Preparation of tert-butyl (S)-4-(3-(trifluoromethyl)phenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide:


[0397]



[0398] The general reaction scheme as per Example 2A above was performed with tert-butyl (S)-(1-(3-fluorophenyl)-2-hydroxyethyl)carbamate (1.00 g, 3.28 mmol, 100 mol%) to yield tert-butyl (S)-4-(3-(trifluoromethyl)phenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (0.528 g, 44% yield) as a white solid. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 92.0-92.6 °C; FTIR (neat, cm^{-1}) 2989, 1720, 1463, 1373, 1325; ^1H NMR (400 MHz, CDCl_3): δ 7.70-7.61 (m, 3H), 7.61-7.55 (m, 1H), 5.35 (dd, $J = 6.8, 4.0$ Hz, 1H), 4.92 (dd, $J = 9.2, 6.8$ Hz, 1H), 4.41 (dd, $J = 9.2, 3.6$ Hz, 1H), 1.46 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 148.2, 138.2, 131.8 (q, $^2J_{\text{CF}} = 32$ Hz), 130.1, 129.4, 126.1 (q, $^3J_{\text{CF}} = 4$ Hz), 123.7 (q, $^1J_{\text{CF}} = 270$ Hz), 123.4 (q, $^3J_{\text{CF}} = 4$ Hz), 86.2, 71.4, 60.2, 27.8; ^{19}F NMR (CDCl_3 , 376 MHz): δ -62.8.

Example 2K: Preparation of tert-butyl (S)-4-phenyl-1,2,3-oxathiazinane-3-carboxylate 2,2-dioxide:

[0399]



[0400] The general reaction scheme as per Example 2A above was performed with tert-butyl (S)-(3-hydroxy-1-phenylpropyl)carbamate (2.00 g, 7.96 mmol, 100 mol%) to yield tert-butyl (S)-4-phenyl-1,2,3-oxathiazinane-3-carboxylate 2,2-dioxide (1.26 g, 51% yield) as a white solid. Column Gradient: 0 to 50% *i*PrOAc. mp: 128.6-129.9 °C; FTIR (neat, cm^{-1}) 2986, 1727, 1449, 1367; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) (94:6 mixture of rotamers): δ 7.48-7.35 (m, 4H), 7.35-7.23 (m, 1H), 5.65 (dd, $J = 4.4, 4.4$ Hz, 0.94H), 5.53 (dd, $J = 11.2, 4.4$ Hz, 0.06H), 4.70 (ddd, $J = 10.4, 7.2, 2.8$ Hz, 0.94H), 4.51 (ddd, $J = 8.8, 4.4, 4.4$ Hz, 0.06H), 4.40 (ddd, $J = 10.4, 10.4, 6.8$ Hz, 1H), 2.80-2.68 (m, 1H), 2.66-2.56 (m, 1H), 1.41 (s, 8.46H), 1.12 (s, 0.54H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) (rotamers): δ 155.1, 143.0, 128.4(128.6), 127.0(127.3),

126.3(125.4), 78.1(84.3), 73.6(70.9), 50.5(60.0), 36.0, 28.2(27.4).

Example 3: Indole alkylation


[0401] Chiral tryptamines are frequently encountered in pharmacology due to their significant biological activities in the central nervous system. In addition, the chiral tryptamine moiety serves as synthetic precursor of many medicinally important indole alkaloids and found in numerous biologically active natural products and pharmaceuticals.

[0402] In particular, stereocontrolled synthesis of tetrahydro- β -carboline-containing compounds such as those described herein often relies on diastereoselective Pictet-Spengler reaction, which in turn requires enantiomerically pure tryptamines as starting material. The latter is typically prepared in a non-stereoselective manner via a multistep sequence involving hazardous nitroalkane reagents. Therefore, the development of a simple regioselective alkylation approach which involves readily available non-protected indoles as nucleophiles and chiral amine-derived electrophiles, such as chiral aziridines and cyclic sulfamidates, provided a convenient access to these valuable chiral scaffolds.


[0403] Aziridine electrophiles have been applied to provide C³-selective alkylation under Lewis acidic conditions; however, this method is only applicable to the synthesis of β -substituted tryptamines as the displacement occurs preferentially at the more substituted carbon. Discovered herein, low order cuprate of indoles in combination with chiral cyclic sulfamidates successfully provided a practical access to both α - and β -substituted chiral tryptamines having high regioselectivity.

[0404] It was found that the reaction with chiral aziridines as the electrophile always led to a mixture of α - and β -substituted tryptamines while reaction with cyclic sulfamidates as the electrophile unambiguously reacted to displace C-O bond.

[0405] Contrary to the literature precedence suggesting preferential alkylation at the C³-position when Grignard reagents were used as base, initial attempts resulted in lower than expected yields and poor site-selectivity. Without being bound by any particular theory, softer indole nucleophiles were tried as they would likely prefer to react as carbon-centered nucleophile. Various additives including Cu and Zn salts were surveyed. Interestingly, a mixed halide system such as MeMgCl in combination with CuBr or CuI, or MeMgBr in combination with CuCl was much less efficient than the chloride-only system. Other copper salts, such as CuCl₂, CuCN, CuTC, or Cu(SCN) were also inferior to CuCl.



entry	base / additive / temperature	isolated yield	C ³ / N ¹ ^b
1	MeMgCl / None / -10 °C	14%	31 / 69
2	MeMgCl / CuCl / -10 °C	66%	95 / 5
3	MeMgCl / CuBr / -10 °C	37%	95 / 5
4	MeMgCl / CuI / -10 °C	26%	90 / 10
5 ^c	MeMgCl / CuCl / -10 °C	38%	67 / 33
6	MeMgCl / ZnCl ₂ / -10 °C	38%	22 / 78
7	MeMgCl / ZnBr ₂ / -10 °C	28%	37 / 63
8	MeMgBr / CuCl / -10 °C	26%	45 / 54
9	PhMgCl / CuCl / -10 °C	66%	96 / 4
10	MeLi / CuCl / -10 °C	N.D.	11 / 89
11	MeMgCl / CuCl / -40 °C	N.D.	42 / 58
12 ^d	MeMgCl / CuCl / -20 °C	76%	97 / 3
13	MeMgCl / CuCl / 0 °C	65%	95 / 5



[0406] A catalytic amount of CuCl was not tolerated and resulted in significant decrease in yield and selectivity. Notably, a reversal of regioselectivity was observed when zinc halides were used instead of CuCl (see above). The same was true when MeLi was used as base instead of MeMgCl, while the use of other Grignard reagent was well-tolerated. The effect of reaction temperature was evaluated and established that the displacement reaction performed optimally at around -20 °C. At below -30 °C, a significant drop in regioselectivity was observed presumably due to the incomplete cuprate formation.


[0407] The Cu-mediated indole alkylation tolerated a variety of substitution both on the indole nucleophile as well as on the cyclic sulfamidate providing the C³-alkylated indole products in moderate to good yield and excellent regioselectivity. For instance, indoles with either electron-donating or electron-withdrawing substituents participated in the reaction well (Exemplary compounds **6b-6g** above), and sterically demanding substrates also worked reasonably well (Exemplary compounds **6h** and **6i** above).

[0408] On the other hand, azaindoles appear to be possible poor substrates. Under standard reaction conditions, 6-azaindole provided only 8% of the alkylation product albeit with comparable regioselectivity (see e.g. exemplary compound **6j** above). Other azaindoles such as indazole and 7-azaindole failed to produce any desired alkylated products

[0409] A variety of sulfamidates were tested in the reaction successfully. Both aryl- and alkyl-substituted sulfamidates, prepared in two step sequence from the corresponding amino alcohols, were converted smoothly to the respective α -substituted chiral tryptamines. Similarly, 6-membered cyclic sulfamidate also participated in the alkylation well producing a homologated tryptamine in good yield and regioselectivity. This alkylation process can be also applied to get access to β -substituted, and α,β -disubstituted tryptamines (see exemplary compounds **6s** and **6t** above). In these cases, the indole nucleophile added to the corresponding cyclic sulfamidate with the inversion of stereochemistry at the carbon bearing oxygen with complete stereospecificity. The utility of this alkylation process was demonstrated as provided herein.


Example 3: Indole alkylation general reaction scheme

[0410] Indole alkylation was done according to the following general reaction scheme where R_A and R_B correspond to the various functional groups in the following Example 3 reactions, and wherein the asterisks represent a chiral center:




Example 3A: Preparation of tert-butyl (R)-(1-(1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0411]



[0412] General reaction scheme 3 above was performed as follows. To a cold ($0\text{ }^\circ\text{C}$) mixture of indole (280 mg, 2.39 mmol, 150 mol%), and CuCl (193 mg, 1.95 mmol, 130 mol%) in CH_2Cl_2 (3.0 mL) was added MeMgCl (3.0 M in THF, 0.65 mL, 1.95 mmol, 130 mol%) over 10 min at $0\text{ }^\circ\text{C}$. The reaction mixture was stirred at $0\text{ }^\circ\text{C}$ for 1 h and cooled to $-20\text{ }^\circ\text{C}$. A solution of tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (500 mg, 1.60 mmol, 100 mol%) in CH_2Cl_2 (2.0 mL) was added into the reaction mixture over 30 min at $-20\text{ }^\circ\text{C}$. The reaction mixture was then stirred at $-20\text{ }^\circ\text{C}$ for 18 h, quenched with 10% aqueous citric acid (5.0 mL) at $0\text{ }^\circ\text{C}$, filtered, extracted with CH_2Cl_2 (10.0 mL x 2), washed with saturated brine (20.0 mL x 2), dried (Na_2SO_4), filtered, purified by chromatography on SiO_2 . Specific gradient used for each sample is included in the characterization data. All the yields reported are corrected based on residual solvent from ^1H NMR. Reaction 3A yielded tert-butyl (R)-(1-(1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate (424 mg, 76% yield) as a white solid. The C^3/N^1 ratio was 97:3. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: $152.1\text{--}153.2\text{ }^\circ\text{C}$; FTIR (neat, cm^{-1}) 3418, 3402, 3376, 2974, 2911, 1684, 1522; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) (85:15 mixture of rotamers) : δ 10.78 (br, 1H), 7.46 (d, $J = 8.0\text{ Hz}$, 1H), 7.32 (d, $J = 8.0\text{ Hz}$, 1H), 7.28–7.21 (m, 2H), 7.19–7.10 (m, 4H), 7.05 (ddd, $J = 8.4, 7.2, 1.2\text{ Hz}$, 1H), 6.95 (ddd, $J = 8.0, 6.8, 1.2\text{ Hz}$, 1H), 6.76 (d, $J = 8.4\text{ Hz}$, 0.85H), 6.34 (d, $J = 9.2\text{ Hz}$, 0.15H), 3.97–3.83 (m, 1H), 2.90–2.65 (m, 4H), 1.29 (s, 7.65H), 1.12 (s, 1.35H); ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$) (rotamers): δ 155.1, 139.6, 136.1, 129.0, 128.0, 127.5, 125.8, 123.2, 120.8, 118.3, 118.1, 111.4, 111.3, 77.3, 52.6, 39.9, 30.4, 28.2(27.8).

Example 3B: Preparation of tert-butyl (S)-(2-(1H-indol-3-yl)-1-phenylethyl)carbamate:**[0413]**

[0414] The general reaction as per Example 3A was performed between indole (294 mg, 2.51 mmol, 150 mol%) and tert-butyl (S)-4-phenyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (500 mg, 1.67 mmol, 100 mol%) to yield tert-butyl (S)-(2-(1H-indol-3-yl)-1-phenylethyl)carbamate (398 mg, 71% yield) as a white solid. The C³/N¹ ratio was 97:3. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 134.6-135.2 °C; FTIR (neat, cm⁻¹) 3416, 3401, 3371, 2980, 2909, 1683, 1524; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.74 (br, 1H), 7.54 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 8.4 Hz, 1H), 7.38-7.25 (m, 5H), 7.20 (dd, *J* = 6.8, 7.2 Hz, 1H), 7.06 (dd, *J* = 7.6, 7.2 Hz, 1H), 7.02 (s, 1H), 6.98 (dd, *J* = 7.6, 7.2 Hz, 1H), 4.89-4.74 (m, 1H), 3.08 (dd, *J* = 14.8, 8.8 Hz, 1H), 2.99 (dd, *J* = 14.4, 6.0 Hz, 1H), 1.31 (s, 7.65H), 1.08 (s, 1.35H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 155.0, 144.5, 136.0, 128.0, 127.3, 126.5, 126.4, 123.2, 120.8, 118.3, 118.2, 111.3, 111.3, 77.6, 55.0, 32.8, 28.2.


Example 3C: Preparation of tert-butyl (S)-(2-(1H-indol-3-yl)-1-(4-methoxyphenyl)ethyl)carbamate:**[0415]**

[0416] The general reaction as per Example 3A was performed between indole (264 mg, 2.25 mmol, 150 mol%) and tert-butyl (S)-4-(4-methoxyphenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (495 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (S)-(2-(1H-indol-3-yl)-1-(4-methoxyphenyl)ethyl)carbamate (378 mg, 69% yield) as a white solid. The C³/N¹ ratio was 98:2. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 173.3-176.5 °C; FTIR (neat, cm⁻¹)

3402, 3326, 2979, 2925, 2904, 1690, 1611, 1506; ^1H NMR (400 MHz, $\text{DMSO-}d_6$) (85:15 mixture of rotamers): δ 10.72 (br, 1H), 7.54 (d, $J = 8.0$ Hz, 1H), 7.39-7.28 (m, 2H), 7.24 (d, $J = 8.8$ Hz, 2H), 7.05 (ddd, $J = 8.4, 7.2, 1.2$ Hz, 1H), 7.02-6.94 (m, 2H), 6.84 (d, $J = 8.4$ Hz, 2H), 4.88-4.57 (m, 1H), 3.72 (s, 3H), 3.06 (dd, $J = 14.8, 8.4$ Hz, 1H), 2.96 (dd, $J = 14.8, 6.4$ Hz, 1H), 1.31 (s, 7.65H), 1.11 (s, 1.35H); ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): δ 157.9, 154.9, 136.4, 136.0, 127.5, 127.3, 123.2, 120.7, 118.3, 118.1, 113.4, 111.4, 111.2, 77.5, 55.0, 54.4, 32.8, 28.2.

Example 3D: Preparation of tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate:


[0417]



[0418] The general reaction as per Example 3A was performed between indole (370 mg, 3.16 mmol, 150 mol%) and tert-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (500mg, 2.11 mmol, 100 mol%) to yield tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate (406 mg, 70% yield) as a white solid. The C^3/N^1 ratio was 99:1. Column Gradient: 0 to 50% iPrOAc in Heptane. mp: 82.2-84.3 °C; FTIR (neat, cm^{-1}) 3416, 3401, 3366, 2974, 2963, 1684, 1524; ^1H NMR (400 MHz, $\text{DMSO-}d_6$): δ 10.77 (br, 1H), 7.56 (d, $J = 7.6$ Hz, 1H), 7.33 (ddd, $J = 8.0, 1.2, 0.8$ Hz, 1H), 7.10 (d, $J = 2.4$ Hz, 1H), 7.05 (ddd, $J = 8.0, 6.8, 1.2$ Hz, 1H), 6.97 (ddd, $J = 8.0, 6.8, 1.2$ Hz, 1H), 6.71 (d, $J = 8.4$ Hz, 1H), 3.82-3.66 (m, 1H), 2.87 (dd, $J = 14.0, 6.0$ Hz, 1H), 2.65 (dd, $J = 14.0, 7.6$ Hz, 1H), 1.38 (s, 9H), 1.01 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$): δ 155.0, 136.1, 127.5, 123.1, 120.7, 118.4, 118.1, 111.6, 111.2, 77.3, 46.8, 32.2, 28.3, 20.1.

Example 3E: Preparation of tert-butyl (R)-(1-(1H-indol-3-yl)-3-methylbutan-2-yl)carbamate:

[0419]




[0420] The general reaction as per Example 3A was performed between indole (331 mg, 2.83

mmol, 150 mol%) and tert-butyl (R)-4-isopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (500 mg, 1.88 mmol, 100 mol%) to yield tert-butyl (R)-(1-(1H-indol-3-yl)-3-methylbutan-2-yl)carbamate (335 mg, 59% yield) as a white solid. The C³/N¹ ratio was 94:6. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 145.1-146.9 °C; FTIR (neat, cm⁻¹) 3417, 3402, 3362, 2978, 1686, 1526; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.72 (br, 1H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.31 (ddd, *J* = 8.0, 1.2, 0.8 Hz, 1H), 7.07 (d, *J* = 2.0 Hz, 1H), 7.04 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.96 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.59 (d, *J* = 9.2 Hz, 0.85H), 6.15 (d, *J* = 10.0 Hz, 0.15H), 3.64-3.53 (m, 1H), 2.80 (dd, *J* = 14.8, 5.2 Hz, 1H), 2.68 (dd, *J* = 14.8, 8.8 Hz, 1H), 1.77-1.65 (m, 1H), 1.32 (s, 7.65H), 1.12 (s, 1.35H), 0.95-0.82 (m, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 155.6, 136.1, 127.5, 122.7, 120.7, 118.3, 118.0, 112.0, 111.2, 77.1, 55.6, 31.4, 28.3, 27.1, 19.4, 17.7.

Example 3F: Preparation of tert-butyl (S)-(1-cyclopropyl-2-(1H-indol-3-yl)ethyl)carbamate:


[0421]



[0422] The general reaction as per Example 3A was performed between indole (264 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-cyclopropyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (395 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (S)-(1-cyclopropyl-2-(1H-indol-3-yl)ethyl)carbamate (292 mg, 65% yield) as a white solid. The C³/N¹ ratio was 99:1. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 128.8-130.5 °C; FTIR (neat, cm⁻¹) 3414, 3400, 3362, 2981, 2937, 1683, 1525; ¹H NMR (400 MHz, DMSO-*d*₆) (9:1 mixture of rotamers): δ 10.73 (br, 1H), 7.52 (d, *J* = 7.6 Hz, 1H), 7.31 (ddd, *J* = 8.0, 1.2, 0.8 Hz, 1H), 7.07 (d, *J* = 2.4 Hz, 1H), 7.04 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.95 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.66 (d, *J* = 8.4 Hz, 0.9H), 6.26 (s, 0.1H), 3.30-3.18 (m, 1H), 2.91 (dd, *J* = 14.4, 5.6 Hz, 1H), 2.85 (dd, *J* = 14.4, 8.0 Hz, 1H), 1.33 (s, 8.1H), 1.17(s, 0.9H), 0.96-0.83 (m, 1H), 0.42-0.32 (m, 1H), 0.32-0.24 (m, 2H), 0.15-0.01 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 155.3, 136.0, 127.7, 122.9, 120.6, 118.4, 118.0, 111.6, 111.2, 77.2, 54.2, 30.4, 28.2, 16.0, 3.0, 1.9.

Example 3G: Preparation of tert-butyl ((1S,2S)-2-(1H-indol-3-yl)-2,3-dihydro-1H-inden-1-yl)carbamate:


[0423]



[0424] The general reaction as per Example 3A was performed between indole (282 mg, 2.41 mmol, 150 mol%) and tert-butyl (3aR,8aS)-8,8a-dihydroindeno[1,2-d][1,2,3]oxathiazole-3(3aH)-carboxylate 2,2-dioxide (500 mg, 1.61 mmol, 100 mol%) to yield tert-butyl ((1S,2S)-2-(1H-indol-3-yl)-2,3-dihydro-1H-inden-1-yl)carbamate (304 mg, 54% yield) as a white solid. The C³/N¹ ratio was 95:5. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 155.7-157.1 °C; FTIR (neat, cm⁻¹) 3387, 3351, 2980, 2938, 1691, 1500; ¹H NMR (400 MHz, DMSO-*d*₆) (87:13 mixture of rotamers): 10.85 (br, 1H), 7.61 (d, *J* = 8.0 Hz, 1H), 7.42-7.32 (m, 2H), 7.30-7.20 (m, 4H), 7.17 (dd, *J* = 8.0, 4.0 Hz, 1H), 7.07 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.97 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 5.18 (dd, *J* = 9.2, 9.2 Hz, 1H), 3.69 (dd, *J* = 18.4, 9.6 Hz, 0.87H), 3.64-3.53 (m, 0.13H), 3.37 (dd, *J* = 15.2, 8.0 Hz, 0.87H), 3.30-3.23 (m, 0.13H), 3.13-3.04 (m, 0.13H), 2.99 (dd, *J* = 15.2, 10.4 Hz, 0.87H), 1.38 (s, 7.83H), 1.12 (s, 1.17H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.9, 144.5, 141.3, 136.6, 127.2, 127.1, 126.4, 124.4, 123.3, 121.7, 120.9, 119.1, 118.2, 115.5, 111.4, 77.7, 60.8, 44.0, 37.2, 28.2 (27.7).

Example 3H: Preparation of tert-butyl (S)-(2-(1H-indol-3-yl)propyl)carbamate:

[0425]




[0426] The general reaction as per Example 3A was performed between indole (264 mg, 2.25 mmol, 150 mol%) and tert-butyl (S)-5-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (356 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (S)-(2-(1H-indol-3-yl)propyl)carbamate (319 mg, 78% yield) as a colorless liquid. The C³/N¹ ratio was 93:7. Column Gradient: 0 to 50% *i*PrOAc in Heptane. FTIR (neat, cm⁻¹) 3412, 3327, 2971, 2930, 1685, 1508, 1456; ¹H NMR (400 MHz, DMSO-*d*₆) (9:1 mixture of rotamers): δ 10.78 (br, 1H), 7.58 (d, *J* = 7.6 Hz, 1H), 7.33 (ddd, *J* = 8.0, 1.2, 0.8 Hz, 1H), 7.10 (d, *J* = 2.0 Hz, 1H), 7.05 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.96 (ddd, *J* = 8.0, 6.8, 1.2 Hz, 1H), 6.86 (dd, *J* = 6.0, 6.0 Hz, 0.9H), 6.51 (s, 0.1H), 3.29 (ddd, *J* = 13.2, 5.6, 5.6 Hz, 1H), 3.18-3.05 (m, 1H), 2.97 (ddd, *J* = 13.2, 8.8, 6.0 Hz, 1H), 1.41 (s, 0.9H), 1.38 (s, 8.1H), 1.25 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ

155.7, 136.4, 126.6, 121.0, 120.8, 118.6, 118.1, 117.9, 111.4, 77.4, 46.8, 30.9, 28.3(28.2), 18.4(18.2).

Example 31: Preparation of tert-butyl (R)-(1-(3-fluorophenyl)-2-(1H-indol-3-yl)ethyl)carbamate:


[0427]



[0428] The general reaction as per Example 3A was performed between indole (263 mg, 2.25 mmol, 150 mol%) and tert-butyl (S)-4-(3-fluorophenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (475 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-(3-fluorophenyl)-2-(1H-indol-3-yl)ethyl)carbamate (382 mg, 72% yield) as a white solid. The C³/N¹ ratio was 99:1. Column Gradient: 0 to 50% iPrOAc in Heptane. mp: 148.6-151.2 °C; FTIR (neat, cm⁻¹) 3414, 3398, 3363, 3055, 2981, 1682, 1591, 1527; ¹H NMR (400 MHz, DMSO-*d*₆) (9:1 mixture of rotamers): δ 10.75 (br, 1H), 7.53 (d, *J* = 7.6 Hz, 1H), 7.44 (d, *J* = 8.4 Hz, 1H), 7.37-7.26 (m, 2H), 7.20-7.11 (m, 2H), 7.08-6.93 (m, 4H), 4.85-4.65 (m, 1H), 3.06 (dd, *J* = 14.4, 8.4 Hz, 1H), 2.98 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.30 (s, 8.1H), 1.08 (s, 0.9H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 162.2 (d, ¹*J*_{CF} = 243 Hz), 155.0, 147.5 (d, ³*J*_{CF} = 7 Hz), 136.0, 129.9 (d, ³*J*_{CF} = 8 Hz), 127.2, 123.3, 122.6, 120.8, 118.3, 118.2, 113.3 (d, ²*J*_{CF} = 21 Hz), 113.0 (d, ²*J*_{CF} = 21 Hz), 111.3, 111.0, 77.8, 54.7, 32.5, 28.2; ¹⁹F NMR (DMSO-*d*₆, 376 MHz): δ -113.6.

Example 3J: Preparation of tert-butyl (R)-(2-(1H-indol-3-yl)-1-(3-(trifluoromethyl)phenyl)ethyl) carbamate:


[0429]



[0430] The general reaction as per Example 3A was performed between indole (176 mg, 1.50 mmol, 150 mol%) and tert-butyl (S)-4-(3-(trifluoromethyl)phenyl)-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (368 mg, 1.00 mmol, 100 mol%) to yield tert-butyl (R)-(2-(1H-indol-3-yl)-1-(3-(trifluoromethyl)phenyl)ethyl) carbamate (320 mg, 79% yield) as a white solid. The C³/N¹ ratio was 98:2. Column Gradient: 0 to 50% iPrOAc in Heptane. mp: 99.3-101.4 °C; FTIR (neat, cm⁻¹) 3414, 3399, 3361, 2982, 2936, 1683, 1523; ¹H NMR (400 MHz, DMSO-*d*₆) (88:12 mixture of rotamers): δ 10.76 (br, 1H), 7.80-7.45 (m, 6H), 7.31 (d, *J* = 8.0 Hz, 1H), 7.08-7.00 (m, 2H), 6.96 (ddd, *J* = 8.0, 7.2, 1.2 Hz, 1H), 5.11-4.69 (m, 1H), 3.09 (dd, *J* = 14.4, 8.4 Hz, 1H), 3.00 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.30 (s, 7.9H), 1.07 (s, 1.1H); ¹³C NMR (100 MHz, DMSO-*d*₆): δ 155.1, 145.8, 136.1, 130.8, 129.0, 128.9 (q, ²*J*_{CF} = 32 Hz), 127.3, 124.4 (q, ¹*J*_{CF} = 270 Hz), 123.3 (q, ³*J*_{CF} = 4 Hz), 123.0, 122.8 (q, ³*J*_{CF} = 4 Hz), 120.8, 118.3, 118.2, 111.3, 110.8, 77.9, 55.0, 32.5, 28.2; ¹⁹F NMR (DMSO-*d*₆, 376 MHz): δ -61.0.

Example 3K: Preparation of tert-butyl (S)-(3-(1H-indol-3-yl)-1-phenylpropyl) carbamate:


[0431]



[0432] The general reaction as per Example 3A was performed between indole (264 mg, 2.25 mmol, 150 mol%) and tert-butyl (S)-4-phenyl-1,2,3-oxathiazinane-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (S)-(3-(1H-indol-3-yl)-1-phenylpropyl)carbamate (401 mg, 73% yield) as a white solid. The C³/N¹ ratio was 98:2. Column Gradient: 0 to 50% iPrOAc in Heptane. mp: 121.7-123.2 °C; FTIR (neat, cm⁻¹) 3390, 2979, 2929, 2859, 1681, 1507, 1457, 1364; ¹H NMR (400 MHz, CDCl₃) (80:20 mixture of rotamers): δ 8.04 (br, 0.2H), 7.96 (br, 0.8H), 7.66 (d, *J* = 8.0 Hz, 0.2H), 7.52 (d, *J* = 8.0 Hz, 0.8H), 7.40-7.22 (m, 6H), 7.22-7.15 (m, 1H), 7.15-7.06 (m, 1H), 7.00 (br, 1H), 4.88 (br, 0.8H), 4.75 (br, 1H), 4.53 (br, 0.2H), 3.52-3.35 (m, 0.4H), 3.32-3.20 (m, 0.4H), 2.87-2.67 (m, 1.6H), 2.16 (d, *J* = 8.8 Hz, 1.6 H), 1.42 (s, 9H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.4(156.2), 143.0, 136.4(136.6), 128.6, 127.3, 127.2, 126.4, 121.8(122.0), 121.5(120.8), 119.0(119.2), 118.7, 115.3, 111.2(111.3), 79.5, 54.7(46.7), 37.2(31.6), 28.4(29.7), 21.9(18.7).

Example 3L: Preparation of tert-butyl (R)-(1-(5-methyl-1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0433]



[0434] The general reaction as per Example 3A was performed between 5-methyl-1*H*-indole (295 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield of tert-butyl (R)-(1-(5-methyl-1*H*-indol-3-yl)-3-phenylpropan-2-yl)carbamate (401 mg, 73% yield) as a white solid. The C³/N¹ ratio was 92:8. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 123.0-123.8 °C; FTIR (neat, cm⁻¹) 3413, 3368, 2974, 2927, 1685, 1524; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.63 (br, 1H), 7.30-7.13 (m, 7H), 7.07 (d, *J* = 2.0 Hz, 1H), 6.87 (dd, *J* = 8.4, 1.6 Hz, 1H), 6.74 (d, *J* = 8.8 Hz, 0.85H), 6.34 (d, *J* = 7.6 Hz, 0.15H), 3.96-3.80 (m, 1H), 2.86-2.64 (m, 4H), 2.36 (s, 3H), 1.30 (s, 7.65H), 1.16(s, 1.35H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.1, 139.6, 134.5, 129.1, 128.0, 127.7, 126.4, 125.8, 123.2, 122.3, 117.9, 111.0, 110.9, 77.2, 52.7, 40.0, 30.2, 28.2(27.8), 21.3.

Example 3M: Preparation of tert-butyl (R)-(1-(7-methyl-1*H*-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0435]



[0436] The general reaction as per Example 3A was performed between 7-methyl-1*H*-indole (295 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-(7-methyl-1*H*-indol-3-yl)-3-phenylpropan-2-yl)carbamate (371 mg, 68% yield) as a white solid. The C³/N¹ ratio was 97:3. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 124.4-125.9 °C; FTIR (neat, cm⁻¹) 3417, 3407, 3370, 2974, 2926, 1683, 1524; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.75 (br d, *J* = 2.0 Hz, 1H), 7.34-7.21 (m, 3H), 7.21-7.09 (m, 4H), 6.90-6.83 (m, 2H), 6.74 (d, *J* = 8.8 Hz, 0.85H), 6.34 (d, *J* = 7.6 Hz, 0.15H), 3.98-3.84 (m, 1H), 2.91-2.65 (m, 4H), 2.44 (s, 3H), 1.31 (s, 7.65H), 1.14 (s, 1.35H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.2, 139.6, 135.7, 129.0, 128.0, 127.2, 125.8, 122.9, 121.3, 120.3, 118.4, 115.9, 111.9, 77.3, 52.6, 39.9, 30.5, 28.2(27.8), 16.7.

Example 3N: Preparation of tert-butyl (R)-(1-(5-chloro-1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0437]



[0438] The general reaction as per Example 3A was performed between 5-chloro-1H-indole (341 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-(5-chloro-1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate (345 mg, 60% yield) as a white solid. The C³/N¹ ratio was 96:4. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 70.5-73.9 °C; FTIR (neat, cm⁻¹) 3417, 3368, 2980, 2928, 1684, 1518; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.99 (br, 1H), 7.49 (d, *J* = 2.0 Hz, 1H), 7.34 (d, *J* = 8.8 Hz, 1H), 7.30-7.22 (m, 2H), 7.22-7.13 (m, 4H), 7.04 (dd, *J* = 8.4, 2.0 Hz, 1H), 6.77 (d, *J* = 8.8 Hz, 0.85H), 6.34 (d, *J* = 8.8 Hz, 0.15H), 3.90-3.76 (m, 1H), 2.83-2.62 (m, 4H), 1.27 (s, 7.65H), 1.10 (s, 1.35H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.1, 139.5, 134.6, 129.0, 128.8, 128.0, 125.8, 125.2, 122.9, 120.6, 117.7, 112.8, 111.5, 77.2, 52.9, 40.0, 30.1, 28.2(27.7).

Example 3O: Preparation of tert-butyl (R)-(1-phenyl-3-(5-(trifluoromethyl)-1H-indol-3-yl)propan-2-yl)carbamate:

[0439]




[0440] The general reaction as per Example 3A was performed between 5-(trifluoromethyl)-1H-indole (417 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-phenyl-3-(5-(trifluoromethyl)-1H-indol-3-yl)propan-2-yl)carbamate (297 mg, 47% yield) as a white solid. The C³/N¹ ratio was 96:4. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 130.9-131.8 °C; FTIR (neat, cm⁻¹) 3417, 3368, 2980, 2929, 1684, 1519; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): 11.26 (br, 1H), 7.84 (s, 1H), 7.50 (d, *J* = 8.4 Hz, 1H),

7.37-7.22 (m, 4H), 7.22-7.10 (m, 3H), 6.80 (d, $J = 8.8$ Hz, 0.85H), 6.36 (d, $J = 9.2$ Hz, 0.15H), 3.99-3.77 (m, 1H), 2.86 (d, $J = 6.8$ Hz, 2H), 2.76 (d, $J = 6.8$ Hz, 2H), 1.22 (s, 7.65H), 1.04 (s, 1.35H); ^{13}C NMR (100 MHz, DMSO- d_6) (rotamers): δ 155.1(154.6), 139.5, 137.6(137.7), 129.1, 128.0, 126.9, 125.8, 125.7 (q, $^1J_{\text{CF}} = 269$ Hz), 125.6, 119.1 (q, $^2J_{\text{CF}} = 32$ Hz), 117.1 (q, $^3J_{\text{CF}} = 4$ Hz), 116.1 (q, $^3J_{\text{CF}} = 4$ Hz), 112.9, 111.9, 77.2, 53.0(53.7), 40.3(41.0), 29.9(30.9), 28.1(27.6); ^{19}F NMR (DMSO- d_6 , 376 MHz): δ -58.1.

Example 3P: Preparation of tert-butyl (R)-(1-(5-methoxy-1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0441]



[0442] The general reaction as per Example 3A was performed between 5-methoxy-1H-indole (331 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-(5-methoxy-1H-indol-3-yl)-3-phenylpropan-2-yl)carbamate (523 mg, 92% yield) as a white solid. The C³/N¹ ratio was 97:3. Column Gradient: 0 to 50% iPrOAc in Heptane. mp: 123.2-123.9 °C; FTIR (neat, cm⁻¹) 3368, 2975, 2933, 1692, 1680, 1516; ^1H NMR (400 MHz, DMSO- d_6) (85:15 mixture of rotamers): δ 10.60 (s, 1H), 7.32-7.24 (m, 2H), 7.24-7.13 (m, 4H), 7.08 (d, $J = 2.4$ Hz, 1H), 6.91 (d, $J = 2.4$ Hz, 0.85H), 6.83 (br, 0.15H), 6.77 (d, $J = 8.8$ Hz, 0.85H), 6.70 (dd, $J = 8.8, 2.4$ Hz, 1H), 6.34 (d, $J = 9.2$ Hz, 0.15H), 3.94-3.80 (m, 1H), 3.72 (s, 3H), 2.85-2.65 (m, 4H), 1.29 (s, 7.65H), 1.13 (s, 1.35H); ^{13}C NMR (100 MHz, DMSO- d_6) (rotamers): δ 155.2, 152.9, 139.6, 131.3, 129.1, 128.0, 127.8, 125.8, 123.8, 111.9, 111.3, 110.8, 100.3, 77.3, 55.3, 52.8, 40.2, 30.2, 28.2(27.8).

Example 3Q: Preparation of tert-butyl (R)-(1-phenyl-3-(5-phenyl-1H-indol-3-yl)propan-2-yl)carbamate:


[0443]



[0444] The general reaction as per Example 3A was performed between 5-phenyl-1*H*-indole (435 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-phenyl-3-(5-phenyl-1*H*-indol-3-yl)propan-2-yl)carbamate (523 mg, 82% yield) as a white solid. The C³/N¹ ratio was 97:3. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 68.9-70.8 °C; FTIR (neat, cm⁻¹) 3427, 3411, 3392, 3310, 2977, 2929, 1715, 1696, 1506; ¹H NMR (400 MHz, DMSO-*d*₆) (85:15 mixture of rotamers): δ 10.85 (br, 1H), 7.70 (s, 1H), 7.67-7.60 (m, 2H), 7.48-7.33 (m, 4H), 7.32-7.24 (m, 3H), 7.23-7.15 (m, 4H), 6.80 (d, *J* = 8.8 Hz, 0.85H), 6.39 (d, *J* = 9.2 Hz, 0.15H), 4.03-3.84 (m, 1H), 2.87 (d, *J* = 6.8 Hz, 2H), 2.77 (d, *J* = 6.8 Hz, 2H), 1.24 (s, 7.65H), 1.09 (s, 1.35H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.1, 142.0, 139.6, 135.8, 130.7, 129.1, 128.6, 128.1, 128.0, 126.6, 126.0, 125.8, 124.0, 120.1, 116.6, 112.2, 111.6, 77.2, 53.1, 40.4, 30.1, 28.2(27.8).

Example 3R: Preparation of tert-butyl (R)-(1-(2-methyl-1*H*-indol-3-yl)-3-phenylpropan-2-yl)carbamate:


[0445]



[0446] The general reaction as per Example 3A was performed between 2-methyl-1*H*-indole (295 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-(2-methyl-1*H*-indol-3-yl)-3-phenylpropan-2-yl)carbamate (386 mg, 71% yield) as a white solid. The C³/N¹ ratio was 99:1. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 117.6-119.0 °C; FTIR (neat, cm⁻¹) 3440, 3385, 2984, 2930, 1684, 1524; ¹H NMR (400 MHz, DMSO-*d*₆) (83:17 mixture of rotamers): δ 10.67 (br, 1H), 7.39 (d, *J* = 7.6 Hz, 0.83H), 7.33 (d, *J* = 8.0 Hz, 0.17H), 7.28-7.18 (m, 3H), 7.18-7.07 (m, 3H), 6.96 (ddd, *J* = 8.0, 7.2, 1.2 Hz, 1H), 6.90 (ddd, *J* = 8.0, 8.0, 1.2 Hz, 1H), 6.73 (d, *J* = 8.8 Hz, 0.83H), 6.30 (d, *J* = 8.8 Hz, 0.17H), 3.94-3.74 (m, 1H), 2.85 (dd, *J* = 14.0, 6.4 Hz, 1H), 2.75-2.62 (m, 3H), 2.32 (s, 3H), 1.27 (s, 7.47H), 1.04 (s, 1.53H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): 155.1, 139.8, 135.2, 132.4, 128.9, 128.7, 128.0, 125.7, 119.8, 118.0, 117.5, 110.2, 107.4, 77.2, 53.4, 39.4, 29.9, 28.2(27.7), 11.4.

Example 3S: Preparation of tert-butyl (R)-(1-phenyl-3-(2-phenyl-1*H*-indol-3-yl)propan-2-yl)carbamate:


[0447]



[0448] The general reaction as per Example 3A was performed between 2-phenyl-1*H*-indole (435 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-phenyl-3-(2-phenyl-1*H*-indol-3-yl)propan-2-yl)carbamate (324 mg, 51% yield) as a white solid. The C³/N¹ ratio was 98:2. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 178.5-178.8 °C; FTIR (neat, cm⁻¹) 3380, 3354, 3376, 2981, 2932, 1683, 1508; ¹H NMR (400 MHz, DMSO-*d*₆) (82:18 mixture of rotamers): δ 11.14 (s, 1H), 7.69 (d, *J* = 7.6 Hz, 1H), 7.60 (d, *J* = 7.2 Hz, 2H), 7.44 (dd, *J* = 7.2, 7.2 Hz, 2H), 7.35 (dd, *J* = 7.2, 7.2 Hz, 2H), 7.21 (dd, *J* = 7.2, 7.2 Hz, 2H), 7.17-6.97 (m, 5H), 6.79 (d, *J* = 9.2 Hz, 0.82H), 6.38 (d, *J* = 9.6 Hz, 0.18H), 4.15-3.90 (m, 1H), 3.06 (dd, *J* = 14.0, 6.8 Hz, 1H), 2.96 (dd, *J* = 14.4, 7.2 Hz, 1H), 2.64 (d, *J* = 6.8 Hz, 2H), 1.21 (s, 7.40H), 0.97 (s, 1.60H); ¹³C NMR (100 MHz, DMSO-*d*₆) (rotamers): δ 155.0, 139.5, 136.0, 134.7, 133.0, 129.2, 128.9, 128.6, 128.0, 127.8, 127.1, 125.8, 121.3, 119.2, 118.5, 111.0, 109.2, 77.2, 53.4, 40.2, 28.1(27.6).

Example 3T: Preparation of tert-butyl (R)-(1-phenyl-3-(1*H*-pyrrolo[2,3-*c*]pyridin-3-yl)propan-2-yl)carbamate:

[0449]




[0450] The general reaction as per Example 3A was performed between 1*H*-pyrrolo[2,3-*c*]pyridine (266 mg, 2.25 mmol, 150 mol%) and tert-butyl (R)-4-benzyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (470 mg, 1.50 mmol, 100 mol%) to yield tert-butyl (R)-(1-phenyl-3-(1*H*-pyrrolo[2,3-*c*]pyridin-3-yl)propan-2-yl)carbamate (40 mg, 8% yield) as a white solid. The C³/N¹ ratio was 98:2. Column Gradient: 0 to 50% *i*PrOAc in Heptane. mp: 196.2-197.0 °C; FTIR (neat, cm⁻¹) 3360, 2978, 2931, 1685, 1625, 1524; ¹H NMR (400 MHz, CDCl₃) (90:10 mixture of rotamers): δ 8.56 (br, 1H), 8.28 (s, 1H), 7.57 (d, *J* = 6.4 Hz, 1H), 7.38-7.31 (m, 2H), 7.30-7.25 (m, 2H), 7.25-7.20 (m, 2H), 6.70 (br, 1H), 5.44 (br, 0.10H), 5.24 (d, *J* = 5.6 Hz, 0.90H), 4.52 (dd, *J* = 13.2, 8.4 Hz, 1H), 4.36 (dd, *J* = 13.6, 4.8 Hz, 1H), 4.26-4.14 (m, 1H), 3.00 (dd, *J* =

14.0, 7.6 Hz, 1H), 2.91 (dd, $J = 14.0, 6.8$ Hz, 1H), 1.27 (s, 8.10H), 1.11 (s, 0.90H).

Example 4

Preparation of azetidinyl-amines

[0451] Preparation of compounds of formula (VII) were performed according to the general scheme below:




[0452] R^5 , v , R^6 , and R^{10} are as described herein.

Example 4A:

Preparation of 1-(3-fluoropropyl)azetid-3-amine ethane-1,2-disulfonate

[0453] 1-(3-fluoropropyl)azetid-3-amine ethane-1,2-disulfonate was prepared according to the following reaction scheme:




[0454] Tert-butyl azetid-3-ylcarbamate hydrochloride (109.7kg, 1.0 eq.) was dissolved in MTBE (793.4kg), and 1-bromo-3-fluoropropane (82.3kg) was added. MTBE (14kg) and water (530kg) were added followed by $LiOH \cdot H_2O$ (66.0kg) at $15-25^\circ C$ followed by stirring at $50-60^\circ C$. After reaction completion, an organic phase was separated and then combined 1,2-ethanedithiolic acid dihydrate aqueous solution (247.8kg) at $0-5^\circ C$. The resulting mixture was stirred for 30 min at $15-20^\circ C$. An aqueous phase was separated, and 1,2-ethanedithiolic acid dihydrate (56.85kg) was added to the organic phase. The resulting mixture was stirred for 5h at $35-40^\circ C$ until deprotection was complete. MeOH (884.1kg) was then added at $35-40^\circ C$, and

the mixture was stirred for 2h at 35-40°C. After cooling to room temperature, the reaction mixture was stirred for 4h. Solid was collected by filtration and rinsed with aq. MeOH. The washed solid was dried at 35-42°C under reduced pressure to provide 106.8 kg of 1-(3-fluoropropyl)azetid-3-amine ethane-1,2-disulfonate (63% yield). ¹H NMR (400 MHz, DMSO-d₆) δ 9.92 (s, 0.7H), 9.63 (s, 0.3H), 8.35 (s, 3H), 4.52 (dt, J = 47.1, 5.6 Hz, 2H), 4.43-4.08 (m, 5H), 3.36 (s, 3H), 2.74 (s, 4H), 2.06 - 1.77 (m, 2H).

Example 4B:


[0455] Preparation of 1-(3-fluoropropyl)azetid-3-amine ethane-1,2-disulfonate *via* strain release chemistry.



Preparation of 2,3-dibromopropan-1-amine hydrobromide (3)

[0456] To **1** (20.0 g, 1.0 equiv, 50 wt% in water) was added a solution of K₂CO₃ (29.55 g, 1.0 equiv) in water (100 mL) and a solution of Boc₂O (93.32 g, 2.0 equiv) in EtOAc (100 mL) at 0 °C. The reaction mixture was stirred at 20 °C for 15 h. The organic layer was separated and solvent swapped to EtOH (100 mL) to provide a solution of **2** in EtOH. It was then added into a cold (0 °C) solution of Br₂ (71.74 g, 2.1 equiv) in EtOH (60 mL) at 0 °C. The reaction mixture was stirred at 20 °C for 16 h, filtered, washed with MTBE (60 mL), dried under vacuum at 40 °C for 15 h to provide **3** as a colorless solid (43.01 g, 66%): ¹H NMR (400 MHz, CD₃OD) δ 4.55 - 4.47 (m, 1H), 4.01 (dd, J = 10.9, 4.6 Hz, 1H), 3.86 (dd, J = 10.9, 8.7 Hz, 1H), 3.71 (dd, J = 13.9, 3.2 Hz, 1H), 3.39 - 3.33 (m, 1H).


[0457] Preparation of N,N-dibenzyleazetid-3-amine ethane-1,1 disulfonate.



[0458] To a solution of Bn_2NH (33.12 g, 1.0 equiv) in THF (330 mL) was added $\text{PrMgCl}\cdot\text{LiCl}$ (1.3 M in THF, 130 mL, 1.0 equiv) at 20 °C and stirred at 20 °C for 5 h to provide a solution of $\text{Bn}_2\text{NMgCl}\cdot\text{LiCl}$ in THF. In a separate reactor was charged **3** (50.0 g, 1.0 equiv) in THF (500 mL), cooled to -60 °C. *n*-BuLi (201 mL, 2.5 M solution in *n*-hexane, 3.0 equiv) was added into the suspension at -60 °C and stirred at -60 °C for 2 h to provide a solution of **4** in THF. $\text{Bn}_2\text{NMgCl}\cdot\text{LiCl}$ solution was added into the **4** at -60 °C, warmed to 20 °C and stirred at 20 °C for 12 h to provide a solution of **5** in THF. The reaction mixture was cooled to 0 °C, a solution of Boc₂O (73.28 g, 2.0 equiv) in THF (200 mL) was added at 0 °C, stirred at 20 °C for 2 h, cooled to 0 °C, quenched with a solution of AcOH (20.16 g, 2.0 equiv) in H₂O (383 mL). The organic layer was washed 5% Na₂SO₄ (150 mL). The combined aqueous layer was back extracted with MTBE (100 mL x 2) to provide a solution of **6**. A solution of ethanedisulfonic acid (40.0 g, 1.5 equiv) in MeOH (225 mL) was added into the solution at 20 °C and stirred at 40 °C for 20 h. The slurry was filtered, washed with THF (100 mL), dried under the vacuum (40 °C) for 20 h to provide **7** (69.21 g, 75% yield) as a white solid: ¹H NMR (400 MHz, D₂O) δ 7.57 - 7.44 (m, 10H), 4.66 (t, *J* = 8.2 Hz, 1H), 4.36 (s, 4H), 4.10 - 3.95 (m, 4H), 3.22 (s, 4H). MS ([M+H]⁺) calculated for C₁₇H₂₁N₂ 253.17, found 253.00.

Preparation of *N,N*-dibenzyl-1-(3-fluoropropyl)azetidin-3-amine (**8**)


[0459]



[0460] To a solution of **7** (40.0 g, 1.0 equiv) in MTBE (200 mL) and H₂O (200 mL) was added 1-bromo-3-fluoropropane (17.20 g, 1.5 equiv) and LiOH·H₂O (15.18 g, 4.0 equiv). The reaction mixture was heated to 55 °C and stirred at 55 °C for 20 h, cooled to 20 °C. The organic layer was separated and washed with 5% Na₂SO₄ (120 mL x 2). **8** can be obtained by concentration of MTBE solution under reduced vacuum as a white solid (22.3 g, 83% yield): ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.35 - 7.22 (m, 10H), 4.45 (t, *J* = 6.0 Hz, 1H), 4.33 (t, *J* = 6.0 Hz, 1H), 3.45 - 3.39 (m, 4H), 3.28 - 3.17 (m, 3H), 2.67 - 2.56 (m, 2H), 2.35 (t, *J* = 7.0 Hz, 2H), 1.64 - 1.48 (m, 2H). MS ([M+H]⁺) calculated for C₂₀H₂₆FN₂ 313.21, found 313.10.

Preparation of 1-(3-fluoropropyl)azetidin-3-amine ethane-1,2-disulfonate

[0461]




[0462] To a solution of **8** (2.00 g, 1.0 equiv) in MeOH (20.0 mL) was added a solution of ethanedithionic acid dihydrate (1.23 g, 1.0 equiv) in H_2O (10.0 mL) at 0°C . The reaction mixture was heated to 30°C , passed through charcoal cartridge. Pd/C (0.40 g, 0.20 equiv) was then charged into the reaction mixture, stirred at 45°C with 40 psi H_2 , filtered, concentrated to 5 mL, charged with MeOH (20.0 mL), stirred at 20°C for 3 h, filtered, washed with MeOH (2 mL) to provide the title compound (1.36 g, 93% yield) as a white solid: $^1\text{H NMR}$ (400 MHz, D_2O) δ 4.70 - 4.57 (m, 3H), 4.57 - 4.31 (m, 4H), 3.51 (t, $J = 7.2$ Hz, 2H), 3.24 (s, 4H), 2.11 - 1.97 (m, 2H); $^{19}\text{F NMR}$ (376 MHz, D_2O) δ -219.59.

Example 5

Preparation of (R)-1-(1H-indol-3-yl)propan-2-amine

[0463]



[0464] To a solution of imidazole (102.8 g, 1.51 mol, 1.5 equiv.) in DCM (1.33 L) was added SOCl_2 (179.3 g, 1.51 mol, 1.5 equiv.) dropwise at -5 to 0°C under N_2 over 30 min. The reaction mixture was stirred for 0.5 h at -5 to 0°C . tert-butyl (R)-1-(hydroxypropan-2-yl)carbamate (Boc-alaninol) (177.7 g, 1.01 mol, 1.0 equiv.) in DCM (1.33 L, 7.5 vol.) was added dropwise at -5 to 0°C over 1 h. The reaction mixture was stirred for 0.5 h at -2 to 0°C followed by addition of triethylamine (204 g, 2.02 mol, 2 equiv.) dropwise at -5 to 0°C . The resulting mixture was stirred for 0.5 h or until N-Boc-alaninol was completely consumed as determined by GC analysis. Water was added to the reaction mixture (1.3 L) at 0 – 20°C . The phases were

separated, and the aqueous phase was extracted with DCM (1.3 L). The organic phases were combined and washed with 10 w% citric acid (1.3 L), aq. NaHCO₃ (1.3 L), and brine (1 L) successively. The organic phase was cooled to 0~10°C, followed by addition of water (3.1 L) and RuCl₃·xH₂O (2.66 g), and then followed by addition of Oxone (927.1 g, 1.51 mol, 1.5 equiv.). The reaction mixture was warmed to 22°C gradually, and held for 3.5 h or until the sulfamidite intermediate was completely consumed as determined by GC analysis. The phases were separated, and the aqueous phase was filtered through a pad of Celite (50 g) followed by rinsing with DCM (1.3 L). The filtrate was extracted with the DCM (1.3 L), and the organic phases were combined and then washed with sat. Na₂S₂O₃ (1.3 L) and brine (1 L x 2). The organic phase was dried with Na₂SO₄ (50 g) and then filtered rinsing with DCM (200 mL). The combined filtrate and washes were concentrated under vacuum at 30°C for 1 h, and then further dried under high vacuum to provide 220 g of tert-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide with >99 wt% in 91.8% isolated yield (corrected) over 2 steps. ¹H NMR (400 MHz, CDCl₃): δ 4.66 (dd, *J* = 9.2, 6.0 Hz, 1H), 4.41 (qdd, *J* = 6.4, 6.0, 2.8 Hz, 1H), 4.19 (dd, *J* = 9.2, 2.8 Hz, 1H), 1.54 (s, 9H), 1.50 (d, *J* = 6.4 Hz, 3 H).

[0465] Other oxidation reactions were run generally according to the procedure immediately above where the oxidant, catalyst, and solvent were varied. The results are reported in Table 1 below where "Exp." refers to experiment number, "Product" refers to tert-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide and "SM" refers to Boc-alaninol starting material, "LC A%" refers to area percent purity by liquid chromatography, and "ND" refers to not detected. The reaction temperature for experiments 1 and 3-9 was 0-25°C and the reaction temperature for experiment 2 was 0-40°C. The reaction time for experiments 1-4 and 7 was 4 hours, the reaction time for experiments 5 and 6 was 8 hours, and the reaction time for experiments 8 and 9 was 18 hours.

Table 1: Oxidant Catalyst Conditions and Yields

Exp.	Oxidant/Catalyst	Solvent	Product (LC A%)	SM (LC A%)
6B	mCPBA (1.3 eq)	DCM (10 vol)	3.2	94.4
6C	Oxone (1.5 eq)	MeCN/H ₂ O (1/1, 10 vol)	4.1	46.6
6D	NaClO (5.26 eq)/NiCl ₂ (2.6 mol%)	DCM/H ₂ O (1/1, 10 vol)	0.5	94.6
6E	H ₂ O ₂ (1.3 eq)/FeCl ₃ (5 mol%)	H ₂ O (10 vol)	ND	93.2
6F	NaClO (5.26 eq)/FeCl ₂ (2.6 mol%)	DCM/H ₂ O (1/1, 10 vol)	ND	>95
6G	NaClO (5.26 eq)/FeCl ₃ (2.6 mol%)	DCM/H ₂ O (1/1, 10 vol)	ND	>95
6H	NaClO (5.26 eq)/RuCl ₃ (1.3 wt%)	DCM/H ₂ O (1/1, 10 vol)	83.1	ND

Exp.	Oxidant/Catalyst	Solvent	Product (LC A%)	SM (LC A%)
6I	Oxone (2.0 eq)/RuCl ₃ (1.3 wt%)	DCM/H ₂ O (1/1, 10 vol)	83.8	ND
6J	Oxone (2.0 eq)/RuCl ₃ (1.3 wt%)	MeCN/H ₂ O (1/1, 10 vol)	70.7	ND

[0466] To a mixture of indole (7.4 g, 63.2 mmol, 1.5 equiv.) and CuCl (5.4 g, 54.7 mmol, 1.3 equiv.) in DCM (60 ml) was added MeMgCl (3.0 M in THF, 18.7 mL, 56 mmol, 1.33 equiv.) at -15°C under N₂ over 10 min. The resulting light yellow mixture was stirred at -20°C for 10 min, and then a solution of tert-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide (10.0 g, 42.1 mmol, 1.0 equiv.) in DCM (40 ml) was added dropwise at -10°C under N₂ over 30 min. The mixture was stirred at -10°C for 2 hours or until the reaction was complete as judged by TLC (PE/EA = 5/1, disappearance of tert-butyl (R)-4-methyl-1,2,3-oxathiazolidine-3-carboxylate 2,2-dioxide) or by GC. The reaction was quenched by adding 10% citric acid (100 mL) while maintaining the internal temperature < 5°C. The phases were separated, and the aqueous phase was extracted with DCM (100 mL x 2). The organic phases were combined and washed with brine (100 mL x 2), followed by addition to the organic phase of activated carbon (5 g) and Na₂SO₄ (10 g). The resulting mixture was stirred at room temperature for 30 min then filtered. The cake was washed with DCM (50 mL x 2). The filtrate and wash were combined and concentrated under vacuum at 30°C to provide crude tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate (16.9 g, 40.9 LCA%). Heptane (200 ml) was added to the crude tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate, and the mixture was stirred at room temperature for 1 hour, during which time, off-white solid precipitated gradually. The solid was collected by filtration, and the cake was washed with heptane (17 mL x 3). The solid was dried under high vacuum at 25 °C to provide 7.0 g of tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate (25.5 mmol, 97 A%, 99 wt%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.76 (s, 1H), 7.55 (d, *J* = 7.9 Hz, 1H), 7.32 (dt, *J* = 8.1, 1.0 Hz, 1H), 7.09 (d, *J* = 2.3 Hz, 1H), 7.05 (ddd, *J* = 8.2, 7.0, 1.2 Hz, 1H), 6.96 (ddd, *J* = 8.0, 6.9, 1.1 Hz, 1H), 6.71 (d, *J* = 8.0 Hz, 1H), 3.76-3.69 (m, 1H), 2.86 (dd, *J* = 13.9, 5.8 Hz, 1H), 2.64 (dd, *J* = 14.1, 7.7 Hz, 1H), 1.37 (s, 9H), 1.00 (d, *J* = 6.6 Hz, 3H).

[0467] Other indole alkylation reactions were run generally according to the procedure immediately above where the copper catalyst species and the stoichiometry of the Grignard reagent were varied. The results are reported in Table 2 below where "Exp." refers to experiment number, "CuX" refers to the copper catalyst species, "eq." refers to equivalents, "Prod" refers to tert-butyl (R)-(1-(1H-indol-3-yl)propan-2-yl)carbamate, "N1" refers to a by-product wherein the indole amine is alkylated, "BA" refers to a bisalkyl byproduct, and "AY" refers to LC Assay yield, "A%" refers to area percent purity by liquid chromatography, and "ND" refers to not detected.

Table 2: Indole Alkylation Reaction Conditions and Yields

Entry	CuX	MgMeCl (eq.)	indole (eq.)	Prod (A%)	N1 (A %)	BA (A %)	AY (%)
6K	CuCl	1.1	1.2	82.9	1.2	6.5	72.2
6L	CuCl	2.2	2.4	87.4	3.0	2.8	84.7
6M	CuI	1.1	1.2	66.4	2.3	1.4	27.7
6N	CuI	2.2	2.4	52.1	6.7	nil	12.8
6O	CuTC	1.1	1.2	68.0	0.8	0.9	34.8
6P	CuTC	2.2	2.4	59.3	16.2	0.4	28.1
6Q	CuCN	1.1	1.2	70.2	16.7	1.6	22.3
6R	CuCN	2.2	2.4	58.9	31.3	1.1	46.8
6S	CuSCN	1.1	1.2	57.4	30.1	1.8	40.9
6T	CuSCN	2.2	2.4	61.9	29.4	1.0	54.3

[0468] Other indole alkylation reactions were run generally according to the procedure immediately above where the reaction temperature was varied. The results are reported in Table 3 below where "Exp." refers to experiment number, "G temp" refers to the Grignard reagent addition temperature in °C, "S temp" refers to the sulfamidate addition temperature in °C, "Prod" refers to *tert*-butyl (*R*)-(1-(1H-indol-3-yl)propan-2-yl)carbamate, "N1" refers to a by-product wherein the indole amine is alkylated, "BA" refers to a bisalkyl byproduct, and "AY" refers to LC assay yield, and "A%" refers to area percent purity by liquid chromatography.

Table 3: Variable temperature indole alkylation experiments

Exp	G Temp (°C)	S Temp (°C)	Prod (A%)	N1 (A%)	BA (A%)	AY (%)
6U	15 ~ 20	-10 ~ -15	83.4	1.2	5.7	65.5
6V	-5 ~ 0		87.6	1.6	6.0	72.1
6W	-10 ~ -15		85.8	1.1	6.3	80.4
6X	-20 ~ -25		89.0	1.3	6.4	80.3
6Y	-30 ~ -35		79.9	1.8	7.0	80.2
6Z	-40 ~ -45		82.8	3.2	6.6	78.9
6AA	-10 ~ -15	15 ~ 20	85.4	2.5	3.9	75.7
6AB		5 ~ 10	89.4	2.0	5.7	81.7
6AC		~ 0	86.9	1.5	6.9	80.8
6AD		-10 ~ -15	85.8	1.1	6.3	80.4
6AE		-20 ~ -25	85.8	0.9	5.5	73.7
6AF		-30 ~ -35	88.4	0.8	5.1	70.6


[0469] To a solution of *tert*-butyl (*R*)-(1-(1H-indol-3-yl)propan-2-yl)carbamate (0.5 g, 1.8 mmol, 92 A%) in MeOH (5 mL, 10 vol.) was added HCl in MeOH (10 M, 1.8 mL, 18 mmol) dropwise at 0°C under N₂ atmosphere over a period of 10 min. The resulting solution was stirred at 0°C for 2 hrs, then at 25-30°C for 1 h. The solution was concentrated under vacuum at 30°C, diluted with water (10 mL), and extracted with DCM (10 mL x 2). The pH of the aqueous phase was adjusted to ~13 with 1 M aq. NaOH at 0 -10°C and then extracted with DCM (20 mL x 4). The

organic phase was dried over Na_2SO_4 (10 g). The desiccant was filtered, and the cake was washed with DCM (10 mL). The combined filtrate and wash was concentrated under vacuum at 30°C to afford 0.31 g of crude (R)-1-(1H-indol-3-yl)propan-2-amine with 95.8 LCA% and 99.2% ee purity in 97% isolated yield. ^1H NMR: 400 MHz (CDCl_3): δ 1.20 (d, $J = 6$, 3H), 1.56 (s, br, 2H), 2.69 (dd, $J = 14$, 8, 1H), 2.91 (dd, $J = 14$, 4, 1H), 3.30-3.33 (m, 1H), 7.05 (s, 1H), 7.14 (t, $J = 7$, 1H), 7.22 (t, $J = 7$, 1H), 7.38 (d, $J = 8$, 1H), 7.64 (d, $J = 8$, 1H), 8.27 (s, 1H).

Example 6

Preparation of (R)-3-((1-(1H-indol-3-yl)propan-2-yl)amino)-2,2-difluoropropan-1-ol

[0470]



[0471] To a solution of 2,2-difluoropropane-1,3-diol (51 kg, 90 wt%, 409.5 mol, 1 eq.) in DCM (332 kg, 5 vol.) was added thionyl chloride (58.4 kg, 491.4 mol, 1.2 eq.) at 20 - 25°C over 2 h. The reaction mixture was heated to 30 - 35°C , stirred for 2 h, and ice water (255 kg) was added to quench the reaction. The phases were separated, and the aqueous phase was extracted with DCM. The organic phases were combined and washed with water. To the crude organic phase was added water (255 kg) and FeCl_3 (1.6 kg), and the biphasic mixture was cooled to -3°C . Bleach [NaClO (8.1 wt%), 680 kg, 1556 mol, 3.8 equiv.] was then added dropwise during a period of 5 h at -5 to 1°C , and the reaction mixture was stirred for 1 h at 0°C . The reaction mixture was filtered through Celite. The phases were separated, and the aqueous phase was extracted with DCM. The organic phases were combined and washed with aq. solution of Na_3SO_3 and brine, then dried over Na_2SO_4 . The desiccant was filtered off, and the filtrate was concentrated to ca. 100 L. Heptane (80 kg) was added to the resulting suspension, and the mixture was further concentrated to ca. 100 L. Solids were collected by filtration and dried under vacuum to afford 52.01 kg of 5,5-difluoro-1,3,2-dioxathiane 2,2-dioxide as an off-white crystal with (72% yield, two steps from 2,2-difluoropropane-1,3-diol). ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 5.14 (t, $J = 10.8$ Hz, 1H).

[0472] Various other catalysts were evaluated for the preparation of 5,5-difluoro-1,3,2-

dioxathiane 2,2-dioxide generally according to the procedure immediately above. The results are reported in Table 4 below where "Exp." refers to experiment number, "diol" refers to 2,2-difluoropropane-1,3-diol, "Prod" refers to 90 wt.% 5,5-difluoro-1,3,2-dioxathiane 2,2-dioxide and "Yield" refers to assay yield. Each reaction was quenched with 5 volumes of water.

Table 4: Conditions for preparation of 5,5-difluoro-1,3,2-dioxathiane 2,2-dioxide

Exp	Diol	NaClO (eq.)	Catalyst (mol%)	Prod (Yield)
7B	50 g	5.26	NiCl ₂ (2.8)	78.6%
7C	50 g	5.26	RuCl ₃ (2.8)	67.1%
7D	10g	5.26	RuCl ₃ (2.0)	57.1%
7E	10g	5.26	RuCl ₃ (2.8)	64.3%
7F	10g	5.26	CoCl ₂ (2.8)	65.7%
7G	10 g	5.26	FeCl ₃ (2.8)	71.4%
7H	10 g	5.26	FeCl ₂ (2.8)	75.7%
7I	10 g	5.26	MnCl ₂ (2.8)	60.7%
7J	100 g	5.26	NiCl ₂ (2.8)	78.3%
7K	100 g	5.26	FeCl ₃ (2.8)	79.5%
7L	500 g	5.26	FeCl ₃ (2.8)	76%
7M	50 g	5.26	NiCl ₂ (2.8)	78%
7N	32 g	5.26	FeCl ₃ (2.8)	76%
7O	32 g	5.26	None	63%


[0473] In a 2 L flask were placed (R)-1-(1H-indol-3-yl)propan-2-amine (99 wt%, 100 g, 568.2 mmol, 1 equiv.), 5,5-difluoro-1,3,2-dioxathiane 2,2-dioxide (97.9 wt%, 108 g, 608 mmol, 1.07 equiv.), K₂CO₃ (55 g, 398 mmol, 0.7 equiv.), and acetonitrile (1 L, 10 vol.). The resulting mixture was heated to 80°C and stirred for 4 hrs. The reaction was cooled to 35°C and filtered. The cake was washed with acetonitrile (100 mL x 2), and p-TsOH monohydrate (119 g, 625.6 mmol, 1.1 equiv.) and water (100 mL, 1 vol) were added to the combined filtrate. The resulting biphasic mixture was heated to 80°C and stirred for 3 hrs. The mixture was then poured into 1 L of ice-water, and the pH of the mixture was adjusted to 9 with saturated Na₂CO₃ (350 mL) at < 5°C. The phases were separated, and the aqueous was extracted with i-PrOAc (500 mL x 2). The organic phases were combined and washed with water (500 mL) and brine (500 mL x 2), then dried over Na₂SO₄ (30 g). The desiccant was filtered off, and the cake was washed with i-PrOAc (100 mL x 2). The filtrate and wash were combined concentrated under vacuum at 40 °C. The residue was dissolved in i-PrOAc (200 mL) and heated to 60°C. Heptane (800 mL) was added dropwise at 60°C, and then the resulting mixture was stirred for 10 min. The mixture was cooled to 30-35°C slowly over 2 hrs, and then further cooled to 0°C. After stirring for 10 min, solids were collected by filtration, and the cake was washed with heptane (100 mL x

2). The solid thus obtained was dried under vacuum at 45°C to afford (R)-3-((1-(1H-indol-3-yl)propan-2-yl)amino)-2,2-difluoropropan-1-ol (155 g, 99.2 LCA%, 96 wt%, 97.6% isolated yield). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.79 (s, 1H), 7.51 (dd, *J* = 7.9, 1.2 Hz, 1H), 7.33 (dt, *J* = 8.2, 1.0 Hz, 1H), 7.13 (d, *J* = 2.3 Hz, 1H), 7.05 (ddd, *J* = 8.2, 6.9, 1.3 Hz, 1H), 6.96 (ddd, *J* = 8.0, 6.9, 1.2 Hz, 1H), 5.35 (t, *J* = 6.3 Hz, 1H), 3.61 (td, *J* = 13.8, 5.2 Hz, 2H), 3.03-2.91 (m, 3H), 2.83 (dd, *J* = 14.0, 5.7 Hz, 1H), 2.59 (dd, *J* = 14.0, 7.2 Hz, 1H), 1.69 (s, 1H), 0.96 (d, *J* = 6.2 Hz, 3H).

Example 7

Preparation of 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidin-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (2R,3R)-2,3-dihydroxysuccinate

[0474]



[0475] In step 1, to a flask with stirring apparatus was combined 4-bromo-2,6-difluorobenzaldehyde (131.79 g, 596.35 mmol), (R)-3-((1-(1H-indol-3-yl)propan-2-yl)amino)-2,2-difluoropropan-1-ol (160 g, 596.35 mmol), acetic acid (51.26 mL, 894.52 mmol), and toluene. The reaction was heated to 75°C and held overnight, cooled, and then diluted with toluene. The resulting solution was then quenched with an aqueous potassium carbonate solution, washed with brine and water, and treated with activated charcoal. Following filtration, the solution was concentrated and crystallized from toluene/heptane to afford 3-((1R,3R)-1-(4-bromo-2,6-difluorophenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-

difluoropropan-1-ol in 81% yield as a pale yellow solid. ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 10.59 (s, 1H), 7.45 - 7.36 (m, 3H), 7.20 (d, $J = 8.1$ Hz, 1H), 7.05 - 6.92 (m, 2H), 5.29 (t, $J = 6.1$ Hz, 1H), 5.21 (s, 1H), 3.71 - 3.55 (m, 1H), 3.53 - 3.38 (m, 2H), 3.17 (q, $J = 15.2$ Hz, 1H), 2.89 (ddd, $J = 15.3, 4.8, 1.5$ Hz, 1H), 2.71 - 2.55 (m, 2H), 1.08 (d, $J = 6.5$ Hz, 3H). MS ($[\text{M}+\text{H}]^+$) calculated for $\text{C}_{21}\text{H}_{19}\text{BrF}_4\text{N}_2\text{O}$ 470.06, found 470.80.


[0476] Alternatively, 4-bromo-2,6-difluorobenzaldehyde (67.2 g, 304 mmol), (R)-3-((1-(1H-indol-3-yl)propan-2-yl)amino)-2,2-difluoropropan-1-ol (80.0 g, 298 mmol), acetic acid (25.6 mL, 447 mmol), and methanol were combined in a stirred flask. The mixture was refluxed for 24 hours then cooled. Solids were precipitated from the resulting solution by addition of an aqueous solution of potassium carbonate. The resulting slurry was filtered and washed to afford 3-((1R,3R)-1-(4-bromo-2,6-difluorophenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol in 97% yield as a pale yellow solid.


[0477] In steps 2 and 3, 3-((1R,3R)-1-(4-bromo-2,6-difluorophenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (100 g, 212.2 mmol), 1-(3-fluoropropyl)azetidino-3-amine ethane-1,2-disulfonate (82.09 g, 254.6 mmol), acetonitrile, DBU (1061 mmol 161.5 g 159.4 mL) were combined in a stirred flask followed by addition of Pd-175 (5.304 mmol 4.144 g). The reaction was heated to 75°C for 2 hours, cooled, concentrated, and then diluted with methyl tert-butyl ether. This resulting solution was worked up with an aqueous solution of ammonium chloride, brine, and water, then scavenged with SiliaMetS Thiol. Following filtration, the solution was concentrated, diluted with ethanol, and crystallized with tartaric acid to afford 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidino-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (2R,3R)-2,3-dihydroxysuccinate in a 90% yield as a yellow solid following filtration and washing. ^1H NMR (400 MHz, $\text{DMSO-}d_6$) δ 10.52 (s, 1H), 7.38 (dd, $J = 7.5, 1.3$ Hz, 1H), 7.21 - 7.16 (m, 1H), 7.02 - 6.90 (m, 2H), 6.82 (d, $J = 6.9$ Hz, 1H), 6.18 - 6.08 (m, 2H), 5.07 (s, 1H), 4.53 (t, $J = 5.8$ Hz, 1H), 4.42 (t, $J = 5.9$ Hz, 1H), 4.18 (s, 2H), 4.14 - 4.05 (m, 1H), 3.93 (ddt, $J = 9.1, 7.0, 3.5$ Hz, 2H), 3.74 - 3.60 (m, 1H), 3.51 - 3.32 (m, 2H), 3.21 - 3.02 (m, 3H), 2.88 - 2.73 (m, 3H), 2.70 - 2.51 (m, 2H), 1.83 - 1.66 (m, 2H), 1.09 - 1.05 (m, 3H). MS ($[\text{M}+\text{H}]^+$) calculated for $\text{C}_{27}\text{H}_{31}\text{F}_5\text{N}_4\text{O}$ 522.24, found 523.00.

Example 8

Preparation of 3-((1R,3R)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidino-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2H-pyrido[3,4-b]indol-2-yl)-2,2-difluoropropan-1-ol (2R,3R)-2,3-dihydroxysuccinate

[0478]





[0479] In steps 1-4, a reactor was charged with 4-bromo-2,6-difluorobenzaldehyde (75.0 g, 0.339 mol, 100 mol%), *p*-toluenesulfonic acid monohydrate (162 mg, 0.000849 mol, 0.250 mol%), and toluene (225 mL). Triethyl orthoformate (55.4 g, 62.1 mL, 0.373 mol, 110 mol%) was charged over 15 min at rt, the mixture was stirred at rt for 1 h, and the mixture was distilled to provide a toluene solution of 5-bromo-2-(diethoxymethyl)-1,3-difluorobenzene. Another reactor was charged with 1-(3-fluoropropyl)azetid-3-amine 2-(trioxidanylthio)ethane-1-sulfonate (131.3 g, 0.407 mol, 120 mol%), and acetonitrile (656 mL). DBU (124.0 g, 122.8 mL, 0.814 mol, 240 mol%) was charged over 15 min at rt, and the mixture was stirred at rt for 2 h, distilled with toluene, and filtered to provide a toluene solution of 1-(3-fluoropropyl)azetid-3-amine which was charged, along with NaOtBu (39.14 g, 0.407 mol, 120 mol%), to the toluene solution of 5-bromo-2-(diethoxymethyl)-1,3-difluorobenzene. The mixture was sparged, charged with BrettPhos Pd G3 (3.076 g, 0.00309 mol, 1.00 mol%), sparged, heated at 60°C for 18 h, cooled, quenched with water, and washed with water twice. Siliamets Thiol (20.0 g) was charged and the mixture was heated at 50 C for 2 h, cooled, and filtered to provide a toluene solution of N-(4-(diethoxymethyl)-3,5-difluorophenyl)-1-(3-fluoropropyl)azetid-3-amine. Acetic acid (21.4 mL, 0.373 mol, 110 mol%) and water (300 mL) were added and the mixture was held at rt for 2 h. An aqueous phase was separated, treated with aqueous NaOH (50 wt%, 21.5 mL, 0.407 mol, 120 mol%), and seeded with 2,6-difluoro-4-((1-(3-fluoropropyl)azetid-3-yl)amino) benzaldehyde (923 mg, 0.00339 mol, 1.00 mol%). The resulting solids were filtered, washed with water, and dried to provide 2,6-difluoro-4-((1-(3-fluoropropyl)azetid-3-yl)amino)benzaldehyde (77.9 g, 84.3 % yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 10.05 (dd, *J* = 1.2 Hz, 1H), 6.05-5.97 (m, 2H), 5.03 (d, *J* = 6.4 Hz, 1H), 4.54 (t, *J* = 6.0 Hz, 1H), 4.42 (t, *J* = 6.0 Hz, 1H), 4.14-4.03 (m, 1H), 3.70 (td, *J* = 6.8, 1.6 Hz, 2H), 2.99-2.93 (m, 2H), 2.59 (t, *J* = 7.2 Hz, 2H), 1.83-1.66 (m, 2H). MS: calcd for C₁₃H₁₅F₃N₂O [M+H]⁺ = 273.1, observed = 273.0.

[0480] In step 5, a reactor was charged with (R)-3-((1-(1H-indol-3-yl)propan-2-yl)amino)-2,2-

difluoropropan-1-ol (6.60 kg, 24.60 mol, 100 mol%), 2,6-difluoro-4-((1-(3-fluoropropyl) azetidino-3-yl)amino)benzaldehyde (6.70 kg, 24.60 mol, 100 mol%), *L*-tartaric acid (5.54 kg, 36.90 mol, 150 mol%), and ethanol (39.6 L). The reaction mixture was heated at 70°C for 2 h, seeded with 3-((1*R*,3*R*)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidino-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2*H*-pyrido[3,4-*b*]indol-2-yl)-2,2-difluoropropan-1-ol (2*R*,3*R*)-2,3-dihydroxysuccinate (0.0827 kg, 0.123 mol, 0.5 mol%), agitated at 70 °C for 2 days, quenched with ethanol (26.4 L), cooled, and filtered to collect solid. The solid was washed with ethanol and dried under vacuum to provide 3-((1*R*,3*R*)-1-(2,6-difluoro-4-((1-(3-fluoropropyl)azetidino-3-yl)amino)phenyl)-3-methyl-1,3,4,9-tetrahydro-2*H*-pyrido[3,4-*b*]indol-2-yl)-2,2-difluoropropan-1-ol (2*R*,3*R*)-2,3-dihydroxysuccinate (12.7 kg, 78% yield) as an off white solid.

[0481] Various acids other than tartaric acid, as well as a comparative example with tartaric acid, were evaluated for the step 5 ring closure reaction generally according to the procedure immediately above. The results are reported in Table 5 below where "Exp." refers to experiment number, and "Conv" refers to the conversion to the fused tricyclic ring structure. The tartaric acid salt resulted in substantial increase in the final yield. Further, the tartaric acid salt minimized presence of byproduct oligomers formed during synthesis, resulting in purer product and less epimerization of the final compound product.



Table 5: Acids tested for ring closure for Compound A


Exp.	Acid	Conv.	Comments
9B	Acetic	95%	20% ring opening by-product
9C	Trifluoroacetic	20%	----
9D	Methanesulfonic	43%	----
9E	Benzoic	17%	----
9F	Pivalic	77%	12% ring opening by-product
9G	Diphenylphosphoric	97%	61% isolated yield and 6% oligomers
9H	Triflic	60%	----
9I	Formic	29%	----
9J	Tartaric	97%	89% isolated yield and 0.3% oligomers
9K	Fumaric	92%	70% isolated yield and 6% oligomers

Example 9:

Recrystallization of Compound B.

[0482]






[0483] To a 250 mL reactor was charged crude Compound B (4.00 g, 5.95 mmol) in MeOH (90.0 mL) and EtOH (10.0 mL) mixture. The slurry was heated to 60 °C and became homogeneous. Compound B seeds (200 mg, 0.297 mmol, 5 mol%) were added into the solution. The slurry was distilled with EtOH, cooled to 25 °C over 1 h, held at 25 °C for 1 h, filtered, dried to provide Compound B (3.83 g, 91% yield) as an off white solid.

Example 10:

Recrystallization of Compound B:

[0484]




[0485] Example 10A: To a 30 L reactor was charged crude Compound B (1.26 kg, 1.88 mol, 100 mol%) and MTBE (6.30 L, 5.00 mL/g). A solution of NaOH (154 g, 3.85 mol, 205 mol%) in water (6.30 L, 5.00 mL/g) was added into the slurry. The reaction mixture was stirred at 20 °C for 30 min. The top organic layer was washed with water (2.53 L, 2.00 mL/g) twice, passed through charcoal cartridge, distilled with EtOH to provide Compound B freebase solution in EtOH.

[0486] Example 10B: To a 20 L reactor was charged with *L*-tartaric acid (296 g, 1.97 mol, 105 mol%), EtOH (2.52 L, 2.00 mL/g) and heated to 70 °C. 20% of Compound B freebase solution was transferred into the 20 L reactor. Compound B seeds (25.2 g, 0.0375 mol, 2 mol%) were added into the solution. The rest of Compound B freebase solution was transferred into the 20 L reactor over 1 h, aged at 70 °C for 30 min, cooled to 20 °C over 5 h, aged at 20 °C for a minimum of 2 h, filtered, washed with EtOH (2.52 L, 2.00 mL/g) three times, dried to provide Compound B (1.12 kg, 89%) as an off white solid.

Example 11:


[0487] Compounds provided herein were characterized by mass spectrometry and/or NMR techniques as understood in the art. The compounds of this example, unless otherwise noted, have verified stereochemistry.

Compound (M1)




Determined m/z: 672.64.

Compound (M2)




Determined m/z: 672.64.

Compound (M3)



Determined m/z: 652.63

Compound (M4)



Determined m/z: 540.57 (stereomeric mixture).

Example 12:

Enzymatic transformations:

[0488] The screening of transaminases in the kinetic resolution mode of rac. α -methyltryptamine revealed enzymes possessing sufficient (R)-enantioselectivity for the asymmetric reductive amination approach and sufficient (S)-enantioselectivity for the kinetic resolution approach. Exemplary transaminases are listed in Table 6.

Table 6: Enantioselective transaminases (TA):

(R)-enantioselective TAs	(S)-enantioselectivity TAs
TA-P2-A01 (Codexis)	TA-P1-F03 [Codexis]
ESC-ATA01 (Libragen)	TA-P1-G05 [Codexis]
TA-P2-A07 (Codexis)	ATA47 [c-LEcta]
ATA117 (Codexis)	3FCR_59F_87F_231G*
ESC-ATA03 (Libragen)	3FCR_59W_87F_231A*
	3FCR_59F_86A_87F_152F_231A_234M_382M*
	3HMU_264V*
	ATA12

[0489] Asymmetric reduction applying transaminase TA-P2-A07 was performed in buffer; iso-propyl amine and organic co-solvents such as DMSO or Acetonitrile on 5 g scale. Compound (XXI) was converted to compound (3) as described herein at a degree above 95% with an enantiomeric excess (EE) above 99% after 1 day.

[0490] The reductive amination was also performed in organic solvents containing small amounts of buffer and iso-propyl amine where the transaminase TA-P2-A07 was immobilized on the solid support. Compound (XXI) was converted to compound (3) at a conversion degree above 95% and an EE above 99% after 4 weeks. The organic solvent enabled higher

substrates loadings up to 5% [w/w].

[0491] Kinetic resolution was performed in the presence of transaminase ATA12 in (1) whole cells; (2) cell free lysate or (3) crude lyophilisate in buffer containing pyruvate and organic cosolvent such as acetonitrile. Enantiomeric excess of >99% was achieved. Conversion to compound (3) was achieved at a degree equal to or above 50%. The undesired enantiomer was depleted by oxidative deamination towards the ketone.

[0492] Sequence Identification of select TAs:

3FCR Y59FIY87FIT231G: amino acid (SEQ ID NO:1)

MLKNDQLDQW	DRDNFFHPST	HLAQHARGES	ANRVIKTASG	VFIEDRDGTK	LLDAFAGL <u>FC</u>	60
VNVGYGRQEI	AEAIADQARE	LAYYHS <u>F</u> VGH	GTEASITLAK	MILDRAPKNM	SKVYFGLGGS	120
DANETNVKLI	WYNNILGRP	EKKKIISRWR	GYHGSGLVTG	SLTGLELFHK	KFDLPVEQVI	180
HTEAPYYFRR	EDLNQTEEQF	VAHCVAELEA	LIEREGADTI	AAFIGEPILG	<u>G</u> GGIVPPPAG	240
YWEAIQTVLN	KHDILLVADE	VVTGFGRGLT	MFGSDHYGLE	PDIITIAKGL	TSAYAPLSGS	300
IVSDKVWKVL	EQGTDENGPI	GHWYSAHP	IGAAAGVANL	KLLDELNLVS	NAGEVGAYLN	360
ATMAEALSQH	ANVG DVRGEG	LLCAVEFVKD	RDSRTFFDAA	DKIGPQISAK	LLEQDKIIAR	420
AMPQGDILGF	APPFCLTRAE	ADQVVEGTLR	AVKAVLG			457

3FCR Y59WIY87FIT231A: amino acid (SEQ ID NO:2)

MLKNDQLDQW	DRDNFFHPST	HLAQHARGES	ANRVIKTASG	VFIEDRDGTK	LLDAFAGL <u>WC</u>	60
VNVGYGRQEI	AEAIADQARE	LAYYHS <u>F</u> VGH	GTEASITLAK	MILDRAPKNM	SKVYFGLGGS	120
DANETNVKLI	WYNNILGRP	EKKKIISRWR	GYHGSGLVTG	SLTGLELFHK	KFDLPVEQVI	180
HTEAPYYFRR	EDLNQTEEQF	VAHCVAELEA	LIEREGADTI	AAFIGEPILG	<u>A</u> GGIVPPPAG	240
YWEAIQTVLN	KHDILLVADE	VVTGFGRGLT	MFGSDHYGLE	PDIITIAKGL	TSAYAPLSGS	300
IVSDKVWKVL	EQGTDENGPI	GHWYSAHP	IGAAAGVANL	KLLDELNLVS	NAGEVGAYLN	360
ATMAEALSQH	ANVG DVRGEG	LLCAVEFVKD	RDSRTFFDAA	DKIGPQISAK	LLEQDKIIAR	420
AMPQGDILGF	APPFCLTRAE	ADQVVEGTLR	AVKAVLG			457

3FCR Y59FIS86AIY87FIY152FIT231AII234MIL382M: amino acid (SEQ ID NO:3)

MLKNDQLDQW	DRDNFFHPST	HLAQHARGES	ANRVIKTASG	VFIEDRDGTK	LLDAFAGL <u>FC</u>	60
VNVGYGRQEI	AEAIADQARE	LAYYHS <u>A</u> FVGH	GTEASITLAK	MILDRAPKNM	SKVYFGLGGS	120
DANETNVKLI	WYNNILGRP	EKKKIISRWR	<u>G</u> FHGSGLVTG	SLTGLELFHK	KFDLPVEQVI	180
HTEAPYYFRR	EDLNQTEEQF	VAHCVAELEA	LIEREGADTI	AAFIGEPILG	<u>A</u> GGMVPPPAG	240
YWEAIQTVLN	KHDILLVADE	VVTGFGRGLT	MFGSDHYGLE	PDIITIAKGL	TSAYAPLSGS	300
IVSDKVWKVL	EQGTDENGPI	GHWYSAHP	IGAAAGVANL	KLLDELNLVS	NAGEVGAYLN	360
ATMAEALSQH	ANVG DVRGEG	<u>L</u> MCAVEFVKD	RDSRTFFDAA	DKIGPQISAK	LLEQDKIIAR	420
AMPQGDILGF	APPFCLTRAE	ADQVVEGTLR	AVKAVLG			457

3HMU I264V: amino acid (SEQ ID NO: 4)

MATITNHMPT	AELQALDAAH	HLHPFSANNA	LGEETRIVIT	RARGVWLND	ALAQKLAE <u>L</u> A	60
GLWCVNIGYG	RDELAEVAAR	QMRELPHYNT	FFKTTHVPAI	ALAQKLAE <u>L</u> A	PGDLNHVFFA	120
GGGSEANDTN	IRMVRTYWQN	KGQPEKTVII	SRKNAYHGST	VASSALGGMA	GMHAQSGLIP	180
DVHHINQPNW	WAEGGDMDPE	EFGLARAREL	EEAILELGEN	RVAAFIAEPV	QGAGGVIVAP	240
DSYWPEIQRI	CDKYDILLIA	DEV <u>V</u> CGFGRT	GNWFGTQTMG	IRPHIMPIAK	GLSSGYAPIG	300
GSIVCDEVVAH	VIGKDEFNHG	YTYSGHPVAA	AVALENLRIL	EEENILDHVR	NVAAPYLKEK	360
WEALTDHPLV	GEAKIVGMMMA	SIALTPNKAS	RAKFASEPGT	IGYICRERCF	ANNLIMRHVG	420
DRMIISPLPV	ITPAEIDEMF	VRIRKSLDEA	QAEIEKQGLM	KSA		464

Bolded and underline residues correspond to mutations to natural sequence.

Example 13:

[0493] Freebase, Salt, and Polymorph Characterization of Compound A**Abbreviations;****[0494]**

MIBK

Methyl Isobutyl Ketone

ACN

Acetonitrile

EtOAc

Ethyl Acetate

EtOH

Ethanol

MTBE

Methyl Tert-butyl Ether

IPAc

Isopropyl Acetate

MeTHF

Methyl Tetrahydrofuran

CPME

Cyclopenyl methyl ether

MeOH

Methanol

THF

Tetrahydrofuran

IPA

Isopropyl Alcohol

DMSO

Dimethyl Sulfoxide

DMC

Dimethyl Carbonate

MEK

Methyl Ethyl Ketone

DCM

Dichloromethane

[0495] Free-base Polymorph Screening. 96-well plate automated High-throughput Screening (HTS) using the Symyx CM2 system (Freeslate Inc., CA) was conducted to identify potential polymorphic forms for Compound A free base. Compound A was dispensed in each well (8 mg/well) using an automated powder dispensing accessory following which 800 μ l of solvent

(neat or mixture) was added and the slurry was stirred for 2 hours at 50°C. Compound A was initially dispensed in the slurry plate using either ethyl acetate (for tartrate) or MIBK (for fumarate) to maintain the solid form. From this master plate, the supernatant was filtered and distributed to three separate plates for evaporation, precipitation by anti-solvent addition and controlled cooling over 8-10 hrs from 50-20 °C. In all cases residual solvents were either evaporated or siphoned off, and the solid was examined using polarized light microscopy and XRPD.

Example 14:

[0496] Salt Screening. According to approximate solubility data of freebase and list of desired acids, five solvent systems were used in the screening. Approximately 20 mg of freebase amorphous compound was first dispersed in 0.5 mL of selected solvent in a glass vial and corresponding acid was then added with a 1:1 ratio of molar charge. An extra molar ratio of 2:1 (acid/freebase) was attempted for HCl, due to the two basic functional groups. The mixtures were stirred at RT overnight. Resulted solids were analyzed by XRPD. Clear solutions obtained after stirring were stirred at 5 °C for 2 days, then 0.5 mL of H₂O was added into each of clear solutions in ACN/H₂O (19:1, v/v) while 0.5 mL of n-heptane was added into each of the clear solutions in the other solvent systems followed by stirring at 5 °C for around 3 days, and the final clear solutions were transferred to slow evaporation at RT, in order to identify as many crystalline hits as possible.

[0497] The first round of screening identified the crystalline form hits as provided in Table 7. Tartrate and fumarate were further scaled up to 50 mg-1.5 g scale for further characterization. Testing in different solvent systems and conditions allowed for a) in-depth characterization of the different polymorphic forms obtained as hits in the screening and b) identifying conditions to inhibit and control cis-epimer formation. The epimer content varied from <1%-22% in different free base lots.

Table 7: Salt screening and crystalline form hits for Compound A.

Acid/Solvent	A	B	C	D	E
	Acetone	IPA	EtOAc	THF	ACN/H ₂ O (19:1 v/v)
Blank	Yellow gel	Yellow gel	Yellow gel	Light yellow gel	Yellow gel
Acetic Acid	Yellow gel	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Benzene Sulfonic Acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Citric Acid	Yellow oil	Yellow gel	Amorphous	Yellow gel	Yellow gel
Ethane sulfonic acid	Yellow oil	Yellow gel	Amorphous	Yellow gel	Yellow gel
1,2-	Amorphous	Amorphous	Amorphous	Amorphous	Dark blue

Acid/Solvent	A	B	C	D	E
	Acetone	IPA	EtOAc	THF	ACN/H ₂ O (19:1 v/v)
Ethanedisulfonic acid					gel
Fumaric acid	Yellow oil	Yellow gel	Fumarate Form A	Fumarate Form A	Yellow gel
Formic acid	Yellow gel	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Gentisic acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Glutamic acid	Glutamic acid	Glutamic acid	Glutamic acid	Glutamic acid	Yellow gel
Hydrochloric acid (1:1)	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Hydrochloric acid (2:1)	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Methanesulfonic acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Malonic acid	Yellow oil	Yellow gel	Yellow gel	Malonate Form A	Yellow gel
Maleic acid	Yellow gel	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Phosphoric acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
p-Toluene sulfonic acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Succinic acid	Yellow oil	Yellow gel	Yellow gel	Yellow gel	Yellow gel
Sulfuric acid	Yellow gel	Amorphous	Amorphous	Yellow gel	Yellow gel
L-Tartaric acid	Tartrate Form A	Yellow gel	Tartrate Form B	Tartrate Form C	Yellow gel

Methods:

[0498] Ambient X-ray Powder Diffractometry (XRPD). XRPD patterns were collected using the PANalytical Empyrean powder X-ray diffractometer (PANalytical Inc., Lelyweg, Netherlands). The powder sample was packed in a zero-background silicon holder and run in reflection mode (Bragg Brentano configuration). The instrument was equipped a Cu K α source with tube voltage and current of 45 kV and 40 mA respectively. Data was collected at ambient temperature from 3.0 to 40.0° 2 θ using a step size of 0.0263°, with a revolution speed of 8 sec. The incident beam path was equipped with a 0.02° soller slit, a fixed 1° anti scatter slit, a fixed incident beam mask of 10 mm, and a programmable divergence slit in automatic mode. A beam knife for linear detectors was used. The diffracted beam was equipped with a 0.02°

soller slit, a programmable anti scatter slit in automatic mode, and a nickel K- β filter. A PIXcel 1D detector was used in the scanning line detector (1D) mode. Data was analyzed using commercial software (JADE[®], version 9, Materials Data Inc., Livermore, CA).

[0499] Water sorption analysis. About 5-6 mg of powder sample was placed in the sample pan of an automated water sorption analyzer (Q5000SA, TA instruments, New Castle, DE) at 25°C and a nitrogen flow rate of 200 mL/min. The sample was initially "dried" at 0%RH for a total of 400 minutes (at 60 followed by 25°C), following which it was subjected to progressive increase in RH from 0-90%, in increments of 10% with a dwell time of 200 minutes at every RH. This was followed by a progressive decrease in RH in decrements of 10% back to 0% RH, using the same protocol.

[0500] Differential Scanning Calorimetry (DSC). Approximately 3-8 mg of powder sample was analyzed using a DSC Q2000[™] (TA instruments, New Castle, DE) equipped with a refrigerated cooling accessory. Samples were packed in non-hermetically pans (Tzero[™], aluminum pans) and typically heated from 0-200°C at 10°C/min under dry nitrogen purge. The instrument was calibrated using sapphire (baseline) and indium (temperature and cell constant). The data was analyzed using commercial software (Universal Analysis 2000, version 4.7A, TA Instruments).

[0501] Thermogravimetry (TGA). In a thermogravimetric analyzer (Discovery TGA, TA instruments), 3-5 mg of Compound A samples were heated in an open aluminum pan from RT to 350°C at a heating rate of 10°C/min under dry nitrogen purge. Temperature calibration was performed using Alumel[®] and Nickel. Standard weights of 100mg and 1gm were used for weight calibration.

[0502] Polarized Light Microscopy (PLM). Samples were dispersed in silicon oil and observed under cross polarizers of a video enhanced Leica DM 4000B microscope equipped with a high resolution CCD camera and motorized stage (Clemex Technologies Inc., Longueuil, Quebec, Canada) at 200X magnification. Photomicrographs were acquired using the Clemex Vision PE software (Clemex Technologies Inc., Longueuil, Quebec, Canada).

[0503] Scanning electron microscopy (SEM). Powder sample sputter coated on SEM stub and then examined using a benchtop Phenom SEM (Nanoscience Instruments, Inc., AZ). Micrographs were acquired at different magnifications.

[0504] Particle size distribution (PSD). Particle size analysis was performed using a Malvern Mastersizer 2000 instrument equipped with a Hydros 2000SM wet dispersion attachment (Malvern Instruments Ltd., Malvern, UK). ~40 mg of API was weighed into a vial and 1 mL of 0.1% Span 85 in heptane was added. The vial was sonicated for 10 seconds, about 0.5 mL was added to the sampler at a stir speed of 1500 rpm, and a PSD was performed at an obscuration of 10-20%. Owing to presence of large clusters in the sample that were settling rapidly out of the suspension, the dispersant was changed to 0.2% Span 85 in heptane to

stabilize the suspension. External sonication of 2 followed by two 5 minute duration was applied to break the clusters and PLM images were acquired before and after sonication. The aliquot was mixed in the sampler for 2 minutes prior to data collection to ensure homogeneity. The instrument was rinsed twice with isopropyl alcohol (IPA) and once with heptane before being filled with 0.1% Span 85 in heptane for each sample. After the last sample had run, the instrument was rinsed with IPA once. Data for replicates with 5 minute sonication has been reported.

[0505] BET Surface area analysis. Surface area measurement was conducted using a Micromeritics ASAP 2460 with a Micromeritics Smart VacPrep attachment (Micromeritics Instrument Corp., GA). A sample of 500 mg-1 g was weighed into an empty ASAP 2460 tube and placed on the Smart VacPrep, degassed for 24 hours under ambient conditions and then exposed to Krypton gas adsorption at 25°C and 100 mm Hg hold pressure. An 11 point measurement was made in the relative pressure range of 0.050-0.300 and the data was analyzed using MicroActive software provided by the vendor.

[0506] Solid-state Nuclear Magnetic Resonance Spectroscopy (SSNMR). All ^{13}C (@ 8 kHz spinning speed) SSNMR experiments were conducted using the 500 Mhz Bruker instrument (Bruker BioSpin GmbH, Karlsruhe, Germany) ^{13}C data were acquired using a CP/TOSS sequence. 1-2 K scans were collected for signal averaging. A contact time of 2-ms and a recycle delay of 5 seconds was used. Spinal 64 sequence was used for decoupling with a pulse length of 5.3 microseconds. ^1H 90 degree pulse length of 2.9 microseconds was employed. All ^{19}F (@ 14 kHz spinning speed) SSNMR experiments were conducted using the 500 Mhz Bruker instrument. ^{19}F data were acquired using a CP and direct polarization sequences. 64-256 K scans were collected for signal averaging. A contact time of 750 microseconds and a recycle delay of 7 seconds was used. ^1H 90 degree pulse length of 3.54 microseconds was employed.

Example 15:

[0507] Preliminary characterization of Compound A free base. Compound A free base was found to be amorphous. The starting material free base Compound A was characterized by XRPD, TGA and mDSC before screening.). As shown by the characterization results in FIG. 32 and FIG. 33, Compound A starting material was amorphous with a weight loss of 9.3% up to 220 °C in TGA and no substantial glass transition signal in mDSC. White solids of freebase Compound A were obtained by anti-solvent addition in MeOH/H₂O (3:20, v/v) with shaking and air dried for ~7 days.

[0508] Polymorph Screening. The solubility of freebase Compound A was estimated in 16 solvents at RT. (Table 8). Polymorph screening experiments were performed using different solution crystallization or solid phase transition methods. The methods uses and crystal forms

identified are summarized in Table 9.

Table 8: Approximate solubility of freebase Compound A

Solvent	Solubility (mg/mL)	Solvent	Solubility (mg/mL)
Acetone	S>60.0	MIBK	S>60.0
2-propanol	S>60.0	MTBE	S>60.0
EtOAc	S>60.0	IPAc	S>60.0
ACN	S>60.0	MeTHF	S>60.0
H ₂ O	S<1.9*	CPME	S>44.0*
1,4-Dioxane	S>60.0	n-Heptane	S<2.1*
EtOH	S>60.0	Cyclohexanes	S<1.9*
Toluene	S>60.0	Iso-butanol	S>40.0*

Table 9: Summary of polymorph screening of freebase Compound A

Method	No. of Experiments	Final Results
Solid Vapor Diffusion	13	Amorphous
Slurry at RT	25	Amorphous
Anti-solvent Addition	15	Amorphous
Liquid Vapor Diffusion	14	Amorphous
Slow Evaporation	12	Gel
Total	79	Amorphous

[0509] Solid Vapor Diffusion. Solid vapor diffusion experiments were conducted using 13 different kinds of solvent. For each experiment, about 15 mg of starting material was weighed into a 3-mL vial, which was placed into a 20-mL vial with 2 mL of corresponding volatile solvent. The 20-mL vial was sealed with a cap and kept at RT for 11 days to allow the solvent vapor to interact with the solid sample. The isolated solids were tested by XRPD. As summarized in Table 10, only oil or amorphous was obtained.

Table 10: Summary of solid vapor diffusion experiments

Solvent	Final Results
H ₂ O	Amorphous
DCM	N/A
EtOH	Oil
MeOH	Oil
ACN	Oil
THF	N/A
CHCl ₃	Oil
Acetone	N/A

Solvent	Final Results
DMF	Oil
EtOAc	N/A
1,4-Dioxane	N/A
IPA	Oil
DMSO	Oil

[0510] Slurry Conversion at RT. Slurry conversion experiments were conducted at RT in different solvent systems. About 40 mg of starting material was suspended in 0.3 mL of solvent in a 1.5-mL glass vial. After stirring, all the clear solutions were transferred to 5 °C, followed by slow evaporation at RT after 3 days. Results summarized in indicated that only amorphous, gel or oil was obtained.

Table 11: Summary of slurry conversion experiments for freebase Compound A

Solvent, v:v	Final Results
MeOH	Gel
EtOH	Gel
IPA	Gel
ACN	Gel
Acetone	Gel
THF	Gel
EtOAc	Gel
2-MeTHF	Gel
DCM	Gel
CPME	Gel
Acetic acid	Gel
DMC	Gel
Triethylamine	N/A
MIBK	Gel
MTBE	Gel
Iso-butanol	Gel
Acetone/H ₂ O (1:1)	Oil
ACN/H ₂ O (1:1)	Oil
EtOH/H ₂ O (1:1)	Gel
EtOH/n-heptane (1:1)	Gel
Xylene	N/A*
Cumene	N/A*

Solvent, v:v	Final Results
Cyclohexane	Amorphous**
N-heptane	Amorphous**
H ₂ O	Amorphous**
* no slow evaporation was applied	
** solids were centrifuged and analyzed by XRPD after stirring at RT for 4 days	
N/A: no solid obtained	

[0511] Anti-solvent Addition. A total of 15 anti-solvent addition experiments were carried out. About 30 mg of starting material was weighed into a 20-mL glass vial and dissolved in 0.15 mL of corresponding solvent at RT. Anti-solvent was added stepwise till the precipitation appeared or the total amount of anti-solvent reached 12 mL, with the sample being stirred magnetically. The precipitate was isolated for XRPD analysis. If no solid was observed, the clear solutions were stirred magnetically at 5 °C overnight and then evaporated at RT. Results in Table 12 showed that only amorphous or gel was generated.

Table 12: Summary of anti-solvent additions experiments for freebase Compound A

Solvent	Anti-solvent (v:v)	Final Results
MeOH	H ₂ O	Amorphous
EtOH		Amorphous
Acetone		Gel
THF		Gel
ACN		Gel
EtOH	Cyclohexane	Gel
Acetone		Gel
EtOAc		Gel
THF		Gel
EtOH	n-Heptane	Gel
Acetone		Gel
EtOAc		Gel
THF		Gel
MeOH	MeOH/H ₂ O (1:8)	Amorphous
EtOH	EtOH/H ₂ O (1:8)	Amorphous

[0512] Liquid Vapor Diffusion. Liquid vapor diffusion experiments were conducted under 14 conditions (Table 13). About 30 mg of starting material was weighed into each 3-mL glass vial. The corresponding solvent was added to get a solution. The vial was sealed into the 20-mL glass vial with 3-mL of corresponding anti-solvent and kept at RT, allowing about 11 days for

the vapor to interact with solution. The precipitates were isolated for XRPD analysis. Clear solutions were transferred to evaporation at RT.

Table 13: Summary of liquid vapor diffusion experiments for freebase Compound A

Solvent	Anti-solvent	Final Results
n-Butanol	Cyclohexane	Gel
	n-Heptane	Gel
Xylene	Cyclohexane	Amorphous
	n-Heptane	Amorphous
Cumene	Cyclohexane	Amorphous
	n-Heptane	Amorphous
MEK	Cyclohexane	Gel
	n-Heptane	Gel
IPAc	Cyclohexane	Gel
	n-Heptane	Gel
	H ₂ O	Amorphous
THF	Cyclohexane	Amorphous
	n-Heptane	Amorphous
ACN	H ₂ O	Gel

[0513] Slow Evaporation. Slow evaporation experiments were performed under 12 conditions. For each experiment, around 20 mg of starting material was weighed into a 3-mL glass vial, followed by the addition of corresponding solvent or solvent mixture to get a clear solution. Subsequently, the vial was covered with parafilm with 3-4 pinholes, and kept at 4 °C to allow the solution to evaporate slowly. Only gel was obtained, as summarized in Table 14.

Table 14: Summary of slow evaporation experiments for freebase Compound A

Solvent (v/v)	Final Results
Xylene	Gel
Cumene	Gel
IPAc	Gel
MIBK	Gel
MTBE	Gel
DCM/n-heptane (1:1)	Gel
THF/n-heptane (1:1)	Gel
Acetone/H ₂ O (1:1)	Gel
IPAc/cyclohexane (1:1)	Gel
MTBE/cyclohexane (1:1)	Gel

Solvent (v/v)	Final Results
EtOH/H ₂ O (1:1)	Gel
EtOH/n-heptane (1:1)	Gel

[0514] Table 15 summarizes the forms that were obtained via manual screening as described herein.

Table 15: Polymorphic forms of Compound A

Polymorph screening: Compound A free base	
<i>Form</i>	<i>Type</i>
N/A	No crystalline hits identified
Salt screening: Compound A free base	
<i>Form</i>	<i>Type</i>
A-tartrate	Acetone solvate
B-tartrate	Anhydrate
C-tartrate	THF solvate
A-malonate	THF solvate
A-fumarate	Anhydrate
Polymorph screening: Compound A tartrate	
<i>Form</i>	<i>Type</i>
D	Hydrate
E	DMSO solvate
F	Anhydrate
G	Methanol solvate

Example 16:

[0515] **Characterization of Compound B Form A.** Compound B Form A was prepared at 200-mg scale via solution reactive crystallization in acetone, as evidenced by XRPD (FIG. 1). As shown in FIG. 2a and FIG. 2b, weight loss of about 7.2% up to 125 °C was observed in TGA and DSC result showed one endotherm at 124.3 °C (onset temperature). ¹H NMR showed that its molar ratio was 0.98 (acid/freebase) and 5.6% acetone (a molar ratio of 0.69 to freebase) was detected. Heating experiments were conducted to further identify Form A. No form change was observed after Form A was heated to 90 °C, but the sample turned to amorphous after being heated to 140 °C. A considerable amount of acetone (4.5%) was still observed after heating Form A sample to 90 °C. The Form A sample was composed primarily of fine particles and some aggregates (FIG. 3). Based on the data collected, Compound B

Form A is an acetone solvate and that the loss of solvent is concomitant with melting.

Table 16: Representative XRPD Peaks for Compound B Form A:

2-Theta (°2θ)	d(Å)
4.643	19.0147
8.263	10.6916
9.286	9.5157
11.183	7.9057
11.495	7.6916
11.969	7.3884
12.543	7.0515
13.776	6.4228
14.226	6.2207
14.619	6.0545
15.092	5.8656
15.564	5.689
16.013	5.5302
17.354	5.1058
18.559	4.777
18.847	4.7046
19.321	4.5903
19.821	4.4757
20.265	4.3785
21.343	4.1597
21.631	4.105
21.92	4.0515
22.524	3.9443
22.97	3.8686
23.285	3.8171
23.549	3.7748
23.942	3.7138
24.81	3.5858
25.964	3.4289

[0516] Preparation of Compound B Form A. About 57.5 mg of tartaric acid was weighed into a 5-mL glass vial, and 2.0 mL acetone was added to get a clear solution. The clear acid solution was added to 2.0 mL freebase stock solution in acetone (-100 mg/mL) into the 5-mL vial with a

molar ratio of 1:1, and stirred at RT. About 1 mg of Compound B Form A seed was added, and the solution turned cloudy. The sample was stirred overnight before taking XRPD measurements as described herein. The the pattern conformed to Compound B Form A. The suspension was then stirred at RT for another 24 hours and the cake dried at 50 °C for 1.5 hrs. Yield: 170.4 mg, with a yield of -65.3%.

Example 17:

[0517] Characterization of Compound B Form B. Compound B Form B was prepared at 200-mg scale via solution reactive crystallization in EtOAc, as evidenced by XRPD (FIG. 4). As shown in FIG. 5a and FIG. 5b, a limited weight loss of about 1.3% up to 140 °C was observed in TGA and DSC result showed a melting endothermic peak at 156.7 °C (onset temperature). Stoichiometry was determined to be 1.08 (acid/freebase) and 1.4% EtOAc (a molar ratio of 0.11 to freebase) was detected by ¹H NMR. Based on the characterization data collected, Compound B Form B is an anhydrate.

[0518] The solid-state characterization data indicates that Form B is substantially crystalline (XRPD), and TGA indicates presence of some surface solvent of ~2% by an early onset of weight loss from RT-100°C) with a melting onset of ~163°C. The melting onset, following a shallow endotherm (vaporization of residual solvent) is at 163°C. Based on the derivative weight loss curve, a very early onset of weight loss (2.4% w/w) was detected up to 100°C, which may be attributed to surface solvent. The total weight loss, including that at the melt is 3.5% w/w. Form B was found to be slightly hygroscopic with a moisture pick up of 1.2% w/w at 90% RH, 25°C as shown by the water sorption-desorption profile in FIG. 8. The ¹³C (FIG. 6) and ¹⁹F SSNMR spectra (FIG. 7) further indicated formation of Form B. SEM (FIG. 9a, 500X magnification) and PLM (FIG. 9b, 200 X magnifications) images of Compound B show Form B comprises of dense spherical aggregates.

Table 17: Representative XRPD Peaks for Compound B Form B:

2-Theta (°2θ)	d(Å)
7.684	11.4966
11.491	7.6942
12.54	7.053
14.245	6.2123
15.303	5.7851
15.557	5.6912
16.014	5.5301
16.634	5.3252
17.371	5.1009
18.242	4.8593

2-Theta (°2θ)	d(Å)
19.163	4.6278
19.424	4.5662
19.892	4.4597
20.243	4.3833
21.817	4.0705
22.524	3.9442
22.996	3.8644
23.253	3.8223
23.573	3.7711
24.676	3.6049
25.073	3.5487
25.915	3.4353

[0519] Preliminary stress stability analysis of Compound B Form B. The XRPD patterns of Compound A salts (fumarate and tartrate) exposed to 40°C/ 75% RH for one month under open conditions. The solid form did not change under these conditions.

[0520] Although no correlation was observed between epimer content ranging from 0.56-0.72% and the solid-state properties of the different salt forms, the SSNMR, XRPD and DSC data indicate that the possibility of obtaining a mixture of forms existed from fumarate salts and that form control in either salt required a substantial amount of effort downstream, given the variability in structural data and melting points with change in crystallization conditions. The tartrate salt lacked the impurities found in the fumarate salt sample at 0.18% upon exposure to 40°C/75%RH for one month.

[0521] Compression analysis of Compound B Form B. FIG. 22, FIG. 23, and FIG. 24 demonstrate the effect of compression on Compound B tartrate Form B. 250 mg compacts of neat Compound B Form B were analyzed. Both SSNMR and XRPD (FIG. 23 and FIG. 22, respectively) show that the form remains unchanged upon compression. A lowering of the ¹⁹F T₁ relaxation time was observed upon compression which is attributable to generation of disorder, albeit very minor as evident from ¹⁹F SSNMR spectra of the compressed Compound B Form B in FIG. 23. ¹⁹F T₁ relaxation values are included in Table 18 for as is and compressed Compound B Form B. Compression did not affect the melting point of Compound B Form B, as seen in the DSC thermgram (FIG. 24). Comparative XRPD collected before and after exposure to accelerated stability conditions of 30°C/65% RH and 40°C/75%RH for up to 6 months (open) demonstrated that Compound A Form B remains unchanged upon exposure to these conditions.

Table 18: ¹⁹F T₁ relaxation times for Compound B Form B

Sample	T ₁ (s) for peaks at -96 to -125 ppm	T ₁ (s) for peaks at -214 to -230 ppm
Uncompressed (As-is)	12.5	11
Compressed	10.2	9

[0522] XRPD of one-month stability sample. The XRPD of Compound B Form B exposed to 40 °C/75% RH for one month shows the Compound B Form B remains unchanged under stress stability conditions including elevated temperature and moisture.

[0523] Preparation of Compound B Form B. About 58.0 mg of tartaric acid was weighed into a 5-mL glass vial, and 2.0 mL EtOAc was added. The acid remained undissolved. About 2.0 mL freebase stock solution in EtOAc (-100 mg/mL) was added into the 5-mL vial with a molar ratio of 1:1, and the solution was stirred at RT. About 1 mg of Form B seed was added and the solution remained clear. The solution was stirred overnight and sampled by XRPD. The pattern conformed to Compound B Form B. The suspension was stirred at 50 °C for 2 more days. The suspension was centrifuged and the cake dried at 50 °C for 2 hrs. Yield: 144.5 mg, with a yield of -56.1%.

Example 18:

[0524] Preparation of Compound B Form C. Form C was prepared at 200-mg scale via solution reactive crystallization in THF, as evidenced by XRPD (FIG. 10). As shown in FIG. 11a and FIG. 11b, a weight loss of 6.8% up to 130 °C was observed in TGA and DSC showed one endotherm at 118.1 °C (onset temperature). The stoichiometric ratio was determined to be 1.02 (acid/freebase) and 9.7%

[0525] Preparation of Compound B Form C. About 56.9 mg of tartaric acid was added into a 3-mL glass vial, and added 1.0 mL THF to get a clear solution. The clear acid solution was added to 2.0 mL freebase stock solution in THF (-100 mg/mL) with a molar ratio of 1:1, and stirred at RT. About 1 mg of Form C seed was added, and the solution turned a little turbid. The suspension was stirred overnight and sampled by XRPD. The pattern conformed to Form C. The suspension was stirred at RT for another 24 hours, the cake dried at 50 °C for 1.5 hours. The suspension was centrifuged to collect the solids. Yield: 161.4 mg, with a yield of -62.7%.

Example 19:

[0526] Preparation of Compound B Form D. Form D samples were obtained via fast evaporation in MeOH/DCM and slurry in H₂O at RT, respectively. The XRPD pattern for Form D

is shown in FIG. 12. TGA and DSC results of Form D sample are provided in FIG. 13a and FIG. 13b, respectively, and show a weight loss of 3.5% before 150 °C and an endotherm at 73.0 °C before melting/decomposition at 163.9 °C (onset temperature). The form change to Form F was observed after Form D was heated to 150 °C, cooled to 30 °C under protection of nitrogen, and then exposed to ambient condition. No significant amount of process solvent MeOH or DCM was detected by ¹H NMR and further determined the stoichiometric ratio of L-tartaric acid to freebase to be 1.0. Form D is a hydrate.

[0527] To evaluate the physical stability of Form D under different humidity, DVS data of Form D sample was collected at 25 °C after the sample was equilibrated at ambient humidity (80%RH). A platform from 20%RH (2.25% water uptake) to 80%RH (2.72% water uptake) was observed during desorption in DVS test of Form D, suggesting the dehydration of hydrous Form D occurred when relevant humidity value was less than 20%. Furthermore, as the theoretical water content of a mono-hydrate is 2.6%. Form D is therefore likely to be a mono-hydrate.

Table 19: Representative XRPD Peaks for Compound B Form D:

2-Theta (°2θ)	d(Å)
7.322	12.0634
10.992	8.0426
11.312	7.8161
12.182	7.2597
13.234	6.6847
13.487	6.5601
14.114	6.2697
14.668	6.0342
15.145	5.8453
15.702	5.6393
16.036	5.5225
16.217	5.4611
16.542	5.3546
17.249	5.1369
17.637	5.0246
18.113	4.8935
18.349	4.8311
19.108	4.641
20.209	4.3905
20.583	4.3115
21.163	4.1948

2-Theta (°2θ)	d(Å)
21.472	4.135
21.891	4.0569
22.762	3.9036
23.334	3.8091
23.569	3.7717

Example 20:

[0528] Preparation of Compound B Form E. Form E was obtained via DMSO-mediated crystallization by adding IPAc into DMSO solution, and its XRPD is shown in FIG. 14. TGA and DSC curves (FIG. 15a and FIG. 15b, respectively) showed a considerable weight loss of 8.3% before 140 °C and an endotherm at 126.3 °C before melting/decomposition at 142.6 °C (onset temperature). The ¹H NMR spectrum indicated a 0.7 equivalent of DMSO (~9.4 wt%) and no significant amount of IPAc were detected. The stoichiometric ratio of L-tartaric acid to freebase was determined to be 1.0. Form E is a DMSO solvate.

Example 21:

[0529] Preparation of Compound B Form F. Form F was obtained via heating a Form D sample to 150 °C, cooling to 30 °C under protection of nitrogen, and exposed to ambient conditions. The XRPD pattern of Form F is provided in FIG. 16. DSC analysis (FIG. 18) indicated that Form F was crystalline with an endothermic peak at 164.2 °C (onset temperature). ¹H NMR determined a stoichiometric ratio of L-tartaric acid to freebase to be 1.0 and no significant solvent signal was detected.

Table 20: Representative XRPD Peaks for Compound B Form F:

2-Theta (°2θ)	d(Å)
3.925	22.4935
10.54	8.3864
11.724	7.5419
12.52	7.0646
14.227	6.2205
15.407	5.7466
15.54	5.6976
15.902	5.5687
16.488	5.3721

2-Theta (°2θ)	d(Å)
16.844	5.2594
17.294	5.1235
18.267	4.8526
18.473	4.799
19.399	4.572
19.661	4.5117
20.005	4.4348
20.501	4.3286
20.655	4.2968
21.161	4.1952
21.287	4.1706
21.951	4.0458
22.972	3.8683
23.498	3.783
23.708	3.7499
23.943	3.7137
24.31	3.6584
24.679	3.6044
24.997	3.5594

[0530] To further characterize Form F, variable temperature XRPD (VT-XRPD) was conducted on Form D where the temperature increased to 150 °C and back to 30 °C under protection of nitrogen. Under such conditions, Form F was observed after dehydration of Form D sample at elevated temperatures, indicating Form F was an anhydrate.

[0531] To evaluate the physical stability of Form F under different humidity, dynamic vapor sorption (DVS) data of Form F was collected at 25 °C after the sample was equilibrated at ambient humidity (80%RH). A water uptake of -1.9% was observed up to 80%RH, suggesting Form F is slightly hygroscopic.

[0532] Preparation of Compound B Form G. Form G was obtained by slow evaporation in MeOH at RT. The Form G XRPD is provided in FIG. 19. TGA and DSC results are provided in FIG. 20a and FIG. 20b, respectively, and demonstrated a weight loss of 3.3% before 150 °C and a melting/decomposition peak at 170.4 °C (onset temperature). The ¹H NMR spectrum showed a 0.49 equivalent of MeOH (equal to ~3.0 wt%). The stoichiometric ratio of L-tartaric acid to freebase was determined to be 1.0. Form G is a MeOH solvate.

Table 21: Representative XRPD Peaks of Compound B Form G;

2-Theta (°2θ)	d(Å)
7.654	11.5416
11.462	7.7139
12.517	7.0663
15.275	5.796
15.514	5.7072
16.007	5.5323
17.349	5.1072
18.217	4.8658
19.114	4.6395
19.294	4.5966
19.424	4.5661
19.845	4.4702
20.236	4.3846
21.317	4.1647
21.577	4.1152
21.791	4.0752
22.497	3.9489
22.971	3.8685
23.228	3.8263
24.651	3.6085
25.046	3.5525
25.888	3.4388

Example 22:

[0533] Polymorph Screening of L-Tartrate. The solubility of Compound B Form B material was estimated in 20 solvents at RT. (Table 22). Approximately 2 mg solids were added into a 3-mL glass vial. Solvents in Table 6-7 were then added into the vials stepwise until the solids were dissolved or a total volume of 1 mL was reached. Results of Table 22 were used to guide the solvent selection in polymorph screening. Polymorph screening experiments were performed using different solution crystallization or solid phase transition methods. The methods utilized and crystal forms identified are summarized in Table 23.

Table 22: Compound B solubilities

Solvent	Solubility (mg/mL)	Solvent	Solubility (mg/mL)
DMSO	S>38.0	Anisole	S<1.9
MeOH	6.7<S<20.0	MTBE	S<2.1
THE	2.1<S<7.0	2-MeTHF	S<1.9
Acetone	2.1<S<7.0	1,4-Dioxane	S<2.2
EtOH	S<1.9	CPME	S<2.1
IPA	S<2.1	ACN	S<2.1
MEK	S<2.0	n Heptane	S<2.0
MIBK	S<1.9	Toluene	S<1.9
EtOAc	S<1.9	H ₂ O	S<1.9
IPAc	S<1.9	DCM	S<2.0

Table 23: Summary of polymorph screenings

Method	No. of Experiments	Isolated Solid Forms
Anti-solvent addition	14	Form B, D, E
Slurry conversion	34	Form B, D, G
Slow evaporation	9	Form G
Solid vapor diffusion	11	Form B
Liquid vapor diffusion	12	Form B, Form B+G, Form B+D
Slow cooling	10	N/A
Total	84	Form B, D, E, G, B+G, B+D

Example 23:

[0534] Anti-solvent Addition. A total of 14 anti-solvent addition experiments were carried out. For each experiment, about 15 mg of Compound B Form B was weighed into a 20-mL glass vial, followed by the addition of 0.3-1.0 mL corresponding solvent. The mixture was then magnetically stirred at the speed of 500 RPM to get a clear solution at RT. Subsequently, the corresponding anti-solvent was added to the solution to induce precipitation or until the total amount of anti-solvent reached 10.0 mL. The clear solutions were transferred to slurry at 5 °C. If no precipitation occurs, the solution was then transferred to fast evaporation at RT. The solids were isolated for XRPD analysis. Results summarized in Table 24 showed that Compound B Forms B, D, and E were obtained.

Table 24: Summary of anti-solvent addition experiments

Solvent	Anti-Solvent	Final Results
MIBK	MeOH	Form D*
EtOAc		Form D**

Solvent	Anti-Solvent	Final Results	
DCM		Form D**	
IPA		Form D**	
1,4-Dioxane		Amorphous	
ACN		Amorphous	
THF		Form B*	
Toluene		DMSO	Form E
MEK	N/A		
IPAc	Form E		
2-MeTHF	N/A		
CPME	Yellow oil		
H ₂ O	N/A		
EtOH	N/A		
* solid was obtained via stirring at 5 °C			
** solid was obtained via fast evaporation at RT			
N/A: no solid was obtained			

Example 24:

[0535] Slurry Conversion at RT. Slurry conversion experiments were conducted at RT in different solvent systems. For each experiment, about 20 mg of Compound B Form B was suspended in 0.3 mL corresponding solvent in a 1.5-mL glass vial. After the suspension was magnetically stirred for 17 days at RT, the remaining solids were isolated for XRPD analysis. Results summarized in Table 25 showed that Compound B Forms B, D, and G were obtained.

Table 25: Summary of slurry conversion experiments at RT.

Solvent (v:v)	Final Results
MeOH	Form G
EtOAc	Form B
Acetone	Form B
Anisole	Form B
MTBE	Form B
CPME	Form B
2-MeTHF	Form B
ACN	Form B
Toluene	Form B
DCM	Form B

Solvent (v:v)	Final Results
H ₂ O	Form D*
EtOH	Form B
IPAc/DMSO, 9:1	Form E
MIBK/MeOH, 9:1	Form B
EtOH/H ₂ O, 97:3, a _w =0.2	Form B
EtOH/H ₂ O, 92.7:7.3, a _w =0.4	Form B
EtOH/H ₂ O, 86:14, a _w =0.6	Form B
EtOH/H ₂ O, 71:29, a _w =0.8	Form D

* heating-cooling was performed on the sample after being stirred for 17 days.

Example 25:

[0536] Slurry Conversion at Elevated Temperatures. Slurry conversion experiments were conducted at 50 °C and 70 °C in different solvent systems. For each experiment, about 20 mg of Compound B Form B was suspended in 0.3 mL corresponding solvent in a 1.5-mL glass vial. After the suspension was magnetically stirred for 17 days at 50 °C and 70 °C, the remaining solids were isolated for XRPD analysis. Results summarized in Table 26 indicated that Compound B Form B was obtained.

Table 26: Summary of slurry conversion experiments at elevated temperatures

Solvent, v:v	Temperature, °C	Final Results
H ₂ O	50	Form B
MeOH		Form B
Acetone		Form B
THF		Form B
MEK		Form B
EtOAc		Form B
CHCl ₃		Form B
ACN		Form B
EtOH		Form B
MTBE		Form B
IPA		70
1-Butanol	Form B	
MIBK	Form B	
IPAc	Form B	

Solvent, v:v	Temperature, °C	Final Results
Anisole		Form B
Toluene		Form B

Example 26:

[0537] Slow Evaporation. Slow evaporation experiments were performed under 9 conditions. For each experiment, around 15 mg of Compound B Form B was weighed into a 3-mL glass vial, followed by the addition of corresponding solvent or solvent mixture to get a clear solution. Subsequently, the vial was covered with parafilm with 3-4 pinholes, and kept at RT to allow the solution to evaporate slowly. The isolated solids were tested by XRPD. As summarized in, Compound B Form G was generated.

Table 27: Summary of slow evaporation experiments

Solvent (v:v)	Final Results
MeOH	Form G
Acetone	Yellow gel
EtOH	Yellow gel
2-MeTHF	Yellow gel
THF	Yellow gel
ACN/MeOH, 1:3	Yellow gel
EtOAc/MeOH, 1:3	Form G
DCM/MeOH, 1:3	Form G
Anisole/MeOH, 1:3	Yellow gel

Example 27:

[0538] Solid Vapor Diffusion. Solid vapor diffusion experiments were conducted using 11 solvents. For each experiment, about 15 mg of Compound B Form B was weighed into a 3-mL vial, which was placed into a 20-mL vial with 4 mL of corresponding solvent. The 20-mL vial was sealed with a cap and kept at RT for 14 days to allow the solvent vapor to interact with the solid sample. The isolated solids were tested by XRPD. The results summarized in Table 28 indicated that Compound B Form B was obtained.

Table 28: Summary of solid vapor diffusion experiments.

Solvent	Final Results
H ₂ O	Form B

Solvent	Final Results
DCM	Form B
EtOH	Form B
MeOH	Form B
ACN	Form B
THF	Form B
CHCl ₃	Form B
Acetone	Form B
EtOAc	Form B
1,4-Dioxane	Form B
IPA	Form B

Example 28:

[0539] Liquid Vapor Diffusion. Twelve liquid vapor diffusion experiments were conducted. For each experiment, about 15 mg of Compound B Form B was dissolved in 0.5-1.0 mL of corresponding solvent to obtain a clear solution in a 3-mL vial. Subsequently, the solution was placed into a 20-mL vial with 4 mL of corresponding anti-solvent. The 20-mL vial was sealed with a cap and kept at RT, allowing sufficient time for solvent vapor to interact with the solution. Solids were isolated for XRPD analysis. Results summarized in Table 29 showed that Compound B Form B and a mixture of Form B+G/B+D were obtained.

Table 29: Summary of liquid vapor diffusion experiments.

Anti-solvent	Solvent	Final Results
1,4-Dioxane	MeOH	Form B
MeOH		Form B+G
Acetone		Form B+G
ACN		Form B+G
THF		Form B+D
EtOAc		Form B+G
CHCl ₃		DMSO
MTBE	N/A	
2-MeTHF	N/A	
DMC	N/A	
IPAc	N/A	
Anisole	N/A	

N/A: no solid obtained

Example 29:

[0540] Slow Cooling. Slow cooling experiments were conducted in 10 solvent systems. For each experiment, about 20 mg of Compound B Form B was suspended in 1.0 mL of corresponding solvent in a 3-mL glass vial at RT. The suspension was transferred to slurry at 50 °C with a magnetic stirrer at the speed of 500 RPM. The sample was equilibrated at 50 °C for 2 hrs and filtered using a 0.45 µm Nylon membrane. Subsequently, the filtrate was slowly cooled down from 50 °C to 5 °C at a rate of 0.1 °C/min. The results summarized in indicated that no solid was observed.

Table 30: Summary of slow cooling experiments.

Solvent, v:v	Final Results
MeOH	N/A
EtOH	N/A
Acetone	N/A
MEK	N/A
THE	N/A
2-MeTHF	N/A
H ₂ O	N/A
ACN/DMSO, 4:1	N/A
Toluene/MeOH, 3:1	N/A
EtOAc/MeOH, 3:1	N/A

Example 30:

[0541] Phase transformation. Crystallization of Compound B from solvents such as ethyl acetate, ethanol, acetone or and aqueous mixture of these may result in the formation of one more of the following forms: anhydrous Form B, anhydrous Form F, monohydrate Form D, and acetone solvate Form A. The XRPD patterns of these four solid forms are shown in FIG. 21. It was therefore important to know the physical stability of these forms to determine the complete phase transformation landscape and control crystallization to obtain the desired Form B. Owing to the close proximity of melting points of Forms F and B (160-164°C onset for Form F vs 166-175°C for Form B), form conversion was slow in competitive slurry bridging experiments at RT and a mixture of both forms was obtained. As a result, equilibrium solubility studies were conducted for both forms in ethanol over 17 hrs at 25, 35 and 50°C to determine their

thermodynamic stability relationship.

[0542] Form F was found to have higher solubility at these three temperatures, confirming Form B to be the thermodynamically stable form from 25-50 °C. The Van't Hoff plot of solubility of Forms B and F vs temperature showed that the two anhydrates were related enantiotropically with a transition temperature at ~19°C. In case of hydrate-anhydrate stability, it was found that Form B remained stable at water activity (a_w) $a_w < 0.2$ (RT), above which the hydrate Form D was stable. However, Form B was found to be kinetically stable up to a_w of 0.4 (RT) in slurry bridging experiments for up to 36 days. This suggests that although the critical a_w value for form conversion to hydrate is low, there is a large kinetic barrier for the conversion of Form B to Form D.

Slurry and crystallization samples. Form B + Form A

[0543] 1:1 mixtures of Compound B lots comprising (1) mixture of Form A/acetone solvate and Form B) and (2) Form B were made and added to 100% acetone and aqueous mixtures of acetone water (90% acetone, 95,96,97, 98 and 99% acetone v/v). Samples were slurried at RT for 120hrs, filtered and then analyzed by XRPD.

Form F

[0544] Form F was slurried in 100% ethanol at RT and 50°C overnight. One of the suspensions was seeded at RT with Form B while the other was left unseeded. The suspension stirred at 50°C was seeded with Form B. Samples were filtered and analysed by XRPD. Samples of Form F were slurried separately at RT in 100% DI water, 1:1 acetone/ water and 100% acetone. Slurries of Form F in neat solvents were maintained at RT whereas a solution of Form F was obtained in the 1:1 acetone: water mixture which was thus agitated at 5°C for cooling crystallization/precipitation. After 24 hrs, all samples were filtered and analyzed by XRPD. Form F was also slurried in 95:5 acetone: water and 97:3 acetone: water mixtures at 50°C for 2 hrs after which the slurries were cooled to RT, filtered and analyzed by XRPD. Finally, Form F was recrystallized from 95:5 acetone: water seeded with Form B at 50°C.

Form D

[0545] Form D was slurried at RT for 48hrs in 100% ethanol. The sample was subsequently filtered and analyzed by XRPD.

Results

[0546] Development of calorimetric method to estimate purity. Switching from ethanol to acetone/water as the crystallization solvent saw a marked improvement in purity, with oligomer content dropping by an average of 5% w/w, with acceptable yield. The change in oligomer content as well as yield was a function of slurry solvent composition. An increase in the water content improved purity but lowered the yield. For a given solvent composition, the melting point onset and % oligomer content consistently showed opposing results, indicating that purity is likely to affect the melting point, with greater % oligomer depressing the melting point and *vice versa*. Based on this observation, several slurry lots obtained from acetone/water were analyzed using DSC and tested for oligomer content by SEC and a correlation curve was developed. XRPD data was collected for all these samples to ensure that they conformed to Compound B Form B. Using the exponential fit to the data, purity estimates were made for several lots, including a 20 gm scale up and IPC sample. An R^2 value of 0.96 was obtained for the linear fit between experimental and predicted oligomer content. Given that these were solid /powder samples with inherent issues of sample homogeneity, this high R^2 value provided confidence in the robustness of the correlation determined between melting point onset and purity.

[0547] Compound B Form B was consistently obtained from 100% ethanol (slurry or crystallization). Further conditions were conducted including slurry or crystallizations from ethanol/water and slurry in acetone/water ($\geq 95\%$ acetone). Depending on the slurry and crystallization parameters, several other solid forms of Compound B were obtained, namely Form F (ethanol/water) Form A/acetone solvent ($\geq 95\%$ acetone) and mixtures of Form A and B or Form A and F. In conditions where Form F was obtained, the formation of an intermediate hydrate (Form D) was postulated since a_w values of ethanol/water mixtures employed for these samples were well above 0.2 (RT), where the hydrate is thermodynamically stable. Table 31 shows the crystallization conditions with the form obtained. Since mixtures of forms were obtained in several cases, a form control strategy was implemented to obtain Form B in the final solid phase.

Table 31: Crystallization conditions and corresponding solid forms

Crystallization condition	Form identified (XRPD)
Acetone/water(96:4), 5 vol, 50C for 6 h	Form A + Form B
Acetone/water(95:5)	Form A + Form B
Acetone/water(99:1), 5 vol, 50C for 2 h	Form A + Form B
crystallize GMP batch: 4 vol acetone/water 90:10, 8 vol acetone anti-solvent with 1% Form B seeds	Form A + Form F
Seeded crystallization in ethanol/water 65:35	Form F

[0548] Since acetone/water system was the most effective in purging oligomers from Compound B, a competitive slurry bridging experiment was conducted using 1:1 mixtures of two separate lots of Compound B: one that was predominantly Form A with some Form B and

a second that was slurried in acetone/water mixtures of various compositions at RT for 120 hrs. Acetone solvate was obtained as the stable form at $\geq 95\%$ acetone while a mixture of Form B and Form D hydrate was obtained at 90% acetone. As the desired acetone level was between 96 and 95% (to oligomer levels), there was a possibility of obtaining any of Forms A, B, or D from the final crystallization.

[0549] Form control strategy: conversion of Forms A and F to Form B. FIG. 2b shows the DSC thermograms of Form A acetone solvate. The first endotherm at 124°C indicates solvent loss and vaporization while the second endotherm at 164°C denotes the melting of the corresponding anhydrate. XRPD confirmed this anhydrate to be Form B when Form A was heated to 152°C . Form A was re-slurried in 100% ethanol overnight for conversion to Form B. Thus a two-step slurry process was proposed to aid form control for Compound B to obtain Form B in the event that the first slurry in acetone/water produces either Form A or a mixture for Form A and B, depending on the slurry conditions. This would ensure that irrespective of a change in form during the first slurry, Form B would be the final Compound B form that is isolated after the second slurry step.

[0550] Unlike the acetone solvate, Form F obtained from a 65:35 ethanol:water crystallization seeded with Form B showed only one melting endotherm at 162°C . Owing to near identical solubility values of Forms F and B in ethanol at 25°C (0.23 mg/mL for Form B vs 0.26 mg/mL for Form F), the thermodynamic driving force for conversion of metastable Form F to stable Form B is less and thus Form F \rightarrow Form B conversion rate would be slow, even in the presence of Form B as seeds. Irrespective of seeding and temperature, Form F \rightarrow Form B conversion is slow and a mixture of both forms is obtained after 12 hours.

[0551] To facilitate Form F \rightarrow Form B conversion in shorter timescales, a solvate route was adopted wherein Form F would be converted to Form B via acetone solvate or hydrate formation, which would subsequently be desolvated to Form B. Form F was slurried overnight in neat water, acetone and 1:1 acetone: water mixture, without any seeding. Form F remains unchanged in 100% acetone but converts to hydrate (Form D) in presence of water. When the hydrate is slurried in neat ethanol, a mixture of Form D and Form F results, thus indicating the propensity of the hydrate to desolvate to the metastable anhydrous form. Form F slurried in 95:5 acetone:water and 97:3 acetone:water at 50°C demonstrated that Form B appears but the conversion is incomplete. Recrystallization of Form F from 95:5 acetone:water using Form B seeds at 50°C resulted in complete conversion Form F \rightarrow Form B. FIG. 25 and FIG. 26 provide schematics of form conversion between Forms A, B, D and F.

[0552] Thermodynamic Relationship between Anhydrous Form B and Form F by slurry competition between anhydrous Form B and F. To determine the thermodynamic stability between Form B and F, competitive slurry experiments were conducted in solvent systems of acetone/EtOH/EtOAc at RT ($25 \pm 3^{\circ}\text{C}$) and 50°C as listed in Table 32. A mixture of Form B and F with equal mass ratio was suspended in saturated acetone/EtOH/EtOAc solutions of Form B and then magnetically stirred at targeted temperature. After slurry for about 4-11 days, remaining solids were isolated for XRPD characterization. A mixture of Form B and F was

observed, indicating slow transition between Form B and F.

Table 32: Slurry competition conditions for thermodynamic relationship between Compound B Form B and Form F.

Starting Form	Solvent	Temperature (°C)	Solid Form
Form B + F	Acetone	50	Form B + F (sampled after 5 days)
	EtOH	RT	Form B + F (sampled after 4 days)
		50	Form B + F (sampled after 11 days)
	EtOAc	RT	Form B + F (sampled after 8 days)

[0553] The thermodynamic stability relationship between anhydrous Form B and F was determined via slurry competition and equilibrium solubility measurement. As a result, a mixture of Form B and F was observed in all the slurry competition experiments, indicating slow transition between Form B and F. Without being bound by any particular theory, Form B and Form F low solubility may be caused by their low solubility in tested solvents (acetone/EtOH/EtOAc). Therefore, equilibrium solubility (17 hours) was measured in EtOH at 25 °C, 35 °C and 50 °C, respectively, to determine their thermodynamic stability relationship. Compared with Form B, Form F showed higher solubility under all the three tested temperatures in EtOH, indicating Form B is thermodynamically more stable than Form F from 25 °C to 50 °C.

[0554] The a_w between anhydrous Form B and hydrous Form D was determined via slurry competition under various water activity conditions at RT. Form D was observed in a_w of 0.6 and 0.8 after one week and in water after 36 days. Form B was observed in EtOH ($a_w < 0.2$) after one week. A mixture of Form D and B was observed in a_w of 0.2 and 0.4 systems after stirring at RT for 36 days. To further confirm the thermodynamic stability relationship between Form B and Form D in a_w of 0.2 and 0.4 systems at RT, their equilibrium solubility (24 hours) under the corresponding conditions was collected. Compared with that of samples that started with Form B, lower solubility was observed in samples that started with Form D (crystal forms of the final limited solids were not checked) in both a_w of 0.2 and 0.4 systems.

[0555] Equilibrium Solubility Measurement of Anhydrous Form B and F. To further determine the thermodynamic stability between Form B and F, equilibrium solubility measurement experiments were conducted in EtOH at 25 °C, 35 °C and 50 °C, respectively. Detailed procedures are summarized as follows: solids of Form B and F were suspended in 0.4 mL of EtOH at targeted temperatures and magnetically stirred for 17 hrs (750 rpm). After centrifugation, the concentration and HPLC purity of freebase in the filtrate was tested. The crystal form of remaining solids was checked by XRPD.

[0556] Compared with Form B, Form F showed higher solubility at 25 °C, 35 °C and 50 °C in EtOH (Table 33). Based on the XRPD results, no form change was observed after solubility test, indicating Form B is likely thermodynamically more stable than Form F from 25 °C to 50 °C.

Table 33: Summary of equilibrium solubility measurements of Form B and Form F in EtOH

Starting Form	Temperature (°C)	Final Form	Solubility (mg/mL)*
Form B	25	Form B	0.23
Form F		Form F	0.26
Form B	35	Form B	0.30
Form F		Form F	0.36
Form B	50	Form B	0.49
Form F		Form F	0.82

[0557] Critical Water Activity Determination between Form B and D. To determine critical water activity between anhydrous Form B and hydrous Form D, slurry competition was performed in various water activity conditions at RT as listed in Table 34. A mixture of Form B and D with equal mass ratio was suspended in saturated EtOH-water (with various a_w) solutions of Form B and then magnetically stirred at RT. After slurry for about 7-36 days, remaining solids were isolated for XRPD characterization.

Table 34: Summary of slurry competition of Form B and Form D under various a_w conditions

Starting Form	Solvent (v:v, a_w)	Sample Time (d)	Solid Form
Form B+D	EtOH	7	Form B
	EtOH:H ₂ O (97:3, 0.2)	36	Form D+B
	EtOH:H ₂ O (92.7:7.3, 0.4)	36	Form D+B
	EtOH:H ₂ O (86:14, 0.6)	7	Form D
	EtOH:H ₂ O (71:29, 0.8)	7	Form D
	H ₂ O	36	Form D

[0558] Form D was observed in a_w of 0.6 and 0.8 after slurry for one week and in water after 36 days. Form B was observed in EtOH after slurry for one week. A mixture of Form D and B was observed in a_w of 0.2 and 0.4 systems after stirring at RT for 36 days.

[0559] Solubility Measurement of Form B and D. To further confirm the thermodynamic stability relationship between Form B and Form D under a_w of 0.2 and 0.4 conditions at RT, equilibrium solubility under the conditions set forth in Table 35 was collected.

Table 35: Summary for equilibrium solubility measurement of Form B and D in $a_w=0.2/0.4$

systems

Starting Form	A_w , solvent system	Solubility (mg/mL)
Form B	0.2,	24.4
Form D	EtOH:H ₂ O (97:3, v/v)	21.7
Form B	0.4,	27.5
Form D	EtOH:H ₂ O (92.7:7.3, v/v)	18.6

[0560] Solids of anhydrous Form B and hydrous Form D were suspended in 0.5 mL of target solvent system ($a_w=0.2/0.4$), respectively, and magnetically stirred for 24 hrs (750 rpm). The suspension was filtered and the concentration of freebase in the filtrate was tested. Lower solubility was observed in samples that started with Form D. Hydrous Form D appears thermodynamically more stable than anhydrous Form B when $a_w > 0.2$ at RT (25 ± 3 °C).

Example 31:

Malonate Form M

[0561] One malonate hit with low crystallinity, malonate Form M, was obtained from screening. Its XRPD pattern is shown in FIG 30. A weight loss of 4.1% up to 110 °C was observed in TGA and DSC result showed multiple endotherms (FIG. 31). Stoichiometry was determined to be 0.50 (acid/freebase) and 6.2% THF (a molar ratio of 0.58 to freebase) was detected by ¹H NMR. Since multiple endotherms and considerable amount of THF are observed, malonate Form M is a THF solvate, but no further characterization was conducted due to its low crystallinity.

Example 32:

Fumarate Form 1

[0562] One crystalline fumarate hit, fumarate Form 1, was obtained from screening. A weight loss of 0.9% up to 150 °C was observed in TGA and DSC (FIG. 28) result showed one melting endothermic peak at 164.3 °C (onset temperature). Stoichiometry was determined to be 0.88 (acid/freebase) and 1.5% EtOAc (a molar ratio of 0.11 to freebase) was detected by ¹H NMR.

[0563] Preparation of Fumarate Form 1. About 44.7 mg of fumaric acid was weighed into a 5-mL glass vial, and 2.0 mL EtOAc was added. The acid remained undissolved. About 2.0 mL of freebase stock solution in EtOAc (-100 mg/mL) was added into the 5-mL vial with a molar ratio of 1:1, and stirred at RT. About 3 mg of fumarate Form 1 seed was added, and the solution turned cloudy. The suspension was stirred overnight and sampled by XRPD (FIG. 27) with a confirmed pattern conformed to fumarate Form 1. The suspension was stirred at 50 °C for 3 more days to increase the crystallinity followed by centrifugation. The cake was dried at 50 °C for 4 hrs. Yield: 186.6 mg, with a yield of -76.2%.

Table 36. Representative XRPD Peaks for Compound C Form 1.

2-Theta (°2θ)	d(Å)
7.589	11.6401
10.596	8.3423
11.449	7.7224
11.844	7.4662
12.5	7.0758
14.444	6.1275
15.454	5.729
15.782	5.6107
16.097	5.5017
17.555	5.0479
18.921	4.6865
19.696	4.5038
19.866	4.4657
20.233	4.3853
21.35	4.1585
22.046	4.0288
23.162	3.837
23.897	3.7206
24.238	3.669
24.672	3.6055
25.236	3.5261
25.932	3.4331

Table 37. Representative XRPD Peaks for Compound C Form 2.

2-Theta (°2θ)	d(Å)
11.521	7.6744
11.879	7.4439

2-Theta (°2θ)	d(Å)
15.558	5.691
16.04	5.5211
16.515	5.3633
17.324	5.1146
18.361	4.8281
19.004	4.6662
19.439	4.5628
19.876	4.4634
20.244	4.383
21.355	4.1574
22.035	4.0306
23.238	3.8246
23.917	3.7175
25.439	3.4985
26.039	3.4193

[0564] Summary. Compound A freebase was identified as an amorphous solid form with <0.5% weight loss before decomposition and a melting point of about 87 °C. It's hygroscopicity at 95% RH was <1% and had an epimer content of <1%. It's solubility at 37 °C was about 6.6 mg/mL. The XRPD pattern for the amorphous form is provided in FIG. 32. Compound C Form 1 was a crystalline solid with <0.5% weight loss before decomposition and a melting point of about 165-174 °C. Multiple forms of Compound C were obtained dependent upon the solvent. It's hygroscopicity at 95% RH was <1.5% and had an epimer content of <1%. It's solubility at 37 °C was about 6.6 mg/mL. The Compound B Form B was a crystalline solid with <0.5% weight loss before decomposition and a melting point of about 168 °C. A single form was present in anhydrous Form B. It's hygroscopicity at 95% RH was <1% and had an epimer content of <2.5%. It's solubility at 37 °C was about 5.9 mg/mL. The anhydrous form of Compound B Form B was found to be a stable pure crystalline form with properties more favorable than Forms D, E, and F as described herein.

Example 33:

[0565] Evaluation of safety, pharmacokinetics, and activity of Compound B. Breast cancer is the most frequent cancer diagnosed in women, with an estimated global incidence of 1.67 million new cases reported in 2012 (Ferlay et al. 2013). Breast cancer accounts for approximately 15% (approximately 522,000 cases) of all cancer deaths.

[0566] Approximately 80% of all breast cancers express the estrogen receptor (ER) factor, and the vast majority of these are dependent on ER for tumor growth and progression. Modulation of estrogen activity and/or synthesis is the mainstay of therapeutic approaches in women with ER-positive breast cancer. However, despite the effectiveness of available endocrine therapies such as ER antagonists (e.g., tamoxifen), aromatase inhibitors (e.g., anastrozole, letrozole, and exemestane) and full ER antagonists/degraders (e.g., fulvestrant), many patients ultimately relapse or develop resistance to these agents and therefore require further treatment for optimal disease control.

[0567] Despite becoming refractory to aromatase inhibitors or tamoxifen, growth and survival of resistant tumor cells remain dependent on ER signaling; therefore, patients with ER-positive breast cancer can still respond to second- or third-line endocrine treatment after progression on prior therapy (Di Leo et al. 2010; Baselga et al. 2012). There is growing evidence that in the endocrine resistant state, ER can signal in a ligand-independent manner via input from other signaling pathways (Miller et al. 2010; Van Tine et al. 2011). Without being bound by any particular theory, an agent with a dual mechanism of action (ER antagonism plus degradation) has the potential to target both ligand-dependent and ligand-independent ER signaling and, consequently, improve treatment outcomes in late-stage ER-positive breast cancer. Furthermore, recent studies have identified mutations in *ESR1* (i.e., the gene that encodes for ER α) affecting the ligand-binding domain (LBD) of the ER (Segal and Dowsett 2014). In nonclinical models, mutant ER can drive transcription and proliferation in the absence of estrogen, suggesting that LBD-mutant forms of ER may be involved in mediating clinical resistance to some endocrine therapies (Li et al. 2013; Robinson et al. 2013; Toy et al. 2013). ER antagonists that are efficacious against these ligand-independent, constitutively-active ER-mutated receptors may possess substantial therapeutic benefit.

[0568] As such, there is a need for new ER-targeting therapies with increased anti-tumor activity to further delay disease progression and/or overcome resistance to the currently available endocrine therapies and ultimately prolong survival in women with ER-positive breast cancer.

[0569] Compound B is a potent, orally bioavailable, small-molecule therapeutic agent that is being developed for the treatment of patients with ER-positive breast cancer. As provided herein Compound B, including its solid forms (e.g. Form B) are stable compounds with favorable properties for continued pharmaceutical development. Without being bound by any particular theory, Compound B appears to antagonize the effects of estrogens via competitive binding to the LBD of both wild-type and mutant ER with nanomolar potency. Upon binding, and without being bound by any particular theory, Compound B induces an inactive conformation to the ER LBD, as measured by displacement of co-activator peptides. In addition to its direct antagonist properties, without being bound by any particular theory, the mechanism of action of Compound B includes reducing levels of ER α protein through proteasome-mediated degradation. Degradation of ER is hypothesized to enable full suppression of ER signaling, which is not achieved by first-generation ER therapeutics such as tamoxifen, which display partial agonism. Compound B potently inhibits the proliferation of multiple ER-positive

breast cancer cell lines *in vitro*, including cells engineered to express clinically relevant mutations in ER.

[0570] *In vivo*, Compound B exhibited dose-dependent anti-tumor activity in xenograft models of ER-positive breast cancer, including in a patient-derived xenograft model that harbors an *ESR1* mutation (ER.Y537S). The efficacious dose range was found to be 0.1-10 mg/kg/day, and all doses were well tolerated. Fulvestrant, when dosed according to a clinically relevant dosing scheme, was less efficacious than Compound B in the assessed xenograft models. Thus, Compound B demonstrated robust nonclinical activity in ER-positive breast cancer models of *ESR1*-wildtype- and *ESR1*-mutation-bearing disease.

Example 34:

[0571] *In vitro* and *in vivo* efficacy analysis of Compound B. Compound B displays superior ER degradation & ER pathway suppression when compared to both GDC-0927 and GDC-0810. Further, Compound B has better DMPK properties than both GDC-0927 and GDC-0810 - resulting in same *in vivo* efficacy as GDC-0927 but at 100x lower doses (e.g. 1 mg or 10 mg dose). (See FIG. 34 and FIG. 35).

[0572] Pharmacokinetics and Metabolism. After a single IV administration to rats, dogs, and monkeys, Compound B was found to have a low to moderate clearance, a large volume of distribution, and a terminal elimination half-life of 7-24 hours. Oral bioavailability was moderate in rats and dogs (41%-55%) and low (17%) in monkeys. *In vitro* data showed that plasma protein binding of Compound B was high across all species, ranging from 98% to 99% bound.

[0573] *In vitro* metabolite identification experiments showed that UGT1A4-mediated glucuronidation was the major *in vitro* metabolic pathway of Compound B. The contribution from CYP450 isoforms was minor and included both CYP3A4 and CYP2C9. *In vitro* CYP inhibition studies in human liver microsomes and induction studies in human hepatocytes suggested a low-to-moderate potential for drug-drug interactions. Compound B directly inhibited CYP3A4 with 50% inhibitory concentration (IC_{50}) values of 6.5 μ M (midazolam 1'-hydroxylation) and 26 μ M (testosterone 6 β -hydroxylation); IC_{50} for CYP2B6 and CYP2C8 inhibition were 13 μ M and 21 μ M, respectively. Compound B showed weak metabolism dependent inhibition of CYP2C9.

[0574] Toxicology. Four-week Good Laboratory Practice (GLP) repeat-dose oral toxicity studies in female rats and monkeys with integrated assessments of neurologic (rats, monkey), respiratory (monkey), and cardiovascular (monkey) function were conducted to characterize the nonclinical safety profile of Compound B.

[0575] In the rat study, Compound B was tolerated at the exemplary dose levels (10, 30, and 100 mg/kg) with adverse effects predominantly in the kidneys and liver at 100 mg/kg. In the

monkey study (20, 60, and 200 mg/kg), the maximum tolerated dose (MTD) was considered to be 60 mg/kg as the high dose of 200 mg/kg was not tolerated. Adverse effects were primarily observed at the high dose level of 200 mg/kg, and lack of tolerability was attributed to kidney and liver injury and inanition.

[0576] In both rats and monkeys, there was a dose-dependent PLD observed in numerous organs at exposures that were higher than those anticipated at the human starting dose in Phase I (at least 44-fold and 6-fold based on area under the concentration-time curve [AUC], respectively), with adverse organ effects largely confined to the kidney and liver. In rats, PLD was not noted at 10 mg/kg (18-fold exposure factor), but increased in incidence and severity from 30 to 100 mg/kg. In monkeys, dose-responsive PLD was present at all doses but was limited to minimal changes in the lung at 20 mg/kg (6-fold exposure factor). These exposure multiples provide evidence that the risk of PLD-associated toxicity to humans in the Phase I starting dose is low.

[0577] The translatability of PLD from nonclinical species to patients is not certain but can be reasoned (Reasor et al. 2006). Drugs such as tamoxifen and palbociclib have not demonstrated any clinical concerns in spite of their PLD findings in nonclinical studies. Although Compound B was associated with PLD in multiple tissues in both rats and monkeys, there was no light microscopic evidence of involvement of critical organs such as heart, eyes, or neurons in these studies (Chatman et al. 2009).

[0578] Following 28-day oral administration to rats and monkeys, the increases in systemic exposure of Compound B were dose proportional. Based on the nature and reversibility of clinical signs, clinical pathology, and histopathology findings, the severely toxic dose for 10% of animals (STD₁₀) for rats was defined as 100 mg/kg, with corresponding maximum plasma concentration (C_{max}) and AUC from 0 to 24 hours (AUC₀₋₂₄) values of 6560 ng/mL and 143,000 ng • hr/mL, respectively. In monkeys, the highest non-severely toxic dose was defined as 60 mg/kg/day, with corresponding C_{max} and AUC₀₋₂₄ values of 841 ng/mL and 16,200 ng • hr/mL, respectively, due to the clinical signs and moribundities present at 200 mg/kg/day.

[0579] In summary, results from the nonclinical toxicity and safety pharmacology studies completed to date provide a robust characterization of the toxicity profile of Compound B and support administration to cancer patients in a Phase I trial.

[0580] Administration of Compound B as a Single Agent Monotherapy. Compound B demonstrated robust nonclinical activity in ER-positive breast cancer models of *ESR1*-wildtype and *ESR1*-mutation bearing disease. The safety, pharmacokinetic (PK), pharmacodynamic (PD) activity and preliminary anti-tumor activity of Compound B as a single agent was analyzed in a Phase I/IIb, multicenter, open-label study in patients with locally advanced or metastatic ER-positive breast cancer. Patients were enrolled in a dose-escalation stage with enrollment in an expansion stage to follow. During the single-agent dose escalation, cohorts were evaluated at escalating dose levels to determine the MTD or maximum administered dose (MAD).

[0581] Once the single-agent MTD or MAD has been established, an escalation cohort treated with Compound B (at or below the MTD or MAD) or Compound B in combination with palbociclib may be enrolled. Additionally, patients will be enrolled in the expansion stage and treated at or below the single-agent MTD or MAD of Compound B alone or in combination with palbociclib and/or LHRH agonist. The single-agent dose expansions may evaluate two different dose levels of Compound B with and without a LHRH agonist.

[0582] Patients were monitored for adverse events during a dose-limiting toxicity (DLT) assessment window, defined as Days -7 to 28 of Cycle 1 (single-agent cohorts). For DLT evaluation, toxicity was graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events, Version 4.0 (NCI CTCAE v4.0).

[0583] Patients enrolled in the single-agent Compound B dose-escalation stage, which consists of a screening period, a PK lead-in period, a treatment period, and a safety follow-up period. Continuous once daily dosing was commenced during the treatment period starting on Cycle 1 Day 1. The starting dose of single-agent Compound B was 10 mg, administered by mouth to patients in the first cohort continuously in 28-day cycles. The dose will be increased by up to 200% of the preceding dose level for each successive cohort, until a safety finding of concern (i.e., either a DLT, any patient with Grade \geq 2 clinically significant toxicity or overall adverse event profile inappropriate for 200% increments) is observed. Once safety findings of concern are observed, dose escalation will not exceed 100% increments.

[0584] Compound B has half-life of about 40 hours. As noted above, exposures of Compound B increased proportionally from 10 to 30 mg with normal variability.

[0585] Six patients were initiated in the initial dose of 10 mg QD on 28-day cycles as described herein. As noted in Table 38 below, all treated patients with FES-PET had qualitative near-complete (NC) or complete responses (CR). There were no observed DLTs, SAEs, AESIs, or clinically significant laboratory abnormalities in the treated patients. All related AEs were Grade 1 or Grade 2 events.

Table 38: Patient response to treatment with Compound B:

Patient	Cohort (Dose)	Days (status)	BL disease (SLD)	Response
C3	3 (90 mg)	15 (active)	Bone-only (NM)	NA
C2	3 (90 mg)	36 (active)	Lung, liver, LN, bone (>100 mm)	NA
C1	3 (90 mg)	57 (active)	Bone-only (NM)	NA
B6	2-BF (30 mg)	2 (active)	LN, Bone (87 mm)	NA
B5	2-BF (30 mg)	28 (active)	LN, Bone (35 mm)	NA
B4	2-BF (30 mg)	29 (active)	Bone-only (NM)	NA
B3	2 (30 mg)	62 (PD)	Liver (30 mm)	PD
B2	2 (30 mg)	119 (active)	Pulm/hilar (38 mm)	SD (-9%)
B1	2 (30 mg)	127 (active)	Bone-only (NM)	SD (NM)
A6	1-BF (10 mg)	64 (Clinical PD)	Liver, LN, Bone (62 mm)	SD (NM)
A5	1-BF (10 mg)	21 (Clinical PD)	Liver, LN, Bone (13 mm)	CPD

AS	1-BF (10 mg)	21 (Clinical PD)	Liver, LN, Bone (15 mm)	CRP
A4	1-BF (10 mg)	45 (Clinical PD)	Bone-only (NM)	cPD
A3	1 (10 mg)	183 (active)	Breast (50 mm)	uPR (-40%)
A2	1 (10 mg)	183 (active)	Pulmonary (36mm)	SD (-27%)
A1	1 (10 mg)	64 (PD)	Bone-only (NM)	PD

[0586] One treated patient was diagnosed with ER+PR+ breast cancer. The patient had undergone previous surgery and previous treatment with anti-cancer agents, including SERM therapy and AI therapy prior to enrollment and treatment. The patient was treated with Compound B at 10 mg and showed response to the treatment after 3 cycles as indicated in FIG. 36a and FIG. 36b.

[0587] Another patient was diagnosed with early stage HR+ breast cancer and had received prior surgery and treatment with anti-cancer agents, including cytotoxics, CDK4/6 inhibitors, and AIs. The patient was treated with Compound B at 10 mg and showed response to the treatment after 3 cycles as indicated in FIG. 37a and FIG. 37b. was enrolled on our trial at 10mg in March 2018.

[0588] The treated patients to-date demonstrate that administration of Compound B is welltolerated at 10 mg and 30 mg, with only Grade 1 and 2 AEs. In general, plasma exposures of Compound B increased proportionally with doses from 10 to 30 mg after single dose. The steady state exposures appeared to increase in a higher than dose proportional manner from 10 mg to 30 mg. The estimated half-life of about 40 hours supports once-daily dosing.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [US20160175289A \[0004\] \[0005\]](#)

Non-patent literature cited in the description


- **JONES et al.** Pharmaceutical Cocrystals: An Emerging Approach to Physical Property Enhancement MRS Bulletin, 2006, vol. 37, 875-879 [0007]
- **SINGLETON et al.** DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY. Wiley & Sons 19940000 [0027]
- **SAMBROOK et al.** MOLECULAR CLONING, A LABORATORY MANUAL Cold Springs Harbor Press 19890000 [0027]
- **P. G. M. WUTST. W. GREENE** Greene's Protective Groups in Organic Synthesis Wiley-Interscience 20060000 [0062]
- March's Advanced Organic Chemistry John Wiley & Sons 20010000 vol. 5, [0064]
- **T. W. GREENE** Protective Groups in Organic Synthesis John Wiley & Sons 19910000 [0064]

PATENTKRAV

1. Forbindelse B med strukturen:



- 5 2. Krystalform, der omfatter forbindelse B ifølge krav 1




med et røntgenpulverdiffraktionsmønster, der omfatter toppe ved 19,32, 20,26, 21,63, 23,28 eller $24,81 \pm 0.1^\circ 2\theta$ ($\pm 0.1^\circ 2\theta$).

- 10 3. Krystalform ifølge krav 2, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster med en, to, tre, fire, fem, seks, syv, otte, ni, ti, elleve, tolv, tretten, fjorten, femten, seksten, sytten, atten, nitten eller tyve karakteristiske røntgenpulverdiffraktionstoppe, der er valgt blandt:

2-Theta ($^\circ 2\theta$)	d(Å)
4,643	19,0147
8,263	10,6916
9,286	9,5157
11,183	7,9057
11,495	7,6916
11,969	7,3884
12,543	7,0515

2-Theta (°2θ)	d(Å)
13,776	6,4228
14,226	6,2207
14,619	6,0545
15,092	5,8656
15,564	5,689
16,013	5,5302
17,354	5,1058
18,559	4,777
18,847	4,7046
19,321	4,5903
19,821	4,4757
20,265	4,3785
21,343	4,1597
21,631	4,105
21,92	4,0515
22,524	3,9443
22,97	3,8686
23,285	3,8171
23,549	3,7748
23,942	3,7138
24,81	3,5858
25,964	3,4289

4. Krystalform ifølge krav 2, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster i det væsentlige som vist i fig. 1.
5. Krystalform, der omfatter forbindelse B ifølge krav 1



med et røntgenpulverdiffraktionsmønster, der omfatter toppe ved 11,49, 12,54, 19,16, 19,42 eller $24,67 \pm 0,1^\circ 2\theta$ ($\pm 0,1^\circ 2\theta$).


6. Krystalform ifølge krav 5, hvor krystalformen af forbindelse B har et
5 røntgenpulverdiffraktionsmønster med en, to, tre, fire, fem, seks, syv, otte, ni, ti, elleve, tolv, tretten, fjorten, femten, seksten, sytten, atten, nitten eller tyve karakteristiske røntgenpulverdiffraktionstoppe, der er valgt blandt:

2-Theta ($^\circ 2\theta$)	d(Å)
7,684	11,4966
11,491	7,6942
12,54	7,053
14,245	6,2123
15,303	5,7851
15,557	5,6912
16,014	5,5301
16,634	5,3252
17,371	5,1009
18,242	4,8593
19,163	4,6278
19,424	4,5662
19,892	4,4597
20,243	4,3833
21,817	4,0705
22,524	3,9442
22,996	3,8644
23,253	3,8223

2-Theta ($^{\circ}2\theta$)	d(Å)
23,573	3,7711
24,676	3,6049
25,073	3,5487
25,915	3,4353

7. Krystalform ifølge krav 5, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster i det væsentlige som vist i fig. 4.

8. Krystalform, der omfatter forbindelse B ifølge krav 1



5 med et røntgenpulverdiffraktionsmønster, der omfatter toppe ved 11,31, 15,70, 16,54, 19,10 eller $22,76 \pm 0.1^{\circ}2\theta$.


9. Krystalform ifølge krav 8, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster med en, to, tre, fire, fem, seks, syv, otte, ni, ti, elleve, tolv, tretten, fjorten, femten, seksten, sytten, atten, nitten eller tyve karakteristiske røntgenpulverdiffraktionstoppe, der er valgt blandt:

2-Theta ($^{\circ}2\theta$)	d(Å)
7,322	12,0634
10,992	8,0426
11,312	7,8161
12,182	7,2597
13,234	6,6847
13,487	6,5601
14,114	6,2697
14,668	6,0342
15,145	5,8453

2-Theta (°2θ)	d(Å)
15,702	5,6393
16,036	5,5225
16,217	5,4611
16,542	5,3546
17,249	5,1369
17,637	5,0246
18,113	4,8935
18,349	4,8311
19,108	4,641
20,209	4,3905
20,583	4,3115
21,163	4,1948
21,472	4,135
21,891	4,0569
22,762	3,9036
23,334	3,8091
23,569	3,7717

10. Krystalform ifølge krav 8, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster i det væsentlige som vist i fig. 12.

11. Krystalform, der omfatter forbindelse B ifølge krav 1



5 med et røntgenpulverdiffraktionsmønster, der omfatter toppe ved 12,52, 15,90, 19,66, 20,65 eller $24,99 \pm 0.1^\circ 2\theta$.

12. Krystalform ifølge krav 11, hvor krystalformen af forbindelse B har et røntgenpulverdiffraktionsmønster med en, to, tre, fire, fem, seks, syv, otte, ni, ti, elleve,

tolv, tretten, fjorten, femten, seksten, sytten, atten, nitten eller tyve karakteristiske røntgenpulverdiffraktionstoppe, der er valgt blandt:

2-Theta (°2θ)	d(Å)
3,925	22,4935
10,54	8,3864
11,724	7,5419
12,52	7,0646
14,227	6,2205
15,407	5,7466
15,54	5,6976
15,902	5,5687
16,488	5,3721
16,844	5,2594
17,294	5,1235
18,267	4,8526
18,473	4,799
19,399	4,572
19,661	4,5117
20,005	4,4348
20,501	4,3286
20,655	4,2968
21,161	4,1952
21,287	4,1706
21,951	4,0458
22,972	3,8683
23,498	3,783
23,708	3,7499
23,943	3,7137
24,31	3,6584
24,679	3,6044
24,997	3,5594

13. Krystalform ifølge krav 11, hvor krystalformen af forbindelse B har et

røntgenpulverdiffraktionsmønster i det væsentlige som vist i fig. 16.

14. Farmaceutisk sammensætning, der omfatter forbindelsen ifølge krav 1 og/eller den faste form ifølge krav 7 og mindst én farmaceutisk acceptabel excipients.


15. Forbindelse ifølge krav 1, fast form ifølge krav 7 eller farmaceutisk sammensætning ifølge krav 14 til anvendelse i en fremgangsmåde til behandling af lungecancer, ovariecancer, endometriecancer, prostatacancer, livmodercancer eller brystcancer hos en patient med denne cancer, hvilken fremgangsmåde omfatter indgivelse af en virksom mængde af en forbindelse ifølge krav 1, en fast form ifølge krav 7 eller den farmaceutiske sammensætning ifølge krav 14 til cancerpatienten.

10 16. Forbindelse ifølge krav 1, fast form ifølge krav 7 eller farmaceutisk sammensætning ifølge krav 14 til anvendelse i en fremgangsmåde til behandling af brystcancer hos en brystcancerpatient, hvilken fremgangsmåde omfatter indgivelse af en virksom mængde af en forbindelse ifølge krav 1, et fast stof ifølge krav 7 eller den farmaceutiske sammensætning ifølge krav 14 til cancerpatienten, hvor brystcanceren er
15 hormonreceptorpositiv brystcancer, HER2-positiv brystcancer eller tredobbelt negativ brystcancer.


17. Forbindelse, fast form eller farmaceutisk sammensætning til anvendelse ifølge krav 16, hvor brystcanceren er metastatisk brystcancer.

18. Forbindelse, fast form eller farmaceutisk sammensætning til anvendelse
20 ifølge et hvilket som helst af kravene 16-17, hvor patienten har fået tidligere behandling med et eller flere anticancermidler eller strålebehandling.


19. Proces til fremstilling af forbindelse B,




hvor processen omfatter trinnene:




20. Proces ifølge krav 19, hvor processen yderligere omfatter omkrystallisation af forbindelse B i methanol og ethanol;



5 21. Proces ifølge krav 19, hvor processen yderligere omfatter omkrystallisation af forbindelse B i MTBE, vand, NaOH og ethanol;



22. Proces ifølge et hvilket som helst af kravene 19-21, hvor indolymellemprodukterne syntetiseres i overensstemmelse med trinnene:



DRAWINGS






FIG. 1

**FIG. 2A****FIG. 2B**




FIG. 3




FIG. 4







FIG. 5

**FIG. 6****FIG. 7**

**FIG. 8**




FIG. 9B




FIG. 9A




FIG. 9C




FIG. 10




FIG. 11




FIG. 13




FIG. 12




FIG. 14




FIG. 15




FIG. 17






FIG. 16

**FIG. 18****FIG. 20**




FIG. 19




FIG. 21






FIG. 22

**FIG. 23****FIG. 24**




FIG. 25




FIG. 26




FIG. 27A





FIG. 27B

**FIG. 28**




FIG. 29




FIG. 30




FIG. 31




FIG. 33




FIG. 32




FIG. 34






FIG. 35

FES-PET (PD) Response




Screening FES-PET




Cycle 3 FES-PET (~100%)

FIG. 36B

Radiographic Minor Response



Screening CT



Cycle 3 CT (~27%)

FIG. 36A

Radiographic uPR Response




Screening MRI



Cycle 3 MRI (-40% uPR)

FIG. 37A

FES-PET (PD) Response



Screening FES-PET



Cycle 3 FES-PET (-100%)

FIG. 37B