



(86) Date de dépôt PCT/PCT Filing Date: 2000/12/11
 (87) Date publication PCT/PCT Publication Date: 2001/06/28
 (85) Entrée phase nationale/National Entry: 2002/06/21
 (86) N° demande PCT/PCT Application No.: US 2000/032443
 (87) N° publication PCT/PCT Publication No.: 2001/046199
 (30) Priorités/Priorities: 1999/12/22 (60/171,373) US;
 2000/08/17 (60/226,076) US; 2000/09/22 (60/234,539) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 498/04, A61K 31/47, A61P 43/00,
 A61P 35/00


(71) Demandeurs/Applicants:
 ELL LILLY AND COMPANY, US;
 WANG, QIUPING, US

(72) Inventeurs/Inventors:
 BONJOUKLIAN, ROSANNE, US;
 COHEN, JEFFREY DANIEL, US;
 GRUBER, JOSEPH MICHAEL, US;
 JOHNSON, DOUGLAS WEBB, US;
 JUNGHEIM, LOUIS NICKOLAUS, US;

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : METHODES ET COMPOSES DESTINES A INHIBER MRP1

(54) Title: METHODS AND COMPOUNDS FOR INHIBITING MRP1



(57) Abrégé/Abstract:

The present invention further relates to a method of inhibiting MRP1 in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula (I).

(72) **Inventeurs(suite)/Inventors(continued)**: KROIN, JULIAN STANLEY, US; LANDER, PETER AMBROSE, US;
LIN, HO-SHEN, US; LOHMAN, MARK CHRISTOPHER, US; MUEHL, BRIAN STEPHEN, US;
NORMAN, BRYAN HURST, US; PATEL, VINOD FRANCIS, US; RICHETT, MICHAEL ENRICO, US;
THRASHER, KENNETH JEFF, US; VEPACHEDU, SREENIVASARAO, US; WHITE, WESLEY TODD, US;
XIE, YONGPING, US; YORK, JEREMY SCHULENBURG, US; PARKHURST, BRANDON LEE, US; WANG, QIUPING, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 June 2001 (28.06.2001)

PCT

(10) International Publication Number
WO 01/46199 A1(51) International Patent Classification⁷: C07D 498/04,
A61K 31/47, A61P 35/00, 43/00

(21) International Application Number: PCT/US00/32443

(22) International Filing Date:
11 December 2000 (11.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/171,373 22 December 1999 (22.12.1999) US
60/226,076 17 August 2000 (17.08.2000) US
60/234,539 22 September 2000 (22.09.2000) US(71) Applicants (for all designated States except US): ELI
LILLY AND COMPANY [US/US]; Lilly Corporate
Center, Indianapolis, IN 46285 (US). WANG, Qiuping
[US/US]; 8430 Harrison Parkway, Fishers, IN 46038 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): BONJOUKLIAN,
Rosanne [US/US]; 318 Dominion Drive, Zionsville, IN
46077 (US). COHEN, Jeffrey, Daniel [US/US]; 1411
Shawnee Road, Indianapolis, IN 46260 (US). GRU-
BER, Joseph, Michael [US/US]; 9272 Shady Bend,
Brownsburg, IN 46112 (US). JOHNSON, Douglas,
Webb [US/US]; 235 Saddlebrook Court, Zionsville, IN
46077 (US). JUNGHEIM, Louis, Nickolaus [US/US];
8218 Meadowbrook Dive, Indianapolis, IN 46240 (US).
KROIN, Julian, Stanley [US/US]; 8418 Hilltop Drive,
Indianapolis, IN 46234 (US). LANDER, Peter, Ambrose
[US/US]; 5407 North Capitol Avenue, Indianapolis, IN
46208 (US). LIN, Ho-Shen [US/US]; 8128 Trevellian
Way, Indianapolis, IN 46217 (US). LOHMAN, Mark,
Christopher [US/US]; 1307 Baker Street, Apartment C,Longmont, CO 80501 (US). MUEHL, Brian, Stephen
[US/US]; 940 Country Lane, Indianapolis, IN 46217
(US). NORMAN, Bryan, Hurst [US/US]; 8648 Admi-
rals Bay Drive, Indianapolis, IN 46236 (US). PATEL,
Vinod, Francis [US/US]; 3 Mossy Lane, Bellows Farm,
Acton, MA 01720 (US). RICHETT, Michael, Enrico
[US/US]; 5832 Baron Court, Indianapolis, IN 46250 (US).
THRASHER, Kenneth, Jeff [US/US]; 8660 Count Turf
Court, Indianapolis, IN 46217 (US). VEPACHEDU,
Sreenivasarao [IN/US]; 16 West 630 Mockingbird Lane,
Apartment 20G, Hinsdale, IL 60521 (US). WHITE,
Wesley, Todd [US/US]; 1434 Grube Street, Indianapolis,
IN 46227 (US). XIE, Yongping [CN/US]; 19 Hunting-
ton Circle, Apartment 15, Naperville, IL 60540 (US).
YORK, Jeremy, Schulenburg [US/US]; 3592 North
Tillotson Avenue, Number 87, Muncie, IN 47304 (US).
PARKHURST, Brandon, Lee [US/US]; 144 Jonquill
Drive, Indianapolis, IN 46227 (US).(74) Agents: DAWALT, Elizabeth, A. et al.; Eli Lilly and Com-
pany, Lilly Corporate Center, Indianapolis, IN 46285 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

[Continued on next page]

(54) Title: METHODS AND COMPOUNDS FOR INHIBITING MRP1

(57) Abstract: The present invention
further relates to a method of inhibiting
MRP1 in a mammal which comprises
administering to a mammal in need
thereof an effective amount of a
compound of formula (I).

WO 01/46199 A1



— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 341

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 341

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

METHODS AND COMPOUNDS FOR INHIBITING MRP1

BACKGROUND

5 Along with surgery and radiotherapy, chemotherapy continues to be an effective therapy for many cancers. In fact, several types of cancer, such as Hodgkin's disease, large cell lymphoma, acute lymphocytic leukemia, testicular cancer and early stage breast cancer, are now considered to be curable by chemotherapy. Other cancers such as ovarian cancer, small cell lung and advanced breast cancer, while not yet curable, are exhibiting positive response to combination chemotherapy.

10 One of the most important unsolved problems in cancer treatment is drug resistance. After selection for resistance to a single cytotoxic drug, cells may become cross resistant to a whole range of drugs with different structures and cellular targets, *e.g.*, alkylating agents, antimetabolites, hormones, platinum-containing drugs, and natural products. This phenomenon is known as multidrug resistance (MDR). In some types of cells, this resistance is inherent, while in others, such as small cell lung cancer, it is usually acquired.

15 Such resistance is known to be multifactorial and is conferred by at least two proteins: the 170 kDa P-glycoprotein (MDR1) and the more recently identified 190 kDa multidrug resistance protein (MRP1). Although both MDR1 and MRP1 belong to the ATP-binding cassette superfamily of transport proteins, they are structurally very different molecules and share less than 15% amino acid homology. Despite the structural divergence between the two proteins, by 1994 there were no known consistent differences in the resistance patterns of MDR1 and MRP1 cell lines. However, the association, or lack thereof, of MRP1 and resistance to particular oncolytics is known. See Cole, *et. al.*,
20 "Pharmacological Characterization of Multidrug Resistant MRP-transfected Human Tumor Cells", *Cancer Research*, 54:5902-5910, 1994. Doxorubicin, daunorubicin, epirubicin, vincristine, paclitaxel, mitoxantrone, melphalan, and etoposide are substrates of MRP1, *i.e.*, MRP1 can bind to these oncolytics and redistribute them away from their site of action, the nucleus, and out of the cell. *Id.* and Marquardt, D., and Center, M.S.,
25 *Cancer Research*, 52:3157, 1992.

30 Doxorubicin, daunorubicin, and epirubicin are members of the anthracycline class of oncolytics. They are isolates of various strains of *Streptomyces* and act by inhibiting nucleic acid synthesis. These agents are useful in treating neoplasms of the bone, ovaries, bladder, thyroid, and especially the breast. They are also useful in the treatment of acute

-2-

lymphoblastic and myeloblastic leukemia, Wilm's tumor, neuroblastoma, soft tissue sarcoma, Hodgkin's and non-Hodgkin's lymphomas, and bronchogenic carcinoma.


Vincristine, a member of the vinca alkaloid class of oncolytics, is an isolate of a common flowering herb, the periwinkle plant (*Vinca rosea* Linn). The mechanism of action of vincristine is still under investigation but has been related to the inhibition of microtubule formation in the mitotic spindle. Vincristine is useful in the treatment of acute leukemia, Hodgkin's disease, non-Hodgkin's malignant lymphomas, rhabdomyosarcoma, neuroblastoma, and Wilm's tumor.

Etoposide, a member of the epipodophyllotoxin class of oncolytics, is a semisynthetic derivative of podophyllotoxin. Etoposide acts as a topoisomerase inhibitor and is useful in the therapy of neoplasms of the testis, and lung.

It is presently unknown what determines whether a cell line will acquire resistance via a MDR1 or MRP1 mechanism. Due to the tissue specificity of these transporters and/or in the case where one mechanism predominates or is exclusive, it would be useful to have a selective inhibitor of that one over the other. Furthermore, when administering a drug or drugs that are substrates of either protein, it would be particularly advantageous to coadminister an agent that is a selective inhibitor of that protein. It is, therefore, desirable to provide compounds that are selective inhibitors of MDR1 or MRP1.

SUMMARY OF THE INVENTION

The present invention relates to a compound of formula:



25

where:

A is a C₃-C₈ cycloalkyl, optionally substituted 1-3 times with a C₁-C₄ alkyl;

het is a five (5) membered heterocyclic ring comprising N and a second heteroatom selected from N, O, or S;

-3-

wherein the non-fused carbon atom of the heteroaryl ring may be optionally substituted with R^b: C₁-C₆ alkyl, optionally substituted aryl, optionally substituted heterocycle, an amino acid ester, CH₂OH, CH₂O-heterocycle, halo, CH₂N₃, CH₂SR¹, CH₂NR⁴R⁶, OR¹, SR¹³,
 5 S(CH₂)_k-phenyl, or NR⁴R⁶; provided that when het is pyrazole or imidazole, the saturated nitrogen of the het ring may be optionally substituted with R^a: C₁-C₄ alkyl;

k is 0, 1, 2, 3, or 4;

10 n is 0, 1, or 2;

p is 0 or 1;

q is 0, 1, or 2;

r is 0, 1, or 2;

t is 0, 1, 2, 3, or 4;

15 u is 0, 1, 2, 3, or 4;

Y is -E-C(O)R³, -E-CH=CHR¹³, -E-C(OH)R¹³, -E-NR⁴R⁵, -E-OR², -E-S(O)_qR¹³, -E-SO₂NR⁴R⁶, -C(R¹¹)=NR⁶, or an optionally substituted heterocycle;

20 E is a bond or -C(R¹¹)(R¹¹)-;

R¹ is independently at each occurrence hydrogen or C₁-C₆ alkyl;


25 R² is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, or optionally substituted heterocycle, C(O)-aryl, or (CH₂)₂NR⁴R⁵;

30 R³ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted heterocycle, OR¹³, or NR⁴R⁶;

35 R⁴ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted (C₁-C₆ alkyl)-aryl, optionally substituted aryl, or R⁴ and R⁵, R⁶, R^{6'} combine to form =CR¹R¹⁴;

-4-

R⁵ is independently at each occurrence hydrogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, optionally substituted heterocycle, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl; optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, C(O)C(O)R¹³,

C(O)R⁷, CH₂R⁷, SO₂R⁸, a moiety of the formula , or R⁴ and R⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R⁶ is independently at each occurrence hydrogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, or R⁴ and R⁶, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R^{6'} is independently at each occurrence hydrogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, (C₁-C₄ alkyl)-OR¹³:

wherein the (C₁-C₄ alkyl) of the (C₁-C₄ alkyl)-OR¹³ may be optionally substituted from 1 to 2 times with C₁-C₄ alkyl, optionally substituted aryl, optionally substituted heterocycle; or R⁴ and R^{6'}, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R⁷ is independently at each occurrence optionally substituted C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₁-C₄ alkoxy)-aryl, (C₁-C₄ alkoxy)-heterocycle, (C₁-C₄ alkoxy)-SiCH₃, optionally substituted (C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-(C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, diphenylmethyl, optionally substituted (C₁-C₄ alkyl)-CO-aryl, optionally substituted CO-aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted CH=CH-heterocycle, optionally substituted phenoxy, optionally substituted heterocycle, optionally

-5-

substituted (C₁-C₄ alkyl)-phenoxy, (CH₂)_tS(O)_rR¹, (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵),
 (CH₂)_tC(R¹²)(R⁹)O(R¹⁷), (CH₂)_tC(R¹²)(R⁹)S(R¹⁷), or NR⁴R^{6'};

R⁸ is independently at each occurrence optionally substituted C₁-C₆ alkyl,
 5 optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally
 substituted (C₁-C₄ alkyl)-heterocycle, or optionally substituted heterocycle;

R⁹ is independently at each occurrence hydrogen, optionally substituted C₁-C₆
 alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl,
 10 optionally substituted aryl, optionally substituted heterocycle, (CH₂)_u-(C₁-C₆ alkoxy),
 optionally substituted (CH₂)_u-O-(C₃-C₈ cycloalkyl), optionally substituted (CH₂)_u-
 (C₁-C₄ alkoxy)-aryl, optionally substituted (CH₂)_u-O-aryl, optionally substituted
 (CH₂)_u-O-heterocycle, (C₁-C₄ alkyl)-CO₂-(C₁-C₆ alkyl), optionally substituted (C₁-C₄
 15 alkyl)-CO₂-(C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-CO₂-(C₁-C₄
 alkyl)-aryl, optionally substituted (C₁-C₄ alkyl)-CO₂-aryl, optionally substituted (C₁-C₄
 alkyl)-CO₂-heterocycle, or R⁹ and R¹² can combine to form a C₃-C₈ cycloalkyl;

R¹⁰ is 0 to 4 substituents from the aryl ring independently at each occurrence
 hydrogen, halo, C(O)R³, cyano, optionally substituted heterocycle, optionally substituted
 20 aryl, C≡C-R¹, C₁-C₄ alkoxy, (C₁-C₄ alkyl)-phenyl, NR¹⁹R²⁰, or C₂-C₆ alkenyl;

R¹¹ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally
 substituted heterocycle, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally
 substituted aryl, or optionally substituted (C₁-C₄ alkyl)-aryl;

25

R¹² is independently at each occurrence hydrogen, optionally substituted C₁-C₆
 alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl,
 optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle or optionally
 substituted heterocycle;

30

R¹³ is independently at each occurrence hydrogen, optionally substituted C₁-C₆
 alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl,
 optionally substituted aryl, CO₂CH₂CO₂CH₂CH₃, or optionally substituted heterocycle;

-6-

R¹⁴ is independently at each occurrence C₁-C₆ alkyl or optionally substituted (C₁-C₄ alkyl)-aryl;

5 R¹⁵ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, C(O)OR¹³, SO₂R⁸, C(O)R¹⁸, or a

moiety of the formula $\begin{array}{c} \text{S} \\ \parallel \\ \text{---} \text{NHR}^8 \end{array}$;

10 R¹⁶ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted aryl, optionally substituted heterocycle, SO₂R⁸, or -COR⁸; or R¹⁶ and R¹⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

15 R¹⁷ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, COR¹⁸, optionally substituted heterocycle, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted C₁-C₆ alkoxy, optionally substituted (C₁-C₄ alkoxy)-aryl, optionally substituted (C₁-C₄ alkoxy)-heterocycle, (C₁-
20 C₄ alkyl)-N(R¹)(R¹), or an amino acid ester;

R¹⁸ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted heterocycle, (C₁-C₄ alkyl)-NHCO₂-
25 (C₁-C₄ alkyl), or optionally substituted (C₁-C₄ alkyl)-heterocycle;

R¹⁹ is independently at each occurrence hydrogen, or optionally substituted C₁-C₆ alkyl;

30 R²⁰ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, CH₂OH, CO-(C₁-C₄ alkyl); or a pharmaceutical salt thereof.

-7-

The present invention further relates to a method of inhibiting MRP1 in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula I.

5 In another embodiment, the present invention relates to a method of inhibiting a resistant neoplasm, or a neoplasm susceptible to resistance in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula I in combination with an effective amount of an oncolytic agent.

10 The present invention also relates to a pharmaceutical formulation comprising a compound of formula I in combination with one or more oncolytics, pharmaceutical carriers, diluents, or excipients therefor.

DETAILED DESCRIPTION OF THE INVENTION

15 The current invention concerns the discovery that compounds of formula I are selective inhibitors of multidrug resistant protein (MRP1), and are thus useful in treating MRP1 conferred multidrug resistance (MDR) in a resistant neoplasm and a neoplasm susceptible to resistance.

20 The terms "inhibit" as it relates to MRP1 and "inhibiting MRP1" refer to prohibiting, alleviating, ameliorating, halting, restraining, slowing or reversing the progression of, or reducing MRP1's ability to redistribute an oncolytic away from the oncolytic's site of action, most often the neoplasm's nucleus, and out of the cell. Additionally, these terms refer to (repeat to redistribute) an MRP1 substrate away from the substrate's site of action.

25 As used herein, the term "effective amount of a compound of formula I" refers to an amount of a compound of the present invention which is capable of inhibiting MRP1. The term "effective amount of an oncolytic agent" refers to an amount of oncolytic agent capable of inhibiting a neoplasm, resistant or otherwise.

30 The term "inhibiting a resistant neoplasm, or a neoplasm susceptible to resistance" refers to prohibiting, halting, restraining, slowing or reversing the progression of, reducing the growth of, or killing resistant neoplasms and/or neoplasms susceptible to resistance.

35 The term "resistant neoplasm" refers to a neoplasm, which is resistant to chemotherapy where that resistance is conferred in part, or in total, by MRP1. Such neoplasms include, but are not limited to, neoplasms of the bladder, bone, breast, lung(small-cell), testis, and thyroid and also includes more particular types of cancer such as, but not limited to, acute lymphoblastic and myeloblastic leukemia, Wilm's tumor,

-8-

neuroblastoma, soft tissue sarcoma, Hodgkin's and non-Hodgkin's lymphomas, and bronchogenic carcinoma.

A neoplasm, which is "susceptible to resistance", is a neoplasm where resistance is not inherent nor currently present but can be conferred by MRP1 after chemotherapy
5 begins. Thus, the methods of this invention encompass a prophylactic and therapeutic administration of a compound of formula I.

The term "chemotherapy" refers to the use of one or more oncolytic agents where at least one oncolytic agent is a substrate of MRP1. A "substrate of MRP1" is an oncolytic that binds to MRP1 and is redistributed away from the oncolytic's site of action
10 (the nucleus of the neoplasm) and out of the cell, thus, rendering the therapy less effective. Preferred oncolytic agents are camptosar, vinorelbine, mitoxantrone, doxorubicin, daunorubicin, epirubicin, vincristine, and etoposide.

The terms "treat" or "treating" bear their usual meaning which includes preventing, prohibiting, alleviating, ameliorating, halting, restraining, slowing or
15 reversing the progression, or reducing the severity of MRP1 derived drug resistance in a multidrug resistant tumor.

In the general formulae of the present document, the general chemical terms have their usual meanings. For example, the term "C₁-C₄ alkyl" refers to methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, cyclobutyl, *s*-butyl, and *t*-butyl. The term "C₁-C₆
20 alkyl" refers to a monovalent, straight, branched, or cyclic saturated hydrocarbon containing from 1 to 6 carbon atoms. Additionally, the term "C₁-C₆ alkyl" includes C₁-C₄ alkyl groups and C₃-C₆ cycloalkyls. The term "C₁-C₆ alkyl" includes, but is not limited to, cyclopentyl, pentyl, hexyl, cyclohexyl, and the like.

The term "C₃-C₈ cycloalkyl" refers to cyclopropyl, cyclobutyl, cyclopentyl,
25 cyclohexyl, cycloheptyl, and cyclooctyl. The term "C₅-C₇ cycloalkyl" refers to cyclopentyl, cyclohexyl, and cycloheptyl.

The term "C₆-C₁₀ bicycloalkyl" refers to bicyclo-[2.1.1]hexanyl, [2.2.1]heptanyl, [3.2.1]octanyl, [2.2.2]octanyl, [3.2.2]nonanyl, [3.3.1]nonanyl, [3.3.2]decanyl, and [4.3.1]decanyl ring systems; and benzofused ring systems including 1,2,3,4-
30 tetrahydronaphthalene, indane, 1,2-dihydrocyclobuta[1,2-*a*]benzene, and hydrocyclopropa[1,2-*a*]benzene.

The terms "optionally substituted C₁-C₄ alkyl" and "optionally substituted C₁-C₆ alkyl" refers to a C₁-C₄ alkyl or C₁-C₆ alkyl, respectively, unsubstituted or substituted from 1 to 3 times with halo, C₁-C₄ alkanol, NH₂, or hydroxy.

The terms "C₁-C₄ alkoxy" and "C₁-C₆ alkoxy" refer to moieties of the formula
35 O-(C₁-C₄ alkyl) and O-(C₁-C₆ alkyl) respectively.

-9-

The term "optionally substituted C₃-C₈ cycloalkyl" refers to a C₃-C₈ cycloalkyl unsubstituted or substituted once with a phenyl, substituted phenyl, hydroxy, or CO₂R¹ group.

5 The terms "optionally substituted (C₁-C₄ alkyl)-(C₃-C₈ cycloalkyl)" refers to optionally substituted C₃-C₈ cycloalkyl linked through an optionally substituted C₁-C₄ alkyl.

The term "optionally substituted O-(C₃-C₈ cycloalkyl)" refers to an optionally substituted C₃-C₈ cycloalkyl linked through an oxygen atom.

10 The term "optionally substituted C₆-C₁₀ bicycloalkyl" refers to a C₆-C₁₀ bicycloalkyl unsubstituted or substituted once with a phenyl, substituted phenyl, or CO₂R¹ group.

The term "halo" or "halide" refers to fluoro, chloro, bromo, and iodo.

The term "aryl" refers to phenyl, and naphthyl.

15 The term "optionally substituted aryl" refers to a phenyl and naphthyl group, respectively, unsubstituted or substituted from 1 to 5 times independently with C₁-C₆ alkyl, C₁-C₄ alkoxy, halo, hydroxy, trifluoromethyl, phenyl, phenoxy, SO₂R¹, OR¹¹; NR⁴R⁵, SO₂N(R¹³)₂, NH-Pg, C₁-C₆ alkoxy, benzyloxy, C(O)R¹³, C₅-C₇ cycloalkyl, trifluoromethoxy, SR¹, cyano, or nitro.

20 The term "optionally substituted (C₁-C₄ alkyl)-aryl" refers to optionally substituted aryl linked through an optionally substituted C₁-C₄ alkyl.

The term "optionally substituted O-aryl" refers to an optionally substituted aryl linked through an oxygen atom.

25 The term "optionally substituted phenoxy" refers to a phenoxy group unsubstituted or substituted from 1 to 3 times independently with C₁-C₆ alkyl, halo, hydroxy, trifluoromethyl, NR⁴R⁶, SO₂N(R¹³)₂, NH-Pg, C₁-C₆ alkoxy, benzyloxy, C(O)R¹³, C₅-C₇ cycloalkyl, trifluoromethoxy, cyano, or nitro.

The term "optionally substituted (C₁-C₄ alkyl)-phenoxy" refers to unsubstituted or substituted phenoxy linked through an optionally substituted C₁-C₄ alkyl.

30 The term "heterocycle" is taken to mean stable unsaturated and saturated 3 to 6 membered rings containing from 1 to 4 heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, said rings being optionally benzofused. All of these rings may be substituted with up to three substituents independently selected from the group consisting of halo, C₁-C₄ alkoxy, C₁-C₄ alkyl, cyano, nitro, hydroxy, -S(O)_m-(C₁-C₄ alkyl) and -S(O)_m-phenyl where m is 0, 1 or 2. Saturated rings include, for example, pyrrolidinyl, 35 azetidine, piperidinyl, piperazinyl, tetrahydrofuryl, oxazolidinyl, morpholino, dioxanyl,

-10-

pyranyl, and the like. Benzofused saturated rings include indolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl and the like. Unsaturated rings include furyl, thienyl, pyridinyl, pyrrolyl, N-methylpyrrolyl, oxazolyl, isoxazolyl, pyrazolyl, imidazolyl, triazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, pyrimidinyl, pyrazinyl, thiophenyl, pyridazinyl, and the like. Benzofused unsaturated rings include isoquinolinyl, benzoxazolyl, benzthiazolyl, quinolinyl, benzofuranyl, thionaphthyl, furanopyridine, cinnolinyl, thiophenopyridine, indolyl and the like.

The term "heteroaryl" is taken to mean an unsaturated or benzofused unsaturated heterocycle as defined in the previous paragraph.

The term "optionally substituted heterocycle" refers to a heterocyclic ring unsubstituted or substituted 1 or 3 times independently with a C₁-C₆ alkyl, halo, benzyl, optionally substituted phenyl, SR¹, C₁-C₄ alkoxy, CO₂R¹, nitro, cyano, (C₁-C₄ alkyl)-cyano, heterocycle, NR¹⁹R²⁰, COR¹², C₁-C₆ alkanol, benzyloxy, phenoxy, trifluoromethyl. Heterocyclic rings may be additionally substituted 1 or 2 times with an oxo group.

The term "optionally substituted O-heterocycle" refers to an optionally substituted heterocycle linked through an oxygen atom.

The term "optionally substituted (C₁-C₄ alkyl)-heterocycle" refers to optionally substituted heterocycle linked through an optionally substituted C₁-C₄ alkyl.

The term "N-heterocycle" refers to a nitrogen containing heterocycle linked through a nitrogen atom.

The term "optionally substituted N-heterocycle" refers to a N-heterocycle, optionally substituted 1 or 3 times independently with a C₁-C₆ alkyl, halo, benzyl, optionally substituted phenyl, SR¹, C₁-C₄ alkoxy, CO₂R¹, nitro, cyano, (C₁-C₄ alkyl)-cyano, heterocycle, NR¹⁹R²⁰, COR¹², C₁-C₆ alkanol, benzyloxy, phenoxy, trifluoromethyl; and additionally substituted 1 or 2 times with an oxo group.

The term "amino acid ester" as used in this specification refers to an amino acid, where the carboxy group is substituted with a C₁-C₆ alkyl or benzyl group. That is, the alkyl group when taken together with the carboxy group forms a C₁-C₆ alkyl ester. A skilled artisan would appreciate that some amino acids have two carboxy groups that may be substituted with a C₁-C₆ alkyl group, for example, aspartic acid and glutamic acid. This invention contemplates the possibility of amino acid mono- or diesters in these circumstances.

The term "amino acid" refers to a chemical unit made up of both a basic amino group and an acidic carboxyl group. Examples of amino acids include alanine,

-11-

asparagine, cysteine, glutamine, glycine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, aspartic acid, glutamic acid, arginine, histidine, and lysine.

5 The term "protecting group" (Pg) refers to an amino protecting group or a hydroxy protecting group. The species of protecting group will be evident from whether the "Pg" group is attached to a nitrogen atom (amino protecting group) or attached to an oxygen atom (hydroxy protecting group).

10 The term "amino protecting group" as used in this specification refers to a substituent(s) of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting groups include the formyl group, the trityl group, the phthalimido group, the acetyl group, the trichloroacetyl group, the chloroacetyl, bromoacetyl, and iodoacetyl groups, urethane-type blocking groups such as benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl ("Fmoc"), and the like; and like amino protecting groups.

15 The species of amino protecting group employed is not critical so long as the derivitized amino group is stable to the condition of subsequent reaction(s) on other positions of the molecule and can be removed at the appropriate point without disrupting the remainder of the molecule. Similar amino protecting groups used in the cephalosporin, penicillin, and peptide arts are also embraced by the above terms. Further examples of groups referred

20 to by the above terms are described by T.W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1991, Chapter 7 hereafter referred to as "Greene". A preferred amino protecting group is *t*-butyloxycarbonyl.

The term "hydroxy protecting group" denotes a group understood by one skilled in the organic chemical arts of the type described in Chapter 2 of Greene. Representative

25 hydroxy protecting groups include, for example, ether groups including methyl and substituted methyl ether groups such as methyl ether, methoxymethyl ether, methylthiomethyl ether, *tert*-butylthiomethyl ether, (phenyldimethylsilyl)methoxy-methyl ether, benzyloxymethyl ether, *p*-methoxybenzyloxy-methyl ether, and *tert*-butoxymethyl ether; substituted ethyl ether groups such as ethoxyethyl ether, 1-(2-chloroethoxy)-ethyl

30 ether, 2,2,2-trichloroethoxymethyl ether, and 2-(trimethylsilyl)ethyl ether; isopropyl ether groups; phenyl and substituted phenyl ether groups such as phenyl ether, *p*-chlorophenyl ether, *p*-methoxyphenyl ether, and 2,4-dinitrophenyl ether; benzyl and substituted benzyl ether groups such as benzyl ether, *p*-methoxybenzyl ether, *o*-nitrobenzyl ether, and 2,6-dichlorobenzyl ether; and alkylsilyl ether groups such as trimethyl- triethyl- and

35 triisopropylsilyl ethers, mixed alkylsilyl ether groups such as dimethylisopropylsilyl ether,

-12-

and diethylisopropylsilyl ether; and ester protecting groups such as formate ester, benzylformate ester, mono-, di-, and trichloroacetate esters, phenoxyacetate ester, and *p*-chlorophenoxyacetate and the like. The species of hydroxy protecting group employed is not critical so long as the derivatized hydroxy group is stable to the conditions of subsequent reaction(s) on other positions of the intermediate molecule and can be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other hydroxy protecting group(s).

The term "carbonyl activating group" refers to a substituent of a carbonyl that increases the susceptibility of that carbonyl to nucleophilic addition. Such groups include, but are not limited to, alkoxy, aryloxy, nitrogen containing unsaturated heterocycles, or amino groups such as oxybenzotriazole, imidazolyl, nitrophenoxy, pentachloro-phenoxy, N-oxysuccinimide, N,N'-dicyclohexylisoure-O-yl, N-hydroxy-N-methoxyamino, and the like; acetates, formates, sulfonates such as methanesulfonate, ethanesulfonate, benzenesulfonate, or *p*-toluenylsulfonate, and the like; and halides especially chloride, bromide, or iodide.

The term "carbonyl activating reagent" refers to a reagent that converts the carbonyl of a carboxylic acid group to one that is more prone to nucleophilic addition and includes, but is not limited to, such reagents as those found in "The Peptides", Gross and Meienhofer, Eds., Academic Press (1979), Ch. 2 and M. Bodanszky, "Principles of Peptide Synthesis", 2nd Ed., Springer-Verlag Berlin Heidelberg, 1993, hereafter referred to as "The Peptides" and "Peptide Synthesis" respectively. Specifically, carbonyl activating reagents include thionyl bromide, thionyl chloride, oxalyl chloride, and the like; alcohols such as nitrophenol, pentachlorophenol, and the like; amines such as N-hydroxy-N-methoxyamine and the like; acid halides such as acetic, formic, methanesulfonic, ethanesulfonic, benzenesulfonic, or *p*-tolylsulfonic acid halide, and the like; and compounds such as 1,1'-carbonyldiimidazole, benzotriazole, imidazole, N-hydroxysuccinimide, dicyclohexylcarbodiimide, and the like.

In general, the term "pharmaceutical" when used as an adjective means substantially non-toxic to living organisms. For example, the term "pharmaceutical salt" as used herein, refers to salts of the compounds of formula I which are substantially non-toxic to living organisms. See, *e.g.*, Berge, S.M, Bighley, L.D., and Monkhouse, D.C., "Pharmaceutical Salts", *J. Pharm. Sci.*, 66:1, 1977. Typical pharmaceutical salts include those salts prepared by reaction of the compounds of formula I with an inorganic or organic acid or base. Such salts are known as acid addition or base addition salts respectively. These pharmaceutical salts frequently have enhanced solubility

-13-

characteristics compared to the compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions.

The term "acid addition salt" refers to a salt of a compound of formula I prepared by reaction of a compound of formula I with a mineral or organic acid. For
5 exemplification of pharmaceutical acid addition salts see, *e.g.*, Berge, S.M, Bighley, L.D., and Monkhouse, D.C., *J. Pharm. Sci.*, 66:1, 1977. Since compounds of this invention can be basic in nature, they accordingly react with any of a number of inorganic and organic acids to form pharmaceutical acid addition salts.

The pharmaceutical acid addition salts of the invention are typically formed by
10 reacting the compound of formula I with an equimolar or excess amount of acid. The reactants are generally combined in a mutual solvent such as diethylether, tetrahydrofuran, methanol, ethanol, isopropanol, benzene, and the like. The salts normally precipitate out of solution within about one hour to about ten days and can be isolated by filtration or other conventional methods.

15 Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids, such as *p*-toluenesulfonic acid, methanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid and the like. Examples of such pharmaceutically acceptable salts thus are the sulfate,
20 pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate,
25 dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, β -hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, 1,5-naphthalene-disulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like.

The term "base addition salt" refers to a salt of a compound of formula I prepared
30 by reaction of a compound of formula I with a mineral or organic base. For exemplification of pharmaceutical base addition salts see, *e.g.*, Berge, S.M, Bighley, L.D., and Monkhouse, D.C., *J. Pharm. Sci.*, 66:1, 1977. This invention also contemplates pharmaceutical base addition salts of compounds of formula I. The skilled artisan would appreciate that some compounds of formula I may be acidic in nature and accordingly
35 react with any of a number of inorganic and organic bases to form pharmaceutical base addition salts. Examples of pharmaceutical base addition salts are the ammonium,

-14-

lithium, potassium, sodium, calcium, magnesium, methylamino, diethylamino, ethylene diamino, cyclohexylamino, and ethanolamino salts, and the like of a compound of formula I.

5 While all of the compounds of the present invention are useful, certain of the compounds are particularly interesting and are preferred. The following listing sets out several groups of preferred compounds. It will be understood that each of the listings may be combined with other listings to create additional groups of preferred embodiments.

- 10 i. A is C₅-C₆ cycloalkyl;
 ii. A is cyclopentyl;
 iii. A is cyclohexyl;
 iv. R^b is C₁-C₆ alkyl;
 v. R^b is methyl;
 15 vi. R^b is halo;
 vii. n is 0;
 viii. n is 1;
 ix. p is 0;
 x. p is 1;
 20 xi. Y is -E-C(O)R³, -E-NR⁴R⁵, or an optionally substituted heterocycle;
 xii. Y is -E-C(O)R³;
 xiii. Y is -E-NR⁴R⁵;
 xiv. Y is an optionally substituted heterocycle;
 xv. E is a bond;
 25 xvi. When Y is -E-C(O)R³, R³ is OR¹³, or NR⁴R⁶;
 xvii. When Y is -E-C(O)R³, R³ is OR¹³, R¹³ is optionally substituted C₁-C₆ alkyl;
 xviii. When Y is -E-C(O)R³, R³ is OR¹³, R¹³ is methyl;
 xix. When Y is -E-C(O)R³, R³ is OR¹³, R¹³ is optionally substituted (C₁-C₄ alkyl)-aryl;
 30 xx. When Y is -E-C(O)R³, R³ is OR¹³ and R¹³ is benzyl;
 xxi. When Y is -E-C(O)R³, R³ is OR¹³ and R¹³ is optionally substituted aryl;
 xxii. When Y is -E-C(O)R³, R³ is OR¹³ and R¹³ is phenyl;
 xxiii. When Y is -E-C(O)R³, R³ is NR⁴R⁶ and R⁴ is hydrogen;
 xxiv. When Y is -E-C(O)R³, R³ is NR⁴R⁶, R⁴ is hydrogen, and R⁶ is C₁-C₆ alkyl,
 35 optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀

-15-

bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, or optionally substituted heterocycle;

- 5 xxv. When Y is -E-C(O)R³, R³ is NR⁴R⁶, R⁴ and R⁶, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;
- xxvi. When Y is -E-NR⁴R⁵, R⁴ is hydrogen;
- xxvii. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, and R⁵ is C(O)R⁷;
- 10 xxviii. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, and R⁵ is C(O)R⁷, R⁷ is C₁-C₆ alkoxy, optionally substituted (C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-(C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, diphenylmethyl, optionally substituted (C₁-C₄ alkyl)-CO-aryl, optionally substituted CO-aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle,
 15 (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), or (CH₂)_tC(R¹²)(R⁹)O(R¹⁷);
- xxix. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, and R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵);
- xxx. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), R¹² is hydrogen, and R⁹ is phenyl;
- 20 xxxi. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), R¹² is hydrogen, R⁹ is phenyl, and R¹⁶ is hydrogen, then R¹⁵ is hydrogen, C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, C(O)OR¹³, SO₂R⁸, or
 25 C(O)R¹⁸;
- xxxii. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), R¹² is hydrogen, R⁹ is phenyl, and R¹⁶ is hydrogen, then R¹⁵ is optionally substituted phenyl;
- 30 xxxiii. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), R¹² is hydrogen, R⁹ is phenyl, and R¹⁶ is hydrogen, then R¹⁵ is optionally substituted heterocycle
- xxxiv. When Y is -E-NR⁴R⁵, R⁴ is hydrogen, R⁵ is C(O)R⁷, R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵), R¹² is hydrogen, and R⁹ is phenyl, then R¹⁶

-16-

and R¹⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

xxxv. When Y is -E-NR⁴R⁵, R⁴ and R⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

5 xxxvi. R⁴ is hydrogen

xxxvii. R⁴ is C₁-C₆ alkyl;

xxxviii. R⁵ is hydrogen;

xxxix. R⁵ is optionally substituted heterocycle;

xl. R⁵ is optionally substituted C₆-C₁₀ bicycloalkyl;

10 xli. R⁵ is optionally substituted (C₁-C₄ alkyl)-aryl;

xlii. R⁵ is SO₂R⁸;

xliii. R⁵ is a moiety of the formula $\begin{array}{c} \text{S} \\ \parallel \\ \text{---} \text{NHR}^8 \end{array}$;

xliv. R⁶ is hydrogen;

xlv. R⁶ is C₁-C₆ alkyl;

15 xlvi. R⁶ is optionally substituted C₃-C₈ cycloalkyl;

xlvii. R⁶ is optionally substituted C₆-C₁₀ bicycloalkyl;

xlviii. R⁶ is optionally substituted (C₁-C₄ alkyl)-aryl;

xlix. R⁶ is optionally substituted aryl;

l. R⁶ is optionally substituted (C₁-C₄ alkyl)-heterocycle;

20 li. R⁶ is optionally substituted heterocycle;

lii. R^{6'} is C₁-C₆ alkyl;

liii. R^{6'} is optionally substituted C₃-C₈ cycloalkyl;

liv. R^{6'} is optionally substituted C₆-C₁₀ bicycloalkyl;

lv. R^{6'} is optionally substituted (C₁-C₄ alkyl)-aryl;

25 lvi. R^{6'} is optionally substituted aryl;

lvii. R⁷ is optionally substituted C₁-C₆ alkyl;

lviii. R⁷ is C₁-C₆ alkoxy;

lix. R⁷ is (C₁-C₄ alkoxy)-aryl;

lx. R⁷ is optionally substituted (C₃-C₈ cycloalkyl);

30 lxi. R⁷ is optionally substituted (C₁-C₄ alkyl)-(C₃-C₈ cycloalkyl);

lxii. R⁷ is optionally substituted (C₁-C₄ alkyl)-aryl;


lxiii. R⁷ is optionally substituted aryl;

-17-

- lxiv. R⁷ is diphenylmethyl;
- lxv. R⁷ is optionally substituted (C₁-C₄ alkyl)-CO-aryl;
- lxvi. R⁷ is optionally substituted CO-aryl;
- lxvii. R⁷ is optionally substituted (C₁-C₄ alkyl)-heterocycle;
- 5 lxviii. R⁷ is optionally substituted phenoxy;
- lxix. R⁷ is optionally substituted heterocycle;
- lxx. R⁷ is (CH₂)_tS(O)_rR¹;
- lxxi. R⁷ is (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵);
- lxxii. R⁷ is (CH₂)_tC(R¹²)(R⁹)O(R¹⁷);
- 10 lxxiii. R⁷ is (CH₂)_tC(R¹²)(R⁹)S(R¹⁷);
- lxxiv. R⁷ is NR⁴R^{6'};
- lxxv. R⁸ is optionally substituted C₁-C₆ alkyl;
- lxxvi. R⁸ is optionally substituted aryl;
- lxxvii. R⁸ is optionally substituted (C₁-C₄ alkyl)-aryl;
- 15 lxxviii. R⁹ is hydrogen;
- lxxix. R⁹ is optionally substituted C₁-C₆ alkyl;
- lxxx. R⁹ is optionally substituted C₃-C₈ cycloalkyl;
- lxxxii. R⁹ is optionally substituted aryl;
- lxxxii. R⁹ is optionally substituted heterocycle;
- 20 lxxxiii. R⁹ is optionally substituted (CH₂)_u-(C₁-C₄ alkoxy)-aryl;
- lxxxiv. R⁹ and R¹² combine to form a C₃-C₈ cycloalkyl;
- lxxxv. R¹⁰ is 1 to 4 substituents selected from the group consisting of hydrogen, halo, cyano, optionally substituted heterocycle, or NR¹⁹R²⁰;
- lxxxvi. R¹⁰ is monosubstituted with chloro;
- 25 lxxxvii. R¹⁰ is monosubstituted with cyano;
- lxxxviii. R¹⁰ is monosubstituted with NH₂;
- lxxxix. R¹¹ is hydrogen;
- xc. R¹² is hydrogen;
- xc. R¹² is optionally substituted C₁-C₆ alkyl;
- 30 xcii. R¹² is optionally substituted aryl;
- xciii. R¹³ is hydrogen;
- xciv. R¹³ is optionally substituted C₁-C₆ alkyl;
- xcv. R¹³ is optionally substituted (C₁-C₄ alkyl)-aryl;
- xcvi. R¹³ is optionally substituted aryl;







-18-

- xcvii. R^{13} is $\text{CO}_2\text{CH}_2\text{CO}_2\text{CH}_2\text{CH}_3$;
- xcviii. R^{15} is hydrogen;
- xcix. R^{15} is $\text{C}_1\text{-C}_6$;
- 5 c. R^{15} is optionally substituted $\text{C}_6\text{-C}_{10}$ bicycloalkyl;
- ci. R^{15} is optionally substituted aryl;
- cii. R^{15} is optionally substituted heterocycle;
- ciii. R^{15} is $\text{C}(\text{O})\text{OR}^{13}$;
- civ. R^{15} is SO_2R^8 ;
- cv. R^{15} is $\text{C}(\text{O})\text{R}^{18}$;
- 10 cvi. R^{16} is hydrogen;
- cvii. R^{16} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl;
- cviii. R^{16} and R^{15} , together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;
- cix. R^{17} is hydrogen;
- 15 cx. R^{17} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl;
- cxii. R^{17} is optionally substituted aryl;
- cxiii. R^{17} is optionally substituted heterocycle;
- cxiiii. R^{18} is hydrogen;
- cxv. R^{18} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl;
- 20 cxvi. R^{18} is optionally substituted aryl;
- cxvii. R^{18} is optionally substituted heterocycle;
- cxviii. R^{18} is $(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-NHCO}_2\text{-(C}_1\text{-C}_4 \text{ alkyl)}$;
- cxix. R^{19} is hydrogen;
- cxix. R^{19} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl;
- 25 cxx. R^{20} is hydrogen;
- cxxi. R^{20} is optionally substituted $\text{C}_1\text{-C}_6$ alkyl;
- cxxii. R^{20} is CH_2OH ;
- cxxiii. R^{20} is $\text{CO-(C}_1\text{-C}_4 \text{ alkyl)}$;



cxxiv. Het is

-19-

- cxxv. Het is ;
- cxxvi. Het is ;
- cxxvii. Het is ;
- cxxviii. Het is ;
- 5 cxxix. Het is ;
- cxxx. Het is ;
- cxxxii. R^b is chloro;
- cxxxiii. R^a is hydrogen;
- cxxxiii. R^a is *t*-butyl;
- 10 cxxxiv. The compound is a pharmaceutical salt; and
- cxxxv. The compound is the hydrochloride salt.

The following group, including the racemic trans and cis, and the isolated enantiomers, is illustrative of compounds contemplated within the scope of this invention:

-20-

- a) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-acetamide
- b) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-3-methyl-butyramide
- 5 c) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-hydroxy-3-phenyl-propionamide
- d) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-hydroxy-3-phenyl-propionamide
- e) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-
10 hydroxy-3-methyl-butyramide
- f) 1-Hydroxy-cyclopropanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide
- g) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-methyl}-methyl-carbamic acid tert-butyl ester
- 15 h) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methylamino-acetamide hydrochloride
- i) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-acetamide
- j) {1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester
- 20 k) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methyl-propionamide hydrochloride
- l) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-methyl}-carbamic acid tert-butyl ester
- 25 m) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide hydrochloride
- n) 2-Hydroxy-hexanoic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide
- o) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-
30 hydroxy-benzamide
- p) 4-{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl}-piperazine-1-carboxylic acid tert-butyl ester
- q) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-piperazin-1-yl-acetamide dihydrochloride
- 35 r) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl}-methyl-carbamic acid tert-butyl ester

-21-

- s) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methylamino-2-phenyl-acetamide hydrochloride
- t) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-2-phenyl-acetamide
- 5 u) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-(4-pyridin-2-yl-piperazin-1-yl)-acetamide
- v) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-piperidin-1-yl-acetamide
- w) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(4-10 methyl-piperazin-1-yl)-acetamide
- x) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-diethylamino-acetamide
- y) 2-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-methoxy-isonicotinamide
- 15 z) 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexanecarboxylic acid (2-hydroxy-1-phenyl-ethyl)-amide
- aa) 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexanecarboxylic acid (2-hydroxy-1-phenyl-ethyl)-amide
- bb) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-20 (methyl-phenyl-amino)-acetamide
- cc) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-(pyridin-3-yloxy)-acetamide
- dd) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-3-yloxy)-acetamide
- 25 ee) {1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-cyclohexyl}-carbamic acid tert-butyl ester.
- ff) 1-Amino-cyclohexanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide hydrochloride
- gg) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-30 morpholin-4-yl-acetamide
- hh) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(4-hydroxy-piperidin-1-yl)-acetamide
- ii) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-2-yloxy)-acetamide
- 35 jj) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-4-yloxy)-acetamide

-22-

- kk) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-4-ylsulfanyl)-acetamide
- ll) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-cyclohexyl-methyl}-carbamic acid tert-butyl ester
- 5 mm) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-cyclohexyl-acetamide hydrochloride
- nn) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-cyclohexyl-acetamide hydrochloride
- oo) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-cyclohexyl-methyl}-carbamic acid tert-butyl ester
- 10 pp) thieno[3,2-b]pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide
- qq) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2-chloro-pyridin-4-yloxy)-acetamide
- 15 rr) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(quinolin-3-yloxy)-acetamide
- ss) 2-tert-Butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide
- tt) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-2-ylsulfanyl)-acetamide
- 20 uu) (2-aminoindan-2-yl)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carboxamide
- vv) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-cyclohexylpropionamide
- 25 ww) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-(3-chlorophenyl)-propionamide
- xx) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-pyrid-1-yl-propionamide
- yy) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-fur-3-yl-propionamide
- 30 zz) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2-chlorophenyl)amino-2-(3-chlorophenyl)-propionamide
- aaa) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-diphenylamino-2-phenylpropionamide
- 35 bbb) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclobutyl]-2-hydroxy-acetamide

-23-

- ccc) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-hydroxy-3-methyl-butyramide
- ddd) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cycloheptyl]-3-hydroxy-3-phenyl-propionamide
- 5 eee) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclobutyl]-3-hydroxy-3-phenyl-propionamide
- fff) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-hydroxy-3-methyl-butyramide
- ggg) 1-Hydroxy-cyclopropanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cycloheptyl]-amide
- 10 hhh) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylcarbamoyl]-methyl}-methyl-carbamic acid tert-butyl ester
- iii) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-methylamino-acetamide hydrochloride
- 15 jjj) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-dimethylamino-acetamide
- kkk) {1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester
- lll) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-methyl-propionamide hydrochloride
- 20 mmm) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylcarbamoyl]-methyl}-carbamic acid tert-butyl ester
- nnn) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-acetamide hydrochloride
- 25 ooo) 2-Hydroxy-hexanoic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-amide
- ppp) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-hydroxy-benzamide
- qqq) 4- {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylcarbamoyl]-phenyl-methyl}-piperazine-1-carboxylic acid tert-butyl ester
- 30 rrr) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-phenyl-2-piperazin-1-yl-acetamide dihydrochloride
- sss) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylcarbamoyl]-phenyl-methyl}-methyl-carbamic acid tert-butyl ester
- 35 ttt) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-methylamino-2-phenyl-acetamide hydrochloride

-24-

- uuu) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-2-phenyl-acetamide
- vvv) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-phenyl-2-(4-pyridin-2-yl-piperazin-1-yl)-acetamide
- 5 www) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-piperidin-1-yl-acetamide
- xxx) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(4-methyl-piperazin-1-yl)-acetamide
- yyy) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
10 2-diethylamino-acetamide
- zzz) 2-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentylmethyl]-6-methoxy-isonicotinamide
- aaaa) 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexanecarboxylic acid (2-hydroxy-1-phenyl-ethyl)-amide
- 15 bbbb) 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexanecarboxylic acid (2-hydroxy-1-phenyl-ethyl)-amide
- cccc) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(methyl-phenyl-amino)-acetamide
- dddd) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
20 2-phenyl-2-(pyridin-3-yloxy)-acetamide
- eeee) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(pyridin-3-yloxy)-acetamide
- ffff) {1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentylcarbonyl]-cyclopentyl}-carbamic acid tert-butyl ester.
- 25 gggg) 1-Amino-cyclohexanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-
isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-amide hydrochloride
- hhhh) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-morpholin-4-yl-acetamide
- iiii)N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-(4-
30 hydroxy-piperidin-1-yl)-acetamide
- jjjj)N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-
(pyridin-2-yloxy)-acetamide
- kkkk) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(pyridin-4-yloxy)-acetamide
- 35 ll)N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-2-
(pyridin-4-ylsulfanyl)-acetamide

-25-


- mmmm) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentylcarbamoyl]-cyclohexyl-methyl}-carbamic acid tert-butyl ester
- nnnn) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentyl]-2-cyclohexyl-acetamide hydrochloride
- 5 oooo) 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexyl]-2-cyclopentyl-acetamide hydrochloride
- pppp) {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentylcarbamoyl]-cyclohexyl-methyl}-carbamic acid tert-butyl ester
- qqqq) thieno[3,2-b]pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-
10 isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-amide
- rrrr) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(2-chloro-pyridin-4-yloxy)-acetamide
- ssss) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(quinolin-3-yloxy)-acetamide
- 15 tttt) 2-tert-Butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentyl]-acetamide
- uuuu) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-(pyridin-2-ylsulfanyl)-acetamide
- vvvv) (2-aminoindan-2-yl)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-
20 5-yl)]cyclopentyl]carboxamide
- wwww) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclopentyl]-2-dimethylamino-2-cyclohexylpropionamide
- xxxx) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-dimethylamino-2-(3-chlorophenyl)-propionamide
- 25 yyyy) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-dimethylamino-2-pyrid-1-yl-propionamide
- zzzz) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-dimethylamino-2-fur-3-yl-propionamide
- aaaa) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
30 2-(2-chlorophenyl)amino-2-(3-chlorophenyl)-propionamide
- bbbb) N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-
2-diphenylamino-2-phenylpropionamide

-26-

The compounds of the present invention can be prepared by a variety of procedures, some of which are illustrated in the Schemes below. The particular order of steps required to produce the compounds of formula I is dependent upon the particular compound being synthesized, the starting compound, and the relative lability of the substituted moieties.

Compounds of formula I(a), wherein R^b , the substituent attached to the non-fused carbon, is C_1 - C_6 alkyl, optionally substituted aryl, optionally substituted heterocycle, CH_2OH , or CH_2O -heterocycle may be prepared from compounds of formula II(a) as illustrated in Scheme 1 below where Y, A, R^{10} , het, n, and p are as described *supra*.

Scheme 1



Compounds of formula I(a) may be prepared by dissolving or suspending a compound of formula II(a) in a suitable solvent, preferably dimethylformamide, and adding a suitable base, including potassium methoxide, potassium *tert*-butoxide, potassium carbonate, sodium hexamethyldisilazane, and preferably potassium hexamethyldisilazane. The base is typically employed in a one to one ratio. However, as the skilled artisan would appreciate, a slight molar excess, usually in about a 1.1 to about a 3 fold molar excess relative to the compound of formula II(a), is acceptable.

The reactants are typically combined at a temperature from about 0 °C to about 100 °C. When het is isoxazole, oxazole, or imidazole, the reactants are preferably combined at room temperature and the resulting solution is typically mixed for from about 5 minutes to about 18 hours, preferably from about 15 minutes to about 3 hours. When het is pyrazole the reactants are preferably combined at room temperature and the

-27-

resulting solution is typically heated to about 100 °C for about 30 minutes to about 18 hours.

It is preferred, when A is cyclopentyl and the base employed is potassium hexamethyldisilazane, to combine the reactants from about 60°C to about 100°C for about 3 to about 6 minutes.


Any protecting groups remaining in the cyclized compound of formula I may be removed as taught in Greene to provide additional compounds of formula I. Preferred choices of protecting groups and methods for their removal may be found in the Preparations and Examples sections below.

10

Compounds of formula I(b), wherein het is substituted with halo may be prepared from compounds of formula II(b) as illustrated in Scheme 2 below where Y, A, R¹⁰, het, n, and p are as described *supra*.

15

Scheme 2



20

Compounds of formula I(b) may be prepared by dissolving or suspending a compound of formula II(b) in a suitable solvent and adding a suitable base, in an inert atmosphere, preferably under N₂. Typically a preferred and convenient solvent is dimethylformamide. A preferred base is sodium trimethylsilylanolate. The base is typically employed in a slight molar excess, usually in about a 1.05 fold molar excess relative to the compound of formula II(b). The reactants are typically combined dropwise at room temperature over a period of time from about 2 hours to about 4 hours.

25

The skilled artisan would appreciate that if other bases are used in the reaction of Scheme 2, the substituent of the het functionality may change. For example if sodium

-28-


methylthiolate is used as the preferred base, the compound of formula II(b) will be converted into the compound of formula I wherein the het functionality is substituted with -SCH₃.

5 Additionally, the compound of formula I(b) can be prepared according to Scheme 1 wherein the reactants are combined at about 0°C and mixed at about -10°C for approximately three hours. The solution is then warmed to room temperature and mixed for an additional 2 to 3 hours.

10 Certain compounds of formula I(c) are useful MRP1 inhibitors and are also useful intermediates for the preparation of other compounds of formula I. As is shown in Scheme 3, when R^b is CH₂O-Pg, a derivative of formula I(c) wherein Pg is a protecting group, the compound of formula I(c) may be further reacted by methods known in the art to produce compounds of formula I(d) where R^{b'} is CH₂OH, CH₂N₃, CH₂SR¹, or CH₂NR⁴R⁶, and Y, A, het, p, n, R¹, R⁴, R⁶, and R¹⁰ are as described *supra*.

15

Scheme 3



20 Compounds of formula I(d) wherein het is substituted with CH₂OH may be prepared by dissolving or suspending a compound of formula I(c) in a suitable solvent and adding a suitable acid. Typically a preferred and convenient solvent is methanol/dichloromethane (2:1). A preferred acid is *p*-toluenesulfonic acid hydrate. The acid is typically employed in a slight molar excess, usually in about a 1.05 fold molar excess relative to the compound of formula I(c). The reactants are typically combined at
25 room temperature and mixed from about 1 hour to about 3 hours.


-29-

The skilled artisan would appreciate that the alcohol can be further converted to compounds of formula I(d) where het is substituted with CH_2N_3 , $\text{CH}_2\text{SR}^{13}$, or $\text{CH}_2\text{NR}^4\text{R}^6$ by methods well known in the art. For general examples of these procedures, see the Preparations and Example section.

5

Compounds of formula I(e) when het is substituted with chloro are useful MRP1 inhibitors and are also useful intermediates for the preparation of other compounds of formula I. As is shown in Scheme 4, the compound of formula I(e) may be further reacted with a nucleophile by methods known in the art to produce compounds of formula I(q) where het is substituted with an amino acid ester, OR^1 , SR^{13} , $\text{S}(\text{CH}_2)_k\text{-phenyl}$, NR^4R^6 , or an optionally substituted heterocycle attached via a heteroatom, and Y, A, p, n, k, het, R^1 , R^4 , R^6 , R^{10} , and R^{13} are as described *supra*.

15



20

Compounds of formula I(q) may be prepared by dissolving or suspending a compound of formula I(e) in a suitable solvent and adding an appropriate nucleophile, in an inert atmosphere, preferably under N_2 . Typically a preferred and convenient solvent is dimethylformamide. The nucleophile is typically employed in a molar excess, usually in about a 2 to about a 4 fold molar excess relative to the compound of formula I(e).


25

The reactants are preferably combined at room temperature and the resulting solution is typically mixed for about 30 minutes to about 3 hours, until the reaction is complete as measured by the consumption of the substrate. The skilled artisan would appreciate that the reaction, depending on the nucleophile used, may require more time to react and may, also require heating. In these instances, it is preferred to mix the reactants

-30-

for approximately 15 to approximately 20 hours, then heat the solution from about 50°C to about 80°C and mix for an additional 3 hours or until the reaction is complete as measured by the consumption of the compound of formula I(e).

- 5 Compounds of formula I(p) where Y is C(R^{11'})(R¹¹)NR⁴R⁵ may be prepared from compounds of formula I(n) as illustrated in Scheme 5 below where R³ and R¹¹ are equivalent, R^{11'} is H, and het, n, p, A, R⁴, R⁵, and R¹⁰ are as described *supra*.




The compounds of formula I(n) may be reductively aminated to form the compounds of formula I(p). Reductive aminations are well known transformations, see, *e.g.*, Larock, "Comprehensive Organic Transformations", pg. 421, VCH Publishers, New York, N.Y., 1989, hereafter referred to as "Larock".

- 15 Amines of formula III may be dissolved or suspended in a suitable solvent, preferably methanol, optionally in the presence of a suitable base, preferably N-methyl morpholine or triethylamine. When the compound of formula III is an acid addition salt, it is preferred to convert the salt to its free amine form. A compound of formula I(n) is then added to the mixture. Optionally, a Lewis acid catalyst, such as titanium(IV)
- 20 isopropoxide, may be employed. Once the compound of formula I(n) has been substantially consumed, the intermediate is typically reacted *in situ* with a suitable reducing agent to provide the compounds of formula I(p). The overall conversion may be performed at about 0°C to the boiling point of the mixture, but room temperature is a preferred reaction temperature. The formation of the compounds of formula I(p) may
- 25 take from 15 minutes to 24 hours as measure by the consumption of the substrate.

-31-

Suitable reducing agents include, but are not limited to, hydrogen over palladium or platinum on carbon, borane or complexes of borane, *e.g.*, borane-pyridine, borane-*t*-butylamine, and borane-dimethylamine complex; and borohydride reducing agents such as sodium borohydride or sodium cyanoborohydride. Sodium cyanoborohydride is a preferred reducing agent.

Compounds of formula I(r) may be prepared from compounds of formula I(n) as illustrated below where p , n , R^3 , R^7 , and R^{10} are as described *supra*.



The compounds of formula I(n) may be converted to compounds of formula I(r). Such reactions are well known transformations, see, *e.g.*, Larock, "Comprehensive Organic Transformations", pg. 421, VCH Publishers, New York, N.Y., 1989, hereafter referred to as "Larock".

The compound of formula I(n) may be reduced to the corresponding alcohol, activated (*i.e.* mesylation), reacted with an appropriate azide, reduced, and then acylated. Additionally, the Curtius rearrangement may be employed to give the compound of formula I(r). See the preparation/Example section for specifics.

20 The Curtius rearrangement is performed by converting the activated compound, in an appropriate solvent, with an appropriate azide, then an appropriate alcohol, to provide a compound of formula I(r). As the skilled artisan would appreciate, the activated compound, dissolved in an appropriate solvent, is first treated with an appropriate azide and optionally a catalyst to provide the intermediate. The intermediate is treated with an appropriate alcohol to obtain the compound of formula I(r). Once the reaction is complete, as measured by the consumption of the substrate, the resulting compound of formula I(r) may be isolated by standard extractions and filtrations. If desired, the

25

-32-

resulting compound of formula I(r) may be further purified by chromatography or crystallization as appropriate.

Appropriate solvents must be capable of dissolving a sufficient amount of the activated compound and the azide for the reaction to proceed. Useful organic solvents include hexamethylphosphoramide, dimethylformamide, and preferably toluene.

The skilled artisan would appreciate that the Curtius rearrangement may be performed via a number of azides and that reaction conditions may vary depending upon the azide used. For example if sodium azide, potassium azide, and the like are used the compound must first be converted to the activated acid with an appropriate activating agent, such as ethyl chloroformate or sulfuric acid. The substrate may need to be pretreated with the activating agent, such as the case with ethyl chloroformate, or may need to be added simultaneously. The skilled artisan would appreciate the potential for reaction at an ester site of the substrate, if the molecule is treated with the azide first as is the case in these circumstances.

Preferably, diphenylphosphoryl azide is used in the process of the present invention without an activating agent.

Appropriate alcohols are lower alkyl alcohols such as methanol, ethanol, propanol, isopropanol, butanol, benzyl, t-butanol, TMS-ethanol, and the like.

The reaction may be carried out over a large range of concentrations, from about 0.001 molar to about 2.0 molar of the azide, dependent upon the solubility of the particular azide in the chosen solvent. The reaction may also be performed on slurries of the azide so long as a sufficient amount of the azide is soluble in the solvent for the reaction to proceed. Preferably the process is performed at a concentration from about 0.1 molar to about 1.0 molar. A concentration of about 0.3 to about 0.4 molar is most preferred.

Reactions may be performed between about 80°C and about 130°C, preferably between about 100°C and about 120°C. Most preferably the reactants are combined at temperature of about 20°C to about 30°C, then heated to about 80°C and about 120°C, the azide is then added, and the reactants are stirred for about 0.5 to about 1.5 hours at reflux. An appropriate alcohol is then added and heated to about 70°C to about 90°C for about 3


-33-

to about 24 hours, preferably from about 75°C to about 85°C for about 8 to about 12 hours.

Compounds of formula I where Y is C(R¹¹)(R¹¹)NR⁴R⁵ or OR², R² is (CH₂)₂NR⁴R⁵, and R⁵ is C₁-C₆ alkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, or COR⁷, may be prepared from compounds of formula I(f) and I(h) as illustrated in Scheme 6 below where X is halide and het, n, p, R¹¹, R⁴, R¹⁰, and R¹² are as described *supra*.

10

Scheme 6



The compounds of formulas I(f) and I(h) may be reductively alkylated to form the corresponding compounds of formulas I(g) and I(i), respectively. Reductive alkylation of primary amines are well known transformations, see, *e.g.*, Larock, pg. 434-435.

15

Once it is determined that the compound of formula IV has been substantially consumed, the intermediate is typically reacted *in situ* with a suitable reducing agent to provide the compounds of formula I(g) and I(i), respectively. The overall conversion may be performed at about 0°C to the boiling point of the mixture but room temperature is a preferred reaction temperature. The formation of the compounds of formulas I(g) and I(i) may take from 15 minutes to 24 hours as measured by the consumption of the substrate.


20

-34-

A base is typically employed when the compound of formula I(f) or I(h) is an acid addition salt in order to convert the salt to its free amine form. Preferred bases for this purpose are N-methylmorpholine and triethylamine. A preferred Lewis acid catalyst is titanium(IV) isopropoxide. Suitable reducing agents include, but are not limited to, borane or complexes of borane, *e.g.*, borane-pyridine, borane-*t*-butylamine, and borane-dimethylamine complex; and lithium aluminum hydride.

Compounds of formulas I(f) and I(h) may be converted to other compounds of the invention by methods well known in the chemical arts. Additional compounds of formula I may be prepared as follows, where Y is $-E-NR^4R^5$ or $-E-OR^2$, R^2 is $(CH_2)_2NR^4R^5$, R^5 is COR^7 , and R^7 is NR^4R^6' or R^5 is a moiety of the formula $\begin{matrix} \text{S} \\ \parallel \\ \text{---} \end{matrix} NHR^8$; R^{30} is R^6' or R^8 ; and het, n, p, A, R^4 , R^6' , R^8 , R^{10} , and R^{11} are as described *supra*.

Scheme 7




The primary amines of formulas I(h) and I(f) may be reacted, by methods well known in the art, with the isocyanates or isothiocyanates of formula VII to prepare the corresponding ureas and thioureas of formulas I(k) and I(o), see, generally, March, pages 802-803.

-35-

Compounds of formula I where Y is $C(R^{11})(R^{11})NR^4R^5$, and R^5 is SO_2R^8 can be prepared as described in Scheme 8, wherein het, n, p, A, R^{11} , R^4 , R^8 , and R^{10} are as described *supra*.

5

Scheme 8




Compounds of formula I(f) may be converted to other compounds of the invention via standard combinatorial synthetic techniques. For example, a compound of formula I(f) dissolved or suspended in a suitable solvent, optionally in the presence of a base, may be treated with a compound of formula VI to provide a compound of formula I(j) where R^5 is SO_2R^8 . Typically a preferred and convenient solvent is dichloromethane. When a base is employed, triethylamine is typically preferred. Furthermore, when a base is employed, the base and compound of formula VI are typically employed in a slight stoichiometric excess. For example a 1.01 to 1.40 molar excess, relative to the compound of formula I(f), is generally employed. About 1.15 to about 1.25 fold molar excess is typically preferred. When a base is not employed, the compound of formula VI is typically employed in a relatively larger stoichiometric excess. For example, about a 1.5 to about a 3 molar excess, relative to the compound of formula I(f), is usually employed. About 1.8 to about 2.2 fold molar excess is typically preferred. The reaction is usually performed at a temperature range of about 0°C to about the reflux temperature of the solvent for from 10 minutes to 18 hours. Preferably, the reaction is performed at about 15°C to about 40°C for from 5 minutes to about 1 hour.

Compounds of formula I where Y is $C(O)R^3$ and R^3 is OR^{13} or NR^4R^6 may be prepared from compounds of formula I(l) as illustrated in Scheme 9 below wherein R^{31} is NR^4R^6 or OR^{13} ; and het, n, p, A, R^4 , R^6 , and R^{13} are as described *supra*.

25

-36-

Scheme 9



Compounds of formula I(l), prepared as described in Scheme 1, may also be converted to other compounds of the invention via solution or solid phase synthetic techniques. For example, acids of formula I(l) may be treated with activating agents to form the activated carboxylic acid derivatives of formula I(l) by methods well known in the chemical arts. See, *e.g.*, The Peptides, Peptide Synthesis and the Examples and Preparations sections below.

Generally, preparation of compounds of formula I(m) where R³¹ is NR⁴R⁶ is performed in a manner similar to the reaction of compounds of formula I(f) or I(h) described in Scheme 6. Specifically, such compounds of formula I(m) may be prepared by dissolving or suspending a compound of the activated carboxylic acid derivatives of formula I(l) in a suitable solvent, optionally in the presence of a suitable base, and adding an amine of formula III. Typically a preferred and convenient solvent is dichloromethane. Preferred bases are triethylamine and piperidinylmethylpolystyrene resin. The amine is typically employed in a molar excess. For example, about a 1.5 to about a 3 molar excess, relative to the compound of the activated carboxylic acid derivatives of formula I(l) is usually employed. About 1.8 to about 2.2 fold molar excess is typically preferred. The reaction is usually performed in a temperature range of about

-37-


0°C to about the reflux temperature of the solvent for from 10 minutes to 18 hours. Preferably, the reaction is performed at about 15°C to about 40°C for from 5 minutes to about 2.5 hours.

The compounds of formula I(m) where R³¹ is OR¹³ may be prepared by methods well known in the chemical arts. For instruction on the conversion of activated carboxylic acid derivatives to esters see, *e.g.*, Larock at 978-979. Alternatively, these compounds of formula I(m) may be prepared directly from the acids of formula I(l) as taught in the Larock reference at pages 966-972.

The starting materials and compounds of the present invention may be obtained by a number of routes. For example, compounds of formula II may be prepared according to the routes shown in Schemes 8 and 9.

Where het, n, p, Y, A, and R¹⁰ are as described *supra*, compounds of formula II may be prepared according to Scheme 10.

15



Compounds of formula XI may be converted to the corresponding acid halide by methods well known to one skilled in the art. Compounds of formula II may be prepared by dissolving or suspending an acid halide of a compound of formula XI in a suitable solvent and adding a compound of formula XII in a suitable solvent. Triethylamine or dimethylformamide is a convenient solvent and is typically preferred for the compound of formula XI. A 1:1 mixture of DMF and dichloromethane is a convenient solvent and is

20

-38-

typically preferred for the amine of formula XII. This amide forming reaction is also preferably run in the presence of 4-dimethylaminopyridine (DMAP).

For compounds in which het is pyrazole, the addition of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI) to the reaction is preferred. The compound of formula XI is preferably the corresponding carboxylic acid and is employed in an equimolar amount, relative to the compound of formula XII, but a slight excess (about a 0.05 to about 0.15 molar excess) is acceptable. DMAP is employed in a catalytic fashion. For example, about 5 molar percent to about 15 molar percent, relative to the compound of formula XII, is typically employed. A 10 molar percent is usually preferred.


Compounds of formula XII where Y is $-E-C(O)R^3$, $-E-NR^4Pg$, or $-E-O(CH_2)_2NR^4-Pg$, which are used to prepare compounds of formula I(l), I(n), I(f), and I(h) are well known in the art and to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

15

Compounds of formula XII(a) where Y is $-E-NR^4R^5$ and p, n, A, E, R^4 , and R^5 are as described supra can be prepared as illustrated in Scheme 11 from compounds of formula (i) wherein Y is $-E-C(O)R^3$ and R^3 is hydrogen.

Scheme 11

20




Compounds of formula XII(a) may be prepared from compounds of formula (i) in a manner similar to that as taught in the Larock reference at pages 421-430. For an example of this transformation, see Preparation 124. Compounds of formula (i) are well known in the art and to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

25

-39-

Compounds of formula XII(c), XII(d), and XII(e) where Y is $-E-S(O)_qR^3$, and R^3 , p, A, and E are as described *supra* can be prepared as illustrated in Scheme 12.

Scheme 12



5

Compounds of formula XII(c) may be prepared from compounds of formula (iv) by methods well known to the skilled artisan. One manner is to combine the compound of formula (iv) with sodium hydride in an appropriate solvent, preferably dimethylformamide, and combining with the appropriate thiol. The compound of formula XII(c) may be further oxidized to form the compounds of formulas XII(d) and XII(e) in a manner similar to that as taught in Preparation 175 and Examples 368, and 467-468. Compounds of formula (iii) can be converted to compounds of formula (iv) in a manner similar to that of Preparation 169. Compounds of formula (iii) are well known in the art

10

-40-


and to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

Furthermore, the transformations described in Schemes 3 - 8 may be performed before the cyclization described in Scheme 1 and 2 to provide the compounds of formula XI with a fully elaborated R substituent. Additionally, the skilled artisan would appreciate that the transformations of Schemes 11-13 may be performed after the cyclization described in Schemes 1 and 2.

Additionally, compounds of formula XII may be prepared according to the following routes where, unless otherwise provided, R^b is the substituent from the carbon atom of het, R^a is the substituent from the saturated nitrogen of het, the second heteroatom is an oxygen or sulfur, and other variables are as described *supra*.

Compounds of formula XI(a), where het is [4,3-c]isoxazole, may be prepared in a manner similar to that described in the literature, for example, see Chen YP, et. al, *Heterocycles*, 1995, 41, 175, and Chantegrel B, et. al, *J. Org. Chem.*, 1984, 49, 4419-4424.

Route 1



-41-

Compounds of formula XVI may be prepared by dissolving or suspending a compound of formula XV and a suitable base in a suitable solvent and adding a compound of formula XIV in a suitable solvent, dropwise. Toluene is a convenient solvent and is typically preferred. Triethylamine is the preferred base. The compound of formula XIV is typically and preferably employed in an equimolar amount, relative to the compound of formula XV, but a slight excess is acceptable.

The reactants are preferably combined at about 0°C and the resulting solution is typically warmed to room temperature and mixed for from about 18 hours to about 24 hours.


The compound of formula XVI may then be converted to the compound of formula XIII by dissolving or suspending a compound of formula XVI in a suitable acidic solvent and adding hydroxylamine hydrochloride. Glacial acetic acid is a convenient acidic solvent and is typically preferred. The ester group is then hydrolyzed to the corresponding carboxylic acid of formula XI(a) through standard procedures commonly employed in the art, see for example, Larock, pgs 981-985.

The reactants are preferably combined at about room temperature then heated to reflux for from about 30 minutes to about 60 minutes. Preferably the reaction is heated to reflux from about 40 to 45 minutes.

Compounds of formula XIV and XV are known in the art and, to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

-42-

Compounds of formula XI(b) and (c), wherein het is [4,5-c]isoxazole, may be prepared according to route 2.

Route 2

5 Generically, the compound of formula XVIII and hydroxylamine hydrochloride are suspended or dissolved in a suitable solvent and a suitable base is added. After the reaction is complete, the solution is acidified with a suitable acid and the resulting oxime is purified by known methods. Typically a preferred and convenient solvent is water/methanol. Typically a preferred and convenient base is sodium hydroxide.

10 The reactants are preferably combined at about 0°C and the resulting solution is typically mixed for about 1 hour at about 25-30°C, until the reaction is complete. After the reaction is complete, the solution is acidified with a suitable acid, preferably hydrochloric acid, and the resulting oxime is purified by known methods.

-43-

The purified oxime is dissolved or suspended in a suitable solvent, preferably DMF, and is then reacted with *N*-chlorosuccinimide (NCS). Preferably, NCS is added in small portions to control the expected exotherm. The initial NCS addition results in a slight temperature decrease. If the reaction does not self-initiate within about 10 minutes, as indicated by a slight temperature rise, hydrogen chloride may be bubbled into the DMF solution. If the reaction does not begin within about 10 to 15 minutes, heating the solution to about 45-60°C is desirable. Once the reaction begins, the temperature is preferably maintained below about 35°C for benzaldoximes with electron-donating substituents and below about 50°C for strong electron-withdrawing substituents.

10 Completion of the reaction is indicated by cessation of the exotherm.

The compound of formula XIX is then converted to the compound of formula XX by methods well known to the skilled artisan. The compound of formula XIX and an appropriate methyl-2-butynoate are dissolved or suspended in a suitable solvent, preferably ethyl ether, and Et₃N is added. The reactants are combined at about room temperature and mixed from about 12 to 24 hours, until the reaction is complete.

The ester group of the compounds of formula XX is then hydrolyzed to the corresponding carboxylic acid of formula XI(b) through standard procedures commonly employed in the art, see for example, Larock, pgs 981-985.

20 Compounds of formula XIX may be prepared in a manner similar to that described in the literature, for example, see Liu K, Shelton BR, Howe, RK, *J. Org. Chem.*, 1980, 45, 3916-3918.

Compounds of formula XXI may be prepared in a manner similar to that described in the literature, for example, see Stevens RL, Albizati KF, *Tetrahedron Lett.* 1984, 25, 4587. For an example of this transformation, see Preparation 3.

25 Compounds of formula XI(c) may be prepared from compounds of formula XXI in a manner similar to that described in the literature, for example, see Micetich RG, Chin CG, *Can. J. Chem.* 1970, 48, 1371. For an example of this transformation, see Preparation 4.


30 Compounds of formula XVIII are known in the art and, to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

-44-

The isoxazole compounds of formula XI may be converted to the isothiazole by methods well known in the art, for example see McGregor DN, Corbin U, Swigor JE, and Cheney LC, "Synthesis of isothiazoles: The transformation of isoxazoles into isothiazoles," *Tetrahedron*, 1969, 25, 389-395.

5 Compounds of formula XI(d) where het is pyrazolyl may be prepared as illustrated below.

Route 3



10 Compounds of formula XXIII may be prepared by combining the compound of formula XXII with ethyl trifluoroacetyl vinyl ether in a manner similar to that described in the literature, for example, see Kamitori et al, *J. Het. Chem.*, 1993, 30, 389. For examples of this transformation, see Preparations 7-8.


15 Additionally, compounds of formula XXIII may be prepared from aldehyde hydrazones and ethyl propiolate as is further described by Kamitori et al, *Heterocycles*, 1994, 38, 21.

20 The trifluoromethyl ketone of formula XXIII may be converted to the corresponding carboxylic acid of formula XI(d) in a manner similar to that described in the literature, for example, see Delgado A, Clardy J, *Tetrahedron Lett.* 1992, 33, 2789-2790.

Compounds of formula XXII are known in the art and, to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

-45-

Compounds of the invention where het is oxazolyl or imidazolyl may be prepared as illustrated below.

Route 4

-46-

Compounds of formula XI(e) may be prepared in a manner similar to that described in the literature, for example, see Nunami et al, *J. Org. Chem.* 1994, 59, 7635-7642.


5 The compound of formula XXV is converted to the compound of formula XXVIII by methods well known to the skilled artisan. The compound of formula XXV is dissolved or suspended in a suitable solvent, preferably tetrahydrofuran, and a base is added, preferably Et₃N. The reactants are combined at about room temperature, preferably under an inert atmosphere, and mixed from about 30 minutes to 2 hours, until the reaction is complete.

10 The ester group of the compounds of formula XXVIII is then hydrolyzed to the corresponding carboxylic acid formula XI(f) through standard procedures commonly employed in the art, see for example, Larock, pgs 981-985.

15 Compounds of formula XIV and XXIV are known in the art and, to the extent not commercially available, are readily synthesized by standard procedures commonly employed in the art.

-47-

Compounds of the invention where het is imidazolyl may, additionally, be prepared as illustrated below:

Route 5

The compound of formula XI(e)' may be prepared by methods well known in the art. For specific conditions, see the preparation section (preparations 400-405).

The pharmaceutical salts of the invention are typically formed by reacting a
 10 compound of formula I with an equimolar or excess amount of acid or base. The reactants are generally combined in a mutual solvent such as diethylether, tetrahydrofuran, methanol, ethanol, isopropanol, benzene, and the like for acid addition salts, or water, an alcohol or a chlorinated solvent such as dichloromethane for base addition salts. The salts normally precipitate out of solution within about one hour to
 15 about ten days and can be isolated by filtration or other conventional methods.

Acids commonly employed to form pharmaceutical acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as *p*-toluenesulfonic, methanesulfonic acid, ethanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic acid,
 20 carbonic acid, succinic acid, citric acid, tartaric acid, benzoic acid, acetic acid, and the

-48-

like. Preferred pharmaceutical acid addition salts are those formed with mineral acids such as hydrochloric acid, hydrobromic acid, and sulfuric acid, and those formed with organic acids such as maleic acid, tartaric acid, and methanesulfonic acid.

5 Bases commonly employed to form pharmaceutical base addition salts are inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt
10 forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is not of a critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

15 The optimal time for performing the reactions of Schemes 1 – 8 and Routes 1-5 can be determined by monitoring the progress of the reaction via conventional chromatographic techniques. Furthermore, it is preferred to conduct the reactions of the invention under an inert atmosphere, such as, for example, argon, or, particularly, nitrogen. Choice of solvent is generally not critical so long as the solvent employed is
20 inert to the ongoing reaction and sufficiently solubilizes the reactants to effect the desired reaction. The compounds are preferably isolated and purified before their use in subsequent reactions. Some compounds may crystallize out of the reaction solution during their formation and then collected by filtration, or the reaction solvent may be removed by extraction, evaporation, or decantation. The intermediates and final products
25 of formula I may be further purified, if desired by common techniques such as recrystallization or chromatography over solid supports such as silica gel or alumina.

The skilled artisan will appreciate that not all substituents are compatible with all reaction conditions. These compounds may be protected or modified at a convenient point in the synthesis by methods well known in the art.

-49-

Preparation 1

2-((1R,3R)-3-Hydroxycyclohexyl)malonic acid dibenzyl ester

To a -78 °C solution of 2-(1(R)-3-oxo-cyclohexyl)malonic acid dibenzyl ester (see
5 Arai, T.; Yamada, Y. M. A.; Yamamoto, N.; Sasai, H.; Shibasaki, M. *Chem. Eur. J.* 1996,
2, 1368-1372.) (3.80 g, 10.0 mmol) in THF (50 mL) under N_2 was added L-selectride
(11.0 mL at 1.0 M, 11.0 mmol) dropwise and the mixture was stirred for 2 h at -78 °C.
EtOAc (100 mL) and H_2O (100 mL) were added and the mixture was warmed to room
temperature. Dilution with more EtOAc followed by washing with 1N NaOH, saturated
10 NH_4Cl , brine, drying ($MgSO_4$), and column chromatography (silica gel, hexanes:EtOAc
gradient) gave title compound (2.92 g, 76 %). Mass spectrum (ES $^-$)(m/z) 381.2 [M-1].

Preparation 2

(1R,3R)-(3-Hydroxycyclohexyl)acetic acid benzyl ester

15

A solution of a compound from preparation 1 (2.30 g, 6.02 mmol), LiCl (0.511 g,
12.0 mmol), H_2O (0.217 mL, 12.0 mmol), and DMSO (20 mL) was lowered into a 165
°C oil bath for 2 h. The reaction was cooled to room temperature and diluted with
EtOAc. Washing with H_2O and brine, drying ($MgSO_4$), and column chromatography
20 (silica gel, hexanes: EtOAc gradient) gave title compound (1.12 g, 75 %).
 1H NMR ($CDCl_3$, 400 MHz) δ 7.34-7.25 (m, 5H), 5.06 (s, 2H), 4.02 (m, 1H), 2.21 (m,
2H), 1.71-1.59 (m 4H), 1.50-1.40 (m, 4H), 1.30 (m, 1H), 1.00 (m, 1H) ppm.

Preparation 3

25

(1R,3S)-(3-Azidocyclohexyl)acetic acid benzyl ester

To a solution of a compound from preparation 2 (1.05 g, 4.24 mmol),
triphenylphosphine (1.33 g, 5.07 mmol), and hydrazoic acid (6.4 mL, at 1.0 M, 6.4 mmol)
in toluene (7 mL) was added DEAD (1.0 mL, 6.4 mmol) dropwise and stirred for 20 min.
30 The mixture was diluted with EtOAc, washed with 0.1N NaOH, H_2O , and brine, dried
($MgSO_4$), and chromatographed (silica gel, hexanes: EtOAc gradient) to give the title
compound (0.916 g, 80%).

-50-

Preparation 4

2-((1R,3S)-3-Azidocyclohexyl)-N-(3,4,5-trimethoxyphenyl)acetamide

5 A solution of a compound from preparation 3 (215 mg, 0.788 mmol) and NaOH (0.32 mL at 5M, 1.6 mmol) in MeOH (5 mL) was heated at 50-55 °C for 3 h then cooled to room temperature. The reaction was diluted with H₂O and acidified with aqueous HCl to pH 2. Extraction with EtOAc and subsequent washing (brine) and drying (Na₂SO₄) gave the crude carboxylic acid. A mixture of the crude acid, 3,4,5-trimethoxyaniline (288
10 mg, 1.58 mmol), EDCI (262 mg, 1.58 mmol), DMAP (20 mg, 0.16 mmol) in CH₂Cl₂ (5 mL) was stirred for 17 h. The mixture was diluted with CH₂Cl₂ and extracted with 1N HCl and brine then dried (MgSO₄). Column chromatography (silica gel, hexanes: EtOAc gradient) gave the title compound (255 mg, 93%). Mass spectrum (ES+) (m/z) 349.2 [M+1]. MS (ES+): [M+1] 349.2 m/z.

15

Preparation 5

2-((3S,1R)-3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)-N-3,4,5-trimethoxyphenyl)acetamide

20 A mixture of a compound from preparation 4 (250 mg, 0.72 mmol), 10% palladium on carbon (76 mg, 0.07 mmol), and EtOH (5 mL) was stirred under an atmosphere of hydrogen (balloon). After 2 h the reaction was filtered through a pad of celite with the aid of MeOH and concentrated. Benzene was added to the resulting residue followed by concentration. The residue was dissolved in CH₂Cl₂ (5 mL) followed
25 by addition of 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (236 mg, 0.87 mmol) and DMAP (175 mg, 1.44 mmol) and stirred for 16 h. The mixture was applied to a silica gel column and elution (hexanes: EtOAc gradient) gave the title compound (256 mg, 64%). Mass spectrum (ES+) (m/z) 560.2 [M+1].

-51-

Preparation 6

N-(3,4,5-Trimethoxyphenyl)-2-(3-oxocyclopentyl)acetamide

To a solution of 1.0g (7 mmol) of 3-oxocyclopentyl acetic acid in dichloromethane
5 (18 mL) and DMF (1 mL), was added 0.68 mL (7.7 mmol) oxalyl chloride. After 30 min,
the solvent was removed *in vacuo*. The residue was dissolved in dichloromethane (20
mL), cooled to 0-5 °C, and 1.29 g of 3,4,5-trimethoxyaniline (0.63 mmol) was added
followed by 0.68 mL of pyridine (8.4 mmol). After 15 min, the reaction was allowed to
warm to ambient temperature. After 2 h, the reaction was rinsed with 1N HCl, followed
10 by aq. sodium bicarbonate, and water (3X). The organic layer was dried, the solvent
removed *in vacuo* and the residue chromatographed on a silica gel with 2% methanol in
methylene chloride to yield 0.25 g of the title compound. MS(ES+)m/z=308.

Preparation 7

15 N-(3,4,5-Trimethoxyphenyl)-2-(3-hydroxycyclopentyl)acetamide

A solution of 1.07 g (3.5 mmol) of a compound from preparation 6 and 0.136 g of
sodium borohydride (3.5 mmol) in methanol (60 mL) was stirred 12 h. The solvent was
removed *in vacuo*, the residue dissolved in ethyl acetate and rinsed with 1N HCl followed
20 by water. The organic layer was dried and concentrated *in vacuo*. The residue was
chromatographed on silica gel and eluted with 5% methanol in dichloromethane to yield
0.45 g of the title compound. MS(ES+)m/z=310. Cis/trans racemic

Preparation 8

25 N-(3,4,5-Trimethoxyphenyl)-2-(3-methylsulfonyloxycyclopentyl)acetamide

A solution of 0.45 g (1.46 mmol) of a compound from preparation 7 and 0.135 mL
(1.75 mmol) of methanesulfonyl chloride and 0.27 mL (1.9 mmol) of triethylamine in
dichloromethane (15 mL) was stirred 2 h. The solvent was removed *in vacuo* and replaced
30 with ethyl acetate. The organic layer was rinsed with 1N HCl followed by aq. sodium
bicarbonate, then brine and dried to yield 0.115 g of the title compound.
MS(ES+)m/z=387.9.

-52-

Preparation 9

N-(3,4,5-Trimethoxyphenyl)-2-(3-azidocyclopentyl)acetamide

5 A solution of 0.6 g (1.55 mmol) of a compound from preparation 8, and 0.21 g (4.3 mmol) of lithium azide in DMF (12 mL) was stirred 12 h. Water and ice were added to the reaction, and the mixture was extracted with ethyl acetate. The organic layer was washed 5 times with water, dried, concentrated *in vacuo*, and the residue was chromatographed on silica gel and eluted with hexane/ethyl acetate 3:2 to yield 0.13 g of
10 *cis* and 0.21 g of *trans* of the title compound. MS(ES+) m/z=335.

Preparation 10*cis*-N-(3,4,5-Trimethoxyphenyl)-2-(3-aminocyclopentyl)acetamide

15 To a solution of 0.13 g (0.39 mmol) of a *cis* compound from preparation 9 in ethyl acetate (30 mL) was added 0.026 g of 10% palladium on carbon and the mixture was hydrogenated at 50 psi over 1.5 h. The catalyst was removed by filtration over celite and the solvent was removed *in vacuo* to yield 0.128 g of the title compound. MS(ES+) m/z=309. Cis (racemic)

20

Preparation 11*trans*-N-(3,4,5-trimethoxyphenyl)-2-(3-aminocyclopentyl)acetamide

25 In the same manner as the preparation 10, 0.21 g of a *tans* compound from preparation 9 was converted to 0.20 g of the title compound. MS(ES+)m/z=309. Trans (racemic)

-53-

Preparation 12

2-(3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclopentyl)-
N-(3,4,5-trimethoxyphenyl)acetamide

5 A solution of 0.123 g (0.42 mmol) of a compound from preparation 10, 0.03 g
(0.46 mmol) of 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride, and
0.07 mL (0.5 mmol) of triethylamine in dichloromethane (10 mL) was stirred for 12 h.
The solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was
washed with 1N HCl, then aq. NaHCO₃, followed by brine and dried. The solvent was
10 removed *in vacuo* to yield 0.255 g of the title compound. MS(ES+) m/z=545.8. *cis*
(racemic)

Preparation 13

15 2-(3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclopentyl)-
N-(3,4,5-trimethoxyphenyl)acetamide

In the same manner as preparation 12, 0.2 g (0.65 mmol) of a compound from
preparation 11 was converted to 0.324 g of the title compound. MS(ES+)m/z=545.8.
Racemic

20

Preparation 14

N-((1S,3R)-3-Cyanomethylcyclopentyl)(*t*-butoxy)carboxamide

To a solution of 8.2 g (0.0222 mol) of N-[*t*-butoxycarbonyl]-2-(*p*-
25 toluenesulfonyloxymethyl)cyclopentylamine in DMF (250 mL) was added 3.3 g (0.067
mmol) of NaCN and the suspension was heated at 80 °C for 4 h. The solvent was
removed *in vacuo* at 40 °C and the residue was slurried in brine. The aqueous layer was
extracted with ethyl acetate, dried and the solvent was removed *in vacuo* to yield 5.7 g of
the title compound. MS(ES+)m/z=225. (one enantiomer)

30

-54-

Preparation 152-{(3S,1R)-3-[(*t*-Butoxy)carbonylamino]cyclopentyl}acetic acid

5 A suspension of 2.0 g (8.9 mmol) of a compound from preparation 14 in 3N KOH (32 mL) and EtOH (12 mL) was heated at reflux for 6 h (solution). The alcohol was removed *in vacuo* and the reaction was cooled and neutralized with 3N HCl. The product was separated by filtration, rinsed with water and dried *in vacuo* at 40 °C to yield 1.33 g of the title compound.

MS(ES+)m/z=244.

10

Preparation 162-{(3S,1R)-3-[(*t*-Butoxy)carbonylamino]cyclopentyl-N-(3,4,5-trimethoxyphenyl)acetamide

15 To a solution of 0.1 g (0.4 mmol) of a compound from preparation 15 in THF/DMF 1:1 (5 mL) was added 0.061 g (0.44 mmol) of 1-hydroxy-7-azabenzotriazole and 0.086 g (0.44 mmol) of EDCI. After 40 min, 0.083 g (0.44 mmol) of 3,4,5-trimethoxyaniline was added. After 3.25 h, the solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1N HCl, aq. NaHCO₃, then
20 3X with water. The organic layer was dried and the solvent was removed *in vacuo* to yield 0.144 g of the title compound. MS(ES+)m/z=409.

Preparation 17

2-((3S,1R)-3-Aminocyclopentyl)-N-(3,4,5-trimethoxyphenyl)acetamide

25

A solution of 0.48 g (0.362 mmol) of a compound from preparation 16 in TFA (10 mL) was stirred for 1 h, after which the TFA was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1N NaOH followed by brine and dried. The organic layer was removed *in vacuo* to yield 0.3 g of the title compound.

30 MS(ES+)m/z=309.

-55-

Preparation 18

2-((3S,1R)-3-[[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino]cyclopentyl)-N-(3,4,5-trimethoxyphenyl)acetamide

5 In the same manner as preparation 12, 0.32 g (1.04 mmol) of a compound from preparation 17 yielded 0.52 g of the title compound. MS(ES+)m/z=546.

Preparation 19

N-(*t*-Butoxycarbonyl)-3-aza-2-oxabicyclo[3.2.1]nonane

10

A solution of 0.59 g (4.8 mmols) of 2-oxo-3-azabicyclo-[3.2.1]nonane, 2.07 g (9.6 mmol) of (BOC)₂O and 1.173 g (9.6 mmol) of DMAP in CH₂Cl₂ (15 mL) was stirred 12 h. The solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1N HCl followed by brine, dried and the solvent removed *in vacuo* to yield 0.495 g of the title compound. MS(ES+)m/z=225.9

15

Preparation 20

cis-3-[[(*t*-Butoxy)carbonylamino]methyl]cyclopentanecarboxylic acid

20 To a solution of 2.22 g (0.98 mmol) of a compound from preparation 19 in THF (113 mL) was added H₂O (32 mL) and the solution cooled to 5 °C. Water (30%, 11.95 mL, 9.8 mmol) was added followed by 2.08 g (4.9 mmol) of LiOH and the reaction was allowed to warm to ambient temperature. After 2 h, the reaction was quenched with sodium sulfite solution (10%) and the THF was removed *in vacuo*. With cooling, the reaction was acidified with 6N HCl and extracted with ethyl acetate, dried and the solvent removed *in vacuo* to yield 1.583 g of the title compound. MS(ES+)m/z=243.9.

25

Preparation 21

cis-N-(3-[[(*t*-Butoxy)carbonylamino]methyl]cyclopentyl)(phenylmethoxy)carboxamide

30

To warm solution of 0.244 g (1.0 mmol) of a compound from preparation 20 in toluene (8 mL) was added 0.2 mL (1.4 mmol) of Et₃N followed by 0.3 mL (1.4 mmol) of

-56-

diphenylphosphoryl azide. The reaction was stirred at ambient temperature for 2.5 h and refluxed for 1.5 h. The solvent was removed *in vacuo* and replaced with ethyl acetate which was rinsed with aq. NaHCO₃, aq. KHSO₄ and brine. The solvent was removed *in vacuo* and the residue was chromatographed on silica gel with 0.5-1% MeOH/CH₂Cl₂ to yield 0.264 g of the title compound. MS(ES+)m/z=349.

Preparation 22

cis-N-[(3-Aminocyclopentyl)methyl](*t*-butoxy)carboxamide

10 A suspension of 0.264 g (0.76 mmol) of a compound from preparation 21 and 0.034 g of Pd/C 10% in MeOH (40 mL) was reduced at 1.0 atm. of H₂ over 2.5 h. The catalyst was removed by filtration and the solvent was removed *in vacuo* to yield 0.187 g of the title compound. MS(ES+)m/z=215.

Preparation 23

cis-N-[(3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclopentyl)methyl](*t*-butoxy)carboxamide

15 To a suspension of 0.187 g (0.873 mmol) of a compound from preparation 22 in dichloromethane (10 mL) was added 0.263 g (0.96 mmol) of 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride followed by 0.2 mL (1.4 mmol) of Et₃N. After 70 min, the solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1N HCl followed by aq. NaHCO₃, dried and the solvent was removed *in vacuo* to give a residue which was chromatographed on silica gel with hexane/ethyl acetate 3-1 to yield 0.156 g of the title compound. MS(ES+)m/z=452.

Preparation 24

cis-N-[[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclopentyl]methyl](*t*-butoxy)carboxamide

30

To a solution of 0.136 g (0.3 mmol) of a compound from preparation 23 in DMF (7.0 mL) at 0-5 °C was added 0.6 mL (0.3 mmol) of KHMDS (0.5N in THF). After 2 min,

-57-

cooling was removed. After 3 additional min, 1N HCl was added and the mixture was extracted with ethyl acetate. The organic layer was rinsed with water, dried and concentrated to a foam *in vacuo* which was chromatographed on silica gel with hexane/ethyl acetate 3:1 to yield 0.057 g of the title compound. MS(ES+)m/z=432.

5

Preparation 25

2-(3-Aminocyclopentyl)ethanenitrile

In the same manner as preparation 17, 0.3 g (4.5 mmol) of a compound from preparation 14 was deprotected with TFA to yield 0.087 g of the title compound. MS(ES+)m/z=125.

10

Preparation 26

N-[(1R,3S)-3-(cyanomethyl)cyclopentyl][3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

15

In the same manner as preparation 18, 0.08 g (0.65 mmol) of a compound from preparation 26 was acylated with 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride to yield a residue which after chromatography on silica gel with hexane/ethyl acetate 1:1 yielded 0.177 g of the title compound. MS(ES+)m/z=362.

20

Preparation 27

2-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclopentyl]ethanenitrile

25

In the same manner as preparation 24, 0.166 g (0.46 mmol) of a compound from preparation 27 was cyclized to a residue which upon chromatography on silica gel with hexane/ethyl acetate 2:1 yielded 0.099 g of the title compound. MS(ES+)m/z=342.

-58-

Preparation 28

2-[(3S,1R)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclopentyl]acetic acid

5 A suspension of 0.086 g (0.25 mmol) of a compound from preparation 27 in HCl conc. (40 mL) was refluxed for 6 h, after which the reaction was concentrated to a residue *in vacuo* at 40 °C. The residue was taken into diethyl ether, the insolubles filtered and discarded, and the filtrate was extracted with 1N NaOH. The basic layer was acidified with 1N HCl, extracted with ethyl acetate, dried and concentrated *in vacuo* to yield 0.059
10 g of the title compound. MS(ES+)m/z=361.

Preparation 29

N-[3(1S,3S)-3-(2-aminoethyl)cyclopentyl](*t*-butoxy)carboxamide

15 0.5 g (2.2 mmols) of a compound from preparation 14 was reduced with 0.065 g of PtO₂ in EtOH (50 mL) at 60 psi of H₂ at 40 °C over 12 h. The catalyst was removed by vacuum filtration, and the solvent was removed *in vacuo* to yield 0.421 g of the title compound. MS(ES+)m/z=229.

Preparation 30

20 N-[(1S,3S)-3-[2-(Phenylcarbonylamino)ethyl]cyclopentyl](*t*-butoxy)carboxamide

In a manner similar to the acylation of a compound from preparation 24, 0.172g (0.75 mmol) of a compound from preparation 29 was acylated with benzoyl chloride and
25 the residue formed was chromatographed on silica gel with hexane/ethyl acetate 2:1 to yield 0.127 g of the title compound. MS(ES+)m/z=333.

-59-

Preparation 31

N-((1S,3S)-3-[2-(Phenylcarbonylamino)ethyl]cyclopentyl)[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

5 In a manner similar to preparation 17, 0.127 g (0.36 mmol) of a compound from preparation 30 was deprotected, followed by acylation with 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride in the same manner as the preparation of a compound from preparation 12 to yield 0.15 g of the title compound. MS(ES+)m/z=470.

10

Preparation 32

N-((1S,3S)-3-{2-[(3,4,5-Trimethoxyphenyl)carbonylamino] ethyl}cyclopentyl)(*t*-butoxy)carboxamide

15 In the same manner as preparation 30, 0.1 g (0.44 mmol) of a compound from preparation 29 was acylated with 3,4,5-trimethoxybenzoyl chloride to yield 0.16 g of the title compound. MS(ES+)m/z=423.

Preparation 33

20 N-((1S,3S)-3-[2-((3,4,5-Trimethoxyphenyl)carbonylamino)ethyl]cyclopentyl)[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

In a manner similar to preparation 31, 0.16 g (0.38 mmol) of a compound from preparation 32 yielded 0.174 g of the title compound. MS(ES+)m/z=560.

25

Preparation 34

((3S,1R)-3-[[*t*-Butoxy]carbonylamino]methyl)cyclopentyl)-N-(3,4,5-trimethoxyphenyl)carboxamide

30 In a manner similar to the preparation of a compound from preparation 16, 0.224 g (0.92 mmols) of a compound from preparation 20 was converted to 0.34 g of the title compound. MS(ES+)m/z=409.

-60-

Preparation 35

[(3S,1R)-3-(aminomethyl)cyclopentyl]-N-(3,4,5-trimethoxyphenyl)carboxamide

In a manner similar to preparation 17, 0.049 g (0.12 mmol) of a compound from preparation 34 was deprotected with TFA to yield 0.013 g of the title compound.

MS(ES+)m/z=308.

Preparation 36

[(3S,1R)-3-({[3-2-chloro-6-fluorophenyl]-5-methylisoxazol-4-yl]carbonylamino)methyl)cyclopentyl]-N-(3,4,5-trimethoxyphenyl)carboxamide

In a manner similar to preparation 18, 0.013 g (0.042 mmol) of a compound from preparation 35 was converted to a residue which was chromatographed on silica gel with dichloromethane 100% to dichloromethane/MeOH 2% to yield 0.018 g of the title compound. MS(ES+)m/z=545.8.

Preparation 37

Ethyl 2-(3-({[4-(2-chloro-6-fluorophenyl)-2-methyl-3-furyl]carbonylamino}cyclohexyl)acetate

To a solution of 3-nitrophenylacetic acid, 1.0 g (5.5 mmol) in 90 mL of ethanol was added 1.0 g of PtO₂ and 2 mL of 5N HCl. The mixture was hydrogenated overnight at 60 °C and 60 psi, filtered through celite and concentrated to dryness. To a solution of the residue in 15 mL of dichloromethane was added 3.0 mL (21.6 mmol) of triethylamine and 2.0 g (7.2 mmol) of 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride. The solution was stirred overnight at ambient temperature. Chloroform was added and the solution was washed with 0.5N HCl, saturated sodium bicarbonate, brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was chromatographed on silica gel using 5% acetone/dichloromethane as eluent to yield 2.0 g (66%) of the desired mixture of isomers as a white solid. ¹H-NMR is consistent with structure. MS (ion spray) 423.2 (M+1).

-61-

Preparation 38

Ethyl 2-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]acetate

To a solution of a compound from preparation 37, 1.0 g (2.4 mmol) in 20 mL of
5 N,N-dimethylformamide was added 0.35 g (3.1 mmol) of potassium *t*-butoxide. After 75
min, the mixture was poured into 1N HCl/ice and extracted with ethyl acetate. The
combined organics were washed with brine, dried over sodium sulfate, filtered and
concentrated to dryness. The residue was chromatographed on silica gel using 5%
acetone/dichloromethane as eluent to yield 0.65 g (67%) of the desired mixture of isomers
10 as a viscous yellow oil. ¹H-NMR is consistent with structure. MS (ion spray) 403.3
(M+1).

Preparation 39

2-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]acetic acid

15

To a solution of a compound from preparation 38, 0.65 g (1.5 mmol) in 10 mL of
tetrahydrofuran and 10 mL of ethanol was added 3.0 mL (15.0 mmol) of 5N sodium
hydroxide. After 40 min at ambient temperature, the reaction was concentrated to
dryness. The residue was partitioned between ethyl acetate and water, was acidified to pH
20 3.0 with 1N HCl, and was extracted with ethyl acetate. The combined organics were
washed with brine, dried over sodium sulfate, filtered and concentrated to dryness to yield
0.53 g (93%) of the desired mixture of isomers as a white foam. ¹H-NMR is consistent
with structure. MS (ion spray) 375.2 (M+1).

25

Preparation 40

Methyl 2-(3-nitrocyclohexyl)acetate

To a solution of *m*-nitrophenylacetic acid, 14.2 g (78.0 mmol) in 200 mL of
methanol at 0 °C was added 20 mL of acetyl chloride dropwise. The reaction mixture was
30 stirred overnight at ambient temperature and concentrated to dryness. The residue was
chromatographed on silica gel using 50% ethyl acetate/hexanes as eluent to yield 14.7 g

-62-

(97%) of the desired mixture of isomers as a yellow oil. $^1\text{H-NMR}$ is consistent with structure. MS (FD) 194.8 (M-1).

Preparation 41

5 Methyl 2-(3-{[4-(2-chloro-6-fluorophenyl)-2-methyl-3-furyl]carbonylamino}cyclohexyl)acetate

To a solution of methyl 2-(3-nitrocyclohexyl)acetate (13.8 g, 71.0 mmol) in 200 mL of acetic acid was added 6.9 g of PtO_2 . The mixture was hydrogenated overnight at ambient temperature and 60 psi, then was filtered through celite and concentrated to dryness. To a solution of the residue, 8.8 g (38.0 mmol) in 200 mL of dichloromethane was added 27 mL (190 mmol) of triethylamine and 10.4 g (38.0 mmol) of 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride. The mixture was stirred overnight at ambient temperature then chloroform was added. The organics were washed with 0.5N HCl, saturated sodium bicarbonate, brine, dried over sodium sulfate, filtered and concentrated to dryness. The resulting residue was chromatographed on silica gel using 5% acetone/dichloromethane to yield 4.43 g (29%) of the desired mixture of isomers as a colorless, viscous oil. $^1\text{H-NMR}$ is consistent with structure. MS (ion spray) 409.2 (M+1).

20

Preparation 42

2-(3-{[4-(2-Chloro-6-fluorophenyl)-2-methyl-3-furyl]carbonylamino}cyclohexyl)acetic acid

25 To a solution of a compound from preparation 41, 1.2 g (3.0 mmol) in 20 mL of absolute ethanol and 20 mL of tetrahydrofuran was added 6.0 mL (30.0 mmol) of 5N sodium hydroxide. The mixture was stirred 4 h, then the organics were concentrated off. The remaining aqueous phase was acidified to pH 3.0 with 1N HCl, then extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was dissolved in chloroform and concentrated to dryness to yield 1.0 g (84%) of the desired mixture of isomers as a white foam. $^1\text{H-NMR}$ is consistent with structure. MS (ion spray) 393.2 (M-1).

30

-63-

Preparation 43

[4-(2-Chloro-6-fluorophenyl)-2-methyl(3-furyl)]-N-{3-[2-(4-methylpiperidyl)-2-oxoethyl]cyclohexyl}carboxamide

5

To a solution of a compound from preparation 42, 1.0 g (2.6 mmol) in 10 mL of N,N-dimethylformamide was added 0.34 mL (2.9 mmol) of 4-methylpiperidine, 0.43 g (3.1 mmol) of 1-hydroxy-7-azabenzotriazole and 0.6 g (3.1 mmol) of 1-(3-(dimethyl-amino)propyl)-3-ethylcarbodiimide. The mixture was stirred overnight at ambient
10 temperature and concentrated to dryness. The resulting residue was partitioned between ethyl acetate and water and was extracted with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was chromatographed on silica gel using a gradient of 30% ethyl acetate/hexanes to 5% methanol/chloroform as eluent to yield 0.9 g (70%) of the desired mixture of
15 isomers as a white foam. ¹H-NMR is consistent with structure. MS (ion spray) 476.2 (M+1).

Preparation 44

5-[(3S,1R)-3-(Aminomethyl)cyclohexyl]-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-
20 4-one

A solution of a compound from preparation 54 (enantiomer 2) (6.6 g (14.8 mmol) in 230 mL of acetic acid saturated with hydrochloric acid was stirred for 2 h at ambient temperature, then was concentrated to dryness. The resulting residue was partitioned
25 between 20% isopropanol/chloroform and saturated sodium bicarbonate and was extracted with 20% isopropanol/chloroform. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness to yield 4.68 g (92%) of the desired product as a yellow solid. ¹H-NMR is consistent with structure. MS (ion spray) 346.1 (M+1).

30

-64-

Preparation 45

Phenylmethyl-2-((3S,1R)-3-[(*t*-butoxy)carbonylamino]cyclohexyl)acetate

To a solution of phenylmethyl 2-((3S,1R)-3-azidocyclohexyl)acetate, 5.0 g (18.3
5 mmol) in 350 mL of ethyl acetate under a nitrogen atmosphere was added 8.0 g (36.6
mmol) of BOC-anhydride and 2.0 g of Lindlar's catalyst. The nitrogen was purged with
hydrogen and the reaction mixture was stirred overnight under a hydrogen balloon, then
concentrated to dryness. The resulting residue was filtered through celite and
chromatographed on silica gel using ethyl acetate/hexanes as eluent to yield 6.25 g (98%)
10 of the desired product as a clear oil which solidifies upon standing. ¹H-NMR is
consistent with structure. MS (ion spray) 347 (M⁺).

Preparation 46

Phenylmethyl 2-((3S,1R)-3-[[4-(2-chloro-6-fluorophenyl)-2-methyl(3-
15 furyl)]carbonylamino]cyclohexyl)acetate

To a solution of a compound from preparation 45, 5.05 g (20.4 mmol) in
80 mL of dichloromethane at 0 °C was added 3.7 mL (24.5 mmol) of triethylamine and 10
mg of 4-dimethylaminopyridine, followed by 6.15 g (22.4 mmol) of 3-(2-chloro-6-
20 fluorophenyl)-5-methylisoxazole-4-carbonyl chloride. The reaction mixture was stirred
overnight while warming to ambient temperature, and was then washed with 0.1N
hydrochloric acid, washed with brine, dried over sodium sulfate, filtered and concentrated
to dryness. The residue was chromatographed on silica gel using ethyl acetate/hexanes as
eluent, concentrated to dryness, dissolved in chloroform and concentrated to dryness to
25 yield 5.74 g (58%) of the desired compound as a white solid. ¹H-NMR is consistent with
structure. MS (ion spray) 485.5 (M+1).

-65-

Preparation 47

2-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]acetic acid

5 To a solution of a compound from Example 491 (3.91 g, 8.4 mmol) in 70 mL of dioxane was added 70 mL (350 mmol) of 5N sodium hydroxide. The mixture was refluxed for 4 h and was cooled to ambient temperature and acidified to pH 1 with 5N hydrochloric acid. The resulting mixture was extracted with 20% isopropanol/
10 chloroform. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was triturated in ether, filtered and dried under vacuum to yield 1.8 g (57%) of the desired product as a tan solid. ¹H-NMR is consistent with structure. MS (ion spray) 374.8 (M+).

Preparation 48

15 *cis*-N-[3-Aminomethylcyclohexyl]-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

A slurry of a compound from preparation 52, 1.2 g (2.7 mmol) in 20 mL of acetic acid saturated with hydrochloric acid was stirred for 4 h at ambient temperature, then concentrated to dryness. The residue was dissolved in toluene and concentrated to
20 dryness. The residue was slurried in ether/hexanes, concentrated to dryness, and dried under vacuum to yield 750 mg (80%) of the title compound as a tan solid. ¹H-NMR is consistent with structure. MS (ion spray) 346.2 (M+1).

Preparation 49

25 (*t*-Butoxy)-N-[(3-nitrophenyl)methyl]carboxamide

To a suspension of 5.00 g (26.5 mmol) of 3-nitrobenzylamine hydrochloride in 100 mL CH₂Cl₂ at room temperature was added 5.79 g (26.5 mmol) of di-*t*-butyl dicarbonate. To this was added 8.13 mL (58.3 mmol) of triethylamine, dropwise over 15
30 min. The reaction was stirred for 3 h at room temperature, after which it was diluted with 300 mL of ethyl acetate. The resulting organic solution was washed three times with 1N

-66-

HCl solution, dried over sodium sulfate and concentrated *in vacuo* to give 6.2 g (93%) of a white solid, which was characterized as the title compound. MS (FIA) $m/z = 253$.

Preparation 50

5 (t-Butoxy)-N-[(3-aminocyclohexyl)methyl]carboxamide

To a solution containing 12.0 g (47.7 mmol) of a compound from preparation 49 in 300 mL ethanol was added 6.0 g of rhodium on carbon. The reaction was subjected to hydrogenation (60 psi) at 60 °C for 18 h, after with the catalyst was removed by vacuum
10 filtration and the solvent removed *in vacuo*, giving 9.5 g (87%) of a clear oil. This material was characterized as the title compound and used without further purification.

Preparation 51

15 N-(3{[t-Butoxy)carbonylamino]methyl}cyclohexyl)[4-(2-chloro-6-fluorophenyl)-2-methyl(3-furyl)]carboxamide

To a solution of 9.5 g (41.6 mmol) of a compound from preparation 50 in 200 mL of CH₂Cl₂ was added 11.4 g (41.6 mmol) of 3-(2-chloro-6-fluorophenyl)-5-methyl-isoxazole-4-carbonyl chloride, followed by 11.6 mL (83.2 mmol) of triethylamine at room
20 temperature. The reaction was stirred at room temperature for 15 h, and concentrated *in vacuo*. The crude solid was dissolved in 200 mL of ethyl acetate and the organic solution was washed twice with 1N HCl solution, dried over sodium sulfate and concentrated *in vacuo* to give a yellow solid. This material was recrystallized from methanol to give 12.5
25 g (64%) of a white crystalline solid, which was characterized as the title compound, as an inseparable mixture of *cis* and *trans* isomers. MS(FIA) $m/z = 466.2$.

Preparation 52 and 53

30 (t-Butoxy)-N-{[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl))cyclohexyl]methyl}carboxamide

To a stirred solution of 7.00 g (15.0 mmol) of a compound from preparation 51 in 200 mL of DMF at room temperature was added 30.0 mL (15 mmol) of

-67-

a 0.5 M toluene solution of potassium bis(trimethylsilyl) amide over 10 min. The resulting dark red reaction was stirred an additional 5 min at room temperature and added to 200 mL of 1N HCl. The two phase solution was diluted with 400 mL of ethyl acetate and the organic layer was separated. The organic solution was washed four times with
5 brine, dried over sodium sulfate and concentrated *in vacuo* to give an orange solid. This material was recrystallized from toluene to give 3.4 g of a light yellow solid, which was characterized as pure racemic *cis* material, 52: MS(FIA) $m/z = 446.1$. Purification of the racemic *trans* material (53) was accomplished by concentrating the mother liquor and
10 subjecting this material to flash chromatography on silica gel, using 50% hexane-ethyl acetate as the eluant. The major fractions were combined to give a light yellow solid, which was characterized as pure racemic *trans* material: MS(FIA) $m/z = 446.1$.

Preparation 54

15 N-{3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl}(*t*-butoxy)carboxamide

The racemic *cis* material from a compound from preparation 52 was separated into its enantiomers by chiral HPLC chromatography using a Chiralpak AD column and 10% ethyl alcohol-heptane as the eluant at a flow of 1.0 mL/min.

20 Retention Time (Enantiomer 1) = 10.501 min.

Retention Time (Enantiomer 2) = 12.576 min.

Preparation 55

Methyl 3-[4-methoxyphenyl)methoxy]benzoate

25

To a solution containing 5.00 g (32.9 mmol) of methyl 3-hydroxybenzoate in 100 mL of DMF was added 4.77 mL (32.9 mmol) of *p*-methoxybenzyl chloride and 4.54 g (32.9 mmol) of anhydrous potassium carbonate. The reaction was stirred at room temperature for 16 h, followed by 1 h at 100 °C. The solution was diluted with 200 mL of
30 ethyl acetate and 200 mL of 1N HCl. The organic phase was separated and washed five times with brine, dried over sodium sulfate and concentrated *in vacuo* to give 8.99 g of a

-68-

white amorphous solid, which was characterized as the title compound. This material was used without further purification.

Preparation 56

5 3-[(4-Methoxyphenyl)methoxy]benzoic acid

To a solution containing 2.75 g (10.0 mmol) of a compound from preparation 55 in 50 mL of MeOH was added 10 mL of 1N NaOH solution. To aid solubility, 20 mL of THF was added and the reaction was heated at reflux for 3 h. The reaction was cooled to
10 room temperature and diluted with 100 mL of ethyl acetate and 100 mL of 1N HCl solution. The organic layer was separated, dried over sodium sulfate and concentrated *in vacuo* to give 2.45 g of a white amorphous solid, which was characterized as the title compound. This material was used without further purification.

15 Preparation 57

(-)-(1R,*cis*)-Pinonic acid

A solution of 1S-(-)-pinene (11.95 mL, 75 mmol) in methanol (50 mL) and chloroform (50 mL) at -78 °C was treated with ozone for 2 h via a gas dispersion tube.
20 After purging with nitrogen for 1 h, methyl sulfide (16.5 mL, 225 mmol) was added, and the reaction was allowed to warm to room temperature overnight. The solvent was removed *in vacuo* and the residue dissolved in diethyl ether, washed with water, saturated aqueous brine solution, dried over magnesium sulfate and concentrated to approximately 11 g of yellow liquid. The liquid was dissolved in ethyl ether and treated with oxygen for
25 2 h, followed by stirring under an O₂ balloon overnight. The reaction was concentrated to give 12.42 g as a slightly yellow liquid, 90% yield. ¹H NMR: consistent with structure. MS (ion spray) 183 (M⁻ - 1).

-69-

Preparation 58

(3-Acetylamino-2,2-dimethylcyclobutyl)acetic acid

To a solution of a compound from preparation 57 (12.37 g, 67.1 mmol) in acetic acid (270 mL) was added hydroxylamine-O-sulfonic acid (11.74 g, 100.7 mmol), and the reaction was heated to reflux under nitrogen for 20 h. The reaction was concentrated and the residue dissolved in 2N HCl, followed by extraction with ethyl acetate (x4). The combined organics were washed with brine, dried over magnesium sulfate, and concentrated. The brown solution was treated with decolorizing charcoal, filtered through celite, and concentrated to 5.85 g as a light yellow oil, 44% yield. ¹H NMR: consistent with structure. MS (ion spray) 200 (M⁺ + 1), 198 (M⁺ - 1).

Preparation 59

(3-Acetylamino-2,2-dimethylcyclobutyl)acetic acid methyl ester

15

To a 0 °C flask of methanol (150 mL) was added thionyl chloride (10.7 mL, 146.8 mmol) dropwise via an addition funnel. To this solution was added a compound from preparation 58 (5.85 g, 29.4 mmol) dropwise as a solution in methanol (100 mL). The cooling bath was removed and the reaction allowed to stir at room temperature for 2 h. The reaction was concentrated to a brown oil, dissolved in ethyl acetate, washed with saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate and concentrated to a 5.5 g of a brown oil. Purification by column chromatography on silica gel (eluting with 50-100% ethyl acetate/hexane) gave 3.1 g as a brown oil, 50% yield. ¹H NMR: consistent with structure. MS (ion spray) 213 (M⁺).

25

Preparation 60

(3-Amino-2,2-dimethylcyclobutyl)acetic acid methyl ester

A compound from preparation 59 (3.09 g, 14.5 mmol) was dissolved in 2N HCl (100 mL) and heated to reflux for 18 h, then concentrated. To a 0 °C flask of methanol (100 mL) was added thionyl chloride (5.3 mL, 72.4 mmol) dropwise via an addition funnel. To this solution was added the crude material dropwise as a solution in methanol

-70-

(15 mL). The clear solution was allowed to warm to room temperature overnight. The solution was concentrated, and purification by SCX column (eluting with tetrahydrofuran/water then ammonia/methanol) gave 2.52 g as a brown solid, 100% yield. ^1H NMR: consistent with structure. MS (ion spray) 172 ($\text{M}^+ + 1$).

5

Preparation 61

(3-{[3-(2-Chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl]amino}-2,2-dimethyl-cyclobutyl)acetic acid methyl ester

10 To a solution of a compound from preparation 60 (566 mg, 3.31 mmol) in dichloromethane (20 mL) was added 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (1.09 g, 3.97 mmol), 4-dimethylaminopyridine (40 mg, 0.33 mmol), and triethylamine (1.4 mL, 9.92 mmol) dropwise via syringe. The solution was stirred under nitrogen at room temperature overnight. The solution was washed with 1.0N HCl (x3),
15 saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 20-40% ethyl acetate/hexane) gave 1.28 g as a yellowish oil, 95% yield. ^1H NMR: consistent with structure. MS (ion spray) 409 (M^+).

20

Preparation 62

[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-2,2-dimethyl-cyclobutyl]-acetic acid methyl ester

To a 0 °C solution of a compound from preparation 61 (2.44 g, 5.97 mmol) in
25 dimethylformamide (100 mL) was added potassium *t*-butoxide (0.80 g, 7.16 mmol). After 15 min, the reaction was quenched with 1.0N HCl and extracted with 20% isopropanol/ chloroform. The organic layer was then washed with 1.0N HCl (x2), saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 15-20%
30 ethyl acetate/hexane) gave 739 mg (32% yield) as a white solid. ^1H NMR: consistent with structure. MS (ion spray) 389 (M^+).

-71-

Preparation 63

[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-2,2-dimethyl-cyclobutyl]acetic acid

5 A solution of a compound from preparation 62 (737 mg, 1.89 mmol) in 5N NaOH (25 mL) was heated to reflux for 1 h. The solution was concentrated, dissolved in 20% isopropanol/chloroform, acidified to pH 2 with 5N HCl, washed with 1.0N HCl (x2), dried over magnesium sulfate and concentrated to give 519 mg as a light orange foam, 73% yield. ¹H NMR: consistent with structure.

10

Preparation 64

[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-2,2-dimethyl-cyclobutylmethyl]carbamic acid 2-trimethylsilanylethyl ester

15 To a mixture of a compound from preparation 63 (318.7 mg, 0.85 mmol) in toluene (15 mL) was added triethylamine (130 μL, 0.94 mmol), diphenylphosphorylazide (202 μL, 0.94 mmol), and the mixture heated to reflux for 3 h. To the reaction was added 2-(trimethylsilyl)-ethanol (183 μL, 1.28 mmol), and the mixture heated at reflux overnight. The solution was concentrated and dissolved in dichloromethane, washed with
20 1.0N HCl, brine, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 0-30% ethyl acetate/hexane) gave 144 mg as a white foam, 35% yield. ¹H NMR: consistent with structure. MS (ion spray) 490 (M⁺).

Preparation 65

25 N-3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl(*t*-butoxy)carboxamide

To a 0°C solution of 1,3-diaminocyclohexane (50.56 g, 442.8 mmol) in chloroform (850 mL) was added a solution of di-*t*-butyl-dicarbonate (20.3 mL, 88.6
30 mmol) in chloroform (250 mL) dropwise via an addition funnel over 25 minutes. The mixture was stirred under nitrogen at room temperature overnight. The reaction was washed with saturated aqueous sodium bicarbonate solution (3x), brine, dried over

-72-

magnesium sulfate, and concentrated to give 17.58 g of a white oil. To a solution of the crude material (17.58 g, 82.0 mmol) in dichloromethane (500 mL) was added 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (23.65 g, 86.1 mmol), dimethylaminopyridine (1.00 g, 8.2 mmol), and triethylamine (34.3 mL, 246.1 mmol) dropwise via syringe. The solution was stirred under nitrogen at room temperature overnight. The solution was washed with 0.1N HCl (2x), saturated aqueous sodium bicarbonate solution (2x), brine, dried over magnesium sulfate and concentrated. The orange solids were treated with ethyl ether, sonicated for 15 minutes, and filtered. The resulting white solids were crushed, treated with ethyl ether, sonicated for 15 minutes, filtered out, and dried on a vacuum overnight to give 16.20 g of the title racemic cis compound as a white solid, 87% yield based on a 1:1 mixture of cis:trans. This material was then separated by chiral LC on a Chiralcel OD column to yield 9 g of isomer 1: (3S,1R) and 7 g of isomer 2: (1S,3R). ¹H NMR: consistent with structure. MS (ion spray) 352 (M⁺ - BOC + 1).

15

Preparation 66

N-(1S,3R)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl](*t*-butoxy)carboxamide

To a solution of a compound from preparation 65 (isomer 1) (500 mg, 1.11 mmol) in dimethylformamide (20 mL) was added KHMDS (3.3 mL, 1.66 mmol, 0.5M in toluene) rapidly via syringe. After 15 minutes, ethyl acetate (50 mL) was added to the reaction, and the solution added to a separatory funnel containing water (50 mL) and brine (30 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (x2). The combined organics were washed with brine, dried over magnesium sulfate, and concentrated to give 709 mg red solid. Purification using flash chromatography on silica gel (eluting with 20-30% ethyl acetate/hexane) gave 205 mg of the title compound as a yellow solid, 43% yield. ¹H NMR: consistent with structure. MS (ion spray) 332 (M⁺ - BOC).

30

-73-

Preparation 67

N-(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl](*t*-butoxy)carboxamide

5 To a solution of a compound from preparation 65 (isomer 2) (500 mg, 1.11 mmol) in dimethylformamide (20 mL) was added KHMDS (3.3 mL, 1.66 mmol, 0.5M in toluene) rapidly via syringe. After 4 minutes, ethyl acetate (50 mL) was added to the reaction, and the solution added to a separatory funnel containing water (50 mL) and brine (30 mL). The layers were separated, and the aqueous layer was extracted with ethyl
10 acetate (2x). The combined organics were washed with brine, dried over magnesium sulfate, and concentrated to give 700 mg red solid. Purification using flash chromatography on silica gel (eluting with 0-0.5% methanol/chloroform) gave 372 mg of the title compound as a yellow solid, 78% yield. ¹H NMR: consistent with structure. MS (ion spray) 332 (M⁺ - BOC).

15

Preparation 68

5-((3S,1R)-3-Aminocyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one hydrochloride

20 To a compound from preparation 66 (200 mg, 0.46 mmol) was added acetic acid saturated with HCl(g) (10 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 1 h. The reaction was concentrated, followed by addition of acetonitrile and concentrated to assist in the removal of acetic acid. The resulting white solid was treated with ethyl ether, sonicated, and filtered to yield 159 mg of the title compound as a
25 white solid, 93% yield. ¹H NMR: consistent with structure. MS (ion spray) 369 (M⁺).

Preparation 69

5-((1S,3R)-3-aminocyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one hydrochloride

30

To a compound from preparation 67 (368 mg, 0.85 mmol) was added acetic acid saturated with HCl(g) (15 mL, ~3N in HCl) and the solution stirred vigorously at room

-74-

temperature for 1 h. The reaction was concentrated, followed by addition of acetonitrile and concentrated to assist in the removal of acetic acid. The resulting white solid was treated with ethyl ether, sonicated, and filtered to yield 249 mg of the title compound as a white solid, 79% yield. $^1\text{H NMR}$: consistent with structure. MS (ion spray) 369 (M^+).

5

Preparation 70

Ethyl 4-aminohexanecarboxylate

Thionyl chloride (3.0 mL, 0.041 mol) was added dropwise to ethanol (125 mL) at room temperature and the mixture stirred for an additional 10 min. Then, 4-amino-1-cyclohexane-carboxylic acid (5.0 g, 0.035 mol) was added and the mixture stirred overnight at ambient temperature. The mixture was then concentrated *in vacuo* and the resulting solid dried overnight *in vacuo* which resulted in the isolation of 6.84 g (94%) of the desired ester hydrochloride. MS(ES): ($\text{M}+1$) $^+$ 172.2 m/z.

15

Preparation 71

Ethyl 4-{{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexane carboxylate

A compound from preparation 70 (0.414 g, 0.002 mol) was combined with 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.548 g, 0.002 mol) and triethylamine (0.70 mL, 0.005 mol) in CH_2Cl_2 (6.0 mL) and the mixture stirred for 5 h at ambient temperature. The mixture was diluted with CH_2Cl_2 , washed with 2x water, and dried over sodium sulfate. The mixture was concentrated *in vacuo*. The resulting residue chromatographed over silica gel using $\text{CH}_2\text{Cl}_2/\text{MeOH}$ as eluant which allowed for isolation of 0.790 g (97%) of the desired amide. MS(ES): ($\text{M}+1$) $^+$ 409.3, 410.3, 411.2 m/z.

25

-75-

Preparation 72

Ethyl 4-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane
carboxylate

5 A compound from preparation 71 (0.70 g, 0.0017 mol) was dissolved in DMF (10
mL) and potassium-*t*-butoxide (0.230 g, 0.0020 mol) was added and the mixture stirred
for 2 h at ambient temperature. Water (100 mL) and aq 1N HCl (5 mL) was added and
the aqueous mixture extracted with ethyl acetate. The combined extracts were dried over
sodium sulfate and then concentrated *in vacuo*. The resulting residue was
10 chromatographed over silica gel using CH₂Cl₂/MeOH as eluant which allowed for
isolation of 0.540 g (81%) of product as a light yellow solid. MS(ES): (M+1)⁺ 389.2,
391.2 m/z.

Preparation 73

15 3-[(*t*-Butoxy)carbonylamino]cyclohexanecarboxylic acid

To 3-aminocyclohexane carboxylic acid (3.00 g, 0.021 mol) in a THF/H₂O
mixture (50 mL/50 mL) was added potassium carbonate (2.90 g, 0.021 mol) and di-*t*-butyl
dicarbonate (4.58 g, 0.021 mol) and the resultant mixture stirred overnight at ambient
20 temperature. Then aqueous 5N HCl (10 mL) was added and the mixture extracted with
ethyl acetate. The combined extracts were dried over sodium sulfate and concentrated *in*
vacuo which allowed for isolation of 4.92 g (96%) of desired product as a white solid.
MS(ES): (M+1)⁺ 244.5 m/z.

25 Preparation 74

(*t*-Butoxy)-N-[3-(hydroxymethyl)cyclohexyl]carboxamide

A compound from preparation 73 (4.00 g, 0.0165 mol) was dissolved in dry THF
(50 mL) and the mixture cooled in an ice bath under a nitrogen atmosphere. Then borane-
30 tetrahydrofuran (1.0 M, 20 mL, 0.02 mol) was added *via* syringe and the mixture stirred
overnight while warming to room temperature. The reaction mixture was quenched into
ice water and solid NaCl was added. This mixture was extracted with ethyl acetate and

-76-

the combined extracts were dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was chromatographed over silica gel using a CH₂Cl₂/THF mixture as eluant which allowed for isolation of 2.40 g (63%) of the desired alcohol as a viscous oil. MS(ES): (M+1)⁺ 230.2 m/z.

5

Preparation 75

(*t*-Butoxy)-N-[3-(*p*-toluenesulfonyloxymethyl)cyclohexyl] carboxamide

A compound from preparation 74 (0.15 g, 0.00066 mol) was combined with *p*-toluenesulfonyl chloride (0.125 g, 0.00066 mol), triethylamine (0.18 mL, 0.0013 mol), and DMAP (0.005 g, cat.) in dichloromethane (5.0 mL) and the resultant mixture stirred overnight at room temperature. The mixture was then concentrated *in vacuo* and the residue chromatographed over silica gel using a mixture of CH₂Cl₂/THF as eluant which allowed for the isolation of the desired product (0.195 g, 77%) as a thick oil that solidified upon standing. MS(ES): (M+1)⁺ 384.2 m/z.

15

Preparation 76

(*t*-Butoxy)-N-[3-(phenylthiomethyl)cyclohexyl]carboxamide

Thiophenol (0.27 mL, 0.0026 mol) was dissolved in DMF (10 mL) at ambient temperature under a nitrogen atmosphere and sodium hydride (60%, 0.115 g, 0.0029 mol) was added. After stirring for 15 min., a compound from preparation 75 was added and the mixture stirred overnight at ambient temperature. The mixture was concentrated *in vacuo* and the residue chromatographed over silica gel with CH₂Cl₂/THF as eluant which allowed for isolation of 0.74 g (88%) of desired product as a white solid. MS(ES): (M+1)⁺ 322.2 m/z.

20

25

Preparation 77

3-(Phenylthiomethyl)cyclohexylamine

30

A compound from preparation 76 (0.50 g, 0.0016 mol) was dissolved in CH₂Cl₂ (10 mL) and TFA (1.0 mL) was added. After 1.5 h, deprotection was incomplete and

-77-

additional TFA (1.0 mL) was added and the mixture stirred at ambient temperature for an additional 4 h. The mixture was concentrated *in vacuo* and the residue treated with aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were concentrated *in vacuo* and the residue eluted over a short plug of silica gel (CH₂Cl₂/MeOH) which
5 allowed for isolation of the desired amine (0.265 g, 77%). MS(ES): (M+1)⁺ 222.2 m/z.

Preparation 78

[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-[3-(phenylthiomethyl)cyclohexyl]carboxamide

10

A compound from preparation 77 (0.26 g, 0.0012 mol) was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (0.322 g, 0.0012 mol) and triethylamine (0.33 mL, 0.0024 mol) in CH₂Cl₂ (10 mL) and the mixture stirred overnight at ambient temperature. The reaction mixture was chromatographed over silica
15 gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.495 g (91%) of the desired amide as a white solid. MS(ES): (M+1)⁺ 459.4 m/z.

Preparation 79

(*t*-Butoxy)-N-[3-(phenylmethylthiomethyl)cyclohexyl] carboxamide

20

Benzyl mercaptan (0.31 mL, 0.0026 mol) was dissolved in DMF (10 mL) at ambient temperature under a nitrogen atmosphere and sodium hydride (60%, 0.115 g, 0.0029 mol) was added. After stirring for 15 min., the compound from preparation 75 (1.00 g, 0.0026 mol) was added and the mixture stirred overnight at ambient temperature.
25 The mixture was concentrated *in vacuo* and the residue chromatographed over silica gel with CH₂Cl₂/THF as eluant which allowed for isolation of 0.80 g (92%) of desired product as a thick oil which solidified upon standing. MS(ES): (M+1)⁺ 336.2 m/z.

-78-

Preparation 80

[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-[3-(phenylmethylthiomethyl)cyclohexyl]carboxamide

5 The compound from preparation 73 (0.40 g, 0.0012 mol) was dissolved in CH₂Cl₂ (8.0 mL) and TFA (2.0 mL) and the mixture stirred at ambient temperature for 3 h. The mixture was concentrated *in vacuo* and the residue treated with aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and concentrated *in vacuo* which allowed for the recovery of 0.19 g (68%) of crude amine.

10 This amine (0.19 g, 0.0008 mol) was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl chloride (0.22 g, 0.0008 mol) and triethylamine (0.23 mL, 0.0016 mol) in CH₂Cl₂ (10 mL) and the mixture stirred overnight at ambient temperature. Water (~30 mL) was added and the mixture extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and then concentrated *in vacuo*. The

15 resulting residue was chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.29 g (77%) of the desired amide as a white solid. MS(ES): (M+1)⁺ 473.1, 475.1 m/z.

Preparation 81

20 (*t*-Butoxy)-N-[3-{(benzylsulfonyl)methyl}cyclohexyl] carboxamide

 A compound from preparation 79 (0.467 g, 0.0014 mol) was dissolved in a methanol/acetone (10 mL)/(10 mL) mixture and NaHCO₃ (0.467 g, 0.0056 mol) and water (10 mL) were added. Oxone® (1.71 g, 0.0028 mol) was then added and the mixture

25 stirred at ambient temperature for 2.5 h. The mixture was concentrated *in vacuo* and the residue taken up in water and extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.420 g (82%) of the desired sulfone as a white solid. MS(FD): M⁺ 367.2 m/z.

30

-79-

Preparation 82

[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-(3-{benzylsulfonyl}methyl)cyclohexyl)carboxamide

5 The compound from preparation 81 (0.40 g, 0.0011 mol) and TFA (2.0 mL) were dissolved in CH₂Cl₂ (10.0 mL) and the mixture stirred at ambient temperature for 2 h. The mixture was concentrated *in vacuo* and the residue treated with aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and concentrated *in vacuo* which allowed for recovery of 0.21 g (71%) of crude amine. This
10 amine (0.21 g, 0.0008 mol) was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (0.22 g, 0.0008 mol), triethylamine (0.22 mL, 0.0016 mol) and DMAP (0.005 g, cat.) in CH₂Cl₂ (10 mL) and the mixture stirred overnight at ambient temperature. The mixture was concentrated *in vacuo* and the resulting residue chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation
15 of 0.35 g (89%) of the desired amide. MS(ES): (M+1)⁺ 505.1 m/z.

Preparation 83

{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}methyl benzoate

20 The compound from preparation 74 (1.00 g, 0.0044 mol) was combined with benzoyl chloride (0.56 mL, 0.0048 mol), triethylamine (1.70 mL, 0.012 mol), and DMAP (0.020 g, cat.) in CH₂Cl₂ (10 mL) and the mixture stirred overnight at ambient temperature. The reaction mixture was concentrated *in vacuo* and the residue chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation
25 of 1.05 g (72%) of the desired ester as an oil. MS(ES): (M+1)⁺ 334.1 m/z.

Preparation 84

(3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)methyl benzoate

30

The compound from preparation 83 (1.00 g, 0.003 mol) and TFA (2.0 mL) were dissolved in CH₂Cl₂ (10 mL) and the mixture treated in a manner similar to preparation

-80-

82, which allowed for isolation of 0.69 g (98%) of crude amine. This amine (0.69 g, 0.0029 mol) was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (0.81 g, 0.0029 mol) and triethylamine (0.82 mL, 0.0059 mol) in CH₂Cl₂ (10 mL) and again treated in a manner similar to preparation 81. Purification
5 resulted in the recovery of 1.30 g (95%) of desired amide as a white foam. MS(ES): (M+1)⁺ 471.1 m/z.

Preparation 85

9-Chloro-5-[3-(hydroxymethyl)cyclohexyl]-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one
10

The compound from Example 489 (0.33 g, 0.00073 mol) was added to aqueous sodium hydroxide (1N, 1.0 mL), ethanol (2.0 mL) and THF (2.0 mL) and the mixture stirred at ambient temperature for 3 h. The mixture was diluted with water and extracted with ethyl acetate. The combined extracts were washed with 2 X sat'd aq NaHCO₃ and
15 dried over sodium sulfate. Concentration *in vacuo* netted 0.225 g (88%) of desired alcohol as a tan foam. MS(ES): 347.2, 349.3 m/z.

Preparation 86

2-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid
20

A compound from preparation 45 (1.0 g; 2.77 mmol) was dissolved in tetrahydrofuran (4 mL) and ethanol (4 mL) under a dry nitrogen atmosphere at room temperature. This cloudy white solution became clear and colorless after mixing with 2N NaOH_(aq) (15 mL; 19.4 mmol; 11.1 equiv) for 2 h. After rotary evaporation to dryness,
25 the white solid was dissolved in water (20 mL) and the resulting solution extracted with diethyl ether (twice). Acidification of the aqueous layer to pH 2 with 1N HCl_(aq) produced a white solid that was extracted into ethyl acetate (thrice). The ethyl acetate was washed with saturated NaCl_(aq), dried with Na₂SO_{4(s)}, and concentrated to dryness by rotary evaporation. The resulting white solid (700 mg; 98% yield) was used in
30 subsequent reactions without further purification. MS(ES) calc'd: [M+Na]⁺ = 280.2 m/z, [M-H]⁻ = 256.2 m/z. Found: 280.1 m/z; 256.2 m/z.

-81-

Preparation 872-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}-*N*-ethoxy-*N*-methyl acetamide

A compound from preparation 86 (690 mg; 2.68 mmol) was dissolved in
5 anhydrous dichloromethane (10 mL) under a dry nitrogen atmosphere at room
temperature. Addition of 1,1'-carbonyldiimidazole (535 mg; 3.22 mmol; 1.2 equiv) to the
clear, colorless starting material solution immediately generated a gas. After stirring the
resulting solution for 30 min, triethyl amine (746 μ L; 5.36 mmol; 2 equiv) and
10 *N,O*-dimethylhydroxylamine hydrochloride (570 mg; 5.90 mmol; 2.2. equiv) were added
and the solution stirred for 15 h. Water was added to the solution and then the solution
was extracted with dichloromethane (thrice). The organic layer was washed with
saturated NaCl_(aq), dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary
evaporation. The resulting oil (850 mg) was purified by radial chromatography on a 4
mm thick silica gel rotor with a 5% tetrahydrofuran/ dichloromethane (v/v) mobile phase.
15 A clear, colorless oil (650 mg; 80%) was obtained. MS(ES) calc'd: [M+Na]⁺ = 323.2
m/z. Found: 323.2 m/z.

Preparation 88*(t*-Butoxy)-*N*-[3-(2-oxo-3-phenylpropyl)cyclohexyl]carboxamide

20

A compound from preparation 87 (4.11 g; 13.7 mmol) was dissolved in anhydrous
tetrahydrofuran (100 mL) under a dry nitrogen atmosphere. The clear, colorless solution
became clear and brown upon addition of 0.5 M benzylmagnesium chloride in
tetrahydrofuran (15 mL; 30.1 mmol; 2.2 equiv). The reaction was quenched with water
25 (100 mL) after 0.5 h and the product was extracted from the quenched solution with ethyl
acetate. The organic layer was washed with saturated NaCl_(aq) (once), dried with
Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The resulting oil
was purified by silica gel chromatography on a 6 \times 15 cm column with a 2% acetonitrile/
dichloromethane (v/v) mobile phase. A white solid (3.17 g; 70%) was obtained after
30 concentration of appropriate fractions. TOF-MS(ES) calc'd: [M+Na]⁺ = 354.2045 m/z.
Found: 354.2037 m/z.

-82-

Preparation 89

[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-[3-(2-oxo-3-phenylpropyl)cyclohexyl]carboxamide

5 A compound from preparation 88 (3.0 g; 9.05 mmol) was dissolved in excess, neat acetic acid saturated with HCl(g) (20 mL). After 30 min of stirring at room temperature, in air, the solution was concentrated to dryness by rotary evaporation. Acetic acid was removed from the resultant solid by consecutive dissolution in, and drying from, acetonitrile (thrice) and then diethyl ether (once). The resulting white powder was
10 dissolved in anhydrous dichloromethane (400 mL) under a nitrogen atmosphere. First was added 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (2.74 g; 9.99 mmol; 1.1 equiv), and then 4-*N,N*-dimethylaminopyridine (114 mg; 0.93 mmol; 0.1 equiv) and finally triethylamine (5 mL; 36.3 mmol; 4 equiv). The reaction solution was stirred 17 h and then quenched with saturated NaHCO₃(aq). The organic layer was
15 washed with saturated NaHCO₃(aq) (twice), saturated NaCl(aq) (once), dried with Na₂SO₄(s), filtered, and concentrated to dryness by rotary evaporation. The resulting oil was dissolved in dichloromethane and purified by column chromatography on a 6 × 15 cm silica gel bed in dichloromethane. Dichloromethane (1 L) was followed by a 2% tetrahydrofuran/dichloromethane (v/v) mobile phase. A white solid (3.93 g; 95%) was
20 obtained. TOF-MS(ES) calc'd: [M+H]⁺ = 469.1694 m/z. Found: 469.1692 m/z.

Preparation 90

2-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid

25 A compound from preparation 87 (845 mg; 2.81 mmol) was dissolved in anhydrous tetrahydrofuran (30 mL) under a dry nitrogen atmosphere. A 1 M solution of DIBAL-H in toluene (6.2 mL; 6.18 mmol; 2.2 equiv) was added dropwise to the -78 °C starting material solution over 5 min. After an additional 15 min of stirring, water quenched the cold reaction solution. Extraction with ethyl acetate produced an emulsion
30 that was destroyed by concentration of the solution to near dryness. The resulting material was dissolved in water/diethyl ether and extracted with diethyl ether without further complication. The organic layer was washed with saturated NaCl(aq) (once),

-83-

dried with $\text{Na}_2\text{SO}_4(\text{s})$, filtered, and concentrated to dryness by rotary evaporation. The product was purified by radial chromatography on a 4 mm thick silica gel rotor with 300 mL of 1% tetrahydrofuran/dichloromethane (v/v) mobile phase followed by 300 mL of 10% tetrahydrofuran/dichloromethane (v/v) mobile phase. A white solid (535 mg; 79%) was obtained after concentration of the desired fractions. MS(ES) calc'd: $[\text{M}+\text{H}]^+ = 264.1$ m/z, $[\text{M}-\text{Boc}+\text{H}]^+ = 142.1$ m/z. Found: 264.1 m/z; 142.1 m/z.

Preparation 91

N-{(1S,3S)-3-[2-(Phenylamino)ethyl]cyclohexyl}(t-butoxy)carboxamide

10

A nitrogen sparged round bottom flask was charged with denatured ethanol (5 mL), $\text{Ti}(i\text{-PrO})_4$ (337 μL ; 1.14 mmol; 2.2 equiv), and aniline (95 μL ; 1.04 mmol; 2 equiv). A compound from preparation 90 (125 mg; 0.518 mmol; 1 equiv) was then added to the clear, colorless reaction solution. The reaction solution became cloudy and then progressed to clear and light yellow over 3.3 h. NaBH_4 (30 mg; 0.793 mmol; 1.5 equiv) was then added and allowed to react overnight at room temperature. The clear, colorless reaction solution was quenched with 2 M $\text{NH}_3(\text{aq})$ (20 mL). After a failed attempt at Celite[®] filtration, the product and Celite[®] mixture were suspended in water and extracted with CHCl_3 (thrice). This organic layer was washed with saturated $\text{NaCl}(\text{aq})$ (once), dried with $\text{Na}_2\text{SO}_4(\text{s})$, filtered, and concentrated to dryness by rotary evaporation. The aqueous layer was decanted from the Celite[®] and the Celite[®] was suspended in 2 M $\text{NH}_3(\text{aq})$ (120 mL). This suspension was extracted with CHCl_3 (thrice) and the organic layer was treated similarly to the first organic layer. The combined extracts were purified by radial chromatography on a 2 mm thick silica gel rotor. The first chromatography with a 2% tetrahydrofuran/ dichloromethane (v/v) mobile phase did not completely purify the product and was followed by a second and third radial chromatography with a 1% tetrahydrofuran/ dichloromethane (v/v) mobile phase. A white solid (55 mg; 33%) was obtained after concentration of the desired fractions. MS(ES) calc'd: $[\text{M}+\text{H}]^+ = 319.2$ m/z; $[\text{M}+\text{C}_2\text{H}_3\text{O}_2]^- = 317.2$ m/z. Found: 319.2 m/z; 377.2 m/z.

30

-84-

Preparation 92

N-{(1S,3S)-3-[2-(Phenylamino)ethyl]cyclohexyl}[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

5 A compound from preparation 91 (53 mg; 0.166 mmol) was dissolved in excess, neat acetic acid saturated with HCl_(g) (2 mL). After 30 min of stirring at room temperature, in air, the solution was concentrated to dryness by rotary evaporation. Acetic acid was removed from the resultant solid by consecutive dissolution in, and drying from, acetonitrile (thrice) and then diethyl ether (twice). The resulting white powder was
10 dissolved in anhydrous dichloromethane (8 mL) under a nitrogen atmosphere. First was added 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (50.2 mg; 0.183 mmol; 1.1 equiv), and then 4-*N,N*-dimethylaminopyridine (2 mg; 0.017 mmol; 0.1 equiv) and finally triethylamine (94 μ L; 0.667 mmol; 4 equiv). The reaction solution was stirred overnight, then quenched with saturated NaHCO_{3(aq)}, then extracted with
15 dichloromethane (twice). The organic layer was washed with saturated NaCl_(aq) (once), dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The resulting product was dissolved in dichloromethane and purified by radial chromatography on a 2 mm thick silica gel rotor. A dichloromethane mobile phase was followed by a 5% tetrahydrofuran/dichloromethane (v/v) mobile phase. An off-white
20 solid (66 mg; 87%) was obtained. TOF-MS(ES) calc'd: [M+H]⁺ = 456.1854 m/z. Found: 456.1850 m/z.

Preparation 93

Methyl 2-(3-{[4-(2-chloro-6-fluorophenyl)-2-methyl-3-furyl]carbonylamino}cyclohexyl)-
25 acetate

Part a: To a solution containing 15.0 g (83 mmol) of *m*-nitrophenylacetic acid in 200 mL ethanol was added 7.5 g 5% rhodium on carbon. The reaction was subjected to hydrogenation (60 psi) at 60⁰ C for 10 hours, after which the catalyst was removed by
30 vacuum filtration and the solvent removed *in vacuo*, yielding 3.5 g (27%) of an oil. This material was used without further purification.

-85-

Part b: 1 mL acetyl chloride was added to 30 mL methanol. To this solution was added 3.5 g (22.3 mmol) of a compound from part a and the reaction was stirred for 1 hour. Solvents were removed *in vacuo*, yielding 3.24 g (70%) of an oil. This material was used without further purification.

5 Part c: To a solution of 1.05 g (5.06 mmol) of a compound from part b in 10 mL of methylene chloride was added 1.53 g (5.57 mmol) of 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride, followed by 2.11 mL (15.2 mmol) of triethylamine at room temperature. The reaction was stirred overnight, then concentrated *in vacuo*. The crude solid was dissolved in 150 mL ethyl acetate, washed 3X with 1N HCl, 3X with
10 NaHCO₃, 3X with brine, dried over sodium sulfate and concentrated *in vacuo* yielding 1.8 g (87%) of a white solid. This material was an inseparable mixture of *cis* and *trans* isomers. MS(FIA) m/z= 409.3

Part d: To a solution of 1.78 g (4.35 mmol) of a compound from part c in 10 mL of DMF at room temperature was added 5 mL of 2N NaOH in MeOH. The reaction was
15 stirred for 1 hour and 0.537 g (4.78 mmol) of potassium *t*-butoxide was added. The resulting dark red reaction was stirred for 6 hours. The reaction was then added to 50 mL of 1N HCl and diluted with 150 mL of ethyl acetate. The organic layer was separated and washed 3X with brine, dried over sodium sulfate and concentrated *in vacuo*. This crude material was purified by flash chromatography, eluting with 5% MeOH/CH₂Cl₂. The
20 major fractions were combined and concentrated *in vacuo* to give 1.15 g (71%) of a clear oil, which was characterized as the title compound. MS(FIA) m/z= 375.2

Preparation 94

25 2-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl] acetic acid

To a solution of 1.78 g (4.35 mmol) of a product of preparation 93, in 10 mL of DMF at room temperature was added 5 mL of 2N NaOH in MeOH. The reaction was stirred for 1 h and 0.537 g (4.78 mmol) of potassium *t*-butoxide was added. The resulting dark red reaction was stirred for 6 h. The reaction was then added to 50 mL of 1N HCl
30 and diluted with 150 mL of ethyl acetate. The organic layer was separated and washed 3X with brine, dried over sodium sulfate and concentrated *in vacuo*. This crude material was purified by flash chromatography, eluting with 5% MeOH/CH₂Cl₂. The major fractions

-86-

were combined and concentrated *in vacuo* to give 1.15 g (71%) of a clear oil, which was characterized as the title compound. MS(FIA) $m/z=375.2$.

Preparation 95

5 2-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl] acetyl
chloride

To an ice bath cooled solution containing 0.75 g (2.0 mmol) a compound from preparation 94 in 10 mL methylene chloride and catalytic DMF (3 drops) was added 0.262
10 mL (3.0 mmol) of oxalyl chloride in one portion. The ice bath was removed and the reaction stirred for 3 h. The solvents were removed *in vacuo* and the resulting oil azeotroped with toluene. This material was characterized as the title compound and used without further purification.

15

Preparation 96

Methyl 3-(azidomethyl)benzoate

To a solution of methyl 3-(bromomethyl)benzoate in DMF (55 ml) under N_2 was added sodium azide (10.39 g, 160 mmol) and stirred for 3 h. The solution was diluted
20 with EtOAc, washed (water twice, then brine), dried ($MgSO_4$), filtered, and concentrated to give the title compound. This material was used without further purification. MS (EI+) (m/z)=101.1 [M^++1]

Preparation 97

25

Methyl 3-{{(*t*-butoxy)carbonylaminomethyl}benzoate

A mixture of a compound from preparation 95 (8.75 g, 45.8 mmol), $(BOC)_2O$ (11.98 g, 55 mmol), 10% Pd/C catalyst (4.76 g), and EtOAc (175 mL) was stirred under an atmosphere of hydrogen gas (balloon) for 18 h, filtered through celite, and
30 concentrated. Column chromatography (silica gel, hexanes/EtOAc gradient) gave the title compound (5.8 g, 48%). Mass Spectrum (ES+) (m/z) 166.0 [$M-BOC$].

-87-

Preparation 98

Methyl 3-{[(*t*-butoxy)carbonylamino]methyl}cyclohexane carboxylate

5 A mixture of a compound from preparation 97 (5.57 g, 20 mmol), 5% Rh/C (2.78 g), and MeOH (140 mL) was shaken in a PARR apparatus at 60 psi and 60 °C for 18 h, filtered through celite, and concentrated. Column chromatography (silica gel, hexanes/EtOAc gradient) gave the title compound (3.5 g, 62%). Mass Spectrum (ES+) (m/z) 166.0 [M-BOC].

10

Preparation 99

Methyl 3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-5-yl]carbonylamino}cyclohexane carboxylate

15 To a solution of a compound from preparation 98 (0.5 g, 1.845 mmol) in dichloromethane (10 mL) was added TFA (7 mL) and stirred for 45 min. The solution was concentrated using benzene to azeotrope. The residue was dissolved in dichloromethane (20 mL) and Et₃N (0.77 mL, 5.54 mmol) and 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.56 g, 2.03 mmol) were added under N₂. The solution was stirred overnight. The reaction was diluted with EtOAc, 20 washed (0.1N HCl, H₂O, and brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (0.52 g, 69%). Mass Spectrum (FIA) (m/z) 409.3 [M+1].

Preparation 100

25 Methyl 3(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]-quinolin-5-yl)cyclohexane carboxylate

30 A compound from preparation 99 (0.445 g, 1.09 mmol) and *t*-BuOK (0.135 g, 1.2 mmol), in DMF (15 mL) were allowed to react for 15 min under N₂. Ice water was added and extracted twice with EtOAc. The combined EtOAc layers were washed, dried, filtered, and concentrated. Column chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (0.329 g, 78%). Mass Spectrum (FIA) (m/z) 389.2 [M+1].

-88-

Preparation 101

Methyl 3(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane
carboxylate

5

A compound from preparation 100 (5.17 g, 12.7 mmol) and *t*-BuOK (1.56 g, 13.97 mmol), in DMF (130 mL) were allowed to react for 20 min under N₂. Ice water was added and extracted twice with EtOAc. The combined EtOAc layers were washed, dried, filtered, and concentrated. Column chromatography (silica gel,
10 acetone/dichloromethane gradient) gave *cis* (0.28 g, 7%), *trans* (1.08 g, 25 %), and unseparated title compound (2.91 g, 68 %).

Mass Spectrum (FIA) (m/z) 389.2 [M+1].

Mass Spectrum (FIA) (m/z) 389.2 [M+1].

15

Preparation 102

3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane carboxylate
racemic

A *cis* compound from preparation 101 (1.08 g, 2.78 mmol), 1N NaOH (7 mL, 6.94
20 mmol), MeOH (20 mL), and THF (20 mL) were allowed to react for 5 h at 50 °C. The reaction was cooled to room temperature, diluted with water, and acidified (conc. HCl) to less than pH 3. The mixture was extracted with EtOAc (2x) and the combined extracts were washed, dried, filtered, and concentrated to give the title compound (1.03 g, 98%).

Mass Spectrum (ES+) (m/z) 375.1 [M+1].

25

Preparation 103

2-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid

To a solution of a compound from preparation 45 (5.0 g, 14.4 mmol),
30 dioxane (96 ml), and 1.0 N NaOH (28.8 ml, 28.8 mmol) was heated at 50 °C for 3 h. The reaction was cooled to room temperature, diluted with H₂O, and extracted with Et₂O. The aqueous layer was acidified (conc. HCl) to less than pH 3, and extracted with EtOAc

-89-

(2x). The combined extracts were washed (brine), dried (MgSO₄), filtered, and concentrated to give the title compound.

Preparation 104

5 N-((1S,3R)-3-{[(Phenylmethoxy)carbonylamino]methyl} cyclohexyl)(*t*-
butoxy)carboxamide

To a solution of a compound from preparation 103 (3.43 g, 13.35 mmol), Et₃N (3.75 mL, 26.96 mmol) in toluene (86 mL) under N₂ was added DPPA (5.8 mL, 26.96
10 mmol) and benzyl alcohol (4.28 mL, 41.38 mmol). The solution was heated to reflux overnight. The reaction was cooled to room temperature, diluted with EtOAc, washed (1.0N NaOH then brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (3.05 g, 63%). Mass Spectrum (ES+) (m/z) 263.1 [M-BOC].

15

Preparation 105

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}(*t*-butoxy)carboxamide

A mixture of a compound from preparation 104 (0.25 g, 0.69 mmol), benzoic
20 anhydride (0.187 g, 0.828 mmol), 10% Pd/C catalyst (0.1 g), and EtOAc (10 mL) was stirred under an atmosphere of hydrogen gas (balloon) for 17 h. Added more benzoic anhydride (0.187 g, 0.828 mmol), and DMAP (0.025 g, 0.23 mmol), stirred for 2.5 h, filtered through celite, and concentrated. Column chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (0.208 g, 91%). Mass Spectrum (FIA) (m/z)
25 333.2 [M+1].

Preparation 106

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[3-(2-chloro-6-fluorophenyl)-5-
30 methylisoxazol-4-yl]carboxamide

In a fashion similar to that described for preparation 46, a compound from preparation 105 (0.1 g, 0.3 mmol), TFA (2.5 mL), dichloromethane (4 mL), 3-(2,6-

-90-

difluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (0.13 g, 0.5 mmol), and Et₃N (0.125 mL, 0.9 mmol) gave the title compound (0.109 g, 80%) after column chromatography (silica gel, hexanes/ EtOAc gradient). Mass Spectrum (ES+) (m/z) 454.2 [M+1].

5

Preparation 107

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[3-(2-chloro-6-fluorophenyl)-5-phenylisoxazol-4-yl]carboxamide

10 In a fashion similar to that described for preparation 46, a compound from preparation 103 (0.1 g, 0.3 mmol), TFA (2.5 mL), dichloromethane (4 mL), 3-(2-chloro-6-fluoro-phenyl)-5-phenyl-isoxazole-4-carbonyl chloride (0.15 g, 0.45 mmol), and Et₃N (0.125 mL, 0.9 mmol) gave the title compound (0.14 g, 88%) after column chromatography (silica gel, acetone/dichloromethane gradient). Mass Spectrum (FIA)
15 (m/z) 532.1 [M+1].

Preparation 108

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[5-(2-fluorophenyl)(1,3-oxazol-4-yl)]carboxamide

20

A solution of a compound from preparation 105 (0.1 g, 0.3 mmol) in TFA (2 mL) was stirred for 1.5 h. The solution was concentrated using benzene to azeotrope, diluted with EtOAc, washed (1.0N NaOH), dried (Na₂SO₄), filtered, and concentrated. The residue was dissolved in dichloromethane (1.5 mL) and added to a mixture of EDCI
25 (0.086 g, 0.45 mmol), DMAP (0.007 g, 0.06 mmol), 5-(2-fluoro-phenyl)-oxazole-4-carboxylic acid (0.087 g, 0.36 mmol) in dichloromethane (1.5 mL) under N₂. The reaction was stirred for 65 h, diluted with dichloromethane, washed (1.0N HCl, 2.0N NaOH, H₂O, and brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, acetone/ dichloromethane gradient) gave the title compound
30 (0.111 g, 81%). Mass Spectrum (FIA) (m/z) 456.2 [M+1].

-91-

Preparation 109

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[3-(2-fluoro-3-iodophenyl)-5-methylisoxazol-4-yl]carboxamide

5 In a fashion similar to that described for preparation 108, a compound from preparation 105(0.1 g, 0.3 mmol), TFA (2 mL), dichloro-methane (1.5 mL), EDCI (0.086 g, 0.45 mmol), DMAP (0.007 g, 0.06 mmol), 3-(2-fluoro-3-iodo-phenyl)-5-methyl-isoxazole-4-carboxylic acid (0.125 g, 0.36 mmol), and dichloromethane (1.5 mL) gave the title compound (0.104 g, 62%) after column chromatography (silica gel,
10 acetone/dichloromethane gradient). Mass Spectrum (FIA) (m/z) 562.1 [M+1].

Preparation 110

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[3-(2-fluoro-3-iodophenyl)-5-iodophenyl-4-yl]carboxamide

15 In a fashion similar to that described for preparation 108, a compound from preparation 105 (0.1 g, 0.3 mmol), TFA (2 mL), dichloro-methane (1.5 mL), EDCI (0.125 g, 0.45 mmol), DMAP (0.007 g, 0.06 mmol), 3-(2-fluoro-5-iodo-phenyl)-5-methyl-isoxazole-4-carboxylic acid (0.125 g, 0.36 mmol), and dichloromethane (1.5 mL) gave the
20 title compound (0.088 g, 53%) after column chromatography (silica gel, acetone/dichloromethane gradient). Mass Spectrum (FIA) (m/z) 562.1 [M+1].

Preparation 111

N-{(1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl}[3-(2,4-difluorophenyl)-5-methylisoxazol-4-yl]carboxamide

25 In a fashion similar to that described for preparation 108, a compound from preparation 105 (0.1 g, 0.3 mmol), TFA (2 mL), dichloromethane (1.5 mL), EDCI (0.086 g, 0.45 mmol), DMAP (0.007 g, 0.06 mmol), 3-(2,4-difluoro-phenyl)-5-methyl-isoxazole-
30 4-carboxylic acid (0.086 g, 0.36 mmol), and dichloromethane (1.5 mL) gave the title compound (0.123 g, 90%) after column chromatography (silica gel, acetone/dichloromethane gradient). Mass Spectrum (FIA) (m/z) 454.2 [M+1].

-92-

Preparation 112

N-((1S,3R)-3-[(phenylcarbonylamino)methyl]cyclohexyl)[3-(2-chloro-6-fluorophenyl)-5-hexylisoxazol-4-yl]carboxamide

5

In a fashion similar to that described for preparation 108, a compound from preparation 105 (0.1 g, 0.3 mmol), TFA (2 mL), dichloromethane (1.5 mL), EDCI (0.086 g, 0.45 mmol), DMAP (0.007 g, 0.06 mmol), 3-(2-chloro-6-fluoro-phenyl)-5-hexylisoxazole-4-carboxylic acid (0.117 g, 0.36 mmol), and dichloromethane (1.5 mL) gave the title compound (0.112 g, 69%) after column chromatography (silica gel, acetone/dichloromethane gradient). Mass Spectrum (FIA) (m/z) 540.5 [M+1].

10

Preparation 113

Phenylmethyl 2-((3S,1R)-3-{[5-methyl-3-(6-fluoro-2-iodophenyl)isoxazol-4-yl]carbonylamino}cyclohexyl)acetate

15

In a fashion similar to that described for preparation 99, a compound from preparation 45 (5.1 g, 14.7 mmol), dichloromethane (30 mL), TFA (30 mL), dichloromethane (150 mL), 3-(2-fluoro-6-iodo-phenyl)-5-methylisoxazole-4-carboxylic acid (6.39 g, 17.49 mmol), and Et₃N (6.03 mL, 43.4 mmol) gave the title compound (7.15 g, 86%) after column chromatography (silica gel, hexanes/ EtOAc gradient). Mass Spectrum (ES+) (m/z) 577.0 [M+1].

20

Preparation 114

Phenylmethyl 2-[(3S,1R)-3-(9-iodo-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]acetate

25

A compound from Example 499 (1.87 g, 3.35 mmol), 1.0N NaOH (6.7 mL, 6.7 mmol), and dioxane (20 mL) were allowed to react for 2 h at 50 °C in a fashion similar to that of preparation 29 to give the title compound (1.57 g, 100%). Mass Spectrum (ES+) (m/z) 467.0 [M+1].

30

-93-

Preparation 115

N-((3S,1R)-3-aminocyclohexyl)methyl) benzamide

5 A solution of a compound from preparation 105 (5.3 g, 15.96 mmol) in TFA (50 mL) was stirred for 2 h. The reaction was concentrated using benzene azeotrope, diluted with EtOAc, washed (1.0N NaOH), dried (Na₂SO₄), filtered, and concentrated to give the title compound (3.41 g, 92%). Mass Spectrum (ES+) (m/z) 233.1 [M+1].

Preparation 116

10 N-[(1S,3S)-3-(3-oxo-3-phenylpropyl)cyclohexyl][5-methyl-3-(6-fluoro-2-iodophenyl)isoxazol-4-yl]carboxamide

15 To a solution of a compound from preparation 115 (3.4 g, 14.7 mmol), Et₃N (6.14 mL, 44.1 mmol) in dichloromethane (150 mL) was added 3-(2-fluoro-6-iodo-phenyl)-5-methylisoxazole-4-carboxylic acid (6.97 g, 19.4 mmol) under N₂ and stirred for 4 h. The reaction was diluted with dichloromethane, washed (0.1N HCl then brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, acetone/dichloromethane gradient) gave the title compound (4.89 g, 59%). Mass Spectrum (ES+) (m/z) 562.0 [M+1].

20

Preparation 117

4-Amino-1-ethylcyclohexanecarboxylate hydrochloride

25 Thionyl chloride (2.55 mL, 35 mmol) was added dropwise to absolute ethanol (100 mL) under a N₂ atmosphere. The resulting solution was stirred at room temperature for 10 min then 4-amino-1-cyclohexanecarboxylic acid (4.29 g, 30 mmol, *cis, trans* mixture) was added as a solid. The resulting solution was stirred at room temperature overnight. The solvent was removed in vacuo, the residue dissolved in ethanol (25 mL) and this solution was slowly added to a flask containing rapidly stirred ether (700 mL).
30 The precipitate that formed was collected by filtration and dried in vacuo to give 5.75 g, 92% as a white solid. MS (ion spray) 172 (M+1).

-94-

Preparation 118

Ethyl 4-([3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino)cyclohexanecarboxylate

5 To a solution of a compound from preparation 117 (1.04 g, 5 mmol) in methylene chloride (15 mL) was added triethyl amine (0.8 mL, 5.7 mmol) followed by 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (1.37 g, 5 mmol) in methylene chloride (10 mL), and an additional aliquot of triethyl amine (0.75 mL, 5.3 mmol). After an initial exotherm the mixture was stirred at room temperature for 4 h. The mixture was
10 diluted with methylene chloride, washed with water, 1N aqueous hydrochloric acid solution, and brine, dried (MgSO₄), filtered, and concentrated in vacuo. The residue was purified by flash chromatography, eluting with 4:1 toluene: ethyl acetate, to give 2.0 g 98% of the title compound as a white solid. ¹H-NMR was consistent with the desired structure. MS (ion spray) 409.3 (M+1).

15

Preparation 119

Ethyl 4-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]-quinolin-5-yl)cyclohexanecarboxylate

20 To a room temperature solution of a compound from preparation 118 (1.75 g, 4.28 mmol) in anhydrous DMF (20 mL) was added potassium *t*-butoxide (560 mg, 5 mmol). The resulting solution was stirred for 50 min, at which time TLC (1:1 hexane: ethyl acetate) showed the starting material had been consumed. The addition of 1N hydrochloric acid solution (5 mL) and water (95 mL) quenched the reaction. The mixture
25 was extracted with ethyl acetate, the extracts were washed with water, and brine, dried (MgSO₄), filtered, and concentrated in vacuo. The residue was purified by flash chromatography, eluting with 9:1 methylene chloride: THF, to give 1.43 g 85% of the title compound, approximately 2:1 mixture of diastereoisomers, as a yellow tinted solid. ¹H-NMR was consistent with the desired structure. MS (ion spray) 389 (M+1).

30

-95-

Preparation 120

4-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane carboxylic acid

5 To a solution of a compound from preparation 119 (390 mg, 1 mmol) in THF (7 mL) was added 0.5 M LiOH solution (5 mL). The mixture was stirred at room temperature for 3 days then acidified with of 0.1N hydrochloric acid solution (5 mL) and extracted with ethyl acetate. The extracts were washed with brine, dried (MgSO₄), filtered, and concentrated in vacuo to give 370 mg of the title compound as a yellow
10 foam. NMR was consistent with the desired product.

Preparation 121

(3-Aminocyclohexyl)(phenylsulfonyl)amine

15 A solution of 1,3-diaminocyclohexane (1.14 g, 10 mmol, undetermined isomer mixture) in methylene chloride (50 mL) was cooled to 0 °C and phenylsulfonyl chloride (0.64 mL, 5 mmol) was added dropwise. The mixture was stirred at 0 °C for 4 h, then 1N sodium hydroxide solution (10 mL) was added and the mixture stirred an additional 5 min. The phases were separated, the organic phase was dried (K₂CO₃), filtered, and
20 concentrated in vacuo. Toluene was added and the mixture re-concentrated to remove any remaining diamine, to give 0.77 g of a semi solid. ¹H-NMR was consistent with the desired structure. MS (ion spray) 255 (M+1).

Preparation 122

25 [3-(2-Chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-{3-[(phenylsulfonyl)amino]cyclohexyl}carboxamide

To a solution of a compound from preparation 121 (750 mg, 2.95 mmol theoretical) in methylene chloride (30 mL) was added triethylamine (0.6 mL, 4.3 mmol)
30 followed by 3-(2-chloro-6-fluoro-phenyl)5-methylisoxazole-4-carbonyl chloride (822 mg, 3 mmol). The resulting mixture was stirred at room temperature for 3 h. The mixture was diluted with ethyl acetate, washed with 1N hydrochloric acid solution, saturated aqueous

-96-

sodium bicarbonate solution, brine, then dried (MgSO_4), filtered, and concentrated in vacuo. Radial chromatography eluting with 95:5 methylene chloride: THF gave the less polar *cis* isomer (780 mg), mixed fractions (130 mg), and the more polar *trans* isomer (410 mg). The *cis* isomer: $^1\text{H-NMR}$ was consistent with the desired structure, plus a small amount of the bis-sulfonylated diamine. MS (ion spray) 492 ($\text{M}+1$). For the *trans* isomer: $^1\text{H-NMR}$ was consistent with the desired structure. MS (ion spray) 492 ($\text{M}+1$).

Preparation 123

2-{(3S,1R)-3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid

10

A compound from preparation 45 (3.30 g, 0.0095 mol) was combined with aqueous 2N sodium hydroxide (35 mL), tetrahydrofuran (10 mL), and ethanol (10 mL) and the mixture stirred at room temperature for 2 h. The mixture was then concentrated *in vacuo* to remove volatile organics and additional water was added to the aqueous mixture. This basic aqueous mixture was then extracted with diethylether followed by careful acidification using 1N HCl. The acidic aqueous mixture was then extracted with ethyl acetate. The combined ethyl acetate extracts were dried over sodium sulfate followed by concentration *in vacuo* which provided the title compound (2.40 g, 98 %) as a white solid. MS(ES): ($\text{M}-1$)⁻ 256.2 m/z.

15
20

Preparation 124

N-[(1S,3R)-3-(2-Hydroxyethyl)cyclohexyl](*t*-butoxy) carboxamide

The compound from preparation 123 (2.10 g, 0.0082 mol) was dissolved in dry THF (35 mL) and the mixture cooled in an ice bath under a nitrogen atmosphere. Then borane-tetrahydrofuran (1.0 M, 12.25 mL, 0.0122 mol) was added *via* syringe and the mixture stirred overnight while slowly warming to room temperature. The reaction mixture was quenched into ice water and solid NaCl was added. This mixture was extracted with ethyl acetate and the combined extracts were dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was chromatographed over silica gel using a $\text{CH}_2\text{Cl}_2/\text{THF}$ mixture as eluant which allowed for isolation of 1.35 g (67%) of the desired alcohol. MS(ES): ($\text{M}+1$)⁺ 244.5 m/z.

25
30

-97-

Preparation 125N-[(1S,3R)-3-(2-*p*-toluenesulfonyl)ethyl]cyclohexyl(*t*-butoxy)carboxamide

5 The compound from preparation 124 (1.30 g, 0.0054 mol) was combined with *p*-toluenesulfonyl chloride (1.12 g, 0.0059 mol), triethylamine (1.49 mL, 0.011 mol) and DMAP (0.02 g, cat.) in dichloromethane (30 mL) and the resultant mixture stirred overnight at room temperature. The mixture was then concentrated *in vacuo* and the residue chromatographed over silica gel using a mixture of CH₂Cl₂/THF as eluant which
10 allowed for the isolation of the desired product (1.92 g, 90%). MS(ES): (M+1)⁺ 398.2 m/z.

Preparation 126N-[(1S,3R)-3-[2-(3-methylphenylthio)ethyl]cyclohexyl](*t*-butoxy)carboxamide

15

3-Methylthiophenol (0.15 mL, 0.0013 mol) was dissolved in DMF (10 mL) at ambient temperature under a nitrogen atmosphere and sodium hydride (60%, 0.050 g, 0.0013 mol) was added. After stirring for 15 min., the compound from preparation 125 (0.50 g, 0.0013 mol) was added and the mixture stirred overnight at ambient temperature.
20 The mixture was concentrated *in vacuo* and the residue chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.42 g (95 %) of the desired product. MS(ES): (M+1)⁺ 350.5 m/z.

Preparation 12725 N-[(1S,3R)-3-{2-[3-methylphenyl)sulfonyl]ethyl}cyclohexyl] (*t*-butoxy)carboxamide

The compound from preparation 126 (0.40 g, 0.00115 mol) was dissolved in a methanol/acetone (10 mL)/(10 mL) mixture and NaHCO₃ (0.40 g, 0.0048 mol) and water (5 mL) were added. Oxone® (1.46 g, 0.0024 mol) was then added and the mixture stirred
30 at ambient temperature for 3.5 h. The mixture was concentrated *in vacuo* and the residue taken up in aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was

-98-

chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.420 g (96%) of the desired sulfone. MS(FD): M⁺ 381.2, 382.2 m/z.

Preparation 128

5 N-((1S,3R)-3-{2-[(3-methylphenyl)sulfonyl]ethyl}cyclohexyl) [3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl] carboxamide

The compound from preparation 127 (0.385 g, 0.0010 mol) and TFA (1.0 mL) were dissolved in CH₂Cl₂ (5.0 mL) and the mixture stirred at ambient temperature for 2
10 h. The mixture was concentrated *in vacuo* and the residue treated with aqueous NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were dried over sodium sulfate and concentrated *in vacuo* which allowed for recovery of crude amine. This amine was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methylisoxazole-4-carbonyl chloride (0.302 g, 0.0011 mol), triethylamine (0.31 mL, 0.0022 mol) and DMAP (0.010 g, cat.) in
15 CH₂Cl₂ (6.0 mL) and the mixture stirred overnight at ambient temperature. The mixture was concentrated *in vacuo* and the resulting residue chromatographed over silica gel using CH₂Cl₂/THF as eluant which allowed for isolation of 0.463 g (88%) of the desired amide. MS(ES): (M+1)⁺ 519.2, 521.2 m/z.

20

Preparation 129

N-[(1S,3R)-3-(2-phenylthioethyl)cyclohexyl](*t*-butoxy)carboxamide

Thiophenol (0.32 mL, 0.0031 mol) was combined with sodium hydride (60%, 0.15 g, 0.0038 mol) in DMF (15 mL) and a compound from preparation 125 (1.25 g, 0.0031
25 mol) subsequently added in a manner similar to preparation 126. Workup and purification allowed for isolation of 1.0 g (98%) of the desired product. MS(ES): (M+1)⁺ 335.1 m/z.

-99-

Preparation 130

N-[(1S,3R)-3-(2-phenylthioethyl)cyclohexyl][3-(2-chloro-6-fluorophenyl)-5-methyl]isoxazol-4-yl]carboxamide

5 The compound from preparation 129 (0.50 g, 0.0015 mol) was combined with TFA (2 mL) in CH₂Cl₂ (10 mL) and treated in a manner similar to preparation 128. The resulting crude amine (0.213 g, 0.00097 mol) was combined with 3-(2-chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl chloride (0.27 g, 0.001097 mol) and triethylamine (0.27 mL, 0.0019 mol) in CH₂Cl₂ (10 mL) again in a manner similar to preparation 128.
10 Workup and purification allowed for isolation of 0.413 g (58%) of the desired amide. MS(ES): (M+1)⁺ 473.1, 475.1 m/z.

Preparation 131

N-((1S,3R)-3-{2-[benzylsulfonyl]ethyl}cyclohexyl)(*t*-butoxy)carboxamide

15

Benzyl mercaptan (0.15 mL, 0.0013 mol) was combined with sodium hydride (60%, 0.05 g, 0.0013 mol) in DMF (10 mL) and the compound from preparation 125 (0.50 g, 0.0013 mol) subsequently added in a manner similar to preparation 126. Workup and purification allowed for isolation of 0.27 g (61%) of the desired sulfide. This
20 compound (0.27 g, 0.0008 mol) was dissolved in a methanol/acetone (5 mL)/(5 mL) mixture and NaHCO₃ (0.27 g, 0.0032 mol) and water (5 mL) were added. Oxone® (0.98 g, 0.0016 mol) was then added and the mixture treated in a similar manner to preparation 127. Workup and purification allowed for isolation of 0.092 g (30%) of desired sulfone intermediate. MS(ES): (M+1)⁺ 382.5 m/z.

25

Preparation 132

N-((1S,3R)-3-{2-[Benzylsulfonyl]ethyl}cyclohexyl[3-(2-chloro-6-fluorophenyl)-5-methyl]isoxazol-4-yl]carboxamide

30

The compound from preparation 131 (0.090 g, 0.00024 mol) was combined with TFA (1 mL) in CH₂Cl₂ (3 mL) and treated in a similar manner to preparation 127. The resulting crude amine was combined with 3-(2-chloro-6-fluoro-phenyl)-5-

-100-

methylisoxazole-4-carbonyl chloride (0.067 g, 0.00024 mol), triethylamine (4 drops), and DMAP (0.005 g, cat.) in CH₂Cl₂ (3 mL) again in a manner similar to preparation 126.

Workup and purification allowed for isolation of 0.10 g (79%) of the desired amide.

MS(ES): (M+1)⁺ 519.2, 521.1 m/z.

5

Preparation 133

N-((1S,3R)-3-[2-(3-Fluorophenylthio)ethyl]cyclohexyl)(*t*-butoxy)carboxamide

3-Fluorothiophenol (0.14 mL, 0.0013 mol) was combined with sodium hydride (60%, 0.055 g, 0.0014 mol) in DMF (6 mL) and the compound from preparation 125 (0.50 g, 0.0013 mol) subsequently added in a manner similar to preparation 126. Workup and purification allowed for isolation of 0.40 g (89%) of the desired product. MS(ES): (M+1)⁺ 354 m/z.

15

Preparation 134

N-((1S,3R)-3-{2-[(3-Fluorophenyl)sulfonyl]ethyl}cyclohexyl)(*t*-butoxy)carboxamide

The compound from preparation 133 (0.38 g, 0.0011 mol) was dissolved in a methanol/acetone (10 mL)/(10 mL) mixture and NaHCO₃ (0.36 g, 0.0043 mol) and water (5 mL) were added. Oxone® (1.30 g, 0.0021 mol) was then added and the mixture treated in a manner similar to preparation 125. Workup and purification allowed for isolation of 0.40 g (98%) of the desired sulfone intermediate. MS(ES): (M+1)⁺ 386.3 m/z.

20

Preparation 135

N-((1S,3R)-3-[2-(3-Fluorophenylthioethyl)cyclohexyl][3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

25

The compound from preparation 134 (0.030 g, 0.00078 mol) was combined with TFA (1.5 mL) in CH₂Cl₂ (6 mL) and treated in a manner similar to preparation 126. The resulting crude amine was combined with 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.213 g, 0.00078 mol), triethylamine (0.22 mL, 0.0016 mol) and DMAP (0.015 g, cat.) in CH₂Cl₂ (10 mL) again in a manner similar to preparation 126.

30

-101-

Workup and purification allowed for isolation of 0.38 g (93%) of the desired amide.

MS(ES): (M+1)⁺ 523.1, 525.2 m/z.

Preparation 136

5 N-((1S,3R)-3-[2-(4-Fluorophenylthio)ethyl]cyclohexyl)(*t*-butoxy)carboxamide

4-Fluorothiophenol (0.16 mL, 0.0015 mol) was combined with sodium hydride (60%, 0.059 g, 0.0015 mol) in DMF (15 mL) and a compound from preparation 125 (0.59 g, 0.0015 mol) subsequently added in a manner similar to preparation 133. Workup and
10 purification allowed for isolation of 0.52 g (98%) of the desired product. MS(ES): (M+1)⁺ 353.6 m/z.

Preparation 137

15 N-((1S,3R)-3-{2-[(3-Fluorophenyl)sulfonyl]ethyl}cyclohexyl)(*t*-butoxy)carboxamide

A compound from preparation 136 (0.50 g, 0.0014 mol) was dissolved in a methanol/acetone (10 mL)/(10 mL) mixture and NaHCO₃ (0.50 g, 0.0060 mol) and water (8 mL) were added. Oxone® (1.83 g, 0.0030 mol) was then added and the mixture treated in a manner similar to preparation 125. Workup and purification allowed for isolation of
20 0.50 g (92%) of the desired sulfone intermediate. MS(ES): (M+Na)⁺ 408.1 m/z.

Preparation 138

25 N-((1S,3R)-3-{2-[(4-Fluorophenyl)sulfonyl]ethyl}cyclohexyl)-[3-(6-chloro-2-fluorophenyl)-5-methylisoxazol-4-yl]-carboxamide

The compound from preparation 137 (0.45 g, 0.0012 mol) was combined with TFA (2 mL) in CH₂Cl₂ (10 mL) and treated in a manner similar to preparation 126. The resulting crude amine (0.30 g, 0.0011 mol) was combined with 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.29 g, 0.0011 mol), triethylamine (0.29 mL, 0.0021 mol) and DMAP (0.010 g, cat.) in CH₂Cl₂ (20 mL) again in a manner
30 similar to preparation 126. Workup and purification allowed for isolation of 0.506 g (92%) of the desired amide. MS(ES): (M+1) 523.1, 525.1 m/z.

-102-

Preparation 139

Diphenylmethyl 2-((1R)-3-oxocyclohexyl)propane-1,3-dioate

5 To a flame dried flask at 0 °C was added lithium aluminum hydride (3.0 mL, 3.0 mmol, 1.0M in tetrahydrofuran) followed by (R)-(+)-1,1'-bi-2-naphthol (1.71 g, 6.0 mmol) in anhydrous tetrahydrofuran (24 mL). After 30 min, the solution was allowed to warm to room temperature. A solution of sodium hydride (108 mg, 2.7 mmol, 60% in mineral oil) and dibenzylmalonate (675 μ L, 2.7 mmol) in anhydrous tetrahydrofuran (30 mL) was then
10 added to the reaction dropwise over approximately 2 min. To this mixture were then added 2-cyclohexen-1-one (2.91 mL, 30.0 mmol) and dibenzylmalonate (7.5 mL, 30 mmol), and the reaction was then stirred at room temperature, under a nitrogen atm, overnight. To the reaction was added 1N hydrochloric acid (60 mL, 60 mmol, 1N in water), followed by extraction with ethyl acetate (3 x 200 mL). The combined organic
15 layers were then washed with saturated sodium bicarbonate solution, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 5-7.5% acetone/hexane) gave 10.25 g of the title compound as a opaque solid, 90% yield. ^1H NMR: consistent with structure. MS (ion spray) 381 ($\text{M}^+ + 1$).

20

Preparation 140*cis*-Diphenylmethyl 2-(3-hydroxycyclohexyl)propane-1,3-dioate

To a solution of a compound from preparation 139 (6.13 g, 16.1 mmol) in tetrahydrofuran (120 mL), methanol (10 mL), and water (10 mL) was added sodium
25 borohydride (1.22 g, 32.2 mmol). After 10 min., the reaction was poured into 150 mL of a saturated ammonium chloride solution, and extracted with ethyl acetate. The ethyl acetate layer was washed with additional saturated ammonium chloride solution, saturated sodium bicarbonate solution, dried over magnesium sulfate, and concentrated. Purification by flash chromatography on silica gel (eluting with 5-13% acetone/hexane)
30 gave 5.38 g of the title compound as a clear oil, 87% yield. ^1H NMR: consistent with structure. MS (ion spray) 383 ($\text{M}^+ + 1$).

-103-

Preparation 141*cis*-Phenylmethyl 2-(3-hydroxycyclohexyl)acetate

To a solution of a compound from preparation 140 (5.38 g, 14.07 mmol) in
5 anhydrous methylsulfoxide was added water (0.51 mL, 28.13 mmol) and lithium chloride
(1.19 g, 28.13 mmol). The reaction was placed in an oil bath at 175 °C and stirred
vigorously for 2.5 h. The reaction was added to water and extracted with ethyl acetate
twice. The organic layer was washed with water (x2), saturated sodium bicarbonate
solution (x2), dried over magnesium sulfate and concentrated. Purification by flash
10 chromatography on silica gel (eluting with 13% acetone/hexane) gave 1.81 g of the title
compound as a yellow oil, 52% yield. ¹H NMR: consistent with structure.

Preparation 142*cis*-Phenylmethyl 2-(3-azidocyclohexyl)acetate

15

To a solution of a compound from preparation 141 (1.81 g, 7.29 mmol) in
anhydrous toluene (70 mL) was added triphenylphosphine (3.82 g, 14.58 mmol),
Zn(N₃)₃·pyridine₂ (1.68 g, 5.47 mmol), followed by dropwise addition of diethyl
azodicarboxylate (2.30 mL, 14.58 mmol) over approximately 15 min., and the reaction
20 was stirred at room temperature overnight under a nitrogen atmosphere. Filtered through
Celite® to remove zinc salts, dilute with ethyl acetate, wash with 0.1N sodium hydroxide
(x3), water, saturated sodium bicarbonate solution, dry over magnesium sulfate and
concentrated. Purification by flash chromatography on silica gel (eluting with 3.5% ethyl
acetate/hexane) gave 652 mg of the title compound as an oil, 33% yield. ¹H NMR:
25 consistent with structure.

Preparation 143*cis*-Phenylmethyl 2-{3-[(*t*-butoxy)carbonylamino]cyclohexyl}acetate

30

To a solution of a compound from preparation 142 (652 mg, 2.4 mmol) and *t*-
butoxycarbonyl anhydride (1.10 mL, 4.8 mmol) in ethyl acetate (50 mL) under a nitrogen
atmosphere was added Lindlar's catalyst (270 mg, 5% by weight). The mixture was

-104-

subjected to a hydrogen atmosphere at room temperature overnight. The mixture was filtered over Celite® to remove the catalyst and concentrated. Purification by flash chromatography on silica gel (eluting with 15% ethyl acetate/hexane) gave 820 mg of the title compound as a clear oil, 100% yield.

5 ¹H NMR: consistent with structure.

Preparation 144

cis-Phenylmethyl 2-(3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)acetate

10

A solution of a compound from preparation 143 in trifluoroacetic acid was stirred at room temperature for 30 min. The solution was concentrated, followed by azeotropic removal of the remaining solvent with acetonitrile (x3). The crude product was dissolved in ethyl acetate, washed with 1N sodium hydroxide (x2), dried over magnesium sulfate and concentrated. To a solution of the resulting oil in dichloromethane (20 mL) was added 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (748 mg, 2.72 mmol), triethylamine (1.03 mL, 7.43 mmol), and dimethylamino pyridine (30 mg, 0.25 mmol), followed by triethylamine (1.03 mL, 7.43 mmol) dropwise via syringe. The reaction was stirred overnight under nitrogen at room temperature. The solution was washed with 0.1N HCl (x3), water, saturated sodium bicarbonate solution, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 0-30% ethyl acetate/hexane) gave 1.13 g of the title compound as a clear oil, 94% yield. ¹H NMR: consistent with structure.

15
20
25

Preparation 145

2-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]acetic acid

To a solution of a compound from Example 494 (291 mg, 0.63 mmol) in dioxane (5 mL) was added 5N NaOH (5 mL) and the solution heated to reflux for 1 h. Upon cooling the reaction was quenched with 1N hydrochloric acid, and extracted with 20% isopropanol/chloroform (x3), dried over magnesium sulfate and concentrated. The

30

-105-

resulting brown oil was treated with ethyl ether, sonicated, and a brown solid filtered out, to give 164 mg of the title compound, 70% yield. ^1H NMR: consistent with structure.

Preparation 146

5 *trans*-5-[3-(Aminomethyl)cyclohexyl]-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one hydrochloride

To *trans*-racemic [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]carbamic acid *t*-butyl ester (1.22 g, 2.74 mmol) was added acetic acid
10 saturated with $\text{HCl}_{(\text{g})}$ (30 mL, ~3N in HCl) and the solution stirred vigorously at room temperature for 10 min. The reaction was concentrated, followed by addition of acetonitrile and concentrated to assist in the removal of acetic acid. The resulting white solid was treated with ethyl ether, sonicated, and filtered to yield 985 mg of the title
15 compound as a white solid, 94% yield. ^1H NMR: consistent with structure. MS (ion spray) 346 ($\text{M}^+ + 1$).

Preparation 147

2-Fluoro-3-iodobenzaldehyde

20 To a solution of diisopropylamine (10.42 mL, 74.36) in THF (135 mL) under N_2 in an ice bath at 0-5 °C was added dropwise over 10 min *n*-BuLi (42.25 mL, 67.6 mmol) and stirred at this temperature for 10 min. Cooled the reaction mixture to -78 °C in a dry ice/acetone bath and dropwise added 1-fluoro-2-iodobenzene (7.88 mL, 67.6 mmol) over
25 5 min. Stirred at this temperature for 1 h, dropwise added DMF (6.26 mL, 74.36 mL) over 5 min, and stirred for 10 min. Added Acetic Acid (13.5 mL) followed by H_2O and extracted with Et_2O (2x). The combined organic solution was washed (0.1 N HCl then brine), dried (MgSO_4), filtered and concentrated. The crude benzaldehyde (16.9 g) was taken onto the next step without purification. ^1H NMR was consistent with structure.

-106-

Preparation 148

2-Fluoro-3-iodobenzaldoxime

To a mixture of a compound from preparation 147 (16.9 g, 67.6 mmol) in H₂O (35 mL), EtOH (35 mL), and ice (25 g) was added hydroxylamine hydrochloride (4.8 g, 74.4 mmol). Then, 169 mmol of 50% NaOH (6.76 g in 6.76 mL H₂O) was added with stirring. Enough ice to keep the temperature at 25-30 °C was added. Stirred for 2.5 h, acidified with conc. HCl to pH 4 (ice was added to keep the temperature at 25-30 °C), and extracted with Et₂O (2x). The combined organic solution was washed (brine), dried (MgSO₄), filtered, and concentrated. The residue was chromatographed on silica gel (EtOAc/dichloromethane gradient) to give the title compound (8.02g, 45% over 2 steps). Mass Spectrum (FIA) (m/z) 264.0 [M-1].

Preparation 149

15

2-Fluoro-3-iodobenzohydroximinoyl chloride

To a stirred solution of a compound from preparation 148 (8.02 g, 30.2 mmol) in DMF (45 mL) at 25-30 °C was added about 1/5 of 30.2 mmol (4.03 g) of NCS. The initial NCS addition results in a slight temperature decrease. If the reaction does not self-initiate within 10 min., as indicated by a slight temperature rise, 5 pipettes of gas from the headspace of a conc. HCl reagent bottle was bubbled into the DMF solution. Carefully added rest of NCS and temperature rose to 45-55 °C. Once reaction cooled to r.t. (about 1 h), added ice H₂O and extracted with Et₂O (2x). The combined organic solution was washed with (brine), dried (MgSO₄), filtered, and concentrated. The title compound (9.04 g) was taken on to the next step without purification. ¹H NMR was consistent with structure.

Preparation 150

30

Ethyl 3-(6-fluoro-2-iodophenyl)-5-methylisoxazole-4-carboxylate

To a solution of a compound from preparation 149 (30.2 mmol) and ethyl-2-butynoate (4.23 mL, 36.24 mmol) in Et₂O (120 mL) under N₂ stirring at 0-5 °C was

-107-

added dropwise a solution of Et₃N (5.46 mL, 39.26 mmol) in Et₂O (17 mL) over 1 h. Allowed to warm to room temperature and stirred overnight. Added Et₂O and washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. The crude residue was chromatographed on silica gel (dichloromethane) to give the title compound (6.92 g, 61%). Mass Spectrum (FIA) (m/z) 376.15 [M+1].

Preparation 153

N-*t*-Butyl-N'-(2-chloro-6-fluorobenzylidene)hydrazine hydrochloride

A mixture of *t*-butyl hydrazine hydrochloride (1.24 g, 10 mmol) and 2-chloro-6-fluorobenzaldehyde (1.1 mL, 10 mmol) dissolved in acetic acid (5 mL) was stirred at 50 °C for half hour. Hexanes (10 mL) was added, filtered and washed the white solid with hexanes (20 mL). Yield: 61%. ESMS: 229 (M+1).

Preparation 156

3-Methyl-5-(2-chloro-6-fluorophenyl)-4-isoxazolecarboxylic acid ethyl ester

To a solution of ethyl 3-aminomethyl crotonate (4.79 g, 33.5 mmol) in toluene (10 mL), was added triethylamine (3.73 g, 37 mmol). The solution was chilled using an ice bath, and then 2-chloro-6-fluorobenzoyl chloride (6.47 g, 33.5 mmol) was added dropwise over a 20 min period. The reaction was allowed to warm slowly to r.t. and stirred for 24 hr. The resulting suspension was then filtered, and the filtrate diluted with ethyl acetate (100 mL) and transferred to a separatory funnel. The organic layer was sequentially washed with water, brine, dried (sodium sulfate), and the volatiles removed under reduced pressure to provide the title compound (9.46 g) as a golden solid, and primarily one geometrical isomer. MS (ES) m/z 299.9 (M+H)⁺.

The crude adduct was then redissolved in glacial HOAc (50 mL) to which was added NH₂OH.HCl (1.8 g, 1.1 eq). The solution was then heated to reflux for 40-45 min to effect isoxazole formation. The reaction mixture was concentrated to an oil, diluted with ether, and transferred to a sep. funnel. The organic phase was washed with saturated bicarbonate, brine, then dried. Filtration and concentration afforded crude isoxazole ethyl ester (7.5 g), which was used without further purification. MS (+ ES) m/z 283.9 (M+H)⁺.

-108-

Preparation 157

5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carboxylic acid

5 Hydrolysis of the ethyl ester was accomplished by dissolving the crude ester (7.5 g, approx. 0.027 mol) in THF (250 mL), and adding aq. LiOH (1.344 g in 100 mL, 2 eq). After stirring overnight at r.t., the solution was concentrated to 2/3rd volume, diluted with EtOAc (200 mL) and 50 mL water, transferred to a separatory funnel, and the aqueous phase collected. The organic phase was washed twice, and the combined aqueous phase
10 was then acidified with 5N HCl. Back extraction with three washings of EtOAc was then followed with a brine wash of the combined organics. After drying over Na₂SO₄, filtration and concentration, clean isoxazole acid was obtained (2.94 g). NMR (CDCl₃) δ 7.13, 7.32 7.46 (3m, 3H), 2.58 (s, CH₃). MS (-ES) m/z 253.8, 255.8 (M-H)⁻.

15

Preparation 158

2-(3-{[5-(2-chloro-6-fluorophenyl)-3-methylisoxazol-4-yl]carbonylamino}cyclohexyl)-N-(3,4,5-trimethoxyphenyl)acetamide

To a compound from preparation 157 (60 mg, 0.234 mmol) in benzene (4 mL) containing catalytic amount of pyridine (20 μ l) was added oxalyl chloride (23 mL). After
20 heating to reflux for 1 hr, an aliquot was concentrated under vacuum and analyzed by ¹H-NMR (CDCl₃) for completion of acid chloride formation; ¹H-NMR (CDCl₃) 7.53 (d of t, 1H), 7.37 (d, 1H), 7.17 (t, 1 H). (racemic)

Solvent was removed under vacuum. To a solution of d,l-cis-1-amino,3-(N-3,4,5-trimethoxyphenyl)cyclohexyl-acetamide (60 mg, 0.185 mmol) in dry MeCl₂ (4 mL) containing catalytic DMAP (10%) and pyridine (20 μ l, 0.24 mmol) was added the crude acid chloride in 0.5 mL MeCl₂ at r.t. Reaction was allowed to proceed for 4 hr at which time amide formation was complete by TLC. The reaction was diluted with MeCl₂ and transferred to a sep funnel, washed with dil. HCl (pH 1), and the aqueous fraction was
25 separated and reextracted two more times in the same manner. The combined organics were then washed sequentially with sat'd bicarbonate, brine, dried over Na₂SO₄, filtered
30 and concentrated under vacuum to afford crude product (69 mg). Purification was

-109-

accomplished via prep TLC (Chromatotron®, 1 mm SI plate) using 70% - 90% EtOAc in hexane to obtain 24 mg (24% from amine). MS (+ES) m/z 559.9 (M+H), 576.9 (M+NH₃).

5

Preparation 159

(3-{{[5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexyl)acetic acid ethyl ester (*cis*)

A compound from preparation 157 (254 mg, 1 mmol) was dissolved in DMF (15 mL). EDC.HCl (211 mg, 1.1 eq) was then added, followed by DMAP (10 mg), and then *cis*-1,3-1(S)aminocyclohexyl-3(R)-acetic acid, methyl ester (60%ee, 188 mg, 1.1 eq) at r.t. After 18 hr, starting material was not yet fully consumed (TLC, NMR of aliquot), so the mixture was heated to 80° for 2 hr. The reaction was then cooled to r.t., and diluted with EtOAc and 0.1 N HCl. The aqueous phase was reextracted twice, and the combined organics washed with sat'd bicarbonate, brine, dried over Na₂SO₄. Filtration and concentration afforded crude product (207 mg). Purification using a Bond-Elut SI column (10g, 4:3 hex/EtOAc) provided 34 mg (8%) of the title compound. MS (+ES) 411.0 (M+H)+.

20

Preparation 160

(3-{{[5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexylmethyl)carbamic acid *t*-butyl ester

Using the acid chloride method described for preparation 158, 3 mmol (0.689 g) of a *cis/trans* mixture of 1-amino, 3-Boc-aminomethylene cyclohexane was dissolved in CH₂Cl₂ (10 mL), followed by addition of Et₃N (0.415 mL, 6 mmol), DMAP (cat, 20 mg), at rt. The above isoxazole acid chloride of preparation 157 (3 mmol in 10 mL CH₂Cl₂) was then added dropwise over 10 min. Reaction was allowed to proceed overnight. The reaction mixture was then transferred to a sep funnel, diluted with additional CH₂Cl₂, and washed sequentially with 5% citric acid (aq, pH 3-4), sat'd bicarbonate, brine, then dried. Filtration and concentration yielded crude product (brown

-110-

oil). Purification by filtration through a pad of Silica Gel 60 in a 60 mL sintered glass funnel using 2:1 hexane/EtOAc afforded clean cis/trans product (0.830 g, 59% overall). NMR (CDCl₃) δ 7.2 – 7.6 (3 m, 3H), 5.55 and 5.2 (trans/cis amide NH's), 4.5 and 4.2 (cis/trans Boc NH's), 3.8 and 2.95 (trans/cis aminomethine (CH)), 2.58 and 2.55 (trans/cis CH₃), 0.6 – 2.9 (m's), 1.4 (2s, cis/trans *t*-butyl). MS (+ES); 465.9 (M+H)⁺; (-ES); 463.9 (M-H)⁻, 499.9 (M+Cl-H)⁻, 524 (M+Oac-H)⁻.

Preparation 161

(*cis*-3(S)-Aminocyclohexylmethyl)carbamic acid benzyl ester

10

S-amino enantiomer of N-({3-[(*t*-butoxy)carbonyl-amino]cyclohexyl}methyl)(phenylmethoxy)carboxamide (1.0 g, 2.76 mmol) was treated with TFA (5 mL) under N₂. After 20 min of stirring at r.t. the reaction was complete. The crude was then concentrated to an oil which was purified on a Varian Bond-Elut SCX column (10 g). The column was eluted consecutively with CHCl₃, MeOH, and ammonia (2.0M in MeOH). The pure product was recovered from the ammonia fractions. The solvent was removed *in vacuo* to afford 0.632 g (87%) as a colorless oil. MS (ES+) *m/z* 263.0 (M+H)⁺.

20

Preparation 162

(3-{{5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexylmethyl)carbamic acid benzyl ester

A solution from preparation 157 (1.0 g, 3.9 mmol) in toluene (30 mL) was treated with a catalytic amount of pyridine (0.1 mL) and cooled to 0 °C. The solution was then treated with oxalyl chloride (0.545 g, 4.3 mmol) and stirred at r.t. for 2 hr. ¹H NMR showed the completion of the acid chloride formation; 7.52 (d, 1H), 7.37 (d, 1H), 7.15 (t, 1H). The solvent was removed *in vacuo*. A solution from preparation 161 (0.550 g, 2.09 mmol) and triethylamine (0.422 g, 4.18 mmol) in dry DMF (25 mL) was stirred at r.t. This solution was then treated with the acid chloride from preparation 158 (0.859 g, 3.135 mmol) which was added dropwise over two min. The reaction was then catalyzed with DMAP (0.025 g, 0.21 mmol) and allowed to stir o.n. The reaction

30

-111-

was diluted in EtOAc (250 mL), transferred to a sep funnel, and washed with a 5% citric acid solution (5 x 50 mL), sat. sodium bicarbonate solution (2 x 50 mL), brine (2 x 50 mL) and was dried over sodium sulfate. The EtOAc solution was filtered and the solvent removed *in vacuo* to yield an orange solid. The solid was purified using silica gel column chromatography. The column was prepared with CHCl₃ and the product was eluted with 10% EtOAc in CHCl₃. The solvent was removed *in vacuo* to afford 0.584 g (56%) of the title compound as a white solid. MS (ES+) m/z 500.1 (M+H)⁺.

Preparation 163

10

cis-Octahydroquinolin-2-one

To a solution of 2-nitrocinnamic acid in 120 mL of acetic acid in a Parr shaker, was added PtO₂ (10 g). Hydrogen (3 atm) was then introduced, and the contents of the shaker were heated to 60 °C for 24 h. After the mixture was cooled to r.t. and filtered through Celite®, the solvent was removed *in vacuo*, and the residue was dissolved in EtOAc, and transferred to a sep. funnel. Treatment with sat'd HCO₃⁻, brine, and then drying over Na₂SO₄, filtration and concentration afforded predominantly *cis* lactam (<7% trans by NMR) in 55% yield (6 g). MS (+ES) m/z 254 (M+1)⁺.

20

Preparation 164

trans-Octahydroquinolin-2-one

To a solution of 3,4,5,6,7,8-hexahydro-1*H*-quinoline-2-one (0.1 g, 0.66 mmol) in dioxane (10 mL) containing a catalytic quantity (40 µL) was added NaCNBH₃ (0.67 g). The reduction was left to stir for 20 hrs at r.t., and then diluted with EtOAc. The organic phase was treated with dilute HCl (3x), and the combined organics were washed with saturated HCO₃⁻, brine, dried over Na₂SO₄, filtered and concentrated to afford the mainly *trans* product (98 mg, 95%, 15-20% *cis* isomer). MS (+ES) m/z 254 (M+1)⁺.

-112-

Preparation 165

cis-3-(2-{[5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexyl)propionic acid methyl ester

5 The acid chloride (0.4 mmol) of preparation 157 was prepared as described previously, and added to a MeCl₂ solution (10 mL) of methyl 3-((1S,2S)-2-aminocyclohexyl)propanoate (222 mg, 1.2 eq) followed by Et₃N (133 μl, 0.96 mmol) and DMAP (10%) at r.t. Acylation was allowed to proceed for 18 hr. Crude product was
10 obtained by dilution of the reaction mixture with MeCl₂ and 0.1 N HCl, transferring to a sep funnel. The organic phase was then washed with sat'd bicarbonate, brine, then dried over Na₂SO₄. Filtration and concentration provided crude product, which was then purified using a Bond-Elut Si column (1 g, 1:1 hex/EtOAc) to generate pure methyl ester, (86 mg, 51%). MS (+ES) m/z 422.9/424.9 (M+H).

15

Preparation 166

cis-3-(2-{[5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexyl)propionic acid

 To a solution of a compound from preparation 165 (86 mg, 0.2 mmol) in THF (5
20 mL) was added 2 mL 0.5M aq. LiOH (5 eq), dropwise at r.t. Hydrolysis was complete at 2.5 hr. After dilution with water and EtOAc, the contents were transferred to a sep funnel, where addition of enough 1N HCl was added to maintain pH 2. The aqueous phase was back extracted 3 times with additional solvent, and the combined organics were then washed with brine, dried over Na₂SO₄. Filtration and concentration afforded clean acid,
25 80 mg (97%), which was used without further purification. MS (+ES) m/z 408.9/410.9 (M+H).

-113-

Preparation 167

5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carboxylic acid-{2-[2-(3,4,5-trimethoxyphenylcarbamoyl)ethyl]cyclohexyl}amide

5 To a compound from preparation 166 (40 mg, 0.1 mmol) in DMF (2 mL) was added catalytic amount of DMAP (10%, 1 mg), followed by EDC.HCl (20.6 mg, 0.11 mmol) and 3,4,5-trimethoxyaniline (20.1 mg, 0.11 mmol) at r.t. After 18 hr, the reaction was transferred to a sep funnel and diluted with EtOAc and 1N NaOH (aq. pH 9-10). The aqueous phase was extracted 3 times, and combined EtOAc fractions were then washed
10 sequentially with 1 N HCl, sat'd bicarbonate, brine, and then dried over Na₂SO₄. After filtration and concentration, crude amide was obtained (33 mg). Starting material (7 mg) was also recovered from the aqueous fraction upon acidification and standard workup. The crude product was then chromatographically purified (Bond-Elut Silica column, 1g, 2:1 EtOAc/hexanes) to afford clean amide (48 mg, 50%). MS (ES+) 573.9, 575.9
15 (M+H)⁺.

Preparation 168

trans-3-(2-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}cyclohexyl)propionic acid

20 A solution of methyl 3-(2-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)propanoate (0.127 g, 0.3 mmol) in THF (5 mL) was treated with a 0.5 M sol'n of LiOH in water (3 mL). The sol'n was stirred at r.t. for 5 hr. The reaction was diluted with water (3 mL) and washed with EtOAc (2 x 5 mL). The pH of
25 the aqueous was adjusted to ~3 with 0.1 M HCl and extracted with EtOAc (4 x 5 mL). The organic extractions were then washed with brine (2 x 5 mL), dried over sodium sulfate and the solvent removed to afford 0.095 g (80%) as a white solid. MS (ES+) m/z 409.0 (M+H)⁺, (ES-) m/z 407.0 (M-H)⁻.

-114-

Preparation 169

trans-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid {2-[2-(3,4,5-trimethoxyphenylcarbonyl)ethyl]cyclohexyl} amide

5 A sol'n of a compound from preparation 168 (0.095 g, 0.23 mmol) and EDC (0.115 g, 0.6 mmol) in DMF (25 mL) was treated with 3,4,5-trimethoxyaniline (0.11 g, 0.6 mmol). A catalytic amount of DMAP (4 mg, 0.03mmol) was also added. The reaction mixture was stirred at r.t. overnight. The reaction was diluted in EtOAc (100 mL) and washed with 5% citric acid sol (3 x 50 mL), water (2 x 50 mL), and dried over sodium sulfate. The solvent was removed *in vacuo* to afford a crude brown oil which was purified by a Varian Bond Elut SI column (10 g) with an eluting solvent of 1:1 Hexanes: EtOAc. The solvent was removed *in vacuo* to afford 0.072 g (54%) as an off white solid. MS (ES+) m/z 574.0 (M+H)⁺, (ES-) m/z 572.1 (M-H)⁻, 632.1 (M+CH₃COO)⁻.

15

Preparation 170

{(3S,1R)-3-[(*t*-butoxy)carbonylamino]cyclopentyl}-N-benzylcarboxamide

A mixture of N-BOC-1R, 3S-1 amino cyclopentane-3-carboxylic acid (229 mg, 1 mmol), EDC (286 mg, 1.5 mmol), benzyl amine (160 mg, 1.5 mmol) and DMAP (5 mg, catalytic) dissolved in DMF (20 mL) was stirred overnight at rt. The reaction mixture was diluted with ethyl acetate (200 mL) and washed with 0.2M HCl (2x50 mL), water (2x50 mL), brine (2x50 mL), dried over Na₂SO₄, filtered, and evaporated to yield a white solid (160 mg, 50%). ESMS: 319 (M+1)⁺.

25

Preparation 171

(1R,3S)-1-Amino-N'-phenylmethylcyclopentane-3-carboxamide

A compound from preparation 170 (100 mg, 0.31 mmol) was dissolved in TFA reagent (9.25 mL TFA, 0.25 mL anisole, 0.25 mL TIS, 0.25 mL water) and stirred at r.t for 30 m. TFA reagent was evaporated and the residue was dissolved in chloroform (10 mL) and filtered through SCX column and eluted with ammonia (10 mL, 2 M solution in

-115-

methanol) and evaporated to yield title compound (68 mg, quantitative). ESMS: 219 (M+1)⁺, 277 (M+Ac)⁺.

Preparation 172

5 {(3S,1R)-3-[N-benzylcarbonyl]cyclopentyl}[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

A compound from preparation 171 was dissolved in DMF (25 mL) and 2-chloro-6-fluorophenyl isoxazolyl chloride, triethylamine and DMAP were added and stirred
10 overnight at r.t. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with HCl (1M, 2x 50 mL), water (2x50 mL), brine (2x50 mL), dried over sodium sulfate, filtered, and evaporated to yield title compound (248 mg, 80%, crude). ESMS: 456 (M)⁺.

Preparation 173

15 (3S,1R)(3-Hydroxymethylcyclopentyl)carbamic acid *t*-butyl ester

(1R,3S)-3-*t*-Butoxycarbonylaminocyclopentanecarboxylic acid (229 mg, 1 mmol) was dissolved in THF (10 mL) and borane-THF (1.5 mL, 1M solution) was added dropwise at 0 °C and stirred overnight at r.t. The reaction mixture was poured into ice-cold
20 water (10 mL) and extracted with ethyl acetate (2 x 50 mL). The ethyl acetate extract was washed with brine, dried over sodium sulfate, filtered and evaporated to yield a semi solid (160 mg, 74%). ESMS (+): 216 (M+1)⁺, 250 (M+Cl)⁺.

Preparation 174

25 (3R,1S)-Toluene-4-sulfonic acid 3-*t*-butoxycarbonylaminocyclopentylmethyl ester

A compound from preparation 173 (150 mg, 0.69 mmol) was dissolved in methylene chloride (20 mL) and *p*-toluenesulfonyl chloride (131 mg, 0.69 mmol), triethylamine (268 μL, 1.86 mmol) and DMAP (10 mg) were added and stirred overnight.
30 The methylene chloride solution was washed with 1M HCl (2 x 10 mL), water (2 x 10 mL), brine (2 x 10 mL), dried over sodium sulfate, filtered and evaporated to yield the title compound (223 mg, 86%). ESMS: 370 (M+1)⁺, M+Ac)⁺.

-116-

Preparation 175(1R,3S)(3-Azidomethyl-cyclopentyl)-carbamic acid *t*-butyl ester

5 To a solution of a compound of preparation 174 (96 mg, 0.26 mmol) dissolved in DMF (10 mL) sodium azide was added and stirred at 80 °C overnight. The reaction mixture was diluted with ethyl acetate (50 mL) and washed with water (2 x 25 mL), brine (2 x 25 mL), dried over sodium sulfate, filtered and evaporated to yield the title compound (55 mg, 88% crude). HNMR (CDCl₃): 1.00-1.12 (m, 1H), 1.2-1.50 (m, 2H),
10 1.40 (s, 9H), 1.52-1.70 (m, 1H), 1.71 (1.82 (m, 1H), 1.89-2.00 (m, 1H), 2.10-2.30 (m, 2H), 3.26-3.28 (d, 2H), 4.80 (br S, 1H).

Preparation 176(1R,3S)(3-Aminomethylcyclopentyl)carbamic acid *t*-butyl ester

15

To a solution of a compound from preparation 175 (55 mg, 0.23 mmol) dissolved in methanol (10 mL) palladium/C (10 mg, 10%) was added and stirred under hydrogen (balloon) at for 2 h at rt. The solution was filtered off through Celite® and evaporated to yield a white solid (44 mg, 89%). ESMS: 215 (M+1)⁺.

20

Preparation 177(1R,3S)-[3-(Benzoylaminomethyl)cyclopentyl]carbamic acid *t*-butyl ester

To a solution of a compound from preparation 176 (44 mg, 0.2 mmol) dissolved in
25 DMF (10 mL), benzoyl chloride (35 mg, 0.25 mmol), triethylamine (0.5 mL) and DMAP (5 mg) was added and stirred overnight at rt. The reaction mixture was diluted with ethyl acetate (50 mL), washed with 1M HCl (2x20 mL), water (2 x 20 mL), brine (2 x 20 mL), dried over sodium sulfate, filtered and evaporated to get title compound (70 mg, crude quantitative). ESMS: 319 (M+1)⁺, 317 (M-1)⁻, 353 (M+Cl)⁻, 317 (M+Ac)⁻.

30

-117-

Preparation 178

(1R,3S)-N-(3-Aminocyclopentylmethyl)benzamide

5 A compound from preparation 177 (70 mg, 0.22 mmol) was dissolved in TFA (5 mL) and stirred for 1/2 h at rt. TFA was evaporated under vacuum and the residue was dissolved in chloroform (5 mL) passed through SCX column (Varian Bond Elut, 6cc) and eluted with 2M ammonia (methanol) to obtain title compound (34 mg, 72%). ESMS: 219 (M+1)⁺, 277 (M+Ac)⁻.

10

Preparation 179

(1R,3S)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid [3-(benzoylaminomethyl)cyclopentyl]amide

15 To a solution of a compound from preparation 178 (34 mg, 0.15 mmol) in DMF (5 mL), 2-chloro-6-fluorophenyl isoxazolyl chloride (85 mg, 0.3 mmol), triethylamine (1 mL), DMAP (5 mg) were added and stirred overnight. The reaction mixture was diluted with ethyl acetate (50 mL), washed with 1M HCl (2 x 20 mL), water (2 x 25 mL), brine (2x 20 mL), dried over sodium sulfate, filtered, evaporated and chromatographed (silica gel, 50% ethyl acetate in hexanes) to yield title compound (36 mg, 52%). ESMS: 456
20 (M)⁺, 514 (M+Ac)⁻.

Preparation 180

(1S,4R)(4-Hydroxymethyl-cyclopent-2-enyl)-carbamic acid *t*-butyl ester

25 To a solution of 4-*t*-butoxycarbonylaminocyclopent-2-enecarboxylic acid (1 g, 4.4 mmol) dissolved in THF (100 ml) and added with borane-THF (4.5 ml, 4.5 mmol) at 0 °C and stirred overnight at r.t. The reaction mixture was poured into ice-cold water (100 ml) and extracted with ethyl acetate. The ethyl acetate extract was washed with brine, dried over sodium sulfate and evaporated to obtain the title compound (655 mg, 69%). ESMS:
30 214 (M+1)⁺, 236 (M+23)⁺, 248 (M+35)⁻, 272 (M+59)⁻.

-118-

Preparation 181

(1S,4R)-Toluene-4-sulfonic acid 4-*t*-butoxycarbonylamino-cyclopent-2-enylmethyl ester

5 A solution of a compound from preparation 180 (655 mg, 3 mmol), *p*-tosyl chloride (760 mg, 4mmol), triethyl amine (0.4 ml) and DMAP (100 mg) dissolved in dichloromethane (50 ml) was stirred at rt, overnight. The reaction mixture was washed with dilute hydrochloric acid (2x), water (2x), brine (2x), dried over sodium sulfate and evaporated to obtain the title compound (1.0 g, 90%). ESMS: 368 (M+1)⁺, 426 (M+59)⁻.

10

Preparation 182

(1S,4R)(4-Azidomethylcyclopent-2-enyl)carbamic acid *t*-butyl ester

15 To a solution of a compound from preparation 181 (1g, 2.7 mmol) in DMF (50 mL), sodium azide (200 mg, 3 mmol) was added and stirred at 80 °C overnight. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with water (2x50 mL), brine (2x50 mL), dried over sodium sulfate and evaporated to yield the title compound (quantitative). IR (chloroform): 2096.6 cm⁻¹ (azide). ¹HNMR (CDCl₃): δ 1.27-1.30 (m, 2H), 1.43 (s, 9H), 2.51-2.56 (m, 2H), 3.27-3.35 (m, 2H), 5.76 (s, 2H).

20

Preparation 183

(1S,3R)-(3-Aminomethylcyclopentyl)carbamic acid *t*-butyl ester

25 To a solution of a compound from preparation 182 (300 mg, 1.25 mmol) in methanol (10 mL), Pd/C (10%, 50 mg) was added and the reaction mixture was hydrogenated (1 atm) overnight at rt. The reaction mixture was filtered through celite and evaporated to yield the title compound (222 mg, 89%). ESMS: 215 (M+1)⁺.

Preparation 184

(1S,3R)[3-(Benzoylaminoethyl)cyclopentyl]carbamic acid *t*-butyl ester

30

A mixture of a compound from preparation 183 (222 mg, 1.03 mmol), benzoyl chloride (210 mg, 1.5 mmol), triethylamine (1 mL), and DMAP (10 mg, catalytic) in

-119-

DMF (50 mL) was stirred overnight. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with dilute hydrochloric acid (1M, 2x50 mL), water (2x50 mL), brine (2x50 mL), dried over sodium sulfate, filtered and evaporated to yield the title compound (395 mg). ESMS: 319 (M+1)⁺, 377 (M+59)⁻.

5

Preparation 185

(1S,3R)-N-(3-Aminocyclopentylmethyl)benzamide

A compound from preparation 184 (crude 390 mg) was dissolved in trifluoroacetic acid (10 mL) and stirred for half an hour at rt. The reaction mixture was evaporated and chromatographed (Varian Bond Elut 60 cc, SCX column, 2M ammonia solution in methanol) to obtain the title compound (233 mg, 87%). ESMS: 219 (M+1)⁺.

10

Preparation 186

15 (1S,3R)-3-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carboxylic acid [3-(benzoylaminoethyl)cyclopentyl]amide

A mixture of a compound from preparation 185 (230 mg, 1.05 mmol), 3-(2-chloro, 6-fluorophenyl)-5-methyl-4-isoxazolyl chloride (328 mg, 1.2 mmol), triethyl amine (1mL), dimethyl amino pyridine (DMAP, 50 mg) dissolved in DMF (20mL) was stirred overnight at rt,. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with dilute HCl (1M, 2 x 50 mL), water (2 x 50 mL), brine (2 x 50 mL), dried over sodium sulfate, filtered, evaporated and then chromatographed (Varian Bond Elut, Si, 60 cc, 50% EtOAc in Hexane) to obtain the title compound (100 mg, 20%). ESMS: 456 (M)⁺, 515 (M+59)⁻.

20

25

Preparation 187

(1S,3R){3-[(4-Fluorobenzoylamino)methyl]cyclopentyl}carbamic acid *t*-butyl ester

30

To a solution of a compound from preparation 183 (68 mg, 0.31 mmol) and 4-fluorobenzoylchloride (60 mg, 0.38 mmol) dissolved in DMF (2 mL), triethyl amine (0.5 mL) and DMAP (10 mg) were added and stirred for four hours at rt. The reaction mixture

-120-

was diluted with ethyl acetate (20 mL) and washed with dilute hydrochloric acid (2x20 mL, 0.1 M), water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain 492323 (127 mg). ESMS: 337 (M+1)⁺, 395 (M+59)⁻.

5

Preparation 188

(1S,3R)-N-(3-Aminocyclopentylmethyl)-4-fluorobenzamide

A compound from preparation 187 (127 mg) was dissolved in trifluoroacetic acid (2 mL) and stirred for half an hour at rt. The reaction mixture was evaporated under vacuum and the residue was dissolved in methylene chloride, filtered through SCX column (Bond Elut, 60 cc) and eluted with ammonia (2M in methanol) to obtain the title compound (81 mg, 91%). ESMS: 237 (M+1)⁺, 295 (M+59)⁻.

10

Preparation 189

15 (1R,3S)-3-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carboxylic acid {3-[(4-fluorobenzoylamino)methyl]cyclopentyl}amide

A solution of a compound from preparation 188 (81 mg, 0.34 mmol), chloro fluoro phenyl isoxazolyl chloride (113 mg, 0.41 mmol), triethyl amine (0.5 mL) and DMAP (10 mg) in DMF (5 mL) was stirred overnight at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed with dilute hydrochloric acid (2x20 mL, 0.1 M), water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (80 mg, crude, 50%, used directly in the next step). ESMS: 474 (M)⁺, 475 (M+1)⁺, 533 (M+59)⁻.

20

25

Preparation 190

(1S,3R)(3-{[(Biphenyl-4-carbonyl)amino]methyl}-cyclopentyl)carbamic acid *t*-butyl ester

A solution of a compound from preparation 183 (68 mg, 0.31 mmol), 4-phenyl benzoyl chloride (82 mg, 0.38 mmol), triethyl amine (0.5 mL), and DMAP (10 mg) dissolved in DMF (2 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed with dilute hydrochloric acid (2x20 mL,

30

-121-

0.1 M), water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (170 mg, crude, used without purification).

ESMS:395 (M+1)⁺, 453 (M+59)⁻.

5

Preparation 191

(1S,3R)-Biphenyl-4-carboxylic acid (3-amino-cyclopentylmethyl)amide

A compound from preparation 190 (170 mg, crude) was dissolved in trifluoroacetic acid (2 mL) and stirred for half an hour. The reaction mixture was evaporated under vacuum and the residue was dissolved in methylene chloride and filtered through scx column (Bond Elut, 60 cc) and eluted with methanol-ammonia solution (2M) to obtain the title compound (63 mg). ESMS: 295 (M+1)⁺, 353 (M+59)⁻

10

Preparation 192

15

(1S,3R)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid (3-
{[(biphenyl-4-carbonyl)amino]methyl}-cyclopentyl)amide

20

A solution a compound from preparation 191 (63 mg, 1.21 mmol), 2-chloro-6-fluoro phenyl isoxazolyl chloride (88 mg, 0.32 mmol), triethyl amine (0.5 mL), and DMAP (10mg) dissolved in DMF (5 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed with dilute hydrochloric acid (2x20 mL, 0.1 M), water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (61 mg, 54%, used without purification). ESMS: 532 (M)⁺, 533 (M+1)⁺, 591 (M+59)⁻.

25

Preparation 193

(1S,3R)(3-{[(Pyridine-3-carbonyl)amino]methyl}cyclopentyl)-carbamic acid *t*-butyl ester

30

A solution of a compound from preparation 183 (68 mg, 0.31 mmol), nicotinoyl chloride (54 mg, 0.38 mmol), triethyl amine (0.5 mL), and DMAP (10 mg) dissolved in DMF (2 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed with water (2x20 mL), brine (2x20mL), dried over sodium

-122-

sulfate, filtered and evaporated to obtain the title compound (58 mg, 59%, used without purification). ESMS: 318 (M-1)⁻, 378 (M+59)⁺.

Preparation 194

5 (1S,3R)-N-(3-Aminocyclopentylmethyl)nicotinamide

A compound from preparation 193 (58 mg, crude) was dissolved in trifluoroacetic acid (2 mL) and stirred for half an hour. The reaction mixture was evaporated under vacuum and the residue was dissolved in methylene chloride and filtered through scx
10 column (Bond Elut, 60 cc) and eluted with methanol-ammonia solution (2M) to obtain the title compound (quantitative). ESMS: 220 (M+1)⁺, 278 (M+59)⁻.

Preparation 195

15 (1S,3R)-N-(3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}cyclopentylmethyl)-nicotinamide

A solution of a compound from preparation 194 (112 mg, 0.51 mmol), 2-chloro-6-fluoro phenyl isoxazolyl chloride (210 mg, 0.76 mmol), triethyl amine (0.5 mL), and DMAP (10mg) dissolved in DMF (5 mL) was stirred for four hours at rt. The reaction
20 mixture was diluted with ethyl acetate (20 mL) and washed water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (104 mg, 28%, used without purification). ESMS: 455 (M-1)⁻, 515 (M+59)⁻.

Preparation 196

25 (1S,3R)(3-{[(Furan-2-carbonyl)amino]methyl}cyclopentyl)-carbamic acid *t*-butyl ester

A solution of a compound from preparation 183 (68 mg, 0.31 mmol), 2-furoyl chloride (50 mg, 0.38 mmol), triethyl amine (0.5 mL), and DMAP (5 mg) dissolved in DMF (2 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl
30 acetate (20 mL) and washed with water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (105 mg, crude, used without purification) ESMS: 309 (M+1)⁺, 343 (M+35)⁻.

-123-

Preparation 197

(1S,3R)-Furan-2-carboxylic acid (3-aminocyclopentylmethyl)-amide

5 A compound from preparation 196 (105 mg, crude) was dissolved in trifluoroacetic acid (2 mL) and stirred for half an hour. The reaction mixture was evaporated under vacuum and the residue was dissolved in methylene chloride and filtered through SCX (Bond Elut, 60 cc) column and eluted with methanol-ammonia solution to obtain the title compound (94 mg, quantitative). ESMS: 209 (M+1)⁺.

10

Preparation 198

(1S,3R)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid (3-[(furan-2-carbonyl)amino]methyl)-cyclopentyl)amide

15 A solution of a compound from preparation 197 (94 mg, 0.51 mmol), 2-chloro-6-fluoro phenyl isoxazolyl chloride (185 mg, 0.67 mmol), triethyl amine (0.5 mL), and DMAP (10mg) dissolved in DMF (5 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound
20 (147 mg, crude, quantitative, used without purification). ESMS: 446 (M+1)⁺, 504 (M+59)⁻.

Preparation 199(1S,3R){3-[(3,4,5-Trimethoxybenzoylamino)methyl]cyclopentyl}carbamic acid *t*-butyl
25 ester

A solution of a compound from preparation 183 (68 mg, 0.31 mmol), 3,4,5-trimethoxy benzoyl chloride (88 mg, 0.38 mmol), triethyl amine (0.5 mL), and DMAP (10 mg) dissolved in DMF (2 mL) was stirred for four hours at rt. The reaction mixture was
30 diluted with ethyl acetate (20 mL) and washed with water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (130 mg, crude, used without purification). ESMS:409 (M+1)⁺, 443 (M+35)⁻.

-124-

Preparation 200

(1S,3R)-N-(3-Aminocyclopentylmethyl)-3,4,5-trimethoxybenzamide

5 A compound from preparation 199 (130 mg, crude) was dissolved in trifluoroacetic acid (2 mL) and stirred for half an hour at rt. The reaction mixture was evaporated under vacuum and the residue was dissolved in methylene chloride and filtered through SCX (Bond Elut, 60 cc) column and eluted with methanol-ammonia solution (2M) to obtain the title compound (82 mg, 83%). ESMS: 309 (M+1)⁺, 367
10 (M+59)⁻.

Preparation 201

(1S,3R)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid {3-[(3,4,5-trimethoxybenzoylamino)methyl]cyclopentyl} amide

15

A solution of a compound from preparation 200 (82 mg, 0.26 mmol), 2-chloro-6-fluoro phenyl isoxazolyl chloride (109 mg, 0.39 mmol), triethylamine (0.5 mL), and DMAP (10mg) dissolved in DMF (5 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed water (2x20 mL), brine
20 (2x20mL), dried over sodium sulfate, filtered and evaporated to obtain the title compound (89 mg, 62%, used without further purification). ESMS: 546 (M+1)⁺, 604 (M+59)⁻.

Preparation 202

(1S,3R)[3-(Benzyloxycarbonylamino)methyl]cyclopentyl]carbamic acid *t*-butyl ester

25

A solution of a compound from preparation 183 (1.3 g, 6 mmol), CBZ chloride (1.2 g, 7 mmol), triethyl amine (1 mL), and DMAP (50 mg) dissolved in DCM (50 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed with water (2x20 mL), brine (2x20mL), dried over sodium sulfate,
30 filtered and evaporated to obtain the title compound (2.0 g, crude). ESMS: 349 (M+1)⁺, 407 (M+59)⁻.

-125-

Preparation 203

(1S,3R)(3-Aminocyclopentylmethyl)carbamic acid benzyl ester

5 A compound from preparation 202 (2.00 g, crude) was dissolved in trifluoroacetic acid (10 mL) and DCM (10 mL) and stirred for half an hour at rt. The reaction mixture was evaporated under vacuum and the residue was dissolved in DCM and filtered through SCX (Bondesil, 20 g) column and eluted with methanol-ammonia solution (2M) to obtain the title compound (550 mg, 38%). ESMS: 249 (M+1)⁺.

10

Preparation 204

(1S,3R)(3-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}cyclopentylmethyl)carbamic acid benzyl ester

15 A solution of a compound from preparation 203 (500 mg, 2 mmol), 2-chloro-6-fluorophenyl isoxazoyl chloride (600 mg, 2.2 mmol), triethylamine (1 mL), and DMAP (100 mg) dissolved in DMF (10 mL) was stirred for four hours at rt. The reaction mixture was diluted with ethyl acetate (20 mL) and washed water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered, evaporated and chromatographed (Si column, 12x2.5 inches, gradient -10% to 50% ethyl acetate in hexanes) to obtain the title compound (355
20 mg, 36%). ESMS: 486 (M+1)⁺, 544 (M+59)⁻.

Preparation 205

25 (1S,3R)(3-{[2-(3,4,5-Trimethoxyphenyl)acetylamino]methyl}-cyclopentyl)carbamic acid
t-butyl ester

A solution of a compound from preparation 183 (1 g, 4.6 mmol), 3,4,5-trimethoxy phenylacetic acid (1.5 g, 6.9 mmol), EDC (1.3 g, 6.9 mmol) and DMAP (100 mg) dissolved in DCM (20 mL) was stirred overnight at rt. The reaction mixture was washed
30 with water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered, and evaporated to obtain the title compound (1.5 g, crude). ESMS: 423 (M+1)⁺, 481 (M+59)⁻.

-126-

Preparation 206

(1S,3R)-N-(3-Aminocyclopentylmethyl)-2-(3,4,5-trimethoxy-phenyl)acetamide

5 A compound from preparation 205 (1.5 g) was dissolved and stirred in a solution of hydrogen chloride in acetic acid (20 mL, 1M) for half an hour at rt. The reaction mixture was evaporated and the residue was triturated with ether to obtain the title compound (1.1 g, used without further purification). ESMS: 323(M+1)⁺, 357 (M+35)⁻.

Preparation 207

10 (1S,3R)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid (3-{[2-(3,4,5-trimethoxyphenyl)acetyl-amino]methyl}cyclopentyl)amide

15 A solution of a compound from preparation 206 (1 g, 3.1 mmol), 2-chloro-6-fluoro phenyl isoxazolyl chloride (850 mg, 3.1mmol), triethylamine (1 mL), and DMAP (100 mg) dissolved in DCM (20 mL) was stirred for four hours at rt. The reaction mixture was washed with water (2x20 mL), brine (2x20mL), dried over sodium sulfate, filtered, evaporated and chromatographed (Si column, 15x6 cm, 50% ethyl acetate in hexane) to obtain the title compound (1.2 g, 70%). ESMS: 486 (M+1)⁺, 544 (M+59)⁻.

20

Preparation 208

(1S,3R)[3-({[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}methyl)cyclopentyl]carbamic acid *t*-butyl ester

25 A mixture of a compound from preparation 183 (215 mg, 1mmol), 2-chloro, 6-fluoro phenyl isoxazolyl chloride (300 mg, 1.1 mmol), triethyl amine (0.1 mL) and DMAP (20 mg) dissolved in DMF (5 mL) was stirred overnight at r.t. The reaction mixture was diluted with ethyl acetate (25 mL) and washed with HCl (1M, 2x 10 mL), water (2x10 mL), brine (2x10 mL), dried over sodium sulfate, filtered, and evaporated to yield the title compound (400 mg, 88%, crude). ESMS: 452 (M+1)⁺.

30

-127-

Preparation 209

[3-({[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}methyl)cyclopentyl]carbamic acid *t*-butyl ester

- 5 A mixture of a compound from preparation 176 (640 mg, 2.9 mmol), 2-chloro, 6-fluoro phenyl isoxazolyl chloride (1.22 g, 4.4 mmol), triethyl amine (0.5 mL) and DMAP (50 mg) dissolved in DCM (50 mL) was stirred overnight at rt. The reaction mixture was washed with HCl (1M, 2x 10 mL), water (2x10 mL), brine (2x10 mL), dried over sodium sulfate, filtered, and evaporated to yield the title compound (1.5 g, crude, quantitative).
- 10 ESMS: 452 (M+1)⁺.

Preparation 210

N-(3-aminocyclohexyl)(*t*-butoxy)carboxamide

- 15 To 1,3 diaminocyclohexyl (20.0g, 175 mmol) in 400 mL CHCl₃ at 0 °C was added di-*t*-butyldicarbonate (7.65 g, 35.0 mmol), which was dissolved in 100 mL CHCl₃, poured rapidly. The ice bath was removed and after 0.5 h, the reaction mixture was washed with saturated NaHCO₃ (3x150 mL) and washed with brine (2x150 mL). The organics were concentrated to give (*cis* and *trans*, 2:1) a crude clear oil (7.50 g, 100%). The crude
- 20 product was used without purification.

Preparation 211 and 212

- N-((1S,3S)-3-{{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl})(*t*-butoxy) carboxamide &
- 25 N-((3S,1R)-3-{{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl})(*t*-butoxy) carboxamide

- To the crude oil from preparation 210 (7.50 g, 35.0 mmol) in 75 mL anhydrous CH₂Cl₂ at 0 °C was added Et₃N (7.32 mL, 52.5 mmol) followed by 3-(6-chloro-2-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (10.01 g, 38.5 mmol). After 5 min.
- 30 the ice bath was removed and the reaction stirred for an additional 25 minutes. The reaction was diluted with 300 mL CH₂Cl₂ and washed with saturated NaHCO₃ (100 mL).

-128-

The organics were dried over Na₂SO₄, filtered and concentrated until a precipitate formed. The precipitate was filtered and the filtrate was concentrated a little more to give more precipitate, which was filtered. A total of 8.5g (54%) of preparation 209 was obtained. The filtrate was concentrated and chromatographed with 20-30% EtOAc in hexanes to
5 give 3.78 g (24%) of preparation 210.

ESIMS *m/e* 451 ³⁵Cl (M⁺+1) and 453 ³⁷Cl (M⁺+1).

ESIMS *m/e* 452 ³⁵Cl (M⁺+1) and 454 ³⁷Cl (M⁺+1).

Preparation 213

10 *Cis*-3-(amino)-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane
hydrochloride

To Example 495 (0.500 g, 1.16 mmol) at 0 °C was added 4M HCl in dioxane (5 mL) to give a suspension. The reaction was sonicated at r.t. for 2 min., then stirred at r.t.
15 for 45 minutes. A yellow solid was obtained (0.411 g, 96%). ESIMS *m/e* 332 ³⁵Cl (M⁺+1) and 334 ³⁷Cl (M⁺+1).

Preparation 214

d,l-trans-Octahydroquinolin-2-one

20

Following the literature procedure of Morzycki et al (Heterocycles, (1995), 41
(12), 2729-2736) 3,4,5,6,7,8-hexahydro-1*H*-quinolin-2-one (100 mg, 0.66 mmol) in
dioxane (10 ml) with trifluoroacetic acid (0.042 ml, catalytic) was treated with NaCNBH₃
(0.67 g, 10.8 mmol), and stirred for 18 hr at RT. The mixture was then diluted with
25 EtOAc and washed with dilute HCl (0.1 N). Additional EtOAc extractions (2x) were
then combined with the first, and then treated with sat'd bicarbonate, brine, dried over
Na₂SO₄, filtered and concentrated to afford the mainly trans product (98 mg, ~5:1).
MS (+ES) 154 (M+H)+.

-129-

Preparation 2153-(2-Amino-*trans*-cyclohexyl)propionic acid methyl ester hydrochloride

5 A compound from preparation 214 (1 g, 6.4 mmol) was dissolved in methanol (100 mL) and conc. HCl (10 ML) and refluxed for 20 h. The solvents were removed under vacuum to obtain the title compound (1.05 g, 88%). ESMS: 186 (M)⁺, 187 (M+1)⁺.

Preparation 216

10 3-(2-{{3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl}amino}*trans*-cyclohexyl)propionic acid methyl ester

15 To a solution of a compound from preparation 215 (1g, 4.5 mmol) dissolved in DMF (50 mL), was added 2-chloro-6-fluorophenyl isoxazolyl chloride (1.23 g, 4.5 mmol), triethyl amine (10 mL) and DMAP (100 mg) and stirred overnight at rt. The reaction mixture was diluted with ethyl acetate (100 mL) and washed with 1M HCl (2x 50 mL), water (2x50 mL), brine (2x50 mL), dried over sodium sulfate, filtered and evaporated to yield 1.9 g. ESMS: 423 (M+1)⁺, 481 (M+Ac)⁻.

Preparations 217 and 218

20 Separation of Preparation 216 into enantiomers

25 Preparative separation into enantiomers was achieved by column chromatography over a Chiralpak AD column using 40% isopropyl alcohol in heptane and 0.2% dimethylethylamine. Thus, the first enantiomer that eluted was designated preparation 217 (Isomer 1), and the second, preparation 218 (Isomer 2).

Preparation 219

30 3-(2-{{3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl}amino}*trans*-cyclohexyl)propionic acid

A solution from preparation 217 (0.127 g, 0.3 mmol) in THF (5 mL) was treated with a 0.5 M soln of LiOH in water (3 mL). The sol'n was stirred at r.t. for 5 hr. The

-130-

reaction was diluted with water (3 mL) and washed with EtOAc (2 x 5 mL). The pH of the aqueous was adjusted to ~3 with 0.1 M HCl and extracted with EtOAc (4 x 5 mL). The organic extractions were then washed with brine (2 x 5 mL), dried over sodium sulfate and the solvent removed to afford 0.095 g (80%) as a white solid. MS (ES+) m/z 409.0 (M+H)⁺, (ES-) m/z 407.0 (M-H)⁻.

Preparation 220

3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid {2-[2-(3,4,5-trimethoxyphenylcarbamoyl)ethyl]*trans*-cyclohexyl}amide

10

A sol'n from preparation 219 (0.095 g, 0.23 mmol) and EDC (0.115 g, 0.6 mmol) in DMF (25 mL) was treated with 3,4,5 trimethoxyaniline (0.11 g, 0.6 mmol). A catalytic amount of DMAP (4 mg, 0.03 mmol) was also added. The reaction mixture was stirred at r.t. overnight. The reaction was diluted in EtOAc (100 mL) and washed with 5% citric acid sol'n (3 x 50 mL), water (2 x 50 mL), and dried over sodium sulfate. The solvent was removed *in vacuo* to afford a crude brown oil which was purified by a Varian Bond Elut SI column (10 g) with an eluting solvent of 1:1 hexanes:EtOAc. The solvent was removed *in vacuo* to afford 0.072 g (54%) as an off white solid. MS (ES+) m/z 574.0 (M+H)⁺, (ES-) m/z 572.1 (M-H)⁻, 632.1 (M⁺CH₃COO⁻)⁻.

15
20

Preparation 221

d,l-cis-Octahydroquinolin-2-one

2-Nitrocinnamic acid (17 g, 82 mmol) was dissolved in acetic acid (120 ml) and treated with PtO₂ (10 g) and H₂ (3 atm) for 24 hr at 60 °C. The crude reaction was then cooled to r.t., filtered and concentrated. The material was then diluted with EtOAc, and sat'd bicarbonate. After reextraction of the aqueous phase with EtOAc (2x), the combined organics were washed with brine, dried over Na₂SO₄, filtered and concentrated to afford a 13:1 mixture of *cis/trans* product (10 g, 80%). Clean *cis* isomer eventually crystallized from residual solvent. MS (+ES) 154 (M+H)⁺.

25
30

-131-

Preparation 2223-(2-Amino-*cis*-cyclohexyl)propionic acid methyl ester hydrochloride

5 A solution of a compound from preparation 221 (4.4 mg, 28.9 mmol) in methanol (50 mL) was treated with conc. HCl (10 mL) and refluxed overnight. The reaction mixture was evaporated to yield the title compound (7g, crude, quantitative). ESMS+: 186 (M+1).

Preparation 223

10 3-(2-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}-*cis*-cyclohexyl)propionic acid methyl ester

15 A compound from preparation 222 (7g, 0.038 mmol) was dissolved in CH₂Cl₂ (100 mL) and treated with methyl isoxazolyl chloride (11 g, 0.04 mmol) in presence of triethylamine (20 mL) at room temperature for 4 h. The reaction mixture was washed with 1 N HCl (2 x 100 mL), water (2 x 100 mL), brine (2 x 100 mL), dried over sodium sulfate, filtered and evaporated to give the title compound (crude gum, 10 g, 62 %). ESMS+: 430 (M), 424 (M+1).

Preparation 224

20 3-(2-{[3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl]amino}-*cis*-cyclohexyl)propionic acid

25 A solution of a compound from preparation 223 (8.5 g, 20 mmol) dissolved in THF (200 mL) and LiOH (100 mL, 0.5 M solution) was stirred for 2 h at r.t. The reaction was diluted with water (50 mL) and washed with EtOAc (2 x 100 mL). The alkaline aqueous solution was acidified to pH 1 with conc. HCl and extracted with EtOAc (3 x 200 mL). The organic extract was washed with brine, dried over sodium sulfate, filtered and evaporated to yield a white solid (8 g, 97%). ESMS+: 409 (M+1).

-132-

Preparation 225

3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid {2-[2-(3,4,5-trimethoxyphenylcarbamoyl)ethyl]-*cis*-cyclohexyl} amide

5 A solution of a compound from preparation 224 (1 g, 2.4 mmol), EDC (955 mg, 5 mmol), trimethoxyaniline (915 mg, 5 mmol), and DMAP (50 mg, catalytic) in DMF (50 mL) was stirred at r.t. overnight. The reaction mixture was diluted with EtOAc (200 mL) and washed with dil. HCL (1 M, 3 x 100 mL), water (3 x 100 mL), brine (2 x 100 mL), dried over sodium sulfate, filtered and evaporated to give the title compound (1.3 g, 97%).
10 ESMS +: 574 (M), 576 (M+2).

Preparation 227

3-(2-{[5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carbonyl]amino}cyclohexyl)propionic acid methyl ester

15

5-(6-Chloro-2-fluorophenyl)-3-methylisoxazole-4-carboxylic acid (0.4 mmol) was prepared as described previously, and added to a MeCl₂ solution (10 ml) of a compound from preparation 222 (222 mg, 1.2 eq) followed by Et₃N (133 ml, 0.96 mmol) and DMAP (10%) at r.t. Acylation was allowed to proceed for 18 h. Crude product was obtained by
20 dilution of the reaction mixture with MeCl₂ and 0.1 N HCl, transferring to a separatory funnel. The organic phase was then washed with sat'd bicarbonate, brine, then dried over Na₂SO₄. Filtration and concentration provided crude product, which was then purified using a Bond-Elut Si column (1 g, 1:1 hex/EtOAc) to generate the title compound (86 mg, 51%). MS (+ES) m/z 422.9/424.9 (M+H).

25

Preparation 227

3-(2-{[5-(2-Chloro-6-fluoro-phenyl)-3-methyl-isoxazole-4-carbonyl]-amino}-cyclohexyl)-propionic acid

30

To a solution of the compound from 226 (86 mg, 0.2 mmol) in THF (5 ml) was added 2 ml 0.5M aq. LiOH (5 eq), dropwise at r.t. Hydrolysis was complete at 2.5 hr. After dilution with water and EtOAc, the contents were transferred to a separatory funnel,

-133-

where addition of enough 1N HCl was added to maintain pH 2. The aqueous phase was back extracted 3 times with additional solvent, and the combined organics were then washed with brine, dried over Na₂SO₄. Filtration and concentration afforded clean acid, 80 mg (97%), which was used without further purification. MS (+ES) m/z 408.9/410.9 (M+H).

Preparation 228

5-(2-Chloro-6-fluorophenyl)-3-methylisoxazole-4-carboxylic acid-{2-[2-(3,4,5-trimethoxyphenylcarbamoyl)ethyl]-cyclohexyl}amide

10

To the acid from preparation 227 (40 mg, 0.1 mmol) in DMF (2 ml) was added catalytic amount of DMAP (10%, 1 mg), followed by EDC.HCl (20.6 mg, 0.11 mmol) and 3,4,5-trimethoxyaniline (20.1 mg, 0.11 mmol) at r.t. After 18 hr, the reaction was transferred to a separatory funnel and diluted with EtOAc and 1N NaOH (aq. pH 9-10). The aqueous phase was extracted 3 times, and combined EtOAc fractions were then washed sequentially with 1 N HCl, sat'd bicarbonate, brine, and then dried over Na₂SO₄. After filtration and concentration, crude amide was obtained (33 mg). Starting material (7 mg) was also recovered from the aqueous fraction upon acidification and standard work-up. The crude product was then chromatographically purified (Bond-Elut Silica column, 1g, 2:1 EtOAc/hexanes) to afford clean amide (48 mg, 50%). MS (ES+) 573.9, 575.9 (M+H)+.

15
20

Preparation 229

Methyl 3-(2-aminocyclohexyl)propanoate hydrochloride

25

A solution of a compound from preparation 163 (4.4 mg, 28.9 mmol) in methanol (50 mL) was treated with conc. HCl (10 mL) and refluxed overnight. The reaction mixture was evaporated to yield the title compound (7g, crude, quantitative). ESMS⁺: 186 (M+1).

-134-

Preparation 230

Methyl 3-(2-{[3-(6-chloro-2-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)propanoate

5 A compound from preparation 229 (7g, 0.038 mmol) was dissolved in methylene chloride (100 mL) and treated with methyl isoxazolyl chloride (11 g, 0.04 mmol) in presence of triethylamine (20 mL) at room temperature for four hours. The reaction mixture was washed with 1 N HCl (2 x 100 mL), water (2 x 100 mL), brine (2 x 100 mL), dried over sodium sulfate, filtered and evaporated to give the title compound (crude gum,
10 10 g, 62 %). ESMS+: 430 (M), 424 (M+1).

Preparation 231

3-(2-{[3-(6-chloro-2-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)propanoic acid

15 A solution of the ester from preparation 230 (8.5 g, 20 mmol) dissolved in THF (200 mL) and LiOH (100 mL, 0.5 M solution) was stirred for 2 h at rt. LiOH solution was diluted with water (50 mL) and washed with ethyl acetate (2 x 100 mL). The alkaline aqueous solution was acidified to pH 1 with conc. HCl and extracted with ethyl acetate (3
20 x 200 mL). The ethyl acetate extract was washed with brine, dried over sodium sulfate, filtered and evaporated to yield a white solid (8 g, 97%). ESMS+: 409 (M+1).

Preparation 232

25 3-(2-{[3-(6-chloro-2-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino}cyclohexyl)-N-(3,4,5-trimethoxyphenyl)propanamide

A solution of acid from preparation 231 (1 g, 2.4 mmol), EDC (955 mg, 5 mmol), trimethoxyaniline (915 mg, 5 mmol), and DMAP (50 mg, catalytic) in DMF (50 mL) was stirred at r.t. overnight. The reaction mixture was diluted with ethyl acetate (200 mL) and
30 washed with dil. HCL (1 M, 3 x 100 mL), water (3 x 100 mL), brine (2 x 100 mL), dried over sodium sulfate, filtered and evaporated to give the title compound (1.3 g, 97%). ESMS +: 574 (M), 576 (M+2).

-135-

Preparation 233

6-Fluoropyridine-2-carboxylic acid

To a heterogeneous solution of 2-fluoro-6-methyl-pyridine (9.65 g, 86.8 mmol) in water (400 mL) was added potassium permanganate (31.6 g, 200 mmol), and the reaction was heated to approximately 100°C. After 30 minutes, additional potassium permanganate (16.5 g, 104 mmol) was added, and the reaction heated at 100°C overnight. The reaction was filtered through celite to remove manganese salts. The mother liquor was washed with ethyl ether (200 mL x3), neutralized to pH 7, and concentrated down to approximately 100 mL total volume. The aqueous solution was acidified to pH 2 with concentrated HCl and extracted with ethyl acetate, followed by 20% i-PrOH/CHCl₃ (x3). The combined organic layers were dried over MgSO₄ and concentrated. The solids were suspended in chloroform, sonicated, then filtered to remove remaining side products. The mother liquor was concentrated to give 5.47 g as a white crystalline solid, 45% yield. ¹H NMR: consistent with structure. MS (ion spray) 140 (M).

Preparation 234

6-Methoxy-pyridine-2-carboxylic acid

To 6-hydroxypicolinic acid (1.39 g, 10.0 mmol) in toluene (35 mL) was added silver oxide (2.43 g, 10.5 mmol) and the mixture was stirred for 30 minutes. To the reaction was added iodomethane (1.31 mL, 21.0 mmol), and the reaction was heated to reflux overnight. The reaction was filtered over celite and concentrated to give 1.45 g of the methyl ester as a yellow solid. The resulting solid was dissolved in tetrahydrofuran (50 mL), and added to a solution of 5N NaOH (20 mL, 100 mmol) and water (5 mL). After one hour, the reaction was acidified to pH 3 with 5N HCl and concentrated. To the resulting white solids was added 20% MeOH/CHCl₃ and the mixture was sonicated for 20 minutes. The mixture was filtered, and the mother liquor was dried over MgSO₄ and concentrated to give 1.33 g of the title compound as a white solid, 87% yield. ¹H NMR: consistent with structure. MS (ion spray) 152 (M).

-136-

Preparation 235

5-Methoxy-nicotinic acid

To sodium hydride (0.42 g, 10.5 mmol, washed with hexanes x3) in
5 dimethylformamide (15 mL) was added 5-hydroxy-nicotinic acid methyl ester (1.53 g,
10.0 mmol) in dimethylformamide (10 mL). After three minutes, iodomethane (0.65 mL,
10.5 mmol) was added dropwise while stirring at room temperature. After 1h, the
reaction was quenched with methanol and concentrated to a brown oil. The solution was
dissolved in 20% isopropanol/chloroform, washed with saturated aqueous sodium
10 bicarbonate solution, brine (x3), dried over magnesium sulfate and concentrated to give
1.46 g of 6-methoxynicotinic acid methyl ester. The resulting oil was dissolved in
tetrahydrofuran (50 mL), and added to a solution of 5N NaOH (20 mL, 100 mmol) and
water (5 mL). After 30 minutes, an additional 25 mL of 5N NaOH was added to the
solution. After one hour, the reaction was acidified to pH 3 with 5N HCl and
15 concentrated. To the resulting white solids was added 20% MeOH/CHCl₃ and the
mixture was sonicated for 20 minutes. The mixture was filtered, and the mother liquor
was dried over MgSO₄ and concentrated to give 1.38 g as a light yellow solid, 90% yield.
¹H NMR: consistent with structure. MS (ion spray) 152 (M⁻).

20

Preparation 236N-{3-[3-(1-Aminoethylidene)-5-chloro-2,4-dioxo-3,4-dihydro-2H-quinolin-1-yl]-
cyclohexylmethyl}benzamide

A compound from Example 125 (0.05 g, 0.1 mmol) and Mo(CO)₆ (0.3 g, 1.1
25 mmol) were combined in a solution of ACN (5 mL) and water (1 mL). The reaction
mixture was heated to 60°C while stirring. After 3 hr of stirring the reaction was
complete. The reaction was concentrated to a dark brown solid under vacuum. The solid
was diluted in CH₂Cl₂ (1 mL) and purified by passing through a Varian Bond Elut SI
column (5 g). The product was eluted with 2% MeOH in CH₂Cl₂. The solvent was
30 removed *in vacuo* to afford 0.041 g (82%) of the title compound as a light brown solid.
MS (ES⁺) m/z 452.0 (M+H)⁺, (ES⁻) 450.0 (M-H)⁻.

-137-

Preparation 237

{3-[3-(1-Aminoethylidene)-5-chloro-2,4-dioxo-3,4-dihydro-2H-quinolin-1-yl]-cyclohexylmethyl}carbamic acid benzyl ester

5 A compound from Example 498 (0.25 g, 0.5 mmol) and Mo(CO)₆ (1.35 g, 5.2 mmol) were combined in a solution of ACN (25 mL) and water (5 mL). The reaction mixture was heated to 60°C while stirring. After stirring overnight the reaction was complete. The reaction was concentrated to a dark brown solid under vacuum. The solid was diluted in CH₂Cl₂ (10 mL) and the solid material was filtered by passing through
10 Celite. The yellow filtrate was purified by silica gel column chromatography using 50% EtOAc in CH₂Cl₂ to elute the product. The solvent was removed *in vacuo* to afford 0.24 g (98%) of the title compound as a light yellow solid. MS (ES+) m/z 481.9 (M+H)⁺, (ES-) 479.9 (M-H)⁻.

15

Preparation 238

N-{3-[3-(1-Aminoethylidene)-5-chloro-2,4-dioxo-3,4-dihydro-2H-quinolin-1-yl]-cyclohexylmethyl}nicotinamide

A compound from Example 126 (0.02 g, 0.04 mmol) and Mo(CO)₆ (0.13
20 g, 0.5 mmol) were combined in a solution of ACN (5 mL) and water (1 mL). The reaction mixture was heated to 60°C and stirred for 2 hr. The reaction was concentrated to a dark brown solid under vacuum. The solid was diluted in CH₂Cl₂ (10 mL) and purified by passing directly through a Bond-Elut cation exchange column. The product was eluted with 2M Ammonia in MeOH. The brown liquid was filtered using a Gelman Nylon
25 Acrodisc to afford a yellow solution. The solvent was removed *in vacuo* to afford 0.018 g (90%) of the title compound as a white solid. MS (ES+) m/z 452.9 (M+H)⁺, (ES-) m/z 450.8 (M-H)⁻, 510.9 (M+CH₃COO)⁻.

-138-

Preparation 239

N-{3-[3-(1-Aminoethylidene)-5-chloro-2,4-dioxo-3,4-dihydro-2H-quinolin-1-yl]-
cyclohexylmethyl}-6-fluoronicotinamide

5 A compound from Example 149 (0.05 g, 0.1 mmol) and Mo(CO)₆ (0.3 g, 1.1 mmol) were combined in a solution of ACN (5 mL) and water (1 mL). The reaction mixture was heated to 60°C while stirring. After 3 hr of stirring the reaction was complete. The reaction was concentrated to a dark brown solid under vacuum. The solid was diluted in CH₂Cl₂ (1 mL) and purified by passing through a Varian Bond Elut SI
10 column (5 g). The product was eluted with 2% MeOH in CH₂Cl₂. The solvent was removed *in vacuo* to afford 0.041 g (82%) of the title compound as a light brown solid. MS (ES+) m/z 452.0 (M+H)⁺, (ES-) 450.0 (M-H)⁻.

Preparation 240

15 N-{3-[5-Chloro-3-(1-hydroxy-ethylidene)-2,4-dioxo-3,4-dihydro-2H-quinolin-1-yl]-
cyclohexylmethyl}benzamide

A compound from preparation 236 (0.05 g, 0.11 mmol) was stirred in a solution of acetonitrile (10 mL) and water (1 mL). The solution was treated with p-toluenesulfonic acid (0.01 g, 0.05 mmol). The reaction was heated to reflux and stirred overnight. The
20 reaction was concentrated to a solid and diluted in EtOAc (50 mL). The organic was washed with sat'd sodium bicarbonate (2 x 10 mL), brine (2 x 10 mL), and dried over sodium sulfate. The solvent was removed to yield a crude solid that was purified using a chromatotron with a silica gel plate. The product was eluted with 50% EtOAc in
25 CH₂Cl₂. The solvent was removed to afford 0.032 g (64%) of the title compound as an off-white solid. MS (ES+) m/z 452.9 (M+H)⁺, (ES-) 450.9 (M-H)⁻.

Preparation 241

(3-Aminocyclohexylmethyl)carbamic acid benzyl ester

30

S-Amino enantiomer of a compound from preparation 104 (0.28 g, 0.77 mmol) was treated with HCl in HOAc (1.0M, 2 mL) under N₂. After 20 min of stirring at r.t., the

-139-

reaction was complete. The crude was then concentrated to a solid using CAN to azeotrope the solvent. The white solid was taken up in ether and filtered. The white solid weighed 0.19 g (95%) and was in good purity. MS (ES+) m/z 263.0 (M+H)+.

5

Preparation 242*t*-Butylaminophenylacetic acid methyl ester

t-Butylamine (9.30 mL, 88.0 mmol) was combined with α -bromophenylacetic acid (5.0 g, 22.0 mmol) and triethylamine (3.40 mL, 22.0 mmol) in tetrahydrofuran (200 mL) and the mixture refluxed overnight. The mixture was then concentrated *in vacuo* and the residue taken up in 20% isopropanol/chloroform and washed 1 x aqueous NaHCO₃, x2 brine, and dried over sodium sulfate. Concentration left a residue which was loaded onto a silica gel column and eluted with methanol/chloroform to yield pure amine (1.34 g, 28%) as an oil. MS(ES): (M+1)⁺ 222.2, 223.2 m/z.

15

Preparation 243*t*-Butylaminophenylacetic acid

A compound from preparation 242 (1.32 g, 6.0 mmol) was combined with aqueous 2N sodium hydroxide (20 mL), tetrahydrofuran (5 mL), and ethanol (5 mL) and the mixture stirred overnight at ambient temperature. The mixture was then adjusted to approx. pH 2.5 with aqueous hydrochloric acid and concentrated to dryness *in vacuo*. The resulting solids were then slurried and washed with 20% ethanol/ethyl acetate. Concentration of the filtrate left crude amino acid derivative (0.94 g, 75%) as an off white solid which was used without further purification. MS(ES): (M+1)⁺ 208.2 m/z.

25

Preparation 244

(2,2-Dimethylpropylamino)phenylacetic acid methyl ester

30

Neopentylamine (5.13 mL, 43.6 mmol) was combined with α -bromophenylacetic acid (5.00 g, 22.0 mmol) and triethylamine (3.65 mL, 26.2 mmol) in tetrahydrofuran (60 mL) and the mixture overnight at ambient temperature. The mixture was then diluted

-140-

with ethyl acetate, washed with water, and dried over sodium sulfate. Concentration left a residue, which was loaded onto a silica gel column and eluted with methanol/chloroform to yield 3.38 g (66%) of pure amine as an oil. MS(ES): (M+1)⁺ 236.1, 237.1 m/z.

5

Preparation 245

(2,2-Dimethylpropylamino)phenylacetic acid

An amine from preparation 244 (3.20 g, 13.6 mmol) was combined with aqueous 2N sodium hydroxide (30 mL), tetrahydrofuran (10 mL), and ethanol (5 mL) and the mixture stirred overnight at ambient temperature. The mixture was then treated in a manner similar to that in preparation 11 which left crude amino acid derivative (0.337g, 11%) as a white solid and was used without further purification. MS(ES): (M+1)⁺ 222.1 m/z.

15

Preparation 246

Pyrimidine-5-carboxylic acid methyl ester

A mixture of 5-bromopyrimidine (3.0 g, 18.8 mmol), 1.86 g of 1,1'-bis-(diphenylphosphino)ferrocene, and 1.15 g of Pd(II)acetate in 40 mL of methanol and 40 mL of N,N-dimethylformamide was stirred under 60 psi of carbon monoxide at 85°C for 16 hours. The reaction mixture was filtered and partitioned between ether and brine. The aqueous layer was extracted with ether. Combined organics were washed with brine, dried over sodium sulfate and was concentrated to dryness. The residue was dissolved in chloroform, filtered through celite and concentrated to dryness to yield 1.89 g (73%) of the desired crude product as a tan solid. Crude product was carried on without further purification. MS (ion spray) 123.3 (M-methyl)..

25

Preparation 247

Pyrimidine-5-carboxylic acid

30

To a solution of a compound from preparation 246, 1.8 g (13.0 mmol) in 60 mL of tetrahydrofuran, was added a solution of 3.0 g (130 mmol) of lithium hydroxide in 60 mL

-141-

of water. The reaction mixture was stirred for five hours at ambient temperature and was then washed with ether and acidified to pH = 1 with 5 N HCl. The resulting solution was extracted with 20% isopropanol/chloroform. Combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness to yield 460 mg (28%) of the desired product as a crude tan oil. Crude product was carried on without further purification. MS (ion spray) 123.3 (M-) for desired product.

Preparation 248

2-Methoxyisonicotinic acid methyl ester

10

To a solution of 2-methoxyisonicotinic acid methyl ester, 0.32 g (1.9 mmol) in 20 mL of tetrahydrofuran was added 1.9 mL (9.5 mmol) of 5 N sodium hydroxide in 10 mL of water. The reaction mixture was stirred for 2.5 hours at ambient temperature, acidified to pH = 3.6 with 5 N HCl and concentrated to dryness. The residue was slurried in 20% isopropanol/chloroform and filtered. The solid was slurried in ethyl acetate and filtered. The combined organics were dried over sodium sulfate, filtered and concentrated to dryness to give a quantitative yield of the crude desired product as a white solid. MS (ion spray) 154.1 (M+). Crude product was carried on without further purification.

20

Preparation 249

2-Cyano-4-methoxypyridine

To a slurry of 4-methoxypyridine N-oxide hydrate, 50 g, in 250 mL of chloroform was added 20 g of magnesium sulfate. The slurry was stirred for 1 hour at ambient temperature, then filtered and concentrated to dryness. To a solution of the residue, 4-methoxypyridine N-oxide, 43 g (350 mmol) in 350 mL of dichloromethane was added 65 mL (490 mmol) of trimethylsilyl cyanide. The solution was cooled to 10°C and 45 mL (490 mmol) of N,N-dimethylcarbonyl chloride was added dropwise such that the reaction warmed to a gentle reflux. Upon completion of the addition, the reaction mixture was stirred overnight at ambient temperature, then carefully quenched at 0°C dropwise with 10% aqueous potassium carbonate. The layers were separated and the aqueous layer was extracted with dichloromethane. The combined organics were washed with brine, dried

-142-

over sodium sulfate, filtered and concentrated to dryness. The residue was triturated with ether. Filtration of the solid provided 31.4 g (67%) of the desired product as a white solid. ¹H-NMR is consistent with structure; MS (ion spray) 135.1 (M+).

5

Preparation 250

4-Methoxypicolinic acid hydrochloride

A solution of a compound from preparation 249, 15.0 g (112 mmol) in 100 mL of 5 N HCl was refluxed overnight, cooled to ambient temperature and concentrated to dryness to yield 22 g the desired product as a crude, white solid. MS (ion spray) 153.9 (M+). Crude product was carried on without further purification.

10

Preparation 251

6-Methoxynicotinic acid

15

To a solution of methyl 6-methoxynicotinate (Avacado), 0.1 g (0.6 mmol) in 5 mL of tetrahydrofuran was added a solution of 1.2 mL of 5 N sodium hydroxide in 5 mL of water. The mixture was stirred for four hours at ambient temperature, acidified to pH = 3 with 5 N HCl and concentrated to dryness to yield the crude desired compound. MS (ion spray) 154.0 (M+1). The crude product was carried on without further purification.

20

Preparation 252

Dimethylaminophenylacetic acid methyl ester

To a solution of α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 120 mL of tetrahydrofuran was added 12 mL (88.0 mmol) of triethylamine and 5.4 g (66.0 mmol) of dimethylamine hydrochloride. The reaction mixture was heated at 60°C in a sealed tube for three hours, cooled to ambient temperature and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by chromatography using a methanol/chloroform gradient as eluent and concentrated to

30

-143-

dryness to yield 1.86 g (44%) of the desired isomer as a yellow oil. MS (ion spray) 193.9 (M+).

Preparation 253

5

Dimethylaminophenylacetic acid

To a solution of a compound from preparation 252, 60 mg (0.84 mmol) in 10 mL of tetrahydrofuran was added a solution of 0.5 mL (2.52 mmol) of 5 N sodium hydroxide in 5 mL of water. The mixture was stirred for two hours at ambient temperature then an additional 1 mL (5.04 mmol) of 5 N sodium hydroxide was added. The reaction mixture was stirred an additional four hours, acidified to pH = 3 with 5 N HCl, and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. Crude product was carried on without further purification. MS (ion spray) 202.1 (M+23(Na)).

15

Preparation 254

Phenylthiomorpholin-4-ylacetic acid methyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 120 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of triethylamine and 6.7 g (66.0 mmol) of thiomorpholine. The reaction mixture was refluxed for four hours, cooled to ambient temperature and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by chromatography using a methanol/chloroform gradient as eluent. A second purification was performed using chloroform as eluent and was concentrated to dryness to yield 5.36 g (97%) of the desired mixture of isomers as a yellow oil. MS (ion spray) 252.1 (M+).

-144-

Preparation 255

Phenylthiomorpholin-4-ylacetic acid

To a solution of a compound from preparation 254, 140 mg (0.56 mmol) in 5 mL of tetrahydrofuran was added a solution of 0.6 mL (2.8 mmol) of 5 N sodium hydroxide in 5 mL of water. The mixture was stirred for two hours at ambient temperature then an additional 0.5 mL (2.3 mmol) of 5 N sodium hydroxide was added in mL of water. The reaction mixture was stirred an additional two hours, acidified to pH = 3 with 5 N HCl, and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. Crude product was carried on without further purification. MS (ion spray) 236.1 (M+23(Na)).

Preparation 256

(4-Methylpiperazin-1-yl)phenylacetic acid methyl ester

15

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of triethylamine and 7.3 g (66.0 mmol) of 1-methylpiperazine. The reaction mixture was refluxed for three hours, cooled to ambient temperature and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by chromatography using a methanol/chloroform gradient as eluent. A second purification was performed using methanol/chloroform as eluent and was concentrated to dryness to yield 5.21 g (95%) of the desired mixture of isomers as a yellow oil. MS (ion spray) 249.1 (M+).

20

25

Preparation 257

(4-Methylpiperazin-1-yl)phenylacetic acid

30

To a solution of racemic a compound from preparation 256, 200 mg (0.84 mmol) in 10 mL of dioxane was added a solution of 200 mg (8.4 mmol) of lithium hydroxide 8 mL of water. The mixture was stirred for six hours at ambient temperature, was acidified

-145-

to pH = 3 with 5 N HCl, and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. Crude product was carried on without further purification.

MS (ion spray) 235.14 (M+).

5

Preparation 258

(4-Acetylpiperazin-1-yl)phenylacetic acid methyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of triethylamine and 8.5 g (66.0 mmol) of 1-acetylpiperazine. The reaction mixture was refluxed for three hours, cooled to ambient temperature and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by chromatography using a methanol/chloroform gradient as eluent and was concentrated to dryness to give a quantitative yield of the desired mixture of isomers as a yellow, viscous oil. MS (ion spray) 277.2 (M+).

20

Preparation 259

(4-Acetylpiperazin-1-yl)phenylacetic acid

To a solution of a racemic compound from preparation 258, 232 mg (0.84 mmol) in 10 mL of dioxane was added a solution of 200 mg (8.4 mmol) of lithium hydroxide 8 mL of water. The mixture was stirred for six hours at ambient temperature, was acidified to pH = 3 with 5 N HCl, and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. Crude product was carried on without further purification.

25

MS (ion spray) 269.2 (M+Li).

30

-146-

Preparation 259

(Indan-2-ylamino)phenylacetic acid methyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0
5 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of triethylamine and
8.8 g (66.0 mmol) of 2-aminoindan. The reaction mixture was refluxed for four hours,
cooled to ambient temperature and concentrated to dryness. The residue was partitioned
between 20% isopropanol/chloroform and water. The mixture was washed with saturated
sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and
10 concentrated to dryness. The residue was purified by chromatography using chloroform
as eluent and was concentrated to dryness to give a quantitative yield of the desired
mixture of isomers as a tan, viscous oil. MS (ion spray) 282.2 (M+).

Preparation 260

15 (Indan-2-ylamino)phenylacetic acid

To a solution of a racemic compound from preparation 259, 236 mg (0.84 mmol)
in 10 mL of dioxane was added a solution of 200 mg (8.4 mmol) of lithium hydroxide 8
mL of water. The mixture was stirred for four hours at ambient temperature, was
20 acidified to pH = 3 with 5 N HCl, and concentrated to dryness to give a quantitative yield
of the crude desired mixture of isomers. Crude product was carried on without further
purification. MS (ion spray) 268.2 (M+1).

Preparation 261

25 Hydroxy-m-tolyl-acetic acid

To m-methylbenzaldehyde, 21 mL (176 mmol) was added 5 mg of zinc (II) iodide
and 23.4 mL (176 mmol) of trimethylsilyl cyanide dropwise. The resulting solution was
stirred overnight at ambient temperature. To this solution was added 75 mL of 9 N
30 hydrochloric acid. The mixture was stirred overnight at ambient temperature and was
concentrated to dryness. The residue was partitioned between water and 20%
isopropanol/chloroform and washed with brine, dried over sodium sulfate, filtered and

-147-

concentrated to dryness. Crude product was recrystallized from chloroform to provide 15 g of a tan solid. NMR analysis indicates a 2:1 mixture of product to cyanohydrin intermediate. This mixture was carried on as is without further purification. MS (ion spray) 165.1 (M-1).

5

Preparation 262

(2-Fluoro-phenyl)-hydroxy-acetic acid

To o-fluorobenzaldehyde, 18.5 mL (176 mmol) was added 5 mg of zinc (II) iodide and 23.4 mL (176 mmol) of trimethylsilyl cyanide dropwise. The resulting solution was stirred overnight at ambient temperature. To this solution was added 100 mL of 9 N hydrochloric acid. The mixture was stirred overnight at ambient temperature, then refluxed overnight. The mixture was then concentrated to dryness. The residue was partitioned between water and 20% isopropanol/chloroform and washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. Crude product was recrystallized from chloroform to provide 4.57 g (15%) of a tan solid. MS (ion spray) 169.2 (M-).

10

15

Preparation 263

[3-(3-Acetyl-4-amino-5-chloro-2-oxo-2H-quinolin-1-yl)-cyclohexylmethyl]-carbamic acid benzyl ester

20

A solution of [3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-carbamic acid benzyl ester (0.5 g, 1.04 mmol) and Mo(CO)₆ (3.0 g, 0.01 mmol) in Acetonitrile (50 mL) and water (5 mL) was heated to 60°C and stirred overnight. The reaction was concentrated to a solid, diluted in CH₂Cl₂ and filtered through Celite to remove the black insoluble material. The filtrate was concentrated to a solid and purified using silica gel column chromatography. The product was eluted with 10% EtOAc in CH₂Cl₂. The solvent was removed to afford 0.520 g (quantitative) of product as a light brown oil. MS (ES+) m/z 482.0 (M+H)⁺, (ES-) m/z 479.9 (M-H)⁻, 540.0 (M+CH₃COO)⁻.

25

30

-148-

Preparation 264

N-[3-(3-Acetyl-4-amino-5-chloro-2-oxo-2H-quinolin-1-yl)-cyclohexylmethyl]-6-fluoro-nicotinamide

5 A solution of Example 420 (0.07 g, 0.15 mmol) and Mo(CO)₆ (0.43 g, 1.6 mmol) in Acetonitrile (5 mL) and water (1 mL) was heated to 60°C and stirred overnight. The reaction was concentrated to a solid, diluted in CHCl₃ and filtered through Celite to remove the black insoluble material. The filtrate was concentrated to a solid and purified using silica gel column chromatography. The product was eluted with 80% EtOAc in
10 CHCl₃. The solvent was removed to afford 0.04 g (57%) of product as a white solid. MS (ES+) m/z 471.1 (M+H)⁺, (ES-) m/z 469.1 (M-H)⁻, 529.1 (M+CH₃COO)⁻.

Preparation 265

(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid ethyl ester

15

To a mixture of ethyl 3-(2-fluoro-6-iodo-phenyl)-5-methyl-isoxazole-4-carboxylate (0.25 g, 0.67 mmol), tetrakis(triphenylphosphine)palladium (0.08 g, 0.067 mmol), and triethylamine (5 ml) under N₂ was added trimethylsilyl cyanide and heated at reflux overnight. The reaction was cooled to room temperature, diluted with H₂O, and
20 extracted with EtOAc (2x). The combined extracts were washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (0.15 g, 82%). Mass Spectrum (ES+) (m/z) 275.0 [M+1].

25

Preparation 266

3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid

A mixture of a compound from preparation 265 (0.14 g, 0.51 mmol), EtOH (2.5 ml), and 5 N NaOH (0.3 ml, 1.5 mmol) was heated at 50 °C for 1.5h. The reaction was
30 cooled to room temperature, diluted with H₂O, and acidified (conc. HCl) to less than pH 3. The mixture was extracted with EtOAc (2x) and the combined extracts were washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated to give the title compound

-149-

(0.1 g, 80%). This material was used without further purification. Mass Spectrum (ES+) (m/z) 247.0 [M+1].

Preparation 267

5 [3-(2-cyano-6-fluorophenyl)-5-methylisoxazol-4-yl]-N-{3-[(phenylcarbonylamino)-methyl]cyclohexyl}carboxamide

To a solution of a compound from preparation 265 (0.1 g, 0.41 mmol) in dichloromethane (10 ml) under N₂ was added oxalyl chloride (0.07 ml, 0.82 mmol) then
10 DMF (1 drop) and stirred for 2 h. The solution was concentrated, redissolved in dichloromethane (5 ml) under N₂, and N-(3-Amino-cyclohexylmethyl)-benzamide (0.114 g, 0.49 mmol) was added. The reaction vessel was submerged in a water bath and triethylamine (0.17 ml, 1.23 mmol) was added dropwise. After 2.5 h of stirring, the mixture was diluted with CH₂Cl₂, washed (1.0 NaOH then brine), dried (MgSO₄),
15 filtered, and concentrated. Flash chromatography (silica gel, Acetone/CH₂Cl₂ gradient) gave the title compound (0.119 g, 63%). Mass Spectrum (ES+) (m/z) 461.2 [M+1].

Preparation 268

(3-Amino-cyclohexylmethyl)-carbamic acid benzyl ester

20 A solution of [3-(benzyloxycarbonylamino-methyl)-cyclohexyl]-carbamic acid tert-butyl ester (6 g, 16.6 mmol) in TFA (30 ml) was stirred for 1.5 h. The solution was concentrated using benzene to azeotrope, diluted with EtOAc, washed (1.0 N NaOH), dried (Na₂SO₄), filtered, and concentrated to afford the title compound (4.2 g, 97%) as a crude solid. Mass Spectrum (ES+) (m/z) 263.1 [M+1].

25

Preparation 269

(3-{[3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino}-cyclohexylmethyl)-carbamic acid benzyl ester

30 In a fashion similar to that described for preparation 267, a compound from preparation 266 (3.05 g, 12.4 mmol), oxalyl chloride (2.16 ml, 24.8 mmol), dichloromethane (50 ml), DMF (0.05 ml), a compound from preparation 268 (4.2 g, 16.1

-150-

mmol), dichloromethane (125 ml), and triethylamine (5.17 ml, 37.2 mmol) gave the title compound (4.6 g, 76%) after flash chromatography (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 491.2 [M+1].

5

Preparation 270

Diphenylmethyl 2-(3-oxocyclohexyl)propane-1,3-dioate (cis racemic)

To a solution of LiAlH₄ (6 ml, 6 mmol) under N₂ at 0°C was added a solution of (S)-1,1'-binaphthol (3.43 g, 12 mmol) in THF (48 ml) and stirred for 30 min. The reaction was warmed to RT. To this was added a solution of sodium salt benzyl malonate in THF which was prepared by reacting benzyl malonate (14.99 ml, 60 mmol), NaH (60% by wt., 0.216 g, 5.4 mmol), and THF (60 ml) until all bubbling ceased. Next, cyclohexenone (5.82 ml, 60 mmol) and dibenzylmalonate (1.35 ml, 5.4 mmol) were added and the reaction mixture was stirred overnight. 1.0 N HCl was added and extracted with EtOAc (3x). The combined EtOAc layers were washed (brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, acetone/ hexanes gradient) gave the title compound (16.17 g, 71%). Mass Spectrum (FIA) (m/z) 381.3 [M+1].

20

Preparation 271

Diphenylmethyl 2-(3-oxocyclohexyl)propane-1,3-dioate (trans racemic)

To a -78 °C solution of a compound from preparation 270 (see Arai, T.; Yamada, Y. M. A.; Yamamoto, N.; Sasai, H.; Shibasaki, M. *Chem. Eur. J.* 1996, 2, 1368-1372.) (16.15 g, 42.5 mmol) in THF (212.5 ml) under N₂ was added L-selectride (46.75 ml at 1.0 M, 46.75 mmol) dropwise and the mixture was stirred for 3.5 h at -78 °C. EtOAc and H₂O were added and the mixture was warmed to room temperature. Dilution with more EtOAc followed by washing with 1N NaOH, saturated NH₄Cl, brine, drying (MgSO₄), and flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (14.29 g, 88%). Mass Spectrum (FIA) (m/z) 383.3 [M+1].

30

-151-

Preparation 272

Phenylmethyl 2-(3-hydroxycyclohexyl)acetate (trans racemic)

5 A solution of a compound from preparation 271 (19.19 g, 50.2 mmol, LiCl (4.27 g, 100.5 mmol, H₂O (1.81 ml, 100.5 mmol), and DMSO (135 ml) was lowered into a 165 °C oil bath for 2 h then heated at 175 °C for 1.5 h. The reaction was cooled to room temperature and diluted with EtOAc. Washing with H₂O and brine, drying (MgSO₄), and flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (9.71 g, 78 %). ¹H NMR: consistent with structure.

10

Preparation 273

Phenylmethyl 2-(3-azidocyclohexyl)acetate (cis racemic)

15 To a solution of a compound from preparation 272 (9.71 g, 39.2 mmol), triphenylphosphine (12.32 g, 41.04 mmol), and hydrazoic acid (30.9 ml, at 1.9 M, 58.7 mmol) in toluene (120 ml) was added DEAD (9.24 ml, 58.7 mmol) dropwise and stirred for 72 h. The mixture was diluted with EtOAc, washed with 0.1 N NaOH, H₂O, and brine, dried (MgSO₄), and chromatographed (silica gel, CH₂Cl₂/hexanes gradient) to give the title compound (10.0 g, 93%). ¹H NMR: consistent with structure.

20

Preparation 274

Phenylmethyl 2-{3-[(tert-butoxy)carbonylamino]cyclohexyl}acetate (cis racemic)

25 A mixture of a compound from preparation 273 (10 g, 36.6 mmol), (BOC)₂O (9.57 g, 43.9 mmol), Lindlar's catalyst (3.7 g), and EtOAc (200 ml) was stirred under an atmosphere of hydrogen gas (balloon) for 24 h, filtered through celite, and concentrated. Flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (7.03 g, 55%). Mass Spectrum (FD+) (m/z) 347.3 [M⁺].

-152-

Preparation 2752-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid (cis racemic)

5 In a fashion similar to that described for preparation 266, a compound from preparation 274 (7 g, 20.2 mmol), 2 N NaOH (25 ml, 50 mmol), Dioxane (100 ml) were reacted for 5 h to give the title compound (5.2 g, 100%). Mass Spectrum (FD+) (m/z) 257.3 [M⁺].

Preparation 276

10 (*t*-Butoxy)-N-(3-[(phenylmethoxy)carbonylamino]methyl)cyclohexyl)carboxamide
(cis racemic)

15 To a solution of a compound from preparation 275 (2.6 g, 10.1 mmol), Et₃N (2.84 ml, 20.4 mmol) in toluene (100 ml) under N₂ was added DPPA (4.39 ml, 20.4 mmol) and benzyl alcohol (3.13 ml, 30.3 mmol). The solution was heated to reflux overnight. The reaction was cooled to room temperature, diluted with EtOAc, washed (1.0 N NaOH then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (2.92 g, 80%). ¹H NMR: consistent with structure.

20

Preparation 277N-[3-(Aminomethyl)cyclohexyl](*t*-butoxy)carboxamide (cis racemic)

25 A mixture of a compound from preparation 276 (1 g, 2.76 mmol), 10% Pd/C catalyst (0.5 g), and EtOAc (30 ml) was stirred under an atmosphere of hydrogen gas (balloon) for 18 h, filtered through celite, and concentrated to give the title compound (0.48 g, 76%), which was taken on without further purification. Mass Spectrum (ES+) (m/z) 229.1 [M+1].

30

Preparation 278N-({3-[(*t*-Butoxy)carbonylamino]cyclohexyl}methyl)[3-(6-chloro-2-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide

-154-

Preparation 281

[3-(6-Chloro-2-fluorophenyl)-5-pyrrolidinylisoxazol-4-yl]-N-{3-
[(phenylcarbonylamino)methyl]cyclohexyl}carboxamide

5 In a fashion similar to that described for preparation 280, a compound from preparation 279 (0.085 g, 0.17 mmol), DMF (2.5 ml), pyrrolidine (0.14 ml, 1.7 mmol) gave the title compound (0.089 g, 100%) which was taken on as a crude residue. Mass Spectrum (ES+) (m/z) 525.2 [M+1].

10 Preparation 282

[3-(6-Chloro-2-fluorophenyl)-5-(ethylamino)isoxazol-4-yl]-N-{3-[(phenylcarbonyl-
amino)methyl]cyclohexyl}carboxamide

15 In a fashion similar to that described for preparation 280, a compound from preparation 61 (0.085 g, 0.17 mmol), DMF (2.5 ml), ethylamine (0.85 ml, 1.7 mmol) gave the title compound (0.085 g, 100%) which was taken on as a crude residue. Mass Spectrum (ES+) (m/z) 499.2 [M+1].

20 Preparation 283

[3-(6-Chloro-2-fluorophenyl)-5-ethylthioisoxazol-4-yl]-N-{3-[(phenylcarbonyl-
amino)methyl]cyclohexyl}carboxamide

25 In a fashion similar to that described for preparation 280, a compound from preparation 279 (0.1 g, 0.2 mmol), DMF (2.5 ml), sodium ethanethiolate (0.103 g, 1.0 mmol) gave the title compound (0.85 g, 100%) which was taken on as a crude residue. ¹H NMR: consistent with structure.

30 Preparation 284

4-(Methoxycarbonyl-phenyl-methyl)-piperazine-1-carboxylic acid tert-butyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of Et₃N and 12.3 g

-155-

(66.0 mmol) of BOC-piperazine. The reaction mixture was refluxed for 3.5 hours, cooled to rt. and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue
5 was purified by chromatography using chloroform as eluent and was concentrated to dryness to yield 7.73 g (100%) of the desired mixture of isomers as a colorless oil. MS (ion spray) 335.2 (M+).

Preparation 285

10 4-(Carboxy-phenyl-methyl)-piperazine-1-carboxylic acid tert-butyl ester

To a solution of the compound from preparation 284, 560 mg (1.68 mmol) in 15 mL of dioxane was added a solution of 400 mg (16.8 mmol) of lithium hydroxide in 15 mL of water. The solution was stirred for five hours at rt. then acidified to pH = 3.0 with
15 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. MS (ion spray) 319.2 (M-). Crude product was carried on without further purification.

Preparation 286

20 [4-(2-Hydroxy-ethyl)-piperazin-1-yl]-phenyl-acetic acid methyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of Et₃N and 8 mL (66.0 mmol) of 1-(hydroxyethyl)piperazine. The reaction mixture was refluxed for three
25 hours, cooled to rt. and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give a quantitative yield of the desired mixture of isomers as a yellow oil. MS (ion spray) 279.1 (M+).

30

-156-

Preparation 287

[4-(2-Hydroxy-ethyl)-piperazin-1-yl]-phenyl-acetic acid

To a solution of the compound from preparation 286, 470 mg (1.68 mmol) in 20 mL of dioxane was added a solution of 400 mg (16.8 mmol) of lithium hydroxide in 20 mL of water. The solution was stirred for 2.5 hours at rt. then acidified to pH = 3.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired mixture of isomers. Crude product was carried on without further purification. MS (ion spray) 319.2 (M-).

10

Preparation 288

Phenyl-(4-pyridin-2-yl-piperazin-1-yl)-acetic acid methyl ester

To a solution of racemic α -bromophenylacetic acid methyl ester, 5.0 g (22.0 mmol) in 200 mL of tetrahydrofuran was added 3.4 mL (22.0 mmol) of Et₃N and 10 mL (66.0 mmol) of 1-(2-pyridyl)-piperazine. The reaction mixture was refluxed for five hours, cooled to rt. and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by column chromatography using MeOH/chloroform as eluant and concentrated to dryness to give a quantitative yield of the desired mixture of isomers as a yellow oil. MS (ion spray) 312.2 (M+).

20

Preparation 289

25

Piperidin-1-yl-acetic acid ethyl ester

To a solution of bromoethylacetate, 3.3 g (30.0 mmol) in 100 mL of tetrahydrofuran was added 4.2 mL (30.0 mmol) of Et₃N and 9 mL (90.0 mmol) of piperidine. The reaction mixture stirred overnight at rt. and was concentrated to dryness. The residue was partitioned between chloroform and saturated aqueous NaHCO₃. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give 4.6 g (89%) of the desired product as a tan oil.

30

-157-

MS (ion spray) 172.2 (M+).

Preparation 290

Piperidin-1-yl-acetic acid

5

To a solution of the compound from preparation 289, 143 mg (0.84 mmol) in 5 mL of dioxane was added a solution of 66 mg (8.4 mmol) of lithium hydroxide in 5 mL of water. The solution was stirred for three hours at rt. then acidified to pH =2.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product.

10 MS (ion spray) 144.1 (M+). Crude product was carried on without further purification.

Preparation 291

(4-Methyl-piperazin-1-yl)-acetic acid ethyl ester

15

To a solution of bromoethylacetate, 3.3 g (30.0 mmol) in 100 mL of tetrahydrofuran was added 4.14 g (30.0 mmol) of potassium carbonate and 3.3 mL (30.0 mmol) of 4-methylpiperazine. The reaction mixture stirred overnight at rt. and was concentrated to dryness. The residue was partitioned between 20% isopropanol/ chloroform and saturated aqueous NaHCO₃. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give 1.23 g (22%) of the desired product as a tan oil. MS (ion spray) 187.1 (M+).

20

Preparation 292

(4-Methyl-piperazin-1-yl)-acetic acid

25

To a solution of the compound from preparation 291, 160 mg (0.84 mmol) in 5 mL of dioxane was added a solution of 66 mg (8.4 mmol) of lithium hydroxide in 5 mL of water. The solution was stirred for two hours at rt. then acidified to pH =3.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product.

30 MS (ion spray) 159.1 (M+). Crude product was carried on without further purification.

-158-

Preparation 293

(Methyl-phenyl-amino)-acetic acid ethyl ester

To a solution of bromoethylacetate, 3.3 mL (30.0 mmol) in 100 mL of
5 tetrahydrofuran was added 4.14 g (30.0 mmol) of potassium carbonate and 3.25 mL of N-
methylaniline. The reaction mixture stirred overnight at rt. and was concentrated to
dryness. The residue was partitioned between 20% isopropanol/chloroform and saturated
aqueous NaHCO₃. The mixture was washed with saturated NaHCO₃, washed with brine,
dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by
10 column chromatography using EtOAc/hexanes as eluent and concentrated to dryness to
give 4.74 g (82%) of the desired product as a tan oil. MS (ion spray) 194.1 (M+1).

Preparation 294

Phenyl-(pyridin-3-yloxy)-acetic acid methyl ester

15

To a slurry of 3-hydroxypyridine sodium salt, 4.6 g (26.4 mmol) in 200 mL of
tetrahydrofuran was added 5.0 g (22.0 mmol) of alpha-bromophenylacetic acid. The
reaction mixture was refluxed for two hours, cooled to rt. and concentrated to dryness.
The residue was partitioned between 20% isopropanol/chloroform and water. The
20 mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄,
filtered and concentrated to dryness. The residue was purified by column chromatography
using MeOH/chloroform as eluent and concentrated to dryness to give 2.8 g of the desired
mixture of isomers as a dark oil. MS (ion spray) 244.1 (M+).

25

Preparation 295

(Pyridin-3-yloxy)-acetic acid ethyl ester

To a solution of 3-hydroxypyridine sodium salt, 6.4 g (36.0 mmol) in 200 mL of
DMF was added 3.3 mL (30.0 mmol) of bromoethylacetate. The reaction mixture stirred
30 overnight at rt. and was concentrated to dryness. The residue was partitioned between
20% isopropanol/chloroform and saturated aqueous NaHCO₃. The mixture was washed
with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and

-159-

concentrated to dryness. The residue was purified by column chromatography using MeOH/chloroform as eluent and concentrated to dryness to give 3.8 g (70%) of the desired product as a tan oil. MS (ion spray) 182.1 (M+).

5

Preparation 296

(Pyridin-3-yloxy)-acetic acid

To a solution of the compound from preparation 295, (pyridin-3-yloxy)-acetic acid ethyl ester, 300 mg (1.68 mmol) in 15 mL of dioxane was added a solution of 400 mg
10 (16.8 mmol) of lithium hydroxide in 15 mL of water. The solution was stirred for three hours at rt. then acidified to pH =1.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product. MS (ion spray) 154.1 (M+). Crude product was carried on without further purification.

15

Preparation 297

1-tert-Butoxycarbonylamino-cyclohexanecarboxylic acid

To a slurry of 1-amino-1-cyclohexane-carboxylic acid, 5.0 g (35.0 mmol) in 50 mL of tetrahydrofuran and 50 mL of water was added 14.5 g (105.0 mmol) of potassium
20 carbonate, 7.7 g (35.0 mmol) of BOC-anhydride and 5 mg of DMAP. The mixture was stirred overnight at rt. and the organics were evaporated off in vacuo. The aqueous layer was acidified to pH = 1 with 5 N HCl and extracted with 20% isopropanol/chloroform. The combined organics were washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to yield 1.83 g (21%) of the desired product as a white oil that
25 solidifies upon standing. MS (IS) 244.2 (M+).

Preparation 298

Morpholin-4-yl-acetic acid ethyl ester

30

To a solution of bromoethylacetate, 3.3 mL (30.0 mmol) in 100 mL of tetrahydrofuran was added 4.14 g (30.0 mmol) of potassium carbonate and 2.6 mL (30.0 mmol) of morpholine. The reaction mixture stirred overnight at rt. and was concentrated

-160-

to dryness. The residue was partitioned between 20% isopropanol/chloroform and saturated aqueous NaHCO₃. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give 2.75 g (53%) of the desired product as a tan oil. MS (ion spray) 174.2 (M+).

5

Preparation 299

Morpholin-4-yl-acetic acid

To a solution of the compound from preparation 298, 150 mg (0.84 mmol) in 10 mL of dioxane was added a solution of 198 mg (8.4 mmol) of lithium hydroxide in 10 mL of water. The solution was stirred for two hours at rt. then acidified to pH =2.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product. MS (ion spray) 146.0 (M+). Crude product was carried on without further purification.

15

Preparation 300

(4-Hydroxy-piperidin-1-yl)-acetic acid ethyl ester

To a solution of bromoethylacetate, 3.3 mL (30.0 mmol) in 100 mL of tetrahydrofuran was added 4.2 g (30.0 mmol) of potassium carbonate and 3.0 mL (30.0 mmol) of 4-hydroxypiperidine. The reaction mixture stirred overnight at rt. and was concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and saturated aqueous NaHCO₃. The mixture was washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness to give 4.65 g (83%) of the desired product as a tan oil. MS (ion spray) 188.0 (M+).

25

Preparation 301

(4-Hydroxy-piperidin-1-yl)-acetic acid

To a solution of the compound from preparation 300, 158 mg (0.84 mmol) in 10 mL of dioxane was added a solution of 198 mg (8.4 mmol) of lithium hydroxide in 10 mL of water. The solution was stirred for 1.5 hours at rt. then acidified to pH =2.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product.

-161-

MS (ion spray) 160.1 (M+). Crude product was carried on without further purification.

Preparation 302

(2-Oxo-2H-pyridin-1-yl)-acetic acid ethyl ester

5

To a slurry of sodium hydride, 1.73 g (43.2 mmol) in 100 mL of DMF was added 3.4 g (36.0 mmol) of 2-hydroxypyridine. The mixture was stirred 10 minutes then added to a solution of 3.3 mL (30.0 mmol) of bromoethylacetate in 100 mL DMF. The reaction mixture was stirred 72 hours at rt. then concentrated to dryness. The residue was

10 partitioned between 20%isopropanol/chloroform and water and washed with saturated aqueous NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by column chromatography using MeOH/chloroform as eluent and concentrated to dryness to yield 3.52 g (65%) of the desired product as a colorless oil. ¹H-NMR is consistent with structure. MS (ion spray) 182.1 (M+).

15

Preparation 303

(2-Oxo-2H-pyridin-1-yl)-acetic acid

To a solution of the compound from preparation 302, 152 mg (0.84 mmol) in 10

20 mL of dioxane was added a solution of 198 mg (8.4 mmol) of lithium hydroxide in 10 mL of water. The solution was stirred for two hours at rt. then acidified to pH =1.0 with 5 N HCl and concentrated to dryness to give a quantitative yield of the crude desired product. MS (ion spray) 154.1 (M+). Crude product was carried on without further purification.

25

Preparation 304

(Pyridin-4-yloxy)-acetic acid ethyl ester

To a slurry of sodium hydride, 1.73 g (43.2 mmol) in 100 mL of DMF was added 3.4 g (36.0 mmol) of 4-hydroxypyridine. The mixture was stirred 10 minutes then added

30 to a solution of 3.3 mL (30.0 mmol) of bromoethylacetate in 100 mL DMF . The reaction mixture was stirred 72 hours at rt. then concentrated to dryness. The residue was partitioned between 20%isopropanol/chloroform and water and washed with saturated

-162-

aqueous NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by radial chromatography using MeOH/chloroform as eluent and concentrated to dryness to yield 510 mg (9.4%) of the desired product as a tan oil. ¹H-NMR is consistent with structure. MS (ion spray) 182.1 (M+).

5

Preparation 305

tert-Butylamino-acetic acid

To a solution of tert-Butylamino-acetic acid methyl ester (Maybridge), 122 mg
10 (0.84 mmol) in 10 mL of dioxane was added a solution of 60 mg (2.52 mmol) of lithium hydroxide in 10 mL of water. The reaction mixture was stirred for two hours, acidified to pH = 1 with 5 N HCl and concentrated to dryness to yield the crude desired product which was carried on without further purification. MS (IS) 132.1 (M+1).

15

Preparation 306

(6-Methoxy-pyridin-3-ylamino)-acetic acid

To a solution of (6-methoxy-pyridin-3-ylamino)-acetic acid methyl ester, 113 mg
20 (0.58 mmol) in 10 mL of dioxane was added a solution of 70 mg (2.9 mmol) of lithium hydroxide in 10 mL of water. The reaction mixture was stirred for three hours, acidified to pH = 1 with 5 N HCl and concentrated to dryness to yield the crude desired product which was carried on without further purification. MS (IS) 183.1 (M+1).

25

Preparation 307

2-(Pyridin-3-yloxy)-propionic acid

To a solution of 2-(pyridin-3-yloxy)-propionic acid ethyl ester, 320 mg (1.6 mmol)
30 in 10 mL of dioxane was added a solution of 190 mg (8.0 mmol) of lithium hydroxide in 10 mL of water. The reaction mixture was stirred for three hours, acidified to pH = 1 with 5 N HCl and concentrated to dryness to yield the crude desired product which was carried on without further purification. MS (IS) 168.1 (M+1).

-163-

Preparation 308

1-Benzoyl-pyrrolidine-2-carboxylic acid methyl ester

5 Benzoyl chloride (1.40 mL, 12.1 mmol) was added in a dropwise manner to a mixture of L-proline methyl ester hydrochloride (2.00 g, 12.1 mmol) and Et₃N (4.20 mL, 30.2 mmol) in CH₂Cl₂ (40 mL) and the resulting mixture stirred overnight at rt. The mixture was concentrated *in vacuo* and the residue treated with water and extracted with EtOAc and the combined extracts dried over Na₂SO₄. Concentration left a residue which was loaded onto a silica gel column and eluted with MeOH/CH₂Cl₂ which allowed for
10 isolation of 2.76 g (94%) of the title compound as a white solid. MS(ES): (M+1)⁺ 234.2 m/z.

Preparation 309

1-Benzoyl-pyrrolidine-2-carboxylic acid

15

The compound from preparation 308 (1.00 g, 4.3 mmol) was combined with aqueous 2N sodium hydroxide (4.0 mL, 8.0 mmol), tetrahydrofuran (2.0 mL) and MeOH (2.0 mL) and the mixture stirred at rt. until hydrolysis was complete. Organic solvents were removed *in vacuo* and the mixture was diluted with water and adjusted to pH 2.0-3.0
20 with aqueous hydrochloric acid. The mixture was then concentrated to dryness *in vacuo*. The resulting solids were then slurried (and washed) with an EtOAc/ethanol mixture followed by filtration. Concentration of the filtrate left desired acid of acceptable purity (0.52 g, 55%) as a white solid. MS(ES): (M+1)⁺ 220.3 m/z.

25

Preparation 310

1-Phenylacetyl-pyrrolidine-2-carboxylic acid methyl ester

Phenacetyl chloride (1.60 mL, 12.1 mmol) was added to a mixture of L-proline methyl ester hydrochloride (2.00 g, 12.1 mmol) and Et₃N (4.20 mL, 30.2 mmol) in
30 CH₂Cl₂ (40 mL) and the resulting mixture stirred overnight at rt. The mixture was treated in a manner similar to that in preparation 103, which resulted in recovery of 1.74 g (58%) of the desired ester as a clear oil. MS(ES): (M+1)⁺ 248.2, 249.2 m/z.

-164-

Preparation 311

1-Phenylacetyl-pyrrolidine-2-carboxylic acid

5 The compound from preparation 310 (1.00 g, 4.0 mmol) was combined with aqueous 2N sodium hydroxide (5.0 mL, 10.0 mmol), tetrahydrofuran (2.0 mL) and MeOH (2.0 mL) and the mixture stirred at rt. until hydrolysis was complete. The mixture was then treated in a manner similar to preparation 104, however when the pH was adjusted the desired acid began to precipitate. Filtration and drying of the precipitate netted 0.70 g
10 (74%) of the title compound as a white powdery solid. MS(ES): (M+1)⁺ 234.3.

Preparation 312

N-Benzyl-L-proline

15 N-Benzyl-L-proline ethyl ester (1.00 g, 4.3 mmol) was combined with aqueous 2N sodium hydroxide (5.0 mL, 10.0 mmol), tetrahydrofuran (2.0 mL) and MeOH (2.0 mL) and the mixture stirred at rt. until hydrolysis was complete. The mixture was then treated in a manner similar to preparation 104. Concentration of the filtrate netted 0.79 g (90%) of crude acid as a light yellowish solid. MS(ES): (M+1)⁺ 206.3, 207.3.

20

Preparation 314

3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid ethyl ester

To a mixture of ethyl 3-(2-Fluoro-6-iodo-phenyl)-5-methyl-isoxazole-4-
25 carboxylate (0.25 g, 0.67 mmol), tetrakis(triphenylphosphine)palladium (0.08 g, 0.067 mmol), and Et₃N (5 mL) under N₂ was added TMSCN and heated at reflux overnight. The reaction was cooled to room temperature, diluted with H₂O, and extracted with EtOAc (2x). The combined extracts were washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, EtOAc/hexanes gradient)
30 gave the title compound (0.15 g, 82%). Mass Spectrum (ES+) (m/z) 275.0 [M+1].

-165-

Preparation 315

3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid

5 A mixture of a compound from preparation 314, (0.14 g, 0.51 mmol), EtOH (2.5 mL), and 5 N NaOH (0.3 mL, 1.5 mmol) was heated at 50 °C for 1.5h. The reaction was cooled to rt., diluted with H₂O, and acidified (conc. HCl) to less than pH 3. The mixture was extracted with EtOAc (2x) and the combined extracts were washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated to give the title compound (0.1 g, 80%). This material was used without further purification. Mass Spectrum (ES+) (m/z) 247.0 [M+1].

10

Preparation 316

3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid [3-(benzoylamino-methyl)-cyclohexyl]-amide

15 To a solution of a compound from preparation 315, (0.1 g, 0.41 mmol) in CH₂Cl₂ (10 mL) under N₂ was added oxalyl chloride (0.07 mL, 0.82 mmol) then DMF (1 drop) and stirred for 2 h. The solution was concentrated, redissolved in CH₂Cl₂ (5 mL) under N₂, and N-(3-amino-cyclohexylmethyl)-benzamide (0.114 g, 0.49 mmol) was added. The reaction vessel was submerged in a water bath and Et₃N (0.17 mL, 1.23 mmol) was added
20 dropwise. After 2.5 h of stirring, the mixture was diluted with CH₂Cl₂, washed (1.0 NaOH then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, Acetone/CH₂Cl₂ gradient) gave (0.119 g, 63%). Mass Spectrum (ES+) (m/z) 461.2 [M+1].

25

Preparation 317

(3-Amino-cyclohexylmethyl)-carbamic acid benzyl ester

A solution of [3-(benzyloxycarbonylamino-methyl)-cyclohexyl]-carbamic acid tert-butyl ester (6 g, 16.6 mmol) in TFA (30 mL) was stirred for 1.5 h. The solution was
30 concentrated using benzene to azeotrope, diluted with EtOAc, washed (1.0 N NaOH), dried (Na₂SO₄), filtered, and concentrated to afford (4.2 g, 97%) as a crude solid. Mass Spectrum (ES+) (m/z) 263.1 [M+1].

-166-

Preparation 318

(3-{[3-(2-Cyano-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino}-
cyclohexylmethyl)-carbamic acid benzyl ester

5

In a fashion similar to that described for preparation 316, a compound from preparation 315 (3.05 g, 12.4 mmol), oxalyl chloride (2.16 mL, 24.8 mmol), CH₂Cl₂ (50 mL), DMF (0.05 mL), (3-amino-cyclohexylmethyl)-carbamic acid benzyl ester (4.2 g, 16.1 mmol), CH₂Cl₂ (125 mL), and Et₃N (5.17 mL, 37.2 mmol) gave the title compound
10 (4.6 g, 76%) after flash chromatography (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 491.2 [M+1].

Preparation 319

2-(3-Oxo-cyclohexyl)-malonic acid dibenzyl ester

15

To a solution of LiAlH₄ (6 mL, 6 mmol) under N₂ at 0°C was added a solution of (S)-1,1'-binaphthol (3.43 g, 12 mmol) in THF (48 mL) and stirred for 30 min. The reaction was warmed to RT. To this was added a solution of sodium salt benzyl malonate in THF which was prepared by reacting benzyl malonate (14.99 mL, 60 mmol), NaH
20 (60% by wt., 0.216 g, 5.4 mmol), and THF (60 mL) until all bubbling ceased. Next, cyclohexenone (5.82 mL, 60 mmol) and dibenzylmalonate (1.35 mL, 5.4 mmol) were added and the reaction mixture was stirred overnight. 1.0 N HCl was added and extracted with EtOAc (3x). The combined EtOAc layers were washed (brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, acetone/hexanes gradient)
25 gave the title compound (16.17 g, 71%). Mass Spectrum (FIA) (m/z) 381.3 [M+1].

Preparation 320

2-(3-Hydroxy-cyclohexyl)-malonic acid dibenzyl ester

30

To a -78 °C solution of a compound from preparation 319 (see Arai, T.; Yamada, Y. M. A.; Yamamoto, N.; Sasai, H.; Shibasaki, M. *Chem. Eur. J.* 1996, 2, 1368-1372.) (16.15 g, 42.5 mmol) in THF (212.5 mL) under N₂ was added L-selectride (46.75 mL at

-167-

1.0 M, 46.75 mmol) dropwise and the mixture was stirred for 3.5 h at $-78\text{ }^{\circ}\text{C}$. EtOAc and H_2O were added and the mixture was warmed to rt. Dilution with more EtOAc followed by washing with 1N NaOH, saturated NH_4Cl , brine, drying (MgSO_4), and flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (14.29 g, 88%). Mass Spectrum (FIA) (m/z) 383.3 [M+1].

Preparation 321

(3-Hydroxy-cyclohexyl)-acetic acid benzyl ester

10 A solution of a compound from preparation 320 (19.19 g, 50.2 mmol, LiCl (4.27 g, 100.5 mmol, H_2O (1.81 mL, 100.5 mmol), and DMSO (135 mL) was lowered into a $165\text{ }^{\circ}\text{C}$ oil bath for 2 h then heated at $175\text{ }^{\circ}\text{C}$ for 1.5 h. The reaction was cooled to rt. and diluted with EtOAc. Washing with H_2O and brine, drying (MgSO_4), and flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (9.71 g, 15 78%). ^1H NMR: consistent with structure.

Preparation 322

(3-Azido-cyclohexyl)-acetic acid benzyl ester

20 To a solution of a compound from preparation 321 (9.71 g, 39.2 mmol), Ph_3P (12.32 g, 41.04 mmol), and hydrazoic acid (30.9 mL, at 1.9 M, 58.7 mmol) in toluene (120 mL) was added DEAD (9.24 mL, 58.7 mmol) dropwise and stirred for 72 h. The mixture was diluted with EtOAc, washed with 0.1 N NaOH, H_2O , and brine, dried (MgSO_4), and chromatographed (silica gel, CH_2Cl_2 /hexanes gradient) to give the title 25 compound (10.0 g, 93%). ^1H NMR: consistent with structure.

Preparation 323

(3-tert-Butoxycarbonylamino-cyclohexyl)-acetic acid benzyl ester

30 A mixture of a compound from preparation 322 (10 g, 36.6 mmol), $(\text{BOC})_2\text{O}$ (9.57 g, 43.9 mmol), Lindlar's catalyst (3.7 g), and EtOAc (200 mL) was stirred under an atmosphere of hydrogen gas (balloon) for 24 h, filtered through celite, and concentrated.

-168-

Flash chromatography (silica gel, EtOAc/hexanes gradient) gave the title compound (7.03 g, 55%). Mass Spectrum (FD+) (m/z) 347.3 [M⁺].

Preparation 324

5 (3-tert-Butoxycarbonylamino-cyclohexyl)-acetic acid

In a fashion similar to that described for preparation 315, a compound from preparation 323 (7 g, 20.2 mmol), 2 N NaOH (25 mL, 50 mmol), Dioxane (100 mL) were reacted for 5 h to give the title compound (5.2 g, 100%). Mass Spectrum (FD+) (m/z)
10 257.3 [M⁺].

Preparation 325

[3-(Benzyloxycarbonylamino-methyl)-cyclohexyl]-carbamic acid tert-butyl ester

15 To a solution of a compound from preparation 324 (2.6 g, 10.1 mmol), Et₃N (2.84 mL, 20.4 mmol) in toluene (100 mL) under N₂ was added DPPA (4.39 mL, 20.4 mmol) and benzyl alcohol (3.13 mL, 30.3 mmol). The solution was heated to reflux overnight. The reaction was cooled to room temperature, diluted with EtOAc, washed (1.0 N NaOH then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel,
20 hexanes/ EtOAc gradient) gave the title compound (2.92 g, 80%). ¹H NMR: consistent with structure.

Preparation 326

(3-Aminomethyl-cyclohexyl)-carbamic acid tert-butyl ester

25

A mixture of a compound from preparation 325 (1 g, 2.76 mmol), 10% Pd/C catalyst (0.5 g), and EtOAc (30 mL) was stirred under an atmosphere of hydrogen gas (balloon) for 18 h, filtered through celite, and concentrated to give (0.48 g, 76%) which was taken on without further purification. Mass Spectrum (ES+) (m/z) 229.1 [M+1].

30

-169-

Preparation 327

[3-([3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino)-methyl]-cyclohexyl]-carbamic acid tert-butyl ester

5 To a solution of a compound from preparation 326 (0.48 g, 2.1 mmol) and 3-(2-chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.75 g, 2.74 mmol) in CH₂Cl₂ (20 mL) under N₂ was added dropwise Et₃N (0.73 mL, 5.25 mmol) and stirred for 4 h. The reaction was diluted with CH₂Cl₂, washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, hexanes/EtOAc
10 gradient) gave the title compound (0.726 g, 74%). Mass Spectrum (ES+) (m/z) 366.1 [M-BOC].

Preparation 328

15 5-Chloro-3-(2-chloro-6-fluoro-phenyl)-isoxazole-4-carboxylic acid [3-(benzoylamino-methyl)cyclohexyl]amide

In a fashion similar to that described for preparation 316, (0.22 g, 0.84 mmol), CH₂Cl₂ (2 mL), oxalyl chloride (0.15 mL, 1.68 mmol), DMF (0.01 mL), CH₂Cl₂ (5 mL), N-(3-Amino-cyclohexylmethyl)-benzamide (0.15 g, 0.65 mmol), and Et₃N (0.35 mL,
20 2.52 mmol) gave the title compound (0.29 g, 90%) after flash chromatography (silica gel, Acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 490.1 [M+1].

Preparation 329

25 3-(2-Chloro-6-fluoro-phenyl)-5-diethylamino-isoxazole-4-carboxylic acid {3-[(2-propenyl-penta-2,4-dienoylamino)-methyl]-cyclohexyl}-amide

To a solution of a compound from preparation 328 (0.1 g, 0.2 mmol) in DMF (2.5 mL) under N₂ was added diethylamine (0.065 mL, 0.63 mmol) and stirred for 2 h. The reaction was diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and
30 concentrated. (0.1 N HCl and brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, acetone/CH₂Cl₂ gradient) gave the title compound (0.1 g, 93%). Mass Spectrum (ES+) (m/z) 527.2 [M+1].

-170-

Preparation 330

3-(2-Chloro-6-fluoro-phenyl)-5-pyrrolidin-1-yl-isoxazole-4-carboxylic acid [3-(benzoylaminomethyl)-cyclohexyl]-amide

5

In a fashion similar to that described for preparation 329, a compound from preparation 328 (0.085 g, 0.17 mmol), DMF (2.5 mL), pyrrolidine (0.14 mL, 1.7 mmol) gave (0.089 g, 100%) which was taken on as a crude residue. Mass Spectrum (ES+) (m/z) 525.2 [M+1].

10

Preparation 331

3-(2-Chloro-6-fluoro-phenyl)-5-ethylamino-isoxazole-4-carboxylic acid [3-(benzoylaminomethyl)-cyclohexyl]-amide

15

In a fashion similar to that described for preparation 329, a compound from preparation 328 (0.085 g, 0.17 mmol), DMF (2.5 mL), ethylamine (0.85 mL, 1.7 mmol) gave the title compound (0.085 g, 100%) which was taken on as a crude residue. Mass Spectrum (ES+) (m/z) 499.2 [M+1].

20

Preparation 332

3-(2-Chloro-6-fluoro-phenyl)-5-ethylsulfanyl-isoxazole-4-carboxylic acid [3-(benzoylamino-methyl)-cyclohexyl]-amide

25

In a fashion similar to that described for preparation 329, a compound from preparation 328 (0.1 g, 0.2 mmol), DMF (2.5 mL), sodium ethanethiolate (0.103 g, 1.0 mmol) gave the title compound (0.85 g, 100%) which was taken on as a crude residue. ¹H NMR: consistent with structure.

-171-

Preparation 333

[3-({[3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino}-methyl)-cyclohexyl]-carbamic acid tert.-butyl ester

5 A mixture of a compound from preparation 326 (0.85 g, 2.35 mmol), 10% Pd/C catalyst (0.5 g), and EtOAc (25 mL) was stirred under an atmosphere of hydrogen gas (balloon) for 18 h, filtered through celite, and concentrated. This residue was mixed with 3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride (0.772 g, 2.82 mmol) in CH₂Cl₂ (7 mL) under N₂ and to this solution was added dropwise Et₃N (1.0 mL, 7.05
10 mmol). After stirring 48h, the mixture was diluted with CH₂Cl₂, washed (0.1 N HCl then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, EtOAc/Hexanes gradient) gave the title compound (0.64 g, 58%). Mass Spectrum (ES+) (m/z) 366.3 [M-BOC].

15

Preparation 334

5-[3-(Benzoylamino-methyl)-cyclohexyl]-3-methyl-4-oxo-4,5-dihydro-isoxazolo[4,3-c]quinolin-9-yl}-carbamic acid 2-trimethylsilyl-ethyl ester

To a solution of 3-methyl-4-oxo-5-{3-[(phenylcarbonylamino)methyl]-
20 cyclohexyl}-5-hydroisoxazolo[4,3-c]quinoline-9-carboxylic acid (0.25 g, 0.55 mmol), Et₃N (0.118 mL, 0.845 mmol) in toluene (3.5 mL) under N₂ was added DPPA (0.18 mL, 0.845 mmol) and 2-(trimethylsilyl) ethanol (0.232 mL, 1.63 mmol). The solution was heated to reflux for 4 h. The reaction was cooled to rt., diluted with EtOAc, washed (1.0 N NaOH then brine), dried (MgSO₄), filtered, and concentrated. Column chromatography
25 (silica gel, acetone/CH₂Cl₂ gradient) gave the title compound (0.285 g, 90%). Mass Spectrum (ES-) (m/z) 573.3 [M-1]

Preparation 335

[3-(3,4,5-Trimethoxy-phenylcarbonyl)-cyclopentylmethyl]-carbamic acid tert-butyl ester

30

To a solution of 0.224g (0.92 mmols) of 3-[(*t*-butoxy)amino]cyclopentane-carboxylic acid was added 0.44 mmol of 1-hydroxy-7-azabenzotriazole and 0.44 mmol of

-172-

EDCI. After 40 minutes, 0.44 mmol of 3,4,5-trimethoxyaniline was added. After 3.25 hours, the solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1 N HCl, aq. NaHCO₃, and then x3 with water. The organic layer was dried and the solvent was removed *in vacuo* to yield 0.34g of the title compound.

5 MS(ES+)m/z=409.

Preparation 336

3-Aminomethyl-cyclopentanecarboxylic acid (3,4,5-trimethoxy-phenyl)-amide

10 A solution of 0.049g (0.12 mmol) of a compound from preparation 335 in TFA (10ml) was stirred for 1 hour, after which the TFA was removed *in vacuo* and replaced with ethyl acetate. The organic layer was rinsed with 1N NaOH followed by brine and dried. The organic layer was removed *in vacuo* to yield 0.013g of the title compound.
MS(ES+)m/z=308.

15

Preparation 337

3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid [3-(3,4,5-trimethoxy-phenylcarbamoyl)-cyclopentylmethyl]-amide

20 A solution of 0.013g (0.042 mmol) of a compound from preparation 336, 0.03g (0.46 mmol) of 3-(6-chloro-2-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride, and 0.07 ml (0.5 mmol) of triethylamine in dichloromethane (10 ml) was stirred for 12 hours. The solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was washed with 1N HCl, then aq. NaHCO₃, followed by brine and dried. The solvent was
25 removed *in vacuo* then chromatographed on silica gel with CH₂Cl₂ 100% to CH₂Cl₂/MeOH 2% to yield 0.018g of the title compound. MS(ES+)m/z=545.8.

-173-

Preparation 338

(3-{[3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino}-cycloheptyl)-
acetic acid methyl ester

5 A solution of 4.4g (24 mmol) of methyl 2-(3-aminocycloheptyl)acetate, 0.03g
(0.46 mmol) of 3-(6-chloro-2-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride, and
0.07 ml (0.5 mmol) of triethylamine in dichloromethane (10 ml) was stirred for 12 hours.
The solvent was removed *in vacuo* and replaced with ethyl acetate. The organic layer was
washed with 1N HCl, then aq. NaHCO₃, followed by brine and dried. The solvent was
10 removed *in vacuo* then chromatographed on silica gel with hexane/EtOAc 3/1 to yield
1.7g of the title compound. MS(ES+)m/z=423.

Preparation 339

15 5-(3-Aminomethyl-cycloheptyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

To a solution of 0.43g (1.02 mmol) of N-{[3-(9-chloro-3-methyl-4-oxo(5-
hydroisoxazolo[4,3-c]quinolin-5-yl))cycloheptyl]methyl}methoxycarboxamide in
CH₂Cl₂ (10 mL) was added 0.35 mL of trimethylsilyl iodide(2.1 mmol) and the solution
was stirred for 15 hours at ambient temperature. MeOH (1.5 mL) was added and the
20 reaction was stirred for 15 min. before the solvent was removed *in vacuo*. The residue was
dissolved in CH₂Cl₂ and rinsed with 1N NaOH followed by 50% brine/water. After
drying over Na₂SO₄, the solvent was removed to yield 0.397g of the title compound.
MS(ES+)m/z=360.1.

25

Preparation 340

tert-butylamino-pyridin-3-yl-acetic acid ethyl ester

A typical synthesis of bromo-pyridin-3-yl-acetic acid ethyl ester is described: A
solution of fresh lithium diisopropylamide (LDA) was prepared at -10 °C from
30 diisopropylamine (8.54 mL; 60.5 mmol) and *n*-butyllithium (nBuLi) (37.8 mL of a 1.6 M
hexanes solution; 12.1 mmol) and stirred for 10 min. After chilling the fresh LDA to -78
°C, ethyl 3-pyridylacetate (9.21 mL; 60.5 mmol) was added and the solution was stirred

-174-

another 10 min. TMSCl (7.68 mL; 60.5 mmol) was added to the resulting opaque yellow slurry and the solution was stirred 5 min. Finally a solution of 4-(dimethylamino) pyridinium tribromide (22.0 g; 60.5 mmol) in tetrahydrofuran was added and the reaction solution was stirred 10 min. After warming to rt. the reaction solution was quenched with saturated NH₄Cl (aq) and extracted twice with ethyl acetate. The combined organic layer was washed once with saturated NaCl (aq), dried with Na₂SO₄, filtered, and concentrated in vacuo. The resulting bromo-pyridin-3-yl-acetic acid ethyl ester was an unstable brown oil and was therefore used immediately by preparing aliquots of product in dichloromethane and proceeding.

10 Bromo-pyridin-3-yl-acetic acid ethyl ester (2.4g; 9.8 mmol) was reacted in refluxing CH₂Cl₂, overnight with tert-butylamine (6.2 mL; 55.8 mmol; 6 equiv) and Et₃N (2.8 mL; 19.6 mmol; 2 equiv). The reaction solution was evaporated to dryness, dissolved in CH₂Cl₂, washed six times with saturated NaHCO₃ (aq), washed once with saturated NaCl (aq), dried with Na₂SO₄, filtered, and concentrated in vacuo. The 15 resulting brown oil was purified with silica gel chromatography in a 6×8 cm glass column using a 30% EtOAc/hexanes (v/v) mobile phase. A total of 955 mg desired product was isolated (41% yield). MS(ES) calc'd: [M+H]⁺ = 237.1 m/z. Found: 237.1 m/z.

Preparation 341

20 (2,2-dimethyl-propylamino)-pyridin-3-yl-acetic acid ethyl ester

Bromopyridin-3-yl-acetic acid ethyl ester (2.54g; 10.2 mmol), prepared in a manner similar to preparation 340, was reacted in refluxing CH₂Cl₂, overnight with neopentylamine (4.83 mL; 41.0 mmol; 4 equiv) and Et₃N (5.8 mL; 41.0 mmol; 4 equiv). 25 The reaction solution was evaporated to dryness, dissolved in CH₂Cl₂, washed five times with saturated NaHCO₃(aq), washed once with saturated NaCl(aq), dried with Na₂SO₄, filtered, and concentrated in vacuo. The resulting brown oil was purified with silica gel chromatography in a 3×25 cm glass column using a 0.5% methanol/chloroform (v/v) mobile phase. A second column purification with 100% chloroform followed by 1% 30 methanol/chloroform (v/v) produced pure product as a yellow oil (1.0 g; 40% yield). MS(ES) calc'd: [M+H]⁺ = 250.1 m/z. Found: 250.1 m/z.

-175-

Preparation 342

Dimethylamino-pyridin-3-yl-acetic acid ethyl ester

5 Bromopyridin-3-yl-acetic acid ethyl ester (2.54g; 10.2 mmol) was reacted in a sealed tube, at rt., in CH₂Cl₂, overnight with *N,N*-dimethylamine hydrochloride (3.34 g; 41.0 mmol; 4 equiv) and Et₃N (5.8 mL; 41.0 mmol; 4 equiv). The reaction solution was evaporated to dryness, dissolved in CH₂Cl₂, washed five times with saturated NaHCO₃ (aq), washed once with saturated NaCl (aq), dried with Na₂SO₄, filtered, and concentrated in vacuo. The resulting brown oil was purified with silica gel chromatography in a 3×25
10 cm glass column using a 0.5% methanol/chloroform (v/v) mobile phase. The desired product was isolated as a yellow oil (1.12 g; 53% yield). MS(ES) calc'd: [M+H]⁺ = 208.1 m/z. Found: 208.1 m/z.

Preparation 343

15 (4-methyl-piperazin-1-yl)-pyridin-3-yl-acetic acid ethyl ester

Bromo-pyridin-3-yl-acetic acid ethyl ester (2.54g; 10.2 mmol), was reacted in refluxing dichloromethane, overnight with 1-methylpiperazine (4.54 mL; 41.0 mmol; 4 equiv) and Et₃N (5.8 mL; 41.0 mmol; 4 equiv). The reaction solution was evaporated to
20 dryness, dissolved in CH₂Cl₂, washed five times with saturated NaHCO₃ (aq), washed once with saturated NaCl (aq), dried with Na₂SO₄, filtered, and concentrated in vacuo. The resulting brown oil was purified with silica gel chromatography in a 3×17 cm glass column using a 2% methanol/chloroform (v/v) mobile phase. A second column eluted with 30% EtOAc/hexanes (v/v) followed by a step gradient of 0.5 – 3%
25 methanol/CH₂Cl₂ (v/v) produced the desired product as a yellow oil (756 mg; 28% yield). MS(ES) calc'd: [M+H]⁺ = 263.1 m/z. Found: 263.1 m/z.

Preparation 34430 2-{3-[(*t*-Butoxy)carbonylamino]cyclohexyl}acetic acid

A compound from preparation 45 (1.0 g, 2.77 mmol) was dissolved in tetrahydrofuran (4 mL) and ethanol (4 mL) under a dry nitrogen atmosphere at room

-176-

temperature. This cloudy white solution became clear and colorless after mixing with 2N NaOH(aq) (15 mL; 19.4 mmol, 11.1 equiv) for 2 h. After rotary evaporation to dryness, the white solid was dissolved in water (20 mL) and the resulting solution extracted with diethyl ether (twice). Acidification of the aqueous layer to pH 2 with 1N HCl(aq) produced a white solid that was extracted into ethyl acetate (thrice). The EtOAc was washed with saturated NaCl(aq), dried with Na₂SO₄(s), and concentrated to dryness by rotary evaporation. The resulting white solid (700 mg, 98% yield) was used in subsequent reactions without further purification. MS(ES) calc'd: [M+Na]⁺ = 280.2 m/z, [M-H]⁻ = 256.2 m/z. Found: 280.1 m/z; 256.2 m/z.

10

Preparation 345

N-((1S,3R)-3-[[phenylmethoxy]carbonylamino]methyl) cyclohexyl)(*t*-butoxy)carboxamide

15

To a solution of a compound from preparation 344 (3.43 g, 13.35 mmol), Et₃N (3.75 mL, 26.96 mmol) in toluene (86 mL) under N₂ was added DPPA (5.8 mL, 26.96 mmol) and benzyl alcohol (4.28 mL, 41.38 mmol). The solution was heated to reflux overnight. The reaction was cooled to rt., diluted with EtOAc, washed (1.0N NaOH then brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (3.05 g, 63%). Mass Spectrum (ES+) (m/z) 263.1 [M-BOC].

20

Preparation 346

N-(*cis*-3(S)-Aminocyclohexylmethyl)(phenylmethoxy)carboxamide

25

A compound from preparation 345 (1.0 g, 2.76 mmol) was treated with TFA (5 mL) under N₂. After 20 min of stirring at rt. the reaction was complete. The crude was then concentrated to an oil which was purified on a Varian Bond-Elut SCX column (10 g). The column was eluted consecutively with CHCl₃, MeOH, and ammonia (2.0M in MeOH). The pure product was recovered from the ammonia fractions. The solvent was removed *in vacuo* to afford 0.632 g (87%) as a colorless oil. MS (ES+) m/z 263.0 (M+H)⁺.

30

-177-

Preparation 347

(3-{[3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carbonyl]-amino}-
cyclohexylmethyl)-carbamic acid benzyl ester

5

To a mechanically stirred solution of crude compound from preparation 346 (20.1 g, 76.6 mmol) and TEA (46.9 g, 459.4 mmol) in 185 ml of dichloromethane was added a compound from preparation 344 (23.1 g, 84.2 mmol) in one portion. The solution warmed slightly and a precipitate began to form. The reaction mixture was stirred at ambient temperature for 19 hours. TLC analysis (20% EtOAc/dichloromethane) showed complete consumption of the starting material. The reaction mixture was diluted with 200 ml of dichloromethane and washed sequentially with 2x 200 ml of 1N hydrochloric acid, 2 x 200 ml of saturated aqueous sodium bicarbonate solution, and 200 ml of half-saturated aqueous NaCl solution. The solution was dried over magnesium sulfate and concentrated to a pale yellow oil, which solidified upon standing. The solid was triturated with 150 ml of diethyl ether, stirred for one hour, filtered, washed with a small amount of diethyl ether, and dried *in vacuo* at 25 °C to provide 29.3 g (76%) of the title compound as a white solid. ¹H NMR (300 MHz, CDCl₃) 7.48 (m, 1H), 7.35 (m, 6H), 7.17 (dt, *J* = 8.5 Hz, 0.9 Hz, 1H), 5.08 (s, 2H), 4.98 (d, *J* = 7.9 Hz, 1H), 4.72 (m, 1H), 3.73 (m, 1H), 3.01 (m, 2H), 2.76 (s, 3H), 1.90 (m, 1H), 1.67 (m, 3H), 1.52 (m, 1H), 1.28 (m, 1H), 0.69 (m, 2H), 0.43 (q, *J* = 12.0 Hz, 1H).

20

Preparation 348

[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-
carbamic acid benzyl ester

25

To a mechanically stirred solution of compound from preparation 347 (27.6 g, 55.2 mmol) in 450 ml of DMF was added a 0.5M solution of potassium bis(trimethylsilyl)amide in toluene (133 ml, 66.3 mmol) over 15 minutes. The resulting purple solution was stirred at ambient temperature for 5 minutes at which time TLC analysis (20% EtOAc/ dichloromethane) indicated that all of the starting material had been consumed. The reaction was diluted with 1 L of EtOAc and quenched with 500 ml of

30

-178-

water. The two layers were stirred together for 15 minutes and separated. The aqueous layer was extracted twice with 250 ml of EtOAc and the combined organic layers were washed twice with 500 ml of water and once with 500 ml of half-saturated NaCl solution. The solution was dried over magnesium sulfate and concentrated to an orange solid. The solid was slurried in a minimal amount of EtOAc, filtered, washed with a small amount of ethyl acetate, and dried in vacuo to afford 17.9 g of the title compound as an off-white solid. An additional 4.4 grams of the title compound were isolated by flash chromatography of the filtrate. Total yield was 22.3 g (84%). ¹H NMR (300 MHz, CDCl₃) 7.40 (m, 2H), 7.33 (m, 6H), 5.06 (s, 2H), 4.85 (m, 1H), 3.16 (m, 2H), 2.89 (s, 3H), 2.56 (br m, 1H), 2.37 (br m, 1H), 1.98 (m, 1H), 1.81 (m, 5H), 1.46 (m, 1H), 1.07 (m, 1H).

Preparation 349

5-(3-Aminomethyl-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one hydroiodide

To a 0°C solution of compound from preparation 348 (5.0 g, 10.4 mmol) in dichloromethane (100 mL) was added iodotrimethylsilane (3.6 mL, 25.0 mmol) dropwise, and the solution allowed to warm to room temperature overnight. Methanol (6.1 mL, 150.0 mmol) was added dropwise over two minutes, and the reaction stirred for 30 minutes. The reaction was concentrated and suspended in ethyl ether. After 15 minutes of sonication, the solids were removed by filtration and washed with ethyl ether. The tan powder was dried on a vacuum pump to give 4.61 g of the title compound as the 1.2 HI salt, 89% yield. MS (ion spray) 346 (M⁺).

Preparation 350

(*t*-Butoxy)-N-[(3-nitrophenyl)methyl]carboxamide

To a suspension of 5.00 g (26.5 mmol) of 3-nitrobenzylamine hydrochloride in 100 mL CH₂Cl₂ at rt. was added 5.79 g (26.5 mmol) of di-*t*-butyl dicarbonate. To this was added 8.13 mL (58.3 mmol) of triethylamine, dropwise over 15 min. The reaction was stirred for 3 h at rt., after which it was diluted with 300 mL of EtOAc. The resulting

-179-

organic solution was washed three times with 1N HCl solution, dried over sodium sulfate and concentrated *in vacuo* to give 6.2 g (93%) of a white solid, which was characterized as the title compound. MS (FIA) $m/z = 253$.

5

Preparation 351

(*t*-Butoxy)-N-[(3-aminocyclohexyl)methyl]carboxamide

To a solution containing 12.0 g (47.7 mmol) of a compound from preparation 350 in 300 mL ethanol was added 6.0 g of rhodium on carbon. The reaction was subjected to hydrogenation (60 psi) at 60 °C for 18 h, after with the catalyst was removed by vacuum filtration and the solvent removed *in vacuo*, giving 9.5 g (87%) of a clear oil. This material was characterized as the title compound and used without further purification. ¹H-NMR is consistent with structure.

15

Preparation 352

N-(3{[*t*-Butoxy)carbonylamino]methyl}cyclohexyl)[4-(2-chloro-6-fluorophenyl)-2-methyl(3-furyl)]carboxamide

To a solution of 9.5 g (41.6 mmol) of a compound from preparation 351 in 200 mL of CH₂Cl₂ was added 11.4 g (41.6 mmol) of a compound from preparation 344, followed by 11.6 mL (83.2 mmol) of TEA at rt. The reaction was stirred at room temperature for 15 h, and concentrated *in vacuo*. The crude solid was dissolved in 200 mL of ethyl acetate and the organic solution was washed twice with 1N HCl solution, dried over sodium sulfate and concentrated *in vacuo* to give a yellow solid. This material was recrystallized from methanol to give 12.5 g (64%) of a white crystalline solid, which was characterized as the title compound, as an inseparable mixture of *cis* and *trans* isomers. MS(FIA) $m/z = 466.2$.

-180-

Preparation 353

(*t*-Butoxy)-N-{[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]methyl}carboxamide

5 To a stirred solution of 7.00 g (15.0 mmol) of a compound from preparation 352 in 200 mL of DMF at rt. was added 30.0 mL (15 mmol) of a 0.5 M toluene solution of potassium bis(trimethylsilyl) amide over 10 min. The resulting dark red reaction was stirred an additional 5 min at rt. and added to 200 mL of 1N HCl. The two phase solution was diluted with 400 mL of ethyl acetate and the organic layer was separated. The
10 organic solution was washed four times with brine, dried over sodium sulfate and concentrated *in vacuo* to give an orange solid. This material was recrystallized from toluene to give 3.4 g of a light yellow solid, which was characterized as pure racemic *cis* material, (a). MS(FIA) *m/z* = 446.1. Purification of the racemic *trans* material (b) was accomplished by concentrating the mother liquor and subjecting this material to flash
15 chromatography on silica gel, using 50% hexane-EtOAc as the eluent. The major fractions were combined to give a light yellow solid, which was characterized as pure racemic *trans* material. MS(FIA) *m/z* = 446.1.

Preparation 354

20 N-{3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]methyl}(*t*-butoxy)carboxamide

The racemic material of a compound from preparation 353 was separated into its enantiomers by chiral HPLC chromatography using a Chiralpak AD column and 10%
25 ethyl alcohol-heptane as the eluent at a flow of 1.0 mL/min.

Retention Time (Enantiomer 1) = 10.501 min.

Retention Time (Enantiomer 2) = 12.576 min.

-181-

Preparation 355

N-[3-(aminomethyl)cyclohexyl]-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one

5 A slurry of a compound from 354 (1.2 g, 2.7 mmol) in 20 mL of acetic acid saturated with hydrochloric acid was stirred for 4 h at rt., then concentrated to dryness. The residue was dissolved in toluene and concentrated to dryness. The residue was slurried in ether/hexanes, concentrated to dryness, and dried under vacuum to yield 750 mg (80%) of the desired mixture of isomers as a tan solid. MS (ion spray) 346.2 (M+1).

10

Preparation 357

1-Methyl-piperidine-3-carboxylic acid

15 To a solution of ethyl 1-methylnipecotate, 0.3 mL (1.65 mmol) in 10 mL of dioxane was added a solution of 118 mg (4.95 mmol) of lithium hydroxide in 10 mL of water. The solution was stirred for 1.5 hours at rt. then was acidified to pH = 1 with 5N HCl. The mixture was concentrated to dryness and carried on without further purification. IS (MS) 144.1 (M+1).

20

Preparation 358

Piperidine-1,3-dicarboxylic acid 1-tert-butyl ester

25 To a solution of nipecotic acid, 2.0 g (16.0 mmol) in 20 mL of THF and 20 mL of water was added 4.7 g (33.6 mmol) of potassium carbonate and 3.5 g (16.0 mmol) of Boc-anhydride. The resulting solution was stirred overnight at ambient temperature then concentrated to dryness. The residue was dissolved in water, washed with ether and acidified to pH = 2 with 5N HCl. The acidic aqueous layer was extracted with 20% isopropanol/chloroform. The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness to yield 3.2 g (86%) of the desired mixture of isomers as a white solid. MS (ion spray) 228.2 (M-1).

30

-182-

Preparation 359

{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-methyl}-carbamic acid tert-butyl ester

5 To a solution of a compound from preparation 69, 130 mg (0.28 mmol) in 10 mL of DMF was added 59 mg (0.34 mmol) of N-t-butoxycarbonylglycine, 46 mg (0.34 mmol) of 1-hydroxy-7-azabenzotriazole, 66 mg (0.34 mmol) of EDC, 5 mg of DMAP and 0.120 mL (0.84 mmol) of Et₃N. The reaction mixture was stirred overnight at rt. and was concentrated to dryness. The residue was dissolved in 20% isopropanol/chloroform, washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by radial chromatography using a MeOH/chloroform gradient and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to yield 97 mg (71%) of the desired isomer as a white foam. MS (ion spray) 489.1 (M+).

15

Preparation 360

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide hydrochloride

20 A solution of the compound from preparation 359, 49 mg (0.10 mmol) in 10 mL of HCl-saturated acetic acid was stirred 3 hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to give a quantitative yield of the desired isomer as a tan foam. MS (ion spray) 389.1 (M+).

25

Preparation 361

2-(Pyridin-3-yloxy)-hexanoic acid ethyl ester

To a solution of ethyl-3-bromohexanoate, 5.0 g (22.4 mmol) in 200 mL of tetrahydrofuran was added 4.6 g (26.9 mmol) of 3-hydroxypyridine sodium salt. The mixture was refluxed for four hours and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water. The organics were washed

30

-183-

with saturated aqueous sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered, and concentrated to dryness. The residue was purified by column chromatography using a methanol/chloroform gradient as eluent and concentrated to dryness to yield 1.82 g (34%) of the desired mixture of isomers as a brown liquid. MS (ion spray) 238.2 (M+1).

Preparation 362

2-(Pyridin-3-yloxy)-hexanoic acid

10 A solution of lithium hydroxide, 200 mg (8.82 mmol) in 10 mL of water was added to a solution of compound from the preparation 361, 700 mg (2.94 mmol) in 10 mL of dioxane. The resulting solution was stirred at ambient temperature for 3 hours, acidified to pH = 1 with 5 N HCl and concentrated to dryness. Crude MS (ion spray) 210.0 (M+1) for desired mix of isomers. Crude product was carried on as is without
15 further purification.

Preparation 363

(Pyrimidin-2-yloxy)-acetic acid ethyl ester

20 To a slurry of sodium hydride (60% dispersion in mineral oil), 280 mg (5.76 mmol) in 10 mL of DMF at 0°C was added 0.52 mL (4.8 mmol) of ethyl glycolate. After 10 minutes, 0.92 g (5.76 mmol) of 2-bromopyrimidine was added. The reaction mixture was stirred 15 min at 0°C then ice bath was removed. After four hours, reaction mixture
25 was quenched with methanol and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and water, then extracted with 20% isopropanol/chloroform. The combined organics were washed with saturated aqueous sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered, and concentrated to dryness. The residue was purified by radial chromatography using a
30 methanol/chloroform gradient as eluent and concentrated to dryness to yield 0.45 g (52%) of the desired product as a colorless oil. MS (ion spray) 183.1 (M+1).

-184-

Preparation 364

(Pyrimidin-2-yloxy)-acetic acid

5 A solution of lithium hydroxide, 260 mg (11.0 mmol) in 5 mL of water was added to a solution of 400 mg (2.2 mmol) of compound from preparation 363 in 5 mL of dioxane. The mixture was stirred for four hours at ambient temperature, then was acidified to pH = 1 with 5 N HCl and concentrated to dryness. The residue was carried on without further purification. MS (ion spray) 155.1 (M+1).

10

Preparation 365

3-Methyl-5-(2-chloro-6-fluorophenyl)-4-isoxazolecarboxylic acid ethyl ester

15 To a solution of ethyl 3-aminomethyl crotonate (4.79 g, 33.5 mmol) in toluene (10 mL), was added triethylamine (3.73 g, 37 mmol). The solution was chilled using an ice bath, and then 2-chloro-6-fluorobenzoyl chloride (6.47 g, 33.5 mmol) was added dropwise over a 20 min period. The reaction was allowed to warm slowly to r.t., and stirred for 24 hr. The resulting suspension was then filtered, and the filtrate diluted with ethyl acetate (100 mL) and transferred to a separatory funnel. The organic layer was sequentially washed with water, brine, dried (sodium sulfate), and the volatiles removed under reduced pressure to provide 2-(2-chloro-6-fluorobenzoyl)-3-methylamino-but-2-enoic acid ethyl ester (9.46 g) as a golden solid, and primarily one geometrical isomer. MS (ES) m/z 299.9 (M+H)⁺.

20

25 Crude adduct was then redissolved in glacial HOAc (50 mL) to which was added NH₂OH.HCl (1.8 g, 1.1 eq). The solution was then heated to reflux for 40-45 min to effect isoxazole formation. The reaction mixture was concentrated to an oil, diluted with ether, and transferred to a sep. funnel. The organic phase was washed with saturated bicarbonate, brine, then dried. Filtration and concentration afforded crude isoxazole ethyl ester (7.5 g), which could be used without further purification. MS (+ ES) m/z 283.9 (M+H)⁺.

30

-185-

Preparation 366

4-Isoxazolecarboxylic acid, 3-Methyl, 5-(2-chloro-6-fluorophenyl)

Hydrolysis of a compound from preparation 365 was accomplished by dissolving
5 the crude ester (7.5 g, approx. 0.027 mol) in THF (250 mL), and adding aq. LiOH (1.344
g in 100 mL, 2 eq). After stirring overnight at r.t., the solution was concentrated to 2/3rd
volume, diluted with EtOAc (200 mL) and 50 mL water, transferred to a separatory
funnel, and the aqueous phase collected. The organic phase was washed twice, and the
combined aqueous phase was then acidified with 5N HCl. Back extraction with three
10 washings of EtOAc was then followed with a brine wash of the combined organics. After
drying over Na₂SO₄, filtration and concentration, clean isoxazole acid was obtained (2.94
g). MS (-ES) m/z 253.8, 255.8 (M-H)⁻.

Preparation 367

15 5-(2-Chloro-6-fluoro-phenyl)-3-methyl-isoxazole-4-carbonyl chloride

To a solution of a compound from preparation 366 (60 mg, 0.234 mmol) in
benzene (4 mL) containing catalytic amount of pyridine (20 μl) was added oxalyl chloride
(23 mL). After heating to reflux for 1 hr, an aliquot was concentrated under vacuum. 1H-
20 NMR (CDCl₃) 7.53 (d of t, 1H), 7.37 (d, 1H), 7.17 (t, 1 H). (racemic)

Preparation 368

1,3-cyclohexanedicarboxylic acid

25 To a suspension of isophthalic acid (500 g, 3 mol) in methanol (2.8 l) was added
5% Rhodium-on-alumina catalyst (50 g) and acetic acid (150 ml). The reaction mixture
was shaken under hydrogen (50 psi) at room temperature overnight. The mixture was
filtered through celite. To this solution was added fresh 5% Rhodium-on-alumina catalyst
(25 g), and the mixture was shaken under 50 psi of hydrogen for another 24 hours. The
30 final reaction mixture was filtered through celite. The solution was concentrated under
vacuum to give 493 g of the title compound as a white powder (96.3 % yield). m.p. 163-
165 °C.

-186-

Preparation 369

3-Oxabicyclo[3.3.1]nonane-2,4-dione

5 A solution of dicyclohexylcarbodiimide (200 g, 1.16 mol) in CH₂Cl₂ (1000 ml) was added dropwise to a suspension of compound from preparation 368 (257 g, 1.25 mol) in CH₂Cl₂ (550 ml), and the mixture was stirred at room temperature for 4 hours. The precipitated dicyclohexylurea was filtered and washed several times with cold CH₂Cl₂ (200 ml x 3). The combined organic layer was concentrated to give a white solid, which
10 was suspended in MTBE (900 ml). This solid was collected by filtration, washed with MTBE (250 ml), and dried under house vacuum to give the title compound (137 g). The filtrate was concentrated to a residue, which was suspended in MTBE (250 ml) to give another 31 g anhydride. The total yield was 168 g (94%). m.p. 138-140°C.

15

Preparation 370

cis-1,3-Cyclohexanedicarboxylic acid diethyl ester

To a solution of compound from preparation 369 (31 g, 0.2 mol) in ethanol (anhydrous, 310 ml) was added p-toluenesulfonic acid monohydrate (1.9 g, 10 mmol, 0.05
20 equiv.) and triethyl orthoformate (50 ml, 0.3 mol). The reaction mixture was stirred at 60°C overnight. The volatiles were stripped and the residue was diluted with ethyl acetate (250 ml), washed with water (120 ml) and brine (100 ml), and dried over MgSO₄. After filtration and evaporation, the residue was purified by chromatography. Eluting the column with 10% ethyl acetate in hexane afforded the title compound (40 g, 87.7% yield).
25 ¹H NMR: (500 MHz, CDCl₃) δ 4.11 (q, J=7.0 Hz, 4H), 2.29 (dt, 2H), 2.11 (dd, 1H), 1.97 (m, 2H), 1.98 (m, 1 H), 1.53 (q, J=12.5 Hz, 2H), 1.30-1.40 (m, 2H), 1.25 (t, J=7.0 Hz, 6H).

-187-

Preparation 371

1,3-Cyclohexanedicarboxylic acid, monoethyl ester (1R, 3S)

To a suspension of compound from preparation 370 (40 g, 17.5 mmol) in pH 7.2 phosphate buffer [0.2 M] (1.2 l) was added lipase AY30 (Amano, 16.7 g). The mixture was stirred vigorously at room temperature for 30 hours. The mixture was acidified with 10-15% HCl to pH < 2, and extracted with ethyl acetate (500 ml x 2). The combined organic solution was washed with aqueous 10% Na₂CO₃ (100 ml x 2) and water (100 ml x 2). The combined aqueous layers were washed again with ethyl acetate (150 ml) and then acidified with 10-15% HCl to pH < 2. The acidified aqueous was then extracted with ethyl acetate (150 ml x 3). The combined organic solution was dried over MgSO₄. After filtration and concentration the title compound (35.6 g, 100% yield) was obtained. ¹H NMR (500 MHz, CDCl₃) δ 4.12 (q, J=7.0 Hz, 2H), 2.20-2.40 (m, 3H), 1.85-2.05 (m, 3H), 1.5 (q, 2H), 1.35 (m, 2 H), 1.24 (t, J=7.0 Hz, 3H), [α]_D +3.2 °, [α]₃₆₅ +10.4 ° (c, 0.434; CHCl₃). [α]_D +3.0 °, [α]₃₆₅ +9.6 ° (c, 0.532; CH₃OH).

Preparation 372

Ethyl-[3-N-(methylcarbamate)-cyclohexyl]-carboxylate (1R, 3S)

A solution of a compound from preparation 371 (73 g, 365 mmol) in toluene (750 ml) was heated to reflux using a Dean-Stark trap to separate trace amounts of water. After collecting about 10 ml of water, the mixture was cooled down to about 40-50°C. To this mixture was added triethylamine (56 ml, 0.4 mol), and diphenylphosphoryl azide (86.5 ml, 0.4 mol). The reaction mixture was stirred at 110 °C for 60 min, cooled to 70 °C, and methanol (64 g, 2 mol) was added with stirring. After addition, the final reaction mixture was then heated to 85°C overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate (700 ml) and washed with water (500 ml). The aqueous layer was extracted with ethyl acetate (500 ml x 2). The combined organic solution was washed again with water (500 ml) and brine (500 ml). After drying over MgSO₄ and concentration under reduced pressure, the title compound was obtained as a colorless oil (86 g, 100 %). ¹H NMR: (300 MHz, CDCl₃) δ 4.60 (sb, 1H), 4.13 (q, 2H),

-188-

3.65 (s, 3H), 3.50 (sb, 1H), 2.38 (t, 1H), 2.23 (d, 1H), 2.00-1.80 (m, 3H), 1.24 (t, 3H), 1.12-0.95 (m, 1H).

Preparation 373

5 Ethyl-((1R, 3S)-3-{{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonyl-amino}cyclohexyl)-carboxylate

To a solution of compound from preparation 372 (86 g, 365 mmol) in CH₂Cl₂ (750 ml) was added iodotrimethylsilane (100 g, 500 mmol) in one portion, at room
10 temperature. The reaction mixture was stirred for 2 hours at ambient temperature, cooled to 0-5 °C, and methanol (50 ml) was added. After stirring 15 minutes, the solution was concentrated under reduced pressure. The residue was dissolved in THF (1l). To this solution was added water (0.5 l), potassium carbonate (138 g, 1 mol), and a solution of a
15 compound from preparation 27 (110 g, 0.4 mol) in 250 ml THF, dropwise. After the addition, the reaction mixture was heated to room temperature and stirred for 12 hours. THF was removed under house vacuum, water (250 ml) was added, and the mixture was extracted with ethyl acetate (500 ml x 3). The combined organic solution was washed with saturated sodium thiosulfate (150 ml), water (500 ml), brine (500 ml) and then dried
20 over MgSO₄. After filtration and evaporation under vacuum, the residue was purified by recrystallization from MTBE (250 ml). Repeating this recrystallization procedure three times provided the title compound (122.7 g, 82.5% yield) as a white powder. M.S. m/z 409 (M⁺, 100 %).

Preparation 374

25 Ethyl [(1R,3S)-3 -(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quindin-5-yl)-cyclohexyl]-carboxylate

To a solution of compound from preparation 373 (78 g, 190 mmol) in DMF (750 ml) was added a solution of KHMDS ([0.5M], 400 ml, 200 mmol). The temperature was
30 kept at 25 °C by using an ice-bath. After the addition was complete, the reaction mixture was analyzed by TLC (silica gel, 50 % EtOAc in hexane) and found to be complete. Water (1 l) was added and the mixture was extracted with EtOAc (800 ml x 3). The

-189-

combined organic solution was washed with 1N HCl (250 ml), water (250 ml), brine (250 ml), dried over MgSO₄ and concentrated to give a residue which was purified by recrystallization from MTBE (500 ml) to afford 66 g of the title compound as a light yellow powder (89.0% yield). M.S. m/z 389 (M⁺+1, 100 %).

5

Preparation 375

(1R, 3S) 3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl
carboxylic acid

10 To a solution of compound from preparation 374 (62 g, 160 mmol) in THF (600 ml) was added 5N aqueous sodium hydroxide (120 ml) at room temperature. The reaction mixture was heated to 60 °C for 15 hours with stirring. After cooling to room temperature, water (750 ml) was added and the mixture was washed with ethyl acetate (500 ml). The aqueous phase was separated and acidified with 15% HCl to pH < 2. The
15 aqueous phase was then extracted with methylene chloride (1000 ml x 3). The combined organic extracts were washed again with water (500 ml), brine (500 ml), and dried over MgSO₄. After filtration and evaporation under vacuum, the dark brown residue was suspended in MTBE (1000 ml), and refrigerated overnight. The mixture was filtered to afford 55.45 g (96.4%) of bright yellow product. M.S. m/e 361 (M⁺, 50 %), 225 (100 %).

20

Preparation 376

2-methyl [(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4, 3-c]quinolin-5-yl)-
cyclohexyl] carbamate

25 To a suspension of compound from preparation 375 (55.4 g, 154 mmol) in toluene (1l) was added triethylamine (23.7 ml, 170 mmol), and diphenylphosphoryl azide (36.5 ml, 170 mmol). The reaction mixture was stirred at 110 °C for 2 hours during which time a solution formed. The solution was cooled to 80 °C and methanol (25 g, 0.77 mol) was added with stirring. The solution was warmed to 85 °C for 22 hours. After cooling to
30 room temperature, the toluene was removed under reduced pressure and the residue was dissolved in dichloromethane (3l) and washed with water (1l). The aqueous phase was extracted with dichloromethane (1l x 2) and the combined organic solution was washed

-190-

again with water (500 ml) and brine (500 ml). After drying over MgSO_4 , the solution was concentrated under vacuum. The crude product was purified by crystallization ($\text{CH}_2\text{Cl}_2/\text{MTBE}$, 0.5 l/2 l) to afford the title compound (46.6 g, 78.2 %). $[\alpha]_{\text{D}} +49.2^\circ$, $[\alpha]_{365} +263.3^\circ$ (c, 0.56; CHCl_3). The filtrate was concentrated to a residue, which was purified by chromatography to obtain a second crop of product (5.1 g). The total yield was 86.8 %. M.S. m/z 476 (M^+ , 25%).

Preparation 377

10 [3-(3-Acetyl-4-amino-5-chloro-2-oxo-2H-quinolin-1-yl)-cyclohexyl]-carbamic acid methyl ester

A compound from preparation 376 (0.1 g, 0.26 mmol) and $\text{Mo}(\text{CO})_6$ (0.13 g, 0.51 mmol) were combined in a solution of acetonitrile (5 mL) and water (1 mL). The reaction mixture was heated to reflux while stirring. After stirring for 2 hr the reaction was complete. A 10% aqueous EDTA solution was added and stirred for an additional 30 minutes. The reaction was then transferred to a separatory funnel and washed with water (5 x 10 mL). The organic solution was filtered through celite and concentrated to a dark brown solid under vacuum. The solid was diluted in EtOAc and was purified by silica gel column chromatography using 50% EtOAc in Hexanes to elute the product. The solvent was removed *in vacuo* to afford 0.078 g of desired product as a white solid. MS (ES+) m/z 392.2 ($\text{M}+\text{H}^+$), (ES-) 390.2 ($\text{M}-\text{H}^-$).

Preparation 378

25 [3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-carbamic acid methyl ester

A solution of a compound from preparation 377 (0.05 g, 0.13 mmol) in acetic acid (5 mL) was treated with hydroxylamine hydrochloride (13 mg, 0.19 mmol). The solution was heated to reflux and stirred 5 hr. The reaction was then diluted in EtOAc (50 mL) and washed with water (3 x 10 mL), sat'd sodium bicarbonate (3 x 10 mL), and brine (2 x 10 mL). The organic was dried over sodium sulfate and the solvent removed. The crude product was purified by silica gel column chromatography using 50% EtOAc in Hexanes

-191-

to elute the product. The solvent was removed *in vacuo* to afford 0.028 g (56%) of title compound as a white solid. MS (ES+) m/z 390.1 (M+H)⁺, (ES-) 388.2 (M-H)⁻.

Preparation 380

5 5-(3-Amino-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

A solution of a compound from preparation 378 (0.38 g, 1.0 mmol) in CH₂Cl₂ (25 mL) was treated with TMSI (0.29 g, 1.5 mmol). The solution was stirred for 2 hr at room temperature. The solution was then concentrated to a solid and the solid was diluted in
10 EtOAc (100 mL). The organic solution was washed with 1.0 N HCl (5 x 20 mL) and the aqueous was adjusted to pH~12 with 5.0 N NaOH. The aqueous was then extracted with CH₂Cl₂ and the solvent was removed to afford 0.26 g (80%) of the title compound as a white solid. MS (ES+) m/z 332.1 (M+H)⁺.

15 Preparation 390

9-Chloro-3-methyl-5-(3-methylaminomethyl-cyclohexyl)-5H-isoxazolo[4,3-c]quinolin-4-one

To a solution of a compound from the preparation 48 (50 mg, 0.12 mmol) in chloroform (0.6 mL) was added trifluoroacetic acid (0.6 mL) and triethylsilane (57 μL, 0.35 mmol)
20 dropwise. The solution was stirred at rt. under nitrogen overnight. The solution was concentrated and dissolved in chloroform (1 mL) and aqueous 1N HCl solution (1 mL). To this mixture was added dichloromethane (10 mL) and saturated sodium bicarbonate solution (10 mL). The layers were separated, and the aqueous layer was extracted with dichloromethane (x3). The combined organic layers were dried over magnesium sulfate
25 and concentrated to give 13.7 mg of the title compound as a brown solid, 32% yield. ¹H NMR: consistent with structure. MS (ion spray) 360 (M⁺).

Preparation 391

Hydroxy-pyridin-2-yl-acetic acid methyl ester

30

To a dry flask was added 2-pyridine-carboxaldehyde (14.4 mL, 150 mmol) and zinc iodide (5 mg, 0.02 mmol), followed by dropwise addition of trimethylsilyl cyanide

-192-

(20.7 mL, 152 mmol) with rapid stirring. The solution was stirred under nitrogen at rt. overnight. The solution was treated with 9N HCl (50 mL, aqueous solution), and stirred under nitrogen at reflux overnight. The solution was partitioned between water and 20% isopropanol/chloroform, and the aqueous layer concentrated to an orange solid. Methanol was added, the mixture was sonicated, and placed in a 0°C-freezer overnight. Ammonium chloride was removed by filtration (x2), and the solution concentrated. The solution was diluted with water (500 mL), and basified to pH 8 with aqueous 5N NaOH. The aqueous layer was extracted with 20% isopropanol/chloroform (x3), and the organic layers were washed with brine, dried over magnesium sulfate and concentrated to give 14.02 g of yellow solid. The material was sonicated as a suspension in diethyl ether and filtered to remove baseline impurities. Purification by silica gel chromatography (eluting with 0-1% methanol/chloroform) gave 4.10g of the title compound as a yellow solid, 16% yield. ¹H NMR: consistent with structure. MS (ion spray) 167 (M⁺).

15

Preparation 392

(tert-Butyl-dimethyl-silanyloxy)-pyridin-2-yl-acetic acid methyl ester

To a solution of a compound from preparation 391 (4.10 g, 24.5 mmol) in DMF (35 mL) was added imidazole (3.34 g, 49.0 mmol) and t-butyldimethylsilyl chloride (7.39 g, 49.0 mmol). The reaction was stirred at rt. overnight under nitrogen. The reaction was concentrated and dissolved in chloroform, washed with saturated aqueous sodium bicarbonate solution, brine, dried over magnesium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 10-20% ethyl acetate/hexane) gave 6.07 g of the title compound as a clear oil, 88% yield. ¹H NMR: consistent with structure. MS (ion spray) 281 (M⁺).

25

Preparation 393

Hydroxy-pyridin-3-yl-acetic acid methyl ester

To a dry flask in an ice bath was added 3-pyridine-carboxaldehyde (14.5 mL, 150 mmol) and zinc iodide (5 mg, 0.02 mmol), followed by dropwise addition of trimethylsilyl cyanide (20.7 mL, 152 mmol). The solution was stirred under nitrogen at rt.

30

-193-

overnight. The solution was treated with 9N HCl (50 mL, in water), and stirred under nitrogen at reflux overnight. The solution was partitioned between an aqueous 1N HCl solution and chloroform, and the aqueous layer concentrated. Ethanol was added, the mixture was heated to 45°C, and ammonium chloride was removed by filtration, and the solution concentrated. To a 0°C flask was added methanol (300 mL) followed by dropwise addition of thionyl chloride (40 ml, 548 mmol). After ten minutes, the concentrated reaction mixture was added dropwise as a solution in methanol (300 mL), and the reaction was allowed to warm to rt. overnight. The solution was concentrated, diluted with water (500 mL), and basified to pH 9 with a saturated aqueous sodium bicarbonate solution. The aqueous solution was extracted with 20% isopropanol/chloroform (x5), and the organics dried over magnesium sulfate and concentrated to give 18 g of red oil. Purification by silica gel chromatography (eluting with 0-2% methanol/chloroform) gave 11.51g of the title compound a yellow oil, 46% yield. ¹H NMR: consistent with structure. MS (ion spray) 167 (M⁺).

15

Preparation 394

1-tert-Butoxycarbonylamino-cyclopentanecarboxylic acid

To a suspension of 1-amino-1-cyclopentane carboxylic acid (4.52 g, 35.0 mmol) in THF (50 mL) and water (50 mL) was added potassium carbonate (14.51 g, 105.0 mmol), di-t-butyl-dicarbonate (7.72 g, 35.4 mmol), and DMAP (5 mg, 0.04 mmol). The reaction was stirred at rt. under nitrogen overnight. The THF was removed *in vacuo*, and the remaining aqueous solution was acidified to pH 5.5 with an aqueous 5N HCl solution. The aqueous solution was extracted with 20% isopropanol/chloroform (x6), dried over magnesium sulfate, and concentrated to give 1.01g of the title compound as a white solid, 13% yield. ¹H NMR: consistent with structure. MS (ion spray) 229 (M⁺).

25

Preparation 395

(4-Acetyl-piperazin-1-yl)-pyridin-3-yl-acetic acid ethyl ester

30

N-acetyl-piperazine (3.7 g; 28.8 mmol; 2.8 equiv) was combined with bromo-pyridin-3-yl-acetic acid ethyl ester (approx. 2.5 g; 10 mmol). Purification on a 3×16 cm

-194-

silica column using a 30% then 40% ethyl acetate/hexanes (v/v) mobile phase followed by a 2% methanol/dichloromethane (v/v) mobile phase resulted in 1.84 g light yellow oil (approx. 60% yield). MS(ES) calc'd: $[M+H]^+ = 292.2$ m/z; $[M+Na]^+ = 314.2$ m/z. Found: 292.1 m/z; 314.1 m/z.

5

Preparation 396

Pyridin-3-yl-(pyridin-2-yloxy)-acetic acid ethyl ester

Sodium hydride (520 mg; 21.8 mmol; 1.4 equiv) was stirred with 2-hydroxypyridine (1.73 g; 18.2 mmol; 1.2 equiv) in DMF (30 mL) for 15 min. This solution was added to an aliquot of bromo-pyridin-3-yl-acetic acid ethyl ester (approx. 3.6 g; 15 mmol) in DMF (20 mL). After stirring overnight, at rt., the resulting opaque black solution was concentrated in vacuo, dissolved in 20% isopropyl alcohol/chloroform, washed 5 times with saturated $\text{NaHCO}_{3(aq)}$, washed twice with saturated $\text{NaCl}_{(aq)}$, and dried over $\text{Na}_2\text{SO}_{4(s)}$. After filtration, the solution was concentrated in vacuo and purified by SiO_2 column chromatography on two consecutive columns with a 1% methanol/chloroform mobile phase and then an ethyl acetate/hexanes mobile phase (30%, 40%, 50% step gradient). A light yellow oil (134 mg; approx. 3% yield) was isolated. MS(ES) calc'd: $[M+H]^+ = 259.2$ m/z. Found: 259.1 m/z.

20

Preparation 397

Pyridin-3-yl-(pyridin-3-yloxy)-acetic acid ethyl ester

The product was prepared in a manner similar to preparation 396 using the pre-formed sodium salt of the 3-hydroxypyridine (2.13 g; 18.2 mmol; 1.2 equiv) and an aliquot of bromo-pyridin-3-yl-acetic acid ethyl ester (approx. 3.6 g; 15 mmol) in dimethylformamide (20 mL). Column purification over SiO_2 (0.5% then 1.5% methanol/chloroform) produced a yellow/orange oil (2.02 g; approx. 50% yield). MS(ES) calc'd: $[M+H]^+ = 259.2$ m/z; $[M-H]^- = 257.2$ m/z. Found: 259.1 m/z; 257.1 m/z.

30

-195-

Preparation 398

Pyridin-3-yl-(pyridin-4-yloxy)-acetic acid ethyl ester

The product was formed in a manner similar to preparation 396 using the in situ-
5 generated sodium salt of the 4-hydroxypyridine (2.13 g; 18.2 mmol; 1.2 equiv) and an
aliquot of bromo-pyridin-3-yl-acetic acid ethyl ester (approx. 3.6 g; 15 mmol) in DMF (20
mL). Column chromatographic purification over SiO₂ (1% methanol/chloroform)
produced a yellow/orange oil (320 mg; approx. 8% yield). MS(ES) calc'd: [M+H]⁺ =
259.2 m/z. Found: 259.1 m/z.

10

Preparation 399

5-(3-Amino-cyclohexyl)-9-chloro-3-methyl-2,5-dihydro-pyrazolo[4,3-c]quinolin-4-one

A solution of a compound from preparation 379 (0.150 g, 0.39 mmol) in CH₂Cl₂
15 (20 mL) was treated with TMSI (0.116 g, 0.58 mmol) and stirred at rt. for 6 hr. The
reaction was then treated with MeOH (5 mL) and stirred for an additional 20 min. The
solution was concentrated to an oil and then ether was added. The solid was filtered to
afford 0.095 g (74%) of the title compound as an orange solid. MS (ES+) m/z 331.2
(M+H)⁺.

20

Preparation 4003-(2-Chloro-6-fluoro-phenyl)-3-oxo-2-(triphenyl-15-phosphanylidene)-propionic acid
ethyl ester

To a solution of 20.0 g (0.115 mol) of 2-chloro-6-fluorobenzoic acid in 150 mL
25 methylene chloride and 0.50 mL of DMF was added 11.0 mL (0.126 mol) oxalyl chloride,
dropwise at rt. over 15 minutes. The reaction was stirred at rt. for one hour and
concentrated *in vacuo*. The resulting acid chloride was redissolved in 50 mL of toluene.
In a separate flask, 34.0 mL (0.138 mol) of bis(trimethylsilyl) acetamide was dissolved in
30 300 mL of toluene. To this was added 40.0 g (0.115 mol) of (carbethoxymethylene)-
triphenylphosphorane. The resulting suspension was cooled to 10°C and stirred as 3-(2-
chloro-6-fluorophenyl)-5-methylisoxazole-4-carbonyl chloride in 50 mL of toluene was

-196-

added dropwise. The cold bath was removed and the reaction was allowed to stir at rt. for 18 hours. The mixture was concentrated *in vacuo* and to the resulting solid was added 50 mL of toluene. The white solid was collected by vacuum filtration to give 49.0 g (84%) of the title compound. MS(ES+) (m/z) 505.1 [M+1].

5

Preparation 401

3-(2-Chloro-6-fluoro-phenyl)-2,3-dioxo-propionic acid ethyl ester

A solution of 25.0 g (49.5 mmol) of the compound from preparation 400 in 125 mL of methylene chloride was cooled to -78°C . Into the cooled solution was passed ozone, prepared using an ozone generator (Griffin Tech. Corp.) and oxygen at 1.8 amps. The ozone was bubbled through the cooled solution using a gas dispersion tube. The reaction was monitored visually, and determined to be completed after 45 minutes, due to the characteristic blue color that is associated with an ozone saturated solution of methylene chloride. The reaction was purged with nitrogen gas and warmed to rt. The organic solution was washed twice with brine, dried over sodium sulfate and concentrated *in vacuo*. This crude material was purified by flash chromatography (silica gel, 50% hexanes-ethyl acetate) to give 13.0 g (98%) of the title compound as a yellow oil. Anal. Calcd. for $\text{C}_{11}\text{H}_8\text{O}_4\text{FCl}$: C, 51.08; H, 3.12. Found: C, 50.67; H, 3.12.

20

Preparation 402

5-(2-Chloro-6-fluoro-phenyl)-3H-imidazole-4-carboxylic acid ethyl ester

A solution containing 6.80 g (26.0 mmol) of a compound from preparation 401 in 50 mL of glacial acetic acid was added to a slurry of 20.3 g (260 mmol) of ammonium acetate in 100 mL of glacial acetic acid. To this mixture was added 3.95 g (130 mmol) of paraformaldehyde. The mixture was stirred and heated at 80°C for one hour and concentrated *in vacuo*. The resulting crude material was taken up in ethyl acetate and washed with saturated sodium bicarbonate (3X). The organic solution was dried over sodium sulfate and concentrated *in vacuo* to give an oily yellow solid. This crude material was precipitated out of 50% ethyl acetate-hexane to give 3.35 g (48%) of the title compound as a light yellow amorphous solid. MS (ES+) (m/z) 269.0 [M+1].

30

-197-

Preparation 403 & 404

5-(2-Chloro-6-fluorophenyl)-3-methyl-3H-imidazole-4-carboxylic acid ethyl ester
(isomer 1)

5 5-(2-Chloro-6-fluorophenyl)-1-methyl-1H-imidazole-4-carboxylic acid ethyl ester
(isomer 2)

To a solution containing 2.90 g (10.8 mmol) of the compound from preparation 402 in 15 mL of DMF was added 2.98 g (21.6 mmol) of solid anhydrous potassium carbonate, followed by 1.34 mL (21.6 mmol) of methyl iodide, with stirring, at rt. The reaction was stirred at rt. for one hour, diluted with 100 mL of ethyl acetate and washed four times with brine. The resulting crude material was purified by flash chromatography (silica gel, hexanes-ethyl acetate gradient) and two main products were separated. The major product contained 1.62 g (53%) and was identified as the title compound (isomer 1). The isomeric minor product, which was identified as the title compound (isomer 2) was found to contain 933 mg (31%).

isomer 1: Mass spectrum (ES+) (m/z) 283.0 [M+1].

Anal. Calcd. for C₁₃H₁₂N₂O₂FCl: C, 55.23; H, 4.28; N, 9.91. Found: C, 55.14; H, 4.26; N, 9.75.

20 isomer 2: Mass spectrum (ES+) (m/z) 283.0 [M+1].

Anal. Calcd. for C₁₃H₁₂N₂O₂FCl: C, 55.23; H, 4.28; N, 9.91. Found: C, 55.08; H, 3.96; N, 9.77.

Preparation 405

25 5-(2-Chloro-6-fluoro-phenyl)-3-methyl-3H-imidazole-4-carboxylic acid

To a solution of 95 mg (0.336 mmol) of a compound from preparation 403. in 5 mL methanol was added 5 mL of 1N sodium hydroxide solution. The reaction was stirred at rt. for 18 hours. The pH of the reaction was adjusted to pH ~ 1 with 1N HCl and this material was concentrated *in vacuo* by adding 100 mL of toluene to aid in the azeotropic removal of water. The subsequent white solid, which contained the free acid. The crude title compound was used without further purification.

-198-

General Procedures, unless otherwise specified:a) for condensation of a primary amine with a carboxylic acid

To "Starting Material A," the primary amine of interest (1 eq.), in anhydrous DMF at 0 °C was added "Starting Material B," the carboxylic acid (3 eq.), collidine (3 eq.) and BOP (3 eq.). The cold bath was removed and after 2 h. The reaction was diluted with EtOAc (25 mL) and saturated NaHCO₃ (25 mL). The organic layer was washed with distilled H₂O (2x25 mL), dried over Na₂SO₄, and chromatographed on silica gel.

10 b) for acylation of primary amines

To a solution of "Starting Material A," the primary amine (1 eq.), in dichloromethane was added "Starting Material B," the acylating reagent (1 eq.), triethylamine (1 eq.), and 4-dimethylaminopyridine (10% mol.eq.). The mixture was stirred overnight at ambient temperature and chloroform was added. The solution was washed with 1N hydrochloric acid, saturated sodium bicarbonate, then brine; dried over sodium sulfate, filtered, and the solvent was removed *in vacuo*. The residue was chromatographed on silica gel to afford the desired compound.

20 c) for coupling of primary amines and carboxylic acids, using 1-(3-(dimethylamino)-propyl)3-ethylcarbodiimide HCl (EDC)

To a solution of "Starting Material A," the carboxylic acid (1 eq.), in N,N-dimethylformamide was added "Starting Material B," the primary amine (1 eq.), along with EDC (1 eq.), 1-hydroxy-7-azabenzotriazole (1 eq.), 4-dimethylaminopyridine (10% mol. eq.), and triethylamine (1 eq.). The mixture was stirred overnight at ambient temperature and concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and 1N hydrochloric acid. The mixture was washed with 1N hydrochloric acid, saturated sodium bicarbonate, brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was chromatographed on silica gel.

30 d) for cyclization of 2-((1R,3S)-3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonylamino }cycloalkyl compounds using potassium *t*-butoxide (KOtBu)

-199-

A solution of "Starting Material" (1 eq.) and 1 eq. of 1.0 N (in THF) KOtBu in DMF was stirred for 3 hours. The reaction was diluted with ethyl acetate and washed 5 times with water; dried and concentrated *in vacuo* to a residue, which was chromatographed on silica gel.

5

e) for cyclization of 2-(3-{[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]-carbonylamino}cycloalkyl compounds using potassium bis(trimethylsilyl)amide (KHMDS)

10 A solution of "Starting Material" (1 eq.) dissolved in dry DMF was stirred and cooled to 0 °C in an ice bath. The reaction was then treated with KHMDS (1.5 eq., 0.5M in THF) and was allowed to warm to room temperature. After 30 minutes the reaction was diluted in EtOAc, transferred to a separatory funnel, and washed consecutively with 1N HCl, saturated sodium bicarbonate, brine and dried over sodium sulfate. The solvent was removed *in vacuo* and the residue was chromatographed on silica gel to afford the desired
15 compound.

f) for Arylation of a primary amine

To a stirred solution of "Starting Material A," the primary amine (1 eq.), in DMF was added K₂CO₃ (2 eq.) and "Starting Material B," the arylhalide (3 eq.). The reaction
20 mixture was stirred at 50°C for 8 hours. It was diluted with ethyl acetate, washed (brine), dried (Na₂SO₄), filtered and concentrated. Column chromatography on silica gel for purification was preformed.

g) Enantiomeric separation of the 1R,3S and 1S,3R isomer configuration of the
25 cycloalkyl Examples

The racemic material was separated into its enantiomers by chiral HPLC, using a Chiralpak AD column and 30% isopropyl alcohol-heptane as the eluent at a flow of 1.0 mL/min.

a) Table for condensation using collidine:

Ex. #	Product	Starting Material A	Starting Material B	Physical Data
1	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(phenylmethyl)carbonyl]amino]cyclohexane	prep 69	phenylacetic acid	ESIMS <i>m/e</i> 450 ³⁵ Cl (M ⁺ +1) and 452 ³⁷ Cl (M ⁺ +1)
2	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(3-methoxyphenyl)methyl]carbonyl]amino]cyclohexane	prep 69	3-methoxy-phenylacetic acid	ESIMS <i>m/e</i> 480 ³⁵ Cl (M ⁺ +1) and 482 ³⁷ Cl (M ⁺ +1)
3	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(4-fluorophenyl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(4-fluorophenyl)-acetic acid	ESIMS <i>m/e</i> 468 ³⁵ Cl (M ⁺ +1) and 470 ³⁷ Cl (M ⁺ +1)
4	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(4-methoxyphenyl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(4-methoxyphenyl)-acetic acid	ESIMS <i>m/e</i> 480 ³⁵ Cl (M ⁺ +1) and 482 ³⁷ Cl (M ⁺ +1)
5	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(2-methoxyphenyl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(2-methoxyphenyl)-acetic acid	ESIMS <i>m/e</i> 480 ³⁵ Cl (M ⁺ +1) and 482 ³⁷ Cl (M ⁺ +1)
6	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(2-fluorophenyl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(2-fluorophenyl)-acetic acid	ESIMS <i>m/e</i> 468 ³⁵ Cl (M ⁺ +1) and 470 ³⁷ Cl (M ⁺ +1)
7	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(3-fluorophenyl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(3-fluorophenyl)-acetic acid	ESIMS <i>m/e</i> 502 ³⁵ Cl (M ⁺ +1) and 504 ³⁷ Cl (M ⁺ +1)
8	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(pyrid-2-yl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(2-pyridyl)acetic acid	ESIMS <i>m/e</i> 451 ³⁵ Cl (M ⁺ +1) and 453 ³⁷ Cl (M ⁺ +1)
9	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[(pyrid-3-yl)methyl]carbonyl]amino]cyclohexane	prep 69	2-(3-pyridyl)acetic acid	ESIMS <i>m/e</i> 451 ³⁵ Cl (M ⁺ +1) and 453 ³⁷ Cl (M ⁺ +1)

-201-

10	(1 <i>S</i> ,3 <i>R</i>)-1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[[[(2 <i>S</i>)-2-[(1,1-dimethylethoxy)carbonyl]amino]-2-(phenyl)acetyl]amino]cyclohexane	prep 68	(2 <i>S</i>)-2-[(<i>t</i> -butoxy)-carbonylamino]-2-phenylacetic acid	ESIMS <i>m/e</i> 565 ³⁵ Cl (<i>M</i> ⁺ +1) and 567 ³⁷ Cl (<i>M</i> ⁺ +1)
11	(1 <i>R</i> ,3 <i>S</i>)-1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[[[(2 <i>R</i>)-2-[(1,1-dimethylethoxy)carbonyl]amino]-2-(phenyl)acetyl]amino]cyclohexane	prep 69	(2 <i>R</i>)-2-[(<i>t</i> -butoxy)-carbonylamino]-2-phenylacetic acid	ESIMS <i>m/e</i> 565 ³⁵ Cl (<i>M</i> ⁺ +1) and 567 ³⁷ Cl (<i>M</i> ⁺ +1)
12	(1 <i>R</i> ,3 <i>S</i>)-1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[[[(2 <i>S</i>)-2-[(1,1-dimethylethoxy)carbonyl]amino]-2-(phenyl)acetyl]amino]cyclohexane	prep 69	(2 <i>S</i>)-2-[(<i>t</i> -butoxy)-carbonylamino]-2-phenylacetic acid	ESIMS <i>m/e</i> 565 ³⁵ Cl (<i>M</i> ⁺ +1) and 567 ³⁷ Cl (<i>M</i> ⁺ +1)
13	(1 <i>S</i> ,3 <i>R</i>)-1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[[[(2 <i>R</i>)-2-[(1,1-dimethylethoxy)carbonyl]amino]-2-(phenyl)acetyl]amino]cyclohexane	prep 68	(2 <i>R</i>)-2-[(<i>t</i> -butoxy)-carbonylamino]-2-phenylacetic acid	ESIMS <i>m/e</i> 565 ³⁵ Cl (<i>M</i> ⁺ +1) and 567 ³⁷ Cl (<i>M</i> ⁺ +1)

b) Table for Acylation:

Ex. #	Product	Starting Material A	Starting Material B	Physical Data	Comments
14	N-[[[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl]cyclohexyl]carboxamide	prep 44	cyclohexylcarbonyl chloride	MS (ion spray) 456.2 (<i>M</i> +1)	
15	N-[[[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl](4-phenylphenyl)carboxamide	prep 44	biphenylcarbonyl chloride	MS (ion spray) 526.2 (<i>M</i> +)	
16	N-[[[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]acetamide	prep 44	acetic anhydride	MS (ion spray) 388 (<i>M</i> +1)	

17	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]methanesulfonamide	prep 44	methanesulfonyl chloride	MS (ion spray) 424.0 (M+1)	
18	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]methoxycarboxamide	prep 44	methyl chloroformate	MS (ion spray) 404.0 (M+1).	
19	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(methylaminothioxomethylaminomethyl)cyclohexyl]-9-chloro-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 44	methyl isothiocyanate	MS (ion spray) 419.0 (M+1)	
20	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-2-methylpropanamide	prep 44	isobutyryl chloride	MS (ion spray) 416.1 (M+1)	
21	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-3-methylthiopropylpropanamide	prep 44	3-methylthiopropionyl chloride	MS (ion spray) 448.1 (M+1)	
22	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]pentanamide	prep 44	valeryl chloride	MS (ion spray) 430.2 (M+1)	
23	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-2-phenylacetamide	prep 44	phenacetyl chloride	MS (ion spray) 463.9 (M+1)	
24	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-3-phenylpropanamide	prep 44	hydrocinnamoyl chloride	MS (ion spray) 478.1 (M+1)	
25	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -pyridin-3-ylacetamide	3-aminopyridine	prep 47	MS (ion spray) 451 (M+1)	prep. 47 was converted to the acid chloride using oxalyl chloride in DCM

26	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -pyridin-4-ylacetamide	4-aminopyridine	prep 47	MS (ion spray) 451.2 (M+1)	prep. 47 was converted to the acid chloride using oxalyl chloride in DCM
27	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -thiazol-2-ylacetamide	2-aminothiazole	prep 47	MS (ion spray) 457.1 (M+1)	prep. 47 was converted to the acid chloride using oxalyl chloride in DCM
28	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -(3-hydroxyphenyl)acetamide	3-aminophenol	prep 47	MS (ion spray) 466.2 (M+1)	prep. 47 was converted to the acid chloride using oxalyl chloride in DCM
29	<i>N</i> -[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]acetamide	prep 48	acetic anhydride	MS (ion spray) 388 (M+1)	
30	<i>N</i> -[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-2,2-dimethylpropionamide	prep 48	pivaloyl chloride	MS (ion spray) 430.1 (M+1)	
31	<i>N</i> -[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-cyclobutyl carboxamide	prep 48	cyclobutylcarbonyl chloride	MS (ion spray) 428 (M+1)	
32	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-isoxazol-5-yl carboxamide	prep 48	isoxazolecarbonyl chloride	MS (ion spray) 441.1 (M+1)	
33	5-((1 <i>S</i> ,3 <i>R</i>)-3-[(Methylsulfonyl)amino]methyl)-9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 48	methanesulfonyl chloride	MS (ion spray) 424.1 (M+1)	

34	5-((1 <i>S</i> ,3 <i>R</i>)-3-[(Phenylsulfonyl)amino]methyl)-9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 48	benzenesulfonyl chloride	MS (ion spray) 486.4 (M+1)	
35	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]phenoxycarboxamide	prep 48	phenyl chloroformate	MS (ion spray) 466.2 (M+1)	
36	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl](phenyl-amino)carboxamide	prep 48	phenylisocyanate	MS (ion spray) 464.9 (M+)	
37	5-[(1 <i>S</i> ,3 <i>R</i>)-3-(Phenylaminothiomethylaminomethyl)-cyclohexyl]-9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 48	phenylthiocyanate	MS (ion spray) 481.02 (M+1)	
38	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]morpholin-4-yl carboxamide	prep 48	morpholine carbonyl chloride	MS (ion spray) 459.2 (M+1)	
39	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]cyclohexyl carboxamide	prep 48	cyclohexyl-carbonyl chloride	MS (ion spray) 456.3 (M+1)	
40	[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]pyridin-2-yl carboxamide	prep 48	picolinoyl chloride HCl	MS (ion spray) 450.9 (M+1)	
41	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]nicotinamide	prep 48	nicotinoyl chloride HCl	MS (ion spray) 451.1 (M+1)	
42	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]isonicotinamide	prep 48	isonicotinoyl chloride HCl	MS (ion spray) 451.1 (M+1)	
43	N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]benzamide	prep 54	benzoyl chloride	MS(FIA) m/z = 450	prep. 54 was converted to the primary amine using TFA in DCM

44	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3,4,5-trimethoxybenzamide	prep 54	3,4,5-Tri-methoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 540.4	prep. 54 was converted to the primary amine using TFA in DCM
45	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-4-methoxy-benzamide	prep 54	4-Methoxy-benzoyl chloride	MS(FIA) <i>m/z</i> = 480.1	prep. 54 was converted to the primary amine using TFA in DCM
46	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3-methoxy-benzamide	prep 54	3-Methoxy-benzoyl chloride	MS(FIA) <i>m/z</i> = 480.2	prep. 54 was converted to the primary amine using TFA in DCM
47	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-4-nitrobenzamide	prep 54	4-Nitrobenzoyl chloride	MS(FIA) <i>m/z</i> = 493.2	prep. 54 was converted to the primary amine using TFA in DCM
48	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3-nitrobenzamide	prep 54	3-Nitrobenzoyl chloride	MS(FIA) <i>m/z</i> = 495.1	prep. 54 was converted to the primary amine using TFA in DCM

49	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-4-carbomethoxybenzamide	prep 54	methyl 4-(chlorocarbonyl)benzoate	MS(FIA) <i>m/z</i> = 508.2	prep 54 was converted to the primary amine using TFA in DCM
50	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3-carbomethoxybenzamide	prep 54	methyl 3-(chlorocarbonyl)benzoate	MS(FIA) <i>m/z</i> = 508.2	prep 54 was converted to the primary amine using TFA in DCM
51	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3-carboxybenzamide	Ex 49	--	MS(FIA) <i>m/z</i> = 492.2	prepared by basic hydrolysis
52	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]-3-carboxybenzamide	Ex 50	--	MS(FIA) <i>m/z</i> = 492.2	prepared by basic hydrolysis
53	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]benzamide	prep 54	Benzoyl chloride	MS(FIA) <i>m/z</i> = 450	prep. 54 was converted to the primary amine using TFA in DCM
54	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]-quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-methoxybenzamide	prep 54	2-Methoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 480.1	prep. 54 was converted to the primary amine using TFA in DCM

55	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-methoxybenzamide	prep 54	3-Methoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 480.2	prep. 54 was converted to the primary amine using TFA in DCM
56	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-methoxybenzamide	prep 54	4-Methoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 480.1	prep. 54 was converted to the primary amine using TFA in DCM
57	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-ethoxybenzamide	prep 54	3-Ethoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 494.3	prep. 54 was converted to the primary amine using TFA in DCM
58	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-trifluoromethoxybenzamide	prep 54	4-Trifluoromethoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 534.2	prep. 54 was converted to the primary amine using TFA in DCM
59	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-ethoxybenzamide	prep 54	4-Ethoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 494.3	prep. 54 was converted to the primary amine using TFA in DCM

60	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3,4-dimethoxy-benzamide	prep 54	3,4-Dimethoxybenzoyl chloride	MS(FIA) <i>m/z</i> = 570.3	prep. 54 was converted to the primary amine using TFA in DCM
61	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3,4-methylenedioxybenzamide	prep 54	3,4-methylenedioxybenzoyl chloride	MS(FIA) <i>m/z</i> = 494.2	prep. 54 was converted to the primary amine using TFA in DCM
62	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-fluorobenzamide	prep 54	2-Fluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 468.2	prep. 54 was converted to the primary amine using TFA in DCM
63	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-fluorobenzamide	prep 54	3-Fluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 468.2	prep. 54 was converted to the primary amine using TFA in DCM
64	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-fluorobenzamide	prep 54	4-Fluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 468.2	prep. 54 was converted to the primary amine using TFA in DCM

65	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2,3-difluorobenzamide	prep 54	2,3-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.4	prep. 54 was converted to the primary amine using TFA in DCM
66	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2,4-difluorobenzamide	prep 54	2,4-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.5	prep. 54 was converted to the primary amine using TFA in DCM
67	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2,5-difluorobenzamide	prep 54	2,5-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.4	prep. 54 was converted to the primary amine using TFA in DCM
68	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2,6-difluorobenzamide	prep 54	2,6-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.6	prep. 54 was converted to the primary amine using TFA in DCM
69	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3,4-difluorobenzamide	prep 54	3,4-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.4	prep. 54 was converted to the primary amine using TFA in DCM

70	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3,5-difluorobenzamide	prep 54	3,5-Difluorobenzoyl chloride	MS(FIA) <i>m/z</i> = 486.4	prep. 54 was converted to the primary amine using TFA in DCM
71	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3,4,5-trifluorobenzamide	prep 54	3,4,5-Trifluorobenzoyl chloride	MS(ES+) <i>m/z</i> = 504.1	prep. 54 was converted to the primary amine using TFA in DCM
72	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-chlorobenzamide	prep 54	2-Chlorobenzoyl chloride	MS(FIA) <i>m/z</i> = 484.4	prep. 54 was converted to the primary amine using TFA in DCM
73	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-chlorobenzamide	prep 54	3-Chlorobenzoyl chloride	MS(FIA) <i>m/z</i> = 484.4	prep. 54 was converted to the primary amine using TFA in DCM
74	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-chlorobenzamide	prep 54	4-Chlorobenzoyl chloride	MS(FIA) <i>m/z</i> = 484.4	prep. 54 was converted to the primary amine using TFA in DCM

75	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-bromobenzamide	prep 54	3-Bromobenzoyl chloride	MS(FIA) <i>m/z</i> = 530.0	prep. 54 was converted to the primary amine using TFA in DCM
76	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-bromobenzamide	prep 54	4-Bromobenzoyl chloride	MS(FIA) <i>m/z</i> = 530.0	prep. 54 was converted to the primary amine using TFA in DCM
77	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-iodobenzamide	prep 54	4-Iodobenzoyl chloride	MS(FIA) <i>m/z</i> = 576.1	prep. 54 was converted to the primary amine using TFA in DCM
78	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-methylbenzamide	prep 54	2-Methylbenzoyl chloride	MS(FIA) <i>m/z</i> = 464.2	prep. 54 was converted to the primary amine using TFA in DCM
79	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-methylbenzamide	prep 54	3-Methylbenzoyl chloride	MS(FIA) <i>m/z</i> = 464.2	prep. 54 was converted to the primary amine using TFA in DCM

80	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-methylbenzamide	prep 54	4-Methylbenzoyl chloride	MS(FIA) <i>m/z</i> = 464.2	prep. 54 was converted to the primary amine using TFA in DCM
81	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-trifluoromethylbenzamide	prep 54	3-Trifluoromethylbenzoyl chloride	MS(FIA) <i>m/z</i> = 518.2	prep. 54 was converted to the primary amine using TFA in DCM
82	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-trifluoromethylbenzamide	prep 54	4-Trifluoromethylbenzoyl chloride	MS(FIA) <i>m/z</i> = 518.2	prep. 54 was converted to the primary amine using TFA in DCM
83	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-phenylbenzamide	prep 54	4-phenylbenzoyl chloride	MS(ES+) <i>m/z</i> = 526.0	prep. 54 was converted to the primary amine using TFA in DCM
84	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4- <i>t</i> -butylbenzamide	prep 54	4- <i>t</i> -Butylbenzoyl chloride	MS(FIA) <i>m/z</i> = 506.2	prep. 54 was converted to the primary amine using TFA in DCM

-213-

85	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-cyanobenzamide	prep 54	4-Cyanobenzoyl chloride	MS(ES+) m/z = 474.9	prep. 54 was converted to the primary amine using TFA in DCM
86	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2,6-dimethylbenzamide	prep 54	2,6-Dimethylbenzoyl chloride	MS(FIA) m/z = 478.2	prep. 54 was converted to the primary amine using TFA in DCM
87	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-dimethylaminobenzamide	prep 54	3-Dimethylamino benzoyl chloride	MS(FD) m/z = 492.2	prep. 54 was converted to the primary amine using TFA in DCM
88	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-4-dimethylaminobenzamide	prep 54	4-Dimethylaminobenzoyl chloride	MS(FIA) m/z = 493.2	prep. 54 was converted to the primary amine using TFA in DCM
89	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-1-naphthylcarboxamide	prep 54	naphthalene-1-carbonyl chloride	MS(FIA) m/z = 500.1	prep. 54 was converted to the primary amine using TFA in DCM

90	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-naphthylcarboxamide	prep 54	naphthalene-2-carbonyl chloride	MS(FIA) <i>m/z</i> = 500.1	prep. 54 was converted to the primary amine using TFA in DCM
91	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-furylcarboxamide	prep 54	2-Furoyl chloride	MS(FIA) <i>m/z</i> = 440.2	prep. 54 was converted to the primary amine using TFA in DCM
92	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-2-thiophenylcarboxamide	prep 54	2-Thiophene carbonyl chloride	MS(FIA) <i>m/z</i> = 456.3	prep. 54 was converted to the primary amine using TFA in DCM
93	<i>N</i> -[3 <i>S</i> -(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)- <i>R</i> -cyclohexylmethyl]-3-thiophenylcarboxamide	prep 54	3-Thiophene carbonyl chloride	MS(FIA) <i>m/z</i> = 456.3	prep. 54 was converted to the primary amine using TFA in DCM
94	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-3-acetylamino benzamide	Ex 567	acetyl chloride	MS(FIA) <i>m/z</i> = 507.2	
95	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo-[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-3- <i>bis</i> -(methanesulfonyl)-aminobenzamide	Ex 567	methanesulfonyl chloride	MS(FIA) <i>m/z</i> = 621.4	

96	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl-methyl]benzamide	prep 64	benzoyl chloride	MS (ion spray) 450 (M ⁺), 448 (M ⁻ - 1)	prep 64 was converted to the amine by TBAF in DCM at rt.
97	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl-methyl]-2-phenylacetamide	prep 64	phenylacetyl chloride	MS (ion spray) 464 (M ⁺), 462 (M ⁻ - 1)	prep 64 was converted to the amine by TBAF in DCM at rt.
98	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl-methyl]-3,4,5-trimethoxybenzamide	prep 64	3,4,5-trimethoxybenzoyl chloride	MS (ion spray) 540 (M ⁺), 548 (M ⁻ - 1)	prep 64 was converted to the amine by TBAF in DCM at rt.
99	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexyl]benzamide	prep 68	benzoyl chloride	MS (ion spray) 436 (M ⁺)	
100	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexyl]benzamide	prep 69	benzoyl chloride	MS (ion spray) 436 (M ⁺)	
101	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	prep 68	phenacetyl chloride	MS (ion spray) 450.1 (M+1)	
102	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	prep 69	phenacetyl chloride	MS (ion spray) 450.1 (M+1)	
103	Phenyl-carbamic acid 3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl ester	prep 85	phenyl isocyanate	MS(ES):(M+1) ⁺ 466.2 m/z	
104	N-[2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexyl]ethyl]-2,2-dimethyl-N-phenylpropionamide	prep 93	trimethylacetyl chloride	MS(ES) [M+H] ⁺ = 520.2367 m/z	

105	9-Chloro-5-{3-[3-chloro-1-oxo-butylamino)-methyl]-cyclohexyl}-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one	prep 48	4-chlorobutyl chloride	MS (FIA) (m/z) 450.1 (M+1)	
106	N-[3-(9-Iodo-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	Ex 594	benzoyl chloride	MS (FIA) (m/z) 542.3 [M+1]	Ex 594 was converted to the primary amine using TFA in DCM
107	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzenesulfonamide	prep 146	phenylsulfon chloride	MS (ion spray) 486 (M ⁺), 484 (M ⁻ - 1)	
108	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-cyclohexylmethyl]nicotinamide	prep 146	nicotinoyl chloride HCl	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)	
109	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-methoxybenzamide	prep 146	m-anisoyl chloride	MS (ion spray) 480 (M ⁺), 478 (M ⁻ - 1)	
110	Pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]amide	prep 146	picolinoyl chloride HCl	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)	
111	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-4-fluorobenzamide	prep 146	4-fluorobenzoyl chloride	MS (ion spray) 468 (M ⁺), 466 (M ⁻ - 1)	
112	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-fluorobenzamide	prep 146	3-fluorobenzoyl chloride	MS (ion spray) 468 (M ⁺), 466 (M ⁻ - 1)	

113	Thiophene-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]amide	prep 146	2-thiophene-carbonyl chloride	MS (ion spray) 456 (M ⁺), 454 (M ⁻ - 1)
114	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo-[4,3-c]quinolin-5-yl)cyclohexylmethyl]isonicotinamide	prep 146	isonicotinoyl chloride HCl	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)
115	N-[(1 <i>R</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]methyl}-2-furylcarboxamide	prep 146	2-furoyl chloride	MS (ion spray) 440 (M ⁺), 438 (M ⁻ - 1)
116	N-[(1 <i>R</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-methyl}-2-phenylacetamide	prep 146	phenylacetyl chloride	MS (ion spray) 440 (M ⁺), 438 (M ⁻ - 1)
117	<i>cis</i> -3-(benzoylamino)-1-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexane	prep 146	benzoylchloride	ESIMS <i>m/e</i> 436 35Cl (M ⁺ +1) & 438 37Cl (M ⁺ +1)
118	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo-[4,5-c]quinolin-5-yl)cyclohexylmethyl]benzamide	Ex 612	benzoyl chloride	MS (+ES)m/z 450.0 (M+H) ⁺
119	(1 <i>S</i> ,3 <i>R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexylmethyl]-4-fluorobenzamide	Ex 613	4-fluorobenzoyl chloride	MS (ES+) m/z 468.0 (M+H) ⁺
120	(1 <i>R</i> ,3 <i>S</i>)-Thiophene-2-carboxylic acid[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)cyclohexylmethyl]-amide	Ex 613	2-Thiophene-carbonyl chloride	MS (ES+) m/z 456.0 (M+H) ⁺
121	(1 <i>S</i> ,3 <i>R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)cyclohexylmethyl]benzamide	Ex 614	benzoyl chloride	MS (ES+) m/z 450.0 (M+H) ⁺
122	(1 <i>S</i> ,3 <i>R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)cyclohexylmethyl]-4-fluorobenzamide	Ex 614	4-Fluorobenzoyl chloride	MS (ES+) m/z 468.0 (M+H) ⁺

123	(<i>1S,3R</i>)-Thiophene-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-amide	Ex 614	2-thiophene-carbonyl chloride	MS (ES+) <i>m/z</i> 455.9 (M+H) ⁺	
124	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]nicotinamide	Ex 614	nicotinoyl chloride	MS (ES+) <i>m/z</i> 451.0 (M+H) ⁺	
125	(<i>1R,3S</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]benzamide	Ex 615	benzoyl chloride	MS (ES+) <i>m/z</i> 500.1 (M+H) ⁺	
126	(<i>1R,3S</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]nicotinamide	Ex 615	nicotinoyl chloride	MS (ES+) <i>m/z</i> 451.0 (M+H) ⁺	
127	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]nicotinamide	Ex 616	nicotinoyl chloride	MS (ES+) <i>m/z</i> 451.0 (M+H) ⁺	
128	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]nicotinamide	Ex 617	nicotinoyl chloride	MS (ES+) <i>m/z</i> 451.0 (M+H) ⁺	
129	<i>N</i> -[(<i>1S,3R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]-3,4-difluorobenzamide	Ex 619	3,4-difluoro-benzoyl chloride	ESMS: 472 (<i>m</i> +1) ⁺	
130	<i>N</i> -[(<i>1S,3R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]-3-methoxybenzamide	Ex 620	3-methoxy-benzoyl chloride	ESMS: 466 (M+1) ⁺	
131	Thiophene-2-carboxylic acid [(<i>1S,3R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]-amide	Ex 620	2-thiophene-carbonyl chloride	ESMS: 442 (M+1) ⁺	
132	(<i>1S,3R</i>)-3-(2-Chloro-6-fluorophenyl)-5-methylisoxazole-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]amide	Ex 620	2-Chloro-6-fluorobenzoyl chloride	ESMS: 569 (M+1) ⁺	
133	Pyrazine-2-carboxylic acid [(<i>1S,3R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]-amide	Ex 620	2-pyrazine-carbonyl chloride	ESMS: 438 (M+1) ⁺	

134	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]carbamoylphenylmethyl-carbamic acid <i>t</i> -butyl ester	Ex 620	(2 <i>S</i>)-2-[(<i>t</i> -butoxy)carbonylamino]-2-phenylacetic acid	ESMS: 565 (M+1)+	
135	2-Amino-N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]-2-phenylacetamide	Ex 620	(2 <i>S</i>)-2-amino-2-phenylacetic acid	ESMS: 465 (M+1)+	
136	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)cyclopentyl]benzamide	Ex 620	benzoyl chloride	ESMS: 436 (M+1)+	
137	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-methyl]cyclopentyl[(3,4,5-trimethoxyphenyl)carboxamide]	Ex 620	3,4,5-trimethoxybenzoyl chloride	ESMS: 526 (M+1)+	
138	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)methyl]cyclopentyl]amide	Ex 621	2-chloro-5-fluoro phenyl isoxazoyl chloride	ESMS: 570 (M+1)+	
139	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)methyl]cyclopentyl[(3,4,5-trimethoxyphenyl)carboxamide]	Ex 621	3,4,5-trimethoxybenzoyl chloride	ESMS: 526 (M+1)+	
140	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)cyclopentyl]-4-fluorobenzamide	Ex 621	4-fluorobenzoyl chloride	ESMS: 454 (M+1)+	
141	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)-cyclopentyl]-3,4-difluorobenzamide	Ex 621	3,4-difluorobenzoyl chloride	ESMS: 472 (M+1)+	

142	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)cyclopentyl]nicotinamide	Ex 621	pyridine-3-carbonyl chloride	ESMS: 437 (M+1)+	
143	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)cyclopentyl]-3-methoxybenzamide	Ex 621	3-methoxybenzoyl chloride	ESMS: 466 (M+1)+	
144	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)-cyclopentyl]phenylmethyl} carbamic acid <i>t</i> -butyl ester	Ex 621	L-N-trifluoroacetyl phenyl glycine	ESMS: 561 (M+1)+	
145	1-Acetyl-2,3-dihydro-1 <i>H</i> -indole-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]amide		acetic anhydride	MS (ion spray) 519 (M ⁺).	
146	N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexyl]- <i>C</i> -phenylmethane sulfonamide	Ex 624	α -toluenesulfonyl chloride	MS (ion spray) 486 (M ⁺).	
147	(<i>cis</i>) {N-[(<i>cis</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-carbonyl} phenylmethyl pyridine-3-carboxylate	Ex 470	nicotinoyl HCl	MS (ion spray) 571 (M ⁺).	
148	(<i>trans</i>) {N-[(<i>cis</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-carbonyl} phenylmethyl pyridine-3-carboxylate	Ex 471	nicotinoyl HCl	MS (ion spray) 571 (M ⁺).	
149	N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-6-fluoro-nicotinamide	Ex 632	6-Fluoro-nicotinic acid	MS m/z (ES+) 468.8 (M+H) ⁺	6-Fluoro-nicotinic acid was converted to the acid chloride by oxalyl chloride in DCM.

150	6-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)cyclohexylmethyl]-nicotinamide	Ex 632	6-Chloro-nicotinic acid	MS m/z (ES+) 468.8 (M+H) ⁺	6-Chloronicotinic acid was converted to the acid chloride by oxalyl chloride in DCM.
151	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)cyclohexyl]-2-methanesulfonylamino-2-phenylacetamide	Ex 633	methanesulfonyl chloride	MS(ES): (M+1) ⁺ 543.0, 545.0 m/z	
152	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)-cyclohexylmethyl]-5-diacetylamino-nicotinamide	Ex 251	acetic anhydride	MS (ion spray) 550.2 (M+).	
153	6-Acetylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-nicotinamide	Ex 245	acetic anhydride	MS (ion spray) 508.2 (M+)	
154	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)cyclohexylmethyl]-3-phenylpropionamide	Ex 632	hydrocinnyl chloride	MS (ion spray) 478.1 (M+).	
155	[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)cyclohexyl]carbamic acid phenyl ester	Ex 634	phenyl chloroformate	MS (ion spray) 451.96 (M+).	
156	1-Benzyl-3-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]urea	Ex 634	benzyl isocyanate	MS (ion spray) 465.0 (M+)	
157	1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-3-phenylthiourea	Ex 634	phenyl isothiocyanate	MS (ion spray) 467.0 (M+)	
158	1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-3-phenylurea	Ex 634	phenyl isocyanate	MS (ion spray) 451.96 (M+).	

159	<i>N</i> -{[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl}-2,2-dimethylpropionamide	prep 243	pivaloyl chloride	MS(ES) exact mass [M] = 548.2190 m/z	
160	<i>N</i> -[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-2-methanesulfonylamino-2-phenyl-acetamide	Ex 638	methanesulfonyl chloride	MS(ES) [M+H] ⁺ = 543.1 m/z	
161	Morpholine-4-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl}amide	Ex 638	morpholine-4-carbonyl chloride	MS(ES) [M] = 577.2092 m/z.	
162	<i>N</i> -{[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylcarbonyl]pyridin-3-ylmethyl}-2,2-dimethylpropionamide (isomer 1)	Ex 640 (isomer 1)	pivaloyl chloride	MS(ES) [M+H] ⁺ = 550.2253 m/z.	
163	<i>N</i> -{[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylcarbonyl]pyridin-3-ylmethyl}-2,2-dimethylpropionamide (isomer 2)	Ex 640 (isomer 2)	pivaloyl chloride	MS(ES) exact mass [M+H] ⁺ = 550.2247 m/z.	
164	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-6-fluoronicotinamide	Ex 632	6-Fluoronicotinic acid	MS m/z (ES+) 468.8 (M+H) ⁺	6-Fluoronicotinic acid was converted to the acid chloride by oxalyl chloride in DCM.
165	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-5-fluoronicotinamide	Ex 632	5-Fluoronicotinic acid	MS m/z (ES+) 468.8 (M+H) ⁺	5-Fluoronicotinic acid was converted to the acid chloride by oxalyl chloride in DCM.

166	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)methyl)cyclohexyl]benzamide	Ex 525	benzoyl chloride	MS(ES+) (m/z) 450.1 [M+1]	Ex 525 was converted to the amine by TFA in DCM.
167	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-ylethyl)cyclohexyl]-4-fluorobenzamide	Ex 525	4-fluorobenzoyl chloride	MS(ES+) (m/z) 468.1 [M+1]	Ex 525 was converted to the amine by TFA in DCM.
168	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)methyl)cyclohexyl]benzamide	Ex 525	3,4-difluorobenzoyl chloride	MS(ES+) (m/z) 486.1 [M+1].	Ex 525 was converted to the amine by TFA in DCM.
169	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)methyl)cyclohexyl]benzamide	Ex 533	benzoyl chloride	MS(ES+) (m/z) 450.1 [M+1].	Ex 533 was converted to the amine by TFA in DCM.
170	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)methyl)-cyclohexyl]-4-fluoro-benzamide	Ex 533	4-fluorobenzoyl chloride	MS(ES+) (m/z) 468.1 [M+1].	Ex 533 was converted to the amine by TFA in DCM.
171	<i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)methyl)cyclohexyl]-3,4-difluorobenzamide	Ex 533	3,4-difluorobenzoyl chloride	MS(ES+) (m/z) 486.1 [M+1].	Ex 533 was converted to the amine by TFA in DCM.
172	<i>N</i> -[3-(9-Cyano-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]-quinolin-5-yl)cyclohexylmethyl]nicotinamide	Ex 532	nicotinoyl chloride HCl	MS(ES+) (m/z) 442.3 [M+1].	

173	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)cyclohexyl]benzamide	Ex 540	benzoyl chloride	MS(ES+) (m/z) 450.3 [M+1].	Ex 540 was converted to the amine by TFA in DCM.
174	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-ylmethyl)cyclohexyl]-4-fluorobenzamide	Ex 540	4-fluorobenzoyl chloride	MS(ES+) (m/z) 468.0 [M+1].	Ex 540 was converted to the amine by TFA in DCM.
175	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)cyclohexyl]-3,4-difluorobenzamide	Ex 540	3,4-difluorobenzoyl chloride	MS(ES+) (m/z) 486.2 [M+1].	Ex 540 was converted to the amine by TFA in DCM.
176	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)cyclohexyl]bis(2-methanesulfonyl)amino-acetamide	prep 360	mesyl chloride	MS (ion spray) 545.1 (M+)	
177	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]-quinolin-5-yl)cyclohexyl]benzamide	prep 380	benzoyl chloride	MS (ES+) m/z 436.1 (M+H) ⁺ , (ES-) 434.1 (M-H) ⁻ .	
178	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3,4-difluoro-phenyl)-acetamide	3,4-difluoroaniline	Prep 47	MS(ES+) m/z = 485.8	Prep 47 was converted to the acid chloride by oxalyyl chloride in DCM.
179	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3,5-difluoro-phenyl)-acetamide	3,5-difluoroaniline	Prep 47	MS(ES+) m/z = 486.1	Prep 47 was converted to the acid chloride by oxalyyl chloride in DCM.

180	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(2-fluoro-phenyl)-acetamide	2-difluoroaniline	Prep 47	MS(ES+) m/z = 468.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
181	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-Benzyl-acetamide	benzylamine	Prep 47	MS(ES+) m/z = 464.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
182	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3,4,5-trimethoxy-benzyl)-acetamide	3,4,5-Trimethoxy-benzylamine	Prep 47	MS(ES+) m/z = 554.2	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
183	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(2-methoxy-phenyl)-acetamide	2-Methoxy-phenylamine	Prep 47	MS(ES+) m/z = 480.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
184	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(4-methoxy-phenyl)-acetamide	4-Methoxy-phenylamine	Prep 47	MS(ES+) m/z = 480.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
185	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(2-methyl-phenyl)-acetamide	2-Methyl-phenylamine	Prep 47	MS(ES+) m/z = 464.2	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.

186	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3-methyl-phenyl)-acetamide	3-Methyl-phenylamine	Prep 47	MS(ES+) m/z = 464.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
187	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(4-methyl-phenyl)-acetamide	4-Methyl-phenylamine	Prep 47	MS(ES+) m/z = 464.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
188	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(2,6-dimethyl-phenyl)-acetamide	2,6-Methyl-phenylamine	Prep 47	MS(ES+) m/z = 478.2	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
189	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(4-carbomethoxy-phenyl)-acetamide	4-Amino-benzoic acid methyl ester	Prep 47	MS(ES+) m/z = 508.2	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
190	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(6-methoxy-quinoline)-acetamide	6-Methoxy-quinolin-8-ylamine	Prep 47	MS(ES+) m/z = 531.0	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
191	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-phenyl-N-methyl-acetamide	Methyl-phenyl-amine	Prep 47	MS(ES+) m/z = 464.0	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.

-227-

192	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-benzyl-N-methyl-acetamide	Methyl-benzyl-amine	Prep 47	MS(ES+) m/z = 478.1	Prep 47 was converted to the acid chloride by oxalyl chloride in DCM.
193	(3-{[5-(2-Chloro-6-fluoro-phenyl)-3-methyl-3H-imidazole-4-carbonyl]-amino}-cyclohexylmethyl)-carbamic acid benzyl ester	Ex 615	prep 405	MS (ES+) (m/z) 499.2 [M+1]	Prep 405 was converted to the acid chloride by oxalyl chloride in DCM.
194	N-[3-(9-Chloro-3-methyl-4-oxo-3,4-dihydro-imidazo[4,5-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	Ex 193	benzoyl chloride	MS (ES+) (m/z) 449.2 [M+1]	Example 193 was converted to the primary amine by TMSI in DCM.
195	1R, 3S-N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]-benzamide	Ex 621	benzoyl chloride	ESMS: 436 (M+1) ⁺	
196	1R, 3S-N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]-2-phenyl-acetamide	Ex 621	phenylacetamide	ESMS: 450 (M+1) ⁺	
197	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-benzamide	prep 399	benzoyl chloride	MS (ES+) m/z 436.1 (M+H) ⁺ , (ES-) 434.1 (M-H) ⁻	
198	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-N-methyl-nicotinamide	prep 390	nicotinoyl chloride HCl	MS (ion spray) 465 (M ⁺)	
199	R(-)Pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]-phenylmethyl ester	Ex 471	picolinoyl chloride HCl	MS (ion spray) 571 (M ⁺)	
200	R(-)Isonicotinic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenylmethyl ester	Ex 471	isonicotinoyl chloride HCl	MS (ion spray) 571 (M ⁺)	

201	S(+)-Pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl ester	Ex 470	picolinoyl chloride HCl	MS (ion spray) 571 (M ⁺)	
202	S(+)-Isonicotinic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl ester	Ex 470	isonicotinoyl chloride HCl	MS (ion spray) 571 (M ⁺)	

c) EDC Coupling:

Ex. #	Product	Starting Material A	Starting Material B	Physical Data	Comments
203	9-Chloro-5-(3-{2-[4-(4-fluorobenzoyl)piperidin-1-yl]-2-oxoethyl}cyclohexyl)-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one	prep 39	4-(4-fluorobenzoyl)piperidine HCl	MS (ion spray) 564.1 (M ⁺)	
204	9-Chloro-3-methyl-5-[3-(2-morpholin-4-yl-2-oxoethyl)-cyclohexyl]-5H-isoxazolo[4,3-c]quinolin-4-one	prep 39	Morpholine	MS (ion spray) 444.2 (M+1)	
205	9-Chloro-3-methyl-5-{3-[2-oxo-2-(4-phenylpiperazin-1-yl)oxoethyl]cyclohexyl}-5H-isoxazolo[4,3-c]quinolin-4-one	prep 39	1-phenylpiperazine	MS (ion spray) 519.2 (M ⁺)	
206	5-{3-[2-(4-Acetylpiperazin-1-yl)-2-oxoethyl]cyclohexyl}-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one	prep 39	1-acetylpiperazine	MS (ion spray) 485.3 (M+1)	
207	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cyclopropylacetamide	prep 39	cyclopropylamine	MS (ion spray) 414.2 (M+1)	
208	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cyclobutylacetamide	prep 39	cyclobutylamine	MS (ion spray) 428.2 (M+1)	
209	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cyclopentylacetamide	prep 39	Cyclopentylamine	MS (ion spray) 442.2 (M+1)	

210	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cyclohexylacetamide	prep 39	Cyclohexylamine	MS (ion spray) 456.3 (M+1)	
211	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cycloheptylacetamide	prep 39	Cycloheptyl-amine	MS (ion spray) 470.2 (M+1)	
212	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-cyclohexyl-N-methylacetamide	prep 39	N-methyl,N-cyclohexyl-amine	MS (ion spray) 470.02 (M+1)	
213	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-indan-2-yl-acetamide	prep 39	2-aminoindane	MS (ion spray) 490.2 (M+1)	
214	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-naphthalen-1-yl-acetamide	prep 39	1-naphthylamine	MS (ion spray) 500.2 (M+)	
215	9-Chloro-3-methyl-5-[3-(2-oxo-2-piperidin-1-yl-ethyl)-cyclohexyl]-5H-isoxazolo[4,3-c]quinolin-4-one	prep 39	Piperidine	MS (ion spray) 442.2 (M+1)	
216	N,N-Dibenzyl-2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide	prep 39	Dibenzylamine	MS (ion spray) 554.3 (M+1)	
217	N-Benzyl-2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-(3,4,5-trimethoxybenzyl)acetamide	prep 39	N-benzyl,N-(3,4,5-trimethoxybenzyl)amine	MS (ion spray) 644.3 (M+1)	
218	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-pyridin-2-ylacetamide	prep 39	2-aminopyridine	MS (ion spray) 451.1 (M+1)	
219	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-pyridin-3-ylacetamide	prep 39	3-aminopyridine	MS (ion spray) 451.1 (M+1)	
220	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-pyridin-4-ylacetamide	prep 39	4-aminopyridine	MS (ion spray) 451.1 (M+1)	

221	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-3-cyclohexylpropanamide	cyclohexane-propionic acid	prep 44	MS (ion spray) 484.2 (M+1)	
222	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]pyrazin-2-yl carboxamide	2-pyrazine-carboxylic acid	prep 44	MS (ion spray) 452.1 (M+1)	
223	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-2-thiophen-2-ylacetamide	2-thiophene-acetic acid	prep 44	MS (ion spray) 470.1 (M+1)	
224	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-2-(1-methyl-1 <i>H</i> -imidazol-4-yl)-acetamide	1-methyl-4-imidazoleacetic acid HCl	prep 44	MS (ion spray) 468.1 (M+1)	
225	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-4-phenoxy-benzamide	4-phenoxy-benzoic acid	prep 44	MS (ion spray) 541.8 (M+)	
226	4-Benzoyl-N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide	4-benzoyl-benzoic acid	prep 44	MS (ion spray) 554.2 (M+)	
227	6-Fluoropyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]amide	prep 1	Ex 632	MS (ion spray) 469 (M ⁺)	
228	N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-2-(4-fluorophenyl)-2-hydroxyacetamide	D-(-)-mandelic acid	Ex 634	MS (ion spray) 484 (M ⁺)	flash chrom. (silica gel; 0-0.5% MeOH/chloroform)g ave both isomers

229	<i>Cis</i> -N-[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl}(5-fluoro(2-pyridyl))-carboxamide	4-fluoro-picolinic acid	Ex 632	MS (ion spray) 469 (M ⁺).	
230	<i>Cis</i> -N-[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl}(6-methoxy(2-pyridyl))carboxamide	6-methoxy-pyridine-2-carboxylic acid	Ex 632	MS (ion spray) 481 (M ⁺).	
231	<i>Cis</i> -N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-5-methoxynicotinamide	prep 3	Ex 632	MS (ion spray) 481 (M ⁺).	
232	<i>cis</i> -N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-2-(2,3-difluorophenyl)-2-hydroxyacetamide	(<i>d/l</i>)-2,3-difluoromandelic acid	Ex 634	MS (ion spray) 502 (M ⁺).	flash chrom. (silica gel: 0-0.5% MeOH/chloroform)gave both isomers
234	<i>cis</i> -N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-2-(3,4-difluorophenyl)-2-hydroxyacetamide	(<i>d/l</i>)-3,4-difluoromandelic acid	Ex 634	MS (ion spray) 502 (M ⁺).	flash chrom. (silica gel: 0-0.5% MeOH/chloroform)gave both isomers
235	<i>cis</i> {[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylcarbamoyl}phenylmethyl} carbamic acid <i>t</i> -butyl ester	N-(<i>t</i> -butoxycarbonyl)-D-phenyl-glycine	Ex 634	MS(ES): (M+Na) ⁺ 587.1 m/z, (M-BOC) ⁺ 465.1 m/z	silica gel column eluted with 30% ethylacetate/hexanes: isomer 1: R _t = 33.67 min. isomer 2: R _t = 23.81 min. Chiracel AD.

236	N-{{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl}}-2,2-dimethyl-propionamide	trimethyl-acetic acid	Ex 633	MS(ES): (M+1) ⁺ 549.1, 551.1 m/z.	
237	N-{{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl}}nicotinamide	nicotinic acid	Ex 633	MS(ES): (M+1) ⁺ 570.0, 572.0 m/z	
238	Pyridine-2-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]-phenylmethyl}amide	picolinic acid	Ex 633	MS(ES): (M+1) ⁺ 570.2, 572.3 m/z	
239	N-{{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl}}isonicotinamide	isonicotinic acid	Ex 633	MS(ES): (M+1) ⁺ 570.3, 572.3 m/z	
240	2- <i>t</i> -Butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	prep 11	Ex 634	MS(ES): (M+1) ⁺ 521.2, 522.2, 523.2 m/z	Isomer 1 38%
241	2- <i>t</i> -Butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	prep 11	Ex 634	MS(ES): (M+1) ⁺ 521.2, 522.2, 523.2 m/z	Isomer 2 30%
242	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(2,2-dimethylpropylamino)-2-phenylacetamide	prep 13	Ex 634	MS(ES): (M+1) ⁺ 535.3, 536.3, 537.3 m/z.	Isomer 1 29%
243	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(2,2-dimethylpropylamino)-2-phenylacetamide	prep 13	Ex 634	MS(ES): (M+1) ⁺ 535.3, 536.3, 537.3 m/z.	Isomer 2 44%
244	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-1-oxynicotinamide	nicotinic acid-N-oxide	Ex 619	MS (ion spray) 467.2 (M+).	

245	6-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	4-amino-nicotinic acid	Ex 619	MS (ion spray) 466.3 (M+).
246	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-2-pyridin-3-yl-acetamide	3-pyridylacetic acid HCl	Ex 619	MS (ion spray) 465.0 (M+).
247	Quinoline-3-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]amide	3-quinoline-carboxylic acid	Ex 619	MS (ion spray) 501 (M+).
248	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-pyrazol-1-yl-nicotinamide	6-(1H-pyrazol-1-yl)-nicotinic acid	Ex 619	MS (ion spray) 517.2 (M+).
249	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-pyridin-3-yl-propionamide	3-pyridylpropionic acid	Ex 619	MS (ion spray) 479.1 (M+).
250	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-trifluoromethyl-nicotinamide	6-trifluoromethylnicotinic acid	Ex 619	MS (ion spray) 501 (M+).
251	5-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	5-aminonicotinic acid	Ex 619	MS (ion spray) 466.2 (M+).
252	Pyrimidine-5-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-amide	prep 15	Ex 619	MS (ion spray) 452.2 (M+).
253	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-4-methylaminobenzamide	4-(methyl-amino)benzoic acid	Ex 619	MS (ion spray) 479.0 (M+).
254	6-Methyl-pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]amide	6-methylpicolinic acid	Ex 619	MS (ion spray) 465.0 (M+).

255	4-Chloro-pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]amide	4-chloro-picolinic acid (TCI-US)	Ex 619	MS (ion spray) 486.9 (M+).	
256	2,6-Dichloro-N-[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-isonicotinamide	2,6-dichloro-isonicotinic acid	Ex 619	MS (ion spray) 518.9 (M+).	
257	N-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-2-fluoroisonicotinamide	2-fluoro-isonicotinic acid	Ex 619	MS (ion spray) 469.0 (M+).	
258	Furo[3,2- <i>b</i>]pyridine-5-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-amide	Furo[3,2- <i>b</i>]pyridine-5-carboxylic acid	Ex 619	MS (ion spray) 491.2 (M+).	
259	6-Chloro-pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]-amide	6-chloro-2-pyridinecarboxylic acid (SALOR)	Ex 619	MS (ion spray) 485.2 (M+).	
260	6-Methoxy-pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]amide	prep 16	Ex 619	MS (ion spray) 481.1 (M+).	
261	4-Methoxypyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexylmethyl]amide	4-methoxy-picolinic acid HCl	Ex 619	MS (ion spray) 481.2 (M+).	
262	5,6-Dichloro-N-[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	5,6-dichloro-nicotinic acid	Ex 619	MS (ion spray) 519.0 (M+).	

263	2-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-6-methylisonicotinamide	2-chloro-6-methylpyridine-4-carboxylic acid	Ex 619	MS (ion spray) 498.8, 500.87 (M+).	
264	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-5-methyl-nicotinamide	5-methyl-nicotinic acid	Ex 619	MS (ion spray) 465.1 (M+).	
265	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-methoxy-nicotinamide	prep 19	Ex 619	MS (ion spray) 481.0 (M+).	
266	5-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	5-chloro-nicotinic acid	Ex 619	MS (ion spray) 485.0 (M+).	
267	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-2-phenylacetamide	phenylacetic acid	Ex 619	MS (ion spray) 501 (M+).	
268	4-Fluoronaphthalene-1-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-amide	4-fluoro-1-naphthoic acid	Ex 619	MS (ion spray) 518.1 (M+).	
269	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-hydroxy-2-methylbenzamide	2-methyl-3-hydroxybenzoic acid	Ex 619	MS (ion spray) 479.9 (M+).	
270	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methoxy-2-phenylacetamide	(S)-(+)- α -methoxyphenyl acetic acid	Ex 634	MS (ion spray) 480.1(M+).	
271	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-methoxy-2-phenylacetamide	(R)-(-)- α -methoxyphenyl acetic acid	Ex 634	MS (ion spray) 480.1(M+).	

272	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-3,3,3-trifluoro-2-methoxy-2-phenylpropionamide	(S)-(-)- α -methoxy- α -(trifluoromethyl)phenylacetic acid	Ex 634	MS (ion spray) 548.1 (M+).	
273	Acetic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl ester	(R)-(-)-O-acetylmandelic acid	Ex 634	MS (ion spray) 508.1 (M+).	
274	Acetic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl ester	(S)-(+)-O-acetylmandelic acid	Ex 634	MS (ion spray) 508.1 (M+).	
275	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenoxy-2-phenylacetamide	phenoxyphenylacetic acid	Ex 634	MS (ion spray) 542.1 (M+).	
276	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl} carbamic acid benzyl ester	CBZ-D(-)-phenylglycine	Ex 634	MS (ion spray) 599.2 (M+).	
277	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]phenylmethyl} carbamic acid benzyl ester	Z-Phg-OH	Ex 634	MS (ion spray) 599.2 (M+)	
278	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-morpholin-4-yl-2-phenylacetamide	Morpholin-4-yl-phenylacetic acid	Ex 634	MS (ion spray) 535.2 (M+)	
279	2-(Acetyl-methyl-amino)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	(acetyl-methyl-amino)phenylacetic acid	Ex 634	MS (ion spray) 521.2 (M+)	

280	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenyl-2-phenylaminoacetamide	phenylphenyl-aminoacetic acid	Ex 634	MS (ion spray) 541.1 (M+)	
281	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-phenyl-acetamide	dimethylamino-phenylacetic acid	Ex 634	MS (ion spray) 493.0 (M+)	diastereomers were separated
282	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenyl-2-thiomorpholin-4-ylacetamide	phenylthiomorpholin-4-ylacetic acid	Ex 634	MS (ion spray) 550.9 (M+)	
283	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(4-methylpiperazin-1-yl)-2-phenylacetamide	(4-methylpiperazin-1-yl)phenylacetic acid	Ex 634	MS (ion spray) 548.2 (M+)	
284	2-(4-Acetylpiperazin-1-yl)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide	(4-acetyl-piperazin-1-yl)-phenylacetic acid	Ex 634	MS (ion spray) 576.2 (M+)	
285	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(indan-2-ylamino)-2-phenylacetamide	(indan-2-ylamino)-phenylacetic acid	Ex 634	MS (ion spray) 581.0 (M+)	diastereomers were separated
286	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-2-phenylpropionamide	(S)-(+)-2-hydroxy-2-phenylpropionic acid	Ex 634	MS (ion spray) 480.13 (M+)	

287	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-propionamide	(R)-(-)-2-phenylpropionic acid	Ex 634	MS (ion spray) 480.1 (M+)	
288	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-2-(3-methoxyphenyl)acetamide	m-methoxy-mandelic acid	Ex 634	MS (ion spray) 496.1 (M+)	diastereomers were separated
289	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-2-(4-methoxyphenyl)acetamide	p-methoxy-mandelic acid	Ex 634	MS (ion spray) 496.1 (M+)	diastereomers were separated
290	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-2-(4-trifluoromethylphenyl)acetamide	p-trifluoromethylmandelic acid	Ex 634	MS (ion spray) 534.0 (M+)	diastereomers were separated
291	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-3-phenylpropionamide	D-3-phenyl-lactic acid	Ex 634	MS (ion spray) 480.2 (M+)	
292	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-3-phenyl-propionamide	L-3-phenyl-lactic acid	Ex 634	MS (ion spray) 480.0 (M+)	
293	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-hydroxy-2-phenyl-propionamide	DL-tropic acid	Ex 634	MS (ion spray) 480.1 (M+).	diastereomers were separated
294	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(3-chloro-phenyl)-2-hydroxyacetamide	(R)-(-)-3-chloro-mandelic acid	Ex 634	MS (ion spray) 500.1 (M+)	

295	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(3-chloro-phenyl)-2-hydroxyacetamide	racemic 3-chloromandelic acid	Ex 634	MS (ion spray) 500.1 (M+)	diastereomers were separated by radial HPLC
296	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-m-tolylacetamide	hydroxy-m-tolylacetic acid	Ex 634	MS (ion spray) 480.0 (M+)	diastereomers were separated
297	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2-fluoro-phenyl)-2-hydroxyacetamide	(2-Fluoro-phenyl)-hydroxy-acetic acid	Ex 634	MS (ion spray) 480.0 (M+)	diastereomers were separated
298	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2,2-diphenylacetamide	benzyllic acid	Ex 634	MS (ion spray) 542 (M+)	
299	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylisobutyramide	dimethylphenyl acetic acid	Ex 634	MS (ion spray) 478.1 (M+)	
300	1-Phenylcyclopropanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide	1-phenyl-1-cyclopropyl-acetic acid	Ex 634	MS (ion spray) 476.1 (M+)	
301	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-fluoro-2-phenylacetamide	α -fluoro-phenylacetic acid	Ex 634	MS (ion spray) 468.1 (M+)	
302	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2,2-difluoro-2-phenylacetamide	difluorophenyl-acetic acid	Ex 634	MS (ion spray) 486.1 (M+)	
303	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-phenylsulfanyl-acetamide	phenylphenylthioacetic acid (Lancaster)	Ex 634	MS (ion spray) 558.0 (M+)	

304	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-propionamide	(R)-(-)-2-phenyl-propionic acid	Ex 634	(ion spray) 464.2 (M+)	
305	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylpropionamide	(S)-(+)-2-phenylpropionic acid	Ex 634	MS (ion spray) 464.2 (M+)	
306	1-Phenylcyclohexanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide	1-phenyl-1-cyclohexanecarboxylic acid (Acros)	Ex 634	MS (ion spray) 518.2 (M+)	
307	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-methylsulfanyl-2-phenylacetamide	α -S-methyl-phenylacetic acid	Ex 634	MS (ion spray) 495.9 (M+)	
308	Bicyclo[4.2.0]octa-1(6),2,4-triene-7-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide	1-benzocyclobutenecarboxylic acid	Ex 634	MS (ion spray) 461.9 (M+)	
309	{2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]carbonyl}indan-2-yl}carbamic acid <i>t</i> -butyl ester	N-BOC-2-aminoindane-2-carboxylic acid	Ex 634	MS (ion spray) 491 (M-BOC)	
310	3-Oxoindan-1-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide	3-oxo-1-indancarboxylic acid	Ex 634	MS (ion spray) 489.9 (M+)	
311	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid <i>t</i> -butyl ester	N-t-BOC-L-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Sigma)	Ex 634	MS (ion spray) 591.2 (M+)	

312	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid <i>t</i> -butyl ester	N-t-BOC-D-1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (BAChem)	Ex 634	MS (ion spray) 591.3 (M+)	
313	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-6-fluoronicotinamide	6-fluoro-nicotinic acid	Ex 634	MS (ion spray) 455.0 (M+)	
314	3-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]-azetidine-1-carboxylic acid <i>t</i> -butyl ester	BOC-azetidine-3-carboxylic acid	Ex 634	MS (ion spray) 415.08 (M+-BOC).	
315	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylcarbonyl]pyrrolidine-1-carboxylic acid <i>t</i> -butyl ester	N-t-BOC-D-proline	Ex 634	MS (ion spray) 529.2 (M+).	
316	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-3,3-dimethyl-butylamide	(<i>S</i>)-(+)-2-hydroxy-3,3-dimethylbutyric acid	Ex 634	MS (ion spray) 446.2 (M+)	
317	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-cyclohexyl-2-hydroxyacetamide	(<i>L</i>)-(+)-hexahydro-mandelic acid	Ex 634	MS (ion spray) 472.2 (M+)	
318	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-cyclohexyl-2-hydroxyacetamide	(<i>R</i>)-(-)-hexahydro-mandelic acid	Ex 634	MS (ion spray) 472.2 (M+)	
319	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3,3,3-trifluoro-2-hydroxy-propionamide	3,3,3-trifluoro-lactic acid	Ex 634	MS (ion spray) 480.0 (M+)	diastereomers were separated

320	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-methyl-propionamide	2-methyl-lactic acid	Ex 634	MS (ion spray) 417.9 (M+)	
321	[1-({[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-carbamoyl]-phenylmethyl)-carbamoyl]-1-methylethyl]-carbamic acid <i>t</i> -butyl ester	<i>N</i> - <i>t</i> -butyl- α -aminoisobutyric acid	Ex 638	MS(ES) [M+H] ⁺ = 650.2 m/z	
322	Pyridine-2-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenylmethyl}-amide	pyridine-2-carboxylic acid	Ex 638	MS(ES) [M+H] ⁺ = 570.2 m/z	
323	Pyridine-3-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenylmethyl}-amide	pyridine-3-carboxylic acid	Ex 638	MS(ES) [M+H] ⁺ = 570.2 m/z	
324	Pyridine-4-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenylmethyl}-amide	pyridine-4-carboxylic acid	Ex 638	MS(ES) [M+H] ⁺ = 570.2 m/z	
325	<i>N</i> -[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-[1,2,4]triazol-1-yl-isobutyramide	2-Methyl-2-[1,2,4]triazol-1-yl-propionic acid	Ex 634	MS(ES) exact mass calc'd: [M+H] ⁺ = 469.1755 m/z.	
326	{[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-pyridin-3-yl-methyl}-carbamic acid <i>t</i> -butyl ester	<i>t</i> -butoxycarbonylaminopyridin-3-ylacetic acid	Ex 634	MS(ES) calc'd: [M+H] ⁺ = 565.20 m/z; [M+Na] ⁺ = 588.20 m/z.	
327	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-acetamide	glycolic acid	Ex 634	MS (ion spray) 389.9 (M+) racemic cis	

328	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-3-methyl-butylamide	D-alpha-hydroxyisovaleric acid	Ex 634	MS (ion spray) 431.9 (M+) racemic cis	
329	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-hydroxy-3-phenyl-propionamide	(S)-3-hydroxy-3-phenylpropionic acid	Ex 634	MS (ion spray) 480.1 (M+) racemic cis	
330	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-hydroxy-3-phenyl-propionamide	(R)-3-hydroxy-3-phenylpropionic acid	Ex 634	MS (ion spray) 480.1 (M+) racemic cis	
331	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-3-methylbutylamide	(S)-(+)-2-hydroxy-3-methylbutyric acid	Ex 634	MS (ion spray) 431.9 (M+) racemic cis	
332	1-Hydroxy-cyclopropanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide	1-hydroxy-1-cyclopropanecarboxylic acid	Ex 634	MS (ion spray) 416.0 (M+) racemic cis	
333	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbonyl}-methyl}-methyl-carbamic acid <i>t</i> -butyl ester	N-t-BOC-sarcosine	Ex 634	MS (ion spray) 503.2 (M+) racemic cis	
334	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-acetamide	N,N-dimethylglycine	Ex 634	MS (ion spray) 416.9 (M+) racemic cis	
335	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbonyl}-1-methyl-ethyl}-carbamic acid <i>t</i> -butyl ester	N-t-BOC- α -aminoisobutyric acid	Ex 634	MS (ion spray) 516.9 (M+) racemic cis	

336	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-methyl}-carbamic acid <i>t</i> -butyl ester	N-t-butoxycarbonylglycine	Ex 634	MS (ion spray) 489.1 (M+) racemic cis	diastereomers were separated
337	2-Hydroxy-hexanoic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide	2-hydroxycaproic acid	Ex 634	MS (ion spray) 446.2 (M+) racemic cis	diastereomers were separated
338	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxybenzamide	salicylic acid	Ex 634	MS (ion spray) 452.0 (M+) racemic cis	
339	4-{{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl}-piperazine-1-carboxylic acid <i>t</i> -butyl ester	prep 285	Ex 634	MS (ion spray) 633.9 (M+) racemic cis	
340	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl}-methyl-carbamic acid <i>t</i> -butyl ester	BOC-N-Me-Phg-OH	Ex 634	MS (ion spray) 578.9 (M+) racemic cis	
341	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-[4-(2-hydroxy-ethyl)-piperazin-1-yl]-2-phenyl-acetamide	prep 287	Ex 634	MS (ion spray) 578.1 (M+) racemic cis	diastereomers were separated
342	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-(4-pyridin-2-yl-piperazin-1-yl)-acetamide	prep 288	Ex 634	MS (ion spray) 610.9 (M+) racemic cis	diastereomers were separated
343	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-piperidin-1-yl-acetamide	prep 290	Ex 634	MS (ion spray) 457.1 (M+) racemic cis	

344	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(4-methyl-piperazin-1-yl)-acetamide	prep 292	Ex 634	MS (ion spray) 472.0 (M+) racemic cis	
345	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-diethylamino-acetamide	N,N-diethyl-glycine sodium salt	Ex 634	MS (ion spray) 445.0 (M+)	
346	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(methyl-phenyl-amino)-acetamide	prep 293	Ex 634	MS (ion spray) 479.0 (M+) racemic cis	
347	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-(pyridin-3-yloxy)-acetamide	prep 294	Ex 634	MS (ion spray) 543.0 (M+)	
348	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-3-yloxy)-acetamide	prep 296	Ex 634	MS (ion spray) 467.1 (M+)	
349	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbonyl}-cyclohexyl} carbamic acid <i>t</i> -butyl ester	prep 297	Ex 634	MS (ion spray) 557.2 (M+)	
350	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-morpholin-4-yl-acetamide	prep 299	Ex 634	MS (ion spray) 459.0 (M+)	
351	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(4-hydroxy-piperidin-1-yl)-acetamide	prep 301	Ex 634	MS (ion spray) 473.1 (M+)	
352	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2-oxo-2H-pyridin-1-yl)-acetamide	prep 303	Ex 634	MS (ion spray) 467.1 (M+)	

353	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-4-yloxy)-acetamide	prep 304	Ex 634	MS (ion spray) 467.0 (M+)	
354	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-4-ylsulfanyl)-acetamide	(4-pyridylthio)-acetic acid	Ex 634	MS (ion spray) 483.1 (M+)	
355	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-cyclohexyl-methyl}-carbamic acid <i>t</i> -butyl ester	N- <i>t</i> -BOC-D- α -cyclohexyl-glycine	Ex 634	MS (ion spray) 571.2 (M+)	
356	{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-cyclohexyl-methyl}-carbamic acid <i>t</i> -butyl ester	N- <i>t</i> -BOC-L- α -cyclohexyl-glycine	Ex 634	MS (ion spray) 571.2 (M+)	
357	Thieno[3,2-b]pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide	thieno[3,2-b]-pyridine-2-carboxylic acid	Ex 634	MS (ion spray) 493.0 (M+)	
358	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2-chloro-pyridin-4-yloxy)-acetamide	thieno[3,2-b]-pyridine-2-carboxylic acid	Ex 634	MS (ion spray) 501.0 (M+)	
359	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(quinolin-3-yloxy)-acetamide	(quinolin-3-yloxy)acetic acid HCl (SALOR)	Ex 634	MS (ion spray) 517.2 (M+)	
360	2- <i>t</i> -Butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide	prep 305	Ex 634	MS (ion spray) 445.1 (M+)	

361	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-2-ylsulfanyl)-acetamide	(pyridin-2-ylsulfanyl)acetic acid (Maybridge)	Ex 634	MS (ion spray) 483.1 (M+)
362	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-pyridin-3-yl-acrylamide	trans-3-(3-pyridyl)acrylic acid	Ex 634	MS (ion spray) 463.1 (M+)
363	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-pyridin-2-yl-acrylamide	trans-3-Pyridin-2-yl-acrylic acid	Ex 634	MS (ion spray) 463.1 (M+)
364	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-3-pyridin-4-yl-acrylamide	trans-3-(4-pyridyl)acrylic acid	Ex 634	MS (ion spray) 463.0 (M+)
365	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbamoyl}-2-hydroxy-ethyl}-carbamic acid <i>t</i> -butyl ester	N-t-BOC-L-Serine	Ex 634	MS (ion spray) 519.0 (M+)
366	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbamoyl}-2-hydroxy-ethyl}-carbamic acid <i>t</i> -butyl ester	N-t-BOC-D-Serine	Ex 634	MS (ion spray) 519.0 (M+)
367	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(6-methoxy-pyridin-3-ylamino)-acetamide	prep 306	Ex 634	MS (ion spray) 496.2 (M+)
368	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-3-yloxy)-propionamide	prep 307	Ex 634	MS (ion spray) 481.2 (M+)
369	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(pyridin-2-yloxy)-acetamide	(pyridin-2-yloxy) acetic acid	Ex 634	MS (ion spray) 467.0 (M+)

370	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-2-phenylethyl}-carbamic acid <i>t</i> -butyl ester	N-BOC-L-phenyl-alanine	Ex 634	MS (ion spray) 579.2 (M+)	
371	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2,2-diphenyl-acetamide	benzyllic acid	Ex 634	MS (ion spray) 542 (M+)	
372	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(9H-fluoren-9-yl)-2-hydroxy-acetamide	9-hydroxy-9-fluorene-carboxylic acid	Ex 634	MS (ion spray) 540.1 (M+)	
373	N-[3 <i>R</i> -(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)] <i>R</i> -cycloheptylmethyl]-6-fluoro-nicotinamide	6-fluoro-nicotinic acid	prep 137	MS(ES+)m/z=4 83.1	
374	2-(Benzenesulfonyl-pyridin-2-yl-amino)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-acetamide	(benzenesulfonyl)pyridin-2-yl-amino acetic acid (Bionet)	prep 137	MS (ion spray) 606.1 (M+)	
375	2-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-methoxy-isonicotinamide	prep 355	2-chloro-6-methoxy-pyridine-4-carboxylic acid	MS (ion spray) 514.8 (M+)	
376	1-Methyl-piperidine-3-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-amide	prep 355	prep 357	MS (ion spray) 471.2 (M+)	
377	3-[[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-carbonyl]-piperidine-1-carboxylic acid <i>tert</i> -butyl ester	prep 355	prep 358	MS (ion spray) 557.2 (M+)	

378	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(1-methyl-1 <i>H</i> -indol-3-yl)-acetamide	prep 69	1-methyl-3-indoleacetic acid	MS (ion spray) 503.3 (M+)	
379	2-(Pyridin-3-yloxy)-hexanoic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide	prep 362	prep 357	MS (ion spray) 523.3 (M+)	
380	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	prep 380	<i>R</i> -Madelic acid	MS (ES+) m/z 466.1 (M+H) ⁺ , (ES-) 464.2 (M-H) ⁻	
381	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-2-(3-fluorophenyl)acetamide	prep 380	3-Fluoro-phenylacetic acid	MS (ES+) m/z 468.1 (M+H) ⁺ , (ES-) 466.2 (M-H) ⁻	
382	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-4-phenyl-butanamide	phenylbutyric acid	Prep 44	MS (ion spray) 492.2 (M+)	
383	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-4-oxo-4-phenyl-butanamide	3-benzoyl-propionic acid	Prep 44	MS (ion spray) 506.2 (M+)	
384	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-(4-hydroxy-phenyl)-propanamide	3-(4-hydroxy-phenyl)propionic acid	Prep 44	MS (ion spray) 494.2 (M+1)	
385	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-3-(1 <i>H</i> -indol-3-yl)-propanamide	3-indole-propionic acid	Prep 44	MS (ion spray) 516.9 (M+)	
386	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-2-dimethylamino-acetamide	N,N-dimethyl-glycine HCL	Prep 44	MS (ion spray) 431.2 (M+1)	

387	(2R)-N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-2-[<i>t</i> -butoxycarbonyl-amino]-3-(phenylmethoxy)propanamide	(D)-BOC-benzylxyserine	Prep 44	MS (ion spray) 523.2 (M-100)(M-BOC)
388	(2S)-N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-2-[<i>t</i> -butoxycarbonyl-amino]-3-(phenylmethoxy)propanamide	(L)-BOC-benzylxyserine	Prep 44	MS (ion spray) 523.0 (M-100)(M-BOC)
389	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-cyclohexyl-acetamide	Prep 47	cyclohexylamine	MS (ion spray) 456.2 (M+1)
390	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-phenyl-acetamide	Prep 47	aniline	MS (ion spray) 448.1 (M+1)
391	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-pyridin-2-yl-acetamide	Prep 47	2-aminopyridine	MS (ion spray) 451.2 (M+1)
392	2-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3-acetyl-phenyl)acetamide	Prep 47	3-aminoacetophenone	MS (ion spray) 492.2 (M+1)
393	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-N-(3-methanesulfonylphenyl)acetamide	Prep 47	3-methanesulfonylaniline HCl	MS (ion spray) 528.2 (M+)
394	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(4-fluoro-phenyl)-acetamide	Prep 47	4-fluoroaniline	MS (ion spray) 468.2 (M+1)
395	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3-fluoro-phenyl)-acetamide	Prep 47	3-fluoroaniline	MS (ion spray) 468.2 (M+1)
396	2-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3-methoxy-phenyl)-acetamide	Prep 47	3-methoxyaniline	MS (ion spray) 480.2 (M+1)

397	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3,5-dimethoxy-4-methylbenzamide	(3,5-dimethoxy-4-methyl)benzoic acid	Prep 48	MS (ion spray) 524.3 (M+1)	
398	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-1-methylpiperid-4-ylcarboxamide	1-methylpiperidine-4-carboxylic acid HCl	Prep 48	MS (ion spray) 471.2 (M+1)	
399	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-[1,2,3]thiadiazol-4-ylcarboxamide	1,2,3-thiadiazole-4-carboxylic acid	Prep 48	MS (ion spray) 458.1 (M+1)	
400	6-Chloro-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	6-chloronicotinic acid	Prep 48	MS (ion spray) 485.1 (M+)	
401	6-Methyl-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-nicotinamide	6-methylnicotinic acid	Prep 48	MS (ion spray) 465.1 (M+1)	
402	2-Methyl-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]nicotinamide	2-methylnicotinic acid	Prep 48	MS (ion spray) 465.1 (M+1)	
403	2-Fluoro-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]nicotinamide	2-fluoronicotinic acid	Prep 48	MS (ion spray) 469.1 (M+1)	
404	N-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-5-fluoro-nicotinamide	5-fluoronicotinic acid	Prep 48	MS (ion spray) 469.1 (M+1)	
405	N-[(1R,3S)-3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-2-methylsulfanylnicotinamide	2-(methylthio)nicotinic acid	Prep 48	MS (ion spray) 497.0 (M+)	
406	2-Chloro-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]isonicotine	2-chloroisonicotinic acid	Ex 615	MS (ion spray) 485.1, 487.1 (M+)	

407	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-pyrazin-2-yl carboxamide	pyrazinecarboxylic acid	Ex 615	MS (ion spray) 452.1 (M+1)
408	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-cinnolin-4-yl carboxamide	cinnoline-4-carboxylic acid	Ex 615	MS (ion spray) 502.2 (M+1)
409	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)cyclohexylmethyl]-1-methyl-1H-pyrrol-3-yl carboxamide	M-methyl-pyrrole-2-carboxylic acid	Ex 615	MS (ion spray) 453.2 (M+1)
410	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-1H-pyrazol-4-yl carboxamide	4-pyrazole-carboxylic acid	Ex 615	MS (ion spray) 440.1 (M+1)
411	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-1H-indol-3-yl carboxamide	indole-3-carboxylic acid	Ex 615	MS (ion spray) 489.1 (M+1)
412	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-1H-indol-2-yl carboxamide	indole-2-carboxylic acid	Ex 615	MS (ion spray) 488.1 (M+)
413	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-1-methyl-1H-indol-2-yl carboxamide	1-methyl-indole-2-carboxylic acid	Ex 615	MS (ion spray) 503.1 (M+)
414	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-1H-indol-4-yl carboxamide	indole-4-carboxylic acid	Ex 615	MS (ion spray) 489.1 (M+)
415	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)cyclohexylmethyl]-2-(1-methyl-1H-imidazol-4-yl)acetamide	1-methyl-4-imidazole acetic acid hydrochloride	Ex 615	MS (ion spray) 468 (M ⁺), 466 (M ⁻ - 1)
416	3-Benzoyl-N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazol-4,3-c/quinolin-5-yl)-cyclohexylmethyl]-benzamide	3-benzoylbenzoic acid	Ex 615	MS (ion spray) 554 (M ⁺), 552 (M ⁻ - 1)

417	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-phenoxy-benzamide	3-phenoxybenzoic acid	Ex 615	MS (ion spray) 542 (M ⁺), 540 (M ⁻ - 1)	
418	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-4-methoxythiophen-3-yl carboxamide	4-methoxythiophene-3-carboxylic acid	Ex 615	MS (ion spray) 486 (M ⁺), 484 (M ⁻ - 1)	
419	1,2,5-Trimethyl-1H-pyrrole-3-carboxylic acid [(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-amide	1,2,5-trimethylpyrrole-3-carboxylic acid	Ex 615	MS (ion spray) 481 (M ⁺)	
420	N-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-6-fluoronicotinamide	6-Fluoro-nicotinic acid	Ex 615	MS (ion spray) 469 (M ⁺)	
421	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl]-N-(3,4,5-trimethoxyphenyl)-acetamide	Prep 63	3,4,5-trimethoxy aniline	MS (ion spray) 450 (M ⁺), 448 (M ⁻ - 1)	
422	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl]-N-phenylacetamide	Prep 63	aniline	MS (ion spray) 540 (M ⁺), 538 (M ⁻ - 1)	
423	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-2,2-dimethylcyclobutyl]-N-pyridin-3-ylacetamide	Prep 63	3-aminopyridine	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)	
424	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(3-fluorophenyl)acetamide	3-fluorophenyl-acetic acid	Prep 68	MS (ion spray) 468 (M ⁺), 466 (M ⁻ - 1)	
425	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(3-fluorophenyl)acetamide	3-fluorophenyl-acetic acid	prep 69	MS (ion spray) 468 (M ⁺), 466 (M ⁻ - 1)	

426	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-pyridin-3-ylacetamide	3-pyridylacetic acid HCl	prep 69	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)	
427	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-pyridin-4-ylacetamide	4-pyridylacetic acid HCl	prep 69	MS (ion spray) 451 (M ⁺), 449 (M ⁻ - 1)	
428	N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(1-methyl-3H-imidazol-4-yl)acetamide	1-methyl-4-imidazoleacetic acid HCl	prep 69	MS (ion spray) 454 (M ⁺), 452 (M ⁻ - 1)	
429	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]carbonyl]-2,3-dihydroindole-1-carboxylic acid <i>t</i> -butyl ester	N-BOC-(S)-(-)-indoline-2-carboxylic acid	prep 69	MS (ion spray) 477 (M ⁺ -BOC + 1)	
430	[[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]carbonyl]-(4-fluorophenyl)methyl]carbamic acid <i>t</i> -butyl ester	N-BOC-L-4-fluorophenylglycine	prep 69	MS (ion spray) 483 (M ⁺ -BOC + 1)	
431	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2,2-diphenylacetamide	diphenylacetic acid	prep 69	MS (ES) 526.2 m/z[M+H] ⁺	
432	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-N-3,4,5-trimethoxyphenyl]acetamide	prep 94	3,4,5-trimethoxy aniline	MS (FIA) m/z= 540.5	diastereomers were separated
433	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclohexanecarboxylic acid cyclobutylamide	prep 102	cyclobutylamine	MS (ES+) (m/z) 428.1 [M+1]	
434	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclohexanecarboxylic acid cyclohexylamide	prep 102	cyclohexylamine	Mass Spectrum (FIA) (m/z) 456.3 [M+1]	
435	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclohexanecarboxylic acid phenylamide	prep 102	aniline	MS (FIA) (m/z) 448.1 [M-1]	
436	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclohexanecarboxylic acid pyridin-2-ylamide	prep 102	3-aminopyridine	Mass Spectrum (FIA) (m/z) 451.1 [M+1]	

437	N-(3,4,5-trimethoxyphenyl)[3-methyl-4-oxo-9-chloro-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide	prep 120	trimethoxyaniline	MS (ion spray) 526.2 (M+1)	diastereomers were separated
438	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-phenyl-acetamide	prep 145	aniline	MS (ion spray) 450 (M ⁺)	
439	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-pyridin-3-yl-acetamide	prep 145	3-aminopyridine	MS (ion spray) 450 (M ⁺)	
440	2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-N-(3-methoxy-phenyl)-acetamide	prep 145	3-Methoxy-phenylamine	MS (ion spray) 480 (M ⁺)	
441	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	Hydroxy-phenyl-pyridin-3-yl-acetic acid	prep 48	MS (ion spray) 543.1 (M+)	
442	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	R-Madelic acid	prep 380	MS (ES+) m/z 466.1 (M+H) ⁺ , (ES-) 464.2 (M-H) ⁻	
443	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	S-Madelic acid	prep 380	MS (ES+) m/z 466.1 (M+H) ⁺ , (ES-) 464.2 (M-H) ⁻	
444	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-cyclohexyl]-2-(3-fluoro-phenyl)-acetamide	3-F-Phenylacetic acid	prep 380	MS (ES+) m/z 468.1 (M+H) ⁺ , (ES-) 466.2 (M-H) ⁻	
445	N-[3-(9-Chloro-3-methyl-4-oxo-2,4-dihydro-pyrazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	R-Madelic acid	prep 399	MS (ES+) m/z 465.2 (M+H) ⁺	
446	N-[3-(9-Chloro-3-methyl-4-oxo-2,4-dihydro-pyrazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-acetamide	S-Madelic acid	prep 399	MS (ES+) m/z 465.2 (M+H) ⁺	

447	6-Hydroxy-pyridine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]-amide	6-hydroxy-picolinic acid	prep 48	MS (ion spray) 467 (M ⁺)	
448	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-pyridin-2-yl-acetamide	Ex 615	prep 69	MS (ion spray) 467 (M ⁺)	Prep 69 was hydro-lized by LiOH in dioxane at rt.
449	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-pyridin-3-yl-acetamide	Ex 615	prep 69	MS (ion spray) 467 (M ⁺)	Prep 69 was hydro-lized by LiOH in dioxane at rt.
450	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbonyl}-cyclopentyl}-carbamic acid tert-butyl ester	Ex 615	prep 69	MS (ion spray) 543 (M ⁺)	
451	{1-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]carbonyl}-cyclopropyl}-carbamic acid tert-butyl ester	Ex 615	prep 69	MS (ion spray) 515 (M ⁺)	
452	2-Amino-bicyclo[2.2.1]heptane-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide	Ex 615	prep 69	MS (ion spray) 469 (M ⁺)	
453	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-4-dimethylamino-2-phenoxy-2-phenyl-butylamide	Ex 615	prep 69	MS (ion spray) 613 (M ⁺)	Prep 69 was hydro-lized by LiOH in dioxane at rt.
454	S(+)-1R,3S-N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]-2-hydroxy-2-phenyl-acetamide	(S)(+) Mandelic acid	Ex 621	ESMS: 466 (M+1) ⁺	
455	R(-)-1R,3S-N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)cyclopentyl]-2-hydroxy-2-phenyl-acetamide	(R)(-) Mandelic acid	Ex 621	ESMS: 466 (M+1) ⁺	

456	S(+)-tert-Butoxycarbonylamino-acetic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl ester	S(+)-N-t-BOC-glycine	prep 69	MS (ion spray) 623 (M ⁺)	
457	R(-)-tert-Butoxycarbonylamino-acetic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl ester	R(-)-N-t-BOC-glycine	prep 69	MS (ion spray) 623 (M ⁺)	
458	2-(4-Acetyl-piperazin-1-yl)-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-pyridin-3-yl-acetamide	prep 395	prep 69	MS(exact mass) [M+H] ⁺ = 577.2340 m/z	Prep 69 was hydro-lized by LiOH in dioxane at rt.
459	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-pyridin-3-yl-2-(pyridin-2-yloxy)-acetamide	prep 396	prep 69	MS(ES) [M+H] ⁺ = 544.4 m/z; [M-H] ⁻ = 542.3 m/z	Prep 69 was hydro-lized by LiOH in dioxane at rt.
460	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-pyridin-3-yl-2-(pyridin-3-yloxy)-acetamide	prep 397	prep 69	MS(ES) [M+H] ⁺ = 544.3 m/z; [M-H] ⁻ = 542.5 m/z	Prep 69 was hydro-lized by LiOH in dioxane at rt.
461	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-pyridin-3-yl-2-(pyridin-4-yloxy)-acetamide	prep 398	prep 69	MS(ES) [M+H] ⁺ = 544.3 m/z; [M-H] ⁻ = 542.4 m/z	Prep 69 was hydro-lized by LiOH in dioxane at rt.
462	4-tert-Butoxycarbonylamino-butyric acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl ester	Ex 471	prep 69	MS(ES)[M+H] ⁺ = 651.2 m/z	
463	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl-2-pyridin-3-yl-acetamide	Hydroxy-phenyl-pyridin-3-yl-acetic acid	prep 69	MS(ion spray) 543.1 (M ⁺)	
465	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(4-fluorophenyl)acetamide	4-fluoro-phenylacetic acid	Ex 634	MS(ion spray) 468 (M ⁺)	no DMAP

466	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-oxo-2-phenylacetamide	benzoylformic acid	Ex 634	MS(ion spray) 464 (M ⁺).	
467	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexanecarboxylic acid benzylamide	Prep 43	benzylamine	MS(ion spray) 450 (M ⁺)	
468	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexanecarboxylic acid (pyridin-3-ylmethyl)amide	Prep 43	3-(aminoethyl)-pyridine	MS(ion spray) 451 (M ⁺)	
469	3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexanecarboxylic acid (pyridazin-3-ylmethyl)amide	Prep 43	3-(aminoethyl)-pyridine	MS(ion spray) 452 (M ⁺)	
470	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenylacetamide	L-(+)-mandelic acid	Ex 634	MS(ion spray) 466 (M ⁺)	
471	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenylacetamide	D-(-)-mandelic acid	Ex 634	MS(ion spray) 466 (M ⁺)	
472	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-phenyl-2-piperidin-1-ylacetamide	phenyl-piperidin-1-yl-acetic acid	Ex 634	MS(ion spray) 533 (M ⁺)	isomer 1 = 15.1mg isomer 2 = 8.8 mg
473	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-phenyl-propionamide	d/l-2-dimethyl-amino-3-phenyl-propionic acid	Ex 634	MS(ion spray) 507 (M ⁺)	
474	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methoxy-2-phenylpropionamide	2-methoxy-2-phenylpropionic acid	Ex 634	MS(ion spray) 494 (M ⁺)	

d) Table for Cyclization using KOtBu

Ex. #	Product	Starting Material	Physical Data
475	2-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-N-(3,4,5-trimethoxyphenyl)acetamide	prep 5	Racemic-68% yield
476	2-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]-N-(3,4,5-trimethoxyphenyl)acetamide	prep 12	MS(ES+)m/z =525.8 Racemic
477	2-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]-N-(3,4,5-trimethoxyphenyl)acetamide	prep 13	MS(ES+)m/z=525.9. Racemic
478	9-Chloro-3-methyl-5-{3-[2-(4-methyl-piperidin-1-yl)-2-oxo-ethyl]-cyclohexyl}-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 43	MS (ion spray) 456.2 (M+1) Racemic
479	<i>Cis</i> -2-[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexyl]-N-(3,4,5-trimethoxyphenyl)-acetamide	prep 158	MS (-ES) m/z 536.9 (M-H), 597.9 (M+OAc).

5 e) Table for Cyclization using KHMDS

Ex. #	Product	Starting Material	Physical Data	Comments
480	2-[(<i>1R,3S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]-N-(3,4,5-trimethoxyphenyl)acetamide	prep 18	MS(ES+)m/z=526. (single enantiomer)	
481	2-[(<i>1R,3S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]-N-(phenyl)acetamide	prep 28	MS(ES+)m/z=436. Single enantiomer	
482	2-[(<i>1R,3S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]-N-(4-fluorophenyl)acetamide	prep 28	MS(ES+)m/z=454. Single enantiomer	

483	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentyl]- <i>N</i> -(pyrid-3-yl)acetamide	prep 28	MS(ES+)m/z=437. Single enantiomer
484	<i>N</i> -{2-[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentyl]ethyl}benzamide	prep 31	MS(ES+)m/z=450 Single enantiomer
485	<i>N</i> -[(1 <i>S</i> ,3 <i>S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylethyl]- <i>N</i> -(3,4,5-trimethoxyphenyl)-acetamide	prep 33	MS(ES+)m/z=540. Single enantiomer
486	{3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-ylmethyl)cyclopentylethyl}- <i>N</i> -(3,4,5-trimethoxyphenyl)carboxamide	prep 36	MS(ES+)m/z=525.8 Racemic
487	9-Chloro-3-methyl-5 <i>H</i> -(3-phenylsulfanyl)methylcyclohexyl)-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	prep 78	MS(FD): M ⁺ 438.3 m/z Racemic
488	9-Chloro-3-methyl-5-(3-phenylmethanesulfonylmethylcyclohexyl)-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one Racemic	prep 82	MS(ES): (M+1) ⁺ 485.5, 487.5 m/z
489	[3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl benzoate Racemic	prep 84	MS(ES): (M+1) ⁺ 450.4, 542.4 m/z
490	9-Chloro-3-methyl-5-[3-(2-oxo-3-phenylpropyl)cyclohexyl]-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (<i>cis</i>)	prep 89	MS(ES): [M+H] ⁺ = 449.1 m/z, [M-H] ⁻ = 447.1 m/z
491	[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexyl]acetic acid benzyl ester (<i>cis</i>)	prep 84	MS (FIA) (m/z) 465.2 [M+1]
492	<i>N</i> -[3-(9-Fluoro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 106	MS (FIA) (m/z) 434.3 [M+1]
493	<i>N</i> -[3-(9-Chloro-4-oxo-3-phenyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 107	MS (FIA) (m/z) 512.4 [M+1]
494	<i>N</i> -[3-(9-Chloro-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 108	MS (ES+) (m/z) 436.2 [M+1]

495	<i>N</i> -[3-(6-Iodo-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 109	MS (FIA) (m/z) 540.3 [M-1]
496	<i>N</i> -[3-(8-Iodo-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 110	MS (FIA) (m/z) 542.2 [M+1]
497	<i>N</i> -[3-(7-Fluoro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 111	MS (FIA) (m/z) 434.3 [M+1]
498	<i>N</i> -[3-(9-Chloro-3-hexyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]benzamide (<i>cis</i>)	prep 112	MS (FIA) (m/z) 520.3 [M+1]
499	phenylmethyl 2-[3-(9-iodo-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]acetate (<i>cis</i>)	prep 113	MS (FIA) (m/z) 557.0 [M+1]
500	9-Chloro-3-methyl-5-[3-[2-(toluene-3-sulfonyl)-ethyl]-cyclohexyl]-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (<i>cis</i>)	prep 128	MS (ES): (M+1) ⁺ 499.2, 501.2 m/z
501	9-Chloro-3-methyl-5-[(3-phenylthioethyl)-cyclohexyl]-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (<i>cis</i>)	prep 131	MS (FD): M ⁺ 452.4 m/z
502	9-Chloro-3-methyl-5-[3-(2-phenylmethanesulfonyl-ethyl)-cyclohexyl]-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (<i>cis</i>)	prep 132	MS (ES): (M+1) ⁺ 499.2, 501.2 m/z
503	9-Chloro-5-[3-[2-(4-fluoro-benzenesulfonyl)-ethyl]-cyclohexyl]-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (<i>cis</i>)	prep 138	MS (ES): (M+1) ⁺ 503.1, 505.1 m/z
504	phenylmethyl 2-((3 <i>R</i> ,1 <i>R</i>)-3-[[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carbonyl-amino]cyclohexyl)acetate (<i>cis</i>)	prep 144	MS (ion spray) 465 (M ⁺)
505	<i>cis</i> -1-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-3-[[1,1-dimethylethoxy]carbonyl]amino]cyclohexane	prep 160	ESIMS <i>m/e</i> 432 35Cl (M ⁺ +1);434 37Cl (M ⁺ +1)
506	[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]carbamic acid <i>t</i> -butyl ester (<i>trans</i>)	prep 160	MS (ES+) m/z 446.1 (M+H) ⁺

507	[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]carbamic acid <i>t</i> -butyl ester (<i>cis</i>)	prep 160	MS (ES+) <i>m/z</i> 446.1 (<i>M</i> + <i>H</i>) ⁺
508	[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]carbamic acid benzyl ester (<i>cis</i>)	prep 162	MS (ES+) <i>m/z</i> 500.1 (<i>M</i> + <i>H</i>) ⁺
509	3-[2-(9-Chloro-3-methyl)-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclohexyl]- <i>N</i> -(3,4,5-trimethoxy-phenyl)propionamide	prep 167	MS (ES+) <i>m/z</i> 554.0 (<i>M</i> + <i>H</i>) ⁺
510	(<i>1R,3S</i>)-3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentanecarboxylic acid benzylamide	prep 170	ESMS: 436 (<i>M</i>) ⁺ , 437 (<i>M</i> +2) ⁺
511	(<i>1R,3S</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]benzamide	prep 179	ESMS: 436 (<i>M</i>) ⁺ , 437 (<i>M</i> +1) ⁺
512	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]benzamide	prep 179	ESMS: 436 (<i>M</i>) ⁺ , 437 (<i>M</i> +1) ⁺
513	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]-4-fluorobenzamide	prep 189	ESMS: 454 (<i>M</i>) ⁺ , 455 (<i>M</i> +1) ⁺ , 513 (<i>M</i> +59) ⁻
514	(<i>1S,3R</i>)-Biphenyl-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]amide	prep 192	ESMS: 512 (<i>M</i> +1) ⁺ , 570 (<i>M</i> +59) ⁻
515	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]-nicotinamide	prep 195	ESMS: 435 (<i>M</i> -1) ⁻ , 495 (<i>M</i> +59) ⁻
516	(<i>1S,3R</i>)-Furan-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]amide	prep 198	ESMS: 426 (<i>M</i> +1) ⁺ , 460 (<i>M</i> +35) ⁻ , 484 (<i>M</i> +59) ⁻
517	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclopentylmethyl]-3,4,5-trimethoxybenzamide	prep 201	ESMS: 526 (<i>M</i> +1) ⁺ , 584 (<i>M</i> +59) ⁻
518	(<i>1S,3R</i>)- <i>N</i> -[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)-cyclopentylmethyl]carbamic acid benzyl ester	prep 204	ESMS: 466 (<i>M</i> +1) ⁺ , 524 (<i>M</i> +59) ⁻

519	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentylmethyl]-2-(3,4,5-trimethoxyphenyl)acetamide	prep 207	ESMS: 540 (M+1)+, 574 (M+35)-,598 (M+59)-	
520	N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]carbamic acid <i>t</i> -butyl ester	prep 208	ESMS: 432 (M+1)+	
521	[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]carbamic acid <i>t</i> -butyl ester	prep 209	ESMS: 432 (M+1)+	
522	<i>trans</i> -[1-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]-quinolin-5-yl)-3-[[1,1-dimethylethoxy]carbonyl]amino]]-cyclohexane	prep 22	ESIMS <i>m/e</i> 432 ³⁵ Cl (M ⁺ +1); 434 ³⁷ Cl (M ⁺ +1)	
523	N-[3-(9-Cyano-3-methyl-4-oxo-5H-isoxazole[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 267	MS(ES+) (m/z) 441.2 [M+1]	
524	[3-(9-Cyano-3-methyl-4-oxo-5H-isoxazole[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-carbamic acid benzyl ester	prep 269	MS(ES+) (m/z) 471.2 [M+1]	
525	[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazole[4,3-c]quinolin-5-ylmethyl)-cyclohexyl]-carbamic acid <i>tert</i> -butyl ester	prep 278	MS(ES+) (m/z) 346.1 [M-BOC].	
526	<i>N</i> -[3-(9-Chloro-3-diethylamino-4-oxo-5H-isoxazole[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 280	MS(ES+) (m/z) 507.2 [M+1]	
527	<i>N</i> -[3-(9-Chloro-4-oxo-3-pyrroldin-1-yl-5H-isoxazole[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 281	MS(ES+) (m/z) 505.1 [M+1]	
528	<i>N</i> -[3-(9-Chloro-3-ethylamino-4-oxo-5H-isoxazole[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 280	MS(ES+) (m/z) 479.1 [M+1]	

529	<i>N</i> -[3-(9-Chloro-3-ethylsulfanyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 283	MS(ES+) (m/z) 496.1 [M+1].	
530	<i>N</i> -[3-(9-Chloro-4-oxo-3-phenylamino-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 278	MS(ES+) (m/z) 527.2 [M+1].	
531	<i>N</i> -[3-(9-Cyano-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 316	MS(ES+) (m/z) 441.2 [M+1].	
532	[3-(9-Cyano-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-carbamic acid benzyl ester	prep 318	MS(ES+) (m/z) 471.2 [M+1].	
533	[3-(9-Chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-ylmethyl)-cyclohexyl]-carbamic acid <i>tert</i> -butyl ester	prep 327	MS(ES+) (m/z) 346.1 [M-BOC].	
534	<i>N</i> -[3-(9-Chloro-3-diethylamino-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 329	MS(ES+) (m/z) 507.2 [M+1].	
535	<i>N</i> -[3-(9-Chloro-4-oxo-3-pyrrolidin-1-yl-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 330	MS(ES+) (m/z) 505.1 [M+1].	
536	<i>N</i> -[3-(9-Chloro-3-ethylamino-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 331	MS(ES+) (m/z) 479.1 [M+1].	
537	<i>N</i> -[3-(9-Chloro-3-ethylsulfanyl-4-oxo-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 332	MS(ES+) (m/z) 496.1 [M+1].	
538	<i>N</i> -[3-(9-Chloro-4-oxo-3-phenylsulfanyl-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 328	MS(ES+) (m/z) 544.2 [M+1].	Prep 328 was initially treated with Na thiophenoxide (5eq) in DMF.
539	<i>N</i> -[3-(9-Chloro-4-oxo-3-phenylamino-5 <i>H</i> -isoxazole[4,3- <i>c</i>]quinolin-5-yl)-cyclohexylmethyl]-benzamide	prep 328	MS(ES+) (m/z) 527.2 [M+1].	Prep 328 was initially treated with aniline (20eq) in DMF.

540	[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclohexyl]-carbamic acid tert.-butyl ester	Prep 333	MS(ES+) (m/z) 346.3 [M-BOC].
541	N-{2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-ethyl}-3,4,5-trimethoxy-benzamide	Prep 33	MS(ES+)m/z=540. Single enantiomer
541a	{3-[(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)methyl]cyclopentyl}-N-(3,4,5-trimethoxyphenyl)-carboxamide	Prep 337	MS(ES+)m/z=525.8
542	[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cycloheptyl]-acetic acid methyl ester	Prep 338	MS(ES+)m/z=402.9.
543	N-{2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-ethyl}-benzamide	Prep 31	MS(ES+)m/z=450.
544	2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclopentyl]-N-(3,4,5-trimethoxyphenyl)-acetamide	Prep 337	MS(ES+)m/z=526. (Single enantiomer)
545	[3-(9-Chloro-3-methyl-4-oxo-3,4-dihydro-imidazo[4,5-c]quinolin-5-yl)-cyclohexylmethyl]-carbamic acid benzyl ester	Ex 405	MS (ES+) (m/z) 479.3 [M+1]

f) Table for Arylation:

Ex. #	Product	Starting Material A	Starting Material B	Physical Data
546	9-Chloro-3-methyl-5-[3-(quinolin-2-ylaminomethyl)-cyclohexyl]-5H-isoxazolo[4,3-c]quinolin-4-one	Ex 619	2-chloroquinoline	MS(FLA) (m/z) 473.1 (M+1)
547	9-Chloro-3-methyl-5-[3-[(5-nitro-pyridin-2-ylamino)-methyl]-cyclohexyl]-5H-isoxazolo[4,3-c]quinolin-4-one	Ex 619	2-chloro-6-nitropyridine	MS(FLA) (m/z) 466.1 (M-1)
548	2-[[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexylmethyl]amino]-nicotinonitrile	Ex 619	2-chloro-3-nitrilepyridine	MS(FLA) (m/z) 448.1 (M+1)

549	9-Chloro-3-methyl-5-{3-[(5-nitropyridin-2-ylamino)methyl]-cyclohexyl}-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	Ex 619	2-chloropyrimidine	MS(FIA) (m/z) 424.1 (M+1)
550	9-Chloro-5-{3-[4-chloropyrimidin-2-ylamino)-methyl]cyclohexyl}-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one and 9-Chloro-5-{3-[2-chloropyrimidin-4-ylamino)-methyl]-cyclohexyl}-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one (racemic)	Ex 619	2,4-dichloropyrimidine	MS(FIA) (m/z) 458.4 (M+1) isomers were separated
551	9-Chloro-5-{3-[(6-chlorobenzothiazol-2-ylamino)methyl]-cyclohexyl}-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	Ex 619	2,6-dichlorobenzothiazole	MS(FIA) (m/z) 515.2 (M+1)
552	5-[3-(Benzothiazol-2-ylaminomethyl)-cyclohexyl]-9-chloro-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	Ex 619	2-chlorobenzothiazole	MS(FIA) (m/z) 479.1 (M+1)
553	5-[3-(Benzooxazol-2-ylaminomethyl)cyclohexyl]-9-chloro-3-methyl-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-4-one	Ex 619	2-chlorobenzooxazole	MS(FIA) (m/z) 463.3 (M+1)

g) Table for Chiral Separation:

Ex. #	Product	Starting Material	Physical Data
554	2-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-N-(3,4,5-trimethoxyphenyl)acetamide	Ex 432	Retention Time = 8.56 minutes
555	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl]-N-(3,4,5-trimethoxyphenyl)acetamide	Ex 432	Retention Time = 16.08 minutes
556	N-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo(5-hydroisoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl)methyl]benzamide	Ex 116	Retention Time = 9.232 min
557	N-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,3- <i>c</i>]quinolin-5-yl)cyclohexyl)methyl]benzamide	Ex 116	Retention Time = 12.704 min

-267-

558	2-[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -(3,4,5-trimethoxyphenyl)acetamide	Ex 479	MS (-ES) <i>m/z</i> 536.9 (M-H), 597.9 (M+OAc)
559	2-[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -(3,4,5-trimethoxyphenyl)acetamide	Ex 479	MS (-ES) <i>m/z</i> 536.9 (M-H), 597.9 (M+OAc)
560	<i>N</i> -{[(1 <i>S</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl}(<i>t</i> -butoxy)carboxamide	Ex 506	MS (+ES) <i>m/z</i> 446 (M+H)+, 463 (M+NH3)+, (-ES) 480 (M-H+Cl)-, 504 (M-H+OAc)-
561	<i>N</i> -{[(1 <i>R</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl}(<i>t</i> -butoxy)carboxamide	Ex 506	MS (+ES) <i>m/z</i> 446 (M+H)+, 463 (M+NH3)+, (-ES) 480 (M-H+Cl)-, 504 (M-H+OAc)-
562	<i>N</i> -{[(1 <i>R</i> ,3 <i>S</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl}(<i>t</i> -butoxy)carboxamide	Ex 507	MS (+ES) <i>m/z</i> 446 (M+H)+, 463 (M+NH3)+, (-ES) 480 (M-H+Cl)-, 504 (M-H+OAc)-
563	<i>N</i> -{[(1 <i>S</i> ,3 <i>R</i>)-3-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]methyl}(<i>t</i> -butoxy)carboxamide	Ex 507	MS (+ES) <i>m/z</i> 446 (M+H)+, 463 (M+NH3)+, (-ES) 480 (M-H+Cl)-, 504 (M-H+OAc)-
564	3-[(1 <i>S</i> ,2 <i>S</i>)-2-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -(3,4,5-trimethoxyphenyl)propanamide	Ex 618	MS (+ES) 553.9, 555.9 (M+H)+
565	3-[(2 <i>S</i> ,1 <i>R</i>)-2-(9-chloro-3-methyl-4-oxo-5 <i>H</i> -isoxazolo[4,5- <i>c</i>]quinolin-5-yl)cyclohexyl]- <i>N</i> -(3,4,5-trimethoxyphenyl)propanamide	Ex 618	MS (+ES) 553.9, 555.9 (M+H)+

-268-

Example 566

(2R)-N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclo-hexylmethyl]-2-amino-3-(phenylmethoxy)propanamide

5 A solution of (2R)-N-{[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]methyl}-2-[(*t*-butoxy)carbonylamino]-3-(phenylmethoxy)-propanamide, 80 mg (0.13 mmol) in 10 mL of acetic acid saturated with hydrochloric acid was stirred for 2 h at ambient temperature and concentrated to dryness. The residue was dissolved in toluene and concentrated to dryness and dried
10 under vacuum to yield a quantitative yield of the desired product as a white foam. ¹H-NMR is consistent with structure. MS (ion spray) 523.2 (M+).

Example 567

15 N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-3-aminobenzamide

To a solution containing 620 mg (1.25 mmol) of a compound from Example 48 in 15 mL of THF was added 848 mg (3.75 mmol) of SnCl₂·2H₂O at room temperature. To the resulting suspension was added 1.0 mL of concentrated HCl and
20 the reaction was stirred for 15 hours at room temperature. The reaction was diluted with 100 mL of ethyl acetate and washed once with brine. The organic solution was separated, dried and concentrated to give a white foam. This crude material was purified by flash chromatography, using ethyl acetate as the eluent. The major fractions were combined and concentrated *in vacuo* to give 400 mg of a white
25 amorphous solid. MS(FIA) m/z = 465.2.

-269-

Example 568

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-4-aminobenzamide

5 The title compound was prepared from the compound from Example 110 according to the conditions for the preparation of Example 567. MS(FIA) *m/z* = 465.2.

Example 569

10 *N*-[3*S*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-*R*-cyclohexylmethyl]-3-(4-methoxy)benzyloxybenzamide

To a suspension of 150 mg (0.581 mmol) of a compound from preparation 56 in 10 mL of CH₂Cl₂ was added 2 drops of DMF, followed by the addition of 51 μL
15 (0.581 mmol) of oxalyl chloride. The reaction was stirred at room temperature and evolution of CO gas was noted. The reaction was stirred an additional 30 minutes at room temperature and then concentrated *in vacuo*. The resulting acid chloride was dissolved in 10 mL of CH₂Cl₂. In a separate flask, 250 mg (0.528 mmol) of the compound from Example 302, was suspended in 10 mL of CH₂Cl₂. To this was
20 added 110 μL of triethylamine and the resulting solution was added to the flask containing the acid chloride, followed by the addition of an additional 110 μL of triethylamine. The reaction was stirred at room temperature for 15 hours, diluted with 50 mL of ethyl acetate and washed twice with 1N HCl, twice with 1N NaOH, dried over sodium sulfate and concentrated *in vacuo* to give a white amorphous solid. This
25 material was recrystallized from toluene to give 190 mg of a white amorphous solid. MS(ES+) *m/z* = 586.19.

-270-

Example 570

N-[3*S*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-*R*-cyclohexylmethyl]-3-hydroxybenzamide

5 To a solution containing 50 mg (0.0853 mmol) of a compound from Example 569 in 2 mL of CH₂Cl₂ was added 2 mL of TFA. The reaction was stirred at room temperature for 18 hours and concentrated *in vacuo*. The resulting crude solid was recrystallized from toluene to give a white amorphous solid. MS(ES+) *m/z* = 466.1.

10

Example 571

N-[3*S*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-*R*-cyclohexylmethyl]-4-(4-methoxy)benzyloxybenzamide

15 The title compound was prepared from methyl 4-hydroxybenzoate according to the conditions for the preparation of Example 569. MS(ES+) *m/z* = 586.19.

Example 572

N-[3*S*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-*R*-cyclohexylmethyl]-4-hydroxybenzamide

20

The title compound was prepared from a compound of Example 571 according to the conditions for the preparation of Example 570. MS(ES+) *m/z* = 466.1.

Example 573

25 *N*-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-benzenesulfonamide

30 *N*-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]benzenethioamide (0.78 g; 1.59 mmol) was dissolved in anhydrous dimethylformamide (16 mL), chilled to -20 °C for 10 min, and mixed with 0.5 M potassium bis(trimethylsilyl)amide in toluene (7.0 mL; 3.5 mmol; 2.2 equiv). The

-271-

reaction solution was stirred 10 min at $-20\text{ }^{\circ}\text{C}$ and another 35 min at room temperature. After quenching the reaction with 1 M HCl (50 mL), the solution was extracted with ethyl acetate (twice). The organic layer was washed with saturated NaCl_(aq) (twice), then dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The resulting white solid was purified by radial chromatography (three consecutive times) on a 4 mm thick silica gel rotor with a 5% tetrahydrofuran/dichloromethane (v/v) mobile phase and then twice with a 50% ethyl acetate/hexanes (v/v) mobile phase. Concentration of product-containing fractions produced 11 mg (2% yield) of a white solid. MS(ES) calc'd: $[\text{M}+\text{H}]^{+} = 472.0\text{ m/z}$, $[\text{M}-\text{H}]^{-} = 470.0\text{ m/z}$. Found: 471.9 m/z, 470.0 m/z.

Example 574

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]benzenesulfonamide

15

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]benzenethioamide (0.41 g; 0.83 mmol) was dissolved in anhydrous dimethylformamide (8 mL), chilled to $-20\text{ }^{\circ}\text{C}$ for 10 min, and mixed with 0.5 M potassium bis(trimethylsilyl)amide in toluene (4 mL; 1.83 mmol; 2.2 equiv). The reaction solution was stirred 10 min at $-20\text{ }^{\circ}\text{C}$ and another 35 min at room temperature. After quenching the reaction with 1 M HCl (50 mL), the solution was extracted with ethyl acetate (twice). The organic layer was washed with saturated NaCl_(aq) (twice), then dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The resulting white solid was purified by radial chromatography on a 4 mm thick silica gel rotor with a 40% ethyl acetate/hexanes (v/v) mobile phase. Product containing fractions were combined, concentrated and further purified by radial chromatography (twice) on a 2 mm thick silica gel rotor with a 50% ethyl acetate/hexanes (v/v) mobile phase. Concentration of product-containing fractions produced 53 mg (14% yield) of a white solid. MS(ES) calc'd: $[\text{M}+\text{H}]^{+} = 472.0\text{ m/z}$, $[\text{M}-\text{H}]^{-} = 470.0\text{ m/z}$. Found: 472.0 m/z, 470.2 m/z.

30

-272-

Example 575

4-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-
cyclohexanecarboxylic acid 3,4,5-trimethoxybenzylamide

5 The compound from preparation 72 (0.40 g, 0.00098 mol) was combined with
THF (5 mL), MeOH (2 mL), water (5 mL) and 2N NaOH (5 mL) and the mixture
stirred at ambient temperature until hydrolysis was complete. The mixture was
concentrated *in vacuo* and the residue taken up in water and carefully acidified using
aq 1N HCl. The aqueous mixture was extracted with ethyl acetate and the combined
10 extracts were dried over sodium sulfate. Concentration *in vacuo* left the crude acid
which was combined with 1-(3-dimethyl-aminopropyl-3-ethylcarbodiimide
hydrochloride (0.186 g, 0.00097 mol), 1-hydroxy-7-azabenzotriazole (0.133 g,
0.00098 mol) and 3,4,5-trimethoxybenzylamine (0.193 g, 0.00098 mol) in DMF (15
mL) and the resulting mixture stirred overnight at ambient temperature. The mixture
15 was concentrated *in vacuo* and the residue taken up in water. The aqueous mixture
was extracted with CH₂Cl₂ and the combined extracts dried over sodium sulfate and
concentrated *in vacuo*. The resulting residue was chromatographed over silica gel
using CH₂Cl₂/THF as eluent which allowed for isolation of 0.120 g (40%) of the
desired product as a yellowish solid. MS(ES): (M+1)⁺ 540.4 m/z.

20

Example 576 - 577

5-(3-Benzenesulfinylmethylcyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
c]quinolin-4-one &

25

5-(3-Benzenesulfonylmethylcyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
c]quinolin-4-one

30

The compound from Example 487 (0.20 g, 0.00046 mol) was dissolved in
CH₂Cl₂ (30 mL) and the mixture cooled under a nitrogen atmosphere in a dry
ice/acetone bath. Then 3-chloroperoxybenzoic acid (50%, 0.16 g, 0.00046 mol) was
added and the mixture stirred for 2 h while warming near ambient temperature. The
mixture was concentrated and the residue chromatographed over silica gel using

-273-

CH₂Cl₂/MeOH as eluent which allowed for isolation of both the sulfoxide (0.132 g, 63%). MS(ES): (M+1)⁺ 455.1, 457.3 m/z, and the sulfone (0.078 g, 36%) MS(ES): (M+1)⁺ 471.2 m/z.

5

Example 578

9-Chloro-5-[3-(2-methoxyimino-3-phenylpropyl)cyclohexyl]-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

To a solution of the compound from Example 490 in denatured ethanol (6 mL) was added a solution of methoxyamine hydrochloride (74.5 mg; 0.892 mmol; 4 equiv) and sodium acetate (73.1 mg; 0.892 mmol; 4 equiv) in water (1 mL). The cloudy reaction mixture became clear and colorless when heated to reflux. After 1.6 h the reaction solution was concentrated to dryness by rotary evaporation. The resulting solid was dissolved in dichloromethane and the organic layer was washed with water (once), washed with saturated NaCl_(aq) (once), dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The product was isolated as a mixture of oxime isomers by radial chromatography on a 2 mm thick silica gel rotor with a 2% acetonitrile/dichloromethane (v/v) mobile phase. A clear, colorless oil (92 mg; 86%) was obtained after concentration of the appropriate fractions. TOF-MS(ES) calc'd: [M+H]⁺ = 478.1897 m/z. Found: 478.1891 m/z.

20

Example 579

2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-N-4-methoxyphenylacetamide

25

To a 7 mL scintillation vial containing 0.10 g (0.254 mmol) of a compound from preparation 95 in 3 mL of methylene chloride was added 0.086 g (0.509 mmol) *p*-anisidine followed by 0.10 mL of triethylamine. The reaction was shaken overnight, passed through a 2 g SCX column, eluting with methylene chloride and the racemic *cis* product crystallized out of one of the fractions giving 0.054 g (44%). MS(FIA) m/z= 480.1.

30

-274-

Example 580

2-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-*N*-2-methoxy-5-nitrophenylacetamide

5

The title compound was prepared from a compound of preparation 95 and 2-methoxy-5-nitroaniline triethylamine in a manner similar to that of Example 579 to yield 0.051 g (38%). MS(FIA) m/z = 525.2.

10

Example 581

9-Chloro-3-methyl-5-[3-(2-oxo-pyrrolidin-1-ylmethylcyclohexyl)-5*H*-isoxazolo[4,3-*c*]quinolin-4-one

To a stirred solution of the compound of Example 105 (148 mg, 0.33 mmol) in DMF (3 mL) and THF (4 mL) was added NaH (41 mg, 1.0 mmol, 60% in mineral oil) and stirred at r.t. for 1.5 hours. The reaction mixture was diluted with ethyl acetate, washed (brine), dried (Na₂SO₄), filtered and concentrated. Column chromatography (silica gel, ethyl acetate) gave 82 mg, 60%. MS(FIA) (m/z) 414.2 (M+1).

20

Example 582

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclopentyl-2-(3,4,5-trimethoxyphenylmethyl)]acetamide

To a stirred solution of *N*-[(1*S*,4*R*)-4-(9-chloro-3-methyl-4-oxo(5-hydroisoxazolo[4,3-*c*]quinolin-5-yl))cyclopent-2-enyl]-2-(3,4,5-trimethoxyphenyl)-acetamide (55 mg, 0.11 mmol) in DMF (5 mL) was added Rh/C (5%) and treated with H₂ and stirred at r.t. for 18 hours. The mixture was filtered. The filtrate was diluted with EtOAc, washed (brine), dried (Na₂SO₄), filtered and concentrated. Column chromatography (silica gel, hexanes/ethyl acetate, gradient) gave the title compound 38 mg, 69%. MS(FIA) (m/z) 526.3 (M+1).

30

-275-

Example 583

N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclopentyl-2-(3-fluorophenylmethyl)]acetamide

5 To a stirred solution of N-[(1*S*,4*R*)-4-(9-chloro-3-methyl-4-oxo(5-hydroisoxazolo[4,3-*c*]quinolin-5-yl))cyclopent-2-enyl]-2-(3,4,5-trimethoxyphenyl)-acetamide (49 mg, 0.11 mmol) in EtOAc (5 mL) was added Rh/C (5%, catalytic amount) and treated with H₂ and stirred at r.t. for 18 hours. The mixture was filtered. The filtrate was diluted with EtOAc, washed (brine), dried (Na₂SO₄), filtered and
10 concentrated. Column chromatography (silica gel, hexanes/ethyl acetate, gradient) gave the title compound 48 mg, 97%. MS(FIA) (m/z) 454.1 (M+1).

Example 584 and 585

15 3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-ylmethyl)cyclohexane-carboxylic acid (3,4,5-trimethoxyphenyl)

A compound from preparation 101 (0.2 g, 0.54 mmol), 1 N NaOH (1.32 ml, 1.32 mmol), and MeOH (8 ml) were heated at 50 °C for 2 h. The reaction was cooled to r.t., acidified to pH < 3, and concentrated using benzene to azeotrope. The residue,
20 EDCI (0.2 g, 1.07 mmol), 3,4,5-trimethoxyaniline (0.119 g, 0.65 mmol), DMAP (0.013 g, 0.11 mmol), and dichloromethane (5.5 ml) were mixed under N₂. The mixture was diluted with EtOAc, washed, dried, filtered, and concentrated to give 585 (0.05 g, 17%) and 584 (0.07 g, 24%) after radial chromatography (silica gel, 5% acetone/ dichloromethane). 585: MS(FIA) (m/z) 540.4 [M+1]. 584: MS(FIA) (m/z)
25 540.4 [M+1].

-276-

Example 586

3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-ylmethyl)cyclohexane-carboxylic acid (2-methoxy-5-nitrophenyl)

5 In a fashion similar to that described for Example 585, methyl 3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexane carboxylate (0.2 g, 0.54 mmol), 1 N NaOH (1.32 ml, 1.32 mmol), MeOH (8 ml), EDCI (0.2 g, 1.07 mmol), 2-methoxy-5-nitroaniline (0.11 g, 0.65 mmol), DMAP (0.013 g, 0.11 mmol), dichloromethane (5.5 ml) gave b (0.052 g, 18%) and a (0.068 g, 24%) after radial
10 chromatography (silica gel, 2% acetone/ dichloromethane).

b: MS(FIA) (m/z) 523.1 [M-1].

a: MS(FIA) (m/z) 523.1 [M-1].

Example 587

15 9-Chloro-5-[3-(4,4-dimethyl-4,5-dihydro-oxazol-2-ylmethyl)cyclohexyl]-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one

The compound from preparation 43 (0.2 g, 0.54 mmol) was dissolved in dichloromethane (3.6 ml) followed by addition of oxalyl chloride (0.108 ml, 1.08
20 mmol) and DMF (0.005 ml). After 1 h of stirring, the volatiles were removed. The residue was dissolved in dichloromethane (1.5 ml) and 2-amino-2-methyl-1-propanol (0.1 ml, 1.08 mmol) was added. The reaction was stirred for 3 h then concentrated. The reaction was dissolved in thionyl chloride (2 ml) and stirred for 40 min. The
25 reaction was concentrated, added 0.1 N NaOH, and extracted with EtOAc (3x). The combined organic layers were washed (brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, hexanes/EtOAc gradient) gave the title compound (0.06 g, 26%). MS(FIA) (m/z) 428.2 [M+1].

-277-

Example 588 & 589

5-(3-Benzooxazol-2-ylmethyl-cyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]-
quinolin-4-one &
5 2-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-*N*-(2-
hydroxyphenyl)acetamide

In a fashion similar to that described for Example 587, compound from
preparation 43 (0.159 g, 0.423 mmol), oxalyl chloride (0.073 ml, 0.84 mmol),
10 dichloromethane (2.8 ml), DMF (0.005 ml), 2-aminophenol (0.2 g, 1.07 mmol),
dichloromethane (1.5 ml), and thionyl chloride (2 ml) gave the title compounds 5-(3-
benzooxazol-2-ylmethyl-cyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-
4-one, 588, (0.089 g, 47%) and 2-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinolin-5-yl)-cyclohexyl]-*N*-(2-hydroxy-phenyl)-acetamide, 589, (0.031 g, 16%)
15 after column chromatography (silica gel, acetone/dichloromethane gradient).
588: MS(FIA) (m/z) 448.1 [M+1]. 589: MS(FIA) (m/z) 466.2 [M+1].

Example 590

9-Chloro-3-methyl-5-[3-(4-phenyl-4,5-dihydro-oxazol-2-ylmethyl)cyclohexyl]-5*H*-
20 isoxazolo[4,3-*c*]quinolin-4-one

In a fashion similar to that described for Example 587, a compound from
preparation 43 (0.159 g, 0.423 mmol), oxalyl chloride (0.073 ml, 0.84 mmol),
dichloromethane (2.8 ml), DMF (0.005 ml), R-(-)-2-phenylglycinol (0.115 g, 0.84
25 mmol), dichloromethane (1.5 ml), and thionyl chloride (2 ml) gave the title compound
(0.105 g, 52%) after column chromatography (silica gel, acetone/ dichloromethane
gradient). MS(FIA) (m/z) 476.1 [M+1].

-278-

Example 591

9-Chloro-3-methyl-5-[3-(4-phenyl-4,5-dihydro-oxazol-2-ylmethyl)cyclohexyl]-5H-isoxazolo[4,3-c]quinolin-4-one

5 In a fashion similar to that described for Example 587, compound from preparation 43 (0.159 g, 0.423 mmol), oxalyl chloride (0.073 ml, 0.84 mmol), dichloromethane (2.8 ml), DMF (0.005 ml), S-(+)-2-phenylglycinol (0.115 g, 0.84 mmol), dichloromethane (1.5 ml), and thionyl chloride (2 ml) gave the title compound (0.07 g, 35%) after column chromatography (silica gel, acetone/ dichloromethane
10 gradient). MS(FIA) (m/z) 476.1 [M+1].

Example 592

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-4,5-dihydroisoxazolo[4,3-c]-quinoline-8-carboxylic acid methyl ester

15 To a mixture of Example 496 (0.034 g, 0.063 mmol), acetonitrile (8 ml), MeOH (3 ml), Et₃N (0.15 ml, 0.2 mmol) submerged in a 65 °C oil bath was added PdCl₂(dppf) (0.003 g, 0.003 mmol) under CO atmosphere (balloon). The reaction was stirred for 5 h. The reaction was cooled to room temperature and concentrated.
20 The residue was diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, acetone/dichloromethane gradient) gave the title compound (0.021 g, 70%). MS(ES+) (m/z) 474.2 [M+1].

-279-

Example 593

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-4,5-dihydroisoxazolo[4,3-c]-quinoline-6-carboxylic acid methyl ester

5 In a fashion similar to that described for Example 592, *N*-[3-(6-Iodo-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]-benzamide (0.048 g, 0.089 mmol), acetonitrile (10 ml), MeOH (5 ml), Et₃N (0.04 ml, 0.267 mmol), and PdCl₂(dppf) (0.004 g, 0.004 mmol) gave the title compound (0.035 g, 83%) after column chromatography (silica gel, acetone/dichloromethane gradient).

10 MS(FIA) (m/z) 472.3 [M-1].

Example 594

[3-(9-Iodo-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-5-yl)cyclohexylmethyl]-carbamic acid *t*-butyl ester

15

To a solution of phenylmethyl 2-[(1*R*,3*S*)-3-(9-iodo-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]acetate (1 g, 2.14 mmol) and *t*-BuOH (15 ml) submerged in a 50 °C oil bath was added Et₃N (0.386 ml, 2.78 mmol) and DPPA (0.6 ml, 2.78 mmol) under N₂. This solution was heated at reflux overnight. The
20 solvents were removed in vacuum and the residue was chromatographed (silica gel, EtOAc/hexanes gradient) to give the title compound (0.47 g, 40%). MS(FIA) (m/z) 538.3 [M+1], 438.0 [M-BOC].

-280-

Example 595

N-[3-(3-Methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]benzamide

5 To a mixture of Example 106 (0.07 g, 0.129 mmol) and tetrakis(triphenylphosphine)palladium (0.015 g, 0.0129 mmol) in toluene (1.3 ml) under N₂ was added tributyltin hydride (0.042 ml, 0.155 mmol) and heated at 80 °C for 17 h. Added more tributyltin hydride (0.069 ml, 0.258 mmol) to the mixture and heated at this temperature for 24 h. The reaction was cooled to room temperature, applied to a
10 silica gel column, and elution (acetone/dichloromethane gradient) gave the title compound (0.02 g, 38%).

MS(ES+) (m/z) 416.2 [M+1].

Example 596

15 *N*-{3-[9-(4-Methoxyphenyl)-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]benzamide

To a mixture of Example 106 (0.1 g, 0.185 mmol), tetrakis(triphenylphosphine)palladium (0.043 g, 0.037 mmol), and 4-methoxyphenylboronic acid (0.7
20 g, 0.46 mmol) in dioxane (2 ml) under N₂ was added Na₂CO₃(aq) (0.185 ml, 0.37 mmol) and heated at reflux for 72 h. The reaction was cooled to room temperature, diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Radial chromatography (silica gel, 33.3% EtOAc/hexanes) gave the title compound (0.057 g, 59%). MS(FIA) (m/z) 522.2 [M+1].

25

-281-

Example 597

5-[3-(Benzoylamino-methyl)-cyclohexyl]-3-methyl-4-oxo-5-hydro-isoxazolo[4,3-c]quinoline-9-carboxylic acid methyl ester

5 In a fashion similar to that described for Example 272, a compound from Example 106 (0.88 g, 1.63 mmol), acetonitrile (60 ml), MeOH (20 ml), Et₃N (0.68 ml, 4.89 mmol), and PdCl₂(dppf) (0.066 g, 0.08 mmol) gave the title compound (0.687 g, 89%) after column chromatography (silica gel, acetone/dichloromethane gradient).

10 MS(FIA) (m/z) 474.1 [M+1].

Example 598

5-[3-(Benzoylamino-methyl)cyclohexyl]-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinoline-9-carboxylic acid

15

A compound from Example 597 (0.35 g, 0.74 mmol), 5.0 N NaOH (2.4 ml, 11.84 mmol), and dioxane (7 ml) were allowed to react for 3 h at 90 °C in a fashion similar to that of Example 472, give the title compound (0.28 g, 85%). MS(ES+) (m/z) 460.2 [M+1].

20

Example 599

5-[3-(Benzoylamino-methyl)cyclohexyl]-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinoline-9-carboxylic acid amide

25

To a solution of Example 598, (0.05 g, 0.11 mmol), EDCI (0.032 g, 0.165 mmol), and HOBt (0.023 g, 0.165 mmol) in DMF (1.5 ml) under N₂ was added NH₄Cl (0.009 g, 0.165 mmol) and diisopropylamine (0.046 ml, 0.33 mmol). The reaction was stirred overnight. The mixture was diluted with EtOAc, washed (1.0 N HCl saturated with NaCl), dried (MgSO₄), filtered, and concentrated. Column

30

chromatography (silica gel, hexanes/ EtOAc gradient) gave the title compound (0.024 g, 48%). MS(ES+) (m/z) 459.1 [M+1].

-282-

Example 600

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinoline-9-carboxylic acid diethylamide

5

In a fashion similar to that described for Example 596, a compound from Example 598 (0.05 g, 0.11 mmol), EDCI (0.032 g, 0.165 mmol), HOBT (0.023 g, 0.165 mmol), DMF (1.5 ml), and diethylamine (0.017 ml, 0.165 mmol) gave the title compound (0.039 g, 68%) after column chromatography (silica gel,

10 acetone/dichloromethane gradient).

MS(ES+) (m/z) 515.2 [M+1].

Example 601

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinoline-9-carboxylic acid ethyl ester

15

To a solution of Example 598, (0.05 g, 0.11 mmol) in DMF (1 ml) under N₂ was added iodoethane (0.062 ml, 0.33 mmol) and Cs₂CO₃ (0.036 g, 0.11 mmol). The mixture was stirred overnight, diluted with EtOAc, washed (H₂O then brine), dried
20 (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, acetone/
dichloromethane gradient) gave the title compound (0.043 g, 80%). MS(FIA) (m/z)
488.4 [M+1].

-283-

Example 602

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-9-carboxylic acid benzyl ester

5 In a fashion similar to that described for Example 601, a compound from Example 598 (0.05 g, 0.11 mmol), benzyl bromide (0.039 ml, 0.33 mmol), Cs₂CO₃ (0.036 g, 0.11 mmol), and DMF (1ml) gave the title compound (0.051 g, 85%) after column chromatography (silica gel, acetone/dichloromethane gradient). MS(FIA) (m/z) 548.3 [M-1].

10

Example 603

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-9-carboxylic acid ethoxycarbonylmethyl ester

15 In a fashion similar to that described for Example 601, a compound from Example 598 (0.05 g, 0.11 mmol), ethyl bromoacetate (0.036 ml, 0.33 mmol), Cs₂CO₃ (0.036 g, 0.11 mmol), and DMF (1 ml) gave the title compound (0.054 g, 90%) after column chromatography (silica gel, acetone/dichloromethane gradient). MS(FIA) (m/z) 546.2 [M+1].

20

Example 604

5-[3-(Benzoylaminomethyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-9-carboxylic acid isopropyl ester

25 In a fashion similar to that described for Example 601, a compound from Example 598 (0.05 g, 0.11 mmol), 2-iodopropane (0.033 ml, 0.33 mmol), Cs₂CO₃ (0.036 g, 0.11 mmol), and DMF (1 ml) gave the title compound (0.021 g, 38%) after column chromatography (silica gel, acetone/dichloromethane gradient). MS(ES+) (m/z) 502.2 [M+1].

30

-284-

Example 605

5-[3-(Benzoylamino-methyl)-cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-9-carboxylic acid ethylamide

- 5 In a fashion similar to that described for Example 599, a compound from Example 598 (0.05 g, 0.11 mmol), EDCI (0.032 g, 0.165 mmol), HOBt (0.023 g, 0.165 mmol), DMF (1.5 ml), ethylamine (0.55 ml, 1.1 mmol), and diisopropylethylamine (0.06 ml, 0.33 mmol) gave the title compound (0.003 g, 6%) after column chromatography (silica gel, acetone/dichloromethane gradient).
- 10 MS(ES+) (m/z) 487.3 [M+1].

Example 606

5-[3-(Benzoylamino-methyl)cyclohexyl]-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinoline-9-carboxylic acid methyl amide

- 15 To a solution of Example 598 (0.05 g, 0.11 mmol) in toluene (1.5 ml) under N₂ was added thionyl chloride (0.5 ml) and heated at reflux for 3 h. The solution was concentrated using toluene to azeotrope. The residue was dissolved in dichloromethane (2 ml) under N₂ and added methylamine (0.55 ml, 1.1 mmol) and
- 20 DMAP (0.025 g, 0.22 mmol). The reaction was stirred overnight, diluted with dichloromethane, washed (0.1 N HCl, H₂O, and brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica gel, acetone/CH₂Cl₂ gradient) gave the title compound (0.029 g, 56%). MS(ES+) (m/z) 473.2 [M+1].

-285-

Example 607

N-{3-[9-(4,5-Dihydro-oxazol-2-yl)-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl]-cyclohexylmethyl}benzamide

5 To a solution of Example 598 (0.1 g, 0.22 mmol) in toluene (4 ml) under N₂ was added thionyl chloride (1 ml) and heated at reflux for 3 h. The solution was concentrated using toluene to azeotrope and the residue was dissolved in dichloromethane (4 ml) under N₂. Ethanolamine (0.131 ml, 2.2 mmol) and DMAP were added to the reaction and stirred overnight. The mixture was diluted with
10 dichloromethane, washed (5% NaHCO₃, H₂O, and brine), dried (MgSO₄), filtered, and concentrated. The residue was dissolved in thionyl chloride (2 ml), stirred overnight, and concentrated. The crude material was purified by column chromatography (silica gel, acetone/CH₂Cl₂ gradient) to give the title compound (0.01 g, 10%). MS(ES+) (m/z) 485.2 [M+1], 521.2 [M+HCl].

15

Example 608

N-[3-(9-Hydroxymethyl-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]benzamide

20 To a solution of Example 598 (0.01 g, 0.022 mmol) in toluene (0.4 ml) under N₂ was added thionyl chloride (0.1 ml) and heated at reflux for 3 h. The solution was concentrated using toluene to azeotrope and the residue was dissolved in dichloromethane (0.5 ml). To this solution was added NaBH₄ (0.008 g, 0.22 mmol) in THF (0.5 ml) and stirred overnight. The mixture was diluted with MeOH (1 ml)
25 and EtOAc, washed (0.1 N NaOH, H₂O, and brine), dried (MgSO₄), filtered, and concentrated to give the title compound (0.005 g, 53%) without further purification. MS(ES+) (m/z) 446.2 [M+1].

-286-

Example 609

N-[3-(3-Methyl-4-oxo-9-propionyl-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-
cyclohexylmethyl]benzamide

5 To a solution of Example 598 (0.05 g, 0.011 mmol) in toluene (2 ml) under N₂
was added thionyl chloride (1 ml) and heated at reflux for 3 h. The solution was
concentrated using toluene to azeotrope. To a solution of diethylzinc (1 ml, 1 mmol)
under N₂ stirring at -20 °C was added a solution of CuCN (0.093 g, 1.04 mmol) and
LiCl (0.086 g, 2.02 mmol) in THF (1.5 ml). This mixture was gradually allowed to
10 warm to 0 °C, stirred for 15 min at this temperature, and cooled to -25 °C. To this
solution was added the acid chloride residue in THF (0.5 ml) and stirred for 1.5 h
gradually allowing to warm to 0 °C. The reaction was diluted with NH₄Cl(aq) and
extracted with dichloromethane (2x). The combined organic layers were washed
(brine), dried (MgSO₄), filtered, and concentrated. Column chromatography (silica
15 gel, acetone/ dichloromethane gradient) gave the title compound (0.013 g, 28%).
MS(ES+) (m/z) 472.2 [M+1].

Example 610 & 611

5-[3-(2-Benzenesulfinyl-ethyl)-cyclohexyl]-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
20 c]quinolin-4-one &
5-[3-(2-Benzenesulfonyl-ethyl)-cyclohexyl]-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
c]quinolin-4-one

A compound from Example 501 (0.20 g, 0.00044 mol) was dissolved in
25 CH₂Cl₂ (30 mL) and the mixture cooled under a nitrogen atmosphere in an ice bath.
Then 3-chloroperoxy-benzoic acid (50%, 0.15 g, 0.00044 mol) was added and the
mixture stirred for 6 h while warming to ambient temperature. The mixture was
quenched with aq NaHCO₃ and extracted with CH₂Cl₂. The combined extracts were
dried over sodium sulfate and concentrated *in vacuo*. The resulting residue was
30 chromatographed over silica gel using CH₂Cl₂/THF as eluent which allowed for

-287-

isolation of both the pure sulfoxide (0.075 g, 36%), MS(ES): (M+1)⁺ 469.1, 471.2 m/z, and pure sulfone (0.025 g, 12%). MS(ES): (M+1)⁺ 485.4, 487.3 m/z.

Example 612

5 5-(3-Aminomethyl-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one (d,l)

To 41.5 mg (0.093 mmol) of Example 507 was added 0.75 ml trifluoroacetic acid. After 10 min stirring at rt, starting material (via TLC) was consumed. The
10 crude reaction mixture was diluted with EtOAc and sat'd aq. bicarbonate and transferred to a sep funnel. The aqueous phase was reextracted two add'l times with EtOAc, and the combined organics were washed with brine, dried over Na₂SO₄, filtered and concentrated to afford clean amine (28 mg, 87%) which was used without further purification. MS (+ES) m/z 345.9 (M+H)⁺.

15

Example 613

(1R,3S)-5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

20 The product from Example 562 (0.069 g, 0.16 mmol) was treated with TFA (2 mL) and stirred at room temperature for 15 min. The reaction was added dropwise to sat HCO₃⁻ (10 mL) and the product was extracted with CH₂Cl₂ (5 x 10 mL). The solvent was removed *in vacuo* to afford 0.05 g (94%) of the title compound as a white solid. MS (ES⁺) m/z 346.0 (M+H)⁺.

25

Example 614

(1S,3R)-5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

30 The product of Example 553 (0.08 g, 0.18 mmol) was treated with TFA (2 mL) and stirred at room temperature for 20 min. The reaction was added dropwise to

-288-

sat HCO_3^- (10 mL) and the product was extracted with CH_2Cl_2 (5 x 10 mL). The solvent was removed *in vacuo* to afford 0.047 g (76%) of title compound as a white solid. MS (ES+) m/z 346.0 (M+H)⁺.

5

Example 615

(1R,3S)-5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

A solution of the product from Example 508 (0.40 g, 0.83 mmol) in dry
10 CH_2Cl_2 (5 mL) was treated with iodotrimethylsilane (0.25 g, 1.25 mmol) and stirred at r.t. After 3 hr the reaction was quenched with MeOH (2 mL) and stirred for an additional 30 min. The solution was then concentrated to an orange solid and taken up in EtOAc (20 mL) and 1 N HCl (20 mL) and transferred to a sep funnel. The organic layer was extracted and the aqueous layer was washed with additional EtOAc
15 (2 x 10 mL). The pH of the aqueous layer was then adjusted to ph~12 with 5 M NaOH. The product was extracted with EtOAc (3 x 25 mL). The EtOAc extractions were dried over sodium sulfate, filtered, and the solvent removed *in vacuo* to afford 0.240 g (84%) as a white solid which was used without further purification. MS (ES+) m/z 346.0 (M+H)⁺.

20

Example 616

5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

The product from Example 560 (30 mg, 0.067 mmol) was treated with HCl-
25 1.0 M in HOAc (2 mL). The reaction was stirred at r.t. for 20 minutes. Acetonitrile (2 mL) was added and the solution was concentrated to a white solid. The white solid was suspended in ether and filtered to afford 15 mg (70 %) as a white solid. MS (ES+) m/z 346.0 (M+H)⁺.

-289-

Example 617

5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,5-c]quinolin-4-one

The product of Example 561 (30 mg, 0.067 mmol) was treated with HCl-1.0 M in HOAc (2 mL). The reaction was stirred at r.t. for 20 minutes. Acetonitrile (2 mL) was added and the solution was concentrated to a white solid. The white solid was suspended in ether and filtered to afford 13 mg (60 %) as a white solid. MS (ES+) m/z 346.0 (M+H)⁺.

10

Example 618

3-[2-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)cyclohexyl]-N-(3,4,5-trimethoxyphenyl)propionamide

The amide from preparation 165 (24 mg, 0.042 mmol) was dissolved in anhydrous DMF (0.5 ml). NaHMDS (1 M in THF, 0.046 ml, 1.1 eq) was added dropwise. After 1hr, sm still present, and additional base (1.1 eq) was added, followed by a third volume (1.1 eq) an hour later. The reaction was then allowed to proceed o.n. The reaction was then quenched with 0.1 N HCl and EtOAc, transferred to a sep funnel, and the aqueous phase was reextracted two additional times. the combined organics were washed with sat'd bicarbonate, then with brine, dried over Na₂SO₄, filtered and concentrated to afford 21 mg of crude product. Purification over silica (1 g Bond-Elut cartridge, 2:1 EtOAc/hexanes) afforded 21 mg of product (40%). MS (+ES) 553.9, 555.9 (M+H)⁺.

25

Example 619

5-(3-Aminomethylcyclopentyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

A mixture of Example 518 (240 mg, 0.5 mmol) and trimethyl silyl iodide (154, 077 mmol) dissolved in methylene chloride (10 mL) was stirred overnight at rt. The reaction mixture was passed through Bond Elut SCX column (20cc, 5g), eluted

30

-290-

with ammonia solution in methanol (1M, 20 mL) and evaporated to obtain the title compound (150 mg, 90%). ESMS: 332 (M+1)⁺.

Example 620

5 5-(((1R,3S)-3-Aminocyclopentyl)methyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
c]quinolin-4-one

N-((1S,3R)-3-((9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinolin-5-yl)methyl)cyclopentyl)(tert-butoxy)carboxamide was converted to the amine by TFA
10 in DCM at rt. for three hours. ESMS: 332 (M+1)⁺.

Example 621

5-(((1S,3R)-3-Aminocyclopentyl)methyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-
c]quinolin-4-one

15

N-((1R,3S)-3-((9-chloro-3-methyl-4-oxo-5*H*-isoxazolo-[4,3-
c]quinolin-5-yl)methyl)cyclopentyl)(tert-butoxy)carboxamide was converted to the amine as
described in Example 620 . ESMS: 332 (M+1)⁺.

20

Example 622

cis-1-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinolin-5-yl)-3-(3-phenyluriedo)-cyclohexane

25

To the suspension of *cis*-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-
c]quinolin-5-yl)-3-amino-cyclohexane (0.0446 g, 0.121 mmol) in 1 mL anhydrous
THF at RT was added 2N NaOH (0.60 mL, 0.121 mmol). After 5 min. the
phenylisocyanate (0.013 mL, 0.121 mmol). After 20 min. the reaction was diluted
with EtOAc, washed with water and concentrated to near dryness. After sonication
and filtration a white solid (0.0412, 75%) was obtained. ESIMS *m/e* 451 ³⁵Cl (M⁺+1)
30 and 453 ³⁷Cl (M⁺+1).

-291-

Example 623

cis-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-[3-(phenylmethyl)-uriedo]cyclohexane

5 A procedure similar to that for Example 622 was used to prepare the title compound as a yellow solid (0.110 g, 87%). ESIMS *m/e* 465 ³⁵Cl (M⁺+1) and 467 ³⁷Cl (M⁺+1).

Example 624

10 *cis*-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(3-cyclohexyluriedo) cyclohexane

A procedure similar to that for Example 622 was used to prepare the title compound as a yellow solid (0.106 g, 86%). ESIMS *m/e* 457 ³⁵Cl (M⁺+1) and 459 ³⁷Cl (M⁺+1).

15

Example 625

Preparation of racemic *cis*-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(3-ethyluriedo) cyclohexane

20

A procedure similar to that for Example 622 was used to prepare the title compound as a yellow solid (0.102 g, 94%). ESIMS *m/e* 403 ³⁵Cl (M⁺+1) and 405 ³⁷Cl (M⁺+1).

25

Example 626

cis-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(3-propyluriedo)cyclohexane

A procedure similar to that for Example 622 was used to prepare the title compound as a yellow solid (0.106 g, 94%). ESIMS *m/e* 417 ³⁵Cl (M⁺+1) and 419 ³⁷Cl (M⁺+1).

30

-292-

Example 627

(1*S*,3*R*)-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(((2*S*)-2-amino-2-phenylacetyl)amino]cyclohexane hydrochloride

5 To the product from Example 11 (0.015 g, 0.027 mmol) in 1 mL CH₂Cl₂ was added 1 mL 4M HCl in dioxane. After 3 h the mixture was stripped to dryness to give the title compound as a white solid (0.0131 g, 98%). ESIMS *m/e* 465 ³⁵Cl (M⁺+1) and 467 ³⁷Cl (M⁺+1).

10

Example 628

(1*R*,3*S*)-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(((2*S*)-2-amino-2-phenylacetyl)amino]cyclohexane hydrochloride

To the product from Example 11 (0.013 g, 0.023 mmol) in 1 mL CH₂Cl₂ was
15 added 1 mL 4M HCl in dioxane. After 3 h the mixture was stripped to dryness to give a white solid (0.0115 g, 100%). ESIMS *m/e* 465 ³⁵Cl (M⁺+1) and 467 ³⁷Cl (M⁺+1).

Example 629

(1*S*,3*R*)-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(((2*R*)-2-
20 amino-2-phenylacetyl) amino]cyclohexane hydrochloride

To the product from Example 11 (0.013 g, 0.019 mmol) in 1 mL CH₂Cl₂ was added 1 mL 4M HCl in dioxane. After 3 h the mixture was stripped to dryness to give a white solid (0.0098 g, 100%). ESIMS *m/e* 465 ³⁵Cl (M⁺+1) and 467 ³⁷Cl (M⁺+1).

25

-293-

Examples 630 and 631

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-(5-methyltetrazol-2-yl)-2-phenylacetamide &

5 N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-(5-methyltetrazol-1-yl)-2-phenylacetamide

To a 0°C solution of N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-2-hydroxy-2-phenyl (30 mg, 0.06 mmol) in toluene (4 mL) was added triphenylphosphine (20 mg, 0.08 mmol), 5-methyltetrazole (8 mg, 0.10 mmol), and diethylazodicarboxylate (16 µL, 0.10 mmol) dropwise. After 30 minutes the reaction was allowed to warm to room temperature overnight.

Purification gave 4.4 mg and 10.2 mg, respectively as white solids, 43% combined yield. ¹H NMR: consistent with structures. MS (ion spray) 532 (M⁺).

15

Example 632

5-(3-Aminomethylcyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one hydroiodide

To a 0°C solution of [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]-carbamic acid be (5.0 g, 10.4 mmol) in dichloromethane (100 mL) was added iodotrimethylsilane (3.6 mL, 25.0 mmol) dropwise, and the solution allowed to warm to room temperature overnight. Methanol (6.1 mL, 150.0 mmol) was added dropwise over two minutes, and the reaction stirred for 30 minutes. The reaction was concentrated and suspended in ethyl ether. After 15 minutes of sonication, the solids were removed by filtration and washed with ethyl ether. The tan powder was dried on a vacuum pump to give 4.61 g as the 1.2 HI salt, 89% yield. ¹H NMR: consistent with structure. MS (ion spray) 346 (M⁺).

-294-

Example 633

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-phenylacetamide

5 A compound from Example 235 (180 mg, 0.32 mmol) was combined with trifluoroacetic acid (0.75 mL) in dichloromethane (4.0 mL) and the mixture stirred at ambient temperature until deprotection was complete. The mixture was then concentrated *in vacuo* and the residue treated with aqueous NaHCO₃ and extracted with ethyl acetate. Concentration left a residue, which was chromatographed over
10 silica (methanol/dichloro-methane) allowing for isolation of the desired product (145 mg, 97%) as a white solid. MS(ES): (M+1)⁺ 465.2, 467.2 m/z.

Example 634

15 5-(3-Aminocyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one hydroiodide

To a solution of [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]carbamic acid methyl ester, 2.0 g (5.1 mmol) in 50 mL of dichloromethane was added 1.74 mL (12.2 mmol) of trimethylsilyliodide. The
20 reaction mixture was stirred overnight at ambient temperature and was quenched dropwise with 3.0 mL (73.4 mmol) of methanol. The mixture was stirred 30 minutes at ambient temperature and was concentrated to dryness. The residue was triturated with ether, filtered and dried to provide a quantitative yield of the desired isomer as a tan solid. ¹H-NMR is consistent with structure. MS (ion spray) 332.1 (M+ for free
25 base).

Example 635

30 Pyridazine-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]amide

4-Methylpyridazine, 4.0 mL (44.6 mmol) was dissolved in 200 mL of water and 16.2 g (103 mmol) of potassium permanganate was added. The resulting slurry

-295-

was refluxed for 30 min, then an additional 8.5 g (54 mmol) of potassium permanganate was added. The resulting slurry was refluxed for 24 h, then filtered hot through celite. The resulting solution was washed with ether, acidified to pH = 7 with 5 N HCl, and concentrated to a volume of 80 mL. This solution was acidified to pH = 2 with 1 N HCl and extracted with ethyl acetate and with 20% isopropanol/chloroform. The combined organics were dried over sodium sulfate, filtered and concentrated to dryness to yield 320 mg of a tan foam. ¹H NMR indicated a 2:1 mixture of 4-pyridazine carboxylic acid: starting material 4-methylpyridazine.

10 This mixture, 40 mg, was added to a solution of 66 mg (0.14 mmol) of a compound from Example 632 in 5 mL of N,N-dimethylformamide. To this solution was added 23 mg (0.17 mmol) of 1-hydroxy-7-azabenzotriazole, 33 mg (0.17 mmol) of 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride, 5 mg of 4-dimethylaminopyridine and 60 μL (0.42 mmol) of triethylamine. Yield=33 mg (53%)

15 of the desired isomer as a white foam. ¹H-NMR is consistent with structure. MS (ion spray) 452.2 (M+).

Example 636

1,2,3,4-Tetrahydroisoquinoline-3-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]amide hydrochloride

20

A solution of 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester, 74 mg (0.125 mmol) in 5 mL of acetic acid saturated with HCl gas was stirred for three hours at ambient temperature, then was concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness to give a quantitative yield of the desired isomer as a white solid. ¹H-NMR is consistent with structure. MS (ion spray) 491.0 (M+).

25

-296-

Example 637

1,2,3,4-Tetrahydro-isoquinoline-3-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide hydrochloride

5 A solution of 3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester, 74 mg (0.125 mmol) in 5 mL of acetic acid saturated with HCl gas was stirred for four hours at ambient temperature, then was concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness to give a quantitative yield of the
10 desired isomer as a white solid. ¹H-NMR is consistent with structure. MS (ion spray) 491.3 (M+).

Example 638

15 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide

{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenyl-methyl}-carbamic acid *tert*-butyl ester (1.24 g; 2.2 mmol) was dissolved in excess, neat acetic acid saturated with HCl_(g) (20 mL). After
20 stirring 30 min at room temperature, the solution was concentrated to dryness by rotary evaporation. Acetic acid was removed from the resulting solid by consecutive dissolution in, and drying from, acetonitrile (thrice) and then diethyl ether (once). The solid was dissolved in 25% (v/v) isopropyl alcohol in chloroform, washed with saturated NaHCO_{3(aq)}, and dried by rotary evaporation to yield the desired product in
25 97% isolated yield.

MS(ES) calc'd: [M+H]⁺ = 465.2 m/z. Found: 465.2 m/z.

-297-

Example 639

2-Amino-*N*-{[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylcarbamoyl]-phenylmethyl}-2-methylpropionamide hydrochloride

5 A compound from Example 321 was deprotected in a manner similar to Example 638 and kept as the hydrochloride salt. MS(ES) calc'd: $[M+H]^+ = 550.2$ m/z; $[M-H]^- = 548.2$ m/z; $[M+Cl]^- = 584.2$ m/z. Found: 550.0 m/z; 548.0 m/z; 584.0 m/z.

10 Example 640

2-Amino-*N*-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-pyridin-3-ylacetamide

15 A compound from Example 326 was deprotected in a manner similar to Example 638. The free base was separated into individual diastereomers by radial chromatography on a 2 mm thick silica gel rotor with a 2% methanol/dichloromethane (v/v) mobile phase (300 mL) followed by increasing methanol to 10% in step gradients (each mobile phase now containing 1% triethylamine). The desired product was isolated as 13 mg of isomer 1, 18 mg of isomer 2, and 7 mg mixed product.

20 Isomer 1 MS(ES) exact mass calc'd: $[M+H]^+ = 466.1646$ m/z. Found: 466.1648 m/z.
Isomer 2 MS(ES) exact mass calc'd: $[M+H]^+ = 466.1646$ m/z. Found: 466.1663 m/z.

Example 641

25 *N*-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,5-*c*]quinolin-5-yl)cyclohexylmethyl]-thiobenzamide

A solution of *cis-N*-{[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,5-*c*]quinolin-5-yl)cyclohexyl]methyl}benzamide (0.1 g, 0.2 mmol) in pyridine (10 mL) was treated with P_2S_5 (0.37 g, 1.7 mmol) and heated to reflux. The reaction was

30 stirred for 1 hour at reflux and then allowed to cool to r.t. The reaction was quenched with water (150 mL) A white precipitate formed and was filtered to afford a light

-298-

yellow solid. This solid was taken up in CH₂Cl₂ (15 mL) and purified by silica gel column chromatography. The product was eluted with 1% MeOH in CH₂Cl₂. The solvent was removed to afford 0.082 g (81%) of a yellow solid. MS m/z (ES+) 465.8 (M+H)⁺, (ES-) 463.8 (M-H)⁻.

5

Example 642

N-[3-(9-Chloro-3-methyl-4-oxo-2,4-dihydropyrazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]benzamide

10 Cis-N-({3-[3-(aminoethylidene)-5-chloro-2,4-dioxohydroquinolyl]-cyclohexyl}methyl)benzamide (0.15 g, 0.3 mmol) and hydrazine hydrate-85% (0.016 g, 0.49 mmol) in EtOH (20 mL). The reaction was heated to reflux and stirred for 6 hr. The reaction was then concentrated to a solid and taken up in CH₂Cl₂. This solution was purified by silica gel column chromatography using 50% EtOAc in
15 CH₂Cl₂ to elute the product. The solvent was removed to afford 0.095 g (64%) as a yellow solid. MS (ES+) m/z 448.9 (M+H)⁺, (ES-) m/z 446.9 (M-H)⁻, 506.9 (M+CH₃COO)⁻.

Example 643

20 N-[3-(9-Chloro-3-methyl-4-oxo-2,4-dihydropyrazolo[4,3-c]quinolin-5-yl)-cyclohexylmethyl]-6-fluoronicotinamide

cis-N-({3-[3-(Aminoethylidene)-5-chloro-2,4-dioxohydroquinolyl]-cyclohexyl}methyl)(6-fluoro(3-pyridyl))carboxamide (0.92 g, 0.2 mmol) and
25 hydrazine hydrate-85% (0.063 g, 0.3 mmol) in EtOH (20 mL). The reaction was heated to reflux and stirred for 1 hr. The reaction was then concentrated to a solid and taken up in CHCl₃. This solution was purified by silica gel column chromatography using 5% MeOH in CHCl₃ to elute the product. The solvent was removed to afford
0.02 g (22%) as a white solid. MS (ES+) m/z 468.1 (M+H)⁺, (ES-) m/z 466.1 (M-H)⁻,
30 526.2 (M+CH₃COO)⁻.

-299-

Example 644

[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,5-*c*]quinolin-5-yl)cyclohexylmethyl]-
carbamic acid benzyl ester

5

A solution of N-{[3-(3-acetyl-4-amino-5-chloro-2-oxohydroquinolyl)-
cyclohexyl]-methyl}(phenylmethoxy)carboxamide (0.02 g, 0.04 mmol) in acetic acid
(2 mL) was treated with hydroxylamine hydrochloride (3 mg, 0.046 mmol). The
solution was heated to reflux and stirred 4 hr. The reaction was then diluted in
10 CH₂Cl₂ (50 mL) and washed with 5M NaOH (3 x 10 mL) and brine (2 x 10 mL).
The organic was dried over sodium sulfate and the solvent removed. The crude
product was purified by silica gel column chromatography using 10% EtOAc in
CH₂Cl₂ to elute the product. The solvent was removed *in vacuo* to afford 0.014 g
(70%) of the title compound as an off-white solid.
15 MS (ES+) m/z 500.1 (M+H)⁺.

Example 645

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,5-*c*]quinolin-5-
yl)cyclohexylmethyl]-6-fluoronicotinamide

20

A solution of N-{[3-(3-acetyl-4-amino-5-chloro-2-
oxohydroquinolyl)cyclohexyl]-methyl}(6-fluoro(3-pyridyl))carboxamide (0.035 g,
0.07 mmol) in acetic acid (5 mL) was treated with hydroxylamine hydrochloride (7.8
mg, 0.11 mmol). The solution was heated to reflux and stirred 3 hr. The reaction was
25 then diluted in CHCl₃ (50 mL) and washed with sat'd sodium bicarbonate (3 x 10
mL) and brine (2 x 10 mL). The organic was dried over sodium sulfate and the
solvent removed. The crude product was purified by silica gel column
chromatography using 50% EtOAc in CHCl₃ to elute the product. The solvent was
removed *in vacuo* to afford 0.025 g (72%) of the title compound as a white solid. MS
30 m/z (ES+) 468.8 (M+H)⁺, (ES-) 466.8 (M-H)⁻, 526.8 (M+CH₃COO)⁻.

-300-

Example 646

N-[3-(9-Cyano-3-methyl-4-oxo-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)-
cyclohexylmethyl]-4-fluoro-benzamide

5 To a solution of a compound from Example 524 (0.106 g, 0.23 mmol) in
CH₂Cl₂ (10 ml) under N₂ was added TMSI (0.36 ml, 2.52 mmol) and stirred
overnight. Added MeOH (5 ml) and stirred for 1 h to quench the reaction. The
mixture was concentrated, triturated with toluene and dried to obtain a crude solid.
This was mixed with 4-fluorobenoyl chloride (0.093 ml, 0.8 mmol) in CH₂Cl₂ (5
10 ml) under N₂ and triethylamine (0.22 ml, 1.6 mmol) was added dropwise. After
stirring overnight, the mixture was diluted with CH₂Cl₂, washed (0.1N HCl then
brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel,
acetone/CH₂Cl₂ gradient) gave the title compound (0.085 g, 88%). Mass Spectrum
(ES+) (m/z) 459.1 [M+1].

15

Example 647

N-[3-(9-Cyano-3-methyl-4-oxo-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-
3,4-difluorobenzamide

20 In a fashion similar to that described for Example 527, a compound from
Example 524 (0.106 g, 0.23 mmol), CH₂Cl₂ (10 ml), TMSI (0.36 ml, 2.52 mmol),
3,4-difluorobenzoyl chloride (0.093 ml, 0.8 mmol), CH₂Cl₂ (5 ml), and triethylamine
(0.22 ml, 1.6 mmol) gave the title compound (0.087 g, 87%) after flash
chromatography (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z)
25 477.1 [M+1].

-301-

Example 648

N-[3-(9-Chloro-4-oxo-3-phenylsulfanyl-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)-
cyclohexylmethyl]benzamide

5 To a solution of *N*-{[3-(3-acetyl-4-amino-5-chloro-2-oxohydroquinolyl)-
cyclohexyl]methyl}(6-fluoro(3-pyridyl))carboxamide (0.1 g, 0.2 mmol) in DMF (2.5
ml) under N₂ was added sodium thiophenoxide (0.132 g, 1.0 mmol) and stirred for 2
h. The reaction was diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄),
filtered, and concentrated. The residue was dissolved in DMF (2 ml) under N₂,
10 cooled to 0 degrees C, and KHMDS (0.566 ml, 0.28 mmol) was added over 15 min.
After 30 min, the solution was diluted with EtOAc, washed (H₂O and brine), dried
(MgSO₄), filtered, and concentrated. Flash chromatography (silica gel,
Acetone/CH₂Cl₂ gradient) gave the title compound (0.045 g, 41%). Mass Spectrum
(ES+) (m/z) 544.2 [M+1].

15

Example 649

N-[3-(9-Chloro-3-ethoxy-4-oxo-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-
benzamide

20 To a solution of 278 (0.1 g, 0.2 mmol) in DMF (2.5 ml) under N₂ was added
sodium ethoxide (0.33 ml at 21% wt., 1.0 mmol) and stirred for 2 h. The reaction was
diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and
concentrated. Purification by flash chromatography gave the title compound (0.025 g,
26%) (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 480.2
25 [M+1].

-302-

Example 650

N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-methylamino-acetamide hydrochloride

5 A solution of the compound from Example 333, 66 mg (0.13 mmol) in 10 mL of HCl-saturated acetic acid was stirred 4 hours at rt., then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness to give a quantitative yield of the desired isomer as a white foam. ¹H-NMR is consistent with structure. MS (ion spray) 403.2 (M+).

10

Example 651

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-methyl-propionamide hydrochloride

15 A solution of the compound from Example 335, 64 mg (0.12 mmol) in 10 mL of HCl-saturated acetic acid was stirred 3 hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to give a quantitative yield of the desired isomer as a white foam. MS (ion spray) 417.1 (M+).

20

Example 652

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-acetamide hydrochloride

25 A solution of the compound from Example 336, 49 mg (0.10 mmol) in 10 mL of HCl-saturated acetic acid was stirred 3 hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to give a quantitative yield of the desired isomer as a tan foam. MS (ion spray) 389.1 (M+).

30

-303-

Example 653

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-phenyl-2-piperazin-1-ylacetamide dihydrochloride

5 A solution of the compound from Example 339, 171 mg (0.27 mmol) in 10 mL of HCl-saturated acetic acid was stirred 2 hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to give a quantitative yield of the desired mixture of isomers as a tan foam. ¹H-NMR is
10 consistent with structure. MS (ion spray) 389.1 (M⁺). The isomers were separated by chiral chromatography in a similar fashion to that described for Example 554 to yield 31.5 mg (19%) of isomer 1 as a white foam and 32.9 mg (20%) of isomer 2 as a white foam. I.S. (534.2) M⁺.

15

Example 654

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-methylamino-2-phenylacetamide hydrochloride

 A solution of the compound from Example 340, 130 mg (0.22 mmol) in 10
20 mL of HCl-saturated acetic acid was stirred 2 hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to give a quantitative yield of the desired mixture of isomers as a white foam. MS (ion spray) 479.2 (M⁺).

25

Example 655

1-Aminocyclohexanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]amide hydrochloride

30

 A solution of the compound from preparation 229a, 198 mg (0.35 mmol) in 15 mL of HCl-saturated acetic acid was stirred three hours at rt. then concentrated to dryness. The residue was slurried 3x in acetonitrile and concentrated to dryness to

-304-

give 160 mg (93%) of the desired isomer as a white solid. MS (ion spray) 457.2 (M+).

Example 656

5 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-cyclohexylacetamide hydrochloride

A solution of the compound from Example 355, 80 mg (0.14 mmol) in 30 mL of HCl-saturated acetic acid was stirred four hours at rt. then concentrated to dryness.
10 The residue was slurried 3x in toluene and concentrated to dryness to give 60 mg (85%) of the desired isomer as a tan solid. MS (ion spray) 471.2 (M+).

Example 657

15 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-cyclohexylacetamide hydrochloride

A solution of the compound from Example 356, 100 mg (0.17 mmol) in 30 mL of HCl-saturated acetic acid was stirred four hours at rt. then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness to give
20 47 mg (55%) of the desired isomer as a tan solid. ¹H-NMR is consistent with structure. MS (ion spray) 471.2 (M+).

Example 658

25 2-Aminoindan-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]amide hydrochloride

A solution of a compound from Example 634, 220 mg (0.37 mmol) in 40 mL of HCl-saturated acetic acid was stirred three hours at rt., then concentrated to dryness. The residue was slurried 3x in toluene and concentrated to dryness give a
30 quantitative yield of the desired isomer as a white solid. MS (ion spray) 491.2 (M+).

-305-

Example 659

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-3-phenylpropionamide

5 A solution of the compound from Example 370, 84 mg (0.14 mmol) in 10 mL of HCl-saturated acetic acid was stirred four hours at rt., then concentrated to dryness. The residue was slurried 3x in acetonitrile and concentrated to dryness give a quantitative yield of the desired isomer as a tan foam. MS (ion spray) 479.1 (M+).

10 Example 660

5-(3-Amino-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

A compound from Example 634 (350 mg, 0.95 mmol) was treated with excess aqueous NaHCO₃ and the mixture extracted with EtOAc. The combined extracts
15 were dried over Na₂SO₄ and concentrated *in vacuo* to yield 309 mg (98%) of the title compound.

Example 661

N-[3-(9-Cyano-3-methyl-4-oxo-5H-isoxazole[4,3-c]quinolin-5-yl)cyclohexylmethyl]-
4-fluorobenzamide

20 To a solution of a compound from Example 532 (0.106 g, 0.23 mmol) in CH₂Cl₂ (2 mL) under N₂ was added TMSI (0.046 mL, 0.32 mmol) and stirred overnight. Added MeOH (1 mL) and stirred for 1 h to quench the reaction. The mixture was concentrated, triturated with toluene and dried to obtain a crude solid.
25 This was mixed with 4-fluorobenoyl chloride (0.093 mL, 0.8 mmol) in CH₂Cl₂ (5 mL) under N₂ and Et₃N (0.22 mL, 1.6 mmol) was added dropwise. After stirring overnight, the mixture was diluted with CH₂Cl₂, washed (0.1N HCl then brine), dried (MgSO₄), filtered, and concentrated. Flash chromatography (silica gel, Acetone/CH₂Cl₂ gradient) gave the title compound (0.085 g, 88%). Mass Spectrum
30 (ES+) (m/z) 459.1 [M+1].

-306-

Example 662

N-[3-(9-Cyano-3-methyl-4-oxo-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-
3,4-difluorobenzamide

5

In a fashion similar to that described for Example 542, a compound from Example 413 (0.1 g, 0.21 mmol), CH₂Cl₂ (2 mL), TMSI (0.046 mL, 0.32 mmol), 3,4-difluorobenoyl chloride (0.093 mL, 0.8 mmol), CH₂Cl₂ (5 mL), and Et₃N (0.22 mL, 1.6 mmol) gave the title compound (0.087 g, 87%) after flash chromatography (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 477.1 [M+1].

10

Example 663

N-[3-(9-Chloro-3-ethoxy-4-oxo-5*H*-isoxazole[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-
benzamide

15

To a solution of a compound from preparation 328 (0.1 g, 0.2 mmol) in DMF (2.5 mL) under N₂ was added sodium ethoxide (0.33 mL at 21% wt., 1.0 mmol) and stirred for 2 h. The reaction was diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and concentrated. Purification by flash chromatography gave the title compound (0.025 g, 26%) (silica gel, acetone/CH₂Cl₂ gradient). Mass Spectrum (ES+) (m/z) 480.2 [M+1].

20

Example 664

N-[3-(9-Cyano-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-
6-fluoronicotinamide

25

To a solution of *N*-{[3-(9-cyano-3-methyl-4-oxo(5-hydroisoxazolo[4,3-*c*]quinolin-5-yl))cyclohexyl]methyl}(phenylmethoxy)carboxamide (0.27 g, 0.58 mmol) in CH₂Cl₂ (7 mL) under N₂ was added TMSI (0.2 mL, 1.4 mmol) and stirred overnight. Added MeOH (3 mL) and stirred for 1 h to quench the reaction. The

30

-308-

Example 667

N-[3-(9-Amino-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexylmethyl]-benzamide

5 To a solution of a compound from preparation 333 (0.24 g, 0.42 mmol) in THF (4.2 mL) under N₂ was added TBAF (0.84 mL, 0.84 mmol, 1.0M in THF) dropwise and stirred for 4 h. The reaction was applied directly to a silica gel column and eluted with EtOAc which gave the title compound (0.18 g, 99%). Mass Spectrum (ES-) (m/z) 429.3 [M-1]

10

Examples 668 & 669

9-Chloro-5-[3-(2-hydroxy-2-phenylethylamino)cyclohexyl]-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one &

9-Chloro-5-[3-(2-hydroxy-1-phenylethylamino)cyclohexyl]-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one

15

To a stirred solution of a compound from Example 634 (440mg, 1.33 mmol) in abs. EtOH (60 mL) was added *S*-styrene oxide (159.2 mg, 1.33 mmole) under N₂ atmosphere. The reaction mixture was refluxed for 22 h. The solvent was removed on *Buchi* and the crude was chromatographed by Elution solution systems (40L column, NH₃ in MeOH: EtOAc, gradient). The title compounds were yielded (A, 201 mg; B, 41 mg, 40.3% yield). Mass Spectrum(FIA); A(m/z) 452.2 (M+1); B(m/z) 452.2 (M+1).

20

-309-

Examples 670 & 671

9-Chloro-5-[3-(2-hydroxy-2-phenylethylamino)cyclohexyl]-3-methyl-5H-
isoxazolo[4,3-c]quinolin-4-one &

5 9-Chloro-5-[3-(2-hydroxy-1-phenylethylamino)cyclohexyl]-3-methyl-5H-
isoxazolo[4,3-c]quinolin-4-one

To a stirred solution of a compound from Example 634 (441mg, 1.33 mmole) in abs. EtOH (60 mL) was added *R*-styrene oxide (160 mg, 1.33 mmole) under N₂ atmosphere. The reaction mixture was refluxed for 202 h. The solvent was removed
10 on *Buchi* and the crude was chromatographed by Elution solution systems (40L column, NH₃ in MeOH: EtOAc, gradient). The title compounds were yielded (A, 161 mg; B, 40 mg, 33.4 % yield). Mass Spectrum (FIA) A(m/z) 452.2 (M+1); B(m/z) 452.2 (M+1).

15

Example 672

9-Chloro-3-methyl-5-[3-(2-oxo-5-phenyloxazolidin-3-yl)cyclohexyl]-5H-
isoxazolo[4,3-c]quinolin-4-one

To a stirred solution of a compound from Example 668 (41mg, 0.091 mmole) in THF (2mL) was added DCC (15 mg, 0.091 mmole) and Et₃N (0.013 mL, 0.091 mmole). The reaction mixture was stirred at rt. under N₂ atmosphere for 18 h. It was diluted with EtOAc, washed with brine, dried over Na₂SO₄ and concentrated. The resulted crude oil was chromatographed (gradient, hexanes: EtOAc) and the desired product was yielded as white foam solid (23g, 54 %). Mass Spectrum: m/z calcd.
25 478.1533, Found 478.1544

-310-

Example 673

9-Chloro-3-methyl-5-[3-(2-oxo-5-phenyloxazolidin-3-yl)cyclohexyl]-5H-
isoxazolo[4,3-c]quinolin-4-one

5 To a stirred solution of a compound from Example 670 (50mg, 0.11 mmol) in
THF (2mL) was added DCC (18 mg, 0.11 mmol), Et₃N (0.016 mL, 0.11 mmol) and
DMAP (catalytic amount). The reaction mixture was stirred at rt. under N₂
atmosphere for 48 h. It was diluted with EtOAc, washed with brine, dried over
Na₂SO₄ and concentrated. The resulted crude oil was chromatographed (gradient,
10 hexanes: EtOAc) and the desired product was yielded as white foam solid (21g, 40
%). Mass Spectrum: m/z calcd. 478.1533, Found 478.1565.

Example 674

15 9-Chloro-3-methyl-5-[3-(2-oxo-4-phenyloxazolidin-3-yl)cyclohexyl]-5H-
isoxazolo[4,3-c]quinolin-4-one

To a stirred solution of a compound from Example 669 (97mg, 0.21 mmole) in
THF (4mL) was added DCC (34.1 mg, 0.21 mmole), Et₃N (0.03 mL, 0.21 mmole)
and DMAP (catalytic amount). The reaction mixture was stirred at rt. under N₂
20 atmosphere for 18 h. It was diluted with EtOAc, washed with brine, dried over
Na₂SO₄ and concentrated. The resulted crude oil was chromatographed (gradient,
hexanes: EtOAc) and the desired product was yielded as white foam solid (40g, 40
%). Mass Spectrum(FIA)(m/z) 478.1 (M+1).

25

Example 675

9-Chloro-3-methyl-5-[3-(2-oxo-4-phenyloxazolidin-3-yl)cyclohexyl]-5H-
isoxazolo[4,3-c]quinolin-4-one

To a stirred solution of a compound from Example 671 (64mg, 0.14 mmole) in
30 CH₂Cl₂ (5mL) was added DCC (115 mg, 0.7 mmole), Et₃N (0.02 mL, 0.14 mmole)

-311-

and DMAP (catalytic amount). The reaction mixture was stirred at rt. under N₂ atmosphere for 48 h. It was diluted with CH₂Cl₂, washed with brine, dried over Na₂SO₄ and concentrated. The resulted crude oil was chromatographed (gradient, hexanes: EtOAc) and the desired product was yielded as white foam solid (51g, 76
5 %). Mass Spectrum (FIA) (m/z) 478.0 (M+1).

Example 676

9-Chloro-5-{3-[2-(6-chloropyridin-3-yl)-2-hydroxyethylamino]cyclohexyl}-3-methyl-
5*H*-isoxazolo[4,3-*c*]quinolin-4-one

10

To a stirred solution of 6-chloronicotinic acid (900mg, 5.7 mmol) in THF (7 mL) was added BH₃ (16 mL, 17.1 mmol, 1M in THF) under N₂ atmosphere. The reaction mixture was stirred for 5 h. It was quenched with methanol (5 mL) and then concentrated. The crude was dissolved in EtOAc (30 mL), washed with 1N NaOH
15 (15mLx3), brine, dried and concentrated. Yield: 615 mg (75%) of 6-chloro-pyridin-3-yl-methanol. To this alcohol (615 mg, 4.28 mmol) solution in CH₂Cl₂ (8 mL) was added Dess-Martin reagent (5.4 mg, 12.8 mmol). It was stirred at rt. for 2 h and then diluted with CH₂Cl₂. The resulting solution was washed with NaOH (1N, 2x5mL),
brine, dried and concentrated. 516 mg (85%) of yellow solid as the desired 6-chloro-
20 pyridine-3-carbaldehyde. To a solution of above aldehyde (401 mg, 2.82 mmol) in CH₂Cl₂ (10 mL) was added 50% NaOH solution (9.3 mL), trimethylsulfonium iodide (1.15 g, 5.64 mmol) and tetrabutylammonium iodide (11mg). It was refluxed for 48 h and poured onto ice water. The organic layer was extracted with CH₂Cl₂, washed
with brine, dried and concentrated. The crude epoxide was chromatographed
25 (gradient, hexanes: EtOAc) and 190 mg (43%) yellow oil was yielded as desired 2-Chloro-5-oxiranyl-pyridine. To the solution of 2-chloro-5-oxiranyl-pyridine (118.4 mg, 0.76 mmol) in absolute ethanol (40 mL) was added a compound from Example 660 (252 mg, 0.76 mmol). The reaction mixture was refluxed for 24 h and
concentrated.

30

-312-

Example 677

[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cycloheptylmethyl]-
carbamic acid 2-trimethylsilanylethyl ester

5 To a warm solution of 0.2g (0.52 mmol) of a compound from Example 680 in
toluene (4 mL) was added 0.154 mL (0.72 mmol) of DPPA followed by 0.1 mL (0.72
mmol) of Et₃N. After heating the solution at 80 °C for 2 hours, 0.11 mL (0.78 mmol)
of the silyl alcohol was added and the reaction was heated at 85 °C for 12 hours. The
reaction was cooled, diluted with EtOAc, rinsed with water followed by brine, and
10 dried over Na₂SO₄. The solvent was removed *in vacuo* and the residue was
chromatographed on silica gel with 2/1 hexanes/EtOAc to yield 0.127g of the title
compound.

MS(ES+)m/z=504.

15

Example 678

N-[3*S*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)*R*-
cycloheptylmethyl]benzamide

A solution of 0.052g (0.2 mmol) of a compound from Example 677 in THF (6
20 mL), and 4.5 mL of TBAF (1.0M in THF)(4.5 mmol) was heated at 60 °C for 4 hours.
The solution was cooled to ambient temperature and water (2mL) then 0.028g of
anhydrous K₂CO₃ (0.2 mmol) were added followed by 0.016g (0.11 mmol) of
benzoyl chloride. The reaction was stirred 12 hours, after which the THF was
replaced with EtOAc and rinsed with 1N HCl and dried over Na₂SO₄. The organic
25 layer was concentrated *in vacuo* to a residue, which was chromatographed on silica
gel with 2/1 hexanes/EtOAc to yield 0.011g of the title compound.

MS(ES+)m/z=464.

-313-

Example 679

N-[3S-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-R-cycloheptylmethyl]-4-fluorobenzamide

5 In a manner similar to the preparation of Example 678, 0.052g (0.1 mmol) of a compound from Example 678 yielded 0.012g of the title compound.

MS(ES+)m/z=482.

Example 680

10 [3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cycloheptyl]acetic acid

A suspension of 0.82g (2.0 mmol) of a compound from Example 424 in 1N NaOH (45 mL) and MeOH (125 mL) was stirred for 16 hrs. at ambient temperature.

15 The MeOH was removed *in vacuo* and the reaction was cooled and neutralized with 3N HCl. The acidic aqueous layer was extracted with EtOAc, rinsed with water and concentrated *in vacuo* at 40° C to yield 0.78g of the title compound.

MS(ES+)m/z=388.9.

20

Example 681

[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cycloheptylmethyl]-carbamic acid methyl ester

25 In a manner similar to the preparation of Example 677 (except the intermediate isocyanate was trapped with MeOH instead of TMS ethanol) 0.68g (1.8 mmols) of a compound from Example 680 was converted to 0.43g of the title compound.

MS(ES+)m/z=418.1.

-314-

Example 682

N-[3*R*-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)*R*-cycloheptylmethyl]benzamide

5 A suspension of 0.036g (0.1 mmol) of a compound from Example 632, 0.028g (2.0 mmol) of anhydrous K₂CO₃, and 0.013 mL (1.1 mmol) of benzoyl chloride in 3 mL of THF/H₂O 2/1 was stirred 15 hours. The solvent was removed *in vacuo* and the residue was taken into EtOAc and rinsed with NaHCO₃ sat., 1N HCl, and brine. The organic layer was dried over anhyd. Na₂SO₄ and the solvent was removed *in vacuo*.
10 The residue was crystallized from MeOH to yield 0.28g of the title compound.
MS(ES+)m/z=464.1.

Example 683

15 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-3-phenylpropionamide

A solution of {1-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylcarbonyl]-2-phenyl-ethyl}-carbamic acid tert-butyl ester, 82 mg (0.14 mmol) in 15 mL of HCl-saturated acetic acid was stirred three hours at rt., then
20 concentrated to dryness. The residue was slurried 3x in acetonitrile and concentrated to dryness to give a quantitative yield of the title compound as a tan foam. MS (ion spray) 479.2. (M+).

Example 684

25 2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-3-hydroxypropionamide

A solution of {1-[3-(9-Chloro-3-methyl-4-oxo-4*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylcarbonyl]-2-hydroxy-ethyl}-carbamic acid tert-butyl ester, 73 mg
30 (0.14 mmol) in 8 mL of 25% TFA in CH₂Cl₂ was stirred rt. for three hours then concentrated to dryness. The residue was partitioned between 20%

-315-

isopropanol/chloroform and saturated aqueous NaHCO₃. The organics were washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by radial chromatography using MeOH/chloroform as eluent and concentrated to dryness to yield 40 mg (69%) of the title compound as a white solid. ¹H-NMR is consistent with structure. MS (IS) 419.1 (M+).

Example 685

2-Amino-N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-3-hydroxypropionamide

A solution of {1-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylcarbamoyl]-2-hydroxy-ethyl}-carbamic acid tert-butyl ester, 92.3 mg (0.18 mmol) in 8 mL of 25% TFA in CH₂Cl₂ was stirred rt. for three hours then concentrated to dryness. The residue was partitioned between 20% isopropanol/chloroform and saturated aqueous NaHCO₃. The organics were washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by radial chromatography using MeOH/chloroform as eluent and concentrated to dryness to yield 37 mg (50%) of the title compound as a white solid. MS (IS) 419.2 (M+).

Example 686

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]-6-dimethylaminonicotinamide

A solution of compound from Example 245, 50 mg, (0.10 mmol) in 2.0 mL (4.0 mmol) of 2*N* *N,N*-dimethylamine in THF was stirred 24 H in a sealed tube at 90°C then concentrated to dryness. The residue was dissolved in 20% isopropanol/chloroform and washed with saturated aqueous sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by radial chromatography using a methanol/chloroform

-316-

gradient as eluent and was concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to yield 48 mg (96%) of the desired isomer as a white foam.

¹H-NMR is consistent with structure.

5 MS (ion spray) 494.2 (M+).

Example 687

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-
cyclohexylmethyl]-6-methylaminonicotinamide

10

A solution of a compound from Example 245, 50 mg, (0.10 mmol) in 2.0 mL (4.0 mmol) of 2N methylamine in tetrahydrofuran was stirred 24 H in a sealed tube at 100°C then concentrated to dryness. Some mechanical loss occurred. The residue was dissolved in 20% isopropanol/chloroform and washed with saturated aqueous sodium bicarbonate, washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was purified by radial chromatography using a methanol/chloroform gradient as eluent and was concentrated to dryness. The residue was slurried in ether/hexanes and concentrated to dryness to yield 30 mg (62%) of the desired isomer as a white foam. MS (ion spray) 480.1 (M+).

15
20

Example 688

N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclohexyl]-2-(pyridin-3-yloxy)propionamide (isomers 1 & 2)

25

The isomers of N-[3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-2-(pyridin-3-yloxy)-propionamide (mixture of isomers, 142 mg (0.29 mmol) was separated via chiral chromatography to yield 63.5 mg of crude isomer 1 and 60.2 mg of crude isomer 2. Isomer 1 was purified by radial chromatography (methanol/chloroform gradient) and concentrated to dryness to yield 55.7 mg (39%) of the desired isomer as a tan foam. ¹H-NMR is consistent with structure.

30

MS (ion spray) 481.2 M+.

-317-

Isomer 2 was purified by radial chromatography (methanol/chloroform gradient) and concentrated to dryness to yield 51.3 mg (36%) of the desired isomer as a white foam.

¹H-NMR is consistent with structure.

MS (ion spray) 481.2 M+.

5

Example 689

N-{{[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclopentyl]methyl}benzamide

10 In the same manner as preparation 17, N-(*t*-butoxycarbonyl)-2-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)cyclopentylamine (0.06 g, 0.14 mmol) of was deprotected to yield 0.044 g (0.133 mmol) of the free amine MS(ES+)m/z=332, which was dissolved in dichloromethane (2.5 mL). To the solution was added 0.0222 mL (0.16 mmol) of Et₃N followed by 0.019 mL (0.16 mmol) of
15 benzoyl chloride. After 45 min, the solvent was removed *in vacuo*, replaced with ethyl acetate and the organic layer was rinsed with 1N HCl followed by 1N NaOH, then water. The organic layer was dried and the solvent was removed *in vacuo* to give a residue which was chromatographed on silica gel with ethyl acetate/hexane 1:1 to yield 0.035 g of the title compound. MS(ES+)m/z=436. Cis (racemic)

20

-318-

Example 690

Phenylmethyl 2-[(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]acetate

5 To a solution of phenylmethyl 2-((1R,3S)-3-{[4-(2-chloro-6-fluorophenyl)-2-methyl(3-furyl)]carbonylamino} cyclohexyl)acetate, 5.71 g (11.8 mmol) in 100 mL of N,N-dimethylformamide at ambient temperature was added 47 mL (23.6 mmol) of 0.5 M potassium bis(trimethylsilyl)amide in toluene dropwise. After 10 min., the reaction was quenched with saturated ammonium chloride and extracted with ethyl acetate.

10 The combined organics were washed with brine, dried over sodium sulfate, filtered and concentrated to dryness. The residue was chromatographed on silica gel using ethyl acetate/hexanes as eluent to yield 3.8 g (70%) of the desired product as a green oil.

¹H-NMR is consistent with structure.

15 MS (ion spray) 465.2 (M+1).

Example 691

5-((1S,3S)-3-[2-(phenylamino)ethyl]cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

20

N-((1S,3S)-3-[2-(phenylamino)ethyl]cyclohexyl)[3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl]carboxamide (54 mg; 0.118 mmol) was dissolved in anhydrous DMF (2 mL) under a dry nitrogen atmosphere at room temperature. When potassium bis(trimethylsilyl)amide (0.5 M in toluene; 572 μ L; 2.2 equiv) was added dropwise to this solution, a clear orange color formed. After 5 min, the reaction was chilled in an ice bath, quenched with 1N HCl_(aq) (10 mL), and extracted with ethyl acetate (twice). The organic layer was washed with saturated NaCl_(aq) (once), dried with Na₂SO_{4(s)}, filtered, and concentrated to dryness by rotary evaporation. The product was purified by radial chromatography on a 2 mm thick silica gel rotor. A dichloromethane mobile phase was ineffective at purifying the product. Re-chromatography with a 100% hexanes mobile phase followed by a 20% ethyl acetate/hexanes (v/v) mobile phase separated product from impurities.

-319-

After drying from diethyl ether, an off-white foam (43 mg; 84%) was obtained.

TOF-MS(ES) calc'd: $[M+H]^+ = 436.1792$ m/z. Found: 436.1797 m/z.

Example 692

5 5-[3-(2-Aza-bicyclo[2.2.1]hept-5-en-2-ylmethyl)-cyclohexyl]-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one

To a suspension of a compound from preparation 48 (500 mg, 1.0 mmol) in water (5 mL) was added formaldehyde (0.24 mL, 3.0 mmol) dropwise, and the
10 mixture stirred rapidly under nitrogen. After 5 minutes, freshly cracked cyclopentadiene (0.41 mL, 5.0 mmol) was added dropwise. The solution was stirred under nitrogen at rt. overnight. The solution was diluted with a saturated aqueous sodium bicarbonate solution, extracted with dichloromethane (x3), and the organics dried over magnesium sulfate and concentrated. Purification by flash
15 chromatography on silica gel (eluting with 1-4% methanol/chloroform/0.5% NH_4OH) gave 183.7 mg of the title compound as a white foam, 43% yield. MS (ion spray) 424 (M^+).

Example 693

20 1-Amino-cyclopentanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-4H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide hydrochloride

A compound from Example 460 (89.9 mg, 0.17 mmol) was dissolved in $HCl_{(g)}/AcOH$ (5 mL, ~3N) and stirred at rt.. After 1h, additional $HCl_{(g)}/AcOH$ (5 mL,
25 ~3N) was added. After 5h, the mixture was concentrated, followed by azeotropic removal of water with acetonitrile *in vacuo* (x3). To the white solid was added diethyl ether, and the mixture was sonicated, and filtered. The resulting material was dried on a vacuum pump to give 52.9 mg of the title compound as a white solid, 67% yield. 1H NMR: consistent with structure. MS (ion spray) 479 (M^+).

-320-

Example 694

1-Amino-cyclopropanecarboxylic acid [3-(9-chloro-3-methyl-4-oxo-4H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide hydrochloride

5

A compound from Example 461 (94.3 mg, 0.18 mmol) was dissolved in HCl_(g)/AcOH (5 mL, ~3N) and stirred at rt.. After 1h, additional HCl_(g)/AcOH (5 mL, ~3N) was added. After 5h, the mixture was concentrated, followed by azeotropic removal of water with acetonitrile *in vacuo* (x3). To the white solid was added diethyl ether, and the mixture was sonicated, and filtered. The resulting material was dried on a vacuum pump to give 70.2 mg of the title compound as a white solid, 85% yield. ¹H NMR: consistent with structure. MS (ion spray) 451 (M⁺).

10

Example 695

15 N-[3-(4,10-Dichloro-5-oxo-5H-benzo[h][1,6]naphthyridin-6-yl)-cyclohexylmethyl]-benzamide

To 0.045g (0.1 mmols) of N-{[(1R,3S)-3-(3-acetyl-4-amino-5-chloro-2-oxohydroquinolyl)cyclohexyl]methyl}benzamide and 0.03 ml (0.3mmols) of POCl₃ were dissolved in DMF (2 ml) at 0-5⁰C. The reaction was allowed to warm to rt. and stirred for 30 minutes. The reaction was then heated at 45⁰C over 12 hours. The solvent was removed *in vacuo* and the residue was chromatographed on silica and eluted with EtOAc/ Hexane 1:1 to 2:1 to yield 0.002g of the title compound. MS(ES+)m/z=480.

20
25Example 696

N-[3-(10-Chloro-4-methoxy-5-oxo-5H-benzo[h][1,6]naphthyridin-6-yl)-cyclohexylmethyl]-benzamide

30

To 0.045g (0.1 mmols) of N-{[(1R,3S)-3-(3-acetyl-4-amino-5-chloro-2-oxohydroquinolyl)cyclohexyl]methyl}benzamide and 0.03 ml (0.3mmols) of POCl₃ were dissolved in DMF (2 ml) at 0-5⁰C. The reaction was allowed to warm to rt. and stirred for 30 minutes. The reaction was then heated at 45⁰C over 12 hours. The

-321-

reaction was cooled to rt. and NaOMe in MeOH (0.4 mmols) was added. The reaction was heated at 60°C for 6 hours. The solvent was removed *in vacuo*. The residue was dissolved in CH₂Cl₂ and rinsed 1 time with 1N HCl, followed by 3 times with water. The solvent was removed in vacuo and the residue was chromatographed on silica and eluted with EtOAc/ Hexane 1:1 to 2:1 to yield 0.011g of the title compound.
MS(ES+)m/z=476.

Example 697

N-[3-(10-Chloro-4-methylamino-5-oxo-5H-benzo[h][1,6]naphthyridin-6-yl)-cyclohexylmethyl]-benzamide

To 0.005g (0.01 mmols) of a compound from Example 696 was combined with 1.0 ml of aqueous methylamine 40% (excess) in ethanol (5 ml). The reaction was refluxed for 30 minutes. MS(ES+)m/z=475.

Example 698

R(-)Amino-acetic acid [3-(9-chloro-3-methyl-4-oxo-4H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenyl-methyl ester hydrochloride

To a compound from Example 472 (96.7 mg, 0.16 mmol) was added acetic acid saturated with HCl_(g) (10 mL, ~3N in HCl) and the solution stirred vigorously at rt. for 1 hour. The reaction was concentrated, followed by addition of acetonitrile and concentration to assist in the removal of acetic acid (x2). The resulting white solid was treated with ethyl ether, sonicated, and filtered to yield 68.4 mg of the title compound as a white solid, 79% yield. ¹H NMR: consistent with structure. MS (ion spray) 523 (M⁺).

Example 699

S(+)-Amino-acetic acid [3-(9-chloro-3-methyl-4-oxo-4H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbamoyl]-phenyl-methyl ester hydrochloride

To a compound from Example 467 (70.0 mg, 0.11 mmol) was added acetic acid saturated with HCl_(g) (10 mL, ~3N in HCl) and the solution stirred vigorously at

-322-

rt. for 1 hour. The reaction was concentrated, followed by addition of acetonitrile and concentration to assist in the removal of acetic acid (x2). The resulting white solid was treated with ethyl ether, sonicated, and filtered to yield 37.84 mg of the title compound as a white solid, 60% yield. ^1H NMR: consistent with structure. MS (ion spray) 523 (M^+).

-323-

Example 700

{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentylcarbamoyl]-phenyl-methyl}-carbamic acid tert-butyl ester

5 A compound from Example 621 was converted to obtain the title compound with L-N-boc phenyl glycine as described for Example 515. ESMS: 565 (M+1)+.

Example 701

10 {[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentylcarbamoyl]-phenyl-methyl}-trifluoroacetamide

A compound from Example 621 was converted to obtain the title compound with L-N-trifluoroacetyl phenyl glycine as described for Example 515. ESMS: 561 (M+1)+.

15

Example 703

cis-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-(3-phenylureido)cyclohexane

20 To the suspension of *cis*-1-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-3-amino-cyclohexane (0.0446 g, 0.121 mmol) in 1 mL anhydrous THF at RT was added 2N NaOH (0.60 mL, 0.121 mmol). After 5 min. the phenylisocyanate (0.013 mL, 0.121 mmol). After 20 min. the reaction was diluted with EtOAc, washed with water and concentrated to near dryness. After sonication and filtration a white solid (0.0412, 75%) was obtained.

25

ESIMS *m/e* 451 ³⁵Cl (M⁺+1) and 453 ³⁷Cl (M⁺+1).

-324-

Example 704

2-[(3S,1R)-3-(9-Chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexylmethyl]-5*H*-isoquinoline-1,3-dione

5 To 5-[(1S,3R)-3-(aminomethyl)cyclohexyl]-9-chloro-3-methyl-5-hydroisoxazolo[4,3-*c*]quinolin-4-one (60 mg, 0.17 mmol) and homophthalic acid (38 mg, 0.21 mmol) was added xylenes (10 mL), and the reaction heated to reflux under nitrogen overnight with a Dean-Stark trap attached to effect removal of water. The solution was concentrated and dissolved in 20% isopropanol/chloroform, washed with
10 1.0N HCl (x2), saturated aqueous sodium bicarbonate solution, brine, dried over sodium sulfate and concentrated. Purification by flash chromatography on silica gel (eluting with 20-25% ethyl acetate/hexane) gave 26 mg of the title compound as a clear oil, 31% yield. ¹H NMR: consistent with structure. MS (ion spray) 490 (M⁺), 488 (M⁻ - 1).

15

Example 705

3-(2-Chloro-6-fluoro-phenyl)-5-methyl-isoxazole-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-ylmethyl)-cyclopentyl]-amide

20 A compound from Example 620 was acylated with 2-chloro-5-fluoro phenyl isoxazolyl chloride to obtain the title compound as described for Example 515. ESMS: 570 (M+1)+.

Example 706

25 1-tert-Butyl-3-(2-chloro-6-fluoro-phenyl)-1*H*-pyrazole-4-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-ylmethyl)-cyclopentyl]-amide

A compound from Example 620 was converted to obtain the title compound with 4-(2-chloro,5-fluoro phenyl)-1-*t*-butyl- pyrazole-3-carboxylic acid as described
30 for Example 515. ESMS: 610 (M+1)+.

-325-

Example 707

{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentylcarbamoyl]-phenyl-methyl}-carbamic acid tert-butyl ester

5 A compound from Example 621 was converted to obtain the title compound with L-N-boc phenyl glycine as described for Example 515. ESMS: 565 (M+1)+.

Example 708

10 1R, 3S-N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-ylmethyl)-cyclopentyl]-2-(3,4,5-trimethoxy-phenyl)-acetamide

15 A mixture of 1R, 3S-5-(3-amino-cyclopentylmethyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one (15 mg, 0.045 mmol), 3,4,5-trimethoxyphenyl acetic acid (14 mg, 0.06 mmol), EDC (12 mg, 0.06 mmol) and DMAP (2 mg) dissolved in DCM (10 mL) was stirred overnight at rt. The reaction mixture was diluted with additional DCM (10 mL), washed with aq. NaHCO₃ (sat'd, 2x 20 mL), water (2x20 mL), brine (2x20 mL), dried over sodium sulfate, filtered, evaporated, and chromatographed (Bond Elut, Si, 60 cc, 50% EtOAc in hexanes) to yield the desired product (10 mg, 41%). ESMS: 540 (M+1)+.

20

Example 709

N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl-cyclopentyl]-2-(3,4,5-trimethoxy-phenyl)-acetamide

25 To a stirred solution of racemic 5-(3-aminocyclopentyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one (81 mg, 0.26 mmol) in CH₂Cl₂ (3 mL) was added 3,4,5-trimethoxy phenyl acetic acid (114 mg, 0.52 mmol), Et₃N (0.1 mL, 0.50 mmol), HOBt (41 mg, 0.3 mmol), DMAP (catalytic amount) and EDCI (72 mg, 0.38 mmol). The reaction mixture was stirred at r.t. for 18 hours. The mixture was diluted with
30 CH₂Cl₂, washed (brine), dried (Na₂SO₄), filtered and concentrated. Column chromatography (silica gel, hexanes/ethyl acetate, gradient) gave the racemic title compound (89 mg, 66%). Mass Spectrum (FIA) (m/z) 454.0 (M+1)

-326-

Example 710

2-[3S-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5yl)-R-cycloheptyl]-N-(3,4,5-trimethoxyphenyl)-acetamide

5

To a solution of 0.038g (0.1 mmol) of 2-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cycloheptyl]acetic acid in THF/DMF 1-1 (5 mL) was added 0.061g (0.44 mmol) of 1-hydroxy-7-azabenzotriazole and 0.86g (0.44 mmol) of EDCI. After 40 minutes, 0.083g (0.44 mmol) of 3,4,5-trimethoxyaniline was added.

10 After 3.25 hours, the solvent was removed in vacuo and replaced with EtOAc. The organic layer was rinsed with 1N HCl, aq. NaHCO₃, then x3 with water. The organic layer was dried and the solvent was removed in vacuo to yield 0.026g of the title compound. MS(ES+)m/z=554.

15

Example 711

2-Acetylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-phenylacetamide

20 A compound as described in Example 633 (75 mg, 0.16 mmol) was combined with 1,3-dicyclohexylcarbodiimide (50 mg, 0.24 mmol), 1-hydroxybenzotriazole hydrate (33 mg, 0.24 mmol), and glacial acetic acid (0.015 mL, 0.24 mmol) in tetrahydrofuran (6 mL) and the mixture stirred over the weekend at ambient temperature. The mixture was then concentrated *in vacuo* and the residue loaded onto a silica gel column and eluted with methanol/dichloromethane, which allowed for
25 isolation of 82 mg (99%) of product as an off white solid. MS(ES): (M+1)⁺ 507.1, 509.1 m/z.

-327-

Example 712

2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexylcarbamoyl]-pyrrolidine-1-carboxylic acid benzyl ester

5 A compound as described in Example 660 (60 mg, 0.18 mmol) was combined
with DCC (56 mg, 0.27 mmol), HOBT (37 mg, 0.27 mmol), and N-carbobenzyloxy-
L-proline (68 mg, 0.27 mmol) in tetrahydrofuran (6 mL) and the mixture stirred
overnight at rt. The mixture was then concentrated *in vacuo* and the residue loaded
onto a silica gel column and eluted with MeOH/CH₂Cl₂, which allowed for isolation
10 of 71 mg (70%) of product as a white solid. MS(ES): (M+1)⁺ 563.0, 565.0 m/z.

Example 713

2-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]
quinolin-5-yl)-cyclohexylcarbamoyl]-pyrrolidine-1-carboxylic acid benzyl ester

15

A compound as described in Example 660 (40 mg, 0.12 mmol) was combined
with DCC (37 mg, 0.18 mmol), HOBT (24 mg, 0.18 mmol), and N-carbobenzyloxy-
D-proline (45 mg, 0.18 mmol) in tetrahydrofuran (4 mL) and the mixture stirred
overnight at rt. The mixture was then concentrated *in vacuo* and the residue loaded
20 onto a silica gel column and eluted with MeOH/CH₂Cl₂, which allowed for isolation
of 27 mg (40%) of product as a white solid. MS(ES): (M+1)⁺ 563.0, 565.0 m/z.

Example 714

1-Benzoyl-pyrrolidine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-
isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide

25

A compound as described in Example 660 (40 mg, 0.12 mmol) was combined
with EDC, HOBT (25 mg, 0.18 mmol), and the compound from preparation 104 (40
mg, 0.18 mmol) in tetrahydrofuran (6 mL) and the mixture stirred overnight at rt. The
30 mixture was then concentrated *in vacuo* and the residue loaded onto a silica gel
column and eluted with EtOAc, which allowed for isolation of 58 mg (90%) of
product as a white solid. MS(ES): (M+1)⁺ 533.3, 5535.3 m/z.

-328-

Example 715

1-Phenylacetyl-pyrrolidine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-
isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide

5

A compound as described in Example 660 (40 mg, 0.12 mmol) was combined with EDC (37 mg, 0.18 mmol), HOBT (25 mg, 0.18 mmol), and the compound from preparation 8R (42 mg, 0.18 mmol) in tetrahydrofuran (5 mL) and the mixture stirred over the weekend at rt. The mixture was then concentrated *in vacuo* and the residue
10 loaded onto a silica gel column and eluted with EtOAc, which allowed for isolation of 54 mg (82%) of product as a white solid. MS(ES): (M+1)⁺ 547.1 m/z.

Example 716

1-Benzyl-pyrrolidine-2-carboxylic acid [3-(9-chloro-3-methyl-4-oxo-5H-
isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-amide

15

A compound as described in Example 660 (40 mg, 0.12 mmol) was combined with DCC (37 mg, 0.18 mmol), HOBT (25 mg, 0.18 mmol), and the compound from preparation 107 (49 mg, 0.24 mmol) in tetrahydrofuran (5 mL) and the mixture stirred
20 over the weekend at rt. The mixture was then concentrated *in vacuo* and the residue loaded onto a silica gel column and eluted with MeOH/CH₂Cl₂. A repeat of the chromatography yielded the title compound (56 mg, 89%). MS(ES): (M+1)⁺ 519.3, 520.3 m/z.

25

Example 717

N-{{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexylcarbonyl]-methyl}-2-phenyl-acetamide

5-(3-Amino-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-
30 one hydroiodide (70 mg, 0.19 mmol) was combined with EDC (47 mg, 0.25 mmol), 1-hydroxy-7-azabenzotriazole (34 mg, 0.25 mmol), N,N-diisopropylethyl amine (0.10 mL, 0.58 mmol), DMAP (5 mg, cat.), and N-benzylglycine hydrochloride (50

-329-

mg, 0.25 mmol) in DMF(6 mL) and the mixture stirred overnight at rt. The mixture was then concentrated *in vacuo* and the residue taken up in chloroform/MeOH and the organic solution washed with aqueous NaHCO₃. The organic solution was dried over Na₂SO₄ and concentrated in vacuo. The residue was loaded onto a silica gel column and eluted with MeOH/CH₂Cl₂, which allowed for the recovery of 32 mg (35%) of the free base. This material was dissolved in minimal EtOAc and treated with excess diethyl ether/hydrochloric acid. Concentration of this mixture to dryness allowed for quantitative recovery of the hydrochloride salt as an off white solid. MS(ES): (M+1)⁺ 479.1, 481.2.

10

Example 718

1-methyl-piperidine-4-carboxylic acid {[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexylcarbonyl]-phenyl-methyl}-amide

15

A product from Example 638 (50 mg; 0.108 mmol) was dissolved in anhydrous dimethylformamide (10 mL) under a nitrogen atmosphere, mixed with 1-methyl-piperidine-4-carboxylic acid hydrochloride (58.0 mg; 0.323 mmol; 3 equiv), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (61.8 mg; 0.323 mmol; 3 equiv), 2,4,6-trimethylpyridine (86 μL; 0.645 mmol; 6 equiv), and 1-hydroxy-7-azabenzotriazole (43.9 mg; 0.323 mmol; 3 equiv), and stirred overnight at room temperature. The reaction solution was diluted with 2 volumes ethyl acetate, 10 volumes water, and 1 volume saturated NaHCO_{3(aq)}. The organic layer was separated and then washed with saturated NaCl_(aq), dried over Na₂SO_{4(s)}, filtered, and concentrated in vacuo. The resulting material was purified by radial chromatography on a 2 mm thick silica gel rotor with a 2% then 4% methanol/dichloromethane (v/v) mobile phase and finally a 2% methanol/0.5% triethylamine/dichloromethane (v/v/v) mobile phase. The desired product was isolated as a white solid in 74% yield (47 mg).

25

-330-

Example 719

N-[3-(9-Acetylamino-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexylmethyl]-benzamide

5 To a solution of a compound from Example 667 (0.025 g, 0.06 mmol) and
AcOH (0.02 mL, 0.36 mmol) in DMF (1 mL) under N₂ was added EDCI (0.023 g,
0.12 mmol) and DMAP (0.002 g, 0.01 mmol). The solution was stirred overnight,
diluted with EtOAc, washed (H₂O then brine), dried (MgSO₄), filtered, and
concentrated. Flash chromatography (silica gel, acetone/ CH₂Cl₂ gradient) gave the
10 title compound (0.023 g, 82%). Mass Spectrum (ES+) (m/z) 473.3 [M+1]

Example 720

N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-
phenylamino-acetamide

15

Anhydrous dimethylformamide (10 mL) in a nitrogen atmosphere was used to
dissolve 5-(3-aminocyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-
one (50 mg; 0.151 mmol), N-phenylglycine (29.6 mg; 0.196 mmol; 1.3 equiv), 1-
hydroxy-7-azabenzotriazole (26.7 mg; 0.196 mmol; 1.3 equiv), 1-[3-
20 (dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (37.6 mg; 0.196 mmol;
1.3 equiv), and 2,4,6-trimethylpyridine (199 μ L; 1.51 mmol; 10 equiv). After
overnight stirring at room temperature, the reaction solution was treated in a manner
similar to Example 159. The resulting material was purified by three consecutive
radial chromatography runs on a 2 mm thick silica gel rotor with a 1%
25 methanol/dichloromethane (v/v) mobile phase, a 1% methanol/0.25%
triethylamine/dichloromethane (v/v/v) mobile phase, and a 5%
acetonitrile/dichloromethane mobile phase. The desired product was isolated as a
white solid 56% yield (30 mg). MS(ES) calc'd: [M+H]⁺ = 465.16 m/z; [M-H]⁻ =
463.16 m/z; [M+OAc]⁻ = 523.16 m/z. Found: 465.1 m/z; 463.2 m/z; 523.2 m/z.

30

-331-

Example 721

2-[3*R*-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-*c*]quinolin-5-yl)-*R*-cycloheptyl]-
N-(3,4,5-trimethoxyphenyl)-acetamide

5 In a manner similar to the preparation of Example 710, 0.055g (0.14 mmol) of a compound from Example 680 was converted to a residue which was chromatographed on silica with EtOAc/hexanes 1/1 to yield 0.044g of the title compound. MS(ES+)m/z = 553.9.

10

Example 722

6-Chloro-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-*c*]quinolin-5-yl)-
cyclohexylmethyl]-nicotinamide

A solution of cis-5-[3-(aminomethyl)cyclohexyl]-9-chloro-3-methyl-5-
15 hydroisoxazolo[4,5-*c*]quinolin-4-one (0.188 g, 0.54 mmol) and 6-chloro-nicotinic acid (0.103 g, 0.65 mmol) in DMF (10 mL) was treated with EDCI (0.125 g, 0.65 mmol) and HOAt (0.089 g, 0.65 mmol). The reaction was then treated with excess Et₃N (0.167 g, 1.62 mmol) and DMAP (6 mg, 0.05 mmol) and stirred at room
20 temperature overnight. The reaction was concentrated to a solid and then taken up in 20% isopropanol in CHCl₃ (100 mL). This solution was transferred to a separatory funnel and washed with a saturated sodium bicarbonate solution (3 x 50 mL) and brine (2 x 50 mL). The organic solution was dried over sodium sulfate, filtered, and the solvent removed to afford a crude yellow solid. The solid was purified using silica gel column chromatography. The product was eluted using 50% EtOAc in
25 CHCl₃. The solvent was removed *in vacuo* to afford 0.150 g (57%) of product as an off white solid. MS m/z (ES+) 484.8 (M+H)⁺, (ES-) 482.8 (M-H)⁻, 542.8 (M+CH₃COO)⁻.

-332-

Example 723

2-tert-butylamino-N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-pyridin-3-yl-acetamide

5 A compound from preparation 341 was saponified over 2 h with LiOH (1.5 equiv in water subsequently mixed with 1,4-dioxane). The carboxylic acid was isolated by rotary evaporation of the solution to dryness, acidification to pH 2 with 1 N HCl (aq), rotary evaporation of the solution to dryness, and overnight drying at 0.1 torr. This material (1.3 equiv) was mixed with the free base of material from
10 Preparation 210 (100 mg; 0.301 mmol) in anhydrous DMF. Diisopropylethylamine (262 μ L; 0.392 mmol; 5 equiv), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (75.1 mg; 0.392 mmol; 1.3 equiv), and 1-hydroxy-7-azabenzotriazole (53.3 mg; 0.392 mmol; 1.3 equiv) were then added and the solution was stirred overnight, at rt. EtOAc (70 mL) and saturated NaHCO₃ (aq)(10 volumes) were added
15 and the organic layer was separated, washed with saturated NaCl (aq)(once), dried with Na₂SO₄ filtered, and concentrated in vacuo. After purification by radial chromatography, 34 mg off-white solid was isolated (22% yield). MS(ES) calc'd: [M+H]⁺ = 522.2 m/z; [M-H]⁻ = 520.2 m/z. Found: 522.2 m/z; 520.3 m/z.

20

Example 724

N-[3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl]-2-(2,2-dimethyl-propylamino)-2-pyridin-3-yl-acetamide

A compound from preparation 341 was used in a manner similar to Example
25 723 (substituting triethylamine for diisopropylethylamine) to prepare the title compound (24 mg white solid; 15% yield). MS(ES) calc'd: [M+H]⁺ = 536.2 m/z. Found: 536.2 m/z.

-333-

Example 725

5 *N*-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-2-(4-methyl-piperazin-1-yl)-2-pyridin-3-yl-acetamide

A compound from preparation 343 was used in a manner similar to Example 723 (substituting Et₃N for diisopropylethylamine) to prepare the desired product. A 1% then 2% then 3% methanol/chloroform/trace Et₃N (v/v/v) mobile phase on a 2 mm chromatotron rotor was used to purify the desired product (129 mg white solid; 78% yield). MS(exact mass) calc'd: [M+H]⁺ = 594.2381 m/z. Found: 594.2385 m/z.

Example 726

15 2-(Benzenesulfonyl-pyridin-2-yl-amino)-*N*-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-acetamide

To a solution of 5-(3-amino-cyclohexyl)-9-chloro-3-methyl-5*H*-isoxazolo[4,3-*c*]quinolin-4-one hydroiodide, 130 mg (0.28 mmol) in 10 mL of DMF was added 82 mg (0.28 mmol) of (benzenesulfonyl-pyridin-2-yl-amino)-acetic acid (Bionet), 46 mg (0.34 mmol) of 1-hydroxy-7-azabenzotriazole, 66 mg (0.34 mmol) of EDC, 5 mg of DMAP and 120 μL (0.84 mmol) of TEA. The reaction mixture was stirred six days at rt. and was concentrated to dryness. The residue was dissolved in 20% isopropanol/chloroform, washed with saturated NaHCO₃, washed with brine, dried over Na₂SO₄, filtered and concentrated to dryness. The residue was purified by radial chromatography using a MeOH/chloroform gradient and concentrated to dryness. 25 The residue was slurried in ether/hexanes and concentrated to dryness to yield 79 mg (47%) of the title compound as a tan foam. MS (ion spray) 606.1 (M⁺).

-334-

Example 727

N-[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,5-c]quinolin-5-yl)-
cyclohexyl]-2-hydroxy-2-phenyl-acetamide

5 A solution of a compound from preparation 380 (0.08 g, 0.24 mmol), *R*-Mandelic acid (0.04 g, 0.29 mmol), EDCI (0.06 g, 0.29 mmol), HOAt (0.04 g, 0.29 mmol) in DMF (10 mL) was treated with Et₃N (0.07 g, 0.7 mmol) and DMAP (3.0 mg, 0.02 mmol) and stirred overnight at room temperature. The solution was then
10 diluted in EtOAc (50 mL) and washed with 5% citric acid (3 x 10 mL), sat'd sodium bicarbonate (3 x 10 mL), and brine (2 x 10 mL). The solution was dried over sodium sulfate and purified by silica gel column chromatography. The solvent was removed *in vacuo* to afford 0.08 g (74%) as a white solid.

Example 728

15 N-{[3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexylcarbonyl]-methyl}-nicotinamide

5-(3-Amino-cyclohexyl)-9-chloro-3-methyl-5H-isoxazolo[4,3-c]quinolin-4-one hydroiodide (70 mg, 0.19 mmol) was combined with EDC (47 mg, 0.25 mmol),
20 1-hydroxy-7-azabenzotriazole (34 mg, 0.25 mmol), *N,N*-diisopropylethyl amine (0.10 mL, 0.58 mmol), (5 mg, cat.), and nicotinic acid (44 mg, 0.24 mmol) in DMF (6 mL) and the mixture stirred overnight at rt. The mixture was then concentrated *in vacuo* and the residue taken up in water and extracted with EtOAc. The combined extracts were dried over Na₂SO₄ and concentrated *in vacuo*. The
25 residue was loaded onto a silica gel column and eluted with MeOH/CH₂Cl₂ which allowed for the recovery of 87 mg (92%) of the title compound as a white solid.
MS(ES): (M+1)⁺ 494.1, 496.1.

-335-

Example 729

N-[3-(9-chloro-3-methyl-4-oxo-5*H*-isoxazolo[4,3-*c*]quinolin-5-yl)-cyclohexyl]-2-dimethylamino-2-pyridin-3-yl-acetamide

5 A compound from preparation 342 was used in a manner similar to Example 723 (substituting Et₃N for diisopropylethylamine) to prepare the desired product. A 1% methanol/chloroform/trace triethylamine (v/v/v) mobile phase on a 2 mm chromatotron rotor was used to purify the desired product (112 mg white solid; 76% yield). MS(exact mass) calc'd: [M+H]⁺ = 494.1959 m/z. Found: 494.1978 m/z.

10

Preparation 406

Pyridin-3-yl-(pyridin-2-yloxy)-acetic acid ethyl ester

15 A typical synthesis of bromopyridin-3-ylacetic acid ethyl ester is described: A solution of fresh lithium diisopropylamide (LDA) was prepared at -10 °C from diisopropylamine (8.54 mL; 60.5 mmol) and *n*-butyllithium (37.8 mL of a 1.6 M hexanes solution; 12.1 mmol) and stirred for 10 min. After chilling the fresh LDA to -78 °C, ethyl 3-pyridylacetate (9.21 mL; 60.5 mmol) was added and the solution was stirred another 10 min. Chlorotrimethylsilane (7.68 mL; 60.5 mmol) was added to the
20 resulting opaque yellow slurry and the solution was stirred 5 min. Finally a solution of 4-(dimethylamino)pyridinium tribromide (22.0 g; 60.5 mmol) in tetrahydrofuran was added and the reaction solution was stirred 10 min. After warming to room temperature the reaction solution was quenched with saturated NH₄Cl_(aq) and extracted twice with ethyl acetate. The combined organic layer was washed once with saturated
25 NaCl_(aq), dried with Na₂SO_{4(s)}, filtered, and concentrated in vacuo. The resulting bromopyridin-3-ylacetic acid ethyl ester was an unstable brown oil and was therefore used immediately.

30 Silver carbonate (2.79 g; 10.1 mmol; 0.5 equiv) was refluxed in toluene (40 mL) for 0.5 h with 2-hydroxypyridine (1.92 g; 20.2 mmol; 1.0 equiv) and an aliquot of bromo-pyridin-3-yl-acetic acid ethyl ester (approx. 4.9 g; 20.2 mmol), prepared as described above. After cooling to room temperature and filtration of the reaction mixture, the supernatant was washed with saturated NaHCO_{3(aq)} (1x), water (1x), and

-336-

saturated $\text{NaCl}_{(\text{aq})}$ (1x). The organic layer was dried over $\text{Na}_2\text{SO}_{4(\text{s})}$, filtered, and concentrated in vacuo. Silica chromatography with a 35% ethyl acetate/hexanes mobile phase produced a yellow oil (1.14 g; 22% yield) that solidified upon standing. MS(ES) calc'd: $[\text{M}+\text{H}]^+ = 259.2$ m/z; $[\text{M}-\text{H}]^- = 257.2$ m/z. Found: 259.1 m/z; 257.2 m/z. REFERENCE: U. Schöllkopf, I. Hoppe, Justus Liebigs Ann. Chem. (1972) 765, 153-170.

Example 730

Preparation a

10 1,3-cyclohexanedicarboxylic acid

To a suspension of isophthalic acid (500 g, 3 mol) in methanol (2.8 l) was added 5% Rhodium-on-alumina catalyst (50 g) and acetic acid (150 ml). The reaction mixture was shaken under hydrogen (50 psi) at room temperature overnight. The mixture was filtered through celite. To this solution was added fresh 5% Rhodium-on-alumina catalyst (25 g), and the mixture was shaken under 50 psi of hydrogen for another 24 hours. The final reaction mixture was filtered through celite. The solution was concentrated under vacuum to give 493 g of the title compound as a white powder (96.3 % yield). m.p. 163-165 °C.

20

Preparation b

3-Oxabicyclo[3.3.1]nonane-2,4-dione

A solution of dicyclohexylcarbodiimide (200 g, 1.16 mol) in CH_2Cl_2 (1000 ml) was added dropwise to a suspension of compound from preparation a (257 g, 1.25 mol) in CH_2Cl_2 (550 ml), and the mixture was stirred at room temperature for 4 hours. The precipitated dicyclohexylurea was filtered and washed several times with cold CH_2Cl_2 (200 ml x 3). The combined organic layer was concentrated to give a white solid, which was suspended in MTBE (900 ml). This solid was collected by filtration, washed with MTBE (250 ml), and dried under house vacuum to give the title

30

-337-

compound (137 g). The filtrate was concentrated to a residue, which was suspended in MTBE (250 ml) to give another 31 g anhydride. The total yield was 168 g (94%). m.p. 138-140°C.

5

Preparation c

cis-1,3-Cyclohexanedicarboxylic acid diethyl ester

To a solution of compound from preparation b (31 g, 0.2 mol) in ethanol (anhydrous, 310 ml) was added p-toluenesulfonic acid monohydrate (1.9 g, 10 mmol, 10 0.05 equiv.) and triethyl orthoformate (50 ml, 0.3 mol). The reaction mixture was stirred at 60°C overnight. The volatiles were stripped and the residue was diluted with ethyl acetate (250 ml), washed with water (120 ml) and brine (100 ml), and dried over MgSO₄. After filtration and evaporation, the residue was purified by chromatography. Eluting the column with 10% ethyl acetate in hexane afforded the 15 title compound (40 g, 87.7% yield).

¹H NMR: (500 MHz, CDCl₃) δ 4.11 (q, J=7.0 Hz, 4H), 2.29 (dt, 2H), 2.11 (dd, 1H), 1.97 (m, 2H), 1.98 (m, 1 H), 1.53 (q, J=12.5 Hz, 2H), 1.30-1.40 (m, 2H), 1.25 (t, J=7.0 Hz, 6H).

20

Preparation d

1,3-Cyclohexanedicarboxylic acid, monoethyl ester (1R, 3S)

To a suspension of compound from preparation c (40 g, 17.5 mmol) in pH 7.2 phosphate buffer [0.2 M] (1.2 l) was added lipase AY30 (Amano, 16.7 g). The 25 mixture was stirred vigorously at room temperature for 30 hours. The mixture was acidified with 10-15% HCl to pH < 2, and extracted with ethyl acetate (500 ml x 2). The combined organic solution was washed with aqueous 10% Na₂CO₃ (100 ml x 2) and water (100 ml x 2). The combined aqueous layers were washed again with ethyl acetate (150 ml) and then acidified with 10-15% HCl to pH < 2. The acidified 30 aqueous was then extracted with ethyl acetate (150 ml x 3). The combined organic solution was dried over MgSO₄. After filtration and concentration the title compound (35.6 g, 100% yield) was obtained.

-338-

¹H NMR (500 MHz, CDCl₃) δ 4.12 (q, J=7.0 Hz, 2H), 2.20-2.40 (m, 3H), 1.85-2.05 (m, 3H), 1.5 (q, 2H), 1.35 (m, 2 H), 1.24 (t, J=7.0 Hz, 3H).

Preparation e

5 Ethyl-[3-N-(methylcarbamate)-cyclohexyl]-carboxylate (1R, 3S)

A solution of a compound from preparation d (73 g, 365 mmol) in toluene (750 ml) was heated to reflux using a Dean-Stark trap to separate trace amounts of water. After collecting about 10 ml of water, the mixture was cooled down to about
10 40-50°C. To this mixture was added triethylamine (56 ml, 0.4 mol), and diphenylphosphoryl azide (86.5 ml, 0.4 mol). The reaction mixture was stirred at 110 °C for 60 min, cooled to 70 °C, and methanol (64 g, 2 mol) was added with stirring. After addition, the final reaction mixture was then heated to 85°C overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate (700 ml) and
15 washed with water (500 ml). The aqueous layer was extracted with ethyl acetate (500 ml x 2). The combined organic solution was washed again with water (500 ml) and brine (500 ml). After drying over MgSO₄ and concentration under reduced pressure, the title compound was obtained as a colorless oil (86 g, 100 %).

¹H NMR: (300 MHz, CDCl₃) δ 4.60 (sb, 1H), 4.13 (q, 2H), 3.65 (s, 3H), 3.50 (sb,
20 1H), 2.38 (t, 1H), 2.23 (d, 1H), 2.00-1.80 (m, 3H), 1.24 (t, 3H), 1.12-0.95 (m, 1H).

Preparation f

Ethyl-((1R, 3S)-3-{{3-(2-chloro-6-fluorophenyl)-5-methylisoxazol-4-yl}carbonyl-amino}cyclohexyl)-carboxylate
25

To a solution of compound from preparation e (86 g, 365 mmol) in CH₂Cl₂ (750 ml) was added iodotrimethylsilane (100 g, 500 mmol) in one portion, at room temperature. The reaction mixture was stirred for 2 hours at ambient temperature, cooled to 0-5 °C, and methanol (50 ml) was added. After stirring 15 minutes, the
30 solution was concentrated under reduced pressure. The residue was dissolved in THF (1l). To this solution was added water (0.5 l), potassium carbonate (138 g, 1 mol), and a solution of 3-(6-fluoro-2-chlorophenyl)-5-methylisoxazole-4-carboxyryl

-339-

chloride (110 g, 0.4 mol) in 250 ml THF, dropwise. After the addition, the reaction mixture was heated to room temperature and stirred for 12 hours. THF was removed under house vacuum, water (250 ml) was added, and the mixture was extracted with ethyl acetate (500 ml x 3). The combined organic solution was washed with saturated sodium thiosulfate (150 ml), water (500 ml), brine (500 ml) and then dried over MgSO₄. After filtration and evaporation under vacuum, the residue was purified by recrystallization from MTBE (250 ml). Repeating this recrystallization procedure three times provided the title compound (122.7 g, 82.5% yield) as a white powder. IR: ν_{\max} (film) 3429, 3011, 2940, 1725, 1662, 1187 cm⁻¹.

10

Preparation g

Ethyl [(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quindin-5-yl)-cyclohexyl]-carboxylate

15

To a solution of compound from preparation f (78 g, 190 mmol) in DMF (750 ml) was added a solution of KHMDS ([0.5M], 400 ml, 200 mmol). The temperature was kept at 25 °C by using an ice-bath. After the addition was complete, the reaction mixture was analyzed by TLC (silica gel, 50 % EtOAc in hexane) and found to be complete. Water (1 l) was added and the mixture was extracted with EtOAc (800 ml x 3). The combined organic solution was washed with 1N HCl (250 ml), water (250 ml), brine (250 ml), dried over MgSO₄ and concentrated to give a residue which was purified by recrystallization from MTBE (500 ml) to afford 66 g of the title compound as a light yellow powder (89.0% yield).

20

IR: ν_{\max} (film) 3030, 1720, 1670, 1220 cm⁻¹.

25

Preparation h

(1R, 3S) 3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl carboxylic acid

30

To a solution of compound from preparation g (62 g, 160 mmol) in THF (600 ml) was added 5N aqueous sodium hydroxide (120 ml) at room temperature. The reaction mixture was heated to 60 °C for 15 hours with stirring. After cooling to room

-340-

temperature, water (750 ml) was added and the mixture was washed with ethyl acetate (500 ml). The aqueous phase was separated and acidified with 15% HCl to pH < 2. The aqueous phase was then extracted with methylene chloride (1000 ml x 3). The combined organic extracts were washed again with water (500 ml), brine (500 ml),
5 and dried over MgSO₄. After filtration and evaporation under vacuum, the dark brown residue was suspended in MTBE (1000 ml), and refrigerated overnight. The mixture was filtered to afford 55.45 g (96.4%) of bright yellow product.
IR: ν_{\max} (Film) 3200, 2936, 1726, 1643, 1595, 1173 cm⁻¹.

10

Preparation i

2-methyl [(1R,3S)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-
cyclohexyl] carbamate

To a suspension of compound from preparation 8 (55.4 g, 154 mmol) in
15 toluene (1l) was added triethylamine (23.7 ml, 170 mmol), and diphenylphosphoryl azide (36.5 ml, 170 mmol). The reaction mixture was stirred at 110 °C for 2 hours during which time a solution formed. The solution was cooled to 80 °C and methanol (25 g, 0.77 mol) was added with stirring. The solution was warmed to 85 °C for 22 hours. After cooling to room temperature, the toluene was removed under reduced
20 pressure and the residue was dissolved in dichloromethane (3l) and washed with water (1l). The aqueous phase was extracted with dichloromethane (1l x 2) and the combined organic solution was washed again with water (500 ml) and brine (500 ml). After drying over MgSO₄, the solution was concentrated under vacuum. The crude product was purified by crystallization (CH₂Cl₂/MTBE, 0.5 l/2 l) to afford the title
25 compound (46.6 g, 78.2 %). $[\alpha]_D +49.2^\circ$, $[\alpha]_{365} +263.3^\circ$ (c, 0.56; CHCl₃). The filtrate was concentrated to a residue, which was purified by chromatography to obtain a second crop of product (5.1 g). The total yield was 86.8 %.
IR ν_{\max} (Film) 3410, 3020, 2950, 1710, 1670, 1220 cm⁻¹.

-341-

Preparation j

(1R, 3S) 3-(9-Chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl
methyl alcohol

5 To a solution of compound from preparation h (4.4 g, 12.2 mmol) in THF (45 ml) was added borane-methyl sulfide complex (2.0 M solution in THF, 12.5 ml, 25 mmol) dropwise at 0 °C. When the addition was complete, the reaction mixture was allowed to warm to room temperature and was stirred for one hour. Methanol (10 ml) was added slowly (gas generated) with stirring. The reaction mixture was then poured
10 into ice-water (60 ml) and extracted with ethyl acetate (100 ml x 3). The combined organic solution was washed with 1N HCl (50 ml), brine (50 ml), and dried over MgSO₄. After filtration and evaporation under vacuum, the yellow title compound (4.34 g, 100%) was obtained. M.S.: m/z 347 (M⁺+1).

15

Preparation k

(1R, 3S) 3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)-cyclohexyl
methyl alcohol mesylate

To a stirred solution of compound from preparation j (4.32 g, 12.5 mol) in
20 pyridine (25 ml) was added DMAP (10 mg) and methanesulfonyl chloride (1.16 ml, 15 mmol, 1.2 equiv) and the resulting mixture was stirred at room temperature for 1.5 h. Water (100 ml) was added, and the mixture was extracted with ethyl acetate (150 ml x2). The combined organic extracts were washed again with brine (100 ml), dried (MgSO₄) and concentrated. The residue was suspended in MTBE (25 ml) and filtered
25 to obtain the title compound as a solid (4.65 g, 87.8%).
[α]_D +8.71°, [α]₃₆₅ +58.7° (c, 0.358; CHCl₃).

DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME 1 DE 2
CONTENANT LES PAGES 1 À 341

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME 1 OF 2
CONTAINING PAGES 1 TO 341

NOTE: For additional volumes, please contact the Canadian Patent Office


NOM DU FICHER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

- 363 -

We Claim:

1. A compound of formula I:



where:

A is a C₃-C₈ cycloalkyl, optionally substituted 1-3 times with a C₁-C₄ alkyl;

10 het is a five (5) membered heterocyclic ring comprising N and a second heteroatom selected from N, O, or S;

wherein the non-fused carbon atom of the heteroaryl ring may be optionally substituted with R^b: C₁-C₆ alkyl, optionally substituted aryl, optionally substituted heterocycle, an amino acid ester, CH₂OH, CH₂O-heterocycle, halo, CH₂N₃, CH₂SR¹, CH₂NR⁴R⁶, OR¹, SR¹³, S(CH₂)_k-phenyl, or NR⁴R⁶; provided that when het is pyrazole or imidazole, the saturated nitrogen of the het ring may be optionally substituted with R^a: C₁-C₄ alkyl;

20 k is 0, 1, 2, 3, or 4;

n is 0, 1, or 2;

p is 0 or 1;

q is 0, 1, or 2;

r is 0, 1, or 2;

- 364 -

t is 0, 1, 2, 3, or 4;

u is 0, 1, 2, 3, or 4;

Y is $-E-C(O)R^3$, $-E-CH=CHR^{13}$, $-E-C(OH)R^{13}$, $-E-NR^4R^5$, $-E-OR^2$, $-E-$
 5 $S(O)_qR^{13}$, $-E-SO_2NR^4R^6$, $-C(R^{11})=NR^6$, or an optionally substituted heterocycle;

E is a bond or $-C(R^{11})(R^{11})-$;

R^1 is independently at each occurrence hydrogen or
 10 C_1-C_6 alkyl;


R^2 is independently at each occurrence hydrogen, C_1-C_6 alkyl, optionally
 substituted C_3-C_8 cycloalkyl, optionally substituted (C_1-C_4) alkyl-aryl, optionally
 substituted aryl, or optionally substituted heterocycle, $C(O)$ -aryl, or $(CH_2)_2NR^4R^5$;
 15

R^3 is independently at each occurrence hydrogen, C_1-C_6 alkyl, optionally
 substituted C_3-C_8 cycloalkyl, optionally substituted (C_1-C_4) alkyl-aryl, optionally
 substituted aryl, optionally substituted heterocycle, OR^{13} , or NR^4R^6 ;

20 R^4 is independently at each occurrence hydrogen, C_1-C_6 alkyl, optionally
 substituted (C_1-C_6) alkyl-aryl, optionally substituted aryl, or R^4 and R^5 , R^6 , R^6' combine
 to form $=CR^1R^{14}$;

R^5 is independently at each occurrence hydrogen, C_1-C_6 alkyl, C_1-C_4 alkoxy,
 25 optionally substituted heterocycle, optionally substituted C_3-C_8 cycloalkyl, optionally
 substituted C_6-C_{10} bicycloalkyl, optionally substituted (C_1-C_4) alkyl-aryl, optionally
 substituted aryl, optionally substituted (C_1-C_4) alkyl-heterocycle, $C(O)C(O)R^{13}$,

- 365 -

C(O)R⁷, CH₂R⁷, SO₂R⁸, a moiety of the formula , or R⁴ and R⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

5 R⁶ is independently at each occurrence hydrogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, or R⁴ and R⁶, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R^{6'} is independently at each occurrence hydrogen, C₁-C₆ alkyl, C₁-C₄ alkoxy, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, (C₁-C₄ alkyl)-OR¹³.

wherein the (C₁-C₄ alkyl) of the (C₁-C₄ alkyl)-OR¹³ may be optionally substituted from 1 to 2 times with C₁-C₄ alkyl, optionally substituted aryl, optionally substituted heterocycle;

20 or R⁴ and R^{6'}, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R⁷ is independently at each occurrence optionally substituted C₁-C₆ alkyl, C₁-C₆ alkoxy, (C₁-C₄ alkoxy)-aryl, (C₁-C₄ alkoxy)-heterocycle, (C₁-C₄ alkoxy)-SiCH₃, optionally substituted (C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-(C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl,

- 366 -

diphenylmethyl, optionally substituted (C₁-C₄ alkyl)-CO-aryl, optionally substituted CO-aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted CH=CH-heterocycle, optionally substituted phenoxy, optionally substituted heterocycle, optionally substituted (C₁-C₄ alkyl)-phenoxy, (CH₂)_tS(O)_rR¹, (CH₂)_tC(R¹²)(R⁹)N(R¹⁶)(R¹⁵),
 5 (CH₂)_tC(R¹²)(R⁹)O(R¹⁷), (CH₂)_tC(R¹²)(R⁹)S(R¹⁷), or NR⁴R^{6'};

R⁸ is independently at each occurrence optionally substituted C₁-C₆ alkyl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, or optionally substituted heterocycle;

10

R⁹ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted heterocycle, (CH₂)_u-(C₁-C₆ alkoxy), optionally substituted (CH₂)_u-O-(C₃-C₈ cycloalkyl), optionally substituted (CH₂)_u-
 15 (C₁-C₄ alkoxy)-aryl, optionally substituted (CH₂)_u-O-aryl, optionally substituted (CH₂)_u-O-heterocycle, (C₁-C₄ alkyl)-CO₂-(C₁-C₆ alkyl), optionally substituted (C₁-C₄ alkyl)-CO₂-(C₃-C₈ cycloalkyl), optionally substituted (C₁-C₄ alkyl)-CO₂-(C₁-C₄ alkyl)-aryl, optionally substituted (C₁-C₄ alkyl)-CO₂-aryl, optionally substituted (C₁-C₄ alkyl)-CO₂-heterocycle, or R⁹ and R¹² can combine to form a C₃-C₈ cycloalkyl;

20

R¹⁰ is 0 to 4 substituents from the aryl ring independently at each occurrence hydrogen, halo, C(O)R³, cyano, optionally substituted heterocycle, optionally substituted aryl, C≡C-R¹, C₁-C₄ alkoxy, (C₁-C₄ alkyl)-phenyl, NR¹⁹R²⁰, or C₂-C₆ alkenyl;

25

R¹¹ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted heterocycle, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted aryl, or optionally substituted (C₁-C₄ alkyl)-aryl;

- 367 -

R¹² is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle or optionally substituted heterocycle;

R¹³ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, CO₂CH₂CO₂CH₂CH₃, or optionally substituted heterocycle;

R¹⁴ is independently at each occurrence C₁-C₆ alkyl or optionally substituted (C₁-C₄ alkyl)-aryl;

R¹⁵ is independently at each occurrence hydrogen, C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₆-C₁₀ bicycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted heterocycle, C(O)OR¹³, SO₂R⁸, C(O)R¹⁸, or a

moiety of the formula  ;

R¹⁶ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted aryl, optionally substituted heterocycle, or -COR⁸; or R¹⁶ and R¹⁵, together with the nitrogen to which they are attached, combine to form an optionally substituted N-heterocycle;

R¹⁷ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl,

- 368 -

optionally substituted aryl, COR¹⁸, optionally substituted heterocycle, optionally substituted (C₁-C₄ alkyl)-heterocycle, optionally substituted C₁-C₆ alkoxy, optionally substituted (C₁-C₄ alkoxy)-aryl, optionally substituted (C₁-C₄ alkoxy)-heterocycle, (C₁-C₄ alkyl)-N(R¹)(R¹), or an amino acid ester;

5

R¹⁸ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted (C₁-C₄ alkyl)-aryl, optionally substituted aryl, optionally substituted heterocycle, (C₁-C₄ alkyl)-NHCO₂-(C₁-C₄ alkyl), or optionally substituted (C₁-C₄ alkyl)-heterocycle;


10

R¹⁹ is independently at each occurrence hydrogen, or optionally substituted C₁-C₆ alkyl;


R²⁰ is independently at each occurrence hydrogen, optionally substituted C₁-C₆ alkyl, CH₂OH, CO-(C₁-C₄ alkyl); or a pharmaceutical salt thereof.

15

2. The compound of Claim 1 where het is



3. The compound of Claim 1 where het is



20

4. The compound of any one of Claims 1-3 where A is 1,3-cyclohexyl.

5. The compound of any one of Claims 1-4 where n is 0.

- 369 -

6. The compound of any one of Claims 1-5 where o is 0 or 1.
7. The compound of any one of Claims 1-6 where Y is E-NR⁴R⁵.
- 5
8. The compound of Claim 7 where R⁵ is COR⁷.
9. The compound of Claim 8 where R⁷ is optionally substituted heterocycle.
- 10
10. The compound of Claim 8 where R⁷ is optionally substituted CO-aryl.
11. The compound of Claim 8 where R⁷ is optionally substituted CO-heteroaryl.
- 15
12. The compound of Claim 8 where R⁷ is (CH₂)_t C(R¹²)(R⁹)N(R¹⁶)(R¹⁵).
13. The compound of any one of Claims 1-12 where R^b is C₁-C₆ alkyl.
14. The compound of Claim 13 where R^b is methyl.
- 20
15. The compound of any one of Claims 1-14 where R¹⁰ is halo.
16. The compound of Claim 15 where R¹⁰ is chloro.
- 25
17. The compound of Claim 16 where R¹⁰ is 9-chloro.
18. The compound of Claim 17 selected from the group consisting of N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-piperidylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-

- 370 -


c]quinolin-5-yl))cyclohexyl]-2-(2-chloro(4-pyridyloxy))acetamide, N-{[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-imidazo[5,4-c]quinolin-5-yl))cyclohexyl]methyl}benzamide, N-{[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]methyl}benzamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-hydroxy-2-phenylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-fluorophenyl)-2-hydroxyacetamide, N-{[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]methyl}-3-pyridylcarboxamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide, and N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide.

19. A method of inhibiting MRP1 in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof.

20. The method according to Claim 19 where the mammal is a human.

20

21. The method of any one of Claims 19-20 where het is



22. The method of any one of Claims 19-20 where het is



23. The method of any one of Claims 19-22 where A is 1,3-cyclohexyl.

25

24. The method of any one of Claims 19-23 where n is 0.

- 371 -

25. The method of any one of Claims 19-24 where o is 0 or 1.
26. The method of any one of Claims 19-25 where Y is E-NR⁴R⁵.
- 5 27. The method of Claim 26 where R⁵ is COR⁷.
28. The method of Claim 27 where R⁷ is optionally substituted heterocycle.
- 10 29. The method of Claim 27 where R⁷ is optionally substituted CO-aryl.
30. The method of Claim 27 where R⁷ is optionally substituted CO-heteroaryl.
31. The method of Claim 27 where R⁷ is (CH₂)_t C(R¹²)(R⁹)N(R¹⁶)(R¹⁵).
- 15 32. The method of any one of Claims 19-31 where R^b is C₁-C₆ alkyl.
33. The method of Claim 32 where R^b is methyl.
- 20 34. The method of any one of Claims 19-33 where R¹⁰ is halo.
35. The method of Claim 34 where R¹⁰ is chloro.
36. The method of Claim 35 where R¹⁰ is 9-chloro.
- 25 37. The method of Claim 36 selected from the group consisting of N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-piperidylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(2-chloro(4-pyridyloxy))acetamide, N-[(3S,1R)-3-(9-

- 372 -

chloro-3-methyl-4-oxo-5H-imidazo[5,4-c]quinolin-5-yl)cyclohexyl)methyl}benzamide, N-([(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl)methyl}benzamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-hydroxy-2-phenylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(4-fluorophenyl)-2-hydroxyacetamide, N-([(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl)methyl}-3-pyridylcarboxamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide, and N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl)cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide.


38. A method of inhibiting a resistant neoplasm, or a neoplasm susceptible to resistance, in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof; in combination with an effective amount of one or more oncolytic agents.

39. The method according to Claim 38 where the mammal is a human.


40. The method according to Claim 39 where the oncolytic(s) is selected from: camptosar, melphalan, paclitaxel, vinorelbine, mitoxantrone, doxorubicin, daunorubicin, epirubicin, vincristine, and etoposide.

41. The method according to Claim 39 where the neoplasm is of the Wilm's type, bladder, bone, breast, lung(small-cell), testis, or thyroid or the neoplasm is associated with acute lymphoblastic and myeloblastic leukemia, neuroblastoma, soft tissue sarcoma, Hodgkin's and non-Hodgkin's lymphomas, and bronchogenic carcinoma.

42. The method of any one of Claims 39-41 where het is



- 373 -



43. The method of any one of Claims 39-41 where het is
44. The method of any one of Claims 39-43 where A is 1,3-cyclohexyl.
- 5
45. The method of any one of Claims 39-44 where n is 0.
46. The method of any one of Claims 39-45 where o is 0 or 1.
- 10
47. The method of any one of Claims 39-46 where Y is E-NR⁴R⁵.
48. The method of Claim 47 where R⁵ is COR⁷.
49. The method of Claim 48 where R⁷ is optionally substituted heterocycle.
- 15
50. The method of Claim 48 where R⁷ is optionally substituted CO-aryl.
51. The method of Claim 48 where R⁷ is optionally substituted CO-heteroaryl.
- 20
52. The method of Claim 48 where R⁷ is (CH₂)_t C(R¹²)(R⁹)N(R¹⁶)(R¹⁵).
53. The method of any one of Claims 39-52 where R^b is C₁-C₆ alkyl.
54. The method of Claim 53 where R^b is methyl.
- 25
55. The method of any one of Claims 39-54 where R¹⁰ is halo.

- 374 -

56. The method of Claim 55 where R¹⁰ is chloro.

57. The method of Claim 56 where R¹⁰ is 9-chloro.

5 58. The method of Claim 57 selected from the group consisting of N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-piperidylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(2-chloro(4-pyridyloxy))acetamide, N-{[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-imidazolo[5,4-c]quinolin-5-yl))cyclohexyl]methyl}benzamide,
10 N-{[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]methyl}benzamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-hydroxy-2-phenylacetamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-fluorophenyl)-2-hydroxyacetamide, N-{[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]methyl}-3-pyridylcarboxamide, N-[(3S,1R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide, and N-[(1S,3R)-3-(9-chloro-3-methyl-4-oxo-5H-isoxazolo[4,3-c]quinolin-5-yl))cyclohexyl]-2-(4-acetylpiperazinyl)-2-phenylacetamide.

20 59. A pharmaceutical formulation comprising a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof; in combination with one or more pharmaceutical carriers, diluents, or excipients therefor.

25 60. A pharmaceutical formulation comprising:

(a) a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof;

(b) one or more oncolytic agents; and

30 (c) one or more pharmaceutical carriers, diluents, or excipients therefor.

- 375 -

61. The formulation according to Claim 60 where the oncolytic(s) is selected from: camptosar, melphalan, paclitaxel, vinorelbine, mitoxantrone, doxorubicin, daunorubicin, epirubicin, vincristine, and etoposide.

5

62. A use of a compound of formula I, as defined in Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for inhibiting a resistant neoplasm, or a neoplasm susceptible to resistance in a mammal.

10

63. A use of a compound of formula I, as defined in Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for inhibiting MRP1.

15

64. A use of a compound of formula I, as defined in Claim 1, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for inhibiting MRP1 conferred MDR in a resistant neoplasm, or a neoplasm susceptible to resistance in a mammal.

20

65. A use of a compound of formula I, as defined in Claim 1, in therapy.

66. A pharmaceutical composition for inhibiting MRP1 in a mammal which comprises an effective amount of a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof.

25

67. The composition according to Claim 66 where the mammal is a human.

30

68. A pharmaceutical composition for inhibiting a resistant neoplasm, or a neoplasm susceptible to resistance, in a mammal which comprises administering to a mammal in need thereof an effective amount of a compound of formula I, as defined in Claim 1, or a pharmaceutical salt thereof; in combination with an effective amount of one or more oncolytic agents.

- 376 -

69. The composition according to Claim 68 where the mammal is a human.

70. The composition according to Claim 69 where the oncolytic(s) is selected
from: camptosar, melphalan, paclitaxel, vinorelbine, mitoxantrone, doxorubicin,
5 daunorubicin, epirubicin, vincristine, and etoposide.

71. The composition according to Claim 69 where the neoplasm is of the
Wilm's type, bladder, bone, breast, lung(small-cell), testis, or thyroid or the neoplasm is
associated with acute lymphoblastic and myeloblastic leukemia, neuroblastoma, soft
10 tissue sarcoma, Hodgkin's and non-Hodgkin's lymphomas, and bronchogenic carcinoma.

