



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|--|
| (51) International Patent Classification⁵ : C12Q 1/68, C07H 15/12, 17/00 | A1 | (11) International Publication Number: WO 92/17484 (43) International Publication Date: 15 October 1992 (15.10.92) |
| (21) International Application Number: PCT/US92/02480 (22) International Filing Date: 26 March 1992 (26.03.92) (30) Priority data: 675,843 27 March 1991 (27.03.91) US (71) Applicant: RESEARCH CORPORATION TECHNOLOGIES, INC. [US/US]; 6840 East Broadway Boulevard, Tucson, AZ 85710 (US). (72) Inventor: KOOL, Eric, T. ; 39 Brighton Street, Apt. 1, Rochester, NY 14600 (US). (74) Agent: SCOTT, Anthony, C.; Scully, Scott, Murphy & Presser, 400 Garden City Plaza, Garden City, NY 11530 (US). | | (81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), MC (European patent), NL (European patent), NO, SE (European patent). Published <i>With international search report.</i> |
| (54) Title: SINGLE-STRANDED CIRCULAR OLIGONUCLEOTIDES | | |
| (57) Abstract <p>The present invention provides single-stranded circular oligonucleotides each with a parallel binding (P) domain and an anti-parallel binding (AP) domain separated from each other by loop domains. Each P and AP domain has sufficient complementarity to bind to one strand of a defined nucleic acid target wherein the P domain binds in a parallel manner to the target and the AP domain binds in an anti-parallel manner to the target. Moreover, the present single-stranded circular oligonucleotides can bind to both single-stranded and double-stranded target nucleic acids. The present invention also provides methods of using these oligonucleotides as well as pharmaceutical compositions containing these oligonucleotides.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AT | Austria | ES | Spain | MG | Madagascar |
| AU | Australia | FI | Finland | ML | Mali |
| BB | Barbados | FR | France | MN | Mongolia |
| BE | Belgium | GA | Gabon | MR | Mauritania |
| BF | Burkina Faso | GB | United Kingdom | MW | Malawi |
| BG | Bulgaria | GN | Guinea | NL | Netherlands |
| BJ | Benin | GR | Greece | NO | Norway |
| BR | Brazil | HU | Hungary | PL | Poland |
| CA | Canada | IT | Italy | RO | Romania |
| CF | Central African Republic | JP | Japan | RU | Russian Federation |
| CG | Congo | KP | Democratic People's Republic of Korea | SD | Sudan |
| CH | Switzerland | KR | Republic of Korea | SE | Sweden |
| CI | Côte d'Ivoire | LI | Liechtenstein | SN | Senegal |
| CM | Cameroon | LK | Sri Lanka | SU | Soviet Union |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| DE | Germany | MC | Monaco | TG | Togo |
| DK | Denmark | | | US | United States of America |

-1-

1 SINGLE-STRANDED CIRCULAR OLIGONUCLEOTIDES

The present application is a continuation-in-part of copending U.S. Serial No. 675,843 filed March 5 27, 1991. Moreover, the subject matter of the present application relates to subject matter contained in Disclosure Document number 234,794 received by the United States Patent and Trademark Office on September 5, 1989.

10 This invention was made with United States government support under grant number GM-46625 awarded by the National Institutes of Health. The United States government has certain rights in the invention.

15 FIELD OF THE INVENTION:

The present invention provides single-stranded circular oligonucleotides capable of binding to a target DNA or RNA and thereby regulating DNA replication, RNA transcription, protein translation, and other processes involving nucleic acid templates. Furthermore, circular oligonucleotides can be labeled for use as probes to detect or isolate a target nucleic acid. Circular oligonucleotides can also displace one strand of a duplex nucleic acid without prior denaturation of the 25 duplex. Moreover, circular oligonucleotides are resistant to exonucleases and bind to a target with higher selectivity and affinity than do linear oligonucleotides.

30 BACKGROUND OF THE INVENTION:

An oligonucleotide binds to a target nucleic acid by forming hydrogen bonds between bases in the

35

1 target and the oligonucleotide. Common B DNA has
conventional adenine-thymine (A-T) and guanine-cytosine
(G-C) Watson and Crick base pairs with two and three
hydrogen bonds, respectively. Conventional
5 hybridization technology is based upon the capability of
sequence-specific DNA or RNA probes to bind to a target
nucleic acid via Watson-Crick hydrogen bonds. However,
other types of hydrogen bonding patterns are known
wherein some atoms of a base which are not involved in
10 Watson-Crick base pairing can form hydrogen bonds to
another nucleotide. For example, thymine (T) can bind
to an A-T Watson-Crick base pair via hydrogen bonds to
the adenine, thereby forming a T-AT base triad.
Hoogsteen (1959, Acta Crystallography 12: 822) first
15 described the alternate hydrogen bonds present in T-AT
and C-GC base triads. More recently, G-TA base triads,
wherein guanine can hydrogen bond with a central
thymine, have been observed (Griffin et al., 1989,
Science 245: 967-971). If an oligonucleotide could bind
20 to a target with both Watson-Crick and alternate
hydrogen bonds an extremely stable complex would form
that would have a variety of in vivo and in vitro
utilities. However, to date there has been no
disclosure of an oligonucleotide with the necessary
25 structural features to achieve stable target binding
with both Watson-Crick and alternate hydrogen bonds.

Oligonucleotides have been observed to bind by
non-Watson-Crick hydrogen bonding in vitro. For
example, Cooney et al., 1988, Science 241: 456 disclose
30 a 27-base single-stranded oligonucleotide which bound to
a double-stranded nucleic acid via non-Watson-Crick
hydrogen bonds. However, triple-stranded complexes of

1 this type are not very stable, because the
oligonucleotide is bound to its target only with less
stable alternate hydrogen bonds, i.e., without any
Watson-Crick bonds.

5 Oligonucleotides have been used for a variety
of utilities. For example, oligonucleotides can be used
as probes for target nucleic acids that are immobilized
onto a filter or membrane, or are present in tissues.
Sambrook et al. (1989, Molecular Cloning: A Laboratory
10 Manual, Vols. 1-3, Cold Spring Harbor Press, NY) provide
a detailed review of hybridization techniques.

Furthermore, there has been great interest
recently in developing oligonucleotides as regulators of
cellular nucleic acid biological function. This
15 interest arises from observations on naturally occurring
complementary, or antisense, RNA used by some cells to
control protein expression. However, the development of
oligonucleotides for in vivo regulation of biological
processes has been hampered by several long-standing
20 problems, including the low binding stability and
nuclease sensitivity of linear oligonucleotides.

For example, transcription of the human c-myc
gene has been inhibited in a cell free, in vitro assay
system by a 27-base linear oligonucleotide designed to
25 bind to the c-myc promoter. Inhibition was only
observed using a carefully controlled in vitro assay
system wherein lower than physiological temperatures
were employed, and many cellular enzymes had been
removed or inactivated. These conditions were necessary
30 because linear oligonucleotides bind with low affinity
and are highly susceptible to enzymes which degrade
linear pieces of DNA (Cooney et al.). Splicing of a

35

1 pre-mRNA transcript essential for Herpes Simplex virus
replication has also been inhibited with a linear
oligonucleotide which was complementary to an acceptor
splice junction. In this instance, a methylphosphonate
5 linkage was employed in the linear oligonucleotide to
increase its nuclease resistance. Addition of this
chemically-modified oligonucleotide to the growth medium
caused reduction in protein synthesis and growth of
uninfected cells, most likely because of toxicity
10 problems at high concentrations (Smith et al., 1986,
Proc. Natl. Acad. Sci. USA 83: 2787-2791).

In another example, linear oligonucleotides
were used to inhibit human immunodeficiency virus
replication in cultured cells. Linear oligonucleotides
15 complementary to sites within or near the terminal
repeats of the retrovirus genome and within sites
complementary to certain splice junctions were most
effective in blocking viral replication. However, these
experiments required large amounts of the linear
20 oligonucleotides before an effect was obtained,
presumably because of the low binding stability and
vulnerability of these linear oligonucleotides to
nucleases (Goodchild et al., 1988, Proc. Natl. Acad.
Sci. USA 85: 5507-5511).

25 Accordingly, oligonucleotides that are useful
as regulators of biological processes preferably possess
certain properties. First, the oligonucleotide should
bind strongly enough to its complementary target nucleic
acid to have the desired regulatory effect. Second, it
30 is generally desirable that the oligonucleotide and its
target be sequence specific. Third, the oligonucleotide
should have a sufficient half-life under in vivo

1 conditions for it to be able to accomplish its desired
regulatory action in the cell. Hence, the
oligonucleotide should be resistant to enzymes that
degrade nucleic acids, e.g. nucleases. Fourth, the
5 oligonucleotide should be able to bind to single- and
double-stranded targets.

While linear oligonucleotides may satisfy the
requirement for sequence specificity, linear
oligonucleotides are sensitive to nucleases and
10 generally require chemical modification to increase
biological half-life. Such modifications increase the
cost of making an oligonucleotide and may present
toxicity problems. Furthermore, linear oligonucleotides
bind to form a two-stranded complex like those present
15 in cellular nucleic acids. Consequently, cellular
enzymes can readily manipulate and dissociate a linear
oligonucleotide bound in a double-stranded complex with
target. The low binding strength and nuclease
sensitivity of linear oligonucleotides can thus
20 necessitate administration of high concentrations of
oligonucleotide, in turn making such administration
toxic or costly. Moreover, while linear
oligonucleotides can bind to a double-stranded target
via alternate hydrogen bonds (e.g. Hoogsteen binding),
25 linear oligonucleotides cannot readily dissociate a
double-stranded target to replace one strand and thereby
form a more stable Watson-Crick bonding pattern.

Furthermore, increased binding strength
increases the effectiveness of a regulatory
30 oligonucleotide. Therefore, an oligonucleotide with
high binding affinity can be used at lower dosages.
Lower dosages decrease costs and reduce the likelihood

1 Accordingly, the present invention provides
single-stranded circular oligonucleotides which, by
nature of the circularity of the oligonucleotide and the
domains present on the oligonucleotide, are nuclease
5 resistant and bind with strong affinity and high
selectivity to their targeted nucleic acids. Moreover,
the present circular oligonucleotides can dissociate and
bind to a double-stranded target without prior
denaturation of that target.

10 Some types of single-stranded circles of DNA
or RNA are known. For example, the structures of some
naturally occurring viral and bacteriophage genomes are
single-stranded circular nucleic acids. Single-stranded
circles of DNA have been studied by Erie et al. (1987,
15 Biochemistry 26: 7150-7159 and 1989, Biochemistry 28:
268-273). However, none of these circular molecules are
designed to bind a target nucleic acid. Hence, the
present invention represents an innovation characterized
by a substantial improvement relative to the prior art
20 since the subject circular oligonucleotides exhibit high
specificity, low or no toxicity and more resistance to
nucleases than linear oligonucleotides, while binding to
single- or double-stranded target nucleic acids more
strongly than conventional linear oligonucleotides.

25 SUMMARY OF THE INVENTION:

The present invention provides a single-
stranded circular oligonucleotide having at least one
parallel binding (P) domain and at least one anti-
30 parallel binding (AP) domain, and having a loop domain
between each binding domain to form the circular
oligonucleotide. Each P and corresponding AP domain has

1 sufficient complementarity to bind detectably to one
strand of a defined nucleic acid target with the P
domain binding in a parallel manner to the target, and
the AP domain binding in an anti-parallel manner to the
5 target. Sufficient complementarity means that a
sufficient number of base pairs exists between the
target nucleic acid and the P and/or AP domains of the
circular oligonucleotide to achieve stable, i.e.
detectable, binding.

10 Another aspect of the present invention
provides the subject single-stranded circular
oligonucleotides derivatized with a reporter molecule to
provide a probe for a target nucleic acid, or with a
drug or other pharmaceutical agent to provide cell
15 specific drug delivery, or with agents which can cleave
or otherwise modify the target nucleic acid or,
furthermore, with agents that can facilitate cellular
uptake or target binding of the oligonucleotide.

An additional aspect of the present invention
20 provides single-stranded circular oligonucleotides
linked to a solid support for isolation of a nucleic
acid complementary to the oligonucleotide.

Another aspect of the present invention
provides a compartmentalized kit for detection or
25 diagnosis of a target nucleic acid including at least
one first container providing any one of the present
circular oligonucleotides.

A further aspect of the present invention
provides a method of detecting a target nucleic acid
30 which involves contacting a single-stranded circular
oligonucleotide with a sample containing the target
nucleic acid, for a time and under conditions sufficient

1 to form an oligonucleotide-target complex, and detecting
the complex. This detection method can be by
fluorescent energy transfer.

A still further aspect of the present
5 invention provides a method of regulating biosynthesis
of a DNA, an RNA or a protein. This method includes
contacting at least one of the subject circular
oligonucleotides with a nucleic acid template for the
DNA, the RNA or the protein under conditions sufficient
10 to permit binding of the oligonucleotide to a target
sequence contained in the template, followed by binding
of the oligonucleotide to the target, blocking access to
the template and thereby regulating biosynthesis of the
DNA, the RNA or the protein.

15 An additional aspect of the present invention
provides pharmaceutical compositions for regulating
biosynthesis of a nucleic acid or protein containing a
biosynthesis regulating amount of at least one of the
subject circular oligonucleotides and a pharmaceutically
20 acceptable carrier.

A further aspect of the present invention
provides a method of preparing a single-stranded
circular oligonucleotide which includes binding a linear
precircle to an end-joining-oligonucleotide, joining the
25 two ends of the precircle and recovering the circular
oligonucleotide product.

Another aspect of the present invention
provides a method of strand displacement in a double-
stranded nucleic acid target by contacting the target
30 with any one of the present circular oligonucleotides
for a time and under conditions effective to denature
the target and to bind the circular oligonucleotide.

35

1 BRIEF DESCRIPTIONS OF THE DRAWINGS:

Fig. 1A depicts the bonding patterns of Watson-Crick (anti-parallel domain) AT and GC base pairs. Fig. 1B depicts T-AT, C+GC and G-TA base triads 5 that can form between P, target and AP nucleotides.

Fig. 2 schematically illustrates a circularization reaction for synthesis of single-stranded circular oligonucleotides. A linear precircle oligonucleotide is bound to an oligonucleotide having 10 the same sequence as the target, i.e. an end-joining-oligonucleotide, to form a precircle complex. After ligation, the circularized oligonucleotides are separated from the end-joining-oligonucleotide.

Fig. 3 depicts the sequence of linear 15 precursors to circular oligonucleotides, i.e. precircles (1-3 having SEQ ID NO: 5, SEQ ID NO: 6 and SEQ ID NO: 7), targets (4,5 having SEQ ID NO: 8 and SEQ ID NO: 9), circular oligonucleotides (6,7,8 and 13 having SEQ ID NO: 5-7 and 14), and linear oligonucleotides (9-12 and 20 14 having SEQ ID NO: 10-13 and 15) described in the Examples.

Fig. 4 depicts the structure of a linear precircle complexed with an end-joining-oligonucleotide before ligation.

25 Fig. 5 depicts the effect of pH on circular oligonucleotide:target complex formation as measured by T_m . Filled circles represent the stability at different pH values for a 6:4 complex while filled squares depict the stability of a 7:5 complex. The sequences of 30 circular oligonucleotides 6 and 7 and targets 4 and 5 are presented in Fig. 3.

1 Fig. 6A depicts the effect of loop size on
complex formation, with a comparison between binding to
two targets: a simple (dA)₁₂ target (squares) and a 36
nucleotide oligonucleotide target (circles). Fig. 6B
5 depicts the effect of target and binding domain length
on complex formation.

Fig. 7 depicts a complex formed between a
circular oligonucleotide and a target where the P and AP
binding domains are staggered on the target.

10 Fig. 8 depicts replacement of one strand of a
fluorecently labeled double stranded target (SEQ ID NO:
11) by either a linear oligonucleotide having SEQ ID NO:
8 (dotted line) or a circular oligonucleotide having SEQ
ID NO: 5 (solid line). Strand replacement was measured
15 by an increase in fluorescein fluorescence intensity (Y-
axis) as a function of time (X-axis).

Fig. 9 depicts a plot of observed pseudo-first
order rate constant, K_{obs} for duplex target (SEQ ID NO:
5) at several concentrations. Uncertainty in rate
20 constants are no more than $\pm 10\%$. The depicted curve is
a rectangular hyperbola generated as a best fit. A
double reciprocal plot of the data, i.e., [circular
oligonucleotide]⁻¹ vs (K_{obs})⁻¹ is linear with a slope of
 $8.95 \times 10^{-6} \text{ sec} \cdot \text{M}^{-1}$ and a y-intercept of 39.8 sec.

25 DETAILED DESCRIPTION OF THE INVENTION:

The present invention relates to single-
stranded circular oligonucleotides, i.e. circles, which
can bind to nucleic acid targets with higher affinity
and selectivity than a corresponding linear
30 oligonucleotide. Moreover, since the present circles
can open up two strands of a double-stranded nucleic


1 acid and bind thereto, both single- and double-stranded
nucleic acids can be targets for binding by the present
circular oligonucleotides.

Furthermore, the strong, selective binding of
5 these circles to either single- or double-stranded
targets provides a variety of uses, including methods of
regulating such biological processes as DNA replication,
RNA transcription, RNA splicing and processing, protein
translation and the like. Similarly, the ability of
10 these circles to dissociate double-stranded nucleic
acids and to selectively and stably bind to targeted
nucleic acids makes them ideal as diagnostic probes or
as markers to localize, for example, specific sites in a
chromosome or other DNA or RNA molecules. Additionally,
15 the present circles are useful for isolation of
complementary nucleic acids or for sequence-specific
delivery of drugs or other molecules into cells.

In particular, the single-stranded circular
oligonucleotides of the present invention have at least
20 one parallel binding (P) domain and at least one anti-
parallel binding (AP) domain and have a loop domain
between each binding domain, so that a circular
oligonucleotide is formed. Moreover, each P and AP
domain exhibits sufficient complementarity to bind to
25 one strand of a defined nucleic acid target with the P
domain binding to the target in a parallel manner and
the AP domain binding to the target in an anti-parallel
manner.

The schematic illustration set forth below
30 shows the circular arrangement of one set of P and AP
oligonucleotide domains relative to each other as well
as when bound to a target (T, as indicated below).

-12-



The arrows indicate the 5' to 3' orientation of each strand with the 5' end of each domain at the tail and the 3' end at the arrowhead. Hence as used herein binding of nucleic acids in a parallel manner means that the 5' to 3' orientation is the same for each strand or nucleotide in the complex. This is the type of binding present between the target and the P domain. As used herein, binding of nucleic acids in an anti-parallel manner means that the 5' to 3' orientations of two strands or nucleotides in a complex lie in opposite directions, i.e. the strands are aligned as found in the typical Watson-Crick base pairing arrangement of double helical DNA.

When more than one P or AP binding domain is present, such binding domains are separated from other P and AP domains by loop domains whose lengths are sufficient to permit binding to multiple targets. Moreover, when a circle has multiple AP and P domains, the corresponding targets need not be linked on one nucleic acid strand. Furthermore, a loop domain of a circular oligonucleotide bound to a given target can be an AP or P domain for binding to a second target when the circular oligonucleotide releases from the first target.

35

1 In accordance with this invention, the
nucleotide sequences of the P and AP domains can be
determined from the defined sequence of the nucleic acid
target by reference to the base pairing rules provided
5 hereinbelow. A target can be either single- or double-
stranded and is selected by its known functional and
structural characteristics. For example, some preferred
targets can be coding regions, origins of replication,
reverse transcriptase binding sites, transcription
10 regulatory elements, RNA splicing junctions, or ribosome
binding sites, among others. A target can also be
selected by its capability for detection or isolation of
a DNA or RNA template. Preferred targets are rich in
purines, i.e. in adenines and guanines.

15 The nucleotide sequence of the target DNA or
RNA can be known in full or in part. When the target
nucleotide sequence is completely known the sequences of
the P and AP domains are designed with the necessary
degree of complementarity to achieve binding, as
20 detected by known procedures, for example by a change in
light absorption or fluorescence. In some instances,
the target sequence can be represented by a consensus
sequence or be only partially known. For example,
circular oligonucleotides (circles) which bind to an
25 entire class of targets represented by a consensus
sequence can be provided by designing the P and AP
domains from the target consensus sequence. In this
instance some of the targets may match the consensus
sequence exactly and others may have a few mismatched
30 bases, but not enough mismatch to prevent binding.
Likewise, if a portion of a target sequence is known,
one skilled in the art can refer to the base pairing

1 rules provided hereinbelow to design circles which bind
to that target with higher affinity than a linear
oligonucleotide that has a sequence corresponding to
that of the circle.

5 Thus, the present invention is also directed
to circles having P and AP domains which are
sufficiently complementary to bind to a nucleic acid
target wherein a sufficient number, but not necessarily
all, nucleotide positions in the P and AP domains are
10 determined from the target sequence in accordance with
the base pairing rules of this invention. The number of
determined (i.e. known) positions is that number of
positions which are necessary to provide sufficient
complementarity for binding of the subject
15 oligonucleotides to their targets, as detected by
standard procedures including a change in light
absorption upon binding or melting.

The base pairing rules of the present
invention provide for the P domain to bind to the target
20 by forming base pairs wherein the P domain and target
nucleotides have the same 5' to 3' orientation. In
particular, these rules are satisfied to the extent
needed to achieve binding of a circular oligonucleotide
to its nucleic acid target, i.e. the degree of
25 complementarity need not be 100% so long as binding can
be detected. Hence, the general rules for determining
the sequence of the P domain are thus:

when a base for a position in the target is
guanine or a guanine analog, then P has cytosine, or a
30 suitable analog thereof, in a corresponding position;

when a base for a position in the target is
adenine, or an adenine analog then P has thymine or

1 uracil, or suitable analogs thereof, in a corresponding
position;

when a base for a position in the target is
thymine, or a thymine analog, then P has cytosine or
5 guanine, or suitable analogs thereof, in a corresponding
position;

when a base for a position in the target is
cytosine, or a cytosine analog, then P has cytosine,
thymine or uracil, or suitable analogs thereof, in a
10 corresponding position; and

when a base for a position in the target is
uracil, or a uracil analog, then P has cytosine,
guanine, thymine, or uracil, or suitable analogs
thereof, in a corresponding position.

15 The base pairing rules of the present
invention provide for the AP domain to bind to the
target by forming base pairs wherein the AP domain and
target nucleotides are oriented in opposite directions.
In particular these rules are satisfied to the extent
20 necessary to achieve detectable binding of a circular
oligonucleotide to its nucleic acid target, i.e. the
degree of complementarity can be less than 100%. Hence,
the base pairing rules can be adhered to only insofar as
is necessary to achieve sufficient complementarity for
25 binding to be detected between the circular
oligonucleotide and its target.

Thus, the general rules for determining the
sequence of the AP domain are as follows:

when a base for a position in the target is
30 guanine, or a guanine analog, then AP has cytosine or
uracil, or suitable analogs thereof, in a corresponding
position;

1 when a base for a position in the target is adenine, or an adenine analog, then AP has thymine or uracil, or suitable analogs thereof, in a corresponding position;

5 when a base for a position in the target is thymine, or a thymine analog, then AP has adenine, or a suitable analog thereof, in a corresponding position; and

10 when a base for a position in the target is cytosine, or a cytosine analog, then AP has a guanine, or a suitable analog thereof, in corresponding position;

15 when a base for a position in the target is uracil, or a uracil analog, then AP has adenine or guanine, or suitable analogs thereof, in a corresponding position.

In a preferred embodiment, the P, AP and loop domains are not complementary to each other.

20 Table 1 summarizes which nucleotides can form anti-parallel base pairs or parallel base pairs with a defined target nucleotide.

Table 1

| Target Nucleotide ^a | Anti-Parallel Domain Nucleotide ^a | Parallel Domain Nucleotide ^a |
|--------------------------------|--|---|
| G | C or U | C |
| A | T or U | T or U |
| T | A | C or G |
| C | G | C, T or U |
| U | A or G | C, G, T or U |

30 ^a Or a suitable analog

35

1 Two complementary single-stranded nucleic
acids form a stable double helix (duplex) when the
strands bind, or hybridize, to each other in the typical
Watson-Crick fashion, i.e. via anti-parallel GC and AT
5 base pairs. For the present invention, stable duplex
formation and stable triplex formation is achieved when
the P and AP domains exhibit sufficient complementarity
to the target sequence to achieve stable binding between
the circular oligonucleotide and the target molecule.
10 Stable binding occurs when an oligonucleotide remains
detectably bound to target under the required
conditions.

 Complementarity between nucleic acids is the
degree to which the bases in one nucleic acid strand can
15 hydrogen bond, or base pair, with the bases in a second
nucleic acid strand. Hence, complementarity can
sometimes be conveniently described by the percentage,
i.e. proportion, of nucleotides which form base pairs
between two strands or within a specific region or
20 domain of two strands. For the present invention
sufficient complementarity means that a sufficient
number of base pairs exist between a target nucleic acid
and the P and/or AP domains of the circular
oligonucleotide to achieve detectable binding.
25 Moreover, the degree of complementarity between the P
domain and the target and the AP domain and the target
need not be the same. When expressed or measured by
percentage of base pairs formed, the degree of
complementarity can range from as little as about 30-40%
30 complementarity to full, i.e. 100%, complementarity. In
general, the overall degree of complementarity between
the P or AP domain and the target is preferably at least

35

1 about 50%. However, the P domain can sometimes have
less complementarity with the target than the AP domain
has with the target, for example the P domain can have
about 30% complementarity with the target while the AP
5 domain can have substantially more complementarity, e.g.
50% to 100% complementarity.

Moreover, the degree of complementarity that
provides detectable binding between the subject circular
oligonucleotides and their respective targets, is
10 dependent upon the conditions under which that binding
occurs. It is well known that binding, i.e.
hybridization, between nucleic acid strands depends on
factors besides the degree of mismatch between two
sequences. Such factors include the GC content of the
15 region, temperature, ionic strength, the presence of
formamide and types of counter ions present. The effect
that these conditions have upon binding is known to one
skilled in the art. Furthermore, conditions are
frequently determined by the circumstances of use. For
20 example, when a circular oligonucleotide is made for use
in vivo, no formamide will be present and the ionic
strength, types of counter ions, and temperature
correspond to physiological conditions. Binding
conditions can be manipulated in vitro to optimize the
25 utility of the present oligonucleotides. A thorough
treatment of the qualitative and quantitative
considerations involved in establishing binding
conditions that allow one skilled in the art to design
appropriate oligonucleotides for use under the desired
30 conditions is provided by Beltz et al., 1983, Methods
Enzymol. 100: 266-285 and by Sambrook et al.

1 Thus for the present invention, one of
ordinary skill in the art can readily design a
nucleotide sequence for the P and AP domains of the
subject circular oligonucleotides which exhibits
5 sufficient complementarity to detectably bind to its
target sequence. As used herein "binding" or "stable
binding" means that a sufficient amount of the
oligonucleotide is bound or hybridized to its target to
permit detection of that binding. Binding can be
10 detected by either physical or functional properties of
the target:circular oligonucleotide complex.

Binding between a target and an
oligonucleotide can be detected by any procedure known
to one skilled in the art, including both functional or
15 physical binding assays. Binding may be detected
functionally by determining whether binding has an
observable effect upon a biosynthetic process such as
DNA replication, RNA transcription, protein translation
and the like.

20 Physical methods of detecting the binding of
complementary strands of DNA or RNA are well known in
the art, and include such methods as DNase I or chemical
footprinting, gel shift and affinity cleavage assays and
light absorption detection procedures. For example, a
25 method which is widely used, because it is so simple and
reliable, involves observing a change in light
absorption of a solution containing an oligonucleotide
and a target nucleic acid at 220 to 300 nm as the
temperature is slowly increased. If the oligonucleotide
30 has bound to its target, there is a sudden increase in
absorption at a characteristic temperature as the
oligonucleotide and target dissociate or melt.

35

1 The binding between an oligonucleotide and its
target nucleic acid is frequently characterized by the
temperature at which 50% of the oligonucleotide is
melted from its target. This temperature is the melting
5 temperature (T_m). A higher T_m means a stronger or more
stable complex relative to a complex with a lower T_m .
The stability of a duplex increases with increasing G:C
content since G:C base pairs have three hydrogen bonds
whereas A:T base pairs have two. The circular
10 oligonucleotides of the present invention provide
additional hydrogen bonds and hence more stability since
two binding domains are available for bonding to a
single target nucleic acid, i.e. the P domain and the AP
domain. Hence, the triplex formed by a circular
15 oligonucleotide bound to its target nucleic acid should
melt at a higher T_m than the duplex formed by a linear
oligonucleotide and a target.

Circular oligonucleotides bind to a nucleic
acid target through hydrogen bonds formed between the
20 nucleotides of the binding domains and the target. The
AP domain can bind by forming Watson-Crick hydrogen
bonds (Fig. 1). The P domain can bind to the target
nucleotides by forming non-Watson-Crick hydrogen bonds
(e.g., Fig. 1 and Table I). When two nucleotides from
25 different strands of DNA or RNA hydrogen bond by the
base pairing rules defined herein, a base pair or duplex
is formed. When a nucleotide from AP and a nucleotide
from P both bind to the same target nucleotide, a base
triad is formed.

30 Parallel domain base pairing with a
complementary target strand of nucleic acid, is
thermodynamically less favorable than Watson-Crick base

1 pairing; however, when both parallel and antiparallel
pairing modes are present in a single molecule, highly
stable complexes can form. Thus, two opposing domains
of a circular oligomer form a complex with a central
5 target, giving a triplex structure, or a triple helical
complex, bounded by the two looped ends of the circle.
For example, this arrangement can allow formation of up
to four hydrogen bonds when two thymines bind to a
target adenine and up to five hydrogen bonds when two
10 cytosines bind to a target guanine.

Furthermore, because of the binding
characteristics of the P and AP domains, the present
circular oligonucleotides have a higher selectivity for
a particular target than do corresponding linear
15 oligonucleotides. At least two factors can contribute
to this high selectivity. First, circular
oligonucleotides of this invention bind twice to the
same central target strand. Hence two domains are
involved in selecting a target. Second, protonation of
20 cytosine in a C+G-C triad is favored only when this
triad forms and the additional proton gives the triad a
positive charge. This positive charge can lessen the
negative charge repulsions arising from the
juxtapositioning of three phosphodiester backbones.

25 Unlike linear oligonucleotides, the present
circular oligonucleotides can displace one strand of a
double-stranded target under conditions where
denaturation of the double-stranded target is
thermodynamically unfavorable. Linear oligonucleotides
30 do not have this capacity to displace a strand of a
duplex. For example, the half-life of a double-stranded
target in the presence of a complementary linear

1 oligonucleotide is about 58 min i.e. so long that the
linear oligonucleotide has little utility for displacing
one strand of the duplex target. However, a double-
stranded target has a half-life of only 30 sec in the
5 presence of the present circular oligonucleotides.
Therefore, the circular oligonucleotides of the present
invention have utility not only for binding single-
stranded targets, but also for binding to double-
stranded targets. Accordingly, since both single- and
10 double-stranded nucleic acids are available as targets
for the present circular oligonucleotides, these
circular oligonucleotides can have greater utility than
linear oligonucleotides. For example, the present
circular oligonucleotides are better regulators of
15 biological processes in vivo and better in vitro
diagnostic probes than corresponding linear
oligonucleotides.

When the nucleic acid template extends beyond
the central triple-stranded target:circle complex, a P
20 or an AP domain may bind as duplex on either side of the
triple standard complex. Hence a target:circular
oligonucleotide complex can be partially two stranded
and partially three-stranded, wherein two-stranded
portions can be P:target duplexes, without bound AP
25 nucleotides, or AP:target duplexes, without bound P
nucleotides. This binding arrangement is a staggered
binding arrangement.

Each P domain, AP domain and target can
independently have about 2 to about 200 nucleotides with
30 preferred lengths being about 4 to about 100
nucleotides. The most preferred lengths are 6 to 36
nucleotides.

1 The P and AP domains are separated by loop
domains which can independently have from about 2 to
about 2000 nucleotides. A preferred loop length is from
about 3 to about 8 nucleotides with an especially
5 preferred length being about 5 nucleotides.

 According to the present invention, the loop
domains do not have to be composed of nucleotide bases.
Non-nucleotide loops can make the present circular
oligonucleotides cheaper to produce. More
10 significantly, circular oligonucleotides with non-
nucleotide loops are more resistant to nucleases and
therefore have a longer biological half-life than
linear oligonucleotides. Furthermore, loops having no
charge, or a positive charge, can be used to promote
15 binding by eliminating negative charge repulsions
between the loop and target. In addition, circular
oligonucleotides having uncharged or hydrophobic non-
nucleotide loops can penetrate cellular membranes better
than circular oligonucleotides with nucleotide loops.

20 As contemplated herein, non-nucleotide loop
domains can be composed of alkyl chains, polyethylene
glycol or oligoethylene glycol chains or other chains
providing the necessary steric or flexibility properties
which are compatible with oligonucleotide synthesis.
25 The length of these chains is equivalent to about 2 to
about 2000 nucleotides, with preferred lengths
equivalent to about 3 to about 8 nucleotides. The most
preferred length for these chains is equivalent to about
5 nucleotides.

30 Preferred chains for non-nucleotide loop
domains are polyethylene glycol or oligoethylene glycol
chains. In particular, oligoethylene glycol chains

35

1 having a length similar to a 5 nucleotide chain, e.g. a
pentaethylene glycol, a hexaethylene glycol or a
heptaethylene glycol chain, are preferred.

The circular oligonucleotides are single-
5 stranded DNA or RNA, with the bases guanine (G), adenine
(A), thymine (T), cytosine (C) or uracil (U) in the
nucleotides, or with any nucleotide analog that is
capable of hydrogen bonding in a parallel or anti-
parallel manner. Nucleotide analogs include
10 pseudocytidine, isopseudocytidine, 3-aminophenyl-
imidazole, 2'-O-methyl-adenosine, 7-deazadenosine, 7-
deazaguanosine, 4-acetylcytidine, 5-(carboxy-
hydroxymethyl)-uridine, 2'-O-methylcytidine, 5-
carboxymethylaminomethyl-2-thioridine, 5-
15 carboxymethylamino-methyluridine, dihydrouridine, 2'-O-
methyluridine, 2'-O-methyl-pseudouridine, beta,D-
galactosylqueosine, 2'-O-methylguanosine, inosine, N6-
isopentenyladenosine, 1-methyladenosine, 1-methyl-
pseudouridine, 1-methylguanosine, 1-methylinosine, 2,2-
20 dimethylguanosine, 2-methyladenosine, 2-methylguanosine,
3-methylcytidine, 5-methylcytidine, 5-methyluridine, N6-
methyl-adenosine, 7-methylguanosine, 5-methylamino-
methyluridine, 5-methoxyaminomethyl-2-thiouridine, beta-D-
mannosylqueosine, 5-methoxycarbonylmethyluridine, 5-
25 methoxyuridine, 2-methyl-thio-N6-isopentenyladenosine,
N-(9-beta-D-ribofuranosyl-2-methylthiopurine-6-yl)-
carbamoyl)threonine, N-(9-beta-D-ribofuranosylpurine-6-
yl)-N-methylcarbamoyl)threonine. When possible, either
ribose or deoxyribose sugars can be used with these
30 analogs. Nucleotides bases in an alpha-anomeric
conformation can also be used in the circular
oligonucleotides of the present invention.

1 Preferred nucleotide analogs are unmodified G,
A, T, C and U nucleotides; pyrimidine analogs with lower
alkyl, lower alkoxy, lower alkylamine, phenyl or lower
alkyl substituted phenyl groups in the 5 position of the
5 base and purine analogs with similar groups in the 7 or
8 position of the base. Especially preferred nucleotide
analogues are 5-methylcytosine, 5-methyluracil,
diaminopurine, and nucleotides with a 2'-O-methylribose
moiety in place of ribose or deoxyribose. As used
10 herein lower alkyl, lower alkoxy and lower alkylamine
contain from 1 to 6 carbon atoms and can be straight
chain or branched. These groups include methyl, ethyl,
propyl, isopropyl, butyl, isobutyl, tertiary butyl,
amyl, hexyl and the like. A preferred alkyl group is
15 methyl.

Circular oligonucleotides can be made first as
linear oligonucleotides and then circularized. Linear
oligonucleotides can be made by any of a myriad of
procedures known for making DNA or RNA oligonucleotides.
20 For example, such procedures include enzymatic synthesis
and chemical synthesis.

Enzymatic methods of DNA oligonucleotide
synthesis frequently employ Klenow, T7, T4, Tag or E.
coli DNA polymerases as described in Sambrook et al.
25 Enzymatic methods of RNA oligonucleotide synthesis
frequently employ SP6, T3 or T7 RNA polymerase as
described in Sambrook et al. Reverse transcriptase can
also be used to synthesize DNA from RNA (Sambrook
et al.). To prepare oligonucleotides enzymatically
30 requires a template nucleic acid which can either be
synthesized chemically, or be obtained as mRNA, genomic
DNA, cloned genomic DNA, cloned cDNA or other

1 recombinant DNA. Some enzymatic methods of DNA
oligonucleotide synthesis can require an additional
primer oligonucleotide which can be synthesized
chemically. Finally, linear oligonucleotides can be
5 prepared by PCR techniques as described, for example, by
Saiki et al., 1988, Science 239:487.

Chemical synthesis of linear oligonucleotides
is well known in the art and can be achieved by solution
or solid phase techniques. Moreover, linear
10 oligonucleotides of defined sequence can be purchased
commercially or can be made by any of several different
synthetic procedures including the phosphoramidite,
phosphite triester, H-phosphonate and phosphotriester
methods, typically by automated synthesis methods. The
15 synthesis method selected can depend on the length of
the desired oligonucleotide and such choice is within
the skill of the ordinary artisan. For example, the
phosphoramidite and phosphite triester method produce
oligonucleotides having 175 or more nucleotides while
20 the H-phosphonate method works well for oligonucleotides
of less than 100 nucleotides. If modified bases are
incorporated into the oligonucleotide, and particularly
if modified phosphodiester linkages are used, then the
synthetic procedures are altered as needed according to
25 known procedures. In this regard, Uhlmann et al. (1990,
Chemical Reviews 90: 543-584) provide references and
outline procedures for making oligonucleotides with
modified bases and modified phosphodiester linkages.

Synthetic, linear oligonucleotides may be
30 purified by polyacrylamide gel electrophoresis, or by
any of a number of chromatographic methods, including
gel chromatography and high pressure liquid

1 chromatography. To confirm a nucleotide sequence,
oligonucleotides may be subjected to DNA sequencing by
any of the known procedures, including Maxam and Gilbert
sequencing, Sanger sequencing, capillary electrophoresis
5 sequencing the wandering spot sequencing procedure or by
using selective chemical degradation of oligonucleotides
bound to Hybond paper. Sequences of short
oligonucleotides can also be analyzed by plasma
desorption mass spectroscopy or by fast atom bombardment
10 (McNeal, et al., 1982, J. Am. Chem. Soc. 104: 976;
Viari, et al., 1987, Biomed. Environ. Mass Spectrom. 14:
83; Grotjahn et al., 1982, Nuc. Acid Res. 10: 4671).
Sequencing methods are also available for RNA
oligonucleotides.

15 The present invention provides several methods
of preparing circular oligonucleotides from linear
precursors (i.e. precircles), including a method wherein
a precircle is synthesized and bound to an end-joining-
oligonucleotide and the two ends of the precircle are
20 joined. Any method of joining two ends of an
oligonucleotide is contemplated by the present
invention, including chemical methods employing, for
example, known coupling agents like BrCN, N-
cyanoimidazole $ZnCl_2$, 1-ethyl-3-(3-
25 dimethylaminopropyl)carbodiimide and other carbodimides
and carbonyl diimidazoles. Furthermore, the ends of a
precircle can be joined by condensing a 5' phosphate and
a 3' hydroxy, or a 5' hydroxy and a 3' phosphate.

In accordance with the present invention, a
30 simple one-step chemical method is provided to construct
the subject circular oligonucleotides, or circles, from
precircles. An oligonucleotide is constructed which has

1 the same sequence as the target nucleic acid; this is
the end-joining oligonucleotide. A DNA or RNA linear
precircle is chemically or enzymatically synthesized and
phosphorylated on its 5' or 3' end, again by either
5 chemical or enzymatic means. The precircle and the end-
joining oligonucleotide are mixed and annealed, thereby
forming a complex in which the 5' and 3' ends of the
precircle are adjacent, as depicted in Fig. 2. It is
preferred that the ends of the precircle fall within a
10 binding domain, not within a loop, and preferably within
the anti-parallel binding domain rather than the
parallel domain. Moreover, it is preferred that a
precircle have a 3'-phosphate rather than a 5'-
phosphate. After complex formation, the ends undergo a
15 condensation reaction in a buffered aqueous solution
containing divalent metal ions and BrCN at about pH 7.0.
In a preferred embodiment the buffer is imidazole-Cl at
pH 7.0 with a divalent metal such as Ni, Zn, Mn, or Co.
Ni is the most preferred divalent metal. Condensation
20 occurs after about 6-48 hr. of incubation at 4-37°C.
Other divalent metals, such as Cu, Pb, Ca and Mg, can
also be used.

One method for RNA circularization
incorporates the appropriate nucleotide sequences,
25 preferably in a loop domain, into an RNA oligonucleotide
to promote self splicing, since a circular product is
formed under the appropriate conditions (Sugimoto
et al., 1988, *Biochemistry*: 27: 6384-6392).

Enzymatic circle closure is also possible
30 using DNA ligase or RNA ligase under conditions
appropriate for these enzymes.

1 Circular oligonucleotides can be separated
from the template by denaturing gel electrophoresis or
melting followed by gel electrophoresis, size selective
chromatography, or other appropriate chromatographic or
5 electrophoretic methods. The recovered circular
oligonucleotide can be further purified by standard
techniques as needed for its use in the methods of the
present invention.

The present invention also contemplates
10 derivatization or chemical modification of the subject
oligonucleotides with chemical groups to facilitate
cellular uptake. For example, covalent linkage of a
cholesterol moiety to an oligonucleotide can improve
cellular uptake by 5- to 10- fold which in turn improves
15 DNA binding by about 10- fold (Boutorin et al., 1989,
FEBS Letters 254: 129-132). Other ligands for cellular
receptors may also have utility for improving cellular
uptake, including, e.g. insulin, transferrin and others.
Similarly, derivatization of oligonucleotides with poly-
20 L-lysine can aid oligonucleotide uptake by cells
(Schell, 1974, Biochem. Biophys. Acta 340: 323, and
Lemaitre et al., 1987, Proc. Natl. Acad. Sci. USA 84:
648). Certain protein carriers can also facilitate
cellular uptake of oligonucleotides, including, for
25 example, serum albumin, nuclear proteins possessing
signals for transport to the nucleus, and viral or
bacterial proteins capable of cell membrane penetration.
Therefore, protein carriers are useful when associated
with or linked to the circular oligonucleotides of this
30 invention. Accordingly, the present invention
contemplates derivatization of the subject circular
oligonucleotides with groups capable of facilitating

1 cellular uptake, including hydrocarbons and non-polar
groups, cholesterol, poly-L-lysine and proteins, as well
as other aryl or steroid groups and polycations having
analogous beneficial effects, such as phenyl or naphthyl
5 groups, quinoline, anthracene or phenanthracene groups,
fatty acids, fatty alcohols and sesquiterpenes,
diterpenes and steroids.

The present invention further contemplates
derivatization of the subject oligonucleotides with
10 agents that can cleave or modify the target nucleic acid
or other nucleic acid strands associated with or in the
vicinity of the target. For example, viral DNA or RNA
can be targeted for destruction without harming cellular
nucleic acids by administering a circular
15 oligonucleotide complementary to the targeted nucleic
acid which is linked to an agent that, upon binding, can
cut or render the viral DNA or RNA inactive. Nucleic
acid destroying agents that are contemplated by the
present invention as having cleavage or modifying
20 activities include, for example, RNA and DNA nucleases,
ribozymes that can cleave RNA, azidoproflavine,
acridine, EDTA/Fe, chloroethylamine, azidophenacyl and
phenanthroline/Cu. Uhlmann et al. (1990, Chemical
Reviews 90: 543-584) provide further information on the
25 use of such agents and methods of derivatizing
oligonucleotides that can be adapted for use with the
subject circular oligonucleotides.

Derivatization of the subject circular
oligonucleotides with groups that facilitate cellular
30 uptake or target binding, as well as derivatization with
nucleic acid destroying agents or drugs, can be done by
any of the procedures known to one skilled in the art.

1 Moreover, the desired groups can be added to nucleotides
before synthesis of the oligonucleotide. For example,
these groups can be linked to the 5-position of T or C
and these modified T and C nucleotides can be used for
5 synthesis of the present circular oligonucleotides. In
addition, derivatization of selected nucleotides permits
incorporation of the group into selected domains of the
circular oligonucleotide. For example, in some
instances it is preferable to incorporate certain groups
10 into a loop where that group will not interfere with
binding, or into an AP or P domain to facilitate
cleavage or modification of the target nucleic acid.

In accordance with the present invention,
modification in the phosphodiester backbone of circular
15 oligonucleotides is also contemplated. Such
modifications can aid uptake of the oligonucleotide by
cells or can extend the biological half-life of such
nucleotides. For example, circular oligonucleotides may
penetrate the cell membrane more readily if the negative
20 charge on the internucleotide phosphate is eliminated.
This can be done by replacing the negatively charged
phosphate oxygen with a methyl group, an amine or by
changing the phosphodiester linkage into a
phosphotriester linkage by addition of an alkyl group to
25 the negatively charged phosphate oxygen. Alternatively,
one or more of the phosphate atoms which is part of the
normal phosphodiester linkage can be replaced. For
example, NH-P, CH₂-P or S-P linkages can be formed.
Accordingly, the present invention contemplates using
30 methylphosphonates, phosphorothioates,
phosphorodithioates, phosphotriesters and phosphorus-
boron (Sood et al., 1990, J. Am. Chem. Soc. 112: 9000)

1 linkages. The phosphodiester group can be replaced with
siloxane, carbonate, acetamidate or thioether groups.
These modifications can also increase the resistance of
the subject oligonucleotides to nucleases. Methods for
5 synthesis of oligonucleotides with modified
phosphodiester linkages are reviewed by Uhlmann et al.

Circular oligonucleotides with non-nucleotide
loops can be prepared by any known procedure. For
example, Durand *et al.* (1990, *Nucleic Acids Res.* 18:
10 6353-6359) provides synthetic procedures for linking
non-nucleotide chains to DNA. Such procedures can
generally be adapted to permit an automated synthesis of
a linear oligonucleotide precursor which is then used to
make a circular oligonucleotide of the present
15 invention. In general, groups reactive with nucleotides
in standard DNA synthesis, e.g. phosphoramidite, H-
phosphonate, dimethoxytrityl, monomethoxytrityl and the
like, can be placed at the ends of non-nucleotide chains
and nucleotides corresponding to the ends of P and AP
20 domains can be linked thereto.

Additionally, different nucleotide sugars can
be incorporated into the oligonucleotides of this
invention. For example, RNA oligonucleotides can be
used since RNA:DNA hybrids are more stable than DNA:DNA
25 hybrids. Additional binding stability can also be
provided by using 2'-O-methyl ribose in the present
circular oligonucleotides. Phosphoramidite chemistry
can be used to synthesize RNA oligonucleotides as
described (Reese, C. B. In Nucleic Acids & Molecular
30 Biology; Springer-Verlag: Berlin, 1989; Vol. 3, p. 164;
and Rao, et al., 1987, *Tetrahedron Lett.* 28: 4897).

1 The synthesis of RNA 2'-O-methyl-
oligoribonucleotides and DNA oligonucleotides differ
only slightly. RNA 2'-O-methyloligonucleotides can be
prepared with minor modifications of the amidite, H-
5 phosphonate or phosphotriester methods (Shibahara et al,
1987, Nucleic Acids Res. 15: 4403; Shibahara et al.,
1989, Nucleic Acids Res. 17: 239; Anoue et al., 1987,
Nucleic Acids Res. 15: 6131).

In another embodiment the present invention,
10 circular oligonucleotides can accelerate the
dissociation of a double-stranded nucleic acid target.
Therefore the double-stranded nucleic acid target does
not have to be subjected to denaturing conditions before
binding of the present circular oligonucleotides. Thus,
15 the circular oligonucleotides can bind to both single-
and double-stranded nucleic acid targets under a wider
variety of conditions, and particularly under
physiological conditions. The present circular
oligonucleotides are several orders of magnitude faster
20 at accelerating duplex nucleic acid strand displacement
than are the corresponding linear oligonucleotides.

The present invention therefore provides a
means to displace one strand of a double-stranded
nucleic acid target with one of the subject circular
25 oligonucleotides without the necessity of prior
denaturation of the double-stranded nucleic acid target.
Thus, the present invention provides a method of strand
displacement in a double-stranded nucleic acid target by
contacting the target with one of the subject circular
30 oligonucleotides for a time and under conditions
effective to denature said target and permit the
circular oligonucleotide to bind to the target. The

1 target for the present circular oligonucleotides can be
a double-stranded nucleic acid, either RNA or DNA, which
has not undergone denaturation by, for example, heating
or exposure to alkaline pH.

5 As used herein, the nucleic acids for strand
displacement can be present in an organism or present in
a sample which includes an impure or pure nucleic acid
preparation, a tissue section, a prokaryotic or
eukaryotic cell smear, a chromosomal squash and the
10 like. Moreover, the nucleic acid targets for strand
displacement by the present circular oligonucleotides
include viral, bacterial, fungal or mammalian nucleic
acids.

According to the present invention, conditions
15 effective to denature the target by strand displacement
and thereby permit binding, include having a suitable
circular oligonucleotide to target nucleic acid ratio.
Moreover, as used herein a suitable ratio of circular
oligonucleotide to target is about 1 to about 100, and
20 is preferably about 1 to about 50.

Moreover, as used herein a time effective to
denature a double-stranded nucleic acid by strand-
displacement with an oligonucleotide of the present
invention is about 1 minute to about 16 hours.

25 A circular oligonucleotide can associate with
a duplex target by first binding in the P domain. Such
P domain binding juxtaposes the AP domain nucleotides to
compete for Watson-Crick binding to target nucleotide.
This P domain pre-association followed by AP domain
30 nucleotide competition for Watson-Crick binding may form
the basis for the observed acceleration in strand
displacement by circular oligonucleotides.

1 In summary, the subject circular
oligonucleotides have three important features which
enable duplex strand displacement. First, the circular
oligonucleotide has the ability to preassociate, which
5 results in a high local concentration. Second, the
circular oligonucleotide contains a second (AP) binding
domain, which competes for binding to a complementary
strand of the duplex. Finally, the circular
oligonucleotide binds with higher affinity than the
10 displaced strand of the duplex, thereby driving the
reaction to completion.

The present invention contemplates a variety
of utilities for the subject circular oligonucleotides
which are made possible by their selective and stable
15 binding properties with both single- and double-stranded
targets. Some utilities include, but are not limited
to: use of circular oligonucleotides of defined
sequence, bound to a solid support, for affinity
isolation of complementary nucleic acids; use of the
20 subject oligonucleotides to provide sequence specific
stop signals during polymerase chain reaction (PCR);
covalent attachment of a drug, drug analog or other
therapeutic agent to circular oligonucleotides to allow
cell type specific drug delivery; labeling circular
25 oligonucleotides with a detectable reporter molecule for
localizing, quantitating or identifying complementary
target nucleic acids; and binding circular
oligonucleotides to a cellular or viral nucleic acid
template and regulating biosynthesis directed by that
30 template.

The subject circular oligonucleotides can be
attached to a solid support such as silica, cellulose,

1 nylon, and other natural or synthetic materials that are
used to make beads, filters, and column chromatography
resins. Attachment procedures for nucleic acids to
solid supports of these types are well known; any known
5 attachment procedure is contemplated by the present
invention. A circular oligonucleotide attached to a
solid support can then be used to isolate a
complementary nucleic acid. Isolation of the
complementary nucleic acid can be done by incorporating
10 the oligonucleotide:solid support into a column for
chromatographic procedures. Other isolation methods can
be done without incorporation of the
oligonucleotide:solid support into a column, e.g. by
utilization of filtration procedures. Circular
15 oligonucleotide:solid supports can be used, for example,
to isolate poly(A)⁺ mRNA from total cellular or viral
RNA by making a circular oligonucleotide with P and AP
domain poly(dT) or poly(U) sequences. Circular
oligonucleotides are ideally suited to applications of
20 this type because they are nuclease resistant and bind
target nucleic acids so strongly.

Further utilities are available for the
subject oligonucleotides in the field of polymerase
chain reaction (PCR) technology. PCR technology
25 provides methods of synthesizing a double-standard DNA
fragment encoded in a nucleic acid template between two
known nucleic acid sequences which are employed as
primer binding sites. In some instances it is desirable
to produce a single-stranded DNA fragment before or
30 after having made some of the double stranded fragment.
This can be done by, for example, binding a circular
oligonucleotide of the present invention to one of the

1 primer binding sites or to a site lying between the
primer binding sites.

The present invention also contemplates using
the subject circular oligonucleotides for targeting
5 drugs to specific cell types. Such targeting can allow
selective destruction or enhancement of particular cell
types, e.g. inhibition of tumor cell growth can be
attained. Different cell types express different genes,
so that the concentration of a particular mRNA can be
10 greater in one cell type relative to another cell type,
such an mRNA is a target mRNA for cell type specific
drug delivery by circular oligonucleotides linked to
drugs or drug analogs. Cells with high concentrations
of target mRNA are targeted for drug delivery by
15 administering to the cell a circular oligonucleotide
with a covalently linked drug that is complementary to
the target mRNA.

The present invention also contemplates
labeling the subject circular oligonucleotides for use
20 as probes to detect a target nucleic acid. Labelled
circular oligonucleotide probes have utility in
diagnostic and analytical hybridization procedures for
localizing, quantitating or detecting a target nucleic
acid in tissues, chromosomes or in mixtures of nucleic
25 acids. Circular oligonucleotide probes of this invention
represent a substantial improvement over linear nucleic
acid probes because the circular oligonucleotides can
replace one strand of a double-stranded nucleic acid,
and because the present oligonucleotides have two
30 binding domains which not only provide increased binding
stability but also impart a greater sequence selectivity

1 (or specificity) for the target:oligonucleotide
interaction.

Labeling of a circular oligonucleotide can be
done by incorporating nucleotides linked to a "reporter
5 molecule" into the subject circular oligonucleotides. A
"reporter molecule", as defined herein, is a molecule or
atom which, by its chemical nature, provides an
identifiable signal allowing detection of the circular
oligonucleotide. Detection can be either qualitative or
10 quantitative. The present invention contemplates using
any commonly used reporter molecule including
radionuclides, enzymes, biotins, psoralens,
fluorophores, chelated heavy metals, and luciferin. The
most commonly used reporter molecules are either
15 enzymes, fluorophores or radionuclides linked to the
nucleotides which are used in circular oligonucleotide
synthesis. Commonly used enzymes include horseradish
peroxidase, alkaline phosphatase, glucose oxidase and β -
galactosidase, among others. The substrates to be used
20 with the specific enzymes are generally chosen because a
detectably colored product is formed by the enzyme
acting upon the substrate. For example, p-nitrophenyl
phosphate is suitable for use with alkaline phosphatase
conjugates; for horseradish peroxidase, 1,2-
25 phenylenediamine, 5-aminosalicylic acid or toluidine
are commonly used. The probes so generated have utility
in the detection of a specific DNA or RNA target in, for
example, Southern analysis, Northern analysis, in situ
hybridization to tissue sections or chromosomal squashes
30 and other analytical and diagnostic procedures. The
methods of using such hybridization probes are well

1 known and some examples of such methodology are provided
by Sambrook et al.

The present circular oligonucleotides can be
used in conjunction with any known detection or
5 diagnostic procedure which is based upon hybridization
of a probe to a target nucleic acid. Moreover, the
present circular oligonucleotides can be used in any
hybridization procedure which quantitates a target
nucleic acid, e.g., by competitive hybridization between
10 a target nucleic acid present in a sample and a labeled
tracer target for one of the present oligonucleotides.
Furthermore, the reagents needed for making a circular
oligonucleotide probe and for utilizing such a probe in
a hybridization procedure can be marketed in a kit.

15 The kit can be compartmentalized for ease of
utility and can contain at least one first container
providing reagents for making a precircle precursor for
a circular oligonucleotide, at least one second
container providing reagents for labeling the precircle
20 with a reporter molecule, at least one third container
providing reagents for circularizing the precircle, and
at least one fourth container providing reagents for
isolating the labeled circular oligonucleotide.

Moreover the present invention provides a kit
25 for isolation of a template nucleic acid. Such a kit
has at least one first container providing a circular
oligonucleotide which is complementary to a target
contained within the template. For example, the
template nucleic acid can be cellular and/or viral
30 poly(A)⁺ mRNA and the target can be the poly(A)⁺ tail.
Hence circular oligonucleotides of the present invention

1 which have utility for isolation of poly(A)+ mRNA have P
and AP domain sequences of poly(dT) or poly(U).

Furthermore, the present invention provides
kits useful when diagnosis of a disease depends upon
5 detection of a specific, known target nucleic acid.
Such nucleic acid targets can be, for example, a viral
nucleic acid, an extra or missing chromosome or gene, a
mutant cellular gene or chromosome, an aberrantly
expressed RNA and others. The kits can be
10 compartmentalized to contain at least one first
container providing a circular oligonucleotide linked to
a reporter molecule and at least one second container
providing reagents for detection of the reporter
molecule.

15 One aspect of the present invention provides a
method of regulating biosynthesis of a DNA, an RNA or a
protein by contacting at least one of the subject
circular oligonucleotides with a nucleic acid template
for that DNA, that RNA or that protein in an amount and
20 under conditions sufficient to permit the binding of the
oligonucleotide(s) to a target sequence contained in the
template. The binding between the oligonucleotide(s)
and the target blocks access to the template, and
thereby regulates biosynthesis of the nucleic acid or
25 the protein. Blocking access to the template prevents
proteins and nucleic acids involved in the biosynthetic
process from binding to the template, from moving along
the template, or from recognizing signals encoded within
the template. Alternatively, when the template is RNA,
30 regulation can be accomplished by allowing selective
degradation of the template. For example, RNA templates
bound by the subject circular oligonucleotides are

1 susceptible to degradation by RNase H and RNase H
degradation of a selected RNA template can thereby
regulate use of the template in biosynthetic processes.

As used herein, biosynthesis of a nucleic acid
5 or a protein includes cellular and viral processes such
as DNA replication, DNA reverse transcription, RNA
transcription, RNA splicing, RNA polyadenylation, RNA
translocation and protein translation, and of which can
lead to production of DNA, RNA or protein, and involve a
10 nucleic acid template at some stage of the biosynthetic
process.

As used herein, regulating biosynthesis
includes inhibiting, stopping, increasing, accelerating
or delaying biosynthesis. Regulation may be direct or
15 indirect, i.e. biosynthesis of a DNA, RNA or protein may
be regulated directly by binding a circular
oligonucleotide to the template for that DNA, RNA or
protein; alternatively, biosynthesis may be regulated
indirectly by oligonucleotide binding to a second
20 template encoding a protein that plays a role in
regulating the biosynthesis of the first DNA, RNA or
protein.

The nucleic acid templates can be RNA or DNA
and can be single-stranded or double-stranded. While
25 the present circular oligonucleotides bind to only one
strand of a target present in the template, double-
stranded templates are opened during biosynthetic
processes and thereby become available for binding.
Furthermore, the P domain of the present circular
30 oligonucleotides can bind to a double-stranded target
and place AP domain nucleotides in a position to compete
for Watson-Crick binding to target nucleotides.

1 DNA replication from a DNA template is
mediated by proteins which bind to an origin of
replication where they open the DNA and initiate DNA
synthesis along the DNA template. To inhibit DNA
5 replication in accordance with the present invention,
circular oligonucleotides are selected which bind to one
or more targets in an origin of replication. Such
binding blocks template access to proteins involved in
DNA replication. Therefore initiation and procession of
10 DNA replication is inhibited. As an alternative method
of inhibiting DNA replication, expression of the
proteins which mediate DNA replication can be inhibited
at, for example, the transcriptional or translational
level.

15 DNA replication from an RNA template is
mediated by reverse transcriptase binding to a region of
RNA also bound by a nucleic acid primer. To inhibit DNA
replication from an RNA template, reverse transcriptase
or primer binding can be blocked by binding a circular
20 oligonucleotide to the primer binding site, and thereby
blocking access to that site. Moreover, inhibition of
DNA replication can occur by binding a circular
oligonucleotide to a site residing in the RNA template
since such binding can block access to that site and to
25 downstream sites, i.e. sites on the 3' side of the
target or binding site.

To initiate RNA transcription, RNA polymerase
recognizes and binds to specific start sequences, or
promoters, on a DNA template. Binding of RNA polymerase
30 opens the DNA template. There are also additional
transcriptional regulatory elements that play a role in
transcription and are located on the DNA template.

1 These transcriptional regulatory elements include
enhancer sequences, upstream activating sequences,
repressor binding sites and others. All such promoter
and transcriptional regulatory elements, singly or in
5 combination, are targets for the subject circular
oligonucleotides. Oligonucleotide binding to these
sites can block RNA polymerase and transcription factors
from gaining access to the template and thereby
regulating, e.g., increasing or decreasing, the
10 production of RNA, especially mRNA and tRNA.
Additionally, the subject oligonucleotides can be
targeted to the coding region or 3'-untranslated region
of the DNA template to cause premature termination of
transcription. One skilled in the art can readily
15 design oligonucleotides for the above target sequences
from the known sequence of these regulatory elements,
from coding region sequences, and from consensus
sequences.

RNA transcription can be increased by, for
20 example, binding a circular oligonucleotide to a
negative transcriptional regulatory element or by
inhibiting biosynthesis of a protein that can repress
transcription. Negative transcriptional regulatory
elements include repressor sites or operator sites,
25 wherein a repressor protein binds and blocks
transcription. Oligonucleotide binding to repressor or
operator sites can block access of repressor proteins to
their binding sites and thereby increase transcription.

The primary RNA transcript made in eukaryotic
30 cells, or pre-mRNA, is subject to a number of
maturation processes before being translocated into the
cytoplasm for protein translation. In the nucleus,

1 introns are removed from the pre-mRNA in splicing
reactions. The 5' end of the mRNA is modified to form
the 5' cap structure, thereby stabilizing the mRNA.
Various bases are also altered. The polyadenylation of
5 the mRNA at the 3' end is thought to be linked with
export from the nucleus. The subject circular
oligonucleotides can be used to block any of these
processes.

A pre-mRNA template is spliced in the nucleus
10 by ribonucleoproteins which bind to splice junctions and
intron branch point sequences in the pre-mRNA.
Consensus sequences for 5' and 3' splice junctions and
for the intron branch point are known. For example,
inhibition of ribonucleoprotein binding to the splice
15 junctions or inhibition of covalent linkage of the 5'
end of the intron to the intron branch point can block
splicing. Maturation of a pre-mRNA template can,
therefore, be blocked by preventing access to these
sites, i.e. by binding circular oligonucleotides of this
20 invention to a 5' splice junction, an intron branch
point or a 3' splice junction. Splicing of a specific
pre-mRNA template can be inhibited by using circular
oligonucleotides with sequences that are complementary
to the specific pre-mRNA splice junction(s) or intron
25 branch point. In a further embodiment, a collection of
related splicing of pre-mRNA templates can be inhibited
by using a mixture of circular oligonucleotides having a
variety of sequences that, taken together, are
complementary to the desired group of splice junction
30 and intron branch point sequences.

Polyadenylation involves recognition and
cleavage of a pre-mRNA by a specific RNA endonuclease at

1 specific polyadenylation sites, followed by addition of
a poly(A) tail onto the 3' end of the pre-mRNA. Hence,
any of these steps can be inhibited by binding the
subject oligonucleotides to the appropriate site.

5 RNA translocation from the nucleus to the
cytoplasm of eukaryotic cells appears to require a
poly(A) tail. Thus, a circular oligonucleotide is
designed in accordance with this invention to bind to
the poly(A) tail and thereby block access to the poly
10 (A) tail and inhibit RNA translocation. For such an
oligonucleotide, both the P and AP domains can consist
of about 10 to about 50 thymine residues, and preferably
about 20 residues. Especially preferred P and AP domain
lengths for such an oligonucleotide are about 6 to about
15 12 thymine residues.

Protein biosynthesis begins with the binding
of ribosomes to an mRNA template, followed by initiation
and elongation of the amino acid chain via translational
"reading" of the mRNA. Protein biosynthesis, or
20 translation, can thus be blocked or inhibited by
blocking access to the template using the subject
circular oligonucleotides to bind to targets in the
template mRNA. Such targets contemplated by this
invention include the ribosome binding site (Shine-
25 Delgarno sequence), the 5' mRNA cap site, the initiation
codon, and sites in the protein coding sequence. There
are also classes of protein which share domains of
nucleotide sequence homology. Thus, inhibition of
protein biosynthesis for such a class can be
30 accomplished by targeting the homologous protein domains
(via the coding sequence) with the subject circular
oligonucleotides.

1 Regulation of biosynthesis by any of the
aforementioned procedures has utility for many
applications. For example, genetic disorders can be
corrected by inhibiting the production of mutant or
5 over-produced proteins, or by increasing production of
an under-expressed proteins; the expression of genes
encoding factors that regulate cell proliferation can be
inhibited to control the spread of cancer; and virally
encoded functions can be inhibited to combat viral
10 infection.

Some types of genetic disorders that can be
treated by the circular oligonucleotides of the present
invention include Alzheimer's disease, some types of
arthritis, sickle cell anemia and others. Many types of
15 viral infections can be treated by utilizing the
circular oligonucleotides of the present invention,
including infections caused by influenza, rhinovirus,
HIV, herpes simplex, papilloma virus, cytomegalovirus,
Epstein-Barr virus, adenovirus, vesticular stomatitis
20 virus, rotavirus and respiratory syncytial virus among
others. According to the present invention, animal and
plant viral infections may also be treated by
administering the subject oligonucleotides.

The c-myc gene is one example of a gene which
25 can have a role in cell proliferation. Inhibition of c-
myc expression has been demonstrated in vitro using a
linear oligonucleotide complementary to a target 115 bp
upstream of the c-myc transcription start site (Cooney
et al., 1988, Science 241: 456-459). Circular
30 oligonucleotides of SEQ ID NO:1, and SEQ ID NO:2, as
depicted below, are complementary to the c-myc promoter
at nucleotides -131 to -120 and -75 to -62,

1 respectively, and are provided to inhibit c-myc
 expression in accordance with the present invention. As
 used in these depictions of SEQ ID NO:1 and SEQ ID NO:2,
 N can be any nucleotide or nucleotide analog.

5 SEQ ID NO:1

```

          1
        N C T C C C C G C C C T C N
       N
      N
     N
    N C T C C C C A C C C T C N
  
```

10

SEQ ID NO:2

```

          1
        N T C T T T T T C T T T C N
       N
      N
     N
    N T C T T T T T C T T T C N
  
```


15

Human immunodeficiency virus (HIV) is a
 20 retrovirus causing acquired immunodeficiency syndrome
 (AIDS). The circular oligonucleotides of this invention
 provide a means of blocking the replication of the virus
 without deleteriously affecting normal cellular
 replication in humans infected with HIV. The retroviral
 25 genome is transcribed as a single, long transcript, part
 of which is spliced to yield RNA encoding viral envelope
 proteins. Inhibition of HIV infection can be
 accomplished by designing oligonucleotides to bind to a
 number of regions within the HIV genome, including
 30 coding regions for functions that replicate the genome
 (i.e., the pol or reverse transcriptase function) or
 functions that control gene expression (e.g. the tat,

35

1 rev or other functions). However, previous work with
 linear oligonucleotides has suggested that splice sites,
 poly(A) addition signals, cap or initiator codon sites,
 and sites implicated in ribosome assembly can be
 5 particularly effective for inhibiting eucaryotic protein
 expression. Furthermore, the terminal structures of the
 retroviral genome are also excellent targets for
 inhibiting retrovirus production not only because these
 structures encode control regions which mediate the rate
 10 of transcription and replication, but also because these
 structures are repeated, allowing an oligonucleotide to
 bind and block access to each repeat.

Accordingly, the present invention provides
 two circular oligonucleotides, set forth in SEQ ID NO:3
 15 and SEQ ID NO:4 wherein N is any nucleotide or
 nucleotide analog and Y is a pyrimidine or a pyrimidine
 analog. SEQ ID NO:3 is complementary to an HIV-1 splice
 junction (nucleotides 6039-52), while SEQ ID NO:4 is
 complementary to part of the tat gene (nucleotides 5974-
 20 88). The circular form of SEQ ID NO:3 is depicted
 below, wherein nucleotide number 1 is the first
 nucleotide in the P domain, i.e., the first T on the top
 line corresponds to base 1.



The circular form of SEQ ID NO:4 is depicted below
 30 wherein nucleotide number 1 is the first nucleotide of
 the P domain.

35

1 antigens can be monitored to assess the effect of
circular oligonucleotides on HIV replication and thereby
to follow the course of the disease state.

Alternatively, human volunteers with AIDS or ARC can be
5 administered with the subject circular oligonucleotides
since the oligonucleotides do not appear to be
cytotoxic. The disease status of these volunteers can
then be assessed to determine the efficacy of the
subject oligonucleotides in treating and preventing AIDS
10 infection.

A further aspect of this invention provides
pharmaceutical compositions containing the subject
circular oligonucleotides with a pharmaceutically
acceptable carrier. In particular, the subject
15 oligonucleotides are provided in a therapeutically
effective amount of about 0.1 μg to about 100 mg per kg
of body weight per day, and preferably of about 0.1 μg
to about 10 mg per kg of body weight per day, to bind to
a nucleic acid in accordance with the methods of this
20 invention. Dosages can be readily determined by one of
ordinary skill in the art and formulated into the
subject pharmaceutical compositions.

As used herein, "pharmaceutically acceptable
carrier" includes any and all solvents, dispersion
25 media, coatings, antibacterial and antifungal agents,
isotonic and absorption delaying agents, and the like.
The use of such media and agents for pharmaceutical
active substances is well known in the art. Except
insofar as any conventional media or agent is
30 incompatible with the active ingredient, its use in the
therapeutic compositions is contemplated. Supplementary

1 active ingredients can also be incorporated into the
compositions.

The subject oligonucleotides may be
administered topically or parenterally by, for example,
5 intravenous, intramuscular, intraperitoneal
subcutaneous or intradermal route, or when suitably
protected, the subject oligonucleotides may be orally
administered. The subject oligonucleotides may be
incorporated into a cream, solution or suspension for
10 topical administration. For oral administration,
oligonucleotides may be protected by enclosure in a
gelatin capsule. Oligonucleotides may be incorporated
into liposomes or liposomes modified with polyethylene
glycol for parenteral administration. Incorporation of
15 additional substances into the liposome, for example,
antibodies reactive against membrane proteins found on
specific target cells, can help target the
oligonucleotides to specific cell types.

Topical administration and parenteral
20 administration in a liposomal carrier is preferred.

The following examples further illustrate the
invention.

25

30

35

1

EXAMPLE 1Circularization of Oligonucleotides Using an
End Joining Oligonucleotide

5 According to the present invention, a simple one-step chemical method has been developed to construct circles from linear precursors (precircles). A DNA oligonucleotide was constructed which had the same sequence as the eventual target, this is the end-
10 joining-oligonucleotide. A precircle oligonucleotide was then constructed and chemically phosphorylated on the 5'-end or 3'-end. As depicted in Fig. 2, the precircle and end-joining-oligonucleotide were mixed and allowed to form a complex in which the ends were
15 adjacent. Cyanogen bromide, imidazole buffer, and a divalent metal were added. After incubation for 6-48 hr, the mixture was dialyzed, lyophilized, and the products were separated by denaturing 20% polyacrylamide gel electrophoresis. UV shadowing revealed major bands
20 which comigrated with the precircle and the end-joining-oligonucleotide, along with one new product which migrated slightly more slowly than the precircle. No product was observed without added end-joining-oligonucleotide or in the absence of a 5'- or 3'-
25 phosphate group on the precircle. The major bands were excised and eluted from the gel, dialyzed to remove salts and quantitated by absorbance at 260 nm. For reactions with precircles 1 and 2 (SEQ ID NO: 5 and SEQ ID NO: 6, respectively), using end-joining-
30 oligonucleotides 4 and 5 (SEQ ID NO: 8 and SEQ ID NO: 9, respectively), the circles 6 and 7 were obtained in 40% and 58% yields, respectively. The sequences of each of

35

1 these molecules and other oligonucleotides are depicted
in Fig. 3.

The circular structure of products 6 and 7 was
confirmed by resistance to 3' exonuclease digestion and
5 to 5' dephosphorylation under reaction conditions in
which a linear precircle was completely destroyed or
dephosphorylated. Accordingly, the 3' exonuclease
activity of T4 DNA polymerase cleaved linear precircles
1 and 2, but not circles 6 and 7. The linear precircles
10 were also 5'-end labeled with ^{32}P and then circularized.
After reaction, the circular products were inert to calf
alkaline phosphatase whereas the precircles completely
released labeled ^{32}P . The slightly slower gel mobility
of the circles relative to the precircles was consistent
15 with the occurrence of circularization.

Optimal Circularization Conditions

Many parameters were optimized to increase
yields of the circular product, including
oligonucleotide and precircle concentrations,
20 temperature, reaction time, metal, metal concentration,
BrCN concentration and pH. Improved circularization
conditions provided an at least two-fold higher yield of
circles compared to prior art conditions wherein two
single-stranded oligonucleotides were joined (Luebke
25 et al., 1989, J. Am. Chem. Soc. 111: 8733 and Kanaya
et al., 1986, Biochemistry 25: 7423).

30

35

1 These improved conditions were:

- 50 μM precircle
- 55 μM end-joining-oligonucleotide
- 100 mM NiCl_2
- 5 200 mM imidazole HCl (pH 7.0)
- 125 mM BrCN
- 25°C, 36 hr.

However circle closure was also effective under the following conditions:

- 10 3-200 μM precircle
- 3-200 μM end-joining-oligonucleotide
- 10-500 mM NiCl_2
- 50-500 mM imidazole-HCl
- 20-200 mM BrCN
- 15 4-37°C, 6-48 hr.
- Other metals (Zn^{2+} , Mn^{2+} , Co^{2+} , Cu^{2+} , Pb^{2+} , Ca^{2+} , Mg^{2+}) also work in place of Ni^{2+} . Additionally, the reaction is pH sensitive.

Closure in AP and P Domains

- 20 Closure of a circle in the AP domain was superior to closure in the P domain. Comparison of the circularization of precircles 2 and 3 (SEQ ID NO: 6 and SEQ ID NO: 7, respectively) around the same end-joining-oligonucleotide (i.e. 5, SEQ ID NO: 9) indicated that
- 25 circle 7 (having SEQ ID NO: 6) was formed with a 58% yield when closed in the AP domain (i.e. using precircle 2) and only a 35% yield when closed in the P domain (i.e. using precircle 3).

Condensing Reagents

- 30 Two reagents have been commonly used for chemical ligation of DNA and RNA, BrCN/imidazole/ NiCl_2 and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC)

1 (Kanaya et al. 1986 Biochemistry 25: 7423 and Ashley et
al. 1991 Biochemistry 30: 2927). Therefore, these
reagents were directly compared for efficacy in
ligating a precircle to circular oligonucleotide 6 (Fig.
5 3 and SEQ ID NO: 5) using a dA₁₂ (SEQ ID NO: 8) end-
joining-oligonucleotide.

BrCN/imidazole/NiCl₂ was used under the
established optimal conditions except that ligation
efficiency was observed at both 4°C and 25°C. EDC was
10 used at 200 mM with 20 mM MgCl₂, 50 mM MES (pH 6.0) at
4°C or 25°C with incubation for 4 days.

At 4°C BrCN was more efficient, yielding 95%
circular product while EDC yielded only 55% product.
However, at 25°C both EDC and BrCN yielded 95% product.
15 Therefore, BrCN is more effective at lower temperatures
but either EDC or BrCN can be used with equal success at
25°C. However, BrCN has an additional advantage over
EDC since ligation with BrCN requires 24 hr or less
while ligation with EDC requires about 4 days.


20 Use of a 5'- or 3'-Phosphate

Under different ligation conditions joining a
3'-phosphate with a 5'-OH yielded more ligated product
than joining a 5'-phosphate with a 3'-OH (Ashley, et
al.).

25 Therefore, the percent conversion to circular
oligonucleotide 6 (Fig. 3) by a 5'-phosphate or by a 3'-
phosphate precircles was compared:

30

35



15 Circularization reactions were performed using
 a dA₁₂ end-joining-oligonucleotide and the established
 optimal conditions, except that 5 nmoles of precircle
 and end-joining-oligonucleotide were used. Products
 were visualized under UV light after separation by
 20 denaturing gel electrophoresis.

Conversion to circular product was 60% (+5%)
 when a 5'-phosphate was present and 95% when a 3'-
 phosphate was present. No increase in yield was
 observed when increased reaction times or increased
 25 reagent concentrations were used.

Accordingly, use of a 3'-phosphate rather than
 a 5'-phosphate improves circularization.

30

35

1

EXAMPLE 2Circular Oligonucleotides Bind Target
Nucleic Acids with Higher Affinity
Than Do Linear Oligonucleotides

5 The binding affinities of circles 6 and 7 (SEQ
ID NO: 5 and SEQ ID NO: 6, respectively) for their
targets were measured by comparison of the melting
temperatures of the circular and linear complexes.
Solutions contained 1:1 ratios of oligonucleotide and
10 target (3 μM each) in 100 mM NaCl, 10 mM MgCl_2 , and 10
mM Tris-HCl (pH 7.0). Mixing curves measured at 260 nm
confirmed that 1:1 complexes were formed. The free
energies ($-\Delta G^\circ_{37}$) of the complexes were derived from
the melting data using a two-state curve-fitting method
15 (Petersheim, et al., 1983, *Biochemistry* 22: 256).

The results showed that the circular
oligonucleotides bound to their targets more strongly
than did linear precircles or Watson-Crick complementary
target-sized oligonucleotides (Table 2). For example,
20 target 4 (SEQ ID NO: 8) formed a duplex with its target-
sized Watson-Crick complement having a T_m of 37.1°C
while the precircle 1:target 4 complex (i.e. SEQ ID NO:
5 bound to SEQ ID NO: 8) had a T_m of 44.7°C. By
comparison, circle 6, having the same sequence as
precircle 1, bound to target 4 with a T_m of 57.5°C and a
25 free energy of binding that was 8.6 kcal/mol more
favorable than the corresponding Watson-Crick duplex.
The corresponding association constant at 37°C is $6 \times$
 10^{11} M^{-1} , which is more than six orders of magnitude
greater than for the Watson-Crick duplex. A similar
30 effect was observed for the binding of circle 7 (SEQ ID
NO: 6) to target 5 (SEQ ID NO: 9); this complex had a T_m

35

1 of 62.3°C, whereas the corresponding Watson-Crick duplex
melted at 43.8°C. These data indicate that the binding
of circular oligonucleotides is stronger than the
binding of a linear oligonucleotide to a target.

5 To determine the binding characteristics when
the target sequence was embedded within a longer
sequence, a 36 nucleotide oligonucleotide was
synthesized with a 12 base target sequence (equivalent
to target 4) in the middle. Melting studies revealed
10 that circle 6 bound to this longer oligonucleotide more
strongly than it did to a target having the same size as
the binding domains of the circle: the T_m of circle 6
with target 4 was 59.8°C whereas with the 36 base
oligonucleotide containing an embedded target the T_m was
15 63.4°C. Therefore the binding strength of circles with
embedded targets was higher than that with binding-
domain-sized-targets.

The binding affinity of circle 6 for an RNA
target was tested by synthesizing oligoribonucleotide
20 rA_{12} and determining the T_m of circle 6 with rA_{12} . The
 T_m of circle 6 with rA_{12} was 58.3°C compared with 57.8°C
with dA_{12} . The data indicate that circles bind to RNA
targets as strongly or more strongly than as to DNA
targets.

25

30

35

1

TABLE II

| oligonucleotide: target | complex | T _m , °C | - G° ₃₇ (kcal/mol) |
|----------------------------|--|---------------------|-------------------------------|
| | 3'-TTTTTTTTTTTTT 5'-AAAAAAAAAAAAA | 37.1 | 8.1 |
| 10 | 3'-TTCTTTTCTTTC 5'-AAGAAAAGAAAG | 43.8 | 10.3 |
| 15 | <pre> TTTTTTTTTTTTT C C A A C AAAAAAAAAAA C A A C C TTTTTT TTTTTT OPO₃⁻ </pre> | 44.7 | 10.5 |
| 20 | <pre> TTCTTTTCTTTC C C A A C AAGAAAAGAAAG C A A C C TTCTTT TCTTTC OPO₃⁻ </pre> | 47.0 | 10.8 |
| 25 | <pre> TTTTTTTTTTTTT C C A A C AAAAAAAAAAA C A A C C TTTTTT TTTTTT </pre> | 57.4 | 16.7 |
| 30 | <pre> TTCTTTTCTTTC C C A A C AAGAAAAGAAAG C A A C C TTCTTTTCTTTC </pre> | 62.3 | 16.4 |
| 35 | | | |

-60-

1

EXAMPLE 3
Circular Oligonucleotides Bind Target More
Selectively Than Linear Oligonucleotides

5 In order to measure the sequence selectivity of
circular oligonucleotides, a set of target
oligonucleotides with one variable base was constructed.
Binding energies for a circle complexed with these
10 targets were measured; the selectivity was defined by the
free energy difference between the correct sequence and
mismatched sequences. The selectivity obtained with the
circular structure was then directly compared to the
selectivity of an analogous linear oligonucleotide.

 DNA oligonucleotides were machine synthesized
15 using the β -cyanoethyl phosphoramidite method. Circular
oligonucleotide 8 was prepared from a linear precircle
having SEQ ID NO: 7:

 5'-pTCTTTCCACACCTTTCTTTTCTTCACACTTCTTT
and was cyclized by assembly around an end-joining
20 oligonucleotide having the sequence 5'-AAGAAAAGAAAG (SEQ
ID NO: 9) using BrCN/imidazole to close the final bond,
as described in Example 1. The circular structure was
confirmed by its resistance to a 3'-exonuclease and 5'-
phosphatase.

25 The sequence selectivity of circle 8 was
measured by hybridizing it with targets which contained a
single mismatched base and determining the strength
(ΔG°_{37}) of the resulting complexes by thermal
denaturation. Eight targets were synthesized which were
30 complementary to circle 8 and linear oligonucleotide 9
except for a single centrally positioned variable base (X

35

-61-

1

or Y = A, G, C, T). Four targets have a variable base X which is matched with two opposing T's in the circle, resulting in a T-X-T triad. In the remaining four

5 targets, the variable base Y is matched with two opposing C's in the circle, giving a C-Y-C triad. For comparison to this circle complex, a linear oligonucleotide 9 was used; resulting in a duplex with a central T-X pair in the first four experiments or a C-Y pair in the remaining
10 four.

| | <u>complex (X,Y = A,T,G,C)</u> | <u>expt. no.</u> |
|----|---------------------------------|------------------|
| | 3' - T T C T T T T C T T T C | |
| | 5' - A A G A X A A G A A A G | 1-4 |
| 15 | A C T T C T T T T C T T T C C A | |
| | C A A G A X A A G A A A G C | 5-8 |
| | A C T T C T T T T C T T T C C A | |
| 20 | 3' - T T C T T T T C T T T C | |
| | 5' - A A G A A A A Y A A A G | 9-12 |
| | A C T T C T T T T C T T T C C A | |
| 25 | C A A G A A A A Y A A A G C | 13-16 |
| | A C T T C T T T T C T T T C C A | |

30 Thermal denaturation of the sixteen complexes was carried out in the presence of 10 mM MgCl₂, 100 mM NaCl, and 10 mM Tris•HCl (pH 7.0), with target and circular or linear oligonucleotide concentrations at 3 μM

35

-62-

1

each. Experiments were carried out in duplicate and the results averaged. Oligonucleotide:target complex melting was monitored at 260 nm. The temperature vs. absorbance curves so generated showed a single transition from bound to free oligonucleotide. Free energies of association were obtained by fitting the data with a two-state curve-fitting method. The results were checked in two cases by measuring the association energies by the van't Hoff method, good agreement was seen between the two methods. Selectivities are defined as the difference in free energies (ΔG) of complexation between matched and mismatched oligomers.

Table III displays the results of the mismatch experiments. Experiments 1-4 show the effects of a T-X target mismatch on a DNA duplex. As expected, the true match ($X = A$) gives the most favorable complex ($-\Delta G^\circ_{37} = 10.3$ kcal/mol); the mismatches ($X = G, C, T$) result in a loss of 3.2-4.4 kcal/mol in binding energy, in good agreement with published mismatch studies. Experiments 5-8, by comparison, show the effects of a T-X-T mismatch on circle complex binding strength. Once again, the true match ($X = A$) gives the most favorable three stranded complexes ($-\Delta G^\circ_{37} = 16.4$ kcal). However, target mismatches ($X = G, T, C$) result in a considerably larger loss of binding energy (6.2-7.6 kcal/mol) for a circular oligonucleotide than for a linear oligonucleotide.

Similarly, experiments 9-12 give the effects of a C-Y mismatch on the two stranded duplex. The matched base ($Y = G$) gives a free energy of duplex association of -10.3 kcal/mol. The mismatches ($Y = A, T, C$) result in a

35

1

loss of 5.2 to 5.8 kcal/mol of binding energy, in reasonable agreement with published data. By contrast, the effects of a C-Y-C mismatch are greater in a three
5 stranded complex (experiments 13-16): the match (Y = G) gives a binding energy of -16.4 kcal/mol, and the mismatches (Y = A, T, C) are less stable by 7.1 to 7.5 kcal/mol.

Thus, in all the cases studied, the circular
10 ligand shows greater selectivity for its correctly matched sequence than does the standard linear oligomer. The selectivity advantage ranges from 1.3 to 2.2 kcal/mol for the C-Y-C series to 3.0 to 3.4 kcal/mol for the T-X-T series. These are quite significant differences,
15 considering they arise from a single base change; in the T-X-T series, the circular oligonucleotide is nearly twice as selective as the linear oligonucleotide. This selectivity difference corresponds to one to two orders of magnitude in binding constant at 37°C.

20 There are two factors which may explain this high selectivity. First, because two domains of the circular oligonucleotide bind the central target strand, the circular oligonucleotide, in effect, checks the sequence twice for correct matching. Secondly,
25 protonation of cytosine within a C+G-C triad may also be a factor in increasing selectivity. This protonation is likely to be favored only when there is base triad formation wherein guanine can share the positive charge; evidence suggests that the pKa of cytosine within a base
30 triad is 2-3 units higher than that of free deoxycytosine. The addition of this positive charge may

35

-64-

1

lessen the negative charge repulsions arising from the high density of phosphates in the complex and thereby increase binding stability.

5

Therefore, circular oligonucleotides, as described herein, to have both higher binding affinity and higher selectivity than can be achieved with Watson-Crick duplexes alone.

10

15

20

25

30

35

-65-

1

TABLE III

| | | expt. # | variable base | $T_m, ^\circ\text{C}$ | $-\Delta G_{37}^\circ$ (kcal/mol) | Selectivity (kcal/mol) |
|----|----------------|---------|---------------|-----------------------|--------------------------------------|---------------------------|
| 10 | duplex | 1 | X=A | 43.8 | 10.3 | -- |
| | | 2 | X=G | 33.8 | 7.1 | 3.2 |
| | | 3 | X=C | 28.3 | 5.9 | 4.4 |
| | | 4 | X=T | 31.1 | 6.4 | 3.9 |
| | | 5 | X=A | 62.3 | 16.4 | -- |
| 15 | circle complex | 6 | X=G | 44.2 | 10.2 | 6.2 |
| | | 7 | X=C | 39.8 | 8.8 | 7.6 |
| | | 8 | X=T | 40.8 | 9.1 | 7.3 |
| | | 9 | Y=A | 26.2 | 5.1 | 5.2 |
| 20 | duplex | 10 | Y=G | 43.8 | 10.3 | -- |
| | | 11 | Y=C | 22.2 | 4.5 | 5.8 |
| | | 12 | Y=T | 27.0 | 5.0 | 5.3 |
| | | 13 | Y=A | 39.9 | 9.0 | 7.4 |
| 25 | circle complex | 14 | Y=G | 62.3 | 16.4 | -- |
| | | 15 | Y=C | 41.3 | 9.3 | 7.1 |
| | | 16 | Y=T | 39.6 | 8.9 | 7.5 |

30

35

SUBSTITUTE SHEET

-66-

1

EXAMPLE 4Factors Effecting Complex Formation

5 1) Solution effects. The effects of NaCl, Mg²⁺, spermine, and pH on circle:target complexes were examined. Circles with cytosines in the binding domains are sensitive to pH, and exhibited greater stability at lower pH values. However, these and other circle:target complexes are quite stable at the physiological pH of 10 7.0-7.4 (Fig. 5). The complexes show salt concentration sensitivity comparable to duplexes; however, small amounts of Mg²⁺ or spermine increase the complex stability markedly. For example, in a concentration of 1 mM Mg⁺⁺ at pH 7.0, with no added salts, a stable 7:5 15 circle:target complex formed having a T_m of 58°C. When a solution of 20 μm spermine containing no added salts was used the 7:5 complex again formed stably with a T_m of 56°C. Both Mg⁺⁺ and spermine are present in at least these concentrations in mammals, and so circle:target 20 complexes will be stable under physiological conditions.

2) Loop size. The optimum number of nucleotides for the loop domain of a circle was determined by observing complex formation between a target and circles with different loop sizes. Precircle 25 linear oligonucleotides similar to precircle 1 were synthesized with 2, 3, 4, 5, 6 and 10 base loops using an arbitrary sequence of alternating C and A residues. Each of these precircles was designed to bind to the A₁₂ template (i.e. target 4 (SEQ ID NO: 8)). The T_m's for 30 circles with 4, 5, 6 and 10 base loops showed that a five-nucleotide loop size was optimum for the circle

35

SUBSTITUTE SHEET

-67-

1

binding either to template A₁₂ or to a longer 36mer sequence containing the A₁₂ binding site (see Fig. 6A).

3) Binding Domain length. The effect of
5 circular oligonucleotide binding domain length on circle:target complex melting temperature was compared to melting of duplexes having the same length. Circles with various size binding domains were constructed and complexed with single-stranded dA_n targets for n equal to
10 4, 8, 12 and 18 nucleotides. Fig. 6B illustrates that considerably higher T_m's were observed for circle:target complexes relative to Watson-Crick duplexes having the same length as the binding domains (determined in 0.1 M NaCl, pH 7). For example, a 12-base circular complex
15 melted at about the same temperature as a 24-base duplex. The 4-base circular complex melted at 34°C, whereas the corresponding Watson-Crick duplex T_m was less than 0°C.

4) Methylation. It has been known for some time that methylation at the C-5 position of cytosine,
20 forming the naturally-occurring base m⁵C, raises the T_m of duplex DNA in which it occurs, relative to unmethylated sequences (Zmudzka et al., 1969, Biochemistry 8: 3049). In order to investigate whether addition of this methyl group would stabilize
25 circle:target complexes, two analogs of circle 7 (having SEQ ID NO: 6) were synthesized. In one circle, the six C's in the binding domains were methylated leaving the loop unmethylated (Me₆). In the second circle, all twelve C's were methylated (Me₁₂). Melting temperatures
30 for the complexes of these methylated circle with target 5 were measured. The Me₆ complex had a T_m of 71.1°C

35

SUBSTITUTE SHEET

-68-

1

(compared to 61.8°C for the unmethylated circle), and the Me₁₂ circle had a T_m of 72.4°C. Thus, use of the natural base m⁵C in place of C increased stability substantially, and in one case resulted in a 12-base complex which melted 10.6°C higher than an unmethylated circle and 28.6°C higher than the corresponding unmethylated Watson-Crick duplex.

10

15

20

25

30

35

1

EXAMPLE 5Replacement of Nucleotide Loop Domains
with Non-Nucleotide Loop Domains

5

The loop domains of circular oligonucleotides were replaced with polyethylene or oligoethylene glycol chains of different lengths and the effect of such synthetic loops upon circular oligonucleotide binding and nuclease resistance was assessed.

10

Methods

Circular oligonucleotides were synthesized having tetra-, penta-, or hexa-ethylene glycol chain loop domains. In each case the ethylene glycol chain was synthetically prepared for automated DNA synthetic procedures using the method of Durand *et al.* (1990, Nucleic Acids Res. 18: 6353-6359). Briefly, a phosphoramidite was placed on a hydroxy group at one end of the ethylene glycol chain and a dimethoxytrityl (DMT) moiety was placed on the other terminal ethylene glycol hydroxy group. This derivatized ethylene glycol chain was then added to the growing linear oligonucleotide at the appropriate step of automated DNA synthesis. Circularization steps were performed by procedures described in Example 1. A linear oligonucleotide precircle having a tetraethylene loop domain was not efficiently circularized. This result indicates that a tetraethylene loop domain may be too short for optimal binding to a target.

Two types of linear oligonucleotides were used as target binding domains for the circular oligonucleotides: Target I was a 12-base oligonucleotide

35

-70-

1

having no non-target nucleotides and Target II was a 36-base oligonucleotide having a 12-base target within it. The target sequences utilized were 5'-AAGAAAAGAAAAG-3' (SEQ ID NO: 9) and 5'-AAAAAAAAAAAA-3' (SEQ ID NO: 8), the latter is termed a poly(dA)₁₂ target sequence.

5

The melting temperatures (T_m) of circular oligonucleotides with polyethylene loops were observed at pH 7.0 (10 mM Tris-HCl) in 10 mM MgCl₂ and 100 mM NaCl. Each linear target and each circular oligonucleotide was present at a 3 μM concentration.

10

Results

The T_m of a circular oligonucleotide having a CACAC nucleotide loop sequence and a poly(dT)₁₂ sequence for both P and AP domains was 57.8°C when bound to a poly(dA)₁₂ target sequence. The T_m of a circular oligonucleotide having the same P and AP domain sequences but hexaethylene glycol loop domains was 51.4 °C when bound to the same target.

15

20

A comparison of T_m values observed for circular oligonucleotides having pentaethylene glycol (PEG) and hexaethylene glycol (HEG) loop domains is depicted in Table IV.

25

30

35

1

TABLE IV

| Complex | | Target I T _m | Target II T _m |
|---------|---|----------------------------|-----------------------------|
| 10 | p T T C T T T T C T T T C p PEG A A G A A A A G A A A G PEG p T T C T T T T C T T T C p | 51.5 | 47.5 |
| 15 | p T T C T T T T C T T T C p HEG A A G A A A A G A A A G HEG p T T C T T T T C T T T C p | 58.0 | 51.1 |
| 20 | p T T T T T T T T T T T T p HEG A A A A A A A A A A A A HEG p T T T T T T T T T T T T p | 51.4 | 46.5 |

25 The T_m value observed for a circular oligonucleotide having a HEG loop is about 4.5°C higher than that of a circular oligonucleotide with a PEG loop. Therefore, circular oligonucleotides with hexaethylene glycol loop domains bind with greater stability than do circular oligonucleotides with tetra- or penta-ethylene glycol loops.

30

35

-72-

1

Nuclease Resistance

5 Circular oligonucleotides were tested for nuclease resistance when unbound and when bound to a target oligonucleotide. All circular oligonucleotides, whether bound or unbound, were completely resistant to exonucleases. Endonuclease sensitivity was assessed using S1 nuclease according to the manufacturer's suggestions.

10

A comparison of the resistance of bound and unbound circular oligonucleotides to S1 nuclease is depicted in Table V.

15

20

25

30

35

1

TABLE V

5

| Oligonucleotide Cleavage | Time For 50% S1 |
|-----------------------------|--------------------|
|-----------------------------|--------------------|

10

| | |
|---|--------|
| p T T C T T T T C T T T C p HEG HEG | 1 min. |
| p T T C T T T T C T T T C p | |

15

| | |
|--|--------|
| p T T C T T T T C T T T C p HEG A A G A A A G A A A G HEG | > 24 h |
| p T T C T T T T C T T T C p | |

20

| | |
|---|--------|
| A C T T C T T T T C T T T C C A C C | 1 min. |
| A C T T C T T T T C T T T C C A | |

25

| | |
|--|---------|
| A C T T C T T T T C T T T C C A C A A G A A A G A A A G C | 40 min. |
| A C T T C T T T T C T T T C C A | |

30

35

-74-

1

These data indicate that unbound circular oligonucleotides are vulnerable to S1 nuclease. However, when bound to a target, a circular oligonucleotide having a polyethylene loop domain is much more resistant to S1 nuclease, at least 36-fold more resistant, than a circular oligonucleotide with a nucleotide loop domain.

The nuclease resistance of circular and linear oligonucleotides was also compared when these oligonucleotides were incubated in human plasma for varying time periods. Circular oligonucleotide 7 and the precursor to this circle, linear oligonucleotide 2, were incubated at a 50 μ M concentration in plasma at 37°C. Aliquots were removed at various time points and cleavage products were separated by gel electrophoresis. Nuclease resistance was assessed by observing whether degradation products were evident on the gels.

When incubated in human plasma the half-life of linear oligonucleotide 2 was 20 min. In contrast, circular oligonucleotide 7 underwent no measurable nuclease degradation during a 48 hr incubation. Accordingly, the half-life of a circular oligonucleotide is greater than 48 hr in human plasma, i.e. more than 140 times longer than a linear oligonucleotide having an equivalent sequence.

30

35

1

EXAMPLE 6
CIRCULAR OLIGONUCLEOTIDES CAN SELECTIVELY
BIND TO RNA

5

Experiments described in this Example indicate that, unlike linear oligonucleotides, circular oligonucleotides can preferentially bind to an RNA, rather than a DNA, target.

10

Two linear deoxyoligonucleotides were prepared as targets, a "T" (SEQ ID NO.: 11) target and a "dU" (SEQ. ID. No.: 12) target:


T target: 5'-A A G A A T A G A A A G-3'; and

15

dU target: 5'-A A G A A U A G A A A G-3'.

A circular oligonucleotide having SEQ ID NO.: 14 was also prepared:

20



For comparison, a linear oligonucleotide complementary to the T and dU targets was also synthesized (i.e. the linear oligonucleotide, SEQ ID NO.: 13):

25

5'C T T T C T A T T C T T 3'.

The melting temperatures (T_m) values observed for the circular vs linear oligonucleotide binding to each of the targets is presented in Table VI.

30

35

-76-

1

TABLE VI

5

| <u>T_m Values for Oligonucleotides</u> | | |
|--|---------------|-----------------|
| <u>Targets</u> | <u>Linear</u> | <u>Circular</u> |
| T target | 42.9° C | 41.1° C |
| dU target | 40.9° C | 42.9° C |

10

The linear oligonucleotide binds more strongly to the T target than to the dU target, by an amount which is significantly larger than experimental error limits.

15 This difference in T_m values corresponds to a difference in free energy of binding of 1.7 kcal/mole.

However, in contrast to the linear oligonucleotide, the circular oligonucleotide binds more strongly to the U target. Therefore, the circular
20 oligonucleotide can exhibit a preference for an RNA target relative to the corresponding DNA target.

Moreover, the increase in binding strength for a circular oligonucleotide to the RNA target corresponds to a free energy difference of 0.8 kcal/mole which
25 indicates that at 37 °C an RNA target would be preferred by about 3:1 over a corresponding DNA target.

30

35

-77-

1

EXAMPLE 7Strand Replacement By Circular Oligonucleotides

5

Circular oligonucleotide 6 (Fig. 3) bound to a dA₁₂ target with 9 kcal/mole greater stability than did a linear dT₁₂ oligonucleotide (Example 2). This increase in stability demonstrates that a circular-oligonucleotide:target complex is thermodynamically favored over a linear-oligonucleotide:target. In addition, a circular oligonucleotide can actually accelerate (or catalyze) dissociation of duplex DNA target sequences to form a complex with one strand of the duplex.

15

To test whether a circular oligonucleotide can readily dissociate duplex DNA and displace one strand of a duplex DNA target, the kinetics of strand displacement were observed for a duplex DNA target in the presence of a complementary linear or circular oligonucleotide.

20

A DNA duplex target with a fluorescein group on one strand and a tetramethylrhodamine group on the other strand was prepared using published procedures (Cardullo *et al.* 1988 Proc. Natl. Acad. Sci. USA 85: 8790; Cooper *et al.* 1990 Biochemistry 29: 9261). The structure of the duplex target (SEQ ID NO.: 15) was as follows:

25

5'-fluorescein-A A A A A A A A A A A A
3'-rhodamine-T T T T T T T T T T T T.

The T_m of this labeled duplex target was normal, therefore the fluorescent substituents had no significant effect upon association kinetics. Moreover, the emission maxima of the fluorescein-dA₁₂ strand was 523 nm while the

35

-78-

1

emission maxima of the rhodamine-dT₁₂ strand was 590 nm, allowing the association kinetics of the two strands could be separately monitored.

5

Strand displacement reactions were done at 10°C in a 1 cm fluorescence cuvette. Reaction conditions were 100 mM NaCl, 10 mM Mg Cl₂ and 10 mM Tris-HCl, pH 7.0 with a reaction volume of 3 ml. Labeled duplex was allowed to equilibrate for at least 1 hr at 10°C before addition of a 40-fold excess of linear or circular oligonucleotide (final concentration 0.01 μM). A Spex Fluorolog F 111A fluorescence instrument with 5 mm slit widths was used. An excitation wavelength of 450 nm and a monitored emission wavelength of 523 nm was used. The results were independent of both excitation and monitored emission wavelengths. Reactions were followed for at least 5 half-lives.

10

Addition of rhodamine-dT₁₂ to fluorescein-dA₁₂ caused a decrease in fluorescein fluorescence and an increase in rhodamine fluorescence. Such effects are due to energy transfer between the fluorescent moieties (Cardullo *et al.*).

15

The association rate constant of the two fluorescently-labeled strands was determined by mixing the strands under pseudo-first order conditions and monitoring the rate of decrease in fluorescein emission. At 10 °C the observed association constant was 3.2 X 10⁶ M⁻¹ sec⁻¹, which agrees well with published rates of association for DNA oligonucleotides (Nelson *et al.* 1982 Biochemistry 21: 5289; Turner *et al.* 1990 in Nucleic

20

25

30

35

-79-

1

Acids (subvolume C), W. Saenger, Ed. Springer-Verlag, Berlin: 201-227).

5 To compare the rates at which a single linear strand (SEQ ID NO.: 8) or a circular oligonucleotide having SEQ ID NO.: 5 (i.e. circular oligonucleotide 6) exchanged with strands in a duplex DNA, an excess of an unlabeled linear or circular oligonucleotide was mixed with the fluorescently-labeled duplex DNA target. The
10 increase in fluorescein emission was then observed at a temperature significantly below the T_m of the duplex target as a measure of duplex target strand dissociation.

Fig. 8 depicts a typical kinetic assay for the dissociation of duplex target by a 40-fold excess of
15 unlabeled $dA_{1,2}$ (dotted line) or circular oligonucleotide 6 (solid line) at 10 °C. As depicted, duplex target dissociation by the circular oligonucleotide is considerably faster than is the dissociation by the linear oligonucleotide. The first order rate constant
20 for dissociation by the linear oligonucleotide is $2.0 \times 10^{-4} \text{ sec}^{-1}$ whereas the first order rate constant for dissociation by the circular oligonucleotide is $2.3 \times 10^{-2} \text{ sec}^{-1}$, almost two orders of magnitude faster. This difference is even more apparent when the half-lives for
25 the target duplex in the presence of linear vs circular oligonucleotides are calculated. At 10 °C, the duplex has a half-life for dissociation of 58 min in the presence of the linear oligonucleotide but only 30 sec in the presence of the circular oligonucleotide.

30 Unlike the rate of reaction between linear oligonucleotide and duplex, the rate of reaction between

35

-80-

1

the circular oligonucleotide and duplex is dependent on the concentration of added circular oligonucleotide at low concentrations, and shows Michaelis-Menten type saturation behavior at higher concentrations (Fig. 9).

5

The dissociation rate of labeled duplex at 10°C can be derived from the duplex association rate constant and ΔG°_{10} values. This rate constant, $8.5 \times 10^{-10} \text{ sec}^{-1}$, is consistent with rates derived from predicted thermodynamic parameters for a duplex complex (Breslauer et al. 1986 Proc. Natl. Acad. Sci. USA 83: 3746) although this rate is significantly slower than the rate constant for strand displacement by a linear oligonucleotide. An increase in duplex dissociation upon addition of a linear oligonucleotide has been noted in other cases (Chamberlin et al. 1965 J. Mol. Biol. 12: 410). Comparison of the rate for the circular oligonucleotide-catalyzed reaction over that of the unassisted duplex dissociation reveals a rate enhancement of about 10^7 fold (Sigler et al. 1962 J. Mol. Biol. 5: 709) .

10

15

20

A double reciprocal plot of $1/[\text{circular oligonucleotide}]$ vs. $1/k_{\text{obs}}$ is linear and yields a k_{cat} of $0.024 \pm 0.005 \text{ sec}^{-1}$ and a K_M of $2.2 \times 10^{-7} \text{ M}$. The k_{cat} is 100-fold greater than the observed rate constant obtained for the reaction of the duplex with either dA_{12} or dT_{12} single strands.

25

30

The observed saturation behavior (Fig. 9) suggests that a complex forms between the circle and the double-stranded target. Using the above K_M value and assuming that $k_{\text{cat}} \ll k_{-1}$, where k_{-1} is the dissociation rate constant for this complex, the free energy of

35

-81-

1

association is $-8.6 \text{ kcal}\cdot\text{mol}^{-1}$ at 10°C . This value is similar to an estimated value of about $-9 \text{ kcal}\cdot\text{mol}^{-1}$ for the P domain in a 12-base triple helix consisting of T·A-
5 T base triads, as derived from the thermodynamic parameters of Pilch et al. (1990 Nucleic Acids Res. 18: 5743).

10

15

20

25

30

35

-82-

1

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Kool, Eric T.

(ii) TITLE OF INVENTION: SINGLE-STRANDED, CIRCULAR
OLIGONUCLEOTIDES

(iii) NUMBER OF SEQUENCES: 15

10

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Scully, Scott, Murphy & Presser
(B) STREET: 400 Garden City Plaza
(C) CITY: Garden City
(D) STATE: New York
(E) COUNTRY: USA
(F) ZIP: 11530

15

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version

#1.25

20

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

25

(A) NAME: McNulty, William E.
(B) REGISTRATION NUMBER: 22,606
(C) REFERENCE/DOCKET NUMBER: 8085Z

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (516) 742-4343
(B) TELEFAX: (516) 742-4366

30 (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

35

-83-

1

(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: circular

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCCCCGCC TCNNNNNCTC CCACCCCTCN NNNN
34

10 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: circular

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCTTTTTTCT TTCNNNNNC TTTTCTTTTT TCTNNNNN
38

20 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: circular

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TTTCYTCGTT CGTCNNNNNC TACTTACTGC TTTNNNNN
38

30 (2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

35

SUBSTITUTE SHEET

-84-

1

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: circular

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCCTTCTTCY CCTCTNNNNN TCTCCGCTTC TTCCTNNNNN
40

10 (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: both

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TTTTTTCACA CTTTTTTTTT TTTCACACTT TTTT
34

20 (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: both

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TCTTTCCACA CCTTTCTTTT CTTCACACTT CTTT
34

30 (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:

35

-85-

1

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: both

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TTTCTTCACA CTTCTTTTCT TTCCACACCT TTCT
34

10 (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AAAAAAAAAA AA
12

20 (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

AAGAAAAGAA AG
12

30 (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:

35

SUBSTITUTE SHEET

-86-

1

- (A) LENGTH: 12 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CTTTCTTTTC TT
12

10 (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

AAGAATAGAA AG
12

20 (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AAGAAUAGAA AG
12

30 (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:

35

-87-

1

- (A) LENGTH: 12 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CTTTCTATTTC TT
12

10 (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 34 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: both

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TTCTTCTCTT TCCACACCTT TCTATTCTTC ACAC
34

20 (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AAAAAAAAAA AA
12

30

35

-88-

1

WHAT IS CLAIMED:

5 1. A single-stranded circular oligonucleotide comprising at least one parallel binding (P) domain and at least one anti-parallel binding (AP) domain having a loop domain between each binding domain to form said circular oligonucleotide; each P and corresponding AP domain having sufficient complementarity to bind detectably to one strand of a defined nucleic acid target wherein said P domain binds in a parallel manner to said target, and said corresponding AP domain binds in an anti-parallel manner to said target.

10 2. The oligonucleotide of Claim 1 wherein said target comprises a known nucleotide sequence from which a nucleotide sequence for a sufficient number of positions in said P domain and in said corresponding AP domain is determined from the sequence of said target for said P domain:

15 when a base for a position in said target is guanine or a guanine analog, then P has cytosine, or a suitable analog thereof, in a corresponding position;

20 when a base for a position in said target is adenine, or an adenine analog then P has thymine or uracil, or suitable analogs thereof, in a corresponding position;

25 when a base for a position in said target is thymine, or a thymine analog, then P has cytosine or guanine, or suitable analogs thereof, in a corresponding position;

30 when a base for a position in said target is cytosine, or a cytosine analog, then P has cytosine,

35

-89-

1

thymine or uracil, or suitable analogs thereof, in a corresponding position; and

5 when a base for a position in said target is uracil, or a uracil analog, then P has cytosine, guanine, thymine or uracil, or suitable analogs thereof, in a corresponding position;

and for said AP domain:

10 when a base for a position in said target is guanine, or a guanine analog, then AP has cytosine or uracil, or suitable analogs thereof, in a corresponding position;

15 when a base for a position in said target is adenine, or an adenine analog, then AP has thymine or uracil, or suitable analogs thereof, in a corresponding position;

when a base for a position in said target is thymine, or a thymine analog, then AP has adenine, or a suitable analog thereof, in a corresponding position;

20 when a base for a position in said target is cytosine, or a cytosine analog, then AP has a guanine, or a suitable analog thereof, in corresponding position; and

25 when a base for a position in said target is uracil, or a uracil analog, then AP has adenine or guanine, or suitable analogs thereof, in a corresponding position;

30 wherein said sufficient number of positions is that number of positions to provide sufficient complementarity for said oligonucleotide to bind detectably to said target.

35

1

3. The oligonucleotide of Claim 1 wherein said P domain comprises a nucleotide sequence which is determined from a known nucleotide sequence of said target:

5

when a base for a position in said target is guanine or a guanine analog, then P has cytosine, or a suitable analog thereof, in a corresponding position;

10

when a base for a position in said target is adenine, or an adenine analog then P has thymine or uracil, or suitable analogs thereof, in a corresponding position;

15

when a base for a position in said target is thymine, or a thymine analog, then P has cytosine or guanine, or suitable analogs thereof, in a corresponding position;

20

when a base for a position in said target is cytosine, or a cytosine analog, then P has cytosine, thymine or uracil, or suitable analogs thereof, in a corresponding position;

25

when a base for a position in said target is uracil, or a uracil analog, then P has cytosine, guanine, thymine or uracil, or suitable analogs thereof, in a corresponding position;

and further wherein said AP domain comprises a nucleotide sequence which is determined from said sequence of said target as follows:

30

when a base for a position in said target is guanine, or a guanine analog, then AP has cytosine or uracil, or suitable analogs thereof, in a corresponding position;

35

1

when a base for a position in said target is adenine, or an adenine analog, then AP has thymine or uracil, or suitable analogs thereof, in a corresponding
5 position;

when a base for a position in said target is thymine, or a thymine analog, then AP has adenine, or a suitable analog thereof, in a corresponding position;

10 when a base for a position in said target is cytosine, or a cytosine analog, then AP has a guanine, or a suitable analog thereof, in corresponding position; and

when a base for a position in said target is uracil, or a uracil analog, then AP has adenine or guanine, or suitable analogs thereof, in a corresponding
15 position.

4. The oligonucleotide of Claim 1, 2 or 3 wherein said target, said P domain and said AP domain independently comprise from about 2 to about 200
nucleotides.

20 5. The oligonucleotide of Claim 4 wherein said target, said P domain and said AP domain independently comprise from about 6 to about 36 nucleotides.

6. The oligonucleotide of Claim 1, 2 or 3 wherein each loop domain independently comprises from
25 about 2 to about 2000 nucleotides.

7. The oligonucleotide of Claim 6 wherein each loop domain independently comprises from about 3 to about 8 nucleotides.

30 8. The oligonucleotide of Claim 1, 2 or 3 wherein said target is single stranded or double stranded.

35

1

9. The oligonucleotide of Claim 1, 2 or 3 wherein said target is RNA or DNA.

5 10. The oligonucleotide of Claim 1, 2 or 3 wherein said target is a domain contained in a nucleic acid template.

11. The oligonucleotide of Claim 1, 2 or 3 wherein said P domain and said AP domain bind to said target in a staggered binding arrangement.

10 12. The oligonucleotide of Claim 1 or 2 wherein sufficient complementarity is less than 100% complementarity.

13. The oligonucleotide of Claim 12 wherein sufficient complementarity is about 30% to about 40% complementarity.

14. The oligonucleotide of Claim 1, 2 or 3 wherein said oligonucleotide is DNA or RNA.

15 20 15. The oligonucleotide of Claim 2 or 3 wherein a suitable analog of cytosine is 5-methylcytosine.

16. The oligonucleotide of Claim 2 or 3 wherein a suitable analog of uracil is 5-methyluracil.

17. The oligonucleotide of Claim 2 or 3 wherein a suitable analog of adenine is diaminopurine.

25 18. The oligonucleotide of Claim 1, 2 or 3 wherein nucleotides have a 2'-O-methylribose in place of ribose or deoxyribose.

19. The oligonucleotide of Claim 1, 2 or 3 wherein said oligonucleotide is taken up in a cell.

30 20. The oligonucleotide of Claim 19, wherein said oligonucleotide further comprises a ligand for a

35

1

cellular receptor, cholesterol group, an aryl group, a steroid group or a polycation.

5

21. The oligonucleotide of Claim 1, 2 or 3 wherein said oligonucleotide further comprises a drug or a drug analog.

22. The oligonucleotide of Claim 1, 2 or 3 wherein said loop domains comprise non-nucleotide loop domains.

10

23. The oligonucleotide of Claim 22 wherein said non-nucleotide loop domains are polyethylene glycol.

24. The oligonucleotide of Claim 23 wherein said polyethylene glycol is pentaethylene glycol, hexaethylene glycol or heptaethylene glycol.

15

25. The oligonucleotide of Claim 1, 2 or 3 wherein said oligonucleotide further comprises at least one methylphosphonate, phosphorothioate, phosphorodithioate, phosphotriester, siloxane, carbonate, acetamidate, thioether or phosphorus-boron linkage.

20

26. The oligonucleotide of Claim 1, 2 or 3 wherein said oligonucleotide further comprises a reporter molecule.

27. A compartmentalized kit for detection or diagnosis of a target nucleic acid, comprising:

25

- at least one first container providing a circular oligonucleotide of any one of Claims 1-3.

30

28. A compartmentalized kit for isolation of a template nucleic acid, comprising at least one first container providing a circular oligonucleotide of Claim 1, 2 or 3, wherein said oligonucleotide is complementary to a target contained within said template.

35

1

29. The kit of Claim 28 wherein said template is poly (A)⁺ mRNA.

5 30. A method of regulating biosynthesis of a DNA, an RNA or a protein which comprises:

contacting at least one oligonucleotide of any one of Claims 1 to 3 with a nucleic acid template for said DNA, said RNA or said protein, under conditions sufficient to permit binding of said at least one
10 oligonucleotide to a target sequence contained within said template;

binding said oligonucleotide to said target;
blocking access to or allowing degradation of said template and thereby regulating biosynthesis of said
15 DNA, said RNA or said protein.

31. The method of Claim 30 wherein said template comprises a double-stranded nucleic acid target, and wherein said conditions are effective to denature said target by strand displacement and thereby permit
20 binding.

32. The method of Claim 30 wherein said biosynthesis comprises at least one of DNA replication, DNA reverse transcription, RNA transcription, RNA splicing, RNA polyadenylation, RNA translocation and
25 protein translation.

33. The method of Claim 32 wherein said template for said DNA replication is an RNA template or a DNA template.

34. The method of Claim 33 wherein said target
30 of said oligonucleotide for regulating said DNA

35

1

replication is an origin of replication or a primer binding site.

5 35. The method of Claim 32 wherein said target of said oligonucleotide for regulating said DNA reverse transcription is a primer binding site, a site in a retroviral genome, or a site in an mRNA.

10 36. The method of Claim 32 wherein said target of said oligonucleotide for regulating said RNA transcription is a promoter, a repressor binding site, an operator, an enhancer, a transcription regulatory element or a site in an mRNA encoding region.

15 37. The method of Claim 32 wherein said target of said oligonucleotide for regulating said RNA splicing is at least one of a 5' splice junction, an intron branch point or a 3' splice junction.

38. The method of Claim 32 wherein said target of said oligonucleotide for regulating said RNA polyadenylation is a polyadenylation site.

20 39. The method of Claim 32 wherein said target of said oligonucleotide for regulating said RNA translocation is a poly(A) tail.

25 40. The method of Claim 32 wherein said template for said protein translation is an mRNA template.

41. The method of Claim 40 wherein said target of said template is a ribosome binding site, a 5' mRNA cap or a site in a protein coding region.

30 42. The method of Claim 30 wherein said template is a viral DNA or RNA template.

35

1

43. The method of Claim 42 wherein said oligonucleotide has a nucleotide sequence of SEQ ID NO: 3 or SEQ ID NO: 4.

5

44. A method of strand displacement in a double-stranded nucleic acid target which comprises contacting said target with a circular oligonucleotide of any one of Claims 1-3 for a time and under conditions effective to denature said target and to bind said circular oligonucleotide.

10

45. The method of Claim 44 wherein said conditions effective to denature said target comprise having said circular oligonucleotide and said target present in a ratio of about 1 to about 100.

15

46. The method of Claim 45 wherein said time effective to denature said target ranges from about 1 minute to about 16 hours.

47. The method of Claim 44 wherein said double-stranded nucleic acid target comprises a viral, a bacterial, a fungal or a mammalian nucleic acid.

20

48. The method of Claim 47 wherein said double-stranded nucleic acid target is an origin of replication, a promoter, a repressor binding site, an operator, an enhancer, a transcription regulatory element or a site in an mRNA encoding region.

25

49. The method of Claim 44 wherein said double-stranded nucleic acid target is present in a pure or impure nucleic acid sample, a tissue section, a cell smear or a chromosomal squash.

30

35

1

50. The method of Claim 49 wherein said oligonucleotide is covalently linked to a reporter molecule.

5

51. A pharmaceutical composition for regulating biosynthesis of a nucleic acid or protein comprising a biosynthesis regulating amount of at least one of the oligonucleotides of Claims 1 to 3 and a pharmaceutically acceptable carrier.

10

52. A method of preparing the single-stranded circular oligonucleotide of Claim 1, 2 or 3 comprising binding a linear precircle to an end-joining-oligonucleotide, joining two ends of said precircle and recovering said single-stranded circular oligonucleotide.

15

53. The method of Claim 52 wherein said linear precircle has a 3'-phosphate.

54. The method of Claim 53 wherein said two ends comprise two nucleotides corresponding to AP nucleotides of said single-stranded circular oligonucleotide.

20

55. The method of Claim 54 wherein said joining is performed with BrCN, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide or N-cyanoimidazole $ZnCl_2$.

25

56. A complex formed between the oligonucleotide of Claim 1, 2 or 3 and a target.

57. A method of specific cell type drug delivery comprising:

a) administering to an animal a drug covalently linked to an oligonucleotide of Claim 1, 2 or 3;

30

3;

35

1

b) binding said oligonucleotide to a target mRNA present in said cell type; and

5 c) thereby delivering said drug to said specific cell type.

58. A method of detecting a target nucleic acid which comprises:

10 contacting a circular oligonucleotide of any one of Claims 1 to 3 with a sample to be tested for containing said nucleic acid for a time and under conditions sufficient to form an oligonucleotide-target complex; and

detecting said complex.

15 59. The method of Claim 58 wherein said nucleic acid comprises a double-stranded nucleic acid target, and wherein said conditions are effective to denature said target by strand displacement and thereby permit binding of said oligonucleotide to form said oligonucleotide-target complex.

20 60. The method of Claim 58 wherein said sample comprises a pure or impure nucleic acid sample, a tissue section, a cell smear or a chromosomal squash.

25 61. The method of Claim 58 wherein said conditions effective to denature said target comprise having said circular oligonucleotide and said target present in a ratio of about 1 to about 100.

30 62. The method of Claim 58 wherein said time effective to denature said target ranges from about 1 minute to about 16 hours.

35

35

1

63. The method of Claim 58 wherein said complex is detected by a fluorescence energy transfer assay.

5

64. The oligonucleotide of Claim 12 wherein sufficient complementarity is at least about 50%.

10

15

20

25

30

35

FIG. IA 1 / 8




FIG. IB




FIG. 2





FIG.3


Precircles (1-3), Targets (4-5), Linear Oligonucleotides (9) and Circles (6-8) used in Experiments

- 1 5'-TTTTTTCACACTTTTTTTTTTTTTCACACTTTTTTT (SEQ ID NO:5)
 2 5'-TCTTCCACACCTTCTTTTCTTTCACACTTCTTT (SEQ ID NO:6)
 3 5'-TTTCTTTCACACTTCTTTTCTTTCACACCTTCT (SEQ ID NO:7)
 4 5'-AAAAAAAAAAAAA (SEQ ID NO:8)
 5 5'-AAGAAAAGAAAG (SEQ ID NO:9)

6 (SEQ ID NO:5)



7 (SEQ ID NO:6)



8 (SEQ ID NO:7)




FIG. 3
CONT.





FIG. 4



pH Effects on T_m

FIG. 5




FIG. 6B




FIG. 6A




FIG. 7




FIG. 8





FIG. 9



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/02480

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(5) : G120, 1/68; C07H 15/12, 17/00
 US CL : 435/6; 538/27, 28, 29
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 U.S. :
 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| Y | US, A. 4,766,062 (Diamond et al) 23 August 1988, see entire document. | 1-64 |

Further documents are listed in the continuation of Box C. See patent family annex.

| | |
|---|--|
| * Special categories of cited documents: | "I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "A" document defining the general state of the art which is not considered to be part of particular relevance | "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "E" earlier document published on or after the international filing date | "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "&" document member of the same patent family |
| "O" document referring to an oral disclosure, use, exhibition or other means | |
| "P" document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search: 23 July 1992
 Date of mailing of the international search report: 28 JUL 1992

Name and mailing address of the ISA/ Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231
 Authorized officer: L. YUAN 