




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ³ : A61K 37/00, 37/48; C07C103/52	A1	(11) International Publication Number: WO 83/ 03971 (43) International Publication Date: 24 November 1983 (24.11.83)
<p>(21) International Application Number: PCT/US83/00723</p> <p>(22) International Filing Date: 12 May 1983 (12.05.83)</p> <p>(31) Priority Application Number: 377,386</p> <p>(32) Priority Date: 12 May 1982 (12.05.82)</p> <p>(33) Priority Country: US</p> <p>(71) Applicant: PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US).</p> <p>(72) Inventor: MURPHY, John, R. ; 35 Ashfield Street, Roslindale, MA 02131 (US).</p> <p>(74) Agent: FURLONG, Robert, W.; Fish & Richardson, 28 State Street, Suite 2100, Boston, MA 02109 (US).</p>	<p>(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), FI, FR (European patent), GB (European patent), JP, KP, LU (European patent), NL (European patent), NO, SE (European patent).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: HYBRID PROTEINS



(57) Abstract

A hybrid protein including protein fragments joined together by peptide bonds, the hybrid protein including, in sequential order, beginning at the amino terminal end of the hybrid protein, (a) the enzymatically active Fragment A of diphtheria toxin, (b) a fragment including the cleavage domain l_1 adjacent Fragment A, (c) a fragment including at least the portion of Fragment B of diphtheria toxin encoded by the portion of the Fragment B encoding gene fragment of the *tox* operon between l_1 and the position about 90 base pairs upstream from the position on the *tox* operon of the NRU I site of the *tox*²²⁸ allele, and (d) a fragment including a portion of a cell-specific polypeptide ligand, such portion including at least a portion of the binding domain of the polypeptide ligand, such portion of the binding domain being effective to cause the hybrid protein to bind selectively to a predetermined class of cells to be attacked by enzymatically active Fragment A.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	LI	Liechtenstein
AU	Australia	LK	Sri Lanka
BE	Belgium	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Republic	MG	Madagascar
CG	Congo	MR	Mauritania
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SE	Sweden
FR	France	SN	Senegal
GA	Gabon	SU	Soviet Union
GB	United Kingdom	TD	Chad
HU	Hungary	TG	Togo
JP	Japan	US	United States of America
KP	Democratic People's Republic of Korea		

HYBRID PROTEINS

This invention was made in the course of work supported by the U.S. government, which has certain rights in the invention.

This application is a continuation-in-part of Murphy U.S. Patent Application S.N. 377,386.

5 This invention relates to the use of recombinant DNA techniques to make hybrid protein molecules, and to the use of such molecules in the treatment of medical disorders.

The literature contains many examples of fused
10 genes which code for hybrid proteins. For example, Villa-Komaroff et al. (1978) P.N.A.S. U.S.A. 75, 3727-3731 describes a fused gene made up of a eukaryotic structural gene fused to a non-cytoplasmic bacterial gene. The fused gene codes for a hybrid protein which
15 is transported out of the cytoplasm.

Hybrid proteins have been also made by methods, e.g. the coupling of two different protein molecules, which do not involve recombinant DNA techniques. For example, it has been proposed to form, by coupling,
20 therapeutic hybrid proteins consisting of a toxin coupled to a ligand capable of binding specifically to a selected class of cells. One attempt to make such a hybrid protein, reported in Chang et al. (1977) J. Biol. Chem. 252, 1515-1522, resulted in a hybrid consisting of
25 the diphtheria toxin A chain coupled to human placental lactogen hormone by cross-linking through a disulfide bond. The hybrid protein, although it bound to cells containing lactogen receptors, did not inhibit protein synthesis in those cells.



- 2 -

A hybrid protein consisting of ricin A toxin coupled to the β chain of human chorionic gonadotropin hormone by similarly cross-linking through a disulfide bond has also been reported; although said to have
5 specificity, its binding capacity has not been reported, and extremely high concentrations were required to significantly inhibit protein synthesis in rat Leydig tumor cells, making it difficult to distinguish between
10 "non-specific" entry caused by endocytosis and "specific" entry caused by transport of the toxic portion of the hybrid across the cytoplasmic membrane of the target cells. Oeltman et al. (1979) J. Biol. Chem., 254, 1028-1032. The same shortcoming was found in a hybrid consisting of diphtheria A coupled to insulin
15 using cystamine as the cross-linking agent. Miskimins et al. (1979) Biochem. Biophys. Res. Commun., 91, 143-151. A hybrid consisting of ricin A coupled to epidermal growth factor (EGF) by means of a heterobifunctional cross-linker has also been made, but
20 the binding characteristics provided by the EGF are not limited to specific cells, but encompass a wide variety of cell types. Cawley et al. (1980) Cell, 22, 563-570.

It has now been found that a superior diphtheria toxin/hormone hybrid protein can be made in
25 which the protein is synthesized as a single unit; i.e., fragments are joined together not by cross-linking but by peptide bonds. The invention will be best understood by referring to the drawings, in which

30 Fig. 1 is a diagrammatic representation of the diphtheria toxin molecule;



- 3 -

Fig. 2 is a diagrammatic representation of a hybrid protein molecule of the invention;

Fig. 3 is a restriction map showing the location and orientation of the diphtheria tox operon on the 3.9 kb BamH-I restriction fragment of corynephage β^{tox} (including a site, NRU I, not found on the wild-type tox allele, but only on the mutant tox²²⁸ allele, as will be explained in more detail below); and

Fig. 4 is a diagrammatic representation of a fused gene of the invention, encoding a hybrid protein of the invention (the gene fragments are labeled in terms of the encoded protein fragments).

Referring to Fig. 1, the diphtheria toxin molecule consists of several functional "domains" which can be characterized, starting at the amino terminal end of the molecule, as hydrophobic leader signal sequence, enzymatically active Fragment A, the fourteen amino acid exposed protease sensitive disulfide loop (DSL) l_1 , containing a cleavage domain, and Fragment B, which includes hydrophobic domain h, DSL l_2 , and carboxy terminal end a make up a.

The process by which diphtheria toxin intoxicates sensitive eukaryotic cells involves at least the following steps: (i) diphtheria toxin binds to specific receptors on the surface of a sensitive cell; (ii) while bound to its receptor, the toxin molecule is internalized in an endocytic vesicle; (iii) either prior to internalization, or within the endocytic vesicle, the toxin molecule is cleaved (or processed) at a site in the region of 47,000 daltons from the N-terminal end;



- 4 -

(iv) as the pH of the endocytic vesicle decreases to below 6, the structural intermediate form of toxin, while still bound to its receptor, is inserted into the membrane; (v) once embedded in the membrane the hydrophobic domain h forms a pore; (vi) a proteolytic cleavage in l_1 , between Fragment A and B, occurs; (vii) thereafter, Fragment A, or a polypeptide containing Fragment A, is released into the cytosol; (viii) the catalytic activity of Fragment A, i.e., the nicotinamide adenine dinucleotide - adenosine diphosphate ribosylation of Elongation Factor 2, causes the death of the intoxicated cell. It is apparent that a single molecule of Fragment A introduced into the cytosol is sufficient to kill the cell.

The hybrid proteins of the invention include, in sequential order, beginning at the amino terminal end of the hybrid protein, the following peptide fragments, joined together by peptide bonds:

a) the enzymatically active Fragment A of diphtheria toxin (without the leader Fragment s, which is clipped during secretion of the protein),

b) a fragment including the cleavage domain l_1 adjacent said Fragment A of diphtheria toxin,

c) a fragment comprising at least the portion of Fragment B of diphtheria toxin encoded by the portion of the Fragment B encoding gene fragment of the tox operon between l_1 and the position about 90 base pairs upstream from the position on the tox operon of the NRU I site of the tox²²⁸ allele, and



- 5 -

d) a fragment comprising a portion of a cell-specific polypeptide ligand, the portion including at least a portion of the binding domain of the polypeptide ligand, the portion of the binding domain being effective to cause the hybrid protein to bind selectively to a predetermined class of cells to be attacked by enzymatically active Fragment A of diphtheria toxin. The necessary portion of the toxin molecule included is depicted as the portion of the molecule between lines y and x in Fig. 1. Preferably, the hybrid protein also includes protease sensitive DSL₁₂ of the toxin molecule, i.e. the portion of the molecule between lines x and z is also included. Line z is preferably at the point 47 amino acids from the carboxy terminal end of B', i.e., at the end of l₂, not closer, to ensure that the generalized eukaryotic binding site of Fragment B is excluded, so that binding will be controlled by the binding domain of the cell-specific ligand. It has been demonstrated that a little more than one-half of Fragment B must be provided for the molecule to act as an effective toxin.

Referring to Fig. 2, a diagrammatic representation of a hybrid protein molecule of the invention, the y-z portion of the diphtheria toxin molecule is joined, by a peptide bond, to Fragment r of a cell-specific polypeptide ligand; i.e. a polypeptide which selectively binds to a predetermined class of cells which are to be attacked by enzymatically active



- 6 -

Fragment A of the diphtheria toxin molecule. Fragment r can consist of the entire ligand, or a portion of the ligand which includes the entire binding domain of the ligand, or an effective portion of the binding domain.

When the ligand being used is large, it is
5 desirable that as little of the non-binding portion as possible of the ligand be included, so that the binding domain of the molecule is positioned close to the hydrophobic domain h of Fragment B. It is also desirable to include all or most of the binding domain
10 of the ligand molecule. In the case of α melanocyte stimulating hormone (MSH), which is a small peptide of thirteen amino acids, or β MSH, which contains seventeen amino acids, the portion of the molecule consisting of nine amino acids at the carboxy terminal
15 end of the molecule can be used, or the entire molecule can be used.

The regions within cell-specific ligands in which the binding domain is located are now known for a number of such ligands. Furthermore, recent advances in
20 solid phase polypeptide synthesis can enable those skilled in this technology to determine the binding domain of practically any such ligand, by synthesizing various fragments of the ligand and testing them for the ability to bind to the class of cells to be attacked.

25 The hybrid protein molecules of the invention, are virtually non-toxic to all mammalian cells except the cells of the specific class to which the ligand



- 7 -

binding domain binds. Thus, the hybrid proteins of the invention are much more specific than many other therapeutic agents, e.g., general cytotoxic anti-cancer drugs.

5 The hybrid proteins of the invention, in which fragments are joined via peptide bonds, are also superior to cross-linked hybrids because the proteins of the invention can be provided in a homogeneous sample, in which all of the identical molecules are effective and selective for a particular class of cells.

10 The specific class of cells which are bound and attacked by the hybrid proteins of the invention is determined by the specific polypeptide ligand which supplies the binding domain of the hybrid molecule. Any cell-specific polypeptide ligand can be used which has a
15 binding domain which is specific for a particular class of cells which are to be attacked. Polypeptide hormones are useful such ligands. Hybrids made using a portion of the binding domain of α or β MSH, for example, selectively bind to melanocytes, rendering the hybrids
20 useful in the treatment of melanoma. Other ligands provide different specificities; e.g., the binding domain of substance P recognizes receptors on the surfaces of neurons involved in the transmission of pain, so that hybrids made using substance P can be used
25 to destroy such neurons to relieve chronic pain. These hybrids can also be used to map areas of the nervous system containing substance P receptors. Other specific-binding ligands which can be used include somatostatin, interleukin I, interleukin II, and



- 8 -

interleukin III. Interleukin II is of particular importance because of its role in allergic reactions and autoimmune diseases such as lupus, involving activated T cells. In addition, since all of the interleukins are specific for T cells, hybrids made with them could be used to treat cancers involving the immune system, and to inhibit the rejection of transplanted organs. Other useful polypeptide ligands having cell-specific binding domains are follicle stimulating hormone (specific for ovarian cells); luteinizing hormone (specific for ovarian cells); thyroid stimulating hormone (specific for thyroid cells); vasopressin (specific for uterine cells, as well as bladder and intestinal cells); prolactin (specific for breast cells); and growth hormone (specific for certain bone cells).

The hybrid proteins of the invention are preferably prepared using recombinant DNA techniques involving forming the desired fused gene coding for the hybrid protein, and then expressing the fused gene, using conventional procedures. Referring to Fig. 3, the location and orientation of the diphtheria tox operon on the 3.9 kb BamH-I restriction fragment of ϕ tox⁺ allows the tox operon to be cleaved at a desired location, and the desired portion of the operon to be fused with the desired portion of the gene for a selected polypeptide ligand. A more detailed description of the tox operon, and a description of the cloning of Fragment A, are contained in Leong et al. (1983) Science 220, 515, hereby incorporated by reference. Fragment A, cloned as described therein to make plasmid pDT201, was deposited in the American Type Culture Collection, Rockville, MD on May , 1983, and has been given ATCC Accession No. _____.



- 9 -

Referring to Figs. 3 and 4, the portion of the diphtheria tox operon (Fig. 3) used to make the fused gene (Fig. 4) encoding the hybrid proteins of the invention is preferably the portion indicated by the dotted lines delineating Fragment D; i.e. the portion of the tox gene from the first Sau 3AI site to the SPH I site. Fragment D thus includes the 831 base pair (bp) gene fragment encoding Fragment A (including the 177 bp sequence ahead of the structural gene which includes the promoter and the portion encoding the signal sequence, S in Fig. 1); the portion of the gene fragment encoding the hydrophobic domain of Fragment B; and the gene fragment encoding DSL 1₂.

As shown in Fig. 3, the first 177 bp of the gene fragment encoding Fragment A includes, ahead of the tox promoter, some DNA which is not part of the tox operon. This DNA is irrelevant and is not transcribed; it is included only because the Sau 3AI site at the start of the Fragment A encoding gene fragment is the most convenient restriction site near the tox promoter (which is depicted as the line just to the left of Hind III).

It should be possible to obtain Fragment D by simply excising it from the tox operon via cleavage at Sau 3AI and SPH I. Alternatively, gene fragments can be fused together as follows. First the gene fragment encoding Fragment A is obtained. Next, the gene fragment encoding most of Fragment B, from Sau 3AI to Sau 3AI (B' of Fig. 3) is obtained. The gene fragment encoding Fragment B' has been cloned in plasmid pUC8, to make plasmid pDT301, and was deposited in the American Type Culture Collection, Rockville, MD on May , 1983, and was given ATCC Accession No. _____. Fragment B'



- 10 -

is cut back, using enzyme Bal31, about 200 bp, to the position of the NRU I site on tox²²⁸ (the wild-type tox allele does not have an NRU I site), to give B'' (Fig. 3). Fragments A and B'' are fused, a fragment encoding l₂ is fused to B'', and a fragment encoding the desired portion of the polypeptide ligand is fused to l₂.

In the above scheme, the Fragment B' encoding gene fragment is preferably cut back to the precise location of NRU I, but can also be cut back to any location between the location 90 bp ahead of NRU I, and the end of the l₂ encoding region, (i.e., within the 208 bp region between the location 90 bp ahead of NRU I, and the beginning of the l₂ encoding region). Or, the B' encoding fragment could be cut back to the carboxy terminal end of the l₂ encoding region (129 bp downstream from NRU I), in which case fusion of a synthetic l₂ encoding region is unnecessary. (It should be evident that, when the B' encoding region is cut back and a synthetic l₂ encoding fragment is fused to it, there generally will be a small number of base pairs normally found between NRU I and SPH I which will not be present in the fused gene, so that, strictly speaking, not all of D of Fig. 3, or all of y-z of Figs. 1 and 2, will necessarily be included.)

An alternative method, less preferred than the scheme above, is to fuse the ligand-encoding gene fragment directly to B'', without employing the l₂ encoding fragment.

The fused gene, either including or omitting the l₂-encoding fragment, can alternatively be made using the B'' encoding gene fragment from the mutant



- 11 -

tox²²⁸ allele, rather than the wild-type allele. The tox²²⁸ allele, containing the NRU I site, is easily processed to yield the B'' encoding fragment. The tox²²⁸ allele is described in Uchida et al. (1973) Jour. Biolog. Chem. 248, 3831, hereby incorporated by reference.

In more detail, fused genes encoding hybrid proteins of the invention can be made as follows.

Vectors

10 The preferred vectors are plasmids pUC7, pUC8, pUC9, and pBR322. The pUC plasmids, described in Viera et al. (1982) Gene 19, 259 (hereby incorporated by reference) are particularly well-suited for double digest cloning, a procedure which permits the fused
15 genes to be made unambiguously.

Fused gene

Below is a flow chart for constructing diphtheria toxin-MSH fused genes containing the protease sensitive loop l_2 between the tox sequences and the
20 ligand (in this case, MSH) sequences.

- (i) pDT201 $\xrightarrow{\text{Sau3A1 digest}}$ purify Sau3A1-2 (Fragment A)
- (ii) pDT301 $\xrightarrow{\text{HindIII digest}}$ $\xrightarrow{\text{Bal31 cutback}}$ $\xrightarrow{\text{Pst1 linkers}}$ $\xrightarrow{\text{Sau3A1 digest}}$
- 25 $\xrightarrow{\text{reclone Sau3A1-Pst1 in pUC8}}$ Select blue colonies on X-G for proper reading frame $\xrightarrow{\text{purify Sau3A1-Pst1}}$ (Fragment B'')
- 30 (iii) in vitro synthesis of protease sensitive loop l_2 (Pst1-loop-EcoR1) \rightarrow clone into Pst1-EcoR1 sites on pUC9 & purify fragment
- (iv) clone (ii) Sau3A1-Pst1 and (iii) Pst1-EcoR1 into the BamH1-EcoR1 sites on pCUB (Sau3A1-Fragment B''-loop-EcoR1)



- 12 -

- (v) clone (iv) Sau3AI-EcoRI and MSH sequence into the BamHI site on pUC8 (Sau3AI-Fragment B''-loop-MSH-BamHI)
- (vi) clone (i) Sau3AI-Fragment A-Sau3AI and (v) Sau3AI-Fragment B''-loop-MSH-BamHI into the BamHI site on pUC8 (Sau3AI-Fragment A-Fragment B''-loop-MSH-BamHI)

Referring to the above flow chart, in step (i), the Fragment A encoding gene fragment is first obtained and purified, as described in Science, Id. In step (ii) the Fragment B'' encoding fragment is obtained by cutting back a larger Fragment B' encoding region using the enzyme Bal31.

As shown in Fig. 3 the gene fragment encoding Fragment B' is the 1,023 bp region between two Sau3AI sites, the first of which is at the DNA sequence encoding the third arginine in l₁, and the second of which is 49 bp before the end of the tox structural gene. The gene fragment encoding B' has, as is mentioned above, been cloned in pUC8. The plasmid carrying this fragment in the same orientation as the lac Z gene has been designated pDT301; (the lac Z and B' genes are out of frame on pDT301).

Referring again to the above flow chart, pDT301 is opened via HindIII digestion and then the B' encoding fragment is cut back by exposure to the exonuclease Bal31 for varying time periods. The ends of the resulting shortened gene fragments (of varying lengths) are then blunt-end ligated with EcoRI and PstI linkers and the fragments are then digested with SauRI. This results in a heterogeneous (in terms of size) population of gene fragments encoding part of B'.

These fragments are then recloned in either the BamHI-PstI or the BamHI-EcoRI sites of pUC8. Cloning in these sites allows selection only of clones having BamHI (Sau3AI) on one end and PstI or EcoRI on the other.



- 14 -

 β -MSH

Asp-Glu-Gly-Pro-Tyr-Met-Glu-His-Phe-
 GAT-GAA-GGT-CCA-TAT-ATG-GAG-CAC-TTT
 C G X X C A T C

5 Arg-Trp-Gly-Ser-Pro-Pro-Lys-Asp
 AGA-TGG-GGT-TCT-CCG-CCG-AAA-GAT
 G X X X X TC C

EcoRI
 linker

stop codon &
BamHI linker

10 The unique PstI and EcoRI sites of pUC8 permit the subcloning of either of the above synthetic MSH sequences, downstream from the l_2 encoding fragment.

15 Finally (step vi) the Fragment A encoding gene fragment is fused to the gene fragment encoding B''- l_2 -MSH, to complete the gene fusion as illustrated in Fig. 4 (labeled in terms of encoded protein fragments), which codes for a hybrid protein which selectively binds to and attacks a selected class of cells (in this case, melanocytes).

20 Another example of a suitable polypeptide ligand is substance P, the utility of which is described above. A fused gene containing the substance P gene, rather than the α or β MSH gene, is made basically as outlined above. The substance P gene is synthesized using conventional solid phase oligonucleotide
 25 synthesis. The substance P gene sequence is:
 CGTCCTAAACCTCAGCAGTTCTTCGGTCTGATG.

30 As is clear from the above, the portion of the genetic sequence for the polypeptide ligand must be sufficient to enable the corresponding amino acid sequence to cause the hybrid protein to bind to the predetermined class of cells. Preferably, the gene for the polypeptide hormone will include all or most of the genetic sequence for the binding domain of the ligand.



- 15 -

Generally, as in the above examples, the manipulative operations are carried out using cloning vectors; e.g., phages or plasmids. The genetic material coding for the binding domain of the polypeptide ligand can be either cloned DNA or a synthetic oligonucleotide sequence, whichever is more convenient for the particular ligand gene employed. Generally the fused gene will reside on a cloning vector, e.g., a plasmid or a phage, which is used to transform cultured microorganisms. The hybrid protein is then harvested from the culture media of the cells using conventional techniques.

The hybrid proteins of the invention are administered to a mammal, e.g., a human, suffering from a medical disorder, e.g., cancer, characterized by the presence of a class of unwanted cells to which a polypeptide ligand can selectively bind. The amount of protein administered will vary with the type of disease, extensiveness of the disease, and size and species, of the mammal suffering from the disease. Generally, amounts will be in the range of those used for other cytotoxic agents used in the treatment of cancer, although in certain instances lower amounts will be needed because of the specificity of the hybrid proteins.

The hybrid proteins can be administered using any conventional method; e.g., via injection, or via a timed-release implant. In the case of MSH hybrids, topical creams can be used to kill primary cancer cells, and injections or implants can be used to kill metastatic cells. The hybrid proteins can be combined with any non-toxic, pharmaceutically-acceptable carrier substance.



- 16 -

What is claimed is:

1. A hybrid protein comprising protein fragments joined together by peptide bonds, said hybrid protein comprising, in sequential order, beginning at the amino terminal end of said hybrid protein,
 - 5 a) the enzymatically active Fragment A of diphtheria toxin,
 - b) a fragment including the cleavage domain l_1 adjacent said Fragment A of diphtheria toxin,
 - c) a fragment comprising at least the portion
10 of Fragment B of diphtheria toxin encoded by the portion of the Fragment B encoding gene fragment of the tox operon between l_1 and the position about 90 base pairs upstream from the position on said tox operon of the NRU I site of the tox ²²⁸ allele, and
 - 15 d) a fragment comprising a portion of a cell-specific polypeptide ligand, said portion including at least a portion of the binding domain of said polypeptide ligand, said portion of said binding domain being effective to cause said hybrid protein to bind
20 selectively to a predetermined class of cells to be attacked by said enzymatically active Fragment A.
2. The hybrid protein of claim 1 wherein said
25 fragment c) is encoded by the portion of the Fragment B encoding gene fragment between l_1 and said position of the NRU I site.
3. The hybrid protein of claim 1 or claim 2, further comprising fragment l_2 between fragments c) and d).



- 17 -

4. The hybrid protein of claim 1 wherein said hybrid protein is coded for by a fused gene comprising regions coding for said protein fragments.

5 5. The hybrid protein of claim 1 wherein said polypeptide ligand is a hormone.

10 6. The hybrid protein of claim 5 wherein said portion of said polypeptide hormone is a portion of α or β melanocyte stimulating hormone effective to cause said hybrid protein to bind to malignant melanocyte cells.

7. The hybrid protein of claim 5 wherein said portion of said polypeptide hormone is a portion of substance P effective to cause said hybrid protein to bind to pain receptor neurons.

15 8. The hybrid protein of claim 5 wherein said portion of said polypeptide hormone is a portion of interleukin I effective to cause said hybrid protein to bind to T cells.

20 9. The hybrid protein of claim 5 wherein said portion of said polypeptide hormone is a portion of interleukin II effective to cause said hybrid protein to bind to T cells.

25 10. The hybrid protein of claim 5 wherein said portion of said polypeptide hormone is a portion of interleukin III effective to cause said hybrid protein to bind to T cells.



- 18 -

11. A fused gene comprising a fragment of the gene coding for diphtheria toxin fused to a fragment of a gene coding for a cell-specific polypeptide ligand, said fused gene coding for a hybrid protein, said fragment of said diphtheria toxin gene including the fragment coding for the hydrophobic leader fragment of said diphtheria toxin, the region coding for the enzymatically active Fragment A of said diphtheria toxin, the region coding for protease sensitive loop l_1 adjacent said Fragment A of said diphtheria toxin, and at least a portion of the Fragment B encoding fragment about 90 base pairs upstream from the position on the diphtheria tox operon of the NRU I site of the tox²²⁸ allele,
- said fragment of a gene coding for said polypeptide ligand encoding a portion of said ligand effective to cause said hybrid protein to bind selectively to a predetermined class of cells.

12. The fused gene of claim 11 wherein said polypeptide ligand is α or β melanocyte stimulating hormone.

13. A method of treating a mammal suffering from a medical disorder characterized by the presence of a class of unwanted cells, said method comprising administering to said mammal the hybrid protein of claim 1 wherein said binding domain is specific for said class of unwanted cells, said hybrid protein being administered in an amount effective to damage at least some of said cells.



1/1




FIG 1




FIG 2




FIG 3




FIG 4



INTERNATIONAL SEARCH REPORT

International Application No PCT/US83/00723

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ³		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int. CL. ³ A61K 37/00,	A61K 37/48,	CO7C 103/52
US CL. 424/177,	260/112.5R	
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
US	435/68, 435/70, 435/91, 435/172, 435/253 435/317, 536/27	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁵		
Chemical Abstracts and Biological Abstract, 1970 to present		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁵ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
P	US, A 4,394,443 issued July 19, 1983 Weissman et al.	11&12
P	US, A 4,336,336 issued June 22, 1982 Silhavy et al.	11&12
P	US, A 4,350,626 issued September 22, 1983 Masuho et al.	1-10 & 13
X	US, A 4,302,386 issued November 24, 1981 Stevens	1-10 & 13
T	US, A 4,383,995 issued May 24, 1983 Stevens	1-10 & 13
X	US, A 4,275,000 issued June 23, 1981 Ross	1-10 & 13
P	US, A 4,379,145 issued April 5, 1983 Masuho et al.	1-10 & 13
<p>¹⁵ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ¹	Date of Mailing of this International Search Report ²	
29 Aug 1983	20 SEP 1983	
International Searching Authority ¹	Signature of Authorized Officer ²⁰	
ISA/US	<i>Dellert R. Phillips</i>	

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹⁰

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. Claim numbers because they relate to subject matter ¹² not required to be searched by this Authority, namely:
2. Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

VI. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ¹¹

This International Searching Authority found multiple inventions in this international application as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- The additional search fees were accompanied by applicant's protest.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM THE FIRST SHEET
(Not for publication)

- | | | | |
|---|----|---------------------------------------------------------------------------------------------------------------------------|------|
| X | N, | The Journal of Biological Chemistry
Vol. 252, No. 4, Issued February 25, 1977
pages 1505-1514, Chang, et al. | 1-13 |
| X | N, | The Journal of Biological Chemistry
Vol. 256, No. 24, Issued December 25, 1981
pages 12731-12739, Gilliland, et al. | 1-13 |
| X | N, | Proceeding National Acad. Science
Vol. 75, No. 8, Issued August 1978,
pages 3727-3731, Komaroff, et al. | 1-13 |