

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2014320757 B2**

(54) Title
Production of oligosaccharides

(51) International Patent Classification(s)
C12P 19/14 (2006.01) **C12P 19/26** (2006.01)
C07H 1/06 (2006.01)

(21) Application No: **2014320757** (22) Date of Filing: **2014.07.04**

(87) WIPO No: **WO15/036138**

(30) Priority Data

(31) Number	(32) Date	(33) Country
13183670.2	2013.09.10	EP

(43) Publication Date: **2015.03.19**

(44) Accepted Journal Date: **2018.04.19**

(71) Applicant(s)
Jennewein Biotechnologie GmbH

(72) Inventor(s)
Jennewein, Stefan

(74) Agent / Attorney
FB Rice Pty Ltd, L 23 44 Market St, Sydney, NSW, 2000, AU

(56) Related Art
US 20120208181 A1
US 20040121431 A1
US 20110300584 A1



- (51) International Patent Classification:
C12P 19/14 (2006.01) C07H 1/06 (2006.01)
C12P 19/26 (2006.01)
- (21) International Application Number:
PCT/EP2014/064280
- (22) International Filing Date:
4 July 2014 (04.07.2014)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
13183670.2 10 September 2013 (10.09.2013) EP
- (71) Applicant: JENNEWEIN BIOTECHNOLOGIE GMBH
[DE/DE]; Maarweg 32, 53619 Rheinbreitbach (DE).
- (72) Inventor: JENNEWEIN, Stefan; Am Burgberg 108,
52080 Aachen (DE).
- (74) Agent: WITTE, WELLER & PARTNERPATENTAN-
WÄLTE MBB / ZUSAMMENSCHLUSS EPA NR. 314;
Postfach 10 54 62, 70047 Stuttgart (DE).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

- with international search report (Art. 21(3))

[Continued on next page]

(54) Title: PRODUCTION OF OLIGOSACCHARIDES

Fig. 9



(57) Abstract: The present invention relates to the use of one or more glycosidases in the process for the production and/or purification of a produced desired oligosaccharide. The process is preferably a microbial fermentation process using a host microorganism, which may also comprise nucleic acids expressing sugar catabolic pathway proteins suitable for the degradation of saccharides otherwise hindering the purification of the desired oligosaccharide.

— *with sequence listing part of description (Rule 5.2(a))*

Production of oligosaccharides

[0001] The present invention generally relates to the processes for the production of oligosaccharides or polysaccharides, such as, e.g., the production of oligosaccharides or polysaccharides via microbial fermentation, and in particular to the use of hydrolases in such processes.

[0002] The commercial production of oligosaccharides has – together with the development of novel manufacturing biotechnologies – become of increasing interest over the past years. Today, oligosaccharides are used, e.g., as functional food ingredients, nutritional additives, or as nutraceuticals. Traditionally, oligosaccharides are defined as polymers of 2 or more (usually up to 10) monosaccharides, however, also polymers up to 20 to 25 are often assimilated with them. In particular prebiotic oligosaccharides are of high interest, since they represent noncariogenic and nondigestible compounds stimulating the growth and development of human gastrointestinal microflora.

[0003] Currently, oligosaccharides are either synthesized by chemical glycosylation and *de novo* using glycosyltransferases, or they can derive from chemical, physical or biological degradation of polysaccharides.

[0004] Today, fructooligosaccharides and galactooligosaccharides synthesized or isolated from plant polysaccharides represent the most abundantly produced oligosaccharides. However, due to superior health benefits the interest in Human Milk Oligosaccharides (HMO) as nutraceuticals has strongly increased over the past years. It is well established that HMOs contribute to the human body's defense mechanism against human pathogens, the establishment of a particular intestinal flora (microbiome), and postnatal stimulation of the immune system. Although HMOs are consumed by infants, it is accepted that the beneficial effects of these oligosaccharides observed during the postnatal period can be also found later in life.

[0005] It has been long known that human mother's milk, besides lactose, comprises a complex mixture of oligosaccharides called Human Milk Oligosaccharides (HMO) represents a complex mixture of oligosaccharides being unique with respect to composition and quantity. Today, more than 80 HMO compounds have been structurally characterized, and, with few exceptions, they are characterized by a lactose moiety at the reducing end and often containing a fucose and/or sialic acid at the non-reducing end. Generally the monosaccharides from which HMOs are derived are D-glucose, D-galactose, *N*-acetylglucosamine, L-fucose and sialic acid.

[0006] The most prominent oligosaccharides are 2'-fucosyllactose and 3'-fucosyllactose which together can contribute up to 1/3 of the total HMO fraction. Further prominent HMOs present in human milk are lacto-*N*-tetraose, lacto-*N*-neotetraose and the lacto-*N*-fucopentaoses. Besides these neutral oligosaccharides, also acidic HMOs can be found in human milk, like for e.g. 3'-sialyllactose, 6'-sialyllactose and 3-fucosyl-3'-sialyllactose, disialyl-lacto-*N*-tetraose etc. These structures are closely related to epitopes of epithelial cell surface glycoconjugates, the Lewis histoblood group antigens such as Lewis x (LeX), and the structural homology of HMO to epithelial epitopes accounts for protective properties against bacterial pathogens.

[0007] Besides the mentioned local effects in the intestinal tract, HMOs have also been shown to elicit systemic effects in infants by entering the systemic circulation. Also, the impact of HMOs on protein-carbohydrate interactions, e.g., selectin-leukocyte binding, can modulate immune responses and reduce inflammatory responses.

[0008] Due to the well-studied beneficial properties of prebiotic oligosaccharides, in particular of HMOs, connected with their limited availability, an effective commercial, i.e. large scale production is highly desirable.

[0009] The main drawback today, however, is in most cases the lack of effective oligosaccharide production. As mentioned above, oligosaccharides can be generated from oligomer engineering with either synthesis, using enzymatic or chemical engineering, or polysaccharide depolymerization, using physical, chemical or enzymatic methods.

[0010] Chemical synthesis of oligosaccharides has proved to be challenging due to the presence of several hydroxyl groups of similar chemical reactivity. Thus, in chemical synthesis of oligosaccharides saccharide building blocks have firstly to be selectively protected in order to control the reactivity of the chemically similar groups, then coupled, and finally de-protected to obtain the desired oligosaccharide. Even if the small-scale synthesis of a desired oligosaccharide is possible, the developed synthetic routes are in general time-consuming, technically challenging and often prohibitively expensive. Enzymatic synthesis or the combination of chemical and enzymatic synthesis offer significant advantages over pure chemical synthetic routes. Several human milk oligosaccharides such as 2'-fucosyllactose, lacto-*N*-tetraose lacto-*N*-neotetraose or 3'-sialyllactose were already obtained by enzymatic synthesis. Besides the enzymatic synthesis also fermentative approaches to oligosaccharides proved successful, and several oligosaccharides – including HMOs – have been made available in considerable yields by fermentative means.

[0011] Further, with a chemical or biochemical synthesis of oligosaccharidic structures, which, as mentioned before, is much more difficult than, e.g. the synthesis of other biopolymers such as peptides and nucleic acids, often oligosaccharides mixtures

are produced containing non-desired oligosaccharides which need to be separated or removed from the desired oligosaccharide. Same applies for oligosaccharides generated via microbial fermentation, since during those processes a high amount of by-products, intermediate products or even starting substrates is to be found in the reaction mixture/cell medium containing the to be fermented microorganisms.

[0012] Against this background, there exists a great need for effective methods and processes for the production of a desired oligosaccharide, in particular of HMOs, the efficient production bearing high medical relevance and as well commercial impact.

[0013] According to one embodiment the description provides for by the present invention by the use of one or more glycosidases in the purification of a produced desired oligosaccharide from a mixture containing the produced desired oligosaccharide and, where applicable, non-desired oligosaccharides or metabolic saccharide products generated during production of said desired oligosaccharide, and/or unused saccharide substrates used in the production of said oligosaccharide, for degradation of the non-desired oligosaccharides or metabolic products or unused substrates.

[0014] In another embodiment the description provides a process for producing an oligosaccharide using a host microorganism, wherein said oligosaccharide is not naturally occurring in said host cell, the process comprising the steps of a) cultivating a host microorganism suitable for the production of a desired oligosaccharide under conditions and in a medium permissive for the production of said oligosaccharide, whereby the oligosaccharide and, where applicable, biosynthetic intermediates and/or side products are produced; b) using a glycosidase in the medium the host microorganism is cultivated in, in order to degrade biosynthetic saccharide intermediates and/or saccharide side products and/or unused saccharide substrates, and c) recovering the desired oligosaccharide.

[0015] According to the invention, a glycosidase is applied in the process for the production of a oligosaccharide, wherein the glycosidase is used for degrading hindering and/or undesired side-products, unused starting substrates and intermediate products generated during the production of the desired oligosaccharide; thus, according

to the invention, the glycosidase is employed for purification of the desired oligosaccharide from a mixture containing the desired oligosaccharide and other unwanted carbohydrate moieties.

[0016] According to the invention, the glycosidase can be used in fermentation processes producing a desired oligosaccharide as well as for the purification or resolution of oligosaccharide mixtures obtained by *in vitro* oligosaccharide synthesis reactions or by chemical synthesis or combinations thereof.

[0017] By means of the glycosidase it can be achieved that, e.g. other (oligo-) saccharides - besides the desired oligosaccharide- , which other (oligo-)saccharides are produced in the microorganism during the synthesis of the desired oligosaccharide, and which other oligosaccharides interfere with the purification step of the desired oligosaccharide, can be metabolised.

[0018] The use of a glycosidase in a process for the production of a desired oligosaccharide has not been described so far. To the contrary: In the processes currently applied, i.e. for the fermentation of HMO, usually lactose is added as a substrate for the synthesis. To prevent the degradation of the added lactose beta-galactosidase, as well as other glycosidase reactions are strongly avoided in fermentation strains. Therefore, beta-galactosidase-deficient stains were applied in the fermentation (see, e.g. Dumon *et al.*, (2004) Biotechnol. Prog. 20, 412-419) or the beta-galactosidase gene of the strain has been specifically inactivated (Dumon *et al.*, (2006) ChemBioChem. 7, 359-365)). Generally, the presences of any glycosidase activity are seen as contra productive to the fermentation of oligosaccharides.

[0019] According to the invention, an "oligosaccharide" is to be understood to be short polymers of monosaccharides, comprising at least 2 sugar-subunits, as already mentioned at the outset. The oligosaccharides may either be branched or form a linear chain of subunits. Moreover, the sugar subunits of oligosaccharides may feature a number of chemical modifications. Accordingly, the oligosaccharides according to the present invention may comprise one or more non-sugar moieties.

[0020] Also, presently and generally in the relevant field, a “glycosidase” – also designated as “glycoside hydrolases” or “glycosyl hydrolases” catalyze the hydrolysis of the glycosidic linkage to release smaller sugars. They can be classified as enzymes catalyzing the hydrolysis of O- or S-glycosides, and they are typically named after the substrate that they act upon. Accordingly, glucosidases catalyze the hydrolysis of glucosides and xylanases catalyze the cleavage of xylan. Thus, a glycosidase according to the invention catalyses the hydrolysis of glycosidic bonds to liberate monosaccharides and oligosaccharides of lower molecular weight than the native simple as well complex carbohydrate substrates.

[0021] According to the invention, in the use and the processes disclosed herein a glycosidase is employed to degrade hindering and/or undesired side-products, unused starting substrates and intermediate products generated during the production of the desired oligosaccharide which undesired side-products, unused starting substrates and intermediate products generally have a higher molecular weight compared to the monosaccharides and oligo- or disaccharides liberated by the action of the glycosidase during degradation of the undesired side-products, unused starting substrates and intermediate products.

[0022] Thus, presently, a “liberated monosaccharide” is to be understood as a monosaccharide that has been generated during the glycosidase-mediated hydrolysis of a glycosidic bond of a carbohydrate substrate where monosaccharides and oligosaccharides of lower molecular weight than the hydrolyzed carbohydrate substrate are formed.

[0023] According to the invention, at least one glycosidase, or a combination of two or more may be applied in the use and the process according to the invention.

[0024] According to one aspect of the invention, the glycosidase is used in a microbial fermentation process employed for the production of the desired oligosaccharide and using a host microorganism, wherein said oligosaccharide and/or said glycosidase are not naturally occurring in the microorganism to said host cell.

[0025] Presently and as generally understood in the related field, a “microbial fermentation” is to be understood as a – generally large-scale – industrial metabolic process where during cultivation of microorganisms such as bacteria, fungi and moulds, enzymatic decomposition and utilization of nutrients, particularly carbohydrates, and the conversion of compounds into other compounds takes place. As such, industrial or large-scale microbial fermentation is the process of controlling microorganisms, i.e. bacteria, yeast, and moulds, to modify food, producing a desired product.

[0026] According to another aspect of the invention, the glycosidase is used for resolution of oligosaccharide mixtures obtained by in vitro oligosaccharide synthesis reactions or by chemical synthesis or by combinations thereof.

[0027] Presently, and generally in the relevant field, a “microorganism” presently designates and encompasses any microscopic organism that comprises either a single cell, cell clusters, or multicellular relatively complex organisms, which is suitable to be employed in the process according to the invention, and particularly includes bacteria and yeast. A microorganism as employed according to the invention can be cultivated in a liquid medium, and generally needs a carbon source in the medium to grow and replicate.

[0028] Presently, and throughout the invention, “recombinant” means genetically engineered DNA prepared by transplanting or splicing genes from one species into the cells of a host microorganism of a different species. Such DNA becomes part of the host's genetic makeup and is replicated.

[0029] Consequently, “a host microorganism” is designated to mean any microorganism containing nucleic acid sequences or expressed proteins foreign to/not naturally occurring in the thus recombinant host microorganism and wherein the foreign/not naturally in said microorganism occurring nucleic acid sequence is integrated in the genome of the host microorganism cell. Thereby, “not naturally occurring” means that the nucleic acid sequence/protein (such as, e.g. an enzyme), is foreign to said host microorganism cell, i.e. the nucleic acid sequences/proteins are heterologous with respect to the microorganism host cell. The heterologous sequence may be stably introduced, e.g. by

transfection, transformation, or transduction, into the genome of the host microorganism cell, wherein techniques may be applied which will depend on the host cell the sequence is to be introduced. Various techniques are known to a person skilled in the art and are, e.g., disclosed in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Thus, the host microorganism the heterologous sequence has been introduced in will produce the heterologous proteins the nucleic acid sequences according to the invention are coding for.

[0030] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof and the nucleic acid sequences of the invention. Introduction of a nucleic acid sequence into the host microorganism cell can be effected by methods described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods in Molecular Biology*, (1986), and Sambrook *et al.*, 1989, *supra*.

[0031] Thus, the nucleic acid sequences according to the invention, may, e.g., be comprised in a vector which is to be stably transformed/transfected or otherwise introduced into host microorganism cells.

[0032] A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and to synthesize a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *supra*.

[0033] Presently, a “desired oligosaccharide” designates an oligosaccharide that shall specifically and intentionally be produced with the applied process. Accordingly, a “non-desired oligosaccharide” represents an oligosaccharide not intended to be produced or generated during the production of the desired oligosaccharide or an oligosaccharide formed by the used organism unrelated to the desired oligosaccharide. A “metabolic product” or “metabolic” saccharide product” or “saccharide intermediate” is a saccharide, i.e. carbohydrate product generated during production of a desired oligosaccharide, and an “unused saccharide substrate” represents a starting molecule or moiety used for/during the production of the desired oligosaccharide.

[0034] As mentioned at the outset, the invention also relates to a process for producing an oligosaccharide using a host microorganism, wherein said oligosaccharide is not naturally occurring in said host cell, the process comprising the steps of a) cultivating a host microorganism suitable for the production of a desired oligosaccharide under conditions and in a medium permissive for the production of said oligosaccharide, whereby the oligosaccharide and, where applicable, biosynthetic saccharide intermediates and/or saccharide side products are produced; b) using a glycosidase in the medium the host microorganism is cultivated in, in order to degrade biosynthetic intermediates and/or side products and/or unused substrates, and c) recovering the desired oligosaccharide.

[0035] With the process according to the invention, an effective manufacturing method has been provided, by means of which a desired oligosaccharide can be produced in a form where non-desired oligosaccharides or otherwise hindering intermediate products or unused substrates are essentially no longer present.

[0036] As used herein, the term “cultivating” means growing a microorganism in a medium and under conditions permissive and suitable for the production of the desired oligosaccharide. A couple of suitable host microorganisms as well as the mediums and conditions for their cultivation will be readily available for one skilled in the art upon reading the disclosure of this invention in connection with the skilled person’s technical and expert background.

[0037] As used herein, the term "recovering" means isolating, harvesting, purifying, collecting or otherwise separating from the host microorganism culture the oligosaccharide produced by the host microorganism according to the invention.

[0038] According to one aspect of the invention, the glycosidase is added to the medium the microorganism is cultivated in. "Adding" hereby means the direct addition of the enzyme to the medium, i.e. contrary to a provision of the glycosidase that would be endogenously produced by the microorganism used for the production. As such, the enzyme is – in its active form – readily and commercially available on the market and can be added in amounts necessary to degrade the undesired oligosaccharides/unused substrates/intermediates. The amount to be applied will depend from the amount of the cultivated microorganism and also from amount of the produced desired oligosaccharide, the amount of substrate and likely to be expected intermediates.

[0039] The glycosidase(s) are, according to this embodiment, externally added to the medium/the supernatant at the end of the production process according to the invention, which is particularly preferred when endogenous genes of the host microorganism encoding glycosidases have been inactivated or deleted. In doing so, undesired oligo- and or monosaccharides cannot accumulate and do not interfere with the recovering of the desired oligosaccharide.

[0040] E.g., to a solution containing a desired oligosaccharide and a non-desired oligosaccharide, such as fermentation derived lacto-*N*-tetraose and lactose, beta-galactosidase is added; the amount of beta-galactosidase to be employed will depend on the (actual or expected) amount of oligosaccharides. E.g. with lacto-*N*-tetraose and lactose both at 10 mM concentration an amount of 50 units/ml (final concentration) of beta-galactosidase (e.g. from Sigma-Aldrich Chemie GmbH, Munich Germany catalog No. G6008) can achieve good results in order to cleave the lactose in glucose and galactose for better chromatographic separation of the saccharides (e.g. by size dependent gel-filtration chromatography). In respective experiments conducted by the inventors, it could be shown that within seconds the lactose was cleaved into its monosaccharides galactose and glucose, whereas the lacto-*N*-tetraose remained as such (see also Figure 1 and the respective description below). These experiments proved the highly selective action of the

purified *E. coli* beta-galactosidase, which selectively only hydrolyzes the Gal(beta1-4)Glc glycosidic bond and not the Gal(beta1-3)GlcNAc glycosidic bond present at the non-reducing end of the lacto-*N*-tetraose. It will be apparent from the present disclosure how much of the respective glycosidase is likely needed to be used.

[0041] E.g., suitable microorganisms for producing, e.g., fucosylated oligosaccharides in fermentation proceedings are known from EP 2 479 263 A1 or from EP 2 439 264 or from WO 2010/142305, the content of which is herewith explicitly referred to and made subject matter of this invention.

[0042] According to another aspect of the invention, the glycosidase is endogenously produced by said host microorganism, wherein the nucleic acid sequences encoding the endogenously produced glycosidase is not naturally occurring in said host cell, and wherein said host cell has been stably transformed to express the not naturally occurring glycosidase, and wherein the expression of the glycosidase in the host microorganism is inducible.

[0043] According to this embodiment, the oligosaccharide producing microorganism endogenously, i.e. contained in its genome, expresses the glycosidase, e.g. beta-galactosidase, upon external induction, e.g., via temperature or substrate-induced expression, which is otherwise deregulated. This means that during synthesis of the desired oligosaccharide, the glycosidase is deregulated and may be induced, e.g. by temperature or adding a inductor, e.g. tetracyclin, at the end of the fermentation process. The expression of the glycosidase will be induced after a sufficient and/or essentially maximum amount of oligosaccharide has been produced during cultivation of the host microorganism. Subsequently, the expressed glycosidase will degrade undesired saccharide intermediates, substrates, etc., rendering the medium essentially free of the saccharide intermediates or substrates that would otherwise hinder or complicate the purification of the desired oligosaccharide. A couple of suitable inducible expression tools are known in the prior art (see, e.g. Sambrook *et. al*, 1989, *supra*), and one skilled will be able to apply a respectively suitable one for the desired oligosaccharide.

[0044] “Regulated” within the present context with reference to a gene is generally understood as a gene, whose expression can be regulated in a controlled fashion, e.g. down- or up-regulated, i.e. the quantity of the synthesised protein encoded by the regulated gene is different, e.g. de-/downregulated or upregulated, from the otherwise unregulated gene.

[0045] According to another aspect of the invention, the glycosidase is produced by a second microorganism additionally added to the medium and expressing the glycosidase.

[0046] In this embodiment, the glycosidase expressed by the microorganism is either a naturally occurring glycosidase or a glycosidase that is encoded by a nucleic acid sequence that has been stably integrated into the genome of the second microorganism. This embodiment is particularly suitable in a continuous fermentation process for the production of the oligosaccharide, where, e.g., two separate fermentation vessels or containers are provided, whereas one vessel/container is used for the oligosaccharide synthesis reaction and the second vessel/container is essentially employed for the degradation of undesired saccharides.

[0047] Accordingly, and according to certain aspects of the invention, the process according to the invention is a batch or a continuous process.

[0048] Thus, according to one aspect of the invention, i.e. in a continuous process, the glycosidase is constantly added to the medium during the cultivating step of the host microorganism, e.g. by the provision of a strain expressing a glycosidase, whereas the glycosidic and catabolic reactions can be spatially separated.

[0049] According to another aspect, i.e. in a batch or feed-batch process, the hydrolytic cleavage reactions are restricted to particular periods.

[0050] According to another aspect of the invention, the oligosaccharide is recovered from supernatant of the cultivated host microorganism, which supernatant is

obtained by centrifuging the cultivated host microorganism to obtain a supernatant and a host microorganism pellet.

[0051] With the newly provided process, it is possible to retrieve the produced oligosaccharide from the medium the host microorganism is cultivated in, since the oligosaccharide which is produced in a microorganism cell is preferably transported into the medium, thus making it effortlessly possible to recover the oligosaccharide from the supernatant, once the cells of the microorganism have been separated from the cultivation medium.

[0052] Also, according to another aspect of the invention, prior to the adding of the glycosidase, a supernatant may be produced separating the host microorganisms from the preferably liquid medium the host microorganism is cultivated in, wherein the produced oligosaccharide is contained in the supernatant. To this supernatant, the glycosidase may be added.

[0053] According to another aspect, the desired oligosaccharide is selected from 2'-fucosyllactose (2'-FL), 3-fucosyllactose (3-FL), difucosyllactose (DF-L), 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), 3-fucosyl-3'-sialyllactose (F-SL), lacto-*N*-tetraose ((LNT), lacto-*N*-neotetraose (LN*neo*T), lacto-*N*-fucopentaose(s) (LNFP-I, II, III, V), lacto-*N*-difucohexaose(s) (LNDH-I and II), lacto-*N*-sialylpentaose(s) (LSTa, b and c), fucosyl-lacto-*N*-sialylhexaose (F-LSTa, b and c), disialyl-lacto-*N*-hexaose (DS-LNT) and in particular selected from those shown in table 1, or derivatives thereof. In this regard, for the production of the oligosaccharides, it is explicitly referred to EP 2 479 263 A1 or from EP 2 439 264 or from WO 2010/142305, the content of which is herewith explicitly referred to and made subject matter of this invention.

Table 1: list of oligosaccharides that can be produced according to the invention (Fig. 11)

No.	Name	Abbreviation	Oligosaccharide
1	2'-Fucosyllactose	2'-FL	Fuc(α 1-2)Gal(β 1-4)Gluc
2	3-Fucosyllactose	3-FL	Gal(β 1-4)Gluc Fuc(α 1-3)
3	2',3-Difucosyllactose	DF-L	Fuc(α 1-2)Gal(β 1-4)Gluc Fuc(α 1-3)
5	Lacto- <i>N</i> -triose II	LNT II	GlcNAc(β 1-3)Gal(β 1-4)Gluc
6	Lacto- <i>N</i> -tetraose	LNT	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc
7	Lacto- <i>N</i> -neotetraose	LNnT	Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc
8	Lacto- <i>N</i> -fucopentaose I	LNFP I	Fuc(α 1-2)Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc
9	Lacto- <i>N</i> -neofucopentaose I	LNnFP I	Fuc(α 1-2)Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc
10	Lacto- <i>N</i> -fucopentaose II	LNFP II	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-4)
11	Lacto- <i>N</i> -fucopentaose III	LNFP III	Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-3)
12	Lacto- <i>N</i> -fucopentaose V	LNFP V	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-3)
13	Lacto- <i>N</i> -neofucopentaose V	LNnFP V	Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-3)
14	Lacto- <i>N</i> -difucohexaose I	LNDH I	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-2) Fuc(α 1-4)
15	Lacto- <i>N</i> -difucohexaose II	LND	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)Gluc Fuc(α 1-4) Fuc(α 1-3)
16	6'-Galactosyllactose	6'-GL	Gal(β 1-6)Gal(β 1-4)Gluc
17	3'-Galactosyllactose	3'-GL	Gal(β 1-3)Gal(β 1-4)Gluc
18	Lacto- <i>N</i> -hexaose	LNH	Gal(β 1-4)GlcNAc(β 1-6)Gal(β 1-4)Gluc Gal(β 1-3)GlcNAc(β 1-3)
19	Lacto- <i>N</i> -neohexaose	LNnH	Gal(β 1-4)GlcNAc(β 1-6)Gal(β 1-4)Gluc Gal(β 1-4)GlcNAc(β 1-3)
20	<i>para</i> -Lacto- <i>N</i> -hexaose	<i>para</i> LNT	Gal(β 1-3)GlcNAc(β 1-3)Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc
21	<i>para</i> -Lacto- <i>N</i> -neohexaose	<i>para</i> LNnH	Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)GlcNAc(β 1-3)Gal(β 1-4)Gluc
22	Difucosyl-lacto- <i>N</i> -neohexaose	DF-LNnH	Fuc(α 1-3)

[0055] In this regard, it is preferred if the glycosidase is selected from one or more of the group comprising beta-galactosidase, beta-glucosidase, beta-*N*-acetylhexoamidase, alpha-fucosidase, beta-fucosidase, alpha-glucosidase alpha-galactosidase, beta-mannosidase, alpha-mannosidase, neuramidase, and it is particularly preferred if the glycosidase is a beta-galactosidase, preferably an *E. coli* beta-galactosidase.

[0056] “Beta-galactosidase” as it is used herein and as generally understood within the field of the invention, is a hydrolase enzyme that catalyzes the hydrolysis of beta-galactosides into monosaccharides.

[0057] Beta-galactosidases, are regularly employed in the genetics, molecular biology and biotechnical engineering techniques, and are *per se* known to one skilled in the art.

[0058] According to a preferred embodiment, the microorganism – used in the process according to the invention and claimed therein – is selected from a bacterium or a yeast, and more preferably, the host microorganism is an *Escherichia coli* strain, a *Corynebacterium* strain, in particular *Corynebacterium glutamicum*, a *Lactobacillus* species, or a *Saccharomyces sp.* strain.

[0059] The bacteria *Escherichia coli*, *Lactobacillus sp.*, and *Corynebacterium glutamicum* and the yeast *Saccharomyces sp.* have the advantage that these microorganisms can be grown easily and inexpensively in laboratory settings, and the bacteria and yeast have been intensively investigated for over 60 years.

[0060] Accordingly, in a preferred embodiment, the host microorganism used in the process according to the invention and otherwise claimed therein is selected from the group consisting of bacteria and yeast, and is preferably an *Escherichia coli* strain.

[0061] According to another aspect of the invention, in the process or the use according to the invention, the host microorganism employed, and preferably expressing a

glycosidase, is further expressing proteins for sugar catabolic pathways otherwise not present in the microorganism, which proteins are selected from at least one of the following: galactose catabolic pathway proteins, such as encoded by the gal operon when using a galactosidase, fucose catabolic pathway when using a fucosidase, N-acetylglucosamine catabolic pathway using beta-N-acetylhexosaminidase. By using the beta-N-acetylhexosaminidase for the degradation of terminal N-acetylglucosamine containing oligosaccharides it proved beneficial to overexpress enzymes involved in the catabolism of the monosaccharide N-acetylglucosamine. Likewise, if the liberated monosaccharide is L-fucose, it is advantageous to express the L-fucose catabolic pathway. Same holds true for sialic acid etc.

[0062] This embodiment has the advantage that with the deregulation of sugar catabolic pathways the monosaccharides liberated by the action of the glycosidases can be efficiently and effectively removed from the fermentation medium. Alternatively, also monosaccharides added as precursors can be equally removed from the fermentation medium by this means. Thus, the host microorganism expresses, besides the glycosidase, proteins for sugar catabolic pathways, such as the galactose catabolic pathway, in order to prevent the accumulation of the degradation product galactose. However, the glycosidase and the sugar catabolic pathway do not have to be expressed by an individual organism, they can be also expressed by two different co-cultured strains or the glycosidase(s) is added to a suitable strain capable of the desired monosaccharide catabolism.

[0063] According to another embodiment, and alternatively to expressing a sugar catabolic pathway for the liberated monosaccharide, the monosaccharide liberated during degradation of the undesired side-products, unused starting substrates and intermediate products can be conferred into its nucleotide activated form by expressing its salvage pathway and reused for oligosaccharide synthesis. Thus, e.g. Fucose can be conferred into GDP-Fucose, or sialic acid into CMP-Neu5Ac.

[0064] E.g., if fucose is generated as liberated monosaccharide, its salvage pathway can be used (i.e. overexpressed (highly regulated) in the host microorganism fucose has been generated) to confer the fucose into GDP-fucose.

[0065] In the so called "fucose salvage pathway" fucose is first phosphorylated to fucose-1-phosphate by the enzyme fucose kinase. The fucose-1-phosphate is then converted to GDP-fucose by the action of the enzyme fucose-1-P- guanylyltransferase. Thus, according to one aspect, a microorganism can be employed which is genetically modified to express a fucose kinase and a guanylyltransferase. According to one embodiment, the bacterial enzyme Fkp is used which represents the first identified bifunctional enzyme with both fucose kinase and L-fucose-1-P-guanylyltransferase. A detailed description of generating such genetically modified microorganisms can found in WO 2010/070104 A1, the disclosure of which is explicitly referred to and whose content is explicitly incorporated by reference.

[0066] For the fermentation of more complex oligosaccharides the combination of glycosidases is employed, i.e. expressed in the host microorganism, such as beta-galactosidase and a beta-*N*-acetylhexosaminidase, the latter of which specifically degrades lacto-*N*-triose II (LNT-2) into *N*-acetylglucosamine and lactose. The released *N*-acetylglucosamine and galactosamine can then equally efficient removed from the fermentation medium by deregulating or introducing specific monosaccharide importers and the monosaccharide specific catabolic pathway.

[0067] According to another embodiment of the process and use of the invention, the host microorganism to be employed is wild-type expressing proteins for sugar catabolic pathways, which proteins are selected from at least one of the following: galactose catabolic pathway proteins when using a galactosidase, fucose catabolic pathway when using a fucosidase, *N*-acetylglucosamine catabolic pathway using beta-*N*-acetylhexosaminidase, sialic acid, wherein an overexpression of such proteins is induced in said host microorganism during the process or the use.

[0068] This embodiment has the advantage that a host microorganism already comprising and expressing the desired proteins for the sugar catabolic pathways, can be employed, wherein the overexpression of such protein(s) is induced. The overexpression can be achieved, e.g., by putting the nucleic acid sequence coding for these proteins under control of an inducible promoter, so that the expression of the gene/polynucleotide can be specifically targeted, and, overexpressed in that way.

[0069] The expression system constructs thus contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express proteins and/or to express a protein in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., see above.

[0070] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry and nucleic acid chemistry and hybridization described above and below are those well known and commonly employed in the art.

[0071] Further advantages follow from the description of the embodiments and the attached drawings.

[0072] It goes without saying that the abovementioned features and the features which are still to be explained below can be used not only in the respectively specified combinations, but also in other combinations or on their own, without departing from the scope of the present invention.

[0073] Several embodiments of the invention are illustrated in the figures and explained in more detail in the following description. In the figures:

[0074] Fig. 1 figure shows the resolution of lacto-*N*-tetraose and lactose using a beta-galactosidase. **A.** shows the superimposed HPLC chromatograms of 10 mM lactose and 10 mM lacto-*N*-tetraose (authentic standards). **B.** HPLC chromatogram of beta-galactosidase reaction containing 10 mM lactose and 10 mM lacto-*N*-tetraose immediately taken after enzyme addition. **C.** HPLC chromatogram of beta-galactosidase reaction containing

10 mM lactose and 10 mM lacto-*N*-tetraose taken 3 hours post enzyme addition;

- [0075]** Fig. 2 shows a diagram displaying a continuous fermentation set-up suitable for the production of an oligosaccharide, such as 2'-fucosyllactose, 3-fucosyllactose, 2'3-difucosyllactose, 3'-sialyllactose, 6'-sialyllactose, according to one embodiment of the invention, with a first fermenter and a second fermenter;
- [0076]** Fig. 3 shows a HPLC analysis of a cell free medium sample prior to the addition of a beta-galactosidase expressing *E. coli* strain to the fermentation. The strain also expressed the galactose catabolic pathway in order to prevent galactose accumulation;
- [0077]** Fig. 4 shows a HPLC analysis of a cell free medium sample post addition of a second beta-galactosidase expressing *E. coli* strain to the fermentation;
- [0078]** Fig. 5 shows a diagram displaying a continuous fermentation set-up useful for the production of 2'-fucosyllactose according to one embodiment of the invention. The first fermenter vessel is used for the continuous synthesis of the synthesis of 2'-fucosyllactose whereas the second fermentation vessel besides the synthesis of 2'fucosyllactose is employed for the removal of excess substrate (lactose) and other saccharide side products;
- [0079]** Fig. 6 shows a HPLC analysis of the product stream taken from fermenter 1. The analysis of the cell free fermentation broth shows the presence of both lactose and 2'-fucosyllactose;
- [0080]** Fig. 7 shows a HPLC analysis of the product stream obtained from fermenter 2. The HPLC analysis shows the lack of the substrate

lactose which as metabolized by the strain through the expression of the *E. coli lacZ* gene encoding a beta-galactosidase and the gal operon for the metabolism of the liberated galactose;

- [0081]** Fig. 8 shows a HPLC analysis of a cell free medium sample prior to the addition of a beta-galactosidase and beta-*N*-acetylhexosaminidase expressing *E. coli* strain to the fermentation. The strain also expressed the galactose catabolic pathway in order to prevent galactose accumulation;
- [0082]** Fig. 9 shows a sample taken after glycosidase and beta-*N*-acetylhexosaminidase treatment: The comparison of the HPLC chromatograms clearly shows the disappearance of the lactose and Lacto-*N*-triose II present in the sample before beta-galactosidase and beta-*N*-acetylhexosaminidase treatment;
- [0083]** Fig. 10 shows the sequence of the gal operon used in the generation of the recombinant *E. coli* strains; and
- [0084]** Fig. 11 shows a list of oligosaccharides that can be produced according to the invention.
- [0085]** Detailed Description of preferred embodiments

[0086] To a solution containing fermentation derived lacto-*N*-tetraose and lactose, both at 10 mM concentration, beta-galactosidase was added; 50 units/ml (final concentration) of beta-galactosidase (Sigma-Aldrich Chemie GmbH, Munich Germany catalog No. G6008)) were added in order to cleave the lactose in glucose and galactose for better chromatographic separation of the saccharides (e.g. by size dependent gel-filtration chromatography). It could be shown that within seconds the lactose was cleaved into its monosaccharides galactose and glucose, whereas the lacto-*N*-tetraose remained as such (see also Figure 1). These experiments proved the highly selective action of the

purified *E. coli* beta-galactosidase, which selectively only hydrolyzes the Gal(beta1-4)Glc glycosidic bond and not the Gal(beta1-3)GlcNAc glycosidic bond present at the non-reducing end of the lacto-*N*-tetraose. Accordingly, fig. 1A shows the superimposed HPLC chromatograms of 10 mM lactose and 10 mM lacto-*N*-tetraose (authentic standards), and fig. 1B shows the HPLC chromatogram of beta-galactosidase reaction containing 10 mM lactose and 10 mM lacto-*N*-tetraose immediately taken after enzyme addition. Fig. 1C shows the HPLC chromatogram of the beta-galactosidase reaction containing 10 mM lactose and 10 mM lacto-*N*-tetraose taken 3 hours post enzyme addition.

[0087] Fig. 2 shows a continuous fermentation set-up useful for the production of oligosaccharides such as 2'-fucosyllactose, 3-fucosyllactose, 2',3-difucosyllactose, 3'-sialyllactose or 6'-sialyllactose and derivatives thereof, according to one embodiment of the invention.

[0088] The first fermenter vessel ("Fermenter 1") is used for the continuous synthesis of the desired oligosaccharide whereas the second fermentation vessel ("Fermenter 2") containing a microbial strain expressing a suitable glycosidase, such as beta-galactosidase, is used for the degradation of excess monosaccharides, such as lactose and other saccharides present in the medium interfering with the subsequent purification of the desired oligosaccharide product. Fermenter 1 and Fermenter 2 are connected with each other suitable conduits and a pump effects the transferral of the fermentation slurry from Fermenter 1 to Fermenter 2. Another pump removes the stream containing the desired oligosaccharide product from Fermenter 2.

[0089] Resolution of saccharide mixtures by glycosidase treatment in the production of 2'-fucosyllactose

[0090] A 2'-fucosyllactose feed-batch fermentation employing a recombinant 2'-fucosyllactose synthesizing *E. coli* strain (*E. coli* BL21(DE3) Δ lacZ, containing a genomic integration 2'-fucosyltransferase, encoded by the *wbgL* gene (EP 11 1151 571.4), and having an additional copy of the *E. coli* *lacY*, *manB*, *manC*, *gmd* and *fcl* all under the control of a strong constitutive tetracyclin promoter, containing a functional gal operon

comprising the gens *galM*, *galK*, *galT* and *galE*; see fig. 10; forward primer P-TTACTCAGCAATAAACTGATATTCCGTCAGGCTGG (SEQ ID No. 2); reverse primer P-TTGATAATCTCGCGCTCTTCAGCAGTCAGACTTTCCATATAGAGCGTAATTTCCGTTAACGTCGGTAGTGCTGACCTTGCCGGAGG (SEQ ID. No. 3)) grown in a defined salt medium (7 g l⁻¹ NH₄H₃PO₄, 7 g l⁻¹ K₂HPO₄, 2 g l⁻¹ KOH, 0.37g l⁻¹ citric acid, 1 ml l⁻¹ antifoam (Struktol J673, Schill + Seilacher), 1 mM CaCl₂, 4 mM MgSO₄, Trace-Elements consisted of 0.101 g l⁻¹ nitrilotriacetic acid pH 6.5, 0.056 g l⁻¹ ammonium ferric citrate, 0.01 g l⁻¹ MnCl₂ x 4 H₂O, 0.002 g l⁻¹ CoCl₂ x 6 H₂O, 0.001g l⁻¹ CuCl₂ x 2 H₂O, 0.002 g l⁻¹ boric acid, 0.009 g l⁻¹ ZnSO₄ x 7 H₂O, 0.001 g l⁻¹ Na₂MoO₄ x 2 H₂O, 0.002 g l⁻¹ Na₂SeO₃, 0.002 g l⁻¹ NiSO₄ x 6 H₂O with 2% glycerol as carbon source); Glycerol-feed consisted of glycerol 800 g l⁻¹, MgSO₄ 2,64 g l⁻¹ and trace element solution 4 ml l⁻¹. For 2'-fucosyllactose formation a lactose feed of 216 g l⁻¹ was employed. The pH was controlled by using ammonia solution (25% v/v), which also served as a nitrogen-source. Lactose was supplied as a precursor for the 2'-fucosyllactose production. Feed batch was cultured at 30°C under constant aeration and agitation for 90 hours. At 90 hours after the start of the fermentation most of the added lactose was converted into 2'-fucosyllactose.

[0091] In order to remove most or all lactose still present in the fermentation supernatant, a second bacterial strain was added to the fermentation vessel 90 hour post fermentation start (see set-up in Fig. 2). The added second bacterial strain was genetically identical to the first employed bacteria strain, differing, however, only in the expression of a genome integrated beta-galactosidase and expressing a functional gal operon (see fig. 10) (for D-galactose degradation). Incubation of the added secondary bacterial strain resulted in the disappearance of the residual lactose within 5 hours. Approximately 25 ml starter culture of the second beta-galactosidase expressing bacterial strain was added per 1l fermentation broth.

[0092] For the HPLC analysis, a cell free sterile filtered medium sample was used, and desalted by passing a 350 µl sample over a desalting column (Strata ABW (55µm, 70A) phenomenex, Aschaffenburg, Germany). For HPLC analysis, a ReproSil Carbohydrate 5 µm, 250 x 4.6 mm column (Dr. Maisch GmbH, Ammerbuch-Entringen, Germany) was used with the following conditions: eluent acetonitrile/ddH₂O (68:32), isocratic condition with flow rate of 1.4 ml/min (column oven was set to 35°C); as an

internal standard for quantification sucrose was used; for identification of lactose and 2'-fucosyllactose authentic standards were employed. For the detection a refractive index detector (RID) was employed. The inclusion of the glycosidase resolution step at the end of the fermentative process led to a significant reduction in the complexity of the resulting fermentation filtrate. Figure 3 shows the analysis of cell free fermentation medium taken before glycosidase treatment and Figure 4 shows a sample taken after glycosidase treatment: The comparison of the HPLC chromatograms clearly shows the disappearance of the lactose present in the sample before beta-galactosidase treatment. The strain also expressed the galactose catabolic pathway in order to prevent D-galactose accumulation.

[0093] Resolution of saccharide mixtures by glycosidase treatment in the continuous production of 2'-fucosyllactose

[0094] Continuous synthesis of 2'-Fucosyllactose was achieved by using two fermentation vessels, fermenter 1 containing a 2'-fucosyllactose fermenting strain deficient in lactose degradation (*E. coli* BL21(DE3) $\Delta lacZ$, containing a genomic integration 2'-fuosyltransferase, encoded by the *wbgL* gene (EP 11 1151 571.4), and having an additional copy of the *E. coli lacY*, *manB*, *manC*, *gmd* and *fcl* all under the control of a strong constitutive tetracyclin promoter, containing a functional gal operon (see fig. 10) comprising the gens *galM*, *galK*, *galT* and *galE*). Fermenter 2 contained a starter culture genetically similar to the used 2'-fucosyllactose fermenting strain of fermenter 1 except that this strain expressed in addition the *E. coli lacZ* gene encoding a beta-galactosidase (see set-up displayed in Figure 5). For both fermenters a defined salt medium containing 7 g l⁻¹ NH₄H₃PO₄, 7 g l⁻¹ K₂HPO₄, 2 g l⁻¹ KOH, 0.37g l⁻¹ citric acid, 1 ml l⁻¹ antifoam (Struktol J673, Schill + Seilacher), 1 mM CaCl₂, 4 mM MgSO₄, Trace-Elements consisted of 0.101 g l⁻¹ nitrilotriacetic acid pH 6.5, 0.056 g l⁻¹ ammonium ferric citrate, 0.01 g l⁻¹ MnCl₂ x 4 H₂O, 0.002 g l⁻¹ CoCl₂ x 6 H₂O, 0.001g l⁻¹ CuCl₂ x 2 H₂O, 0.002 g l⁻¹ boric acid, 0.009 g l⁻¹ ZnSO₄ x 7 H₂O, 0.001 g l⁻¹ Na₂MoO₄ x 2 H₂O, 0.002 g l⁻¹ Na₂SeO₃, 0.002 g l⁻¹ NiSO₄ x 6 H₂O with 10 mM lactose as substrate and 2% glycerol as carbon source was used.

[0095] As illustrated, to fermenter 1 a constant feed of medium (7 g l⁻¹ NH₄H₃PO₄, 7 g l⁻¹ K₂HPO₄, 2 g l⁻¹ KOH, 0.37g l⁻¹ citric acid, 1 ml l⁻¹ antifoam, 1 mM CaCl₂,

4 mM MgSO₄, Trace-Elements consist of 0.101 g l⁻¹ nitrilotriacetic acid pH 6.5, 0.056 g l⁻¹ ammonium ferric citrate, 0.01 g l⁻¹ MnCl₂ x 4 H₂O, 0.002 g l⁻¹ CoCl₂ x 6 H₂O, 0.001 g l⁻¹ CuCl₂ x 2 H₂O, 0.002 g l⁻¹ boric acid, 0.009 g l⁻¹ ZnSO₄ x 7 H₂O, 0.001 g l⁻¹ Na₂MoO₄ x 2 H₂O, 0.002 g l⁻¹ Na₂SeO₃, 0.002 g l⁻¹ NiSO₄ x 6 H₂O) containing 40 mM lactose and 5% glycerol was supplied. Using a New Brunswick parallel fermentation system (Bio-Flow/CelliGen 115) with a working volume of 1 l, a continuous flow of 20 ml/h feed solution was applied. The action of three synchronous working pumps led then to the similar production of a product stream of 20 ml/h containing almost exclusively 2'-fucosyllactose as sole saccharide product.

[0096] Fig. 6 shows a HPLC analysis of the product stream taken from fermenter 1. The analysis of the cell free fermentation broth shows the presence of both lactose and 2'-fucosyllactose. In contrast, fig. 7 shows a HPLC analysis of the product stream obtained from fermenter 2. The HPLC analysis shows the lack of the substrate lactose which was metabolized by the strain through the expression of the *E. coli lacZ* gene encoding a beta-galactosidase and the gal operon for the metabolism of the liberated galactose.

[0097] Resolution of saccharide mixtures by glycosidase treatment in the production of lacto-*N*-tetraose

[0098] In the fermentation of lacto-*N*-tetraose from lactose besides the added substrate also lacto-*N*-triose II (LNT-2) can accumulate. In order to obtain a more economical fermentation filtrate for purification of lacto-*N*-tetraose, the specific degradation of lactose and LNT-2 proved desirable.

[0099] For this purpose a strain was constructed which, besides the beta-galactosidase, expressed a beta-*N*-acetylhexosaminidase (encoded by the *Bifidobacterium bididum* JCM1254 *bbhl* gene) for the specific degradation of LNT-2 into *N*-acetylglucosamine and lactose. The resulting lactose was further cleaved into glucose and galactose. Whereas the resulting glucose was efficiently metabolized by the organisms the employed *E. coli* BL21 (DE3) strain was incapable of catabolizing galactose due to the

DE3 prophage integration into the gal operon. For the efficient degradation of galactose the gal operon from the *E. coli* K strain JM 109 was amplified and transformed into *E. coli* BL21 (DE3) by simultaneous expression of the Lambda Red recombination protein (from an episomal vector system). *E. coli* BL21 transformants capable of galactose degradation were isolated by screening them on galactose containing minimal medium agar plates and McConkey agar plates containing galactose.

[00100] After completion of the lacto-*N*-tetraose fermentation - it is noted that alternatively the oligosaccharide mixture could have been obtained via an enzymatic or chemical synthesis reaction - the engineered glycosidase strain was added to the batch fermentation. Similar to the above described 2'-Fucosyllactose fermentation the degradation of the undesired oligosaccharides were completed within a few minutes to several hours depending on the amounts of culture used for the resolution step.

[00101] The inclusion of the glycosidase resolution step at the end of the fermentative process led to a significant reduction in the complexity of the resulting fermentation filtrate. Fig. 8 shows the analysis of cell free fermentation medium taken before glycosidase and beta-*N*-acetylhexosaminidase treatment and Fig. 9 shows a sample taken after glycosidase and beta-*N*-acetylhexosaminidase treatment: The comparison of the HPLC chromatograms clearly shows the disappearance of the lactose and Lacto-*N*-triose II present in the sample before beta-galactosidase and beta-*N*-acetylhexosaminidase treatment.

[00102] As shown above, the use and the process according to the invention proved particularly useful for the purification of human milk oligosaccharides (HMOs) from microbial fermentation reaction. Almost all human milk oligosaccharides contain a Gal-beta-1,4-Glc disaccharide subunit at the reducing end. The lactose moiety (Gal-beta-1,4-Glc) subunit can either be obtained by total fermentation or lactose is added as a substrate to the microbial fermentation. It is often advantageous for attaining maximum volumetric yields to add excess amounts of lactose to the fermentation.

[00103] According to the invention, and in order to remove lactose or unwanted side products thereof, a glycosidase such as a beta-galactosidase is added to the fermentation at a specific time point to the fermentation leading to the specific degradation of the lactose and side products.

[00104] As mentioned above, alternatively, a transcription-regulated (or otherwise activity regulated) glycosidase can be induced in the production strain or an additional strain expressing the desired activity can be added to the fermentation at a specific time point.

[00105] As also mentioned and shown above, the process of the present invention further involves deregulation of sugar catabolic pathways in order to efficiently and effectively remove from the fermentation medium the monosaccharides liberated by the action of the glycosidases. Alternatively, also monosaccharides added as precursors can be equally removed from the fermentation medium by this means.

[00106] Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated step or element or integer or group of steps or elements or integers but not the exclusion of any other step or element or integer or group of elements or integers.

[00107] Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority date of each of the appended claims.

Claims

1. Use of one or more glycosidases in the purification of a produced desired human milk oligosaccharide (HMO) from a mixture containing the produced desired HMO and one or more selected from the group consisting of non-desired oligosaccharides, metabolic saccharide products generated during production of said desired HMO, and unused saccharide substrates used in the production of said HMO, for degradation of the non-desired oligosaccharides or metabolic saccharide products or unused substrates, characterized in that the glycosidase is used in a fermentation process employed for the production of a not naturally occurring HMO in a host microorganism, wherein the glycosidase is used for degradation of the non-desired oligosaccharide, metabolic products or unused saccharide substrates.
2. The use of claim 1, characterized in that the glycosidase is not naturally occurring in the microorganism of said host microorganism.
3. The use of claim 1 or 2, characterized in that, where applicable, naturally expressed endogenous genes encoding glycosidases of the host microorganism have been inactivated or deleted.
4. The use of any of claims 1 to 3, characterized in that the glycosidase is used for the resolution of oligosaccharide mixtures in order to obtain 2'-fucosyllactose.
5. Process for producing an human milk oligosaccharide (HMO) using a host microorganism, wherein said HMO is not naturally occurring in said host cell, the process comprising the steps of:
 - a) cultivating a host microorganism suitable for the production of a desired HMO under conditions and in a medium permissive for the production of said HMO, whereby the HMO and, where applicable, biosynthetic saccharide intermediates and/or side products are produced;

- b) using a glycosidase in the medium the host microorganism is cultivated in, in order to degrade biosynthetic saccharide intermediates and/or saccharide side products and/or unused saccharide substrates, and
 - c) recovering the desired HMO.
6. The process of claim 5, characterized in that the glycosidase is added to the medium.
 7. The process of claim 5, characterized in that the glycosidase is endogenously produced of said host microorganism, wherein the nucleic acid sequences encoding the endogenously produced glycosidase is not naturally occurring in said host cell, and wherein said host cell has been stably transformed to express the not naturally occurring glycosidase, and wherein the expression of the glycosidase in the host microorganism is inducible.
 8. The process of claim 5, characterized in that the glycosidase is produced by a second microorganism additionally added to the medium and expressing the glycosidase.
 9. The process of any of claims 5 to 8, characterized in that, the process is a batch or a continuous process.
 10. The process of any of claims 5 to 9, characterized in that the HMO is recovered from supernatant of the cultivated host microorganism, which supernatant is obtained by centrifuging the cultivated host microorganism to obtain a supernatant and a host microorganism pellet.
 11. The process of any of the preceding claims, characterized in that the desired HMO is selected from 2'-fucosyllactose, 3-fucosyllactose, 2'3-difucosyllactose, 3'-sialyllactose, 6'-sialyllactose, 3-fucosyl-3'-sialyllactose, lacto-*N*-tetraose, lacto-*N*-neotetraose, lacto-*N*-fucopentaose I, lacto-*N*-fucopentaose II, lacto-*N*-fucopentaose III, lacto-*N*-fucopentaose V, lacto-*N*-difucosylhexose I, lacto-*N*-

difucosylhexaose II, lacto-*N*-sialylpentaose LSTa, LSTb, LSTc, and in particular from one of the oligosaccharides shown in table 1 of Fig. 11 or derivatives thereof.

12. The process of any of claims 5 to 11, or the use of any of claims 1 to 4, characterized in that the glycosidase is selected from one or more of the group comprising, galactosidases, mannosidases, fucosidases, sialidases (neuraminidase), glucosidase, *N*-acetylglucoamidases, *N*-acetylhexoamidase..
13. The process of any of claims 5 to 11 or the use of any of claims 1 to 4, characterized in that the glycosidase is selected from one or more of the group comprising beta-galactosidase, alpha-galactosidase, beta-*N*-acetylglucoamidase, beta-*N*-acetylhexoamidase, beta-mannosidase, alpha-mannosidase, alpha-fucosidase, beta-fucosidase, beta-glucosidase, alpha-glucosidase, neuraminidase
14. The process of any of claims 6 to 13 or the use of any of claims 1 to 4, characterized in that the host microorganism is selected from a bacterium and a yeast.
15. The process of any of claims 6 to 14 or the use of any of claims 1 to 4, characterized in that the host microorganism is an *Escherichia coli* strain, *Lactobacillus* species or a *Corynebacterium glutamicum* strain or a *Saccharomyces sp.* strain.
16. The process of any of claims 6 to 15 or the use of any of claims 1 to 4, characterized in that the host microorganism employed is expressing proteins for sugar catabolic pathways otherwise not present in the microorganism, which proteins are selected from at least one of the following: galactose catabolic pathway proteins when using a galactosidase, fucose catabolic pathway when using a fucosidase, *N*-acetylglucosamine catabolic pathway using beta-*N*-acetylhexosaminidase.
17. The process of any of claims 6 to 15 or the use of any of claims 1 to 4, characterized in that the in the host microorganism employed a salvage pathway for a monosaccharide liberated during degradation is overexpressed, in order to convert the monosaccharide

18. The process of any of claims 6 to 17 or the use of any of claims 1 to 4, characterized in that the host microorganism employed is wild-type expressing proteins for sugar catabolic pathways, which proteins are selected from at least one of the following: galactose catabolic pathway proteins when using a galactosidase, fucose catabolic pathway when using a fucosidase, N-acetylglucosamine catabolic pathway using beta-*N*-acetylhexosaminidase, wherein an overexpression of such proteins is induced in said host microorganism during the process or the use.





Fig. 1

**Fig. 2**




Fig. 3





Fig. 4

4 / 12

**Fig. 5**




Fig. 6





Fig. 7

7 / 12

**Fig. 8**




Fig. 9

Fig. 10

Sequence of the gal operon fragment (obtained from *E. coli* K strain JM109) and inserted into the employed *E. coli* BL21 (DE3) strain.

Sequence of used gal-operon fragment (SEQ ID No. 1)

```

T T A C T C A G C A A T A A A C T G A T A T T C C G T C A G G C T G G A A T A C T C T T C G C C A G G A C G C A G G A A G C A G T C C G G T T G C G G C C A T T C
A G G G T G G T T C G G G C T G T C C G G T A G A A A C T C G C T T T C C A G A G C C A G C C C T T G C C A G T C G G C G T A A G G T T C G G T T C C C C G C G
A C G G T G T G C C G C C G A G G A A G T T G C C G G A G T A G A A T T G C A G A G C C G G A G C G G T G G T G T A G A C C T T C A G C T G C A A T T T T T C A
T C T G T G T A C C A G A C A T G C G C C C C A C T T T C T T G C C A T C G C C T T T G G C C T G T A A C A A G A A T G C G T G A T C G T A A C C T T T C A C T T
T G C G C T G A T C G T C G T C G G C A A G A A A C T A C T G G C G A T G A T T T T G G C G C T G C G G A A T C A A A G A C G T T C C G G C G A C A G A T
T T C A G G C C G T C G T G C G G A A T G C C G C C T T C A T C A A C C G G C A G A T A T T C G T C C G C C A G A A T C T G C A A C T T G T G A T T G C G C A C G
T C A G A C T G C T C G C C G T C A A G A T T G A A A T A G A C G T G A T T A G T C A T A T T C A C C G G G C A A G G T T T A T C A A C T G T G G C G C G A T A A
G T A A T G G A G A T A C G G T T A T C G T C G G T C A G A C G A T A T T G C A C C G T C G C G C G A G A T T A C C C G G A A G C C C T G A T C A C C A T C
A T C T G A A C T C A G G G C A A A C A G C A C C T G A C G A T C G T T C T G G T T C A C A A T C T G C C A G C G A C G T T T G T C G A A C C C T T C C G G C C
G C C G T G C A G C T G G T T A A C G C C C T G A C T T G G C G A A A G C G T C A C G T T T C A C C G T C A A A G G T A T A A C G G C T A T T G G C G A T A C
G G T T G G C A T A A C G A C C A A T A G A G G C C C C A G A A A C G C G G C C T G A T C C T G A T A G C A T T C C G G G C T G G C A C A G C C G A G C A G
C G C C T C G C G G A C G C T G C C A T C G G A A A G C G G A A T A C G G G C G G A A A G T A A A G T C G C A C C C C A G T C C A T C A G C G T G A C T A C C
A T C C C T G C G T T G T T A C G C A A A G T T A A C A G T C G G T A C G G C T G A C C A T C G G G T G C C A G T G C G G G A G T T T C G T T C A G C A C T G T C
C T G C T C C T T G T G A T G G T T A C A A A C G T A A A A A G T C T C T T T A A T A C C T G T T T T G C T T C A T A T T G T T C A G C G A C A G C T T G C T G T
A C G G C A G G C A C C A G C T T T C C G G G A T C A G C G C G A C G A T A C A G C C G C C A A T C C G C C G C C G T C A T G C G T A C G C C A C C T T
T G T C G C C A A T C A C A G C T T T G A C G A T T T C T A C C A G A G T G T C A A T T T G C G G C A C G G T G A T T T C G A A A T C A T C G C G C A T A G A G G
C A T G A G A C T C C G C C A T C A A C T C G C C C A T A C G T T T C A G G T C G C C T T G C T C C A G C G C G C T G G C A G C T T C A A C G G T G C G G G C G
T T T T C A G T C A G T A T A T G A C G C A C G C G T T T T G C C A C G A T C G G G T C C A G T T C A T G C G C A A C A G C G T T G A A C T T T C A A T G G T G A
C A T C A C G C A G G G C T G G C T G C T G G A A G A A A C G C G C A C C G T T T C G C A C T G T T C A C G A C G G G T G T T G A T T C G C T G C C A A C C
A G G G T A C G T T T G A A G T T A C T G T T G A T G A T G A C G A C A G C C A C A C C T T T G G G C A T G G A A A C T G C T T T G G T C C C C A G T G A G C G G
C A A T C G A T C A G C A A G G C A T G A T C T T T C T T G C C G A G C G C G G A A A T T A G C T G A T C C A T G A T C C C G C A G T T A C A G C C T A C A A A C
T G G T T T T C T G C T T C C T G A C C G T T A A G C G C G A T T T G T G C G C C G T C C A G C G G C A G A T G A T A A A G C T G C T G C A A T A C G G T T C C G
A C C G C G A C T T C C A G T G A A G C G G A A G A A C T T A A C C C G G C A C C C T G C G G C A C A T T G C C G C T G A T C A C C A T G T C C A C G C C G C C
G A A G C T G T T G T T A C G C A G T T G C A G A T G T T T C A C C A C G C C A C G A A C G T A G T T A G C C C A T T G A T A G T T T T C A T G T G C G A C A A T G
G G C G C A T C G A G G G A A A A C T C G T C G A G C T G A T T T T C A T A A T C G G C T G C C A T C A C G C G A A C T T T A C G G T C A T C G C G T G G T G C A
C A A C T G A T C A C G G T T T G A T A A T C A A T C G C G C A G G G C A G A A C G A A A C C G T C G T T G T A G T C G G T G T G T T C A C C A A T C A A A T T C
A C G C G G C C A G G C G C C T G A A T G G T G T G A G T G G C A G G G T A G C C A A A T G C G T T G G C A A A C A G A G A T T G T G T T T T T T C T T T C A G A
C T C A T T T C T T A C A C T C C G G A T T C G C G A A A A T G G A T A T C G C T G A C T G C G C G C A A A C G C T C T G C T G C C T G T T C T G C G G T C A G G
T C T C G C T G G G T C T C T G C C A G C A T T T C A T A A C C A A C C A T A A A T T T A C G T A C G G T G G C G G A G C G C A G C A G A G G C G G A T A A A A G
T G C G C G T G C A G C T G C C A G T G T T G A T T C T C T T C G C C A T T A A A T G G C G C G C C G T G C C A G C C C A T A G A G T A G G G G A A G G A G C A
C T G G A A G A G G T T G T C A T A A C G A C T G G T C A G C T T T T T C A A C G C C A G C G C C A G A T C G C T G C G C T G G G C G T C G G T C A A A T C G G
T G A T C C G T A A A A C G T G G G C T T T G G G C A G C A G T A G C G T T T C G A A C G G C C A G G C A G C C C A G T A A G G C A C G A C G G C T A A C C A G
T G T T C G G T T T C G A C A A C G G T A C G G C T A C C G T C T G C C A G C T C G C G C T G A A C A T A A T C C A C C A G C A T T G G T G A T T T C T G T T C G
G C A A A A T A T T C T T T T T G C A G G C G G T C T T C G C G C T C A G C T T C G T T A G G C A G G A A G C T A T T T G C C C A A A T C T G A C C G T G C G G A
T G C G G G T T A G A G C A G C C C A T C G C C G C G C C T T T G T T T T C A A A A C C T G C A C C C A T G G G T A C G T T T T C C C C A G T T C T G C G G T T
T G C T C C T G C C A G G T T T T G A C G A T T T C C G T C A A T G C T G C A A C G C T G A G C T C T G G C A G C G T T T T A C T G T G A T C C G G T G A A A A G
C A G A T C A C C C G G C T G G T G C C G C G C G C G C T C T G G C A A C G C A T C A G C G G A T C G T G A C T T T C T G G C G C A T C T G G C G T G T C A G
A C A T C A A A G C C G C A A A G T C A T T A G T G A A A C G T A A G T C C C G G T G T A A T C G G G G T T T T A T C G C C T G T C A C C C G C A C A T T A C C

```

Fig. 10 (continued)

TGCGCAGAGGAAGCAATCTGGATCGTGCGCAGGTAACACCTGTTTGGCTGGCGTTTCCTGCGCCCCCTGCCAGGGGCGC
TTAGCGCGGTGCGGTGAAACCAGAATCCATTGCCCGGTGAGCGGGTTGTAGCGGCGATGTGGATGATCAACGGGATTA
TTGCGTCATGGTCGTTCCCTTAATCGGGATATCCCTGTGGATGGCGTGACTIONGCCAGTGCCAGGTGCCTGCGCCATTT
CAGTGTGCGCGTTACGCGCCAGTTCAGTTCACGGTCGGCTTTGCTGGCGTCCGCCCAGTAGGCCGGAAGTCCGCCCTCG
CGACGCGGTGCAAAATGATAATTAACCGTTTGGCCGAGGCTTTGCTGAAGGCATTAACCACGTCCAGCACGCTGTTGCCT
ACGCCAGCGCCGAGGTTGTAGATGTGTACGCCTGGCTTGTTCGCCAGTTTTTCCATCGCCACGACGTGACCGTCCGCCAG
ATCCATTACGTGGATGTAATCGCGTACGCCAGTACCATCTTCGGTCCGATAATCGTTACCAAAAATCGCCAGCGAGTCGCG
ACGGCCTACAGCAACCTGGGCGATGTATGGCATCAGGTTATTCCGAATGCCTTGGCGATCTTCGCCCATATCGCCCGACG
GATGCGCGCCAACCGGTTGAAGTAGCGCAGCAGGGCAATGCTCCAGTCCGGCTGGGCTTTTTGCAGATCGGTGAGGAT
CTGTTCCACCATCAGCTTGCTTTTCCGTAAGGGCTTTGCGGTGTGCCGGTCGGGAAGCTTTCAACGTATGGAATTTGGG
CTGATCGCCATAAACGGTGGCGGAGGAGCTAAAAATAAAGTTTTTGACGTTAGCGGCGCGCATGGCGCTAATCAGGCGCA
GAGTGCCGTTGACATTGTTGTCGTAATATCCAGCGGTTTTTGTACCGATTCCGCCACGGCTTTCAGCCCGGCGAAGTGGA
TCACGGTGTGATAGCGTATCGTGCAGGATCTCGGTTCATCAACGCTTCCGTTACGAATATCGCCTTCAACAAACGTTGGAT
GTTTGCCGCCTAACGCTCGATAACAGGCAGTACGCTGCGCTTACTGTTACAGAGGTTATCAAGAATGATGACATCATGAC
CGTTTTGCAGTAATTGCACACAGGTATGACTTCCAATGTAACCGCTACCACCGGTAACCAGAACTCTCATAATTCGCTCCAT
TAGGCTTATGGTATGAAATAACCATAGCATAACAAAGATGCGAAAAGTGTGACATGGAATAAATTAGTGAATCGTTTACAC
AAGAATTTAGCCGTTTTTTATGCGCGATTAAGTGATTATAAACAGAGGGTTTATGAATGATTGCGCTTTTTATCTGAAAAA
GACGCGGTTTCATGCCTGCATGCGTCGAACCGTTGGCCGAGAGGGTGCTAAGGCCGCTCCGGCAAGGTCAGCACTAC
CGACGTTAACGGAAATTACGCTCTATATGGAAGTCTGACTGCTGAAGAGCGCGAGATTATCAA

Fig. 11 (continued)

20	<i>para</i> -Lacto- <i>N</i> -hexaose	paraLNT	Gal(β1-3)GlcNAc(β1-3)Gal(β1-4)GlcNAc(β1-3)Gal(β1-4)Gluc
21	<i>para</i> -Lacto- <i>N</i> -neohexaose	paraLNnH	Gal(β1-4)GlcNAc(β1-3)Gal(β1-4)GlcNAc(β1-3)Gal(β1-4)Gluc
22	Difucosyl-lacto- <i>N</i> -neohexaose	DF-LNnH	$\begin{array}{c} \text{Fuc}(\alpha 1-3) \\ \\ \text{Gal}(\beta 1-4)\text{GlcNAc}(\beta 1-6)\text{Gal}(\beta 1-4)\text{Glc} \\ \\ \text{Gal}(\beta 1-4)\text{GlcNAc}(\beta 1-3) \\ \\ \text{Fuc}(\alpha 1-3) \end{array}$
23	3'-Sialyllactose	3'-SL	Neu5Ac(α2-3)Gal(β1-4)Gluc
24	6'-Sialyllactose	6'-SL	Neu5Ac(α2-6)Gal(β1-4)Gluc
25	Lacto- <i>N</i> -sialylpentaose a	LST-a	Neu5Ac(α2-3)Gal(β1-3)GlcNAc(β1-3)Gal(β1-4)Gluc
26	Lacto- <i>N</i> -sialylpentaose b	LST-b	$\begin{array}{c} \text{Gal}(\beta 1-3)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Neu5Ac}(\alpha 2-6) \end{array}$
27	Lacto- <i>N</i> -sialylpentaose c	LST-c	Neu5Ac(α2-6)Gal(β1-4)GlcNAc(β1-3)Gal(β1-4)Gluc
28	Fucosyl-lacto- <i>N</i> -sialylpentaose a	F-LST-a	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-3)\text{Gal}(\beta 1-3)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Fuc}(\alpha 1-4) \end{array}$
29	Fucosyl-lacto- <i>N</i> -sialylpentaose b	F-LST-b	$\begin{array}{c} \text{Fuc}(\alpha 1-2)\text{Gal}(\beta 1-3)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Neu5Ac}(\alpha 2-6) \end{array}$
30	Fucosyl-lacto- <i>N</i> -sialylpentaose c	F-LST-c	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-3)\text{Gal}(\beta 1-3)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Fuc}(\alpha 1-3) \end{array}$
31	Disialyl-lacto- <i>N</i> -tetraose	DS-LNT	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-3)\text{Gal}(\beta 1-4)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Neu5Ac}(\alpha 2-6) \end{array}$
32	Disialyl-lacto- <i>N</i> -fucopentaose	DS-LNFP V	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-3)\text{Gal}(\beta 1-4)\text{GlcNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \qquad \qquad \qquad \\ \text{Neu5Ac}(\alpha 2-6) \qquad \qquad \text{Fuc}(\alpha 1-3) \end{array}$
33	3-Fucosyl-3'-sialyllactose	3F-3'-SL	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-3)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Fuc}(\alpha 1-3) \end{array}$
34	3-Fucosyl-6'-sialyllactose	3F-6'-SL	$\begin{array}{c} \text{Neu5Ac}(\alpha 2-6)\text{Gal}(\beta 1-4)\text{Gluc} \\ \\ \text{Fuc}(\alpha 1-3) \end{array}$
35	Lacto- <i>N</i> -neodifucohexaose I	LNnDFH I	$\begin{array}{c} \text{Gal}(\beta 1-4)\text{GalNAc}(\beta 1-3)\text{Gal}(\beta 1-4)\text{Glc} \\ \qquad \qquad \qquad \\ \text{Fuc}(\alpha 1-3) \qquad \qquad \text{Fuc}(\alpha 1-3) \end{array}$

eol f-seq1 . txt
SEQUENCE LISTING

<110> Jennewei n Bi otechnol ogi e GmbH
<120> Producti on of Oli gosacchari des
<130> 2827P108W0
<160> 3
<170> PatentIn versi on 3.5

<210> 1
<211> 4579
<212> DNA
<213> Escheri chi a col i

<400> 1
ttactcagca ataaactgat attccgtcag gctggaatac tcttcgccag gacgcaggaa 60
gcagtccggt tgcggccatt caggggtggt cgggctgtcc ggtagaaact cgctttccag 120
agccagccct tgccagtcgg cgtaaggttc ggttccccgc gacggtgtgc cgccgaggaa 180
gttgccggag tagaattgca gagccggagc ggtggtgtag accttcagct gcaatttttc 240
atctgctgac cagacatgcg ccgccacttt cttgccatcg cctttggcct gtaacaagaa 300
tgcgtgatcg taacctttca ctttgcgctg atcgtcgtcg gcaagaaact cactggcgat 360
gattttggcg ctgcgaaaat caaaagacgt tccggcgaca gatttcaggc cgtcgtgagg 420
aatgccgcct tcatcaaccg gcagatattc gtccgccaga atctgcaact tgtgattgag 480
cacgtcagac tgctcggcgt caagattgaa atagacgtga ttagtcatat tcaccgggca 540
aggtttatca actgtggcgc gataagtaat ggagatacgg ttatcgtcgg tcagacgata 600
ttgcaccgct gcgccgagat taccggggaa gccctgatca ccatcatctg aactcagggc 660
aaacagcacc tgacgatcgt tctggttcac aatctgccag cgacgtttgt cgaacccttc 720
cggcccggcg tgcagctggt taacgccctg acttggcgaa agcgtcacgg tttcaccgct 780
aaaggataaa cggctattgg cgatacgggt ggcataacga ccaatagagg cccccagaaa 840
cgcggcctga tcctgatagc attccgggct ggcacagccg agcagcgcct cgcggacgct 900
gcatcggaa agcgggaatac gggcgaaaag taaagtgcga cccagtcca tcagcgtgac 960
taccatccct gcgttggttac gcaaagttaa cagtcggtac ggctgaccat cgggtgccag 1020
tgcgggagtt tcgttcagca ctgtcctgct cttgtgatg gtttacaac gtaaaaagtc 1080
tctttaatac ctgtttttgc ttcattttgt tcagcgacag cttgctgtac ggcaggcacc 1140
agctcttccg ggatcagcgc gacgatacag ccgccaatc cgccgccggt catgctgacg 1200
ccacctttgt cgccaatcac agctttgacg atttctacca gagtgtcaat ttgcggcagc 1260
gtgatttcga aatcatcgcg catagaggca tgagactccg ccatcaactc gccatacgt 1320
ttcaggctgc cttgctccag cgcgctggca gcttcaacgg tgcgggaggt ttcagtcagt 1380

eol f-seq1 . txt

atatgacgca	cgcgttttgc	cacgatcggg	tccagttcat	gcgcaacagc	gttgaactct	1440
tcaatggtga	catcacgcag	ggctggctgc	tggaagaaac	gcgaccgggt	ttcgactgt	1500
tcacgacggg	tgttgtattc	gctgccaacc	agggtacgtt	tgaagttact	gttgatgatg	1560
acgacagcca	cacctttggg	catggaaact	gctttggtcc	ccagtgagcg	gcaatcgatc	1620
agcaaggcat	gatctttctt	gccgagcgcg	gaaattagct	gatccatgat	cccgcagtta	1680
cagcctacaa	actggttttc	tgcttcctga	ccgtaagcg	cgatttggtc	gccgtccagc	1740
ggcagatgat	aaagctgctg	caatacgggt	ccgaccgcga	cttccagtga	agcgggaagaa	1800
cttaaccggg	caccctgcgg	cacattgccg	ctgatcacca	tgtccacgcc	gccgaagctg	1860
ttgttacgca	gttgcatgat	tttcaccacg	ccacgaacgt	agttagccca	ttgatagttt	1920
tcatgtgcga	caatgggctc	atcgagggaa	aactcgtcga	gctgattttc	ataatcggct	1980
gccatcacgc	gaactttacg	gtcatcgcgt	ggtgcacaaac	tgatcacgggt	ttgataatca	2040
atcgcgcagg	gcagaacgaa	accgtcgttg	tagtcggtgt	gttcaccaat	caaattcacg	2100
cggccaggcg	cctgaatggt	gtgagtggca	gggtagccaa	atgcgttggc	aacagagat	2160
tgtgtttttt	ctttcagact	catttcttac	actccggatt	cgcgaaaatg	gatatcgctg	2220
actgcgcgca	aacgctctgc	tgctgttct	gcggtcaggt	ctcgctgggt	ctctgccagc	2280
atttcataac	caaccataaa	tttacgtacg	gtggcggagc	gcagcagagg	cggataaaaag	2340
tgcgctgca	gctgccagtg	ttgattctct	tcgccattaa	atggcgcgcc	gtgccagccc	2400
atagagtagg	ggaaggagca	ctggaagagg	ttgtcataac	gactggtcag	ctttttcaac	2460
gccagcgcca	gatcgctgcg	ctgggcgtcg	gtcaaatcgg	tgatccgtaa	aacgtgggct	2520
ttgggcagca	gtagcgtttc	gaacggccag	gcagcccagt	aaggcacgac	ggctaaccag	2580
tgttcggttt	cgacaacgggt	acggctaccg	tctgccagct	cgcgctgaac	ataatccacc	2640
agcattggtg	atttctgttc	ggcaaaaat	tctttttgca	ggcgtcttc	gcgctcagct	2700
tcgttaggca	ggaagctatt	tgcccaaatac	tgaccgtgcg	gatgcggggt	agagcagccc	2760
atcgccgcgc	ctttgttttc	aaaaacctgc	acccatgggt	acgttttccc	cagttctgcg	2820
gtttgctcct	gccaggtttt	gacgatttcc	gtcaatgctg	caacgctgag	ctctggcagc	2880
gttttactgt	gatccgggtga	aaagcagatc	acccggctgg	tgccgcgcgc	gctctggcaa	2940
cgcatcagcg	gatcgtgact	ttctggcgca	tctggcgtgt	cagacatcaa	agccgcaaag	3000
tcattagtga	aaacgtaagt	cccgggtgaa	tcggggtttt	tatcgccctgt	cacccgcaca	3060
ttacctgcgc	agaggaagca	atctggatcg	tgcgcaggta	acacctgttt	ggctggcgtt	3120
tcctgcgccc	cctgccaggg	gcgcttagcg	cggtgcggtg	aaaccagaat	ccattgcccg	3180
gtgagcgggt	tgtagcggcg	atgtggatga	tcaacgggat	taaattgcgt	catggtcgtt	3240

ccttaatcgg gatatccctg tggatggcgt gactgccagt gccagggtgtc ctgcccatt 3300
tcatcgagtg tgcgcgttac gcgccagttc agttcacggt cggctttgct ggcgcccgc 3360
cagtaggccg gaaggtcgcc ctgcgcgacgc ggtgcaaaat gataattaac cggtttgccg 3420
caggctttgc tgaaggcatt aaccacgtcc agcacgctgt tgcctacgcc agcggccgagg 3480
ttgtagatgt gtacgcctgg cttgttcgcc agtttttcca tcgccacgac gtgaccgtcc 3540
gccagatcca ttacgtggat gtaatcgcgt acgccagtac catcttcggt cggataatcg 3600
ttaccaaaaa tcgccagcga gtcgcgacgg cctacagcaa cctgggcgat gtatggcatc 3660
aggttattcg gaatgccttg cggatcttcg cccatatcgc ccgacggatg cgcgccaacc 3720
gggttgaagt agcgcagcag ggcaatgctc cagtccggct gggctttttg cagatcggtg 3780
aggatctgtt ccaccatcag cttgcttttg ccgtaagggc tttgcggtgt gccggtcggg 3840
aagctttcaa cgtatggaat tttgggctga tcgccataaa cggtggcgga ggagctaaaa 3900
ataaagtttt tgacgttagc ggcgcgcatg gcgctaataa ggcgcagagt gccgttgaca 3960
ttgttgcgt aatattccag cggtttttg accgattcgc ccacggcttt cagcccggcg 4020
aagtggatca cgggtgcatg agcgtgatcg tgcaggatct cggatcatcaa cgcttcgtta 4080
cgaatatcgc cttcaacaaa cgttggatgt ttgccgccta aacgctcgat aacaggcagt 4140
acgctgcgct tactgttaca gaggttatca agaattgatga catcatgacc gttttgcagt 4200
aattgcacac aggtatgact tccaatgtaa ccgctaccac cggtaaccag aactctcata 4260
attcgtcca ttaggcttat ggtatgaaat aaccatagca taacaaagat gcgaaaagt 4320
tgacatggaa taaattagtg gaatcgttta cacaagaatt tagccgtttt ttatgcgcga 4380
ttaagtgatt ataaaacaga gggtttatga atgattgcgc tttttatctg aaaaaagacg 4440
cggtttcatg cctgcatcgc tcgaaccgtt ggccggagag ggtgctaagg ccgcctccgg 4500
caaggtcagc actaccgacg ttaacggaaa ttacgctcta tatggaaagt ctgactgctg 4560
aagagcgcga gattatcaa 4579

<210> 2
<211> 35
<212> DNA
<213> Arti fi ci al sequence

<220>
<223> forward primer

<400> 2
ttactcagca ataaactgat attccgtcag gctgg 35

<210> 3
<211> 86
<212> DNA
<213> Arti fi ci al sequence

eol f-seq1 . txt

<220>

<223> reverse primer

<400> 3

ttgataatct cgcgctcttc agcagtcaga ctttccatat agagcgtaat ttccgttaac 60

gtcggtagtg ctgaccttgc cggagg 86