



- (51) International Patent Classification:
C12Q 1/68 (2006.01) *G06F 19/00* (2011.01)
- (21) International Application Number:
PCT/IB2012/000015
- (22) International Filing Date:
5 January 2012 (05.01.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/430,032 5 January 2011 (05.01.2011) US
61/475,632 14 April 2011 (14.04.2011) US
- (71) Applicant (for all designated States except US): **THE CHINESE UNIVERSITY OF HONG KONG** [CN/CN];
Shatin, New Territories, Hong Kong (CN).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LO, Yuk, Ming, Denis** [GB/CN]; 4th Floor, 7 King Tak Street Homantin, Kowloon, Hong Kong (CN). **CHIU, Wai, Kwun, Rossa** [AU/CN]; House 31, Double Haven 52 Ma Lok Path, Shat-

in, New Territories, Hong Kong (CN). **CHAN, Kwan, Chee** [CN/CN]; Flat A, 13/F, Block 34, Broadway Street, Mei Foo Sun Chuen, Kowloon, Hong Kong (CN). **TSUI, Bo, Yin** [CN/CN]; Rm H, 2/F, Block 3, Bayshore Towers, Ma On Shan, New Territories, Hong Kong (CN).

- (74) Agent: **GRIFFITH HACK**; GPO Box 1285, Melbourne, VIC 3001 (AU).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU,

[Continued on next page]

(54) Title: NONINVASIVE PRENATAL GENOTYPING OF FETAL SEX CHROMOSOMES




FIG. 1

(57) Abstract: Methods, apparatuses, and system are provided for analyzing a maternal sample to determine whether a male fetus of a pregnant female has inherited an X-linked mutation from the mother. A percentage of fetal DNA in the sample is obtained, and cutoff values for the two possibilities (fetus inherits mutant or normal allele) are determined. A proportion of mutant alleles relative to a normal allele on the X-chromosome can then be compared to the cutoff values to make a classification of which allele is inherited. Alternatively, a number of alleles from a target region on the X-chromosome can be compared to a number of alleles from a reference region on the X-chromosome to identify a deletion or amplification. The fetal DNA percentage can be computed by counting reactions with a fetal-specific allele, and correcting the number to account for a statistical distribution among the reactions.

WO 2012/093331 A1

TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

NONINVASIVE PRENATAL GENOTYPING OF FETAL SEX CHROMOSOMES

CROSS-REFERENCES TO RELATED APPLICATIONS

5 [0001] The present application claims priority from and is a non-provisional application of U.S. Provisional Application No. 61/430,032, entitled "Noninvasive Prenatal Genotyping Of Fetal Sex Chromosomes" by Lo et al. (008300US), filed January 5, 2011; and U.S. Provisional Application No. 61/475,632, entitled "Noninvasive Prenatal Genotyping Of Fetal Sex Chromosomes" by Lo et al. (008301US), filed April 14, 2011, the entire contents of
10 which are herein incorporated by reference for all purposes.

[0002] This application is related to commonly owned U.S. Patent Application No. 12/178,116 entitled "Determining a Nucleic Acid Sequence Imbalance" by Lo et al. (005210US), filed July 23, 2008, the disclosure of which is incorporated by reference in its entirety.

15

BACKGROUND

[0003] Hemophilias A and B are caused by heterogeneous mutations in the genes on chromosome X that encode for the *coagulation factor VIII (F8)* (Kemball-Cook G, Tuddenham EG, *Nucleic Acids Res.*, 25:128-132 (1997)) and *coagulation factor IX (F9)* (Giannelli F, Green PM, Sommer SS, et al., *Nucleic Acids Res.*, 26:265-268 (1998)),
20 respectively. There is a 25% chance for a pregnant hemophilia carrier to have an affected male fetus in each pregnancy. Prenatal diagnosis is an important aspect of reproductive choices for women in families with hemophilia (Lee CA, Chi C, Pavord SR, et al., *Haemophilia.*, 12:301-336 (2006)). In addition, it is also beneficial for appropriate obstetric management during labor and delivery as prolonged labor, invasive monitoring techniques
25 and instrumental deliveries should be avoided in affected fetuses to minimize potential fetal and neonatal hemorrhagic complications (Lee CA, Chi C, Pavord SR, et al., *Haemophilia.*, 12:301-336 (2006)). Therefore, the development of a noninvasive prenatal diagnostic approach for hemophilia is beneficial to both obstetricians and hemophilia families.

[0004] Current prenatal diagnostic methods for sex-linked diseases are typically invasive
30 and pose a risk to the fetus. The discovery of cell-free fetal DNA in maternal plasma has offered new opportunities for noninvasive prenatal diagnosis (Lo YMD et al., *Lancet.*,

350:485-487 (1997); Lo YMD, Chiu RWK, *Nat Rev Genet.*, 8:71-77 (2007)). A number of promising clinical applications have been developed based on the detection of paternally inherited genetic traits in maternal plasma. For example, the noninvasive detection of fetal sex and *RHD* status are useful for the clinical management of sex-linked diseases and RhD incompatibility (Bustamante-Aragones A et al., *Haemophilia.*, 14:593-598 (2008); Finning K et al., *BMJ.*, 336:816-818 (2008)). For monogenic diseases such as achondroplasia and β -thalassemia, the detection of the presence or absence of paternally inherited mutations in maternal plasma would allow one to diagnose autosomal dominant diseases or exclude autosomal recessive diseases of the fetuses, respectively (Saito H et al., *Lancet.*, 356:1170 (2000); Chiu RWK et al., *Lancet.*, 360:998-1000 (2002); Ding C et al., *Proc Natl Acad Sci U S A.*, 101:10762-10767 (2004)).

[0005] Despite the rapid development of the field, it has remained difficult to detect fetal alleles that are inherited from mothers who are carriers for the mutations. The difficulty is caused by the coexistence of fetal and maternal DNA in maternal plasma, and the maternally inherited fetal allele is indistinguishable from the background maternal DNA (Lo YMD, Chiu RWK, *Nat Rev Genet.*, 8:71-77 (2007)).

[0006] Therefore, it is desirable to provide accurate and efficient methods for determining whether a male fetus has inherited an X-linked mutation.

BRIEF SUMMARY

[0007] Methods, apparatuses, and system are provided for analyzing a maternal sample to determine whether a male fetus of a pregnant female has inherited an X-linked mutation from the mother. A percentage of fetal DNA in the sample is obtained, and cutoff values for the two possibilities (fetus inherits mutant or normal allele) are determined. A proportion of mutant alleles relative to a normal allele on the X-chromosome can then be compared to the cutoff values to make a classification of which allele is inherited. Alternatively, a number of alleles from a target region on the X-chromosome can be compared to a number of alleles from a reference region on the X-chromosome to identify a deletion or amplification. The fetal DNA percentage can be computed by counting reactions with a fetal-specific allele, and correcting the number to account for a statistical distribution among the reactions.

[0008] According to one embodiment, a method is provided for determining whether a male fetus of a pregnant female has an X-linked mutation. The pregnant female is heterozygous for a mutant and a normal allele at a locus on the X chromosome. Data is

received from a plurality of reactions, each involving one or more nucleic acid molecules from a biological sample. The biological sample includes nucleic acid molecules from the pregnant female and from the male fetus. The data includes a first set of quantitative data indicating a first amount of the mutant allele at the locus and a second set of quantitative data indicating a second amount of the normal allele at the locus. A parameter is determined from the first amount and the second amount, where the parameter represents a relative amount between the first and second amounts. A percentage Pf of fetal nucleic acid molecules in the biological sample is obtained. A first cutoff value for determining whether the fetus has inherited the mutant allele at the locus is calculated, where the first cutoff value is derived at least from a first proportion of $k/(1+k-Pf)$, where k is a number of mutant alleles on a mutant chromosome of the pregnant female, k being an integer equal to or greater than one. A second cutoff value for determining whether the fetus has inherited the normal allele at the locus is calculated, where the second cutoff value is derived at least from a second proportion of $[k(1-Pf)]/[1+k-kPf]$. The parameter is compared to at least one of the first and second cutoff values to determine a classification of whether the fetus has inherited the mutant allele or the normal allele.

[0009] According to another embodiment, a method is provided for determining whether a male fetus of a pregnant female has an X-linked mutation. The pregnant female is heterozygous for a mutation and a normal allele at a target region on the X chromosome. The mutation is a deletion or an amplification of the target region. Data from a plurality of reactions is received. Each reaction involves one or more nucleic acid molecules from a biological sample. The biological sample includes nucleic acid molecules from the pregnant female and from the male fetus. The data includes a first set of quantitative data indicating a first amount of the nucleic acid molecules that are from the target region and a second set of quantitative data indicating a second amount of the nucleic acid molecules that are from a reference region on the X chromosome. A parameter is determined from the first amount and the second amount, where the parameter represents a relative amount between the first and second amounts. A percentage Pf of fetal nucleic acid molecules in the biological sample is obtained. A first cutoff value for determining whether the fetus has inherited the mutation is calculated. The first cutoff value is dependent on the percentage Pf. A second cutoff value for determining whether the fetus has inherited the normal allele is calculated. The second cutoff value is dependent on the percentage Pf. The parameter is compared to at least one of the first and second cutoff values to determine a classification of whether the fetus has inherited the mutation or the normal allele.

[0010] According to another embodiment, a method of obtaining a percentage Pf of fetal nucleic acid molecules in a biological sample from a female pregnant with a fetus. Data is received from a plurality of reactions. Each reaction involves a plurality of nucleic acid molecules from a biological sample, which includes nucleic acid molecules from the pregnant female and from the fetus. A first allele is detected in the reactions. The first allele is shared by the mother and fetus at a locus where the pregnant female is homozygous and the fetus is either heterozygous or hemizygous. A corrected concentration Px of the first allele is calculated based on a number of reactions positive for the first allele, where Px is corrected for an expected statistical distribution of the first allele in the plurality of reactions. A second allele is detected in the reactions, where the second allele is specific to the fetus. A corrected concentration Py of the second allele is calculated based on a number of reactions positive for the second allele. Py is corrected for an expected statistical distribution of the second allele in the plurality of reactions. The fetal percentage Pf is then calculated using $[(2Py)/(Px+Py)]$.

[0011] Other embodiments are directed to systems, and computer readable media associated with methods described herein.

[0012] A better understanding of the nature and advantages of the present invention may be gained with reference to the following detailed description and the accompanying drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

[0013] FIG. 1 is a flowchart illustrating a method 100 for analyzing a maternal biological sample to diagnose an X-linked disorder in a fetus according to embodiments of the present invention.

[0014] FIG. 2A illustrates the two possibilities of the fetus inheriting the mutant allele or the normal allele. FIG. 2B shows a plot of cutoff values for classifying a sample as obtained using sequential probability ratio test (SPRT) according to embodiments of the present invention.

[0015] FIG. 3 is a flowchart illustrating a method 300 for determining whether a male fetus of a pregnant female has an X-linked mutation according to embodiments of the present invention.

[0016] FIG. 4 illustrates a method 400 for determining whether a male fetus has inherited an X-linked mutation according to embodiments of the present invention.

- 5 [0017] FIG. 5A shows a table 500 illustrating a dosage imbalance between mutant and wild-type alleles for mutations on chromosome X. FIG. 5B illustrates a first scenario for detecting an amplification when the pregnant subject is heterozygous at the locus of interest. FIG. 5C illustrates a second scenario for detecting an amplification when the pregnant subject is homozygous at the locus of interest.
- [0018] FIG. 6 is a flowchart illustrating a method 600 for determining whether a male fetus of a pregnant female has an X-linked mutation.
- [0019] FIG. 7 is a table 700 showing a dosage imbalance between the target and the reference loci for deletion and duplication mutations on chromosome X.
- 10 [0020] FIG. 8 is a flowchart illustrating a method 800 for obtaining a percentage P_f of fetal nucleic acid molecules in a biological sample from a female pregnant with a fetus according to embodiments of the present invention.
- [0021] FIG. 9 shows a table 900 with clinical information of the seven pregnant women who are carriers of hemophilia mutations.
- 15 [0022] FIG. 10 is a table 1000 showing oligonucleotide sequences and real-time PCR conditions for the allele-discriminative assays.
- [0023] FIG. 11 is a table 1100 showing fetal genotyping for rs6528633 in maternal plasma by digital RMD.
- [0024] FIG. 12 shows the validation of digital RMD assays with artificial DNA mixtures.
- 20 [0025] FIG. 13 is a table 1300 showing non-invasive detection of fetal hemophilia mutations in maternal plasma by digital RMD.
- [0026] FIG. 14 shows plots of SPRT analysis for fetal hemophilia mutations in maternal plasma samples. Case numbers are indicated at the top of the graphs. P_r , proportion of positive wells containing the mutant allele.
- 25 [0027] FIG. 15 shows digital RMD result for maternal plasma samples from normal pregnancies.
- [0028] FIG. 16 shows a block diagram of an example computer system 1600 usable with system and methods according to embodiments of the present invention.

DEFINITIONS

[0029] The term "*biological sample*" as used herein refers to any sample that is taken from a subject (e.g., a human, such as a pregnant woman) and contains one or more nucleic acid molecule(s) of interest.

5 [0030] The term "*nucleic acid*" or "*polynucleotide*" refers to a deoxyribonucleic acid (DNA) or ribonucleic acid (RNA) and a polymer thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogs of natural nucleotides that have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless
10 otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions), alleles, orthologs, SNPs, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-
15 base and/or deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* **19**:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* **260**:2605-2608 (1985); and Rossolini *et al.*, *Mol. Cell. Probes* **8**:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, small noncoding RNA, micro RNA (miRNA), Piwi-interacting RNA, and short hairpin RNA (shRNA) encoded by a gene or locus.

20 [0031] The term "*gene*" means the segment of DNA involved in producing a polypeptide chain. It may include regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

[0032] The term "*reaction*" as used herein refers to any process involving a chemical, enzymatic, or physical action that is indicative of the presence or absence of a particular
25 polynucleotide sequence of interest. An example of a "*reaction*" is an amplification reaction such as a polymerase chain reaction (PCR). Another example of a "*reaction*" is a sequencing reaction, either by synthesis, ligation, hybridization or degradation. An "*informative reaction*" is one that indicates the presence of one or more particular polynucleotide sequence of interest, and in one case where only one sequence of interest is present. The term "*well*" as
30 used herein refers to a reaction at a predetermined location within a confined structure, e.g., a well-shaped vial, cell, chamber in a PCR array, a droplet in an emulsion, a particle, a nanopore or an area on a surface.

[0033] The term "*overrepresented nucleic acid sequence*" as used herein refers to the nucleic acid sequence among two sequences of interest (e.g., a clinically relevant sequence and a background sequence) that is in more abundance than the other sequence in a biological sample.

5 [0034] The term "*based on*" as used herein means "*based at least in part on*" and refers to one value (or result) being used in the determination of another value, such as occurs in the relationship of an input of a method and the output of that method. The term "*derive*" as used herein also refers to the relationship of an input of a method and the output of that method, such as occurs when the derivation is the calculation of a formula.

10 [0035] The term "*quantitative data*" as used herein means data that are obtained from one or more reactions and that provide one or more numerical values. For example, the number of wells that show a fluorescent marker for a particular sequence would be quantitative data.

[0036] The term "*parameter*" as used herein means a numerical value that characterizes a quantitative data set and/or a numerical relationship between quantitative data sets. For
15 example, a ratio (or function of a ratio) between a first amount of a first nucleic acid sequence and a second amount of a second nucleic acid sequence is a parameter.

[0037] As used herein, the term "*locus*" or its plural form "*loci*" is a location or address of any length of nucleotides (or base pairs) which has a variation across genomes. The term "*alleles*" refers to alternative DNA sequences at the same physical genomic locus, which may
20 or may not result in different phenotypic traits. In any particular diploid organism, with two copies of each chromosome (except the sex chromosomes in a male human subject), the genotype for each gene comprises the pair of alleles present at that locus, which are the same in homozygotes and different in heterozygotes. A population or species of organisms typically includes multiple alleles at each locus among various individuals. A genomic locus
25 where more than one allele is found in the population is termed a polymorphic site. Allelic variation at a locus is measurable as the number of alleles (i.e., the degree of polymorphism) present, or the proportion of heterozygotes (i.e., the heterozygosity rate) in the population. As used herein, the term "*polymorphism*" refers to any inter-individual variation in the human genome, regardless of its frequency. Examples of such variations include, but are not limited
30 to, single nucleotide polymorphisms, simple tandem repeat polymorphisms, insertion-deletion polymorphisms, mutations (which may be disease causing) and copy number variations.

[0038] The term "*cutoff value*" as used herein means a numerical value whose value is used to arbitrate between two or more states (e.g. diseased and non-diseased) of classification for a biological sample. For example, if a parameter is greater than the cutoff value, a first classification of the quantitative data is made (e.g. diseased state); or if the parameter is less than the cutoff value, a different classification of the quantitative data is made (e.g. non-diseased state).

[0039] The term "*imbalance*" as used herein means any significant deviation as defined by at least one cutoff value in a quantity of the clinically relevant nucleic acid sequence from a reference quantity. For example, the reference quantity could be a ratio of 3/5, and thus an imbalance would occur if the measured ratio is 1:1.

[0040] The term "*sequenced tag*" as used herein refers to a string of nucleotides sequenced from any part or all of a nucleic acid molecule. For example, a sequenced tag may be a short string of nucleotides sequenced from a nucleic acid fragment, a short string of nucleotides at both ends of a nucleic acid fragment, or the sequencing of the entire nucleic acid fragment that exists in the biological sample. A nucleic acid fragment is any part of a larger nucleic acid molecule. A fragment (e.g. a gene) may exist separately (i.e. not connected) to the other parts of the larger nucleic acid molecule.

DETAILED DESCRIPTION

[0041] Current prenatal diagnostic methods for sex-linked diseases are typically invasive and pose a risk to the fetus. Cell-free fetal DNA analysis in maternal plasma provides a noninvasive means of assessing fetal sex in such pregnancies. However, the disease status of male fetuses remains unknown if mutation-specific confirmatory analysis is not performed. Here we have developed a noninvasive tests to diagnose if the fetus has inherited a causative mutation for sex-linked disease from its mother. One strategy is based on a relative mutation dosage (RMD) approach which we have previously established for determining the mutational status of fetuses for autosomal disease mutations. The RMD method is used to deduce if a fetus has inherited a sex-linked mutation on chromosome X by detecting if the concentration of the mutant or wild-type allele is overrepresented in the plasma of heterozygous women carrying male fetuses.

[0042] Embodiments provide the application of the RMD approach in prenatal diagnosis of X-linked disorders, e.g., hemophilia. A difference between the RMD analyses for autosomal diseases and X-linked diseases is that for the former there are three possible fetal genotypes

(i.e. homozygous normal, homozygous mutant, and heterozygous) while for the latter there are only two possible fetal genotypes. In the context of X-linked diseases, a male fetus possesses only one chromosome X and thus it would be of either mutant or wild-type genotype. The two outcomes for X-linked diseases, as compared with the three outcomes for autosomal diseases, can make the RMD approach more robust for X-linked diseases for a given degree of analytical precision. Embodiments can also be used for other sex-linked diseases, including but not limited to Duchenne muscular dystrophy, X-linked adrenoleukodystrophy, Becker muscular dystrophy, choroideremia, Hunter syndrome, Lesch Nyhan syndrome, Norrie's syndrome and ornithine transcarbamylase deficiency.

5 [0043] We illustrate the concept using hemophilia, a X-linked bleeding disorder, as an example. We correctly detected fetal genotypes for hemophilia mutations in all of the 12 studied maternal plasma samples obtained from pregnancies at-risk of hemophilia (a sex-linked disease) from as early as the 11th week of gestation. This development would make the decision to undertake prenatal testing less traumatic and safer for at-risk families.

15 I. DETERMINING SEX-LINKED MUTATION

[0044] FIG. 1 is a flowchart illustrating a method 100 for analyzing a maternal biological sample to diagnose an X-linked disorder in a fetus according to embodiments of the present invention. Method 100 is noninvasive and can use DNA circulating in the maternal biological sample.

20 [0045] In step 110, a pregnant subject with a known mutation on an X chromosome is identified. The mutation may be of any type as described herein, such as hemophilia. The mutation may be determined in a variety of ways, such as DNA sequencing, Southern blot analysis, PCR (including allele-specific PCR), melting curve analysis, etc. The mutation is such that only one of the X chromosomes of the pregnant subject has the mutation, i.e., the pregnant subject is heterozygous at a locus associated with the mutation. Embodiments can also be applied for the noninvasive prenatal diagnosis of other sex-linked disorders involving point mutations or sequence deletion, duplication or inversion, for examples, choroideremia and Norrie's syndrome.

25 [0046] In step 120, a biological sample of the pregnant subject is received. The sample may be any biological sample that contains fetal nucleic acids, such as plasma, urine, serum, and saliva. For example, maternal plasma sample can be collected from a pregnant carrier receiving obstetric care.

[0047] In step 130, the sex of the fetus is determined. The sex can be determined by detecting X and Y chromosomes. Through the detection of chromosome Y DNA sequences in maternal plasma, male fetuses could be identified with an accuracy of greater than 97% from the 7th week of gestation onwards. Unnecessary invasive testing could be avoided for female fetuses, as they are either unaffected or are disease carriers.

[0048] In step 140, the fetus is determined to be female, and then no further analysis is performed at step 145. Female fetuses are affected as carriers, except rare scenarios like skewed X-inactivation.

[0049] In step 150, the fetus is determined to be male, and then in step 155, DNA fragments on the X chromosome are analyzed. In one embodiment, a fetal mutation detection is performed by a relative mutation dosage (RMD) technique, which is described in more detail below. In another embodiment, a fetal mutation of a deletion or amplification is detected by comparing an amount of alleles at a target region (which includes the mutation in the mother) to an amount of alleles at a reference region, which is normal in the mother.

[0050] In step 157, a determination that the fetus did not inherit the mutated X chromosome of the maternal subject can be made. In step 159, a determination that the fetus did inherit the mutated X chromosome of the maternal subject can be made. The classification could be confirmed, if necessary, by a second maternal plasma sample taken at a later stage of pregnancy when fetal DNA percentages are higher (Lun FMF et al., *Clin Chem.*, 54:1664-1672 (2008)), allowing for more robust testing.

II. CLASSIFICATION BETWEEN NORMAL AND MUTANT

[0051] The analysis in step 155 of method 100 analyzes DNA fragments in the maternal sample. As the maternal sample also contains fetal DNA, a genotype of the X chromosome of the male fetus can be determined. For any mutation on chromosome X, there is always an allelic imbalance between the concentrations of the mutant and the wild-type alleles in the plasma of heterozygous women carrying male fetuses. The overrepresented allele is the one inherited by the fetus. In one embodiment, the genotype of the fetus can be determined by the RMD technique, which can include comparing a number of mutant alleles to a number of normal alleles in the maternal sample.

[0052] FIG. 2A illustrates the two possibilities of the fetus inheriting the mutant allele or the normal allele. The maternal DNA 210 is shown for a particular locus on the X chromosomes. The locus 215 is heterozygous with one allele being normal N (wild type) and

the other allele being mutant M. The mutation can be of various types, such as a different sequence, a deletion, an insertion, and an inversion. Each of these mutations can be identified as a different allele than the normal allele at locus 215.

5 [0053] The fetal DNA 220 is shown with the two possibilities. Since the male fetus has only one X chromosome, only one of the X chromosomes of maternal DNA 210 will be inherited by the male fetus. Possibility 222 shows the male fetus inheriting the mutant allele M. Possibility 224 shows the male fetus inheriting the normal allele N. The Y chromosome, which is smaller than the X chromosome, is also shown for each possibility.

10 [0054] The maternal sample (e.g. plasma) 230 will have a different proportion of mutant alleles to normal alleles depending on whether the fetus inherits the mutant or normal alleles. For possibility 222, the maternal sample will have more mutant alleles M since the male fetus had inherited the mutant allele M. This is because the fetal DNA would only contribute the mutant allele M, while the maternal DNA would contribute roughly equal parts of mutant allele M and normal allele N when a statistically significant amount of DNA is analyzed. For
15 possibility 224, the maternal sample will have more normal alleles N since the male fetus had inherited the normal allele N.

[0055] The number of DNA fragments showing the normal and mutant alleles can be counted in various ways, such as digital PCR, sequencing (including Sanger sequencing, massively parallel sequencing and single molecule sequencing), and other methods that
20 would allow the analysis of single DNA molecules or amplified groups of DNA molecules (e.g. clusters on a solid surface). Once the number of N and M alleles are counted, various techniques can be used to perform a classification, such as affected or unaffected (e.g. a diagnosis of whether the fetus has hemophilia or is normal). For instance, a parameter (e.g. a ratio or a difference) can be determined from the number of N and M alleles, and the
25 parameter can be compared against one or more cutoff values. The cutoff value(s) can be obtained through various statistical techniques, such as sequential probability ratio test (SPRT) (Zhou W, Galizia G, Lieto E, et al., *Nat Biotechnol.*, 19:78-81 (2001); Zhou W, Goodman SN, Galizia G, et al., *Lancet.*, 359:219-225 (2002)).

30 [0056] FIG. 2B shows a plot 250 of cutoff values for classifying a sample as obtained using SPRT according to embodiments of the present invention. The Y-axis shows the proportion P_r (an example of a parameter) of alleles that are mutant. The X-axis shows the number of alleles for locus 215 that are counted. The two curves correspond to the cutoff values for determining whether the fetus has the mutation (e.g. hemophilia), is normal, or is

unclassifiable. Samples with mutant allele proportion (P_r) above the upper boundary and below the lower boundary are classified as mutant and wild-type, respectively. Samples with P_r in between the two curves are unclassifiable and require additional digital analysis (e.g., data from additional PCR wells).

5 [0057] The particular cutoff values to use depends on the number of alleles counted. When only a few alleles are counted, there can be a large statistical variation, and thus the cutoff values require extreme values in P_r to confidently classify the sample as mutant or normal. As is described in more detail below, digital PCR may be used (where the Y-axis can be the proportion of positive wells containing the mutant allele and the X-axis can be the number of
10 positive wells). The position of the curves on the Y-axis can change depending on how the parameter is calculated, e.g., the unclassifiable area could be centered at 1.0 if the parameter was the number of N alleles divided by the number of M alleles.

[0058] In another implementation, where the mutation is a deletion or amplification, a comparison between a number of fragment at a target region (e.g. locus 215) where one of the
15 maternal X chromosomes has a deletion/amplification and a reference region (not having an amplification or deletion) can be used to identify the deletion/amplification. Such an implementation does not depend on an identification of a heterozygous locus, thus the pregnant subject can be homozygous at the target region. For a deletion, one would expect fewer fragments from the target region than from the reference region. For an amplification,
20 one would expect more fragments from the target region than from the reference region. The cutoff values can also be determined using SPRT or similar techniques.

III. RMD METHOD

[0059] FIG. 3 is a flowchart illustrating a method 300 for determining whether a male fetus of a pregnant female has an X-linked mutation according to embodiments of the present
25 invention. The pregnant female is heterozygous for a mutant and a normal allele at a locus on the X chromosome. Method 300 uses a relative amount of the mutant and normal allele to make a disease classification.

[0060] In step 310, data from a plurality of reactions is received. Each reaction involves one or more nucleic acid molecules from a biological sample, which includes nucleic acid
30 molecules from the pregnant female and from the male fetus. The reactions can be of various types, such as digital PCR reactions in various wells. Other embodiments can use other reactions, such as sequencing reactions (for example by a massively parallel sequencing

platform, including but not limited to the Illumina Genome Analyzer, Roche 454, Life Technologies SOLiD, Pacific Biosciences single molecule real-time sequencing or Ion Torrent), primer extension reactions, mass spectrometry, analysis using a nanopore, optical methods or hybridization to a fluorescent or other probe. Thus, the data can include

5 fluorescent signals from digital PCR wells, sequenced tags obtained from sequencing at least a portion of the DNA molecules in the wells, or other data resulting from such reactions.

[0061] The data from the reactions includes a first set of quantitative data indicating a first amount of the mutant allele at the locus, and a second set of quantitative data indicating a second amount of the normal allele at the locus. The amount for a particular allele at the
10 locus can be measured in various ways, such as by a total number of wells that are positive for a particular allele, counting the number of sequenced tags that include the particular allele and align to the locus (using a reference genome), and the number of sequenced nucleotides (basepairs) or the accumulated lengths of sequenced nucleotides (basepairs) that include the particular allele and align to the locus.

15 [0062] In step 320, a parameter is determined from the first amount and the second amount. The parameter represents a relative amount between the first and second amounts. The parameter may be, for example, a simple ratio of the first amount to the second amount, or the first amount to the second amount plus the first amount. In one aspect, each amount could be an argument to a function or separate functions, where a ratio may be then taken of
20 these separate functions. One skilled in the art will appreciate the number of different suitable parameters. For example, the parameter can be a ratio of the number of mutant alleles to the total number of mutant and wild-type alleles, denoted by P_r , present in a plasma sample.

[0063] In step 330, a percentage P_f of fetal nucleic acid molecules in the biological sample
25 is obtained. The percentage P_f provides a measurement of how much fetal DNA is in the maternal sample relative to the maternal DNA. If the percentage P_f is higher, then the overrepresentation of the inherited allele will become larger. The percentage can be expressed as a fraction between 0 and 1, with 1 being 100%.

[0064] In step 340, a first cutoff value for determining whether the fetus has inherited the
30 mutant allele at the locus is calculated. The first cutoff value is derived at least from a first proportion of $1/(2-P_f)$. Depending on how the parameter from step 320 is formulated, the proportion $1/(2-P_f)$ can be equal to the expected ratio of the first and second amounts if the mutant allele was inherited. The expected value can be input into a statistical function to

determine the cutoff. The cutoff value may be determined using many different types of methods, such as SPRT, false discovery, confidence interval, and receiver operating characteristic (ROC) curve analysis.

5 [0065] In step 350, a second cutoff value for determining whether the fetus has inherited the normal allele at the locus is calculated. The second cutoff value is derived at least from a second proportion of $(1-P_f)/(2-P_f)$.

10 [0066] In step 360, the parameter is compared to at least one of the first and second cutoff values to determine a classification of whether the fetus has inherited the mutant allele or the normal allele. As mentioned above, the classifications can include affected (mutation inherited) and unaffected (normal inherited), and also may include unclassified. A probability of accuracy may also be included with the classification, e.g., the accuracy may be determined by how much the parameter exceeds (above or below) a cutoff. In one implementation, the classification may be a score that is to be interpreted at a later date, for example, by a doctor.

15 [0067] The data that indicates an amount of an allele can be from a linked allele. Thus, an allele that is linked to either the mutant or the normal allele can be used instead of the normal and mutant alleles. For example, an allele at a polymorphic site linked to the mutant nucleic acid sequence can be an allele located on the same maternal haplotype as the mutant nucleic acid sequence, where the probability of recombination between the polymorphic site and the mutant nucleic acid sequence is less than a certain value, e.g. 1%. Thus, the polymorphic site can provide the same or similar quantitative data as measuring the mutant allele directly. As another example, an allele at a polymorphic site linked to the normal nucleic acid sequence can be an allele located on the same maternal haplotype as the normal nucleic acid sequence, where the probability of recombination between the polymorphic site and the mutant nucleic acid sequence is less than a certain value, e.g. 1%.

20

25

A. Example Using PCR With Plasma

[0068] As mentioned above, digital PCR can be used as the method for identifying DNA fragments that include the mutant or normal allele. In digital PCR, a sample is separated into a plurality of compartments (e.g., wells and beads). On average, each compartment contains less than one of any of the two alleles. Thus, a positive well can be counted as a single instance of a fragment containing the allele.

30

[0069] FIG. 4 illustrates a method 400 for determining whether a male fetus has inherited an X-linked mutation according to embodiments of the present invention. Digital PCR is used to determine a mutant allele proportion and the fetal DNA percentage. The fetal DNA percentage is used to determine a cutoff value to which the mutant allele proportion is compared, thereby providing a classification of whether the male fetus has inherited the mutation. As the mutant allele proportion is determined, embodiments can be referred to as the RMD method.

[0070] As illustrated, for each maternal plasma DNA sample, both the mutant DNA proportion (P_r) and the fetal DNA percentage P_f are determined by digital PCR, although other reactions that can identify certain sequences may be used. Steps for determining P_r is provided on the left (process 401), and steps for determining the fractional fetal DNA concentration P_f are on the right (process 402). As shown, P_r is determined using a real-time PCR assay targeting the mutation carried by the mother, while the fetal DNA percentage P_f is determined using the real-time PCR assay for the homologous *ZFY* and *ZFX* gene regions.

[0071] In step 410, the PCR mixture is prepared. As shown, the mixtures are different for the two measurements. For the P_r measurement (process 401), the mixture contains PCR primers to amplify a region on the X chromosome that includes the locus to be tested. The mixture also contains a fluorescent probe to identify the existence of a DNA fragment with the wild-type allele, and a fluorescent probe to identify the existence of a DNA fragment with the mutant allele. For the P_f measurement (process 402), the mixture contains primers for the *ZFY* and *ZFX* gene regions. The mixture also includes fluorescent probe to identify the existence of a DNA fragment containing a sequence from the *ZFX* gene, and a fluorescent probe to identify the existence of a DNA fragment containing a sequence from the *ZFY* gene.

[0072] In step 420, the reaction mixtures are loaded into a PCR machine. In one embodiment, the digital PCR is carried out in a microfluidics Digital Array (Fluidigm), which consists of 12 panels with each panel further partitioned into 765 reaction chambers. Each DNA sample (i.e. one for P_r and one for P_f) is analyzed using 6 panels, i.e., $765 \times 6 = 4590$ chambers. The PCR mixture can be first manually added into the sample inlet of each panel. The mixture is next aliquoted into 765 chambers in each panel automatically by an Integrated Microfluidics Circuit Controller (Fluidigm). Each chamber contains a final reaction volume of 6 nL. The cell-free DNA concentration in maternal plasma is typically very low such that there is less than one template molecule per chamber on average. Hence, the distribution of template molecules to the chambers follows the Poisson distribution. For other samples, one

may need to dilute the DNA sample before analysis. It will also be obvious to those of skill in the art that the digital PCR can be performed using methods well-known to those of skill in the art, e.g. microfluidics chips, nanoliter PCR microplate systems, emulsion PCR (including the RainDance platform), polony PCR, rolling-circle amplification, primer extension and mass spectrometry.

[0073] As shown for the P_r measurement, wells (chambers) containing a DNA fragment with the wild-type allele are shown in blue, and wells containing a DNA fragment with the mutant allele are shown in red. Wells that do not contain a template DNA molecule (i.e. no allele for which there is a probe) are shown simply as white. Similarly for the P_f

measurement, wells containing the ZFX gene are shown in blue, and wells containing the ZFY gene are shown in red.

[0074] In step 430, real-time PCR is performed, e.g., on the BioMark System (Fluidigm). Each well is carried through a series of cycles that amplify DNA regions that correspond to the primers in the corresponding mixture. Since most of the chambers contain zero or one template DNA molecule, the amplified products from a well originate from one template DNA molecule.

[0075] In step 440, the number of chambers with positive PCR amplifications are counted. For the process 401, the number of chambers that are positive for the wild-type allele can be counted and the number of chambers for the mutant allele can be counted. For process 402, the number of chambers that are positive for the ZFX gene can be counted and the number of chambers for the ZFY gene can be counted. In each process, the number of chambers that are positive for both of the alleles can also be identified. The detection of a positive chamber can be performed in various ways, such as detecting a fluorescent signal (e.g. each allele will emit a different color signal). For example, chambers containing the ZFX gene can emit a blue fluorescent signal, and wells containing the ZFY gene can emit a red fluorescent signal.

[0076] In step 450, the mutant DNA proportion (P_r) and the fetal DNA percentage P_f are calculated using the corresponding numbers counted in step 440. For example, the mutant allele proportion could be calculated as the number of chambers positive for the mutant allele divided by the total number of positive wells. As other examples, the denominator could be the total number of chambers that are positive only for one allele. Instead of a ratio involving the raw number of counts, the values could be concentrations themselves, effectively dividing the numerator and the denominator by any of the values above. Similar values can be used to calculate the fetal DNA percentage P_f using the equation $[(2Y)/(X+Y)]*100\%$, where Y is the

measured amount for the *ZFY* gene (e.g., count of positive chambers or proportion of positive chambers), and X is the measured amount for the *ZFX* gene.

5 [0077] Since there was less than one template molecule per reaction well, the actual number of template molecules distributed to each reaction chamber followed the Poisson distribution. Hence, the number of chambers for any allele can be Poisson-corrected using the equation $[-\ln((N-P)/N)] * N$, where N is the total number of reaction chambers analyzed, P is the number of chambers positive for the allele, and \ln is the natural logarithm. The Poisson-corrected values can then be used in a similar manner as mentioned above to determine the proportion P_r and the fetal DNA percentage P_f .

10 [0078] In step 460, the mutant DNA proportion (P_r) and the fetal DNA percentage P_f are used to perform a classification of whether the male fetus had inherited the mutation or not. As for method 300, cutoff values can be determined from the fetal DNA percentage P_f , e.g., as in steps 340 and 350. The cutoff may also be derived from (which includes equal to) an average reference template concentration (m_r), e.g., the experimentally measured percentage
15 of positive chambers for the wild-type allele can be used to determine the cutoff value used in step 460. This strategy can further minimize the amount of testing required before confident classification could be made. This is of particular relevance to plasma nucleic acid analysis where the template amount is often limiting.

B. SPRT

20 [0079] SPRT is a method which allows two probabilistic hypotheses to be compared as data accumulate. In other words, it is a statistical method to classify the results of digital PCR as being suggestive of the skewing towards either the mutant or the normal allele. It has the advantage of minimizing the number of wells to be analyzed to achieve a given statistical power and accuracy.

25 [0080] In an exemplary SPRT analysis, the experimental results would be tested against two alternative hypotheses. The first alternative hypothesis is accepted when the mutant allele is over-represented. The second alternative hypothesis is accepted when the mutant allele is under-represented. The measured P_r would be compared with at least one of the two cutoff values to accept the first or the second alternative hypotheses. If neither hypothesis is
30 accepted, the sample would be marked as unclassified which means that the observed digital PCR result is not sufficient to classify the sample with the desired statistical confidence. More data can be collected to obtain the desired statistical confidence.

[0081] A pair of curves, which depend on the amount of data collected, can define the probabilistic boundaries (cutoffs) for accepting or rejecting the hypotheses (Zhou W, Galizia G, Lieto E, et al., *Nat Biotechnol.*, 19:78-81 (2001); Zhou W, Goodman SN, Galizia G, et al., *Lancet.*, 359:219-225 (2002)). The SPRT curves delineated the required P_r (y -axis) for a given total number of positive reactions (x -axis) for classifying a fetal genotype. Hypothesis (i) or (ii) are accepted if the experimental P_r fell above the upper boundary or below the lower boundary, respectively. The equations for calculating the SPRT boundaries can be determined with varying levels of statistical confidence (e.g. adjusted to a threshold likelihood ratio of 8). In one aspect, the cutoff values of the SPRT curves are sample-specific. The cutoff values are dependent on the fractional fetal DNA concentration (fetal DNA percentage) as described above. The cutoff values can also depend on an average reference template concentration per PCR well (m_r) for a given set of reactions (Lo YMD et al., *Proc Natl Acad Sci U S A.* 2007;104:13116-13121 (2007); Lun FMF, Tsui NBY, Chan KCA, et al., *Proc Natl Acad Sci U S A.*, 105:19920-19925 (2008)). The reference template can refer to the allele that showed the lesser positive amplification counts in the sample.

[0082] SPRT can offer an advantage that a smaller amount of testing is required for a given level of confidence than other statistical methods. In practical terms, SPRT allows the acceptance or rejection of either of the hypotheses as soon as the required amount of data has been accumulated and thus minimizes unnecessary additional analyses. This feature is of particular relevance to the analysis of plasma nucleic acids which are generally present at low concentrations where the number of available template molecules is limiting. In addition to a strict classification, the classification may also include a percent accuracy. For example, a classification resulting from a comparison with a cutoff value may provide that a sample shows a likelihood of a nucleic acid sequence imbalance with a certain percentage, or equivalently that a determined imbalance is accurate to a certain percentage or other value.

[0083] For embodiments using SPRT, one may use the equations for calculating the upper and lower boundaries of the SPRT curves from El Karoui et al (El Karoui N, Zhou W, Whittemore AS, *Stat Med.* 25:3124-3133 (2006)). Furthermore, the level of statistical confidence preferred for accepting the first or second hypothesis could be varied through adjusting the threshold likelihood ratio in the equations. A threshold likelihood ratio of 8 has been shown to provide satisfactory performance to discriminate samples with and without allelic imbalance in the context of cancer detection. Thus, in one embodiment, the equations for calculating the upper and lower boundaries of the SPRT curves are:

Upper boundary = $[(\ln 8)/N - \ln \delta]/\ln \gamma$

Lower boundary = $[(\ln 1/8)/N - \ln \delta]/\ln \gamma$

5 where, $\delta = (1 - \theta_1)/(1 - \theta_2)$, $\gamma = \frac{\theta_1(1 - \theta_2)}{\theta_2(1 - \theta_1)}$, \ln is a mathematical symbol representing the natural logarithm, i.e. \log_e , N = total number of molecules (i.e. the sum of mutant and normal molecules analyzed),

10 θ_1 = proportion of mutant molecules to the total number of mutant and normal molecules if the first alternative hypothesis is true (i.e., the fetus has inherited the mutant allele); and

15 θ_2 = proportion of mutant molecules to the total number of mutant and wild-type molecules if the second alternative hypothesis is true (i.e., the fetus has inherited the normal allele).

[0084] For the determination of θ_1 for accepting the first alternative hypothesis, the sample is assumed to be obtained from a pregnant woman carrying a male fetus which has inherited the mutant (M) allele. θ_1 is determined to be $1/(2 - Pf)$, where Pf is the percentage of fetal DNA in the sample. Pf can be corrected for a statistical distribution, such as the Poisson distribution, as is described herein.

[0085] For the determination of θ_2 for accepting the second alternative hypothesis, the sample is assumed to be obtained from a pregnant woman carrying a male fetus which has inherited the normal (N) allele. θ_2 is determined to be $(1 - Pf)/(2 - Pf)$.

25 [0086] After an experimental determination of the numbers of mutant and wild-type molecules, the proportion of mutant molecules to the total number of mutant and wild-type molecules (Pr) can be calculated. The value of Pr can then be compared with the cutoff values to determine if the mutant or the wild-type alleles are overrepresented in the maternal plasma.

C. Poisson correction of cutoff values

30 [0087] In one embodiment using digital PCR, the average concentration per well (reaction or reaction mixture) is determined, and the expected number of wells showing that sequence may be calculated. This amount may be expressed as a percentage, a fractional value, or an integer value. In one implementation, a Poisson distribution is assumed for the distribution of the normal (N) allele, or the mutant allele, among the reaction mixtures of the wells of the measurement procedure, such as digital PCR. In other implementations, other distribution functions are used, such as a binomial distribution.

35

[0088] The Poisson equation is: $P(n) = \frac{m^n e^{-m}}{n!}$ where, n = number of template molecules

per well; $P(n)$ = probability of n template molecules in a particular well; and m = average number of template molecules in one well in a particular digital PCR experiment.

Accordingly, the probability of any well not containing any molecule of the normal allele at
5 an average normal-allele concentration of 0.5 would be:

$$P(0) = \frac{0.5^0 e^{-0.5}}{0!} = e^{-0.5} = 0.6065.$$

[0089] Hence, the probability of any well containing at least one molecule of the normal allele would be: $1 - 0.6065 = 0.3935$. Therefore, ~39% of the wells would be expected to
10 contain at least one molecule of the normal allele. In one embodiment, $P(0)$ for mutant or wild-type can be determined from an experimentally derived proportion of negative wells (e.g. using digital PCR). $P(0)$ can then be used to calculate the average number of molecules per well (m). The parameter can then be calculated from the average number of molecules per well, e.g., mutant average divided by the sum of the averages for the mutant and normal
15 alleles. Given this relationship between the number of positive wells and the number of molecules, an alternative is to correct the number of positive wells to provide the number of molecules (as described above via equation $[-\ln((N-P)/N)] * N$, where N is the total number of reaction chambers analyzed and P is the number of chambers positive for the allele).

[0090] The measurement of m_r may be performed through a variety of mechanisms as
20 known or will be known to one skilled in the art. In one embodiment, the value of m_r is determined during the experimental process of digital PCR analysis. As the relationship between the value of m_r and the total number of wells being positive for the reference allele can be governed by a distribution (e.g. the Poisson distribution), m_r can be calculated from the number of wells being positive for the reference allele using this formula:

25 $m_r = -\ln(1 - \text{proportion of wells being positive for the reference allele})$

This approach provides a direct and precise estimation of m_r in the DNA sample used for the digital PCR experiment.

[0091] This method may be used to achieve a desired concentration. For example, the extracted nucleic acids of a sample may be diluted to a specific concentration, such as one
30 template molecule per reaction well. In an embodiment using the Poisson distribution, the expected proportion of wells with no template may be calculated as e^{-m} , where m is the

average concentration of template molecules per well. For example, at an average concentration of one template molecule per well, the expected proportion of wells with no template molecule is given by e^{-1} , i.e., 0.37 (37%). The remaining 63% of wells will contain one or more template molecules. Typically, the number of positive wells in a digital PCR run would then be counted. The definition of informative wells and the manner by which the digital PCR data are interpreted depends on the application.

[0092] In other embodiments, the average concentration per well, m_r , is measured by another quantification method, for example, quantitative real-time PCR, semi-quantitative competitive PCR, and real-competitive PCR using mass spectrometric methods.

10 [0093] In one implementation, the proportion of the mutant allele to the normal allele can be calculated using corrected concentrations. The concentration m for each allele can be calculated as described above. The concentration for each allele can then be determined, and a proportion Pr of the concentrations can be used as the experimentally derived and distribution-corrected proportion to compare to the expected proportion for each hypothesis
15 (e.g. mutant or wild-type inheritance). For example, the experimentally determined Pr of a tested sample can be calculated using the equation: (concentration of mutant allele) / (concentration of mutant + wild-type alleles). In another implementation, the proportion of the number of wells for each allele is used. The expected proportion (cutoff value) can also be corrected based on a statistical distribution.

20 *D. Illustration*

[0094] FIG. 5A shows a table 500 illustrating a dosage imbalance between mutant and wild-type alleles for mutations on chromosome X according to embodiments of the present invention. To illustrate the calculation, a maternal plasma sample containing a total of 100 genomic equivalents (GE) of DNA with 10% fetal DNA was used. For the maternal genome,
25 one GE contains two copies of the alleles, i.e., one copy each of the M and the N allele. This provides 90 copies each of the mutant and normal alleles. For the fetal genome, one GE contains one copy of the X-linked allele, i.e., one copy of either the mutant (M) or the normal (N) allele. This provides 0 or 10 copies of each allele depending on which allele is inherited by the fetus.

30 [0095] In table 500, the upper row corresponds to the fetus inheriting the normal allele, and thus the ratio of mutant to normal alleles is less than 1. In the lower row, the fetus inherited the mutant allele, and thus the ratio of mutant to normal alleles is greater than 1.

E. Deletions, Amplifications, Insertions, and Inversions

[0096] Methods 300 and 400 can be applied in additional situations besides a standard SNP. Embodiment can be further applied to noninvasive detection of fetal mutations involving deletion, amplification (e.g. duplication), insertion, and inversion, e.g., of a large DNA segment. Examples of such mutations are relevant to X-linked diseases such as Duchenne muscular dystrophy, Becker muscular dystrophy and ornithine transcarbamylase deficiency. The approach is to detect the mutant allele by targeting the junctions of the rejoining sequences of the deletion, between the amplified (e.g. duplicated) DNA segments, or between the inverted and the adjacent normal DNA segments. The fetal genotype could then be deduced by the dosage imbalance between the normal and the mutant alleles with the methods described herein.

[0097] FIG. 5B illustrates a first scenario for detecting an amplification when the pregnant subject is heterozygous at the locus of interest. For amplifications on a first chromosome, where the amplified allele B is different than the non-amplified allele A, there will be different junctions for the various copies B1 and B2 of the amplified allele B. This is because the amplified copies B1 and B2 will be at different locations on the first chromosome. If one of the junctions is unique (e.g., the junction at the start of B or at the end of B2 is unique, while the junctions between B-B1 and B1-B2 are the same), the unique junction can be used as the mutant allele for comparison to the normal allele on the other chromosome. In this manner, the cutoff values can be derived in the same manner as in steps 340 and 350. Alternatively, all of the instances of the amplified allele B (i.e. is B, B1, and B2) can be used, regardless of location in the first chromosome. In such an embodiment, $\theta_1 = (1+n)/(2+n-Pf)$, and $\theta_2 = [(1+n)(1-Pf)]/[2+n-Pf(1+n)]$, where n is the number of additional copies ($n=2$ as shown), where n is an integer equal to or greater than zero. These formulas can also be written as $\theta_1 = k/(1+k-Pf)$ and $\theta_2 = [k(1-Pf)]/[1+k-kPf]$, where k is the number of copies of the mutant allele (which can be a newly formed junction) on the mutant chromosome, where k is an integer equal to or greater than one.

[0098] Junctions can also be used in a similar manner for RMD analysis for mutations on autosomes, but the values of θ_1 and θ_2 would need to be adjusted. For example, if the fetus inherited the amplification mutation, the sample would have the same ratio as the mother, assuming the chromosome inherited from the father is the normal chromosome. In this scenario, the value of θ_1 would be $k/(k+1)$, where k is the number of additional junctions created by the amplification mutation, and the additional junction is used as the mutant allele (thus for a duplication or a deletion, there is one mutant allele and for a triple amplification

there are two mutant alleles, and so on). If the fetus inherited the normal chromosome from the mother, then the value of θ_2 would be $k(1-Pf)/[k+1+(1-k)Pf]$.

[0099] FIG. 5C illustrates a second scenario for detecting an amplification when the pregnant subject is homozygous at the locus of interest. When the amplified allele and the non-amplified allele are the same (A as shown), two junctions 510 will be the same (for the two alleles at the normal location), and the additional (new) junction(s) 520 of the additional copies of the allele will be different, since these additional alleles will be at a different genomic location. The additional junctions can be used as the mutant allele, and the normal junction 510 can be used as the normal allele. One can use just one of the additional junctions 520 for the additional allele(s) (there would be only one for a duplication). In such an embodiment, $\theta_1 = 1/(3-Pf)$; and $\theta_2 = (1-Pf)/(3-2Pf)$. Note that the amount of additional copies is not used in such formulas since just one additional junction is used.

[0100] If there are more than one additional copy of A, the additional junction that is used should be chosen to be unique (e.g. the junction after the last amplified copy of A). Or, one could sum all (or some number more than 1) of the additional junctions and compare to the junctions of the two alleles at the normal location. In such an embodiment, $\theta_1 = n/(n+2-Pf)$; and $\theta_2 = n(1-Pf)/[n+2-Pf(n+1)]$, where n is the number of new junctions 520 that are used. Note that the amount of additional copies is used in such formulas since just more than one additional junction is used. Junctions can also be used in a similar manner for RMD analysis for mutations on autosomes, but the values of θ_1 and θ_2 would need to be adjusted. For example, if the fetus inherited the amplification mutation (amplification), the sample would have the same ratio (e.g., 1:2 for a duplication) as the mother, assuming the chromosome inherited from the father does not have the mutation. In this scenario, the value of θ_1 would be $n/(n+2)$, where n is the number of additional junctions created by the amplification mutation. If the fetus inherited the normal chromosome from the mother, then the value θ_2 would be $n(1-Pf)/(n+2-nPf)$. Another approach for detecting deletions and amplifications is described below.

IV. TARGET REGION VS REFERENCE REGION

[0101] In the RMD method described above, different junctions can be used as the alleles when the mutation is a deletion, amplification, insertions, or inversion. Another approach, which is applicable to deletion and amplification (e.g. duplication) mutations, is to compare the amount of molecules arising from the target region (i.e. the region that is deleted or amplified) to the amount of molecules arising from a reference region. Any genomic locus on

chromosome X not affected by the deletion (or amplification) can be used as a reference locus/region, for example, the *ZFX* gene if it is not deleted or amplified.

[0102] The ratio (R) of the number of molecules from the target region to the number of molecules from the reference region (or some other parameter representing a relative amount) can be used to determine whether the mutation is inherited. In a non-pregnant woman who is carrying the deletion mutation, the expected value of R would be 0.5 because only half of the X chromosomes (those carrying the normal allele) would contribute to the amount of target molecules in the plasma. When a woman carrying this deletion mutation is pregnant with a male fetus, the expected value of R would deviate from 0.5 due to the contribution of the DNA from the one extra X chromosome from the male fetus. The expected deviation of R would depend on whether the mutation is a deletion or an amplification.

[0103] FIG. 6 is a flowchart illustrating a method 600 for determining whether a male fetus of a pregnant female has an X-linked mutation. The pregnant female is heterozygous for a mutation and a normal allele at a target region on the X chromosome. The mutation is a deletion or an amplification of the target region.

[0104] In step 610, data from a plurality of reactions is received. The data may be of the same type as received in step 310 of method 300. Each reaction involves one or more nucleic acid molecules from a biological sample, which includes nucleic acid molecules from the pregnant female and from the male fetus. The data includes a first set of quantitative data indicating a first amount of the nucleic acid molecules that are from the target region, and a second set of quantitative data indicating a second amount of the nucleic acid molecules that are from a reference region on the X chromosome. The amounts may be computed in various ways, e.g., as described above for step 310.

[0105] In step 620, a parameter is determined from the first amount and the second amount. The parameter represents a relative amount between the first and second amounts. In one embodiment, the parameter is a ratio T of the first amount to the second amount. Other embodiments can use parameters as described herein, such the first amount divided by a sum of the first amount and the second amount.

[0106] In step 630, a percentage Pf of fetal nucleic acid molecules in the biological sample is obtained. The percentage Pf can be calculated as described herein. The percentage Pf can also be determined from a distribution corrected (e.g. Poisson-corrected) values for counting fetal specific molecules.

[0107] In step 640, a first cutoff value for determining whether the fetus has inherited the mutation is calculated. The first cutoff value is dependent on the percentage Pf. The specific equations for calculating the first cutoff value depends on whether the mutation is a deletion or an amplification.

5 [0108] In step 650, a second cutoff value for determining whether the fetus has inherited the normal allele is calculated. The second cutoff value is dependent on the percentage Pf. The specific equations for calculating the first cutoff value depends on whether the mutation is a deletion or an amplification.

10 [0109] In step 660, the parameter is compared to at least one of the first and second cutoff values to determine a classification of whether the fetus has inherited the mutant or the normal allele. The classifications can be of the same type as step 360, such as affected, unaffected, or unclassified (or a raw score).

15 [0110] FIG. 7 is a table 700 showing a dosage imbalance between the target and the reference loci for deletion and duplication mutations on chromosome X. Table 700 illustrates the calculation of the degree of allelic imbalance. An increase or decrease of R when compared with R of a non-pregnant woman carrying the same deletion mutation would indicate a normal or affected fetus, respectively. Conversely, in a non-pregnant woman who is carrying the segmental amplification, such as a duplication as shown in table 700, the expected value of R would be 1.5 due to the contribution of a doubled dose of target
20 molecules from the mutant allele. When a woman carrying this duplication mutation is pregnant, an increase or decrease of R when compared with R of a non-pregnant woman carrying the same duplication mutation would indicate an affected or normal fetus, respectively.

25 [0111] The degree of increase or decrease of R in each scenario is dependent on the fractional fetal DNA concentration (Pf) in a sample. In one embodiment, SPRT analysis can be used to determine if R is statistically significantly increased or decreased compared to the non-pregnant women carrying the same mutation. The equations for calculating the upper and lower boundaries (cutoff values) of the SPRT can have a similar structure of:

30 Upper boundary = $[(\ln 8)/N - \ln \delta]/\ln \gamma$;

Lower boundary = $[(\ln 1/8)/N - \ln \delta]/\ln \gamma$

where $\delta = (1 - \theta_1)/(1 - \theta_2)$; $\gamma = \frac{\theta_1(1 - \theta_2)}{\theta_2(1 - \theta_1)}$; \ln is a mathematical symbol representing the natural logarithm, i.e. \log_e ; N = total number of mutant and reference molecules;

5 θ_1 = ratio (R_1) of target molecules to the reference molecules if the first alternative hypothesis is true (i.e., R_1 is increased when compared with the value of R of a non-pregnant woman carrying the same mutation)

θ_2 = ratio (R_2) of target molecules to reference molecules if the second alternative hypothesis is true (i.e., R_2 is decreased when compared with the value of R of a non-pregnant woman carrying the same mutation)

10

[0112] θ_1 describes the situation in which the ratio of the amount of target molecules to the amount of reference molecules is increased when compared to the corresponding ratio of a non-pregnant woman carrying the same mutation, e.g., a normal case for a deletion mutation, or a mutant case for a duplication mutation. Similarly, θ_2 can describe the situation in which
15 the ratio of the amount of target molecules to the amount of reference molecules is decreased when compared to the corresponding ratio from a non-pregnant woman carrying the same mutation, e.g., a mutant case for a deletion mutation, or a normal case for a duplication mutation.

[0113] In one embodiment, for a deletion mutation, θ_1 is calculated as the sample is
20 assumed to be obtained from a pregnant woman carrying a male fetus that has inherited the normal (N) allele. θ_1 is determined to be $1/(2-Pf)$. θ_2 is calculated as the sample is assumed to be obtained from a pregnant woman carrying a male fetus that has inherited the mutation (e.g. the chromosome X with the deletion mutation). θ_2 is determined to be $(1-Pf)/(2-Pf)$.

[0114] In another embodiment, for duplication mutation, θ_1 is calculated as the sample is
25 assumed to be obtained from a pregnant woman carrying a male fetus that has inherited the mutation (i.e. the chromosome X with the duplication mutation). θ_1 is determined to be $(3-Pf)/(2-Pf)$. θ_2 is calculated as the sample is assumed to be obtained from a pregnant woman carrying a male fetus that has inherited the normal (N) allele. θ_2 is determined to be $(3-2xPf)/(2-Pf)$. The generalized formulas for any level of amplification is: θ_1 is $(n+2-Pf)/(2-Pf)$,
30 and θ_2 is $[n+2-Pf(n+1)]/(2-Pf)$, where n is the number of additional copies of amplified segments.

V. DETERMINING FETAL PERCENTAGE

[0115] As mentioned above, probabilities $P(n)$ for certain alleles (e.g. specific to chromosome X and a fetal-specific sequence) can be used to adjust the percentage (Pf) of

fetal DNA in the sample. This adjusted Pf can then be used to calculate the cutoffs for determining whether the mutant or the wild-type allele is inherited.

5 [0116] FIG. 8 is a flowchart illustrating a method 800 for obtaining a percentage Pf of fetal nucleic acid molecules in a biological sample from a female pregnant with a fetus according to embodiments of the present invention. The biological sample includes nucleic acid molecules from the pregnant female and from the fetus.

10 [0117] In step 810, data is received from a plurality of reactions. Each reaction involves a plurality of nucleic acid molecules from a biological sample. In one aspect, the reactions may be of any type where a reaction is considered positive for a particular allele if one or more of the alleles are present in the reaction.

[0118] In step 820, a first allele is detected in the reactions. The first allele is shared by the mother and fetus at a locus where the pregnant female is homozygous and the fetus is either heterozygous or hemizygous. In one embodiment, the first allele is the X chromosome.

15 [0119] In step 830, a corrected concentration Px of the first allele is calculated based on a number of reactions positive for the first allele. Px is corrected for an expected statistical distribution of the first allele in the plurality of reactions. For example, Px can be corrected based on the Poisson distribution. In one embodiment, a first corrected concentration for a first allele shared by the mother and fetus where the mother is homozygous and the fetus is either heterozygous or hemizygous is calculated, e.g., as $[-\ln((N-P1)/N)]*N$, where N is the total number of reaction chambers analyzed, P1 is the number of chambers positive for the first allele, and ln is the natural logarithm.

20 [0120] In step 840, a second allele that is specific to the fetus is detected. In one embodiment, the second allele is on the Y chromosome, where the fetus is a male fetus. In another embodiment, the fetal-specific allele is a paternally-inherited allele on an autosome. In yet another embodiment, the fetal-specific allele includes a methylation marker specific to the fetus.

25 [0121] In step 850, a corrected concentration Py of the second allele is calculated based on a number of reactions positive for the second allele. Py is corrected for an expected statistical distribution of the second allele in the plurality of reactions. For example, Py can be corrected based on the Poisson distribution. In one embodiment, a second corrected concentration for a fetal-specific allele which the fetus is heterozygous or hemizygous can be calculated as $[-\ln((N-P2)/N)]*N$, where N is the total number of reaction chambers analyzed,

P2 is the number of chambers positive for the fetal-specific allele, and \ln is the natural logarithm.

[0122] In step 860, the percentage Pf of fetal nucleic acid molecules in the biological sample is calculated using $[(2P_y)/(P_x+P_y)]$, which can provide a fractional value. The fetal DNA percentage can be calculated using the equation $[(2P_2)/(P_1 + P_2)]*100\%$.

VI. EXAMPLES

[0123] Seven women who were carriers of hemophilia (three carriers of hemophilia A, four carriers of hemophilia B) and pregnant with male fetuses were recruited from the Royal Free Hospital, London, UK. We also recruited 20 pregnant women (non-carriers of hemophilia) each pregnant with a singleton healthy male fetus. Ten of them were recruited from the Royal Free Hospital, London, UK and the other ten were recruited from the Prince of Wales Hospital, Hong Kong. Clinical information of the cases is shown in table 900 of FIG. 9, which shows clinical information of the seven pregnant women who are carriers of hemophilia mutations.

[0124] All women were recruited with informed consent. Ethical approvals were granted by the respective institutional boards. Ten milliliters of peripheral blood samples was collected into EDTA tubes from the pregnant women. For five of the pregnant hemophilia carriers, peripheral blood samples were taken on two occasions during their pregnancies (table 900). None of the pregnant hemophilia carriers in this study had invasive prenatal testing. Fetal sex and hemophilia status were confirmed following delivery. For the ten unaffected pregnant women recruited in Hong Kong, placental tissues were also collected following deliveries.

[0125] We centrifuged the blood samples at 1600g for 10 min at 4°C. The plasma portion was recentrifuged at 16000g for 10 min at 4°C. Maternal plasma and buffy coat samples were stored at -20°C until further processing. All samples collected in the UK were processed and stored frozen locally and were shipped on dry ice to Hong Kong. We extracted DNA from maternal plasma with the QIAamp DSP DNA Blood Mini Kit (Qiagen) following the manufacturer's instructions. Buffy coat DNA was extracted using the Illustra DNA Extraction Kit (GE Healthcare) following the manufacturer's protocol.

30 *Genotyping of rs6528633 SNP and hemophilia mutations*

[0126] To assess the feasibility of the RMD approach, we studied a SNP (rs6528633) on chromosome X. This SNP was chosen for illustration purposes and other SNPs can be used.

The fetal and maternal SNP genotypes were determined using DNA obtained from the placental and maternal buffy coat samples, respectively. Genotyping was performed using MassARRAY homogenous MassEXTEND (hME) assays (Sequenom) as previously described (Tsui NBY, Chiu RWK, Ding C, et al., *Clin Chem.*, 51:2358-2362 (2005); Tsui NBY, Chiu RWK, Ding C, et al., *Clin Chem.*, 51:2358-2362 (2005)). Genomic DNA obtained from the peripheral blood samples of the pregnant hemophilia carriers was used for hemophilia mutation detection. PCRs were performed for all exons covering coding regions, intron/exon boundaries, promoter and 3' UTR. Cycle sequencing was carried out using Big Dye Terminators V1.1 (Applied Biosystems) and analyzed on an Applied Biosystems 3100 Avant Genetic Analyser.

Digital RMD reactions for maternal plasma analyses

[0127] The experimental workflow of digital RMD is illustrated in FIG. 4 according to certain embodiments of the present invention. We measured the fractional fetal DNA concentrations in the maternal plasma samples using the previously described digital *ZFY/X* assay, which quantified the homologous *ZFY* and *ZFX* gene loci located on chromosomes Y and X, respectively (Lun FMF et al., *Clin Chem.*, 54:1664-1672 (2008); Lun FMF, Tsui NBY, Chan KCA, et al., *Proc Natl Acad Sci U S A.*, 105:19920-19925 (2008)). For the rs6528633 SNP, a real-time PCR assay with two allele-specific TaqMan probes (Applied Biosystems) was designed to distinguish the two SNP alleles. For the mutations of the pregnant cases at risk for hemophilia, a real-time PCR assay for allelic discrimination was designed for each mutation. Each assay contained two allele-specific TaqMan probes for the mutant and the wild-type alleles. The primer and probe sequences are listed in table 1000 in FIG. 10, which shows oligonucleotide sequences and real-time PCR conditions for the allelic-discriminative assays. In other embodiments, the fractional fetal DNA concentration can be determined by using a sequence that is differentially methylated between the fetal and maternal DNA in maternal plasma (for examples, see Chim SS et al., *Proc Natl Acad Sci USA.*, 102: 14753-14758 (2005); Chan KCA et al., *Clin Chem.*, 52: 2211-2218 (2006)).

[0128] We performed digital PCR analyses on the BioMark System (Fluidigm) using the 12.765 Digital Arrays (Fluidigm) (Lun FMF et al., *Clin Chem.*, 54:1664-1672 (2008)). Six of the 12 panels on the Digital Array were used for each DNA sample, which corresponded to 4590 individual PCRs. The reaction for one sample (6 panels) was set up using 2X TaqMan Universal PCR Master Mix (Applied Biosystems) in a reaction volume of 52 μ L. The reactions were set up according to the manufacturer's protocol with the primer and probe

compositions listed in table 1000 of FIG. 10. Each reaction mix contained 18.2 μ L of the DNA sample. The reaction mixture was automatically loaded onto the Digital Array by the NanoFlex IFC Controller (Fluidigm). The reactions were carried out on the BioMark System (Fluidigm). The reactions were initiated at 50°C for 2 minutes, followed by 95°C for 10
5 minutes, and 45 cycles of 95°C for 15 seconds and assay-specific annealing temperatures (FIG. 10 TABLE 3) for 1 minute. For a sample that remained unclassified by the RMD with data from one 4590-well digital PCR set, additional 4590-well digital PCR sets were carried out until a genotype call could be made.

Results

10 *Principle of digital RMD for X-linked polymorphisms*

[0129] Embodiments can use digital PCR to measure the concentration difference between the total amount (maternal- plus fetal-derived) of mutant and wild-type alleles in the plasma of heterozygous pregnant women carrying male fetuses. Since a male fetus possesses a single chromosome X, the relative concentration between the wild-type and the mutant allele is
15 always in dosage imbalance (FIG. 2A). An over- or under-representation of the mutant allele represents an affected or normal fetus, respectively. We used SPRT to test for dosage imbalance. A pair of SPRT curves was constructed (FIG. 2B). Samples with data points above the upper curve or below the lower curve were classified as affected or normal, respectively. Samples with data points in between the two curves were not classified because
20 of insufficient statistical power and additional digital PCRs would be performed.

Noninvasive determination of the fetal genotype for a SNP on chromosome X

[0130] We used a SNP, rs6528633 (A/T polymorphism), on chromosome X as a model to assess the practical feasibility of the RMD approach for determining the fetal genotype of a locus on chromosome X. The current RMD analysis is relevant to at-risk pregnant cases, i.e.,
25 pregnant women who are heterozygous for mutations on chromosome X and are carrying male fetuses. Hence, we studied the plasma samples from ten pregnant women who were heterozygous for the SNP on chromosome X and were carrying male fetuses. We developed an allele-discriminative digital real-time PCR assay to measure the concentrations of the A- and T-allele in each sample. We further measured the fractional fetal DNA concentrations
30 with the *ZFY/X* assay. The digital RMD result is shown in table 1100 of FIG. 11, which shows fetal genotyping for rs6528633 in maternal plasma by digital RMD.

[0131] For all of the cases, the fetal SNP genotypes were concordant with the SPRT classification. The fractional fetal DNA concentrations (fetal % in table 1100) ranged from 5% to 24%. The result hence confirmed the feasibility of the digital RMD strategy.

Digital RMD for hemophilia mutation detection in DNA mixtures

5 [0132] We next applied the digital RMD approach for hemophilia mutation detection. We developed seven duplex digital real-time PCR assays to detect three mutations in the *F8* gene, four mutations in the *F9* gene and their corresponding wild-type counterparts. We evaluated the performance of the digital PCR assays by constructing artificial DNA mixtures that simulated the composition of maternal plasma samples with a minority male fetal DNA
10 component amongst a majority maternal DNA background. We mixed 10% or 20% of placental DNA obtained from an unaffected male fetus with blood cell DNA obtained from women heterozygous for the corresponding mutations. FIG. 12 shows the validation of digital RMD assays with artificial DNA mixtures. The artificial mixtures were constructed to simulate the fetal and maternal DNA compositions in maternal plasma. As shown in table
15 1200 of FIG. 12, the genotypes of the placental DNA, which mimicked the fetal DNA in maternal plasma, were correctly detected in all of the DNA mixtures by digital RMD analysis.

Detection of fetal hemophilia mutations in maternal plasma

[0133] We tested the digital RMD method for detecting fetal genotypes for the hemophilia
20 mutations through maternal plasma DNA analysis. We carried out digital PCR on 12 plasma samples obtained from seven pregnant women heterozygous for the causative mutations (TABLE 900). All of the cases involved male fetuses. We also measured the fractional fetal DNA concentrations in the maternal plasma samples by the *ZFY/X* assay. The digital RMD results are shown in table 1300 of FIG. 13, which shows non-invasive detection of fetal
25 hemophilia mutations in maternal plasma by digital RMD.

[0134] The fetal genotypes were correctly classified in all studied cases by the SPRT algorithm (FIG. 14). For three of the cases (H26a, H25a and H12a), the fetal DNA proportions were less than 10%. Hence, the degree of quantitative difference between the amount of mutant and the wild-type alleles was too small to be classified with data from one
30 4590-well digital PCR set. Additional 4590-well digital PCR sets were therefore performed until classifications could be made.

[0135] As controls, we also studied five maternal plasma samples obtained from normal pregnant women using each of the mutation-specific assays. FIG. 15 shows digital RMD result for maternal plasma samples from normal pregnancies. As shown in table 1500 of FIG. 15, no mutant alleles were detected in most of the cases. For six of the 35 studied maternal plasma cases, the positive wells containing the mutant alleles constituted less than 0.3% of the total number of positive wells in the experiments. These positive signals might have resulted from cross hybridizations of the fluorescent probes during PCR. Nonetheless, such low numbers of mutant-positive wells would not skew the allelic ratio between mutant and wild-type alleles to an extent that would alter the RMD classification by SPRT.

10 Discussion

[0136] In this study, we have developed noninvasive prenatal diagnostic strategies to directly detect causative mutations carried by male fetuses in pregnancies at-risk of X-linked diseases, using hemophilia as an example. By using the digital RMD approach for genetic loci on chromosome X, we have accurately identified the mutant or the wild-type alleles inherited by the male fetuses in all of the 12 studied maternal plasma samples from seven pregnant carriers of hemophilia (table 1300). The fetal genotypes could be detected as early as the 11th week of gestation (table 900), demonstrating the potential for early diagnostic use of the method. The approach using a target region and a reference region on chromosome X can also be used.

20 [0137] This noninvasive prenatal mutation detection method could be combined with the existing noninvasive fetal sex determination test to further minimize the number of at-risk pregnant cases that would require invasive diagnostic testing. The identification of affected fetuses could also facilitate subsequent obstetric management for pregnant women who would not otherwise consider invasive prenatal testing. Three to four percent of infants with hemophilia experience a cranial bleed (Kulkarni R, Lusher JM., *J Pediatr Hematol Oncol.*, 21:289-295 (1999)) that occurs during labor and delivery. Prolonged labor and difficult instrumental deliveries are the main risk factors for this complication (Kadir RA et al., *Haemophilia.*, 6:33-40 (2000); Chi C et al., *Haemophilia.*, 14:56-64 (2008)) and should be avoided for delivery of affected fetuses (Lee CA, Chi C, Pavord SR, et al., *Haemophilia.*, 12:301-336 (2006)). It is also recommended that affected fetuses are delivered in a tertiary unit with an affiliated hemophilia center to ensure availability of necessary expertise and resources for their management (Lee CA, Chi C, Pavord SR, et al., *Haemophilia.*, 12:301-336 (2006)). Recently, prenatal diagnosis by third trimester amniocentesis has been suggested to

help appropriate planning of the mode and place of delivery for parents who are unwilling to accept the risk of fetal loss associated with earlier prenatal testing (Chi C, Kadir RA., *Obstetric Management. In: Lee CA, Kadir RA, Kouides PA, eds. Inherited Bleeding Disorders in Women*, Chichester, West Sussex, UK: Wiley-Blackwell, 122-148 (2009)). If a fetus is unaffected, labor and delivery can be managed without any restrictions in local maternity units. However, third trimester amniocentesis is also an invasive procedure and associated with potential risks and complications (Hodor JG, Poggi SH, Spong CY, et al., *Am J Perinatol.*, 23:177-180 (2006); O'Donoghue K et al., *Prenat Diagn.*, 27:1000-1004 (2007)). Fetal DNA concentration is the highest during the third trimester of pregnancy (Lun FMF et al., *Clin Chem.*, 54:1664-1672 (2008)), thus embodiments can offer an accurate noninvasive alternative to third trimester amniocentesis for this purpose.

VII. COMPUTER SYSTEM

[0138] Any of the computer systems mentioned herein may utilize any suitable number of subsystems. Examples of such subsystems are shown in FIG. 16 in computer apparatus 1600. In some embodiments, a computer system includes a single computer apparatus, where the subsystems can be the components of the computer apparatus. In other embodiments, a computer system can include multiple computer apparatuses, each being a subsystem, with internal components.

[0139] The subsystems shown in FIG. 16 are interconnected via a system bus 1675. Additional subsystems such as a printer 1674, keyboard 1678, fixed disk 1679, monitor 1676, which is coupled to display adapter 1682, and others are shown. Peripherals and input/output (I/O) devices, which couple to I/O controller 1671, can be connected to the computer system by any number of means known in the art, such as serial port 1677. For example, serial port 1677 or external interface 1681 can be used to connect computer system 1600 to a wide area network such as the Internet, a mouse input device, or a scanner. The interconnection via system bus 1675 allows the central processor 1673 to communicate with each subsystem and to control the execution of instructions from system memory 1672 or the fixed disk 1679, as well as the exchange of information between subsystems. The system memory 1672 and/or the fixed disk 1679 may embody a computer readable medium. Any of the values mentioned herein can be output from one component to another component and can be output to the user.

[0140] A computer system can include a plurality of the same components or subsystems, e.g., connected together by external interface 1681 or by an internal interface. In some

embodiments, computer systems, subsystem, or apparatuses can communicate over a network. In such instances, one computer can be considered a client and another computer a server, where each can be part of a same computer system. A client and a server can each include multiple systems, subsystems, or components.

5 [0141] It should be understood that any of the embodiments of the present invention can be implemented in the form of control logic using hardware and/or using computer software in a modular or integrated manner. Based on the disclosure and teachings provided herein, a person of ordinary skill in the art will know and appreciate other ways and/or methods to implement embodiments of the present invention using hardware and a combination of
10 hardware and software.

[0142] Any of the software components or functions described in this application may be implemented as software code to be executed by a processor using any suitable computer language such as, for example, Java, C++ or Perl using, for example, conventional or object-oriented techniques. The software code may be stored as a series of instructions or
15 commands on a computer readable medium for storage and/or transmission, suitable media include random access memory (RAM), a read only memory (ROM), a magnetic medium such as a hard-drive or a floppy disk, or an optical medium such as a compact disk (CD) or DVD (digital versatile disk), flash memory, and the like. The computer readable medium may be any combination of such storage or transmission devices.

20 [0143] Such programs may also be encoded and transmitted using carrier signals adapted for transmission via wired, optical, and/or wireless networks conforming to a variety of protocols, including the Internet. As such, a computer readable medium according to an embodiment of the present invention may be created using a data signal encoded with such programs. Computer readable media encoded with the program code may be packaged with
25 a compatible device or provided separately from other devices (e.g., via Internet download). Any such computer readable medium may reside on or within a single computer program product (e.g. a hard drive, a CD, or an entire computer system), and may be present on or within different computer program products within a system or network. A computer system may include a monitor, printer, or other suitable display for providing any of the results
30 mentioned herein to a user.

[0144] Any of the methods described herein may be totally or partially performed with a computer system including a processor, which can be configured to perform the steps. Thus, embodiments can be directed to computer systems configured to perform the steps of any of

the methods described herein, potentially with different components performing a respective steps or a respective group of steps. Although presented as numbered steps, steps of methods herein can be performed at a same time or in a different order. Additionally, portions of these steps may be used with portions of other steps from other methods. Also, all or portions of a
5 step may be optional. Additionally, any of the steps of any of the methods can be performed with modules, circuits, or other means for performing these steps.

[0145] The specific details of particular embodiments may be combined in any suitable manner without departing from the spirit and scope of embodiments of the invention. However, other embodiments of the invention may be directed to specific embodiments
10 relating to each individual aspect, or specific combinations of these individual aspects.

[0146] The above description of exemplary embodiments of the invention has been presented for the purposes of illustration and description. It is not intended to be exhaustive or to limit the invention to the precise form described, and many modifications and variations are possible in light of the teaching above. The embodiments were chosen and described in
15 order to best explain the principles of the invention and its practical applications to thereby enable others skilled in the art to best utilize the invention in various embodiments and with various modifications as are suited to the particular use contemplated.

[0147] A recitation of "a", "an" or "the" is intended to mean "one or more" unless specifically indicated to the contrary.

20 [0148] All patents, patent applications, publications, and descriptions mentioned above are herein incorporated by reference in their entirety for all purposes. None is admitted to be prior art.

WHAT IS CLAIMED IS:

- 1 1. A method for determining whether a male fetus of a pregnant female
2 has an X-linked mutation, wherein the pregnant female is heterozygous for a mutant and a
3 normal allele at a locus on the X chromosome, the method comprising:
4 receiving data from a plurality of reactions, each involving one or more
5 nucleic acid molecules from a biological sample, the biological sample including nucleic acid
6 molecules from the pregnant female and from the male fetus, wherein the data includes:
7 a first set of quantitative data indicating a first amount of the mutant
8 allele at the locus; and
9 a second set of quantitative data indicating a second amount of the
10 normal allele at the locus;
11 determining a parameter from the first amount and the second amount,
12 wherein the parameter represents a relative amount between the first and second amounts ;
13 obtaining a percentage Pf of fetal nucleic acid molecules in the biological
14 sample;
15 calculating a first cutoff value for determining whether the fetus has inherited
16 the mutant allele at the locus, wherein the first cutoff value is derived at least from a first
17 proportion of $k/(1+k-Pf)$, where k is a number of mutant alleles on a mutant chromosome of
18 the pregnant female, k being an integer equal to or greater than one;
19 calculating a second cutoff value for determining whether the fetus has
20 inherited the normal allele at the locus, wherein the second cutoff value is derived at least
21 from a second proportion of $[k(1-Pf)]/[1+k-kPf]$; and
22 comparing the parameter to at least one of the first and second cutoff values to
23 determine a classification of whether the fetus has inherited the mutant allele or the normal
24 allele.
- 1 2. The method of claim 1, wherein the parameter is compared to the first
2 and second cutoff values.
- 1 3. The method of claim 2 wherein the classifications include disease
2 state, non-disease state, and non-classifiable.
- 1 4. The method of any one of claims 1 to 3, wherein obtaining the
2 percentage Pf includes:

3 correcting an experimentally derived percentage of fetal nucleic acid
4 molecules in the biological sample with an expected statistical distribution of molecules in
5 the plurality of reactions.

1 5. The method of any one of claims 1 to 4, wherein obtaining the
2 percentage Pf includes:

3 detecting a first allele in the reactions, wherein the first allele is shared by the
4 mother and fetus at a locus where the pregnant female is homozygous and the fetus is either
5 heterozygous or hemizygous;

6 calculating a Poisson-corrected concentration P_x with the equation $[-\ln((N-$
7 $P1)/N)]*N$, where N is the total number of reactions analyzed, $P1$ is the number of reactions
8 positive for the first allele, and \ln is the natural logarithm;

9 detecting a second allele in the reactions, wherein the second allele is specific
10 to the fetus; and

11 calculating a Poisson-corrected concentration P_y with the equation $[-\ln((N-$
12 $P2)/N)]*N$, where N is the total number of reactions analyzed, and $P2$ is the number of
13 reactions positive for the second allele.

1 6. The method of claim 5, wherein the second allele is on chromosome Y.

1 7. The method of claim 5, wherein the first allele is on chromosome X.

1 8. The method of claim 5, wherein the fetal-specific allele is a paternally-
2 inherited allele on an autosome.

1 9. The method of claim 5, wherein the fetal-specific allele includes a
2 methylation marker specific to the fetus.

1 10. The method of claim 5, further comprising:
2 calculating Pf as $[(2P_y)/(P_x+P_y)]*100\%$.

1 11. The method of any one of claims 1 to 10, wherein the first and second
2 cutoff values are determined using a sequential probability ratio test (SPRT) to determine
3 whether the fetus has inherited the mutant or the normal nucleic acid sequence.

1 12. The method of any one of claims 1 to 11, wherein an allele at a
2 polymorphic site linked to the mutant nucleic acid sequence is an allele located on the same
3 maternal haplotype as the mutant nucleic acid sequence, and wherein the probability of

4 recombination between the polymorphic site and the mutant nucleic acid sequence is less
5 than 1%.

1 13. The method of any one of claims 1 to 11, wherein an allele at a
2 polymorphic site linked to the normal nucleic acid sequence refers is an allele located on the
3 same maternal haplotype as the normal nucleic acid sequence, and wherein the probability of
4 recombination between the polymorphic site and the mutant nucleic acid sequence is less
5 than 1%.

1 14. The method of any one of claims 1 to 13, wherein the reactions include
2 any one or more of the following: sequencing reactions, optical analysis, and hybridization
3 using a fluorescent probe, or nanopore sequencing.

1 15. The method of any one of claims 1 to 13, wherein a reaction is an
2 amplification reaction.

1 16. The method of claim 15, wherein the reactions include polymerase
2 chain reactions.

1 17. The method of claim 17, wherein the average concentration is less than
2 one template molecule per reaction, and wherein a Poisson distribution is used in determining
3 the percentage Pf of fetal nucleic acid molecules in the biological sample.

1 18. The method of any one of claims 1 to 17 wherein the biological sample
2 is plasma, serum, or whole blood from a pregnant woman.

1 19. A method for determining whether a male fetus of a pregnant female
2 has an X-linked mutation, the method comprising:

3 receiving data from a plurality of reactions, each involving one or more
4 nucleic acid molecules from a biological sample, the biological sample including nucleic acid
5 molecules from the pregnant female and from the male fetus,

6 wherein the pregnant female is homozygous for an allele at a locus on the X
7 chromosome, has a mutation of an amplification of the allele on a mutant X chromosome, the
8 mutant X chromosome having a normal copy of the allele at the locus and one or more
9 additional copies of the allele, and has a normal X chromosome having a normal copy of the
10 allele at the locus,

11 wherein the data includes:

12 a first set of quantitative data indicating a first amount of an additional
13 junction created by the one or more additional copies of the allele; and
14 a second set of quantitative data indicating a second amount of a
15 normal junction created by the normal copy of the allele on both X chromosomes;
16 determining a parameter from the first amount and the second amount,
17 wherein the parameter represents a relative amount between the first and second amounts ;
18 obtaining a percentage Pf of fetal nucleic acid molecules in the biological
19 sample;
20 calculating a first cutoff value for determining whether the fetus has inherited
21 the mutant X chromosome, wherein the first cutoff value is derived at least from a first
22 proportion of $n/(n+1-Pf)$, where n is the number of additional copies of the allele, n being an
23 integer equal to or greater than one;
24 calculating a second cutoff value for determining whether the fetus has
25 inherited the normal X chromosome, wherein the second cutoff value is derived at least from
26 a second proportion of $[n(1-Pf)/[n+2-Pf(n+1)]]$; and
27 comparing the parameter to at least one of the first and second cutoff values to
28 determine a classification of whether the fetus has inherited the mutant X chromosome or the
29 normal X chromosome.

1 20. A method for determining whether a male fetus of a pregnant female
2 has an X-linked mutation, wherein the pregnant female is heterozygous for a mutation and a
3 normal allele at a target region on the X chromosome, wherein the mutation is a deletion or
4 an amplification of the target region, the method comprising:
5 receiving data from a plurality of reactions, each involving one or more
6 nucleic acid molecules from a biological sample, the biological sample including nucleic acid
7 molecules from the pregnant female and from the male fetus, wherein the data includes:
8 a first set of quantitative data indicating a first amount of the nucleic
9 acid molecules that are from the target region; and
10 a second set of quantitative data indicating a second amount of the
11 nucleic acid molecules that are from a reference region on the X chromosome;
12 determining a parameter from the first amount and the second amount,
13 wherein the parameter represents a relative amount between the first and second amounts;
14 obtaining a percentage Pf of fetal nucleic acid molecules in the biological
15 sample;

16 calculating a first cutoff value for determining whether the fetus has inherited
17 the mutation, the first cutoff value being dependent on the percentage Pf;
18 calculating a second cutoff value for determining whether the fetus has
19 inherited the normal allele, the second cutoff value being dependent on the percentage Pf; and
20 comparing the parameter to at least one of the first and second cutoff values to
21 determine a classification of whether the fetus has inherited the mutation or the normal allele.

1 21. The method of claim 20, wherein the mutation is an amplification,
2 wherein the first cutoff value is determined based on the assumption that a ratio of the first
3 amount to the second amount is increased when compared with a corresponding ratio of a
4 non-pregnant woman carrying the same amplification mutation, and the second cutoff value
5 is based on the assumption that the ratio of the first amount to second amount is decreased
6 when compared with the corresponding ratio of a non-pregnant woman carrying the same
7 amplification mutation.

1 22. The method of claim 20, wherein the mutation is a deletion, wherein
2 the second cutoff value is determined based on the assumption that a ratio of the first amount
3 to the second amount is increased when compared with a corresponding ratio of a non-
4 pregnant woman carrying the same deletion mutation, and the first cutoff value is based on
5 the assumption that the ratio of the first amount to the second amount is decreased when
6 compared with the corresponding ratio of a non-pregnant woman carrying the same deletion
7 mutation.

1 23. The method of claim 20, wherein the mutation is a deletion, wherein
2 the second cutoff value is derived at least from a first proportion of $1/(2-Pf)$, and wherein the
3 first cutoff value is derived at least from a second proportion of $(1-Pf)/(2-Pf)$.

1 24. The method of claim 20, wherein the mutation is a duplication,
2 wherein the second cutoff value is derived at least from a first proportion of $(3-Pf)/(2-Pf)$, and
3 wherein the first cutoff value is derived at least from a second proportion of $(3-2Pf)/(2-Pf)$.

1 25. The method of any one of claims 20 to 24, wherein obtaining the
2 percentage Pf includes:
3 correcting an experimentally derived percentage of fetal nucleic acid
4 molecules in the biological sample with an expected statistical distribution of molecules in
5 the plurality of reactions.

1 26. A method of obtaining a percentage P_f of fetal nucleic acid molecules
2 in a biological sample from a female pregnant with a fetus, the method comprising:
3 receiving data from a plurality of reactions, each involving a plurality of
4 nucleic acid molecules from a biological sample, the biological sample including nucleic acid
5 molecules from the pregnant female and from the fetus;
6 detecting a first allele in the reactions, wherein the first allele is shared by the
7 mother and fetus at a locus where the pregnant female is homozygous and the fetus is either
8 heterozygous or hemizygous;
9 calculating a corrected concentration P_x of the first allele based on a number
10 of reactions positive for the first allele, where P_x is corrected for an expected statistical
11 distribution of the first allele in the plurality of reactions;
12 detecting a second allele in the reactions, wherein the second allele is specific
13 to the fetus;
14 calculating a corrected concentration P_y of the second allele based on a
15 number of reactions positive for the second allele, where P_y is corrected for an expected
16 statistical distribution of the second allele in the plurality of reactions; and
17 calculating P_f using $[(2P_y)/(P_x+P_y)]$.

1 27. The method of claim 26, wherein P_f equals $[(2P_y)/(P_x+P_y)] * 100\%$.

1 28. The method of any one of claims 26 and 27, wherein the statistical
2 distribution is Poisson, and wherein the Poisson-corrected concentration P_x uses the equation
3 $[-\ln((N-P_1)/N)] * N$, where N is the total number of reactions analyzed, P_1 is the number of
4 wells positive for the first allele, and \ln is the natural logarithm, and wherein the Poisson-
5 corrected concentration P_y uses the equation $[-\ln((N-P_2)/N)] * N$, where N is the total number
6 of reactions analyzed, and P_2 is the number of wells positive for the second allele.

1 29. The method of any one of claims 26 to 28, wherein the data includes:
2 a first set of quantitative data indicating a first amount of the mutant nucleic
3 acid sequence or an allele at a polymorphic site linked to the mutant nucleic acid sequence;
4 and
5 a second set of quantitative data indicating a second amount of the normal
6 nucleic acid sequence or an allele at a polymorphic site linked to the normal nucleic acid
7 sequence, the method further comprising:
8 determining a parameter from the two data sets;

9 determining a first cutoff value for determining whether the fetus has inherited
10 a mutant nucleic acid sequence, wherein the first cutoff value is determined based on the
11 percentage Pf;
12 determining a second cutoff value for determining whether the fetus has
13 inherited the normal nucleic acid sequence, wherein the second cutoff value is determined
14 based on the percentage Pf;
15 comparing the parameter to at least one of the first and second cutoff values;
16 and
17 based on the comparison, determining a classification of whether the fetus has
18 inherited the mutant or the normal nucleic acid sequence.

1 30. A computer program product comprising a non-transitory computer
2 readable medium storing a plurality of instructions for controlling a processor to perform an
3 operation, the instructions comprising any of the above methods.

1 31. A system comprising:
2 one or more processors configured to perform any of the above methods.

1 32. A system comprising modules configured to perform any of the above
2 methods.




FIG. 1





FIG. 2

300







FIG. 3

FIG. 4

DNA extracted from the plasma of a high-risk pregnant woman




500

Disease type	Maternal genotype	Fetal genotype	Maternal plasma DNA						M:N ratio
			Maternal-derived alleles		Fetal-derived alleles		(maternal + fetal) alleles		
			M (copies)	N (copies)	M (copies)	N (copies)	M (copies)	N (copies)	
X-linked diseases			Total DNA alleles in maternal plasma = 100 GE ¹ Fetal DNA percentage = 10% ¹						0.90 : 1
			Total maternal alleles = 90 GE		Total fetal alleles = 10 GE				
			M (copies)	N (copies)	M (copies)	N (copies)	M (copies)	N (copies)	
			90 ²	90 ²	0 ³	10 ³	90	100	0.90 : 1
			90	90	10	0	100	90	1.11 : 1

¹ To illustrate the calculation, a maternal plasma sample containing a total of 100GE of DNA with 10% fetal DNA was used.
² For the maternal genome, 1GE contains 2 copies of the alleles, i.e., one copy each of the M and the N allele.
³ For the fetal genome, 1GE contains 1 copy of the X-linked allele, i.e., one copy of either the M or the N allele.
 GE. Genome-equivalent; M, mutant allele; N, wild-type allele.

FIG. 5A



Normal Chromosome

Mutant Chromosome

Dosage comparison: [hatched box] vs [striped box]




FIG. 5B

FIG. 5C

7/17

600




FIG. 6

		Maternal plasma DNA				FIG. 7	
		Total DNA alleles in maternal plasma = 100 GE ¹ Fetal DNA proportion = 10% ¹					
Mutation type	Maternal genotype	Fetal genotype	Maternal-derived alleles ² Total maternal alleles = 90 GE		Fetal-derived alleles ³ Total fetal alleles = 10 GE		T : ZFX
			T (copies)	ZFX (copies)	T (copies)	ZFX (copies)	
Deletion			90	180	10	10	0.53 : 1
			90	180	0	10	
Duplication			270	180	10	10	1.47 : 1
			270	180	20	10	

¹ To illustrate the calculation, a maternal plasma sample containing a total of 100GE of DNA with 10% fetal DNA was used.
² For the maternal genome, 1GE contains 2 copies of ZFX, as well as 1 copy of the target allele in the deleted region, or 3 copies of the target allele in the duplicated region.
³ For the fetal genome, 1GE contains 1 copy of ZFX, and 1 copy of the target allele for the normal fetus. For fetus inherits the deleted or the duplicated locus, 1 GE corresponds to 0 and two copies of the target allele, respectively.
 GE, genome-equivalent; T, the target molecules.

9/17

800




FIG. 8

10/17

900

Sample	Gestation (weeks)	Affected gene	Mutation ¹	Severity of hemophilia	Fetal information	
					Sex	Hemophilia status
H9	36	<i>F9</i>	c.874delC (p.Gln292Lysfs)	Severe	Male	Affected
H12a ²	18	<i>F8</i>	c.6278A>G (p.Asp2093Gly)	Mild	Male	Affected
H12b	34					
H15a ²	34	<i>F9</i>	c.1144T>C (p.Cys382Arg)	Severe	Male	Affected
H15b	38					
H17	28	<i>F8</i>	c.826G>A (p.Val276Met)	Mild	Male	Unaffected
H25a ²	23	<i>F9</i>	c.802T>A (p.Cys268Ser)	Moderate/ Severe	Male	Affected
H25b	32					
H26a ²	11	<i>F8</i>	c.1171C>T (p.Arg391Cys)	Severe	Male	Unaffected
H26b	23					
H30a ²	32	<i>F9</i>	c.1069G>A (p.Gly357Arg)	Moderate/ Severe	Male	Unaffected
H30b	40					

¹ Mutation nomenclature is based on the guidelines of the Human Genome Variation Society. The reference sequences for hemophilia A and B mutations are *F8* mRNA variant 1 (Genbank accession NM_000132.3) and *F9* mRNA (NM_000133.3), respectively. Nucleotide position +1 corresponds to the A nucleotide of the ATG translation initiation codon. Amino acid changes deduced from the DNA mutations are shown in the brackets. The reference sequences correspond to coagulation factor VIII isoform a precursor (NP_000123.1) and coagulation factor IX preproprotein (NP_000124.1), respectively. The translation initiator methionine is numbered as position +1.

²Peripheral blood samples were taken on two occasions from the same women during their pregnancies.

FIG. 9

1000

Locus		Sequence	Final concentration, nM	Annealing temperature
ZFYX	F-primer	5'-CAAGTGCTGGACTCAGATGTAAGT-3'	900	57°C
	R-primer	5'-TGAAGTAATGTCAGAAGCTAAAACATCA-3'	900	
	X-probe	5'-(VIC)TCTTTAGCACATTGCA(MGBNFQ)-3'	100	
	Y-probe	5'-(FAM)TCTTTACCACACTGCAC(MGBNFQ)-3'	100	
Rs6528633	F-primer	5'-GGAAGACCAAAAAGGGATAAAGG-3'	900	57°C
	R-primer	5'-CACCTACTCCCAGCCAATTT-3'	900	
	T-probe	5'-(VIC)TGAGATATGATATGGTCATG(MGBNFQ)-3'	200	
	A-probe	5'-(FAM)TGAGATATGATAAGGTCATG(MGBNFQ)-3'	179	
F8 c.826G>A	F-primer	5'-TGGATGCCACAGGAAATCAG-3'	900	58°C
	R-primer	5'-CTTCAGGAGTGGTGCCATT-3'	900	
	G-probe	5'-(VIC)CTATTGGCATGTGATTG(MGBNFQ)-3'	200	
	A-probe	5'-(FAM)CTATTGGCATATGATTG(MGBNFQ)-3'	179	
F8 c.1171C>T	F-primer	5'-TGGATGTGGTCAGGTTTGTATGA-3'	900	58°C
	R-primer	5'-TTTTAGGATGCTTCTTGGCAACT-3'	900	
	C-probe	5'-(FAM)CTGAGCGAATTTGGATA(MGBNFQ)-3'	150	
	T-probe	5'-(VIC)CTGAGCAAATTTGGAT(MGBNFQ)-3'	200	
F8 c.6278A>G	F-primer	5'-TTTCAGGAGGTAGCACATACATTT-3'	900	56°C
	R-primer	5'-TGCCGTGAATAATCATTGGT-3'	900	
	A-probe	5'-(VIC)CAACAGATCCACCTAC(MGBNFQ)-3'	300	
	G-probe	5'-(FAM)AACAGACCCACCTAC(MGBNFQ)-3'	269	
F9 c.802T>A	F-primer	5'-TCTGTGGAGGCTCTATCGTTAATG-3'	900	58°C
	R-primer	5'-ACCTGCGACAACGTGAATTTTAACAC-3'	900	
	T-probe	5'-(VIC)TGCCCACTGTGTTGA(MGBNFQ)-3'	220	
	A-probe	5'-(FAM)CTGCCACAGTGTGTTG(MGBNFQ)-3'	100	
F9 c.874delC	F-primer	5'-TGTCGCAGGTGAACATAATATTGA-3'	900	58°C
	R-primer	5'-GGTGAGGAATAATTCGAATCACATT-3'	900	
	C-probe	5'-(VIC)ACATACAGAGCAAAAG(MGBNFQ)-3'	120	
	del-probe	5'-(FAM)CATACAGAGAAAAGC(MGBNFQ)-3'	120	
F9 c.1069G>A	F-primer	5'-CCTCAAATTTGGATCTGGCTA-3'	900	58°C
	R-primer	5'-GCTGATCTCCCTTTGTGGAA-3'	900	
	G-probe	5'-(VIC)ACTCTTCCCAGCCAC(MGBNFQ)-3'	179	
	A-probe	5'-(FAM)ACTCTTCTCCAGCCACT(MGBNFQ)-3'	200	
F9 c.1144T>C	F-primer	5'-CAGTACCTTAGAGTTCCACTTGTTGAC-3'	900	58°C
	R-primer	3'	900	
	T-probe	5'-(VIC)CCACATGTCTTCG(MGBNFQ)-3'	200	
	C-probe	5'-(FAM)AGCCACACGTCTTC(MGBNFQ)-3'	130	

VIC and FAM, fluorescent reporter dyes; MGBNFQ, minor groove binding nonfluorescent quencher.

FIG. 10

12/17

1100

Sample	Gestation (weeks)	SNP genotype ¹		Digital PCR result			Fetal % ³	m_r ⁴	SPRT Classification ⁵
		Mother	Fetus	Total wells	A count ²	T count ²			
M5193P	17 3/7	AT	A	4590	410	335	7.5	0.09	A
M5280P	17 4/7	AT	A	4590	204	144	10.7	0.05	A
M5269P	19 2/7	AT	T	4590	337	409	11.2	0.08	T
M5297P	19 3/7	AT	A	13770 ⁶	1482	1368	5.4	0.11	A
M5244P	22 6/7	AT	T	4590	139	163	8.5	0.03	T
M5240P	38 2/7	AT	T	4590	396	484	23.7	0.09	T
M5241P	38 2/7	AT	T	4590	706	920	13.9	0.17	T
M4817P	38 4/7	AT	A	4590	502	425	17.2	0.12	A
M4847P	39 3/7	AT	A	4590	379	312	14.9	0.09	A
M4846P	40	AT	A	4590	700	516	17.5	0.17	A

¹SNP genotypes were determined by mass spectrometry.

²A count, number of wells positive for the A allele. T count, number of wells positive for the T allele.

³Fetal DNA percentages were determined by the digital ZFY/X assay.

⁴Average reference template concentration per PCR well. The reference template referred to the allele with the lesser count in each sample.

⁵Classification of fetal genotypes by SPRT.

⁶M5297P was unclassifiable by SPRT with data from the first 4590-well digital PCR set. Two additional 4590-well digital PCR sets were performed after which classification could be made.

FIG. 11

13/17

1200

Mutation	Fetal DNA percentage ¹	Digital PCR result			m_r ³	SPRT Classification ⁴
		Total wells	Mutant count ²	Wild-type count ²		
F8	20%	4590	1495	1721	0.47	Wild-type
c.826G>A	10%	4590	2917	3006	1.06	Wild-type
F8	20%	4590	406	507	0.12	Wild-type
c.1171C>T	10%	4590	435	483	0.11	Wild-type
F8	20%	4590	485	593	0.14	Wild-type
c.6278A>G	10%	4590	857	892	0.22	Wild-type
F9	20%	4590	88	109	0.02	Wild-type
c.30117T>A	10%	4590	1640	1740	0.48	Wild-type
F9	20%	4590	375	463	0.11	Wild-type
c.874delC	10%	4590	388	438	0.10	Wild-type
F9	20%	4590	308	365	0.05	Wild-type
c.1144T>C	10%	4590	380	408	0.04	Wild-type
F9	20%	4590	673	835	0.20	Wild-type
c.1069G>A	10%	4590	966	1046	0.26	Wild-type

¹Fetal DNA was mixed in the specified percentages with maternal DNA. Fetal DNA was obtained from the placenta of a normal male fetus. Maternal DNA was obtained from the blood cells of pregnant women heterozygous for the corresponding mutations.

²Mutant count, number of wells positive for the mutant allele. Wild-type count, number of wells positive for the wild-type allele.

³Average reference template concentration per PCR well. The reference template referred to the allele with the lesser count in each sample.

⁴SPRT classification of "fetal genotypes", which was mimicked by the normal placental DNA in the artificial mixtures.

FIG. 12

14/17

1300

Plasma sample	Fetal hemophilia status	Digital PCR result			Fetal % ²	m_r ³	SPRT Classification ⁴
		Total wells	Mutant count ¹	Wild-type count ¹			
H9	Affected	4590	1022	801	14.8	0.19	Mutant
H26a	Unaffected	9180	1590	1710	3.8	0.21	Wild-type
H26b	Unaffected	4590	590	650	6.8	0.15	Wild-type
H15a	Affected	4590	573	435	10.5	0.10	Mutant
H15b	Affected	4590	2506	1956	10.7	0.56	Mutant
H17	Unaffected	4590	329	342	14.0	0.08	Wild-type
H30a	Unaffected	4590	611	780	18.2	0.19	Wild-type
H30b	Unaffected	4590	1839	2017	11.4	0.58	Wild-type
H25a	Affected	9180	1160	1108	4.6	0.13	Mutant
H25b	Affected	4590	223	166	15.0	0.04	Mutant
H12a	Affected	9180	511	464	9.0	0.05	Mutant
H12b	Affected	4590	293	230	25.1	0.05	Mutant

¹Mutant count, number of wells positive for the mutant allele. Wild-type count, number of wells positive for the wild-type allele.

²Fetal DNA percentages were determined by the digital ZFYX assay.

³Average reference template concentration per PCR well. The reference template referred to the allele with the lesser count in each sample.

⁴Classification of fetal genotypes by SPRT.

FIG. 13




FIG. 14

16/17

1500

Mutation	Plasma sample	Gestation (weeks)	Digital PCR result			Fetal % ³
			Total wells	Mutant count ^{1,2}	Wild-type count ¹	
<i>F8</i> c.826G>A	N1P	26	4590	0	589	12.8
	N10P	36	4590	0	1541	17.9
	N11P	28	4590	0	587	10.0
	N17P	39	4590	0	2476	16.9
	N21P	30	4590	0	763	10.3
<i>F8</i> c.1171C>T	N1P	26	4590	0	575	12.8
	N10P	36	4590	0	1346	17.9
	N11P	28	4590	0	583	10.0
	N17P	39	4590	1	2202	16.9
	N21P	30	4590	0	640	10.3
<i>F8</i> c.6278A>G	N2P	37	4590	0	1584	14.0
	N3P	35	4590	0	737	18.3
	N11P	28	4590	0	406	10.0
	N13P	42	4590	0	587	29.8
	N19P	36	4590	0	623	18.6
<i>F9</i> c.802T>A	N1P	26	3825	0	261	12.8
	N10P	36	4590	0	814	17.9
	N11P	28	4590	0	410	10.0
	N17P	39	4590	0	1406	16.9
	N21P	30	4590	0	549	10.3
<i>F9</i> c.874delC	N1P	26	4590	0	457	12.8
	N10P	36	4590	0	1244	17.9
	N11P	28	4590	0	551	10.0
	N17P	39	4590	0	1767	16.9
	N21P	30	4590	0	606	10.3
<i>F9</i> c.1069G>A	N2P	37	4590	0	2124	14.0
	N3P	35	4590	0	990	18.3
	N11P	28	4590	0	483	10.0
	N17P	39	4590	1	2112	16.9
	N21P	30	4590	1	708	10.3
<i>F9</i> c.1144T>C	N1P	26	4590	1	613	12.8
	N10P	36	4590	0	1132	17.9
	N11P	28	4590	1	602	10.0
	N17P	39	4590	0	1808	16.9
	N21P	30	4590	2	597	10.3

¹Mutant count, number of wells positive for the mutant allele. Wild-type count, number of wells positive for the wild-type allele.

²Some of the cases showed 1 or 2 wells positive for the mutant alleles. This might be due to cross hybridizations of the mutant fluorescent probes against the wild-type amplicons.

³Fetal DNA percentages were determined by the digital ZFY/X assay.

FIG. 15




FIG. 16

1675

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2012/000015

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl.

C12Q 1/68 (2006.01)

G06F 19/00 (2011.01)

Action Date: 26 April 2012

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 MEDLINE, CAPLUS, BIOSIS, BIOTECHABS, WPI, EPODOC (pregnant, impregnated, prenatal, maternal, antenatal, fetus, X-linked, X chromosome, sex-linked, classified, quantitative, percentage, ratio, statistical, calculate, nucleic acid, DNA, RNA, mutation, allele, loci, polymorphism, genotype, heterozygous, hemizygous, homozygous, cell-free, extra-cellular, circulating, plasma, serum, relative mutation dosage, hemophilia, muscular dystrophy; and other synonyms and like terms)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LUN, F. M. F. et al., "Noninvasive prenatal diagnosis of monogenic diseases by digital size selection and relative mutation dosage on DNA in maternal plasma", Proceedings of the National Academy of Sciences of the USA, 16 December 2008, Vol. 105, No. 50, pages 19920-19925 See whole document especially Abstract	
A	CHIU, R. W. K. and LO, Y. M. D., "Application of fetal DNA in maternal plasma for noninvasive prenatal diagnosis", Expert Review of Molecular Diagnostics, January 2002, Vol. 2, No. 1, pages 32-40 See whole document	

 Further documents are listed in the continuation of Box C See patent family annex

* Special categories of cited documents:		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search
26 April 2012Date of mailing of the international search report
27 April 2012Name and mailing address of the ISA/AU
AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
E-mail address: pct@ipaaustralia.gov.au
Facsimile No. +61 2 6283 7999Authorized officer
DAMIAN TRIFFETT
AUSTRALIAN PATENT OFFICE
(ISO 9001 Quality Certified Service)
Telephone No : +61 2 6283 2845

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2012/000015

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	TSUI, N. B. Y. et al., "Noninvasive prenatal diagnosis of hemophilia by microfluidics digital PCR analysis of maternal plasma DNA", Blood, 31 March 2011, Vol. 117, No. 13, pages 3684-3691 See whole document especially Abstract	1-32