

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199880170 B2
(10) Patent No. 739192

(54) Title
Fungicidal mixtures

(51)⁶ International Patent Classification(s)
A01N 047/24 A01N 037/50
A01N 037/06

(21) Application No: **199880170** (22) Application Date: **1998.05.13**

(87) WIPO No: **WO98/52417**

(30) Priority Data

(31) Number	(32) Date	(33) Country
19721402	1997.05.22	DE
19721401	1997.05.22	DE

(43) Publication Date : **1998.12.11**

(43) Publication Journal Date : **1999.02.04**

(44) Accepted Journal Date : **2001.10.04**

(71) Applicant(s)
BASF Aktiengesellschaft

(72) Inventor(s)
Klaus Schelberger; Maria Scherer; Hubert Sauter; Bernd Muller; Erich Birner; Joachim Leyendecker ; Eberhard Ammermann; Gisela Lorenz; Siegfried Strathmann; Reinhold Saur

(74) Agent/Attorney
WATERMARK PATENT and TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN VIC 3122

(56) Related Art
WO 97/40688
WO 98/01033
WO 96/01258

OPI DATE 11/12/98 APPLN. ID 80170/98
 AOJP DATE 04/02/99 PCT NUMBER PCT/EP98/02820



AU9880170


<p>(51) Internationale Patentklassifikation ⁶ : A01N 47/24, 37/50, 37/06 // (A01N 47/24, 37:50, 37:06) (A01N 37/50, 47:24, 37:06)</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 98/52417 (43) Internationales Veröffentlichungsdatum: 26. November 1998 (26.11.98)</p>
<p>(21) Internationales Aktenzeichen: PCT/EP98/02820 (22) Internationales Anmeldedatum: 13. Mai 1998 (13.05.98) (30) Prioritätsdaten: 197 21 402.9 22. Mai 1997 (22.05.97) DE 197 21 401.0 22. Mai 1997 (22.05.97) DE (71) Anmelder (für alle Bestimmungsstaaten ausser US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE). (72) Erfinder; und (75) Erfinder/Anmelder (nur für US): SCHELBERGER, Klaus [AT/DE]; Traminerweg 2, D-67161 Gönheim (DE). SCHERER, Maria [DE/DE]; Hermann-Jürgens-Strasse 30, D-76829 Landau (DE). SAUTER, Hubert [DE/DE]; Neckarpromenade 20, D-68167 Mannheim (DE). MÜLLER, Bernd [DE/DE]; Jean-Ganss-Strasse 21, D-67227 Frankenthal (DE). BIRNER, Erich [DE/DE]; Hauptstrasse 78, D-67317 Altleiningen (DE). LEYEN-DECKER, Joachim [DE/DE]; Stahlbühlring 79, D-68526 Ladenburg (DE). AMMERMANN, Eberhard [DE/DE]; Von-Gagem-Strasse 2, D-64646 Heppenheim (DE). LORENZ, Gisela [DE/DE]; Erlenweg 13, D-67434 Neustadt (DE). STRATHMANN, Siegfried [DE/DE]; Don-</p>	<p>nersbergstrasse 9, D-67117 Limburgerhof (DE). SAUR, Reinhold [DE/DE]; Königsberger Strasse 9, D-67459 Böhl-Iggelheim (DE). (74) Gemeinsamer Vertreter: BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE). (81) Bestimmungsstaaten: AL, AU, BG, BR, BY, CA, CN, CZ, GE, HU, ID, IL, JP, KR, KZ, LT, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, curasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Veröffentlicht Mit internationalem Recherchenbericht.</p>	

(54) Title: FUNGICIDAL MIXTURES

(54) Bezeichnung: FUNGIZIDE MISCHUNGEN


(57) Abstract

The invention relates to a fungicidal mixture containing a synergistically active quantity of a) a phenyl-benzyl ether derivative of formula (I.a), (I.b) or (I.c) or I.c, and/or a₂ a carbamate of formula (I.d), in which X is CH and N, n is 0, 1 or 2 and R is a halogen, C₁-C₄ alkyl and C₁-C₄ halogen alkane, whereby the R radicals can be different if n equals 2, and one of its salts or adducts, as well as b) a dinitrophenol derivative of formula (II.a) or (II.b), in which n is 0, 1 or 2.



(57) Zusammenfassung

Fungizide Mischung, enthaltend a₁) ein Phenyl-benzyletherderivat der Formel (I.a), (I.b) oder (I.c), und/oder a₂) ein Carbamat der Formel (I.d), in der X CH und N bedeutet, n für 0, 1 oder 2 steht und R Halogen, C₁-C₄-Alkyl und C₁-C₄-Halogenalkyl bedeutet, wobei die Reste R verschieden sein können, wenn n für 2 steht, eines seiner Salze oder Addukte, sowie b) ein Dinitrophenolderivat der Formel (II.a) oder (II.b), in der n für 0, 1 oder 2 steht, in einer synergistisch wirksamen Menge.





Fungicidal mixtures

5 Abstract

A fungicidal mixture comprises

- a₁) a phenyl benzyl ether derivative of the formula I.a, I.b or I.c


10



and/or

- 25 a₂) a carbamate of the formula Id

30



where X is CH or N, n is 0, 1 or 2 and R is halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl, it being possible for the radicals R to be different if n is 2, or one of its salts or adducts,



35

and

- 40 b) a dinitrophenol derivative of the formula II.a or II.b

45





10 where n is 0, 1 or 2
 in a synergistically effective amount.

15

20

25

30

35

40

45




Fungicidal mixtures

The present invention relates to a fungicidal mixture which comprises


5

- a₁) a phenyl benzyl ether derivative of the formula I.a, I.b or I.c



and/or


- a₂) a carbamate of the formula Id



where X is CH or N, n is 0, 1 or 2 and R is halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl, it being possible for the radicals R to be different if n is 2, or one of its salts or adducts, and [sic]

35 and

- b) a dinitrophenol derivative comprising compounds of the formula II.a and/or II.b



where n is 0, 1 or 2



in a synergistically effective amount.

The invention further relates to methods for controlling harmful fungi using mixtures of the compounds I (I.a, I.b or I.c) and II 5 (II.a or II.b) and to the use of the compound I and the compound II for preparing such mixtures.

The compounds of the formula Ia-c, their preparation and their activity against harmful fungi are known from the literature 10 (EP-A 253 213; EP-A 254 426; EP-A 398 692).

The compounds of the formula Id, their preparation and their activity are known from WO-A 93/15046, WO-A 96/01256 and WO-A 96/01258.

15

Also known is the mixture of the compounds II (II.a and II.b) (common name: Dinocap), their preparation and their activity against harmful fungi and arachnids (US-A 2,526,660; US-A 2,810,767).

20

It is an object of the present invention to provide mixtures which have an improved activity against harmful fungi combined with a reduced total amount of active ingredients applied (synergistic mixtures), with a view to reducing the application 25 rates and to improving the activity spectrum of the known compounds I and II.

We have found that this object is achieved by the mixture defined at the outset. In addition, we have found that, by 30 applying the compound I and the compound II simultaneously, that is to say separately as well as together, or by applying the compound I and the compounds II in succession, better control of harmful fungi is possible than when the individual compounds are used.

35


The formula Id in particular represents carbamates where the combination of the substituents corresponds to a row of the table below:

40

45



3



(I)

5

No.	X	R _n
I.1	N	2-F
I.2	N	3-F
10 I.3	N	4-F
I.4	N	2-Cl
I.5	N	3-Cl
I.6	N	4-Cl
15 I.7	N	2-Br
I.8	N	3-Br
I.9	N	4-Br
I.10	N	2-CH ₃
I.11	N	3-CH ₃
20 I.12	N	4-CH ₃
I.13	N	2-CH ₂ CH ₃
I.14	N	3-CH ₂ CH ₃
I.15	N	4-CH ₂ CH ₃
I.16	N	2-CH(CH ₃) ₂
25 I.17	N	3-CH(CH ₃) ₂
I.18	N	4-CH(CH ₃) ₂
I.19	N	2-CF ₃
I.20	N	3-CF ₃
I.21	N	4-CF ₃
30 I.22	N	2,4-F ₂
I.23	N	2,4-Cl ₂
I.24	N	3,4-Cl ₂
I.25	N	2-Cl, 4-CH ₃
I.26	N	3-Cl, 4-CH ₃
35 I.27	CH	2-F
I.28	CH	3-F
I.29	CH	4-F
I.30	CH	2-Cl
I.31	CH	3-Cl
40 I.32	CH	4-Cl
I.33	CH	2-Br
I.34	CH	3-Br
I.35	CH	4-Br
I.36	CH	2-CH ₃
45 I.37	CH	3-CH ₃
I.38	CH	4-CH ₃
I.39	CH	2-CH ₂ CH ₃
I.40	CH	3-CH ₂ CH ₃



No.	X	R _n
I.41	CH	4-CH ₂ CH ₃
I.42	CH	2-CH(CH ₃) ₂
I.43	CH	3-CH(CH ₃) ₂
I.44	CH	4-CH(CH ₃) ₂
I.45	CH	2-CF ₃
I.46	CH	3-CF ₃
I.47	CH	4-CF ₃
I.48	CH	2,4-F ₂
I.49	CH	2,4-Cl ₂
I.50	CH	3,4-Cl ₂
I.51	CH	2-Cl, 4-CH ₃
I.52	CH	3-Cl, 4-CH ₃

15 Particular preference is given to the compounds I.12, I.23, I.32 and I.38.

20 When preparing the mixtures, it is preferred to employ the pure active ingredients I and II, to which further active ingredients against harmful fungi or other pests, such as insects, arachnids or nematodes, or else herbicidal or growth-regulating active ingredients or fertilizers can be admixed.

25 The mixtures of the compounds I and II, or the simultaneous joint or separate use of the compounds I and II, have outstanding action against a wide range of phytopathogenic fungi, in particular from the classes of the Ascomycetes, Basidiomycetes, Phycomycetes and Deuteromycetes. Some of them act systemically and can therefore also be employed as foliar and soil-acting fungicides.

35 They are especially important for controlling a large number of fungi in a variety of crop plants, such as cotton, vegetable species (eg. cucumbers, beans, tomatoes, potatoes and cucurbits), barley, grass, oats, bananas, coffee, maize, fruit species, rice, rye, soya, grapevine, wheat, ornamentals, sugar cane, and a variety of seeds.

40 They are particularly suitable for controlling the following phytopathogenic fungi: Erysiphe graminis (powdery mildew) in cereals, Erysiphe cichoracearum and Sphaerotheca fuliginea in cucurbits, Podosphaera leucotricha in apples, Uncinula necator in grapevines, Puccinia species in cereals, Rhizoctonia species in cotton, rice and lawns, Ustilago species in cereals and sugar cane, Venturia inaequalis (scab) in apples, Helminthosporium species in cereals, Septoria nodorum in wheat, Botrytis cinerea (gray mold) in strawberries, vegetables, ornamentals and



grapevines, *Cercospora arachidicola* in groundnuts, *Pseudocercospora herpotrichoides* in wheat and barley, *Pyricularia oryzae* in rice, *Phytophthora infestans* in potatoes and tomatoes, *Plasmopara viticola* in grapevines
 5 *Pseudocercospora* species in hops and cucumbers, *Alternaria* species in vegetables and fruit, *Mycosphaerella* species in bananas, and *Fusarium* and *Verticillium* species.

Furthermore, they can be used in the protection of materials
 10 (eg. in the protection of wood), for example against *Paecilomyces variotii*.

The compounds I and II can be applied simultaneously, that is either together or separately, or in succession, the sequence,
 15 in the case of separate application, generally not having any effect on the result of the control measures.

The compounds I and II are usually in a weight ratio of 10:1 to 0.01:1, preferably 5:1 to 0.05:1, in particular 1:1 to 0.05:1.
 20

Depending on the nature of the desired effect, the application rates of the mixtures according to the invention are, in particular, in agricultural crops, from 0.01 to 8 kg/ha, preferably 0.1 to 5 kg/ha, in particular 0.5 to 3.0 kg/ha.
 25

The application rates are, in the case of the compounds I, from 0.01 to 2.5 kg/ha, preferably 0.05 to 2.5 kg/ha, in particular 0.05 to 1.0 kg/ha.

30 In the case of the compounds II, the application rates are from 0.01 to 10 kg/ha, preferably 0.05 to 5 kg/ha, in particular 0.1 to 2.0 kg/ha.

For seed treatment, the application rates of the mixture are
 35 generally from 0.001 to 250 g/kg of seed, preferably 0.01 to 100 g/kg, in particular 0.01 to 50 g/kg.

If phytopathogenic harmful fungi are to be controlled, the separate or joint application of the compounds I and II or of
 40 the mixtures of the compounds I and II is effected by spraying or dusting the seeds, the plants or the soils before or after sowing of the plants, or before or after plant emergence.

The fungicidal synergistic mixtures according to the invention,
 45 or the compounds I and II, can be formulated for example in the form of ready-to-spray solutions, powders and suspensions or in the form of highly concentrated aqueous, oily or other



suspensions, dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading or granules, and applied by spraying, atomizing, dusting, spreading or pouring. The use form depends on the intended purpose; in any case, it should
5 guarantee as fine and uniform as possible a distribution of the mixture according to the invention.

The formulations are prepared in a manner known per se, eg. by adding solvents and/or carriers. It is usual to admix inert
10 additives, such as emulsifiers or dispersants, to the formulations.

Suitable surfactants are the alkali metal salts, alkaline earth metal salts and ammonium salts of aromatic sulfonic acids, eg.
15 ligno-, phenol-, naphthalene- and dibutyl-naphthalenesulfonic acid, and of fatty acids, alkyl- and alkylarylsulfonates, alkyl, lauryl ether and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and octadecanols, or of fatty alcohol glycol ethers, condensates of sulfonated naphthalene and its
20 derivatives with formaldehyde, condensates of naphthalene or of the naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene octylphenol ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl polyether alcohols, isotridecyl
25 alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers or polyoxypropylene [lacuna], lauryl alcohol polyglycol ether acetate, sorbitol esters, lignosulfite waste liquors or methylcellulose.

30 Powders, materials for spreading and dusts can be prepared by mixing or jointly grinding the compounds I and II or the mixture of the compounds I and II with a solid carrier.

Granules (eg. coated granules, impregnated granules or
35 homogeneous granules) are usually prepared by binding the active ingredient, or active ingredients, to a solid carrier.

Fillers or solid carriers are, for example, mineral earths, such as silica gel, silicic acids, silica gels, silicates, talc,
40 kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, and fertilizers, such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal,
45 tree bark meal, wood meal and nutshell meal, cellulose powders or other solid carriers.



The formulations generally comprise 0.1 to 95% by weight, preferably 0.5 to 90% by weight, of one of the compounds I or II or of the mixture of the compounds I and II. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR or HPLC spectra).

The compounds I and II, or the mixtures, or the corresponding formulations, are applied by treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with a fungicidally effective amount of the mixture, or of the compounds I and II in the case of separate application.

15 Application can be effected before or after infection by the harmful fungi.

Use Example

20 The synergistic action of the mixtures according to the invention was demonstrated by the following experiments:

The active ingredients, separately or together, were formulated as a 10% emulsion in a mixture of 70% by weight of cyclohexanone, 20% by weight of Nekanil[®] LN (Lutensol[®] AP6, wetting agent having emulsifying and dispersing action, based on ethoxylated alkylphenols) and 10% by weight of Emulphor[®] EL (Emulan[®] EL, emulsifier based on ethoxylated fatty alcohols) and diluted with water to give the desired concentration.

30 Example 1

Activity against powdery mildew of wheat

Leaves of wheat seedlings cv. "Frühgold" which had been grown in pots were sprayed to runoff point with an aqueous preparation of active ingredient which had been made from a stock solution comprising 10% of active compound, 63% of cyclohexanone and 27% of emulsifier, and, 24 hours after the spray coating had dried on, were dusted with spores of powdery mildew of wheat (*Erysiphe graminis forma specialis tritici*). The test plants were subsequently placed in a greenhouse at from 20 to 24°C and a relative atmospheric humidity of 70 to 90%. After 7 days, the extent of mildew development was determined visually in % infection of the total leaf area.



Evaluation was carried out by determining the infected leaf areas in percent. These percentages were converted into degrees of action. The efficacy (E) was calculated as follows using Abbot's formula:

5

$$E = (1 - \alpha) \cdot 100 / \beta$$

- α corresponds to the fungal infection of the treated plants in % and
 10 β corresponds to the fungal infection of the untreated (control) plants in %

A degree of action of 0 means that the infection level of the treated plants corresponds to that of the untreated control plants; a degree of action of 100 means that the treated plants were not infected.

15

The expected degrees of action of the mixtures of the active ingredients were determined using Colby's formula [R.S. Colby, Weeds 15, (1967) 20-22] and compared with the observed degrees of action.

20

$$\text{Colby's formula: } E = x + y - x \cdot y / 100$$

25

- E expected degree of action, expressed in % of the untreated control, when using the mixture of the active ingredients A and B at the concentrations a and b
 x degree of action, expressed in % of the untreated control, when using active ingredient A at a concentration of a
 30 y degree of action, expressed in % of the untreated control, when using active ingredient B at a concentration of b

The results of the activity against powdery mildew of wheat are shown in the tables that follow.

35

Ex.	Active ingredient	Concentration of active ingredient in the spray liquor in ppm	Efficacy in % of the untreated control
40 1V	Control (untreated)	(100% infestation)	0
2V	Ia	0.6 0.3	0 0
45 3V	Ib	0.6 0.3	15 15



4V	Compound I.32 (Id)	0.63 0.31	75 40
5V	IIa	12.5 6.3 3.1	10 0 0

Table 3

Ex.	Mixture according to the invention	Observed efficacy	Calculated efficacy ^{*)}
6	0.6 ppm Ia + 12.5 ppm IIa	40	10
7	0.3 ppm Ia + 12.5 ppm IIa	40	10
8	0.6 ppm Ib + 12.5 ppm IIa	85	23.5
9	0.3 ppm Ib + 12.5 ppm IIa	80	23.5
10	0.63 ppm Id + 6.3 ppm IIa	95	75
11	0.31 ppm Id + 3.1 ppm IIa	80	40

^{*)} calculated using Colby's formula

The test results show that the observed efficacy in all mixing ratios is higher than the efficacy that had been calculated beforehand using Colby's formula.

35

"Comprises/comprising" when used in this specification is taken to specify the presence of stated features, integers, steps or components but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof".

40


45



We claim:


1. A fungicidal mixture comprising

a₁) a phenyl benzyl ether derivative of the formula I.a, I.b or I.c,



and/or


a₂) a carbamate of the formula Id



where X is CH or N, n is 0, 1 or 2 and R is halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl, it being possible for the radicals R to be different if n is 2, or one of its salts or adducts,

and

b) a dinitrophenol derivative comprising compounds of the formula IIa and/or IIb.



where n is 0, 1 or 2
in a synergistically effective amount.

2. A fungicidal mixture according to claim 1 wherein the dinitrophenol derivative comprises a mixture of IIa and IIb.
3. A fungicidal mixture as claimed in claim 1 or 2 wherein the weight ratio of the compound I to the compound II is from 10:1 to 0.01:1.
4. A fungicidal mixture comprising a synergistically effective amount of phenyl benzyl ethyl derivative and/or a carbamate and a dinitrophenyl derivative, which mixture is substantially as herein described with reference to any one of Examples 6 to 11.
5. A method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with a compound of the formula I as set forth in claim 1 and a compound of the formula II as set forth in claim 1 in a synergistically effective amount.
6. A method as claimed in claim 5, wherein a compound I as set forth in claim 1 and a compound II as set forth in claim 1 are applied simultaneously, or in succession.
7. A method as claimed in claim 6, wherein the simultaneous application of compound I and compound II includes application separately, or together.
8. A method as claimed in any one of claims 5 to 7, wherein the compound I as set forth in claim 1 is applied in an amount of from 0.01 to 2.5 kg/ha.
9. A method as claimed in any one of claims 5 to 8, wherein the compound II as set forth in claim 1 is applied in an amount of from 0.01 to 10 kg/ha.



9. A method as claimed in any one of claims 5 to 8, wherein the compound II as set forth in claim 1 is applied in an amount of from 0.01 to 10 kg/ha.

10. A method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with a fungicidal mixture of claim 4.

11. The use of a compound I as set forth in claims 1 for preparing a fungicidally effective synergistic mixture as claimed in claim 1 or 2.

12. The use of a compound II as set forth in claim 1 for preparing a fungicidally effective synergistic mixture as claimed in claim 1 or 2.

13. A composition as claimed in claim 1 comprising two parts, one part comprising a compound of the formula I as set forth in claim 1 in a solid or liquid carrier and the other part comprising a compound of the formula II as set forth in claim 1 in a solid or liquid carrier, when used for controlling harmful fungi as claimed in any one of claims 5 to 9.

DATED this 10th Day of August, 2001

BASF AKTIENGESELLSCHAFT

WATERMARK PATENT & TRADEMARK ATTORNEYS
290 BURWOOD ROAD
HAWTHORN VICTORIA 3122
AUSTRALIA

P16257AU00/ LCG:KML:HB

