

(19) AUSTRALIAN PATENT OFFICE

- (54) Title
Food products comprising long chain polyunsaturated fatty acids and methods for preparing the same
- (51)⁶ International Patent Classification(s)
A23D 9/00 (2006.01) 20060101AFI2007120
A23D 9/00 7BHS
PCT/US2007/066471
- (21) Application No: 2007238131 (22) Application Date: 2007.04.11
- (87) WIPO No: W007/121273
- (30) Priority Data
- | | | |
|-------------|------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/791,358 | 2006.04.11 | US |
- (43) Publication Date : 2007.10.25
- (71) Applicant(s)
Martek Biosciences Corporation
- (72) Inventor(s)
Crandell, Michelle, Abril, Jesus Ruben
- (74) Agent/Attorney
Allens Arthur Robinson Patent & Trade Marks Attorneys, Level 27 530 Collins Street,
Melbourne, ACT, 3000
- (56) Related Art
US 5882703
US 5023100
US 6461662
US 5518753

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2007 (25.10.2007)

PCT

(10) International Publication Number
WO 2007/121273 A3

- (51) International Patent Classification:
A23D 9/00 (2006.01)
- (21) International Application Number:
PCT/US2007/066471
- (22) International Filing Date: 11 April 2007 (11.04.2007)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/791,358 11 April 2006 (11.04.2006) US
- (71) Applicant (for all designated States except US):
MARTEK BIOSCIENCES CORPORATION [US/US];
6480 Dobbins Road, Columbia, Massachusetts 21045 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ABRIL, Jesus Ruben [US/US]; 6638 West 98th Drive, Westminster, Colorado 80021 (US). CRANDELL, Michelle [US/US]; 5121 Williams Fork Trail, #201, Boulder, Colorado 80301 (US).
- (74) Agent: YOERG, Darla G., Sheridan Ross P.C., 1560 Broadway, Suite 1200, Denver, Colorado 80202 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LI, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
24 July 2008

WO 2007/121273 A3

(54) Title: FOOD PRODUCTS COMPRISING LONG CHAIN POLYUNSATURATED FATTY ACIDS AND METHODS FOR PREPARING THE SAME

(57) Abstract: The present invention includes a food oil composition comprising a blend of a first oil comprising an LC PUFA and a second oil comprising substantially no LC PUFA. The first oil can preferably comprise an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof. The present invention also provides methods of food preparation, more particularly, methods for skillet-frying, deep-frying, methods for preparing edible lipid-containing food sauces, methods for preparing extruded food products, and methods for enhancing the LC PUFA content of a food product, particularly previously cooked food products, and food products prepared in accordance with such methods. Such compositions and methods are useful, for example, for increasing intake of LC PUFAs.

FOOD PRODUCTS COMPRISING LONG CHAIN POLYUNSATURATED FATTY ACIDS AND METHODS FOR PREPARING THE SAME

FIELD OF THE INVENTION

5 The invention relates to food oil compositions, methods for food preparation, and food products comprising long chain polyunsaturated fatty acids, and particularly, omega-3 long chain polyunsaturated fatty acids, omega-6 long chain polyunsaturated fatty acids, and mixtures thereof.

BACKGROUND

10 It is desirable to increase the dietary intake of the beneficial omega-3 polyunsaturated fatty acids (omega-3 PUFA), and omega-3 long chain polyunsaturated fatty acids (LC PUFA). Other beneficial nutrients are omega-6 long chain polyunsaturated fatty acids. As used herein, reference to a long chain polyunsaturated fatty acid or LC PUFA, refers to a polyunsaturated fatty acid having 20 or more carbons.

15 Omega-3 PUFAs are recognized as important dietary compounds for preventing arteriosclerosis and coronary heart disease, for alleviating inflammatory conditions, cognitive impairment and dementia related diseases and for retarding the growth of tumor cells. One important class of omega-3 PUFAs is omega-3 LC PUFAs. Omega-6 LC-PUFAs serve not only as structural lipids in the human body, but also as precursors

20 for a number of factors in inflammation such as prostaglandins, and leukotrienes.

Fatty acids are carboxylic acids and are classified based on the length and saturation characteristics of the carbon chain. Short chain fatty acids have 2 to about 6 carbons and are typically saturated. Medium chain fatty acids have from about 6 to about 18 carbons and may be saturated or unsaturated. Long chain fatty acids have from

25 20 to 24 or more carbons and may also be saturated or unsaturated. In longer chain fatty acids there may be one or more points of unsaturation, giving rise to the terms "monounsaturated" and "polyunsaturated," respectively. Long chain PUFAs (LC PUFAs) are of particular interest in the present invention.

LC PUFAs are categorized according to the number and position of double bonds

30 in the fatty acids according to a well understood nomenclature. There are two series or families of LC PUFAs, depending on the position of the double bond closest to the methyl end of the fatty acid: the ω -3 (or n-3 or omega-3) series contains a double bond at the third carbon, while the ω -6 (or n-6 or omega-6) series has no double bond until the sixth carbon. Thus, docosahexaenoic acid ("DHA") has a chain length of 22 carbons

with 6 double bonds beginning with the third carbon from the methyl end and is designated "22:6 n-3". Another important LC PUFA is eicosapentaenoic acid ("EPA") which is designated "20:5 n-3".

De novo or "new" synthesis of the omega-3 and omega-6 fatty acids such as
5 DHA and ARA does not occur in the human body; however, the body can convert shorter chain fatty acids to LC PUFAs such as DHA and ARA although at very low efficiency. Both omega-3 and omega-6 fatty acids must be part of the nutritional intake since the human body cannot insert double bonds closer to the omega end than the seventh carbon atom counting from that end of the molecule. Thus, all metabolic
10 conversions occur without altering the omega end of the molecule that contains the omega-3 and omega-6 double bonds. Consequently, omega-3 and omega-6 acids are two separate families of essential fatty acids since they are not interconvertible in the human body.

Over the past twenty years, health experts have recommended diets lower in
15 saturated fats and higher in polyunsaturated fats. While this advice has been followed by a number of consumers, the incidence of heart disease, cancer, diabetes and many other debilitating diseases has continued to increase steadily. Scientists agree that the type and source of polyunsaturated fats is as critical as the total quantity of fats. The most common polyunsaturated fats are derived from vegetable matter and are lacking in long
20 chain fatty acids (most particularly omega-3 LC-PUFAs). In addition, the hydrogenation of polyunsaturated fats to create synthetic fats has contributed to the rise of certain health disorders and exacerbated the deficiency in some essential fatty acids. Indeed, many medical conditions have been identified as benefiting from an omega-3 supplementation. These include acne, allergies, Alzheimer's, arthritis, atherosclerosis,
25 breast cysts, cancer, cystic fibrosis, diabetes, eczema, hypertension, hyperactivity, intestinal disorders, kidney dysfunction, leukemia, and multiple sclerosis. Of note, the World Health Organization has recommended that infant formulas be enriched with omega-3 and omega-6 fatty acids.

The polyunsaturates derived from meat contain significant amounts of omega-6
30 but little or no omega-3. While omega-6 and omega-3 fatty acids are both necessary for good health, they must be consumed in a balance of about 4:1. Today's Western diet has created a serious imbalance with current consumption on average of 20 times more omega-6 than omega-3. Concerned consumers have begun to look for health food

supplements to restore the equilibrium. Principal sources of omega-3 are flaxseed oil and fish oils. The past decade has seen rapid growth in the production of flaxseed and fish oils. Both types of oil are considered good dietary sources of omega-3 polyunsaturated fats. Flaxseed oil contains no EPA, DHA, or DPA but rather contains
5 linolenic acid--a building block that can be elongated by the body to build longer chain PUFAs. There is evidence, however, that the rate of metabolic conversion can be slow and unsteady, particularly among those with impaired health. Fish oils vary considerably in the type and level of fatty acid composition depending on the particular species and their diets. For example, fish raised by aquaculture tend to have a lower level of omega-
10 3 fatty acids than fish from the wild. In light of the health benefits of such omega-3 and omega-6 LC-PUFAs, it would be desirable to supplement foods with such fatty acids.

Due to the scarcity of sources of omega-3 LC PUFAs, typical home-prepared and convenience foods are low in both omega-3 PUFAs and omega-3 LC PUFAs (chain length greater than 20), such as docosahexaenoic acid, docosapentaenoic acid, and
15 eicosapentaenoic acid. In light of the health benefits of such omega-3 LC PUFAs (chain length greater than 20), it would be desirable to supplement foods with such fatty acids.

While foods and dietary supplements prepared with LC PUFAs may be healthier, they also have an increased vulnerability to rancidity. Rancidity in lipids, such as unsaturated fatty acids, is associated with oxidation off-flavor development. The
20 oxidation off-flavor development involves food deterioration affecting flavor, aroma, and the nutritional value of the particular food. A primary source of oxidation off-flavor development in lipids, and consequently the products that contain them, is the chemical reaction of lipids with oxygen. The rate at which this oxidation reaction proceeds has generally been understood to be affected by factors such as temperature, degree of
25 unsaturation of the lipids, oxygen level, ultraviolet light exposure, presence of trace amounts of pro-oxidant metals (such as iron, copper, or nickel), lipoxidase enzymes, and so forth.

The susceptibility and rate of oxidation of the unsaturated fatty acids can rise dramatically as a function of increasing degree of unsaturation in particular. In this
30 regard, EPA and DHA contain five and six double bonds, respectively. This high level of unsaturation renders the omega-3 fatty acids readily oxidizable. The natural instability of such oils gives rise to unpleasant odor and unsavory flavor characteristics even after a relatively short period of storage time.

PUFAs may be extracted from microbial sources for use in nutritional and/or pharmaceutical products. For example, DHA-rich microbial oil is manufactured from the dinoflagellate *Cryptocodinium cohnii* and ARA-rich oil is manufactured from the filamentous fungus *Mortierella alpina*, both for use as nutritional supplements and in food products such as infant formula. Similarly, DHA-rich microbial oil from *Schizochytrium* is manufactured for use as a nutritional supplement or food ingredient. Typically, the LC PUFAs are extracted from biomass and purified. The extracted and purified oils can be further processed to achieve specific formulations for use in food products (such as a dry powder or liquid emulsion).

In light of the desirability of supplementing foods with omega-3 LC PUFAs and/or omega-6 LC PUFAs, and in view of the shortcomings of the prior art in providing these foods, there is a need for methods for enriching foods with omega-3 LC PUFAs and/or omega-6 LC PUFAs and also for food oil compositions and food products comprising omega-3 LC PUFAs and/or omega-6 LC PUFAs. These and other needs are answered by the present invention.

SUMMARY OF THE INVENTION

According to a first aspect of the present invention, there is provided a food oil composition comprising a blend of:

- a first oil comprising an LC PUFA, and
- a second oil comprising substantially no LC PUFA and that is liquid at room temperature,

wherein the blend comprises about 0.01% to 0.5% LC PUFA.

According to a second aspect of the present invention, there is provided a topical food oil composition comprising a blend of a first oil comprising an LC PUFA, a second oil comprising substantially no LC PUFA and that is liquid at room temperature, and an antioxidant, wherein the LC PUFA comprises about 0.01% to 0.5% of the food oil composition.

According to a third aspect of the present invention, there is provided a food product comprising the composition according to either of the first or second aspects of the present invention.

According to a fourth aspect of the present invention, there is provided a method of food preparation for a food item capable of being skillet-fried, comprising:

- (a) placing the food item and an oil onto the skillet; and
- 5 (b) applying heat to the skillet sufficient to heat the food item, wherein the oil comprises a composition according to either of the first or second aspects of the present invention.

According to a fifth aspect of the present invention, there is provided a method of food preparation, comprising topically applying a composition
10 according to either of the first or second aspects of the present invention to a food product, wherein the topically applying is selected from the group consisting of spraying, dipping and brushing.

According to a sixth aspect of the present invention, there is provided a food preparation for a food item capable of being deep-fried comprising:

- 15 (a) immersing the food item in an oil; and
- (b) applying heat to the oil sufficient to heat the food item, wherein the oil comprises a composition according to either of the first or second aspects of the present invention.

Embodiments of the present invention are directed toward food oil
20 compositions and their uses in food products. The food oil compositions generally include a blend of a first oil having LC PUFAs and preferably, an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof and a second oil that includes substantially no LC PUFAs, and preferably, substantially no omega-3 LC PUFA and substantially no omega-6 LC-PUFA and that is liquid at room temperature.

25 In one embodiment, the food oil composition includes a blend of a first oil comprising an LC PUFA, and preferably an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof and a second oil comprising substantially no LC PUFAs, and preferably, substantially no omega-3 LC PUFA, wherein the second oil is liquid at room temperature. In an alternate embodiment the food oil composition includes
30 a blend of a first oil comprising an LC PUFA, and preferably an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof and a second oil comprising substantially

no LC PUFAs, and substantially no omega-6 LC PUFA, wherein the second oil is liquid at room temperature. In these embodiments, the blend comprises between about 0.01% and about 5% of the LC PUFAs. In a further embodiment, the blend can comprise between about 0.08% and about 3% LC PUFAs or between about 5 0.1% and about 0.5% LC PUFAs. This first embodiment of the invention is particularly useful for preparing skillet-fried food products. Such products can include between about 5mg and about 150mg omega-3 LC PUFAs, omega-6 LC PUFAs or mixtures thereof per food product or serving. A further aspect of this embodiment is a method for food preparation of a food item capable of being 10 skillet fried. This method includes placing the food item and an oil on to a skillet. The oil includes the first food oil composition embodiment described above. Heat is applied to the skillet sufficient to heat the food item, thereby frying the food. In an alternate embodiment, this food oil composition is useful for preparing deep-fried food products, such as tempura or fries, as well as methods for food 15 preparation of a food item capable of being deep-fried. This method includes immersing the food item in an oil. The oil includes the first food oil composition embodiment described above. Heat is applied to the oil sufficient to heat the food item, thereby deep-frying the food.

A further embodiment of the present invention includes a blend of a first oil 20 comprising an LC PUFA and preferably, an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof and a second oil comprising substantially no LC PUFAs and preferably substantially no omega-3 LC PUFAs and substantially no omega-6 LC PUFAs, and wherein the second oil is liquid at room temperature. In this embodiment, the LC PUFA content of the blend is between about 1% and about 25 30%. In this embodiment, the LC PUFA content of the oil blend can also be between about 10% and about 20%, or between about 1% and about 5%. The second food oil composition embodiment can be used in a method for preparing a food product that includes contacting an oil with additional food components. Such food products can include any edible lipid-containing food sauce, such as 30 salad dressings, marinades, remoulades, vegetable sauces, fruit sauces, fish sauces, and meat sauces, such as poultry sauces, beef sauces, veal sauces, and lamb sauces.

Another embodiment of the present invention includes a topical food oil composition that includes a blend of a first oil having an LC PUFA and preferably, an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof, a second oil comprising substantially no LC PUFAs and preferably, no omega-3 LC PUFAs and
5 substantially no omega-6 LC PUFAs, and that is liquid at room temperature and an antioxidant. In this embodiment, the blend comprises between about 0.25% and about

10

15

[This region of the page is intentionally left blank]

10% LC PUFA. In this embodiment, the LC PUFA content of the blend can also be between about 1% and about 5%. A further embodiment of the present invention is a food product comprising the third food oil composition embodiment. The food product can be selected from a previously cooked food product, such as one that was previously
5 baked, fried, or deep-fried. The food product can be selected from baked goods, salted snacks, specialty snacks, confectionary snacks, and naturally occurring snack foods. For example, the food product can be selected from cookies, crackers, sweet goods, muffins, cereals, snack cakes, pies, granola/snack bars, toaster pastries, potato chips, corn chips, wheat chips, sorghum chips, soy chips, extruded snacks, popcorn, pretzels, potato crisps,
10 dried fruit snacks, meat snacks, pork rinds, health food bars, rice cakes, corn cakes, candy, nuts, dried fruits and vegetables.

A further embodiment of the present invention is a method of food preparation that includes topically applying the third food oil composition embodiment to a food product. The step of topically applying can be selected from spraying, dipping and
15 brushing. This method can further include packaging the food product after application of the food oil composition. The step of packaging can include packaging the food product in an inert atmosphere. Such an atmosphere can include nitrogen or can include nitrogen and carbon dioxide.

All of the food oil composition embodiments of the present invention can further
20 include an antioxidant, which can be selected from Vitamin E, BHT, BHA, TBHQ, propyl gallate, Vitamin C, phospholipids and natural antioxidants and combinations thereof. Preferred antioxidants include BHA, BHT, TBHQ, a blend of BHA/BHT, and combinations thereof, and particularly, TBHQ. In preferred embodiments, the antioxidant can be present in the oil blend in an amount between about 0.01% and about
25 1% and alternatively between about 0.1% and about 0.5%.

In various embodiments of the food oil compositions, the second oil can be selected from borage oil, black currant seed oil, corn oil, coconut oil, canola oil, soybean oil, safflower oil, high oleic safflower oil, sunflower oil, high oleic sunflower oil, olive oil, evening primrose oil, cottonseed oil, rice bran oil, grapeseed oil, flaxseed oil, garlic
30 oil, peanut oil, almond oil, walnut oil, wheat germ oil, sesame oil, animal fat, animal oil, marine fat, marine oil, microbial oil, a hydrogenated oil of any of the foregoing, and mixtures of the foregoing. The omega-3 LC PUFA and/or omega-6 LC PUFA in various embodiments of the present invention can be selected from docosahexaenoic acid,

eicosapentaenoic acid, docosapentaenoic acid, and arachidonic acid (ARA). In various embodiments, the first oil can be from a microbial source, such as algae, protists, bacteria and fungi. The microbial source can be an oleaginous microorganism. The microbial source can be selected from microorganisms of the genus *Thraustochytrium*, microorganisms of the genus *Schizochytrium*, microorganisms of the genus *Althornia*, microorganisms of the genus *Aplanochytrium*, microorganisms of the genus *Japonochytrium*, microorganisms of the genus *Elina*, microorganisms of the genus *Cryptocodinium*, and microorganisms of the genus *Mortierella*. In preferred embodiments, the microorganism is selected from microorganisms of the genus *Schizochytrium*, microorganisms of the genus *Cryptocodinium*, and microorganisms of the genus *Mortierella*.

The first oil can also be from a plant source, such as plants that have been genetically modified to produce LC PUFAs, wherein the plant is selected from soybean, corn, safflower, sunflower, canola, flax, peanut, mustard, rapeseed, chick pea, cotton, lentil, white clover, olive, palm, borage, evening primrose, linseed and tobacco.

In a further embodiment, the first oil can be from an animal source, which can be selected from aquatic animals, lipids extracted from animal tissues and animal products. Further, the first oil can include at least about 20% omega-3 LC PUFAs and/or omega-6 LC PUFAs or at least about 60% omega-3 LC PUFAs and/or omega-6 LC PUFAs.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 illustrates the results of consumer testing of food products (fried potatoes, omelets and fried French toast) of the present invention.

Fig. 2 illustrates the effect on OSI induction period of blending various vegetable oils with an omega-3 LC PUFA-containing oil.

Fig. 3 illustrates the effect on OSI induction period of blending various vegetable oils with an omega-3 LC PUFA-containing oil, with and without antioxidants.

Fig. 4 illustrates the effect on production of primary oxidation products of blending corn oil with an omega-3 LC PUFA-containing oil, with and without antioxidants.

Fig. 5 illustrates the effect on production of secondary oxidation products of blending corn oil with an omega-3 LC PUFA-containing oil, with and without antioxidants.

Fig. 6 illustrates the effect on OSI induction period of blending corn oil with an omega-3 LC PUFA-containing oil.

Fig. 7 illustrates the effect on OSI induction period of blending soybean oil with an omega-3 LC PUFA-containing oil.

5 Fig. 8 illustrates the effect on OSI induction period of blending canola oil with an omega-3 LC PUFA-containing oil.

Fig. 9 illustrates the effect on OSI induction period of blending safflower oil with an omega-3 LC PUFA-containing oil.

10 Fig. 10 illustrates the effect on OSI induction period of blending sunflower oil with an omega-3 LC PUFA-containing oil.

Fig. 11 illustrates the peroxide values of a blend of corn oil and an omega-3 LC PUFA-containing oil over time.

Fig. 12 illustrates the alkenal values of a blend of corn oil and an omega-3 LC PUFA-containing oil over time.

15 Fig. 13 illustrates the DHA content of a blend of corn oil and an omega-3 LC PUFA-containing oil over time.

DETAILED DESCRIPTION

20 The food oil and food product compositions, methods for food preparation, and methods for enhancing the LC PUFA content and preferably, the omega-3 LC PUFA and/or omega-6 LC PUFA content of previously prepared food products, as taught by the present invention, provide for increased intake of LC PUFAs and particularly omega-3 LC PUFAs and/or omega-6 LC PUFAs. This improvement can provide health benefits to those consuming such products. The present invention also provides methods to 25 minimize the oxidative degradation of LC PUFAs in the food products and food oil compositions.

In various embodiments, the present invention includes a food oil composition comprising a blend of a first oil comprising an LC PUFA and preferably, an omega-3 LC PUFA and/or omega-6 LC PUFA and a second oil comprising substantially no LC PUFA 30 and preferably, substantially no omega-3 LC PUFA and substantially no omega-6 LC PUFA that is liquid at room temperature. In a first embodiment, the blend comprises between about 0.01% and about 5% LC PUFAs. This embodiment of the food oil composition is particularly useful for fast frying food, such as in a skillet, to impart LC

PUFAs and preferably, omega-3 LC PUFAs, omega-6 LC PUFAs or mixtures thereof into a diet. In an alternate embodiment, this food oil composition is useful for preparing deep-fried food products, such as tempura or fries, in which the food item is immersed in the oil. In a second food oil composition embodiment, the LC PUFA and preferably, the omega-3 LC PUFA and/or omega-6 LC PUFA content of the blend is between about 1% and about 30%. This second food oil composition embodiment is particularly useful in food products such as edible lipid-containing food sauces, such as salad dressings, marinades, remoulades, vegetable sauces, fruit sauces, fish sauces, and meat sauces. In a third food oil composition embodiment, the blend comprises between about 0.25% and about 10% LC PUFA and preferably, omega-3 LC PUFA and/or omega-6 LC PUFA and the composition further includes an antioxidant. This third embodiment is particularly useful for topical application of the composition to foods, such as baked goods, salted snacks, specialty snacks, confectionary snacks, and naturally occurring snack foods. Such foods with topical applications of the oil composition are typically packaged products and are packaged in an inert atmosphere.

A food oil preferably contains greater than about 90% fatty acids by weight, whereas a product such as margarine and butter is typically an emulsion of fat and water having a fatty acid content of between about 80% by weight and about 95% by weight. As used herein, all percentages are given by weight unless explicitly stated otherwise.

The oil blend of the present invention includes a first oil that comprises an LC PUFA, and preferably an omega-3 LC PUFA, an omega-6 LC PUFA or mixtures thereof. Preferred omega-3 LC PUFAs include, for example, docosahexaenoic acid C22:6(n-3) (DHA), eicosapentaenoic acid C20:5(n-3)(EPA), and docosapentaenoic acid C22:5(n-3) (DPA). DHA is particularly preferred. Preferred omega-6 LC-PUFAs include arachidonic acid C20:4(n-6) (ARA). The PUFAs can be in any of the common forms found in natural lipids including but not limited to triacylglycerols, diacylglycerols, phospholipids, free fatty acids, esterified fatty acids, or in natural or synthetic derivative forms of these fatty acids (e.g. calcium salts of fatty acids, ethyl esters, etc). Reference to a first oil comprising an omega-3 LC PUFA and/or omega-6 LC PUFA, as used in the present invention, can refer to either an oil comprising only a single omega-3 LC PUFA and/or omega-6 LC PUFA such as DHA or an oil comprising a mixture of omega-3 LC PUFAs and/or omega-6 LC PUFA such as DHA and EPA, or DHA and ARA.

A preferred source of oils that comprise LC PUFAs and preferably, omega-3 LC PUFAs and/or omega-6 LC PUFAs, in the compositions and methods of the present invention includes a microbial source. Microbial sources and methods for growing microorganisms comprising nutrients and/or LC PUFAs are known in the art (*Industrial*
5 *Microbiology and Biotechnology*, 2nd edition, 1999, American Society for Microbiology). Preferably, the microorganisms are cultured in a fermentation medium in a fermentor. The methods and compositions of the present invention are applicable to any industrial microorganism that produces any kind of nutrient or desired component such as, for example algae, protists, bacteria and fungi (including yeast).

10 Microbial sources can include microorganisms such as algae, bacteria, fungi and/or protists. Preferred organisms include those selected from the group consisting of golden algae (such as microorganisms of the kingdom Stramenopiles), green algae, diatoms, dinoflagellates (such as microorganisms of the order Dinophyceae including members of the genus *Cryptocodinium* such as, for example, *Cryptocodinium cohnii*),
15 yeast, and fungi of the genera *Mucor* and *Mortierella*, including but not limited to *Mortierella alpina* and *Mortierella* sect. *schmuckeri*. Members of the microbial group Stramenopiles include microalgae and algae-like microorganisms, including the following groups of microorganisms: Hamatores, Proteromonads, Opalines, Develpayella, Diplophrys, Labrinthulids, Thraustochytrids, Bioseccids, Oomycetes,
20 Hypochytridiomycetes, Commation, Reticulosphaera, Pelagomonas, Pelagococcus, Ollicola, Aurcococcus, Parmals, Diatoms, Xanthophytes, Phaeophytes (brown algae), Eustigmatophytes, Raphidophytes, Synurids, Axodines (including Rhizochromulinaales, Pedinellales, Dictyochales), Chrysomeridales, Sarcinochrysidales, Hydrurales, Hibberdiales, and Chromulinales. This detailed description of the invention will discuss
25 processes for growing microorganisms which are capable of producing lipids comprising omega-3 and/or omega-6 polyunsaturated fatty acids, in particular microorganisms that are capable of producing DHA (or closely related compounds such as DPA, EPA or ARA). Additional preferred microorganisms are algae, such as Thraustochytrids of the order Thraustochytriales, more specifically Thraustochytriales, including
30 *Thraustochytrium*, *Schizochytrium* and *Ulkenia*, and including Thraustochytriales which are disclosed in commonly assigned U.S. Patent Nos. 5,340,594 and 5,340,742, both issued to Barclay, all of which are incorporated herein by reference in their entirety, in addition to microorganisms of the genus *Althornia*, genus *Aplanochytrium*, genus

Japonochytrium, and genus *Elina* and mixtures thereof. More preferably, the microorganisms are selected from the group consisting of microorganisms having the identifying characteristics of ATCC number 20888, ATCC number 20889, ATCC number 20890, ATCC number 20891 and ATCC number 20892, strains of *Mortierella schmuckeri* and *Mortierella alpina*, strains of *Cryptocodium cohnii*, mutant strains derived from any of the foregoing, and mixtures thereof. It should be noted that many experts agree that *Ulkenia* is not a separate genus from the genus *Thraustochytrium*. Accordingly, as used herein, the genus *Thraustochytrium* will include *Ulkenia*. Oleaginous microorganisms are also preferred. As used herein, "oleaginous microorganisms" are defined as microorganisms capable of accumulating greater than 20% of the weight of their cells in the form of lipids. Genetically modified microorganisms that produce LC PUFAs are also suitable for the present invention. These can include naturally LC PUFA-producing microorganisms that have been genetically modified as well as microorganisms that do not naturally produce LC PUFAs (including yeast, bacteria, fungi, algae and/or protists) but that have been genetically engineered to do so.

Suitable organisms may be obtained from a number of available sources, including by collection from the natural environment. For example, the American Type Culture Collection currently lists many publicly available strains of microorganisms identified above. As used herein, any organism, or any specific type of organism, includes wild strains, mutants, or recombinant types. Growth conditions in which to culture or grow these organisms are known in the art, and appropriate growth conditions for at least some of these organisms are disclosed in, for example, U.S. Patent No. 5,130,242, U.S. Patent No. 5,407,957, U.S. Patent No. 5,397,591, U.S. Patent No. 5,492,938, and U.S. Patent No. 5,711,983, all of which are incorporated herein by reference in their entirety.

Another preferred source of oils comprising LC PUFAs includes a plant source, such as oilseed plants. Since plants do not naturally produce LC PUFAs, plants producing LC PUFAs are those genetically engineered to express genes that produce LC PUFAs. Such genes can include genes encoding proteins involved in the classical fatty acid synthase pathways, or genes encoding proteins involved in the PUFA polyketide synthase (PKS) pathway. The genes and proteins involved in the classical fatty acid synthase pathways, and genetically modified organisms, such as plants, transformed with

such genes, are described, for example, in Napier and Sayanova, *Proceedings of the Nutrition Society* (2005), 64:387-393; Robert et al., *Functional Plant Biology* (2005) 32:473-479; or U.S. Patent Application Publication 2004/0172682. The PUFA PKS pathway, genes and proteins included in this pathway, and genetically modified
5 microorganisms and plants transformed with such genes for the expression and production of PUFAs are described in detail in: U.S. Patent No. 6,566,583; U.S. Patent Application Publication No. 20020194641, U.S. Patent Application Publication No. 20040235127A1, and U.S. Patent Application Publication No. 20050100995A1, each of which is incorporated herein by reference in its entirety.

10 Preferred oilseed crops include soybeans, corn, safflower, sunflower, canola, flax, peanut, mustard, rapeseed, chickpea, cotton, lentil, white clover, olive, palm oil, borage, evening primrose, linseed, and tobacco that have been genetically modified to produce LC PUFA as described above.

Genetic transformation techniques for microorganisms and plants are well-known
15 in the art. Transformation techniques for microorganisms are well known in the art and are discussed, for example, in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press. A general technique for transformation of dinoflagellates, which can be adapted for use with *Cryptocodinium cohnii*, is described in detail in Lohuis and Miller, *The Plant Journal* (1998) **13**(3): 427-435. A general
20 technique for genetic transformation of Thraustochytrids is described in detail in U.S. Patent Application Publication No. 20030166207, published September 4, 2003. Methods for the genetic engineering of plants are also well known in the art. For instance, numerous methods for plant transformation have been developed, including biological and physical transformation protocols. See, for example, Miki et al.,
25 "Procedures for Introducing Foreign DNA into Plants" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 67-88. In addition, vectors and *in vitro* culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber et al., "Vectors for Plant Transformation" in *Methods in Plant Molecular Biology and*
30 *Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pp. 89-119. See also, Horsch et al., *Science* **227**:1229 (1985); Kado, C.I., *Crit. Rev. Plant. Sci.* **10**:1 (1991); Moloney et al., *Plant Cell Reports* **8**:238 (1989); U.S. Patent No. 4,940,838; U.S. Patent No. 5,464,763; Sanford et al., *Part. Sci. Technol.* **5**:27

(1987); Sanford, J.C., *Trends Biotech.* **6**:299 (1988); Sanford, J.C., *Physiol. Plant* **79**:206 (1990); Klein et al., *Biotechnology* **10**:268 (1992); Zhang et al., *Bio/Technology* **9**:996 (1991); Deshayes et al., *EMBO J.*, **4**:2731 (1985); Christou et al., *Proc Natl. Acad. Sci. USA* **84**:3962 (1987); Hain et al., *Mol. Gen. Genet.* **199**:161 (1985); Draper et al., *Plant Cell Physiol.* **23**:451 (1982); Donn et al., In Abstracts of VIIIth International Congress on Plant Cell and Tissue Culture IAPTC, A2-38, p. 53 (1990); D'Halluin et al., *Plant Cell* **4**:1495-1505 (1992) and Spencer et al., *Plant Mol. Biol.* **24**:51-61 (1994).

When oilseed plants are the source of LC PUFAs, the seeds can be harvested and processed to remove any impurities, debris or indigestible portions from the harvested seeds. Processing steps vary depending on the type of oilseed and are known in the art. Processing steps can include threshing (such as, for example, when soybean seeds are separated from the pods), dehulling (removing the dry outer covering, or husk, of a fruit, seed, or nut), drying, cleaning, grinding, milling and flaking. After the seeds have been processed to remove any impurities, debris or indigestible materials, they can be added to an aqueous solution preferably water, and then mixed to produce a slurry. Preferably, milling, crushing or flaking is performed prior to mixing with water. A slurry produced in this manner can be treated and processed the same way as described for a microbial fermentation broth. Size reduction, heat treatment, pH adjustment, pasteurization and other known treatments can be used in order to improve quality (nutritional and sensory).

Another preferred source of oils that comprise LC PUFAs includes an animal source. Examples of animal sources include aquatic animals (e.g., fish, marine mammals, and crustaceans such as krill and other euphausiids) and animal tissues (e.g., brain, liver, eyes, etc.) and animal products (e.g., eggs and milk). Techniques for recovery of LC PUFA containing oils from such sources are known in the art.

Preferably, the first oil comprises at least about 20% LC PUFA, at least about 30% LC PUFA, at least about 40% LC PUFA, at least about 50% LC PUFA, at least about 60% LC PUFA, at least about 70% LC PUFA, and at least about 80% LC PUFA.

The oil blend of the present invention includes a second oil that can include any oil known in the art. Such oils include, for example, oils derived from plants, such as borage, black currant seed, corn, coconut, canola, soybean, safflower, high oleic safflower, sunflower, high oleic sunflower, olive, evening primrose, cottonseed, rice bran, grapeseed, flaxseed, garlic, peanuts, almonds, walnuts, wheat germ, and sesame. Such vegetable sources naturally produce fatty acids only to about 18 carbons.

Additional oils suitable as the second oil of the oil composition includes animal fats and oils, marine fats and oils, microbiological oils, and combinations of any of these oils and fats. Most preferably, the balance of the oil composition in the first oil composition comprises the following oils/fats: corn oil, soy oil, canola oil, cottonseed oil, sunflower
5 oil, high oleic sunflower oil, safflower oil, high oleic safflower oil, and olive oil. Hydrogenated oils may also be used as the second oil of the oil composition, however, hydrogenated oils are not as preferred as non-hydrogenated oils. Without intending to be bound by theory, in various embodiments of the present invention, the blending of the second oil with the first oil increases the oxidative stability of the first oil (e.g., as
10 measured by increases in the OSI induction period and/or the production of primary and/or secondary oxidation products under mild accelerated conditions). Particularly, in the instance in which the second oil is corn oil, soybean oil or canola oil, the oxidative stability of the first oil can be improved.

In some embodiments, the composition is stable to oxidation for a period of time
15 when stored at room temperature. The period of time can be at least about one month, about two months, about three months, about four months, about five months, about six months, about seven months, about eight months, about nine months, about ten months, about eleven months and about twelve months. By stable, it is meant that the levels of oxidation products, such as peroxides and/or alkenals, do not increase appreciably in the
20 time interval. For example, the level of an oxidation products measured as peroxides, typically will be less than about 3.0 meq/kg fat, less than about 2.5 meq/kg fat, less than about 2.0 meq/kg fat, less than about 1.5 meq/kg fat, less than about 1.0 meq/kg fat, less than about 0.5 meq/kg fat, or less than about 0.25 meq/kg fat over the various time frames referenced above.

In addition to oxidative stability, the LC-PUFA level in the composition is stable
25 for a period of time when stored at room temperature. The period of time can be at least about one month, about two months, about three months, about four months, about five months, about six months, about seven months, or about eight months. By stable, it is meant that the levels of LC-PUFA do not decrease appreciably in the time interval. For
30 example, the level of LC-PUFA that can be recovered after the various time frames referenced above is at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, and at least about 99%.

In addition, the sensory characteristics of the composition remain constant over a period of time when stored at room temperature. By constant, it is meant that the sensory characteristic measured (e.g., green/beany, fishy, painty, herbal, or other) does not change significantly over a period of time. The period of time can be at least about
5 one month, about two months, about three months, about four months, about five months, about six months, about seven months, or about eight months. For example, a negative sensory characteristic typically will increase less than about 100%, less than about 75%, less than about 50%, less than about 40%, less than about 30%, less than about 25%, less than about 20%, less than about 15%, less than about 10%, or less than about
10 5%, over the various time frames referenced above.

The second oil comprises substantially no LC PUFA and preferably substantially no omega-3 LC PUFA and substantially no omega-6 LC PUFA. Generally, reference to substantially no LC PUFA includes oils having less than about 5% LC PUFA, less than about 3% LC PUFA, less than about 1% LC PUFA, less than about 0.1% LC PUFA, or
15 less than about 0.01% LC PUFA. The second oil is preferably also liquid at room temperature (approximately 21°C -23°C).

Blending the first oil and the second oil can be performed by any method known in the art. Blending can be done by: 1) batch or 2) continuous in-line mixing. Batch mixing can include using a stainless steel container, with an agitator, and if possible, the
20 container is blanketed with nitrogen during the blending operation. The second oil (substantially no LC PUFA) is typically added first with agitation at a speed that does not create vortices until a steady state is attained. The first oil (with LC PUFA) is then added until completely mixed in. Agitation preferably can continue for about 3 to 5 minutes (times may vary for different sized containers) until a mixture, homogeneous in
25 appearance, is obtained. Synthetic antioxidants can be added to enhance dispersion and dissolution of the first and second oils, for example, at the start of agitation. In the case of continuous mixing, the first oil (typically, lower volume) can be added at mixing point, depending on the in-line mixing equipment used, and will be added at a rate to produce a desired composition. Synthetic antioxidants can be pre-dissolved in the
30 second oil, preferentially, but can also be introduced into the first oil assuming the desired amounts of antioxidants can be dissolved completely.

The food oil compositions and food products comprising the food oil compositions of the present invention can have an LC PUFA content such that an

individual serving of a food product comprising the food oil compositions of the present invention has an appropriate amount of LC PUFA per serving. Appropriate amounts of LC PUFAs and preferably, omega-3 LC PUFA and/or omega-6 LC PUFA per serving are known in the art. For example, preferred amounts of omega-3 LC PUFA and/or
5 omega-6 LC PUFA per serving include amounts of omega-3 LC PUFA and/or omega-6 LC PUFA between about 5 mg per serving and about 150 mg per serving; between about 10 mg per serving and about 100 mg per serving; between about 25 mg per serving and about 75 mg per serving; and between about 35 mg per serving and about 50 mg per serving.

10 The final concentration of LC PUFA in the blend can vary depending on the use or purpose of the oil and the amount of LC PUFA desired per serving. For example, a food product comprising a significant percentage by weight of oil, thus resulting in a relatively greater amount of oil per serving, will require an oil composition that has a relatively smaller percentage of LC PUFA. Knowing the approximate amount of LC
15 PUFA desired per serving and the amount of oil per serving, the skilled artisan can make the necessary calculations to determine the appropriate percentage of LC PUFA in the oil blend.

In the first embodiment, such as where the oil blend is to be used as a fast frying oil, the blend can have an LC PUFA and preferably, an omega-3 LC PUFA and/or
20 omega-6 LC PUFA content in an amount between about 0.01% and about 5%, between about 0.8% and about 3%, and between about 0.1% and about 0.5%. When the oil blend is to be used in food products, for example, as an edible lipid-containing food sauce, preferably, the blend can have an LC PUFA and preferably, an omega-3 LC PUFA and/or omega-6 LC PUFA content in an amount between about 1% and about 30%,
25 between about 10% and about 20%, and between about 1% and about 5%. When the oil blend is to be used in food products, for example, in an embodiment wherein the oil blend is sprayed onto a food product as a topical oil, the blend can have an LC PUFA and preferably, an omega-3 LC PUFA and/or omega-6 LC PUFA between about 0.25% and about 10% and between about 1% and about 5%.

30 In preferred embodiments, the blend comprises oil components that are not LC PUFAs and specifically, not omega-3 LC PUFA nor omega-6 LC PUFA having 20 or more carbons in an amount of at least about 70%, at least about 80%, at least about 90%, and at least about 95%.

In preferred embodiments, the food oil compositions and food products of the present invention comprise an antioxidant, and methods for food preparation comprise the addition of an antioxidant. In particular, in the embodiment of a topical food oil composition of the present invention, an antioxidant is part of the composition. In other
5 embodiments, antioxidants can be used, but are optional. Any antioxidant suitable for food oils or fats preservation known in the art is compatible with the present invention, and include vitamin E, butylhydroxytoluene (BHT), butylhydroxyanisole (BHA), tert-butylhydroquinone (TBHQ), propyl gallate (PG), vitamin C (as used herein, reference to
10 vitamin C includes derivatives thereof), phospholipids, and natural antioxidants such as rosemary extract, and combinations thereof. Preferred antioxidants include BHA, BHT, TBHQ, a blend of BHA/BHT, and combinations thereof, and particularly, TBHQ. Amounts of antioxidant to include in the composition will vary depending on the application as determined by one skilled in the art. For example, food oil compositions of the present invention comprising relatively greater amounts of LC PUFAs and
15 preferably, omega-3 LC PUFAs and/or omega-6 LC PUFAs (having 20 or more carbons) preferably contain higher amounts of antioxidant, such as, for example, amounts up to the maximum allowed by current United States law. Antioxidants may be added to or blended with the oil by any method known in the art. Preferred amounts of antioxidant in the oil compositions of the present invention include amounts between
20 about 0.01% and about 1%, and between about 0.1% and about 0.5%.

In preferred embodiments, the food oil compositions and food products of the present invention are stored under appropriate conditions to minimize oxidative degradation. Many methods to effect such storage conditions are known in the art and are suitable for use with the present invention, such as, for example, replacement of
25 ambient air with an inert gas atmosphere. A preferred method by which to reduce or minimize oxidative degradation is to store food oil compositions and food products under a nitrogen (N₂) atmosphere or mixed nitrogen and carbon dioxide atmosphere. Preferably, packaged food oil compositions and food products are packaged under nitrogen. Methods for producing a nitrogen gas atmosphere into a food container are
30 known in the art.

In another embodiment, the present invention includes a method for food preparation for a food item capable of being skillet-fried, comprising placing the food item and an oil blend of the present invention onto the skillet, and applying heat to the

skillet sufficient to heat the food item. Suitable food items include any food item that is capable of being skillet-fried with an oil, and includes, for example, meats, eggs (e.g., omelets), fish, vegetables, starchy tubers such as potatoes, rice, doughs, batters, breads, batter-coated breads (e.g., French toast), corn products, and mixtures of the foregoing.

5 The term "skillet" refers to any cooking utensil that is suitable for heating food items, and more particularly refers to a wide metal or tempered-glass vessel. A suitable proportion of food and oil for use in the invention may be determined by one skilled in the art. This embodiment additionally includes a skillet-fried food product comprising an oil blend of the present invention.

10 In another embodiment, the present invention includes a method for preparing an oil blend of the present invention, wherein the oil blend is contacted with other food components to make a variety of products such as any edible lipid-containing food sauce, such as salad dressings, marinades, remoulades, vegetable sauces, fruit sauces, fish sauces, and meat sauces, such as poultry sauces, beef sauces, veal sauces, and lamb
15 sauces. This method includes mixing a first oil comprising an LC PUFA and preferably, an omega-3 LC PUFA and/or omega-6 LC PUFA with additional components conventionally found in those products such as spices, flavorings, thickeners, and emulsifiers. Suitable recipes and methods of combining the first oil and additional components are known in the art.

20 In another embodiment, the present invention includes a method for enhancing the LC PUFA and preferably, the omega-3 PUFA and/or omega-6 LC PUFA content of a food product, comprising applying an oil blend of the present invention to the food product. By this method, the LC PUFA content of the food product is enhanced, without
25 subjecting the LC PUFAs to harsh thermal processes during cooking. Such a method can produce food products having a shelf life of approximately 6 months or more. A preferred food product is a previously cooked food product. Preferred previously cooked food products include food products that were previously baked, fried, or deep-fried. The oil blend of the present invention may be applied to the food product by any method
30 known in the art, such as spraying the food product with the oil, dipping the food product into the oil, and brushing the oil onto the surface of the food product. Preferably, the oil is sprayed onto the surface of the food product. Preferred food products include baked goods such as cookies, crackers, sweet goods, muffins, cereals, snack cakes, pies, granola/snack bars, and toaster pastries; salted snacks such as potato chips, corn chips,

tortilla chips, extruded snacks, popcorn, pretzels, potato crisps, and nuts; specialty snacks such as dried fruit snacks, meat snacks, pork rinds, health food bars, rice cakes and corn cakes; confectionary snacks such as candy; and naturally occurring snack foods such as nuts, dried fruits and vegetables. Preferred food products include cookies, crackers, 5 potato chips, corn chips, wheat chips, sorghum chips, soy chips and nuts. This embodiment additionally includes a previously cooked food product comprising an oil blend of the present invention.

In another embodiment, the present invention includes a method for enhancing the PUFA content of a food product, comprising applying an oil blend of the present 10 invention to a food product intended for consumption by infants or toddlers. For instance, snack foods containing ARA are suitable for consumption by children that are still consuming infant formula, but who are also starting to eat solid foods. In some of these embodiments, ratios of DHA:ARA in oils of the present invention are from about 1:0.5 to about 1:5. Additional ratios are at about 1:1.5, at about 1:2 and at about 1:3.

15 The following examples and test results are provided for the purposes of illustration and are not intended to limit the scope of the invention.

EXAMPLES

Example 1

20 The example illustrates an embodiment of the present invention in which a blend of oils is used for frying various foods.

800 g of a blend of oils was prepared by mixing 799.2 g of commercially available corn oil with 0.8 g of DHASCO[®]-S oil (Martek Biosciences Corporation, Columbia, MD). DHASCO[®]-S comprises approximately 35% by weight DHA, resulting in an omega-3 content of about 0.035%. Fried potatoes (French fried style), omelets and fried 25 French toast were prepared using this oil blend and tested for consumer acceptability by a consumer panel of nine or twelve people. The oil blend was stored for one month at room temperature and then was re-tested preparing the same foods as before. The results of the consumer testing are shown in Fig. 1. The amount of DHA per serving of food product, as well as the oil before and after deep frying, was analyzed and is shown below 30 in Table 1.

Table 1

Sample	DHA mg/serving
French fry (100 g serving)	4.6
French toast (50 g serving)	10.9
Egg (100 g serving)	56
Oil Before Frying (1 g)	0.5
Oil After Frying (1 g)	0.5

From the results in Fig. 1, it can be seen that the French toast and omelet had 100% or almost 100% "Like" response in the consumer acceptability testing both at time zero and after one month, indicating that after one month, the oil was still in excellent condition. In the case of the French fries, although >50% provided a "Dislike" response after one month, the majority of the comments were neutral (e.g., related to an oily flavor or a different flavor), with <50% of the total attributing the "Dislike" rating to a negative comment (e.g., fishiness). Therefore, the "Dislike" ratings can be attributed to the corn oil aging rather than the omega-3 portion of the oil being degraded and providing off flavors.

Example 2

This example examines the effect on the oxidative stability of an oil containing an omega-3 LC PUFA of blending a vegetable oil containing substantially no omega-3 LC PUFA and substantially no omega-6 LC PUFA.

An oil containing about 35% by weight DHA (DHASCO[®]-S, Martek Biosciences Corporation, Columbia, MD) was diluted with 20% of various vegetable oils and 30% corn oil as shown in Fig. 2. The DHASCO[®]-S oil and the blended oils were tested for the time to the OSI induction period, measured in hours. The oils were kept at 80°C with air bubbled through and evaluated for the time until the oil begins to oxidize.

The results of this testing are shown in Fig. 2 in which it is shown that at 20% corn oil and soybean oil and 30% corn oil, an increase in the OSI induction period was achieved.

Example 3

This example examines the effect on the oxidative stability of an oil containing an omega-3 LC PUFA of blending corn oil, with and without added antioxidants.

5 An oil containing about 32% by weight DHA (DHA-HM, Martek Biosciences Corporation, Columbia, MD) was diluted with 30% or 40% of corn oil, with and without the addition of 400ppm or 600ppm of an antioxidant blend of ascorbyl palmitate and tocopherols (Grindox™, Danisco) as shown in Fig. 3. The DHA-HM oil and the corn oil blends were tested for the time to the OSI induction period, measured in hours. The oils
10 were kept at 80°C with air bubbled through and evaluated for the time until the oil begins to oxidize.

The results of this testing are shown in Fig. 3 in which it is shown that all of the corn oil blends, with and without antioxidant increased the OSI induction period.

Example 4

15 This example examines the effect on the oxidative stability of an oil containing an omega-3 LC PUFA of blending corn oil, with and without added antioxidants.

An oil containing about 32% by weight DHA (DHA-HM, Martek Biosciences Corporation, Columbia, MD) was diluted with 30% or 40% of corn oil, with and without the addition of 400ppm or 600ppm of an antioxidant blend of ascorbyl palmitate and
20 tocopherols (Grindox™, Danisco) as shown in Figs. 4 and 5. The DHA-HM oil and the corn oil blends were stored at 40°C over a period of weeks and tested for the production of peroxides (primary oxidation products) and alkenals (secondary oxidation products).

The results of this testing are shown in Figs. 4 and 5 in which it is shown that all of the corn oil blends, with and without antioxidant, delayed the occurrence of primary
25 and secondary oxidation products.

Example 5

This example examines the effect on the oxidative stability of an oil containing an omega-3 LC PUFA of blending vegetable oils containing substantially no omega-3 LC PUFA and substantially no omega-6 LC PUFA.

30 Five commonly used vegetable oils were combined with an oil containing about 35% by weight DHA (DHASCO®-S, Martek Biosciences Corporation, Columbia, MD) at five dilution levels. The DHASCO®-S oil, the vegetable oil and the blended oils were tested for the time to the OSI induction period, measured in hours. The oils were kept at

80°C with air bubbled through and evaluated for the time until the oil begins to oxidize. The oils, dilution levels and OSI values are shown below in Tables 2-6. The results are also shown in Figs. 6-10.

5

Oil	OSI Value
100.00% Corn Oil	99.9
99.99 % Corn Oil	87.5
99.75% Corn Oil	87.5
95.00% Corn Oil	59.3
90.00% Corn Oil	53.15
70.00% Corn Oil	41.025
100.00% DHASCO [®] -S	36.2

5

Oil	OSI Value
100.00% Soy Oil	36.9
99.99% Soy Oil	37.225
99.75% Soy Oil	38.725
95.00% Soy Oil	38.7
90.00% Soy Oil	23.25
70.00% Soy Oil	19.3
100.00% DHASCO [®] -S	36.2

10

Oil	OSI Value
100.00% Canola Oil	67.4
99.99% Canola Oil	67.95
99.75% Canola Oil	66.275
95.00% Canola Oil	56.075
90.00% Canola Oil	50.05
70.00% Canola Oil	39.725
100.00% DHASCO [®] -S	36.2

10

Oil	OSI Value
100.00% Safflower Oil	28.5
99.99% Safflower Oil	28.95
99.75% Safflower Oil	29.75
95.00% Safflower Oil	24.925
90.00% Safflower Oil	22.8
70.00% Safflower Oil	19.35
100.00% DHASCO [®] -S	36.2

10

Oil	OSI Value
100.00% Sunflower Oil	48.6
99.99% Sunflower Oil	44.9
99.75% Sunflower Oil	45.35
95.00% Sunflower Oil	34.4
90.00% Sunflower Oil	27.45

70.00% Sunflower Oil	20.6
100.00% DHASCO®-S	36.2

The results show that an increase in the OSI induction period can be achieved with blends of DHASCO®-S and vegetable oils.

5 Example 6

This example examines the sensory qualities of corn oil containing 0.5% w/w DHASCO®-S oil.

Sensory qualities of the blended oil composition were determined. The oil was then stored in a metal container at room temperature for six months and the sensory
 10 qualities of the oil were again determined. Characteristics were determined on a scale of 1-15, with 15 being the worst. Results are shown in Table 7.

Table 7. Sensory Results

Attributes	Time	
	T=0	T=6 months
AROMA		
Total Impact	3.5	4
Green/Beany	2.5	1.5
Fishy	0	0
Painty	0	2
Herbal	0	0
Other	1	2
AROMATICS		
Total Impact	4.5	5.5
Green/Beany	1	1
Fishy	0	0
Painty	4 (painty/other)*	4 (painty/other)*
Herbal	0	0
Other		
Aftertaste		painty

15

* This is a characteristic of rancid corn oil, and is not related to DHA. This result indicates some inherent problem with the quality of the corn oil.

These results indicate that the sensory characteristic of oil blends of the present invention remain stable over storage times.

Example 7

5 Corn oil containing 0.5% w/w DHASCO[®]-S oil from Example 6 was analyzed for peroxide content, alkenal content, and DHA level. Peroxide content is a measure of oxidation of the oil. Peroxide content is shown in Figure 11. After eight months, there is no increase in the amount of peroxides in the oil. There is therefore no apparent oxidation of the oil. Alkenals are secondary products of oxidation. Alkenal content is shown in Figure 12. There is no increase in the amount of alkenals in the oil, over time. Secondary products of oxidation typically increase as oxidation progresses or remain constant if there is no oxidation. DHA levels did not decrease over time, as shown in Figure 13.

15 Example 8

This example evaluates the use of corn oil containing 0.5% w/w DHASCO[®]-S oil from Example 6 that had been stored for six months to prepare French toast, French fries, and scrambled eggs.

20 Once the foods were cooked they were evaluated by a small taste panel. No off notes were detected in any of the foods.

 Cooked foods were also evaluated for DHA content, to determine the amount of DHA transferred to the food from the oil. Samples of each food were freeze dried in preparation for analysis. Once dry, the samples were ground to a fine powder. DHA was determined by FAME analysis. Duplicate analyses were performed for each sample using a three-point internal standard (C23:0) calibration curve to quantitate DHA. The DHA results are summarized in Table 8. The eggs used to prepare the scrambled eggs naturally contained between 10 and 20 mg DHA per serving, and therefore the difference between 66.4 mg and the naturally occurring amount of 10-20 mg of DHA is due to DHA transfer from the fortified corn oil.

30

Table 8: DHA content of various foods cooked in DHA-fortified corn oil.

Sample Description	Solids (%)	DHA (mg DHA Free Fatty Acids/g food)	Fried Food DHA (mg DHA Free Fatty Acids/serving) ²
French toast	54.61	0.96	57.1
French toast	44.02	0.41	24.9
Scrambled eggs	35.22	1.89	66.4

¹ Reported on a dry weight basis

² Reported on an "as received" basis

5 The principles, preferred embodiments and modes of operation of the present invention have been described in the foregoing specification. The invention which is intended to be protected herein should not, however, be construed as limited to the particular forms disclosed, as these are to be regarded as illustrative rather than restrictive. Variations and changes may be made by
10 those skilled in the art without departing from the spirit of the present invention. Accordingly, the foregoing best mode of carrying out the invention should be considered exemplary in nature and not as limiting to the scope and spirit of the invention as set forth in the appended claims.

15 The above discussion of documents, acts, materials, devices, articles and the like is included in the specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.

20 It is to be understood that, throughout the description and claims of the specification the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

The claims defining the invention are as follows:

1. A food oil composition comprising a blend of:
a first oil comprising an LC PUFA, and
a second oil comprising substantially no LC PUFA and that is liquid at
5 room temperature,
wherein the blend comprises about 0.01% to 0.5% LC PUFA.
2. A topical food oil composition comprising a blend of a first oil comprising
an LC PUFA, a second oil comprising substantially no LC PUFA and that is
liquid at room temperature, and an antioxidant, wherein the LC PUFA
10 comprises about 0.01% to 0.5% of the food oil composition.
3. The composition according to claim 1 or claim 2, wherein the blend
comprises about 0.1% to 0.5% LC PUFAs.
4. The composition according to claim 1 or claim 3, further comprising an
antioxidant.
- 15 5. The composition according to claim 2 or claim 4, wherein the antioxidant is
selected from the group consisting of vitamin E, butylhydroxytoluene
(BHT), butylhydroxyanisole (BHA), tertbutylhydroquinone (TBHQ), propyl
gallate (PG), vitamin C, phospholipids, and natural antioxidants, and
combinations thereof.
- 20 6. The composition according to claim 4 or claim 5, wherein the antioxidant is
present in the oil blend in an amount of about 0.01% to about 1%.
7. The composition according to any one of claims 4 to 6, wherein the
antioxidant is present in the oil blend in an amount of about 0.1% to about
0.5%.
- 25 8. The composition according to any one of claims 1 to 7, wherein the second
oil is selected from the group consisting of borage oil, black currant seed
oil, corn oil, coconut oil, canola oil, soybean oil, safflower oil, high oleic
safflower oil, sunflower oil, high oleic sunflower oil, olive oil, evening
primrose oil, cottonseed oil, rice bran oil, grapeseed oil, flaxseed oil, garlic
30 oil, peanut oil, almond oil, walnut oil, wheat germ oil, sesame oil, animal

- fat, animal oil, marine fat, marine oil, microbial oil, a hydrogenated oil of any of the foregoing, and mixtures of the foregoing.
9. The composition according to any one of claims 1 to 8, wherein the LC PUFA is selected from the group consisting of an omega-3 LC PUFA, an omega-6 LC PUFA, and mixtures thereof.
10. The composition according to claim 9, wherein the LC PUFA is selected from the group consisting of docosahexaenoic acid, eicosapentaenoic acid, docosapentaenoic acid, and arachidonic acid.
11. The composition according to any one of claims 1 to 10, wherein the first oil is from a microbial source.
12. The composition according to claim 11, wherein the microbial source comprises a microorganism selected from the group consisting of algae, protists, bacteria and fungi.
13. The composition according to claim 11, wherein the microbial source is an oleaginous microorganism.
14. The composition according to claim 12, wherein the microbial source is a microorganism selected from the group consisting of microorganisms of the genus *Thraustochytrium*, microorganisms of the genus *Schizochytrium*, microorganisms of the genus *Altbornia*, microorganisms of the genus *Aplanochytrium*, microorganisms of the genus *Japonochytrium*, microorganisms of the genus *Elina*, microorganisms of the genus *Cryptobcodinium*, microorganisms of the genus *Mortierella*, and mixtures thereof.
15. The composition according to claim 12, wherein the microorganism is selected from the group consisting of microorganisms of the genus *Schizochytrium*, microorganisms of the genus *Cryptobcodinium*, microorganisms of the genus *Mortierella*, and mixtures thereof.
16. The composition according to any one of claims 1 to 10, wherein the first oil is from a plant source.
17. The composition according to claim 16, wherein the plant source has been genetically modified to produce long chain polyunsaturated fatty acids,

wherein the plant is selected from the group consisting of soybean, corn, safflower, sunflower, canola, flax, peanut, mustard, rapeseed, chickpea, cotton, lentil, white clover, olive, palm, borage, evening primrose, linseed and tobacco.

- 5 18. The composition according to any one of claims 1 to 10, wherein the first oil is from an animal source.
19. The composition according to claim 18, wherein the animal source is selected from the group consisting of aquatic animals, animal tissues and animal products.
- 10 20. The composition according to any one of claims 1 to 19, wherein the first oil comprises at least about 20% omega-3 LC PUFAs, omega-6 LC PUFAs, or mixtures thereof.
21. The composition according to any one of claims 1 to 20, wherein the first oil comprises at least about 60% omega-3 LC PUFAs, omega-6 LC PUFAs, or mixtures thereof.
- 15 22. The composition according to any one of claims 1 to 21, wherein the blend is stable to oxidation for at least about three months when stored at room temperature; the LCPUFA level is stable for at least about three months when stored at room temperature; and the sensory characteristics of the composition remain constant for at least about three months when stored at room temperature.
- 20 23. A food product comprising the composition according to any one of claims 1 to 22.
24. The food product according to claim 23, wherein the food product comprises about 5 mg to about 150 mg omega-3 LC PUFA and/or omega-6 LC PUFA.
- 25 25. The food product according to claim 23, wherein the food product is selected from the group consisting of salad dressings, marinades, remoulades, vegetable sauces, fruit sauces, fish sauces and meat sauces.
- 30 26. The food product according to claim 23, wherein the food product comprises about 5 mg omega-3 LC PUFA, omega-6 LC PUFA, or mixtures

- thereof, to about 150 mg omega-3 LC PUFA, omega-6 LC PUFA, or mixtures thereof.
27. The food product according to claim 23, wherein the food product comprises about 10 mg omega-3 LC PUFA, omega-6 LC PUFA, or mixtures thereof, to about 100 mg omega-3 LC PUFA, omega-6 LC PUFA, or mixtures thereof.
28. The food product according to claim 23, wherein the food product is a previously cooked food product.
29. The food product according to claim 28, wherein the previously cooked food product is selected from the group consisting of a previously baked food product, a previously fried food product, and a previously deep-fried food product.
30. The food product according to claim 23, wherein the food product is selected from the group consisting of baked goods, salted snacks, specialty snacks, confectionery snacks, and naturally occurring snack foods.
31. A method of food preparation for a food item capable of being skillet-fried, comprising:
- (a) placing the food item and an oil onto the skillet; and
 - (b) applying heat to the skillet sufficient to heat the food item, wherein the oil comprises a composition according to any one of claims 1 to 22.
32. A method for preparing a food product comprising: contacting an oil with at least one food component, wherein the oil comprises a composition according to any one of claims 1 to 22.
33. The method according to claim 32, wherein the food product is selected from the group consisting of salad dressings, marinades, remoulades, vegetable sauces, fruit sauces, fish sauces and meat sauces.
34. The method according to claim 32, wherein the food product is selected from the group consisting of cookies, crackers, sweet goods, muffins, cereals, snack cakes, pies, granola/snack bars, toaster pastries, potato chips, corn chips, wheat chips, sorghum chips, soy chips, extruded snacks,

popcorn, pretzels, potato crisps, dried fruit snacks, meat snacks, pork rinds, health food bars, rice cakes, corn cakes, candy, nuts, dried fruits and vegetables.

- 5 35. The method according to claim 32, further comprising packaging the food product.
36. The method according to claim 35, wherein the packaging comprises packaging the food product in an inert atmosphere, wherein the atmosphere is selected from the group consisting of:
- 10 an atmosphere comprising nitrogen; and
an atmosphere comprising nitrogen and carbon dioxide.
37. A method of food preparation, comprising topically applying a composition according to any one of claims 1 to 22 to a food product, wherein the topically applying is selected from the group consisting of spraying, dipping and brushing.
- 15 38. A method of food preparation for a food item capable of being deep-fried, comprising:
- (a) immersing the food item in an oil; and
- (b) applying heat to the oil sufficient to heat the food item, wherein the oil comprises a composition according to any one of claims 1 to 22.
- 20 39. A food oil composition according to any one of claims 1 to 22, or a topical food oil composition according to claim 2, or a food product according to any one of claims 23 to 30, or a method of food preparation according to any one of claims 31 to 38, substantially as hereinbefore described and with reference to any one of the examples.




Fig. 1




Fig. 2

3/13




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig.8




Fig. 9




Fig. 10

11/13




Fig. 11

12/13




Fig. 12

13/13




Fig. 13