



AU9468278

---

**(12) PATENT ABRIDGMENT (11) Document No. AU-B-68278/94**  
**(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 691810**

---

- (54) Title  
**METHODS FOR TREATING MUSCLE DISEASES AND DISORDERS**
- International Patent Classification(s)  
(51)<sup>5</sup> **A61K 037/10 A61K 037/36**  
(51)<sup>6</sup> **G01N 033/53 G01N 033/68**
- (21) Application No. : **68278/94** (22) Application Date : **06.05.94**
- (87) PCT Publication Number : **WO94/26298**
- (30) Priority Data
- | (31) Number   | (32) Date       | (33) Country                       |
|---------------|-----------------|------------------------------------|
| <b>059022</b> | <b>06.05.93</b> | <b>US UNITED STATES OF AMERICA</b> |
| <b>209204</b> | <b>08.03.94</b> | <b>US UNITED STATES OF AMERICA</b> |
- (43) Publication Date : **12.12.94**
- (44) Publication Date of Accepted Application : **28.05.98**
- (71) Applicant(s)  
**CAMBRIDGE NEUROSCIENCE**
- (72) Inventor(s)  
**ROBERT SKLAR; MARK MARCHIONNI; DAVID I GWYNNE**
- (74) Attorney or Agent  
**DAVIES COLLISON CAVE , 1 Little Collins Street, MELBOURNE VIC 3000**
- (56) Prior Art Documents  
**AU 47694/93**
- (57) **For treatment of smooth muscle disease, Becker's dystrophy, cardiac muscle disorder arterial sclerosis, vascular lesion, acetylcholine receptor insufficiency.**
- Claim

1. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide encoded by pGGF2HBS11, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).

38. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 35 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the rat I-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

68278/94

**CORRECTED  
VERSION\***

**PCT**



pages 86-165, description, replaced by new pages 86-152; pages 166-180, claims, renumbered as pages 153-167; pages 1/57-11/57, 20/57, 26/57-27/57 and 55/57, drawings, replaced by new pages 1/55-9/55, 18/55, 24/55-25/55 and 53/55; pages 12/57-19/57, 21/57-25/57, 28/57-54/57, 56/57 and 57/57, renumbered as pages 10/55-17/55, 19/55-23/55, 26/55-52/55, 54/55 and 55/55; due to late

INTERNATIONAL APP transmittal by the receiving Office **RATION TREATY (PCT)**

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 37/10, 37/36</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/26298</b>
			<b>(43) International Publication Date:</b> 24 November 1994 (24.11.94)
<b>(21) International Application Number:</b> PCT/US94/05083		<b>(81) Designated States:</b> AU, CA, JP, KR, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
<b>(22) International Filing Date:</b> 6 May 1994 (06.05.94)		<b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(30) Priority Data:</b> 08/059,022      6 May 1993 (06.05.93)      US 08/209,204      8 March 1994 (08.03.94)      US			
<b>(71) Applicant (for all designated States except US):</b> CAMBRIDGE NEUROSCIENCE [US/US]; Building 700, One Kendall Square, Cambridge, MA 02139 (US).			
<b>(72) Inventors; and</b>			
<b>(75) Inventors/Applicants (for US only):</b> SKLAR, Robert [US/US]; 36 Spiers Road, Newton, MA 02159 (US). MARCHIONNI, Mark [US/US]; 24 Twin Circle Drive, Arlington, MA 02174 (US). GWYNNE, David, I. [GB/US]; 77 Grover Street, Beverly, MA 01915 (US).			
<b>(74) Agent:</b> CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110 (US).			
<b>(54) Title:</b> METHODS FOR TREATING MUSCLE DISEASES AND DISORDERS			
<b>(57) Abstract</b>  The invention relates to methods of treating diseases and disorders of the muscle tissues in a vertebrate by the administration of compounds which bind the p185 <sup>erbB2</sup> receptor. These compounds are found to cause increased differentiation and survival of cardiac, skeletal and smooth muscle.			

69 18 10

AUSTRALIAN INDUSTRIAL  
26 AUG 1996  
PROPERTY ORGANISATION

\* (Referred to in PCT Gazette No.37/1996, Section II)

**METHODS FOR TREATING MUSCLE DISEASES AND DISORDERS****Background of the Invention**

The invention relates to prophylactic or affirmative  
5 treatment of diseases and disorders of the musculature by  
administering polypeptides found in vertebrate species,  
which polypeptides are growth, differentiation and survival  
factors for muscle cells.

Muscle tissue in adult vertebrates will regenerate  
10 from reserve myoblasts called satellite cells. Satellite  
cells are distributed throughout muscle tissue and are  
mitotically quiescent in the absence of injury or disease.  
Following muscle injury or during recovery from disease,  
satellite cells will reenter the cell cycle, proliferate and  
15 1) enter existing muscle fibers or 2) undergo  
differentiation into multinucleate myotubes which form new  
muscle fiber. The myoblasts ultimately yield replacement  
muscle fibers or fuse into existing muscle fibers, thereby  
increasing fiber girth by the synthesis of contractile  
20 apparatus components. This process is illustrated, for  
example, by the nearly complete regeneration which occurs in  
mammals following induced muscle fiber degeneration; the  
muscle progenitor cells proliferate and fuse together  
regenerating muscle fibers.

25 Several growth factors which regulate the  
proliferation and differentiation of adult (and embryonic)  
myoblasts in vitro have been identified. Fibroblast growth  
factor (FGF) is mitogenic for muscle cells and is an  
inhibitor of muscle differentiation. Transforming growth  
30 factor  $\beta$  (TGF $\beta$ ) has no effect on myoblast proliferation, but  
is an inhibitor of muscle differentiation. Insulin-like  
growth factors (IGFs) have been shown to stimulate both  
myoblast proliferation and differentiation in rodents.  
Platelet derived growth factor (PDGF) is also mitogenic for  
35 myoblasts and is a potent inhibitor of muscle cell

- 2 -

differentiation see: Florini and Magri, 1989:256:C701-C711).

In vertebrate species both muscle tissue and neurons are potential sources of factors which stimulate myoblast proliferation and differentiation. In diseases affecting the neuromuscular system which are neural in origin (i.e., neurogenic), the muscle tissue innervated by the affected nerve becomes paralyzed and wastes progressively. During peripheral nerve regeneration and recovery from neurologic and myopathic disease, neurons may provide a source of growth factors which elicit the muscle regeneration described above and provide a mechanism for muscle recovery from wasting and atrophy.

A recently described family of growth factors, the neuregulins, are synthesized by motor neurons (Marchionni et al. *Nature* 362:313, 1993) and inflammatory cells (Tarakhovsky et al., *Oncogene* 6:2187-2196 (1991)). The neuregulins and related p185<sup>erbB2</sup> binding factors have been purified, cloned and expressed (Benveniste et al., *PNAS* 82:3930-3934, 1985; Kimura et al., *Nature* 348:257-260, 1990; Davis and Stroobant, *J. Cell. Biol.* 110:1353-1360, 1990; Wen et al., *Cell* 69:559, 1992; Yarden and Ullrich, *Ann. Rev. Biochem.* 57:443, 1988; Holmes et al., *Science* 256:1205, 1992; Dobashi et al., *Proc. Natl. Acad. Sci.* 88:8582, 1991; Lupu et al., *Proc. Natl. Acad. Sci.* 89:2287, 1992). Recombinant neuregulins have been shown to be mitogenic for peripheral glia (Marchionni et al., *Nature* 362:313, 1993) and have been shown to influence the formation of the neuromuscular junction (Falls et al., *Cell* 72:801, 1993). Thus the regenerating neuron and the inflammatory cells associated with the recovery from neurogenic disease and nerve injury provide a source of factors which coordinate the remyelination of motor neurons and their ability to form the appropriate connection with their target. After



- 3 -

muscle has been reinnervated the motor neuron may provide factors to muscle, stimulating muscle growth and survival.

5 Currently, there is no useful therapy for the promotion of muscle differentiation and survival. Such a therapy would be useful for treatment of a variety of neural and muscular diseases and disorders.

#### Summary of the Invention

10 We have discovered that increased mitogenesis differentiation and survival of muscle cells may be achieved using proteins heretofore described as glial growth factors, acetylcholine receptor inducing activity (ARIA), heregulins, neu differentiation factor, and, more generally, neuregulins. We have discovered that these  
15 compounds are capable of inducing both the proliferation of muscle cells and the differentiation and survival of myotubes. These phenomena may occur in cardiac and smooth muscle tissues in addition to skeletal muscle tissues. Thus, the above compounds, regulatory compounds  
20 which induce synthesis of these compounds, and small molecules which mimic these compounds by binding to the receptors on muscle or by stimulating through other means the second messenger systems activated by the ligand-receptor complex are all extremely useful as prophylactic  
25 and affirmative therapies for muscle diseases.

A novel aspect of the invention involves the use of the above named proteins as growth factors to induce the mitogenesis, survival, growth and differentiation of muscle cells. Treating of the muscle cells to achieve  
30 these effects may be achieved by contacting muscle cells with a polypeptide described herein. The treatments may be provided to slow or halt net muscle loss or to

- 4 -

increase the amount or quality of muscle present in the vertebrate.

These factors may be used to produce muscle cell mitogenesis, differentiation, and survival in a vertebrate (preferably a mammal, more preferably a human) by administering to the vertebrate an effective amount of a polypeptide or a related compound. Neuregulin effects on muscle may occur, for example, by causing an increase in muscle performance by inducing the synthesis of particular isoforms of the contractile apparatus such as the myosin heavy chain slow and fast isoforms; by promoting muscle fiber survival via the induction of synthesis of protective molecules such as, but not limited to, dystrophin; and/or by increasing muscle innervation by, for example, increasing acetylcholine receptor molecules at the neuromuscular junction.

The term muscle cell as used herein refers to any cell which contributes to muscle tissue. Myoblasts, satellite cells, myotubes, and myofibril tissues are all included in the term "muscle cells" and may all be treated using the methods of the invention. Muscle cell effects may be induced within skeletal, cardiac and smooth muscles.

Mitogenesis may be induced in muscle cells, including myoblasts or satellite cells, of skeletal muscle, smooth muscle or cardiac muscle. Mitogenesis as used herein refers to any cell division which results in the production of new muscle cells in the patient. More specifically, mitogenesis in vitro is defined as an increase in mitotic index relative to untreated cells of 50%, more preferably 100%, and most preferably 300%, when the cells are exposed to labelling agent for a time equivalent to two doubling times. The mitotic index is the fraction of cells in the culture which have labelled

- 5 -

nuclei when grown in the presence of a tracer which only incorporates during S phase (i.e., BrdU) and the doubling time is defined as the average time required for the number of cells in the culture to increase by a factor of two).

An effect on mitogenesis in vivo is defined as an increase in satellite cell activation as measured by the appearance of labelled satellite cells in the muscle tissue of a mammal exposed to a tracer which only incorporates during S phase (i.e., BrdU). In useful therapeutic is defined in vivo as a compound which increases satellite cell activation relative to a control mammal by at least 10%, more preferably by at least 50%, and most preferably by more than 200% when the mammal is exposed to labelling agent for a period of greater than 15 minutes and tissues are assayed between 10 hours and 24 hours after administration of the mitogen at the therapeutic dose. Alternatively, satellite cell activation in vivo may be detected by monitoring the appearance of the intermediate filament vimentin by immunological or RNA analysis methods. When vimentin is assayed, the useful mitogen is defined as one which causes expression of detectable levels of vimentin in the muscle tissue when the therapeutically useful dosage is provided.

Myogenesis as used herein refers to any fusion of myoblasts to yield myotubes. Most preferably, an effect on myogenesis is defined as an increase in the fusion of myoblasts and the enablement of the muscle differentiation program. The useful myogenic therapeutic is defined as a compound which confers any increase in the fusion index in vitro. More preferably, the compound confers at least a 2.0-fold increase and, most preferably, the compound confers a 3-fold or greater

- 6 -

increase in the fusion index relative to the control. The fusion index is defined as the fraction of nuclei present in multinucleated cells in the culture relative to the total number of nuclei present in the culture.

5 The percentages provided above are for cells assayed after 6 days of exposure to the myogenic compound and are relative to an untreated control. Myogenesis may also be determined by assaying the number of nuclei per area in myotubes or by measurement of the levels of muscle  
10 specific protein by Western analysis. Preferably, the compound confers at least a 2.0-fold increase in the density of myotubes using the assay provided, for example, herein, and, most preferably, the compound confers a 3-fold or greater increase.

15 The growth of muscle may occur by the increase in the fiber size and/or by increasing the number of fibers. The growth of muscle as used herein may be measured by A) an increase in wet weight, B) an increase in protein content, C) an increase in the number of muscle fibers,  
20 or D) an increase in muscle fiber diameter. An increase in growth of a muscle fiber can be defined as an increase in the diameter where the diameter is defined as the minor axis of ellipsis of the cross section. The useful therapeutic is one which increases the wet weight,  
25 protein content and/or diameter by 10% or more, more preferably by more than 50% and most preferably by more than 100% in an animal whose muscles have been previously degenerated by at least 10% and relative to a similarly treated control animal (i.e., an animal with degenerated  
30 muscle tissue which is not treated with the muscle growth compound). A compound which increases growth by increasing the number of muscle fibers is useful as a therapeutic when it increases the number of fibers in the diseased tissue by at least 1%, more preferably at least

- 7 -

20%, and most preferably, by at least 50%. These percentages are determined relative to the basal level in a comparable untreated undiseased mammal or in the contralateral undiseased muscle when the compound is administered and acts locally.

The survival of muscle fibers as used herein refers to the prevention of loss of muscle fibers as evidenced by necrosis or apoptosis or the prevention of other mechanisms of muscle fiber loss. Survival as used herein indicates an decrease in the rate of cell death of at least 10%, more preferably by at least 50%, and most preferably by at least 300% relative to an untreated control. The rate of survival may be measured by counting cells stainable with a dye specific for dead cells (such as propidium iodide) in culture when the cells are 8 days post-differentiation (i.e., 8 days after the media is changed from 20% to 0.5% serum).

Muscle regeneration as used herein refers to the process by which new muscle fibers form from muscle progenitor cells. The useful therapeutic for regeneration confers an increase in the number of new fibers by at least 1%, more preferably by at least 20%, and most preferably by at least 50%, as defined above.

The differentiation of muscle cells as used herein refers to the induction of a muscle developmental program which specifies the components of the muscle fiber such as the contractile apparatus (the myofibril). The therapeutic useful for differentiation increases the quantity of any component of the muscle fiber in the diseased tissue by at least 10% or more, more preferably by 50% or more, and most preferably by more than 100% relative to the equivalent tissue in a similarly treated control animal.

- 8 -

Atrophy of muscle as used herein refers to a significant loss in muscle fiber girth. By significant atrophy is meant a reduction of muscle fiber diameter in diseased, injured or unused muscle tissue of at least 10% relative to undiseased, uninjured, or normally utilized tissue.

Methods for treatment of diseases or disorders using the polypeptides or other compounds described herein are also part of the invention. Examples of muscular disorders which may be treated include skeletal muscle diseases and disorders such as myopathies, dystrophies, myoneural conductive diseases, traumatic muscle injury, and nerve injury. Cardiac muscle pathologies such as cardiomyopathies, ischemic damage, congenital disease, and traumatic injury may also be treated using the methods of the invention, as may smooth muscle diseases and disorders such as arterial sclerosis, vascular lesions, and congenital vascular diseases. For example, Duchennes muscular dystrophy, Beckers' dystrophy, and Myasthenia gravis are but three of the diseases which may be treated using the methods of the invention.

The invention also includes methods for the prophylaxis or treatment of a tumor of muscle cell origin such as rhabdomyosarcoma. These methods include administration of an effective amount of a substance which inhibits the binding of one or more of the polypeptides described herein and inhibiting the proliferation of the cells which contribute to the tumor.

The methods of the invention may also be used to treat a patient suffering from a disease caused by a lack of a neurotrophic factor. By lacking a neurotrophic factor is meant a decreased amount of neurotrophic factor relative to an unaffected individual sufficient to cause

- 9 -

detectable decrease in neuromuscular connections and/or muscular strength. The neurotrophic factor may be present at levels 10% below those observed in unaffected individuals. More preferably, the factor is present at  
5 levels 20% lower than are observed in unaffected individuals, and most preferably the levels are lowered by 80% relative to unaffected individuals under similar circumstances.

The methods of the invention make use of the fact  
10 that the neuregulin proteins are encoded by the same gene. A variety of messenger RNA splicing variants (and their resultant proteins) are derived from this gene and many of these products show binding to P185<sup>erbB2</sup> and activation of the same. Products of this gene have been  
15 used to show muscle cell mitogenic activity (see Examples 1 and 2, below), differentiation (Examples 3 and 6), and survival (Examples 4 and 5). This invention provides a use for all of the known products of the neuregulin gene (described herein and in the references listed above)  
20 which have the stated activities as muscle cell mitogens, differentiation factors, and survival factors. Most preferably, recombinant human GGF2 (rhGGF2) is used in these methods.

The invention also relates to the use of other,  
25 not yet naturally isolated, splicing variants of the neuregulin gene. Fig. 29 shows the known patterns of splicing. These patterns are derived from polymerase chain reaction experiments (on reverse transcribed RNA), analysis of cDNA clones (as presented within), and  
30 analysis of published sequences encoding neuregulins (Peles et al., Cell 69:205 (1992) and Wen et al., Cell 69:559 (1992)). These patterns, as well as additional patterns disclosed herein, represent probable splicing variants which exist. The splicing variants are fully

- 10 -

described in Goodearl et al., USSN 08/036,555, filed March 24, 1993, incorporated herein by reference.

More specifically, cell division, survival, differentiation and growth of muscle cells may be  
 5 achieved by contacting muscle cells with a polypeptide defined by the formula

## WBAZCX

wherein WBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 134, 135,  
 10 137-139, 156); wherein W comprises the polypeptide segment F, or is absent; wherein Z comprises the polypeptide segment G or is absent; and wherein X comprises the polypeptide segment C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL,  
 15 C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL and/or by contacting muscle cells with a polypeptide defined by the formula

20

## YBAZCX

wherein YBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 133-135, 156, 159); wherein Y comprises the polypeptide segment E, or is absent; wherein Z comprises the polypeptide segment G or is absent; and wherein X comprises the polypeptide  
 25 segment C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL,  
 30 or C/D C/D' D' HKL.

Generally, the N-terminus of the above-described polypeptides begins with either the F or E polypeptide



- 11 -

segments. When the F polypeptide is present it may be cleaved upon maturation of the protein to yield the mature polypeptide. When the E sequence is present the first 50 amino acids which represent the N-terminal signal sequence may be absent from the polypeptides.

Furthermore, the invention includes a method of treating muscle cells by the application to the muscle cell of a

-30 kD polypeptide factor isolated from the MDA-MB 231 human breast cell line; or

-35 kD polypeptide factor isolated from the rat I-~~B~~ transformed fibroblast cell line to the glial cell or

-75 kD polypeptide factor isolated from the SKBR-3 human breast cell line; or

-44 kD polypeptide factor isolated from the rat I-EJ transformed fibroblast cell line; or

-25 kD polypeptide factor isolated from activated mouse peritoneal macrophages; or

-45 kD polypeptide factor isolated from the MDA - MB 231 human breast cell; or

-7 to 14 kD polypeptide factor isolated from the ATL-2 human T-cell line to the glial cell; or

-25 kD polypeptide factor isolated from the bovine kidney cells; or

-42 kD ARIA polypeptide factor isolated from brain; -46-47 kD polypeptide factor which stimulates 0-2A glial progenitor cells; or

-43-45 kD polypeptide factor, GGFI11,175

U.S. patent application Serial No. 07/931,041, filed August 17, 1992, incorporated herein by reference.

The invention further includes methods for the use of the EGFL1, EGFL2, EGFL3, EGFL4, EGFL5, and EGFL6 polypeptides, Fig. 37 to 42 and SEQ ID Nos. 150 to 155,

- 12 -

respectively, for the treatment of muscle cells in vivo and in vitro.

Also included in the invention is the administration of the GGF2 polypeptide whose sequence is shown in Fig. 44 for the treatment of muscle cells.

An additional important aspect of the invention are methods for treating muscle cells using:

(a) a basic polypeptide factor also known to have glial cell mitogenic activity, in the presence of fetal calf plasma, a molecular weight of from about 30 kD to about 36 kD, and including within its amino acid sequence any one or more of the following peptide sequences:

	F K G D A H T E	(SEQ ID NO: 1)
	A S L A D E Y E Y M X K	(SEQ ID NO: 2)
15	T E T S S S G L X L K	(SEQ ID NO: 3)
	A S L A D E Y E Y M R K	(SEQ ID NO: 7)
	A G Y F A E X A R	(SEQ ID NO: 11)
	T T E M A S E Q G A	(SEQ ID NO: 13)
	A K E A L A A L K	(SEQ ID NO: 14)
20	F V L Q A K K	(SEQ ID NO: 15)
	E T Q P D P G Q I L K K V P M V I G A Y T	(SEQ ID NO: 165)
	E Y K C L K F K W F K K A T V M	(SEQ ID NO: 17)
	E X K F Y V P	(SEQ ID NO: 19)
25	K L E F L X A K	(SEQ ID NO: 32); and

(b) a basic polypeptide factor for use in treating muscle cells which is also known to stimulate glial cell mitogenesis in the presence of fetal calf plasma, has a molecular weight of from about 55 kD to about 63 kD, and including within its amino acid sequence any one or more of the following peptide sequences:

- 13 -

	V H Q V W A A K	(SEQ ID NO: 33)
	Y I F F M E P E A X S S G	(SEQ ID NO: 34)
	L G A W G P P A F P V X Y	(SEQ ID NO: 35)
	W F V V I E G K	(SEQ ID NO: 36)
5	A S P V S V G S V Q E L Q R	(SEQ ID NO: 37)
	V C L L T V A A L P P T	(SEQ ID NO: 38)
	K V H Q V W A A K	(SEQ ID NO: 48)
	K A S L A D S G E Y M X K	(SEQ ID NO: 49)
	D L L L X V	(SEQ ID NO: 39)

10           Methods for the use of the peptide sequences set  
out above, derived from the smaller molecular weight  
polypeptide factor, and from the larger molecular weight  
polypeptide factor, are also aspects of this invention.  
Monoclonal antibodies to the above peptides are  
15 themselves useful investigative tools and therapeutics.

Thus, the invention further embraces methods of  
using a polypeptide factor having activities useful for  
treating muscle cells and including an amino acid  
sequence encoded by:

- 20           (a) a DNA sequence shown in any one of Figs. 27A,  
27B or 27C, ~~SEQ~~ ID Nos. 129-131, respectively;
- (b) a DNA sequence shown in Fig. 21, SEQ ID No.  
85;
- (c) the DNA sequence represented by nucleotides  
25 281-557 of the sequence shown in Fig. 27A, SEQ ID No.  
129; or
- (d) a DNA ~~sequence~~ hybridizable to any one of the  
DNA sequences according to (a), (b) or (c).

Following factors as muscle cell mitogens:

- 30           (a) a basic polypeptide factor which has, if  
obtained from bovine pituitary material, an observed  
molecular weight, whether in reducing conditions or not,  
of from about 30kD to about 36kD on SDS-polyacrylamide

- 14 -

gel electrophoresis which factor has muscle cell mitogenic activity including stimulating the division of myoblasts, and when isolated using reversed-phase HPLC retains at least 50% of said activity after 10 weeks  
5 incubation in 0.1% trifluoroacetic acid at 4°C; and

(b) a basic polypeptide factor which has, if obtained from bovine pituitary material, an observed molecular weight, under non-reducing conditions, of from about 55 kD to about 63 kD on SDS-polyacrylamide gel  
10 electrophoresis which factor the human equivalent of which is encoded by DNA clone GGF2HBS5 and which factor has muscle cell mitogenic activity and when isolated using reversed-phase HPLC retains at least 50% of the activity after 4 days incubation in 0.1% trifluoroacetic  
15 acid at 4°C.

Thus other important aspects of the invention are the use of:

(a) A series of human and bovine polypeptide factors having cell mitogenic activity including  
20 stimulating the division of muscle cells. These peptide sequences are shown in Figs. 30, 31, 32 and 33, SEQ ID Nos. 132-133, respectively.

(b) A series of polypeptide factors having cell mitogenic activity including stimulating the division of  
25 muscle cells and purified and characterized according to the procedures outlined by Lupu et al. *Science* 249: 1552 (1990); Lupu et al. *Proc. Natl. Acad. Sci USA* 89: 2287 (1992); Holmes et al. *Science* 256: 1205 (1992); Peles et al. 62: 205 (1992); Yarden and Peles *Biochemistry* 30:  
30 3543 (1991); Dobashi et al. *Proc. Natl. Acad. Sci.* 88: 8582 (1991); Davis et al. *Biochem. Biophys. Res. Commun.* 179: 1536 (1991); Beaumont et al., patent application PCT/US91/03443 (1990); Bottenstein, U.S. Patent No.

- 15 -

5,276,145, issued 1/4/94; and Greene et al. patent application PCT/US91/02331 (1990).

(c) A polypeptide factor (GGFBPP5) having glial cell mitogenic activity including stimulating the  
5 division of muscle cells. The amino acid sequence is shown in Fig. 31, SEQ ID No. 144.

Methods for stimulating mitogenesis of a myoblast by contacting the myoblast cell with a polypeptide defined above as a muscle cell mitogen *in vivo* or *in*  
10 *vitro* are included as features of the invention.

Muscle cell treatments may also be achieved by administering DNA encoding the polypeptide compounds described above in an expressible genetic construction. DNA encoding the polypeptide may be administered to the  
15 patient using techniques known in the art for delivering DNA to the cells. For example, retroviral vectors, electroporation or liposomes may be used to deliver DNA.

The invention includes the use of the above named family of proteins as extracted from natural sources  
20 (tissues or cell lines) or as prepared by recombinant means.

Other compounds in particular, peptides, which bind specifically to the p185<sup>erbB2</sup> receptor can also be used according to the invention as muscle cell mitogens.  
25 A candidate compound can be routinely screened for p185<sup>erbB2</sup> binding, and, if it binds, can then be screened for glial cell mitogenic activity using the methods described herein.

The invention includes use of any modifications or  
30 equivalents of the above polypeptide factors which do not exhibit a significantly reduced activity. For example, modifications in which amino acid content or sequence is altered without substantially adversely affecting activity are included. The statements of effect and use

- 16 -

contained herein are therefore to be construed accordingly, with such uses and effects employing modified or equivalent factors being part of the invention.

5           The human peptide sequences described above and presented in Figs. 30, 31, 32 and 33, SEQ ID Nos. 132-146, respectively, represent a series of splicing variants which can be isolated as full length complementary DNAs (cDNAs) from natural sources (cDNA  
10 libraries prepared from the appropriate tissues) or can be assembled as DNA constructs with individual exons (e.g., derived as separate exons) by someone skilled in the art.

          The invention also includes a method of making a  
15 medicament for treating muscle cells, i.e., for inducing muscular mitogenesis, myogenesis, differentiation, or survival, by administering an effective amount of a polypeptide as defined above. Such a medicament is made by administering the polypeptide with a pharmaceutically  
20 effective carrier.

          Another aspect of the invention is the use of a pharmaceutical or veterinary formulation comprising any factor as defined above formulated for pharmaceutical or veterinary use, respectively, optionally together with an  
25 acceptable diluent, carrier or excipient and/or in unit dosage form. In using the factors of the invention, conventional pharmaceutical or veterinary practice may be employed to provide suitable formulations or compositions.

30           Thus, the formulations to be used as a part of the invention can be applied to parenteral administration, for example, intravenous, subcutaneous, intramuscular, intraorbital, ophthalmic, intraventricular, intracranial, intracapsular, intraspinal, intracisternal,

- 17 -

intraperitoneal, topical, intranasal, aerosol, scarification, and also oral, buccal, rectal or vaginal administration.

5 The formulations of this invention may also be administered by the transplantation into the patient of host cells expressing the DNA encoding polypeptides which are effective for the methods of the invention or by the use of surgical implants which release the formulations of the invention.

10 Parenteral formulations may be in the form of liquid solutions or suspensions; for oral administration, formulations may be in the form of tablets or capsules; and for intranasal formulations, in the form of powders, nasal drops, or aerosols.

15 Methods well known in the art for making formulations are to be found in, for example, "Remington's Pharmaceutical Sciences." Formulations for parenteral administration may, for example, contain as excipients sterile water or saline, polyalkylene glycols  
20 such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes, biocompatible, biodegradable lactide polymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the present factors. Other potentially useful parenteral  
25 delivery systems for the factors include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain as excipients, for example, lactose, or may be aqueous solutions containing, for  
30 example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration,

- 18 -

methoxysalicylate for rectal administration, or citric acid for vaginal administration.

The present factors can be used as the sole active agents, or can be used in combination with other active ingredients, e.g., other growth factors which could facilitate neuronal survival in neurological diseases, or peptidase or protease inhibitors.

The concentration of the present factors in the formulations of the invention will vary depending upon a number of issues, including the dosage to be administered, and the route of administration.

In general terms, the factors of this invention may be provided in an aqueous physiological buffer solution containing about 0.1 to 10% w/v compound for parenteral administration. General dose ranges are from about 1 mg/kg to about 1 g/kg of body weight per day; a preferred dose range is from about 0.01 mg/kg to 100 mg/kg of body weight per day. The preferred dosage to be administered is likely to depend upon the type and extent of progression of the pathophysiological condition being addressed, the overall health of the patient, the make up of the formulation, and the route of administration.

The polypeptide factors utilized in the methods of the invention can also be used as immunogens for making antibodies, such as monoclonal antibodies, following standard techniques. These antibodies can, in turn, be used for therapeutic or diagnostic purposes. Thus, conditions perhaps associated with muscle diseases resulting from abnormal levels of the factor may be tracked by using such antibodies. *In vitro* techniques can be used, employing assays on isolated samples using standard methods. Imaging methods in which the antibodies are, for example, tagged with radioactive isotopes which can be imaged outside the body using



- 19 -

techniques for the art of tumor imaging may also be employed.

A further general aspect of the invention is the use of a factor of the invention in the manufacture of a medicament, preferably for the treatment of a muscular disease or disorder. The "GGF2" designation is used for all clones which were previously isolated with peptide sequence data derived from GGF-II protein (i.e., GGF2HBS5, GGF2BPP3) and, when present alone (i.e., GGF2 or rhGGF2), to indicate recombinant human protein encoded by plasmids isolated with peptide sequence data derived from the GGF-II protein (i.e., as produced in insect cells from the plasmid HBS5). Recombinant human GGF from the GGFHBS5 clone is called GGF2, rhGGF2 and GGF2HBS5 polypeptide.

Treating as used herein means any administration of the compounds described herein for the purpose of increasing muscle cell mitogenesis, survival, and/or differentiation, and/or decreasing muscle atrophy and degeneration. Most preferably, the treating is for the purpose of reducing or diminishing the symptoms or progression of a disease or disorder of the muscle cells. Treating as used herein also means the administration of the compounds for increasing or altering the muscle cells in healthy individuals. The treating may be brought about by the contacting of the muscle cells which are sensitive or responsive to the compounds described herein with an effective amount of the compound, as described above. Inhibitors of the compounds described herein may also be used to halt or slow diseases of muscle cell proliferation.

- 20 -

### Brief Description of the Drawings

The drawings will first be described.

#### Drawings

5 Fig. 1 is a graph showing the results of rhGGF2 in a myoblast mitogenesis assay.

Fig. 2 is a graph showing the effect of rhGGF2 on the number of nuclei in myotubes.

Fig. 3 is a graph of a survival assay showing the effect of rhGGF2 on survival of differentiated myotubes.

10 Fig. 4 is a graph of survival assays showing the effect of rhGGF2 on differentiated myotubes relative to human platelet derived growth factor, human fibroblast growth factor, human epidermal growth factor, human leucocyte inhibitory factor, and human insulin-like growth factors I and II.

Fig. 5 is a graph showing the increased survival on Duchenne muscular dystrophy cells in the presence of rhGGF2.

20 Fig. 6 is a graph of increasing human growth hormone (hGH) expression in C2 cells from an hGH reporter gene under control of the AchR delta subunit transcriptional control elements. This increase is tied to the addition of GGF2 to the media.

Fig. 7 is a graph of increasing hGH reporter synthesis and bungarotoxin (BTX) binding to AchRs following the addition of increasing amounts of GGF2 to C2 cells.

Figs. 8, 9, 10 and 11 are the peptide sequences derived from GGF-I and GGF-II, SEQ ID Nos. 1-20, 22-29, 32-50 and 165, (see Examples 11-13 hereinafter).

30 Fig. 9, Panel A, is the sequences of GGF-I peptides used to design degenerate oligonucleotide probes and degenerate PCR primers are listed (SEQ ID Nos. 1, 17 and 22-29). Some of the sequences in Panel A were also used to design synthetic peptides. Panel B is a listing

- 21 -

of the sequences of novel peptides that were too short (less than 6 amino acids) for the design of degenerate probes or degenerate PCR primers (SEQ ID Nos. 17 and 32);

Fig. 11, Panel A, is a listing of the sequences of GGF-II peptides used to design degenerate oligonucleotide probes and degenerate PCR primers (SEQ ID Nos. 42-49). Some of the sequences in Panel A were used to design synthetic peptides. Panel B is a listing of the novel peptide that was too short (less than 6 amino acids) for the design of degenerate probes or degenerate PCR primers (SEQ ID No. 50);

Figs. 12, 13A, 13B, 14, 15, 16, 17, 18, and 19 relate to Example 8, below, and depict the mitogenic activity of factors of the invention;

Figs. 20, 21, 22, 23, 24, 25, 26, and 27 relate to Example 10, below and are briefly described below:

Fig. 20 is a listing of the degenerate oligonucleotide probes (SEQ ID Nos. 51-84) designed from the novel peptide sequences in Figure 7, Panel A and Figure 9, Panel A;

Fig. 21 (SEQ ID No. 85) depicts a stretch of the putative bovine GGF-II gene sequence from the recombinant bovine genomic phage GGF2BG1, containing the binding site of degenerate oligonucleotide probes 609 and 650 (see Figure 18, SEQ ID NOS. 66 and 69, respectively). The figure is the coding strand of the DNA sequence and the deduced amino acid sequence in the third reading frame. The sequence of peptide 12 from factor 2 (bold) is part of a 66 amino acid open reading frame (nucleotides 75272);

Fig. 22 is the degenerate PCR primers (Panel A, SEQ IS Nos. 86-104) and unique PCR primers (Panel B, SEQ ID Nos. 105-115) used in experiments to isolate segments of the bovine GGF-II coding sequences present in RNA from posterior pituitary;

- 22 -

Fig. 23 depicts of the nine distinct contiguous bovine GGF-II cDNA structures and sequences that were obtained in PCR amplification experiments. The top line of the Figure is a schematic of the coding sequences which contribute to the cDNA structures that were characterized;

Fig. 24 is a physical map of bovine recombinant phage of GGF2BG1. The bovine fragment is roughly 20 kb in length and contains two exons (bold) of the bovine GGF-II gene. Restriction sites for the enzymes XbaI, SpeI, NdeI, EcoRI, KpnI, and SstI have been placed on this physical map. Shaded portions correspond to fragments which were subcloned for sequencing;

Fig. 25 is a schematic of the structure of three alternative gene products of the putative bovine GGF-II gene. Exons are listed A through E in the order of their discovery. The alternative splicing patterns 1, 2 and 3 generate three overlapping deduced protein structures (GGF2BPP1, 2, and 3), which are displayed in the various Figs. 27A, 27B, 27C (described below);

Fig. 26 (SEQ ID Nos. 116-128) is a comparison of the GGF-I and GGF-II sequences identified in the deduced protein sequences shown in Figs. 27A, 27B, 27C (described below) with the novel peptide sequences listed in Figs. 9 and 11. The Figure shows that six of the nine novel GGF-II peptide sequences are accounted for in these deduced protein sequences. Two peptide sequences similar to GGF-I sequences are also found;

Fig. 27 (SEQ ID No. 129) is a listing of the coding strand DNA sequence and deduced amino acid sequence of the cDNA obtained from splicing pattern number 1 in Figure 25. This partial cDNA of the putative bovine GGF-II gene encodes a protein of 206 amino acids in length. Peptides in bold were those identified from

- 23 -

the lists presented in Figs. 9 and 11. Potential glycosylation sites are underlined (along with polyadenylation signal AATAAA);

5 Fig. 27 (SEQ ID No. 130) is a listing of the coding strand DNA sequence and deduced amino acid sequence of the cDNA obtained from splicing pattern number 2 in Fig. 25. This partial cDNA of the putative bovine GGF-II gene encodes a protein of 281 amino acids in length. Peptides in bold are those identified from  
10 the lists presented in Figs. 7 and 9. Potential glycosylation sites are underlined (along with polyadenylation signal AATAAA);

Fig. 27 (SEQ ID No. 131) is a listing of the coding strand DNA sequence and deduced amino acid  
15 sequence of the cDNA obtained from splicing pattern number 3 in Fig. 25. This partial cDNA of the putative bovine GGF-II gene encodes a protein of 257 amino acids in length. Peptides in bold are those identified from the lists in Figs. 9 and 11. Potential glycosylation  
20 sites are underlined (along with polyadenylation signal AATAAA).

Fig. 28, which relates to Example 16 hereinafter, is an autoradiogram of a cross hybridization analysis of putative bovine GGF-II gene sequences to a variety of  
25 mammalian DNAs on a southern blot. The filter contains lanes of EcoRI-digested DNA (5  $\mu$ g per lane) from the species listed in the Figure. The probe detects a single strong band in each DNA sample, including a four kilobase fragment in the bovine DNA as anticipated by the physical  
30 map in Fig. 24. Bands of relatively minor intensity are observed as well, which could represent related DNA sequences. The strong hybridizing band from each of the other mammalian DNA samples presumably represents the GGF-II homologue of those species.

- 24 -

Fig. 29 is a diagram of representative splicing variants. The coding segments are represented by F, E, B, A, G, C, C/D, C/D', D, D', H, K and L. The location of the peptide sequences derived from purified protein  
5 are indicated by "o".

Fig. 30 (SEQ ID Nos. 136-143, 156, 157, 169-178) is a listing of the DNA sequences and predicted peptide sequences of the coding segments of GGF. Line 1 is a listing of the predicted amino acid sequences of bovine  
10 GGF, line 2 is a listing of the nucleotide sequences of bovine GGF, line 3 is a listing of the nucleotide sequences of human GGF (heregulin) (nucleotide base matches are indicated with a vertical line) and line 4 is a listing of the predicted amino acid sequences of human  
15 GGF/hereregulin where it differs from the predicted bovine sequence. Coding segments E, A' and K represent only the bovine sequences. Coding segment D' represents only the human (heregulin) sequence.

Fig. 31 (SEQ ID No. 144) is the predicted GGF2  
20 amino acid sequence and nucleotide sequence of BPP5. The upper line is the nucleotide sequence and the lower line is the predicted amino acid sequence.

Fig. 32 (SEQ ID No. 145) is the predicted amino acid sequence and nucleotide sequence of GGF2BPP2. The  
25 upper line is the nucleotide sequence and the lower line is the predicted amino acid sequence.

Fig. 33 (SEQ ID No. 146) is the predicted amino acid sequence and nucleotide sequence of GGF2BPP4. The  
30 upper line is the nucleotide sequence and the lower line is the predicted amino acid sequence.

Fig. 34 (SEQ ID Nos. 147-149) depicts the alignment of two GGF peptide sequences (GGF2BPP4 and GGF2BPP5) with the human EGF (hEGF). Asterisks indicate positions of conserved cysteines.

- 25 -

Fig. 35 depicts the level of GGF activity (Schwann cell mitogenic assay) and tyrosine phosphorylation of a ca. 200kD protein (intensity of a 200 kD band on an autoradiogram of a Western blot developed with an antiphosphotyrosine polyclonal antibody) in response to increasing amounts of GGF.

Fig. 36 is a list of splicing variants derived from the sequences shown in Fig. 30.

Fig. 37 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL1 (SEQ ID No. 150).

Fig. 38 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL2 (SEQ ID No. 151).

Fig. 39 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL3 (SEQ ID No. 152).

Fig. 40 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL4 (SEQ ID No. 153).

Fig. 41 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL5 (SEQ ID No. 154).

Fig. 42 is the predicted amino acid sequence, bottom, and nucleic sequence, top, of EGFL6 (SEQ ID No. 159).

Fig. 43 is a scale coding segment map of the clone. T3 refers to the bacteriophage promoter used to produce mRNA from the clone. R = flanking EcoRI restriction enzyme sites. 5' UT refers to the 5' untranslated region. E, B, A, C, C/D', and D refer to the coding segments. O = the translation start site.  $\Lambda$  = the 5' limit of the region homologous to the bovine E

- 26 -

segment (see Example 17) and 3' UT refers to the 3' untranslated region.

Fig. 44 is the predicted amino acid sequence (middle) and nucleic sequence (top) of GGF2HBS5 (SEQ ID No. 21). The bottom (intermittent) sequence represents peptide sequences derived from GGF-II preparations (see Figs. 8, 9).

Fig. 45 (A) is a graph showing the purification of rGGF on cation exchange column by fraction; Fig. 45 (B) is a photograph of a Western blot using fractions as depicted in (A) and a GGFII specific antibody.

Fig. 46 is the sequence of the GGFHBS5, GGFHFB1 and GGF BPP5 polypeptides (SEQ ID NOS: 166, 167, and 168).

Fig. 47 is a map of the plasmid pCDHRFpolyA.

15

#### Detailed Description

The invention pertains to the use of isolated and purified neuregulin factors and DNA sequences encoding these factors, regulatory compounds which increase the extramuscular concentrations of these factors, and compounds which are mimetics of these factors for the induction of muscle cell mitogenesis, differentiation, and survival of the muscle cells *in vivo* and *in vitro*.

It is evident that the gene encoding GGF/p185<sup>erbB2</sup> binding neuregulin proteins produces a number of variably-sized, differentially-spliced RNA transcripts that give rise to a series of proteins. These proteins are of different lengths and contain some common peptide sequences and some unique peptide sequences. The conclusion that these factors are encoded by a single gene is supported by the differentially-spliced RNA sequences which are recoverable from bovine posterior pituitary and human breast cancer cells (MDA-MB-231). Further support for this conclusion derives from the size

20  
25  
30



- 27 -

range of proteins which act as both mitogens for muscle tissue (as disclosed herein) and as ligands for the p185<sup>erbB2</sup> receptor (see below).

Further evidence to support the fact that the  
5 genes encoding GGF/p185<sup>erbB2</sup> binding proteins are homologous comes from nucleotide sequence comparison. Holmes et al., (Science 256:1205-1210, 1992) demonstrate the purification of a 45-kilodalton human protein (Heregulin- $\alpha$ ) which specifically interacts with the  
10 receptor protein p185<sup>erbB2</sup>. Peles et al. (Cell 69:205 (1992)) and Wen et al. (Cell 69:559 (1992)) describe a complementary DNA isolated from rat cells encoding a protein called "neu differentiation factor" (NDF). The translation product of the NDF cDNA has p185<sup>erbB2</sup> binding  
15 activity. Several other groups have reported the purification of proteins of various molecular weights with p185<sup>erbB2</sup> binding activity. These groups include Lupu et al. ((1992) Proc. Natl. Acad. Sci. USA 89:2287); Yarden and Peles ((1991) Biochemistry 30:3543); Lupu et  
20 al. ((1990) Science 249:1552)); Dobashi et al. ((1991) Biochem. Biophys. Res. Comm. 179:1536); and Huang et al. ((1992) J. Biol. Chem. 257:11508-11512).

We have found that p185<sup>erbB2</sup> receptor binding proteins stimulate muscle cell mitogenesis and hence,  
25 stimulates myotube formation (myogenesis). This stimulation results in increased formation of myoblasts and increased formation of myotubes (myogenesis). The compounds described herein also stimulate increased muscle growth, differentiation, and survival of muscle  
30 cells. These ligands include, but are not limited to the GGF's, the neuregulins, the heregulins, NDF, and ARIA. As a result of this mitogenic activity, these proteins, DNA encoding these proteins, and related compounds may be administered to patients suffering from traumatic damage

- 28 -

or diseases of the muscle tissue. It is understood that all methods provided for the purpose of mitogenesis are useful for the purpose of myogenesis. Inhibitors of these ligands (such as antibodies or peptide fragments) may be administered for the treatment of muscle derived tumors.

These compounds may be obtained using the protocols described herein (Examples 9-17) and in Holmes et al., *Science* 256: 1205 (1992); Peles et al., *Cell* 69:205 (1992); Wen et al., *Cell* 69:559 (1992); Lupu et al., *Proc. Natl. Acad. Sci. USA* 89:2287 (1992); Yarden and Peles, *Biochemistry* 30:3543 (1991); Lupu et al., *Science* 249:1552 (1990); Dobashi et al., *Biochem. Biophys. Res. Comm.* 179:1536 (1991); Huang et al., *J. Biol. Chem.* 257:11508-11512 (1992); Marchionni et al., *Nature* 362:313, (1993); and in the GGF-III patent, all of which are incorporated herein by reference. The sequences are provided and the characteristics described for many of these compounds. For sequences see Figs. 8-11, 20-27C, 29-34, 36-44, and 46. For protein characteristics see Figs. 12-19, 28 35, 45A and 45B.

Compounds may be assayed for their usefulness *in vitro* using the methods provided in the examples below. *In vivo* testing may be performed as described in Example 1 and in Sklar et al., *In Vitro Cellular and Developmental Biology* 27A:433-434, 1991.

#### Other Embodiments

The invention includes methods for the use of any protein which is substantially homologous to the coding segments in Fig. 30 (SEQ ID Nos. 132-143, 156, 1576-147, 160, and 161) as well as other naturally occurring GGF polypeptides for the purpose of inducing muscle mitogenesis. Also included are the use of: allelic

- 29 -

variations; natural mutants; induced mutants; proteins encoded by DNA that hybridizes under high or low stringency conditions to a nucleic acid naturally occurring (for definitions of high and low stringency see  
5 Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1989, 6.3.1 - 6.3.6, hereby incorporated by reference); and the use of polypeptides or proteins specifically bound by antisera to GGF polypeptides. The term also includes the use of chimeric polypeptides that  
10 include the GGF polypeptides comprising sequences from Fig. 28 for the induction of muscle mitogenesis.

As will be seen from Example 8, below, the present factors exhibit mitogenic activity on a range of cell types. The general statements of invention above in  
15 relation to formulations and/or medicaments and their manufacture should clearly be construed to include appropriate products and uses.

A series of experiments follow which provide additional basis for the claims described herein. The  
20 following examples relating to the present invention should not be construed as specifically limiting the invention, or such variations of the invention, now known or later developed.

The examples illustrate our discovery that  
25 recombinant human GGF2 (rhGGF2) confers several effects on primary human muscle culture. rhGGF2 has significant effects in three independent biological activity assays on muscle cultures. The polypeptide increased  
mitogenesis as measured by proliferation of subconfluent  
30 quiescent myoblasts, increased differentiation by confluent myoblasts in the presence of growth factor, and increased survival of differentiated myotubes as measured by loss of dye exclusion and increased acetylcholine receptor synthesis. These activities indicate efficacy

- 30 -

of GGF2 and other neuregulins in inducing muscle repair, regeneration, and prophylactic effects on muscle degeneration.

#### EXAMPLE 1

##### 5 Mitogenic Activity of rhGGF on Myoblasts

Clone GGF2HBS5 was expressed in recombinant Baculovirus infected insect cells as described in Example 14, *infra*, and the resultant recombinant human GGF2 was added to myoblasts in culture (conditioned  
 10 medium added at 40  $\mu$ l/ml). Myoblasts (057A cells) were grown to preconfluence in a 24 well dish. Medium was removed and replaced with DMEM containing 0.5% fetal calf serum with or without GGF2 conditioned medium at a concentration of 40  $\mu$ l/ml. Medium was changed after 2  
 15 days and cells were fixed and stained after 5 days. Total nuclei were counted as were the number of nuclei in myoblasts (Table 1).

TABLE 1

Treatment	Total Number of Nuclei/mm <sup>2</sup>	Nuclei in Myotubes	Fusion Index
20 Control	395 $\pm$ 28.3	204 $\pm$ 9.19	0.515 $\pm$ 0.01
GGF 40 $\mu$ l/ml	636 $\pm$ 8.5	381 $\pm$ 82.7	0.591 $\pm$ 0.15

GGF treated myoblasts showed an increased number of total nuclei (636 nuclei) over untreated controls (395 nuclei) indicating mitogenic activity. rhGGF2 treated myotubes  
 25 had a greater number of nuclei (381 nuclei) than untreated controls (204 nuclei). Thus, rhGGF2 enhances the total number of nuclei through proliferation and increased cell survival. rhGGF2 is also likely to enhance the formation of myotubes.

- 31 -

The mitogenic activity of rhGGF2 may be measured *in vivo* by giving a continuous supply of GGF2 and [<sup>3</sup>H]thymidine to rat muscle via an osmotic mini pump. The muscle bulk is determined by wet weight after one and  
5 two weeks of treatment. DNA replication is measured by counting labeled nuclei in sections after coating for autoradiography (Sklar et al., *In Vitro Cellular and Developmental Biology* 27A:433-434, 1991) in sham and  
10 rhGGF2-treated muscle. Denervated muscle is also examined in this rat animal model via these methods and this method allows the assessment of the role of rhGGF2 in muscle atrophy and repair. Mean fiber diameter can also be used for assessing effects of FGF on prevention of atrophy.

15

EXAMPLE 2Effect of rhGGF2 on Muscle Cell Mitogenesis

Quiescent primary clonal human myoblasts were prepared as previously described (Sklar, R., Hudson, A., Brown, R., *In vitro Cellular and Developmental Biology*  
20 1991; 27A:433-434). The quiescent cells were treated with the indicated agents (rhGGF2 conditioned media, PDGF with and without methylprednisolone, and control media) in the presence of 10 $\mu$ M BrdU, 0.5% FCS in DMEM. After  
25 two days the cells were fixed in 4% paraformaldehyde in PBS for 30 minutes, and washed with 70% ethanol. The cells were then incubated with an anti-BrdU antibody, washed, and antibody binding was visualized with a peroxidase reaction. The number of staining nuclei were then quantified per area. The results show that GGF2  
30 induces an increase in the number of labelled nuclei per area over controls (see Table 2).

TABLE 2

## Mitogenic Effects of GGF on Human Myoblasts

Treatment	Labelled Nuclei/cm <sup>2</sup>	T-Test p value
Control	120 ± 22.4	
5 Infected Control	103 ± 11.9	
GGF 5 µl/ml	223 ± 33.8	0.019
PDGF 20 ng/ml	418 ± 45.8	0.0005
IGFI 30 ng/ml	280 ± 109.6	0.068
Methylprednisolone 1.0 µM	142 ± 20.7	0.293

10 Platelet derived growth factor (PDGF) was used as a positive control. Methylprednisolone (a corticosteroid) was also used in addition to rhGGF2 and showed no significant increase in labelling of DNA.

rhGGF2 purified to homogeneity (>95% pure) is also  
15 mitogenic for human myoblasts (Fig. 1).

Recombinant human GGF2 also causes mitogenesis of primary human myoblasts (see Table 2 and Fig. 1). The mitogenesis assay is performed as described above. The mitotic index is then calculated by dividing the number  
20 of BrdU positive cells by the total number of cells.

EXAMPLE 3Effect of rhGGF2 on Muscle Cell Differentiation

The effects of purified rhGGF2 (95% pure) on muscle culture differentiation were examined (Fig. 2).  
25 Confluent myoblast cultures were induced to differentiate by lowering the serum content of the culture medium from 20% to 0.5%. The test cultures were treated with the indicated concentration of rhGGF2 for six days, refreshing the culture medium every 2 days. The cultures

- 33 -

were then fixed, stained, and the number of nuclei counted per millimeter. The data in Fig. 2 demonstrate a large increase in the number of nuclei in myotubes when rhGGF2 is present, relative to controls.

5

**EXAMPLE 4****Effect of rhGGF2 on the Survival of Differentiated Myotubes**

The survival of differentiated myotubes was significantly increased by rhGGF2 treatment. Muscle  
10 cultures were differentiated in the presence of rhGGF2 and at various times the number of dead myotubes were counted by propidium iodide staining. As can be seen in Fig. 3, the number of dead myotubes is lower in the rhGGF2 treated culture at 4, 5, 6, and 8 days of  
15 differentiation. The number of nuclei in myotubes was significantly increased by GGF2 treatment compared to untreated cultures after 8 days of differentiation. Specifically, the control showed 8.6 myonuclei/mm<sup>2</sup>, while rhGGF2 treated cultures showed 57.2 myonuclei/mm<sup>2</sup>  
20 (p=0.035) when counted on the same plates after geimsa staining.

The survival assay was also performed with other growth factors which have known effects on muscle culture. The rhGGF2 effect was unique among the growth  
25 factors tested (Fig. 4). In this experiment cultures were treated in parallel with the rhGGF2 treated plates with the indicated concentrations of the various growth factors. Survival of myotubes was measured as above at 8 days of differentiation of 057A myoblast cells.  
30 Concentrations of factors were as follows: rhGGF2: 100ng/ml; human platelet derived growth factor: 20ng/ml; human basic fibroblast growth factor: 25ng/ml; human epidermal growth factor: 30ng/ml; human leucocyte

- 34 -

inhibitory factor: 10ng/ml; human insulin like growth factor I: 30ng/ml; human insulin like growth factor II: 25ng/ml.

5 The observed protection of differentiated myotubes from death indicates that rhGGF2 has promise as a therapy for intervention of muscle degeneration characterized by numerous muscle diseases. Thus, agents which increase the extramuscular concentration of neuregulins may have a prophylactic effect or slow the progress of muscle-  
10 wasting disorders and increase rates of muscle differentiation, repair, conditioning, and regeneration.

#### EXAMPLE 5

##### rhGGF2 Promotes Survival of Differentiated Myotubes with a Genetic Defect at the Duchenne Muscular Dystrophy Locus

15 The positive effects of rhGGF2 on myotube survival could reflect potential efficacy in degenerative disorders. These effects on myotube survival were tested on a clonally-derived primary Duchenne myoblast to determine if the response observed in normal muscle  
20 culture could also be demonstrated in cultures derived from diseased individuals. The data presented in Fig. 5 was obtained using the same muscle culture conditions (Example 4, above) used for normal individual. rhGGF2 significantly decreased the number of dead myotubes in  
25 the differentiated Duchenne muscle culture, compared to controls ( $p=0.032$ ). Concentrations were as follows: GGF2: 100ng/ml; human platelet derived growth factor: 20ng/ml; human insulin like growth factor I: 30ng/ml.

30 This example demonstrates that rhGGF2 can also promote survival of differentiated Duchenne myotubes and provides strong evidence that rhGGF2 may slow or prevent the course of muscle degeneration and wasting in mammals.



- 35 -

**EXAMPLE 6****rhGGF2 Effect on the Differentiation Program: Induction of MHC Slow and Dystrophin Proteins**

The effects of purified rhGGF2 on muscle culture differentiation was also examined by Western analysis of culture lysates. The levels of muscle specific proteins were determined in triplicate treated and untreated cultures. These cultures were prepared and treated as above except that the plate size was increased to 150 mm and the muscle culture layer was scraped off for Western analysis as described in Sklar, R., and Brown, R. (*J. Neurol. Sci.* 101:73-81, 1991). The results presented in Table A indicate that rhGGF2 treatment increases the levels of several muscle specific proteins, including dystrophin, myosin heavy chain (MHC, adult slow and fast isoforms), but does not increase the levels of HSP72 or MHC neonate isoform to a similar level per amount of protein loaded on the Western. The levels of muscle specific proteins induced by rhGGF2 were similar to the quantitative increases in the number of myonuclei/mm<sup>2</sup> (Table 3).

TABLE 3

value	Control $\pm$ SD	rhGGF2 Treat- ment $\pm$ SD	p	
5	Total Protein ( $\mu$ g)	554 $\pm$ 38.4	798 $\pm$ 73.6	0.007
	Myonuclei/mm <sup>2</sup>	29.0 $\pm$ 12.2	106 $\pm$ 24.1	0.008
	MHC fast/ $\mu$ g protein	1.22 $\pm$ 0.47	4.00 $\pm$ 0.40	0.001
	MHC slow/ $\mu$ g protein	0.17 $\pm$ 0.13	1.66 $\pm$ 0.27	0.001
	MHC neonate/ $\mu$ g protein	0.30 $\pm$ 0.27	0.55 $\pm$ 0.04	0.199
10	dystrophin/ $\mu$ g protein	6.67 $\pm$ 0.37	25.5 $\pm$ 11.0	0.042
	HSP 72/ $\mu$ g protein	3.30 $\pm$ 0.42	4.54 $\pm$ 0.08	0.008

The rhGGF2 dependent increase in the adult myosin heavy chain isoforms (slow is found in type I human muscle fibers; fast is found in type 2A and 2B human muscle fibers) may represent a maturation of the myotubes, as the neonatal isoform was not significantly increased by rhGGF2 treatment. During rat muscle development MHC isoforms switch from fetal to neonatal forms followed by a switch to mature adult slow and fast MHC isoforms (Periasamy et al. *J. Biol. Chem.* 259:13573-13578, 1984; Periasamy et al. *J. Biol. Chem.* 260:15856-15862, 1985; Wieczorek et al. *J. Cell Biol.* 101:618-629, 1985). While muscle can autonomously undergo some of these isoform transitions in the absence of neural cells or tissue, mouse muscle explants express the adult fast MHC isoform only when cultured in the presence of mouse spinal cord (Ecob-Prince et al. *J. Cell Biol.* 103:995-1005, 1986). Additional evidence that MHC isoform transitions are influenced by nerve was established by Whalen et al. (*Deve. Biol.* 141:24-40, 1990); after regeneration of notexin treated rat soleus muscles only

- 37 -

the adult fast MHC isoform was produced in the new denervated muscle, but innervated regenerated muscle made both fast and slow adult MHC isoforms. Thus the demonstration in Table 3 that rhGGF2 increases the synthesis of adult MHC isoforms indicates that rhGGF2 may induce a developmental maturation of muscle which may mimic neuronal innervation.

#### EXAMPLE 7

Neuregulins, including rhGGF2, induce the synthesis of acetylcholine receptors in muscle.

The expression of acetylcholine receptor (AChR) subunit proteins can be induced by exposing muscle cells to neuregulins. More specifically, we have shown that contacting muscle cells with rhGGF2 can induce the synthesis of AChR subunit proteins. This induction following rhGGF2 exposure was observed in two ways: first, we detected increased expression of human growth hormone via the product of a reporter gene construct and second we detected increased binding of alpha-bungarotoxin to cells.

In the following example a mouse myoblast cell line C2 was used. C2 cells were transfected with a transgene that contained the 5' regulatory sequences of the AChR delta subunit gene of mouse linked to a human growth hormone full-length cDNA (Baldwin and Burden, 1988. J. Cell Biol. 107:2271-2279). This reporter construct allows the measurement of the induction of AChR delta gene expression by assaying the quantity of growth hormone secreted into the media. The line can be induced to form myotubes by lowering serum concentration in the media from 20% to 0.5%.

Specifically, mouse C2 myoblasts transfected with an AChR-human growth hormone reporter construct and were

- 38 -

assayed for expression of hGH following treatment with rhGGF2. The results of two separate experiments are summarized in Table 4 and in Figures 6 (hGH expression) and 7 (hGH expression and alpha-bungarotoxin binding).

5 Shown are the dose response curves for secreted human growth hormone and for bungarotoxin binding from muscle cultures treated with rhGGF2.

TABLE 4

10 Effects of rhGGF2 on the expression of AChR delta subunit/hGH transgene and the synthesis of AChR

	Exp 1		Exp 2	
15	GGF (ul)	hGH (ng/ml)	hGH (ng/ml)	AChR (cpm/mg protein)
	0	9.3 + 2.1	5.7 + 2.1	822 + 170
	0.1	-	6.8 + 1.5	891 + 134
	0.5	-	12.0 + 0.9	993 + 35
	1.0	-	9.7 + 2.3	818 + 67
20	5.0	17.5 + 2.8	14.7 + 3.5	1300 + 177
	10.0	14.3 + 3.2	14.1 + 3.3	1388 + 137
	15.0	22.0 + 1.4	-	-
25	-	-	-	-

30 C2 myotubes were treated with cold  $\alpha$ -BTX (20 nM) for 1 hr. at 37°C, washed with culture medium twice and then treated with GGF2. Culture medium was adjusted with bovine serum albumin at the concentration of 1 mg/ml. 24 hours later, culture medium was removed and saved for hGH

- 39 -

assay. Muscle cultures were treated with  $^{125}\text{I}$ - $\alpha$ -BTX (20 nM) for 1 hour at 37°C, washed and scraped in PBS containing 1% SDS. Non-specific binding was determined in the presence of cold  $\alpha$ -BTX (40 nM). The cell  
5 homogenate was counted for radioactivity and assayed for total protein amount.

The presence of rhGGF2 led to a greater than 2-fold increase in hGH gene expression, thereby indicating that rhGGF2 induced the synthesis of the delta subunit of  
10 the acetylcholine receptor. Furthermore, increased bungarotoxin binding is consistent with assembly of these subunit proteins into functional acetylcholine receptors. To strengthen the interpretation of these data the analysis was repeated on cultures that had the hGH  
15 reporter linked to a metallothionein promoter, which should not be responsive to rhGGF2. The results of that control experiment showed that the hGH response was mediated through transcriptional activation of the AchR delta subunit gene control elements.

20 These results indicate that rhGGF2 could be useful in replenishing AchRs as part of the therapy for the autoimmune disease Myasthenia gravis. This activity may also be beneficial in treatment of peripheral nerve regeneration and neuropathy by stimulating a key step in  
25 re-innervation of muscle.

#### EXAMPLE 8

##### Additional Mitogenic Activities of Purified GGF-I and GGF-II

The mitogenic activity of a highly purified sample  
30 containing both GGFs I and II was studied using a quantitative method, which allows a single microculture to be examined for DNA synthesis, cell morphology, cell number and expression of cell antigens. This technique

- 40 -

has been modified from a method previously reported by Muir et al., Analytical Biochemistry 185, 377-382, 1990. The main modifications are: 1) the use of uncoated microtiter plates, 2) the cell number per well, 3) the use of 5% Foetal Bovine Plasma (FBP) instead of 10% Foetal Calf Serum (FCS), and 4) the time of incubation in presence of mitogens and bromodeoxyuridine (BrdU), added simultaneously to the cultures. In addition the cell monolayer was not washed before fixation to avoid loss of cells, and the incubation time of monoclonal mouse anti-BrdU antibody and peroxidase conjugated goat anti-mouse immunoglobulin (IgG) antibody were doubled to increase the sensitivity of the assay. The assay, optimized for rat sciatic nerve Schwann cells, has also been used for several cell lines, after appropriate modifications to the cell culture conditions.

#### I. Methods of Mitogenesis Testing

On day 1, purified Schwann cells were plated onto uncoated 96 well plates in 5% FBP/Dulbecco's Modified Eagle Medium (DMEM) (5,000 cells/well). On day 2, GGFs or other test factors were added to the cultures, as well as BrdU at a final concentration of 10  $\mu$ M. After 48 hours (day 4) BrdU incorporation was terminated by aspirating the medium and cells were fixed with 200  $\mu$ l/well of 70% ethanol for 20 min at room temperature. Next, the cells were washed with water and the DNA denatured by incubation with 100  $\mu$ l 2N HCl for 10 min at 37°C. Following aspiration, residual acid was neutralized by filling the wells with 0.2 M borate buffer, pH 9.0, and the cells were washed with phosphate buffered saline (PBS). Cells were then treated with 50  $\mu$ l of blocking buffer (PBS containing 0.1% Triton X 100 and 2% normal goat serum) for 15 min at 37°C. After

- 41 -

aspiration, monoclonal mouse anti-BrdU antibody (Dako Corp., Santa Barbara, CA) (50  $\mu$ l/well, 1.4  $\mu$ g/ml diluted in blocking buffer) was added and incubated for two hours at 37°C. Unbound antibodies were removed by three washes  
5 in PBS containing 0.1% Triton X-100 and peroxidase-conjugated goat anti-mouse IgG antibody (Dako Corp., Santa Barbara, CA) (50  $\mu$ l/well, 2  $\mu$ g/ml diluted in blocking buffer) was added and incubated for one hour at 37°C. After three washes in PBS/Triton and a final rinse  
10 in PBS, wells received 100  $\mu$ l/well of 50 mM phosphate/citrate buffer, pH 5.0, containing 0.05% of the soluble chromogen o-phenylenediamine (OPD) and 0.02%  $H_2O_2$ . The reaction was terminated after 5-20 min at room temperature, by pipetting 80  $\mu$ l from each well to a clean  
15 plate containing 40  $\mu$ l/well of 2N sulfuric acid. The absorbance was recorded at 490nm using a plate reader (Dynatech Labs). The assay plates containing the cell monolayers were washed twice with PBS and immunocytochemically stained for BrdU-DNA by adding 100  
20  $\mu$ l/well of the substrate diaminobenzidine (DAB) and 0.02%  $H_2O_2$  to generate an insoluble product. After 10-20 min the staining reaction was stopped by washing with water, and BrdU-positive nuclei observed and counted using an inverted microscope. occasionally, negative nuclei were  
25 counterstained with 0.001% Toluidine blue and counted as before.

## II. Cell lines used for Mitogenesis Assays

*Swiss 3T3 Fibroblasts:* Cells, from Flow Labs, were maintained in DMEM supplemented with 10% FCS,  
30 penicillin and streptomycin, at 37°C in a humidified atmosphere of 10%  $CO_2$  in air. Cells were fed or subcultured every two days. For mitogenic assay, cells were plated at a density of 5,000 cells/well in complete

- 42 -

medium and incubated for a week until cells were  
confluent and quiescent. The serum containing medium was  
removed and the cell monolayer washed twice with serum  
free-medium. 100  $\mu$ l of serum free medium containing  
5 mitogens and 10 $\mu$ M of BrdU were added to each well and  
incubated for 48 hours. Dose responses to GGFs and serum  
or PDGF (as a positive control) were performed.

*BHK (Baby Hamster Kidney) 21 C13 Fibroblasts:*

Cells from European Collection of Animal Cell Cultures  
10 (ECACC), were maintained in Glasgow Modified Eagle Medium  
(GMEM) supplemented with 5% tryptose phosphate broth, 5%  
FCS, penicillin and streptomycin, at 37°C in a humidified  
atmosphere of 5% CO<sub>2</sub> in air. Cells were fed or  
subcultured every two to three days. For mitogenic  
15 assay, cells were plated at a density of 2,000 cell/well  
in complete medium for 24 hours. The serum containing  
medium was then removed and after washing with serum free  
medium, replaced with 100  $\mu$ l of 0.1% FCS containing GMEM  
or GMEM alone. GGFs and FCS or bFGF as positive controls  
20 were added, coincident with 10 $\mu$ M BrdU, and incubated for  
48 hours. Cell cultures were then processed as described  
for Schwann cells.

*C6 Rat Glioma Cell Line:* Cells, obtained at  
passage 39, were maintained in DMEM containing 5% FCS, 5%  
25 Horse serum (HS), penicillin and streptomycin, at 37°C in  
a humidified atmosphere of 10% CO<sub>2</sub> in air. Cells were  
fed or subcultured every three days. For mitogenic  
assay, cells were plated at a density of 2,000 cells/well  
in complete medium and incubated for 24 hours. Then  
30 medium was replaced with a mixture of 1:1 DMEM and F12  
medium containing 0.1% FCS, after washing in serum free  
medium. Dose responses to GGFs, FCS and  $\alpha$ FGF were then



- 43 -

performed and cells were processed through the ELISA as previously described for the other cell types.

**PC12 (Rat Adrenal Pheochromocytoma Cells):** Cells from ECACC, were maintained in RPMI 1640 supplemented with 10% HS, 5% FCS, penicillin and streptomycin, in collagen coated flasks, at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> in air. Cells were fed every three days by replacing 80% of the medium. For mitogenic assay, cells were plated at a density of 3,000 cells/well in complete medium, on collagen coated plates (50 µl/well collagen, Vitrogen Collagen Corp., diluted 1 : 50, 30 min at 37°C) and incubated for 24 hours. The medium was then placed with fresh RPMI either alone or containing 1 mM insulin or 1% FCS. Dose responses to FCS/HS (1:2) as positive control and to GGFs were performed as before. After 48 hours cells were fixed and the ELISA performed as previously described.

**III. Results of Mitogenesis Assays:** All the experiments presented in this Example were performed using a highly purified sample from a Sepharose 12 chromatography purification step containing a mixture of GGF-I and GGF-II (GGFs).

First, the results obtained with the BrdU incorporation assay were compared with the classical mitogenic assay for Schwann cells based on [<sup>125</sup>I]-UdR incorporation into DNA of dividing cells, described by J. P. Brookes (*Methods Enzymol.* 147:217, 1987).

Fig. 12 shows the comparison of data obtained with the two assays, performed in the same cell culture conditions (5,000 cells/well, in 5% FBP/DMEM, incubated in presence of GGFs for 48 hrs). As clearly shown, the results are comparable, but BrdU incorporation assay

- 44 -

appears to be slightly more sensitive, as suggested by the shift of the curve to the left of the graph, i.e. to lower concentrations of GGFs.

As described under the section "Methods of  
5 Mitogenesis Testing", after the immunoreactive BrdU-DNA has been quantitated by reading the intensity of the soluble product of the OPD peroxidase reaction, the original assay plates containing cell monolayers can undergo the second reaction resulting in the insoluble  
10 DAB product, which stains the BrdU positive nuclei. The microcultures can then be examined under an inverted microscope, and cell morphology and the numbers of BrdU-positive and negative nuclei can be observed.

In Fig. 13A and Fig. 13B the BrdU-DNA  
15 immunoreactivity, evaluated by reading absorbance at 490 nm, is compared to the number of BrdU-positive nuclei and to the percentage of BrdU-positive nuclei on the total number of cells per well, counted in the same cultures. Standard deviations were less than 10%. The two  
20 evaluation methods show a very good correlation and the discrepancy between the values at the highest dose of GGFs can be explained by the different extent of DNA synthesis in cells detected as BrdU-positive.

The BrdU incorporation assay can therefore provide  
25 additional useful information about the biological activity of polypeptides on Schwann cells when compared to the (125) I-UdR incorporation assay. For example, the data reported in Fig. 15 show that GGFs can act on Schwann cells to induce DNA synthesis, but at lower doses  
30 to increase the number of negative cells present in the microculture after 48 hours.

The assay has then been used on several cell lines of different origin. In Fig. 15 the mitogenic responses of Schwann cells and Swiss 3T3 fibroblasts to GGFs are

- 45 -

compared; despite the weak response obtained in 3T3 fibroblasts, some clearly BrdU-positive nuclei were detected in these cultures. Control cultures were run in parallel in presence of several doses of FCS or human recombinant PDGF, showing that the cells could respond to appropriate stimuli (not shown).

The ability of fibroblasts to respond to GGFs was further investigated using the BHK 21 C13 cell line. These fibroblasts, derived from kidney, do not exhibit contact inhibition or reach a quiescent state when confluent. Therefore the experimental conditions were designed to have a very low background proliferation without compromising the cell viability. GGFs have a significant mitogenic activity on BHK21 C13 cells as shown by Fig. 16 and Fig. 17. Fig. 16 shows the BrdU incorporation into DNA by BHK 21 C13 cells stimulated by GGFs in the presence of 0.1% FCS. The good mitogenic response to FCS indicates that cell culture conditions were not limiting. In Fig. 17 the mitogenic effect of GGFs is expressed as the number of BrdU-positive and BrdU-negative cells and as the total number of cells counted per well. Data are representative of two experiments run in duplicates; at least three fields per well were counted. As observed for Schwann cells in addition to a proliferative effect at low doses, GGFs also increase the numbers of nonresponding cells surviving. The percentage of BrdU positive cells is proportional to the increasing amounts of GGFs added to the cultures. The total number of cells after 48 hours in presence of higher doses of GGFs is at least doubled, confirming that GGFs induce DNA synthesis and proliferation in BHK21 C13 cells. Under the same conditions, cells maintained for 48 hours in the presence

- 46 -

of 2% FCS showed an increase of about six fold (not shown).

C6 glioma cells have provided a useful model to study glial cell properties. The phenotype expressed  
5 seems to be dependent on the cell passage, the cells more closely resembling an astrocyte phenotype at an early stage, and an oligodendrocyte phenotype at later stages (beyond passage 70). C6 cells used in these experiments were from passage 39 to passage 52. C6 cells are a  
10 highly proliferating population, therefore the experimental conditions were optimized to have a very low background of BrdU incorporation. The presence of 0.1% serum was necessary to maintain cell viability without significantly affecting the mitogenic responses, as shown  
15 by the dose response to FCS (Fig. 18).

In Fig. 19 the mitogenic responses to aFGF (acidic Fibroblast growth factor) and GGFS are expressed as the percentages of maximal BrdU incorporation obtained in the presence of FCS (8%). Values are averages of two  
20 experiments, run in duplicates. The effect of GGFS was comparable to that of a pure preparation of aFGF. aFGF has been described as a specific growth factor for C6 cells (Lim K. et al., Cell Regulation 1:741-746, 1990) and for that reason it was used as a positive control.  
25 The direct counting of BrdU positive and negative cells was not possible because of the high cell density in the microcultures. In contrast to the cell lines so far reported, PC12 cells did not show any evident responsiveness to GGFS, when treated under culture  
30 conditions in which PC12 could respond to sera (mixture of FCS and HS as used routinely for cell maintenance). Nevertheless the number of cells plated per well seems to affect the behavior of PC12 cells, and therefore further experiments are required.

- 47 -

EXAMPLE 9Amino acid sequences of purified GGF-I and GGF-II

Amino acid sequence analysis studies were performed using highly purified bovine pituitary GGF-I and GGF-II. The conventional single letter code was used to describe the sequences. Peptides were obtained by lysyl endopeptidase and protease V8 digests, carried out on reduced and carboxymethylated samples, with the lysyl endopeptidase digest of GGF-II carried out on material eluted from the 55-65 RD region of a 11% SDS-PAGE (MW relative to the above-quoted markers).

A total of 21 peptide sequences (see Fig. 8, SEQ ID Nos. 1-20, 165) were obtained for GGF-I, of which 12 peptides (see Fig. 9, SEQ ID Nos. 1, 22-29, 17, 19, and 32) are not present in current protein databases and therefore represent unique sequences. A total of 12 peptide sequences (see Fig. 10, SEQ ID Nos. 42-50 and 161-163) were obtained for GGF-II, of which 10 peptides (see Fig. 11, SEQ ID Nos. 42-50) are not present in current protein databases and therefore represent unique sequences (an exception is peptide GGF-II 06 which shows identical sequences in many proteins which are probably of no significance given the small number of residues). These novel sequences are extremely likely to correspond to portions of the true amino acid sequences of GGFs I and II.

Particular attention can be drawn to the sequences of GGF-I 07 and GGF-II 12, which are clearly highly related. The similarities indicate that the sequences of these peptides are almost certainly those of the assigned GGF species, and are most unlikely to be derived from contaminant proteins.

In addition, in peptide GGF-II 02, the sequence X S S is consistent with the presence of an N linked

- 48 -

carbohydrate moiety on an asparagine at the position denoted by X.

In general, in Figs. 8 and 10, X represents an unknown residue denoting a sequencing cycle where a single position could not be called with certainty either because there was more than one signal of equal size in the cycle or because no signal was present. An asterisk denotes those peptides where the last amino acid called corresponds to the last amino acid present in that peptide. In the remaining peptides, the signal strength after the last amino acid called was insufficient to continue sequence calling to the end of that peptide. The right hand column indicates the results of a computer database search using the GCG package FASTA and TFASTA programs to analyze the NBRF and EMBL sequence databases. The name of a protein in this column denotes identity of a portion of its sequence with the peptide amino acid sequence called allowing a maximum of two mismatches. A question mark denotes three mismatches allowed. The abbreviations used are as follows:

HMG-1	High Mobility Group protein-1
HMG-2	High Mobility Group protein-2
LH-alpha	Luteinizing hormone alpha subunit
LH-beta	Luteinizing hormone beta subunit

25

**EXAMPLE 10****Isolating and Cloning of Nucleotide Sequences encoding proteins containing GGF-I and GGF-II peptides**

Isolation and cloning of the GGF-II nucleotide sequences was performed as outlined herein, using peptide sequence information and library screening, and was performed as set out below. It will be appreciated that the peptides of Figs. 10 and 11 can be used as the

30

- 49 -

starting point for isolation and cloning of GGF-I sequences by following the techniques described herein. Indeed, Fig. 20, SEQ ID Nos. 51-84) shows possible degenerate oligonucleotide probes for this purpose, and  
5 Fig. 22, SEQ ID Nos. 86-115, lists possible PCR primers. DNA sequence and polypeptide sequence should be obtainable by this means as with GGF-II, and also DNA constructs and expression vectors incorporating such DNA sequence, host cells genetically altered by incorporating  
10 such constructs/vectors, and protein obtainable by cultivating such host cells. The invention envisages such subject matter.

#### I. Design and Synthesis of oligonucleotide Probes and Primers

15 Degenerate DNA oligomer probes were designed by backtranslating the amino acid sequences (derived from the peptides generated from purified GGF protein) into nucleotide sequences. Oligomers represented either the coding strand or the non-coding strand of the DNA  
20 sequence. When serine, arginine or leucine were included in the oligomer design, then two separate syntheses were prepared to avoid ambiguities. For example, serine was encoded by either TCN or AGY as in 537 and 538 or 609 and 610. Similar codon splitting was done for arginine or  
25 leucine (e.g. 544, 545). DNA oligomers were synthesized on a Biosearch 8750 4-column DNA synthesizer using  $\beta$ -cyanoethyl chemistry operated at 0.2 micromole scale synthesis. Oligomers were cleaved off the column (500 angstrom CpG resins) and deprotected in concentrated  
30 ammonium hydroxide for 6-24 hours at 55-60°C. Deprotected oligomers were dried under vacuum (Speedvac) and purified by electrophoresis in gels of 15% acrylamide (20 mono : 1 bis), 50 mM Tris-borate-EDTA buffer

- 50 -

containing 7M urea. Full length oligomers were detected in the gels by UV shadowing, then the bands were excised and DNA oligomers eluted into 1.5 ml H<sub>2</sub>O for 4-16 hours with shaking. The eluate was dried, redissolved in 0.1 ml H<sub>2</sub>O and absorbance measurements were taken at 260nm.

Concentrations were determined according to the following formula:

$$(A_{260} \times \text{units/ml}) (60.6/\text{length} = x \mu\text{M})$$

All oligomers were adjusted to 50  $\mu\text{M}$  concentration by addition of H<sub>2</sub>O.

Degenerate probes designed as above are shown in Fig. 20, SEQ ID Nos. 54-88.

PCR primers were prepared by essentially the same procedures that were used for probes with the following modifications. Linkers of thirteen nucleotides containing restriction sites were included at the 5' ends of the degenerate oligomers for use in cloning into vectors. DNA synthesis was performed at 1 micromole scale using 1,000 angstrom CpG resins and inosine was used at positions where all four nucleotides were incorporated normally into degenerate probes. Purifications of PCR primers included an ethanol precipitation following the gel electrophoresis purification.

## 25 II. Library Construction and Screening

A bovine genomic DNA library was purchased from Stratagene (Catalogue Number: 945701). The library contained  $2 \times 10^6$  15-20kb Sau3A1 partial bovine DNA fragments cloned into the vector lambda DashII. A bovine total brain cDNA library was purchased from Clontech (Catalogue Number: BL 10139). Complementary DNA libraries were constructed (In Vitrogen; Stratagene) from mRNA prepared from bovine total brain, from bovine



- 51 -

pituitary and from bovine posterior pituitary. In Vitrogen prepared two cDNA libraries: one library was in the vector lambda g10, the other in vector pCDNAI (a plasmid library). The Stratagene libraries were prepared  
5 in the vector lambda unizap. Collectively, the cDNA libraries contained 14 million primary recombinant phage.

The bovine genomic library was plated on *E. coli* K12 host strain LE392 on 23 x 23 cm plates (Nunc) at 150,000 to 200,000 phage plaques per plate. Each plate  
10 represented approximately one bovine genome equivalent. Following an overnight incubation at 37°C, the plates were chilled and replicate filters were prepared according to procedures of Maniatis et al. (2:60-81). Four plaque lifts were prepared from each plate onto  
15 uncharged nylon membranes (Pall Biodyne A or MSI Nitropure). The DNA was immobilized onto the membranes by cross-linking under UV light for 5 minutes or, by baking at 80°C under vacuum for two hours. DNA probes were labelled using T4 polynucleotide kinase (New England  
20 Biolabs) with gamma 32P ATP (New England Nuclear; 6500 Ci/mmol) according to the specifications of the suppliers. Briefly, 50 pmols of degenerate DNA oligomer were incubated in the presence of 600 µCi gamma 32P-ATP and 5 units T4 polynucleotide kinase for 30 minutes at  
25 37°C. Reactions were terminated, gel electrophoresis loading buffer was added and then radiolabelled probes were purified by electrophoresis. 32P labelled probes were excised from gel slices and eluted into water. Alternatively, DNA probes were labelled via PCR  
30 amplification by incorporation of α-32P-dATP or α-32P dCTP according to the protocol of Schowalter and Sommer, Anal. Biochem 177:90-94 (1989). Probes labelled in PCR reactions were purified by desalting on Sephadex G-150 columns.

- 52 -

Prehybridization and hybridization were performed in GMC buffer (0.52 M NaPi, 7% SDS, 1% BSA, 1.5 mM EDTA, 0.1 M NaCl 10 mg/ml tRNA). Washing was performed in oligowash (160 ml 1 M Na<sub>2</sub>HPO<sub>4</sub>, 200 ml 20% SDS, 8.0 ml 0.5 M EDTA, 100 ml 5M NaCl, 3632 ml H<sub>2</sub>O). Typically, 20 filters (400 sq. centimeters each) representing replicate copies of ten bovine genome equivalents were incubated in 200 ml hybridization solution with 100 pmols of degenerate oligonucleotide probe (128-512 fold degenerate). Hybridization was allowed to occur overnight at 5°C below the minimum melting temperature calculated for the degenerate probe. The calculation of minimum melting temperature assumes 2°C for an AT pair and 4°C for a GC pair.

Filters were washed in repeated changes of oligowash at the hybridization temperatures four to five hours and finally, in 3.2M tetramethylammonium chloride, 1% SDS twice for 30 min at a temperature dependent on the DNA probe length. For 20mers, the final wash temperature was 60°C. Filters were mounted, then exposed to X-ray film (Kodak XAR5) using intensifying screens (Dupont Cronex Lightening Plus). Usually, a three to five day film exposure at minus 80°C was sufficient to detect duplicate signals in these library screens. Following analysis of the results, filters could be stripped and reprobated. Filters were stripped by incubating through two successive cycles of fifteen minutes in a microwave oven at full power in a solution of 1% SDS containing 10mM EDTA pH8. Filters were taken through at least three to four cycles of stripping and reprobating with various probes.

- 53 -

### III. Recombinant Phage Isolation, Growth and DNA Preparation

These procedures followed standard protocol as described in Recombinant DNA (Maniatis et al 2:60-2:81).

### 5 IV. Analysis of Isolated Clones Using DNA Digestion and Southern Blots

Recombinant Phage DNA samples (2 micrograms) were digested according to conditions recommended by the restriction endonuclease supplier (New England Biolabs).  
10 Following a four hour incubation at 37°C, the reactions products were precipitated in the presence of 0.1M sodium acetate and three volumes of ethanol. Precipitated DNA was collected by centrifugation, rinsed in 75% ethanol and dried. All resuspended samples were loaded onto  
15 agarose gels (typically 1% in TAE buffer; 0.04M Tris acetate, 0.002M EDTA). Gel runs were at 1 volt per centimeter from 4 to 20 hours. Markers included lambda Hind III DNA fragments and/or  $\phi$ X174HaeIII DNA fragments (New England Biolabs). The gels were stained with 0.5  
20 micrograms/ml of ethidium bromide and photographed. For southern blotting, DNA was first depurinated in the gel by treatment with 0.125 N HCl, denatured in 0.5 N NaOH and transferred in 20x SSC (3M sodium chloride, 0.03 M sodium citrate) to uncharged nylon membranes. Blotting  
25 was done for 6 hours up to 24 hours, then the filters were neutralized in 0.5 Tris HCl pH 7.5, 0.15 M sodium chloride, then rinsed briefly in 50 mM Tris-borate EDTA.

For cross-linking, the filters were wrapped first in transparent plastic wrap, then the DNA side exposed  
30 for five minutes to an ultraviolet light. Hybridization and washing was performed as described for library screening (see section 2 of this Example). For hybridization analysis to determine whether similar genes

- 54 -

exist in other species slight modifications were made. The DNA filter was purchased from Clontech (Catalogue Number 7753-1) and contains 5 micrograms of EcoRI digested DNA from various species per lane. The probe  
5 was labelled by PCR amplification reactions as described in section 2 above, and hybridizations were done in 80% buffer B (2 g polyvinylpyrrolidone, 2 g Ficoll-400, 2 g bovine serum albumin, 50 ml 1M Tris-HCl (pH 7.5) 58 g NaCl, 1 g sodium pyrophosphate, 10 g sodium dodecyl  
10 sulfate, 950ml H<sub>2</sub>O) containing 10% dextran sulfate. The probes were denatured by boiling for ten minutes then rapidly cooling in ice water. The probe was added to the hybridization buffer at 10<sup>6</sup> dpm <sup>32</sup>P per ml and incubated overnight at 60°C. The filters were washed at 60°C first  
15 in buffer B followed by 2X SSC, 0.1% SDS then in 1x SSC, 0.1% SDS. For high stringency, experiments, final washes were done in 0.1 x SSC, 1% SDS and the temperature raised to 65°C.

Southern blot data were used to prepare a  
20 restriction map of the genomic clone and to indicate which subfragments hybridized to the GGF probes (candidates for subcloning).

#### V. Subcloning of Segments of DNA Homologous to Hybridization Probes

25 DNA digests (e.g. 5 micrograms) were loaded onto 1% agarose gels then appropriate fragments excised from the gels following staining. The DNA was purified by adsorption onto glass beads followed by elution using the protocol described by the supplier (Bio 101). Recovered  
30 DNA fragments (100-200 ng) were ligated into linearized dephosphorylated vectors, e.g. pT3T7 (Ambion), which is a derivative of pUC18, using T4 ligase (New England Biolabs). This vector carries the *E. coli*  $\beta$  lactamase

- 55 -

gene, hence, transformants can be selected on plates containing ampicillin. The vector also supplies  $\beta$ -galactosidase complementation to the host cell, therefore non-recombinants (blue) can be detected using  
5 isopropylthiogalactoside and Bluogal (Bethesda Research Labs). A portion of the ligation reactions was used to transform *E. coli* K12 XL1 blue competent cells (Stratagene Catalogue Number: 200236) and then the transformants were selected on LB plates containing 50  
10 micrograms per ml ampicillin. White colonies were selected and plasmid mini preps were prepared for DNA digestion and for DNA sequence analysis. Selected clones were retested to determine if their insert DNA hybridized with the GGF probes.

#### 15 VI. DNA Sequencing

Double stranded plasmid DNA templates were prepared from 5 ml cultures according to standard protocols. Sequencing was by the dideoxy chain termination method using Sequenase 2.0 and a  
20 dideoxynucleotide sequencing kit (US Biochemical) according to the manufacturers protocol (a modification of Sanger et al. PNAS; USA 74:5463 (1977)). Alternatively, sequencing was done in a DNA thermal  
cycler (Perkin Elmer, model 4800) using a cycle  
25 sequencing kit (New England Biolabs; Bethesda Research Laboratories) and was performed according to manufacturers instructions using a 5'-end labelled primer. Sequence primers were either those supplied  
with the sequencing kits or were synthesized according to  
30 sequence determined from the clones. Sequencing reactions were loaded on and resolved on 0.4mm thick sequencing gels of 6% polyacrylamide. Gels were dried and exposed to X-Ray film. Typically, 35S was

- 56 -

incorporated when standard sequencing kits were used and a 32P end labelled primer was used for cycle sequencing reactions. Sequences were read into a DNA sequence editor from the bottom of the gel to the top (5' direction to 3') and data were analyzed using programs supplied by Genetics Computer Group (GCG, University of Wisconsin).

#### VII. RNA Preparation and PCR Amplification

Open reading frames detected in the genomic DNA and which contained sequence encoding GGF peptides were extended via PCR amplification of pituitary RNA. RNA was prepared from frozen bovine tissue (Pelfreeze) according to the guanidine neutral-CsCl procedure (Chirgwin et. al. Biochemistry 18:5294(1979).) Polyadenylated RNA was selected by oligo-dT cellulose column chromatography (Aviv and Leder PNAS (USA) 69:1408 (1972)).

Specific DNA target sequences were amplified beginning with either total RNA or polyadenylated RNA samples that had been converted to cDNA using the Perkin Elmer PCR/RNA Kit Number: N808-0017. First strand reverse transcription reactions used 1  $\mu$ g template RNA and either primers of oligo dT with restriction enzyme recognition site linkers attached or specific antisense primers determined from cloned sequences with restriction sites attached. To produce the second strand, the primers either were plus strand unique sequences as used in 3' RACE reactions (Frohman et. al., PNAS (USA) 85:8998 (1988)) or were oligo dT primers with restriction sites attached if the second target site had been added by terminal transferase tailing first strand reaction products with dATP (e.g. 5' race reactions, Frohman et. al., *ibid*). Alternatively, as in anchored PCR reactions

- 57 -

the second strand primers were degenerate, hence, representing particular peptide sequences.

The amplification profiles followed the following general scheme: 1) five minutes soak file at 95°C; 2) thermal cycle file of 1 minute, 95°C; 1 minute ramped down to an annealing temperature of 45°C, 50°C or 55°C; maintain the annealing temperature for one minute; ramp up to 72°C over one minute; extend at 72°C for one minute or for one minute plus a 10 second auto extension; 3) extension cycle at 72°C, five minutes, and; 4) soak file 4°C for infinite time. Thermal cycle files (#2) usually were run for 30 cycles. A sixteen  $\mu$ l sample of each 100  $\mu$ l amplification reaction was analyzed by electrophoresis in 2% Nusieve 1% agarose gels run in TAE buffer at 4 volts per centimeter for three hours. The gels were stained, then blotted to uncharged nylon membranes which were probed with labelled DNA probes that were internal to the primers.

Specific sets of DNA amplification products could be identified in the blotting experiments and their positions used as a guide to purification and reamplification. When appropriate, the remaining portions of selected samples were loaded onto preparative gels, then following electrophoresis four to five slices of 0.5 mm thickness (bracketing the expected position of the specific product) were taken from the gel. The agarose was crushed, then soaked in 0.5 ml of electrophoresis buffer from 2-16 hours at 40°C. The crushed agarose was centrifuged for two minutes and the aqueous phase was transferred to fresh tubes.

Reamplification was done on five microliters (roughly 1% of the product) of the eluted material using the same sets of primers and the reaction profiles as in the original reactions. When the reamplification

- 58 -

reactions were completed, samples were extracted with chloroform and transferred to fresh tubes. Concentrated restriction enzyme buffers and enzymes were added to the reactions in order to cleave at the restriction sites present in the linkers. The digested PCR products were purified by gel electrophoresis, then subcloned into vectors as described in the subcloning section above. DNA sequencing was done described as above.

#### VIII. DNA Sequence Analysis

DNA sequences were assembled using a fragment assembly program and the amino acid sequences deduced by the GCG programs GelAssemble, Map and Translate. The deduced protein sequences were used as a query sequence to search protein sequence databases using WordSearch. Analysis was done on a VAX Station 3100 workstation operating under VMS 5.1. The database search was done on SwissProt release number 21 using GCG Version 7.0.

#### IX. Results of Cloning and Sequencing of genes encoding GGF-I and GGF-II

As indicated above, to identify the DNA sequence encoding bovine GGF-II degenerate oligonucleotide probes were designed from GGF-II peptide sequences. GGF-II (SEQ ID No. 44), a peptide generated via lysyl endopeptidase digestion of a purified GGF-II preparation (see Figs. 16 and 17) showed strong amino acid sequence homology with GGF-I 07 (SEQ ID No. 39), a tryptic peptide generated from a purified GGF-I preparation. GGF-II 12 was thus used to create ten degenerate oligonucleotide probes (see oligos 609, 610 and 649 to 656 in Fig. 20, SEQ ID Nos. 66, 67, 68 and 75, respectively). A duplicate set of filters were probed with two sets (set 1=609, 610; set 2=649-656) of probes encoding two



- 59 -

overlapping portions of GGF-II 12. Hybridization signals were observed, but, only one clone hybridized to both probe sets. The clone (designated GGF2BG1) was purified.

Southern blot analysis of DNA from the phage clone  
5 GGF2BG1 confirmed that both sets of probes hybridized with that bovine DNA sequence, and showed further that both probes reacted with the same set of DNA fragments within the clone. Based on those experiments a 4 kb Eco  
10 RI sub-fragment of the original clone was identified, subcloned and partially sequenced. Fig. 21 shows the nucleotide sequence, SEQ ID No. 89) and the deduced amino acid sequence of the initial DNA sequence readings that included the hybridization sites of probes 609 and 650, and confirmed that a portion of this bovine genomic DNA  
15 encoded peptide 12 (KASLADSGEYM).

Further sequence analysis demonstrated that GGF-II 12 resided on a 66 amino acid open reading frame (see below) which has become the starting point for the isolation of overlapping sequences representing a  
20 putative bovine GGF-II gene and a cDNA.

Several PCR procedures were used to obtain additional coding sequences for the putative bovine GGF-II gene. Total RNA and oligo dT-selected (poly A containing) RNA samples were prepared from bovine total  
25 pituitary, anterior pituitary, posterior pituitary, and hypothalamus. Using primers from the list shown in Fig. 22, SEQ ID Nos. 109-119, one-sided PCR reactions (RACE) were used to amplify cDNA ends in both the 3' and 5' directions, and anchored PCR reactions were performed  
30 with degenerate oligonucleotide primers representing additional GGF-II peptides. Fig. 29 summarizes the contiguous DNA structures and sequences obtained in those experiments. From the 3' RACE reactions, three alternatively spliced cDNA sequences were produced,

- 60 -

which have been cloned and sequenced. A 5' RACE reaction led to the discovery of an additional exon containing coding sequence for at least 52 amino acids. Analysis of that deduced amino acid sequence revealed peptides GGF-II-6 and a sequence similar to GGF-I-18 (see below). The anchored PCR reactions led to the identification of (cDNA) coding sequences of peptides GGF-II-1, 2, 3 and 10 contained within an additional cDNA segment of 300 bp. The 5' limit of this segment (i.e., segment E, see Fig. 30) is defined by the oligonucleotide which encodes peptide GGF-II-1 and which was used in the PCR reaction (additional 5' sequence data exists as described for the human clone in Example 11). Thus this clone contains nucleotide sequences encoding six out of the existing total of nine novel GGF-II peptide sequences.

The cloned gene was characterized first by constructing a physical map of GGF2BG1 that allowed us to position the coding sequences as they were found (see below, Fig. 30). DNA probes from the coding sequences described above have been used to identify further DNA fragments containing the exons on this phage clone and to identify clones that overlap in both directions. The putative bovine GGF-II gene is divided into at least 5 coding segments. Coding segments are defined as discrete lengths of DNA sequence which can be translated into polypeptide sequences using the universal genetic code. The coding segments described in Fig. 36 and referred to in the present application are: 1) particular exons present within the GGF gene (e.g. coding segment a), or 2) derived from sets of two or more exons that appear in specific sub-groups of mRNAs, where each set can be translated into the specific polypeptide segments as in the gene products shown. The polypeptide segments

- 61 -

referred to in the claims are the translation products of the analogous DNA coding segments. Only coding segments A and B have been defined as exons and sequenced and mapped thus far. The summary of the contiguous coding sequences identified is shown in Fig. 31. The exons are listed (alphabetically) in the order of their discovery. It is apparent from the intron/exon boundaries that exon B may be included in cDNAs that connect coding segment E and coding segment A. That is, exon B cannot be spliced out without compromising the reading frame. Therefore, we suggest that three alternative splicing patterns can produce putative bovine GGF-II cDNA sequences 1, 2 and 3. The coding sequences of these, designated GGF2BPP1.CDS, GGF2BPP2.CDS and GGF2BPP3.CDS, respectively, are given in Figs. 27A (SEQ ID No. 129), 27B (SEQ ID No. 130), and 27C (SEQ ID No. 131), respectively. The deduced amino acid sequence of the three cDNAs is also given in Figs. 27A, (SEQ ID No. 129), 27B (SEQ ID No. 130), and 27C (SEQ ID No. 131).

The three deduced structures encode proteins of lengths 206, 281 and 257 amino acids. The first 183 residues of the deduced protein sequence are identical in all three gene products. At position 184 the clones differ significantly. A codon for glycine GGT in GGF2BPP1 also serves as a splice donor for GGF2BPP2 and GGF2BPP3, which alternatively add on exons C, C/D, C/D' and D or C, C/D and D, respectively, and shown in Fig. 32, SEQ ID No. 145). GGF2BPP1 is a truncated gene product which is generated by reading past the coding segment A splice junction into the following intervening sequence (intron). This represents coding segment A' in Fig. 30 (SEQ ID No. 136). The transcript ends adjacent to a canonical AATAAA polyadenylation sequence, and we suggest that this truncated gene product represents a

- 62 -

bona fide mature transcript. The other two longer gene products share the same 3' untranslated sequence and polyadenylation site.

5 All three of these molecules contain six of the nine novel GGF-II peptide sequences (see Fig. 11) and another peptide is highly homologous to GGF-I-18 (see Fig. 26). This finding gives a high probability that this recombinant molecule encodes at least a portion of bovine GGF-II. Furthermore, the calculated isoelectric points for the three peptides are consistent with the physical properties of GGF-I and II. Since the molecular size of GGF-II is roughly 60 kD, the longest of the three cDNAs should encode a protein with nearly one-half of the predicted number of amino acids.

15 A probe encompassing the B and A exons was labelled via PCR amplification and used to screen a cDNA library made from RNA isolated from bovine posterior pituitary. One clone (GGF2BPP5) showed the pattern indicated in Fig. 29 and contained an additional DNA coding segment (G) between coding segments A and C. The entire nucleic acid sequence is shown in Fig. 31 (SEQ ID No. 144). The predicted translation product from the longest open reading frame is 241 amino acids. A portion of a second cDNA (GGF2BPP4) was also isolated from the bovine posterior pituitary library using the probe described above. This clone showed the pattern indicated in Fig. 29. This clone is incomplete at the 5' end, but is a splicing variant in the sense that it lacks coding segments G and D. BPP4 also displays a novel 3' end with regions H, K and L beyond region C/D. The sequence of BPP4 is shown in Fig. 33 (SEQ ID No. 146).

#### EXAMPLE 11

#### GGF Sequences in Various Species

- 63 -

The GGF proteins are the members of a new superfamily of proteins. In high stringency cross hybridization studies (DNA blotting experiments) with other mammalian DNAs we have shown, clearly, that DNA probes from this bovine recombinant molecule can readily detect specific sequences in a variety of samples tested. A highly homologous sequence is also detected in human genomic DNA. The autoradiogram is shown in Fig. 28. The signals in the lanes containing rat and human DNA represent the rat and human equivalents of the GGF gene, the sequences of several cDNA's encoded by this gene have been recently reported by Holmes et al. (Science 256: 1205 (1992)) and Wen et al. (Cell 69: 559 (1992)).

#### EXAMPLE 12

##### 15 Isolation of a Human Sequence Encoding Human GGF2

Several human clones containing sequences from the bovine GGFII coding segment E were isolated by screening a human cDNA library prepared from brain stem (Stratagene catalog #935206). This strategy was pursued based on the strong link between most of the GGF2 peptides (unique to GGF2) and the predicted peptide sequence from clones containing the bovine E segment. This library was screened as described in Example 8, Section II using the oligonucleotide probes 914-919 listed below.

25	914TCGGGCTCCATGAAGAAGATGTA	(SEQ ID NO: 179)
	915TCCATGAAGAAGATGTACCTGCT	(SEQ ID NO: 180)
	916ATGTACCTGCTGTCCTCCTTGA	(SEQ ID NO: 181)
	917TTGAAGAAGGACTCGCTGCTCA	(SEQ ID NO: 182)
	918AAAGCCGGGGGCTTGAAGAA	(SEQ ID NO: 183)
30	919ATGARGTGTGGGCGCGAAA	(SEQ ID NO: 184)

Clones detected with these probes were further analyzed by hybridization. A probe derived from coding segment A (see Fig. 30), which was produced by labeling a

- 64 -

polymerase chain reaction (PCR) product from segment A, was also used to screen the primary library. Several clones that hybridized with both A and E derived probes were selected and one particular clone, GGF2HBS5, was selected for further analysis. This clone is represented by the pattern of coding segments (EBACC/D'D as shown in Fig. 30). The E segment in this clone is the human equivalent of the truncated bovine version of E shown in Fig. 30. GGF2HBS5 is the most likely candidate to encode GGF-II of all the "putative" GGF-II candidates described. The length of coding sequence segment E is 786 nucleotides plus 264 bases of untranslated sequence. The predicted size of the protein encoded by GGF2HBS5 is approximately 423 amino acids (approximately 45 kilodaltons, see Fig. 44, SEQ ID NO: 21), which is similar to the size of the deglycosylated form of GGF-II (see Example 20). Additionally, seven of the GGF-II peptides listed in Fig. 26 have equivalent sequences which fall within the protein sequence predicted from region E. Peptides II-6 and II-12 are exceptions, which fall in coding segment B and coding segment A, respectively. RNA encoding the GGF2HBS5 protein was produced in an in vitro transcription system driven by the bacteriophage T7 promoter resident in the vector (Bluescript SK [Stratagene Inc.] see Fig. 47) containing the GGF2HBS5 insert. This RNA was translated in a cell free (rabbit reticulocyte) translation system and the size of the protein product was 45 Kd. Additionally, the cell-free product has been assayed in a Schwann cell mitogenic assay to confirm biological activity. Schwann cells treated with conditioned medium show both increased proliferation as measured by incorporation of <sup>125</sup>I-Uridine and phosphorylation on tyrosine of a protein in the 185 kilodalton range.

- 65 -

Thus the size of the product encoded by GGF2HBS5 and the presence of DNA sequences which encode human peptides highly homologous to the bovine peptides shown in Fig. 11 confirm that GGF2HBS5 encodes the human equivalent of bovine GGF2. The fact that conditioned media prepared from cells transformed with this clone elicits Schwann cell mitogenic activity confirms that the GGFIHBS5 gene product (unlike the BPP5 gene product) is secreted. Additionally the GGFIIBPP5 gene product seems to mediate the Schwann cell proliferation response via a receptor tyrosine kinase such as p185<sup>erbB2</sup> or a closely related receptor (see Example 19).

#### EXAMPLE 13

#### Expression of Human Recombinant GGF2 in Mammalian and Insect Cells

The GGF2HBS5 cDNA clone encoding human GGF2 (as described in Example 12 and also referred to herein as HBS5) was cloned into vector pcDL-SR $\alpha$ 296 and COS-7 cells were transfected in 100 mm dishes by the DEAE-dextran method. Cell lysates or conditioned media from transiently expressing COS cells were harvested at 3 or 4 days post-transfection. To prepare lysates, cell monolayers were washed with PBS, scraped from the dishes lysed by three freeze/thaw cycles in 150  $\mu$ m of 0.25 M Tris-HCl, pH8. Cell debris was pelleted and the supernatant recovered. Conditioned media samples (7 mls.) were collected, then concentrated and buffer exchanged with 10 mM Tris, pH 7.4 using Centiprep-10 and Centricon-10 units as described by the manufacturer (Amicon, Beverly, MA). Rat nerve Schwann cells were assayed for incorporation of DNA synthesis precursors, as described. Conditioned media or cell lysate samples were tested in the Schwann cell proliferation assay as described in Marchionni et al., *Nature* 362:313 (1993).

- 66 -

The cDNA, GGF2HBS5, encoding GGF2 directed the secretion of the protein product to the medium. Minimal activity was detectable inside the cells as determined by assays using cell lysates. GGF2HFB1 and GGFBPP5 cDNA's failed to direct the secretion of the product to the extracellular medium. GGF activity from these clones was detectable only in cell lysates.

Recombinant GGF2 was also expressed in CHO cells. The GGF2HBS5 cDNA encoding GGF2 was cloned into the EcoRI site of vector pcdhfrpolyA and transfected into the DHFR negative CHO cell line (GG44) by the calcium phosphate coprecipitation method. Clones were selected in nucleotide and nucleoside free  $\alpha$  medium (Gibco) in 96-well plates. After 3 weeks, conditioned media samples from individual clones were screened for expression of GGF by the Schwann cell proliferation assay as described in Marchionni et al., Nature 362:313 (1993). Stable clones which secreted significant levels of GGF activity into the medium were identified. Schwann cell proliferation activity data from different volume aliquots of CHO cell conditioned medium were used to produce the dose response curve shown in Fig. 46 (Graham and Van Der Eb, Virology 52:456, 1973). This material was analyzed on a Western blot probed with polyclonal antisera raised against a GGF2 specific peptide. A band of approximately 65 Kd (the expected size of GGF2 extracted from pituitary) is specifically labeled (Fig. 48, lane 12).

Recombinant GGF2 was also expressed in insect cells using the Baculovirus expression. Sf9 insect cells were infected with baculovirus containing the GGF2HBS5 cDNA clone at a multiplicity of 3-5 ( $10^6$  cells/ml) and cultured in Sf900-II medium. Schwann cell mitogenic activity was secreted into the extracellular medium.



- 67 -

Different volumes of insect cell conditioned medium were tested in the Schwann cell proliferation assay in the absence of forskolin and the data used to produce a dose response curve.

5           This material was also analyzed on a Western blot (Fig. 45B) probed with the GGF II specific antibody described above.

          The methods used in this example were as follows:

          Schwann cell mitogenic activity of recombinant  
10 human and bovine glial growth factors was determined as follows: Mitogenic responses of cultured Schwann cells were measured in the presence of 5  $\mu$ M forskolin using crude recombinant GGF preparations obtained from transient mammalian expression experiments.  
15 Incorporation of [<sup>125</sup>I]-Urd was determined following an 18-24 hour exposure to materials obtained from transfected or mock transfected cos cells as described in the Methods. The mean and standard deviation of four sets of data are shown. The mitogenic response to  
20 partially purified native bovine pituitary GGF (carboxymethyl cellulose fraction; Goodearl et al., submitted) is shown (GGF) as a standard of one hundred percent activity.

          cDNAs (Fig. 46, SEQ ID NOs. 166-168) were cloned  
25 into pcDL-SR $\alpha$ 296 (Takebe et al., Mol. Cell Biol. 8:466-472 (1988)), and COS-7 cells were transfected in 100 mm dishes by the DEAE-dextran method (Sambrook et al., In *Molecular Cloning. A Laboratory Manual*, 2nd. ed. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY,  
30 1989)). Cell lysates or conditioned media were harvested at 3 or 4 days post-transfection. To prepare lysates, cell monolayers were washed with PBS, scraped from the dishes, and lysed by three freeze/thaw cycles in 150  $\mu$ l of 0.25 M Tris-HCl, pH 8. Cell debris was pelleted and

- 68 -

the supernate recovered. Conditioned media samples (7  
mls) were collected, then concentrated and buffer  
exchanged with 10 mM Tris, pH 7.4 using Centriprep-10 and  
Centricon-10 units are described by the manufacturers  
5 (Amicon, Beverly, MA). Rat sciatic nerve Schwann cells  
were assayed for incorporation of DNA synthesis  
precursors, as described (Davis and Stroobant, J. Cell  
Biol. 110:1353-1360 (1990); Brockes et al., Brain Res.  
165:105-118 (1979)).

10 Western blot of recombinant CHO cell conditioned  
medium were performed as follows: A recombinant CHO  
clone was cultured in MCDB302 protein-free for 3 days. 2  
ml of conditioned medium was harvested, concentrated,  
buffered exchanged against 10 mM Tris-HCl, pH 7.4 and  
15 lyophilized to dryness. The pellet was resuspended in  
SDS-PAGE sample buffer, subjected to reducing SDS gel  
electrophoresis and analyzed by Western blotting with a  
GGF peptide antibody. A CHO control was done by using  
conditioned medium from untransfected CHO-DG44 host and  
20 the CHO HBS5 levels were assayed using conditioned medium  
from a recombinant clone.

#### EXAMPLE 14

##### Identification of Functional Elements of GGF

The deduced structures of the family of GGF  
25 sequences indicate that the longest forms (as represented  
by GGF2BPP4) encode transmembrane proteins where the  
extracellular part contains a domain which resembles  
epidermal growth factor (see Carpenter and Wahl in  
Peptide Growth Factors and Their Receptors I pp. 69-133,  
30 Springer-Verlag, NY 1991). The positions of the cysteine  
residues in coding segments C and C/D or C/D' peptide  
sequence are conserved with respect to the analogous  
residues in the epidermal growth factor (EGF) peptide

- 69 -

sequence (see Fig. 32, SEQ ID Nos. 147-149). This suggests that the extracellular domain functions as receptor recognition and biological activation sites. Several of the variant forms lack the H, K, and L coding segments and thus may be expressed as secreted, diffusible biologically active proteins. GGF DNA sequences encoding polypeptides which encompass the EGF-like domain (EGFL) can have full biological activity for stimulating glial cell mitogenic activity.

10 Membrane bound versions of this protein may induce Schwann cell proliferation if expressed on the surface of neurons during embryogenesis or during nerve regeneration (where the surfaces of neurons are intimately associated with the surfaces of proliferating Schwann cells).

15 Secreted (non membrane bound) GGFs may act as classically diffusible factors which can interact with Schwann cells at some distance from their point of secretion. Other forms may be released from intracellular sources via tissue injury and cell disruption. An example of a secreted GGF is the protein encoded by GGF2HBS5; this is the only GGF known which has been found to be directed to the exterior of the cell. Secretion is probably mediated via an N-terminal hydrophobic sequence found only in region E, which is the N-terminal domain contained within recombinant GGF2 encoded by GGF2HBS5.

20  
25 Other GGF's appear to be non-secreted. These GGFs may be injury response forms which are released as a consequence of tissue damage.

30 Other regions of the predicted protein structure of GGF2 (encoded by GGF2HBS5) and other proteins containing regions B and A exhibit similarities to the human basement membrane heparan sulfate proteoglycan core protein. The peptide ADSGEY, which is located next to the second cysteine of the C2 immunoglobulin fold in

- 70 -

these GGF's, occurs in nine of twenty-two C-2 repeats found in that basal lamina protein. This evidence strongly suggests that these proteins may associate with matrix proteins such as those associated with neurons and glia, and may suggest a method for sequestration of glial growth factors at target sites.

#### EXAMPLE 15

##### Purification of GGFs from Recombinant Cells

In order to obtain full length or portions of GGFs to assay for biological activity, the proteins can be overproduced using cloned DNA. Several approaches can be used. A recombinant *E. coli* cell containing the sequences described above can be constructed. Expression systems such as pNH8a or pHH16a (Stratagene, Inc.) can be used for this purpose by following manufacturers procedures. Alternatively, these sequences can be inserted in a mammalian expression vector and an overproducing cell line can be constructed. As an example, for this purpose DNA encoding a GGF, clone GGF2BPP5 has been expressed in COS cells and can be expressed in Chinese hamster ovary cells using the pMSXND expression vector (Lee and Nathans, J. Biol. Chem. 263, 3521-3527, (1981)). This vector containing GGF DNA sequences can be transfected into host cells using established procedures.

Transient expression can be examined or G418-resistant clones can be grown in the presence of methotrexate to select for cells that amplify the *dhfr* gene (contained on the pMSXND vector) and, in the process, co-amplify the adjacent GGF protein encoding sequence. Because CHO cells can be maintained in a totally protein-free medium (Hamilton and Ham, In Vitro 13, 537-547 (1977)), the desired protein can be purified

- 71 -

from the medium. Western analysis using the antisera produced in Example 17 can be used to detect the presence of the desired protein in the conditioned medium of the overproducing cells.

5           The desired protein (rGGF2) was purified from the medium conditioned by transiently expressing cos cells as follows. rGGF II was harvested from the conditioned medium and partially purified using Cation Exchange  
10           Chromatography (POROS-HS). The column was equilibrated with 33.3 mM MES pH 6.0. Conditioned media was loaded at flow rate of 10 ml/min. The peak containing Schwann cell proliferation activity and immunoreactive (using the polyclonal antisera was against a GGF2 peptide described above) was eluted with 50 mM Tris, 1M NaCl pH 8.0.

15           rhGGF2 is also expressed using a stable Chinese Ovary Hamster cell line. rGGF2 from the harvested conditioned media was partially purified using Cation Exchange Chromatograph (POROS-HS). The column was equilibrated with PBS pH 7.4. Conditioned media was  
20           loaded at 10 ml/min. The peak containing the Schwann Cell Proliferative activity and immunoreactivity (using GGF2 polyclonal antisera) was eluted with 50 mM Hepes, 500 mM NaCl pH 8.0. An additional peak was observed at 50 mM Hepes, 1M NaCl pH 8.0 with both proliferation as  
25           well as immunoreactivity (Fig. 45).

          rhGGF2 can be further purified using Hydrophobic Interaction Chromatography as a high resolution step; Cation exchange/Reserve phase Chromatography (if needed as second high resolution step); A viral inactivation  
30           step and a DNA removal step such as Anion exchange chromatography.

          Schwann Cell Proliferation Activity of recombinant GGF2 peak eluted from the Cation Exchange column was determined as follows: Mitogenic responses of the

- 72 -

cultured Schwann cells were measured in the presence of 5 M Forskolin using the peak eluted by 50 mM Tris 1 M NaCl pH 8.0. The peak was added at 20  $\mu$ l, 10  $\mu$ l (1:10) 10  $\mu$ l and (1:100) 10  $\mu$ l. Incorporation of  $^{125}$ I-Uridine was  
5 determined and expressed as (CPM) following an 18-24 hour exposure.

An immunoblot using polyclonal antibody raised against a peptide of GGF2 was carried out as follows: 10  $\mu$ l of different fractions were ran on 4-12% gradient gels.  
10 The gels were transferred on to Nitrocellulose paper, and the nitrocellulose blots were blocked with 5% BSA and probed with GGF2-specific antibody (1:250 dilution).  $^{125}$ I protein A (1:500 dilution, Specific Activity = 9.0/Ci/g) was used as the secondary antibody. The immunoblots were  
15 exposed to Kodax X-Ray films for 6 hours. The peak fractions eluted with 1 M NaCl showed an immunoreactive band at 69K.

GGF2 purification on cation exchange columns was performed as follows: CHO cell conditioned media  
20 expressing rGGFII was loaded on the cation exchange column at 10 ml/min. The column was equilibrated with PBS pH 7.4. The elution was achieved with 50 mM Hepes 500 mM NaCl pH 8.0 and 50 mM Hepes 1M NaCl pH 8.0 respectively. All fractions were analyzed using the  
25 Schwann cell proliferation assay (CPM) described herein. The protein concentration (mg/ml) was determined by the Bradford assay using BSA as the standard.

A Western blot using 10  $\mu$ l of each fraction was performed and immunoreactivity and the Schwann cell  
30 activity were observed to co-migrate.

The protein may be assayed at various points in the procedure using a Western blot assay. Alternatively, the Schwann cell mitogenic assay described herein may be used to assay the expressed product of the full length

- 73 -

clone or any biologically active portions thereof. The full length clone GGF2BPP5 has been expressed transiently in COS cells. Intracellular extracts of transfected COS cells show biological activity when assayed in the Schwann cell proliferation assay described in Example 8. In addition, the full length close encoding GGF2HBS5 has been expressed transiently in COS cells. In this case both cell extract and conditioned media show biological activity in the Schwann cell proliferation assay described in Example 8. Any member of the family of splicing variant complementary DNA's derived from the GGF gene (including the Heregulins) can be expressed in this manner and assayed in the Schwann cell proliferation assay by one skilled in the art.

Alternatively, recombinant material may be isolated from other variants according to Wen et al. (Cell 69:559 (1992)) who expressed the splicing variant Neu differentiation factor (NDF) in COS-7 cells. cDNA clones inserted in the pJT-2 eukaryotic plasmid vector are under the control of the SV40 early promoter, and are 3'-flanked with the SV40 termination and polyadenylation signals. COS-7 cells were transfected with the pJT-2 plasmid DNA by electroporation as follows:  $6 \times 10^6$  cells (in 0.8 ml of DMEM and 10% FBS) were transferred to a 0.4 cm cuvette and mixed with 20  $\mu$ g of plasmid DNA in 10  $\mu$ l of TE solution (10 mM Tris-HCl (pH 8.0), 1 mM EDTA). Electroporation was performed at room temperature at 1600 V and 25  $\mu$ F using a Bio-Rad Gene Pulser apparatus with the pulse controller unit set at 200 ohms. The cells were then diluted into 20 ml of DMEM, 10% FBS and transferred into a T75 flask (Falcon). After 14 hr. of incubation at 37°C, the medium was replaced with DMEM, 1% FBS, and the incubation continued for an additional 48 hr. Conditioned medium containing recombinant protein

- 74 -

which was harvested from the cells demonstrated biological activity in a cell line expressing the receptor for this protein. This cell line (cultured human breast carcinoma cell line AU 565) was treated with recombinant material. The treated cells exhibited a morphology change which is characteristic of the activation of the erbB2 receptor. Conditioned medium of this type also can be tested in the Schwann cell proliferation assay.

10

**EXAMPLE 16****N-terminal sequence analysis**

The cDNA encoding hGGF2 was cloned into the amplifiable vector pcdhfrpolyA and transfected into CHO-DG44 cells for stable expression. rhGGF2 is secreted into the conditioned media. The ability of the recombinant GGF2 to be secreted is presumably mediated through the N-terminal hydrophobic stretch (signal sequence). A signal sequence, once having initiated export of a growing protein chain across the rough endoplasmic reticulum, is cleaved from the mature protein at a specific site. N-terminal sequence analysis of the expressed and purified rhGGF2 indicates the site of cleavage as shown below. The sequence of the first 50 amino acid residues at the N-terminus of the protein was



- 75 -

confirmed by N-terminal sequence analysis (Table 5),  
below.

TABLE 5

## N-terminal sequence analysis of rhGGF2

Cycle #	Primary Sequence	pMoles
1	Gly (G)	210.6
2	Asn (N)	163
3	GLU (E)	149
4	Ala (A)	220
5	Ala (A)	180
6	Pro (P)	173
7	Ala (A)	177
8	Gly (G)	154.9
9	Ala (A)	162.4
10	Ser (S)	65.4
11	Val (V)	132.7
12	Val (V) *(Cys)	11.7
13	Tyr (Y)	112.7
14	Ser (S)	47.6
15	Ser (S)	27.1

The N-terminal sequence analysis is performed by Edman Degradation Process

\*Cys residues are destroyed by the Edman Degradation Process and cannot be detected

The following sequence (SEQ ID NO: 185) represents the amino acid sequence of hGGF2. The shaded area indicates the cleaved signal sequence.

~~MTTRRAPPRS GRGGPRAQRF GSAARSEPPV PLEPMDDEIG TAAIAPGAA~~  
GNEAAPAGAS VCYSSPPSVG SVQELAQRRA VVIEGKVHPQ RRQOGALDRK  
AAAAGEAGA WGGDREPPAA GPRALGPPAE EPLLANGTV PSWPTAPVPS

- 76 -

AGEPGEEAPY LVKVHQVWAV KAGGLKKDSL LTVRLGTWGH FAFPPSCGRLK  
EDSRYIFFME PDANSTSRAP AAFRASFPPL ETGRNLKKEV SRVLCKRCAL  
PPQLKEMKSQ ESAAGSKLVL RCETSSEYSS LRFKWFKNGN ELNRKKNKPN  
IKIQKPKGKS ELRINKASLA DSGEYMCKVI SKLGNDSASA NITIVESNAT  
5 STSTTGTSHL VKCAEKETF CVNGGECFMV KDLSNPSRYL CKCPNEFTGD  
RCQNYVMASF YSTSTPFLSL PE (SEQ ID NO: 185)

The shaded area represents experimentally determined 15  
amino acid residues at the N-terminal of the rhGGF2,  
indicating A<sub>50</sub>-G<sub>51</sub> bond to be the cleavage site for the  
10 signal sequence.

#### EXAMPLE 17

##### Isolation of a Further Splicing Variant

Methods for updating other neuregulins described  
in U.S. patent application Serial No. 07/965,173, filed  
15 October 23, 1992, incorporated herein by reference,  
produced four closely related sequences (heregulin  $\alpha$ ,  $\beta$ 1,  
 $\beta$ 2,  $\beta$ 3) which arise as a result of splicing variation.  
Peles et al. (Cell 69:205 (1992)), and Wen et al. (Cell  
69:559 (1992)) have isolated another splicing variant  
20 (from rat) using a similar purification and cloning  
approach to that described in Examples 1-9 and 11  
involving a protein which binds to p185<sup>erbB2</sup>. The cDNA  
clone was obtained as follows (via the purification and  
sequencing of a p185<sup>erbB2</sup> binding protein from a  
25 transformed rat fibroblast cell line).  
A p185<sup>erbB2</sup> binding protein was purified from conditioned  
medium as follows. Pooled conditioned medium from three  
harvests of 500 roller bottles (120 liters total) was  
cleared by filtration through 0.2  $\mu$  filters and  
30 concentrated 31-fold with a Pelicon ultrafiltration  
system using membranes with a 20kd molecular size cutoff.  
All the purification steps were performed by using a

- 77 -

Pharmacia fast protein liquid chromatography system. The concentrated material was directly loaded on a column of heparin-Sepharose (150 ml, preequilibrated with phosphate-buffered saline (PBS)). The column was washed  
5 with PBS containing 0.2 M NaCl until no absorbance at 280 nm wavelength could be detected. Bound proteins were then eluted with a continuous gradient (250 ml) of NaCl (from 0.2 M to 1.0 M), and 5 ml fractions were collected. Samples (0.01 ml of the collected fractions were used for  
10 the quantitative assay of the kinase stimulatory activity. Active fractions from three column runs (total volume = 360 ml) were pooled, concentrated to 25 ml by using a YM10 ultrafiltration membrane (Amicon, Danvers, MA), and ammonium sulfate was added to reach a  
15 concentration of 1.7 M. After clearance by centrifugation (10,000 x g, 15 min.), the pooled material was loaded on a phenyl-Superose column (HR10/10, Pharmacia). The column was developed with a 45 ml gradient of  $(\text{NH}_4)_2\text{SO}_4$  (from 1.7 M to no salt) in 0.1 M  
20  $\text{Na}_2\text{PO}_4$  (pH 7.4), and 2 ml fractions were collected and assayed (0.002 ml per sample) for kinase stimulation (as described in Example 19). The major peak of activity was pooled and dialyzed against 50 mM sodium phosphate buffer (pH 7.3). A Mono-S cation-exchange column (HR5/5,  
25 Pharmacia) was preequilibrated with 50 mM sodium phosphate. After loading the active material (0.884 mg of protein; 35 ml), the column was washed with the starting buffer and then developed at a rate of 1 ml/min. with a gradient of NaCl. The kinase stimulatory activity  
30 was recovered at 0.45-0.55 M salt and was spread over four fractions of 2 ml each. These were pooled and loaded directly on a  $\text{Cu}^{+2}$  chelating columns (1.6 ml, HR2/5 chelating Superose, Pharmacia). Most of the proteins adsorbed to the resin, but they gradually eluted with a

- 78 -

30 ml linear gradient of ammonium chloride (0-1 M). The activity eluted in a single peak of protein at the range of 0.05 to 0.2 M  $\text{NH}_4\text{Cl}$ . Samples from various steps of purification were analyzed by gel electrophoresis  
5 followed by silver staining using a kit from ICN (Costa Mesa, CA), and their protein contents were determined with a Coomassie blue dye binding assay using a kit from Bio-Rad (Richmond, CA).

The p44 protein (10  $\mu\text{g}$ ) was reconstituted in 200  
10  $\mu\text{l}$  of 0.1 M ammonium bicarbonate buffer (pH 7.8). Digestion was conducted with L-1-tosyl-amide 2-phenylethyl chloromethyl ketone-treated trypsin (Serva) at 37°C for 18 hr. at an enzyme-to-substrate ratio of 1:10. The resulting peptide mixture was separated by  
15 reverse-phase HPLC and monitored at 215 nm using a Vydac C4 micro column (2.1 mm i.d. x 15 cm, 300 Å) and an HP 1090 liquid chromatographic system equipped with a diode-array detector and a workstation. The column was equilibrated with 0.1% trifluoroacetic acid (mobile phase  
20 A), and elution was effected with a linear gradient from 0%-55% mobile phase B (90% acetonitrile in 0.1% trifluoroacetic acid) over 70 min. The flow rate was 0.2 ml/min. and the column temperature was controlled at 25°C. One-third aliquots of the peptide peaks collected  
25 manually from the HPLC system were characterized by N-terminal sequence analysis by Edman degradation. The fraction eluted after 27.7 min. (T27.7) contained mixed amino acid sequences and was further rechromatographed after reduction as follows: A 70% aliquot of the peptide  
30 fraction was dried in vacuo and reconstituted in 100  $\mu\text{l}$  of 0.2 M ammonium bicarbonate buffer (pH 7.8). DTT (final concentration 2 mM) was added to the solution, which was then incubated at 37°C for 30 min. The reduced peptide mixture was then separated by reverse-phase HPLC

- 79 -

using a Vydac column (2.1 mm i.d. x 15 cm). Elution conditions and flow rate were identical to those described above. Amino acid sequence analysis of the peptide was performed with a Model 477 protein sequencer (Applied Biosystems, Inc., Foster City, CA) equipped with an on-line phenylthiohydantoin (PTH) amino acid analyzer and a Model 900 data analysis system (Hunkapiller et al. (1986) In Methods of Protein Microcharacterization, J.E. Shively, ed. (Clifton, New Jersey: Humana Press p. 223-247). The protein was loaded onto a trifluoroacetic acid-treated glass fiber disc precycled with polybrene and NaCl. The PTH-amino acid analysis was performed with a micro liquid chromatography system (Model 120) using dual syringe pumps and reverse-phase (C-18) narrow bore columns (Applied Biosystems, 2.1 mm x 250 mm). RNA was isolated from Rat1-EJ cells by standard procedures (Maniatis et al., Molecular Cloning: A Laboratory Manual (Cold Spring Harbor, New York (1982) and poly (A)<sup>+</sup> was selected using an mRNA Separator kit (Clontech Lab, Inc., Palo Alto, CA). cDNA was synthesized with the Superscript kit (from BRL Life Technologies, Inc., Bethesda, MD). Column-fractionated double-strand cDNA was ligated into an Sall- and NotI-digested pJT-2 plasmid vector, a derivative of the pCD-X vector (Okayama and Berg, Mol. Cell Biol. 3: 280 (1983)) and transformed into DH10B E. coli cells by electroporation (Dower et al., Nucl. Acids Res. 16: 6127 (1988)). Approximately  $5 \times 10^5$  primary transformants were screened with two oligonucleotide probes that were derived from the protein sequences of the N-terminus of NDF (residues 5-24) and the T40.4 tryptic peptide (residues 7-12). Their respective sequences were as follows (N indicates all 4 nt):

- 80 -

(1) 5'-ATA GGG AAG GGC GGG GGA AGG GTC NCC CTC NGC

A T

AGG GCC GGG CTT GCC TCT GGA GCC TCT-3'

(2) 5'-TTT ACA CAT ATA TTC NCC-3'

5 C G G C

(1: SEQ ID No. 163; 2: SEQ ID No. 164)

The synthetic oligonucleotides were end-labeled with [ $\gamma$ - $^{32}$ P]ATP with T4 polynucleotide kinase and used to screen replicate sets of nitrocellulose filters. The hybridization solution contained 6 x SSC, 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 2 x Denhardt's solution, 50  $\mu$ g/ml salmon sperm DNA, and 20% formamide (for probe 1) or no formamide (for probe 2). The filters were washed at either 50°C with 0.5 x SSC, 0.2% SDS, 2 mM EDTA (for probe 1) or at 37°C with 2 x SSC, 0.2% SDS, 2 mM EDTA (for probe 2). Autoradiography of the filters gave ten clones that hybridized with both probes. These clones were purified by replating and probe hybridization as described above.

The cDNA clones were sequenced using an Applied Biosystems 373A automated DNA sequencer and Applied Biosystems Taq DyeDeoxy<sup>™</sup> Terminator cycle sequencing kits following the manufacture's instructions. In some instances, sequences were obtained using [ $^{35}$ S]dATP (Amersham) and Sequenase<sup>™</sup> kits from U.S. Biochemicals following the manufacturer's instructions. Both strands of the cDNA clone 44 were sequenced by using synthetic oligonucleotides as primers. The sequence of the most 5' 350 nt was determined in seven independent cDNA clones.

The resultant clone demonstrated the pattern shown in Fig. 27 (NDF).

- 89 -

Xaa Ile Lys Gly Glu His Pro Gly Leu Ser Ile Gly Asp Val Ala Lys  
 1 5 10 15

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 9:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine and Xaa in position 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Xaa Met Ser Glu Tyr Ala Phe Phe Val Gln Thr Xaa Arg  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 10:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Xaa Ser Glu His Pro Gly Leu Ser Ile Gly Asp Thr Ala Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 11:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine; Xaa in position 8 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Xaa Ala Gly Tyr Phe Ala Glu Xaa Ala Arg  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 12:

SUBSTITUTE SHEET (RULE 26)

- 90 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine; Xaa in position 7 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Xaa Lys Leu Glu Phe Leu Xaa Ala Lys  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 13:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Xaa Thr Thr Glu Met Ala Ser Glu Gln Gly Ala  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 14:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Xaa Ala Lys Glu Ala Leu Ala Ala Leu Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 15:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear





(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: Xaa in position 2 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Glu Xaa Lys Phe Tyr Val Pro  
 1 5

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Glu Leu Ser Phe Ala Ser Val Arg Leu Pro Gly Cys Pro Pro Gly Val  
 1 5 10 15

Asp Pro Met Val Ser Phe Pro Val Ala Leu  
 20 25

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2003
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: N in positions 31 and 32 could be either A or G.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

GGAATTCCTT TTTTTTTTTT TTTTTTCTT MNTTTTTTTT TGCCCTTATA CCTCTTCGCC	60
TTTCTGTGGT TCCATCCACT TCTTC <del>CC</del> CCCT CCTCCTCCCA TAAACAAC TCCTACCCCT	120
GCACCCCAA TAAATAAATA AAAGGAGGAG GGCAAGCGGG GAGGAGGAGG AGTGGTGCTG	180
CGAGGGGAAG GAAAGGGGAG GCAGCGCGAG AAGAGCCGGG CAGAGTCCGA ACCGACAGCC	240
AGAAGCCCGC ACGCACCTCG CACC ATG AGA TGG CGA CGC GCC CCG CGC CGC	291
Met Arg Trp Arg Arg Ala Pro Arg Arg	
1 5	
TCC GCG CGT CCC GGC CCC CGG GCC CAG CGC CCC GGC TCC GCC GCC CGC	339
Ser Gly Arg Pro Gly Pro Arg Ala Gln Arg Pro Gly Ser Ala Ala Arg	
10 15 20 25	

TCG TCG CCG CCG CTG CCG CTG CTG CCA CTA CTG CTG CTG CTG GGG ACC	387
Ser Ser Pro Pro Leu Pro Leu Leu Pro Leu Leu Leu Leu Leu Gly Thr	
30 35 40	
GCG GCC CTG GCG CCG GGG GCG GCG GCC GGC AAC GAG GCG GCT CCC GCG	435
Ala Ala Leu Ala Pro Gly Ala Ala Ala Gly Asn Glu Ala Ala Pro Ala	
45 50 55	
GGG GCC TCG GTG TGC TAC TCG TCC CCG CCC AGC GTG GGA TCG GTG CAG	483
Gly Ala Ser Val Cys Tyr Ser Ser Pro Pro Ser Val Gly Ser Val Gln	
60 65 70	
GAG CTA GCT CAG CGC GCC GCG GTG GTG ATC GAG GGA AAG GTG CAC CCG	531
Glu Leu Ala Gln Arg Ala Ala Val Val Ile Glu Gly Lys Val His Pro	
75 80 85	
CAG CCG CCG CAG CAG GGG GCA CTC GAC AGG AAG GCG GCG GCG GCG GCG	579
Gln Arg Arg Gln Gln Gly Ala Leu Asp Arg Lys Ala Ala Ala Ala Ala	
90 95 100 105	
GGC GAG GCA GGG GCG TGG GGC GGC GAT CGC GAG CCG CCA GCC GCG GGC	627
Gly Glu Ala Gly Ala Trp Gly Gly Asp Arg Glu Pro Pro Ala Ala Gly	
110 115 120	
CCA CCG GCG CTG GGG CCG CCC GCC GAG GAG CCG CTG CTC GCC GCC AAC	675
Pro Arg Ala Leu Gly Pro Pro Ala Glu Glu Pro Leu Leu Ala Ala Asn	
125 130 135	
GGG ACC GTG CCC TCT TGG CCC ACC GCC CCG GTG CCC AGC GCC GGC GAG	723
Gly Thr Val Pro Ser Trp Pro Thr Ala Pro Val Pro Ser Ala Gly Glu	
140 145 150	
CCC GGG GAG GAG GCG CCC TAT CTG GTG AAG GTG CAC CAG GTG TGG GCG	771
Pro Gly Glu Glu Ala Pro Tyr Leu Val Lys Val His Gln Val Trp Ala	
155 160 165	
GTG AAA GCC GGG GGC TTG AAG AAG GAC TCG CTG CTC ACC GTG GCG CTG	819
Val Lys Ala Gly Gly Leu Lys Lys Asp Ser Leu Leu Thr Val Arg Leu	
170 175 180 185	
GGG ACC TGG GGC CAG CCC GCC TTC CCC TCC TGC GGG AGG CTC AAG GAG	867
Gly Thr Trp Gly His Pro Ala Phe Pro Ser Cys Gly Arg Leu Lys Glu	
190 195 200	
GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC GAC GCC AAC AGC ACC AGC	915
Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Asp Ala Asn Ser Thr Ser	
205 210 215	
GCG GCG CCG GCC GCC TTC GGA GCC TCT TTC CCC CCT CTG GAG ACG GGC	963
Arg Ala Pro Ala Ala Phe Arg Ala Ser Phe Pro Pro Leu Glu Thr Gly	
220 225 230	
CGG AAC CTC AAG AAG GAG GTC AGC CCG GTG CTG TGC AAG CCG TGC GCC	1011
Arg Asn Leu Lys Lys Glu Val Ser Arg Val Leu Cys Lys Arg Cys Ala	
235 240 245	
TTG CCT CCC CAA TTG AAA GAG ATG AAA AGC CAG GAA TCG GCT GCA GGT	1059
Leu Pro Pro Gln Leu Lys Glu Met Lys Ser Gln Glu Ser Ala Ala Gly	
250 255 260 265	
TCC AAA CTA GTC CTT CCG TGT GAA ACC AGT TCT GAA TAC TCC TCT CTC	1107
Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu	
270 175 180	
AGA TTC AAG TGG TTC AAG AAT GGG AAT GAA TTG AAT CGA AAA AAC AAA	1155
Arg Phe Lys Trp Phe Lys Asn Gly Asn Glu Leu Asn Arg Lys Asn Lys	
185 190 195	

CCA CAA AAT ATC AAG ATA CAA AAA AAG CCA GGG AAG TCA GAA CTT CGC Pro Gln Asn Ile Lys Ile Gln Lys Lys Pro Gly Lys Ser Glu Leu Arg 200 205 210	1203
ATT AAC AAA GCA TCA CTG GCT GAT TCT GGA GAG TAT ATG TGC AAA GTG Ile Asn Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val 215 220 225	1251
ATC AGC AAA TTA GGA AAT GAC AGT GCC TCT GCC AAT ATC ACC ATC GTG Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val 230 235 240 245	1299
GAA TCA AAC GCT ACA TCT ACA TCC ACC ACT GGG ACA AGC CAT CTT GTA Glu Ser Asn Ala Thr Ser Thr Ser Thr Gly Thr Ser His Leu Val 250 255 260	1347
AAA TGT GCG GAG AAG GAG AAA ACT TTC TGT GTG AAT GGA GGG GAG TGC Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys 265 270 275	1395
TTC ATG GTG AAA GAC CTT TCA AAC CCC TCG AGA TAC TTG TGC AAG TGC Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys 280 285 290	1443
CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser 295 300 305	1491
TTC TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT GAA Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro Glu 400 405 410	1530
TAGGAGCATG CTCAGTTGGT GCTGCTTTCT TGTTGCTGCA TCTCCCCTCA GATTCCACCT	1590
AGAGCTAGAT GTGTCTTACC AGATCTAATA TTGACTGCCT CTGCCTGTCTG CATGAGAACA	1650
TTAACAAAAG CAATTGTATT ACTTCCTCTG TTCGCGACTA GTTGGCTCTG AGATACTAAT	1710
AGGTGTGTGA GGCTCCGGAT GTTCTGGAA TTGATATTGA ATGATGTGAT ACAAATTGAT	1770
AGTCAATATC AAGCAGTGAA ATATGATAAT AAAGGCATTT CAAAGTCTCA CTTTTATTGA	1830
TAAAATAAAA ATCATTCTAC TGAACAGTCC ATCTTCTTTA TACAATGACC ACATCCTGAA	1890
AAGGGTGTG CTAAGCTGTA ACCGATATGC ACTTGAATG ATGGTAAGTT AATTTTGATT	1950
CAGAATGTGT TATTTGTCAC AAATAACAT AATAAAAGGA AAAAAAAAAA AAA	2003

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: Xaa in position 11 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Ala Ser Leu Ala Asp Glu Tyr Glu Tyr Met Xaa Lys  
1 5 10



- 96 -

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 27:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

Ala Lys Glu Ala Leu Ala Ala Leu Lys  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 28:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Phe Val Leu Gln Ala Lys Lys  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 29:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

Glu Thr Gln Pro Asp Pro Gly Gln Ile Leu Lys Lys Val Pro Met Val  
 1 5 10 15  
 Ile Gly Ala Tyr Thr  
 20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 30:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in positions 1, 3, 17 and 19 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

Xaa Glu Xaa Lys Glu Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Glu  
 1 5 10 15

- 97 -

Xaa Gly Xaa Gly Lys  
20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 31:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu  
1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 32:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 6 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

Lys Leu Glu Phe Leu Xaa Ala Lys  
1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 33:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

Xaa Val His Gln Val Trp Ala Ala Lys  
1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 34:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear







- 100 -

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 41:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

Cys Cys Thr Cys Gly Cys Thr Cys Cys Thr Thr Cys Thr Thr Cys Thr  
 1 5 10 15  
 Thr Gly Cys Cys Cys Thr Thr Cys  
 20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 42:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 8  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

Val His Gln Val Trp Ala Ala Lys  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 43:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 10 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

Tyr Ile Phe Phe Met Glu Pro Glu Ala Xaa Ser Ser Gly  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 44:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

Leu Gly Ala Trp Gly Pro Pro Ala Phe Pro Val Xaa Tyr  
 1 5 10

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

Trp Phe Val Val Ile Glu Gly Lys  
 1 5

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 46:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

Ala Ser Pro Val Ser Val Gly Ser Val Gln Glu Leu Val Gln Arg  
 1 5 10 15

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 47:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

Val Cys Leu Leu Thr Val Ala Ala Leu Pro Pro Thr  
 1 5 10

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 48:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

Lys Val His Gln Val Trp Ala Ala Lys  
 1 5

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 49:

(i) SEQUENCE CHARACTERISTICS:

- 102 -

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Xaa Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 50:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 5 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

Asp Leu Leu Leu Xaa Val  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 51:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

TTYAARGGNG AYGCNCAYAC

20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 52:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

CAIRTAAYTCR TAYTCRTCNG C

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 53:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20

- 103 -

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

TGYTCNGANG CCATYTCNGT

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

TGYTCRCCTNG CCATYTCNGT

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 55:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

CCDATNACCA TNGGNACYTT

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

GCNGCCCANNA CYTGRTGNAC

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 57:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

GCYTCNGGYT CCATRAARA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 58:

## (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

CCYTCDATNA CNACRAACCA

20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 59:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

TCNGCRAART ANCCNGC

17

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 60:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

GCNGCNAGNG CYTCYTTNGC

20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 61:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

GCNGCYAANG CYTCYTTNGC

20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 62:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

TTYTTNGCYT GNAGNACRAA

20

- 105 -

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 63:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

TTYTTNGCYT GYAANACRAA

20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 64:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

TGNACNAGYT CYTGNAC

17

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 65:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

TGNACYAAYT CYTGNAC

17

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 66:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

CATRTAYTCN CCNGARTCNG C

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 67:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

CATRTAYTCN CCRCTRTCNG C

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 68:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

NGARTCNGCY AANGANGCYT T

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 69:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

NGARTCNGCN AGNGANGCYT T

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 70:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

RCCTRTCNGCY AANGANGCYT T

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 71:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

RCCTRTCNGCN AGNGANGCYT T

21

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 72:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear



- 107 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:  
 NGARTCNGCY AARCTNGCYT T 21
- (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 73:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:  
 NGARTCNGCN AGRCTNGCYT T 21
- (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 74:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:  
 RCTRTCNGCY AARCTNGCYT T 21
- (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 75:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:  
 RCTRCTNGCN AGRCTNGCYT T 21
- (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 76:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:  
 ACNACNGARA TGGCTCNGA 20
- (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 77:  
 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20  
 (B) TYPE: nucleic acid

- 108 -

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

ACNACNGARA TGGCAGYNGA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 78:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

CAYCARGTNT GGGCNGCNAA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 79:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

TTYGTNGTNA THGARGGNAA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 80:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

AARGGNGAYG CNCAYACNGA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 81:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

GARGCNYTNG CNGCNYTNAA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 82:

(i) SEQUENCE CHARACTERISTICS:

**SUBSTITUTE SHEET (RULE 26)**

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

GTNGGNTCNG TNCARGARYT 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 83:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

GTNGGNAGYG TNCARGARYT 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 84:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

NACYTTYTTN ARDATYTGNC C 21

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 85:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 417  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: Xaa in positions 14, 23, 90, 100, 126, and 135 is a stop codon.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

TCTAA AAC TAC AGA GAC TGT ATT TTC ATG ATC ATC ATA GTT CTG TGA AAT ATA 53  
 Asn Tyr Arg Asp Cys Ile Phe Met Ile Ile Ile Val Leu Xaa Asn Ile  
 1 5 10 15

CTT AAA CCG CTT TGG TCC TGA TCT TGT AGG AAG TCA GAA CTT CGC ATT 101  
 Leu Lys Pro Leu Trp Ser Xaa Ser Cys Arg Lys Ser Glu Leu Arg Ile  
 20 25 30

AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA TAT ATG TCC AAA GTG ATC 149  
 Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Ser Met Cys Lys Val Ile  
 35 40 45

- 110 -

AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Arg Ile Val Glu 80 55 60	197
TCA AAC GGT AAG AGA TGC CTA CTG CGT GCT ATT TCT CAG TCT CTA AGA Ser Asn Gly Lys Arg Cys Leu Leu Arg Ala Ile Ser Gln Ser Leu Arg 65 70 75 80	245
GGA GTG ATC AAG GTA TGT GGT CAC ACT TGA ATC ACG CAG GTG TGT GAA Gly Val Ile Lys Val Cys Gly His Thr Xaa Ile Thr Gln Val Cys Glu 85 90 95	293
ATC TCA TTG TGA ACA AAT AAA AAT CAT GAA AGG AAA ACT CTA TGT TTG Ile Ser Cys Xaa Thr Asn Lys Asn His Glu Arg Lys Thr Leu Cys Leu 100 105 110	341
AAA TAT CTT ATG GGT CCT CCT GTA AAG CTC TTC ACT CCA TAA GGT GAA Lys Tyr Leu Met Gly Pro Pro Val Lys Leu Phe Thr Pro Xaa Gly Glu 115 120 125	389
ATA GAC CTG AAA TAT ATA TAG ATT ATT T Ile Asp Leu Lys Tyr Ile Xaa Ile Ile 130 135	417

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 86:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at positions 19, 25, and 31 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

CCGAATTCTG CAGGARACNC ARCCNGAYCC NGG 33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 87:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at positions 14, 20, 23, 29, and 35 is Inosine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

AAGGATCCTG CAGNGTRTAN GCNCCDATNA CCGATNGG 37

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 88:

## (i) SEQUENCE CHARACTERISTICS:

- 111 -

(A) LENGTH: 34  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at positions 16, 21, and 24 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:

CCGAATTCTG CAGGCNGAYT CNGGNGARTA YATG

34

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 89:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at positions 16 and 25 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

CCGAATTCTG CAGGCNGAYA GYGGNGARTA YAT

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 90:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at positions 14, 15, 16, 26, and 29 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

AAGGATCCTG CAGNNNCATR TAYTCNCNG ARTC

34

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 91:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

- 112 -

(D) OTHER INFORMATION: N at positions 14, 15, 16,  
and 26 is Inosine. Y can  
be cytidine or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

AAGGATCCTG CAGNNNCATR TAYTCNCCRC TRTC

34

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 92:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at positions 21, 28, and  
31 is Inosine. Y can be  
cytidine or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

CCGAATTCTG CAGCCTCARG TNTGGGCNGC NAA

33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 93:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at position 31 is  
Inosine. Y can be cytidine  
or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

CCGAATTCTG CAGATHTTYT TYATGGARCC NGARG

35

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 94:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at positions 18, 21, 24,  
27, and 33 is Inosine. Y  
can be cytidine or  
thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

CCGAATTCTG CAGGGGNCC NCCNGCNTTY CCGT

35

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 95:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at positions 21 and 24 is  
 Inosine. Y can be cytidine  
 or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

CCGAATTCTG CAGTGGTTYG TNGTNATHGA RGG

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 96:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at positions 17, 20, and  
 26 is Inosine. Y can be  
 cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96

AAGGATCCTG CAGYTTNGCN GCCCANACYT GRTG

34

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 97:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N at position 19 is  
 Inosine. Y can be cytidine  
 or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

AAGGATCCTG CAGGCYTCNG GYTCCATRAA RAA

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 98:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid

- 114 -

- (C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at positions 16, 22, 25, 28, and 31 is Inosine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

AAGGATCCTG CAGACNGGRA ANGCNNGNGG NCC

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 99:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at positions 17, 26, and 29 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

AAGGATCCTG CAGYTTNCCY TCDATNACNA CRAAC

35

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 100:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at position 18 is Inosine. Y can be cytidine or thymidine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

CATRTAYTCR TAYTCTCNGC AAGGATCCTG CAG

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 101:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ix) FEATURE:

- (D) OTHER INFORMATION: N at position 19, 25, and 31 is Inosine. Y can be cytidine or thymidine.



- 115 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

CCGAATTCTG CAGAARGGNG AYGCNCAYAC NGA

33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 102:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at position 3 and 18 is  
 Inosine. Y can be cytidine  
 or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:

GCNGCYAANG CYTCYTTNGC AAGGATCCTG CAG

33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 103:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at position 3, 6, 9, and  
 18 is Inosine. Y can be  
 cytidine or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

GCNGCNAGNG CYTCYTTNGC AAGGATCCTG CAG

33

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 104:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N at position 3, 12, and 15  
 is Inosine. Y can be  
 cytidine or thymidine.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

TCNGCRAART ANCCNGCAAG GATCCTGCAG

30

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 105:

- 116 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:

CATCGATCTG CAGGCTGATT CTGGAGAATA TATGTGCA 38

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 106:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:

AAGGATCCTG CAGCCACATC TCGAGTCGAC ATCGATT 37

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 107:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

CCGAATTCTG CAGTGATCAG CAAACTAGGA AATGACA 37

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 108:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

CATCGATCTG CAGCCTAGTT TGCTGATCAC TTTGCAC 37

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 109:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:

AAGGATCCTG CAGTATATTC TCCAGAATCA GCCAGTG 37

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 110:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

AAGGATCCTG CAGGCACGCA GTAGGCATCT CTTA

34

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 111:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

CCGAATTCTG CAGCAGAACT TCGCATTAGC AAAGC

35

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 112:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

CATCCCGGGA TGAAGAGTCA GGAGTCTGTG GCA

33

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 113:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

ATACCCGGGC TGCAGACAAT GAGATTCAC ACACCTGCG

39

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 114:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

AAGGATCCTG CAGTTTGGAA CCGCCACAG ACTCCT

36

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 115:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 39  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

ATACCCGGGC TGCAGATGAG ATTCACACA CCTGCGTGA

39

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 116:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

His Gln Val Trp Ala Ala Lys Ala Ala Gly Leu Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 117:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

Gly Gly Leu Lys Lys Asp Ser Leu Leu Thr Val Arg Leu Gly Ala Asn  
 1 5 10 15

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 118:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

Leu Gly Ala Trp Gly Pro Pro Ala Phe Pro Val Xaa Tyr  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 119:

- 119 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

Leu Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser  
 1 5 10 15

Cys Gly Arg Leu Lys Glu Asp  
 20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 120:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 10 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

Tyr Ile Phe Phe Met Glu Pro Glu Ala Xaa Ser Ser Gly  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 121:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Glu Ala Asn Ser  
 1 5 10 15

Ser Gly Gly Pro Gly Arg Leu  
 20

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 122:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

Val Ala Gly Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 123:

- 120 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 123:

Glu Tyr Lys Cys Leu Lys Phe Lys Trp Phe Lys Lys Ala Thr Val Met  
 1 5 10 15

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 124:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 124:

Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu Lys Phe Lys Trp Phe Lys  
 1 5 10 15

Asn Gly Ser Glu Leu Ser Arg Lys Asn Lys  
 20 25

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 125:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 125:

Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Xaa Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 126:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 126:

Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met  
 1 5 10 15

Cys Lys Val Ile Ser Lys Leu  
 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 127:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 127:

Ala Ser Leu Ala Asp Glu Tyr Glu Tyr Met Arg Lys  
 1 5 10

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 128:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 128:

Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys  
 1 5 10 15  
 Lys Val Ile Ser Lys Leu  
 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 129:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 744  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 129:

CCTGCAG CAT CAA GTG TGG GCG GCG AAA GCC GGG GGC TTG AAG AAG GAC TCG CTG 55  
 His Gln Val Trp Ala Ala Lys Ala Gly Gly Leu Lys Lys Asp Ser Leu  
 1 5 10 15  
 CTC ACC GTG CGC CTG GGC GCC TGG GGC CAC CCC GCC TTC CCC TCC TGC 103  
 Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser Cys  
 20 25 30  
 GGG CGC CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC GAG 151  
 Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Glu  
 35 40 45  
 GCC AAC AGC AGC GGC GGG CCC GGC CGC CTT CCG AGC CTC CTT CCC CCC 199  
 Ala Asn Ser Ser Gly Gly Pro Gly Arg Leu Pro Ser Leu Leu Pro Pro  
 50 55 60  
 TCT CGA GAC GGG CCG GAA CCT CAA GAA GGA GGT CAG CCG GGT GCT GTG 247  
 Ser Arg Asp Gly Pro Glu Pro Gln Glu Gly Gly Gln Pro Gly Ala Val  
 65 70 75 80  
 CAA CGG TGC GCC TTG CCT CCC CGC TTG AAA GAG ATG AAG AGT CAG GAG 295  
 Gln Arg Cys Ala Leu Pro Pro Arg Leu Lys Glu Met Lys Ser Gln Glu  
 85 90 95

TCT GTG GCA GGT TCC AAA CTA GTG CTT CGG TGC GAG ACC AGT TCT GAA Ser Val Ala Gly Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu 100 105 110	343
TAC TCC TCT CTC AAG TTC AAG TGG TTC AAG AAT GGG AGT GAA TTA AGC Tyr Ser Ser Leu Lys Phe Lys Trp Phe Lys Asn Gly Ser Glu Leu Ser 115 120 125	391
CGA AAG AAC AAA CCA GAA AAC ATC AAG ATA CAG AAA AGG CCG GGG AAG Arg Lys Asn Lys Pro Glu Asn Ile Lys Ile Gln Lys Arg Pro Gly Lys 130 135 140	439
TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA TAT Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Ser Gly Glu Tyr 145 150 155 160	487
ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn 165 170 175	535
ATC ACC ATT GTG GAG TCA AAC GGT AAG AGA TGC CTA CTG CGT GCT ATT Ile Thr Ile Val Glu Ser Asn Gly Lys Arg Cys Leu Leu Arg Ala Ile 180 185 190	583
TCT CAG TCT CTA AGA GGA GTG ATC AAG GTA TGT GGT CAC ACT Ser Gln Ser Leu Arg Gly Val Ile Lys Val Cys Gly His Thr 195 200 205	625
TGAATCACGC AGGTGTGTGA AATCTCATTG TGAACAAATA AAAATCATGA AAGGAAAAAA	685
AAAAAATAA AATCGATGTC GACTCGAGAT GTGGCTGCAG GTCGACTCTA GAGGATCCC	744

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 130:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1193
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 130:

CCTGCAG CAT CAA GTG TGG GCG GCG AAA GCC GGG GGC TTG AAG AAG GAC TCG CTG 55 His Gln Val Trp Ala Ala Lys Ala Gly Gly Leu Lys Lys Asp Ser Leu 1 5 10 15	
CTC ACC GTG CGC CTG GGC GCC TGG GGC CAC CCC GCC TTC CCC TCC TGC 103 Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser Cys 20 25 30	
GGG CGC CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC GAG 151 Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Glu 35 40 45	
GCC AAC AGC AGC GGC GGG CCC GGC CGC CTT CCG AGC CTC CTT CCC CCC 199 Ala Lys Ser Ser Gly Gly Pro Gly Arg Leu Pro Ser Leu Leu Pro Pro 50 55 60	
TCT CGA GAC GGG CCG GAA CCT CAA GAA GGA GGT CAG CCG GGT GCT GTG 247 Ser Arg Asp Gly Pro Glu Pro Gln Glu Gly Gly Gln Pro Gly Ala Val 65 70 75 80	



CAA CGG TGC GCC TTG CCT CCC CGC TTG AAA GAG ATG AAG AGT CAG GAG Gln Arg Cys Ala Leu Pro Pro Arg Leu Lys Glu Met Lys Ser Gln Glu 85 90 95	295
TCT GTG GCA GGT TCC AAA CTA GTG CTT CGG TGC GAG ACC AGT TCT GAA Ser Val Ala Gly Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu 100 105 110	343
TAC TCC TCT CTC AAG TTC AAG TGG TTC AAG AAT GGG AGT GAA TTA AGC Tyr Ser Ser Leu Lys Phe Lys Trp Phe Lys Asn Gly Ser Glu Leu Ser 115 120 125	391
CGA AAG AAC AAA CCA GAA AAC ATC AAG ATA CAG AAA AGG CCG GGG AAG Arg Lys Asn Lys Gly Gly Asn Ile Lys Ile Gln Lys Arg Pro Gly Lys 130 135 140	439
TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA TAT Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr 145 150 155 160	487
ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn 165 170 175	535
ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG ACA Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr Ser Thr Ala Gly Thr 180 185 190	583
AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT Ser His Leu Val Lys Ser Ala Glu Lys Glu Lys Thr Phe Cys Val Asn 195 200 205	631
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr 210 215 220	679
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn 225 230 235 240	727
GTG CCC ATG AAA GTC CAA ACC CAA GAA AGT GCC CAA ATG AGT TTA CTG Val Pro Met Lys Val Gln Thr Gln Glu Ser Ala Gln Met Ser Leu Leu 245 250 255	775
GTG ATC GCT GCC AAA ACT ACG TAATGGCCAG CTTCTACAGT ACGTCCACTC Val Ile Ala Ala Lys Thr Thr 260	826
CCTTTCTGTC TCTGCCTGAA TAGCGCATCT CAGTCGGTGC CGCTTTCTTG TTGCCGCATC	886
TCCCCTCAGA TTCCTCCTAG AGCTAGATGC GTTTTACCAG GTCTAACATT GACTGCCTCT	946
GCCTGTGCGCA TGAGAACATT AACACAAGCG ATTGTATGAC TTCCTCTGTC CGTGACTAGT	1006
GGGCTCTGAG CTA CTCTCGTAG GTGCGTAAGG CTCCAGTGTT TCTGAAATTG ATGTTGAATT	1066
ACTGTGATAC GACATGATAG TCCCTCTCAC CCAGTGCAAT GACAATAAAG GCCTTGAAAA	1126
GTCAAAAAAA AAAAAAATAA AAAAAATCGA TGTCGACTCG AGATGTGGCT GCAGGTGCGAC	1186
TCTAGAG	1193

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 131:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1108  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 131:

CCTGCAG	CAT	CAA	GTG	TGG	GCG	GCG	AAA	GCC	GGG	GGC	TTG	AAG	AAG	GAC	TCG	CTG	55
	His	Gln	Val	Trp	Ala	Ala	Lys	Ala	Gly	Gly	Leu	Lys	Lys	Asp	Ser	Leu	
	1				5					10					15		
CTC	ACC	GTG	CGC	CTG	GGC	GCC	TGG	GGC	CAC	CCC	GCC	TTC	CCC	TCC	TGC		103
Leu	Thr	Val	Arg	Leu	Gly	Ala	Trp	Gly	His	Pro	Ala	Phe	Pro	Ser	Cys		
		20					25					30					
GGG	CGC	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	ATG	GAG	CCC	GAG		151
Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	Met	Glu	Pro	Glu		
		35					40					45					
GCC	AAC	AGC	AGC	GGC	GGG	CCC	GGC	CGC	CTT	CCG	AGC	CTC	CTT	CCC	CCC		199
Ala	Asn	Ser	Ser	Gly	Gly	Pro	Gly	Arg	Leu	Pro	Ser	Leu	Leu	Pro	Pro		
	50					55					60						
TCT	CGA	GAC	GGG	CCG	GAA	CCT	CAA	GAA	GGA	GGT	CAG	CCG	GGT	GCT	GTG		247
Ser	Arg	Asp	Gly	Pro	Glu	Pro	Gln	Glu	Gly	Gly	Gln	Pro	Gly	Ala	Val		
					70					75					80		
CAA	CGG	TGC	GCC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG		295
Gln	Arg	Cys	Ala	Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu		
				85					90					95			
TCT	GTG	GCA	GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA		343
Ser	Val	Ala	Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu		
			100				105						110				
TAC	TCC	TCT	CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC		391
Tyr	Ser	Ser	Leu	Lys	Phe	Lys	Trp	Phe	Lys	Asn	Gly	Ser	Glu	Leu	Ser		
		115					120					125					
CGA	AAG	AAC	AAA	CCA	GAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GGG	AAG		439
Arg	Lys	Asn	Lys	Pro	Glu	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Pro	Lys		
	130					135					140						
TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	GAA	TAT		487
Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly	Glu	Tyr		
	145				150					155					160		
ATG	TGC	AAA	GTG	ATC	AGC	AAA	CTA	GGA	AAT	GAC	AGT	GCC	TCT	GCC	AAC		535
Met	Cys	Lys	Val	Ile	Ser	Lys	Leu	Gly	Asn	Asp	Ser	Ala	Ser	Ala	Asn		
				165					170					175			
ATC	ACC	ATT	GTG	GAG	TCA	AAC	GCC	ACA	TCC	ACA	TCT	ACA	GCT	GGG	ACA		583
Ile	Arg	Ile	Val	Glu	Ser	Asn	Ala	Thr	Ser	Thr	Ser	Thr	Ala	Gly	Thr		
			180				185						190				
AGC	CAT	CTT	GTC	AAG	TGT	GCA	GAG	AAG	GAG	AAA	ACT	TTC	TGT	GTG	AAT		631
Ser	His	Leu	Val	Lys	Cys	Ala	Glu	Lys	Glu	Lys	Thr	Phe	Cys	Val	Asn		
		195				200						205					
GGA	GCC	GAG	TGC	TTC	ATG	GTG	AAA	GAC	CTT	TCA	AAT	CCC	TCA	AGA	TAC		679
Gly	Gly	Glu	Cys	Phe	Met	Val	Lys	Asp	Leu	Ser	Asn	Pro	Ser	Arg	Tyr		
	210					215					220						
TTG	TGC	AAG	TGC	CCA	AAT	GAG	TTT	ACT	GGT	GAT	CGC	TGC	CAA	AAC	TAC		727
Leu	Cys	Lys	Cys	Pro	Asn	Glu	Phe	Thr	Gly	Asp	Arg	Cys	Gln	Asn	Tyr		
	225				230					235					240		

GTA ATG GCC AGC TTC TAG AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT 775  
 Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro  
 245 250 255

GAA TAGCGCATCT CAGTCGGTGC CGCTTTCTTG TTGCCGCATC TCCCCTCAGA TTCCGCCTAG 838  
 Glu

AGCTAGATGC GTTTTACCAG GTCTAACATT GACTGCCTCT GCCTGTCCGA TGAGAACATT 898

AACACAAGCG ATTGTATGAC TTCCTCTGTC CGTGACTAGT GGGCTCTGAG CTA CTCTCGTAG 958

GTCCGTAAGG CTCCAGTGTT TCTGAAATTG ATCTTGAATT ACTGTGATAC GACATGATAG 1018

TCCCTCTCAC CCAGTGCAAT GACAATAAAG GCCTTGAAAA GTCAAAAAA AAAAAAAAAA 1078

AAAAATCGAT GTCGACTCGA GATGTGGCTG 1108

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 132:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 559
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: N in position 214 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 132:

AGTTTCCCCC CCCAACTTGT CGGA ACTCTG GGCTCGCGCG CAGGGCAGGA GCGGAGGGGC 60

GGCGGCTGCC CAGGCGATGC GAGCGCGGGC CGGACGGTAA TCGCCTCTCE CTCCTCGGGC 120

TGCGAGCGCG CCGGACCGAG GCAGCGACAG GAGCGGACCG CGGCGGGAAC CGAGGACTCC 180

CCAGCGGGCG GCCAGCAGGA GCCACCCCGC GAGNCGTGCG ACCGGGACGG AGCGCCCCGC 240

AGTCCCAGGT GGCCCGGACC GCACGTTGG TCCCGCGCT CCCC GCCGGC GACAGGAGAC 300

GCTCCCCCCC ACGCCGCGCG CGCCTCGGCC CGGTGCTGG CCCGCCTCCA CTCGGGGAC 360

AAACTTTTCC CGAAGCCGAT CCCAGCCCTC GGACCCAAAC TTGTGCGCGG TCGCCTTCGC 420

CGGGAGCCGT CCGCGCAGAG CGTGCACTTC TCGGGCGAG ATG TCG GAG CGC AGA 474  
 Met Ser Glu Arg Arg  
 1 5

GAA GGC AAA GGC AAG GGG AAG GGC GGC AAG AAG GAC CGA GGC TCC GGG 522  
 Glu Gly Lys Gly Lys Gly Lys Gly Gly Lys Lys Asp Arg Gly Ser Gly  
 10 15 20

AAG AAG CCC GTG CCC GCG GCT GGC GGC CCG AGC CCA G 559  
 Lys Lys Pro Val Pro Ala Ala Gly Gly Pro Ser Pro Ala  
 25 30

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 133:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 252
- (B) TYPE: nucleic acid

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ix) FEATURE:

(D) OTHER INFORMATION: N in position 8 could be either A or G.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 133:

CC CAT CAN GTG TGG GCG GCG AAA GCC GGG GGC TTG AAG AAG GAC TCG	47
His Gln Val Trp Ala Ala Lys Ala Gly Gly Leu Lys Lys Asp Ser	
1 5 10 15	
CTG CTC ACC GTG CGC CTG GGC GCC TGG GGC CAC CCC GCC TTC CCC TCC	95
Leu Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser	
20 25 30	
TGC GGG CGC CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC	143
Cys Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro	
35 40 45	
GAG GCC AAC AGC AGC GGC GGC CCC GGC CGC CTT CCG AGC CTC CTT CCC	191
Glu Ala Asn Ser Ser Gly Gly Pro Gly Arg Leu Pro Ser Leu Leu Pro	
50 55 60	
CCC TCT CGA GAC GGG CCG GAA CCT CAA GAA GGA GGT CAG CCG GGT GCT	239
Pro Ser Arg Asp Gly Pro Glu Pro Gln Glu Gly Gly Gln Pro Gly Ala	
65 70 75	
GTG CAA CGG TGC G	252
Val Gln Arg Cys	
80	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 134:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 178  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 134:

CCT TGC CTC CCC GCT TGA AAG AGA TGA AGA GTC AGG AGT CTG TGG CAG	48
Leu Pro Pro Arg Leu Lys Glu His Lys Ser Gln Glu Ser Val Ala Gly	
1 5 10 15	
GTT CCA AAC TAG TGC TTC GGT GCG AGA CCA GTT CTG AAT ACT CCT CTC	96
Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu	
20 25 30	
TCA AGT TCA AGT GGT TCA AGA ATG GGA GTC AAT TAA GCC GAA AGA ACA	144
Lys Phe Lys Trp Phe Lys Asn Gly Ser Glu Leu Ser Arg Lys Asn Lys	
35 40 45	
AAC CAC AAA ACA TCA AGA TAC AGA AAA GGC CGG G	178
Pro Gly Asn Ile Lys Ile Gln Lys Arg Pro Gly	
50 55	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 135:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 122  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 135:

G AAG TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA 46  
 Lys Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly  
 1 5 10 15

GAA TAT ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT 94  
 Glu Tyr Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser  
 20 25 30

GCC AAC ATC ACC ATT GTG GAG TCA AAC G 122  
 Ala Asn Ile Thr Ile Val Glu Ser Asn Ala  
 35

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 136:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 417  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 136:

TCTAAACTA CAGAGACTGT ATTTTCATGA TCATCATAGT TCTGTGAAAT ATACTTAAAC 60

CGCTTTGGTC CTGATCTTGT AGG AAG TCA GAA CTT CGC ATT AGC AAA GCG 110  
 Lys Ser Glu Leu Arg Ile Ser Lys Ala  
 1 5

TCA CTG GCT GAT TCT GGA GAA TAT ATG TGC AAA GTG ATC AGC AAA CTA 158  
 Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys Leu  
 10 15 20 25

GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG TCA AAC GGT 206  
 Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn Gly  
 30 35 40

AAG AGA TGC CTA CTG CGT GCT ATT TCT CAG TCT CTA AGA GGA GTG ATC 254  
 Lys Arg Cys Leu Leu Arg Ala Ile Ser Gln Ser Leu Arg Gly Val Ile  
 45 50 55

AAG GTA TGT GGT CAC ACT TGAATCACGC AGGTGTGTGA AATCTCATTG 302  
 Lys Val Cys Gly His Thr  
 60

TGAACAAATA AAAATCATGA AAGGAAACT CTATGTTTGA AATATCTTAT GGGTCCTCCT 362

GTAAAGCTCT TCACTCCATA AGGTGAAATA GACCTGAAAT ATATATAGAT TATTT 417

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 137:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 137:

AG ATC ACC ACT GGC ATG CCA GCC TCA ACT GAG ACA GCG TAT GTG TCT	47
Glu Ile Thr Thr Gly Met Pro Ala Ser Thr Glu Thr Ala Tyr Val Ser	
1 5 10 15	
TCA GAG TCT CCC ATT AGA ATA TCA GTA TCA ACA GAA GGA ACA AAT ACT	95
Ser Glu Ser Pro Ile Arg Ile Ser Val Ser Thr Glu Gly Thr Asn Thr	
20 25 30	
TCT TCA T	102
Ser Ser Ser	
35	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 138:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 138:

AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT GTG CCC	48
Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn Val Pro	
1 5 10 15	
ATG AAA GTC CAA ACC CAA GAA	69
Met Lys Val Gln Thr Gln Glu	
20	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 139:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 60
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 139:

AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC GTA ATG	48
Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met	
1 5 10 15	
GCC AGC TTC TAC	60
Ala Ser Phe Tyr	
20	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 140:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 140:

AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT GAA TAG 36  
 Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro Glu  
 1 5 10

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 141:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 141:

AAG CAT CTT GCG ATT GAA TTT ATG GAG 27  
 Lys His Leu Gly Ile Glu Phe Met Glu  
 1 5

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 142:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 569
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 142:

AAA GCG GAG GAG CTC TAC CAG AAG AGA GTG CTC ACC ATT ACC GGC ATT 48  
 Lys Ala Glu Glu Leu Tyr Gln Lys Arg Val Leu Thr Ile Thr Gly Ile  
 1 5 10 15

TGC ATC GCG CTC CTC GTG GTT GGC ATC ATG TGT GTG GTG GTC TAC TGC 96  
 Cys Ile Ala Leu Leu Val Val Gly Ile Met Cys Val Val Val Tyr Cys  
 20 25 30

AAA ACC AAG AAA CAA CCG AAA AAG CTT CAT GAC CGG CTT CGG CAG ACC 144  
 Lys Thr Lys Lys Gln Arg Lys Lys Leu His Asp Arg Leu Arg Gln Ser  
 35 40 45

CTT CGG TCT GAA AGA AAC ACC ATG ATG AAC GTA GCC AAC GGG CCC CAC 192  
 Leu Arg Ser Glu Arg Asn Thr Met Met Asn Val Ala Asn Gly Pro His  
 50 55 60

CAG CCC AAT CCG CCC CCC GAG AAC GTG CAG CTG GTG AAT CAA TAC GTA 240  
 His Pro Asn Pro Pro Pro Glu Asn Val Gln Leu Val Asn Gln Tyr Val  
 65 70 75 80

TCT AAA AAT GTC ATC TCT AGC GAG CAT ATT GTT GAG AGA GAG GCG GAG 288  
 Ser Lys Asn Val Ile Ser Ser Glu His Ile Val Glu Arg Glu Ala Glu  
 85 90 95

AGC TCT TTT TCC ACC AGT CAC TAC ACT TCG ACA GCT CAT CAT TCC ACT 336  
 Ser Ser Phe Ser Thr Ser His Tyr Thr Ser Thr Ala His His Ser Thr  
 100 105 110

ACT GTC ACT CAG ACT CCC AGT CAC AGC TGG AGC AAT GGA CAC ACT GAA 384  
 Thr Val Thr Gln Thr Pro Ser His Ser Trp Ser Asn Gly His Thr Glu  
 115 120 125

AGC ATC ATT TCG GAA AGC CAC TCT GTC ATC GTG ATG TCA TCC GTA GAA 432  
 Ser Ile Ile Ser Glu Ser His Ser Val Ile Val Met Ser Ser Val Glu  
 130 135 140

AAC AGT AGG CAC AGC AGC CCG ACT GGG GGC CCG AGA GGA CGT CTC AAT	480
Asn Ser Arg His Ser Ser Pro Thr Gly Gly Pro Arg Gly Arg Leu Asn	
145 150 155 160	
GGC TTG GGA GGC CCT CGT GAA TGT AAC AGC TTC CTC AGG CAT GCC AGA	528
Gly Leu Gly Gly Pro Arg Glu Cys Asn Ser Phe Leu Arg His Ala Arg	
165 170 175	
GAA ACC CCT GAC TCC TAC CGA GAC TCT CCT CAT AGT G AAAG	569
Glu Thr Pro Asp Ser Tyr Arg Asp Ser Pro His Ser	
180 185	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 143:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 735
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 143:

G TAT GTA TCA GCA ATG ACC ACC CCG GCT CGT ATG TCA CCT GTA GAT	46
Tyr Val Ser Ala Met Thr Thr Pro Ala Arg Met Ser Pro Val Asp	
1 5 10 15	
TTC CAC ACG CCA AGC TCC CCC AAG TCA CCC CCT TCG GAA ATG TCC CCG	94
Phe His Thr Pro Ser Ser Pro Lys Ser Pro Pro Ser Glu Met Ser Pro	
20 25 30	
CCC GTG TCC AGC ACG ACG GTC TCC ATG CCC TCC ATG GCG GTC AGT CCC	142
Pro Val Ser Ser Thr Thr Val Ser Met Pro Ser Met Ala Val Ser Pro	
35 40 45	
TTC GTG GAA GAG GAG AGA CCC CTG CTC CTT GTG ACG CCA CCA CGG CTG	190
Phe Val Glu Glu Glu Arg Pro Leu Leu Leu Val Thr Pro Pro Arg Leu	
50 55 60	
CGG GAG AAG TAT GAC CAC CAC GCC CAG CAA TTC AAC TCG TTC CAC TGC	238
Arg Glu Lys Tyr Asp His His Ala Gln Gln Phe Asn Ser Phe His Cys	
65 70 75	
AAC CCC GCG CAT GAG AGC AAC AGC CTG CCC CCC AGC CCC TTG AGG ATA	286
Asn Pro Ala His Glu Ser Asn Ser Leu Pro Pro Ser Pro Leu Arg Ile	
80 85 90 95	
GTG GAG GAT GAG GAA TAT GAA ACG ACC CAG GAG TAC GAA CCA GCT CAA	334
Val Glu Asp Glu Glu Tyr Glu Thr Thr Gln Glu Tyr Glu Pro Ala Gln	
100 105 110	
GAG CCG GTT AAG AAA CTC ACC AAC AGC AGC CCG CCG GCC AAA AGA ACC	382
Glu Pro Val Lys Lys Leu Thr Asn Ser Ser Arg Arg Ala Lys Arg Thr	
115 120 125	
AAG CCC AAT GGT CAC ATT GCC CAC AGG TTG GAA ATG GAC AAC AAC ACA	430
Lys Pro Asn Gly His Ile Ala His Arg Leu Glu Met Asp Asn Asn Thr	
130 135 140	
GGC GCT GAC AGC AGT AAC TCA GAG AGC GAA ACA GAG GAT GAA AGA GTA	478
Gly Ala Asp Ser Ser Asn Ser Glu Ser Glu Thr Glu Asp Glu Arg Val	
145 150 155	
GGA GAA GAT ACG CCT TTC CTG GCC ATA CAG AAC CCC CTG GCA GCC AGT	526
Gly Glu Asp Thr Pro Phe Leu Ala Ile Gln Asn Pro Leu Ala Ala Ser	
160 165 170 175	



CTC GAG GCG GCC CCT GCC TTC CGC CTG GTC GAC AGC AGG ACT AAC CCA 574  
 Leu Glu Ala Ala Pro Ala Phe Arg Leu Val Asp Ser Arg Thr Asn Pro  
 180 185 190

ACA GGC GGC TTC TCT CCG CAG GAA GAA TTG CAG GCC AGG CTC TCC GGT 622  
 Thr Gly Gly Phe Ser Pro Gln Glu Glu Leu Gln Ala Arg Leu Ser Gly  
 195 200 205

GTA ATC GCT AAC CAA GAC CCT ATC GCT GTC TAAACCGAA ATACACCCAT 672  
 Val Ile Ala Asn Gln Asp Pro Ile Ala Val  
 210 215

AGATTCACCT GTAAACTTT ATTTTATATA ATAAAGTATT CCACCTTAAA TTAACAAA 730

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 144:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1654
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 144:

AGTTTCCCCC CCCAACTTGT CGGAACTCTG GGCTCGCGCG CAGGGCAGGA GCGGAGCGGC 60

GGCGGCTGCC CAGGCGATGC GAGCGCGGGC CGGACGGTAA TCGCCTCTCC CTCCTCGGGC 120

TGCGAGCGCG COGACCGAG GCAGCGACAG GAGCGGACCG CGGCGGGAAC CGAGGACTCC 180

CCAGCGGGGC GCCAGCAGGA GCCACCCCGC GAGCGTGCGA CCGGGACGGA GCGCCCGCCA 240

GTCCCAGGTG GCCCGGACCG CACGTTGGT CCCCGCGCTC CCCGCCGGCG ACAGGAGACG 300

CTCCCCCCCA CGCGCGCGCC GCTCGGCCC GGTCGCTGGC CCGCCTCCAC TCCGGGGACA 360

AACTTTTCCC GAAGCCGATC CCAGCCCTCG GACCCAACT TGTCGCGCGT CGCCTTCGCC 420

GGGAGCCGTC CGCGCAGAGC GTGCACTTCT CGGGCGAG ATG TCG GAG CGC AGA 473  
 Met Ser Glu Arg Arg  
 1 5

GAA GGC AAA GGC AAG GGG AAG GGC GGC AAG AAG GAC CGA GGC TCC GGG 521  
 Glu Gly Lys Gly Lys Gly Lys Gly Gly Lys Lys Asp Arg Gly Ser Gly  
 10 15 20

AAG AAG CCC GTG CCC GCG GCT GGC GGC CCG AGC CCA GCC TTG CCT CCC 569  
 Lys Lys Pro Val Pro Ala Ala Gly Gly Pro Ser Pro Ala Leu Pro Pro  
 25 30 35

CGC TTG AAA GAG ATG AAG ATG CAG GAG TCT GTG GCA GGT TCC AAA CTA 617  
 Arg Leu Lys Glu Met Lys Ser Gln Glu Ser Val Ala Gly Ser Lys Leu  
 40 45 50

GTG CTT CGG TGC GAG ACC AGT TCT GAA TAC TCC TCT CTC AAG TTC AAG 665  
 Val Leu Arg Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu Lys Phe Lys  
 55 60 65

TGG TTC AAG AAT GGG AGT GAA TTA AGC CGA AAG AAC AAA CCA CAA AAC 713  
 Trp Phe Lys Asn Gly Ser Glu Leu Ser Arg Lys Asn Lys Pro Gln Asn  
 70 75 80 85

ATC AAG ATA CAG AAA AGG CCG GGG AAG TCA GAA CTT CGC ATT AGC AAA 761  
 Ile Lys Ile Gln Lys Arg Pro Gly Lys Ser Glu Leu Arg Ile Ser Lys  
 90 95 100

GCG TCA CTG GCT GAT TCT GGA GAA TAT ATG TGC AAA GTG ATC AGC AAA Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys 105 110 115	809
CTA GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG TCA AAC Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn 120 125 130	857
GAG ATC ACC ACT GGC ATG CCA GCC TCA ACT GAG ACA GCG TAT GTG TCT Glu Ile Thr Thr Gly Met Pro Ala Ser Thr Glu Thr Ala Tyr Val Ser 135 140 145	905
TCA GAG TCT CCC ATT AGA ATA TCA GTA TCA ACA GAA GGA ACA AAT ACT Ser Glu Ser Pro Ile Arg Ile Ser Val Ser Thr Glu Gly Thr Asn Thr 150 155 160 165	953
TCT TCA TCC ACA TCC ACA TCT ACA GCT GGG ACA AGC CAT CTT GTC AAG Ser Ser Ser Thr Ser Thr Ser Thr Ala Gly Thr Ser His Leu Val Lys 170 175 180	1001
TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT GGA GGC GAG TGC TTC Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys Phe 185 190 195	1049
ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC TTG TGC AAG TGC CCA Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys Pro 200 205 210	1097
AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe 215 220 225	1145
TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT GAA TAGGCGCATG Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro Glu 230 235 240	1191
CTCAGTCGGT GCCGCTTCT TGTGCGCA TCTCCCCTCA GATTCAACCT AGAGCTAGAT	1251
GCGTTTTACC AGGTCTAACA TTGACTGCCT CTGCCTGTCTG CATGAGAACA TTAACACAAG	1311
CGATTGTATG ACTTCTCTG TCCGTGACTA GTGGGCTCTG AGCTACTCGT AGGTGCGTAA	1371
GGCTCCAGTG TTTCTGAAAT TGATCTTGAA TTACTGTGAT ACGACATGAT AGTCCCTCTC	1431
ACCCAGTGCA ATGACAATAA AGGCCTTGAA AAGTCTCACT TTTATTGAGA AAATAAAAAT	1491
CGTTCCACGG GACAGTCCCT CTTCTTTATA AAATGACCCT ATCCTTGAAA AGGAGGTGTG	1551
TTAAGTTGTA ACCAGTACAC ACTTGAAATG ATGGTAAGTT CGCTTCGGTT CAGAATGTGT	1611
TCTTTCTGAC AAATAAACAG AATAAAAAA AAAAAAAAAA A	1652

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 145:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1140
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 145:

CAT CAN GTG TGG GCG GCG AAA GCC GGG GGC TTG AAG AAG GAC TCG CTG His Gln Val Trp Ala Ala Lys Ala Gly Gly Leu Lys Lys Asp Ser Leu 1 5 10 15	48
CTC ACC GTG CGC CTG GGC GCC TGG GGC CAC CCC GCC TTC CCC TCC TGC Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser Cys 20 25 30	96
GGG CGC CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC GAG Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Glu 35 40 45	144
GCC AAC AGC AGC GGC GGG CCC GGC CGC CTT CCG AGC CTC CTT CCC CCC Ala Asn Ser Ser Gly Gly Pro Gly Arg Leu Pro Ser Leu Leu Pro Pro 50 55 60	192
TCT CGA GAC GGG CCG GAA CCT CAA GAA GGA GGT CAG CCG GGT GCT GTG Ser Arg Asp Gly Pro Glu Pro Gln Glu Gly Gln Pro Gly Ala Val 65 70 75 80	240
CAA CGG TGC GCC TTG CCT CCC CGC TTG AAA GAG ATG AAG AGT CAG GAG Gln Arg Cys Ala Leu Pro Pro Arg Leu Lys Glu Met Lys Ser Gln Glu 85 90 95	288
TCT GTG GCA GGT TCC AAA CTA GTG CTT CGG TGC GAG ACC AGT TCT GAA Ser Val Ala Gly Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu 100 105 110	336
TAC TCC TCT CTC AAG TTC AAG TGG TTC AAG AAT GGG AGT GAA TTA AGC Tyr Ser Ser Leu Lys Phe Lys Trp Phe Lys Asn Gly Ser Glu Leu Ser 115 120 125	384
CGA AAG AAC AAA CCA GAA AAC ATC AAG ATA CAG AAA AGG CCG GGG AAG Arg Lys Asn Lys Pro Glu Asn Ile Lys Ile Gln Lys Arg Pro Gly Lys 130 135 140	432
TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA TAT Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr 145 150 155 160	480
ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn 165 170 175	528
ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG ACA Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr Ser Thr Ala Gly Thr 180 185 190	576
AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn 195 200 205	624
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr 210 215 220	672
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn 225 230 235 240	720
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr 245 250 255	768
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC TAC AGT ACG TCC Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser 260 265 270	816

ACT CCC TTT CTG TCT CTG CCT GAA TAGCGCATCT CAGTCGGTGC CGCTTCTTG 870  
 Thr Pro Phe Leu Ser Leu Pro Glu  
 275 280

TGCCCGCATC TCCCCTCAGA TTCNCCTAG AGCTAGATGC GTTTTACCAG GTCTAACATT 930

GACTGCCTCT GCCTGTGCA TGAGAACATT AACACAAGCG ATTGTATGAC TTCCTCTGTC 990

CGTGACTAGT GGGCTCTGAG CTACTCGTAG GTGCGTAAGG CTCCAGTGTT TCTGAAATTG 1050

ATCTGAATT ACTGTGATAC GACATGATAG TCCCTCTCAC CCAGTGCAAT GACAATAAAG 1110

GCCTTGAAA GTCAAAAAA AAAAAAAAAA 1140

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 146:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1764
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 146:

G AAG TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA 49  
 Lys Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu  
 1 5 10 15

TAT ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC 97  
 Tyr Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala  
 20 25 30

AAC ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG 145  
 Asn Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr Thr Ala Gly  
 35 40 45

ACA AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG 193  
 Thr Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val  
 50 55 60

AAT GGA GGC GAC TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA 241  
 Asn Gly Gly Asp Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg  
 65 70 75 80

TAC TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG 289  
 Tyr Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu  
 85 90 95

AAT GTG CCC ATG AAA GTC CAA ACC CAA GAA AAA GCG GAG GAG CTC TAC 337  
 Asn Val Pro Met Lys Val Gln Thr Thr Gln Glu Lys Ala Glu Glu Leu Tyr  
 100 105 110

CAG AAG AGA GTG CTC ACC ATT ACC GGC ATT TGC ATC GCG CTG CTC GTG 385  
 Gln Lys Arg Val Leu Thr Ile Thr Gly Ile Cys Ile Ala Leu Leu Val  
 115 120 125

GTT GGC ATC ATG TGT GTG GTG GTC TAC TGC AAA ACC AAG AAA CAA CGG 433  
 Val Gly Ile Met Cys Val Val Val Tyr Cys Lys Thr Lys Lys Gln Arg  
 130 135 140

AAA AAG CTT CAT GAC CGG CTT CGG CAG AGC CTT CGG TCT GAA AGA AAC 481  
 Lys Lys Leu His Asp Arg Leu Arg Gln Ser Leu Arg Ser Glu Arg Asn  
 145 150 155 160

ACC ATG ATG AAC GTA GCC AAC GGG CCC CAC CAC CCC AAT CCG CCC CCC	529
Thr Met Met Asn Val Ala Asn Gly Pro His His Pro Asn Pro Pro Pro	
165 170 175	
GAG AAC GTG CAG CTG GTG AAT CAA TAC GTA TCT AAA AAT GTC ATC TCT	577
Glu Asn Val Gln Leu Val Asn Gln Tyr Val Ser Lys Asn Val Ile Ser	
180 185 190	
AGC GAG CAT ATT GTT GAG AGA GAG GCG GAG AGC TCT TTT TCC ACC AGT	625
Ser Glu His Ile Val Glu Arg Glu Ala Glu Ser Ser Phe Ser Thr Ser	
195 200 205	
CAC TAC ACT TCG ACA GCT CAT CAT TCC ACT ACT GTC ACT CAG ACT CCC	673
His Tyr Thr Ser Thr Ala His His Ser Thr Thr Val Thr Gln Thr Pro	
210 215 220	
AGT CAC AGC TGG AGC AAT GGA CAC ACT GAA AGC ATC ATT TCG GAA AGC	721
Ser His Ser Trp Ser Asn Gly His Thr Glu Ser Ile Ile Ser Glu Ser	
225 230 235 240	
CAC TCT GTC ATC GTG ATG TCA TCC GTA GAA AAC AGT AGG CAC AGC AGC	769
His Ser Val Ile Val Met Ser Ser Val Glu Asn Ser Arg His Ser Ser	
245 250 255	
CCG ACT GGG GGC CCG AGA GGA CGT CTC AAT GGC TTG GGA GGC CCT CGT	817
Pro Thr Gly Gly Pro Arg Gly Arg Leu Asn Gly Leu Gly Gly Pro Arg	
260 265 270	
GAA TGT AAC AGC TTC CTC AGG CAT GCC AGA GAA ACC CCT GAC TCC TAC	865
Glu Cys Asn Ser Phe Leu Arg His Ala Arg Glu Thr Pro Asp Ser Tyr	
275 280 285	
CGA GAC TCT CCT CAT AGT GAA AGA CAT AAC CTT ATA GCT GAG CTA AGG	913
Arg Asp Ser Pro His Ser Glu Arg His Asn Leu Ile Ala Glu Leu Arg	
290 295 300	
AGA AAC AAG GCC CAC AGA TCC AAA TGC ATG CAG ATC CAG CTT TCC GCA	961
Arg Asn Lys Ala His Arg Ser Lys Cys Met Gln Ile Gln Leu Ser Ala	
305 310 315 320	
ACT CAT CTT AGA GCT TCT TCC ATT CCC CAT TGG GCT TCA TTC TCT AAG	1009
Thr His Leu Arg Ala Ser Ser Ile Pro His Trp Ala Ser Phe Ser Lys	
325 330 335	
ACC CCT TGG CCT TTA GGA AGG TAT GTA TCA GCA ATG ACC ACC CCG GCT	1057
Thr Pro Trp Pro Leu Gly Arg Tyr Val Ser Ala Met Thr Thr Pro Ala	
340 345 350	
CGT ATG TCA CCT GTA GAT TTC CAC ACG CCA AGC TCC CCC AAG TCA CCC	1105
Arg Met Ser Pro Val Asp Phe His Thr Pro Ser Ser Pro Lys Ser Pro	
355 360 365	
CCT TCG GAA ATG TCC CCG CCC GTG TCC AGC ACG ACG GTC TCC ATG CCC	1153
Pro Ser Glu Met Ser Pro Pro Val Ser Ser Thr Thr Val Ser Met Pro	
370 375 380	
TCC ATG GCG GTC AGT CCC TTC GTG GAA GAG GAG AGA CCC CTG CTC CTT	1201
Ser Met Ala Val Ser Pro Phe Val Glu Glu Glu Arg Pro Leu Leu Leu	
385 390 395 400	
GTG ACG CCA CCA CCG CTG CCG GAG AAG TAT GAC CAC CAC GCC CAG CAA	1249
Val Thr Pro Pro Arg Leu Arg Glu Lys Tyr Asp His His Ala Gln Gln	
405 410 415	
TTC AAC TCG TTC CAC TGC AAC CCC GCG CAT GAG AGC AAC AGC CTG CCC	1297
Phe Asn Ser Phe His Cys Asn Pro Ala His Glu Ser Asn Ser Leu Pro	
420 425 430	

CCC AGC CCC TTG AGG ATA GTG GAG GAT GAG GAA TAT GAA ACG ACC CAG Pro Ser Pro Leu Arg Ile Val Glu Asp Glu Glu Tyr Glu Thr Thr Gln 435 440 445	1345
GAG TAC GAA CCA GCT CAA GAG CCG GTT AAG AAA CTC ACC AAC AGC AGC Glu Tyr Glu Pro Ala Gln Glu Pro Val Lys Lys Leu Thr Asn Ser Ser 450 455 460	1393
CGG CGG GCC AAA AGA ACC AAG CCC AAT GGT CAC ATT GCC CAC AGG TTG Arg Arg Ala Lys Arg Thr Lys Pro Asn Gly His Ile Ala His Arg Leu 465 470 475 480	1441
GAA ATG GAC AAC AAC ACA GGC GCT GAC AGC AGT AAC TCA GAG AGC GAA Glu Met Asp Asn Asn Thr Gly Ala Asp Ser Ser Asn Ser Glu Ser Glu 485 490 495	1489
ACA GAG GAT GAA AGA GTA GGA GAA GAT ACG CCT TTC CTG GCC ATA CAG Thr Glu Asp Glu Arg Val Gly Glu Asp Thr Pro Phe Leu Ala Ile Gln 500 505 510	1537
AAC CCC CTG GCA GCC AGT CTC GAG GCG GCC CCT GCC TTC CGC CTG GTC Asn Pro Leu Ala Ala Ser Leu Glu Ala Ala Pro Ala Phe Arg Leu Val 515 520 525	1585
GAC AGC AGG ACT AAC CCA ACA GGC GGC TTC TCT CCG CAG GAA GAA TTG Asp Ser Arg Thr Asn Pro Thr Gly Gly Phe Ser Pro Gln Glu Glu Leu 530 535 540	1633
CAG GCC AGG CTC TCC GGT GTA ATC GCT AAC CAA GAC CCT ATC GCT GTC Gln Ala Arg Leu Ser Gly Val Ile Ala Asn Gln Asp Pro Ile Ala Val 545 550 555 560	1681
TAAAACCGAA ATACACCCAT AGATTCACCT GTAAACTTT ATTTTATATA ATAAAGTATT	1741
CCACCTTAAA TTAACAAAA AAA	1764

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 147:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 147:

Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys  
1 5 10 15  
Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys  
20 25 30  
Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser  
35 40 45  
Phe Tyr  
50

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 148:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50
- (B) TYPE: amino acid

(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 148:

Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys  
 1 5 10 15  
 Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys  
 20 25 30  
 Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn Val Pro Met Lys  
 35 40 45  
 Val Gln  
 50

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 149:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 46  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 149:

Glu Cys Leu Arg Lys Tyr Lys Asp Phe Cys Ile His Gly Glu Cys Lys  
 1 5 10 15  
 Tyr Val Lys Glu Leu Arg Ala Pro Ser Cys Lys Cys Gln Gln Glu Tyr  
 20 25 30  
 Phe Gly Glu Arg Cys Gly Glu Lys Ser Asn Lys Thr His Ser  
 35 40 45

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 150:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 198  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 150:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
1 5 10 15	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
20 25 30	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	144
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
35 40 45	
GTA ATG GCC AGC TTC TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT	192
Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro	
50 55 60	

- 81 -

EXAMPLE 19Purification and Assay of Other Proteins which bind p185<sup>erbB2</sup> ReceptorI. Purification of gp30 and p70

5 Lupu et al. (Science 249, 1552 (1990)) and Lippman and Lupu (patent application number PCT/US91/03443 (1990)), hereby incorporated by reference, have purified a protein from conditioned media of a human breast cancer cell line MDA-MB-231.

10 Lupu et al. (Proc. Natl. Acad. Sci. 89, 2287 (1992)) purified another protein which binds to the p185<sup>erbB2</sup> receptor. This particular protein, p75, was purified from conditioned medium used for the growth of SKBr-3 (a human breast cancer cell line) propagated in  
15 improved Eagle's medium (IMEM: GIBCO) supplemented with 10% fetal bovine serum (GIBCO).

II. Other p185<sup>erbB2</sup> ligands

Peles et al. (Cell 69, 205 (1992)) have also purified a 185<sup>erbB2</sup> stimulating ligand from rat cells.  
20 Holmes et al. (Science 256, 1205 (1992)) have purified Heregulin  $\alpha$  from human cells which binds and stimulates 185<sup>erbB2</sup> (see Example 5). Tarakovsky et al. Oncogene 6:218 (1991) have demonstrated binding of a 25 kD polypeptide isolated from activated macrophages to the  
25 Neu receptor, a p185<sup>erbB2</sup> homology, herein incorporated by reference.

III. NDF Isolation

Yarden and Peles (Biochemistry 30, 3543 (1991)) have identified a 35 kilodalton glycoprotein which will  
30 stimulate the 185<sup>erbB2</sup> receptor.



- 82 -

In other publications, Davis et al. (Biochem. Biophys. Res. Commun. 179, 1536 (1991), Proc. Natl. Acad. Sci. 88, 8582 (1991) and Greene et al., PCT patent application PCT/US91/02331 (1990)) describe the  
5 purification of a protein from conditioned medium of a human T-cell (ATL-2) cell line.

Huang et al. (1992, J. Biol. Chem. 257:11508-11512), hereby incorporated by reference, have isolated  
10 an additional neu/erb B2 ligand growth factor from bovine kidney. The 25 kD polypeptide factor was isolated by a procedure of column fractionation, followed by sequential column chromatography on DEAE/cellulose (DE52), Sulfadex (sulfated Sephadex G-50), heparin-Sepharose 4B, and Superdex 75 (fast protein liquid chromatography). The  
15 factor, NEL-GF, stimulates tyrosine-specific autophosphorylation of the neu/erb B2 gene product.

#### IV. Purification of acetylcholine receptor inducing activity (ARIA)

ARIA, a 42 kD protein which stimulates  
20 acetylcholine receptor synthesis, has been isolated in the laboratory of Gerald Fischbach (Falls et al., (1993) Cell 72:801-815). ARIA induces tyrosine phosphorylation of a 185 Kda muscle transmembrane protein which resembles p185<sup>erbB2</sup>, and stimulates acetylcholine receptor synthesis  
25 in cultured embryonic myotubes. ARIA is most likely a member of the GGF/erbB2 ligand group of proteins, and this is potentially useful in the glial cell mitogenesis stimulation and other applications of, e.g., GGF2 described herein.

- 83 -

EXAMPLE 19Protein tyrosine phosphorylation mediated by GGF

Rat Schwann cells, following treatment with sufficient levels of Glial Growth Factor to induce proliferation, show stimulation of protein tyrosine phosphorylation. Varying amounts of partially purified GGF were applied to a primary culture of rat Schwann cells according to the procedure outlined in Example 9. Schwann cells were grown in DMEM/10% fetal calf serum/5  $\mu$ M forskolin/0.5 $\mu$ g per mL GGF-CM (0.5mL per well) in poly D-lysine coated 24 well plates. When confluent, the cells were fed with DMEM/10% fetal calf serum at 0.5mL per well and left in the incubator overnight to quiesce. The following day, the cells were fed with 0.2mL of DMEM/10% fetal calf serum and left in the incubator for 1 hour. Test samples were then added directly to the medium at different concentrations and for different lengths of time as required. The cells were then lysed in boiling lysis buffer (sodium phosphate, 5mM, pH 6.8; SDS, 2%,  $\beta$ -mercaptoethanol, 5%; dithiothreitol, 0.1M; glycerol, 10%; Bromophenol Blue, 0.4%; sodium vanadate, 10mM), incubated in a boiling water bath for 10 minutes and then either analyzed directly or frozen at -70°C. Samples were analyzed by running on 7.5% SDS-PAGE gels and then electroblotting onto nitrocellulose using standard procedures as described by Towbin et al. (1979) Proc. Natl. Acad. Sci. USA 76:4350-4354. The blotted nitrocellulose was probed with antiphosphotyrosine antibodies using standard methods as described in Kamps and Selton (1988) Oncogene 2:305-315. The probed blots were exposed to autoradiography film overnight and developed using a standard laboratory processor. Densitometric measurements were carried out using an Ultrascan XL enhanced laser densitometer (LKB).

- 84 -

Molecular weight assignments were made relative to prestained high molecular weight standards (Sigma). The dose responses of protein phosphorylation and Schwann cell proliferation are very similar (Fig. 33). The  
5 molecular weight of the phosphorylated band is very close to the molecular weight of p185<sup>erbB2</sup>. Similar results were obtained when Schwann cells were treated with conditioned media prepared from COS cells translates with the GGF2HBS5 clone. These results correlate well with  
10 the expected interaction of the GGFs with and activation of 185<sup>erbB2</sup>.

This experiment has been repeated with recombinant GGF2. Conditioned medium derived from a CHO cell line stably transformed with the GGF2 clone (GGF2HBS5)  
15 stimulates protein tyrosine phosphorylation using the assay described above. Mock transfected CHO cells fail to stimulate this activity.

#### EXAMPLE 20

##### N-glycosylation of GGF

20 The protein sequence predicted from the cDNA sequence of GGF-II candidate clones GGF2BPP1,2 and 3 contains a number of consensus N-glycosylation motifs. A gap in the GGFII02 peptide sequence coincides with the asparagine residue in one of these motifs, indicating  
25 that carbohydrate is probably bound at this site.

N-glycosylation of the GGFs was studied by observing mobility changes on SDS-PAGE after incubation with N-glycanase, an enzyme that cleaves the covalent linkages between carbohydrate and asparagine residues in  
30 proteins.

N-Glycanase treatment of GGF-II yielded a major band of MW 40-42 kDa and a minor band at 45-48 kDa.

- 85 -

Activity single active deglycosylated species at ca 45-50 kDa.

Activity elution experiments with GGF-I also demonstrate an increase in electrophoretic mobility when  
5 treated with N-Glycanase, giving an active species of MW 26-28 kDa. Silver staining confirmed that there is a mobility shift, although no N-deglycosylated band could be assigned because of background staining in the sample used.

- 86 -

## (1) GENERAL INFORMATION:

- (i) APPLICANTS: Robert Sklar, Mark Marchionni,  
David I. Gwynne
- (ii) TITLE OF INVENTION: METHODS FOR ALTERING  
MUSCLE CONDITION
- (iii) NUMBER OF SEQUENCES: 185
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Fish & Richardson  
(B) STREET: 225 Franklin Street  
(C) CITY: Boston  
(D) STATE: Massachusetts  
(F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Diskette, 5.25 inch, 360  
kb storage  
(B) COMPUTER: IBM  
(C) OPERATING SYSTEM: PC-DOS  
(D) SOFTWARE: Wordperfect
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER: 94/05083 A  
(B) FILING DATE: 06-MAY-94  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: 08/209,204  
(B) FILING DATE: 08-MAR-94
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: 08/059,022  
(B) FILING DATE: 06-May-93
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Clark, Paul T.  
(B) REGISTRATION NUMBER: 30,162  
(C) REFERENCE/DOCKET NUMBER: 04585/028WO1
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (617) 542-5070  
(B) TELEFAX: (617) 542-8906  
(B) TELEX: 200154

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Phe Lys Gly Asp Ala His Thr Glu  
1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 2:

SUBSTITUTE SHEET (RULE 26)

- 87 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine  
 or Arginine; Xaa in position  
 12 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Xaa Ala Ser Leu Ala Asp Glu Tyr Glu Tyr Met Xaa Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 3:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine  
 or Arginine; Xaa in  
 position 10 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Xaa Thr Glu Thr Ser Ser Ser Gly Leu Xaa Leu Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 4:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or  
 Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Xaa Lys Leu Gly Glu Met Trp Ala Glu  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 5:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

- 88 -

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Xaa Leu Gly Glu Lys Arg Ala  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 6:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Xaa Ile Lys Ser Glu His Ala Gly Leu Ser Ile Gly Asp Thr Ala Lys  
 1 5 10 15

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 7:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Xaa Ala Ser Leu Ala Asp Glu Tyr Glu Tyr Met Arg Lys  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 8:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is Lysine or Arginine.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GAA TAG  
Glu  
65

198

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 151:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 192  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 151:

AGC	CAT	CTT	GTC	AAG	TGT	GCA	GAG	AAG	GAG	AAA	ACT	TTC	TGT	GTG	AAT	48
Ser	His	Leu	Val	Lys	Cys	Ala	Glu	Lys	Glu	Lys	Thr	Phe	Cys	Val	Asn	
1				5					10					15		
GGA	GGC	GAG	TGC	TTC	ATG	GTG	AAA	GAC	CTT	TCA	AAT	CCC	TCA	AGA	TAC	96
Gly	Gly	Glu	Cys	Phe	Met	Val	Lys	Asp	Leu	Ser	Asn	Pro	Ser	Arg	Tyr	
			20					25					30			
TTG	TGC	AAG	TGC	CAA	CCT	GGA	TTC	ACT	GGA	GCG	AGA	TGT	ACT	GAG	AAT	144
Leu	Cys	Lys	Cys	Gln	Pro	Gly	Phe	Thr	Gly	Ala	Arg	Cys	Thr	Glu	Asn	
			35					40						45		
GTG	CCC	ATG	AAA	GTC	CAA	ACC	CAA	GAA	AAA	GCG	GAG	GAG	CTC	TAC	TAA	192
Val	Pro	Met	Lys	Val	Gln	Thr	Gln	Glu	Lys	Ala	Glu	Glu	Leu	Tyr		
			50				55							60		

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 152:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 183  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 152:

AGC	CAT	CTT	GTC	AAG	TGT	GCA	GAG	AAG	GAG	AAA	ACT	TTC	TGT	GTG	AAT	48
Ser	His	Leu	Val	Lys	Cys	Ala	Glu	Lys	Glu	Lys	Thr	Phe	Cys	Val	Asn	
1				5					10					15		
GGA	GGC	GAG	TGC	TTC	ATG	GTG	AAA	GAC	CTT	TCA	AAT	CCC	TCA	AGA	TAC	96
Gly	Gly	Glu	Cys	Phe	Met	Val	Lys	Asp	Leu	Ser	Asn	Pro	Ser	Arg	Tyr	
			20					25					30			
TTG	TGC	AAG	TGC	CCA	AAT	GAG	TTT	ACT	GGT	GAT	CGC	TGC	CAA	AAC	TAC	144
Leu	Cys	Lys	Cys	Pro	Asn	Glu	Phe	Thr	Gly	Asp	Arg	Cys	Gln	Asn	Tyr	
			35					40						45		
GTA	ATG	GCC	AGC	TTC	TAC	AAA	GCG	GAG	GAG	CTC	TAC	TAA				183
Val	Met	Ala	Ser	Phe	Tyr	Lys	Ala	Glu	Gln	Leu	Tyr					
			50				55					60				

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 153:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 210  
(B) TYPE: nucleic acid



(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 153:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
1 5 10 15	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
20 25 30	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	144
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
35 40 45	
GTA ATG GCC AGC TTC TAC AAG CAT CTT GGG ATT GAA TTT ATG GAG AAA	192
Val Met Ala Ser Phe Tyr Lys His Leu Gly Ile Glu Phe Met Glu Lys	
50 55 60	
GCG GAG GAG CTC TAC TAA	210
Ala Glu Glu Leu Tyr	
65	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 154:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 267  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 154:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
1 5 10 15	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
20 25 30	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	144
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
35 40 45	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT	192
Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr	
50 55 60	
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC TAC AGT ACG TCC	240
Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser	
65 70 75 80	
ACT CCC TTT CTG TCT CTG CCT GAA TAG	267
Thr Pro Phe Leu Ser Leu Pro Glu	
85	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 155:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 252

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 155:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
1 5 10 15	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
20 25 30	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	144
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
35 40 45	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT	192
Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr	
50 55 60	
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC ACC TTC TAC AAA GCG GAG	240
Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Lys Ala Glu	
65 70 75 80	
GAG CTC TAC TAA	252
Glu Leu Tyr	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 156:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 128  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 156:

CC ACA TCC ACA TCT ACA GCT GGG ACA AGC CAT CTT GTC AAG TGT GCA	47
Thr Ser Thr Ser Thr Ala Gly Thr Ser His Leu Val Lys Cys Ala	
1 5 10 15	
GAG AAG GAG AAA ACT TTC TGT GTG AAT GGA GGC GAG TGC TTC ATG GTG	95
Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys Phe Met Val	
20 25 30	
AAA GAC CTT TCA AAT CCC TCA AGA TAC TTG T GC	128
Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu	
35 40	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 157:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 141  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 157:

A CAT AAC CTT ATA GCT GAG CTA AGG AGA AAC AAG GCC CAC AGA TCC 46  
 His Asn Leu Ile Ala Glu Leu Arg Arg Asn Lys Ala His Arg Ser  
 1 5 10 15

AAA TGC ATG CAG ATC CAG CTT TCC GCA ACT CAT CTT AGA GCT TCT TCC 94  
 Lys Cys Met Gln Ile Gln Leu Ser Ala Thr His Leu Arg Ala Ser Ser  
 20 25 30

ATT CCC CAT TGG GCT TCA TTC TCT AAG ACC CCT TGG CCT TTA GGA AG 141  
 Ile Pro His Trp Ala Ser Phe Ser Lys Thr Pro Trp Pro Leu Gly Arg  
 35 40 45

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 158:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24
- (B) TYPE: amino acid
- (C) STRANDEDNESS: linear
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: Xaa in positions 15 and 22 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 158:

Ala Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Xaa Phe  
 1 5 10 15

Met Val Lys Asp Leu Xaa Asn Pro  
 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 159:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 745
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

ATG AGA TGG CGA CGC GCC CCG CGC CGC TCC GGG CGT CCC GGC CCC CGG 48  
 Met Arg Trp Arg Arg Ala Pro Arg Arg Ser Gly Arg Pro Gly Pro Arg  
 1 5 10 15

GCC CAG CGC CCC GGC TCC GCC GGC ~~CGC~~ TCG TCG CCG CCG CTG CCG CTG 96  
 Ala Gln Arg Pro Gly Ser Ala Ala Arg Ser ~~CGC~~ Pro Pro Leu Pro Leu  
 20 25 30

CTG CCA CTA CTG CTG CTG CTG GGG ACC GCG ~~GCC~~ CTG GCG CCG GGG GCG 144  
 Leu Pro Leu Leu Leu Leu Leu Gly Thr Ala Ala Leu Ala Pro Gly Ala  
 35 40 45

GCG GCC GGC AAC GAG GCG GCT CCC GCG GGG GCC TCG GTG TGC TAC TCG 192  
 Ala Ala Gly Asn Glu Ala Ala Pro Ala Gly Ala Ser Val Cys Tyr Ser  
 50 55 60

TCC CCG CCC AGC GTG GGA TCG GTG CAG GAG CTA GCT CAG CGC GCC CCG 240  
 Ser Pro Pro Ser Val Gly Ser Val Gln Glu Leu Ala Gln Arg Ala Ala  
 65 70 75 80

GTG GTG ATC GAG GGA AAG GTG CAC CCG CAG CGG CGG CAG CAG GGG GCA	288
Val Val Ile Glu Gly Lys Val His Pro Gln Arg Arg Gln Gln Gly Ala	
85 90 95	
CTC GAC AGG AAG GCG GCG GCG GCG GCG GGC GAG GCA GGG GCG TGG GGC	336
Leu Asp Arg Lys Ala Ala Ala Ala Ala Gly Glu Ala Gly Ala Trp Gly	
100 105 110	
GGC GAT CGC GAG CCG CCA GCC GCG GGC CCA CGG GCG CTG GGG CCG CCC	384
Gly Asp Arg Glu Pro Pro Ala Ala Gly Pro Arg Ala Leu Gly Pro Pro	
115 120 125	
GCC GAG GAG CCG CTG CTC GCC GCC AAC GGG ACC GTG CCC TCT TGG CCC	432
Ala Glu Glu Pro Leu Leu Ala Ala Asn Gly Thr Val Pro Ser Trp Pro	
130 135 140	
ACC GCC CCG GTG CCC AGC GCC GGC GAG CCC GGG GAG GAG GCG CCC TAT	480
Thr Ala Pro Val Pro Ser Ala Gly Glu Pro Gly Glu Glu Ala Pro Tyr	
145 150 155 160	
CTG GTG AAG GTG CAC CAG GTG TGG GCG GTG AAA GCC GGG GGC TTG AAG	528
Leu Val Lys Val His Gln Val Trp Ala Val Lys Ala Gly Gly Leu Lys	
165 170 175	
AAG GAC TCG CTG CTC ACC GTG CGC CTG GGG ACC TGG GGC CAC CCC GCC	576
Lys Asp Ser Leu Leu Thr Val Arg Leu Gly Thr Trp Gly His Pro Ala	
180 185 190	
TTC CCC TCC TGC GGG AGG CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC	624
Phe Pro Ser Cys Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe	
195 200 205	
ATG GAG CCC GAC GCC AAC AGC ACC AGC CGC GCG CCG GCC GCC TTC CGA	672
Met Glu Pro Asp Ala Asn Ser Thr Ser Arg Ala Pro Ala Ala Phe Arg	
210 215 220	
GCC TCT TTC CCC CCT CTG GAG ACG GGC CGG AAC CTC AAG AAG GAG GTC	720
Ala Ser Phe Pro Pro Leu Glu Thr Gly Arg Asn Leu Lys Lys Glu Val	
225 230 235 240	
AGC CGG GTG CTG TGC AAG CGG TGC G	745
Ser Arg Val Leu Cys Lys Arg Cys	
245	

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 160:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ix) FEATURE:

- (D) OTHER INFORMATION: Xaa in position 1 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 160:

Xaa Ala Leu Ala Ala Ala Gly Tyr Asp Val Glu Lys  
 1 5 10

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 161:

- 143 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in position 1 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 161:

Xaa Leu Val Leu Arg  
 1 5

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 162:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: Xaa in positions 1, 2, and 3 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 162:

Xaa Xaa Xaa Tyr Pro Gly Gln Ile Thr Ser Asn  
 1 5 10

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 163:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 60  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N in positions 25 and 36 is unknown.

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 163:

ATAGGGAAGG GCGGGGAAG GGTCCNCCCTC NGCAGGGCCG GGCTTGCCTC TGGAGCCTCT 60

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 164:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (ix) FEATURE:

(D) OTHER INFORMATION: N in position 16 is unknown.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 164:

TTTACACATA TATTCC

18

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 165:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 165:

Glu Thr Gln Pro Asp Pro Gly Gln Ile Leu Lys Lys Val Pro Met Val  
 1 5 10 15  
 Ile Gly Ala Tyr Thr  
 20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 166:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 422  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 166:

Met Arg Trp Arg Arg Ala Pro Arg Arg Ser Gly Arg Pro Gly Pro Arg  
 1 5 10 15  
 Ala Gln Arg Pro Gly Ser Ala Ala Arg Ser Ser Pro Pro Leu Pro Leu  
 20 25 30  
 Leu Pro Leu Leu Leu Leu Leu Gly Thr Ala Ala Leu Ala Pro Gly Ala  
 35 40 45  
 Ala Ala Gly Asn Glu Ala Ala Pro Ala Gly Ala Ser Val Cys Tyr Ser  
 50 55 60  
 Ser Pro Pro Ser Val Gly Ser Val Gln Glu Leu Ala Gln Arg Ala Ala  
 65 70 75 80  
 Val Val Ile Glu Gly Lys Val His Pro Gln Arg Arg Gln Gln Gly Ala  
 85 90 95  
 Leu Asp Arg Lys Ala Ala Ala Ala Ala Gly Glu Ala Gly Ala Trp Gly  
 100 105 110  
 Gly Asp Arg Glu Pro Pro Ala Ala Gly Pro Arg Ala Leu Gly Pro Pro  
 115 120 125  
 Ala Glu Glu Pro Leu Leu Ala Ala Asn Gly Thr Val Pro Ser Trp Pro  
 130 135 140  
 Thr Ala Pro Val Pro Ser Ala Gly Glu Pro Gly Glu Glu Ala Pro Tyr  
 145 150 155 160

- 145 -

Leu Val Lys Val His Gln Val Trp Ala Val Lys Ala Gly Gly Leu Lys  
 165 170 175

Lys Asp Ser Leu Leu Thr Val Arg Leu Gly Thr Trp Gly His Pro Ala  
 180 185 190

Phe Pro Ser Cys Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe  
 195 200 205

Met Glu Pro Asp Ala Asn Ser Thr Ser Arg Ala Pro Ala Ala Phe Arg  
 210 215 220

Ala Ser Phe Pro Pro Leu Glu Thr Gly Arg Asn Leu Lys Lys Glu Val  
 225 230 235 240

Ser Arg Val Leu Cys Lys Arg Cys Ala Leu Pro Pro Gln Leu Lys Glu  
 245 250 255

Met Lys Ser Gln Glu Ser Ala Ala Gly Ser Lys Leu Val Leu Arg Cys  
 260 265 270

Glu Thr Ser Ser Glu Tyr Ser Ser Leu Arg Phe Lys Trp Phe Lys Asn  
 275 280 285

Gly Asn Glu Leu Asn Arg Lys Asn Lys Pro Gln Asn Ile Lys Ile Gln  
 290 295 300

Lys Lys Pro Gly Lys Ser Glu Leu Arg Ile Asn Lys Ala Ser Leu Ala  
 305 310 315 320

Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp  
 325 330 335

Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr  
 340 345 350

Ser Thr Thr Gly Thr Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys  
 355 360 365

Thr Phe Cys Val Asn Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser  
 370 375 380

Asn Pro Ser Arg Tyr Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp  
 385 390 395 400

Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro  
 405 410 415

Phe Leu Ser Leu Pro Glu  
 420

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 167:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 69
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 167:

Met Ser Glu Arg Lys Glu Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys  
 1 5 10 15

Glu Arg Gly Ser Gly Lys Lys Pro Glu Ser Ala Ala Gly Ser Gln Ser  
 20 25 30  
 Pro Arg Glu Ile Ile Thr Gly Met Pro Ala Ser Thr Glu Gly Ala Tyr  
 35 40 45  
 Val Ser Ser Glu Ser Pro Ile Arg Ile Ser Val Ser Thr Glu Gly Ala  
 50 55 60  
 Asn Thr Ser Ser Ser  
 65

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 168:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 168:

Arg Lys Gly Asp Val Pro Gly Pro Arg Val Lys Ser Ser Arg Ser Thr  
 1 5 10 15  
 Thr Thr Ala

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 169:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 231
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 169:

CGCGAGCGCC TCAGCGCGGC CGCTCGCTCT CCCCTCGAG GGACAAACTT TTCGCAAACC 60  
 CGATCCGAGC CCTTGGACCA AACTCGCCTG CGCCGAGAGC CGTCCGCGTA GAGCGCTCCG 120  
 TCTCCGGCGA GATGTCCGAG CGCAAAGAAG GCAGAGGCCAA AGGGAAGGGC AAGAAGAAGG 180  
 AGCCGAGGCTC CGGCAAGAAG CCGGAGTCCG CGGCGGGCAG CCAGAGCCCA G 231

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 170:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 178
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 170:

CCTTGCCTCC CCGATTGAAA GAGATGAAA GCCAGGAATC GGCTGCAGGT TCCAAACTAG 60  
 TCCTTCGGTG TGAACCAGT TCTGAATACT CCTCTCTCAG ATTCAAGTGG TTCAAGAATG 120  
 GGAATGAATT GAATCGAAA AACAAACCAC AAAATATCAA GATACAAAAA AAGCCAGG 178



## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 171:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 122  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 171:

GAAGTCAGAA CTTCGCATTA ACAAAGCATC ACTGGCTGAT TCTGGAGAGT ATATGTGCAA 60  
 AGTGATCAGC AAATTAGGAA ATGACAGTGC CTCTGCCAAT ATCACCATCG TGGAAATCAA 120  
 CG 122

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 172:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 172:

AGATCATCAC TGGTATGCCA GCCTCAACTG AAGGAGCATA TGTGTCTTCA GAGTCTCCCA 60  
 TTAGAATATC AGTATCCACA GAAGGAGCAA ATACTTCTTC AT 102

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 173:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 128  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 173:

CTACATCTAC ATCCACCACT GGGACAAGCC ATCTTGTAAG ATGTGCGGAG AAGGAGAAAA 60  
 CTTTCTGTGT GAATGGAGGG GAGTGCTTCA TGGTAAAAGA CCTTCAAAC CCCTCGAGAT 120  
 ACTTGTGC 128

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 174:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 69  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 174:

AAGTGCCAAC CTGGATTCAC TGGAGCAAGA TGTACTGAGA ATGTGCCCAT GAAAGTCCAA 60  
 AACCAAGAA 69

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 175:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 60  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 175:

AAGTGCCCAA ATGAGTTTAC TGGTGATCGC TGCCAAACT ACGTAATGGC CAGCTTCTAC 60

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 176:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 176:

AGTACGTCCA CTCCCTTCT GTCTCTGCCT GAATAG 36

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 177:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 569  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 177:

AAGGCGGAGG AGCTGTACCA GAAGAGAGTG CTGACCATAA CCGGCATCTG CATCGCCCTC 60  
 CTGTGGTTCG GCATCATGTG TGTGGTGGCC TACTGCCAAA CCAAGAAACA GCCGAAAAG 120  
 CTGCATGACC GTCTTCGGCA GAGCCTTCGG TCTGAACGAA ACAATATGAT GAACATTGCC 180  
 AATGGGCCTC ACCATCCTAA CCCACCCCCC GAGAATGTCC AGCTGGTGAA TCAATACGTA 240  
 TCTAAAACG TCATCTCCAG TGAGCATATT GTTGAGAGAG AAGCAGAGAC ATCCTTTTCC 300  
 ACCAGTCACT ATACTTCCAC AGCCCATCAC TCCACTACTG TCACCCAGAC TCCTAGCCAC 360  
 AGCTGGAGCA ACGGACACAC TGAAAGCATC CTTTCCGAAA GCCACTCTGT AATCGTGATG 420  
 TCATCCGTAG AAAACAGTAG GCACAGCAGC CCAACTGGGG GCCCAAGAGG ACGTCTTAAT 480  
 GGCACAGGAG GCCCTCGTGA ATGTAACAGC TTCCTCAGGC ATGCCAGAGA AACCCCTGAT 540  
 TCCTACCGAG ACTCTCCTCA TAGTGAAAG 569

## (2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 178:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 730  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single

- 149 -

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 178:

GTATGTGTCA GCCATGACCA CCCC GGCTCG TATGTCACCT GTAGATTTC ACACGCCAAG 60  
 CTCCCCCAAA TCGCCCCCTT CGGAAATGTC TCCACCCGTG TCCAGCATGA CGGTGTCCAT 120  
 GCCTTCCATG GCGGTCAGCC CCTTCATGGA AGAAGAGAGA CCTCTACTTC TCGTGACACC 180  
 ACCAAGGCTG CGGGAGAAGA AGTTTGACCA TCACCCTCAG CAGTTCAGCT CCTTCCACCA 240  
 CAACCCCGCG CATGACAGTA ACAGCCTCCC TGCTAGCCCC TTGAGGATAG TGGAGGATGA 300  
 GGAGTATGAA ACGACCCAAG AGTACGAGCC AGCCCAAGAG CCTGTTAAGA AACTCGCCAA 360  
 TAGCCGGCGG GCCAAAAGAA CCAAGCCCAA TGGCCACATT GCTAACAGAT TGGAAAGTGA 420  
 CAGCAACACA AGCTCCCGA GCAGTAACTC AGAGAGTGAA ACAGAAGATG AAAGAGTAGG 480  
 TGAAGATACG CCTTTCCTGG GCATACAGAA CCCCCTGGCA GCCAGTCTTG AGGCAACACC 540  
 TGCCTTCCGC CTGGCTGACA GCAGGACTAA CCCAGCAGGC CGCTTCTCGA CACAGGAAGA 600  
 AATCCAGGCC AGGCTGTCTA GTGTAATTGC TAACCAAGAC CCTATTGCTG TATAAACCT 660  
 AAATAAACAC ATAGATTCAC CTGTAAACT TTATTTTATA TAATAAAGTA TTCCACCTTA 720  
 AATTAAACAA 730

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 179:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 179:

TCGGGCTCCA TGAAGAAGAT GTA 23

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 180:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 180:

TCCATGAAGA AGATGTACCT GCT 23

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 181:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 181:

ATGTACCTGC TGCTCCTT GA

22

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 182:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 182:

TTGAAGAAGG ACTCGCTGCT CA

22

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 183:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 183:

AAAGCCGGGG GCTTGAAGAA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 184:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 184:

ATGARGTGTG GGCGGCGAAA

20

(2) INFORMATION FOR SEQUENCE IDENTIFICATION NUMBER: 185:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 422  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 185:

Met Arg Trp Arg Arg Ala Pro Arg Arg Ser Gly Arg Pro Gly Pro Arg  
1 5 10 15Ala Gln Arg Pro Gly Ser Ala Ala Arg Ser Ser Pro Pro Leu Pro Leu  
20 25 30Leu Pro Leu Leu Leu Leu Gly Thr Ala Ala Leu Ala Pro Gly Ala  
35 40 45

Ala Ala Gly Asn Glu Ala Ala Pro Ala Gly Ala Ser Val Cys Tyr Ser  
 50 55 60  
 Ser Pro Pro Ser Val Gly Ser Val Gln Glu Leu Ala Gln Arg Ala Ala  
 65 70 75 80  
 Val Val Ile Glu Gly Lys Val His Pro Gln Arg Arg Gln Gln Gly Ala  
 85 90 95  
 Leu Asp Arg Lys Ala Ala Ala Ala Ala Gly Glu Ala Gly Ala Trp Gly  
 100 105 110  
 Gly Asp Arg Glu Pro Pro Ala Ala Gly Pro Arg Ala Leu Gly Pro Pro  
 115 120 125  
 Ala Glu Glu Pro Leu Leu Ala Ala Asn Gly Thr Val Pro Ser Trp Pro  
 130 135 140  
 Thr Ala Pro Val Pro Ser Ala Gly Glu Pro Gly Glu Glu Ala Pro Tyr  
 145 150 155 160  
 Leu Val Lys Val His Gln Val Trp Ala Val Lys Ala Gly Gly Leu Lys  
 165 170 175  
 Lys Asp Ser Leu Leu Thr Val Arg Leu Gly Thr Trp Gly His Pro Ala  
 180 185 190  
 Phe Pro Ser Cys Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe  
 195 200 205  
 Met Glu Pro Asp Ala Asn Ser Thr Ser Arg Ala Pro Ala Ala Phe Arg  
 210 215 220  
 Ala Ser Phe Pro Pro Leu Glu Thr Gly Arg Asn Leu Lys Lys Glu Val  
 225 230 235 240  
 Ser Arg Val Leu Cys Lys Arg Cys Ala Leu Pro Pro Gln Leu Lys Glu  
 245 250 255  
 Met Lys Ser Gln Glu Ser Ala Ala Gly Ser Lys Leu Val Leu Arg Cys  
 260 265 270  
 Glu Thr Ser Ser Glu Tyr Ser Ser Leu Arg Phe Lys Trp Phe Lys Asn  
 275 280 285  
 Gly Asn Glu Leu Asn Arg Lys Asn Lys Pro Gln Asn Ile Lys Ile Gln  
 290 295 300  
 Lys Lys Pro Gly Lys Ser Glu Leu Arg Ile Asn Lys Ala Ser Leu Ala  
 305 310 315 320  
 Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp  
 325 330 335  
 Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr  
 340 345 350  
 Ser Thr Thr Gly Thr Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys  
 355 360 365  
 Thr Phe Cys Val Asn Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser  
 370 375 380  
 Asn Pro Ser Arg Tyr Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp  
 385 390 395 400

Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro  
                  405                  410                  415  
Phe Leu Ser Leu Pro Glu  
                  420

What is claimed is:

- 153 -

1. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide encoded by pGGF2HBS5 deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347) with a pharmaceutical carrier.

2. A method of making a medicament for the treating of muscle cells of a mammal, said method comprising admixing a polypeptide encoded by the E sequence (SEQ ID Nos. 133 and 159) and at least a portion of the peptide encoded by the DNA sequences flanking the E encoding sequence on clone pGGF2HBS5, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).

3. A method of making a medicament for the treating of muscle cells of a mammal, said method comprising admixing a polypeptide defined by the formula

YBAZCX

wherein YBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 133-135, 156, 159); wherein Y comprises polypeptide segment E, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL with a pharmaceutical carrier.

- 154 -

4. A method of making a medicament for the treating of muscle cells of a mammal, said method comprising admixing a polypeptide defined by the formula

WBAZCX

5            wherein WBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 134, 135, 137-139, 156); wherein W comprises polypeptide segment F, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide  
10 segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL with a pharmaceutical carrier.

15            5. The method of any one of claims 1-3, wherein 50 N-terminal amino acids are cleaved from said peptide comprising the E sequence (SEQ ID Nos. 133 and 159).

6. The method of claim 3 or 4, wherein X is C/D  
HKL.

20            7. The method of claim 3 or 4, wherein X is C/D  
H.

8. The method of claim 3 or 4, wherein X is C/D  
HL.

25            9. The method of claim 3 or 4, wherein X is C/D  
D.

10. The method of claim 3 or 4, wherein X is C/D'  
HL.



- 155 -

11. The method of claim 3 or 4, wherein X is C/D'  
HKL.
12. The method of claim 3 or 4, wherein X is C/D'  
H.
- 5 13. The method of claim 3 or 4, wherein X is C/D'  
D.
14. The method of claim 3 or 4, wherein X is C/D  
C/D' HKL.
- 15 15. The method of claim 3 or 4, wherein X is C/D  
10 C/D' H.
16. The method of claim 3 or 4, wherein X is C/D  
C/D' HL.
17. The method of claim 3 or 4, wherein X is C/D  
C/D' D.
- 15 18. The method of claim 3 or 4, wherein X is C/D  
D' H.
19. The method of claim 3 or 4, wherein X is C/D  
D' HL.
- 20 20. The method of claim 3 or 4, wherein X is C/D  
20 D' HKL.
21. The method of claim 3 or 4, wherein X is C/D'  
D' H.

- 156 -

22. The method of claim 3 or 4, wherein X is C/D'  
D' HL.
23. The method of claim 3 or 4, wherein X is C/D'  
D' HKL.
- 5 24. The method of claim 3 or 4, wherein X is C/D  
C/D' D' H.
25. The method of claim 3 or 4, wherein X is C/D  
C/D' D' HL.
- 10 26. The method of claim 3 or 4, wherein X is C/D  
C/D' D' HKL.
27. A method of making a medicament for the  
treating of muscle cells of a mammal said method  
comprising admixing a polypeptide comprising FBA  
polypeptide segments having the amino acid sequences  
15 shown in Fig. 30 (SEQ ID Nos. 132, 134, 135) with a  
pharmaceutically acceptable carrier.
28. A method of making a medicament for the  
treating of muscle cells of a mammal said method  
comprising admixing a polypeptide comprising FBA'  
20 polypeptide segments having the amino acid sequences  
shown in Fig. 30 (SEQ ID Nos. 132, 134, 136) with a  
pharmaceutically acceptable carrier.
29. A method of making a medicament for the  
treating of muscle cells of a mammal said method  
25 comprising admixing a polypeptide comprising FEBA  
polypeptide segments having the amino acid sequences  
shown in Fig. 30 (SEQ ID Nos. 132, 135, 159) with a  
pharmaceutically acceptable carrier.

- 157 -

30. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising FEBA' polypeptide segments having the amino acid sequences  
5 corresponding to polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132-134, 136, 159) to muscle cells with a pharmaceutically acceptable carrier.

31. A method of making a medicament for the treating of muscle cells of a mammal said method  
10 comprising admixing GGF2 polypeptide with a pharmaceutically acceptable carrier.

32. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a compound which specifically binds  
15 the p185<sup>erbB2</sup> receptor of muscle cells with a pharmaceutically acceptable carrier.

33. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising EGFL1,  
20 having the amino acid sequence shown Fig. 37, Seq. ID No. 150, with a pharmaceutically acceptable carrier.

34. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising EGFL2,  
25 having the amino acid sequence shown in Fig. 38, Seq. ID No. 151, with a pharmaceutically acceptable carrier.

- 158 -

35. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising EGFL3, with the amino acid sequence shown in Fig. 39, Seq. ID No. 5 152, with a pharmaceutically acceptable carrier.

36. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising EGFL4, with the amino acid sequence shown in Fig. 40, Seq. ID No. 10 153, with a pharmaceutically acceptable carrier.

37. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide comprising EGFL5, with the amino acid sequence shown in Fig. 41, Seq. ID No. 15 154, to muscle cells, with a pharmaceutically acceptable carrier.

38. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a polypeptide, comprising EGFL6, with 20 the amino acid sequence shown Fig. 42, Seq. ID No. 155, with a pharmaceutically acceptable carrier.

39. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 35 kD polypeptide factor isolated 25 from the rat I-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

- 159 -

40. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 75 kD polypeptide factor isolated from the SKBR-3 human breast cell line to said muscle  
5 cells, with a pharmaceutically acceptable carrier.

41. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 44 kD polypeptide factor isolated from the rat I-EJ transformed fibroblast cell line to  
10 said muscle cells, with a pharmaceutically acceptable carrier.

42. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 45 kD polypeptide factor isolated  
15 from the MDA - MB 231 human breast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

43. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 7 to 14 kD polypeptide factor  
20 isolated from the ATL-2 human T-cell line to said muscle cells, with a pharmaceutically acceptable carrier.

44. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 25 kD polypeptide factor isolated  
25 from activated mouse peritoneal macrophages to said muscle cells, with a pharmaceutically acceptable carrier.

- 160 -

45. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 25 kD polypeptide factor isolated from bovine kidney to said muscle cells, with a  
5 pharmaceutically acceptable carrier.

46. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a ARIA polypeptide to said muscle cells, with a pharmaceutically acceptable carrier.

10 47. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing a 46-47 kD polypeptide factor which stimulates 0-2A glial progenitor cells to said muscle cells, with a pharmaceutically acceptable carrier.

15 48. A method of making a medicament for the treating of muscle cells of a mammal said method comprising admixing GGF-III to said muscle cells, with a pharmaceutically acceptable carrier.

- 161 -

49. A method of making a medicament for the treating of muscle cells of a mammal, said method comprising admixing with a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the  
5 formula

YBAZCX

wherein YBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 133-135, 156, 159); wherein Y comprises polypeptide segment E, or is  
10 absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D'  
15 HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA in an expressible genetic construction.

- 162 -

50. A method of making a medicament for the treating of muscle cells of a mammal, said method comprising admixing with a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the  
5 formula

## WBAZCX

wherein WBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 134, 135, 137-139, 156); wherein W comprises polypeptide segment F,  
10 or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D'  
15 H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA in an expressible genetic construction.

51. A method of making a medicament for the prophylaxis or treatment of pathophysiological condition  
20 of the musculature in a mammal in which said condition involves a muscle cell type which is sensitive or responsive to a polypeptide as defined in any one of claims 1, 3, 4, and 31, said method comprising admixing an effective amount of said polypeptide with a  
25 pharmaceutically acceptable carrier.

52. A method of making a medicament for the treatment of a condition which involves muscle damage in a mammal, said method comprising admixing an effective amount of a polypeptide, as defined in any one of claims  
30 1, 3, 4, and 31 with a pharmaceutically acceptable carrier.



- 163 -

53. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for decreasing the atrophy of said muscle cells.

54. The method of any one of claims 1, 3, 4, and 5 31, wherein said medicament is for increasing the muscle fibers present in said mammal.

55. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for increasing muscle cell survival in a said mammal.

10 56. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for increasing muscle growth in a said mammal.

57. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for increasing muscle 15 regeneration in a said mammal.

58. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for stimulating muscle cell mitogenesis.

59. The method of any one of claims 1, 3, 4, and 20 31, wherein said medicament is for increasing acetylcholine receptor synthesis.

60. The method of any one of claims 1, 3, 4, and 31, wherein said medicament is for treating a patient lacking a neurotrophic factor.

- 164 -

61. A method of claims 1, 3, 4, and 31, wherein said medicament is for treating a muscle cell which is a myoblast.

5 62. A method of claims 1, 3, 4, and 31, wherein said medicament is for treating a muscle cell which is a satellite cell.

63. A method of claims 1, 3, 4, and 31 wherein said medicament is for treating a muscle cell in skeletal muscle.

10 64. A method of claims 1, 3, 4, and 31 wherein said medicament is for treating a muscle cell in cardiac muscle.

15 65. A method of claims 1, 3, 4, and 31 wherein said medicament is for treating a muscle cell in smooth muscle.

66. A method of claims 1, 3, 4, and 31, wherein said medicament is for treating a muscle cell in a patient with a skeletal muscle disease.

20 67. A method of claim 66, wherein said skeletal muscle disease is a myopathy.

68. A method of claim 66, wherein said skeletal muscle disease is a dystrophy.

69. A method of claim 68, wherein said dystrophy is Duchennes muscular dystrophy.

- 165 -

70. A method of claim 68, wherein said dystrophy is Beckker's dystrophy.

71. A method of claim 66, wherein said skeletal muscle disease is a result of a neural condition.

5 72. A method of claim 66, wherein said skeletal muscle disease is an injury.

73. A method of claim 66, wherein said skeletal muscle disease is resulting from a nerve injury.

10 74. A method of claim 66, wherein said skeletal muscle disease is resulting from a neuropathy.

75. A method of claims 1, 3, 4, and 31, wherein said medicament is for treating a muscle cell in a patient with a cardiac muscle disorder.

15 76. A method of claim 75, wherein said cardiac disorder is cardiomyopathy.

77. A method of claim 75, wherein said cardiac disorder is ischemic damage.

78. A method of claim 75, wherein said cardiac disorder is a congenital disease.

20 79. A method of claim 75, wherein said cardiac disorder is cardiac trauma.

80. A method of claims 1, 3, 4, and 31, wherein said medicament is for treating a muscle cell in a patient with a smooth muscle disorder.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide encoded by pGGF2HBS11, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).
2. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide encoded by the E sequence (SEQ ID Nos. 133 and 159) and at least a portion of the peptide encoded by the DNA sequences flanking the E encoding sequence on clone pGGF2HBS11, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).
3. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide defined by the formula



WYBAZCX

wherein WYBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 133-135, 156, 159); wherein W comprises polypeptide segment F, or is absent; wherein Y comprises polypeptide segment E, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL with a pharmaceutical carrier.

4. The method of any one of claims 1-3, wherein 50 N-terminal amino acids are cleaved from said peptide comprising the E sequence (SEQ ID Nos. 133 and 159).



5. The method of claim 3, wherein X is C/D HKL.
6. The method of claim 3, wherein X is C/D H.
7. The method of claim 3, wherein X is C/D HL.
8. The method of claim 3, wherein X is C/D D.
9. The method of claim 3, wherein X is C/D' HL.
10. The method of claim 3, wherein X is C/D' HKL.
11. The method of claim 3, wherein X is C/D' H.
12. The method of claim 3, wherein X is C/D' D.
13. The method of claim 3, wherein X is C/D C/D' HKL.
14. The method of claim 3, wherein X is C/D C/D' H.
15. The method of claim 3, wherein X is C/D C/D' HL.
16. The method of claim 3, wherein X is C/D C/D' D.
17. The method of claim 3, wherein X is C/D D' H.
18. The method of claim 3, wherein X is C/D D' HL.
19. The method of claim 3, wherein X is C/D D' HKL.



20. The method of claim 3, wherein X is C/D' D' H.
21. The method of claim 3, wherein X is C/D' D' HL.
22. The method of claim 3, wherein X is C/D' D' HKL.
23. The method of claim 3, wherein X is C/D C/D' D' H.
24. The method of claim 3, wherein X is C/D C/D' D' HL.
25. The method of claim 3, wherein X is C/D C/D' D' HKL.
26. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising FBA polypeptide segments having the amino acid sequences shown in Fig. 30 (SEQ ID Nos. 132, 134, 135) with a pharmaceutically acceptable carrier.
27. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising FBA' polypeptide segments having the amino acid sequences shown in Fig. 30 (SEQ ID Nos. 132, 134, 136) with a pharmaceutically acceptable carrier.
28. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising FEBA polypeptide segments having the amino acid sequences shown in Fig. 30 (SEQ ID Nos. 132, 135, 159) with a pharmaceutically acceptable carrier.



29. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising FEBA' polypeptide segments having the amino acid sequences corresponding to polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132-134, 136, 159) to muscle cells with a pharmaceutically acceptable carrier.

30. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal GGF2 polypeptide with a pharmaceutically acceptable carrier.

31. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a compound which specifically binds the P185<sup>erbB2</sup> receptor of muscle cells with a pharmaceutically acceptable carrier.

32. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL1, having the amino acid sequence shown in Fig. 37, SEQ ID No. 150, with a pharmaceutically acceptable carrier.

33. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL2, having the amino acid sequence shown in Fig. 38, SEQ ID No. 151, with a pharmaceutically acceptable carrier.

34. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL3 (SEQ ID No. 152), with a pharmaceutically acceptable carrier.



35. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL4 (SEQ ID No. 153), with a pharmaceutically acceptable carrier.

36. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL5 (SEQ ID No. 154), to muscle cells, with a pharmaceutically acceptable carrier.



37. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a polypeptide comprising EGFL6 (SEQ ID No. 155), with a pharmaceutically acceptable carrier.



38. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 35 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the rat 1-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

39. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 75 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the SKBR-3 human breast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

40. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 44 kD polypeptide factor having muscle cell mitogenesis differentiation and/or





survival inducing properties and isolated from the rat I-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

41. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 45 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the MDA - MB 231 human breast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

42. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 7 to 14 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the ATL-2 human T-cell line to said muscle cells, with a pharmaceutically acceptable carrier.

43. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 25 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from activated mouse peritoneal macrophages to said muscle cells, with a pharmaceutically acceptable carrier.

44. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 25 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from bovine kidney to said muscle cells, with a pharmaceutically acceptable carrier.

45. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a ARIA polypeptide having muscle cell mitogenesis differentiation and/or



survival inducing properties to said muscle cells, with a pharmaceutically acceptable carrier.

46. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a 46-47 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and which stimulates 0-2A glial progenitor cells to said muscle cells, with a pharmaceutically acceptable carrier.

47. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal GGF-III to said muscle cells, with a pharmaceutically acceptable carrier.

48. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the formula

YBAZCX


wherein YBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 133-135, 156, 159); wherein Y comprises polypeptide segment E, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA is an expressible genetic construction.

49. A method of treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal, said method comprising administering to said mammal a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the formula




**WBAZCX**

wherein WBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 134, 135, 137-139, 156); wherein W comprises polypeptide segment F, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA is an expressible genetic construction.



50. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide encoded by pGGF2HBS11, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).



51. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide encoded by the E sequence (SEQ ID Nos. 133 and 159) and at least a portion of the peptide encoded by the DNA sequences flanking the E encoding sequence on clone pGGF2HBS11, deposited with the A.T.C.C. November 6, 1992 (A.T.C.C. Deposit No. 75347).



52. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide defined by the formula

**WYBAZCX**

wherein WYBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 133-135, 156, 159); wherein W comprises polypeptide segment F, or is absent; wherein Y comprises polypeptide segment E, or is absent; wherein Z comprises



polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL with a pharmaceutical carrier.

53. The composition according to any one of claims 50-52, wherein 50 N-terminal amino acids are cleaved from said peptide comprising the E sequence (SEQ ID Nos. 133 and 159).

54. The composition according to claim 52, wherein X is C/D HKL.

55. The composition of claim 52, wherein X is C/D H.

56. The composition of claim 52, wherein X is C/D HL.

57. The composition of claim 52, wherein X is C/D D.

58. The composition of claim 52, wherein X is C/D' HL.

59. The composition of claim 52, wherein X is C/D' HKL.

60. The composition of claim 52, wherein X is C/D' H.

61. The composition of claim 52, wherein X is C/D' D.

62. The composition of claim 52, wherein X is C/D C/D' HKL.

63. The composition of claim 52, wherein X is C/D C/D' H.



64. The composition of claim 52, wherein X is C/D C/D' HL.
65. The composition of claim 52, wherein X is C/D C/D' D.
66. The composition of claim 52, wherein X is C/D D' H.
67. The composition of claim 52, wherein X is C/D D' HL.
68. The composition of claim 52, wherein X is C/D D' HKL.
69. The composition of claim 52, wherein X is C/D' D' H.
70. The composition of claim 52, wherein X is C/D' D' HL.
71. The composition of claim 52, wherein X is C/D' D' HKL.
72. The composition of claim 52, wherein X is C/D C/D' D' H.
73. The composition of claim 52, wherein X is C/D C/D' D' HL.
74. The composition of claim 52, wherein X is C/D C/D' D' HKL.
75. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising FBA polypeptide segments having the amino acid sequences shown in Fig. 30 (SEQ ID Nos. 132, 134, 135) with a pharmaceutically acceptable carrier.
76. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising FBA' polypeptide segments having the amino acid sequences shown in Fig.



30 (SEQ ID Nos. 132, 134, 136) with a pharmaceutically acceptable carrier.

77. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising FEBA polypeptide segments having the amino acid sequences shown in Fig. 30 (SEQ ID Nos. 132, 135, 159) with a pharmaceutically acceptable carrier.

78. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising FEBA' polypeptide segments having the amino acid sequences corresponding to polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132-134, 136, 159) to muscle cells with a pharmaceutically acceptable carrier.

79. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a GGF2 polypeptide with a pharmaceutically acceptable carrier.

80. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a compound which specifically binds the P185<sup>erbB2</sup> receptor of muscle cells with a pharmaceutically acceptable carrier.

81. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL1, having the amino acid sequence shown in Fig. 37, SEQ ID No. 150, with a pharmaceutically acceptable carrier.


82. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL2, having the amino acid sequence shown in Fig. 38, SEQ ID No. 151,




with a pharmaceutically acceptable carrier.

83. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL3 (SEQ ID No. 152), with a pharmaceutically acceptable carrier.


84. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL4 (SEQ ID No. 153), with a pharmaceutically acceptable carrier.



85. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL5 (SEQ ID No. 154), to muscle cells, with a pharmaceutically acceptable carrier.



86. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a polypeptide comprising EGFL6 (SEQ ID No. 155), with a pharmaceutically acceptable carrier.



87. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 35 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the rat I-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

88. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 75 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the SKBR-3 human breast cell line to said muscle cells, with a pharmaceutically acceptable carrier.



89. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 44 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the rat I-EJ transformed fibroblast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

90. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 45 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the MDA - MB 231 human breast cell line to said muscle cells, with a pharmaceutically acceptable carrier.

91. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 7 to 14 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from the ATL-2 human T-cell line to said muscle cells, with a pharmaceutically acceptable carrier.

92. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 25 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from activated mouse peritoneal macrophages to said muscle cells, with a pharmaceutically acceptable carrier.

93. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 25 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and isolated from bovine kidney to said muscle cells, with a pharmaceutically acceptable carrier.





94. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising an ARIA polypeptide having muscle cell mitogenesis differentiation and/or survival inducing properties to said muscle cells, with a pharmaceutically acceptable carrier.

95. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a 46-47 kD polypeptide factor having muscle cell mitogenesis differentiation and/or survival inducing properties and which stimulates 0-2A glial progenitor cells to said muscle cells, with a pharmaceutically acceptable carrier.

96. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising GGF-III to said muscle cells, with a pharmaceutically acceptable carrier.

97. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the formula



wherein YBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 133-135, 156, 159); wherein Y comprises polypeptide segment E, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA is an expressible genetic construction.



98. A composition when used for treating muscle cells to increase mitogenesis, differentiation and/or survival of said muscle cell in a mammal comprising a pharmaceutically acceptable carrier, a DNA sequence encoding a polypeptide of the formula

**WBAZCX**

wherein WBAZCX is composed of the polypeptide segments shown in Fig. 30 (SEQ ID Nos. 132, 134, 135, 137-139, 156); wherein W comprises polypeptide segment F, or is absent; wherein Z comprises polypeptide segment G or is absent; and wherein X comprises polypeptide segments C/D HKL, C/D H, C/D HL, C/D D, C/D' HL, C/D' HKL, C/D' H, C/D' D, C/D C/D' HKL, C/D C/D' H, C/D C/D' HL, C/D C/D' D, C/D D' H, C/D D' HL, C/D D' HKL, C/D' D' H, C/D' D' HL, C/D' D' HKL, C/D C/D' D' H, C/D C/D' D' HL, or C/D C/D' D' HKL, said DNA is an expressible genetic construction.

99. A method of prophylaxis or treatment of pathophysiological condition of the musculature in a mammal in which said condition involves a muscle cell type which is sensitive or responsive to a polypeptide as defined in any one of claims 1, 3, and 30, said method comprising administering to said mammal an effective amount of said polypeptide with a pharmaceutically acceptable carrier.

100. A method of treatment of a condition which involves muscle damage in a mammal, said method comprising administering to said mammal an effective amount of a polypeptide, as defined in any one of claims 1, 3, and 30 with a pharmaceutically acceptable carrier.

101. The method of any one of claims 1, 3, or 30, wherein said method is for decreasing the atrophy of said muscle cells.



102. The method of any one of claims 1, 3, or 30, wherein said method is for increasing the muscle fibers present in said mammal.

103. The method of any one of claims 1, 3, or 30, wherein said method is for increasing muscle cell survival in a said mammal.

104. The method of any one of claims 1, 3, or 30, wherein said method is for increasing muscle growth in a said mammal.

105. The method of any one of claims 1, 3, or 30, wherein said method is for increasing muscle regeneration in a said mammal.

106. The method of any one of claims 1, 3, or 30, wherein said method is for stimulating muscle cell mitogenesis.

107. The method of any one of claims 1, 3, or 30, wherein said method is for increasing acetylcholine receptor synthesis.

108. The method of any one of claims 1, 3, or 30, wherein said method is for treating a patient lacking a neurotrophic factor.

109. A method of claim 3, wherein said method is for treating a muscle cell which is a myoblast.

110. A method of claim 3, wherein said method is for treating a muscle cell which is a satellite cell.

111. A method of claim 3, wherein said method is for treating a muscle cell in skeletal muscle.



112. A method of claim 3, wherein said method is for treating a muscle cell in cardiac muscle.

113. A method of claim 3, wherein said method is for treating a muscle cell in smooth muscle.

114. A method of claim 3, wherein said method is for treating a muscle cell in a patient with a skeletal muscle disease.

115. A method of claim 114, wherein said skeletal muscle disease is a myopathy.

116. A method of claim 114, wherein said skeletal muscle disease is a dystrophy.

117. A method of claim 116, wherein said dystrophy is Duchennes muscular dystrophy.

118. A method of claim 116, wherein said dystrophy is Beckker's dystrophy.

119. A method of claim 114, wherein said skeletal muscle disease is a result of a neural condition.

120. A method of claim 114, wherein said skeletal muscle disease is an injury.

121. A method of claim 114, wherein said skeletal muscle disease is resulting from a nerve injury.

122. A method of claim 114, wherein said skeletal muscle disease is resulting from a neuropathy.

123. A method of claim 3, wherein said method is for treating a muscle cell in a patient with a cardiac muscle disorder.



- 183 -

124. A method of claim 123, wherein said cardiac disorder is cardiomyopathy.
125. A method of claim 123, wherein said cardiac disorder is ischemic damage.
126. A method of claim 123, wherein said cardiac disorder is a congenital disease.
127. A method of claim 123, wherein said cardiac disorder is cardiac trauma.
128. A method of claim 3, wherein said method is for treating a muscle cell in a patient with a smooth muscle disorder.
129. A method of claim 128, wherein said disorder is arterial sclerosis.
130. A method of claim 128, wherein said disorder is a vascular lesion.
131. A method of claim 128, wherein said disorder is a congenital vascular disease.
132. A method of claim 3, wherein said method is for treating a muscle cell which has insufficient functional acetylcholine receptors.
133. A method of claim 132, wherein said muscle cell lacking sufficient acetylcholine receptor is a muscle cell in a patient with myasthenia gravis.
134. A method as claimed in claim 132, wherein said condition involves muscular damage.
135. A method prophylaxis or treatment of a muscular tumor in a patient, said method comprising administering an effective amount of a substance which inhibits the binding of a polypeptide as defined in any one of claims 1, 3, and 30 to a receptor therefor with a pharmaceutically acceptable carrier.



136. A method of treatment of a mammal suffering from a disease of muscle cell proliferation, said method comprising administering an antibody which binds to a polypeptide of any of one of claims 1, 3, and 30 with a pharmaceutically acceptable carrier.

137. A method of identifying a nucleic acid sequence encoding a polypeptide, said polypeptide capable of binding the p185<sup>erbB2</sup> receptor and coding for a molecule having muscle cell mitogenic activity, said method comprising contacting a cell containing sample with a muscle cell mitogen specific antibody to determine expression of said mitogen in said sample and isolating said nucleic acid sequence from the cells exhibiting said expression.

138. The method of claim 30, wherein said GGF2 is human recombinant GGF2.

139. A method of stimulating myogenesis of a muscle cell said method comprising contacting said muscle cell with a polypeptide which specifically binds the p185<sup>erbB2</sup> receptor of muscle cells.

Dated this 3rd day of April, 1998

**Cambridge Neuroscience**  
by their Patent Attorneys  
DAVIES COLLISON CAVE






FIG. 1




FIG. 2





FIG. 3





2 298 6078




FIG. 4




FIG. 5




FIG. 6




FIG. 7

955  
92  
4

## FIG. 8

	N-terminus		
GGF-I 01	F K G D A H T E	(SEQ ID NO: 1)	
	Trypsin peptides		
GGF-I 02	K/R A S L A D E Y E Y M X K *	(SEQ ID NO: 2)	
GGF-I 03	K/R T E T S S S G L X L K *	(SEQ ID NO: 3)	
GGF-I 04	K/R K L G E M W A E	(SEQ ID NO: 4)	HMG-1
GGF-I 05	K/R L G E K R A	(SEQ ID NO: 5)	HMG-1?
GGF-I 06	K/R I K S E H A G L S I G D T A K *	(SEQ ID NO: 6)	HMG-2
GGF-I 07	K/R A S L A D E Y E Y M R K *	(SEQ ID NO: 7)	
GGF-I 08	K/R I K G E H P G L S I G D V A K *	(SEQ ID NO: 8)	HMG-1
GGF-I 09	K/R M S E Y A F F V Q T X R *	(SEQ ID NO: 9)	HMG-2
GGF-I 10	K/R S E H P G L S I G D T A K *	(SEQ ID NO: 10)	HMG-1
GGF-I 11	K/R A G Y F A E X A R *	(SEQ ID NO: 11)	
GGF-I 12	K/R K L E F L X A K *	(SEQ ID NO: 12)	
GGF-I 13	K/R T T E M A S E Q G A	(SEQ ID NO: 13)	
GGF-I 14	K/R A K E A L A A L K *	(SEQ ID NO: 14)	
GGF-I 15	K/R F V L Q A K K *	(SEQ ID NO: 15)	
GGF-I 16	K/R L G E M W	(SEQ ID NO: 16)	HMG-1
	Protease V8 peptides		
GGF-I 17	E T Q P D P G Q I L K K V P M V I G A Y T	(SEQ ID NO: 165)	
GGF-I 18	E Y K C L K F K W F K K A T V M	(SEQ ID NO: 17)	
GGF-I 19	E A K Y F S K X D A	(SEQ ID NO: 18)	LH-alpha
GGF-I 20	E X K F Y V P	(SEQ ID NO: 19)	
GGF-I 21	E L S F A S V R L P G C P P G V D P M V S F P V A L	(SEQ ID NO: 20)	LH-beta

6/72

# FIG. 9

## A

GGF-I 01	F K G D A H T E	(SEQ ID NO: 1)
GGF-I 02	A S L A D E Y E Y M X K	(SEQ ID NO: 22)
GGF-I 03	T E T S S S G L X L K	(SEQ ID NO: 23)
GGF-I 07	A S L A D E Y E Y M R K	(SEQ ID NO: 24)
GGF-I 11	A G Y F A E X A R	(SEQ ID NO: 25)
GGF-I 13	T T E M A S E Q G A	(SEQ ID NO: 26)
GGF-I 14	A K E A L A A L K	(SEQ ID NO: 27)
GGF-I 15	F V L Q A K K	(SEQ ID NO: 28)
GGF-I 17	E T Q P D P G Q I L K K V P M V I G A Y T	(SEQ ID NO: 29)
GGF-I 18	E Y K C L K F K W F K K A T V M	(SEQ ID NO: 17)

## B

GGF-I 20	E X K F Y V P	(SEQ ID NO: 19)
GGF-I 12	K L E F L X A K	(SEQ ID NO: 32)

**FIG. 10**

Trypsin peptides		
GGF-II 01	K/R V H Q V W A A K *	(SEQ ID NO: 42)
GGF-II 02	K/R Y I F F M E P E A X S S G	(SEQ ID NO: 43)
GGF-II 03	K/R L G A W G P P A F P V X Y	(SEQ ID NO: 44)
GGF-II 04	K/R W F V V I E G K *	(SEQ ID NO: 45)
GGF-II 05	K/R A L A A A G Y D V E K *	Histone H1 (SEQ ID NO: 160)
GGF-II 06	K/R L V L R *	(SEQ ID NO: 161)
GGF-II 07	K/R X X Y P G Q I T S N	Trypsin (SEQ ID NO: 162)
GGF-II 08	K/R A S P V S V G S V Q E L V Q R *	(SEQ ID NO: 46)
GGF-II 09	K/R V C L L T V A A P P T	(SEQ ID NO: 47)
GGF-II 10	K/R D L L L X V	(SEQ ID NO: 50)
Lysyl Endopeptidase-C peptides		
GGF-II 11	K V H Q V W A A K *	(SEQ ID NO: 48)
GGF-II 12	K A S L A D S G E Y M X K*	(SEQ ID NO: 49)

2 200 6070

## FIG. 11

A

GGF-II 01	V H Q V W A A K	(SEQ ID NO: 42)
GGF-II 02	Y I F F M E P E A X S S G	(SEQ ID NO: 43)
GGF-II 03	L G A W G P P A F P V X Y	(SEQ ID NO: 44)
GGF-II 04	W F V V I E G K	(SEQ ID NO: 45)
GGF-II 08	A S P V S V G S V Q E L V Q R	(SEQ ID NO: 46)
GGF-II 09	V C L L T V A A P P T	(SEQ ID NO: 47)
GGF-II 11	K V H Q V W A A K	(SEQ ID NO: 48)
GGF-II 12	K A S L A D S G E Y M X K	(SEQ ID NO: 49)

B


Novel Factor II Peptides - others

GGF-II 10	D L L L X V	(SEQ ID NO: 50)
-----------	-------------	-----------------

9/72


**FIG. 12**

**Comparison of BrdU-ELISA and [<sup>125</sup>I]UdR Counting Method for the DNA Synthesis Assay in Schwann Cell Cultures**






**FIG. 13A**  
**Comparison of Br-UdR Immunoreactivity**  
**and Br-UdR Labelled Cell Number**



**FIG. 13B**  
**Comparison of Br-UdR Immunoreactivity**  
**and Br-UdR Labelled Cell Number**




2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50

12/72


**FIG. 14**

**Mitogenic Response of Rat Sciatic Nerve Schwann cell to GGFs**




**FIG. 15**


**DNA Synthesis in Rat Sciatic Nerve Schwann Cells and 3T3 Fibroblasts in the presence of GGFs**




**FIG. 16**  
**Mitogenic Response of**  
**BHK 21 C13 Cells to FCS and GGFs**



**FIG. 17**  
**Survival and Proliferation of BHK21 C13 Cell**  
**Microcultures After 48 Hours in Presence of GGFs**





**FIG. 18**  
Mitogenic Response  
of C6 Cells to FCS



2529  
299  
2

FIG. 19

**Mitogenic response of  
C6 Cells to aFGF & GGFs**




15/78

## FIG. 20

### Degenerate Oligonucleotide Probes for Factor I & Factor II

Oligo	Sequence	Peptide	
535	TTYAARGGNGAYGCNCAYAC!	GGFI-1	(SEQ ID NO: 51)
536	CATRTAYTCRTAYTCRTCNGC!	GGFI-2	(SEQ ID NO: 52)
537	TGYTCNGANGCCATYTCNGT!	GGFI-13	(SEQ ID NO: 53)
538	TGYTCRCTNGCCATYTCNGT!	GGFI-13	(SEQ ID NO: 54)
539	CCDATNACCATNNGNACYTT!	GGFI-17	(SEQ ID NO: 55)
540	GCNGCCCANACYTGRTGNAC!	GGFII-1	(SEQ ID NO: 56)
541	GCYTCNGGYTCCATRAARAA!	GGFII-2	(SEQ ID NO: 57)
542	CCYTCDATNACNACRAACCA!	GGFII-4	(SEQ ID NO: 58)
543	TCNGCRAARTANCCNGC!	GGFI-11	(SEQ ID NO: 59)
544	GCNGCNAGNGCYTCYTTNGC!	GGFI-14	(SEQ ID NO: 60)
545	GCNGCYAANGCYTCYTTNGC!	GGFI-14	(SEQ ID NO: 61)
546	TTYTTNGCYTGNAGNACRAA!	GGFI-15	(SEQ ID NO: 62)
551	TTYTTNGCYTGAAANACRAA!	GGFI-15	(SEQ ID NO: 63)
568	TGNACNAGYTCYTGNAAC!	GGFII-8	(SEQ ID NO: 64)
569	TGNACYAAAYTCYTGNAAC!	GGFII-8	(SEQ ID NO: 65)
609	CATRTAYTCNCCNGARTCNGC!	GGFII-12	(SEQ ID NO: 66)
610	CATRTAYTCNCCRCRTRCNGC!	GGFII-12	(SEQ ID NO: 67)
649	NGARTCNGCYAANGANGCYTT!	GGFII-12	(SEQ ID NO: 68)
650	NGARTCNGCNAGNGANGCYTT!	GGFII-12	(SEQ ID NO: 69)
651	RCTRTRCNGCYAANGANGCYTT!	GGFII-12	(SEQ ID NO: 70)
652	RCTRTRCNGCNAGNGANGCYTT!	GGFII-12	(SEQ ID NO: 71)
653	NGARTCNGCYAARCTNGCYTT!	GGFII-12	(SEQ ID NO: 72)
654	NGARTCNGCNAGRCTNGCYTT!	GGFII-12	(SEQ ID NO: 73)
655	RCTRTRCNGCYAARCTNGCYTT!	GGFII-12	(SEQ ID NO: 74)
656	RCTRTRCNGCNAGRCTNGCYTT!	GGFII-12	(SEQ ID NO: 75)
659	ACNACNGARATGGCTCNNGA!	GGFI-13	(SEQ ID NO: 76)
660	ACNACNGARATGGCAGYNGA!	GGFI-13	(SEQ ID NO: 77)
661	CAYCARGTNTGGGCNGCNA!	GGFII-1	(SEQ ID NO: 78)
662	TTYGTNGTNATHGARGGNAA!	GGFII-4	(SEQ ID NO: 79)
663	AARGGNGAYGCNCAYACNGA!	GGFI-1	(SEQ ID NO: 80)
664	GARGCNYTNGCNGCNYTNAA!	GGDI-14	(SEQ ID NO: 81)
665	GTNGGNTCNGTNCARGARYT!	GGFII-8	(SEQ ID NO: 82)
666	GTNGGNGYGTNCARGARYT!	GGFII-8	(SEQ ID NO: 83)
694	NACYTTYTTNARHATYTGNC!	GGFI-17	(SEQ ID NO: 84)



2 296 6078

# FIG. 21

## Putative Bovine Factor II Gene Sequences

SEQ ID NO: 85:

TCTAA AAC TAC AGA GAC TGT ATT TTC ATG ATC ATC ATA GTT CTG TGA AAT ATA	53
Asn Tyr Arg Asp Cys Ile Phe Met Ile Ile Ile Val Leu Xaa Asn Ile	
CTT AAA CCG CTT TGG TCC TGA TCT TGT AGG AAG TCA GAA CTT CGC ATT	101
Leu Lys Pro Leu Trp Ser Xaa Ser Cys Arg Lys Ser Glu Leu Arg Ile	
AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA TAT ATG TGC AAA GTG ATC	149
Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu Ser Met Cys Lys Val Ile	
AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG	197
Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Arg Ile Val Glu	
TCA AAC GGT AAG AGA TCC CTA CTG CGT GCT ATT TCT CAG TCT CTA AGA	245
Ser Asn Gly Lys Arg Cys Leu Leu Arg Ala Ile Ser Gln Ser Leu Arg	
GGA GTG ATC AAG GTA TGT GGT CAC ACT TGA ATC ACG CAG GTG TGT GAA	293
Gly Val Ile Lys Val Cys Gly His Thr Xaa Ile Thr Gln Val Cys Glu	
ATC TCA TTG TGA ACA AAT AAA AAT CAT GAA AGG AAA ACT CTA TGT TTG	341
Ile Ser Cys Xaa Thr Asn Lys Asn His Glu Arg Lys Thr Leu Cys Leu	
AAA TAT CTT ATG GGT CCT CCT GTA AAG CTC TTC ACT CCA TAA GGT GAA	389
Lys Tyr Leu Met Gly Pro Pro Val Lys Leu Phe Thr Pro Xaa Gly Glu	
ATA GAC CTG AAA TAT ATA TAG ATT ATT T	417
Ile Asp Leu Lys Tyr Ile Xaa Ile Ile	

177A

**FIG. 22 1/2**  
**PCR Primers for Factor I & Factor II**

**Degenerate PCR Primers**

Oligo Sequence	Peptide	
657 CCGAATTCTGCAGGARACNCARCCNGAYCCNGG!	GGFI-17	(SEQ ID NO: 86 )
658 AAGGATCCTGCAGNGTRTANGCNCCHATNACCATNGG!	GGFI-17	(SEQ ID NO: 87 )
667 CCGAATTCTGCAGGCNGAYTCNGGNGARTAYATG!	GGFII-12	(SEQ ID NO: 88 )
668 CCGAATTCTGCAGGCNGAYATYGGNGARTAYAT!	GGFII-12	(SEQ ID NO: 89 )
669 AAGGATCCTGCAGNNNCATRTAYTCNCCNGARTC!	GGFII-12	(SEQ ID NO: 90 )
670 AAGGATCCTGCAGNNNCATRTAYTCNCCRTRRTC!	GGFII-12	(SEQ ID NO: 91 )
671 CCGAATTCTGCAGCAYCARGTNTGGGCNGCNA!	GGFII-1	(SEQ ID NO: 92 )
672 CCGAATTCTGCAGATRTTYTTYATGGARCCNGARG!	GGFII-2	(SEQ ID NO: 93 )
673 CCGAATTCTGCAGGGGGNCCNCCNGCNTTYCCNGT!	GGFII-3	(SEQ ID NO: 94 )
674 CCGAATTCTGCAGTGGTTYGTNGTNATHGARGG!	GGFII-4	(SEQ ID NO: 95 )
677 AAGGATCCTGCAGYTTNGCNGCCCANACYTGRTG!	GGFII-1	(SEQ ID NO: 96 )
678 AAGGATCCTGCAGGCYTCNGGYTCCATRAARAA!	GGFII-2	(SEQ ID NO: 97 )
679 AAGGATCCTGCAGACNGGRAANGCNGGNGGNCC!	GGFII-3	(SEQ ID NO: 98 )
680 AAGGATCCTGCAGYTTNCCYTCDATNACNACRAAC!	GGFII-4	(SEQ ID NO: 99 )
681 CATRTAYTCRTAYTCTCNGCAAGGATCCTGCAG!	GGFI-2	(SEQ ID NO: 100 )
682 CCGAATTCTGCAGAARGGNGAYGCNCAYACNGA!	GGFI-1	(SEQ ID NO: 101 )
683 GCNGCYAANGCYRCYTTNGCAAGGATCCTGCAG!	GGFI-14	(SEQ ID NO: 102 )
684 GCNGCNAGNGCYTCYTTNGCAAGGATCCTGCAG!	GGFI-14	(SEQ ID NO: 103 )
685 TCNGCRAARTANCCNGCAAGGATCCTGCAG!	GGFII-1	(SEQ ID NO: 104 )

18/72




**FIG. 22 2/2**  
**PCR Primers for Factor I & Factor II**

**Unique PCR Primers for Factor II**

Oligo Sequence	Comment	
711 CATCGATCTGCAGGCTGATTCTGGAGAATATATGTGCA!	3' RACE	(SEQ ID NO: 105)
712 AAGGATCCTGCAGCCACATCTCGAGTCGACATCGATT!	3' RACE	(SEQ ID NO: 106)
713 CCGAATTCTGCAGTGATCAGCAAAGTAGGAAATGACA!	3' RACE	(SEQ ID NO: 107)
721 CATCGATCTGCAGCCTAGTTTGGCTGATCACTTTGCAC!	5' RACE	(SEQ ID NO: 108)
722 AAGGATCCTGCAGTATATTCTCCAGAATCAGCCAGTG!	5' RACE; ANCHORED	(SEQ ID NO: 109)
725 AAGGATCCTGCAGGCACGCAGTAGGCATCTCTTA!	EXON A	(SEQ ID NO: 110)
726 CCGAATTCTGCAGCAGAACTTCGCATTAGCAAAGC!	EXON A	(SEQ ID NO: 111)
771 CATCCCGGGATGAAGAGTCAGGAGTCTGTGGCA!	EXONS B+A	(SEQ ID NO: 112)
772 ATACCCGGGCTGCAGACAATGAGATTTACACACACCTGCG!		(SEQ ID NO: 113)
773 AAGGATCCTGCAGTTTGGAACTGCCACAGACTCCT!	ANCHORED	(SEQ ID NO: 114)
776 ATACCCGGGCTGCAGATGAGATTTACACACACCTGCGTGA!	EXONS B+A	(SEQ ID NO: 115)


19/72

**FIG. 23**  
Summary of Contiguous GGF-II  
cDNA Structures & Sequences




2 298 6078

FIG. 24



21/78

**FIG. 25**  
**Alternative Gene Products of Putative Bovine GGF-II**



22/73

**FIG. 26****GGF-II Peptides Identified in Deduced Amino Acid Sequences of Putative Bovine GGF-II Proteins**

Peptide	Pos.	Sequence match	ID Sequences
II-1	1:	VHQVWAAK HQVWAAK AAGLK	(SEQ ID NO: 116)
II-10	14:	DLLLXV GGLKK dsl1tv RLGAW	(SEQ ID NO: 117)
II-03	21:	LGAWGPPAFPVXY LLTVR lgawghpafpsc RLKED	(SEQ ID NO: 118 ) (SEQ ID NO: 119 )
II-02	41:	YIFFMEPEAXSSG KEDSR YIFFMEPEANSSG GPGRL	(SEQ ID NO: 120 ) (SEQ ID NO: 121 )
II-6	103:	LVLRL VAGSK LVLRL CETSS	(SEQ ID NO: 122 )
I-18	112:	EYKCLKFKWFKKATVM CETSS eysslkfkfwkngsel SRKNK	(SEQ ID NO: 123 ) (SEQ ID NO: 124 )
II-12	151:	KASLADSGEYMXK ELRIS KASLADSGEYMCK VISKL	(SEQ ID NO: 125 ) (SEQ ID NO: 126 )
I-07	152:	ASLADEYEYMRK LRISK asladsgeymck VISKL	(SEQ ID NO: 127 ) (SEQ ID NO: 128 )

**FIG. 27** 1/5

SEQ ID NO: 129 :

CCTGCAG	CAT	CAA	GTG	TGG	GCG	GCG	AAA	GCC	GGG	GGC	TTG	AAG	AAG	GAC	TCG	CTG	55
	His	Gln	Val	Trp	Ala	Ala	Lys	Ala	Gly	Gly	Leu	Lys	Lys	Asp	Ser	Leu	
CTC	ACC	GTG	CGC	CTG	GGC	GCC	TGG	GGC	CAC	CCC	GCC	TTC	CCC	TCC	TGC		103
Leu	Thr	Val	Arg	Leu	Gly	Ala	Trp	Gly	His	Pro	Ala	Phe	Pro	Ser	Cys		
GGG	CGC	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	ATG	GAG	CCC	GAG		151
Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	Met	Glu	Pro	Glu		
GCC	AAC	AGC	AGC	GGC	GGG	CCC	GGC	CGC	CTT	CCG	AGC	CTC	CTT	CCC	CCC		199
Ala	<u>Asn</u>	<u>Ser</u>	<u>Ser</u>	Gly	Gly	Pro	Gly	Arg	Leu	Pro	Ser	Leu	Leu	Pro	Pro		
TCT	CGA	GAC	GGG	CCG	GAA	CCT	CAA	GAA	GGA	GGT	CAG	CCG	GGT	GCT	GTG		247
Ser	Arg	Asp	Gly	Pro	Glu	Pro	Gln	Glu	Gly	Gly	Gln	Pro	Gly	Ala	Val		
CAA	CGG	TGC	GCC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG		295
Gln	Arg	Cys	Ala	Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu		
TCT	GTG	GCA	GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA		343
Ser	Val	Ala	Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu		
TAC	TCC	TCT	CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC		391
Tyr	Ser	Ser	Leu	Lys	Phe	Lys	Trp	Phe	Lys	<u>Asn</u>	<u>Gly</u>	<u>Ser</u>	Glu	Leu	Ser		
CGA	AAG	AAC	AAA	CCA	GAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GGG	AAG		439
Arg	Lys	Asn	Lys	Pro	Glu	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Gly	Lys		
TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	GAA	TAT		487
Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly	Glu	Tyr		
ATG	TGC	AAA	GTG	ATC	AGC	AAA	CTA	GGA	AAT	GAC	AGT	GCC	TCT	GCC	AAC		535
Met	Cys	Lys	Val	Ile	Ser	Lys	Leu	Gly	<u>Asn</u>	<u>Asp</u>	<u>Ser</u>	Ala	Ser	Ala	<u>Asn</u>		
ATC	ACC	ATT	GTG	GAG	TCA	AAC	GGT	AAG	AGA	TGC	CTA	CTG	CGT	GCT	ATT		583
<u>Ile</u>	<u>Thr</u>	Ile	Val	Glu	Ser	Asn	Gly	Lys	Arg	Cys	Leu	Leu	Arg	Ala	Ile		
TCT	CAG	TCT	CTA	AGA	GGA	GTG	ATC	AAG	GTA	TGT	GGT	CAC	ACT				625
Ser	Gln	Ser	Leu	Arg	Gly	Val	Ile	Lys	Val	Cys	Gly	His	Thr				
TGAATCACGC	AGGTGTGTGA	AATCTCATTG	TGAACAAATA	AAAATCATGA	AAGGAAAAAA												685
AAAAA	AAAAA	AATCGATGTC	GACTCGAGAT	GTGGCTGCAG	GTCGACTCTA	GAGGATCCC											744

24/73

2 298 8278

## FIG. 27 2/5

### Nucleotide Sequences & Deduced Amino Acid Sequences of GGF2BPP2

SEQ ID NO: 130:

CCTGCAG	CAT	CAA	GTG	TGG	GCG	GCG	AAA	GCC	GGG	GGC	TTG	AAG	AAG	GAC	TCG	CTG	55
	His	Gln	Val	Trp	Ala	Ala	Lys	Ala	Gly	Gly	Leu	Lys	Lys	Asp	Ser	Leu	
CTC	ACC	GTG	CGC	CTG	GGC	GCC	TGG	GGC	CAC	CCC	GCC	TTC	CCC	TCC	TGC		103
Leu	Thr	Val	Arg	Leu	Gly	Ala	Trp	Gly	His	Pro	Ala	Phe	Pro	Ser	Cys		
GGG	CGC	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	ATG	GAG	CCC	GAG		151
Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	Met	Glu	Pro	Glu		
GCC	AAC	AGC	AGC	GGC	GGG	CCC	GGC	CGC	CTT	CCG	AGC	CTC	CTT	CCC	CCC		199
Ala	<u>Lys</u>	<u>Ser</u>	<u>Ser</u>	Gly	Gly	Pro	Gly	Arg	Leu	Pro	Ser	Leu	Leu	Pro	Pro		
TCT	CGA	GAC	GGG	CCG	GAA	CCT	CAA	GAA	GGA	GGT	CAG	CCG	GGT	GCT	GTG		247
Ser	Arg	Asp	Gly	Pro	Glu	Pro	Gln	Glu	Gly	Gly	Gln	Pro	Gly	Ala	Val		
CAA	CGG	TGC	GCC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG		295
Gln	Arg	Cys	Ala	Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu		
TCT	GTG	GCA	GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA		343
Ser	Val	Ala	Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu		
TAC	TCC	TCT	CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC		391
Tyr	Ser	Ser	Leu	Lys	Phe	Lys	Trp	Phe	Lys	<u>Asn</u>	<u>Gly</u>	<u>Ser</u>	Glu	Leu	Ser		
CGA	AAG	AAC	AAA	CCA	GAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GGG	AAG		439
Arg	Lys	Asn	Lys	Gly	Gly	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Gly	Lys		
TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	GAA	TAT		487
Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly	Glu	Tyr		
ATG	TGC	AAA	GTG	ATC	AGC	AAA	CTA	GGA	AAT	GAC	AGT	GCC	TCT	GCC	AAC		535
Met	Cys	Lys	Val	Ile	Ser	Lys	Leu	Gly	<u>Asn</u>	<u>Asp</u>	<u>Ser</u>	Ala	Ser	Ala	<u>Asn</u>		

25/73

**FIG. 27 3/5**

**Nucleotide Sequences & Deduced Amino Acid Sequences of GG2BPP2**

ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG ACA	583
<u>Ile Thr</u> Ile Val Glu Ser <u>Asn Ala Thr</u> Ser Thr Ser Thr Ala Gly Thr	
AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	631
Ser His Leu Val Lys Ser Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	679
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	727
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AGT GCC CAA ATG AGT TTA CTG	775
Val Pro Met Lys Val Gln Thr Gln Glu Ser Ala Gln Met Ser Leu Leu	
GTG ATC GCT GCC AAA ACT ACG TAATGGCCAG CTTCTACAGT ACGTCCACTC	826
Val Ile Ala Ala Lys Thr Thr	
CCTTCTGTGTC TCTGCCTGAA TAGCGCATCT CAGTCGGTGC CGCTTCTTG TTGCCGCATC	886
TCCCCTCAGA TTCCTCCTAG AGCTAGATGC GTTTTACCAG GTCTAACATT GACTGCCTCT	946
GCCTGTGCGCA TGAGAACATT AACACAAGCG ATTGTATGAC TTCCTCTGTC CGTGACTAGT	1006
GGGCTCTGAG CTA CTCTCGTAG GTGCGTAAGG CTCCAGTGTT TCTGAAATTG ATCTTGAATT	1066
ACTGTGATAC GACATGATAG TCCCTCTCAC CCAGTGCAAT <u>GACAATAAAG</u> GCCTTGAAAA	1126
GTCAAAAAAA AAAAAAAAAA AAAAAATCGA TGTCGACTCG AGATGTGGCT GCAGGTCGAC	1186
TCTAGAG	1193



**FIG. 27 4/5**

**Nucleotide Sequences & Deduced Amino Acid Sequences of GGF2BPP3**

SEQ ID NO: 131:

CCTGCAG	CAT	CAA	GTG	TGG	GCG	GCG	AAA	GCC	GGG	GGC	TTG	AAG	AAG	GAC	TCG	CTG	55
	His	Gln	Val	Trp	Ala	Ala	Lys	Ala	Gly	Gly	Leu	Lys	Lys	Asp	Ser	Leu	
CTC	ACC	GTG	CGC	CTG	GGC	GCC	TGG	GGC	CAC	CCC	GCC	TTC	CCC	TCC	TGC		103
Leu	Thr	Val	Arg	Leu	Gly	Ala	Trp	Gly	His	Pro	Ala	Phe	Pro	Ser	Cys		
GGG	CGC	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	ATG	GAG	CCC	GAG		151
Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	Met	Glu	Pro	Glu		
GCC	AAC	AGC	AGC	GGC	GGG	CCC	GGC	CGC	CTT	CCG	AGC	CTC	CTT	CCC	CCC		199
Ala	<u>Asn</u>	<u>Ser</u>	<u>Ser</u>	Gly	Gly	Pro	Gly	Arg	Leu	Pro	Ser	Leu	Leu	Pro	Pro		
TCT	CGA	GAC	GGG	CCG	GAA	CCT	CAA	GAA	GGA	GGT	CAG	CCG	GGT	GCT	GTG		247
Ser	Arg	Asp	Gly	Pro	Glu	Pro	Gln	Glu	Gly	Gly	Gln	Pro	Gly	Ala	Val		
CAA	CGG	TGC	GCC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG		295
Gln	Arg	Cys	Ala	Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu		
TCT	GTG	GCA	GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA		343
Ser	Val	Ala	Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu		
TAC	TCC	TCT	CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC		391
Tyr	Ser	Ser	Leu	Lys	Phe	Lys	Trp	Phe	Lys	<u>Asn</u>	<u>Gly</u>	<u>Ser</u>	Glu	Leu	Ser		
CGA	AAG	AAC	AAA	CCA	GAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GGG	AAG		439
Arg	Lys	Asn	Lys	Pro	Glu	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Pro	Lys		
TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	GAA	TAT		487
Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly	Glu	Tyr		

2772


**FIG. 27 5/5**

**Nucleotide Sequences & Deduced Amino Acid Sequences of GGF2BPP3**


ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC AAC	535
Met Cys Lys Val Ile Ser Lys Leu Gly <u>Asn Asp Ser</u> Ala Ser Ala <u>Asn</u>	
ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG ACA	583
<u>Ile Arg Ile</u> Val Glu Ser <u>Asn Ala Thr</u> Ser Thr Ser Thr Ala Gly Thr	
AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	631
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	679
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser <u>Asn Pro Ser</u> Arg Tyr	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	727
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
GTA ATG GCC AGC TTC TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT	775
Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro	
GAA TAGCGCATCT CAGTCGGTGC CGCTTTCTTG TTGCCGCATC TCCCCTCAGA TTCCGCCTAG	838
Glu	
AGCTAGATGC GTTTTACCAG GTCTAACATT GACTGCCTCT GCCTGTCGCA TGAGAACATT	898
AACACAAGCG ATTGTATGAC TTCCTCTGTC CGTGACTAGT GGGCTCTGAG CTACTCGTAG	958
GTGCGTAAGG CTCCAGTGTT TCTGAAATTG ATCTTGAATT ACTGTGATAC GACATGATAG	1018
TCCCTCTCAC CCAGTGCAAT <u>GACAATAAAG</u> GCCTTGAAAA GTCAAAAAAA AAAAAAAAAA	1078
AAAAATCGAT GTCGACTCGA GATGTGGCTG	1108



2 298 6078



**FIG. 29**



30/78

**FIG. 30 1/16**

**Coding Segments  
of Glial Growth  
Factor/Heregulin  
Gene**

CODING SEGMENT F: (SEQ ID NO: 132)

AGTTTCCCCC	CCCAACTTGT	CGGAACTCTG	GGCTCGCGCG	CAGGGCAGGA	GCGGAGCGGC	60
GGCGGCTGCC	CAGGCGATGC	GAGCGCGGGC	CGGACGGTAA	TCGCCTCTCC	CTCCTCGGGC	120
TGCGAGCGCG	CCGGACCGAG	GCAGCGACAG	GAGCGGACCG	CGGCGGGAAC	CGAGGACTCC	180
CCAGCGGCGC	GCCAGCAGGA	GCCACCCCGC	GAGNCGTGCG	ACCGGGACGG	AGCGCCCGCC	240
AGTCCCAGGT	GGCCCGGACC	GCACGTTGCG	TCCCCGCGCT	CCCCGCCGGC	GACAGGAGAC	300
GCTCCCCCCC	ACGCCGCGCG	CGCCTCGGCC	CGGTGCGCTGG	CCCGCCTCCA	CTCCGGGGAC	360
	CGCGAG	CGCCTCAGCG	CGGCCGCTCG	CTCTC..CCC	CTCGAGGGAC	
AAACTTTTCC	CGAAGCCGAT	CCCAGCCCTC	GGACCCAAAC	TTGTGCGCGG	TCGCCTTCGC	420
AAACTTTTCC	CAAACCCGAT	CCGAGCCCTT	GGACCAA...	.....C	TCGCCTGCGC	
				Met Ser Glu Arg Arg		
CGGGAGCCGT	CCGCGCAGAG	CGTGCACTTC	TCGGGCGAG	ATG TCG GAG CGC AGA		474
CGAGAGCCGT	CCGCGTAGAG	CGTC.CGTC	TCCGGCGAG	ATG TCC GAG CGC AAA		
				K		
Glu Gly Lys Gly Lys Gly Lys Gly Gly Lys Lys Asp Arg Gly Ser Gly						
GAA GGC AAA GGC AAG GGG AAG GGC GGC AAG AAG GAC CGA GGC TCC GGG						522
GAA GGC AGA GGC AAA GGG AAG GGC AAG AAG AAG GAG CGA GGC TCC GGC						
	R			K	E	
Lys Lys Pro Val Pro Ala Ala Gly Gly Pro Ser Pro Ala						
AAG AAG CCC GTG CCC GCG GCT GGC GGC CCG AGC CCA G						559
AAG AAG CCG GAG TCC GCG GCG GGC AGC CAG AGC CCA G						
	E	S				

3172


2 299 6070

# FIG. 30 2/16

CODING SEGMENT E: (SEQ ID NO: 133)

CC CAT CAN GTG TGG GCG GCG AAA GCC GGG GGC TTG AAG AAG GAC TCG	47
His Gln Val Trp Ala Ala Lys Ala Gly Gly Leu Lys Lys Asp Ser	
CTG CTC ACC GTG CGC CTG GGC GCC TGG GGC CAC CCC GCC TTC CCC TCC	95
Leu Leu Thr Val Arg Leu Gly Ala Trp Gly His Pro Ala Phe Pro Ser	
TGC GGG CGC CTC AAG GAG GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC	143
Cys Gly Arg Leu Lys Glu Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro	
GAG GCC AAC AGC AGC GGC GGG CCC GGC CGC CTT CCG AGC CTC CTT CCC	191
Glu Ala Asn Ser Ser Gly Gly Pro Gly Arg Leu Pro Ser Leu Leu Pro	
CCC TCT CGA GAC GGG CCG GAA CCT CAA GAA GGA GGT CAG CCG GGT GCT	239
Pro Ser Arg Asp Gly Pro Glu Pro Gln Glu Gly Gly Gln Pro Gly Ala	
GTG CAA CGG TGC G	252
Val Gln Arg Cys	

3272



# FIG. 30 3/16

CODING SEGMENT B: (SEQ ID NO: 134)

Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu	Ser	Val	Ala		
CC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG	TCT	GTG	GCA	47
CC	TTG	CCT	CCC	CGA	TTG	AAA	GAG	ATG	AAA	AGC	CAG	GAA	TCG	GCT	GCA	
				Q											A	
Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu	Tyr	Ser	Ser	
GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA	TAC	TCC	TCT	95
GGT	TCC	AAA	CTA	GTC	CTT	CGG	TGT	GAA	ACC	AGT	TCT	GAA	TAC	TCC	TCT	
Leu	Lys	Phe	Lys	Trp	Phe	Lys	Asn	Gly	Ser	Glu	Leu	Ser	Arg	Lys	Asn	
CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC	CGA	AAG	AAC	143
CTC	AGA	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AAT	GAA	TTG	AAT	CGA	AAA	AAC	
	R								N			N				
Lys	Pro	Gln	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Gly					
AAA	CCA	CAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GG					178
AAA	CCA	CAA	AAT	ATC	AAG	ATA	CAA	AAA	AAG	CCA	GG					
																K

33/72

2 298 6078

### FIG. 30 4/16

CODING SEGMENT A: (SEQ ID NO: 135)

Lys	Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly		
G	AAG	TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	46
G	AAG	TCA	GAA	CTT	CGC	ATT	AAC	AAA	GCA	TCA	CTG	GCT	GAT	TCT	GGA	
							N									
Glu	Tyr	Met	Cys	Lys	Val	Ile	Ser	Lys	Leu	Gly	Asn	Asp	Ser	Ala	Ser	
GAA	TAT	ATG	TGC	AAA	GTG	ATC	AGC	AAA	CTA	GGA	AAT	GAC	AGT	GCC	TCT	94
GAG	TAT	ATG	TGC	AAA	GTG	ATC	AGC	AAA	TTA	GGA	AAT	GAC	AGT	GCC	TCT	
Ala	Asn	Ile	Thr	Ile	Val	Glu	Ser	Asn	Ala							
GCC	AAC	ATC	ACC	ATT	GTG	GAG	TCA	AAC	G							122
GCC	AAT	ATC	ACC	ATC	GTG	GAA	TCA	AAC	G							

34/72



2 298 6078

### FIG. 30 5/16

CODING SEGMENT A': (SEQ ID NO: 136)

TCTAAAATA CAGAGACTGT ATTTTCATGA TCATCATAGT TCTGTGAAAT ATACTTAAAC	60
CGCTTTGGTC CTGATCTTGT AGG AAG TCA GAA CTT CGC ATT AGC AAA GCG	110
Lys Ser Glu Leu Arg Ile Ser Lys Ala	
TCA CTG GCT GAT TCT GGA GAA TAT ATG TGC AAA GTG ATC AGC AAA CTA	158
Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys Leu	
GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG TCA AAC GGT	206
Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn Gly	
AAG AGA TGC CTA CTG CGT GCT ATT TCT CAG TCT CTA AGA GGA GTG ATC	254
Lys Arg Cys Leu Leu Arg Ala Ile Ser Gln Ser Leu Arg Gly Val Ile	
AAG GTA TGT GGT CAC ACT TGAATCACGC AGGTGTGTGA AATCTCATTG	302
Lys Val Cys Gly His Thr	
TGAACAAATA AAAATCATGA AAGGAAAACCT CTATGTTTGA AATATCTTAT GGGTCCTCCT	362
GTAAAGCTCT TCACTCCATA AGGTGAAATA GACCTGAAAT ATATATAGAT TATTT	417

35/72

2 298 6078

# FIG. 30 6/16

CODING SEGMENT G: (SEQ ID NO: 137)

Glu	Ile	Thr	Thr	Gly	Met	Pro	Ala	Ser	Thr	Glu	Thr	Ala	Tyr	Val	Ser	
AG	ATC	ACC	ACT	GGC	ATG	CCA	GCC	TCA	ACT	GAG	ACA	GCG	TAT	GTG	TCT	47
AG	ATC	ATC	ACT	GGT	ATG	CCA	GCC	TCA	ACT	GAA	GGA	GCA	TAT	GTG	TCT	
		I								G						

Ser	Glu	Ser	Pro	Ile	Arg	Ile	Ser	Val	Ser	Thr	Glu	Gly	Thr	Asn	Thr	
TCA	GAG	TCT	CCC	ATT	AGA	ATA	TCA	GTA	TCA	ACA	GAA	GGA	ACA	AAT	ACT	95
TCA	GAG	TCT	CCC	ATT	AGA	ATA	TCA	GTA	TCC	ACA	GAA	GGA	GCA	AAT	ACT	
													A			

Ser	Ser	Ser														
TCT	TCA	T														102
TCT	TCA	T														

36/72

2 298 8078

# FIG. 30 7/16

CODING SEGMENT C: (SEQ ID NO: 156)

Thr	Ser	Thr	Ser	Thr	Ala	Gly	Thr	Ser	His	Leu	Val	Lys	Cys	Ala		
CC	ACA	TCC	ACA	TCT	ACA	GCT	GGG	ACA	AGC	CAT	CTT	GTC	AAG	TGT	GCA	47
CT	ACA	TCT	ACA	TCC	ACC	ACT	GGG	ACA	AGC	CAT	CTT	GTA	AAA	TGT	GCG	
						T										
Glu	Lys	Glu	Lys	Thr	Phe	Cys	Val	Asn	Gly	Gly	Glu	Cys	Phe	Met	Val	
GAG	AAG	GAG	AAA	ACT	TTC	TGT	GTG	AAT	GGA	GGC	GAG	TGC	TTC	ATG	GTG	95
GAG	AAG	GAG	AAA	ACT	TTC	TGT	GTG	AAT	GGA	GGG	GAG	TGC	TTC	ATG	GTG	
Lys	Asp	Leu	Ser	Asn	Pro	Ser	Arg	Tyr	Leu	Cys						
AAA	GAC	CTT	TCA	AAT	CCC	TCA	AGA	TAC	TTG	TGC						128
AAA	GAC	CTT	TCA	AAC	CCC	TCG	AGA	TAC	TTG	TGC						

3772

2 200 6070

# FIG. 30 8/16

CODING SEGMENT C/D: (SEQ ID NO: 138)


Lys	Cys	Gln	Pro	Gly	Phe	Thr	Gly	Ala	Arg	Cys	Thr	Glu	Asn	Val	Pro	
AAG	TGC	CAA	CCT	GGA	TTC	ACT	GGA	GCG	AGA	TGT	ACT	GAG	AAT	GTG	CCC	
AAG	TGC	CAA	CCT	GGA	TTC	ACT	GGA	GCA	AGA	TGT	ACT	GAG	AAT	GTG	CCC	

48

Met	Lys	Val	Gln	Thr	Gln	Glu	
ATG	AAA	GTC	CAA	ACC	CAA	GAA	
ATG	AAA	GTC	CAA	AAC	CAA	GAA	
				N			

69

38/72



**FIG.30** 9/16 CODING SEGMENT C/D' (SEQ ID NO: 139)

Lys	Cys	Pro	Asn	Glu	Phe	Thr	Gly	Asp	Arg	Cys	Gln	Asn	Tyr	Val	Met	
AAG	TGC	CCA	AAT	GAG	TTT	ACT	GGT	GAT	CGC	TGC	CAA	AAC	TAC	GTA	ATG	48
AAG	TGC	CCA	AAT	GAG	TTT	ACT	GGT	GAT	CGC	TGC	CAA	AAC	TAC	GTA	ATG	

Ala	Ser	Phe	Tyr													
GCC	AGC	TTC	TAC													60
GCC	AGC	TTC	TAC													

CODING SEGMENT D: (SEQ ID NO: 140)

Ser	Thr	Ser	Thr	Pro	Phe	Leu	Ser	Leu	Pro	Glu	*					
AGT	ACG	TCC	ACT	CCC	TTT	CTG	TCT	CTG	CCT	GAA	TAG					36
AGT	ACG	TCC	ACT	CCC	TTT	CTG	TCT	CTG	CCT	GAA	TAG					

CODING SEGMENT D': (SEQ ID NO: 141)

Lys	His	Leu	Gly	Ile	Glu	Phe	Met	Glu								
AAG	CAT	CTT	GGG	ATT	GAA	TTT	ATG	GAG								27

2 298 8878

FIG. 30

10/16

CODING SEGMENT H: (SEQ ID NO: 142)

Lys	Ala	Glu	Glu	Leu	Tyr	Gln	Lys	Arg	Val	Leu	Thr	Ile	Thr	Gly	Ile		48
AAA	GCG	GAG	GAG	CTC	TAC	CAG	AAG	AGA	GTG	CTC	ACC	ATT	ACC	GGC	ATT		
AAG	GCG	GAG	GAG	CTG	TAC	CAG	AAG	AGA	GTG	CTG	ACC	ATA	ACC	GGC	ATC		
Cys	Ile	Ala	Leu	Leu	Val	Val	Gly	Ile	Met	Cys	Val	Val	Val	Tyr	Cys		96
TGC	ATC	GCG	CTG	CTC	GTG	GTT	GGC	ATC	ATG	TGT	GTG	GTG	GTC	TAC	TGC		
TGC	ATC	GCC	CTC	CTT	GTG	GTC	GGC	ATC	ATG	TGT	GTG	GTG	GCC	TAC	TGC		
												A					
Lys	Thr	Lys	Lys	Gln	Arg	Lys	Lys	Leu	His	Asp	Arg	Leu	Arg	Gln	Ser		144
AAA	ACC	AAG	AAA	CAA	CGG	AAA	AAG	CTT	CAT	GAC	CGG	CTT	CGG	CAG	AGC		
AAA	ACC	AAG	AAA	CAG	CGG	AAA	AAG	CTG	CAT	GAC	CGT	CTT	CGG	CAG	AGC		
Leu	Ser	Ser	Glu	Arg	Asn	Thr	Met	Met	Asn	Val	Ala	Asn	Gly	Pro	His		192
CTT	CGG	TCT	GAA	AGA	AAC	ACC	ATG	ATG	AAC	GTA	GCC	AAC	GGG	CCC	CAC		
CTT	CGG	TCT	GAA	CGA	AAC	AAT	ATG	ATG	AAC	ATT	GCC	AAT	GGG	CCT	CAC		
						N				I							
His	Pro	Asn	Pro	Pro	Pro	Glu	Asn	Val	Gln	Leu	Val	Asn	Gln	Tyr	Val		240
CAC	CCC	AAT	CCG	CCC	CCC	GAG	AAC	GTG	CAG	CTG	GTG	AAT	CAA	TAC	GTA		
CAT	CCT	AAC	CCA	CCC	CCC	GAG	AAT	GTC	CAG	CTG	GTG	AAT	CAA	TAC	GTA		
Ser	Lys	Asn	Val	Ile	Ser	Ser	Glu	His	Ile	Val	Glu	Arg	Glu	Ala	Glu		288
TCT	AAA	AAT	GTC	ATC	TCT	AGC	GAG	CAT	ATT	GTT	GAG	AGA	GAG	GCG	GAG		
TCT	AAA	AAC	GTC	ATC	TCC	AGT	GAG	CAT	ATT	GTT	GAG	AGA	GAA	GCA	GAG		

40/72

2 298 6078

FIG. 30 11/16

Ser	Ser	Phe	Ser	Thr	Ser	His	Tyr	Thr	Ser	Thr	Ala	His	His	Ser	Thr	
AGC	TCT	TTT	TCC	ACC	AGT	CAC	TAC	ACT	TCG	ACA	GCT	CAT	CAT	TCC	ACT	336
ACA	TCC	TTT	TCC	ACC	AGT	CAC	TAT	ACT	TCC	ACA	GCC	CAT	CAC	TCC	ACT	
T																
Thr	Val	Thr	Gln	Thr	Pro	Ser	His	Ser	Trp	Ser	Asn	Gly	His	Thr	Glu	
ACT	GTC	ACT	CAG	ACT	CCC	AGT	CAC	AGC	TGG	AGC	AAT	GGA	CAC	ACT	GAA	384
ACT	GTC	ACC	CAG	ACT	CCT	AGC	CAC	AGC	TGG	AGC	AAC	GGA	CAC	ACT	GAA	
Ser	Ile	Ile	Ser	Glu	Ser	His	Ser	Val	Ile	Val	Met	Ser	Ser	Val	Glu	
AGC	ATC	ATT	TCG	GAA	AGC	CAC	TCT	GTC	ATC	GTG	ATG	TCA	TCC	GTA	GAA	432
AGC	ATC	CTT	TCC	GAA	AGC	CAC	TCT	GTA	ATC	GTG	ATG	TCA	TCC	GTA	GAA	
		L														
Asn	Ser	Arg	His	Ser	Ser	Pro	Thr	Gly	Gly	Pro	Arg	Gly	Arg	Leu	Asn	
AAC	AGT	AGG	CAC	AGC	AGC	CCG	ACT	GGG	GGC	CCG	AGA	GGA	CGT	CTC	AAT	480
AAC	AGT	AGG	CAC	AGC	AGC	CCA	ACT	GGG	GGC	CCA	AGA	GGA	CGT	CTT	AAT	
Gly	Leu	Gly	Gly	Pro	Arg	Glu	Cys	Asn	Ser	Phe	Leu	Arg	His	Ala	Arg	
GGC	TTG	GGA	GGC	CCT	CGT	GAA	TGT	AAC	AGC	TTC	CTC	AGG	CAT	GCC	AGA	528
GGC	ACA	GGA	GGC	CCT	CGT	GAA	TGT	AAC	AGC	TTC	CTC	AGG	CAT	GCC	AGA	
		T														
Glu	Thr	Pro	Asp	Ser	Tyr	Arg	Asp	Ser	Pro	His	Ser	Glu	Arg			
GAA	ACC	CCT	GAC	TCC	TAC	CGA	GAC	TCT	CCT	CAT	AGT	GAA	AG			569
GAA	ACC	CCT	GAT	TCC	TAC	CGA	GAC	TCT	CCT	CAT	AGT	GAA	AG			

4172

2 293 6078

**FIG. 30 12/16**

CODING SEGMENT K: (SEQ ID NO: 157)

A	CAT	AAC	CTT	ATA	GCT	GAG	CTA	AGG	AGA	AAC	AAG	GCC	CAC	AGA	TCC	46
	His	Asn	Leu	Ile	Ala	Glu	Leu	Arg	Arg	Asn	Lys	Ala	His	Arg	Ser	
AAA	TGC	ATG	CAG	ATC	CAG	CTT	TCC	GCA	ACT	CAT	CTT	AGA	GCT	TCT	TCC	94
Lys	Cys	Met	Gln	Ile	Gln	Leu	Ser	Ala	Thr	His	Leu	Arg	Ala	Ser	Ser	
ATT	CCC	CAT	TGG	GCT	TCA	TTC	TCT	AAG	ACC	CCT	TGG	CCT	TTA	GGA	AG	141
Ile	Pro	His	Trp	Ala	Ser	Phe	Ser	Lys	Thr	Pro	Trp	Pro	Leu	Gly	Arg	

42/78



2 298 6078

CODING SEGMENT L: (SEQ ID NO: 143)

FIG. 30

13/16

Tyr Val Ser Ala Met Thr Thr Pro Ala Arg Met Ser Pro Val Asp  
 G TAT GTA TCA GCA ATG ACC ACC CCG GCT CGT ATG TCA CCT GTA GAT 46  
 | ||| || ||| || ||| ||| ||| ||| ||| ||| ||| ||| ||| |||  
 G TAT GTG TCA GCC ATG ACC ACC CCG GCT CGT ATG TCA CCT GTA GAT

Phe His Thr Pro Ser Ser Pro Lys Ser Pro Pro Ser Glu Met Ser Pro  
 TTC CAC ACG CCA AGC TCC CCC AAG TCA CCC CCT TCG GAA ATG TCC CCG 94  
 ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||  
 TTC CAC ACG CCA AGC TCC CCC AAA TCG CCC CCT TCG GAA ATG TCT CCA

Pro Val Ser Ser Thr Thr Val Ser Met Pro Ser Met Ala Val Ser Pro  
 CCC GTG TCC AGC ACG ACG GTC TCC ATG CCC TCC ATG GCG GTC AGT CCC 142  
 ||| ||| ||| ||| | | ||| ||| ||| ||| ||| ||| ||| ||| |||  
 CCC GTG TCC AGC ATG ACG GTG TCC ATG CCT TCC ATG GCG GTC AGC CCC  
 M

Phe Val Glu Glu Glu Arg Pro Leu Leu Leu Val Thr Pro Pro Arg Leu  
 TTC GTG GAA GAG GAG AGA CCC CTG CTC CTT GTG ACG CCA CCA CGG CTG 190  
 ||| || ||| || ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||  
 TTC ATG GAA GAA GAG AGA CCT CTA CTT CTC GTG ACA CCA CCA AGG CTG  
 N

Arg Glu Lys - Tyr Asp His His Ala Gln Gln Phe Asn Ser Phe His  
 CGG GAG AAG ... TAT GAC CAC CAC GCC CAG CAA TTC AAC TCG TTC CAC 238  
 ||| ||| ||| | | ||| ||| ||| | ||| ||| ||| ||| ||| ||| |||  
 CGG GAG AAG AAG TTT GAC CAT CAC CCT CAG CAG TTC AGC TCC TTC CAC  
 K F P

Cys Asn Pro Ala His Glu Ser Asn Ser Leu Pro Pro Ser Pro Leu Arg  
 TGC AAC CCC GCG CAT GAG AGC AAC AGC CTG CCC CCC AGC CCC TTG AGG 286  
 | ||| ||| ||| ||| ||| ||| ||| ||| ||| | ||| ||| ||| ||| |||  
 CAC AAC CCC GCG CAT GAC AGT AAC AGC CTC CCT GCT AGC CCC TTG AGG  
 N D A

43/73

2 298 6878

FIG. 30 14/16


Ile	Val	Glu	Asp	Glu	Glu	Tyr	Glu	Thr	Thr	Gln	Glu	Tyr	Glu	Pro	Ala	
ATA	GTG	GAG	GAT	GAG	GAA	TAT	GAA	ACG	ACC	CAG	GAG	TAC	GAA	CCA	GCT	334
ATA	GTG	GAG	GAT	GAG	GAG	TAT	GAA	ACG	ACC	CAA	GAG	TAC	GAG	CCA	GCC	
Gln	Glu	Pro	Val	Lys	Lys	Leu	Thr	Asn	Ser	Ser	Arg	Arg	Ala	Lys	Arg	
CAA	GAG	CCG	GTT	AAG	AAA	CTC	ACC	AAC	AGC	AGC	CGG	CGG	GCC	AAA	AGA	382
CAA	GAG	CCT	GTT	AAG	AAA	CTC	GCC	AA.	..T	AGC	CGG	CGG	GCC	AAA	AGA	
							A									
Thr	Lys	Pro	Asn	Gly	His	Ile	Ala	His	Arg	Leu	Glu	Met	Asp	Asn	Asn	
ACC	AAG	CCC	AAT	GGT	CAC	ATT	GCC	CAC	AGG	TTG	GAA	ATG	GAC	AAC	AAC	430
ACC	AAG	CCC	AAT	GGC	CAC	ATT	GCT	AAC	AGA	TTG	GAA	GTG	GAC	AGC	AAC	
							N					V		S		
Thr	Gly	Ala	Asp	Ser	Ser	Asn	Ser	Glu	Ser	Glu	Thr	Glu	Asp	Glu	Arg	
ACA	GGC	GCT	GAC	AGC	AGT	AAC	TCA	GAG	AGC	GAA	ACA	GAG	GAT	GAA	AGA	478
ACA	AGC	TCC	CAG	AGC	AGT	AAC	TCA	GAG	AGT	GAA	ACA	GAA	GAT	GAA	AGA	
	S	S	Q													

44/72

FIG.30 15/16

Val	Gly	Glu	Asp	Thr	Pro	Phe	Leu	Ala	Ile	Gln	Asn	Pro	Leu	Ala	Ala	
GTA	GGA	GAA	GAT	ACG	CCT	TTC	CTG	GCC	ATA	CAG	AAC	CCC	CTG	GCA	GCC	526
GTA	GGT	GAA	GAT	ACG	CCT	TTC	CTG	GGC	ATA	CAG	AAC	CCC	CTG	GCA	GCC	
								G								
Ser	Leu	Glu	Ala	Ala	Pro	Ala	Phe	Arg	Leu	Val	Asp	Ser	Arg	Thr	Asn	
AGT	CTC	GAG	GCG	GCC	CCT	GCC	TTC	CGC	CTG	GTC	GAC	AGC	AGG	ACT	AAC	574
AGT	CTT	GAG	GCA	ACA	CCT	GCC	TTC	CGC	CTG	GCT	GAC	AGC	AGG	ACT	AAC	
				T						A						
Pro	Thr	Gly	Gly	Phe	Ser	Pro	Gln	Glu	Glu	Leu	Gln	Ala	Arg	Leu	Ser	
CCA	ACA	GGC	GGC	TTC	TCT	CCG	CAG	GAA	GAA	TTG	CAG	GCC	AGG	CTC	TCC	622
CCA	GCA	GGC	CGC	TTC	TCG	ACA	CAG	GAA	GAA	ATC	CAG	GCC	AGG	CTG	TCT	
	A		R			T				I						
Gly	Val	Ile	Ala	Asn	Gln	Asp	Pro	Ile	Ala	Val	*					
GGT	GTA	ATC	GCT	AAC	CAA	GAC	CCT	ATC	GCT	GTC	TAA	AAC	CGA	AAT	ACA	672
AGT	GTA	ATT	GCT	AAC	CAA	GAC	CCT	ATT	GCT	GTA	TAA	AAC	CTA	AAT	AAA	
															S	
CCC	ATA	GAT	TCA	CCT	GTA	AAA	CTT	TAT	TTT	ATA	TAA	TAA	AGT	ATT	CCA	718
CAC	ATA	GAT	TCA	CCT	GTA	AAA	CTT	TAT	TTT	ATA	TAA	TAA	AGT	ATT	CCA	
CCT	TAA	ATT	AAA	CAA												733
CCT	TAA	ATT	AAA	CAA												

45/72



**FIG. 30 16/16**

HUMAN CODING SEGMENT E:  
(SEQ ID NO: 159)

ATG	AGA	TGG	CGA	CGC	GCC	CCG	CGC	CGC	TCC	GGG	CGT	CCC	GGC	CCC	CGG	48
Met	Arg	Trp	Arg	Arg	Ala	Pro	Arg	Arg	Ser	Gly	Arg	Pro	Gly	Pro	Arg	
GCC	CAG	CGC	CCC	GGC	TCC	GCC	GCC	CGC	TCG	TCG	CCG	CCG	CTG	CCG	CTG	96
Ala	Gln	Arg	Pro	Gly	Ser	Ala	Ala	Arg	Ser	Ser	Pro	Pro	Leu	Pro	Leu	
CTG	CCA	CTA	CTG	CTG	CTG	CTG	GGG	ACC	GCG	GCC	CTG	GCG	CCG	GGG	GCG	144
Leu	Pro	Leu	Leu	Leu	Leu	Leu	Gly	Thr	Ala	Ala	Leu	Ala	Pro	Gly	Ala	
GCG	GCC	GGC	AAC	GAG	GCG	GCT	CCC	GCG	GGG	GCC	TCG	GTG	TGC	TAC	TCG	192
Ala	Ala	Gly	Asn	Glu	Ala	Ala	Pro	Ala	Gly	Ala	Ser	Val	Cys	Tyr	Ser	
TCC	CCG	CCC	AGC	GTG	GGA	TCG	GTG	CAG	GAG	CTA	GCT	CAG	CGC	GCC	GCG	240
Ser	Pro	Pro	Ser	Val	Gly	Ser	Val	Gln	Glu	Leu	Ala	Gln	Arg	Ala	Ala	
GTG	GTG	ATC	GAG	GGA	AAG	GTG	CAC	CCG	CAG	CGG	CGG	CAG	CAG	GGG	GCA	288
Val	Val	Ile	Glu	Gly	Lys	Val	His	Pro	Gln	Arg	Arg	Gln	Gln	Gly	Ala	
CTC	GAC	AGG	AAG	GCG	GCG	GCG	GCG	GCG	GGC	GAG	GCA	GGG	GCG	TGG	GGC	336
Leu	Asp	Arg	Lys	Ala	Ala	Ala	Ala	Ala	Gly	Glu	Ala	Gly	Ala	Trp	Gly	
GGC	GAT	CGC	GAG	CCG	CCA	GCC	GCG	GGC	CCA	CGG	GCG	CTG	GGG	CCG	CCC	384
Gly	Asp	Arg	Glu	Pro	Pro	Ala	Ala	Gly	Pro	Arg	Ala	Leu	Gly	Pro	Pro	
GCC	GAG	GAG	CCG	CTG	CTC	GCC	GCC	AAC	GGG	ACC	GTG	CCC	TCT	TGG	CCC	432
Ala	Glu	Glu	Pro	Leu	Leu	Ala	Ala	Asn	Gly	Thr	Val	Pro	Ser	Trp	Pro	
ACC	GCC	CCG	GTG	CCC	AGC	GCC	GGC	GAG	CCC	GGG	GAG	GAG	GCG	CCC	TAT	480
Thr	Ala	Pro	Val	Pro	Ser	Ala	Gly	Glu	Pro	Gly	Glu	Glu	Ala	Pro	Tyr	
CTG	GTG	AAG	GTG	CAC	CAG	GTG	TGG	GCG	GTG	AAA	GCC	GGG	GGC	TTG	AAG	528
Leu	Val	Lys	Val	His	Gln	Val	Trp	Ala	Val	Lys	Ala	Gly	Gly	Leu	Lys	
AAG	GAC	TCG	CTG	CTC	ACC	GTG	CGC	CTG	GGG	ACC	TGG	GGC	CAC	CCC	GCC	576
Lys	Asp	Ser	Leu	Leu	Thr	Val	Arg	Leu	Gly	Thr	Trp	Gly	His	Pro	Ala	
TTC	CCC	TCC	TGC	GGG	AGG	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	624
Phe	Pro	Ser	Cys	Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	
ATG	GAG	CCC	GAC	GCC	AAC	AGC	ACC	AGC	CGC	GCG	CCG	GCC	GCC	TTC	CGA	672
Met	Glu	Pro	Asp	Ala	Asn	Ser	Thr	Ser	Arg	Ala	Pro	Ala	Ala	Phe	Arg	
GCC	TCT	TTC	CCC	CCT	CTG	GAG	ACG	GGC	CGG	AAC	CTC	AAG	AAG	GAG	GTC	720
Ala	Ser	Phe	Pro	Pro	Leu	Glu	Thr	Gly	Arg	Asn	Leu	Lys	Lys	Glu	Val	
AGC	CGG	GTG	CTG	TGC	AAG	CGG	TGC	G								745
Ser	Arg	Val	Leu	Cys	Lys	Arg	Cys									

46/72

2 298 6078


**FIG. 31 1/2**

**GGF2BPP5 Nucleotide Sequence & Deduced Protein Sequence**

SEQ ID NO: 144:

AGTTTCCCCC	CCCAACTTGT	CGGAACTCTG	GGCTCGCGCG	CAGGGCAGGA	GCGGAGCGGC	60
GGCGGCTGCC	CAGGCGATGC	GAGCGCGGGC	CGGACGGTAA	TCGCCTCTCC	CTCCTCGGGC	120
TGCGAGCGCG	CCGGACCGAG	GCAGCGACAG	GAGCGGACCG	CGGCGGGAAC	CGAGGACTCC	180
CCAGCGGCGC	GCCAGCAGGA	GCCACCCCGC	GAGCGTGCGA	CCGGGACGGA	GCGCCCGCCA	240
GTCCCAGGTG	GCCCCGACCG	CACGTTGCGT	CCCCGCGCTC	CCCGCCGGCG	ACAGGAGACG	300
CTCCCCCCCA	CGCCGCGCGC	GCCTCGGCCC	GGTCGCTGGC	CCGCCTCCAC	TCCGGGGACA	360
AACTTTTCCC	GAAGCCGATC	CCAGCCCTCG	GACCCAAACT	TGTCGCGCGT	CGCCTTCGCC	420
GGGAGCCGTC	CGCGCAGAGC	GTGCACTTCT	CGGGCGAG	ATG TCG GAG CGC AGA		475
				Met Ser Glu Arg Arg		
GAA GGC AAA GGC AAG GGG AAG GGC GGC AAG AAG GAC CGA GGC TCC GGG						523
Glu Gly Lys Gly Lys Gly Lys Gly Gly Lys Lys Asp Arg Gly Ser Gly						
AAG AAG CCC GTG CCC GCG GCT GGC GGC CCG AGC CCA GCC TTG CCT CCC						571
Lys Lys Pro Val Pro Ala Ala Gly Gly Pro Ser Pro Ala Leu Pro Pro						
CGC TTG AAA GAG ATG AAG ATG CAG GAG TCT GTG GCA GGT TCC AAA CTA						619
Arg Leu Lys Glu Met Lys Ser Gln Glu Ser Val Ala Gly Ser Lys Leu						
GTG CTT CGG TGC GAG ACC AGT TCT GAA TAC TCC TCT CTC AAG TTC AAG						667
Val Leu Arg Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu Lys Phe Lys						
TGG TTC AAG AAT GGG AGT GAA TTA AGC CGA AAG AAC AAA CCA CAA AAC						715
Trp Phe Lys Asn Gly Ser Glu Leu Ser Arg Lys Asn Lys Pro Gln Asn						
ATC AAG ATA CAG AAA AGG CCG GGG AAG TCA GAA CTT CGC ATT AGC AAA						763
Ile Lys Ile Gln Lys Arg Pro Gly Lys Ser Glu Leu Arg Ile Ser Lys						
GCG TCA CTG GCT GAT TCT GGA GAA TAT ATG TGC AAA GTG ATC AGC AAA						811
Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val Ile Ser Lys						

47/72




## FIG. 31 2/2

### GGF2BPP5 Nucleotide Sequence & Deduced Protein Sequence

CTA GGA AAT GAC AGT GCC TCT GCC AAC ATC ACC ATT GTG GAG TCA AAC	859
Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val Glu Ser Asn	
GAG ATC ACC ACT GGC ATG CCA GCC TCA ACT GAG ACA GCG TAT GTG TCT	907
Glu Ile Thr Thr Gly Met Pro Ala Ser Thr Glu Thr Ala Tyr Val Ser	
TCA GAG TCT CCC ATT AGA ATA TCA GTA TCA ACA GAA GGA ACA AAT ACT	955
Ser Glu Ser Pro Ile Arg Ile Ser Val Ser Thr Glu Gly Thr Asn Thr	
TCT TCA TCC ACA TCC ACA TCT ACA GCT GGG ACA AGC CAT CTT GTC AAG	1003
Ser Ser Ser Thr Ser Thr Ser Thr Ala Gly Thr Ser His Leu Val Lys	
TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT GGA GGC GAG TGC TTC	1051
Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys Phe	
ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC TTG TGC AAG TGC CCA	1099
Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys Pro	
AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC	1147
Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe	
TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT GAA TAGGCGCATG	1193
Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro Glu	
CTCAGTCGGT GCCGCTTTCT TGTTGCCGCA TCTCCCCTCA GATTCAACCT AGAGCTAGAT	1253
GCGTTTTACC AGGTCTAACA TTGACTGCCT CTGCCTGTCTG CATGAGAACA TTAACACAAG	1313
CGATTGTATG ACTTCCTCTG TCCGTGACTA GTGGGCTCTG AGCTACTCGT AGGTGCGTAA	1373
GGCTCCAGTG TTTCTGAAAT TGATCTTGAA TTACTIONGAT ACGACATGAT AGTCCCTCTC	1433
ACCCAGTGCA ATGACAATAA AGGCCTTGAA AAGTCTCACT TTTATTGAGA AAATAAAAAT	1493
CGTTCCACGG GACAGTCCCT CTCTTTTATA AAATGACCCT ATCCTTGAAA AGGAGGTGTG	1553
TTAAGTTGTA ACCAGTACAC ACTTGAAATG ATGGTAAGTT CGCTTCGGTT CAGAATGTGT	1613
TCTTTCTGAC AAATAAACAG AATAAAAAAA AAAAAAAA A	1653

48/73



## FIG. 32 1/2

### GGF2BPP2 Nucleotide Sequence & Deduced Protein Sequence

SEQ ID NO: 145:

CAT	CAN	GTG	TGG	GCG	GCG	AAA	GCC	GGG	GGC	TTG	AAG	AAG	GAC	TCG	CTG	48
His	Gln	Val	Trp	Ala	Ala	Lys	Ala	Gly	Gly	Leu	Lys	Lys	Asp	Ser	Leu	
CTC	ACC	GTG	CGC	CTG	GGC	GCC	TGG	GGC	CAC	CCC	GCC	TTC	CCC	TCC	TGC	96
Leu	Thr	Val	Arg	Leu	Gly	Ala	Trp	Gly	His	Pro	Ala	Phe	Pro	Ser	Cys	
GGG	CGC	CTC	AAG	GAG	GAC	AGC	AGG	TAC	ATC	TTC	TTC	ATG	GAG	CCC	GAG	144
Gly	Arg	Leu	Lys	Glu	Asp	Ser	Arg	Tyr	Ile	Phe	Phe	Met	Glu	Pro	Glu	
GCC	AAC	AGC	AGC	GGC	GGG	CCC	GGC	CGC	CTT	CCG	AGC	CTC	CTT	CCC	CCC	192
Ala	Asn	Ser	Ser	Gly	Gly	Pro	Gly	Arg	Leu	Pro	Ser	Leu	Leu	Pro	Pro	
TCT	CGA	GAC	GGG	CCG	GAA	CCT	CAA	GAA	GGA	GGT	CAG	CCG	GGT	GCT	GTG	240
Ser	Arg	Asp	Gly	Pro	Glu	Pro	Gln	Glu	Gly	Gly	Gln	Pro	Gly	Ala	Val	
CAA	CGG	TGC	GCC	TTG	CCT	CCC	CGC	TTG	AAA	GAG	ATG	AAG	AGT	CAG	GAG	288
Gln	Arg	Cys	Ala	Leu	Pro	Pro	Arg	Leu	Lys	Glu	Met	Lys	Ser	Gln	Glu	
TCT	GTG	GCA	GGT	TCC	AAA	CTA	GTG	CTT	CGG	TGC	GAG	ACC	AGT	TCT	GAA	336
Ser	Val	Ala	Gly	Ser	Lys	Leu	Val	Leu	Arg	Cys	Glu	Thr	Ser	Ser	Glu	
TAC	TCC	TCT	CTC	AAG	TTC	AAG	TGG	TTC	AAG	AAT	GGG	AGT	GAA	TTA	AGC	384
Tyr	Ser	Ser	Leu	Lys	Phe	Lys	Trp	Phe	Lys	Asn	Gly	Ser	Glu	Leu	Ser	
CGA	AAG	AAC	AAA	CCA	GAA	AAC	ATC	AAG	ATA	CAG	AAA	AGG	CCG	GGG	AAG	432
Arg	Lys	Asn	Lys	Pro	Glu	Asn	Ile	Lys	Ile	Gln	Lys	Arg	Pro	Gly	Lys	
TCA	GAA	CTT	CGC	ATT	AGC	AAA	GCG	TCA	CTG	GCT	GAT	TCT	GGA	GAA	TAT	480
Ser	Glu	Leu	Arg	Ile	Ser	Lys	Ala	Ser	Leu	Ala	Asp	Ser	Gly	Glu	Tyr	
ATG	TGC	AAA	GTG	ATC	AGC	AAA	CTA	GGA	AAT	GAC	AGT	GCC	TCT	GCC	AAC	528
Met	Cys	Lys	Val	Ile	Ser	Lys	Leu	Gly	Asn	Asp	Ser	Ala	Ser	Ala	Asn	

49/73

**FIG. 32 2/2**

**GGF2BPP2 Nucleotide Sequence & Deduced Protein Sequence**

ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG ACA	576
Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr Ser Thr Ala Gly Thr	
AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	624
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	672
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	720
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT	768
Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr	
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC TAC AGT ACG TCC	816
Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser	
ACT CCC TTT CTG TCT CTG CCT GAA TAGCGCATCT CAGTCGGTGC CGCTTTCTTG	870
Thr Pro Phe Leu Ser Leu Pro Glu	
TTGCCGCATC TCCCCTCAGA TTCCNCCTAG AGCTAGATGC GTTTTACCAG GTCTAACATT	930
GACTGCCTCT GCCTGTCGCA TGAGAACAATT AACACAAGCG ATTGTATGAC TTCCTCTGTC	990
CGTGACTAGT GGGCTCTGAG CTACTCGTAG GTGCGTAAGG CTCCAGTGTT TCTGAAATTG	1050
ATCTTGAATT ACTGTGATAC GACATGATAG TCCCTCTCAC CCAGTGCAAT GACAATAAAG	1110
GCCTTGAAAA GTCAAAAAAA AAAAAAAAAA	1140

50/72



2 29 6078

# FIG. 33 1/3

## GGF2BPP4 Nucleotide Sequence & Deduced Protein Sequence

SEQ ID NO: 146:

G AAG TCA GAA CTT CGC ATT AGC AAA GCG TCA CTG GCT GAT TCT GGA GAA	49
Lys Ser Glu Leu Arg Ile Ser Lys Ala Ser Leu Ala Asp Ser Gly Glu	
TAT ATG TGC AAA GTG ATC AGC AAA CTA GGA AAT GAC AGT GCC TCT GCC	97
Tyr Met Cys Lys Val Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala	
AAC ATC ACC ATT GTG GAG TCA AAC GCC ACA TCC ACA TCT ACA GCT GGG	145
Asn Ile Thr Ile Val Glu Ser Asn Ala Thr Ser Thr Ser Thr Ala Gly	
ACA AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG	193
Thr Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val	
AAT GGA GGC GAC TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA	241
Asn Gly Gly Asp Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg	
TAC TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG	289
Tyr Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu	
AAT GTG CCC ATG AAA GTC CAA ACC CAA GAA AAA GCG GAG GAG CTC TAC	337
Asn Val Pro Met Lys Val Gln Thr Gln Glu Lys Ala Glu Glu Leu Tyr	
CAG AAG AGA GTG CTC ACC ATT ACC GGC ATT TGC ATC GCG CTG CTC GTG	385
Gln Lys Arg Val Leu Thr Ile Thr Gly Ile Cys Ile Ala Leu Leu Val	
GTT GGC ATC ATG TGT GTG GTG GTC TAC TGC AAA ACC AAG AAA CAA CGG	433
Val Gly Ile Met Cys Val Val Val Tyr Cys Lys Thr Lys Lys Gln Arg	
AAA AAG CTT CAT GAC CGG CTT CGG CAG AGC CTT CGG TCT GAA AGA AAC	481
Lys Lys Leu His Asp Arg Leu Arg Gln Ser Leu Arg Ser Glu Arg Asn	
ACC ATG ATG AAC GTA GCC AAC GGG CCC CAC CAC CCC AAT CCG CCC CCC	529
Thr Met Met Asn Val Ala Asn Gly Pro His His Pro Asn Pro Pro Pro	
GAG AAC GTG CAG CTG GTG AAT CAA TAC GTA TCT AAA AAT GTC ATC TCT	577
Glu Asn Val Gln Leu Val Asn Gln Tyr Val Ser Lys Asn Val Ile Ser	

S172

**FIG. 33 2/3**

**GGF2BPP4 Nucleotide Sequence & Deduced Protein Sequence**

AGC GAG CAT ATT GTT GAG AGA GAG GCG GAG AGC TCT TTT TCC ACC AGT	625
Ser Glu His Ile Val Glu Arg Glu Ala Glu Ser Ser Phe Ser Thr Ser	
CAC TAC ACT TCG ACA GCT CAT CAT TCC ACT ACT GTC ACT CAG ACT CCC	673
His Tyr Thr Ser Thr Ala His His Ser Thr Thr Val Thr Gln Thr Pro	
AGT CAC AGC TGG AGC AAT GGA CAC ACT GAA AGC ATC ATT TCG GAA AGC	721
Ser His Ser Trp Ser Asn Gly His Thr Glu Ser Ile Ile Ser Glu Ser	
CAC TCT GTC ATC GTG ATG TCA TCC GTA GAA AAC AGT AGG CAC AGC AGC	769
His Ser Val Ile Val Met Ser Ser Val Glu Asn Ser Arg His Ser Ser	
CCG ACT GGG GGC CCG AGA GCA CGT CTC AAT GGC TTG GGA GGC CCT CGT	817
Pro Thr Gly Gly Pro Arg Gly Arg Leu Asn Gly Leu Gly Gly Pro Arg	
GAA TGT AAC AGC TTC CTC AGG CAT GCC AGA GAA ACC CCT GAC TCC TAC	865
Glu Cys Asn Ser Phe Leu Arg His Ala Arg Glu Thr Pro Asp Ser Tyr	
CGA GAC TCT CCT CAT AGT GAA AGA CAT AAC CTT ATA GCT GAG CTA AGG	913
Arg Asp Ser Pro His Ser Glu Arg His Asn Leu Ile Ala Glu Leu Arg	
AGA AAC AAG GCC CAC AGA TCC AAA TGC ATG CAG ATC CAG CTT TCC GCA	961
Arg Asn Lys Ala His Arg Ser Lys Cys Met Gln Ile Gln Leu Ser Ala	
ACT CAT CTT AGA GCT TCT TCC ATT CCC CAT TGG GCT TCA TTC TCT AAG	1009
Thr His Leu Arg Ala Ser Ser Ile Pro His Trp Ala Ser Phe Ser Lys	
ACC CCT TGG CCT TTA GGA AGG TAT GTA TCA GCA ATG ACC ACC CCG GCT	1057
Thr Pro Trp Pro Leu Gly Arg Tyr Val Ser Ala Met Thr Thr Pro Ala	
CGT ATG TCA CCT GTA GAT TTC CAC ACG CCA AGC TCC CCC AAG TCA CCC	1105
Arg Met Ser Pro Val Asp Phe His Thr Pro Ser Ser Pro Lys Ser Pro	
CCT TCG GAA ATG TCC CCG CCC GTG TCC AGC ACG ACG GTC TCC ATG CCC	1153
Pro Ser Glu Met Ser Pro Pro Val Ser Ser Thr Thr Val Ser Met Pro	

**FIG. 33 3/3**

**GGF2BPP4 Nucleotide Sequence & Deduced Protein Sequence**

TCC ATG GCG GTC AGT CCC TTC GTG GAA GAG GAG AGA CCC CTG CTC CTT	1201
Ser Met Ala Val Ser Pro Phe Val Glu Glu Glu Arg Pro Leu Leu Leu	
GTG ACG CCA CCA CGG CTG CGG GAG AAG TAT GAC CAC CAC GCC CAG CAA	1249
Val Thr Pro Pro Arg Leu Arg Glu Lys Tyr Asp His His Ala Gln Gln	
TTC AAC TCG TTC CAC TGC AAC CCC GCG CAT GAG AGC AAC AGC CTG CCC	1297
Phe Asn Ser Phe His Cys Asn Pro Ala His Glu Ser Asn Ser Leu Pro	
CCC AGC CCC TTG AGG ATA GTG GAG GAT GAG GAA TAT GAA ACG ACC CAG	1345
Pro Ser Pro Leu Arg Ile Val Glu Asp Glu Glu Tyr Glu Thr Thr Gln	
GAG TAC GAA CCA GCT CAA GAG CCG GTT AAG AAA CTC ACC AAC AGC AGC	1393
Glu Tyr Glu Pro Ala Gln Glu Pro Val Lys Lys Leu Thr Asn Ser Ser	
CGG CGG GCC AAA AGA ACC AAG CCC AAT GGT CAC ATT GCC CAC AGG TTG	1441
Arg Arg Ala Lys Arg Thr Lys Pro Asn Gly His Ile Ala His Arg Leu	
GAA ATG GAC AAC AAC ACA GGC GCT GAC AGC AGT AAC TCA GAG AGC GAA	1489
Glu Met Asp Asn Asn Thr Gly Ala Asp Ser Ser Asn Ser Glu Ser Glu	
ACA GAG GAT GAA AGA GTA GGA GAA GAT ACG CCT TTC CTG GCC ATA CAG	1537
Thr Glu Asp Glu Arg Val Gly Glu Asp Thr Pro Phe Leu Ala Ile Gln	
AAC CCC CTG GCA GCC AGT CTC GAG GCG GCC CCT GCC TTC CGC CTG GTC	1585
Asn Pro Leu Ala Ala Ser Leu Glu Ala Ala Pro Ala Phe Arg Leu Val	
GAC AGC AGG ACT AAC CCA ACA GGC GGC TTC TCT CCG CAG GAA GAA TTG	1633
Asp Ser Arg Thr Asn Pro Thr Gly Gly Phe Ser Pro Gln Glu Glu Leu	
CAG GCC AGG CTC TCC GGT GTA ATC GCT AAC CAA GAC CCT ATC GCT GTC	1681
Gln Ala Arg Leu Ser Gly Val Ile Ala Asn Gln Asp Pro Ile Ala Val	
TAAAACCGAA ATACACCCAT AGATTCACCT GTAAACTTT ATTTTATATA ATAAAGTATT	1741
CCACCTTAAA TTAAACAAAA AAA	1764

53/72

2 298 6078


**FIG. 34**

\* \* \* \* \*

**GGF2bpp5** (SEQ ID NO: 147) KCAEKEKTFCVNGGECFMVKDLSNPSRYLCKCPNEFTGDRCQNYVMASFY  
**GGF2bpp4** (SEQ ID NO: 148) KCAEKEKTFCVNGGDCFMVKDLSNPSRYLCKCQPGFTGARCTENVPMKVQ  
**hEGF** (SEQ ID NO: 149) ECLRKYKDFCIH-GECKYVKELRAPS---CKCQOEYFGERCGEKSNKTHS



54/73

**FIG. 35**  
200 kDa Tyrosine Phosphorylation  
Compared with Mitogenic Activity



# FIG. 36 GGF/Heregulin Splicing Variants

F-B-A'	1/2	F-E-B-A'
F-B-A-C-C/D-D		F-E-B-A-C-C/D-D
F-B-A-C-C/D-H		F-E-B-A-C-C/D-H
F-B-A-C-C/D-H-L		F-E-B-A-C-C/D-H-L
F-B-A-C-C/D-H-K-L		F-E-B-A-C-C/D-H-K-L
F-B-A-C-C/D-D'-H		F-E-B-A-C-C/D-D'-H
F-B-A-C-C/D-D'-H-L		F-E-B-A-C-C/D-D'-H-L
F-B-A-C-C/D-D'-H-K-L		F-E-B-A-C-C/D-D'-H-K-L
F-B-A-C-C/D'-D		F-E-B-A-C-C/D'-D
F-B-A-C-C/D'-H		F-E-B-A-C-C/D'-H
F-B-A-C-C/D'-H-L		F-E-B-A-C-C/D'-H-L
F-B-A-C-C/D'-H-K-L		F-E-B-A-C-C/D'-H-K-L
F-B-A-C-C/D'-D'-H		F-E-B-A-C-C/D'-D'-H
F-B-A-C-C/D'-D'-H-L		F-E-B-A-C-C/D'-D'-H-L
F-B-A-C-C/D'-D'-H-K-L		F-E-B-A-C-C/D'-D'-H-K-L
F-B-A-C-C/D-C/D'-D		F-E-B-A-C-C/D-C/D'-D
F-B-A-C-C/D-C/D'-H		F-E-B-A-C-C/D-C/D'-H
F-B-A-C-C/D-C/D'-H-L		F-E-B-A-C-C/D-C/D'-H-L
F-B-A-C-C/D-C/D'-H-K-L		F-E-B-A-C-C/D-C/D'-H-K-L
F-B-A-C-C/D-C/D'-D'-H		F-E-B-A-C-C/D-C/D'-D'-H
F-B-A-C-C/D-C/D'-D'-H-L		F-E-B-A-C-C/D-C/D'-D'-H-L
F-B-A-C-C/D-C/D'-D'-H-K-L		F-E-B-A-C-C/D-C/D'-D'-H-K-L
F-B-A-G-C-C/D-D		F-E-B-A-G-C-C/D-D
F-B-A-G-C-C/D-H		F-E-B-A-G-C-C/D-H
F-B-A-G-C-C/D-H-L		F-E-B-A-G-C-C/D-H-L
F-B-A-G-C-C/D-H-K-L		F-E-B-A-G-C-C/D-H-K-L
F-B-A-G-C-C/D-D'-H		F-E-B-A-G-C-C/D-D'-H
F-B-A-G-C-C/D-D'-H-L		F-E-B-A-G-C-C/D-D'-H-L
F-B-A-G-C-C/D-D'-H-K-L		F-E-B-A-G-C-C/D-D'-H-K-L
F-B-A-G-C-C/D'-D		F-E-B-A-G-C-C/D'-D
F-B-A-G-C-C/D'-H		F-E-B-A-G-C-C/D'-H
F-B-A-G-C-C/D'-H-L		F-E-B-A-G-C-C/D'-H-L
F-B-A-G-C-C/D'-H-K-L		F-E-B-A-G-C-C/D'-H-K-L
F-B-A-G-C-C/D'-D'-H		F-E-B-A-G-C-C/D'-D'-H
F-B-A-G-C-C/D'-D'-H-L		F-E-B-A-G-C-C/D'-D'-H-L
F-B-A-G-C-C/D'-D'-H-K-L		F-E-B-A-G-C-C/D'-D'-H-K-L
F-B-A-G-C-C/D-C/D'-D		F-E-B-A-G-C-C/D-C/D'-D
F-B-A-G-C-C/D-C/D'-H		F-E-B-A-G-C-C/D-C/D'-H
F-B-A-G-C-C/D-C/D'-H-L		F-E-B-A-G-C-C/D-C/D'-H-L
F-B-A-G-C-C/D-C/D'-H-K-L		F-E-B-A-G-C-C/D-C/D'-H-K-L
F-B-A-G-C-C/D-C/D'-D'-H		F-E-B-A-G-C-C/D-C/D'-D'-H
F-B-A-G-C-C/D-C/D'-D'-H-L		F-E-B-A-G-C-C/D-C/D'-D'-H-L
F-B-A-G-C-C/D-C/D'-D'-H-K-L		F-E-B-A-G-C-C/D-C/D'-D'-H-K-L



**FIG. 36 2/2**  
**GGF/Heregulin**  
**Splicing Variants**

E-B-A'

E-B-A-C-C/D-D

E-B-A-C-C/D-H

E-B-A-C-C/D-H-L

E-B-A-C-C/D-H-K-L

E-B-A-C-C/D-D'-H

E-B-A-C-C/D-D'-H-L

E-B-A-C-C/D-D'-H-K-L

E-B-A-C-C/D'-D

E-B-A-C-C/D'-H

E-B-A-C-C/D'-H-L

E-B-A-C-C/D'-H-K-L

E-B-A-C-C/D'-D'-H

E-B-A-C-C/D'-D'-H-L

E-B-A-C-C/D'-D'-H-K-L

E-B-A-C-C/D-C/D'-D

E-B-A-C-C/D-C/D'-H

E-B-A-C-C/D-C/D'-H-L

E-B-A-C-C/D-C/D'-H-K-L

E-B-A-C-C/D-C/D'-D'-H

E-B-A-C-C/D-C/D'-D'-H-L

E-B-A-C-C/D-C/D'-D'-H-K-L

E-B-A-G-C-C/D-D

E-B-A-G-C-C/D-H

E-B-A-G-C-C/D-H-L

E-B-A-G-C-C/D-H-K-L

E-B-A-G-C-C/D-D'-H

E-B-A-G-C-C/D-D'-H-L

E-B-A-G-C-C/D-D'-H-K-L

E-B-A-G-C-C/D'-D

E-B-A-G-C-C/D'-H

E-B-A-G-C-C/D'-H-L

E-B-A-G-C-C/D'-H-K-L

E-B-A-G-C-C/D'-D'-H

E-B-A-G-C-C/D'-D'-H-L

E-B-A-G-C-C/D'-D'-H-K-L

E-B-A-G-C-C/D-C/D'-D

E-B-A-G-C-C/D-C/D'-H



E-B-A-G-C-C/D-C/D'-H-L

E-B-A-G-C-C/D-C/D'-H-K-L

E-B-A-G-C-C/D-C/D'-D'-H

E-B-A-G-C-C/D-C/D'-D'-H-L

E-B-A-G-C-C/D-C/D'-D'-H-K-L



2 29 0078

**FIG. 37**  
**EGFL1**

SEQ ID NO: 150:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	144
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
GTA ATG GCC AGC TTC TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT	192
Val Met Ala Ser Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro	
GAA TAG	198
Glu	

58/72



2 298 6070

**FIG. 38**  
**EGFL2**

SEQ ID NO: 1 51:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	144
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAA GCG GAG GAG CTC TAC TAA	192
Val Pro Met Lys Val Gln Thr Gln Glu Lys Ala Glu Glu Leu Tyr	

59/72

2 298 6070

**FIG. 39**  
**EGFL3**

SEQ ID NO: 152:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	144
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
GTA ATG GCC AGC TTC TAC AAA GCG GAG GAG CTC TAC TAA	183
Val Met Ala Ser Phe Tyr Lys Ala Glu Glu Leu Tyr	

60/72

2 29 6078

# FIG. 40

## EGFL4

SEQ ID NO: 153:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC	144
Leu Cys Lys Cys Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr	
GTA ATG GCC AGC TTC TAC AAG CAT CTT GGG ATT GAA TTT ATG GAG AAA	192
Val Met Ala Ser Phe Tyr Lys His Leu Gly Ile Glu Phe Met Glu Lys	
GCG GAG GAG CTC TAC TAA	210
Ala Glu Glu Leu Tyr	

61/72

2 298 68078

# FIG. 41

## EGFL5

SEQ ID NO: 154:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	144
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT	192
Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr	
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC TAC AGT ACG TCC	240
Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Ser Thr Ser	
ACT CCC TTT CTG TCT CTG CCT GAA TAG	267
Thr Pro Phe Leu Ser Leu Pro Glu	

62/72

2 298 6078

# FIG. 42

## EGFL6


SEQ ID NO: 155:

AGC CAT CTT GTC AAG TGT GCA GAG AAG GAG AAA ACT TTC TGT GTG AAT	48
Ser His Leu Val Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn	
GGA GGC GAG TGC TTC ATG GTG AAA GAC CTT TCA AAT CCC TCA AGA TAC	96
Gly Gly Glu Cys Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr	
TTG TGC AAG TGC CAA CCT GGA TTC ACT GGA GCG AGA TGT ACT GAG AAT	144
Leu Cys Lys Cys Gln Pro Gly Phe Thr Gly Ala Arg Cys Thr Glu Asn	
GTG CCC ATG AAA GTC CAA ACC CAA GAA AAG TGC CCA AAT GAG TTT ACT	192
Val Pro Met Lys Val Gln Thr Gln Glu Lys Cys Pro Asn Glu Phe Thr	
GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC TTC TAC AAA GCG GAG	240
Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser Phe Tyr Lys Ala Glu	
GAG CTC TAC TAA	252
Glu Leu Tyr	

63/72

2 299 6578

**FIG. 43**  
GGF2HBS5



64/72


**FIG. 44 1/4**

**Nucleotide Sequence & Deduced Acid Sequence of GGF2HBS5**

SEQ ID NO: 21:

GGAATTCCTT TTTTTTTTTT TTTTTTCTT NNTTTTTTTTTT TGCCCTTATA CCTCTTCGCC	60
TTTCTGTGGT TCCATCCACT TCTTCCCCCT CCTCCTCCCA TAAACAACATC TCCTACCCCT	120
GCACCCCAA TAAATAAATA AAAGGAGGAG GGCAAGGGGG GAGGAGGAGG AGTGGTGCTG	180
CGAGGGGAAG GAAAAGGGAG GCAGCGCGAG AAGAGCCGGG CAGAGTCCGA ACCGACAGCC	240
AGAAGCCCGC ACGCACCTCG CACC ATG AGA TGG CGA CGC GCC CCG CGC CGC	291
Met Arg Trp Arg Arg Ala Pro Arg Arg	
TCC GGG CGT CCC GGC CCC CGG GCC CAG CGC CCC GGC TCC GCC GCC CGC	339
Ser Gly Arg Pro Gly Pro Arg Ala Gln Arg Pro Gly Ser Ala Ala Arg	
TCG TCG CCG CCG CTG CCG CTG CTG CCA CTA CTG CTG CTG CTG GGG ACC	387
Ser Ser Pro Pro Leu Pro Leu Leu Pro Leu Leu Leu Leu Leu Gly Thr	
Val Cys Leu Leu Thr Val	
GGF-II 09	
GCG GCC CTG GCG CCG GGG GCG GCG GCC GGC AAC GAG GCG GCT CCC GCG	435
Ala Ala Leu Ala Pro Gly Ala Ala Ala Gly Asn Glu Ala Ala Pro Ala	
Ala Ala Leu Pro Pro	
GGG GCC TCG GTG TGC TAC TCG TCC CCG CCC AGC GTG GGA TCG GTG CAG	483
Gly Ala Ser Val Cys Tyr Ser Ser Pro Pro Ser Val Gly Ser Val Gln	
Ala Ser Pro Val Ser Val Gly Ser Val Gln	
GGF-II 08	
GAG CTA GCT CAG CGC GCC GCG GTG GTG ATC GAG GGA AAG GTG CAC CCG	531
Glu Leu Ala Gln Arg Ala Ala Val Val Ile Glu Gly Lys Val His Pro	
Glu Leu Val Gln Arg Trp Phe Val Val Ile Glu Gly Lys	
GGF-II 04	

65/72



# FIG. 44 2/4

## Nucleotide Sequence & Deduced Acid Sequence of GGF2HBS5

CAG CGG CGG CAG CAG GGG GCA CTC GAC AGG AAG GCG GCG GCG GCG GCG	579
Gln Arg Arg Gln Gln Gly Ala Leu Asp Arg Lys Ala Ala Ala Ala Ala	
GGC GAG GCA GGG GCG TGG GGC GGC GAT CGC GAG CCG CCA GCC GCG GGC	627
Gly Glu Ala Gly Ala Trp Gly Gly Asp Arg Glu Pro Pro Ala Ala Gly	
CCA CGG GCG CTG GGG CCG CCC GCC GAG GAG CCG CTG CTC GCC GCC AAC	675
Pro Arg Ala Leu Gly Pro Pro Ala Glu Glu Pro Leu Leu Ala Ala Asn	
GGG ACC GTG CCC TCT TGG CCC ACC GCC CCG GTG CCC AGC GCC GGC GAG	723
Gly Thr Val Pro Ser Trp Pro Thr Ala Pro Val Pro Ser Ala Gly Glu	
CCC GGG GAG GAG GCG CCC TAT CTG GTG AAG GTG CAC CAG GTG TGG GCG	771
Pro Gly Glu Glu Ala Pro Tyr Leu Val Lys Val His Gln Val Trp Ala	
Lys Val His Glu Val Trp Ala	
GGF-II 01 & GGF-II 11	
GTG AAA GCC GGG GGC TTG AAG AAG GAC TCG CTG CTC ACC GTG CCC CTG	819
Val Lys Ala Gly Gly Leu Lys Lys Asp Ser Leu Leu Thr Val Arg Leu	
Ala Lys Asp Leu Leu Leu Xaa Val Leu	
GGF-II 10	
GGG ACC TGG GGC CAC CCC GCC TTC CCC TCC TGC GGG AGG CTC AAG GAG	867
Gly Thr Trp Gly His Pro Ala Phe Pro Ser Cys Gly Arg Leu Lys Glu	
Gly Ala Trp Gly Pro Pro Ala Phe Pro Val Xaa Tyr	
GGF-II 03	
GAC AGC AGG TAC ATC TTC TTC ATG GAG CCC GAC GCC AAC AGC ACC AGC	915
Asp Ser Arg Tyr Ile Phe Phe Met Glu Pro Asp Ala Asn Ser Thr Ser	
Tyr Ile Phe Phe Met Glu Pro Glu Ala Xaa Ser Ser Gly	
GGF-II 02	

66/72



**FIG. 44 3/4**

**Nucleotide Sequence & Deduced Acid Sequence of GGF2HBS5**


CGC GCG CCG GCC GCC TTC CGA GCC TCT TTC CCC CCT CTG GAG ACG GGC	963
Arg Ala Pro Ala Ala Phe Arg Ala Ser Phe Pro Pro Leu Glu Thr Gly	
CGG AAC CTC AAG AAG GAG GTC AGC CGG GTG CTG TGC AAG CGG TGC GCC	1011
Arg Asn Leu Lys Lys Glu Val Ser Arg Val Leu Cys Lys Arg Cys Ala	
TTG CCT CCC CAA TTG AAA GAG ATG AAA AGC CAG GAA TCG GCT GCA GGT	1059
Leu Pro Pro Gln Leu Lys Glu Met Lys Ser Gln Glu Ser Ala Ala Gly	
TCC AAA CTA GTC CTT CGG TGT GAA ACC AGT TCT GAA TAC TCC TCT CTC	1107
Ser Lys Leu Val Leu Arg Cys Glu Thr Ser Ser Glu Tyr Ser Ser Leu	
Leu Val Leu Arg	
GGF-II 06	
AGA TTC AAG TGG TTC AAG AAT GGG AAT GAA TTG AAT CGA AAA AAC AAA	1155
Arg Phe Lys Trp Phe Lys Asn Gly Asn Glu Leu Asn Arg Lys Asn Lys	
CCA CAA AAT ATC AAG ATA CAA AAA AAG CCA GGG AAG TCA GAA CTT CGC	1203
Pro Gln Asn Ile Lys Ile Gln Lys Lys Pro Gly Lys Ser Glu Leu Arg	
ATT AAC AAA GCA TCA CTG GCT GAT TCT GGA GAG TAT ATG TGC AAA GTG	1251
Ile Asn Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Cys Lys Val	
Lys Ala Ser Leu Ala Asp Ser Gly Glu Tyr Met Xaa Lyx	
GGF-II 12	
ATC AGC AAA TTA GGA AAT GAC AGT GCC TCT GCC AAT ATC ACC ATC GTG	1299
Ile Ser Lys Leu Gly Asn Asp Ser Ala Ser Ala Asn Ile Thr Ile Val	
GAA TCA AAC GCT ACA TCT ACA TCC ACC ACT GGG ACA AGC CAT CTT GTA	1347
Glu Ser Asn Ala Thr Ser Thr Ser Thr Thr Gly Thr Ser His Leu Val	

**FIG. 44 4/4**

**Nucleotide Sequence & Deduced Acid Sequence of GGF2HBS5**

AAA TGT GCG GAG AAG GAG AAA ACT TTC TGT GTG AAT GGA GGG GAG TGC Lys Cys Ala Glu Lys Glu Lys Thr Phe Cys Val Asn Gly Gly Glu Cys	1395
TTC ATG GTG AAA GAC CTT TCA AAC CCC TCG AGA TAC TTG TGC AAG TGC Phe Met Val Lys Asp Leu Ser Asn Pro Ser Arg Tyr Leu Cys Lys Cys	1443
CCA AAT GAG TTT ACT GGT GAT CGC TGC CAA AAC TAC GTA ATG GCC AGC Pro Asn Glu Phe Thr Gly Asp Arg Cys Gln Asn Tyr Val Met Ala Ser	1491
TTC TAC AGT ACG TCC ACT CCC TTT CTG TCT CTG CCT GAA Phe Tyr Ser Thr Ser Thr Pro Phe Leu Ser Leu Pro Glu	1530
TAGGAGCATG CTCAGTTGGT GCTGCTTTCT TGTTGCTGCA TCTCCCCTCA GATTCCACCT	1590
AGAGCTAGAT GTGTCTTACC AGATCTAATA TTGACTGCCT CTGCCTGTCTG CATGAGAACA	1650
TTAACAAAAG CAATTGTATT ACTTCCTCTG TTCGCGACTA GTTGGCTCTG AGATACTAAT	1710
AGGTGTGTGA GGCTCCGGAT GTTTCTGGAA TTGATATTGA ATGATGTGA ACAAATTGAT	1770
AGTCAATATC AAGCAGTGAA ATATGATAAT AAAGGCATTT CAAAGTCTCA CTTTTATTGA	1830
TAAAATAAAA ATCATTCTAC TGAACAGTCC ATCTTCTTTA TACAATGACC ACATCCTGAA	1890
AAGGGTGTG CTAAGCTGTA ACCGATATGC ACTTGAAATG ATGGTAAGTT AATTTTGATT	1950
CAGAATGTGT TATTTGTCAC AAATAAACAT AATAAAAGGA AAAAAAAAA AAA	2003

**FIG. 45A**  
rGGF Purification on Cation Exchange Column







**FIG. 47**

