

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
26 February 2004 (26.02.2004)

PCT

(10) International Publication Number
WO 2004/016760 A2

- (51) International Patent Classification⁷: C12N
- (21) International Application Number:
PCT/US2003/025625
- (22) International Filing Date: 15 August 2003 (15.08.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/404,063 16 August 2002 (16.08.2002) US
60/456,368 21 March 2003 (21.03.2003) US
60/458,853 27 March 2003 (27.03.2003) US
60/458,696 27 March 2003 (27.03.2003) US

(71) Applicant (for all designated States except US): GENENCOR INTERNATIONAL, INC. [US/US]; 925 Page Mill Road, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DAY, Anthony, G. [GB/US]; 551 Monterey Boulevard, San Francisco, CA 94127 (US). GOEDEGEBUUR, Frits [NL/NL]; Rozenlaan 128, NL-3135XV Vlaardingen (NL). GUALFETTI,

Peter [US/US]; 129 Arkansas Street #1, San Francisco, CA 94107 (US). MITCHINSON, Colin [GB/US]; 381 Myrtle Street, Half Moon Bay, CA 94019 (US). NEEFFE, Paulien [NL/NL]; Basalt 48, NL-2719 TN Zoetermeer (NL). SANDGREN, Mats [SE/SE]; Rabeniusvagen 20, S-SE-75655 Uppsala (SE). SHAW, Andrew [GB/US]; 2560 Hyde Street, San Francisco, CA 94109 (US). STAHLBERG, Jerry [SE/SE]; Malma Skogsvag 3A, S-SE-75645 Uppsala (SE).

(74) Agent: BOYD, Victoria, L.; Genencor International, INC., 925 Page Mill Road, Palo Alto, CA 94304 (US).


(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: NOVEL VARIANT *HYPROCREA JECORINA* CBH1 CELLULASES

Replicative expression pRAXdesCBH1 vector of CBH1 genes under the control of the glucoamylase promoter.



(57) Abstract: Described herein are variants of *H. jecorina* CBH I, a Cel7 enzyme. The present invention provides novel cellobiohydrolases that have improved thermostability and reversibility.

WO 2004/016760 A2



European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished upon receipt of that report*

NOVEL VARIANT

***HYPROCREA JECORINA* CBH1 CELLULASES**

CROSS-REFERENCE TO RELATED APPLICATIONS

5 [01] This application claims priority to U.S. Provisional Application No. 60/404,063, filed August 16, 2002 (Attorney Docket No. GC772P), to U.S. Provisional Application No. 60/458,853 filed March 27, 2003 (Attorney Docket No. GC772-2P), to U.S. Provisional Application No. 60/456,368 filed March 21, 2003 (Attorney Docket No. GC793P) and to
10 U.S. Provisional Application No. 60/458,696 filed March 27, 2003 (Attorney Docket No. GC793-2P), all herein incorporated by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY 15 SPONSORED RESEARCH AND DEVELOPMENT

[02] Portions of this work were funded by Subcontract No. ZCO-0-30017-01 with the National Renewable Energy Laboratory under Prime Contract No. DE-AC36-99GO10337 with the U.S. Department of Energy. Accordingly, the United States Government may
20 have certain rights in this invention.

FIELD OF THE INVENTION

[03] The present invention relates to variant cellobiohydrolase enzymes and isolated nucleic acid sequences which encode polypeptides having cellobiohydrolase activity. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the
25 nucleic acid sequences as well as methods for producing recombinant variant CBH polypeptides.

REFERENCES

1. Sheehan and Himmel *Biotechnology Progress* 15, pp 817-827 (1999)
- 30 2. Matti Linko Proceedings of the Second TRICEL Symposium on *Trichoderma reesei* Cellulases and Other Hydrolases pp 9-11 (1993)
3. Tuula T. Teeri *Trends in Biotechnology* 15, pp 160-167 (1997)
4. T.T. Teeri et al. Spec. Publ. - R. Soc. Chem., 246 (Recent Advances in Carbohydrate Bioengineering), pp 302-308. (1999)
- 35 5. PDB reference **2OVW**: Sulzenbacher, G., Schulein, M., Davies, G. J. *Biochemistry* 36 pp. 5902 (1997)
PDB reference **1A39**: Davies, G. J., Ducros, V., Lewis, R. J., Borchert, T. V., Schulein, M. *Journal of Biotechnology* 57 pp. 91 (1997)
- 40 7. PDB reference **6CEL**: Divne, C., Stahlberg, J., Teeri, T. T., Jones, T. A. *Journal of Molecular Biology* 275 pp. 309 (1998)

- 2 -

8. PDB reference **1EG1**: Kleywegt, G. J., Zou, J. Y., Divne, C., Davies, G. J., Sinning, I., Stahlberg, J., Reinikainen, T., Srisodsuk, M., Teeri, T. T., Jones, T. A. *Journal of Molecular Biology* 272 pp. 383 (1997)

9. PDB reference **1DY4 (8CEL)**: J. Stahlberg, H. Henriksson, C. Divne, R. Isaksson, G. Pettersson, G. Johansson, T. A. Jones

BACKGROUND OF THE INVENTION

[04] Cellulose and hemicellulose are the most abundant plant materials produced by photosynthesis. They can be degraded and used as an energy source by numerous microorganisms, including bacteria, yeast and fungi, that produce extracellular enzymes capable of hydrolysis of the polymeric substrates to monomeric sugars (Aro *et al.*, *J. Biol. Chem.*, vol. 276, no. 26, pp. 24309-24314, June 29, 2001). As the limits of non-renewable resources approach, the potential of cellulose to become a major renewable energy resource is enormous (Krishna *et al.*, *Bioresource Tech.* 77:193-196, 2001). The effective utilization of cellulose through biological processes is one approach to overcoming the shortage of foods, feeds, and fuels (Ohmiya *et al.*, *Biotechnol. Gen. Engineer. Rev.* vol. 14, pp. 365-414, 1997).

[05] Cellulases are enzymes that hydrolyze cellulose (beta-1,4-glucan or beta D-glucosidic linkages) resulting in the formation of glucose, cellobiose, celooligosaccharides, and the like. Cellulases have been traditionally divided into three major classes: endoglucanases (EC 3.2.1.4) ("EG"), exoglucanases or cellobiohydrolases (EC 3.2.1.91) ("CBH") and beta-glucosidases ([beta] -D-glucoside glucohydrolase; EC 3.2.1.21) ("BG"). (Knowles *et al.*, *TIBTECH* 5, 255-261, 1987; Shulein, *Methods Enzymol.*, 160, 25, pp. 234-243, 1988). Endoglucanases act mainly on the amorphous parts of the cellulose fibre, whereas cellobiohydrolases are also able to degrade crystalline cellulose (Nevalainen and Penttila, *Mycota*, 303-319, 1995). Thus, the presence of a cellobiohydrolase in a cellulase system is required for efficient solubilization of crystalline cellulose (Suurnakki, *et al. Cellulose* 7:189-209, 2000). Beta-glucosidase acts to liberate D-glucose units from cellobiose, cello-oligosaccharides, and other glucosides (Freer, *J. Biol. Chem.* vol. 268, no. 13, pp. 9337-9342, 1993).

[06] Cellulases are known to be produced by a large number of bacteria, yeast and fungi. Certain fungi produce a complete cellulase system capable of degrading crystalline forms of cellulose, such that the cellulases are readily produced in large quantities via fermentation. Filamentous fungi play a special role since many yeast, such as *Saccharomyces cerevisiae*, lack the ability to hydrolyze cellulose. See, e.g., Aro *et al.*,

2001; Aubert *et al.*, 1988; Wood *et al.*, Methods in Enzymology, vol. 160, no. 9, pp. 87-116, 1988, and Coughlan, *et al.*, "Comparative Biochemistry of Fungal and Bacterial Cellulolytic Enzyme Systems" Biochemistry and Genetics of Cellulose Degradation, pp. 11-30 1988..

5 [07] The fungal cellulase classifications of CBH, EG and BG can be further expanded to include multiple components within each classification. For example, multiple CBHs, EGs and BGs have been isolated from a variety of fungal sources including *Trichoderma reesei* which contains known genes for 2 CBHs, *i.e.*, CBH I and CBH II, at least 8 EGs, *i.e.*, EG I, EG II, EG III, EGIV, EGV, EGVI, EGVII and EGVIII, and at least 5 BGs, *i.e.*, BG1, BG2,
10 BG3, BG4 and BG5.

[08] In order to efficiently convert crystalline cellulose to glucose the complete cellulase system comprising components from each of the CBH, EG and BG classifications is required, with isolated components less effective in hydrolyzing crystalline cellulose (Filho *et al.*, Can. J. Microbiol. 42:1-5, 1996). A synergistic relationship has been observed
15 between cellulase components from different classifications. In particular, the EG-type cellulases and CBH- type cellulases synergistically interact to more efficiently degrade cellulose. See, *e.g.*, Wood, Biochemical Society Transactions, 61th Meeting, Galway, vol. 13, pp. 407-410, 1985.

[09] Cellulases are known in the art to be useful in the treatment of textiles for the
20 purposes of enhancing the cleaning ability of detergent compositions, for use as a softening agent, for improving the feel and appearance of cotton fabrics, and the like (Kumar *et al.*, Textile Chemist and Colorist, 29:37-42, 1997).

[10] Cellulase-containing detergent compositions with improved cleaning performance (US Pat. No. 4,435,307; GB App. Nos. 2,095,275 and 2,094,826) and for use in the
25 treatment of fabric to improve the feel and appearance of the textile (US Pat. Nos. 5,648,263, 5,691,178, and 5,776,757; GB App. No. 1,358,599; The Shizuoka Prefectural Hammamatsu Textile Industrial Research Institute Report, Vol. 24, pp. 54-61, 1986), have been described.

[11] Hence, cellulases produced in fungi and bacteria have received significant
30 attention. In particular, fermentation of *Trichoderma spp.* (*e.g.*, *Trichoderma longibrachiatum* or *Trichoderma reesei*) has been shown to produce a complete cellulase system capable of degrading crystalline forms of cellulose.

[12] Although cellulase compositions have been previously described, there remains a need for new and improved cellulase compositions for use in household detergents,

stonewashing compositions or laundry detergents, etc. Cellulases that exhibit improved performance are of particular interest.

BRIEF SUMMARY OF THE INVENTION

[13] The invention provides an isolated cellulase protein, identified herein as variant
5 CBH I, and nucleic acids which encode a variant CBH I.

[14] In one embodiment the invention is directed to a variant CBH I cellulase, wherein
said variant comprises a substitution or deletion at a position corresponding to one or
more of residues S8, Q17, G22, T41, N49, S57, N64, A68, A77, N89, S92, N103, A112,
S113, E193, S196, M213, L225, T226, P227, T246, D249, R251, Y252, T255, D257,
10 D259, S278, S279, K286, L288, E295, T296, S297, A299, N301, E325, T332, F338, S342,
F352, T356, Y371, T380, Y381, V393, R394, S398, V403, S411, G430, G440, T445,
T462, T484, Q487, and P491 in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2). In first
aspect, the invention encompasses an isolated nucleic acid encoding a polypeptide having
cellobiohydrolase activity, which polypeptide is a variant of a glycosyl hydrolase of family
15 7, and wherein said nucleic acid encodes a substitution at a residue which is sensitive to
temperature stress in the polypeptide encoded by said nucleic acid, wherein said variant
cellobiohydrolase is derived from *H. jecorina* cellobiohydrolase. In second aspect, the
invention encompasses an isolated nucleic acid encoding a polypeptide having
cellobiohydrolase activity, which polypeptide is a variant of a glycosyl hydrolase of family
20 7, and wherein said nucleic acid encodes a substitution at a residue which effects
enzyme processivity in the polypeptide encoded by said nucleic acid, wherein said
variant cellobiohydrolase is derived from *H. jecorina* cellobiohydrolase. In third aspect, the
invention encompasses an isolated nucleic acid encoding a polypeptide having
cellobiohydrolase activity, which polypeptide is a variant of a glycosyl hydrolase of family
25 7, and wherein said nucleic acid encodes a substitution at a residue which effects
product inhibition in the polypeptide encoded by said nucleic acid, wherein said variant
cellobiohydrolase is derived from *H. jecorina* cellobiohydrolase.

[15] In a second embodiment the invention is directed to a variant CBH I cellulose
comprising a substitution at a position corresponding to one or more of residues S8P,
30 Q17L, G22D, T41I, N49S, S57N, N64D, A68T, A77D, N89D, S92T, N103I, A112E,
S113(T/N/D), E193V, S196T, M213I, L225F, T226A, P227(L/T/A), T246(C/A), D249K,
R251A, Y252(A/Q), T255P, D257E, D259W, S278P, S279N, K286M, L288F, E295K,
T296P, S297T, A299E, N301(R/K), E325K, T332(K/Y/H), F338Y, S342Y, F352L, T356L,
Y371C, T380G, Y381D, V393G, R394A, S398T, V403D, S411F, G430F, G440R, T462I,

- 5 -

T484S, Q487L and/or P491L in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2). In one aspect of this embodiment the variant CBH I cellulase further comprises a deletion at a position corresponding to T445 in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2). In a second aspect of this embodiment the variant CBH I cellulase further comprises the
 5 deletion of residues corresponding to residues 382-393 in CBH I of *Hypocrea jecorina* (SEQ ID NO: 2).

[16] In a third embodiment the invention is directed to a variant CBH I cellulase, wherein said variant comprises a substitution at a position corresponding to a residue selected from the group consisting of S8P, N49S, A68T, A77D, N89D, S92T, S113(N/D),
 10 L225F, P227(A/L/T), D249K, T255P, D257E, S279N, L288F, E295K, S297T, A299E, N301K, T332(K/Y/H), F338Y, T356L, V393G, G430F in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2).

[17] In a fourth embodiment the invention is directed to a variant CBH I consists essentially of the mutations selected from the group consisting of

- 15 i. A112E/T226A;
- ii. S196T/S411F;
- iii. E295K/S398T;
- iv. T246C/Y371C;
- v. T41I plus deletion at T445
- 20 vi. A68T/G440R/P491L;
- vii. G22D/S278P/T296P;
- viii. T246A/R251A/Y252A;
- ix. T380G/Y381D/R394A;
- x. T380G/Y381D/R394A plus deletion of 382-393, inclusive;
- 25 xi. Y252Q/D259W/S342Y;
- xii. S113T/T255P/K286M;
- xiii. P227L/E325K/Q487L;
- xiv. P227T/T484S/F352L;
- xv. Q17L/E193V/M213I/F352L;
- 30 xvi. S8P/N49S/A68T/S113N;
- xvii. S8P/N49S/A68T/S113N/P227L;
- xviii. T41I/A112E/P227L/S278P/T296P;
- xix. S8P/N49S/A68T/A112E/T226A;
- xx. S8P/N49S/A68T/A112E/P227L;
- 35 xxi. S8P/T41I/N49S/A68T/A112E/P227L;

- 6 -

- xxii. G22D/N49S/A68T/P227L/S278P/T296P;
- xxiii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P;
- xxiv. G22D/N49S/A68T/N103I/S113N/P227L/S278P/T296P;
- xxv. G22D/N49S/A68T/N103I/A112E/P227L/S278P/T296P;
- 5 xxvi. G22D/N49S/N64D/A68T/N103I/S113N/S278P/T296P;
- xxvii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P;
- xxviii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P/N301R;
- 10 xxix. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P/N301R;
- xxx. S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/T296P/N301R;
- xxxi. S8P/T41I/N49S/S57N/A68T/S113N/P227L/D249K/S278P/T296P/N301R;
- 15 xxxii. S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/N301R;
- xxxiii. S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I;
- xxxiv. S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I;
- 20 xxxv. S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F;
- xxxvi. S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F;
- xxxvii. S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T2578P/S278P/T296P/N301R/E325K/S411F;
- xxxviii. S8P/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T2578P/T296P/N301R/E325K/V403D/S411F/T462I;
- 25 xxxix. S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T2578P/S278P/T296P/N301R/E325K/V403D/S411F/T462I;

in CBH I from *Hypocrea jecorina* (SEQ ID NO:2).

[18] In a fifth embodiment the invention is directed to a vector comprising a nucleic acid encoding a variant CBH I. In another aspect there is a construct comprising the nucleic acid of encoding the variant CBH I operably linked to a regulatory sequence.

[19] In a sixth embodiment the invention is directed to a host cell transformed with the vector comprising a nucleic acid encoding a CBH I variant.

[20] In a seventh embodiment the invention is directed to a method of producing a CBH I variant comprising the steps of:

35

- 7 -

- (a) culturing a host cell transformed with the vector comprising a nucleic acid encoding a CBH I variant in a suitable culture medium under suitable conditions to produce CBH I variant;
- (b) obtaining said produced CBH I variant.

5 [21] In an eighth embodiment the invention is directed to a detergent composition comprising a surfactant and a CBH I variant. In one aspect of this embodiment the detergent is a laundry detergent. In a second aspect of this embodiment the detergent is a dish detergent. In third aspect of this invention, the variant CBH I cellulase is used in the treatment of a cellulose containing textile, in particular, in the stonewashing or indigo dyed
10 denim.

[22] In a ninth embodiment the invention is directed to a feed additive comprising a CBH I variant.

[23] In a tenth embodiment the invention is directed to a method of treating wood pulp comprising contacting said wood pulp with a CBH I variant.

15 [24] In a eleventh embodiment the invention is directed to a method of converting biomass to sugars comprising contacting said biomass with a CBH I variant.

[25] In an embodiment, the cellulase is derived from a fungus, bacteria or Actinomycete. In another aspect, the cellulase is derived from a fungus. In a most preferred embodiment, the fungus is a filamentous fungus. It is preferred the filamentous
20 fungus belong to Euecomycete, in particular, *Aspergillus spp.*, *Gliocladium spp.*, *Fusarium spp.*, *Acremonium spp.*, *Myceliophthora spp.*, *Verticillium spp.*, *Myrothecium spp.*, or *Penicillium spp.* In a further aspect of this embodiment, the cellulase is a cellobiohydrolase.

[26] Other objects, features and advantages of the present invention will become
25 apparent from the following detailed description. It should be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the scope and spirit of the invention will become apparent to one skilled in the art from this detailed description.

30 BRIEF DESCRIPTION OF THE DRAWINGS

[27] Figure 1 is the nucleic acid (lower line; SEQ ID NO: 1) and amino acid (upper line; SEQ ID NO: 2) sequence of the wild type Cel7A (CBH I) from *H. jecorina*.

[28] Figure 2 is the 3-D structure of *H. jecorina* CBH I.

[29] Figure 3 shows the amino acid alignment of the Cel7 family members for which there were crystal structures available. The sequences are: 2OVW - *Fusarium oxysporum* Cel7B, 1A39 - *Humicola insolens* Cel7B, 6CEL - *Hypocrea jecorina* Cel7A, 1EG1 - *Hypocrea jecorina* Cel7B.

5 [30] Figure 4 illustrates the crystal structures from the catalytic domains of these four Cel7 homologues aligned and overlaid as described herein.

[31] Figure 5 A-M is the nucleic acid sequence and deduced amino acid sequence for eight single residue mutations and five multiple mutation variants.

[32] Figure 6 A-D is the nucleic acid sequence for pTrex2.

10 [33] Figure 7 A & B depicts the construction of the expression plasmid pTEX.

[34] Figure 8 A-J is the amino acid alignment of all 42 members of the Cel7 family.

[35] Figure 9A is a representation of the thermal profiles of the wild type and eight single residue variants. Figure 9B is a representation of the thermal profiles of the wild type and five variants. Legend for Figure 9B: Cel7A = wild-type *H. jecorina* CBH I; N301K = N301K variant; 334 = P227L variant; 340 = S8P/N49S/A68T/S113N variant; 350 = S8P/N49S/A68T/S113N/ P227L variant; and 363 = S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P variant.

[36] Figure 10 is the pRAX1 vector. This vector is based on the plasmid pGAPT2 except a 5259bp HindIII fragment of *Aspergillus nidulans* genomic DNA fragment AMA1
20 sequence (*Molecular Microbiology* 1996 19:565-574) was inserted. Base 1 to 1134 contains *Aspergillus niger* glucoamylase gene promoter. Base 3098 to 3356 and 4950 to 4971 contains *Aspergillus niger* glucoamylase terminator. *Aspergillus nidulans* pyrG gene was inserted from 3357 to 4949 as a marker for fungal transformation. There is a multiple cloning site (MCS) into which genes may be inserted.

25 [37] Figure 11 is the pRAXdes2 vector backbone. This vector is based on the plasmid vector pRAX1. A Gateway cassette has been inserted into pRAX1 vector (indicated by the arrow on the interior of the circular plasmid). This cassette contains recombination sequence attR1 and attR2 and the selection marker catH and ccdB. The vector has been made according to the manual given in Gateway™ Cloning Technology: version 1 page
30 34-38 and can only replicate in *E. coli* DB3.1 from Invitrogen; in other *E. coli* hosts the ccdB gene is lethal. First a PCR fragment is made with primers containing attB1/2 recombination sequences. This fragment is recombined with pDONR201 (commercially available from Invitrogen); this vector contains attP1/2 recombination sequences with catH and ccdB in between the recombination sites. The BP clonase enzymes from Invitrogen
35 are used to recombine the PCR fragment in this so-called ENTRY vector, clones with the

PCR fragment inserted can be selected at 50µg/ml kanamycin because clones expressing ccdB do not survive. Now the att sequences are altered and called attL1 and attL2. The second step is to recombine this clone with the pRAXdes2 vector (containing attR1 and attR2 catH and ccdB in between the recombination sites). The LR clonase enzymes from Invitrogen are used to recombine the insert from the ENTRY vector in the destination vector. Only pRAXCBH1 vectors are selected using 100µg/ml ampicillin because ccdB is lethal and the ENTRY vector is sensitive to ampicillin. By this method the expression vector is now prepared and can be used to transform *A. niger*.

[38] Figure 12 provides an illustration of the pRAXdes2cbh1 vector which was used for expression of the nucleic acids encoding the CBH1 variants in *Aspergillus*. A nucleic acid encoding a CBH1 enzyme homolog or variant was cloned into the vector by homologous recombination of the att sequences.

DETAILED DESCRIPTION

[39] The invention will now be described in detail by way of reference only using the following definitions and examples. All patents and publications, including all sequences disclosed within such patents and publications, referred to herein are expressly incorporated by reference.

[40] Unless defined otherwise herein, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY, 2D ED., John Wiley and Sons, New York (1994), and Hale & Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY, Harper Perennial, NY (1991) provide one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, nucleic acids are written left to right in 5' to 3' orientation; amino acid sequences are written left to right in amino to carboxy orientation, respectively. Practitioners are particularly directed to Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., 1989, and Ausubel FM *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, 1993, for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary.

[41] The headings provided herein are not limitations of the various aspects or embodiments of the invention which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification as a whole.

5 [42] All publications cited herein are expressly incorporated herein by reference for the purpose of describing and disclosing compositions and methodologies which might be used in connection with the invention.

I. DEFINITIONS

10 [43] The term "polypeptide" as used herein refers to a compound made up of a single chain of amino acid residues linked by peptide bonds. The term "protein" as used herein may be synonymous with the term "polypeptide" or may refer, in addition, to a complex of two or more polypeptides.

15 [44] "Variant" means a protein which is derived from a precursor protein (e.g., the native protein) by addition of one or more amino acids to either or both the C- and N-terminal end, substitution of one or more amino acids at one or a number of different sites in the amino acid sequence, or deletion of one or more amino acids at either or both ends of the protein or at one or more sites in the amino acid sequence. The preparation of an enzyme variant is preferably achieved by modifying a DNA sequence which encodes for the native protein, transformation of that DNA sequence into a suitable host, and expression of the modified DNA sequence to form the derivative enzyme. The variant CBH I enzyme of the invention includes peptides comprising altered amino acid sequences in comparison with a precursor enzyme amino acid sequence wherein the variant CBH enzyme retains the characteristic cellulolytic nature of the precursor enzyme but which may have altered properties in some specific aspect. For example, a variant CBH enzyme may have an increased pH optimum or increased temperature or oxidative stability but will retain its characteristic cellulolytic activity. It is contemplated that the variants according to the present invention may be derived from a DNA fragment encoding a cellulase variant CBH enzyme wherein the functional activity of the expressed cellulase derivative is retained. For example, a DNA fragment encoding a cellulase may further include a DNA sequence or portion thereof encoding a hinge or linker attached to the cellulase DNA sequence at either the 5' or 3' end wherein the functional activity of the encoded cellulase domain is retained.

30 [45] "Equivalent residues" may also be defined by determining homology at the level of tertiary structure for a precursor cellulase whose tertiary structure has been determined by x-ray crystallography. Equivalent residues are defined as those for which the atomic

35

- 11 -

coordinates of two or more of the main chain atoms of a particular amino acid residue of a cellulase and *Hypocrea jecorina* CBH (N on N, CA on CA, C on C and O on O) are within 0.13nm and preferably 0.1nm after alignment. Alignment is achieved after the best model has been oriented and positioned to give the maximum overlap of atomic coordinates of non-hydrogen protein atoms of the cellulase in question to the *H. jecorina* CBH I. The best model is the crystallographic model giving the lowest R factor for experimental diffraction data at the highest resolution available.

$$R\ factor = \frac{\sum_h |Fo(h)| - |Fc(h)|}{\sum_h |Fo(h)|}$$

[46] Equivalent residues which are functionally analogous to a specific residue of *H. jecorina* CBH I are defined as those amino acids of a cellulase which may adopt a conformation such that they either alter, modify or contribute to protein structure, substrate binding or catalysis in a manner defined and attributed to a specific residue of the *H. jecorina* CBH I. Further, they are those residues of the cellulase (for which a tertiary structure has been obtained by x-ray crystallography) which occupy an analogous position to the extent that, although the main chain atoms of the given residue may not satisfy the criteria of equivalence on the basis of occupying a homologous position, the atomic coordinates of at least two of the side chain atoms of the residue lie with 0.13nm of the corresponding side chain atoms of *H. jecorina* CBH I. The crystal structure of *H. jecorina* CBH I is shown in Figure 2.

[47] The term "nucleic acid molecule" includes RNA, DNA and cDNA molecules. It will be understood that, as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding a given protein such as CBH I may be produced. The present invention contemplates every possible variant nucleotide sequence, encoding CBH I, all of which are possible given the degeneracy of the genetic code.

[48] A "heterologous" nucleic acid construct or sequence has a portion of the sequence which is not native to the cell in which it is expressed. Heterologous, with respect to a control sequence refers to a control sequence (*i.e.* promoter or enhancer) that does not function in nature to regulate the same gene the expression of which it is currently regulating. Generally, heterologous nucleic acid sequences are not endogenous to the cell or part of the genome in which they are present, and have been added to the cell, by infection, transfection, transformation, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding

sequence combination that is the same as, or different from a control sequence/DNA coding sequence combination found in the native cell.

[49] As used herein, the term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

[50] Accordingly, an "expression cassette" or "expression vector" is a nucleic acid construct generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a target cell. The recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA, virus, or nucleic acid fragment. Typically, the recombinant expression cassette portion of an expression vector includes, among other sequences, a nucleic acid sequence to be transcribed and a promoter.

[51] As used herein, the term "plasmid" refers to a circular double-stranded (ds) DNA construct used as a cloning vector, and which forms an extrachromosomal self-replicating genetic element in many bacteria and some eukaryotes.

[52] As used herein, the term "selectable marker-encoding nucleotide sequence" refers to a nucleotide sequence which is capable of expression in cells and where expression of the selectable marker confers to cells containing the expressed gene the ability to grow in the presence of a corresponding selective agent, or under corresponding selective growth conditions.

[53] As used herein, the term "promoter" refers to a nucleic acid sequence that functions to direct transcription of a downstream gene. The promoter will generally be appropriate to the host cell in which the target gene is being expressed. The promoter together with other transcriptional and translational regulatory nucleic acid sequences (also termed "control sequences") are necessary to express a given gene. In general, the transcriptional and translational regulatory sequences include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences.

[54] "Chimeric gene" or "heterologous nucleic acid construct", as defined herein refers to a non-native gene (*i.e.*, one that has been introduced into a host) that may be composed of parts of different genes, including regulatory elements. A chimeric gene construct for transformation of a host cell is typically composed of a transcriptional regulatory region (promoter) operably linked to a heterologous protein coding sequence,

or, in a selectable marker chimeric gene, to a selectable marker gene encoding a protein conferring antibiotic resistance to transformed cells. A typical chimeric gene of the present invention, for transformation into a host cell, includes a transcriptional regulatory region that is constitutive or inducible, a protein coding sequence, and a terminator sequence. A chimeric gene construct may also include a second DNA sequence encoding a signal peptide if secretion of the target protein is desired.

[55] A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA encoding a secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors, linkers or primers for PCR are used in accordance with conventional practice.

[56] As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, that may or may not include regions preceding and following the coding region, e.g. 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

[57] In general, nucleic acid molecules which encode the variant CBH I will hybridize, under moderate to high stringency conditions to the wild type sequence provided herein as SEQ ID NO:1. However, in some cases a CBH I-encoding nucleotide sequence is employed that possesses a substantially different codon usage, while the protein encoded by the CBH I-encoding nucleotide sequence has the same or substantially the same amino acid sequence as the native protein. For example, the coding sequence may be modified to facilitate faster expression of CBH I in a particular prokaryotic or eukaryotic expression system, in accordance with the frequency with which a particular codon is utilized by the host. Te'o, *et al.* (FEMS Microbiology Letters 190:13-19, 2000), for example, describes the optimization of genes for expression in filamentous fungi.

[58] A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one another under moderate to high stringency hybridization and wash conditions. Hybridization

conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe. For example, "maximum stringency" typically occurs at about $T_m - 5^\circ\text{C}$ (5° below the T_m of the probe); "high stringency" at about $5-10^\circ$ below the T_m ; "moderate" or "intermediate stringency" at about $10-20^\circ$ below the T_m of the probe; and "low stringency" at about $20-25^\circ$ below the T_m . Functionally, maximum stringency conditions may be used to identify sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to identify sequences having about 80% or more sequence identity with the probe.

[59] Moderate and high stringency hybridization conditions are well known in the art (see, for example, Sambrook, *et al*, 1989, Chapters 9 and 11, and in Ausubel, F.M., *et al.*, 1993, expressly incorporated by reference herein). An example of high stringency conditions includes hybridization at about 42°C in 50% formamide, 5X SSC, 5X Denhardt's solution, 0.5% SDS and 100 $\mu\text{g/ml}$ denatured carrier DNA followed by washing two times in 2X SSC and 0.5% SDS at room temperature and two additional times in 0.1X SSC and 0.5% SDS at 42°C .

[60] As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

[61] As used herein, the terms "transformed", "stably transformed" or "transgenic" with reference to a cell means the cell has a non-native (heterologous) nucleic acid sequence integrated into its genome or as an episomal plasmid that is maintained through multiple generations.

[62] As used herein, the term "expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation.

[63] The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

[64] It follows that the term "CBH I expression" refers to transcription and translation of the *cbh I* gene, the products of which include precursor RNA, mRNA, polypeptide, post-translationally processed polypeptides, and derivatives thereof, including CBH I from related species such as *Trichoderma koningii*, *Hypocrea jecorina* (also known as
5 *Trichoderma longibrachiatum*, *Trichoderma reesei* or *Trichoderma viride*) and *Hypocrea schweinitzii*. By way of example, assays for CBH I expression include Western blot for CBH I protein, Northern blot analysis and reverse transcriptase polymerase chain reaction (RT-PCR) assays for CBH I mRNA, and endoglucanase activity assays as described in Shoemaker S.P. and Brown R.D.Jr. (Biochim. Biophys. Acta, 1978, 523:133-146) and
10 Schulein (Methods Enzymol., 160, 25, pp. 234-243, 1988).

[65] The term "alternative splicing" refers to the process whereby multiple polypeptide isoforms are generated from a single gene, and involves the splicing together of nonconsecutive exons during the processing of some, but not all, transcripts of the gene. Thus a particular exon may be connected to any one of several alternative exons to form
15 messenger RNAs. The alternatively-spliced mRNAs produce polypeptides ("splice variants") in which some parts are common while other parts are different.

[66] The term "signal sequence" refers to a sequence of amino acids at the N-terminal portion of a protein which facilitates the secretion of the mature form of the protein outside the cell. The mature form of the extracellular protein lacks the signal sequence which is
20 cleaved off during the secretion process.

[67] By the term "host cell" is meant a cell that contains a vector and supports the replication, and/or transcription or transcription and translation (expression) of the expression construct. Host cells for use in the present invention can be prokaryotic cells, such as *E. coli*, or eukaryotic cells such as yeast, plant, insect, amphibian, or mammalian
25 cells. In general, host cells are filamentous fungi.

[68] The term "filamentous fungi" means any and all filamentous fungi recognized by those of skill in the art. A preferred fungus is selected from the group consisting of *Aspergillus*, *Trichoderma*, *Fusarium*, *Chrysosporium*, *Penicillium*, *Humicola*, *Neurospora*, or alternative sexual forms thereof such as *Emericella*, *Hypocrea*. It has now been
30 demonstrated that the asexual industrial fungus *Trichoderma reesei* is a clonal derivative of the ascomycete *Hypocrea jecorina*. See Kuhls et al., PNAS (1996) 93:7755-7760.

[69] The term "cellooligosaccharide" refers to oligosaccharide groups containing from 2-8 glucose units and having β -1,4 linkages, e.g., cellobiose.

[70] The term "cellulase" refers to a category of enzymes capable of hydrolyzing
35 cellulose polymers to shorter cello-oligosaccharide oligomers, cellobiose and/or glucose.

Numerous examples of cellulases, such as exoglucanases, exocellobiohydrolases, endoglucanases, and glucosidases have been obtained from cellulolytic organisms, particularly including fungi, plants and bacteria.

[71] CBH I from *Hypocrea jecorina* is a member of the Glycosyl Hydrolase Family 7 (hence Cel 7) and, specifically, was the first member of that family identified in *Hypocrea jecorina* (hence Cel 7A). The Glycosyl Hydrolase Family 7 contains both Endoglucanases and Cellobiohydrolases/exoglucanases, and that CBH I is the latter. Thus, the phrases CBH I, CBH I-type protein and Cel 7 cellobiohydrolases may be used interchangeably herein.

[72] The term "cellulose binding domain" as used herein refers to portion of the amino acid sequence of a cellulase or a region of the enzyme that is involved in the cellulose binding activity of a cellulase or derivative thereof. Cellulose binding domains generally function by non-covalently binding the cellulase to cellulose, a cellulose derivative or other polysaccharide equivalent thereof. Cellulose binding domains permit or facilitate hydrolysis of cellulose fibers by the structurally distinct catalytic core region, and typically function independent of the catalytic core. Thus, a cellulose binding domain will not possess the significant hydrolytic activity attributable to a catalytic core. In other words, a cellulose binding domain is a structural element of the cellulase enzyme protein tertiary structure that is distinct from the structural element which possesses catalytic activity. Cellulose binding domain and cellulose binding module may be used interchangeably herein.

[73] As used herein, the term "surfactant" refers to any compound generally recognized in the art as having surface active qualities. Thus, for example, surfactants comprise anionic, cationic and nonionic surfactants such as those commonly found in detergents. Anionic surfactants include linear or branched alkylbenzenesulfonates; alkyl or alkenyl ether sulfates having linear or branched alkyl groups or alkenyl groups; alkyl or alkenyl sulfates; olefinsulfonates; and alkanesulfonates. Ampholytic surfactants include quaternary ammonium salt sulfonates, and betaine-type ampholytic surfactants. Such ampholytic surfactants have both the positive and negative charged groups in the same molecule. Nonionic surfactants may comprise polyoxyalkylene ethers, as well as higher fatty acid alkanolamides or alkylene oxide adduct thereof, fatty acid glycerine monoesters, and the like.

[74] As used herein, the term "cellulose containing fabric" refers to any sewn or unsewn fabrics, yarns or fibers made of cotton or non-cotton containing cellulose or cotton or non-

cotton containing cellulose blends including natural cellulosics and manmade cellulosics (such as jute, flax, ramie, rayon, and lyocell).

[75] As used herein, the term "cotton-containing fabric" refers to sewn or unsewn fabrics, yarns or fibers made of pure cotton or cotton blends including cotton woven
5 fabrics, cotton knits, cotton denims, cotton yarns, raw cotton and the like.

[76] As used herein, the term "stonewashing composition" refers to a formulation for use in stonewashing cellulose containing fabrics. Stonewashing compositions are used to modify cellulose containing fabrics prior to sale, *i.e.*, during the manufacturing process. In contrast, detergent compositions are intended for the cleaning of soiled garments and are
10 not used during the manufacturing process.

[77] As used herein, the term "detergent composition" refers to a mixture which is intended for use in a wash medium for the laundering of soiled cellulose containing fabrics. In the context of the present invention, such compositions may include, in addition to cellulases and surfactants, additional hydrolytic enzymes, builders, bleaching agents,
15 bleach activators, bluing agents and fluorescent dyes, caking inhibitors, masking agents, cellulase activators, antioxidants, and solubilizers.

[78] As used herein, the term "decrease or elimination in expression of the *cbh1* gene" means that either that the *cbh1* gene has been deleted from the genome and therefore cannot be expressed by the recombinant host microorganism; or that the *cbh1* gene has
20 been modified such that a functional CBH1 enzyme is not produced by the host microorganism.

[79] The term "variant *cbh1* gene" or "variant CBH1" means, respectively, that the nucleic acid sequence of the *cbh1* gene from *H. jecorina* has been altered by removing, adding, and/or manipulating the coding sequence or the amino acid sequence of the
25 expressed protein has been modified consistent with the invention described herein.

[80] As used herein, the term "purifying" generally refers to subjecting transgenic nucleic acid or protein containing cells to biochemical purification and/or column chromatography.

[81] As used herein, the terms "active" and "biologically active" refer to a biological
30 activity associated with a particular protein and are used interchangeably herein. For example, the enzymatic activity associated with a protease is proteolysis and, thus, an active protease has proteolytic activity. It follows that the biological activity of a given protein refers to any biological activity typically attributed to that protein by those of skill in the art.

- 18 -

[82] As used herein, the term "enriched" means that the CBH is found in a concentration that is greater relative to the CBH concentration found in a wild-type, or naturally occurring, fungal cellulase composition. The terms enriched, elevated and enhanced may be used interchangeably herein.

5 [83] A wild type fungal cellulase composition is one produced by a naturally occurring fungal source and which comprises one or more BGL, CBH and EG components wherein each of these components is found at the ratio produced by the fungal source. Thus, an enriched CBH composition would have CBH at an altered ratio wherein the ratio of CBH to other cellulase components (i.e., EGs, beta-glucosidases and other endoglucanases) is
10 elevated. This ratio may be increased by either increasing CBH or decreasing (or eliminating) at least one other component by any means known in the art.

[84] Thus, to illustrate, a naturally occurring cellulase system may be purified into substantially pure components by recognized separation techniques well published in the literature, including ion exchange chromatography at a suitable pH, affinity
15 chromatography, size exclusion and the like. For example, in ion exchange chromatography (usually anion exchange chromatography), it is possible to separate the cellulase components by eluting with a pH gradient, or a salt gradient, or both a pH and a salt gradient. The purified CBH may then be added to the enzymatic solution resulting in an enriched CBH solution. It is also possible to elevate the amount of CBH I produced by
20 a microbe using molecular genetics methods to overexpress the gene encoding CBH, possibly in conjunction with deletion of one or more genes encoding other cellulases.

[85] Fungal cellulases may contain more than one CBH component. The different components generally have different isoelectric points which allow for their separation via ion exchange chromatography and the like. Either a single CBH component or a
25 combination of CBH components may be employed in an enzymatic solution.

[86] When employed in enzymatic solutions, the homolog or variant CBH1 component is generally added in an amount sufficient to allow the highest rate of release of soluble sugars from the biomass. The amount of homolog or variant CBH1 component added depends upon the type of biomass to be saccharified which can be readily determined by
30 the skilled artisan. However, when employed, the weight percent of the homolog or variant CBH1 component relative to any EG type components present in the cellulase composition is from preferably about 1, preferably about 5, preferably about 10, preferably about 15, or preferably about 20 weight percent to preferably about 25, preferably about 30, preferably about 35, preferably about 40, preferably about 45 or preferably about 50
35 weight percent. Furthermore, preferred ranges may be about 0.5 to about 15 weight

percent, about 0.5 to about 20 weight percent, from about 1 to about 10 weight percent, from about 1 to about 15 weight percent, from about 1 to about 20 weight percent, from about 1 to about 25 weight percent, from about 5 to about 20 weight percent, from about 5 to about 25 weight percent, from about 5 to about 30 weight percent, from about 5 to about 35 weight percent, from about 5 to about 40 weight percent, from about 5 to about 45 weight percent, from about 5 to about 50 weight percent, from about 10 to about 20 weight percent, from about 10 to about 25 weight percent, from about 10 to about 30 weight percent, from about 10 to about 35 weight percent, from about 10 to about 40 weight percent, from about 10 to about 45 weight percent, from about 10 to about 50 weight percent, from about 15 to about 20 weight percent, from about 15 to about 25 weight percent, from about 15 to about 30 weight percent, from about 15 to about 35 weight percent, from about 15 to about 30 weight percent, from about 15 to about 45 weight percent, from about 15 to about 50 weight percent.

II. HOST ORGANISMS

[87] Filamentous fungi include all filamentous forms of the subdivision Eumycota and Oomycota. The filamentous fungi are characterized by vegetative mycelium having a cell wall composed of chitin, glucan, chitosan, mannan, and other complex polysaccharides, with vegetative growth by hyphal elongation and carbon catabolism that is obligately aerobic.

[88] In the present invention, the filamentous fungal parent cell may be a cell of a species of, but not limited to, *Trichoderma*, e.g., *Trichoderma longibrachiatum*, *Trichoderma viride*, *Trichoderma koningii*, *Trichoderma harzianum*; *Penicillium sp.*; *Humicola sp.*, including *Humicola insolens* and *Humicola grisea*; *Chrysosporium sp.*, including *C. lucknowense*; *Gliocladium sp.*; *Aspergillus sp.*; *Fusarium sp.*, *Neurospora sp.*, *Hypocrea sp.*, and *Emericella sp.* As used herein, the term "*Trichoderma*" or "*Trichoderma sp.*" refers to any fungal strains which have previously been classified as *Trichoderma* or are currently classified as *Trichoderma*.

[89] In one preferred embodiment, the filamentous fungal parent cell is an *Aspergillus niger*, *Aspergillus awamori*, *Aspergillus aculeatus*, or *Aspergillus nidulans* cell.

[90] In another preferred embodiment, the filamentous fungal parent cell is a *Trichoderma reesei* cell.

III. CELLULASES

[91] Cellulases are known in the art as enzymes that hydrolyze cellulose (beta-1,4-glucan or beta D-glucosidic linkages) resulting in the formation of glucose, cellobiose,

cellooligosaccharides, and the like. As set forth above, cellulases have been traditionally divided into three major classes: endoglucanases (EC 3.2.1.4) ("EG"), exoglucanases or cellobiohydrolases (EC 3.2.1.91) ("CBH") and beta-glucosidases (EC 3.2.1.21) ("BG"). (Knowles, *et al.*, TIBTECH 5, 255-261, 1987; Schulein, 1988).

5 [92] Certain fungi produce complete cellulase systems which include exo-cellobiohydrolases or CBH-type cellulases, endoglucanases or EG-type cellulases and beta-glucosidases or BG-type cellulases (Schulein, 1988). However, sometimes these systems lack CBH-type cellulases and bacterial cellulases also typically include little or no CBH-type cellulases. In addition, it has been shown that the EG components and CBH
10 components synergistically interact to more efficiently degrade cellulose. See, *e.g.*, Wood, 1985. The different components, *i.e.*, the various endoglucanases and exocellobiohydrolases in a multi-component or complete cellulase system, generally have different properties, such as isoelectric point, molecular weight, degree of glycosylation, substrate specificity and enzymatic action patterns.

15 [93] It is believed that endoglucanase-type cellulases hydrolyze internal beta -1,4-glucosidic bonds in regions of low crystallinity of the cellulose and exo-cellobiohydrolase-type cellulases hydrolyze cellobiose from the reducing or non-reducing end of cellulose. It follows that the action of endoglucanase components can greatly facilitate the action of exo-cellobiohydrolases by creating new chain ends which are recognized by exo-
20 cellobiohydrolase components. Further, beta-glucosidase-type cellulases have been shown to catalyze the hydrolysis of alkyl and/or aryl β -D-glucosides such as methyl β -D-glucoside and p-nitrophenyl glucoside as well as glycosides containing only carbohydrate residues, such as cellobiose. This yields glucose as the sole product for the microorganism and reduces or eliminates cellobiose which inhibits cellobiohydrolases and
25 endoglucanases.

[94] Cellulases also find a number of uses in detergent compositions including to enhance cleaning ability, as a softening agent and to improve the feel of cotton fabrics (Hemmpel, ITB Dyeing/Printing/Finishing 3:5-14, 1991; Tyndall, Textile Chemist and Colorist 24:23-26, 1992; Kumar *et al.*, Textile Chemist and Colorist, 29:37-42, 1997).
30 While the mechanism is not part of the invention, softening and color restoration properties of cellulase have been attributed to the alkaline endoglucanase components in cellulase compositions, as exemplified by U.S. Patent Nos. 5,648,263, 5,691,178, and 5,776,757, which disclose that detergent compositions containing a cellulase composition enriched in a specified alkaline endoglucanase component impart color restoration and improved
35 softening to treated garments as compared to cellulase compositions not enriched in such

a component. In addition, the use of such alkaline endoglucanase components in detergent compositions has been shown to complement the pH requirements of the detergent composition (e.g., by exhibiting maximal activity at an alkaline pH of 7.5 to 10, as described in U.S. Patent Nos. 5,648,263, 5,691,178, and 5,776,757).

5 [95] Cellulase compositions have also been shown to degrade cotton-containing fabrics, resulting in reduced strength loss in the fabric (U.S. Patent No. 4,822,516), contributing to reluctance to use cellulase compositions in commercial detergent applications. Cellulase compositions comprising endoglucanase components have been suggested to exhibit reduced strength loss for cotton-containing fabrics as compared to
10 compositions comprising a complete cellulase system.

[96] Cellulases have also been shown to be useful in degradation of cellulase biomass to ethanol (wherein the cellulase degrades cellulose to glucose and yeast or other microbes further ferment the glucose into ethanol), in the treatment of mechanical pulp (Pere *et al.*, 1996), for use as a feed additive (WO 91/04673) and in grain wet milling.

15 [97] Most CBHs and EGs have a multidomain structure consisting of a core domain separated from a cellulose binding domain (CBD) by a linker peptide (Suurnakki *et al.*, 2000). The core domain contains the active site whereas the CBD interacts with cellulose by binding the enzyme to it (van Tilbeurgh *et al.*, 1986; Tomme *et al.*, Eur. J. Biochem. 170:575-581, 1988). The CBDs are particularly important in the hydrolysis of crystalline
20 cellulose. It has been shown that the ability of cellobiohydrolases to degrade crystalline cellulose clearly decreases when the CBD is absent (Linder and Teeri, J. Biotechnol. 57:15-28, 1997). However, the exact role and action mechanism of CBDs is still a matter of speculation. It has been suggested that the CBD enhances the enzymatic activity merely by increasing the effective enzyme concentration at the surface of cellulose
25 (Stahlberg *et al.*, Bio/Technol. 9:286-290, 1991), and/or by loosening single cellulose chains from the cellulose surface (Tormo *et al.*, EMBO J. vol. 15, no. 21, pp. 5739-5751, 1996). Most studies concerning the effects of cellulase domains on different substrates have been carried out with core proteins of cellobiohydrolases, as their core proteins can easily be produced by limited proteolysis with papain (Tomme *et al.*, 1988). Numerous
30 cellulases have been described in the scientific literature, examples of which include: from *Trichoderma reesei*: Shoemaker, S. et al., Bio/Technology, 1:691-696, 1983, which discloses *CBHI*; Teeri, T. et al., Gene, 51:43-52, 1987, which discloses *CBHII*. Cellulases from species other than *Trichoderma* have also been described e.g., Ooi *et al.*, Nucleic Acids Research, vol. 18, no. 19, 1990, which discloses the cDNA sequence coding for
35 endoglucanase F1-CMC produced by *Aspergillus aculeatus*; Kawaguchi T *et al.*, Gene

- 22 -

173(2):287-8, 1996, which discloses the cloning and sequencing of the cDNA encoding beta-glucosidase 1 from *Aspergillus aculeatus*; Sakamoto *et al.*, *Curr. Genet.* 27:435-439, 1995, which discloses the cDNA sequence encoding the endoglucanase CMCCase-1 from *Aspergillus kawachii* IFO 4308; Saarilahti *et al.*, *Gene* 90:9-14, 1990, which discloses an endoglucanase from *Erwinia carotovora*; Spilliaert R, *et al.*, *Eur J Biochem.* 224(3):923-30, 1994, which discloses the cloning and sequencing of bglA, coding for a thermostable beta-glucanase from *Rhodothermus marinus*; and Halldorsdottir S *et al.*, *Appl Microbiol Biotechnol.* 49(3):277-84, 1998, which discloses the cloning, sequencing and overexpression of a *Rhodothermus marinus* gene encoding a thermostable cellulase of glycosyl hydrolase family 12. However, there remains a need for identification and characterization of novel cellulases, with improved properties, such as improved performance under conditions of thermal stress or in the presence of surfactants, increased specific activity, altered substrate cleavage pattern, and/or high level expression *in vitro*.

[98] The development of new and improved cellulase compositions that comprise varying amounts CBH-type, EG-type and BG-type cellulases is of interest for use: (1) in detergent compositions that exhibit enhanced cleaning ability, function as a softening agent and/or improve the feel of cotton fabrics (e.g., "stone washing" or "biopolishing"); (2) in compositions for degrading wood pulp or other biomass into sugars (e.g., for bio-ethanol production); and/or (3) in feed compositions.

IV. MOLECULAR BIOLOGY

[99] In one embodiment this invention provides for the expression of variant CBH I genes under control of a promoter functional in a filamentous fungus. Therefore, this invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and Ausubel *et al.*, eds., *Current Protocols in Molecular Biology* (1994)).

A. Methods for Identifying Homologous CBH1 Genes

[100] The nucleic acid sequence for the wild type *H. jecorina* CBH1 is shown in Figure 1. The invention, in one aspect, encompasses a nucleic acid molecule encoding a CBH1 homolog described herein. The nucleic acid may be a DNA molecule.

[101] Techniques that can be used to isolate CBH I encoding DNA sequences are well known in the art and include, but are not limited to, cDNA and/or genomic library screening with a homologous DNA probe and expression screening with activity assays or antibodies against CBH I. Any of these methods can be found in Sambrook, *et al.* or in CURRENT
5 PROTOCOLS IN MOLECULAR BIOLOGY, F. Ausubel, *et al.*, ed. Greene Publishing and Wiley-Interscience, New York (1987) ("Ausubel").

B. Methods of Mutating CBH I Nucleic Acid Sequences

[102] Any method known in the art that can introduce mutations is contemplated by the present invention.

10 [103] The present invention relates to the expression, purification and/or isolation and use of variant CBH1. These enzymes are preferably prepared by recombinant methods utilizing the *cbh* gene from *H. jecorina*.

[104] After the isolation and cloning of the *cbh1* gene from *H. jecorina*, other methods known in the art, such as site directed mutagenesis, are used to make the substitutions,
15 additions or deletions that correspond to substituted amino acids in the expressed CBH1 variant. Again, site directed mutagenesis and other methods of incorporating amino acid changes in expressed proteins at the DNA level can be found in Sambrook, *et al.* and Ausubel, *et al.*

[105] DNA encoding an amino acid sequence variant of the *H. jecorina* CBH1 is
20 prepared by a variety of methods known in the art. These methods include, but are not limited to, preparation by site-directed (or oligonucleotide-mediated) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared DNA encoding the *H. jecorina* CBH1.

[106] Site-directed mutagenesis is a preferred method for preparing substitution variants.
25 This technique is well known in the art (see, e.g., Carter et al. Nucleic Acids Res. 13:4431-4443 (1985) and Kunkel et al., Proc. Natl. Acad.Sci.USA 82:488 (1987)). Briefly, in carrying out site-directed mutagenesis of DNA, the starting DNA is altered by first hybridizing an oligonucleotide encoding the desired mutation to a single strand of such starting DNA. After hybridization, a DNA polymerase is used to synthesize an entire
30 second strand, using the hybridized oligonucleotide as a primer, and using the single strand of the starting DNA as a template. Thus, the oligonucleotide encoding the desired mutation is incorporated in the resulting double-stranded DNA.

[107] PCR mutagenesis is also suitable for making amino acid sequence variants of the starting polypeptide, i.e., *H. jecorina* CBH1. See Higuchi, in PCR Protocols, pp.177-183
35 (Academic Press, 1990); and Vallette et al., Nuc. Acids Res. 17:723-733 (1989). See,

also, for example Cadwell et al., PCR Methods and Applications, Vol 2, 28-33 (1992). Briefly, when small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs
5 from the template sequence only at the positions where the primers differ from the template.

[108] Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells et al., Gene 34:315-323 (1985). The starting material is the plasmid (or other vector) comprising the starting polypeptide DNA to be mutated. The
10 codon(s) in the starting DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the starting polypeptide DNA. The plasmid DNA is cut at these sites to linearize it. A double-stranded
15 oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures, wherein the two strands of the oligonucleotide are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 5' and 3' ends that are compatible with the
20 ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated DNA sequence.

[109] Alternatively, or additionally, the desired amino acid sequence encoding a variant CBH I can be determined, and a nucleic acid sequence encoding such amino acid sequence variant can be generated synthetically.

[110] The variant CBH I(s) so prepared may be subjected to further modifications, oftentimes depending on the intended use of the cellulase. Such modifications may involve further alteration of the amino acid sequence, fusion to heterologous polypeptide(s) and/or covalent modifications.

V. *cbh1* Nucleic Acids And CBH1 Polypeptides.

A. Variant *cbh*-type Nucleic acids

[111] The nucleic acid sequence for the wild type *H. jecorina* CBH I is shown in Figure 1. The invention encompasses a nucleic acid molecule encoding the variant cellulases described herein. The nucleic acid may be a DNA molecule.

[112] After the isolation and cloning of the CBH I, other methods known in the art, such
35 as site directed mutagenesis, are used to make the substitutions, additions or deletions

- 25 -

that correspond to substituted amino acids in the expressed CBH I variant. Again, site directed mutagenesis and other methods of incorporating amino acid changes in expressed proteins at the DNA level can be found in Sambrook, *et al.* and Ausubel, *et al.*

[113] After DNA sequences that encode the CBH1 variants have been cloned into DNA constructs, the DNA is used to transform microorganisms. The microorganism to be transformed for the purpose of expressing a variant CBH1 according to the present invention may advantageously comprise a strain derived from *Trichoderma sp.* Thus, a preferred mode for preparing variant CBH1 cellulases according to the present invention comprises transforming a *Trichoderma sp.* host cell with a DNA construct comprising at least a fragment of DNA encoding a portion or all of the variant CBH1. The DNA construct will generally be functionally attached to a promoter. The transformed host cell is then grown under conditions so as to express the desired protein. Subsequently, the desired protein product is purified to substantial homogeneity.

[114] However, it may in fact be that the best expression vehicle for a given DNA encoding a variant CBH1 may differ from *H. jecorina*. Thus, it may be that it will be most advantageous to express a protein in a transformation host that bears phylogenetic similarity to the source organism for the variant CBH1. In an alternative embodiment, *Aspergillus niger* can be used as an expression vehicle. For a description of transformation techniques with *A. niger*, see WO 98/31821, the disclosure of which is incorporated by reference in its entirety.

[115] Accordingly, the present description of a *Trichoderma spp.* expression system is provided for illustrative purposes only and as one option for expressing the variant CBH1 of the invention. One of skill in the art, however, may be inclined to express the DNA encoding variant CBH1 in a different host cell if appropriate and it should be understood that the source of the variant CBH1 should be considered in determining the optimal expression host. Additionally, the skilled worker in the field will be capable of selecting the best expression system for a particular gene through routine techniques utilizing the tools available in the art.

B. Variant CBH1 Polypeptides

[116] The amino acid sequence for the wild type *H. jecorina* CBH I is shown in Figure 1. The variant CBH I polypeptides comprises a substitution or deletion at a position corresponding to one or more of residues S8, Q17, G22, T41, N49, S57, N64, A68, A77, N89, S92, N103, A112, S113, E193, S196, M213, L225, T226, P227, T246, D249, R251, Y252, T255, D257, D259, S278, S279, K286, L288, E295, T296, S297, A299, N301, E325, T332, F338, S342, F352, T356, Y371, T380, Y381, V393, R394, S398, V403, S411,

- 26 -

G430, G440, T445, T462, T484, Q487, and P491 in CBH I from *Hypocrea jecorina*.

Furthermore, the variant may further comprises a deletion of residues corresponding to residues 382-393 in CBH I from *Hypocrea jecorina*.

[117] The variant CBH I's of this invention have amino acid sequences that are derived
5 from the amino acid sequence of a precursor CBH I. The amino acid sequence of the
CBH I variant differs from the precursor CBH I amino acid sequence by the substitution,
deletion or insertion of one or more amino acids of the precursor amino acid sequence. In
a preferred embodiment, the precursor CBH I is *Hypocrea jecorina* CBH I. The mature
amino acid sequence of *H. jecorina* CBH I is shown in Figure 1. Thus, this invention is
10 directed to CBH I variants which contain amino acid residues at positions which are
equivalent to the particular identified residue in *H. jecorina* CBH I. A residue (amino acid)
of an CBH I homolog is equivalent to a residue of *Hypocrea jecorina* CBH I if it is either
homologous (*i.e.*, corresponding in position in either primary or tertiary structure) or is
functionally analogous to a specific residue or portion of that residue in *Hypocrea jecorina*
15 CBH I (*i.e.*, having the same or similar functional capacity to combine, react, or interact
chemically or structurally). As used herein, numbering is intended to correspond to that of
the mature CBH I amino acid sequence as illustrated in Figure 1. In addition to locations
within the precursor CBH I, specific residues in the precursor CBH I corresponding to the
amino acid positions that are responsible for instability when the precursor CBH I is under
20 thermal stress are identified herein for substitution or deletion. The amino acid position
number (*e.g.*, +51) refers to the number assigned to the mature *Hypocrea jecorina* CBH I
sequence presented in Figure 1.

[118] The variant CBH1's of this invention have amino acid sequences that are derived
from the amino acid sequence of a precursor *H. jecorina* CBH1. The amino acid
25 sequence of the CBH1 variant differs from the precursor CBH1 amino acid sequence by
the substitution, deletion or insertion of one or more amino acids of the precursor amino
acid sequence. The mature amino acid sequence of *H. jecorina* CBH1 is shown in Figure
1. Thus, this invention is directed to CBH1 variants which contain amino acid residues at
positions which are equivalent to the particular identified residue in *H. jecorina* CBH1. A
30 residue (amino acid) of an CBH1 variant is equivalent to a residue of *Hypocrea jecorina*
CBH1 if it is either homologous (*i.e.*, corresponding in position in either primary or tertiary
structure) or is functionally analogous to a specific residue or portion of that residue in
Hypocrea jecorina CBH1 (*i.e.*, having the same or similar functional capacity to combine,
react, or interact chemically or structurally). As used herein, numbering is intended to
35 correspond to that of the mature CBH1 amino acid sequence as illustrated in Figure 1. In

addition to locations within the precursor CBH1, specific residues in the precursor CBH1 corresponding to the amino acid positions that are responsible for instability when the precursor CBH1 is under thermal stress are identified herein for substitution or deletion. The amino acid position number (e.g., +51) refers to the number assigned to the mature
5 *Hypocrea jecorina* CBH1 sequence presented in Figure 1.

[119] Alignment of amino acid sequences to determine homology is preferably determined by using a "sequence comparison algorithm." Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* **2**:482 (1981), by the homology alignment algorithm
10 of Needleman & Wunsch, *J. Mol. Biol.* **48**:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l Acad. Sci. USA* **85**:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), by visual inspection or MOE by Chemical Computing Group, Montreal
15 Canada.

[120] An example of an algorithm that is suitable for determining sequence similarity is the BLAST algorithm, which is described in Altschul, *et al.*, *J. Mol. Biol.* **215**:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<www.ncbi.nlm.nih.gov>). This algorithm involves
20 first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. These initial neighborhood word hits act as starting points to find longer HSPs containing them. The word hits are expanded in both directions along each of the two sequences being
25 compared for as far as the cumulative alignment score can be increased. Extension of the word hits is stopped when: the cumulative alignment score falls off by the quantity X from a maximum achieved value; the cumulative score goes to zero or below; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word
30 length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* **89**:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M'5, N'-4, and a comparison of both strands.

[121] The BLAST algorithm then performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* **90**:5873-5787
35 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum

probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, an amino acid sequence is considered similar to a protease if the smallest sum probability in a comparison of the test amino acid sequence to a protease amino acid sequence is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[122] Additional specific strategies for modifying stability of CBH1 cellulases are provided below:

[123] (1) Decreasing the entropy of main-chain unfolding may introduce stability to the enzyme. For example, the introduction of proline residues may significantly stabilize the protein by decreasing the entropy of the unfolding (*see, e.g., Watanabe, et al., Eur. J. Biochem.* **226**:277-283 (1994)). Similarly, glycine residues have no β -carbon, and thus have considerably greater backbone conformational freedom than many other residues. Replacement of glycines, preferably with alanines, may reduce the entropy of unfolding and improve stability (*see, e.g., Matthews, et al., Proc. Natl. Acad. Sci. USA* **84**: 6663-6667 (1987)). Additionally, by shortening external loops it may be possible to improve stability. It has been observed that hyperthermophile produced proteins have shorter external loops than their mesophilic homologues (*see, e.g., Russel, et al., Current Opinions in Biotechnology* **6**:370-374 (1995)). The introduction of disulfide bonds may also be effective to stabilize distinct tertiary structures in relation to each other. Thus, the introduction of cysteines at residues accessible to existing cysteines or the introduction of pairs of cysteines that could form disulfide bonds would alter the stability of a CBH1 variant.

[124] (2) Decreasing internal cavities by increasing side-chain hydrophobicity may alter the stability of an enzyme. Reducing the number and volume of internal cavities increases the stability of enzyme by maximizing hydrophobic interactions and reducing packing defects (*see, e.g., Matthews, Ann. Rev. Biochem.* **62**:139-160 (1993); Burley, et al., *Science* **229**:23-29 (1985); Zuber, *Biophys. Chem.* **29**:171-179 (1988); Kellis, et al., *Nature* **333**:784-786 (1988)). It is known that multimeric proteins from thermophiles often have more hydrophobic sub-unit interfaces with greater surface complementarity than their mesophilic counterparts (Russel, et al., *supra*). This principle is believed to be applicable to domain interfaces of monomeric proteins. Specific substitutions that may improve stability by increasing hydrophobicity include lysine to arginine, serine to alanine and threonine to alanine (Russel, et al., *supra*). Modification by substitution to alanine or proline may increase side-chain size with resultant reduction in cavities, better packing

and increased hydrophobicity. Substitutions to reduce the size of the cavity, increase hydrophobicity and improve the complementarity the interfaces between the domains of CBH1 may improve stability of the enzyme. Specifically, modification of the specific residue at these positions with a different residue selected from any of phenylalanine, tryptophan, tyrosine, leucine and isoleucine may improve performance.

5 [125] (3) Balancing charge in rigid secondary structure, *i.e.*, α -helices and β -turns may improve stability. For example, neutralizing partial positive charges on a helix N-terminus with negative charge on aspartic acid may improve stability of the structure (*see, e.g., Eriksson, et al., Science 255:178-183 (1992)*). Similarly, neutralizing partial negative charges on helix C-terminus with positive charge may improve stability. Removing positive charge from interacting with peptide N-terminus in β -turns should be effective in conferring tertiary structure stability. Substitution with a non-positively charged residue could remove an unfavorable positive charge from interacting with an amide nitrogen present in a turn.

15 [126] (4) Introducing salt bridges and hydrogen bonds to stabilize tertiary structures may be effective. For example, ion pair interactions, *e.g.*, between aspartic acid or glutamic acid and lysine, arginine or histidine, may introduce strong stabilizing effects and may be used to attach different tertiary structure elements with a resultant improvement in thermostability. Additionally, increases in the number of charged residue/non-charged residue hydrogen bonds, and the number of hydrogen-bonds generally, may improve thermostability (*see, e.g., Tanner, et al., Biochemistry 35:2597-2609 (1996)*). Substitution with aspartic acid, asparagine, glutamic acid or glutamine may introduce a hydrogen bond with a backbone amide. Substitution with arginine may improve a salt bridge and introduce an H-bond into a backbone carbonyl.

25 [127] (5) Avoiding thermolabile residues in general may increase thermal stability. For example, asparagine and glutamine are susceptible to deamidation and cysteine is susceptible to oxidation at high temperatures. Reducing the number of these residues in sensitive positions may result in improved thermostability (*Russel, et al., supra*). Substitution or deletion by any residue other than glutamine or cysteine may increase stability by avoidance of a thermolabile residue.

30 [128] (6) Stabilization or destabilization of binding of a ligand that confers modified stability to CBH1 variants. For example, a component of the matrix in which the CBH1 variants of this invention are used may bind to a specific surfactant/thermal sensitivity site of the CBH1 variant. By modifying the site through substitution, binding of the component to the variant may be strengthened or diminished. For example, a non-aromatic residue in

the binding crevice of CBH1 may be substituted with phenylalanine or tyrosine to introduce aromatic side-chain stabilization where interaction of the cellulose substrate may interact favorably with the benzyl rings, increasing the stability of the CBH1 variant.

[129] (7) Increasing the electronegativity of any of the surfactant/ thermal sensitivity ligands may improve stability under surfactant or thermal stress. For example, substitution with phenylalanine or tyrosine may increase the electronegativity of D (aspartate) residues by improving shielding from solvent, thereby improving stability.

C. Anti-CBH Antibodies

[130] The present invention further provides anti-CBH antibodies. The antibodies may be polyclonal, monoclonal, humanized, bispecific or heteroconjugate antibodies.

[131] Methods of preparing polyclonal antibodies are known to the skilled artisan. The immunizing agent may be an CBH polypeptide or a fusion protein thereof. It may be useful to conjugate the antigen to a protein known to be immunogenic in the mammal being immunized. The immunization protocol may be determined by one skilled in the art based on standard protocols or routine experimentation.

[132] Alternatively, the anti-CBH antibodies may be monoclonal antibodies. Monoclonal antibodies may be produced by cells immunized in an animal or using recombinant DNA methods. (See, e.g., Kohler *et al.*, *Nature*, vol. 256, pp. 495-499, August 7, 1975; U.S. Patent No. 4,816,567).

[133] An anti-CBH antibody of the invention may further comprise a humanized or human antibody. The term "humanized antibody" refers to humanized forms of non-human (e.g., murine) antibodies that are chimeric antibodies, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding partial sequences of antibodies) which contain some portion of the sequence derived from non-human antibody. Methods for humanizing non-human antibodies are well known in the art, as further detailed in Jones *et al.*, *Nature* 321:522-525, 1986; Riechmann *et al.*, *Nature*, vol. 332, pp. 323-327, 1988; and Verhoeyen *et al.*, *Science*, vol. 239, pp. 1534-1536, 1988. Methods for producing human antibodies are also known in the art. See, e.g., Jakobovits, A, *et al.*, *Annals New York Academy of Sciences*, 764:525-535, 1995 and Jakobovits, A, *Curr Opin Biotechnol* 6(5):561-6, 1995.

VI. Expression Of Recombinant CBH1 Variants

[134] The methods of the invention rely on the use cells to express variant CBH I, with no particular method of CBH I expression required.

- 31 -

[135] The invention provides host cells which have been transduced, transformed or transfected with an expression vector comprising a variant CBH-encoding nucleic acid sequence. The culture conditions, such as temperature, pH and the like, are those previously used for the parental host cell prior to transduction, transformation or
5 transfected and will be apparent to those skilled in the art.

[136] In one approach, a filamentous fungal cell or yeast cell is transfected with an expression vector having a promoter or biologically active promoter fragment or one or more (*e.g.*, a series) of enhancers which functions in the host cell line, operably linked to a DNA segment encoding CBH, such that CBH is expressed in the cell line.

10 A. Nucleic Acid Constructs/Expression Vectors.

[137] Natural or synthetic polynucleotide fragments encoding CBH I ("CBH I-encoding nucleic acid sequences") may be incorporated into heterologous nucleic acid constructs or vectors, capable of introduction into, and replication in, a filamentous fungal or yeast cell. The vectors and methods disclosed herein are suitable for use in host cells for the
15 expression of CBH I. Any vector may be used as long as it is replicable and viable in the cells into which it is introduced. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. Cloning and expression vectors are also described in Sambrook *et al.*, 1989, Ausubel FM *et al.*, 1989, and Strathern *et al.*, *The Molecular Biology of the Yeast Saccharomyces*, 1981, each of which is
20 expressly incorporated by reference herein. Appropriate expression vectors for fungi are described in van den Hondel, C.A.M.J.J. et al. (1991) In: Bennett, J.W. and Lasure, L.L. (eds.) *More Gene Manipulations in Fungi*. Academic Press, pp. 396-428. The appropriate DNA sequence may be inserted into a plasmid or vector (collectively referred to herein as "vectors") by a variety of procedures. In general, the DNA sequence is inserted into an
25 appropriate restriction endonuclease site(s) by standard procedures. Such procedures and related sub-cloning procedures are deemed to be within the scope of knowledge of those skilled in the art.

[138] Recombinant filamentous fungi comprising the coding sequence for variant CBH I may be produced by introducing a heterologous nucleic acid construct comprising the
30 variant CBH I coding sequence into the cells of a selected strain of the filamentous fungi.

[139] Once the desired form of a variant *cbh* nucleic acid sequence is obtained, it may be modified in a variety of ways. Where the sequence involves non-coding flanking regions, the flanking regions may be subjected to resection, mutagenesis, etc. Thus, transitions, transversions, deletions, and insertions may be performed on the naturally
35 occurring sequence.

[140] A selected variant *cbh* coding sequence may be inserted into a suitable vector according to well-known recombinant techniques and used to transform filamentous fungi capable of CBH I expression. Due to the inherent degeneracy of the genetic code, other nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence may be used to clone and express variant CBH I. Therefore it is appreciated that such substitutions in the coding region fall within the sequence variants covered by the present invention. Any and all of these sequence variants can be utilized in the same way as described herein for a parent CBH I-encoding nucleic acid sequence.

[141] The present invention also includes recombinant nucleic acid constructs comprising one or more of the variant CBH I-encoding nucleic acid sequences as described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation.

[142] Heterologous nucleic acid constructs may include the coding sequence for variant *cbh*: (i) in isolation; (ii) in combination with additional coding sequences; such as fusion protein or signal peptide coding sequences, where the *cbh* coding sequence is the dominant coding sequence; (iii) in combination with non-coding sequences, such as introns and control elements, such as promoter and terminator elements or 5' and/or 3' untranslated regions, effective for expression of the coding sequence in a suitable host; and/or (iv) in a vector or host environment in which the *cbh* coding sequence is a heterologous gene.

[143] In one aspect of the present invention, a heterologous nucleic acid construct is employed to transfer a variant CBH I-encoding nucleic acid sequence into a cell *in vitro*, with established filamentous fungal and yeast lines preferred. For long-term, production of variant CBH I, stable expression is preferred. It follows that any method effective to generate stable transformants may be used in practicing the invention.

[144] Appropriate vectors are typically equipped with a selectable marker-encoding nucleic acid sequence, insertion sites, and suitable control elements, such as promoter and termination sequences. The vector may comprise regulatory sequences, including, for example, non-coding sequences, such as introns and control elements, *i.e.*, promoter and terminator elements or 5' and/or 3' untranslated regions, effective for expression of the coding sequence in host cells (and/or in a vector or host cell environment in which a modified soluble protein antigen coding sequence is not normally expressed), operably linked to the coding sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, many of which are commercially available and/or are described in Sambrook, *et al.*, (*supra*).

- 33 -

[145] Exemplary promoters include both constitutive promoters and inducible promoters, examples of which include a CMV promoter, an SV40 early promoter, an RSV promoter, an EF-1 α promoter, a promoter containing the tet responsive element (TRE) in the tet-on or tet-off system as described (ClonTech and BASF), the beta actin promoter and the metallothionine promoter that can upregulated by addition of certain metal salts. A promoter sequence is a DNA sequence which is recognized by the particular filamentous fungus for expression purposes. It is operably linked to DNA sequence encoding a variant CBH I polypeptide. Such linkage comprises positioning of the promoter with respect to the initiation codon of the DNA sequence encoding the variant CBH I polypeptide in the disclosed expression vectors. The promoter sequence contains transcription and translation control sequence which mediate the expression of the variant CBH I polypeptide. Examples include the promoters from the *Aspergillus niger*, *A. awamori* or *A. oryzae* glucoamylase, alpha-amylase, or alpha-glucosidase encoding genes; the *A. nidulans* *gpdA* or *trpC* Genes; the *Neurospora crassa* *cbh1* or *trp1* genes; the *A. niger* or *Rhizomucor miehei* aspartic proteinase encoding genes; the *H. jecorina* (*T. reesei*) *cbh1*, *cbh2*, *egl1*, *egl2*, or other cellulase encoding genes.

[146] The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well known in the art. Typical selectable marker genes include *argB* from *A. nidulans* or *T. reesei*, *amdS* from *A. nidulans*, *pyr4* from *Neurospora crassa* or *T. reesei*, *pyrG* from *Aspergillus niger* or *A. nidulans*. Additional exemplary selectable markers include, but are not limited to *trpc*, *trp1*, *olic31*, *niaD* or *leu2*, which are included in heterologous nucleic acid constructs used to transform a mutant strain such as *trp-*, *pyr-*, *leu-* and the like.

[147] Such selectable markers confer to transformants the ability to utilize a metabolite that is usually not metabolized by the filamentous fungi. For example, the *amdS* gene from *H. jecorina* which encodes the enzyme acetamidase that allows transformant cells to grow on acetamide as a nitrogen source. The selectable marker (e.g. *pyrG*) may restore the ability of an auxotrophic mutant strain to grow on a selective minimal medium or the selectable marker (e.g. *olic31*) may confer to transformants the ability to grow in the presence of an inhibitory drug or antibiotic.

[148] The selectable marker coding sequence is cloned into any suitable plasmid using methods generally employed in the art. Exemplary plasmids include pUC18, pBR322, pRAX and pUC100. The pRAX plasmid contains AMA1 sequences from *A. nidulans*, which make it possible to replicate in *A. niger*.

- 34 -

[149] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, Sambrook *et al.*, 1989; Freshney, *Animal Cell Culture*, 1987; Ausubel, *et al.*, 1993; and Coligan *et al.*, *Current Protocols in Immunology*, 1991.

B. Host Cells and Culture Conditions For CBH I Production

(i) Filamentous Fungi

[150] Thus, the present invention provides filamentous fungi comprising cells which have been modified, selected and cultured in a manner effective to result in variant CBH I production or expression relative to the corresponding non-transformed parental fungi.

[151] Examples of species of parental filamentous fungi that may be treated and/or modified for variant CBH I expression include, but are not limited to *Trichoderma*, *e.g.*, *Trichoderma reesei*, *Trichoderma longibrachiatum*, *Trichoderma viride*, *Trichoderma koningii*; *Penicillium sp.*, *Humicola sp.*, including *Humicola insolens*; *Aspergillus sp.*, *Chrysosporium sp.*, *Fusarium sp.*, *Hypocrea sp.*, and *Emericella sp.*

[152] CBH I expressing cells are cultured under conditions typically employed to culture the parental fungal line. Generally, cells are cultured in a standard medium containing physiological salts and nutrients, such as described in Pourquie, J. *et al.*, *Biochemistry and Genetics of Cellulose Degradation*, eds. Aubert, J. P. *et al.*, Academic Press, pp. 71-86, 1988 and Ilmen, M. *et al.*, *Appl. Environ. Microbiol.* 63:1298-1306, 1997. Culture conditions are also standard, *e.g.*, cultures are incubated at 28°C in shaker cultures or fermenters until desired levels of CBH I expression are achieved.

[153] Preferred culture conditions for a given filamentous fungus may be found in the scientific literature and/or from the source of the fungi such as the American Type Culture Collection (ATCC; "<http://www.atcc.org/>"). After fungal growth has been established, the cells are exposed to conditions effective to cause or permit the expression of variant CBH I.

[154] In cases where a CBH I coding sequence is under the control of an inducible promoter, the inducing agent, *e.g.*, a sugar, metal salt or antibiotics, is added to the medium at a concentration effective to induce CBH I expression.

[155] In one embodiment, the strain comprises *Aspergillus niger*, which is a useful strain for obtaining overexpressed protein. For example *A. niger* var *awamori* dgr246 is known to secrete elevated amounts of secreted cellulases (Goedegebuur *et al.*, *Curr. Genet* (2002) 41: 89-98). Other strains of *Aspergillus niger* var *awamori* such as GCDAP3,

- 35 -

GCDAP4 and GAP3-4 are known Ward et al (Ward, M, Wilson, L.J. and Kodama, K.H., 1993, Appl. Microbiol. Biotechnol. 39:738-743).

[156] In another embodiment, the strain comprises *Trichoderma reesei*, which is a useful strain for obtaining overexpressed protein. For example, RL-P37, described by Sheir-
5 Neiss, et al., Appl. Microbiol. Biotechnol. 20:46-53 (1984) is known to secrete elevated amounts of cellulase enzymes. Functional equivalents of RL-P37 include *Trichoderma reesei* strain RUT-C30 (ATCC No. 56765) and strain QM9414 (ATCC No. 26921). It is contemplated that these strains would also be useful in overexpressing variant CBH1.

[157] Where it is desired to obtain the variant CBH I in the absence of potentially
10 detrimental native cellulolytic activity, it is useful to obtain a *Trichoderma* host cell strain which has had one or more cellulase genes deleted prior to introduction of a DNA construct or plasmid containing the DNA fragment encoding the variant CBH I. Such strains may be prepared by the method disclosed in U.S. Patent No. 5,246,853 and WO 92/06209, which disclosures are hereby incorporated by reference. By expressing a
15 variant CBH I cellulase in a host microorganism that is missing one or more cellulase genes, the identification and subsequent purification procedures are simplified. Any gene from *Trichoderma sp.* which has been cloned can be deleted, for example, the *cbh1*, *cbh2*, *egl1*, and *egl2* genes as well as those encoding EG III and/or EGV protein (see e.g., U.S. Patent No. 5,475,101 and WO 94/28117, respectively).

[158] Gene deletion may be accomplished by inserting a form of the desired gene to be
20 deleted or disrupted into a plasmid by methods known in the art. The deletion plasmid is then cut at an appropriate restriction enzyme site(s), internal to the desired gene coding region, and the gene coding sequence or part thereof replaced with a selectable marker. Flanking DNA sequences from the locus of the gene to be deleted or disrupted, preferably
25 between about 0.5 to 2.0 kb, remain on either side of the selectable marker gene. An appropriate deletion plasmid will generally have unique restriction enzyme sites present therein to enable the fragment containing the deleted gene, including flanking DNA sequences, and the selectable marker gene to be removed as a single linear piece.

[159] A selectable marker must be chosen so as to enable detection of the transformed
30 microorganism. Any selectable marker gene that is expressed in the selected microorganism will be suitable. For example, with *Aspergillus sp.*, the selectable marker is chosen so that the presence of the selectable marker in the transformants will not significantly affect the properties thereof. Such a selectable marker may be a gene that encodes an assayable product. For example, a functional copy of a *Aspergillus sp.* gene

- 36 -

may be used which if lacking in the host strain results in the host strain displaying an auxotrophic phenotype. Similarly, selectable markers exist for *Trichoderma sp.*

[160] In one embodiment, a *pyrG*⁻ derivative strain of *Aspergillus sp.* is transformed with

a functional *pyrG* gene, which thus provides a selectable marker for transformation. A

5 *pyrG*⁻ derivative strain may be obtained by selection of *Aspergillus sp.* strains that are resistant to fluoroorotic acid (FOA). The *pyrG* gene encodes orotidine-5'-monophosphate decarboxylase, an enzyme required for the biosynthesis of uridine. Strains with an intact

pyrG gene grow in a medium lacking uridine but are sensitive to fluoroorotic acid. It is possible to select *pyrG*⁻ derivative strains that lack a functional orotidine monophosphate

10 decarboxylase enzyme and require uridine for growth by selecting for FOA resistance. Using the FOA selection technique it is also possible to obtain uridine-requiring strains

which lack a functional orotate pyrophosphoribosyl transferase. It is possible to transform

these cells with a functional copy of the gene encoding this enzyme (Berges & Barreau,

Curr. Genet. 19:359-365 (1991), and van Hartingsveldt et al., (1986) Development of a

15 homologous transformation system for *Aspergillus niger* based on the *pyrG* gene. *Mol.*

Gen. Genet. 206:71-75). Selection of derivative strains is easily performed using the FOA

resistance technique referred to above, and thus, the *pyrG* gene is preferably employed as a selectable marker.

[161] In a second embodiment, a *pyr4*⁻ derivative strain of *Hyprocrea sp.* (*Hyprocrea sp.*

20 *Trichoderma sp.*) is transformed with a functional *pyr4* gene, which thus provides a selectable marker for transformation. A *pyr4*⁻ derivative strain may be obtained by

selection of *Hyprocrea sp.* (*Trichoderma sp.*) strains that are resistant to fluoroorotic acid (FOA). The *pyr4* gene encodes orotidine-5'-monophosphate decarboxylase, an enzyme required for the biosynthesis of uridine. Strains with an intact *pyr4* gene grow in a medium

25 lacking uridine but are sensitive to fluoroorotic acid. It is possible to select *pyr4*⁻ derivative strains that lack a functional orotidine monophosphate decarboxylase enzyme and require

uridine for growth by selecting for FOA resistance. Using the FOA selection technique it is also possible to obtain uridine-requiring strains which lack a functional orotate

pyrophosphoribosyl transferase. It is possible to transform these cells with a functional

30 copy of the gene encoding this enzyme (Berges & Barreau, *Curr. Genet.* 19:359-365

(1991)). Selection of derivative strains is easily performed using the FOA resistance

technique referred to above, and thus, the *pyr4* gene is preferably employed as a selectable marker.

[162] To transform *pyrG*⁻ *Aspergillus sp.* or *pyr4*⁻ *Hyprocrea sp.* (*Trichoderma sp.*) so as

35 to be lacking in the ability to express one or more cellulase genes, a single DNA fragment

comprising a disrupted or deleted cellulase gene is then isolated from the deletion plasmid and used to transform an appropriate *pyr⁻ Aspergillus* or *pyr⁻ Trichoderma* host.

Transformants are then identified and selected based on their ability to express the *pyrG* or *pyr4*, respectively, gene product and thus complement the uridine auxotrophy of the
5 host strain. Southern blot analysis is then carried out on the resultant transformants to identify and confirm a double crossover integration event that replaces part or all of the coding region of the genomic copy of the gene to be deleted with the appropriate *pyr* selectable markers.

[163] Although the specific plasmid vectors described above relate to preparation of *pyr⁻*
10 transformants, the present invention is not limited to these vectors. Various genes can be deleted and replaced in the *Aspergillus sp.* or *Hyprocrea sp.* (*Trichoderma sp.*) strain using the above techniques. In addition, any available selectable markers can be used, as discussed above. In fact, any host, e.g., *Aspergillus sp.* or *Hyprocrea sp.*, gene that has been cloned, and thus identified, can be deleted from the genome using the above-
15 described strategy.

[164] As stated above, the host strains used may be derivatives of *Hyprocrea sp.* (*Trichoderma sp.*) that lack or have a nonfunctional gene or genes corresponding to the selectable marker chosen. For example, if the selectable marker of *pyrG* is chosen for *Aspergillus sp.*, then a specific *pyrG⁻* derivative strain is used as a recipient in the
20 transformation procedure. Also, for example, if the selectable marker of *pyr4* is chosen for a *Hyprocrea sp.*, then a specific *pyr4⁻* derivative strain is used as a recipient in the transformation procedure. Similarly, selectable markers comprising *Hyprocrea sp.* (*Trichoderma sp.*) genes equivalent to the *Aspergillus nidulans* genes *amdS*, *argB*, *trpC*, *niaD* may be used. The corresponding recipient strain must therefore be a derivative
25 strain such as *argB⁻*, *trpC⁻*, *niaD⁻*, respectively.

[165] DNA encoding the CBH I variant is then prepared for insertion into an appropriate microorganism. According to the present invention, DNA encoding a CBH I variant comprises the DNA necessary to encode for a protein that has functional cellulolytic activity. The DNA fragment encoding the CBH I variant may be functionally attached to a
30 fungal promoter sequence, for example, the promoter of the *glaA* gene in *Aspergillus* or the promoter of the *cbh1* or *egl1* genes in *Trichoderma*.

[166] It is also contemplated that more than one copy of DNA encoding a CBH I variant may be recombined into the strain to facilitate overexpression. The DNA encoding the CBH I variant may be prepared by the construction of an expression vector carrying the
35 DNA encoding the variant. The expression vector carrying the inserted DNA fragment

- 38 -

encoding the CBH I variant may be any vector which is capable of replicating autonomously in a given host organism or of integrating into the DNA of the host, typically a plasmid. In preferred embodiments two types of expression vectors for obtaining expression of genes are contemplated. The first contains DNA sequences in which the promoter, gene-coding region, and terminator sequence all originate from the gene to be expressed. Gene truncation may be obtained where desired by deleting undesired DNA sequences (*e.g.*, coding for unwanted domains) to leave the domain to be expressed under control of its own transcriptional and translational regulatory sequences. A selectable marker may also be contained on the vector allowing the selection for integration into the host of multiple copies of the novel gene sequences.

[167] The second type of expression vector is preassembled and contains sequences required for high-level transcription and a selectable marker. It is contemplated that the coding region for a gene or part thereof can be inserted into this general-purpose expression vector such that it is under the transcriptional control of the expression cassettes promoter and terminator sequences.

[168] For example, in *Aspergillus*, pRAX is such a general-purpose expression vector. Genes or part thereof can be inserted downstream of the strong *glaA* promoter.

[169] For example, in *Hypocrea*, pTEX is such a general-purpose expression vector. Genes or part thereof can be inserted downstream of the strong *cbh1* promoter.

[170] In the vector, the DNA sequence encoding the CBH I variant of the present invention should be operably linked to transcriptional and translational sequences, *i.e.*, a suitable promoter sequence and signal sequence in reading frame to the structural gene. The promoter may be any DNA sequence that shows transcriptional activity in the host cell and may be derived from genes encoding proteins either homologous or heterologous to the host cell. An optional signal peptide provides for extracellular production of the CBH I variant. The DNA encoding the signal sequence is preferably that which is naturally associated with the gene to be expressed, however the signal sequence from any suitable source, for example an exo-cellobiohydrolase or endoglucanase from *Trichoderma*, is contemplated in the present invention.

[171] The procedures used to ligate the DNA sequences coding for the variant CBH I of the present invention with the promoter, and insertion into suitable vectors are well known in the art.

[172] The DNA vector or construct described above may be introduced in the host cell in accordance with known techniques such as transformation, transfection, microinjection, microporation, biolistic bombardment and the like.

[173] In the preferred transformation technique, it must be taken into account that the permeability of the cell wall to DNA in *Hyprocrea sp. (Trichoderma sp.)* is very low. Accordingly, uptake of the desired DNA sequence, gene or gene fragment is at best minimal. There are a number of methods to increase the permeability of the *Hyprocrea*
5 *sp. (Trichoderma sp.)* cell wall in the derivative strain (*i.e.*, lacking a functional gene corresponding to the used selectable marker) prior to the transformation process.

[174] The preferred method in the present invention to prepare *Aspergillus sp.* or *Hyprocrea sp. (Trichoderma sp.)* for transformation involves the preparation of protoplasts from fungal mycelium. See Campbell *et al.* Improved transformation efficiency of *A.niger*
10 using homologous *niaD* gene for nitrate reductase. Curr. Genet. 16:53-56; 1989. The mycelium can be obtained from germinated vegetative spores. The mycelium is treated with an enzyme that digests the cell wall resulting in protoplasts. The protoplasts are then protected by the presence of an osmotic stabilizer in the suspending medium. These stabilizers include sorbitol, mannitol, potassium chloride, magnesium sulfate and the like.
15 Usually the concentration of these stabilizers varies between 0.8 M and 1.2 M. It is preferable to use about a 1.2 M solution of sorbitol in the suspension medium.

[175] Uptake of the DNA into the host strain, (*Aspergillus sp.* or *Hyprocrea sp. (Trichoderma sp.)*), is dependent upon the calcium ion concentration. Generally between about 10 mM CaCl_2 and 50 mM CaCl_2 is used in an uptake solution. Besides the need for
20 the calcium ion in the uptake solution, other items generally included are a buffering system such as TE buffer (10 Mm Tris, pH 7.4; 1 mM EDTA) or 10 mM MOPS, pH 6.0 buffer (morpholinepropanesulfonic acid) and polyethylene glycol (PEG). It is believed that the polyethylene glycol acts to fuse the cell membranes thus permitting the contents of the medium to be delivered into the cytoplasm of the host cell, by way of example either
25 *Aspergillus sp.* or *Hyprocrea sp.* strain, and the plasmid DNA is transferred to the nucleus. This fusion frequently leaves multiple copies of the plasmid DNA tenderly integrated into the host chromosome.

[176] Usually a suspension containing the *Aspergillus sp.* protoplasts or cells that have been subjected to a permeability treatment at a density of 10^5 to 10^6 /mL, preferably 2×10^5 /mL are used in transformation. Similarly, a suspension containing the *Hyprocrea sp. (Trichoderma sp.)* protoplasts or cells that have been subjected to a permeability
30 treatment at a density of 10^8 to 10^9 /mL, preferably 2×10^8 /mL are used in transformation. A volume of 100 μL of these protoplasts or cells in an appropriate solution (*e.g.*, 1.2 M sorbitol; 50 mM CaCl_2) are mixed with the desired DNA. Generally a high concentration of
35 PEG is added to the uptake solution. From 0.1 to 1 volume of 25% PEG 4000 can be

- 40 -

added to the protoplast suspension. However, it is preferable to add about 0.25 volumes to the protoplast suspension. Additives such as dimethyl sulfoxide, heparin, spermidine, potassium chloride and the like may also be added to the uptake solution and aid in transformation.

5 [177] Generally, the mixture is then incubated at approximately 0°C for a period of between 10 to 30 minutes. Additional PEG is then added to the mixture to further enhance the uptake of the desired gene or DNA sequence. The 25% PEG 4000 is generally added in volumes of 5 to 15 times the volume of the transformation mixture; however, greater and lesser volumes may be suitable. The 25% PEG 4000 is preferably
10 about 10 times the volume of the transformation mixture. After the PEG is added, the transformation mixture is then incubated either at room temperature or on ice before the addition of a sorbitol and CaCl₂ solution. The protoplast suspension is then further added to molten aliquots of a growth medium. This growth medium permits the growth of transformants only. Any growth medium can be used in the present invention that is
15 suitable to grow the desired transformants. However, if *Pyr*⁺ transformants are being selected it is preferable to use a growth medium that contains no uridine. The subsequent colonies are transferred and purified on a growth medium depleted of uridine.

[178] At this stage, stable transformants may be distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a
20 smooth, rather than ragged outline on solid culture medium lacking uridine. Additionally, in some cases a further test of stability may be made by growing the transformants on solid non-selective medium (*i.e.* containing uridine), harvesting spores from this culture medium and determining the percentage of these spores which will subsequently germinate and grow on selective medium lacking uridine.

25 [179] In a particular embodiment of the above method, the CBH I variant(s) are recovered in active form from the host cell after growth in liquid media either as a result of the appropriate post translational processing of the CBH I variant.

(ii) Yeast

[180] The present invention also contemplates the use of yeast as a host cell for CBH I
30 production. Several other genes encoding hydrolytic enzymes have been expressed in various strains of the yeast *S. cerevisiae*. These include sequences encoding for two endoglucanases (Penttila *et al.*, Yeast vol. 3, pp 175-185, 1987), two cellobiohydrolases (Penttila *et al.*, Gene, 63: 103-112, 1988) and one beta-glucosidase from *Trichoderma reesei* (Cummings and Fowler, Curr. Genet. 29:227-233, 1996), a xylanase from
35 *Aureobasidium pullulans* (Li and Ljungdahl, Appl. Environ. Microbiol. 62, no. 1, pp. 209-

- 41 -

213, 1996), an alpha-amylase from wheat (Rothstein *et al.*, Gene 55:353-356, 1987), etc. In addition, a cellulase gene cassette encoding the *Butyrivibrio fibrisolvens* endo- [beta] - 1,4-glucanase (END1), *Phanerochaete chrysosporium* cellobiohydrolase (CBH1), the *Ruminococcus flavefaciens* cellodextrinase (CEL1) and the *Endomyces fibrilizer*
5 cellobiase (Bgl1) was successfully expressed in a laboratory strain of *S. cerevisiae* (Van Rensburg *et al.*, Yeast, vol. 14, pp. 67-76, 1998).

C. Introduction of an CBH I-Encoding Nucleic Acid Sequence into Host Cells.

[181] The invention further provides cells and cell compositions which have been
10 genetically modified to comprise an exogenously provided variant CBH I -encoding nucleic acid sequence. A parental cell or cell line may be genetically modified (*i.e.*, transduced, transformed or transfected) with a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc, as further described above.

15 [182] The methods of transformation of the present invention may result in the stable integration of all or part of the transformation vector into the genome of the filamentous fungus. However, transformation resulting in the maintenance of a self-replicating extra-chromosomal transformation vector is also contemplated.

[183] Many standard transfection methods can be used to produce *Trichoderma reesei*
20 cell lines that express large quantities of the heterologous protein. Some of the published methods for the introduction of DNA constructs into cellulase-producing strains of *Trichoderma* include Lorito, Hayes, DiPietro and Harman, 1993, Curr. Genet. 24: 349-356; Goldman, VanMontagu and Herrera-Estrella, 1990, Curr. Genet. 17:169-174; Penttila, Nevalainen, Ratto, Salminen and Knowles, 1987, Gene 6: 155-164, for *Aspergillus* Yelton,
25 Hamer and Timberlake, 1984, Proc. Natl. Acad. Sci. USA 81: 1470-1474, for *Fusarium* Bajar, Podila and Kolattukudy, 1991, Proc. Natl. Acad. Sci. USA 88: 8202-8212, for *Streptomyces* Hopwood *et al.*, 1985, The John Innes Foundation, Norwich, UK and for *Bacillus* Brigidi, DeRossi, Bertarini, Riccardi and Matteuzzi, 1990, FEMS Microbiol. Lett. 55: 135-138).

30 [184] Other methods for introducing a heterologous nucleic acid construct (expression vector) into filamentous fungi (*e.g.*, *H. jecorina*) include, but are not limited to the use of a particle or gene gun, permeabilization of filamentous fungi cells walls prior to the transformation process (*e.g.*, by use of high concentrations of alkali, *e.g.*, 0.05 M to 0.4 M CaCl₂ or lithium acetate), protoplast fusion or agrobacterium mediated transformation. An
35 exemplary method for transformation of filamentous fungi by treatment of protoplasts or

spheroplasts with polyethylene glycol and CaCl₂ is described in Campbell, E.I. et al., *Curr. Genet.* 16:53-56, 1989 and Penttila, M. et al., *Gene*, 63:11-22, 1988.

[185] Any of the well-known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, biolistics, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see, e.g., Sambrook *et al.*, *supra*). Also of use is the *Agrobacterium*-mediated transfection method described in U.S. Patent No. 6,255,115. It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing the heterologous gene.

[186] In addition, heterologous nucleic acid constructs comprising a variant CBH I-encoding nucleic acid sequence can be transcribed *in vitro*, and the resulting RNA introduced into the host cell by well-known methods, e.g., by injection.

[187] The invention further includes novel and useful transformants of filamentous fungi such as *H. jecorina* and *A. niger* for use in producing fungal cellulase compositions. The invention includes transformants of filamentous fungi especially fungi comprising the variant CBH I coding sequence, or deletion of the endogenous *cbh* coding sequence.

[188] Following introduction of a heterologous nucleic acid construct comprising the coding sequence for a variant *cbh 1*, the genetically modified cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying expression of a variant CBH I-encoding nucleic acid sequence. The culture conditions, such as temperature, pH and the like, are those previously used for the host cell selected for expression, and will be apparent to those skilled in the art.

[189] The progeny of cells into which such heterologous nucleic acid constructs have been introduced are generally considered to comprise the variant CBH I-encoding nucleic acid sequence found in the heterologous nucleic acid construct.

[190] The invention further includes novel and useful transformants of filamentous fungi such as *H. jecorina* for use in producing fungal cellulase compositions. The invention includes transformants of filamentous fungi especially fungi comprising the variant *cbh 1* coding sequence, or deletion of the endogenous *cbh* coding sequence.

[191] Stable transformants of filamentous fungi can generally be distinguished from unstable transformants by their faster growth rate and the formation of circular colonies with a smooth rather than ragged outline on solid culture medium. Additionally, in some

cases, a further test of stability can be made by growing the transformants on solid non-selective medium, harvesting the spores from this culture medium and determining the percentage of these spores which will subsequently germinate and grow on selective medium.

5 **VII. Analysis For CBH1 Nucleic Acid Coding Sequences and/or Protein Expression.**

[192] In order to evaluate the expression of a variant CBH I by a cell line that has been transformed with a variant CBH I-encoding nucleic acid construct, assays can be carried out at the protein level, the RNA level or by use of functional bioassays particular to
10 cellobiohydrolase activity and/or production.

[193] In one exemplary application of the variant *cbh 1* nucleic acid and protein sequences described herein, a genetically modified strain of filamentous fungi, *e.g.*, *Trichoderma reesei*, is engineered to produce an increased amount of CBH I. Such genetically modified filamentous fungi would be useful to produce a cellulase product with
15 greater increased cellulolytic capacity. In one approach, this is accomplished by introducing the coding sequence for *cbh 1* into a suitable host, *e.g.*, a filamentous fungi such as *Aspergillus niger*.

[194] Accordingly, the invention includes methods for expressing variant CBH I in a filamentous fungus or other suitable host by introducing an expression vector containing
20 the DNA sequence encoding variant CBH I into cells of the filamentous fungus or other suitable host.

[195] In another aspect, the invention includes methods for modifying the expression of CBH I in a filamentous fungus or other suitable host. Such modification includes a decrease or elimination in expression of the endogenous CBH.

25 [196] In general, assays employed to analyze the expression of variant CBH I include, Northern blotting, dot blotting (DNA or RNA analysis), RT-PCR (reverse transcriptase polymerase chain reaction), or *in situ* hybridization, using an appropriately labeled probe (based on the nucleic acid coding sequence) and conventional Southern blotting and autoradiography.

30 [197] In addition, the production and/or expression of variant CBH I may be measured in a sample directly, for example, by assays for cellobiohydrolase activity, expression and/or production. Such assays are described, for example, in Becker et al., *Biochem J.* (2001) 356:19-30 and Mitsuishi et al., *FEBS* (1990) 275:135-138, each of which is expressly incorporated by reference herein. The ability of CBH I to hydrolyze isolated soluble and
35 insoluble substrates can be measured using assays described in Srisodsuk et al., J.

Biotech. (1997) 57:49-57 and Nidetzky and Claeysens Biotech. Bioeng. (1994) 44:961-966. Substrates useful for assaying cellobiohydrolase, endoglucanase or β -glucosidase activities include crystalline cellulose, filter paper, phosphoric acid swollen cellulose, cellooligosaccharides, methylumbelliferyl lactoside, methylumbelliferyl cellobioside, 5 orthonitrophenyl lactoside, paranitrophenyl lactoside, orthonitrophenyl cellobioside, paranitrophenyl cellobioside.

[198] In addition, protein expression, may be evaluated by immunological methods, such as immunohistochemical staining of cells, tissue sections or immunoassay of tissue culture medium, *e.g.*, by Western blot or ELISA. Such immunoassays can be used to 10 qualitatively and quantitatively evaluate expression of a CBH I variant. The details of such methods are known to those of skill in the art and many reagents for practicing such methods are commercially available.

[199] A purified form of a variant CBH I may be used to produce either monoclonal or polyclonal antibodies specific to the expressed protein for use in various immunoassays. 15 (See, *e.g.*, Hu *et al.*, Mol Cell Biol. vol.11, no. 11, pp. 5792-5799, 1991). Exemplary assays include ELISA, competitive immunoassays, radioimmunoassays, Western blot, indirect immunofluorescent assays and the like. In general, commercially available antibodies and/or kits may be used for the quantitative immunoassay of the expression level of cellobiohydrolase proteins.

20 VIII. Isolation And Purification Of Recombinant CBH1 Protein.

[200] In general, a variant CBH I protein produced in cell culture is secreted into the medium and may be purified or isolated, *e.g.*, by removing unwanted components from the cell culture medium. However, in some cases, a variant CBH I protein may be produced in a cellular form necessitating recovery from a cell lysate. In such cases the variant CBH 25 I protein is purified from the cells in which it was produced using techniques routinely employed by those of skill in the art. Examples include, but are not limited to, affinity chromatography (Tilbeurgh *et al.*, FEBS Lett. 16:215, 1984), ion-exchange chromatographic methods (Goyal *et al.*, Bioresource Technol. 36:37-50, 1991; Fliess *et al.*, Eur. J. Appl. Microbiol. Biotechnol. 17:314-318, 1983; Bhikhabhai *et al.*, J. Appl. Biochem. 6:336-345, 1984; Ellouz *et al.*, J. Chromatography 396:307-317, 1987), including 30 ion-exchange using materials with high resolution power (Medve *et al.*, J. Chromatography A 808:153-165, 1998), hydrophobic interaction chromatography (Tomaz and Queiroz, J. Chromatography A 865:123-128, 1999), and two-phase partitioning (Brumbauer, *et al.*, Bioseparation 7:287-295, 1999).

- 45 -

[201] Typically, the variant CBH I protein is fractionated to segregate proteins having selected properties, such as binding affinity to particular binding agents, e.g., antibodies or receptors; or which have a selected molecular weight range, or range of isoelectric points.

[202] Once expression of a given variant CBH I protein is achieved, the CBH I protein thereby produced is purified from the cells or cell culture. Exemplary procedures suitable for such purification include the following: antibody-affinity column chromatography, ion exchange chromatography; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; and gel filtration using, e.g., Sephadex G-75. Various methods of protein purification may be employed and such methods are known in the art and described e.g. in Deutscher, Methods in Enzymology, vol. 182, no. 57, pp. 779, 1990; Scopes, Methods Enzymol. 90: 479-91, 1982. The purification step(s) selected will depend, e.g., on the nature of the production process used and the particular protein produced.

IX. Utility of *cbh1* and CBH1

[203] It can be appreciated that the variant *cbh* nucleic acids, the variant CBH I protein and compositions comprising variant CBH I protein activity find utility in a wide variety applications, some of which are described below.

[204] New and improved cellulase compositions that comprise varying amounts BG-type, EG-type and variant CBH-type cellulases find utility in detergent compositions that exhibit enhanced cleaning ability, function as a softening agent and/or improve the feel of cotton fabrics (e.g., "stone washing" or "biopolishing"), in compositions for degrading wood pulp into sugars (e.g., for bio-ethanol production), and/or in feed compositions. The isolation and characterization of cellulase of each type provides the ability to control the aspects of such compositions.

[205] Variant (or mutant) CBHs with increased thermostability find uses in all of the above areas due to their ability to retain activity at elevated temperatures.

[206] Variant (or mutant) CBHs with decreased thermostability find uses, for example, in areas where the enzyme activity is required to be neutralized at lower temperatures so that other enzymes that may be present are left unaffected. In addition, the enzymes may find utility in the limited conversion of cellulotics, for example, in controlling the degree of crystallinity or of cellulosic chain-length. After reaching the desired extent of conversion the saccharifying temperature can be raised above the survival temperature of the destabilized CBH I. As the CBH I activity is essential for hydrolysis of crystalline cellulose, conversion of crystalline cellulose will cease at the elevated temperature.

[207] Variant (or mutant) CBHs with increased reversibility, i.e., enhanced refolding and retention of activity, also find use in similar areas. Depending upon the conditions of thermal inactivation, reversible denaturation can compete with, or dominate over, the irreversible process. Variants with increased reversibility would, under these conditions, exhibit increased resistance to thermal inactivation. Increased reversibility would also be of potential benefit in any process in which an inactivation event was followed by a treatment under non-inactivating conditions. For instance, in a Hybrid Hydrolysis and Fermentation (HHF) process for biomass conversion to ethanol, the biomass would first be incompletely saccharified by cellulases at elevated temperature (say 50°C or higher), then the temperature would be dropped (to 30°C, for instance) to allow a fermentative organism to be introduced to convert the sugars to ethanol. If, upon decrease of process temperature, thermally inactivated cellulase reversibly re-folded and recovered activity then saccharification could continue to higher levels of conversion during the low temperature fermentation process.

[208] In one approach, the cellulase of the invention finds utility in detergent compositions or in the treatment of fabrics to improve the feel and appearance.

[209] Since the rate of hydrolysis of cellulosic products may be increased by using a transformant having at least one additional copy of the *cbh* gene inserted into the genome, products that contain cellulose or heteroglycans can be degraded at a faster rate and to a greater extent. Products made from cellulose such as paper, cotton, cellulosic diapers and the like can be degraded more efficiently in a landfill. Thus, the fermentation product obtainable from the transformants or the transformants alone may be used in compositions to help degrade by liquefaction a variety of cellulose products that add to the overcrowded landfills.

[210] Separate saccharification and fermentation is a process whereby cellulose present in biomass, e.g., corn stover, is converted to glucose and subsequently yeast strains convert glucose into ethanol. Simultaneous saccharification and fermentation is a process whereby cellulose present in biomass, e.g., corn stover, is converted to glucose and, at the same time and in the same reactor, yeast strains convert glucose into ethanol. Thus, in another approach, the variant CBH type cellulase of the invention finds utility in the degradation of biomass to ethanol. Ethanol production from readily available sources of cellulose provides a stable, renewable fuel source.

[211] Cellulose-based feedstocks are comprised of agricultural wastes, grasses and woods and other low-value biomass such as municipal waste (e.g., recycled paper, yard clippings, etc.). Ethanol may be produced from the fermentation of any of these cellulosic

- 47 -

feedstocks. However, the cellulose must first be converted to sugars before there can be conversion to ethanol.

[212] A large variety of feedstocks may be used with the inventive variant CBH and the one selected for use may depend on the region where the conversion is being done. For example, in the Midwestern United States agricultural wastes such as wheat straw, corn stover and bagasse may predominate while in California rice straw may predominate. However, it should be understood that any available cellulosic biomass may be used in any region.

[213] A cellulase composition containing an enhanced amount of cellobiohydrolase finds utility in ethanol production. Ethanol from this process can be further used as an octane enhancer or directly as a fuel in lieu of gasoline which is advantageous because ethanol as a fuel source is more environmentally friendly than petroleum derived products. It is known that the use of ethanol will improve air quality and possibly reduce local ozone levels and smog. Moreover, utilization of ethanol in lieu of gasoline can be of strategic importance in buffering the impact of sudden shifts in non-renewable energy and petro-chemical supplies.

[214] Ethanol can be produced via saccharification and fermentation processes from cellulosic biomass such as trees, herbaceous plants, municipal solid waste and agricultural and forestry residues. However, the ratio of individual cellulase enzymes within a naturally occurring cellulase mixture produced by a microbe may not be the most efficient for rapid conversion of cellulose in biomass to glucose. It is known that endoglucanases act to produce new cellulose chain ends which themselves are substrates for the action of cellobiohydrolases and thereby improve the efficiency of hydrolysis of the entire cellulase system. Therefore, the use of increased or optimized cellobiohydrolase activity may greatly enhance the production of ethanol.

[215] Thus, the inventive cellobiohydrolase finds use in the hydrolysis of cellulose to its sugar components. In one embodiment, a variant cellobiohydrolase is added to the biomass prior to the addition of a fermentative organism. In a second embodiment, a variant cellobiohydrolase is added to the biomass at the same time as a fermentative organism. Optionally, there may be other cellulase components present in either embodiment.

[216] In another embodiment the cellulosic feedstock may be pretreated. Pretreatment may be by elevated temperature and the addition of either of dilute acid, concentrated acid or dilute alkali solution. The pretreatment solution is added for a time sufficient to at least partially hydrolyze the hemicellulose components and then neutralized.

- 48 -

[217] The major product of CBHI action on cellulose is cellobiose which is available for conversion to glucose by BG activity (for instance in a fungal cellulase product). Either by the pretreatment of the cellulosic biomass or by the enzymatic action on the biomass, other sugars, in addition to glucose and cellobiose, can be made available from the biomass. The hemi-cellulose content of the biomass can be converted (by hemi-cellulases) to sugars such as xylose, galactose, mannose and arabinose. Thus, in a biomass conversion process, enzymatic saccharification can produce sugars that are made available for biological or chemical conversions to other intermediates or end-products. Therefore, the sugars generated from biomass find use in a variety of processes in addition to the generation of ethanol. Examples of such conversions are fermentation of glucose to ethanol (as reviewed by M.E. Himmel *et al.* pp2-45, in "Fuels and Chemicals from Biomass", ACS Symposium Series 666, ed B.C. Saha and J. Woodward, 1997) and other biological conversions of glucose to 2,5-diketo-D-gluconate (US Patent No. 6,599,722), lactic acid (R. Datta and S-P. Tsai pp224-236, *ibid*), succinate (R.R. Gokarn, M.A. Eiteman and J. Sridhar pp237-263, *ibid*), 1,3-propanediol (A-P. Zheng, H. Biebl and W-D. Deckwer pp264-279, *ibid*), 2,3-butanediol (C.S. Gong, N. Cao and G.T. Tsao pp280-293, *ibid*), and the chemical and biological conversions of xylose to xylitol (B.C. Saha and R.J. Bothast pp307-319, *ibid*). See also, for example, WO 98/21339.

[218] The detergent compositions of this invention may employ besides the cellulase composition (irrespective of the cellobiohydrolase content, i.e., cellobiohydrolase -free, substantially cellobiohydrolase -free, or cellobiohydrolase enhanced), a surfactant, including anionic, non-ionic and ampholytic surfactants, a hydrolase, building agents, bleaching agents, bluing agents and fluorescent dyes, caking inhibitors, solubilizers, cationic surfactants and the like. All of these components are known in the detergent art. The cellulase composition as described above can be added to the detergent composition either in a liquid diluent, in granules, in emulsions, in gels, in pastes, and the like. Such forms are well known to the skilled artisan. When a solid detergent composition is employed, the cellulase composition is preferably formulated as granules. Preferably, the granules can be formulated so as to contain a cellulase protecting agent. For a more thorough discussion, see US Patent Number 6,162,782 entitled "Detergent compositions containing cellulase compositions deficient in CBH I type components," which is incorporated herein by reference.

[219] Preferably the cellulase compositions are employed from about 0.00005 weight percent to about 5 weight percent relative to the total detergent composition. More

- 49 -

preferably, the cellulase compositions are employed from about 0.0002 weight percent to about 2 weight percent relative to the total detergent composition.

[220] In addition the variant *cbh I* nucleic acid sequence finds utility in the identification and characterization of related nucleic acid sequences. A number of techniques useful for determining (predicting or confirming) the function of related genes or gene products include, but are not limited to, (A) DNA/RNA analysis, such as (1) overexpression, ectopic expression, and expression in other species; (2) gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing (VIGS, see Baulcombe, 100 Years of Virology, Calisher and Horzinek eds., Springer-Verlag, New York, NY 15:189-201, 1999); (3) analysis of the methylation status of the gene, especially flanking regulatory regions; and (4) in situ hybridization; (B) gene product analysis such as (1) recombinant protein expression; (2) antisera production, (3) immunolocalization; (4) biochemical assays for catalytic or other activity; (5) phosphorylation status; and (6) interaction with other proteins via yeast two-hybrid analysis; (C) pathway analysis, such as placing a gene or gene product within a particular biochemical or signaling pathway based on its overexpression phenotype or by sequence homology with related genes; and (D) other analyses which may also be performed to determine or confirm the participation of the isolated gene and its product in a particular metabolic or signaling pathway, and help determine gene function.

[221] All patents, patent applications, articles and publications mentioned herein, are hereby expressly incorporated herein by reference.

EXAMPLES

[222] The present invention is described in further detail in the following examples which are not in any way intended to limit the scope of the invention as claimed. The attached Figures are meant to be considered as integral parts of the specification and description of the invention. All references cited are herein specifically incorporated by reference for all that is described therein.

EXAMPLE 1

Alignment of known Cel7A cellulases

[223] The choice of several of the mutations was determined by first aligning *Hypocrea jecorina* Cel7A to its 41 family members using structural information and a modeling program. The alignment of the primary amino acid sequence of all 42 family members is shown in Figure 8.

- 50 -

[224] For four of the members (i.e., 20VW.1, 1A39, 6CEL and 1EG1.1), the crystal structure had been previously determined. The 4 aligned proteins for which there were published structures had their alignment locked for all residues whose backbone atoms were within a specific RMS deviation (RMS less than or equal to 2.0 Å). The tertiary structural alignment of the four sequences was performed using MOE version 2001.01 by

5 structural alignment of the four sequences was performed using MOE version 2001.01 by Chemical Computing Group, Montreal Canada. The overlapping structural elements were used to freeze the primary structures of the four sequences. The remaining 38 sequences then had their primary amino acid structure aligned with the frozen four using MOE with secondary structure prediction on and other parameters set to their default settings.

10 [225] Based on the alignments, various single and multiple amino acid mutations were made in the protein by site mutagenesis.

[226] Single amino acid mutations were based on the following rationale (see also Table 1): After examining the conservation of amino acids between the homologues, sites were picked in the *H. jecorina* sequence where a statistical preference for another amino acid was seen amongst the other 41 sequences (e.g.: at position 77 the Ala, only present in *H. jecorina* and 3 other homologues, was changed to Asp, present in 22 others). The effect of each substitution on the structure was then modeled.

Table 1: Cel7A Variants and Rationale for Change

<u>Cel7A Variants and Rationale for Change</u>	T _m	ΔT _m
Wild Type <i>H. jecorina</i>	62.5	
(4)A77D(22) 3 possible H-bonds to Q7 and I80	62.2	-0.3
(7)S113D(18) numerous new H-bonds to backbone to stabilize turn	62.8	0.3
(8)L225F(13) better internal packing	61.6	-0.9
25 (5)L288F(17) better internal packing	62.4	-0.1
(1)A299E(24) extra ligand to cobalt atom observed in crystal structure	61.2	-1.3
(4)N301K(11) salt bridges to E295 and E325	63.5	1.0
(5)T356L(20) better internal packing	62.6	0.1
(2)G430F(17) better surface packing	61.7	-0.8

30 [227] Multiple amino acid mutations were based on a desire to affect the stability, processivity, and product inhibition of the enzyme. The following multiple site changes in the *H. jecorina* sequence were constructed:

- 1) Thr 246 Cys + Tyr 371 Cys
- 35 2) Thr 246 Ala + Arg 251 Ala + Tyr 252 Ala
- 3) Thr 380 Gly + Tyr 381 Asp + Arg 394 Ala + deletion of Residues 382 to 393, inclusive
- 4) Thr 380 Gly + Tyr 381 Asp + Arg 394 Ala
- 5) Tyr 252 Gln + Asp 259 Trp + Ser 342 Tyr

40

[228] The T246A/R251A/Y252A and the other triple + deletion mutant are both predicted to decrease the product inhibition of the enzyme. The Thr246Cys + Tyr371Cys is predicted to increase the stability of the enzyme and increase the processivity of it. The D259W/Y252Q/S342Y variant is predicted to affect the product inhibition of the enzyme.

5 [229] Other single and multiple mutations were constructed using methods well known in the art (see references above) and are presented in Table 2.

Table 2: *H. jecorina* CBH I variants

Mutations
S8P
N49S
A68T
A77D
N89D
S92T
S113N
S113D
L225F
P227A
P227L
D249K
T255P
D257E
S279N
L288F
E295K
S297T
A299E
N301K
T332K
T332Y
T332H
T356L
F338Y
V393G
G430F
T41I (plus deletion of Thr @ 445)
V403D/T462I
S196T/S411F
E295K/S398T
A112E/T226A
T246C/Y371C
G22D/S278P/T296P
S8P/N103I/S113N
S113T/T255P/K286M
P227L/E325K/Q487L
P227T/T484S/F352L
T246A/R251A/Y252A

- 52 -

Mutations
T380G/Y381D/R394A
Y252Q/D259W/S342Y
A68T/G440R/P491L
Q17L/E193V/M213I/F352L
S8P/N49S/A68T/S113N
A112E/P227L/S278P/T296P
S8P/N49S/A68T/N103I/S113N
S8P/N49S/A68T/S278P/T296P
G22D/N49S/A68T/S278P/T296P
G22D/N103I/S113N/S278P/T296P
S8P/N49S/A68T/S113N/P227L
S8P/N49S/A68T/A112E/T226A
S8P/N49S/A68T/A112E/P227L
T41I/A112E/P227L/S278P/T296P
S8P/T41I/N49S/A68T/S113N/P227L
S8P/T41I/N49S/A68T/A112E/P227L
G22D/N49S/A68T/P227L/S278P/T296P
G22D/N49S/A68T/N103I/S113N/S278P/T296P
G22D/N49S/A68T/N103I/S113N/P227L/S278P/ T296P
G22D/N49S/A68T/N103I/A112E/P227L/S278P/ T296P
G22D/N49S/N64D/A68T/N103I/S113N/S278P/ T296P
S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F
S8P/G22D/T41I/N49S/A68T/N103I/S113N/S278P/T296P
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P
S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/N301R
S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F
S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P/N301R
S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/T296P/N301R
S8P/T41I/N49S/S57N/A68T/S113N/P227L/D249K/S278P/T296P/N301R
S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P/N301R
S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/S278P/T296P/N301R/ E325K/S411F
S8P/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/S278P/T296P/N301R/E325K/ V403D/S411F/T462I
S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/S278P/T296P/N301R/ E325K/V403D/S411F/T462I

EXAMPLE 2

Cloning and Expression of CBHI variants in *H. jecorina*

5

A. Construction of the *H. jecorina* general-purpose expression plasmid-PTEX.

[230] The plasmid, pTEX was constructed following the methods of Sambrook et al. (1989), *supra*, and is illustrated in FIG. 7. This plasmid has been designed as a multi-purpose expression vector for use in the filamentous fungus *Trichoderma longibrachiatum*.

The expression cassette has several unique features that make it useful for this function. Transcription is regulated using the strong CBH I gene promoter and terminator sequences for *T. longibrachiatum*. Between the CBH I promoter and terminator there are unique PmeI and SstI restriction sites that are used to insert the gene to be expressed.

5 The *T. longibrachiatum* pyr4 selectable marker gene has been inserted into the CBH I terminator and the whole expression cassette (CBH I promoter-insertion sites-CBH I terminator-pyr4 gene-CBH I terminator) can be excised utilizing the unique NotI restriction site or the unique NotI and NheI restriction sites.

[231] This vector is based on the bacterial vector, pSL1180 (Pharmacia Inc., Piscataway, N.J.), which is a PUC-type vector with an extended multiple cloning site. One skilled in the art would be able to construct this vector based on the flow diagram illustrated in FIG. 7.

[232] The vector pTrex2L was constructed from pTrex2, a derivative of pTEX. The sequence for pTrex2 is given in Figure 6.

[233] The exact plasmid used is not that important as long as the variant protein is expressed at a useful level. However, maximizing the expression level by forcing integration at the *cbh1* locus is advantageous.

B. Cloning

[234] Using methods known in the art a skilled person can clone the desired CBH I variant into an appropriate vector. As noted above, the exact plasmid used is not that important as long as the variant protein is expressed at a useful level. The following description of the preparation of one of the inventive variant CBH I enzymes can be utilized to prepare any of the inventive variants described herein.

[235] The variant *cbh 1* genes were cloned into the pTrex2L vector.

[236] Construction of plasmid pTrex2L was done as follows: The 6 nucleotides between the unique Sac II and Asc I sites of pTrex2 were replaced with a synthetic linker containing a BstE II and BamH I sites to produce plasmid Trex2L. The complementary synthetic linkers

21-mer synthetic oligo CBHlink1+: GGTTTGGATCCGGTCACCAGG

and

27-mer synthetic oligo CBHlink-: CGCGCCTGGTGACCGGATCCAAACCGC

were annealed.

[237] The pTrex2 was digested with Sac II and Asc I. The annealed linker was then ligated into pTrex2 to create pTrex2L. The plasmid was then digested with an appropriate restriction enzyme(s) and a wild type CBH I gene was ligated into the plasmid.

- 54 -

[238] Primers were used to introduce the desired mutations into the wild-type gene. It will be understood that any method that results in the introduction of a desired alteration or mutation in the gene may be used. Synthetic DNA primers were used as PCR templates for mutant constructions. It is well within the knowledge of the skilled artisan to design the primers based on the desired mutation to be introduced.

[239] The mutagenic templates were extended and made double stranded by PCR using the synthetic DNA oligonucleotides. After 25 PCR cycles the final product was primarily a 58 bp double stranded product comprising the desired mutation. The mutagenic fragments were subsequently attached to wild-type CBH I fragments and ligated into the plasmid using standard techniques.

C. Transformation and Expression

[240] The prepared vector for the desired variant was transformed into the uridine auxotroph version of the double or quad deleted *Trichoderma* strains (see Table 3; see also U.S. Patent Nos. 5,861,271 and 5,650,322) and stable transformants were identified.

Table 3: Transformation/Expression strain

CBH I Variant	Expression Strain
A77D	quad-delete strain (1A52)
S113D	double-delete strain
L225F	double-delete strain
L288F	double-delete strain
A299E	quad-delete strain (1A52)
N301K	quad-delete strain (1A52)
T356L	double-delete strain
G430F	quad-delete strain (1A52)
T246C/Y371C	quad-delete strain (1A52)
T246A/R251A/Y252A	quad-delete strain (1A52)
Y252Q/D259W/S342Y	quad-delete strain (1A52)
T380G/Y381D/R394A	quad-delete strain (1A52)
T380G/Y381D/R394A plus deletion of 382-393	quad-delete strain (1A52)

"double-delete" (Δ CBHI & Δ CBHII) and the "quad-delete" (Δ CBHI & Δ CBHII, Δ EGI & Δ EGII) *T.reesei* host strains

- 55 -

[241] To select which transformants expressed variant CBH I, DNA was isolated from strains following growth on Vogels+1% glucose and Southern blot experiments performed using an isolated DNA fragment containing only the variant CBH I. Transformants were isolated having a copy of the variant CBH I expression cassette integrated into the genome of the host cell. Total mRNA was isolated from the strains following growth for 1 day on Vogels+1% lactose. The mRNA was subjected to Northern analysis using the variant CBH I coding region as a probe. Transformants expressing variant CBH I mRNA were identified.

[242] One may obtain any other novel variant CBH I cellulases or derivative thereof by employing the methods described above.

EXAMPLE 3

Expression of CBH1 variants in *A. niger*

[243] The PCR fragments were obtained using the following primers and protocols

[244] The following DNA primers were constructed for use in amplification of homologous CBH1 genes from genomic DNA's isolated from various microorganisms. All symbols used herein for protein and DNA sequences correspond to IUPAC IUB Biochemical Nomenclature Commission codes.

[245] Homologous 5' (FRG192) and 3' (FRG193) primers were developed based on the sequence of CBH1 from *Trichoderma reesei*. Both primers contained Gateway cloning sequences from Invitrogen® at the 5' of the primer. Primer FRG192 contained attB1 sequence and primer FRG193 contained attB2 sequence.

Sequence of FRG192 without the attB1:
ATGTATCGGAAGTTGGCCG (signal sequence of CBH1 *H. jecorina*) (SEQ ID NO: 3)

Sequence of FRG193 without the attB2:
TTACAGGCACTGAGAGTAG (cellulose binding module of CBH1 *H. jecorina*) (SEQ ID NO: 4)

[246] The *H. jecorina* CBH I cDNA clone served as template.

[247] PCR conditions were as follows: 10 µL of 10X reaction buffer (10X reaction buffer comprising 100mM Tris HCl, pH 8-8.5; 250 mM KCl; 50 mM (NH₄)₂SO₄; 20 mM MgSO₄); 0.2 mM each of dATP, dTTP, dGTP, dCTP (final concentration), 1 µL of 100 ng/µL genomic DNA, 0.5 µL of PWO polymerase (Boehringer Mannheim, Cat # 1644-947) at 1 unit per µL, 0.2µM of each primer, FRG192 and FRG193, (final concentration), 4µl DMSO and water to 100 µL.

- 56 -

[248] Various sites in *H. jecorina* CBH1 may be involved in the thermostability of the variants and the *H. jecorina* CBH1 gene was therefore subjected to mutagenesis.

[249] The fragments encoding the variants were purified from an agarose gel using the Qiagen Gel extraction KIT. The purified fragments were used to perform a clonase
5 reaction with the pDONR™201 vector from Invitrogen® using the Gateway™ Technology instruction manual (version C) from Invitrogen®, hereby incorporated by reference herein. Genes were then transferred from this ENTRY vector to the destination vector (pRAXdes2) to obtain the expression vector pRAXCBH1.

[250] Cells were transformed with an expression vector comprising a variant CBH I
10 cellulase encoding nucleic acid. The constructs were transformed into *A. niger var. awamori* according to the method described by Cao *et al* (Cao Q-N, Stubbs M, Ngo KQP, Ward M, Cunningham A, Pai EF, Tu G-C and Hofmann T (2000) Penicillopepsin-JT2 a recombinant enzyme from *Penicillium janthinellum* and contribution of a hydrogen bond in subsite S3 to *kcat* *Protein Science* 9:991-1001).

[251] Transformants were streaked on minimal medium plates (Ballance DJ, Buxton FP,
15 and Turner G (1983) Transformation of *Aspergillus nidulans* by the orotidine-5'-phosphate decarboxylase gene of *Neurospora crassa* *Biochem Biophys Res Commun* 112:284-289) and grown for 4 days at 30°C. Spores were collected using methods well known in the art (See <<http://www.fgsc.net/fgn48/Kaminskyj.htm>>). *A. nidulans* conidia are harvested in
20 water (by rubbing the surface of a conidiating culture with a sterile bent glass rod to dislodge the spores) and can be stored for weeks to months at 4°C without a serious loss of viability. However, freshly harvested spores germinate more reproducibly. For long-term storage, spores can be stored in 50% glycerol at -20°C, or in 15-20% glycerol at -80°C. Glycerol is more easily pipetted as an 80% solution in water. 800µl of aqueous conidial
25 suspension (as made for 4°C storage) added to 200µl 80% glycerol is used for a -80°C stock; 400 µl suspension added to 600 µl 80% glycerol is used for a -20°C stock. Vortex before freezing. For mutant collections, small pieces of conidiating cultures can be excised and placed in 20% glycerol, vortexed, and frozen as -80°C stocks. In our case we store them in 50% glycerol at -80°C.

[252] *A. niger var awamori* transformants were grown on minimal medium lacking uridine
30 (Ballance et al. 1983). Transformants were screened for cellulase activity by inoculating 1cm² of spore suspension from the sporulated grown agar plate into 100ml shake flasks for 3 days at 37°C as described by Cao et al. (2000).

[253] The CBHI activity assay is based on the hydrolysis of the nonfluorescent 4-
35 methylumbelliferyl-β-lactoside to the products lactose and 7-hydroxy-4-methylcoumarin,

- 57 -

the latter product is responsible for the fluorescent signal. Pipette 170 μ l 50 mM NaAc buffer pH 4.5 in a 96-well microtiter plate (MTP) (Greiner, Fluotrac 200, art. nr. 655076) suitable for fluorescence. Add 10 μ l of supernatant and then add 10 μ l of MUL (1 mM 4-methylumbelliferyl- β -lactoside (MUL) in milliQ water) and put the MTP in the Fluostar Galaxy (BMG Labtechnologies; D-77656 Offenburg). Measure the kinetics for 16 min. (8 cycles of 120s each) using $\lambda_{320 \text{ nm}}$ (excitation) and $\lambda_{460 \text{ nm}}$ (emission) at 50°C. Supernatants having CBH activity were then subjected to Hydrophobic Interaction Chromatography.

EXAMPLE 4

Stability of CBH 1 variants

[254] CBH I cellulase variants were cloned and expressed as above (see Examples 2 and 3). Cel7A wild type and variants were then purified from cell-free supernatants of these cultures by column chromatography. Proteins were purified using hydrophobic interaction chromatography (HIC). Columns were run on a BioCAD® Sprint Perfusion Chromatography System using Poros® 20 HP2 resin both made by Applied Biosystems.

[255] HIC columns were equilibrated with 5 column volumes of 0.020 M sodium phosphate, 0.5 M ammonium sulfate at pH 6.8. Ammonium sulfate was added to the supernatants to a final concentration of approximately 0.5 M and the pH was adjusted to 6.8. After filtration, the supernatant was loaded onto the column. After loading, the column was washed with 10 column volumes of equilibration buffer and then eluted with a 10 column volume gradient from 0.5 M ammonium sulfate to zero ammonium sulfate in 0.02 M sodium phosphate pH 6.8. Cel7A eluted approximately mid-gradient. Fractions were collected and pooled on the basis of reduced, SDS-PAGE gel analysis.

[256] The melting points were determined according to the methods of Luo, *et al.*, *Biochemistry* **34**:10669 and Gloss, *et al.*, *Biochemistry* **36**:5612. See also Sandgren *et al.* (2003) *Protein Science* **12**(4) pp848.

[257] Data was collected on the Aviv 215 circular dichroism spectrophotometer. The native spectra of the variants between 210 and 260 nanometers were taken at 25°C. Buffer conditions were 50 mM Bis Tris Propane/50 mM ammonium acetate/glacial acetic acid at pH 5.5. The protein concentration was kept between 0.25 and 0.5 mgs/mL. After determining the optimal wavelength to monitor unfolding, the samples were thermally denatured by ramping the temperature from 25°C to 75°C under the same buffer conditions. Data was collected for 5 seconds every 2 degrees. Partially reversible unfolding was monitored at 230 nanometers in a 0.1 centimeter path length cell. While at 75°C, an unfolded spectra was collected as described above. The sample was then

cooled to 25°C to collect a refolded spectra. The difference between the three spectra at 230nm was used to assess the variants reversibility.

[258] The thermal denaturation profiles are shown in Figure 9A and 9B for wildtype CBH I and various variant CBH I's. See also Table 4.

5

Table 4: Thermal Stability of Variant CBH I cellulases

<i>H. jecorina</i> CBH I Residue Substitution	T _m	delta T _m	% rev 230nm
Wild type	62.5		23
S8P	63.1	0.6	
N49S	63.7	1.2	
A68T	63.7	1.2	32
A77D	62.2	-0.3	
N89D	63.6	1.1	50
S92T	64.4	1.9	25
S113D	62.8	0.3	
S113N	64.0	1.5	
L225F	61.6	-0.9	
P227A	64.8	2.3	49
P227L	65.2	2.7	45
D249K	64.0	1.5	39
T255P	64.4	1.9	35
S279N	62.4	-0.1	~95
E295K	64.0	1.5	~95
T332K	63.3	0.8	37
T332Y	63.3	0.8	37
T332H	62.7	0.2	64
F338Y	60.8	-1.7	~95
G430F	61.7	-0.8	
L288F	62.4	-0.1	
A299E	61.2	-1.3	
N301K	63.5	1.0	
T356L	62.6	0.1	
D257E	61.8	-0.7	45
V393G	61.7	-0.8	43
S297T	63.3	0.8	31
T41I plus deletion @ T445	64.2	1.7	
T246C/Y371C	65.0	2.5	
S196T/S411F	65.3	2.8	27
E295K/S398T	63.9	1.4	36
V403D/T462I	64.5	2	53
A112E/T226A	63.5	1.0	
A68T/G440R/P491L	63.1	0.6	32
G22D/S278P/T296P	63.6	1.1	
T246A/R251A/Y252A	63.5	1.0	
T380G/Y381D/R394A	58.1	-4.4	
Y252Q/D259W/S342Y	59.9	-2.6	50
S113T/T255P/K286M	63.8	1.3	16

- 59 -

<i>H. jecorina</i> CBH I Residue Substitution	Tm	delta Tm	% rev 230nm
P227L/E325K/Q487L	64.5	2.0	22
P227T/T484S/F352L	64.2	1.7	45
Q17L/E193V/M213I/F352L	64.0	1.5	34
S8P/N49S/A68T/S113N	64.5	2.0	90
S8P/N49S/A68T/S113N/P227L	66.0	3.5	86
T41I/A112E/P227L/S278P/T296P	66.1	3.6	48
S8P/N49S/A68T/A112E/T226A	64.6	2.1	46
S8P/N49S/A68T/A112E/P227L	65.2	2.7	32
S8P/T41I/N49S/A68T/A112E/P227L	67.6	5.1	40
G22D/N49S/A68T/P227L/S278P/T296P	65.9	3.4	26
G22D/N49S/A68T/N103I/S113N/P227L/S278P/T296P	65.3	2.8	72
G22D/N49S/A68T/N103I/A112E/P227L/S278P/T296P	65.1	2.6	20
G22D/N49S/N64D/A68T/N103I/S113N/S278P/T296P	61.4	-1.1	75
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/ T296P	68.8	6.3	56
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/ S278P/T296P	69.0	6.5	71
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/ T296P/N301R	68.7	6.2	70
S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/ S278P/T296P/N301R	68.8	6.3	74
S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/ T296P/N301R	69.9	7.4	88
S8P/T41I/N49S/S57N/A68T/S113N/P227L/D249K/S278P/ T296P/N301R	68.9	6.4	~100
S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/ N301R	68.7	6.2	92
S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/ T462I	68.8	6.3	~100
S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/ V403D/T462I	68.5	6.0	~100
S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F	68.6	6.1	~100
S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/ S411F	69.5	7.0	~100
S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/ D249K/T255P/S278P/T296P/N301R/E325K/S411F	70.7	8.2	~100
S8P/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/ T255P/S278P/T296P/N301R/E325K/V403D/S411F/T462I	71.0	8.5	~100
S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/ D249K/T255P/S278P/T296P/N301R/E325K/V403D/S411F/ T462I	70.9	8.4	~100

[259] Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed

should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the claims.

CLAIMS

1. A variant CBH I cellulase, wherein said variant comprises a substitution or deletion at a position corresponding to one or more of residues S8, Q17, G22, T41, N49, S57, N64, A68, A77, N89, S92, N103, A112, S113, E193, S196, M213, L225, T226, P227, T246, D249, R251, Y252, T255, D257, D259, S278, S279, K286, L288, E295, T296, S297, A299, N301, E325, T332, F338, S342, F352, T356, Y371, T380, Y381, V393, R394, S398, V403, S411, G430, G440, T445, T462, T484, Q487, and P491 in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2).
2. A variant CBH I cellulase according to Claim 1, wherein said variant comprises a substitution at a position corresponding to one or more of residues S8P, Q17L, G22D, T41I, N49S, S57N, N64D, A68T, A77D, N89D, S92T, N103I, A112E, S113(T/N/D), E193V, S196T, M213I, L225F, T226A, P227(L/T/A), T246(C/A), D249K, R251A, Y252(A/Q), T255P, D257E, D259W, S278P, S279N, K286M, L288F, E295K, T296P, S297T, A299E, N301(R/K), E325K, T332(K/Y/H), F338Y, S342Y, F352L, T356L, Y371C, T380G, Y381D, V393G, R394A, S398T, V403D, S411F, G430F, G440R, T462I, T484S, Q487L and/or P491L in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2).
3. A variant CBH I cellulose according to Claim 2, further comprising a deletion at a position corresponding to T445 in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2).
4. A variant CBH I cellulase, wherein said variant comprises a substitution at a position corresponding to a residue selected from the group consisting of S8P, N49S, A68T, A77D, N89D, S92T, S113(N/D), L225F, P227(A/L/T), D249K, T255P, D257E, S279N, L288F, E295K, S297T, A299E, N301(R/K), T332(K/Y/H), F338Y, T356L, V393G, G430F in CBH I from *Hypocrea jecorina* (SEQ ID NO: 2).
5. A variant CBH I cellulase, wherein said variant CBH I consists essentially of the mutations selected from the group consisting of
 - xi. A112E/T226A;
 - xli. S196T/S411F;
 - xlii. E295K/S398T;
 - xliv. T246C/Y371C;
 - xliv. V403D/T462I
 - xlvi. T41I plus deletion at T445
 - xlvi. A68T/G440R/P491L;
 - xlvi. G22D/S278P/T296P;
 - xlvi. T246A/R251A/Y252A;

- 62 -

- xlix. T380G/Y381D/R394A;
- l. Y252Q/D259W/S342Y;
- li. S113T/T255P/K286M;
- lii. P227L/E325K/Q487L;
- liii. P227T/T484S/F352L;
- liv. Q17L/E193V/M213I/F352L;
- lv. S8P/N49S/A68T/S113N;
- lvi. S8P/N49S/A68T/S113N/P227L;
- lvii. T41I/A112E/P227L/S278P/T296P;
- lviii. S8P/N49S/A68T/A112E/T226A;
- lix. S8P/N49S/A68T/A112E/P227L;
- lx. S8P/T41I/N49S/A68T/A112E/P227L;
- lxi. G22D/N49S/A68T/P227L/S278P/T296P;
- lxii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P;
- lxiii. G22D/N49S/A68T/N103I/S113N/P227L/S278P/T296P;
- lxiv. G22D/N49S/A68T/N103I/A112E/P227L/S278P/T296P;
- lxv. G22D/N49S/N64D/A68T/N103I/S113N/S278P/T296P;
- lxvi. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P
;
- lxvii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/S278P/T296P/N301R
;
- lxviii. S8P/G22D/T41I/N49S/A68T/N103I/S113N/P227L/D249K/S278P/T296P
/N301R
- lxix. S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/T296P/N301
R;
- lxx. S8P/T41I/N49S/S57N/A68T/S113N/P227L/D249K/S278P/T296P/N301
R;
- lxxi. S8P/G22D/T41I/N49S/A68T/S113N/P227L/D249K/S278P/N301R;
- lxxii. S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I;
- lxxiii. S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/V403D/T462I;
- lxxiv. S8P/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F;
- lxxv. S8P/G22D/T41I/N49S/A68T/S92T/S113N/P227L/D249K/S411F;
- lxxvi. S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/
S278P/T296P/N301R/E325K/S411F;

- 63 -

lxxvii. S8P/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/S278P
/T296P/N301R/E325K/V403D/S411F/T462I;

lxxviii. S8P/G22D/T41I/N49S/A68T/S92T/S113N/S196T/P227L/D249K/T255P/
S278P/T296P/N301R/E325K/V403D/S411F/T462I

in CBH I from *Hypocrea jecorina* (SEQ ID NO:2).

6. A nucleic acid encoding a CBH I variant according to claim 1.
7. A nucleic acid encoding a CBH I variant according to claim 4.
8. A nucleic acid encoding a CBH I variant according to claim 5.
9. A vector comprising a nucleic acid encoding a CBH I variant of claim 6.
10. A vector comprising a nucleic acid encoding a CBH I variant of claim 7.
11. A vector comprising a nucleic acid encoding a CBH I variant of claim 8.
12. A host cell transformed with the vector of claim 9.
13. A host cell transformed with the vector of claim 10.
14. A host cell transformed with the vector of claim 11.
15. A method of producing a CBH I variant comprising the steps of:
 - (a) culturing the host cell according to claim 12 in a suitable culture medium under suitable conditions to produce CBH I variant;
 - (b) obtaining said produced CBH I variant.
16. A method of producing a CBH I variant comprising the steps of:
 - (a) culturing the host cell according to claim 13 in a suitable culture medium under suitable conditions to produce CBH I variant;
 - (b) obtaining said produced CBH I variant.
17. A method of producing a CBH I variant comprising the steps of:
 - (a) culturing the host cell according to claim 14 in a suitable culture medium under suitable conditions to produce CBH I variant;
 - (b) obtaining said produced CBH I variant.
18. A detergent composition comprising a surfactant and a CBH I variant, wherein said CBH I variant comprises a CBH I variant according to claim 1.
19. The detergent according to claim 18, wherein said detergent is a laundry detergent.
20. The detergent according to claim 18, wherein said detergent is a dish detergent.
21. A feed additive comprising a CBH I variant according to claim 1.
22. A method of treating wood pulp comprising contacting said wood pulp with a CBH I variant according to claim 1.
23. A method of converting biomass to sugars comprising contacting said biomass with a CBH I variant according to claim 1.

Figure 1: Amino Acid and Nucleic Acid Sequences of *Hypocrea jecorina* Cel7A

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCGGCCT GCACTCTCCA ATCGGAGACT CACCCGGCTC TGACATGGCA GAAATGCTCG TCTGGTGGCA CTTCGACTCA ACAGACAGGC TCCGTGGTCA

101 · IASpAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCGGTGG ACTCACGCTA CGAACAGCAG CAGCAACTGC TACGATGGCA ACACCTGGAG CTCGACCCCTA TGTCCTGACA ACGAGACCTG

201 · AlalysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGGCGC CTACGGCTCC ACCTACGGAG TTACCACGAG CGGTAAACAGC CTCCTCATTTG GCTTTGTTCAC CCAGTCTGGC

301 GlnLysAsnVal GlyAlaArg LeuTyrLeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAGAAGAC TTGGCGCTCG CCTTTACCTT ATGGCGAGCG ACAGACCTA CCAGGAATC ACCCTGCTTG GCAACGATT CTCCTTCGAT GTTGAIGTTT

401 · SGLLeuPro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlalysTyr
 CGCAGCTGCC GTGGCGGTGG AACGGAGCTC TCTACTTCTGT GTCCATGGAC CGGGAATGGT GCGTGAGCAA GTATCCACC AACACCCGCTG GCGCCAAATA

501 · GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleAsn GlyGlnAla AsnValGlu GlyTrpGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGCGAATCTG AAGTTCATCA ATGGCCAGGC CAACCTTGAG GGCTGGGAGC CGTCAATCAA CAACCGCGAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGlu AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
 ACGGCGATTG GAGACACGG AAGCTGTCTG TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTACCCTC CCACCCCTGC AGGACTGTGC

701 · GGLnGluIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
 GCCAGGAGAT CTCGGAGGCT GATGGGTGGC GCGGAACCTA CTCGGATAAC AGATATGGCG GCACCTGGCA TCCCGATGGC TCGCACTGGA ACCCATACCG

801 · LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGTTCT ACGGCCCTGG CTCAAAGCTT ACCCTCGATA CCACCAAGAA ATTGACCGTT GTCACCCAGT TCGAGACGTC GGGTGGCCATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGlu LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGlu
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCCAGCAGC CCAACGCCGA GCTTGGTAGT TACTCTGGCA ACGAGCTCAA CGATGATTAAC TGCACAGCTG

1001 · GGLuAlaGlu PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValLeuValMet SerLeuTrp
 ACGAGGCAGA ATTCGGCGGA TCCTCTTTCT CAGACAAGG CGGCCTGACT CAGTTCAAGA AGGCTACCTC TGGGGGCATG GTTCTGTGCA TGAGTCTGTG

1101 · AspAspTyr TyrAlaAsnMet LeuTrpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTAAC TAGCCCAA CA TGCTGTGGCT GGACTCCACC TACCCGACAA ACGAGACCTC CTCACACCC GGTGCCCTGC GCGGAAGCTG CTCACCCAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCCGGTGTCC CTGCTCAGT CGAATCTCAG TCTCCCAA CGAAGTCAAC CTCCTCCAAC ATCAAGTTCG GACCCATGG CAGCACCGGC AACCCTAGCG

1301 · GGLyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GCGGCAACC TCCGGGGGGA AACCGCCCTG GCACCCACC CACCCGCCG CACCCGCGG CCACTGGAAG CTCCTCCGGA CCTACCCAGT CTCACTACCG

1401 · GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysLeu
 CCAGTGGCGC GGTATTGGCT ACAGCGGCC CACGGTCTGC GCCAGCGGCA CAACTTGCCA GGTCTCTAAC CCTTACTACT CTCAGTGCCT G

Figure 2



Hypocrea jecorina Cel7A

497 amino acids

1 -431 in the catalytic module

432-461 in the linker region

462-497 in the cellulose binding module

12 disulfide bonds--10 in the catalytic module

E212 and E217 are the active site residues

Figure 3

```

20VW.A F. oxysporum Cel7B .....PDE-AR C K ETY TK S --G--K
1A39 H. insolens Cel7B .....KGE-TR V Q TT TEG --G--K
6CEL H. jecorina Cel7A .....ACTLOS T P TQ SSG --T--T
1EG1.A H. jecorina Cel7B .....QDGTSTP V K TTY TEG --G--V
consensus .....tpgt-tkSvBPhLttYrCtkag--g--ck

20VW.A KQNTI A AGIHGIRQE---NGAGCGDWG KPNATA EA V A ILS --DBNA K
1A39 PATNF L SLWHWHR A EGLGPGCGDWG PPDG VE A IE --PD--B
6CEL QQTGE ANRWYTHATNSBTHCYDGT---WSTL NE A CL --A--A A
1EG1.A AQTSL L WNRW H-DANYNSCTVHG---WTL BA --PFE --D A
consensus -qtosIFIDanw-wih-----n-cgqgdwqg-pnstlCPDe-sCaRNCilaG-n---aIa

20VW.A -A E ME RL-QQ N---Q VSD L ENK EMLH T -T FE
1A39 -Y T M TS RL-QH PD---G LPS LDI E EMLH T -F FE
6CEL STY T E MS S DF TQSA---HVG A A D TT QPFT L -F FE
1EG1.A A-S V E SS T -DI P S S C G T -S V S P L D S Q -G E V M L E N -S L S
consensus n-yGvTTsGnsLrl-qqlips-----n-rlvspRv-YLld-tk-kkIenbLtg-nBfsFD

20VW.A E E E M GA LSE WQD G-E---STB NSEK Y-----I
1A39 DAKK M SA LSE HPT E---S---IHP Y-----T
6CEL DTSQ GA FVS DAD G-V---SKYP THTA K-----T
1EG1.A DUSA E GS LEQ DEN G-A---NQ THTA N-----F
consensus VdastLPCGmGalfIsamd-dGg-k-----s-r--yntaGly-----IGTG

20VW.A A IV-TPFI G IK-----GV NEE W A RAHHA P
1A39 A FV-TPFI G IE-----GK S NEM W A RAHHA T
6CEL S PRDLPI C EGWEPSSHNANTGIGCH S SQM W A ISEALT P
1EG1.A A FV-QT R TL TS-----HQ F NEM L RAHALT S
consensus YCDaQ Cpv-tpfIHGvgNie-----gqGcCCnandIwBaNSrst-l-PHpC

20VW.A EPGL-YGC TDECG-----SSG K-A GW-ENR NTFD R-KQ KFD
1A39 HNRCL-Y C-EG EC-----EG K-N GW-NIR NTFD R-EEPKTM
6CEL TTVCQ-EG C-EGDCCGTYSD---NR GGT P-D DW-PIR GNTS P-SBPTD
1EG1.A TA-----TAE B-A G-PIGSGIES P--DTTD
consensus tk-gl-ylc-egdecg-----f-gICDk-sCCGw-Hpyr1-rt-ITG--G--f-wd


20VW.A TKEF VT AN---QGD-L E-H H DNKVIESA VVNISGPPK-IFINDKIC
1A39 TLEPF VV AN---EGK-LB-EIH V DGVIESFTTNKEGVPV-TNMIDD C
6CEL TTEKL VV ETS-----G-AIN V NGVTFQCPAE-LGSYS-GHE NDTIC
1EG1.A TKEF VT HTDNGSPSGN-L-SIT K Q NGVDIPSA P-----G-GDTI SSC--
consensus ttkkfTvtQFv-nk---g--li--ihRfivQ-g-wiesan-n--g-p--gn-indeye

20VW.A ARTG---NE--YML CTKQNGD SR AN WSGDF A QG-----
1A39 EATG---SE--YML CTQNGCE L R AN I WCGN E EG-----
6CEL TAEBAEPCSS---SK ELTQPER TEG VM DDTI N L STIPTNETS-B
1EG1.A-----PS--ASAY GLATNGF LSS VF I NC SQT N SG-----
consensus -atg---a-s-y-elGg--qngkAIsrGMVL-nSiWvdqggmN-WLDsg-----

20VW.A .....FA D DATE D KNITK NDE TFS IRI E
1A39 .....EA D AEGE A ENITQTE FPE TIRW E
6CEL TPGATR E S S S V A E S S Q S NAK TFS I P
1EG1.A .....NA D S E T E N S N I A N H T H V F S I R W
consensus .....wGpCstteG-Paniv-wqPnpvVtfsNirwGgIG-ST

20VW.A .....
1A39 .....YDEL-Q
6CEL .....GKPS-G
1EG1.A .....
consensus .....sq

```

4/39
Figure 4

C- α trace of the crystal structures from the catalytic domains of four Cel7 homologues aligned and overlaid as described.

Red = α -helix,

Cyan = disordered, Blue/Green = turns

Figure 5A: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant L225F

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValAlaIle
 CAGTCGGGCTT GCACTCTCCG ATCGGAGACT CACCCGGCTTC TGACATGGCA GAATAGCTCC TCTGGTGGCA CTTGCACTCA ACAGACAGGC TCCGGTGTCA
 101 .IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCGCTGG ACTCACGCTA CGAACAGCAG CACGAAGTGC TACGATGGCA ACACCTGGAG CTCGACCCTA TGTCTCCACA ACGAAGACCTG
 201 .AlaLysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGTGTGTCTGG ACGGTGCCGC CTACGCGTCC ACGTACGGAG TTACCACGAG CGGTACAGAG CTCCTCCATTG GCTTGTCCAC CCAAGTCGCG
 301 GlnLysAsnVal GlyAlaAlaArg LeuTyrLeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAAGAACG TTGGCGCTCC CTTTACCCTT ATGGCGAGCG ACACGACCTTA CCAAGAAATTC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGATGTTT
 401 .SglnLeuPro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr
 CGCAGCTGCC GTGCGGGCTTG AACGGAGCTC TCTACTTCTT GTCCATGGAC GCGGATGGTG GCGTGAAGCA GTATCCCAAC AACACGGCTG GGGCCAAAGTA
 501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleAsn GlyGlnAla AsnValGln GlyTyrGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGGATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGTGGGAGC GGTTCATCCAA CAACGCCAAC
 601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGluAla PheThrPro HisProCys ThrThrValGly
 ACGGGCATTTG GAGGACACCGG AAGTGTCTGC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTTTTACCCC CCACCCTTGC ACGACTGTGC
 701 .GlnGluIle CysGluIle AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
 GCCAGGAGAT CTGCGAGGGT GATGGGTGCG GCGGAATCTTA CTCGATTAAC AGATATGGCG GCACTTGGCA TCCCAGATGG TCGCAGTGGG ACCCATACCG
 801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CTTGGGCAAC ACCAAGCTTCT ACGGCCCTTG CTCAAGCTTT ACCCTCGATA CCACCAAGAA ATTAGCCGTT GTCAACCAGT TCGAGAGCTC GGGTGCATTC
 901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGln
 AACCGATACT AHTGCCAAGA TGGGCTCACT TTCCAGCAGC CCACAGCCCGA GCTTGGTAGT TACTCTGGCA ACGAGCTCAA CGATGATTAAC TGCACAGACTG
 1001 .GluAlaGln PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValLeuValMet SerLeuTrp
 AGGAGGACGA ATTCCGGCGGA TCTCTTCTTCT CAGACAAGGG CGGCTGACT CAGTTCAAGA AGGCTACCTC TGGCCGGATG GTTCTGGTCA TGAATCTGTG
 1101 .AspAspTyr TyrAlaAsnMet LeuTrpLeu AspSerThr TyrProThrAsn GlnThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTAAC TACGCCCAACA TGTGTGTGCT GGAATCCACC TACCAGACAA ACGAAGCCTC CTCACACACC GATGCCGTGC GCGGAAGCTG CTCCACCAAGC
 1201 SerGlyValPro AlaGlnVal GlnSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCCGGTGTCC TGTCTCAGGT CGAATCTCAG TCTCCCAACG CCAAGTCCAC CTTTCCCAAC ATCAAGTTGG GACCCAFITGG CAGCACCGGC AACCTTAGGG
 1301 .GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProgly ProThrGlnSer HisTyrGly
 GCGGCAACCC TCCCGGGCGGA AACCCGCTGG GCACCAACAC CACCCGCGCG CCAAGCACTA CCACGGAAG CTCTCCCGGA CCTACCCAGT CTCACACTAGG
 1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysLeu
 CCAAGTGGGG GGTATTTGGT ACAAGCGGCC CACGGTCTGC GCCAAGGGA CAACITGGCA GGTCTGAAC CCTTACTACT CTCAGTGGCT G

Figure 5B: Amino Acid and Nucleic Acid Sequence for *Hypocrea jecorina* Cel7A mutant S113D

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProIleu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCGGCCT GCACTCTCCA ATCGAGAGACT CACCGGCCCTC TGACATGGCA GAAATGCTCG TCTGTGGGCA CTGGCACTCA ACAGACAGGGC TCCGTTGGTCA

101 -IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnGly TyrAspGlyAsn ThrTrpSer SerThrIleu CysProIleAsn GluThrCys
 TCGACGCCAA CTGGCGCTGG ACTCAAGCTA GAAACAGCAG CACCAACTGC TACCAATGGCA ACACCTGGAG CTGCAGCCCTA TGTCTGTGACA ACCGACCTG

201 -AlaLysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGTCTGTCTGG ACCGTGCCCG CTACCGCTCC ACGTACGGAG TTACCAAGAG CGGTAAACAG CTCTCCATGTG GCTTTGTCAAC CCAAGTCTGGG

301 GlnLysAsnVal GlyAlaArg LeuTyrLeu MetAlaAspAsp ThrThrTyr GlnGlnPhe ThrLeuLeuGly AsnGlnPhe SerPheAsp ValAspValSer
 CAGAAAGAAC TTGGCGCTCG CCTTTAACCCTT ATGGCGGAGC ACACCACTTA CCAAGAAATC ACCCTGTGTG GCAACGAGTT CTCTTTGGAT GTTGAATGTTT

401 -SGlnLeuPro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerIys TyrProThr AsnThrAlaGly AlaLysTyr
 CGCAGCTGGC GTGGGGCTTG AACGGAAGCTC TCTACTTCCG GTCCATGGAC GCGGATGGTG GCGTGAAGCA GTATCCCAAC AACACCGCTG GCGCCAAAGTA

501 -GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleAsn GlyGlnAla AsnValGln GlyTyrGlnPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGCAATCTG AAGTTGATCA ATGGCCAGGC CAACGTTGAG GGCTGGGAGC CGTCAATCCA CAACGGCAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
 ACGGGCAATG GAGGACACCG AAGCTGTGTC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCAGG CTCTTACCCC CAACCCCTTG ACACCTGTG

701 -GGlnGlnIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrAsp
 GCCAAGAAAT CTGGAGAGGGT GATGGGTGCG GCGGAACCTTA CTCCGAAATAC AGATATGGCG GCACTTGCGA TCCCGATGGC TCGACATGGA ACCCATACCG

801 -LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrIlysIys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTGCATA CCAACCAAGAA ATTTGACCCTT GTCAACCCAGT TCGAGACCTG GGGTGGCCATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGln
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCCAGCAGC CCAACGCCGA GCTTGGTAAT TACTTGGCA ACGAGCTCAA CGATGATTAAC TGCACAGGCTG

1001 -GGluAlaGlu PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysIys AlaThrSer GlyGlyMet ValLeuValMet SerLeuTrp
 AAGAGGCAGA ATTCGGCGGA TCCCTCTTCT CAGACAAAGG CCGCCTGACT CAGTTCAAGA AGGCTACTCT TGGCGGGCANT GTTCTGTGCA TGAATCTGTG

*1101 -AspAspTyr TyrAlaAsnMet LeuThrIleu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTAAC TACGCCAACA TGTCTGTGCT GGACTGCACC TACCCGACAA ACGAGACTCT CTCCACACCC GGTGGCCGTG GCGGAAGCTG CTCCACACAGC

1201 SerGlyValPro AlaGlnVal GlnSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCCGGGTGTC CTGCTCAGGT CGAATCTCAG TCTCCCAAG CCAAGGTAC CTCTCCCAAC ATCAAGTTG GACCCATTTG GACCAACCGGC AACCTTAGCG

1301 -GlyLysPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GGGGCAACCC TCCGGGGCGA AACCGCCTG GCACCAACAC CACCCGGCCG CCAAGCCACTA CCACTGGAG CTCTCCCGGA CCAACCCAGT CTCACTACCG

1401 -GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysIleu
 CCAGTGCGGC GGTATTGGCT ACAGGGGCC CACGCTGTC GCCAGGGGA CAACTTGGCA GGTTCCTGAAC CCTTACTACT CTCAATGCTT G

Figure 5C: Amino Acid and Nucleic Acid Sequence for *Hypocrea jecorina* Cell7A mutant A77D

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProIeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCGGCCCT GCACTCTCCA ATGGAGAGACT CACCCCGCCCTC TGACATGGCA GAATAGCTCG TCTGGTGGCA CTTGGACTCA ACAGACAGGC TCCGTTGTCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrIeu CysProAspAsn GluThrCys.
 TCGACGCCAA CTGGCGCTGG ACTCAACGCTA CGAACACAGC CACGAAGTGC TACGATGGCA ACACCTGGAG CTCGACCCTA TGTCTTGACA ACGAGACTCG

201 .AlaLysasn CysCysLeuasp GlyAlaLasp TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAGAAC TGCCTGTCTGG ACGGTGCCGA CTACGCGTTC ACGTACGGAG TTACCACGAG CGGTAAACAG CTCCTCAATTG GCTTGTTCAC CGAGTCTGCG

301 GlnLysasnVal GlyAlaArg LeuTyrIeu MetAlaSerasp ThrThrTyr GlnGluPhe ThrIleuGly AsnGluPhe SerPheasp ValAspValSer.
 CAGAAGAACG TTGGCGCTCG CCTTTAACCTT ATGGCGAGCG ACACGACCTA CCAGGAATTC ACCCTGCTTG GCAACGAGTT CTCTTTGAT GTTGATGTTT

401 .SglnIeuPro CysGlyIeu AsnGlyAlaIeu TyrPheVal SerMetasp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr.
 CGCAGCTGCC GTGCGGCTTG AACGAGACTC TCTACTTCTGT GCCATGGAC GCGGATGGTG GCGTGAACAA GTATCCCAAC AACACCGCTG GCGCCAAATTA

501 .GlyThrGly TyrCysaspSer GlnCysPro ArgAspIeu LysPheIleasn GlyGlnAla AsnValGln GlyTyrGluPro SerSerasn AsnAlaasn
 CGGCACCGGG TACTGTGACA GCCAGTGTCC CCGCATCTG AAGTTATCA ATGGCCAGC CAACGTTGAG GCGTGGAGC GGTTCATCCA CAACGCGAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetasp IleTrpGln AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly.
 ACGGCAATTG GAGGACACCG AAGCTGTGTC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCGAGG CTCTTACCCC CCACCCTTGC ACGACTGTGC

701 .GlnGlnIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysasp ProaspGly CysAspTrpAsn ProTyrArg.
 GCCAGAGAAAT CTGGGAGGGT GATGGGTGCG GCGGAATTA CTCCGATAAC AGATATGGCG GCACTGGCA TCCCGATGGC TGGCACTGGA ACCCAITACCG

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrIleAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTCGATA CCACCAAGAA ATTAGCCGTT GTCAACCCAGT TCGAGACGCTC GGGTGGCAATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluIleAsn AspAspTyr CysThrAlaGln.
 AACCCGATACT ATGTCCAGAA TGGCGTCACT TTCCAGCAGC CCACCGCCGA GCTTGGTAAAT TACTCTGGCA ACGAAGCTCAA CGATGATTAC TGCACAGCTG

1001 .GglnAlaGln PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp.
 AGGAGGCAAG ATTCGGCGCGA TCCCTTTTCT CAGACAAAGG CGGCTGACT CAGTTCAGA AGGCTACTC TGGCGGCAAG GTTCTGTGTA TGAATCTGTG

1101 .AspAspTyr TyrAlaAsnMet LeuTrpIeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGAATGATTAC TAGCCACACA TGCCTGGGCT GACTCTCAC TACCCGCAAA ACGAGACTC CTCCACACCCC GGTGCCGTGC GCGGAAGCTG CTCCACACAGC

1201 SerGlyValPro AlaGlnVal GlnSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly.
 TCCGGTGTCC CTGCTCAGGT CGAATCTCAG TCTCCCAACG CCAAGTCAAC TTCTCTCAAC ATCAAGTTTCG GACCACATTCG CAGCACCGGC AACCTTAGCG

1301 .GgLYAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly.
 GCGGCAACCC TCCCGGCGGA AACCCGCGCTG GACACACAGC CACCCCGCCG CCAAGCCACTA CCACTGGAAG CTCTCCCGGA CCAACCCAGT CTCACTAAGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuAsn ProTyrTyrSer GlnCysIeu
 CGAGTGGCGG GATATTGGCT ACAGCGGCC CACGGTCTGC GCCAGCGGA CAACTTGCCA GGTCTGAAAC CCTTACTACT CTCAGTGCCT G

Figure 5D: Amino Acid and Nucleic Acid Sequences of *Hypocrea jecorina* Cel7A mutant L288f

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProlLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTGGCCCT GCACTCTCCA ATCCGAGACT CACCCCGCTC TGAATGGCA GAATAGCTCG TCTGGTGGCA CTTGCACTCA ACAAGACAGG CCGGTGGTCA

101 .IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCGCTGG ACTCACGCTA CGAACAGCAG CACAACCTGC TACCAATGGCA ACACCTGGAG CTCGACCCCTA TGTCTGACA ACGAGACTCG

201 .AlaLysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCGC CTACCGCTCC ACGTACGAG TTACCACAGAG CCGTACAGC CTCTCCATTG GCTTGTGAC CCACTCGCG

301 GlnLysAsnVal GlyAlaArg LeuTyrLeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAAGAAG TTGGCGCTCG CTTTACTT ATGGCCAGCG ACACGACCTA CCAGGAAATC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGATGTTT

401 .SerIleuPro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr
 CGCAGCTGCC GTGGCGCTTG AACGAGCTC TCTACTTCCGT GTCCATGGAC GCGGATGGTG GCGTGAAGCA GATCCGACC AACACCGCTG GGGCCAAAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleAsn GlyGlnAla AsnValGlu GlyTrpGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTACA GCCAGTGTCC CCGGATCTG AAGTTCATCA ATGGCCAGG CAAGTTGAG GGCTGGAGC CGTCATCCA CAACCGCAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGlu AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
 ACGGCAATG GAGGACACAGG AAGCTGTCC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTAACCC CCACCCCTTG ACGACTGTCG

701 .GlnGluIle CysGlnGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
 GCCAGGAGAT CTGGCAGGGT GATGGGTGG GCGGAACCTTA CTCCGATTAAC AGATATGGCG GCACCTGGCA TCCCGATGGC TGGACTGGA ACCCATACCG

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys PheThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTCGATA CCACCAAGAA ATTACCCGTT GTCAACCAGT TCGAGACGTC GGGTGCATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGlu LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGlu
 AACCGATACT ATGTCCAGAA TGGGTCTACT TTCAGCAGC CCAAGCCGA GCTTGGTAGT TACTCTGGCA ACGAGCTCA CGATGATTA TGCACAGCTG

1001 .GlnAlaGlu PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValLeuValMet SerLeuTrp
 AGGAGGCAAG ATTCGGCGGA TCCTCTTCT CAGCAAGGG CGGCTGACT CAGTCAAGA AGGCTACCTC TGGCGGCATG GTTCTGGTCA TGAAGTCTGG

1101 .AspAspTyr TyrAlaAsnMet LeuTrpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTA TACGCCAACA TGTGTGGGT GGACTCCACC TACCAGCAAA ACGAAGCTTC CTCCACACCC GGTGCCGTGC GCGGAAGCTG CTCCACCAAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProlleGly SerThrGly AsnProSerGly
 TCCGGTCTCC CTGCTCAGGT CGAATCTCA GTCGCCAAG CCAAGGTCA CTTCTCCAAC ATCAAGTTGG GACCCATTTG CAGCACCGGC AACCCTAGG

1301 .GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GCGGCAACCC TCCCGCGCGA AACCCCGCTG GCACCAACAC CACCCCGCC CAGCCACTA CCACGTGAAG CTCTCCCGGA CCTACCCAGT CTCACTAGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysLeu
 CCAAGTGGGC GGTATTGGCT ACGACGGCCC CAGGTTCTGC GCCAAGGGA CAACTGGCA GGTCTGAAC CCTTACTACT CTCAGTGGCT G

Figure 5E: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant A299E

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCGGCCCT GCACCTCTCCA ATCCGAGACT CACCCCGCTC TCACATGACA GAATGCTCG TCTGGTGGA CCTGCACATCA ACAGACAGGC TCCGTGGTCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCGCTGG ACTCACGCTA CGAACAGCG CAGCACTGC TACGATGGCA ACACCTTGGAG CTCCACCCCTA TGTCTGACA ACGAGACCTG

201 .AlaIysasn CysCysIleuasp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGTGTGTGG ACGGTGCCCG CTACGCGTCC ACGTAACGGAG TTTACACAGG CCGTAACAGC CTCTCCATTG GCTTTGTAC CCAGTCTGGC

301 GlnIysasnVal GlyAlaArg LeuTyrLeu MetAlaSerasp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheasp ValaspValSer
 CAGAGAAGC TTGGCGCTCG CCTTACCTT ATGGCGAGGG ACAACACTTA CCAGGAATTC ACCCTGCTTG GCAAGAGATT CTCTTTGGAT GTTGAATGTT

401 .SGlnLeupro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetasp AlaaspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaIysTyr
 CGCAGCTGCC GTGCGGCTTG AACGAGGCTC TCTACTTGGT GTCATGAGC GCGGATGGT GCGTGAACA GTATCCACACC AACACCGCTG GCGCCAAATA

501 .GlyThrGly TyrCysaspser GlnCysPro ArgAspLeu LysPheIleasn GlyGlnAla AsnValGln GlyTrpGluPro SerSerAsn AsnAlaasn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGCATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGTGGGAGC GGTCAATCCA CAACGGGAAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetasp IleTrpGln AlaasnSer IleSerGlnAla LeuThrPro HisProCys ThrThrValGly
 ACGGCGATTG GAGGACACGG AAGCTGCTGC TCTGAGATGG AATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTAACCC CCACCCTTG ACGACTGTGC

701 .GlnGlnIle CysGlnGly AspGlyCysGly GlyThrTyr Seraspasn ArgTyrGlyGly ThrCysasp ProaspGly CysaspTrpasn ProTyrArg
 GCCAGGAGAT CTGCCAGGGT GATGGGTGG GGGGAACCTA CTCCGATAC AGAATGGCG GCACCTGGCA TCCCGATGGC TGGCACTGGA ACCCATACCG

801 .LeuGlyasn ThrSerPheTyr GlyProGly SerSerPhe ThrIleuaspThr ThrIlyLys LeuThrVal ValThrGlnPhe GluThrSer GlyGlnIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTCGATA CCACCAAGAA ATTGAACGTT GTACCCAGT TCGAGACGTC GGGTGAATC

901 AsnArgTyrTyr ValGlnasn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyasn GluIleuasn AspaapTyr CysThrAlaGlu
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCACAGCAGC CCAACGCCGA GCTTGGTATG TACTGTGGCA ACGAGCTCAA CGATGATTAC TGACACAGCTG

1001 .GlnIleGln PheGlyGly SerSerPheSer AspIlyGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp
 AAGAGGCGAG ATTCGGCGGA TCCTCTTCTT CAGACAAAGGG CGGCTGACT CAGTTCAGA AGGCTACCTC TGGCGGCATG GTTCTGGTCA TGAGTCTGTG

1101 .AspaapTyr TyrAlaasnMet LeuTrpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTAC TACGCCAACA TGTGTGGGT GACTCCACC TACCCGACAA ACGAACCTC CTCCACACC GGTGCCGTGC GCGGAAGCTG CTCCACCCAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCCGGTGTCC CTGCTCAGGT CGAATCTGAG TCTCCCAAG CCAAGGTAC CTTCACCAAC ATCAAGTTTG GACCCATTGG CAGCACCGGC AACCCATTGG

1301 .GlyIasnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProIleThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GCGGCAACC TCCCGGGCGA AACCCGCTTG GCACCAACCAC CACCCGCGC CAGCCACATA CCACGTGGAAG CTCTCCCGGA CCTACCCAGT CTCACATACGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuasn ProTyrTyrSer GlnCysLeu
 CCAATGCGGC GGTATTGGCT ACAAGCGGCC CACGCTGTC GCCACGGGCA CAACTGGCCA GGTCTGAAAC CCTTACTACT CTCAGNGCCT G

Figure 5F: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant N301K

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProProLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTGGGCT GCACTCTCCA AFGGAGAGACT CACCCGCCCTC TGAATGGCA GAATGCTCG TCTGTGGCA CTTGCACTCA ACAGACAGGC TCCGTGGTCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnGly TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCCGCTGG ACTCAAGCTCA CGAACAAGCAG CACGAACCTG TACCATTGGCA ACACCTTGGAG CTCGACCCCTA TGTCTTGACA AGAGAAGCTG

201 .Alalysasn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly pheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCCG CTACGGCTCC ACGTACGGAG TTACCACGAG CGGTACAGC CTCTCCATTG GCTTGTTCAC CCAAGTCTGG

301 GlnLysAsnVal GlyAlaArg LeuTyrLeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAAAGAAC TTGGCGGCTCG CTTTAACTT ATGGCGAGCG ACACGACCTA CCAGGAATTC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGATGTTT

401 .SglnLeupro CysGlyLeu AsnGlyAlaLeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlalysTyr
 CGCAGCTGCC GTGCGGCTTG ACGGAGCTC TCTACTTCTGT GTCCAHGGAC GCGGATGGTG GCGTGAAGCA GTATCCCAAC AACACCGCTG GGGCCAAAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleasn GlyGlnAla AsnValGlu GlyTyrGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGCGATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGCTGGGAGC CGTCAATCCA CAACGCCAAG

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGlu AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
 ACGGGCAATG GAGGACACCG AAGTGTCTCG TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTACCCC CCAACCTTGC ACGACTGTC

701 .GglnGluIle CysGlnGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
 GCCAGGAGAT CTGGGAGGGT GATGGGTGGC GCGGAACCTA CTCCGATTAAC AGATATGGCG GCACTTGGGA TCCCGATGGC TGGCACTGGA ACCCAATACC

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTGGATA CCACCAAGAA ATTGACCGTT GTCAACCCAGT TCGAAGACCTC GGGTGGCCATC

901 LysArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGlu LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGlu
 AAGCGAFACT ATGCCACAGA TGGCGTCACT TTCCAGCAGC CCAACGCCGA GCTTGGTATG TACTCTGGCA ACGAGCTCAA CGATGATTAC TGCAACAGCTG

1001 .GglnAlaGlu PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValLeuValMet SerLeuTrp
 AAGAGGCAAG ATTGGCGCGA TCCCTTTTCT CAGACAAAGG CGGCTGACT CAGTTCAGA AAGCTTACCTC TGGCGGCATG GTTCTGTGTC TGAATCTGTG

1101 .AspAspTyr TyrAlaAsnMet LeuTrpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTAC TACGCCAACA TGCTGTGGCT GGACTCCAAC TACCCEGACA AGAAGACTCC CTCACACACC GGTGCCGTGC GCGGAAGCTG CTCCACACAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCGGGTGTCC CTGTCAAGT CGAATCTCAG TCTCCCAAG CCAAGTCAAC CTCTCCAAC ATCAAGTTGC GACCCATTTG CAGCAACGGC AACCTTAGCG

1301 .GglnAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GCGGCAACCC TCCCGGCGGA AACCGGCTG GCAACACCAAC CACCCGCGGC CAGGCCACTA CCACTGGAAG CTCTCCCGGA CTTACCCAGT CTCACTAACG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysLeu
 CCAAGTGGCG GGTATTTGGT ACAGCGGCC CACGGTCTGC GCCAGCGGCA CAACTTGCCA GGTCCCTGAC CTTACTACT CTCAAGTGGCT G

Figure 5G: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant T3561

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProPheLeu ThrTirpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTGGGCGCT GCACTCTCCA ATCCGAGACT CACCCTCCCTC TGACATGGCA GAATAGCTCG TCTGGTGGCA CTTGGCACTCA ACAGACAGGC TCCGTGGTCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrIleu CysProAspAsn GluThrCys.
 TCGACCGCCA CTGGCCGCTGG ACTCACGCTA CGAACAGCAG CACGAACTGC TACGATGGCA ACACCTGGAG CTCGACCCTA TGTCTTGACA ACGAGACTCG

201 .AlaIysasn CysCysIleuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCCG CTACCGCTCC ACGTACGGAG TTACCAAGAG CCGTACAGC CTCTCCATTTG GCTTTGTAC CCAGTCTGGG

301 GlnIysasnVal GlyAlaArg LeuTyrIleu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuIleGly AsnGluPhe SerPheAsp ValAspValSer.
 CAGAAAGAAC TTGGCGGCTGG CCTTTAACCTT ATGGGAGAGC ACACGACCTA CAGGAATTC ACCCTGGCTTG GCAACGAGTT CTCTTTGGAT GTTGATGTTT

401 .SglnLeuPro CysGlyIleu AsnGlyAlaIleu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaIysTyr.
 CGGACGCTGCC GTGGCGGCTTG AACGAGAGCTC TCTACTCTGT GTCCATGGAC GCGGATGGTG GCGTGAAGCA GTATCCGACC AACACCGGCTG GCGCCAAAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspIleu LysPheIleasn GlyGlnAla AsnValGlu GlyTirpGluPro SerSerAsn AsnAlaAsn
 CGGCAACGGGG TACTGTGACA GCCAGTGTCC CCGCAATCTG AAGTTCAATCA ATGGCCAGGC CAACGTTGAG GGCCTGGGAGC CGTCATTCGA CAACGGGAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGlnMetAsp IleTirpGlu AlaAsnSer IleSerGlnAla LeuThrPro HisProCys ThrThrValGly.
 ACGGGGCAITG GAGGACACCGG AAGCTGCTGC TCTGAGATGG AATCTGGGA GGGCAACTCC ATCTCCGAGG CTCTTACCCTC CCACCTTTGC ACGACTGTGC

701 .GlnGlnIle CysGlnGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTirpAsn ProTyrArg.
 GCCAGGAGAT CTGCCAAGGGT GATGGGTGG GGGGAAGCTTA CTCGGAATAC AAGTATGGGG GCACTTGGCA TCCCGATGGC TGGCACTGGA ACCCAATACCG

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAAGCTTT ACCCTCGATA CCAACCAAGAA ATTGAACCGTT GTCAACCCAGT TCGAGACGTC GGGTGGCATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGlu LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGlu.
 AACGATLACT ATGTCCAGAA TGGCGTCACT TTCACAGCAGC CCAACGCCGA GCTTGGTATG TACTCTGGCA ACGAGCTCA CGATGATTTAC TGCACAGACTG

1001 .GlnAlaGlu PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaLeuSer GlyGlyMet ValLeuValMet SerLeuTirp.
 AAGAGGCAGA ATTCGGCGGA TCCCTCTTCT CAGACAAAGG CCGGCTGACT CAGTTCAGA AGGCTCTCTC TGGCGGCATG GTTCTGGTCA TGAATCTGTG

1101 .AspAspTyr TyrAlaAsnMet LeuTirpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTTAC TACGCCACA TGTGTGTGGT GGACTCCACC TACCAGACA ACGAGACTTC CTCACACACC GGTGCCGTGC GCGGAGACTG CTCACACCAAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly SerThrSerGly.
 TCCGGTGTCC CTGCTCAGGT CGAATCTGAG TCTGCCAACG CCAAGTTCAC CTCTCCAAC ATCAAGTTCG GACCCATTTGG CAGCACCGGC AACCTTAGCG

1301 .GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly.
 GCGGCAACCC TCCCGGCGGA AACCCGCTG GACCCACCAAC CACCCGCGC CAGCCACTA CCACITGGAAG CTCTCCGGA CCTAACCAAGT CTCACACTAGCG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValLeuAsn ProTyrTyrSer GlnCysIleu
 CCAAGTGGGC GTTATTTGCT ACAAGCGGCC CACGGTCTGC GCCAGGGGA CAACTGGCA GGTCTGAAC CCTTACTACT CTCAGTGGCT G

Figure 5H: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant G430F

12/39

1
GlnSerAlaCys ThrLeuGln SerGluThr HisProPheLeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
CAGTCGGCCT GCACTCTCCA ATGGGAGACT CACCCGCCCTC TGACATGGCA GAAMAGCTCG TCTGGTGGCA CTTGGCACTCA ACAGACAGGC TCCGTTGGCA

101
-IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys
TCGACCGCCA CTGGCGCTGG ACTCAAGCTA CGAACAGCAG CACGAACTGC TACGATGGCA ACACCTTGGAG CTCGACCCTA TGCTCTGACA ACGAGACCTG

201
-AlaLysasn CysCysLeuasp GlyAlaIle TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
CGCGAAGAAC TGCTGTCTGG ACGGTGCCCG CTACGCCGTC ACGTAACGAG TTACCAACGAG CGGTACAGC CTCCTCATTTG GCTTTGTCA CCAGTCTGGC

301
GlnLysasnVal GlyAlaArg LeuTyrLeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheasp ValaspValSer
CAGAAGAAG TTGGCGCTCG CCTTACCTT ATGGCGAGCG ACACGACCTA CCAGGAATTC ACCCTGCTTG GCAACGAGTT CTCTTTGAT GTTGAATGTT

401
-SglnLeuPro CysGlyLeu AsnGlyAlaIleu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr
CGCAGCTGCC GTCGGGCTTG AACGAGCTC TCTACTTCTG GTCCATGGAC GCGGATGGTG GCGTGAACAA GATTCGCAAC AACACCGCTG GCGCCAAAGTA

501
-GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleasn GlyGlnAla AsnValGln GlyTrpGluPro SerSerAsn AsnAlaAsn
CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGGATCTG AAGTTGATCA ATGGCCAGGC CAACCTTGAAG GAGCTGGAGC GGTCAATCCA CAACGCGAAC

601
ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
ACGGGCATTTG GAGGACACCG AAGCTGCTGC TCTGAGATGG ATATCTGGGA GGCACACTCC ATCTCCGAGG CTCTTACCCC CAACCTTGGC ACGACTGTCG

701
-GlnGlnIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
GCCAGAGAT CTGGGAGGGT GATGGGTGCG GCGGAACCTA CTCGGAATAC AGATATGGCG GCACTTGGCA TCCCGATGGC TGGGACTGGA ACCCAATACCG

801
-LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAGACTTT ACCCTCGATA CCACCAAGAA ATTTGACCGTT GTCAACCCAGT TCGAGACCTC GGGTGGCART

901
AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGln
AACCGATCT ATGTCCAGAA TGGCGTCACT TTCCAGCAGC CCAACGCCGA GCTTGGTAT TACTCTGGCA ACGAGCTCA CGATGATTA TGCACAGCTG

1001
-GluAlaGln PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp
AGGAGGCAGA ATTGGGCGGA TCCTCTTCTT CAGACAAAGG GGGCTGACT CAGTTCAAGA AGGCTACTCTC TGGCGGCARTG GTTCTGTGTA TGAATCTGTG

1101
-AspAspTyr TyrAlaAsnMet LeuThrLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
GGATGATTA TACGCCAACA TGCTGTGGCT GACTCCACC TACCCGCAAA ACGAGACCTC CTCACACACC GGTGCCGTG GCGGAAGCTG CTCACACAGC

1201
SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrPhe AsnProSerGly
TCGGGTGTC CTGTCAAGT CGAATCTCAG TCTCCCAAG CCAAGGTCA CTTCTCCAAC ATCAAGTTG GACCACATTTG CAGCAACTTC AACCTAAGC

1301
-GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
GGGGCAACC TCCCGGCGGA AACCCGCTG GCAACCAACC CACCCGCCCG CCAAGCCACTA CCACTGGAAG CTCTCCCGGA CTAACCCAGT CTCACTAAGG

1401
-GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuAsn ProTyrTyrSer GlnCysLeu
CCAGTGGCGC GGFATTTGGT ACAAGCGCC CACGTTGCG GCCAGCGGA CAACTTGCCA GGTCCCTGAAC CCTTACTACT CTCAGTGGCT G

Figure 5I: Amino Acid and Nucleic Acid Sequence for *Hypocrea jecorina* Cel7A mutant T246C/371C

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProPoleu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle-
 CAGTGGGCTT GCACCTCTCCA ATCCGAGACT CACCCGGCTTC TGAATGAGGA GAATATGCTCG TCTGGTGGCA CTTCGACTCA ACAGACAGGC TCCGGTGTCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys-
 TCGACGCCAA CTGGCGCTGG ACTCACGGCTA CGAACAAGCA CAGAACCTGC TACGATGGCA ACACCTGGAG CTCCACCCCTA TGTCTTGACA ACGAGACCTG

201 .AlaIysasn CysCysLeuasp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValIThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGGCCG CTAAGCGCTGC ACGTACGGAG TTACCAACGAG CGGTAAACAG CTCTCCATTG GCTTGTTCAC CCAAGTCTGGC

301 GlnIysasnVal GlyAlaArg LeuTyrLeu MetAlaSerasp ThrThrTyr GlnGlnIphe ThrIleuLeuGly AsnGluIphe SerPheasp ValAspValSer-
 CAGAGAAGC TTGGCGCTCG CCTTACCTT ATGGCGAGCG ACACGACCTA CCAAGAAATTC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGAATGTTT

401 .SglnIleupro CysGlyIleu AsnGlyAlaIleu TyrPheVal SerMetasp AlaAspGlyGly ValSerIys TyrProThr AsnThrAlaGly AlaIysTyr-
 CGCAGCTGCC GTGGCGCTTG AACGAGACTC TCTACTTCTGT GTCCATGGAC GCGAATGGTG GCGTGAAGCAA GTATCCACACC AACACCGCTG GCGCCAAAGTA

501 .GlyThrGly TyrCysaspSer GlnCysPro ArgAspLeu LysPheIleasn GlyGlnAla AsnValGln GlyTyrGlnPro SerSerAsn AsnAlaasn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGCAATCTG AAGTTGATCA ATGGCCAGGC CAACGTTGAG GGCTGGGAGC CGTCAATCCA CAACGCCAAGC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetasp IleTrpGln AlaasnSer IleSerGlnAla LeuThrPro HisProCys ThrThrValGly-
 ACGGGCAATTG GAGGACACCG AAGCTGCTGC TCTGAATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTACCCC CCAACCTTGC ACGACTGTGC

701 .GlnGlnIle CysGlnGly AspGlyCysGly GlyCysTyr SerAspAsn ArgTyrGlyGly ThrCysasp ProAspGly CysAspTrpAsn ProTyrArg-
 GCCAAGAAAT CTGGGAGGT GATGGGTGC GCGGATGTTA CTCCGATVAC AAGATATGGCG GCACTTGGGA TCCCGATGGC TCGAAGTGG AACCATAACCG

801 .LeuGlyasn ThrSerPheTyr GlyProGly SerSerPhe ThrIleuAspThr ThrIlyIys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTCGATA CCAACCAAGAA ATTGACCGTT GTCAACCAAGT TCGAGACCTC GGGTGGCCATC

901 AsnArgTyrTyr ValGlnasn GlyValIThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluIleuAsn AspAspTyr CysThrAlaGln-
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCACGACAGC CCAACGCCGA GCTTGGTATG TACTCTGGCA ACGAGCTCA CGATGATVAC TGCACAGACTG

1001 .GlnAlaGln PheGlyGly SerSerPheSer AspIlyGly GlyLeuThr GlnPheIlyIys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp-
 AGAGGCAGA ATTCGGCGCGA TCCCTCTTCT CAGACAAAGG CGGCTTCACT CAGTTCAAGA AGGCTAACCTC TGGCGGCATG GTTCTGTGTA TGAGTCTGTG

1101 .AspAspTyr CysAlaAsnMet LeuTrpLeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATVAC TGGCGCAACA TGCTGTGGCT GGAATCCACC TACCCGACAA ACGAGACCTC CTCCACACCC GGTGCCGTGC GCGAAAGCTG CTCCACCAAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValIThr PheSerAsn IleIysPheGly ProIleGly SerThrGly AsnProSerGly-
 TCCGGTGTCC CTGCTCAGGT CGAATCTGAG TCTCCCAAGC CCAAGGTCACT CTCTCCAC ATCAAGTTCG GACCCATTTG CAGCACCGGC AACCCATAGCG

1301 .GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly-
 GCGGCAACCC TCCCGGCGGA AACCGCGCTG GCACCAACAC CACCGGCCCG CCAAGCACTA CCACGGGAAG CTCTCCCGGA CCTAACCAAG CTCACTACGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuAsn ProTyrTyrSer GlnCysIleu
 CCAAGTCCGGC GGTATTTGCT ACAAGCGGCC CACGGTCTGC GCCAAGGCA CAACTTGCCA GTCTCTGAAC CCTTACTACT CTCAAGTGGCT G

Figure 5J: Amino Acid and Nucleic Acid Sequence for *Hypocrea jecorina* Cel7A mutant T246A/R251A/Y252A

1 GlnSerAlaCys ThrIleuGln SerGluThr HisProProIleu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTGGGCTT GCACTCTCCA ATCCGAGACT CACCCGCCCTC TGAATGCTCG TCTGGTGGCA CTGCACTCA ACAGACAGGC TCCGTTGGTCA

101 .IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrIleu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGCGCTGG ACTCACGCTTA CGAACAGCAG CACGAATCTC TACGAATGGCA ACACCTGGAG CTCCAGCCCTA TGTCTTGACA ACGAGACTGG
 CCGAAGAAGC TGCCTGTCTGG ACGGATGCCG CTACGCCCTCC ACGTACGGAG TTACCAAGAG CCGTAAACAGC CTCTCCATTTG GCTTTGTAC CCAAGTCTGG

201 .AlaIysAsn CysCysIleuAsp GlyAlaAla TyrAlaSer ThrTrpGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 GlnIysAsnVal GlyAlaArg LeuTyrIleu MethIaSerAsp ThrThrTyr GlnGluPhe ThrIleuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAAAGACG TTGGCGCTTCG CCTTTACCTT ATGGGAGCG ACACGACCTA CCAGGAATTC ACCCTGGCTTG GCAACGAGTT CTCTTTGGAT GTTGATGTTT

301 .GlnIleuPro CysGlyIleu AsnGlyAlaIleu TyrPheVal SerMetAsp AlaAspGlyGly ValSerIys TyrProThr AsnThrAlaGly AlaIysTyr
 CGCAGCTGGC GTGGCGCTTG AACGAGACTC TCTACTTCTG TTCCAATGGAC GCGGATGGTG GCGTGAACAA GTATCCGACC AACACCGCTG GCGCCAAAGTA

401 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspIleu LysPheIleAsn GlyGlnAla AsnValGln GlyTrpGluPro SerSerAsn AsnAlaAsn
 CCGCACCGGG TACTGTGACA GCCAGTGTCC CCGCGATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGCTGGGAGC CGTTCATCCA CAACGCCAAGC

501 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly
 ACGGGCATTG GAGGACACCG AAGCTGTGTC TCTGAATGAG ATATCTGGGA GCCCACTCC ATCTCCAGG CTCTTACCC CCACCCCTGC ACGACTGTC

601 .GGlnGluIle CysGluGly AspGlyCysGly GlyAlaTyr SerAspAsn AlaAlaGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg
 GCCAGGAGAT CTGCCAGGGT GATGGGTGCC GCGGAGCTTA CTCCGATPAC GAGAGCTGGC GCACTTGCCA TCCCGATGGC TGGCACTGGA ACCCATPACC

701 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrIleuAspThr ThrIleuVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTCGATA CCACCAAGAA ATTGACCCTT GTCACCCAGT TCGAAGACTC GGGTGCATC

801 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluIleuAsn AspAspTyr CysThrAlaGln
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCACAGCAGC CCNACGCCGA GCTTGGTATG TACTCTGGCA ACGAGCTCA CGATGATTAAC TGCACAGACTG

901 .GluAlaGln PheGlyGly SerSerPheSer AspIysGly GlyLeuThr GlnPheIysAla ThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 AGGAGGCAGA ATTCGGCGGA TCCTCTTCTT CAGACAAAGG CCGGCTGACT CAGTTCAAGA AGGCTACCTC TGGGGGCATG GTTCTGGTCA TGAGTCTGTG
 GGATGATTAAC TACGCCAACA TECTGTGGCT GGAATCCACC TACCCGACA ACGAGACTC CTCACACACC GGTGCCGTGC GCGGAGACTG CTCACACAGC

1101 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly
 TCCGCTGTCC CTGCTCAAGT CGAATCTGAG TCTCCCAAG CCAAGTAC CTCTCCAAAC ATCAAGTTCC GACCATTGG CAGCACCGGC AACCTHAGC

1201 .GlyAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProIaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly
 GCGGCAACCC TCCCGGGGGA AACCCGCTCG GCAACACAC CACCCGCCG CCAGCCACTA CCACGGAAG CTCTCCCGGA CCTACCCAGT CTCACATACG

1301 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuAsn ProTyrTyrSer GlnCysIleu
 CCAGTGGGCG GGTATTGGCT ACGAGCGGCC CACGCTCTGC GCCAGCGGA CACTTGGCA GGTCTGAAC CCTACTACT CTCAGTGCCT G

1401

Figure 5K: Amino Acid and Nucleic Acid Sequence of Hypocrea jecorina Cel7A mutant T380G/Y381D/R394A

1 GlnSerAlaCys ThrLeuGln SerGluThr HisProPoleu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTGGGCGCT GCACCTCTCCA ATCGGAGACT CACCCCGCCTC TGCATGAGGA GAATATGCTCG TCGTGGTGGCA CTTGCACTCA ACAGACAGGC TCCGTRGATCA

101 .IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrLeu CysProAspAsn GluThrCys.
 TCGAGGCCAA CTGGCGCTGG ACTCACGCTA CGAACAAGCA CAGGAACCTGC TACCATGGCA ACACCTTGGAG CTGCACCCCTA TGTCTTGACA ACGAGCCTG

201 .AlaIysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValIThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCGC CTACGCGCTCC ACCTACGGAG TTAACAAGAG CGGTRACAGC CTCACCATG GCTTGTCTAC CCAATCTGGCG

301 GlnIysAsnVal GlyAlaArg LeuTyrIleu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrLeuLeuGly AsnGluPhe SerPheAsp ValAspValSer.
 CAGAGAAGAC TTGGCGGCTCG CCTTTACCTT ATGGCGAGCG ACACGACCTA CCAAGAAATTC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGAATGTTT

401 .SglnIleuPro CysGlyLeu AsnGlyAlaIleu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaIysTyr.
 CGCAGCTGCC GTGGCGGCTTG ACGGAGACTC TCTACTTCTT GTCCATGGAC GCGGATGGTG GCGTGAAGCAA GTATCCACACC AACACCGGCTG GCGCCAAAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspLeu LysPheIleAsn GlyGlnAla AsnValGln GlyTyrGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAGTGTCC CCGGATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGCCTGGAGC GGTCAATCCA CAACGCCAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGluAla LeuThrPro HisProCys ThrThrValGly.
 ACGGGCAATG GAGGACACCG AAGCTGTGC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTACC CCACCTTGC AGCATGTGCG

701 .GlnGlnIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProTyrArg.
 GCCAAGAAAT CTGGGAGGGT GATGGGTGCG GCGGAACCTTA CTCGAAATAC AAGATATGGCG GCACTTGGCA TCCCGAATGC TCGGACTGGA ACCCATACCG

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrIlysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCCAAC ACCAGCTTCT ACGGCCCTGG CTCAAGCTTT ACCCTGCATA CCAACAGAA ATTGACCGTT GTCACCCAGT TCGAGACGCTC GGGTGGCCATC

901 .AsnArgTyrTyr ValGlnAsn GlyValIThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGln.
 AACCGAATACT ATGTCCAGAA TGGCGTCACT TTCCAGGACG CCAACGCCGA GCTTGGTATG TACTCTGGCA ACGAGCTCAA CGAATGATTAC TGCACAGCTG

1001 .GlnValGln PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp.
 AAGAGGCAGA ATTGGCGGGA TCCCTCTTCT CAGACAAAGG CGGCCTGACT CAGTTCAAGA AAGCTTACTTC TGGCGGGCATG GTTCTGTGTA TGAATCTGTG

101 .AspAspTyr TyrAlaAsnMet LeuTrpLeu AspSerGly AspProThrAsn GluThrSer SerThrPro GlyAlaValAla GlySerCys SerThrSer
 GGATGATTAC TACGCCAACA TGCTGTGGCT GAACCTCCGGC GACCCGACAA ACGAGACCTC CTCACACACC GGTGCCGTGG CCGGAAGCTG CTCACACAGC

1201 SerGlyValPro AlaGlnVal GluSerGln SerProAsnAla LysValIThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly.
 TCCGGTGTCC CTGTCTCAGG GAATCTCAG TCTCCCAAG CCAAGGTCA CTTCTCCAAC ATCAAGTTCG GACCCATGG CAGCAACGGC AACCTTAGCG

1301 .GlyIysAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly.
 GCGGCAACCC TCCCGCGGGA AACCCGCTCG GCAACCAACA CACCCGCGC CAGCACACTA CCACTGGAG CTTCCCGGA CCTACCAGT CTCACATACGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIleuAsn ProTyrTyrSer GlnCysLeu
 CCAATGCGGC GATATGACT ACAAGCGCC CAGGATGCG CCAAGCGGCA CAACTTGCCA GGTCTGAAC CCTTACTACT CTCAATGCT G

Figure 5L: Amino Acid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant T380G/Y381D/R394A with residues 381 through 393 deleted

1 GlnSerAlaCys ThrIleuGln SerGluThr HisProProIleu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCGGCCCT GCACCTCTCCA ATCCGAGACT CACCCGGCCTC TGACATGCGA GAAATGCTCG TCTGGTGGCA CTTGCACCTCA ACAAGACAGGC TCCGTGGTCA

101 .IAspAlaAsn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrIleu CysProAspAsn GluThrCys
 TCGACGCCAA CTGGGGCTGG ACTCAGCCTA CGAACAGCAG CACGAACCTGC TACGATGGCA ACACTTGGAG CTGCACCCCTA TGTCTTGACA ACAAGACCTG

201 .AlaLysAsn CysCysLeuAsp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer LeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCCG CTACGCTGCC ACGTACGGAG TTAACACAGAG CCGTAACAGC CTCTCCATFG GCTTTGTTCAC CCAGTCTGCC

301 GlnLysAsnVal GlyAlaArg LeuYrIleu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrIleuLeuGly AsnGluPhe SerPheAsp ValAspValSer
 CAGAAGAACG TTGGCGCTCG CCTTACCTT ATGGCCGAGCG ACAAGACCTA CCAAGGAAATC ACCCTGCTTG GCAACGAGTT CTCTTTCGAT GTTGATGTTT

401 .SglnLeuPro CysGlyLeu AsnGlyAlaIleu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr
 CGCAGCTGCC GTGGCGCTTG AACGAGGCTC TCTACTTGGT GTCCATGGAC GCGGATGGNG GCGTGAAGCA GTATCCACC ACCACCGCTG GCGCCAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspIleu LysPheIleAsn GlyGlnAla AsnValGln GlyTrpGluPro SerSerAsn AsnAlaAsn
 CGGCACGGGG TACTGTGACA GCCAAGTACC CCGCGATCTG AAGTTCATCA ATGGCCAGGC CAACGTTGAG GGCTGGGAGC CGTCATCCAA CAACGCCAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGlnMetAsp IleTrpGln AlaAsnSer IleSerGlnAla LeuThrPro HisProCys ThrThrValGly
 ACGGGCATTG GAGGACACGG AAGCTGCTGC TCTGAGATGG ATATCTGGGA GGCCAACTCC ATCTCCGAGG CTCTTACCCC CCACCTTTGC ACGACTGTGC

701 .GlnGluIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgTyrGlyGly ThrCysAsp ProAspGly CysAspTrpAsn ProYrArg
 GCCAGAGAT CTGCGAGGGT GATGGGTGG GCGGAACCTTA CTCCGATTAC AGATATGGCG GCACTTGCGA TCCCGATGGC TGCAGACTGGA ACCCATACCG

801 .LeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrIleuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CCTGGGCCAC ACCAGCTTCT ACGGCCCTTG CTCAAGCTTT ACCCTGATA CCACCAAGAA ATTGACCGTT GTCAACCCAGT TCGAAGACGTC GGTGGCCATC

901 AsnArgTyrTyr ValGlnAsn GlyValIleThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluIleuAsn AspAspTyr CysThrAlaGlu
 AACCGATTACT ATGTCCAGAA TGGCGTCACT TTCCAGCAGC CCACGCCCGA GCTTGGTAGT TACTCTGGCA ACGAGCTTCA CGATGATTAC TGCACAGCTG

1001 .GlnValGln PheGlyGly SerSerPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIleuValMet SerLeuTrp
 AGGAGGCAGA ATTCGGCCGG TCTCTTTCT CAGACAAAGG CCGCCTGACT CAGTTCAGA AGGCTTACCTC TGGCCGGCANG GTTCTGTCA TAGTCTGTG

1101 .AspAspTyr TyrAlaAsnMet LeuThrIleu AspSerGly AspAlaGlySer CysSerThr SerSerGly ValProAlaGln ValGluSer GlnSerPro
 GGATGATTAC TACGCCAACA TGCTGTGGCT GACTCTCGGC GACGCCGGA GGTGCTCCAC CAGCTCCGGT GTCCCTGTCTC AGGTGGAATC TCAGTCTCCC

1201 AsnAlaLysVal ThrPheSer AsnIleLys PheGlyProIle GlySerThr GlyAsnPro SerGlyGlyAsn ProProGly GlyAsnPro ProGlyThrThr
 AACGCCAAGG TCACCTTCTC CAACATCAAG TTGGGAGCCA TTGGCAGAGAC GGGCAACCTT AGGAGCCGGA ACCCTCCCAG CCGAAACCCG CTTGGCACCA

1301 .ThrThrArg ArgProAla ThrThrThrGly SerSerPro GlyProThr GlnSerHisTyr GlyGlnCys GlyGlyIle GlyTyrSerGly ProThrVal
 CCACCACCCG CCGCCAGACC ACTACCACTG GAAAGTCTCC CGGAACCTACC CAGTCTCACT ACCGCCAAGNG CCGCGGTATT GGTACACAGC GCCCCACGGT

1401 .CysAlaSer GlyThrThrCys GlnValIleu AsnProTyr TyrSerGlnCys Leu
 CTGGGCCAGC GGCACAACTT GCCAGGTCTT GAACCTTAC TACTCTCAAT GCTTG

Figure 5M: AminoAcid and Nucleic Acid Sequence of *Hypocrea jecorina* Cel7A mutant Y252Q/D259W/S342Y

1 GlnSer1Acys ThrLeuGln SerGluThr HisProProIeu ThrTrpGln LysCysSer SerGlyGlyThr CysThrGln GlnThrGly SerValValIle
 CAGTCCGGCCT GCACTCTCCA ATCCGAGACT CACCCGCCCTC TGACATGGCA GAATCTCTCG TCTGGTGGCA CTTGCACTCA ACAGACAGGC TCCCGTGTGCA

101 .IAspAlaasn TrpArgTrp ThrHisAlaThr AsnSerSer ThrAsnCys TyrAspGlyAsn ThrTrpSer SerThrIeu CysProAspAsn GluThrCys.
 TCGACCGCCA CTGGCGCTGG ACTCAACGCTA CGAACAAGCAG CACGAACTGC TACGATGGCA ACACCTGGAG CTCGACCCTA TGTCTGTACA ACGAGACTGT

201 .AlaLysasn CysCysLeuasp GlyAlaAla TyrAlaSer ThrTyrGlyVal ThrThrSer GlyAsnSer IeuSerIleGly PheValThr GlnSerAla
 CGCGAAGAAC TGCTGTCTGG ACGGTGCCCG CTACCGCTCC ACGTACGGAG TTAACAAGAG CCGTAAACAG CTCCTCAATTG GCTTTGTAC CCAAGTCTGGG

301 GlnLysasnVal GlyAlaArg LeuTyrIeu MetAlaSerAsp ThrThrTyr GlnGluPhe ThrIeuLeuGly AsnGluPhe SerPheAsp ValAspValSer.
 CAGAAGAAGC TTGGCGCTCG CCTTAACTT ATGGGAGCG ACACGACCTA CCAGGAATTC ACCCTGCTTG GCAACGAGTT CTCTTGGAT GTTGATGTTT

401 .SglnLeupro CysGlyIeu AsnGlyAlaIeu TyrPheVal SerMetAsp AlaAspGlyGly ValSerLys TyrProThr AsnThrAlaGly AlaLysTyr.
 CGCAGCTGGCC GTGGGGCTTG AACGGAGCTC TCTACTTCTG GTCCATGGAC GCGGATGGTG GGGTANGCA GATCCCAAC AACACCGCTG GCGCCAAGTA

501 .GlyThrGly TyrCysAspSer GlnCysPro ArgAspIeu LysPheIleAsn GlyGlnAla AsnValGln GlyTrpGluPro SerSerAsn AsnAlaAsn
 CGGCACCGGG TACTGTGACA GCCAGTGTCC CCGCGATCTG AAGTTGATCA ATGGCCAGGC CAACGTTGAG GGGCTGGAGC CGTCATCCA CAACGCGAAC

601 ThrGlyIleGly GlyHisGly SerCysCys SerGluMetAsp IleTrpGln AlaAsnSer IleSerGlnAla LeuThrPro HisProCys ThrThrValGly.
 ACGGCGATTG GAGGACACCG AACCTGCTGC TCTGAGATGG ATATCTGGGA GGCCAACTCC AICTCCAGAG CTCTTACCCT CCACCTTTC ACCGACTGTC

701 .GglnGlnIle CysGluGly AspGlyCysGly GlyThrTyr SerAspAsn ArgGlnGlyGly ThrCysAsp ProTrpGly CysAspTrpAsn ProTyrArg.
 GCCAAGAGAT CTGGGAGGGT GATGGGTGCG GCGGAACCTA CTCGGAATAC AGACAAGGCG GCACTTGGCA TCCCTGGGGC TGGGACTGGA ACCCATACCG

801 .IeuGlyAsn ThrSerPheTyr GlyProGly SerSerPhe ThrLeuAspThr ThrLysLys LeuThrVal ValThrGlnPhe GluThrSer GlyAlaIle
 CTGGGGCAAC ACCAGCTTCT ACGGGCCCTGG CTCAGCTTT ACCCTGCATA CCACCAAGAA ATTGACCCTT GTCACCCAGT TCGAGACGTC GGGTGCATC

901 AsnArgTyrTyr ValGlnAsn GlyValThr PheGlnGlnPro AsnAlaGln LeuGlySer TyrSerGlyAsn GluLeuAsn AspAspTyr CysThrAlaGln.
 AACCGATACT ATGTCCAGAA TGGCGTCACT TTCCAGAGAC CCAACGCCGA GCTTGTGAT TACTCTGGCA ACGAGCTGA CGATGATTA TGCACAGCTG

1001 .GglnAlaGln PheGlyGly SerTyrPheSer AspLysGly GlyLeuThr GlnPheLysLys AlaThrSer GlyGlyMet ValIeuValMet SerLeuTrp.
 AAGAGGCAGA ATTCGGCGGA TCCCTATTCT CAGACAAGGG CGGCTGACT CAGTTCAGA AGGCTACTC TGGCGGCATG GTTCTGTCA TGAATCTGTG

1101 .AspAspTyr TyrAlaAsnMet LeuTrpIeu AspSerThr TyrProThrAsn GluThrSer SerThrPro GlyAlaValArg GlySerCys SerThrSer
 GGATGATTA TACGCCAACA TGCTGTGGCT GAACTCAAC TACCCGACAA ACGAGACTTC CTCCACACCC GGTGCCGTGC GCGGAAGCTG CTCACACAGC

1201 SerGlyValPro AlaGlnVal GlnSerGln SerProAsnAla LysValThr PheSerAsn IleLysPheGly ProIleGly SerThrGly AsnProSerGly.
 TCCGGTGTCC CTGCTCAGGT CGAATCTCAG TCTCCCAAG CCAAGGTAC CTCTCCAAC ATCAAGTTCC GACCATTGG CAGCACCGGC AACCTTAGCG

1301 .GglnAsnPro ProGlyGly AsnProProGly ThrThrThr ThrArgArg ProAlaThrThr ThrGlySer SerProGly ProThrGlnSer HisTyrGly.
 GCGGCAACCC TCCCGCGCGA AACCCGCGCTG GACACCAAC CACCCGCGCC CCAAGCCACTA CCATGGAAG CTCTCCCGGA CTAACCAAGT CTACTACGG

1401 .GlnCysGly GlyIleGlyTyr SerGlyPro ThrValCys AlaSerGlyThr ThrCysGln ValIeuAsn ProTyrTyrSer GlnCysIeu
 CCAATGGCGC GATATTGGCT ACAGCGGCC CACGATGCG GCAAGCGGA CAACTTGCCA GGTCTGAAC CCTTACTACT CTCAGTGGCT G

FIGURE 6A: pTEX2

AAGCTTAAGG	TGCACGGCCC	ACGTGGCCAC	TAGTACTTCT	CGAGCTCTGT	50
ACATGTCCGG	TCGCGACGTA	CGCGTATCGA	TGGCGCCAGC	TGCAGGCGGC	100
CGCCTGCAGC	CACTTGCAGT	CCCGTGG AAT	TCTCACGGTG	AATGTAGGCC	150
TTTTGTAGGG	TAGGAATTGT	CACTCAAGCA	CCCCAACCT	CCATTACGCC	200
TCCCCATAG	AGTCCCAAT	CAGTGAGTCA	TGGCACTGTT	CTCAAATAGA	250
TTGGGGAGAA	GTTGACTTCC	GCCCAGAGCT	GAAGGTCGCA	CAACCGCATG	300
ATATAGGGTC	GGCAACGGCA	AAAAAGCACG	TGGCTCACCG	AAAAGCAAGA	350
TGTTTTGCGAT	CTAACATCCA	GGAACCTGGA	TACATCCATC	ATCACGCACG	400
ACCACTTTGA	TCTGCTGGTA	AACTCGTATT	CGCCCTAAAC	CGAAGTGCCT	450
GGTAAATCTA	CACGTGGGCC	CCTTTCGGTA	TACTGCGTGT	GTCTTCTCTA	500
GGTGCCATTC	TTTTCCCTTC	CTCTAGTGT	GAATTGTTTG	TGTTGGAGTC	550
CGAGCTGTAA	CTACCTCTGA	ATCTCTGGAG	AATGGTGGAC	TAACGACTAC	600
CGTGCACCTG	CATCATGTAT	ATAATAGTGA	TCCTGAGAAG	GGGGGTTTGG	650
AGCAATGTGG	GACTTTGATG	GTCATCAAAC	AAAGAACGAA	GACGCCTCTT	700
TTGCAAAGTT	TTGTTTCGGC	TACGGTGAAG	AACTGGATAC	TTGTTGTGTC	750
TTCTGTGTAT	TTTTGTGGCA	ACAAGAGGCC	AGAGACAATC	TATTCAAACA	800
CCAAGCTTGC	TCTTTTGAGC	TACAAGAACC	TGTGGGGTAT	ATATCTAGAG	850
TTGTGAAGTC	GGTAATCCCG	CTGTATAGTA	ATACGAGTCG	CATCTAAATA	900
CTCCGAAGCT	GCTGCGAACC	CGGAGAATCG	AGATGTGCTG	GAAAGCTTCT	950
AGCGAGCGGC	TAAATTAGCA	TGAAAGGCTA	TGAGAAATTC	TGGAGACGGC	1000
TTGTTGAATC	ATGGCGTTCC	ATTCTTCGAC	AAGCAAAGCG	TTCCGTGCGA	1050
GTAGCAGGCA	CTCATTCCCG	AAAAA ACTCG	GAGATTCCTA	AGTAGCGATG	1100
GAACCGGAAT	AATATAATAG	GCAATACATT	GAGTTGCCTC	GACGGTTGCA	1150
ATGCAGGGGT	ACTGAGCTTG	GACATAACTG	TTCCGTACCC	CACCTCTTCT	1200
CAACCTTTGG	CGTTTCCCTG	ATTCAGCGTA	CCCGTACAAG	TCGTAATCAC	1250
TATTAACCCA	GACTGACCGG	ACGTGTTTTG	CCCTTCATTT	GGAGAAATAA	1300
TGTCATTGCG	ATGTGTAATT	TGCCTGCTTG	ACCGACTGGG	GCTGTTGCGA	1350
GCCCGAATGT	AGGATTGTTA	TCCGAACTCT	GCTCGTAGAG	GCATGTTGTG	1400
AATCTGTGTC	GGGCAGGACA	CGCCTCGAAG	GTTACGCGCA	AGGGAAACCA	1450
CCGATAGCAG	TGTC'TAGTAG	CAACCTGTAA	AGCCGCAATG	CAGCATCACT	1500
GGAAAATACA	AACCAATGGC	TAAAAGTACA	TAAGTTAATG	CCTAAAGAAG	1550
TCATATACCA	GCGGCTAATA	ATTGTACAAT	CAAGTGGCTA	AACGTACCGT	1600
AATTTGCCAA	CGGCTTGTGG	GGTTGCAGAA	GCAACGGCAA	AGCCCCACTT	1650
CCCCACGTTT	GTTTCTTCAC	TCAGTCCAAT	CTCAGCTGGT	GATCCCCCAA	1700
TTGGGTGCGT	TGTTTGTTC	GGTGAAGTGA	AAGAAGACAG	AGGTAAGAAT	1750
GTCTGACTCG	GAGCGTTTTG	CATACAACCA	AGGGCAGTGA	TGGAAGACAG	1800
TGAAATGTTG	ACATTCAAGG	AGTATTTAGC	CAGGGATGCT	TGAGTGTATC	1850
GTGTAAGGAG	GTTTGTCTGC	CGATACGACG	AATACTGTAT	AGTCACTTCT	1900
GATGAAGTGG	TCCATATTGA	AATGTAAGTC	GGCACTGAAC	AGGCAAAAGA	1950
TTGAGTTGAA	ACTGCCTAAG	ATCTCGGGCC	CTCGGGCCTT	CGGCCTTTGG	2000
GTGTACATGT	TTGTGCTCCG	GGCAAATGCA	AAGTGTGGTA	GGATCGAACA	2050
CACTGCTGCC	TTTACCAAGC	AGCTGAGGGT	ATGTGATAGG	CAAATGTTCA	2100
GGGGCCACTG	CATGGTTTCG	AATAGAAAGA	GAAGCTTAGC	CAAGAACAAT	2150
AGCCGATAAA	GATAGCCTCA	TTAAACGGAA	TGAGCTAGTA	GGCAAAGTCA	2200
GCGAATGTGT	ATATATAAAG	GTTTCGAGTC	CGTGCCTCCC	TCATGCTCTC	2250
CCCATCTACT	CATCAACTCA	GATCCTCCAG	GAGACTTGTA	CACCATCTTT	2300
TGAGGCACAG	AAACCCAATA	GTCAACCGCG	GTTTAGGCGC	GCCAGCTCCG	2350
TGCGAAAGCC	TGACGCACCG	GTAGATTCTT	GGTGAGCCCG	TATCATGACG	2400
GCGGCGGGAG	CTACATGGCC	CCGGGTGATT	TATTTTTTTT	GTATCTACTT	2450

FIGURE 6B: pTEX2

CTGACCCTTT	TCAAATATAC	GGTCAACTCA	TCTTTCCTG	GAGATGCGGC	2500
CTGCTTGGTA	TTGCGATGTT	GTCAGCTTGG	CAAATTGTGG	CTTTCGAAAA	2550
CACAAAACGA	TTCCTTAGTA	GCCATGCATT	TTAAGATAAC	GGAATAGAAG	2600
AAAGAGGAAA	TTAAAAAAA	AAAAAAAACA	AACATCCCGT	TCATAACCCG	2650
TAGAATCGCC	GCTCTTCGTG	TATCCAGTA	CCAGTTTAAA	CGGATCTCAA	2700
GCTTGCATGC	AAAGATACAC	ATCAATCGCA	GCTGGGGTAC	AATCATCCAT	2750
CATCCCAACT	GGTACGTCAT	AACAAAAATC	GACAAGATGG	AAAAAGAGGT	2800
CGCCTAAATA	CAGCTGCATT	CTATGATGCC	GGGCTTTGGA	CAAGAGCTCT	2850
TTCTCAGCTC	CGTTTGTCTT	CCCTCCCTTT	TCCCCCTTCT	TGCTAAATGC	2900
CTTCTTTTAC	TTCTTTCTTC	CCTTCCCTCC	CCTATCGCAG	CAGCCTCTCG	2950
GTGTAGGCTT	TCCACGCTGC	TGATCGGTAC	CGCTCTGCCT	CCTCTACGGG	3000
GTCTGAGGCC	TTGAGGATGC	CCCGGCCAC	AATGGCAATG	TCGCTGCCGG	3050
CGATGCCAAT	CAGCTTGTGC	GGCGTGTGT	ACTGCTGGCC	CTGGCCGTCT	3100
CCACCGACCG	ATCCGTTGGT	CTGCTGGTCC	TCGTCTTCGG	GGGGCAGCTG	3150
GCAGCCGGGC	GTCATGTGGA	TAAAGGCATC	GTCGGGCTCG	GTGTTGAGCG	3200
TCTCCTGCGA	GATGAAGCCC	ATGACAAAGT	CCTTGTGCTC	CCGGGCGGCC	3250
TCGACGCAGG	CCTGCGTGTA	CTCCTTGTTC	ATGAAGTTGC	CCTGGCTGGA	3300
CATTTGGGCG	AGGATCAGGA	GGCCTCGGCT	CAGCGGCGCC	TCCTCGATGC	3350
CCGGGAAGAG	CGACTCGTCG	CCCTCGGCGA	TGGCCTTTGT	TAACCGGGGC	3400
GAGGAGACGG	ACTCGTACTG	CTGGGTGACG	GTGGTGATGG	AGACGATGCT	3450
GCCCTTGCGG	CCGTGCGCCG	ACCGGTTCGA	GTAGATGGGC	TTGTCCAGGA	3500
CGCCAATGGA	GCCCATGCCG	TTGACGGCGC	CGGCGGGCTC	GGCGTCCCTG	3550
GAGTCGGCGT	CGTCGTCAA	CGAGTCCATG	GTGGGCGTGC	CGACGGTGAC	3600
GGACGTCTTG	ACCTCGCAGG	GGTAGCGCTC	GAGCCAGCGC	TTGGCGCCCT	3650
GGGCCAGCGA	GGCCACCGAC	GCCTTGCCGG	GCACCATGTT	GACGTTGACA	3700
ATGTGCGCCC	AGTCGATGAT	GCGCGCCGAC	CCGCCCGTGT	ACTGCAGCTC	3750
GACGGTGTGG	CCAATGTGCG	CAAAC TTGCG	GTCCTCGAAG	ATGAGGAAGC	3800
CGTGCTTGCG	CGCCAGCGAC	GCCAGCTGGG	CTCCCGTGCC	CGTCTCCGGG	3850
TGGAAGTCCC	AGCCCCGAGAC	CATGTCGTAG	TGCGTCTTGA	GCACGACAAT	3900
CGACGGGCCA	ATCTTGTGCG	CCAGTACAG	CAGCTCGCGC	GCTGTCGGCA	3950
CGTCGGCGCT	CAGGCACAGG	TTGGACGCCT	TGAGGTCCAT	GAGCTTGAAC	4000
AGGTAAGCCG	TCAGCGGGTG	CGTCGCCGTC	TCGCTCCTGG	CCGCGAAGGT	4050
GGCCTTGAGC	GTCGGGTGTG	GTGCCATGGC	TGATGAGGCT	GAGAGAGGCT	4100
GAGGCTGCGG	CTGGTTGGAT	AGTTTAACCC	TTAGGGTGCC	GTTGTGGCGG	4150
TTAGAGGGG	GGGAAAAAA	AGAGAGAGAT	GGCACAATTC	TGCTGTGCGA	4200
ATGACGTTGG	AAGCGCGACA	GCCGTGCGGG	AGGAAGAGGA	GTAGGAACTG	4250
TCGGCGATTG	GGAGAATTTT	GTGCGATCCG	AGTCGTCTCG	AGGCGAGGGA	4300
GTTGCTTTAA	TGTCGGGCTC	GTCCCCTGGT	CAAATTTCTA	GGGAGCAGCG	4350
CTGGCAACGA	GAGCAGAGCA	GCAGTAGTCG	ATGCTAGAAA	TCGATAGATC	4400
CACGATGCCA	AAAAGCTTGT	TCATTTCCGG	TAGCCCGTGA	TCCTGGCGCT	4450
TCTAGGGCTG	AAACTGTGTT	GTTAATGTAT	TATTGGCTGT	GTAAGTACTG	4500
TGAATGGGGA	ATGAGGAGCG	CGATGGATTG	GCTTGCATGT	CCCCTGGCCA	4550
AGACGAGCCG	CTTTGGCGGT	TTGTGATTCG	AAGGTGTGTC	AGCGGAGGCG	4600
CCAGGGCAAC	ACGCACTGAG	CCAGCCAACA	TGCATTGCTG	CCGACATGAA	4650
TAGACACCG	CCGAGCAGAC	ATAGGAGACG	TGTTGACTGT	AAAAATTCTA	4700
CTGAATATTA	GCACGATG	TCTCAATAAG	AGCAATAGGA	ATGCTTGCCA	4750
ATCATAAGTA	CGTATGTGCT	TTTTCTGCA	AATGGTACGT	ACGGACAGTT	4800
CATGTTGTCT	GTCATCCCCC	ACTCAGGCTC	TCATGATCAT	TTTATGGGAC	4850
TGGGGTTTTG	CTGACTGAAT	GGATTGAGCC	GCACGAAACA	AATTGGGGGG	4900


FIGURE 6C: pTEX2

CATGCAGAAG	GGAAGCCCCC	CCAGCCCCCT	G TTCATAATT	TGTTAAGAGT	4950
CGGAGAGCTG	CCTAGTATGA	AGCAGCAATT	GATAACGTTG	ACTTTGCGCA	5000
TGAGCTCTGA	AGCCGGGCAT	ATGTATCACG	TTTCTGCCTA	GAGCCGCACG	5050
GGACCCAAGA	AGCTCTTGTC	ATAAGGTATT	TATGAGTGTT	CAGCTGCCAA	5100
CGCTGGTTCT	ACTTTGGCTC	AACCGCATCC	CATAAGCTGA	ACTTTGGGAG	5150
CTGCCAGAAT	GTCTCTTGAT	GTACAGCGAT	CAACAACCGT	GCGCCGGTCG	5200
ACAAC TGTTT	ACCGATCAGG	GACGCGAAGA	GGACCCAATC	CCGGTTAACG	5250
CACCTGCTCC	GAAGAAGCAA	AAGGGCTATG	AGGTGGTGCA	GCAAGGAATC	5300
AAAGAGCTCT	ATCCACTTGA	CAAGGCCAAT	GTCGCTCCCG	ATCTGGAGTA	5350
AGTCAACCCT	GAAGTGGAAG	TTTGCTTCTC	TGATTAGTAT	GTAGCATCGT	5400
GTTTGTCCCA	GGACTGGGTG	CAAATCCCGA	AGACAGCTGG	AAGTCCAGCA	5450
AGACCGACTT	CAATTGGACC	ACGCATACAG	ATGGCCTCCA	GAGAGACTTC	5500
CCAAGAGCTC	GGTTGCTTCT	GTATATGTAC	GACTCAGCAT	GGACTGGCCA	5550
GCTCAAAGTA	AAACAATFCA	TGGGCAATAT	CGCGATGGGG	CTCTTGGTTG	5600
GGCTGAGGAG	CAAGAGAGAG	GTAGGCCAAA	CGCCAGACTC	GAACCGCCAG	5650
CCAAGTCTCA	AACTGACTGC	AGGCGGCCGC	CATATGCATC	CTAGGCCTAT	5700
TAATATTCCG	GAGTATACGT	AGCCGGCTAA	CGTTAACAAAC	CGGTACCTCT	5750
AGAACTATAG	CTAGCATGCG	CAAATTTAAA	GCGCTGATAT	CGATCGCGCG	5800
CAGATCCATA	TATAGGGCCC	GGGTTATAAT	TACCTCAGGT	CGACGTCCCA	5850
TGGCCATTCC	AATTCGTAAT	CATGGTCATA	GCTGTTTCCT	GTGTGAAATT	5900
GTTATCCGCT	CACAATTCCA	CACAACATAC	GAGCCGGAAG	CATAAAGTGT	5950
AAAGCCTGGG	GTGCCTAATG	AGTGAGCTAA	CTCACATTAA	TTGCGTTGCG	6000
CTCACTGCC	GCTTTCCAGT	CGGGAAACCT	GTCGTGCCAG	CTGCATTAAT	6050
GAATCGGCCA	ACGCGCGGGG	AGAGGCGGTT	TGCGTATTGG	GCGCTCTTCC	6100
GCTTCCTCGC	TCACTGACTC	GCTGCGCTCG	GTCGTTCCGC	TGCGGCGAGC	6150
GGTATCAGCT	CACTCAAAGG	CGGTAATACG	GTTATCCACA	GAATCAGGGG	6200
ATAACGCAGG	AAAGAACATG	TGAGCAAAAAG	GCCAGCAAAA	GGCCAGGAAC	6250
CGTAAAAGG	CCGCGTTGCT	GGCGTTTTTC	CATAGGCTCC	GCCCCCTGA	6300
CGAGCATCAC	AAAAATCGAC	GCTCAAGTCA	GAGGTGGCGA	AACCCGACAG	6350
GACTATAAAG	ATACCAGGCG	TTTCCCCCTG	GAAGCTCCCT	CGTGCGCTCT	6400
CCTGTTCCGA	CCCTGCCGCT	TACCGGATAC	CTGTCCGCCT	TTCTCCCTTC	6450
GGGAAGCGTG	GCGCTTCTC	ATAGCTCACG	CTGTAGGTAT	CTCAGTTCGG	6500
TGTAGGTCGT	TCGCTCCAAG	CTGGGCTGTG	TGCACGAACC	CCCCGTTCAG	6550
CCCGACCGCT	GCGCCTTATC	CGGTAACTAT	CGTCTTGAGT	CCAACCCGGT	6600
AAGACACGAC	TTATCGCCAC	TGGCAGCAGC	CACTGGTAAC	AGGATTAGCA	6650
GAGCGAGGTA	TGTAGGCGGT	GCTACAGAGT	TCTTGAAGTG	GTGGCCTAAC	6700
TACGGCTACA	CTAGAAGAAC	AGTATTTGGT	ATCTGCGCTC	TGCTGAAGCC	6750
AGTTACCTTC	GGAAAAGAG	TTGGTAGCTC	TTGATCCGGC	AAACAAACCA	6800
CCGCTGGTAG	CGGTGGTTTT	TTTGTTTGCA	AGCAGCAGAT	TACGCGCAGA	6850
AAAAAAGGAT	CTCAAGAAGA	TCCTTTGATC	TTTTCTACGG	GGTCTGACGC	6900
TCAGTGGAAC	GAAAACTCAC	GTTAAGGGAT	TTTGGTCATG	AGATTATCAA	6950
AAAGGATCTT	CACCTAGATC	CTTTTAAATT	AAAAATGAAG	TTTTAAATCA	7000
ATCTAAAAGTA	TATATGAGTA	AACTTGGTCT	GACAGTTACC	AATGCTTAAT	7050
CAGTGAGGCA	CCTATCTCAG	CGATCTGTCT	ATTTCTGTTCA	TCCATAGTTG	7100
CCTGACTCCC	CGTCGTGTAG	ATAACTACGA	TACGGGAGGG	CTTACCATCT	7150
GGCCCCAGTG	CTGCAATGAT	ACCGCGAGAC	CCACGCTCAC	CGGCTCCAGA	7200
TTTATCAGCA	ATAAACCAGC	CAGCCGGAAG	GGCCGAGCGC	AGAAGTGGTC	7250
CTGCAACTTT	ATCCGCCTCC	ATCCAGTCTA	TTAATTGTTG	CCGGGAAGCT	7300
AGAGTAAGTA	GTTCCGCCAGT	TAATAGTTTG	CGCAACGTTG	TTGCCATTGC	7350

FIGURE 6D: pTEX2

TACAGGCATC	GTGGTGTAC	GCTCGTCGTT	TGGTATGGCT	TCATTCAGCT	7400
CCGGTTCCCA	ACGATCAAGG	CGAGTTACAT	GATCCCCCAT	GTTGTGCAA	7450
AAAGCGGTTA	GCTCCTTCGG	TCCTCCGATC	GTTGTCAGAA	GTAAGTTGGC	7500
CGCAGTGTTA	TCACTCATGG	TTATGGCAGC	ACTGCATAAT	TCTCTTACTG	7550
TCATGCCATC	CGTAAGATGC	TTTTCTGTGA	CTGGTGAGTA	CTCAACCAAG	7600
TCATTCTGAG	AATAGTGTAT	GCGGCGACCG	AGTTGCTCTT	GCCC GGCGTC	7650
AATACGGGAT	AATACCGCGC	CACATAGCAG	AACTTTAAAA	GTGCTCATCA	7700
TTGAAAACG	TTCTTCGGGG	CGAAA ACTCT	CAAGGATCTT	ACCGCTGTTG	7750
AGATCCAGTT	CGATGTAACC	CACTCGTGCA	CCCAACTGAT	CTTCAGCATC	7800
TTTTACTTTC	ACCAGCGTTT	CTGGGTGAGC	AAAAACAGGA	AGGCAAAATG	7850
CCGCAAAAAA	GGGAATAAGG	GCGACACGGA	AATGTTGAAT	ACTCATACTC	7900
TTCTTTTTTC	AATATTATTG	AAGCATTTAT	CAGGGTTATT	GTCTCATGAG	7950
CGGATACATA	TTTGAATGTA	TTTAGAAAAA	TAAACAAATA	GGGGTTCCGC	8000
GCACATTTCC	CCGAAAAGTG	CCACCTGACG	TCTAAGAAAC	CATTATTATC	8050
ATGACATTAA	CCTATAAAAA	TAGGCGTATC	ACGAGGCCCT	TTCGTCTCGC	8100
GCGTTTCGGT	GATGACGGTG	AAAACCTCTG	ACACATGCAG	CTCCC GGAGA	8150
CGGTCACAGC	TTGTCTGTAA	GCGGATGCCG	GGAGCAGACA	AGCCC GTCAG	8200
GGCGCGTCAG	CGGGTGTGG	CGGGTGTCCG	GGCTGGCTTA	ACTATGCGGC	8250
ATCAGAGCAG	ATTGTA CTGA	GAGTGCACCA	TAAAATTGTA	AACGTTAATA	8300
TTTTGT TAAA	ATTCGCGTTA	AATTTTGTGTT	AAATCAGCTC	ATTTTTTAAC	8350
CAATAGGCCG	AAATCGGCAA	AATCCCTTAT	AAATCAAAAG	AATAGCCCGA	8400
GATAGGGTTG	AGTGTTGTTC	CAGTTTGGAA	CAAGAGTCCA	CTATTAAAGA	8450
ACGTGGACTC	CAACGTCAAA	GGGCGAAAAA	CCGTCTATCA	GGGCGATGGC	8500
CCACTACGTG	AACCATCACC	CAAATCAAGT	TTTTTGGGGT	CGAGGTGCCG	8550
TAAAGCACTA	AATCGGAACC	CTAAAGGGAG	CCCCCGATTT	AGAGCTTGAC	8600
GGGAAAGCC	GGCGAACGTG	GCGAGAAAGG	AAGGGAAGAA	AGCGAAAGGA	8650
GCGGGCGCTA	GGGCGCTGGC	AAGTGTAGCG	GTCACGCTGC	GCGTAACCAC	8700
CACACCCGCC	GCGCTTAATG	CGCCGCTACA	GGGCGCGTAC	TATGGTTGCT	8750
TTGACGTATG	CGGTGTGAAA	TACCGCACAG	ATGCGTAAGG	AGAAAATACC	8800
GCATCAGGCG	CCATTGCGCA	TTCAGGCTGC	GCAACTGTTG	GGAAGGGCGA	8850
TCGGTGC GGG	CCTCTTCGCT	ATTACGCCAG	CTGGCGAAAG	GGGGATGTGC	8900
TGCAAGGCCA	TTAAGTTGGG	TAACGCCAGG	GTTTTCCAG	TCACGACGTT	8950
GTAAAACGAC	GGCCAGTGCC				8970

FIG. 7A




```

20VW.A          -----TPDK-AKEQHPKLETY-----TKAS--G--KKQ
1A39            -----KPGE-TKEVHPQLTTF-----TKRG--G--KPA
6CEL           -----SACTLQSETHPPLTWQK-----SGG--T--QQ
1EG1.A         -----QFGTSTPEVHPKLTYYK-----TKSG--G--VAQ
P_chrysosporium M----RTALA-L-ALA AFS---AVSAQQAGTITAETHPPLTIQQ TQSG--G--APL
P_chrysospori_5 M----RTALA-L-ALA AFS---AVSAQQAGTITAETHPPLTIQQ TQSG--G--APL
P_chrysospori_6 M----VDIQI-ATFLLLGIV---GVAAQQVGTYPENHPLLATQS TQSG--G--APL
P_chrysospori_7 M----RTA--T-AFTAA---MVF--QQVGTNTAENHRTLSQK TKSG--G--NL
P_chrysospori_8 M----RTA--T-AFTAA---MVF--QQVGTNTAENHRTLSQK TKSG--G--NL
P_chrysospori_9 M----RTA--T-AFTAA---MVF--QQVGTNTARSHPALTSQK TKSG--G--NL
P_chrysospor_10 M----RAA--A-AFTCA---MVS--QQAGTNTAENHPQLQSQQ TTSG--G--KPL
P_chrysospor_11 M----RAA--A-AFTCA---MVS--QQAGTNTAENHPQLQSQQ TTSG--G--KPL
P_chrysospor_12 M----RAA--A-AFTCA---MVS--QQAGTNTAENHPQLQSQQ TTSG--G--KPL
I_lacteus_45863 M----RKA--A-AFSFA---IAH--QQVGTNQAENHPSLPSQK TQSG--G--NL
I_lacteus_45_14 M----HKA--V-AFSFA---IVH--QQAGTQTAENHPQLSQQ TAGG--S--SA
I_lacteus_45_15 M----PKA--S-AALSFA---AVY--QQVGTQMAEVHPKLPSSL TKSG--G--NL
A_alternata_617 -----TWQS-----TAKG--S--NK
L_maculans_7804 M----RS--L-FATSAS---LAK--QLVG-----NL--K--G--S--AK
C_parasitica_39 M----SKF--A-TGSLAS---AVNAQQVGTQQTETHPQLTWQS TSPS--S--TN
C_carbonum_3913 M----RTL--AFASLSYS---AARAQQVGTSTAENHPKLTWQT TGTG--GTN--NK
H_grisea_134622 M----RTAK--FATLAAFA---SAAAQQAC--LTTERRHPSL--WKK TAGG--Q--QTV
H_grisea_950686 M----RTAK--FATLAAFA---SAAAQQAC--LTTERRHPSL--WKK TAGG--Q--QTV
F_oxysporum_117 M----RIV--ATAS-AFA---AARAQQVC--LNTETKPALTWSK TSSG--G--DV
C_purpurea_1906 M----HPSLQ-T-LSAFT---TAHAQQAC--SKPETHPPL--WS--RSG-----RSV
H_thermoidea_40 MQIK-S-IQY--LAAALP--S---SVAAQQAGTITAENHPR--TWK--GPG--N--QTV
H_thermoidea_74 MQIK-S-IQY--LAAALP--S---SVAAQQAGTITAENHPR--TWK--GPG--N--QTV
L_maculans_7_26 M----LSASK-AAAILAFCA--HTASAWVVDQQTETHPKLNWQ--TGKGRSS--NV
N_crassa_729649 M----RASL--LAFSLA-AA---VAG--QQAGTLTAKRHPSL--TWQK--TRGG-----PTL
A_aculeatus_391 M--VD-S-SIYK-TA--LLS--FA---TSNAQQVGTYYTAETHPSL--TWQT--GSG--S--TT
A_niger_6164684 M--S-S-QIYR-AA--LLS--FA---TANAQQVGTYYTETHPSL--TWQT--TSDG--S--TN
P_janthinellum MKGSIS-QIYKGA--LSA--N---SVSAQQVGTTLTAETHPAL--TWSK--TA-G--X--QV
A_niger_6164682 M----HORA--L-FS-AFA---AVRAQQAGTLTEEVHPSL--TWQK--TSEG--S--EQ
H_ceramica_1218 M----RKL--A-S-AFA---TARAQSACTLQSETHPPL--TWQK--SGG--T--QQ
H_jecorina_2238 -----ESACTLQSETHPPL--TWQK--SGG--T--QQ
T_viride_121854 M----QKL--A-S-AFA---TARAQSACTLQAEHPPL--TWQK--SGG--T--QQ
T_viride_406299 M----QKL--A-S-AFA---TARAQSACTLQAEHPPL--TWQK--SGG--T--QQ
T_harzianum_710 M----RKL--A-S-AFA---AARAQQVCTQQAETHPPL--TWQK--TASG-----PQ
A_bisporus_3913 M----PRS--I-SALS--TA---VAL--QQVGTNMAENHPSL--TWQK--TSSG-----QNV
V_volvacea_5231 M----RA--S-AFS--N---SAA--QQAGTLQTKNHPSL--TSQK--ROGG-----PQV
T_longibrachiat M----APSAT-LP-TTA--AIARLVAAQQPGTSTPEVHPKLTYYK--TTSG--G--VAQ
H_jecorina_1217 M----APSAT-LP-TTA--AIARLVAAQQPGTSTPEVHPKLTYYK--TKSG--G--VAQ
A_oryzae_246737 M----IWTL--APFV-A--P---LVTAQQVGT--TA--AHPRLTYYK--TSQN--G--RQ
consensus      m----f-----il--mva-----aqq-gt-tae-hp-ltwqkCt--g-----Ct--

```

Figure 8A

```

20VW.A      TNYVVA AGIHG RQK ---NGAGCGDWGQKPNATA PDEAS AKN ILS -MDSNAYK-N
1A39        TNFVVL SLWHW HRAEGLGPGGCGDWGNPEPKD PDVE AKN I -IPD--Y-Q
6CEL        TSVVVD ANWRWTHATNSSTNCYDGN---WS TL PDNET AKN CLD -A---AYAST
1EG1.A     DTAVVL WN RW HDAN NSCTVNGG---VN TL PDEAT AKN F -V---DYA-A
P_chrysosporium TTKVVL VNWRW HSTTG TNCY GNT---WDAIL PDPVT AAN ALD -A---DY T
P_chrysospori_5 TTKVVL VNWRW HSTTG TNCY GNT---WDAIL PDPVT AAN ALD -A---DY T
P_chrysospori_6 SSKVVL ANRRW HSTLGTTSCLT NG---WDPTL PDGIT ANY ALD -V---SY ST
P_chrysospori_7 NTKVVL ANWRW HSTSG TNCYTGNO---WDATL PDGKT AAN ALD -A---DY T
P_chrysospori_8 NTKVVL ANWRW HSTSG TNCYTGNO---WDATL PDGKT AAN ALD -A---DY T
P_chrysospori_9 NTKVVL ANWRW HSTSG TNCYTGNO---WDATL PDGKT AAN ALD -A---DY T
P_chrysospor_10 STKVVL SNWRW HSTSG TNCYTGNE---WN TL PDGKT AAN ALD -A---DY T
P_chrysospor_11 STKVVL SNWRW HSTSG TNCYTGNE---WD TL PDGKT AAN ALD -A---DY T
P_chrysospor_12 STKVVL SNWRW HSTSG TNCYTGNE---WD TL PDGKT AAN ALD -A---DY T
I_lacteus_45863 STAVVL ANWRW HSTTG TNCYTGNT---WDA PDGVT AKA ALD -A---DY T
I_lacteus_45_14 STAVVL SNWRW HSTTG TNCYTGNT---WDA SDPV AQN ALD -A---DYA T
I_lacteus_45_15 NTAVVL ANWRW HSTTG TNCYTGNS---WDATL PDATT AQN ALD -A---DY T
A_alternata_617 NTKVVL ANWRW HKKGG DNCYTGNE---WDATA AAN ALD -A---DY T
L_maculans_7804 NTKVVL ANWRW HVKGG TNCYTGNE---WDATA PDNK ATN ALD -A---DYR-R
C_parasitica_39 QEVV SNWRW HDKDG VNCYTGNT---WN TL PDDKT AAN VLD -A---DY ST
C_carbonum_3913 SIVVL SNNRWAHNVGG TNCYTGNS---WS QY PDGD TKN ALD -A---DY T
H_grisea_134622 QAVTL SNNRWTHQVSGSTNCYTGNK---WD TDAK AQN CLD -A---DY ST
H_grisea_950686 QAVTL SNNRWTHQVSGSTNCYTGNK---WD TDAK AQN CLD -A---DY ST
F_oxysporum_117 KAVV ANWRWTHQTSGSTNCYTGNK---WD TDGKT AEK CLD -A---DY T
C_purpurea_1906 QAVT ANWLWT TVDGSQNCYTGNR---WD SSEKT SES CLD -A---DYA T
H_thermoidea_40 QEVV ANNRW HN-NG-QNCYEGNK---WT Q- SSATD AQR ALD -A---NYQST
H_thermoidea_74 QEVV ANNRW HN-NG-QNCYEGNK---WT Q- SSATD AQR ALD -A---NYQST
L_maculans_7_26 NEVV ANNRW AHRSG TNCYTGSE---WNO A PNNEA TKN ALD -S---DYA T
N_crassa_729649 NTAVVL ANNRWTHATSGSTKCYTGNK---WQATL PDGK AAN ALD -A---DY T
A_aculeatus_391 SIVV ANNRW HEVGG TNCY GNT---WD STDTT ASE AL -A---TYEST
A_niger_6164684 DEVV ANNRW HSTSSATNCYTGNE---WD TDDVT AAN ALD -A---TYE T
P_janthinellum SIVV ANWPX HSTVSGSTNCYTGNT---WDATL PDDVT AAN ALD -A---RRQ-H
A_niger_6164682 SIVV SNNRWTHSVNDSTNCYTGNT---WDATL PDDVT AAN ALD -A---DYEST
H_ceramica_1218 TAVV ANNRWTHATNSSTNCYDGN---WS TL PDNET AKN CLD -A---AYAST
H_jecorina_2238 TAVV ANNRWTHATNSSTNCYDGN---WS TL PDNET AKN CLD -A---AY SA
T_viride_121854 TAVV ANNRWTHATNSSTNCYDGN---WS TL PDNET AKN CLD -A---AYAST
T_viride_406299 TAVV ANNRWTHATNSSTNCYDGN---WS TL PDNET AKN CLD -A---AYAST
T_harzianum_710 QAVVL ANNRWTHDTKSTTNCYDGN---WS TL PDDAT AKN CLD -A---NY T
A_bisporus_3913 NTKVTL ANNRWTHRIND TNCYTGNE---WD PDGVT AEN ALD -A---DYA T
V_volvacea_5231 NTAVVL ANNRWTHSTSGSTNCYTGNT---WQATL PDGKT AAN ALD -A---DY T
T_longibrachiat DTAVVL WN RW HDAN NSCTVNGG---VN TL PDEAT AKN Y -V---DYA
H_jecorina_1217 DTAVVL WN RW HDAN NSCTVNGG---VN TL PDEAT AKN F -V---DYA
A_oryzae_246737 NTAVVL AATH HKKGTQTSCTNSNG---LD A PDKQT ADN V D IT---DYA-
consensus -gsvvlDanwrwih-t-gytncytn---wdstlCpd--tCa-nCaldG-a---dysgt

```

Figure 8B

```

20VW.A      AGITTSGNKLRRL-QQI-N---N-QLVSP  -Y  EENK-KK  EMLHLTG-T  FF  D
1A39       YGTTNGTSLRRL-QHLPD---G-RVPSP  -Y  DKTK-RR  EMLHLTG-F  FT  D
6CEL       YGTTSGNSLRL-DFVTQSA---Q-KNVGA  -Y  MA-SD-T  QEFTLLG-  F  D
1EG1.A     SGTTSGSSLRL-NQYMPS-SGGY-SSVSP  -Y  DSD-GE  VMLKLNGL-  L  D
P_chrysosporium  GILPSGTSRL-RPVD-G-----GL  -Y  MA-DD-H  QMFQLLN-K  FT  D
P_chrysospori_5  GILPSGTSRL-RPVD-G-----GL  -Y  MA-DD-H  QMFQLLN-K  FT  D
P_chrysospori_6  YGTTSGSALRRL-QFVT-G-----N-  -Y  MA-DD-H  RTFQLLN-  LA  D
P_chrysospori_7  YGITASGSSLKL-QFVT-G-----NVGS  -Y  MA-DD-H  QMFQLLN-  FT  D
P_chrysospori_8  YGITASGSSLKL-QFVT-G-----NVGS  -Y  MA-DD-H  QMFQLLN-  FT  D
P_chrysospori_9  YGITASGSSLKL-QFVT-G-----NVGS  -Y  MA-DD-H  QMFQLLN-  FT  D
P_chrysospor_10  YGITSGTALRL-KFVT-G-----NVGS  -Y  MA-DD-H  QLKLLN-  FT  D
P_chrysospor_11  YGITSGTALRL-KFVT-G-----NVGS  -Y  MA-DD-H  QLKLLN-  FT  D
P_chrysospor_12  YGITSGTALRL-KFVT-G-----NVGS  -Y  MA-DD-H  QLKLLN-  FT  D
I_lacteus_45863  YGITTSGNALRL-QFVK-G-----NVGS  -Y  MQ-DA-N  QMFQLLN-  FT  D
I_lacteus_45_14  YGITTSGDALRL-KFVT-G-----NVGS  -Y  ME-DE-N  QMFQLLN-  FT  D
I_lacteus_45_15  YGITTSGNALRL-KFVK-G-----NVGS  -Y  MQ-TD-A  QMFQLLN-  FT  D
A_alternata_617  YGITAGSNSLKL-KFVTKG-Y--S-TN-  T-Y  MK-DD-T  QMFKFTGN-  FT  D
L_maculans_7804  LRHYCERQLLGT-EVHHQGLY--S-TN-  T-Y  MQ-DD-T  QFKFTGS-  FT  D
C_parasitica_39  YGITTSGNALRL-QFVTQS---G-KN-  T-Y  ME-SS-K  HFDLIG-  FA  D
C_carbonum_3913  YGITTSNNALRL-KFVTKG-F--S-SN-  T-Y  ME-TD-K  QMFNLN-K  FT  D
H_grisea_134622  YGITTINGDSLRL-KFVTKGQH--S-TN-  T-Y  MD-G-DK  QTFELLG-  FT  D
H_grisea_950686  YGITTINGDSLRL-KFVTKGQY--S-TN-  T-Y  MD-G-DK  QTFELLG-  FT  D
F_oxysporum_117  YGITTSGNQLRL-GFVTNGPY--S-KN-  T-Y  ME-N-NT  QMFQLLG-  FT  D
C_purpurea_1906  YGITTSGDALRL-KFVQGGPY--S-KN-  -Y  MK-D-R  EMFTLLG-  FT  D
H_thermoidea_40  YGATSGDSLRL-KFVTKHEY--G-TN-  F-Y  MA-NQ-NK  QMFTLN-  FA  D
H_thermoidea_74  YGATSGDSLRL-KFVTKHEY--G-TN-  F-Y  MA-NQ-NK  QMFTLN-  FA  D
L_maculans_7_26  YGITTSGNQRLN-KFVTKRPY--S-TN-  T-Y  MK-D-QN  EMFQLG-  FT  D
N_crassa_729649  YGITSGWSLRL-QFVT-D-----NVGA  A-Y  MA-DD-Q  QMLELLN-  LW  D
A_aculeatus_391  YGTTSGSSLRL-NFVTT---Q-KN-  -Y  MA-DD-T  ETFKLFN-R  FT  D
A_niger_6164684  YGTTSGSELRL-NFVTQG---S-KN-  -Y  MS-DD-N  EFKLLG-  FT  D
P_janthinellum  LRTTSGNSLRL-NFVTT---Q-KN-  -Y  E-ND-T  QKFNLN-  FT  D
A_niger_6164682  YGTTDGDSLRL-KFVT-G-----NVGS  -Y  MDTSD-EG  QTFNLLD-A  FT  D
H_ceramica_1218  YGTTSGNSLRL-GFVTQSA---Q-KN-  -Y  MA-SD-T  QEFTLLG-  F  D
H_jecorina_2238  YSSZPGGGGGV--IFFK-----NVGA  -Y  MA-SD-T  QEFTLLG-  F  D
T_viride_121854  YGTTSDSLRL-GFVTQSA---Q-KN-  -Y  MA-SD-T  QEFTLLG-  F  D
T_viride_406299  YGTTSDSLRL-GFVTQSA---Q-KN-  -Y  MA-SD-T  QEFTLLG-  F  D
T_harzianum_710  YGTTSGDALRL-QFVT-G-----NVGS  -Y  MA-ND-T  QEFTLSG-  F  D
A_bisporus_3913  YGTTSGTALRL-KFVTESQ---Q-KN-  -Y  MA-DD-N  EFNLLN-K  FT  D
V_volvacea_5231  YGTTSGNSLRL-QFVT-Q-----NVGA  -Y  MA-DD-T  QMFNLLN-  FW  D
T_longibrachiat  -GTTASGSLRLNQPSSG--GYSSVSP  -Y  -PD-GE  VMLKLNGL-  L  D
H_jecorina_1217  -GTTSGSSLRLNQPSSG--GYSSVSP  -Y  D-SD-GE  VMLKLNGL-  L  D
A_oryzae_246737  YGTTQTKNDRLRLQQTGNA--T-KS-  -Y  A-DGEN  SMLKLN-  FT  D
consensus  ygittsg-slsl--fvt-gs-----nvgsRv-yLma-dd-t-Yqmf-lln-neftFDVd

```

Figure 8C

20VW.A	MEKLP	MNGALYLSE	PQDGG-K	-----	STSR	---	NSKAG	Y	-----	YGA	Y
1A39	AKLP	MNSALYLSE	HPTGA-K	-----	S--K	---	YNPAG	Y	-----	YGT	Y
6CEL	SALP	LNGALYFVS	DADGG-	-----	SKYP	---	TNTAG	K	-----	YGT	Y
1EG1.A	SALP	ENGSLYLSQ	DENGG-A	-----	NO	---	YNTAG	N	-----	YG	Y
P_chrysosporium	PNMR	SSGALHLTA	DADGG-	-----	AKYP	---	GNQAG	K	-----	YGT	Y
P_chrysospori_5	PNMR	SSGALHLTA	DADGG-	-----	AKYP	---	GNQAG	K	-----	YGT	Y
P_chrysospori_6	SKLP	LNGALYFVA	DADGG-K	-----	SKYP	---	GNAG	K	-----	YGT	Y
P_chrysospori_7	SNLP	LNGALYLSA	DADGG-	-----	AKYP	---	TNKAG	K	-----	YGT	Y
P_chrysospori_8	SNLP	LNGALYLSA	DADGG-	-----	AKYP	---	TNKAG	K	-----	YGT	Y
P_chrysospori_9	SNLP	LNGALYLSA	DADGG-	-----	AKYP	---	TNKAG	K	-----	YGT	Y
P_chrysospor_10	SNLP	LNGALYLSA	DADGG-	-----	SKYP	---	GNKAG	K	-----	YGT	Y
P_chrysospor_11	SNLP	LNGALYLSA	DADGG-	-----	SKYP	---	GNKAG	K	-----	YGT	Y
P_chrysospor_12	SNLP	LNGALYLSA	DADGG-	-----	SKYP	---	GNKAG	K	-----	YGT	Y
I_lacteus_45863	SNLP	LNGALYLSQ	DQDGG-	-----	SAP	---	TNTAG	K	-----	YGT	Y
I_lacteus_45_14	SNLP	LNGALYFVQ	DQDGG-T	-----	SKP	---	NNKAG	K	-----	YGT	Y
I_lacteus_45_15	SNLP	LNGALYLSQ	DQDGG-	-----	SKP	---	TNKAG	K	-----	YGT	Y
A_alternata_617	SNLP	FNGALYFVS	DADGG-	-----	KKYS	---	TNKAG	K	-----	YGT	Y
L_maculans_7804	SNLP	LNGALYFVS	DADGG-	-----	KKYP	---	TNKAG	K	-----	YGT	Y
C_parasitica_39	SKLP	LNGALYFVT	DADGG-	-----	AKYS	---	TNTAG	E	-----	YGT	Y
C_carbonum_3913	SKLP	LNGALYFVE	AADGG-	-----	GK-G	---	NNKAG	K	-----	YGT	Y
H_grisea_134622	SNIG	LNGALYFVS	DADGG-	-----	SYP	---	GNKAG	K	-----	YGT	Y
H_grisea_950686	SNIG	LNGALYFVS	DADGG-	-----	SYP	---	GNKAG	K	-----	YGT	Y
F_oxysporum_117	SGLG	LNGAPHFVS	DEDGG-K	-----	AKYS	---	GNKAG	K	-----	YGT	Y
C_purpurea_1906	SKLG	LNGALYFVS	DEDGG-	-----	KP	---	MNKAG	K	-----	YGT	Y
H_thermoidea_40	SKER	MNSALYFVA	EDGG-	-----	ASYP	---	SNAG	K	-----	YGT	Y
H_thermoidea_74	SKER	MNSALYFVA	EDGG-	-----	ASYP	---	SNAG	K	-----	YGT	Y
L_maculans_7_26	S-R	MNGALYFVS	PQKGQ-	-----		---	GAPG	K	-----	YGT	Y
N_crassa_729649	SNP	LNGALYLSA	DADGG-	-----	RKYP	---	TNKAG	K	-----	Y	Y
A_aculeatus_391	SNLP	LNGALYFVS	DADGG-	-----	SP	---	TNKAG	K	-----	YGT	Y
A_niger_6164684	SNLP	LNGALYFVA	DADGG-T	-----	SEYS	---	GNKAG	K	-----	YGT	Y
P_janthinellum	SNLP	LNGALYFVD	DADGG-	-----	AKYP	---	TNKAG	K	-----	YGT	Y
A_niger_6164682	SNLP	LNGALYFTA	DADGG-	-----	SKYP	---	ANKAG	K	-----	YGT	Y
H_ceramica_1218	SALP	LNGALYFVS	DADGG-	-----	SKYP	---	TNTAG	K	-----	YGT	Y
H_jecorina_2238	SALP	LNGALYFVS	DADGG-	-----	SKYP	---	TNTAG	K	-----	YGT	Y
T_viride_121854	SALP	LNGALYFVS	DADGG-	-----	SKYP	---	TNTAG	K	-----	YGT	Y
T_viride_406299	SALP	LNGALYFVS	DADGG-	-----	SKYP	---	TNTAG	K	-----	YGT	Y
T_harzianum_710	SALP	LNGALYFVS	DADGG-Q	-----	SKYP	---	GNAAG	K	-----	YGT	Y
A_bisporus_3913	SKLP	LNGALYFSE	AADGG-	-----	SS	---	TNTAG	K	-----	YGT	Y
V_volvacea_5231	SNLP	LNGALYFSA	ARTW	PMVVC	ASTPLISTR	ST	LLR	LPVPPRSRYGR	I		
T_longibrachiat	SALP	ENGSLYLSQ	DENGG-A	-----	NOY	---	NTAG	N	-----	YG	Y
H_jecorina_1217	SALP	ENGSLYLSQ	DENGG-A	-----	NOY	---	NTAG	N	-----	YG	Y
A_oryzae_246737	ASTLV	MNGALYLSE	ASGG-K	-----	SSL	---	NOAG	K	-----	YGT	Y
consensus	msnlp	CGlngalyfv	Mdadgg	-v	---	skyp	---	nkag	AK	-----	YGTGyC

Figure 8D

```

20VW.A      DA YV-TFFIN V  TK-----GQ V  NE DIW A SRATHIAP P  K
1A39        DA FV-TFFIN LG  E-----GK  NEMDIW A SRASHVAP T  NK
6CEL        DS PRD KFIN QA VEGWEPS NNANTGIGGH  QMDIW A S SEALTP P  T
1EG1.A     DA PV-QT R N TL TS-----HQ F  NEMDIL  SRANALTP S  TA
P_chrysosporium SA P G KFIN QA VEGWLG A--TTGTGFF  A W A DNSAS AP P  T
P_chrysospori_5 SA P G KFIN QA VEGWLG A--TTGTGFF  A W A DNSAS AP P  T
P_chrysospori_6 DS PRD QFIN QA VQGWNA A--TTGTGSY  E DIW A SNAAALTP T  TN
P_chrysospori_7 DS PRD KFIN EA VEGWNA A--NAGTGNY  EMDIW A NDAAA TP P  T
P_chrysospori_8 DS PRD KFIN EA VEGWNA A--NAGTGNY  EMDIW A NDAAA TP P  T
P_chrysospori_9 DS PRD KFIN EA VEGWNA A--NAGTGNY  EMDIW A NDAAA TP P  T
P_chrysospor_10 DS P D KFIN EA VGNWTE GS--NTGTGSY  EMDIW A NDAAA TP P  T
P_chrysospor_11 DS P D KFIN EA VGNWTE GS--NTGTGSY  EMDIW A NDAAA TP P  T
P_chrysospor_12 DS P D KFIN EA VGNWTE GS--NTGTGSY  EMDIW A NDAAA TP P  T
I_lacteus_45863 DS PRD KFIN EA VEGWTGS TDSNSGTGNY  EMDIW A S AAA TP P  V
I_lacteus_45_14 DS PQD KFIN EA VDWTASAGDANS GTGSF  QEMDIW A S SAA TP P  TV
I_lacteus_45_15 DS PHD KFIN MA VAGWAGSASDPNAGSGTL  EMDIW A NDAAA TP P  V
A_alternata_617 DA PRD KFIN E  VEGWKPS NDANAGVGGH  AEMDIW A S STAVTP S
L_maculans_7804 DA PRD KFIN E  VEGWQPSKNDQNAVGGH  AEMDIW A S STAVTP S
C_parasitica_39 DS PRD KFIN Q  VEGWTPS NDANAGVGGH  EMD W A S DMA TP P  E
C_carbonum_3913 DS PHD KFIN KA VEGWNPSDADPNAGGAGKI A  PEMDIW A S STA TP P  RG
H_grisea_134622 DA PRD KFIN EA EGWTGS NDPNAGAGRY  EMDIW A NATA TP P  TI
H_grisea_950686 DA PRD KFIN EA EGWTGS NDPNAGAGRY  EMDIW A NATA TP P  TI
F_oxysporum_117 DA PRD KFIN VA SEGWKPS SDVNAGVGNL  PEMDIW A S STA TP P  TK
C_purpurea_1906 DS PRD KFIN MA SKDWIPSKSDANAGIGSL A  REMDIW A N ASA TP P  KN
H_thermoidea_40 DA ARD KFIG KA EGWRPS NDPNAGVGPMA  A E D W S AYAYA TP A  G
H_thermoidea_74 DA ARD KFIG KA EGWRPS NDPNAGVGPMA  A E D W S AYAYA TP A  G
L_maculans_7_26 DA ARD KFIN SA AEGWTKSASDPNSGVGKK A  AQMD W A SAATALTP S  OP
N_crassa_729649 DA PRD KFIN IA VEGWTPS ND-ANGIGDH  EMDIW A K STA TP P  T
A_aculeatus_391 DS PRD KFIN QA EGWEPS TDVNAGTGNH  PEMDIW A S SSA TA P  D
A_niger_6164684 DS PRD KFIN EA CGWEPS NNVNTGVGDH  AEMD W A S SNA TA P  D
P_janthinellum DS PRD KFIN QA V GWTPSKNDVNSGIGNH  AEMDIW A S SNAVTP P  D
A_niger_6164682 DS PRD KFIN QA V GWEPS NNDNTGIGNH  PEMDIW A K STALTP P  D
H_ceramica_1218 DS PRD KFIN QA VEGWEPS NNANTGIGGH  EMDIW A S SEALTP P  T
H_jecorina_2238 DS PRD KFIN QA VEGWEPS NNANTGIGGH  EMDIW A S SEALTP P  T
T_viride_121854 DS PRD KFIN QA VEGWEPS NNANTGIGGH  EMDIW A S SEALTP P  T
T_viride_406299 DS PRD KFIN QA VEGWEPS NNANTGIGGH  EMDIW A S SEALTP P  T
T_harzianum_710 DS PRD KFIN QA VEGWEPS NNANTGVGGH  EMDIW A S SEALTP P  E
A_bisporus_3913 DS PRD KFIN EA SEGWEPSNDVNAGTGNF A  GEMDIW A S SSA TP P  RE
V_volvacea_5231 DS PRD KFIN EA VQGWQPS PNDTNAGTGNY A  NKMD W A S STA TP P  TQ
T_longibrachiat DA PVQ-T R N TL TSG-----Q  F  NEMDIL  SRANALTP S  TA
H_jecorina_1217 DA PVQ-T R N TL TS-----HQ  F  NEMDIL  SRANALTP S  TA
A_oryzae_246737 DA YTT-B I N E  TE-----SV  QEMDIW A ARAT LTP P  N
consensus dsQCprdlkfinG-aNvegw--ss--n-g-g--GsCCsemdiwEaNsia-aftpHpCtt

```

Figure 8E

```

20VW.A      PQL-YGC-TGECG-----SSGI  K-A  CGF  HNR NVTD FYGR  KQYKVD
1A39        KQL-YLC-EGEC-----EGV   K-N  CGF  NYR NVTD FYGR  EEFKNTL
6CEL        VEQ-EIC-EGGCGGTYS--NRYGGT P-D  CDF  PYR GNT FYGP  SSFT DT
1EG1.A      -----TA  S-A  CGF  PYGSGYK FYGP  --DTVDT
P_chrysosporium NSQ-RC-SGSDC-----TADSGL  A-D  CNF  SRMGNT FYGA  --MVDT
P_chrysospori_5 NSQ-RC-SGSDC-----TADSGL  A-D  CNF  SRMGNT FYGA  --MVDT
P_chrysospori_6 NQ-RC-SGSNC-----TSNTGF  A-D  CDF  SR GNT FLGA  --MVDT
P_chrysospori_7 NQ-RC-SGSDC-----TRDTGL  A-D  CDF  SRMGDQ FLGK  --LTVDT
P_chrysospori_8 NQ-RC-SGSDC-----TRDTGL  A-D  CDF  SRMGDQ FLGK  --LTVDT
P_chrysospori_9 NQ-RC-SGSDC-----TRDTGL  A-D  CDF  SRMGDQ FLGK  --LTVDT
P_chrysospor_10 TQ-RC-SGSDC-----ARNTGL  HGD  CDF  SRMGDK FLGK  --MTVDT
P_chrysospor_11 TQ-RC-SGSDC-----ARNTGL  G-D  CDF  SRMGDK FLGK  --MTVDT
P_chrysospor_12 TQ-RC-SGSDC-----ARNTGL  G-D  CDF  SRMGDK FLGK  --MTVDT
I_lacteus_45863 NQQ-RC-TGADCGQ--GD--RYDGV  P-D  CDF  SRMGDQ FLGK  --LTVDT
I_lacteus_45_14 TEQ-RC-SGSDCGQ--GS--RNGI  P-D  CDF  SRMGNT FYGK  --LTVDT
I_lacteus_45_15 DEQ-QC-SGTQCGD--DD--RYSGL  K-D  CDF  SRMGDK FLGK  --MTVDT
A_alternata_617 IEQ-RC-DGCGGTYS--RYAGV  P-D  CDF  SYRMGVKDFYGK  --KTVDT
L_maculans_7804 IEQ-RC-DGCGGTYS--RYAGV  P-D  CDF  SYRMGVKDFYGK  --KTVDT
C_parasitica_39 A-Q-HSC-NCECGGTYS--SRYAGD  P-D  CDF  PYRMGNKDFYGS  --DTVDT
C_carbonum_3913 V-L-QECSDA SCGD--GS--NRYDGO  K-D  CDF  SYRMGVKDFYGP  --ATDT
H_grisea_134622 IQ-RC-EGSCGGTYSN--RYAGV  P-D  CDF  SYROGK FYGK  --MTVDT
H_grisea_950686 IQ-RC-EGSCGGTYSN--RYAGV  P-D  CDF  SYROGK FYGK  --MTVDT
F_oxysporum_117 LTQ-HSC-TGSCGGTYS--RYGGT  A-D  CDF  AYROGK FYGP  SNFN DT
C_purpurea_1906 SY-HSC-TGCGGTYSK--NRYSGD  P-D  CDF  SYR GNT FYGP  PKFT DT
H_thermoidea_40 KNRYHIC-ETNCCGGTYS--RYAGY  A-N  CDF  PYRMGNKDFYGK  --KTVDTN
H_thermoidea_74 KNRYHIC-ETNCCGGTYS--RYAGY  A-N  CDF  PYRMGNKDFYGK  --KTVDTN
L_maculans_7_26 A-Y-V-EDTNCGGTYSE--RYAGT  A-N  CDF  PYRMGVKDFYGK  --KTVDT
N_crassa_729649 IEQ-HMC-EGSCGGTYS--RYGVL  A-D  CDF  SYRMGNT FYGE  --KTVDT
A_aculeatus_391 VQQ-HMC-TGTCGGTYSDTT--RYSGT  P-D  CDF  PYRFGNTNFYGP  --KTVDN
A_niger_6164684 VSQ-HMC-DGSCGGTYSASG--RYSGT  P-D  CDF  PYR GNTDFYGP  --LTVDTN
P_janthinellum PSQ-HMC-TGQRCGGTYST--RYGGT  P-D  CDF  PYRMGVTNFYGP  --ETDTK
A_niger_6164682 SEQ-HMC-EGNDCGGTYS--RYGGT  P-D  CDF  PYRMGND FYGP  --KTDTG
H_ceramica_1218 VEQ-EIC-EGGCGGTYS--NRYGGT  P-D  CDF  PYR GNT FYGP  SSFT DT
H_jecorina_2238 VEQ-EIC-EGGCGGTYS--NRYGGT  P-D  CDF  PYR GNT FYGP  SSFT DT
T_viride_121854 VEQ-EIC-EGSCGGTYSG--RYGGT  P-D  CDF  PYR GNT FYGP  SSFT DT
T_viride_406299 VEQ-EIC-DGSCGGTYSG--RYGGT  P-D  CDF  PYR GNT FYGP  SSFT DT
T_harzianum_710 VEQ-HMC-SGSCGGTYSN--RYGGT  P-D  CDF  PYR GNT FYGP  SSFA DT
A_bisporus_3913 PQL-QRC-EGNTCS--VN--RYATE  P-D  CDF  SRMGDK FYGP  --MTVDTN
V_volvacea_5231 PQL-VRC-SGTACGG--GS--NRYGSI  H-D  LGFQ  LGMGRTRVR RV  RVKQFNR
T_longibrachiat T-----S-A  CGF  PYGSGYPN GP  --DTVDT
H_jecorina_1217 T-----S-A  CGF  PYGSGYK FYGP  --DTVDT
A_oryzae_246737 TQL-YEC-SGSGCG-----SGV  K-A  CGF  PYG GAKD FYGY  --LKVNTN
consensus  -gq-t-c--gd-cggtys--dry-g-CD--dGcdf-N-yrmgn-sfyg--G--tvdt
    
```

Figure 8F

20VW.A KFTVVTQFVANK---QGDLI-EIHRHYDNVIESAVVNISGPPK-INFINKCAAT
1A39 KPFTVVVTQFANR---RGKLE-KIHRFYVDGVIESFYTNKEGOPY-TNMDIFCEAT
6CEL KKLTVVVTQFES-----G-AINRYVNGVTFQPNAE-LGSYS-GNEINACTAE
1EG1.A KTFTTQFN DNGSPSGNLV-SITRKYONGVDIPSAQP-----G-GDITSC-----
P_chrysosporium KLETVVVTQFSDNTMGALV-EIHRLYNGQVIQNSVVNIPGNP-ATSITLCAQE
P_chrysospori_5 KLETVVVTQFSDNTMGALV-EIHRLYNGQVIQNSVVNIPGNP-ATSITLCAQE
P_chrysospori_6 KTFTVVVTQFSDNTTGNLT-EIHRLYVNGNVIPNSVVNITGG-VNSITPFCSQQ
P_chrysospori_7 KPFTVVVTQFNDGTAGTLT-EIHRLYVNGVIONSSVKIPGDP-VNSITNFCSQQ
P_chrysospori_8 KPFTVVVTQFNDGTAGTLT-EIHRLYVNGVIONSSVKIPGDL-VNSITNFCSQQ
P_chrysospori_9 KPFTVVVTQFNDGTAGTLT-EIHRLYVNGVIONSSVKIPGDP-VNSITNFCSQQ
P_chrysospor_10 KPFTDVTQFNDNTTGTLS-EIHRLYVNGVIONSVANIPGDP-VNSITNFCAQQ
P_chrysospor_11 KPFTVVVTQFNDNTTGTLS-EIHRLYVNGVIONSVANIPGDP-VNSITNFCAQQ
P_chrysospor_12 KPFTVVVTQFNDNTTGTLS-EIHRLYVNGVIONSVANIPGDP-VNSITNFCAQQ
I_lacteus_45863 KFTVVTQFDDGTSGNLA-EIHRFYVDGNVIPNSKVSIAGD-VNSITGFCTQQ
I_lacteus_45_14 OKFTVVTQFDDGTADGNLA-EIHRFYVNGVIVIPNSVVQITGDP-VNSITGFCTQQ
I_lacteus_45_15 KFTVVTQFDDGTANGDLH-EIHRLYVDGVIONSVVSIPGD-VDSITNFCAQQ
A_alternata_617 KKFTVVVTQF-----GTGDAM-EIHRFYVNGVTIAPASAPGEG-NSITTKFCDQQ
L_maculans_7804 KKFTVVVTQF-----GSGDAM-EIHRFYVNGVTIPDPSTIPGTG-NSITTFCDQAQ
C_parasitica_39 QKFTVVVTQFHG-----GSSLT-EISQYVGGTKIQPNSTWPTTG-YNSITPFCKAQ
C_carbonum_3913 KKMTVVVTQFG-----GSSLS-EIHRFYVNGVYKNSQSAAGTG-NSITSFCTAQ
H_grisea_134622 KPFTVVVTQFKD---ANGDLG-EIHRFYVDGLIPNSESTIPGEG-NSITQCDRQ
H_grisea_950686 KKITVVVTQFKD---ANGDLG-EIHRFYVDGLIPNSESTIPGEG-NSITQCDRQ
F_oxysporum_117 KKMTVVVTQFHKG---NGRSL-EITRLYVNGVIANSESKIAGNPG-SSITQFCSKQ
C_purpurea_1906 KIVVVVTQFKG---RDGSLR-EIHRFYVNGVIPNSVSRIRGPG-NSITQGFCAQ
H_thermoidea_40 KFTVVRF-----ERNR---SQFYVDGKIEVPPPTWPGPN-SADITPLCDAQ
H_thermoidea_74 KFTVVRF-----ERNR---SQFYVDGKIEVPPPTWPGPN-SADITPLCDAQ
L_maculans_7_26 KKMTVVVTQFG-----GNQLS-EIHRFYVDGVIANPEPTIPG-----CNTQ
N_crassa_729649 SKFTVVVTQFKD---AGDLA-EIHRFYVNGVVIENSQSNIDGSG-NSITQSFCKSQ
A_aculeatus_391 KPFTVVVTQFHDGTDGTLT-EIHRLYVNGVVIENGSTYTPASG-NSITSFCKAE
A_niger_6164684 SPFTVVVTQFDDGTSGTTLT-EIHRLYVNGEVIANGASTYSSNG-SSITSAFCESE
P_janthinellum SPFTVVVTQFNDGTGTLS-EIHRLYVGGVIGNPQSTIVGSG-NSITSFCAQ
A_niger_6164682 SKMTVVVTQFID---GSGSL-EIHRFYVNGNVIANADSNISGATG-NSITTFCTAQ
H_ceramica_1218 KKLTVVVTQF---E---SGA---INRYVNGVTFQPNAEGSYSG-NEINACTAE
H_jecorina_2238 KKLTVVVTQF---E---SGA---INRYVDGVTFOPNAEGSYSG-NEINACTAE
T_viride_121854 KKLTVVVTQF---E---SGA---INRYVNGVTFQPNAEGDYSG-NSIDCAA
T_viride_406299 KKLTVVVTQF---E---SGA---INRYVNGVTFQPNAEGDYSG-NSIDCAA
T_harzianum_710 KKLTVVVTQFA-----DGS---ISRHYVNGVKFQPNAGSYSG-NINTCAA
A_bisporus_3913 QPITVVVTQFDNGSDNGLQ-EIHRFYVNGQVIQNSNVNIPGDS-GNSIFAFCDDQA
V_volvacea_5231 SVVEPISTKQTTLHLGNLPKWSADCNVNGVIONSKVNIIPGPTMDSITFCNAQ
T_longibrachiat KTFTTQFN DNGSPSGNLV-SITRKYRNGVDIPSAK---PG--G--DITSC--CPSA
H_jecorina_1217 KTFTTQFN DNGSPSGNLV-SITRKYONGVDIPSAQ---PG--G--DITSC--CPSA
A_oryzae_246737 ETFTVVVTQFNDNTSGQLS-EIHRLYVNGQVIQNAAVTSGGKTV--DSITKPFCSGE
consensus kktftvvtqfvfvt-----s-g-l--eirrfyvQngkvi-n-----ipgv-g--nsitdefc--q

Figure 8G

```

20VW.A      G----ANE--MRLGGTKQMGDA SRGM AM WWSSEDFMA QG-----
1A39        G----SRK--MELG TQGMGEALTRGM AM IWWDQENME HG-----
6CEL        EAEFGG S--FSDKGGLTQFKKATSGGM VM WDDYANML STYPTNETS-STPG
1EG1.A     -----PS--ASAYGGLATMGKALSSGM VF IWNDNSQYMN SG-----
P_chrysosporium NAAFGG SS-FAQHGGLAQ GEALRSGM A IVNS-AADTL SNYPADADP-SAPG
P_chrysospori_5 NAAFGG SS-FAQHGGLAQ GEALRSGM A IVNS-AADTL SNYPADADP-SAPG
P_chrysospori_6 KKAFI NY-FAQHGGLAQ GQALRTGM AF ISDDPANHML SNPPSANP-AVPG
P_chrysospori_7 KTAFGD NY-FAQHGGLKQ GEALRTGM A IWDDYANML SNYPTNKDP-STPG
P_chrysospori_8 KTAFGD NY-FAQHGGLKQ GEALRTGM A IWDDYANML SNYPTNKDP-STPG
P_chrysospori_9 KTAFGD NY-FAQHGGLKQ GEALRTGM A IWDDYANML SNYPTNKDP-STPG
P_chrysospor_10 KTAFGD NW-FAQKGGLKQMGGEALGNMG A IWDDHAANML SDYPTDKDP-SAPG
P_chrysospor_11 KTAFGD NW-FAQKGGLKQMGGEALGNMG A IWDDHAANML SDYPTDKDP-SAPG
P_chrysospor_12 KTAFGD NW-FAQKGGLKQMGGEALGNMG A IWDDHAANML SDYPTDKDP-SAPG
I_lacteus_45863 KTAFGD NR-FAAQGGLKQMGGAALKSGM A WDDHAANML SDYPTTADA-SNPG
I_lacteus_45_14 KTVFGD NN-FAAKGGLKQMGGAALKSGM A WDDYAA ML SDYPTTADP-SQPG
I_lacteus_45_15 KSVFGD NY-FATLGGGLKMGGAALKSGM AM WDDHAASMQ SNYPADGDA-KPG
A_alternata_617 KAVFGD YT-FKDKGG A M KALANGM VM WDDHYSNML STYPTDKNPD-DLG
L_maculans_7804 KKAFGD KYT-FKDKGG A M PSTCNGM VM WDDHYSNML STYPTDKNPD-DAG
C_parasitica_39 KVEFND DV-FSEKGGLAQMGGA ADGM VM WDDHYANML STYPTDADA-SSPG
C_carbonum_3913 KKAFGD SS-FAALGGLNEMGASLARGH A WGDHAVNML STYPTDADP-SKPG
H_grisea_134622 KVAFGD IDD-FNRKGG KQMGKALAGPM VM IWDDHASNML STPVDAAG--KPG
H_grisea_950686 KVAFGD IDD-FNRKGG KQMGKALAGPM VM IWDDHASNML STPVDAAG--KPG
F_oxysporum_117 KSVFGD IDD-FSKKGGWNGMSDAL SAPM VM WDDHHSNML STYPTDST---KVG
C_purpurea_1906 KKMFGAHES-FNAKGG KGMSAA SKPM VM WDDHHSNML STYPTNSR---QRG
H_thermoidea_40 FRVFDDRNR-FAETGGFDA NEALTIPM VM IWDDHHSNML SSYPPEKAG--LPG
H_thermoidea_74 FRVFDDRNR-FAETGGFDA NEALTIPM VM IWDDHHSNML SSYPPEKAG--LPG
L_maculans_7_26 KKVFQ EAYPFNEFGG ASMSE SQGM VM WDDHYANML SNPREADP-AKPG
N_crassa_729649 KTAFGD IDD-FNKKGGLKQMGKALAQ M VM IWDDHAANML STYFPV---P-KVPG
A_aculeatus_391 KTLFGD NV-FETHGGLSAMDALGDGM V WDDHAADM L SDYPTTSCA-SSPG
A_niger_6164684 KTLFGD ENV-FDKHGGLGEGMGEA AKGM V WDDYAADML SDYPVNSSA-STPG
P_janthinellum KSAFGD NE-FSKHGG MAGMGA LADGM VM WDDHASDML STYPTNATS--TPG
A_niger_6164682 KKAFGDEDI-FAEHNGLAG SDA SS-M WDDYYASME SDYPENATA--DPG
H_ceramica_1218 EAEFGG S--FSDKGGLTQFKKATSGGM VM WDDYANML STYPTNETS-STPG
H_jecorina_2238 EAEFGG S--FSDKGGLTQFKKATSGGM VM WDDYANML STYPTNETS-STPG
T_viride_121854 EAEFGG S--FSDKGGLTQFKKATSGGM VM WDDYANML STYPTNETS-STPG
T_viride_406299 EAEFGG S--FSDKGGLTQFKKATSGGM VM WDDYANML STYPTNETS-STPG
T_harzianum_710 QTAFGG S--FTDKGGLAQ NKAFOGGM VM WDDYAVNML STYPTNATA-STPG
A_bisporus_3913 KEAFGDERS-FQDRGGLSGMGSALDRGM V IWDDHAVNML SDYPLDASP-SQPG
V_volvacea_5231 KTAFNDF S-FQOKGG A M SEALRRGM V IWDDHAANML SITSAACR-STPS
T_longibrachiat SA-----GGLATMGKALSSGM VF IWNDNSQYMN S-----G
H_jecorina_1217 SA-----GGLATMGKALSSGM VF IWNDNSQYMN S-----G
A_oryzae_246737 -----GSA--FNRLGGLLEEMGHALGRGM A IWNDAS S F M Q S-----G
consensus k--fgds--f--ggl-qmg-al--gmVLvmSiwddhaanmlWLDs-ypt-----s-pg

```

Figure 8H


```

20VW.A      -VAGPCDATEGDPKNVVKVQPNPE TFSNI I EIG-ST5
1A39        -EAGPCAKGEGAPSNVQVEPFPE TSNIG EIG-ST5-----QELQ
6CEL        AVRG CST SGVPAQVES SPNAK TFSNIKF PIG-ST5-----NP G
1EG1.A      -NAGPCS TEGNPSN LA NPNTH VFSNI DIG-STT
P_chrysosporium VARG CPQDS---AS P-EAPTPS VFSNIKL DIG-TT --G---AG AL-----FS
P_chrysospori_5 VARG CPQDS---AS P-EAPTPS VFSNIKL DIG-TT --G---AG AL-----F-
P_chrysospori_6 VARG MCSITSGNPADV GILNPSPY SFLNIKF SIG-TT -----RPA
P_chrysospori_7 VARG CATTSGVPAQ EA SPNAY VFSNIKF DIN-TT T-GT--VSS SV-----SS
P_chrysospori_8 VARG CATTSGVPAQ EA SPNAY VFSNIKF DIN-TT T-GT--VSS SV-----SS
P_chrysospori_9 VARG CATTSGVPAQ EA SPNAY VFSNIKF DIN-TT T-GT--VSS SV-----SS
P_chrysospor_10 VARG CATTSGVPSDVES VPNSQ VFSNIKF DIG-ST S-G---TS P-----NP
P_chrysospor_11 VARG CATTSGVPSDVES VPNSQ VFSNIKF DIG-ST S-G---TS P-----NP
P_chrysospor_12 VARG CATTSGVPSDVES VPNSQ VFSNIKF DIG-ST S-G---TS P-----NP
I_lacteus_45863 VARG CPTTSGFPRDVES SGSAT T SNIKF DIN-ST T-GTL-TTP GS-----SS
I_lacteus_45_14 VARG CPTTSGVPSQVEG EGSSS SNIKF DIN-ST T-GTLNPS PA-----GP
I_lacteus_45_15 VARG CSADSG PTNVES SASAS TFSNIKF DIN-ST T-G---TG T-----SP
A_alternata_617 TGRGECET SGVPADVES HADAT V SNIKF PIN-ST --G
L_maculans_7804 SGRGECAITSGVPADVES HPDAS SNIKF PIN-TT --G
C_parasitica_39 KQRG CATTSGVPADVES SDASAT SNIKF PIG-AT
C_carbonum_3913 AARG CPTTSGKPEDVEK SPDAT VFSNIKF PIG-ST --G-----QPA
H_grisea_134622 AERGACPTTSGVPAEVEAEAPNSN VFSNI F PIG-STVA-GLPGAGNGGN-----NG
H_grisea_950686 AERGACPTTSGVPAEVEAEAPNSN VFSNI F PIG-STVA-GLPGAGNGGN-----NG
F_oxysporum_117 SQRG CATTSGKPSD ERDVPNSK SFSNIKF PIG-ST --G-----KSDG-----TT
C_purpurea_1906 SKRG CPA SGRPTDVESAPDST VFSNIKF PIG-ST --G-----SRGK
H_thermoidea_40 GDRGPCPTTSGVPAEVEA YPNAQ V SNI F PIG-STV-----NV
H_thermoidea_74 GDRGPCPTTSGVPAEVEA YPDAQ V SNI F PIG-STV-----NV
L_maculans_7_26 VARRDCPTGGKPSEVEAANPNAQ VFSNIKF PIG-ST --G-----AHAA
N_crassa_729649 AYRG GPTTSGVPAEVEA APNSK AFSNIKF HGISP S-GGS-SGTPPS-----NP
A_aculeatus_391 VARG CPTTGNATYVEA YPNSY T SNIKF T N-ST S-GTSSGGS SSSTTLTKA
A_niger_6164684 VARG CSTDSGVPATVEAESPNAY T SNIKF PIG-ST SSGS--SSG GS-----SS
P_janthinellum AKRG CDI RR-PNTVESTYPNAY SNIKT PIN-ST T-GGTTSS TT-----TT
A_niger_6164682 VARG CD ESGVPATVEGAHPDSS TFSNIKF PIN-ST --G-----SA A
H_ceramica_1218 AVRG CST SGVPAQVES SPNAK TFSNIKF PIG-ST --G---NP GG-----NP
H_jecorina_2238 AVRG CST SGVPAQVES SPNAK TFSNIKF PIG-ST --G---NP GG-----NP
T_viride_121854 AVRG SST SGVPAQ ES SPNAK V SNIKF PIG-ST --G---NP GG-----NP
T_viride_406299 AVRG CST SGVPAQ ES SPNAK V SNIKF PIG-ST --G---NS GG-----NP
T_harzianum_710 AKRG CST SGVPAQVEA SPNSK V SNI F PIG-ST --G---GN GS-----NP
A_bisporus_3913 ISRG CSRDSGKPEDVEA AGGVQ V SNIKF DIN-ST --N---NNGG-----
V_volvacea_5231 EVH PLRESQ RSSHRSR T--RY T SNIKF PFN-ST --G---TTYT-----TG
T_longibrachiat -RAGPCS TEGNPSN LA NPGTH V SNI DIG-STTN-ST--GGNPP-----PP
H_jecorina_1217 -NAGPCS TEGNPSN LA NPNTH VFSNI DIG-STT -N---STAPP-----PP
A_oryzae_246737 GSAGPCNATEGNPAL EKLYPDTH KFSKI DIG-ST --G-----RH
consensus --rgsc-ttsgvpa-ve-q-pn--VvfsnikfGpig-sty--g-----s-----
    
```

Figure 8I

20VW.A
 1A39
 6CEL
 1EG1.A
 P_chrysosporium -GRSPPGPVPGS-APAS--S---ATA---TAPP--QCGGLGAGPTGVCPSPYTCQA
 P_chrysospori_5 -SGRS
 P_chrysospori_6
 P_chrysospori_7 SHSSTSTSSSHS-SSS--PPTQPTGV--TVPQ--QCGGIGGPT-TCASPYTCHV
 P_chrysospori_8 SHSSTSTSSSHS-SSS--PPTQPTGV--TVPQ--QCGGIGGPT-TCASPYTCHV
 P_chrysospori_9 SHSSTSTSSSHS-SSS--PPTQPTGV--TVPQ--QCGGIGGPT-TCASPYTCHV
 P_chrysospor_10 PGGST-TSSPVT-TSP--PP--PTGP--TVPQ--QCGGIGGPT-TCASPYTCHV
 P_chrysospor_11 PGGST-TSSPVT-TSP--PP--PTGP--TVPQ--QCGGIGGPT-TCASPYTCHV
 P_chrysospor_12 PGGST-TSSPVT-TSP--PP--PTGP--TVPQ--QCGGIGGPT-TCASPYTCHV
 I_lacteus_45863 PSSPASTSGSST-SAS--SASVPTQS--GTVAQ--QCGGIGGAT-TCVSPYTCHV
 I_lacteus_45_14 PVTSSPSEPSQS-TQS--QPAQPTQPA-GTAAQ--QCGGMGGPT-VCASPFTCHV
 I_lacteus_45_15 SSPAGFVSSSTS-VASQ--PT-QPAQG--TVAQ--QCGGTGGPT-VCASPFTCHV
 A_alternata_617
 L_maculans_7804
 C_parasitica_39
 C_carbonum_3913
 H_grisea_134622 GNPPPPPTTTTSS-APA--TTTASAGP---KAGR--Q-QCGGIGGPT-QCEEPYICTK
 H_grisea_950686 GNPPPPPTTTTSS-APA--TTTASAGP---KAGR--Q-QCGGIGGPT-QCEEPYTCTK
 F_oxysporum_117 PNPPASSSTTGS-STP--NP--PAG----SVDQ--QCGGQNGPT-TCKSPFTCKK
 C_purpurea_1906
 H_thermoidea_40
 H_thermoidea_74
 L_maculans_7_26
 N_crassa_729649 SSSASPTSSTAKPSST--TASNPSGT---GAAH--QCGGIGGPT-TCPEPYTCAK
 A_aculeatus_391 STSTTSSKTTTT-TSK--STTSSSST---NVAQL--QCGGQGPT-TCASG-TCTK
 A_niger_6164684 SSSSTTTKATST-TLK--TSTTSSGSSSTSAAQA--QCGGQGPT-TCVSGYTCTY
 P_janthinellum TSKSTSTSSSSK-TTT--VTTTTTSSGSS-GTGARD--QCGGNGGPT-TCVSPYTCTK
 A_niger_6164682
 H_ceramica_1218 PGGNR-GTTTTTR-RPA--TTGSSSPGP---TQSH--QCGGIGGPT-VCASGTTTCQV
 H_jecorina_2238 PGGNPPGTTTTT-TTS--SZ-PPPG---AHRR--QCGGIGGPT-VCASGTTTCQV
 T_viride_121854 PGGNPPGTTTT-P-RPA--STGSSSPGP---TQTH--QCGGIGGPT-VCASGSTTCQV
 T_viride_406299 PGGNPPGTTTTTR-RPA--STGSSSPGP---TQTH--QCGGIGGPT-VCASGSTTCQV
 T_harzianum_710 PGTSTTRAPPSS-TGS-----PTA---TQTH--QCGGTGGPT-RCASGYTCQV
 A_bisporus_3913 -GGGNPSPTTTR-----PNSP---AOTM--QCGGQGPT-ACQSPSTCHV
 V_volvacea_5231 SVPTTSTSTGTT-GSS--PP-QPTGV---TVPQ--QCGGIGGPT-TCASPTTCHV
 T_longibrachiat PPPASSTTFSTT-RRS--TTSSSPSC---TQTH--QCGGIGGCK-TCTSGTTCQY
 H_jecorina_1217 PASSTTFSTTRR-SST--SS--SPSC---TQTH--QCGGIGGCK-TCTSGTTCQY
 A_oryzae_246737
 consensus -----s-----fg-qcgg-gytg-t--c-s--tc--

Figure 8J

20VW.A
 1A39
 6CEL
 1EG1.A
 P_chrysosporium LNIYYSQ-CI
 P_chrysospori_5
 P_chrysospori_6
 P_chrysospori_7 LNPYYSQ-CY
 P_chrysospori_8 LNPYYSQ-CY
 P_chrysospori_9 LNPYYSQ-CY
 P_chrysospor_10 LNPYYSQ-CY
 P_chrysospor_11 LNPYYSQ-CY
 P_chrysospor_12 LNPCEISILSLQRSNADQYLQTTTSATKRRRLDTALQPRK
 I_lacteus_45863 VNAIYYSQ-CY
 I_lacteus_45_14 LNPYYSQ-CY
 I_lacteus_45_15 VNPYYSQ-CY
 A_alternata_617
 L_maculans_7804
 C_parasitica_39
 C_carbonum_3913
 H_grisea_134622 LNDIYYSQ-CL
 H_grisea_950686 LNDIYYSQ-CL
 F_oxysporum_117 INDYYSQ-CQ
 C_purpurea_1906
 H_thermoidea_40
 H_thermoidea_74
 L_maculans_7_26
 N_crassa_729649 DHDIYYSQ-CV
 A_aculeatus_391 QNDIYYSQ-CL
 A_niger_6164684 ENAYYSQ-CL
 P_janthinellum QNDIYYSQ-CL
 A_niger_6164682
 H_ceramica_1218 LNPYYSQ-CL
 H_jecorina_2238 LNPYYSQ-CL
 T_viride_121854 LNPYYSQ-CL
 T_viride_406299 LNPYYSQ-CL
 T_harzianum_710 LNPYYSQ-CL
 A_bisporus_3913 INDYYSQ-CF
 V_volvacea_5231 LNPYYSQ-CY
 T_longibrachiat GNDIYYSQ-CL
 H_jecorina_1217 SNDIYYSQ-CL
 A_oryzae_246737
 consensus -n-yysq-c-

Figure 8K

Figure 9A





Figure 9B

Thermal Melts of *H. jecorina* Cel7A and Variants






Figure 10: pRAX1

Figure 11: Destination vector pRAXdes2 for expression in *A. niger*




Figure 12: Replicative expression pRAXdesCBH1 vector of CBH1 genes under the control of the glucoamylase promoter.

