

**(12) STANDARD PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. **AU 2013399895 B2**

(54) Title  
**Synergistic stain removal through novel chelator combination**

(51) International Patent Classification(s)  
**C11D 3/06** (2006.01)                      **C11D 7/12** (2006.01)  
**C11D 3/10** (2006.01)                      **C11D 7/16** (2006.01)  
**C11D 3/33** (2006.01)                      **C11D 7/32** (2006.01)

(21) Application No: **2013399895**                      (22) Date of Filing: **2013.09.09**

(87) WIPO No: **WO15/032447**

(43) Publication Date: **2015.03.12**

(44) Accepted Journal Date: **2016.10.13**

(71) Applicant(s)  
**Ecolab USA Inc.**

(72) Inventor(s)  
**Foster, Tobias;Gohl, David;Klose, Sven;Kullwitz, Dirk;Mansergh, John;Meier, Timothy;Pathicheril, Beana**

(74) Agent / Attorney  
**Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000**

(56) Related Art  
**WO 2011/100344 A1**



(51) International Patent Classification:

*C11D 3/06* (2006.01)      *C11D 7/12* (2006.01)  
*C11D 3/10* (2006.01)      *C11D 7/16* (2006.01)  
*C11D 3/33* (2006.01)      *C11D 7/32* (2006.01)

(21) International Application Number:

PCT/EP2013/068611

(22) International Filing Date:

9 September 2013 (09.09.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant: **ECOLAB USA INC.** [US/US]; 370 Wabasha Street N, St. Paul, MN 55102-1390 (US).

(72) Inventors; and

(71) Applicants (*for US only*): **FOSTER, Tobias** [DE/DE]; Schirmerstr. 20, 50823 Köln (DE). **GOHL, David** [US/US]; 685 Stonewood Road, Eagan, MN 55123 (US). **KLOSE, Sven** [DE/DE]; Am Ehrenmahl 53, 45277 Essen (DE). **KULLWITZ, Dirk** [DE/DE]; Pappelallee 40, 47918 Tönisvorst (DE). **MANSERGH, John** [US/US]; 7176 Hidden Valley Cove S., 55016 Cottage Grove, MN 55016 (US). **MEIER, Timothy** [US/US]; 1351 Eagle Bluff Cir, Hastings, MN 55033 (US). **PATHICHERIL, Beana** [DE/DE]; Stephaniestr. 30, 40211 Düsseldorf (DE).

(74) Agent: **GODEMEYER BLUM LENZE PARTNER-SCHAFT, PATENTANWÄLTE - WERKPATENT**; An den Gärten 7, 51491 Overath (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*



**WO 2015/032447 A1**

(54) Title: SYNERGISTIC STAIN REMOVAL THROUGH NOVEL CHELATOR COMBINATION

(57) Abstract: The invention relates to a concentrated detergent composition comprising an alkali metal carbonate, methylglycinediacetic acid, glutamic acid N,N-diacetic acid, and alkali metal triphosphate. The composition is particularly suited to remove tea and coffee soil in warewashing applications.

## Synergistic stain removal through novel chelator combination

The present invention relates to concentrated detergent compositions comprising a mixture of chelators (complexing agents) for warewashing, especially adapted  
5 for the removal of tea and coffee soil.

It is known in the field of detergent chemistry that calcium and magnesium ions usually present in hard water can react with components of detergent compositions to form insoluble precipitates. This is a highly unfavorable effect as it  
10 causes the formation of scale on cleaned goods and negatively effects the detergent's capacity to remove soil.

Detergents therefore commonly comprise complexing agents that bind to metal ions and thereby reduce the concentration of free metal ions in aqueous systems.  
15 Most complexing agents act as polydentate ligands to form chelate complexes with the metal ions. Commonly used complexing agents are, for example, phosphates, citric acid, gluconic acid, methylglycinediacetic acid (MGDA), nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), diethylenetriaminepentaacetic acid (DTPA), hydroxyethylenediaminetriacetic acid  
20 (HEDTA), or iminodisuccinate (IDS).

By binding free magnesium or calcium ions, complexing agents reduce water hardness and prevent scale from forming. Complexing agents can also even help to redissolve scale by sequestering magnesium or calcium ions that are bound to  
25 and stabilize precipitated scale. Complexing agents thus serve a dual role by both reducing water hardness and redissolving scale. Complexing agents further may prevent metal ions from participating in typical chemical reactions, for example the chemical decomposition of peroxide compounds catalyzed by manganese, iron and copper ions. Complexing agents are therefore particularly used to enhance  
30 the performance of cleaning compositions comprising peroxide bleaches.

It is known that the quantity of complexing agents required to sequester a given concentration of metal ions depends on the binding stoichiometry of complexing agent to metal ion and on the dissociation constant of the binding equilibrium.

Complexing agents for use as water softeners are commonly characterized by their calcium binding capacity, which is a measure for the amount of calcium bound by a given amount of complexing agent at a given pH and temperature. For mixtures of complexing agents it is assumed that the total binding capacity of the mixture is the sum of the individual binding capacities. The total amount of  
5 complexing agent required for a detergent application can therefore be calculated as a function of the known calcium binding capacity and the water hardness.

Complexing agents are selected based on their calcium binding capacity, metal  
10 binding capacity in general, and their cost. In addition properties such as toxicology, detergent compatibility, and environmental restrictions have also to be considered. To make the use of complexing agents as cost efficient as possible, it is desirable to minimize the amount of complexing agent needed for a given application. There is therefore the need to increase the efficiency of the  
15 complexing agents.

The present invention deals with mild alkaline detergent compositions for the removal of tea and coffee soil in warewashing applications. Mild alkaline detergents are formulated on the basis of alkali carbonate as an alkaline source,  
20 in particular sodium carbonate. Tea and coffee soil is thought to comprise oxidized polyphenols (e.g. tannins) bridged by calcium silicate. This type of soil has proven to be particularly difficult to dissolve. It is therefore the object of the present invention to provide a highly efficient detergent composition for the removal of tea and coffee soil in warewashing applications.

25

It has surprisingly been found that the combination of the complexing agents methylglycinediacetic acid (MGDA), glutamic acid N,N-diacetic acid (GLDA) and sodium tripolyphosphate (STPP) exhibits synergy in a carbonate based detergent composition with regard to the removal of tea and coffee stains. By synergy it is  
30 meant that the total concentration of the three complexing agents required to achieve a cleaning effect is lower than what would be expected based on each agent's individual calcium binding capacity. This allows to minimize the amount of complexing agents used in a detergent composition.

According to a first aspect of the present invention there is provided a concentrated detergent composition comprising  
alkali metal carbonate,  
methylglycinediacetic acid,  
glutamic acid N,N-diacetic acid, and  
alkali metal tripolyphosphate.

In general, the concentrated detergent composition comprises an effective amount of alkali metal carbonate. In the context of the present invention, an effective amount of the alkali metal carbonate is an amount that provides a use solution having a pH of at least 6, preferably a pH of at least 8, more preferably a pH of 9.5 to 11, most preferably 10 to 10.3 measured at room temperature (20°C). For the purpose of determining the pH of the use solution, this use solution is defined as a solution of 1 g of the concentrated detergent composition dissolved in 1 liter distilled water.

To provide the required alkalinity, the concentrated detergent composition typically comprises at least 5 percent by weight alkali metal carbonate, preferably the composition comprises 10 to 80 percent by weight, more preferably 15 to 70 percent by weight, most preferably 20 to 60 percent by weight alkali metal carbonate.

Suitable alkali metal carbonates are for example sodium or potassium carbonate, sodium or potassium bicarbonate, sodium or potassium sesquicarbonate, and mixtures thereof.

Due to the use of an alkali metal carbonate as alkaline source, other alkaline sources such as alkali metal hydroxides are not required. Preferably, the concentrated detergent composition therefore does not comprise alkali metal hydroxides.

The concentrated detergent composition comprises the complexing agents methylglycinediacetic acid (MGDA), glutamic acid N,N-diacetic acid (GLDA), and an alkali metal tripolyphosphate. In the context of the present invention,

methylglycinediacetic acid and glutamic acid N,N-diacetic acid may be used as free acids or as salts. Commonly, the sodium salts of the mentioned compounds will be included in the detergent compositions. The alkali metal tripolyphosphate preferably is sodium tripolyphosphate (STPP).

5

The complexing agents are readily available to the person skilled in the art. For example, the trisodium salt of methylglycinediacetic acid is sold under the trademark Trilon M by BASF, and the tetrasodium salt of glutamic acid N,N-diacetic acid is available under the trademark Dissolvine GL from AkzoNobel.

10

The concentration of the three complexing agents is usually adjusted based on the amount of alkali metal carbonate present, such that upon dilution of the concentrated composition suitable working concentrations of both the alkali metal carbonate and the complexing agents are obtained. Preferably, the molar ratio of the sum of glutamic acid N,N-diacetic acid, methylglycinediacetic acid and alkali metal tripolyphosphate to alkali metal carbonate is 0.01 to 0.5, more preferably 0.05 to 0.12, most preferably 0.07 to 0.12.

15

The relative amounts of the three complexing agents may be adjusted in order to maximize the cleaning efficiency. Preferably, the molar ratio of methylglycinediacetic acid to alkali metal tripolyphosphate therefore is 0.14 to 14.3, more preferably 0.5 to 5, most preferably 1.35 to 1.7. In addition, the molar ratio of glutamic acid N,N-diacetic acid to the sum of methylglycinediacetic acid and alkali metal tripolyphosphate preferably is 0.03 to 29, more preferably 0.05 to 2, most preferably 0.08 to 0.45.

20  
25

In another preferred embodiment the total concentration of glutamic acid N,N-diacetic acid, methylglycinediacetic acid and alkali metal tripolyphosphate is 1 to 50 % by weight based on the total weight of the concentrated detergent composition, more preferably 14 to 28 % by weight, most preferably 18 to 26 % by weight. The amount of glutamic acid N,N-diacetic acid preferably is 1 to 30 % by weight based on the total weight of the concentrated detergent composition, more preferably 1 to 23 % by weight, most preferably 2 to 8 % by weight. The amount of methylglycinediacetic acid preferably is 1 to 30 % by weight based on the total

30

weight of the concentrated detergent composition, more preferably 2 to 22 % by weight, most preferably 8 to 10 % by weight. The amount of alkali metal tripolyphosphate preferably is 1 to 30 % by weight based on the total weight of the concentrated detergent composition, more preferably 2 to 23 % by weight, most  
5 preferably 8 to 10 % by weight.

The concentrated detergent composition of the present invention may further comprise at least one of the compounds selected from the list consisting of surfactants, bleaching agents, activating agents, chelating/sequestering agents,  
10 silicates, detergent fillers or binding agents, defoaming agents, anti-redeposition agents, enzymes, dyes, odorants, catalysts, threshold polymers, soil suspension agents, antimicrobials, and mixtures thereof.

A variety of surfactants can be used in the present composition, such as anionic,  
15 nonionic, cationic, and zwitterionic surfactants. The concentrated detergent composition can comprise 0.5 to 20 % by weight surfactant based on the total weight of the concentrated detergent composition, preferably 1.5 to 15 % by weight.

20 Suitable anionic surfactants are, for example, carboxylates such as alkylcarboxylates (carboxylic acid salts) and polyalkoxycarboxylates, alcohol ethoxylate carboxylates, nonylphenol ethoxylate carboxylates; sulfonates such as alkylsulfonates, alkylbenzenesulfonates, alkylarylsulfonates, sulfonated fatty acid esters; sulfates such as sulfated alcohols, sulfated alcohol ethoxylates, sulfated  
25 alkylphenols, alkylsulfates, sulfosuccinates, alkylether sulfates; and phosphate esters such as alkylphosphate esters. Exemplary anionic surfactants include sodium alkylarylsulfonate, alpha-olefinsulfonate, and fatty alcohol sulfates.

Suitable nonionic surfactants are, for example, those having a polyalkylene oxide  
30 polymer as a portion of the surfactant molecule. Such nonionic surfactants include, for example, chlorine-, benzyl-, methyl-, ethyl-, propyl-, butyl- and other like alkyl-capped polyethylene glycol ethers of fatty alcohols; polyalkylene oxide free nonionics such as alkyl polyglycosides; sorbitan and sucrose esters and their ethoxylates; alkoxyated ethylene diamine; alcohol alkoxyates such as alcohol

ethoxylate propoxylates, alcohol propoxylates, alcohol propoxylate ethoxylate propoxylates, alcohol ethoxylate butoxylates, and the like; nonylphenol ethoxylate, polyoxyethylene glycol ethers and the like; carboxylic acid esters such as glycerol esters, polyoxyethylene esters, ethoxylated and glycol esters of fatty acids, and  
5 the like; carboxylic amides such as diethanolamine condensates, monoalkanolamine condensates, polyoxyethylene fatty acid amides, and the like; and polyalkylene oxide block copolymers including an ethylene oxide/propylene oxide block copolymer such as those commercially available under the trademark Pluronic (BASF), and other like nonionic compounds. Silicone surfactants can  
10 also be used.

Suitable cationic surfactants include, for example, amines such as primary, secondary and tertiary monoamines with C<sub>18</sub> alkyl or alkenyl chains, ethoxylated alkylamines, alkoxylates of ethylenediamine, imidazoles such as a 1-(2-  
15 hydroxyethyl)-2-imidazoline, 2-alkyl-1-(2-hydroxyethyl)-2-imidazoline; and quaternary ammonium salts, as for example, alkylquaternary ammonium chloride surfactants such as n-alkyl(C<sub>12</sub>-C<sub>18</sub>)dimethylbenzyl ammonium chloride, n-tetradecyldimethylbenzylammonium chloride monohydrate, naphthylene-substituted quaternary ammonium chloride such as dimethyl-1-  
20 naphthylmethylammonium chloride. The cationic surfactant can be used to provide sanitizing properties.

Suitable zwitterionic surfactants include, for example, betaines, imidazolines, amine oxides, and propinates.

25

If the concentrated detergent composition is intended to be used in an automatic dishwashing or warewashing machine, the surfactants selected, if any surfactant is used, can be those that provide an acceptable level of foaming when used inside a dishwashing or warewashing machine. It should be understood that  
30 warewashing compositions for use in automatic dishwashing or warewashing machines are generally considered to be low-foaming compositions.

Suitable bleaching agents include, for example, peroxygen compounds, such as alkali metal percarbonates, in particular sodium percarbonate, alkali metal



perborates, alkali metal persulfates, urea peroxide, hydrogen peroxide; and hypochlorites, such as sodium hypochlorite or calcium hypochlorite. These compounds may be used, for example, as sodium lithium, potassium, barium, calcium, or magnesium salts. In another embodiment, the peroxygen source is an organic peroxide or hydroperoxide compound. According to a further embodiment, the peroxygen source is hydrogen peroxide prepared in situ using an electrochemical generator or other means of generating hydrogen peroxide in-situ.

10 Alkali metal percarbonates are particularly preferred bleaching agents. The bleaching agent may be present in an amount of 5 to 60 % by weight based on the total weight of the concentrated detergent composition, preferably 5 to 50 % by weight, most preferably 10 to 40 % by weight.

15 If the detergent composition includes a peroxygen compound, an activating agent may be included to further increase the activity of the peroxygen compound. Suitable activating agents include sodium-4-benzoyloxy benzene sulphonate (SBOBS); N,N,N',N'-tetraacetyl ethylene diamine (TAED); sodium-1-methyl-2-benzoyloxy benzene-4-sulphonate; sodium-4-methyl-3-benzoyloxy benzoate; 20 SPCC trimethyl ammonium toluoyloxy benzene sulphonate; sodium nonanoyloxybenzene sulphonate, sodium 3,5,5,-trimethyl hexanoyloxybenzene sulphonate; penta acetyl glucose (PAG); octanoyl tetra acetyl glucose and benzoyl tetracetyl glucose. The concentrated detergent composition may comprise an activating agent or a mixture of activating agents at a concentration of 1 to 8 % by weight based on the total weight of the concentrated detergent 25 composition, preferably 2 to 5 % by weight.

The detergent composition may comprise further chelating/sequestering agents in addition to the complexing agents mentioned above. Suitable additional 30 chelating/sequestering agents are, for example, citrate, aminocarboxylic acid, condensed phosphate, phosphonate, and polyacrylate. A chelating agent in the context of the present invention is a molecule capable of coordinating (i.e., binding) the metal ions commonly found in natural water to prevent the metal ions from interfering with the action of the other deterative ingredients of a cleaning

composition. Chelating/sequestering agents can generally be referred to as a type of builder. The chelating/sequestering agent may also function as a threshold agent when included in an effective amount. The concentrated detergent composition can include 0.1 to 70 % by weight of a chelating/sequestering agent based on the total weight of the concentrated detergent composition, preferably 5 to 60 % by weight, more preferably 5 to 50 % by weight, most preferably 10 to 40 % by weight.

Suitable aminocarboxylic acids include, for example, N-hydroxyethyliminodiacetic acid, nitrilotriacetic acid (NTA), ethylenediaminetetraacetic acid (EDTA), N-hydroxyethyl-ethylenediaminetriacetic acid (HEDTA), and diethylenetriaminepentaacetic acid (DTPA).

Examples of condensed phosphates include sodium and potassium orthophosphate, sodium and potassium pyrophosphate, sodium hexametaphosphate, and the like. A condensed phosphate may also assist, to a limited extent, in solidification of the composition by fixing the free water present in the composition as water of hydration.

The composition may include a phosphonate such as 1-hydroxyethane-1,1-diphosphonic acid  $\text{CH}_3\text{C}(\text{OH})[\text{PO}(\text{OH})_2]_2$  (HEDP); amino tri(methylenephosphonic acid)  $\text{N}[\text{CH}_2\text{PO}(\text{OH})_2]_3$ ; aminotri(methylenephosphonate), sodium salt  $(\text{NaO})(\text{HO})\text{P}(\text{OCH}_2\text{N}[\text{CH}_2\text{PO}(\text{ONa})_2]_2)$ ; 2-hydroxyethyliminobis(methylenephosphonic acid)  $\text{HOCH}_2\text{CH}_2\text{N}[\text{CH}_2\text{PO}(\text{OH})_2]_2$ ; diethylenetriaminepenta(methylenephosphonic acid)  $(\text{HO})_2\text{POCH}_2\text{N}[\text{CH}_2\text{CH}_2\text{N}[\text{CH}_2\text{PO}(\text{OH})_2]_2]_2$ ; diethylenetriaminepenta(methylenephosphonate), sodium salt  $\text{C}_9\text{H}_{(28-x)}\text{N}_3\text{Na}_x\text{O}_{15}\text{P}_5$  ( $x=7$ ); hexamethylenediamine(tetramethylenephosphonate), potassium salt  $\text{C}_{10}\text{H}_{(28-x)}\text{N}_2\text{K}_x\text{O}_{12}\text{P}_4$  ( $x=6$ ); bis(hexamethylene)triamine(pentamethylenephosphonic acid)  $(\text{HO})_2\text{POCH}_2\text{N}[(\text{CH}_2)_6\text{N}[\text{CH}_2\text{PO}(\text{OH})_2]_2]_2$ ; and phosphorus acid  $\text{H}_3\text{PO}_3$ .

Preferred phosphonates are 1-Hydroxy Ethylidene-1,1-Diphosphonic Acid (HEDP), aminotris(methylenephosphonic acid) (ATMP) and Diethylenetriamine penta(methylene phosphonic acid) (DTPMP).

- 5 A neutralized or alkaline phosphonate, or a combination of the phosphonate with an alkali source prior to being added into the mixture such that there is little or no heat or gas generated by a neutralization reaction when the phosphonate is added is preferred. The phosphonate can comprise a potassium salt of an organo phosphonic acid (a potassium phosphonate). The potassium salt of the
- 10 phosphonic acid material can be formed by neutralizing the phosphonic acid with an aqueous potassium hydroxide solution during the manufacture of the solid detergent. The phosphonic acid sequestering agent can be combined with a potassium hydroxide solution at appropriate proportions to provide a stoichiometric amount of potassium hydroxide to neutralize the phosphonic acid.
- 15 A potassium hydroxide having a concentration of from about 1 to about 50 wt % can be used. The phosphonic acid can be dissolved or suspended in an aqueous medium and the potassium hydroxide can then be added to the phosphonic acid for neutralization purposes.
- 20 The chelating/sequestering agent may also be a water conditioning polymer that can be used as a form of builder. Exemplary water conditioning polymers include polycarboxylates. Exemplary polycarboxylates that can be used as water conditioning polymers include polyacrylic acid, maleic/olefin copolymer, acrylic/maleic copolymer, polymethacrylic acid, acrylic acid-methacrylic acid
- 25 copolymers, hydrolyzed polyacrylamide, hydrolyzed polymethacrylamide, hydrolyzed polyamide-methacrylamide copolymers, hydrolyzed polyacrylonitrile, hydrolyzed polymethacrylonitrile, and hydrolyzed acrylonitrile-methacrylonitrile copolymers.
- 30 The concentrated detergent composition may include the water conditioning polymer in an amount of 0.1 to 20 % by weight based on the total weight of the concentrated detergent composition, preferably 0.2 to 5 % by weight.

Silicates may be included in the concentrated detergent composition as well. Silicates soften water by the formation of precipitates that can be easily rinsed away. They commonly have wetting and emulsifying properties, and act as buffering agents against acidic compounds, such as acidic soil. Further, silicates  
5 can inhibit the corrosion of stainless steel and aluminium by synthetic detergents and complex phosphates. A particularly well suited silicate is sodium metasilicate, which can be anhydrous or hydrated. The concentrated detergent composition may comprise 1 to 10 % by weight silicates based on the total weight of the concentrated detergent composition.

10

The composition can include an effective amount of detergent fillers or binding agents. Examples of detergent fillers or binding agents suitable for use in the present composition include sodium sulfate, sodium chloride, starch, sugars, and C<sub>1</sub>-C<sub>10</sub> alkylene glycols such as propylene glycol. The detergent filler may be  
15 included an amount of 1 to 20 % by weight based on the total weight of the concentrated detergent composition, preferably 3 to 15 % by weight.

A defoaming agent for reducing the stability of foam may also be included in the composition to reduce foaming. The defoaming agent can be provided in an  
20 amount of 0.01 to 20 % by weight based on the total weight of the concentrated detergent composition.

Suitable defoaming agents include, for example, ethylene oxide/propylene block copolymers such as those available under the name Pluronic N-3, silicone  
25 compounds such as silica dispersed in polydimethylsiloxane, polydimethylsiloxane, and functionalized polydimethylsiloxane, fatty amides, hydrocarbon waxes, fatty acids, fatty esters, fatty alcohols, fatty acid soaps, ethoxylates, mineral oils, polyethylene glycol esters, defoaming emulsions and alkyl phosphate esters such as monostearyl phosphate.

30

The composition can include an anti-redeposition agent for facilitating sustained suspension of soils in a cleaning solution and preventing the removed soils from being redeposited onto the substrate being cleaned. Examples of suitable anti-redeposition agents include fatty acid amides, fluorocarbon surfactants, complex

phosphate esters, styrene maleic anhydride copolymers, and cellulosic derivatives such as hydroxyethyl cellulose, hydroxypropyl cellulose, and the like. The anti-redeposition agent can be included in an amount of 0.01 to 25 % by weight based on the total weight of the concentrated detergent composition, preferably 1 to 5 %  
5 by weight.

The composition may include enzymes that provide desirable activity for removal of protein-based, carbohydrate-based, or triglyceride-based soil. Although not limiting to the present invention, enzymes suitable for the cleaning composition  
10 can act by degrading or altering one or more types of soil residues encountered on crockery thus removing the soil or making the soil more removable by a surfactant or other component of the cleaning composition. Suitable enzymes include a protease, an amylase, a lipase, a gluconase, a cellulase, a peroxidase, a catalase, or a mixture thereof of any suitable origin, such as vegetable, animal,  
15 bacterial, fungal or yeast origin. The concentrated detergent composition may comprise 0.01 to 30 % by weight enzymes based on the total weight of the concentrated detergent composition, preferably 0.01 to 15 % by weight, more preferably 0.01 to 10 % by weight, most preferably 0.01 to 8 % by weight.

20 Examples of proteolytic enzymes which can be employed in the cleaning composition of the invention include (with trade names) Savinase®; a protease derived from *Bacillus lentus* type, such as Maxacal®, Opticlean®, Durazym®, and Properase®; a protease derived from *Bacillus licheniformis*, such as Alcalase®, Maxatase®, Deterzyme®, or Deterzyme PAG 510/220; a protease derived from  
25 *Bacillus amyloliquefaciens*, such as Primase®; and a protease derived from *Bacillus alcalophilus*, such as Deterzyme APY. Exemplary commercially available protease enzymes include those sold under the trade names Alcalase®, Savinase®, Primase®, Durazym®, or Esperase® by Novo Industries A/S (Denmark); those sold under the trade names Maxatase®, Maxacal®, or  
30 Maxapem® by Gist-Brocades (Netherlands); those sold under the trade names Purafect®, Purafect OX, and Properase by Genencor International; those sold under the trade names Opticlean® or Optimase® by Solvay Enzymes; those sold

under the tradenames Deterzyme®, Deterzyme APY, and Deterzyme PAG 510/220 by Deerland Corporation, and the like.

Preferred proteases will provide good protein removal and cleaning performance, will not leave behind a residue, and will be easy to formulate with and form stable products. Savinase®, commercially available from Novozymes, is a serine-type endo-protease and has activity in a pH range of 8 to 12 and a temperature range from 20 °C to 60 °C. Savinase is preferred when developing a liquid concentrate. A mixture of proteases can also be used. For example, Alcalase®, commercially available from Novozymes, is derived from *Bacillus licheniformis* and has activity in a pH range of 6.5 to 8.5 and a temperature range from 45 °C to 65 °C. And Esperase®, commercially available from Novozymes, is derived from *Bacillus* sp. and has an alkaline pH activity range and a temperature range from 50 °C to 85 °C. A combination of Esperase and Alcalase is preferred when developing a solid concentrate because they form a stable solid. In some embodiments, the total protease concentration in the concentrate product is from about 1 to about 15 % by weight, from about 5 to about 12 % by weight, or from about 5 to about 10 % by weight. In some embodiments, there is at least 1-6 parts of Alcalase for every part of Esperase (e.g., Alcalase:Esperase of 1:1, 2:1, 3:1, 4:1, 5:1, or 6:1).

20

Detergent proteases are described in patent publications including: GB 1,243,784, WO 9203529 A (enzyme/inhibitor system), WO 9318140 A, and WO 9425583 (recombinant trypsin-like protease) to Novo; WO 9510591 A, WO 9507791 (a protease having decreased adsorption and increased hydrolysis), WO 95/30010, WO 95/30011, WO 95/29979, to Procter & Gamble; WO 95/10615 (*Bacillus amyloliquefaciens* subtilisin) to Genencor International; EP 130,756 A (protease A); EP 303,761 A (protease B); and EP 130,756 A. A variant protease is preferably at least 80% homologous, preferably having at least 80% sequence identity, with the amino acid sequences of the proteases in these references.

30

Mixtures of different proteolytic enzymes may be incorporated into the disclosed compositions. While various specific enzymes have been described above, it is to be understood that any protease which can confer the desired proteolytic activity to the composition may be used.

The disclosed compositions can optionally include different enzymes in addition to the protease. Exemplary enzymes include amylase, lipase, cellulase, and others.

5 Exemplary amylase enzymes can be derived from a plant, an animal, or a microorganism. The amylase may be derived from a microorganism, such as a yeast, a mold, or a bacterium. Exemplary amylases include those derived from a *Bacillus*, such as *B. licheniformis*, *B. amyloliquefaciens*, *B. subtilis*, or *B. stearothermophilus*. The amylase can be purified or a component of a microbial  
10 extract, and either wild type or variant (either chemical or recombinant).

Exemplary amylase enzymes include those sold under the trade name Rapidase by Gist-Brocades® (Netherlands); those sold under the trade names Termamyl®, Fungamyl® or Duramyl® by Novo; those sold under the trade names Purastar  
15 STL or Purastar OXAM by Genencor; those sold under the trade names Thermozyme® L340 or Deterzyme® PAG 510/220 by Deerland Corporation; and the like. A mixture of amylases can also be used.

Exemplary cellulase enzymes can be derived from a plant, an animal, or a  
20 microorganism, such as a fungus or a bacterium. Cellulases derived from a fungus include the fungus *Humicola insolens*, *Humicola* strain DSM1800, or a cellulase 212-producing fungus belonging to the genus *Aeromonas* and those extracted from the hepatopancreas of a marine mollusk, *Dolabella Auricula* Solander. The cellulase can be purified or a component of an extract, and either  
25 wild type or variant (either chemical or recombinant).

Examples of cellulase enzymes include those sold under the trade names Carezyme® or Celluzyme® by Novo; under the tradename Cellulase by Genencor; under the tradename Deerland Cellulase 4000 or Deerland Cellulase  
30 TR by Deerland Corporation; and the like. A mixture of cellulases can also be used.

Exemplary lipase enzymes can be derived from a plant, an animal, or a microorganism, such as a fungus or a bacterium. Exemplary lipases include those

derived from a *Pseudomonas*, such as *Pseudomonas stutzeri* ATCC 19.154, or from a *Humicola*, such as *Humicola lanuginosa* (typically produced recombinantly in *Aspergillus oryzae*). The lipase can be purified or a component of an extract, and either wild type or variant (either chemical or recombinant).

5

Exemplary lipase enzymes include those sold under the trade names Lipase P "Amano" or "Amano-P" by Amano Pharmaceutical Co. Ltd., Nagoya, Japan or under the trade name Lipolase® by Novo, and the like. Other commercially available lipases include Amano-CES, lipases derived from *Chromobacter viscosum*, e.g. *Chromobacter viscosum* var. *lipolyticum* NRRLB 3673 from Toyo Jozo Co., Tagata, Japan; *Chromobacter viscosum* lipases from U.S. Biochemical Corp., U.S.A. and Disoynt Co., and lipases derived from *Pseudomonas gladioli* or from *Humicola lanuginosa*. A preferred lipase is sold under the trade name Lipolase® by Novo. A mixture of lipases can also be used.

15

Additional suitable enzymes include a cutinase, a peroxidase, a gluconase, and the like. Exemplary cutinase enzymes are described in WO 8809367 A to Genencor. Exemplary peroxidases include horseradish peroxidase, ligninase, and haloperoxidases such as chloro- or bromo-peroxidase. Exemplary peroxidases are also disclosed in WO 89099813 A and WO 8909813 A to Novo.

20

These additional enzymes can be derived from a plant, an animal, or a microorganism. The enzyme can be purified or a component of an extract, and either wild type or variant (either chemical or recombinant). Mixtures of different additional enzymes can be used.

25

Various dyes, odorants including perfumes, and other aesthetic enhancing agents can be included in the composition. Dyes may be included to alter the appearance of the composition, as for example, Direct Blue 86 (Miles), Fastsol Blue (Mobay Chemical Corp.), Acid Orange 7 (American Cyanamid), Basic Violet 10 (Sandoz), Acid Yellow 23 (GAF), Acid Yellow 17 (Sigma Chemical), Sap Green (Keystone Aniline and Chemical), Metanil Yellow (Keystone Aniline and Chemical), Acid Blue 9 (Hilton Davis), Sandolan Blue/Acid Blue 182 (Sandoz), Hisol Fast Red (Capitol Color and Chemical), Fluorescein (Capitol Color and Chemical), and Acid Green 25 (Ciba-Geigy).

30



Fragrances or perfumes that may be included in the compositions include, for example, terpenoids such as citronellol, aldehydes such as amyl cinnamaldehyde, a jasmine such as C1S-jasmine or jasmal, and vanillin.

5

The concentrated detergent composition may be provided, for example, in the form of a solid, a powder, a liquid, a gel or a paste. Preferably, the concentrated detergent composition is provided in the form of a solid or a powder.

- 10 The components used to form the concentrated detergent composition can include an aqueous medium such as water as an aid in processing. It is expected that the aqueous medium will help provide the components with a desired viscosity for processing. In addition, it is expected that the aqueous medium may help in the solidification process when is desired to form the concentrated
- 15 detergent composition as a solid. When the concentrated detergent composition is provided as a solid, it can, for example, be provided in the form of a block or pellet. It is expected that blocks will have a size of at least about 5 grams, and can include a size of greater than about 50 grams. It is expected that the concentrated detergent composition will include water in an amount of 0.001 to 50 % by weight
- 20 based on the total weight of the concentrated detergent composition, preferably 2 to 20 % by weight.

- When the components that are processed to form the concentrated detergent composition are processed into a block, it is expected that the components can be
- 25 processed by a known solidification technique, such as for example extrusion techniques or casting techniques. In general, when the components are processed into a block, the amount of water present in the concentrated detergent composition should be 0.001 to 40 % by weight based on the total weight of the concentrated detergent composition, preferably 0.001 to 20 % by weight. If the
- 30 components are processed by extrusion techniques, it is believed that the concentrated detergent composition can include a relatively smaller amount of water as an aid for processing compared with the casting techniques. In general, when preparing the solid by extrusion, it is expected that the concentrated detergent composition can contain 0.001 to 20 % by weight water based on the

total weight of the concentrated detergent composition. When preparing the solid by casting, it is expected that the amount of water is 0.001 to 40 % by weight based on the total weight of the concentrated detergent composition.

Also disclosed herein is a use solution of the concentrated detergent composition. The use solution is an aqueous solution of 0.1 to 10 g concentrated detergent composition per liter of the aqueous solution, preferably 0.5 to 5 g/l, most preferably 1 to 1.5 g/l.

According to a second aspect of the present invention there is provided an aqueous solution comprising 0.1 to 10 g/l of the concentrated detergent composition according to the first aspect of the invention.

Due to the synergy achieved by the inventive combination of complexing agents it is possible to formulate a use solution on the basis of hard water. The term "hard water" used herein is defined based on the concentration of  $\text{CaCO}_3$ . According to the US Geological Survey, water having a concentration of at least 61 mg/l  $\text{CaCO}_3$  is qualified as moderately hard water, a concentration of at least 121 mg/l  $\text{CaCO}_3$  is qualified as hard water, and a concentration of at least 181 mg/l  $\text{CaCO}_3$  as very hard water.

Generally the present invention is not limited to the case of hard water. In a preferred embodiment, however, the water used to prepare the use solution has a hardness of at least 50 mg/l  $\text{CaCO}_3$ , more preferably at least 61 mg/l  $\text{CaCO}_3$ , even more preferably at least 85 mg/l, most preferably at least 121 mg/l.

Also disclosed herein is the use of a concentrated detergent composition as described above as a warewashing detergent for the removal of tea and coffee soil. This soil is characterized by the presence of oxidized polyphenols and calcium silicates. The concentrated detergent composition may therefore be generally be used as a warewashing detergent for the removal of solid comprising oxidized polyphenols and calcium silicates.

According to a third aspect of the present invention there is provided use of a concentrated detergent composition according to the first aspect of the invention as a warewashing detergent for the removal of soil comprising oxidized polyphenols and calcium silicates.

Preferably, the concentrated detergent composition is diluted at a concentration of 0.1 to 10 g of concentrated detergent composition per liter of the final solution, preferably 0.5 to 5 g/l, most preferably 1 to 1.5 g/l to provide a use solution. Importantly, the present invention allows to use hard water for dilution of the detergent composition. In a preferred embodiment, the concentrated detergent

composition is therefore diluted with water having a hardness of at least 50 mg/l CaCO<sub>3</sub>, more preferably at least 61 mg/l CaCO<sub>3</sub>, even more preferably at least 85 mg/l, most preferably at least 121 mg/l to provide a use solution.

5

## Examples

The following examples illustrate the invention by testing the removal of tea and coffee soil from ceramic tiles.

- 10 Ceramic tiles (5.1 x 15.2 cm white, glazed ceramic tiles ) were stained with tea soil (Lipton brand tea) according to the following procedure. Hard water having a hardness of > 249.9 mg/l CaCO<sub>3</sub> was heated to > 71 °C. The tea was then mixed into the hot hard water. The ceramic tiles were then immersed into the tea for 1 min and then taken out for 1 min to dry. This procedure was repeated until a stain  
15 was formed, which was typically after 25 cycles. The tiles were then cured for 48 hrs at room temperature. At this time the tiles are ready for testing.

- Cleaning test were carried out in a standard automatic dishwasher. The cleaning efficiency was evaluated by visually comparing the amount of soil left on the tiles  
20 after one full cleaning cycle to the amount of soil on the tiles before the cleaning procedure. The results were rated according to table 1:

**Table 1**

Rating	% of stain removal
1	100
2	80 – 99.9
3	20 – 79.9
4	< 20
5	no removal

- 25 A rating of 1 was considered to be an excellent result. A rating of 2 (at least 80 % stain removal) was considered to be an acceptable cleaning performance.

The complexing agents shown in table 2 were tested for their effect on cleaning efficiency. For each complexing agent the theoretical concentration of the 100 % active compound required to cover 85.5 mg/l CaCO<sub>3</sub> of water hardness was calculated on the basis of the calcium binding capacity and the activity of the raw materials. The concentration given relates to the respective sodium salts. It should be noted that the calcium binding capacities in table 2 give the amount of CaCO<sub>3</sub> bound by a given amount of the raw material having an activity that can be lower than 100%, as specified in table 2.

10 **Table 2**

<b>Complexing agent</b>	<b>Trade name</b>	<b>Ca-binding capacity of raw material (mg/g)</b>	<b>Activity of raw material</b>	<b>Amount of 100% active material required for 85.5 mg/l CaCO<sub>3</sub> (mg/l)</b>
methylglycinediacetic acid (MGDA)	Trilon M Granules SG	310	83 %	229
sodium tripolyphosphate (STPP)		232	100 %	369
glutamic acid N,N-diacetic acid (GLDA)	Dissolvine 47S	138.2	47 %	291
iminodisuccinate (IDS)	Baypure CX100	111.8	34 %	260

The cleaning efficiency of different detergent formulations containing 1000 mg/l sodium carbonate and varying amounts of complexing agents was tested. All formulations were prepared in water having a hardness of 85.5 mg/l CaCO<sub>3</sub>. The concentrations given relate to the concentrations of the 100% active compounds in the use solution.

A first series of tests involved a combination of varying amounts of MGDA, STPP, and GLDA. Based on the calcium binding capacities and activities of the raw materials given in table 2, the theoretical amount of water hardness (expressed in mg/l CaCO<sub>3</sub>) covered was calculated for each formulation and was compared to the cleaning effect achieved by the formulation. The test data are shown in table 3.

**Table 3**

<b>Example</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>
GLDA (mg/l)	75	100	100	75	50	25	100	75	50	25
MGDA (mg/l)	100	100	125	125	125	125	125	125	125	125
STPP (mg/l)	100	100	125	125	125	125	100	100	100	100
CaCO <sub>3</sub> covered theoretically (mg/l)	82,6	89,9	105,0	97,7	90,4	83,0	99,2	91,9	84,6	77,2
Rating	2	1	1	1	1	1	1	1	1	2

The results showed that with the inventive combination of complexing agents in a carbonate based cleaning solution, acceptable to excellent cleaning results can be achieved even at a total concentration of complexing agents less than what would be theoretical required to cover 85.5 mg/l CaCO<sub>3</sub> of water hardness (examples 1, 6, 9 and 10). The combination of GLDA, MGDA, and STPP in a carbonate based detergent composition therefore exhibits synergy with respect to the cleaning effect. This allow to minimize the total amount of complexing agents used in a detergent composition.

A second series involved the combination of MGDA, STPP, and IDS (table 4).

<b>Example</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>10</b>
IDS (mg/l)	75	100	100	75	50	25	100	75	50	25
MGDA (mg/l)	100	100	125	125	125	125	125	125	125	125
STPP (mg/l)	100	100	125	125	125	125	100	100	100	100
CaCO <sub>3</sub> covered theoretically (mg/l)	85,2	93,4	108,6	100,3	92,1	83,9	102,8	94,5	86,3	78,1
Rating	3	2	1	1	1	2	1	2	2	3

The results of the second series showed that the overall cleaning performance of the combination of IDS, MGDA, and STPP is lower than for the combination of GLDA, MGDA, and STPP, even though the total concentrations of complexing agents are the same. This is unexpected as the theoretically required amount of IDS to cover 85.5 mg/l CaCO<sub>3</sub> of water hardness is lower than for GLDA and hence IDS should be more effective than GLDA (table 2). Further, as the total concentration of complexing agents drops below the amount required to fully cover 85.5 mg/l CaCO<sub>3</sub> of water hardness (examples 1 and 10) the cleaning efficiency becomes unacceptable, in contrast to the combination of GLDA, MGDA, and STPP. The combination of IDS, MGDA, and STPP therefore does not exhibit synergy.

**CLAIMS:**

1. A concentrated detergent composition comprising alkali metal carbonate, methylglycinediacetic acid, glutamic acid N,N-diacetic acid, and alkali metal tripolyphosphate.
2. The concentrated detergent composition according to any one of the preceding claims, wherein the molar ratio of methylglycinediacetic acid to alkali metal tripolyphosphate is 0.14 to 14.3.
3. The concentrated detergent composition according to any one of the preceding claims, wherein the molar ratio of glutamic acid N,N-diacetic acid to the sum of methylglycinediacetic acid and alkali metal tripolyphosphate is 0.03 to 29.
4. The concentrated detergent composition according to any one of the preceding claims, wherein the molar ratio of the sum of glutamic acid N,N-diacetic acid, methylglycinediacetic acid and alkali metal tripolyphosphate to alkali metal carbonate is 0.01 to 0.5.
5. The concentrated detergent composition according to any of the preceding claims, wherein the composition comprises at least 5 % by weight alkali metal carbonate.
6. The concentrated detergent composition according to any one of the preceding claims, wherein the alkali metal carbonate is sodium or potassium carbonate, sodium or potassium bicarbonate, sodium or potassium sesquicarbonate, or a mixture thereof.
7. The concentrated detergent composition according to any one of the preceding claims, wherein the alkali metal tripolyphosphate is sodium tripolyphosphate.
8. The concentrated detergent composition according to any one of the preceding claims, wherein the composition provides a pH of at least 6 when diluted in distilled water at a concentration of 1 g/l and measured at a temperature of 20°C.
9. The concentrated detergent composition according to any one of the preceding claims, wherein the composition further comprises at least one of the compounds selected from the list consisting of surfactants, bleaching agents, activating agents, chelating/sequestering agents, silicates, detergent fillers or binding agents, defoaming agents, anti-redeposition agents,



enzymes, dyes, odorants, catalysts, threshold polymers, soil suspension agents, antimicrobials and mixtures thereof.

10. The concentrated detergent composition according to any one of the preceding claims, wherein the composition is provided in the form of a solid, a powder, a liquid, a gel, or a paste.

11. An aqueous solution comprising 0.1 to 10 g/l of the concentrated detergent composition according to any one of claims 1 to 10.

12. Use of a concentrated detergent composition according to any one of claims 1 to 10 as a warewashing detergent for the removal of soil comprising oxidized polyphenols and calcium silicates.

13. The use according to claim 12, wherein the concentrated detergent composition is diluted to provide a use solution with a concentration of 0.1 to 10 g/l.

14. The use according to claim 13, wherein the concentrated detergent composition is diluted with water having a hardness of at least 50 mg/l CaCO<sub>3</sub>.

15. The use according to any one of claims 12 to 14, wherein the warewashing detergent is used for the removal of tea and coffee soil.

**Ecolab USA Inc.**

**Patent Attorneys for the Applicant/Nominated Person**

**SPRUSON & FERGUSON**