

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2007253903 B2**

(54) Title
Use of non-agrobacterium bacterial species for plant transformation

(51) International Patent Classification(s)
C12N 15/82 (2006.01)

(21) Application No: **2007253903** (22) Date of Filing: **2007.05.16**

(87) WIPO No: **WO07/137075**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/800,872	2006.05.16	US

(43) Publication Date: **2007.11.29**

(44) Accepted Journal Date: **2012.07.19**

(71) Applicant(s)
Monsanto Technology LLC

(72) Inventor(s)
Williams, Edward;Shen, Junjiang;Ye, Xudong

(74) Agent / Attorney
Pizzeys, Level 2, Woden Plaza Offices Woden Town Square Woden, Canberra, ACT, 2606

(56) Related Art
US 2005/0289672 A1

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2007 (29.11.2007)

PCT

(10) International Publication Number
WO 2007/137075 A3

(51) International Patent Classification:
C12N 15/82 (2006.01)

(21) International Application Number:

PCT/US2007/069053

(22) International Filing Date: 16 May 2007 (16.05.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/800,872 16 May 2006 (16.05.2006) US

(71) Applicant (for all designated States except US): MON-SANTO TECHNOLOGY LLC [US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): YE, Xudong [US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). SHEN, Junjiang [CN/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US). WILLIAMS, Edward [US/US]; 800 North Lindbergh Blvd., St. Louis, MO 63167 (US).

(74) Agent: HANSON, Robert, E.; Sonnenschein Nath & Rosenthal LLP, P.O. Box 061080, Wacker Drive Station, Chicago, IL 60606 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(88) Date of publication of the international search report:
24 April 2008

(48) Date of publication of this corrected version:
17 July 2008

(15) Information about Correction:
see Notice of 17 July 2008

(54) Title: USE OF NON-AGROBACTERIUM BACTERIAL SPECIES FOR PLANT TRANSFORMATION

(57) Abstract: The invention relates to methods for Rhizobia-mediated genetic transformation of plant cells, including soybean, canola, corn, and cotton cells. These include both VirD2 -dependent and VirD2 -independent methods. Bacterial species utilized include strains of *Rhizobium* sp., *Sinorhizobium* sp., and *Mesorhizobium* sp. Vectors for use in such transformation are also disclosed.



WO 2007/137075 A3

DESCRIPTION

USE OF NON-*AGROBACTERIUM* BACTERIAL SPECIES FOR PLANT TRANSFORMATION

5 BACKGROUND OF THE INVENTION

This application claims the priority of United States Provisional Patent Application 60/800,872, filed May 16, 2006, the disclosure of which is incorporated herein by reference in its entirety.

1. Field of the Invention

10 The present invention relates to the field of plant biotechnology. In particular, the invention relates to methods for producing transgenic plants and plant cells by using non-*Agrobacterium* bacterial species..

2. Description of Related Art

15 *Agrobacterium* spp., members of the Rhizobiales, are common soil bacteria, along with *Rhizobium* spp., *Mesorhizobium* spp., *Sinorhizobium* spp., and related species and genera. A number of wild-type and disarmed (non-pathogenic) strains of *Agrobacterium tumefaciens* and *Agrobacterium rhizogenes* harboring Ti or Ri plasmids can be used for gene transfer into plants. Phytohormone synthesis genes located in the T-DNA of wild type *Agrobacteria* harboring a Ti or Ri plasmid are
20 expressed in plant cells following transformation, and cause tumor formation or a hairy root phenotype depending on the *Agrobacterium* strain or species. Importantly, T-DNA of *Agrobacteria* can be engineered to replace many of its virulence and pathogenicity determinants with “genes of interest” while retaining the ability to be transferred into a plant cell and integrated into a plant genome. Strains containing
25 such “disarmed” Ti plasmids are widely used for plant transformation.

The mechanism of T-DNA transfer to plant cells by *Agrobacterium* has been well documented. Briefly, the T-DNA is delimited by two border regions, referred to as right border (RB) and left border (LB). The borders are nicked by virulence protein VirD2 which produces single stranded transferred DNA (the “T-strand”) with
30 covalent attachment of the VirD2 on its 5’ end. The protein-DNA complex, also

including *Agrobacterium* VirE2 protein, exits *Agrobacterium* cells through the so-called Type 4 secretion system (T4SS, both virulence protein and ssDNA transporter), and is transferred into plant cells and integrated in the plant genome with the help of both *Agrobacterium* virulence proteins and plant factors. The use of *Agrobacterium*-mediated vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described by Fraley *et al.*, (1985), Rogers *et al.*, (1987) and U.S. Patent No. 5,563,055, specifically incorporated herein by reference in its entirety.

Agrobacterium-mediated transformation is efficient in many dicotyledonous plants including *Arabidopsis*, tobacco, and tomato. Methods for *Agrobacterium*-mediated transformation of other species have also been devised (*e.g.* U.S. Patent No. 6,384,301, relating to soybean transformation). While *Agrobacterium*-mediated transformation was at first only used with dicotyledonous plants, advances in *Agrobacterium*-mediated transformation techniques made the technique applicable to monocotyledonous plants as well. For example, *Agrobacterium*-mediated transformation techniques have been applied to rice (Hiei *et al.*, 1997; Zhang *et al.*, 1997; U.S. Patent No. 5,591,616, specifically incorporated herein by reference in its entirety), wheat (McCormac *et al.*, 1998), barley (Tingay *et al.*, 1997; McCormac *et al.*, 1998), and maize (Ishida *et al.*, 1996). However, a number of plant species are recalcitrant to *Agrobacterium*-mediated transformation, and efficiency is low in others. Additionally, since *A. tumefaciens* enters plant tissues at wound sites, and does not naturally infect unwounded tissues, the use of certain tissues as transformation targets is not available.

Besides the T4SS-dependent T-strand delivery system, *Agrobacterium* has additional plasmid mobilization systems that can also transfer and integrate plasmids, such as the IncQ plasmid pRSF1010, between bacterial cells and into the plant genome with lower frequency via conjugal transfer (Buchanan-Wollaston *et al.* 1987, Shadenkov *et al.* 1996; Chen *et al.*, 2002). For example, the conjugal transfer protein MobA, in conjunction with MobB and MobC proteins of the RSF1010 plasmid, cleaves the *oriT* (origin of transfer) site, attaches to the 5' end and transfers the ssDNA into cells independent of the T4SS system (Bravo-Angel *et al.* 1999 and references therein).

Conjugal transfer systems are widely present in bacteria, resulting in exchange of genetic information between bacterial cells. In *Rhizobium*, phylogenetically related but distinct from *Agrobacterium* (Spaink, *et al.*, (ed.), 1998; Farrand *et al.*, 2003), the conjugal transfer system has been partially characterized in some species (Freiberg *et al.*, 1997; Turner *et al.* 2002, Tun-Garrido *et al.* 2003, Perez-Mendoza *et al.* 2004).
5 The conjugal system requires an *oriT* as the nicking site and TraA or Mob as a nicking enzyme, which is different from the conventional elements used in T-DNA mobilization (VirD2 and RB and LB sites, respectively). Unlike VirD2, which was found to have plant NLS (nuclear localization signal) at its C-terminus for plant
10 nuclear targeting, the TraA or Mob doesn't have an obvious NLS. The precise mechanism and site of integration of DNA in plants by TraA remains unclear.

Members of the Rhizobiales other than *Agrobacterium* sp., such as *Rhizobium* spp., are known to symbiotically associate with plant roots in specialized nitrogen-fixing nodules (*e.g.* Long, 2001). In addition to host-specific nodulation of plant
15 roots, especially of legumes, some plant growth promoting effects by members of the Rhizobiales are known in the absence of nodulation (*e.g.* Noel *et al.*, 1996). Recently, reports have been published describing transformation of plants by bacteria other than *Agrobacterium* sp. (*e.g.* Broothaerts *et al.*, 2005; U.S. Patent Application Publications 20050289667; 20050289672; Weller *et al.*, 2004; Weller *et al.*, 2005).

20 Broothaerts *et al.*, reported transformation by *Rhizobium* sp., *Mesorhizobium loti*, and *Sinorhizobium meliloti* strains that was limited to *Arabidopsis*, tobacco, and rice. Weller *et al.* (2004, 2005) reported that several bacteria, including strains of *Rhizobium* sp. and *Ochrobactrum* sp. that harbored Ri plasmids apparently transformed hydroponically grown cucumber and tomato plants, leading to a hairy
25 root phenotype. However the presence of *Agrobacteria* was not ruled out as a possibility in some inoculated plants, complicating the analysis. Transfer of DNA to soybean, corn, cotton, or canola plant cells by non-*Agrobacterium* bacterial strains has not been reported. In addition, transformation efforts in rice, tobacco, and *Arabidopsis* with non-*Agrobacterium* bacterial strains have to date been hampered by
30 low transformation efficiencies. There is, therefore, a great need in the art for the development of improved methods allowing the transformation of important crop

species using non-*Agrobacterium* bacterial strains, and improving transformation efficiencies in general.

BRIEF DESCRIPTION OF THE FIGURES

The following drawings are part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to the drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1: Schematic map of pMON96033.

FIG. 2: Schematic map of pMON96036.

10 **FIG. 3:** Schematic map of pMON101316.

FIG. 4: Transient GUS assay of *Rhizobium*-mediated transformation in soybean with *Mesorhizobium loti* (ML), *Rhizobium leguminosarum* (RL), *Sinorhizobium fredii* (SF), *Sinorhizobium meliloti* (SM) with either disarmed Ti-plasmid (pTiBo542G or pTi4404kan). RL4404: *R. leguminosarum* strain Madison with pTi4404kan; ML542G: *M. loti* USDA3471 with pTiBo542G; ML4404: *M. loti* USDA3471 with pTi4404kan; 2370LBA: *R. leguminosarum* USDA2370 with pTi4404kan; 2370G: *R. leguminosarum* USDA2370 with pTiBo542G; SF4404: *S. fredii* USDA205 with pTi4404kan; SM542C: *S. meliloti* USDA1002 with pTiBo542G; ABI: *A. tumefaciens* ABI strain control.

20 **FIG. 5:** Germline transmission of *gus* transgene in soy produced through *Rhizobium*-mediated transformation.

FIG. 6: Schematic map of pMON96913.

FIG. 7: Schematic map of pMON96914.

FIG. 8: Schematic map of pMON96026.

25 **FIG. 9:** Rhizobia-mediated transformation of canola with several strains as shown by GUS transient assay. A) ML542C (22.4%); B) RL2370G (33.3%); C) RL2370LBA (20%); D) SF542C (30.5%); E) SF4404 (20.6%); and F) SM542C (13%). The % of explants with GUS positive sectors are shown in parentheses.

30 **FIG. 10:** Stable transgenic canola calli transformation with several strains of Rhizobia. A) ML542C (50%); B) RL2370G (21%); C) RL2370LBA (67%); D)

SF542C (36%); and E) SM542C (73%). The % of explants with GUS positive sectors are shown in parentheses.

FIG. 11: Southern blot detection of the CP4 transgene in canola plants derived from *Rhizobium*-mediated transformation. Lane 1: BN_A22 line; lane 2: BN_A24
5 line; lane 3: BN_A28 line; and lane 4: BN_A35 line.

FIG. 12. Cotton transformation by Rhizobia containing pMON101316: A) ML542C (47.8%); B) RL2370G (56%); C) RL2370LBA (31.4%); D) SF542C(23.2%); E) SF4404(31.5%); and F) SM542C (44.4%). RL2370 was used as a negative control; *Agrobacterium tumefaciens* ABI strain was used as a positive
10 control. The percentage of GUS staining positive explants are written in parentheses above.

FIG. 13: Stable transformation of cotton calli by several Rhizobia strains: A) ML542C; B) SF542C; C) SM542C; D) SF4404; E) RL2370LBA; and F) RL2370G.

FIG. 14: Detection of the *gus* transgene by Southern hybridization in cotton
15 calli derived from *Rhizobium*-mediated transformation. RL2370LBA: *R. leguminosarum* 2370 with LBA4404 Ti helper plasmid; SF542: *Sinorhizobium fredii* 205 with pTiBo542 helper plasmid from AGL0 strain; and SF4404: *Sinorhizobium fredii* 205 with LBA4404 Ti helper plasmid.

FIG. 15: Rhizobia-mediated corn transformation as shown by transient
20 expression of a *gus* gene in corn immature embryos. ABI: *A. tumefaciens*; RL2370LBA: *Rhizobium leguminosarum* USDA2370 with LBA4404 Ti plasmid; SM542C: *Sinorhizobium meliloti* USDA1002 with pTiBo542; ML542G: *Mesorhizobium loti* with pTiBo542; SF4404: *Sinorhizobium fredii* USDA205 with LBA4404 Ti plasmid; SF542C: *Sinorhizobium fredii* USDA205 with pTiBo542. All
25 strains contained pMON96036 and were induced in ATA medium pH5.4.

FIG. 16: Corn calli expressing the *gfp* marker after transformation with Rhizobia strains.

FIG. 17: Southern hybridization analysis of transgene integration in corn plants derived from *Rhizobium*-mediated transformation. DIG-labeled *gus* probe was
30 used to detect the transgene. Lane 1-2 and 11-12: lines derived after transformation with *M. loti* ML542G/pMON96036; lane 3-9: lines derived after transformation with

A. tumefaciens ABI control; lane 13-17: lines derived after transformation with *S. fredii* SF4404/pMON96033; Lane 18-19: lines derived after transformation with *S. fredii* SF542C/pMON96036.

SUMMARY OF THE INVENTION

5 In one aspect, the invention provides a method for transforming a plant cell, comprising: (a) contacting at least a first plant cell with a bacterium other than *Agrobacterium* sp. comprising: (i) a first nucleic acid comprising a *vir* gene region of a Ti plasmid wherein the *vir* gene region acts to introduce a nucleic acid coding for a sequence of interest into the plant cell in a VirD2-dependent manner; and (ii) a second
10 nucleic acid comprising one or more T-DNA border sequence(s) operably linked to a nucleic acid of interest; and (b) selecting at least a first plant cell transformed with the nucleic acid of interest, wherein the plant cell is a soybean, canola, corn, or cotton plant cell.

 In another aspect, the invention provides a method for transforming a plant
15 cell, comprising: (a) contacting at least a first plant cell with a bacterium other than *Agrobacterium* comprising (i) a first nucleic acid required for conjugative transfer of DNA sequences independent of VirD2 function, and (ii) a second nucleic acid comprising a nucleic acid of interest; wherein the plant cell is a soybean, canola, corn, or cotton plant cell and wherein polypeptides encoded by the nucleic acid required for
20 conjugative transfer act to transfer the nucleic acid of interest into the plant cell; and (b) selecting at least a first plant cell transformed with the nucleic acid of interest. In such a method, the conjugative transfer may be *traA*, *traI*, or *mobA*-dependent, and the first nucleic acid comprises *oriT*. The first nucleic acid may lack left and right T-DNA border sequences.

25 In a method of the invention, the bacterium may be Rhizobia cell. In certain embodiments, the Rhizobia is grown under suitable conditions to minimize polysaccharide production by the Rhizobia cells. The Rhizobia cell may be grown in the presence of acetosyringone or other compound, such as a phenolic compound, that induces *vir* gene function prior to contacting the plant cell. The Rhizobia cell may be
30 selected from the group consisting of: *Rhizobium* spp., *Sinorhizobium* spp., *Mesorhizobium* spp., *Phyllobacterium* spp. *Ochrobactrum* spp. and *Bradyrhizobium* spp. In specific embodiments, the Rhizobia cell is a *Rhizobium leguminosarum* cell

and may further be a cell of *R. leguminosarum* *bv. trifolii*, *R. leguminosarum* *bv. phaseoli* or *Rhizobium leguminosarum. bv. viciae*.

In another aspect of a transformation method provided by the invention, a plant cell that is transformed may be comprised in an explant from a plant seed, for example, from a seedling, callus, cell suspension, cotyledon, meristem, leaf, root, or stem. The explant may comprise an embryonic meristem explant; callus; cell suspension; cotyledon; or tissue from leaves, roots, or stems.

A bacterium used for transformation in accordance with the invention may comprise nucleic acids introduced, for example, by electroporation. The sequences may comprise nucleic acid required for conjugative transfer independent of VirD2 function. The nucleic acids may include first and second nucleic acids.

In another aspect of the invention, a transformation method provided herein may comprise selecting a plant cell transformed with a nucleic acid of interest in the absence of a selection agent. Selecting a plant cell transformed with a nucleic acid of interest may comprise culturing the plant cell in the presence of a selection agent, wherein the nucleic acid of interest confers tolerance to the selection agent or is operably linked to a further nucleic acid that confers tolerance to the selection agent. Examples of such selection agents include glyphosate, kanamycin, bialaphos or dicamba. In one embodiment, the nucleic acid of interest or further nucleic acid encodes EPSP synthase and in a still further embodiment encodes the EPSP synthase protein CP4. In another embodiment, the selection agent is glyphosate. In yet other embodiments, the sequence of interest may be defined as not physically linked to a selectable marker gene. For example, the marker gene and nucleic acid of interest may genetically segregate in progeny of a plant regenerated from the plant cell transformed with the nucleic acid of interest.

A bacterium in accordance with the invention may comprise at least a third nucleic acid comprising a further nucleic acid of interest, wherein the plant cell is transformed with the third nucleic acid. In a method of the invention, a plant may be regenerated a transgenic plant cell, wherein the plant comprises the sequence of interest. Regenerating a plant may comprise inducing formation of one or more shoots from an explant comprising the plant cell and cultivating at least a first shoot into a whole fertile plant. In certain embodiments, the plant may be a corn or cotton

plant. In further embodiments, regeneration occurs by organogenesis. In other embodiments, the plant is a soybean or canola plant.

In another aspect, the invention provides a Rhizobia cell selected from the group consisting of: *Rhizobium* spp., *Sinorhizobium* spp., *Mesorhizobium* spp.,
5 *Phyllobacterium* spp. *Ochrobactrum* spp. and *Bradyrhizobium* spp., the cell comprising (i) a first nucleic acid comprising a *vir* gene region of a Ti plasmid wherein the *vir* gene region acts to introduce a nucleic acid coding for a sequence of interest into a plant cell in a VirD2-dependent manner; and (ii) a second nucleic acid comprising one or more T-DNA border sequence(s) operably linked to a nucleic acid
10 coding for a sequence of interest. In one embodiment, the cell is further defined as comprising a selectable marker. In another embodiment, the Rhizobia cell is selected from the group consisting of: *Rhizobium* sp., *Rhizobium* sp. NGR234, *Rhizobium leguminosarum* Madison, *R. leguminosarum* USDA2370, *R. leguminosarum* USDA2408, *R. leguminosarum* USDA2668, *R. leguminosarum* 2370G, *R.*
15 *leguminosarum* 2370LBA, *R. leguminosarum* 2048G, *R. leguminosarum* 2048LBA, *R. leguminosarum* bv. *phaseoli*, *R. leguminosarum* bv. *phaseoli* 2668G, *R. leguminosarum* bv. *phaseoli* 2668LBA, *R. leguminosarum* RL542C, *R. leguminosarum* bv. *viciae*, *R. leguminosarum* bv. *trifolii*, *Rhizobium etli* USDA 9032, *R. etli* bv. *phaseoli*, *Rhizobium tropici*, *Mesorhizobium* sp., *Mesorhizobium loti*
20 ML542G, *M. loti* ML4404, *Sinorhizobium* sp., *Sinorhizobium meliloti* SD630, *S. meliloti* USDA1002, *Sinorhizobium fredii* USDA205, *S. fredii* SF542G, *S. fredii* SF4404, *S. fredii* SM542C, *Bradyrhizobium* sp., *Bradyrhizobium japonicum* USDA 6, and *B. japonicum* USDA 110. In specific embodiments, the cell is a *Rhizobium leguminosarum* cell and may further be. For example, a *R. leguminosarum* bv. *trifolii*,
25 *R. leguminosarum* bv. *phaseoli* or *Rhizobium leguminosarum* bv. *viciae* cell.

In yet another aspect of the invention, a DNA construct is provided competent for virD2-independent transfer from Rhizobia and lacking T-DNA border sequence, the construct comprising an *oriT* sequence and *traA* or *mob* sequence operably linked to a nucleic acid of interest. The invention further provides a Rhizobia cell
30 transformed with such a DNA construct of claim 35, wherein the Rhizobia is selected from the group consisting of: *Rhizobium* spp., *Sinorhizobium* spp., *Mesorhizobium* spp., *Phyllobacterium* spp. *Ochrobactrum* spp. and *Bradyrhizobium* spp. In one

embodiment, the Rhizobia cell is selected from the group consisting of: *Rhizobium* sp., *Rhizobium* sp. NGR234, *Rhizobium leguminosarum* Madison, *R. leguminosarum* USDA2370, *R. leguminosarum* USDA2408, *R. leguminosarum* USDA2668, *R. leguminosarum* 2370G, *R. leguminosarum* 2370LBA, *R. leguminosarum* 2048G, *R. leguminosarum* 2048LBA, *R. leguminosarum* bv. *phaseoli*, *R. leguminosarum* bv. *phaseoli* 2668G, *R. leguminosarum* bv. *phaseoli* 2668LBA, *R. leguminosarum* RL542C, *R. leguminosarum* bv. *viciae*, *R. leguminosarum* bv. *trifolii*, *Rhizobium etli* USDA 9032, *R. etli* bv. *phaseoli*, *Rhizobium tropici*, *Mesorhizobium* sp., *Mesorhizobium loti* ML542G, *M. loti* ML4404, *Sinorhizobium* sp., *Sinorhizobium meliloti* SD630, *S. meliloti* USDA1002, *Sinorhizobium fredii* USDA205, *S. fredii* SF542G, *S. fredii* SF4404, *S. fredii* SM542C, *Bradyrhizobium* sp., *Bradyrhizobium japonicum* USDA 6, and *B. japonicum* USDA 110. In specific embodiments, the cell is a *Rhizobium leguminosarum* cell, and in still further embodiments, may be a *R. leguminosarum* bv. *trifolii*, *R. leguminosarum* bv. *phaseoli* or *Rhizobium leguminosarum*. bv. *viciae* cell.

DETAILED DESCRIPTION OF THE INVENTION

The following is a detailed description of the invention provided to aid those skilled in the art in practicing the present invention. Those of ordinary skill in the art may make modifications and variations in the embodiments described herein without departing from the spirit or scope of the present invention.

The present invention provides methods and compositions for the efficient genetic transformation of plant cells of important crop species by Rhizobia. The invention overcomes substantial limitations in the art, including limited transformation efficiency and failure to describe techniques amenable to transformation of important crop plants by use of non-*Agrobacterium* strains. For example, while use of bacteria other than *Agrobacterium* has been discussed for several plant varieties, transformation frequencies have been low. In the case of rice, transformation frequencies of 0.6% and lower have been reported, with only one transformed plant obtained from 687 inoculated calli (Broothaerts *et al.*, 2005). This contrasts to 50-80% transformation frequencies using *Agrobacterium*. Even using model organisms easily transformed by *Agrobacterium*, transformation frequencies were only a fraction of those obtained by *Agrobacterium*-mediated transformation.

To date considerable research had been required in many instances to apply even well developed transformation procedures such as *Agrobacterium*-mediated transformation to different plant species. Plants of different species often exhibit substantial physiological differences that effect amenability to genetic transformation.

5 Methods for transformation of one species of plant therefore often do not work effectively, if at all, with other plants and the ability to transform a plant is not necessarily predictive of the ability to transform even related species using that procedure. This is particularly true for bacterial transformation, which involves complex biochemical interactions between the bacterial strains used and target plant

10 cells. Rhizobia interact with plants in the native environment and therefore can exhibit host-specificities, the impact of which is unknown for many crop species.

Thus, identifying plants amenable to Rhizobia-mediated transformation, and developing procedures allowing increased transformation efficiencies is of great interest. Efficient transformation in particular is important because the extent to

15 which any given transformation event is expressed can vary substantially depending upon the integration site in the plant genome. The ability to select transformation events having a suitable expression profile is thus dependent upon the ability to efficiently produce transformants. As explained in the working examples below, transient transformation frequencies of as high as 5% were obtained by the inventors

20 for the transformation of soybeans (FIG. 4) and frequencies approaching 56% and 33% were obtained in the case of cotton (FIG. 12) and canola (FIG. 9), respectively.

The present invention overcomes limitations in the art by providing, in one embodiment, techniques for the use of Rhizobia to transform important crop plants that were not previously known to be transformable by Rhizobia, including canola,

25 corn, cotton, and soybean. The invention also provides techniques for the efficient transformation of plants using Rhizobia, including those already known to be amenable to transformation by Rhizobia at a low frequency. The invention also provides methods for the transformation of tissue targets differing from those of *Agrobacterium*. For example, while *Agrobacterium* typically requires a wound site to

30 infect plants, some other members of the Rhizobiales, including *Rhizobiaceae* such as *Rhizobium*, naturally infect plant roots via infection threads that penetrate plant tissues, allowing for use of non-wounded tissue as a transformation target.

The following definitions are provided in order to aid those skilled in the art in understanding the detailed description of the present invention.

As used herein, “plant growth regulator” or “plant hormone” refers to compounds that affect plant growth. Plant growth regulators include, but are not limited to, auxins, cytokinins, ABA, gibberellins, ethylene, brassinosteroids, and polyamines. Auxins affect the elongation of shoots and roots at low concentration but inhibit growth at higher levels. Commonly used auxins include picloram (4-amino-3,5,6-trichloropicolinic acid), 2,4-D (2,4-dichlorophenoxyacetic acid), IAA (indole-3-acetic acid), NAA (α -naphthaleneacetic acid), and dicamba (3,6-dichloroanistic acid). Cytokinins cause cell division, cell differentiation, and shoot differentiation. Commonly used cytokinins include kinetin, BA (6-benzylaminopurine), 2-ip (2-isopentenyladenine), BAP (6-benzylaminopurine), thidiazuron (TDZ), zeatin riboside, and zeatin.

“Coding sequence”, “coding region” or “open reading frame” refers to a region of continuous sequential nucleic acid triplets encoding a protein, polypeptide, or peptide sequence.

“Dicot” or “dicotyledonous” refers to plants having two cotyledons. Examples include, without limitation, plants such as alfalfa, beans, broccoli, cabbage, canola, carrot, cauliflower, celery, cotton, cucumber, eggplant, lettuce, melon, pea, pepper, potato, pumpkin, radish, rapeseed, spinach, soybean, squash, tomato, and watermelon.

“Endogenous” refers to materials originating from within the organism or cell.

“Exogenous” refers to materials originating from outside of the organism or cell. As used herein, exogenous is intended to refer to any nucleic acid from a source other than the recipient cell or tissue, regardless of whether a similar (but not identical) nucleic acid may already be present in the recipient cell or tissue.

“Explant” refers to a plant part that is capable of being transformed and subsequently regenerated into a transgenic plant. Examples include embryos, callus, cell suspensions, cotyledons, meristems, seedlings, seeds, leaves, stems or roots.

“Monocot” or “monocotyledonous” refers to plants having a single cotyledon. Examples include, without limitation, onions, corn, rice, sorghum, wheat, rye, millet, sugarcane, oat, triticale, barley and turfgrass.

“Nucleic acid” refers to deoxyribonucleic acid (DNA) or ribonucleic acid
5 (RNA).

“Phenotype” refers to a trait exhibited by an organism resulting from the expression (or lack of expression) of nucleic acids in the genome (including non-genomic DNA and RNA such as plasmids and artificial chromosomes) and/or organelles of the organism.

10 The term “plant” encompasses any higher plant and progeny thereof, including monocots (*e.g.*, corn, rice, wheat, barley, *etc.*), dicots (*e.g.*, soybean, cotton, tomato, potato, *Arabidopsis*, tobacco, *etc.*), gymnosperms (pines, firs, cedars, *etc.*) and includes parts of plants, including reproductive units of a plant (*e.g.*, seeds, bulbs, tubers, meristematic tissues, or other parts or tissues from that the plant can be
15 reproduced), fruits and flowers.

“Polyadenylation signal” or “polyA signal” refers to a nucleic acid sequence located 3' to a coding region that promotes the addition of adenylate nucleotides to the 3' end of an mRNA transcribed from the coding region.

“Promoter” or “promoter region” refers to a nucleic acid sequence, usually
20 found 5' to a coding sequence, that alters expression of the coding sequence by providing a recognition site for RNA polymerase and/or other recognition sites for other transcription-related factors utilized to produce RNA and/or initiate transcription at the correct site on the DNA.

“Recombinant nucleic acid vector” or “vector” or “construct” refers to any
25 agent such as a plasmid, cosmid, virus, autonomously replicating sequence, phage, or linear or circular single- or double-stranded DNA or RNA nucleotide segment, derived from any source, capable of genomic integration or autonomous replication, comprising a nucleic acid molecule in which one or more nucleic acid sequences have been linked in a functionally operative manner. Such recombinant nucleic acid
30 vectors or constructs typically comprise a 5' regulatory sequence or promoter region and a coding sequence encoding for a desired gene product. The vectors are typically

designed such that once delivered into a cell or tissue, the coding sequence is transcribed into mRNA, which is optionally translated into a polypeptide or protein.

“Regeneration” refers to the process of growing a plant from a plant cell or tissue.

5 “Rhizobia” refers without limitation to bacterial genera, species, and strains that may be assigned to the order Rhizobiales other than *Agrobacterium* bacterial strains comprising the taxonomic families Rhizobiaceae (e.g. *Rhizobium* spp., *Sinorhizobium* spp.); Phyllobacteriaceae (e.g. *Mesorhizobium* spp., *Phyllobacterium* spp.); Brucellaceae (e.g. *Ochrobactrum* spp.); Bradyrhizobiaceae (e.g.
10 *Bradyrhizobium* spp.), and Xanthobacteraceae (e.g. *Azorhizobium* spp.), among others. For the purposes of the present application, “Rhizobia” does not include, biovars, or species.

Taxonomic assignment may be done as is known in the art, for instance by comparison of 16S rDNA sequences or other classification methods. Wild type
15 strains of many *Rhizobium* species are typically able to induce formation of nitrogen fixing nodules in root tissues of host plants such as leguminous plants (Fabaceae). However, the ability to nodulate roots of a given plant species is not required for *Rhizobium*-mediated DNA transfer into cells of the plant species.

“Selectable marker” or “screenable marker” refers to a nucleic acid sequence
20 whose expression confers a phenotype facilitating identification of cells, tissues, or plants containing the nucleic acid sequence.

“Transcription” refers to the process of producing an RNA copy from a DNA template.

“Transformation” refers to a process of introducing an exogenous nucleic acid
25 sequence into a cell or tissue. The transformation may be transient or stable. In stable transformations, part or all of the exogenous nucleic acid is incorporated (e.g., integrated or stably maintained) in the nuclear genomic DNA, plastid DNA, or is capable of autonomous replication in the nucleus or plastid.

“Transgenic” refers to organisms into which an exogenous nucleic acid
30 sequence has been stably transformed.

In designing a vector for the transformation process, one or more genetic components are selected that will be introduced into the plant cell or tissue. Genetic components can include any nucleic acid that is introduced into a plant cell or tissue using the method according to the invention. Genetic components can include non-plant DNA, plant DNA or synthetic DNA.

In one embodiment, the genetic components are incorporated into a DNA composition such as a recombinant, double-stranded plasmid or vector molecule comprising at least one or more of following types of genetic components: (a) a promoter that functions in plant cells to cause the production of an RNA sequence, (b) a structural DNA sequence that causes the production of an RNA sequence that encodes a product of agronomic utility, and (c) a 3' non-translated DNA sequence that functions in plant cells to cause the addition of polyadenylated nucleotides to the 3' end of the RNA sequence.

The vector may contain a number of genetic components to facilitate transformation of the plant cell or tissue and to regulate expression of the structural nucleic acid sequence. In one preferred embodiment, the genetic components are oriented so as to express a mRNA, that in an optional embodiment can be translated into a protein. The expression of a plant structural coding sequence (a gene, cDNA, synthetic DNA, or other DNA) that exists in double-stranded form involves transcription of messenger RNA (mRNA) from one strand of the DNA by RNA polymerase enzyme and subsequent processing of the mRNA primary transcript inside the nucleus. This processing involves a 3' non-translated region that adds polyadenylated nucleotides to the 3' ends of the mRNA.

Means for preparing plasmids or vectors containing the desired genetic components are well known in the art. Vectors typically consist of a number of genetic components, including but not limited to regulatory elements such as promoters, leaders, introns, and terminator sequences. Regulatory elements are also referred to as cis- or trans-regulatory elements, depending on the proximity of the element to the sequences or gene(s) they control.

Transcription of DNA into mRNA is regulated by a region of DNA usually referred to as the "promoter". The promoter region contains a sequence of bases that signals RNA polymerase to associate with the DNA and to initiate the transcription

into mRNA using one of the DNA strands as a template to make a corresponding complementary strand of RNA.

A number of promoters that are active in plant cells have been described in the literature. Such promoters would include but are not limited to the nopaline synthase (NOS) and octopine synthase (OCS) promoters that are carried on Ti plasmids of *Agrobacterium tumefaciens*, the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S and 35S promoters and the Figwort mosaic virus (FMV) 35S promoter, and the enhanced CaMV35S promoter (e35S). A variety of other plant gene promoters that are regulated in response to environmental, hormonal, chemical, and/or developmental signals, also can be used for expression of any DNA construct in plant cells, including, for instance, promoters regulated by (1) heat (Callis *et al.*, 1988, (2) light (*e.g.*, pea RbcS-3A promoter, Kuhlemeier *et al.*, (1989); maize RbcS promoter, Schaffner *et al.*, (1991); (3) hormones, such as abscisic acid (Marcotte *et al.*, 1989, (4) wounding (*e.g.*, Wuni, Siebertz *et al.*, 1989); or other signals or chemicals. Tissue specific expression is also known. As described below, it is preferred that the particular promoter selected should be capable of causing sufficient expression to result in the production of an effective amount of the gene product of interest. Examples describing such promoters include without limitation U.S. Patent 6,437,217 (maize RS81 promoter), U.S. Patent 5,641,876 (rice actin promoter, OsAct1), U.S. Patent 6,426,446 (maize RS324 promoter), U.S. Patent 6,429,362 (maize PR-1 promoter), U.S. Patent 6,232,526 (maize A3 promoter), U.S. Patent 6,177,611 (constitutive maize promoters), U.S. Patents 5,322,938, 5,352,605, 5,359,142 and 5,530,196 (35S promoter), U.S. Patent 6,433,252 (maize L3 oleosin promoter), U.S. Patent 6,429,357 (rice actin 2 promoter as well as a rice actin 2 intron), U.S. Patent 5,837,848 (root specific promoter), U.S. Patent 6,294,714 (light inducible promoters), U.S. Patent 6,140,078 (salt inducible promoters), U.S. Patent 6,252,138 (pathogen inducible promoters), U.S. Patent 6,175,060 (phosphorus deficiency inducible promoters), U.S. Patent 6,635,806 (gamma-coixin promoter), and U.S. Patent 7,151,204 (maize chloroplast aldolase promoter). Additional promoters that may find use are a nopaline synthase (NOS) promoter (Ebert *et al.*, 1987), the octopine synthase (OCS) promoter (which is carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*), the caulimovirus promoters such as the cauliflower mosaic virus (CaMV) 19S promoter (Lawton *et al.*, 1987), the CaMV 35S promoter

(Odell *et al.*, 1985), the figwort mosaic virus 35S-promoter (Walker *et al.*, 1987; U.S. Patents 6,051,753; 5,378,619), the sucrose synthase promoter (Yang *et al.*, 1990), the R gene complex promoter (Chandler *et al.*, 1989), and the chlorophyll a/b binding protein gene promoter, PCISV (US Patent 5,850,019). In the present invention,
5 CaMV35S with enhancer sequences (e35S; U.S. Patent Nos. 5,322,938; 5,352,605; 5,359,142; and 5,530,196), FMV35S (U.S. Patents 6,051,753; 5,378,619), peanut chlorotic streak caulimovirus (PCISV; U.S. Patent 5,850,019), At.Act 7 (Accession # U27811), At.ANT1 (U.S. Patent Application 20060236420), FMV.35S-EF1a (U.S. Patent Application Publication 2005/0022261), eIF4A10 (Accession # X79008) and
10 AGRtu.nos (GenBank Accession V00087; Depicker *et al.*, 1982; Bevan *et al.*, 1983), rice cytosolic triose phosphate isomerase (OsTPI; U.S. Patent No. 7,132,528), and rice actin 15 gene (OsAct15; U.S. Patent Application Publication 2006/0162010) promoters may be of particular benefit. In some instances, *e.g.* OsTPI and OsAct 15, a promoter may include a 5'UTR and/or a first intron.

15 Promoter hybrids can also be constructed to enhance transcriptional activity (U.S. Patent No. 5,106,739), or to combine desired transcriptional activity, inducibility and tissue specificity or developmental specificity. Promoters that function in plants include but are not limited to promoters that are inducible, viral, synthetic, constitutive as described, and temporally regulated, spatially regulated, and
20 spatio-temporally regulated. Other promoters that are tissue-enhanced, tissue-specific, or developmentally regulated are also known in the art and envisioned to have utility in the practice of this invention.

The promoters used in the DNA constructs (*i.e.* chimeric/recombinant plant genes) of the present invention may be modified, if desired, to affect their control
25 characteristics. Promoters can be derived by means of ligation with operator regions, random or controlled mutagenesis, etc. Furthermore, the promoters may be altered to contain multiple "enhancer sequences" to assist in elevating gene expression.

The mRNA produced by a DNA construct of the present invention may also contain a 5' non-translated leader sequence. This sequence can be derived from the
30 promoter selected to express the gene and can be specifically modified so as to increase or decrease translation of the mRNA. The 5' non-translated regions can also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic

gene sequence. Such “enhancer” sequences may be desirable to increase or alter the translational efficiency of the resultant mRNA. The present invention is not limited to constructs wherein the non-translated region is derived from both the 5' non-translated sequence that accompanies the promoter sequence. Rather, the non-translated leader
5 sequence can be derived from unrelated promoters or genes (see, for example U.S. Patent No. 5,362,865). Examples of non-translation leader sequences include maize and petunia heat shock protein leaders (U.S. Patent No. 5,362,865), plant virus coat protein leaders, plant rubisco leaders, GmHsp (U.S. Patent 5,659,122), PhDnaK (U.S. Patent No. 5,362,865), AtAnt1, TEV (Carrington and Freed, 1990), OsAct1 (U.S.
10 Patent No. 5,641,876), OsTPI (U.S. Patent No. 7,132,528), and OsAct15 (U.S. Publication No. 20060162010), and AGRtu.nos (GenBank Accession V00087; Bevan *et al.*, 1983). Other genetic components that serve to enhance expression or affect transcription or translational of a gene are also envisioned as genetic components.

Intron sequences are known in the art to aid in the expression of transgenes in
15 monocot plant cells. Examples of introns include the corn actin intron (U.S. Patent 5,641,876), the corn HSP70 intron (ZmHSP70; U.S. Patent 5,859,347; U.S. Patent 5,424,412), and rice TPI intron (OsTPI; U.S. Patent No. 7,132,528) and are of benefit in practicing this invention.

Termination of transcription may be accomplished by a 3' non-translated DNA
20 sequence operably linked to a recombinant transgene (*e.g.* the gene of interest, the identification sequence comprising a screenable gene, or the plant selectable marker gene). The 3' non-translated region of a recombinant DNA molecule contains a polyadenylation signal that functions in plants to cause the addition of adenylate nucleotides to the 3' end of the RNA. The 3' non-translated region can be obtained
25 from various genes that are expressed in plant cells. The nopaline synthase 3' untranslated region (Fraley *et al.*, 1983), is commonly used in this capacity. Polyadenylation molecules from a *Pisum sativum* RbcS2 gene (Ps.RbcS2-E9; Coruzzi *et al.*, 1984), AGRtu.nos (Genbank Accession E01312), E6 (Accession # U30508), rice glutelin (Okita *et al.*, 1989), and TaHsp17 (wheat low molecular weight heat
30 shock protein gene; GenBank Accession # X13431) in particular may be of benefit for use with the invention.

In one embodiment, the vector contains a selectable, screenable, or scoreable marker gene. These genetic components are also referred to herein as functional genetic components, as they produce a product that serves a function in the identification of a transformed plant, or a product of agronomic utility. The DNA that serves as a selection or screening device may function in a regenerable plant tissue to produce a compound that would confer upon the plant tissue resistance to an otherwise toxic compound. A number of screenable or selectable marker genes are known in the art and can be used in the present invention. Examples of selectable markers and genes providing resistance against them are disclosed in Miki and McHugh, 2004. Genes of interest for use as a selectable, screenable, or scoreable marker would include but are not limited to *gus*, *gfp* (green fluorescent protein), luciferase (LUX), genes conferring tolerance to antibiotics like kanamycin (Dekeyser *et al.*, 1989), neomycin, kanamycin, paromomycin, G418, aminoglycosides, spectinomycin, streptomycin, hygromycin B, bleomycin, phleomycin, sulfonamides, streptothricin, chloramphenicol, methotrexate, 2-deoxyglucose, betaine aldehyde, S-aminoethyl L-cysteine, 4-methyltryptophan, D-xylose, D-mannose, benzyladenine-N-3-glucuronidase, genes that encode enzymes that give tolerance to herbicides like glyphosate (*e.g.* 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS): Della-Cioppa *et al.*, 1987; U.S. Patent 5,627,061; U.S. Patent 5,633,435; U.S. Patent 6,040,497; U.S. Patent 5,094,945; WO04074443, and WO04009761; glyphosate oxidoreductase (GOX; U.S. Patent 5,463,175); glyphosate decarboxylase (WO05003362 and US Patent Application 20040177399; or glyphosate N-acetyltransferase (GAT): Castle *et al.*, U.S. Patent Publication 20030083480), dalapon (*e.g.* *dehI* encoding 2,2-dichloropropionic acid dehalogenase conferring tolerance to 2,2-dichloropropionic acid (Dalapon; WO9927116)), bromoxynil (haloarylnitrilase (Bxn) for conferring tolerance to bromoxynil (WO8704181A1; US 4,810,648; WO8900193A)), sulfonyl herbicides (*e.g.* acetohydroxyacid synthase or acetolactate synthase conferring tolerance to acetolactate synthase inhibitors such as sulfonylurea, imidazolinone, triazolopyrimidine, pyrimidyloxybenzoates and phthalide; (US 6,225,105; US 5,767,366, U.S. 4,761,373; U.S. 5,633,437; U.S. 6,613,963; US 5,013,659; US 5,141,870; US 5,378,824; US 5,605,011)); encoding ALS, GST-II), bialaphos or phosphinothricin or derivatives (*e.g.* phosphinothricin acetyltransferase (*bar*) conferring tolerance to phosphinothricin or glufosinate (US 5,646,024, US

5,561,236, EP 275,957; US 5,276,268; US 5,637, 489; US 5,273, 894), atrazine (encoding GST-III), dicamba (dicamba monooxygenase (DMO); US Patent Applications 20030115626, 20030135879), or sethoxydim (modified acetyl-coenzyme A carboxylase for conferring tolerance to cyclohexanedione (sethoxydim) and aryloxyphenoxypropionate (haloxyfop) (U.S. 6,414,222)), among others. Other selection procedures can also be implemented including positive selection mechanisms (e.g. use of the *manA* gene of *E. coli*, allowing growth in the presence of mannose) and would still fall within the scope of the present invention (see also Miki and McHugh (2004)).

10 The present invention can be used with any suitable plant transformation plasmid or vector containing a selectable or screenable marker and associated regulatory elements as described, along with one or more nucleic acids expressed in a manner sufficient to confer a particular desirable trait. Examples of suitable structural genes of agronomic interest envisioned by the present invention would include but are not limited to genes for disease, insect, or pest tolerance, herbicide tolerance, genes for quality improvements such as yield, nutritional enhancements, environmental or stress tolerances, or any desirable changes in plant physiology, growth, development, morphology or plant product(s) including starch production (U.S. Patents 6,538,181; 15 6,538,179; 6,538,178; 5,750,876; 6,476,295), modified oils production (U.S. Patents 6,444,876; 6,426,447; 6,380,462), high oil production (U.S. Patents 6,495,739; 5,608,149; 6,483,008; 6,476,295), modified fatty acid content (U.S. Patents 6,828,475; 6,822,141; 6,770,465; 6,706,950; 6,660,849; 6,596,538; 6,589,767; 6,537,750; 6,489,461; 6,459,018), high protein production (U.S. Patent 6,380,466), fruit ripening (U.S. Patent 5,512,466), enhanced animal and human nutrition (U.S. 25 Patents 6,723,837; 6,653,530; 6,5412,59; 5,985,605; 6,171,640), biopolymers (U.S. Patents RE37,543; 6,228,623; 5,958,745 and U.S. Patent Publication No. US20030028917). Also environmental stress resistance (U.S. Patent 6,072,103), pharmaceutical peptides and secretable peptides (U.S. Patents 6,812,379; 6,774,283; 6,140,075; 6,080,560), improved processing traits (U.S. Patent 6,476,295), improved 30 digestibility (U.S. Patent 6,531,648) low raffinose (U.S. Patent 6,166,292), industrial enzyme production (U.S. Patent 5,543,576), improved flavor (U.S. Patent 6,011,199), nitrogen fixation (U.S. Patent 5,229,114), hybrid seed production (U.S. Patent 5,689,041), fiber production (U.S. Patent 6,576,818; 6,271,443; 5,981,834; 5,869,720)

and biofuel production (U.S. Patent 5,998,700). Any of these or other genetic elements, methods, and transgenes may be used with the invention as will be appreciated by those of skill in the art in view of the instant disclosure.

Alternatively, the DNA sequences of interest can affect these phenotypes by
5 the inhibition of expression of an endogenous gene via gene silencing technologies such as cosuppression, antisense, RNAi, expression of miRNAs (natural or engineered), expression of trans-acting siRNAs, and expression of ribozymes (see *e.g.*, U.S. Patent Application Publication 20060200878).

Exemplary nucleic acids that may be introduced by the methods encompassed
10 by the present invention include, for example, DNA sequences or genes from another species, or even genes or sequences that originate with or are present in the same species, but are incorporated into recipient cells by genetic engineering methods rather than classical reproduction or breeding techniques. However, the term “exogenous” is also intended to refer to genes that are not normally present in the cell
15 being transformed, or perhaps simply not present in the form, structure, etc., as found in the transforming DNA segment or gene, or genes that are normally present yet that one desires, *e.g.*, to have over-expressed. Thus, the term “exogenous” gene or DNA is intended to refer to any gene or DNA segment that is introduced into a recipient cell, regardless of whether a similar gene may already be present in such a cell. The
20 type of DNA included in the exogenous DNA can include DNA that is already present in the plant cell, DNA from another plant, DNA from a different organism, or a DNA generated externally, such as a DNA sequence containing an antisense message of a gene, or a DNA sequence encoding a synthetic or modified version of a gene.

In light of this disclosure, numerous other possible selectable or screenable
25 marker genes, regulatory elements, and other sequences of interest will be apparent to those of skill in the art. Therefore, the foregoing discussion is intended to be exemplary rather than exhaustive.

For Rhizobia-mediated transformation, after the construction of the plant transformation vector or construct, the nucleic acid molecule, prepared as a DNA
30 composition *in vitro*, is introduced into a suitable host such as *E. coli* and mated into another suitable host such as Rhizobia, including *Rhizobium*, or directly transformed (*e.g.* electroporated) into competent Rhizobia. The Ti or Ri plasmid may be naturally

transferred into nitrogen-fixing *Rhizobium* and may induce tumors or hairy roots, respectively (Hooykaas *et al.* 1977, Weller *et al.* 2004). Such Ti or Ri plasmid may alternatively be “disarmed”, and unable to cause plant cell proliferation. Since *Rhizobium* and *Agrobacterium* have differing infection mechanisms, deep infection by
5 *Rhizobium* or other Rhizobia through its infection thread may increase the frequency of germ line transformation of a gene of interest during soybean transformation once the Ti or Ri helper plasmid is introduced.

The present invention encompasses the use of bacterial strains to introduce one or more genetic components into plants. In one embodiment, the hosts contain
10 disarmed Ti or Ri plasmids that do not contain the oncogenes that cause tumorigenesis or rhizogenesis, derivatives of which are used as the vectors and contain the genes of interest that are subsequently introduced into plants. In another embodiment, the bacteria transfer DNA into plant cells by means of a T4SS-independent mechanism, namely *oriT*-mediated conjugal transfer. Functions needed
15 for T4SS-independent DNA transfer may reside on the plasmid containing the DNA to be transferred, or may reside on the chromosome or another plasmid, including a Ti or Ri plasmid, also present in such a bacterial cell.

Bacterial species and strains include but are not limited to *Rhizobium* sp., *Rhizobium* sp. NGR234, *Rhizobium leguminosarum* Madison, *R. leguminosarum*
20 USDA2370, *R. leguminosarum* USDA2408, *R. leguminosarum* USDA2668, *R. leguminosarum* 2370G, *R. leguminosarum* 2370LBA, *R. leguminosarum* 2048G, *R. leguminosarum* 2048LBA, *R. leguminosarum* bv. *phaseoli*, *R. leguminosarum* bv. *phaseoli* 2668G, *R. leguminosarum* bv. *phaseoli* 2668LBA, *R. leguminosarum* RL542C, *R. leguminosarum* bv. *viciae*, *R. leguminosarum* bv. *trifolii*, *Rhizobium etli*
25 USDA 9032, *R. etli* bv *phaseoli*, *Rhizobium tropici*, *Mesorhizobium* sp., *Mesorhizobium loti* ML542G, *M. loti* ML4404, *Sinorhizobium* sp., *Sinorhizobium meliloti* SD630, *S. meliloti* USDA1002, *Sinorhizobium fredii* USDA205, *S. fredii* SF542G, *S. fredii* SF4404, *S. fredii* SM542C, *Bradyrhizobium* sp., *Bradyrhizobium japonicum* USDA 6, *B. japonicum* USDA 110.

30 Any suitable plant culture medium can be used to develop or maintain a plant tissue culture, supplemented as appropriate with additional plant growth regulators including but not limited to auxins such as picloram (4-amino-3,5,6-trichloropicolinic

acid), 2,4-D (2,4-dichlorophenoxyacetic acid) and dicamba (3,6-dichloroanistic acid); cytokinins such as BAP (6-benzylaminopurine) and kinetin; ABA; and gibberellins. Other media additives can include but are not limited to amino acids, macro elements, iron, microelements, inositol, vitamins and organics, carbohydrates, undefined media components such as casein hydrolysates, with or without an appropriate gelling agent such as a form of agar, such as a low melting point agarose or Gelrite if desired. Those of skill in the art are familiar with the variety of tissue culture media, which when supplemented appropriately, support plant tissue growth and development and are suitable for plant transformation and regeneration. These tissue culture media can either be purchased as a commercial preparation, or custom prepared and modified. Examples of such media would include but are not limited to Murashige and Skoog (1962), N6 (Chu *et al.*, 1975), Linsmaier and Skoog (1965), Uchimiya and Murashige (1962), Gamborg's media (Gamborg *et al.*, 1968), D medium (Duncan *et al.*, 1985), McCown's Woody plant media (McCown and Lloyd, 1981), Nitsch and Nitsch (1969), and Schenk and Hildebrandt (1972) or derivations of these media supplemented accordingly. Those of skill in the art are aware that media and media supplements such as nutrients and growth regulators for use in transformation and regeneration and other culture conditions such as light intensity during incubation, pH, and incubation temperatures that can be optimized for the particular variety of interest.

After a transformable plant tissue is isolated or developed in tissue culture, or transformable plant tissue is identified and/or prepared *in planta*, the next step of the method is introducing the genetic components into the plant tissue. This process is also referred to herein as "transformation." The plant cells are transformed and optionally subject to a selection step. The independent transformants are referred to as transgenic events. A number of methods utilizing *Agrobacterium* strains have been reported and can be used to insert genetic components into transformable plant tissue. However, non-*Agrobacterium* spp. had not typically been utilized to transform plants.

Those of skill in the art are aware of the typical steps in the plant transformation process. The Rhizobia to be used can be prepared either by inoculating a liquid medium such as TY or YEM media (Beringer *et al.*, 1974) directly from a glycerol stock or streaking the bacteria onto a solidified media from a

glycerol stock, allowing the bacteria to grow under the appropriate selective conditions. The Rhizobia may be “pre-induced” by growth under nutritional or cultural conditions including the presence of acetosyringone in an amount that facilitates transformation. Those of skill in the art are familiar with procedures for growth and suitable culture conditions for bacteria as well as subsequent inoculation procedures. The density of the bacterial culture used for inoculation and the ratio of the number of bacterial cells to amount of explant tissue can vary from one system to the next, and therefore optimization of these parameters for any transformation method is expected.

10 The next stage of the transformation process is the inoculation. In this stage the suitably prepared plants, plant tissues, or explants, and the bacterial cell suspension are mixed together. The duration and condition of the inoculation and bacterial cell density will vary depending on the plant transformation system. Growth or inoculation of transforming bacteria may occur in the presence of acetosyringone, 15 or other known inducer of expression of the virulence genes located on Ti or Ri plasmids. In certain embodiments, growing of the bacterium other than *Agrobacterium* sp. is done under conditions to minimize polysaccharide production during growth in induction medium. In particular embodiments, the carbon source used to minimize polysaccharide production during Rhizobia growth in induction 20 medium is glucose in AB-TY medium, or L-arabinose and potassium gluconate in ATA medium.

After inoculation any excess bacterial suspension can be removed and the bacteria and target plant material are co-cultured. The co-culture refers to the time post-inoculation and prior to transfer to an optional delay or selection medium. Any 25 number of plant tissue culture media can be used for the co-culture step. Plant tissues after inoculation with bacteria may be cultured in a liquid or semi-solid media. The co-culture is typically performed for about one to four days.

After co-culture with bacteria, the inoculated plant tissues or explants can optionally be placed directly onto selective media. Alternatively, after co-culture with 30 bacteria, they could be placed on media without the selective agent and subsequently placed onto selective media. Those of skill in the art are aware of the numerous modifications in selective regimes, media, and growth conditions that can be varied

depending on the plant system and the selective agent. Typical selective agents include but are not limited to antibiotics such as geneticin (G418), kanamycin and paromomycin, or the herbicides glyphosate, glufosinate, and DICAMBA. Additional appropriate media components can be added to the selection or delay medium to
5 inhibit bacterial growth. Such media components can include, but are not limited to, antibiotics such as carbenicillin or cefotaxime.

The cultures are subsequently transferred to a medium suitable for the recovery of transformed plantlets. Those of skill in the art are aware of the number of methods to recover transformed plants. A variety of media and transfer requirements
10 can be implemented and optimized for each plant system for plant transformation and recovery of transgenic plants. Consequently, such media and culture conditions disclosed in the present invention can be modified or substituted with nutritionally equivalent components, or similar processes for selection and recovery of transgenic events, and still fall within the scope of the present invention.

Once the transformable plant tissue is inoculated, plant cells in the tissue may be transformed, and independently transformed plant cells are selected. The independent transformants are referred to as transgenic events. *Agrobacterium*-mediated transformation and regeneration systems for many monocot and dicot plant species are known in the art (e.g. Komari *et al.*, 1998; Zhou *et al.* 1995; Hiei *et al.*,
15 1994. Plant J.; 6:271–282; Ishida *et al.* 1996; Rogers *et al.*, 1987; Schrammeijer *et al.*, 1990; U.S. Patent 6,384,301), although use of Rhizobia for plant cell transformation has been reported only for tobacco, *Arabidopsis*, and rice (Broothaerts *et al.*, 2005). Following transformation and regeneration, transgenic plants are identified. Finally, one of skill in the art will recognize that after the expression cassette is stably
20 incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

The transformants produced, and their progeny, may subsequently be analyzed to determine the presence or absence of a particular nucleic acid of interest contained
30 on the transformation vector. Molecular analyses can include but are not limited to Southern blots (Southern, 1975), PCR (polymerase chain reaction) analyses, analysis of enzymatic activities, immunodiagnostic approaches, and field evaluations and the

like (Also see, for example, Sambrook *et al.*, 1989). These and other well known methods can be performed to confirm the stability of the transformed plants produced by the methods disclosed.

5 The above-described techniques may be suitable for any plant and is especially useful for plants such as alfalfa, barley, beans, beet, broccoli, cabbage, carrot, canola, cauliflower, celery, Chinese cabbage, corn, cotton, cucumber, dry bean, eggplant, fennel, garden beans, gourd, leek, lettuce, melon, oat, okra, onion, pea, pepper, pumpkin, peanut, potato, pumpkin, radish, rice, sorghum, soybean, spinach, squash, sweet corn, sugarbeet, sunflower, tomato, watermelon, and wheat.

10

EXAMPLES

Those of skill in the art will appreciate the many advantages of the methods and compositions provided by the present invention. The following examples are included to demonstrate the preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples 15 that follow represent techniques discovered by the inventors to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and 20 scope of the invention. All references cited herein are incorporated herein by reference to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, or compositions employed herein.

Example 1

***Rhizobium* and *Agrobacterium* strains**

25 *Agrobacterium tumefaciens* AGL0 was obtained from ATCC (ATCC Number: BAA-100™, Lazo *et al.*, 1991). *Rhizobium leguminosarum* strain Madison and *Sinorhizobium meliloti* SD630 were isolated from weed clover in a home garden in Madison, WI, USA, and confirmed by sequencing the PCR product of a 16S rRNA amplified with the following primers: 5' GAGAGTTTGATCCTGGCTCAG 3' 30 (Xd578; SEQ ID NO:1) and 5' AAGGAGGTGATCCAGCCGCAG 3' (Xd579; SEQ ID NO:2). Other *Rhizobium* strains were obtained from USDA *Rhizobium* collection center (Table 1). *Rhizobium* strains were grown in TY or MAG medium and

Agrobacterium in LB medium. Strains are shown below and 16s rRNA sequences amplified in strains isolated are provided as SEQ ID NOs:24-30.

Table 1: *Agrobacterium* and *Rhizobium* strains

Strain Name	Ti plasmid	Source
<i>A. tumefaciens</i> AGL0	pTiBo542	ATCC; Lazo <i>et al.</i> , 1991
<i>A. tumefaciens</i> LBA4404	pAL4404	Hoekema <i>et al.</i> , 1983
<i>A. tumefaciens</i> AGL0C	pTiBo542C (kanR)	This study
<i>A. tumefaciens</i> AGL0G	pTiBo542G (kanR)	This study
<i>A. tumefaciens</i> 4404TIK	pTi4404kan (kanR)	This study
<i>A. tumefaciens</i> ABI	pTiC58 (kanR, gentR)	Monsanto
<i>Rhizobium leguminosarum</i> <i>Madison</i>	None	This study
<i>Sinorhizobium meliloti</i> <i>SD630</i>	None	This study
<i>Rhizobium leguminosarum</i> <i>USDA2370</i>	None	USDA <i>Rhizobium</i> collection center
<i>Rhizobium leguminosarum</i> <i>bv. trifolii</i> <i>USDA2048</i>	None	USDA <i>Rhizobium</i> collection center
<i>Rhizobium leguminosarum</i> <i>bv. phaseoli</i> <i>USDA2668</i>	None	USDA <i>Rhizobium</i> collection center
<i>Sinorhizobium fredii</i> <i>USDA205</i>	None	USDA <i>Rhizobium</i> collection center
<i>Sinorhizobium meliloti</i> <i>USDA1002</i>	None	USDA <i>Rhizobium</i> collection center
<i>Mesorhizobium loti</i> <i>USDA</i> <i>3471</i>	None	USDA <i>Rhizobium</i> collection center
<i>Bradyrhizobium japonicum</i> <i>USDA 6</i>	None	USDA <i>Rhizobium</i> collection center
<i>Bradyrhizobium japonicum</i> <i>USDA 110</i>	None	USDA <i>Rhizobium</i> collection center
<i>Rhizobium etli</i> <i>USDA 9032</i> <i>(CFN42)</i>	None	USDA <i>Rhizobium</i> collection center
<i>R. leguminosarum</i> <i>2370G</i>	pTiBo542G	This study
<i>R. leguminosarum</i> <i>2370LBA</i>	pTi4404kan	This study
<i>R. leguminosarum</i> <i>bv. trifolii</i> <i>2048G</i>	pTiBo542G	This study
<i>R. leguminosarum</i> <i>bv. trifolii</i> <i>2048LBA</i>	pTi4404kan	This study
<i>R. leguminosarum</i> <i>bv.</i> <i>phaseoli</i> <i>2668G</i>	pTiBo542G	This study
<i>R. leguminosarum</i> <i>bv.</i> <i>phaseoli</i> <i>2668LBA</i>	pTi4404kan	This study
<i>R. leguminosarum</i> <i>RL542C</i>	pTiBo542C	This study
<i>S. fredii</i> <i>SF542G</i>	pTiBo542G	This study
<i>S. fredii</i> <i>SF4404</i>	pTi4404kan	This study
<i>S. meliloti</i> <i>SM542C</i>	pTiBo542C	This study
<i>M. loti</i> <i>ML542G</i>	pTiBo542G	This study
<i>M. loti</i> <i>ML4404</i>	pTi4404kan	This study

Example 2

Transformation of *Agrobacterium*

The *Agrobacterium* competent cells were prepared by washing a log phase culture in LB medium with chilled deionized water and 10% glycerol, and stored at -
5 80°C. Fifty microliters of thawed competent cells were mixed with 1 or 2 µl DNA on ice and electroporated in 1 mm gap curvet with 200 ohm resistance, 25 µF capacity and 1.8 kv using a BIO-RAD Gene Pulser® II device (BIO-RAD, Hercules, CA).

Example 3

Construction of Ti plasmids with an antibiotic selectable marker gene

10 To select Ti plasmids in *Rhizobium* spp., a homologous sequence was amplified from a corresponding Ti plasmid and inserted into a kanamycin resistance vector. The homologous sequence was used to integrate the kanamycin resistance gene into the Ti plasmid by homologous recombination.

To construct the pTiBo542C plasmid, the entire *virC* gene (Genbank accession
15 number AB027257) from the AGL0 *Agrobacterium* strain was amplified with PCR using the following primers 5' ACAATAATGTGTGTTGTTAAGTCTTGTTGC 3' (Xd683 SEQ ID NO:3) and 5' CTCAAACCTACACTCAATATTTGGTGAG 3' (Xd684 SEQ ID NO:4) and *Pfu* polymerase (STRATAGENE, La Jolla, CA) and inserted into the TOPO cloning blunt vector (Invitrogen Carlsbad, CA) giving rise to
20 an intermediate vector pMON67402. The intermediate vector was further ligated to a *trfA* fragment from pCGN11206 digested with PvuII/MscI, which resulted in construct pMON96913 (FIG. 6). The vector was then introduced into the AGL0 *Agrobacterium* strain by standard electroporation as outlined above, and plated on Kanamycin 50 mg/l LB medium to select for a single crossover event. Since the
25 integration vector is not maintained in AGL0 cells, the resistant colonies were presumably due to integration of the vector into a Ti plasmid by homologous recombination. The resulting strain was designated AGL0C.

Similarly, to construct the pTi542G plasmid, the entire *virG* sequence (Genbank accession number AB027257) from the AGL0 *Agrobacterium* strain was
30 PCR amplified using 5' AGATCTGGCTCGCGGCGGACGCAC 3' (Xd681; SEQ ID NO:5) and 5' CGCTCGGTCATTCTTTGCTGGAG 3' (Xd682; SEQ ID NO:6) with *Pfu* polymerase and inserted into pMON67402, which resulted in construct

pMON96026 (FIG. 8). This vector was introduced into the AGL0 strain by standard electroporation and plated on kanamycin 50 mg/l LB medium to select for a single crossover event. The resulting strain was designated AGL0G.

In order to construct the pTi4404kan helper plasmid, the *traI* and *trbCE* region
5 of octopine Ti plasmid pAL4404 (Genbank accession number NC_002377) from LBA4404 was PCR amplified with primers 5' TCAGCAGGATGACGCCGTTATCG 3' (Xd695; SEQ ID NO:7) and 5' TCTCGCCCGACCAATACCAACAC 3' (Xd696; SEQ ID NO:8) (sequence from Genbank AF242881) with *Pfu* polymerase, and inserted into pMON67402. The intermediate vector was further ligated to a *trfA*
10 fragment from pCGN11206 digested with PvuII/MscI, which resulted in construct pMON96914 (FIG. 7). This plasmid vector was introduced into LBA4404 by electroporation and selected on LB medium with kanamycin 50 mg/l to select for a single crossover event.

After a three day culture on solid medium, the kanR resistant colonies were
15 transferred into 2 mls of liquid LB medium with kanamycin 50 mg/l. One microliter of overnight culture was directly amplified with YieldAce® Taq polymerase following manufacturer instruction (Stratagene) with the following primers: 5' GCTGACGGCCCGGATGAATGTCAGCTACTG 3' (Xd715; SEQ ID NO:9) and 5' GCTCTAGAAATTGTAAGCGTTAATAATTCAGAAGAACTCGTC 3' (Xd716; SEQ ID NO:10) and integration of the kanamycin resistance gene into Ti plasmids
20 was confirmed. The resulting strain was designated as 4404TIK.

Example 4

Extraction of Ti plasmids from *Agrobacterium*

The modified Ti plasmids, pTiBo542C, pTiBo542G, pTi4404kan and pTiC58
25 (ABI), were extracted from the modified *Agrobacterium* strains AGL0C, AGL0G, 4404TIK and ABI, containing the respective plasmids. Five mls of overnight culture in LB with kanamycin 50 mg/l was spun down, resuspended in 400 µl of P1 buffer, mixed with 400 µl of P2 buffer, neutralized with 400 µl P3 buffer (buffers from QIAGEN maxi-prep kit). After 5 min incubation at room temperature, the mixture
30 was spun for 10 min at 12g at 4°C. Approximately 1200 µl of supernatant was mixed with 800 µl of isopropanol and spun for 10 min at 4°C. The pellet was washed with

70% ethanol once and resuspended in 200 μ l of TE without drying. The mega plasmids were subsequently stored at 4°C.

Example 5

Rhizobia competent cells and introduction of modified Ti plasmids into Rhizobia

5 Rhizobia competent cells were prepared according to Garg *et al.* 1999 with modification. Briefly, one loopful of Rhizobia (*e.g. Rhizobium*) cells from a frozen glycerol stock was grown in 10 mls of TY broth for 24 hours, and then transferred into 500 ml TY broth at 30°C with vigorous shaking and allowed to grow to mid-logarithmic phase ($OD_{600} = 0.4-0.6$). The culture was transferred into two 250
10 centrifuge tubes, chilled on ice for 15-30 min and centrifuged at 9,000 rpm for 10 min at 4°C to harvest cells. The cell pellet was washed with cold sterile deionized water, and with 10% cold glycerol and resuspended in 10% cold glycerol. The cell suspension was aliquoted at 50 μ l/tube for immediate use or frozen in liquid nitrogen and stored at -80°C.

15 Electroporation of the modified Ti plasmids into Rhizobia strains: Fifty microliters of the competent cells were thawed on ice, mixed with 1 or 2 μ ls of the prepared Ti plasmid, and kept on ice for 30 min. The mixture was transferred into a chilled 1 mm-gap electroporation cuvette. The electroporation parameters (BIO-RAD Gene Pulser® II) were set as follows: 2 KV/400 Ω resistance/25 μ F capacity or 1.5
20 KV/400 Ω resistance/25 μ F capacity or 1.5 KV/800 Ω resistance/10 μ F capacity. After electroporation, the cuvette was kept on ice for 5-10 min before adding 1 ml of TY or MAG medium and transferring into a 14-ml Falcon tube. The tube was cultured for 3 hours at 30°C, plated onto TY or MAG solid medium with 50 mg/l kanamycin and cultured at 30°C for three days to recover resistant colonies.

25 Confirmation of Rhizobia transformed with Ti plasmids: The kanamycin resistant colonies were transferred into 3 mls of liquid TY or MAG medium with 50 mg/l kanamycin and cultured overnight. One microliter of culture was directly amplified with YieldAce® Taq polymerase following manufacturer's instructions (Stratagene).

30 To detect pTiBo542C or pTiBo542G in Rhizobia strains, the *virC* primers 5' ACAATAATGTGTGTTGTTAAGTCTTGTTC 3' (Xd683; SEQ ID NO:3) and 5'

CAATTGCATTTGGCTCTTAATTATCTGG 3' (Xd684a; SEQ ID NO:11) or virG primers 5' AGATCTGGCTCGCGGCGGACGCAC 3' (Xd681; SEQ ID NO:5) and 5' CGCTCGCGTCATTCTTTGCTGGAG 3' (Xd682; SEQ ID NO:6) were used to amplify a 2.35 kb or a 1.2 kb fragment, respectively.

5 For the pTi4404kan plasmid, the following primers were used: 5' GCATGCCCCGATCGCGCTCAAGTAATC 3' (Xd699; SEQ ID NO:12) and 5' TCTAGGTCCCCCGCGCCCATCG 3' (Xd700; SEQ ID NO:13)) amplifies a 1274 bp *virD2* coding sequence for the octopine Ti plasmid; 5' CCATGGATCTTTCTGGCAATGAGAAATC 3' (Xd701; SEQ ID NO:14) and 5' 10 GTCAAAGCTGTTGACGCTTTGGCTACG 3' (Xd702; SEQ ID NO:15) amplifies a 1602 bp *virE2* fragment; 5' ACGGGAGAGGCGGTGTTAGTTGC 3' (Xd703; SEQ ID NO:16) and 5' CGATAGCGACAATGCCGAGAACG 3' (Xd704; SEQ ID NO:17) amplifies approximately a 0.9 kb *virB1* fragment.

In order to identify the pTiC58 plasmid from the ABI strain, three pairs of 15 primers were used: 5' ATGCCCCGATCGAGCTCAAGTTATC 3' (Xd685; SEQ ID NO:18) and 5' TGAAAGGACACCTCTCCGTTGCTG 3' (Xd686; SEQ ID NO:19) amplifies a 1247 bp *virD2* fragment; 5' CCATGGATCCGAAGGCCGAAGGCAATG 3' (Xd687; SEQ ID NO:20) and 5' CTACAGACTGTTTACGGTTGGGC 3' (Xd688; SEQ ID NO:21) amplifies a 20 1670bp *virE2* entire coding sequence; 5' GTGAGCAAAGCCGCTGCCATATC 3' (Xd689; SEQ ID NO:22) and 5' TAGAGCGTCTGCTTGGTTAAACC 3' (Xd690; SEQ ID NO:23) amplifies a 1102bp partial *repA* fragment.

Example 6

Media for bacterial growth

25 Media used for Rhizobia growth in the Rhizobia-mediated transformation protocol employed to develop transformed plants were prepared using standard methods known to one skilled in the art. Media formulations are as follows:

TY medium

30	Bactotryptone	per L
	Yeast extract	5g/L
	CaCl ₂ .2H ₂ O	3g/L
	pH 7.0	0.87 g/L

For solid TY medium, add 15 g/l Bacto-Agar before autoclaving.

MAG Medium Liquid
(From USDA *Rhizobium* collection center)

5	HEPES	1.3 g/L
	MES	1.1 g/L
	Yeast extract	1 g/L
	L-Arabinose	1 g/L
	Potassium gluconate	1 g/L
10	KH ₂ PO ₄	0.22 g/L
	Na ₂ SO ₄	0.25 g/L
	pH 6.6 with KOH	

Autoclave and add the following filter sterilized stock solution:

15	NH ₄ Cl (16 g/100 ml)	2 ml/l
	FeCl ₃ (0.67g/100 ml, FS)	1.0 ml/l
	CaCl ₂ (1.5 g/100 ml)	1.0 ml/l
	MgSO ₄ (18 g/100 ml)	1.0 ml
	NaMoO ₄ · 2H ₂ O (1 g/100 ml)	1.0 ml/l
20	NiCl ₂ · 6H ₂ O (2.2 g/100 ml)	0.1 ml/l
	For solid MAG medium, add 15 g/l Bacto-Agar before autoclaving.	

	LB medium	Per liter
	Bacto-tryptone	10 g
25	Bacto-yeast extract	5 g
	NaCl	10 g
	Adjust pH to 7.5 with sodium hydroxide.	
	After autoclaving, distribute into culture plate (25 ml/plate)	
	For solid LB medium, add 15 g/l Bacto-Agar before autoclaving	

30 **AB Minimal Medium:**

20x AB Buffer:
 K₂HPO₄ 60 g/l
 NaH₂PO₄ 20 g/l
 Autoclave separately

35	20x AB Salts (Filter sterilized, keep in dark):	
	NH ₄ Cl	20 g/l
	MgSO ₄ · 7H ₂ O	6 g/l
	KCl	3 g/l
40	CaCl ₂	0.2 g/l
	FeSO ₄ · 7H ₂ O	50 mg/l
	pH to 7 before autoclaving	

45 Combine 50 ml AB Buffer and 50 ml AB Salts with 900 ml sucrose-water (final concentration of sucrose in one liter is 0.5%).

1x AB-TY Induction medium:

	Glucose	5 g
	20x AB buffer	50 ml
	20X AB Salt stock	50 ml
5	TY medium	20 ml
	Add sterile water to	1000 ml
	Adjust to pH 5.4 with 100 mM MES	

Add 100 μ M acetosyringone (1 M stock in DMSO; add 1 μ l stock/10 ml medium after suspending bacteria)

10 ATA Medium for vir Induction in *Rhizobium*:

The ATA (AB minimal medium +TY +Arabinose) medium was modified from AB-TY medium using arabinose and potassium gluconate to replace glucose. The growth rate of all *Rhizobia* is almost doubled in this medium. The bacteria produced much less polysaccharide in this medium and the bacterial pellets were much tighter.

15

	L-Arabinose	1 g/L
	Potassium gluconate	1 g/L
	20x AB buffer	50 ml
	20x AB salt stock	50 ml
20	TY medium	20 ml
	Add sterile water to	1000 ml
	pH 5.4 with 100 mM MES	
	Add 200 μ M acetosyringone (1 M stock in DMSO) after resuspending bacteria.	

Example 7**25 Crop transformation vectors for use with modified *Rhizobia* strains**

Rhizobium transformation vectors were constructed using standard molecular techniques known to those skilled in the art. Plasmid constructs pMON96033 (FIG. 1; for soybean and canola transformation), pMON96036 (FIG. 2; for corn transformation), or pMON101316 (FIG. 3; for cotton transformation) were employed.

30 All three constructs contain a pVS1 replication origin, and either GUS, GFP, or both reporter genes. Recombinant plasmids were transferred into various modified *Rhizobia* strains by electroporation and confirmed by restriction enzyme digestion of miniprep DNA.

The FMV CP4 gene used in constructing the plasmids has a promoter from

35 Figwort Mosaic Virus (FMV) followed by the CP4syn gene, a synthetic gene encoding CP4 EPSP synthase. See, U.S. Patent No. 5,633,435, which is incorporated by reference herein. EPSP synthase, when expressed, confers a substantial degree of glyphosate resistance upon the plant cell and plants generated there from. The e35s GUS gene is a β -glucuronidase gene, which is typically used as a histochemical

marker, behind the e35S promoter. The FMV GUS gene is the FMV promoter with GUS. The NOS NPTII gene has a neomycin phosphotransferase gene, which confers resistance to kanamycin, behind the promoter for the nopaline synthase gene (NOS). The Act 1 GFP gene has an actin promoter from rice and the gene for green
5 fluorescence protein, which is a screenable marker. The e35s GFP gene is the gene for green fluorescence protein behind the e35S promoter. Overnight cultures of a Rhizobia strain containing the plasmid used were grown to log phase and then diluted to a final optical density of 0.3 to 0.6.

Example 8

10

Rhizobia- mediated soybean transformation

Transformation was performed using an organogenesis process, as described by Martinell *et al.* (U.S. Patent 7,002,058), with modifications. pMON96033 containing the GUS and CP4 genes were transferred into various modified Rhizobia strains (*e.g. Rhizobium sp., Mesorhizobium sp., Sinorhizobium sp.*) by electroporation.
15 Single colonies were recovered on MAG or TY medium with 50 mg/l spectinomycin and 50 mg/l kanamycin and inoculated in 20-50 mls of liquid TY medium with the same selection in a shaker at 30°C at 200 rpm. The presence of plasmid in the Rhizobia culture was verified by restriction enzyme digestion of mini-prepared plasmid from 10 ml culture. The remaining liquid culture was mixed with glycerol to
20 a final concentration of 20%, aliquoted and stored at -80°C as seed cultures.

To prepare Rhizobia inoculum, 0.25-1 ml frozen seed culture was inoculated into 250 or 500 mls of TY medium with the same antibiotic selection as above and grown overnight at 28°C with shaking at 200 rpm to mid-log growth phase. The culture was spun down and directly suspended in an inoculation medium (INO
25 medium) at the concentration of OD₆₆₀ about 0.3.

Induced Rhizobia culture was also used in soybean transformation. To induce the Rhizobia, the overnight culture was resuspended in AB-TY medium at an OD₆₆₀ of about 0.3 and acetosyringone was added to a final concentration of 100 µM. The culture was further shaken overnight at 28°C, spun down and re-suspended in the
30 inoculation medium (INO medium) to a concentration of OD₆₆₀ about 0.3.

Soybean cultivar A3525 (U.S. Patent 7,002,058) was used for Rhizobia-mediated transformation. The method was modified for Rhizobia-mediated

transformation as follows. Soybean seeds were germinated at room temperature in BGM medium and meristem explants from soy mature seeds were excised by machine (U.S. Application 20050005321). Soybean meristem explants in a PLANTCON lid were mixed with Rhizobia suspension in INO medium and sonicated in a W-113 Sonicator (Honda Electronics Co., Ltd, Aichi, Japan). After sonication, the explants were co-cultured in the same PLANTCON for 1-11 days at 23°C with a 16/8 hour light-dark photo period. The explants were then transferred onto the surface of the WPM selection medium containing 75 µM glyphosate. After 2 weeks, explants were transferred again to 75 µM glyphosate solid WPM medium. Shoots with fully expanded trifolia were recovered after 6-10 weeks post-inoculation and rooted in BRM medium (optionally with fungicide) containing 0.1 mg/l IAA and 25 µM glyphosate selection. The rooted plantlets were transferred to the greenhouse for maturity.

Table 2: Media components for soy transformation.

BGM medium for soybean seed germination	
amount/L	Compound
0.505 g	Potassium nitrate
0.24 g	Ammonium nitrate
0.493 g	Magnesium sulfate
0.176 g	Calcium chloride
27.2 mg	Potassium phosphate monobasic
1.86 mg	Boric acid
5.07 mg	Manganese sulfate
2.58 mg	Zinc sulphate
0.249 mg	Potassium iodide
0.216 mg	Sodium Molybdate
0.0008 mg	Copper sulphate
0.0008 mg	Cobalt chloride stock
3.36 mg	Disodium EDTA
2.49 mg	Ferrous sulphate
1.34 mg	Thiamine HCl
0.5 mg	Nicotinic acid
0.82 mg	Pyridoxine HCl
20 g/L	Sucrose (Ultra Pure)
125 mg	Cefotaxime
pH	5.6
INO medium for soy co-culture	
amount/L	Compound
1/10x of Gamborg B5 medium micronutrient and vitamin components; 2/5x of macronutrients	

1 g	Potassium Nitrate (KNO ₃)
30 g	Glucose
3.9 g	MES (pH 5.4)
After autoclaving, lipoic acid added to inoculum to a final concentration of 250 μM	
SOY WPM shooting medium	
amount/L	Compound
2.41 g	WPM Powder (PhytoTech Laboratories)
20 g	Sucrose (Ultra Pure)
1.29 g	Calcium Gluconate (Sigma)
4.0 g	AgarGel (pH 5.6)
mL/L	Post-autoclaving ingredients
4 mL	Cefotaxime (50 mg/mL)
1 ml	Ticarcillin (100 mg / ml)
5 mL	Carbenicillin (40 mg/mL)
0.15 mL	Glyphosate (0.5 FS Stock) (0.075 mM)
BRM rooting medium	
amount/L	Compound
2.15 g	MS Powder (Phytotech)
0.1 g	myo-Inositol
2 mg	Glycine
0.5 mg	Nicotinic acid
0.5 mg	Pyridoxine HCl
0.1 mg	Thiamine HCl
30 g	Sucrose (Ultra Pure)
10 ml	L-Cysteine (10 mg/ml)
8 g	Washed Agar
mL/L	Post-autoclaving ingredients
5.0	IAA (0.033 mg/ml in 1 mM KOH)
1 mL	Ticarcillin (100 mg / ml)
0.05 mL	Glyphosate (0.5 FS Stock) (0.025 mM)

The binary vector pMON96033 was transferred into Rhizobia strains and co-cultivated with soybean meristem explants, and GUS positive results were observed (Table 3 and FIG. 4). *S. meliloti*, *S. fredii*, *M. loti* and one *R. leguminosarum* showed T-DNA delivery into soybean explants demonstrated by small blue spots of GUS activity. Transgenic soybean plants were obtained from Rhizobia-mediated transformation experiments with various strains (Table 4). The transgenic nature of

these soybean plants were confirmed by transgene copy number assay, where most of the transformants revealed 1-2 copy simple integration pattern. (Table 5).

Table 3: Transient Expression of *gus* Gene with Rhizobia-mediated T-DNA Delivery in soybean meristem explants.*

Strain	Origin	Transient GUS assay
RL4404	<i>Rhizobium leguminosarum</i> strain Madison+pAL4404	+
2370LBA	<i>Rhizobium leguminosarum</i> strain USDA2370+pTiBo542C	+
2370G	<i>Rhizobium leguminosarum</i> strain USDA2370+pTiBo542G	+
SF4404	<i>Sinorhizobium fredii</i> USDA205+pAL4404	+
SF542C	<i>Sinorhizobium fredii</i> USDA205+pTiBo542C	+
SM542C	<i>Mesorhizobium loti</i> USDA3471+pAL4404	+
ML4404	<i>Mesorhizobium loti</i> USDA3471+pAL4404	+
ML542G	<i>Mesorhizobium loti</i> USDA3471+pTiBo542G	+

5 * Transient assays were performed after 4 day co-culture period.

Table 4: Rhizobia-mediated soy transformation summary.

Strains	Soy Explants	Rooted Plants	TF
RL 4404	3705	2	0.05%
SF 4404	5553	2	0.04%
SM 542C	2555	1	0.04%

10 **Table 5: Copy Number Assay of Transgenic Plants from Rhizobia-mediated Transformation***

NOS copy	RL4404	SF4404	SM542C
0 copy	0	0	1
1-2 copy	1	2	0
>2 copy	0	0	0
Total Plant	1	2	1

*Copy number was analyzed by INVADER method (Third Wave Technologies, Madison, WI) using a *nos* probe and compared with an internal genome control.

To test if the *gus* transgene was transmitted to the seed progeny, seeds of two soy transgenic lines derived from Rhizobia-mediated transformation were stained in GUS solution (FIG. 5). The GM_A9196D line was found to have one copy of the linked *nos* gene as assayed by the INVADER method. Twelve R1 seeds from this line were assayed for GUS by histochemical staining after imbibition and removal of seed coat, and 9 were GUS positive, indicating a segregation ratio of 3:1 for one copy insert.

Example 9

Rhizobia- mediated canola transformation

A. *Rhizobium* inoculum preparation:

Rhizobia strains with pMON96033 were used for canola transformation. The
5 Rhizobia strains with the vector from a glycerol stock were inoculated into 10 mls of
TY medium with 50 mg/l kanamycin and 50 mg/l spectinomycin in a 50 ml Falcon
tube and shaken at 28°C overnight at 200 rpm. The overnight Rhizobia culture was
pelleted by centrifugation and resuspended in MG/L liquid medium. (MG/L broth:
Mannitol 5 g/l, L-glutamic acid 1 g/l, KH₂PO₄ 250 mg/l, NaCl 100 mg/l,
10 MgSO₄·7H₂O 100 mg/l, biotin 1 µg/l, yeast extract 2.5 g/l, pH7.0). The OD₆₀₀ was
between 0.05-0.1.

B. Canola explant preparation and co-cultivation:

Canola transformation was done according to U.S. Patent 5,750,871 and
Radke *et al.*, 1992. About 0.25g of canola seed, cv. Ebony, was transferred into a
15 1.5-ml Eppendorf tube and wetted with 95% ethanol. To sterilize the seeds, 1 ml of
1% sodium hypochlorite solution was added for 30 min. The bleaching solution was
replaced with distilled water and the seeds were rinsed several times. The seeds were
spread onto 1/10 MS germination medium and kept in a Percival incubator at 24°C
with a 16 hour light photo period.

20 **Seed Germination Medium (1/10 MS medium):** 1/10X MS minimal
organics medium (Gibco BRL; final sucrose 0.3%), pyridoxine 50µg/l, nicotinic acid
50µg/l, glycine 200 µg/l, PHYTAGAR (Gibco Invitrogen) 6 g/l, pH 5.8; 20).
Etiolated seedlings from 7-14 days old cultures were used as the explant source.

25 Explants were inoculated in 1 x 10⁸ bacteria/ml. *Rhizobium* suspension was
drawn off, and the inoculated explants were placed onto co-cultivation plates on top
of filter paper, and incubated for about 2 days at 24°C in continuous light. Co-
cultivated explants were assayed for *gus* expression and found to contain blue spots
indicating transformation of canola cells (FIG. 9).

30 **Co-cultivation Medium (MS-1):** MS salts (Caisson Laboratories, Logan,
UT), myo-inositol 100 mg/l, thiamine-HCl 1.3 mg/l, KH₂PO₄ 200 mg/l, 2,4-D 1 mg/l,
sucrose 3%, PHYTAGAR (Gibco Invitrogen) 7 g/l, pH 5.8.

C. Callus Induction:

Co-cultivated explants were transferred to Callus Induction (B5-1) medium. for 6 days at 24°C in continuous light at ~ 100 $\mu\text{E}/\text{m}^2/\text{s}$. Five strains from *M. loti*, *R. leguminosarum*, *S. fredii* and *S. meliloti* showed efficient gene transfer into canola
5 explants with frequency of *gus* positive explants ranging from 21% to 73% (FIG. 10).

Callus Induction Medium (B5-1): Gamborg's B5 salts (Caisson Labs), B5 vitamins (1 mg/l nicotinic acid, 1 mg/l pyridoxine-HCl, 10 mg/l thiamine-HCl), 100 mg/l inositol, 1 mg/l 2,4-D, sucrose 3%, carbenicillin (PhytoTechnology, Shawnee Mission, KS) adjusted to final potency of 325 mg/l, 50 mg/l Timentin, 7 g/l
10 PHYTAGAR (Gibco Invitrogen), pH5.8.

D. Shoot Regeneration and Selection:

Explants having callus were transferred to Shoot Regeneration medium (B5BZ) with AgNO_3 and incubated at 24°C in continuous light of 100 $\mu\text{E}/\text{m}^2/\text{s}$ for 14 days. Explants were next transferred to Shoot Regeneration medium (B5BZ) without
15 AgNO_3 . Shoots regenerated from glyphosate-selected calli were harvested ~ every two weeks. An example of early shoots showing *gus* expression is shown in FIG. 11.

Shoot Regeneration Medium with Silver Nitrate (B5BZ + 3Ag): Gamborg's B5 salts (Caisson Labs), B5 vitamins (1 mg/l nicotinic acid, 1 mg/l pyridoxine-HCl, 10 mg/l thiamine-HCl), 100 mg/l inositol, BAP 3 mg/l (Sigma),
20 zeatin 1 mg/l (Sigma), AgNO_3 , 3 mg/l (Sigma), 45 mg/l glyphosate (Monsanto, 96.5% dry acid,), sucrose 1%, carbenicillin (PhytoTechnology) with potency adjusted to 325 mg/l, 50 mg/l Timentin, PHYTAGAR (Gibco Invitrogen) 7 mg/l, pH 5.8.

Shoot Regeneration Medium (B5BZ): Gamborg's B5 salts (Caisson Labs), B5 vitamins (1 mg/l nicotinic acid, 1 mg/l pyridoxine-HCl, 10 mg/l thiamine-HCl),
25 100 mg/l inositol, BAP 3 mg/l (Sigma), zeatin 1 mg/l (Sigma), 45 mg/l glyphosate, sucrose 1%, carbenicillin (PhytoTechnology) with potency adjusted to 325 mg/l, Timentin 50 mg/l, PHYTAGAR (Gibco Invitrogen) 7 mg/l, pH 5.8.

E. Shoot Harvest:

Green shoots at least 0.5 cm in length were trimmed to isolate the main axis.
30 Trimmed shoots were placed on Shoot Harvest medium (B5-0). Shoots were transferred to Rooting medium after 2 weeks.

Shoot Harvest Medium (B5-0): Gamborg's B5 salts (Caisson Labs), B5 vitamins (1 mg/l nicotinic acid, 1 mg/l pyridoxine-HCl, 10 mg/l thiamine-HCl), 100 mg/l inositol, carbenicillin (PhytoTechnology) with potency adjusted to 195 mg/l, sucrose 1%, PHYTAGAR (Gibco Invitrogen) 6 g/l, pH 5.8.

5 **F. Shoot Growth and Rooting:**

Green shoots were transferred to Rooting medium (B5-0 + 2IBA). Shoots remained on Rooting medium until they formed roots. Shoots were maintained at 24°C, 16 hours light/day, ~100 uE/m²/s.

10 **Rooting Medium (B5-0 + 2IBA):** Gamborg's B5 salts (Caisson Labs), B5 vitamins (1 mg/l nicotinic acid, 1 mg/l pyridoxine-HCl, 10 mg/l thiamine-HCl), 100 mg/l inositol, IBA 2 mg/l (indole-3-butyric acid, Sigma) 150 mg/l cefotaxime (PhytoTechnology), sucrose 1%, PHYTAGAR (Gibco Invitrogen) 6 g/l, pH 5.8.

G. Transgene detection and transformation frequency:

15 Total genomic DNA was extracted from greenhouse grown canola plants, digested with a single cutter *Bgl*III, and hybridized with DIG-labeled CP4 probe. Four lines were confirmed to be transgenic (FIG. 11). The transformation frequency is summarized in the Table 6.

Table 6: Canola transformation frequency (TF) with *Rhizobium* strains

Strains	Explants	GUS/CP4 positive	TF
RL 2370G	150	2	1.33
SF 542	120	1	0.83
SM542C	120	1	0.83

20

Example 10

Rhizobia-mediated cotton transformation through embryogenesis

A. Rhizobia inoculum preparation:

25 pMON101316 was electroporated into Rhizobia strains, verified by restriction digestion of mini-prepared DNA and stored at -80°C. The Rhizobia strains with the vector from the glycerol stock were inoculated into 10 mls of TY medium with kanamycin (50 mg/l) and spectinomycin (50 mg/l) in a 50 ml Falcon tube and shaken at 28°C overnight at 200 rpm. The overnight Rhizobia culture was pelleted by

centrifugation, resuspended in 20 mls of MS0 liquid medium and centrifuged again. The pellet was resuspended in 20 mls of MS0 medium. The washed Rhizobia was diluted in MS0 to an OD₆₆₀ of about 1.0 for inoculation.

B. Explant Preparation:

5 Cotton transformation was done essentially according to U.S. Publ. 2004087030. Seven days after seedlings were germinated, etiolated cotton seedlings from cultivar Coker were removed from a dark Percival incubator. The hypocotyls from the PHYTATRAYS were harvested and placed in a sterile Petri dish containing sterile MS0 to prevent the tissue from drying out. Hypocotyls were cut into small
10 explants.

C. Inoculation and co-cultivation:

Using a sterile forceps, explants were transferred to sterile Petri dishes, and Rhizobia inoculum was added. Explants were left in a sterile hood for 20 minutes, with swirling to ensure good contact of all explants with the Rhizobia inoculum. The
15 Rhizobia inoculum solution was then aspirated out and explants were gently blotted with sterile filter paper. The inoculated hypocotyl pieces were placed onto culture plates. Co-culture of the plates of explants, covered with a plastic bag, was performed in a Percival incubator set at about 22-24°C, with a 10 hour light/14 hour dark photoperiod for 2 days.

20 **D. Stable transformation through embryogenesis:**

Two days post inoculation, cotton explant pieces were stained with X-gluc to test for GUS transient expression (FIG. 12). Blue spots in hypocotyls indicate the expression of the *gus* gene and transformation of cotton cells. In order to obtain stable transformed plants, hypocotyl explants were transferred onto a plate containing
25 UMSEL1629 selection medium, containing the appropriate selection agent. The plates were then covered with PARAFILM and cultured 28°C with a 16/8 hr. (day/night) photo period.

The stably transformed calli were confirmed by X-Gluc staining for *gus* expression after 4 weeks on the selection medium (FIG. 13). Four weeks after the
30 initial transfer to selection medium, all the hypocotyls were transferred to UMSEL 1788 medium, PARAFILMed and cultured for 7 days. Then the explants were

transferred back onto UMSEL1629 for 4 weeks at 28°C with a 16/8 hr. (day/night) photo period.

Approximately 4 weeks after the second transfer on UMSEL1629, depending upon the growth rate of the callus, clumps of calli were recovered and were transferred to UMO plates. Individual plates are then labeled, covered with PARAFILM, and cultured at 28°C in continuous dark.

Six to eight weeks after the callus has been on UMO the calli are subcultured onto fresh UMO medium and cultured at 28°C in continuous dark. After 6-10 weeks on UMO, embryogenic callus (EC) is ready to be harvested from independent callus lines on UMO and is transferred to TRP+ medium. Every 3-5 weeks, for approximately 3 months, actively growing tissue and small embryos on TRP+ plates are transferred to fresh TRP+ medium and cultured at 28°C in continuous dark. Embryos are transferred to SHSU medium in Petri plates and covered with PARAFILM. Plates are cultured at 28°C in a Percival with a 16/8 (day/night) photo period with maximum lighting (shelf and side lights). The embryos may be subcultured on same medium one more time until germination. Plantlets recovered are cultured in a Percival or warm room at 28°C with a 16/8 (day/night) photo period.

E. Molecular analysis of transgenic nature of cotton calli derived from *Rhizobium*-mediated transformation:

Genomic DNA was extracted from callus tissue, digested with a single cutter *Bam*HI, fractionated in 1% agarose gel, and transferred onto Hybond™ membrane (e.g. Appligen-Oncor, Illkirch, France; or Amersham-Pharmacia Biotech). A DIG labeled *gus* probe was used to detect the presence of the transgene as an indicator of transformation. Six lines of cotton calli, derived from *S. meliloti*, *S. fredii* and *R. leguminosarum* transformation with Ti helper plasmid, were found to contain the *gus* gene (FIG. 14).

F. Media for cotton culture:

Recipe for 1 L of UMSEL - 4.33 g MS salts, 2 ml 500X B5 vitamins, 0.1 ml 2,4-D (1 mg/ml), 1 ml kinetin (0.5 mg/ml), 30 g glucose, pH 5.8, 2.5 g PHYTAGEL, 1.7 ml carbenicillin (250 mg/ml), 1 ml cefotaxime (100 mg/ml), plus selection agent: kanamycin 40 mg/L final concentration. Carbenicillin, cefotaxime and selective agents were added post-autoclaving.

Recipe for 1 L of UMSEL1788: - 4.33 g MS salts, 2 ml 500X B5 vitamins, 0.1 ml 2,4-D (1 mg/ml), 1 ml kinetin (0.5mg/ml), 30 g glucose, pH 5.8, 2.5 g PHYTAGEL, 1.7 ml (250 mg/ml) carbenicillin, 1 ml (100 mg/ml) cefotaxime, plus selection agent: kanamycin 40 mg/L final concentration and 0.1 g sucrose dissolved in
5 100 ml water. Carbenicillin, cefotaxime and selective agents were added post-autoclaving.

Recipe for 1 L of UMSEL1629: - 4.33 g MS salts, 2 ml 500X B5 vitamins, 0.1 ml 2,4-D (1 mg/ml), 1ml kinetin (0.5 mg/ml), 30 g glucose, pH 5.8, 2.5g PHYTAGEL, 1.7 ml (250 mg/ml) carbenicillin, 1ml (100 mg/ml) cefotaxime, plus
10 selection agent: kanamycin 40 mg/L final concentration. Carbenicillin, cefotaxime and selective agents were added post-autoclaving.

Recipe for 1 L of UMO - 4.33g MS salts, 2 ml 500X B5 vitamins, 30 g glucose, pH 5.8, 3.5 g GELRITE, 1.7 ml (250 mg/ml) carbenicillin, 1 ml (100mg/ml) cefotaxime, 100 mg/l ascorbic acid, plus selection agent: kanamycin 50 mg/L final
15 concentration.

Recipe for 1 L of TRP+ - 4.33 g/l MS salts, 2 ml 500X B5 vitamins, 1.9 g/l KNO₃, 30 g/l glucose, 0.1 g/l casein hydrolysate, 3.5 g GELRITE, pH 5.8.

Recipe for 1 L of SHSU – 100 ml Stewart & Hsu Majors (10x), 10 ml Stewart & Hsu Minors (100x), 1.5 ml iron (100x), 10 ml Stewart & Hsu Organics
20 (100x), 5 g glucose, 50 mg/l benlate, 2.2 g GELRITE, pH 6.8 (Stewart & Hsu, 1977).

Example 11

Rhizobia-mediated corn transformation

A. *Rhizobium* inoculum preparation and media composition:

pMON96036 containing CP4, GUS and *gfp* expression cassettes was used for
25 corn transformation. The vector was electroporated into various modified Rhizobia strains, verified, and stored at -80°C. Rhizobia containing the vector in a glycerol stock were streaked out on solid TY medium supplemented with antibiotics (kanamycin 40 mg/L and spectinomycin 31 mg/L), and incubated at 28°C for 2 days.

Two days before Rhizobia inoculation of the maize immature embryos, one
30 loopful of cells from a Rhizobia culture plate was inoculated into 25 mL of liquid TY medium supplemented with 62 mg/L of spectinomycin and 40 mg/L of kanamycin in

a 250 mL flask. The flask was placed on a shaker at approximately 150-200 rpm and 27-28°C overnight. The Rhizobia culture was then diluted (1 to 5) in the same liquid medium and put back on the shaker. Several hours later, one day before inoculation, the Rhizobia cells were spun down at 3500 rpm for 15 min. The bacterial cell pellet
5 was re-suspended in AB-TY or ATA induction broth with 200 µM of acetosyringone and 50 mg/L spectinomycin and 25 mg/L kanamycin and the cell density was adjusted to 0.2 at OD₆₆₀. The bacterial cell culture (50 mL in each 250 mL flask) was then put back on the shaker and grown overnight. On the morning of inoculation day, the bacterial cells were spun down and washed with liquid 1/2 MS VI medium (U.S.
10 Publ. 20040244075) supplemented with 200 µM of acetosyringone. The bacterial culture was centrifuged and the cell pellet was re-suspended in 1/2 MS PL medium (U.S. Publ. 20040244075) with 200 µM of acetosyringone and the cell density was adjusted to 1.0 at OD₆₆₀ for inoculation. Reagents are commercially available and can be purchased from a number of suppliers (see, for example Sigma Chemical Co., St.
15 Louis, Mo.).

B. Corn embryo isolation and *Rhizobium* co-cultivation:

For Rhizobia-mediated transformation, ears containing immature corn (*Zea mays*) embryos were isolated and transformed by bacterial co-culture as generally described by Cai *et al.* (U.S. Patent Application Publication 20040244075), except
20 that the immature embryos were isolated from surface sterilized ears and directly dropped into the prepared Rhizobia cell suspension. After the Rhizobia cell suspension was removed, the immature embryos were transferred onto the co-culture medium (U.S. Publ. 20040244075).

To investigate GFP transient expression, the co-cultivated corn embryos were
25 directly placed under a microscope with fluorescence light for GFP observation. Alternatively, 10 randomly picked embryos after co-cultivation were transferred into 1.5 ml Eppendorf tube and stained with X-gluc solution overnight at 37°C for *gus* transient expression. FIG. 15 represents GUS transient expression of corn immature embryos transformed with five *Rhizobium* strains of *R. leguminosarum*, *M. loti*, *S.*
30 *fredii* and *S. meliloti* compared to *Agrobacterium tumefaciens* strain ABI using ATA induction medium.

It was noted that routine AB minimal medium used for *Agrobacterium* growth and induction does not efficiently support Rhizobia growth. Rhizobia inoculums did not show significant growth in AB minimal medium without any selection after one week shaking with 220 rpm at 28°C. Inclusion of 20 ml TY medium in AB minimal medium dramatically improves Rhizobia growth rate; while replacing glucose in AB-TY medium with L-arabinose and potassium gluconate in ATA medium decreases polysaccharide production, resulting into tighter pellets. The change of carbon source in induction medium significant improves *gus* transient expression with Rhizobia strains in corn.

10 **C. Callus induction and regeneration of transgenic plants:**

After co-cultivation, transformation was continued essentially as described in U.S. Publ. 20040244075, with modifications of selection conditions as appropriate. The embryos were transferred onto a modified MS medium (U.S. Publ. 20040244075) supplemented with 250 mg/L carbenicillin and 0.1 mM glyphosate. Stably transformed calli with *gfp* expression were observed from the *M. loti*, *S. fredii* and *S. meliloti* stains used in transient assay at this stage (FIG. 16). Representative transformation frequencies are shown in Table 7.

Table 7: Corn transformation frequency with different Rhizobia strains.

Induction medium	Rhizobia Strain	Embryos	Transgenic plants	TF
AB induction	ML4404	177	3	1.69%
	ML542G	154	2	1.30%
	ABI	83	7	8.40%
ATA Induction	ML542G	112	2	1.78%
	SF4404	116	6	5.17%
	SF542C	124	2	1.61%
	SM542C	134	1	0.75%
	2370LBA	139	0	0%
	ABI	100	1	1%

20 **D. Molecular analysis of transgenic plants derived from Rhizobia-mediated transformation:**

Total genomic DNA was isolated from greenhouse grown corn plants and digested with a single cutter *Bam*HI to estimate transgene copy number. A DIG-

labeled *gus* probe was used to hybridize with the genomic DNA. The transgenic nature of putatively transformed tissues was confirmed for all lines but one (FIG. 17).

E. Germline transmission of transgenes in the transgenic corn plants:

The flowering transgenic corn plants were either selfed or outcrossed with the parental line of the corn genotype used for transformation (line LH244). Dry seeds were imbibed in water for 1 day for *gus* staining or 2 days for *gfp* counting. *gus* or *gfp* expression and segregation in the transgenic R1 seeds were confirmed (Table 8).

Table 8: Transgene expression in the transgenic corn R1 seeds.

Transgenic line	seed #	GFP + seeds / seeds assayed	Gus + seeds / Seeds assayed	Rhizobia strain
ZM_A8232	112	9 / 20	6/7	ML542G
ZM_A8232/LH244	112	4 / 20		ML542G
ZM_A8234	48	0 / 10	0 / 20	ML542G
ZM_A8246	133	12 / 20	5/5	ML542G
ZM_A8247	148	4 / 20	1/3	SF4404
LH244/ZM_A8247	150		13 / 20	SF4404
ZM_A8248	75	0 / 20	4/6	SF4404
LH244/ZM_A8249	104		13 / 20	SF4404
LH244/ZM_A8249	275		9 / 20	SF4404
LH244/ZM_A8251	148		7 / 20	SF4404
LH244/ZM_A8251	123		5 / 20	SF4404
LH244/ZM_A8251	108		4 / 20	SF4404
ZM_A8252	137	9 / 20	4/6	SF4404
ZM_A8255	78	10 / 20		SM542C
LH244/ZM_A8245	31		2 / 10	ML542G
LH244/ZM_A8253	138		10 / 20	SF542C
LH244/ZM_A8253	67		5 / 20	SF542C
LH244/ZM_A8254	72		8 / 20	SF542C
LH244/ZM_A8254	100		6 / 20	SF542C
LH244/ZM_A8235	81		12 / 20	ABI
LH244/ZM_A8235	116		13 / 20	ABI
LH244/ZM_A8235	161		8 / 20	ABI
ZM_A8237/LH244	294	7 / 20	5/7	ABI
ZM_A8244/LH244	250	4 / 20		ABI
ZM_A8240	14		3/4	ABI
ZM_A8240/LH244	86	7 / 20		ABI
ZM_A8238	31		10 / 10	ABI
LH244/ZM_A8238	90		10 / 20	ABI
ZM_A8256	53	3 / 10		ABI
LH244/ZM_A8256	55		8 / 20	ABI

Example 12**Rhizobia-mediated crop transformation through conjugal transfer system**

An *oriT*-dependent plasmid conjugal transfer system in *Rhizobium* and related species may also be used to deliver a gene of interest (GOI) into plant cells and subsequently be integrated into the plant genome. A homogenous or heterogeneous conjugal transfer system could be used for the gene transfer. Transgenic plants could then be regenerated with selectable markers through an established tissue culture system. Rhizobia strains may include *Sinorhizobium* spp., *Mesorhizobium loti*, *Rhizobium leguminosarum* and *Rhizobium* sp. NGR234, among others.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Patent 4,761,373; U.S. Patent 4,810,648; U.S. Patent 5,013,659; U.S. Patent 5,094,945; U.S. Patent 5,106,739; U.S. Patent 5,107,065; U.S. Patent 5,141,870; U.S. Patent 5,229,114; U.S. Patent 5,273,894; U.S. Patent 5,276,268; U.S. Patent 5,322,938; U.S. Patent 5,352,605; U.S. Patent 5,359,142; U.S. Patent 5,362,865; U.S. Patent 5,378,619; U.S. Patent 5,378,824; U.S. Patent 5,424,412; U.S. Patent 5,463,175; U.S. Patent 5,512,466; U.S. Patent 5,530,196; U.S. Patent 5,543,576; U.S. Patent 5,561,236; U.S. Patent 5,563,055; U.S. Patent 5,591,616; U.S. Patent 5,605,011; U.S. Patent 5,608,149; U.S. Patent 5,627,061; U.S. Patent 5,633,435; U.S. Patent 5,633,437; U.S. Patent 5,637,489; U.S. Patent 5,641,876; U.S. Patent 5,646,024; U.S. Patent 5,659,122; U.S. Patent 5,689,041; U.S. Patent 5,750,871; U.S. Patent 5,750,876; U.S. Patent 5,767,366; U.S. Patent 5,837,848; U.S. Patent 5,850,019; U.S. Patent 5,859,347; U.S. Patent 5,869,720; U.S. Patent 5,958,745; U.S. Patent 5,981,834; U.S. Patent 5,985,605; U.S. Patent 5,998,700; U.S. Patent 6,011,199; U.S. Patent 6,040,497; U.S. Patent 6,051,753; U.S. Patent 6,072,103; U.S. Patent 6,080,560; U.S. Patent 6,140,075; U.S. Patent 6,140,078; U.S. Patent 6,166,292; U.S. Patent 6,171,640; U.S. Patent 6,175,060; U.S. Patent 6,177,611; U.S. Patent 6,225,105; U.S. Patent 6,228,623; U.S. Patent 6,232,526; U.S. Patent 6,252,138; U.S. Patent 6,271,443; U.S. Patent 6,294,714; U.S. Patent 6,380,462; U.S. Patent 6,380,466; U.S. Patent 6,384,301; U.S. Patent 6,414,222; U.S. Patent 6,426,446; U.S. Patent 6,426,447; U.S. Patent 6,429,357; U.S. Patent 6,429,362; U.S. Patent 6,433,252; U.S. Patent 6,437,217; U.S. Patent 6,444,876; U.S. Patent 6,459,018; U.S. Patent 6,476,295; U.S. Patent 6,483,008; U.S. Patent 6,489,461; U.S. Patent 6,495,739; U.S. Patent 6,506,559; U.S. Patent 6,531,648; U.S. Patent 6,537,750; U.S. Patent 6,538,178; U.S. Patent 6,538,179; U.S. Patent 6,538,181; U.S. Patent

6,541,259; U.S. Patent 6,576,818; U.S. Patent 6,589,767; U.S. Patent 6,596,538; U.S. Patent 6,635,806; U.S. Patent 6,613,963; U.S. Patent 6,653,530; U.S. Patent 6,660,849; U.S. Patent 6,706,950; U.S. Patent 6,723,837; U.S. Patent 6,770,465; U.S. Patent 6,774,283; U.S. Patent 6,812,379; U.S. Patent 6,822,141; U.S. Patent 6,828,475; U.S. Patent 7,002,058; U.S. Patent 7,132,528; U.S. Patent 7,151,204; U.S. Patent RE37,543

U.S. Patent Publ. 20020168707; U.S. Patent Publ. 20030028917; U.S. Patent Publ. 20030083480; U.S. Patent Publ. 20030115626; U.S. Patent Publ. 20030135879; U.S. Patent Publ. 20040177399; US Patent Publ. 20040244075; U.S. Patent Publ. 20050005321; U.S. Patent Publ. 20050022261; U.S. Patent Publ. 20050289667; U.S. Patent Publ. 20050289672; U.S. Patent Publ. 20060200878; U.S. Patent Publ. 20060162010; U.S. Patent Publ. 20060236420;

Beringer, *J. Gen. Microbiol.*, 84:188-198, 1974.

Bevan *et al.*, *Nature*, 304:184-187, 1983.

Bird *et al.*, *Biotech Gen. Eng. Rev.*, 9:207-227, 1991.

Bravo-Angel *et al.*, *J. Bacteriol.*, 181:5758-5765, 1999.

Broothaerts *et al.*, *Nature*, 433:629-633, 2005.

Buchanan-Wollaston, *Nature*, 328:172-175, 1987.

Callis *et al.*, *Plant Physiol.*, 88:965-968, 1988.

Carrington and Freed, *J. Virol.*, 64(4):1590-1597, 1990.

Chandler *et al.*, *Plant Cell*, 1:1175-1183, 1989.

Chen *et al.*, *J. Bacteriol.* 184:4838-4845, 2002.

Chu *et al.*, *Scientia Sinica* 18:659, 1975.

Corruzziet *et al.*, *EMBO J.* 3:1671-1679, 1984.

Dekeyser *et al.*, *Plant Physiol.*, 90:217-223, 1989.

Della-Cioppa *et al.*, *Bio/Technology*, 5:579-584, 1987.

Depicker *et al.*, *J. Mol. Appl. Genet.* 1:561-573, 1982.

Dube and Thomson, *Plasmid.*, 50:1-11, 2003.

Dube *et al.*, *Plant Mol. Biol.*, 55(4):531-539, 2004.

Duncan *et al.*, *Planta*, 165:322-332, 1985.

Ebert *et al.*, *Proc. Nat. Acad. Sci. USA* 84:5745-5749, 1987.

- European Appln. 0385 962;
European Appln. 275 957
Farrand *et al.*, *Int. J. Syst. Evol. Microbiol.*, 53:1681-1687, 2003.
Fraley *et al.*, *Bio/Technology*, 3:629-635, 1985.
Fraley *et al.*, *Proc. Natl. Acad. Sci. U S A.*, 80:4803-4807, 1983.
Freiberg *et al.*, *Nature*, 387:394-401, 1997.
Gamborg *et al.*, *Exp. Cell Res.*, 50:151, 1968.
Garg *et al.*, *Appl. Env. Microbiol.*, 65:2802-2804, 1999.
Gibson and Shillitoe, *Mol. Biotech.* 7:125-137, 1997.
Hiei *et al.*, *Plant J.*; 6:271-282, 1994.
Hiei *et al.*, *Plant. Mol. Biol.*, 35(1-2):205-218, 1997.
Hoekema *et al.*, *Nature*, 303:179, 1983.
Hooykaas *et al.* *J. Gen. Microbiol.* 98:477-484, 1977.
Ishida *et al.*, *Nature Biotechnol.*, 14:745-750, 1996.
Komari *et al.*, *Curr. Opin. Plant Biol.*, 1:161-165, 1998.
Kuhlemeier *et al.*, *Plant Cell*, 1:471-478, 1989.
Lawton *et al.*, *Plant Mol. Biol.* 9:315-324, 1987.
Lazo *et al.*, *Bio/Technol.*, 9:963-967, 1991.
Linsmaier and Skoog, *Physiol. Plant.*, 18:100, 1965.
Long, *Plant Physiol.*, 125:69-72, 2001.
Marcotte *et al.*, *Plant Cell*, 1:969-976, 1989.
McCormac *et al.*, *Euphytica*, 99(1):17-25, 1998.
McCown and Lloyd, *Hort. Science*, 16:453, 1981.
Miki and McHugh, *J. Biotechnol.* 107 193-232, 2004.
Murashige and Skoog, *Physiol. Plant*, 15:473-497, 1962.
Nitsch and Nitsch, *Science*, 163:85-87, 1969.
Noel *et al.*, *Can. J. Microbiol.* 42:279-283, 1996.
Odell *et al.*, *Nature* 313:810-812, 1985.
Okita *et al.*, *J Biol Chem.* 264:12573, 1989.
PCT Appln. WO 87/04181; PCT Appln. PCT Appln. WO 89/00193; PCT Appln. WO
99/027116; PCT Appln. WO 04/009761; PCT Appln. WO 04/074443; PCT
Appln. WO 05/003362
Perez-Mendoza *et al.*, *J. Bacteriol.*, 186(17):5753-5761, 2004.

- Radke *et al.*, *Plant Cell Rep.* 11:499-505, 1992.
- Rogers *et al.*, *Methods Enzymol.* 153:253-277, 1987.
- Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, 1989
- Schaffner *et al.*, *Plant Cell*, 3:997-1012, 1991.
- Schenk and Hildebrandt, *Can. J. Bot.*, 50:199-204, 1972.
- Schrammeijer *et al.*, *Plant Cell Reports*, 9: 55-60, 1990.
- Shadenkov *et al.*, *Mol. Biol.*, 30:272-275, 1996.
- Southern, *Mol. Biol.*, 98:503-517, 1975
- Spaink, H. P. *et al.*, (ed.), *The Rhizobiaceae*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1998.
- Stewart & Hsu, *Planta*, 137:113-117, 1977.
- Tingay *et al.*, *Plant J.*, 11(6):1369-1376, 1997.
- Tun-Garrido *et al.*, *J. Bacteriol.*, 185(5):1681-1692, 2003.
- Turner *et al.*, *FEMS Microbiol. Ecol.*, 42(2):227-234, 2002.
- Uchimiya and Murashige, *Plant Physiol.*, 15:473, 1962.
- Walker *et al.*, *Proc. Natl. Acad. Sci. USA*, 84:6624, 1987.
- Weller *et al.*, *Pl. Pathol.* 54:799-805, 2005.
- Weller *et al.*, *Appl. Env. Microbiol.*, 70:2779-2785, 2004.
- Wuni *et al.*, *Plant Cell*, 1:961-968, 1989.
- Yang *et al.* *Proc. Natl. Acad. Sci. USA*, 87:4144-4148, 1990.
- Zhang and Meyer, *Mol. Microbiol.*, 25:509-516, 1997.
- Zhou *et al.*, *Plant Cell Rep.*, 15:159-163, 1995.

CLAIMS

1. A method for transforming a corn plant cell, comprising:
 - (a) contacting at least a first corn plant cell with a Rhizobiales bacterium other than *Agrobacterium* sp, comprising: (i) a first nucleic acid comprising a *vir* gene region of a Ti plasmid wherein the *vir* gene region acts to introduce a nucleic acid of interest into the plant cell in a VirD2-dependent manner; and (ii) a second nucleic acid comprising one or more T-DNA border sequence(s) operably linked to a nucleic acid of interest; and
 - (b) selecting at least a first corn plant cell transformed with the nucleic acid of interest.
2. The method of claim 1, wherein the bacterium is a Rhizobia cell.
3. The method of claim 2, wherein the Rhizobia cell is grown in the presence of acetosyringone or other compound that induces *vir* gene function prior to contacting the plant cell.
4. The method of claim 2, wherein the Rhizobia cell is selected from the group consisting of: *Rhizobium* spp., *Sinorhizobium* spp., *Mesorhizobium* spp., *Phyllobacterium* spp. *Ochrobactrum* spp. and *Bradyrhizobium* spp.
5. The method of claim 4, wherein the Rhizobia cell is *Rhizobium leguminosarum*.
6. The method of claim 5, wherein the Rhizobia cell is *R. leguminosarum* *bv. trifolii*, *R. leguminosarum* *bv. phaseoli* or *Rhizobium leguminosarum. bv. viciae*.
7. The method of claim 1, wherein the plant cell is comprised in an explant from a plant seed, seedling, callus, cell suspension, cotyledon, meristem, leaf, root, or stem; and the explant is contacted with the bacterium.
8. The method of claim 7, wherein the explant comprises an embryonic meristem; callus; cell suspension; cotyledon; or tissue from leaves, roots, or stems.

9. The method of claim 1, wherein the first and second nucleic acids are introduced into the bacterium by electroporation.
10. The method of claim 1, wherein selecting a plant cell transformed with the nucleic acid of interest is carried out in the absence of a selection agent.
11. The method of claim 1, wherein selecting a plant cell transformed with the nucleic acid of interest comprises culturing the plant cell in the presence of a selection agent, wherein the nucleic acid of interest confers tolerance to the selection agent or is operably linked to a further nucleic acid that confers tolerance to the selection agent.
12. The method of claim 11, wherein the selection agent is glyphosate, kanamycin, bialaphos or dicamba.
13. The method of claim 12, wherein the nucleic acid of interest or further nucleic acid encodes EPSP synthase.
14. The method of claim 13, wherein the EPSP synthase protein is CP4.
15. The method of claim 11, wherein the selection agent is glyphosate.
16. The method of claim 1, wherein the nucleic acid of interest is not physically linked to a selectable marker gene.
17. The method of claim 16, wherein the marker gene and the nucleic acid of interest genetically segregate in progeny of a plant regenerated from the plant cell transformed with the nucleic acid of interest.

18. The method of claim 1, wherein the bacterium comprises at least a third nucleic acid comprising a further nucleic acid of interest and wherein the plant cell is transformed with the third nucleic acid.


19. The method of claim 1, further comprising regenerating a plant from the plant cell, wherein the plant comprises the nucleic acid of interest.

20. The method of claim 19, wherein regenerating a plant from the plant cell comprises inducing formation of one or more shoots from an explant comprising the plant cell and cultivating at least a first shoot into a whole fertile plant.

21. The method of claim 19, wherein regeneration occurs by organogenesis.

22. The method of claim 1, further comprising growing the bacterium other than *Agrobacterium* sp. in a medium comprising carbon source(s) which minimize polysaccharide production during growth in induction medium, wherein the carbon source(s) used to minimize polysaccharide production during growth in induction medium is glucose in AB-TY medium, or L-arabinose and potassium gluconate in ATA medium..

FIG. 1







Fig. 4



RL4404: *R. leguminosarum* strain Madison with pTi4404kan
 ML542G: *M. loti* USDA3471 with pTiBo542G
 ML4404: *M. loti* USDA3471 with pTi4404kan
 2370LBA: *R. leguminosarum* usda2370 with pTi4404kan
 2370G: *R. leguminosarum* USDA2370 with pTiBo542G
 SF4404: *S. fredii* USDA205 with pTi4404kan
 SM542C: *S. meliloti* USDA1002 with pTiBo542G
 ABI: *A. tumefaciens* ABI strain control

FIG. 5




FIG. 6




FIG. 7





FIG. 8



9 / 17

FIG. 9




FIG. 10



10 / 17




FIG. 11





FIG. 12



13 / 17

FIG. 13






FIG. 14

FIG. 15






FIG. 16

A) SF4404/pMON96036



B) ML542G/pMON96036



C) SM542C/pMON96036





FIG. 17



mons100wo-seq-listing.txt
SEQUENCE LISTING

<110> YE, XUDONG
SHEN, JUNJIANG
WILLIAMS, EDWARD

<120> USE OF NON-AGROBACTERIUM BACTERIAL SPECIES FOR PLANT
TRANSFORMATION

<130> MONS:100US

<140> UNKNOWN

<141> 2007-05-16

<150> 60/800,872

<151> 2006-05-16

<160> 30

<170> PatentIn Ver. 2.1

<210> 1

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 1

gagagtttga tcctggctca g

21

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 2

aaggaggtga tccagccgca g

21

<210> 3

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 3

acaataatgt gtgttgtaa gtcttggtgc

30

<210> 4

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Primer

<400> 4
ctcaaacctactcaatat ttggtgag 28

<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 5
agatctggct cgcgccggac gcac 24

<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 6
cgctcgcgct attctttgct ggag 24

<210> 7
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 7
tcagcaggat gacgccgta tcg 23

<210> 8
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 8
tctcgccga ccaatacaa cac 23

<210> 9
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
Primer

<400> 9
gctgacgggc ccggatgaat gtcagctact g 31

<210> 10
 <211> 42
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 10
 gctctagaaa ttgtaagcgt taataattca gaagaactcg tc 42

 <210> 11
 <211> 28
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 11
 caattgcatt tggctcttaa ttatctgg 28

 <210> 12
 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 12
 gcatgcccga tcgcgctcaa gtaatc 26

 <210> 13
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 13
 tctaggtccc cccgccccca tcg 23

 <210> 14
 <211> 28
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 14
 ccatggatct ttctggcaat gagaaatc 28

 <210> 15

<211> 28
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 15
 gtcaaaagct gttgacgctt tggctacg 28

 <210> 16
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 16
 acgggagagg cgggtgtagt tgc 23

 <210> 17
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 17
 cgatagcgac aatgccgaga acg 23

 <210> 18
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 18
 atgcccgatc gagctcaagt tatc 24

 <210> 19
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic
 Primer

 <400> 19
 tgaaaggaca cctctccggt gctg 24

 <210> 20
 <211> 27
 <212> DNA
 <213> Artificial Sequence

```

<220>
<223> Description of Artificial Sequence: Synthetic
      Primer

<400> 20
ccatggatcc gaaggccgaa ggcaatg                27

<210> 21
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
      Primer

<400> 21
ctacagactg tttacggttg ggc                    23

<210> 22
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
      Primer

<400> 22
gtgagcaaag ccgctgccat atc                    23

<210> 23
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial sequence: Synthetic
      Primer

<400> 23
tagagcgtct gcttggttaa acc                    23

<210> 24
<211> 1490
<212> DNA
<213> A. tumefaciens

<400> 24
tagaaaggag gtgatccagc cgcaggttcc cctacggcta ccttgttacg acttcacccc 60
agtcgctgac cctaccgtgg ttagctgcct ccttgcggtt agcgcactac cttcgggtaa 120
aaccaactcc catggtgtga cgggcggtgt gtacaaggcc cggaacgta ttcaccgcag 180
catgctgatc tgcgattact agcgattcca acttcatgca ctcgagttgc agagtgcaat 240
ccgaactgag atggcttttg gagattagct cgacatcgct gtctcgctgc ccaactgtcac 300
caccattgta gcacgtgtgt agcccagccc gtaagggcca tgaggacttg acgtcatccc 360
caccttcctc tcggcttata accggcagtc cccttagagt gcccaactaa atgctggcaa 420
ctaagggcga gggttgcgct cgttgcggga cttaaccxaa catctcacga cacgagctga 480
cgacagccat gcagcacctg ttctggggcc agcctaactg aaggacatcg tctccaatgc 540
ccataccccg aatgtcaaga gctggtaagg ttctgcgctg tgcttcgaat taaccacat 600
gctccaccgc ttgtgcgggc ccccgtaact tcctttgagt ttaaatcttg cgaccgtact 660
ccccaggcgg aatgtttaat gcgtttagctg cgccaccgaa cagtatactg cccgacggct 720
aacattcatc gtttacggcg tggactacca gggtatctaa tcctgtttgc tccccacgct 780
ttcgcacctc agcgtcagta atggaccagt aagccgcctt cgccactggt gttcctccga 840

```

mons100wo-seq-listing.txt

```

atatctacga atttcacctc tacactcggg attccactta cctcttccat actcaagata 900
cccagtatca aaggcagttc cgcagttgag ctgcgggatt tcaccctga cttaaatc 960
cgcttacgtg cgctttacgc ccagtaattc cgaacaacgc tagccccctt cgtattaccg 1020
cggctgctgg cacgaagtta gccggggctt cttctccgac taccgtcatt atcttcatcg 1080
gtgaaagagc tttaacaacc taaggccttc atcactcacg cggcatggct ggatcaggct 1140
tgcgcccatt gtccaatatt ccccactgct gcctcccgta ggagtttggg ccgtgtctca 1200
gtcccaatgt ggctgatcat cctctcagac cagctatgga tcgtcgcctt ggtaggcctt 1260
taccaccaca actagtaaat ccaacgcggg ctcatccatc cccgataaat ctttcccccg 1320
tagggcgat tagcgtattaa ttccagtttc ccagagctat tccgcagaga tgggtagatt 1380
cccacgcggt actcacccgt ctgccactcc ctttgcgggg cgttcgactt gcatgtgtta 1440
agcctgccgc cagcgttcgt tctgagccag gatcaaacct tcaagttgag 1490

```

<210> 25
 <211> 1366
 <212> DNA
 <213> Rhizobium leguminosarum

```

<400> 25
cttcgccag tcgctgacc kascgkgggt agctgsctcc kkgcggttag cgcactacct 60
tcgggtaaaa ccaactccca tgggtgtgag ggcgggtgtg acaaggcccg ggaacgtatt 120
caccgcggca tgctgatccg cgattactag cgattccaac ttcatgcact cgagttgcag 180
agtgaatcc gaactgarat ggcttttggg gattagctca cactcgcgtg ctcgctgccc 240
actgtcacca ccattgtagc acgtgtgtag cccagcccgt aagggccatg aggacttgac 300
gtcatmcca cttctctc ggcttatcac cggcagtccc cttagagtgc ccaactgaat 360
gttggaact aaggcgagg gttgcgctc ttgcgggact taaccaaca tctcacgaca 420
cgagctgacg acagccatgc agcacctgtg tcccggctcc cgaagggaac cttgcatctc 480
tgcaagtagc cgggcatgtc aagggtgtg aaggttctgc gcgttgcttc gaattaaacc 540
acatgctcca ccgcttgctc gggccccctg caattccttt gagttttaat cttgcgaccg 600
tactccccag gcggaatgtt taatgcgta gctgcgccac cgaacagtat actgcccgac 660
ggtaaacatt catcgtttac ggcgtggact accagggtat ctaatcctgt ttgctcccc 720
cgctttcgca cctcagctc agtaatggac cagtgcgccc ccttcgccac tgggttctct 780
ccgaatatct acgaatttca cctctacact cgggaattcca ctcacctctt ccatactcca 840
gatcgacagt atcaaaggca gttccagggt tgagccctgg gatttcaccc ctgactgatc 900
gatccgccta cgtgcgcttt acgcccagta attccgaaca acgctagccc ccttcgtatt 960
accgcggctg ctggcacgaa gttagccggg gcttcttctc cggataccgt cattatcttc 1020
tccggtgaaa gagctttaca acctagggc cttcatcact cacgcggcat ggcgtgata 1080
ggcttgcgcc cattgtccaa tattccccac tgctgcctcc cgtaggagt 1140
ctcagtccca atgtggctga tcctctctc agaccagcta tggatcgtcg ctttggtagg 1200
cctttacccc accaactagc taatccaacg cgggctcatc cttgaccgat aaatcttct 1260
cccgaaggac acatacggt ttagcacaag tttccctgct ttattccgta gtcaagggta 1320
gattcccacg cgttactcac ccgtctgccc ctccccctgc ggggcg 1366

```

<210> 26
 <211> 1323
 <212> DNA
 <213> Rhizobium leguminosarum

```

<400> 26
agcgcactac ctctcgggta aaaccaactc ccatggtgtg acgggcggtg tgtacaaggc 60
ccgggaacgt attcaccgcg gcatgctgat ccgcgattac tagcgattcc aacttcatgc 120
actcgagttg cagagtgcaa tccgaactga gatggctttt ggagattagc tcacactcgc 180
gtgctcgctg cccactgtca ccaccattgt agcacgtgtg tagcccagcc cgtaagggcc 240
atgaggactt gagctawca ccaccttct ctcggcttat caccggcagt ccccttagag 300
tgcccaactg aatgctggca actaagggcg agggttgctc tcgttgcggg acttaacca 360
acatctcacg acacgagctg acgacagcca tgcagacct gtgtcccggg ccccgaggg 420
accttgcatc tctgcaagta gcgggcatgt caagggtggt taaggttctg cgcgttgctt 480
cgaattaaac cacatgctcc accgcttgtg cgggcccccg tcaattcctt gagttttaat 540
cttgcgaccg tactccccag gcggaatgtt taatgcgta gctgcgccac cgaacagtat 600
actgcccagc ggtaaacatt catcgtttac ggcgtggact accagggtat ctaatcctgt 660
ttgctcccc  cgctttcgca cctcagctc agtaatggac cagtgcgccc ccttcgccac 720
tgggtgttct ccgaatatct acgaatttca cctctacact cgggaattcca ctcacctctt 780
ccatactcca gatcgacagt atcaaaggca gttccagggt tgagccctgg gatttcaccc 840
ctgactgatc gatccgccta cgtgcgcttt acgcccagta attccgaaca acgctagccc 900
ccttcgtatt accgcggctg ctggcacgaa gttagccggg gcttcttctc cggataccgt 960
cattatcttc tccggtgaaa gagctttaca acctagggc cttcatcact cacgcggcat 1020
ggctkgatca ggcttgcgcc cattgtccaa tattccccac tgctgcctcc cgtaggagt 1080

```

mons100wo-seq-listing.txt

```
tgggccgtgt ctcagtccca atgtggctga tcatccttc agaccagcta tggatcgctg 1140
ccttggtagg cctttacccc accaactagc taatccaacg cgggctcatc cttgaccgat 1200
aatctttct cccgaaggac acatacggta ttagcacaag tttccctgcg ttattccgta 1260
gtcaagggta gattcccacg cgttactcac ccgtctgccg cttccccttg cggggcgcat 1320
acg 1323
```

<210> 27
 <211> 1341
 <212> DNA
 <213> Rhizobium leguminosarum

```
<400> 27
ggtagctgc ctcttgccg ttagcgcact accttcgggt aaaaccaact cccatgggtg 60
gacgggagg gtgtacaagg cccgggaacg tattcaccgc ggcagtgctga tccgcgatta 120
cgaccgattc caacttctag cactcgagtt gcagagtga atccgaactg agatggcttt 180
tggagattag ctacactcg cgtgctcgct gccactgtc accaccattg tagcacgtgt 240
gtagcccagc ccgtaagggc catgaggact tgacgtcatc cccaccttcc tctcggctta 300
tcaccggcag tccccttaga gtgcccaact gaatgctggc aactaagggc gagggttgcg 360
ctcgttgccg gacttaacc aacatctcac gacacgagct gacgacagcc atgcagacc 420
tgtgtcccgg tccccgaagg gaaccttgca tctctgcaag tagccgggca tgtcaagggc 480
tggtaaagtt ctgctgctg cttcraatta aaccacatgc tccaccgctt gtgcgggccc 540
ccgtcaattc ctttgagttt taatcttgcg accgtactcc ccaggcggaa tgtttaatgc 600
gtagctgctg ccaccgaaca gtatactgcc cgacggctaa cattcatcgt ttacggcgctg 660
gactaccagg gtatctaate ctgtttgctc cccacgcttt cgcacctcag cgtcagtaat 720
ggaccagtga gccgccttcg ccactgggtg tcctccgaat atctacgaat ttcacctcta 780
cactcggaat tccactcacc tcttccatac tccagatcga cagtatcaaa ggcagttcca 840
gggttgagcc ctgggatttc acccctgact gatcgatccg cctacgtgcg ctttacgccc 900
agtaattccg aacaacgcta gcccccttcg tattaccgcg gctgctggca cgaagttagc 960
cggggcttct tctccggata ccgtcattat cttctccggt gaaagagctt tacaacccta 1020
gggcttcat cactcacgcy gcatggcttg atcaggcttg cgcccattgt ccaatattcc 1080
ccactgctgc ctcccgtagg agtttgggcc gtgtctcagt cccaatgtgg ctgatcatcc 1140
tctcagacca gctatggatc gtcgcttgg taggccttta ccccaccaac tagctaatec 1200
aacgcgggcc gatcctttac cgataaatct ttccccgaa gggcacatac ggtattagca 1260
caagtttccc tgcgttattc cgtagtaaag ggtacgttcc cacgcgttac tcaccctctc 1320
gccgctcccc ttgctgaggc c 1341
```

<210> 28
 <211> 1341
 <212> DNA
 <213> Mesorhizobium loti

```
<400> 28
ccttgcgggt agcacagcgc cttcgggtaa aaccaactcc catggtgtga cgggcgggtg 60
gtacaaggcc cgggaacgta ttcaccgcyg catgctgata cgcgattact agcgattcca 120
acttcatgca ctcgagttgc agagtgcaat ccgaactgag atggcttttg gagattagct 180
cgacctcgcg gtctcgtcgc ccactgtcac caccattgta gcaggtgtgt agcccagccc 240
gtaagggcca tgagacttg acgtcatccc caccttctc tcggcttata accggcagtc 300
cccttagagt gcccaactga atgtggcaa ctaagggcga gggttgcgct cgttgcggga 360
cttaacccaa catctcacga cacgagctga cgacagccat gcagcacctg tcaccggtcc 420
agccgaactg aaggtatcca tctctgaaa ccgcgaccgg gatgtcaagg gctggtaagg 480
tctgctcgtg tgcttcaaat taaccacat gctccaccgc ttgtgcgggc ccccgtcaat 540
tcctttgagt ttaaatcttg cgaccgtact ccccaggcgg gaagcttaat gcgttagctg 600
cgccaccgac aagtaaactt gccaacggct agcttccatc gtttacggcg tggactacca 660
gggtatctaa tctgtttgc tccccacgct ttgcacctc agcgtcagta ccggaccagt 720
gagccgcctt cgccactggt gttcctccga atatctacga atttcacctc tactctcggg 780
attccactca cctcttccgg actcgagata cccagtatca aaggcagttc cggggttgag 840
ccccgggatt tcacccttga cttaaagtac cgcctacgtg cgctttacgc ccagtaattc 900
cgaacaacgc tagccccctt cgtattaccg cgggtgcttg cacgaagtta gccggggctt 960
cttctacggt atcttcaccg atcttcaccg ttgaaagagc tttacaacc tagggccttc 1020
atcactcacg cggcatggct ggatcaggct tgcgccatt gtccaatatt ccccactgct 1080
gcctcccgta ggagtctggg ccgtgtctca gtcccagttg ggctgatcat cctctcagac 1140
cagctatgga tctctgcctt ggtaggccat taccaccaca actagctaate ccaacgcggg 1200
ctcatccatc tccgataaat ctttctccaa aaggacgtat acggtattag ctccagtttc 1260
ccggagttgt tccgtagaga tgggtagatt cccacgcgtt actcaccctg ctgccgctcc 1320
ccttgcgggg cacgcgcctt t 1341
```

mons100wo-seq-listing.txt

<210> 29
 <211> 1339
 <212> DNA
 <213> S. Fredii

<400> 29
 agccttgagg ttagcgcact accttcgggt agaaccaact cccatggtgt gacgggcggt 60
 gtgtamaagg cccgggaacg tattcaccgc agcatgctga tctgcgatta ctagcgattc 120
 caacttcatg cactcgagtt gcagagtgca atccgaactg agatggcttt tggagattag 180
 ctcgacctcg cggctcgcet gcccactgtc accaccattg tascacgtgt gtagcccagc 240
 ccgtaagggc catgaggact tgacgtcatc cccaccttc tctcggctta tcaccggcag 300
 tccccctaga gtgaccaact aaatgctggc aactaagggc gaggggttgcg ctcggttgcgg 360
 gacttaacc aacatctcac gacacgagct gacgacagcc atgcagcacc tgtctccgat 420
 ccagccgaac tgaaggatac gatctctcgt atccgcgacg gggatgtcaa gggctggtaa 480
 ggttctgctc gttgcttcga attaaaccac atgctccacc gcttgtgctg gcccccgta 540
 attcctttga gttttaatct tgcgaccgta ctccccaggc ggaatgttta atgctgtagc 600
 tgcgccaccg aacagtaaac tgcccagcgg ctaacattca tcgtttacgg cgtggactac 660
 cagggatctt aatcctgttt gctccccacc ctttcgcacc tcagcgtcag taccggacca 720
 gtgagccgct ttcgccactg gtgttcctcc gaatatctac gaatttcacc tctacactcg 780
 gaattccact cacctcttcc ggactctaga caccagat caaaggcagt tccggggttg 840
 agccccggga tttaccacct gacttaaatg tccgcctacg tgcgctttac gccagtaat 900
 tccgaacaac gtagcccc ttcgtattac cgcggtgctt ggcacgaagt tagccggggc 960
 ttcttctccg gttaccgtca ttatcttcac cggtgaaaga gctttacaac ctagggcct 1020
 tcatcactca cgcggcatgg ctggatcagg ctgtgcacca ttgtccaata tccccactg 1080
 ctgcttccc taggagtttg ggccgtgtct cagtcccaat gtggctgatc atcctctcag 1140
 accagctatg gatcgtcgc ttggtaggcc tttaccacc caactagcta atccaacgcg 1200
 ggctcatcct tccccgataa atctttcccc cgaagggctc atacggtatt agcacaagtt 1260
 tccctgcgtt attccgtaga aaagggtaga ttcccacgcg ttactcaccm atctgccgcy 1320
 cccccgcrgc gcgctccc 1339

<210> 30
 <211> 1368
 <212> DNA
 <213> Bradyrhizobium sp.

<400> 30
 gaaggtaccg tggccggctg cctacctagc gggttagcgc accgtcttca ggtaaaacca 60
 actcccattg tgtgacgggc ggtgtgtaca agggccggga acgtattcac cgtggcgtgc 120
 tgatccacga ttactagcga ttccaacttc atgggctcga gttgcagagc ccaatccgaa 180
 ctgagacggc tttttgagat ttgcgaaggg tcgcccctta gcatccatt gtcaccgcca 240
 ttgtagcac tgtgtagccc agcccgtaag ggccatgagg acttgacgtc atccccacct 300
 tctcgcggc ttatcaccgg cagtctcctt agagtgtca actaaatggt agcaactaag 360
 gacggggggt gcgctcgttg cgggacttaa cccaacatct cacgacacga gctgacgaca 420
 gcatgcagca cctgtgctcc aggtccgaa gagaaggtea catctctgct accggctctg 480
 gacatgtcaa gggctggtaa ggttctgctc gttgcgtcga attaaaccac atgctccacc 540
 gcttgtgctg gccccgtaa attccttgag ttttaattct gcgaccgtac tccccaggcg 600
 gaatgcttaa agcgttaget gcgccactag tgagtaaac cactaacggc tggcattcat 660
 cgtttacggc gtggactacc agggatctta atctgtttg ctccccacgc tttcgtgcct 720
 cagcgtcagt atcgggccag tgagccgctc tcgccactgg tgttcttgct aatatctacg 780
 aattcacct ctacactcgc agttccactc acctctccc aactcaagat cctcagtatc 840
 aaaggcagtt ctggagttga gctccaggat ttcaccctg acttaaagac ccgcctacgc 900
 accctttacg cccagtgatt ccgagcaacg ctagccccct tcgtattacc gcggctgctg 960
 gcacgaagtt agccggggct tattcttgct gtaccgtcat tatcttccc cacaaaagag 1020
 cttacaacc ctagggcctt catcactcac gcggcatggc tggatcaggg ttgccccat 1080
 tgccaatat tccccactgc tgcttcccgt aggagtttgg gccgtgtctc agtcccaatg 1140
 tggctgatca tctctcaga ccagctactg atcgtcgcct tggtagacca ttacctacc 1200
 aactagctaa tcagacgagg gccgatcttt cggcgataaa tctttcccc taagggtta 1260
 tccggtatta gcacaagttt cctgtgttg ttccgaacca aaaggtacgt tcccacgcgt 1320
 tactacccc tctgcccgtg acgtatgctw cgcccgcgcg cckcct 1368